

GREAT BASIN NATURALIST MEMOIRS

Number 13

Brigham Young University

1992



A Catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index Volume B



HARVARD UNIVERSITY



Library of the
Museum of
Comparative Zoology

GREAT BASIN NATURALIST MEMOIRS


Number 13

Brigham Young University

1992



A Catalog of Scolytidae and Platypodidae (Coleoptera), Part 2: Taxonomic Index Volume B



LIBRARY
OF BRIGHAM YOUNG UNIVERSITY
1992

MCZ
LIBRARY

NOV 15 1993

HARVARD
UNIVERSITY

Copyright © 1992 by Brigham Young University. All rights reserved.
Printed in the United States of America.

ISBN 0-8425-2310-3

CONTENTS

Part 2, Volume A

ABSTRACT	1
INTRODUCTION (English)	1
INTRODUCTION (French)	7
EINLEITUNG (German)	13
INTRODUCCION (Spanish)	19
LIST OF MUSEUM TYPE REPOSITORIES	25
KEY TO SUBJECT AREAS	27
FAMILY SCOLYTIDAE	28
SUBFAMILY HYLESININAE	28
Tribe Hylastini	29
Tribe Hylesinini	64
Tribe Tomicini	103
Tribe Phrixosomini	189
Tribe Hyorrhynchini	191
Tribe Diamerini	193
Tribe Bothrosternini	205
Tribe Phloeotribini	217
Tribe Phloeosomini	236
Tribe Hypoborini	271
Tribe Polygraphini	283
SUBFAMILY SCOLYTINAE	313
Tribe Scolytini	313
Tribe Ctenophorini	383
Tribe Scolytoplatypodini	401
Tribe Micracini	408
Tribe Cactopinini	435
Tribe Carphodicticini	437
Tribe Ipinini	438
Tribe Dryocoetini	539
Tribe Crypturgini	615
Tribe Xyloterini	632
Tribe Xyleborini	651

Part 2, Volume B

Tribe Xyloctonini	835
Tribe Cryphalini	842
Tribe Corthylini	949
Subtribe Pityophthorina	949
Subtribe Corthylina	1034
GENERA INCERTAE SEDIS	1082
FAMILY PLATYPODIDAE	1084
SUBFAMILY COPTONOTINAE	1084
Tribe Coptonotinae	1084
Tribe Mecopelmini	1086
Tribe Schedlarini	1087
SUBFAMILY TESSEROCERINI	1088
Tribe Diapodini	1088
Tribe Tesserocerini	1097
SUBFAMILY PLATYPODINAE	1113
Tribe Platypodini	1113
HOST LIST AND SPECIES REPORTED TO ATTACK THEM	1241
SUPPLEMENT TO THE 1987 BIBLIOGRAPHY	1349
ADDENDUM	1440
INDEX FOR PLATYPODIDAE	1441
INDEX FOR SCOLYTIDAE	1460

Tribe Xyloctonini Eichhoff

Xyloctonidae

References: Eichhoff 1878b: 171.

Xyloctoninae

References: Hopkins 1915c: 224.

Xyloctonini

References: Wood, S. L. 1978a: 114, 1982b: 63, 1986a: 84–85.

Genus *Cryphalomimus* Eggers

CRYPHALOMIMUS EGGER 1927a: 174. Type-species:

Cryphalomimus striatus Eggers, monobasic.

References: (ay) Nobuchi 1969a: 55. (hb) Wood, S. L. 1986a: 85. (ds) Wood, S. L. 1986a: 85. (tx) Eggers 1927a: 174, 1932c: 23; Schedl 1939i: 379, 1961k: 441; Wood, S. L. 1986a: 85.

ater Numberg 1952: 23. Holotype, sex?; Repub. of the Congo: Garamba Nat. Park; Inst. Parcs Nationaux du Congo Collection.

Distribution: Africa (Zaire).

References: (ds) Numberg 1952: 23. (tx) Numberg 1952: 23, 1965b: pl. 1.

grandis Schedl 1971e: 10. Holotype, sex?; Ehemaliger Kongo Belge, Yangambi; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

References: (tx) Schedl 1971e: 10, 1979c: 108.

striatus Eggers 1927a: 174. Holotype, sex?; Ost-Africa (Muansa); Methner Collection.

Distribution: Africa (Tanzania/Zaire).

References: (ds) Schedl 1961k: 442. (tx) Eggers 1927a: 174; Schedl 1939i: 379, 1961k: 442.

Genus *Glostatus* Schedl

GLOSTATUS SCHEDL 1939i: 386. Type-species: *Glostatus declividepressus* Schedl, monobasic.

Ctonocryphus Schedl 1941d: 398. Type-species:

Ctonocryphus xyloctonus Schedl, monobasic.

Synonymy: Wood 1986: 85.

References: (tx) Schedl 1941d: 398, 1957e: 865–883, 1961k: 638.

Apoglostatus Schedl 1957b: 155. Type-species:

Apoglostatus acaciae Schedl, monobasic. Synonymy: Schedl 1964m: 305.

References: (tx) Schedl 1957b: 155, 1961k: 642, 1964m: 305.

Paraglostatus Schedl 1964m: 304. Type-species:

Ctonocryphus uigricestis Schedl, original designation. Synonymy: Wood 1984b: 225.

References: (tx) Schedl 1964m: 304; Wood, S. L. 1984b: 225.

Rhopalocryphus Numberg 1967: 320. Type-species: *Rhopalocryphus seydeli* Numberg, monobasic. Synonymy: Schedl 1980d: 117.

References: (tx) Numberg 1967: 320; Schedl 1980d: 117.

References: (hb) Wood, S. L. 1986a: 85. (ds)

Wood, S. L. 1986a: 85. (tx) Schedl 1939i: 386,

1957b: 155, 1958a: 557–560, 1961k: 638–639, 1964m: 305; Wood, S. L. 1986a: 85.

acaciae (Schedl) 1957b: 155 (*Apoglostatus*). Holotype, sex?; Muguga, Kenya, and Londiani, Kenya; BMNH, London.

Distribution: Africa (Kenya/Tanzania).

Hosts: *Acacia decurrens*, A. sp., *Ficus* sp.,

Juniperus procera, *Maba* sp., *Rhus* sp.

Notes: (1) Schedl 1964m: 305 (to *Glostatus*).

References: (ds) Gardner 1957: 30; Schedl 1961k: 642. (tx) Schedl 1957b: 150, 155, 1961k: 642–643, 1964m: 305, 1979c: 9.

assimilis Schedl 1957e: 871. Holotype, sex?; Kenya, Jilore; BMNH, London.

Distribution: Africa (Kenya).

References: (ds) Browne 1973a: 279; Schedl 1961k: 640. (tx) Powell, W. 1980: 29; Schedl 1957e: 871, 1958a: 558, 1959c: 705, 1961k: 640, 1979c: 29.

bispinosus Beaver 1955: 77. Holotype ♂; Zambia: Kitwe, Chati; BMNH, London.

Distribution: Africa (Zambia).

References: (hb) Loytyniemi, Beaver, & Loytyniemi 1984. (tx) Beaver 1955: 77.

carinifer Schedl 1957e: 871. Holotype, sex?; Tanganyika, Lushoto; BMNH, London.

Distribution: Africa (Tanzania).

Hosts: *Gologolo* sp., *Syzygium* sp.

References: (ds) Schedl 1961k: 640. (tx) Schedl 1957e: 871, 1961k: 640.

declividepressus Schedl 1939i: 386. Holotype, sex?; Usambara, Ost-Afrika; Schedl Collection in NHMW, Wien.

Distribution: Africa (Kenya/Tanzania).

References: (tx) Schedl 1939i: 386, 1957b: 154, 1961k: 641, 1979c: 76.

delicatus (Schedl) 1962k: 1072 (*Paraglostatus*). Holotype, sex?; Zululand; Eshowe; Schedl Collection in NHMW, Wien.

Distribution: Africa (South Africa).

References: (ds) Schedl 1962k: 1072, 1965h: 111, 1975k: 277.

gracilior (Schedl) 1957e: 872 (*Apoglostatus*). Holotype, sex?; Shagai, Tanganyika; BMNH, London.

Distribution: Africa (Tanzania).

Notes: (1) Schedl 1964m: 305 (to *Glostatus*).

References: (tx) Schedl 1957e: 872, 1961k: 644, 1964m: 305, 1979c: 107.

hirsutus Schedl 1962k: 1070. Holotype, sex?; D.O.A.: Usambara; Schedl Collection in NHMW, Wien.

Distribution: Africa (Kenya).

References: (tx) Schedl 1962k: 1070, 1979c: 117.

kenyae Schedl 1957b: 154. Lectotype, sex?; Kenya, Forest de Nairobi, 1700 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 132.

Distribution: Africa (Kenya).

References: (tx) Schedl 1957b: 150, 154, 1957e: 871, 1961k: 641, 1979c: 131–132.

leprosus Browne 1973a: 290. Holotype ♀; Nigeria: Adamawa, Cangoro Forest Reserve at 4000 feet; MRCB, Tervuren.

Distribution: Africa (Nigeria/ Zaire).

References: (tx) Browne 1973a: 290.

multispinosus (Schedl) 1957e: 873 (*Ctonocryphus*). Holotype, sex?; Kenya, Kilifi; BMNH, London.

Distribution: Africa (Kenya/ Zambia).

Hosts: *Julbernardia paniculata*.

Notes: (1) Schedl 1964n: 305 (to *Glostatus*).

References: (hb) Loyttyniemi, Beaver, & Loyttyniemi 1984. (ds) Beaver & Loyttyniemi 1985a: 78.

(tx) Schedl 1957e: 873, 1961k: 638, 1964m: 305, 1979c: 161.

nigrivestis (Schedl) 1957e: 873 (*Ctonocryphus*). Holotype, sex?; Tanganyika, Ngorogoro; BMNH, London.

Distribution: Africa (Tanzania).

Notes: (1) Schedl 1964m: 305 (to *Paraglostatus*, = *Glostatus*).

References: (tx) Schedl 1957e: 873, 1961k: 638, 1964m: 305, 1979c: 167–168.

perplexus Schedl 1952: 281. Holotype, sex?; Cape Province, South Africa; Schedl Collection in NHMW, Wien.

Distribution: Africa (South Africa).

References: (tx) Schedl 1952: 281.

pondoanus Schedl 1958a: 558. Holotype, sex?; South Africa: Pondoland, Port St. John; BMNH, London.

Distribution: Africa (South Africa).

References: (tx) Schedl 1958a: 558, 1961k: 642, 1979c: 197.

seydeli (Nunberg) 1967b: 321 (*Rhopalocryphus*). Holotype, sex?; Elisabethville, Congo; MRCB, Tervuren.

Distribution: Africa (Zaire/ Zambia).

Hosts: *Brachystegia utilis*.

Notes: (1) Wood 1986a: 85 (to *Glostatus*).

References: (hb) Loyttyniemi, Beaver, & Loyttyniemi 1984. (ds) Beaver & Loyttyniemi 1985a: 76. (tx) Nunberg 1967a: 336–337, 1967b: 321; Wood, S. L. 1986a: 85.

spinarinatus Beaver in Beaver & Loyttyniemi 1985a: 78. Holotype ♂; Zambia: Chati, Kitwe; BMNH, London.

Distribution: Africa (Zaire).

References: (hb) Loyttyniemi, Beaver, & Loyttyniemi 1984. (tx) Beaver & Loyttyniemi 1985a: 78.

squamosus Schedl 1962k: 1071. Holotype, sex?; Natal; Schedl Collection in NHMW, Wien.

Distribution: Africa (South Africa).

References: (ds) Schedl 1965h: 111, 1969d: 5. (tx) Schedl 1962k: 1071, 1979c: 236.

vrydaghi (Nunberg) 1973: 10 (*Apoglostatus*). Holotype, sex?; north Ruwenzori, Kalonge, Zaire; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Nunberg 1973: 10.

xyloctonus (Schedl) 1941d: 398 (*Ctonocryphus*). Holotype, sex?; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon).

References: (tx) Schedl 1941d: 398, 1961k: 639, 1979c: 270.

Genus *Ctonoxylon* Hagedorn

CTONXYLON HAGEDORN 1910b: 4. Type-species: *Ctonoxylon auratum* Hagedorn, subsequent designation by Hopkins 1914: 119.

References: (ay) Nobuchi 1969a: 55. (hb) Schedl 1958d: 184, 1961k: 426, 1977b: 35; Wood, S. L. 1986a: 85. (ds) Hagedorn 1910d: 81; Schedl 1961k: 426, 1977b: 35; Wood, S. L. 1986a: 85. (tx) Eggers 1923a: 130, 1935a: 76; Hagedorn 1910a: 136–137, 1910b: 4–5; Hopkins 1914: 119, 132, 1915c: 226; Schedl 1939i: 380, 1941d: 398, 1952g: 50, 1957d: 1–162, 1961k: 426, 1977b: 35; Wood, S. L. 1986a: 85.

acuminatum Schedl 1957d: 45. Holotype, sex?; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Clitandra* cf. *staultii*.

References: (cc) Schedl 1961k: 426. (hb) Schedl 1961k: 426. (ds) Schedl 1961k: 426, 1982: 280. (tx) Schedl 1957d: 45, 1961k: 426, 1979c: 11.

alutaceus (Schaufuss) 1897b: 102 (*Stephanoderes*). Holotype, sex?; Tanga in Ostafrika; Hamburg Museum, lost.

Distribution: Africa (Tanzania).

Notes: (1) Eggers 1922b: 173 (to *Ctonoxylon*).

References: (ds) Hagedorn 1910d: 40; Kleine 1913b: 121, 1914b: 322. (tx) Eggers 1922b: 173; Hagedorn 1910a: 85; Hopkins 1915b: 24; Schaufuss 1897b: 102; Schedl 1961k: 547.

amanicum Hagedorn 1912b: 42. Holotype, sex?; Amani; Hamburg Museum, lost.

Distribution: Africa (Tanzania).

References: (bv) Aulmann 1911: 441–442; Aulmann & La Baume 1912: 65. (cn) Hagedorn 1913a: 14; Kleine 1932a: 303; Morstatt 1937: 26. (hb) Aulmann 1911: 441–442; Hagedorn 1913a: 14; Kleine 1932a: 303; Kolbe 1911: 508; Schedl 1961k: 428. (ds) Hagedorn 1913c; Kleine 1913b: 149, 1914b: 322, 1932a: 303, 1934a: 168, 630; Roba 1935: 338. (tx) Aulmann 1911; Eggers 1920: 39, 42–43; Hagedorn 1912b: 42; Schedl 1961k: 428.

atrum Browne 1965a: 190. Holotype, sex?; Nigeria: Idanre; BMNH, London.

Distribution: Africa (Nigeria).

Hosts: *Canthium* sp.

References: (tx) Browne 1965a: 190; Roberts 1969: 123.

- auratum Hagedorn** 1910b: 4. Holotype, sex?; Kamerun; MNB, Berlin.
Distribution: Africa (Cameroon/ Zaire).
References: (ds) Hagedorn 1910d: 81; Kleine 1913b: 149, 1914b: 312. (tx) Eggers 1927a: 198; Hagedorn 1910a: 137, 1910b: 4, 1912b: 42; Hopkins 1914: 119; Schedl 1961k: 428.
- bosqueiae Schedl** 1962h: 66. Holotype, sex?; Africa: Ghana, Bobiri, Kumasi; BMNH, London.
Distribution: Africa (Ghana/ South Africa).
Hosts: *Bosqueia angolense*.
References: (hb) Browne 1963a: 235. (ds) Browne 1963a: 235; Roberts 1969: 123; Schedl 1971g: 190, 1982: 279. (tx) Schedl 1962h: 66, 1962k: 1067, 1979c: 44.
- camerunum Hagedorn** 1910b: 4. Holotype, sex?; Kamerun; MNB, Berlin.
Distribution: Africa (Cameroon/ Equatorial Guinea/ Gabon/ Ivory Coast/ Nigeria/ Zaire).
Hosts: Liana.
References: (cn) Mayne & Donis 1951: 331. (ec) Mayne & Donis 1951: 331. (hb) Roberts 1969: 123. (ds) Hagedorn 1910d: 81; Kleine 1913b: 149, 1914b: 312; Schedl 1961k: 429. (tx) Eggers 1920: 37–38, 1927a: 198, 1943d: 246; Hagedorn 1910a: 137, 1910b: 4; Schedl 1939g: 171, 1950d: 6, 1961k: 429.
- fuscum Hagedorn** 1910b: 5. Holotype, sex?; Kamerun; MNB, Berlin.
Notes: (1) Eggers 1920: 37 (a variety of *camerunum*).
References: (ds) Hagedorn 1910d: 81; Schedl 1962k: 1067, 1964f: 618. (tx) Eggers 1920: 37–38; Hagedorn 1910a: 137, 1910b: 5; Schedl 1961k: 429, 1962k: 1067.
- capensis Schedl** 1971e: 8. Holotype, sex?; Umgeb. Cape Town; Schedl Collection in NHMW, Wien.
Distribution: Africa (South Africa).
References: (tx) Schedl 1971e: 8, 1979c: 51.
- caudatum Schedl** 1971e: 8. Holotype, sex?; Congo Belge, Stanleyville; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: (tx) Schedl 1971e: 8–9, 1979c: 54.
- conradti Schedl** 1939g: 171. Holotype, sex?; Ukerewi Island; Schedl Collection in NHMW, Wien.
Distribution: Africa (Angola/ Tanzania).
References: (ds) Ferreira 1965: 1112; Schedl 1959p: 16, 1961k: 430. (tx) Schedl 1939g: 171, 1961k: 430, 1979c: 63.
- cornutum Eggers** 1943d: 246. Holotype ♂; Kamerun; Strohmeier Collection.
Distribution: Africa (Cameroon).
References: (tx) Eggers 1943d: 246; Schedl 1961k: 430.
- crenatum Hagedorn** 1910b: 5. Holotype, sex?; Kamerun; MNB, Berlin.
Distribution: Africa (Cameroon).
References: (hb) Roberts 1969: 123. (ds) Hagedorn 1910d: 81; Kleine 1913b: 149, 1914b: 312; Schedl 1966c: 224. (tx) Eggers 1920: 39; Hagedorn 1910a: 137, 1910b: 5; Schedl 1961k: 430.
- dentigerum Schedl** 1941d: 388. Holotype, sex?; Spanish Guinea; Schedl Collection in NHMW, Wien.
Distribution: Africa (Equatorial Guinea).
References: (tx) Schedl 1941d: 388, 1961k: 430, 1979c: 78.
- festivum Schedl** 1941d: 389. Holotype, sex?; Kamerun, Soppo, 800 m; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon).
References: (tx) Schedl 1941d: 389, 1961k: 431, 1979c: 96.
- flavescens Hagedorn** 1910b: 4. Syntypes, sex?; Kamerun; MNB, Berlin.
Distribution: Africa (Cameroon/ Gabon/ Ghana/ Guinea/ Tanzania/ Zaire), Madagascar.
Hosts: *Allanblackia stuhlmannii*, *Carpodinus subrepanda*, *Landolphia robusta*, *Triposchiton scleroxylon*.
Notes: (1) Eggers 1920: 38 (elevated to species rank from a variety of *camerunum*).
References: (bv) Aulmann 1911: 411. (hb) Aulmann 1911: 411; Roberts 1969: 124. (ds) Hagedorn 1910d: 81; Kleine 1913b: 149, 1914b: 312; Numborg 1952: 17, 1965b: 17; Schedl 1961k: 431, 1964j: 40, 1965e: 351, 1965f: 4, 1965g: 19, 1966a: 275. (tx) Aulmann 1911: 411; Eggers 1920: 38, 1922b: 170; Hagedorn 1910a: 137, 1910b: 4; Numborg 1952: 17; Schedl 1954e: 49, 1955d: 268, 1961k: 431, 1965f: 4.
- camerunum hirsutum Hagedorn** 1910b: 4. Syntypes, sex?; Kamerun; MNB, Berlin, 1 in Eggers Collection, in NHMW, Wien.
Notes: (3) Schedl 1961k: 431 (treated as a variety of *flavescens*).
References: (ds) Hagedorn 1910d: 81; Kleine 1913b: 149; Roberts 1969: 124; Schedl 1964f: 618, 1972e: 280; Thompson 1963: 37. (tx) Eggers 1920: 38; Hagedorn 1910a: 137, 1910b: 4; Schedl 1954e: 49, 70, 1961k: 431, 1979c: 50.
- flavescens usambaricum Eggers** 1920: 38. Syntypes, sex?; Mkulumusiberg (1000 m), bei Sigi in Ostafrika; 1 in Eggers Collection, in NHMW, Wien.
References: (ce) Schedl 1961k: 433. (hb) Schedl 1961k: 433, 1977b: 35. (ds) Schedl 1961k: 433, 1977b: 35. (tx) Eggers 1920: 38; Schedl 1961k: 433, 1977b: 35, 1979c: 97.
- griseum Schedl** 1941d: 389. Holotype, sex?; Brit. Ost-Afrika, Kikuyu; Schedl Collection in NHMW, Wien.
Distribution: Africa (Kenya/ South Africa).
Hosts: *Olea capensis*, *O. hochstetteri*.
References: (cn) Anonymous 1970c: 13. (ds) Anon-

- amous 1970c: 13; Gardner 1957a: 30; Schedl 1961k: 435, 1965f: 4. **(tx)** Browne 1965: 190; Eggers 1920: 35; Schedl 1941d: 389, 1955i: 211, 1961k: 435, 1965f: 4, 1979c: 113.
- hamatum** Schedl 1941d: 390. Holotype, sex?; Nairobi; Schedl Collection in NHMW, Wien.
Distribution: Africa (Kenya/ South Africa/ Tanzania).
Hosts: *Olea hochstetteri*, *O. welwitschii*.
References: **(ds)** Gardner 1957a: 30. **(tx)** Browne 1969: 190; Powell 1980: 29; Schedl 1941d: 390, 1957b: 150, 1961k: 436, 1979c: 115.
- hirtellum** Schedl 1971e: 9. Holotype, sex?; Congo Belge, Yangambi; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: **(tx)** Schedl 1971e: 9, 1979c: 117.
- intermedium** Schedl 1971e: 10. Holotype, sex?; Kamerun; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon).
References: **(tx)** Schedl 1971e: 10, 1979c: 126.
- kivuensis** Schedl 1957d: 44. Holotype, sex?; Congo Belge; Kivu, Mulungu; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Popowia* sp.
References: **(ds)** Schedl 1961k: 437. **(tx)** Schedl 1957d: 44, 1961k: 437, 1979c: 133.
- longipilum** Eggers 1935c: 308. Holotype ♀; Brit. Ostafrika (Mulanje); USNM, Washington.
Distribution: Africa (Ghana/ Ivory Coast/ Kenya/ Tanzania). Madagascar.
References: **(ds)** Schedl 1972e: 280, 1977b: 36, 1977d: 275, 1982: 281. **(tx)** Anderson, W. H. & Anderson 1971: 15; Eggers 1935c: 308, 1940c: 236; Nunberg 1969: 380; Schedl 1953d: 67, 1961k: 437, 1977b: 36.
- methneri** Eggers 1922b: 170. Holotype, sex?; Utzungwe-Berge, 1300–1600 m; Methner Collection.
Distribution: Africa (Kenya/ South Africa/ Tanzania).
References: **(tx)** Browne 1965: 22; Eggers 1922b: 170; Schedl 1961k: 435, 1962k: 1067, 1965g: 19.
- montanum** Eggers 1922b: 170. Holotype ♀; Buca, Kamerun; USNM, Washington.
Distribution: Africa (Cameroon).
References: **(tx)** Anderson, W. H. & Anderson 1971: 21; Eggers 1922b: 170, 1935c: 308; Schedl 1961k: 435.
- nodosum** Eggers 1940c: 236. Holotype, sex?; Congostaat: Mongwalu (Kilo), im Osten an der Grenze gegen Brit. Ostafrika; MRCB, Tervuren.
Distribution: Africa (Cameroon/ Ghana/ Nigeria/ Uganda/ Zaire).
References: **(ds)** Browne 1973a: 279; Roberts 1969: 124; Schedl 1962h: 58, 1962k: 1067, 1965f: 4. **(tx)** Eggers 1940c: 236; Schedl 1961k: 435, 1962k: 1067, 1965f: 4.
- pygmaeum** Eggers 1920: 39. Syntypes, sex?; Soppo in Kamerun, 800 m; MNB, Berlin, and Eggers Collection, 2 Eggers syntypes in NHMW, Wien.
Distribution: Africa (Cameroon).
References: **(tx)** Eggers 1920: 39; Schedl 1961k: 439, 1962h: 66, 1979c: 207.
- setifer** Eggers 1920: 39. Holotype, sex?; Amani; Hamburg Museum, lost.
Distribution: Africa (Kenya/ Tanzania/ Zaire).
References: **(ds)** Gardner 1957a: 30; Schedl 1961k: 439. **(tx)** Eggers 1920: 39, 1922b: 170, 1927a: 195; Schedl 1941d: 390, 1951e: 39, 1952g: 50, 1961k: 439, 1979c: 226.
- spathifer** Schedl 1951e: 39. Syntypes, sex?; Cote d'Ivoire, Reserve du Banco; MNHN, Paris, and Schedl Collection in NHMW, Wien.
Distribution: Africa (Ivory Coast/ Tanzania).
Hosts: *Olea welwitschii*, *Pachylobus deliciosus*, *Strombosia pustulata*.
References: **(ds)** Cachan 1957: 15; Schedl 1961k: 440, 1979b: 415. **(tx)** Schedl 1951e: 39, 1954d: 570, 1961k: 440, 1979c: 233.
- spinifer** Eggers 1920: 39. Lectotype ♀; Kamerun; USNM, Washington, designated by Anderson & Anderson 1971: 31.
Distribution: Africa (Cameroon/ Ivory Coast).
References: **(ds)** Cachan 1957: 15; Roberts 1969: 124; Schedl 1961k: 440, 1964j: 40, 1967e: 210. **(tx)** Anderson, W. H. & Anderson 1971: 30–31; Eggers 1920: 39; Schedl 1951e: 35, 1954d: 570, 1961k: 441, 1965h: 114.
- uniserialatum** Schedl 1965h: 114. Holotype, sex?; S.W. Africa; Schedl Collection in NHMW, Wien.
Distribution: Africa (South West Africa).
References: **(tx)** Schedl 1965h: 114, 1979c: 261.

Genus *Scolytomimus* Blandford

- Scolytomimus* BLANDFORD 1895a: 319. Type-species: *Scolytomimus dilutus* Blandford, monobasic.
- Neoxylotomus* Eggers 1923a: 143. Type-species: *Neoxylotomus philippinensis* Eggers, monobasic. Synonymy: Schedl 1951i: 53.
References: **(tx)** Eggers 1923a: 143; Schedl 1951i: 53.
- Scolytocleptes* Schedl 1962q: 490. Type-species: *Scolytomimus maculatus* Beeson, original designation. Synonymy: Browne 1970: 557.
References: **(tx)** Browne 1970: 557; Schedl 1962q: 490.
- Keys: Browne 1958a: 487.
References: **(hb)** Browne 1961c: 79–81; Wood, S. L. 1986a: 85. **(ds)** Browne 1961c: 79–81; Wood, S. L. 1986a: 85. **(tx)** Beeson 1938b: 290; Blandford 1895a: 319; Browne 1958: 487–497, 1970: 557; Schedl 1951i: 53; Wood, S. L. 1960a: 1–73, 1986a: 85.
- andamanensis** Wood 1988c: 199. Holotype ♂; Andaman Islands; FRI, Dehra Dun.
Distribution: Asia (Andaman Islands in India).

Hosts: *Sideroxylon longepetiolatum*.

Notes: (3) Schedl 1951i: 53 (*Xyloctonus andamanensis* Beeson, nomen nudum).

References: (tx) Wood, S. L. 1988c: 199–200.

assamensis Schedl 1951i: 54. Holotype, sex?; Assam, Lushai Hills, India; Schedl Collection in NHMW, Wien.

Distribution: Asia (Bangladesh/ Assam and Bihār in India/ Sri Lanka), Indonesia (Sarawak in Borneo).

Hosts: *Bassia latifolia*, *B. butyraces*, *Bauhinia tomentosa*, *Camellia sinensis*, *Dichopsis polyantha*, *Isonandra polyantha*, *Palaquium pseudo-rostratum*.

References: (cn) Mathur & Singh 1960b: 73. (ds) Mathew 1981, 1987: 188; Mathur & Singh 1960b: 73; Schedl 1962p: 203. (tx) Schedl 1951i: 54, 1962p: 203, 1979c: 29.

bassiae Schedl 1951i: 54. Holotype, sex?; Indien, Bihār, Balaghat; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992:(in press).

References: (ds) Bhasin, Roonwal, & Singh 1958. (tx) Browne 1958a: 491; Schedl 1951i: 54, 1979c: 35.

quadrioculatus Browne 1958: 490. Holotype, sex?; Sarawak: Setapok, near Kuching; BMNH, London. Synonymy: Schedl 1962p: 203.

References: (ds) Bhasin, Roonwal, & Singh 1958. (tx) Browne 1958: 490; Schedl 1962p: 203.

baloghi Schedl 1969e: 157. Holotype, sex?; New Guinea, Mt. Kaindi; NIIMB, Budapest.

Distribution: New Guinea.

References: (tx) Schedl 1969e: 157, 1979c: 33.

bicolor Wood 1960a: 18. Holotype ♀; E. Ngatpang, Babelthuap, Palau Island; USNM, Washington.

Distribution: Micronesia (Palau Island).

References: (tx) Wood, S. L. 1960a: 17–18.

costipennis Schedl 1953c: 291. Syntypes 2, sex?; Malaya, Pahang, Ginting Kial; BMNH, London, and Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

References: (ds) Browne 1961c: 81. (tx) Browne 1970: 557; Schedl 1953c: 291, 1962q: 490, 1979c: 67.

dilutus Blandford 1895a: 320. Syntypes 2, sex?; Ceylon, Bogawantalawa; BMNH, London.

Distribution: Asia (Sri Lanka).

Hosts: Kiribade.

Notes: (3) Schedl 1951i: 53 (*darwinii*, nomen nudum).

References: (ds) Kleine 1913b: 149; Schedl 1959a: 474, 1969c: 48. (tx) Beeson 1929: 224, 1938: 290; Blandford 1895a: 320; Browne 1958a: 491; Hopkins 1914: 129; Lucas 1920: 572; Schedl 1951i: 53, 1953c: 291, 1954a: 138, 1959a: 474. (ms) Lucas 1920: 572.

maculatus Beeson 1929: 223. Holotype, sex?; Upolu: Apia, Vailima; BMNH, London.

Distribution: Samoan Islands.

Hosts: *Planchonella samoensis*.

References: (cc) Roberts 1976: 382. (hb) Beaver 1976b: 536; Roberts 1976: 382. (ds) Beaver 1976b: 532; Beeson 1938b: 290; Browne 1974a: 65, 1980c: 484, 1981b: 598, 1984a: 151; Olmo et al. 1987: 87; Roberts 1976: 382. (tx) Beeson 1929: 223; Browne 1970: 557; Schedl 1950f: 36, 1951i: 53, 1951k: 132, 1962q: 490, 1979c: 144; Wood, S. L. 1960a: 18.

mimusops Wood 1988c: 200. Holotype ♀; Palangama, Anuradhapura, Ceylon; FRI, Dehra Dun.

Distribution: Asia (Sri Lanka).

Hosts: *Bassia latifolia*, *Mimusops elengi*.

Notes: (3) Schedl 1951i: 53 (*Xyloctonus mimusops* Beeson, nomen nudum).

References: (tx) Wood, S. L. 1988c: 200.

philippinensis (Eggers) 1923: 143 (*Neoxyloctonus*). Syntypes, sex?; Butuan auf Mindanao, Philippines; SMTD, Dresden.

Distribution: Australia, Philippine Islands (Mindanao).

Hosts: *Palmctia pinnata*.

References: (ds) Browne 1966: 245; Schedl 1966b: 16, 1972a: 143. (tx) Eggers 1923: 143, 1927b: 406, 1937b; Nobuchi 1983: 299; Schedl 1951i: 53.

pusillus (Eggers) 1927c: 88 (*Neoxyloctonus*). Lectotype, sex?; Philippines: Mindanao, Prov. Lanao, Kolambugan; USNM, Washington, designated by Anderson & Anderson 1971: 27.

Distribution: Asia (Andaman Islands in India/ Malaya), Indonesia (Borneo, Java), New Guinea, Philippine Islands (Mindanao).

Hosts: *Achras sapota*, *Mimusops littoralis*, *Palaquium mainingayi*, *P. stellatum*.

References: (hb) Beaver & Browne 1978: 599; Browne 1961c: 80; Kalshoven 1958b: 180. (ds) Beaver & Browne 1978: 599; Browne 1961c: 80, 1966: 245, 1980b: 381, 1980c: 484, 1984a: 151; Ohno, Yoneyama, & Nakazawa 1987b: 93; Ohno, Yoshioka et al. 1988a: 91, 1989: 60; Schedl 1966b: 16, 1975i: 346. (tx) Anderson, W. H. & Anderson 1971: 27; Browne 1961: 80; Eggers 1927c: 88; Nobuchi 1983: 300; Schedl 1940b: 435, 1951i: 50, 53, 1953c: 288, 1962p: 203, 1979c: 206.

kalshoveni Eggers 1940d: 132. Holotype, sex?; Java: Kalshoven Collection. Synonymy: Wood 1989: 175.

References: (hb) Kalshoven 1958: 180. (tx) Browne 1958a: 491; Eggers 1940d: 132; Schedl 1942d: 1, 1951i: 53, 1953c: 288, 1979c: 131; Wood, S. L. 1989: 175.

brunigi Browne 1958: 488. Holotype, sex?; Sarawak: Bako National Park; BMNH, London. Synonymy: Schedl 1962p: 203.

References: (ds) Browne 1961a: 304. (tx) Browne 1958: 488; Schedl 1962p: 203.

- menoni* Browne 1958: 457. Holotype, sex?; Malaya: Selangor, Kepong; BMNH, London. Synonymy: Schedl 1962p: 203. References: **(ds)** Browne 1961c: 81, **(tx)** Browne 1958: 457, 1961c: 81; Schedl 1962p: 203.
- insularis* Schedl 1962q: 491 (*Scolytocleptes*). Holotype, sex?; Andanan Islands; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 175. Notes: (1) Browne 1970: 557 (to *Scolytominus*). (3) Beeson 1941: 295 (*insularis*, nomen nudum). References: **(ds)** Beeson 1961: 295, **(tx)** Browne 1958a: 491, 1970: 557; Schedl 1951i: 53, 1955b: 287, 1962q: 491, 1979c: 126; Wood, S. L. 1989: 175.
- quadridens* Wood 1988c: 200. Holotype ♂; New Guinea (NE), Hiron Peninsula, Finschafen; Wood Collection. Distribution: New Guinea. References: **(tx)** Wood, S. L. 1988c: 200–201.
- rectus* Wood 1988c: 201. Holotype ♂; 40 km NE Polonnaruwa, Pol. Distr., Sri Lanka; USNM, Washington. Distribution: Asia (Sri Lanka). Hosts: *Manikara hexandra*. References: **(tx)** Wood, S. L. 1988c: 201.
- tennis* Browne 1984: 77. Holotype, sex?; Fiji: Viti Levu, Namosi Road, Savura Creek; BMNH, London. Distribution: Fiji Islands. References: **(tx)** Browne 1984: 77.
- woodi* Browne 1958: 489. Holotype, sex?; Malaya: Selangor, Kepong; BMNH, London. Distribution: Asia (Malaya). Hosts: *Palaquium maingayi*, *P. stellatum*. Notes: (3) Schedl 1962p: 203 (incorrectly cited as a synonym of *pusillus*). References: **(hb)** Browne 1961c: 80, **(ds)** Browne 1961c: 80, **(tx)** Browne 1958: 489, 1961c: 80; Schedl 1962p: 203.

Genus *Xyloctonus* Eichhoff

- Xyloctonus* EICHHOFF 1872a: 134. Type-species: *Xyloctonus scolytoides* Eichhoff, monobasic. Keys: Menier 1974: 635. References: **(hb)** Schedl 1958d: 184, 1961k: 418, 1977b: 31; Wood, S. L. 1986a: 85, **(ds)** Hagedorn 1910d: 81; Schedl 1961k: 418, 1977b: 31; Wood, S. L. 1986a: 85, **(tx)** Blandford 1893d: 441, 1895a: 320, 1904: 412; Eggers 1939a: 14; Eichhoff 1872a: 134, 1878b: 335; Hagedorn 1904: 412, 1910a: 135–136, 1910b: 4; Hopkins 1914: 132, 135, 1915c: 226; Menier 1974: 653–666; Schedl 1939e: 340, 1939i: 350, 1941d: 398, 1957d: 1–162, 1961k: 418, 1965e: 349–379, 1977b: 31; Wood, S. L. 1986a: 85.
- aethiops* Schedl 1953d: 77. Lectotype, sex?; Madagascar, Ankorika; MNHN, Paris, designated by Menier 1974: 660. Distribution: Madagascar.

Hosts: *Savia hildebrandti*.

References: **(tx)** Menier 1974: 660; Schedl 1953d: 77, 1977b: 31, 1979c: 14.

stenographus Schedl 1961e: 130. Holotype, sex?; Madagascar, Perinet; MNHN, Paris. Synonymy: Menier 1974: 660.

References: **(hb)** Schedl 1977b: 34, **(ds)** Schedl 1977b: 34, **(tx)** Menier 1974: 660; Schedl 1961e: 130, 1977b: 31–34, 1979c: 237.

bimarginatus Eggers 1939a: 17. Holotype, sex?; Congostaat (Kundelungus); MRCB, Tervuren.

Distribution: Africa (Zaire).

References: **(tx)** Eggers 1939a: 17; Menier 1974: 656; Schedl 1961k: 419.

biseriatus Schedl 1953d: 76. Lectotype, sex?; Madagascar, Region de l'Androy, Ambovombe; MNHN, Paris, designated by Menier 1974: 661.

Distribution: Madagascar.

References: **(tx)** Menier 1974: 656, 661; Schedl 1953d: 76, 1977b: 31, 1979c: 41.

latus Eggers 1939a: 14. Lectotype, sex?; Abessinia (Djem-djem Forest); BMNH, London, designated by Menier 1974: 661.

Distribution: Africa (Ethiopia/ South Africa/ Zambia).

References: **(ds)** Schedl 1965e: 351, 1965h: 111, 1969d: 5, **(tx)** Eggers 1939a: 14–15; Menier 1974: 661; Schedl 1961k: 419, 1965e: 365, 1979c: 137.

maculatus Schedl 1965h: 113. Holotype, sex?; Cape Prov., Port Elizabeth; Transvaal Museum.

Distribution: Africa (South Africa).

Hosts: *Sideroxylon incurv.*

References: **(tx)** Schedl 1965g: 19, 1965h: 113, 1979c: 114.

mauritanus Menier 1974: 662. Holotype ♂?; Corps de garde, Mauritius; MNHN, Paris.

Distribution: Mauritius Island.

References: **(tx)** Menier 1974: 662, 663.

niger Schedl 1938d: 452. Lectotype, sex?; Uganda: Entebbe; BMNH, London, designated by Menier 1974: 663.

Distribution: Africa (Uganda).

Hosts: *Conopharyngia holzkii*.

References: **(ds)** Gardner 1957a: 33; Schedl 1938d: 452, **(tx)** Eggers 1939b: 14; Menier 1974: 663; Schedl 1938d: 452, 1961k: 419, 1979c: 179.

opacus Schedl 1957d: 43. Holotype, sex?; Ruanda: Ihembe; MRCB, Tervuren.

Distribution: Africa (Ruanda in Zaire).

Hosts: *Chrysophyllum boivinianum*.

References: **(cc)** Schedl 1958d: 190–191, 1961k: 420, **(hb)** Schedl 1958d: 191, 1961k: 420, 1977b: 32, **(ds)** Mayne & Donis 1962: 323; Schedl 1961k: 420, 1977b: 32, 1979b: 415, **(tx)** Schedl 1957d: 43, 1961k: 420, 1977b: 32, 1979c: 179.

pubifer Schedl 1965e: 365. Holotype, sex?; Port Elisabeth; Schedl Collection in NHMW, Wien.

Distribution: Africa (South Africa/ Zambia).

References: **(hb)** Loyttniemi, Beaver, & Loyttniemi 1984. **(ds)** Beaver & Loyttniemi 1985a: 79; Schedl 1982: 281. **(tx)** Schedl 1965c: 365, 1979c: 203.

punctipennis Eggers 1939a: 16. Holotype, sex?; Somaliland (Basso Genana); USNM, Washington. Distribution: Africa (Somalia).

References: **(tx)** Anderson, W. H. & Anderson 1971: 27; Eggers 1939a: 16; Schedl 1961k: 422, 1979c: 205.

quadricinctus Schedl 1941d: 387. Holotype, sex?; Usambara; Schedl Collection in NHMW, Wien. Distribution: Africa (Ghana/ Tanzania).

Hosts: *Chrysophyllum albidum*.

References: **(hb)** Thompson, G. H. 1963: 73. **(ds)** Roberts 1969: 134; Schedl 1961k: 423; Thompson, G. H. 1963: 73. **(tx)** Menier 1974: 656, 665; Schedl 1941d: 387, 1954e: 49, 1957d: 43, 1961k: 422, 1979c: 208.

quadridens Schedl 1953d: 77. Lectotype ♂; Madagascar, Mt. d'Ambre; MNHN, Paris, designated by Menier 1974: 665.

Distribution: Madagascar.

Hosts: *Filicium abbreviatum*.

References: **(hb)** Schedl 1977b: 33. **(ds)** Schedl 1977b: 33. **(tx)** Menier 1974: 665; Schedl 1953d: 77, 1977b: 33, 1979c: 208.

scolytoides Eichhoff 1872a: 134. Lectotype, sex?; Port Natal; IRSNB, Brussels, designated by Menier 1974: 665.

Distribution: Africa (Ghana/ Ivory Coast/ Kenya/ Nigeria/ South Africa/ Sudan/ Tanzania/ Zaire).

Hosts: *Butyrospermum parkii*, *Illepe* (*Bassia*) *latifolia*, *Minusops caffra*.

References: **(cn)** Anonymous 1970c: 14. **(cc)** Beeson 1922c: 500. **(hb)** Browne 1963a: 236;

Roberts 1969: 134. **(ds)** Anonymous 1970c: 14; Beeson 1922c: 500, 1961: 310; Browne 1963a: 236; Gemminger & Harold 1872: 2684; Hagedorn 1910d: 81; Kleine 1913b: 148, 1914b: 319, 1934a: 167–168; Schedl 1938d: 450, 1961k: 423, 1962h: 53, 1962k: 1066, 1964j: 40, 1972c: 280, 1975h: 351, 1977d: 178, 1982: 279. **(tx)** Blandford 1895a: 320; Eggers 1922b: 173–174, 1927a: 198, 1936b: 28–33, 1939b: 14; Eichhoff 1872a: 134, 1878b: 171–172; Hagedorn 1910a: 136; Hopkins 1914: 132, 135, 1915c: pl. 9, fig. 6; Lucas 1920: 675; Menier 1974: 665; Schedl 1939g: 170, 1941d: 380, 1951i: 53–54, 1957b: 150, 1957d: 43, 1961k: 423, 1962k: 1066, 1965g: 17. **(ms)** Lucas 1920: 675.

cmarginatus Eggers 1939a: 16. Holotype, sex?; Brit. Ostafrika (W. Nile); USNM, Washington. Synonymy: Menier 1974: 665.

References: **(tx)** Anderson, W. H. & Anderson 1971: 12; Eggers 1939a: 15–16; Menier 1974: 665; Schedl 1961k: 419, 1979c: 90.

striatus Eggers 1939a: 15. Holotype, sex?; Mozambique (Sangadze, Moulima); MNHN, Paris.

Distribution: Africa (Mozambique).

Hosts: *Acacia* sp.

References: **(ds)** Schedl 1961k: 425. **(tx)** Eggers 1939a: 18; Schedl 1961k: 425, 1965e: 365, 1979c: 238.

subcostatus Eggers 1939a: 15. Holotype, sex?; Deutsch Ost Africa (Bez. Tabora; Ngulu); USNM, Washington.

Distribution: Africa (Guinea/ Mozambique/ Sudan/ Tanzania).

References: **(ds)** Schedl 1961k: 425. **(tx)** Anderson, W. H. & Anderson 1971: 32; Eggers 1939a: 15–16; Schedl 1955d: 268, 1961k: 425, 1979c: 240.

Tribe Cryphalini Lindemann

Cryphaloidea

References: Lindemann 1876: 165.

Cryphalidae

References: Eichhoff 1878b: 107, 1881a: 169.

Cryphalinae

References: Hagedorn 1909: 162, 1910a: 24, 80, 1910d: 40; Hopkins 1915b, 1915c: 224; Leng 1920: 339; Lucas 1920: 19; Nusslin 1911: 430; Tredl 1907: 12.

Cryphalini

References: Leng 1920: 339; Reiter 1913a: 28; Wood, S. L. 1978a: 114, 1982b: 69, 1986a: 85–93.

Cryphalina

References: Balachowsky 1949a: 200; Barbey 1901: 21; Beal & Massey 1945: 119; Bedel 1888b: 397, 413.

Cryphali

References: Blandford 1904: 225.

Trypophloeinae

References: Nusslin 1911: 373, 431.

Ernoporinae

References: Nusslin 1911: 375, 429.

Eidophelinae

References: Murayama 1954: 200.

Genus *Trypophloeus* Fairmaire

TRYPOPHLOEUS FAIRMAIRE 1865: 105. Type-species:

Bostrichus binodulus Ratzeburg, monobasic.

Glyptoderes Eichhoff 1878b: 34, 44, 137. Type-species: *Bostrichus binodulus* Ratzeburg, subsequent designation by Hopkins 1914: 122. Synonymy: Bedel 1888b: 397.

References: (tx) Balachowsky 1949a: 213; Eichhoff 1878b: 34, 44, 137; Göz 1985: 278; Wood, S. L. 1954a: 998.

Keys: Wood 1954a: 989, 1982b: 851 for North America; Balachowsky 1949a: 214, Reitter 1913a: 69, Stark 1952: 281 for Asia and Europe.

References: (hb) Wood, S. L. 1982b: 850, 1986a: 89. (ds) Wood, S. L. 1982b: 850, 1986a: 89. (tx) Balachowsky 1949a: 213; Bright 1976d: 108; Dodge 1935: 18; Fairmaire 1864: 121–127, 1865: 105; Hansen, V. 1955: 169–176; Hopkins 1914: 131–132, 1915b: 8, 36; Karaman 1972: 122; Klimesch 1913: 105–116, 1914: 213–219, 231–241; Kostin 1973: 253; Krivolitskaya 1956: 152; Nimberg 1956: 141; Peyerimhoff 1935: 194–195; Pfeffer 1889a: 83; Reitter 1913a: 65, 69–71; Rudnev 1958: 372; Saalas 1914: 73, 83; Schedl 1962p: 204, 1981b: 72; Scheerpeltz & Winkler 1930: 257; Schimitschek 1937: 50–51; Sokanovskii 1954: 21–22; Spessivtsev 1922: 470, 1931: 88; Stresemann et al. 1989: 353; Swaine 1918a: 45, 90; Trappen 1935: 142; Wood, S. L. 1954a: 988, 1967a: 122, 1982b: 850, 1986a: 89. (ms) Kangas 1969.

alni (Lindemann) 1875a: 136 (*Cryphalus*). Syn-types, sex?; Moskau; not located.

Figures: Grune 1979: 98.

Distribution: Asia (Xinjiang in China), Europe (Austria/ Czechoslovakia/ Finland/ Germany/ N Italy/ Norway/ Poland/ W USSR).

Hosts: *Alnus glutinosa*, *A. incana*.

References: (ay) Klimesch 1914: 218. (bv) Grune 1979: 99. (cn) Grandi 1951: Kontkanen 1932: 63; Koppen 1882: 252; Nestertschuk 1930: 176; Nusslin 1913: 291; Pierce, W. D. 1917: 12; Rhumbler 1922: 334, 1927: 351. (ec) Balazy & Michalski 1964b; Kangas 1942b; Kleine 1908c: 187, 1944: 70; Michalski & Ratajczak 1989; Nusslin 1927: 351. (hb) Altum 1881c: 322; Dombrowsky 1887, 1889; Eichhoff 1881a: 46, 189; Gornostaev 1916: 314; Grandi 1951; Henschel 1895a: 170; Kangas 1940b: 41–50; Klimesch 1915: 11; Kostin 1960: 132; Lengerken 1954b: 84; Lindemann 1875a: 136; Palm 1956: 63; Nusslin 1913: 291, 1927: 351; Rhumbler 1922: 334, 1927: 351; Rupertsberger 1879: 231, 1880: 229; Schedl 1981b: 73; Stark 1926a: 330, 1952: 283. (ds) Gornostaev 1917; Grune 1979: 99; Hagedorn 1910d: 40; Hansen, V. 1939; Hellen 1947; Henschel 1895a: 170; Heyden, Reitter, & Weise 1883: 181, 1891: 671, 1906: 711; Kangas 1942b; Klefbeck & Sjöberg 1960: 203; Kleine 1913a: 35, 1913b: 123, 1934a: 147; Kontkanen 1932: 63; Koppen 1882: 252; Kraatz 1876c: 107; Lindemann 1884b: 264; Linnamuri 1949a: 185, 1949b: 34; Lucht 1987: 278; Nimberg 1954: 50; Nuorteva 1971; Palm 1956: 63; Pierce, W. D. 1917: 12; Pittioni 1943: 176; Postner 1974: 428; Reitter 1889a: 93, 1894a: 72; Schanfuss 1915: 1233; Schedl 1980a: 15, 1981b: 73; Seidlitz 1891b: 563, 608; Sokanovskii 1960; Stark 1926a: 330, 1926j: 125, 1931a: 25, 1952: 283; Stein & Weise 1877: 164; Strand 1946: 600, 1953: 62; Swaine 1912a: 349; Tredl 1907: 13; Wichmann 1909b: 173; Yarovskii & Tegshzhargal 1985: 409; Zinovjev 1955: 191. (tx) Balachowsky 1949a: 218; Dombrowsky 1887, 1889; Eichhoff 1878b: 44, 139, 1881a: 46, 189, 1883a: 110, 133; Grune 1979: 98, 99; Hagedorn 1910a: 85; Henschel 1895a: 170; Hopkins 1915b: 36; Lindemann 1875a: 136, 1877a: 161; Lucht 1987: 278; Nimberg 1954: 50; Pfeffer 1955a: 184; Reitter 1889a: 93, 1894a: 72, 1913a: 71; Rhumbler 1922: 334, 1927: 351; Rupertsberger 1879: 231, 1880: 229; Schedl 1934f: 1639, 1980a: 15, 1981b: 73; Sedlaczek 1912: 307; Seidlitz 1891a: 563, 1891b: 608; Sokanovskii 1954: 22, 1960: 676; Spessivtsev 1922a: 471, 492, 1931: 47; Stark 1935: 152, 1952: 203; Strand 1953: 62. (ms) Kraatz 1876c: 107.

holdhausi Wichmann 1912c: 186. Holotype ♀; Campo grosso [N Italy]; NHMW, Wien. Synonymy: Balachowsky 1949a: 218.

References: (ds) Kleine 1913b: 123; Schanfuss 1915: 1233. (tx) Balachowsky 1949a: 218;

Eggers 1914; Reitter 1913a: 70–71; Schedl 1934f: 1639; Wichmann 1912c: 186.

***binodulus* (Ratzeburg) 1837: 163 (*Bostrichus*).**

Syntypes, sex?, apparently Germany; not located. Distribution: Africa (Algeria/ Morocco), Europe (Austria/ Belgium/ Bulgaria/ Corsica/ Czechoslovakia/ Denmark/ England/ Finland/ France/ Germany/ Hungary/ Italy/ Netherlands/ Norway/ Poland/ Romania/ Spain/ Sweden/ Switzerland). Hosts: *Populus pyramidalis*, *P. tremula*, *Salix caprea*, *S. fragilis*, *S. sp.*

Notes: (3) Most references (except Gyllenhal 1813: 363) to *Bostrichus asperatus* published prior to Wood 1972: 41 refer to *binodulus*; the species designated by the name *asperatus* is in *Cryphalus*.

References: (ay) Escherich 1923b: 482; Klimesch 1914: 216–217, 237; Nusslin 1911a: 112. (cn) Barbey 1925: 566; Escherich 1923b: 482; Esterberg 1959; Kontkanen 1932: 63; Nester-tschuk 1930: 176; Nusslin 1913: 291; Palm 1951: 231; Rhumbler 1922: 334, 1927: 351; Schwerdtfeger 1944a: 175, 1957a: 182; Wachtl 1901: 381. (cc) Brammanis 1938, 1940; Elliot & Morley 1907; Heqvist 1967: 69; Kleine 1908c: 187, 1909a: 47; Nosek 1959a: 118, 1959b: 86; Nusslin 1927: 351; Rondani 1873: 146; Schwerdtfeger 1944a: 175, 1957a: 182. (hb) Altum 1881c: 322; Barbey 1901: 74, 1925: 566; Chapman 1869a: 198; Dombrowsky 1887, 1889; Eichhoff 1881a: 46, 188; Escherich 1923b: 482; Everts 1903: 756; Gyorfi 1957; Hagedorn 1903a, 1904f; Henschel 1895a: 170; Karpinski & Strawinski 1948: 155; Kellner 1862: 280; Kemner 1919: 175; Klimesch 1915: 11; Lengerken 1939: 63, 1954: 84; Munro 1926: 62; Nordlinger 1856: 29; Nosek 1959a: 118, 1959b: 86; Nunberg 1929a: 105; Nusslin 1895: 280, 1913: 291, 1927: 351; Palm 1950: 142, 1951: 231; Peyerimhoff 1926: 387; Pfeffer 1942a: 15; Ratzeburg 1837: 163, 1839: 199; Rhumbler 1922: 334, 1927: 351; Rupertsberger 1879: 231, 1880: 229; Saalas 1913a: 69, 83; Schwerdtfeger 1944a: 175, 1957a: 182; Stark 1926a: 334, 1952: 289; Wachtl 1876a: 458, 1901: 381. (ds) Aclouque 1896; Allen, A. 1954, 1958: 216; Arnoldi et al. 1955: 694; Bedel 1888b: 397, 414; Bielz 1887; Blanchere & Robert 1889; Borchert 1951; Calver 1884, 1893; Carpentier & Delaby 1908; Crotch 1863; Eggers 1904; Endrodi 1958b; Ericson & Sandin 1893; Escherich 1923b: 482, 1932b; Esterberg 1959; Everts 1922: 642; Fowler 1891; Fuchs 1905a; Fuss 1874; Gaubil 1849: 126; Gemminger & Harold 1872: 2683; Gozis 1875: 80, 1885: 278; Gredler 1875: 115; Grill 1895: 309; Hagedorn 1903a, 1910d: 41; Hansen, V. 1939, 1863; Hellen 1947; Henschel 1895a: 170; Heyden 1876: 299, 1879: 140; Heyden, Reitter, & Weise 1883: 181, 1891: 671, 1906: 711; Horion 1951; Jansson 1935: 77; Jazentkovsky 1912: 288; Kangas 1946a: 36;

Karpinski 1925: 216, 1931: 20, 39, 1932b: 54; Karpinski & Strawinski 1948: 155; Keler 1925b: 272; Kellner 1862: 280; Kemner 1919: 175; Kestercanek 1881a: 12; Kleine 1912a: 263, 267, 1913a: 35, 1913b: 123, 1934a: 147; Kloft & Hinks 1945: 218; Koltze 1901: 153; Kontkanen 1932: 63; Kozikowsky & Kuntze 1925: 20; Kraatz 1869: 59; Lacordaire 1866: 379; Langhoffer 1915c: 158; Lindemann 1884b: 264; Linder 1953: 71; Lindgren 1945: 77; Lomnicki 1886a: 242, 1913b: 148; Matthews & Fowler 1883: 42; Nunberg 1928b: 88, 106, 1954: 50; Nusslin 1895: 280; Palm 1950: 142, 1959: 24, 350; Perris 1876a: 255, 1877a: 415; Peyerimhoff 1926: 387; Pfeffer 1931b: 73, 1936: 90; Pittioni 1943: 176; Poppus 1900: 108; Power 1865: 212; Prossen 1913: 83; Rapp 1934: 728; Ratzeburg 1837: 163, 1839: 199; Redtenbacher 1858: 832, 1874: 375; Reitter 1869b: 154, 1894a: 72, 1916: 289; Rohrig 1955: 37; Ronbal 1941: 264; Saalas 1913a: 69, 83; Sahlberg 1900: 105; Sainte-Claire 1914: 472; Sainte-Claire & Mequignon 1938: 445; Schaufuss 1915: 1233; Schann 1859: 96, 1862: 101; Scheerpeltz & Winkler 1930: 257; Schilsky 1909: 188; Seidlitz 1872: 393, 1891a: 563, 1891b: 608; Sharp & Fowler 1893: 34; Stark 1926a: 334, 1926b: 103, 1926j: 125, 1931a: 25, 1952: 289; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 438; Thomson 1868: 219; Tredl 1907: 13; Wachtl 1876a: 458; Westhoff 1882: 238; Wichmann 1927a: 68. (tx) Aclouque 1896; Bach 1854; Balachowsky 1949: 215; Barbey 1901: 74; Bedel 1888b: 397, 414; Bertolini 1872; Carpentier & Delaby 1908; Doehner 1860; Dombrowsky 1887, 1889; Duffy 1953; Eggers 1927: 121–122; Eichhoff 1864b: 35, 1878b: 44, 139, 1881a: 46, 188, 1883a: 110, 133; Endrodi 1957b: 416; Escherich 1923b: 482; Everts 1903: 756, 1922: 642; Fairmaire 1864: 105; Fauvel 1889; Ferrari 1867a: 11, 1867b: 114; Fleischer 1927; Formanek 1907: 36; Hagedorn 1904a: 230; Hansen, V. 1955, 1956: 179; Henschel 1895a: 170; Hopkins 1914: 122, 131, 1915b: 36–37; Houllbert 1922a: pl. 1; Jacquelin du Val & Fairmaire 1868: 105; Karpinski & Strawinski 1948: 155; Klimesch 1913: 112–113, 1914: 231–241; Kuhn 1913: 1055; Lacordaire 1866: 379; Letzner 1891: 375; Nordlinger 1856: 29; Nunberg 1954: 50, 1956c: 160, 168; Nusslin 1911a: 112; Perris 1877a: 415; Pfeffer 1932b: 18, 1942a: 15, 1955a: 186; Portevin 1935: 325; Quaschik 1953: 35; Ratzeburg 1837: 163, 1839: 199; Redtenbacher 1849a: 791, 1849b: 26, 1858: 832, 1874: 375; Reitter 1894a: 72, 1913a: 69, 1916: 289; Rhumbler 1922: 334, 1927: 351; Rupertsberger 1879: 231, 1880: 229; Saalas 1913a: 69, 83; Schedl 1934f: 1639, 1952f: 87, 1964a: 99; Seidlitz 1872: 393, 1891a: 563, 1891b: 608; Sharp 1879: 101; Sokanovskii 1954: 17; Spesivtsev 1922: 471, 491, 1925a: 171, 1931: 47–48; Stark 1952: 289; Stierlin 1898: 438; Thomson 1868: 219, 1871: 393. (ms) Escherich 1932b.

grothii Hagedorn 1904a: 232 (*Cryphalus*). Syn-
types, sex?; Deutschl. [Osdorf near Hamburg];
Hamburg Museum, lost. Synonymy: Reitter
1913a: 69.

Notes: (3) Hansen 1955: 180 (treated this as a
good species).

References: (ay) Eichelbaum 1905: 248–250;
Klimesch 1914: 236, 238; Nürnberg 1928a:
140; Nusslin 1911a: 58, 89, 278, 337, 378. (cn)
Barbey 1925: 566. (cc) Kleine 1908c: 187,
1909a: 46, 1944: 82. (hb) Barbey 1925: 566;
Hagedorn 1904a: 232; Palm 1956: 63. (ds)
Bejer-Petersen & Jorinn 1977: 24; Bucking
1932; Hagedorn 1910d: 43; Hansen, V. 1956:
178, 1964: 462; Heyden, Reitter, & Weise
1906: 711; Horion 1960: 162; Jazentkovsky
1912: 288; Klefbeck & Sjöberg 1960: 230;
Kleine 1912a: 263, 267, 1913b, 1934a: 146;
Klimesch 1913: 106–116; Kozikowsky &
Kuntze 1925: 20; Lundgren 1988; Mahler
1987: 232; Munster 1922b; Negru 1965: 114;
Palm 1956: 63, 1959: 24, 350; Pfeffer 1989a:
85; Rohrig 1955: 37; Rydh 1977; Schilsky 1909:
188; Strand 1963: 91. (tx) Balachowsky 1949a:
215; Endrodi 1957b: 416; Hagedorn 1904a:
232, 1910a; Hansen, V. 1955: 180, 1956: 178,
1964: 462; Klimesch 1913: 106; Lundgren
1988; Nürnberg 1928a: 140; Nusslin 1911a: 58,
89, 278, 337, 378; Reitter 1913a: 69; Schedl
1934f: 1639.

spiculatus Eggers 1927d: 122. Lectotype ♂;
Klana (Kräin); USNM, Washington, desig-
nated by Anderson & Anderson 1971: 30. Syn-
onymy: Sokanovskii 1954: 17.

References: (bv) Grune 1979: 101. (cn)
Grandi 1951. (hb) Grandi 1951; Stark 1952:
285. (ds) Arnoldi et al. 1955: 695; Grune 1979:
101; Horion 1951; Kleine 1934a: 147; Postner
1974: 428; Reitter 1916: 350; Sainte-Claire &
Mequignon 1938: 446; Stark 1952: 285. (tx)
Anderson, W. H. & Anderson 1971: 30; Bala-
chowsky 1949a: 217; Eggers 1927d: 122; Grune
1979: 101; Nürnberg 1956c: 159–160, 168;
Pfeffer 1955a: 185, 188; Postner 1974: 428;
Reitter 1916: 350; Schedl 1934f: 1639, 1979c:
234; Sokanovskii 1954: 17; Stark 1952: 285.

berezinae Stark 1952: 285. Holotype, sex?; Arch-
angel [USSR]; IZL, Leningrad. Synonymy:
Sokanovskii 1954: 17.

References: (bv) Grune 1979: 101. (hb) Stark
1952: 285. (ds) Grune 1979: 101; Postner
1974: 428; Stark 1952: 285. (tx) Grune 1979:
101; Hansen, V. 1956: 179; Pfeffer 1955a: 185;
Postner 1974: 428; Sokanovskii 1954: 17; Stark
1952: 285.

dejeri Stark 1936: 152. Lectotype, sex?; Asiatic
Russia: Sejung Mt., Margassan R.; IZL, Leningrad,
designated by Michalski 1969a: 892.

Distribution: Asia (Sakhalin Island, Siberia in E
USSR).

Hosts: *Salix sachalinensis*.

Notes: (3) Original publication not seen by us;
Stark 1952: 284 spelled the name *deeri*.

References: (hb) Kurenzov & Kononov 1961: 600.
(ds) Krivolutskaya 1983; Kurenzov & Kononov
1961: 595; Stark 1936c: 152; Yanovskii & Tegsh-
zhiargal 1985: 409; Zinovjev 1955: 187. (tx) Krivo-
lutskaia 1955: 153; Kurenzov 1941a: 232;
Michalski 1969b: 567, 1970: 892; Pfeffer 1944:
131; Schedl 1952k: 158; Sokanovskii 1954: 17,
1960: 676; Stark 1936: 152, 1951: 299, 1952: 284.

dejeri Eggers 1942c: 31. Synatypes 2, sex?;
Baikalgebiet (Sajon), Siberia, USSR; Eggers
Collection, both in NHMW, Wien. Synonymy:
Pfeffer 1944a: 131.

References: (hb) Stark 1952: 284. (ds) Stark
1952: 284. (tx) Eggers 1942c: 31; Pfeffer
1944a: 131; Schedl 1952k: 158; Stark 1952:
284–299. (ms) Pfeffer 1944a: 131, 1979c: 76.

grandis Schedl 1964a: 99. Holotype ♀; Alger;
MNHN, Paris.

Distribution: Africa (Algeria, Morocco).

References: (tx) Schedl 1964a: 99.

granulatus (Ratzeburg) 1837: 164 (*Bostrichus*).
Synatypes, sex?; Passau, Linz, in Sweden; not
located.

Figures: Balachowsky 1949a: 210, 216, Grune
1979: 102.

Distribution: Africa (Egypt), Europe (Bulgaria/
Czechoslovakia/ Denmark/ England/ Finland/
France/ Germany/ Greece/ Hungary/ Italy/
Norway/ Poland/ Spain/ Sweden/ W USSR).

Hosts: *Populus alba*, *P. pyramidalis*, *P. tremula*.

References: (ay) Escherich 1923b: 483; Heeger
1866; Klimesch 1914: 218, 237. (bv) Grune 1979:
103; Hellen 1921. (cn) Chararas 1972; Escherich
1923b: 483; Grandi 1951; Hellen 1921; Nusslin
1913: 291; Rhumbler 1922: 334, 1927: 351;
Schwerdtfeger 1944a: 175, 1957a: 182; Wachtl
1901: 381. (cc) Balazy & Michalski 1960; Kleine
1908c; Krivolutskaya 1974; Krivosheina 1974;
Michalski & Ratajczak 1989; Nosek 1959a: 118,
1959b: 86; Nusslin 1927: 351; Pfeffer 1923a: 331,
1928b: 2; Ruhn 1956b: 4; Schwerdtfeger 1944a:
175, 1957a: 182. (hb) Altum 1881c: 322; Budge
1949; Dombrowsky 1887, 1889; Eichhoff 1881a:
46, 188; Escherich 1923b: 483; Grandi 1951;
Gyorfi 1957; Heeger 1866; Henschel 1895a: 170;
Karpinski 1933b: 28; Karpinski & Strawinski
1948: 155; Klimesch 1915: 6; Kostin 1960: 132;
Lengerken 1939: 63, 1954: 84; Munro 1926: 62;
Nosek 1959a: 118, 1959b: 86; Nusslin 1913: 291,
1927: 351; Palm 1949: 240, 1956: 63; Pfeffer
1942a: 15; Ratzeburg 1837: 164, 1839: 199;
Rhumbler 1922: 334, 1927: 351; Saalas 1913a: 69,
83; Schedl 1981b: 73; Schwerdtfeger 1944a: 175,
1957a: 182, 1981: 188; Stark 1926a: 334, 1952:

- 288; Wachtl 1876a: 458, 1901: 381. (**ds**) Allen, A. A. 1954, 1977; Arnoldi 1955: 694; Balazy & Michalski 1960; Barthe 1896; Buresh & Lazarov 1956; Calver 1884, 1893; Eggers 1904; Endrodi 1958b; Ericson & Sandin 1893; Escherich 1923: 483, 1932b; Forster 1849: 439; Fowler 1891; Gaubil 1849: 126; Gemminger & Harold 1872: 2683; Gozis 1875: 80; Grill 1895: 309; Grune 1979: 103; Hagedorn 1904e, 1904f, 1910d: 43; Hansen, V. 1939, 1956: 183; Hellen 1947; Henschel 1895a: 170; Heyden, Reitter, & Weise 1883: 181, 1891: 671, 1906: 711; Horion 1951; Kangas 1946a: 36; Karpinski 1931: 20, 27, 1932b: 54, 1933b: 28, 1948b: 230; Karpinski & Strawinski 1948: 155; Kestercanek 1881a: 12; Kleine 1912a: 263, 267, 1913a: 35, 1913b: 123, 1934a: 147; Kloft & Hinks 1945: 218; Kraatz 1869: 59; Kurenzov 1936b: 350; Lacordaire 1866: 379; Langhoffer 1915c: 158; Leclercq 1971; Lomnicki 1913b: 148; Lucht 1987: 278; Lundblad 1950c: 115; Matthews & Fowler 1883: 42; Munster 1925: 289; Negru 1965: 114, 1968c: 91; Nunberg 1928b: 88, 107, 1954: 50, 1956c: 157, 1960b: 156; Palm 1949: 240, 1950: 143, 1956: 63; Perris 1876a: 254, 1877a: 414; Pfeffer 1923b: 106, 1924a: 96, 1924b: 471, 1928b: 2, 1931b: 73, 1936: 90, 1989a: 85; Pittioni 1943: 176; Postner 1974: 428; Ratzeburg 1837: 164, 1839: 199; Redtenbacher 1858: 832, 1874: 375; Reitter 1894a: 72, 1916: 289; Rohrig 1955: 37; Roubal 1941: 264; Saalas 1913a: 69, 83, 1931: 67; Sainte-Claire & Mequignon 1938: 446; Schanfuss 1915: 1233; Schumm 1859: 96, 1862: 101; Schedl 1980a: 16, 1981b: 73; Scheerpeltz & Winkler 1930: 257; Schilsky 1890: 196, 1909: 188; Schwerdtfeger 1981: 188; Seidlitz 1872: 393, 1891a: 563; Sharp 1871: 84; Sharp & Fowler 1893: 34; Sokanovskii 1960; Stark 1926a: 334, 1926j: 126, 1952: 288; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 438; Teocchi 1965b; Tredl 1907: 13; Wachtl 1876a: 458; West 1938: 184; Wichmann 1927a: 68; Winter, T. G. 1983: 28; Yanovskii & Tegshzhargal 1985: 410. (**tx**) Allen, A. A. 1965b; Bach 1854; Balachowsky 1949a: 215; Berger 1917: 226–248; Doeblner 1860; Dombrowsky 1887, 1889; Duffy 1953; Eggers 1915c: 188, 1927d: 121, 1942c: 31; Eichhoff 1878b: 44, 137, 1881a: 46, 188, 1883a: 110, 133; Endrodi 1957b; Escherich 1923b: 483; Fauvel 1889; Ferrari 1967b: 113; Fleischer 1927; Formanek 1907: 36; Grune 1979: 102–103; Hagedorn 1910a: 86; Hansen, V. 1955, 1956; Henschel 1895a: 170; Hopkins 1915b: 36–37; Karpinski & Strawinski 1948: 155; Klimesch 1913: 106, 1914: 231–241, 1915: 6–13; Kihnt 1913: 1056; Kurenzov 1941: 232; Lacordaire 1866: 379; Lucht 1987: 278; Nunberg 1954: 50; Perris 1877a: 414; Pfeffer 1932b: 18, 1942a: 15, 1955a: 187; Portevin 1935: 325; Postner 1974: 428; Quaschik 1953: 35; Ratzeburg 1837: 164, 1839: 199; Redtenbacher 1849a: 792, 1849b: 26, 1858: 832, 1874: 375; Reitter 1894a: 72, 1913a: 67, 69–70, 1916: 289; Rhumbler 1922: 334, 1927: 351; Saalas 1913a: 69, 83, 1914: 23, 305; Schedl 1934f: 1639, 1964a: 99, 1980a: 16, 1981b: 73; Seidlitz 1872: 393, 1891a: 563, 1891b: 608; Sokanovskii 1954: 17, 1960: 671; Spessivtsev 1922a: 471, 491, 1925a: 172, 1931: 48; Stark 1952: 288; Stierlin 1898: 438; Thomson 1871: 393. (**ms**) Escherich 1932b; Hellen 1921. *tredli* Reitter 1908: 55. Syntypes, sex?, Agypien: Cairo; NHMB, Budapest. Synonymy: Hagedorn 1904f: 232, Reitter 1913: 69. References: (**hb**) Hagedorn 1904f: 232. (**ds**) Negru 1965: 114; Schilsky 1909: 188. (**tx**) Balachowsky 1949a: 216; Reitter 1908: 55, 1913a: 67; Schedl 1934f: 1639. *bispimulus* Eggers 1927d: 121. Syntypes, sex?, Finland: 2 in Eggers Collection, in NHMW, Wien. Synonymy: Sokanovskii 1954: 17, Nunberg 1956c: 159. Notes: (3) Hansen 1955: 182 (treated as a good species). References: (**bv**) Grune 1979: 103. (**cn**) Saalas 1949: 342, 372. (**cc**) Kangas 1940a, 1941, 1942a; Nuorteva 1957b: 68; Saalas 1949: 342, 372. (**hb**) Karpinski & Strawinski 1948: 155; Palm 1949: 239, 1950: 142; Saalas 1949: 342, 372; Stark 1952: 288. (**ds**) Arnoldi et al. 1955: 694; Grune 1979: 103; Hansen, V. 1939, 1956: 182; Hellen 1947; Kangas 1940: 32–34, 1941: 359–365, 1946a: 36, 1966d; Karpinski 1948a: 173; Karpinski & Strawinski 1948: 155; Klefbeck & Sjoberg 1960: 230; Lundberg 1981: 151; Nuorteva 1971: 71; Palm 1949: 239, 1950: 142, 1959: 24, 350; Pfeffer 1955a: 185; Postner 1974: 428; Stark 1952: 288; Yanovskii 1977a. (**tx**) Eggers 1927d: 121–122; Grune 1979: 103; Hansen, V. 1955, 1956; Karpinski & Strawinski 1948: 155; Nunberg 1956c: 157, 159, 167; Postner 1974: 428; Saalas 1949: 342, 372; Schedl 1934f: 1639, 1979c: 41; Sokanovskii 1954: 17; Stark 1952: 288. *klimeschi* Eggers 1915c: 188. Holotype ♂: Alai im Ferganagebiet (Russisch-Turkestan); Eggers Collection, in NHMW, Wien. Distribution: Asia (Turkestan in W USSR). Hosts: *Elaeagnus* sp., *Populus diversifolia*, *P. pruinosa*, *P. suaveolens*, *Tamarix* sp. References: (**cn**) Yagdyev 1984. (**hb**) Stark 1952: 287. (**ds**) Kadyrov 1988: 43, 1989; Stark 1952: 287. (**tx**) Eggers 1915c: 188, 1933f: 6; Schedl 1934f: 1639, 1979c: 134; Sokanovskii 1954: 22; Stark 1952: 287. *kurenzovi* Nunberg 1956d: 208. Lectotype, sex?, Ussuri, USSR; ISBN, Novosibirsk, USSR, and IZL, Leningrad/St. Petersburg, automatic, designated by Michalski 1969a: 893. Distribution: Asia (Japan/ Siberia, Ussuri in E USSR). Hosts: *Populus tremula*.

Notes: (1) Original spelling was *kurenzovi*; corrected here to conform to Dr. Kurenzov's spelling of his own name.

References: (tx) Krivolutsкая 1953; Michalski 1969b: 568; Nobuchi 1955c: 11; Numberg 1956d: 208; Schedl 1959r: 42, 1962p: 204.

populi Kurenzov 1941a: 30, 164. Lectotype, sex?; Ussuri, USSR; ISBN, Novosibirsk, USSR, and IZL, Leningrad/St. Petersburg, preoccupied by Hopkins 1915, designated by Michalski 1969a: 893.

References: (hb) Kurenzov 1948b: 113; Stark 1952: 286; Wood, S. L. 1953: 63, 1954a: 993. (ds) Keen 1929c: 29; Kleine 1934a: 147; Nobuchi 1966: 54; Stark 1952: 286; Wood, S. L. 1948: 45, 1951a: 128. (tx) Keen 1929c: 29; Kurenzov 1941a: 30, 163–164, 1948b: 113; Michalski 1969a: 893, 1969b: 567; Numberg 1956: 208; Schedl 1959r: 42; Stark 1952: 286; Wood, S. L. 1953: 63, 1954a: 993.

kurenzovi Schedl 1959r: 42. Lectotype, sex?; Ussuri, USSR; ISBN, Novosibirsk, USSR, and IZL, Leningrad/St. Petersburg, preoccupied by Numberg 1956, designated by Michalski 1969a: 893.

References: (tx) Michalski 1969a: 893; Schedl 1959r: 42.

niger Stark 1936e: 152. Lectotype, sex?; Far East: Vladivostok, Schkotov; IZL, Leningrad, designated by Michalski 1969a: 893.

Distribution: Asia (Siberia in E USSR).

Hosts: *Salix* sp.

References: (cc) Kurenzov 1934a: 50. (hb) Kurenzov 1935a: 19, 30, 1948b: 113; Stark 1952: 282. (ds) Kurenzov 1934a: 50, 1935a: 19, 30; Stark 1936e: 152, 1952: 282. (tx) Kurenzov 1941a: 163–164, 1948b: 113; Michalski 1969b: 568, 1970: 542; Pfeffer 1944a: 131; Sokanovskii 1954: 17; Stark 1936e: 152, 1952: 282. (ms) Pfeffer 1944a: 131.

palmi Hansen 1956: 183. Holotype ♂; Sweden: Ostergotland, Uppland, Jamtland; Hansen Collection.

Distribution: Europe (Finland/ Sweden).

Hosts: *Populus tremula*.

Notes: (1) This is probably a synonym of *granulatus*.

References: (hb) Palm 1956: 63. (ds) Hansen, V. 1956: 183; Kangas 1966d: 135; Karppinen 1966: 331; Klefbeck & Sjöberg 1960: 203; Nuorteva 1971: 71; Palm 1956: 63, 1959: 19, 24, 350. (tx) Hansen, V. 1955, 1956: 183.

populi Hopkins 1915b: 37. Holotype ♀; Williams, Arizona [USA]; USNM, Washington.

Figures: Krivolutsкая & Kupyanskaya 1970, Nobuchi 1966e: pl. 2.

Distribution: North America (Manitoba, New Brunswick, Saskatchewan in Canada/ Arizona, Colorado, Idaho, Montana, Nevada, Utah in USA).

Hosts: *Populus acuminata*, *P. angustifolia*, *P. tremuloides*, *P. trichocarpa*.

References: (bv) Petty, J. L. 1977. (cn) Doane et al. 1936. (cc) Tomalak, Welch, & Galloway 1989: 11. (hb) Doane et al. 1936; Furniss, M. M. & Johnson 1987: 380; Gast et al. 1989: 384; Harrison 1959: 80; Petty, J. L. 1977; Stewart, D. A., Booth, & Petty 1979; Wood, S. L. 1982b: 553. (ds) Atkinson & Equihua 1985c: 356; Bright 1976d: 109; Chamberlin 1939: 323; Furniss, M. M. & Johnson 1987: 380; Furniss, R. L. & Carolin 1977: 378; Gast et al. 1989: 384; Krivolutsкая & Kupyanskaya 1970; Nobuchi 1966c: 414, pl. 2; Wood, S. L. 1972a: 414, 1982b: 553. (tx) Bright 1976d: 109; Chamberlin 1939: 323; Hopkins 1915b: 36–37; Wood, S. L. 1951a: 128, 1954a: 990, 993, 1972a: 414, 1982b: 553. (ms) Stewart, D. A., Booth, & Petty 1979.

rybinskii Reitter 1894a: 72. Syntypes, sex?; Galizien: Tarnow; N11MB, Budapest.

Figures: Grune 1979: 98.

Distribution: Europe (Austria/ Corsica/ Czechoslovakia/ France/ Hungary/ Italy/ Poland/ Spain/ W USSR).

Hosts: *Alnus suaveolens*, *A. viridis*, *Salix silesiaca*.

References: (ay) Klimesch 1914: 218, 237. (bv) Grune 1979: 99. (cn) Nosek 1952b: 100; Nusslin 1913: 291; Rhumbler 1922: 334, 1927: 351; Wachtl 1901: 351. (cc) Kleine 1908c: 187, 1909a: 46, 76, 1944: 70, 1952b: 100, 1959a: 118; Nusslin 1927: 351; Pfeffer 1932a: 19, 1943b: 181; Rybinski 1897: 61. (hb) Formanek 1899; Karpinski & Strawinski 1948: 155; Klimesch 1915: 11; Lengerken 1939: 63, 1954: 84; Nosek 1959a: 118; Nusslin 1913: 291, 1927: 351; Pfeffer 1942a: 15; Rhumbler 1922: 334, 1927: 351; Stark 1952: 283; Wachtl 1901: 351. (ds) Endrodi 1958b; Escherich 1932b; Grune 1979: 99; Hagedorn 1910d: 45; Heyden, Reitter, & Weise 1906: 711; Horion 1951; Karpinski & Strawinski 1948: 155; Kleine 1913a: 35, 1913b: 123, 1934a: 147; Lomnicki 1913b: 148; Lucht 1987: 278; Numberg 1928b: 88, 1954: 50; Pfeffer 1930b: 119, 1931b: 73, 1935: 158, 1989a: 84; Pittioni 1943: 176; Reitter 1894a: 72, 1916: 289; Ronbal 1941: 264; Schaufuss 1915: 1233; Schedl 1980a: 16, 1981b: 73; Schilsky 1909: 188; Stark 1952: 283; Tredl 1907: 13. (tx) Balachowsky 1949a: 215–216; Endrodi 1957a: 307, 1957b; Fleischer 1927; Formanek 1907: 36; Grune 1979: 98–99; Hagedorn 1910a: 87; Karpinski & Strawinski 1948: 155; Lucht 1987: 278; Michalski 1969a: 893; Numberg 1954: 50; Pfeffer 1932b: 19, 1942a: 15, 1955a: 185; Postner 1974: 428; Reitter 1894a: 72, 1913a: 70, 1916: 289; Rhumbler 1922: 334, 1927: 351; Schedl 1934f: 1639, 1980a: 16, 1981b: 73; Schmidt, G. 1980: 26; Sokanovskii 1954: 17; Stark 1952: 283. (ms) Escherich 1932b; Schimitschek 1930b: 407. *corsicus* Eggers 1912b: 113. Holotype, sex?; Mt.

Renoso auf Corsica; Leonhard Collection in IPKE, Eberswalde. Synonymy: Balachowsky 1949a: 216.

References: **(ds)** Kleine 1913b: 123, 1934a: 147; Sainte-Claire 1914: 472; Sainte-Claire & Mequignon 1938: 446. **(tx)** Balachowsky 1949a: 216; Eggers 1912b: 113; Portevin 1935: 325; Reitter 1913a: 70; Schedl 1934f: 1639; Wichmann 1912c: 186.

rybinskii salicis Stark 1952: 283. Syntypes, sex?; USSR; IZL, Leningrad, preoccupied by Hopkins 1915.

References: **(tx)** Stark 1952: 283.

salicis Hopkins 1915b: 36. Holotype ♀; Del Monte, California [USA]; USNM, Washington.

Distribution: North America (California, Oregon, Washington in USA).

Hosts: *Alnus* sp., *Salix* sp.

References: **(cn)** Doane et al. 1936. **(hb)** Wood, S. L. 1953: 60, 1954a: 991. **(ds)** Bright 1976d: 109; Chamberlin 1939: 323, 1958: 131; Furniss, R. L. & Carolin 1977: 378; Keen 1934a: 147; Wood, S. L. 1972a: 414, 1982b: 552. **(tx)** Bright 1976d: 109; Chamberlin 1939: 323, 1958: 131; Hopkins 1915b: 36; Wood, S. L. 1953: 60, 1954a: 959, 979, 990–991, 1972a: 414, 1982b: 552.

concentralis Hopkins 1915b: 36. Holotype ♂; Easton, Washington [USA]; USNM, Washington. Synonymy: Wood 1954a: 991.

References: **(hb)** Chamberlin 1939: 324; Wood, S. L. 1953: 60. **(ds)** Chamberlin 1939: 324; Patterson & Hatch 1945: 152. **(tx)** Chamberlin 1939: 324; Hopkins 1915b: 36; Wood, S. L. 1953: 60, 1954a: 991.

striatulus (Mannerheim) 1853: 235 (*Cryphalus*). Holotype, sex?; Kenai Peninsula, Alaska; not in Helsinki Museum, presumably lost.

Figures: Bright 1976d: 202, 209 (adult).

Distribution: North America (Alaska/ Newfoundland, Nova Scotia, Quebec, Yukon in Canada/ Colorado, Idaho, Minnesota, Oregon, Utah in USA).

Hosts: *Alnus crispa*, *A. rugosa*, *Salix scouleriana*. References: **(cn)** Felt 1901: 516, 1906: 376, 650, 673; Swaine 1918a: 90. **(hb)** Chamberlin 1939: 315; Schwarz 1894b: 255; Swaine 1918a: 90; Wood, S. L. 1954a: 996, 1982b: 551. **(ds)** Blatchley & Leng 1916: 606; Bright 1976d: 109; Chamberlin 1925, 1939: 315; Elias 1985: 39; Furniss, R. L. & Carolin 1977: 378; Gemminger & Harold 1872: 2683; Hagedorn 1910d: 45; Hamilton 1894c: 35; Henshaw 1885: 148; Kleine 1913b: 118, 1914b: 397, 1934a: 142; Lacordaire 1866: 379; Swaine 1909: 92; Wood, S. L. 1982b: 551. **(tx)** Blatchley & Leng 1916: Bright 1976d: 109, 202, 209; Chamberlin 1939: 315; Eichhoff 1878b: 147; Ferrari 1867a: 17; Hagedorn 1910a: 99; Hopkins 1915b: 33; Lacordaire 1866: 379; LeConte 1868: 177, 1876: 362; Mannerheim 1853: 235; Swaine 1909: 92–93, 1912a: 349, 1918a: 89–90;

Wood, S. L. 1954a: 989–990, 996, 1969c: 115, 1973c: 186, 1982b: 551.

nitidus Swaine 1912a: 349. Lectotype, sex?; Weymouth, Nova Scotia [Canada]; CNCI, Ottawa, designated by Bright 1967b: 679. Synonymy: Wood 1969c: 115.

References: **(cn)** Swaine 1918a: 90. **(hb)** Chamberlin 1939: 323, 1958: 131; Swaine 1918a: 90; Wood, S. L. 1953: 57, 1954a: 990. **(ds)** Beaulne 1956: Blatchley & Leng 1916: 605; Chamberlin 1939: 323, 1958: 131; Dodge 1983: Kleine 1913b: 123, 1914b: 399, 1934a: 147; Wood, S. L. 1972a: 414. **(tx)** Blatchley & Leng 1916: 605; Bright 1967b: 679; Chamberlin 1939: 323, 1958: 131; Dodge 1983: 16, 39–40; Hoebeke 1978: de Ruette 1970: 114; Swaine 1912a: 349–350, 1918a: 90; Wood, S. L. 1953: 57, 1954a: 959, 990, 1969c: 115, 1972a: 414, 1973c: 186.

punctipennis Hopkins 1915b: 37. Holotype ♀; Alta, Utah [USA]; USNM, Washington. Synonymy: Wood 1954a: 990.

References: **(hb)** Chamberlin 1939: 323; Wood, S. L. 1953: 58. **(ds)** Chamberlin 1939: 323; Wood, S. L. 1948: 44, 1951a: 128. **(tx)** Chamberlin 1939: 323; Hopkins 1915b: 36–37; Wood, S. L. 1951a: 126, 1953: 58, 1954a: 990.

thatcheri (Wood) 1954a: 990, 994 (*Cryphalus*). Holotype ♀; 2 miles NW Blue Lake, Lassen County, California [USA]; SMUK, Lawrence.

Distribution: North America (British Columbia in Canada/ California, N Idaho, Oregon in USA).

Hosts: *Populus tremuloides*, *P. trichocarpa*.

References: **(hb)** Bright & Stark 1973: 66; Wood, S. L. 1953: 66, 1954a: 994. **(ds)** Bright 1976d: 110; Bright & Stark 1973: 66; Furniss, R. L. & Carolin 1977: 389; Wood, S. L. 1982b: 554. **(tx)** Bright 1976d: 110; de Ruette 1970: 100; Wood, S. L. 1953: 66, 1954a: 990, 994, 1982b: 554.

tremulae Stark 1952: 287. Lectotype, sex?; Crimea, USSR; IZL, Leningrad, designated by Michalski 1969a: 893.

Distribution: Europe (Crimea in W USSR).

Hosts: *Populus tremula*.

Notes: (3) Sokanovskii 1954: 17 (treated as a synonym of *granulatus*).

References: **(bv)** Grune 1979: 101. **(hb)** Stark 1952: 287. **(ds)** Grune 1979: 101; Pfeffer 1989a: 85; Postner 1974: 428; Stark 1952: 289. **(tx)** Grune 1979: 101; Michalski 1969b: 568, 1969a: 893; Pfeffer 1955a: 185; Postner 1974: 428; Sokanovskii 1954: 17; Stark 1952: 287.

Genus *Procryphalus* Hopkins

PROCYPHALUS HOPKINS 1915b: 7, 33. Type-species:

Procryphalus populi Hopkins, original designation.

Keys: Hopkins 1915b: 33, Wood 1954a: 982, 1982b: 559 for North America.

References: **(hb)** Bright & Stark 1973: 67; Wood,

S. L. 1986a: 89. (**ds**) Bright & Stark 1973: 67; Wood, S. L. 1982b: 859, 1986a: 89. (**tx**) Arnett 1960: 1043, 1965: 1043; Balachowsky 1949a: 201; Bright 1976d: 111; Hopkins 1915b: 7, 33–34; Swaine 1918a: 45, 90; Wood, S. L. 1954a: 981, 1982b: 859, 1986a: 89.

fraxini (Berger) 1916: 235 (*Ernoporius*). Lectotype, sex?, Ussuri, USSR; IZL, Leningrad, designated by Michalski 1969a: 892.

Figures: Tsai & Li 1959: 92.

Distribution: Asia (Heilongjiang, Jilin in China/ Ussuri in E USSR).

Hosts: *Fraxinus mandshurica*.

References: (**cn**) Kurenzov 1935c: 189, 1951c: 79. (**ce**) Kurenzov 1934a: 50, 1951d: 24. (**hb**) Kurenzov 1935a: 19, 33, 1948b: 128, 1951d: 24; Stark 1952: 273. (**ds**) Kleine 1934a: 147; Kurenzov 1934a: 50, 1935a: 19, 33, 1935c: 189, 1936a: 110, 1936b: 350, 1935a: 59, 1940: 77; Stark 1952: 273; Yin, Huang, & Li 1984: 118. (**tx**) Berger 1916: 235; Kurenzov 1941a: 156, 1948b: 128; Michalski 1969a: 892; Schedl 1934f: 1639, 1962r: 91; Stark 1952: 273; Tsai & Li 1959: 92.

mucronatus (LeConte) 1879: 518 (*Cryphalus*). Holotype ♀; La Veta Pass, Colorado [USA]; MCZ, Cambridge.

Figures: Kusch 1967: 10.

Distribution: North America (Alaska/ Alberta, British Columbia in Canada/ Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah in USA).

Hosts: *Populus tremuloides*.

References: (**bv**) Petty, J. L. 1977. (**hb**) Chamberlin 1939: 322, 1955: 136; Furniss, M. M. & Johnson 1987: 379; Gast et al. 1989: 384; Petty, J. L. 1977; Wood, S. L. 1953: 46, 1954a: 984, 1982b: 859. (**ds**) Bright 1976d: 111; Chamberlin 1939: 322, 1958: 136; Furniss, M. M. & Johnson 1987: 379; Furniss, R. L. & Carolin 1977: 377; Gast et al. 1989: 384; Hagedorn 1910d: 44; Henshaw 1882: 269, 1885: 148; Kleine 1913b: 118; Kusch 1967; Schwarz 1886: 42; Swaine 1909: 92; Wickham 1896a: 309; Wood, S. L. 1972a: 414. (**tx**) Bright 1976d: 111; Chamberlin 1939: 322, 1958: 136; Hagedorn 1910a: 87; Hopkins 1915b: 33; Kusch 1967: 10; LeConte 1879: 518; Schwarz 1886: 42; Swaine 1909: 92; Wood, S. L. 1953: 46, 1954a: 959, 984, 1982b: 859.

idahoensis Hopkins 1915b: 34. Holotype ♀; Beaver Canyon, Idaho [USA]; USNM, Washington. Synonymy: Wood 1954a: 985.

References: (**hb**) Chamberlin 1939: 321. (**ds**) Chamberlin 1939: 321. (**tx**) Chamberlin 1939: 321; Hopkins 1915b: 33–34; Wood, S. L. 1954a: 959, 984.

populi Hopkins 1915b: 34. Holotype ♂; Tercio, Colorado [USA]; USNM, Washington. Synonymy: Wood 1954a: 985.

References: (**cn**) Doane et al. 1936. (**hb**)

Chamberlin 1939: 321; Doane et al. 1936. (**ds**) Chamberlin 1939: 321; Keen 1929c: 29; Kleine 1934a: 148; Wood, S. L. 1945: 42, 1951a: 128. (**tx**) Balachowsky 1949a: 201; Chamberlin 1939: 321; Hopkins 1915b: 33–34; Keen 1929: 29; Wood, S. L. 1951a: 128, 1954a: 959, 982–984.

utahensis Hopkins 1915b: 33. Holotype ♀; Alta, Utah [USA]; USNM, Washington.

Figures: Bright 1976d: 202, 209 (adult).

Distribution: North America (Alaska/ British Columbia, Quebec in Canada/ California, Idaho, Oregon, South Dakota, Utah in USA).

Hosts: *Salix scouleriana*, *S.* sp.

Notes: Hopkins 1915b: 33 (reported from *Acer macrophyllum*, an error); breeds only in *Salix*.

References: (**hb**) Bright & Stark 1973: 67; Chamberlin 1939: 321, 1958: 136; Wood, S. L. 1953: 44, 1954a: 983. (**ds**) Bright 1976d: 111; Bright & Stark 1973: 67; Chamberlin 1939: 321, 1958: 136; Furniss, R. L. & Carolin 1977: 378; Keen 1929c: 29; Wood, S. L. 1951a: 128, 1972a: 414, 1982b: 859. (**tx**) Bright 1976d: 111, 202, 209; Chamberlin 1939: 321, 1958: 136; Hopkins 1915b: 33; Keen 1929c: 29; Wood, S. L. 1951a: 128, 1953: 44, 1954a: 959, 979, 983, 1972a: 414, 1979a: 22, 1982b: 859. *salicis* Hopkins 1915b: 33. Holotype ♀; Black Hills, South Dakota [USA]; USNM, Washington. Synonymy: Wood 1954a: 983.

References: (**cn**) Doane et al. 1936. (**hb**) Chamberlin 1939: 321; Doane et al. 1936. (**ds**) Chamberlin 1939: 321. (**tx**) Chamberlin 1939: 321; Hopkins 1915b: 33; Wood, S. L. 1954a: 983. *aceris* Hopkins 1915b: 33. Holotype ♀; Albany, Oregon [USA]; USNM, Washington. Synonymy: Wood 1975a: 22.

References: (**cn**) Doane et al. 1936. (**hb**) Chamberlin 1939: 321, 1958: 136; Doane et al. 1936; Wood, S. L. 1953: 42, 1954a: 982. (**ds**) Chamberlin 1917: 355, 1939: 321, 1958: 136; Keen 1929c: 29; Kleine 1934a: 148; Wood, S. L. 1972a: 414. (**tx**) Chamberlin 1939: 321, 1958: 136; Hopkins 1915b: 33; Keen 1929c: 29; Wood, S. L. 1953: 42, 1954a: 979, 982, 1972a: 414, 1975a: 22.

Genus *Ernoporiscus* Berger

ERNOPORISCUS BERGER 1917: 242. Type-species:

Ernoporiscus spessitzzevi Berger, monobasic.

Eocryphalus Kurenzov 1941a: 161, 230. Type-species: *Eocryphalus semenovi* Kurenzov, monobasic. Synonymy: Schedl 1962r: 93.

References: (**tx**) Krivolutskaya 1958: 148; Kurenzov 1941a: 161, 230; Schedl 1962r: 93–94.

Ernopocerus Balachowsky 1949a: 211. Type-species: *Ernoporiscus caucasicus* Lindemann, subsequent designation by Wood 1954a: 986. Synonymy: Wood 1980c: 93.

References: (**tx**) Balachowsky 1949a: 211;

- Nunberg 1958: 485; Schedl 1958k: 142, 1962r: 90; Wood, S. L. 1954a: 980.
- Keys: Balachowsky 1949a: 211, Reitter 1913a: 68, Stark 1952: 271 for Asia and Europe.
- Notes: (3) Kurenzov specimens of *Hypothenemus corni* deposited in the Wood Collection are of *Ernoporicus*.
- References: (ay) Nobuchi 1969a: 57. (hb) Wood, S. L. 1982b: 558, 1986a: 89. (ds) Wood, S. L. 1982b: 558, 1986a: 89. (tx) Arnett 1960: 1043, 1968: 1043; Balachowsky 1949a: 201, 211–213; Berger 1917: 242; Kurenzov 1941a: 161–162, 231; Nunberg 1958: 485; Pfeffer 1989a: 82; Schedl 1960e: 104, 1962r: 95, 1981b: 75; Sokanovskii 1959: 277; Wood, S. L. 1982b: 558, 1986a: 89.
- africanus** Schedl 1977c: 396 (*Stephanorhopalus*). Holotype, sex?; Sudafrika: Natal; Schedl Collection in NHMW, Wien.
- Distribution: Africa (Natal in South Africa).
- Notes: (1) Wood 1953 (notes, p. 13, to *Ernoporicus*, type examined).
- References: (tx) Schedl 1977c: 396.
- aluiphagus** Nobuchi 1975: 43. Holotype, sex?; Konseitoge, Gunma pref., Japan; Nobuchi Collection, Iwabaki.
- Figures: Nobuchi 1975: pl. 1, fig. 3.
- Distribution: Asia (Japan).
- Hosts: *Alnus hirsutus*.
- References: (tx) Nobuchi 1975: 43.
- ater** Nobuchi 1975: 43. Holotype, sex?; Sogoya, Chichibu, Saitama pref., Japan; Nobuchi Collection, Iwabaki.
- Figures: Nobuchi 1975: pl. 1, fig. 2.
- Distribution: Asia (Japan).
- Hosts: *Pterocarya rhoifolia*.
- References: (tx) Nobuchi 1975: 42–43.
- capucinus** (Schedl) 1971a: 283 (*Cryphalops*). Holotype, sex?; Central Prov.: Hantane Estate, Kandy; Schedl Collection in NHMW, Wien.
- Distribution: Asia (Sri Lanka).
- References: (tx) Schedl 1971a: 283, 1979c: 52.
- caucasicus** (Lindemann) 1875c: 373 (*Ernoporus*). Syntypes, sex?; Piatigorsk, nordlichen Kaukasus; not located.
- Figures: Grune 1979: 108, Pfeffer 1989a: pl. 6.
- Distribution: Asia (Turkey/ E USSR), Europe (Austria/ Belgium/ Czechoslovakia/ Denmark/ England/ France/ Germany/ Greece/ Hungary/ Italy/ Poland/ Spain/ Sweden/ W USSR).
- Hosts: *Tilia intermedia*, *T. parvifolia*, *T. rubra*, *T. tomentosa*, rare in *Hibiscus syriacus*, *Ulmus montana*.
- Notes: (1) Balachowsky 1949a: 209 (to *Ernoporicus*).
- References: (bv) Grune 1979: 109. (cn) Koppen 1882: 252; Nusslin 1913: 291; Pierce, W. D. 1917: 141; Rhumbler 1922: 334, 1927: 350; Wachtl 1901: 351. (ec) Kleine 1908c: 185; Nosek 1959a: 118, 1959b: 87; Nusslin 1927: 350. (hb) Fuchs 1904a; Györfi 1957; Karpinski & Strawinski 1948: 155; Lengerken 1939: 63, 1954: 83; Nosek 1959a: 118, 1959b: 87; Nusslin 1913: 291, 1927: 350; Pfeffer 1942a: 18; Postner 1974: 431; Rhumbler 1922: 334, 1927: 350; Stark 1952: 276; Vinogradov-Nikitin 1911: 1472–1477; Wachtl 1901: 351; Wichmann 1910b: 210. (ds) Allen, A. A. 1970; Bakke 1963a; Bedel 1888b: 398, 413; Bejer-Petersen & Jorun 1977: 23; Borchert 1951; Brakman 1966b: 206; Endrodi 1958b; Ermisch 1953; Escherich 1932b; Fuchs 1904a; Grune 1979: 109; Hagedorn 1904e, 1910d: 41; Hansen, V. 1939, 1956, 1964: 462; Hellen 1947; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1951; Karpinski & Strawinski 1948: 155; Klefbeck & Sjöberg 1960: 230; Kleine 1912a: 303, 1913a: 34, 1934a: 147; Koppen 1882: 282; Leclercq 1971; Lomnicki 1913b: 148; Lovendal 1890c: 210; Lucht 1987: 278; Mahler 1987: 232; Nunberg 1928b: 88, 106, 1954: 54; Palm 1959: 29, 130, 1960: 350; Pfeffer 1924b: 471, 1931b: 74, 1935: 160, 1989a: 83; Pierce, W. D. 1917: 141; Pittioni 1943: 175; Pjatnitskii 1930a: 165; Postner 1974: 431; Reitter 1894a: 71, 1916: 289; Roubal 1935b: 72, 1941: 264; Sainte-Claire & Mequignon 1938: 445; Schaufuss 1915: 1232; Schedl 1961b: 187, 1980a: 15, 1981b: 75; Schilsky 1909: 188; Stark 1927b: 88, 1952: 276; Tredl 1907: 12; West 1983: 184; Wichmann 1910b: 210, 1927a: 66; Winter, T. G. 1983: 11. (tx) Allen, A. A. 1970: 246; Bakke 1964: 121; Balachowsky 1949a: 211–212; Bedel 1887: 191, 1888b: 398, 414; Eggers 1923b: 137; Endrodi 1957a: 307, 1957b; Fleischer 1927; Formanek 1907: 35; Grune 1979: 108–109; Hagedorn 1910a: 85; Hansen, V. 1956, 1964: 462; Karpinski & Strawinski 1948: 155; Kulmt 1913: 1055; Lindemann 1875c: 321; Lovendal 1890c: 210, 1898: 128; Lucht 1987: 278; Nunberg 1954: 54; Pfeffer 1932b: 18, 1942a: 18, 1955a: 181, 1989a: pl. 6; Pjatnitskii 1932: 295–302; Postner 1974: 431; Reitter 1894a: 71, 1913a: 68–69, 1916: 289; Rhumbler 1922: 334, 1927: 350; Schedl 1934f: 1639, 1958k: 142, 1962r: 90, 1980a: 15, 1981b: 75; Semenov 1902: 271; Spessittsev 1922a: 473, 491, 1931: 50; Stark 1952: 276; Wood, S. L. 1954a: 986. (ms) Eggers 1910b; Escherich 1932b.
- schreineri** Eichhoff 1881a: 185 (*Cryphalus*). Syntypes, sex?; Hinterpommern; Hamburg Museum, lost. Synonymy: Bedel 1887: 191, Eggers 1923b: 137.
- References: (hb) Dombrowsky 1887; Eichhoff 1881a: 46, 185; Hagedorn 1903a; Henschel 1895a: 168. (ds) Barthe 1896; Grill 1895: 310; Hagedorn 1903a, 1910d: 45; Henschel 1895a: 168; Heyden, Reitter, & Weise 1883: 182, 1891: 671; Kleine 1913a: 85, 1934a: 147; Lovendal 1890c: 210; Pfeffer 1935: 160; Schilsky 1909: 188; Seidlitz 1891a: 562, 1891b: 608. (tx) Balachowsky 1949a: 213; Bedel 1887: 191; Dombrowsky 1887; Eggers 1923b: 137; Eichhoff 1881a: 46, 185, 1883a: 109, 133;

Fauvel 1857, 1859; Hagedorn 1910a: 58; Henschel 1895a: 168; Hopkins 1915b: 35; Loven-
dal 1890c: 210; Schedl 1934f: 1639, 1979c: 222;
Seidlitz 1891a: 562, 1891b: 605. (**ms**) Eggers
1910a, 1910b.

ceylonicus (Schedl) 1959a: 475 (*Ptilopodius*). Syn-
types 2, sex?; Ceylon: Sabargamiva, Millawitiya
Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka).
References: (**tx**) Schedl 1959a: 475, 1979c: 56.

eximius (Schedl) 1942d: 9 (*Erischidius*). Holotype,
sex?; Ost-Java; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Java).
Notes: (3) Schedl 1963i: 64 (cited in *Ernoporus*).
References: (**tx**) Schedl 1942d: 9, 1963i: 64,
1979c: 93.

fagi (Fabricius) 1798: 157 (*Apatē*). Holotype, sex?;
Saxoniae; UZMC, Copenhagen, missing from its
pin, apparently lost.
Figures: Postner 1974: 431 (adult), Pfeffer
1959a: pl. 6.
Distribution: Asia (Turkey), Europe (Austria/ Bel-
gium/ Bulgaria/ Czechoslovakia/ Denmark/
England/ France/ Germany/ Greece/ Hungary/ N
Italy/ Netherlands/ Norway/ Poland/ Romania/
Sweden/ Switzerland/ W USSR/ Yugoslavia).
Hosts: *Fagus orientalis*, *F. sylvatica*, less common
in *Betula* spp., *Carpinus betulus*, *Quercus cerris*.
References: (**ay**) Burrini et al. 1888; Escherich
1923b: 482, 510; Reichenbach-Klinke 1953: 338.
(**bv**) Grune 1979: 109. (**cn**) Barbey 1925: 508;
Chorbadzhievo 1929; Escherich 1923b: 482, 510;
Georgescu et al. 1957: 357, 407; Gradojevic 1940;
Grandi 1951; Koppen 1882: 252; Kovacevic 1924:
21–22; Marcu 1926c: 64; Nosek 1951: 107, 1952b:
98; Nusslin 1913: 291; Pierce, W. D. 1917: 40;
Puecher Passavalli 1931b: 657; Rhumbler 1922:
334, 1927: 350; Schimitschek 1937c: 52, 1955a:
139, 1955c: 83; Schwerdtfeger 1944a: 174, 1957a:
182; Wachtl 1901: 381; Wichmann 1927b: 361.
(**ce**) Apel 1983; Balazy & Michalski 1964b; Benick
1952; Cooreman 1963: 45; Elliot & Morley 1907;
Gillanders 1906; Hirschmann 1960, 1971a, 1971b;
Hirschmann & Wisniewski 1982, 1983; Kleine
1908c: 185, 1909a: 46, 77, 1944: 73; Michalski &
Ratajczak 1989; Nosek 1951: 107, 1952b: 98,
1959b: 87; Novak, P. 1952: 414; Nusslin 1927:
350; Pfeffer 1923a: 331, 1928b: 9; Ratzeburg
1869a: 176; Rondani 1873: 146; Ruhm 1956b: 4;
Schimitschek 1930a: 281, 1955a: 139; Schwerdt-
feger 1944a: 174, 1957a: 182; Sedlacek 1915a:
45; Wanka 1920: 202–203; Westerboer 1963: 349.
(**hb**) Altum 1881c: 322; Apel 1983; Bach 1864;
Barbey 1901: 23, 73, 1925: 508; Beffa 1961;
Bnddenberg 1885: 97; Cecconi 1906, 1924; Chor-
badzhievo 1929; Dallimore & Munro 1922;
Dombrowsky 1887; Eichhoff 1881a: 46, 184;
Escherich 1923b: 482, 510; Everts 1903: 756;
Fuchs 1904a; Gillanders 1906, 1908; Grandi 1951;

Gyorfi 1957; Hagedorn 1903a; Henschel 1895a:
167; Karpinski & Strawinski 1948: 155; Karsch
1883: 142; Kemner 1919: 176; Knotek 1894a:
558; Lengerken 1939: 63, 1954: 83; Masutti 1964;
Munro 1926: 62; Nordlinger 1856: 26; Nosek
1959b: 87; Nunberg 1929: 105; Nusslin 1898:
280, 1906b: 14, 1913: 291, 1927: 350; Palm 1954a:
184; Pfeffer 1942a: 6; Postner 1974: 431;
Rhumbler 1922: 334, 1927: 350; Schedl 1951b:
75; Schimitschek 1930a: 281, 1955a: 139;
Schwerdtfeger 1944a: 174, 1957a: 182, 1981:
188; Stark 1952: 276; Tschorbadjiev 1929: 166;
Wachtl 1876a: 458, 1901: 381; Wichmann 1927b:
361. (**ds**) Acloque 1896; Andras & Schaefer 1957;
Bangsholt 1975: 95; Barthe 1896; Bau 1888;
Bejer-Petersen & Jorun 1977: 24; Bennett 1902,
1902b: 76; Blanchere & Robert 1889; Borchert
1951; Brakman 1966b: 206; Brancsik 1906; Buck
1955b: 191; Buresh & Lazarov 1956; Calver 1884,
1893; Cecconi 1897, 1906; Chapuis & Candeze
1853; Chorbadzhievo 1924d, 1929; Crotch 1863;
Dallimore & Munro 1922: 189–193; Dejean 1821,
1825, 1837; Duftschmidt 1825; Eggers 1904;
Endrodi 1958a, 1958b; Ericson & Sandin 1893;
Escherich 1923b: 482, 510, 1932b; Everts 1922:
642; Fleischer 1888; Fowler 1891; Fricken 1889:
343; Fuchs 1904a, 1905a; Gaubil 1849: 126; Gem-
minger & Harold 1872: 2682–2683; Gorham
1860; Cozis 1875: 80; Grill 1895: 310; Hagedorn
1910d: 42; Hansen, V. 1939, 1956, 1964: 462;
Hellen 1947; Henschel 1895a: 167; Heyden 1879:
140; Heyden, Reitter, & Weise 1883: 182, 1891:
671, 1906: 711; Holdhaus & Deubel 1910: 145;
Horion 1951; Jazentkovsky 1912: 288; Kaltenbach
1874: 624; Karpinski 1948b: 230; Karpinski &
Strawinski 1948: 155; Kemner 1919: 176; Kersten
1933: 74; Klefbeck & Sjoberg 1960: 230; Kleine
1912a: 263, 267, 1913a: 34, 1934a: 147; Kloft &
Hinks 1945: 218; Knotek 1892a: 37, 1894a: 558;
Koltze 1901: 153; Koppen 1882: 252; Kozikowsky
1921: 150; Kraatz 1869: 59; Kunr 1947c: 31;
Lacordaire 1866: 379; Leclercq 1971; Lomnicki
1913b: 148; Lucht 1987: 278; Lundberg 1988;
Mahler 1987: 232; Marcu 1926c: 64; Matthews &
Fowler 1883: 42; Munro 1920a: 257; Negrn
1966b: 401, 1968a: 456; Nosek 1952: 414; Novak,
P. 1952: 414, 1964; Nunberg 1928b: 88, 1954: 54,
1960b: 156; Nusslin 1898: 280; Orest 1926c: 64;
Palm 1959: 32, 130, 1960: 350; Perris 1876a: 254,
1877a: 414; Pfeffer 1924b: 472, 1928b: 9, 1931b:
74, 1950b: 76, 1989a: 83; Pierce, W. D. 1917: 40;
Pittioni 1943: 175; Postner 1974: 431; Prossen
1913: 83; Rapp 1934: 725; Redtenbacher 1874:
376; Reitter 1869b: 154, 1894a: 71, 1911: 69,
1916: 289; Roubal 1941: 264; Schaufuss 1915:
1232; Schumm 1859: 96, 1862: 101; Schedl 1971b:
530, 1980a: 15, 1981b: 75; Schilsky 1909: 188;
Schiodte 1873: 102; Schwerdtfeger 1981: 188;
Seidlitz 1872: 394, 1891a: 562, 1891b: 608; Sharp
& Fowler 1893: 34; Stark 1952: 276; Stein 1868:

114; Stein & Weise 1877: 164; Stierlin 1898: 439; Stierlin & Gautard 1871: 292; Sturm 1826: 102; Thomson 1865: 360, 1868: 220; Tredl 1907: 12; Tschorbadjev 1929: 166; Wachtl 1876a: 458; West 1938: 184; Westhoff 1882: 235; Wichmann 1927a: 66; Winter, T. G. 1983: 11. (**tx**) Acloque 1896; Allen, A. A. 1970; Apel 1983; Bach 1854, 1864; Balachowsky 1949a: 211; Barbey 1901: 23, 73; Belfa 1961; Bertolini 1872; Chapuis & Candèze 1853; Chorbadzhievo 1924d; Dejean 1821, 1825; Doebner 1860, 1868: 368; Dombrowsky 1887; Duftschmidt 1825; Eggers 1911a, 1929e, 1939b: 5; Eichhoff 1864: 35, 1868d: 422, 1876a: 378, 1878b: 132, 1881a: 46, 184, 1883a: 109, 133; Endrodi 1957b; Erichson 1836: 62; Escherich 1923b: 482, 510; Everts 1903: 756, 1922: 642; Fabricius 1798: 157, 1801: 383; Fairmaire 1864: 159; Fauvel 1887; Ferrari 1867a: 11, 13, 15, 1867b: 114, 1868: 255–256; Fleischer 1905, 1927; Formanek 1907: 35; Fricken 1889: 343; Gebien 1907: 197; Gillanders 1908; Grune 1979: 109; Hagedorn 1910a: 86; Hansen, V. 1956, 1964: 462; Henschel 1895a: 167; Hopkins 1915b: 35, 1915c: 221; Houlbert 1922a: pl. 1; Jacquelin du Val & Fairmaire 1868: 104; Karpinski & Strawinski 1948: 155; Knotek 1892a: 37; Lacordaire 1866: 379; Letzner 1891: 375; Lindemann 1875c: 321; Lovendal 1889b: 52, 1898: 126; Lucht 1987: 2781; Lundberg 1988; Negrin 1966b: 401; Nordlinger 1847: 21, 1848: 242, 1856: 26; Numberg 1954: 54; Perris 1877a: 414; Pfeffer 1932b: 18, 1942a: 6, 1955a: 181, 1989: pl. 6; Portevin 1935: 325; Quaschik 1953: 35; Redtenbacher 1849a: 851, 1849b: 26, 1874: 387; Reitter 1894a: 71, 1913a: 69, 1916: 289; Rhumbler 1922: 334, 1927: 350; Schedl 1934f: 1639, 1952f: 87, 1958k: 142, 1962r: 90, 1980a: 15, 1981b: 75; Schimitschek 1937c: 52, 1955c: 83; Schwerdtfeger 1981: 188; Seidlitz 1872: 394, 1891a: 562, 1891b: 608; Spessivtsev 1922a: 472, 491, 1925a: 174, 1931: 47, 50; Stark 1952: 276; Stierlin 1898: 439; Thomson 1865: 360, 1868: 220. (**ms**) Eichhoff 1868d: 422; Escherich 1932b.

serratus Panzer 1795a: 288 (*Bostrichus*). Syn-types, sex?; Germania: Panzer Collection, lost. Synonymy: Eggers 1911a: 73.

References: (**hb**) Wachtl 1876a: 458. (**ds**) Beck 1817; Calwer 1884; Hagedorn 1910d: 116; Kleine 1913b: 169; Sainte-Claire & Mequignon 1938: 445; Wachtl 1876a: 458. (**tx**) Balachowsky 1949a: 211; Bechstein et al. 1805: 104; Eggers 1911a: 73; Hagedorn 1910a; Panzer 1795a: 288.

thomsoni Ferrari 1867a: 14 (*Ernoporus*). Syn-types, sex?; Europa; presumably NHMW, Wien, not found. Synonymy: Ferrari 1868: 255. References: (**ds**) Bedel 1888b: 398, 414; Heyden 1876: 299; Kraatz 1869: 59; Stein & Weise 1877: 164. (**tx**) Balachowsky 1949a: 211; Bedel 1888b: 394, 414; Dallimore 1868;

Doebner 1868: 368; Eichhoff 1868d: 422, 1876a: 378–379; Ferrari 1867a: 12, 14, 1867b: 114, 1868: 255; Letzner 1891: 375.

kanachae (Hopkins) 1915b: 35 (*Ernoporus*). Holotype ♂; Kanawah Station [Wood Co., West Virginia]; USNM, Washington.

Distribution: North America (Wood County, West Virginia in USA).

Notes: (1) Wood 1954a: 987, 1982b: 858 (re-described).

References: (**hb**) Chamberlin 1939: 317; Wood, S. L. 1953: 51, 1954a: 987. (**ds**) Blatchley & Leng 1916: 605; Chamberlin 1939: 317; Wood, S. L. 1982b: 858. (**tx**) Blatchley & Leng 1916: 605; Chamberlin 1939: 317; Hopkins 1915b: 35; Schedl 1962r: 90; Wood, S. L. 1953: 51, 1954a: 986–987, 1982b: 858.

longus (Eggers) 1926b: 136 (*Ernoporus*). Lectotype, sex?; Tomakomai, Japan; USNM, Washington, designated by Anderson & Anderson 1971: 18. Distribution: Asia (Hokkaido in Japan/Kamchatka, Siberia in E USSR).

Hosts: *Alnus fruticosa*, *A. kamshatica*, *A. maximowitzi*.

References: (**cn**) Kurenzov 1935c: 188. (**cc**) Kurenzov 1934a: 54. (**hb**) Kurenzov 1935a: 20, 34, 1948b: 115; Kurenzov & Kononov 1961: 600; Stark 1952: 275. (**ds**) Kurenzov 1934a: 54, 1935a: 20, 34, 1935c: 188, 1936b: 351, 1938a: 64, 1951b: 17, 1965, 1967; Kurenzov & Kononov 1961: 595–600; Stark 1952: 275. (**tx**) Anderson, W. H. & Anderson 1971: 18; Eggers 1926b: 136; Krivolutsкая 1955; Kurenzov 1941a: 158, 1948b: 115; Nobuchi 1985c: 11; Schedl 1934f: 1639, 1979c: 142; Sokanovskii 1954: 17; Stark 1952: 275.

semenovi (Kurenzov) 1941a: 231 (*Eocryphalus*). Lectotype, sex?; mountainous fir and broad-leaved woods in the upper course of the river Suputinka (Ussuri Region); ISBN, Novosibirsk, USSR, and IZL, Leningrad/St. Petersburg, designated by Michalski 1969a: 892.

Distribution: Asia (Sakhalin Island, Ussuri in E USSR).

Hosts: *Kalopanax ricinifolium*.

References: (**hb**) Krivolutsкая 1973: 137; Kurenzov 1948b: 125; Stark 1952: 279. (**ds**) Krivolutsкая 1965a: 235, 1973: 137; Stark 1952: 279. (**tx**) Krivolutsкая 1958: 148; Kurenzov 1941a: 161, 231, 1948b: 125; Michalski 1969a: 892, 1969b: 567; Stark 1952: 279.

spessivtzeri Berger 1916: 243. Holotype, sex?; Sedanka, Ussuri, USSR; IZL, Leningrad.

Distribution: Asia (Ussuri in E USSR).

Hosts: *Fraxinus mandschurica*.

References: (**cn**) Kurenzov 1935c: 193, 1951c: 79. (**cc**) Kurenzov 1934a: 51, 1951d: 24. (**hb**) Krivolutsкая 1973: 136; Kurenzov 1948b: 128, 1951d: 24; Stark 1952: 277. (**ds**) Kleine 1934a: 147; Krivolutsкая 1965a: 235, 1973: 136; Kurenzov

1934a: 51, 1935c: 193, 1936a: 110, 1936b: 350; Stark 1952: 277. (tx) Balachowsky 1949a: 201; Berger 1916: 243; Kurenzov 1941a: 160–161, 1948b: 128; Michalski 1969a: 892, 1969b: 567; Schedl 1934f: 1639; Stark 1952: 277.

takahashii Nobuchi 1975: 43. Holotype, sex?, Mt. Daisetsu, Hokkaido, Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. I, fig. 4.
Distribution: Asia (Hokkaido in Japan).
Hosts: *Sorbus commixta*.
References: (tx) Nobuchi 1975: 43–44.

zachvatkini (Krivolutskaya) 1958: 149 (*Eocryphalus*). Holotype, sex?, Sakhalin Island, S. Uglegorsk region, near mouth of Kotan River, USSR; IZM, Moscow.
Distribution: Asia (Sakhalin Island in E USSR).
References: (hb) Krivolutskaya 1973: 137. (ds) Krivolutskaya 1973: 137. (tx) Krivolutskaya 1958: 149; Schedl 1962r: 93–94.

Genus *Allernoporus* Kurenzov

ALLERNOPORUS KURENZOV 1941a: 159. Type-species: *Allernoporus euonymi* Kurenzov, monobasic.
Notes: (3) When more thorough studies are made, it is probable that this genus will become a synonym of *Ernoporiscus*.

References: (hb) Wood, S. L. 1986a: 89. (ds) Wood, S. L. 1986a: 89. (tx) Kurenzov 1941a: 159–160; Schedl 1962r: 93–94; Wood, S. L. 1986a: 89.

euonymi Kurenzov 1941a: 159, 230. Lectotype, sex?, mountainous woods in South of Soviet Far East: ISBN, Novosibirsk, USSR, and IZL, Leningrad/St. Petersburg, Michalski 1969a: 892.
Distribution: Asia (Sakhalin Island, Siberia in E USSR).

Hosts: *Euonymus* sp.

References: (hb) Kurenzov 1948b: 123; Stark 1952: 278. (ds) Krivolutskaya 1983; Stark 1952: 278. (tx) Krivolutskaya 1955: 140; Kurenzov 1941a: 159, 230, 1948b: 123; Michalski 1969a: 892, 1969b: 567; Schedl 1962r: 92–94; Stark 1952: 278.

Genus *Stegomerus* Wood

STEGOMERUS WOOD 1967c: 129. Type-species: *Stegomerus vulgaris* Wood, original designation.

Keys: Wood 1967c: 130, 1982b: 855.

References: (hb) Wood, S. L. 1986a: 90. (ds) Wood, S. L. 1986a: 90. (tx) Wood, S. L. 1967c: 129, 1982b: 854–858, 1986a: 90.

chiriquensis Wood 1967c: 131. Holotype ♂; Cerro Punta, Chiriqui, Panama; Wood Collection.
Distribution: North America (Panama).

Hosts: *Canavalia villosa*.

References: (tx) Wood, S. L. 1967c: 131, 1982b: 856.

mexicanus Wood 1967c: 133. Holotype ♂; 21 km W Morelia, Michoacan, Mexico; Wood Collection.
Figures: Atkinson & Equihua 1988: 101 (adult).

Distribution: North America (Michoacan, Puebla in Mexico).

Hosts: *Serjania* sp.

References: (ds) Atkinson & Equihua 1985c: 356, 1988: 101. (tx) Atkinson & Equihua 1988: 101; Wood, S. L. 1967c: 133, 1982b: 857.

mirandus Wood 1971: 32. Holotype ♀; El Laurel Experimental Farm, 12 km SW Caracas, Miranda, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: Vine (creeper).

References: (tx) Wood, S. L. 1971: 32–33.

montanus Wood 1967c: 132. Holotype ♂; Volcan Irazu, Cartago Prov., Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).

Hosts: *Mucenbeckia tannifolia*.

References: (tx) Wood, S. L. 1967c: 132.

pygmaeus Wood 1967c: 130. Holotype ♂; Los Corchos, Nayarit, Mexico; Wood Collection.

Distribution: North America (Costa Rica/ Honduras/ Jalisco, Nayarit in Mexico).

Hosts: *Canavalia villosa*, *Cestrum scandans*, *Dioclea megacarpa*.

References: (hb) Atkinson & Equihua 1985c: 356. (ds) Atkinson & Equihua 1985c: 356. (tx) Wood, S. L. 1967c: 130, 1982b: 855–856.

vulgaris Wood 1967c: 134. Holotype ♂; Volcan Zunil, Quezaltenango, Guatemala; Wood Collection.
Distribution: North America (Costa Rica/ Guatemala/ Honduras).

Hosts: *Serjania mexicana*, *S. triquetra*, *S. sp.*

References: (tx) Wood, S. L. 1967c: 133–134.

Genus *Neocryphus* Nunberg

NEOCRYPHUS NUNBERG 1956a: 139. Type-species: *Neocryphus argentinensis* Nunberg, original designation.

Notes: (3) Schedl 1979: 35 (erroneously placed in synonymy with *Phacrylus*).

References: (hb) Wood, S. L. 1986a: 90. (ds) Wood, S. L. 1986a: 90. (tx) Nunberg 1956a: 139; Wood, S. L. 1986a: 90.

argentinensis Nunberg 1956a: 141. Holotype, sex?, Argentina, Cordova; IZW, Warsaw.

Distribution: South America (Argentina).

References: (tx) Nunberg 1956a: 141.

cristatus (Schedl) 1979e: 61 (*Phacrylus*). Holotype, sex?, S. Luis, San Geronimo, Argentina; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

References: (tx) Schedl 1979e: 61.

Genus *Acorthylus* Brethes

ACORTHYLUS BRETHES 1922: 304. Type-species: *Acorthylus asperatus* Brethes, monobasic.

Phacrylus Schedl 1935i: 24. Type-species: *Phacrylus bosqui* Schedl, monobasic. Synonymy: Wood 1983a: 647.

References: **(tx)** Nimberg 1956a: 141; Schedl 1935i: 24; Wood, S. L. 1983: 647.

References: **(hb)** Wood, S. L. 1986a: 90. **(ds)** Wood, S. L. 1986a: 90. **(tx)** Brethes 1922: 304; Wood, S. L. 1983a: 647, 1986a: 90.

asperatus Brethes 1922: 305. Holotype, sex?; Buenos Aires, Argentina; Brethes Collection.

Distribution: South America (Argentina).

References: **(ds)** Blackwelder 1947: 784. **(tx)** Brethes 1922: 305.

bosqui (Schedl) 1935i: 24 (*Phacrylus*). Lectotype, sex?; Argentinien, Prov. de Jujuy; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 44.

Distribution: South America (Argentina).

References: **(ds)** Blackwelder 1947: 778; Schedl 1966f: 84, 1967d: 2, 1972d: 141. **(tx)** Schedl 1935i: 24, 1951d: 17, 1952a: 453, 1964m: 304, 1972d: 141, 1979c: 44.

squamulosus Eggers 1943a: 356 (*Ernoporus*). Holotype, sex?; Bolivia (Cochabamba); Eggers Collection, not mentioned by either Anderson & Anderson 1971 or Schedl 1979c, Eggers cotype in NHMW, Wien. Synonymy: Schedl 1964m: 304.

References: **(tx)** Eggers 1943a: 356; Schedl 1964m: 304, 1979c: 236.

gracilis (Schedl) 1972g: 58 (*Phacrylus*). Holotype, sex?; Brasil, Jacareacanga, Para; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: **(ds)** Schedl 1973a: 367. **(tx)** Schedl 1972g: 58, 1979c: 108.

pruni (Wood) 1971: 33 (*Phacrylus*). Holotype ♀; La Carbonera Experimental Forest, 50 km NW Merida, Merida, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Prunus sphaerocarpa*.

References: **(tx)** Wood, S. L. 1971: 33.

robustus (Schedl) 1952a: 453 (*Phacrylus*). Lectotype ♂; Argentina, Cordoba, Dep. Calamuchita, El Sauce, La Cienaga, Belen, Catamarca; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina/ Brazil).

References: **(hb)** Viana 1964: 122. **(ds)** Viana 1964: 122. **(tx)** Schedl 1951h: 284, 1951m: 73, 1952a: 453, 1979c: 212.

Genus *Stephanopodius* Schedl

STEPHANOPODIUS SCHEDL 1941d: 396. Type-species: *Stephanoderes dispar* Eggers, subsequent designation by Schedl 1961k: 633.

Cryphalomimus Browne 1962g: 75. Type-species: *Hypocryphalus ghanaensis* Schedl, original designation, preoccupied by Eggers 1927. Synonymy: Wood 1980c: 95.

References: **(tx)** Browne 1962g: 75, 1963a: 242; Schedl 1963j: 478; Wood, S. L. 1980c: 95.

Cryphalomimetus Browne 1963b: 242. Type-spe-

cies: *Hypocryphalus ghanaensis* Schedl, automatic. Synonymy: Wood 1980c: 95.

References: **(tx)** Browne 1963b: 242; Schedl 1964m: 305; Wood, S. L. 1980c: 95.

References: **(hb)** Wood, S. L. 1986a: 90. **(ds)** Wood, S. L. 1986a: 90. **(tx)** Schedl 1941d: 396, 1961k: 633; Wood, S. L. 1986a: 90.

dispar (Eggers) 1936c: 35 (*Stephanoderes*). Holotype, sex?; S. Rhodesia; Salisbury; BMNH, London.

Figures: Schedl 1961k: 634.

Distribution: Africa (South Africa/ Tanzania/ Zimbabwe).

References: **(hb)** Loytyniemi, Beaver, & Loytyniemi 1984; Schedl 1961k: 633. **(ds)** Beaver & Loytyniemi 1985a: 81; Schedl 1961k: 633. **(tx)** Eggers 1936c: 35; Schedl 1941d: 396, 1951j: 20, 1961i: 224, 1961k: 633–634, 1979c: 82.

ghanaensis (Schedl) 1962h: 66 (*Hypocryphalus*). Holotype, sex?; Ghana, Kumasi; BMNH, London. Distribution: Africa (Ghana).

Hosts: *Bussa occidentalis*, *Hymenostegia afzelii*.

References: **(hb)** Browne 1963a: 235. **(ds)** Browne 1963a: 235. **(tx)** Browne 1962g: 74–75, 1963b: 242; Schedl 1962h: 66, 1962k: 1069, 1979c: 103.

giganteus Schedl 1950d: 26. Holotype, sex?; Rhodesien; Schedl Collection in NHMW, Wien. Distribution: Africa (South Africa/ Zimbabwe). References: **(tx)** Schedl 1950d: 26, 1961k: 634, 1979c: 104.

mkulumusius (Eggers) 1919: 241 (*Stephanoderes*). Syntypes, sex?; Mkulumusiberg bei Sigi; Amani; Hamburg Museum, lost, and Eggers Collection ♂ ♀, in NHMW, Wien.

Distribution: Africa (Tanzania).

Notes: (1) Schedl 1979c: 157 (neotype designation invalid).

References: **(tx)** Eggers 1919: 241, 1922b: 168; Schedl 1950d: 3, 1961k: 635, 1979c: 157.

squamosus Nunberg 1973: 9. Holotype ♀; Congo Belge, P.N.A., expl. sect. Nord Massif Ruwenzori, Kalonge; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: **(tx)** Nunberg 1973: 9.

usambaricus Schedl 1941d: 396. Holotype, sex?; Usambara, Tanganyika; Schedl Collection in NHMW, Wien.

Distribution: Africa (Tanzania).

References: **(tx)** Schedl 1941d: 396, 1961k: 635, 1979c: 262.

Genus *Coriacephilus* Schedl

CORIACEPHILUS SCHEDL 1939e: 339. Type-species: *Stephanoderes coriaceus* Eichhoff, original designation.

References: **(hb)** Wood, S. L. 1986a: 90. **(ds)** Browne 1961c: 67; Schedl 1966b: 27; Wood, S. L.

1986a: 90. (tx) Schedl 1939e: 339–340; Wood, S. L. 1986a: 90.

coriaceus (Eichhoff) 1878b: 494 (*Stephanoderes*). Holotype ♀; Asia (Siam); Schedl Collection in NHMW, Wien.

Distribution: Asia (Burma/Thailand).

References: (ds) Beaver & Browne 1975: 287; Hagedorn 1910d: 41; Kleine 1913b: 121, 1914b: 280. (tx) Eichhoff 1878b: 494; Hagedorn 1910a: 86; Hopkins 1915b: 23; Schedl 1939e: 339–340, 1940d: 591, 1979c: 65.

birmanus Eggers 1925: 153 (*Cryphalus*). Syntypes, sex?; Tenasserim; Prague Museum, and 1 in Eggers Collection, in NHMW, Wien. Synonymy: Schedl 1940d: 591.

References: (tx) Eggers 1925: 153, 156; Schedl 1940d: 591, 1979c: 40.

cribripennis Schedl 1943b: 40. Lectotype, sex?; Kalambangan, Mindanao, Philippines; Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands (Mindanao).

References: (tx) Schedl 1943b: 40, 1979c: 69.

proximus (Eggers) 1925: 156 (*Cryphalus*). Holotype, sex?; Sud-Mysore (Vorder-Indien); Eggers Collection, in NHMW, Wien.

Distribution: Asia (Mysore in India).

Notes: (1) Schedl 1950f: 36 (to *Coriacephilus*).

References: (tx) Eggers 1925: 156; Schedl 1950f: 36–37, 1979c: 199.

xyloctonoides Schedl 1939e: 340. Lectotype, sex?; Malaya, Selangor, Sungai Buloh For. Res.; Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

Hosts: *Kucma furfuracea*.

References: (cn) Mathur & Singh 1960b: 84. (ds) Beeson 1961: 286; Browne 1961c: 67; Mathur & Singh 1960b: 84. (tx) Browne 1961c: 67; Schedl 1939e: 340, 1979c: 270.

Genus *Ernopor* Thomson

ERNOPORUS THOMSON 1859: 147. Type-species: *Bostrichus tiliac* Panzer, original designation.

Cryphalops Reitter 1889a: 94. Type-species: *Cryphalus lederi* Reitter = *Bostrichus tiliac* Panzer, monobasic. Synonymy: Reitter 1894: 71.

References: (tx) Balachowsky 1949a: 201; Hopkins 1915b: 8; Reitter 1889a: 94, 1894a: 71.

Stephanorhopalus Hopkins 1915b: 35. Type-species: *Stephanorhopalus melodori* Hopkins, original designation, spelling of type species amended by Schedl 1966: 19. Synonymy: Wood 1980c: 93.

References: (tx) Browne 1961c: 70; Hopkins 1915b: 35; Schedl 1962r: 93–94, 1966b: 19; Wood, S. L. 1980c: 93.

Euptilius Schedl 1940d: 589. Type-species:

Ernopor *concentralis* Eggers, original designation. Synonymy: Wood 1980c: 93.

References: (tx) Schedl 1940d: 589, 1962r: 90; Wood, S. L. 1980c: 93.

References: (ay) Heymons 1920: 103; Nobuchi 1969a: 56. (hb) Escherich 1923: 460–482; Palm 1959: 34, 44, 57; Schedl 1977a: 41; Wood, S. L. 1986a: 90. (ds) Bertolini 1872: 201; Brimblecombe 1953: 23; Browne 1961c: 70; Feytaud 1950: 9; Schedl 1934f: 1639, 1966b: 19, 1977a: 41; Scheerpeltz & Winkler 1930: 257; Wood, S. L. 1986a: 90. (tx) Allen, A. A. 1970: 246; Balachowsky 1949a: 208; Beeson 1938b: 290; Berger 1916: 227–228; Blandford 1904: 226; Blatchley & Leng 1916: 593, 604; Browne 1961c: 70, 79; Chamberlin 1939: 316–317; Choo 1983: 69; Choo, Woo, & Nobuchi 1988b; Costa Lima 1956: 282; Endrodi 1957b: 416; Fauvel 1887: 276; Ferrari 1867a: 10, 16; Hopkins 1914: 121, 135, 1915b: 8, 35, 194–195, 221, 226; Kurenzov 1941: 159–162, 230–231; Kurenzov & Kononov 1961: 595; Lacordaire 1886: 378; Lekander 1962: 431; Lindemann 1876: 148–167, 320–380; Luigioni 1929: 997; Munro 1946: 19, 32, 42, 61; Murayama 1954: 205; Nurnberg 1928: 145, 1954: 10, 48, 52–53, 1956: 141; Nusslin 1898: 276, 1911, 1912; Peyerimhoff 1935: 194–195; Pfeffer 1989a: 82; Reitter 1913a: 65, 68–69, 1916: 287, 289; Schedl 1958d: 188, 1958k: 142, 1959a: 475, 1962r: 93–94, 1969i: 130, 1977a: 41, 1977b: 41, 1980b: 75; Schimitschek 1937: 50–52; Sokanovskii 1954: 17; Spessivtsov 1923: 422, 1931: 88; Stresemann et al. 1989: 354; Swaine 1909: 92; Thomson 1859: 147, 1865: 360; Trappen 1935: 142; Wood, S. L. 1954a: 988, 1960a: 13, 18, 1986a: 90.

***acanthopanaxi* (Niisima)** 1913a: 4 (*Cryphalus*). Syntypes, sex?; Sapporo, Japan; Nobuchi Collection, Ibaraki.

Distribution: Asia (Hokkaido in Japan).

Hosts: *Acanthopanax ricinifolium*.

References: (ds) Kleine 1934a: 147; Nobuchi 1985c: 11. (tx) Murayama 1953a: 35; Niisima 1913a: 4; Schedl 1934f: 1639, 1979c: 9.

***antennarius* (Schedl)** 1974d: 461 (*Ernopor*). Holotype ♀; Upper Manki logging area, Bulolo, Morobe District; CSIRO, Canberra.

Distribution: New Guinea.

References: (tx) Schedl 1974d: 461, 1979c: 22.

papuans Browne 1984f: 70 (*Euptilius*). Holotype, sex?; New Guinea: Morobe District, Mount Kaindi, 2500 m; BMNH, London, preoccupied by Schedl 1973. Synonymy: Wood 1989: 172.

References: (tx) Browne 1984f: 70; Wood, S. L. 1989: 172

***armatus* (Browne)** 1981a: 129 (*Euptilius*). Holotype, sex?; Bangkok (Thailand) to Yokohama (Japan), imported; BMNH, London.

Distribution: Asia (Thailand).

- Hosts: *Pterocarpus* sp.
References: (tx) Browne 1981a: 129.
- concentralis** Eggers 1936d: 629. Holotype, sex?; Vorderindien, Mysore: Sakalaspur; BMNH, London.
Distribution: Asia (Mysore in India).
Hosts: *Dalbergia latifolia*, *Pterocarpus marsupium*.
References: (ds) Beeson 1961: 287. (tx) Eggers 1936d: 629; Schedl 1940d: 589–590.
- dispar** (Schedl) 1972i: 49 (*Cryphalops*). Holotype ♂; New Ireland Dist., Karbil; CSIRO, Canberra.
Distribution: New Ireland Island.
Hosts: *Intsia bijuga*.
References: (tx) Schedl 1972i: 49, 1979c: 82.
- exiguus** (Browne) 1984c: 451 (*Euptilius*). Holotype, sex?; Tolitoli (Celebes) to Omaezaki (Japan), imported; BMNH, London.
Distribution: Indonesia (Celebes).
References: (ds) Browne 1984c: 451, 1986a: 89.
- inermis** (Schedl) 1939e: 343 (*Stephanorhopalus*). Lectotype, sex?; Malaya, Selangor: Port Swettenham; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 124.
Distribution: Asia (Malaya).
Hosts: *Rhizophora mucronata*.
References: (ds) Beeson 1961: 299; Browne 1961c: 70. (tx) Schedl 1939e: 343, 1979c: 124.
- japonicus** Nobuchi 1966e: 52. Holotype ♀; Arashiyama, Kyoto City; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1966e: 52.
Distribution: Asia (Honshu in Japan).
References: (ds) Nobuchi 1955c: 11. (tx) McNamara 1977: 196; Nobuchi 1966e: 52.
- melodori** (Hopkins) 1915b: 36 (*Stephanorhopalus*). Holotype ♂; Calapan, P. I.; USNM, Washington.
Distribution: Philippine Islands (Mindoro).
Hosts: *Melodorum fulgens*.
Notes: (1) Schedl 1966b: 16 (*nulodori*, lapsus calami, corrected to *melodori*).
References: (ds) Kleine 1934a: 148; Schedl 1966b: 19. (tx) Hopkins 1915b: 35–38; Nobuchi 1983: 300; Schedl 1966b: 16, 19.
- papuanus** (Schedl) 1973e: 87 (*Margadillius*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
References: (tx) Schedl 1973e: 87, 1979c: 183–184.
- quadridens** (Schedl) 1971a: 284 (*Cryphalops*). Holotype, sex?; Central Prov.: Hantane Estate, Kandy; Schedl Collection in NHMW, Wien.
Figures: Schedl 1971a: 283.
Distribution: Asia (Sri Lanka).
References: (tx) Schedl 1971a: 283–284, 1979c: 208.
- shimanensis** Murayama 1953a: 36. Holotype ♂; Kakinoki vill. (Suzuno-ohitani), Kanoashi County, Shimane pref., Japan; Murayama Collection in USNM, Washington.
- Distribution: Asia (Honshu in Japan).
Hosts: *Fagus crenata*.
References: (ds) Murayama 1954b: 185; Nobuchi 1985c: 11. (tx) Murayama 1953a: 36, 1954b: 185.
- thailandicus** (Schedl) 1967a: 127 (*Euptilius*). Holotype, sex?; Thailand via Hong Kong to Yokohama (Japan), imported; PPST, Tokyo.
Distribution: Asia (Thailand).
Hosts: Kwararin wood.
References: (ds) Beaver & Browne 1975: 290. (tx) Schedl 1967a: 127, 1979c: 252.
- tiliae** (Panzer) 1793: 14 (*Apate*). Syntypes, sex?; Brungsvigiae; not located, presumed lost.
Figures: Balachowsky 1963a: 1280, Grune 1979: 110, Pfeffer 1989a: pl. 6, Postner 1974: 432 (adult).
Distribution: Asia (Korea/ Turkey/ E USSR), Europe (Austria/ Bulgaria/ Corsica/ Czechoslovakia/ Denmark/ England/ Finland/ France/ Germany/ Greece/ Hungary/ N Italy/ Norway/ Poland/ Romania/ Spain/ Sweden/ Switzerland/ W USSR).
Hosts: *Carpinus betulus*, *Fagus sylvatica*, *Hibiscus syriacus*, *Tilia amurensis*, *T. cordata*, *T. intermedia*, *T. parvifolia*.
References: (ay) Escherich 1923b: 482; Imhoff 1856: 228; Lekander 1959b: 60; Numberg 1928a: 140; Nusslin 1911a: 58, 155, 277, 337, 378. (bv) Grune 1979: 111; Hellen 1921. (cn) Acatay 1943a: 65; Barbey 1925: 587; Chorbadzhiev 1929; Escherich 1923b: 482; Esterberg 1959; Georgescu et al. 1957: 357, 406; Grandi 1951; Hellen 1921; Kamp 1956b: 471; Kholodkovskii 1912: 307; Koppen 1882: 251; Kovacevic 1924: 21–22; Marcu 1926c: 63; Nosek 1951: 106, 1952b: 98; Nusslin 1913: 290; Picard & Lichtenstein 1917: 17; Pierce, W. D. 1917: 141; Rhumbler 1922: 334, 1927: 350; Schimitschek 1937c: 52, 1955a: 146, 1955c: 83; Schuster 1918: 97; Schwerdtfeger 1944a: 175, 1957a: 182; Wachtl 1901: 381; Wichmann 1927b: 361. (cc) Balazy & Michalski 1964b; Elliot & Morley 1907; Gillanders 1906; Kielczewski 1976; Kielczewski, Moser, & Wisniewski 1983; Kielczewski & Wisniewski 1978, 1980a, 1983; Kleine 1908c: 186, 1909a: 46, 1944: 73; Kostenko 1929; Kurenzov 1934a: 57; Michalski & Ratajczak 1989; Nosek 1951: 106, 1952b: 98, 1959a: 118, 1959b: 87; Numberg & Wiackowski 1958: 130; Nusslin 1927: 350; Palmén 1944: 60; Perris 1856a: 244; Pfeffer 1923a: 331; Rondani 1873: 146; Saalas 1930: 118; Schimitschek 1955a: 146; Schuster 1918: 97; Schwerdtfeger 1944a: 175, 1957a: 182; Sedlaczek 1935a: 162; Wisniewski 1979b. (hb) Acatay 1943a: 65; Altum 1881c: 322; Balachowsky 1963a: 1279; Barbey 1901: 23, 72, 1925: 587; Bèffa 1961; Bukowsky 1930; Chorbadzhiev 1929; Dombrowsky 1887; Eichhoff 1881a: 45, 181, 1882a: 245, 1882c: 322, 1892b: 99; Escherich 1923b: 482; Fuchs 1904a, 1907: 48; Gillanders 1906, 1908; Gornostaev 1916: 314; Grandi 1951;

- Györfi 1957; Hagedorn 1903a; Henschel 1895a: 168; Karpinski & Stravinski 1948: 155; Karsch 1883: 142; Keller 1913: 242; Kemmer 1919: 176; Kharltonov 1924: 199–204; Kholodkovskii 1912: 307; Kurenzov 1935a: 34; Lekander 1959b: 60; Lengerken 1954: 83; Marcu 1941: 403; Marie 1955: 78; Munro 1926: 61; Nordlinger 1856: 30; Nosek 1959a: 118, 1959b: 87; Numberg 1929c: 119; Nusslin 1898: 280, 1906b: 14, 1913: 290, 1927: 350; Perris 1856a: 244; Postner 1974: 432; Ratzeburg 1837: 164, 1839: 199; Rhumbler 1922: 334, 1927: 350; Rupertsberger 1879: 231, 1880: 229; Schedl 1981b: 76; Schimitschek 1955a: 146; Schwerdtfeger 1944a: 175, 1957a: 182, 1981: 188; Sedlacek 1935a: 102; Spessivtsev 1913a: 90; Stark 1926a: 335, 1952: 272; Tschorbadjiev 1929: 166; Vinogradov-Nikitin 1911: 1472–1477; Wachtl 1876a: 453, 1901: 381; Wichmann 1927b: 361; Zvierezhomb-Zubovsky & Rostoff 1918: 1–36. (ds)
- Acatay 1943: 65; Aclouque 1896; Allen, A. A. 1970; Ammann & Knabl 1913; Andersch 1851; Arnoldi et al. 1955: 693; Andras & Schaeffer 1957; Bakke 1963a: 121; Balachowsky 1963a: 1279; Barthe 1896; Bau 1888; Bedel 1888b: 398, 413–414; Bejer-Petersen & Jørm 1977: 23; Bielz 1851, 1887; Blanchere & Robert 1889; Borchert 1951; Brakman 1966b: 206; Brukowsky 1930; Buresh & Lazarov 1956; Calver 1884, 1893; Carpentier & Delaby 1908; Cecconi 1897; Cho 1957; Choo 1983: 69; Choo & Woo 1985: 164; Chorbadzhiyev 1929; Dejean 1837; Duftschmidt 1825; Eder 1934; Eggers 1904; Endrodi 1958a, 1958b; Ericson & Sandin 1893; Escherich 1923b: 482, 1932b; Esterberg 1928, 1959; Fjellberg 1966: 154; Fowler 1891; Fuchs 1904a, 1905a, 1907: 48; Gaubil 1849: 126; Gemminger & Harold 1872: 2683; Gornostaev 1917; Gozis 1875: 80; Grill 1895: 309; Grune 1979: 111; Gyllenhal 1827: 624; Hagedorn 1903a, 1910d: 46; Hansen, V. 1939, 1956, 1964: 462; Heinemann 1908a; Hellen 1947; Henschel 1895a: 168; Heyden 1876: 299, 1879: 140; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1951; Jazentkovsky 1912: 288; Kamp 1956b: 471; Karpinski 1926: 82, 1931: 27, 1948b: 230; Karpinski & Stravinski 1948: 155; Keler 1925b: 272; Kemmer 1919: 176; Kersten 1933: 74; Kestercanek 1881a: 12; Klefbeck & Sjöberg 1960: 2, 230; Kleine 1912a: 262, 267, 1913a: 34, 1934a: 143; Kloft & Hinks 1945: 218; Ko 1969: 276; Kolbe, W. 1918: 211; Koltze 1901: 153; Koppén 1882: 251; Kostenko 1929; Kozikowsky 1921: 180; Kraatz 1869: 59; Kurenzov 1934a: 57, 1935a: 34, 1936a: 110, 1936b: 350–351, 1938a: 59; Kurir 1947c: 27; Lacordaire 1866: 379; Langhoffer 1915c: 158; Leclercq 1971; Lengerken 1939: 39; Lentz 1857: 139; Liegel 1886: 43; Lindemann 1884b: 264; Lokaj 1868: 64; Lomnicki 1886a: 242, 1913b: 148; Lucht 1987: 278; Lundberg 1981: 151; Lundblad 1950b: 72; Mahler 1987: 232; Marcu 1926c: 63; Matthews & Fowler 1883: 42; Murayama 1929b: 2, 1930b: 10, 1937b: 375; Negru 1966b: 402, 1968a: 456; Negru & Rosca 1967: 141; Numberg 1928b: 85, 106, 1954: 53; Nusslin 1898: 280; Orest 1926c: 63; Palm 1959: 29, 60, 1986; Palmén 1944: 60; Perris 1876a: 254, 1877a: 414; Pfeffer 1924b: 471, 1931b: 74, 1950b: 76, 1989a: 82; Pierce, W. D. 1917: 141; Pittioni 1943: 175; Power 1865: 212; Postner 1974: 432; Prossen 1913: 83; Rapp 1934: 728; Ratzeburg 1837: 164, 1839: 199; Redtenbacher 1858: 832, 1874: 375; Reitter 1869b: 154, 1889a: 94, 1894a: 71, 1916: 289; Roubal 1941: 264; Ruskov 1928e: 61; Saalas 1930: 118, 1931: 67; Sainte-Claire 1914: 472; Sainte-Claire & Mequignon 1938: 445; Schaufuss 1915: 1232; Schaum 1859: 96, 1862: 101; Schedl 1980a: 15, 1981b: 76; Schilsky 1909: 188; Schiodte 1873: 101; Schwerdtfeger 1981: 188; Seidlitz 1891b: 607; Sharp & Fowler 1893: 34; Siebke 1875: 283; Stark 1926a: 335, 1926j: 126, 1927b: 88, 1936c: 142, 1952: 272; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 439; Stierlin & Gautard 1871: 292; Sturm 1826: 102, 1843: 230; Thomson 1859: 147, 1865: 350, 1868: 218; Tredl 1907: 12; Tschorbadjiev 1929: 166; Villa & Villa 1833: 26; Wachtl 1876a: 453; Westhoff 1882: 238; Wichmann 1927a: 6; Winter, T. G. 1983: 11; Zinovjev 1955: 157. (tx) Aclouque 1896; Allen, A. A. 1970; Bach 1854; Balachowsky 1949a: 209, 1963a: 1279–1280; Barbey 1901: 23, 72; Bechstein et al. 1805: 108; Bedel 1888b: 398, 413–414; Beffa 1961; Berger 1917: 226–248; Bertolini 1872; Calver 1888; Carpentier & Delaby 1908; Castelnau 1840; Choo 1983: 69; Doebner 1860; Dombrowsky 1887; Duftschmidt 1825; Eggers 1929c: 41, 1936d: 629, 1942c: 29, 1943a: 355; Eichhoff 1864b: 35, 1878b: 130, 1881a: 45, 181, 1883a: 109, 133; Endrodi 1957b; Erichson 1836: 62; Escherich 1923b: 482; Fabricius 1798: 157, 1801: 383; Fauvel 1889; Ferrari 1867a: 12, 1867b: 106, 114; Fleischer 1905, 1927; Formanek 1907: 35; Gebien 1907: 197; Gillanders 1908; Grune 1979: 110–111; Gyllenhal 1813: 369, 1827: 624; Hagedorn 1910a: 88; Hansen, V. 1956, 1964: 462; Henschel 1895a: 168; Hopkins 1914: 121, 1915b: 35, 1915c: 221; Kalina 1970: 129; Karpinski & Stravinski 1948: 155; Kulmt 1913: 1055; Lacordaire 1866: 379; Lekander 1959b: 60; Letzner 1891: 375; Lindemann 1875c: 321; Lovendal 1889b: 52, 1898: 126; Lucht 1987: 278; Murayama 1930b: 10, 14, 30, 1937b: 375; Negru 1966b: 402; Nordlinger 1848: 245, 1856: 30; Numberg 1928a: 140, 1929c: 119, 1954: 53; Nusslin 1911a: 58, 155, 277, 337, 378; Panzer 1793: 14, 1795a: 284; Perris 1877a: 414; Pfeffer 1932b: 18, 1955a: 180, 1989a:pl. 6; Portevin 1935: 324; Postner 1974: 432; Quaschik 1953: 35; Ratzeburg 1837: 164, 1839: 199; Redtenbacher 1849a: 359, 1849b: 26, 1858: 832, 1874: 375; Reitter 1889a: 94, 1894a: 71, 1913a: 68, 1916: 289; Rhumbler 1922:

334, 1927: 350; Rupertsberger 1879: 231, 1880: 229; Sahlberg 1836: 154; Sampson 1919: 114; Schedl 1934f: 1639, 1952f: 87, 1958k: 142, 1967a: 127, 1980a: 15, 1981b: 76; Schimitschek 1937c: 52, 1955c: 83; Seidlitz 1891b: 607; Semenov 1902: 271; Sokanovskii 1954: 17; Spessivtsev 1913a: 90, 1922a: 472, 1925a: 173, 1931: 48–50; Stark 1952: 272; Stierlin 1898: 439; Thomson 1859: 147, 1865: 360, 1865: 219. (**ms**) Escherich 1932b; Heinemann 1908a; Hellen 1921.

cinereus Herbst 1793: 116 (*Bostrichus*). Syntypes, sex?; Deutschland; not located. Synonymy: Balachowsky 1949a: 209.

References: (**tx**) Balachowsky 1949a: 209; Herbst 1793: 116; Iablokoff-Khnzorian 1961: 105.

ratzeburgi Ferrari 1867a: 11 (*Cryphalus*). Syntypes, sex?; Europe; not located. Synonymy: Balachowsky 1949a: 209.

References: (**ds**) Blanchere & Robert 1889; Heyden 1876: 299; Kestercanek 1881a: 12; Kraatz 1869: 59; Redtenbacher 1874: 375; Schilsky 1909: 155; Seidlitz 1872: 393; Stein 1868: 114; Stein & Weise 1877: 164. (**tx**) Balachowsky 1949a: 209; Bertolini 1872; Ferrari 1867a: 11, 1867b: 114; Letzner 1891: 375; Redtenbacher 1874: 375; Sampson 1923a: 270; Seidlitz 1872: 393.

lederi Reitter 1889a: 93 (*Cryphalus*). Syntypes 2, sex?; Circassia; NHMB, Budapest. Synonymy: Balachowsky 1949a: 209.

References: (**ds**) Heyden, Reitter, & Weise 1891: 671; Reitter 1889a: 93. (**tx**) Balachowsky 1949a: 209; Hopkins 1914: 119; Reitter 1889a: 93; Schedl 1934f: 1639.

eggerti Stark 1936: 142. Syntypes, sex?; Soviet Far East; IZL, Leningrad, 1 Eggertsyntype, in NHMW, Wien. Synonymy: Sokanovskii 1954: 17. Notes: (1) Original publication not seen, cited in Stark 1941: 155, 159.

References: (**hb**) Kurenzov 1948b: 124; Muranov 1941: 155, 159; Stark 1952: 273. (**ds**) Arnoldi et al. 1955: 692; Balachowsky 1963a: 1280; Kri-volutskaya 1983; Sokanovskii 1954: 17; Stark 1952: 273. (**tx**) Kurenzov 1941a: 155, 1948b: 124; Pfeffer 1944a: 131; Schedl 1952k: 158; Sokanovskii 1954: 17; Stark 1936: 142, 1952: 273. (**ms**) Pfeffer 1944a: 131.

starki Eggers 1942c: 31. Holotype, sex?; Ussuri; Eggers Collection, in NHMW, Wien. Synonymy: Schedl 1952k: 158.

References: (**tx**) Eggers 1942c: 31; Pfeffer 1944a: 131; Schedl 1952k: 158, 1979c: 236. (**ms**) Pfeffer 1944a: 131.

tuberculatus (Browne) 1981a: 128 (*Euptilius*). Holotype, sex?; Rangoon (Burma) to Kobe (Japan), imported; BMNH, London.

Distribution: Asia (Burma).

Hosts: *Pterocarpus* sp.

References: (**tx**) Browne 1981a: 128.

Genus *Ernocladius* Wood

ERNOCLADIUS WOOD 1980c: 93. Type-species: *Cryphalus corpulentus* Sampson, original designation.

References: (**hb**) Wood, S. L. 1986a: 90. (**ds**) Wood, S. L. 1986a: 90. (**tx**) Wood, S. L. 1980c: 93, 1986a: 90.

corpulentus (Sampson) 1919: 113 (*Cryphalus*). Holotype ♂; India, Nilgiri Hills; BMNH, London. Distribution: Asia (Bengal, Tamil Nadu in India/Malaya), Indonesia (Java).

Hosts: *Hibiscus macrophyllus*, *Kydia calycina*, *Thespesia populnea*, rare in *Ficus religiosa*.

References: (**en**) Mathur & Singh 1960b: 86, 1961b: 51. (**hb**) Browne 1961c: 71; Kalshoven 1955b: 166–168. (**ds**) Browne 1961c: 71; Kleine 1934a: 147; Mathur & Singh 1961b: 51; Sampson 1919: 113. (**tx**) Eggers 1925: 153; Nobuchi 1959c: 22, 1966c: 53; Sampson 1919: 113–114; Schedl 1940d: 590–591, 1942a: 169, 1942d: 2; Wood, S. L. 1980c: 93.

corpulentus sundri Schedl 1969c: 48 (*Margadillius*). Holotype, sex?; India: Sundarbans (Bengal); FRI, Dehra Dun, not present. Synonymy: Wood 1989: 172.

Notes: (1) The holotype never reached the Forest Research Institute and is probably represented by Schedl's "paratype" that was examined by us. (3) This name has also been treated in *Ernoporus* and *Margadillius*.

References: (**tx**) Schedl 1969c: 48, 1979c: 65–66; Wood, S. L. 1989: 172.

guiboutiae (Schedl) 1957d: 53 (*Miocryphalus*). Holotype, sex?; Congo Belge: Mayumbe, Jombo de Bombo, 86 km Nord-Ouest de Tshela; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Guibourtia arnoldiana*.

References: (**tx**) Schedl 1957d: 53, 1961k: 538, 1979c: 114.

Genus *Cryphalogenes* Wood

CRYPHALOGENES WOOD 1980c: 91. Type-species: *Cryphalogenes euphorbiae* Wood, original designation.

References: (**hb**) Wood, S. L. 1986a: 90. (**ds**) Wood, S. L. 1986a: 90. (**tx**) Wood, S. L. 1980c: 91, 1986a: 90.

euphorbiae Wood 1980c: 91. Holotype ♂; 30 km SE Puttalam, Sri Lanka; USNM, Washington.

Distribution: Asia (Sri Lanka).

Hosts: *Euphorbia antiquorum*.

References: (**tx**) Wood, S. L. 1980c: 91.

exiguus Wood 1980c: 92. Holotype ♂; 30 km SE Puttalam, Sri Lanka; USNM, Washington.

Distribution: Asia (Sri Lanka).

Hosts: *Euphorbia antiquorum*.

References: (**tx**) Wood, S. L. 1980c: 92.

Genus *Scolytogenes* Eichhoff

Scolytogenes EICHHOFF 1878b: 475, 497. Type-species: *Scolytogenes darwini* Eichhoff, monobasic.

Lepicerus Eichhoff 1878b: 476, 501. Type-species: *Lepicerus aspericollis* Eichhoff, monobasic, preoccupied. Synonymy: Wood 1980e: 95. References: (tx) Eggers 1929e: 53; Eichhoff 1878b: 476, 501; Hinton 1936: 473; Hopkins 1915b: 8; Schedl 1940d: 555–557, 1952d: 344; Wood, S. L. 1954a: 996, 1980e: 95.

Cryphalomorphus Schauffuss 1891: 12. Type-species: *Cryphalomorphus communis* Schauffuss, monobasic. Synonymy: Schedl 1940d: 587.

References: (ay) Gardner 1934: 10–11, (hb) Browne 1961c: 67, (ds) Blackwelder 1947: 778; Brimblecombe 1953: 21, 39; Hagedorn 1910d: 46, (tx) Chamberlin 1939: 279, 315–316; Eichhoff 1878b: 476, 501; Hagedorn 1910a: 80, 83, 1913b: 253; Hopkins 1914: 119, 133, 1915b: 7, 1915c: 226; Kurenzov 1941: 71; Schauffuss 1891: 12; Schedl 1938f: 41, 1939e: 337–338, 1940d: 555–557, 1950d: 2, 1952d: 344, 1954a: 138, 1957b: 152, 1957d: 59, 1959a: 476, 1961k: 523, 1962p: 203, 1963b: 265, 1963i: 62, 1965e: 349–379, 1965f: 3–15, 1970b: 358, 1971e: 11, 1977b: 44; Wood, S. L. 1954a: 980, 996–997, 1960a: 13.

Letznerella Reitter 1913a: 68. Type-species: *Bostrichus jalapae* Letzner, monobasic. Synonymy: Schedl 1940d: 587.

References: (tx) Balachowsky 1949a: 20; Chamberlin 1939: 316; Reitter 1913a: 68; Schedl 1938f: 41, 583–587, 1952d: 344; Swaine 1918a: 90; Wood, S. L. 1954a: 996, 1960a: 37.

Hypothenoides Hopkins 1915b: 7, 11. Type-species: *Hypothenoides parvus* Hopkins, original designation. Synonymy: Schedl 1954a: 135.

References: (tx) Hopkins 1915b: 7, 11; Schedl 1954a: 138; Wood, S. L. 1960a: 27.

Ernoporides Hopkins 1915b: 8, 34. Type-species: *Ernoporides floridensis* Hopkins, original designation. Synonymy: Schedl 1940d: 587.

References: Chamberlin 1939: 315; Hopkins 1915b: 8, 34; Schedl 1938f: 41, 1940d: 584–587, 1952d: 344; Wood, S. L. 1954a: 990, 1960a: 27.

Neocryphalus Eggers 1922b: 169. Type-species: *Neocryphalus usagarius* Eggers, monobasic. Synonymy: Schedl 1940d: 587.

References: (tx) Eggers 1922b: 169; Schedl 1940d: 587, 1950d: 2, 1952d: 344, 1959a: 476; Wood, S. L. 1954a: 996, 1986: 90.

Negritus Eggers 1923a: 141. Type-species: *Negritus ater* Eggers, subsequent designation by Wood 1982b: 861. Synonymy: Schedl 1957b: 152.

References: (tx) Eggers 1923a: 141–142; Schedl 1939e: 339, 1957b: 152, 1959a: 476; Wood, S. L. 1982b: 861.

Cylindrotomicus Eggers 1936d: 633. Type-species: *Cylindrotomicus squamulosus* Eggers, monobasic. Synonymy: Schedl 1957b: 152.

References: (tx) Eggers 1936d: 633; Schedl 1957b: 152, 1959a: 476.

Lepicerinus Hinton 1936: 473. Type-species: *Lepicerus aspericollis* Eichhoff, automatic. Synonymy: Wood 1982b: 861.

References: (tx) Hinton 1936: 473; Schedl 1950d: 2, 1952d: 344; Wood, S. L. 1954a: 996, 1982b: 861.

Xylocryptus Schedl 1975f: 352. Type-species: *Xylocryptus papuanus* Schedl, original designation. Synonymy: Wood 1986: 90.

Cryphalophilus Schedl 1970b: 358. Type-species: *Cryphalophilus afer* Schedl, monobasic. Synonymy: Wood 1984b: 228, 1992b: 81.

References: (tx) Schedl 1970b: 358; Wood, S. L. 1984b: 228, 1986a: 61, 1992b: 81.

References: (tx) Schedl 1975f: 352; Wood, S. L. 1986a: 90.

Keys: Wood 1960a: 27, 1982b: 862.

References: (ay) Nobuchi 1969a: 57, (hb) Beaver 1987a: 19; Schedl 1961k: 523, 1977a: 44; Wood, S. L. 1982b: 861, 1986a: 90, (ds) Beaver 1987a: 19; Browne 1961c: 67; Schedl 1961k: 523, 1966b: 22, 1977a: 44; Wood, S. L. 1982b: 861, 1986a: 90, (tx) Arnett 1960: 1043, 1968: 1043; Choo, Woo, & Nobuchi 1988b; Eggers 1929e: 53; Eichhoff 1878b: 475, 497; Schedl 1939e: 337–338, 1961k: 523, 1962p: 203, 1977a: 44; Wood, S. L. 1982b: 861–866, 1986a: 90.

absonus (Schedl) 1975f: 344 (*Cryphalomorphus*). Holotype, sex?, Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 344, 1979c: 9.

afer (Schedl) 1970b: 359 (*Cryphalophilus*). Holotype, sex?, Mozambique, Lourenco-Marques to Yokahama (Japan), imported; PPST, Tokyo. Distribution: Africa (Mozambique).

References: (tx) Schedl 1970b: 359, 1979c: 14.

alternans (Schedl) 1975f: 345 (*Cryphalomorphus*). Holotype, sex?, Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 345, 1979c: 16.

amanicus (Eggers) 1919: 239 (*Cryphalus*). Holotype, sex?, Amami (Ostafrika); Hamburg Museum, lost.

Distribution: Africa (Kenya/Tanzania).

References: (tx) Eggers 1919: 239–240, 1931c: 185, 1932c: 23; Schedl 1940d: 587, 1961k: 525.

ankius (Schedl) 1979g: 96 (*Cryphalomorphus*). Holotype, sex?, Papua [New Guinea], Bulolo, Morobe Dist., Upper Manki L. A.; Schedl Collection in NHMW, Wien.

- Distribution: New Guinea.
References: (tx) Schedl 1979g: 96.
- apicalis** (Schedl) 1971e: 11 (*Cryphalomorphus*).
Holotype, sex?; Umgebung Brazzaville; Schedl Collection in NHMW, Wien.
Distribution: Africa (Congo).
References: (tx) Schedl 1971e: 11, 1979c: 23.
- approximatus** (Schedl) 1975f: 346 (*Cryphalomorphus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 346, 1979c: 23.
- aspericollis** (Eichhoff) 1878b: 501 (*Lepicerus*).
Holotype, sex?; Hindostan Asiae (Birma); Hamburg Museum, lost.
Distribution: Africa (Ethiopia/ South Africa), Asia (Burma/ India/ Malaya), Europe (intercepted in Italy), Indonesia (Sumba).
Hosts: *Ipomoea turpethum*.
References: (cc) Schedl 1961k: 525–526. (hb) Emden 1925a: 215–216, 1925b: 89; Zacher 1927: 161. (ds) Browne 1950d: 493; Hagedorn 1910d: 69; Kleine 1913b: 121, 139, 1914b: 277–278, 1934a: 148–149; Schedl 1934f: 1643, 1961c: 71, 1961k: 525, 1965f: 4, 1971c: 363, 1975e: 447. (tx) Eggers 1925: 153, 159, 1927b: 397, 1929e: 53; Eichhoff 1878a: 388, 1878b: 501–502; Hagedorn 1904d, 1910a: 91; Hinton 1936: 473; Hopkins 1914: 124; Lucas 1920: 357; Schedl 1934f: 1643, 1940d: 584–587, 1951i: 55–56, 1961c: 71, 1961k: 525, 1965f: 4; Wood, S. L. 1954a: 996.
- stierlini** Eggers 1911a: 121 (*Cryphalus*). Lectotype, sex?; Italia: Livorno (aus dem indischen Gebiet eingeschleppt); USNM, Washington, designated by Anderson & Anderson 1971: 31. Synonymy: Eggers 1929e: 53.
References: (cn) Emden 1925b. (ds) Emden 1925b; Kleine 1913b: 118; Schaufuss 1915: 1232. (tx) Anderson, W. H. & Anderson 1971: 31; Eggers 1911a: 121, 1925: 153, 1927b: 397, 1929e: 53; Reitter 1913a: 68; Schedl 1934f: 1643, 1940d: 585, 587, 1961k: 525.
- ater** (Eggers) 1923a: 142 (*Negritus*). Lectotype, sex?; Pt. Princessa (Insel Palawan); USNM, Washington, designated by Anderson & Anderson 1971: 5.
Distribution: Indonesia (E Borneo), Philippine Islands (Palawan), Australia.
References: (ds) Schedl 1966b: 23. (tx) Anderson, W. H. & Anderson 1971: 5; Eggers 1923a: 142, 1927b: 406; Nobuchi 1953: 300; Schedl 1939e: 337–338, 1940b: 434, 1950g: 894, 1957b: 152, 1965g: 23, 1979c: 29.
- australis** (Schedl) 1942c: 175 (*Lepicerinus*). Holotype, sex?; Australien; Schedl Collection in NHMW, Wien.
Distribution: Australia.
- References: (tx) Schedl 1942c: 175, 1950f: 43, 1952d: 344, 1979c: 31.
- badius** (Nobuchi) 1975: 44 (*Cryphalomorphus*).
Holotype, sex?; Inugakitage, Fukuoka pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1, fig. 5.
Distribution: Asia (Japan).
Hosts: *Wisteria floribunda*.
References: (ds) Nobuchi 1985c: 12. (tx) Nobuchi 1975: pl. 1, 44.
- bangensis** (Eggers) 1927c: 75 (*Cryphalomorphus*).
Lectotype, sex?; Philippinen: Mindanao, Provinz Zamboanga, Port Banga; USNM, Washington, designated by Anderson & Anderson 1971: 5.
Distribution: Philippine Islands (Mindanao).
Notes: (3) Schedl 1951i: 56 (described male).
References: (ds) Schedl 1966b: 23. (tx) Anderson, W. H. & Anderson 1971: 5; Eggers 1927c: 75–76, 1929e: 53; Nobuchi 1953: 300; Schedl 1940d: 587, 1951i: 50, 55–56, 1979c: 33.
- basilaris** (Wood) 1960a: 30 (*Cryphalomorphus*).
Holotype ♀; Caroline Islands (Koror, Palau); USNM, Washington.
Distribution: Micronesia (Koror Island in Palau Islands in Caroline Islands).
References: (tx) Wood, S. L. 1960a: 30.
- braderi** (Browne) 1965a: 191 (*Cryphalomorphus*).
Holotype, sex?; Ivory Coast: Adiopodoume; RNH, Leiden.
Distribution: Africa (Ghana/ Ivory Coast).
Hosts: *Triplochiton scleroxylon*.
References: (tx) Browne 1965a: 191; Wood, S. L. 1992b: 84.
- orientalis** Schedl 1971e: 11 (*Cryphalomorphus*).
Holotype, sex?; Ghana, Bekwai; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992b: 84.
References: (ds) Schedl 1972e: 281. (tx) Schedl 1971e: 11–12, 1979c: 180; Wood, S. L. 1992b: 84.
- brimblecombei** (Schedl) 1972a: 146 (*Cryphalomorphus*). Holotype, sex?; Queensland: Emm Vale; Queensland Museum, Brisbane.
Distribution: Australia (Queensland).
References: (tx) Schedl 1972a: 146, 1979c: 47.
- buruensis** (Eggers) 1926a: 300 (*Cryphalomorphus*). Holotype, sex?; Buru-Insel; Eggers Collection, in NHMW, Wien.
Distribution: Indonesia (Buru Island).
References: (tx) Eggers 1926a: 300, 1927c: 75, 1929e: 53; Schedl 1940d: 587, 1952d: 344, 1979c: 49.
- camelliae** (Nobuchi) 1975: 45 (*Cryphalomorphus*).
Holotype, sex?; Tainaijōya, Nigata pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1, fig. 6.
Distribution: Asia (Japan).
Hosts: *Camellia japonica rusticana*.

- References: (ds) Nobuchi 1955c: 12. (tx) Nobuchi 1975: 45–46.
- candidus** (Nobuchi) 1975: 47 (*Cryphalomorphus*). Holotype, sex?; Gyokuto, Kumamoto pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1, fig. 8.
Distribution: Asia (Kyushu in Japan).
References: (ds) Nobuchi 1955c: 12. (tx) Nobuchi 1975: 47.
- ceylonicus** (Schedl) 1959a: 477 (*Cryphalomorphus*). Holotype, sex?; Ceylon: Sabargamuva, Millawitiya Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Andaman Islands, Bengal, Uttar Pradesh in India/ Sri Lanka).
Hosts: *Lindera pulcherrima*.
References: (ds) Schedl 1959a: 477. (tx) Schedl 1959a: 477, 1979c: 56.
- cicatricosus** (Schedl) 1942c: 176 (*Lepicerinus*). Holotype, sex?; Neu-Guinea; Schedl Collection in NHMW, Wien.
Distribution: Fiji, New Caledonia, New Guinea.
References: (tx) Schedl 1942c: 176, 1952d: 343–344, 1978d: 73, 1979c: 57, 1979f: 103.
- coccotrypanoides** (Schedl) 1939e: 343 (*Lepicerinus*). Lectotype, sex?; Malaya, Selangor: Kuala Lumpur; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 59.
Distribution: Asia (Malaya).
References: (hb) Browne 1961c: 67. (ds) Browne 1961c: 67. (tx) Schedl 1939e: 343, 1940d: 588, 1952d: 343–344, 1979c: 59.
- communis** (Schaufuss) 1891: 12 (*Cryphalomorphus*). Holotype, sex?; Madagascar; Hamburg Museum, lost.
Distribution: Africa (Seychelles Islands), Madagascar.
Notes: (1) Schedl 1979c: 60 (reported 2 cotypes in his collection; if correct, these came from Eggers as a loan).
References: (ds) Alluaud 1900: 439–442; Anonymous 1892: 165; Eggers 1926g; Fairmaire 1892b; Hagedorn 1910d: 41, 46; Kleine 1913b: 121, 126, 1914b: 326; Kolbe 1910: 40; Sampson 1914: 386. (tx) Anderson, W. H. & Anderson 1971: 9; Eggers 1926a: 300; Fairmaire 1892b; Hagedorn 1910a: 83, 1913b: 254; Hopkins 1914: 119, 133, 1915b: 14; Michalski 1969b: 570; Sampson 1914: 386; Schaufuss 1891: 11–12, 1897a: 210; Schedl 1940d: 584, 588, 1961k: 523, 1969d: 11, 1971e: 11, 1977a: 45, 1977b: 70, 1979c: 60; Wood, S. L. 1954a: 996, 1960a: 27.
- concentralis** (Schedl) 1975f: 342 (*Cryphalophilus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 342, 1979c: 60.
- confragosus** (Sampson) 1914: 386 (*Cryphalomorphus*). Holotype, sex?; Seychelles, Mahe: Cascade Estate, 800–1000 feet; BMNH, London.
Distribution: Africa (Seychelles Islands).
References: (ds) Beaver 1987a: 20; Sampson 1914: 386; Schedl 1969d: 11. (tx) Sampson 1914: 386; Schedl 1952d: 344.
- corpulentus** (Schedl) 1965c: 54 (*Cryphalomorphus*). Holotype, sex?; Madagascar, Montagne d'Ambre; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
References: (tx) Schedl 1965c: 54, 1977a: 46, 1979c: 65.
- corrugatus** (Schedl) 1950d: 19 (*Stephanorhopalus*). Holotype ♀; Mauritius; Schedl Collection in NHMW, Wien.
Distribution: Africa (Mauritius Island).
References: (ds) Schedl 1969d: 11. (tx) Schedl 1950d: 19, 1962r: 94, 1969d: 11, 1977a: 46, 1979c: 66.
- creber** (Schedl) 1975f: 346 (*Cryphalomorphus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 346, 1979c: 68.
- crenatus** (Sampson) 1914: 385 (*Cryphalomorphus*). Syntypes, sex?; Seychelles, Mahe: from high damp forest between Trois Freres and Morne Seychellois, 1500–2000 feet; BMNH, London?
Distribution: Africa (Seychelles Islands).
Hosts: *Wormia ferruginea*.
References: (ds) Beaver 1987a: 20; Sampson 1914: 385; Schedl 1969d: 11. (tx) Sampson 1914: 385; Schedl 1952d: 344.
- cylindricus** (Schedl) 1959a: 476 (*Cryphalomorphus*). Syntypes, sex?; Ceylon: Sabargamuva, Millawitiya Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka/ Vietnam).
References: (ds) Schedl 1959a: 476, 1962b: 186, 1971a: 280. (tx) Schedl 1959a: 476, 1979c: 73.
- darwinii** Eichhoff 1878b: 497. Holotype, sex?; Hindostan (Birma); deposited in Dohrn Collection, now in Schedl Collection in NHMW, Wien.
Distribution: Asia (Burma/ Yunnan in China/ Uttar Pradesh in India/ Sri Lanka/ Vietnam), Indonesia (Java), New Guinea, Philippine Islands.
Hosts: *Cryptolepis buchanani*.
References: (ds) Hagedorn 1910d: 81; Kleine 1913b: 149, 1914b: 278. (tx) Eggers 1925: 159; Eichhoff 1878b: 387, 497; Hagedorn 1910a: 133; Hopkins 1914: 129; Lucas 1920: 572; Schedl 1939e: 337–338. (ms) Lucas 1920: 572.
- similis* Eggers 1923a: 142 (*Nigrites*). Lectotype, sex?; Java; USNM, Washington, designated by

- Anderson & Anderson 1971: 30. Synonymy: Wood 1985: 269.
References: **(bv)** Gray, B. 1974c. **(hb)** Gray, B. 1974c. **(tx)** Anderson, W. H. & Anderson 1971: 30; Eggers 1923a: 142; Schedl 1939c: 337–338, 1957b: 152, 1979c: 229; Wood, S. L. 1985: 269.
- major* Eggers 1927c: 69 (*Nigritus*). Lectotype, sex?; Philippines: Mindanao, Provinz Surigao, Surigao; USNM, Washington, designated by Anderson & Anderson 1971: 19. Synonymy: Wood 1985: 269.
References: **(cn)** Mathur & Singh 1960a: 36. **(hb)** Kalshoven 1958b: 180. **(ds)** Beeson 1961: 295; Mathur & Singh 1960a: 36; Schedl 1964c: 305, 1966b: 15. **(tx)** Anderson, W. H. & Anderson 1971: 19; Eggers 1927c: 69, 1929e: 53; Nobuchi 1983: 299; Schedl 1939c: 337–338, 1942d: 1, 1951i: 55, 1974c: 262, 1979c: 146; Wood, S. L. 1985: 269.
- cryptolepis* Schedl 1951i: 55. Holotype, sex?; Nakronda, Dehra Dun, U.P., India; Schedl Collection in NHMW, Wien. Synonymy: Wood 1985: 269.
References: **(tx)** Schedl 1951i: 55, 1955i: 214, 1979c: 71; Wood, S. L. 1985: 269.
- devius* (Schedl) 1975k: 280 (*Cryphalophilus*). Holotype, sex?; Transvaal: Barberton: TMP, Pretoria. Figures: Schedl 1975k: 281.
Distribution: Africa (South Africa).
References: **(tx)** Schedl 1975k: 280–281, 1979: 79.
- eggersi* (Schedl) 1962q: 490 (*Cryphalomorphus*). Lectotype, sex?; Philippines: Luzon, Manila; USNM, Washington, designated by Anderson & Anderson 1971: 20, automatic.
Distribution: Philippine Islands (Luzon).
References: **(tx)** Schedl 1962q: 490, 1979c: 154.
- minor* Eggers 1927c: 76 (*Cryphalomorphus*). Lectotype, sex?; Philippines: Luzon, Manila; USNM, Washington, designated by Anderson & Anderson 1971: 20, preoccupied by Eggers 1927c: 69.
References: **(tx)** Anderson, W. H. & Anderson 1971: 20; Eggers 1927c: 76, 1929e: 53; Schedl 1939c: 338, 1962q: 490, 1979c: 154.
- excellens* (Schedl) 1979g: 97 (*Cryphalomorphus*). Holotype, sex?; Papua New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1979g: 97.
- expers* (Blandford) 1894d: 85 (*Hypothenemus*). Syntypes 2, sex?; Kumamoto and Nagasaki, Japan; BMNH, London.
Figures: Nakane et al. 1963: 382.
Distribution: Asia (Japan).
Notes: (1) Wood 1966b: 22 (to *Scolytogenes*).
References: **(ds)** Hagedorn 1910d: 42; Kleine 1913b: 124, 1914b: 257; Nakane et al. 1963: 382; Nobuchi 1985c: 12. **(tx)** Blandford 1894d: 85; Hagedorn 1910a: 86; Hopkins 1915b: 24; Murayama 1961: 31; Nakane et al. 1963: 382; Schedl 1934f: 1640; Wood, S. L. 1966b: 22.
- fijianus* (Schedl) 1950f: 42 (*Lepicerinus*). Syntypes, sex?; Fiji: Viti Levu, several localities; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.
Distribution: Fiji Islands.
Hosts: *Plerandra grayi* leaf petioles.
Notes: (1) Schedl 1979c: 96 (citation of holotype invalid).
References: **(ds)** Beaver 1991: 54; Browne 1974a: 63. **(tx)** Schedl 1950f: 42, 1952d: 343–344, 1979c: 96.
- fugax* (Schedl) 1975f: 347 (*Cryphalomorphus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975f: 347, 1979c: 101.
- fujisanus* (Nobuchi) 1975: 46 (*Cryphalomorphus*). Holotype, sex?; Mt. Fuji, Yamanashi pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1, fig. 7.
Distribution: Asia (Japan).
Hosts: *Pterocarya rhoifolia*.
References: **(ds)** Nobuchi 1985c: 12. **(tx)** Nobuchi 1975: 46–47.
- fulgens* (Schedl) 1975f: 348 (*Cryphalomorphus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975f: 348, 1979c: 101.
- fulgidus* (Schedl) 1975f: 348 (*Cryphalomorphus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975f: 348, 1979c: 101.
- fulvipennis* (Nobuchi) 1975: 47 (*Cryphalomorphus*). Holotype, sex?; Yona, Okinawa pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1, fig. 9.
Distribution: Asia (Okinawa in Japan/Taiwan).
References: **(ds)** Nobuchi 1985c: 12. **(tx)** Nobuchi 1975: 47–48.
- ghanaensis* (Schedl) 1977d: 281 (*Cryphalomorphus*). Holotype, sex?; Ghana, Volta Region, N 6 degrees 70', E 0 degrees 03'; Schedl Collection in NHMW, Wien.
Distribution: Africa (Ghana).
References: **(tx)** Schedl 1977d: 281.
- gracilis* (Schedl) 1950f: 44 (*Lepicerinus*). Holotype, sex?; Fiji Islands; Eggers Collection, in NHMW, Wien.
Distribution: Fiji Islands, New Caledonia.
Hosts: *Garcinia* sp.

- References: (ds) Anonymous 1974a: 64. (tx) Schedl 1950f: 44, 1952d: 343–344, 1979c: 107.
- gramulatus (Wood)** 1960a: 31 (*Cryphalomorphus*). Holotype ♂; Kalabera area, Saipan, S. Mariana Islands; FMNH, Chicago.
Distribution: Micronesia (Saipan Island in Mariana Islands).
Hosts: *Cerbera manghas*.
References: (tx) de Ruette 1970: 99; Wood, S. L. 1960a: 31.
- grobleri (Schedl)** 1962h: 68 (*Cryphalomorphus*). Holotype, sex?, Cape Province, Alexandria; NICP, Pretoria.
Distribution: Africa (South Africa).
Hosts: *Strychnos decussata*.
References: (tx) Schedl 1962h: 68, 1962k: 1069, 1979c: 113.
- hirtus (Wood)** 1974a: 18 (*Cryphalomorphus*). Holotype ♀; 16 km S Oaxaca, Oaxaca, Mexico; Wood Collection.
Distribution: North America (Oaxaca, Puebla in Mexico).
Hosts: *Ipomoea* sp.
References: (ds) Atkinson et al. 1986: 66; Wood, S. L. 1982b: 864. (tx) Wood, S. L. 1974a: 18, 1982b: 864.
- hobohmi (Schedl)** 1955i: 215 (*Cryphalomorphus*). Lectotype, sex?, S.W. Africa, Otjéwarongo; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 119.
Figures:
Distribution: Africa (Namibia).
References: (tx) Schedl 1955i: 211, 215, 1961k: 526, 1965c: 54, 1979c: 119.
- hylesinopsis (Schedl)** 1975f: 349 (*Cryphalomorphus*). Holotype, sex?, New Guinea, Wan, McAdam Park; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 349, 1979c: 120.
- indicus Wood** 1989: 184. Holotype, sex?, Amarkantak, Rewah State, C. I. 3500 ft.; FRI, Dehra Dun.
Distribution: Asia (Burma/ Kashmir, Madhya, Uttar Pradesh in India).
Hosts: *Hedera helix*, *Moringa pterygosperma*, *Stereospermum suaveolens*, *Valleria haynei*.
Notes: (3) This is *indicus* Beeson 1941 (1961: 240), nomen nudum. Schedl 1979c: 127 (cited it as a synonym of *varius* Schedl; however, 2 distinct species are represented by these names).
References: (tx) Schedl 1939e: 343, 1940d: 588, 1959a: 477, 1979c: 123; Wood, S. L. 1989: 184.
- insularis (Nobuchi)** 1975: 48 (*Cryphalomorphus*). Holotype, sex?, Kawada, Miyake Is., Tokyo, Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 1.
Distribution: Asia (Mikura Island in Japan).
- References: (ds) Nobuchi 1985c: 12. (tx) Nobuchi 1975: 48, pl. 1.
- jalapae (Letzner)** 1848b: 99 (*Bostrichus*). Syntypes, sex?, taken from Jalapa root in Mexico; not located.
Distribution: North America (Mexico), transported through commerce to many parts of the world but not established there.
Hosts: Presumably *Exogonium jalapa*.
References: (av) Imhoff 1856: 228. (cn) Hagedorn 1913a; Kleine 1932a: 297; Mumford 1960: 22; Swaine 1918a: 90. (cc) Wichmann 1955a: 96. (hb) Blandford 1894b; Chamberlin 1939: 316; Dombrowsky 1887; Eichhoff 1881a: 46, 187; Hagedorn 1903a, 1913a; Kleine 1932a: 297; Lengerken 1939: 63, 1954: 83; Swaine 1918a: 90; Wachtl 1876a: 458; Wood, S. L. 1953: 73, 1954a: 997, 1982b: 866. (ds) Anonymous 1926c: 518; Blackwelder 1947: 778; Blandford 1894b: 261, 1897b, 1904: 226; Calwer 1884, 1893; Chamberlin 1939: 316; Cola 1973; Ferrer 1942; Gemminger & Harold 1872: 2683; Hagedorn 1903a, 1910d: 43, 1913a: 46–47; Hamilton 1889: 158, 1894b: 406; Henshaw 1885: 148, 1895: 44; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1935, 1951; Kleine 1914b: 349, 1932a: 297, 1934a: 142; Koltze 1901: 153; Lacordaire 1866: 383; Leonard 1928: 518; Lomnicki 1913b: 148; Mumford 1960: 22; Nunberg 1964a: 235; Pittioni 1943: 176; Reitter 1894a: 70, 1916: 289, 350; Schedl 1980a: 17; Schwarz 1883: 40, 140, 1886: 42, 1901: 432; Seidlitz 1891a: 562, 1891b: 608; Stein 1868: 114; Stein & Weise 1877: 164; Swaine 1909: 92; Townsend 1889: 235; Wachtl 1876a: 458; Wichmann 1927a: 79, 1955a: 96; Wood, S. L. 1977a: 69, 1982b: 866. (tx) Balachowsky 1949a: 201; Berger 1917: 226–248; Blandford 1894b: 261, 1904: 226; Chamberlin 1939: 316; Dombrowsky 1887; Eichhoff 1878b: 134, 1881a: 46, 74, 187, 1883a: 110, 133; Fauvel 1889; Ferrari 1867a: 12, 14, 16; Hagedorn 1910a: 86; Hopkins 1915b: 35; Kuhn 1913: 1055; Lacordaire 1866: 383; Letzner 1844, 1848b: 99; Lindemann 1875c: 321; Portevin 1935: 324; Reitter 1894a: 70, 1908a: 56, 1913a: 68, 1916: 289, 350; Schedl 1934f: 1639, 1938f: 41, 1940a: 341, 1940d: 584, 588, 1959a: 476, 1965i: 368, 1980a: 17; Schwarz 1886: 42; Seidlitz 1891a: 562, 1891b: 608; Swaine 1909: 92, 1918a: 90; Wood, S. L. 1953: 73, 1954a: 997, 1982b: 866.
- knabi (Hopkins)** 1915b: 34 (*Eruopirides*). Holotype ♀; Cordoba, Mexico; USNM, Washington.
Figures: Bright 1972d: 59.
Distribution: Antilles Islands (Cuba/ Dominican Republic/ Guadalupe/ Jamaica/ Tortola in British West Indies, introduced?), Asia (Japan, introduced?), North America (Guatemala/ Honduras/ Chiapas, Nayarit, Oaxaca, Veracruz in Mexico/

Florida in USA, introduced?), South America (Brazil/ Venezuela? introduced?).

Hosts: *Calonciton tannifolium*, *Candiosperma holacobiium*, *Ipomoea pres-carpa*, *Serjania* spp., and other lianas.

Notes: Although known primarily from tropical America where it has no near relatives, this species was probably introduced; its occurrence in Japan supports that supposition.

References: (hb) Burgos & Sancedo 1982b: 865; Wood, S. L. 1982b: 865. (ds) Atkinson & Equihua 1985c: 356, 1985: 100; Atkinson et al. 1986: 66; Bright 1972d: 61, 1976d: 61, 1985c: 175; Burgos & Sancedo 1983: 103; Ferrer 1942; Schedl 1940a: 342, 1940d: 588, 1960d: 75; Wood, S. L. 1982b: 865. (tx) Bright 1972d: 59, 1976d: 61, 1985c: 175; Hopkins 1915b: 34; Schedl 1940a: 342, 1940d: 588, 1959a: 477, 1960a: 75; Wood, S. L. 1966b: 23, 1977c: 385, 1982b: 865.

floridensis Hopkins 1915b: 34 (*Ernoporides*). Holotype ♀; Biscayne, Florida [USA]; USNM, Washington. Synonymy: Wood 1966b: 23.

References: (hb) Chamberlin 1939: 315; Wood, S. L. 1954a: 998. (ds) Blatchley & Leng 1916: 604; Chamberlin 1939: 315. (tx) Balachowsky 1949a: 201; Blatchley & Leng 1916: 604; Chamberlin 1939: 315; Hopkins 1915b: 34; Schedl 1940d: 588; Wood, S. L. 1954a: 979, 998, 1966b: 22.

ritchiei Sampson 1918: 295 (*Hypothenemus*). Syntypes ♀; Jamaica; BMNH, London. Synonymy: Wood 1966b: 23.

References: (ds) Cowdery 1926: 27. (tx) Eggers 1940a: 108; Sampson 1918: 295; Schedl 1940d: 588; Wood, S. L. 1966b: 23.

carabicus Schedl 1951m: 96 (*Cryphalomorphus*). Syntypes, sex?; Guadeloupe, Delaney; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 23.

References: (tx) Schedl 1951m: 96, 1979c: 53; Wood, S. L. 1966b: 23.

minutissimus Schedl 1951m: 97 (*Cryphalomorphus*). Holotype ♂; Guadeloupe, Env. de Trois-Rivieres; Schedl Collection in NHMW, Wien. Synonymy: Wood 1977c: 385.

References: (tx) Schedl 1951m: 97, 1979c: 156; Wood, S. L. 1977c: 385.

subtriatus Schedl 1952d: 360 (*Cryphalomorphus*). Holotype, sex?; Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 23.

References: (tx) Schedl 1951m: 97, 1952d: 360, 1979c: 244; Wood, S. L. 1966b: 23.

alienus Schedl 1976a: 65 (*Cryphalomorphus*). Holotype, sex?; Corcovado, Guanabara, Brasil; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 175.

References: (tx) Schedl 1976a: 65; Wood, S. L. 1989: 175.

landolphiae (Schedl) 1961e: 133 (*Cryphalomorphus*). Holotype ♀; Madagascar, Perinet; IRSM, Madagascar.

Distribution: Madagascar.

References: (hb) Schedl 1977a: 45. (tx) Schedl 1961e: 133, 1965c: 56, 1977a: 46, 1979c: 136.

leprosulus (Browne) 1974a: 66 (*Cryphalomorphus*). Holotype, sex?; Fiji: Viti Levu, Nandarivatu; BMNH, London.

Distribution: Fiji Islands.

Hosts: *Garcinia* sp.

References: (ec) Roberts 1976: 377. (hb) Roberts 1976: 377. (ds) Browne 1974a: 63, 66; Roberts 1976: 377. (tx) Browne 1974a: 66.

longipennis (Eggers) 1936b: 30 (*Cryphalomorphus*). Holotype, sex?; Abyssinia: Mt. Chillalo in the Forest, circa 9000 feet; BMNH, London.

Distribution: Africa (Ethiopia).

References: (tx) Browne 1965: 191; Eggers 1936b: 30; Schedl 1940d: 588, 1961k: 527.

mauritanus (Schedl) 1965c: 56 (*Cryphalomorphus*). Holotype, sex?; Mauritius; Schedl Collection in NHMW, Wien.

Distribution: Africa (Mauritius Island).

References: (ds) Schedl 1969d: 10. (tx) Schedl 1965c: 10, 56, 1969d: 10, 1977a: 47, 1979c: 150.

micans (Eggers) 1927b: 396 (*Stephanoderes*). Holotype, sex?; Java (Tenggergebirge); Eggers Collection, in NHMW, Wien.

Distribution: Indonesia (Java).

References: (tx) Eggers 1927b: 396, 1940b: 62; Schedl 1952d: 343–344, 1979c: 152.

minor (Eggers) 1927c: 69 (*Negritus*). Holotype, sex?; Philippines: Mindanao, Provinz Zamboanga, Port Banga; Eggers Collection [not mentioned by Anderson & Anderson 1971 or Schedl 1979c].

Distribution: Asia (Malaya/ Vietnam), Philippine Islands (Mindanao).

References: (ds) Schedl 1962b: 186, 1966b: 15, 23, 1971c: 363. (tx) Eggers 1927c: 69, 76, 1929e: 53; Schedl 1939e: 337, 1940d: 588, 1942c: 175, 1942d: 2, 1950f: 44, 1951i: 50, 1962q: 490.

mus (Schedl) 1975f: 349 (*Cryphalomorphus*). Holotype, sex?; New Guinea, War; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 349, 1979c: 162.

nanulus (Wood) 1960a: 29 (*Cryphalomorphus*). Holotype, sex?; Mt. Tagpochau, Saipan; FMNH, Chicago.

Distribution: Micronesia (Saipan, Tinian in Caroline Islands).

References: (tx) de Rnette 1970: 99; Wood, S. L. 1960a: 29.

nigellatus (Schedl) 1950f: 44 (*Lepicerinus*). Holotype, sex?; Fiji; Eggers Collection, in NHMW, Wien.

- Distribution: Fiji Islands.
References: (ds) Browne 1974a: 63. (tx) Schedl 1950f: 44, 1951m: 97, 1952d: 343–344, 1979c: 166.
- nubilus** (Wood) 1960a: 30 (*Cryphalomorplus*).
Holotype ♀: Ine I., Arno Atoll, Marshall Islands; USNM, Washington.
Distribution: Micronesia (Caroline Islands/ Arno Atoll in Marshall Islands).
Hosts: *Scaevola frutescens*.
References: (tx) de Ruelle 1970: 99; Wood, S. L. 1960a: 30.
- ocularis** (Schedl) 1965e: 365 (*Cryphalomorplus*).
Holotype, sex?; Congo ex belge-Leopoldville; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: (tx) Schedl 1965e: 365, 1979c: 177.
- onyanganus** (Schedl) 1941d: 391 (*Letznerella*).
Lectotype, sex?; Mt. Cameroon, Onyanga, 8100 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 175.
Distribution: Africa (Cameroon/ Kenya/ Nigeria/ South Africa/ Tanzania/ Uganda).
Hosts: *Acacia decurrens*, *Celtis* sp., *Euclea lanceolata*.
References: (tx) Browne 1973a: 290; Schedl 1940d: 585, 1941d: 391–392, 1952d: 344, 1961k: 527, 1979c: 175.
similaris Schedl 1965f: 8 (*Cryphalomorplus*).
Holotype, sex?; Uganda, Mpanga; BMNH, London. Synonymy: Browne 1973a: 290.
References: (ds) Browne 1970: 540; Schedl 1969d: 5, 1982: 275. (tx) Browne 1973a: 290; Schedl 1965c: 55, 1965f: 8, 1979c: 229.
- opacus** (Schedl) 1959a: 477 (*Cryphalomorplus*).
Holotype, sex?; Ceylon: Sabargamuva, Millawitiya Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka).
References: (ds) Schedl 1959a: 477. (tx) Schedl 1959a: 477, 1979c: 179.
- pacificus** (Schedl) 1942c: 176 (*Lepicerinus*). Lectotype, sex?; Piti, Guam; Schedl Collection in NHMW, Wien.
Distribution: Micronesia (Caroline Islands/ Guam, Truk in Mariana Islands).
Hosts: Breadfruit.
References: (ds) Wood, S. L. 1960a: 27. (tx) Schedl 1942c: 176, 1950f: 44, 1952d: 343–345, 1960a: 27–28, 1979c: 182.
- papuanus** (Schedl) 1974d: 459 (*Cryphalomorplus*).
Holotype, sex?; Panganda logging area, Walnut Valley, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Erodia* sp., *Nothofagus* sp.
References: (tx) Schedl 1974d: 459, 1979c: 183.
- papuensis** Wood 1989: 179. Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
- References: Wood, S. L. 1989: 179, 1992b: 79–80.
papuanus Schedl 1975f: 352 (*Xylocryptus*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1974.
References: (tx) Schedl 1975f: 352, 1979c: 184; Wood, S. L. 1989: 179, 1992b: 79–80.
- paradoxus** Wood 1992b: 80. Holotype, sex?; Papua New Guinea; Schedl Collection in NHMW, Wien, automatic.
Distribution: New Guinea.
References: (tx) Wood, S. L. 1992b: 80.
- papuanus* Schedl 1979g: 97 (*Cryphalomorplus*).
Holotype, sex?; Papua New Guinea; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1974.
References: (tx) Schedl 1979g: 97; Wood, S. L. 1992b: 80.
- parvatis** (Wood) 1974a: 17 (*Cryphalomorplus*).
Holotype ♀; La Lima, Cortez, Honduras; Wood Collection.
Distribution: North America (Costa Rica/ Honduras).
Hosts: *Cestrum scandens*.
References: (tx) Wood, S. L. 1974a: 17, 1982b: S62–S63.
- parvus** (Hopkins) 1915b: 11 (*Hypothenoides*).
Holotype, sex?; Calapan, Philippine Islands; USNM, Washington.
Distribution: Philippine Islands.
Hosts: *Lemon silvestre*.
References: (ds) Kleine 1934a: 148; Schedl 1966b: 23. (tx) Hopkins 1915b: 11; Nobuchi 1983: 300.
- pityophthorinus** (Schedl) 1943b: 39 (*Lepicerinus*).
Holotype, sex?; Balbolan [=Balbalan], Philippines; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands.
Notes: (1) Schedl 1979c: 195 (Errors in spelling both generic and specific names in the original publication are considered lapsus calami and were corrected).
References: (ds) Schedl 1966b: 24. (tx) Nobuchi 1983: 300; Schedl 1943b: 39, 1952d: 345, 1979c: 195.
- pleiocarpae** (Schedl) 1957d: 51 (*Cryphalomorplus*). Holotype, sex?; Congo Belge: Yangambi; MRCB, Tervuren.
Figures: Schedl 1961k: 528.
Distribution: Africa (Angola/ Ghana/ Zaire).
Hosts: *Pleiocarpa micrantha*, *P. tubicina*.
References: (cc) Schedl 1958d: 188, 193, 1961k: 527, 1972c: 67. (hb) Schedl 1961k: 527. (ds) Ferreira 1965: 1114; Schedl 1959p: 17, 1961k: 527, 1964j: 40. (tx) Schedl 1957d: 51, 1961e: 133, 1961k: 527–528, 1965f: 8, 1979c: 196.

- praeda (Browne)** in Beaver & Browne 1978: 589 (*Ptilopodius*). Holotype, sex?; Malaysia; BMNH, London.
Distribution: Asia (Malaya).
References: (tx) Beaver 1991: 53; Beaver & Browne 1978: 604; Browne 1979: 589.
- puerarae Choo & Woo** 1989: 58. Holotype ♀; Korea: Sacheon, Gyeongnam province; not given.
Figures: Choo & Woo 1989: 60.
Distribution: Asia (Korea).
Hosts: *Pueraria thuyergiana*.
References: (tx) Choo & Woo 1989: 58–60.
- pumilionides (Schedl)** 1977f: 500 (*Cryphalomorphus*). Holotype, sex?; India: Kerala; Schedl Collection in NHMW, Wien.
Distribution: Asia (Kerala in India).
References: (tx) Schedl 1977f: 500.
- pumilus (Wood)** 1960a: 29 (*Cryphalomorphus*). Holotype ♀; northwest slope, Mt. Lasso, Timian; FMNH, Chicago.
Distribution: Micronesia (Timian Island).
References: (tx) Wood, S. L. 1960a: 28–29.
- punctatulus Nobuchi** 1976: 72. Holotype, sex?; Mt. Inugaki, Fukuoka pref., Japan; Nobuchi Collection, Ibaraki, automatic.
Figures: Nobuchi 1975: pl. 1, fig. 11.
Distribution: Asia (Japan).
Hosts: *Trachelospermum asiaticum*.
References: (ds) Nobuchi 1985c: 12. (tx) Nobuchi 1976: 72.
- punctatus Nobuchi** 1975: 49 (*Cryphalomorphus*). Holotype, sex?; Inugaki, Fukuoka pref., Japan; Nobuchi Collection, Ibaraki, preoccupied by Schedl 1951.
References: (tx) Nobuchi 1975: 49, 1976: 72, 1985c: 12.
- nobuchii Schedl** 1980: 118 (*Cryphalomorphus*). Holotype, sex?; Mt. Inugaki, Fukuoka pref., Japan; Nobuchi Collection, Ibaraki, automatic.
Notes: (1) This is an unneeded replacement name.
References: (tx) Choo & Woo 1989a; Schedl 1980: 118.
- punctatus (Schedl)** 1951i: 55 (*Cryphalomorphus*). Holotype, sex?; Himalaya, Darjeeling; Schedl Collection in NHMW, Wien.
Figures: Nobuchi 1975: pl. 1.
Distribution: Asia (Bengal in India).
References: (tx) Nobuchi 1975: pl. 1, 1976: 72; Schedl 1951i: 55, 1952d: 345, 1979c: 204.
- puncticollis (Schedl)** 1950f: 43 (*Lepicerinus*). Syn-types, sex?; Fiji: Viti Levu, several localities; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.
Distribution: Fiji Islands.
Notes: (1) Schedl 1979c: 205 (citation of holotype invalid).
- References: (ds) Browne 1974a: 63. (tx) Beaver 1991: 93; Schedl 1950f: 43, 1952d: 345, 1979c: 205.
- grosseopunctatus Browne** 1974a: 66 (*Cryphalomorphus*). Holotype, sex?; Fiji: Viti Levu, Nausori Highlands; BMNH, London. Synonymy: Beaver 1991: 93.
Notes: (3) Schedl 1980d: 118 (treated as a synonym of *gracilis*).
References: (ec) Roberts 1976: 376. (hb) Roberts 1976: 376. (ds) Browne 1974a: 63; Roberts 1976: 376. (tx) Beaver 1991: 93; Browne 1974a: 66; Schedl 1980d: 118.
- pygmaeolus (Schedl)** 1971e: 12 (*Cryphalomorphus*). Holotype, sex?; Cote d'Ivoire, Forêt du Banco; Schedl Collection in NHMW, Wien.
Distribution: Africa (Ivory Coast).
References: (tx) Schedl 1971e: 12, 1979c: 207.
- quadridens (Browne)** 1984f: 71 (*Hypothenemus*). Holotype, sex?; New Guinea: Morobe District, Wan, 1700 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Macaranga aleurutoides*.
References: (tx) Beaver 1991: 54; Browne 1984f: 71.
- robustus (Schedl)** 1955c: 32 (*Cryphalomorphus*). Lectotype, sex?; Somalia It: Belet Uen-Cabredarre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 212.
Distribution: Africa (Somalia).
References: (tx) Schedl 1955c: 32, 1961k: 533, 1979c: 212.
- rusticus (Wood)** 1974a: 18 (*Cryphalomorphus*). Holotype ♀; 33 km N Juchitlan, Jalisco, Mexico; Wood Collection.
Figures: Atkinson et al. 1986: 73.
Distribution: North America (Chiapas, Guajuato, Jalisco, Michoacan in Mexico).
Hosts: *Ipomoea* sp.
References: (ec) Equihua & Atkinson 1986: 633. (hb) Atkinson & Equihua 1985c: 356; Atkinson et al. 1986: 66; Burgos & Sancedo 1983: 101; Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 864. (ds) Atkinson & Equihua 1985c: 356; Atkinson et al. 1986: 66; Burgos & Sancedo 1983: 101; Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 864. (tx) Atkinson et al. 1986: 73; Wood, S. L. 1974a: 18, 1982b: 864–865.
- samoanus (Browne)** 1977c: 61 (*Cryphalomorphus*). Holotype, sex?; Samoa: Upolu; BMNH, London.
Distribution: Samoan Islands.
Hosts: *Psidium guajava*, *Theobroma cacao*.
References: (hb) Beaver 1976b: 534. (ds) Beaver 1976d: 534. (tx) Browne 1977c: 61.
- scolytomimoides (Nobuchi)** 1975: 50 (*Cryphalomorphus*). Holotype, sex?; Asakawa, Tokyo, Japan.
Figures: Nobuchi 1975: pl. 1.
Distribution: Asia (Japan).
Hosts: *Trachelospermum asiaticum*.

- References: (ds) Nobuchi 1953c: 12. (tx) Nobuchi 1975: 50, pl. 1.
- separandus** (Schedl) 1965c: 56 (*Cryphalomorphus*). Holotype, sex?, Madagascar; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
References: (tx) Schedl 1965c: 56, 1977a: 47, 1979c: 225.
- setifer** (Wood) 1974a: 18 (*Cryphalomorphus*). Holotype ♀; Volcan Pacaya, Esquinla, Guatemala; Wood Collection.
Distribution: North America (Guatemala).
Hosts: Liana.
References: (ds) Wood, S. L. 1982b: 863. (tx) Wood, S. L. 1974a: 18, 1982b: 863–864.
- sodalis** (Schedl) 1965c: 55 (*Cryphalomorphus*). Holotype, sex?, Madagascar, Perinet; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
Hosts: *Landolphia* sp.
References: (tx) Schedl 1965c: 55, 1977a: 47, 1979c: 231.
- spirostachius** (Schedl) 1958a: 557 (*Ptilopodius*). Holotype, sex?, Zululand: W. Umfolosi Riv.; BMNH, London.
Distribution: Africa (South Africa).
Hosts: *Spirostactys* sp.
References: (ds) Schedl 1961k: 455, 1965h: 111. (tx) Schedl 1958a: 557, 1961k: 455, 1979c: 235.
- splendens** (Schedl) 1975f: 350 (*Cryphalomorphus*). Holotype, sex?, Bulolo, Compt. 5, Taun logging area, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 350, 1979c: 235.
- squamitilis** (Schedl) 1977f: 500 (*Cryphalomorphus*). Holotype, sex?, Vietnam: Dinh by NE of Hanoi; Schedl Collection in NHMW, Wien.
Distribution: Asia (Vietnam).
References: (tx) Schedl 1977f: 500.
- squamosus** (Schedl) 1942c: 175 (*Lepicerinus*). Holotype, sex?, New-Guinea, Kapa-Kapa; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (bv) Gray, B. 1974c. (hb) Gray, B. 1974c. (ds) Browne 1961c: 79. (tx) Schedl 1942c: 175, 1943b: 40, 1951m: 96, 1952d: 345, 1979c: 236.
- squamulosus** (Eggers) 1936d: 633 (*Cylindrotomicus*). Holotype, sex?, Vorderindien (Mysore: Jakkur); BMNH, London, 1 Eggers cotype, in NHMW, Wien.
Distribution: Asia (Karnataka in India).
Hosts: Wild creeper.
References: (tx) Eggers 1936d: 633; Schedl 1957b: 152, 1979c: 236.
- sumatranus** (Schedl) 1970b: 358 (*Chiloxylon*). Holotype, sex?, Sumatra, Palembang to Akita (Japan), imported; PPST, Tokyo.
Distribution: Indonesia (Sumatra).
Hosts: Pulau log.
References: (ds) Browne 1980d: 491; Ohno, Yoneyama, & Nakazawa 1982b: 7. (tx) Schedl 1970b: 358, 1979c: 246.
- tonsus** (Schedl) 1969c: 49 (*Cryphalomorphus*). Holotype, sex?, N. Vietnam: Hoa Binh, Tonkin.
Distribution: Asia (Vietnam).
References: (tx) Schedl 1969c: 49, 1979c: 254.
- tricolor** (Lea) 1910: 141 (*Cryphalus*). Holotype, sex?, Queensland: Cairns; SAM, Adelaide.
Distribution: Australia (Queensland).
Hosts: *Millettia megasperma*.
References: (ds) Brimblecombe 1953: 21; Schedl 1972a: 144. (tx) Lea 1910: 141; Schedl 1938f: 34, 41, 1940d: 588.
- trucis** (Wood) 1974a: 19 (*Cryphalomorphus*). Holotype ♀; 16 km S Oacaca, Oaxaca, Mexico; Wood Collection.
Distribution: North America (Oaxaca, Puebla in Mexico).
Hosts: *Ipomoea* sp.
References: (ds) Wood, S. L. 1982b: 866. (tx) Wood, S. L. 1974a: 19, 1982b: 866.
- uncatus** (Schedl) 1971c: 373 (*Cryphalophilus*). Holotype, sex?, Ins. Kei (Planten); MNHN, Paris.
Distribution: Indonesia (Kei Island), Niue Island, Tahiti.
References: (tx) Schedl 1971c: 373, 1979c: 260.
- ater** Schedl 1972a: 146 (*Cryphalomorphus*). Holotype, sex?, Queensland: Yarraman; Queensland Museum, Brisbane, preoccupied by Eggers 1923. Synonymy: Beaver 1991: 93.
References: (tx) Beaver 1991: 93; Schedl 1940b: 434, 1950g: 894, 1972a: 146, 1972b: 267, 1979c: 29.
- usagarius** (Eggers) 1922b: 169 (*Neocryphalus*). Holotype, sex?, Sudost-Usagra in Deutsch-Ostafrika; Methner Collection in MNH, Berlin, 1 Eggers cotype, in NHMW, Wien.
Distribution: Africa (Kenya/ Tanzania).
References: (hb) Roberts 1969: 123. (ds) Schedl 1971e: 5. (tx) Eggers 1922b: 169; Schedl 1950d: 2, 1952d: 345, 1955i: 215, 1961k: 534, 1979c: 262.
- varius** (Schedl) 1975f: 350 (*Cryphalomorphus*). Holotype, sex?, Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 350, 1979c: 265.

Genus *Hemicryphalus* Schedl

- HEMICRYPHALUS SCHEDL 1963h: 264. Type-species: *Eidophelus argutus* Wood, original designation.
Keys: Wood 1960a: 32.
Notes: (3) Schedl 1965a: 341 (used *Taphropteris*, nomen nudum, to cite a member of this genus).
References: (hb) Wood, S. L. 1986a: 90. (ds)

Wood, S. L. 1986a: 90. (**tx**) Schedl 1963h: 264, 1965a: 341; Wood, S. L. 1986a: 90.

argutus (Wood) 1960a: 33 (*Eidophelus*). Holotype ♀; Saipan, S. Mariana Is.; FMNH, Chicago.

Figures: Wood 1960a: 33.

Distribution: Micronesia (Saipan in Mariana Islands).

References: (**tx**) de Ruette 1970: 101; Schedl 1963h: 264; Wood, S. L. 1960a: 33.

atomus (Wood) 1960a: 34 (*Eidophelus*). Holotype ♂; Kolonia (Yaptown), Yap, Yap Is.; USNM, Washington.

Distribution: Micronesia (Yap Islands).

References: (**tx**) Schedl 1963h: 264; Wood, S. L. 1960a: 34.

incomptus (Wood) 1960a: 32 (*Eidophelus*). Holotype ♀; Iwo Jima, Volcano Is.; FMNH, Chicago. Distribution: Micronesia (Iwo Jima in Volcano Islands).

References: (**tx**) Schedl 1963h: 264; Wood, S. L. 1960a: 32.

Genus *Eidophelus* Eichhoff

EIDOPHELUS EICHHOFF 1875: 200. Type-species: *Eidophelus imitans* Eichhoff, monobasic.

Phellodendrophagus Krivolutsкая 1958: 150. Type-species: *Phellodendrophagus elegans* Krivolutsкая = *Eidophelus imitans* Eichhoff, monobasic. Synonymy: Schedl 1962q: 493.

References: (**tx**) Krivolutsкая 1958: 150; Schedl 1962q: 493.

References: (**hb**) Wood, S. L. 1986a: 90. (**ds**) Wood, S. L. 1986a: 90. (**tx**) Choo 1983: 76; Choo, Woo, & Nobuchi 1985b; Eichhoff 1875: 200; Hopkins 1915b: 8; Murayama 1930: 21–22; Niisima 1917: 1; Schedl 1935a: 158, 1962q: 493; Wood, S. L. 1960a: 31, 1986a: 90.

borneensis Browne 1984a: 153. Holotype, sex?; Baram (Saravak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Saravak in Borneo).

Hosts: *Terminalia* sp.

References: (**tx**) Browne 1984a: 153.

gracilis Browne 1984a: 152. Holotype, sex?; Lasibu (New Britain) to Nagoya (Japan), imported; BMNH, London.

Distribution: New Britain Island.

Hosts: *Labiella* log.

References: (**ds**) Ohno, Yoshioka, et al. 1989: 60. (**tx**) Browne 1984a: 152.

imitans Eichhoff 1875: 201. Syntypes, sex?; Japan; 1 in BMNH, London.

Figures: Krivolutsкая 1958: 150, Kusch 1967: 10.

Distribution: Asia (Uttar Pradesh in India/ Japan/ Korea/ Nepal/ Sri Lanka/ Sakhalin Island in E USSR).

Hosts: *Phellodendron sachalinensis*, *Quercus acutissima*, *Toxicodendron trichocarpum*.

References: (**cn**) Shiraki 1952. (**ds**) Blandford 1894c; Cho 1957; Choo & Woo 1985: 165; Hagedorn 1910d: 69; Kleine 1913b: 139, 1914b: 258, 1934a: 158; Ko 1969: 276; Murayama 1929b: 2, 1930b: 17, 1937b: 375, 1954b: 200; Nobuchi 1967: 20; Shiraki 1952. (**tx**) Choo 1983: 76; Eichhoff 1875: 201, 1877a: 118; Hagedorn 1910a: 90; Hopkins 1914: 121; Murayama 1930b: 17, 21–22, 31, 1937b: 375, 1954b: 200; Nobuchi 1981c: 29; Schedl 1934f: 1643, 1962b: 191.

elegans Krivolutsкая 1958(?): (p. 2, 150) (*Phellodendrophagus*). Holotype, sex?; Sakhalin Island; Zoological Institute, Academy of Science, Vladivostok. Synonymy: Wood 1989: 172. Notes: Krivolutsкая 1958: 150 apparently is not the original description of this name.

References: (**hb**) Krivolutsкая 1973: 137.

(**ds**) Krivolutsкая 1965a: 236, 1973: 137; Krivolutsкая & Kupyanskaya 1970. (**tx**) Krivolutsкая 1958: 150; Schedl 1962q: 493; Wood, S. L. 1989: 172.

nitidus Schedl 1959a: 475 (*Ptilopodius*). Holotype ♀; Ceylon: Sabargamuva, Millawitiya Estate; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 172.

References: (**ds**) Evans, D., Lowe, & Hunt 1978; Kusch 1967; Schedl 1959a: 475, 1965g: 20. (**tx**) Kusch 1967: 10; Schedl 1959a: 475, 1979c: 170; Wood, S. L. 1989: 172.

minutissimus Schedl 1962b: 191. Holotype ♀; Tonkin: Saigon; MNHN, Paris.

Distribution: Asia (Vietnam).

Notes: (1) Schedl 1979c: 156 (suggested a transfer to *Acanthotomicus* [not seen by us]).

References: (**tx**) Schedl 1962b: 191, 1979c: 156.

minutus Blandford 1894d: 88. Holotype, sex?; Chiusenji, Japan; BMNH, London.

Distribution: Asia (Japan).

References: (**ds**) Blandford 1894c: 579; Hagedorn 1910d: 69; Kleine 1913b: 139, 1914b: 258; Murayama 1954b: 165. (**tx**) Blandford 1894d: 88; Hagedorn 1910a: 90; Lucas 1920: 256; Murayama 1954b: 165; Nobuchi 1981c: 29; Schedl 1934f: 1643. (**ms**) Lucas 1920: 256.

samoanus Schedl 1972b: 268. Holotype, sex?; Samoa, Upolu; Schedl Collection in NHMW, Wien.

Distribution: Samoan Islands.

References: (**ds**) Beaver 1976b: 536. (**tx**) Schedl 1972b: 268, 1979c: 218.

sumatranus Schedl 1961c: 72. Holotype, sex?; Sumatra's Ostkuste, Bandar Baroe, 850 m; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

Notes: (1) Schedl 1979c: 246 (suggested a transfer to *Acanthotomicus* [not seen by us]).

References: (**tx**) Schedl 1961c: 72, 1979c: 246.

Genus *Hypocryphalus* Hopkins

HYPOCRYPHALUS HOPKINS 1915b: 8, 41. Type-species: *Hypocryphalus rotundus* Hopkins, original designation.

Dacryphalus Hopkins 1915b: 8, 42. Type-species: *Dacryphalus obesus* Hopkins, original designation. Synonymy: Schedl 1938f: 48.

References: (tx) Hopkins 1915b: 8, 42; Schedl 1938f: 48, 1960g: 142; Wood, S. L. 1954a: 999.

References: (ay) Nobuchi 1969a: 58. (hb) Beaver 1987a: 18; Browne 1961c: 67; Schedl 1961k: 542, 1977a: 57; Wood, S. L. 1982b: 871. (ds) Beaver 1987a: 18; Beeson 1938: 288; Blackwelder 1947: 778; Brimblecombe 1953: 19; Browne 1961c: 67; Kalshoven 1945: 8; Schedl 1961k: 542, 1966b: 25, 1977a: 57; Wood, S. L. 1982b: 871. (tx) Arnett 1960: 1043, 1968: 1043; Hopkins 1915b: 8, 41; Schedl 1938f: 48, 1939i: 382, 1959a: 483, 1961k: 542, 1962h: 67, 1962m: 697–699, 1963j: 478, 1964i: 246–249, 1964m: 305, 1977a: 57; Wood, S. L. 1954a: 991, 1960a: 13, 1961a: 45, 1982b: 871, 1986a: 90.

aciculatus Schedl 1939e: 341. Lectotype, sex?; Malaya, Pahang; Semangkok For. Res., 2000 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 10.

Distribution: Asia (Malaya).

References: (tx) Schedl 1939e: 341, 1962r: 106, 1979c: 10.

afiamalus Schedl 1951k: 148. Lectotype, sex?; Upolu, Afiamalu: 2100 feet; Schedl Collection in NHMW, Wien.

Distribution: Samoan Islands.

References: (ds) Beaver 1976b: 534. (tx) Schedl 1951k: 134, 148, 1979c: 14.

angustior (Eggers) 1927b: 300 (*Cryphalus*). Holotype ♂; Snd-Sumatra: Buru; USNM, Washington.

Distribution: Indonesia (Sumatra).

References: (tx) Anderson, W. H. & Anderson 1971: 4; Eggers 1927b: 300; Schedl 1950g: 36–37, 1979c: 21.

asper (Broun) 1881: 742 (*Tomicus*). Syntypes, sex?; Wellington, New Zealand; not located.

Distribution: New Zealand.

References: (ds) Hagedorn 1910d: 48; Kleine 1913b: 127. (tx) Broun 1881: 742; Schedl 1938f: 48, 1942c: 163.

bakeri (Eggers) 1927c: 77 (*Stephanoderes*). Syntypes, sex?; Philippinen: Catanduanes, Prov. Albay, Virac; Eggers Collection [not mentioned by either Anderson & Anderson 1971 or Schedl 1979c].

Distribution: Philippine Islands.

References: (ds) Schedl 1966b: 25. (tx) Eggers 1927c: 77; Nobuchi 1983: 300; Schedl 1951i: 50, 1962h: 66.

basihirtus (Beeson) 1929: 227 (*Cryphalus*). Syntypes 2, sex?; [Samoa] Upolu: Malololelei; BMNH, London.

Distribution: Samoan Islands.

Hosts: *Myristica* sp.

Notes: (3) Schedl 1951k: 145 (redescription, male described).

References: (cn) Kalshoven 1954: 8. (ds) Beaver 1976b: 534; Beeson 1938b: 288. (tx) Beeson 1929: 227; Schedl 1951k: 134, 145.

bidentatus Browne 1980b: 383. Holotype, sex?; Wasag (Philippine) to Kinunra (Japan), imported; BMNH, London.

Distribution: Philippine Islands.

References: (tx) Browne 1980b: 383; Nobuchi 1983: 301.

brevior Schedl 1943b: 40. Lectotype, sex?; St. Theodoro, Mindoro [Island], Philippinen; USNM, Washington, designated by Anderson & Anderson 1971: 8.

Distribution: Philippine Islands (Mindoro).

References: (ds) Schedl 1966b: 26. (tx) Anderson, W. H. & Anderson 1974: 7; Nobuchi 1983: 301; Schedl 1943b: 40, 1979c: 46.

caplandicus Schedl 1965h: 115. Holotype, sex?; Cape Province, Grootvadersbosch; TMP, Pretoria.

Distribution: Africa (South Africa).

Hosts: *Podocarpus latifolius*, *Virgilia oroboides*.

References: (tx) Schedl 1965h: 115, 1979c: 52.

corpulentus Schedl 1942d: 22. Lectotype, sex?; Java, Mount Gede, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 65.

Distribution: Asia (Malaya), Indonesia (Java).

Hosts: *Artocarpus scortechini*, *Hibiscus tiliaceus*.

References: (cn) Mathur & Singh 1960b: 51. (hb) Browne 1961c: 67. (ds) Browne 1961c: 67; Mathur & Singh 1960b: 51. (tx) Schedl 1942a: 176, 1942d: 22, 1953c: 289, 1979c: 65.

cylindripennis Schedl 1959a: 483. Holotype ♀; Ceylon: Millawitiya Estate; Schedl Collection in NHMW, Wien.

Distribution: Asia (Sri Lanka).

References: (ds) Schedl 1959a: 483. (tx) Schedl 1959a: 483, 1979c: 74.

cylindrus (Browne) 1950b: 647 (*Dacryphalus*). Holotype ♀; Malaya: Selangor, Kepong; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Palaquium stellatum*.

References: (tx) Browne 1950b: 647, 1961c: 68; Schedl 1959a: 483.

densepilosus Schedl 1942d: 21. Lectotype ♀; Java, Mount Gede, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 77.

Distribution: Indonesia (Java).

References: (tx) Schedl 1942d: 21, 1948g: 27, 1979c: 77.

- discrepans** Schedl 1965c: 58. Holotype, sex?; Madagascar, Montagne d'Ambre, 1000 m; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
References: (tx) Schedl 1965c: 58, 1977a: 59, 1979c: 82.
- dubiosus** Schedl 1970h: 177. Holotype, sex?; Alexandria (C.P.); NICP, Pretoria.
Figures: Schedl 1970h: 181.
Distribution: Africa (South Africa).
Hosts: *Ekebergia capensis*, *Eleocharis* sp.
References: (cn) Anonymous 1970c: 13. (ds) Anonymous 1970c: 13. (tx) Schedl 1970h: 177, 181, 1979c: 98.
- fici** Browne 1986a: 90. Holotype, sex?; Solomon Islands (Bougainville Island); Dios to Nagoya (Japan), imported; BMNH, London.
Distribution: Solomon Islands (Bougainville).
Hosts: *Ficus* sp. log.
References: (tx) Browne 1986a: 90.
- formosanus** Schedl 1952c: 62. Holotype, sex?; Formosa, Riran; Schedl Collection in NHMW, Wien.
Distribution: Asia (Taiwan).
References: (ds) Nobuchi 1967: 24. (tx) Schedl 1952c: 62, 1979c: 98.
- glabratus** Schedl 1959a: 484. Holotype ♀; Ceylon; Millawitiya Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka).
References: (ds) Schedl 1959a: 484. (tx) Schedl 1959a: 484, 1979c: 105.
- granulatus** Schedl 1942a: 176. Lectotype, sex?; Malaya, Selangor; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 110.
Distribution: Asia (Malaya).
Hosts: *Myristica furfuracea*.
References: (ds) Browne 1961c: 67. (tx) Schedl 1942a: 176, 1979c: 110.
- imitans** Schedl 1951k: 148. Lectotype ♂; Upolu, Tapatapao, 800 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 121.
Distribution: Samoan Islands.
References: (ds) Beaver 1976b: 534. (tx) Schedl 1951k: 134, 148, 1979c: 121.
- interponens** Schedl 1953b: 126. Lectotype ♀; Saigon, Jardin Botanique; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka/Vietnam).
References: (tx) Schedl 1953b: 126, 1979c: 127.
- kalambanganus** Schedl 1943b: 39. Lectotype, sex?; Kalambangan, Mindanao, Philippines; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 131.
Distribution: Philippine Islands (Mindanao).
References: (ds) Schedl 1966b: 26. (tx) Nobuchi 1983: 301; Schedl 1943b: 39, 1979c: 131.
- laevis** Browne 1980a: 374. Holotype, sex?; Batania (New Guinea) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: *Calophyllum* sp.
References: (tx) Browne 1980a: 374.
- laticollis** Browne 1974a: 67. Holotype, sex?; Fiji; Viti Levu, Colo-i-Suva; BMNH, London.
Distribution: Fiji Islands.
Notes: (3) Schedl 1980d: 118 (treated as a synonym of *tutuilaensis*). Beaver 1991: 91 (a good species).
References: (ec) Roberts 1976: 380. (hb) Roberts 1976: 380. (ds) Browne 1974a: 64; Roberts 1976: 380. (tx) Beaver 1991: 91; Browne 1974a: 64, 67; Schedl 1980d: 118.
- longipilis** Browne 1981a: 128. Holotype, sex?; Bacau (Moluccas) to Omaezaki (Japan), imported. Distribution: Indonesia (Moluccas), New Britain Island.
References: (tx) Browne 1981a: 128.
- malayensis** Schedl 1942a: 176. Lectotype ♀; Malaya, Pahang, Kemasul; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 146.
Distribution: Asia (Malaya), Indonesia (Borneo).
Hosts: Burceraceae sp., *Canarium littorale*, *Dacryodes rostrata*.
References: (hb) Browne 1961c: 67. (ds) Browne 1961c: 67. (tx) Schedl 1942a: 176, 1953c: 289, 1979c: 146.
- mangiferae** (Stebbing) 1914: 542 (*Cryphalus*). Lectotype ♀; Eastern Dm, northern India; BMNH, London, designated by Wood 1982b: 871, name conserved by plenary powers.
Distribution: Africa (Ghana/ Guinea/ Kenya/ Ivory Coast/ Mauritius/ Nigeria/ Tanzania/ Uganda/ Zaire), Antilles Islands (Barbados/ Guadeloupe), Asia (Burma/ Nicobar Islands in India/ Malaya/ Sri Lanka/ Thailand), Australia (Queensland), Cook Islands (Rarotonga), Hawaii, Indonesia (Java), Madagascar, Micronesia (Caroline Islands, Palau), North America (Costa Rica/ Honduras/ Veracruz in Mexico/ Panama/ Florida in USA), Reunion Island, Samoan Islands, South America (Brazil/ Venezuela).
Hosts: *Mangifera indica*, *M. odorata*.
Notes: (1) Tubbs 1986: 245 (priority given *mangiferae* under plenary powers). (3) Schedl 1952: 62 (*africanus* Eggers, nomen nudum, synonymy). *Cryphalus robustus* Eichhoff may also be a synonym.
References: (ay) Browne 1962g: 77; Costa Lima 1929: 109–110; Gardner 1934: 11. (cn) Costa Lima 1956; Ebeling 1950, 1959; Pyenson 1935: 29–30. (ec) Silvas, Gayeo, & Castro 1959: 13. (hb) Beeson 1929: 226, 1938: 255, 1940: 198, 1941: 374; Browne 1968: 350–351; Carvalho 1938; Castro da Silveira 1960: 54; Costa Lima 1956, 1957: 276–290; Ebeling 1950, 1951: 680; Kalsho-

ven 1954: 8; Medeiros & Rosetto 1966: 30–38; Paulian 1951: 27; Rao 1953: 52; Roberts 1969: 125; Tucker 1952: 347; Wichmann 1954: 503–510; Wood, S. L. 1953: 77. **(ds)** Beaver 1989a: 2, 1990a: 280; Beaver & Maddison 1990: 1371; Beeson 1940, 1961: 288; Blackwelder 1947; Blackwelder & Blackwelder 1948; Bright 1985c: 174; Browne 1964: 188, 1965: 188, 1970: 540; Costa Lima 1936: 355, 1956; Dumbleton 1954: 28, 134; Ebeling 1950, 1959; Hopkins 1927: 23–32; Kalshoven 1958: 164; Kleine 1934a: 148; Mathur & Singh 1961a: 45, 51, 1961b: 51; Nunberg 1958: 479, 507; Paulian 1951: 27; Schedl 1959a: 483, 1960a: 76, 79, 1961k: 542–544, 1966f: 84, 1967d: 2, 1967e: 211; Swezey 1949: 445; Tucker 1952: 347. **(tx)** Beeson 1929: 226, 1940; Bright 1985c: 174; Costa Lima 1929, 1956; Eggers 1928c: 85, 1931c: 185; Hopkins 1927: 28; Kalshoven 1958: 164, 169; Schedl 1942a: 176, 1942d: 2, 1948g: 25–29, 1950g: 894, 1951j: 19, 1951k: 134, 1952c: 62, 1958k: 153, 1961k: 543–544, 1970e: 81, 1977b: 59; Stebbing 1914: 542–543; Swezey 1949: 445; Thirumala Rao 1953: 52; Wood, S. L. 1953: 77, 1954a: 962–966, 999–1000, 1087, 1960a: 24–27.

inops Eichhoff 1872a: 131 (*Cryphalus*). Holotype ♀; Guadeloupe; IRSNB, Brussels, name suppressed by plenary powers. Synonymy: Wood 1982b: 871.

References: **(ds)** Beeson 1929: 225; Blackwelder 1947: 778; Fleutiaux & Salle 1890: 457; Hagedorn 1910d: 43; Kleine 1913b: 118, 1914b: 379. **(tx)** Eichhoff 1872a: 131; Hagedorn 1910a: 86; Tubbs 1986: 245; Wood, S. L. 1982b: 871.

griseus Blackburn 1885: 194 (*Hypothenemus*). Holotype ♀; plains of Honolulu, Oahu, Hawaiian Islands; BMNH, London, name suppressed by plenary powers. Synonymy: Wood 1982b: 871.

References: **(ds)** Beeson 1938b: 289; Hagedorn 1910d: 43; Kleine 1913b: 124, 1914b: 302; Swezey 1949: 445. **(tx)** Blackburn 1885: 194; Hagedorn 1910a: 86; Hopkins 1915b: 24; Perkins 1900: 180; Sharp 1892: 194; Swezey 1949: 445–446; Wood, S. L. 1982b: 871.

mangiferae Eggers 1928c: 85. Lectotype ♂; Brazil; USNM, Washington, designated by Anderson & Anderson 1971: 19. Synonymy: Eggers 1931: 185.

References: **(cn)** Anonymous 1961k, 1962g, 1962y; Estrada & Atkinson 1988: 209; Kalshoven 1954: 8; Stebbing 1914: 542; Thirumala Rao 1953: 52. **(cc)** Abrahao & Wegmuller 1974; Castro 1960a; Equihua & Atkinson 1986: 630; Stebbing 1914: 542. **(hb)** Abrahao & Wegmuller 1969; Beaver 1976b: 534; Beaver & Browne 1978: 588; Beeson 1929; Castro 1960a; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Kalshoven 1958b: 164;

Schedl 1961k: 543, 1977a: 59; Stebbing 1914: 542; Wood, S. L. 1954a: 999, 1982b: 871. **(ds)** Anonymous 1961k, 1962g, 1962y; Atkinson & Equihua 1985c: 356; Beaver 1976b: 534; Beaver & Browne 1978: 588; Beeson 1929, 1938b, 1961: 288; Blackwelder 1947; Bright 1985c: 174; Browne 1970: 540; Castro 1960a; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Estrada & Atkinson 1988: 209; Hopkins 1927: 28; Mathew 1982, 1987: 188; Nunberg 1958a: 480; Pedrosa-Macedo & Schonherr 1985: 45; Schedl 1959a: 483, 1960d: 76, 1961k: 543, 1977a: 59; Thirumala Rao 1953: 52; Wood, S. L. 1977a: 70, 1982b: 871. **(tx)** Anderson, W. H. & Anderson 1971: 19; Beeson 1929; Bright 1985c: 174; Castro 1960a; Eggers 1928c: 85, 1931c: 185; Nunberg 1958a: 480; Pedrosa-Macedo & Schonherr 1985: 45; Schedl 1959a: 483, 1961k: 543, 1977a: 59; Stebbing 1914: 542; Wood, S. L. 1954a: 999, 1982b: 871, 1984a.

subcylindricus Schedl 1942d: 16 (*Cryphalus*). Lectotype ♀; Java, Semarang, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 240. Synonymy: Schedl 1958k: 153.

References: **(hb)** Browne 1961c: 70. **(ds)** Browne 1961c: 70; Schedl 1959a: 482. **(tx)** Schedl 1942a: 169–171, 1942d: 16–17, 1953c: 289, 1958k: 153, 1959a: 482, 1979c: 240.

mimicus Schedl 1942d: 17 (*Cryphalus*). Lectotype, sex?; Java; Pasoeroean; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 153. Synonymy: Kalshoven 1958: 164.

References: **(ds)** Schedl 1959a: 481. **(tx)** Kalshoven 1958: 164; Schedl 1942c: 171, 1942d: 17, 1959a: 481, 1979c: 153.

opacus Schedl 1942d: 20. Lectotype, sex?; Java, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 179. Synonymy: Kalshoven 1958: 164, Wood 1989: 173.

References: **(ds)** Maiti & Saha 1986: 157. **(tx)** Kalshoven 1958: 164; Schedl 1942a: 176, 1942d: 20, 1943b: 39, 1979c: 179; Wood, S. L. 1989: 173.

mindoroensis Schedl 1943b: 39. Lectotype ♀; Philippinen, San Theodoro [Mindoro]; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 153.

Distribution: Philippine Islands.

References: **(ds)** Schedl 1966b: 26. **(tx)** Anderson, W. H. & Anderson 1971: 20; Nobuchi 1983: 301; Schedl 1942a: 176, 1943b: 39, 1979c: 153.

minor Schedl 1943b: 40. Holotype, sex?; Port Banga, Mindanao; Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands (Mindanao).

References: **(hb)** Browne 1961c: 71. **(ds)** Browne

- 1961c: 71; Schedl 1966b: 26. (**tx**) Nobuchi 1983: 301; Schedl 1943b: 40, 1979c: 154.
- minutus** Browne 1980d: 495. Holotype, sex[?]; Surabaya (Java) to Osaka (Japan), imported; BMNH, London.
Distribution: Indonesia (Java).
Hosts: Inggis log.
References: (**tx**) Browne 1980d: 495.
- montanus** Schedl 1974d: 460. Holotype ♀; Mt. Kaindi, 2250 m, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Dryadodaphne* sp.
References: (**tx**) Schedl 1974d: 460, 1979c: 158.
- moorei** Schedl 1964i: 247. Holotype ♂; Somersby, N.S.W., Australia; Australian Museum, Sydney.
Distribution: Australia (New South Wales).
Hosts: *Hakea sericea*.
References: (**hb**) Moore 1964: 300. (**tx**) Schedl 1964i: 247, 1979c: 159.
- nigrosetosus** Schedl 1948g: 27. Lectotype ♀; Queensland, Kalpowar; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 168.
Distribution: Australia (Queensland).
Hosts: *Capparis uobilis*.
References: (**ds**) Brimblecombe 1953: 19. (**tx**) Schedl 1948g: 27, 1951k: 148, 1979c: 168.
- nitidicollis** Schedl 1975g: 219. Holotype, sex[?]; New Guinea, Bismarck-Geb.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (**tx**) Schedl 1975g: 219, 1979c: 168.
- obesus** (Hopkins) 1915b: 42 (*Dacryphalus*). Holotype ♀; Pagbilao; USNM, Washington.
Distribution: Philippine Islands (Luzon).
Hosts: Margadilao log.
References: (**ds**) Kleine 1934a: 148; Schedl 1966b: 26. (**tx**) Hopkins 1915b: 42; Nobuchi 1983: 301.
- obscurus** Hopkins 1915b: 41. Holotype ♀; Calapan, P.I.; USNM, Washington.
Distribution: Philippine Islands (Mindoro).
Hosts: *Cacao silvestre*.
Notes: (3) Schedl 1963j: 480 (*setulosum* Eggers, nomen nudum[?], cited as a possible synonym of *obscurus* Hopkins).
References: (**ds**) Kleine 1934a: 148; Schedl 1966b: 26. (**tx**) Hopkins 1915b: 41, 42; Nobuchi 1983: 301; Schedl 1963j: 480.
- oralicollis** Schedl 1942c: 177. Holotype, sex[?]; Australien, Tambourin Mountain; Schedl Collection in NHMW, Wien.
Distribution: Australia.
References: (**tx**) Schedl 1942c: 177, 1979c: 181.
- perminimus** (Schedl) 1942d: 13 (*Cryphalus*). Lectotype, sex[?]; Java, Semarang teak forest; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 90.
Figures: Numberg 1961b: 626.
- Distribution: Asia (Ryukyu Islands/ Malaya), Fiji Islands, Indonesia (Java), New Britain Island, New Guinea, Philippine Islands.
Hosts: *Artocarpus* spp., *Ficus* sp., *Pterocarpus indicus*.
Notes: (1) Schedl 1958k: 153 (to *Hypocryphalus*).
References: (**ec**) Roberts 1976: 381. (**hb**) Browne 1961c: 68; Kalshoven 1958b: 164; Roberts 1976: 381. (**ds**) Browne 1961c: 68, 1966: 244, 1974a: 64, 1980a: 371; Roberts 1976: 381; Schedl 1966b: 26. (**tx**) Nobuchi 1983: 301; Schedl 1942a: 169–170, 1942c: 170, 1942d: 13, 1951k: 145, 1958b: 99, 1958k: 153, 1962p: 697, 1979c: 190.
- constrictus** Schedl 1942d: 22. Lectotype, sex[?]; Java, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 64. Synonymy: Schedl 1958k: 153.
References: (**tx**) Schedl 1942c: 174, 1942d: 22, 1950f: 49, 1958k: 153, 1979c: 64.
- constrictus** Schedl 1942c: 174 (*Cryphalus*). Holotype, sex[?]; New-Guinea; Schedl Collection in NHMW, Wien. Synonymy: Kalshoven 1958: 164.
References: (**tx**) Kalshoven 1958: 164; Schedl 1942c: 174, 1942d: 22, 1950f: 49, 1954a: 139, 1958k: 153, 1979c: 64.
- froggatti** Numberg 1961b: 611. Holotype, sex[?]; New Britain, Rabaul; BMNH, London. Synonymy: Schedl 1962n: 697.
References: (**tx**) Numberg 1961b: 611, 626; Schedl 1962n: 697.
- pilifer** Schedl 1979g: 98. Holotype, sex[?]; New Guinea, N. E., Wau, Mt. Kaindi, Nami Creek; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (**tx**) Schedl 1979g: 98.
- piliger** Schedl 1975g: 219. Holotype, sex[?]; New Guinea; Wau; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (**tx**) Schedl 1975g: 219, 1979c: 194.
- polynesiae** Schedl 1979g: 105. Holotype, sex[?]; Samoa; Schedl Collection in NHMW, Wien.
Distribution: Samoan Islands.
References: (**tx**) Schedl 1979g: 105.
- reflexus** Browne 1980b: 383. Holotype, sex[?]; Ulanona (New Britain) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Britain Island.
References: (**tx**) Browne 1980b: 383.
- rotundus** Hopkins 1915b: 41. Holotype, sex[?]; Pagbilao, P.I.; USNM, Washington.
Distribution: Philippine Islands (Luzon).
Hosts: *Dipterocarpus grandiflorus*.
References: (**ds**) Beeson 1938b: 289; Kleine 1934a: 148; Schedl 1966b: 27. (**tx**) Hopkins 1915b: 41; Nobuchi 1983: 301; Wood, S. L. 1954a: 999.

- sandakanensis** (Schedl) 1937e: 545 (*Cryphalus*).
 Lectotype ♀; N. Borneo, Bettotan, near Sandakan; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 219.
 Distribution: Indonesia (Borneo)
 Hosts: *Garcinia* sp.
 References: (tx) Schedl 1937e: 545, 1939f: 37, 1942a: 169–170, 1979c: 219.
maculatus Browne 1961a: 303. Holotype, sex?; Sarawak; Bako National Park; BMNH, London. Synonymy: Wood 1989: 173.
 References: (hb) Browne 1961a: 303. (tx) Browne 1961a: 303; Schedl 1942a: 169; Wood, S. L. 1989: 173.
- spathulatus** Schedl 1935f: 49. Lectotype, sex?; Cairns district; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 233.
 Distribution: Australia (Queensland).
 References: (hb) Ymus & Hua 1980: 225. (ds) Brimblecombe 1953: 19; Schedl 1966b: 27; Ymus & Hua 1980: 225. (tx) Nobuchi 1983: 301; Schedl 1935f: 49–50, 1979c: 233.
- striatus** Hopkins 1915b: 41. Holotype ♀; Calapan, P.I.; USNM, Washington.
 Distribution: Asia (Malaya), Philippine Islands (Mindoro).
 Hosts: *Canarium* sp., *Parinarium* sp.
 References: (ds) Beeson 1961: 287–288; Kleine 1934a: 145. (tx) Hopkins 1915b: 41–42; Schedl 1939c: 325, 1942a: 176.
- sumatranus** (Schedl) 1939f: 38 (*Dacryphalus*).
 Lectotype, sex?; Solok, Sumatra; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 246.
 Distribution: Indonesia (Sumatra).
 Hosts: *Myristica fragrans*.
 References: (cn) Kalshoven 1954: 5. (hb) Kalshoven 1955b: 164. (ds) Browne 1980d: 492. (tx) Schedl 1939f: 38, 1979c: 246.
- tongaensis** Schedl 1979g: 104. Holotype, sex?; Tonga, Tongatapu; Schedl Collection in NHMW, Wien.
 Distribution: Tongan Islands.
 References: (tx) Schedl 1979g: 104.
- triangularis** Schedl 1975f: 351. Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District.
 Distribution: New Guinea.
 References: (tx) Schedl 1975f: 351, 1979c: 255.
- tutuilaensis** Schedl 1951k: 147. Lectotype ♂; Samoa: Tutuila, Fagatogo, 800 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 259.
 Distribution: Samoan Islands.
 Hosts: *Calophyllum* sp.
 References: (ds) Beaver 1976b: 534. (tx) Schedl 1951k: 134, 147, 1979c: 259.

Genus *Cryphalus* Erichson

- CRYPHALUS** ERICHSON 1836: 61. Type-species: *Bostrichus asperatus* Gyllenhal, subsequent designation by Thomson 1859: 147, lectotype designated for type-species by Wood 1972f: 41.
- Pseudocryphalus* Ferrari 1868b: 252. Type-species: *Pseudocryphalus siducyanus* Ferrari, monobasic. Synonymy: Wood 1986a: 91.
 References: (tx) Eggers 1931c: 155; Ferrari 1868b: 252; Wood, S. L. 1986a: 91.
- Taenioglyptes* Bedel 1885b: 398. Type-species: *Bostrichus abietis* Ratzeburg = *Bostrichus asperatus* Gyllenhal, original designation. Synonymy: Wood 1972f: 40.
 References: (tx) Balachowsky 1949a: 205; Bedel 1885b: 398; Bright & Stark 1973: 67; Wood, S. L. 1954a: 960, 1001, 1972f: 40.
- Cryptarthrum* Blandford 1896b: 200. Type-species: *Cryptarthrum walkeri* Blandford, monobasic. Synonymy: Schedl 1963i: 62.
 References: (tx) Blandford 1896b: 200; Hopkins 1915b: 8; Schedl 1963i: 62.
- Allarthrum* Hagedorn 1912c: 355. Type-species: *Allarthrum kolbei* Hagedorn, monobasic. Synonymy: Wood 1986: 91.
 References: (tx) Hagedorn 1912c: 355; Wood, S. L. 1986: 91.
- Ericryphalus* Hopkins 1915b: 8, 38. Type-species: *Ericryphalus hawaii* Hopkins = *Hypothenemus sylvicola* Perkins, original designation. Synonymy: Schedl 1962r: 92.
 References: (tx) Beeson 1938: 288; Hopkins 1915b: 8, 38; Schedl 1939i: 353, 1942d: 14, 1957d: 1–162, 1958k: 142, 1961k: 455, 1962r: 92, 1963i: 62; Wood, S. L. 1960a: 22.
- Piperius* Hopkins 1915b: 8, 39. Type-species: *Piperius pinii* Hopkins = *Hypothenemus sylvicola* Perkins, original designation. Synonymy: Schedl 1962r: 92.
 References: (tx) Hopkins 1915b: 8, 39; Schedl 1962r: 92, 1963i: 62.
- Erucryphalus* Murayama 1958: 934. Type-species: *Erucryphalus hirosimensis* Murayama, original designation. Synonymy: Schedl 1963i: 62.
 References: (tx) Murayama 1958: 934; Schedl 1962p: 203, 1963i: 62.
- Acryphalus* Tsai & Li 1963: 604, 622. Type-species: *Cryphalus lipingensis* Tsai & Li, subsequent designation by Wood 1984b: 224. Synonymy: Wood 1984b: 224.
 References: (tx) Tsai & Li 1963: 604, 622; Wood, S. L. 1984b: 224.
- Jugocryphalus* Tsai & Li 1963: 602, 622. Type-species: *Cryphalus piceus* Eggers, subsequent designation by Wood 1984b: 224. Synonymy: Wood 1984b: 224.
 References: (tx) Tsai & Li 1963: 602, 605, 622; Wood, S. L. 1984b: 224.
- Keys: Wood 1954a: 1002, 1960: 23, 1982b: 867 for

North America and Micronesia; Balachowsky 1949a: 206, Reitter 1913a: 66, Stark 1952: 254 for Europe and parts of Asia.

References: (ay) Fisher 1937: 118; Fuchs 1912: 14–33; Hymons 1921: 103; Nusslin 1910: 289–298, 1911, 1912; Nobuchi 1969a: 55, 57; Tsai & Li 1963: 597. (cn) Beaver 1987a; Beeson 1916: 1–5. (cc) Nickle 1976b. (hb) Barbey 1901: 69; Bargmann 1906: 315; Beaver 1987a: 17; Beeson 1941: 372; Blackman 1950; Bright & Stark 1973: 66; Brimblecombe 1953: 15; Browne 1961c: 68; Escherich 1923: 446–476; Fextand 1950: 9; Judeich & Nitsche 1895: 448, 451; Mayne 1917: 1–80; Munro 1946: 5–60; Nusslin 1907: 613; Schedl 1958d: 188, 1961k: 457, 1981b: 74; Tragardh 1930c: 469–480, 1931: 58; Wood, S. L. 1982b: 867, 1986a: 91. (ds) Beaver 1987a: 17; Bright & Stark 1973: 68; Browne 1961c: 68; Gaubil 1849: 126; Hagedorn 1910d: 40; Niisima 1908: 89; Schedl 1961k: 455, 1966b: 19, 1981b: 74; Scheerpeltz & Winkler 1930: 257; Tsai & Li 1963: 597; Wichmann 1927: 356–362; Wood, S. L. 1982b: 867, 1986a: 91. (tx) Arnett 1960: 1043, 1968: 1043; Balachowsky 1949a: 200, 205; Beal & Massey 1945: 59, 119; Bedel 1888b: 396–397; Beeson 1935b: 257; Berger 1916: 227–228; Bertolini 1872: 201; Blandford 1894b: 201, 1904: 225, 234; Blatchley & Leng 1916: 605–606; Bright 1976d: 112; Bright & Stark 1973: 66; Browne 1965: 188; Chamberlin 1918: 12–13, 1939: 316–327, 1958: 130–132; Chatterjee 1917: 1–4; Choo 1983: 70; Choo, Woo, & Nobuchi 1988b; Dodge 1938: 48; Duffy 1953: 6; Eggers 1943c: 68; Eichhoff 1864: 34–35, 45–46, 1868e: 391, 1872a: 132, 134, 1878a: 384, 1878b: 34–457, 1881a: 45, 172, 1883: 107, 109; Endrodi 1957b: 417; Erichson 1836: 61; Fauvel 1887: 276; Ferrari 1867a: 10–18, 1867b: 113; Gardner 1934: 4, 11; Gozis 1886: 278; Hagedorn 1910a: 84–85; Hopkins 1914: 119, 130–135, 1915b: 8, 39–40, 1915c: 180–226; Inouye & Nobuchi 1957: 46–48; Jacquelin du Val & Fairmaire 1868: 104–105; Kalina 1970: 125; Karaman 1972: 118; Krivolutsкая 1958: 107, 140; Kurenzov 1941: 159–162, 230–231; Lacordaire 1866: 373; LeConte 1865: 151–152, 1876: 349, 362; LeConte & Horn 1883: 518; Lindemann 1876: 148–168, 345; Lovendal 1889: 17–18; Luigioni 1929: 996; Murayama 1930: 14, 1958: 935; Niisima 1909: 141; Numborg 1954: 48–52; Nusslin 1898: 276, 280; Peyerimhoff 1935: 194, 196; Pfeffer 1989a: 81; Prell 1930: 638; Ratzeburg 1837: 139, 163; Redtenbacher 1845: 95, 151, 162; Reitter 1889: 94, 1894: 69–75, 1906: 11, 1913a: 65–66, 1916: 287–288; Saalas 1914: 73, 83; Schedl 1934f: 638, 1938i: 24, 1957d: 1–162, 1960h: 105, 1960i: 105, 1961k: 455, 458, 1962r: 91–93, 1963h: 266, 1963i: 62, 1967f: 154–155, 1968e: 261–270, 1969i: 129, 1977b: 38, 1981b: 74; Schimitschek 1937: 50–52; Seidlitz 1891a: 561–563, 1891b: 151, 607–608; Sharp 1879: 101;

Sokanovskii 1954: 16; Spessivtsev 1922: 471, 1931: 88; Stark 1936e: 142–144, 1952: 15–280; Stebbing 1914: 528; Stresemann et al. 1989: 351; Swaine 1909: 91–92, 1918a: 87–89; Thomson 1859: 147, 1865: 354; Trappen 1935: 142; Tsai & Li 1963: 597, 600; Wollaston 1960: 364–365; Wood, S. L. 1954a: 980, 1960a: 23, 1961a: 45, 1967a: 121–122, 1972f: 41, 1974: 171, 1982b: 867–871, 1986a: 91; Yin & Huang 1981: 561; Yin, Huang & Li 1984: 97; Zivojinovic 1948: 65–73.

abbreviatus Schedl 1943b: 35. Lectotype ♀; Catbalogan, Ins. Samar, Philippines; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 8.

Distribution: Asia (Okinawa in Japan/ Ryuku Islands), Philippine Islands (Samar).

References: (cn) Egorov 1958: 1492; Ekici 1971. (hb) Ekici 1971. (ds) Krivolutsкая 1965: 234; Nobuchi 1985c: 13; Schedl 1971c: 363. (tx) Nobuchi 1959c: 22; Schedl 1943b: 35, 1951i: 50, 57, 1979c: 8.

alni Krivolutsкая 1958: 145. Holotype, sex?; Sakhalin Island: Gornoravsk region, Valley of Kuznetsovsk and Ribatsko rivers; Zoological Institute, Academy of Science, Vladivostok.

Distribution: Asia (E USSR).

Hosts: *Alnus* sp.

References: (hb) Krivolutsкая 1973: 136. (ds) Krivolutsкая 1964a: 234, 1973: 136. (tx) Krivolutsкая 1958: 141–145.

aquilinus (Nobuchi) 1975: 52 (*Taenioglyptes*). Holotype ♀; Nigorikawa, Hokkaido, Japan; Nobuchi Collection. Ibaraki.

Figures: Nobuchi 1975: pl. 2, fig. 14.

Distribution: Asia (Japan).

Hosts: *Carpinus cordata*.

References: (ds) Nobuchi 1985c: 13. (tx) Nobuchi 1975: 52.

araucariae Schedl 1969b (1970): 214. Holotype, sex?; New Guinea, Bulolo, Morobe Dist.; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Araucaria cunninghamii*.

References: (tx) Schedl 1969b (1970): 214, 1979c: 24.

armatus Schedl 1974d: 459. Holotype, sex?; Wan, Morobe District; BPBM, Honolulu.

Distribution: New Guinea.

References: (tx) Schedl 1974d: 459, 1979c: 26.

artestriatus (Browne) 1970: 553 (*Taenioglyptes*). Holotype ♀; Uganda: Zika; BMNH, London.

Distribution: Africa (Uganda).

References: (tx) Browne 1970: 553.

artocarpus (Schedl) 1939e: 342 (*Ericryphalus*). Syntypes, sex?; Malaya, Selangor: Kepong; BMNH, London, and Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

Hosts: *Artocarpus kunstleri*.

Notes: (3) Beeson 1941 (1961: 257) (*artocarpus*, nomen nudum).

References: (**hb**) Yunus & Hua 1950: 225. (**ds**) Beeson 1941 (1961: 257); Browne 1961c: 65, 1954a: 150, 1954b: 257; Yunus & Hua 1950: 225. (**tx**) Schedl 1939c: 342, 1979c: 27.

asperatus (**Cyllenhal**) 1813: 363 (*Bostrichus*). Lectotype, sex?; Sweden: University of Uppsala Museum, designated by Wood 1972f: 41.

Figures: Balachowsky 1949a: 210, 216.

Distribution: Africa (Algeria/ Morocco), Asia (Honslu in Japan/ Turkey), Europe (Austria/ Belgium/ Bulgaria/ Czechoslovakia/ Denmark/ England/ Finland/ France/ Germany/ Greece/ Hungary/ Italy/ Luxemburg/ Netherlands/ Norway/ Poland/ Romania/ Spain/ Sweden/ Switzerland/ W USSR/ Yugoslavia).

Hosts: *Abies pectinata*, *Picea excelsa*, *Larix* sp., *Pseudotsuga douglasii*.

Notes: (1) Due to an ancient error in identification, that has been perpetuated, numerous citations of this name actually refer to *Trypophloeus binodulus*, a species that breeds in *Populus* spp.

References: (**ay**) Heeger 1866: 533–537. (**bv**) Chararas 1950b: 563; Grune 1979: 101. (**cn**) Chararas 1972; Escherich 1917: 97–115; Karpinski 1933: 52–56; Wachtl 1901: 381. (**hb**) Eichhoff 1881a: 45, 179, 1882a: 241; Heeger 1866: 533–537; Henschel 1855b, 1895a: 166; Ratzeburg 1839: 197; Schwerdtfeger 1981: 188; Wachtl 1901: 381, 1876a: 458. (**ds**) Allen, A. A. 1977; Barthe 1896; Brakman 1966b: 206; Buresh & Lazarov 1956; Ericson & Sandin 1893; Gemminger & Harold 1872: 2653; Gozis 1875: 50; Grill 1895: 309; Cronzelle 1905; Grune 1979: 101; Cyllenhal 1827: 624; Hagedorn 1910d: 41; Henschel 1895a: 166; Heyden, Reitter, & Weise 1853: 182; Kaltenback 1874: 655; Kamp 1979; Kangas 1964d, 1966d; Kestercanek 1881a: 12; Klefbeck & Sjoberg 1960: 230; Kraatz 1869: 59; Lacordaire 1866: 379; Leclercq 1971; Licht 1987: 278; Negru 1966b: 401; Nuorteva 1971: 71; Ortzen 1886: 279; Pfeffer 1959a: 84; Postner 1974: 428; Ratzeburg 1839: 197; Redtenbacher 1855: 832, 1874: 376; Reitter 1869b: 154; Sahlberg 1900: 105; Schaum 1859: 96, 1862: 101; Schedl 1964f; Schwerdtfeger 1981: 188; Seidlitz 1872: 393, 1891a: 562, 1891b: 607; Siebke 1875: 283; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 439; Stierlin & Cantard 1871: 292; Sturm 1843: 230; Thomson 1859: 146, 1865: 359, 1868: 219; Wachtl 1876a: 458; Westhoff 1882: 238; Winter, T. G. 1983: 28. (**tx**) Bach 1854; Balachowsky 1949a: 215; Beeson 1938b: 287; Bertolini 1872; Eichhoff 1864b: 35, 1866: 276, 1876a: 378, 1878b: 125, 1881a: 45, 179, 1883a: 109, 132; Erichson 1836: 62; Fauvel 1859; Ferrari 1867a: 11, 15, 80, 1867b: 114; Gebien 1907: 197; Grune 1979: 100–101; Cyllenhal 1813: 365, 1827: 624;

Hagedorn 1904: 230, 1910a: 85; Henschel 1895a: 166; Hopkins 1914: 118, 1915b: 39, 1915c: 221, pls. 9–10, fig. 1; Kalina 1969; Lacordaire 1866: 379; Letzner 1844: 68, 1891: 375; Licht 1987: 278; Negru 1966b: 401; Postner 1974: 428; Ratzeburg 1839: 197; Redtenbacher 1849a: 791, 1849b: 26, 1858: 832, 1874: 376; Sahlberg 1836: 153; Schedl 1934f: 1639; Seidlitz 1872: 393, 1891a: 562, 1891b: 607; Stierlin 1898: 439; Thomson 1859: 146, 1865: 359, 1868: 219; Wood, S. L. 1954a: 988, 1972f: 42.

abietis Ratzeburg 1837: 163 (*Bostrichus*). Syn-types, sex?; Ober-Schleien, Ostprussen, dem Thuringer Walde und dem Harze; MNB, Berlin. Synonymy: Wood 1972f: 41.

References: (**ay**) Chararas 1973c; Escherich 1923b: 481, 597; Feytaud 1950a; Ritchie 1919: 171–199; Scherlb 1971. (**bv**) Byers, Anderbrant, & Lofqvist 1989; Chararas 1973c; Chararas, Desveaux, & Kogane-Charles 1978a; Chararas et al. 1982: 1094; Grune 1979: 107; Kevdina 1897: 108; Magema, Gaspar, Severin 1982; Namann-Etienne 1978a; Nuorteva 1956c: 94; Nuorteva & Nuorteva 1965; Tragardh 1930b: 108; Winter, K. 1980. (**cn**) Anonymous 1978i; Barbey 1906a; Borcea 1924: 221–260; Bruneau 1950; Byers, Anderbrant, & Lofqvist 1989; Chararas 1961b: 69, 1961c: 92, 1978; Chorbadzhiev 1929; Eckstein 1926: 578; Escherich 1923b: 481, 597, 1936; Esterberg 1959; Feytaud 1946, 1950a; Forbes 1910; Gabler 1955; Georgescu et al. 1957: 357, 460; Hanson 1937; Hartig 1861: 330; Hess & Beck 1914: 265, 1927: 321; Inouye & Yamaguchi 1955a: 235; Joly 1976; Judeich & Nitsche 1895: 526; Kailidis & Georgiivits 1972; Kalandra 1944; Kamp 1950; Keller 1903b: 49; Kholodkovskii 1912: 278, 307; Koch 1913: 100; Komarek et al. 1931: 1–256; Kozikowsky 1929: 254; Kunstler 1864: 783; Lekander 1955b: 17; Lozovoi 1961: 91–113; Lozovoi & Tropin 1965, 1967; MacDougall 1917: 1–38; Marcu 1926c: 63; Mokrzejcki 1925: 1–7; Muller 1912: 185; Nestertschnk 1930: 176; Nosek 1951: 107, 1952b: 98; Nuorteva & Nuorteva 1965; Nusslin 1913: 227; Pfeffer 1928a: 467, 1950c: 3; Pierce, W. D. 1917: 69; Rlumibler 1922: 299, 1927: 312; Ritchie 1916: 301–302, 1919: 171; Ruhl 1958: 287; Schimitschek 1930d: 136, 1937c: 52, 1938b: 114, 1955a: 32, 36, 1955c: 82, 1964a; Schwerdtfeger 1944a: 180, 1957a: 186; Stewart 1915: 70; Tragardh 1919: 237–248, 1938a: 12; Wachtl 1901: 381; Weber, H. 1926: 578; Wichmann 1927b: 374; Zolk 1937: 147–172. (**ce**) Apfelbeck 1916b; Balazy & Michalski 1960, 1964a; Barbey 1906a, 1927; Capek 1957; Chararas 1959g; Chararas, Desveaux, & Kogane-Charles 1978a; Eidmann 1974b; Galoux 1947c, 1948a; Gillanders 1906;

- Heqvist 1963: 152; Hierholzer 1954b: 357; Inouye & Yamaguchi 1955a: 235; Kielczewski 1976; Kielczewski & Wisniewski 1978, 1983; Kleine 1905c: 186, 1909a: 46, 1944: 74; Kraemer 1950b: 379; Lozovoi & Tropin 1965, 1967; Lundberg 1984; Michalski & Ratajczak 1989; Naumann-Etienne 1978a; Nosek 1951: 107, 1952b: 98; Nuorteva 1957b: 68; Nuorteva & Nuorteva 1968; Nusslin 1927: 300; Perris 1856a: 244; Pfeffer 1923a: 331, 1932a: 5, 1943b: 179, 1950c: 3, 1955b: 84, 1959: 2; Poinar 1975: 151; Ratzeburg 1869a: 79; Ruhm 1955c: 176, 1956b: 3; Saalas 1917a: 18, 1930: 118; Schimitschek 1955a: 32, 36, 1964a; Schwerdtfeger 1944a: 180, 1957a: 186; Sedlaczek 1908: 46, 1935a: 163; Sitowski 1930: 4; Thompson, W. R. 1943: 37; Wichmann 1959: 412; Winter, K. 1980; Wisniewski 1979b; Zuur 1955b. (hb) Altum 1881c: 321; Apfelbeck 1916b, 1917; Barbey 1901: 23, 71, 1925: 164; Beffa 1949, 1961; Boas 1923: 360; Borcea 1924; Budge 1949; Cecconi 1924; Chararas 1961b: 69, 1961c: 92, 1962c: 251; Charvat 1950; Chorbadzhievo 1929; Dallimore & Munro 1922; Eckstein 1897, 1926: 578; Eichhoff 1881a: 45, 176, 1882a: 241; Eidmann 1974b; Escherich 1923b: 451, 597; Everts 1903: 755; Feytaud 1946, 1950a; Fuchs 1904a, 1906c: 209, 1907: 48; Gabler 1955; Galoux 1947a, 1948a; Gillanders 1906, 1908; Gornostaev 1916: 314; Györfi 1957; Hagedorn 1903a; Hartig 1861: 330; Heinemann 1909; Henschel 1876a: 34, 240, 1895a: 166; Hess & Beck 1914: 265, 1927: 321; Hufnagl 1887: 512; Joly 1976; Judeich & Nitsche 1895: 526; Kahlich 1865b; Karpinski 1933b: 28; Karpinski & Strawinski 1948: 155; Karsch 1883: 142; Kholodkovskii 1912: 275, 307; Kraemer 1950b: 379; Lengerken 1939: 62, 1954: 83; Lozovoi & Tropin 1965, 1967; Lumardoni & Leonardi 1889: 458; MacDougall 1917: 127; Magma, Gaspar, & Severin 1982; Masutti 1964; Munro 1926: 60; Naumann-Etienne 1978a; Nordlinger 1856: 28; Numberg 1929: 102, 1947c: 106, 1948: 1-30; Nuorteva 1956c: 94; Nusslin 1898: 280, 1906b: 14, 1913: 227, 1927: 300; Palm 1954a: 185; Perris 1856a: 244; Pfeffer 1941b: 1, 1941c: 5, 1989a; Postner 1974: 430; Ratzeburg 1837: 135, 163, 1839: 165, 198; Rhumbler 1922: 299, 1927: 312; Ritchie 1919: 171; Rupertsberger 1879: 231, 1880: 229; Saalas 1913a: 69, 83; Schimitschek 1955a: 32, 36; Schwerdtfeger 1944a: 180, 1957a: 156, 1981: 192; Sedlaczek 1935a: 163; Spessivtsev 1913a: 91; Stark 1926a: 335, 1952: 265; Thum 1885: 24; Tragardh 1929a: 313, 1930b: 108, 1930c: 474, 1931: 57, 1939b: 147, 198; Tredl 1908a: 30; Tschorbadjiev 1929: 166; Wachtl 1876a: 458, 1901: 381; Weber, H. 1926: 578; Wichmann 1927b: 374; Winter, K. 1980. (ds) Ammann & Knabl 1913, 1923; Andersch 1851; Anonymous 1978i; Arnoldi et al. 1955: 687; Arru, Covassi, & de Bellis 1966: 32; Balazy & Michalski 1960; Barthe 1896; Bau 1888; Bedel 1888b: 398; Beffa 1949; Bejer-Petersen & Jorum 1977: 23; Benick 1921; Benz 1985; Bielz 1887; Boas 1923: 360; Borcea 1924; Borchert 1951; Brakman 1966b: 206; Braucsik 1871; Bucking 1932; Buresh & Lazarov 1956; Butovitsch & Heqvist 1947; Cecconi 1897; Chapuis & Candezze 1853; Chararas 1962c: 251; Charvat 1950; Chorbadzhievo 1924d, 1929; Crotch 1863; Dallimore & Munro 1922: 189-193; Dejean 1821, 1825, 1837; Eder 1934; Eggers 1904; Eidmann 1974b; Endrodi 1958b; Ermsch 1953; Escherich 1923b: 451, 597, 1932b; Esterberg 1925, 1959; Everts 1922: 642; Favre 1890; Florov 1949: 85; Forster 1849: 439; Fowler 1891; Fuchs 1904a, 1905a, 1906c: 209, 1907: 48; Gaubil 1849: 126; Gornostaev 1917: 308-315; Gozis 1875: 80; Grill 1895: 309; Gronzelle 1905; Grune 1979: 107; Gussmann 1919; Hagedorn 1903a, 1910d: 40; Hansen, V. 1939, 1956, 1964: 462; Heinemann 1908a; Hellen 1947; Hennig 1954: 257; Henschel 1895a: 166; Heyden 1879: 140; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1951; Inouye & Nobuchi 1957: 49; Jannicky 1960a; Jazentkovsky 1912: 288; Joly 1960; Judeich & Nitsche 1895: 526; Kahlich 1865b; Kailidis & Georgivits 1972; Kaltenbach 1874: 685; Karpinski 1931: 27, 1933b: 28, 1948b: 230; Karpinski & Strawinski 1948: 155; Keler 1925b: 272; Kersten 1933: 74; Klefbeck & Sjöberg 1969: 230; Kleine 1912a: 262, 264, 267, 1913a: 35, 1913b: 118, 1934a: 142; Kloft & Hinks 1945: 218; Koltze 1901: 153; Kono & Tamamki 1939: 88, 91; Koschitsky 1900: 83; Kozikowsky 1921: 180; Kraatz 1869: 59; Kurir 1947c: 15; Lacordaire 1866: 379; Langhoffer 1915c: 157; Leclercq 1971; Leiler & Prutz 1977; Lekander 1955b: 17; Leng 1918: 211; Lentz 1857: 139; Lindemann 1884b: 264; Lokaj 1868: 64; Lomnicki 1913b: 148; Lucht 1987: 278; Lumardoni & Leonardi 1889: 458; Lundberg 1979: 31, 1981: 151; Lundblad 1950c: 115; Marcu 1926c: 63; Matthews & Fowler 1883: 42; Mequignon 1936: 48; Michalski 1957: 164; Munro 1921: 57, 1926: 4-77; Munster 1922a: 155, 1928: 289; Murayama 1948: 2, 1949a: 12, 1951c: 3, 1954b: 162; Nash, D. R. 1978; Negrin 1966b: 402, 1968a: 456; Nobuchi 1955c: 13; Numberg 1927a: 213, 1928b: 88, 105, 1954: 52; Nusslin 1898: 280; Orest 1926c: 63; Pacher 1865: 152; Palm 1959: 351; Perris 1876a: 254, 1877a: 414; Pfeffer 1924b: 471, 1931b: 74, 1947e: 2, 1950b: 75; Pierce, W. D. 1917: 69; Pittioni 1943: 175; Postner 1974: 430; Power 1865: 212; Rapp

- 1934: 727; Ratzeburg 1837: 135, 163, 1839: 165, 198; Redtenbacher 1855: 832; Reitter 1869b: 154, 1889a: 94, 1894a: 70, 1916: 285; Ronbal 1935b: 72, 1941: 263; Saalas 1913a: 69, 83, 1917a: 18, 1930: 118, 1931: 68; Sahlberg 1900: 105; Sainte-Claire & Mequignon 1938: 445; Schaschl 1854: 133; Schaufuss 1915: 1231; Schann 1859: 96, 1862: 101; Schedl 1959b: 100, 1961b: 184, 1967c: 70, 1980a: 14, 1981b: 75; Schilsky 1909: 185; Schimitschek 1935b: 114; Schiodte 1873: 101; Schwerdtfeger 1981: 192; Seidlitz 1872: 393, 1891a: 562, 1891b: 607; Sharp & Fowler 1893: 34; Stark 1926a: 335, 1926b: 103, 1926j: 126, 1936e: 143, 1952: 265; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 439; Stierlin & Cantard 1906: 205; Sturm 1843: 230; Swaine 1909: 93; Tragardh 1939b: 147, 198; Tredl 1907: 12; Tschorbadjiev 1929: 166; Verhoff 1891: 22; Wachtl 1876a: 458; Wessel 1877: 390; Westhoff 1882: 238; Wichmann 1927a: 67; Winter, T. G. 1983: 5; Yanovskii & Tegshzhargal 1955: 415. **(tx)** Allen, A. A. 1965b: 47; Bach 1854; Balachowsky 1949a: 207; Barbey 1901: 23, 71; Bedel 1888b: 395; Beffa 1949, 1961; Boas 1923: 360; Brancsik 1871; Chapuis & Candeze 1853; Chararas 1962c: 252; Charvat 1950; Chorbadzhievo 1924d; Crotch 1966; Csiki 1910; Dejean 1821, 1825; Doebner 1860; Duffy 1953; Eggers 1920: 241, 1921; Eichhoff 1864b: 35, 1866: 275–276, 1878b: 124, 1881a: 45, 176, 1883a: 109, 132; Endrodi 1957b; Escherich 1923b: 481, 592; Everts 1903: 755, 1922: 642; Ferrari 1911; Ferrari 1867a: 15, 79, 80, 1867b: 106; Fleischer 1927; Formanek 1907: 35; Gabler 1955; Gillanders 1908; Grune 1979: 106–107; Hagedorn 1910a: 85; Hansen, V. 1956, 1964: 462; Henry 1892: 14; Henschel 1876a: 34, 240, 1895a: 166; Hopkins 1915b: 40; Inouye & Nobuchi 1957a; Joly 1976; Jendeli & Nitsche 1895: 526; Kalina 1970: 125; Karpinski 1949: 129; Karpinski & Strawinski 1948: 155; Koch 1913: 100, 1928: 104, 1932: 146; Kuhn 1913: 1055; Lacordaire 1866: 379; Letzner 1844: 68, 1891: 375; Lindemann 1875c: 362; Lovendal 1889b: 51, 1898: 124; Lucht 1987: 278; Lunardoni & Leonardi 1889: 458; Murayama 1954b: 162; Negri 1966b: 402; Niisima 1910a: 8, 1929: 376–381; Nordlinger 1848: 244, 1856: 28; Numberg 1954: 52; Perris 1877a: 414; Pfeffer 1932b: 17, 1941b: 1, 1941c: 5, 1947c: 2, 1955a: 177; Portevin 1935: 324; Postner 1974: 430; Quaschnik 1953: 35; Ratzeburg 1837: 135, 163, 1839: 165, 198; Redtenbacher 1849a: 791, 1849b: 26, 1858: 832; Reitter 1889a: 94, 1894a: 70, 1913a: 67; Rhumbler 1922: 299, 1927: 231; Rupertsberger 1879: 231, 1880: 229; Saalas 1913a: 69, 83, 1914: 304–306, 1916: 110–116; Schedl 1934f: 1635, 1952f: 87, 1959: 100, 1980a: 14, 1981b: 75; Scherb 1971; Schimitschek 1937c: 52, 1955c: 82; Seidlitz 1872: 393, 1891a: 562, 1891b: 607; Simmel 1918: 288–291; Sokanovskii 1954: 16; Spessivtsev 1913a: 90–91, 1922a: 472, 1925a: 36, 173, 1931: 49; Stark 1936c: 143, 1952: 265; Stierlin 1898: 439; Swaine 1909: 93, 1918a: 89; Wood, S. L. 1972f: 41–42. **(ms)** Escherich 1932b; Heinemann 1908a; Kozikowsky 1929: 254; Sedlacek 1902a: 126; Wichmann 1959: 412.
- asperulus** Schedl 1948g: 26. Lectotype ♀; Queensland, Imbil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 28.
Distribution: Australia (Queensland).
Hosts: *Cryptocarya erythroxylon*, *Grevillea robusta*.
References: **(ds)** Brimblecombe 1953: 16. **(tx)** Schedl 1948g: 26, 1979c: 28.
- ater** Browne 1984d: 91. Holotype, sex?, New Guinea: Gumi; BMNH, London.
Distribution: New Guinea.
Hosts: *Garcinia* sp.
References: **(tx)** Browne 1984d: 91.
- babai** Murayama 1961b: 29. Holotype ♂; Noo, south part of Niigata prefecture, Japan; Murayama Collection in USNM, Washington.
Distribution: Asia (Honsu in Japan).
References: **(ds)** Nobuchi 1985c: 13. **(tx)** Murayama 1961b: 29.
- balanopselaphus** Eggers 1920: 121. Holotype, sex?, Amani, Ostafrika; Hamburg Museum, lost.
Distribution: Africa (Tanzania).
Hosts: *Rubus* sp.
References: **(tx)** Eggers 1920: 121–122; Sampson 1922a: 141; Schedl 1957d: 49, 1961k: 460.
- bellus** Schedl 1957d: 47. Holotype, sex?, Congo Belge; Yangambi; MRCB, Tervuren.
Distribution: Africa (Ivory Coast/ Zaïre).
Hosts: *Ficus thonningis*, *Guarea cedrata*, *Musanga cercropioides*.
References: **(hb)** Schedl 1961k: 460. **(ds)** Browne 1973a: 279; Schedl 1961k: 460. **(tx)** Schedl 1957d: 47, 1961j: 349, 1961k: 460, 1979c: 36.
- bicarinatus** (Nobuchi) 1975: 53 (*Taenioglyptus*). Holotype ♂; Arakawa, Okinawa; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 2, fig. 16.
Distribution: Asia (Okinawa in Japan).
References: **(ds)** Nobuchi 1985c: 13. **(tx)** Nobuchi 1975: 53–54.
- bicolor** (Browne) 1984f: 69 (*Cryphalomorphus*). Holotype, sex?, New Guinea: Morobe District, Wau, Kunai Creek, 1500 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Castanopsis* sp.
References: **(tx)** Beaver 1991b: 53; Browne 1984f: 69.

- birosimensis** (Murayama) 1958: 935 (*Ernocryphalus*). Holotype ♂; Isl. Biro, Kagoshima pref., Japan; USNM, Washington.
Distribution: Asia (Japan).
Hosts: *Pittosporum tobira*.
Notes: (1) Schedl 1963i: 62 (to *Cryphalus*). Nobuchi 1985c: 12 (treated in *Scolytogenes*).
References: (ds) Nobuchi 1985c: 12. (tx) Murayama 1958: 935; Schedl 1962p: 203, 1963i: 62.
- boettcheri** Schedl 1951i: 57. Syntypes, sex?; Philippinen. Mindoro, San Theodoro; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands (Mindoro).
References: (ds) Schedl 1966b: 19. (tx) Nobuchi 1983: 300; Schedl 1951i: 57, 1979c: 42.
- borneensis** Schedl 1943b: 38. Holotype, sex?; Borneo; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya), Indonesia (Borneo).
Hosts: *Bruguiera gymnorhiza*, *Elaeocarpus* sp.
References: (hb) Browne 1961c: 68–69. (ds) Browne 1961c: 68–69; Schedl 1962b: 186. (tx) Schedl 1943b: 38, 1958b: 99, 1979c: 43.
- brasiliensis** Schedl 1976a: 65. Holotype, sex?; Botafofo, Guanabara, Brasil; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
Notes: (3) This holotype is either mislabeled or it represents an introduction from another area of the world.
References: (tx) Schedl 1976a: 65.
- brevipilosus** Schedl 1942c: 173. Holotype, sex?; New-Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1942c: 173, 1979c: 46.
- brimblecombei** Schedl 1948g: 26. Lectotype ♀; Queensland, Emu Vale; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 47.
Distribution: Australia (Queensland).
Hosts: *Cryptocarya erythroxylon*.
References: (ds) Brimblecombe 1953: 17; Schedl 1972a: 143. (tx) Schedl 1948g: 26, 1950f: 51, 1979c: 47.
- brownei** Wood 1992b: 79. Holotype, sex?; Sarawak, Semengoh; BMNH, London, automatic.
Distribution: Indonesia (Sarawak in Borneo).
Hosts: *Artocarpus* sp.
References: (tx) Wood, S. L. 1992b: 79.
- artocarpus** Schedl 1958h: 498. Holotype, sex?; Sarawak, Semengoh; BMNH, London, preoccupied by Schedl 1939.
References: (tx) Schedl 1958h: 498, 1979c: 27; Wood, S. L. 1992b: 79.
- brunneus** Browne 1984d: 93. Holotype, sex?; New Guinea; Sawmill L. A.; BMNH, London.
Distribution: New Guinea.
Hosts: *Araucaria hunsteinii*.
References: (tx) Browne 1984d: 93.
- buloloensis** Browne 1984e: 67. Holotype, sex?; New Guinea; Morobe District, Bulolo, Snake Creek, 1000 m; BMNH, London.
Distribution: New Guinea.
References: (tx) Browne 1984c: 67.
- capucinicollis** Schedl 1950f: 47. Syntypes 2, sex?; Viti Levu; Navai Mill, near Nandarivatu, 2500 ft.; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.
Distribution: Fiji Islands.
References: (ds) Browne 1974a: 64; Roberts 1976: 377. (tx) Schedl 1950f: 47, 1979c: 52.
- capucinoides** Eggers 1939b: 4. Holotype, sex?; Nordost-Birma (Kambaiti, 7000 Fuss); NHR, Stockholm.
Distribution: Asia (Burma).
References: (tx) Eggers 1939b: 4–5.
- capucinomorphus** Schedl 1950f: 48. Syntypes ♂ ♀; Viti Levu; Tholo-i-Suva, 500 ft.; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.
Distribution: Fiji Islands.
References: (cc) Roberts 1976: 378. (hb) Roberts 1976: 378. (ds) Browne 1979a: 64; Roberts 1976: 378. (tx) Schedl 1950f: 38, 47–48, 1953f: 83, 1955b: 284, 1979c: 52.
- capucinus** Schedl 1938g: 425. Lectotype ♂; Luzon, Laguna Province, Mount Maquiling; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 52.
Distribution: Asia (Burma/ "China"), Philippine Islands.
Hosts: *Cassia javanica*.
Notes: (3) This is *parkeri* Beeson 1941: 591, nomen nudum.
References: (ds) Schedl 1938g: 425, 1966b: 20. (tx) Eggers 1939b: 4; Nobuchi 1983: 300; Schedl 1938g: 425, 1940d: 591, 1950f: 47, 1979c: 52.
- carpini** Berger 1916: 234. Lectotype ♀; Ussuri, USSR; IZL, Leningrad, designated by Michalski 1969a: 891.
Distribution: Asia (Japan/ Korea/ Siberia, Ussuri in E USSR).
Hosts: *Carpinus cordata*, *C. laxifolia*.
Notes: (3) Choo 1983: 70 (redescribed).
References: (cn) Kurenzov 1935c: 157. (cc) Kurenzov 1934a: 52. (hb) Kurenzov 1935a: 19, 30, 1948b: 114; Stark 1952: 261. (ds) Choo 1983: 70; Choo & Woo 1985: 164; Kleine 1934a: 142; Krivolitskaya 1983; Kurenzov 1934a: 52, 1935a: 19, 30, 1935c: 157, 1936b: 350; Nobuchi 1985c: 13; Stark 1936c: 143, 1952: 261. (tx) Berger 1916: 234; Choo 1983: 70; Eggers 1929b: 11, 1942c: 29; Kurenzov 1941a: 140–142, 1948b: 114; Michalski 1969a: 891, 1969b: 567; Nobuchi 1964: 133; Schedl 1934f: 1635, 1979c: 53; Stark 1936c: 143, 1952: 261.

- carpinivorus** Murayama 1930b: 14. Holotype, sex?; Korvo (Korea); Murayama Collection in USNM, Washington.
Distribution: Asia (Korea).
Hosts: *Carpinus laxiflora*.
References: (ds) Cho 1957; Choo & Woo 1955: 164; Kleine 1934a: 142; Ko 1969: 275; Murayama 1930b: 10, 1937b: 375. (tx) Murayama 1930b: 10, 14–16, 1937b: 375; Schedl 1934f: 1639.
- chamaecipariae** Niisima 1910a: 10. Syntypes, sex?; Tokio, Japan (Garten der Tokioter Universitat); Nobuchi Collection, Ibaraki.
Distribution: Asia (Honshu in Japan).
Hosts: *Chamaeciparus obtusa*.
Notes: (3) Nobuchi 1955c: 15 (treated in *Hypothenemus*).
References: (ds) Kleine 1934a: 143; Murayama 1954b: 162; Nobuchi 1955c: 15. (tx) Murayama 1954b: 162; Niisima 1910a: 10; Schedl 1934f: 1640.
- chulingensis** Tsai & Li 1963: 605, 612, 623. Holotype ♂; Szechuan, south of Yangtze River, China; Institute of Zoology, Academia Sinica.
Figures: Li, Dang, & Shi 1977: 192, Li & Zhou 1980: 192, Yin, Huang, & Li 1984: 108.
Distribution: Asia (Shanxi, Sichuan in China).
Hosts: *Pinus armandi*.
References: (hb) Li, Dang & Shi 1977: 72; Li & Zhou 1980: 72; Yang 1989c. (ds) Li, Dang, & Shi 1977: 72; Yin, Huang, & Li 1984: 107. (tx) Tsai & Li 1963: 605, 612, 623; Yin, Huang, & Li 1984: 107.
- ciliatipes** Blandford 1896f: 242. Syntypes ♂; Noumea a Kone (DeLauney), New Caledonia; BMNH, London.
Distribution: New Caledonia Island.
References: (ds) Beeson 1938b: 287; Hagedorn 1910d: 41; Kleine 1913b: 118, 1914b: 300. (tx) Blandford 1896f: 242; Hagedorn 1910a: 85; Schedl 1953c: 296, 1957d: 12, 1979c: 57.
- cinereotestaceus** (Motschulsky) 1866: 403 (*Hypoborus*). Holotype ♀; Des Montagnes de Nura-Ellia, Ceylon; IZM, Moscow.
Distribution: Asia (Sri Lanka).
Notes: (3) Wood 1969c: 118 (near *paganus* or *pallidus*).
References: (ds) Gemminger & Harold 1872: 2685; Hagedorn 1910d: 29; Kleine 1912b: 186, 1914b: 273. (tx) Hagedorn 1910a: 61; Motschulsky 1866: 403; Wood, S. L. 1969c: 118.
- compactus** Lea 1910: 139. Syntypes, sex?; Queensland: Port Denison; SAM, Adelaide?.
Distribution: Australia (Queensland).
References: (ds) Brimblecombe 1953: 17. (tx) Lea 1910: 139; Schedl 1938f: 46–47.
- confusus** Eggers 1927b: 395. Holotype ♀; Sud Sumatra; Eggers Collection, in NHMW, Wien.
Distribution: Indonesia (Sumatra), New Guinea.
- Notes: (3) Schedl 1969b: 215 (described male).
References: (tx) Eggers 1927b: 395; Schedl 1943b: 35, 1969b: 215, 1979c: 62.
- coryli** Stark 1936: 144. Syntypes, sex?; publication not seen; IZL, Leningrad.
Distribution: Asia (E USSR).
Notes: (3) Doubtful species, could = *mandshuricus* Eggers.
References: (hb) Kurenzov 1948b: 113; Stark 1952: 269. (ds) Krivolutskaya 1983; Stark 1936e: 144, 1952: 269. (tx) Eggers 1942c: 28; Kurenzov 1948b: 113; Michalski 1969a: 591, 1969b: 567; Schedl 1979c: 66; Sokanovskii 1954: 17; Stark 1936e: 144, 1950: 229, 1952: 269.
- cryptomeriae** Niisima 1908a: 91. Syntypes 4, sex?; Kumamoto in der Provinz Higo, Japan; Nobuchi Collection, Ibaraki.
Distribution: Asia (Japan).
Hosts: *Cryptomeria japonica*.
References: (ds) Kleine 1913b: 118, 1914b: 257, 1934a: 142; Murayama 1954b: 199; Nobuchi 1955c: 13. (tx) Murayama 1954b: 199; Niisima 1908a: 91; Schedl 1934f: 1639.
- cylindricus** Browne 1980b: 384. Holotype, sex?; Pagai Island (Mentawai Islands, Indonesia) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Mentawai).
References: (ds) Browne 1980b: 384, 1981b: 597.
- cylindrus** Browne 1984d: 92. Holotype, sex?; New Guinea: Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Auracaria hunsteinii*.
References: (tx) Browne 1984d: 92.
- densepilosus** Schedl 1943b: 36. Holotype, sex?; Mt. Makiling, Laguna, Luzon, Philippines; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands (Luzon).
Hosts: *Ficus nimialhasae*.
References: (ds) Schedl 1966b: 20. (tx) Nobuchi 1983: 300; Schedl 1943b: 36–37, 1953c: 295, 1979c: 77.
- dipterocarpi** Wood 1989: 179. Holotype ♀; Margherita Factory, Lakhimpur, Assam, India; FRI, Dehra Dun.
Distribution: Asia (Birma/ Assam in India).
Hosts: *Dipterocarpus alatus* D. pilosus.
Notes: (1) This is *dipterocarpi* Beeson 1941 (1961: 288), nomen nudum.
References: (ds) Beeson 1961: 288. (tx) Schedl 1940d: 591, 1942a: 169–170, 1943b: 39; Wood, S. L. 1989: 179.
- dissimilis** (Nobuchi) 1975: 57 (*Tacnioglyptes*). Holotype ♀; Mt. Fuji, Yamagashi pref., Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 2, fig. 21.
Distribution: Asia (Japan).
Hosts: *Abies homolepis*.

References: **(ds)** Nobuchi 1955c: 13. **(tx)** Nobuchi 1975: 57–58.

diversicolor Browne 1984d: 90. Holotype, sex?; New Guinea: Sawmill L. A.; BMNH, London.

Distribution: New Guinea.

Hosts: *Arancaria lunsteinii*.

References: **(tx)** Browne 1984d: 90.

dorsalis (Motschulsky) 1866: 403 (*Hypoborus*).
Syntypes 1 ♂, 2 ♀; India: IZM, Moscow.

Distribution: Asia (Burma/ Hainan Island in China/ Andaman Islands, Assam in India/ Vietnam), Indonesia (Java, Sumatra), Philippines (Luzon, Mindanao).

Hosts: *Artocarpus lakoocha*, *Ficus glomerata*, *F. rumphii*.

References: **(ds)** Gemminger & Harold 1872: 2654; Hagedorn 1910d: 29; Kleine 1912b: 186, 1914b: 273. **(tx)** Hagedorn 1910a: 61; Motschulsky 1866: 403; Wood, S. L. 1969c: 119.

nebulosus Motschulsky 1866: 403 (*Hypoborus*).

Syntypes 2, sex?; India: IZM, Moscow. Synonymy: Wood 1969c: 119.

References: **(ds)** Gemminger & Harold 1872: 2654; Hagedorn 1910d: 29; Kleine 1912b: 186, 1914b: 268. **(tx)** Hagedorn 1910a: 61; Motschulsky 1866: 403; Wood, S. L. 1969c: 119.

sericeus Motschulsky 1866: 402 (*Hylesinus*).

Holotype ♀; Ceylon: IZM, Moscow; preoccupied by Mannerheim 1843. Synonymy: Wood 1969c: 118.

References: **(ds)** Kleine 1912b: 171, 1914b: 273. **(tx)** Gemminger & Harold 1872: 2675; Hagedorn 1910a: 49; Motschulsky 1866: 402; Wood, S. L. 1969c: 118.

indicus Eichhoff 1878b: 489. Holotype ♂; Hindostan (Birma); Hamburg Museum, lost. Synonymy: Wood 1969c: 118.

References: **(cn)** Roonwal 1954: 89. **(hb)** Beaver & Browne 1975: 555; Browne 1935a, 1961c: 69; Kalshoven 1958b: 164; Yunn & Hua 1980: 228. **(ds)** Beaver & Browne 1975: 288, 1978: 555; Beeson 1961: 287; Browne 1935a, 1961c: 69; Hagedorn 1910d: 43; Kleine 1913b: 118, 1914b: 276, 278, 1934a: 142; Roonwal 1954: 89; Schedl 1938g: 424, 1962b: 186, 1966b: 21, 1971c: 376; Yunn & Hua 1980: 228. **(tx)** Eggers 1925: 152, 1927b: 396; Eichhoff 1878a: 384, 1878b: 489; Hagedorn 1910a: 87; Nobuchi 1983: 300; Schedl 1938g: 424, 1942a: 169, 1942d: 2, 22, 1951i: 50, 1965g: 22; Stebbing 1906: 403, 1914: 538; Wood, S. L. 1969c: 118.

dubiosus Schedl 1963i: 66. Holotype, sex?; Saigon; Schedl Collection in NHMW, Wien.

Distribution: Asia (Vietnam).

References: **(tx)** Schedl 1963i: 66, 1979c: 84.

duplosquamosus Schedl 1942d: 15. Lectotype, sex?; Java, Britenzorg; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java).

Hosts: *Canarium commune*.

References: **(ds)** Kalshoven 1958: 164. **(tx)** Schedl 1942d: 15, 1979c: 86.

elaboratus Schedl 1950f: 51. Syntypes 2, sex?; Vaua Mbadavi: Loma Loma, 200–500 ft., Myana, 200 ft.; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.

Distribution: Fiji Islands.

References: **(ds)** Browne 1974a: 64. **(tx)** Schedl 1950f: 51, 1979c: 88.

elongatus (Nobuchi) 1975: 42 (*Ericryphalus*).
Holotype, sex ?; Mt. Takao, Tokyo, Japan; Nobuchi, Ibaraki. Possible homonym.

Distribution: Asia (Japan).

Hosts: *Cornus controversa*.

Notes: (3) The position of this species is doubtful; it was named in an invalid genus and was said to be near *Ernioporicus caucasicus* (SLW).

References: **(tx)** Nobuchi 1975: 42, 1985c: 11.

elongatus Schedl 1962r: 105. Holotype, sex?; Fiji Inseln; Schedl Collection in NHMW, Wien.

Figures: Nobuchi 1975: pl. 1.

Distribution: Fiji Islands.

References: **(ds)** Browne 1974a: 64. **(tx)** Nobuchi 1975: pl. 1; Schedl 1962r: 105, 1979c: 89.

erraticum Schedl 1979g: 96. Holotype, sex?; Papua, Bulolo, Morobe District, Upper Manki L. A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: **(tx)** Schedl 1979g: 96.

exiguus Blandford 1894d: 82. Holotype, sex?; Fukushima (Japan); BMNH, London.

Figures: Yin, Huang, & Li 1984: 112.

Distribution: Asia (Anhui, Guizhou, Hebei, Jiangsu, Shandong, Sichuan, Zhejiang in China/ Japan/ Korea/ Kuril Islands/ Taiwan).

Hosts: *Broussonetia kazinoki*, *Celtis sinensis*, *Diospyros kaki*, *Exodia rutacarpa*, *Ficus* spp., *Morus alba*, *M. bombycis*, *Salix* sp.

Notes: (3) Choo 1983: 71 (redescribed).

References: **(cn)** Anonymous 1980g; Clausen 1931; Shiraki 1952. **(hb)** Krivolitskaya 1965a: 233, 1973: 134. **(ds)** Anonymous 1980g; Blandford 1894c; Cho 1957; Choo 1983: 71; Choo & Woo 1985: 164; Clausen 1931; Hagedorn 1910d: 42; Kleine 1913b: 118, 1914b: 256, 1934a: 142; Ko 1969: 275; Krivolitskaya 1965a: 233, 1973: 134; Ku 1964; Murayama 1930b: 10, 1933b: 2, 1936a: 126, 1937b: 375, 1949c: 100, 1950b: 1292, 1952a: 22, 1953c: 151, 1954b: 162; Shiraki 1952; Tsai & Li 1963: 603, 606; Yin, Huang, & Li 1984: 112. **(tx)** Blandford 1894d: 82; Choo 1983: 71; Murayama 1930b: 10, 14, 30, 1933: 14, 1936a: 126, 1937b: 375, 1950: 1292, 1952a: 22, 1954b: 162; Niiisima 1905a: 89–90, 1909: 140, 142, 1910a: 8–9; Nobuchi 1959c: 23; Schedl 1934f: 1639; Yin, Huang, & Li 1984: 112.

- pilosus* Sasaki 1899: 235. Syntypes?: Japan, publication not seen. Synonymy: Niisima 1909: 144. References: (tx) Niisima 1909: 146; Sasaki 1899: 235.
- felis* Wood 1859: 180.** Holotype ♀; Mussoorie, Uttar Pradesh, India; FRI, Dehra Dun.
Distribution: Asia (Uttar Pradesh in India).
Hosts: *Quercus* sp., *Vitis* sp.
Notes: (1) This is *felis* Beeson, nomen nudum.
References: (tx) Schedl 1940d: 591, 1943b: 37; Wood, S. L. 1959: 180.
- ficivorus* Murayama 1958: 933.** Syntypes 4, sex?; Muroto, Kochi pref., Shikoku, Japan; Murayama Collection in USNM, Washington.
Distribution: Asia (Shikoku in Japan).
Hosts: *Ficus wightiana*.
References: (ds) Nobuchi 1985c: 13. (tx) Murayama 1958: 933.
- formosanus* Schedl 1942a: 175.** Holotype, sex?; Formosa; Schedl Collection in NHMW, Wien.
Distribution: Asia (Taiwan).
References: (ds) Nobuchi 1967: 24. (tx) Murayama 1934c: 288; Schedl 1942a: 174–175, 1943b: 35, 1979c: 95.
- fugax* Schedl 1973e: 87.** Holotype, sex?; Rambuso, Sudest Island, 0–100 m. Milne Bay District; AMNH, New York.
Distribution: New Guinea.
References: (tx) Schedl 1973e: 87, 1979c: 101.
- fuliginosus* Blandford 1895a: 319.** Holotype, sex?; Ceylon, Bogawantalawa; BMNH, London.
Distribution: Asia (Sri Lanka).
Hosts: Liana.
References: (ds) Hagedorn 1910d: 43; Kleine 1913b: 118, 1914b: 273; Schedl 1959a: 481. (tx) Blandford 1895a: 319; Hagedorn 1910a: 86; Schedl 1959a: 481.
- fulmineus* Wood 1859: 180.** Holotype, sex?; Tharal, Carlwal, Uttar Pradesh, India; FRI, Dehra Dun.
Distribution: Asia (Sichuan in China/ Uttar Pradesh in India).
Hosts: *Ahus nitida*.
Notes: (1) This is *dumincus* Beeson 1941 (1961: 286), nomen nudum.
References: (ds) Roonwal 1954: 59. (tx) Schedl 1940d: 591; Wood, S. L. 1959: 180.
- fulvus* Niisima 1905a: 92.** Syntypes, sex?; Yatsuo-Berg in der Provinz Ohmi, Japan; Nobuchi Collection, Ibaraki.
Figures: Nakane et al. 1963: 382, Nobuchi 1966d: pl. 3.
Distribution: Asia (Liaoning, "Manchuria" in China/ Japan/ Korea/ Ryukyu Islands).
Hosts: *Pinus hiansiana*, *P. densiflora*, *P. koraiensis*, *P. rigida*, *P. tabulaeformis* var. *mukdensis*, *P. thumbergii*.
Notes: (3) Choo 1983: 72 (redescribed).
References: (ay) Sasakawa & Sasakawa 1981; Sasakawa & Yoshiyasu 1983. (bv) Nagasawa, Asano, & Makita 1968: 153–158, 1969: 80–85, 1970: 22–26; Nobuchi 1969b, 1969c; Sasakawa & Katayama 1975; Sasakawa & Kawaguchi 1987; Sasakawa & Negishi 1973; Sasakawa, Ohta, & Negishi 1976; Sasakawa & Sasakawa 1981, 1984; Sumimoto, Kondo, & Kaniyama 1974; Sumimoto et al. 1975. Yasunaga, Oshina, & Kuwatsuka 1962: 197–200; Yoshikawa et al. 1986. (cn) Anonymous 1980g; Godha et al. 1964; Inouye 1949a: 13, 110, 1949b; Ko 1984; Ko & Morimoto 1985; Nagasawa & Asano 1975; Nagasawa, Asano, & Fushimi 1968a, 1968b; Oda, Kato, & Nobuchi 1964; Sakai 1966: 120–129; Sakai et al. 1964: 61–68; Shiraki 1952; Yoshida & Fukuma 1972; Yoshida et al. 1974. (cc) Asano, Nagasawa, & Fushimi 1968; Heqvist 1967: 65; Kamijo 1981; Kishi 1969, 1970, 1972; Koyama 1963; Sasakawa & Negishi 1973; Sasakawa, Ohta, & Negishi 1976; Yasumatsu & Watanabe 1965: 68; Yoshikawa 1987a, 1987b; Yoshikawa et al. 1986. (hb) Asano, Nagasawa, & Fushimi 1968: 46–61; Nagasawa, Asano, & Makita 1969; Nagasawa et al. 1968; Sasakawa 1975; Sasakawa & Katayama 1975; Sasakawa & Negishi 1973; Sasakawa & Sasakawa 1984; Takahashi 1989: 403; Takahashi & Ito 1951: 159–161; Yoshikawa et al. 1986. (ds) Anonymous 1980g; Browne 1984b: 287; Cho 1955, 1957; Choo 1983: 72; Choo, Woo, & Nobuchi 1988a: 134; Choo, Woo, & Park 1983: 175; Inouye 1949b; Kleine 1913b: 118, 1914b: 257, 1934a: 142; Ko 1969: 275; Ku 1964; Murayama 1929b: 2, 1929d: 2, 1930a: 7, 1930b: 11, 1936b: 115, 1937b: 375, 1940a: 232, 1942a: 54, 1948: 2, 1949a: 11, 1949c: 100, 1950b: 1292, 1951a: 6, 1951c: 3, 1952a: 22, 1953a: 9, 1953c: 151, 1954b: 163, 1955: 98, 103; Nakane et al. 1963: 382; Nobuchi 1966d: 22; Nobuchi & Ono 1973: 181; Nohira & Ogawa 1986; Shiraki 1952; Tsai & Li 1963: 603; Yin, Huang, & Li 1984: 110. (tx) Choo 1983: 72; Eggers 1923b: 139; Hagedorn 1910a: 86; Murayama 1930b: 11, 15, 30, 1934c: 298, 1937b: 375, 1939: 144, 1940a: 232, 1950b: 1292, 1952a: 22, 1953a: 9, 1954b: 163, 1955: 92, 98, 103; Nakane et al. 1963: 382; Niisima 1908a: 92, 1910a: 8; Nobuchi 1959c: 25, 1966d: 22, pl. 3; Schedl 1934f: 1639, 1979c: 101; Sokanovskii 1954: 17; Tsai & Li 1963: 605; Yin, Huang, & Li 1984: 110.
- pini* Eggers 1921: 39. Lectotype ♂; Kiautschou (China); USNM, Washington, designated by Anderson & Anderson 1971: 139. Synonymy: Eggers 1923b: 139.
References: (ds) Schedl 1966b: 19. (tx) Anderson, W. H. & Anderson 1971: 25; Eggers 1921: 39, 1923b: 139; Nobuchi 1983: 300; Schedl 1934f: 1639.
- furukawai* Murayama 1934a: 59** (reprint p. 3). Holotype, sex?; Chochin, south Korea; USNM, Washington.

Distribution: Asia (Japan/ Korea).

Hosts: *Alnus sibirica*, *Ficus wightiana*, *Pinus maximowiczii*, *P. thunbergii*.

Notes: (3) Nobuchi 1985c: 16 (treated in *Hypothenemus* [not seen by us]).

References: (ds) Cho 1957; Choo 1983: 72; Choo & Woo 1985: 164; Ko 1969: 275; Murayama 1936a: 127, 1937b: 375, 1938: 126, 1952a: 19, 1953c: 151, 1954b: 163; Nobuchi 1966d: 23, 1985c: 16. (tx) Choo 1983: 72; Murayama 1934a: 59 (reprint p. 3), 1936a: 127, 1937b: 375, 1952a: 19, 1954b: 163; Nobuchi 1966d: 23.

garambaensis Nunberg 1965b: 25. Holotype, sex?; Repub. of the Congo: Garamba Nat. Park, near 2801; Inst. Parcs Nat. Congo Coll.

Figures: Nunberg 1965a: 35, 1965b: pl. 1–2.

Distribution: Africa (Congo).

References: (ds) Nunberg 1952: 23. (tx) Nunberg 1952: 23, 1965a: 31, 35, 1965b: 25, pl. 1–2.

giganteus Schedl 1950d: 22. Holotype, sex?; Natal; Schedl Collection in NHMW, Wien.

Distribution: Africa (Natal in South Africa).

References: (tx) Schedl 1950d: 22, 1961k: 462, 1979c: 104.

gigas Schedl 1975g: 218. Holotype ♀; New Guinea, Wau; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 351, 1975g: 218, 1979c: 104.

helopioides Schedl 1953c: 295. Lectotype, sex?; Malaya Peninsula, Kelantan, Bt. Mengkebang; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 116.

Distribution: Asia (Malaya).

References: (ds) Browne 1961c: 69; Schedl 1971c: 363. (tx) Schedl 1953c: 295, 1979c: 116.

hirsutus (Nobuchi) 1975: 52 (*Taenioglyptes*). Holotype ♀; Komano-yu, Nigata pref., Japan; Nobuchi Collection, Ibaraki.

Figures: Nobuchi 1975: pl. 2, fig. 15.

Distribution: Asia (Japan).

References: (ds) Nobuchi 1985c: 13. (tx) Nobuchi 1975: 52–53.

horridus Eichhoff 1878b: 488. Syntypes², sex?; Asiae India orientalis; Hamburg Museum, lost.

Distribution: Asia (Burma).

References: (hb) Schedl 1961k: 462. (ds) Hagedorn 1910d: 43; Kleine 1913b: 118, 1914b: 269; Schedl 1961k: 462. (tx) Eichhoff 1878a: 384, 1878b: 488; Hagedorn 1910a: 86; Schedl 1960h: 104, 1961k: 462.

infimus Schedl 1972e: 287. Holotype, sex?; Ghana, Ashanti Region, Kwadaso, 320 m, N 6 degrees 42, W 1 degree 39; NHMB, Budapest.

Distribution: Africa (Ghana).

References: (tx) Schedl 1972e: 287.

intermedius Ferrari 1867a: 79. Holotype, sex?; Germania borealis (Norddeutschland); NHMW, Wien?

Figures: Grune 1979: 104. Pfeffer 1989a: pl. 6.

Distribution: Europe (Austria/ Czechoslovakia/ Germany/ Hungary/ Italy/ Poland/ Romania).

References: (ay) Escherich 1923b: 481, 616; Nusslin 1911a: 129. (bv) Grune 1979: 105; Kinelski 1958: 73. (cn) Escherich 1923b: 481, 616; Koch 1913: 95; Nusslin 1913: 257; Pierce, W. D. 1917: 84; Rhumbler 1922: 299, 1927: 312; Schimitschek 1937c: 52, 1955c: 82, 1955a: 44, 91, 1961a: 154; Schwerdtfeger 1944a: 183, 1957a: 188; Wichmann 1927b: 378. (cc) Hirschmann & Wisniewski 1982, 1983; Karpinski 1932a: 95; Kielczewski, Moser, & Weisniewski 1983; Kielczewski & Wisniewski 1978, 1980a, 1983; Kleine 1908c: 186; Michalski & Ratajczak 1989; Nusslin 1927: 300; Pfeffer 1960: 345; Ruhm 1955c: 176; Schedl 1958d: 187; Schimitschek 1955a: 44, 91; Schwerdtfeger 1944a: 183, 1957a: 188; Wisniewski 1979a, 1979b. (hb) Dombrowsky 1887; Eichhoff 1881a: 45, 180; Escherich 1923b: 481, 616; Györfi 1957; Henschel 1885b, 1895a: 167; Karpinski & Strawinski 1948: 155; Lengerken 1939, 1954: 83; Nunberg 1929: 103; Nusslin 1913: 257, 1927: 300; Pfeffer 1941b: 18; Postner 1974: 430; Rhumbler 1922: 299, 1927: 312; Schimitschek 1955a: 44, 91; Schwerdtfeger 1944a: 183, 1957a: 188, 1981: 194; Stark 1952: 263; Tredl 1908a: 30–31; Wachtl 1876a: 458; Wichmann 1927b: 378. (ds) Calver 1884, 1893; Eggers 1904; Endrodi 1958b; Escherich 1923b: 481, 616, 1932b; Fuchs 1905a; Gemminger & Harold 1872: 2683; Grune 1979: 104–105; Hagedorn 1910d: 43; Henschel 1895a: 167; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1951; Karpinski 1932a: 95; Karpinski & Strawinski 1948: 155; Kiefer et al. 1942: 528; Kinelski 1958: 73; Kleine 1912a: 267, 1913a: 35, 1913b: 118, 1934a: 142; Kraatz 1869: 59; Kurir 1947c: 11; Lucht 1987: 278; Nunberg 1954: 52; Pfeffer 1926a: 11, 1931b: 75, 1950b: 75, 1960: 345, 1989a: 82; Pierce, W. D. 1917: 84; Pittioni 1943: 176; Postner 1974: 430; Prossen 1913: 83; Redtenbacher 1874: 376; Reitter 1894a: 70, 1916: 288; Rouhal 1941: 263; Schaufuss 1915: 1232; Schedl 1980a: 14, 1981b: 74; Schilsky 1909: 188; Schremmer 1959: 137; Schwerdtfeger 1981: 194; Seidlitz 1872: 393, 1891a: 562; Stark 1936a: 143, 1952: 263; Stein 1868: 114; Stein & Weise 1877: 164; Tredl 1907: 12; Wachtl 1876a: 458; Wichmann 1927a: 67–68. (tx) Dombrowsky 1887; Eggers 1929b: 10; Eichhoff 1868d: 49, 1878b: 126, 1881a: 45, 180, 1883a: 109, 132; Endrodi 1957a: 307, 1957b; Escherich 1923b: 481, 616; Ferrari 1867a: 79–80, 1867b: 114; Fleischer 1927; Grune 1979: 104–105; Hagedorn 1910a: 87; Henschel 1895a: 167; Karpinski & Strawinski 1948: 155; Koch 1913: 95, 1932: 146; Kuhn 1913:

- 1055; Lucht 1957: 278; Niisima 1909: 144; Numborg 1930, 1954: 52; Nusslin 1911a: 129; Pfeffer 1932b: 18, 1941b: 18, 1955a: 179, 1959a: pl. 6; Postner 1974: 430; Quaschik 1953: 35; Redtenbacher 1874: 376; Reitter 1894a: 70, 1913a: 67, 1916: 255; Rhumbler 1922: 299, 1927: 312; Schedl 1934f: 1639, 1980a: 14, 1981b: 74; Schimitschek 1937c: 52, 1955c: 82; Seidlitz 1872: 393, 1891a: 562; Stark 1936e: 143, 1952: 263. (**ms**) Escherich 1932b.
- jeholensis** Murayama 1939: 143. Syntypes, sex?; Shotoku, Manchukuo; Murayama Collection in USNM, Washington.
Figures: Nakane et al. 1963: 352, Nobuchi 1966d: pl. 3.
Distribution: Asia (Hebei, "Manchuria" in China/Japan/Korea).
Hosts: *Abies firma*, *Pinus densiflora*, *P. nigra*, *P. tabulaeformis*, *P. thunbergii*, *Smilax china*.
Notes: (3) Inouye & Nobuchi 1957: 50 (re-described).
References: (**ds**) Choo 1953: 73; Choo & Woo 1955: 164; Choo, Woo, & Nobuchi 1955a: 134; Murayama 1939: 140, 143, 1942a: 54, 1948: 2, 1949a: 12, 1951c: 3, 1952a: 20, 1953a: 10, 1953c: 151, 1954b: 163; Nakane et al. 1963: 352; Nobuchi 1966d: 23, 1955c: 14; Sawamoto 1940b: 141, 145; Tsai & Li 1963: 603, 605; Yin, Huang, & Li 1954: 110. (**tx**) Choo 1953: 73; Inouye & Nobuchi 1957a: 49; Murayama 1939: 140, 143, 1952a: 20, 1953a: 10, 1954b: 163; Nakane et al. 1963: 352; Nobuchi 1966d: 23, pl. 3; Sawamoto 1940b: 141–146; Yin, Huang, & Li 1954: 110.
- jezoensis** Inouye & Nobuchi 1957: 49. Syntypes 13, sex?; Soumkyo, Daisetsu-zan National Park, Hokkaido, Japan; Nobuchi Collection, Ibaraki.
Distribution: Asia (Hokkaido in Japan).
Hosts: *Picea jezoensis*.
References: (**ds**) Nobuchi 1955c: 14. (**tx**) Inouye & Nobuchi 1952a, 1957: 49.
- juglansi** Niisima 1913a: 3. Syntypes, sex?; Prov. Kushiro (Hokkaido); Nobuchi Collection, Ibaraki.
Distribution: Asia (Hokkaido, Honshu in Japan).
Hosts: *Juglans sieboldiana*.
References: (**cn**) Anonymous 1980g. (**ds**) Anonymous 1980g; Kleine 1934a: 142; Murayama 1954b: 199; Nobuchi 1955c: 14. (**tx**) Murayama 1954b: 199; Niisima 1913a: 3; Schedl 1934f: 1639, 1979c: 130.
- kagoshimensis** (Nobuchi) 1975: 56 (*Tacnioglyptes*). Holotype ♀; Satamisaki, Kagoshima, Japan; Nobuchi Collection, Ibaraki.
Figures: Nobuchi 1975: pl. 2, fig. 19.
Distribution: Asia (Japan).
References: (**ds**) Nobuchi 1955c: 14. (**tx**) Nobuchi 1975: 56.
- kesiyae** Browne 1975: 288 in Beaver & Browne. Holotype, sex?; Thailand; Chiang Mai, Doi Pui, 1600 m; BMNH, London.
Distribution: Asia (Thailand).
Hosts: *Pinus kesiya*.
References: (**tx**) Beaver & Browne 1975: 288; Browne 1980a: 370.
- kiruensis** Schedl 1957d: 48. Holotype, sex?; Congo Belge; Kivu, Hembe Bitale; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Ficus* cf. *capensis*.
References: (**ds**) Schedl 1961k: 464. (**tx**) Schedl 1957d: 45, 1961k: 464, 1979c: 132.
- kolbei** (Hagedorn) 1912c: 355 (*Allarthrum*). Syntypes, sex?; Peterhafen, Deutsch-Neuguinea; MNB, Berlin.
Distribution: New Guinea.
References: (**ds**) Kleine 1913b: 145, 1914b: 296; Krivolutsкая 1965a: 232, 1983. (**tx**) Hagedorn 1912c: 355; Hopkins 1914: 116; Michalski 1969b: 567; Schedl 1979c: 134.
- kurenzori** Stark 1936: 142, 150. Lectotype, sex?; Maykhe R. Valley, Far East Terr.; IZL, Leningrad, designated by Michalski 1969a: 591.
Distribution: Asia (Sakhalin Island, Siberia, Ussuri in E USSR).
Hosts: *Abies holophylla*, *A. nephrolepis*.
References: (**cn**) Kurenzov 1935c: 187. (**cc**) Kurenzov 1934a: 52. (**hb**) Kurenzov 1935a: 20, 1948b: 102, 1950d: 145; Stark 1952: 256. (**ds**) Kurenzov 1934a: 52, 1935a: 20, 1935c: 187, 1936b: 351, 1938a: 61, 1950: 145; Stark 1936e: 142, 150, 1952: 256. (**tx**) Eggers 1942c: 28–30; Krivolutsкая 1958: 141–143; Kurenzov 1941a: 142–143, 1948b: 102; Michalski 1969a: 591, 1969b: 567; Murayama 1961: 30; Nobuchi 1965: 130; Nobuchi & Takahashi 1965: 3; Pfeffer 1944a: 131; Schedl 1952k: 158; Sokanovskii 1954: 16; Stark 1936e: 142, 150, 1951: 229, 1952: 256. (**ms**) Pfeffer 1944a: 131.
- punctulatus** Eggers 1942c: 29. Holotype, sex?; Ussuri, USSR; Eggers Collection (not mentioned by Anderson & Anderson 1971 or Schedl 1979c). Synonymy: Pfeffer 1944a: 131.
References: (**tx**) Eggers 1942c: 29; Pfeffer 1944a: 131; Schedl 1952k: 158; Sokanovskii 1958: 38; Stark 1951: 229. (**ms**) Pfeffer 1944a: 131.
- ussuriensis** Eggers 1942c: 29. Holotype, sex?; Ussuri; Eggers Collection, in NHMW, Wien.
Synonymy: Schedl 1952k: 158.
References: (**hb**) Stark 1952: 269. (**ds**) Krivolutsкая 1983; Pfeffer 1944a: 131; Stark 1952: 269. (**tx**) Eggers 1942c: 29–30; Pfeffer 1944a: 131; Schedl 1952k: 158, 1979c: 262; Stark 1952: 269.
- kurilensis** Krivolutsкая 1968: 53. Holotype, sex?; Kuril Islands; Institute of Zoology, Academy of Science, Vladivostok.
Figures: Krivolutsкая 1968: 53.
Distribution: Asia (Kuril Islands).
Hosts: *Toxicodendron trichocarpum*.

- References: (tx) Krivolutskaya 1968: 53–56, 1973: 136.
- kyotoensis Nobuchi** 1966d: 53. Holotype ♀; Mizorogaike, Kyoto City, Japan; Nobuchi Collection, Ibaraki.
 Figures: Nobuchi 1966c: pls. 1–2.
 Distribution: Asia (Honsu in Japan).
 Hosts: *Alnus firma*.
 References: (ds) Nobuchi 1955c: 14, (tx) McNamara 1977: 196; Nobuchi 1966c: pls. 1–2, 1966d: 53.
- laevis Browne** 1984b: 288. Holotype, sex?; Malaysia: Kudat (Sabah) to Nagoya (Japan), imported; BMNH, London.
 Distribution: Asia (Malaya).
 Hosts: Keruing log.
 References: (tx) Browne 1984b: 288.
- laricis Niisima** 1909: 142. Syntypes, sex?; Nopporo in der Prov. Ishikari, Japan; Nobuchi Collection, Ibaraki.
 Figures: Nakane et al. 1963: 382, Nobuchi 1966d: pl. 3.
 Distribution: Asia (Japan/ Korea).
 Hosts: *Abies holophylla*, *A. sachaliensis*, *Larix leptolepis*, *Picea jezoensis*, *Pinus densiflora*, *P. koraiensis*, *P. silvestris*, *P. thumbergii*.
 Notes: (3) Inouye & Nobuchi 1957: 50 (re-described).
 References: (cn) Anonymous 1980g, (cc) Nishiguchi 1957: 75–78, 1959: 271, (ds) Anonymous 1980g: Choo 1983: 73; Choo & Woo 1985: 164; Kleine 1913b: 118, 1914b: 256, 1934a: 142; Kono 1938b: 65–68; Krivolutskaya 1983; Murayama 1948: 3, 1949a: 12, 1953a: 10, 1954b: 199; Nakane et al. 1963: 382; Nobuchi 1966d: 24, 1985c: 14; Sawamoto 1940a: 95–101, (tx) Choo 1983: 73; Hagedorn 1910a: 99; Inouye & Nobuchi 1957a: 49–50; Murayama 1953a: 10, 1954b: 199; Nakane et al. 1963: 382; Niisima 1909: 142; Nobuchi 1966d: 24, pl. 3; Sawamoto 1940a: 95, 100; Schedl 1934f: 1693, 1979c: 136.
- laticollis Browne** 1984b: 288. Holotype, sex?; Pomio (New Britain) to Nagoya (Japan), imported; BMNH, London.
 Distribution: New Guinea.
 Hosts: *Garcinia* sp.
 Notes: (3) Beaver 1991: 91 (cited in *Hypocryphalus* [not seen by us]).
 References: (ds) Ohno, Yoshioka, et al. 1989: 60, (tx) Beaver 1991: 91; Browne 1974a: 67, 1984b: 288.
- latus Eggers** 1929b: 10. Holotype, sex?; Ussuri, USSR; USNM, Washington, 2 Eggers cotypes in NHMW, Wien.
 Figures: Yin, Huang, & Li 1984: 103.
 Distribution: Asia (Heilongjiang in China/ Ussuri in E USSR).
 Hosts: *Abies* sp., *Larix dahurica*, *L. gmelinii*, *L. sp.*, *Picea* sp.
 References: (cn) Kurenzov 1935c: 188, (cc) Kurenzov 1934a: 57, (hb) Kurenzov 1935a: 19, 32, 1948b: 103; Stark 1952: 264, (ds) Krivolutskaya 1983; Kurenzov 1934a: 57, 1935a: 19, 32, 1935c: 188, 1936a: 113, 1938a: 65, 1951b, 1967; Schedl 1969a: 202; Stark 1936e: 143, 1952: 264; Tsai & Li 1963: 602; Yanovskii & Tegshzhargal 1985: 415; Yin, Huang, & Li 1984: 102, (tx) Anderson, W. H. & Anderson 1971: 17; Eggers 1929b: 10; Kurenzov 1941a: 143–144, 1948b: 103; Murayama 1943: 97; Schedl 1934f: 1639, 1979c: 137; Sokanovskii 1954: 16; Stark 1936e: 143, 1952: 264; Tsai & Li 1963: 605; Yin, Huang, & Li 1984: 102.
- lepocrinus Tsai & Li** 1963: 606, 619, 624. Holotype ♂; Szechuan, China; IZAS, Beijing.
 Figures: Tsai & Li 1963: 620; Yin, Huang, & Li 1984: 117 (adult).
 Distribution: Asia (Sichuan in China).
 Hosts: *Abies* sp.
 References: (ds) Yin, Huang, & Li 1984: 116, (tx) Tsai & Li 1963: 606, 609, 619–620, 624; Yin, Huang, & Li 1984: 116.
- leprosulus Browne** 1980e: 774. Holotype, sex?; Zaire: Riviere Samliki, 690 m; BMNH, London.
 Distribution: Africa (Zaire).
 References: (tx) Browne 1980e: 774.
- lipingensis Tsai & Li** 1959: 90. Holotype, sex?; North China (publication not seen); IZAS, Beijing.
 Figures: Li, Dang, & Shi 1977: 192, Li & Zhou 1980: 192, Tsai & Li 1959: 91, 1963: 616.
 Distribution: Asia (Shanxi, Sichuan in China).
 Hosts: *Pinus armandi*.
 References: (hb) Li, Dang, & Shi 1977: 73; Li & Zhou 1980: 73; Yang 1959c, (ds) Li, Dang, & Shi 1977: 73; Yin, Huang, & Li 1984: 113, (tx) Tsai & Li 1959: 90–91, 1963: 606, 616, 624; Yin, Huang, & Li 1984: 113.
- longior Browne** 1984d: 91. Holotype, sex?; New Guinea: Kaisenik; BMNH, London.
 Distribution: New Guinea.
 Hosts: *Castanopsis* sp.
 References: (tx) Browne 1984d: 91.
- longipennis (Browne)** 1970: 552 (*Taenioglyptes*). Holotype ♀; New Zealand: Greymouth; BMNH, London.
 Distribution: New Zealand.
 References: (tx) Browne 1970: 552–553.
- longipilus Schedl** 1943b: 34. Lectotype ♀; Sabaan, Ins. Mindoro, Philippinen; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 141.
 Distribution: Indonesia (Sumatra), Micronesia (Palau), Philippine Islands (Mindoro).
 References: (ds) Schedl 1966b: 21, (tx) Nobuchi 1983: 300; Schedl 1943b: 34, 1962r: 91, 1979c: 141; Wood, S. L. 1960a: 25.
- longisetosus (Nobuchi)** 1975: 57 (*Taenioglyptes*). Holotype ♀?; Hirogawara, Yamanashi pref., Japan; Nobuchi Collection, Ibaraki.
 Figures: Nobuchi 1975: pl. 2, fig. 20.

Distribution: Asia (Japan).

Hosts: *Tsuga diversifolia*.

References: (ds) Nobuchi 1955c: 14. (tx) Nobuchi 1975: 57.

major Stebbing 1903a: 270. Holotype ♀; India, Tono Valley; FRI, Dehra Dun.

Distribution: Asia (Himachal Pradesh, Punjab, Uttar Pradesh in India).

Hosts: *Picea morinda*, *Pinus excelsa*, *P. roxburghii*. References: (cn) Beeson 1915a: 317–325; Matlur & Singh 1960b: 51; Pierce, W. D. 1917: 74; Stebbing 1903a: 270, 1914: 530; Troup 1916: 1–126. (cc) Stebbing 1914: 530. (hb) Beeson 1915a: Stebbing 1903a: 270, 1914: 530. (ds) Beeson 1961: 286; Hagedorn 1910d: 44; Kleine 1913b: 118, 1914b: 276, 1934a: 142; Matlur & Singh 1960b: 51; Pierce, W. D. 1960b: 51; Stebbing 1903a: 270. (tx) Hagedorn 1910a: 87; Stebbing 1903a: 270, 1914: 530; Wood, S. L. 1959: 171.

morinda Stebbing 1903: 265. Syntypes 2, sex?; Sakahar, Bashar, Punjab, India; FRI, Dehra Dun. Synonymy: Wood 1959: 171.

References: (cn) Pierce, W. D. 1917: 81; Stebbing 1903a: 265, 1914: 535. (cc) Chararas 1959c; Stebbing 1914: 535. (hb) Stebbing 1903a: 265, 1914: 535. (ds) Hagedorn 1910d: 44; Kleine 1913b: 118, 1914b: 269, 1934a: 142; Pierce, W. D. 1917: 81; Stebbing 1903a: 265. (tx) Hagedorn 1910a: 87; Stebbing 1903: 265, 1914: 535; Wood, S. L. 1959: 171.

malloti Schedl 1943b: 37. Lectotype, sex?; Baguio, Luzon, Philippines; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 147.

Distribution: Philippine Islands (Luzon).

Hosts: *Mallotus ricinoides*, *Pinus insularis*.

References: (ds) Schedl 1966b: 21. (tx) Nobuchi 1953: 300; Schedl 1943b: 37, 1952b: 363, 1953c: 295, 1979c: 147.

malus Niisima 1909: 144. Syntypes, sex?; Hakodate; Nobuchi Collection, Ibaraki.

Distribution: Asia (Liaoning in China/ Japan/ Korea).

Hosts: *Armanica* sp., *Prunus armeniaca*, *P. mume*, *P. persica*, *P. pseudo-cerasus*, *P. salicina*, *P. sargentii*, *Pyrus malus*, *Sorbus commixta*.

References: (cn) Clausen 1931. (ds) Choo 1983: 74; Choo & Woo 1985: 165; Choo, Woo, & Nobuchi 1983: 171, 1985a: 134; Clausen 1931; Kleine 1913b: 118, 1914b: 257, 1934a: 142; Murayama 1954b: 199, 1955: 103; Nobuchi 1955c: 14; Tsai, Huang, & Li 1984: 104; Tsai & Li 1963: 602. (tx) Hagedorn 1910a: 88; Murayama 1954b: 199, 1955: 123; Niisima 1909: 144; Schedl 1934f: 1639; Tsai, Huang, & Li 1984: 104; Tsai & Li 1963: 605–606, 623.

mandschuricus Eggers 1929c: 10. Lectotype ♂; Ussuri; USNM, Washington, designated by Anderson & Anderson 1971: 19.

Figures: Yin, Huang, & Li 1984: 113 (gallery).

Distribution: Asia (Heilongjiang in China/ E USSR).

Hosts: *Corylus mandshurica*, *Juglans mandshurica*. Notes: (3) Tsai & Li 1963: 25 (*coryli* Stark, nomen nudum, synonymy).

References: (cn) Kurenzov 1935c: 191. (cc) Kurenzov 1934a: 57. (hb) Kurenzov 1935a: 19, 30, 1945b: 113; Stark 1952: 262; Tsai & Li 1963: 599; Yin, Huang, & Li 1984: 113. (ds) Kleine 1934a: 142; Krivolutsкая 1983; Kurenzov 1934a: 51, 1935a: 19, 30; 1935c: 191, 1936a: 111, 1936b: 350, 1938a: 61; Murayama 1939: 137, 1942a: 49, 55; Stark 1936c: 143, 1952: 262; Tsai & Li 1963: 599; Yin, Huang, & Li 1984: 113. (tx) Anderson, W. H. & Anderson 1971: 19; Eggers 1929c: 10; 1942c: 28; Kurenzov 1941a: 145–146, 1945b: 113; Murayama 1939: 137, 140; Schedl 1934f: 1639, 1979c: 148; Stark 1936c: 143, 1952: 262; Tsai & Li 1963: 606; Yin, Huang, & Li 1984: 113.

markangensis Tsai & Li 1963: 606, 615, 624. Holotype ♂; Szechuan (Markan); IZAS, Beijing.

Figures: Tsai & Li 1963: 618; Yin, Huang, & Li 1984: 116 (head, antenna).

Distribution: Asia (Sichuan in China).

Hosts: *Abies faxoniana*.

References: (ds) Yin, Huang, & Li 1984: 115. (tx) Tsai & Li 1963: 606, 615, 624; Yin, Huang, & Li 1984: 115.

massonianus Tsai & Li 1963: 605, 613, 626. Holotype ♂; Nanking, China; IZAS, Beijing.

Distribution: Asia (Jiangsu in China).

Hosts: *Pinus massoniana*.

Notes: Tsai & Li 1963: 613 (This species was treated as *Cryphalus piceus* in Tsai & Li 1959: 55–59; *C. piceus* is a different species).

References: (ds) Yin, Huang, & Li 1984: 109. (hb) Tsai & Li 1963: 599, 605, 613, 623, 626; Yin, Huang, & Li 1984: 109.

meridionalis (Nobuchi) 1975: 55 (*Tacnioglyptes*). Holotype ♀; Yona, Okinawa pref., Japan; Nobuchi Collection, Ibaraki.

Figures: Nobuchi 1975: pl. 2, fig. 19.

Distribution: Asia (Okinawa in Japan).

References: (ds) Nobuchi 1955c: 14. (tx) Nobuchi 1975: 55–56.

minimus Eggers 1927c: 76. Lectotype, sex?; Philippines: Luzon, Provinz Camarines Sur, Mount Isarog; USNM, Washington, designated by Anderson & Anderson 1971: 20.

Distribution: Micronesia (Caroline Islands, Palau), Philippine Islands (Luzon).

References: (ds) Schedl 1938g: 425; Wood, S. L. 1960a: 22. (tx) Anderson, W. H. & Anderson 1971: 20; Eggers 1927c: 76–77; Schedl 1938g: 425, 1943b: 34–35, 1953c: 289, 293, 1962r: 91, 1979c: 153; Wood, S. L. 1960a: 22, 25.

miyalopiceus Tsai & Li 1963: 602, 604, 608, 622. Holotype, sex?; China; IZAS, Beijing.

- Figures: Tsai & Li 1963: 608, Yin, Huang, & Li 1984: 102 (adult).
 Distribution: Asia (Sichuan in China).
 Hosts: *Picea asperata*.
 References: (ds) Yin, Huang, & Li 1984: 101. (tx) Tsai & Li 1963: 602, 604, 608, 618, 622; Yin, Huang, & Li 1984: 101.
- mollis Schedl** 1955b: 288. Holotype ♀; Fiji, Suva; BMNH, London.
 Distribution: Fiji Islands, Niue Island, Samoan Islands.
 Hosts: *Artocarpus* sp.
 References: (ds) Beaver 1976b: 534, 1987b: 64; Beaver & Maddison 1991: 1371; Browne 1974a: 64; Roberts 1976: 379; Schedl 1955b: 284. (tx) Beaver 1979c: 158, 1987b: 67; Schedl 1955b: 284, 288, 1979c: 158.
- montanus Nobuchi** 1964: 129. Holotype ♂; Shirotoge, Okunikko, Gumma pref., Japan; Nobuchi Collection, Ibaraki.
 Figures: Nobuchi 1964: 129.
 Distribution: Asia (Honshu in Japan).
 Hosts: *Abies mariesii*.
 References: (cn) Anonymous 1980g, (ds) Anonymous 1980g; Nobuchi 1985c: 14. (tx) McNamara 1977: 196; Nobuchi 1964: 129; Schedl 1979c: 158.
- neglectus Schedl** 1962p: 106. Holotype, sex?; Bengal: Sunderbans; Schedl Collection in NHMW, Wien.
 Distribution: Asia (Bengal in India).
 Hosts: *Hibiscus tiliaceus*.
 Notes: (3) This is *Erierythralus neglectus* Beeson 1941: 373 (1961: 287), nomen nudum. The type was sent by Beeson to Schedl for comparison, but was never returned to the Forest Research Institute.
 References: (cn) Mathur & Singh 1960b: 51. (ds) Beeson 1961: 278; Mathur & Singh 1960b: 51; Schedl 1971a: 275. (tx) Schedl 1940d: 591, 1950f: 49, 1953c: 289, 1962p: 106, 1979c: 164.
- negrosensis Browne** 1979: 85. Holotype, sex?; Philippines: Negros, Dewey Island; BMNH, London.
 Distribution: Philippine Islands (Negros).
 Hosts: Bucasian seed, *Rhizophora* sp.
 References: (tx) Browne 1979: 85; Nobuchi 1983: 300.
- niger Schedl** 1942c: 172. Holotype ♂; Australien, Queensland; Schedl Collection in NHMW, Wien.
 Distribution: Australia (Queensland).
 References: (hb) Schedl 1977a: 37. (tx) Schedl 1942c: 172–173, 1977a: 37, 1979c: 166.
- nigericus Browne** 1973a: 289. Holotype ♀; Nigeria: Bongo; MRCB, Tervuren.
 Distribution: Africa (Nigeria).
 Hosts: *Albizia glaberrima*.
 References: (tx) Browne 1973a: 289.
- nigricans Schedl** 1943b: 35. Lectotype ♀; Baguio, Luzon, Philippinen; Schedl Collection in NHMW, Wien.
 Distribution: Asia (Sri Lanka), Philippine Islands (Luzon).
 Hosts: *Myristica dactyloides*.
 References: (ds) Schedl 1966b: 22. (tx) Nobuchi 1983: 300; Schedl 1942c: 172, 1943b: 35, 1979c: 167.
- niponensis Inouye & Nobuchi** 1957a: 51. Syntypes 4, sex?; Somkyo, Daisetsu-zan National Park, Hokkaido, Japan; Nobuchi Collection, Ibaraki.
 Distribution: Asia (Hokkaido in Japan).
 Hosts: *Picea jezoensis*.
 References: (ds) Nobuchi 1985c: 14. (tx) Inouye & Nobuchi 1957a: 51.
- nitens Browne** 1980: 494. Holotype, sex?; Togian Isl. (Celebes) to Osaka (Japan), imported; BMNH, London.
 Distribution: Indonesia (Celebes).
 Hosts: *Gauna* sp.
 References: (ds) Browne 1985a: 191, 1986c: 663. (tx) Browne 1980: 494.
- nitidipennis Browne** 1984d: 89. Holotype, sex?; New Guinea: Stony L. A.; BMNH, London.
 Distribution: New Guinea.
 Hosts: *Protium* sp.
 References: (ds) Browne 1983a: 555, 1984d: 89. (tx) Browne 1984d: 89.
- nothofagi Browne** 1984c: 68. Holotype, sex?; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London.
 Distribution: New Guinea.
 Hosts: *Nothofagus* sp.
 References: (tx) Browne 1984c: 68.
- numidicus Eichhoff** 1878b: 487. Syntypes, sex?; Europae Graecia; Hamburg Museum, lost.
 Distribution: Africa (Algeria/ Egypt/ Libya/ Morocco/ Tunisia), Asia (Turkey), Europe (Bulgaria/ Corsica/ Greece/ Italy/ Spain/ Switzerland).
 Hosts: *Abies numidica*.
 References: (ay) Chararas 1973c. (bv) Chararas 1973c; Chararas, Desveaux, & Kogane-Charles 1978a; Grune 1979: 105. (cn) Chararas 1977c; Strohmeier 1930: 2. (cc) Chararas, Desveaux, & Kogane-Charles 1978a; Kleine 1908c: 186; Peyerimhoff 1934: 52. (hb) Barbey 1934; Dombrowsky 1887; Eichhoff 1881a: 45, 176; Henschel 1895a: 167; Lepiney & Mimeur 1932: 45; Postner 1974: 431; Strohmeier 1930: 2. (ds) Champion 1894; Grune 1979: 105; Hagedorn 1910d: 44; Henschel 1895a: 167; Heyden, Reitter, & Weise 1883: 181, 1891: 671, 1906: 711; Kleine 1913b: 119, 1934a: 142; Ortzen 1886: 279; Peyerimhoff 1919: 284, 1933b: 361, 1934: 52; Pfeffer 1935: 158, 1947d: 128; Pittioni 1943: 176; Postner 1974: 431; Reitter 1889a: 94, 1894a: 70; Schaufuss 1915: 1232; Schedl 1967c: 70; Stein & Weise 1877: 164; Tredl 1907: 12. (tx) Balachowsky 1949a: 207; Dombrowsky 1887; Eggers 1940g: 37; Eichhoff 1868b: 404, 1878a: 385, 1878b: 124, 487, 1881a: 45, 176, 1883a: 109, 132; Grune 1979: 105; Hagedorn 1910a: 57; Henschel

- 1895a: 167; Hopkins 1915b: 37, 39; Pfeffer 1955a: 176; Postner 1974: 431; Reitter 1889a: 94, 1894a: 70, 1913a: 66; Schedl 1934f: 1639; Strohmeier 1930: 1-7.
- nyalubombae** Schedl 1957d: 49. Holotype, sex?; Congo Belge: Kivu, Mt. Kahizi: MRCB, Tervuren. Figures: Schedl 1961k: 465. Distribution: Africa (Zaire). Hosts: *Xymalos monospora*. References: (ds) Schedl 1961k: 464, (tx) Schedl 1957d: 49, 1961k: 464-465, 1979c: 173.
- oblongus** Niisima 1910a: 9. Holotype, sex?; Tokio, Japan; Nobuchi Collection, Ibaraki. Distribution: Asia (Honshu in Japan). Hosts: *Pinus densiflora*. Notes: (3) Nobuchi 1985c: 16 (treated in *Hypothenemus* [not seen by us]). References: (ds) Kleine 1934a: 143; Murayama 1945: 2, 1949a: 12, 1951c: 3, 1953a: 10, 1954b: 164; Nobuchi 1966b: 22, 1985c: 16. (tx) Murayama 1953a: 10, 1954b: 164; Niisima 1910a: 9; Nobuchi 1966b: 22; Schedl 1934f: 1640.
- orientalis** Eggers 1911a: 122. No mention of types; Herzegovina: Metallon Pass; not given. Distribution: Europe (Yugoslavia). Notes: (1) Originally named as a subspecies of *piccae*, elevated to species rank by Stark 1936c: 143. References: (ec) Krivosheina 1974; Nikitsky 1978; Poinar 1975: 151. (hb) Stark 1952: 260. (ds) Arnoldi et al. 1955: 689; Pfeffer 1924b: 472; Pjatkitskii 1930a: 165; Stark 1936c: 143, 1952: 260. (tx) Eggers 1911a: 122-123; Iablokoff-Khmzorian 1961: 157; Pfeffer 1932b: 17; Pjatkitskii 1932: 295-302; Schedl 1934f: 1639; Sokanovskii 1954: 17; Stark 1936c: 143, 1952: 260.
- paganus** Eichhoff 1878b: 43, 129-130, 474. Holotype, sex?; Insula principalis prope oram Guineensem Africanam; Dohrn Collection at Stettin Museum, now in Schedl Collection in NHMW, Wien. Distribution: Africa (Equatorial Guinea/ Ghana/ Guinea/ Principe Island/ Sao Tome Island/ Zaire). References: (ds) Hagedorn 1910d: 44; Kleine 1913b: 119, 1914b: 315; Schedl 1961k: 465, 1962h: 55, 1962k: 1065, 1964e: 65, 1971g: 190. (tx) Browne 1965: 188; Eggers 1943c: 68; Eichhoff 1878b: 43, 129-130, 474; Hagedorn 1910a: 87; Schedl 1961k: 465, 1962k: 1065, 1979c: 182.
- palawanus** Schedl 1942a: 174. Syntypes, sex?; N. Palawan, Binaluan, Philippinen; Schedl Collection in NHMW, Wien. Distribution: Asia (Sri Lanka), Philippine Islands (Palawan). Notes: (1) Schedl 1979c: 182 (citation of holotype invalid). References: (ds) Schedl 1959a: 481, 1966b: 22. (tx) Nobuchi 1953: 300; Schedl 1942a: 174, 1959a: 481, 1979c: 182.
- pallidus** Eichhoff 1872a: 131. Holotype ♂; Madagascar: IRSNB, Brussels. Distribution: Africa (Burkina Faso/ Mauritius/ Seychelles Islands), Madagascar. Hosts: *Ficus melleri*, *F. thomningii*. References: (hb) Beaver 1957a: 17; Paulian 1951: 27; Schedl 1961k: 466. (ds) Alluaud 1900: 439; Anonymous 1892: 165; Beaver 1957a: 17; Fairmaire 1892b; Gemminger & Harold 1872: 2683; Hagedorn 1910d: 44; Kleine 1913b: 119; Lee 1971: 31; Lepesme 1945: 145; Paulian 1951: 27; Sampson 1914: 382; Schedl 1961k: 466. (tx) Duffy 1953: 13; Eichhoff 1872a: 131, 1878b: 127-128, 530; Fairmaire 1892b; Hagedorn 1910a: 87, 1913b: 254; Lepesme 1945: 145; Powell, W. 1980: 29; Sampson 1914: 382; Schedl 1951j: 19, 1961k: 466, 1969d: 8, 11, 1977a: 40, 1979c: 182.
- parculus** Niisima 1910a: 8. Holotype, sex?; Tokio, Japan; Nobuchi Collection, Ibaraki. Distribution: Asia (Honshu in Japan). References: (ds) Murayama 1954b: 199; Nobuchi 1985c: 14. (tx) Murayama 1954b: 199; Niisima 1910a: 8; Schedl 1934f: 1640.
- parvus** Browne 1984a: 152. Holotype, sex?; Viru Harbour (Solomon Islands) to Shimizu (Japan), imported; BMNH, London. Distribution: Solomon Islands. References: (tx) Browne 1984a: 152.
- pexus** Schedl 1979f: 104. Holotype, sex?; Samoa, Upolu, Tapatapao; Schedl Collection in NHMW, Wien. Distribution: Samoan Islands. References: (tx) Schedl 1979f: 104.
- piccae** (Ratzburg) 1837: 163 (*Bostrichus*). Syn-types, sex?; Oberschlesien und Baiern; not located. Figures: Balachowsky 1949a: 16, 204, 206, 208. Chararas 1962c: 242, Grune 1979: 104, Hasek 1961: 5, Nobuchi 1966d: pl. 3, Pfeffer 1989a: pl. 6, Postner 1974: 429. Distribution: Africa (Algeria), Asia ("Manchuria" in China/ Japan/ Korea/ Kuril Islands/ Turkey/ Sakhalin Island, Siberia in E USSR), Europe (Austria/ Bulgaria/ England/ France/ Germany/ Greece/ Hungary/ Italy/ Poland/ Romania/ Spain/ Switzerland/ W USSR/ Yugoslavia). Hosts: *Abies cephalonica*, *A. firma*, *A. holophylla*, *A. mariesii*, *A. pectinata*, *A. sachalinensis*, *Picea excelsa*, *P. glehnii*, *P. jezoensis*, *Pinus densiflora*. Notes: (3) Inouye & Nobuchi 1957: 52 (re-described). Ferrari 1867a: 12 misidentified *asperatus* var. *abietis*; it was actually *piccae*. References: (ay) Chararas 1973c; Escherich 1923b: 462, 481, 611; Hadorn 1933; Nunberg 1928a: 140; Nusslin 1910: 293, 1911a: 5, 255, 277, 337, 378; Ritchie 1919: 171; Scherb 1971. (bv) Chararas 1973c; Chararas & Stephanopoulos 1975a; Chararas et al. 1982: 1094; Grune 1979: 105; Krol & Zabecky 1976; Prell 1931: 364;

- Schneider-Orelli 1947c: 94; Schwerdtfeger 1981: 193; Wichmann 1953a: 107, 1967. (**cn**) Androic 1966: 48; Barbey 1901: 69, 1906a, 1925: 162; Bezares 1921, 1929: 38–42; Borcea 1924: 221–260, 1930: 271; Braun 1941b: 375; Bruneau 1950; Capecki 1982; Capek et al. 1957; Chararas 1959c: 113–129, 1959f: 3612–3614, 1961b: 69, 1961c: 92, 1978; Chorbadzhievo 1929; Eckstein 1915, 1926: 578; Elmstrom et al. 1974; Escherich 1923b: 462, 481, 611, 1936; Fankhauser 1896; Felt 1906: 753; Fice 1961: 173–204; Fitze 1953: 1–28; Gabler 1955; Georgescu et al. 1957: 357, 448; Georgijevic 1966: 3–48; Gradojevic 1920; Grandi 1951; Hasek 1955, 1961: 5; Hess 1898: 360; Hess & Beck 1914: 265, 1927: 321; Hierholzer 1954a: 338; Hopkins 1899: 444; Hrubik 1973; Inouye 1955; Inouye & Yamaguchi 1955a: 235; Jacentkovsky 1933: 270; Joly 1949b: 253, 1976b; Judeich & Nitsche 1895: 492, 526; Kahl 1896; Kailidis 1964a: 41–54, 1966a: 81–85, 1968; Kailidis & Georgevits 1968: 1–64, 1972; Kailidis & Markalas 1988; Keller 1903b: 58; Koch 1913: 95; Kovacevic 1957: 67; Krol & Zabecki 1976; Maksimovic & Milanovic 1964, 1966; Maksymov 1950: 502; Marcu 1926c: 63, 1930: 327–336; Marie 1926: 167–171; Martin & Cobos 1986; Merker 1954b: 209; Mizuno 1963: 272–277, 1966: 93–97; Muller 1912: 185; Murayama 1954a: 14; Nosek 1951: 105, 1952b: 98; Novak, V., Hrozinka, & Stary 1976: 69; Nusslin 1913: 223; Pfeffer 1924b: 472; Pierce, W. D. 1917: 69; Regula 1955: 120; Rhumbler 1922: 299, 1927: 312; Ritchie 1919: 171; Scheidter 1919: 84, 1920: 209; Schimitschek 1937b: 9, 1937c: 52, 1947g: 188, 1949b: 180, 1950: 53, 1952a: 194, 1955a: 43, 85, 1955b: 102, 1955c: 82, 1956: 342, 1961a: 154; Schmidt 1881: 34; Schneeberg 1925: 495; Schneider-Orelli & Kuhn 1948: 533; Schwappach et al. 1929: 186; Schwerdtfeger 1944a: 182, 1950b: 68, 1957a: 188; Sedlaczek 1921: 335, 1933: 307, 1936: 200; Shiraki 1952; Sinreich 1958: 196–198, 1961: 166, 1962; Slander 1948: 10; Spaic 1955: 454; Stockl 1941: 177; Strohmeier 1930: 2; Sturm 1843: 230; Thalenhorst 1950: 90; Tamanuki 1933: 1–54; Tomie 1957: 207–210; Trinchieri 1921: 1–12; Wachtl 1901: 381; Wahl & Muller 1913: 1–70, 1919: 168–170; Wardle 1929: 325; Weber, H. 1926: 578; Wilke 1931: 636; Wohlmann 1936: 41; Wolff & Krausse 1922: 88; Yamagisawa 1952: 106; Zivojinovic 1950: 299–310. (**cc**) Apfelbeck 1916b; Balazy & Michalski 1964b; Barbey 1906a; Belanowski 1930; Capecki 1982; Capek 1957; Capek & Capecki 1979; Chararas 1958b, 1959a, 1959c, 1959d, 1959e, 1959f, 1960a: 31; Chararas et al. 1982: 1094; Fuchs 1914a, 1929a, 1930; Gaulle 1906: 237; Györfi 1952b; Haeselbarth 1962: 283, 1967; Hierholzer 1954a: 338, 1954b: 387; Hirschmann & Wisniewski 1982: 1983; Hubandt 1923a; Inouye 1954: 171; Inouye & Yamaguchi 1955a: 235; Inouye et al. 1955: 58; Jannicky 1957c; Joly 1949a: 7; Karpinski 1932a: 95; Kielczewski, Moser, & Wisniewski 1983; Kielczewski & Wisniewski 1978, 1983; Kleine 1908c: 186, 1909a: 46, 1944: 80; Kraemer 1950b: 354; Krol & Zabecki 1976; Merker 1954b: 209; Michalski & Ragajczak 1989; Nishiguchi 1959: 271; Nosek 1951: 105, 1952b: 98; Numberg 1930: 203; Nusslin 1927: 300; Perris 1856a: 244; Pfeffer 1923a: 331, 1928b: 7, 1932a: 18, 1943b: 179, 1955b: 84, 1960: 345; Poinar 1975: 151; Rondani 1873: 146; Ruhm 1955c: 76, 1956b: 3; Scheidter 1919: 69–90, 1920, 1936: 235; Schimitschek 1930a: 326, 1950: 53, 1952a: 194, 1955a: 43, 85, 1964e; Schwerdtfeger 1944a: 182, 1950b: 68, 1957a: 188; Sedlaczek 1900: 503, 1908: 46, 1933: 307, 1935a: 163; Szymusik 1923: 7; Tenkacova & Mituch 1987; Thompson, W. R. 1943: 37; Wichmann 1953a: 107, 1967; Wilke 1931: 636; Yasumatsu & Watanabe 1965: 68. (**hb**) Altum 1881c: 322; Apfelbeck 1916b, 1917; Barbey 1901: 10, 22, 69, 1913, 1925: 162, 1934, 1942; Bargmann 1897a, 1900, 1906; Beffa 1949, 1961; Borcea 1924; Braun 1941c; Cecconi 1906, 1924; Chamberlin 1939: 315; Chararas 1961b: 69, 1961c: 92, 1962c: 242; Charvat 1950; Chorbadzhievo 1929; Dombrowsky 1887; Eckstein 1897, 1915, 1926: 578; Eichhoff 1881a: 45, 172; Escherich 1923b: 462, 481, 611; Fuchs 1904a, 1905c: 339; Gabler 1955; Gillanders 1906, 1908; Grandi 1951; Györfi 1957; Hadorn 1933; Heimbacher 1924; Hennings 1908b, 1908c: 215; Henschel 1876a: 240, 1895a: 167; Hess 1898: 360; Hess & Beck 1914: 265, 1927: 321; Inouye 1954: 171; Inouye et al. 1954: 129, 1955: 58; Joly 1949a: 7, 1976b; Judeich & Nitsche 1895: 526; Kalandra 1948: 107–116; Kalina 1970: 123; Karpinski & Strawinski 1948: 155; Kraemer 1950b: 354; Kurenzov 1935a: 20; Lengerken 1939: 39, 1954: 82; Maksymov 1950: 502; Masutti 1964; Mizuno 1963; Nordlinger 1856: 29; Novak, V., Hrozinka, & Stary 1976: 69; Numberg 1929: 102; Nusslin 1898: 280, 1906b: 14, 1907: 613, 1913: 223, 1927: 300; Perris 1856a: 244; Pfeffer 1941b: 16; Postner 1974: 429; Prell 1931: 364; Ratzeburg 1837: 157, 1839: 197; Reh 1900a: 94; Rhumbler 1922: 299, 1927: 312; Ritchie 1919: 171; Rupertsberger 1880: 229; Schimitschek 1929: 83, 1930a: 326, 1955a: 43, 85; Schmidt 1881: 34; Schneider-Orelli 1947c: 94; Schwappach et al. 1929: 186; Schwerdtfeger 1944a: 182, 1957a: 188, 1951: 193; Sedlaczek 1900: 503, 1921: 335, 1935a: 163; Spessivtsev 1913a: 91; Stark 1952: 260; Strohmeier 1930: 2; Szajek 1955: 242; Wachtl 1901: 381; Weber, H. 1926: 578; Wolff & Krausse 1922: 88. (**ds**) Acloune 1896; Allen, A. A. 1965b: 47; Anmann & Knabl 1923; Androic 1966: 48–49; Anonymous 1928c: 202; Arnoldi et al. 1955: 689; Arri, Covassi, & de Bellis 1966: 32; Andras & Schaefer 1957; Bach 1854: 136; Barthe 1896; Ban 1888; Bandisch 1899; Bedel 1888b: 398, 414; Beffa 1949; Bielz 1887;

- Blanchere & Robert 1889; Blatchley & Leng 1916: 606; Borcea 1924; Borchert 1951; Brakman 1966b: 206; Brancsik 1906; Bucking 1932; Buresh & Lazarov 1956; Capek et al. 1957; Cecconi 1897, 1906; Chamberlin 1939: 315; Chapuis & Candeze 1853; Charvat 1950; Cho 1955, 1957; Choo 1983: 74; Choo & Woo 1985: 165; Chorbadzhievo 1924d, 1929; Crotch 1863; Eggers 1904; Endrodi 1955b; Escalera 1919; Fowler 1891; Fuchs 1904a, 1905a, 1905c: 339; Fuss 1874; Gabler 1949b; Gaubil 1849: 126; Gozis 1875: 80; Gredler 1866: 373, 1875: 115; Grune 1979: 105; Hagedorn 1910d: 44; Henschel 1895a: 167; Heyden 1879: 140; Heyden, Reitter, & Weise 1883: 181, 1891: 671, 1906: 711; Holdhaus & Deubel 1910: 145; Horion 1935, 1951; Hubault 1923b; Jacentkovsky 1933: 270, 1939: 76; Jazentkovsky 1912: 288; Joly 1949b: 253, 1976b; Judeich & Nitsche 1895: 526; Kailidis 1964a, 1966: 55, 1985; Kailidis & Georgevits 1972; Kailidis & Markalas 1988; Kaltenbach 1874: 685; Karpinski 1925: 216, 1926: 82, 1932a: 95, 1948b: 230; Karpinski & Strawinski 1948: 155; Keler 1925b: 272; Kestercanek 1881a: 12; Kiefer et al. 1942: 528; Kleine 1912a: 161, 263–264, 267, 1913a: 35, 1913b: 119, 1914b: 256, 1934a: 142; Kloft & Hinks 1945: 218; Knotek 1892a: 37, 1894a: 555; Ko 1969: 275; Kocher 1953: 133; Kono 1938b: 65; Kono & Tamamki 1939: 88, 92–93; Kovacevic 1957: 67; Kraatz 1869: 59; Krivolutskaia 1983; Ku 1964; Kugela 1946a: 339; Kurenzov 1935a: 20; Kurir 1947c: 17; Lacordaire 1866: 379; Langhoffer 1915c: 157; Leng 1918: 211; Lokaj 1868: 64; Lomnicki 1913b: 145; Lucht 1987: 278; Marcu 1926c: 63; Matthews & Fowler 1883: 42; Michalski 1957: 164; Munro 1926: 4–77; Murayama 1929e: 44, 1930b: 7, 1949a: 12, 1949c: 101, 1950b: 1291, 1953a: 10, 1953c: 151, 1954a: 14, 1954b: 164; Negri 1966b: 402, 1968a: 456; Nobuchi 1966d: 24, 1985c: 14; Numberg 1928b: 88, 105, 1954: 52, 1964a: 235; Nusslin 1898: 280; Orest 1926c: 63; Perris 1876a: 254, 1877a: 414; Pfeffer 1924a: 96, 1928b: 7, 1931b: 75, 1947d: 125, 1960: 345, 1989a: 81; Pierce, W. D. 1917: 69; Pittioni 1943: 175; Postner 1974: 429; Prossen 1913: 83; Rapp 1934: 727; Ratzeburg 1837: 157, 1839: 197; Redtenbacher 1858: 832, 1874: 376; Reitter 1869a: 154, 1888b: 280, 1889a: 94, 1894a: 70, 1916: 288, 350; Ronbal 1941: 263; Sainte-Claire & Mequignon 1938: 445; Schanfuss 1915: 1231; Schann 1859: 96, 1862: 101; Schedl 1959b: 100, 1964a, 1967c: 70, 1980a: 14, 1981b: 74; Scheerpeltz & Winkler 1930: 257; Schilsky 1909: 188; Schwerdtfeger 1981: 193; Seidl 1876: 4; Seidlitz 1891a: 562, 1891b: 607; Sharp & Fowler 1893: 34; Shiraki 1952; Solla 1893: 217; Stark 1926b: 103, 1927b: 88, 1936c: 143, 1952: 260; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 439; Stierlin & Cantard 1871: 292, 1906: 205; Swaine 1909: 92; Szujecki 1955: 242; Thomson 1865: 359; Tredl 1907: 12; Wanka 1920: 202–213; Wichmann 1924: 15–17, 1927a: 67; Wilke 1931: 636; Winter, T. G. 1983: 8. (tx) Acloque 1896; Allen, A. A. 1965b; Bach 1854; Balachowsky 1949a: 16, 204, 206, 208; Barbey 1901: 10, 22, 69; Bedel 1888b: 398, 414; Beffa 1949, 1961; Bertolini 1872; Blatchley & Leng 1916: 606; Browne 1970: 552; Ceballos 1945; Chapuis & Candeze 1853; Chararas 1962c: 242; Charvat 1950; Choo 1983: 74; Chorbadzhievo 1924d; Csiki 1910; Doebner 1860, 1862; Dobrowsky 1887; Duffy 1953; Eggers 1911a: 121, 1921: 39–40, 1926b: 136, 1940g: 37; Eichhoff 1864b: 35, 1876a: 379, 1878a: 378–379, 1878b: 122, 1881a: 45, 172–176, 1883a: 109, 132; Endrodi 1957b; Escherich 1923b: 462, 481, 611; Fanvel 1889; Ferrant 1911; Ferrari 1867a: 12, 15; Fleischer 1905, 1927; Formanek 1907: 34; Gabler 1949b, 1955; Gebien 1907: 197; Gillanders 1906, 1908; Grune 1979: 104–105; Hagedorn 1910a: 87; Hasek 1961: 5; Henschel 1876a: 240, 1878a, 1895a: 167; Hopkins 1914: 130, 1915b: 39; Iablokoff-Khnzorian 1961: 157; Inouye & Nobuchi 1957a: 52; Joh 1976, 1976b; Judeich & Nitsche 1895: 526; Kailidis 1966a; Kalina 1970: 124; Karpinski 1949: 129; Karpinski & Strawinski 1948: 155; Kama 1970: 278; Knotek 1892a: 37; Koch 1913: 95; Kuhn 1913: 1055; Lacordaire 1866: 379; Letzner 1891: 375; Lucht 1987: 278; Murayama 1950b: 1291, 1953a: 10, 1954b: 164; Negri 1966b: 402; Niisima 1908a: 90, 1909: 141, 1910a: 8, 1929: 376–381; Nobuchi 1966d: 24, pl. 3; Nordlinger 1848: 244, 1856: 29; Novak, V., Hrozinka, & Stary 1976: 69; Numberg 1928a: 140, 1930: 200–208, 1948a: 1–30, 1954: 52; Nusslin 1911a: 5, 255, 277, 337, 378; Perris 1877a: 414; Pfeffer 1932b: 17, 1941b: 16, 1955a: 176, 1989a: pl. 6; Portevin 1935: 324; Postner 1974: 429; Quaschik 1953: 35; Ratzeburg 1837: 157, 163, 1839: 197; Redtenbacher 1849a: 791, 1849b: 26, 1858: 832, 1874: 376; Reitter 1889a: 94, 1894a: 70, 1913a: 66, 1916: 288, 350; Rhumbler 1922: 299, 1927: 312; Rupertsberger 1880: 229; Schedl 1934f: 1639, 1952f: 87, 1980a: 14, 1981b: 74; Scherb 1971; Schmittschek 1936a: 565–602, 1937b: 1–25, 1937c: 52, 1955c: 82; Seidlitz 1891a: 562, 1891b: 607; Spessivtsev 1913a: 91, 1931: 49, 1936: 143; Stark 1952: 260; Stierlin 1898: 439; Swaine 1909: 92–93; Thomson 1865: 359; Wood, S. L. 1954a: 1002–1004. (ms) Chararas 1959d; Fankhauser 1885: 185; Mizano 1963; Reh 1900a: 94; Ritter 1929: 555; Schmittschek 1955b: 102; Sedlaczek 1936: 200; Sinreich 1962. *hattorii* Kono 1938b: 67. Holotype, sex?; Nukahira; Kono Collection. Synonymy: Inouye & Nobuchi 1957a: 52. References: (cn) Inouye & Yamaguchi 1955a: 235, (cc) Inouye & Yamaguchi 1955a: 235. (ds) Kono 1938b: 65, 67–68; Murayama 1954b: 199. (tx) Inouye & Nobuchi 1957a: 52; Kono 1938b: 65, 67–68; Murayama 1954b: 199.

- subdepressus* Eggers 1940g: 37. Holotype, sex?; Kleinasien (Ayancik); Eggers Collection, in NHMW, Wien. Synonymy: Wood 1992a: 89. References: (cn) Schimitschek 1944: 172. (ec) Schimitschek 1941a: 305. (hb) Schimitschek 1944: 172. (tx) Eggers 1940g: 37–38; Schedl 1979c: 241; Wood, S. L. 1992a: 89.
- piceus* Eggers 1926b: 136. Lectotype, sex?; Hokkaido (Jozankei); USNM, Washington, designated by Anderson & Anderson 1971: 25. Figures: Tsai & Li 1959: 88. Distribution: Asia (Heilongjiang in China/ N Japan/ Sakhalin Island, Siberia in E USSR). Hosts: *Picea excelsa*, *P. glehnii*, *P. jezoensis*, *P. obovata*, *Pinus koraiensis*, *P. montana*. References: (cn) Anonymous 1980g; Inouye 1955; Inouye & Yamaguchi 1955a: 235; Kurenzov 1935c: 187, 1950a: 6, 1956a: 90, 1956b: 29. (ec) Inouye & Yamaguchi 1955a: 235; Kono & Watanabe 1935: 69; Kurenzov 1934a: 57; Nishiguchi 1960c: 64–73. (hb) Inouye, Koizumi, & Takai 1954: 129; Krivolutsкая 1956: 832, 1973: 134; Kurenzov 1935a: 32, 1948b: 109, 1950d: 146; Lunardon & Leonardi 1889: 457; Munro 1926: 60. (ds) Anonymous 1980g; Krivolutsкая 1956: 832, 1965a: 232, 1973: 134; Kurenzov 1934a: 57, 1935a: 32, 1935c: 187, 1936a: 111, 1936b: 351, 1938a: 59, 1957b: 17, 1965; Lunardon & Leonardi 1889: 457; Murayama 1951c: 4, 1954b: 199; Stark 1936e: 144, 1952: 268; Tsai & Li 1963: 603; Yin, Huang, & Li 1984: 108. (tx) Anderson, W. H. & Anderson 1971: 25; Beeson 1935: 107; Eggers 1926b: 136, 1942c: 29; Inouye & Nobuchi 1957a: 53; Koch 1932: 147; Krivolutsкая 1956: 832, 1958: 142; Kurenzov 1941a: 146–147, 1948b: 109, 1950: 146; Lunardon & Leonardi 1889: 457; Murayama 1954b: 199, 1958: 934; Schedl 1934f: 1639, 1979c: 193; Stark 1936e: 144, 151, 1952: 268; Tsai & Li 1959: 88, 1963: 605; Yin, Huang, & Li 1984: 108.
- pilifer* Eggers 1927b: 394. Syntypes 1 ♂, 1 ♀; Sumatra (Si Rambe); MCG, Genova, and 1 in Eggers Collection, in NHMW, Wien. Distribution: Indonesia (Sumatra). References: (tx) Eggers 1927b: 394; Schedl 1979c: 193.
- pilosellus* Erichson 1842: 212. Holotype, sex?; Vandiemensland; MNB, Berlin. Distribution: Australia (Tasmania). Hosts: *Bursaria spinosa*. Notes: (3) Eichhoff 1878b: 135 (re-described). References: (ds) Brimblecombe 1953: 17; Gemminger & Harold 1872: 2683; Hagedorn 1910d: 44; Kleine 1913b: 119, 1914b: 299. (tx) Eichhoff 1878b: 135; Erichson 1842: 212; Ferrari 1867a: 17; Hagedorn 1910a: 87; Lea 1910a: 143; Schedl 1936g: 527; 1943b: 36.
- pilosulus* Browne 1980b: 355. Holotype, sex?; Kimi (New Guinea) to Yatsushiro (Japan), imported; BMNH, London. Distribution: New Guinea. References: (tx) Browne 1980b: 355.
- pilosus* Tsai & Li 1963: 606, 617, 624. Holotype, sex?; Ssuehuan, China; IZAS, Beijing. Figures: Tsai & Li 1963: 616–617, Yin, Huang, & Li 1984: 115 (adult). Distribution: Asia (Sichuan in China). Hosts: *Abies faxoniana*. References: (tx) Tsai & Li 1963: 606, 616–617, 624; Yin, Huang, & Li 1984: 114.
- planicollis* Schedl 1978d: 74. Holotype, sex?; Nunkaledonien, Col. d'Amieu; Schedl Collection in NHMW, Wien. Distribution: New Caledonia Island. References: (tx) Schedl 1978d: 74.
- premayaensis* Murayama 1943b: 97. Holotype ♂; Premaya, dans le grand Khingan, Manchuria; Murayama Collection in USNM, Washington. Distribution: Asia ("Manchuria" in China). Hosts: *Larix gmelini*. References: (ds) Murayama 1943b: 97. (tx) Murayama 1943b: 97.
- procerus* Schedl 1953c: 296. Holotype, sex?; Sumatra, Post de Kock, 920 m; Schedl Collection in NHMW, Wien. Distribution: Indonesia (Sumatra). Notes: (3) Schedl 1953c: 296 (*sumatranus* Eggers, nomen nudum, synonymy). References: (tx) Schedl 1953c: 296, 1979c: 199.
- pruni* Eggers 1929c: 11. Holotype, sex?; Ussuri; USNM, Washington. Distribution: Asia (Yunnan in China/ Sakhalin Island, Siberia in E USSR). Hosts: *Prunus triloba*. References: (cn) Kurenzov 1935c: 189. (ec) Kurenzov 1934a: 50. (hb) Kurenzov 1935a: 19, 31, 1948b: 121; Stark 1952: 270. (ds) Kleine 1934a: 142; Krivolutsкая 1958: 144, 1983; Krivolutsкая & Kupyanskaya 1970; Kurenzov 1934a: 50, 1935a: 19, 31, 1935c: 189, 1936a: 110, 1936b: 350; Stark 1936e: 144, 1952: 270. (tx) Anderson, W. H. & Anderson 1971: 26; Eggers 1929c: 11; Kurenzov 1941a: 41, 146–149, 154, 1948b: 121; Schedl 1934f: 1639, 1979c: 200; Stark 1936e: 144, 151, 1942: 270.
- pseudochiulingensis* Tsai & Li 1963: 605, 610, 623. Holotype ♂; Shenhshi (Hsze Chuan), China. Distribution: Asia (Shanxi in China). Hosts: *Pinus armandi*, *P. tabulaeformis*. References: (hb) Yang 1989c. (ds) Yin, Huang, & Li 1984: 110. (tx) Tsai & Li 1963: 605, 610, 623; Yin, Huang, & Li 1984: 110.
- pseudotabulaeformis* Tsai & Li 1963: 605, 613, 623. Holotype ♂; Shenhshi (Li Ping), China; IZAS, Beijing.

- Distribution: Asia (Sichuan in China).
 Hosts: *Pinus tabulaeformis*.
 References: (**ds**) Yin, Huang, & Li 1984: 110 (**tx**) Tsai & Li 1963: 599, 605, 613, 623–624; Yin, Huang, & Li 1984: 110.
- puberulus** Schedl 1942c: 171. Lectotype ♀; Australien, Queensland, Imbil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 202. Distribution: Australia (Queensland).
 Hosts: *Grevillea robusta*.
 References: (**tx**) Schedl 1942c: 171, 1979c: 202
- pubescens** Hopkins 1915b: 39–40. Holotype ♀; Port Williams, Washington [USA]; USNM, Washington.
 Figures: Bright & Stark 1973: 157 (adult).
 Distribution: North America (British Columbia in Canada/ California, Oregon, Washington in USA).
 Hosts: *Abies grandis*, *Picea sitchensis*, *Pinus lambertiana*, *Pseudotsuga menziesii*, *Sequoia sempervirens*.
 References: (**bv**) Daterman, Rudinsky, & Nagel 1965. (**cn**) Doane et al. 1936; Hopkins 1904c: 16; Swaine 1918a: 87. (**cc**) Dextrup & Gara 1978: 274; Kinn 1971; Marsh 1979: 150, 158. (**hb**) Bright & Stark 1973: 67; Chamberlin 1939: 314, 1958: 133; Daterman, Rudinsky, & Nagel 1965; DeLeon 1952: 78; Doane et al. 1936; Hopkins 1904a: 16; Swaine 1918a: 87; Wood, S. L. 1953: 84, 1954a: 1003. (**ds**) Bright 1976d: 112; Bright & Stark 1973: 67; Chamberlin 1925, 1939: 314, 1958: 133; DeLeon 1952; Furniss, R. L. & Carolin 1977: 377; Keen 1929a: 29; Kleine 1934a: 142; Patterson & Hatch 1945: 152; Wood, S. L. 1972a: 415, 1982b: 867. (**tx**) Bright 1976d: 112; Bright & Stark 1973: 157; Chamberlin 1939: 314, 1958: 133; Hopkins 1915b: 39–40; Keen 1929a: 29; Swaine 1918a: 87–88; Wood, S. L. 1953: 84, 1954a: 959–960, 1002–1003, 1972a: 415, 1982b: 867.
- subconcentricus** Hopkins 1915b: 39–40. Holotype ♀; Astoria, Oregon [USA]; USNM, Washington. Synonymy: Wood 1954a: 1003.
 References: (**cn**) Doane et al. 1936; Swaine 1918a: 87–88. (**hb**) Chamberlin 1939: 313; Doane et al. 1936; Swaine 1918a: 87–88. (**ds**) Chamberlin 1917, 1925, 1939: 313; Hopping 1922; Keen 1929a: 29, 1934a: 142. (**tx**) Chamberlin 1939: 313; Hopkins 1915b: 39–40; Keen 1929a: 29; Swaine 1918a: 87–88; Wood, S. L. 1954a: 959, 979, 1003.
- pulchellus** (Nobuchi) 1975: 54 (*Taenioglyptes*). Holotype, sex?; Higashikawa, Nara pref., Japan; Nobuchi Collection, Ibaraki.
 Figures: Nobuchi 1975: pl. 2, fig. 17.
 Distribution: Asia (Japan).
 Hosts: *Aesculus turbinata*.
 References: (**ds**) Nobuchi 1985c: 15. (**tx**) Nobuchi 1975: 54.
- punctatostratus** Schedl 1942a: 175. Holotype, sex?; Singapur; Schedl Collection in NHMW, Wien.
- Distribution: Asia (Singapore in Malaya).
 References: (**ds**) Browne 1961c: 69. (**tx**) Schedl 1942a: 175, 1953c: 296, 1979c: 204.
- punctatus** Browne 1954: 66. Holotype, sex?; New Guinea: Morobe District, Wau, Kunai Creek; BMNH, London.
 Distribution: New Guinea.
 References: (**ds**) Browne 1954c: 66.
- punctipennis** Schedl 1942c: 169. Holotype, sex?; Australien, Queensland; Schedl Collection in NHMW, Wien.
 Distribution: Australia (Queensland).
 References: (**tx**) Schedl 1942c: 169, 1979c: 205.
- pusillimus** Schedl 1942c: 171. Holotype, sex?; New Guinea; Schedl Collection in NHMW, Wien.
 Distribution: New Guinea.
 References: (**tx**) Schedl 1942c: 171, 1979c: 206.
- pusillus** Schedl 1943b: 38. Holotype, sex?; Port Banga, Mindanao, Philippinen; Schedl Collection in NHMW, Wien.
 Distribution: Philippine Islands (Mindanao).
 References: (**ds**) Schedl 1966b: 22. (**tx**) Nobuchi 1983: 300; Schedl 1943b: 38, 1951i: 50, 1979c: 206.
- redikorzevi** Berger 1916: 232. Holotype ♀; S. Ussuri Terr., Sedanka; not located.
 Figures: Yin, Huang, & Li 1984: 100 (adult).
 Distribution: Asia (Sichuan in China/ Sakhalin Island, Siberia in E USSR).
 Hosts: *Abies fabri*, *A. holophylla*.
 References: (**cn**) Kurenzov 1935c: 191. (**cc**) Kurenzov 1934a: 51. (**hb**) Krivolutsкая 1956: 838; Kurenzov 1935a: 19, 32, 1948b: 102, 1950d: 144; Stark 1952: 257. (**ds**) Kleine 1934a: 142; Krivolutsкая 1956: 838, 1983; Kurenzov 1934a: 51, 1935a: 19, 32, 1935c: 191, 1936b: 350; Stark 1936e: 142, 1952: 257; Tsai & Li 1963; Yin, Huang, & Li 1984: 100. (**tx**) Berger 1916: 232; Eggers 1921: 39–40; Krivolutsкая 1956: 838, 1958: 142; Kurenzov 1941a: 149–150, 154, 1948b: 102, 1950: 144, 152; Michalski 1969a: 892, 1969b: 567; Schedl 1934f: 1639, 1952: 257; Sokanovskii 1954: 16; Stark 1936e: 142, 1952: 257; Tsai & Li 1963: 604, 622; Yin, Huang, & Li 1984: 100.
- resiniferi** Schedl 1943b: 36. Lectotype ♀; Benquet, Mt. Sta. Tomas (Philippines); Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 210. Distribution: Philippine Islands (Luzon).
 Hosts: *Pittosporum resiniferum*.
 References: (**ds**) Schedl 1966b: 22. (**tx**) Nobuchi 1983: 300; Schedl 1943b: 36, 1954a: 148, 1979c: 210.
- rhysi** Niisima 1909: 145. Holotype, sex?; Chitose in der Prov. Ishikari; Nobuchi Collection, Ibaraki.
 Distribution: Asia (Japan).
 Hosts: *Rhus toxicodendron* var. *radicans*.
 References: (**ds**) Kleine 1913b: 119, 1914b: 251, 1934a: 142; Murayama 1954b: 164, 1955: 103; Nobuchi 1985c: 15. (**tx**) Hagedorn 1910a: 88;

Murayama 1954b: 164, 1955: 103; Niisima 1909: 145; Schedl 1934f: 1639.

robustus Eichhoff 1872a: 131. Syntypes 5, sex?; Amer. bor. (?), Eichhoff 1878b: 122 gives America septentrionalis; IRSNB, Brussels.

Distribution: America septentrionalis (?).

Notes: The syntypes resemble *Hypocryphalus mangiferae* and are either that species or they were mislabeled or represent an interception. They were apparently among Dejean material with only a blank disk to indicate origin.

References: (ds) Gemminger & Harold 1872: 2683; Hagedorn 1910d: 44–45; Kleine 1913b: 119; Swaine 1909: 93. (tx) Eichhoff 1872a: 131, 1878b: 121; Hagedorn 1910a: 87; LeConte 1876: 362; Swaine 1909: 93.

rubentis Hopkins 1915b: 39–40. Holotype ♀; Pocahontas Co., West Virginia [USA]; USNM, Washington.

Distribution: North America (North Carolina, Pennsylvania, West Virginia in USA).

Hosts: *Picea rubens*.

References: (cn) Blackman 1950; Doane et al. 1936; Hopkins 1904a: 24. (hb) Baker, W. L. 1972: 259; Blackman 1950; Chamberlin 1939: 313; Doane et al. 1936; Hopkins 1904a: 24; Wood, S. L. 1953: 87, 1954a: 1004. (ds) Baker, W. L. 1972: 259; Blackman 1950; Blatchley & Leng 1916: 606; Chamberlin 1925, 1939: 313; Drooz 1985: 363; Kleine 1934a: 142; Wood, S. L. 1952b: 869. (tx) Blackman 1922c: 126–128; Blatchley & Leng 1916: 606; Chamberlin 1939: 313; Hopkins 1915b: 39–40; Wood, S. L. 1953: 87, 1954a: 959, 979, 1004, 1952b: 869.

ruficollis Hopkins 1915b: 39–40. Holotype ♀; Alta, Utah [USA]; USNM, Washington.

Figures: Bright 1976d: 202, 210, Kusch 1967: 10. Distribution: North America (Alaska/ Alberta, British Columbia, Manitoba, New Brunswick, Newfoundland, Nova Scotia, Ontario, Prince Edward Island, Quebec in Canada/ Nuevo Leon in Mexico/ Arizona/ California/ Colorado/ Idaho/ Maine/ Montana/ New Mexico/ New York/ Oregon/ Utah/ Washington/ Wyoming in USA).

Hosts: *Abies amabilis*, *A. balsamea*, *A. fraseri*, *A. grandis*, *A. magnifica*, *A. lasiocarpa*, *Picea engelmannii*, *P. glauca*, *P. rubens*, *Pseudotsuga menziesii*.

References: (bv) Furniss, M. M., Baker, & Hostetler 1976: 1300. (cn) Beckwith 1972; Dewey & Clinton 1980; Hopkins 1904a: 24; Smith, G. J. & Melvin 1974a, 1974b. (cc) Hertert, Miller, & Partridge 1975: 201; Tomalak, Welsh, & Garroway 1989b; Werner & Holsten 1984. (hb) Baker, W. L. 1972: 259; Bright & Stark 1973: 68; Chamberlin 1939: 314, 1958: 133–134; Hopkins 1904a: 24; Wood, S. L. 1953: 92, 1954a: 1005, 1952b: 870. (ds) Baker, W. L. 1972: 256; Beckwith 1972a; Bright 1971a: 125, 1976d: 113;

Bright & Stark 1973: 68; Chamberlin 1939: 314, 1958: 133–134; Drooz 1985: 363; Elias 1982c, 1985: 39; Evans, D., Lowe, & Hunt 1978; Furniss, M. M. 1978; Furniss, R. L. & Carolin 1977: 377; Cast et al. 1989: 385; Gautreau & Melvin 1974: 14; Hertert, Miller, & Partridge 1975: 201; Kusch 1967; Miller, R. F. & Morgan 1982; Miller, R. F., Morgan, & Hicock 1985: 500; Morgan, A. V. & Morgan 1980; Smith, G. J. & Melvin 1974a, 1974b; Still, Tidsbury, & Melvin 1974a: 14; Werner & Holsten 1984; Wood, S. L. 1948: 41, 1972a: 415, 1982b: 870. (tx) Beckwith 1972, 1972a; Bright 1976d: 113, 202, 210; Chamberlin 1939: 314, 1958: 133–134; Hopkins 1915b: 39–41; Kusch 1967: 10; Wood, S. L. 1953: 92, 1954a: 960, 1005, 1972a: 415, 1977d: 512, 1982b: 870, 1989: 171.

approximatus Hopkins 1915b: 40. Holotype ♀; Sandpoint, Idaho [USA]; USNM, Washington. Synonymy: Wood 1954a: 1005.

References: (cn) Doane et al. 1936; Swaine 1918a: 87. (hb) Chamberlin 1939: 315; Doane et al. 1936; Swaine 1918a: 87. (ds) Chamberlin 1925, 1939: 315; Keen 1929a: 29; Kleine 1934a: 142; Wood, S. L. 1951a: 128. (tx) Chamberlin 1939: 315; Hopkins 1915b: 39–41; Keen 1929a: 29; Swaine 1918a: 87; Wood, S. L. 1954a: 959, 1005.

fraseri Hopkins 1915b: 40–41. Holotype ♀; Pisgah Ridge, North Carolina [USA]; USNM, Washington. Synonymy: Wood 1989: 171.

References: (cn) Blackman 1950; Doane et al. 1936; Lindquist, O. H. & Syme 1981: 106. (hb) Baker, W. L. 1972: 259; Beal & Massey 1945: 119–120; Blackman 1950; Chamberlin 1939: 313; Doane et al. 1936; Wood, S. L. 1954a: 110, 1982b: 869. (ds) Amman 1969a; Baker, W. L. 1972: 259; Beal & Massey 1945: 119–120; Blackman 1950; Blatchley & Leng 1916: 607; Chamberlin 1925, 1939: 313; Drooz 1985: 363; Kleine 1934a: 142; Lindquist, O. H. & Syme 1981: 106; Wood, S. L. 1952b: 869. (tx) Beal & Massey 1945: 119–120; Blatchley & Leng 1916: 607; Chamberlin 1939: 313; Hopkins 1915b: 39–41; Lindquist, O. H. & Syme 1981: 106; Wood, S. L. 1954a: 960, 1010, 1952b: 869, 1989: 171.

balsameus Hopkins 1915b: 40. Holotype ♀; Camp Caribou, Maine [USA]; USNM, Washington. Synonymy: Wood 1954a: 1010.

References: (bv) Hosking & Knight 1975. (cn) Blackman 1919: 145, 1950; Doane et al. 1936; Swaine 1918a: 87, 89. (cc) Felt 1906: 673. (hb) Blackman 1950; Chamberlin 1939: 313; Doane et al. 1936; Felt 1906: 673; Swaine 1918a: 87, 89. (ds) Anonymous 1926c: 518; Beaulne 1956; Blackman 1950; Blatchley & Leng 1916: 607; Chamberlin 1925, 1939: 313; Dodge 1938: 15, 39; Hatch 1924; Kleine 1934a: 142; Leonard 1928: 515. (tx) Blackman

- 1922c: 126–128; Blatchley & Leng 1916: 607; Chamberlin 1939: 313; Dodge 1938; Hopkins 1915b: 40–41; Swaine 1918a: 87, 89; Titus, Meikle, & Harrison 1985: 51; Wood, S. L. 1954a: 101.
- grandis* Chamberlin 1917: 323. Lectotype ♀; Corvallis, Oregon [USA]; CNCI, Ottawa, designated by Bright 1967c: 681. Synonymy: Wood 1954a: 1005.
- References: (cn) Doane et al. 1936. (hb) Chamberlin 1939: 315; Doane et al. 1936. (ds) Chamberlin 1917: 323, 1925, 1939: 315; Keen 1929a: 29; Kleine 1934a: 142. (tx) Bright 1967c: 681; Chamberlin 1917: 323, 1939: 315; Keen 1929a: 29; Wood, S. L. 1954a: 1005.
- amabilis* Chamberlin 1917: 321. Lectotype ♀; Elk Lake, Oregon [USA]; CNCI, Ottawa, designated by Bright 1967c: 681. Synonymy: Wood 1954a: 1005.
- References: (cn) Chamberlin 1924; Doane et al. 1936. (cc) Chamberlin 1918a; Schedl 1958d: 190. (hb) Chamberlin 1939: 312, 1958: 134; Doane et al. 1936; Wood, S. L. 1954a: 1005. (ds) Chamberlin 1917, 1918a: 13, 1925, 1939: 312, 1958: 134; Keen 1929a: 29; Kleine 1934a: 142; Patterson & Hatch 1945: 152. (tx) Bright 1967c: 681; Chamberlin 1917: 321–322, 1939: 312, 1958: 134; Keen 1929a: 29; de Ruette 1970: 99; Wood, S. L. 1954a: 959, 1002, 1005, 1008, 1977d: 512.
- canadensis* Chamberlin 1918: 88. Holotype ♀; Roger's Pass, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Wood 1954a: 1006.
- References: (cn) Swaine 1918a: 87–88. (hb) Chamberlin 1939: 314; Swaine 1918a: 87–88. (ds) Chamberlin 1939: 314; Hopping 1922; Keen 1929a: 29; Kleine 1934a: 142. (tx) Chamberlin 1918: 87–88, 1939: 314; Keen 1929a: 29; de Ruette 1970: 99; Swaine 1918a: 87–88; Wood, S. L. 1954a: 1006.
- mainensis* Blackman 1922c: 126. Holotype ♀; Orono, Maine [USA]; USNM, Washington. Synonymy: Wood 1954a: 1006.
- References: (cn) Blackman 1950; Doane et al. 1936. (hb) Blackman 1950; Chamberlin 1939: 314; Doane et al. 1936. (ds) Anonymous 1926c: 518; Blackman 1922c: 126, 1950; Chamberlin 1925, 1939: 314; Leng & Mutchler 1927: 52; Leonard 1928: 518. (tx) Blackman 1922c: 126; Chamberlin 1939: 314; de Ruette 1970: 100; Schedl 1979c: 145; Wood, S. L. 1954a: 960, 1005.
- ruficollis coloradensis* Wood 1954a: 1008. Holotype ♀; 7 miles N Grand Canyon National Park, Arizona [USA]; SMUK, Lawrence. Synonymy: Wood, S. L. 1977d: 512.
- Notes: (3) This is an indistinct geographical race that probably does not warrant subspecies recognition.
- References: (hb) Wood, S. L. 1954a: 1008. (tx) Wood, S. L. 1954a: 1008, 1977d: 512.
- rufopilosus* Schedl 1942a: 174. Lectotype, sex?; Celebes; Schedl Collection in NHMW, Wien.
- Distribution: Indonesia (Celebes).
- References: (tx) Schedl 1942a: 174, 1979c: 215.
- rugosus* (Schedl) 1958b: 102 (*Ericryphalus*). Lectotype, sex?; Malaya, Selangor, Kepong; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 216–217.
- Distribution: Asia (Malaya).
- Hosts: *Shorca gysbertsiana*.
- References: (ds) Browne 1961c: 70. (tx) Schedl 1958b: 102, 1979c: 216–217.
- saltuarius* Weise 1891: 336. Syntypes?, sex?, Caucasus; not located.
- Figures: Grune 1979: 106, Yin, Huang, & Li 1984: 103 (adult).
- Distribution: Asia (Sichuan in China/ E USSR), Europe (Austria/ Bulgaria/ Czechoslovakia/ Denmark/ Finland/ Germany/ Hungary/ Italy/ Poland/ Sweden/ W USSR/ Yugoslavia).
- Hosts: *Picca asperata*, *P. excelsa*, *P. obovata*, *P. orientalis*.
- References: (ay) Escherich 1923b: 481, 597; Nusslin 1911a: 51. (bv) Grune 1979: 107; Nuorteva 1956c: 17; Nuorteva & Nuorteva 1968. (cn) Chorbazdzhevo 1929; Escherich 1923b: 481, 616; Esterberg 1959; Gabler 1955; Juntinen 1960: 24; Kailidis & Georgevits 1972; Koch 1913: 95; Kozikowsky 1929: 254; Kurenzov 1935c: 188; Marcu 1926c: 63; Nestertschuk 1930: 176; Nosek 1951: 106, 1952b: 98; Nuorteva & Nuorteva 1968; Nusslin 1913: 257; Pfeiffer 1949b: 147, 150, 1950c: 2; Pierce, W. D. 1917: 69; Rhumbler 1927: 312; Saalas 1949: 342, 372; Schimitschek 1937c: 52, 1955c: 82; Schwerdtfeger 1944a: 180; Tragardh 1938a: 12; Wachtl 1901: 381; Wardle 1929: 322. (cc) Annala 1977; Borodin 1967b; Heqvist 1963: 153; Kangas 1946b: 22; Karpinski 1932a: 103; Kleine 1908c: 186, 1944: 74; Kraemer 1950b: 379; Kurenzov 1934a: 54; Lundberg 1984; Nosek 1951: 106, 1952b: 98; Nuorteva 1956a: 17, 1957b: 68, 1968a, 1970, 1971; Nuorteva & Nuorteva 1968; Nusslin 1927: 300; Palmen 1946: 194; Pfeiffer 1923a: 331, 1932a: 16, 1949b: 147, 150, 1950c: 2, 1959: 5; Rhumbler 1922: 299; Ruhm 1955c: 176, 1956b: 3; Saalas 1917a: 18, 1928: 652, 1949: 342, 372; Schwerdtfeger 1944a: 180; Wichmann 1959: 412. (hb) Annala 1977; Beffa 1949; Charvat 1950; Chorbazdzhevo 1929; Escherich 1923b: 481, 616; Gabler 1955; Gornostaev 1916: 314; Karpinski & Strawinski 1948: 155; Knotek 1899b: 1, 1901: 565; Kraemer 1950b: 379; Kurenzov 1935a: 20, 32, 1948b: 110; Lengerken 1939: 63; Lindberg 1963: 243; Nuorteva 1956c: 17, 1968a, 1970; Nusslin 1898: 280, 1913: 257, 1927: 300; Pfeiffer 1941c: 5; Postner 1974: 430; Rhumbler 1922: 299, 1927: 312; Saalas

1913a: 69, 82, 1949: 342, 372; Schwerdtfeger 1944a: 180, 1981: 192; Spessivtsev 1923: 207, 1928a: 221; Stark 1952: 267; Tragardh 1939b: 158, 199; Tschorbadjiev 1929: 166; Wachtl 1901: 381. **(ds)** Bakke & Kvanne 1977; Beffa 1949; Borchert 1951; Brakman 1966b: 206; Bucking 1932; Buresh & Lazarov 1956; Butovitsch & Heqvist 1947; Charvat 1950; Chorbadzhievo 1924d, 1929; Eggers 1904; Endrodi 1958b; Escherich 1923b: 481, 597, 1932b; Esterberg 1959; Florov 1949: 84; Fuchs 1905a; Gornostaev 1917; Grune 1979: 107; Hagedorn 1910a: 45; Hansen, V. 1939, 1956; Hellen 1947; Helliesen 1916: 84; Herger 1985c: 91; Heyden, Reitter, & Weise 1891: 671, 1906: 711; Holdhaus & Deubel 1910: 160; Horion 1951; Kailidis & Georgevits 1972; Karpinski 1932a: 103, 1948a: 173, 1948b: 230; Karpinski & Strawinski 1948: 155; Kiefer et al. 1942: 528; Klefbeck & Sjöberg 1960: 230–231; Kleine 1912a: 263, 1913a: 35, 1913b: 119, 1934a: 142; Knotek 1899b: 1, 1901: 565; Krivolutskaya 1983; Kurenzov 1934a: 54, 1935a: 20, 32, 1935c: 188; Langhoffer 1915c: 157; Lindberg & Saris 1952: 59; Lomnicki 1913b: 148; Lucht 1987: 278; Lundberg 1974: 92; Marcu 1926c: 63; Munster 1928: 259; Numberg 1928b: 55, 105, 1954: 52; Nuorteva 1971: 67; Nusslin 1898: 280; Orest 1926c: 63; Palm 1946: 122; Palmén 1946: 194; Pfeffer 1924b: 471, 1931b: 75, 1947e: 5, 1950b: 75, 1989a: 82; Pierce, W. D. 1917: 69; Pittioni 1943: 176; Platonoff 1943: 141; Postner 1974: 430; Prossen 1913: 83; Rapp 1934: 727; Reitter 1894a: 70, 1916: 288; Ronbal 1941: 263; Saalas 1913a: 69, 82, 1917a: 18, 1919: 1–415, 1931: 68; Schaufuss 1915: 1231; Schedl 1967c: 70, 1980a: 15, 1981b: 74; Schilsky 1909: 188; Schwerdtfeger 1981: 192; Stark 1926d: 103, 1926j: 126, 1931a: 25, 1931d: 546, 1936e: 144, 1952: 267; Strand 1946: 599; Tragardh 1938: 10–14, 1939b: 158, 199; Tredl 1907: 12; Tsai & Li 1963: 602; Tschorbadjiev 1929: 166; Wegelius 1960: 106; Wichmann 1927a: 67; Wood, S. L. 1982b: 74; Yanovskii & Tegshzhargal 1985: 415; Yin, Huang, & Li 1984: 103; Zinovjev 1955: 187. **(tx)** Beeson 1941 (1961: 287); Beffa 1949; Charvat 1950; Chorbadzhievo 1924d; Eggers 1914: 188; Endrodi 1957a: 307, 1957b; Escherich 1923b: 481, 597; Fleischer 1927; Formanek 1907: 35; Gabler 1955; Grune 1979: 106–107; Hagedorn 1910a: 88; Hansen, V. 1956; Inouye & Nobuchi 1957a: 49; Kalina 1970: 125; Karpinski & Strawinski 1948: 155; Koch 1913: 95; Kurenzov 1941: 150–151, 1948b: 110; Lucht 1987: 278; Numberg 1954: 52; Nusslin 1911a: 51; Pfeffer 1932b: 17, 1941c: 5, 1947e: 5, 1955a: 178; Postner 1974: 430; Quaschik 1953: 35; Reitter 1894a: 70, 1913a: 67, 1916: 288; Rhumbler 1922: 299, 1927: 312; Saalas 1913a: 69, 82, 1916: 91–95, 110–116, 1949: 342, 372; Schedl 1934f: 1639, 1940d: 591, 1980a: 15, 1981b: 74; Schimitschek 1937c: 52, 1955c: 82; Sokanovskii

1929: 521–526; Spessivtsev 1922a: 472, 1923: 200–214, 1925a: 173, 1925b: 29, 1925d: 102–106, 1931: 49; Stark 1936e: 144, 1952: 267; Tsai & Li 1963: 605; Weise 1891: 336; Yin, Huang, & Li 1984: 103. **(ms)** Escherich 1932b; Kozikowsky 1929: 254; Wichmann 1959: 412, 1961: 332. *scriba* Gozis 1886: 31. Automatic.

Notes: This was presented as a replacement name for *asperatus* Ratzeburg 1837: 163, a misidentification of *asperatus* Gyllenhal that never had standing in nomenclature.

References: **(tx)** Gozis 1886: 31; Hagedorn 1910: 45; Saalas 1914: 304–306.

samoensis Beeson 1929: 224. Holotype ♂; Samoa. Upolu; Malololelei; BMNH, London.

Distribution: Samoan Islands.

Hosts: *Planchonella samoensis*.

References: **(hb)** Beaver 1976b: 534; Beeson 1938b: 257. **(tx)** Beaver 1976b: 534; Beeson 1929: 224–226, 1935: 107, 1938b: 257; Schedl 1951k: 133.

sarawakensis Schedl 1972m: 199. Holotype, sex?; Borneo, Sarawak, Mt. Serrisoon, 3000 ft.; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: **(tx)** Schedl 1972m: 199.

savadai Nobuchi & Takahashi 1965: 3. Holotype ♂; Shirotoge, Gunma pref.; Nobuchi Collection. Ibaraki.

Figures: Nobuchi 1967.

Distribution: Asia (Honshu in Japan).

Hosts: *Abies veitchii*.

References: **(ds)** Nobuchi 1985c: 15. **(tx)** Nobuchi 1967; Nobuchi & Takahashi 1965: 3.

scabricollis Eichhoff 1878b: 491. Holotype, sex?; Hindostan Asiae; Hamburg Museum, lost.

Distribution: Asia (Bangladesh/ Burma/ Hainan Island, Yunnan in China/ Andaman Islands, Assam, Bengal, Bihar, Madhya Pradesh, Maharashtra, Punjab, Tamil Nadu, Uttar Pradesh in India/ Sri Lanka/ Vietnam), Philippine Islands (Luzon).

Hosts: *Azalia bijuga*, *Albizia stipulata*, *Bassia latifolia*, *Bombax malabaricum*, *Buchanania lanzan*, *Canarium euphyllum*, *Excaecaria agallocha*, *Ficus* spp., *Garuga pinnata*, *Mesua ferrea*, *Odina wodier*.

Notes: (3) Beeson 1941 (1961: 287) (*Ericryphalus ficus*, nomen nudum, synonymy), *Cryphalus flabellifer* Sampson, nomen nudum, is this species. Schedl Collection had male *mimicus* Schedl, nomen nudum, under this name.

References: **(ds)** Bhasin, Roonwal, & Singh 1955; Hagedorn 1910d: 45; Kleine 1913b: 119, 1914b: 269, 1934a: 142; Schedl 1959a: 481. **(tx)** Eggers 1925: 152; Eichhoff 1878b: 491; Hagedorn 1910a: 88; Schedl 1958b: 100, 1959a: 481–482, 1962b: 186, 1975e: 446, 1979c: 220; Wood, S. L. 1989: 172, 1992b: 82.

discretus Eichhoff 1878b: 490. Holotype, sex?;

- Hindustan (Birma); Hamburg Museum, lost. Synonymy: Wood 1989: 172.
References: **(cn)** Roonwal 1954: 40. **(hb)** Beaver 1987a: 17; Beaver & Browne 1978: 584. **(ds)** Beaver 1987a: 17; Beaver & Browne 1978: 584; Beeson 1935: 107, 1961: 287; Browne 1961c: 69, 1984c: 449; Hagedorn 1910d: 41, 1913b: 118, 1914b: 278, 1934a: 142; Ohno, Yoshioka et al. 1988a: 91, 1989: 60; Roonwal 1954: 40. **(tx)** Eggers 1925: 152; Eichhoff 1878a: 385, 1878b: 490; Hagedorn 1910a: 86; Schedl 1958b: 99; Wood, S. L. 1989: 172.
- dilutus* Eichhoff 1878b: 490. Holotype, sex?; Hindostan (Birma); Dohrn Collection in Stettin Museum, now in Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 172. References: **(ds)** Beaver & Browne 1975: 288; Browne 1980b: 350, 1984a: 150, 1985a: 191; Hagedorn 1910d: 41; Kleine 1913b: 118, 1914b: 278; Ohno, Yoshioka, et al. 1988a: 91, 1989: 60; Schedl 1960h: 104. **(tx)** Eggers 1925: 152; Eichhoff 1878a: 384, 1878b: 490; Hagedorn 1910a: 86; Nobuchi 1983: 300; Schedl 1960h: 104, 1979c: 81; Wood, S. L. 1989: 172.
- brevisetosus* Schedl 1943b: 36. Lectotype ♀; Bagnio, Luzon, Philippinen; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 47. Synonymy: Wood 1992b: 82. References: **(ds)** Schedl 1959a: 481, 1966b: 19. **(tx)** Nobuchi 1983: 300; Schedl 1942a: 174, 1943b: 36, 1952b: 367, 1955b: 258, 1959a: 481, 1979c: 47; Wood, S. L. 1992b: 82.
- scabripennis* Schedl 1972m: 199. Holotype, sex?; Borneo, Sarawak, Mt. Poi, 4350 ft.; BMNH, London. Distribution: Indonesia (Sarawak in Borneo). References: **(tx)** Schedl 1972m: 199, 1979c: 220.
- scopiger* Berger 1917: 228. Lectotype ♂; S. Ussuri Terr., near Vladivostok; IZL, Leningrad, designated by Michalski 1969a: 892. Figures: Yin, Huang, & Li 1984: 105 (adult). Distribution: Asia (Heilongjiang in China/ Japan/ Korea/ Sakhalin Island, Siberia in E USSR). Hosts: *Padus asiatica*, *P. pubescens*, *Prunus padus*. References: **(cn)** Kurenzov 1935c: 189. **(ec)** Kurenzov 1934a: 50, 1951e: 66. **(hb)** Kurenzov 1935a: 19, 31, 1951e: 66; Stark 1952: 258. **(ds)** Krivolutsкая 1983; Krivolutsкая & Kupyanskaya 1970; Kurenzov 1934a: 50, 1935a: 19, 31, 1935c: 189, 1936a: 110, 1936b: 350, 1938a: 59; Nobuchi 1985c: 15; Stark 1936e: 143, 1952: 258. **(tx)** Berger 1916: 228; Eggers 1942c: 29–30; Kurenzov 1941a: 151–152, 157; Michalski 1969a: 892, 1969b: 567; Nobuchi 1964: 133, 1966e: 54; Schedl 1934f: 1639, 1962p: 203, 1979c: 222; Sokanovskii 1954: 16; Stark 1952: 258.
- padi* Krivolutsкая 1958: 144. Holotype, sex?; Sakhalin Island: Sakhalin region, Suani River valley; not located. Synonymy: Schedl 1962p: 203. References: **(hb)** Krivolutsкая 1973: 135. **(ds)** Krivolutsкая 1965a: 234, 1973: 135; Tsai & Li 1963: 602; Yin, Huang, & Li 1984: 105. **(tx)** Krivolutsкая 1958: 144; Schedl 1962p: 203, 1979c: 182; Tsai & Li 1963: 605, 623; Yin, Huang, & Li 1984: 105.
- securus* (Schedl) 1940b: 436 (*Ericryphalus*). Holotype, sex?; Dutch New Guinea, Cyclops Mts., Sabron, Camp 1, 1200 ft.; BMNH, London. Distribution: New Guinea. References: **(tx)** Schedl 1940b: 435–437, 1962r: 106, 1979c: 223.
- sejugatus* Schedl 1965: 54. Holotype, sex?; Madagascar, Montagne d'Ambre, 1000 m; Schedl Collection in NHMW, Wien. Distribution: Madagascar. References: **(tx)** Schedl 1965c: 54, 1977a: 41, 1979c: 223.
- sichotensis* Kurenzov 1941a: 152. Syntypes, sex?; Kolumbe River (middle Sikhote-Alin), Soviet Far East; ISB, Novosibirsk, USSR. Distribution: Asia (Hokkaido in Japan/ Sakhalin Island, E USSR). Hosts: *Picca jezoensis*. Notes: (3) Sokanovskii 1954: 16 (a possible synonym of *fulvus* Niisima). References: **(hb)** Stark 1952: 266. **(ds)** Arnoldi et al. 1955: 691; Krivolutsкая 1983; Kurenzov 1967; Stark 1952: 266. **(tx)** Kurenzov 1941a: 152–153; Sokanovskii 1954: 16; Stark 1952: 266.
- sidneyanus* (Nordlinger) 1856: 75 (*Bostrichus*). Holotype ♂; Australischen Holz; NHMW, Wien. Distribution: Australia. Notes: (3) Also cited under the generic name *Pseudocryphalus*. References: **(ds)** Gemminger & Harold 1872: 2683; Hagedorn 1910d: 45; Kleine 1913b: 119, 1914b: 298. **(tx)** Eggers 1931c: 185; Eichhoff 1878b: 136; Ferrari 1867a: 17, 75, 1868: 252; Hagedorn 1910a: 88; Nordlinger 1856: 75.
- silvanus* Schedl 1951k: 145. Lectotype ♂; Samoa, Upolu, Apia, Falefa Falls, Tapatapao 800 ft., Afiamalu 2200 ft., Tutuila, Utulei 500 ft., Fagatogo 700 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 229. Distribution: Samoan Islands. References: **(tx)** Beaver 1976b: 534; Schedl 1951k: 133, 145, 1958h: 498–499, 1979c: 229.
- similis* (Eggers) 1922b: 168 (*Stephanoderes*). Syntypes 6, sex?; S.O. Usagara und Nyerengere (Deutsch-Ostafrika); Methner Collection and 2 Eggers syntypes in NHMW, Wien. Distribution: Africa (Tanzania). Notes: (1) Schedl 1939i: 383 (to *Cryphalus*). References: **(tx)** Eggers 1922b: 168; Schedl 1939i: 383, 1961k: 457, 1979c: 230.

- simplex** (Schedl) 1939e: 341 (*Erieryphalus*). Syntypes 2, sex?; Malaya, Selangor: Kepong; BMNH, London, and Schedl Collection in NHMW, Wien. Distribution: Asia (Malaya). Hosts: *Arthrophyllum diversifolia*. References: (ds) Beeson 1961: 287; Browne 1961c: 68. (tx) Schedl 1939e: 341.
- sinoabietis** Tsai & Li 1963: 602, 604, 607, 622. Holotype ♂; Ssu Chuan, China; IZAS, Beijing. Figures: Tsai & Li 1963: 607, Yin, Huang, & Li 1984: 101 (admt). Distribution: Asia (Sichuan in China). Hosts: *Abies faxoniana*. References: (ds) Yin, Huang, & Li 1984: 100. (tx) Tsai & Li 1963: 602, 604, 607, 622; Yin, Huang, & Li 1984: 100.
- sinoabietis** *apienensis* Tsai & Li 1963: 602, 604, 607, 622. Holotype ♂; Ssu Chuan, China; IZAS, Beijing. References: (ds) Yin, Huang, & Li 1984: 101. (tx) Tsai & Li 1963: 602, 604, 607, 622; Yin, Huang, & Li 1984: 101.
- sordidus** (Nobuchi) 1975: 51 (*Taenioglyptes*). Holotype ♀; Mt. Hayachine, Iwate, Japan; Nobuchi Collection, Ibaraki. Figures: Nobuchi 1975: pl. 2, fig. 13. Distribution: Asia (Japan). References: (ds) Nobuchi 1985c: 15. (tx) Nobuchi 1975: 51–52.
- sparsepilosus** Schedl 1942c: 172. Holotype, sex?; Australien; Schedl Collection in NHMW, Wien. Distribution: Australia. References: (tx) Schedl 1942a: 175, 1942c: 172, 1979c: 233.
- squameus** Schedl 1943b: 38. Holotype, sex?; Borneo; Schedl Collection in NHMW, Wien. Distribution: Indonesia (Borneo). References: (tx) Schedl 1943b: 38, 1979c: 235.
- squamulosus** Strohmeier 1911b: 20. Holotype, sex?; Mindoro, Calapan, Philippine Islands; Entomological Collection, Bureau of Science, Manila, P.I., and IPKE, Eberswalde. Distribution: Philippine Islands (Mindoro). References: (ds) Kleine 1913b: 119, 1914b: 290; Schedl 1966b: 22. (tx) Nobuchi 1983: 300; Strohmeier 1911b: 20–21.
- striatulus** Browne 1981a: 127. Holotype, sex?; Buluma (New Britain) to Omazaki (Japan), imported; BMNH, London. Distribution: New Britain Island. References: (ds) Ohno, Yoneyama, & Nakazawa 1987: 93; Ohno, Yoshioka, et al. 1988a: 91, 1989: 60. (tx) Browne 1981a: 127.
- striatus** Browne 1974a: 66. Holotype, sex?; Fiji: Vanna Levu, Dreketi; BMNH, London. Distribution: Fiji Islands. Hosts: *Storekiella* sp.
- References: (ds) Browne 1974a: 64; Roberts 1976: 379. (tx) Browne 1974a: 66.
- strigilatus** Wichmann 1914c: 137. Holotype, sex?; Tana, Ostafrika; NHMW, Wien. Distribution: Africa (Ethiopia/ Kenya). References: (tx) Schedl 1966j: 467; Wichmann 1914c: 137.
- strigipennis** Schedl 1950f: 50. Lectotype, sex?; Viti Levu: Tholo-i-Suva, 500–1000 ft., Navai Mill, near Nandarivatu, 2500 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 235. Distribution: Fiji Islands. References: (ds) Browne 1974a: 64. (tx) Schedl 1950f: 50, 1979c: 235.
- strohmeieri** Stebbing 1914: 538. Syntypes 2, sex?; Jamsar, NW Himalaya, Chakrata, Uttar Pradesh, India; FRI, Dehra Dun, automatic. Distribution: Asia (Sichuan, Xizang [Tibet], Yunnan in China/ Bengal, Kashmir, Uttar Pradesh in India). Hosts: *Abies donsae*, *A. pindrow*, *A. webbiana*, *Pinus excelsa*. References: (cn) Pierce, W. D. 1917: 79; Roonwal 1954: 12; Stebbing 1914: 538. (ce) Stebbing 1914: 538. (hb) Stebbing 1914: 538. (ds) Beeson 1961: 286; Kleine 1934a: 142; Pierce, W. D. 1917: 79; Roonwal 1954: 12; Yin & Huang 1981: 561. (tx) Schedl 1939i: 350; Stebbing 1914: 538; Yin & Huang 1981: 561.
- indicus** Stebbing 1902: 403. Syntypes 2, sex?; Jamsar, NW Himalaya, Chakrata, Uttar Pradesh, India; FRI, Dehra Dun, preoccupied by Eichhoff 1878. References: (ds) Hagedorn 1910d: 43; Kleine 1913b: 118, 1934a: 142. (tx) Hagedorn 1910a: 87; Stebbing 1902: 403, 1914: 538.
- subcompactus** Lea 1910: 140. Syntypes, sex?; N.S. Wales: Galston, Sydney; Australia: SAM, Adelaide? Distribution: Australia (New South Wales, Queensland). Hosts: *Grevillea robusta*. References: (ds) Brimblecombe 1953: 17; Schedl 1972a: 144. (tx) Lea 1910: 140; Schedl 1938f: 47, 1948g: 25, 1979c: 240.
- subgranulatus** Schedl 1943b: 37. Holotype, sex?; S. Theodoro, Ins. Mindoro, Philippines; Schedl Collection in NHMW, Wien. Distribution: Philippine Islands (Mindoro). References: (ds) Schedl 1966b: 22. (tx) Nobuchi 1983: 300; Schedl 1943b: 37, 1979c: 241.
- submuricatus** Eichhoff 1878b: 492. Syntypes, sex?; Asiae India orientalis; Hamburg Museum, lost. Distribution: Asia (Burma/ Andaman Islands in India). References: (ds) Beeson 1929: 255–226; Hagedorn 1910d: 45; Kleine 1913b: 119, 1914b: 269. (tx) Eggers 1925: 152; Eichhoff 1878a: 355, 1878b: 492; Hagedorn 1910a: 85.

- substriatus** Schedl 1942a: 174. Lectotype, sex?; Malaya, Selangor. Sg. Buloh F. R.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 244.
Distribution: Asia (Yunnan in China/ Malaya).
Hosts: *Gluta elegans*, *Shorea sideroxylon*, *Sloetia elongata*.
References: (hb) Beaver & Browne 1978: 555; Browne 1961c: 70. (ds) Beaver & Browne 1978: 555; Browne 1961c: 70, 1984a: 151; Schedl 1959a: 482. (tx) Schedl 1942a: 174, 1953c: 288, 1959a: 482, 1979c: 244.
- subtuberculatus** Schedl 1942c: 168. Holotype, sex?; N.O. Papua, Mt. Lamington; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1942c: 168, 1979c: 245.
- subvestitus** Schedl 1959a: 482. Syntypes ♀; Ceylon: Millawitiya Estate; Schedl Collection in NHMW, Wien.
Distribution: Asia (Sri Lanka).
References: (ds) Schedl 1959a: 482. (tx) Schedl 1959a: 482, 1979c: 245.
- sundaensis** Schedl 1942d: 14. Lectotype, sex?; Java, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 247.
Distribution: Asia (Burma/ Thailand), Indonesia (Java).
References: (hb) Kalshoven 1958b: 165. (ds) Beaver & Browne 1975: 289; Schedl 1969c: 50. (tx) Schedl 1942d: 14, 1943b: 36, 1979c: 247.
- sylvicola** (Perkins) 1900: 181 (*Hypothenemus*). Syntypes, sex?; Lanai and Kauai, Hawaiian Islands; BMNH, London.
Distribution: Cook Islands, Fiji Islands, Hawaiian Islands, Micronesia (Caroline Islands, Guam, Mariana Islands), Nine Island, Philippine Islands, Samoan Islands.
Hosts: *Acacia* sp., *Artocarpus* sp., *Ficus* spp., *Ilex* sp., *Pinus insularis*, *Piper guahamense*.
References: (hb) Beaver 1976b: 534. (ds) Beaver 1976b: 534; Beeson 1938b: 289; Browne 1970: 540, 1974a: 64; Hagedorn 1910d: 45; Kleine 1913b: 124; Krauss 1943: 88; Schedl 1972b: 267; Swezey 1949: 446, 1954: 116; Wood, S. L. 1960a: 23. (tx) Browne 1970: 520; Hagedorn 1910a: 88; Perkins 1900: 181; Schedl 1934g: 177, 179, 1939c: 327, 1941f: 111, 1950f: 48, 1963i: 62; Swezey 1949: 446; Wood, S. L. 1960a: 23, 1989: 172.
- henshawi** Hopkins 1915b: 38 (*Eriocryphalus*). Holotype ♀; Hilo, Hawaii; USNM, Washington. Synonymy: Schedl 1941f: 111.
References: (ds) Beeson 1938: 288, 1939: 446; Ford 1960: 333; Swezey 1949: 446. (tx) Hopkins 1915b: 38; Schedl 1941f: 111, 1963i: 62; Swezey 1949: 446.
- pini** Hopkins 1915b: 39 (*Piperius*). Holotype ♀; Baguio, Philippine Islands; USNM, Washington. Synonymy: Schedl 1963i: 62.
References: (ds) Kleine 1934a: 148. (tx) Eggers 1923b; Hopkins 1915b: 39; Schedl 1963i: 62.
- swezeyi** Schedl 1942b: 147. Lectotype ♀; Guam: Dededo, Yigo, Ritidian, Mt. Alifan, Piti; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 248. Synonymy: Wood 1989: 172.
References: (ds) Swezey 1949: 446, 448. (tx) Schedl 1942b: 147–148, 1943b: 34, 1979c: 248; Wood 1960a: 23–25, 1989: 172.
- dimorphus** Schedl 1950f: 38. Lectotype ♀; Fiji: Vunua Mbalavu: Loma Loma, 200–500 ft., Myana, 200 ft., 700 ft., Ovalau near Vunua, 200 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 81. Synonymy: Beaver 1991: 89.
References: (ds) Browne 1974a: 64. (tx) Beaver 1991: 89; Schedl 1950f: 38, 49, 1979c: 81.
- sylvicola obliquus** Schedl 1950f: 48. Lectotype ♀; Fiji: Vunua Moalavu, Loma Loma; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 248. Synonymy: Wood 1960a: 23, 1989: 172.
References: (tx) Schedl 1950f: 48, 1951k: 133, 1979c: 248; Wood, S. L. 1960a: 23, 1989: 172.
- szechuanensis** Tsai & Li 1963: 605, 614, 624. Holotype ♂; Te Chamg, Ssu Chuan, China; IZAS, Beijing.
Figures: Tsai & Li 1963: 615, Yin, Huang, & Li 1984: 111 (adult).
Distribution: Asia (Sichuan in China).
Hosts: *Pinus yunnanensis*.
References: (ds) Yin, Huang, & Li 1984: 111. (tx) Tsai & Li 1963: 605, 614–615, 624; Yin, Huang, & Li 1984: 111.
- szechuanensis telchangensis** Tsai & Li 1963: 606, 615, 624. Holotype ♂; Te Chamg, Ssu Chuan, China; IZAS, Beijing.
References: (tx) Tsai & Li 1963: 606, 615, 624.
- tabulaeformis** Tsai & Li 1963: 605, 609, 623. Holotype ♂; Chin Huang Tao, Ho Pei, China; IZAS, Beijing.
Figures: Tsai & Li 1963: 609, Yin, Huang, & Li 1984: 106 (adult).
Distribution: Asia (Hebei in China, subspecies in Shanxi).
Hosts: *Pinus tabulaeformis*.
References: (ds) Yin, Huang, & Li 1984: 105. (tx) Tsai & Li 1963: 605, 609, 623; Yin, Huang, & Li 1984: 105.
- tabulaeformis chienzhuangensis** Tsai & Li 1963: 605, 610, 623. Holotype, sex?; China; IZAS, Beijing.
References: (ds) Yin, Huang, & Li 1984: 106. (tx) Tsai & Li 1963: 605, 610, 623; Yin, Huang, & Li 1984: 106.
- taivanus** Schedl 1970b: 359. Holotype, sex?; Formosa, Kaoshiung to Nagoya (Japan), imported; PPST, Tokyo.

- Distribution: Asia (Taiwan).
 Hosts: *Tsuga* sp.
 References: (tx) Schedl 1970b: 359, 1979c: 249.
- tenuis** Schedl 1942d: 16. Lectotype ♀; Java, Mount Gede, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 251.
 Distribution: Asia (Malaya), Indonesia (Java).
 Hosts: *Artocarpus elasticus*, *A. rigidus*, *A. scortechini*, *Campnosperma auriculata*, *Schefflera aromatica*.
 References: (ce) Beaver 1979b: 298. (hb) Beaver 1979b: 298; Beaver & Browne 1978: 585; Browne 1961c: 68. (ds) Beaver & Browne 1978: 585. (tx) Kalshoven 1958: 164; Schedl 1942c: 172, 1942d: 16, 1943b: 36, 1958b: 100, 1979c: 251.
- terminaliae** Browne 1980b: 384. Holotype, sex?; Pulau Adi (New Guinea) to Nagoya (Japan), imported; BMNH, London.
 Distribution: New Guinea.
 Hosts: *Terminalia* sp.
 References: (ds) Browne 1984b: 287; Ohno, Yoshioka, et al. 1988a: 91, 1989: 60. (tx) Browne 1980b: 384.
- tetricus** (Schedl) 1940b: 437 (*Ericryphalus*). Holotype, sex?; Dutch New Guinea, Cyclops Mts., Sabron, Camp 1, 1200 ft.; BMNH, London.
 Distribution: New Guinea.
 References: (ds) Schedl 1971a: 275. (tx) Schedl 1940b: 437–438, 1953b: 124–125, 1958b: 102, 1979c: 252.
- papuanus** Schedl 1942c: 170. Holotype ♂; N.O. Papua, Mt. Lamington; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1979g: 95.
 References: (tx) Schedl 1942c: 170, 1969b: 214, 1972q: 256, 1979c: 183, 1979g: 95.
- grayi** Schedl 1968e: 265. Holotype, sex?; New Guinea, Rille Range, L.A. Bulolo, Inakanda L.A. Bulolo; CSIRO, Canberra. Synonymy: Schedl 1979g: 95.
 References: (tx) Schedl 1968e: 265, 1979c: 112.
- brevis** Browne 1970: 556 (*Ptilopodius*). Holotype, sex?; New Guinea: Dorey; BMNH, London.
 Synonymy: Schedl 1979q: 256.
 References: (tx) Browne 1970: 556–557; Schedl 1972q: 256.
- theobromae** Nunberg 1956d: 208. Syntypes, sex?; West Africa, S. Ashanti; not located, automatic.
 Distribution: Africa (Ghana).
 Hosts: *Theobroma cacao*.
 References: (tx) Nunberg 1956d: 208.
- horridus** Graham 1908: 113. Syntypes, sex?; West Africa, S. Ashanti; not located, preoccupied by Eichhoff 1878.
 References: (tx) Graham 1908: 113; Hagedorn 1910a: 86; Nunberg 1956d: 208; Schedl 1960i: 104, 1961k: 462.
- trypanoides** Beeson 1935: 106. Holotype ♂?; Marquesas Islands: Uapou, Hakabetau Valley, 2600 feet; BMNH, London.
- Distribution: Hawaiian Islands, Marquesas Islands.
 Hosts: *Artocarpus altilis*.
 References: (ds) Beeson 1935b: 285, 1939: 106; Gemminger & Harold 1872: 2692; Van Zwaluwenburg 1956: 9. (tx) Beeson 1935: 106–107.
- trypanus** Sampson 1914: 383. Holotype ♂; Seychelles, Silhouette, from near Mont Pot-a-eau, Mahe, 1500 feet; BMNH, London.
 Distribution: Africa (Seychelles Islands), Asia (Yunnan in China).
 References: (hb) Beaver 1987a: 17. (ds) Beaver 1987a: 17; Sampson 1914: 382; Schedl 1969d: 11. (tx) Beeson 1935: 107; Sampson 1914: 382–383.
- tuberculatus** Schedl 1943b: 37. Holotype, sex?; Tenasserim; Schedl Collection in NHMW, Wien.
 Distribution: Asia (Burma).
 References: (tx) Schedl 1943b: 37, 1979c: 257.
- uapouensis** (Beeson) 1935a: 107 (*Ericryphalus*). Syntypes 3 ♀; Marquesas Islands: Uapou, Hakabetau Valley, 1000 feet; BMNH, London.
 Distribution: Marquesas Islands.
 References: (ds) Beeson 1935b. (tx) Beeson 1935a: 107, 1935b: 288; Schedl 1942c: 173.
- upoluensis** Schedl 1951k: 144. Lectotype ♂; Samoa, Tutuila, Fagasa Trail 800 ft., Matafao Trail 1300 ft., Upolu, Tapatapao; Lamtoo Trail 1200 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 261.
 Distribution: Samoan Islands.
 References: (tx) Beaver 1976b: 534; Schedl 1951k: 133, 144, 1979c: 261.
- variolosus** Schedl 1950f: 49. Syntypes 2, sex?; Viti Levu: Navai Mill, near Nandarivatu 2500 ft., Ovalau: near Vuma 700 ft.; BPBM, Honolulu, and Schedl Collection (2 syntypes) in NHMW, Wien.
 Distribution: Fiji Islands.
 References: (ce) Roberts 1976: 379. (hb) Roberts 1976: 379. (ds) Browne 1974a: 64; Roberts 1976: 379. (tx) Schedl 1950f: 49, 1951i: 57, 1979c: 264.
- vestitus** Blandford 1895a: 318. Holotype, sex?; Ceylon, Bogawantalawa; BMNH, London.
 Distribution: Asia (Sri Lanka).
 Hosts: *Callophyllum cuneifolium*, *Camellia sinensis*.
 References: (ds) Bhasin, Roonwal, & Singh 1958: Hagedorn 1910d: 46; Kleine 1913b: 119, 1914b: 273; Schedl 1959a: 482. (tx) Blandford 1895a: 318; Hagedorn 1910a: 88; Schedl 1959a: 482.
- viburni** Stark 1936: 143. Lectotype, sex?; Far East, Vladivostok, Schkotov; IZL, Leningrad, designated by Michalski 1969a: 892.
 Distribution: Asia (Shanxi in China/ E USSR).
 Hosts: *Viburnum dilatatum*.
 References: (hb) Stark 1952: 259. (ds) Kurenzov 1941: 153–154; Stark 1952: 259; Yin, Huang, & Li 1984: 104. (tx) Pfeffer 1944a: 131; Michalski 1969a: 892; Schedl 1952k: 158; Stark 1936: 143, 151, 1951: 229, 1952: 259; Tsai & Li 1963: 605–

- 606; Yin, Huang, & Li 1984: 104. (**ms**) Pfeffer 1944a: 131.
- viburni* Eggers 1942c: 30. Holotype, sex?; Ussuri; Eggers Collection (2 Eggers cotypes in NHMW, Wien, may include this holotype). Synonymy: Pfeffer 1942: 30.
Notes: (1) Michalski 1969a: 592 (designated a lectotype although Eggers' holotype designation appears valid).
References: (**cn**) Kurenzov 1935c: 189. (**cc**) Kurenzov 1934a: 57. (**hb**) Kurenzov 1935a: 19, 31, 1945b: 126. (**ds**) Krivolitskaya 1983; Kurenzov 1934a: 57, 1935a: 19, 31, 1935c: 189, 1936a: 110, 1936b: 350, 1938a: 59; Stark 1936e: 143, 151; Tsai & Li 1963: 602. (**tx**) Eggers 1942c: 30; Kurenzov 1941a: 153, 1945b: 126; Michalski 1969a: 592, 1969b: 567; Pfeffer 1942: 30, 1944: 131; Schedl 1952k: 155, 1979c: 267; Stark 1951: 155.
- vitiensis* Browne 1974a: 67. Holotype, sex?; Fiji: Viti Levu, Nausori Highlands; BMNH, London.
Distribution: Fiji Islands.
Hosts: *Dysoxylum* sp.
References: (**cc**) Roberts 1976: 350. (**hb**) Roberts 1976: 350. (**ds**) Browne 1974a: 64; Roberts 1976: 350. (**tx**) Browne 1974a: 67.
- walkeri* (Blandford) 1896b: 200 (*Cryptarthrum*). Syntypes 2, sex?; Damma I.; BMNH, London.
Distribution: Asia (Malaya/ Vietnam), Australia (Victoria), Indonesia (Damma, Java, Sumatra), Philippine Islands (Luzon, Mindanao, Mindoro).
Hosts: *Urostigma* sp.
References: (**cn**) Kleine 1932a: 302. (**hb**) Kleine 1932a: 302. (**ds**) Beaver & Browne 1978: 555; Browne 1966: 244, 1968a: 133; Hagedorn 1910d: 46; Kleine 1913b: 125, 1914b: 250, 1932a: 302, 1934a: 148; Schedl 1971f: 146. (**tx**) Blandford 1896b: 200; Browne 1968a: 132; Hagedorn 1910a: 83, 1912a: 340; Hopkins 1914: 119; Lucas 1920: 210; Nobuchi 1983: 300; Schedl 1963i: 62. (**ms**) Lucas 1920: 210.
- hagedorni* Eggers 1908c: 216 (*Coccotrypes*). Lectotype ♀; Java; USNM, Washington, designated by Anderson & Anderson. Synonymy: Schedl 1963i: 62.
Notes: (1) Eggers 1927b: 393 (to *Cryphalus*).
References: (**cn**) Kalshoven 1932: 252. (**hb**) Kalshoven 1955b: 165. (**ds**) Hagedorn 1910d: 68; Kalshoven 1932: 252, 1955b: 165; Kleine 1914b: 288, 1934a: 142; Schedl 1935g: 424, 1966b: 20. (**tx**) Anderson, W. H. & Anderson 1971: 14; Eggers 1908c: 216, 1927b: 393–394; Hagedorn 1910a: 94; Nobuchi 1983: 300; Schedl 1935g: 424, 1942a: 175, 1942d: 2, 13, 1950f: 48, 1951i: 50, 1958i: 214, 1958k: 153, 1963i: 61–62.
- jacanus* Schedl 1954a: 148. Lectotype, sex?; Java: Buitenzorg, 250 m, Semarang, 40 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 129. Synonymy: Schedl 1958k: 153.
References: (**tx**) Schedl 1954a: 139, 148–149, 1958k: 153, 1979c: 129.
- wapleri* Eichhoff 1872a: 131. Holotype, sex?; Australia; IRSNB, Brussels.
Distribution: Australia (Queensland), New Guinea.
Hosts: *Dysoxylum* sp., *Ficus carica*, *F. sp.*, *Malasia scandens*.
References: (**cc**) Wichmann 1955a: 96. (**ds**) Brimblecombe 1953: 17; Browne 1980b: 351, 1985b: 291; Gemminger & Harold 1872: 2683; Hagedorn 1910d: 46; Kleine 1913b: 119, 1914b: 295, 1934a: 142; Kuschel 1972; Ohno, Yoneyama, & Narazawa 1987b: 93; Ohno, Yoshioka, et al. 1988a: 92, 1989: 60; Wichmann 1955a: 96. (**tx**) Eichhoff 1872a: 131, 1878b: 129; Hagedorn 1910a: 88.
- mckeoi* Schedl 1972h: 61. Holotype, sex?; Laviana Village, M. Bay Dist.; CSIRO, Canberra. Synonymy: Schedl 1979j: 126.
References: (**tx**) Schedl 1972h: 61, 1979c: 151, 1979j: 126.
- yamaguchii* Inouye & Nobuchi 1957a: 55. Holotype, sex?; Soumkyo, Daisetsu-zen National Park, Hokkaido, Japan; Nobuchi Collection, Ibaraki.
Distribution: Asia (Hokkaido in Japan).
Hosts: *Abies sachalinensis*.
References: (**ds**) Nobuchi 1985c: 15. (**tx**) Inouye & Nobuchi 1957a: 48, 55.
- zimmermani* Schedl 1950f: 51. Syntypes ♀; Viti Levu: Navai Mill, near Nandarivatu, 2500 ft.; BPBM, Honolulu, and Schedl Collection in NHMW, Wien.
Distribution: Fiji Islands.
References: (**ds**) Browne 1974a: 64. (**tx**) Schedl 1950f: 51, 1979c: 271.

Genus *Margadillius* Hopkins

MARGADILLIUS HOPKINS 1915b: 8, 37. Type-species: *Margadillius margadillaonis* Hopkins, original designation.

Keys: Hopkins 1915b: 37.

Notes: (1) The status of this genus is doubtful; it may not be separable from *Cryphalus*.

References: (**ay**) Nobuchi 1969a: 56. (**hb**) Wood, S. L. 1986a: 91. (**ds**) Browne 1961c: 70–71; Schedl 1961k: 447, 1966b: 17. (**tx**) Hopkins 1915b: 8, 37; Nunberg 1956: 141; Schedl 1961k: 447; Wood, S. L. 1986a: 91.

carinatus Browne 1950b: 382. Holotype, sex?; Buhma (New Britain) to Nagoya (Japan), imported; BMNH, London.

Distribution: New Britain Island.

References: (**tx**) Browne 1980b: 382.

concentralis Schedl 1975f: 344. Holotype, sex?; Papua, upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 344, 1979c: 61.

confusus Hopkins 1915b: 38. Holotype ♀; Pagbilao, P.I.; USNM, Washington.

Distribution: Philippine Islands (Luzon).

Hosts: Tucuen tree.

References: (ds) Kleine 1934a: 148; Schedl 1966b: 17. (tx) Hopkins 1915b: 36–38; Nobuchi 1983: 300.

erythrinae Hopkins 1915b: 38. Holotype ♀; Calapan, P.I.; USNM, Washington.

Distribution: Philippine Islands.

References: (ds) Kleine 1934a: 148; Schedl 1966b: 17. (tx) Hopkins 1915b: 36, 37; Nobuchi 1983: 300.

fulvus Browne 1984b: 289. Holotype, sex?; Fakfak (West Irian) to Nagoya (Japan), imported; BMNH, London.

Distribution: New Guinea.

Hosts: *Terminalia* sp.

References: (tx) Browne 1984b: 289.

loranthus Schedl 1942d: 8. Lectotype, sex?; Java, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 142.

Distribution: Indonesia (Java).

Hosts: Loranthaceae seeds.

References: (hb) Kalshoven 1958b: 166. (ds) Browne 1986b: 334; Ohno, Yoshioka, et al. 1988b: 91. (tx) Schedl 1942d: 8, 1979c: 142.

magnus Browne 1984d: 88. Holotype, sex?; New Guinea: Mount Giluwe; BMNH, London.

Distribution: New Guinea.

Hosts: *Quintonia* sp.

References: (tx) Browne 1984d: 88.

margadilaonis Hopkins 1915b: 38. Holotype ♀; Pagbilao, P.I.; USNM, Washington.

Distribution: Philippine Islands (Luzon).

Hosts: Margadelaio log.

References: (ds) Kleine 1934a: 148; Schedl 1966b: 18. (tx) Hopkins 1915b: 36–38; Nobuchi 1983: 300.

minor Schedl 1942a: 176. Lectotype, sex?; Malaya, Selangor, Kepong; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 155.

Distribution: Asia (Malaya), Indonesia (Java).

Hosts: *Hibiscus macrophyllus*.

References: (hb) Browne 1961c: 71. (tx) Schedl 1942a: 176, 1979c: 155.

minutus Hopkins 1915b: 37. Holotype ♀; Pagbilao, P.I.; USNM, Washington.

Distribution: Philippine Islands.

Hosts: Tucuen tree.

References: (ds) Kleine 1934a: 148; Schedl 1966b: 18. (tx) Hopkins 1915b: 36–37; Nobuchi 1983: 300.

parvulus Eggers 1943e: 75. Holotype, sex?; Zambezi (Nova Choupanga pres Chemba); Eggers Collection, in NHMW, Wien.

Figures: Schedl 1961k: 447.

Distribution: Africa (Mozambique/ South Africa).

Hosts: *Hibiscus* sp.

References: (ds) Schedl 1961k: 447. (tx) Eggers 1943e: 75; Schedl 1961k: 447.

quadrifuberculatus Schedl 1963i: 64. Holotype, sex?; Java, Batoerraden; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java).

References: (tx) Schedl 1963i: 64–65, 1979c: 186.

terminaliae Browne 1984c: 450. Holotype, sex?; Vini Harbour (Solomon Islands) to Nagoya (Japan), imported; BMNH, London.

Distribution: Solomon Islands.

Hosts: *Terminalia* sp.

References: (ds) Browne 1985a: 191; Ohno, Yoneyama, & Nakazawa 1987b: 93; Ohno, Yoshioka, et al. 1988a: 91. (tx) Browne 1984c: 450.

Genus *Ptilopodius* Hopkins

PTILOPODIUS HOPKINS 1915b: 7, 11. Type-species: *Ptilopodius stepheguis* Hopkins, original designation.

Keys: Wood 1960a: 18 for Micronesia.

References: (hb) Wood, S. L. 1986a: 91. (ds) Beeson 1935: 101, 1938: 290; Brimblecombe 1953: 23; Browne 1961c: 79; Schedl 1961k: 454, 1966b: 18; Wood, S. L. 1986a: 91. (tx) Hopkins 1915b: 7, 11–12; Schedl 1953c: 294, 1961k: 454, 1962r: 93; Wood, S. L. 1960a: 18, 1986a: 91.

aitutakii Beaver & Maddison 1990: 1367. Holotype ♀; Cook Islands: Aitutaki Island, Angari islet; Auckland Museum.

Distribution: Cook Islands.

References: (tx) Beaver & Maddison 1990: 1367.

bambusae Browne 1984f: 69. Holotype, sex?; New Guinea: Morobe District, Wau, 1200 m; BMNH, London.

Distribution: New Guinea.

Hosts: Bamboo.

References: (tx) Browne 1984f: 69.

dubiosus Wood 1960a: 19. Holotype ♀; Caroline Islands, Palau Islands, Peleliu; USNM, Washington.

Distribution: Micronesia (Palau Islands in Caroline Islands).

Hosts: *Pandanus* sp.

References: (ds) Browne 1984a: 150; Ohno, Yoshioka, et al. 1987a: 87. (tx) de Rnette 1970: 111; Wood, S. L. 1960a: 19.

formosanus Browne 1981a: 129. Holotype, sex?; Hualien (Formosa) to Yatsushiro (Japan), imported; BMNH, London.

Distribution: Asia (Taiwan).

References: (tx) Browne 1981a: 129.

javanus Schedl 1942d: 10. Lectotype, sex?; Java, Gedeh; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 129.

- Distribution: Indonesia (Java).
References: (hb) Kalshoven 1955b: 166. (tx) Schedl 1942d: 10, 1943b: 34, 1979c: 129.
- madagascariensis** Schedl 1961e: 133. Holotype, sex?; Madagascar, Montagne d'Ambre, Joffreville; IRSM, Madagascar.
Distribution: Madagascar.
References: (hb) Schedl 1977a: 44. (ds) Schedl 1977a: 44. (tx) Schedl 1961e: 133, 1962r: 93, 1977a: 44, 1979c: 144.
- marquesanus** Beeson 1935: 101. Syntypes 26, sex?; Marquesas Is.: Eiao, Hivaoa; Tahauku, Uapou, Hakagetan Valley; BPBM, Honolulu?
Distribution: Cook Islands (Aitutaki), Fiji Islands, Gambier Islands, Marquesas Islands, Samoan Islands (American Samoa, Western Samoa).
Hosts: *Hibiscus tiliaceus*.
References: (ds) Beeson 1935b: 290, 1940: 193; Schedl 1972b: 266. (tx) Beaver 1991: 93; Beeson 1935: 101–102, 1938b: 290, 1940: 193; Schedl 1951k: 143.
- zimmernani** Schedl 1951k: 143. Syntypes, sex?; Tutuila, west side; Afono Trail; Amonli 400 ft.; Fagatogo 700 ft.; Samoa, Upolu; Tapatapao; BPBM, Honolulu, and Schedl Collection in NHMW, Wien. Synonymy: Beaver 1991: 93.
References: (tx) Beaver 1991: 93; Schedl 1951k: 134, 143, 1953c: 294.
- minutissimus** Schedl 1943b: 34. Lectotype, sex?; Mt. Isarog, Luzon, Philippines; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 156.
Distribution: Philippine Islands (Luzon).
References: (hb) Kalshoven 1955b: 166. (ds) Schedl 1966b: 18. (tx) Nobuchi 1983: 300; Schedl 1943b: 34, 1951i: 50, 1954a: 139, 1979c: 156.
- pacificus** Schedl 1941f: 111. Holotype, sex?; Oahu; Waimano; BPBM, Honolulu?
Distribution: Micronesia (Kusaie, Marshall Islands, S Mariana Islands), Hawaiian Islands.
References: (tx) Browne 1970: 556; Schedl 1941f: 111, 1951m: 105, 1979c: 182; Wood, S. L. 1960a: 20.
- ramosus** Beeson 1935: 115. Holotype ♀; India: Bengal, Sunderbans; FRI, Dehra Dun.
Figures: Maiti & Saha 1956: 170 (adult, antenna).
Distribution: Asia (Bengal, Nicobar Islands in India), Mangareva Islands, Micronesia (Caroline Islands, Palau Islands, Ponape, Society Islands, Truk Island, Yap Island).
Hosts: *Hibiscus tiliaceus*.
References: (cn) Mathur & Singh 1960b: 5. (cc) Beeson 1922c: 498. (hb) Beeson 1922c: 498. (ds) Beeson 1922c: 498, 1938b: 290, 1940: 193, 1961: 295; Kleine 1934a: 148; Maiti & Saha 1956: 169; Mathur & Singh 1960b: 5; Wood, S. L. 1960a: 20. (tx) Beeson 1935: 115–116, 1940: 193; Maiti & Saha 1956: 169; Wood, S. L. 1960a: 20.
- rugosus** Schedl 1943b: 34. Lectotype, sex?; Manila, Philippines; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 217.
Distribution: Philippine Islands (Luzon).
References: (ds) Schedl 1966b: 18. (tx) Nobuchi 1983: 300; Schedl 1943b: 34, 1979c: 217.
- shoreae** Schedl 1953c: 293. Lectotype, sex?; Malaya, Selangor, Kepong; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 227.
Distribution: Asia (Malaya).
Hosts: *Shorea macroptera*.
References: (ds) Browne 1961c: 79. (tx) Schedl 1953c: 293, 1979c: 227.
- squamosus** Schedl 1953c: 294. Lectotype, sex?; Malaya, Selangor, Kepong; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 236.
Distribution: Asia (Malaya).
Hosts: *Shorea gylbertiana*, *S. parvifolia*.
References: (hb) Browne 1961c: 79. (tx) Schedl 1953c: 294.
- stephegynis** Hopkins 1915b: 11. Holotype ♀; Calapan, P.I.; USNM, Washington.
Distribution: Philippine Islands.
Hosts: *Stephegyne diversifolia*.
References: (ds) Beeson 1938: 290; Kleine 1934a: 148; Schedl 1959a: 475, 1966b: 18. (tx) Hopkins 1915b: 11; Nobuchi 1983: 300; Schedl 1933d: 201, 1959a: 475, 1961e: 133.
- tarawai** Beaver 1990: 149. Holotype ♀; Kiribati: S. Tarawa I.; Auckland Museum.
Distribution: Tarawa Island.
Hosts: *Scavola* sp.
References: (tx) Beaver 1990: 149.
- venustus** Schedl 1953d: 75. Syntypes 2, sex?; Madagascar, Mt. d'Ambre; MNHN, Paris and Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
References: (tx) Schedl 1953d: 78–79, 1962r: 93, 1977a: 44, 1979c: 265.

Genus *Cosmoderes* Eichhoff

- COSMODERES EICHHOFF 1878b: 495. Type-species: *Cosmoderes monilicollis* Eichhoff, monobasic.
- Erioschidius* Schedl 1935k: 42. Type-species: *Cryphalus setistriatus* Lea, subsequent designation by Wood 1960a: 21. Synonymy: Wood 1950c: 91.
References: (ds) Brimblecombe 1958: 18. (tx) Nunberg 1956a: 141; Schedl 1935k: 42, 1957d: 1–162, 1959a: 474, 1961k: 444, 1962r: 85–107, 1977b: 36; Wood, S. L. 1960a: 21, 1980c: 91.
- Dendriops* Schedl 1953b: 125. Type-species: *Dendriops granulicollis* Schedl, monobasic. Synonymy: Wood 1986a: 91.
References: (tx) Schedl 1953b: 125; Wood, S. L. 1986a: 91.
- Pseudocosmoderes* Nobuchi 1981c: 16. Type-species: *Pseudocosmoderes attenuatus* Nobuchi

probably = *Cosmoderes monilicollis* Eichhoff, monobasic. Synonymy: Wood 1992b: 80–81.

References: (tx) Nobuchi 1981c: 16–17; Wood, S. L. 1992b: 80–81.

Vitaderes Beeson 1941 (1961: 301). Type-species: *Vitaderes huffae*, nomen nudum, = *Cosmoderes monilicollis* Eichhoff, nomen nudum, no status.

References: (tx) Beeson 1941 (1961: 301).

References: (hb) Schedl 1977a: 36; Wood, S. L. 1986a: 91. (ds) Schedl 1961k: 444, 1966b: 17, 1977a: 36; Wood, S. L. 1986a: 91. (tx) Eichhoff 1878b: 475, 495; Hopkins 1915b: 7, 10–11; Nunberg 1956a: 141; Schedl 1961k: 444, 467, 1962r: 88, 1977a: 36; Wood, S. L. 1954a: 1050, 1986a: 91.

anticus (Schedl) 1975f: 343 (*Erioschidias*). Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 343, 1979c: 22.

consobrinus Blandford 1894d: 86. Holotype ♀?; Japan: BMNH, London.

Distribution: Asia (Japan).

References: (ds) Blandford 1894c: 579; Hagedorn 1910d: 46; Kleine 1913b: 125, 1914b: 257; Murayama 1949c: 101, 1961: 31. (tx) Blandford 1894d: 86; Hagedorn 1910a: 81; Schedl 1934f: 1640, 1963i: 61.

corpulentus (Schedl) 1971f: 151 (*Erioschidias*). Holotype, sex?; Philippinen, Palawan: Brooke's Point, Uring Uring; UZMC, Copenhagen.

Distribution: Philippine Islands (Palawan).

References: (tx) Schedl 1971f: 151–152.

cylindricus (Schedl) 1962r: 103 (*Erioschidias*). Holotype, sex?; Fiji Inseln; Schedl Collection in NHMW, Wien.

Distribution: Fiji Islands.

References: (ds) Browne 1974a: 64. (tx) Schedl 1962r: 103, 1979c: 73–74.

donisi (Schedl) 1957d: 46 (*Erioschidias*). Holotype, sex?; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Cynometra hankei*.

References: (tx) Schedl 1957d: 46, 1979c: 83.

elegans Schedl 1975f: 342. Holotype, sex?; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 342, 1979c: 88.

elongatus (Eggers) 1939b: 4 (*Erioschidias*). Holotype, sex?; Nordost-Birma (Kambaiti, 7000 Fuss); NHR, Stockholm.

Distribution: Asia (Burma).

References: (tx) Eggers 1939b: 4–5.

frontalis (Wood) 1960a: 21 (*Erioschidias*). Holotype ♂; Ulehsehel (Aurapushekarn) Is., Palau Islands; USNM, Washington.

Figures: Wood 1960a: 22.

Distribution: Micronesia (Palau Islands, Truk in Caroline Islands).

References: (tx) de Ruetten 1970: 101; Wood, S. L. 1960a: 21–22.

imitatrix (Schedl) 1977f: 499 (*Erioschidias*). Holotype, sex?; Vietnam: Yen So, SW of Hanoi; Schedl Collection in NHMW, Wien.

Distribution: Asia (Vietnam).

References: (tx) Schedl 1977f: 499.

madagascariensis (Schedl) 1963i: 65 (*Enptilius*). Syntypes, sex?; Madagascar: Mont. d'Ambre; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

Notes: (1) Schedl 1979c: 144 (citation of holotype invalid). (3) The syntype in NHMW, Wien, is largely covered by glue; identity uncertain.

References: (tx) Schedl 1963i: 65–67, 1979c: 144.

monilicollis Eichhoff 1878b: 496. Holotype, sex?; Hindostan; Hamburg Museum, lost.

Distribution: Asia (Burma/ Guizhou in China/ India/ Sri Lanka/ Vietnam), Indonesia (Sumatra). Hosts: *Actinidia polygama*, *Akebia trifoliata*, *Lagenaria vulgaris*, *Luffa aegyptiaca*, *Stemtonia hexaphylla*.

References: (ds) Blandford 1895a; Hagedorn 1910d: 46; Kleine 1913b: 125, 1914b: 269; Schedl 1959a: 476. (tx) Eggers 1925: 153; Eichhoff 1878a: 387, 1878b: 496; Hagedorn 1910a: 81; Hopkins 1914: 119; Lucas 1920: 205; Schedl 1959a: 476, 1963i: 60; Wood, S. L. 1992b: 81. (ms) Lucas 1920: 205.

granulicollis Schedl 1953b: 125 (*Dendriops*).

Lectotype, sex?; Indo-China: Saigon; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 111. Synonymy: Schedl 1963i: 60. References: (ds) Schedl 1965a: 339. (tx) Schedl 1953b: 125, 1963i: 60, 1979c: 111.

coriaceus Schedl 1959a: 474 (*Erioschidias*). Syntypes, sex?; Ceylon: Sabargamuva, Millawitiya Estate; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1963i: 60.

Notes: (1) Schedl 1979c: 65 (citation of holotype invalid).

References: (ds) Schedl 1959a: 474, 1971a: 280. (tx) Schedl 1959a: 474, 1961e: 69, 1962r: 103, 1963i: 60, 1979c: 65.

attenuatus Nobuchi 1981c: 17 (*Pseudocosmoderes*). Holotype, sex?; Shimizucho, Wakayama [Japan]; Nobuchi Collection, Ibaraki. Synonymy: Wood 1992b: 81.

Notes: Nobuchi's photograph of the type indicates that it is a female of *monilicollis*; confirmation is needed (SLW).

References: (tx) Nobuchi 1981c: 17–18; Wood, S. L. 1992b: 80–81.

niger (Schedl) 1961e: 131 (*Erioschidias*). Holotype, sex?; Madagascar, Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Erythroxylon sphaeranthrum*.

References: (tx) Schedl 1961e: 131, 1977b: 37, 1979c: 166.

papuanus Schedl 1975f: 341. Holotype, sex?; Peria Creek, Kwagira River, 50 m; AMNH, New York. Distribution: New Guinea.

References: (tx) Schedl 1975f: 341.

pellitus (Schedl) 1953d: 79 (*Erioschidias*). Lectotype, sex?; Madagascar, Mt. d'Ambre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 187.

Distribution: Madagascar.

References: (tx) Schedl 1953d: 79, 1961e: 131, 1977a: 37, 1979c: 187.

philippinensis (Schedl) 1967a: 126 (*Erioschidias*). Holotype, sex?; Bislig, Mindanao, Philippines to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Philippine Islands (Mindanao).

References: (tx) Nobuchi 1983: 300; Schedl 1967a: 126–127, 1979c: 192.

queenslandi (Schedl) 1938f: 43 (*Erioschidias*). Lectotype, sex?; Cairns district; Schedl Collection in NHMW, Wien.

Distribution: Australia (Queensland).

References: (ds) Brimblecombe 1953: 18. (tx) Schedl 1938f: 43, 1979c: 209.

ruandae (Schedl) 1962r: 104 (*Erioschidias*). Holotype, sex?; Ruanda; Ihembe; Schedl Collection in NHMW, Wien.

Distribution: Africa (Ruanda in Zaire).

References: (tx) Schedl 1962k: 1067, 1962r: 104, 1979c: 213.

sericeus (Schedl) 1942d: 10 (*Erioschidias*). Lectotype ♂; Java, Mount Gede, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 225.

Distribution: Indonesia (Java).

References: (hb) Kalshoven 1955b: 166. (tx) Schedl 1942d: 10, 1958k: 153, 1979c: 225.

kalshoveni Schedl 1954a: 149 (*Ptilopodius*). Lectotype ♀; Java, Mt. Gede, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 131. Synonymy: Schedl 1958k: 153.

References: (ds) Schedl 1965g: 23. (tx) Schedl 1954a: 139, 149, 1958k: 153, 1979c: 131.

setistriatus (Lea) 1910: 141 (*Cryphalus*). Syntypes, sex?; W. Australia; Rottnest Island; SAM, Adelaide? Distribution: Australia (Western Australia).

Notes: (1) Schedl 1938f: 42 (to *Erioschidias*, =*Cosmoderes*, redescribed).

References: (ds) Brimblecombe 1953: 18. (tx) Lea 1910: 141; Schedl 1936g: 528, 1938f: 42, 1959a: 474.

solitarius (Schedl) 1957d: 46 (*Erioschidias*). Holotype, sex?; Congo Belge; Yangambi; MRCB, Tervuren.

Figures: Schedl 1961k: 445–446.

Distribution: Africa (Zaire).

Hosts: *Dialium corbisieri*.

References: (hb) Schedl 1961k: 445. (ds) Schedl 1961k: 445. (tx) Schedl 1957d: 46, 1961k: 445–446, 1979c: 232.

Genus *Cryptocareus* Eggers

CRYPTOCAREUS EGGER 1937a: 79. Type-species: *Cryptocareus diadematus* Eggers, original designation.

Tachyderes Blackman 1943b: 35. Type-species: *Tachyderes floridensis* Blackman = *Cryptocareus scriatus* Eggers, original designation. Synonymy: Schedl 1951m: 72.

References: (tx) Blackman 1943b: 35; Schedl 1951m: 72, 1962p: 205; Wood, S. L. 1954a: 1011.

Keys: Wood 1982b: 912 for North and Central America.

References: (ay) Nobuchi 1969a: 57. (bv) Schedl 1960f: 11. (hb) Wood, S. L. 1982b: 911, 1986a: 91. (ds) Schedl 1962k: 23; Wood, S. L. 1982b: 911, 1986a: 91. (tx) Arnett 1960: 1043, 1968: 1043; Bright 1972d: 62; Eggers 1933b: 10, 1937a: 79–80; Schedl 1939a: 343, 1939h: 46–47, 1951m: 72, 1958f: 33–46, 1960f: 11, 34, 1962j: 23, 1962p: 205, 1965e: 349–379; Wood, S. L. 1954a: 981, 1011, 1982b: 911, 1986a: 91.

acaciae Schedl 1958f: 45. Syntypes, sex?; Argentinien: Buenos Aires, Tigre; Schedl Collection in NHMW, Wien, and Viana Collection.

Distribution: South America (Argentina).

Hosts: *Acacia nigra*.

Notes: (1) Schedl 1979c: 9 (citation on holotype invalid).

References: (hb) Viana 1964: 125. (ds) Numberg 1972b: 193; Viana 1964: 125. (tx) Schedl 1951m: 96, 1958f: 45, 1979c: 9.

brasilienis (Schedl) 1951m: 96 (*Miocryphalus*). Holotype, sex?; Brasilien, Bahia; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1951m: 96, 1979c: 45.

brericollis Eggers 1937a: 81. Holotype ♀; Bolivien (Cochabamba); Eggers Collection, in NHMW, Wien.

Distribution: South America (Bolivia/ Brazil/ Venezuela).

Hosts: *Nectandra* sp.

References: (ds) Blackwelder 1947: 781. (tx) Eggers 1937a: 81; Schedl 1939h: 47, 1979c: 46; Wood, S. L. 1976a: 347.

coronatus Wood 1971d: 36. Holotype ♀; 20 km SW El Vigia, Merida, Venezuela; Wood Collection. Synonymy: Wood 1976a: 347.

References: (hb) Wood, S. L. 1971d: 36. (ds) Beaver 1974a. (tx) Wood, S. L. 1971: 36, 1976a: 347.

- diadematus** Eggers 1937a: 80. Holotype ♀; Brasilien (Corumbá, Matto Grosso); USNM, Washington.
Distribution: Antilles Islands (Jamaica), North America (Costa Rica/ Oaxaca in Mexico/ Panama), South America (Brazil/ Colombia/ Venezuela).
Hosts: *Persca americana*, *Saraca indica*, *Serjania* sp.
References: (ds) Beaver 1974; Blackwelder 1947: 781; Pedrosa-Macedo & Schonherr 1985: 49; Schedl 1970e: 82; Wood, S. L. 1982b: 912. (tx) Anderson, W. H. & Anderson 1971: 11; Eggers 1937a: 80; Pedrosa-Macedo & Schonherr 1985: 49; Schedl 1939b: 47, 1979c: 79; Wood, S. L. 1954a: 1012, 1982b: 912.
- harringtoni** (Blackman) 1943b: 38 (*Tachyderes*). Holotype ♀; Yaguajay, Bolivia; USNM, Washington.
Distribution: South America (Bolivia).
References: (ds) Nunberg 1955a: 480. (tx) Blackman 1943b: 38; Nunberg 1955a: 480; Schedl 1962p: 205.
- hevae** (Hagedorn) 1912a: 335 (*Stephanoderes*). Lectotype ♀; Eala, Congo; MRCB, Tervuren, designated by Wood 1982b: 914.
Figures: Schedl 1962j: 24–25.
Distribution: Africa (Ghana/ Zaire), Antilles Islands (Cuba/ Guadeloupe/ Jamaica/ Virgin Islands), North America (Costa Rica/ Colima, Veracruz in Mexico/ Panama/ Florida in USA), South America (Brazil/ Colombia/ Venezuela).
Hosts: *Canavalia villosa*, *Vitis* sp.
References: (ay) Van Ryn-Tournel 1975. (bv) Schedl 1960f: 71. (cn) Auhmann 1912: 55; Ghesquiere 1933a: 33–36, 1933b: 783; Green 1916: 608–636; Mayne & Donis 1951: 334; Schmitz & Crisinel 1957: 12. (cc) Equihua & Atkinson 1986: 630; Mayne & Donis 1951: 334. (hb) Beaver 1974a; Browne 1963a: 235; Equihua & Atkinson 1986: 630; Kalshoven 1963; Schedl 1962k: 24; Schmitz & Crisinel 1957: 12. (ds) Atkinson & Equihua 1985c: 358; Beaver 1974a; Bright 1972d: 62, 1985c: 174; Browne 1963a: 235; Equihua & Atkinson 1986: 630; Estrada & Atkinson 1988: 209; Ghesquiere 1933c; Kalshoven 1963; Kleine 1913b: 121, 1914b: 314; Mayne 1915: 577–596; Pedrosa-Macedo & Schonherr 1985: 50; Schedl 1960f: 34–94, 1962k: 24, 1964j: 41, 1967e: 214, 1970e: 82; Wichmann 1954: 525; Wood, S. L. 1957c: 396, 1977a: 69, 1982b: 914. (tx) Bright 1972d: 62, 1985c: 174; Eggers 1924: 110, 1927a: 197, 1932c: 23, 1937a: 80, 1940h: 61; Hagedorn 1912a: 338, 1913a: 18; Pedrosa-Macedo & Schonherr 1985: 50; Schedl 1938a: 173, 1938i: 26, 1939h: 47, 1962j: 24–25; Wood, S. L. 1954a: 1014, 1957c: 396, 1962: 78, 1975b: 393, 1982b: 914.
- caraibicus** Eggers 1937a: 82. Holotype ♀; Guadeloupe; USNM, Washington. Synonymy: Wood 1975b: 393.
References: (ds) Blackwelder 1947: 781. (tx) Anderson, W. H. & Anderson 1971: 9; Eggers 1937a: 82; Schedl 1938a: 162, 183, 1939h: 47, 1979c: 53; Wood, S. L. 1975f: 392.
- parvus** Blackman 1943b: 36 (*Tachyderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1962: 78.
References: (tx) Blackman 1943b: 36; Schedl 1962p: 205; Wood, S. L. 1962: 78.
- porosus** Wood 1954a: 1014. Holotype ♀; Royal Palm Hammock State Park, Florida [USA]; SMUK, Lawrence. Synonymy: Wood 1957c: 396.
References: (hb) Wood, S. L. 1953: 106, 1954a: 1014. (tx) Wood, S. L. 1953: 106, 1954a: 1014, 1957c: 396, 1962k: 24.
- laevigatus** (Blandford) 1904: 230 (*Hypothenemus*). Lectotype ♀; Los Remedios, Chiriquí, Panama; BMNH, London, designated by Wood 1982b: 914.
Distribution: North America (Nicaragua/ Panama).
References: (ds) Hagedorn 1910d: 44; Kleine 1913b: 121, 1914b: 361; Wood, S. L. 1982b: 914. (tx) Blandford 1904: 229–230; Browne 1970: 557; Hagedorn 1910a: 87, 1912a: 338–339; Wood, S. L. 1982b: 914.
- lepidus** Wood 1971: 36. Holotype ♀; Beverley, Limon Prov., Costa Rica; Wood Collection.
Distribution: North America (Costa Rica/ Guatemala/ Oaxaca in Mexico/ Panama), South America (Brazil/ Colombia).
Hosts: *Canavalia villosa*, *Coffea robusta*, *Protium* sp., *Serjania* sp., *Xeipomia* sp.
Notes: (3) Wood 1982b: 913 (redescribed).
References: (ds) Atkinson & Equihua 1985c: 358, 1988: 88; Beaver 1974a; Wood, S. L. 1971: 36–37, 1982b: 913.
- pilosus** Eggers 1937a: 81. Holotype, sex?; Bolivien (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia/ Brazil).
Hosts: *Protium* sp.
References: (hb) Beaver 1974a. (ds) Beaver 1974a; Blackwelder 1947: 781. (tx) Anderson, W. H. & Anderson 1971: 25; Eggers 1937a: 81; Schedl 1939d: 410, 1939h: 47.
- pubescens** Wood 1986c: 271. Holotype ♀; 69 km N Manaus, Brazil; Wood Collection.
Distribution: South America (Brazil).
References: (tx) Wood, S. L. 1986c: 271.
- punctifrons** Schedl 1939d: 410. Syntypes, sex?; Isla Martín García, Argentina; Schedl Collection in NHMW, Wien, and Viana Collection.
Distribution: South America (Argentina).
Notes: (3) Schedl 1979c: 205 (citation of holotype invalid).
References: (hb) Viana 1964: 125. (ds) Blackwelder 1947: 781; Viana 1964: 125. (tx) Schedl 1939d: 410, 1979c: 205.
- pygmaeus** Schedl 1965e: 370. Holotype, sex?; Congo ex belge, Yangambi; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Hosts: *Caloncobia wehrichtii*.

Notes: (1) The holotype of this species does not belong to this genus; it may be a *Miocryptophagus*.

References: (tx) Schedl 1965c: 370, 1979c: 207.

seriatus Eggers 1933b: 10. Holotype ♀; French Guayana (Nonveau Chantier); MNHN, Paris.

Figures: Atkinson & Aquihua 1988: 91, Bright 1972d: 59, Pedrosa-Macedo & Schonherr 1985: 51.

Distribution: Antilles Islands (Cuba/ Haiti/ Jamaica/ Virgin Islands), North America (Costa Rica/ Honduras/ Colima, Nayarit, Oaxaca, Sinaloa in Mexico/ Florida, S Texas in USA), South America (Bolivia/ Brazil/ Cayenne/ Venezuela).

Hosts: *Araucaria angustifolia*, *Canavalia villosa*, *Chenopodium ambrosioides*, *Conocarpus erecta*, *Dipholis salicifolia*, *Ficus aurca*, *Ipomoea pres-caprae*, *Mangifera indica*, *Metopium toxiferum*, *Persea borbonica*, *Pithecellobium guadeloupense*, *Rhacoma crossopetalum*.

References: (bv) Atkinson, Foltz, & Connor 1988. (ec) Equihua & Atkinson 1986: 630. (hb) Atkinson, Foltz, & Connor 1988; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Wood, S. L. 1953: 104. (ds) Atkinson & Equihua 1985c: 358, 1988: 88; Beaver 1974a; Blackwelder 1947: 781; Bright 1972d: 63, 1985c: 174; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1987a: 66; Estrada & Atkinson 1988: 209; Numberg 1958a: 480, 1972b: 193; Pedrosa-Macedo & Schonherr 1985: 51; Schedl 1960a: 78, 1972g: 39; Schonherr & Pedrosa-Macedo 1981: 52; Wood, S. L. 1977a: 69, 1982b: 913. (tx) Atkinson & Equihua 1988: 91; Bright 1972d: 59, 63, 1985c: 174; Eggers 1933b: 10; Numberg 1958a: 480; Pedrosa-Macedo & Schonherr 1985: 51; Schedl 1939h: 47, 1957a: 194, 1962p: 205, 1979c: 225; Wood, S. L. 1953: 104, 1972e: 193–194, 1975a: 21, 1982b: 913.

adustus Eggers 1933b: 11. Holotype ♀; St. Jean du Maroni, French Guayana; MNHN, Paris. Synonymy: Wood 1972e: 194.

References: (ds) Blackwelder 1947: 781. (tx) Eggers 1933b: 11; Wood, S. L. 1972e: 194.

floridensis Blackman 1943b: 36 (*Tachyderes*). Holotype ♀; Paradise Key, Florida [USA]. Synonymy: Schedl 1962p: 205.

References: (hb) Wood, S. L. 1954a: 1012. (ds) Blackwelder & Blackwelder 1948. (tx) Blackman 1943b: 36; Schedl 1962p: 205; Wood, S. L. 1954a: 959, 979, 1012.

bolivianus Eggers 1943a: 356. Holotype ♀; Cochabamba, Bolivia; USNM, Washington. Synonymy: Wood 1975a: 21.

References: (tx) Anderson, W. H. & Anderson 1971: 7; Eggers 1943a: 356; Schedl 1958f: 46, 1979c: 43; Wood, S. L. 1975a: 21.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Struthanthus* sp.

References: (tx) Wood, S. L. 1986c: 272.

Genus *Hypothenemus* Westwood

HYPOTHEXEMUS WESTWOOD 1836: 34. Type-species: *Hypothenemus eruditus* Westwood, monobasic.

Stephanoderes Eichhoff 1872a: 132. Type-species: *Stephanoderes chapuisi* Eichhoff = *Crypturgus dissimilis* Zimmermann, subsequent designation by Hopkins 1914: 130. Synonymy: Browne 1963: 53.

References: (ay) Van Ryn-Tournel 1975. (bv) Schedl 1960f: 11. (cn) Albuquerque Leao 1941: 319–328; Blackman 1950: 306, 327; Browne 1968: 660; Coleman 1931: 1–26; Dorsey & Leach 1956: 224; Hargreaves 1936: 8–11; Lefevre 1944: 191–200; Luigioni 1929: 997; Mertens 1920: 243–251; Myers 1947: 7; Roque 1946: 30–31; Tucker 1911: 27; Ultee 1925: 1–52; Wolcott 1921: 11. (hb) Browne 1961c: 71–76; Ghesquiere 1933a: 25, 1933b: 774–778; Nusslin 1912: 82; Schedl 1961k: 546, 1977a: 61. (ds) Blackwelder 1947: 777; Browne 1961c: 71; Costa Lima 1956: 292–293; Hagedorn 1910d: 40; Heyden, Reitter, & Weise 1906; Leng 1920; Schedl 1961k: 546, 1977a: 61; Swaine 1909: 116. (tx) Arnett 1960: 1043, 1968: 1043; Balachowsky 1949a: 201; Beal & Massey 1945: 48, 51, 115–117; Beeson 1938: 290; Blackman 1922a: 79, 89; Blandford 1894b: 262, 1894d: 83–85, 1904: 226; Blatchley & Leng 1916: 593, 599; Browne 1963c: 53; Chamberlin 1939: 303; Costa Lima 1928: 117; Dodge 1938: 18; Duffy 1953: 6, 14; Eggers 1940a: 123–141, 1940b: 99–108; Eichhoff 1872a: 132, 1878a: 387, 1878b: 44–46, 142, 475, 496, 1881a: 46, 190, 1883: 110, 134, 1896: 608; Hagedorn 1910a: 84, 1912: 33–43, 1913b: 3; Hopkins 1914: 130, 135, 1915b: 1–75, 1915c: 171–226; Numberg 1960: 287–308; Peyerimhoff 1935: 193–195; Reitter 1894: 73, 1906: 711, 1913a: 5, 71; Roepke 1919: 8; Sampson 1914: 379; Schedl 1934d: 84, 1939a: 342, 1939i: 381, 1941d: 379–424, 1957d: 1–162, 1958f: 33–46, 1959q: 705–710, 1961k: 546, 1962r: 89, 1963: 215–216, 1965f: 3–15, 1977a: 61, 1977b: 61; Stark 1952: 280; Swaine 1918a: 16, 18, 35, 45; Wood, S. L. 1954a: 1011, 1015, 1960a: 35.

Homococryphalus Lindemann 1876: 168. Type-species: *Stephanoderes ehlersii* Eichhoff = *Hypothenemus eruditus* Westwood, monobasic. Synonymy: Fauvel 1884: 315. References: (tx) Blandford 1904: 226; Fauvel 1884: 315, 390; Lindemann 1876: 168; Swaine 1909: 116.

spatulatus Wood 1986c: 272. Holotype ♀; Sta. Maria Chimalpa, Oaxaca, Mexico; Wood Collection.

Adiacretus Hagedorn 1909a: 744. Type-species: *Adiacretus spinosus* Hagedorn = *Stephanoderes elaphus* Eichhoff, monobasic. Synonymy: Schedl 1939: 380.

References: (tx) Hagedorn 1909a: 744; Schedl 1939i: 380, 1961k: 467; Wood, S. L. 1954a: 1050.

Stylotentus Schedl 1939i: 380. *Hypothenemus concolor* Hagedorn, subsequent designation by Schedl 1961k: 448. Synonymy: Wood 1983: 648.

References: (tx) Numberg 1956: 141; Schedl 1939i: 280, 1953c: 294, 1957d: 10, 1961k: 448, 1962r: 89–90, 1977b: 37.

Triarmocerus Eichhoff 1878b: 42, 119. Type-species: *Triarmocerus cryphaloides* Eichhoff, monobasic. Synonymy: Synonymy is based on Eichhoff's placement of *birmanus* in *Triarmocerus* and on a specimen in the Schedl Collection of *cryphaloides* presumably compared with the type.

References: (tx) Eichhoff 1878b: 42, 119; Hagedorn 1910a: 46; Numberg 1956: 141; Schedl 1962r: 89.

Chondronoderes Schedl 1940d: 589. Type-species: *Stephanoderes magnus* Eggers, monobasic. Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1940d: 589, 1961k: 545; Wood, S. L. 1984b: 226.

Archeophalus Schedl 1941d: 392. Type-species: *Archeophalus natalensis* Schedl, monobasic. Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1941d: 392, 1961k: 443; Wood, S. L. 1984b: 226.

Pachynoderes Schedl 1941d: 393. Type-species: *Pachynoderes deprecator* Schedl, monobasic. Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1941d: 393, 1961k: 632; Wood, S. L. 1984b: 226.

Lepiceroides Schedl 1957d: 59. Type-species: *Lepiceroides aterrinus* Schedl, monobasic. Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1957d: 59, 1961k: 635; Wood, S. L. 1984b: 226.

Ernophloeus Numberg 1958a: 484. Type-species: *Ernophloeus costalimai* Numberg = *Stephanoderes fuscicollis* Eichhoff, original designation. Synonymy: Wood 1986c: 266.

References: (tx) Numberg 1958a: 484; Schedl 1960i: 104; Wood, S. L. 1986c: 226.

Epsips Beeson 1941 (1961: 287). Type-species: *Epsips sylvaticum*, nomen nudum. Beeson's series was examined by SLW, but the name was never validated and is not needed.

References: (tx) Beeson 1941 (1961: 287).

Macrocryphalus Nobuchi 1981c: 14. Type-species: *Macrocryphalus oblongus* Nobuchi, probable synonym of *Stephanoderes fuscicollis* Eichhoff. Synonymy: Wood 1992b: 81.

References: (tx) Nobuchi 1981c: 14; Wood, S. L. 1992b: 81.

Keys: Wood 1954a: 1017, 1052, 1982b: 876, Blandford 1904: 227 for North America.

References: (ay) Nobuchi 1969a: 58; Nusslin 1911, 1912; Van Ryn-Toumel 1975. (bv) Schedl 1960f: 11–12. (cn) Blackman 1950: 306, 326–327; Gonçalves 1935: 414–420; Mayne 1917: 1–80; Tulashevili 1930: 189–230. (hb) Beaver 1987a: 18; Bright & Stark 1973: 68; Browne 1963a: 233; Chesquiere 1933a: 25, 1933b: 774–775; Lengerken 1939: 63; Schedl 1958d: 189, 1961k: 467, 1977a: 37; Smith 1890: 54–55; White, R. E. 1983: 33; Wood, S. L. 1986a: 91. (ds) Beaver 1987a: 18; Bright & Stark 1973: 68; Britten 1920: 95–96; Hagedorn 1910d: 40; Schedl 1934f: 1640, 1961k: 467, 1966b: 24, 1977a: 37; Scheerpeltz & Winkler 1930: 257; Swaine 1909: 116; Wachtl 1876: 453; Wood, S. L. 1982b: 875, 1986a: 91. (tx) Arnett 1960: 1043, 1968: 1043; Balachowsky 1949a: 200; Beal & Massey 1945: 60, 117; Beeson 1938: 288, 1941: 374; Blackman 1922b: 76, 82; Blandford 1894b: 260–264, 1894d: 83–86, 1895b: 81–83, 1897a: 183, 1898b: 185, 225, 1904: 226–232, 249; Blatchley & Leng 1916: 592–596; Bright 1972: 48–49; Bright & Stark 1973: 68; Browne 1961: 76–79, 1963: 53–58, 1968: 60; Chamberlin 1939: 12, 37, 102–106, 275, 288, 303; Choo 1983: 65; Choo, Woo, & Nobuchi 1985b; Costa Lima 1928: 118–120; Dodge 1938: 18; Eggers 1934: 27, 1943c: 68; Eichhoff 1864: 34, 45, 56, 1868d: 421, 1896: 608; Erichson 1836: 61; Fauvel 1884: 315; Ferrari 1867a: 4, 7, 1867b: 113; Gozman 1885: 278; Hagedorn 1910a: 84, 86, 138; Hopkins 1914: 116, 123, 133, 1915b: 7, 12–15, 1915c: 117–121; Jacquelin du Val & Fairmaire 1868: 105; Kurenzov 1941: 70; Luigioni 1929: 997; LeConte 1876: 355; LeConte & Horn 1883: 517; Peyerimhoff 1935: 193–196; Reitter 1894: 73–75, 1906: 711, 1913a: 65, 71–74; Sampson 1914: 379; Schedl 1939i: 380, 1940c: 203–208, 1952g: 50, 1957d: 79, 1958f: 33–46, 1961k: 467, 1962r: 88, 1963h: 265, 1963i: 62, 1964k: 215–216, 1977a: 37, 1977b: 48; Sharp 1879: 101–102; Stark 1952: 13–27, 78, 153, 290; Westwood 1836: 34; Wood, S. L. 1954a: 981, 1067, 1960a: 40, 1961: 45, 1982b: 875–911, 1986: 91–93, 1986a: 91.

aberrans Browne 1973a: 287. Holotype ♀; Zaire: Lovanium (Kinshasa); MRCB, Tervuren. Distribution: Africa (Zaire).

References: (hb) Van Ryn-Toumel 1975. (tx) Browne 1973a: 287.

abruptus (Schedl) 1961e: 135 (*Stephanoderes*). Holotype, sex?; Madagascar, Betsatsakry pres Perinet; IRSM, Madagascar. Distribution: Madagascar.

Hosts: *Ravensara* sp.

References: (hb) Schedl 1977a: 62. (ds) Schedl 1977a: 62. (tx) Schedl 1961e: 135, 1977a: 62, 1979c: 9.

acaciae (Eggers) 1920: 120 (*Cryphalus*). Lectotype ♀; Cheren, Abessinien; USNM, Washington, designated by Anderson & Anderson 1971: 3.

Distribution: Africa (Ethiopia).

Hosts: *Acacia catechu*.

References: (hb) Schedl 1961k: 54S. (ds) Schedl 1961k: 54S, 1982b: 27S. (tx) Anderson, W. H. & Anderson 1971: 3; Eggers 1920: 120, 1927: 17S, 1935c: 304, 1936c: 35; Schedl 1961k: 54S, 1979c: 9.

adscitus (Schedl) 1950f: 46 (*Stephanoderes*). Holotype, sex?; Fiji Islands; Eggers Collection, in NHMW, Wien.

Distribution: Fiji Islands.

Hosts: *Acalypha* sp.

References: (ds) Browne 1974a: 64. (tx) Schedl 1950f: 37, 46, 1979c: 12.

aethiops (Schedl) 1965g: 26 (*Lepicicroides*). Holotype, sex?; Congo ex belge, Yangambi; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Hosts: *Dichopetalum thomieri*.

References: (tx) Schedl 1965g: 26, 1979c: 13.

africanus (Hopkins) 1915b: 30 (*Stephanoderes*). Holotype ♀; Capetown, South Africa; USNM, Washington.

Distribution: Africa (South Africa), Antilles Islands (Bahama Islands/ Dominican Republic in Hispaniola/ Jamaica/ Puerto Rico/ Virgin Islands), Asia (Malaya), Indonesia (Java), North America (Belize/ Costa Rica/ Honduras/ Alabama, Louisiana in USA), South America (Brazil/ Venezuela). Hosts: Fruiting pods and twigs of trees, *Pyrus malus*.

Notes: (3) Wood 1967b: 7S (a possible synonym of *setosus*).

References: (hb) Wood, S. L. 1982b: 90S. (ds) Bright 1973: 18, 1985c: 174; Schedl 1961k: 54S; Wood, S. L. 1977a: 70, 1982b: 90S. (tx) Bright 1985c: 174; Hopkins 1915b: 23, 30; Schedl 1961k: 54S; Wood, S. L. 1967b: 7S, 1982b: 90S.

agnatus (Eggers) 1924: 103 (*Stephanoderes*). Lectotype ♂; Elisabethville (Congo); USNM, Washington, designated by Anderson & Anderson 1971: 3.

Distribution: Africa (Equatorial Guinea/ Fernando Po/ Namibia/ South Africa/ Tanzania/ Uganda/ Zaire/ Zambia), Madagascar.

Notes: (1) Schedl 1979c: 15 (designation of lectotype invalid).

References: (ds) Schedl 1961k: 549, 1977a: 62, 1982: 27S. (tx) Anderson, W. H. & Anderson 1971: 3; Eggers 1924: 103, 1927a: 197, 1940b: 102, 1940c: 235–236; Schedl 1939g: 170, 1939i: 3S5, 1961k: 54S–549, 1962h: 67, 1977a: 62, 1979c: 15.

alternatus (Eggers) 1943e: 73 (*Stephanoderes*). Holotype, sex?; Zambeze (Environ de Sone, M. Soussou); MNHN, Paris.

Distribution: Africa (Mozambique).

References: (tx) Eggers 1943e: 73; Schedl 1961k: 549, 1979c: 17.

amakusauus (Murayama) 1934c: 2S7 (*Stephanoderes*). Holotype ♀; Kadoyama in Amakusa Is., Kimmamoto Pref., Kinsu; Murayama Collection in USNM, Washington.

Distribution: Asia (Japan).

Hosts: *Daphniphyllum glaucescens*.

References: (cn) Anonymous 1980g. (ds) Anonymous 1980g; Murayama 1953c: 150, 1954b: 199; Nobuchi 1955c: 15. (tx) Murayama 1934a: 4, 1934c: 2S7–2S8, 1954b: 199.

apicalis Wood 1974a: 19. Holotype ♀; 3 km E Armeria, Colima, Mexico; Wood Collection.

Distribution: North America (Colima, Oaxaca in Mexico).

Hosts: Liana, shrub.

References: (ds) Wood, S. L. 1982b: 8S3. (tx) Wood, S. L. 1974a: 19, 1982b: 8S3.

arecae (Hornung) 1842: 117 (*Bostrichus*). Lectotype ♀; Ostindien nuts intercepted in Germany; MNB, Berlin, designated by Wood 1974d: 2S2.

Figures: Maiti & Saha 1956: 160 (adult, antenna).

Distribution: Africa (Cameroon/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Liberia/ Nigeria/ Sierra Leone/ Uganda/ Zaire), Antilles Islands (Bahama Islands/ Martinique/ Puerto Rico/ Virgin Islands), Asia (Bangladesh/ Burma/ Andaman Islands, Assam, Bengal, Maharashtra, Punjab, Tamil Nadu, Uttar Pradesh in India/ Sri Lanka/ Thailand), Europe (intercepted in Austria, Germany), Hawaiian Islands, Indonesia (Sumatra), Marquesas Islands, Micronesia (Caroline Islands, Mariana Islands, Marshall Islands), New Caledonia, Nine Island, North America (S Florida in USA), Philippine Islands, South America (Brazil).

Hosts: Numerous species of herbs, shrubs, and trees (Browne 1961c: 77–7S); Schedl records are confused with those of *cruditus*. *Bosqueria angolensis*.

Notes: (3) Schedl 1957d: 12 (*mayumbensis* Eggers, nomen nudum, synonymy).

References: (cn) Grandi 1951; Kalshoven 1932: 242; Kleine 1932a: 29S; Mathur & Singh 1960a: 16; Mathur, Singh, & Lal 195S: 32; Schmitz & Crisinel 1957: 10; Sivaram 1980: 5S; Smith, J. B. 1900: 362. (cc) Felt 1906: 702; Wichmann 1955a: 96, 9S. (hb) Blandford 1894b; Browne 1961c: 77–7S; Felt 1906: 702; Grandi 1951; Kleine 1932a: 29S; Lengerken 1939: 64, 1954: 84; Schmitz & Crisinel 1952: 10; Vayssiere 1923: 10S; Yum & Hua 1980: 22S. (ds) Atkinson & Equihua 1985c: 357; Beaver 1987b: 64, 1990a: 2S0; Beeson 1935a: 103, 1935b, 1935b: 2S8, 1961: 2S8; Blandford 1894b; Borchert 1951; Browne 1961c: 7S, 1980b: 3S0; Estrada & Atkinson 198S: 209; Gemminger & Harold 1872: 26S2; Hagedorn 1910d: 40, 1913b: 45–46; Heyden, Reitter, & Weise 1906: 711; Horion 1951; Kalshoven 1932:

242, 1958: 170; Kleine 1913b: 121, 1914b: 269, 288, 300, 307, 1932a: 298, 1934a: 145; LePelley 1968: 113; Lepesme et al. 1948: 645; Maiti & Saha 1980: 159; Mathur & Singh 1960a: 16; Mathur, Singh, & Lal 1958: 32; Pittioni 1943: 176; Reitter 1894a: 75, 1916: 290; Roba 1935: 338; Schedl 1980a: 16; Scheerpeltz & Winkler 1930: 257; Schilsky 1909: 188; Smith, J. B. 1900: 362; Stein & Weise 1877: 164; Sturm 1843: 230; Swaine 1909: 118; Vayssiere 1923: 108; Wichmann 1927a: 79, 1955a: 96, 98; Wood, S. L. 1957e: 1273, 1960a: 41, 1977a: 70, 1982b: 906; Yimus & Hua 1980: 228. **(tx)** Beaver 1991: 92; Beeson 1935a, 1935b; Blandford 1894b: 229–230, 1904: 229; Eggers 1908: 1, 1922a: 86, 1925: 153, 1929e: 55, 1936d: 626; Eichhoff 1878b: 165–166, 1896: 608; Fauvel 1884b: 315, 390; Ferrari 1867a: 17; Hagedorn 1910a: 85; Hopkins 1915b: 14, 1915c: 222; Hornung 1842: 117; Maiti & Saha 1980: 159; Numberg 1956a: 144; Portevin 1935: 325; Reitter 1894a: 75, 1902a: 141, 1913a: 73, 1916: 290; Schedl 1934f: 1640, 1935j: 1, 1942a: 169–170, 1942d: 2, 1950d: 19, 1950g: 895, 1953c: 288, 1954a: 139, 1957d: 12, 1980a: 16; Swaine 1909: 118; Wood, S. L. 1960a: 41, 1974d: 282, 1982b: 906.

vafer Blandford 1896f: 241. Syntypes 4, sex?; Noumea, New Caledonia; BMNH, London. Synonymy: Wood 1974d: 282.

References: **(ds)** Beaver 1976b: 536; Beeson 1935b: 288, 1940: Hagedorn 1910d: 46; Kalshoven 1958: 170; Kleine 1913b: 124, 1914b: 301. **(tx)** Beaver 1976b: 536; Beeson 1935b: 290, 1940: 198; Blandford 1896f: 241; Hagedorn 1910a: 88; Hopkins 1915b: 23; Schedl 1939f: 35; Wood, S. L. 1972f: 51, 1974d: 282.

fungicola Eggers 1908c: 216 (*Stephanoderes*). Holotype ♀; Java; Fiori Collection, Bologna. Synonymy: Eggers 1929e: 52, 55.

References: **(cn)** Kalshoven 1932: 242. **(ds)** Hagedorn 1910d: 43; Kalshoven 1932: 242; Kleine 1913b: 121. **(tx)** Eggers 1908c: 216, 1922a: 86, 1924: 153, 1929e: 55; Hagedorn 1910a: 86; Hopkins 1915b: 14.

polyphagnus Eggers 1924: 104 (*Stephanoderes*). Syntypes, sex?; Congostaat: Mayumbe and Barumba; MRCB, Tervuren, and 4 in Eggers Collection, in NHMW, Wien. Synonymy: Wood 1972f: 52.

Notes: (3) Schedl 1950: 15 (*mayumbensis* Eggers, nomen nudum).

References: **(cn)** Azavedo 1924a; Ghesquiere 1933a: 32–36, 1933b: 782–786; Magnin 1954: 468; Mayne & Donis 1951: 333; Schmitz & Crisinel 1957: 12. **(cc)** Mayne & Donis 1951: 33, 334; Schedl 1958d: 192. **(hb)** Schmitz & Crisinel 1957: 12. **(ds)** Hargreaves 1937: 509. **(tx)** Eggers 1924: 104; Numberg 1960a: 291, 1961b: 609; Schedl 1950c: 204, 1950d: 7, 15, 25, 1952g: 51, 1954e: 50, 70, 1957d: 12, 54,

1959q: 707, 1961k: 610, 1962k: 1070, 1965g: 20, 1979c: 119; Wood, S. L. 1957e: 1273, 1972f: 51.

hispidus Eggers 1925: 156 (*Stephanoderes*). Lectotype, sex?; Birma and India orient, both probably from Tenasserim; Eggers Collection, in NHMW, Wien, designated by Schedl 1979c: 119.

References: **(bv)** Schedl 1960f: 30. **(hb)** Browne 1961c: 78. **(ds)** Bredo 1934: 18; Browne 1961c: 78–79; Ghesquiere 1933a: 26–36, 1933b: 775–782; Hagedorn 1910d: 43; Lepesme 1948: 145; Schedl 1951c: 69, 1959a: 479, 1960f: 30, 1961j: 504, 1962i: 74, 1964c: 305, 1966b: 25. **(tx)** Eggers 1925: 156; Hagedorn 1910a: 86; Nobuchi 1983: 300; Schedl 1950c: 204, 1950d: 3, 17, 1950f: 36, 1953c: 288, 1954d: 871, 1954e: 50, 62, 1959a: 478–479, 1961k: 504, 1979c: 119; Wood, S. L. 1960a: 41–42.

heterolepsis Costa Lima 1928: 117. Syntypes, sex?; Bahia, Brasil; Colleeao Entomologica do Instituto, Brazil. Synonymy: Wood 1972f: 52.

References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956. **(ds)** Blackwelder 1947; Costa Lima 1936, 1956; Kleine 1934a: 145; Schedl 1967d: 2, 1973d: 155. **(tx)** Costa Lima 1928: 117, 1956; Schedl 1948d: 35, 1951m: 71–73, 1952k: 159–160; Wood, S. L. 1972f: 52.

capitalis Beeson 1935: 102. Lectotype ♀; Hakehetau Valley 1000 feet, Uahuka; Vaipall Valley 800 feet, Marquesas Islands; BPBM, Honolulu. Synonymy: Wood 1982b: 906.

References: **(tx)** Beeson 1935: 102–103, 1935b: 289, 1940: 193; Wood, S. L. 1960a: 42, 1982b: 906.

cupolyphagnus Beeson 1940: 193. Holotype ♀; Dehra Dun, United Provinces, India; FRI, Dehra Dun. Synonymy: Wood 1960a: 41.

References: **(cn)** Mathur & Singh 1960b: 86, 1961a: 13, 1961b: 13; Roonwal 1954: 37. **(ds)** Beeson 1935b: 298, 1961: 288; Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1960b: 86, 1961a: 13, 1961b: 13; Roonwal 1954: 37. **(tx)** Beeson 1940: 193–195; Wood, S. L. 1960a: 41–42.

subvestitus Eggers 1940c: 232 (*Stephanoderes*). Holotype ♀; Mosolo Kwenge, Kwango, Congo; MRCB, Tervuren. Synonymy: Wood 1972f: 52.

References: **(cn)** Alvarado 1939b; Ghesquiere 1933a; Schmitz & Crisinel 1957: 12. **(cc)** Bredo 1934: 3–20, 1939: 266–307. **(hb)** Alvarado 1939b; Schmitz & Crisinel 1957: 12. **(ds)** Ghesquiere 1933a; Lepesme 1948: 145; Mayne & Donis 1962: 308; Roba 1935: 340; Thompson, G. H. 1963: 47. **(tx)** Eggers 1940c: 232; Lepesme 1948: 145; Schedl 1950c: 204, 1950d: 3, 17, 1954d: 871, 1954e: 49–50, 1959a: 478, 1961k: 504; Wood, S. L. 1972f: 51.

- baubesaui* Eggers 1940c: 232 (*Stephanoderes*). Holotype ♀; Congostaat: Bambesa: MRCB, Tervuren. 1 Eggers cotype in NHMW, Wien. Synonymy: Wood 1989: 173. References: (tx) Eggers 1940c: 232; Schedl 1950d: 3, 1961k: 469, 1979c: 33; Wood, S. L. 1989: 173.
- martiniquensis* Eggers 1941a: 99 (*Stephanoderes*). Holotype ♀; St. Pierre, Martinique; USNM, Washington. Synonymy: Wood 1972f: 52. References: (ds) Bright 1985c: 175. (tx) Anderson, W. H. & Anderson 1971: 19; Bright 1985c: 175; Eggers 1941a: 99; Schedl 1951m: 72, 1979c: 149; Wood, S. L. 1972f: 51.
- oahuensis* Schedl 1941f: 110. Syntypes ♀; Punaluu, Oahu, Hawaiian Islands; BPBM, Honolulu. Synonymy: Wood 1960a: 41–42. References: (ds) Swezey 1941: 118. (tx) Schedl 1941f: 110, 1951k: 134, 1979c: 36; Wood, S. L. 1960a: 41–42.
- subglabratus* Schedl 1942c: 174. Holotype, sex?; Fiji; Schedl Collection in NHMW, Wien. Synonymy: Beaver 1991: 92. References: (ds) Browne 1974a: 64. (tx) Beaver 1991: 92; Schedl 1942c: 174, 1950f: 35, 1953c: 292, 1979c: 241.
- bauhaniae* Schedl 1950d: 19. Lectotype, sex?; Sierra Leone, Njala; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 36. Synonymy: Wood 1989: 173. References: (hb) Schedl 1961k: 470. (ds) Beaver & Loytyniemi 1955a: 79; Browne 1980b: 382; Schedl 1961k: 470, 1962h: 55, 1962k: 1065. (tx) Schedl 1950d: 19, 1952j: 2, 1957d: 10, 1961k: 470, 1979c: 36; Wood, S. L. 1989: 173.
- occidentalis* Schedl 1954e: 76 (*Stephanoderes*). Lectotype, sex?; Mpraeso [Ghana]; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 177. Synonymy: Wood 1989: 173. References: (hb) Thompson, G. H. 1963: 66. (ds) Schedl 1961k: 605; Thompson, G. H. 1963: 66. (tx) Browne 1965: 188; Schedl 1954e: 49, 76, 1961k: 605, 1979c: 177; Wood, S. L. 1989: 173.
- artocarpi* Browne in Beaver & Browne 1978: 588. Holotype ♀; Malaysia: Penang, Penang Hill, 1500 feet; BMNH, London. Figures: Beaver & Browne 1978: 576. Distribution: Asia (Malaya/Thailand). Hosts: *Artocarpus elastica*, *A. lauceifolius*, *A. rigidus*, *A. scortechinii*, *A. sticus*, *A.* sp., *Campnosperma* sp. References: (cc) Beaver 1979b: 298. (hb) Beaver 1979b: 298; Beaver & Browne 1978: 588. (ds) Beaver 1990a: 280. (tx) Beaver & Browne 1978: 576; Browne 1978: 588.
- arundinis* (Eichhoff) 1878b: 157 (*Stephanoderes*). Holotype, sex?; Italis (Pedemontina); IRSNB, Brussels. Distribution: Europe (Italy). References: (cc) Kleine 1905c: 187. (hb) Eichhoff 1881a: 47, 191. (ds) Hagedorn 1910d: 40–41; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Kleine 1913b: 121; Reitter 1894a: 73; Stein & Weise 1877: 164; Tredl 1907: 13. (tx) Eichhoff 1878a: 356, 1878b: 157, 1881a: 47, 185, 191, 1883a: 110, 134; Hagedorn 1910a: 85; Hopkins 1914b: 23, 1915b: 23; Reitter 1894a: 73, 1913a: 72; Schedl 1934f: 1640.
- ascitus* Wood 1971: 35. Holotype ♀; Puerto Viejo, Heredia Prov., Costa Rica; Wood Collection. Distribution: North America (Costa Rica/Panama). References: (hb) Wood, S. L. 1982b: 898. (ds) Wood, S. L. 1982b: 898. (tx) Wood, S. L. 1971: 35, 1982b: 898.
- ater* (Eggers) 1932c: 31 (*Stephanoderes*). Holotype, sex?; Congostaat (Arawimi: N' Gazi); MRCB, Tervuren. Figures: Numberg 1965b: 31. Distribution: Africa (Congo/ Fernando Po/ Gabon/ Ghana/ Ivory Coast/ Nigeria/ Sierra Leone/ Zaïre). Hosts: *Bauhinia tomentosa*, *Caloucoba welchitschii*, *Carapa procera*, *Chrysophyllum lacourtianum*, *Pascocia* spp., *Peucelethra macrophylla*, *Piptadenia africana*, *Theobroma cacao*. References: (cn) Ghesquiere 1933a: 31–36, 1933b: 782, 786; Gregory 1954. (cc) Schedl 1958d: 192. (hb) Roberts 1969: 129; Schedl 1961k: 448. (ds) Ghesquiere 1933a: 31–36, 1933b: 782, 786; Kleine 1934: 145; Mayne & Donis 1962: 313; Numberg 1965b: 31; Schedl 1961k: 448, 1971g: 192, 1972e: 280. (tx) Eggers 1927b, 1932c: 31–32; Numberg 1965b: 23, 31; Schedl 1939i: 380–381, 1950d: 20, 1961k: 448, 1962p: 204, 1965g: 23.
- aterrimulus* Wood 1989: 178. Holotype, sex?; Ruanda: Ihembe; MRCB, Tervuren, automatic. Distribution: Africa (Ruanda in Zaïre). Hosts: *Microglossa volubilis*. References: (tx) Wood, S. L. 1989: 178, 1992b: 79.
- aterrimus* Schedl 1957d: 59 (*Lepiceroideis*). Holotype, sex?; Ruanda: Ihembe; MRCB, Tervuren, preoccupied by Schedl 1951. References: (cc) Schedl 1958d: 192. (tx) Schedl 1951m: 104, 1957d: 59, 1961k: 636, 1979c: 30; Wood, S. L. 1989: 178, 1992b: 79.
- aterrimus* (Schedl) 1951m: 104 (*Stephanoderes*). Syntypes, sex?; Bolivien, Cochabamba; Schedl Collection in NHMW, Wien. Distribution: South America (Bolivia). Notes: (1) Schedl 1979c: 30 (citation of holotype invalid). References: (hb) Schedl 1961k: 636. (ds) Schedl 1961k: 636. (tx) Schedl 1951m: 104, 1957d: 59, 1961k: 636, 1979c: 30.

atratus (Schedl) 1964j: 45 (*Stephanoderes*). Holotype, sex?; Congo, Yangambi; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Hosts: *Theobroma cacao*.

References: (tx) Schedl 1964j: 45, 1979c: 30–31.

balachowskyi Menier 1973: 141. Holotype ♂; Madagascar Sud, 25 km a l'Est de Tulear sur la route de Tananarive; MNHN, Paris.

Distribution: Madagascar.

Hosts: *Euphorbia oucoclada*.

References: (tx) Menier 1973: 141.

baloghi (Schedl) 1967e: 226 (*Stephanoderes*). Holotype ♀; Kindamba, Meya, Loulo River, Congo; NHMB, Budapest.

Distribution: Africa (Zaire).

References: (tx) Schedl 1967e: 226–227, 1979c: 33.

bambusae Browne 1980: 775. Holotype, sex?; Zaire: Kilindera, 2750 m, face N. Ruwenzori; BMNH, London.

Distribution: Africa (Zaire).

Hosts: Bamboo.

References: (tx) Browne 1980: 775.

bauhaniae Schedl 1950d: 20 (*Stylotenus*). Lectotype, sex?; Sierra Leone, Njala; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 36. Distribution: Africa (Ghana/ Ivory Coast/ Sierra Leone/ Zaire).

Hosts: *Bauhania tomentosa*, *Manihot utilisima*, *Saccharum spontaneum*.

References: (tx) Browne 1965: 188; Schedl 1950d: 19–20, 1952g: 51, 1952j: 2, 1957d: 10, 1961k: 451, 470, 1962k: 1068, 1979c: 36.

bezaziani Peyerimhoff 1935: 192. Syntypes 6, sex?; Perregaux (dep d'Oran), Algeria, Maison-Blanche pres Alger; MNHN, Paris.

Distribution: Africa (Algeria).

Hosts: *Epischlia* sp., *Juglans nigra*.

References: (ds) Peyerimhoff 1935: 192. (tx) Balachowsky 1949a: 205; Peyerimhoff 1935: 192–193.

bicinctus Schedl 1959a: 479. Syntypes ♀; Ceylon; Millawitiya Estate; Schedl Collection in NHMW, Wien.

Distribution: Sri Lanka.

Notes: (1) Schedl 1979c: 37 (citation of holotype invalid).

References: (ds) Schedl 1959a: 479. (tx) Schedl 1959a: 479, 1979c: 37.

bidens Browne 1973a: 285. Holotype ♀; Zaire: Lovanium (Kinshasa); MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Browne 1973a: 285.

birmanus (Eichhoff) 1878b: 486 (*Triarmocerus*). Holotype ♀; Birma; Dohrn Collection at Stettin Museum, now in Schedl Collection in NHMW, Wien.

Figures: Maiti & Saha 1986: 163 (adult).

Distribution: Antilles Islands (Cuba/ Jamaica),

Asia (Bangladesh/ Bonin Islands/ Burma/ Hong Kong in China/ Andaman Islands, Madhya Pradesh, Uttar Pradesh in India/ Japan/ Malaya/ Pakistan/ Sri Lanka/ Thailand/ Vietnam), Australia (Queensland), Cook Islands (Rarotonga), Fiji Islands, Indonesia (Sarawak in Borneo, Celebes, Java, Sumatra), Hawaiian Islands, Madagascar, Micronesia (Guam), New Caledonia, New Guinea, Nine Island, North America (Costa Rica/ Colima in Mexico/ Panama/ Florida in USA), Philippine Islands, Samoan Islands, Society Islands, South America (Galapagos Islands).

Hosts: *Acras sapota*, *Adenanthera pavonia*, *Annona* sp., *Ardesia paniculata*, *Cassia florida*, *Dalbergia gastrophylum*, *Eucalyptus trachyphloia*, *Eugenia buxifolia*, *Ficus aurca*, *Litchi chinensis*, *Mangifera indica*, *Melia azedarack*, *Ocotea catesbyana*, *Persea borbonica*, *Phelocarpus septentrionalis*, *Prunus domestica*, *Quercus* spp., *Rhizophora mangle*, *Swietenia macrophylla*, *Trema floridana*, *Vitis* sp.

Notes: (3) Schedl 1942b: 148 (holotype compared to specimen from Guam).

References: (bv) Gray, B. 1974c. (cn) Anonymous 1966j, 1967m, 1969g; Bigger 1982: 257; Browne 1968: 353; Low Chong Moi 1975. (cc) Equihua & Atkinson 1986: 630. (hb) Beaver 1976b: 534, 1987a: 18; Beaver & Browne 1978: 589; Beaver & Loytyniemi 1985a: 80; Browne 1961c: 71, 1968: 353; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Gray, B. 1974c; Kalkhoven 1958b: 168; Schedl 1977a: 65; Yunus & Hua 1980: 229. (ds) Anonymous 1966j, 1967m, 1969g, 1974j; Atkinson & Equihua 1985c: 356; Beaver 1976b: 534, 1987a: 18, 1987b: 64; Beaver & Browne 1978: 589; Beaver & Loytyniemi 1985a: 80; Beeson 1933: 11, 1938: 288–289, 1940: 197–198, 1941: 357; Blackburn & Sharp 1885: 192–194; Bright 1972d: 53, 1985c: 174; Browne 1961c: 71, 1965b: 188, 1968: 353, 1970: 556, 1974a: 64, 1984a: 353; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Fumasari 1967: 332; Hagedorn 1910d: 44, 46; Kleine 1913b: 126, 1914b: 278; Low 1975; Maiti & Saha 1986: 162; Mathew 1982, 1987: 188; Murayama 1934: 287–288, 1936: 127, 1939: 237; Nobuchi 1985c: 15; Nobuchi & Ono 1973: 181; Perkins 1900: 180; Peter, Bagle, & Balasubramanian 1984; Schedl 1934f: 1640, 1960f: 172, 1962b: 186, 1964c: 304, 1965g: 25, 1966b: 28, 1977a: 65; Swezey 1954: 145; Wood, S. L. 1957c: 402, 1960a: 35, 1977a: 70, 1982b: 886; Yunus & Hua 1980: 229. (tx) Blandford 1894c: 579; Bright 1972d: 53, 1985c: 174; Browne 1970: 556; Costa Lima 1928: 123, 1936: 357; Eggers 1925: 153, 1927: 406, 1935: 153; Eichhoff 1878a: 384, 1878b: 486; Hagedorn 1910a: 81; Hopkins 1915b: 23–24; Maiti & Saha 1986: 162; Michalski 1969b: 570; Nobuchi 1983: 300; Sampson 1914: 379; Schedl 1934g: 177, 1939c: 327, 1939f: 34–35, 1940a: 322, 1941f: 111,

1942b: 147–148, 1951k: 135, 1960c: 172, 1962r: 59, 1965g: 25, 1967f: 163, 1971a: 274, 1971c: 272, 1971f: 148, 1977a: 65, 1979c: 40; Wood, S. L. 1957c: 402, 1960a: 35–37, 1972f: 43, 1982b: 886. *maculicollis* Sharp 1879: 101. Syntypes, sex?;

Oahu, Hawaiian Islands; BMNH, London. Synonymy: Browne 1970: 556, Wood 1972f: 43.

References: (ds) Beeson 1938b: 289; Brimblecombe 1953: 26; Hagedorn 1910d: 44; Kleine 1913b: 124, 1914b: 302; Perkins 1900: 180; Swezey 1941: 119, 1954: 145. (tx) Browne 1970: 556; Hagedorn 1910a: 87; Hopkins 1915b: 23; Schedl 1934g: 177, 1941f: 111, 1951k: 135; Sharp 1879: 101, 1892: 194; Wood, S. L. 1960a: 37, 1972f: 43.

peritus Blandford 1894d: 84. Holotype ♀; Nagasaki, Japan; BMNH, London. Synonymy: Browne 1970: 556.

References: (cn) Ebeling 1959. (ds) Blandford 1894c; Ebeling 1959; Hagedorn 1910d: 44; Kleine 1913b: 121, 1914b: 251; Lever 1943a: 52; Murayama 1936a: 127, 1951c: 3, 1954b: 199; Nobuchi 1966b: 22. (tx) Blandford 1894d: 84; Browne 1970: 556; Hagedorn 1910a: 87; Hopkins 1915b: 24; Murayama 1934c: 287–288, 1936a: 127, 1953: 10, 1954b: 199; Nobuchi 1966b: 22; Schedl 1934f: 1640.

farinosus Blandford 1896f: 241. Syntypes 2, sex?; Noumea [New Caledonia]; BMNH, London. Synonymy: Browne 1970: 556.

References: (ds) Anonymous 1964k, 1965e, 1965g; Beeson 1938b; Hagedorn 1910d: 43; Kleine 1913b: 124, 1914b: 301; Nakao 1963: 30. (tx) Blandford 1896f: 241; Browne 1970: 556; Hagedorn 1910a: 86; Hopkins 1915b: 23.

validus valens Sampson 1914: 355. Holotype ♀; Seychelles, Silhouette; Mare aux Cochons over 100 ft.; BMNH, London.

References: (ds) Sampson 1914: 385. (tx) Browne 1970: 556; Sampson 1914: 385.

perkinsi Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; Honolulu, Hawaii; USNM, Washington. Synonymy: Wood 1972f: 43.

References: (ds) Kleine 1934a: 145. (tx) Hopkins 1915b: 23, 31; Wood, S. L. 1972f: 43.

sterculiæ Hopkins 1915b: 32 (*Stephanoderes*). Holotype ♀; Calapan, P.I.; USNM, Washington. Synonymy: Wood 1972f: 43.

References: (ds) Kleine 1934a: 145; Schedl 1966b: 29. (tx) Beeson 1933: 11; Hopkins 1915b: 24, 32; Nobuchi 1983: 300; Wood, S. L. 1972f: 43.

psidii Hopkins 1915b: 32 (*Stephanoderes*). Holotype ♀; Calapan, P.I.; USNM, Washington. Synonymy: Wood 1972f: 43.

References: (ds) Kleine 1934a: 145; Schedl 1966b: 29. (tx) Beeson 1933: 11; Hopkins 1915b: 24, 32; Nobuchi 1983: 300; Wood, S. L. 1972f: 43.

alter Eggers 1923a: 219 (*Stephanoderes*). Syntypes 10 ♀; New Guinea (Augustafluss),

Rahm, Neu Pommern (New Britannien), Philippinen (Los Banos), Borneo (Sarawak); MNB, Berlin, SMTD, Dresden, MCG, Genova, USNM, Washington. Synonymy: Wood 1960a: 35.

References: (cn) Mathur & Singh 1961a: 79; McIntosh 1949: 46. (hb) Beeson 1933: 11; Yunus & Hua 1950: 229. (ds) Beeson 1933: 11, 1938b: 290, 1961: 295; Mathur & Singh 1961a: 79; Schedl 1959a: 484, 1961c: 71, 1962b: 186; Yunus & Hua 1950: 229. (tx) Beeson 1933: 11; Eggers 1923a: 219, 1927b: 406, 1936d: 624, 625; Michalski 1969b: 570; Schedl 1939e: 327, 329, 1939f: 35, 1939i: 350, 1940b: 434, 1942d: 2–3, 1950f: 37, 45, 1953b: 124, 1958b: 100, 1959a: 484; Wood, S. L. 1960a: 35.

uter Eggers 1923a: 219 (*Stephanoderes*). Syntypes 3, sex?; Neu Guinea (Andai und Waigmina) und Australien (Somerset). Synonymy: Browne 1970: 556.

References: (tx) Beeson 1933: 11; Eggers 1923a: 219, 1936d: 627–628; Schedl 1942d: 19, 1950d: 20.

nibarani Beeson 1933: 10 (*Stephanoderes*). Holotype ♀; Madras: North Salem Division, Jawalagiri; FRI, Dehra Dun. Synonymy: Wood 1959: 174.

References: (cn) Roonwal 1954: 73. (ds) Beeson 1961: 299; Roonwal 1954: 73. (tx) Beeson 1933: 10; Schedl 1940d: 591, 1942c: 178, 1942d: 20; Wood, S. L. 1989: 174.

ampliatus Eggers 1936d: 627 (*Stephanoderes*). Holotype, sex?; Brit. Indien (Mysore: Jakkur); BMNH, London, 1 Eggers cotype, in NHMW, Wien. Synonymy: Wood 1989: 174.

References: (tx) Eggers 1936d: 627–628; Schedl 1979c: 19; Wood, S. L. 1989: 174.

pacificus Beeson 1940: 197 (*Stephanoderes*). Holotype ♀; Henderson Island: northwest side; BPBM, Honolulu. Synonymy: Wood 1960a: 35.

References: (tx) Beeson 1940: 197; Wood, S. L. 1960a: 37.

castaneus Wood 1954a: 1027 (*Stephanoderes*). Holotype ♀; Homestead, Florida [USA]; SMUK, Lawrence. Synonymy: Wood 1960a: 35.

References: (hb) Wood, S. L. 1953: 135, 1954a: 1027. (tx) Wood, S. L. 1953: 135, 1954a: 1027, 1957c: 402, 1960a: 35–37.

biseriatus (Eggers) 1919: 240 (*Stephanoderes*). Lectotype ♀; Kilossa, Deutsch-Ostafrika; USNM, Washington, designated by Anderson & Anderson 1971: 6.

Distribution: Africa (Cameroon/ Ghana/ Kenya/ Mauritius/ Mozambique/ Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaire), Reunion Island. Hosts: *Acacia sieberiana*, *Albizia* spp., *Baphia pubescens*, *Cassia siamea*, *Chlorophora excelsa*, *Chrysophyllum lacourtiannum*, *Dialium pachyphyllum*, *Gilbertiodendron dewevrei*, *Hevea*

brasilensis, *Hylodendron gabunense*, *Musanga cecropioides*, *Paropsia schiebeniana*, *Pen-tachlethra macrophylla*, *Piptadenia africana*, *Tetrapleura tetraptera*, *Trema guineensis*.

References: **(bv)** Schedl 1960f: 75. **(en)** Ghesquiere 1933a: 35, 1933b: 782; Mayne & Donis 1951: 333; Schmitz & Crisinel 1957: 12. **(ce)** Mayne & Donis 1951: 33. **(hb)** Schmitz & Crisinel 1957: 12. **(ds)** Gardiner 1957a: 32; Ghesquiere 1933a: 35, 1933b: 782; Kleine 1934a: 145; Mayne & Donis 1962: 311; Schedl 1959q: 705, 1960f: 31, 1961k: 553, 1967e: 221, 1969d: 7, 1977a: 69; Wichmann 1954: 514. **(tx)** Anderson, W. H. & Anderson 1971: 6; Eggers 1919: 240, 1924: 110, 1932c: 31, 1943c: 74; Schedl 1950c: 204, 1950d: 18, 1952j: 8, 1959q: 705, 1961k: 553, 1967e: 211, 1969d: 7–10, 1977a: 69.

bolivianus (Eggers) 1931b: 29 (*Stephanoderes*). Holotype ♀; Cochabamba, Bolivia; USNM, Washington.

Figures: Pedrosa-Macedo & Schonherr 1985: 46. Distribution: North America (Costa Rica), South America (Bolivia/ Brazil).

Hosts: *Guatteria* sp., *Pouteria* sp., *Protium* sp.

References: **(hb)** Beaver 1974a. **(ds)** Beaver 1974a; Blackwelder 1947: 777; Bright 1972d: 56; Numberg 1958a: 481; Pedrosa-Macedo & Schonherr 1985: 46; Schedl 1966f: 84, 1973d: 154. **(tx)** Anderson, W. H. & Anderson 1971: 7; Bright 1972d: 56; Eggers 1931b: 29, 1934a: 79; Numberg 1958a: 481; Pedrosa-Macedo & Schonherr 1985: 46; Schedl 1951m: 71, 1979c: 43.

brevicollis (Eggers) 1927a: 177 (*Stephanoderes*). Holotype, sex?, Ostafrika: Kikumba Muansa; Eggers Collection, in NHMW, Wien.

Distribution: Africa (Kenya/ Mozambique/ South Africa/ Tanzania), Madagascar.

References: **(ds)** Schedl 1961k: 555, 1977a: 70. **(tx)** Eggers 1927a: 177, 1935a: 304, 1936c: 36; Schedl 1950d: 18, 1961k: 555, 1977a: 70, 1979c: 46.

brevis (Eggers) 1932c: 30 (*Stephanoderes*). Holotype, sex?, Congostaat (Katanga, Prov. Katanga); MRCB, Tervuren.

Distribution: Africa (Zaire).

References: **(tx)** Eggers 1932c: 30; Schedl 1961k: 555.

broiuei Beaver 1991: 53. Holotype ♀; Malaysia: Penang, Georgetown; BMNH, London, automatic.

Figures: Beaver & Browne 1978: 576.

Distribution: Asia (Malaya).

References: **(tx)** Beaver 1991: 53.

striatulus Browne in Beaver & Browne 1978: 584 (*Cryphalomorphus*). Holotype ♀; Malaysia: Penang, Georgetown; BMNH, London, pre-occupied by Schedl 1942.

References: **(tx)** Beaver 1991: 53; Beaver & Browne 1978: 576, 584.

brunneus (Hopkins) 1915b: 31 (*Stephanoderes*). Holotype ♀; Brownsville, Texas [USA]; USNM, Washington.

Figures: Bright 1972d: 59.

Distribution: Antilles Islands (Bahama Islands/ Cuba/ Guadeloupe/ Jamaica/ Puerto Rico/ Trinidad/ Virgin Islands), North America (Honduras/ Colima, Jalisco, Nayarit, Oaxaca, Puebla, Sinaloa, Sonora, Tamaulipas, Veracruz, Yucatán in Mexico/ Panama/ Alabama, Florida, Texas in USA), South America (Galapagos Islands).

Hosts: *Acacia farnesiana*, *Albizia labbekoides*, *Annona* sp., *Ardesia paniculata*, *Bauhinia grandiceps*, *Berria amonilla*, *Cajanus cajan*, *Cantavalia* sp., *Celtis ignana*, *C. laevigata*, *Clematis* sp., *Condalia obtusifolia*, *Dalbergia ecastophyllum*, *Ficus* sp., *Galactia spiciformis*, *Giricidia sepium*, *Grevia asiatica*, *Hovenia dulcis*, *Ichthyomethia communis*, *Leucania glauca*, *Lysiloma bahamensis*, *Mimosa* sp., *Ocotea catesbyana*, *Poinsettia heterophylla*, *Serjania* sp., *Trema floridana*, *Vachellia farnesiana*.

References: **(ce)** Equihua & Atkinson 1986: 630. **(hb)** Chamberlin 1939: 309; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630; Wood, S. L. 1953: 142, 1954a: 1031. **(ds)** Atkinson & Equihua 1985c: 357; Blackwelder 1947: 777; Bright 1972d: 54, 1985c: 174; Chamberlin 1939: 309; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 630, 1988: 209; Ferrer 1942; Kirkendall 1984: 242; Leng 1920: 340; Martorell 1945: 468–469; Wolcott 1936: 317; Wood, S. L. 1977a: 70, 1982b: 910. **(tx)** Bright 1972d: 54, 59, 1985c: 174; Chamberlin 1939: 309; Hopkins 1915b: 23, 31; Schedl 1940a: 342; Wood, S. L. 1953: 142, 1954a: 959, 1018, 1031, 1977b: 208, 1982b: 910.

frontalis Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; Brownsville, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1031.

References: **(hb)** Chamberlin 1939: 309. **(ds)** Blackwelder 1947: 778; Chamberlin 1939: 309; Ferrer 1942; Leng 1920: 340. **(tx)** Chamberlin 1939: 309; Hopkins 1915b: 23, 31; Schedl 1940a: 342; Wood, S. L. 1954a: 959, 1031.

cryphalomorphus Schedl 1939n: 14. Holotype ♀; Trinidad, British West Indies; BMNH, London. Synonymy: Wood 1977b: 208.

References: **(ds)** Blackwelder 1947: 777. **(tx)** Schedl 1939n: 14; Wood, S. L. 1977b: 208.

bituberculatus Eggers 1940a: 126 (*Stephanoderes*). Holotype ♀; Env. de Trois Rivières, Guadeloupe; MNHN, Paris. Synonymy: Wood 1977b: 208.

References: **(tx)** Eggers 1940a: 126, 1979c: 41; Wood, S. L. 1977b: 208.

californicus Hopkins 1915b: 19. Holotype ♀; Pomona, California [USA]; USNM, Washington. Distribution: Africa (Liberia), Asia (Israel/ Korea), North America (Michoacan, Mexico, Tamaulipas in Mexico/ California, District of Columbia, Florida, Illinois, Indiana, Kansas, Kentucky, Maryland, Missouri, New Jersey, North Carolina, South Carolina, Tennessee, Texas, Virginia in USA), South America (Brazil).

Hosts: *Aloe vera*, *Bauhinia alba*, *Bidens pilosa*, *Boehmeria scabra*, *Cajanus cajan*, *Cappria bifolia*, *Galactia spiciformis*, *Ipomoea cathartica*, *I. litoralis*, *Iva imbricata*, *Malvastrum* sp., *Mangifera indica*, *Paspalum vaginatum*, *Quisqualis indica*, *Salix babylonica*, *Uniola paniculata*, *Verbena* sp., *Zea mays*.

Notes: (3) This species was probably introduced to America from Africa or SW Asia; synonymy with other species, possibly *leprieuri*, is expected.

References: (ec) Equihua & Atkinson 1986: 631. (hb) Bright & Stark 1973: 69; Burgos & Saucedo 1983: 106; Chamberlin 1939: 294; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 631; Wood, S. L. 1953: 190, 1954a: 1053, 1982b: 897. (ds) Atkinson & Equihua 1985c: 357; Atkinson et al. 1986: 64; Bright & Stark 1973: 69; Burgos & Saucedo 1983: 106; Chamberlin 1939: 294; Choo 1983: 65; Choo & Woo 1985: 164; Choo, Woo, & Park 1983: 174; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 631; Kirk 1970; Leng 1920: 340; Weber, B. C. & McPherson 1991: 49, 54; Wood, S. L. 1977a: 70, 1982b: 897. (tx) Chamberlin 1939: 294; Choo 1983: 65; Hopkins 1915b: 14, 19; Wood, S. L. 1953: 190, 1954a: 979, 1053, 1972f: 44, 1982b: 897.

tritici Hopkins 1915b: 19. Holotype ♀; Dallas, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1055.

References: (hb) Chamberlin 1939: 295; Wood, S. L. 1953: 192, 1954a: 1055. (ds) Chamberlin 1939: 295; Kleine 1934a: 144; Leng 1920: 340. (tx) Chamberlin 1939: 295; Hopkins 1915b: 14, 19; Wood, S. L. 1953: 192, 1954a: 1055, 1972f: 44.

thoracicus Hopkins 1916: 598. Holotype ♀; Clark Co., Indiana [USA]; USNM, Washington. Synonymy: Wood 1954a: 1055, 1982b: 897.

References: (hb) Chamberlin 1939: 294. (ds) Chamberlin 1939: 294; Leng 1920: 340. (tx) Chamberlin 1939: 294; Hopkins 1916: 598; Wood, S. L. 1954a: 959, 1055.

zcae Schedl 1973d: 169 (*Stephanoderes*). Holotype ♀; Brasilien, Sao Paulo, Pinhal; MZUSP, Sao Paulo. Synonymy: Wood 1989: 174.

References: (ds) Schedl 1973d: 155, 169, 1979c: 270; Wood, S. L. 1989: 174.

camerunus (Eggers) 1922b: 167 (*Stephanoderes*). Holotype, sex?; Joko (Kamerun); Eggers Collection.

Figures: Numberg 1960a: 290, 292, 1960b: 29.

Distribution: Africa (Angola/ Cameroon/ Congo/ Ghana/ Ivory Coast/ Sierra Leone/ Tanzania/ Uganda/ Zaïre).

Hosts: *Acacia* spp., *Albizia* spp., *Baphia pubescens*, *Celtis soyauxii*, *Chrysophyllum* sp., *Dialium* sp., *Gilbertiodendron* sp., *Herce* sp. See also Browne 1963a: 233.

Notes: (1) Neither Anderson & Anderson 1971 nor Schedl 1979c mention this holotype; however, Schedl 1979c: 51 mentions a cotype in NHMW, Wien, although no cotypes were designated. This specimen could be the type on loan to Schedl.

References: (hb) Browne 1963a: 223; Schedl 1961k: 555; Thompson, C. H. 1963: 66. (ds) Browne 1955: 188, 1963a: 233; Ferreira 1965: 1114; Gardner 1957a: 32; Mayne & Donis 1960: 104, 1962: 311; Schedl 1959p: 17, 1961k: 555, 1964j: 40, 1965f: 4, 1967e: 211; Thompson, C. H. 1963: 66. (tx) Eggers 1922b: 167, 1924: 111, 1927a: 198, 1941a: 99; Schedl 1954e: 49–50, 1961k: 555, 1962h: 69, 1962k: 1069, 1979c: 51. *brunneipes* Numberg 1960a: 289 (*Stephanoderes*).

Holotype, sex?; Uganda: Bugiri, 1400 m; MRCB, Tervuren. Synonymy: Schedl 1962k: 1069.

References: (tx) Numberg 1960a: 289–290, 292, 1960b: 29; Schedl 1962h: 64, 1962k: 1069.

carbonarius (Eggers) 1943e: 73 (*Stephanoderes*).

Holotype ♀; Mozambique (Vila Pery); MNHN, Paris, 1 Eggers cotype, in NHMW, Wien.

Distribution: Africa (Mozambique/ South Africa).

Hosts: *Gossypium hirsutum*.

References: (ds) Schedl 1982: 278. (tx) Eggers 1943e: 73; Schedl 1950d: 24, 1957b: 150, 1961k: 557, 1979c: 53.

ceibae Hopkins 1915b: 20. Holotype ♀; Cayamas, Cuba; USNM, Washington.

Distribution: Antilles Islands (Cuba/ Granada/ Puerto Rico).

Hosts: *Ceiba* sp.

References: (ds) Blackwelder 1947: 777; Bright 1985c: 174; Kleine 1934a: 143; Leng & Mutchler 1917: 219. (tx) Bright 1985c: 174; Hopkins 1915b: 15, 20.

colae (Schedl) 1957d: 54 (*Stephanoderes*). Holotype ♀; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaïre).

Hosts: *Cola congolana*, *C. griseiflora*, *Duboscia viridiflora*.

References: (ec) Schedl 1958d: 190–191. (hb) Schedl 1961k: 558. (ds) Schedl 1961k: 558. (tx) Schedl 1957d: 54, 1961k: 558, 1979c: 59.

columbi Hopkins 1915b: 18. Holotype ♀; Columbus, Texas [USA]; USNM, Washington.

Distribution: Antilles Islands (Bahama Islands/ Cuba), North America (Belize/ Costa Rica/ Colima, Nayarit, Sinaloa, Veracruz in Mexico/ Panama/ Florida, Georgia, Louisiana, Mississippi,

- South Carolina, S Texas in USA), South America (Colombia/ Venezuela).
- Hosts: *Bauhinia alba*, *Citrus auratifolia*, *Ficus* sp., *Ichthyomethia communis*, *Morus rubra*, *Quercus* spp., *Salix* sp., *Serjania* sp.
- References: (ce) Equihua & Atkinson 1986: 631. (hb) Chamberlin 1939: 294; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 631; Wood, S. L. 1953: 208, 1954a: 1062. (ds) Atkinson & Equihua 1985c: 357; Bright 1985c: 174; Chamberlin 1939: 294; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 631; Estrada & Atkinson 1988: 209; Kirkendall 1984: 242; Leng 1920: 340; Wood, S. L. 1977a: 70, 1982b: 907. (tx) Bright 1985c: 174; Chamberlin 1939: 294; Hopkins 1915b: 14, 18; Wood, S. L. 1953: 208, 1954a: 959, 1062, 1982b: 907.
- abdominalis* Hopkins 1915b: 18. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1954a: 1062.
- References: (ds) Blackwelder 1947: 777; Leng & Mutchler 1917: 219. (tx) Hopkins 1915b: 14, 18; Wood, S. L. 1954a: 1062.
- brunneipennis* Hopkins 1915b: 18. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1954a: 1062.
- References: (ds) Blackwelder 1947: 777; Leng & Mutchler 1917: 219. (tx) Hopkins 1915b: 14, 18; Wood, S. L. 1954a: 959, 1062.
- rufopalliat* Hopkins 1915b: 18. Holotype ♀; Charleston, South Carolina [USA]; USNM, Washington. Synonymy: Wood 1954a: 1062.
- References: (hb) Chamberlin 1939: 294. (ds) Blatchley & Leng 1916: 598; Chamberlin 1939: 294; Kirk 1969; Kleine 1934a: 144; Leng 1920: 340. (tx) Blatchley & Leng 1916: 598; Chamberlin 1939: 294; Hopkins 1915b: 14, 18; Wood, S. L. 1954a: 959, 1062.
- amplipennis* Hopkins 1915b: 19. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1954a: 1062.
- References: (cn) Souphieff & Scherbinovskaja 1937: 81. (ds) Blackwelder 1947; Leng & Mutchler 1917: 219; Souphieff & Scherbinovskaja 1937: 81. (tx) Hopkins 1915b: 14, 19; Wood, S. L. 1954a: 1062.
- comosus* Bright 1972d: 50. Holotype ♀; Clydesdale, St. Andrew Parish, Jamaica; CNCI, Ottawa.
- Distribution: Antilles Islands (Jamaica).
- References: (ds) Bright 1985c: 174. (tx) Bright 1972d: 50, 1985c: 174; McNamara 1977: 196.
- concolor* Hagedorn 1909a: 744. Syntypes, sex?; Kamerun; MNB, Berlin.
- Figures: Schedl 1961k: 452.
- Distribution: Africa (Cameroon/ Canary Islands/ Congo/ Ghana/ Ivory Coast/ Uganda/ Zaire).
- Hosts: *Aucoumea klaineana*, *Bussea occidentalis*, *Pancovia laurentii*, *Pentaclethra macrophylla*, *Synsepalum longecucucatum*, *Terminalia superba*, *Theobroma cacao*.
- References: (cn) Chesquiere 1933a: 32–36, 1933b: 783, 786; Magnin 1954: 468. (hb) Browne 1963: 234; Roberts 1969: 129; Schedl 1961k: 451. (ds) Browne 1963a: 234, 1970: 540; Chesquiere 1933a: 32–36, 1933b: 783, 786; Hagedorn 1910d: 41; Kleine 1913b: 124, 1914b: 310; Mayne & Donis 1962: 314; Schedl 1961k: 451, 1962l: 58, 1962k: 1068, 1964f: 618. (tx) Eggers 1919: 243, 1924: 102, 1932c: 31–32; Hagedorn 1909a: 744, 1910a: 86; Schedl 1939i: 380–381, 1950d: 20, 1954d: 871, 1961k: 451–452, 634.
- confusus* (Eggers) 1940c: 235 (*Stephanoderes*). Holotype, sex?; S. Rhodesia; Salisbury; Eggers Collection, in NHMW, Wien.
- Distribution: Africa (Namibia/ South Africa/ Tanzania/ Zambia/ Zimbabwe).
- References: (ds) Beaver & Loytyniemi 1989; Schedl 1961k: 560. (tx) Beaver & Loytyniemi 1989; Eggers 1940c: 235; Schedl 1957b: 150, 1961k: 560, 1979c: 62.
- corni* Kurenzov 1941a: 162. Syntypes, sex?; Iman and Khor rivers, Ussuri, USSR; ISBN, Novosibirsk, and IZL, Leningrad.
- Figures: Michalski 1969b: 568.
- Distribution: Asia (Sakhalin Island, E USSR).
- Hosts: *Cornus tataricum*.
- Notes: (1) Two specimens deposited by Kurenzov in the Wood Collection are of *Ernoporicus* [confirmation is needed].
- References: (cn) Kurenzov 1956a: 86, 102. (hb) Stark 1952: 283. (ds) Krivolutskaia 1958: 139, 1983; Kurenzov 1951b: 17; Stark 1952: 283. (tx) Kurenzov 1941a: 162–163; Michalski 1969a: 893, 1969b: 568; Stark 1952: 283.
- cosmoderoides* Murayama 1961b: 30. Holotype ♀; Noo, south part of Niigata prefecture, Japan; Murayama Collection in USNM, Washington.
- Distribution: Asia (Honshu in Japan).
- References: (ds) Nobuchi 1985c: 15. (tx) Murayama 1961b: 30.
- costatus* (Eichhoff) 1878a: 386 (*Stephanoderes*). Syntypes, sex?; America septentrionalis (Venezuela); Hamburg Museum, lost.
- Distribution: South America (Venezuela).
- References: (ds) Blackwelder 1947: 777; Hagedorn 1910d: 41; Kleine 1913b: 121, 1914b: 339. (tx) Eichhoff 1878a: 386, 1878b: 154; Hagedorn 1910a: 86; Hopkins 1915b: 24.
- criticus* (Schedl) 1937b: 398 (*Cryphalus*). Lectotype, sex?; Belgisch-Kongo, Stanleyville; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 70.
- Distribution: Africa (Cameroon/ Ghana/ Uganda/ Zaire).
- Hosts: *Beilschmiedia corbisieri*, *Paropsia schiebeniana*, *Theobroma cacao*, *Tridesmostemon claessensii*.

References: (**hb**) Schedl 1961k: 453–454. (**ds**) Mayne & Donis 1962: 314; Schedl 1961k: 453, 1964f: 61S, 1967e: 210, 1972e: 280. (**tx**) Schedl 1937b: 39S, 1957d: 10, 1961k: 453, 1972p: 15S, 1979c: 70.

alternans Browne 1970: 554. Holotype ♀; Uganda, Entebbe; BMNH, London. Synonymy: Schedl 1972p: 15S.

References: (**tx**) Browne 1970: 554; Schedl 1972p: 15S.

crudiæ (Panzer) 1791: 35 (*Bostrichus*). Syntypes, sex?; India occidentali: lost.

Figures: Schedl 1961k: 611.

Distribution: Africa (Angola/ Cameroon/ Cape Verde Islands/ Congo/ Dahomey/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Nigeria/ Sierra Leone/ South Africa/ Uganda/ Zaïre), Antilles Islands (Cuba/ Grenada/ Puerto Rico/ Trinidad), Asia (Malaya/ Sri Lanka/ Thailand/ Vietnam), Cook Islands (Rarotonga), Hawaiian Islands, Indonesia (Java), Madagascar, Marquesas Islands, Micronesia, North America (Costa Rica/ El Salvador/ Guatemala/ Honduras/ all states in Mexico/ Nicaragua/ Alabama, Arkansas, District of Columbia, Florida, Georgia, Illinois, Indiana, Kansas, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA), South America (Argentina/ Bolivia/ Brazil/ Colombia/ Ecuador/ Guyana/ Paraguay/ Suriname/ Venezuela).

Hosts: *Abutilon mollissimum*, *Acacia farnesiana*, *Achras sapota*, *Adenanthera pavonina*, *Aloc vera*, *Astragalus* sp., *Bauhinia grandiceps*, *B. krugi*, *Betula* sp., *Bidens pilosa*, *Bignonia* sp., *Boehmeria scabra*, *Bucida buceras*, *Busera* sp., *Carica papaya*, *Carya* spp., *Cassia nodosa*, *Cinnamomum camphora*, *Citrus* sp., *Clerodendron squamatum*, *Crataegus* sp., *Dalbergia ecastophyllum*, *Derris* sp., *Dioclea megacarpa*, *Diphyisia robinoides*, *Dolonix regia*, *Ficus* spp., *Glycine max*, *Grevia asiatica*, *Inga* sp., *Juglans nigra*, *Magnolia* sp., *Mangifera indica*, *Morus rubra*, *Passiflora latifolia*, *Phalocarpus septentrionis*, *Pinus taeda*, *Prunus persica*, *Pyrus malus*, *Quercus* spp., *Quisqualis indica*, *Rhizophora mangle*, *Richinus communis*, *Schleichera trifuga*, *Serjania racemosa*, *Sida rhombifolia*, *Smilax* sp., *Theobroma cacao*, *Wisteria* sp., *Yucca* sp.

Notes: (1) Identification of this species was based on specimens compared to the types by Eggers before they were lost. Published reports of this species from India and other parts of Asia are apparently based on misidentifications.

References: (**cn**) Smith, J. B. 1900: 362. (**cc**) Equihua & Atkinson 1956: 631; Halperin & Holzschuh 1956: 26. (**hb**) Beaver & Browne 1975: 290; Burgos & Saucedo 1983: 105; Deyrup & Atkinson 1957a: 66; Equihua & Atkinson 1956:

631; Halperin & Holzschuh 1954: 26; Wood, S. L. 1953: 163, 1954a: 1041, 1952b: 891. (**ds**) Atkinson & Equihua 1955c: 357; Beaver & Browne 1975: 290; Blackwelder 1947: 777; Bright 1955c: 174; Browne 1974a: 64, 1955b: 291; Burgos & Saucedo 1983: 105; Deyrup 1981b: 7; Deyrup & Atkinson 1957a: 66; Equihua & Atkinson 1956: 631, 1958: 209; Hagedorn 1910d: 43; Halperin & Holzschuh 1954: 26; Kirkendall 1954: 241; Smith, J. B. 1900: 362; Weber, B.C. & McPherson 1991: 54; Wood, S. L. 1977a: 71, 1983: 105. (**tx**) Bright 1955c: 174; Eggers 1931c: 184; Eichhoff 1896: 60S; Panzer 1791: 35, 37; Schedl 1935i: 24; Wood, S. L. 1953: 163, 1954a: 1041, 1972f: 44, 1952b: 891.

mucronifer Wollaston 1867: 116 (*Cryphalus*). Syntypes, sex?; S. Antao, S. Vicente, S. Iago, et Fogo, Cape Verde Islands; BMNH, London. Synonymy: Wood 1959: 174.

References: (**cn**) Souphieff & Scherbinovskaja 1937: 36. (**ds**) Gemminger & Harold 1872: 2683; Hagedorn 1910d: 44; Kleine 1913b: 11S, 1914b: 379; Souphieff & Scherbinovskaja 1937: 26. (**tx**) Hagedorn 1910a: 87; Wollaston 1867: 116; Wood, S. L. 1959: 174.

hispidulus LeConte 1868: 156 (*Cryphalus*). Syntypes, sex?; District of Columbia, Georgia, Louisiana [USA]; MCZ, Cambridge. Synonymy: Eichhoff 1896: 60S.

References: (**cn**) Funk 1907: 217; Kleine 1932a: 302. (**hb**) Beal & Massey 1945: 11S–119; Blackman 1922b: 84; Chamberlin 1939: 289; Costa Lima 1928b: 11S–120; Kleine 1932a: 302; Morstatt 1924: 45; Smith, J. B. 1890b: 54. (**ds**) Anonymous 1926c: 51S; Beal & Massey 1945: 11S–119; Blackman 1922b: 84; Blackwelder 1947: 77S; Blatchley & Leng 1916: 596; Britton 1920a: Chamberlin 1939: 289; Costa Lima 1936; Currie 1905; Gemminger & Harold 1872: 268S; Hagedorn 1910d: 43; Hamilton 188S: 15S, 1894: 406; Henshaw 1882: 269; Kleine 1913b: 121, 1914b: 35S, 1932a: 302, 1934a: 145; Leng 1920: 339; Leonard 192S: 51S; Pfeffer 1932: 1–23; Schwarz 1878d: 46S; Smith, J. B. 1890b: 54, 1900: 362, 1910: 401; Swaine 1909: 11S. (**tx**) Beal & Massey 1945: 11S–119; Blackman 1922b: 84; Blandford 1894b: 263; Blatchley & Leng 1916: 596; Chamberlin 1939: 289; Eichhoff 1878b: 136, 1896: 60S, 610; Fauvel 1884, 1887; Hopkins 1915b: 13; LeConte 1868: 156, 1876: 35S, 1878a: 46S; Muesebeck 1942: 97, 1950: 133; Sharp 1879: 102; Swaine 1909: 11S; Wood, S. L. 1954a: 959, 1041, 1972f: 44.

nanus Hagedorn 1909a: 744. Syntypes ♀; Argentina; La Plata Museum. Synonymy: Schedl 1952k: 162.

References: (**hb**) Viana 1964: 122. (**ds**) Blackwelder 1947: 777; Bruch 1914a; Hagedorn 1910d: 44; Kleine 1913b: 124, 1914b: 333; Viana

- 1964: 122. (**tx**) Hagedorn 1909a: 744, 1910a: 87; Hopkins 1915b: 23; Schedl 1935i: 24, 1948d: 36, 1948f: 363, 1951h: 284, 1951m: 71–73, 1952a: 446, 1952k: 159–162, 1955f: 34.
- differens* Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; San Bernardino, Paraguay; USNM, Washington. Synonymy: Wood 1972f: 44.
- References: (**ds**) Blackwelder 1947. (**tx**) Hopkins 1915b: 21, 25; Wood, S. L. 1962: 78, 1967b: 78, 1972f: 44.
- guatemalensis* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Trece Aguas, Alta Verapaz, Guatemala; USNM, Washington. Synonymy: Wood 1954a: 1041.
- References: (**cn**) Anonymous 1928d: 14. (**ds**) Blackwelder 1947: 778; Dampf 1928: 14; Ferrer 1942; Kleine 1934a: 145; Wolcott 1936: 317. (**tx**) Hopkins 1915b: 22, 26; Schedl 1940a: 342, 1950c: 206–207; Wood, S. L. 1954a: 1041.
- brasilensis* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Perambuco, Brazil; USNM, Washington. Synonymy: Wood 1954a: 1041.
- References: (**ds**) Blackwelder 1947: 777; Costa Lima 1936; Martorell 1945: 468; Wolcott 1936: 317, 1948: 380. (**tx**) Blackman 1922b: 95; Hopkins 1915b: 22, 26; Wood, S. L. 1954a: 959, 1041.
- paraguayensis* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; San Bernardino, Paraguay; USNM, Washington. Synonymy: Wood 1972f: 44.
- References: (**ds**) Blackwelder 1947: 778. (**tx**) Hopkins 1915b: 22, 26; Wood, S. L. 1972f: 44.
- lecontei* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Jefferson County, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1041.
- References: (**hb**) Chamberlin 1939: 305. (**ds**) Blatchley & Leng 1916: 600; Chamberlin 1939: 305; Kleine 1934a: 145; Leng 1920: 340. (**tx**) Blatchley & Leng 1916: 600; Chamberlin 1939: 305; Hopkins 1915b: 22, 27; Wood, S. L. 1954a: 959, 1041.
- polyphagus* Costa Lima 1924: 316 (*Stephanoderes*). Syntypes, sex?; publication not seen, automatic. Synonymy: Wood 1972f: 44.
- References: (**bv**) Schedl 1960f: 43. (**cn**) Ghesquiere 1933a. (**cc**) Schedl 1961k: 610. (**hb**) Browne 1963a: 234; Schedl 1961k: 610. (**ds**) Browne 1963a: 234; Ghesquiere 1933a; LePelley 1968: 113; Mayne & Donis 1960: 104, 1962: 312; Nonveiller 1984: 41; Nunberg 1961b: 609; Schedl 1960f: 11, 1961k: 610, 1962h: 58, 1962k: 1070, 1964e: 68, 1964j: 41, 1965g: 20, 1966c: 224, 1967e: 212. (**tx**) Costa Lima 1924: 316, 414, 1925: 194–199, 365–368; Schedl 1961k: 610–611; Wood, S. L. 1972f: 44.
- fallax* Costa Lima 1924: 414 (*Stephanoderes*). Syntypes, sex?; publication not seen, automatic. Synonymy: Wood 1972f: 44.
- References: (**cn**) Azavedo 1924a: 359–360; Bondar 1925: 1–13; Kleine 1932a: 302. (**hb**) Kleine 1932a: 145, 302. (**ds**) Kleine 1932a: 145, 302. (**tx**) Costa Lima 1924: 316, 414, 1925: 197, 1925: 122; Wood, S. L. 1972f: 44.
- largipennis* Toleda Piza Junior 1924b: 354 (*Stephanoderes*). Syntypes, sex?; Brazil; not located. Synonymy: Costa Lima 1925: 197.
- References: (**cn**) Kleine 1932a: 302. (**hb**) Kleine 1932a: 302. (**ds**) Kleine 1932a: 145, 302. (**tx**) Costa Lima 1925: 197, 1925: 122; Toleda Piza Junior 1924b: 354–355.
- misericordatus* Eggers 1924: 103 (*Stephanoderes*). Lectotype ♀; Congostaat (Luebo); USNM, Washington, designated by Anderson & Anderson 1971: 35. Synonymy: Wood 1972f: 44.
- Notes: (3) Schedl 1963j: 479 (*Stephanoderes notatus* Eggers, nomen nudum, synonymy). References: (**bv**) Schedl 1960f: 43. (**cn**) Coleman 1931: 1–26; Corbett 1933; Ghesquiere 1933a: 32–35, 1933b: 783; Mathur & Singh 1960a: 13; Mathur, Singh, & Lal 1958: 31; Mathur et al. 1958: 103; Mayne & Donis 1951: 333–334; Schmitz & Crisinel 1957: 12; Schouteden 1924: 56–58; Thomas, K. M. 1949: 83. (**cc**) Mayne & Donis 1951: 333–334; Schedl 1961k: 627. (**hb**) Browne 1963a: 234; Corbett 1933; Schedl 1961: 627; Schmitz & Crisinel 1957: 12; Schouteden 1924: 56; Thomas, K. M. 1949: 83. (**ds**) Beeson 1941: 389, 1961: 288, 299; Brader 1964: 5; Browne 1963a: 234; Ferreira 1965: 1116; Ghesquiere 1933a; Mathur & Singh 1960a: 13; Mathur, Singh, & Lal 1958: 31, 103; Peyerimhoff 1935: 193; Schedl 1959a: 485, 1959p: 17, 1960f: 11, 1961k: 627, 1962h: 58, 1962k: 1070, 1964e: 68, 1964j: 41, 1967e: 212, 1969c: 50, 1969d: 5, 1977a: 77; Wichmann 1954: 514. (**tx**) Anderson, W. H. & Anderson 1971: 35; Corbett 1933; Eggers 1924: 103, 1936d: 626; Michalski 1969b: 570; Peyerimhoff 1935: 193; Schedl 1934d: 91, 1950d: 17–18, 1951j: 19, 1953d: 68, 1955d: 268, 1959a: 485, 1961k: 627, 1962k: 1070, 1963j: 478–479, 1964e: 68, 1964j: 41, 1967e: 212, 1967f: 151, 1969c: 50, 1969d: 5, 1971a: 280, 1971c: 364, 1977b: 77, 1979c: 261; Wood, S. L. 1972f: 44.
- hivaeca* Beeson 1935b: 105 (*Stephanoderes*). Holotype ♀; Tahauku, Hivaoca, Marquesas Islands; BPBM, Honolulu. Synonymy: Wood 1972f: 44.
- References: (**ds**) Beeson 1935b: 290; Ford 1958: 333; Wood, S. L. 1960a: 35. (**tx**) Beeson 1935b: 105, 1935b: 290, 1939: 105–106; Wood, S. L. 1960a: 38–39, 1972f: 44.
- lebromeci* Beeson 1935b: 104 (*Stephanoderes*). Syntypes, sex?; Tahuata, Hanatuma Valley 150 feet, Uapou; Hakahuetau 500 feet, Marquesas

- Islands; BPBM, Honolulu. Synonymy: Wood 1972f: 44.
References: **(hb)** Yunus & Hua 1980: 229, **(ds)** Beeson 1935b; Swezey 1941: 119; Yunus & Hua 1980: 229, **(tx)** Beeson 1935b: 104, 1938: 290; Schedl 1941f: 111, 1965g: 25; Wood, S. L. 1960a: 38–39, 1972f: 44.
- cryphaloides* (Eichhoff)** 1878b: 119 (*Triarmocerus*). Holotype ♀; Madagascar, insula Africana; Hamburg Museum, lost.
Distribution: Madagascar.
Notes: (3) A female from Clarion Island (Mexico) in the Schedl Collection was determined by Schedl as this species. Although that specimen is actually of *erectus* LeConte, *erectus* was apparently introduced to America from Africa. It is possible that the type of *cryphaloides* is a synonym of *erectus*.
References: **(ds)** Alluand 1900: 439; Hagedorn 1910d: 46; Kleine 1913b, 1914b: 326, **(tx)** Eichhoff 1878a: 384, 1878b: 119; Hagedorn 1910a: 81, 1913b; Hopkins 1914: 131.
- cuneolus* (Schedl)** 1936j: 24 (*Stephanoderes*). Lectotype, sex?; Malay Peninsula, Selangor: Kanching Forest Reserve; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 72.
Distribution: Asia (Malaya).
Hosts: *Balanocarpus heimii*, *Dryobalanops aromatica*, *Shorea leprosula*.
References: **(ds)** Beeson 1961: 298; Browne 1961c: 72, 1970: 554; Schedl 1936j: 24, **(tx)** Schedl 1936j: 24, 1979c: 72.
- curtipennis* (Schedl)** 1950f: 45 (*Stephanoderes*). Holotype, sex?; Fiji, Viti Levu: Belt Road, 30 miles west of Suva, 10–250 feet; BPBM, Honolulu.
Distribution: Fiji Islands, Samoan Islands.
References: **(ds)** Beaver 1976b: 536; Browne 1974a: 64, **(tx)** Schedl 1950f: 45–46, 1951k: 135, 1979c: 72.
- cylindraceus* (Schedl)** 1972c: 287 (*Stephanoderes*). Holotype, sex?; Ghana, Kintampo: Ashanti Region, Ofinso, 259 m, N 6 54, W 1 39; Congo, Brazzaville, Lefinie reservation, Mbeokala forest; Brazzaville, Kindamba, Meya, Bangou forest; Schedl Collection in NHMW, Wien.
Distribution: Africa (Congo/ Ghana).
References: **(ds)** Wood, S. L. 1982b: 905, **(tx)** Schedl 1972c: 287, 1979c: 73; Wood, S. L. 1982b: 905.
- cynometrae* Schedl** 1957d: 50. Holotype, sex?; Congo Belge: Yangambi, Mahiba, 90 km N. Ouest de Tshela; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Cynometra hankui*.
References: **(cc)** Schedl 1961k: 471, **(hb)** Schedl 1961k: 471, **(ds)** Schedl 1961k: 471, **(tx)** Schedl 1957d: 50, 1961k: 471, 1979c: 75.
- delicatus* Schedl** 1964j: 44. Holotype, sex?; Congo: Kivu, Terr. Fizi, Itombwe, 900 m; MRCB, Tervuren.
Distribution: Africa (Zaire).
References: **(tx)** Schedl 1964j: 44, 1979c: 77.
- deprecator* (Schedl)** 1941d: 393 (*Pachynoderes*). Holotype, sex?; Deutsch Ostafrika, Amani; Schedl Collection in NHMW, Wien.
Figures: Schedl 1961k: 632.
Distribution: Africa (Tanzania).
References: **(tx)** Schedl 1941d: 393, 1961k: 632–633, 1979c: 78.
- dexter* (Sampson)** 1922: 141 (*Cryphalus*). Holotype, sex?; Portuguese East Africa: Xinavane; BMNH, London.
Distribution: Africa (Mozambique).
Notes: (1) Browne 1970: 553 (to *Hypothenemus*).
References: **(hb)** Beaver & Loytyniemi 1955a: 80, **(ds)** Beaver & Loytyniemi 1955a: 80; Schedl 1961k: 462, **(tx)** Browne 1970: 553; Sampson 1922: 141; Schedl 1957b: 150, 1961k: 462.
- dimorphus* (Schedl)** 1959c: 168 (*Stephanoderes*). Holotype, sex?; Malaya, Negri Sembilan, Jelelu, Triang For. Res.; Forest Research Institute, Kepong.
Distribution: Asia (Malaya).
Hosts: *Acacia auriculaeformis*.
References: **(cn)** Browne 1968: 353, **(ds)** Browne 1961c: 72, 1968: 353; Schedl 1959c: 168, **(tx)** Schedl 1959c: 168, 1979c: 81.
- dipteroearpi* Hopkins** 1915b: 17. Holotype ♀; Calapan, Mindoro, P.I.; USNM, Washington.
Distribution: Mangareva Islands, Philippine Islands (Mindoro).
Hosts: *Dipteroearpus polosapis*, *Hibiscus tiliaceus*.
References: **(ds)** Browne 1983a: 555, 1983b: 76, 1984c: 449; Kleine 1934a: 143; Olmo, Yoneyama, & Nakazawa 1987b: 93; Schedl 1966b: 24, **(tx)** Hopkins 1915b: 14, 17; Nobuchi 1983: 300; Wood, S. L. 1972f: 45.
- mangarevanus* Beeson** 1940: 196. Holotype ♀; Mangareva Islands: Ankena, northwest side; BPBM, Honolulu. Synonymy: Wood 1972f: 45.
References: **(ds)** Beeson 1935b: 289; Wood, S. L. 1960a: 41, **(tx)** Beeson 1940: 196; Wood, S. L. 1960a: 41, 1972f: 45.
- dissimilis* (Zimmermann)** 1868: 144 (*Crypturgus*). Holotype ♀; North Carolina [USA]; MCZ, Cambridge.
Figures: Blackman 1922b: pl. 5, figs. 29–30.
Distribution: North America (Alabama, Connecticut, District of Columbia, Florida, Georgia, Illinois, Indiana, Louisiana, Kentucky, Maryland, Minnesota, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA).
Hosts: *Accer rubrum*, *Carya* spp., *Cercis canadensis*, *Fagus grandifolia*, *Kalmia latifolia*, *Quercus*

spp., *Ocoeca catesbyana*, *Rhamnus lanceolata*, *Vitis* spp.

References: (**bv**) Kirkendall 1984: 241; Roling & Kearby 1975b, 1977; Turnbow & Franklin 1980. (**cn**) Baker, W. L. 1972: 254; Lintner 1896: 270; Packard 1890: 93; Roling & Kearby 1977; Smith, J. B. 1900: 362. (**cc**) Burks 1979: 856; Bushing 1965: 460; Felt 1906: 702; Roling & Kearby 1977. (**hb**) Baker, W. L. 1972: 254; Beal & Massey 1945: 110–113; Blackman 1922b: 89; Chamberlin 1939: 304; Chittenden 1890, 1893a, 1895: 385, 1897: 393; Deyrup & Atkinson 1987a: 66; Drooz 1985: 367; Felt 1906: 702; Hopkins 1899a: 343; Packard 1890: 93; Pierce, W. D. 1907: 291; Schwarz 1888a: 80; Smith, J. B. 1890b: 54; Wenzel 1905: 124; Wood, S. L. 1953: 123, 1954a: 1021, 1982b: 882. (**ds**) Anonymous 1926c: 518; Beal & Massey 1942, 1945: 110–113; Beaulieu 1956; Blackman 1922b: 89; Blatchley & Leng 1916: 603; Bright 1976d: 114; Britton 1920a; Chamberlin 1939: 304; Chittenden 1890, 1893: 393, 1895b: 385; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Dodge 1938; Drooz 1985: 367; Gemminger & Harold 1872: 2688; Hagedorn 1910d: 41; Hamilton 1895: 346, 378; Henshaw 1885: 148; Hoffmann 1940: 60, 1942: 13; Hopkins 1893a: 133, 1893b: 210; Kirk 1969, 1970; Kleine 1913b: 121, 1914b: 386, 1934a: 145; Leng 1920: 340; Leonard 1928: 518; Lintner 1896: 270; Schwarz 1878d: 468, 1888a: 80; Smith, J. B. 1890a: 276, 1890b: 54, 1900: 362, 1910: 402; Swaine 1909: 116; Turnbow & Franklin 1980; Ulke 1902: 56; Wenzel 1905: 124; Wood, S. L. 1982b: 882. (**tx**) Beal & Massey 1945: 110–113; Blackman 1922b: 89; Blatchley & Leng 1916: 603; Bright 1976d: 114; Chamberlin 1939: 304; Dodge 1938: 39; Eichhoff 1872a: 132, 1878b: 143–144, 1896: 608; Hagedorn 1910a: 86; Hopkins 1915b: 24; LeConte 1868: 154, 1876: 356, 1878a: 468; Schedl 1951m: 101; Sharp 1879: 101; Swaine 1909: 116, 1918a: 40; Wood, S. L. 1953: 123, 1954a: 959, 1017, 1021, 1982b: 882; Zimmerman 1868: 144.

chapuisii Eichhoff 1872a: 132 (*Stephanoderes*). Lectotype ♀; America bor. [USA]; IRSNB, Brussels, designated by Wood 1982b: 882. Synonymy: Eichhoff 1896: 608.

References: (**hb**) Baker, W. L. 1972: 254; Beal & Massey 1945: 112; Blackman 1922b: 90–91; Chamberlin 1939: 305. (**ds**) Beal & Massey 1945: 112; Beeson 1938b: 290; Blackman 1922b: 90–91; Blatchley & Leng 1916: 604; Chamberlin 1939: 305; Drooz 1985: 368; Gemminger & Harold 1872: 2684; Hagedorn 1910d: 41; Kleine 1934a: 145; Knull 1934: 212; Leng 1920: 340; Swaine 1909: 117. (**tx**) Beal & Massey 1945: 112; Blackman 1922b: 90–91; Blatchley & Leng 1916: 604; Chamberlin 1939: 305; Eichhoff 1872a: 132, 1878b: 143, 1896: 608, 610; Hopkins 1914: 130,

1915b: 24; LeConte 1876: 356; Swaine 1909: 117; Wood, S. L. 1954a: 959, 1021.

distinctus Wood 1954a: 1053. Holotype ♀; Union, Missouri [USA]; SMUK, Lawrence. Distribution: North America (Florida, Indiana, Missouri, Oklahoma).

Hosts: *Rhus aromatica*.

References: (**hb**) Wood, S. L. 1953: 214, 1982b: 903. (**ds**) Atkinson et al. 1991: 161; Wood, S. L. 1982b: 903. (**tx**) Wood, S. L. 1953: 214, 1954a: 1053, 1066, 1982b: 903.

dolichocola Hopkins 1915b: 19. Holotype ♀; Canton, China; USNM, Washington. Distribution: Asia (Canton in China).

Hosts: *Dolichos lablab*.

References: (**ds**) Kleine 1934a: 143. (**tx**) Hopkins 1915b: 14, 19.

dolosus Wood 1974a: 21. Holotype ♀; Pandora, Limon, Costa Rica; Wood Collection. Distribution: North America (Costa Rica/ Honduras/ Chiapas, Veracruz in Mexico).

Hosts: *Mimosa* sp., *Swietenia* sp.

References: (**ds**) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 889. (**tx**) Wood, S. L. 1974a: 21, 1982b: 889.

donisi (Schedl) 1957d: 46 (*Eriocryphalus*). Holotype, sex?; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire), Madagascar.

Hosts: *Cynometra hankiei*.

References: (**tx**) Schedl 1957d: 46, 1961k: 455–456; Wood, S. L. 1989: 174.

madagascariensis Schedl 1961e: 131 (*Eriocryphalus*). Holotype, sex?; Madagascar, petite foret pres de Tananarive; IRSM, Madagascar. Synonymy: Wood 1989: 174.

References: (**hb**) Schedl 1977a: 40. (**tx**) Schedl 1961e: 131, 1977a: 40; Wood, S. L. 1989: 174.

dorsosignatus (Schedl) 1950f: 46 (*Stephanoderes*). Holotype, sex?; Fiji Islands, Suva; Eggers Collection, in NHMW, Wien.

Distribution: Fiji Islands.

References: (**ds**) Browne 1974a: 64. (**tx**) Beaver 1991: 92; Schedl 1950f: 46–47, 1979c: 84.

fijianus Schedl 1955b: 289 (*Stephanopodius*). Holotype, sex?; Fiji; Schedl Collection in NHMW, Wien. Synonymy: Beaver 1991: 92.

References: (**ds**) Browne 1974a: 65. (**tx**) Beaver 1991: 92; Schedl 1955b: 289, 1979c: 96.

dubius (Schedl) 1971c: 372 (*Stylotenus*). Holotype, sex?; Ceylon: Rolonnenarawa; Schedl Collection in NHMW, Wien.

Distribution: Asia (Sri Lanka/Vietnam).

References: (**tx**) Schedl 1971c: 372, 1973c: 377, 1979c: 85.

elaphas (Eichhoff) 1872a: 132 (*Stephanoderes*). Holotype, sex?; Isle de France [Eichhoff 1878b: 143 gives St. Mauritius]; IRSNB, Brussels.

- Distribution: Africa (Mauritius Island/ Mozambique/ South Africa/ Tanzania/ Zaire), Madagascar.
References: **(ds)** Alluaud 1900: 439; Gemminger & Harold 1872: 2684; Hagedorn 1910d: 42; Kleine 1913b: 121; Schedl 1961k: 472, 1962b: 58, 1962k: 106S, 1965c: 352. **(tx)** Eggers 1920: 121, 1924: 202, 1932: 31, 1934b: 27, 1943c: 75; Eichhoff 1872a: 132, 1878b: 142–143; Hagedorn 1910a: 52, 86, 1913a: 254; Hopkins 1914: 116, 135, 1915b: 24; Schedl 1939i: 380, 1950c: 203, 1953d: 6S, 1961k: 472–473, 1962h: 58, 1962k: 106S, 1969d: 10, 1977b: 49.
- spinosus* Hagedorn 1909a: 745 (*Adiaeretus*). Syn-types 2, sex?; Caffraria, Transvaal; MNB, Berlin. Synonymy: Schedl 1961k: 472.
References: **(ds)** Hagedorn 1910d: 47; Kleine 1913b, 1914b: 319. **(tx)** Eggers 1924: 102, 1932c: 31, 1934b: 27; Hagedorn 1909a: 745, 1910a: 82; Hopkins 1914: 116; Schedl 1939i: 380, 387, 1961k: 472.
- ° *emmi* (Hagedorn) 1913a: 254 (*Cryphalus*). Syn-types, sex?; Madagascarkopai; Hamburg Museum, lost.
Distribution: Madagascar (fossil in copal).
References: **(ds)** Kleine 1912a: 213; Spahr 1981. **(tx)** Eggers 1919: 240, 1927a: 196; Hagedorn 1913a: 254.
- emmi* Eggers 1919: 240 (*Stephanoderes*). Holotype, sex?; Madagascar-Copal; Hamburg Museum, lost. Synonymy: Eggers 1919: 241.
Notes: (1) Eggers renamed this species from the Hagedorn holotype without realizing that a description had already been published.
References: **(tx)** Eggers 1919: 240–241, 1927a: 196.
- erectus* LeConte 1876: 356. Lectotype ♀; labeled “Texas, [collected by] Belfrage,” presumably from the lower Rio Grande Valley [USA]; MCZ, Cambridge, designated by Wood 1972f: 45.
Figures: Atkinson & Equihua 1988: 95 (adult).
Distribution: “Africa”, Antilles Islands (Cuba/ St. Thomas), North America (Honduras/ Chiapas, Colima, Hidalgo, Jalisco, Nayarit, Nuevo Leon, Puebla, Queretaro, San Luis Potosi, Sinaloa, Tamaulipas, Veracruz in Mexico/ S Texas in USA), South America (Venezuela).
Hosts: *Condalia* sp., *Ficus* sp., *Inga* sp., *Miconia* sp., *Mimosa* spp., *Prosopis* sp., *Rubus* sp., *Serjania* sp., *Toxicodendron* sp., *Verbisina agricolorum*, *Vismia* sp.
Notes: (3) This species was apparently introduced to America from Africa or Madagascar where numerous variants occur.
References: **(cn)** Packard 1890: 93; Smith, J. B. 1900: 362. **(cc)** Equihua & Atkinson 1986: 631; Felt 1906: 702, 740. **(hb)** Burgos & Saucedo 1983: 104; Chamberlin 1939: 310; Chittenden 1890; Equihua & Atkinson 1986: 631; Felt 1906: 702, 740; Packard 1890: 93; Pierce, W. D. 1907: 291; Smith, J. B. 1890b: 54; Wood, S. L. 1953: 132, 1954a: 1026. **(ds)** Atkinson & Equihua 1985c: 356, 1986a: 422, 1988: 92; Blackwelder 1947: 778; Blandford 1904: 227; Blatchley & Leng 1916: 602; Bright 1955c: 174; Burgos & Saucedo 1983: 104; Chamberlin 1939: 310; Chittenden 1890; Equihua & Atkinson 1986: 631; Estrada & Atkinson 1988: 209; Ferrer 1942; Hagedorn 1910d: 42; Henshaw 1882: 269, 1885: 148; Hopkins 1893a: 132, 1893b; Kleine 1913b: 121, 1914b: 350, 386, 1934a: 145; Leng 1920: 340; Smith, J. B. 1890a: 267, 1890b: 54, 1900: 362, 1910: 401; Swaine 1909: 117; Thomas, J. B. 1966; Wood, S. L. 1977a: 71, 1983: 104. **(tx)** Atkinson & Equihua 1988: 95; Blandford 1904: 227; Blatchley & Leng 1916: 602–603; Bright 1955c: 174; Chamberlin 1939: 310; Eichhoff 1878b: 146, 1896: 608; Hagedorn 1910a: 86; Hopkins 1915b: 24; LeConte 1876: 356; Schedl 1940a: 342; Swaine 1909: 117; Wood, S. L. 1953: 132, 1954a: 959, 101S, 1026, 1972f: 45, 1976a: 34S, 1982b: 88S.
- validus* Blandford 1904: 22S. Holotype ♀; Motzorongo, Veracruz, Mexico; BMNH, London. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 778; Ferrer 1942; Hagedorn 1910a: 46; Kleine 1913b: 124, 1914b: 350; Sampson 1914: 385; Wichmann 1955a: 107. **(tx)** Blandford 1904: 22S; Browne 1970: 556; Hagedorn 1910a: 88; Hopkins 1915b: 24; Sampson 1914: 385; Schedl 1940a: 342; Wood, S. L. 1972f: 46.
- puncticollis* Hopkins 1915b: 32 (*Stephanoderes*). Holotype ♀; Tampico, Tamaulipas, Mexico; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 778; Ferrer 1942; Schedl 1963c: 157; Wood, S. L. 1961c: 1. **(tx)** Hopkins 1915b: 25, 32; Schedl 1940a: 342, 1940c: 206; Wood, S. L. 1961c: 1, 1972f: 46.
- cubensis* Hopkins 1915b: 32 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 777; Leng & Mutchler 1917: 219. **(tx)** Eggers 1940a; Hopkins 1915b: 24, 32; Schedl 1951m: 103; Wood, S. L. 1972f: 46.
- brunneicollis* Hopkins 1915b: 33 (*Stephanoderes*). Holotype ♀; Round Mountain, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1026.
References: **(hb)** Chamberlin 1939: 310. **(ds)** Chamberlin 1939: 310; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Chamberlin 1939: 310; Hopkins 1915b: 24, 32–33; Wood, S. L. 1954a: 959, 1026.
- discedens* Schedl 1950d: 23 (*Stephanoderes*). Holotype ♀; St. Thomas; Schedl Collection in NHMW, Wien. Synonymy: Wood 1976a: 34S.
References: **(tx)** Schedl 1950d: 23, 1961k: 562; Wood, S. L. 1976a: 34S.

eruditus Westwood 1836: 34. Syntypes ♀: England?; some in BMNH, London.
 Figures: Balachowsky 1949a: 203–204, Grune 1979: 112, Masutti 1968: 365, Pedrosa-Macedo & Schonherr 1985: 47.
 Distribution: Africa (Algeria/ Angola/ Cameroon/ Canary Islands/ Egypt/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Morocco/ Nigeria/ Seychelles Islands/ Sierra Leone/ South Africa/ Tanzania/ Togo/ Uganda/ Zaire), Antilles Islands (Cuba/ Guadeloupe/ Jamaica/ Puerto Rico/ Trinidad), Asia (Bonin Islands/ Burma/ Andaman Islands, Assam, Punjab, Uttar Pradesh in India/ Iran/ Israel/ Japan/ Malaya/ Sri Lanka/ Thailand/ Vietnam), Australia (Queensland), Cook Islands, Europe (Corsica/ England/ France/ Italy/ Spain), Fiji Islands, Hawaiian Islands, Indonesia (Java/ Sumatra), Madagascar, Marquesas Islands, Micronesia, New Caledonia, Niue Island, North America (Costa Rica/ Guatemala/ Honduras/ Chiapas, Colima, Jalisco, Michoacan, Nayarit, Oaxaca, Puebla, San Luis Potosi, Veracruz, Yucatan in Mexico/ Panama/ Alabama, Arkansas, California, District of Columbia, Florida, Georgia, Illinois, Louisiana, Maryland, Michigan, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA), Philippine Islands.
 Hosts: Numerous hosts recorded by Beaver 1976b: 536, 1979b: 295, Browne 1961c: 76, Schedl 1961k: 488–498, Wood 1954a, 1982b: 902–903.
 Notes: (3) Schedl 1965g: 19 (*Styphnaderes phascoli* Eggers, nomen nudum, synonymy).
 References: (ay) Van Ryn-Tourmel 1975. (bv) Gray, B. 1974c; Grune 1979: 113; Schedl 1960f: 11; Turnbow & Franklin 1980. (cn) Anonymous 1970c: 13; Arruda 1965; Box 1953a; Brader 1964: 5; Browne 1961c: 76, 1968: 353; Chittenden 1893: 250; Corbet 1931: 351–355; Currie 1905: 7; Downe & Williams 1950; Ebeling 1950: 353, 533; Equihua 1988; Hagedorn 1913a; Hammad 1961; Hargreaves 1927: 24–27; Hubbard 1885: 173; Kleine 1932c: 302; Kumar, Jayaraj, & Mnthukrishnan 1979; Murillo Quinche 1959: 300; Petri 1912; Roark 1939: 308; Souphieff & Scherbinovskaja 1937: 81; Tsankov 1977; Vuillaume et al. 1981; Zacher 1913: 131–144. (ec) Ashmead 1883a, 1893: 451; Beaver 1979b: 298; Bushing 1965: 460; Chittenden 1893b; Equihua & Atkinson 1986: 632; Felt 1906: 725; Halperin & Holzschuh 1984: 26; Kleine 1908c: 187; Mendel 1986c: 115; Schedl 1958d: 190; Vuillaume et al. 1981; Wichmann 1955a: 105. (hb) Ashe 1949; Atkinson et al. 1986: 64; Baker, W. L. 1972: 254; Beaver 1974a, 1976b: 536, 1979b: 295, 1987a: 19; Beaver & Browne 1978: 590; Blandford 1894b; Bodenheimer 1930; Bright & Stark 1973: 69; Browne 1961c: 76, 1968: 353; Burgos & Saucedo 1983: 108; Deyrup & Atkinson 1987a: 66;

Equihua & Atkinson 1986: 632; Felt 1906: 725; Gardner 1957: 31; Gray, B. 1974c; Hagedorn 1913a; Halperin & Holzschuh 1984: 26; Hammad 1961; Hubbard 1885: 173; Kalshoven 1932: 242, 1951: 856, 1958b: 170–172, 179, 1963; Kleine 1932a: 302; Lengerken 1939: 64, 1954: 84; Loytyniemi, Beaver, & Loytyniemi 1984; Masutti 1968; Murillo Quinche 1959: 300; Pierce, W. D. 1907: 291; Pollet 1977; Postner 1974: 432; Roberts 1969: 66, 86; Schedl 1960f: 86, 1961k: 474; Schwarz 1889: 139, 1891e: 73–74; Smith, J. B. 1890b: 54; Wachtl 1876a: 453; Wood, S. L. 1953: 198, 1954a: 1058, 1982b: 900; Yunus & Hua 1980: 229; Zacher 1927: 161. (ds) Alkan 1964: 374, 390; Anonymous 1970c: 13; Atkinson & Equihua 1985c: 357, 1986a: 422, 1988: 92; Atkinson et al. 1986: 64; Aulmann 1912: 55–56; Barber 1919: 53–60; Beaver 1974a, 1976b: 536, 1987a: 19, 1987b: 65, 1990a: 280; Beaver & Browne 1975: 590; Beeson 1935: 103, 1938b: 289, 1940: 196–197, 1941: 374; Blackwelder 1947: 777; Blandford 1894b: 261–264, 1897b, 1904: 229–230; Bright 1972d: 50, 1985c: 175; Bright & Stark 1973: 69; Brimley 1935: 247; Browne 1963: 53, 1966: 245, 1968b: 353–354, 1968c: 111, 1974a: 64, 1976, 1984a: 150; Burgos & Saucedo 1983: 108; Bytinski-Salz 1966: 38; Cachan 1957: 15; Calver 1884, 1893; Chobant 1897: 262; Choo 1983: 65; Choo & Woo 1985: 164; Choo, Woo, & Nobuchi 1983: 172; Crotch 1863, 1866: 18; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Drooz 1985: 368; Equihua & Atkinson 1986: 632; Estrada & Atkinson 1988: 210; Fairmaire 1863: 33; Ferreira 1965: 1113; Ferrer 1942; Fowler 1891; Fowler & Donisthorpe 1913: 199; Gardner 1957a; Ganil 1849: 127; Gemminger & Harold 1872: 2679; Ghesquiere 1933a: 32–36, 1933b: 783–786; Gray 1967: 9; Grune 1979: 113; Hagedorn 1903b: 546, 1910d: 42, 1913a: 44–45; Halperin & Holzschuh 1984: 26; Hamad 1961: 150; Hamilton 1859: 158, 1894b: 406, 1895a: 346, 378; Hargreaves 1948; Henshaw 1885: 148; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Hopkins 1893a: 132; Israelson 1972; Kalshoven 1963: 233; Kirk 1970; Kirkendall 1984: 242; Kleine 1913b: 124, 1914a: 15, 22, 1914b: 300, 308, 350, 361, 408, 1932a: 143, 302, 1934: 142–146; Kloft & Hinks 1945: 218; Krauss 1943: 88; Lacordaire 1866: 374; Lepesme 1944: 241, 1947: 645–646; Lintner 1896: 270; Lundblad 1958: 489; Marcu 1934: 57; Martorell 1945: 469; Masutti 1968: 360–370; Matthews & Fowler 1883: 42; Mayne & Donis 1951: 332–333; Miller, N. C. E. 1934; Moore 1964: 300; Munro 1946: 15; Newberry 1910: 83; Nobuchi 1985c: 16; Nobuchi & Ono 1973: 181; Nohira & Ogawa 1986; Nunnberg 1952: 18, 1961b: 609, 1965b: 18, 1971: 58, 1972b: 195; Ohno, Yoneyama, & Nakazawa 1982b: 7; Ohno, Yoskioka et al. 1985a: 92; Pedrosa-Macedo & Schonherr 1985: 47; Peyerimhoff

- 1911: 314, 1912: 173, 1915: 60, 1919: 225, 1935: 195; Pollet 1977; Postner 1974: 432; Redtenbacher 1858: 831, 1874: 374; Reitter 1894a: 75; Ross 1919: 83; Rye 1866a: 198, 1890: 270; Sampson 1914: 385; Sankeral & Rao 1966: 67; Schamm 1859: 96, 1862: 101; Schedl 1934f: 1640, 1959a: 478, 1959p: 16, 1960d: 76, 1960f: 11, 1961b: 187, 1961c: 69, 1961e: 131, 1961j: 349, 1961k: 474–476, 1961m: 84, 1962b: 58, 1962k: 1068, 1963c: 157, 1963f: 53, 1964a, 1964c: 305, 1966b: 24, 1969g: 290; Schedl, Lindberg, & Lindberg 1959: 16; Schonherr & Pedrosa-Macedo 1981: 52; Scudder 1864: 14, 1865: 13–14; Seidlitz 1891: 151; Smith, J. B. 1890b: 54, 1892: 43, 1900: 362, 1910: 401; Souphieff & Scherbinovskaja 1937: 81; Stein & Weise 1877: 164; Stephens 1830: 418, 1839: 209; Sniew 1937: 65–68; Swaine 1909: 117–118; Swezey 1941: 117–118; Terra 1987: 19; Turnbow & Franklin 1980; Ulke 1902: 56; Wachtl 1876a: 453; Wichmann 1927: 79, 1927: 379, 1954: 509–510, 522, 1955a: 105; Wolcott 1945: 379; Wollaston 1864: 256, 1865: 239, 1868: 116–117; Wood, S. L. 1957e: 1273, 1960a: 41, 1977a: 71, 1982b: 900; Yunus & Hua 1980: 229. **(tx)** Balachowsky 1943: 168, 1949a: 202–205; Beal & Massey 1945: 117–118; Blackman 1922b: 6–104; Blandford 1894b: 261–264, 1904: 229–230; Blatchley & Leng 1916: 598; Bright 1972d: 50, 1985c: 175; Castelnau 1840; Chamberlin 1939: 288–289; Choo 1983: 65; Costa Lima 1925: 118, 1936: 356; Dodge 1938: 39; Duffy 1953: 3, 6; Eggers 1919: 241, 1921: 40, 1922: 86, 1924: 110, 1925: 153, 1927: 173, 1929: 55, 1936: 126; Eichhoff 1864b: 34, 1868d: 421, 1878a: 387, 1878b: 165, 1883: 134, 1896: 608; Erichson 1836: 61; Fauvel 1884a: 390, 1884b: 315, 1889, 1897: 66; Ferrari 1867a: 7, 17, 89, 1867b: 113, 1868: 255–256; Grune 1979: 112–113; Hagedorn 1903b: 546, 1909: 744, 1910a: 85–86; Hammad 1961; Hopkins 1914: 123, 133, 1915b: 12, 14, 1915c: 221–222; Houlbert 1922a: 13; Jacquelin du Val & Fairmaire 1868: 105; Lacordaire 1866: 374; LeConte 1876: 355–356; Masutti 1968: 365; Miller, N. C. E. 1934: 27; Murayama 1936: 126; Newberry 1910: 83; Niisima 1913: 1; Nobuchi 1983: 300; Numberg 1952: 18, 1956: 144; Pedrosa-Macedo & Schonherr 1985: 47; Perkins 1900: 173–180; Peyerimhoff 1935: 195–196; Portevin 1935: 325; Postner 1974: 432; Redtenbacher 1858: 831, 1874: 374; Reitter 1887: 194, 1894a: 75, 1902: 141, 1906: 711, 1911: 711, 1913a: 73–74; Sampson 1914: 385; Schedl 1934g: 178, 1935j: 1, 1939e: 328, 1939f: 32, 1939g: 170, 1940a: 322, 342, 1940c: 207, 1941f: 110, 1942a: 170, 1942b, 1942c: 174, 1942d: 12–13, 1950c: 204, 1950d: 3–20, 1950g: 895, 1951f: 40, 1951j: 19, 1951k: 134, 1952a: 449, 1952g: 49–50, 1952j: 2, 1953b: 124, 1953c: 288, 1953d: 68, 1954a: 139, 1954d: 871, 1954e: 50, 70, 1957a: 191–192, 1957b: 150, 1957d: 10–11, 1958b: 99, 1958d: 189–190, 1959a: 478, 1960a: 76, 1961b: 187, 1961k: 474, 1963e: 155, 1963h: 266, 1964a: 96–97, 1964c: 305, 1964e: 65, 1964i: 248, 1964j: 40, 1964k: 222, 1965a: 399, 1965f: 4, 1965g: 19, 22, 1966b: 24, 1966f: 80, 83, 1967e: 210, 1968e: 261–262, 1969d: 8–11, 1969e: 261–262, 1969g: 286, 290, 1969i: 140, 1970e: 80, 91, 1971a: 275, 280, 1971c: 366, 1971e: 2, 1971f: 147, 1977b: 51; Schedl, Lindberg, & Lindberg 1959: 16; Sharp 1879: 101–102; Stark 1952: 292; Stephens 1830: 418, 1839: 209–210; Swaine 1909: 117–118; Terra 1987: 19; Tredl 1907: 13; Westwood 1836: 34–36, 1839: 353, 1840: 157; Wood, S. L. 1953: 198, 1954a: 959, 1058–1061, 1960a: 41–44, 1972f: 46, 1974d: 282, 1977d: 513, 1982b: 900; Wylie & Yule 1977. **(ms)** Eichhoff 1868d: 421. *aspericollis* Wollaston 1860a: 365 (*Cryphalus*).
 Syntypes, sex?, Madera; BMNH, London.
 Synonymy: Schedl 1960h: 16.
 References: **(ay)** Peyerimhoff 1912: 173. **(cn)** Ebeling 1959; Garcia-Tejero 1955: 234; Gentry, J. W. 1975: 105; Grandi 1951; Hammad 1961; Souphieff & Scherbinovskaja 1937: 83. **(hb)** Balachowsky 1963a: 1280; Davatchi 1955; Eichhoff 1878b: 162; Grandi 1951; Hammad 1961; Peyerimhoff 1915: 19. **(ds)** Balachowsky 1963a: 1280; Beeson 1940: 196–197, 1961: 288; Davatchi 1958: 76; Ebeling 1959; Eichhoff 1878b: 162; Escalera 1919; Fauvel 1897; Gemminger & Harold 1872: 2683; Gentry, J. W. 1965: 105; Hagedorn 1910d: 41; Jansson 1940: 63; Lacordaire 1866: 379; Lundblad 1958: 489; Newberry 1910: 84; Peyerimhoff 1912: 173, 1935: 194–195; Schmitz 1898: 157; Sharp 1879: 102; Souphieff & Scherbinovskaja 1937: 83. **(tx)** Balachowsky 1949a: 202, 1963a: 1280; Beeson 1940; Eggers 1921: 40; Eichhoff 1878b: 162; Fauvel 1884; Hagedorn 1910a: 85; Hammad 1961; Hopkins 1915b: 13; Lacordaire 1866: 379; Newberry 1910: 84; Peyerimhoff 1911: 314, 1912: 173–174, 1935: 194; Schedl 1934f: 1640, 1951a: 449, 1960h: 16; Schedl, Lindberg, & Lindberg 1959: 16; Sharp 1879: 102; Sokanovskii 1954: 17; Wollaston 1860a: 365, 1864: 256, 1865: 239. *boieldieu* Perrond 1864: 188 (*Bostrichus*). Holotype ♀; Kanala, New Caledonia; not located. Synonymy: Eichhoff 1878b: 166.
 References: **(ds)** Gemminger & Harold 1872: 2690; Swaine 1909: 118. **(tx)** Blandford 1904: 229–230; Eichhoff 1878b: 166; Fauvel 1884b; Ferrari 1867a: 75; Hopkins 1915c: 222; Perrond 1864: 188; Swaine 1909: 118.
obscurus Ferrari 1867a: 17 (*Cryphalus*). Holotype ♀; Cuba; NHMW, Wien, preoccupied by Fabricius 1801. Synonymy: Wood 1974d: 282. References: **(ds)** Browne 1965: 187; Hagedorn 1910d: 44; Kalshoven 1964: 233; Kleine 1914b: 378. **(tx)** Eggers 1929e: 50–51, 1940a; Eichhoff 1878b: 163; Ferrari 1867a: 17; Hage-

- dorn 1910a: 87; Hopkins 1915b: 13; Wood, S. L. 1974d: 282.
- geruari* Eichhoff 1875b: 159 (*Stephanoderes*). Syntypes, sex?; America borealis (Mexico); Hamburg Museum, lost. Synonymy: Wood 1977d: 513.
- References: (**hb**) Blackman 1922b: 83–84; Chamberlin 1939: 290. (**ds**) Blackman 1922b: 83–84; Blackwelder 1947: 778; Blandford 1904: 127–128; Chamberlin 1939: 290; Ferrer 1942; Hagedorn 1910d: 43; Kleine 1913b: 121, 1914b: 349; Leng & Mutchler 1927: 52. (**tx**) Blackman 1922b: 83–84; Blandford 1904: 127–128; Chamberlin 1939: 290; Eichhoff 1878a: 386, 1878b: 159; Hagedorn 1910a: 86; Hopkins 1915b: 13; Schedl 1940a: 342; Wood, S. L. 1954a: 1058, 1977d: 513.
- myrmedon* Eichhoff 1875b: 160 (*Stephanoderes*). Holotype ♂; Colombia: IRSNB, Brussels. Synonymy: Wood 1977d: 513.
- References: (**ds**) Blackwelder 1947: 778; Hagedorn 1910d: 44; Kleine 1913b: 121, 1914b: 342. (**tx**) Eichhoff 1865, 1878a: 386, 1878b: 160; Hagedorn 1910a: 87; Hopkins 1915b: 21; Reitter 1902a: 141; Wood, S. L. 1977d: 513.
- chlersi* Eichhoff 1878b: 493 (*Stephanoderes*). Syntypes, sex?; Spain; Hamburg Museum, lost. Synonymy: Balachowsky 1949a: 202.
- References: (**cn**) Anonymous 1980g; Widiez 1932: 415–445. (**hb**) Eichhoff 1881a: 47, 192; Peyerimhoff 1915: 60, 1919: 255. (**ds**) Anonymous 1980g; Balachowsky 1943a; Barthe 1896; Beeson 1940: 197; Davatchi 1955: 78; Fauvel 1885; Hagedorn 1910d: 42; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Kleine 1913b: 121, 1914a: 19, 1934a: 145; Murayama 1954b: 162; Peyerimhoff 1919: 255, 1935: 195–196; Pittioni 1943: 176; Ragusa 1924: 116; Reitter 1894a: 74; Sainte-Claire & Mequignon 1938: 446; Stein & Weise 1877: 164; Tredl 1907: 13. (**tx**) Balachowsky 1926: 263–264, 1943a: 165, 1949a: 202; Eggers 1940c: 233–234; Eichhoff 1878a: 387, 1878b: 493–495, 1881a: 47, 192, 1883a: 110, 134; Fauvel 1884a: 390, 1884b: 315, 1885; Hagedorn 1910a: 86; Hopkins 1914: 122, 1915b: 13, 1915c: 222; Murayama 1954b: 162; Niisima 1910a: 9; Portevin 1935: 325; Reitter 1887b: 194, 1894a: 74, 1913a: 73; Schedl 1934f: 1640, 1961k: 472.
- communis* Schaufuss 1891: 11 (*Stephanoderes*). Holotype ♀; Madagascar; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992a: 90.
- Notes: (1) Schedl 1977b: 70 (head of type missing).
- References: (**hb**) Beaver 1987a: 18. (**ds**) Alluaud 1900: 439; Beaver 1987a: 18; Hagedorn 1910d: 41; Kleine 1914b: 326; Sampson 1914: 384; Schedl 1969d: 11, 1977b: 70. (**tx**) Hagedorn 1910a: 86; Sampson 1914: 384; Schaufuss 1891: 11; Schedl 1977b: 70–71; Wood, S. L. 1992a: 90.
- insularis* Perkins 1900: 181. Syntypes, sex?; Kauai, Hawaiian Islands; BMNH, London. Synonymy: Wood 1960a: 42.
- References: (**ds**) Beeson 1938b: 289; Hagedorn 1910d: 43; Kleine 1913b: 124, 1914b: 302; Krauss 1943: 88; Swezey 1940: 169, 1941: 117, 1954: 14, 83. (**tx**) Hagedorn 1910a: 86; Perkins 1900: 181; Schedl 1934g: 177–178, 1939f: 32, 1941f: 109, 1942b: 147, 1951k: 134, 1951m: 71, 1958f: 35, 1966b: 24; Wood, S. L. 1960a: 42–43.
- tectonae* Stebbing 1903a: 263 (*Cryphalus*). Syntypes ♀; India. Berar, Melghat teak forests; FRI, Dehra Dun. Synonymy: Wood 1989: 174.
- References: (**cn**) Mathur & Singh 1961b: 13; Stebbing 1903a: 263. (**hb**) Stebbing 1903a: 263. (**ds**) Beeson 1941 (1961: 288); Hagedorn 1910d: 45; Kleine 1914b: 269, 1934a: 143; Mathur & Singh 1961b: 13; Stebbing 1903a: 263. (**tx**) Beeson 1941 (1961: 288); Hagedorn 1910a: 88; Stebbing 1903a: 263, 1914: 536; Wood, S. L. 1989: 174.
- striatopunctatus* Lea 1910: 142 (*Cryphalus*). Syntypes, sex?; National Park, Sidney, New South Wales, Australia; SAM, Adelaide. Synonymy: Wood 1989: 174.
- References: (**ds**) Schedl 1936f: 527. (**tx**) Lea 1910: 141–142; Schedl 1938f: 45, 1939f: 32; Wood, S. L. 1989: 174.
- tantillus* Lea 1910: 142 (*Cryphalus*). Holotype, sex?; N.S. Wales: Richmond River, Australia; SAM, Adelaide. Synonymy: Wood 1989: 174.
- Notes: (1) Schedl 1938f: 44 (to *Hypothenemus*).
- References: (**ds**) Brimblecombe 1953: 20. (**tx**) Lea 1910: 142; Schedl 1938f: 44, 1942a: 169–170, 1942c: 163, 1948g: 25; Wood, S. L. 1989: 174.
- basjoo* Niisima 1910a: 9 (*Cryphalus*). Syntypes, sex?; Tokyo, Japan; Nobuchi Collection, Ibaraki. Synonymy: Wood 1972f: 46.
- References: (**ds**) Beeson 1938: 288; Kleine 1934a: 142; Murayama 1936a: 126, 1953a: 9, 1954b: 162. (**tx**) Eggers 1929e; Murayama 1936a: 126, 1953a: 9, 1954b: 162; Niisima 1910a: 9, 1929: 55; Schedl 1934f: 1640, 1961k: 474; Wood, S. L. 1960a: 41, 1972f: 46.
- tuberculosis* Hagedorn 1912a: 339. Syntypes, sex?; Congo; Hamburg Museum, lost. Synonymy: Wood 1989: 174.
- Notes: (3) Schedl 1939g: 170 (*pterygosperma*, nomen nudum).
- References: (**cn**) Chesquiere 1933a; Kleine 1932a: 302; Mayne & Donis 1951: 332. (**cc**) Aulmann 1913: 1–126; Mayne 1915: 577–596; Mayne & Donis 1951: 332. (**hb**) Kleine 1932a: 302. (**ds**) Chesquiere 1933a; Kleine 1913b: 124, 1914b: 314, 1932a: 302, 1934a: 144; Lepesme et al. 1948: 645; Thompson, C. H.

- 1963: 45. **(tx)** Eggers 1924: 110, 1927a: 173; Hagedorn 1912a: 339; Schedl 1939g: 170, 1950d: 7, 17, 1951f: 40, 1951j: 19, 1952g: 50, 1952j: 2, 1954d: 871, 1954e: 49–50, 1957d: 10–11, 1961k: 475; Wood, S. L. 1959: 174.
- schwarzii* Hopkins 1915b: 11 (*Cosmoderes*). Holotype ♀; Hav Creek, Florida [USA]; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(bv)** Schedl 1960f: 29, **(hb)** Wood, S. L. 1953: 215, 1954a: 1066. **(ds)** Blatchley & Leng 1916: 593; Leng 1920: 339; Schedl 1960f: 29, 1961k: 502. **(tx)** Blatchley & Leng 1916: 593; Chamberlin 1939: 287; Hopkins 1915b: 11; Schedl 1961k: 502; Wood, S. L. 1953: 215, 1954a: 959, 1066, 1972f: 46.
- bradfordi* Hopkins 1915b: 15. Holotype ♀; Honolulu, Hawaii; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Beeson 1938b: 289, **(tx)** Hopkins 1915b: 13, 15; Wood, S. L. 1972f: 46.
- flavosquamosus* Hopkins 1915b: 15. Holotype ♀; Mount Coffee, Liberia, West Africa; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Kleine 1934a: 143, **(tx)** Hopkins 1915b: 13, 15; Schedl 1961k: 502; Wood, S. L. 1972f: 46.
- asiminae* Hopkins 1915b: 16. Holotype ♀; Plummers Island, Maryland [USA]; USNM, Washington. Synonymy: Wood 1954a: 1058. References: **(hb)** Blackman 1922b: 85; Chamberlin 1939: 291. **(ds)** Blackman 1922b: 85; Blatchley & Leng 1916: 597; Chamberlin 1939: 291; Kleine 1934a: 143; Leng 1920: 340. **(tx)** Blackman 1922b; Blatchley & Leng 1916: 597; Chamberlin 1939: 291; Hopkins 1915b: 13, 16; Schedl 1961k: 475; Wood, S. L. 1954a: 979, 1058.
- hamamelidis* Hopkins 1915b: 16. Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1058. References: **(hb)** Chamberlin 1939: 294. **(ds)** Blatchley & Leng 1916: 597; Chamberlin 1939: 294; Kleine 1934a: 143; Leng 1920: 340. **(tx)** Blatchley & Leng 1916: 597; Chamberlin 1939: 294; Hopkins 1915b: 13, 16; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- myristicac* Hopkins 1915b: 16. Holotype ♀; Buitenzorg, Java; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(hb)** Kalshoven 1958a: 172. **(ds)** Browne 1961c: 78; Kleine 1934a: 143; Schedl 1959a: 479. **(tx)** Hopkins 1915b: 14, 16; Schedl 1939d: 408, 1939f: 31, 1942d: 2, 1958b: 99, 1959a: 479; Wood, S. L. 1972f: 46.
- nigricollis* Hopkins 1915b: 16. Holotype ♀; Capetown, South Africa; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Kleine 1934a: 143. **(tx)** Hopkins 1915b: 13–19; Schedl 1961k: 475; Wood, S. L. 1972f: 46.
- pruni* Hopkins 1915b: 16. Holotype ♀; Tryon, North Carolina [USA]; USNM, Washington. Synonymy: Wood 1954a: 1058. References: **(hb)** Beal & Massey 1945: 119; Chamberlin 1939: 294. **(ds)** Beal & Massey 1945: 119; Blatchley & Leng 1916: 597; Chamberlin 1939: 294; Kleine 1934a: 143; Leng 1920: 340. **(tx)** Beal & Massey 1945: 119; Blatchley & Leng 1916: 597; Chamberlin 1939: 294; Hopkins 1915b: 13, 16; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- runseji* Hopkins 1915b: 16. Holotype ♀; Little Falls, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1058. References: **(hb)** Beal & Massey 1945: 117; Blackman 1922b; Chamberlin 1939: 290. **(ds)** Beal & Massey 1945: 117; Blackman 1922b: 85; Blatchley & Leng 1916: 597; Chamberlin 1939: 290; Hopkins 1915b: 13, 16; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- temis* Hopkins 1915b: 16. Holotype ♀; Trece Aguas, Alta Verapaz, Guatemala; USNM, Washington. Synonymy: Wood 1972f: 47. References: **(ds)** Blackwelder 1947: 777; Kleine 1934a: 144. **(tx)** Hopkins 1915b: 14, 16; Wood, S. L. 1972f: 47.
- kochelei* Hopkins 1915b: 17. Holotype ♀; Brazil; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Blackwelder 1947: 777. **(tx)** Hopkins 1915b: 14, 17; Wood, S. L. 1972f: 46.
- lineatifrons* Hopkins 1915b: 17. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Blackwelder 1947: 777; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 14, 17; Wood, S. L. 1972f: 46.
- mdli* Hopkins 1915b: 17. Holotype ♀; Capetown, South Africa; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Kleine 1934a: 143. **(tx)** Hopkins 1915b: 14, 17; Schedl 1961k: 508; Wood, S. L. 1972f: 46.
- parvus* Hopkins 1915b: 17. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 46. References: **(ds)** Blackwelder 1947: 777; Kleine 1934a: 143; Leng & Mutchler 1917: 219; Martorell 1945: 469. **(tx)** Hopkins 1915b: 14, 17; Wood, S. L. 1972f: 46.
- sacchari* Hopkins 1915b: 17. Holotype ♀; Nevis, West Indies; USNM, Washington. Synonymy: Wood 1972f: 47. References: **(ds)** Blackwelder 1947: 777; Kleine 1934a: 144; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 14, 17; Wood, S. L. 1972f: 47.
- webbi* Hopkins 1915b: 17. Holotype ♀; Calapan,

- Mindoro, Philippine Islands; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Kleine 1934a: 144; Schedl 1966b: 25. **(tx)** Hopkins 1915b: 14, 17; Nobuchi 1983: 300; Wood, S. L. 1972f: 47.
- flavipes* Hopkins 1915b: 18. Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 777; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 14, 18; Wood, S. L. 1972f: 46.
- punctifrons* Hopkins 1915b: 18. Holotype ♀; Lakeland, Florida [USA]; USNM, Washington. Synonymy: Wood 1954a: 1059.
References: **(hb)** Blackman 1922b: 88; Chamberlin 1939: 291. **(ds)** Blackman 1922b: 88; Blatchley & Leng 1916: 598; Chamberlin 1939: 291; Dodge 1938: 39; Hoffmann 1942: 13; Kaston 1938: 240; Kleine 1934a: 144; Leng 1920: 340. **(tx)** Blackman 1922b: 88; Blatchley & Leng 1916: 598; Chamberlin 1939: 291; Dodge 1938: 39; Hopkins 1915b: 14, 18; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- nigripennis* Hopkins 1915b: 19. Holotype ♀; Tallulah, Louisiana [USA]; USNM, Washington. Synonymy: Wood 1954a: 1059.
References: **(hb)** Blackman 1922b: 86–87; Chamberlin 1939: 291. **(ds)** Blackman 1922b: 86–87; Blatchley & Leng 1916: 598; Chamberlin 1939: 291; Kirk 1960a; Kleine 1934a: 143; Leng 1920: 340. **(tx)** Blackman 1922b: 86–87; Blatchley & Leng 1916: 598; Chamberlin 1939: 291; Hopkins 1915b: 14, 19; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058–1059.
- ferrugineus* Hopkins 1915b: 20. Holotype ♀; Trece Agnas, Alta Verapaz, Guatemala; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 777. **(tx)** Blackman 1922b: 87; Hopkins 1915b: 14, 20; Wood, S. L. 1972f: 46.
- heathi* Hopkins 1915b: 20. Holotype ♀; Independencia, Parahyba, Brazil; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Blackwelder 1947: 777. **(tx)** Hopkins 1915b: 14, 20; Schedl 1952e: 450; Wood, S. L. 1972f: 46.
- punctipennis* Hopkins 1915b: 20. Holotype ♀; Capetown, "West" Africa; USNM, Washington. Synonymy: Wood 1972f: 46.
References: **(ds)** Kleine 1934a: 144; Schedl 1961k: 510. **(tx)** Hopkins 1915b: 15, 20; Schedl 1961k: 510; Wood, S. L. 1972f: 46.
- flavicollis* Hopkins 1915b: 24 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Blackwelder 1947: 778; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 21, 24; Wood, S. L. 1972f: 47.
- pygmaeus* Hopkins 1915b: 24 (*Stephanoderes*). Holotype ♀; Pagbilao, P.I.; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Schedl 1966b: 29. **(tx)** Hopkins 1915b: 21, 24; Nobuchi 1983: 300; Wood, S. L. 1972f: 47.
- elongatus* Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Blackwelder 1947: 778; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 21, 25; Wood, S. L. 1972f: 47.
- subconcentralis* Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Blackwelder 1947: 778; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 21, 25; Wood, S. L. 1972f: 46.
- unicolor* Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 47.
References: **(ds)** Blackwelder 1947: 778; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 21, 25; Wood, S. L. 1972f: 47.
- evonymi* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1059.
References: **(hb)** Chamberlin 1939: 303. **(ds)** Chamberlin 1939: 303; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Blatchley & Leng 1916: 600; Chamberlin 1939: 303; Hopkins 1915b: 22, 26; Schedl 1961k: 475; Wood, S. L. 1954a: 1058.
- bicolor* Eggers 1919: 241. Lectotype ♀; Amami, East Africa; USNM, Washington, designated by Anderson & Anderson 1971: 6. Synonymy: Schedl 1957d: 11.
References: **(cn)** Souphieff & Scherbinovskaya 1937: 81. **(ds)** Kleine 1934a: 143; Souphieff & Scherbinovskaja 1937: 81. **(tx)** Anderson, W. H. & Anderson 1971: 6; Eggers 1919: 241, 1924: 110.
- chlersi rottroni* Peyerimhoff 1919: 255. Syntypes, sex?; Sidi-bel-Abbes, Oran, North Africa; MNHN, Paris. Synonymy: Balachowsky: 1949a: 202.
References: **(tx)** Balachowsky 1943: 168. 1949a: 202; Peyerimhoff 1919: 255, 1938: 195; Schedl 1934f: 1640, 1961k: 475.
- juglandis* Blackman 1922b: 88. Syntypes, sex?; Port Gibson, Mississippi [USA]; USNM, Washington. Synonymy: Wood 1954a: 1058.
References: **(hb)** Chamberlin 1939: 292. **(ds)** Chamberlin 1939: 292; Kleine 1934a: 243; Leng & Mutchler 1927: 52. **(tx)** Blackman 1922b: 88; Chamberlin 1939: 292; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- pusillus* Eggers 1927a: 173. Lectotype ♀; Belg.-Congo (Mayumbe); USNM, Washington, designated by Anderson & Anderson 1971: 27. Synonymy: Wood 1989: 174.

- References: **(bv)** Schedl 1960f. **(cn)** Ghesquiere 1933a: 32–36, 1933b: 753–756; Magnin 1954: 468; Mayne & Donis 1951: 331; Schedl 1960f: 77; Souplieff & Scherbinovskaja 1937: 51. **(ec)** Cachan 1957: 15; Entwistle 1963a: 57; Mayne & Donis 1951: 331; Schedl 1955d: 190, 1961k: 511. **(hb)** Jones, Roberts, & Baker 1959: 23; Paulian 1951: 27–31; Schedl 1958d: 190, 1961k: 511. **(ds)** Ghesquiere 1933a: 32–36; Kleine 1934a: 144; Mayne & Donis 1962: 307; Schedl 1960f, 1961k: 511; Souplieff & Scherbinovskaja 1937: 51; Thompson, G. H. 1963: 47. **(tx)** Anderson, W. H. & Anderson 1971: 27; Eggers 1927a: 173; Mayne & Donis 1962: 307; Schedl 1950c: 204, 1950d: 7, 17, 1951f: 40, 1952j: 2, 1953d: 68, 1954d: 570, 1954e: 49, 1955f: 258, 1957c: 324, 1957d: 11, 1961k: 511, 1979c: 206; Wood, S. L. 1989: 174.
- intersetosus* Eggers 1928c: 85 (*Stephanoderes*). Lectotype ♀; Brazil, Sao Paulo (Umgegend der Stadt); USNM, Washington designated by Anderson & Anderson 1971: 16. Synonymy: Wood 1972f: 47. References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956; Viana 1964: 122. **(ds)** Blackwelder 1947; Bright 1972d: 53; Costa Lima 1936, 1956; Kleine 1934a: 145; Schedl 1960a: 76; Viana 1964: 122. **(tx)** Anderson, W. H. & Anderson 1971: 16; Bright 1972d: 53, 1973: 18; Costa Lima 1956; Eggers 1928c: 85; Schedl 1951m: 71–73, 1957a: 192, 1958f: 35; Wood, S. L. 1972f: 47, 1977d: 513, 1979c: 127.
- gracilis* Eggers 1929c: 51 (*Stephanoderes*). Holotype ♀; Cuba; NHMW, Wien, automatic. Synonymy: Wood 1974d: 282. Notes: (1) A replacement name for *obscurus* Ferrari. References: **(cn)** Costa Lima 1956; Schmitz & Crisinel 1957: 11. **(hb)** Costa Lima 1956; Kalshoven 1963; Schmitz & Crisinel 1957: 11. **(ds)** Blackwelder 1947: 778; Costa Lima 1956; Dinther 1960: 111; Kalshoven 1963. **(tx)** Costa Lima 1956; Eggers 1929c: 51, 1940a; Schedl 1951b: 375–376, 1951m: 71–73; Wood, S. L. 1974d: 282.
- lezjavai* Pjatsnitskii 1929a: 15. Syntypes, sex?; Georgia, USSR; not located. Synonymy: Schedl 1961k: 482. References: **(hb)** Davatchi 1955; Lezhava 1930: 563; Stark 1952: 291. **(ds)** Balachowsky 1963a: 1281; Davatchi 1958; Kleine 1934a: 143; Kobakhidze 1957: 175; Pfeffer 1935: 158; Stark 1952: 291. **(tx)** Pjatsnitskii 1929a: 1, 15; Schedl 1934f: 1640, 1961k: 475; Sokanovskii 1954: 17; Stark 1952: 291.
- citri* Ebeling 1935: 21. Holotype ♀; Orange, California [USA]; CAS, San Francisco. Synonymy: Wood 1954a: 1059. References: **(cn)** Ebeling 1950, 1959. **(hb)** Chamberlin 1939: 289; Ebeling 1950. **(ds)** Blackwelder 1939; Chamberlin 1939: 289; Ebeling 1935: 21, 1950, 1959. **(tx)** Chamberlin 1939: 289; Ebeling 1935: 21; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058.
- erythrinae* Eggers 1936d: 628 (*Stephanoderes*). Holotype ♀; Sakalaspur, India; BMNH, London, 1 Eggers cotype in NHMW, Wien. Synonymy: Wood 1972f: 47. References: **(ds)** Beeson 1940: 195, 1961: 288; Brimblecombe 1953: 20; Browne 1961c: 78. **(tx)** Eggers 1936d: 627–628; Schedl 1948g: 25, 1953b: 124, 1955b: 254, 1958b: 99, 1979c: 92; Wood, S. L. 1972: 47.
- bicolor* Schedl 1939f: 32. Lectotype ♀; Java, Pengandaran; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 38, preoccupied by Eggers 1919. Synonymy: Schedl 1957d: 11. References: **(hb)** Kalshoven 1958: 170; Yunus & Hua 1980: 229. **(ds)** Yunus & Hua 1980: 229. **(tx)** Eggers 1919, 1924; Numberg 1960: 609; Schedl 1939c: 326, 328, 1939f: 32, 1942d: 2, 1950g: 896, 1953b: 124, 1953c: 288, 1957d: 11, 1961k: 475–476, 1979c: 38.
- argentincensis* Schedl 1939d: 408. Lectotype, sex?; Corrientes, Buenos Aires; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 24. Synonymy: Wood 1989: 174. References: **(hb)** Viana 1964: 121. **(ds)** Blackwelder 1947; Viana 1964: 121. **(tx)** Schedl 1939d: 408–409, 1948d: 36, 1952a: 446, 1958f: 34–35, 1979c: 24; Wood, S. L. 1989: 174.
- cylindricus* Schedl 1939d: 409. Lectotype, sex?; Isla Martin Garcia [Argentina]; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 74, preoccupied by Hopkins 1915. Synonymy: Wood 1989: 174. References: **(cn)** Ghesquiere 1933a: 32. **(ds)** Blackwelder 1947: 777; Ghesquiere 1933a. **(tx)** Schedl 1939d: 409, 1951m: 71–73, 1952a: 446, 1957d: 12, 1961k: 547, 1979c: 74; Wood, S. L. 1989: 174.
- asaroriensis* Beeson 1940: 195. Holotype ♀; United Provinces: Asarori, Dehra Dun Division; FRI, Dehra Dun. Synonymy: Wood 1989: 174. References: **(cn)** Mathur & Singh 1960b: 9, 1961a: 63. **(ds)** Beeson 1961: 228; Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1960b: 9, 1961a: 63. **(tx)** Beeson 1940: 195; Wood, S. L. 1989: 174.
- dubiosus* Schedl 1940c: 207. Lectotype, sex?; Hamburgfarm, Ebene Limon, Costa Rica; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 84. Synonymy: Wood 1972f: 47. References: **(tx)** Schedl 1940c: 207, 1979c: 84; Wood, S. L. 1972f: 47.
- subcylindricus* Eggers 1940c: 233 (*Stephanoderes*). Holotype ♀; Mosolo Kwenge, Kwango,

- Congo; MRCB, Tervuren. Synonymy: Schedl 1957d: 10.
References: **(tx)** Eggers 1940c: 233–234; Schedl 1950d: 3, 1953d: 68, 1957d: 10–11, 1961k: 476, 1979c: 241.
- mauiensis* Schedl 1941f: 110. Lectotype, sex?; Maui, Iao Valley; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 150. Synonymy: Wood 1989: 174.
References: **(ds)** Ford 1958: 333; Swezey 1941: 119, 1954: 83. **(tx)** Schedl 1941f: 110, 1959a: 480, 1979c: 150; Wood, S. L. 1989: 174.
- glabratus* Schedl 1942a: 175. Lectotype ♀; Kuala Lumpur, Malaya; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 105. Synonymy: Schedl 1961k: 482.
References: **(ds)** Browne 1961c: 78. **(tx)** Schedl 1942a: 175, 1942c: 174, 1953c: 288, 292, 1961k: 476, 482, 1979c: 105.
- calacensis* Eggers 1944b: 94 (*Archicophalus*). Syntypes ♀; Eala, Belgisch-Congo; MRCB, Tervuren. Synonymy: Schedl 1957d: 10.
Notes: (1) Schedl 1961k: 476 (to *Hypothenus*).
References: **(cn)** Mayne & Donis 1951: 333. **(ec)** Mayne & Donis 1951: 333. **(ds)** Mayne & Donis 1962: 311. **(tx)** Eggers 1944b: 94; Schedl 1950c: 204, 1950d: 3, 17, 1957d: 10–11, 1961k: 476, 1979c: 86.
- namulus* Schedl 1948f: 263 (*Stephanoderes*). Lectotype, sex?; Fernando, Noronha; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 163. Synonymy: Wood 1989: 174.
References: **(tx)** Schedl 1948f: 263, 1979c: 163; Wood, S. L. 1989: 174.
- parilis* Schedl 1951m: 100. Lectotype, sex?; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 185. Synonymy: Wood 1989: 174.
References: **(ec)** Schedl 1958d: 190. **(tx)** de Ruette 1970: 102; Schedl 1951m: 100, 1963i: 64, 1979c: 185; Wood, S. L. 1989: 174.
- obscuriceps* Schedl 1952a: 449. Lectotype ♀; Argentina: Buenos Aires, Pilar, Misiones, Dep. Concep., Santa Maria; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 175. Synonymy: Wood 1989: 174.
References: **(hb)** Viana 1964: 121. **(ds)** Viana 1964: 121. **(tx)** Schedl 1951m: 98, 1952a: 449, 1958f: 35–36, 1979c: 175; Wood, S. L. 1989: 174.
- tigrens* Schedl 1952a: 452 (*Stephanoderes*). Lectotype, sex?; Argentina, Buenos Aires, Tigre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 253. Synonymy: Wood 1989: 174.
References: **(hb)** Browne 1961c: 78. **(ds)** Browne 1961c: 78. **(tx)** Schedl 1952a: 452, 1979c: 253; Wood, S. L. 1989: 174.
- glabratus* Schedl 1953c: 292. Lectotype, sex?; Selangor, Kepong, Malaya; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 105. Synonymy: Schedl 1961k: 482.
References: **(hb)** Browne 1961c: 78. **(ds)** Bright 1972d: 52; Browne 1961c: 78. **(tx)** Bright 1972d: 52; Schedl 1953c: 288, 292, 1961k: 476, 482, 1979c: 105.
- parvus* Schedl 1957d: 49. Holotype, sex?; Congo Belge: Yangambi, Mabikwa; MRCB, Tervuren. Synonymy: Wood 1989: 174.
References: **(hb)** Schedl 1961k: 509. **(ds)** Browne 1963a: 556; Schedl 1959a: 479, 1961k: 509. **(tx)** Schedl 1957d: 49, 1959a: 479, 1961k: 509, 1963i: 44, 1979c: 185; Wood, S. L. 1989: 174.
- cylindripennis* Schedl 1957d: 51. Holotype, sex?; Congo Belge: Kivu, Mulungu; MRCB, Tervuren. Synonymy: Wood 1989: 174.
References: **(ds)** Mayne & Donis 1962: 306; Schedl 1961k: 471. **(tx)** Schedl 1957d: 51, 1961k: 471, 1979c: 74; Wood, S. L. 1989: 174.
- viana* Schedl 1958f: 42. Lectotype, sex?; Argentinien: Buenos Aires, Tigre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 267. Synonymy: Wood 1989: 174.
References: **(hb)** Viana 1964: 121. **(ds)** Viana 1964: 121. **(tx)** Schedl 1958f: 42, 1979c: 267; Wood, S. L. 1989: 174.
- mesoleius* Schedl 1959a: 480. Syntypes ♀; Ceylon: Millawitiya Estate; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 174.
References: **(ds)** Schedl 1959a: 480. **(tx)** Schedl 1959a: 480, 1979c: 152.
- minutulus* Schedl 1972j: 225. Holotype, sex?; Ceylon, Central-North Central, Peradeniya, 550 m; MHNG, Geneve. Synonymy: Wood 1989: 174.
References: **(tx)** Schedl 1972j: 225; Wood, S. L. 1989: 174.
- minutus* Schedl 1978c: 299 (*Cryphalus*). Holotype, sex?; Brasilien, Nova Teutonia, 300–500 m, 27 11 Br., 52 23 L.; Schedl Collection in NHMW, Wien, preoccupied by Hopkins 1915. Synonymy: Wood 1989: 177.
References: **(tx)** Schedl 1978c: 299, 1979c: 157; Wood, S. L. 1989: 277.
- euphorbiae* (Schedl) 1961e: 135 (*Stephanoderes*). Holotype, sex?; Madagascar, Faux-Cap; IRSM, Madagascar.
Distribution: Madagascar.
Hosts: *Euphorbia stenoclada*.
References: **(tx)** Schedl 1961e: 135, 1977a: 71, 1979c: 92.
- eximius* Schedl 1951m: 99. Lectotype, sex?; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 94.
Distribution: South America (Brazil).

References: (tx) Schedl 1951m: 99, 1979c: 94.

externedentatus Schedl 1959a: 450. Holotype ♀; Ceylon: Millawitiya Estate; Schedl Collection in NHMW, Wien.

Distribution: Asia (Sri Lanka).

References: (ds) Schedl 1959a: 450, (tx) Schedl 1959a: 450, 1979c: 94.

flavus Hopkins 1915b: 17. Holotype ♀; Island of Java; USNM, Washington.

Distribution: Indonesia (Java).

References: (tx) Hopkins 1915b: 14, 17; Schedl 1942d: 12, 1950g: 896.

fuscicollis (Eichhoff) 1878b: 148 (*Stephanoderes*). Holotype, sex?; Columbia; Hamburg Museum, lost.

Distribution: Africa (Ghana), Antilles Islands (Jamaica), Asia (Punjab in India/ Japan), Indonesia (Java, Sumatra), North America (Belize), South America (Brazil/ Colombia/ Venezuela).

Notes: (1) Identity of this species is based on a specimen in the Schedl Collection (on loan from Eggers) that was compared by Eggers in 1925 to the holotype of *fuscicollis*.

References: (cn) Costa Lima 1956, (hb) Costa Lima 1956, (ds) Blackwelder 1947: 778; Costa Lima 1936, 1956; Hagedorn 1910d: 43; Kleine 1913b: 121, 1914b: 342; LePelly 1968: 113, (tx) Costa Lima 1956; Eichhoff 1878a: 386, 1878b: 148; Hagedorn 1910a: 86; Hopkins 1915b: 23.

sundaensis Eggers 1927b: 396 (*Stephanoderes*). Holotype ♀; Haboko, Sumatra; Eggers Collection, in NHMW, Wien. Synonymy: Wood 1989: 174.

References: (hb) Kalshoven 1958b: 170, (ds) Browne 1961c: 79, 1970: 540, 1984a: 150; Peter, Bagle, & Balasabramanian 1984; Schedl 1959a: 455, 1964c: 304; Wood, S. L. 1982b: 881, (tx) Eggers 1927b; Schedl 1939f: 34, 1942a: 169–170, 1950f: 36–37, 1952c: 63, 1953b: 124, 1959a: 455, 1979c: 247; Wood, S. L. 1982b: 881, 1989: 174.

acqualiclavatus Schedl 1939f: 33. Lectotype ♀; Buitenzorg, Java; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 13. Synonymy: Wood 1982b: 881.

References: (ds) Kalshoven 1958: 170; Wood, S. L. 1977a: 70, (tx) Schedl 1939f: 33–35, 1942d: 2, 1979c: 13; Wood, S. L. 1982b: 881.

costalimai Numberg 1958a: 484 (*Ernophilocus*). Holotype, sex?; Ceara [Brazil]; Numberg Collection, IZW, Warsaw. Synonymy: Wood 1986c: 266.

References: (cn) Azavedo 1924a, (ds) Numberg 1958a: 480, (tx) Numberg 1958a: 480; Wood, S. L. 1986c: 266.

ghanaensis Schedl 1962h: 67. Holotype, sex?; Ghana, Kumasi; BMNH, London. Synonymy: Wood 1989: 174.

References: (tx) Schedl 1962h: 67, 1962k: 1068, 1979c: 103; Wood, S. L. 1989: 174.

glabratus (Schedl) 1957a: 192 (*Stephanoderes*). Syntypes, sex?; Jamaica, Rum Cave vicinity; St. Ann Parish, and Puerto Rico: Rio Piedras; Schedl Collection in NHMW, Wien.

Distribution: Antilles Islands (Jamaica/ Puerto Rico).

Notes: (1) Schedl 1979c: 105 (citation of holotype invalid).

References: (ds) Bright 1972d: 52, 1985c: 175, (tx) Bright 1972d: 52, 1985c: 175; Schedl 1957a: 192, 1979c: 105.

glabripennis (Hopkins) 1915b: 32 (*Stephanoderes*). Holotype ♀; Angat, P.I.; USNM, Washington.

Figures: Maiti & Saha 1986: 165 (adult).

Distribution: Asia ("China"/ Andaman Islands in India/ Malaya/ Thailand), Indonesia (Java, Sumatra), Philippine Islands.

Hosts: *Cassia* sp., *Styrax benzoin*, *Swietenia macrophylla*, *Tamarindus indica*.

References: (hb) Beaver & Browne 1975: 290, 1978: 591; Kalshoven 1958b: 168–169, (ds) Beaver & Browne 1975: 290, 1978: 591; Beeson 1961: 295; Browne 1961c: 72; Maiti & Saha 1986: 164, (tx) Browne 1970: 555; Hopkins 1915b: 24, 32; Maiti & Saha 1986: 164; Nobuchi 1983: 300; Schedl 1942a: 169–170, 1942d: 2, 1953b: 124, 1954a: 139.

gossypii (Hopkins) 1915b: 25 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington.

Distribution: Antilles Islands (Cuba), North America (Hidalgo in Mexico/ S Florida in USA).

Hosts: *Ammonia* sp., *Bidens pilosa*, *Cajanus cajan*, *Capparis bifolia*, *Gossypium* spp., *Ica imbricata*, *Mangifera indica*, *Parmentiera edulis*, *Persca americana*, *Poinsettia heterophylla*, *Sida rhombifolia*, *Tectona grandis*.

References: (cn) Simmonds 1924: 61–62; Souphieff & Scherbinovskaja 1937: 81, (ds) Atkinson & Equihua 1985c: 357; Atkinson et al. 1986: 64; Beeson 1938b: 295; Bright 1985c: 175; Costa Lima 1928: 121; Hargreaves 1948; Kleine 1934a: 145; Leng & Mutchler 1917: 219; Souphieff & Scherbinovskaja 1937: 81; Wood, S. L. 1982b: 903, (tx) Bright 1985c: 175; Hopkins 1915b: 21, 25; Schedl 1950f: 38; Wood, S. L. 1967b: 78, 1982b: 903.

beameri Wood 1954a: 1056. Holotype ♀; Homestead, Florida [USA]; SMUK, Lawrence. Synonymy: Wood 1967b: 78, 1972: 48.

References: (hb) Wood, S. L. 1953: 196, 1954a: 1056, (tx) Wood, S. L. 1953: 196, 1954a: 1056, 1967b: 78, 1972f: 48.

grandis Schedl 1939i: 384. Lectotype, sex?; Ukerewe, Tanganyika Territory; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 109.

Figures: Schedl 1961k: 503.

Distribution: Africa (Tanzania/ Zaire).

Hosts: *Anisopappus* sp., *Coffea* sp., *Dissotis grandiflora*, *Doryalis* sp., *Geniosporum paludosum*, *Indigofera arrecta*, *Kosteletzkya adoensis*, *Lactuca* sp., *Paullinia pinnata*, *Phyllanthus* sp., *Polygonum tomentosum*, *Prunus salasil*, *Theobroma cacao*, *Triunfetta* sp., *Veronia* sp., *Vireta multiflora*.

References: (tx) Schedl 1939g: 170, 1939i: 384, 1961k: 502–503, 1964j: 45, 1979c: 109.

hampei (Ferrari) 1867a: 11–12 (*Cryptalus*). Syn-types ♀; Gallia [in coffee beans]; NHMW, Wien. Figures: Frolich & Rodewald 1969, Le Pelley 1968: 114, Mbondji 1973, Schedl 1961k: 562–563, Yunus & Balsubramaniam 1975: 74. Distribution: Africa (Angola/ Cameroon/ Canary Islands/ Equatorial Guinea/ Central African Republic/ Chad/ Congo/ Fernando Po/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Liberia/ Nigeria/ Sao Tome Island/ Sierra Leone/ South Africa/ Sudan/ Tanzania/ Togo/ Uganda/ Zaire), Antilles Islands (Jamaica/ Puerto Rico), Asia (India/ Iran/ Malaya/ Sri Lanka/ Thailand), Indonesia (Java, Sumatra), Micronesia, New Caledonia, North America (Guatemala/ Honduras), Philippine Islands, Samoan Islands, South America (Argentina/ Brazil/ Colombia/ Suriname), Tahiti Islands. Intercepted worldwide in coffee beans.

Hosts: *Coffea arabica*, *C. liberica*, *C. robusta*.

References: (ay) Brun, Gaudichan, & Marcillan 1989; Edwards et al. 1952, 1953; Ferrao 1951; Herfs 1949: 26; Lhoste & Roche 1960; Mbondji 1973; Van Ryn-Tournal 1975. (bv) Abdul Aziz 1969; Baker, P. 1984; Meixner 1937: 1217; Schedl 1960f: 43. (cn) A. T. 1925: 287–298; Abasa 1975, 1983; Abdul Aziz 1969; Ahrens & Vandenput 1952: 65–72; Almeida & Cavalcante 1964: 85–90; Almeida, Cavalcante, & Holanda 1967; Almeida, Pigatti, & Arruda 1980; Alvarado 1939b; Amante, Cavalcante, & Balut 1971; Amaral 1959, 1962, 1963a, 1963b, 1964; Amaral, Arruda, & Yaya 1965; Amaral Castro 1924: 342–343; Ament 1928: 952; Anderson 1933: 95–110; Andrade 1914: 1–118; Annand 1942: 45–46; Anonymous 1919b: 194, 1921d: 603, 1922a: 456, 1924a: 354, 1926d: 920, 1928d: 14, 1930a: 23, 1937a: 217, 1962q, 1970p, 1980d; Arens 1921a: 314–320, 1921b: 296; Arisz 1923: 40–41, 1924: 1–63, 1925: 1–70; Arruda 1965; Aulmann & La Baume 1912: 48–54; Averna Sacea 1926: 1–87, 1930: 10–24, 195–213; Averoldi 1948; Azevedo 1924a: 85–86, 359–360; Baker, P. 1984; Bakker 1923: 1342; Bally 1921: 1, 1922: 43–61, 1924: 1–24, 1931; Bardner 1978: 459; Begemann 1925: 12–20, 1925: 97, 1925: 1263, 1926: 187, 1926: 194–232, 1926: 364, 1926: 803, 1927a: 329, 1927b: 1–21, 1927c: 194–207, 1928a: 285, 1928b: 573, 1928c: 353–359, 1929: 1–54, 1930, 1930: 11, 1956; Beille 1925: 387–388;

Bemelians 1930: 418–424; Bergamin 1943: 31–72, 1944a, 1944b, 1944d, 1944e: 706–716, 1944g, 1945a, 1945c: 290–293, 1945d, 1945e, 1945f, 1945g, 1945h, 1945i, 1945j, 1946b, 1947a, 1947b, 1957, 1960; Berghe 1922: 937; Bernard 1923: 175–187, 1926: 1–22, 1928: 1–14; Berry 1959; Berthet 1913a: 312–313, 701, 1925: 1–54; Betrem 1931: 799, 1936: 196; Blackman 1950; Bondar 1925: 1–13, 1928: 1–64; Bonnefil 1954: 12; Boocock 1959d; Borgmeier 1927: 279; Bos 1923: 988–999; Bontilly 1900: 1–139; Bowden 1960; Bredo 1934: 8–19, 494–514; Browne 1961c: 73; Brum, Gandichon, & Marcillan 1989; Bunting & Milsum 1930; Busck & Oliveira Filho 1925: 1–19; Bussy 1921, 1925: 249; Buxck, Schmitz, & Crisinel 1954: 297–310; Cachan 1957: 15; Caminha Filho 1926: 122–127; Campos Novaes 1922: 67–70; Cheam & Ung 1973; Chevalier 1947: 1–356; Chiaromonte 1938b: 398–399; Cohic 1951, 1958: 10–14; Coleman 1931: 1–26; Coolhaas 1951; Coquard 1955: 137–142; Corbett 1929: 261–276, 1930: 212–214, 1931: 36–40, 1931: 48–64, 1931: 496–498, 1933: 8–22, 1935: 43–56, 1936: 41–53; Cordeiro Leite 1935: 645–654; Corporaal 1921: 1–5; Correa & Bastos 1960a, 1960b; Costa Lima 1922: 34–35, 1923: 111–114, 1925a: 194–199, 1925b: 368–374, 1925c: 16–19, 141–143, 226–227, 1928b: 117–123, 1928c: 3–49, 1936: 1–460, 1956; Costa Lima & Ravache 1925: 39–42; Cramer 1924: 20–21; Dammerman 1929; Dampf 1928: 14; D'Angremond 1940: 1–76; Davelaar 1921: 1–2; De Albuquerque Leao 1941; Decelle 1954; De Ingunza & Augusto 1964: 96–98; De La Torre et al. 1962a: 1–6, 1962b; Dias 1936: 60–66; Dinther 1960: 111; Do Amaral & Oliveira 1974; Do Amaral, Vaz Arruda, & Do Amaral 1973; Drouillon 1959: 198–207; Dumbleton 1954: 28, 104; Duport 1925: 79; Duval 1949: 85–102; Duval et al. 1948; Edwall 1924: 257–302; Edwards et al. 1952, 1953; Emden 1924: 187–188; Escherich 1926: 593–613, 645–652, 1927: 493–498; Esteves 1959a, 1959b, 1961; Evans, A. C. 1960; Evans, D. E. 1965a, 1965b: 335, 1965c; Ferrao 1951, 1964; Ferreira Lima 1945: 225–231; Figueiredo, Puzzie, & Orlando 1959: 21–24; Filho 1925; Fleutiaux 1901; Flores Ruegg, Lord, & Mesquita 1977; Fonseca 1930: 87, 1934: 267, 1937: 220–229, 1939a: 57–59; Fonseca & Antuori 1932: 1–87; Fonseca & Araujo 1939: 486–504; Fonseca & Morae 1938: 285–291, 325–334, 368–376; Fontes 1961a, 1961b; Forsyth 1960: 280; Franco do Amaral 1956: 39–47; Franco do Amaral & Puzzi 1960; Franz 1949a; Friederichs 1921: 368, 1922b, 1922c, 1922e, 1922f, 1923a, 1923b, 1923c, 1923d, 1923e, 1923f, 1924b, 1924c, 1924d, 1925a, 1925b, 1925c, 1925d, 1925e, 1926d: 36–40, 78, 123, 1927b: 1–14; Friederichs & Bally 1922: 75–80, 1923: 103–147; Frolich & Rodewald 1969: 97; Gallo 1963; Gandrup 1921: 886, 1922b, 1922c,

- 1923a: 2508, 1924a: 219-228, 1925: 124, 1926: 605, 1927: 1-42, 1929: 1842; Gentry, J. W. 1965: 190; Chesquiere 1933a: 26-36, 1933b: 773-786; Giannotti et al. 1965; Gignoux 1949: 183-184; Girard 1935: 96-98; Golding 1946: 1-48; Gomes 1962; Gonzaga et al. 1954: 49-50; Gonzalez 1978; Goot 1928: 2-4, 38-46, 69-72, 1935: 1-79, 1935: 1-94; Cowdley 1910: 7-8, 1912: 1-32, 1914: 36-58, 1916: 48-53, 1917: 32-37; Guenther 1925: 400-414; Haan 1922: 63-75, 1927: 16-30; Hagedorn 1913a: Halauer 1921: 1-2, 1922: 6-11, 1923: 1-41; Hall [van Hall] 1913: 256, 1919a: 291, 1919a: 980, 1919b: 201-205, 1919c: 1099, 1919d: 1-49, 1920: 1-50, 1921: 1-50, 1922: 25, 1923a: 1913-1926, 1924: 1-47, 1924: 1-53, 1925: 1-51, 1926: 75; Hall, Dammerman, & Rutgers 1915; Hall & Rutgers 1922: 81-89; Hambleton 1932: 384, 1944: 383-386, 1948; Hancock 1925: 25-28; Hargreaves 1921: 57-64, 1922: 29-32, 1922: 58, 1923: 15-21, 1923: 172-173, 347-354, 1924: 16, 1926: 51, 1927: 24-27, 1928: 1-11, 1929: 44-45, 1933: 50-54, 1934: 62-72, 1935: 218-224, 1936: 10, 1940: 372-380; Harris 1935: 84-89; Heinrich 1960, 1961: 17-29, 1965; Hempel 1933: 197-212, 1934a: 551-555; Herfs 1949: 26, 1950: 6; Hernandez-Paz & Penagos Dardon 1974; Heuren 1919: 1-66; Heyden, Reitter, & Weise 1883: 182; Hill, D. S. 1983: 70; Hoedt 1924: 248-258, 1929: 147; Huckle 1925: 1-11; Hutson 1936: 379, 1937a: 22-28, 1937b: 1, 1939: 36-41, 1941: 19-20; Ihering 1924: 111-114; Ingram 1965, 1968; Ingunza, S. 1962, 1964; Jack 1927: 23-27; Jepson 1936: 47-53; Jervis 1939: 121-124; Jong 1925: 1-30; Kaatz 1964; Kaden 1930: 1-56; Kalshoven 1932: 253, 1937: 1, 1951: 852, 1953b: 157; Kleine 1913b: 121, 1914b: 288, 323, 1932a: 145, 298; Knaus 1930: 1036; Kuneman 1923: 2321-2323; Kunhi 1930: 1-12; Lambourne 1936; Lankester 1921: 1-26, 1922: 1-26; Lavalre 1958: 124, 1962: 36, 1979; Leeftmans 1920: 645-659, 1922a: 339, 1923: 5, 1923: 191-201, 1923: 1341, 1924a: 9, 1924b: 191-201, 1926: 439-441, 1927: 29-31, 1929: 1-96, 1930: 1-100; Lefevre 1954: 261-263; Leggieri 1952: 85; Lepage & Giannotti 1949: 299-308, 1950: 299-308; Lepage, Giannotti, & Orlando 1948: 17-18; LePelley 1959, 1968: 114, 1973; Lepesme 1944: 239-241, 1947: 644; Leplae 1928a: 271-276, 1928b: 121-123; Leroy 1936: 1-30; Lester-Smith 1937: 39; Lew 1924: 1-26; Liceras & Farge 1974; Ligt 1931: 5-7; Loureiro 1948: 24; Maas 1927: 803; Maas & Boedijn 1927: 1-16, 1927: 233; Maas et al. 1927: 806; Machado 1925: 1-11; Mallamaire 1935: 70-79, 1937: 1-45; Mancion & Alibert 1936: 33-43; Mansingh & Rhodes 1953; Maren 1934: 57; Margarinos Torres 1926: 41-43; Mathur & Singh 1960a: 13; Mathur, Singh, & Lal 1958: 31, 1958: 103; Maurenbacher 1922: 769; Mayne 1914: 596-598, 1924; Mayne & Donis 1951: 333; McCrae 1958: 68; McDonald 1925: 163; McPherson, C. I. 1978; Mendes 1938a: 339-355, 1938b: 405, 1939a: 549-551, 1939b: 574-591, 1949a: 203-214, 1949b: 275-290; Mendes & Franco 1940: 1-33; Mendes et al. 1940: 3; Mertens 1916: 285-301, 1920: 246, 1922: 909; Michelmores 1949: 1-15, 1952: 4-22; Montealegre 1949c: 477; Monti 1954: 817-885; Moreira 1925: 1-26, 1928: 24-25; Morstatt 1911b: 384, 1912: 87, 1914: 137-139, 308, 1937: 26, 1941: 216, 1943: 84; Mumford 1960: 4, 1961: 33, 1962: 35, 1963: 44, 1964: 39, 1965: 33, 1966: 50; Munro 1927: 563; Nanta 1954: 471-473; Neiva 1925: 235-238, 1928: 1-27; Neiva & Acerra-Sacca 1927: 10; Neiva, Costa Lima, & Navarro de Andrade 1924: 235-238, 1925: 1-11; Neiva & Friederichs 1927: 69-74; Neiva, Navarro de Andrade, & Queiroz Telles 1924: 484-503, 1925a: 53, 93, 1925b: 1-11; Newcomer 1950: 5; Oakley 1953: 174-185; Oliveira Filho 1927: 1-95; Ore 1965; Ortiz & Leon 1972a, 1972b; Pamplona 1927: 1-104; Pascual 1939: 753-764; Paxquiere 1932: 223-253; Penados-Robles & Ochoa 1978; Penagos Dardon 1974a; Penagos Dardon & Flores 1974; Petch 1926: 244-271; Pierrand 1962; Pigatti & Pereira 1960: 206-209; Pinto da Fonseca [Fonseca] 1932: 1-87, 1939: 57, 1939: 133-136; Poer, Trench, & Anderson 1930: 1-19; Porteres 1959: 6; Prates 1971; Puzzi 1939: 259-264; Ramachandrarao 1927: 373-374; Rai 1935: 10-25; Rayner 1951: 90-91; Restrepo 1920: 326-327; Rhodes, L. F. & Mansingh 1952; Ribas et al. 1976; Rienoso, C. 1964; Ritchie 1924: 20-22, 1925: 41-44 (141-144?), 1926: 33-36, 1929: 34-40, 1934: 73, 1935: 73-83, 1935: 95-103; Roba 1935: 299-305, 333-347, 371-379; Rocha 1926: 1-37; Rocha Lima 1945: 46; Rodovalho 1925: 1-32; Rodrigo 1941; Rocchoudt 1933: 34-35; Roepke 1909: 365, 1919a: 7; Ross 1919: 85; Rubies 1959: 11; Ruest 1938: 330, 1946: 18; Rutgers 1920: 1-43, 1921: 1-25, 1922: 1-27, 85-89; Sauer 1939: 165-192; Sauer, Duval, & Falanghe 1947: 205-214; Schedl 1961k: 562; Schmeideknecht 1924: 202; Schmidt 1939: 78, 132-150, 187, 239; Schmitz 1958: 261-266; Schmitz & Crisinel 1957: 9-156; Schmutterer 1961: 469; Schonteden 1924: 56-60, 1927: 114; Schwarz 1924: 68-69; Schweizer 1924a: 1387-1390, 1924b: 307-314, 1932: 1-87; Sefer 1961; Seixas 1947: 205, 215, 1948a: 71-89, 1948f: 163-164, 1950: 216-217, 1958a: 13, 1958b: 21; Serratet 1922: 117-118; Siemaszko 1937: 1-83; Silberschmidt 1951: 217; Sivaram 1980; Sladden 1932: 331, 1934: 26, 77; Small 1915: 751, 1921: 41; Smee 1927: 13-20; Smith & Andre Audant 1930: 978; Soares 1945: 138; Solly 1966; Sonza Geribello 1928: 270-271; Sprengel 1926: 78; Stahel 1925: 103, 1926: 79; Staner 1928: 128-140; Stayaert 1935: 1-46; Steinman 1925: 143; Strohmeier 1910f: 187; Sulistyowati 1986; Swain 1953: 166; Taylor, W. E. 1973: 57; Tempany 1930: 14; Thomas, J. M. 1949: 73; Thomas, R. T. S. 1961; Ticheler 1961: 1, 1963: 223-318; Toledo 1942:

- 233–260, 1943: 79–83, 1945: 27–32, 1947: 213–238; Toledo, Duvall, & Sauer 1947: 113–118; Toledo & Sauer 1947: 113; Toledo Piza Junior 1924: 3, 13, 1924b: 354–355, 1928a: 84–85, 1928b: 1–52, 1938: 424; Toledo Piza Junior and Pinto da Fonseca 1935a: 179–199; Toledo Rodovalho [Rodovalho] 1925b: 188–192; Trench 1932: 158–175; Ultee 1919: 1–25, 1921: 1–25, 1923: 22–24, 1924a: 359–367, 1924b: 1–43, 1924: 478, 1925: 1–43, 1926: 1–52, 1927: 18–23, 1928: 10–18, 41–42, 1929: 1–64, 1930: 647, 1931: 1–51, 1931: 1–55; Ultee & Arisz 1924: 229–237; Van der Goot 1928: 39; Van Roehodt 1933: 34; Vayssiere 1923: 107–112, 136–150, 1924: 26–32, 1925: 241–248, 1927: 106–110; Vecht 1953: 272–277; Vonte 1937: 28–34; Vrydagh 1940: 1–4; Vuillet 1914: 19–21; Wallace 1942: 45–46; Waterston 1923: 112–118; Weele [van der Weele] 1910: 1–6, 1910: 308–316; Wildman 1927: 205; Wilkinson 1928: 1–10, 1929: 172–186, 1933: 172–186, 1939: 86–101; Wille 1924: 171–174, 1925: 139–141; Willet 1952: 383; Wood, S. L. 1977a: 71; Wurth 1908a: 63–78, 1908b: 2–20, 1910: 101–105, 1913, 1919: 245–247, 1920: 1–21, 1922: 3–6, 49–52; Wyniger 1962b: 83; Yaya, V. 1965; Yunus & Balasubramaniam 1975: 75; Zacher 1927: 161. (cc) Abasa 1983; Averna-Sacca 1930; Baker, P. 1984; Bally 1922; Begemann 1925, 1926; Bemelmann 1930; Betrem 1961; Bredo 1934, 1939; Browne 1961c: 73; Buyckx et al. 1954; Chiaromonte 1938a; Clausen 1978: 292; Costa & Villacorta 1989; Fonseca & Araujo 1939; Friederichs 1923a, 1923b, 1923c, 1923d, 1923e, 1924a; Fry 1989: 16; Gomes 1962; Gosswald 1938: 434; Hempel 1934a; Koch, J. M. 1973; Leefmans 1923: 5; Madelin 1963: 260; Mayne & Donis 1951: 333; Mbondji 1974; Melville, A. R. 1958; Mendes 1949a: 203; Mumford 1960: 4; Puzzi 1939; Schedl 1961k: 562; Schouteden 1924: 56; Schwarz 1924: 68; Sladden 1934: 26, 77; Tanada 1963: 454; Thompson, W. R. 1943: 112; Thompson, W. R. & Simmonds 1964: 38; Toledo 1942: 234, 1948: 189; Toledo & Sauer 1947: 113; Toledo Piza Junior & Pinto da Fonseca 1935a: 179; Villacorta 1984; Waterston 1923: 112–118; Wichmann 1955a: 96; Yokoyama et al. 1978. (hb) Abasa 1983; Abdul Aziz 1969; Ahrens & Vandenput 1952; Alvarado 1939b; Amante & Balut 1972; Anonymous 1926d: 920; Baltazar & Salazar 1979: 86; Barkmeyer 1927; Bemelmann 1930; Bergamin 1944d; Blackman 1950: 327; Browne 1961c: 73–76; Chevalier 1947; Clausen 1978: 292; Cohic 1958; Corbett 1933; Costa & Villacorta 1989; Costa Lima 1924, 1956; Essig 1942: 605; Evans 1952: 141; Everts 1903: 756; Ferrao 1951; Filho 1928; Friederichs 1924a, 1925a; Frolich & Rodewald 1969: 97; Hagedorn 1913a; Herfs 1949: 26; Hill, D. S. 1983: 70; Hutson 1936: 379; Kalshoven 1937: 1, 1951: 852–855, 1958: 172, 1963; Kleine 1932a: 298; Lengerken 1939: 64; LePelly 1968: 114; Mbondji 1974; Mendes 1949c: 203, 1951a: 213; Montealegre 1949c: 477; Munro 1927: 563; Ortiz & Leon 1972a, 1972b; Penagos Dardon 1974a; Pasquier 1932: 238–247; Prates 1971; Roepke 1919a: 7; Schedl 1961k: 562; Schmitz & Crisinel 1957: 9–156; Sequeira 1961; Sivaram 1980; Sladden 1934: 26, 77; Sulistyowati 1986; Thomas, K. M. 1949: 83; Ticheler 1961, 1963; Wachtl 1876a: 456, 458; Wood, S. L. 1982b: 888; Yunus & Balasubramaniam 1975: 75; Yunus & Hua 1980: 228. (ds) Anonymous 1919b: 194, 1962q, 1968j, 1970p, 1980d, 1989c; Atkinson & Equihua 1985c: 357; Barthe 1896; Beaver 1989a: 2; Beaver & Browne 1975: 291; Beeson 1941: 368, 387–388, 925, 1961: 298; Begemann 1926; Bielz 1887; Blackman 1950; Blackwelder 1947: 778; Brakman 1966b: 206; Bright 1981c: 154, 1985c: 175; Browne 1961c: 73; Choo & Woo 1983; Costa Lima 1922, 1924, 1936, 1956; De Ingunza & Augusto 1964; Dumbleton 1954; Endrodi 1958b; Evans 1952: 141; Everts 1927; Ferreira 1965: 1115; Gemminger & Harold 1872: 2684; Gentry, J. W. 1965: 190; Ghesquiere 1933a; Gonzalez 1978; Hagedorn 1910d: 43, 1912: 41, 1913a: 12; Hambleton 1932, 1947, 1948; Heyden, Reitter, & Weise 1891: 671, 1906: 711; Hill, D. S. 1983: 70, 495, 552; Horion 1951; Johnston, A. 1963b; Kaatz 1961, 1962, 1964; Kalshoven 1932: 253, 1937: 1, 1963; Kleine 1913a: 35, 1932a: 298, 1934: 145; Lee 1971: 31; LePelley 1968: 114; Mathur & Singh 1960a: 13; Mathur, Singh, & Lal 1958: 31; Mayne & Donis 1962: 331; McPherson, G. I. 1978; Mendes 1951a: 213; Mumford 1960: 4, 1961: 33, 1962: 38, 1963: 44, 1964: 39, 1965: 33, 1966: 50; Nonveiller 1984: 40; Nunberg 1952: 19, 1961a: 329, 1965b: 20, 1972b: 197; Ortiz & Leon 1972a, 1972b; Peyerimhoff 1935: 193; Reid, J. C. 1983; Reitter 1894a: 74, 1916: 290; Roba 1935: 340; Schedl 1959a: 485, 1959p: 17, 1960f: 11, 1961k: 562, 662, 1962b: 58, 1962h: 58, 1962k: 1069, 1964j: 40, 1965g: 19, 1967e: 211, 1971d: 425, 1980a: 16; Schmutterer 1961: 469; Sefer 1961; Seidlitz 1891: 608; Sprecher 1934: 150; Stahl 1926: 79; Stein 1868: 114; Stein & Weise 1877: 164; Straby 1970: 6; Strohmeier 1910f: 186–187; Ticheler 1961, 1963; Tymchak 1976: 9, 1977: 10, 1978: 12, 1980: 17; Wachtl 1876a: 456–459; Wichmann 1927a: 79, 1955a: 96; Wood, S. L. 1957e: 1273, 1960a: 35, 1977a: 71, 1982b: 888; Yunus & Hua 1980: 228; Zimmermann 1904: 70. (tx) Bright 1985c: 175; Corbett 1933; Costa Lima 1924, 1956; Eggers 1922a: 85, 1923a: 129–130, 1927a: 177, 1929: 50, 1936d: 626, 1940c: 233; Eichhoff 1872a: 133, 1878b: 153–155; Endrodi 1957b; Essig 1942: 605; Everts 1903: 756; Ferrari 1867a: 11–12, 1867b: 113; Frolich & Rodewald 1969; Hagedorn 1910a: 86, 1910e: 2; Hopkins 1915b: 24; Le Pelley 1968: 114; Lindemann 1877a: 168; Mbondji 1973; Meixner 1937: 1217; Nunberg

1952: 19; Peyerinhoff 1935: 193; Reitter 1894a: 74, 1902: 191, 1906: 707–713, 1913a: 72, 1916: 290; Sampson 1923a: 269–271; Schedl 1934f: 1640, 1941d: 380, 1942d: 2, 1950c: 204, 1950d: 7, 16, 19, 1951b: 376, 1951f: 40, 1953g: 241, 1954d: 871, 1955f: 258, 1957a: 192, 1959a: 485, 1961k: 562–563, 1980a: 16; Wood, S. L. 1957e: 1273, 1960a: 35, 1972f: 48, 1982b: 888; Yunus & Balasubramaniam 1975: 74. (**ms**) Boocock 1959d; Busse 1926; Do Amaral, Vaz Arruda, & Do Amaral 1973; Eggers 1940e: 63; Howard 1930: 360; Kalshoven 1961b: 94; Merino-Rodriguez 1966: 49; Solly 1966; Sprengel 1926: 78.

coffea Hagedorn 1910e: 1 (*Stephanoderes*). Syntypes ♀; Entebbe, Uganda, Zentralafrika, Angola; BMNH, London. Synonymy: Hagedorn 1912: 40.

References: (**cn**) A. A. 1924; A. T. 1925; Alvarado 1939a; Anonymous 1921k: 624, 1924g: 266, 1924m: 265; Azavedo 1924a, 1924b, 1925a, 1925c; Bergamin 1943: 31–72; Conceicao 1943: 895–904; Costa Lima 1924, 1925c; Duclos 1928; Escherich 1927; Gowdey 1912: 1–32, 1913: 29–39, 1914: 36–58, 1917: 32–37, 48–53; Grandi 1951; Hagedorn 1913a; Herfs 1950: 6; Leplae 1928a: 271, 1928b: 121; Mayne 1914: 596; Morstatt 1911a: 69, 1914b: 308, 1914e: 138; Paulian 1943: 359; Rodovalho 1925b: 188; Roepke 1910: 91, 1919a: 7; Schmitz & Crisinel 1957: 9; Schouteden 1924: 56, 1927: 114; Sladden 1934: 27; Small 1915: 71–77, 1915: 751, 1921: 36; Van Roehoudt 1933: 34; Vayssiere 1924: 26; Vuillet 1914: 19–21, 1925: 601; Wille 1924: 172, 1925a: 139. (**cc**) Neger 1911a: 51; Neiva, Costa Lima, and Navarro de Andrade 1925a: 235, 264–273; Schouteden 1924: 56; Sladden 1934: 27; Thompson, W. R. 1943: 112. (**hb**) A. T. 1925; Alvarado 1939a; Costa Lima 1924: 316, 1925c; Grandi 1951; Hagedorn 1913a; Lengerken 1939: 64, 1954: 84; Morstatt 1924: 11, 17, 30; Paulian 1943: 359; Roepke 1919a: 7; Schmitz & Crisinel 1957: 9; Sladden 1934: 27; Vayssiere 1923: 107; Yunus & Hua 1980: 229. (**ds**) Azavedo 1925c; Blackwelder 1947: 778; Costa Lima 1924, 1925c; Hagedorn 1910d: 41, 1913a; Kleine 1913b: 121, 1914b: 288, 1934a: 145; Toledo Piza Junior 1924a: 325; Vayssiere 1923: 107; Yunus & Hua 1980: 229. (**tx**) Costa Lima 1924: 316; Eggers 1922a: 85, 1923a: 129–130; Hagedorn 1910a: 86, 1910e: 1–2, 1912b: 40–41; Lepesme 1944: 238; Sampson 1923a: 270; Schedl 1961k: 563–564; Toledo Piza Junior 1924b: 354.

coffeivorus Weele 1910: 1 (*Xyleborus*). Syntypes, sex?; Java; not located. Synonymy: Strohmeier 1910f: 186.

References: (**cn**) Chevalier 1947; Kalshoven 1932: 253; Roepke 1919a: 7; Strohmeier 1910f: 186–187. (**hb**) Chevalier 1947; Kalsho-

ven 1932: 253; Roepke 1919a: 7. (**ds**) Brakman 1966: 206; Everts 1927; Strohmeier 1910f: 186–187. (**tx**) Hagedorn 1912b: 40–41; Schedl 1962j: 583; Weele 1910: 308.

cooki Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Mount Coffee, Liberia; USNM, Washington. Synonymy: Schedl 1959a: 485.

References: (**tx**) Hopkins 1915b: 22, 27; Schedl 1959a: 111, 1961k: 563.

cofeicola Campos Novaes 1922: 67 (*Xyleborus*). Syntypes, sex?; Brazil; not located. Synonymy: Costa Lima 1924: 316.

References: (**cn**) Bergamin 1945: 290–293. (**hb**) Kalshoven 1963. (**ds**) Blackwelder 1947: 778; Kalshoven 1963; Kleine 1934a: 172; Roba 1935: 340. (**tx**) Campos Novaes 1922: 67–68; Costa Lima 1924: 316–319, 413–416, 1925: 194–199, 365–368; Schedl 1959a: 485, 1961k: 563.

punctatus Eggers 1924: 101 (*Stephanoderes*). Lectotype, sex?; Eala (Congostaat); USNM, Washington, designated by Anderson & Anderson 1971: 27. Synonymy: Wood 1972f: 48. References: (**bv**) Schedl 1960f: 33. (**cn**) Ghesquiere 1933a: 26–36, 1933b: 774–782; Kemmer 1924: 1–33; Mayne & Donis 1951: 333–334. (**cc**) Mayne & Donis 1951: 333–334. (**hb**) Schedl 196k: 618. (**ds**) Ghesquiere 1933a: 26–36; Lepesme 1947: 644–645; Lepesme et al. 1948: 644; Mayne & Donis 1962: 313; Roba 1935: 340; Schedl 1960f: 11, 1961k: 618, 1967e: 212. (**tx**) Anderson, W. H. & Anderson 1971: 27; Eggers 1924: 101; Schedl 1961k: 618; Wood, S. L. 1972f: 48.

glabellus Schedl 1952a: 452 (*Stephanoderes*). Lectotype, sex?; Argentina, Buenos Aires, Isla Martin Cargia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 104. Synonymy: Wood 1989: 174.

References: (**hb**) Viana 1964: 122. (**ds**) Viana 1964: 122. (**tx**) Schedl 1952a: 452, 1979c: 104; Wood, S. L. 1974: 174.

hirsutus (Wood) 1954a: 1020 (*Stephanoderes*). Holotype ♀; Plantation Key, Florida [USA]; SMUK, Lawrence.

Distribution: North America (S Florida in USA). Hosts: *Aceras sapota*, *Ardesia paniculata*, *Eugenia buxifolia*, *Ipomoea cathartica*, *Lysoloma bahamensis*, *Metopium toxiferum*, *Pithecellobium guadeloupense*, *P. nigris-cati*, *Reynosia septentrionalis*. References: (**hb**) Wood, S. L. 1953: 120, 1954a: 1020. (**ds**) Wood, S. L. 1982b: 881. (**tx**) Wood, S. L. 1953: 120, 1954a: 1017, 1020, 1982b: 881.

hirtipennis Schedl 1952a: 450. Lectotype, sex?; Argentine, Buenos Aires, Tigre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 118.

Distribution: South America (Argentina).

References: **(hb)** Viana 1964: 121. **(ds)** Viana 1964: 121. **(tx)** Schedl 1952a: 450, 1979c: 118.

hystrix (Eggers) 1919: 242 (*Adiacretus*). Lectotype ♀; Amami (Deutsch-Ostafrika); USNM, Washington, designated by Anderson & Anderson 1971: 14. Distribution: Africa (Tanzania).

Hosts: *Laurus nobilis*.

References: **(ds)** Kleine 1934a: 149; Schedl 1961k: 507. **(tx)** Anderson, W. H. & Anderson 1971: 14; Eggers 1919: 242–243, 1920: 120–121; Schedl 1939i: 380, 1961k: 507.

incognitus (Schedl) 1967e: 227 (*Stephanoderes*). Holotype ♀; Kindamba, Mexa, Bangou forest [Congo, Brazzaville]; NHMB, Budapest.

Distribution: Africa (Congo).

References: **(tx)** Schedl 1967e: 227–228, 1979c: 123.

indigens Wood 1974a: 20. Holotype ♀; 9 km S Huajuapán, Oaxaca, Mexico; Wood Collection.

Distribution: North America (Jalisco, Nayarit, Oaxaca, Puebla in Mexico).

Hosts: *Psittacanthus* sp., *Serjania* sp., *Toxicodendron* sp.

References: **(ds)** Wood, S. L. 1982b: 883. **(tx)** Wood, S. L. 1974a: 20, 1982b: 883.

inermis Browne 1984d: 94. Holotype, sex?; New Guinea: Kaisenik; BMNH, London.

Distribution: New Guinea.

Hosts: *Nothofagus* sp.

References: **(tx)** Browne 1984d: 94.

ingens (Schedl) 1942d: 18 (*Stephanoderes*). Lectotype, sex?; Java, Semarang, teak forest, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 124.

Distribution: Asia (Vietnam), Indonesia (Java, Sumatra).

References: **(hb)** Kalshoven 1958b: 169. **(ds)** Browne 1970: 540; Schedl 1965a: 339, 1967e: 212. **(tx)** Schedl 1942d: 18, 1953b: 124, 1954a: 139, 1965g: 22, 1979c: 124.

grandis Schedl 1969c: 49 (*Cryphalomorphus*).

Holotype, sex?; Indo-China; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 174.

References: **(tx)** Schedl 1961k: 502, 1969c: 49, 1971e: 10, 1979c: 108; Wood, S. L. 1989: 174.

insignis Browne 1984a: 151. Holotype, sex?; Niah (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: **(tx)** Browne 1984a: 151.

interstitialis (Hopkins) 1915b: 28 (*Stephanoderes*). Holotype ♀; Victoria, Texas [USA]; USNM, Washington.

Distribution: Antilles Islands (Cuba/ Jamaica), North America (Costa Rica/ Guatemala/ Honduras/ Chiapas, Nayarit, Veracruz in Mexico/ Panama/ Alabama, Connecticut, District of Columbia, Florida, Georgia, Illinois, Kansas, Ken-

tucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA), South America (Brazil/ Colombia).

Hosts: *Acer rubrum*, *Aesculus* sp., *Carya* spp., *Cercis canadensis*, *Fagus grandifolia*, *Ficus* sp., *Liquidambar styraciflua*, *Magnolia* sp., *Miconia* sp., *Mimosa* sp., *Morus rubra*, *Ocotea catesbyana*, *Persca borbonica*, *Picea* sp., *Prosopis* sp., *Quercus* spp., *Rhododendron* sp., *Rhus* spp., *Serjania* sp., *Vismia* sp., *Vitis* sp.

References: **(bv)** Roling & Kearby 1975b, 1977. **(cn)** Roling & Kearby 1977. **(ce)** Equihua & Atkinson 1986: 632; Roling & Kearby 1977. **(hb)** Baker, W. L. 1972: 254; Blackman 1922b: 93–94; Chamberlin 1939: 307; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 632; Wood, S. L. 1953: 145, 1954a: 1033. **(ds)** Atkinson & Equihua 1985c: 356, 1988: 92; Blackman 1922b: 93–94; Bright 1985c: 175; Chamberlin 1939: 307; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Drooz 1985: 368; Equihua & Atkinson 1986: 632, 1988: 210; Kirk 1970; Leng 1940: 340; Turnbow & Franklin 1980; Wood, S. L. 1982b: 887. **(tx)** Blackman 1922b: 93–94; Bright 1985c: 175; Chamberlin 1939: 307; Hopkins 1915b: 22, 28; Wood, S. L. 1953: 145, 1954a: 959, 1033, 1972f: 48, 1982b: 887.

interpunctus Hopkins 1915b: 28 (*Stephanoderes*). Holotype ♀; Brownsville, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1033.

References: **(hb)** Beal & Massey 1945: 114; Blackman 1922b: 93; Chamberlin 1939: 307. **(ds)** Beal & Massey 1945: 114; Blackman 1922b: 93; Blackwelder 1947: 778; Chamberlin 1939: 307; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Beal & Massey 1945: 114; Blackman 1922b: 93; Chamberlin 1939: 307; Hopkins 1915b: 22, 28; Schedl 1940a: 342; Wood, S. L. 1954a: 959, 1033.

flavescens Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Tallahassee, Florida [USA]; USNM, Washington. Synonymy: Wood 1954a: 1033.

References: **(hb)** Chamberlin 1939: 309. **(ds)** Blatchley & Leng 1916: 602; Chamberlin 1939: 309; Leng 1920: 340. **(tx)** Blackman 1922b: 92; Blatchley & Leng 1916: 602; Chamberlin 1939: 309; Hopkins 1915b: 23, 29; Wood, S. L. 1954a: 959, 1033.

approximatus Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Columbus, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1033.

References: **(hb)** Blackman 1922b: 93; Chamberlin 1939: 307. **(ds)** Blackman 1922b; Chamberlin 1939: 307; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Blackman 1922b: 93;

- Chamberlin 1939: 307; Hopkins 1915b: 23, 29; Wood, S. L. 1954a: 959, 1033.
- obliquus* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 48. References: (**ds**) Blackwelder 1947: 778; Leng & Mutchler 1917: 219. (**tx**) Hopkins 1915b: 23, 30; Wood, S. L. 1972f: 48.
- opacipennis* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; District of Columbia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1033. References: (**hb**) Chamberlin 1939: 309. (**ds**) Blatchley & Leng 1916: 602; Chamberlin 1939: 309; Leng 1920: 340. (**tx**) Blatchley & Leng 1916: 602; Chamberlin 1939: 309; Hopkins 1915b: 23, 30; Wood, S. L. 1954a: 959, 1033.
- quadridentatus* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1033. References: (**hb**) Blackman 1922b: 91; Chamberlin 1939: 306. (**ds**) Blackman 1922b: 91; Blatchley & Leng 1916: 602; Chamberlin 1925, 1939: 306; Hoffmann 1942: 13; Kleine 1934a: 145; Leng 1920: 340. (**tx**) Blackman 1922b: 91; Blatchley & Leng 1916: 602; Chamberlin 1939: 306; Hopkins 1915b: 23, 30; Wood, S. L. 1954a: 959, 1033.
- intricatus* (Schedl) 1950d: 25 (*Stephanoderes*). Holotype, sex?; Uganda; Schedl Collection in NHMW, Wien. Distribution: Africa (Uganda). Hosts: *Theobroma cacao*. References: (**ds**) Schedl 1961k: 599. (**tx**) Numberg 1960: 291; Schedl 1950d: 25, 1961k: 590, 599, 1979c: 128.
- japonicus* (Niisima) 1910a: 10 (*Cryphalus*). Syn-types, sex?; Japan: Honshu; Nobuchi Collection, Ibaraki. Distribution: Asia (Honshu in Japan). References: (**ds**) Murayama 1954b: 199; Nobuchi 1985c: 16. (**tx**) Murayama 1954b: 199; Niisima 1910a: 10; Schedl 1934f: 1640.
- javanus* (Eggers) 1908c: 215 (*Stephanoderes*). Lectotype ♀; Java; USNM, Washington, designated by Anderson & Anderson 1971: 16. Figures: Maiti & Saha 1986: 167 (adult, antenna). Distribution: Africa (Cameroon/ Congo/ Gabon/ Ghana/ Ivory Coast/ Mozambique/ Sierra Leone/ St. Thome Island/ Zaïre/ Zambia), Antilles Islands (Cuba/ Guadeloupe/ Haiti in Hispaniola/ Martinique), Asia (Andaman Islands, Assam, Kerala, Nicobar Islands in India/ Malaya/ Sri Lanka/ Taiwan/ Thailand), Indonesia (Borneo, Java), North America (Jalisco in Mexico/ Florida in USA), Philippine Islands, South America (Brazil/ Venezuela).
- Hosts: *Banhinia alba*, *B. tomentosa*, *Budiana sessilifolia*, *Cacsalpinia pulcherrima*, *Conocarpus erecta*, *Dipterocarpus zeylanicus*, *Dryobalanops aromatica*, *Dyera costulata*, *Elcagnus pungens fruitlandi*, *Ficus aurea*, *Gossypium hirsutum*, *Hevea brasiliensis*, *Leucania glauca*, *Mangifera indica*, *Ochroma* sp., *Ocotea catesbiana*, *Personia borbonia*, *Rhedia* sp., *Rhizophora mangle*, *Sterculia macrophylla*, *Tamarindus indica*, *Trema floridana*, *Vitis* sp. References: (**cn**) Mathur & Singh 1961a: 70; Sladden 1934: 27. (**cc**) Sladden 1934: 27; Wichmann 1955a: 98. (**hb**) Beaver 1957a: 19; Beaver & Loytyniemi 1955a: 80; Browne 1961c: 72–73; Deyrup & Atkinson 1957a: 66; Kalshoven 1958b: 169; Loytyniemi, Beaver, & Loytyniemi 1954; Sladden 1934: 27; Yunus & Hua 1980: 229. (**ds**) Atkinson & Equihua 1985c: 357; Beaver 1957a: 19, 1990a: 280; Beaver & Loytyniemi 1955a: 80; Beeson 1961: 299; Bright 1955c: 175; Browne 1961c: 73, 1980d: 493; Deyrup & Atkinson 1957a: 66; Hagedorn 1910d: 43; Kleine 1913b: 121, 1914b: 288, 1934a: 145; Maiti & Saha 1986: 167; Mathur & Singh 1961a: 70; Nobuchi 1967: 24; Schedl 1936d: 1, 1959a: 485; Van Swaloenborg 1956: 9; Wichmann 1955a: 98; Wood, S. L. 1957c: 402, 1977a: 71, 1982b: 909; Yunus & Hua 1980: 229. (**tx**) Anderson, W. H. & Anderson 1971: 16; Bright 1955c: 175; Eggers 1908c: 215, 1922b: 167, 1927b: 396, 1936d: 626; Hagedorn 1910a: 86; Hopkins 1915b: 24; Schedl 1939f: 35, 1942d: 3, 1952c: 62, 1954a: 139, 1958k: 143, 1959a: 485, 1959e: 328, 1961k: 603, 1962p: 204, 1979c: 130; Wood, S. L. 1957c: 402, 1960a: 79, 1967b: 79, 1975b: 393, 1976a: 348, 1982b: 909.
- obesus* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1957c: 402. References: (**hb**) Wood, S. L. 1953: 139, 1954a: 1029. (**ds**) Blackwelder 1947: 778; Leng & Mutchler 1917: 219. (**tx**) Hopkins 1915b: 23, 30; Schedl 1962p: 204; Wood, S. L. 1953: 139, 1954a: 1018, 1029, 1957c: 402, 1975b: 393.
- philippinensis* Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; Angat, Philippine Islands; USNM, Washington. Synonymy: Wood 1975f: 393. References: (**tx**) Hopkins 1915b: 23, 31; Wood, S. L. 1957c: 402, 1975b: 393.
- bananensis* Eggers 1922b: 167 (*Stephanoderes*). Syntypes 2, sex?; Banana, Congo; Eggers Collection (USNM, Washington or NHMW, Wien?). Synonymy: Wood 1972f: 51. References: (**bv**) Schedl 1960f: 77. (**cn**) Ghesquiere 1933a: 31–35, 1933b: 782; Schmitz & Crisinel 1957: 12. (**hb**) Schmitz & Crisinel 1957: 12. (**ds**) Ghesquiere 1933a; Kleine 1934a: 145; Schedl 1960f: 31, 1961k: 552. (**tx**) Eggers 1922b: 167–168, 1924: 102, 1927a:

- 197; Schedl 1948c: 664, 1950d: 17, 1954d: 571, 1961k: 547, 552; Wood, S. L. 1967b: 79, 1972f: 51, 1975b: 393.
- kalshoveni* Schedl 1939f: 35 (*Stephanoderes*). Lectotype, sex?; Pasoeroean, Java; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 131. Synonymy: Wood 1972f: 51. References: (hb) Kalshoven 1955: 169. (tx) Schedl 1939f: 35–36; Wood, S. L. 1972f: 51, 1975b: 393, 1979c: 131.
- subagnatus* Eggers 1940b: 101 (*Stephanoderes*). Holotype ♀; Eala, Congo; MRCB, Tervuren. Synonymy: Wood 1972f: 51. References: (cc) Schedl 1961k: 625. (hb) Schedl 1961k: 625. (ds) Mayne & Donis 1962: 313; Schedl 1961k: 625, 1962h: 58, 1962k: 1070. (tx) Eggers 1940b: 101–102; Schedl 1961k: 625, 1979c: 239; Wood, S. L. 1967b: 79, 1972f: 51, 1975b: 393.
- pistor* Schedl 1951m: 102 (*Stephanoderes*). Syn-types 2, sex?; Havana, Cuba; Schedl Collection in NHMW, Wien. Synonymy: Wood 1976a: 348. Notes: (1) Schedl 1979c: 195 (citation of holotype invalid). References: (tx) Schedl 1951m: 102, 1979c: 195; Wood, S. L. 1976a: 348.
- prosper* Schedl 1951m: 103 (*Stephanoderes*). Holotype ♀; Guadeloupe; Schedl Collection in NHMW, Wien. Synonymy: Wood 1976a: 348. References: (ds) Schedl 1960a: 76. (tx) Schedl 1951m: 103, 1979c: 199; Wood, S. L. 1976a: 348.
- kraunhiae* (Murayama) 1950a: 50 (*Cryphalus*). Syntypes, sex?; Fuchu, Tokyo pref., Honshu, Japan; Murayama Collection in USNM, Washington. Distribution: Asia (Honshu in Japan). Hosts: *Kraunhia floribunda*. References: (ds) Murayama 1954b: 164. (tx) Murayama 1950a: 50, 1954b: 164.
- krivolutskayae* Wood 1992b: 79. Holotype, sex?; Eastern USSR; Institute of Zoology, Academy of Science, Vladivostok, automatic. Distribution: Asia (eastern USSR). Hosts: *Botrocarius controversum*. References: (tx) Krivolutskaya 1968: 56–58; Wood, S. L. 1992b: 79.
- insularum* Krivolutskaya 1968: 56. Holotype, sex?; Eastern USSR; Institute of Zoology, Academy of Science, Vladivostok, preoccupied by Perkins 1900. References: (hb) Krivolutskaya 1973: 138. (ds) Krivolutskaya 1965a: 236, 1973: 138. (tx) Krivolutskaya 1968: 56–58; Wood, S. L. 1992b: 79.
- lamuensis* (Eggers) 1935c: 304 (*Stephanoderes*). Holotype, sex?; Kenya Colony (Lamu); BMNH, London. Distribution: Africa (Kenya). References: (tx) Eggers 1935c: 304; Schedl 1950d: 17, 1961k: 601.
- lefevrei* (Schedl) 1952j: 8 (*Stephanoderes*). Lectotype, sex?; Congo Belge: Kivu, Mulungu; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 138. Distribution: Africa (Ruanda/ Zaire). Hosts: *Ficus capensis*, *Millettia ferruginea*, *Myrica salicifolia*. References: (ds) Schedl 1961k: 602. (tx) Schedl 1952j: 8, 1961k: 602, 1979c: 138.
- leprieuri* (Perris) 1866: 194 (*Dryocoetes*). Holotype, sex?; Alger; MNHN, Paris. Distribution: Africa (Algeria), Asia (Israel/ Syria/ Turkey), Europe (Sardinia). Notes: (1) Eggers 1940b: 62 (to *Hypothenemus*). (3) Although verification is needed, *californicus* may be a synonym (SLW). References: (ds) Gemminger & Harold 1872: 2687; Hagedorn 1910d: 67; Kleine 1913b: 136, 1914a: 16; Peyerimhoff 1935: 195; Reitter 1894a: 75; Schedl 1969g: 290. (tx) Balachowsky 1949a: 205; Eggers 1921: 40, 1940b: 62; Eichhoff 1868d: 419; Ferrari 1867a: 31; Hagedorn 1910a: 96; Perris 1866: 194; Peyerimhoff 1935: 195–196; Reitter 1894a: 75; Schedl 1934f: 1640; Wichmann 1913d: 144. (ms) Eichhoff 1986d: 419.
- albipilis* Reitter 1887b: 195 (*Stephanoderes*). Holotype, sex?; Jerusalem; NHMB, Budapest. Synonymy: Eggers 1940b: 62. References: (ds) Hagedorn 1910d: 40; Hariri 1971: 262; Kleine 1913b: 121; Reitter 1894a: 74. (tx) Eggers 1940b: 62; Hagedorn 1910a: 85; Hopkins 1915b: 15; Reitter 1887b: 195, 1894a: 74, 1913a: 72; Schedl 1934f: 1640; Wichmann 1911c: 210.
- kraussei* Wichmann 1911c: 210. Syntypes, sex?; Sardinein; not located. Synonymy: Eggers 1921: 40. References: (tx) Eggers 1912f: 29, 1921: 40, 1940b: 62; Reitter 1913a: 72; Schedl 1934f: 1640; Wichmann 1911c: 210, 1913: 144.
- liberiensis* (Hopkins) 1915b: 31 (*Stephanoderes*). Holotype, sex? Mount Coffee, Liberia, West Africa; USNM, Washington. Distribution: Africa (Ivory Coast/ Liberia/ Tanzania/ Uganda/ Zaire). Hosts: *Coffea* sp., *Delonix regia*, *Prosopis chilensis*, *Theobroma cacao*. References: (cc) Halperin & Holzschuh 1984: 26. (hb) Halperin & Holzschuh 1984: 26. (ds) Beaver & Loytyniemi 1985a: 80; Halperin & Holzschuh 1984: 26; Schedl 1961k: 602. (tx) Hopkins 1915b: 24, 31; Schedl 1961k: 602; Wood, S. L. 1967b: 78.
- theobromae* Eggers 1932c: 32 (*Stephanoderes*). Holotype, sex?; Congostaat (N' Gazi, Aruwimi), und Barumbi; MRCB, Tervuren. Synonymy: Wood 1968b: 78.

- References: **(cn)** Ghesquiere 1933a: 25–36, 1933b: 774–786; Magnin 1954: 465. **(hb)** Schedl 1961k: 626. **(ds)** Ghesquiere 1933a: Kleine 1934a: 145; Schedl 1961k: 625, 1962k: 1070, 1965f: 4, 1967c: 212. **(tx)** Eggers 1932c: 32–33; Schedl 1950d: 17, 1961k: 625–626, 1962k: 1070.
- lineatus (Eggers)** 1927a: 175 (*Stephanoderes*). Holotype ♀; Belg.-Congo: Banana; USNM, Washington.
Distribution: Africa (Tanzania/ Zaire).
References: **(ds)** Schedl 1961k: 507. **(tx)** Anderson, W. H. & Anderson 1971: 18; Eggers 1927a: 175–176; Schedl 1950d: 3, 1961k: 507
- longipennis (Eggers)** 1935c: 305 (*Stephanoderes*). Holotype, sex?; Congostaat (Nialopu); Schedl Collection in NHMW, Wien.
Distribution: Africa (Angola/ Tanzania/ Zaire).
Hosts: *Hagenia abyssinica*, *Persea indica*.
References: **(ds)** Ferreira 1965: 1115; Mayne & Donis 1962: 312; Schedl 1959p: 17, 1961k: 602, 1967c: 212. **(tx)** Eggers 1935c: 305, 1936c: 36; Schedl 1961k: 602–603, 1979c: 141.
- longipilis Schedl** 1952a: 451. Lectotype, sex?; Argentina, Buenos Aires, Tigre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 141.
Distribution: South America (Argentina).
References: **(hb)** Viana 1964: 121. **(ds)** Viana 1964: 121. **(tx)** Schedl 1952a: 451, 1979c: 141.
- macrolobii (Eggers)** 1940c: 234 (*Stephanoderes*). Holotype, sex?; Congostaat; Eggers Collection, in NHMW, Wien.
Distribution: Africa (Angola/ Tanzania/ Zaire), Madagascar.
Hosts: *Anacardium occidentale*, *Gilbertiodendron deccorei*, *Jatropha* sp., *Macrolobium* sp., *Microcos pinnatifida*, *Polyalthia suaveolens*, *Synsepalum subcordatum*.
References: **(cn)** Beccari & Gerini 1968: 129–134. **(hb)** Schedl 1961: 603–604, 1977a: 71. **(ds)** Schedl 1961k: 604, 1977a: 71. **(tx)** Eggers 1940c: 234; Schedl 1950d: 17–18, 1954e: 76, 1957d: 58, 1961k: 604, 1977a: 71, 1979c: 143.
- madagascariensis Schedl** 1953d: 82. Lectotype ♀; Tananarive, Tsimbazaza; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 144.
Distribution: Madagascar.
Notes: (3) The lectotype on which this species is based could be a specimen of *erectus* LeConte or *cryphaloides* Eichhoff.
References: **(tx)** Schedl 1953d: 82, 1961e: 131–132, 1963i: 61–62, 1977b: 40, 56, 1979c: 144.
- maguus (Eggers)** 1924: 102 (*Stephanoderes*). Holotype, sex?; Malela (Congostaat); MRCB, Tervuren. Figures: Schedl 1961k: 545.
Distribution: Africa (Congo).
References: **(tx)** Eggers 1924: 102, 1927a: 198, 1932c: 33; Schedl 1939g: 170, 1939i: 384, 1940d: 589, 1961k: 545.
- major Browne** 1970: 555. Holotype ♀; Sarawak: Quop; BMNH, London.
Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).
References: **(tx)** Browne 1970: 555.
- malayensis (Schedl)** 1977f: 499 (*Lepiceriodes*). Holotype, sex?; Malaysia: Malaya, Selangor, Sungei Buloh, Kuala Lumpur; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
References: **(tx)** Schedl 1950d: 3, 1977f: 499.
- mallyi (Hopkins)** 1915b: 32 (*Stephanoderes*). Holotype ♀; Capetown, South Africa; USNM, Washington.
Distribution: Africa (Congo/ Mozambique/ South Africa).
References: **(tx)** Hopkins 1915b: 24, 32; Schedl 1961k: 604; Wood, S. L. 1972f: 48.
- sonssouensis Eggers** 1943e: 74 (*Stephanoderes*). Holotype ♀; Zambesi (Environ de Sone, M. Soussou); MNHN, Paris, 2 Eggers cotypes in NHMW, Wien. Synonymy: Wood 1972f: 48.
References: **(ds)** Eggers 1943e: 74; Schedl 1965g: 20, 1982: 278. **(tx)** Schedl 1953d: 68, 1961k: 624, 1965c: 59, 1965g: 20, 1965j: 198, 1979c: 232; Wood, S. L. 1972f: 48.
- malus (Schedl)** 1957d: 56 (*Stephanoderes*). Holotype, sex?; Cape Province: Trappes Valley; BMNH, London.
Distribution: Africa (South Africa).
Hosts: *Ficus* sp., *Pyrus malus*, *Rhus lancea*.
References: **(cn)** Anonymous 1970c: 13. **(ds)** Anonymous 1970c: 13; Schedl 1961k: 604. **(tx)** Schedl 1957d: 56, 1961k: 604, 1979c: 147.
- mangororus Schedl** 1961e: 134. Holotype, sex?; Madagascar, Montagne d'Ambre, Joffreville; IRSM, Madagascar.
Distribution: Madagascar.
Hosts: *Mangifera indica*.
References: **(tx)** Schedl 1961e: 134, 1977b: 56, 1979c: 148.
- marshalli (Eggers)** 1936c: 36 (*Stephanoderes*). Holotype, sex?; Natal: Estcourt, 4000 feet; BMNH, London.
Distribution: Africa (South Africa/ Zaire).
Hosts: *Azadirachta indica*, *Scutia myrtina*.
References: **(tx)** Eggers 1936c: 36; Schedl 1961k: 605.
- mateui (Schedl)** 1965j: 198 (*Stephanoderes*). Holotype, sex?; Sahara mer., Ouedi Kobone, Ennedi; MNHN, Paris.
Distribution: Africa (Chad), Asia (Israel).
Hosts: *Acacia raddiana*.
References: **(cc)** Halperin & Holzschuh 1984: 26. **(hb)** Halperin & Holzschuh 1984: 26. **(ds)**

- Halperin & Holzschuh 1984: 26. **(tx)** Schedl 1965j: 198, 1971d: 433, 1979c: 149.
- melanarius (Schedl)** 1953d: 80 (*Cosmoderes*). Lectotype, sex?; Madagascar, Mt. d'Ambre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 151.
Distribution: Madagascar.
Notes: (1) Schedl 1962: 88 (to *Stylotentus*, = *Hypothenemus*).
References: **(hb)** Schedl 1977a: 38. **(ds)** Schedl 1977a: 38. **(tx)** Schedl 1953d: 80–81, 1962r: 88, 1977a: 38, 1979c: 151.
- melasomus (Lea)** 1910: 140 (*Cryphalus*). Syntypes, sex?; N.S. Wales: Clearence River, Wollongong; SAM, Adelaide.
Distribution: Australia (New South Wales/Queensland).
Hosts: *Alphitonia excelsa*, *Casuarina* sp., *Delonix regia*, *Encalyptus acmenioides*.
References: **(ds)** Brimblecombe 1953: 26; Schedl 1936g: 527, 1962i: 74. **(tx)** Lea 1910: 140; Schedl 1936g: 527, 1938f: 46, 1948g: 25, 1958i: 214, 1959c: 168.
- miles (LeConte)** 1878a: 433 (*Cryphalus*). Syntypes 2 ♀; Tampa, Florida [USA]; USNM, Washington, 1 syntype lost before 1952.
Distribution: North America (Florida, Georgia, E Texas in USA).
References: **(hb)** Chamberlin 1939: 288; Deyrup & Atkinson 1987a: 66. **(ds)** Atkinson et al. 1991: 161; Chamberlin 1939: 288; Deyrup & Atkinson 1987a: 66; Hagedorn 1910d: 44; Henshaw 1882: 269, 1885: 148; Kleine 1913b: 118, 1914b: 383, 1934a: 142; Leng 1920: 339; Schwarz 1878d: 468; Swaine 1909: 92; Wood, S. L. 1982b: 904. **(tx)** Chamberlin 1939: 288; Hagedorn 1910a: 87; Hopkins 1915b: 13; LeConte 1878a: 433, 468; Swaine 1909: 92; Wood, S. L. 1953: 211, 1954a: 979, 1053, 1064, 1982b: 904.
- minor (Eggers)** 1927a: 178 (*Adiaeretus*). Holotype, sex?; Ostafrika: Muansa; Methner Collection in MNB, Berlin.
Distribution: Africa (Mozambique/Tanzania).
References: **(tx)** Eggers 1927a: 178; Schedl 1961k: 508.
- modestus (Murayama)** 1940a: 236 (*Cryphalus*). Syntypes, sex?; Seiryu dans Jehol, Manchoukuo, China; Murayama Collection in USNM, Washington.
Distribution: Asia ("Manchoukuo," "Manchuria" in China/Japan/ Korea).
Hosts: *Ulmus pumila*.
References: **(ds)** Murayama 1940a: 232, 236, 1942a: 55, 1953c: 151, 1954b: 164, 1955: 102. **(tx)** Murayama 1940a: 232, 236, 1950a: 51, 1954b: 164, 1955: 102; Nobuchi 1985c: 16; Sokanovskii 1959a: 277.
- morigerus (Schedl)** 1957d: 57 (*Stephanoderes*). Holotype, sex?; Congo Belge: Yangambi, Kivu, Clute de la Rutschuru; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Afzelia bella*, *Scutia myrtina*.
References: **(hb)** Schedl 1961k: 605. **(ds)** Schedl 1961k: 605. **(tx)** Schedl 1957d: 57, 1961k: 605, 1979c: 159.
- morio (Eggers)** 1940b: 101 (*Stephanoderes*). Lectotype ♀; Congostaat (Lusambo); USNM, Washington, designated by Anderson & Anderson 1971: 21.
Distribution: Africa (Angola/ Ghana/ Sierra Leone/ South Africa/ Uganda/ Zaire), Madagascar.
Hosts: *Bauhinia tomentosa*, *Cassia parkii*, *Cynometra alexandri*, *Mangifera indica*, *Spondianthus preussii*, *Theobroma cacao*.
References: **(hb)** Pollet 1977; Schedl 1961k: 605. **(ds)** Ferreira 1965: 1115; Mayne & Donis 1962: 312; Pollet 1977; Schedl 1959p: 17, 1961k: 605, 1962h: 58, 1962k: 1070, 1965f: 4, 1977a: 72. **(tx)** Anderson, W. H. & Anderson 1971: 21; Eggers 1940b: 101; Schedl 1950d: 24, 1961k: 605, 1965g: 19, 1977a: 72, 1979c: 159.
- morosus Schedl** 1965c: 57. Holotype, sex?; Madagascar, Ambodivoangy, Madagascar-Est, det., Sambava, R.N. XII, Marojejy-Onest, 1140 m; IRSM, Madagascar.
Distribution: Madagascar.
References: **(tx)** Schedl 1965c: 57, 1977b: 57, 1979c: 160.
- mozambiquensis Eggers** 1943e: 75. Holotype ♀; Mozambique (Chimoio, Vila Pery); MNHN, Paris, 2 Eggers cotypes in NHMW, Wien.
Distribution: Africa (Mozambique/ Sierra Leone/ South Africa).
Hosts: *Bauhinia tomentosa*.
References: **(ds)** Schedl 1961k: 508. **(tx)** Browne 1970: 553; Eggers 1943e: 75; Schedl 1950d: 18, 1953d: 80, 82, 1961k: 508, 1979c: 160.
- mulongensis (Eggers)** 1940c: 235 (*Stephanoderes*). Syntypes ♀; Congostaat, Mulongo (Niumzu); MRCB, Tervuren.
Distribution: Africa (Ghana/ Zaire).
Hosts: *Cassia siamea*.
References: **(cn)** Magnin 1954: 468. **(hb)** Paulian 1951: 27–28. **(ds)** Mayne & Donis 1962: 306; Paulian 1951: 28; Thompson, G. H. 1963: 47. **(tx)** Eggers 1940c: 235; Schedl 1950d: 3, 1951j: 20, 1952j: 2, 1953d: 68, 1954d: 870, 1954e: 49, 1957d: 11, 1961k: 511, 1979c: 160.
- multidentatulus (Schedl)** 1962r: 95 (*Stephanoderes*). Holotype, sex?; Uganda: Budongo; BMNH, London, automatic.
Distribution: Africa (Uganda).
Notes: (3) This is a probable synonym of *setosus* Eichhoff.
References: **(tx)** Schedl 1962r: 95.
- multidentatus Schedl** 1959q: 707 (*Stephano-*

deres). Holotype, sex?; Uganda: Budongo; BMNH, London, preoccupied by Hopkins 1915. References: (ds) Ferrer 1942; Schedl 1959q: 705, 707. (tx) Schedl 1959q: 707, 1961k: 605, 1962r: 95, 1979c: 161.

multidentatus (Hopkins) 1915b: 28 (*Stephanoderes*). Holotype ♀; Tampico, Tamaulipas, Mexico; USNM, Washington.

Distribution: North America (Guatemala/ Puebla, Tamaulipas in Mexico), South America (Colombia). Hosts: *Coffea* spp., etc.

References: (hb) Wood, S. L. 1982b: 894. (ds) Blackwelder 1947: 778; Browne 1973a: 280; Wood, S. L. 1982b: 894. (tx) Hopkins 1915b: 22, 28; Schedl 1940a: 342, 1961k: 608; Wood, S. L. 1972f: 49, 1982b: 894.

ferrugineus Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Livingston, Guatemala; USNM, Washington, preoccupied by Hopkins 1915. Synonymy: Wood 1972f: 49.

References: (tx) Browne 1963c: 53; Hopkins 1915b: 23, 29; Wood, S. L. 1967b: 78, 1972f: 49.

nitidifrons Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; Tampico, Tamaulipas, Mexico; USNM, Washington. Synonymy: Wood 1972f: 49. References: (ds) Blackwelder 1947: 778; Ferrer 1942. (tx) Hopkins 1915b: 24, 31; Schedl 1940a: 342; Wood, S. L. 1972f: 49.

hopkinsi Browne 1963c: 53. Holotype ♀; Livingston, Guatemala; USNM, Washington, automatic. Synonymy: Wood 1972f: 49.

References: (tx) Browne 1963c: 53; Wood, S. L. 1972f: 49.

multipunctatus (Schedl) 1939f: 36 (*Stephanoderes*). Lectotype, sex?; Buitenzorg, Java; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 161.

Distribution: Indonesia (Java), Samoa.

Notes: (3) A possible synonym of *africanus* Hopkins (SLW, and Beaver 1976b).

References: (hb) Kalshoven 1958b: 170. (ds) Beaver 1976b: 536. (tx) Schedl 1939f: 36, 1951k: 135, 1953b: 124, 1979c: 161.

muticus (Schedl) 1961e: 136 (*Stephanoderes*). Holotype, sex?; Madagascar, Ambila, La Reunion; IRSM, Madagascar.

Distribution: Madagascar, Reunion Island.

Hosts: *Ochrocarpus* sp.

References: (ds) Schedl 1969d: 10, 1977a: 72. (tx) Schedl 1961e: 136, 1969d: 10, 1977a: 72, 1979c: 162.

namosianus Browne 1984e: 78. Holotype, sex?; Viti Levu: Namosi Road [Fiji]; BMNH, London.

Distribution: Fiji Islands.

References: (tx) Browne 1984e: 78.

nanellus Wood 1971: 34. Holotype ♀; Turrialba, Cartago Prov., Costa Rica, 800 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Tree twigs.

References: (hb) Wood, S. L. 1982b: 899. (ds) Wood, S. L. 1982b: 899. (tx) Wood, S. L. 1971: 34, 1982b: 899.

natalensis (Schedl) 1941d: 392 (*Archcophalus*). Holotype ♀; Natal; Schedl Collection in NHMW, Wien.

Distribution: Africa (South Africa).

References: (ds) Schedl 1961k: 442–443. (tx) Schedl 1941d: 392, 1957c: 324, 1961k: 443, 1979c: 163.

nigripiceus (Schedl) 1951j: 20 (*Stephanoderes*). Holotype, sex?; Madagascar, Friedrichs; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

References: (tx) Schedl 1950d: 23, 1951j: 20, 1953d: 68, 1957d: 59, 1977a: 73, 1979c: 168.

novateutonicus (Schedl) 1951m: 105 (*Ptilopodius*). Syntypes, sex?; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien and Plannam Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 172 (citation of holotype invalid).

References: (tx) de Ruelle 1970: 112, 113; Schedl 1951m: 105, 1979c: 172.

oblongus (Nobuchi) 1981c: 14 (*Macrocryphalus*). Holotype, sex?; Takao, Tokyo; Nobuchi Collection, Ibaraki.

Figures: Nobuchi 1981c: 15.

Distribution: Asia (Japan).

Hosts: *Trachelospermum asiaticum*.

Notes: (1) The Nobuchi photographs of the "male" and "female" types appear to represent females of two different species of *Hypothenemus*. The female almost certainly is *fuscicollis* Eichhoff. The "male" photograph is not clear enough to permit species identification; further study is needed to determine specific synonymy (SLW).

References: (tx) Nobuchi 1981c: 14–16; Wood, S. L. 1992: (in press).

obscurus (Fabricius) 1801: 395 (*Hylesinus*). Lectotype ♀; type labeled as Essequibo [Suriname], published as *Americae meridionalis*; UZMC, Copenhagen, designated by Wood 1972f: 49.

Figures: Pedrosa-Macedo & Schonherr 1985: 48. Distribution: Antilles Islands (Cuba/ Dominican Republic in Hispaniola/ Guadeloupe/ Jamaica/ Puerto Rico/ Trinidad/ Virgin Islands), North America (Costa Rica/ "Mexico"/ Panama/ Florida in USA), South America (Brazil/ Colombia/ Suriname/ Venezuela). Intercepted worldwide in Brazil nuts.

Hosts: *Bertholletia excelsa*, *Crotalaria* sp., *Hymenaea courbaril*, *Myristica fragrans*, *Tamarindus indica*, *Theobroma cacao*.

Notes: (3) Eggers 1929e: 50 (*Stephanoderes subopacus*, nomen nudum, synonymy).

References: (cn) Costa Lima 1956; Wood, S. L.

- 1977a: 71. (**cc**) Beaver 1979b: 298. (**hb**) Baker, W. L. 1972: 254; Beaver 1974a, 1979b: 298; Beaver & Browne 1978: 591; Costa Lima 1956; Kalshoven 1963; Wood, S. L. 1982b: 893. (**ds**) Beaver 1974a; Beaver & Browne 1978: 591; Berti-Filho 1979: 41; Blackwelder 1947; Bright 1972d: 57, 1985c: 175; Costa Lima 1956; Drooz 1985: 368; Kalshoven 1963; Kirk 1969, 1970; Kleine 1913b: 118, 1934a: 145; Leng & Mutchler 1917: 219; LePeller 1968: 113; Pedrosa-Macedo & Schonherr 1985: 48; Roba 1935: 340; Schedl 1960a: 78, 1963c: 157, 1963f: 54, 1966f: 84, 1967d: 2, 1970e: 81, 1973d: 155; Schonherr & Pedrosa-Macedo 1981: 52; Wood, S. L. 1977a: 71, 1982b: 893. (**tx**) Benoit 1985: 136; Bright 1972d: 57, 1985c: 175; Costa Lima 1956; Eggers 1929c: 43; Fabricius 1801: 395; Pedrosa-Macedo & Schonherr 1985: 48; Reitter 1902a: 141; Schedl 1939j: 564, 1940c: 206, 1942c: 178, 1951m: 71–73, 1952k: 159–160, 1963i: 60; Wood, S. L. 1954a: 959, 1041, 1972f: 49, 1977d: 513, 1982b: 893.
- asperulus* Eichhoff 1872a: 133 (*Stephanoderes*). Lectotype ♀; northern South America; IRSNB, Brussels, designated by Wood 1982b: 893, preoccupied by LeConte 1868. Synonymy: Wood 1977d: 513, 1982b: 893. References: (**tx**) Eichhoff 1872a: 133; Wood, S. L. 1977: 513, 1982b: 893.
- cassiae* Eichhoff 1878b: 152 (*Stephanoderes*). Lectotype ♀; northern South America; IRSNB, Brussels, designated by Wood 1982b: 893, automatic. Synonymy: Wood 1977d: 513, 1982b: 893. References: (**cn**) Mathur & Singh 1960a: 15, 1960b: 24; Mathur, Singh, & Lal 1958: 24, 103. (**hb**) Beaver & Browne 1975: 290. (**ds**) Beaver & Browne 1975: 290; Beeson 1961: 298; Hagedorn 1910d: 41; Kleine 1913b: 121, 1914b: 247, 269, 1934a: 145; Mathur & Singh 1960a: 15, 1960b: 24; Mathur, Singh, & Lal 1958: 24, 103; Peyerimhoff 1935: 193. (**tx**) Eggers 1936d: 626; Eichhoff 1872a: 133, 1878b: 152; Hagedorn 1910a: 85, 1910e; Hopkins 1915b: 24; Peyerimhoff 1935: 193; Sampson 1923a: 270; Schedl 1934f: 1640; Wood, S. L. 1977d: 513, 1982b: 893.
- kuennemanni* Reitter 1902a: 140. Lectotype ♀; Bremen, Germany [in Brazil nuts]; NHMB, Budapest, designated by Wood 1972f: 49. Synonymy: Wood 1972f: 49. References: (**cn**) Costa Lima 1956; Souphieff & Scherbinovskaja 1937: 81. (**cc**) Wichmann 1955a: 96. (**hb**) Costa Lima 1956; Lengerken 1939: 64, 1954: 84. (**ds**) Blackwelder 1947: 778; Brakman 1966b: 206; Costa Lima 1956; Everts 1922: 642, 1927; Hagedorn 1910d: 44; Horion 1951; Kleine 1913b: 121; Numberg 1958a: 481; Reitter 1916: 290; Schilsky 1909: 188; Souphieff & Scherbinovskaja 1937: 81; Straby 1975: 7; Wichmann 1955a: 96. (**tx**) Costa Lima 1956; Everts 1922: 642; Hagedorn 1910a: 87; Hopkins 1915b: 14; Numberg 1958a: 481; Reitter 1902a: 140, 1913a: 72, 1916: 290; Schedl 1934f: 1640; Wood, S. L. 1972f: 49.
- moschatae* Schaufuss 1905: 8 (*Stephanoderes*). Holotype ♀; Guadeloupe; Hamburg Museum, lost. Synonymy: Wood 1972f: 49. References: (**cn**) Hagedorn 1913a; Souphieff & Scherbinovskaja 1937: 81. (**cc**) Wichmann 1955a: 96. (**hb**) Hagedorn 1913a. (**ds**) Blackwelder 1947: 778; Browne 1970: 540; Hagedorn 1910d: 44, 1913a; Kleine 1913b: 121, 1914b: 379, 1934a: 145; Leng & Mutchler 1917: 219; McFarlane 1961; Souphieff & Scherbinovskaja 1937: 81; Wichmann 1955a: 96. (**tx**) Eggers 1940a; Hagedorn 1910a: 87; Hopkins 1915b: 14; Schaufuss 1905: 8; Schedl 1948f: 261, 1957a: 192; Wood, S. L. 1972f: 49.
- rufescens* Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Allegheny, Pennsylvania [USA]; USNM, Washington. Synonymy: Wood 1972f: 49. References: (**hb**) Chamberlin 1939: 309; Wood, S. L. 1953: 153, 1954a: 1036. (**ds**) Blatchley & Leng 1916: 601; Chamberlin 1939: 309; Leng 1920: 340. (**tx**) Blatchley & Leng 1916: 601; Chamberlin 1939: 309; Hopkins 1915b: 23, 29; Wood, S. L. 1953: 153, 1954a: 1019, 1036, 1966b: 29, 1972f: 49.
- buscki* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; Trinidad, West Indies; USNM, Washington. Synonymy: Wood 1972f: 49. References: (**ds**) Blackwelder 1947: 777; Leonard 1933: 97–137; Martorell 1945: 468–469; Wolcott 1936: 317, 1948: 380. (**tx**) Hopkins 1915b: 23, 30; Wood, S. L. 1972f: 49.
- amazoniens* Eggers 1934a: 78 (*Stephanoderes*). Lectotype ♀; Manaus, Brazil; USNM, Washington, designated by Anderson & Anderson 1971: 4. Synonymy: Wood 1972f: 49. References: (**ds**) Blackwelder 1947: 777; Schedl 1963f: 54. (**tx**) Anderson, W. H. & Anderson 1971: 4; Eggers 1934a: 78; Schedl 1951a: 452; Wood, S. L. 1967b: 78, 1972f: 49, 1979c: 18.
- emarginatus* Schedl 1942d: 11. Lectotype ♀; Buitenzorg, Java; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 90. Synonymy: Wood 1972f: 50. References: (**tx**) Schedl 1942d: 11; Wood, S. L. 1966g: 29, 1972f: 50, 1979c: 90.
- opacus* (Eichhoff) 1872a: 132 (*Stephanoderes*). Lectotype ♀; Nov. Grenada; IRSNB, Brussels, designated by Wood 1982b: 889. Distribution: North America (Costa Rica/Panama), South America (Brazil/Colombia/Venezuela). Hosts: *Calliandra confusa*, *Cassia* sp., *Coffea*

- arabica*, *Inga* sp., *Mimosa* sp., *Nectandra* sp., *Ochroma* sp., *Protium* sp., *Totonita guianensis*.
References: (cn) Costa Lima 1956; Murillo Quinche 1959: 300. (hb) Costa Lima 1956; Murillo Quinche 1959: 300. (ds) Blackwelder 1947: 778; Costa Lima 1936, 1956; Gemminger & Harold 1872: 2684; Hagedorn 1910d: 44; Kleine 1913b: 121, 1914b: 336, 342; LePelley 1968: 113; Wood, S. L. 1982b: 889. (tx) Costa Lima 1956; Eichhoff 1872a: 132–133, 1878b: 151; Hagedorn 1910a: 87; Hopkins 1915b: 24; Wood, S. L. 1982b: 889.
- paradoxus** Schedl 1964j: 45. Holotype, sex?; Congo, Yangambi; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: (tx) Schedl 1964j: 45, 1979c: 184.
- parallelus** (Hopkins) 1915b: 25 (*Stephanoderes*). Holotype ♀; Tampico, Mexico; USNM, Washington. Distribution: Hawaiian Islands, North America (Colima, Tamaulipas in Mexico).
Hosts: "*L. glauca*."
References: (ds) Atkinson & Equihua 1958: 92; Blackwelder 1947: 778; Ferrer 1942; Wood, S. L. 1977a: 71, 1982b: 905. (tx) Hopkins 1915b: 21, 25; Schedl 1940a: 342; Wood, S. L. 1982b: 905.
- parculus** Browne 1984d: 93. Holotype, sex?; New Guinea: Savmill L. A.; BMNH, London.
Distribution: New Guinea.
Hosts: *Evodia* sp.
References: (tx) Browne 1984d: 93.
- perappositus** (Schedl) 1934d: 91 (*Stephanoderes*). Lectotype, sex?; Java, Bandjar; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 188.
Distribution: Indonesia (Java).
Hosts: From grass.
References: (hb) Kalshoven 1958b: 170. (tx) Schedl 1934d: 91, 1979c: 188.
- perhispidus** (Eggers) 1927a: 177 (*Stephanoderes*). Holotype, sex?; Belg.-Congo: Sankuru; MRCB, Tervuren.
Distribution: Africa (Angola/ Ghana/ Nigeria/ Zaire/ Zambia).
Hosts: *Balanites wilsoniana*, *Elaeis guineensis*, *Gilbertiodendron dewevrei*, *Pleiocarpa tubicina*, *Polypodium lycopodioides*, *Randia congolana*, *Theobroma cacao*.
References: (bv) Schedl 1960f: 32. (cn) Chesquiere 1933a: 27–35, 1933b: 776–782; Schmitz & Crisinel 1957: 12. (ce) Schedl 1961k: 608. (hb) Loyttyneimi, Beaver, & Loyttyneimi 1984; Schedl 1961k: 608; Schmitz & Crisinel 1957: 12. (ds) Beaver & Loyttyneimi 1985a: 81; Ferreira 1966: 1115; Chesquiere 1933a; Kleine 1934a: 145; Lepesme et al. 1948: 643–644; Schedl 1959p: 17, 1960f: 32, 1961k: 608, 1964j: 40. (tx) Eggers 1927a: 177; Schedl 1961k: 608.
- perpunctatus** (Eggers) 1940c: 233 (*Stephanoderes*). Holotype, sex?; Congostaat (Rutshuru); Eggers Collection, in NHMW, Wien.
Distribution: Africa (Zaire).
References: (tx) Eggers 1940c: 233; Schedl 1950d: 3, 1961k: 609, 1979c: 190.
- pilosus** Hopkins 1915b: 20. Holotype, sex?; Cayamas, Cuba; USNM, Washington.
Distribution: Antilles Islands (Cuba).
References: (ds) Blackwelder 1947: 777; Bright 1985c: 175; Leng & Mutchler 1917: 219. (tx) Bright 1985c: 175; Hopkins 1915b: 20.
- plumeriae** (Nordlinger) 1856: 74 (*Bostrichus*). Holotype ♀; Venezuela; NHMW, Wien.
Figures: Schedl 1961k: 547.
Distribution: Africa (Liberia/ Zaire), Antilles Islands (Gnadeloupe/ San Domingo/Trinidad), North America (Costa Rica/ Guatemala/ Honduras/ Veracruz in Mexico/ Panama), South America (Brazil/ Colombia/ Venezuela).
Hosts: *Acacia* sp., *Canavalia villosa*, *Cayaponia microdonta*, *Coffea* sp., *Daphnopsis seibertii*, *Dioclea megacarpa*, *Harungana madagascariensis*, *Qualea wittrockii*, *Serjania* sp.
Notes: (3) This species is apparently native to western Africa and was introduced into America.
References: (cn) Box 1953a; Costa Lima 1956; De Albuquerque Leao 1941; Green 1912a; Kleine 1932a: 302; Schmitz & Crisinel 1957: 11. (hb) Costa Lima 1928b: 121–122, 1956; Kleine 1932a: 302; Schmitz & Crisinel 1957: 11. (ds) Alvarenga 1962: 24; Blackwelder 1947: 778; Blandford 1904; Costa Lima 1928: 121–122, 1936, 1956; Ferreira 1947; Ferrer 1942; Gemminger & Harold 1872: 2683; Green 1912a: 2–5, 1916: 608–636; Hagedorn 1910d: 45; Kleine 1913b: 121, 1914b: 336, 339, 361, 379, 1932a: 145, 302; Lacordaire 1866: 383; LePelley 1968: 113; Ran 1935: 10–25; Schedl 1960a: 79; Stahel 1925: 1–3. (tx) Blandford 1904; Costa Lima 1945; Eichhoff 1878b: 164; Ferrari 1867a: 17, 1868: 252; Green 1912a; Hagedorn 1910a: 87; Kirsch 1875: 283; Lacordaire 1866: 383; Nordlinger 1856: 74; Schedl 1940a: 342, 1952k: 159–160; Wood, S. L. 1989: 174.
- pallidus** Hopkins 1915b: 18. Holotype ♀; Mt. Coffee, Liberia; USNM, Washington. Synonymy: Wood 1972f: 45.
References: (bv) Schedl 1960f: 29. (ds) Kleine 1914b: 326; Schedl 1960f: 29, 1961k: 509. (tx) Hopkins 1915b: 14, 18; Schedl 1961k: 509; Wood, S. L. 1972f: 45.
- cylindricus** Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; Trece Aguas, Alta Verapaz, Guatemala; USNM, Washington. Synonymy: Wood 1989: 174.
References: (bv) Schedl 1960f: 110. (hb) Beaver 1974a; Viana 1964: 121. (ds) Beaver 1974a; Blackwelder 1947: 777; Bright 1985c:

174; Schedl 1960f: 110; Viana 1964: 121; Wood, S. L. 1977a: 71. **(tx)** Bright 1985c: 174; Hopkins 1915b: 21, 25; Schedl 1957d; Wood, S. L. 1972f: 45, 1976a: 348, 1989: 174.

transatlanticus Eggers 1941a: 99 (*Stephanoderes*). Holotype ♀; Trois Rivières, Guadeloupe; MNHN, Paris. Synonymy: Wood 1972f: 45. References: **(tx)** Eggers 1941a: 99; Schedl 1951m: 72, 1979c: 254; Wood, S. L. 1972f: 45. *guadeloupensis* Schedl 1951m: 98. Syntypes ♀; Guadeloupe; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 174.

Notes: (1) Schedl 1979c: 114 (citation of holotype invalid).

References: **(tx)** Schedl 1951m: 98, 1979c: 114; Wood, S. L. 1976a: 348, 1989: 174.

ituriensis Schedl 1957d: 55 (*Stephanoderes*). Holotype, sex?; Congo Belge: Kivu, Beni; MRCB, Tervuren. Synonymy: Wood 1989: 174. References: **(cc)** Schedl 1958d: 190. **(hb)** Schedl 1961k: 599. **(ds)** Schedl 1961k: 599. **(tx)** Schedl 1957d: 55, 1961k: 547, 599, 1965g: 20, 1979c: 128; Wood, S. L. 1989: 174.

praecellens (Schedl) 1972e: 288 (*Stephanoderes*). Holotype, sex?; Ghana, Ashanti Region, Ofinso, 259 m, N 6 54, W 1 39; NHMB, Budapest. Distribution: Africa (Ghana).

References: **(tx)** Schedl 1972e: 288, 1979c: 198.

pubescens Hopkins 1915b: 19. Holotype ♀; Key West, Florida [USA]; USNM, Washington. Distribution: Antilles Islands (Puerto Rico), Hawaiian Islands, North America (Durango, Yucatan in Mexico/ Florida, Texas in USA), South America (Argentina/ Brazil).

Hosts: *Andropogon* sp., *Cynodon dactylon*, *Paspalum vaginatum*.

Notes: (3) Breeds in fruiting stems of grass.

References: **(cn)** Anonymous 1965e, 1966m, 1966t. **(hb)** Chamberlin 1939: 295; Wood, S. L. 1953: 206, 1954a: 1061, 1952b: 899. **(ds)** Anonymous 1965e, 1966m, 1966t; Blatchley & Leng 1916: 598; Bright 1985c: 175; Chamberlin 1939: 295; Davis 1965: 36; Leng 1920: 340; Thomas, J. B. 1966; Wood, S. L. 1977a: 71, 1982b: 899. **(tx)** Blatchley & Leng 1916: 598; Bright 1985c: 175; Chamberlin 1939: 295; Hopkins 1915b: 14, 19; Wood, S. L. 1953: 206, 1954a: 1053, 1061, 1972f: 50, 1982b: 899.

subelongatus Hopkins 1915b: 19. Holotype ♀; Victoria, Texas [USA]; USNM, Washington. Synonymy: Wood 1972f: 50.

References: **(hb)** Chamberlin 1939: 295. **(ds)** Chamberlin 1939: 295; Leng 1920: 340. **(tx)** Chamberlin 1939: 295; Hopkins 1915b: 14, 19; Schedl 1961k: 475; Wood, S. L. 1954a: 959, 1058, 1972f: 50.

opacifrons Hopkins 1915b: 25 (*Stephanoderes*). Holotype ♀; Aguadilla, Puerto Rico; USNM, Washington. Synonymy: Wood 1972f: 50.

References: **(ds)** Blackwelder 1947; Gowdey 1926: 27; Leng & Mitchell 1917: 219; Walcott 1948: 351. **(tx)** Hopkins 1915b: 21, 25; Wood, S. L. 1972f: 50.

minutissimus Schedl 1952a: 450. Lectotype, sex?; Argentina, Buenos Aires, Pilar, Córdoba, Dep. Pimilla; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 174.

References: **(hb)** Viana 1964: 121. **(ds)** Viana 1964: 121. **(tx)** Schedl 1952a: 450, 1958f: 35, 1979c: 156; Wood, S. L. 1989: 174.

pubipennis (Eggers) 1935c: 305 (*Stephanoderes*). Holotype, sex?; Congostaat (Mulongo Nimzui); MRCB, Tervuren.

Distribution: Africa (Congo/ Ghana/ Ivory Coast/ Nigeria/ Zaire).

Hosts: *Celtis brieii*, *Citrus* sp., *Cola griseiflora*, *Ficus polita*, *Monodora* cf. *myristica*, *Pterocarpus tinctorius*, *Theobroma cacao*, *Trema guineensis*, *Triplochiton scleroxylon*.

References: **(bv)** Schedl 1960f: 33. **(ec)** Cachan 1957: 15; Schedl 1961k: 616–617. **(hb)** Schedl 1961k: 616; Thompson, G. H. 1963: 67. **(ds)** Mayne & Donis 1962: 312; Schedl 1960f: 33, 1961k: 616; Thompson, G. H. 1963: 67. **(tx)** Eggers 1935c: 305, 1936c: 36; Schedl 1950e: 211, 1954d: 871, 1954e: 50, 1961k: 616, 1965g: 20, 1979c: 203.

rotundicollis (Eichhoff) 1878b: 45 (*Stephanoderes*). Syntypes ♀; America septentrionalis [USA?]; Hamburg Museum, lost.

Distribution: Antilles Islands (Puerto Rico/ Virgin Islands), North America (Michoacan, Nayarit, Oaxaca, Tamaulipas in Mexico/ Arkansas, Florida, Georgia, Illinois, Kansas, Maryland, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Tennessee, Texas, Virginia, West Virginia in USA).

Hosts: *Carya alba*, *C. glabra*, *C. ovata*, *Cercis canadensis*, *Fagus grandifolia*, *Mimosa* sp., *Prosopis* sp., *Quercus* sp., *Rhamnus lanceolata*.

References: **(bv)** Roling & Kearby 1975b. **(cn)** Baker, W. L. 1972: 254. **(hb)** Atkinson et al. 1986: 64; Baker, W. L. 1972: 254; Beal & Massey 1945; Blackman 1922b: 91; Blatchley & Leng 1916: 602; Burgos & Saucedo 1983: 103; Chamberlin 1939: 306; Deyrup & Atkinson 1987a: 66; Hagedorn 1910d: 45; Hopkins 1893a: 133; Kleine 1913b: 121, 1934a: 145; Leng 1920: 340; Swaine 1909: 118; Wood, S. L. 1953: 128, 1954a: 1023. **(ds)** Beal & Massey 1945; Blackman 1922b: 91; Blatchley & Leng 1916: 602; Burgos & Saucedo 1983: 103; Chamberlin 1939: 306; Hagedorn 1910d: 45; Hopkins 1893a: 133; Kleine 1913b: 121, 1934a: 145; Leng 1920: 340; Weber, B. C. & McPherson 1991: 54; Wood, S. L. 1982b: 854. **(tx)** Beal & Massey 1945; Blackman 1922b: 91; Blatchley & Leng 1916: 602; Bright 1985c: 175; Chamberlin 1939: 306; Eichhoff 1878a: 355.

- 1878b: 45, 145, 1896: 608; Hagedorn 1910a: 57; Hopkins 1915b: 24; Swaine 1909: 118; Wood, S. L. 1953: 128, 1954a: 960, 1023, 1982b: 884.
sculpturatus Eichhoff 1878b: 45 (*Stephanoderes*).
 Syntypes, sex?; Tennessee [USA]; Hamburg Museum, lost. Synonymy: Eichhoff 1896: 610. References: (**hb**) Chamberlin 1939: 310. (**ds**) Blatchley & Leng 1916: 603; Chamberlin 1939: 310; Hagedorn 1910d: 42; Leng 1920: 340; Swaine 1909: 117. (**tx**) Blatchley & Leng 1916: 603; Chamberlin 1939: 310; Eichhoff 1878a: 385, 1878b: 45, 146, 1896: 608, 610; Hopkins 1915b: 24; Swaine 1909: 117; Wood, S. L. 1954a: 960, 1023.
- quercus* Hopkins 1915b: 32 (*Stephanoderes*).
 Holotype ♀; Berkeley, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1024.
 References: (**hb**) Baker, W. L. 1972: 254; Beal & Massey 1945: 112; Blackman 1922b: 91; Chamberlin 1939: 306. (**ds**) Beal & Massey 1942, 1945: 112; Blackman 1922b: 91; Blatchley & Leng 1916: 602; Chamberlin 1939: 306; Drooz 1985: 368; Kleine 1934a: 145; Leng 1920: 340. (**tx**) Beal & Massey 1945: 112; Blackman 1922b: 91; Blatchley & Leng 1916: 602; Chamberlin 1939: 306; Hopkins 1915b: 24, 32; Wood, S. L. 1954a: 960, 1023–1024.
- ruficeps* Perkins 1900: 181. Holotype ♀; Oahu; Kaala, Waianae range (2000 ft.); BMNH, London, apparently lost.
 Distribution: Hawaiian Islands
 Notes: (3) The type of *ruficeps* is lost, but this species is probably *crudiae* Panzer.
 References: (**ds**) Hagedorn 1910d: 45; Kleine 1913b: 124, 1914b: 302; Swezey 1935: 21, 1941: 118, 1954: 14. (**tx**) Hagedorn 1910a: 87; Hopkins 1915b: 21; Perkins 1900: 181; Schedl 1934g: 177, 179, 1941f: 110, 1951m: 100.
- rugifer* (Schedl) 1965f: 9 (*Stephanoderes*). Holotype, sex?; Uganda, Waluhumbu; BMNH, London.
 Distribution: Africa (Uganda).
 Hosts: *Prosopis chilensis*.
 References: (**tx**) Schedl 1965f: 9, 1977c: 395.
- ruginosus* Wood 1989: 178. Holotype, ♀; Sudafrica: Potgietersrust, Transvaal; Schedl Collection in NHMW, Wien, automatic.
 Distribution: Africa (South Africa).
 References: (**tx**) Wood, S. L. 1989: 178.
- rugifer* Schedl 1977c: 395 (*Pachynoderes*). Holotype, ♀; Sudafrica: Potgietersrust, Transvaal; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1965.
 References: (**tx**) Schedl 1977c: 395; Wood, S. L. 1989: 178.
- sambesianus* Eggers 1943e: 74. Holotype ♀; Zambeze (Nova Choupanga pres Chemba); MNHN, Paris, 2 Eggers cotypes in NHMW, Wien.
 Distribution: Africa (Mozambique).
 References: (**tx**) Eggers 1943e: 74; Schedl 1961k: 523, 1979c: 215.
- sapporoensis* (Niisima) 1910a: 3 (*Cryphalus*). Holotype ♀; Sapporo; Nobuchi Collection, Ibaraki.
 Distribution: Asia (Hokkaido, Honshu in Japan).
 Hosts: *Tilia cordata* var. *japonica*.
 References: (**ds**) Kleine 1934a: 144; Murayama 1954b: 199; Nobuchi 1955c: 16. (**tx**) Murayama 1954b: 199; Niisima 1910a: 3; Schedl 1934f: 1640, 1979c: 21.
- sassaensis* (Eggers) 1924: 102 (*Stephanoderes*).
 Lectotype, sex?; Congostaat (Region de Sassa); Eggers Collection, in NHMW, Wien.
 Distribution: Africa (Zaire).
 References: (**tx**) Eggers 1924: 102; Schedl 1961k: 619, 1979c: 219.
- schedli* Browne 1963c: 54. Holotype, sex?; Takao, Formosa, Santer; Schedl Collection in NHMW, Wien, automatic.
 Distribution: Asia (Taiwan).
 References: (**tx**) Browne 1963c: 54.
- pubescens* Schedl 1942d: 18 (*Stephanoderes*).
 Holotype, sex?; Takao, Formosa, Santer; Schedl Collection in NHMW, Wien, preoccupied by Hopkins 1915.
 References: (**tx**) Browne 1963c: 54; Schedl 1942d: 18, 1979c: 203.
- scutiae* (Schedl) 1959q: 708 (*Stephanoderes*). Holotype, sex?; Kenya, Mukutano; BMNH, London.
 Distribution: Africa (Kenya).
 Hosts: *Scutia commersonii*.
 Notes: (1) This is probably a *Glostatus* (Schedl's slide mount has the funicle twice broken; only 5 segments show).
 References: (**ds**) Schedl 1959q: 705, 708, 1961k: 619. (**tx**) Schedl 1959q: 708, 1961k: 619, 1979c: 223.
- seoulensis* Choo & Woo 1989: 58. Holotype ♀; Korea; Cheongyangri, Dongdaemun, Seoul; not given.
 Figures: Choo & Woo 1989: 60.
 Distribution: Asia (Korea).
 References: (**tx**) Choo & Woo 1989: 58.
- seriatus* (Eichhoff) 1872a: 133 (*Stephanoderes*).
 Lectotype ♀; New Orleans, Louisiana [USA]; IRSNB, Brussels, designated by Wood 1973c: 177.
 Figures: Numberg 1965b: pl. 1.
 Distribution: Africa (Angola/ Congo/ Egypt/ Equatorial Guinea/ Ghana/ Ivory Coast/ Liberia/ Mozambique/ Nigeria/ Ruanda/ Seychelles Islands/ Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaire), Antilles Islands (Barbados/ Cuba/ Haiti in Hispanola/ Puerto Rico/ Virgin Islands), Asia (Sri Lanka/ Turkey), Australia, Fiji Islands, Hawaiian Islands, Indonesia (Java), Madagascar, Micronesia, Niue Island, North America (Costa

Rica/ El Salvador/ Guatemala/ Honduras/ Chiapas, Jalisco, Nayarit, Oaxaca, Puebla, San Luis Potosi, Tamaulipas, Veracruz, Yucatan in Mexico/ Panama/ Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Texas, Virginia, West Virginia in USA), Philippine Islands, South America (Brazil/ Colombia/ Paraguay).

Hosts: *Acia farnesiana*, *Acras sapota*, *Acrocomia sclerocarpa*, *Aleurites fordii*, *Cassia glauca*, *Bauhinia tomentosa*, *Bursera* sp., *Cajanus cajan*, *Calliandra confusa*, *Callicarpa* sp., *Canavalia* sp., *Carya* spp., *Cecropia* sp., *Citrus aurantifolia*, *Coccolrinax alta*, *Coffea bukovensis*, *Cordia* sp., *Dipholis salicifolia*, *Eleagnus pungens*, *Erythrina* sp., *Eugenia buxifolia*, *Ficus* spp., *Galactia spiciformis*, *Guacca quara*, *Hydracaptans* sp., *Ipomoea cathartica*, *Juglans nigra*, *Liquidambar styraciflua*, *Maclura pomifera*, *Mangifera indica*, *Muonia* sp., *Ochroma* sp., *Ocotea catesbiana*, *Persca americana*, *P. borbonica*, *Philabertella clausa*, *Pinus* spp., *Pithecellobium guadeloupense*, *Pittospermum* sp., *Populus deltoides*, *Prunus persica*, *Quercus* spp., *Rhamnus* sp., *Rhus glabra*, *Saba parviflora*, *Salix* sp., *Serjania* sp., *Sida rhombifolia*, *Tectona grandis*, *Theobroma cacao*, *Thespersia pulpurea*, *Trachylobium narrucosum*, *Trema floridana*, *Trichilia arborea*, *Wisteria* sp., *Urena* sp., *Yucca* sp.

References: (cn) Caminha Filho 1926: 122–127; Costa Lima 1956; Kleine 1932a: 302; Moreira 1928: 24–25; Stokes 1975a. (ec) Equihua & Atkinson 1986: 632. (hb) Atkinson et al. 1986: 66; Beaver 1987a: 19; Burgos & Saucedo 1983: 105; Chamberlin 1939: 103; Costa Lima 1956; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 632; Fonseca 1937: 366–368; Kleine 1932a: 302; Sequira 1961; Wood, S. L. 1982b: 895. (ds) Atkinson & Equihua 1985c: 357, 1986a: 422, 1988: 92; Atkinson et al. 1986: 66; Azavedo 1925b: 85–86; Beaver 1987a: 19, 1987b: 65; Blatchley & Leng 1916: 600; Bright 1985c: 175; Burgos & Saucedo 1983: 105; Chamberlin 1939: 103; Costa Lima 1925: 147, 1928: 117–122, 1956; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 632; Estrada & Atkinson 1988: 210; Gemminger & Harold 1872: 2685; Grunwald 1930: 1–58; Hagedorn 1910d: 45; Hambleton 1947; Kleine 1913b: 121, 1914b: 355, 1932a: 145, 302; Leng 1920: 340; Roba 1935: 340; Swaine 1909: 118; Tucker 1952: 347; Wood, S. L. 1977a: 72, 1983: 105. (tx) Blatchley & Leng 1916: 600; Bright 1985c: 175; Chamberlin 1939: 103; Costa Lima 1956; Eggers 1940a; Eichhoff 1872a: 133, 1878b: 158; Fauvel 1884; Hagedorn 1910a: 85; Hopkins 1915b: 22; LeConte 1876: 356; Reitter 1907: 192; Schedl 1940d: 160, 1948d: 35, 1951m: 104, 1952k: 159–160; Sharp 1879: 102; Swaine 1909: 118; Wood, S. L. 1954a: 959, 1041, 1972f: 50, 1973c: 177, 1977d: 513, 1982b: 895.

pulverulentus Eichhoff 1872a: 133 (*Stephanoderes*). Syntypes ♀; Mexico; Hamburg Museum, lost. Synonymy: Wood 1973c: 177. References: (ds) Blackwelder 1947: 778; Blandford 1904; Browne 1974a: 64; Ferrer 1942; Gemminger & Harold 1872: 2684; Hagedorn 1910d: 45; Kleine 1913b: 121, 1914b: 349. (tx) Blandford 1904; Eichhoff 1872a: 133, 1878b: 161; Hagedorn 1910a: 87; Hopkins 1915b: 22; Reitter 1907: 192; Schaufuss 1891: 12; Schedl 1940a: 342; Wood, S. L. 1972f: 50, 1973c: 177.

vulgaris Schaufuss 1897a: 209 (*Stephanoderes*). Syntypes ♀; La Digue, Sechelles Islands; Hamburg Museum, lost. Synonymy: Wood 1972f: 50.

References: (cn) Anonymous 1961m, 1966j, 1966k; Hammad 1961. (hb) Hammad 1961: 150. (ds) Alhnaud 1900: 439; Anonymous 1961m, 1966j; Ford 1961: 321; Fumasaki 1967: 343; Hagedorn 1910d: 46; Kleine 1913b: 121; Kolbe 1910: 40; Sampson 1914: 384; Schedl 1961k: 629, 1962h: 58, 1962k: 1070, 1969d: 11, 1977a: 78. (tx) Hagedorn 1910a: 85; Hammad 1961: 150; Hammad & El Cherif 1962; Hopkins 1915b: 22; Sampson 1914: 384; Schaufuss 1897a: 209; Schedl 1940d: 586, 1950f: 46, 1951j: 19, 1953d: 68, 1957b: 150, 1961k: 629, 1977a: 78.

aulmanni Hagedorn 1912b: 41 (*Cryphalus*). Syntypes ♀; Daressalam; Hamburg Museum, lost. Synonymy: Wood 1989: 174.

References: (bv) Schedl 1960f: 71. (cn) Ghesquiere 1933a; Hagedorn 1913a; Kleine 1932a: 302; Morstatt 1937: 26; Schmitz & Crisinel 1957: 17. (hb) Hagedorn 1913a; Kleine 1932a: 302; Schedl 1961k: 549, 1977a: 63; Schmitz & Crisinel 1957: 12. (ds) Aulmann & La Baume 1911: 64–65; Beaver & Loytyniemi 1985a: 79; Ghesquiere 1933a; Hagedorn 1913a: 13; Kleine 1913b: 121, 1914b: 322, 1932a: 145, 302; LePelley 1968: 113; Mayne & Donis 1962: 311; Morstatt 1913: 292; Numberg 1952: 19, 1965b: 19; Roba 1935: 338; Schedl 1959a: 484, 1961k: 549, 1965f: 4, 1971d: 432, 1977a: 63; Thompson, G. H. 1963: 66. (tx) Eggers 1922: 173, 1927a: 176, 1936c: 36–37; Hagedorn 1912b: 41–42; Numberg 1952: 19, 1965b: pl. 1; Schedl 1950c: 204, 1952j: 2, 1954e: 49–50, 1959a: 484, 1961k: 549, 1965f: 4, 1977a: 63; Wood, S. L. 1989: 174.

georgiae Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Georgia [USA]; USNM, Washington. Synonymy: Wood 1972f: 50.

References: (cn) Anonymous 1960t, 1961m, 1966j. (ec) Bushing 1965: 460. (hb) Baker, W. L. 1972: 254; Beal & Massey 1945: 114–115; Chamberlin 1939: 303; Kalshoven 1963; Wood, S. L. 1953: 177, 1954a: 1048. (ds) Anonymous 1960t, 1961m, 1966j; Beal &

- Massey 1945; Blackwelder 1947: 778; Blatchley & Leng 1916: 600; Bright 1972d: 58; Chamberlin 1939: 303; Cooper 1935; Ford 1961: 321; Fnnasaki 1967: 333; Kalshoven 1963; Kirk 1970; Knull 1934: 212; Leng 1920: 340; Martorell 1945: 469; Wolcott 1936: 317, 1945: 350; Wood, S. L. 1960a: 35. **(tx)** Beal & Massey 1945: 114–115; Blackman 1922b: 95; Blatchley & Leng 1916: 600; Bright 1972d: 58; Chamberlin 1939: 303; Hopkins 1915b: 22, 26; Wood, S. L. 1953: 177, 1954a: 959, 1048, 1960a: 35, 1972f: 50.
- minutus* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 50. References: **(ds)** Blackwelder 1947: 775; Leng & Mutchler 1917: 219. **(tx)** Hopkins 1915b: 22, 26; Wood, S. L. 1972f: 50.
- texanus* Hopkins 1915b: 26 (*Stephanoderes*). Holotype ♀; Columbus, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Blackman 1922b: 94; Chamberlin 1939: 305. **(ds)** Blackman 1922b: 94; Chamberlin 1939: 305; Kleine 1934a: 145; Leng 1920: 340; Wolcott 1936: 317. **(tx)** Blackman 1922b: 94; Chamberlin 1939: 305; Hopkins 1915b: 22, 26; Wood, S. L. 1954a: 959, 1048.
- fiebrigi* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; San Bernardino, Paraguay; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(ds)** Blackwelder 1947: 778. **(tx)** Hopkins 1915b: 22, 27; Wood, S. L. 1954a: 1048, 1967b: 78, 1972f: 50.
- floridensis* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Haw Creek, Florida [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Chamberlin 1939: 306. **(ds)** Blatchley & Leng 1916: 601; Chamberlin 1939: 306; Leng 1920: 340. **(tx)** Blatchley & Leng 1916: 601; Chamberlin 1939: 306; Hopkins 1915b: 22, 27; Wood, S. L. 1954a: 1048.
- pini* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Kanawha Station, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Chamberlin 1939: 305. **(ds)** Blatchley & Leng 1916: 600; Chamberlin 1939: 305; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Blatchley & Leng 1916: 600; Chamberlin 1939: 305; Hopkins 1915b: 22, 27; Wood, S. L. 1954a: 959, 1048.
- salicis* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Beal & Massey 1945: 113; Bright & Stark 1973: 66; Chamberlin 1939: 305. **(ds)** Beal & Massey 1945: 113; Blatchley & Leng 1916: 600; Bright & Stark 1973: 66; Chamberlin 1939: 305; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Beal & Massey 1945: 113; Blatchley & Leng 1916: 600; Chamberlin 1939: 305; Hopkins 1915b: 22, 27; Wood, S. L. 1954a: 959, 1048.
- tamarindi* Hopkins 1915b: 27 (*Stephanoderes*). Holotype ♀; Manila, Philippine Islands; USNM, Washington. Wood 1972f: 50. References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956. **(ds)** Costa Lima 1956; Kleine 1934a: 145; Schedl 1966b: 30. **(tx)** Costa Lima 1956; Hopkins 1915b: 22, 27; Nobuchi 1953: 300; Wood, S. L. 1972f: 50.
- ficus* Hopkins 1915b: 28 (*Stephanoderes*). Holotype ♀; Alabama [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Beal & Massey 1945: 110–113; Blackman 1922b: 94; Chamberlin 1939: 305; James 1942: 37. **(ds)** Beal & Massey 1945: 110–113; Blackman 1922b: 94; Blatchley & Leng 1916: 602; Chamberlin 1939: 308; Kirk 1969; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Beal & Massey 1945: 110–113; Blackman 1922b: 94; Blatchley & Leng 1916: 601; Chamberlin 1939: 308; Hopkins 1915b: 22, 28; Schedl 1960h: 104; Wood, S. L. 1954a: 959, 1048.
- lucasi* Hopkins 1915b: 28 (*Stephanoderes*). Holotype ♀; southern United States [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Blackman 1922b: 94; Chamberlin 1939: 308. **(ds)** Blackman 1922b: 94; Chamberlin 1939: 308. **(tx)** Blackman 1922b: 94; Chamberlin 1939: 308; Hopkins 1915b: 22, 28; Wood, S. L. 1954a: 959, 1048.
- soltani* Hopkins 1915b: 28 (*Stephanoderes*). Holotype ♀; New Orleans, Louisiana [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Chamberlin 1939: 308. **(ds)** Chamberlin 1939: 308; Leng 1920: 340. **(tx)** Blatchley & leng 1916: 601; Chamberlin 1939: 308; Hopkins 1915b: 22, 28; Wood, S. L. 1954a: 959, 1048.
- virentis* Hopkins 1915b: 28 (*Stephanoderes*). Holotype ♀; Lakeland, Florida [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: **(hb)** Chamberlin 1939: 308. **(ds)** Blatchley & Leng 1916: 602; Chamberlin 1939: 308; Kleine 1934a: 145; Leng 1920: 340. **(tx)** Blatchley & Leng 1916: 602; Chamberlin 1939: 308; Hopkins 1915b: 22, 28; Wood, S. L. 1954a: 959, 1048.
- nitidipennis* Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1972f: 50. References: **(hb)** Wood, S. L. 1953: 150, 1954a: 1035. **(ds)** Blackwelder 1947: 778;

- Leng & Mutchler 1917: 219. (**tx**) Hopkins 1915b: 23, 29; Wood, S. L. 1953: 150, 1954a: 959, 1018, 1035, 1972f: 50.
- nitidulus* Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1977d: 513. References: (**ds**) Blackwelder 1947: 778. (**tx**) Hopkins 1915b: 23, 29; Wood, S. L. 1954a: 959, 1035, 1977d: 513.
- pecanis* Hopkins 1915b: 29 (*Stephanoderes*). Holotype ♀; Orlando, Florida [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: (**hb**) Beal & Massey 1945: 114; Chamberlin 1939: 308. (**ds**) Beal & Massey 1945: 114; Blatchley & Leng 1916: 601; Chamberlin 1939: 308; Leng 1920: 340. (**tx**) Beal & Massey 1945: 114; Blatchley & Leng 1916: 601; Chamberlin 1939: 308; Hopkins 1915b: 23, 29; Wood, S. L. 1954a: 1048.
- subopacicolis* Hopkins 1915b: 30 (*Stephanoderes*). Holotype ♀; Cayamas, Cuba; USNM, Washington. Synonymy: Wood 1977d: 513. References: (**ds**) Blackwelder 1947: 778; Leng & Mutchler 1917: 219. (**tx**) Hopkins 1915b: 23, 30; Wood, S. L. 1954a: 959, 1035, 1977d: 513.
- niger* Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; Brownsville, Texas [USA]; USNM, Washington. Synonymy: Wood 1972f: 50. References: (**hb**) Chamberlin 1939: 309; Wood, S. L. 1953: 157, 1954a: 1038. (**ds**) Chamberlin 1939: 309; Ferrer 1942; Leng 1920: 340. (**tx**) Chamberlin 1939: 309; Hopkins 1915b: 24, 31; Schedl 1940a: 342; Wood, S. L. 1953: 157, 1954a: 1019, 1038, 1972f: 50.
- tonsus* Eggers 1919: 242. Lectotype ♀; Amani (Ostafrika); USNM, Washington, designated by Anderson & Anderson 1971: 34. Synonymy: Eggers 1922b: 173. References: (**tx**) Anderson, W. H. & Anderson 1971: 34; Eggers 1919: 242. 1922b: 173, 1927a: 176; Schedl 1961k: 549.
- robustus* Blackman 1922b: 88. Syntypes ♀; Newton, Mississippi [USA]; USNM, Washington. Synonymy: Wood 1954a: 1048. References: (**hb**) Chamberlin 1939: 293. (**ds**) Chamberlin 1939: 293; Kleine 1934a: 144; Leng & Mutchler 1927: 52. (**tx**) Blackman 1922b: 88; Chamberlin 1939: 293; Wood, S. L. 1954a: 959, 1048.
- cassavaensis* Schedl 1938d: 453. Lectotype ♀; Uganda, Kampala; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 54. Synonymy: Wood 1989: 175. References: (**cc**) Beaver 1979b: 298. (**hb**) Beaver 1976b: 535, 1979b: 298; Beaver & Browne 1978: 590. (**ds**) Beaver 1976b: 535; Beaver & Browne 1978: 590; Brader 1964: 5; Browne 1980c: 485, 1980d: 493; Ferreira 1965: 1113; Paulian 1951: 31; Schedl 1938d: 453, 1959a: 478, 1959p: 16, 1961k: 470. (**tx**) Schedl 1938d: 453–454, 1941d: 397, 1951j: 21, 1951k: 134, 1953d: 81, 1955b: 284, 1957d: 12, 1959a: 478, 1959p: 16, 1961k: 470, 1977b: 49; Wood, S. L. 1989: 175.
- hawaiiensis* Schedl 1941f: 112 (*Stephanoderes*). Lectotype ♀; Oahu, Honolulu; BPBM, Honolulu², designated by Schedl 1979c: 116. Synonymy: Wood 1989: 175. References: (**ds**) Swezey 1941: 120. (**tx**) Schedl 1941f: 112, 1979c: 116; Wood, S. L. 1960a: 39, 1961a: 38, 1989: 175.
- striatulus* Schedl 1942d: 12. Lectotype, sex²; Java, Buitenzorg, Nr. 388 Kalshoven, Walikoekoen; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 237. Synonymy: Wood 1989: 175. References: (**ds**) Blackwelder 1947: 778; Browne 1961c: 79. (**tx**) Schedl 1942d: 12, 1951k: 134, 1953c: 288, 1955b: 284, 1957d: 12, 1959a: 478, 1979c: 237; Wood, S. L. 1989: 175.
- darwinensis* Schedl 1942c: 178 (*Stephanoderes*). Lectotype, sex²; Australia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 75. Synonymy: Wood 1972f: 50. References: (**ds**) Brimblecombe 1953: 25; Schedl 1959d: 67. (**tx**) Schedl 1942c: 178, 1948g: 25, 1950f: 37, 1959d: 67, 1979c: 75; Wood, S. L. 1972f: 50.
- marovoayi* Schedl 1953d: 81. Lectotype, sex²; Madagascar, plaine de Marovoay, Madagascar, plantation du Sambirano; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 149. Synonymy: Wood 1989: 175. References: (**tx**) Schedl 1953d: 81, 1977b: 56, 1979c: 149; Wood, S. L. 1989: 175.
- andersoni* Wood 1954a: 1045 (*Stephanoderes*). Holotype ♀; Coconut Grove, Florida [USA]; USNM, Washington. Synonymy: Wood 1972f: 50. References: (**hb**) Wood, S. L. 1953: 171, 1954a: 1045. (**tx**) Wood, S. L. 1953: 171, 1954a: 1045, 1972f: 50.
- liquidambarae* Wood 1954a: 1046 (*Stephanoderes*). Holotype ♀; Jacksonboro, South Carolina [USA]; SMUK, Lawrence. Synonymy: Wood 1972f: 51. References: (**hb**) Wood, S. L. 1953: 174, 1954a: 1046. (**tx**) Wood, S. L. 1953: 174, 1954a: 1046, 1972f: 50.
- asperatus* Schedl 1967e: 226 (*Stephanoderes*). Holotype ♀; Kindamba, Meya, Bangou forest, Sibiti, IRHO rain forest; NHMB, Budapest. Synonymy: Wood 1989: 175. References: (**tx**) Schedl 1967e: 226, 1979c: 28; Wood, S. L. 1989: 175.
- setosus* (Eichhoff) 1868e: 391 (*Hypoborus*). Syntypes ♀; Guadeloupe; Hamburg Museum, lost except for 1 in USNM, Washington. Distribution: Africa (Cameroon/ Ivory Coast/

Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaire), Antilles Islands (Cuba/ Guadeloupe/ Haiti in Hispaniola/ Jamaica/ Puerto Rico), Asia (Taiwan), Madagascar, North America (Costa Rica/ Honduras/ Chiapas in Mexico/ Panama/ Florida in USA), South America (Brazil/ Colombia/ Venezuela).

Hosts: *Acacia pennatula*, *Bauhinia variegata*, *Cecropia* sp., *Mangifera indica*, *Theobroma cacao*.

References: (**cn**) Aulmann 1912: 54; Aulmann & La Baume 1911: 6, 67; Corporaal 1921: 5; Ghesquiere 1933a: 31, 1933b: 782. (**cc**) Wichmann 1955a: 98. (**hb**) Beaver & Browne 1975: 291, 1978: 591; Eichhoff 1881a: 46, 191; Kalshoven 1963; Wachtl 1876a: 458; Wood, S. L. 1982b: 909. (**ds**) Atkinson & Equihua 1988: 92; Beaver 1990a: 280; Beaver & Browne 1975: 291, 1978: 591; Beeson 1941: 389; Blackwelder 1947: 779; Brader 1964: 5; Bright 1972d: 55, 1985c: 175; Browne 1981b: 600; Estrada & Atkinson 1988: 210; Fleutiaux & Salle 1890: 457; Gemminger & Harold 1872: 2684, 2686; Hagedorn 1910d: 45; Heyden, Reitter, & Weise 1883: 182, 1891: 671, 1906: 711; Horion 1951; Kalshoven 1955: 169–170, 1963: 233; Kleine 1913b: 121; Kraatz 1869: 59; Leng & Mutchler 1917: 219; Lepesme 1947: 645; Nobuchi 1967: 13, 24; Pittioni 1943: 176; Reitter 1894a: 73, 1916: 290; Schedl 1966b: 29, 1977a: 73; Schilsky 1909: 188; Seidlitz 1872: 394, 1891a: 563, 1891b: 608; Stein 1868: 114; Wachtl 1876a: 458; Wichmann 1954: 514, 1955a: 98; Wood, S. L. 1977a: 72, 1982b: 909. (**tx**) Bright 1972d: 55, 1985c: 175; Chamberlin 1939: 303; Eggers 1922b: 174, 1923a: 130, 1924: 102, 1927a: 195, 1936d: 626, 1940a, 1941a: 108; Eichhoff 1868d: 419, 1868e: 391, 1872a: 133, 1877a: 117, 1878b: 149–150, 1881a: 46, 191, 1883a: 110, 134; Ferrari 1867a: 81, 1867b: 81, 114; Hagedorn 1910a: 88, 1910e: 2; Hopkins 1915b: 23; Lindemann 1877a: 168; Nobuchi 1983: 300; Reitter 1894a: 73, 1913a: 71, 1916: 290; Schaufuss 1905: 8; Schedl 1934f: 1640, 1936j: 1, 1939e: 328, 1939f: 35, 1942a: 197, 1942d: 3, 1945c: 664, 1950d: 17, 1951m: 102, 1952c: 62, 1954a: 139, 1954d: 871, 1955k: 143, 1959a: 485, 1962h: 58, 1962p: 204, 1967b: 29, 1971a: 274, 280, 1971c: 363, 1971f: 148, 1977a: 73; Seidlitz 1872: 394, 1891a: 563, 1891b: 608; Wood, S. L. 1954a: 1029, 1957c: 402–403, 1962: 79, 1967b: 79, 1972f: 51, 1975b: 393, 1982b: 909. (**ms**) Eichhoff 1868d: 419.

obscurus Eichhoff 1872a: 133 (*Stephanoderes*). Holotype ♀; Antilles; IRSNB, Brussels, preoccupied by Fabricius 1801. Synonymy: Wood 1975b: 393.

References: (**ds**) Gemminger & Harold 1872: 2684; Hagedorn 1910d: 133. (**tx**) Eichhoff 1872a: 133, 1878b: 155; Ferrari 1867a: 17; Wood, S. L. 1975b: 393.

depressus Eichhoff 1878b: 155 (*Stephanoderes*).

Holotype ♀; Antilles; IRSNB, Brussels, automatic. Synonymy: Wood 1975b: 393.

References: (**ds**) Blackwelder 1947: 777; Bright 1985c: 174; Hagedorn 1910d: 41, 47; Kleine 1913b: 121; Leng & Mutchler 1917: 219. (**tx**) Bright 1985c: 174; Eggers 1929e; Eichhoff 1878b: 155; Hagedorn 1910a: 86; Hopkins 1915b: 21; Wood, S. L. 1975b: 393.

congonus Hagedorn 1912a: 337 (*Stephanoderes*). Lectotype ♀; Eala, Congo; MRCB, Tervuren, designated by Wood 1975b: 393. Synonymy: Wood 1975b: 393.

References: (**bv**) Schedl 1960f: 11. (**cn**) Aulmann 1912: 54–55; Ghesquiere 1933a: 34–36, 1933b: 782–785; Kleine 1932a: 302; Mayne 1915: 577–596; Mayne & Donis 1951: 333. (**cc**) Mayne & Donis 1951: 333. (**hb**) Kleine 1932a: 302; Schedl 1961k: 560. (**ds**) Aulmann 1913: 1–126; Gardner 1933a, 1957a: 32; Green 1916: 608–636; Kleine 1913b: 121, 1914b: 314, 1932a: 145, 302, 1934: 145; Schedl 1960f: 11, 1961k: 560. (**tx**) Eggers 1922b: 174, 1924: 103, 1927a: 197; Hagedorn 1912a: 337, 1913b: 18; Numberg 1960: 291; Schedl 1950d: 17, 1961k: 560, 1965g: 19; Wood, S. L. 1967b: 78, 1975b: 393.

simoni (Reitter) 1887b: 194 (*Stephanoderes*). Holotype, sex?; Syrien: Haifa; NHMB, Budapest. Distribution: Asia (Israel/ Syria).

References: (**ds**) Hagedorn 1910d: 45; Hariri 1971: 262; Kleine 1913b: 124; Reitter 1894a: 74; Schedl 1969g: 290. (**tx**) Hagedorn 1910a: 88; Hopkins 1915b: 15; Reitter 1887b: 194, 1894a: 74, 1913a: 73; Schedl 1934f: 1640.

sobrinus (Schedl) 1965c: 58 (*Stephanoderes*). Holotype, sex?; Madagascar, Montange d'Ambre; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

References: (**tx**) Schedl 1965c: 58, 1977a: 76, 1979c: 231.

socialis (Schedl) 1957d: 57 (*Stephanoderes*). Holotype, sex?; Congo Belge: Yangambi, Kibalituri, Kilometre 518, Ouest d'Irumu; MRCB, Tervuren. Distribution: Africa (Angola/ Congo/ Ghana/ Zaire).

Hosts: *Anthonotha macrocarpa*, *Balanites wilsoniana*, *Cola griseiflora*, *Dialium corbisieri*, *Duboscia viridiflora*, *Gilbertiodendron devicrei*, *Polyalthia suae-colaensis*, *Trema guineense*.

References: (**bv**) Schedl 1960f: 33. (**cc**) Schedl 1961k: 620. (**hb**) Browne 1963a: 234; Schedl 1961k: 620. (**ds**) Browne 1963a: 234; Ferreira 1965: 1115; Schedl 1959p: 17, 1960f: 33, 1961k: 620, 1962h: 58, 1962k: 1070, 1964j: 41. (**tx**) Schedl 1957d: 57, 1959p: 17, 1961k: 620, 1962k: 1070, 1965g: 20.

solitarius (Schedl) 1950d: 24 (*Stephanoderes*). Holotype, sex?; D.O. Africa, Uganda; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Nigeria/ South Africa/ Uganda/ Zaire).

Hosts: *Albizia gummifera*, *Bambus* sp., *Coffea* sp., *Pancovia harmsiana*, *Pterocarpus tinctorius*, *Synsepalum subcordatum*, *Theobroma cacao*.

References: (cn) Gregory 1954; Magnin 1954: 468; Schmitz & Crisinel 1957: 12. (hb) Pollet 1977; Schedl 1961k: 622; Schmitz & Crisinel 1957: 12. (ds) Mayne & Donis 1962: 313; Pollet 1977; Schedl 1961k: 622, 1964e: 68, 1965f: 618. (tx) Schedl 1950d: 24, 1954d: 871, 1961k: 622–623, 1965g: 20, 1979c: 232.

solicis Wood 1974a: 21. Holotype ♀; 3 km W Armeria, Colima, Mexico; Wood Collection.

Distribution: North America (Colima, Nayarit, Sinaloa in Mexico).

Hosts: *Inga* sp., *Phoradendron* sp., *Serjania* sp.

References: (ec) Equihua & Atkinson 1986: 632. (hb) Baker, W. L. 1972: 254; Equihua & Atkinson 1986: 632. (ds) Equihua & Atkinson 1986: 632; Wood, S. L. 1982b: 891. (tx) Wood, S. L. 1974a: 21, 1982b: 891.

sparsedentatus (Schedl) 1942c: 178 (*Stephanoderes*). Holotype, sex?; Witu, Tanagebiet, Denhardt; Schedl Collection in NHMW, Wien.

Distribution: Australia.

References: (tx) Schedl 1942c: 178, 1979c: 232.

sparsus Hopkins 1915b: 20. Holotype ♀; Cohnbus, Texas [USA]; USNM, Washington.

Distribution: North America (Tamaulipas in Mexico/ S Texas in USA).

Hosts: *Celtis pallida*, *Condalia* sp., *Rhamnus* sp. References: (hb) Blackman 1922b: 87–88; Chamberlin 1939: 292; Wood, S. L. 1953: 159, 1954a: 1040, 1982b: 896. (ds) Blackman 1922b: 87–88; Chamberlin 1939: 292; Kleine 1934a: 144; Leng 1920: 340; Wood, S. L. 1982b: 896. (tx) Blackman 1922b: 87–88; Chamberlin 1939: 292; Hopkins 1915b: 14, 20; Wood, S. L. 1953: 159, 1954a: 959, 1040, 1982b: 896.

similis Hopkins 1915b: 20. Holotype ♀; Victoria, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1040.

References: (hb) Chamberlin 1939: 295. (ds) Chamberlin 1939: 295; Leng 1920: 340. (tx) Anderson, W. H. & Anderson 1971: 30; Chamberlin 1939: 295; Hopkins 1915b: 14, 20; Wood, S. L. 1954a: 959, 1040.

tridentatus Hopkins 1915b: 31 (*Stephanoderes*). Holotype ♀; San Diego, Texas [USA]; USNM, Washington. Synonymy: Wood 1954a: 1040.

References: (hb) Chamberlin 1939: 309. (ds) Chamberlin 1939: 309; Leng 1920: 340. (tx) Chamberlin 1939: 309; Hopkins 1915b: 24, 31; Wood, S. L. 1954a: 1040.

spimicollis (Schedl) 1965c: 59 (*Stephanoderes*).

Holotype, sex?; Madagascar, Umgebung Fort Dauphin; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

References: (tx) Schedl 1965c: 59, 1977a: 76, 1979c: 234.

spinus (Schedl) 1979g: 98 (*Lepiceroides*). Holotype, sex?; Papua, Bnlolo, Morobe District. Upper Manki L. A.; Schedl Collection in NHMW, Wien preoccupied by Hagedorn 1909.

Distribution: New Guinea.

Notes: (1) Technically, this is a junior homonym, but the probability that it will become a junior synonym is so high that a replacement name was not considered to be in the best interests of nomenclature.

References: (tx) Schedl 1979g: 98, 1961k: 635; Wood, S. L. 1984b: 226.

squamosus (Hopkins) 1915b: 26 (*Stephanoderes*).

Holotype ♀; Caymanas, Cuba; USNM, Washington.

Distribution: Antilles Islands (Cuba), North America (Colima, Tamaulipas, Veracruz in Mexico/ S Florida, Texas in USA).

Hosts: *Ardisia paniculata*, *Dipholis salicifolia*, *Galactia spiciformis*, *Ichthyomethia communis*, *Lysiloma bahamensis*, *Parthenocissus quinquefolia*, *Phoradendron* sp., *Pithecellobium unguicatum*, *Serjania* sp., *Torruba longifolia*, *Vismia* sp.

References: (ec) Equihua & Atkinson 1986: 633. (hb) Equihua & Atkinson 1986: 633; Wood, S. L. 1954a: 1038. (ds) Atkinson & Equihua 1985c: 357; Atkinson et al. 1991: 161; Blackwelder 1947: 775; Equihua & Atkinson 1986: 633; Estrada & Atkinson 1988: 211; Wood, S. L. 1982b: 890. (tx) Hopkins 1915b: 21, 26; Wood, S. L. 1954a: 1019, 1037–1038, 1982b: 890.

stigmatosus (Schedl) 1951m: 101 (*Stephanoderes*).

Holotype, sex?; Argentinien; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

References: (tx) Schedl 1951l: 284, 1951m: 101, 1979c: 237.

garciac Schedl 1958f: 42 (*Stephanoderes*). Holotype, sex?; Argentinien; Buenos Aires, Isla Martin Garcia; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 175.

References: (tx) Schedl 1958f: 42, 1979c: 102; Wood, S. L. 1989: 175.

styrax (Schedl) 1942a: 177 (*Stephanoderes*). Lectotype, sex?; Malaya, Selangor, Kepong; Schedl Collection in NHMW, Wien, designated by Schedl

1979c: 239.

Distribution: Asia (Malaya).

Hosts: *Styrax benzoin*.

References: (ds) Browne 1961c: 73. (tx) Schedl 1942a: 177, 1979c: 239.

subacuminatus (Schedl) 1942a: 177 (*Stephanoderes*). Lectotype, sex?; Malaya, Pk; Schedl Collection in NHMW, Wien, designated by Schedl

1979c: 239.

Distribution: Asia (Malaya).

References: (ds) Browne 1961c: 73, 1980d: 492. (tx) Schedl 1942a: 177, 1979c: 239.

suspectus Wood 1974a: 22. Holotype ♀; Pandora, Limon, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica/ Nayarit in Mexico/ Panama), South America (Venezuela).

Hosts: *Albizia caribaea*, *Cecropia* sp., *Inga* sp., *Theobroma cacao*.

References: (ds) Wood, S. L. 1982b: 906. (tx) Wood, S. L. 1974a: 22. 1982b: 906.

taihokuensis (Schedl) 1952c: 63 (*Stephanoderes*). Holotype, sex?; Formosa, Taihoku; Schedl Collection in NHMW, Wien.

Figures: Nobuchi 1967: pl. 1.

Distribution: Asia (Taiwan).

References: (ds) Nobuchi 1967: 24. (tx) Nobuchi 1967: 24. pl. 1; Schedl 1952c: 63. 1979c: 249.

teretis Wood 1971: 35. Holotype ♀; Finca La Lola, Limon, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica), South America (Venezuela).

Hosts: *Canavalia villosa*, *Cecropia* sp., *Inga* sp., *Serjania* sp., *Sterculia pruriens*, *Theobroma cacao*.

References: (hb) Wood, S. L. 1982b: 904. (ds) Wood, S. L. 1982b: 904. (tx) Wood, S. L. 1971: 35–36. 1982b: 904.

teteforti (Menier) 1971: 141 (*Stephanoderes*). Holotype ♂; Madagascar Sud, 25 km à l'Est de Tulear, sur la route de Tananarive; MNHN, Paris.

Distribution: Madagascar.

Hosts: *Euphorbia oncoclada*.

References: (tx) Menier 1971: 141; Schedl 1977a: 77.

tredli (Reitter) 1908a: 55 (*Cryptobius*). Holotype, sex?; Agypten; Cairo: NHMB, Budapest.

Distribution: Africa (Egypt).

References: (ds) Kleine 1914a: 15. (tx) Eggers 1940h: 62; Reitter 1908a: 55–56, 1913a: 67; Schedl 1934f: 1639.

trinitatis (Hopkins) 1915b: 28 (*Stephanoderes*). Holotype ♀; Montserrat, Trinidad; USNM, Washington.

Distribution: Antilles Islands (Puerto Rico/ Trinidad).

Hosts: *Guarica trichiloides*.

References: (ds) Blackwelder 1947: 778; Bright 1985c: 175; Martorell 1945: 469; Wolcott 1948: 381. (tx) Bright 1985c: 175; Hopkins 1915b: 22, 28.

tristis (Eichhoff) 1875: 200 (*Stephanoderes*). Holotype, sex?; Japan; IRSNB, Brussels.

Distribution: Asia (Japan).

References: (ds) Blandford 1894c; Hagedorn 1919d: 46; Kleine 1913b: 121, 1914b: 257; Murayama 1954b: 199; Nobuchi 1985c: 16. (tx) Blandford 1894d; Eichhoff 1875: 200, 1877a: 117, 1878b: 150; Hagedorn 1910a: 88; Hopkins 1915b: 24; Murayama 1934c: 298, 1954b: 199; Schedl 1934f: 1640. 1958k: 143.

apatoides Eichhoff 1875: 201 (*Dryocoetes*). Holotype, sex?; Japan; IRSNB, Brussels. Synonymy: Schedl 1958k: 143.

References: (tx) Eichhoff 1875: 201; Schedl 1958k: 143.

trivialis Wood 1974a: 20. Holotype ♀; Santa Ana, San Jose, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica/ Veracruz in Mexico/ Panama), South America (Colombia/ Venezuela).

Hosts: *Cuparia guatemalensis*, *Ficus* sp., *Ochroma* sp., *Serjania* sp., *Vismia guianensis*.

References: (ds) Wood, S. L. 1982b: 887. (tx) Wood, S. L. 1974a: 20. 1982b: 887.

tuberosus (Schedl) 1942d: 19 (*Stephanoderes*). Lectotype, sex?; Java, Buitenzorg; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 258.

Distribution: Indonesia (Java).

References: (tx) Schedl 1942d: 19. 1979c: 258.

tungamcansolus (Schedl) 1939i: 355 (*Stephanoderes*). Lectotype, sex?; Ukerewe, Tanganyika; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 258.

Distribution: Africa (Tanzania/ Zambia).

Hosts: *Bridelia scleroneuroides*.

References: (ds) Beaver & Loytyniemi 1989; Schedl 1961k: 626. (tx) Beaver & Loytyniemi 1989; Schedl 1939g: 170, 1939i: 355–356, 1957d: 56, 1961k: 626, 1979c: 258.

vesculus Wood 1974a: 21. Holotype ♀; Ocosingo Valley, Chiapas, Mexico; Wood Collection.

Distribution: North America (Chiapas in Mexico).

References: (ds) Wood, S. L. 1982b: 898. (tx) Wood, S. L. 1974a: 21. 1982b: 898.

vitis Browne 1970: 554. Holotype ♀; South Africa: Transvaal; BMNH, London.

Distribution: Africa (South Africa).

Hosts: *Vitis* sp.

References: (tx) Browne 1970: 554, 556; Schedl 1972q: 256.

winkleri (Reitter) 1907: 192 (*Stephanoderes*). Holotype ♀; Kamerun; NHMB, Budapest.

Figures: Numberg 1965a: 35, 1965b: pl. 2.

Distribution: Africa (Cameroon).

References: (ds) Hagedorn 1910d: 46; Kleine 1913b: 121, 1914b: 310, 1934a: 145; Numberg 1952: 20, 1965b: 20. (tx) Hagedorn 1910a: 88; Hopkins 1915b: 24; Numberg 1952: 20, 1965a: 35, 1965b: 26, pl. 2; Reitter 1907: 192; Schedl 1959q: 708, 1965e: 367.

xanthophloeae (Schedl) 1957d: 58 (*Stephanoderes*). Holotype, sex?; Kenya: Nairobi, Rift Valley; Schedl Collection in NHMW, Wien.

Distribution: Africa (Kenya).

Hosts: *Acacia xanthophloea*.

References: (cc) Schedl 1961k: 630. (hb) Schedl 1961k: 630. (ds) Schedl 1961k: 630. (tx) Schedl 1957d: 58, 1961k: 630, 1979c: 270.

Genus *Trischidias* Hopkins

TRISCHIDIAS HOPKINS 1915b: 7, 12. Type-species:

Trischidias georgiae Hopkins, original designation.

Keys: Wood 1954a: 1067, 1982b: 872.

References: (hb) Wood, S. L. 1986a: 91. (ds) Wood, S. L. 1986a: 91. (tx) Arnett 1960: 1043, 1968: 1043; Hopkins 1915b: 7, 12; Numberg 1956: 141; Wood, S. L. 1954a: 962, 981, 1067, 1982b: 872, 1986a: 92.

atoma (Hopkins) 1915b: 15 (*Hypothenemus*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington.

Distribution: North America (District of Columbia, Florida, Georgia, Kansas, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, South Carolina, Texas, Tennessee, Virginia, West Virginia in USA), South America (Brazil).

Hosts: *Acer rubrum*, *Asimina trilobata*, *Carya* spp., *Castanea dentata*, *Liriodendron tulipifera*, *Pinus* sp., *Quercus marylandica*, *Rhizophora mangle*, *Rhus toxicodendron*, *Robinia pseudacacia*, *Salix nigra*, *Ulmus rubra*.

References: (hb) Beal & Massey 1945: 118; Chamberlin 1939: 293; Deyrup & Atkinson 1987a: 66; Wood, S. L. 1953: 219, 1954a: 1068, 1982b: 872. (ds) Beal & Massey 1945: 118; Blatchley & Leng 1916: 596; Chamberlin 1939: 293; Deyrup 1981b: 7; Deyrup & Atkinson 1987a: 66; Kleine 1934a: 143; Leng 1920: 339; Smith, J. B. 1910: 401; Wood, S. L. 1982b: 872. (tx) Beal & Massey 1945: 118; Blatchley & Leng 1916: 596; Chamberlin 1939: 293; Hopkins 1915b: 15; Schedl 1962r: 89; Wood, S. L. 1953: 219, 1954a: 960, 976, 1068, 1982b: 872–873.

impressifrons Hopkins 1915b: 15 (*Hypothenemus*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1068.

References: (hb) Chamberlin 1939: 293. (ds) Chamberlin 1939: 293; Kleine 1934a: 143; Leng 1920: 339. (tx) Chamberlin 1939: 293; Hopkins 1915b: 15; Wood, S. L. 1954a: 979, 1068.

marylandica Hopkins 1915b: 15 (*Hypothenemus*). Holotype ♀; Pregnall, South Carolina [USA]; USNM, Washington. Synonymy: Wood 1954a: 960.

References: (hb) Blackman 1922b: 83; Chamberlin 1939: 290. (ds) Blackman 1922b: 83; Blatchley & Leng 1916: 596; Chamberlin 1939: 290; Kirk 1969; Kleine 1934a: 143; Leng 1920: 339. (tx) Blackman 1922b: 83; Blatchley & Leng 1916: 596; Chamberlin 1939: 290; Hopkins 1915b: 15; Schedl 1962r: 89; Wood, S. L. 1954a: 960, 1068.

robiniae Hopkins 1915b: 15 (*Hypothenemus*). Holotype ♀; Chevy Chase, Maryland [USA]; USNM, Washington. Synonymy: Wood 1954a: 1068.

References: (hb) Beal & Massey 1945: 118;

Chamberlin 1939: 293. (ds) Beal & Massey 1945: 118; Blatchley & Leng 1916: 597; Chamberlin 1939: 293; Kleine 1934a: 144; Leng 1920: 339. (tx) Beal & Massey 1945: 118; Blatchley & Leng 1916: 597; Chamberlin 1939: 293; Hopkins 1915b: 15; Schedl 1962r: 89; Wood, S. L. 1954a: 960, 1068.

toxicodendri Hopkins 1915b: 15 (*Hypothenemus*). Holotype ♀; Morgantown, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1954a: 1068.

References: (hb) Blackman 1922b: 83; Chamberlin 1939: 290. (ds) Blackman 1922b: 83; Blatchley & Leng 1916: 597; Chamberlin 1939: 290; Kleine 1934a: 144; Leng & Mutchler 1920: 339. (tx) Blackman 1922b: 83; Blatchley & Leng 1916: 597; Chamberlin 1939: 290; Hopkins 1915b: 15; Schedl 1962r: 89–90; Wood, S. L. 1954a: 960, 1068.

exigua Wood 1986c: 273. Holotype ♀; Campo Experimental, INIF, Escarcega, Campeche, Mexico; Wood Collection.

Distribution: North America (Campeche in Mexico/S Florida in USA).

Hosts: *Belotia campbelli*.

References: (hb) Deyrup 1987; Deyrup & Atkinson 1987a: 66. (ds) Deyrup 1987; Deyrup & Atkinson 1987a: 66; Estrada & Atkinson 1988: 211. (tx) Wood, S. L. 1986c: 273.

georgiae Hopkins 1915b: 12. Holotype ♀; Brunswick, Georgia [USA]; USNM, Washington.

Distribution: North America (Georgia).

References: (hb) Chamberlin 1939: 287; Wood, S. L. 1954a: 1070. (ds) Blatchley & Leng 1916: 594; Chamberlin 1939: 287; Leng 1920: 339; Wood, S. L. 1982b: 974. (tx) Blatchley & Leng 1916: 594; Chamberlin 1939: 287; Hopkins 1915b: 12; Wood, S. L. 1954a: 979, 1068, 1070, 1982b: 874.

minutissima Wood 1954a: 1069. Holotype ♀; Sugar Loaf Key, Florida [USA]; SMUK, Lawrence.

Distribution: North America (S Florida in USA).

Hosts: *Rhizophora mangle* roots.

References: (hb) Wood, S. L. 1954a: 1069, 1982b: 874. (ds) Wood, S. L. 1982b: 874. (tx) Schedl 1962r: 89; Wood, S. L. 1954a: 1068–1069, 1982b: 874.

nigrina (Schedl) 1967d: 7 (*Ernoporus*). Holotype, sex?, Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

Notes: (3) A probable synonym of *atoma* Hopkins.

References: (tx) de Ruette 1970: 101; Schedl 1967d: 7, 1979c: 167.

spinata (Schedl) 1977d: 282 (*Miocryptalus*). Holotype, sex?, Ghana, Volta Region. Ho, N 6 degrees 70', E 0 degrees 03'; NHMB, Budapest.

Distribution: Africa (Ghana).

References: (tx) Schedl 1977d: 282.

Genus *Periocryphalus* Wood

PERIOCRYPHALUS WOOD 1971: 33. Type-species: *Periocryphalus pullus* Wood, original designation.
References: (hb) Wood, S. L. 1986a: 92. (ds) Wood, S. L. 1986a: 92. (tx) Wood, S. L. 1971: 33, 1986a: 92.
pullus Wood 1971: 33. Holotype ♀; 40 km SE Socopo, Barinas, Venezuela; Wood Collection.
Distribution: South America (Brazil/Venezuela).

Hosts: *Liana*.
References: (hb) Wood, S. L. 1971: 33. (ds) Beaver 1974a. (tx) Wood, S. L. 1971: 33.
sobrinus Wood 1974a: 22. Holotype ♀; 260 km N Xavantina, Mato Grosso, Brazil; BMNH, London.
Distribution: South America (Brazil).
Hosts: *Protium* sp., *Siparuna* cf. *guianensis*.
References: (hb) Beaver 1974a. (ds) Beaver 1974a. (tx) Wood, S. L. 1974a: 22.

Tribe Corthylini LeConte

Corthyli

References: Blandford 1904: 248; LeConte 1876: 346–347; LeConte & Horn 1883: 516.

Corthyliidae

References: Eichhoff 1878b: 421.

Corthylinae

References: Beal & Massey 1945: 104; Hagedorn 1909: 162, 1910a: 24, 137, 1910d: 90; Hopkins 1915c: 224.

Corthylini

References: Leng 1920: 339; Lucas 1920: 339; Wood, S. L. 1978a: 114, 1982b: 69, 1986a: 93–100.

Pityophthoridae

References: Eichhoff 1878b: 173.

Araptidae

References: Eichhoff 1878b: 305.

Amphiceranidae

References: Eichhoff 1878b: 460.

Corthylini: Pityophthorina

Pityophthoridae

References: Eichhoff 1878b: 173.

Pityophthorinae

References: Nusslin 1911: 431.

Pityophthorina

References: Balachowsky 1949a: 232; Nunberg 1954: 17; Wood, S. L. 1978a: 115, 1982b: 73, 1986a: 94, 97–98.

Araptidae

References: Eichhoff 1878b: 305.

Genus *Mimiocurus* Schedl

MIMIOCURUS SCHEDL 1957d: 72. Type-species: *Mimiocurus acuminatus* Schedl, monobasic.

Micracidendron Schedl 1957d: 71. Type-species: *Micracidendron montanum* Schedl, monobasic. Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1957d: 71, 1962j: 29, 1977b: 120; Wood, S. L. 1984b: 226.

Mimiophthorus Schedl 1957d: 77. Type-species: *Brachydendrus montanus* Schedl, nomen nudum, an error for *B. congopus* Schedl, original designation (Schedl 1962j: 63). Synonymy: Wood 1984b: 226.

References: (tx) Schedl 1957d: 77, 1962j: 63; Wood, S. L. 1984b: 226.

References: (hb) Wood, S. L. 1986a: 97. (ds) Schedl 1962j: 52, 63, 1977b: 120; Wood, S. L. 1986a: 97. (tx) Schedl 1957d: 72, 1962j: 52, 63, 1977b: 120; Wood, S. L. 1986a: 97.

acuminatus Schedl 1957d: 72. Holotype, sex?; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: Liana.

References: (ce) Schedl 1962j: 53. (hb) Schedl

1962j: 53. (ds) Schedl 1962j: 53. (tx) Schedl 1957d: 72, 1962j: 53, 1979c: 11.

beesoni Wood 1989: 182. Holotype ♂; Amarampalam R., Nilambur, Madras, India; FRI, Dehra Dun.

Distribution: Asia (Tamil Nadu in India).

Hosts: *Tiliacora acuminata*.

References: (tx) Wood, S. L. 1989: 182.

congopus (Schedl) 1952i: 14 (*Brachydendrus*). Holotype ♀; Kivu, Mulungu, Congo Belge; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Asparagus* sp., *Chlorocodon whiteii*, *Leucas* sp.

Notes: (3) Schedl 1962j: 63 (treated in *Mimiophthorus*).

References: (hb) Schedl 1962j: 63. (ds) Mayne & Donis 1962: 302; Schedl 1962j: 63. (tx) Schedl 1952i: 14–15, 1955f: 258, 262, 1957d: 77, 1962j: 63–64, 1963a: 30, 1964m: 317, 1979c: 62.

kikusae (Schedl) 1957d: 77 (*Mimiophthorus*). Holotype, sex?; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.

Distribution: Africa (Ivory Coast/ Zaire).

Hosts: *Salacia* cf. *erecta*.

References: (ce) Schedl 1962j: 67. (hb) Schedl 1958d: 185, 1962j: 67. (ds) Schedl 1962j: 67. (tx) Schedl 1957d: 77, 1961m: 83–84, 1962j: 67, 1979c: 132.

kikusuae occidentalis Schedl 1961m: 83 (*Mimiophthorus*). Lectotype, sex?; Cote d'Ivoire; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 132.

References: (ds) Schedl 1964e: 68, 1972e: 282. (tx) Schedl 1961m: 83, 1962j: 68, 1979c: 132.

montanus (Schedl) 1957d: 71 (*Micracidendron*). Holotype, sex?; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Alchornea hirtella*.

References: (hb) Schedl 1958d: 185–194, 1962j: 29. (ds) Schedl 1962j: 29. (tx) Schedl 1957d: 71, 1962j: 29–30, 1979c: 158.

monticulus Wood 1989: 178. Holotype, sex?; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren, automatic.

Distribution: Africa (Zaire).

References: (ds) Schedl 1962j: 55. (tx) Schedl 1962j: 55; Wood, S. L. 1989: 178.

montanus Schedl 1957d: 73. Holotype, sex?; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren, preoccupied by Schedl 1957.

References: (ce) Schedl 1958d: 185. (tx) Schedl 1957b: 156–157, 1957d: 72–73, 1962j: 55, 1979c: 158; Wood, S. L. 1989: 178.

orientalis (Schedl) 1972i: 51 (*Mimiophthorus*). Holotype ♀; New Guinea (NW), Nabire; BPBM, Honolulu.

- Distribution: New Guinea.
References: (tx) Schedl 1972i: 51, 1979c: 181.
- rosseli (Schedl)** 1973e: 88 (*Taphroterus*). Holotype, sex?; New Guinea: Ahaleti, Rossel Island, Milne Bay District; AMNH, New York.
Distribution: New Guinea.
Notes: (3) The holotype was examined by Wood and referred to *Miniocurus* (Wood 1983 private notes, p. 16).
References: (tx) Schedl 1973e: 88, 1979c: 213.
- rugicollis (Schedl)** 1957d: 78 (*Mimiophthorus*). Holotype, sex?; Congo Belge: Kivu, Hembé-Bitale; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Salacia* cf. *erecta*.
References: (ec) Schedl 1962j: 70. (hb) Schedl 1962j: 70. (ds) Schedl 1962j: 70, 1971g: 191. (tx) Schedl 1957d: 78, 1962j: 70, 1979c: 216.
- ruwenzoriensis (Schedl)** 1963a: 30 (*Mimiophthorus*). Holotype, sex?; Uganda, Ruwenzori Range, Nyangasani Valley, 13,000 ft.; BMNH, London.
Distribution: Africa (Uganda).
Hosts: *Senecio eriocaulon*.
References: (tx) Schedl 1963a: 30, 1979c: 217.
- setifer (Schedl)** 1957b: 156 (*Mimips*). Lectotype, sex?; Tanganyika, Lushoto; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 226.
Distribution: Africa (Tanzania).
Hosts: *Syzygium* sp.
Notes: (1) Schedl 1979c: 226 (to *Mimiophthorus*, = *Miniocurus*). This species has also been cited in *Dendrocranulus*.
References: (hb) Gardner 1957: 31. (ds) Schedl 1962j: 60. (tx) Schedl 1957b: 156, 1962j: 60, 1979c: 226.
- tongatapai (Schedl)** 1979f: 106 (*Mimiophthorus*). Holotype, sex?; Tonga, Tongatapu: Tupon College; New Zealand DSIR, Auckland.
Distribution: Tonga Islands.
References: (tx) Beaver 1987: 68; Schedl 1979f: 106.
- umbratus (Schedl)** 1972e: 290 (*Mimiophthorus*). Holotype ♀; [Ghana] Ashanti Region: Kumasi, Nhasu 330 m, N 6-43, W 1-36; NHMB, Budapest.
Distribution: Africa (Ghana).
References: (tx) Schedl 1972e: 290, 1979c: 260.
- uiseriatius (Schedl)** 1979d: 451 (*Mimiophthorus*). Holotype, sex?; Kamerun, Mukonje; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon).
References: (tx) Schedl 1979d: 451.
- 232; Hagedorn 1910a: 96-97, 1910d: 69; Wood, S. L. 1982b: 916-917, 1986a: 97-98.
- boliviae (Schedl)** 1961i: 224 (*Stephanopodius*). Holotype ♀; Bolivia, Do Santa Cruz, Prov. Jaíta, Buenavista; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (tx) Schedl 1961i: 224, 1979c: 42.
- brasiliensis (Schedl)** 1972g: 58 (*Stephanopodius*). Holotype, sex?; Brasilien, Corcovado, Gramabara; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 58, 1979c: 45.
- granulatum Blandford** 1904: 232. Holotype ♂; Bugaba, Chiriqui, Panama; BMNH, London.
Distribution: North America (Costa Rica/ Campeche in Mexico/ Panama).
Hosts: *Spondias mombin*.
Notes: (3) Wood 1971: 43 (described female).
References: (ec) Equihua & Atkinson 1986: 634; Matthews 1969: 111. (hb) Equihua & Atkinson 1986: 634. (ds) Blackwelder 1947: 782; Equihua & Atkinson 1986: 634; Estrada & Atkinson 1988: 212; Kleine 1913b: 139, 1914b: 362; Wood, S. L. 1982b: 917. (tx) Blandford 1904: 232; Hagedorn 1910a: 97; Hopkins 1914: 130; Lucas 1920: 619; Wood, S. L. 1982b: 917. (ms) Lucas 1920: 619.
- subulatum Wood** 1971: 43. Holotype ♀; 9 km S Barancas, Barinas, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Astronium graveolens*.
References: (hb) Wood 1971: 43.

Genus *Dendroterus* Blandford

- DENDROTERUS BLANDFORD** 1904: 233. Type-species: *Dendroterus mexicanus* Blandford, subsequent designation by Hopkins 1914: 120.
- Plesiophthorus** Schedl 1940a: 343. Type-species: *Plesiophthorus perspectus* Schedl, monobasic. Synonymy: Wood 1959b: 2.
References: (tx) Schedl 1940a: 343, 1964m: 312; Wood, S. L. 1959b: 2, 1986a: 97.
- Xylochilus** Schedl 1956b: 30. Type-species: *Xylochilus insularis* Schedl = *Dendroterus sallaci* Blandford, original designation. Synonymy: Wood 1972c: 191.
References: (tx) Schedl 1956b: 30; Wood, S. L. 1972c: 191, 1986a: 97.
Keys: Wood 1982b: 918.
- References: (hb) Bright & Stark 1973: 96; Wood, S. L. 1982b: 917, 1986a: 97. (ds) Bright & Stark 1973: 96; Estrada & Atkinson 1988: 212; Wood, S. L. 1982b: 917, 1986a: 97. (tx) Arnett 1960: 1045, 1968: 1045; Blandford 1904: 233; Hopkins 1914: 120, 1915b: 9; Wood, S. L. 1982b: 917, 1986a: 97.
- cognatus Wood** 1971: 46. Holotype ♀; 5 km W El Salto, Durango, Mexico; Wood Collection.
Distribution: North America (Nayarit in Mexico).
Hosts: *Bursera* sp.
References: (hb) Wood, S. L. 1982b: 923. (ds)

Genus *Styphlosoma* Blandford

- STYPHILOSOMA BLANDFORD** 1904: 232. Type-species: *Styphlosoma granulatum* Blandford, monobasic.
References: (hb) Wood, S. L. 1986a: 97-98. (ds) Wood, S. L. 1986a: 97-98. (tx) Blandford 1904:

- Wood, S. L. 1982b: 923. **(tx)** Wood, S. L. 1971: 46, 1982b: 923.
- decipiens** Wood 1959b: 5. Holotype ♂; 3 miles NW Tequila, Jalisco, Mexico; SMUK, Lawrence.
Distribution: North America (Jalisco, Oaxaca in Mexico).
Hosts: *Bursera* sp.
References: **(hb)** Atkinson et al. 1986: 68; Burgos & Saucedo 1983: 109; Wood, S. L. 1982b: 926. **(ds)** Atkinson et al. 1986: 68; Burgos & Saucedo 1983: 109; Wood, S. L. 1982b: 926. **(tx)** Wood, S. L. 1959b: 5, 1982b: 926.
- defectus** Wood 1971: 45. Holotype ♀; near Rincon de Osa, Puntarenas, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica/Panama).
Hosts: *Bursera simarubra*.
References: **(hb)** Wood, S. L. 1982b: 923. **(ds)** Wood, S. L. 1982b: 923. **(tx)** Wood, S. L. 1971: 45–46, 1982b: 923.
- eximius** Wood 1971: 44. Holotype ♀; Lago Amatitlan, Esquinilla, Guatemala; Wood Collection.
Distribution: North America (Guatemala).
Hosts: *Bursera simarubra*.
References: **(hb)** Wood, S. L. 1982b: 923. **(ds)** Wood, S. L. 1982b: 923. **(tx)** Wood, S. L. 1971: 44, 1982b: 923.
- fossifrons** Wood 1984e: 114. Holotype ♂; Estacion de Biologia, Jalisco, Mexico; Wood Collection.
Distribution: North America (Jalisco in Mexico).
Hosts: *Bursera* sp.
References: **(ec)** Equihua & Atkinson 1986: 633. **(hb)** Equihua & Atkinson 1986: 633. **(ds)** Equihua & Atkinson 1986: 633. **(tx)** Wood, S. L. 1984e: 114.
- luteolus** (Schedl) 1951m: 111 (*Plesiophthorus*). Holotype ♂; Mexico; Schedl Collection in NHMW, Wien.
Figures: Atkinson et al. 1986: 77.
Distribution: North America (Chiapas, Morelos, Nayarit, Puebla in Mexico).
Hosts: *Bursera* spp.
References: **(ec)** Equihua & Atkinson 1986: 634. **(hb)** Atkinson et al. 1986: 70; Equihua & Atkinson 1986: 634; Wood, S. L. 1954a: 107, 1982b: 926. **(ds)** Atkinson & Equihua 1988: 90; Atkinson et al. 1986: 70; Equihua & Atkinson 1986: 634; Estrada & Atkinson 1988: 211; Wood, S. L. 1982b: 926. **(tx)** Atkinson et al. 1986: 77; Schedl 1951m: 111–112, 1952e: 123, 1979c: 143; Wood, S. L. 1954a: 1071, 1959b: 2, 1972e: 194, 1982b: 926.
- mnudus** Wood 1959b: 3. Holotype ♂; Tehuiztingo, Puebla, Mexico; SMUK, Lawrence. Synonymy: Wood 1972e: 194.
References: **(tx)** Schedl 1964m: 312; Wood, S. L. 1959b: 3, 1972e: 194.
- mexicanus** Blandford 1904: 233. Holotype ♀; Tehuantepec, Oaxaca, Mexico; BMNH, London.
Distribution: North America (Jalisco, Morelos, Nayarit, Oaxaca in Mexico).
Hosts: *Bursera* spp.
References: **(ec)** Equihua & Atkinson 1986: 634. **(hb)** Atkinson & Equihua 1985c: 358; Atkinson et al. 1986: 70; Burgos & Saucedo 1983: 105; Equihua & Atkinson 1986: 634; Wood, S. L. 1982b: 924. **(ds)** Atkinson & Equihua 1985c: 358, 1988: 90; Atkinson et al. 1986: 70; Blackwelder 1947: 782; Burgos & Saucedo 1983: 105; Equihua & Atkinson 1986: 634; Ferrer 1942; Hagedorn 1910d: 69; Kleine 1913b: 139, 1914b: 351; Wood, S. L. 1982b: 924. **(tx)** Blandford 1904: 233; Hagedorn 1910a: 91; Hopkins 1914: 120; Lucas 1920: 230; Schedl 1940a: 343; Wood, S. L. 1959b: 2, 1972e: 194, 1976a: 348, 1982b: 924. **(ms)** Lucas 1920: 230.
- unbratus** Schedl 1937h: 168 (*Conophthoracraulus*). Holotype ♀; Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1976a: 348.
References: **(tx)** Schedl 1937h: 168–169, 1940a: 346, 1964m: 312; Wood, S. L. 1976a: 348.
- confinis** Wood 1959b: 6. Holotype ♂; 23 km NW Magdalena, Jalisco, Mexico; SMUK, Lawrence. Synonymy: Wood 1972e: 196.
References: **(tx)** Schedl 1959b: 6, 1964m: 312; Wood, S. L. 1959b: 6, 1972e: 194.
- modicus** Wood 1984e: 115. Holotype ♀; Volcan Ceboruco, Nayarit, Mexico, 1100 m; Wood Collection.
Distribution: North America (Nayarit in Mexico).
Hosts: *Bursera* sp.
References: **(tx)** Wood, S. L. 1984e: 115.
- parilis** Wood 1971: 44. Holotype ♀; Santa Ana, San Jose, Costa Rica, 1300 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Bursera simarubra*.
References: **(hb)** Wood, S. L. 1982b: 922. **(ds)** Wood, S. L. 1982b: 922. **(tx)** Wood, S. L. 1971: 44, 1982b: 922.
- perspectus** (Schedl) 1940a: 343 (*Plesiophthorus*). Holotype ♀; Omiltilme, Guerrero, Mexico; Schedl Collection in NHMW, Wien.
Distribution: North America (Guerrero in Mexico).
Notes: (1) Wood 1959b: 2 (to *Dendroterus*). (3) Wood 1982b: 924 (redescribed female).
References: **(hb)** Atkinson et al. 1986: 70. **(ds)** Atkinson et al. 1986: 70; Blackwelder 1947: 781; Ferrer 1942. **(tx)** Schedl 1940a: 343, 1952e: 122, 1979c: 191; Wood, S. L. 1959b: 2, 1982b: 924.
- resolutus** Wood 1971: 45. Holotype ♀; Playa del Coco, Guanacaste, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Bursera simarubra*.
References: **(hb)** Wood, S. L. 1982b: 925. **(ds)** Wood, S. L. 1982b: 925. **(tx)** Wood, S. L. 1971: 45, 1982b: 925.

sallaei Blandford 1904: 233. Holotype ♀; Veracruz, Veracruz, Mexico; BMNH, London.

Distribution: North America (Costa Rica/ Chiapas, Jalisco, Nayarit, Oaxaca, Tamaulipas, Tres Marias Islands in Mexico).

Hosts: *Bursera sinarubra*, *B. sp.*

References: (cc) Equihua & Atkinson 1986: 634. (hb) Equihua & Atkinson 1986: 634; Wood, S. L. 1982b: 924. (ds) Atkinson & Equihua 1988: 90; Blackwelder 1947: 782; Equihua & Atkinson 1986: 634; Ferrer 1942; Hagedorn 1910d: 69; Kleine 1913b: 139, 1914b: 351; Wood, S. L. 1982b: 924. (tx) Blandford 1904: 233; Hagedorn 1910a: 91; Lucas 1920: 230; Schedl 1940a: 343; Wood, S. L. 1959b: 2, 1972e: 195, 1982b: 924. (ms) Lucas 1920: 230.

insularis Schedl 1956b: 31 (*Xylochilus*). Holotype ♂; Arroyo Hondo, Maria Madre, Tres Marias Islands, Gulf of California, Mexico; CAS, San Francisco. Synonymy: Wood 1972e: 195.

References: (hb) Wood, S. L. 1982b: 921. (ds) Wood, S. L. 1982b: 921. (tx) Schedl 1956b: 31, 1979c: 126; Wood, S. L. 1972e: 195, 1982b: 921.

sodalis Wood 1971: 44. Holotype ♀; Volcan de Agua, Esquinla, Guatemala; Wood Collection.

Distribution: North America (Guatemala).

Hosts: *Bursera sinarubra*.

References: (tx) Wood, S. L. 1971: 44.

striatus (LeConte) 1868: 156 (*Cryphalus*). Lectotype ♀; Cape San Lucas, Baja California, Mexico; MCZ, Cambridge, designated by Wood 1982b: 927. Distribution: North America (Baja California in Mexico/ S California in USA).

Hosts: *Bursera microphylla*.

Notes: (1) Wood 1954a: 1071 (to *Plesiophthorus*, 1972e: 195 to *Dendroterus*).

References: (hb) Bright & Stark 1973: 96; Wood, S. L. 1954a: 1071, 1982b: 927. (ds) Blatchley & Leng 1916: 631; Bright & Stark 1973: 96; Chamberlin 1939: 295; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 45; Henshaw 1882: 269, 1885: 148; Horn 1894: 359; Leng 1920: 340; Wood, S. L. 1982b: 927. (tx) Blatchley & Leng 1916: 631; Hagedorn 1910a: 88; Hopkins 1915b: 33; LeConte 1868: 156, 1876: 356; Swaine 1909: 119; Wood, S. L. 1954a: 960, 1071, 1959b: 2, 1972e: 195, 1982b: 927. *californicus* Schedl 1952e: 123 (*Plesiophthorus*).

Holotype ♀; Angeles Bay, Gulf of California, Baja California, Mexico; CAS, San Francisco. Synonymy: Wood 1972e: 195.

References: (hb) Wood, S. L. 1954a: 1071. (tx) Schedl 1952e: 122–123, 1979c: 50; Wood, S. L. 1954a: 1071, 1959b: 2, 1972e: 195.

texasus Wood 1959b: 4. Holotype ♂; Presidio, Texas [USA]; USNM, Washington.

Distribution: North America (SW Texas in USA).

Hosts: *Jatropha spatulata*.

References: (hb) Wood, S. L. 1982b: 920. (ds) Wood, S. L. 1982b: 920. (tx) Schedl 1964m: 312; Wood, S. L. 1959b: 4, 1982b: 920.

Genus *Phloeoterus* Wood

PHLOEOTERUS WOOD 1984e: 117. Type-species: *Phloeoterus burserae* Wood, original designation. References: (hb) Wood, S. L. 1986a: 97. (ds) Wood, S. L. 1986a: 97. (tx) Wood, S. L. 1984e: 117, 1986a: 97.

burserae Wood 1984e: 117. Holotype ♀; Estacion de Biologia, Chamela, Jalisco, Mexico; Wood Collection.

Distribution: North America (Jalisco in Mexico).

Hosts: *Bursera instabilis*.

References: (cc) Equihua & Atkinson 1986: 634. (hb) Equihua & Atkinson 1986: 634. (ds) Equihua & Atkinson 1986: 634. (tx) Muskus, A. 1984: 100; Wood, S. L. 1984e: 117.

Genus *Araptus* Eichhoff

ARAPTUS EICHHOFF 1872a: 136. Type-species: *Araptus rufopalliatius* Eichhoff, monobasic.

Neodryocotes Eggers 1933b: 9. Type-species: *Neodryocotes hymenaeae* Eggers, monobasic. Synonymy: Blackman 1942a: 180, Wood 1973c: 170.

References: (tx) Blackman 1942a: 178–181; Bright 1972d: 91; Diakonoff 1938: 12–14; Eggers 1933b: 9, 1936a: 388–390, 1940a: 123–141; Schedl 1937h: 165, 1938a: 160–167, 177, 1939e: 344, 1951m: 72, 1962p: 206, 1964m: 311; Wood, S. L. 1973c: 170, 1986a: 97.

Thamniophthorus Schedl 1938a: 174. Type-species: *Thamniophthorus volastos* Schedl, subsequent designation by Blackman 1942a: 178. Synonymy: Wood 1973c: 170.

References: (tx) Arnett 1960: 1045, 1968: 1045; Blackman 1942a: 178; Schedl 1938a: 174; Wood, S. L. 1961a: 41–48, 1973c: 170, 1986a: 97.

Neopityophthorus Schedl 1938a: 180. Type-species: *Pityophthorus laevigatus* Eggers, subsequent designation by Wood 1982b: 928. Synonymy: Schedl 1951m: 72.

References: (tx) Blackman 1942a: 180; Eggers 1940a: 123–141; Schedl 1937h: 165, 1938a: 160–167, 180, 1951m: 72; Wood, S. L. 1982b: 928, 1986a: 97.

Sphenoceros Schedl 1939j: 565. Type-species: *Sphenoceros limax* Schedl, monobasic. Synonymy: Wood 1973c: 170.

References: (tx) Schedl 1939j: 565–566; Wood, S. L. 1973c: 170, 1986a: 97.

Hypertenus Hagedorn 1950: 164 (in Schedl 1950i: 164). Type-species: *Hypertenus ritteri*, nomen nudum, = *Sphenoceros limax* Schedl. Nomen nudum, no status.

References: (tx) Hagedorn 1950: 164 (in Schedl 1950i); Wood, S. L. 1986a: 97.

Brachydendrus Schedl 1951m: 114. Type-species: *Brachydendrus eggersi* Schedl, monobasic. Synonymy: Wood 1984b: 224.

- References: (tx) Schedl 1951m: 114, 1957d: 1–162; Wood, S. L. 1984b: 224, 1986a: 97.
- Gnathocranus* Schedl 1951m: 116. Type-species: *Gnathocranus novateutonicus* Schedl, monobasic. Synonymy: Wood 1984b: 224.
- References: (tx) Schedl 1951m: 116; Wood, S. L. 1984b: 224, 1986a: 97.
- Gnathoborus* Schedl 1970e: 93. Type-species: *Breviophthorus argentiniae* Schedl, original designation. Synonymy: Wood 1984b: 224.
- References: (tx) Schedl 1970e: 93; Wood, S. L. 1984b: 224, 1986a: 97.
- Keys: Wood 1982b: 928 for North and Central America, Bright 1972d: 91 for Jamaica.
- References: (hb) Wood, S. L. 1982b: 927, 1986a: 97. (ds) Hagedorn 1910d: 77; Wood, S. L. 1982b: 927, 1986a: 97. (tx) Eichhoff 1872a: 136, 1878b: 35, 37, 305; Hagedorn 1909a: 743, 1909b: 412, 1910a: 110; Hopkins 1914: 117, 1915c: 228; Schedl 1962j: 28; Wood, S. L. 1973c: 170, 1982b: 927–928, 1986a: 97.
- accinctus** Wood 1974a: 44. Holotype ♂; 3 km SE Acatlan, Puebla, Mexico; Wood Collection.
- Distribution: North America (Puebla in Mexico).
- Hosts: Liana, in pith.
- References: (hb) Atkinson et al. 1986: 68; Wood, S. L. 1982b: 938. (ds) Atkinson et al. 1986: 68; Wood, S. L. 1982b: 938. (tx) Wood, S. L. 1974a: 44, 1982b: 938.
- amazonicus** (Eggers) 1936a: 391 (*Neodryocoetes*). Holotype ♂; Brasilien, Amazonas, Manaos; Eggers Collection, in NHMW, Wien.
- Distribution: South America (Brazil).
- References: (ds) Blackwelder 1947: 781. (tx) Eggers 1936a: 391–392; Schedl 1938a: 177, 1979c: 18.
- araucariae** (Schedl) 1966f: 109 (*Conophthoracranulus*). Holotype ♀?; Argentinien, Misiones, Dep. Concep., Sta. Maria; Schedl Collection in NHMW, Wien.
- Distribution: South America (Argentina).
- Hosts: *Araucaria brasiliensis*, in seeds.
- References: (tx) Schedl 1966e: 43, 1966f: 109, 1979c: 24.
- araujiae** Brethes 1921: 165 (*Xyleborus*). Syntypes, sex?; La Plata, Argentina; Museo Argentino de Ciencias Naturales, Buenos Aires.
- Figures: Brethes 1921: 166 (adult, antenna).
- Distribution: South America (Argentina/ Brazil).
- Hosts: *Araujia scricifera*.
- Notes: (1) Schedl 1939m: 169 (to *Neodryocoetes*, = *Araptus*).
- References: (cn) Costa Lima 1956. (hb) Costa Lima 1956; Viana 1964: 125. (ds) Blackwelder 1947: 781; Costa Lima 1956; Schedl 1972g: 43, 1973d: 155; Viana 1964: 125. (tx) Brethes 1921: 165; Costa Lima 1956; Schedl 1939d: 412, 1939m: 169, 1951h: 284, 1952a: 446.
- longicollis* Schedl 1938a: 179 (*Neodryocoetes*). Holotype, sex?; Argentinien, Prov. Tucuman; Eggers Collection, in NHMW, Wien. Synonymy: Costa Lima 1956: 288.
- Notes: (3) Schedl 1938a: 25 gives the host as *Araujia* sp.
- References: (ds) Blackwelder 1947: 781. (tx) Costa Lima 1956: 288; Schedl 1938a: 179, 1938i: 25, 1939m: 169, 1961b: 187, 1979c: 140.
- argentiniae** (Schedl) 1958f: 44 (*Breviophthorus*). Syntypes, sex?; Argentinien: Misiones, Dep. Concep., Sta. Maria; Schedl Collection in NHMW, Wien and Viana Collection.
- Distribution: South America (Argentina/ Brazil).
- Notes: (1) Schedl 1970e: 93, 1979c: 25 (citation of holotype invalid).
- References: (ds) Schedl 1958f: 44, 1970e: 82, 93; Schonherr & Pedrosa-Macedo 1981: 53. (tx) Schedl 1958f: 44, 1979c: 25.
- attenuatus** Wood 1975a: 30. Holotype ♀; 48 km W Bajia de los Angeles, Baja California Norte, Mexico; Wood Collection.
- Distribution: North America (Baja California Norte in Mexico).
- Hosts: *Pedialanthus macrocarpus*.
- References: (ds) Wood, S. L. 1982b: 951. (tx) McNamara 1984: 752; Wood, S. L. 1975a: 30, 1982b: 951.
- aztecus** (Wood) 1971: 54 (*Sphenoceros*). Holotype ♀; 6 km W Tepic, Nayarit, Mexico; Wood Collection.
- Distribution: North America (Nayarit, San Luis Potosi in Mexico).
- Hosts: Tree limb, in phloem, in transverse, biramous parental tunnels; *Ficus* sp.
- References: (hb) Wood, S. L. 1982b: 962. (ds) Atkinson & Equihua 1985c: 355; Estrada & Atkinson 1988: 211; Wood, S. L. 1982b: 962. (tx) Wood, S. L. 1971: 54, 1982b: 962.
- blanditus** Wood 1974a: 47. Holotype ♀; Fortin de las Flores, Veracruz, Mexico; Wood Collection.
- Distribution: North America (Veracruz in Mexico/ Panama).
- Hosts: Tree branch.
- References: (ds) Wood, S. L. 1982b: 951. (tx) Wood, S. L. 1974a: 47, 1982b: 951.
- bolivianus** (Schedl) 1951m: 110 (*Neodryocoetes*). Holotype, sex?; Bolivien, Cochabamba (German); Schedl Collection in NHMW, Wien.
- Distribution: South America (Bolivia).
- References: (tx) Schedl 1951h: 291, 1951m: 110, 115, 1979c: 43.
- brasiliensis** (Schedl) 1938a: 178 (*Neodryocoetes*). Holotype ♂; Brasilien; Schedl Collection in NHMW, Wien.
- Distribution: South America (Brazil).
- References: (ds) Blackwelder 1947: 781. (tx) Schedl 1938a: 175, 178, 1938i: 26, 1979c: 45.
- carinifrons** (Blandford) 1904: 244 (*Pityophthorus*). Holotype ♂; Motzorongo, Veracruz, Mexico; BMNH, London.

- Distribution: North America (Veracruz in Mexico/ Panama).
 Hosts: Liana, in pith.
 Notes: (1) Bright 1976c: 425 (to *Araptus*).
 References: (**hb**) Wood, S. L. 1982b: 941. (**ds**) Blackwelder 1947: 781; Hagedorn 1910d: 70; Kleine 1913b: 140, 1914b: 351; Wood, S. L. 1982b: 941. (**tx**) Blandford 1904: 244; Bright 1976c: 425; Hagedorn 1910a: 100; Schedl 1940a: 348; Wood, S. L. 1982b: 941.
- chilensis** (Schedl) 1955c: 258 (*Conophthocranulus*). Syntypes, sex?; Chile, Valparaíso, Algarrobo; Kuschel Collection and 2 ♂ in Schedl Collection in NHMW, Wien.
 Distribution: South America (Chile).
 Notes: (1) Schedl 1979: 56 (citation of holotype invalid).
 References: (**ds**) Schedl 1972d: 144. (**tx**) Schedl 1955c: 256, 258, 1972d: 144, 1979c: 56.
- columbianus** (Schedl) 1938a: 178 (*Neopityophthorus*). Holotype, sex?; Colombian; Schedl Collection in NHMW, Wien.
 Distribution: South America (Colombia).
 Hosts: *Theobroma cacao*, in seeds.
 References: (**ds**) Blackwelder 1947: 781. (**tx**) Blackman 1942a: 183–184; Schedl 1938a: 178, 1979c: 60.
- concentralis** (Schedl) 1951m: 107 (*Neodryocoetes*). Holotype, sex?; Mexico; Schedl Collection in NHMW, Wien.
 Distribution: North America (Jalisco, Oaxaca, Puebla in Mexico).
 Hosts: Shrub (? Compositae), in phloem, in radiate parental tunnels.
 References: (**hb**) Wood, S. L. 1982b: 959. (**ds**) Wood, S. L. 1982b: 959. (**tx**) Schedl 1951m: 107, 1979c: 61; Wood, S. L. 1982b: 959.
- conditus** Wood 1974a: 48. Holotype ♀; Puerto Viejo, Heredia, Costa Rica; Wood Collection.
 Distribution: North America (Costa Rica).
 Hosts: Liana phloem, in radiate parental tunnels.
 References: (**hb**) Wood, S. L. 1982b: 953. (**ds**) Wood, S. L. 1982b: 953. (**tx**) Wood, S. L. 1974a: 48, 1982b: 953.
- confinis** (Blandford) 1904: 241 (*Pityophthorus*). Lectotype ♂; Jalapa, Veracruz, Mexico; BMNH, London, designated by Wood 1975b: 39.
 Distribution: North America (Guatemala/ Honduras/ Chiapas, Veracruz in Mexico/ Panama).
 Hosts: Liana.
 References: (**hb**) Wood, S. L. 1982b: 942. (**ds**) Blackwelder 1947: 781; Ferrer 1942; Hagedorn 1910d: 70; Kleine 1913b: 140, 1914b: 351, 362; Wood, S. L. 1982b: 942. (**tx**) Blandford 1904: 241; Bright 1976b: 185; Eggers 1936a: 390; Hagedorn 1910a: 100; Schedl 1938a: 177, 1940a: 346, 1955g: 18; Wood, S. L. 1975b: 391, 1982b: 942.
glabricollis Schedl 1938a: 181. Holotype ♂; [Teopisca, Chiapas] Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1975b: 391.
- References: (**ds**) Blackwelder 1947: 781; Ferrer 1942. (**tx**) Schedl 1938a: 181, 1940a: 346, 1951m: 108, 1979c: 106; Wood, S. L. 1975b: 391.
- confluens** (Schedl) 1964k: 311 (*Ctenyophthorus*). Holotype ♀; Nova Tentonia, Santa Catarina, Brazil; Schedl Collection in NHMW, Wien, automatic.
 Distribution: South America (Brazil).
 References: (**tx**) Schedl 1964k: 311.
- concentralis** Schedl 1963d: 222 (*Ctenyophthorus*). Holotype, sex?; Nova Tentonia, Santa Catarina, Brazil; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1951.
 References: (**tx**) Schedl 1951m: 107, 1963d: 222, 1964m: 311, 1979c: 61.
- consobrinus** Wood 1975b: 394. Holotype ♂; 6 km N Tepic, Nayarit, Mexico, 1000 m; Wood Collection.
 Distribution: North America (Jalisco, Nayarit in Mexico).
 Hosts: *Ficus* sp.
 References: (**cc**) Equihua & Atkinson 1986: 633. (**hb**) Equihua & Atkinson 1986: 633. (**ds**) Atkinson & Equihua 1985c: 358; Equihua & Atkinson 1986: 633; Kirkendall 1984: 242; Wood, S. L. 1982b: 950. (**tx**) Wood, S. L. 1975b: 394, 1982b: 950.
- corpulentus** (Schedl) 1973a: 371 (*Neodryocoetes*). Holotype, sex?; Bolivien, Guayaramerin (Beni); NHMB, Budapest.
 Distribution: South America (Bolivia).
 References: (**tx**) Schedl 1973a: 371.
- costaricensis** (Schedl) 1938a: 180 (*Neodryocoetes*). Holotype ♀; Costa Rica; Schedl Collection in NHMW, Wien.
 Distribution: North America (Costa Rica/ Panama).
 Hosts: *Cecropia* sp. petiole, and tree twigs.
 Notes: (1) Validated as *insularis* var. *costaricensis*. (3) Allied to *laevigatus*, with similar reproduction and sexual dimorphism.
 References: (**ds**) Wood, S. L. 1982b: 945. (**tx**) Schedl 1938a: 180, 1951m: 73, 1979c: 126; Wood, S. L. 1982b: 945.
- crassulus** Wood 1981: 122. Holotype ♀; 8 km NE Cerro Jefe, Panama Prov., Panama, 700 m; CNCI, Ottawa, automatic.
 Distribution: North America (Panama).
 References: (**ds**) Wood, S. L. 1982b: 947. (**tx**) Wood, S. L. 1981: 122, 1982b: 947.
- crassus** Wood 1977b: 211. Holotype ♀?; 8 km NE Cerro Jefe, Panama Prov., Panama; CNCI, Ottawa, preoccupied by Schedl 1966.
 References: (**tx**) Wood, S. L. 1977b: 211.
- crassus** (Schedl) 1966f: 108 (*Thammophthorus*). Holotype, sex?; Surinam, Moenge; Schedl Collection in NHMW, Wien.
 Distribution: South America (Suriname).

References: (tx) Schedl 1966f: 108, 1979c: 68.

cribricollis (Schedl) 1954b: 36 (*Neodryocoetes*).
Syntypes ♂ ♀; Brasilien, Fortaleza R. G.; Schedl Collection in NHMW, Wien, and Planmann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 69 (citation of holotype invalid).

References: (tx) Schedl 1954b: 36, 1963d: 223, 1979c: 69.

sulcatus Schedl 1959m: 552 (*Breviophthorus*).
Holotype ♀; Matto Grosso: Rio Caraguata; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1963d: 223.

References: (tx) Schedl 1959m: 552, 1963d: 223, 1979c: 245.

cribripenis (Schedl) 1976a: 70 (*Neodryocoetes*).
Holotype ♀?; Xingu, Mato Grosso, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 70.

cubensis (Blackman) 1942a: 191 (*Neodryocoetes*).
Holotype ♀; Cayamas, Cuba; USNM, Washington.
Figures: Blackman 1942a: figs. 8-9.

Distribution: Antilles Islands (Cuba).

References: (ds) Blackwelder 1947: 781; Bright 1985c: 175. (tx) Blackman 1942a: 191-192; Bright 1985c: 175; Schedl 1951m: 107.

decorulus Wood 1974d: 278. Holotype ♂; Rincon de Osa, Puntarenas, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).

Hosts: Tree log, in phloem.

References: (hb) Wood, S. L. 1982b: 948. (ds) Wood, S. L. 1982b: 948. (tx) Wood, S. L. 1974d: 278, 1982b: 948.

decorus Wood 1974a: 47. Holotype ♂; Rincon de Osa, Puntarenas, Costa Rica; Wood Collection, preoccupied by Bright 1972.

References: (tx) Wood, S. L. 1974a: 47, 1974d: 278.

decorus (Bright) 1972d: 96 (*Neodryocoetes*). Holotype ♂; Good Hope, Trelawny Parish, Jamaica; CNCI, Ottawa.

Distribution: Antilles Islands (Jamaica).

References: (ds) Bright 1985c: 176. (tx) Bright 1972d: 96, 1985c: 176; McNamara 1977: 196; Wood, S. L. 1974d: 278.

delicatus Wood 1974a: 44. Holotype ♂; 8 km S La Huerta, Jalisco, Mexico; Wood Collection.
Distribution: North America (Jalisco, Nayarit in Mexico).

Hosts: Liana, in pith.

References: (cc) Equihua & Atkinson 1986: 633. (hb) Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 939. (ds) Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 939. (tx) McNamara 1984: 752; Wood, S. L. 1974a: 44, 1982b: 939.

dentifrons Wood 1974a: 45. Holotype ♂; 3 km SE Acatlan, Puebla, Mexico; Wood Collection.

Distribution: North America (Honduras/ Jalisco, Puebla in Mexico/ Florida in USA).

Hosts: Lianas (in pith), *Sarcostemma clausum*.

References: (hb) Atkinson et al. 1986: 35; Burgos & Saucedo 1983: 110; Wood, S. L. 1982b: 942. (ds) Atkinson & Equihua 1985a: 88; Atkinson et al. 1986: 35, 1991: 162; Burgos & Saucedo 1983: 110; Wood, S. L. 1982b: 942. (tx) Wood, S. L. 1974a: 45, 1982b: 942.

mexicanus Schedl 1963c: 161 (*Breviophthorus*).

Holotype ♂; Mexico; Schedl Collection in NHMW, Wien, preoccupied by Eggers 1936.
Synonymy: Wood 1982b: 942.

References: (tx) Schedl 1963c: 161, 1979c: 152; Wood, S. L. 1982b: 942.

deyrollei (Blandford) 1904: 245 (*Pityophthorus*).
Lectotype ♂; Mexico; BMNH, London, designated by Wood 1974d: 278.

Distribution: North America (Guatemala/ Honduras/ "Mexico").

Hosts: In seeds.

References: (hb) Wood, S. L. 1982b: 937. (ds) Blackwelder 1947: 781; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 351; Wood, S. L. 1982b: 937. (tx) Blandford 1904: 245; Hagedorn 1910a: 100; Schedl 1938a: 174, 1940a: 348; Wood, S. L. 1974d: 278, 1982b: 937.

insinuat Wood 1974a: 43. Holotype ♂; Guatemala seeds intercepted at San Pedro, California; Wood Collection. Synonymy: Wood 1974d: 278.

References: (tx) Wood, S. L. 1974a: 43, 1974d: 278.

dubiosus (Schedl) 1964b: 206 (*Thamnopthorus*).
Holotype, sex?; Jacarepagna, Rio de Janeiro, Argentina; Servicio de Defensa Sanitaria Vegetal, Rio de Janeiro.

Distribution: South America (Argentina).

Hosts: *Nerium oleander*.

References: Schedl 1964b: 206, 1979c: 84.

dubius (Schedl) 1966f: 105 (*Neodryocoetes*). Holotype ♀; Brasilien, Nova Teutonia, Sta. Catarina; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).

References: (ds) Schedl 1966f: 85, 105, 1979c: 85.

eggersi (Schedl) 1951m: 115 (*Brachydendrulus*).
Holotype, sex?; Bolivien, Cochabamba (German); Schedl Collection in NHMW, Wien.

Distribution: South America (Bolivia).

Notes: (3) Schedl 1951m: 115 (cited as *Brachydendrulus laevis* Eggers, nomen nudum).

References: (tx) Schedl 1951m: 115, 1952i: 14, 1954b: 34, 1979c: 86.

eggersianus (Schedl) 1955k: 144 (*Pityophthorus*).
Holotype, sex? Guadeloupe: Trois-Rivieres; Eggers Collection, in NHMW, Wien, automatic.
Distribution: Antilles Islands (Guadeloupe).

- References: (tx) Schedl 1958k: 144, 1960i: 106.
denticulatus Eggers 1940a: 129 (*Pityophthorus*).
 Holotype, sex?; Guadeloupe: Trois-Rivieres; Eggers Collection, in NHMW, Wien, preoccupied by Wichmann 1915.
 Notes: (1) Bright 1981c: 158 (to *Araptus*).
 References: (ds) Bright 1985c: 176. (tx) Bright 1981c: 158, 1985c: 176; Eggers 1940a: 129; Schedl 1938a: 162, 1952d: 347, 1957c: 878, 1958k: 144, 1960h: 106, 1979c: 78.
- guadeloupensis* Nunberg 1956d: 208 (*Pityophthorus*). Holotype, sex?; Guadeloupe: Trois-Rivieres; Eggers Collection, in NHMW, Wien, automatic, preoccupied by Schedl 1951.
 References: (tx) Nunberg 1956d: 208, 1963c: 98; Schedl 1960i: 106.
- elongatus* (Schedl) 1961i: 226 (*Thamnophthorus*). Holotype, sex?; Bolivia, Cochabamba, Taguina; Schedl Collection in NHMW, Wien.
 Distribution: South America (Bolivia).
 References: (ds) Schedl 1960a: 77, 1961i: 226, 1979c: 90.
- eruditus* (Schedl) 1938a: 182 (*Neopityophthorus*). Holotype ♀; [El Cora, Tepic], Mexico; Schedl Collection in NHMW, Wien.
 Figures: Blackman 1942a: fig. 15.
 Distribution: North America (Costa Rica/ Guatemala/ Honduras/ Chiapas, Nayarit, Oaxaca in Mexico/ Panama).
 Hosts: Limb and log, in phloem, in radiate parental tunnels.
 Notes: (1) Wood 1975b: 392 (to *Araptus*).
 References: (hb) Wood, S. L. 1982b: 956. (ds) Atkinson & Equihua 1986a: 422; Blackwelder 1947: 781; Ferrer 1942; Wood, S. L. 1982b: 956. (tx) Schedl 1938a: 182, 1940a: 346, 1951m: 109, 1954b: 36, 1979c: 91; Wood, S. L. 1975b: 392, 1982b: 956.
- buscki* Blackman 1942a: 192 (*Neodryocotes*). Holotype ♀; Cabima, Panama; USNM, Washington. Synonymy: Wood 1975b: 392.
 References: (ds) Blackwelder 1947: 781. (tx) Blackman 1942a: 192; Wood, S. L. 1975b: 392.
- exigialis* Wood 1974a: 50. Holotype ♀; Fort Clayton, Canal Zone, Panama, 30 m; Wood Collection.
 Distribution: North America (Panama).
 Hosts: Liana phloem, in radiate parental tunnels.
 References: (hb) Wood, S. L. 1982b: 957. (ds) Wood, S. L. 1982b: 957. (tx) Wood, S. L. 1974a: 50, 1982b: 957.
- facetis* Wood 1974a: 46. Holotype ♀; Rio Tempisque, Guanacaste, Costa Rica; Wood Collection.
 Distribution: North America (Costa Rica).
 Hosts: *Ficus* sp. twigs.
 References: (hb) Wood, S. L. 1982b: 944. (ds) Wood, S. L. 1982b: 944. (tx) Wood, S. L. 1974a: 46, 1982b: 944.
- fossifrons* Wood 1975a: 30. Holotype ♂; Lago Amatitlan, Guatemala; Wood Collection.
- Figures: Burgos & Saucedo 1983: 111.
 Distribution: North America (Guatemala/ Jalisco, Nayarit, Oaxaca in Mexico).
 Hosts: Seed pods of trees, lianas.
 References: (cc) Equihua & Atkinson 1986: 633. (hb) Atkinson et al. 1986: 68; Burgos & Saucedo 1983: 110; Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 937. (ds) Atkinson et al. 1986: 68; Burgos & Saucedo 1983: 110; Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 937. (tx) Burgos & Saucedo 1983: 111; Wood, S. L. 1975a: 30, 1982b: 937.
- foreifrons* (Schedl) 1963c: 161 (*Thamnophthorus*). Holotype ♂; Mexico: Jalisco, Guadalajara; Schedl Collection in NHMW, Wien.
 Distribution: North America (Guatemala/ Jalisco in Mexico).
 Hosts: Liana, in pith.
 Notes: Wood 1975a: 21 (to *Araptus*).
 References: (hb) Wood, S. L. 1982b: 938. (ds) Wood, S. L. 1982b: 938. (tx) Schedl 1963c: 161, 1979c: 99; Wood, S. L. 1974a: 44, 1975a: 21, 1982b: 938.
- interjectus* Wood 1974a: 44. Holotype ♂; Volcan de Agua, Guatemala; Wood Collection. Synonymy: Wood 1975a: 21.
 References: (tx) Wood, S. L. 1974a: 44, 1975a: 21.
- frenatus* (Schedl) 1939d: 411 (*Thamnophthorus*). Syntypes, sex?; Cordoba, San Javier [Argentina]; Schedl Collection in NHMW, Wien, and Bosq Collection.
 Distribution: South America (Argentina).
 Notes: (1) Schedl 1979c: 100 (citation of holotype invalid).
 References: (hb) Viana 1964: 125. (ds) Blackwelder 1947: 781; Viana 1964: 125. (tx) Schedl 1939d: 411, 1964b: 208, 1979c: 100.
- frontalis* Wood 1974a: 52. Holotype ♀; Volcan Zunil, Qnezahltenango, Guatemala, 1000 m; Wood Collection.
 Distribution: North America (Guatemala).
 Hosts: Compositae shrub, in phloem, in radiate parental tunnels.
 References: (tx) Wood, S. L. 1974a: 52, 1982b: 959, 1989: 177.
- frontis* Wood 1989: 177. Holotype ♀; Brasilien, Encruzilhada, Bahia; Schedl Collection in NHMW, Wien, automatic.
 Distribution: South America (Brazil).
 References: (tx) Wood, S. L. 1989: 117.
- frontalis* Schedl 1978c: 302 (*Gnathocranus*). Holotype ♀; Brasilien, Encruzilhada, Bahia; Schedl Collection in NHMW, Wien, preoccupied by Wood 1974.
 References: (ds) Schedl 1973d: 162. (tx) Schedl 1978c: 302; Wood, S. L. 1989: 117.
- frugalis* Wood 1974a: 49. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica; Wood Collection.

- Distribution: North America (Costa Rica/ Panama).
Hosts: Liana.
References: **(hb)** Wood, S. L. 1982b: 954. **(ds)** Wood, S. L. 1982b: 954. **(tx)** Wood, S. L. 1974a: 49, 1982b: 954.
- furvescens** Wood 1974a: 53. Holotype ♀; Volcan Pacaya, Esquinla, Guatemala; Wood Collection.
Distribution: North America (Guatemala).
Hosts: *Roupala* sp., in phloem.
References: **(ds)** Wood, S. L. 1982b: 961. **(tx)** Wood, S. L. 1974a: 53, 1982b: 961.
- furrus** Wood 1974a: 53. Holotype ♀; Cerro Punta (Volcan Chiriqui), Chiriqui, Panama, 1800 m; Wood Collection.
Distribution: North America (Panama).
Hosts: *Roupala* sp. in phloem, in longitudinal, biramous parental tunnels.
References: **(hb)** Wood, S. L. 1982b: 961. **(ds)** Wood, S. L. 1982b: 961. **(tx)** Wood, S. L. 1974a: 53, 1982b: 961.
- genealis** Wood 1974a: 45. Holotype ♂; Volcan de Agua, Guatemala; Wood Collection.
Distribution: North America (Guatemala).
Hosts: Liana, in pith.
References: **(hb)** Burgos & Saucedo 1983: 112; Wood, S. L. 1982b: 939. **(ds)** Burgos & Saucedo 1983: 112; Wood, S. L. 1982b: 939. **(tx)** Wood, S. L. 1974a: 45, 1982b: 939.
- gracileus** Wood 1976a: 364. Holotype ♂; 6 km N Mazatlan, Sinaloa, Mexico; CNCI, Ottawa.
Distribution: North America (Sinaloa in Mexico).
Hosts: Twig in occupied oriole nest.
References: **(ds)** Wood, S. L. 1982b: 942. **(tx)** McNamara 1984: 752; Wood, S. L. 1976a: 364, 1982b: 942.
- gracilentus** (Schedl) 1972g: 61 (*Neodryocoetes*). Holotype, sex?; Brasilien, Pedra Azul, M. Gerais; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: **(tx)** Schedl 1972g: 61, 1979c: 107.
- gracilis** (Schedl) 1966f: 106 (*Neodryocoetes*). Holotype, sex?; Argentinien, Buenos Aires, Isla Maria Garcia; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
References: **(ds)** Schedl 1966f: 95, 106, 1979c: 108.
- grandis** (Schedl) 1954b: 35 (*Neodryocoetes*). Syn-types, sex?; Brasilien, Matto Grosso, Rio Caraguata; Schedl Collection in NHMW, Wien and Plamann Collection.
Distribution: South America (Brazil).
Notes: (1) Schedl 1979c: 109 (citation of holotype invalid).
References: **(tx)** de Ruetten 1970: 104; Schedl 1954b: 35, 1959m: 550, 1963c: 161, 1979c: 109.
- graulipennis** (Schedl) 1967d: 12 (*Neodryocoetes*). Holotype ♂; Brasilien, Nova Tentonia, Santa Catarina; Schedl Collection in NHMW, Wien.
- Distribution: South America (Brazil).
References: **(ds)** Schedl 1976a: 52. **(tx)** McNamara 1977: 196; Schedl 1967d: 12, 1979c: 111.
- guadeloupanus** Wood 1989: 177. Holotype, sex?; Guadeloupe; Schedl Collection in NHMW, Wien, automatic.
Distribution: Antilles Islands (Guadeloupe).
References: **(tx)** Wood, S. L. 1989: 177.
- guadeloupensis** Schedl 1970e: 91 (*Brachyden-drusus*). Holotype, sex?; Guadeloupe; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1951.
References: **(ds)** Bright 1985c: 176. **(tx)** Bright 1985c: 176; Schedl 1970e: 91, 1979c: 113–114; Wood, S. L. 1989: 117.
- hostilis** (Blackman) 1942a: 189 (*Neodryocoetes*). Holotype ♀; *Erythrina cristagalli* seeds from Paris intercepted at Washington, D.C.; USNM, Washington.
Distribution: South America?
Hosts: *Erythrina cristagalli*, in seeds.
References: **(ds)** Blackwelder 1947: 781. **(tx)** Blackman 1942a: 189.
- hymenaeae** (Eggers) 1933b: 9 (*Neodryocoetes*). Holotype ♂; Gourdonville, French Guiana; MNHN, Paris.
Figures: Blackman 1942a: fig. 14.
Distribution: Antilles Islands (Guadeloupe/ Dominican Republic in Hispanola/ Jamaica/ Santa Lucia/ St. Vincent/ Trinidad), North America (Panama), South America (Brazil/ Cayenne/ Colombia/ Guyana/ Suriname/ Venezuela).
Hosts: *Brounea* sp., *Cajanus cajan*, *Ceratonia* sp., *Cicer arietinum*, *Erythrina* sp., *Hymenaea courbaril*, in seeds.
References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956; Kalshoven 1963; Wood, S. L. 1982b: 936. **(ds)** Blackwelder 1947: 781; Bright 1981c: 152, 1982b: 163, 1985c: 176; Costa Lima 1956; Kalshoven 1961: 233, 1963; Numberg 1958a: 480, 1963c: 97; Schedl 1960a: 78, 1966f: 85, 1970e: 82, 1971f: 147, 1976a: 52; Wood, S. L. 1982b: 936. **(tx)** Blackman 1942a: 179; Bright 1985c: 176; Costa Lima 1956; Eggers 1933b: 9, 1936a: 388, 390, 1938a: 177, 1940a: 127–128; Numberg 1958a: 480; Schedl 1938a: 177, 1979c: 120; Wood, S. L. 1973c: 172, 1974b: 392, 1982b: 936, 1986a: 97.
- insularis** Eggers 1940a: 128 (*Neodryocoetes*). Holotype ♀; Guadeloupe; Flentiaux Collection, MNHN, Paris. Synonymy: Wood 1973c: 172.
References: **(hb)** Kalshoven 1963; **(ds)** Blackwelder 1947: 781; Bright 1972d: 91; Kalshoven 1963: 235; Schedl 1973d: 155. **(tx)** Bright 1972d: 91; Eggers 1940a: 128; Schedl 1951m: 73, 1979c: 125; Wood, S. L. 1973c: 172.
- caribaeus** Blackman 1942a: 185 (*Neodryocoetes*). Holotype ♀; Trinidad, West Indies; USNM, Washington. Synonymy: Wood 1973c: 172.
References: **(cn)** Costa Lima 1956. **(hb)** Costa

- Lima 1956; Kalshoven 1963: 235. (**ds**) Blackwelder 1947: 781; Browne 1970: 541; Costa Lima 1956; Kalshoven 1963. (**tx**) Blackman 1942a: 185; Costa Lima 1956; Schedl 1964k: 221; Wood, S. L. 1973c: 172.
- guianae* Blackman 1942a: 186 (*Neodryocotes*). Holotype ♀; British Guiana; USNM, Washington. Synonymy: Wood 1973c: 172. References: (**cn**) Mumford 1960: 33. (**ds**) Blackwelder 1947: 781; Mumford 1960: 33. (**tx**) Blackman 1942a: 186; Schedl 1963f: 57; Wood, S. L. 1973c: 172.
- hoodi* Blackman 1942a: 187 (*Neodryocotes*). Holotype ♀; Taboga Island, Panama; USNM, Washington. Synonymy: Wood 1973c: 172. References: (**ds**) Blackwelder 1947: 781. (**tx**) Blackman 1942a: 187-188; Wood, S. L. 1973c: 172.
- humilis* Blackman 1942a: 188 (*Neodryocotes*). Holotype ♀; Bonito, Pernambuco, Brazil; USNM, Washington. Synonymy: Wood 1975b: 392. References: (**ds**) Blackwelder 1947: 781. (**tx**) Blackman 1942a: 188; Wood, S. L. 1975b: 392.
- imitatrix* (Schedl) 1972g: 62 (*Neodryocotes*). Holotype ♂?; Brasil, Jacareacanga. Para; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (**tx**) Schedl 1972g: 62, 1979c: 121.
- impensus* (Wood) 1961c: 6 (*Thamnophtorus*). Holotype ♂; Bogota, Colombia; Wood Collection. Distribution: South America (Colombia). Hosts: *Podocarpus raspigiliosii*, in fallen seeds. References: (**tx**) Wood, S. L. 1961c: 6.
- incommodus* (Blandford) 1904: 245 (*Pityophthorus*). Holotype ♀; Cerro Zamil, Guatemala, 1300 m; BMNH, London. Distribution: North America (Guatemala). Notes: (3) Bright 1976c: 426 (to *Araptus*). References: (**ds**) Blackwelder 1947: 782; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 363. (**tx**) Blandford 1904: 245; Bright 1976c: 426; Hagedorn 1910a: 101; Schedl 1955g: 18; Wood, S. L. 1982b: 941.
- incompositus* (Blandford) 1904: 243 (*Pityophthorus*). Holotype ♀; Coatepeque, Guatemala, 1300 feet; BMNH, London. Distribution: North America (Guatemala). Notes: (1) Bright 1976c: 426 (to *Araptus*). References: (**ds**) Blackwelder 1947: 782; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 363; Wood, S. L. 1982b: 954. (**tx**) Blandford 1904: 243; Bright 1976c: 426; Hagedorn 1910a: 101; Schedl 1955g: 18; Wood, S. L. 1982b: 954.
- laevigatus* (Eggers) 1933b: 6 (*Pityophthorus*). Holotype ♀; St. Laurent du Maroni, French Guiana; MNHN, Paris. Distribution: Antilles Islands (Guadeloupe), North America (Costa Rica/ Panama), South America (Brazil/ Cayenne/ Colombia/ Suriname). Hosts: *Brownea* sp., *Cynometra hemitomophylla*, *Daphnopsis seibertii*, *Entada gigas*, *Euterpe oleracea*, in seeds. Notes: (3) Wood 1973c: 173 (to *Araptus*). This species apparently reproduces by arrhenotocous parthenogenesis; flightless males are dwarfed. References: (**hb**) Wood, S. L. 1982b: 944. (**ds**) Blackwelder 1947: 781; Bright 1955c: 176; Schedl 1963f: 56; Wood, S. L. 1982b: 944. (**tx**) Bright 1955c: 176; Eggers 1933b: 6; Schedl 1938a: 180, 1963f: 56, 1979c: 135; Wood, S. L. 1973c: 173, 1982b: 944. *insularis* Eggers 1940a: 130 (*Neopityophthorus*). Holotype ♀; Trois Rivières, Guadeloupe; Eggers Collection, in NHMW, Wien, preoccupied by Eggers 1940a: 128. Synonymy: Wood 1973c: 173. References: (**tx**) Blackman 1942a: 195; Eggers 1936a: 390, 1940a: 130; Schedl 1938a: 180, 1951m: 73, 1979c: 125; Wood, S. L. 1973c: 173.
- guadeloupensis* Schedl 1951m: 73 (*Neodryocotes*). Holotype ♀; Trois Rivières, Guadeloupe; Eggers Collection, in NHMW, Wien, automatic. Synonymy: Wood 1973c: 173. References: (**tx**) Schedl 1951m: 73, 109; Wood, S. L. 1973c: 173.
- laevis* (Schedl) 1938a: 181 (*Neopityophthorus*). Syntypes, sex?, Guadeloupe; Eggers Collection (in USNM, Washington?) and 2 ♀ in Schedl Collection in NHMW, Wien. Distribution: Antilles Islands (Guadeloupe). Notes: (1) Schedl 1979c: 135 (citation of holotype invalid). References: (**ds**) Blackwelder 1947: 781; Bright 1955c: 176. (**tx**) Bright 1955c: 176; Eggers 1940a: 108; Schedl 1938a: 181. 1979c: 135.
- laudatus* Wood 1974a: 49. Holotype ♀; San Isidro del General, San Jose, Costa Rica; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree branch, in phloem. References: (**hb**) Wood, S. L. 1982b: 955. (**ds**) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 955. (**tx**) Wood, S. L. 1974a: 54, 1982b: 955.
- lepidus* Wood 1974a: 54. Holotype ♀; Volcan, Puntarenas, Costa Rica, 1000 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree branch, in phloem. References: (**hb**) Wood, S. L. 1982b: 962. (**ds**) Wood, S. L. 1982b: 962. (**tx**) Wood, S. L. 1974a: 54, 1982b: 962.
- leptus* (Bright) 1972c: 1666 (*Neodryocotes*). Holotype ♀; Ejipantla, 8 km S San Andres Tuxtla, Veracruz, Mexico; CNCI, Ottawa. Distribution: North America (Veracruz in Mexico). References: (**ce**) Equihua & Atkinson 1986: 633. (**hb**) Equihua & Atkinson 1986: 633. (**ds**) Equihua & Atkinson 1986: 633; Wood, S. L.

- 1982b: 950. (tx) Bright 1972c: 1666; McNamara 1977: 196; Wood, S. L. 1982b: 950.
- limax (Schedl)** 1939j: 566 (*Sphenoceros*). Syntypes ♀; Brasilien, Sta. Catarina; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
Notes: (3) Schedl 1950i: 164 (cites *Hypertensus reitteri* Hagedorn, nomen nudum, synonymy).
References: (ds) Blackwelder 1947: 784; Schedl 1966f: 91. (tx) Schedl 1939j: 566–567, 1950i: 164, 1979c: 139.
- linearis (Schedl)** 1938a: 174 (*Thamnophthorus*). Holotype, sex?; Bolivien, Cochabamba; Eggers Collection, in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Blackwelder 1947: 781. (tx) Anderson, W. H. & Anderson 1971: 17; Schedl 1938a: 169, 174, 1961i: 226, 1979c: 139.
- macer (Bright)** 1972c: 1666 (*Neodryocoetes*). Holotype ♀; Ejipantla, 8 km S San Andres Tuxtla, Veracruz, Mexico; CNCI, Ottawa.
Figures: Bright 1972c: 1669.
Distribution: North America (Honduras/Veracruz in Mexico).
Hosts: Shrub, in phloem.
References: (ds) Wood, S. L. 1982b: 960. (tx) Bright 1972c: 1666, 1669; McNamara 1977: 197; Wood, S. L. 1973c: 173, 1982b: 960.
- tuberculatus** Bright 1972c: 1665 (*Neodryocoetes*). Holotype ♀; Lago Catemaco, Veracruz, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 173.
References: (tx) Bright 1972c: 1665, 1669; McNamara 1977: 197; Wood, S. L. 1973c: 173.
- medialis** Wood 1974a: 48. Holotype ♀; San Isidro del General, San Jose, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: Tree branch, in phloem.
References: (hb) Wood, S. L. 1982b: 953. (ds) Wood, S. L. 1982b: 953. (tx) Wood, S. L. 1974a: 48, 1982b: 953.
- mendicus** Wood 1974a: 54. Holotype ♀; Cartago, Cartago, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Persea americana*.
References: (ds) Wood, S. L. 1982b: 953. (tx) Wood, S. L. 1974a: 54, 1982b: 953.
- micaceus** Wood 1975b: 395. Holotype ♂; Los Corchos, Nayarit, Mexico; Wood Collection.
Distribution: North America (Nayarit in Mexico).
Hosts: Liana.
References: (ds) Wood, S. L. 1982b: 947. (tx) Wood, S. L. 1975b: 395, 1982b: 947.
- micropilosus** Wood 1982a: 224. Holotype ♀; Rancho Tepetates, Km 35 on Veracruz-Xalapa highway, Veracruz, Mexico; Wood Collection.
Distribution: North America (Veracruz in Mexico).
References: (tx) Wood, S. L. 1982a: 224.
- minutissimus (Schedl)** 1954b: 35 (*Brachyden-drulius*). Syntypes, sex?; Brasilien, Panama, Rondon, 500 m; Schedl Collection in NHMW, Wien, and Planmann Collection.
Distribution: South America (Brazil).
Notes: (1) Schedl 1979c: 155 (citation of holotype invalid).
References: (ds) Schedl 1966f: 85. (tx) de Ruette 1970: 98; Schedl 1954b: 35, 1979c: 155.
- montanus (Bright)** 1972d: 93 (*Neodryocoetes*). Holotype, sex?; Hardwar Gap, 4000', St. Andrew Parish, Jamaica; CNCI, Ottawa.
Figures: Bright 1972d: 59, 75.
Distribution: Antilles Islands (Jamaica).
References: (ds) Bright 1985c: 176. (tx) Bright 1972d: 59, 75, 93, 1985c: 176; McNamara 1977: 197.
- morigerus** Wood 1982a: 224. Holotype ♀; Cerro Punta (labeled Volcan Chiriqui), 5500 ft., Panama; Wood Collection.
Distribution: North America (Panama).
Hosts: Tree branch, in phloem.
References: (tx) Wood, S. L. 1982a: 224.
- mucunae (Blackman)** 1942a: 181 (*Neodryocoetes*). Holotype ♀; Perene River, Peru; USNM, Washington.
Distribution: South America (Peru).
Hosts: *Mucuna* sp., in seeds.
References: (ds) Blackwelder 1947: 781. (tx) Blackman 1942a: 181–182.
- namulus** Wood 1974a: 54. Holotype ♀; Tampico, Tamaulipas, Mexico; Wood Collection.
Distribution: North America (Tamaulipas in Mexico).
Hosts: *Disholcopsis* gall on *Quercus virginiana*.
References: (ds) Wood, S. L. 1982b: 948. (tx) Wood, S. L. 1974a: 54, 1982b: 948.
- niger (Bright)** 1972d: 93 (*Neodryocoetes*). Holotype, sex?; Hardwar Gap, 4000', St. Andrew Parish, Jamaica; CNCI, Ottawa.
Figures: Bright 1972d: 75.
Distribution: Antilles Islands (Dominican Republic in Hispaniola/Jamaica).
References: (ds) Bright 1981c: 152, 1985c: 176. (tx) Bright 1972d: 75, 93, 1985c: 176; McNamara 1977: 197.
- nigrellus** Wood 1974a: 52. Holotype ♀; 10 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Myrica pubescens*, in phloem, in radiate parental tunnels.
References: (hb) Wood, S. L. 1982b: 960. (ds) Wood, S. L. 1982b: 960. (tx) McNamara 1984: 197; Wood, S. L. 1974a: 52, 1982b: 960.

nitidipennis (Schedl) 1963f: 57 (*Neodryocotes*).

Holotype, sex?; Suriname, Maripahuvel. Oe. v. Dam; Schedl Collection in NHMW, Wien.

Distribution: South America (Suriname).

References: (ds) Schedl 1963f: 57, 1973d: 155, 1979c: 169.

nitidulus (Schedl) 1967d: 11 (*Neodryocotes*).

Holotype, sex?; Brasiliens, Parana, Caioba, 10 m, 25 50, 48 40; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Schedl 1970e: 82, (tx) McNamara 1977: 197; Schedl 1967d: 11, 1979c: 169.

novateutonicus (Schedl) 1951m: 116 (*Gnathocranus*).

Syntypes ♀; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien and Plaumann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 172 (citation of holotype invalid).

References: (ds) Schedl 1976a: 53, 1978c: 291; Schonherr & Pedrosa-Macedo 1981: 53, (tx) Schedl 1951h: 284, 1951m: 116, 1979c: 172.

nudus (Schedl) 1938a: 176 (*Thamnophtorus*).

Holotype, sex?; Brasilien, Sao Paulo; Eggers Collection, in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 781; Schedl 1973d: 155, (tx) Schedl 1938a: 176, 1964b: 207, 1979c: 173.

obesus Wood 1977b: 212. Holotype ♀?; 18 km NE

Oriximina, Brazil; CNCI, Ottawa.

Distribution: South America (Brazil).

References: (tx) Wood, S. L. 1977b: 212.

obscurus (Eggers) 1936a: 390 (*Neodryocotes*).

Holotype, sex?; Brasilien, Sta. Catharina, Blumenau; USNM, Washington [2 cotypes in Eggers Collection, in NHMW, Wien (Schedl 1979c: 175)].

Distribution: South America (Brazil/ Paraguay).

References: (ds) Blackwelder 1947: 781; Schedl 1973a: 367, (tx) Anderson, W. H. & Anderson 1971: 24; Eggers 1936a: 390; Schedl 1938a: 177, 1962p: 206, 1979c: 175.

obsoletus (Blandford) 1904: 242 (*Pityophthorus*).

Holotype ♀; Las Mercedes, Guatemala; BMNH, London.

Distribution: North America (Guatemala/ Jalisco in Mexico).

Hosts: Liana, *Ficus* sp., in phloem.

Notes: (3) Bright 1977: 512 (to *Araptus*).

References: (hb) Wood, S. L. 1982b: 946, (ds) Blackwelder 1947: 781; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 363; Wood, S. L. 1982b: 946, (tx) Blackman 1928b: 148; Blandford 1904: 242; Bright 1977: 512; Hagedorn 1910a: 101; Schedl 1938a: 177, 1955g: 18; Wood, S. L. 1982b: 946.

oleanderi (Schedl) 1961b: 187 (*Neodryocotes*).

Holotype ♂; Turkei bei Gilindire; Schedl Collection in NHMW, Wien.

Distribution: Europe (Turkey, obviously introduced, probably from South America).

Hosts: *Nerium oleander*.

References: (tx) Schedl 1961b: 187, 1979c: 178.

pallidus (Blackman) 1942a: 193 (*Neodryocotes*).

Holotype ♀; Cayamas, Cuba; USNM, Washington.

Distribution: Antilles Islands (Cuba/ Dominican Republic in Hispanola/ Puerto Rico), South America (Brazil).

Hosts: *Callophyllum brasiliensis* var. *antellannum*.

References: (ds) Blackwelder 1947: 781; Bright 1981c: 152, 1985c: 176, (tx) Blackman 1942a: 193; Bright 1985c: 176; Wood, S. L. 1977b: 208.

portoricensis Schedl 1951m: 109 (*Neodryocotes*). Holotype, sex?; Portorico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1977b: 208.

References: (ds) Bright 1985c: 174, (tx) Bright 1985c: 174; Schedl 1951m: 109, 1964m: 311, 1979c: 198; Wood, S. L. 1977b: 208.

devius Schedl 1972g: 61 (*Neodryocotes*).

Holotype, sex?; Cuba, Valle Ancon, Pinar del Rio Prov.; Schedl Collection in NHMW, Wien. Synonymy: Wood 1977b: 208.

References: (tx) Schedl 1972g: 61, 1979c: 79; Wood, S. L. 1977b: 208.

paranae (Schedl) 1954b: 34 (*Brachydendrulus*).

Syntypes ♂ ♀; Brasilien, Parana, Rondon, 500 m; Schedl Collection in NHMW, Wien, and Plaumann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 185 (citation of holotype invalid).

References: (tx) de Ruetle 1970: 98; Schedl 1954b: 34, 1979c: 185.

placetulus Wood 1982a: 225. Holotype ♀;

Uruapan, Michoacan, Mexico, 1600 m; Wood Collection.

Distribution: North America (Michoacan in Mexico).

Hosts: *Persea americana*.

References: (ds) Atkinson & Equihua 1985c: 358; Wood, S. L. 1982a: 225.

plaumannii (Schedl) 1976a: 70 (*Brachydendrulus*).

Holotype, sex?; Nova Teutonia, Brasilien; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 70.

politus (Blandford) 1904: 244 (*Pityophthorus*).

Lectotype ♂; Mexico; BMNH, London, designated by Bright 1976b: 184.

Figures: Blackman 1942a: figs. 3–4.

Distribution: Antilles Islands (Cuba/ Haiti in Hispanola/ Jamaica), North America (Costa Rica/ Veracruz in Mexico/ Florida in USA).

Hosts: *Cola* sp., *Mucuna andreana*, *M. fauvecetti*, in seeds.

Notes: (1) Wood 1973c: 173 (to *Araptus*), 1982b: 940 (lectotype designation invalid).

References: (hb) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 940. (ds) Atkinson & Equihua 1986a: 422; Blackwelder 1947: 781; Bright 1985c: 176; Ferrer 1942; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 351; Schedl 1963c: 157; Wood, S. L. 1977a: 69, 1982b: 940. (tx) Blandford 1904: 244; Bright 1976b: 184, 1985c: 176; Hagedorn 1910a: 101; Schedl 1938a: 177, 1940a: 346, 348; Wood, S. L. 1973c: 173, 1982b: 940.

mexicanus Eggers 1936a: 391 (*Neodryocoetes*). Holotype, sex?; Colonia [Veracruz?], Mexico; Eggers Collection, in NHMW, Wien. Synonymy: Schedl 1938a: 177.

References: (tx) Eggers 1936a: 391; Schedl 1938a: 177, 1940a: 346, 1979c: 152.

hubbardi Blackman 1942a: 182 (*Neodryocoetes*). Holotype ♀; Kingston, Jamaica; USNM, Washington. Synonymy: Wood 1973c: 173.

References: (cn) Judd 1970; Mumford 1961: 28. (ds) Blackwelder 1947: 781; Bright 1972d: 94; Judd 1970; Mumford 1961: 28; Straly 1975: 8. (tx) Blackman 1942a: 182–183; Bright 1972d: 94; Wood, S. L. 1973c: 173.

poricollis (Blandford) 1904: 238 (*Pityophthorus*). Holotype ♂; Cerro Zumil, Quezaltenango, Guatemala; BMNH, London.

Distribution: North America (Guatemala).

Notes: (1) Bright 1977: 512 (to *Araptus*).

References: (ds) Blackwelder 1947: 781; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 363; Wood, S. L. 1982b: 958. (tx) Blandford 1904: 238; Bright 1977: 512; Hagedorn 1910a: 101; Schedl 1938a: 180, 1955g: 18; Wood, S. L. 1982b: 958.

pubescens (Schedl) 1951h: 291 (*Neodryocoetes*). Syntypes, sex?; Argentinien, Cordoba, Punilla Dep., Valle Hermoso; Schedl Collection in NHMW, Wien, and Viana Collection.

Distribution: South America (Argentina).

Notes: (1) Schedl 1979: 203 (citation of holotype invalid).

References: (hb) Viana 1964: 125. (ds) Viana 1964: 125. (tx) Schedl 1951h: 291–292, 1979c: 203.

punctatissimus (Schedl) 1938a: 179 (*Neodryocoetes*). Holotype ♂; Bolivien, Cochabamba; Eggers Collection, in NHMW, Wien.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 781; Schedl 1966f: 86. (tx) Schedl 1938a: 179, 1939m: 169, 1951h: 284–285, 1958f: 35, 1979c: 204.

refertus Wood 1974a: 51. Holotype ♀; Volcan Zumil, Quezaltenango, Guatemala; Wood Collection.

Distribution: North America (Guatemala).

Hosts: Shrub phloem, in radiate parental tunnels.

References: (hb) Wood, S. L. 1982b: 957. (ds)

Wood, S. L. 1982b: 957. (tx) Wood, S. L. 1974a: 51, 1982b: 957.

robustus (Schedl) 1964b: 207 (*Thamnophthorus*).

Holotype, sex?; Belem, Para; Servico de Defesa Sanitaria Vegetal, Rio de Janeiro.

Distribution: South America (Brazil).

References: (tx) Schedl 1964b: 207, 1979c: 213.

*rufopalliatu*s Eichhoff 1872a: 136. Holotype ♀; Nova Granada [now Colombia]; IRSNB, Brussels. Distribution: South America (Colombia/Venezuela). Hosts: *Nectandra* sp., in phloem, in transverse, biramous, parental tunnels.

References: (ds) Blackwelder 1947: 758; Gemminger & Harold 1872: 2687; Hagedorn 1910d: 77; Kleine 1913b: 145, 1914b: 342. (tx) Eichhoff 1872a: 136, 1878b: 305; Hagedorn 1910a: 111; Hopkins 1914: 117.

schedli (Blackman) 1942a: 195 (*Neodryocoetes*).

Holotype ♂; Tampico, Mexico; USNM, Washington. Distribution: North America (Tamaulipas, Veracruz in Mexico).

Hosts: Seeds.

References: (ds) Blackwelder 1947: 781; Wood, S. L. 1982b: 946. (tx) Blackman 1942a: 195; Wood, S. L. 1975b: 392, 1982b: 946.

lenis Blackman 1942a: 198 (*Neodryocoetes*).

Holotype ♂; Cordoba, Veracruz, Mexico; USNM, Washington. Synonymy: Wood 1975b: 392.

References: (ds) Blackwelder 1947: 781. (tx) Blackman 1942a: 198; Wood, S. L. 1975b: 392.

schwarzii (Blackman) 1942a: 178 (*Thamnophthorus*). Holotype ♀; Boquete, Panama; USNM, Washington.

Distribution: North America (Costa Rica/Hidalgo, Quintana Roo in Mexico), South America (Ecuador).

Hosts: *Annona cherimola*, *Persea americana*, *P. pittieri*, in seeds.

References: (hb) Atkinson & Equihua 1985a: 88; Blackwelder 1947: 781; Wood, S. L. 1982b: 952. (ds) Atkinson & Equihua 1985a: 88, 1985c: 358, 1988: 85; Wood, S. L. 1982b: 952. (tx) Blackman 1942a: 178; Wood, S. L. 1982b: 952.

sobrinus Wood 1974d: 287. Holotype ♀; Siquatepec, Honduras; Wood Collection.

Distribution: North America (Honduras).

Hosts: *Pinus oocarpa* rust cones.

References: (ds) Wood, S. L. 1982b: 952. (tx) Wood, S. L. 1974d: 287, 1982b: 952.

sparsepunctatus (Schedl) 1938i: 26 (*Neodryocoetes*). Syntypes, sex?; Argentinien, Prov. Buenos Aires, Tigre; 2 ♂, 7 ♀ in Schedl Collection in NHMW, Wien, and others in Viana Collection.

Distribution: South America (Argentina).

Hosts: *Acacia negra*.

Notes: (1) Schedl 1979c: 233 (citation of holotype invalid).

- References: **(hb)** Viana 1964: 125. **(ds)** Blackwelder 1947: 781; Viana 1964: 125. **(tx)** Schedl 1938i: 26, 1939m: 169, 1951h: 284, 1952a: 446, 1958f: 35, 1979c: 233.
- speciosus** Wood 1980b: 357. Holotype ♀; 8 km S La Huerta, Jalisco, Mexico, 500 m; Wood Collection. Distribution: North America (Jalisco in Mexico). Hosts: *Ficus* sp. twigs, in pith. Reference: **(hb)** Atkinson & Equihua 1985c: 359; Wood, S. L. 1982b: 943. **(ds)** Atkinson & Equihua 1985c: 358; Atkinson et al. 1986: 68; Wood, S. L. 1982b: 943. **(tx)** Wood, S. L. 1980b: 357, 1982b: 943.
- splendidulus** (Schedl) 1966f: 107 (*Neodryocoetes*). Holotype ♀?; Brasilien, Corumba, Matto Grosso; Eggers Collection, in NHMW, Wien. Distribution: South America (Brazil). References: **(ds)** Schedl 1966f: 86. **(tx)** Schedl 1966f: 107, 1979c: 235.
- sulcatus** (Numberg) 1964b: 435 (*Neodryocoetes*). Holotype, sex?; Argentina, Salta Prov., Urundel; NHMB, Budapest, preoccupied by Schedl 1959. Figures: Numberg 1964b: 436. Distribution: South America (Argentina). Notes: (1) Although technically this is a secondary homonym, a new name was not given because of the high probability that synonymy exists. References: **(tx)** Numberg 1964b: 435–436.
- tabogae** (Blackman) 1942a: 184 (*Neodryocoetes*). Holotype ♀; Taboga Island, Panama; USNM, Washington. Distribution: North America (Costa Rica/ Honduras/ Veracruz in Mexico/ Panama). Hosts: Liana, in pith. References: **(cc)** Equihua & Atkinson 1986: 633. **(hb)** Equihua & Atkinson 1986: 633; Wood, S. L. 1982b: 940. **(ds)** Blackwelder 1947: 781; Equihua & Atkinson 1986: 633; Estrada & Atkinson 1988: 211; Wood, S. L. 1982b: 940. **(tx)** Blackman 1942a: 184; Wood, S. L. 1973c: 174, 1982b: 940.
- vincalis** Bright 1972c: 1667 (*Neodryocoetes*). Holotype ♀; Ejipantla, 8 km S San Andres Tuxtla, Veracruz, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 174. References: **(tx)** Bright 1972c: 1667; McNamara 1977: 197; Wood, S. L. 1973c: 174.
- tenellus** (Schedl) 1951m: 109 (*Neodryocoetes*). Holotype ♂; [Chiapas], Mexico; Schedl Collection in NHMW, Wien. Distribution: North America (Campeche, Chiapas, Nayarit in Mexico). Hosts: Tree branch phloem. Notes: (1) Wood 1982b: 945 (to *Araptus*). References: **(hb)** Wood, S. L. 1982b: 945. **(ds)** Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 945. **(tx)** Schedl 1951m: 109, 1979c: 251; Wood, S. L. 1975b: 392, 1982b: 945.
- mexicanus** Schedl 1963c: 162 (*Ctenyophthorus*). Holotype ♀; Mexico: Trampaluz, Escarcaga, Campeche; Schedl Collection in NHMW, Wien, preoccupied by Eggers 1936a: 391. Synonymy: Wood 1975b: 392. References: **(tx)** Schedl 1963c: 162, 1964m: 311, 1979c: 152; Wood, S. L. 1975b: 392.
- granulatus** Schedl 1964m: 311 (*Neodryocoetes*). Holotype ♀; Trampaluz, Escarcaga, Campeche, Mexico; Schedl Collection in NHMW, Wien, automatic. Synonymy: Wood 1975b: 392. Notes: (1) Wood 1975b: 392 (to *Araptus*). References: **(tx)** Schedl 1964m: 311; Wood, S. L. 1975b: 392.
- cuspidis** Wood 1974a: 46. Holotype ♀; 8 km E San Blas, Nayarit, Mexico; Wood Collection. Synonymy: Wood 1975b: 392. References: **(tx)** Wood, S. L. 1974a: 46, 1975b: 392.
- tenuis** (Blackman) 1942a: 197 (*Neodryocoetes*). Holotype ♀; Tampico, Mexico; USNM, Washington. Distribution: North America (Honduras/ Veracruz in Mexico). Hosts: Lianas, in pith. References: **(hb)** Wood, S. L. 1982b: 949. **(ds)** Blackwelder 1947: 781; Wood, S. L. 1982b: 949; **(tx)** Blackman 1942a: 197; Wood, S. L. 1982b: 949.
- placatus** Wood 1974a: 46. Holotype ♂; 5 km W Jaltipan, Veracruz, Mexico; Wood Collection. Synonymy: Wood 1982b: 949. References: **(tx)** Wood, S. L. 1974a: 46, 1982b: 949.
- teres** (Blackman) 1942a: 190 (*Neodryocoetes*). Holotype ♀; Paraiso, Canal Zone, Panama; USNM, Washington. Distribution: North America (Panama). References: **(ds)** Blackwelder 1947: 781; Wood, S. L. 1982b: 949. **(tx)** Blackman 1942a: 190; Wood, S. L. 1982b: 949.
- trepidus** Wood 1974a: 51. Holotype ♀; Volcan de Agua, Esquintla, Guatemala, 1000 m; Wood Collection. Distribution: North America (Guatemala). Hosts: *Meispermia* sp., in stem. References: **(hb)** Atkinson & Equihua 1985c: 358; Wood, S. L. 1982b: 958. **(ds)** Atkinson & Equihua 1985c: 358; Wood, S. L. 1982b: 958. **(tx)** Wood, S. L. 1974a: 51, 1982b: 958.
- umbraticus** (Schedl) 1966f: 108 (*Neodryocoetes*). Holotype ♂; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). Notes: (3) Schedl 1967d: 11 (described female). References: **(ds)** Schedl 1966f: 86, 1967d: 2. **(tx)** Schedl 1966f: 108, 1967d: 11, 1979c: 259.
- resculus** Wood 1974a: 50. Holotype ♀; Guapiles, Limon, Costa Rica, 100 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Liana, in phloem. References: **(hb)** Wood, S. L. 1982b: 957. **(ds)**

Wood, S. L. 1982b: 957. (tx) Wood, S. L. 1974a: 50, 1982b: 957.

cinnulus Wood 1974a: 53. Holotype ♀; San Ignacio de Acosta, San Jose, Costa Rica; Wood Collection. Distribution: North America (Costa Rica).

Hosts: *Roupala complicata*, in phloem, in radiate parental tunnels.

References: (hb) Wood, S. L. 1982b: 961. (ds) Wood, S. L. 1982b: 961. (tx) Wood, S. L. 1974a: 53, 1982b: 961.

volastos (Schedl) 1938a: 175 (*Thamnophtorus*). Holotype, sex?, Bolivien. Cochabamba; USNM, Washington.

Distribution: South America (Bolivia).

References: (tx) Anderson, W. H. & Anderson 1971: 36; Blackman 1942a: 178; Schedl 1938a: 165, 169, 175, 1979c: 268; Wood, S. L. 1961c: 6.

xylotrupes (Eichhoff) 1872a: 135 (*Pityophthorus*). Syntypes, sex?, Bahia, Brazil; IRSNB, Brussels.

Distribution: South America (Argentina/ Brazil).

References: (cn) Costa Lima 1956. (hb) Costa Lima 1956. (ds) Blackwelder 1947: 781; Bruch 1914a; Costa Lima 1936, 1956; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 75; Kleine 1913b: 141. (tx) Costa Lima 1956; Eggers 1936a: 390; Eichhoff 1872a: 135, 1875b: 194; Hagedorn 1910a: 101.

Genus *Conophthorus* Hopkins

CONOPHITHORUS HOPKINS 1915a: 430. Type-species: *Pityophthorus coniperda* Schwarz, original designation.

Keys: Hopkins 1915a: 430, Wood 1982b: 982.

Notes: (3) All species of *Conophthorus* breed in cones of *Pinus* spp. in North America; a few species may also breed in twigs when cones are unavailable.

References: (ay) Nobuchi 1969a: 66. (cn) Shenefelt & Benjamin 1955: 68–69. (cc) McCain, Koehler, & Tjosvold 1957. (hb) Flores & Bright 1987: 181; Herdy 1959: 1–2; Herdy & Thomas 1961: 936; Lyons 1956: 600; Ruckes 1963: 43–50; Schwerdtfeger 1957: 494–508; Thomas & Herdy 1961: 406–408; Wood, S. L. 1986a: 97. (ds) Wood, S. L. 1986a: 97. (tx) Blackman 1928b: 6–16, 1942a: 178–179; Bright 1976d: 169–173; Bright & Stark 1973: 97; Chamberlin 1939: 332–337, 1958; Cibrian-Tovar et al. 1986: 38; Dodge 1938: 18; Hopkins 1915a: 430–433; Keen 1955: 47; Sawamoto 1942: 166; Schedl 1935h: 343, 1938a: 158, 163, 165, 167, 1955g: 4; Swaine 1918a: 46, 92; Thomas 1957: 4, 26; Wood, S. L. 1982b: 982–992, 1986a: 97. (ms) Austara 1979.

apachecae Hopkins 1915a: 432. Holotype ♂; Chiricahua Mountains, Arizona [USA]; USNM, Washington.

Distribution: North America (Arizona in USA).

Hosts: *Pinus engelmannii*.

References: (cn) Cibrian-Tovar et al. 1986: 40;

Doane et al. 1936; Keen 1938: 16, 1955a: 47–48, 159–164. (cc) Keen 1938: 16, 1955a: 47–48, 159–164. (hb) Chamberlin 1939: 335; Doane et al. 1936; Keen 1938: 16, 1955a: 48. (ds) Chamberlin 1939: 335; Cibrian-Tovar et al. 1986: 40; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1955a: 47–48, 159–164; Kleine 1934a: 162; Leng 1920: 340; Wood, S. L. 1982b: 989. (tx) Chamberlin 1939: 335; Hopkins 1915a: 432; Keen 1929: 30; Wood, S. L. 1982b: 989.

conicolens Wood 1977b: 212. Holotype ♀; 13 km W Texmelucan, Puebla, Mexico; Wood Collection.

Figures: Atkinson et al. 1986: 33.

Distribution: North America (Hidalgo, Mexico, Puebla in Mexico).

Hosts: *Pinus leiophylla*.

References: (cn) Cibrian-Tovar et al. 1986: 41. (hb) Atkinson et al. 1986: 35; Cibrian-Tovar et al. 1986: 41. (ds) Atkinson & Equihua 1985a: 90, 1985c: 359, 1988: 86; Atkinson et al. 1986: 35; Cibrian-Tovar et al. 1986: 41; Wood, S. L. 1977b: 212, 1982b: 988. (tx) Atkinson et al. 1986: 33; Muskus, A. 1984: 97; Wood, S. L. 1977b: 212, 1982b: 988.

coniperda (Schwarz) 1895a: 144 (*Pityophthorus*). Lectotype ♂; Marquette, Michigan [USA]; USNM, Washington, designated by Wood 1977c: 354.

Distribution: North America (Nova Scotia, Ontario, Quebec in Canada/ Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Pinus strobus*, accidental or non-persistent in other *Pinus* spp.

References: (ay) Godwin & Odell 1965: 213; Golden & Norris 1964b: 747; Hardy 1959; Henson 1961a, 1962; Thomas, J. B. 1967. (bv) Godwin & Odell 1965; Henson 1960a, 1961b, 1966b; Morgan, F. D. & Mailh 1976. (cn) Anonymous 1956f: 108, 1961b, 1961s, 1962h, 1963v, 1965b, 1965m, 1967f, 1977a: 44; Baker, W. L. 1972: 256; Barber et al. 1979; Beal et al. 1952; Bean 1966: 43; Blackman 1950; Borden 1967: 1187; Britton 1933: 393; Brown 1958: 47, 1959: 20–22; Browne 1968b: 185; Buchanan 1964, 1965: 35; DeBarr, Barber, & Maxwell 1982; Doane et al. 1936; Drooz 1985: 365; Ebel et al. 1976: 24; Ehlers, Hoffard, & Jarrett 1975; Felt 1906: 751; Felt & Rankin 1932: 378, 449; Fogal 1979: 10; Gerhold et al. 1966; Godwin & Odell 1965; Graber 1964: 499–500; de Groot 1985; Hall, K. C., Biggs, & MacLeod 1975: 3; Hensen 1966a; Hitchings 1908: 4; Hoekstra, Merkel, & Powers 1961; Hofacker & Loomis 1952: 26, 1953: 28; Hofacker, Loomis, & Tucker 1954: 45, 1957; Hofacker et al. 1989; Johannsen 1910: 40; Kondo & Moody 1957: 50; Lentz 1929: 424–425; Lindquist, O. H. & Syme 1981: 31; Loomis, Hofacker, &

- Tucker 1986; Lyons 1956: 600; MacLeod & Lawrence 1972: 4; Magasi 1957: 63, 1985: 103; Marquis 1959: 31–41; Martineau 1985: 76; Merkel 1976: 364; Moody 1988: 69; Mook et al. 1959: 36; Morgan, F. D. & Mailu 1976; Odell & Goodwin 1964: 1–7; Pierson 1927: 109–110; Rogers, T. J. 1975c: 25; Santamour 1965: 1–8; Sterner & Davidson 1981: 32; Swaine 1918a: 92; Syme & Nystrom 1988: 33; Wilson, L. F. 1977: 51. **(cc)** Anonymous 1956f: 108, 1965: 9; Felt 1906: 751; Gerhold et al. 1966; Godwin & Odell 1965: 213; Henson 1960a, 1961b, 1962: 524, 1964: 77; Marsh 1966b: 541–549, 1979: 157; Matthews 1970; Muesebeck 1942: 94, 1950: 130; Odera 1971; Santamour 1965. **(hb)** Baker, W. L. 1972: 256; Blackman 1950; Bright 1976d: 170; Browne 1968b: 185; Chamberlin 1939: 335; Clittenden 1899a; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Drooz 1985: 365; Durfee 1958; Durfee et al. 1959: 32–33; Ebel et al. 1972: 24; Felt 1906: 751; Felt & Rankin 1932: 378, 449; Fogal 1979: 9; Godwin & Odell 1965: 213; Harrington 1902a: 116, 1902b; Henson 1960a, 1961b, 1962; Lyons 1956: 600; Martineau 1985: 76; Miller 1975: 695; Morgan, F. D. & Mailu 1976; Odell & Goodwin 1964; Pierson 1927: 109–110; Schwarz 1895a: 144; Swaine 1918a: 92; Thomas & Lindquist 1956: 2; Wilson, L. F. 1977: 81. **(ds)** Allison 1977, 1978; Anderson, R. L. & Barry 1980: 44; Anonymous 1926c: 518, 1963y, 1965b, 1967f, 1977v; Beaulne 1956; Blackman 1950; Blatchley & Leng 1916: 634; Bright 1976d: 170; Browne 1968b: 185; Chamberlin 1925: 1, 30, 1939: 335; Clittenden 1899a; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Dodge 1938: 14, 41; Drooz 1985: 365; Felt & Rankin 1932: 378, 449; Godwin & Odell 1965: 213–219; Hagedorn 1910d: 70; Kleine 1913b: 140, 1934a: 162; Knoll 1932: 66; Leonard 1928: 518; Lindquist, O. H. & Syme 1981: 31; Morgan, A. V. & Morgan 1979, 1980; Proctor 1946: 207; Swaine 1909: 134; Syme & Nystrom 1988: 33; Thomas, J. B. 1957: 4; Wood, S. L. 1982b: 987. **(tx)** Benoit 1985: 76; Blackman 1928b: 148; Blatchley & Leng 1916: 634; Chamberlin 1939: 335; Dodge 1938: 14, 41; Ebel et al. 1976: 24; Godwin & Odell 1965; Hagedorn 1910a: 100; Hopkins 1915a: 430–432; Lindquist, O. H. & Syme 1981: 31; Schedl 1938a: 158, 164; Schwarz 1895a: 144; Swaine 1909: 134, 1918a: 92–93; Syme & Nystrom 1988: 33; Thomas, J. B. 1957: 4, 17–26, 1967; Titus, Meikle, & Harrison 1985: 49; Wood, S. L. 1977c: 384, 1982b: 987. **(ms)** Eckstein 1900c; Hoekstra, Merkel, & Powers 1961; Marquis 1961: 23.
- taedae* Hopkins 1915a: 431. Holotype, sex?; Ft. Monroe, Virginia [USA]; USNM, Washington. Synonymy: Wood 1977c: 385. References: **(cn)** Beal et al. 1952; Blackman 1950; Doane et al. 1936. **(hb)** Baker, W. L. 1972: 258; Blackman 1950; Doane et al. 1936. **(ds)** Baker, W. L. 1972: 258; Blackman 1950; Doane et al. 1936; Kleine 1934a: 163; Leng 1920: 340. **(tx)** Chamberlin 1939: 333–334; Hopkins 1915a: 431; Wood, S. L. 1977c: 384.
- clunicus* Hopkins 1915a: 432. Holotype ♀; “13, 122, *Tomicus clunicus* Fitch det. No. 12 Hopk., Collection Fitch” [USA]; USNM, Washington. Synonymy: Wood 1977c: 384. References: **(hb)** Chamberlin 1939: 335. **(ds)** Chamberlin 1939: 335; Leng 1920: 340. **(tx)** Chamberlin 1939: 335; Hopkins 1915a: 432; Wood, S. L. 1977c: 384.
- echinatae* Wood 1978b: 398. Holotype ♀; Winona, Missouri [USA]; Wood Collection. Distribution: North America (Missouri in USA). Hosts: *Pinus echinata*. References: **(cn)** Yates, H. O. 1981. **(ds)** Wood, S. L. 1982b: 986. **(tx)** Wood, S. L. 1978b: 398, 1982b: 986.
- edulis* Hopkins 1915a: 430. Holotype ♀; Las Vegas Hot Springs, New Mexico [USA]; USNM, Washington. Distribution: North America (Chihuahua in Mexico/ Arizona, Colorado, New Mexico, W Texas, Utah in USA). Hosts: *Pinus cembroides*, *P. discolor*, *P. edulis*. References: **(cn)** Doane et al. 1936; Keen 1938: 16, 1958a: 48; Lyons 1956: 600; Ostmark & Wilford 1956: 12–13. **(cc)** Burks 1979: 781; Bushing 1965: 458; Forcella 1980; Keen 1938: 16, 1958a: 48; Muesebeck 1942: 94, 1950: 130. **(hb)** Chamberlin 1939: 334; Doane et al. 1936; Forcella 1980; Keen 1938: 16, 1958a: 48; Little 1943: 249; Lyons 1956: 600. **(ds)** Chamberlin 1925, 1939: 334; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1958: 48; Kleine 1934a: 162; Leng 1920: 340; Wood, S. L. 1982b: 985. **(tx)** Chamberlin 1939: 334; Hopkins 1915a: 430; Keen 1929: 30; Wood, S. L. 1982b: 985, 1989: 171.
- cembroides* Wood 1972d: 74. Holotype ♂; Miller Canyon, Huachuca Mts., Arizona [USA]; Wood Collection. Synonymy: Wood 1989: 171. References: **(cn)** Cibrian-Tovar et al. 1986: 40. **(hb)** Cibrian-Tovar et al. 1986: 40. **(ds)** Atkinson & Equihua 1985c: 359; Cibrian-Tovar et al. 1986: 40; Furniss, R. L. & Carolin 1977: 377; Wood, S. L. 1982b: 985. **(tx)** Wood, S. L. 1972d: 74, 1982b: 985, 1989: 171.
- mexicanus* Wood 1962: 79. Holotype ♂; Necaxa, Puebla, Mexico; Wood Collection. Distribution: North America (Puebla in Mexico). Hosts: *Pinus* sp. References: **(cn)** Cibrian-Tovar et al. 1986: 42. **(hb)** Cibrian-Tovar et al. 1986: 42. **(ds)** Atkinson & Equihua 1985: 86; Cibrian-Tovar et al. 1986: 42; Wood, S. L. 1982b: 988. **(tx)** Wood, S. L. 1962: 79, 1982b: 988.
- michoacanae* Wood 1980b: 354. Holotype ♂; Urnapan, Michoacan, Mexico; Wood Collection.

Distribution: North America (Michoacan in Mexico).

Hosts: *Pinus michoacana*.

References: **(cn)** Cibrian-Tovar et al. 1986: 43. **(hb)** Cibrian-Tovar et al. 1986: 43. **(ds)** Cibrian-Tovar et al. 1986: 43; Wood, S. L. 1982b: 989. **(tx)** Wood, S. L. 1980b: 354, 1982b: 989.

monophyllae Hopkins 1915a: 433. Holotype ♀; Ventura County, California [USA]; USNM, Washington.

Distribution: North America (California, SW Idaho, Nevada, Utah in USA).

Hosts: *Pinus monophylla*.

References: **(ay)** Thomas, J. B. 1967. **(cn)** Cibrian-Tovar et al. 1986: 44; Keen 1938: 16, 1958a: 48. **(cc)** Bushing 1965: 458; Keen 1938: 16, 1958a: 48; Poinar 1975: 151; Poinar & Caylor 1974; Richerson, J. V. & Borden 1972a. **(hb)** Bright & Stark 1973: 98; Chamberlin 1939: 336; Cibrian-Tovar et al. 1986: 44; Keen 1938: 16, 1958a: 48; Ruckes 1963; Wood, S. L. 1982b: 984. **(ds)** Bright & Stark 1973: 98; Chamberlin 1939: 336; Cibrian-Tovar et al. 1986: 44; Furniss, M. M. & Johnson 1987: 376; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1958a: 48, 159–164; Kleine 1934a: 162; Leng 1920: 340; Ruckes 1963: 47; Wood, S. L. 1982b: 984. **(tx)** Chamberlin 1939: 336; Hopkins 1915a: 433; Keen 1929: 30; Thomas, J. B. 1967; Wood, S. L. 1982b: 984.

ponderosae Hopkins 1915a: 431. Holotype ♀; Ashland, Oregon [USA]; USNM, Washington.

Figures: Evans 1982: 8, 20, Furniss & Carolin 1977: 376, Kinzer, Ridgill, & Watts 1972: 12.

Distribution: North America (British Columbia in Canada/ Durango, Mexico, Michoacan in Mexico/ Arizona, California, Idaho, Montana, Nebraska, Nevada, New Mexico, Oregon, Utah, Washington, Wyoming in USA).

Hosts: *Pinus aristata*, *P. contorta* (rare, mostly in twigs), *P. douglasiana*, *P. durangensis*, *P. flexilis*, *P. jeffreyi*, *P. lambertiana*, *P. leiophylla*, *P. montezumae*, *P. monticola*, *P. ponderosa*, *P. strobiformis*, *P. washoensis*.

Notes: (3) Recent studies (not seen) of epicuticular hydrocarbons suggest that *lambertianae* may be distinct from *ponderosae* (SLW).

References: **(ay)** Kinzer & Ridgill 1972; Thomas, J. B. 1967. **(bv)** Kinzer & Reeves 1976; Kinzer, Ridgill, & Reeves 1972; Kinzer, Ridgill, & Watts 1970. **(cn)** Anonymous 1961h, 1965b, 1966i, 1979g, 1980m: 33; Arno & Hoff 1989; Beatty 1980: 8; Blake, Wagner, & Koerber 1986, 1989; Bodenham & Stevens 1981; Chamberlin 1924; Cibrian-Tovar et al. 1986: 44; Dale & Schenk 1978, 1979; Doane et al. 1936; Essig 1926: 519, 1958: 519; Hagle et al. 1987: 95; Hatch 1938: 194; Haverty & Dell 1984; Hopkins 1899b: 8; Jenkins 1984b; Keen 1929: 54, 1938: 16, 1958a: 48; Kinzer, Ridgill, & Watts 1972: 795–798; Koerber

1968: 229; Lyons 1956: 600; Miller 1914: 4, 1915: 1; Ragenovich 1979; Rogers, T. J. 1981; Schmid, Mitchell, & Mata 1986; Schmidt et al. 1984; Shea, P. J. 1984; Shea, P. J., Jenkins, & Haverty 1984; Stevens, Brewer, & Leatherman 1980: 27, 1982: 27; Swaine 1918a: 92. **(cc)** Bushing 1965: 458; Jenkins 1983, 1984b; Keen 1938: 16, 1958a: 48, 159–164; Kinzer, Ridgill, & Watts 1972: 795–798; Marsh 1979: 270; Muesebeck 1942: 94, 1950: 130, 1957: 54; Ruckes 1956: 184–185; Shea, P. J., Jenkins, & Haverty 1984. **(hb)** Atkinson et al. 1986: 36; Bright & Stark 1973: 98; Chamberlin 1939: 334, 1958: 141–142; Cibrian-Tovar et al. 1986: 44; Dale & Schenk 1979; Doane et al. 1936; Essig 1926: 519, 1958: 519; Hopkins 1899b: 8; Jenkins 1983, 1984a, 1984b; Keen 1929: 54, 1938: 16, 1958a: 48; Kinzer & Reeves 1976; Kinzer, Ridgill, & Watts 1970, 1972; Lyons 1956: 600; Miller 1915: 1; Ruckes 1963: 47; Shea, P. J., Jenkins, & Haverty 1984; Stevens, Brewer, & Leatherman 1980: 27, 1982: 27; Swaine 1918a: 92; Wood, S. L. 1982b: 990. **(ds)** Anonymous 1965b, 1966f, 1979g, 1980m: 33; Atkinson & Equihua 1985a: 90; Atkinson et al. 1986: 36; Bright 1976d: 172; Bright & Stark 1973: 98; Chamberlin 1917, 1925, 1939: 334, 1958: 141–142; Cibrian-Tovar et al. 1986: 44; Essig 1926: 519, 1958: 519; Furniss, R. L. & Carolin 1977: 377; Gast et al. 1989: 385; Hopping 1922; Keen 1929: 50, 54, 1938: 16, 1958a: 48; Kinzer & Reeves 1976; Kleine 1934a: 162; Leng 1920: 340; Ruckes 1963: 47; Schedl 1963c: 157; Thomas, J. B. 1966; Wood, S. L. 1982b: 990. **(tx)** Benoit 1986: 26; Bodenham & Stevens 1981; Chamberlin 1939: 334, 1958: 141–142; Hopkins 1915a: 431; Keen 1929: 30; Kinzer, Ridgill, & Watts 1972: 12; Swaine 1918a: 92; Thomas, J. B. 1967; Wood, S. L. 1962: 79, 1977c: 385, 1982b: 990. **(ms)** Hatch 1938: 194.

scopulorum Hopkins 1915a: 431. Holotype ♀; Flagstaff, Arizona [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.

References: **(cn)** Anonymous 1967t; Doane et al. 1936; Keen 1938: 16, 1958a: 48; Lucht 1966: 30; Pearson 1950: 155; Swaine 1918a: 92. **(cc)** Keen 1938: 16, 1958a: 48, 159–164. **(hb)** Chamberlin 1939: 335; Doane et al. 1936; Keen 1938: 16, 1958a: 48; Swaine 1918a: 92. **(ds)** Anonymous 1967t; Chamberlin 1939: 335; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1958a: 48; Kleine 1934a: 163; Leng 1920: 340. **(tx)** Chamberlin 1939: 335; Hopkins 1915a: 431; Keen 1929: 30; Swaine 1918a: 92; Wood, S. L. 1977c: 385.

contortae Hopkins 1915a: 432. Holotype ♀; Newport, Oregon [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.

References: **(cn)** Clapp 1942: 31; Doane et al. 1936; Keen 1938: 16, 1958a: 48; Lindgren 1980a: 93; Swaine 1918a: 92. **(cc)** Keen 1938:

- 16, 1958a: 48. **(hb)** Bright 1976d: 171; Bright & Stark 1973: 100; Chamberlin 1939: 336, 1958: 141–142; Doane et al. 1936; Keen 1938: 16, 1958a: 48; Lindgren 1980a: 93; Ruckes 1959: 94, 1963; Swaine 1912a: 92. **(ds)** Bright 1976d: 171; Bright & Stark 1973: 100; Chamberlin 1939: 336, 1958: 141–142; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1958a: 48; Kleine 1934a: 162; Leng 1920: 340; Ruckes 1963. **(tx)** Chamberlin 1939: 336, 1958: 141–142; Hopkins 1915a: 432; Keen 1938: 47–48; Swaine 1918a: 92; Wood, S. L. 1977c: 385.
- monticolae* Hopkins 1915a: 432. Holotype ♀; Priest River, Idaho [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.
- References: **(cn)** Andrews 1968: 154; Barnes, Brigham, & Schenk 1962: 1–7; Browne 1968b: 185; Dewey & Clinton 1980; Doane et al. 1936; Evans, D. 1982: 8; Graham 1963; Hatch 1938: 194; Haverty & Wood 1981; Hedlin 1974: 34; Keen 1938: 16, 1958a: 48; Morris, E. V. & Monts 1973a: 15; Robinson, L. A. & Dooling 1978: 11; Ruppel 1967: 27; Schenk & Goyer 1967: 186–187; Silver & Ross 1957: 85; Swaine 1918a: 93. **(cc)** Keen 1938: 16, 1958a: 48; Schenk & Goyer 1967: 186–187; Williamson, D. L., Schenk, & Barr 1966. **(hb)** Barnes, Brigham, & Schenk 1962: 1–7; Bright 1976d: 172; Bright & Stark 1973: 100; Browne 1968b: 185; Chamberlin 1939: 337, 1958: 141–142; Doane et al. 1936; Evans, D. 1982: 8; Graham 1963; Hedlin 1982: 34; Keen 1938: 16, 1958a: 48, 159–164; Ruckes 1959: 94, 1963: 49; Swaine 1918a: 93; Williamson, D. L., Schenk, & Barr 1966. **(ds)** Bright & Stark 1973: 100; Browne 1968b: 185; Chamberlin 1925, 1939: 336, 1958: 141–142; Evans, D. 1983: 31; Furniss, R. L. & Carolin 1977: 377; Hopping 1922; Keen 1929: 30, 1938: 16, 1958a: 48, 159–164; Kleine 1934a: 162; Leng 1920: 340; Patterson & Hatch 1945: 152; Ross 1958: 30; Ruckes 1963: 49; Ruppel 1967: 27. **(tx)** Bright 1976d: 172; Chamberlin 1939: 336, 1958: 141–142; Evans, D. 1982: 8, 20, 1983: 31; Hopkins 1915a: 432; Keen 1929: 30; Schedl 1938a: 161; Swaine 1918a: 92–93; Wood, S. L. 1977c: 385.
- flexilis* Hopkins 1915a: 433. Holotype ♀; Mount Manitou, Colorado [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.
- References: **(ay)** Thomas, J. B. 1967. **(bv)** Kinzer & Reeves 1976. **(cn)** Johnson, D. W. & Minnemeyer 1976; Keen 1938: 16, 1958a: 48. **(cc)** Burks 1979: 781; Bushing 1965: 458; Keen 1938: 16, 1958a: 48. **(hb)** Bright & Stark 1973: 100; Chamberlin 1939: 337, 1958: 142; Keen 1938: 16, 1958a: 48. **(ds)** Atkinson & Equihua 1988: 86; Bright 1964: 170, 1976d: 171; Bright & Stark 1973: 100; Chamberlin 1939: 337, 1958: 142; Furniss, R. L. & Carolin 1977: 377; Keen 1929: 30, 1938: 16, 1958a: 48; Kinzer & Reeves 1976; Kleine 1934a: 162; Leng 1920: 340; McComb et al. 1953: 1; Wood, S. L. 1948: 46, 1951a: 128. **(tx)** Chamberlin 1939: 337, 1958: 142; Hopkins 1915a: 433; Keen 1929: 30; Thomas, J. B. 1967; Wood, S. L. 1951a: 128, 1977c: 385.
- lambertianae* Hopkins 1915a: 433. Holotype ♀; Hilt, California [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.
- References: **(ay)** Thomas, J. B. 1967. **(cn)** Anonymous 1955j: 3, 1958a: 10; Chamberlin 1924; Doane et al. 1936; Essig 1926: 519, 1958: 519; Fowells & Schubert 1956: 27; Hall 1955; Keen 1929: 54, 1958a: 48–55; Lyons 1956: 600; Struble 1947b: 48–50. **(cc)** Anonymous 1960i, 1960j: 12, 1961h, 1961i, 1961s, 1962i, 1963y, 1965b, 1966i, 1979g, 1980a, 1980m: 33; Bedard, W. L. Jr. 1968b; Hoekstra, Plerker, & Powers 1961; Keen 1958a: 48; Muesebeck 1942: 94, 1950: 130; Ruckes 1956: 184–185; Thatcher, T. O. 1961. **(hb)** Bedard 1966b: 152–157, 1968a: 7–17, 1968b: 1–6; Bright & Stark 1973: 99; Chamberlin 1939: 337, 1958: 142; Doane et al. 1936; Essig 1926: 519, 1958: 519; Keen 1929: 54–55, 1958a: 48; Lyons 1956: 600; Miller 1915: 1–12; Ruckes 1957: 367, 1958: 214, 1963: 48; Struble 1947b: 48; Thatcher, T. O. 1961. **(ds)** Anonymous 1958a: 10, 1962i, 1963y, 1965b, 1979g, 1980a, 1980m: 33; Atkinson & Equihua 1988: 86; Bedard 1966b, 1968b; Bright & Stark 1973: 99; Chamberlin 1925, 1939: 337, 1958: 142; Essig 1926: 519, 1958: 519; Furniss, R. L. & Carolin 1977: 377; Hall & Eaton 1960: 8; Hopping 1922; Keen 1929: 30, 54, 1958a: 48; Kleine 1934a: 162; Leng 1920: 340; Ruckes 1963; Thatcher, T. O. 1961. **(tx)** Anonymous 1960j: 12; Blackman 1928b; Chamberlin 1939: 337, 1958: 142; Furniss, R. L. & Carolin 1977: 376; Hopkins 1915a: 433; Keen 1929: 30; Thomas, J. B. 1967; Wood, S. L. 1977c: 385. **(ms)** Hoekstra, Merkel, & Powers 1961.
- radiatae* Hopkins 1915a: 432. Holotype ♀; Pacific Grove, California [USA]; USNM, Washington. Figures: Schaefer 1962: 570.
- Distribution: North America (California in USA). Hosts: *Pinus radiata*.
- References: **(ay)** Schaefer, C. H. 1962, 1964; Thomas, J. B. 1967. **(bv)** Schaefer, C. H. 1962. **(cn)** Keen 1929: 55, 1938: 16, 1958a: 48. **(cc)** Bushing 1965: 458; Keen 1938: 16; Muesebeck 1942: 94, 1950: 130; Odera 1971; Ruhm 1964; Schaefer, C. H. 1962, 1963, 1964. **(hb)** Bright & Stark 1973: 101; Chamberlin 1939: 336; Godwin & Odell 1965: 213–219; Keen 1929: 55, 1938: 16, 1958a: 48; Ruckes 1955: 214, 1963: 49; Schaefer, C. H. 1962, 1964; Wood, S. L. 1982b: 986. **(ds)** Atkinson & Equihua 1988: 86; Bedard 1966b;

- 156; Berryman & Stark 1962a; Bright & Stark 1973: 101; Chamberlin 1925, 1939: 336; Furniss, R. L. & Carolin 1977: 377; Hopping 1922; Keen 1929: 30, 55, 1938: 16, 1958a: 48, 159–164; Kleine 1934a: 163; Leng 1920: 340; McPherson, Wilson, & Stehr 1970a: 1008; Ruckes 1963: 49; Schaefer, C. H. 1962: 569, 1963: 79, 1964: 195; Schedl 1963c: 157; Wood, S. L. 1982b: 986. **(tx)** Chamberlin 1939: 336; Hopkins 1915a: 432; Keen 1929: 30; Schaefer 1962: 570; Thomas, J. B. 1967; Wood, S. L. 1982b: 986.
- resinosae** Hopkins 1915a: 431. Holotype ♂; not given, collected by Harrington [USA]; USNM, Washington.
- Figures: Bright 1976d: 204, 213 (adults), MacLeod, Hook, & Livesay 1975: 2.
- Distribution: North America (Nova Scotia, Ontario, Quebec in Canada/ Michigan, Minnesota, New Hampshire, New Jersey, New York, West Virginia, Wisconsin in USA).
- Hosts: *Pinus resinosa*, uncommon in *P. banksiana* twigs, rare and non-persistent in other *Pinus* spp.
- References: **(ay)** Herdy 1959; McPherson, J. E., Wilson, & Stehr 1970a; Thomas, J. B. 1957: 4, 1967, 1971b. **(bv)** Mattson, Tabashnik, & Miller 1984; McPherson, J. E. 1969b; McPherson, J. E., Wilson, & Stehr 1970b; Tabashnik, Mattson, & Miller 1985. **(cn)** Anonymous 1956f: 108, 1958a: 9, 1964h; Applejohn & Howse 1982b; Baker, W. L. 1972: 257; Berryman 1987; Blackman 1950; Bowser & Weir 1975: 12; Browne 1968b: 185; Doane et al. 1936; Drooz 1985: 365; Fogal 1979: 9; Forbes, Underwood, & Van Sickle 1969: 25; Foster & Applejohn 1975: 7; Hall, K. C., Biggs, & MacLeod 1975: 3; Hall, K. C., Livesay, & Houser 1972; Hard 1964; Howse et al. 1982: 47; Kondo & Moody 1987: 97; Kondo & Taylor 1985: 55; Lindquist, O. H. & Syme 1981: 31; Lyons 1951: 1, 1956: 600, 1957a: 70–79, 1957b: 150, 1957c: 265; MacLeod & Lawrence 1972: 4; Mattson 1968, 1971, 1972a, 1972b, 1978, 1980; Miller, W. E. 1978, 1979; Moody 1988: 66; Prebble 1954: 221; Rauf, Benjamin, & Cecich 1985; Sippell, MacDonald, & Rose 1960: 59; Sterner & Davidson 1981, 1983; Swaine 1918a: 92–93; Syme & Nystrom 1988: 33; Vandenburg 1966: 33; Weir, H. J. et al. 1984a: 105; Weir & Lawrence 1975: 10; Wilson, L. F. 1977: 81. **(ce)** Anonymous 1956f: 108; Berryman 1987; Mattson 1968: 1–2, 1972b, 1980, 1986; Miller, W. E. 1978; Muesebeck 1942: 94; Thomas & Lindquist 1956: 2. **(hb)** Baker, W. L. 1972: 257; Blackman 1950; Bright 1976d: 173; Browne 1968b: 185; Doane et al. 1936; Drooz 1985: 365; Fogal 1979: 9; Godwin & Odell 1965: 213–219; Hard 1964; Lyons 1951: 1, 1956: 599–600, 1957a: 150, 1957b: 264–265; Mattson 1980; Mattson, Tabashnik, & Miller 1984; McPherson, J. E. 1969b: 241–242; McPherson, J. E., Wilson, & Stehr 1970a: 1008–1022; Miller, W. E. 1978; Ruckes 1958: 215; Swaine 1918a: 92, 93; Wilson, L. F. 1977: 81; Wood, S. L. 1982b: 988. **(ds)** Anonymous 1926c: 518, 1958a: 9; Beaulne 1956; Blackman 1950; Bright 1976d: 173; Browne 1968b: 185; Deyrup 1981b: 8; Dodge 1938: 14, 41; Drooz 1985: 365; Kleine 1934a: 163; Leng 1920: 340; Leonard 1928: 518; Lindquist, O. H. & Syme 1981: 31; McPherson, J. E., Wilson, & Stehr 1970a; Miller, W. E. 1978; Syme & Nystrom 1988: 33; Wood, S. L. 1982b: 988. **(tx)** Benoit 1985: 76; Bright 1976d: 204, 213; Chamberlin 1939: 334; Dodge 1938: 14, 41; Hopkins 1915a: 431; Lindquist, O. H. & Syme 1981: 31; McPherson, J. E. 1969b; Swaine 1918a: 92–93; Syme & Nystrom 1988: 33; Thomas, J. B. 1957: 4, 22, 1967, 1971b; Titus, Meikle, & Harrison 1985: 49; Wood, S. L. 1977c: 385, 1982b: 988, 1989: 171. **(ms)** Mattson, Tabashnik, & Miller 1984.
- virginianae** Hopkins 1915a: 431. Holotype ♂; Huttonsville, West Virginia [USA]; USNM, Washington. Synonymy: Wood 1977c: 385.
- References: **(cn)** Beal et al. 1952; Blackman 1950; Doane et al. 1936. **(hb)** Baker, W. L. 1972: 258; Blackman 1950; Chamberlin 1939: 334; Doane et al. 1936. **(ds)** Blackman 1950; Chamberlin 1939: 334; Kleine 1934a: 163; Leng 1920: 340. **(tx)** Chamberlin 1939: 334; Hopkins 1915a: 431; Wood, S. L. 1977c: 385.
- banksianae** McPherson 1970: 1020. Holotype ♂; Fife Lake, Michigan [USA]; Michigan State University Collection, East Lansing. Synonymy: Wood 1989: 171.
- References: **(ay)** McPherson, Wilson, & Stehr 1970a. **(bv)** Anonymous 1979a; Hall, D. J. 1973b; McPherson, J. E. 1969b. **(cn)** Anonymous 1979a, 1989; Drooz 1985: 366; Howse, Gross, & Rose 1981: 61; Howse et al. 1981: 43, 1982: 42, 1983; Kondo & Moody 1987: 91; Kondo & Taylor 1985: 50; Lindquist, O. H. & Syme 1981; MacLeod, Hook, & Livesay 1975; Sippell, Rose, & Gross 1975: 65; Sterner & Davidson 1981, 1983; Syme & Nystrom 1988: 33; Thompson, M. J. & Houser 1975: 2. **(ce)** Hall, D. J. 1975; Hall, D. J. & Wilson 1975. **(hb)** Anonymous 1979a; Bright 1976d: 170; Drooz 1985: 366; Hall, D. J. 1973b; Hall, D. J. & Wilson 1974, 1975; McPherson 1969b; McPherson, Wilson, & Stehr 1970a; Wilson, L. F. 1977: 72; Wood, S. L. 1982b: 987. **(ds)** Bright 1976d: 170; Drooz 1985: 366; Lindquist, O. H. & Syme 1981; McPherson, Wilson, & Stehr 1970a; Syme & Nystrom 1988: 33; Wood, S. L. 1982b: 987. **(tx)** Benoit 1985: 76; Lindquist, O. H. & Syme 1981; MacLeod, Hook, & Livesay 1975: 2; McNamara 1977: 195; McPherson 1969b: 241, 1970: 1020; Syme & Nystrom 1988: 33; Titus, Meikle, & Harrison 1985: 49; Wood, S. L. 1982b: 987, 1989: 171.

teocotum Wood 1950b: 354. Holotype ♂; Uruapan, Michoacan, Mexico; Wood Collection.

Distribution: North America (Michoacan in Mexico).

Hosts: *Pinus teocote*.

References: (cn) Cibrian-Tovar et al. 1986: 46. (hb) Cibrian-Tovar et al. 1986: 46. (ds) Cibrian-Tovar et al. 1986: 46; Wood, S. L. 1982b: 991. (tx) Wood, S. L. 1980b: 354, 1982b: 991.

terminalis Flores & Bright 1987: 181. Holotype ♀; Mexico: Nuevo Leon. La Primavera, Galeana, 1700 m; CNCI, Ottawa.

Distribution: North America (Nuevo Leon in Mexico).

Hosts: *Pinus cembroides*.

References: (tx) Flores & Bright 1987: 181.

Genus *Pityoborus* Blackman

PITYOBORUS BLACKMAN 1922b: 96. Type-species: *Crypturgus comatus* Zimmermann, monobasic.

Keys: Wood 1958a: 47, 1982b: 1147.

Notes: Members of this genus breed in the shaded-out, dying branches of living *Pinus* spp.

References: (hb) Wood, S. L. 1986a: 97. (ds) Wood, S. L. 1986a: 97. (tx) Beal & Massey 1945: 60, 124; Blackman 1922b: 96, 1928b: 3–16, 1942a: 177; Schedl 1938a: 159, 163, 165, 168; Wood, S. L. 1958a: 46–56, 1982b: 1146–1152, 1986a: 97.

comatus (Zimmermann) 1865: 143 (*Crypturgus*). Holotype ♀; South Carolina [USA]; MCZ, Cambridge.

Figures: Blackman 1922b: pl. 10, figs. 51–52 (adult).

Distribution: Antilles Islands (Andros Island in Bahamas Islands), North America (Alabama, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina in USA).

Hosts: *Pinus* spp.

Notes: (3) Blackman 1928b: 145 (redescribed).

References: (ay) Furniss, M. M. et al. 1987. (bv) Atkinson, Foltz, & Connor 1988; Turnbow & Franklin 1980. (cn) Baker, W. L. 1972: 258; Doane et al. 1936; Schuder 1969: 79; Swaine 1918a: 104. (cc) Furniss, M. M. et al. 1987. (hb) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 258; Beal & Massey 1945: 125; Blackman 1922b: 96–98; Chamberlin 1939: 343; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Drooz 1985: 364; Swaine 1918a: 104. (ds) Atkinson et al. 1991: 162; Baker, W. L. 1972: 258; Beal & Massey 1945: 125; Blackman 1922b: 96–98; Blatchley & Leng 1916: 632; Chamberlin 1925, 1939: 343; Deyrup & Atkinson 1987a: 66; Drooz 1985: 364; Hagedorn 1910d: 70; Henshaw 1885: 148; Kirk 1970; Kleine 1913b: 140, 1934a: 163; Leng 1920: 341; Ostmark 1968; Schuder 1969: 79; Swaine 1909: 134; Turnbow & Franklin 1980; Wood, S. L. 1982b: 1148. (tx) Beal & Massey 1945: 125;

Blackman 1922b: 96–98, 1928b: 145–146, 1942a: 203; Blatchley & Leng 1916: 632; Chamberlin 1939: 343; Eichlhoﬀ 1878b: 136; Hagedorn 1910a: 100; LeConte 1868: 157, 1876: 354–355; Schedl 1938a: 159; Swaine 1909: 134, 1918a: 104; Wood, S. L. 1958a: 48, 53, 1982b: 1145; Zimmerman 1865: 143.

seriatus LeConte 1878a: 432 (*Pityophthorus*).

Holotype ♂; Tampa, Florida [USA]; MCZ, Cambridge. Synonymy: Blackman 1928b: 146.

References: (cn) Swaine 1918a: 104. (hb) Swaine 1918a: 104. (ds) Hagedorn 1910d: 70; Henshaw 1882: 269, 1885: 148; Kleine 1913b: 140; Leng 1920: 341; Schwarz 1878d: 468. (tx) Blackman 1928b: 145–146; Chamberlin 1939: 342; Hagedorn 1910a: 101; LeConte 1878a: 432–433, 468; Swaine 1909: 140, 1918a: 104.

frontalis Wood 1971: 49. Holotype ♀; 13 km SE El Cameron, Oaxaca, Mexico; Wood Collection.

Figures: Bright 1972c: 1675.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Pinus oocarpa*, *P. sp.*

References: (ds) Wood, S. L. 1982b: 1151. (tx) Wood, S. L. 1971: 49, 1973c: 182, 1982b: 1151.

severus Bright 1972c: 1676. Holotype ♀; 5 km N Suchixtepec, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 182.

References: (tx) Bright 1972c: 1675–1676; McNamara 1977: 197; Wood, S. L. 1973c: 182, 1982b: 1151.

hirtellus Wood 1958a: 50. Holotype ♀; 23 km NW Guadalajara, Jalisco, Mexico; SMUK, Lawrence. Distribution: North America (Durango, Jalisco in Mexico).

Hosts: *Pinus pseudostrobus*, *P. sp.*

References: (hb) Atkinson & Equihua 1985a: 98; Atkinson et al. 1986: 36. (ds) Atkinson & Equihua 1985a: 98; Atkinson et al. 1986: 36; Wood, S. L. 1985a: 98. (tx) Wood, S. L. 1958a: 47, 50, 1982b: 1150.

hondurensis Wood 1971: 49. Holotype ♀; Yuncaran, Paraiso, Honduras; Wood Collection. Distribution: North America (Belize/ Honduras).

Hosts: *Pinus caribaea*, *P. oocarpa*.

References: (ds) Wood, S. L. 1982b: 1151. (tx) Wood, S. L. 1971: 49, 1982b: 1151.

rubentis Wood 1958a: 51. Holotype ♀; 23 km W Texmelucan, Puebla, Mexico; SMUK, Lawrence. Distribution: North America (Chihuahua, Durango, Michoacan, Puebla, Veracruz in Mexico).

Hosts: *Pinus* spp.

References: (hb) Wood, S. L. 1982b: 1150. (ds) Atkinson & Equihua 1985: 94; Wood, S. L. 1982b: 1150. (tx) McNamara 1977: 197; Wood, S. L. 1958a: 48, 51, 1982b: 1150.

secundus Blackman 1928b: 146. Holotype ♀; La Sal Mts., Utah [USA]; USNM, Washington. Figures: Bright 1972c: 1675.

Distribution: North America (Distrito Federal, Durango, Hidalgo, Michoacan, Puebla, Veracruz in Mexico/ Arizona, New Mexico, Utah in USA).

Hosts: *Pinus ayacahuite*, *P. leiophylla*, *P. oocarpa*, *P. ponderosa*, *P. pseudostrobus*, *P.* spp.

References: (cc) Furniss, R. L. & Carolin 1977: 399. (hb) Atkinson et al. 1986: 36; Chamberlin 1939: 345; Furniss, R. L. & Carolin 1977: 399. (ds) Atkinson & Equihua 1988: 94; Atkinson et al. 1986: 36; Chamberlin 1939: 345; Furniss, R. L. & Carolin 1977: 399; Leng & Mutchler 1933: 52; Wood, S. L. 1960b: 69, 1982b: 1148. (tx) Blackman 1928b: 146, 1942a: 203; Chamberlin 1939: 345; Wood, S. L. 1958a: 48, 53–54, 1960b: 69, 1973c: 182, 1982b: 1148.

tertius Blackman 1942a: 202. Holotype ♀; Chalco, D.F., Mexico; USNM, Washington. Synonymy: Wood 1973c: 182.

References: (ds) Blackwelder 1947: 781; Bright 1972c: 1679. (tx) Blackman 1942a: 202–203; Wood, S. L. 1958a: 48, 58, 1973c: 182.

intonsus Wood 1958a: 54. Holotype ♀; 23 km W Texmelucan, Puebla, Mexico; SMUK, Lawrence. Synonymy: Wood 1973c: 182.

References: (ay) Thomas, J. B. 1967. (ds) Thomas, J. B. 1967. (tx) Thomas, J. B. 1967; Wood, S. L. 1958a: 48, 54, 1973c: 182, 1982b: 1148.

immitus Bright 1972c: 1674. Holotype ♀; 68 km W Durango, Durango, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 182.

References: (tx) Bright 1972c: 1674–1675; McNamara 1977: 197; Wood, S. L. 1973c: 182.

ramosus Bright 1972c: 1677. Holotype ♀; 5.6 km S Suchixtepec, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 182.

References: (tx) Bright 1972c: 1675, 1677; Wood, S. L. 1973c: 182, 1982b: 148.

velutinus Wood 1958a: 48. Holotype ♀; 23 km NW Guadalajara, Jalisco, Mexico; SMUK, Lawrence. Distribution: North America (Jalisco in Mexico). Hosts: *Pinus* sp.

References: (ds) Atkinson & Equihua 1988: 94; Wood, S. L. 1982b: 1151. (tx) Wood, S. L. 1958a: 47–48, 1982b: 1151.

Genus *Dacnophthorus* Wood

DACNOPHTHORUS WOOD 1975b: 394. Type-species: *Gnathophthorus clematis* Wood, original designation.

Keys: Wood 1982b: 1152 for Central America.

Notes: The species of this genus breed in the phloem of lianas.

References: (hb) Wood, S. L. 1986a: 98. (ds) Wood, S. L. 1986a: 98. (tx) Wood, S. L. 1975b: 394, 1982b: 1152–1154, 1986a: 98.

artus (Wood) 1974a: 27 (*Gnathophthorus*). Holotype ♀; ca 260 km N Xavantina, Matto Grosso, Brasil (12 49 S, 41 46 W); BMNH, London.

Distribution: South America (Brazil).

Notes: (1) Wood 1975b: 394 (to *Dacnophthorus*).

References: (tx) Wood, S. L. 1974a: 27, 1975b: 394.

clematis (Wood) 1971: 51 (*Gnathophthorus*). Holotype ♀; Volcan Colima, Jalisco, Mexico; Wood Collection.

Figures: Wood 1971: 52, 1982b: 1153 (female outline, head).

Distribution: North America (Costa Rica/ Honduras/ Jalisco in Mexico/ Panama).

Hosts: *Clematis* sp.

Notes: (1) Wood 1975b: 394 (to *Dacnophthorus*).

References: (hb) Wood, S. L. 1982b: 1153. (ds) Wood, S. L. 1982b: 1153. (tx) Wood, S. L. 1971: 51–52, 1975b: 394, 1982b: 1153, 1986a: 98.

craceus (Wood) 1971: 53 (*Gnathophthorus*). Holotype ♀; La Ceiba, Atlantida, Honduras; Wood Collection.

Distribution: North America (Honduras).

Notes: (1) Wood 1975b: 394 (to *Dacnophthorus*).

References: (ds) Wood, S. L. 1982b: 1154. (tx) Wood, S. L. 1971: 53, 1975b: 394, 1982b: 1154.

pertusus (Wood) 1971: 53 (*Gnathophthorus*). Holotype ♀; 20 km SW El Vigía, Merida, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: Bignoniaceae liana.

References: (tx) Wood, S. L. 1971: 53–54, 1975b: 394.

rallus (Wood) 1971: 53 (*Gnathophthorus*). Holotype ♀; Merida, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Clematis* sp.

Notes: (1) Wood 1975b: 394 (to *Dacnophthorus*).

References: (tx) Wood, S. L. 1971: 53, 1975b: 394.

Genus *Pityotrichus* Wood

PITYOTRICHUS WOOD 1962: 76. Type-species: *Pityophilus barbatus* Blackman, automatic.

Pityophilus Blackman 1928b: 147. Type-species: *Pityophilus barbatus* Blackman, original designation, preoccupied by Brulle 1884.

References: (tx) Arnett 1960: 1045, 1968: 1045; Blackman 1928b: 8–16, 147–148; Chamberlin 1939: 345; Schedl 1938a: 159, 163, 165, 168; Wood, S. L. 1961: 48, 1962: 76, 1982b: 1140, 1986a: 98.

Keys: Bright 1971b: 68–69, Wood 1982b: 1141.

Notes: The species of this genus breed in the shaded-out, dying branches of living *Pinus* spp.

References: (hb) Wood, S. L. 1982b: 1140, 1986a: 98. (ds) Wood, S. L. 1982b: 1140, 1986a: 98. (tx) Bright 1971b: 68–69; Wood, S. L. 1962: 76, 1982b: 1140, 1986a: 98.

barbatus (Blackman) 1928b: 147 (*Pityophilus*). Holotype ♀; Las Vegas H.S. [Hot Springs], New Mexico [USA]; USNM, Washington.

Distribution: North America (Arizona, New Mexico in USA).

Hosts: *Pinus edulis*, *P. cembroides*, uncommon in *P. ponderosa*.

Notes: (3) Blackman 1928b: 147–148 (*Pityophthorus edulis*, nomen nudum, synonymy), 1928b: 147–149 (*Pityophthorus nitidicollis*, nomen nudum, synonymy).

References: (hb) Chamberlin 1939: 345; Wood, S. L. 1952b: 1141. (ds) Chamberlin 1939: 345; Furniss, R. L. & Carolin 1977: 400; Keen 1929a: 34; Kleine 1934a: 163; Leng & Mutchler 1933: 52; Wood, S. L. 1952b: 1141. (tx) Blackman 1928b: 147–148; Bright 1971b: 68; Chamberlin 1939: 345; Keen 1929a: 34; de Ruette 1970: 107; Schedl 1935a: 162; Wood, S. L. 1952b: 1141.

hesperius Bright 1971b: 69. Holotype ♀; Pinaleno Mts., Graham County, Arizona [USA]; CNCI, Ottawa.

Distribution: North America (Arizona, New Mexico in USA).

Hosts: *Pinus flexilis*, *P. strobiformis*.

References: (ds) Furniss, R. L. & Carolin 1977: 401; Wood, S. L. 1952b: 1142. (tx) Bright 1971b: 69–70; McNamara 1977: 198; Wood, S. L. 1952b: 1142.

Genus *Gnatholeptus* Blackman

GNATHOLEPTUS BLACKMAN 1943b: 34. Type-species: *Gnatholeptus mandibularis* Blackman = *Pityophthorus shannoni* Blackman, original designation.

Keys: Wood 1952b: 1142.

Notes: (1) Bright 1981d: 16 (treated as a subgenus of *Pityophthorus*).

References: (ds) Wood, S. L. 1956a: 98. (tx) Blackman 1943b: 34; Bright 1981: 16; Wood, S. L. 1952b: 1142–1146, 1956a: 98.

panamensis Blackman 1943b: 35. Holotype ♀; Barro Colorado Island, Canal Zone, Panama; USNM, Washington.

Figures: Wood 1952b: 1144 (female head).

Distribution: North America (Costa Rica/Panama), South America (Suriname).

Hosts: *Protium* sp.

References: (ds) Blackman 1943b: 35; Bright 1981d: 19; Wood, S. L. 1952b: 1145. (tx) Blackman 1943b: 35; Bright 1981d: 19; Wood, S. L. 1979b: 134, 1952b: 1145.

epistomalis Schedl 1961i: 224 (*Pityophthorus*). Holotype ♀; Barro Colorado Island, Canal Zone, Panama; Cornell University, Ithaca, New York. Synonymy: Wood 1979b: 134.

References: (tx) Bright 1977: 513; Schedl 1961i: 224; Wood, S. L. 1979b: 134, 1952b: 145.

semiermis (Nunberg) 1963c: 98 (*Pityophthorus*). Holotype ♂; Finca La Lola, Limon, Costa Rica; University of Wisconsin Collection, Madison.

Figures: Nunberg 1963c: 102, Wood 1952b: 1144 (female head).

Distribution: North America (Costa Rica), South America (Colombia/Suriname/Venezuela).

Hosts: *Protium* cf. *copal*.

Notes: (1) Bright 1977: 513 (to *Gnatholeptus*).

References: (ds) Bright 1981d: 21; Estrada & Atkinson 1988: 212; Wood, S. L. 1952b: 1145. (tx) Bright 1977: 513, 1981d: 21; Nunberg 1963c: 98, 102; Wood, S. L. 1952b: 1145.

shannoni (Blackman) 1942a: 224 (*Pityophthorus*). Holotype ♀; Cano Saddle at Gatun Lake, Panama; USNM, Washington.

Figures: Bright 1981d: 334, Wood 1952b: 1144 (female head).

Distribution: North America (Panama), South America (Colombia).

Hosts: *Protium* sp.

Notes: (1) Bright 1977: 513 (to *Gnatholeptus*).

References: (ds) Blackwelder 1947: 782; Bright 1981d: 20; Wood, S. L. 1952b: 1144. (tx) Blackman 1942a: 224; Bright 1977: 513, 1981d: 20, 334; Schedl 1961i: 226; Wood, S. L. 1979b: 134, 1952b: 1144.

mandibularis Blackman 1943b: 34. Holotype ♀; Barro Colorado Island, Canal Zone, Panama; USNM, Washington. Synonymy: Wood 1979b: 134.

References: (tx) Blackman 1943b: 34; Wood, S. L. 1979b: 134, 1952b: 1144.

gentilis Schedl 1961i: 225 (*Pityophthorus*). Holotype ♂; Barro Colorado Island, Gatun Lake, Canal Zone, Panama; Cornell University Collection, Ithaca, New York. Synonymy: Bright 1977: 513, Wood 1979b: 134.

References: (tx) Bright 1977: 513; Schedl 1961i: 225, 1979c: 103; Wood, S. L. 1979b: 134, 1952b: 1144.

Genus *Pityodendron* Schedl

PITYODENDRON SCHEDL 1953d: 93. Type-species: *Pityodendron madagascariensis* Schedl, monobasic.

References: (hb) Wood, S. L. 1956a: 98. (ds) Wood, S. L. 1956a: 98. (tx) Schedl 1953d: 93, 1964m: 307, 1977b: 106; Wood, S. L. 1956a: 98.

madagascariensis Schedl 1953d: 93. Lectotype, sex?; Madagascar, Mt. d'Ambre; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 144.

Figures: Schedl 1977a: 105.

Distribution: Madagascar.

References: (tx) Schedl 1953d: 93, 1977b: 105–106, 1979c: 144.

Genus *Saurophtilius* Browne

SAUROPTILIUS BROWNE 1970: 558. Type-species: *Xyleborus sauropterus* Schedl, original designation.

References: **(hb)** Wood, S. L. 1986: 98. **(ds)** Wood, S. L. 1986a: 98. **(tx)** Browne 1970: 558–559; Schedl 1977b: 114; Wood, S. L. 1986a: 98.

sauropterus (Schedl) 1953d: 101 (*Xyleborus*). Holotype ♂; Madagascar, Sikora; Schedl Collection in NHMW, Wien.

Figures: Schedl 1977b: 113.

Distribution: Madagascar.

Hosts: *Mauloutchia* sp.

References: **(tx)** Schedl 1953d: 101–102, 1970d: 558–559, 1977b: 113–114, 1979c: 220; Wood, S. L. 1986a: 98, 1989: 175.

saurophteroides Schedl 1970d: 237 (*Xyleborus*). Holotype ♀; Madagascar, Bemampidy; MNHN, Paris. Synonymy: Wood 1989: 175.

References: **(tx)** Schedl 1970d: 234, 237, 1977b: 114, 1979c: 220; Wood, S. L. 1989: 175.

Genus *Phelloterus* Wood

PHELLOTERUS WOOD 1971: 46. Type-species: *Phelloterus tersus* Wood, original designation.

References: **(hb)** Wood, S. L. 1986a: 98. **(ds)** Wood, S. L. 1986a: 98. **(tx)** Wood, S. L. 1971: 46–47, 1986a: 98.

anaxeus Wood 1971: 47. Holotype ♂; 8 km S Colonia (near Buenaventura), Valle de Cauca, Colombia; Wood Collection.

Distribution: South America (Colombia).

Hosts: *Couma macrocarpa*, *Eshweilera* sp., *Lecythia* sp., *Licania* sp., *Sacoglochia procera*.

References: **(tx)** McNamara 1977: 197; Wood, S. L. 1971: 47.

atrocis Wood 1971: 48. Holotype ♂; Campamento Capote, 27 km NE Montoya, Santander, Colombia; Wood Collection.

Distribution: South America (Colombia).

Hosts: *Eschweilera* sp., *Sacoglochia* sp.

References: **(tx)** Wood, S. L. 1971: 48.

tersus Wood 1971: 47. Holotype ♂; Colonia Tovar, Aragua, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Eschweilera* sp.

References: **(tx)** Wood, S. L. 1971: 47, 1986a: 98.

Genus *Spermophthorus* Costa Lima

SPERMOPHTHORUS COSTA LIMA 1929: 111. Type-species: *Spermophthorus apuleiae* Costa Lima, monobasic.

References: **(hb)** Wood, S. L. 1986: 98. **(ds)** Wood, S. L. 1986: 98. **(tx)** Blackman 1942a: 178; Costa Lima 1929: 111–112; Schedl 1937b: 165, 1938a: 160, 168; Wood, S. L. 1982b: 963–965, 1986a: 98.

aberrans Wood 1968b: 11. Holotype ♂; 6 km S San Vito, Puntarenas, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: 3.5 x 4.5 cm gall found on forest floor.

References: **(hb)** Wood, S. L. 1982b: 964. **(ds)** Wood, S. L. 1982b: 964. **(tx)** Wood, S. L. 1968b: 11, 1982b: 964.

apuleiae Costa Lima 1929: 111. Syntypes ♂ ♀; Campos, Estado do Rio, Brasil; Inst. Oswaldo Cruz, Rio de Janeiro.

Distribution: South America (Argentina/ Brazil).

Hosts: *Apuleia forrea* seeds.

References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956. **(ds)** Costa Lima 1936, 1956. **(tx)** Blackman 1942a: 203–204; Costa Lima 1929: 111, 1956; Schedl 1938a: 162, 1964m: 311; Wood, S. L. 1986a: 98.

vianai Schedl 1938i: 27 (*Conophthocranulus*).

Syntypes ♂ ♀; Argentinien, Prov. Salta; Viana Collection, and Schedl Collection in NHMW, Wien. Synonymy: Schedl 1964m: 311.

Notes: (1) Schedl 1979c: 267 (citation of holotype invalid).

References: **(hb)** Viana 1964: 125. **(ds)** Viana 1964: 125. **(tx)** Schedl 1938i: 27, 1939m: 169, 1964m: 311, 1979c: 267.

caesalpiniae Blackman 1942a: 203. Holotype ♀; Paraguay; USNM, Washington.

Distribution: South America (Brazil/ Paraguay).

Hosts: *Caesalpinia melanocarpa*.

References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956. **(ds)** Costa Lima 1956. **(tx)** Blackman 1942a: 203; Costa Lima 1956.

Genus *Pseudopityophthorus* Swaine

PSEUDOPITYOPHTHORUS SWAINE 1918a: 93. Type-species: *Crypturgus minutissimus* Zimmermann, original designation.

Xenophthorus Wood & Yin 1986: 462. Type-species: *Pseudopityophthorus peregrinus* Wood & Yin, original designation. Used as a subgenus for this Asiatic species.

References: **(tx)** Wood, S. L. & Yin 1986: 462.

Keys: Swaine 1918a: 93, Blackman 1931a: 225, Wood 1982b: 966.

Notes: (3) Blackman 1931a and Wood 1982b: 965–982 (revision of genus).

References: **(ay)** Nobuchi 1969a: 67. **(cn)** Rexrode 1967: 755–757, 1969a: 448–449. **(cc)** Rexrode 1969a: 448–449. **(hb)** Anderson, R. F. 1960: 243; Bright & Stark 1973: 101; Cowlin 1956: 35–36; Griswold & Neiswander 1953: 208; Gruenfeld et al. 1956: 92–93; McGinnies 1955: 47; Orr 1955: 4, 7; Rexrode & Krause 1965: 1340–1341; Stambough 1955: 567–571; Whiteside 1957: 2, 20; Wood, S. L. 1982b: 965, 1986a: 98. **(ds)** Anderson, R. F. 1960: 243; Bright & Stark 1973: 101; Leng & Mutchler 1933: 95; Wood, S. L. 1982b: 965, 1986a: 98. **(tx)** Anderson, R. F.

- 1960: 243; Arnett 1960: 1045; Beal & Massey 1945: 60, 121; Blackman 1922b: 98–99, 1928b: 6–16, 1931a: 223–236, 1942a: 175; Bright 1976d: 176; Bright & Stark 1973: 101; Chamberlin 1939: 346–353; Dodge 1938: 19; Schedl 1938a: 158, 163, 168; Swaine 1918a: 93; Wood, S. L. 1961a: 48, 1967d: 37–57, 1982b: 965–982, 1986a: 98; Wood, S. L. & Yin 1986: 462.
- agrifoliae** Blackman 1931a: 230. Holotype ♂; Golden Gate Park, San Francisco, California [USA]; USNM, Washington.
Distribution: North America (S Arizona, California in USA).
Hosts: *Quercus agrifolia*, *Q. kelloggii*, *Q. wislizenii*, *Q. sp.*
References: (hb) Bright & Stark 1973: 101; Chamberlin 1939: 350; Wood, S. L. 1982b: 975. (ds) Atkinson & Equihua 1985: 100; Bright & Stark 1973: 101; Chamberlin 1939: 350; Leng & Mutchler 1933: 95; Wood, S. L. 1982b: 975. (tx) Blackman 1931a: 230; Chamberlin 1939: 350; Wood, S. L. 1967d: 42, 1982b: 975.
- asperulus** (LeConte) 1868: 155 (*Cryphalus*). Holotype ♂; Virginia [USA]; MCZ, Cambridge.
Figures: Blackman 1922b: pl. 1, fig. 4.
Distribution: North America (Connecticut, District of Columbia, Florida, Georgia, Illinois, Louisiana, Maine, Massachusetts, Mississippi, Missouri, New Jersey, New York, North Carolina, Pennsylvania, Tennessee, E Texas, West Virginia in USA).
Hosts: *Quercus nigra*, *Q. rubra*, *Q. velutina*, *Q. sp.*, rare and not persistent in *Betula populifera*, *Castanea dentata*, *Ostrya virginiana*.
Notes: (1) Blackman 1931a: 224 (to *Pseudopityophthorus*). (3) Blackman 1931a: 226 (re-described).
References: (bv) Atkinson, Foltz, & Connor 1988; Roling & Kearby 1975b, 1977. (cn) Blackman 1950; Doane et al. 1936; Felt 1906: 726; Packard 1890: 720; Roling & Kearby 1977. (ec) Roling & Kearby 1977; Steinhaus 1946: 404. (hb) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 256; Beal & Massey 1945: 124; Blackman 1950; Chamberlin 1939: 349; Chittenden 1890; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Hubbard 1897; Packard 1890: 720; Schwarz 1858a: 80; Wood, S. L. 1982b: 971. (ds) Atkinson et al. 1991: 163; Baker, W. L. 1972: 256; Beal & Massey 1945: 124; Blackman 1950; Blatchley & Leng 1916: 647; Bright 1976d: 177; Chamberlin 1939: 349; Chittenden 1890; Deyrup 1981b: 8, 1987a: 66; Dorsey & Leach 1955: 224; Drooz 1985: 363; Gemminger & Harold 1872: 2684, 2688; Hagedorn 1910d: 96; Henshaw 1885: 147; Hubbard & Schwarz 1878b: 626; Kleine 1913b: 145, 158, 1914b: 394, 1934a: 165; Leng 1920: 340; Schwarz 1886: 40, 1888a: 80; Swaine 1909: 108; Ulke 1902: 56; Weber, B. C. & McPherson 1991: 54; Wood, S. L. 1982b: 971. (tx) Beal & Massey 1945: 124; Blackman 1931a: 224–228, 1931b: 266; Blatchley & Leng 1916: 647; Bright 1976d: 177; Chamberlin 1939: 349; Eichhoff 1878b: 202; Hagedorn 1910a: 147; Hubbard & Schwarz 1878a: 666; LeConte 1868: 155, 1876: 350; Swaine 1909: 108, 1918a: 91–92; Wood, S. L. 1959b: 2, 1982b: 971. *gracilis* Blackman 1921: 6. Lectotype ♂; Natchez, Mississippi [USA]; USNM, Washington, designated by Wood 1982b: 971. Synonymy: Blackman 1931a: 228.
References: (hb) Blackman 1922b: 99–100. (ds) Blackman 1922b: 99–100; Henshaw 1882i: 268; Kleine 1934a: 162; Leng & Mutchler 1927: 52. (tx) Blackman 1921: 6–7, 1922b: 99–100, 1931a: 224–228.
- cincinnatus** (Blandford) 1904: 242 (*Pityophthorus*). Holotype ♀; Quiche Mts., Guatemala, 7–9000 ft.; BMNH, London.
Distribution: North America (Guatemala).
References: (ds) Hagedorn 1910d: 70; Kleine 1913b: 140; Wood, S. L. 1982b: 973. (tx) Blandford 1904: 242; Bright 1981d: 328; Hagedorn 1910a: 100; Schedl 1955g: 17; Wood, S. L. 1982b: 973.
- colombianus** Wood 1971: 51. Holotype ♂; Piedras Blancas, 10 km E Medellin, Antioquia, Colombia; Wood Collection.
Distribution: South America (Colombia).
Hosts: *Quercus humboldtii*.
References: (tx) Wood, S. L. 1971: 51, 1982b: 966.
- comosus** Bright 1972c: 1670. Holotype ♂; Highway 131, 178 km S Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.
Figures: Bright 1972c: 1669, 1675.
Distribution: North America (Oaxaca in Mexico).
Hosts: *Quercus sp.*
References: (ds) Wood, S. L. 1982b: 978. (tx) Bright 1972c: 1669–1670; McNamara 1977: 198; Wood, S. L. 1982b: 978.
- declivis** Wood 1971: 50. Holotype ♀; Laguna Santa Maria, Nayarit, Mexico; Wood Collection.
Distribution: North America (Chiapas, Jalisco, Nayarit, Oaxaca in Mexico).
Hosts: *Quercus sp.*
References: (hb) Wood, S. L. 1982b: 969. (ds) Atkinson & Equihua 1985c: 358; Schedl 1977e: 42; Wood, S. L. 1982b: 969. (tx) Wood, S. L. 1971: 50, 1973c: 184, 1982b: 969.
- truncatus** Bright 1972c: 1673. Holotype ♂; 184 km S Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.
Synonymy: Wood 1973c: 184.
References: (tx) Bright 1972c: 1673–1675; McNamara 1977: 199; Wood, S. L. 1973c: 184.
- curtus** Bright 1972c: 1674. Holotype ♀; 13 km N Ocosingo, Chiapas, Mexico; CNCI, Ottawa.
Synonymy: Wood 1973c: 184.
References: (tx) Bright 1972c: 1674–1675; McNamara 1977: 199; Wood, S. L. 1973c: 184.

denticulus Wood 1977b: 216. Holotype ♀; Boot Springs, Big Bend National Park, Brewster Co., Texas [USA]; CNCI, Ottawa.

Distribution: North America (W Texas in USA).

Hosts: *Quercus* sp.

References: (hb) Atkinson et al. 1986: 41; Burgos & Saucedo 1983: 112. (ds) Atkinson & Equihua 1985c: 358; Atkinson et al. 1986: 41; Burgos & Saucedo 1983: 112; Wood, S. L. 1982b: 971. (tx) Wood, S. L. 1977b: 216, 1982b: 971.

durangoensis Wood 1987: 549. Holotype ♂; 96 km W Durango, Durango, Mexico; Wood Collection. Distribution: North America (Durango in Mexico).

Hosts: *Quercus* sp.

References: (tx) Wood, S. L. 1987: 549.

fagi Blackman 1931a: 228. Holotype ♂; Morgantown, West Virginia [USA]; USNM, Washington. Distribution: North America (Pennsylvania, West Virginia in USA).

Hosts: *Fagus grandifolia*.

References: (ec) Felt 1906: 702. (hb) Chamberlin 1939: 349; Felt 1906: 702. (ds) Chamberlin 1939; Leng & Mutchler 1933: 95; Smith, J. B. 1910: 401; Wood, S. L. 1982b: 975. (tx) Blackman 1931a: 225–229; Chamberlin 1939: 349; Wood, S. L. 1982b: 975.

festivus Wood 1974a: 55. Holotype ♂; 11 km NE Copala, Sinaloa, Mexico; Wood Collection.

Distribution: North America (Sinaloa in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 980. (ds) Wood, S. L. 1982b: 980. (tx) Wood, S. L. 1974a: 55, 1982b: 980.

granulatus Blackman 1931a: 230. Holotype ♂; Prescott National Forest, Arizona [USA]; USNM, Washington.

Distribution: North America (Arizona in USA).

Hosts: *Quercus hypoleucoides*, *Q.* spp.

References: (hb) Chamberlin 1939: 350; Wood, S. L. 1982b: 970. (ds) Chamberlin 1939: 350; Leng & Mutchler 1933: 95; Wood, S. L. 1982b: 970. (tx) Blackman 1931a: 225, 227, 230; Chamberlin 1939: 350; Wood, S. L. 1967d: 42, 1982b: 970.

granulifer Wood 1967d: 42. Holotype ♀; Zamorano, Morazan, Honduras; Wood Collection.

Distribution: North America (Guatemala/ Honduras/ Chiapas in Mexico).

Hosts: *Quercus hondurensis*, *Q. sapotaefolia*.

References: (ec) Bright 1976a. (hb) Wood, S. L. 1982b: 970. (ds) Bright 1972c: 1679; Wood, S. L. 1982b: 970. (tx) Wood, S. L. 1967d: 42–43, 1982b: 970.

hispidus Eggers 1930a: 170. Holotype ♀; labeled "R.d.M. Mexico," published as Valle de Mexico, Mexico; MNB, Berlin.

Distribution: North America (Distrito Federal in Mexico).

References: (ds) Atkinson & Equihua 1985a: 90; Blackwelder 1947: 782; Ferrer 1942; Wood, S. L. 1982b: 968. (tx) Eggers 1930a: 170; Schedl 1940a: 348, 1979c: 118; Wood, S. L. 1982b: 968.

hondurensis Wood 1967d: 42. Holotype ♂; Buenos Aires, Cortes, Honduras; Wood Collection.

Figures: Bright 1972c: 1669.

Distribution: North America (Honduras/ Chiapas in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 972. (ds) Wood, S. L. 1982b: 972. (tx) Wood, S. L. 1967d: 42, 1973c: 184, 1982b: 972.

montanus Bright 1972c: 1667. Holotype ♂; Mt. Tzontehuitz, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 184.

References: (tx) Bright 1972c: 1667, 1669; McNamara 1977: 199; Wood, S. L. 1973c: 184.

limbatus Eggers 1930a: 169. Holotype ♀; "R.d.M.," Mexico; Eggers Collection, in NHMW, Wien.

Figures: Atkinson et al. 1986: 43.

Distribution: North America (Distrito Federal, Durango, Hidalgo, Michoacan, Oaxaca, Tlaxcala in Mexico).

Hosts: *Quercus* sp.

References: (hb) Atkinson & Equihua 1985c: 358; Burgos & Saucedo 1983: 113; Wood, S. L. 1982b: 975. (ds) Atkinson & Equihua 1985a: 89, 1985c: 358, 1988: 100; Blackwelder 1947; Burgos & Saucedo 1983: 113; Ferrer 1942; Wood, S. L. 1982b: 975. (tx) Atkinson et al. 1986: 43; Eggers 1930a: 169; Schedl 1940a: 347, 1979c: 139; Wood, S. L. 1966b: 27, 1975b: 394, 1982b: 976.

micans Wood 1967d: 44. Holotype ♂; 96 km W Durango, Durango, Mexico; Wood Collection. Synonymy: Wood 1975b: 394.

References: (hb) Thomas, J. B. 1967. (tx) de Ruette 1970: 113; Thomas, J. B. 1967; Wood, S. L. 1967d: 44–45, 1973c: 184, 1975b: 394.

minutissimus (Zimmermann) 1868: 143 (*Crypturgus*). Lectotype ♂; Carolina [USA]; MCZ, Cambridge, designated by Wood 1982b: 976.

Figures: Blackman 1922b: pl. 9, fig. 45, Bright 1976d: 204, 213.

Distribution: North America (Arkansas, District of Columbia, Florida, Georgia, Illinois, Iowa, Kansas, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Quercus* spp., rare and apparently not persistent in *Amelanchier* sp., *Betula* sp., *Carpinus* sp., *Carya* sp., *Fagus* sp., *Hamamelis* sp.

Notes: (3) Blackman 1931a: 231 (redescribed).

References: (ay) Hopkins 1894g. (bv) Haack, R. A., Benjamin, & Haack 1983; Rexrode 1969: 306–313; Rexrode & Jones 1972; Roling & Kearby 1975b, 1977; Turnbow & Franklin 1980. (cn)

- Anonymous 1979d; Baker, W. L. 1972: 255; Blackman 1950; Buchanan 1956: 654, 1958a: 9, 1958b: 414–415, 1958c: 646; Clittenden 1898: 78; Currie 1905: 80; Doane et al. 1936; Felt 1905: 257, 295, 1906: 295, 1926: 247–248, 1930a: 272; Felt & Rankin 1932: 359; Fitch 1858; Hopkins 1894: 277, 1898a: 105, 1904a: 24; MacAloney & Ewan 1964; McGinnies 1957: 45–47; Packard 1890: 93, 221, 720; Rexrode & Jones 1971: 108–111; Roling & Kearby 1977; Walsh 1867: 103. (**cc**) Baker, W. L. 1972: 255; Berry & Bretz 1959: 66–67, 1966; Buchanan 1956: 654, 1958a: 9, 1958b: 414–415, 1958c: 646; Burks 1979: 791; Bushing 1965: 465; Clittenden 1901; Drooz 1985: 362; Griswold 1956: 560; Griswold & Bart 1954; Griswold & Neiswander 1953, 1954: 708; Haack, R. A., Benjamin, & Haack 1983; Marsh 1979: 150; Pierce, W. D. 1908: 385; Rexrode 1969, 1976; Rexrode & Jones 1971: 108–111, 1972; Rexrode, Jones, & Jones 1972; Rexrode & Krause 1968: 814–816; Roling & Kearby 1977; Singh 1977: 101; Thewke & Enns 1965: 215–219. (**hb**) Baker, W. L. 1972: 255; Beal & Massey 1945; Blackman 1922b: 99–100, 1950; Bright 1976d: 177; Buchanan 1956, 1958b; Chamberlin 1939: 350; Clittenden 1890; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Drooz 1985: 362; Felt 1906: 295, 1926: 247–248, 1930a: 272; Felt & Rankin 1932: 359; Fitch 1858; Haack, R. A., Benjamin, & Haack 1983; Hopkins 1894g, 1899a: 343, 1904a: 24; MacAloney & Ewan 1964; McMullen et al. 1955: 491; Packard 1890: 221, 720; Pierce, W. D. 1907: 290; Rexrode 1969; Rexrode & Jones 1972; Rexrode, Jones, & Jones 1972; Rexrode & Krause 1968a; Schwarz 1858a: 113; Swaine 1918a: 93–94; Wood, S. L. 1982b: 976. (**ds**) Anonymous 1926c: 518, 1979d; Atkinson et al. 1991: 163; Beal & Massey 1945; Beaulne 1956; Blackman 1922b: 99–100, 1950; Blandford 1895b; Blatchley & Leng 1916: 628; Bright 1976d: 177; Chamberlin 1939: 350; Clittenden 1890; Currie 1905; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Dodge 1938; Drooz 1985: 362; Felt 1926: 247–248, 1930a: 247, 272; Felt & Rankin 1932: 359; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 72; Henshaw 1885: 147; Hopkins 1893a: 129, 149, 1893b: 208; Hubbard & Schwarz 1878a: 665; Kirk 1970; Kleine 1913b: 140, 1914b: 392, 1934a: 162; Leng 1920: 341; Leonard 1928: 518; Rexrode 1969; Rexrode & Jones 1972; Schwarz 1886: 40, 1888a: 113; Smith, J. B. 1900: 362 1910: 401; Swaine 1909: 139; Turnbow & Franklin 1980; Ulke 1902: 56; Weber, B. C. & McPherson 1991: 54; Wickham 1896c: 168; Wood, S. L. 1982b: 976. (**tx**) Beal & Massey 1945; Benoit 1985: 211; Blackman 1921: 6–7, 1922b: 99–100, 1928b: 148, 1931a: 224–232; Blandford 1895b; Blatchley & Leng 1916: 628; Bright 1976d: 204, 213; Chamberlin 1939: 350; Dodge 1938: 16, 41; Eichhoff 1878b: 200; Green 1918: 257; Hagedorn 1910a: 101; LeConte 1868: 154, 157, 1876: 351, 1878b: 665; Schedl 1938a: 155; Schwarz 1886: 40, 1888: 56, 113, 1891a: 168; Swaine 1909: 139–140, 1918a: 93–94; Wood, S. L. 1982b: 976; Zimmermann 1868: 143. (**ms**) Rexrode & Krause 1968a. (?) *pusillus* Harris 1837: 82 (*Tomicus*). Holotype, sex?, eastern U.S.; lost.
- Notes: (1) LeConte 1868: 143 (possible synonymy with *minutissimus*). The probability of this synonymy being correct is remote, but no alternative has been suggested. Because of this uncertainty, the name *minutissimus* is retained as the name for this species.
- References: (**cn**) Fitch 1858; Packard 1890: 717. (**hb**) Fitch 1858; Packard 1890: 717. (**ds**) Gemminger & Harold 1872: 2689; Leng 1920: 341; Melsheimer 1853: 87; Swaine 1909: 140. (**tx**) Blackman 1931a: 225; Blackman & Stage 1918: 55; Eichhoff 1878b: 200; Harris 1837: 82; LeConte 1868: 143, 1876: 351; Swaine 1909: 140.
- opacicolis** Blackman 1931a: 235. Holotype ♂; Santa Catalina Mts., Arizona [USA]; USNM, Washington.
- Figures: Bright 1972c: 1675.
- Distribution: North America (Chiapas, Durango, Hidalgo, Oaxaca, Tlaxcala in Mexico/S Arizona in USA).
- Hosts: *Quercus emoryi*, *Q. spp.*
- References: (**hb**) Atkinson et al. 1986: 42; Burgos & Saucedo 1983: 115; Chamberlin 1939: 353; Wood, S. L. 1982b: 980. (**ds**) Atkinson & Equihua 1985a: 89, 1985c: 359, 1988: 100; Atkinson et al. 1986: 42; Burgos & Saucedo 1983: 115; Chamberlin 1939: 353; Leng & Mutchler 1933: 95; Thomas, J. B. 1966. (**tx**) Blackman 1931a: 226–236; Chamberlin 1939: 353; Wood, S. L. 1959b: 1, 1967d: 44, 1973c: 185, 1982b: 980.
- aesculinus** Bright 1972c: 1672. Holotype ♀; 112 km N Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.
- Synonymy: Wood 1973c: 185.
- References: (**tx**) Bright 1972c: 1672, 1675; McNamara 1977: 198; Wood, S. L. 1973c: 185.
- peregrinus** Wood & Yin 1986: 462. Holotype ♀; Zayu, Xizang (Tibet), China; IZAS, Beijing.
- Distribution: Asia (Xizang [Tibet] in China).
- Hosts: *Quercus* sp.
- Notes: (1) This species is in the subgenus *Xenophthorus*.
- References: (**tx**) Wood, S. L. & Yin 1986: 462.
- pruinus** (Eichhoff) 1878b: 198 (*Pityophthorus*). Lectotype ♂; published as “Carolina,” labeled “Am. bor., No. 11” [USA]; USNM, Washington, designated by Wood 1982b: 979.
- Figures: Blackman 1922b: pl. 9, fig. 46, Bright 1972c: 1675.
- Distribution: North America (Guatemala/ Honduras/ Durango, Michoacan, Oaxaca, Sinaloa, Veracruz in Mexico/ Alabama, Arizona, Arkansas,

District of Columbia, Florida, Georgia, Louisiana, Maryland, Michigan, Mississippi, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA).

Hosts: *Quercus hondurensis*, *Q. hypoleucoides*, *Q. marylandica*, *Q. nigra*, *Q. sapotaefolia*, *Q.* spp.

Notes: (3) Blackman 1931a: 232 (redescribed).

References: (bv) Rexrode 1969; Rexrode & Jones 1972; Roling & Kearby 1975b, 1977; Turnbow & Franklin 1980. (cn) Anonymous 1971e; Blackman 1950; Chamberlin 1924; Doane et al. 1936; Dorsey et al. 1953: 419–420; Griswold 1955: 23–24; Hopkins 1903: 318; Rexrode & Jones 1971; Rexrode, Kulman, & Dorsey 1965; Roling & Kearby 1977; Swaine 1918a: 94. (cc) Burks 1979: 556; Bushing 1965: 465; Funk, D. W., Brooks, & Butler 1973; Griswold 1955: 23–24; Griswold & Bart 1954: 591; Griswold & Neiswander 1953: 208, 1954; Jewell 1956: 255; Marsh 1979: 150; Poinar 1975: 167; Rexrode 1969, 1976; Rexrode & Jones 1971, 1972; Rexrode, Jones, & Jones 1972; Rexrode, Kulman, & Dorsey 1965; Roling & Kearby 1977; Singh 1977: 101; Stambaugh et al. 1955: 867. (hb) Atkinson et al. 1986: 42; Baker, W. L. 1972: 256; Beal & Massey 1945: 122; Blackman 1922b: 99–100, 1950; Bright 1976d: 178; Burgos & Saucedo 1983: 113; Chamberlin 1939: 351; Doane et al. 1936; Hopkins 1904b: 314; Ostmark 1968; Rexrode 1969; Rexrode & Jones 1972; Rexrode, Jones, & Jones 1972; Rexrode & Krause 1968a; Rexrode, Kulman, & Dorsey 1965; Swaine 1918a: 94; Wood, S. L. 1982b: 979. (ds) Anonymous 1926c: 518; Atkinson & Equihua 1985c: 358; Atkinson et al. 1986: 42, 1991: 163; Baker, W. L. 1972: 256; Beal & Massey 1945: 122; Blackman 1922b: 99–100, 1950; Blatchley & Leng 1916: 629; Bright 1976d: 178; Burgos & Saucedo 1983: 113; Chamberlin 1939: 351; Deyrup 1981b: 8; Dorsey & Leach 1956: 220; Hagedorn 1910d: 73; Kirk 1969, 1970; Kleine 1913b: 140, 1914b: 384, 1934a: 162; Leng 1920: 341; Leonard 1928: 518; Norris 1955: 252; Ostmark 1968; Rexrode 1969; Rexrode & Jones 1972; Swaine 1909: 137; Turnbow & Franklin 1980; Wood, S. L. 1982b: 979. (tx) Beal & Massey 1945: 122; Blackman 1922b: 99–100, 1925b: 148, 1931a: 224–233; Blatchley & Leng 1916: 629; Bright 1976d: 178; Chamberlin 1939: 351; Drooz 1985: 363; Eichhoff 1878a: 390, 1878b: 198, 1896: 609–610; Hagedorn 1910a: 101; Schedl 1938a: 155; Swaine 1909: 137, 1918a: 93–94; Wood, S. L. 1973c: 185, 1982b: 979. (ms) Rexrode & Krause 1968a.

tomentosus Eichhoff 1878b: 201 (*Pityophthorus*). Holotype, sex?; America borealis [USA]; Hamburg Museum, lost. Synonymy: Blackman 1931a: 225.

References: (ds) Blatchley & Leng 1916: 630; Hagedorn 1910d: 7; Kleine 1913b: 140; Leng 1920: 341; Swaine 1909: 140. (tx) Blackman

1925b: 148, 1931: 224–233; Blatchley & Leng 1916: 630; Eichhoff 1878a: 390, 1878b: 201; Hagedorn 1910a: 101; Swaine 1909: 140.

querciperda Schwarz 1888a: 56 (*Pityophthorus*). Syntypes, sex?; New York to Florida [USA]; USNM, Washington. Synonymy: Eichhoff 1896: 609.

References: (cn) Packard 1890: 93. (cc) Felt 1906: 702. (hb) Chittenden 1890; Felt 1906: 702; Packard 1890: 93; Pierce, W. D. 1907: 291; Schwarz 1888a: 56. (ds) Chittenden 1890; Henshaw 1887: 8, 1895: 44; Leng 1920: 341; Schwarz 1888a: 56; Smith, J. B. 1900: 362, 1910: 401; Swaine 1909: 137. (tx) Blackman 1931a: 224–231; Eichhoff 1896: 609–610; Schwarz 1888a: 56; Swaine 1909: 137, 1918: 94.

pulvereus Blackman 1931a: 232. Holotype ♂; Chiricahua Reserve, Arizona [USA]; USNM, Washington. Synonymy: Wood 1973c: 185.

References: (hb) Bright & Stark 1973: 102; Chamberlin 1939: 351. (ds) Bright & Stark 1973: 102; Chamberlin 1939: 351; Leng & Mutchler 1933: 95; Wood, S. L. 1960b: 69. (tx) Blackman 1931a: 232; Chamberlin 1939: 351; Wood, S. L. 1959b: 1; 1967d: 43, 1973c: 185.

tropicalis Wood 1967d: 43. Holotype ♂; Zamorano, Morazan, Honduras; Wood Collection. Synonymy: Wood 1967d: 43–44, 1973c: 185. References: (tx) McNamara 1977: 199; Wood, S. L. 1967d: 43, 1973c: 185.

contexus Bright 1972c: 1672. Holotype ♂; 184 km S Oaxaca, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 185.

References: (tx) Bright 1972c: 1672, 1675; McNamara 1977: 199; Wood, S. L. 1973c: 185.

pubescens Blackman 1931a: 229. Holotype ♂; Tryon, North Carolina [USA]; USNM, Washington. Distribution: North America (Maryland, North Carolina, Oklahoma, Virginia in USA).

Hosts: *Quercus alba*, *Q. borealis*, *Q. rubra*, one record from *Castanea dentata*.

References: (bv) Turnbow & Franklin 1980. (cn) Hopkins 1904a: 16. (hb) Baker, W. L. 1972: 256; Beal & Massey 1945. (ds) Atkinson et al. 1991: 163; Baker, W. L. 1972: 256; Beal & Massey 1945; Chamberlin 1939: 349; Drooz 1985: 363; Leng & Mutchler 1933: 95; Rexrode 1969: 307; Turnbow & Franklin 1980; Wood, S. L. 1982b: 974. (tx) Beal & Massey 1945; Blackman 1931a: 229; Chamberlin 1939: 349; McNamara 1984: 759; Wood, S. L. 1982b: 974.

pubipennis (LeConte) 1860: 59 (*Bostrichus*). Lectotype ♀; San Jose, California [USA]; MCZ, Cambridge, designated by Wood 1982b: 981. Figures: Bright 1973: 161.

Distribution: North America (California, Oregon in USA).

Hosts: *Quercus agrifolia*, *Q. garryana*, *Q. kelloggii*.

Notes: (1) Swaine 1915a: 93 (to *Pseudopityophthorus*). (3) Blackman 1931a: 234 (re-described).

References: (cn) Anonymous 1960q, 1961p, 1963q, 1964m, 1964n, 1965b, 1965f; Currie 1905: 71; Doane et al. 1936; Essig 1926: 519, 1958: 519; Herbert 1936: 41; Hopkins 1904a: 16; Keen 1938: 132, 1952c: 169; Packard 1890: 93; Swaine 1915a: 93. (cc) Bushing 1965: 203; Keen 1938: 132; Marsh 1979: 15. (hb) Bright 1976d: 179; Bright & Stark 1973: 102; Chamberlin 1939: 352, 1958: Chittenden 1890; Doane et al. 1936; Essig 1926: 519, 1958: 519; Hopkins 1904a: 16; Keen 1938: 132, 1952c: 169; Packard 1890: 93; Pierce, W. D. 1907: 290; Swaine 1915a: 93. (ds) Anonymous 1961p, 1965b, 1965f; Blackwelder 1947: 782; Blandford 1904; Bright 1976d: 179; Bright & Stark 1973: 102; Chamberlin 1917: 354, 1939: 352, 1958; Chittenden 1890; Currie 1905: 71; Essig 1926: 519, 1958: 519; Furniss, R. L. & Carolin 1977: 402; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 74; Henshaw 1882: 268, 1885: 147; Keen 1929a: 34, 1938: 132, 1949a: 93, 1952c: 169; Kleine 1913b: 140, 1914b: 391, 1934a: 162; Lacordaire 1866: 353; Leng 1920: 341; Patterson & Hatch 1945: 152; Wood, S. L. 1972a: 634, 1982b: 951. (tx) Benoit 1956: 36; Blackman 1928b: 148, 1931a: 224–235; Blandford 1904: 243; Bright & Stark 1973: 161, 1976d: 179; Chamberlin 1939: 352, 1958; Eggers 1930a: 159; Eichhoff 1878b: 197; Ferrari 1867: 17; Hagedorn 1910a: 101; Keen 1929a: 34; Lacordaire 1866: 353; LeConte 1857: 22, 59, 1860: 59, 1865: 156, 1876: 351; Schedl 1935a: 158; Swaine 1909: 137–138, 1915a: 93; Wood, S. L. 1972a: 425, 1982b: 951.

singularis Wood 1971: 50. Holotype ♂; 1 km W Las Vigas, Veracruz, Mexico; Wood Collection.
Distribution: North America (Chiapas, Veracruz in Mexico).

Hosts: *Quercus* sp.

References: (ds) Atkinson & Equihua 1955c: 358; Wood, S. L. 1982b: 973. (tx) Wood, S. L. 1971: 50, 1973c: 185, 1982b: 973.

acuminatus Bright 1972c: 1671. Holotype ♂; 13 km NE San Cristobal de las Casas, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 185.

References: (tx) Bright 1972c: 1669–1671; McNamara 1977: 198; Wood, S. L. 1973c: 185.

squamosus Bright 1972c: 1670. Holotype ♀; 14 km W La Ciudad, Durango, Mexico; CNCI, Ottawa. Figures: Bright 1972c: 1669.

Distribution: North America (Durango, Sinaloa in Mexico).

Hosts: *Quercus* sp.

References: (ds) Atkinson & Equihua 1955c: 358; Wood, S. L. 1982b: 978. (tx) Bright 1972c: 1669–1670; McNamara 1977: 199; Wood, S. L. 1973c: 184, 1982b: 978.

tenuis Wood 1959b: 1. Holotype ♂; 15 km NE Jacala, Hidalgo, Mexico; SMUK, Lawrence.

Figures: Bright 1972c: 1669.

Distribution: North America (Chiapas, Hidalgo, Veracruz in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 974. (ds) Bright 1972c: 1679; Thomas, J. B. 1966; Wood, S. L. 1982b: 974. (tx) Wood, S. L. 1959b: 1–2, 1973c: 185, 1982b: 974.

hirsutus Bright 1972c: 1665. Holotype ♂; 5 km SE Teopisca, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 185.

References: (tx) Bright 1972c: 1668–1669; McNamara 1977: 199; Wood, S. L. 1973c: 185.

virilis Wood 1971: 50. Holotype ♂; 1 km W Las Vigas, Veracruz, Mexico; Wood Collection.

Distribution: North America (Hidalgo, Veracruz in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 971. (ds) Atkinson & Equihua 1955c: 358; Wood, S. L. 1982b: 971. (tx) Wood, S. L. 1971: 50, 1982b: 971.

xalapae Wood 1957: 549. Holotype ♂; Xalapa, Veracruz, Mexico; Wood Collection.

Distribution: North America (Veracruz in Mexico).

Hosts: Presumably from *Quercus* sp.

References: (tx) Wood, S. L. 1957: 549.

yavapaii Blackman 1931a: 233. Holotype ♂; Prescott National Forest, Arizona [USA]; USNM, Washington.

Distribution: North America (Chihuahua in Mexico/ Arizona, W Texas in USA).

Hosts: *Quercus griseus*, *Q. hypoleucoides*, *Q.* spp.
References: (hb) Chamberlin 1939: 352; Wood, S. L. 1982b: 982. (ds) Atkinson et al. 1991: 163; Chamberlin 1939: 352; Furniss, R. L. & Carolin 1977: 403; Leng & Mntchler 1933: 95; Wood, S. L. 1960b: 69, 1982b: 982. (tx) Blackman 1931a: 233; Chamberlin 1939: 352; Wood, S. L. 1959b: 1, 1960b: 69, 1982b: 982.

Genus *Pityophthorus* Eichhoff

PITYOPHTHORUS EICHHOFF 1864b: 39. Type-species: *Bostrichus lichtensteini* Ratzelburg, subsequent designation by Hopkins 1914: 127.

Trigonogenius Hagedorn 1912c: 354. Type-species: *Trigonogenius fallax* Hagedorn, monobasic, preoccupied by Solier 1849. Synonymy: Schedl 1952d: 347.

References: (tx) Hagedorn 1912c: 354; Lucas 1920: 683; Schedl 1952d: 347.

Hagedornius Lucas 1920: 683. Type-species: *Trigonogenius fallax* Hagedorn, automatic. Synonymy: Schedl 1952d: 347.

References: (tx) Lucas 1920: 683; Schedl 1952d: 347.

Myeloborus Blackman 1928b: 16. Type-species:

Pityophthorus ramiperda Swaine, original designation. Synonymy: Bright 1977: 511.

References: (tx) Arnett 1960: 1040, 1045, 1968: 1040, 1045; Blackman 1928b: 3–16, 1942a: 178, 199; Bright 1977: 511, 1978: 167; Bright & Stark 1973: 96; Chamberlin 1939: 338–343, 1958: 143; Craighead 1950: 331; Dodge 1938: 15, 41; Furniss & Carolin 1977: 398; Murayama 1963: 390; Sawamoto 1942: 166; Schedl 1935h: 342, 1937h: 165, 1938a: 159–167, 1959r: 42, 1964m: 311; Wood, S. L. 1961a: 47, 1972a: 424.

Gnathophorus Schedl 1935h: 342. Type-species: *Gnathophorus sparsepilosus* Schedl, preoccupied by Kirby 1837. Synonymy: Wood 1975b: 391.

References: (tx) Schedl 1935h: 342; Wood, S. L. 1962: 76, 1975b: 391.

Conophthorocranulus Schedl 1935h: 343. Type-species: *Conophthorocranulus blackman* Schedl, monobasic. Synonymy: Wood 1982b: 963.

References: (tx) Schedl 1935h: 343, 1937h: 165, 1938a: 160–168, 1950i: 146, 1955e: 258; Wood, S. L. 1982b: 963.

Breviophthorus Schedl 1938a: 176. Type-species: *Breviophthorus brasiliensis* Schedl, monobasic. Synonymy: Wood 1984b: 227.

References: (tx) Schedl 1938a: 160, 166, 176–177, 1958f: 33–46, 1959m: 545–557; Wood, S. L. 1984b: 227.

Pityophthoroides Blackman 1942a: 199. Type-species: *Pityophthoroides pudens* Blackman, original designation. Synonymy: Wood 1977b: 207.

References: (tx) Blackman 1942a: 178, 199; Schedl 1964m: 311; Wood, S. L. 1977b: 207.

Cladoborus Sawamoto 1942: 165. Type-species: *Cladoborus arakii* Sawamoto, monobasic. Synonymy: Bright 1977d: 511.

References: (tx) Bright 1977d: 511; Murayama 1963c: 390; Sawamoto 1942: 165–167; Schedl 1959r: 421.

Neomips Schedl 1954b: 37. Type-species: *Neomips brasiliensis* Schedl = *Pityophthorus dimorphus* Schedl, monobasic. Synonymy: Wood 1984b: 227.

References: (tx) Schedl 1954b: 37; Wood, S. L. 1984b: 227.

Ctenyophthorus Schedl 1955g: 26. Type-species: *Ctenyophthorus glabratus* Schedl, monobasic. Synonymy: Bright 1977d: 511.

References: Bright 1977d: 511; Schedl 1955g: 26, 1963a: 162, 1964m: 311; Scherdtfeger 1957: 494–508.

Gnathophthorus Wood 1962: 76. Type-species: *Gnathophthorus sparsepilosus* Schedl, automatic. Synonymy: Wood 1975b: 391.

References: (tx) Wood, S. L. 1962: 76, 1975b: 391.

Hypopityophthorus Bright 1981d: 14. Type-

species: *Pityophthorus inops* Wood, original designation. Synonymy: Wood 1983a: 648.

References: (tx) Bright 1981d: 14; Wood, S. L. 1983a: 648.

Keys: Blackman 1928b for Canada and USA, Bright 1981d and Wood 1982b: 991 for North and Central America, Blandford 1904: 235 for Mexico and Central America, Pfeffer 1976: 334 for Europe and part of Asia, Yin, Huang, & Li 1984: 119 for China.

References: (ay) Fuchs 1912: 14–33; Nobuchi 1969a: 67; Van Ryn-Tournel 1975. (bv) Chow et al. 1988; Schedl 1960f: 11. (cn) Bess 1944: 14–16; Blackman 1950: 295–330; Caesar 1910: 17; Felt 1906: 372–373, 1924: 272; Furniss, R. L. & Carolin 1977: 401; Haddow & Newman 1942: 1–17; Harper & Lockwood 1959: 116–128; Keen 1928: 53–54; Kondo & Taylor 1985: 42; Parr 1943: 419; Schultz & Kliejunas 1981; Zivojinovic 1948: 65–73; Zolk 1935: 258–294. (cc) Ashraf & Barryman 1969: 13; Balazy & Michalski 1964a; McCain, Koehler, & Tjosvold 1987; Nickle 1976b; Tomalak, Welch, & Garroway 1989b. (hb) Bright 1981d; Bright & Stark 1973: 103; Escherich 1923: 446, 476–477, 555, 619; Fisher 1937: 118; Formanek 1907: 33, 37–38; Judeich & Nitsche 1895: 448, 451; Karpinski 1933: 290–302; Keen 1933: 297, 1953: 37, 154; Lengerken 1939: 68, 75–76; Lloyd 1908: 124; Luggler 1899: 299; Lyons 1957: 163; Muesebeck 1938: 251–287; Munro 1946: 7–8, 43, 63; Rice 1968: 53–56; Schedl 1962j: 1, 1977b: 106; Scheidter 1934; Scherdtfeger 1957: 494; Trappen 1935: 143; Wichmann 1927: 348, 362, 374; Wood, D. L. & Stark 1968: 147; Wood, S. L. 1982b: 991, 1986a: 95. (ds) Beeson 1941: 364, 366, 379; Bertolini 1872: 201; Blackwelder 1947: 781; Bright 1981d, 1982a: 127; Bright & Stark 1973: 103; Chamberlin 1917: 353–356, 1918: 15; Duffy 1953: 9, 14; Fowler 1882: 262; Hagedorn 1910d: 69–75; Jacobson 1927: 463–478; Kostin 1973: 261; Leng 1920: 341; Luigioni 1929: 1000; Martin 1965: 40–43; Pfeffer 1976: 324, 1989a: 62; Rio Mora 1985; Rohwer 1917: 151–176; Schedl 1934f: 1643, 1960f: 11, 34, 1962j: 7, 1966b: 41, 1981b: 76; Scheerpeltz & Winkler 1930: 258; Scherdlin 1914: 255, 1920: 206; Schilsky 1885: 121; Schwarz 1885: 40; Swaine 1909: 82, 133–134; Tredl 1907: 14; Van Dyke 1922: 180; Westhoff 1882: 238; Wood, S. L. 1982b: 991, 1986a: 98. (tx) Arnett 1960: 1040, 1045, 1968: 1040, 1045; Baker, W. L. 1972: 254; Balachowsky 1949a: 232; Barbey 1901: 74; Beal & Massey 1945: 125, 135; Bedel 1885b: 396, 398; Blackman 1922b: 77, 100–102, 1928b: 1–183, 1932: 264–265, 1942a: 177, 1943: 24; Blandford 1891: 15–16, 1894b: 260, 1894d: 55, 57, 1895b: 81, 84–86, 1897a: 183–184, 1904: 231–237; Blatchley & Leng 1916: 627–628; Bright 1968c: 604–608, 1972d: 86–87, 1976d: 179, 1977: 511, 1978, 1981d: 1–377; Bright & Stark 1973: 103;

- Chamberlin 1939: 353–402, 1955: 146–148; Choo 1983: 75; Choo, Woo, & Nobuchi 1988b; Costa Lima 1929: 111–112; Dodge 1938: 15, 19, 42–43; Eggers 1914: 183, 1937: 80, 1940a: 123–141, 1940b: 99–108; Eichhoff 1864: 39, 1872b: 137, 1878b: 35, 47, 49, 173, 476, 1881a: 49, 192, 1883a: 107, 110; Ferrari 1867a: 32–35, 1867b: 114; Feytaud 1950: 9; Hagedorn 1906: 116, 1907: 260, 1909: 135, 1910a: 88–89, 99–101, 1912: 354, 1913a: 29, 1913b: 253; Hopkins 1914: 127, 134, 1915c: 173–226; Hopping 1922: 128–134; Keen 1958: 95, 159–164; Krivolutskaia 1958: 163; Kurenzov 1941: 71, 180; Lacordaire 1866: 373, 381–382; LeConte 1876: 347, 349–352; LeConte & Horn 1883: 517; Lekander 1962: 431; Lindemann 1876: 148–168, 345; Murayama 1953: 11, 1957: 590, 1963c: 375; Niisima 1911: 1, 1917: 1; Numberg 1954: 6, 11, 68; Nusslin 1898: 281, 1911, 1912; Pfeffer 1940: 107–127, 1946: 113, 1962: 240, 244, 1976: 324; Redtenbacher 1874: 379; Reitter 1869: 154, 1894: 68, 75–76, 1906: 711, 1913a: 92, 1916: 297; Rio Mora 1985: Saalas 1914: 74, 86; Sawamoto 1942: 166; Schedl 1930: 195–199, 1931: 163–168, 1935j: 342–343, 1936h: 59, 1938a: 158–183, 1940a: 348, 1951m: 114, 1952d: 347, 1954b: 37, 1955g: 4, 1956a: 31, 1957d: 77, 1958d: 187–193, 1958k: 144, 1959m: 545, 557, 1960i: 106, 1961i: 224, 1962j: 104, 1963h: 264–265, 1964m: 310–311, 1965f: 3–15, 1969e: 157, 1970e: 93, 1977b: 106, 1981b: 76; Schimitschek 1937: 50–55; Sedlaczek 1935: 155, 157; Spessivtsev 1922: 487, 1929: 297–303, 1931: 89; Stark 1952: 340–349; Stebbing 1914: 551; Swaine 1909: 133–134, 1913: 41–43, 87–92, 1918a: 46, 94; Stresemann et al. 1989: 351; Wachtl 1876: 460; Watson 1928: 629; Wood, S. L. 1961a: 45, 1967d: 37–57, 1972a: 435, 1975b: 391, 1977b: 207, 1982b: 991–1140, 1986a: 95.
- abbreviatus** Schedl 1972g: 63. Holotype ♂?; Brasil, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 63.
- abiegnus** Wood 1964: 67. Holotype ♀; 7 km W Rio Frio, Mexico, Mexico; Wood Collection.
Distribution: North America (Jalisco, Mexico, Oaxaca, Puebla, Tlaxcala in Mexico).
Hosts: *Abies religiosa* limbs.
References: (hb) Atkinson et al. 1986: 36; Burgos & Saucedo 1983: 121; Wood, S. L. 1982b: 1097. (ds) Atkinson & Equihua 1985a: 94; Atkinson et al. 1986: 36; Bright 1981d: 165; Burgos & Saucedo 1983: 121; Wood, S. L. 1964: 67, 1982b: 1097. (tx) Bright 1981d: 165; Schedl 1979c: 8; Wood, S. L. 1964: 67.
- abietinus** Wood 1989: 175. Lectotype, sex?, S Coast (Primorsky) Region, Voroshilovsk district; ISBN, Novosibirsk, designated by Michalsky 1969a: 895, automatic.
- Distribution: Asia (E USSR).
Hosts: *Abies holophylla*.
References: (tx) Wood, S. L. 1989: 178–179.
- abietis** Kurenzov 1941: 179, 234. Lectotype, sex?, S Coast (Primorsky) Region, Voroshilovsk district; ISBN, Novosibirsk, designated by Michalsky 1969a: 895, preoccupied by Blackman 1925.
References: (hb) Karpinski 1949: 129; Krivolutskaia 1956: 838; Kurenzov 1948b: 102, 1950d: 228; Stark 1952: 357. (ds) Kleine 1934a: 158; Krivolutskaia 1956: 838; Pfeffer 1976: 340; Stark 1952: 357. (tx) Krivolutskaia 1956: 838; Kurenzov 1941a: 179, 234, 1948b: 102, 1956: 118; Michalski 1969a: 895; Numberg 1956: 208; Pfeffer 1976: 340; Stark 1952: 357; Wood, S. L. 1989: 179.
- sibiricus** Numberg 1956: 208. Lectotype, sex?, S Coast (Primorsky) Region, Voroshilovsk district; ISBN, Novosibirsk, designated by Michalsky 1969a: 895, automatic, preoccupied by Stark 1952.
References: (ds) Krivolutskaia 1958; Yanovskii & Tegshzhargal 1985: 410. (tx) Michalski 1969b: 569; Numberg 1956: 208; Wood, S. L. 1989: 179.
- ablusus** Bright 1985b: 476. Holotype ♀; Mexico: Pachuca, Edo. Hgo; CNCI, Ottawa.
Distribution: North America (Hidalgo in Mexico).
Hosts: *Zexmenia* sp.
References: (tx) Bright 1985b: 476.
- abnormalis** Bright 1972d: 85. Holotype ♀; Duncans, Trelawny Parish, Jamaica; CNCI, Ottawa.
Figures: Bright 1972d: 19, 59.
Distribution: Antilles Islands (Jamaica).
References: (ds) Bright 1985c: 176. (tx) Bright 1972d: 19, 59, 88, 1985c: 176; McNamara 1977: 197.
- absonus** Blackman 1928b: 35. Holotype ♀; Mineral King, California [USA]; USNM, Washington.
Figures: Bright 1981d: 362.
Distribution: North America (Alberta, British Columbia in Canada/ California, Colorado, Montana, Nevada, New Mexico, Utah in USA).
Hosts: *Abies lasiocarpa*, *Pinus albicaulis*, *P. aristata*, *P. balfouriana*, *P. contorta*, *P. flexilis*, *P. monticola*.
References: (cc) Stephen & Dahlsten 1976b: 292. (hb) Bright & Stark 1973: 107; Chamberlin 1939: 357. (ds) Bright 1981d: 230; Bright & Stark 1982b: 1037; Chamberlin 1939: 357; Evans, D. 1983: 34; Furniss, M. M. & Johnson 1987: 377; Gast et al. 1989: 385; Keen 1929a: 32; Kleine 1934a: 158; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 35; Bright 1977: 513, 1981d: 230, 362; Chamberlin 1939: 357; Evans, D. 1983: 34; Keen 1929a: 32; Wood, S. L. 1982b: 1037.
- demissus** Blackman 1928b: 74. Holotype ♀; Glacier National Park, Montana [USA]; USNM, Washington. Synonymy: Bright 1977: 513.

References: **(hb)** Chamberlin 1939: 377, 1958; **(ds)** Chamberlin 1939: 377, 1958; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1948: 52, 1951a: 128, 1972a: 425. **(tx)** Blackman 1928b: 74; Bright 1977: 513; Chamberlin 1939: 377, 1958: 152; Keen 1929a: 32; Wood, S. L. 1972a: 425.

inyoensis Bright 1971b: 65. Holotype ♀; 2 km S Onion Valley, Inyo Co., California [USA]; CNCI, Ottawa. Synonymy: Bright 1977: 513. References: **(hb)** Bright & Stark 1973: 110. **(ds)** Bright & Stark 1973: 110. **(tx)** Bright 1971b: 65–66, 1977: 513; McNamara 1977: 195.

abstrusus Bright 1976c: 427. Holotype ♀; 40 km W Orizaba, Veracruz, Mexico; CNCI, Ottawa. Distribution: North America (Durango, Veracruz in Mexico).

Hosts: *Pinus ayacahuite*, *P. sp.*

References: **(ds)** Bright 1981d: 232; Wood, S. L. 1982b: 1034. **(tx)** Bright 1976c: 427, 1981d: 232; McNamara 1984: 754; Wood, S. L. 1982b: 1034.

aciculatus Bright 1977: 519. Holotype ♀; San Cristobal, Huehuetenango, Guatemala; Wood Collection.

Figures: Bright 1981d: 347.

Distribution: North America (Guatemala/ Mexico in Mexico).

Hosts: *Pinus leiophylla*, *P. pseudostrobus*.

References: **(hb)** Atkinson et al. 1986: 36; Burgos & Saucedo 1983: 117. **(ds)** Atkinson & Equihua 1985a: 94, 1988: 94; Atkinson et al. 1986: 36; Bright 1981d: 139; Burgos & Saucedo 1983: 117; Wood, S. L. 1982b: 1049. **(tx)** Bright 1977: 519, 1981d: 139, 347; McNamara 1984: 754; Wood, S. L. 1982b: 1049.

acuminatus (Schedl) 1940a: 346 (*Neopityophthorus*). Syntypes ♀; Tuxtepec, Oaxaca and Finca la Florida, Chiapas, Mexico; Schedl Collection in NHMW, Wien, and Dampf Collection.

Distribution: North America (Honduras/ Campeche, Chiapas, Oaxaca, Tabasco in Mexico).

Hosts: Log phloem.

Notes: (1) Bright 1981d: 18 (to *Pityophthorus*).

References: **(ds)** Estrada & Atkinson 1988: 212; Wood, S. L. 1982b: 1134. **(tx)** Bright 1981d: 18; Schedl 1940a: 346–347, 1979c: 11; Wood, S. L. 1982b: 1134.

acutus Blackman 1928b: 134. Holotype ♀; Kaibab N.F., Arizona [USA]; USNM, Washington.

Figures: Bright 1981d: 373.

Distribution: North America (Arizona, Colorado, New Mexico, Utah in USA).

Hosts: *Pinus flexilis*, *P. leiophylla*, *P. ponderosae*, *P. strobiformis*, rare in *Pseudotsuga menziesii*.

References: **(cn)** Blackman 1931c. **(ec)** Blackman 1931c. **(hb)** Blackman 1931c; Bright 1981d: 321; Chamberlin 1939: 398. **(ds)** Bright 1981d: 321; Chamberlin 1939: 398; Keen 1929a: 33; Kleine 1934a: 155; Leng & Mutchler 1933: 52. **(tx)**

Blackman 1928b: 116, 134–136; Bright 1981d: 321, 373; Chamberlin 1939: 398; Keen 1929a: 33; Wood, S. L. 1982b: 111.

africanulus Wood 1992b: 79. Holotype, sex?; Congo: Mayumbe; MRCB, Tervuren, automatic.

Distribution: Africa (Mozambique/ Zaire).

Hosts: *Klainedoxa gabonensis*.

References: **(tx)** Schedl 1962k: 1079; Wood 1992b: 79.

africanus Schedl 1962k: 1079 (*Neodryocotes*).

Holotype, sex?; Congo: Mayumbe; MRCB, Tervuren, preoccupied by Eggers 1927.

References: **(tx)** Schedl 1962k: 1079; Wood, S. L. 1992b: 79.

africanus Eggers 1927a: 184. Holotype, sex?; Kirumba bei Muansa, Ost-Africa; Methner Collection in MNB, Berlin.

Distribution: Africa (Tanzania/ Zaire).

Hosts: *Klainedoxa gabonensis*.

References: **(hb)** Schedl 1962j: 4. **(ds)** Schedl 1962j: 4. **(tx)** Eggers 1927a: 184; Schedl 1953d: 69, 1961e: 143, 1962j: 4, 1962k: 1079, 1979c: 15.

alienus Eichhoff 1872b: 135. Holotype, sex?; Brasilia; Hamburg Museum, lost.

Distribution: South America (Brazil).

References: **(ds)** Blackwelder 1947: 781; Gemminger & Harold 1872: 2688; Hagedorn 1910d: 70; Kleine 1913b: 140, 1914b: 336. **(tx)** Eichhoff 1872b: 135, 1878b: 192; Hagedorn 1910a: 100.

alni Blackman 1942a: 209. Holotype ♀; Mexico, Veracruz, Jalapa; USNM, Washington.

Distribution: North America (Hidalgo, Veracruz in Mexico).

Hosts: *Alnus* sp.

References: **(hb)** Atkinson & Equihua 1985c: 359. **(ds)** Atkinson & Equihua 1985c: 359; Blackwelder 1947: 781; Bright 1981d: 93; Wood, S. L. 1982b: 1027. **(tx)** Blackman 1942a: 209–210; Bright 1981d: 93; McNamara 1984: 754; Wood, S. L. 1982b: 1027.

alnicolens Wood 1977b: 213. Holotype ♂?; Highway 131, 184 km S Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Alnus* sp.

References: **(ds)** Bright 1981d: 94; Wood, S. L. 1982b: 1027. **(tx)** Bright 1981d: 94; Wood, S. L. 1977b: 213, 1982b: 1027.

alpinensis Hopping, G. 1960a: 865. Holotype ♀; Highwood Pass, Alberta [Canada]; CNCI, Ottawa.

Figures: Hopping 1960a: 866 (adult). Kusch 1967: 10. Distribution: North America (Alberta in Canada/ Idaho, Montana in USA).

Hosts: *Larix lyallii*.

References: **(cn)** Baranyay & Stevenson 1966: 55; Hopping 1960b: 258. **(hb)** Cast et al. 1989: 384. **(ds)** Baranyay & Stevenson 1966: 55; Bright

- 1976d: 180, 1981d: 132; Cast et al. 1989: 384; Hopping 1960b: 258; Kusch 1967: Wood, S. L. 1982b: 1054. (tx) Bright 1977: 516, 1981d: 132; Hopping 1960a: 865–867; Kusch 1967: 10; Wood, S. L. 1964: 59, 1982b: 1054.
- alvarengai** Schedl 1972g: 63. Holotype ♂; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 63, 1979c: 17.
- amiculus** Wood 1975b: 398. Holotype ♂; Guapiles, Limon, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica/ Veracruz in Mexico).
Hosts: Liana.
References: (ds) Bright 1981d: 97; Wood, S. L. 1982b: 1030. (tx) Bright 1981d: 97; McNamara 1954: 754; Wood, S. L. 1975b: 398, 1982b: 1030.
- amoenus** Blandford 1904: 237. Holotype ♀; Duenas, Guatemala; BMNH, London.
Distribution: North America (Guatemala).
References: (ds) Blackwelder 1947: 781; Bright 1981d: 48; Hagedorn 1910d: 70; Kleine 1913b: 140, 1914b: 362; Wood, S. L. 1982b: 1062. (tx) Blackman 1942a: 218; Blandford 1904: 237; Bright 1981d: 48; Hagedorn 1910a: 100; Schedl 1955g: 17; Wood, S. L. 1982b: 1062.
- amplus** (Blackman) 1928b: 18 (*Myeloborus*). Holotype ♀; Kaibab N.F., Arizona [USA]; USNM, Washington.
Distribution: North America (Arizona, S Nevada, New Mexico in USA).
Hosts: *Pinus flexilis*, *P. ponderosa*, *P. strobiformis*.
Notes: (1) Bright 1977: 511 (to *Pityophthorus*).
References: (cn) Doane et al. 1936. (hb) Chamberlin 1939: 339; Doane et al. 1936. (ds) Bright 1981d: 203; Chamberlin 1939: 339; Furniss, R. L. & Carolin 1977: 398; Keen 1929a: 31; Leng & Mutchler 1933: 53; Wood, S. L. 1982b: 1044. (tx) Blackman 1928b: 3–6, 18; Bright 1977: 511, 1981d: 203; Chamberlin 1939: 339; Keen 1929a: 31; McNamara 1954: 753; Wood, S. L. 1982b: 1044.
- angustus** Blackman 1928b: 83. Holotype ♀; Cranberry Lake, New York [USA]; USNM, Washington.
Distribution: North America (New Brunswick, Nova Scotia, Ontario in Canada/ New York, North Carolina in USA).
Hosts: *Abies balsamea*, *Picea rubens*, *Pinus strobus*.
Notes: (3) Bright 1981d: 182 (This name was treated as a synonym of *balsameus*).
References: (cn) Blackman 1950. (hb) Blackman 1950. (ds) Blackman 1950; Bright 1976d: 185; Chamberlin 1939: 380; Kleine 1934a: 158; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 83–84; Chamberlin 1939: 380; de Ruelle 1970: 107; Wood, S. L. 1982b: 1077.
- annectens** LeConte 1878b: 622. Lectotype ♀; Tampa, Florida [USA]; MCZ, Cambridge, designated by Bright 1976: 185.
Figures: Blackman 1922b: pl. 9, fig. 47, Bright 1981d: 371.
Distribution: Antilles Islands (Andros Island/ Bahamas Islands), North America (Belize/ Chiapas (?), Durango, Hidalgo, Jalisco, Mexico, Michoacan in Mexico/ Alabama, Arizona, Arkansas, Florida, Georgia, Illinois, Louisiana, Mississippi, New Mexico, North Carolina, South Carolina, E Texas, Virginia, West Virginia in USA).
Hosts: *Pinus ayacahuite*, *P. caribaea*, *P. edulis*, *P. elliotii*, *P. engelmannii*, *P. lawsoni*, *P. leiophylla*, *P. palustris*, *P. ponderosa*, *P. resinosa*, *P. taeda*, *P. spp.*
Notes: (3) Blackman 1928b: 138 (redescribed).
References: (bv) Dixon & Payne 1979b; Jacobson 1972; Turnbow & Franklin 1980; Vite 1965: 267; Vite, Gara, & von Scheller 1964: 461–470; Vite & Pitman 1967: 683–701. (cn) Blackman 1950; Packard 1890: 715; Swaine 1918a: 104. (ec) Dixon & Payne 1979b; Felt 1906: 751; Kinn 1983a; Schwarz 1889a: 164. (hb) Atkinson et al. 1986: 36; Baker, W. L. 1972: 255; Beal & Massey 1945: 136; Blackman 1922b: 109–110, 1950; Bright 1981d: 293; Chamberlin 1939: 400; Chittenden 1890; Deyrup & Atkinson 1987a: 66; Felt 1906: 751; Ostmark 1968; Packard 1890: 715; Swaine 1918a: 104. (ds) Atkinson et al. 1986: 36, 1991: 162; Baker, W. L. 1972: 255; Beal & Massey 1945: 136; Blackman 1922b: 109–110, 1950; Blatchley & Leng 1916: 631; Bright 1981d: 293; Chamberlin 1925, 1939: 400; Chittenden 1890; Deyrup & Atkinson 1987a: 66; Drooz 1985: 364; Hagedorn 1910d: 70; Henshaw 1882: 268, 1885: 147; Hubbard & Schwarz 1878b: 622; Kirk 1970; Kleine 1913b: 140, 1914b: 383, 1934a: 158; Leng 1920: 341; Ostmark 1968; Rio Mora 1985: 14; Schwarz 1889a: 164, 1891c: 357; Swaine 1909: 133; Turnbow & Franklin 1980; Weber, B. C. & McPherson 1991: 54; Ulke 1902: 56; Wood, S. L. 1982b: 1101. (tx) Beal & Massey 1945: 136; Blackman 1922b: 109–110, 1928b: 117, 138; Blatchley & Leng 1916: 631; Bright 1976b: 185, 1981d: 293, 371; Chamberlin 1939: 400; Hagedorn 1910a: 100; LeConte 1878b: 622; Schedl 1955g: 25; Schwarz 1889: 163–164, 1891a: 167, 1891c: 357; Swaine 1909: 133–134, 1918a: 104; Wood, S. L. 1977d: 514, 1982b: 1101.
- citus** Blackman 1928b: 137. Holotype ♀; Chiricahua Mts., Arizona [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.
References: (hb) Chamberlin 1939: 399. (ds) Chamberlin 1939: 399; Keen 1929a: 34; Kleine 1934a: 159; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 137; Chamberlin 1939: 399; Keen 1929a: 34; Wood, S. L. 1977d: 514.
- anthracinus** Bright 1976c: 428. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico; Wood Collection.

Distribution: North America (Nuevo Leon, Tlaxcala in Mexico).

Hosts: *Abies religiosa*.

References: (hb) Bright 1981d: 317. (ds) Bright 1981d: 317; Wood, S. L. 1982b: 1087. (tx) Bright 1976c: 428, 1981d: 317; McNamara 1984: 754; Wood, S. L. 1982b: 1087.

auticus Schedl 1976a: 66. Holotype ♂; Brasilien, Rio Negro; Schedl Collection in NHMW, Wien. Figures: Pedrosa-Macedo & Schonherr 1985: 52. Distribution: South America (Brazil).

Hosts: *Araucaria angustifolia*, *Pinus elliottii*.

References: (ds) Pedrosa-Macedo & Schonherr 1985: 52; Schonherr & Pedrosa-Macedo 1981: 53. (tx) Pedrosa-Macedo & Schonherr 1985: 52; Schedl 1976a: 66.

antillicus Bright 1981c: 162. Holotype ♀; Jarabacoa, Dominican Republic, Hopk US 32062-G; USNM, Washington.

Distribution: Antilles Islands (Dominican Republic in Hispanola).

Hosts: *Pinus occidentalis*.

References: (bv) Haack et al. 1989. (cc) Haack et al. 1989. (hb) Haack et al. 1989. (ds) Bright 1985c: 176. (tx) Bright 1981c: 162, 1985c: 176.

apachae Bright 1977: 520. Holotype ♀; Rustler's Park, Chiricahua Mts., Arizona [USA]; Wood Collection.

Distribution: North America (Arizona in USA).

Hosts: *Abies concolor*.

References: (ds) Bright 1981d: 241; Wood, S. L. 1982b: 1070. (tx) Bright 1977: 520, 1981d: 241; McNamara 1984: 754; Wood, S. L. 1982b: 1070.

apicenotatus Schedl 1976a: 66. Holotype, sex?; Cachimbo, BA, Brasil; Schedl Collection in NMHW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 66.

apicipennis Schedl 1976a: 67. Holotype ♂; Serra do Caraca, Minas Gerais, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 67.

apiculatus Schedl 1937h: 167. Holotype ♀; Nova Teutonia [Brazil]; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 781; Schedl 1966f: 86. (tx) Schedl 1937h: 167-168, 1938a: 183, 1979c: 23.

aquilus Blackman 1928b: 33. Holotype ♀; Kaibab N.F., Arizona [USA]; USNM, Washington.

Figures: Bright 1981d: 361.

Distribution: North America (Alberta, British Columbia, Saskatchewan in Canada/Arizona, California, Colorado, Montana, New Mexico, South Dakota, Utah, Wyoming in USA).

Hosts: *Picea engelmannii* (rare), *Pinus albicaulis*, *P. aristata*, *P. contorta*, *P. flexilis*, *P. ponderosa*.

References: (cn) Loomis, Hofacker, & Tucker 1986; Raimo 1985. (hb) Bright 1981d: 224; Chamberlin 1939: 358; Furniss, M. M. & Johnson 1987: 377. (ds) Bright 1981d: 224; Chamberlin 1939: 358; Evans, D., Lowe, & Hunt 1978; Furniss, M. M. & Johnson 1987: 377; Gast et al. 1989: 385; Keen 1929a: 32; Kleine 1934a: 158; Leng & Mutchler 1933: 52; Wood, S. L. 1948: 52, 1951a: 128, 1982b: 1064. (tx) Blackman 1928b: 30-34; Bright 1977: 514, 1981d: 224, 361; Chamberlin 1939: 358; Keen 1929a: 32; de Ruette 1970: 105; Wood, S. L. 1977b: 209, 1982b: 1064.

caelator Blackman 1928b: 78. Holotype ♀; Black Hills, South Dakota [USA]; USNM, Washington. Synonymy: Bright 1977: 514.

References: (cn) Blackman 1931c. (cc) Blackman 1931c. (hb) Blackman 1931c; Chamberlin 1939: 378. (ds) Chamberlin 1939: 378; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 62, 78-79; Bright 1977: 514; Chamberlin 1939: 378; Keen 1929a: 32; de Ruette 1970: 108.

aristatae Bright 1964: 166. Holotype ♀; Crooked Creek, White Mts., Mono County, California [USA]; CAS, San Francisco. Synonymy: Bright 1971b: 67.

References: (tx) Bright 1964: 166, 168, 1971b: 67, 1977: 514.

arakii (Sawamoto) 1942: 167 (*Cladoborus*). Syn-types ♂ ♀; Hokkaido (Kutchan); Entomological Institute, Imperial Hokkaido University and Sawamoto Collection.

Distribution: Asia (Japan).

Hosts: *Picea excelsa*.

Notes: (1) Bright 1977: 511 (to *Pityophthorus*).

References: (ds) Murayama 1954b: 201. (tx) Bright 1977: 511; Murayama 1954b: 201, 1963c: 391; Nobuchi 1981c: 29; Sawamoto 1942: 167; Schedl 1959r: 42.

arcanus Bright 1976c: 429. Holotype ♀; Bear Canyon, Santa Catalina Mts., Santa Cruz Co., Arizona [USA]; CNCI, Ottawa.

Figures: Bright 1981d: 335.

Distribution: North America (Arizona, Nevada, W Texas in USA).

Hosts: *Pinus cembroides*, *P. monophylla*, *P. ponderosa*, 1 series from *Pseudotsuga menziesii*.

References: (ds) Bright 1981d: 35; Wood, S. L. 1982b: 1110. (tx) Bright 1976c: 429, 1981d: 35, 335; McNamara 1984: 754; Wood, S. L. 1982b: 1110.

arceuthobii Wood 1971: 48. Holotype ♀; 96 km W Durango, Durango, Mexico; Wood Collection.

Distribution: North America (Durango, Mexico in Mexico).

Hosts: *Arceuthobium globosum* (parasite on *Pinus montezumae*).

- References: **(ds)** Atkinson & Equihua 1985a: 96; Bright 1981d: 242; Wood, S. L. 1982b: 1046. **(tx)** Bright 1981d: 242; Wood, S. L. 1971: 48, 1982b: 1046.
- argentinensis** Eggers 1951: 150. Lectotype, sex?; Argentinien (Salta); USNM, Washington.
Distribution: South America (Argentina/ Brazil).
References: **(hb)** Viana 1964: 125. **(ds)** Schedl 1967d: 2; Viana 1964: 125. **(tx)** Anderson, W. H. & Anderson 1971: 4; Eggers 1951: 150; Schedl 1938i: 28, 1951h: 285, 1958f: 35, 1979c: 25.
- argentinae** (Schedl) 1958f: 44 (*Breviophthorus*).
Syntypes, sex?; Argentinien: Misiones, Dep. Concep., Sta. Maria; Schedl Collection in NHMW, Wien and Viana Collection.
Distribution: South America (Argentina).
Notes: (1) Schedl 1979c: 25 (citation of holotype invalid).
References: **(tx)** Schedl 1958f: 44, 1970e: 93, 1979c: 25.
- ascendens** Schedl 1972g: 64. Holotype ♀; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: **(tx)** Schedl 1972g: 64, 1979c: 27.
- ashanti** Schedl 1972e: 291. Holotype ♀; Ghana, Ashanti Region, Kwadaso, 320 m, N 6 42, W 1 39; NHMB, Budapest.
Distribution: Africa (Ghana).
References: **(tx)** Schedl 1972e: 291, 1979c: 28.
- assitus** Wood 1977b: 214. Holotype ♀; Lagos des Colores, Chiapas, Mexico, 1300 m; CNCI, Ottawa.
Distribution: North America (Chiapas, Oaxaca in Mexico).
References: **(ds)** Bright 1981d: 50; Wood, S. L. 1982b: 1129. **(tx)** Bright 1981d: 50; Wood, S. L. 1977b: 214, 1982b: 1129.
- aterrimus** Eggers 1931b: 31. Holotype, sex?; Brasilien (Sao Paulo); Eggers Collection [not mentioned by Anderson & Anderson 1971 or Schedl 1979c], not in NHMW, Wien.
Distribution: South America (Brazil).
References: **(ds)** Blackwelder 1947: 781. **(tx)** Eggers 1931b: 31.
- atkinsoni** Bright 1985a: 467. Holotype ♀; Mexico: Cardonal (Cerca Ixmiquilpan), Edo. Hidalgo; CNCI, Ottawa.
Distribution: North America (Hidalgo in Mexico).
Hosts: *Flourensia resinosa*.
References: **(tx)** Bright 1985a: 467.
- atomus** Wood 1964: 61. Holotype ♀; Veracruz, Veracruz, Mexico; Wood Collection.
Distribution: North America (Hidalgo, Jalisco, Michoacan, Oaxaca, Veracruz in Mexico).
Hosts: Shrub.
References: **(ds)** Bright 1981d: 44, 1986b: 682; Wood, S. L. 1982b: 1137. **(tx)** Bright 1981d: 44;
- McNamara 1977: 197; Wood, S. L. 1964: 61, 1982b: 1137.
- attenuatus** Blackman 1942a: 222. Holotype ♀; Mexico; USNM, Washington.
Figures: Bright 1981d: 336.
Distribution: North America (El Salvador/ Guatemala/ Hidalgo, Jalisco, Michoacan in Mexico).
Hosts: Compositae shrub.
References: **(hb)** Atkinson et al. 1986: 36; Burgos & Saucedo 1983: 124. **(ds)** Atkinson et al. 1986: 36; Blackwelder 1947: 781; Bright 1981d: 45; Burgos & Saucedo 1983: 124; Schedl 1977e: 42; Wood, S. L. 1982b: 1136. **(tx)** Blackman 1942a: 222; Bright 1981d: 45, 336; McNamara 1984: 754; Wood, S. L. 1978b: 397, 1982b: 1136.
- pusillus** Wood 1964: 62. Holotype ♀; 14 km S Zimapan, Hidalgo, Mexico; Wood Collection.
Synonymy: Wood 1978b: 397.
References: **(tx)** McNamara 1977: 195; Wood, S. L. 1964: 62, 1978b: 397.
- aztecus** Bright 1977: 520. Holotype ♀; Km 50 Mexico-Puebla [Highway], Mexico; USNM, Washington.
Figures: Bright 1981d: 340.
Distribution: North America (Coahuila, Mexico, Veracruz in Mexico).
Hosts: *Pinus* sp. branches.
References: **(hb)** Atkinson et al. 1986: 37; Burgos & Saucedo 1983: 115. **(ds)** Atkinson & Equihua 1985a: 92, 1988: 94; Atkinson et al. 1986: 37; Bright 1981d: 90; Burgos & Saucedo 1983: 115; Wood, S. L. 1982b: 1046. **(tx)** Bright 1977: 520, 1981d: 90, 340; McNamara 1984: 754; Wood, S. L. 1982b: 1046.
- balcanicus** Pfeffer 1940b: 123. Syntypes, sex?; Bulgaria near Sofia; Serbia, Demir Kapija, Kozuf Planina Boania; Zavidovic, Hercegovina, Albania; Ilgora; Pfeffer Collection in Prague Museum.
Distribution: Europe (Albania/ Bulgaria/ Yugoslavia).
Hosts: *Pinus heldreichii*, *P. nigra*.
References: **(bv)** Grune 1979: 121. **(ds)** Grune 1979: 121; Pfeffer 1976: 340, 1984: 277; Postner 1974: 437. **(tx)** Grune 1979: 121; Pfeffer 1940b: 109, 113, 123, 1955a: 211, 1964m: 312, 1976: 340; Postner 1974: 437.
- balsameus** Blackman 1922c: 119. Holotype ♀; Orono, Maine [USA]; USNM, Washington.
Figures: Blackman 1922c: figs. 20–21, Bright 1981d: 355.
Distribution: North America (New Brunswick, Nova Scotia in Canada/ Maine, Michigan, Minnesota, New Hampshire, New York, North Carolina, West Virginia, Wisconsin in USA).
Hosts: *Abies balsamea*, *Picea glauca*, *P. sp.*, *Pinus banksiana*, *P. resinosa*, *P. strobus*.
References: **(en)** Blackman 1950. **(cc)** Bushing 1965: 464; Marsh 1979: 158; Matthews 1970. **(hb)** Baker, W. L. 1972: 255; Blackman 1950; Chamberlin 1939: 381. **(ds)** Baker, W. L. 1972: 255;

- Blackman 1950; Bright 1976d: 185, 1981d: 181; Chamberlin 1939: 381; Drooz 1985: 364; Kleine 1934a: 158; Leng & Mutchler 1927: 52; Wood, S. L. 1982b: 1076. **(tx)** Blackman 1922c: 119–120, 1928b: 63, 88; Bright 1981d: 181, 355; Chamberlin 1939: 381; Wood, S. L. 1977d: 514, 1982b: 1076. *patchi* Blackman 1922c: 120. Holotype ♀; Orono, Maine [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.
References: **(cn)** Blackman 1950. **(hb)** Baker, W. L. 1972: 255; Blackman 1950; Chamberlin 1939: 379. **(ds)** Baker, W. L. 1972: 255; Blackman 1950; Bright 1976d: 187; Chamberlin 1925, 1939: 379; Kleine 1934a: 161; Leng & Mutchler 1927: 52; Proctor 1946: 208. **(tx)** Blackman 1922c: 120; Chamberlin 1939: 379; de Ruelle 1970: 110; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1977d: 514.
- barberi** Blackman 1928b: 112. Holotype, sex?; New Mexico: Las Vegas H.S. [Hot Springs] [USA]; USNM, Washington.
Figures: Bright 1981d: 339.
Distribution: North America (Durango, Nuevo Leon in Mexico/ Arizona, Colorado, New Mexico, W Texas, Utah in USA).
Hosts: *Pinus cembroides*, *P. edulis*, *P. leiophylla*, *P. ponderosa*.
References: **(hb)** Bright 1981d: 69; Chamberlin 1939: 389. **(ds)** Bright 1981d: 69; Chamberlin 1939: 389; Keen 1929a: 33; Kleine 1934a: 158; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1117. **(tx)** Blackman 1928b: 91, 112–113; Bright 1981d: 69, 339; Chamberlin 1939: 389; Keen 1929a: 33; de Ruelle 1970: 108; Schedl 1937h: 167; Wood, S. L. 1982b: 1117.
- barbifer** Schedl 1964j: 46. Holotype ♂; Ghana: Kintampo; BMNH, London.
Distribution: Africa (Ghana).
Hosts: *Daniellia oliveri*.
References: **(hb)** Roberts 1969: 126. **(ds)** Schedl 1962j: 5–6, 1964j: 46, 1972e: 282. **(tx)** Schedl 1962j: 5, 1964j: 46, 1979c: 35.
- bassetti** Blackman 1920b: 1. Lectotype ♂; Pitkin, Colorado [USA]; USNM, Washington, designated by Bright 1976b: 183.
Figures: Bright 1981d: 319, 373.
Distribution: North America (Alaska/ Northwest Territories, Yukon in Canada/ Chihuahua in Mexico/ Colorado, Utah in USA).
Hosts: *Picea chiluhuanana*, *P. engelmannii*, *P. glauca*, *P. pungens*.
Notes: (3) Blackman 1928b: 136 (redescribed).
References: **(ec)** Werner & Holsten 1984. **(hb)** Blackman 1920b: 1; Chamberlin 1939: 399, 1958. **(ds)** Bright 1981d: 318; Chamberlin 1925, 1939: 399, 1958; Elias 1982d; Keen 1929a: 33; Kleine 1934a: 155; Leng & Mutchler 1927: 52; Werner & Holsten 1984; Wood, S. L. 1948: 55, 1951a: 128, 1972a: 427, 1982b: 1109. **(tx)** Blackman 1920b: 1, 4, 1928b: 117, 136; Bright 1976b: 183, 1981d: 318–319, 373; Chamberlin 1939: 399, 1958; Keen 1929a: 33; de Ruelle 1970: 108; Wood, S. L. 1951a: 128, 1972a: 427, 1982b: 1109.
- bioealis** Blackman 1922c: 122. Holotype ♀; Brunswick, Maine [USA]; USNM, Washington.
Distribution: North America (Nova Scotia, Ontario in Canada/ Maine, New York in USA).
Hosts: *Picea glauca*, *P. mariana*, *P. rubens*, *Pinus strobus*.
References: **(cn)** Baker, W. L. 1972: 255; Blackman 1950. **(hb)** Baker, W. L. 1972: 255; Blackman 1950; Chamberlin 1939: 380. **(ds)** Baker, W. L. 1972: 255; Blackman 1950; Bright 1976d: 255, 1981d: 178; Chamberlin 1925, 1939: 380; Drooz 1985: 364; Kleine 1934a: 158; Leng & Mutchler 1927: 52; Proctor 1946: 208; Wood, S. L. 1982b: 1074. **(tx)** Blackman 1922c: 122–123, 1928b: 63, 85; Bright 1981d: 178; Chamberlin 1939: 380; Wood, S. L. 1982b: 1074.
- blackmani** (Schedl) 1935h: 344 (*Conophthoracranulus*). Holotype ♀; Costa Rica, Turrialba, 800 m; Schedl Collection in NIMW, Wien.
Distribution: North America (Costa Rica).
Notes: (1) Wood 1986a: 98 (to *Pityophthorus*).
References: **(ds)** Blackwelder 1947: 781. **(tx)** Schedl 1935h: 344, 1938a: 162, 1979c: 42; Wood, S. L. 1982b: 963, 1986a: 98.
- blandulus** Schedl 1955g: 19. Lectotype ♀; Guatemala, Las Trojades; Schedl Collection in NIMW, Wien, designated by Bright 1976b: 186.
Figures: Bright 1981d: 363.
Distribution: North America (Guatemala/ Chiapas, Mexico in Mexico).
Hosts: *Pinus hartwegii*, *P. ochoterrenai*, *P. oocarpa*, *P. rudis*.
References: **(hb)** Schwerdtfeger 1957c: 494. **(ds)** Atkinson & Equihua 1985c: 360, 1988: 94; Bright 1981d: 244; Schedl 1963c: 158; Wood, S. L. 1982b: 1068. **(tx)** Bright 1976b: 186, 1981d: 244, 363; Schedl 1955g: 17, 19, 1979c: 42; Wood, S. L. 1982b: 1068.
- blandulus** Blackman 1928b: 107. Holotype ♀; Argus Mts., California [USA]; USNM, Washington.
Figures: Bright 1981d: 363.
Distribution: North America (Arizona, California, Colorado, Nevada, New Mexico, Utah in USA).
Hosts: *Pinus monophylla*, *P. edulis*, 1 pair from *P. ponderosa*.
References: **(hb)** Bright & Stark 1973: 114; Chamberlin 1939: 387. **(ds)** Bright 1981d: 245; Bright & Stark 1973: 114; Chamberlin 1939: 387; Furniss, M. M. & Johnson 1987: 377; Keen 1929a: 23; Kleine 1934a: 158; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 107–108; Bright 1977: 513, 1981d: 245, 363; Chamberlin 1939: 387; Keen 1929a: 33; Wood, S. L. 1982b: 1066.
- singularis** Bright 1966b: 300. Holotype ♀; 19 km W Lone Pine, Inyo Co., California [USA];

CAS, San Francisco. Synonymy: Bright 1977: 513.

References: (hb) Bright & Stark 1973: 111. (ds) Bright & Stark 1973: 111; (tx) Bright 1966b: 300, 1977: 513; de Ruette 1970: 111.

bolivianus Eggers 1943a: 359. Holotype, sex?; Bolivia (Cochabamba); MNHN, Paris. Distribution: South America (Bolivia). References: (tx) Eggers 1943a: 359; Schedl 1951m: 73, 1979c: 43.

borrichiae Wood 1964: 60. Holotype ♀; Key Largo, Florida [USA]; Wood Collection. Distribution: North America (S Florida). Hosts: *Borrichia arborescens*, *B. frutescens*. References: (ds) Bright 1981d: 61; Wood, S. L. 1982b: 1130. (tx) Bright 1981d: 61; McNamara 1977: 197; Wood, S. L. 1964: 60, 1982b: 1130.

boycei Swaine 1925b: 192. Holotype ♂; Cisco, Placer Co., California [USA]; CNCI, Ottawa. Figures: Bright 1981d: 355, Ives & Wong 1985: 54. Distribution: North America (Alberta, British Columbia in Canada/California, Colorado, Idaho, Montana, Oregon, South Dakota, Wyoming in USA).

Hosts: *Pinus aristata*, *P. contorta*, *P. flexilis*, *P. monticola*, *P. ponderosa*.

Notes: (3) Blackman 1928b: 26 (redescribed).

References: (cn) Doane et al. 1936; Evans, D. 1982; Ives & Wong 1985: 55; Lindgren 1980a: 66. (hb) Bright & Stark 1973: 97; Chamberlin 1939: 342, 1958: 145; Doane et al. 1936; Evans, D. 1982; Ives & Wong 1985: 55; Lindgren 1980a: 66. (ds) Bright 1976d: 168, 1980d: 204; Bright & Stark 1973: 97; Chamberlin 1939: 342, 1958: 145; Evans, D. 1983: 35; Evans, D., Lowe, & Hunt 1975; Furniss, R. L. & Carolin 1977: 395; Gast et al. 1989: 355; Keen 1929a: 31; Kleine 1934a: 163; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 424, 1982b: 1043. (tx) Blackman 1928b: 16–17, 26–27; Bright 1967b: 678, 1976c: 426, 1976d: 168, 1981d: 204, 358; Chamberlin 1939: 342, 1958: 145; Evans, D. 1982, 1983: 35; Ives & Wong 1985: 54; Keen 1929a: 31; de Ruette 1970: 108; Swaine 1925b: 192–193; Wood, S. L. 1972a: 424, 1977c: 357, 1982b: 1043.

catulus Blackman 1928b: 21. Holotype ♀; Clark's Fork, Idaho [USA]; USNM, Washington. Synonymy: Bright 1976b: 186, 1976c: 426.

References: (hb) Chamberlin 1939: 341, 1958. (ds) Chamberlin 1939: 341, 1958: 144–145; Keen 1929a: 31; Kleine 1934a: 163; Leng & Mutchler 1933: 53; Wood, S. L. 1972a: 424. (tx) Blackman 1928b: 17, 21–22; Bright 1976b: 186, 1976c: 426; Chamberlin 1939: 341, 1958: 144–145; Keen 1929a: 31; de Ruette 1970: 104; Wood, S. L. 1972a: 424.

iniquus Blackman 1928b: 27. Holotype ♀; Keystone, Wyoming [USA]; USNM, Washington. Synonymy: Bright 1976c: 426.

References: (cn) Doane et al. 1936. (hb) Chamberlin 1939: 343; Doane et al. 1936. (ds) Chamberlin 1939: 343; Keen 1929a: 31; Kleine 1934a: 163; Leng & Mutchler 1933: 53. (tx) Blackman 1928b: 27; Bright 1976c: 426; Chamberlin 1939: 343; Keen 1929a: 31.

stouxcensis Bright 1977: 439. Holotype ♀; Black Hills, South Dakota [USA]; CNCI, Ottawa. Synonymy: Wood 1977c: 357.

References: (tx) Bright 1977: 439; McNamara 1984: 757; Wood, S. L. 1977d: 357.

brasiliensis (Schedl) 1938a: 177 (*Breviophthorus*). Holotype, sex?; Brasilien, Sao Paulo; Schedl Collection in NHMW, Wien.

Figures: Schedl 1938a: 161, 164.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 781. (tx) Schedl 1938a: 161, 164, 177, 1979c: 44.

bravoi Bright 1986b: 679. Holotype ♀; Mexico: Carr. Mex-Popo., Km 85; CNCI, Ottawa.

Distribution: North America (Mexico in Mexico).

References: (tx) Bright 1986b: 679.

brevicomatus Bright 1976c: 430. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico; CNCI, Ottawa.

Distribution: North America (Nuevo Leon in Mexico).

Hosts: *Pinus strobiformis*.

References: (ds) Bright 1981d: 145; Wood, S. L. 1982b: 1094. (tx) Bright 1976c: 430, 1981d: 145; McNamara 1984: 754; Wood, S. L. 1982b: 1094.

brevis Blackman 1928b: 81. Holotype ♀; Kaibab National Forest, Arizona [USA]; USNM, Washington.

Figures: Bright 1981d: 364.

Distribution: North America (Durango in Mexico/Arizona, Colorado, New Mexico, W Texas in USA).

Hosts: *Pinus ayacalhuite*, *P. edulis*, *P. ponderosa*, *P. strobiformis*.

References: (cn) Blackman 1931c. (ce) Blackman 1931c. (hb) Blackman 1931c; Chamberlin 1939: 379. (ds) Atkinson & Equilua 1985c: 360, 1988: 94; Bright 1981d: 247; Chamberlin 1939: 379; Keen 1929a: 32; Kleine 1934a: 158; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1065. (tx) Blackman 1928b: 81; Bright 1981d: 247; Chamberlin 1939: 379; Keen 1929a: 32; de Ruette 1970: 108; Wood, S. L. 1982b: 1065.

brevisetosus Eggers 1933b: 3, 7. Holotype, sex?; Venezuela (Caracas); MNHN, Paris.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 781. (tx) Eggers 1933b: 3, 7–8.

brighti Wood 1989: 178. Holotype ♀; Amecameca, Mexico, Mexico; USNM, Washington.

Distribution: North America (Hidalgo, Mexico in Mexico).

Hosts: *Abies religiosa*.

References: **(tx)** Wood, S. L. 1989: 178.

blackmani Bright 1977: 521. Holotype ♀; Amecameca, Mexico, Mexico; USNM, Washington, preoccupied by Schedl 1935.

References: **(ds)** Atkinson & Equilma 1955a: 96; Bright 1981d: 243. **(tx)** Bright 1977: 521, 1981d: 243; McNamara 1984: 754; Wood, S. L. 1989: 178.

briscoei Blackman 1922c: 123. Holotype ♀; Brunswick, Maine [USA]; USNM, Washington.

Distribution: North America (New Brunswick, Ontario, Quebec in Canada/ Maine, Minnesota, New Hampshire, New York, Pennsylvania in USA). Hosts: *Picea mariana*, *P. rubens*, *Pinus banksiana*, *P. strobus*.

References: **(cn)** Blackman 1950. **(hb)** Blackman 1950; Chamberlin 1939: 380. **(ds)** Blackman 1950; Bright 1976d: 185, 1981d: 184; Chamberlin 1925, 1939: 380; Kleine 1934a: 158; Leng & Mutchler 1927: 52; Wood, S. L. 1982b: 1076. **(tx)** Blackman 1922c: 123–124, 1928b: 63, 84–85; Bright 1981d: 184; Chamberlin 1939: 380; Wood, S. L. 1977d: 514, 1982b: 1076.

mundus Blackman 1928b: 86. Holotype ♀; Littleton, New Hampshire [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.

References: **(cn)** Blackman 1950. **(hb)** Blackman 1950; Chamberlin 1939: 380. **(ds)** Blackman 1950; Bright 1976d: 186; Chamberlin 1939: 380; Dodge 1938; Kleine 1934a: 160; Knoll 1932: 66; Leng & Mutchler 1933: 52; Proctor 1946: 208. **(tx)** Blackman 1928b: 86–87; Chamberlin 1939: 380; Dodge 1938: 43–45; McNamara 1984: 757; Wood, S. L. 1977d: 514.

burserae Wood 1976a: 362. Holotype ♀; Atencuque, Jalisco, Mexico, 1000 m; Wood Collection. Distribution: North America (Jalisco).

Hosts: *Bursera* sp.

References: **(ds)** Bright 1981d: 76; Wood, S. L. 1982b: 1127. **(tx)** Bright 1981d: 76; Wood, S. L. 1976a: 362, 1982b: 1127.

busseae Schedl 1962h: 69. Holotype ♂♀; Ghana, Bobiri Forest Reserve; BMNH, London.

Distribution: Africa (Ghana).

Hosts: *Bussea occidentalis*.

References: **(hb)** Browne 1963a: 236. **(ds)** Browne 1963a: 236; Schedl 1962h: 59, 1962j: 6, 1967e: 214, 1972e: 282. **(tx)** Schedl 1962h: 69, 1962j: 6, 1962k: 1080, 1979c: 49.

buyssoni Reitter 1901a: 101. Syntypes, sex?; Europe; NHMB, Budapest.

Figures: Chararas 1962c: 360.

Distribution: Europe (Bulgaria/ Corsica/ Czechoslovakia/ S France/ Greece/ Italy/ Spain/ Yugoslavia). Hosts: *Pinus nigra*, *P. sylvestris*, rare in *Larix europaea*.

References: **(bv)** Grune 1979: 121. **(cn)** Grandi 1951; Joly 1976. **(ec)** Kleine 1905c: 213; Pfeffer 1960: 346. **(hb)** Chararas 1982c: 363; Grandi

1951; Joly 1976; Pfeffer 1941b: 3; Spessivtsev 1921b: 221. **(ds)** Andras & Schaefer 1957; Barthe 1896; Buresh & Lazarov 1956; Chararas 1962c: 363; Grandi 1979: 121; Hagedorn 1910d: 70; Joly 1976; Karpinski 1931: 30; Kleine 1913b: 140, 1934a: 159; Linder 1953: 71; Maire et al. 1927: 1576; Pfeffer 1947e: 17, 1960: 346, 1976: 340, 1984: 227, 1989a: 65; Pittioni 1943: 176; Postner 1974: 437; Sainte-Claire 1905: 157; Sainte-Claire & Mequignon 1938: 447; Tredl 1907: 14. **(tx)** Balachowsky 1949a: 238; Chararas 1962c: 360; Eggers 1914: 189, 1915b: 13–14, 1923a: 218, 1933i: 7; Grune 1979: 121; Hagedorn 1910a: 100; Joly 1976; Peyerimhoff 1930: 259; Pfeffer 1927: 111–112, 1940b: 109, 113, 121, 1941b: 3, 1947e: 17, 1955a: 219, 1976: 340; Postner 1974: 437; Reitter 1901a: 101, 1913a: 94; Schedl 1934f: 1643, 1960b: 164, 1964m: 312; Spessivtsev 1922: 221, 1926: 50; Wichmann 1913d: 144.

senex Wichmann 1913d: 143. Holotype ♀?; Bistolas, Htes Alpes; not given. Synonymy: Balachowsky 1949a: 238.

References: **(hb)** Spessivtsev 1921b: 221. **(ds)** Karpinski 1931: 30; Kleine 1913b: 141; Sainte-Claire & Mequignon 1938: 448. **(tx)** Balachowsky 1949a: 238–239; Eggers 1915b; Pfeffer 1927: 111–112, 1940b: 109, 113, 122; Schedl 1934f: 1643, 1964m: 312; Spessivtsev 1922: 221, 1926: 50; Wichmann 1913d: 143–144.

angeri Pfeffer 1927: 111. Syntypes, sex?; Vizzavona, Venaco (Korsika); Pfeffer Collection in Prague Museum.

Notes: (1) Pfeffer 1940b: 122 (treated as a subspecies of *buyssoni*).

References: **(ds)** Pfeffer 1947d: 127; Sainte-Claire & Mequignon 1938: 447. **(tx)** Balachowsky 1949a: 239; Pfeffer 1927: 111, 1940b: 109; Schedl 1934f: 1643.

cacuminatus Blandford 1904: 238. Lectotype ♀; San Geronimo, Verapaz, Guatemala; BMNH, London, designated by Bright 1976b: 184.

Distribution: North America (Guatemala/ Honduras/ Jalisco, Michoacan in Mexico).

Hosts: *Pinus oocarpa*, *P. spp.*

References: **(hb)** Atkinson & Equihua 1955c: 360; Atkinson et al. 1956: 37. **(ds)** Atkinson & Equihua 1955c: 360, 1958: 94; Atkinson et al. 1956: 37; Blackwelder 1947: 751; Bright 1981d: 304; Hagedorn 1910d: 70; Kleine 1913b, 1914b: 362; Wood, S. L. 1982b: 1101. **(tx)** Blackman 1942a: 218; Blandford 1904: 238; Bright 1976b: 184, 1981d: 304; Hagedorn 1910a: 100; Schedl 1938a: 183, 1955g: 17; Wood, S. L. 1982b: 1101.

californicus Bright 1976c: 427. Holotype ♀; Bonnie Doone, Santa Cruz Co., California [USA]; USNM, Washington, automatic.

Distribution: North America (California).

Hosts: *Pinus attenuata*, *P. radiata*.

References: **(ds)** Bright 1981d: 249; Wood, S. L.

- 1952b: 1069. **(tx)** Bright 1976c: 427, 1981d: 249; Wood, S. L. 1982b: 1069.
- deleoni* Bright 1966b: 302. Holotype ♀; Bonnie Doone, Santa Cruz Co., California [USA]; USNM, Washington, preoccupied by Blackman 1942.
- References: **(tx)** Bright 1966b: 302, 1976c: 427.
- camerunus* Eggers 1920: 35. Syntypes 2, sex?; Kamerun; Hamburg Museum specimen to Eggers Collection now in NIMW, Wien, 1 in MNB, Berlin.
- Distribution: Africa (Cameroon).
- Notes: (1) Schedl 1979c: 51 (citation of holotype invalid).
- References: **(ds)** Schedl 1965e: 353, 1966c: 225. **(tx)** Eggers 1920: 35, 1927a: 184; Schedl 1938a: 183, 1962j: 6, 1979c: 51.
- carinatus* Bright 1978: 73. Holotype ♀; Sevey, New York [USA]; CNCI, Ottawa.
- Notes: (1) Bright 1978: 73 (subspecies established).
- References: **(ds)** Bright 1981d: 179; Furniss, M. M. 1978; Wood, S. L. 1982b: 1077. **(tx)** Bright 1978: 73, 1981d: 179; McNamara 1984: 754; Wood, S. L. 1982b: 1077.
- carinatus carinatus*:
- Distribution: North America (New Brunswick, Quebec in Canada/ New York in USA).
- Hosts: *Picea* sp., *Pinus strobus*.
- References: **(tx)** Bright 1978: 73.
- carinatus monticolae* Bright 1978: 74 Holotype ♀; Hazelton, British Columbia [Canada]; CNCI, Ottawa.
- Distribution: North America (Alberta, British Columbia in Canada).
- Hosts: *Pinus contorta*.
- References: **(ds)** Bright 1981d: 179; Wood, S. L. 1982b: 1077. **(tx)** Bright 1978: 74, 1981d: 179; McNamara 1984: 754; Wood, S. L. 1982b: 1077.
- cariniceps* LeConte 1876: 353. Holotype ♀; Detroit, Michigan [USA]; MCZ, Cambridge.
- Figures: Bright 1981d: 355.
- Distribution: North America (Alberta, NE British Columbia, Manitoba, New Brunswick, Nova Scotia, Ontario, Quebec, Saskatchewan in Canada/ Connecticut, Indiana, Kentucky, Maine, Massachusetts, Michigan, Minnesota, New Hampshire, New York, North Carolina, Ohio, Pennsylvania, South Carolina, Virginia, Wisconsin in USA).
- Hosts: *Abies balsamea*, *Picea glauca*, *Pinus banksiana*, *P. resinosa*, *P. strobus*, *P. virginiana*.
- Notes: (3) Blackman 1928b: 72 (redescribed).
- References: **(cn)** Blackman 1950; Browne 1968b: 564; Chamberlin 1924; Currie 1905: 80; Doane et al. 1936; Hopkins 1899c: 443, 1901b: 8, 1904a: 24; Lindquist, O. H. & Syme 1981: 88; Martin, J. L. 1964; Schuder 1969: 80; Swaine 1918a: 97, 102; Syme & Nystrom 1988: 84. **(cc)** Bushing 1965: 464; Felt 1906: 674; Frederick, Sloan, & Skowron 1976; Hopkins 1901b: 8; Pierce, W. D. 1908: 357; Thomas & Lindquist 1956: 2. **(hb)** Baker, W. L. 1972: 255; Blackman 1950; Browne 1968b: 564; Chamberlin 1939: 377; Clittenden 1893a: 393; Doane et al. 1936; Felt 1906: 674; Hopkins 1899c: 443, 1901b: 8, 1904a: 24; Pierce, W. D. 1907: 290; Swaine 1918a: 97, 102. **(ds)** Anonymous 1926c: 519; Atkinson et al. 1991: 162; Baker, W. L. 1972: 255; Blackman 1950; Blatchley & Leng 1916: 633; Bright 1976d: 185, 1981d: 174; Browne 1968b: 564; Chamberlin 1925, 1939: 377; Currie 1905; Deyrup & Atkinson 1987b: 67; Dodge 1938; Drooz 1985: 364; Hagedorn 1910d: 70; Henshaw 1882: 268, 1885: 147; Hopkins 1893a: 130, 1893b: 209; Hubbard & Schwarz 1878a: 665; Kleine 1913b: 140, 1934a: 159; Leng 1920: 341; Leonard 1928: 519; Lindquist, O. H. & Syme 1981: 88; Morgan, A. V. & Morgan 1980: 1110; Proctor 1946: 208; Schuder 1969: 80; Schwert et al. 1985; Swaine 1909: 133; Syme & Nystrom 1988: 84; Wichmann 1896a: 309; Wood, S. L. 1982b: 1077. **(tx)** Blackman 1919b: 134–136, 142, 1928b: 72; Blatchley & Leng 1916: 633; Bright 1981d: 174, 355; Chamberlin 1939: 377; Dodge 1938: 43–45; Hagedorn 1910a: 100; LeConte 1876: 353; Lindquist, O. H. & Syme 1988: 84; Swaine 1909: 133–134, 1917: 24, 1918a: 97, 102; Syme & Nystrom 1988: 84; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1977d: 514, 1982b: 1077. **(ms)** Eckstein 1900c.
- canadensis* Swaine 1917: 24. Lectotype ♀; Ste. Anne de Bellevue, Quebec [Canada]; CNCI, Ottawa, designated by Bright 1967b: 678. Synonymy: Blackman 1928b: 72.
- References: **(cn)** Chamberlin 1924; Swaine 1918a: 97, 102. **(hb)** Swaine 1918a: 97, 102. **(ds)** Anonymous 1926c: 519; Beaulne 1956; Britton 1920a; Chamberlin 1925; Hatch 1924; Leng 1920: 341; Leonard 1928: 519. **(tx)** Blackman 1919b: 136–137, 1922c, 1928b: 72; Bright 1967b: 678; de Ruette 1970: 108; Swaine 1917: 24, 1918a: 97, 102.
- cognatus* Blackman 1928b: 69. Holotype ♀; Davidson's R., Asheville, North Carolina, 3200 ft. [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.
- References: **(cn)** Blackman 1950; Schuder 1969: 80. **(hb)** Beal & Massey 1945: 132; Blackman 1950; Chamberlin 1939: 376. **(ds)** Beal & Massey 1945: 132; Blackman 1950; Chamberlin 1939: 376; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Schuder 1969: 80. **(tx)** Beal & Massey 1945: 132; Blackman 1928b: 69; Chamberlin 1939: 376; de Ruette 1970: 108; Wood, S. L. 1977d: 514.
- carinulatus* Swaine 1925b: 193. Holotype ♂; Tres Ritos, New Mexico [USA]; CNCI, Ottawa.
- Distribution: North America (Alaska/ Arizona, Colorado, New Mexico, Utah in USA).

Hosts: *Picea engelmannii*, *P. pungens*, 2 males from Alaska in *Pinus contorta*.

References: **(hb)** Chamberlin 1939: 375. **(ds)** Bright 1981d: 228; Chamberlin 1939: 375; Evans, D. 1983: 35; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1068. **(tx)** Blackman 1928b: 60, 66–67; Bright 1967b: 678, 1977: 514, 1981d: 228; Chamberlin 1939: 375; Evans, D. 1983: 35; LeConte 1874a: 70; de Ruelle 1970: 108; Swaine 1925b: 193; Wood, S. L. 1977b: 209, 1982b: 1068. *opimus* Blackman 1928b: 80. Holotype ♀;

Capitan Mts., New Mexico [USA]; USNM, Washington. Synonymy: Bright 1977: 514.

References: **(cn)** Evans, D. 1982; Lindgren 1980: 69. **(cc)** Stevens 1973. **(hb)** Bright & Stark 1973: 109; Chamberlin 1939: 379; Evans, D. 1982; Lindgren, B. 1980a: 69. **(ds)** Bright 1976d: 183; Bright & Stark 1973: 109; Chamberlin 1939: 379; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 62, 80–81; Bright 1971b: 67, 1977: 514; Chamberlin 1939: 379; Evans, D. 1982; Keen 1929a: 32; McNamara 1984: 757; Wood, S. L. 1951a: 128.

carmeli Swaine 1918a: 100. Holotype ♀; Carmel, California [USA]; CNCI, Ottawa.

Figures: Bright 1981d: 347.

Distribution: North America (California).

Hosts: *Pinus attenuata*, *P. coulteri*, *P. muricata*, *P. radiata*, *P. torreyana*.

Notes: (3) Blackman 1928b: 97 (redescribed). The holotype is missing from its microcard and may be lost (DEB).

References: **(cn)** Anonymous 1964h, 1966p, 1968g, 1975d; Doane et al. 1936; Herrick 1935: 250; Swaine 1918a: 96, 100. **(cc)** Bushing & Bright 1965: 203; Kim 1971; Marsh 1979: 150. **(hb)** Bright 1981d: 137; Bright & Stark 1973: 111; Burke 1932; Chamberlin 1939: 385; Doane et al. 1936; Herrick 1935: 250; Swaine 1918a: 96, 100. **(ds)** Anonymous 1964h, 1966p, 1968g, 1975d; Bright 1981d: 137; Bright & Stark 1973: 111; Burke 1932: 366, 1933: 49–59, 1937: 28; Chamberlin 1925, 1939: 385; Hopping 1922; Keen 1929a: 33; Kleine 1934a: 159; Leng 1920: 341; Wood, S. L. 1982b: 1049. **(tx)** Blackman 1928b: 97; Bright 1964: 166, 1967b: 678, 1981d: 137, 347; Chamberlin 1939: 385; Keen 1929a: 33; de Ruelle 1970: 108; Swaine 1918a: 96, 100; Wood, S. L. 1982b: 1049.

torreyanae Swaine 1918a: 101. Holotype ♀; San Diego, California [USA]; CNCI, Ottawa. Synonymy: Blackman 1928b: 98.

References: **(cn)** Keen 1929: 54; Swaine 1918a: 97, 101. **(hb)** Keen 1929: 54; Swaine 1918a: 97, 101. **(ds)** Chamberlin 1925: Hopping 1922; Keen 1929: 54; Leng 1920: 341. **(tx)** Blackman 1928b: 97; Bright 1970: 679; de Ruelle 1970: 111; Swaine 1918a: 97, 101–102.

carriolicus Wichmann 1910a: 145. Syntypes, sex?; Krain (Austria-Hung.); Adelsberg; not given.

Figures: Grune 1979: 118.

Distribution: Europe (Austria/ Czechoslovakia/ France/ Germany/ Hungary).

Hosts: *Pinus nigra*, *P. sylvestris*.

References: **(bv)** Grune 1979: 119. **(cc)** Nosek 1959b: 87; Pfeffer 1960: 346; Wichmann 1916: 23. **(hb)** Nosek 1959b: 87; Spessivtsev 1921b: 221, 1923: 208; Wichmann 1916: 23. **(ds)** Bovey 1976; Grune 1979: 119; Holzschuh 1966: 61; Horion 1951; Karpinski 1931: 30; Kleine 1913a: 35, 1934a: 159; Lucht 1987: 278; Pfeffer 1960: 346, 1976: 340, 1984: 277, 1989a: 65; Postner 1974: 436; Schedl 1980a: 22, 1981b: 78. **(tx)** Balachowsky 1949a: 238; Eggers 1915b; Grune 1979: 118–119; Lucht 1987: 278; Pfeffer 1927: 111–112, 1940b: 109, 113, 123, 1955a: 220, 1976: 340; Postner 1974: 436; Reitter 1913a: 94; Schedl 1934f: 1643, 1980a: 22, 1981b: 78; Spessivtsev 1922: 221, 1926: 50; Wichmann 1910a: 145, 1913d: 144, 1916: 23–25, 1916: 432–433.

cascoensis Blackman 1928b: 99. Holotype ♀; Peak Island, Maine [USA]; USNM, Washington.

Figures: Bright 1981d: 349.

Distribution: North America (Alberta, Newfoundland, Northwest Territories, Ontario in Canada/ Maine, New York in USA).

Hosts: *Picea glauca*.

References: **(cn)** Blackman 1950. **(hb)** Blackman 1950; Chamberlin 1939: 386. **(ds)** Blackman 1950; Bright 1976d: 186, 1981d: 150; Chamberlin 1939: 386; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1088. **(tx)** Blackman 1928b: 89, 99–100; Bright 1977: 515, 1981d: 150, 349; Chamberlin 1939: 386; McNamara 1984: 754; Wood, S. L. 1977b: 209, 1982b: 1088.

pilifer Schedl 1931b: 166. Holotype ♀; Frater, Ontario [Canada]; CNCI, Ottawa. Synonymy: Bright 1977: 515.

References: **(hb)** Chamberlin 1939: 378. **(ds)** Chamberlin 1939: 378; Kleine 1934a: 161; Leng & Mutchler 1933: 95. **(tx)** Bright 1977: 515; Chamberlin 1939: 378; de Ruelle 1970: 110; Schedl 1931b: 166–168.

cavatus Bright 1978: 74. Holotype ♀; Canoe Lake, Saskatchewan [Canada]; CNCI, Ottawa.

Distribution: North America (Manitoba, Saskatchewan in Canada).

Hosts: *Pinus banksiana*, *P. resinosa*.

References: **(ds)** Bright 1981d: 187; Wood, S. L. 1982b: 1075. **(tx)** Bright 1978: 74, 1981d: 187; McNamara 1984: 754; Wood, S. L. 1982b: 1075.

cedri Wood 1989: 182. Holotype ♀; Kashmir: Buniyar, Jhelum Valley; FRI, Dehra Dun.

Distribution: Asia (Kashmir in India).

Hosts: *Cedrus deodara*, *Pinus gerardiana*.

Notes: (1) Beeson 1941: 292 (*cedri*, nomen nudum).

References: **(tx)** Wood, S. L. 1989: 182.

- celatus (Schedl)** 1967d: 9 (*Breriophthorus*). Holotype, sex?; Nova Tentonia, Santa Catarina, Brasilien; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) McNamara 1977: 194; Schedl 1967d: 9–10, 1979c: 55.
- cephalonicae Pfeffer** 1940b: 119. Syntypes, sex?; Graecia Meridionalis, Montes Chelmos; Pfeffer Collection in Prague Museum. Distribution: Europe (Greece/ Poland). Hosts: *Abies cephalonica*, *A. alba*. References: (bv) Grune 1979: 119, (ec) Michalski & Ratajczak 1989, (hb) Karpinski 1955: 80, (ds) Atkinson et al. 1986: 38; Grune 1979: 119; Karpinski 1955: 80; Nunberg 1954: 71, 1956c: 163, 168; Pfeffer 1940b: 119, 1947: 129, 1976: 340; Postner 1974: 437. (tx) Grune 1979: 119; Nunberg 1954: 71; Pfeffer 1940b: 108, 119, 1955a: 215, 1976: 340; Postner 1974: 437.
- polonicus Karpinski** 1949: 125. Syntypes, sex?; mountain forests of National Park in Pieniny (Carpathian Mts.); Karpinski Collection. Synonymy: Nunberg 1956c: 163. References: (ec) Boucek 1955: 83–92; Graham 1969: 878; Nuorteva 1957b: 69. (hb) Karpinski 1955: 77. (ds) Horion 1951; Karpinski 1949: 125, 1955: 77. (tx) Karpinski 1949: 125; Nunberg 1956c: 161, 163, 168.
- chalconensis Hopkins** 1905b: 73. Holotype ♀; Chalco, Mexico; USNM, Washington. Distribution: North America (Mexico, Michoacan in Mexico). Hosts: *Pinus* sp. (bole). Notes: (1) Hopkins published the species name *chaleocensis* and the type locality as “Chaleco,” but the label on the holotype clearly reads “Chalco.” The error in spelling is an obvious lapsus calami that was corrected in Wood 1982b: 1099. References: (ds) Atkinson & Equihua 1985a: 97, 1985: 96; Blackwelder 1947: 781; Bright 1981d: 279; Ferrer 1942; Hagedorn 1910d: 70; Kleine 1913b: 140, 1914b: 351; Wood, S. L. 1982b: 1099. (tx) Bright 1978: 71, 1981d: 279; Hagedorn 1910a: 100; Hopkins 1905b: 73; Schedl 1940a: 348; Wood, S. L. 1982b: 1099.
- herrarai Hopkins** 1905b: 71. Syntypes, sex?; Chalco, Mexico; USNM, Washington. Synonymy: Bright 1978: 71. References: (ds) Blackwelder 1947: 782; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 351. (tx) Bright 1978: 71; Hagedorn 1910a: 101; Hopkins 1905b: 71; Schedl 1940a: 348.
- chilgoza Wood** 1989: 183. Holotype ♀; Punjab: Kilba, U. Bashar: FR1, Delhra Dim. Distribution: Asia (Punjab in India). Hosts: *Pinus gerardiana*. Notes: (1) Beeson 1941 (1961: 292) (*chilgoza*, nomen nudum). References: (tx) Wood, S. L. 1989: 183.
- ciliatus Blackman** 1942a: 211. Holotype ♀; Jalapa, Veracruz, Mexico; USNM, Washington. Distribution: North America (Veracruz in Mexico). Hosts: *Pinus patula*, *P. sp.* References: (hb) Atkinson & Equihua 1985c: 360. (ds) Atkinson & Equihua 1985c: 360, 1985: 94; Bright 1981d: 250; Thomas, J. B. 1966; Wood, S. L. 1982b: 1069. (tx) Blackman 1942a: 211; Bright 1981d: 250; McNamara 1984: 754; Wood, S. L. 1982b: 1069.
- clarus Blackman** 1928b: 130. Holotype ♀; Kaibab National Forest, Arizona [USA]; USNM, Washington. Distribution: North America (Arizona). Hosts: *Pinus leiophylla*, *P. ponderosa*. References: (cn) Blackman 1931c. (ec) Blackman 1931c. (hb) Blackman 1931c; Chamberlin 1939: 397. (ds) Bright 1981d: 324; Chamberlin 1939: 397; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1108. (tx) Blackman 1928b: 130; Bright 1981d: 324; Chamberlin 1939: 397; Keen 1929a: 33; de Ruelle 1970: 108; Wood, S. L. 1982b: 1108.
- clivus Bright** 1977: 522. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico; CNCI, Ottawa. Distribution: North America (Nuevo Leon in Mexico). Hosts: *Pinus strobiformis*. References: (ds) Bright 1981d: 250; Wood, S. L. 1982b: 1040. (tx) Bright 1977: 522, 1981d: 250; McNamara 1984: 754; Wood, S. L. 1982b: 1040.
- collaris Schedl** 1965c: 67. Holotype ♀; Madagascar, Montagne d'Ambre; Schedl Collection in NHMW, Wien. Distribution: Madagascar. Hosts: Tree twigs. References: (tx) Schedl 1965c: 67, 1977b: 110, 1979c: 59.
- comosus Blackman** 1928b: 65. Holotype ♀; Sacramento Mts., New Mexico [USA]; USNM, Washington. Distribution: North America (Durango in Mexico/ Arizona, New Mexico in USA). Hosts: *Pinus ponderosa*. References: (hb) Chamberlin 1939: 375. (ds) Bright 1981d: 277; Chamberlin 1939: 375; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1045. (tx) Blackman 1928b: 65; Bright 1977: 514, 1981d: 277; Chamberlin 1939: 375; Keen 1929a: 32; McNamara 1984: 754; Wood, S. L. 1982b: 1045.
- foratus Wood** 1967d: 40. Holotype ♀; 16 km W El Salto, Durango, Mexico; CNCI, Ottawa. Synonymy: Bright 1977: 514. References: (tx) Bright 1977: 514; de Ruelle 1970: 109; Wood, S. L. 1967d: 40.
- concarus Blackman** 1928b: 85. Holotype ♀; Seney, Michigan [USA]; USNM, Washington.

Distribution: North America (British Columbia (?), New Brunswick, Nova Scotia, Ontario, Quebec in Canada/ Maine, Michigan, Wisconsin in USA). The British Columbia population could be distinct.

Hosts: *Pinus banksiana*, *P. resinosa*, *Picea glauca*.
References: (cn) Blackman 1950. (hb) Blackman 1950; Chamberlin 1939: 380. (ds) Blackman 1950; Bright 1976d: 156, 1981d: 185; Chamberlin 1939: 380; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1075. (tx) Blackman 1928b: 85; Bright 1981d: 185; Chamberlin 1939: 380; Wood, S. L. 1982b: 1075.

hesperius Bright 1978: 76. Holotype ♀; Hudson Hope, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Wood 1982b: 1075.

Notes: (1) This could possibly be a subspecies; not enough material available for study to evaluate (SLW); should be treated as a species (DEB).

References: (ds) Bright 1981d: 181. (tx) Bright 1978: 76, 1981d: 181; McNamara 1984: 755; Wood, S. L. 1982b: 1075.

concentralis Eichhoff 1878b: 188. Holotype, sex?; Cuba; Hamburg Museum, lost; neotype ♀; Biscayne Bay, Florida [USA]; USNM, Washington, designated by Bright 1978: 72.

Figures: Bright 1981d: 337.

Distribution: Antilles Islands (Bahamas Islands/Cuba), North America (Florida in USA).

Hosts: *Metopium toxiferum*.

References: (cn) Swaine 1918a: 95, 104. (cc) Schwarz 1889a: 163. (hb) Chamberlin 1939: 361; Swaine 1918a: 95, 104. (ds) Beutenmuller 1891: 50; Blackwelder 1947; Blatchley & Leng 1916: 630; Bright 1981d: 56, 1985c: 176; Chamberlin 1939: 361; Hagedorn 1910d: 70; Henshaw 1895: 44, 1899: 132; Kleine 1913b: 140, 1914b: 378, 383, 1934a: 159; Leng 1920: 341; Schwarz 1889a: 163–164; Swaine 1909: 134; Wood, S. L. 1957c: 401, 1982b: 1114. (tx) Blackman 1928b: 37, 40; Blatchley & Leng 1916: 630; Bright 1978: 72, 1981d: 56, 337, 1985c: 176; Chamberlin 1939: 361; Eichhoff 1878a: 389, 1878b: 188; Hagedorn 1910a: 100; Swaine 1909: 134, 1918a: 95, 104; Wood, S. L. 1957c: 401, 1982b: 1114.

lateralis Swaine 1917: 27. Lectotype ♀; Key West, Florida [USA]; Cornell University, Ithaca, New York, designated by Bright 1967b: 678. Synonymy: Wood 1957c: 401.

References: (cn) Swaine 1918a: 95, 104. (hb) Chamberlin 1939: 361; Swaine 1918a: 95, 104. (ds) Chamberlin 1939: 361; Kleine 1934a: 160; Leng 1920: 341. (tx) Blackman 1928b: 37, 40; Bright 1967b: 678; Chamberlin 1939: 361; Hoebeke 1978; de Ruette 1970: 110; Swaine 1917: 27, 1918a: 95, 104; Wood, S. L. 1957c: 401, 1982b: 1114.

concinus Wood 1977b: 214. Holotype ♀; in wood from Plaquepague, Mexico, intercepted at Nogales, Arizona [USA]; USNM, Washington.

Distribution: North America ("Mexico").

References: (ds) Bright 1981d: 34; Wood, S. L. 1982b: 1116. (tx) Bright 1981d: 34; Wood, S. L. 1977b: 214, 1982b: 1116.

confertus Swaine 1917: 27. Lectotype ♀; Adams Lake, British Columbia [Canada]; CNCI, Ottawa, designated by Bright 1966b: 304, 1967b: 678.

Figures: Bright 1981d: 306, 372, Evans 1982: 31.

Distribution: North America (S British Columbia in Canada/ Baja California, Chihuahua, Coahuila, Hidalgo, Nuevo Leon, Puebla in Mexico/ Arizona, California, Colorado, Idaho, W Montana, Nevada, New Mexico, Oregon, W Texas, Utah, Washington, Wyoming in USA).

Hosts: *Pinus albicollis*, *P. cembroides*, *P. contorta*, *P. edulis*, *P. flexilis*, *P. jeffreyi*, *P. monticola*, *P. ponderosa*, *P. strobfornis*.

Notes: (3) Blackman 1928b: 122 (redescribed). Bright 1981d: 307 (subspecies *agnatus* recognized, validity?).

References: (bv) Jacobson 1972; Miller & Keen 1960: 115; Schmitz 1958; Vite 1965: 267; Vite & Pitman 1967: 683–701. (cn) Chamberlin 1924; Evans, D. 1982b: 8; Keen 1929: 53; Lindgren 1980a: 68; Miller & Keen 1960: 115; Ruppel 1967: 78; Salman 1938a: 613–616; Schuder 1969: 78; Swaine 1918a: 98, 103. (ec) Amman, A. G., Amman, & Amman 1974; Chatelain & Schenk 1983; Dahlsten & Stephen 1974: 1213; Deyrup & Gara 1978: 275; Miller & Keen 1960: 115; Rice 1965; Salman 1938a: 613; Schmitz 1985. (hb) Amman, A. G., Amman, & Amman 1974; Bright & Stark 1973: 115; Chamberlin 1939: 395, 1955; Evans, D. 1982: 8; Keen 1929: 53; Lindgren 1980a: 68; Miller & Keen 1960: 115; Schmitz 1985; Swaine 1918a: 98, 103. (ds) Amman, A. G., Amman, & Amman 1974; Atkinson & Equihua 1985c: 361, 1988: 94; Bright 1976d: 181, 1981d: 305; Bright & Stark 1973: 115; Chamberlin 1917: 355, 1925, 1939: 395, 1955; Evans, D. 1983: 35; Furniss, R. L. & Carolin 1977: 402; Hopping 1922; Keen 1929a: 33; 1929c: 53; Kleine 1934a: 159; Leng 1920: 341; McComb et al. 1953: 2; Patterson & Hatch 1945: 153; Ruppel 1967: 78; Schuder 1969: 78; Wood, S. L. 1948: 53, 1951a: 128, 1972a: 427, 1982b: 1107. (tx) Blackman 1928b: 115, 122–123; Bright 1966b: 304, 1967b: 678, 1981d: 305–306, 372; Chamberlin 1939: 395, 1955; Evans, D. 1982: 8, 31, 1983: 35; Hoebeke 1978; Keen 1929a: 33; de Ruette 1970: 109; Swaine 1917: 27, 1918a: 98, 103; Wood, S. L. 1951a: 128, 1972a: 427, 1977d: 514, 1982b: 1107, *agnatus* Blackman 1928b: 125. Holotype ♀; Clondercroft, New Mexico [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.

Notes: (1) Bright 1981d: 305 (treated as a sub-

species, validity doubtful). Wood 1952b: 1107 (this name was based on a minor clinal variation with no taxonomic value).

References: **(hb)** Chamberlin 1939: 396. **(ds)** Bright 1981d: 305; Chamberlin 1939: 396; Keen 1929a: 33; Kleine 1934a: 158; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 115, 125–127; Bright 1981d: 305; Chamberlin 1939: 396; Keen 1929a: 33; de Ruelle 1970: 107; Wood, S. L. 1977d: 514.

comptus Blackman 1928b: 127. Holotype ♀: Santa Catalina Mts., Arizona [USA]; USNM, Washington. Synonymy: Wood 1977d: 514.

References: **(hb)** Chamberlin 1939: 396. **(ds)** Chamberlin 1939: 396; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 127; Chamberlin 1939: 396; Keen 1929a: 33; de Ruelle 1970: 109; Wood, S. L. 1977d: 514.

hurkei Blackman 1928b: 129. Holotype ♀: Meyers, California [USA]; USNM, Washington. Synonymy: Bright 1966b: 304.

References: **(ec)** Bushing 1965: 464; DeLeon 1934a. **(hb)** Chamberlin 1939: 396, 1958. **(ds)** Chamberlin 1939: 396, 1958; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1948: 53, 1951a: 128. **(tx)** Blackman 1928b: 115, 129–130; Bright 1966b: 304; Chamberlin 1939: 396, 1958; Keen 1929a: 33; Wood, S. L. 1951a: 128.

confinis LeConte 1876: 354. Lectotype ♀: Mojave region, California [USA]; MCZ, Cambridge, designated by Bright 1976b: 185.

Figures: Bright 1976d: 204, 213, 1981d: 268, 367. Distribution: North America (British Columbia in Canada/ Arizona, California, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, Wyoming in USA).

Hosts: *Pinus albicaulis*, *P. contorta*, *P. coulteri*, *P. jeffreyi*, *P. lambertiana*, *P. monophylla*, *P. ponderosa*, *P. sabiniana*, *P. strobiformis*.

References: **(ay)** Hopkins 1894. **(bv)** Miller & Keen 1960: 115. **(en)** Anonymous 1964h; Chamberlin 1924; Currie 1905: 71; Doane et al. 1936; Essig 1926: 519, 1958: 519; Evans, D. 1982: 8; Felt 1906: 751; Hopkins 1894: 278, 1904a: 16; Keen 1929c: 53; Lindgren 1980a: 68; Miller & Keen 1960: 115; Sahnan 1938a: 613–618; Schuder 1969: 78; Swaine 1918a: 97, 101. **(ec)** Felt 1906: 751; Miller & Keen 1960: 115; Sahnan 1938a: 613. **(hb)** Bright 1981d: 267; Bright & Stark 1973: 267; Chamberlin 1939: 375, 1958; Doane et al. 1936; Essig 1926: 519, 1958: 519; Evans, D. 1982: 8; Felt 1906: 751; Hopkins 1894g, 1904a: 16; Keen 1929c: 53; Lindgren 1980a: 68; Miller & Keen 1960: 115; Pierce, W. D. 1907: 290; Swaine 1918a: 97, 101. **(ds)** Anonymous 1964h; Atkinson & Equihua 1985: 94; Bright 1976d: 181, 1981d: 267; Bright & Stark

1973: 111; Chamberlin 1925, 1939: 375, 1958; Currie 1905; Essig 1926: 519, 1958: 519; Evans, D. 1983: 35; Ferrer 1942; Furniss, R. L. & Carolin 1977: 402; Gast et al. 1989: 385; Hagedorn 1910d: 70; Henshaw 1882: 268, 1885: 147; Hopkins 1893a: 130, 1893b: 209, 1899: 284, 344, 443; Hopping 1922; Keen 1929a: 32, 1929c: 53; Kleine 1913b: 140, 1914b: 351, 391, 1934a: 159; Leng 1920: 341; Patterson & Hatch 1945: 163; Schuder 1969: 78; Swaine 1909: 135; Wood, S. L. 1972a: 426, 1982b: 1079. **(tx)** Blackman 1928b: 60, 67; Bright 1976b: 185, 1976d: 204, 213, 1981d: 267–268, 367; Chamberlin 1939: 375, 1958; Evans, D. 1982: 8, 1983: 35; Hagedorn 1910a: 100; Keen 1929a: 32; LeConte 1876: 354; Schedl 1940a: 348; Swaine 1909: 135, 1918a: 97, 101; Wood, S. L. 1972a: 426, 1982b: 1079.

confractus Bright 1985c: 179. Holotype ♀: Jamaica: Portland between Hardwar Cap and Green Hills, 1100 m; Carnegie Museum of Natural History, Pittsburgh.

Distribution: Antilles Islands (Jamaica).

References: **(ds)** Bright 1985c: 176, 179. **(tx)** Bright 1985c: 176, 179.

confusus Blandford 1904: 237. Lectotype ♀: San Geronimo, Guatemala; BMNH, London, designated by Bright 1976b: 184.

Figures: Bright 1981d: 370.

Distribution: North America (El Salvador/ Guatemala/ Honduras/ Chiapas, Durango, Guerrero, Jalisco in Mexico/ Nicaragua/ Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, Pennsylvania, South Carolina, Texas, Virginia, West Virginia in USA).

Hosts: *Pinus cchinata*, *P. elliotti*, *P. michoacana*, *P. ochloterenai*, *P. oocarpa*, *P. palustris*, *P. taeda*.

Notes: (3) Bright 1981d: 291 (subspecies *bellus* recognized). Subspecies apparently based on obscure clinal variation; validity very doubtful (SLW).

References: **(bv)** Turnbow & Franklin 1980. **(ec)** Becker 1953, 1955. **(hb)** Becker 1953, 1955; Deyrup & Atkinson 1987a: 66; Schwerdtfeger 1957c: 494. **(ds)** Atkinson & Equihua 1985c: 360, 1988: 96; Becker 1953; Blackwelder 1947: 781; Bright 1981d: 291; Deyrup & Atkinson 1987a: 66; Kleine 1913b: 140, 1914b: 363; Schedl 1963c: 158, 1977e: 42; Turnbow & Franklin 1980; Wood, S. L. 1977a: 72, 1982b: 1100. **(tx)** Blandford 1904: 237; Bright 1976b: 184, 1981d: 291; Hagedorn 1910a: 100; Hopkins 1905b: 74; de Ruelle 1970: 104; Schedl 1938a: 183, 1955g: 18–19; Wood, S. L. 1977d: 515, 1982b: 1100.

bellus Blackman 1928b: 123. Holotype ♀: Monongalia Co., West Virginia [USA]; USNM, Washington. Synonymy: Wood 1977d: 515.

Notes: (3) Bright 1981: 291 (a subspecies of *confusus*, validity?).

References: **(en)** Blackman 1950. **(hb)** Beal & Massey 1945: 129; Blackman 1950; Chamberlin

1939: 396. **(ds)** Beal & Massey 1945: 129; Blackman 1950; Bright 1981d: 291; Chamberlin 1939: 396; Kirk 1969, 1970; Kleine 1934a: 158; Knull 1932: 66; Leng & Mutchler 1933: 52; Smith, J. B. 1910: 401. **(tx)** Beal & Massey 1945: 129; Blackman 1928b: 115, 123–124; Bright 1981d: 291, 370; Chamberlin 1939: 396; Hopkins 1915c: 186; de Ruette 1970: 108; Wood, S. L. 1977d: 515.

congonus Eggers 1927a: 185. Syntypes, sex?, Belg. Congo: MRCB, Tervuren and Eggers Collection, 1 Eggers syntype in NHMW, Wien.

Distribution: Africa (Zaire).

Hosts: *Azelia africanus*, *Albizia gummifera*, *Pentaclethra celtidifera*.

References: **(cn)** Ghesquiere 1933a, **(hb)** Schedl 1962j: 6. **(ds)** Ghesquiere 1933a: 33–35, 1933b: 783; Kleine 1934a: 159; Schedl 1962j: 6. **(tx)** Eggers 1927a: 185; Schedl 1938a: 183, 1962j: 5, 7, 1965f: 10, 1979c: 63.

conscriptus Bright 1986b: 680. Holotype ♀; Mexico: La Herradura, Mpio. Tepoztlán, Mor; CNCI, Ottawa.

Distribution: North America (Morelos in Mexico).

References: **(tx)** Bright 1986b: 680.

consimilis LeConte 1878b: 622. Lectotype ♂; Marquette, Michigan [USA]; MCZ, Cambridge, designated by Bright 1976b: 185.

Figures: Bright 1981d: 369.

Distribution: North America (Alberta, British Columbia, Manitoba, Nova Scotia, Ontario, Quebec, Saskatchewan in Canada/ Florida, Georgia, Kentucky, Louisiana, Maine, Massachusetts, Michigan, Minnesota, Mississippi, New Hampshire, New York, North Carolina, Ohio, South Carolina, Tennessee, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Picea glauca*, *P. sp.*, *Pinus banksiana*, *P. resinosa*, *P. rigida*, *P. strobus*, *P. taeda*, *P. virginiana*, *P. spp.*, *Larix laricina*.

Notes: (3) Blackman 1928b: 131 (redescribed).

References: **(ay)** Thomas, J. B. 1957: 4. **(cn)** Bentenmiller 1891: 50; Blackman 1950; Doane et al. 1936; Felt 1906: 482, 503, 737; Schuder 1969: 80; Swaine 1918a: 104; Syme & Nystrom 1988: 84; Thomas, J. B. 1958: 394. **(ec)** Ashmead 1894: 33; Bushing 1965: 464; Felt 1906: 482; Pierce, W. D. 1908: 385; Riley 1891c: 123; Riley & Howard 1891: 123; Schwarz 1889a: 163; Thomas, J. B. 1958: 394. **(hb)** Beal & Massey 1945; Blackman 1950; Chamberlin 1939: 397; Doane et al. 1936; Felt 1906: 482; Swaine 1918a: 104. **(ds)** Anonymous 1926c: 519; Beal & Massey 1945; Beaulne 1956; Blackman 1950; Blatchley & Leng 1916: 630; Bright 1976d: 186, 1981d: 283; Burks 1979: 781; Chamberlin 1939: 397; Chapin 1917: 29; Deyrup 1951b: 8; Dodge 1938; Evans, D. 1983: 35; Hagedorn 1910d: 71; Henshaw 1882: 268, 1885: 147; Hopkins 1893a: 130, 1893b: 209;

Hubbard & Schwarz 1878a: 665, 1878b: 622; Kleine 1913b: 140, 1914b: 384, 1934a: 159; Leng 1920: 341; Leonard 1928: 519; Notman 1920: 184; Schuder 1969: 80; Schwarz 1889a: 163–164, 1891c: 357; Swaine 1909: 135; Syme & Nystrom 1988: 84; Ulke 1902: 56; Wood, S. L. 1982b: 1104. **(tx)** Beal & Massey 1945; Blackman 1921: 14–16, 1928b: 116, 131–133; Blatchley & Leng 1916: 630; Bright 1976b: 185, 1981d: 283, 369; Chamberlin 1939: 397; Dodge 1938: 43, 45; Evans, D. 1983: 35; Hagedorn 1910a: 100; LeConte 1878b: 622; Schedl 1955g: 25; Schwarz 1891a: 167, 1891c: 357, 1893: 167, 1909: 135; Swaine 1909: 135, 1918a: 104; Syme & Nystrom 1988: 84; Thomas, J. B. 1957: 4; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1977d: 515, 1982b: 1104.

granulatus Swaine 1917: 28. Lectotype ♂; presumably Manitoba, Quebec, or Nova Scotia [Canada]; CNCI, Ottawa, designated by Bright 1967b: 678. Synonymy: Blackman 1928b: 132.

References: **(cn)** Barker 1949: 82; Browne 1968b: 566; Chamberlin 1924; Pettit 1928: 1–2; Swaine 1918a: 98, 103. **(hb)** Blackman 1919a, 1922b; Browne 1968b: 564; Swaine 1918a: 98, 103. **(ds)** Anonymous 1926c: 519; Blackman 1922b; Browne 1968b: 564; Chamberlin 1925; Hatch 1924; Kleine 1934a: 160; Leng 1920: 341; Leonard 1928: 519. **(tx)** Blackman 1919a: 85–96, 1919b: 137–139, 1922b: 101, 107–108, 1928b: 131–133; Bright 1967b: 678; Hoebeke 1978; de Ruette 1970: 109; Swaine 1917: 28, 1918a: 98, 103.

nudus Swaine 1917: 30. Lectotype ♂; Ste. Anne de Bellevue, Quebec [Canada]; CNCI, Ottawa, designated by Bright 1967b: 678. Synonymy: Wood 1977d: 515.

References: **(cn)** Blackman 1950; Chamberlin 1924; Swaine 1918a: 98, 104; Swaine & Craighead 1924: 1–27. **(hb)** Beal & Massey 1945; Blackman 1922b: 102, 108, 1950; Chamberlin 1939: 397; Swaine 1918a: 98, 104; Thomas, J. B. 1958: 394. **(ds)** Anonymous 1926c: 519; Beal & Massey 1945; Beaulne 1956; Blackman 1922b: 102, 108, 1950; Chamberlin 1925, 1939: 397; Dodge 1938; Hatch 1924; Kleine 1934a: 160; Knull 1932: 66; Leng 1920: 341; Leonard 1928: 519. **(tx)** Beal & Massey 1945; Blackman 1919b: 139–140, 1922b: 102, 108, 1928b: 113, 116; Bright 1967b: 678; Chamberlin 1939: 397; Dodge 1938: 43–45; Hoebeke 1978; de Ruette 1970: 110; Swaine 1917: 28, 30, 1918a: 98, 103–106; Wood, S. L. 1977d: 515.

conspectus Wood 1976c: 360. Holotype ♀; Volcan Irazú, Cartago, Costa Rica, 2300 m; Wood Collection. Distribution: North America (Costa Rica).

Hosts: *Quercus* sp.

References: **(ds)** Bright 1981d: 105; Wood, S. L.

- 1952b: 1053. (tx) Bright 1951d: 105; Wood, S. L. 1976c: 360, 1952b: 1053.
- coronarius Blackman** 1942a: 220. Holotype ♀; Jalisco, Mexico; USNM, Washington.
 Figures: Bright 1951d: 334.
 Distribution: North America (Jalisco in Mexico).
 Hosts: *Sambucus* sp.
 References: (hb) Burgos & Saucedo 1983: 122. (ds) Atkinson & Equihua 1955c: 359; Atkinson et al. 1956: 70; Blackwelder 1947: 781; Bright 1951d: 33, 1955a: 472; Burgos & Saucedo 1983: 122; Wood, S. L. 1942a: 220, 1952b: 1115. (tx) Blackman 1942a: 220; Bright 1951d: 33, 334.
- corruptus Wood** 1976c: 363. Holotype ♀; 5 km S Matamoros, Puebla, Mexico, 2000 m; Wood Collection.
 Distribution: North America (Puebla in Mexico).
 Hosts: *Toxicodendron* sp.
 References: (ds) Bright 1951d: 68, 1956a: 645, 1956b: 682. (tx) Bright 1951d: 65; McNamara 1984: 754; Wood, S. L. 1976c: 363, 1952b: 1133.
- cortezii Bright** 1977: 523. Holotype ♀; Ixtaccihuatl-Popocatepetl National Park, Mexico-Puebla border, Mexico; CNCI, Ottawa.
 Figures: Bright 1951d: 352.
 Distribution: North America (Mexico, Puebla in Mexico).
 Hosts: *Pinus hartwegii*.
 References: (hb) Atkinson & Equihua 1955c: 360; Atkinson et al. 1956a: 37. (ds) Atkinson & Equihua 1955a: 95, 1955c: 360, 1955: 96; Atkinson et al. 1956: 37; Bright 1951d: 172; Wood, S. L. 1952b: 1097. (tx) Bright 1977: 523, 1951d: 172, 352; McNamara 1984: 755; Wood, S. L. 1952b: 1097.
- corticalis Eichhoff** 1872a: 135. Holotype, sex?; Chili; Hamburg Museum, lost.
 Distribution: South America (Chile).
 References: (ds) Blackwelder 1947: 781; Gemminger & Harold 1872: 2688; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 332; Schedl 1972d: 144. (tx) Eichhoff 1872a: 135, 1878b: 191; Hagedorn 1910a: 100; Schedl 1938a: 183, 1951d: 17, 1951m: 112, 1972d: 144.
- costabilis Wood** 1976c: 352. Holotype ♀; 16 km W Tizapan, Jalisco, Mexico; Wood Collection.
 Distribution: North America (Jalisco).
 Hosts: *Thevetia* sp.
 References: (cc) Equihua & Atkinson 1956: 634. (hb) Atkinson et al. 1956: 70; Equihua & Atkinson 1956: 634. (ds) Atkinson et al. 1956: 70; Bright 1951d: 74, 1955b: 482; Equihua & Atkinson 1956: 634; Wood, S. L. 1952b: 1121. (tx) Bright 1951d: 74, 1955b: 482; Wood, S. L. 1976c: 352, 1952b: 1121.
- costalimai Blackman** 1942a: 223. Holotype ♀; Brazil; USNM, Washington.
 Figures: Blackman 1942a: fig. 34.
 Distribution: South America (Brazil).
 References: (cn) Costa Lima 1956. (hb) Costa Lima 1956. (ds) Blackwelder 1947; Costa Lima 1956. (tx) Blackman 1942a: 223–224; Costa Lima 1956; Schedl 1938a: 165, 169.
- costatulus Wood** 1976c: 351. Holotype ♀; 29 km W Niltpec, Oaxaca, Mexico; Wood Collection.
 Distribution: North America (Jalisco, Oaxaca in Mexico).
 Hosts: *Thevetia* sp.
 References: (cc) Equihua & Atkinson 1956: 634. (hb) Atkinson et al. 1956: 70; Burgos & Saucedo 1983: 122; Equihua & Atkinson 1956: 634. (ds) Atkinson et al. 1956: 70; Bright 1951d: 73, 1955b: 482; Burgos & Saucedo 1983: 122; Equihua & Atkinson 1956: 634; Wood, S. L. 1952b: 1120. (tx) Bright 1951d: 73; McNamara 1984: 755; Wood, S. L. 1976c: 351, 1952b: 1120.
- costatus Wood** 1975b: 395. Holotype ♀; Tapanti, Cartago, Costa Rica, 1800 m; Wood Collection.
 Distribution: North America (Costa Rica).
 Hosts: Liana.
 References: (ds) Bright 1951d: 47; Wood, S. L. 1952b: 1025. (tx) Bright 1951d: 47; McNamara 1984: 755; Wood, S. L. 1975b: 395, 1952b: 1025.
- costifera Bright** 1955b: 477. Holotype ♀; Mexico: Taxco, Guerrero; CNCI, Ottawa.
 Distribution: North America (Guerrero in Mexico).
 Hosts: Apocynaceae sp.
 References: (tx) Bright 1955b: 477.
- cracentis Bright** 1955b: 477. Holotype ♀; Mexico: Carr. Xoelicalco-Cuentepec, Km 6, Temixco, Moralia; CNCI, Ottawa.
 Distribution: North America (Morelos in Mexico).
 Hosts: Compositae sp.
 References: (tx) Bright 1955b: 477.
- crassus Blackman** 1928b: 67. Holotype ♀; Capitan, New Mexico [USA]; USNM, Washington.
 Figures: Bright 1951d: 368.
 Distribution: North America (Durango, Hidalgo, Mexico, Michoacan, Morelos, Nuevo Leon, Oaxaca, Puebla, Querteraro, Tlaxcala, Veracruz in Mexico; Arizona, Colorado, New Mexico, W Texas, Wyoming in USA).
 Hosts: *Pinus aristata*, *P. arizonica*, *P. ayacahuite*, *P. cembroides*, *P. contorta* (rare), *P. cooperi*, *P. culminicola*, *P. durangensis*, *P. edulis*, *P. engelmannii*, *P. flexilis*, *P. greggii*, *P. hartwegii*, *P. leiophylla*, *P. lumholtzii*, *P. montezumae*, *P. oocarpa*, *P. patula*, *P. ponderosa*, *P. strobiformis*, *P. teocote*.
 References: (hb) Atkinson et al. 1956: 37; Burgos & Saucedo 1983: 120; Chamberlin 1939: 376. (ds) Atkinson & Equihua 1955a: 97, 1955: 96; Atkinson et al. 1956: 37; Bright 1951c: 275; Burgos & Saucedo 1983: 120; Chamberlin 1939: 376; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 62; Rio Mora 1955: 11; Schedl 1963c: 158. (tx) Blackman 1928b: 60, 67–69, 1942a: 212–213;

Bright 1981d: 275, 368; Chamberlin 1939: 376; Keen 1929a: 32; Rio Mora 1985: 11; de Ruette 1970: 109; Wood, S. L. 1982b: 1081.

crinalis Blackman 1928b: 41. Holotype ♀; Washington, D.C. [USA]; USNM, Washington.

Figures: Bright 1981d: 338.

Distribution: North America (Connecticut, District of Columbia, Florida, Georgia, Illinois, Indiana, Kentucky, Maryland, Massachusetts, Michigan, Mississippi, North Carolina, Texas, West Virginia in USA).

Hosts: *Rhus toxicodendron*, *R. vernix*.

References: **(bv)** Turnbow & Franklin 1980. **(hb)**

Baker, W. L. 1972: 255; Beal & Massey 1945: 128; Chamberlin 1939: 362; Deyrup & Atkinson 1987a: 66. **(ds)** Beal & Massey 1945: 128; Bright 1981d: 66; Chamberlin 1939: 362; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Drooz 1985: 363; Favinger & Wade 1973; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Turnbow & Franklin 1980; Wood, S. L. 1982b: 1134. **(tx)** Beal & Massey 1945: 128; Blackman 1928b: 41; Bright 1981d: 66, 338; Chamberlin 1939: 362; McNamara 1984: 755; Weber, B. C., & McPherson 1991: 54; Wood, S. L. 1982b: 1134.

cristatus Wood 1964: 68. Holotype ♀; 14 km N Perote, Veracruz, Mexico; Wood Collection.

Figures: Bright 1981d: 118, 343.

Distribution: North America (Durango, Hidalgo, Mexico, Michoacan, Oaxaca, Veracruz in Mexico/ Arizona in USA).

Hosts: *Pinus ayacahuite*, *P. engelmannii*, *P. leiophylla*, *P. lumholtzii*, *P. spp.*

References: **(hb)** Atkinson & Equihua 1985c: 359; Atkinson et al. 1986: 37; Burgos & Sancedo 1983: 118. **(ds)** Atkinson & Equihua 1985c: 359, 1985: 96; Atkinson et al. 1986: 37; Bright 1971b: 68, 1981d: 114; Burgos & Sancedo 1983: 118; Rio Mora 1985: 13; Thomas, J. B. 1966; Wood, S. L. 1982b: 1067. **(tx)** Bright 1981d: 114, 118, 343; Rio Mora 1985: 13; Wood, S. L. 1964: 68, 1982b: 1067.

crotonis Wood 1977d: 517. Holotype ♂; 30 km E Merida, Merida, Venezuela, 2500 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Croton* sp.

References: **(tx)** Wood, S. L. 1977d: 517.

culminicola Bright 1977: 524. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico; CNCI, Ottawa.

Figures: Bright 1981d: 359.

Distribution: North America (Nuevo Leon in Mexico).

Hosts: *Pinus culminicola*.

References: **(ds)** Bright 1981d: 215; Wood, S. L. 1982b: 1036. **(tx)** Bright 1977: 524, 1981d: 215, 359; McNamara 1984: 755; Wood, S. L. 1982b: 1036.

cuspidatus Blackman 1942a: 217. Holotype ♀; Jacala, Hidalgo, Mexico; USNM, Washington.

Distribution: North America (Durango, Hidalgo, Mexico, Puebla, Veracruz in Mexico/ S Arizona in USA).

Hosts: *Pinus ayacahuite*, *P. lawsonii*, *P. lumholtzii*, *P. ponderosa*, *P. strobiformis*.

References: **(ds)** Atkinson & Equihua 1985c: 98, 1988: 96; Blackwelder 1947: 781; Bright 1981d: 324. **(tx)** Blackman 1942a: 217; Bright 1981d: 324; McNamara 1984: 755.

debilis Wood 1976a: 354. Holotype ♀; San Ignacio de Acosta, San Jose, Costa Rica; Wood Collection. Distribution: North America (Costa Rica/ Chiapas in Mexico).

Hosts: *Mauria glauca*, *Rhus* sp., and a tree branch.

References: **(ds)** Bright 1981d: 15; Wood, S. L. 1982b: 1122. **(tx)** Bright 1981d: 15; McNamara 1984: 755; Wood, S. L. 1976a: 354, 1982b: 1122.

declivisetosus Bright 1977: 525. Holotype ♀; 51 km SE Nochistlan, Oaxaca, Mexico; CNCI, Ottawa.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Pinus pringlei*, *P. sp.*

References: **(ds)** Bright 1981d: 251; Wood, S. L. 1982b: 1062. **(tx)** Bright 1977: 525, 1981d: 251; McNamara 1984: 755; Wood, S. L. 1982b: 1062.

degener Wood 1975b: 397. Holotype ♂; Volcan Chiriqui (near Cerro Punta), Chiriqui, Panama, 1800 m; Wood Collection.

Distribution: North America (Panama).

Hosts: Tree limb.

References: **(ds)** Bright 1981d: 96; Wood, S. L. 1982b: 1029. **(tx)** Bright 1981d: 96; Wood, S. L. 1975b: 397, 1982b: 1029.

deleoni (Blackman) 1942a: 201 (*Myeloborus*). Holotype ♀; El Seco, Puebla, Mexico; USNM, Washington.

Distribution: North America (Nuevo Leon, Puebla in Mexico).

Hosts: *Pinus* sp.

Notes: (1) Bright 1976c: 427 (to *Pityophthorus*).

References: **(hb)** Bright 1981d: 202; Bright & Stark 1973: 109. **(ds)** Blackwelder 1947: 781; Bright 1981d: 202; Bright & Stark 1973: 109; Wood, S. L. 1982b: 1044. **(tx)** Blackman 1942a: 201; Bright 1976c: 427, 1981d: 202; Wood, S. L. 1982b: 1044.

deletus LeConte 1879: 519. Lectotype, sex?, Veta Pass, Colorado [USA]; MCZ, Cambridge, designated by Bright 1976b: 185.

Figures: Bright 1981d: 343.

Distribution: North America (Coahuila, Durango in Mexico/ Arizona, California, Colorado, Nevada, New Mexico, South Dakota, W Texas, Utah, Wyoming in USA).

Hosts: *Pinus cembroides*, *P. edulis*, *P. monophylla*, uncommon in *P. flexilis*, *P. ponderosa*, *P. strobiformis*.

Notes: (3) Blackman 1928b: 43 (redescribed).

References: **(cn)** Swaine 1915a: 104. **(hb)** Cham-

berlin 1939: 364; Furniss, M. M. & Johnson 1987: 377; Swaine 1918a: 104. **(ds)** Atkinson & Equihua 1988: 96; Bright 1981d: 110, 1986b: 682; Chamberlin 1939: 364; Furniss, M. M. & Johnson 1987: 377; Hagedorn 1910d: 71; Henshaw 1882: 268, 1885: 147; Keen 1929a: 32; Kleine 1913b: 140, 1934a: 159; Leng 1920: 341; Swaine 1909: 135; Wickham 1896a: 309; Wood, S. L. 1982b: 1040. **(tx)** Blackman 1928b: 43–46; Bright 1976b: 185, 1977: 515, 1981d: 110, 343; Chamberlin 1939: 364; Hagedorn 1910a: 100; Keen 1929a: 32; LeConte 1879: 519; Swaine 1909: 135, 1918a: 104; Wood, S. L. 1977c: 387, 1982b: 1040.

inquietus Blackman 1928b: 46. Holotype ♀; Las Vegas H.S. [Hot Springs], New Mexico [USA]; USNM, Washington. Synonymy: Wood 1977c: 387.

References: **(hb)** Chamberlin 1939: 364. **(ds)** Chamberlin 1939: 364; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52; Wood, S. L. 1948: 51, 1951a: 128, 1972a: 425. **(tx)** Blackman 1928b: 43–47; Chamberlin 1939: 364; Keen 1929a: 32; de Ruette 1970: 109; Wood, S. L. 1951a: 128, 1972a: 425, 1977c: 387.

monophyllae Blackman 1928b: 47. Holotype ♀; Argus Mts., California [USA]; USNM, Washington. Synonymy: Wood 1977c: 387.

References: **(hb)** Bright & Stark 1973: 107; Chamberlin 1939: 365. **(ds)** Bright & Stark 1973: 107; Chamberlin 1939: 365; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 47; Bright 1971b: 67; Chamberlin 1939: 365; Keen 1929a: 32; Wood, S. L. 1964: 65, 1977c: 387.

socius Blackman 1928b: 48. Holotype ♂; Argus Mts., California [USA]; USNM, Washington. Synonymy: Bright 1971b: 67.

References: **(hb)** Chamberlin 1939: 365. **(ds)** Chamberlin 1939: 365; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 48; Bright 1971b: 67; Chamberlin 1939: 365; Keen 1929a: 32; McNamara 1984: 758.

piceus Bright 1966b: 297. Holotype ♀; Mount Pinos, Ventura Co., California [USA]; CAS, San Francisco. Synonymy: Bright 1971b: 67.

References: **(tx)** Bright 1966b: 297, 1971b: 67.

praealtus Bright 1966b: 303. Holotype ♀; Mt. Shasta Ski Area, Siskiyou Co., California [USA]; CAS, San Francisco. Synonymy: Wood 1977c: 387.

References: **(hb)** Bright & Stark 1973: 106. **(ds)** Bright & Stark 1973: 106. **(tx)** Bright 1966b: 303; de Ruette 1970: 110; Wood, S. L. 1977c: 387.

brucki Bright 1971b: 63. Holotype ♂; Mt. Hawkins, San Bernardino Co., California [USA]; OSUC, Columbus. Synonymy: Wood 1977c: 388.

References: **(hb)** Bright & Stark 1973: 106. **(ds)** Bright & Stark 1973: 106. **(tx)** Bright 1971b: 63–64; Wood, S. L. 1977c: 387–388.

delicatus Wood 1978b: 399. Holotype ♀; 56 km SW El Salto, Durango, Mexico; Wood Collection. Distribution: North America (Guatemala/ Honduras/ Chiapas, Chihuahua, Durango, Mexico, Michoacan in Mexico).

Hosts: *Pinus ayacahuite*, *P. leiophylla*, *P. oocarpa*, *P. pseudostrobus*, *P. tenuifolia*, *P.* spp.

References: **(cn)** Schwerdtfeger 1956b: 48. **(hb)** Schwerdtfeger 1956b: 48. **(ds)** Atkinson & Equihua 1988: 96; Bright 1981d: 323; Wood, S. L. 1982b: 1109. **(tx)** Bright 1981d: 323; Wood, S. L. 1978b: 399, 1982b: 1109.

denticulatus (Wichmann) 1915a: 106 (*Trigonogenius*). Holotype ♂; Deutsch-Ostafrika: MNB, Berlin.

Distribution: Africa (Tanzania).

Hosts: Tree.

References: **(tx)** Eggers 1940a: 30; Lucas 1920: 683; Nurnberg 1956: 205; Schedl 1952d: 347, 1957e: 878, 1958k: 144, 1960i: 106, 1962j: 7–8; Wichmann 1915a: 106–107.

dentifrons Blackman 1922c: 125. Holotype ♀; Orono, Maine [USA]; USNM, Washington.

Figures: Blackman 1922c: figs. 22–23.

Distribution: North America (Alberta, New Brunswick, Newfoundland, Nova Scotia, Ontario, Prince Edward Island, Quebec in Canada/ Maine, Maryland, Minnesota, New Hampshire, New York, North Carolina, West Virginia in USA).

Hosts: *Abies fraseri*, *Picea glauca*, *P. rubens*, *Pinus banksiana*, *P. strobus*.

References: **(cn)** Blackman 1950. **(hb)** Baker, W. L. 1972: 255; Beal & Massey 1945: 134; Blackman 1950; Chamberlin 1939: 379. **(ds)** Anonymous 1926c: 519; Baker, W. L. 1972: 255; Beal & Massey 1945: 134; Blackman 1950; Bright 1971a: 127, 1976d: 186, 1981d: 226; Chamberlin 1925, 1939: 379; Drooz 1985: 364; Kleine 1934a: 159; Leng & Mutchler 1927: 52; Leonard 1928: 519; Wood, S. L. 1982b: 1072. **(tx)** Beal & Massey 1945: 134; Blackman 1922c: 125–126, 1928b: 62, 81; Bright 1981d: 226; Chamberlin 1939: 379; de Ruette 1970: 109; Wood, S. L. 1982b: 1072.

deodara (Stebbing) 1903: 274 (*Cryphalus*). Syn-types ♀; India: N.W. Himalayas, Bashahr Div., Kotgarh, Nagkela Forest, Jaunsar Div., Konain; FRI, Dehra Dun.

Distribution: Asia (Himachal Pradesh, Kashmir, and Uttar Pradesh in India).

Hosts: *Abies pindrow*, *A. webbiana*, *Cedrus deodara*, *Pinus excelsa*, *P. roxburghii*.

Notes: (3) Beeson 1941 (1961: 292) (cited *chir*, nomen nudum, *pindrow*, nomen nudum, synonyms).

References: **(cn)** Pierce, W. D. 1917: 52; Stebbing 1903a: 274, 1914: 528. **(cc)** Stebbing 1914: 528.

- (**hb**) Stebbing 1903a: 274, 1914: 528. (**ds**) Beeson 1961: 292; Hagedorn 1910d: 41; Kleine 1913b: 118, 1914b: 269, 276, 1934a: 159; Pierce, W. D. 1917: 52; Stebbing 1903a: 274. (**tx**) Hagedorn 1910a: 86; Stebbing 1903: 274, 1914: 528.
- himalayensis* Stebbing 1914: 540 (*Cryptahus*).
 Syntypes, sex?: Jamnagar, Simla Forests, North-West Himalaya; FRI, Dehra Dun. Synonymy: Wood 1989: 175.
 References: (**cn**) Pierce, W. D. 1917: 52; Stebbing 1914: 540. (**cc**) Stebbing 1914: 540. (**hb**) Stebbing 1914: 540. (**ds**) Beeson 1961: 292; Pierce, W. D. 1917: 52; Schedl 1974a: 86. (**tx**) Stebbing 1914: 540; Wood, S. L. 1989: 175.
- sampsoni* Stebbing 1914: 551. Syntypes, sex?: Jamnagar, North-West Himalaya; FRI, Dehra Dun, lost. Synonymy: Wood 1989: 175.
 References: (**cn**) Browne 1968: 565; Pierce, W. D. 1917: 74; Stebbing 1914: 551. (**cc**) Beeson 1922c: 497; Stebbing 1914: 551. (**hb**) Beeson 1922c: 497; Browne 1968: 565; Stebbing 1914: 551. (**ds**) Beeson 1922c: 497, 1961: 292; Browne 1968: 565; Pierce, W. D. 1917: 74; Zetlner 1973. (**tx**) Stebbing 1914: 551; Wood, S. L. 1989: 175.
- deprecator* Schaufuss 1891: 15. Syntypes 2, sex?: Madagascar; Hamburg Museum, lost.
 Distribution: Madagascar.
 References: (**ds**) Alluaud 1900: 441; Anonymous 1892: 165; Fairmaire 1892b; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 326; Schedl 1977b: 111. (**tx**) Fairmaire 1892b; Hagedorn 1910a: 100, 1913b: 244; Schaufuss 1891: 15; Schedl 1938a: 183, 1977b: 111.
- desultorius* Bright 1985b: 478. Holotype ♀; Mexico: Oriental, Puebla; CNCI, Ottawa.
 Distribution: North America (Puebla in Mexico).
 Hosts: Compositae sp.
 References: (**tx**) Bright 1985b: 478.
- detectus* Schedl 1972g: 66. Holotype ♀; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (**tx**) Schedl 1972g: 66.
- detentus* Wood 1976a: 352. Holotype ♀; Carapan, Michoacan, Mexico, 2300 m; Wood Collection.
 Distribution: North America (Michoacan, Queretaro in Mexico).
 Hosts: *Toxicodendron* sp., and a liana.
 References: (**ds**) Bright 1981d: 82; Wood, S. L. 1982b: 112. (**tx**) Bright 1981d: 82; McNamara 1984: 755; Wood, S. L. 1976a: 352, 1982b: 1121.
- digestus* (LeConte) 1874a: 71 (*Cryptahus*). Lectotype, sex?: Mojave Desert [USA]; MCZ, Cambridge, designated by Bright 1976b: 185.
 Figures: Bright 1981d: 366.
 Distribution: North America (Alberta, S British Columbia in Canada/ Durango in Mexico/ Arizona, California, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, Wyoming in USA).
 Hosts: *Pinus contorta*, *P. jeffreyi*, *P. ponderosa*, *P. washoensis*.
 Notes: (1) LeConte 1876: 355 (to *Pityophthorus*). (3) Blackman 1928b: 54 (re-described).
 References: (**hb**) Bright & Stark 1973: 108; Chamberlin 1939: 368. (**ds**) Blaisdell 1892: 36; Bright 1981d: 261; Bright & Stark 1973: 108; Chamberlin 1939: 368; Gast et al 1959: 355; Hagedorn 1910d: 71; Henshaw 1882: 269, 1885: 147; Keen 1929a: 32; Kleine 1913b: 140, 1914b: 391, 1934a: 159; Leng 1920: 341; Swaine 1909: 135; Wood, S. L. 1982b: 1032. (**tx**) Blackman 1928b: 53–55; Bright 1976b: 185, 1977: 514, 1981d: 261, 366; Chamberlin 1939: 368; Hagedorn 1910a: 100; Keen 1929a: 32; LeConte 1874a: 71, 1876: 354–355; Swaine 1909: 135; Wood, S. L. 1978b: 395, 1982b: 1032.
- idoneus* Blackman 1928b: 55. Holotype ♀; Centerville, Idaho [USA]; USNM, Washington. Synonymy: Bright 1977: 514.
 References: (**cn**) Anonymous 1964h; Lindgren 1980a: 68. (**hb**) Bright & Stark 1973: 108; Chamberlin 1939: 368, 1955: 149; Lindgren 1980a: 68. (**ds**) Anonymous 1964h; Bright 1976d: 182; Bright & Stark 1973: 108; Chamberlin 1939: 368; Evans, D. 1983: 35; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 425. (**tx**) Blackman 1928b: 55; Bright 1966b: 304, 1977: 514; Chamberlin 1939: 368, 1955: 149; Evans, D. 1983: 35; Keen 1929a: 32; Wood, S. L. 1972a: 425.
- hopkinsi* Blackman 1928b: 56. Holotype ♀; California: Ventura Co. [USA]; USNM, Washington. Synonymy: Bright 1966b: 304.
 References: (**cc**) Marsh 1979: 150. (**hb**) Chamberlin 1939: 368. (**ds**) Chamberlin 1939: 368; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52. (**tx**) Blackman 1928b: 56; Bright 1966b: 304, 1977: 514; Chamberlin 1939: 368; Keen 1929a: 32.
- ponderosae* Blackman 1928b: 57. Holotype ♀; Las Vegas H.S. [Hot Springs], New Mexico [USA]; USNM, Washington. Synonymy: Bright 1966b: 304.
 References: (**cn**) Anonymous 1967f. (**hb**) Chamberlin 1939: 369, 1958: 150. (**ds**) Anonymous 1967f; Chamberlin 1939: 369, 1958: 150; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Smith, G. S. 1930. (**tx**) Blackman 1928b: 57; Bright 1966b: 304, 1977: 514; Chamberlin 1939: 369, 1958: 150; Hoebeke 1978; Keen 1929a: 32; de Ruelle 1970: 110; Schedl 1930: 195; Wood, S. L. 1967d: 39.
- aplanatus* Schedl 1930: 195. Holotype ♀; Athabasca Falls, Alberta [Canada]; CNCI, Ottawa. Synonymy: Wood 1978b: 395.

Notes: (3) Bright 1981d: 212 (a good species). Wood 1952b: 1032 (a character cline from Alberta to eastern Wyoming intergrades with *digestus*).

References: (cn) Lindgren, B. 1980a: 67. (hb) Chamberlin 1939: 369; Lindgren, B. 1980a: 67. (ds) Bright 1976d: 180, 1981d: 212; Chamberlin 1939: 369; Evans, D. 1983: 34; Leng & Mutchler 1933: 52. (tx) Bright 1981d: 212, 352; Chamberlin 1939: 369; Evans, D. 1983: 34; de Ruelle 1970: 108; Schedl 1930: 195; Wood, S. L. 1975: 395.

***diglyphus* Blandford** 1904: 240. Lectotype, sex?; Quiche Mts., Guatemala; BMNH, London, designated by Bright 1976b: 184.

Distribution: North America (Guatemala).

Hosts: *Pinus pseudostrobus*.

References: (hb) Atkinson et al. 1986: 38; Burgos & Sancedo 1983: 118. (ds) Atkinson et al. 1986: 38; Blackwelder 1947: 781; Bright 1981d: 121, 1985b: 683; Burgos & Sancedo 1983: 118; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 362; Wood, S. L. 1982b: 1058. (tx) Blandford 1904: 240; Bright 1976b: 184, 1977: 515, 1981d: 121; Hagedorn 1910a: 100; Schedl 1935a: 183, 1955g: 18; Wood, S. L. 1982b: 1058.

***diligens* Wood** 1976a: 363. Holotype ♀; 16 km E Pachuca, Hidalgo, Mexico; Wood Collection.

Distribution: North America (Hidalgo in Mexico).

Hosts: Desert shrub, *Zexmenia* sp.

References: (ds) Atkinson & Equihua 1985a: 92; Bright 1981c: 77, 1982b: 482, 1985b: 482; Wood, S. L. 1982b: 1132. (tx) Bright 1981d: 77; McNamara 1984: 755; Wood, S. L. 1976a: 363, 1982b: 1132.

***dimidiatus* Blackman** 1942a: 221. Holotype ♀; Mexico [intercepted at USA port of entry]; USNM, Washington.

Distribution: North America ("Mexico").

References: (ds) Blackwelder 1947: 781; Bright 1981d: 41; Wood, S. L. 1982b: 1138. (tx) Blackman 1942a: 221; Bright 1981c: 41; McNamara 1984: 755; Wood, S. L. 1982b: 1138.

***diminuticus* Bright** 1985a: 468. Holotype ♀; Mexico: Estacion de Biologia, Chamela, Edo. Jalisco; CNCI, Ottawa.

Distribution: North America (Jalisco in Mexico).

Hosts: Leguminosae sp.

References: (tx) Bright 1985a: 468.

***dimorphus* Schedl** 1959m: 551. Holotype ♀; Matto Grosso: Rio Caraguata; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1959m: 551, 1979c: 81.

***brasilensis* Schedl** 1954b: 38. Syntypes ♀; Brasilien: Matto Grosso, Rio Caraguata; Schedl Collection in NHMW, Wien, and Plannmann Collection, preoccupied by Schedl

1938. Synonymy: Schedl 1979c: 81, Wood 1984b: 227.

Notes: (1) Schedl 1979c: 45 (citation of holotype invalid).

References: (tx) Schedl 1954b: 38, 1979c: 45, 81; Wood, S. L. 1984b: 227.

***discretus* Wood** 1977c: 394. Holotype ♀; 5 km W El Salto, Durango, Mexico, 2500 m; Wood Collection.

Distribution: North America (Durango, Jalisco, Michoacan, Puebla in Mexico).

Hosts: *Pinus ayacahuite*, *P. sp.*

References: (ds) Bright 1981d: 283; Rio Mora 1985: 14; Wood, S. L. 1982b: 1098. (tx) Bright 1981d: 283; Wood, S. L. 1977c: 394, 1982b: 1098.

***dispar* Bright** 1976c: 431. Holotype ♀; 11 km E San Cristobal, Chiapas, Mexico; CNCI, Ottawa.

Figures: Bright 1981d: 341.

Distribution: North America (Chiapas in Mexico).

Hosts: *Pinus montezuma*, *P. ochoterenai*.

References: (ds) Bright 1981d: 91; Wood, S. L. 1982b: 1037. (tx) Bright 1976c: 431, 1981d: 91, 341; McNamara 1984: 755; Wood, S. L. 1982b: 1037.

***dissolutus* Wood** 1975b: 398. Holotype ♀; 13 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.

Distribution: North America (Costa Rica/ Panama).

Hosts: Liana and a tree seedling.

References: (ds) Bright 1981d: 98; Wood, S. L. 1982b: 1031. (tx) Bright 1981d: 98; McNamara 1984: 755; Wood, S. L. 1975b: 398, 1982b: 1031.

***diversus* Bright** 1972d: 87. Holotype ♂; Cornwall Mountain, Westmoreland Parish, Jamaica; CNCI, Ottawa.

Distribution: Antilles Islands (Jamaica).

References: (ds) Bright 1985c: 176. (tx) Bright 1972d: 87, 1985c: 176; McNamara 1977: 197.

***djuguensis* Eggers** 1940b: 103. Holotype, sex?; Congostaat: Ituri (Djugu); MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Eggers 1940b: 103–104; Schedl 1962j: 8.

***ituriensis* Eggers** 1940b: 104. Holotype, sex?; Congostaat: Ituri (Djugu); MRCB, Tervuren.

Synonymy: Schedl 1962j: 8.

References: (tx) Eggers 1940b: 104; Schedl 1962j: 8.

***dolus* Wood** 1964: 65. Holotype ♀; McCloud, Siskiyou Co., California [USA]; Wood Collection.

Distribution: North America (California in USA).

Hosts: *Pinus lambertiana*, *P. ponderosa*.

Notes: (3) Bright 1981d: 110 (treated this name as a synonym of *deletus*).

References: (hb) Bright & Stark 1973: 107. (ds) Bright 1981d: 110; Bright & Stark 1973: 107; Wood, S. L. 1982b: 1039. (tx) Wood, S. L. 1964: 65, 1982b: 1039.

dorsalis Schedl 1953d: 94. Syntypes ♂ ♀; Madagascar: Mt. d'Ambre; MNHN, Paris and Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

Hosts: *Dichaetanthera cordifolia*.

References: (hb) Schedl 1977b: 111. (ds) Schedl 1977b: 111. (tx) Schedl 1953d: 94, 1965c: 69, 1977b: 111, 1979c: 83.

durus Blackman 1928b: 70. Holotype ♀; Paradise, Arizona [USA]; USNM, Washington.

Figures: Bright 1981d: 364.

Distribution: North America (Chihuahua, Durango, Hidalgo, Mexico, Michoacan, Morelos, Nuevo Leon, Oaxaca, Puebla, Sonora, Veracruz in Mexico/ Arizona in USA).

Hosts: *Pinus lawsoni*, *P. leiophylla*, *P. michoacana*, *P. pringlei*, *P. strobfiformis*, *P. sp.*

References: (hb) Atkinson et al. 1986: 38; Chamberlin 1939: 377. (ds) Atkinson & Equihua 1985a: 96, 1988: 96; Atkinson et al. 1986: 38; Bright 1981d: 252; Chamberlin 1939: 377; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Rio Mora 1985: 13; Thomas, J. B. 1966; Wood, S. L. 1982b: 1060. (tx) Blackman 1928b: 60, 70–71; Bright 1981d: 252, 364; Chamberlin 1939: 377; Rio Mora 1985: 13; Wood, S. L. 1982b: 1060.

eggersi Schedl 1952d: 347. Holotype ♀?; Kamerun; Schreiner Collection in Hamburg Museum to Eggers Collection, now in NHMW, Wien, automatic. Distribution: Africa (Cameroon).

References: (ds) Schedl 1966c: 225. (tx) Schedl 1952d: 347, 1962j: 8.

similis Eggers 1920: 34 (*Trigonogenius*). Holotype ♀?; Kamerun; Schreiner Collection in Hamburg Museum to Eggers Collection, in NHMW, Wien, preoccupied by Eichhoff 1869. References: (ds) Blackwelder 1947: 782; Hagedorn 1910d: 75. (tx) Eggers 1920: 34–35; Hagedorn 1910a: 101; Lucas 1920: 683; Schedl 1952d: 347, 1979c: 230.

elatinus Wood 1964: 66. Holotype ♀; 8 km W Ciudad Hidalgo, Michoacan, Mexico; Wood Collection.

Figures: Atkinson et al. 1986: 33, Bright 1981d: 345.

Distribution: North America (Mexico, Michoacan, Tlaxcala in Mexico).

Hosts: *Abies religiosa*.

References: (hb) Atkinson & Equihua 1985a: 93; Atkinson et al. 1986: 38; Burgos & Saucedo 1983: 119. (ds) Atkinson & Equihua 1985a: 93, 1988: 96; Atkinson et al. 1986: 38; Bright 1981d: 125; Burgos & Saucedo 1983: 119; Wood, S. L. 1982b: 1072. (tx) Atkinson et al. 1986: 33; Bright 1981d: 125, 345; Wood, S. L. 1964: 66, 1982b: 1072.

electus Blackman 1928b: 140. Holotype ♀; Ashland, Oregon [USA]; USNM, Washington.

Distribution: North America (California, Oregon in USA).

Hosts: *Pinus coulteri*, *P. jeffreyi*, *P. ponderosa*.

References: (hb) Chamberlin 1939: 401, 1958: 161. (ds) Bright 1981d: 254; Chamberlin 1939: 401, 1958: 161; Keen 1929a: 34; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 427, 1982b: 1066. (tx) Blackman 1928b: 140; Bright 1981d: 254; Chamberlin 1939: 401, 1958: 161; Keen 1929a: 34; Wood, S. L. 1972a: 427, 1982b: 1066.

elegans Schedl 1935a: 184. Holotype ♂; Guatemala; USNM, Washington.

Distribution: North America (Guatemala).

References: (ds) Blackwelder 1947: 781; Bright 1981d: 32; Wood, S. L. 1982b: 1028. (tx) Anderson, W. H. & Anderson 1971: 12; Bright 1981d: 32; Schedl 1935a: 184–185; Wood, S. L. 1982b: 1028.

elongatulus Schedl 1976a: 67. Holotype ♀; Tapuruquara, Rio Negro [Brazil]; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 67.

equihuai Bright 1985a: 469. Holotype ♀; Mexico: Estacion de Biologia, Chamela, Edo. Jalisco; CNCI, Ottawa.

Distribution: North America (Jalisco in Mexico).

References: (tx) Bright 1985a: 469.

erraticus Schedl 1976a: 68. Holotype ♀; Jacareacanga, Para, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 68.

espinosai Brethes 1925: 202. Syntypes, sex?; Santiago, Chile; Brethes Collection in MACN, Buenos Aires.

Distribution: South America (Chile).

References: (ds) Blackwelder 1947: 781; Schedl 1972d: 144. (tx) Brethes 1925: 202; Porter 1932: 106; Schedl 1972d: 144.

euterpes Bright 1978: 75. Holotype ♂; 35 km W Lazaro Cardenas, Chiapas, Mexico; CNCI, Ottawa.

Distribution: North America (Chiapas in Mexico).

Hosts: *Pinus oocarpa*.

References: (ds) Atkinson & Equihua 1985: 96; Bright 1981d: 52; Wood, S. L. 1982b: 1128. (tx) Bright 1978: 75, 1981d: 52; McNamara 1984: 755; Wood, S. L. 1982b: 1128.

excellens Schedl 1972g: 66. Holotype ♀; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1972g: 66, 1979c: 93.

eximius Schedl 1935a: 184. Holotype ♀; Kongo [Lekimi, Schedl 1962j: 9]; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

References: (tx) Schedl 1938a: 184, 1962j: 9, 1979c: 94.

explicitus Wood 1975b: 399. Holotype ♂; 9 km NE Teziutlan, Puebla, Mexico, 1600 m; Wood Collection.

Distribution: North America (Puebla in Mexico).

Hosts: Liana.

References: (ds) Bright 1981d: 98; Wood, S. L. 1982b: 1031. (tx) Bright 1981d: 98; McNamara 1984: 755; Wood, S. L. 1975b: 399, 1982b: 1031.

exquisitus (Blackman) 1942a: 196 (*Neodryocotes*). Holotype ♀; Mexico: USNM, Washington.

Figures: Blackman 1942a: fig. 21, Bright 1981d: 342.

Distribution: Antilles Islands (Virgin Islands), North America (Jalisco, Michoacan in Mexico).

Hosts: *Sambucus* sp.

Notes: (1) Bright 1981d: 102 (to *Pityophthorus*).

References: (hb) Atkinson et al. 1986: 38; (ds) Atkinson et al. 1986: 38, 72; Blackman 1942a: 196–197; Bright 1981d: 102, 176, 1986b: 683; Wood, S. L. 1982b: 1026. (tx) Blackman 1942a: 196–197; Bright 1981d: 102, 176, 342.

inceptus Wood 1975b: 396. Holotype ♀; 6 km W Quiroga, Michoacan, Mexico; Wood Collection. Synonymy: Bright 1981d: 102, Wood 1982b: 1026.

References: (tx) Bright 1981d: 102; Wood, S. L. 1975b: 396, 1982b: 1026.

exsculptus (Ratzburg) 1837: 162. Syntypes, sex?; Europe; not located.

Distribution: Europe (Austria/ Bulgaria/ Czechoslovakia/ France/ Germany/ Hungary/ Poland/ Switzerland/ W USSR/ Yugoslavia).

Hosts: *Picea excelsa*, rare in *Pinus* spp.

References: (ay) Escherich 1923b: 483, 555, 602; Nusslin 1911a: 129. (bv) Chararas 1977b; Grune 1979: 115; Meixner 1937: 1214. (cn) Eckstein 1926: 579; Escherich 1923b: 483, 555, 602; Gabler 1955; Joly 1976; Judeich & Nitsche 1895: 526; Nosek 1951: 109; Nusslin 1913: 205; Pfeffer 1928a: 468, 1950c: 2; Pierce, W. D. 1917: 81; Rhumbler 1922: 308, 1927: 322; Schimitschek 1937c: 55, 1955a: 59, 1955c: 87; Schwerdtfeger 1944a: 181, 1957a: 186; Wachtl 1901: 381; Weber 1926: 579; Wichmann 1927b: 358. (ce) Balazy & Michalski 1964b; Kleine 1908c: 213; Michalski & Ratajczak 1989; Nosek 1951: 109; Nusslin 1927: 322; Pfeffer 1923a: 331, 1950c: 2, 1959: 2, 1960: 344; Schimitschek 1955a: 59; Schwerdtfeger 1944a: 181, 1957a: 186; Sedlaczek 1935a: 163; Zimm 1985b. (hb) Chararas 1977b; Charvat 1950; Eckstein 1926: 579; Escherich 1923b: 483, 555, 602; Fuchs 1904a; Gabler 1955; Györfi 1957; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski & Strawinski 1948: 156; Lenggerken 1939: 65, 1954: 85; Lindemann 1881a: 236; Nusslin 1913: 205, 1927: 322; Pfeffer 1941a: 11, 1989a: 63; Postner 1974: 433; Ratzburg 1837: 162, 1839: 197;

Rhumbler 1922: 308, 1927: 322; Schimitschek 1955a: 59; Schwerdtfeger 1944a: 181, 1957a: 186, 1981: 193; Sedlaczek 1935a: 163; Spessivtsev 1913a: 87, 1921b: 221; Tredl 1908b: 138, 1915a: 148; Wachtl 1876a: 460, 1901: 381; Weber, H. 1926: 579; Wichmann 1927b: 358. (ds) Ammann & Knabl 1923; Andras & Schaefer 1957; Buresh & Lazarov 1956; Calwer 1884, 1893; Chararas 1977b; Charvat 1950; Endrodi 1958b; Escherich 1923b: 483, 555, 602, 1932b; Fuchs 1904a, 1905a; Gaubil 1849: 126; Gemminger & Harold 1872: 2688; Grune 1979: 115; Hagedorn 1910d: 71; Heyden, Reitter & Weise 1883: 182, 1891: 672, 1906: 712; Horion 1951; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski 1926: 82, 1931: 30, 1933a: 290; Karpinski & Strawinski 1948: 156; Kleine 1912a: 268, 1913a: 35, 1913b: 140, 1934a: 159; Kolbe, W. 1918: 211; Kraatz 1869: 59; Kurir 1947c: 6, 14; Langhoffer 1915c: 158; Leder 1871: 132; Lomnicki 1913b: 148; Lucht 1987: 278; Nunberg 1928b: 88, 112, 1938: 126, 1954: 70, 1960b: 158; Pacher 1865: 152; Pfeffer 1931b: 74, 1935: 159, 1947e: 4, 1950b: 75, 1960: 345, 1976: 340, 1989a: 63; Pierce, W. D. 1917: 81; Pittioni 1943: 176; Postner 1974: 433; Prossen 1913: 83; Ratzburg 1837: 162, 1839: 197; Reitter 1894a: 76, 1916: 298; Roubal 1941: 269; Sainte-Claire & Mequignon 1938: 448; Schaufuss 1915: 1242; Schamm 1859: 95, 1862: 101; Schedl 1980a: 22; Schilsky 1909: 188; Schwerdtfeger 1981: 193; Seidlitz 1872: 394; Stein 1868: 114; Stein & Weise 1877: 164; Tredl 1907: 14, 1908b: 138; Wachtl 1876a: 460; Wichmann 1927a: 74. (tx) Bach 1854; Balachowsky 1949a: 234; Charvat 1950; Eggers 1914, 1915b: 13–14, 1920; Eichhoff 1864: 39; Endrodi 1957a: 307, 1957b; Escherich 1923b: 483, 555, 602; Ferrari 1867: 32, 1867b; Fleischer 1927; Formanek 1907: 38; Gabler 1955; Grune 1979: 115; Hagedorn 1910a: 101; Jansson 1936: 220; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski & Strawinski 1948: 156; Koch 1928: 82, 1932: 122; Kuhn 1913: 1056; Letzner 1891: 376; Lindemann 1881a: 236; Lucht 1987: 278; Meixner 1937: 1214; Nunberg 1938: 126, 1954: 70; Nusslin 1911a: 129; Postner 1974: 433; Quaschik 1953: 35; Pfeffer 1932f: 23, 1940b: 107, 110, 1941c: 11, 1947e: 4, 1955a: 212, 1976: 340; Pjatnitskii 1931: 171–173; Ratzburg 1837: 162, 1839: 197; Redtenbacher 1849a: 790, 1849b: 26; Reitter 1894a: 76, 1913a: 93, 1916: 298; Rhumbler 1922: 308, 1927: 322; Schedl 1934f: 1643, 1980a: 22; Schimitschek 1937c: 55, 1955c: 87; Seidlitz 1872: 394; Spessivtsev 1913a: 87–89, 1922: 221, 1923: 200–214, 1925: 50. (ms) Eggers 1910a, 1910b; Escherich 1932b.

macrographus Eichhoff 1880a: 200. Syntypes, sex?; Silesia; Hamburg Museum, lost. Synonymy: Balachowsky 1949a: 234.

References: (cn) Barbey 1925: 70; Judeich & Nitsche 1895: 525. (ce) Barbey 1927. (hb)

- Barbey 1901: 23, 78, 1925: 70; Bargmann 1906; Dombrowsky 1887, 1891; Eckstein 1897; Eichhoff 1881a: 147, 200; Henschel 1895a: 172; Judeich & Nitsche 1895: 525; Seitner 1911: 99. (**ds**) Gerhardt 1898b; Gronzelle 1905; Hagedorn 1910d: 72; Henschel 1895a: 172; Heyden, Reitter, & Weise 1883: 182, 1891: 672; Judeich & Nitsche 1895: 525; Kleine 1913b: 140; Pfeffer 1935: 159; Schilsky 1909: 188; Seidlitz 1891a: 564, 1891b: 610. (**tx**) Balachowsky 1949a: 234; Barbey 1901: 23, 78; Dombrowsky 1887, 1891; Eichhoff 1881a: 147, 200; Hagedorn 1910a: 101; Henschel 1895a: 172; Judeich & Nitsche 1895: 525; Schedl 1934f: 1643; Seidlitz 1891a: 564, 1891b: 610; Seitner 1887: 44; Spessivtsev 1913: 87–89. (**ms**) Eggers 1910a, 1910b.
- exsectus** Schedl 1972g: 67. Holotype ♀; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (**tx**) Schedl 1972g: 67, 1979c: 94.
- fallax** (Hagedorn) 1912c: 354 (*Trigonogenius*). Holotype, sex?, Amani, Deutsch-Ostafrika; Hamburg Museum, lost.
Distribution: Africa (Tanzania).
Hosts: *Khaya senegalensis*.
References: (**cn**) Hagedorn 1913a: 29. (**hb**) Hagedorn 1913a: 29; Schedl 1962j: 9. (**ds**) Hagedorn 1913a: 29; Kleine 1914b: 322, 1934a: 163; Schedl 1962j: 9. (**tx**) Eggers 1920: 34; Hagedorn 1912c: 354, 1913a: 29; Hopkins 1914: 131; Lucas 1920: 683; Schedl 1952d: 347, 1962j: 9; Wichmann 1915a: 107.
- festus** Wood 1967d: 39. Holotype ♂; 29 km W El Salto, Durango, Mexico; Wood Collection.
Figures: Bright 1981d: 359.
Distribution: North America (El Salvador/ Chiapas, Chihuahua, Durango, Jalisco, Mexico, Oaxaca, Puebla, Veracruz in Mexico/ Arizona in USA).
Hosts: *Pinus ayacahuite*, *P. durangensis*, *P. leiophylla*, *P. oocarpa*, *P. tenuifolia*, *P. sp.*
References: (**ds**) Atkinson & Equihua 1988: 96; Bright 1981d: 213; Rio Mora 1985: 14; Schedl 1977e: 42; Wood, S. L. 1982b: 1033. (**tx**) Bright 1981d: 213, 359; Rio Mora 1985: 14; de Ruette 1970: 109; Wood, S. L. 1967d: 39, 1982b: 1033.
- flavimaculatus** Murayama 1963: 392. Holotype, sex?, Tairei, North Manchuria; Murayama Collection in USNM, Washington.
Distribution: Asia (Manchuria).
Hosts: *Picea obovata*.
References: (**tx**) Murayama 1963: 392.
- formosus** Bright 1972d: 88. Holotype ♂; Barbecue Bottom, Trelawny Parish, Jamaica; CNCI, Ottawa.
Distribution: Antilles Islands (Jamaica).
References: (**ds**) Bright 1985c: 176. (**tx**) Bright 1972d: 88, 1985c: 176; McNamara 1985c: 176.
- franseriae** Wood 1972d: 75. Holotype ♀; 10 km W High Rolls, Lincoln National Forest, New Mexico [USA]; Wood Collection.
Distribution: North America (New Mexico in USA).
Hosts: *Franseria* sp.
References: (**tx**) Bright 1981d: 80; Wood, S. L. 1972d: 75, 1982b: 1124.
- frontalis** (Schedl) 1967d: 8 (*Breviophthorus*). Holotype ♂; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (**tx**) Schedl 1967d: 8, 1979c: 100.
- fulgens** Schedl 1965c: 68. Holotype ♀; Madagascar, Montagne d'Ambre; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
Hosts: Tree twigs.
References: (**tx**) Schedl 1965c: 68, 1977b: 111, 1979c: 101.
- furnissi** Bright 1976c: 433. Holotype ♀; Amecameca, Mexico; USNM, Washington.
Distribution: North America (Mexico in Mexico).
Hosts: *Pinus hartwegii*.
References: (**ds**) Atkinson & Equihua 1985a: 94; Bright 1981d: 144, 1986b: 683; Wood, S. L. 1982b: 1094. (**tx**) Bright 1976c: 433, 1981d: 144; McNamara 1984: 755; Wood, S. L. 1976c: 433, 1982b: 1094.
- fuscus** Blackman 1928b: 32. Holotype ♀; Glacier National Park, Montana [USA]; USNM, Washington.
Distribution: North America (British Columbia in Canada/ Montana in USA).
Hosts: *Pinus contorta*.
References: (**cn**) Lindgren 1980a: 68. (**hb**) Lindgren 1980a: 68. (**ds**) Bright 1976d: 182, 1981d: 211; Chamberlin 1939: 358; Gast et al. 1989: 385; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 425, 1982b: 1053. (**tx**) Blackman 1928b: 30–33; Bright 1981d: 211; Chamberlin 1939: 358; Keen 1929a: 33; Wood, S. L. 1972a: 425, 1977d: 515; 1982b: 1053.
- smithii** Schedl 1931b: 163. Holotype ♀; Copper Mountain, British Columbia [Canada]; USNM, Washington. Synonymy: Wood 1977d: 515.
References: (**cn**) Lindgren 1980a: 69. (**hb**) Chamberlin 1939: 369, 1958: 150; Lindgren 1980a: 69. (**ds**) Bright 1976d: 184; Chamberlin 1939: 369, 1958: 150; Kleine 1934a: 161; Leng & Mutchler 1933: 95. (**tx**) Chamberlin 1939: 369, 1958: 150; de Ruette 1970: 111; Schedl 1931b: 163, 1979c: 231; Wood, S. L. 1977d: 515.
- galeritus** Wood 1976a: 355. Holotype ♀; Rio Damitas, Dota Mts., San Jose, Costa Rica, 250 m; Wood Collection.
Distribution: North America (Costa Rica).

Hosts: *Rhicedia edulis*.

References: (**ds**) Bright 1981d: 75; Wood, S. L. 1982b: 1125. (**tx**) Bright 1981d: 75; Wood, S. L. 1976a: 355, 1982b: 1125.

germanus Bright 1976c: 434. Holotype ♂; 154 km N Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.

Distribution: North America (Oaxaca in Mexico).
References: (**ds**) Bright 1981d: 51; Wood, S. L. 1982b: 1127. (**tx**) Bright 1976c: 434, 1981d: 51; McNamara 1984: 755; Wood, S. L. 1982b: 1127.

ghanaensis Schedl 1972k: 292. Holotype, sex²: Ghana: Ashanti Region, Kwadaso, 320 m, N 6 42, W 1 39; NHMB, Budapest.

Distribution: Africa (Ghana).
References: (**tx**) Schedl 1972k: 292, 1979c: 104.

glabratus (Schedl) 1955g: 26 (*Ctenyophthorus*).
Lectotype ♀; Guatemala, Quezaltenango, 2350 m; Schedl Collection in NHMW, Wien, designated by Bright 1976b: 186.

Figures: Bright 1981d: 344.
Distribution: North America (Guatemala/ Chiapas, Oaxaca in Mexico).

Hosts: *Pinus ayacalhuite*, *P. ocheternae*, *P. pringlei*, *P. rudis*, *P. pseudostrobus*.

Notes: (1) Bright 1976b: 186 (to *Pityophthorus*). Schedl 1979c: 105 (citation of holotype invalid). (3) Schedl 1963c: 162 (cited in *Neodryocotes*).

References: (**hb**) Schwerdtfeger 1957c: 505. (**tx**) Bright 1976b: 186, 1977: 512, 1981d: 117, 359; Schedl 1955g: 18, 26, 1963c: 162, 1964m: 311, 1979c: 105; Wood, S. L. 1982b: 1056.

glabratus Eichhoff 1878b: 179. Holotype, sex²: Berlin, Corsica; Hamburg Museum, lost.

Figures: Grune 1979: 120.
Distribution: Africa (Egypt), Europe (Austria/ Bulgaria/ Corsica/ Czechoslovakia/ Denmark/ France/ Germany/ Greece/ Hungary/ Luxembourg/ Poland/ Spain/ S Sweden/ Switzerland/ W USSR).
Hosts: *Pinus mugo rotundata*, *P. nigra*, *P. sylvestris*, *P. uncinata*, rare in *Larix europaea*.

References: (**ay**) Escherich 1923b: 483, 555; Feytaud 1950a. (**bv**) Grune 1979: 121. (**cn**) Barbey 1924a, 1925: 281; Chorbadzhievo 1929; Eckstein 1926: 579, 1939a: 33–42, 81–92; Escherich 1923b: 483, 555; Feytaud 1950a; Grandi 1951; Joly 1976; Judeich & Nitsche 1895: 526; Maksimovic 1952: 20–40; Maksimovic & Milanovic 1964, 1966; Marcu 1926c: 65; Nusslin 1913: 205; Pierce, W. D. 1917: 75; Rhumbler 1922: 308, 1927: 322; Schimitschek 1937c: 55, 1955a: 29, 1955c: 87; Schwerdtfeger 1944a: 177, 1957a: 183; Wachtl 1883b: 319, 1901: 377, 381; Weber, H. 1926: 579; Wichmann 1927b: 361; Wilke 1931: 663. (**ec**) Barbey 1927; Berland 1934; Karpinski 1932a: 95; Kleine 1908c: 213; Masutti 1959: 268, 299; Michalski & Ratajczak 1989; Nusslin 1927: 322; Peyerimhoff 1928: 110–111; Pfeffer 1923a: 332, 1925b: 8, 1943b: 181, 1960: 346; Roubal 1934a: 86; Ruhm 1956b: 4; Schimit-

schek 1955a: 29; Schwerdtfeger 1944a: 177, 1957a: 183; Sedlaczek 1935a: 163; Seitner & Notzl 1925: 188; Thompson, W. R. 1943: 90; Wilke 1931: 663. (**hb**) Barbey 1901: 23, 76, 1913, 1924a, 1925: 281; Bargmann 1906; Chararas 1962c: 363; Charvat 1950; Chorbadzhievo 1929; Dombrowsky 1887, 1891; Eckstein 1926: 579, 1939a, 1939c; Eckstein, F. 1939; Eggers 1906; Eichhoff 1881a: 47, 196; Escherich 1923b: 483, 555; Feytaud 1950a; Grandi 1951; Györfi 1957; Hagedorn 1903a; Henschel 1895a: 172; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski 1948: 156; Karpinski & Strawinski 1948: 156; Kemner 1919: 175; Lengerken 1954: 86; Masutti 1959: 269, 299, 1964; Nunberg 1929c: 122; Nusslin 1913: 205, 1927: 322; Orest 1926b: 81; Pfeffer 1941b: 3; Postner 1974: 436; Rhumbler 1922: 308, 1927: 322; Schimitschek 1955a: 29; Schwerdtfeger 1944a: 177, 1957a: 183, 1981: 190; Sedlaczek 1935a: 163; Seitner & Notzl 1925: 188; Shatilov 1985; Spessivtsev 1913a: 88, 1921b: 220, 1923: 209; Stark 1926a: 336, 1952: 354; Tschorbadzhiev 1929: 170; Wachtl 1901: 377, 381; Weber, H. 1926: 579; Wichmann 1927b: 361. (**ds**) Ammann & Knabl 1913, 1923; Barthe 1896; Bejer-Petersen & Jorum 1977: 25; Borchert 1951; Brakman 1966b: 206; Buresh & Lazarov 1956; Champion 1894; Chararas 1962c: 363; Charvat 1950; Chorbadzhievo 1929; Eggers 1904, 1906; Endrodi 1958b; Escherich 1923b: 483, 555, 1932b; Everts 1922: 643, 1925; Fjellberg 1966: 154; Fuchs 1905a; Grune 1979: 121; Hagedorn 1903a, 1910d: 71; Hansen, V. 1939, 1956, 1964: 463; Hellen 1947; Henschel 1895a: 172; Heyden, Reitter, & Weise 1883: 182, 1891: 672, 1906: 71; Horion 1951; Jansson 1935: 77, 1936: 220; Jazentkovsky 1912: 288; Joly 1976; Judeich & Nitsche 1895: 526; Kangas 1968; Karpinski 1931: 30, 1932a: 95, 1948a: 173, 1948b: 231; Karpinski & Strawinski 1948: 156; Kemner 1919: 175; Kersten 1933: 76; Kiefer et al. 1942: 528; Klefbeck & Sjöberg 1960: 230; Kleine 1912a: 268, 1913a: 35, 1913b: 140, 1934a: 159; Kozikowsky 1921: 181; Kurir 1947c: 6; Lomnicki 1913b: 148; Lucht 1987: 278; Mahler 1987: 232; Maire et al. 1927: 1516; Marcu 1926c: 65, 1930: 327–336; Mequignon 1936: 54; Mirzorian 1950: 139; Morivee 1961: 141; Murayama 1940a: 232, 1942a: 55; Nunberg 1928b: 88, 112, 1938: 126–131, 1954: 71, 1960b: 159; Orest 1926c: 65, 81–87; Pfeffer 1924b: 471, 1928b: 8, 1931b: 74, 1947e: 12, 1950b: 76, 1960: 346, 1976: 340, 1984: 277, 1989a: 65; Pierce, W. D. 1917: 75; Pittioni 1943: 175; Postner 1974: 436; Prossen 1913: 83; Rapp 1934: 729; Reitter 1894a: 76; Roubal 1935b: 72, 1941: 269; Sainte-Claire 1914: 472; Sainte-Claire & Mequignon 1938: 447; Schanfuß 1915: 1243; Schedl 1971d: 427, 1980a: 22, 1981b: 78; Schilsky 1891: 157, 1909: 188; Schwerdtfeger 1981: 190; Seidlitz 1891a: 564, 1891b: 610; Stark 1926a: 336, 1926b: 105, 1926j:

- 127, 1952: 554; Stein & Weise 1877: 164; Tredl 1907: 14; Tschorbadjiev 1929: 170; Wichmann 1927a: 73; Wilke 1931: 663; Zinovjev 1955: 157. (**tx**) Balachowsky 1949a: 237; Barbey 1901: 23, 76; Charvat 1950; Dombrowsky 1887, 1891; Eggers 1915b: 13–14, 1920: 125–126; Eichhoff 1878b: 179, 1881a: 47, 196, 1883a: 110, 134; Endrodi 1957a: 307, 1957b; Escherich 1923b: 483, 555; Everts 1922: 643; Ferrant 1911; Fleischer 1927; Formanek 1907: 37; Grune 1979: 120–121; Hagedorn 1904e, 1910a: 101; Hansen, V. 1956, 1964: 463; Henschel 1895a: 172; Iablokoff-Khnzorian 1961: 105; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski & Strawinski 1948: 156; Koch 1913: 126, 1932: 121; Kuhn 1913: 1056; Lucht 1987: 278; Michalski 1969b: 569; Murayama 1940a: 232; Numberg 1929c: 122, 1935: 126; Peyerimhoff 1930: 258–259; Pfeffer 1927: 111–112, 1932b: 23, 1940b: 109, 112, 120, 1941b: 3, 1947e: 12, 1955a: 219, 1976: 340; Postner 1974: 436; Quaschik 1953: 35; Reitter 1894a: 76, 1913a: 94; Rhumbler 1922: 38, 1927: 322; Schedl 1934f: 1643, 1938a: 164, 1952f: 87, 1980a: 22, 1981b: 78; Schimitschek 1937c: 55, 1955c: 87; Seidlitz 1891a: 564, 1891b: 610; Seitner 1887: 45; Sokanovskii 1929: 670–672; Spessivtsev 1913a: 88, 1922a: 488, 490, 1922b: 219–221, 1923: 209, 1925a: 183, 1926: 50, 1929c: 298–303, 1931: 64; Stark 1952: 354. (**ms**) Escherich 1932b.
- glutae** Wood 1989: 183. Holotype ♀; Tamil Nadu: Evergreens, Tinnevely (Tirunelveli); FRI, Dehra Dun.
Distribution: Asia (Tamil Nadu in India).
Hosts: *Gluta travancorica*.
References: (**cn**) Mathur & Singh 1960b: 14. (**ds**) Mathur & Singh 1960b: 14. (**tx**) Wood, S. L. 1989: 183.
- grandis** Blackman 1928b: 119. Holotype, sex?; Arizona: Kaibab National Forest [USA]; USNM, Washington.
Figures: Bright 1981d: 370.
Distribution: North America (British Columbia in Canada/ Arizona, California, Colorado, Nebraska, New Mexico, South Dakota, W Texas, Utah, Wyoming in USA).
Hosts: *Pinus flexilis*, *P. leiophylla*, *P. monophylla*, *P. ponderosa*.
References: (**cn**) Blackman 1931c. (**ec**) Blackman 1931c. (**hb**) Blackman 1931c; Bright & Stark 1973: 115; Chamberlin 1939: 395, 1958: 157. (**ds**) Bright 1976d: 182, 1981d: 289; Bright & Stark 1973: 115; Chamberlin 1939: 395, 1958: 157; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 426, 1982b: 1106. (**tx**) Blackman 1928b: 114, 119–121; Bright 1981d: 289, 370; Chamberlin 1939: 395, 1958: 157; Keen 1929a: 33; de Ruette 1970: 109; Wood, S. L. 1972a: 426, 1982b: 1106.
- granulipennis** Schedl 1965c: 69. Holotype ♀; Madagascar, Mt. d'Ambre; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
Hosts: Tree twigs.
References: (**tx**) Schedl 1965c: 69, 1977b: 112, 1979c: 112.
- granulosus** (Schedl) 1972h: 60 (*Breviophthorus*). Holotype, sex?; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (**tx**) Schedl 1972h: 60, 1979c: 112.
- guatemalensis** Blandford 1904: 239. Lectotype ♀; Guatemala City, Guatemala; BMNH, London, designated by Bright 1976b: 184.
Figures: Bright 1981d: 342.
Distribution: North America (Guatemala).
Hosts: Probably *Quercus* sp.
Notes: (3) Bright 1981d: 106 (treated *quercinus* as a synonym; it is at least a subspecies, perhaps a distinct species).
References: (**hb**) Bright 1981d: 106. (**ds**) Blackwelder 1947: 782; Bright 1981d: 106; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 363; Wood, S. L. 1982b: 1084. (**tx**) Blandford 1904: 239; Bright 1976b: 184, 1977: 515, 1981d: 106, 342; Hagedorn 1910a: 101; Schedl 1938a: 183, 1955g: 18; Wood, S. L. 1982b: 1084.
- gunneri** Schedl 1970f: 583. Holotype ♀; Guyane française, Camopi-Oyapock dans nid d'Atta; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
Notes: (1) Below the type label on the holotype is a subsequently placed white label with red margin that says "holotype, *Pityophthorus quadrispinatus* Schedl." Whether this was an error in labeling or a suggestion of synonymy by Schedl was not determined.
References: (**tx**) Schedl 1970f: 583–584, 1979c: 113.
- henscheli** Seitner 1887: 44. Syntypes, sex?; Tyrol, Austria; not located, 1 syntype in NHMW, Wien.
Figures: Grune 1979: 118.
Distribution: Europe (Austria/ Czechoslovakia/ Germany/ Greece/ Hungary/ Italy/ Poland/ Romania/ W USSR/ Yugoslavia).
Hosts: *Pinus cembrae montana*, *P. heldreichii*, *P. mugo*.
Notes: (1) Schedl 1979c: 116 (citation of holotype invalid).
References: (**bx**) Grune 1979: 119; Hellrigl 1985. (**cn**) Schimitschek 1938b: 114, 1955a: 42; Schwerdtfeger 1944a: 177, 1957a: 183; Wachtl 1901: 381; Wichmann 1927b: 378. (**ec**) Hellrigl 1985; Heqvist 1955a: 96; Jahn & Simreich 1960b; Kleine 1908c: 213; Pfeffer 1960: 346; Ruschka 1925: 200; Schimitschek 1955a: 42; Seitner 1913a: 27; Seitner & Notzl 1925: 187; Schwerdtfeger 1944a: 177, 1957a: 183; Thompson, W. R. 1943: 90. (**hb**) Bargmann 1906; Dombrowsky 1891; Henschel

- 1895a: Karpinski & Strawinski 1948: 156; Knotek 1899b: 1, 1901: 565; Lengerken 1939: 65; Numberg 1947c: 106; Schimitschek 1955a: 42; Seitner & Notz 1925: 187; Schwerdtfeger 1944a: 177, 1957a: 183, 1981: 190; Spessivtsev 1921b: 221; Wachtl 1901: 381; Wichmann 1927b: 378. **(ds)** Annamm & Knabl 1923: Bilek 1944: Endrodi 1957a: 307, 1958b; Favre 1890: Fuchs 1905a; Grune 1979: 119; Hagedorn 1910d: 71; Henschel 1895a: 172; Heyden, Reitter, & Weise 1891: 672, 1906: 711; Horion 1951, 1954b: 21; Karpinski 1931: 30; Karpinski & Strawinski 1948: 156; Kleine 1913a: 35, 1913b: 140, 1934a: 160; Knotek 1899b: 1, 1901: 565; Knir 1947c: 9; Lucht 1987: 278; Numberg 1938: 126, 1954: 71; Pfeffer 1924a: 96, 1931b: 76, 1960: 346, 1989a: 66; Pittioni 1943: 176; Postner 1974: 436; Prossen 1913: 83; Reitter 1894a: 76, 1916: 299; Schedl 1967c: 71, 1980a: 22, 1981b: 78; Schilsky 1909: 188; Schimitschek 1938b: 114; Schwerdtfeger 1951: 190; Stierlin 1898: 441; Tredl 1907: 14; Wichmann 1927a: 73. **(tx)** Dombrowsky 1891: Eggers 1911a: 122, 1914: 189; Endrodi 1957a, 1957b; Fanvel 1887, 1889; Fleischer 1927; Grune 1979: 118–119; Hagedorn 1910a: 101; Hellbrigl 1985; Henschel 1895a: 172; Karpinski & Strawinski 1948: 156; Lucht 1987: 278; Numberg 1938: 126–131, 1954: 71; Pfeffer 1932b: 4, 1940b: 109, 113, 123, 1955a: 211; Postner 1974: 436; Reitter 1894a: 76, 1913a: 94, 1916: 299; Schedl 1934f: 1643, 1964m: 312, 1981b: 78; Seitner 1887: 44–45; Spessivtsev 1922: 221, 1926: 49; Stierlin 1898: 441, 1925: 187–196; Wichmann 1910a: 146. **(ms)** Eggers 1911b; Seitner 1913a: 27.
- hermosus Wood** 1976a: 356. Holotype ♀; Yuscaram, Paraiso, Honduras; Wood Collection.
Distribution: North America (Honduras).
Hosts: *Eupatorium dolioides*, *Perynanium grande*.
References: **(ds)** Bright 1981d: 40; Wood, S. L. 1982b: 1139. **(tx)** Bright 1981d: 40; McNamara 1984: 755; Wood, S. L. 1976a: 356, 1982b: 1139.
- hintzi Schedl** 1938h: 459. Syntypes ♂; Portugiesisch Guinea: MNB, Berlin and Schedl Collection in NHMW, Wien.
Distribution: Africa (Ghana/ Guinea-Bissau/ Ivory Coast/ Tanzania/ Zaire).
Hosts: *Khaya* sp., *Mitragyne* sp.
Notes: (1) Schedl 1979c: 117 (citation of holotype invalid).
References: **(ds)** Browne 1975a: 758, 1981a: 126; Mayne & Donis 1962: 309; Schedl 1962j: 10, 1972e: 282, 1977d: 278, 1979b: 415. **(tx)** Schedl 1938h: 459, 1962j: 10, 1970i: 223, 1979c: 117.
- acuminatus Brownie** 1965a: 192. Holotype ♀; Ivory Coast: Adiopodoume; RNIH, Leiden, preoccupied by Schedl 1940a: 346. Synonymy: Schedl 1970i: 223 (possible synonymy).
References: **(tx)** Browne 1965a: 192; Schedl 1970i: 223.
- hispaniolus Bright** 1985c: 181. Holotype ♀; Colonia, Dominican Republic, 1000 m; NHMBS Basel, Switzerland.
Distribution: Antilles Islands (Dominican Republic).
References: **(ds)** Bright 1985c: 176. **(tx)** Bright 1985c: 176, 181.
- hylocuroides Wood** 1964: 69. Holotype ♀; 18 km NE Jacala, Hidalgo, Mexico; Wood Collection.
Distribution: North America (Hidalgo in Mexico).
Hosts: *Rhus* cf. *aromatica*.
References: **(ds)** Bright 1981d: 32, 1986b: 683; Wood, S. L. 1982b: 1120. **(tx)** Bright 1981d: 32; McNamara 1984: 756; Wood, S. L. 1964: 69, 1982b: 1120.
- ignotus Schedl** 1961e: 143. Holotype ♀; Madagascar: Mt. d'Ambre; Schedl Collection in NHMW, Wien.
Distribution: Madagascar.
References: **(tx)** Schedl 1953d: 69, 1961e: 143, 1977b: 112, 1979c: 120.
- ikelaensis Numberg** 1967b: 322. Holotype ♀; Tshunapa, Ikela [Congo]; MRCB, Tervuren.
Figures: Numberg 1967a: 337.
Distribution: Africa (Zaire).
References: **(tx)** Numberg 1967a: 337, 1967b: 322.
- imitans (Eggers)** 1920: 34 (*Trigonogenius*). Lectotype, sex?, Amani (Deutsch-Ostafrika); USNM, Washington, designated by Anderson & Anderson 1971: 15.
Distribution: Africa (Tanzania).
References: **(ds)** Schedl 1962j: 10. **(tx)** Anderson, W. H. & Anderson 1971: 15; Eggers 1920: 34; Lucas 1920: 683; Numberg 1969a: 396; Schedl 1952d: 347, 1979c: 121.
- immanis Blackman** 1928b: 98. Holotype ♀; Chiricahua Mts., Arizona [USA]; USNM, Washington.
Distribution: North America (Chiricahua in Mexico/ Arizona, New Mexico in USA).
Hosts: *Pinus ponderosa*, *P. strobiformis*.
References: **(hb)** Chamberlin 1939: 385. **(ds)** Bright 1981d: 15; Chamberlin 1939: 385; Keen 1929a: 33; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 89, 98–99; Bright 1981d: 159; Chamberlin 1939: 385; Keen 1929a: 33; McNamara 1984: 756; Wood, S. L. 1964: 67, 1977d: 515, 1982b: 1097.
- sulcatus Bright** 1977: 528. Holotype ♀; Mt. Lemon, Pima Co., Arizona [USA]; CNCI, Ottawa. Synonymy: Wood 1977d: 515, 1982b: 1096–1097.
Notes: Bright 1981d: 166 (treated as a distinct species).
References: **(ds)** Bright 1981d: 166. **(tx)** Bright 1977: 528, 1981d: 166, 352; McNamara 1984: 758; Schedl 1979c: 245; Wood, S. L. 1977d: 515, 1982b: 1096.
- impexus Bright** 1978: 76. Holotype ♀; 10 km S Carapan, Michoacan, Mexico; Wood Collection.

Distribution: North America (Durango, Mexico, Michoacan in Mexico).

Hosts: *Pinus ayacaluute*, *P. sp.*

References: (**hb**) Burgos & Saucedo 1983: 120. (**ds**) Atkinson & Equihua 1985a: 95; Bright 1981d: 197; Burgos & Saucedo 1983: 120; Wood, S. L. 1982b: 1086. (**tx**) Bright 1978: 76, 1981d: 197; McNamara 1984: 756; Wood, S. L. 1982b: 1086.

inaequidens Schedl 1976a: 68. Holotype, sex?; Jacareacanga, Para, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (**tx**) Schedl 1976a: 68.

indefessus Bright 1986a: 641. Holotype, sex?; Mexico: Estacion de Biologia, Chamela, Edo. Jalisco; CNCI, Ottawa.

Distribution: North America (Jalisco in Mexico).

References: (**tx**) Bright 1986a: 641.

indigen Wood 1976a: 361. Holotype ♀; 3 km W Armeria, Colima, Mexico, 70 m; Wood Collection. Distribution: North America (Colima, Jalisco, Oaxaca in Mexico).

Hosts: *Bursera* sp.

References: (**cc**) Equihua & Atkinson 1986: 634. (**hb**) Equihua & Atkinson 1986: 634. (**ds**) Bright 1981d: 79; Equihua & Atkinson 1986: 634; Wood, S. L. 1982b: 1126. (**tx**) Bright 1981d: 79; Wood, S. L. 1976a: 361, 1982b: 1126.

indigus Wood 1978b: 398. Holotype ♀; 3 km E Gould, Jackson Co., Colorado [USA]; Wood Collection, automatic.

Distribution: North America (Colorado in USA).

Hosts: *Pinus contorta*

References: (**ds**) Bright 1981d: 128; Wood, S. L. 1982b: 1053. (**tx**) Bright 1981d: 128; Wood, S. L. 1978b: 398, 1982b: 1053.

indigen Wood 1977b: 214. Holotype ♀; 3 km E Gould, Jackson Co., Colorado [USA]; Wood Collection, preoccupied by Wood 1976a: 361.

References: (**tx**) Wood, S. L. 1977b: 214, 1978b: 398.

irritans Schedl 1979j: 127. Holotype ♀; 3 km E Gould, Jackson Co., Colorado [USA]; Wood Collection, automatic.

Notes: (1) This is an unneeded replacement name for *indigen* Wood 1977: 214.

References: (**tx**) Bright 1981d: 128; Schedl 1979j: 127.

ineditus Bright 1976c: 434. Holotype ♀; 53 km S Valle Nacional, Oaxaca, Mexico; CNCI, Ottawa. Distribution: North America (Oaxaca, Veracruz in Mexico).

Hosts: *Pinus* sp.

References: (**ds**) Bright 1981d: 118; Wood, S. L. 1982b: 1057. (**tx**) Bright 1976c: 434, 1981d: 118; McNamara 1984: 756; Wood, S. L. 1982b: 1057.

infirmus Schedl 1972g: 68. Holotype, sex?; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (**tx**) Schedl 1972g: 68, 1979c: 124.

infulatus Blackman 1928b: 103. Holotype ♀; Sacramento Mts., New Mexico [USA]; USNM, Washington.

Distribution: North America (Arizona, New Mexico in USA).

Hosts: *Pinus ponderosa*, *P. strobiformis*.

References: (**hb**) Chamberlin 1939: 357. (**ds**) Atkinson & Equihua 1985c: 360, 1988: 98; Bright 1981d: 160; Chamberlin 1939: 357; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1090. (**tx**) Blackman 1928b: 103; Bright 1977: 515, 1981d: 160; Chamberlin 1939: 357; Keen 1929a: 32; Wood, S. L. 1982b: 1090.

mollis Blackman 1928b: 104. Holotype ♀; Tortillo Canyon, New Mexico [USA]; USNM, Washington. Synonymy: Bright 1977: 514.

References: (**hb**) Chamberlin 1939: 357. (**ds**) Chamberlin 1939: 357; Keen 1929a: 33; Kleine 1934a: 160; Leng & Mutchler 1933: 52. (**tx**) Blackman 1928b: 104; Bright 1977: 514; Chamberlin 1939: 357; Keen 1929a: 33; McNamara 1984: 757.

hubbardi Blackman 1928b: 105. Holotype ♀; Chiricahua Mts., Arizona [USA]; USNM, Washington. Synonymy: Bright 1977: 514.

References: (**hb**) Chamberlin 1939: 357. (**ds**) Chamberlin 1939: 357; Leng & Mutchler 1933: 52. (**tx**) Blackman 1928b: 105, 1942a: 215–216; Bright 1977: 515; Chamberlin 1939: 357; McNamara 1984: 755.

ingens Blackman 1928b: 124. Holotype ♀; Santa Catalina Mts., Arizona [USA]; USNM, Washington. Distribution: North America (Chihuahua, Durango in Mexico/ Arizona in USA).

Hosts: *Pinus ponderosa*, *P. sp.*

References: (**hb**) Chamberlin 1939: 395. (**ds**) Bright 1981d: 281; Chamberlin 1939: 395; Furniss, M. M. 1978; Keen 1929a: 33; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 110. (**tx**) Blackman 1928b: 115, 124–125; Bright 1981d: 281; Chamberlin 1939: 395; Keen 1929a: 33; McNamara 1984: 756; Wood, S. L. 1982b: 1100.

inhabilis Bright 1986a: 642. Holotype ♀; Mexico: Chilapa, Guerrero; CNCI, Ottawa.

Distribution: North America (Guerrero in Mexico).

References: (**tx**) Bright 1986a: 642.

inops Wood 1976c: 353. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Tree limb.

References: (**ds**) Bright 1981d: 15; Wood, S. L. 1982b: 1122. (**tx**) Bright 1981d: 15; Wood, S. L. 1976c: 353, 1982b: 1122, 1983a: 648.

insuetus Bright 1955b: 479. Holotype ♀; Mexico: Huitzilac, Morelos: CNCI, Ottawa.

Distribution: North America (Morelos in Mexico).

Hosts: Compositae sp.

References: (tx) Bright 1955b: 479.

intentus Bright 1975: 77. Holotype ♀; Bear Canyon, Santa Catalina Mts., Arizona [USA]; CNCI, Ottawa.

Figures: Bright 1981d: 369.

Distribution: North America (Durango in Mexico/ Arizona in USA).

Hosts: *Pinus engelmannii*, *P. leiophylla*, *P. ponderosa*, *P. strobiformis*, *Pseudotsuga menziesii*.

References: (ds) Bright 1981d: 286; Wood, S. L. 1982b: 1105. (tx) Bright 1975: 77, 1981d: 286, 369; McNamara 1984: 756; Wood, S. L. 1982b: 1105.

intextus Swaine 1917: 29. Holotype ♂; Athabaska Landing, Alberta [Canada]; CNCI, Ottawa.

Distribution: North America (Alberta, British Columbia, Manitoba, New Brunswick, Newfoundland, Nova Scotia, Ontario, Saskatchewan in Canada/ Maine, Michigan, New Hampshire, New York, West Virginia, Wyoming in USA).

Hosts: *Picea canadensis*, *Pinus resinosa*.

Notes: (3) Blackman 1928b: 101 (redescribed). Bright 1981d: 147 (treated *ornatus* Blackman as a synonym).

References: (cn) Doane et al. 1936; Swaine 1918a: 97, 102. (hb) Chamberlin 1939: 386, 1958: 153–154; Doane et al. 1936; Swaine 1918a: 97, 102. (ds) Bright 1976d: 182, 1981d: 147; Chamberlin 1925, 1939: 386, 1958: 153–154; Drooz 1985: 364; Hopping 1922; Kleine 1934a: 160; Leng 1920: 341; Wood, S. L. 1972a: 426, 1982b: 1089. (tx) Blackman 1928b: 90, 101; Bright 1967b: 678, 1977: 515, 1981d: 147, 349; Chamberlin 1939: 386, 1958: 153–154; Hoebeke 1978; Pardy 1983; de Ruelle 1970: 109; Swaine 1917: 29–30, 1918a: 97, 102; Wood, S. L. 1972a: 426, 1977b: 209, 1982b: 1089.

shepardi Blackman 1922c: 124. Holotype ♀; Chamberlin Lake region, Township 7, R12, Piscataquis Co., Maine [USA]; USNM, Washington. Synonymy: Bright 1977: 515.

References: (cn) Blackman 1950. (hb) Baker, W. L. 1972: 255; Blackman 1950; Chamberlin 1939: 386. (ds) Anonymous 1926c: 519; Baker, W. L. 1972: 255; Blackman 1950; Bright 1971a: 127, 1976d: 188; Chamberlin 1925, 1939: 386; Kleine 1934a: 161; Leng & Mutchler 1927: 52; Leonard 1928: 519. (tx) Blackman 1922c: 124–125, 1928b: 90, 100–101; Bright 1977: 515; Chamberlin 1939: 386.

tonsus Blackman 1928b: 101. Holotype ♀; Grand Island, Michigan [USA]; USNM, Washington. Synonymy: Bright 1977: 515.

References: (cn) Blackman 1950. (hb) Blackman 1950; Chamberlin 1939: 386. (ds) Blackman 1950; Bright 1976d: 188; Chamberlin

1939: 386; Kleine 1934a: 161; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 101; Bright 1977: 515; Chamberlin 1939: 386; de Ruelle 1970: 111.

irregularis Eggers 1931b: 31. Holotype ♀; Brasilien (Sao Paulo); Eggers Collection, in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 782; Eggers 1931b: 31; Schedl 1972g: 44, 1979c: 128.

jeffreyi Blackman 1928b: 113. Holotype ♀; Inyo N.F., Bishop, California [USA]; USNM, Washington. Distribution: North America (California, Oregon in USA).

Hosts: *Pinus aristata*, *P. jeffreyi*, *P. ponderosa*, rare or accidental in other *Pinus* spp.

References: (hb) Bright & Stark 1973: 114; Chamberlin 1939: 389. (ds) Bright 1981d: 71; Bright & Stark 1973: 114; Chamberlin 1939: 389; Keen 1929a: 33; Kleine 1934a: 160; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 91, 113–114; Bright 1981d: 71; Chamberlin 1939: 389; Keen 1929a: 33; de Ruelle 1970: 110; Schedl 1939h: 167; Wood, S. L. 1982b: 1116.

joveri Schedl 1954d: 881. Lectotype ♀; Cote d'Ivoire, Adiopodoume; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 130. Figures: Schedl 1962j: 11.

Distribution: Africa (Equatorial Guinea/ Gabon/ Ghana/ Ivory Coast/ Zaïre).

Hosts: *Fagara macrophylla*, *Phialodiscus bancoensis*.

References: (cc) Schedl 1958d: 190, 1962k: 10. (hb) Schedl 1958d: 191, 1962j: 10. (ds) Schedl 1962h: 59, 1962j: 10, 1966c: 225, 1971g: 191, 1972e: 282. (tx) Schedl 1954d: 881–882, 1962j: 10–11, 1962k: 1080, 1964j: 46, 1979c: 130.

jucundus Blandford 1894d: 87. Syntypes 4, sex?: near Nagasaki, Japan; BMNH, London.

Figures: Nobuchi 1966d: pl. 5.

Distribution: Asia (NE China/ Japan/ Korea/ Sakhalin Island, Siberia in E USSR).

Hosts: *Pinus densiflora*, *P. thunbergii*, *Picea ajanensis*, *Abies* sp.

References: (bv) Choo, Woo, & Park 1988. (cn) Inouye 1949b; Murayama 1957a: 7; Shiraki 1952; Yasunga 1962: 197–200. (cc) Inouye et al. 1955: 80. (hb) Inouye et al. 1955: 80; Stark 1952: 351. (ds) Blandford 1894c; Cho 1957; Choo 1983: 76; Choo & Woo 1985: 165; Choo, Woo, & Park 1988; Hagedorn 1910d: 71; Inouye 1949b; Kleine 1913b: 140, 1914b: 258, 1934a: 160; Ko 1969: 281; Kurenzov 1965; Murayama 1929b: 2, 1930a: 9, 1930b: 14, 1936b: 116, 1937b: 375, 1948: 2, 1949a: 12, 1949c: 101, 1950b: 1293, 1951c: 4, 1954a: 7, 1954b: 169, 1955: 99; Nobuchi 1966d: 33; Pfeffer 1976: 340; Shiraki 1952; Stark 1936c: 142, 1952: 351. (tx) Blandford 1894d: 87; Choo 1983: 76; Hagedorn 1910a: 101; Murayama

1930b: 14, 1934c: 298, 1937b: 375, 1950b: 1293, 1954b: 169, 1955: 99; Nobuchi 1966d: 33, pl. 5; Pfeffer 1940b: 108, 1976: 340; Pjatkitskii 1931: 169; Schedl 1934f: 1643, 1938a: 183; Stark 1952: 351.

juglandis Blackman 1928b: 42. Holotype ♀; Lone Mountain, New Mexico [USA]; USNM, Washington. Distribution: North America (Chihuahua in Mexico/ Arizona, California, New Mexico in USA). Hosts: *Juglans californica*, *J. major*, *J. nigra*. References: (bv) Roling & Kearby 1975b. (cn) Anonymous 1960c: 28, 1961t, 1964h. (hb) Bright & Stark 1973: 106; Chamberlin 1939: 362. (ds) Anonymous 1961t, 1964h; Bright 1981c: 81; Bright & Stark 1973: 106; Chamberlin 1939: 362; Furniss, R. L. & Carolin 1977: 402; Harper 1965: 82, 1969: 93; Keen 1929a: 32; Kleine 1934a: 160; Leng & Mutchler 1933: 52; Warner 1960: 1; Wood, S. L. 1960b: 69, 1977a: 72, 1982b: 1123. (tx) Blackman 1928b: 42; Bright 1981d: 81; Chamberlin 1939: 362; Keen 1929a: 32; McNamara 1984: 756; Wood, S. L. 1960b: 69, 1982b: 1123.

keeni (Blackman) 1928b: 19 (*Myeloborus*). Holotype, sex?; Jacumba, California [USA]; USNM, Washington.

Distribution: North America (Arizona, California, Colorado, New Mexico in USA).

Hosts: *Pinus cembroides*, *P. edulis*, *P. monophylla*.

Notes: (1) Bright 1976c: 427 (to *Pityophthorus*). References: (hb) Bright & Stark 1973: 96; Chamberlin 1939: 340. (ds) Bright 1981d: 209; Bright & Stark 1973: 96; Chamberlin 1939: 340; Keen 1929a: 31; Kleine 1934a: 163; Leng & Mutchler 1933: 53; Wood, S. L. 1982b: 1042. (tx) Blackman 1928b: 19; Bright 1976c: 427, 1981d: 209; Chamberlin 1939: 340; Keen 1929a: 31; McNamara 1984: 753; Wood, S. L. 1982b: 1042.

kenyae Schedl 1955i: 217. Syntypes, sex?; Kenya: Londiani; BMNH, London, and Schedl Collection in NHMW, Wien.

Distribution: Africa (Kenya).

Hosts: *Calodendrum capensis*.

References: (ds) Gardner 1957a: 31; Schedl 1962j: 12. (tx) Schedl 1955i: 217, 1962j: 12, 1979c: 132.

kirgisicus Pjatkitskii 1931: 167. Lectotype, sex?; Kirgisien, USSR; Pjatkitskii Collection, not located, designated by Michalski 1969a: 895.

Figures: Tsai & Li 1959: 94.

Distribution: Europe (W USSR).

Hosts: *Picea schrenkiana*.

References: (cc) Kostin 1964: 111. (hb) Kostin 1960: 133; Stark 1952: 348. (ds) Kadyrov 1989; Kleine 1934a: 160; Marikovskii 1956: 73; Parnfentev 1951: 429; Pfeffer 1976: 340; Stark 1952: 348. (tx) Michalski 1969a: 895, 1969b: 569; Pfeffer 1940b: 108, 111, 117, 1976: 340; Pjatkitskii 1931: 167–169; Schedl 1934f: 1643, 1979c: 132; Stark 1952: 348; Tsai & Li 1959: 94.

kiruensis Schedl 1957d: 61. Holotype, sex?; Congo Belge; Kivu, Mt. Kabuzi versant Ouest, 2310 m; MRCB, Tervuren.

Figures: Schedl 1962j: 14.

Distribution: Africa (Zaire).

Hosts: *Albizia gummifera*, *Persca gratissima*, *Ricinus communis*, *Sapium manneanum*, *Trema orientale*.

References: (cc) Schedl 1962j: 14. (hb) Schedl 1962j: 14. (ds) Mayne & Donis 1962: 309; Schedl 1962j: 14. (tx) Schedl 1955i: 217, 1957d: 61, 1962j: 14, 1979c: 133.

knoteki Reitter 1898b: 356. Syntypes, sex?; Tyrol, Austria; NHMB, Budapest.

Figures: Grune 1979: 116, Pfeffer 1989a: pl. 9.

Distribution: Europe (Austria/ Bulgaria/ Poland/ W USSR).

Hosts: *Pinus cembrae montana*, *P. mugo*, *P. peuce*.

References: (bv) Grune 1979: 117; Hellrigl 1985. (cn) Schimitschek 1938b: 115, 120, 1955a: 42; Wachtl 1901: 351. (cc) Hellrigl 1985; Jahn & Sinreich 1960; Kleine 1908c: 313; Schimitschek 1955a: 42; Seitner 1913a: 27; Thompson, W. R. 1943: 91; Wichmann 1959: 413. (hb) Karpinski & Strawinski 1948: 156; Pfeffer 1941b: 19; Postner 1974: 436; Schimitschek 1955a: 42; Spessivtsev 1921b: 221; Wachtl 1901: 351. (ds) Buresh & Lazarov 1956; Grune 1979: 117; Hagedorn 1910d: 71; Heyden, Reitter, & Weise 1906: 711; Horion 1961; Karpinski 1931: 30; Karpinski & Strawinski 1948: 156; Kleine 1913a: 35, 1913b: 140, 1934a: 160; Marcu 1957b: 213; Numberg 1938: 126; Pfeffer 1936: 90, 1976: 340, 1989a: 64; Postner 1974: 436; Reitter 1916: 248; Schilsky 1909: 188; Schimitschek 1938b: 115, 120; Tredl 1907: 14. (tx) Balachowsky 1949a: 236; Cole & Freude 1972; Eggers 1915b: 13–14; Grune 1979: 116–117; Hagedorn 1910a: 101; Hellrigl 1985; Jansson 1936: 220; Karpinski & Strawinski 1948: 156; Kolnbajiv 1934: 65; Numberg 1938: 126; Pfeffer 1940b: 108, 112, 118, 1941b: 19, 1955a: 210, 1965: 63, 1976: 340, 1989a: pl. 9; Pjatkitskii 1931: 169; Postner 1974: 436; Reitter 1898b: 356, 1913a: 93, 1916: 298; Schedl 1934f: 1643, 1979c: 134; Spessivtsev 1922: 221, 1926: 50. (ms) Eggers 1911b; Seitner 1913a: 27; Wichmann 1959: 413.

kuscheli Schedl 1951d: 19. Syntypes, sex?; Chile; NMHN, Santiago, and 4 in Schedl Collection in NHMW, Wien.

Distribution: South America (Chile).

References: (ds) Schedl 1972d: 145. (tx) Schedl 1951d: 19, 1972d: 145, 1979c: 135.

laetus Wood 1976a: 355. Holotype ♀; Volcan Poas, Heredia, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Tree branch.

References: (ds) Bright 1981d: 103; Wood, S. L. 1982b: 1081. (tx) Bright 1981d: 103; Wood, S. L. 1976a: 355, 1982b: 1081.

languidus Eichhoff 1878b: 186. Holotype, sex?; America meridionalis (Venezuela); Hamburg Museum, lost.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 782; Bruch 1914a; Hagedorn 1910d: 71; Kleine 1913b: 140, 1914b: 339. (tx) Eichhoff 1878a: 389, 1878b: 186; Hagedorn 1910a: 101.

lupponicus Stark 1952: 356. Syntypes, sex?; Kola peninsula, Maritime Prov.; IZL, Leningrad.

Distribution: Asia (Finland/ Ussuri in E USSR).

Hosts: *Picea obovata*, *Pinus cembra siberica*, *P. koraiensis*.

References: (cc) Pfeffer 1960: 345. (hb) Kurenzov 1948b: 108, 1950d: 148; Stark 1952: 356. (ds) Nuorteva 1971: 71; Pfeffer 1960: 345, 1976: 340; Postner 1974: 437; Stark 1952: 356. (tx) Kurenzov 1941a: 177, 1948b: 108; Pfeffer 1976: 340; Postner 1974: 437; Stark 1952: 356.

laticeps Bright 1978: 78. Holotype ♂; 20.5 km N Oaxaca, Oaxaca, Mexico, 300 m; CNCI, Ottawa.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Pinus* sp.

References: (ds) Bright 1981d: 237; Wood, S. L. 1982b: 1060. (tx) Bright 1978: 78, 1981d: 237; McNamara 1984: 756; Wood, S. L. 1982b: 1060.

lautus Eichhoff 1872a: 135. Holotype, sex?; Amer. bor.; Hamburg Museum, lost; neotype ♀; Morgantown, West Virginia [USA], designated by Bright 1976c: 427.

Figures: Blackman 1922b:pl. 1, fig. 5, Bright 1981d: 337.

Distribution: North America (Ontario, Quebec in Canada/ Connecticut, District of Columbia, Florida, Illinois, Indiana, Kansas, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Acer saccharum*, *Cercis canadensis*, *Hammamelis* sp., *Juglans nigra*, *Quercus* sp., *Picea* sp., *Pinus strobus*, *Rhus typhina*, *R. radicans*.

Notes: (3) Blackman 1928b: 37 (redescribed), 1928b: 39 (*rhois acerini*, *rhois hammamelidis*, *rhois swainnei*, *piccae*, nomen nudum, no status).

References: (bv) Kirkendall 1951b. (cn) Felt 1906: 751; Hopkins 1899c: 443; Lindquist, O. H. & Syme 1981: 88; Swaine 1918a: 95; Syme & Nystrom 1988: 84. (cc) Bushing 1965: 465; Felt 1906: 751; Kirkendall 1951b. (hb) Blackman 1922b: 101, 103; Bright 1981d: 58; Chamberlin 1939: 360; Felt 1906: 751; Hopkins 1899c: 443; Kirkendall 1951b; Swaine 1918a: 95. (ds) Atkinson & Equihua 1985a: 92, 1988: 98; Atkinson et al. 1991: 163; Blackman 1922b: 101, 103; Blatchley & Leng 1916: 633; Bright 1981d: 58; Chamberlin 1925, 1939: 360; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Drooz 1985: 363; Gemminger & Harold 1872: 2688; Hagedorn

1910d: 71; Henshaw 1885: 147; Hopkins 1893a: 131, 1893b: 209; Kleine 1913b: 140, 1914b: 386, 1934a: 160; Leng 1920: 341; Lindquist, O. H. & Syme 1981: 88; Swaine 1909: 135; Syme & Nystrom 1988: 84; Weber, B. C., & McPherson; Wood, S. L. 1982b: 1131. (tx) Blackman 1922b: 101, 103, 1928b: 36–38; Blandford 1898c: 6; Blatchley & Leng 1916: 633; Bright 1976c: 427, 1981d: 58, 337; Chamberlin 1939: 360; Eichhoff 1872a: 135, 1878b: 190; Hagedorn 1910a: 101; LeConte 1876: 354; Lindquist, O. H. & Syme 1988: 84; Swaine 1909: 135, 1918a: 95, 104; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1982b: 1131. (ms) Eckstein 1900c.

rhois Swaine 1917: 26. Holotype, sex?; Ithaca, New York [USA]; CNCI, Ottawa. Synonymy: Bright 1976c: 427.

References: (cn) Blackman 1950; Lindquist, O. H. & Syme 1988: 84; Swaine 1918a: 95, 99; Syme & Nystrom 1988: 84. (cc) Loan & Matthews 1973; Marsh 1979: 295. (hb) Baker, W. L. 1972: 255; Beal & Massey 1945: 126–129; Blackman 1922b, 1950; Chamberlin 1939: 360; Swaine 1918a: 95, 99. (ds) Anonymous 1926c: 518; Beal & Massey 1945: 126–129; Beaulne 1956; Blackman 1922b: 103–104, 1950; Bright 1976d: 188; Chamberlin 1939: 360; Dodge 1938; Glick 1939: 37, 71; Hoffmann 1940: 60–61, 1942: 13; Kirk 1969; Kleine 1934a: 161; Leng 1920: 341; Leonard 1928: 518; Lindquist, O. H. & Syme 1981: 88; Syme & Nystrom 1988: 84. (tx) Beal & Massey 1945: 126–129; Blackman 1921: 7–8, 1922b: 103–104, 1928b: 36–39; Bright 1967b: 678, 1976c: 427, 1977: 517; Chamberlin 1939: 360; Dodge 1938: 43; Hoebeke 1978; Lindquist, O. H. & Syme 1981: 88; de Ruette 1970: 111; Swaine 1917: 26, 1918a: 95, 99, 104; Syme & Nystrom 1988: 84; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1967d: 40.

natalis Blackman 1921: 8. Lectotype ♂?; Agricultural College, Mississippi [USA]; USNM, Washington. Synonymy: Bright 1976c: 427.

Notes: (3) Blackman 1928b: 37 (redescribed).

References: (cn) Blackman 1950. (hb) Baker, W. L. 1972: 255; Beal & Massey 1945: 130; Blackman 1922b: 101–105, 1950; Chamberlin 1939: 360. (ds) Beal & Massey 1945: 130; Blackman 1922b, 1950; Chamberlin 1939: 360; Kleine 1934a: 160; Leng & Mutchler 1927: 52. (tx) Beal & Massey 1945: 130; Blackman 1921: 8, 1922b: 101–105, 1928b: 37–40; Bright 1976c: 193, 427, 1977: 517; Chamberlin 1939: 360; de Ruette 1970: 110; Wood, S. L. 1964: 61.

lecontei Bright 1977: 525. Holotype ♀; Kitt Peak, Pima Co., Arizona [USA]; CNCI, Ottawa.

Figures: Bright 1981d: 366.

Distribution: North America (Chihuahua in Mexico/ Arizona, New Mexico in USA).

Hosts: *Pinus cembroides*, *P. edulis*, one pair in *P. ponderosa*.

References: (ds) Bright 1981d: 260; Wood, S. L. 1982b: 1065. (tx) Bright 1977: 525, 1981d: 260, 366; McNamara 1984: 756; Wood, S. L. 1982b: 1065.

leechi Wood 1977b: 215. Holotype ♀; 3 km NNE Angwin on N side of Howell Mt., Napa Co., California [USA]; CAS, San Francisco.

Distribution: North America (California in USA).

Hosts: *Pinus ponderosa*.

References: (ds) Bright 1981d: 134; Wood, S. L. 1982b: 1051. (tx) Bright 1981d: 134; Wood, S. L. 1977b: 215, 1982b: 1051.

leiophyllae Blackman 1942a: 205. Holotype ♀; Chalco, Mexico, Mexico; USNM, Washington.

Distribution: North America (Hidalgo, Mexico, Michoacan, Oaxaca, Puebla, Veracruz in Mexico).

Hosts: *Pinus cembroides*, *P. leiophylla*, *P. teocote*, *P. sp.*

References: (hb) Atkinson & Equihua 1985a: 93; Atkinson et al. 1986: 39. (ds) Atkinson & Equihua 1985a: 93, 1985c: 360; Atkinson et al. 1986: 39; Blackwelder 1947: 782; Bright 1981d: 120; Wood, S. L. 1982b: 1057. (tx) Blackman 1942a: 205–206; Bright 1977: 515, 1981d: 120; Schedl 1955g: 24; Wood, S. L. 1982b: 1057.

auctor Blackman 1942a: 214. Holotype ♀; near Perote, Veracruz, Mexico; USNM, Washington. Synonymy: Bright 1977: 514.

References: (ds) Blackwelder 1947: 781. (tx) Blackman 1942a: 214; Bright 1977: 514; McNamara 1984: 754.

lenis Wood 1976c: 348. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Liana.

References: (ds) Bright 1981d: 101; Wood, S. L. 1982b: 1082. (tx) Bright 1981d: 101; McNamara 1984: 756; Wood, S. L. 1976c: 348, 1982b: 1082.

lepidus Bright 1977: 526. Holotype ♀; 5.6 km S Suchistepec on Highway 175, Oaxaca, Mexico, 2600 m; CNCI, Ottawa.

Figures: Bright 1981d: 356.

Distribution: North America (Oaxaca, Veracruz in Mexico).

Hosts: *Pinus lawsonii*, *P. sp.*

References: (hb) Atkinson et al. 1986: 39. (ds) Atkinson et al. 1986: 39; Bright 1981d: 191; Wood, S. L. 1982b: 1071. (tx) Bright 1977: 526, 1981d: 191, 356; McNamara 1984: 756; Wood, S. L. 1982b: 1071.

leris Wood 1986: 273. Holotype ♀; 25 km NW Flagstaff, Arizona (Hart's Prairie Road) [USA]; Wood Collection.

Distribution: North America (Arizona in USA).

Hosts: *Pinus ponderosa* twigs.

References: (tx) Wood, S. L. 1986: 273.

lichtensteini (Ratzeburg) 1837: 162 (*Bostrichus*).

Syntypes, sex?, Berlin, Germany; not located.

Figures: Grune 1979: 116, Pfeffer 1989a: pl. 9.

Distribution: Asia (Fujian in China/ E USSR), Europe (Austria/ Belgium/ Bulgaria/ Czechoslovakia/ Denmark/ England/ Finland/ France/ Germany/ Greece/ Hungary/ Italy/ Luxembourg/ Netherlands/ Norway/ Poland/ Spain/ Sweden/ Switzerland/ W USSR/ Yugoslavia).

Hosts: *Pinus cembra*, *P. nigra*, *P. peuce*, *P. sylvestris*, *Picea obovata*, rare in *Larix sibericae*.

Notes: (3) Pfeffer 1940: 108, 112 (cites var. *robustus*, synonymy in Pfeffer 1965: 63).

References: (ay) Escherich 1923b: 483; Feytaud 1950a; Lekander 1959b: 36; Numberg 1928a: 141; Nusslin 1911a: 109; Scherb 1971. (bv) Grune 1979: 117; Hellen 1921. (cn) Barbey 1925: 281; Chorbadzhievo 1929; Eckstein 1926: 579; Egorov 1958: 1482; Escherich 1923b: 483; Esterberg 1959; Feytaud 1950a; Greckin 1962b: 707; Hellen 1921; Joly 1976; Judeich & Nitsche 1895: 526; Kangas 1937; Kobakhidze 1960: 1853; Koch 1913: 123; Kontkanen 1932: 59; Maar & Voore 1935: 641–651; Muller 1912: 185; Nestertschuk 1930: 172; Nosek 1952b: 103; Nusslin 1913: 205; Pfeffer 1933: 43; Pierce, W. D. 1917: 75; Rhumbler 1922: 308, 1927: 322; Saalas 1949: 343, 380; Schimitschek 1937c: 55, 1955c: 87; Schwerdtfeger 1944a: 177, 1957a: 183; Seitner 1913a: 27; Wachtl 1883b: 319, 1901: 377, 381; Weber, H. 1926: 579; Wichmann 1927b: 353. (ec) Barbey 1927; Boucek 1974; Carpelan 1921; Jamnicky 1957b: 26; Kangas 1937; Karpinski 1932a: 95; Kleine 1908c: 213; Kobakhidze 1960: 1853; Kostin 1964: 110; Lundberg 1984; Michalski & Ratajczak 1989; Nosek 1952b: 103, 1959a: 118, 1959b: 87; Nuorteva 1956a: 17, 1968a, 1970; Nusslin 1927: 322; Okolow 1963; Pfeffer 1923a: 332, 1928b: 3, 1933: 43, 1960: 345; Roubal 1934a: 86; Rulm 1956b: 4; Saalas 1930: 119, 1949: 343, 380; Schwerdtfeger 1944a: 177, 1957a: 183; Sedlaczek 1935a: 163; Seitner 1913a: 27; Seitner & Notzl 1925: 185; Tragardh 1927a: 210; Wichmann 1959: 413. (hb) Altum 1881c: 322; Barbey 1901: 23, 74, 1925: 281; Bargmann 1906; Bukowsky 1930; Chararas 1962c: 363; Charvat 1950; Chorbadzhievo 1929; Dombrowsky 1887, 1891; Eckstein 1926: 579; Eichhoff 1881a: 47, 193; Escherich 1923b: 483; Esterberg 1959; Feytaud 1950a; Fuchs 1904a; Gerhard 1908: 157–162; Gornostaev 1916: 312; Györfi 1957; Heinemann 1908: 156–164, 1909; Henschel 1895a: 172; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski 1933b: 30; Karpinski & Strawinski 1948: 156; Knotek 1897: 136; Kostin 1960: 134; Krivolitskaya 1973: 140; Lekander 1959b: 36; Lengerken 1939: 65, 1954: 85; Michalski 1959a: 291; Munro 1920, 1926: 63; Nordlinger 1856: 27; Nosek 1959a: 118, 1959b: 87; Numberg 1929a: 111, 1929c: 121; Nuorteva 1968a, 1970; Nusslin 1898: 280, 1913: 205, 1927:

322; Pfeffer 1941b: 3; Postner 1974: 435; Ratzeburg 1839: 197; Rhumbler 1922: 308, 1927: 322; Rinski-Korsakov et al. 1949: 288; Rupertsberger 1879: 231, 1880: 229; Saalas 1919, 1949: 343, 380; Schwerdtfeger 1944a: 177, 1957a: 183, 1981: 190; Sedlacek 1935a: 163; Seitner & Notzl 1925: 188; Spessivtsev 1913a: 59, 1921b: 221; Stark 1926a: 336, 1952: 350; Tschorbadjiev 1929: 170; Wachtl 1876a: 460, 1901: 377, 381; Weber, H. 1926: 579; Wichmann 1927b: 353. (**ds**) Acloque 1896; Allen, A. 1954; Bangsholt 1975: 95; Barthe 1896; Bau 1888; Bejer-Petersen & Jorum 1977: 25; Blandford 1891c; Borchert 1951; Brakman 1966b: 206; Brundin 1934; Bucking 1932; Bukowsky 1930; Buresh & Lazarov 1956; Calwer 1884, 1893; Carpelan 1921; Chapuis & Candeze 1853; Chararas 1962c: 363; Charvat 1950; Chorbazhievo 1924d, 1929; Csiki 1914; Eggers 1904; Endrodi 1958b; Ericson & Saindin 1893; Escherich 1923b: 483, 1932b; Esterberg 1928; Everts 1922: 643, 1925; Favre 1890; Fedorov 1930; Fowler 1891; Fuchis 1904a; Gaubil 1849: 126; Gemminger & Harold 1872: 2688; Gerhardt 1900; Gobbi 1989: 61; Gornostaev 1917; Gozis 1875: 80; Greekin 1962b: 707; Grill 1895: 311; Grune 1979: 117; Gussmann 1919: 62; Hagedorn 1910d: 72; Hansen, V. 1939, 1956, 1964: 463; Heinemann 1908a, 1908b; Hellen 1947; Henschel 1895a: 172; Heyden 1876: 299, 1879: 140; Heyden, Reitter, & Weise 1883: 182, 1891: 672, 1906: 711; Horion 1951; Janovsky & Tegshzhargal 1955: 41; Joly 1976; Judeich & Nitsche 1895: 526; Kaltenbach 1874: 685; Karpinski 1926: 82, 1931: 25, 28, 1932a: 95, 1933b: 30; Karpinski & Strawinski 1948: 156; Kersten 1933: 75; Kiefer et al. 1942: 528; Klefbeck & Sjoberg 1960: 230; Kleine 1912a: 267, 1913a: 35, 1913b: 140, 1934a: 160; Kloft & Hinks 1945: 218; Kolbe, W. 1918: 211; Kontkanen 1932: 59; Kotula 1873b: 80; Kraatz 1869: 59; Krivolitskaya 1965a: 238, 1973: 140, 1983; Krogerus 1921b: 115; Kunnemann 1921: 58; Kurir 1947c: 6; Lacordaire 1866: 382; Leclercq 1971; Lindberg & Saris 1952: 59; Lomnicki 1886a: 242, 1913b: 148; Lucht 1987: 278; Lundberg 1974: 92, 1981: 151; Maren 1957b: 213; Mequignon 1936: 54; Michalski 1957: 165; Mirzoiian 1950: 141; Munster 1928: 289; Negru 1968a: 456; Numberg 1928b: 88, 112, 1935: 126, 1954: 70; Nusslin 1898: 280; Perris 1876a: 255, 1877a: 415; Pfeffer 1924b: 471, 1925b: 3, 1931b: 74, 1935: 158, 1947e: 17, 1960: 346, 1976: 340, 1984: 277, 1989a: 64; Pierce, W. D. 1917: 75; Pittioni 1943: 176; Postner 1974: 435; Ragusa 1924: 116; Rapp 1934: 728; Ratzeburg 1839: 197; Redtenbacher 1858: 830, 1874: 379; Reitter 1869b: 154, 1894a: 75, 1916: 295; Rinski-Korsakov et al. 1949: 288; Rouhal 1941: 269; Ruskov 1928c: 61; Rye 1866: 93; Saalas 1930: 119, 1931: 70; Sainte-Claire & Mequignon 1938: 447; Schaufuss 1915: 1242; Schaum 1859: 95, 1862: 101; Schedl 1980a: 23, 1981b: 78;

Schilsky 1909: 188; Schwerdtfeger 1981: 190; Seidlitz 1872: 394, 1891a: 564, 1891b: 610; Sokanovskii 1960; Stark 1926a: 336, 1926b: 105, 1926j: 127, 1931a: 28, 1952: 350; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 441; Stierlin & Cantard 1871: 292, 1906: 205; Strand 1940: 601; Strand & Hanssen 1935: 70; Thomson 1865: 368, 1868: 223; Trappen 1935: 143; Tredl 1907: 14; Tschorbadjiev 1929: 170; Wachtl 1870: 259, 1876a: 460; Wichmann 1927a: 72; Winter, T. G. 1983: 22; Wiren 1945: 43; Yanovskii 1977a, 1989: 64; Yanovskii & Tegshzhargal 1955: 410; Zinovjev 1955: 187; Zivojinovic 1960: 25. (**tx**) Acloque 1896; Bach 1854; Balachovsky 1949a: 236; Barbey 1901: 23, 74; Blandford 1891c; Chapuis & Candeze 1853; Charvat 1950; Chorbazhievo 1924d; Cola 1972: 12; Cola & Freude 1972; Doebner 1860; Dombrowsky 1887, 1891; Duffy 1953; Eggers 1915b: 13–14, 1931b: 31, 1941b: 222; Eichhoff 1864: 39, 1868d: 422, 1875: 201, 1878b: 176, 1881a: 47, 193, 1883a: 110, 134; Endrodi 1957a: 307, 1957b; Escherich 1923b: 483; Everts 1922: 643; Fairmaire 1864: 33; Fauvel 1889; Ferrant 1911; Ferrari 1867a: 32–33, 1867b: 114; Fleischer 1927; Formanek 1907: 37; Grune 1979: 116–117; Hagedorn 1910a: 101; Hansen, V. 1956, 1964: 463; Henschel 1895a: 172; Hopkins 1914: 127; Iablokoff-Khuzorian 1961: 105; Jacquelin du Val & Fairmaire 1868: 104, 112; Jansson 1936: 220; Joly 1976; Judeich & Nitsche 1895: 526; Karpinski & Strawinski 1948: 156; Koch 1913: 123, 1932: 121; Kuhn 1913: 1056; Lacordaire 1866: 382; Lekaneder 1959b: 36; Letzner 1891: 375; Lovendal 1898: 131; Lucht 1987: 278; Nordlinger 1848: 242, 1856: 27; Numberg 1928a: 141, 1929c: 121, 1938: 126, 1954: 70; Nusslin 1911a: 109; Perris 1877a: 415; Pfeffer 1932b: 23, 1933: 3–54, 1940b: 108, 112, 117, 1941b: 3, 1947e: 17, 1955a: 215, 1965: 63, 1976: 340, 1989a: 9; Pjatnitskii 1931: 169; Portevin 1935: 333; Postner 1974: 435; Quaschik 1953: 35; Ratzeburg 1837: 162, 1839: 197; Redtenbacher 1849a: 790, 1849b: 26, 1858: 830, 1874: 379; Reitter 1894a: 75, 1898b: 356, 1901: 101, 1913a: 93, 1916: 298; Rhumbler 1922: 308, 1927: 322; Rupertsberger 1879: 231, 1880: 229; Saalas 1949: 343, 380; Schedl 1934f: 1643, 1938a: 158, 1952f: 87, 1980a: 23, 1981b: 78; Scherb 1971; Schimitschek 1937c: 55, 1955c: 87; Seidlitz 1872: 394, 1891a: 564, 1891b: 610; Seitner 1887: 44; Sokanovskii 1929: 670–672, 1959b: 93–94, 1960: 677; Spessivtsev 1913a: 89, 1922a: 487, 1922b: 221, 1925a: 182, 1925b: 27, 1926: 50, 1929c: 298, 1931: 61–63; Stark 1952: 350; Stierlin 1898: 441; Thomson 1865: 368, 1868: 223; Wichmann 1918: 106–107. (**ms**) Eichhoff 1868d: 422; Escherich 1932b; Heinemann 1908a; Hellen 1921; Michalski 1951a: 291; Wichmann 1959: 413.

- liquidambarus** Blackman 1921: 14. Lectotype ♀; Mound, Louisiana [USA]; USNM, Washington, designated by Bright 1976b: 183. Figures: Bright 1981d: 338. Distribution: North America (Arkansas, Connecticut, District of Columbia, Florida, Georgia, Illinois, Indiana, Louisiana, Mississippi, North Carolina, West Virginia in USA). Hosts: *Liquidambar styraciflua*. References: (cn) Blackman 1950, (cc) Bright 1981d: 64; Burks 1979: 856; Bushing 1965: 464; Marsh 1977: 150. (hb) Baker, W. L. 1972: 255; Beal & Massey 1945: 131; Blackman 1922b: 101–105, 1950; Bright 1981d: 64; Chamberlin 1939: 361; Ostmark 1968. (ds) Beal & Massey 1945: 131; Blackman 1922b: 101–105, 1950; Bright 1981d: 64; Chamberlin 1939: 361; Deyrup & Atkinson 1987b: 68; Drooz 1955: 363; Kirk 1970; Kleine 1934a: 160; Leng & Mutchler 1927: 52; Ostmark 1968; Weber, B. C., & McPherson 1991: 54; Wood, S. L. 1982b: 1133. (tx) Beal & Massey 1945: 131; Blackman 1921: 14, 1922b: 101–105, 1928b: 37–41; Bright 1976b: 183, 1981d: 64, 338; Chamberlin 1939: 361; de Ruelle 1970: 110; Wood, S. L. 1982b: 1133.
- litos** Bright 1976c: 435. Holotype ♀; 40 km W Orizaba, Veracruz, Mexico; CNCI, Ottawa. Distribution: North America (Veracruz in Mexico). Hosts: *Pinus* sp. References: (ds) Bright 1981d: 136; Wood, S. L. 1982b: 1055. (tx) Bright 1976c: 435, 1981d: 136; McNamara 1984: 756; Wood, S. L. 1982b: 1055.
- longipilus** Schedl 1951m: 112. Holotype ♀; Bolivien, Cochabamba; Eggers Collection, in NHMW, Wien. Distribution: South America (Bolivia). References: (tx) Schedl 1951m: 112, 1979c: 141.
- madagascariensis** Schedl 1951j: 21. Syntypes, sex²; Tananarive, Tsimbazaza; IRSM, Madagascar, and 2 ♂, 2 ♀ in Schedl Collection in NHMW, Wien. Distribution: Madagascar. Hosts: *Vernonia appendiculata*. References: (hb) Paulian 1951: 28. (ds) Paulian 1951: 28; Schedl 1977b: 112. (tx) Schedl 1951j: 2, 1953d: 69, 1965c: 69, 1977b: 112, 1979c: 144.
- malleatus** Bright 1978: 79. Holotype ♀; Walker, Yavapai Co., Arizona [USA]; CNCI, Ottawa. Figures: Bright 1981d: 351. Distribution: North America (Arizona in USA). Hosts: *Pseudotsuga menziesii*. References: (ds) Bright 1981d: 164. (tx) Bright 1978: 79, 1981d: 164, 351; McNamara 1984: 756.
- mandibularis** Schedl 1951m: 113. Syntypes ♂ ♀; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, and Plannmann Collection. Distribution: South America (Brazil). Notes: (1) Schedl 1979c: 148 (citation of holotype invalid).
- References: (ds) Schedl 1966f: 86. (tx) de Ruelle 1970: 110; Schedl 1951m: 112–113, 1979c: 148.
- maroantsetrae** Schedl 1965c: 69. Holotype ♂ ♀; Madagascar, Maroantsetra; Schedl Collection in NHMW, Wien. Distribution: Madagascar. References: (tx) Schedl 1965c: 69, 1977b: 113, 1979c: 149.
- mauretanicus** Peyerimhoff 1930: 259. Syntypes, sex²; Algerie: Djurdjura a Tidjda, 1750 m; MNHN, Paris. Distribution: Africa (Algeria/ Egypt/ Libya/ Morocco/ Tunisia), Europe (France). Hosts: *Pinus nigra mauretanicus*. Notes: (1) Originally named as a subspecies of *glabratus*. Pfeffer 1940: 121 (raised to specific rank). References: (ec) Berland 1934: 215. (hb) Chararas 1962c: 363. (ds) Chararas 1962c: 363; Peyerimhoff 1933b: 370; Pfeffer 1935: 158, 1976: 340, 1984: 277. (tx) Balachowsky 1949a: 239; Peyerimhoff 1930: 255, 259; Pfeffer 1940b: 109, 112, 1955a: 211, 1976: 340; Schedl 1934f: 1643, 1979c: 106.
- medialis** Wood 1976c: 361. Holotype ♀; Volcan Irazu, Cartago, Costa Rica, 2300 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Quercus* sp. References: (ds) Bright 1981d: 106; Wood, S. L. 1982b: 1084. (tx) Bright 1981d: 106; McNamara 1984: 756; Wood, S. L. 1976c: 361, 1982b: 1084.
- megas** Bright 1976c: 436. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico; CNCI, Ottawa. Distribution: North America (Nuevo Leon in Mexico). Hosts: *Pinus cumminicola*. References: (ds) Bright 1981d: 298; Wood, S. L. 1982b: 1113. (tx) Bright 1976c: 436, 1981d: 298; McNamara 1984: 756; Wood, S. L. 1982b: 1113.
- melanurus** Wood 1976c: 364. Holotype ♀ ♀; 8 km W San Cristobal de las Casas, Chiapas, Mexico; CNCI, Ottawa. Distribution: North America (Chiapas in Mexico). Hosts: *Quercus* sp. References: (ds) Bright 1981d: 93; Wood, S. L. 1976c: 364, 1982b: 1026. (tx) Bright 1981d: 3; Wood, S. L. 1976c: 364.
- mendosus** Wood 1975b: 397. Holotype ♂; San Isidro del General, Costa Rica, 1000 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: "Fosforo" leaf petioles. References: (ds) Bright 1981d: 95; Wood, S. L. 1982b: 1029. (tx) Bright 1981d: 95; McNamara 1984: 756; Wood, S. L. 1975b: 397, 1982b: 1029.
- mesembria** Bright 1978: 80. Holotype ♀; Cerro Calel, Quezaltenango, Guatemala; Wood Collection. Distribution: North America (Guatemala). Hosts: *Abies guatemalensis*.

References: (**ds**) Bright 1981d: 318; Wood, S. L. 1982b: 1088. (**tx**) Bright 1978: 80, 1981d: 318; McNamara 1984: 756; Wood, S. L. 1982b: 1088.

mexicanus Blackman 1928b: 121. Holotype ♀; Correen, Coahuila, Mexico; USNM, Washington. Distribution: North America (Coahuila, Jalisco, Tabasco in Mexico).

Hosts: *Parthenium argentatum*.

References: (**cn**) Lloyd 1911: 43. (**hb**) Bright 1981d: 30; Lange 1944: 395; Lloyd 1911: 43. (**ds**) Blackwelder 1947: 782; Bright 1981d: 30; Ferrer 1942; Kleine 1934a: 160; Lloyd 1911: 43; Romney 1946: 670–671; Wood, S. L. 1982b: 1118. (**tx**) Blackman 1928b: 121; Bright 1981d: 30; McNamara 1984: 756; Schedl 1940a: 348; Wood, S. L. 1982b: 1118.

micans Bright 1981d: 188. Holotype ♀; Chiltepec, Oaxaca, Mexico; CNCI, Ottawa.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Pinus montezumae*.

References: (**ds**) Wood, S. L. 1982b: 1070. (**tx**) Bright 1981d: 188; Wood, S. L. 1982b: 1070.

micrographus (Linnaeus) 1758: 355 (*Dermestes*). Syntypes, sex?; Europa; not located.

Figures: Postner 1974: 433 (adult).

Distribution: Asia (Turkey), Europe (Austria/ Bulgaria/ Czechoslovakia/ Denmark/ England/ Finland/ France/ Germany/ Greece/ Hungary/ Italy/ Norway/ Luxemburg/ Poland/ Romania/ Spain/ Sweden/ Switzerland/ W USSR/ Yugoslavia).

Hosts: *Abies bornmuelleriana*, *Picea abies*, *P. excelsa*, *P. obovata*, *P. omorika*, *Larix sibirica*.

References: (**ay**) Escherich 1923b: 463, 483, 601; Hadorn 1933; Nusslin 1911a: 5, 51, 60, 278, 338, 375; Sedlacek 1902b: 244. (**bv**) Grune 1979: 117; Kinelski 1958: 73; Meixner 1937: 1214; Naumann-Etienne 1978a; Nuorteva 1956c: 93; Rozhkov 1970: 143; Tragardh 1930b: 102. (**cn**) Androic 1966: 49; Baisch 1954: 321; Barbey 1906a, 1925: 70; Besceli 1963, 1964: 50–57; Borcea 1924; Borodajewsky 1929: 515–521, 1931: 155–156; Eckstein 1926: 579; Eliescu & St. Negru 1956; Escherich 1923b: 463, 483, 601, 1929, 1930a; Esterberg 1959; Feytaud 1950a; Gabler 1955; Hartig 1861: 330, 1877: 196; Hess 1898: 348; Hess & Beck 1914: 283, 1927: 339; Jacentkovsky 1933: 271; Jaensch 1835: 50; Judeich & Nitsche 1895: 524; Juntinen 1960: 25; Kamp 1950; Kangas 1947: 1–192, 1958d: 164; Keller 1903b: 48; Kholodkovskii 1912: 307; Koch 1913: 125; Komarek et al. 1931: 1–256; Kovacevic 1952: 69, 1957: 67; Kozikowsky 1929: 252; Maksimovic & Barlov 1961: 37; Maksimovic & Milanovic 1964, 1966; Marcu 1926c: 65; Merker 1954b: 209; Muller 1912: 252; Mumford 1960: 37; Nusslin 1913: 205; Ohnesorge 1955: 280; Pfeffer 1928a: 467, 1947c: 204, 1948a: 801, 1949b: 147, 1950c: 3; Pierce, W. D. 1917: 69; Postner 1971; Ratzeburg 1871c: 81; Rhumbler 1927: 321; Rozhkov 1970: 143; Saalas

1949: 343, 379; Scheidter 1919: 88; Schimitschek 1930d: 136, 1935b: 149, 1938b: 115, 1944: 182, 1947g: 191, 1955a: 36–37, 1955c: 87; Schneeberg 1925: 495; Schwerdtfeger 1944a: 181, 1957a: 186; Sedlacek 1933: 307; Sierpinski 1958: 69; Sinreich 1961: 166; Slander 1948: 10; Spaic 1956: 88, 1966: 454; Springer 1936b: 178; Thalenhorst 1950: 90; Tragardh 1938a: 12; Vappula 1965: 153; Wachtl 1883b: 319, 1901: 381; Walsh 1867: 103; Wardle 1929: 322; Weber, H. 1926: 579; Wichmann 1927b: 353; Zieger 1948a: 378. (**cc**) Baisch 1954: 321; Balazy 1962; Balazy & Michalski 1960; Barbey 1906a, 1927; Borodin 1962; Egger, A. 1974; Gauss & Wellenstein 1950; Geschwind 1918; Graham 1969: 877; Györfi 1941b; Heqvist 1956, 1957a, 1963: 154; Kangas 1946b: 21, 1971; Karpinski 1932a: 101; Kielczewski 1976; Kielczewski & Wisniewski 1983; Kleine 1908c: 213, 1909a: 47, 77, 1944: 69–70; Kraemer 1950b: 380; Lundberg 1984; Merker 1954b: 209; Michalski & Ratajczak 1989; Naumann-Etienne 1978a; Numberg & Wiackowski 1958: 130; Nuorteva 1956a: 17, 1957b: 62, 1968a, 1970, 1971; Nusslin 1927: 321; Okolov 1963; Pfeiffer 1923a: 331, 1943b: 179, 1947c: 204, 1949b: 147, 1950c: 3; Poinar 1975: 167; Rühm 1956b: 4; Saalas 1937: 154, 1949: 343, 379, 1951: 16; Schimitschek 1930a: 340, 1955a: 36–37, 1957a: 186; Schwerdtfeger 1944a: 181; Sedlacek 1908: 53, 1933: 307, 1935a: 163; Seitter 1913a: 27; Sitowski 1930: 4; Szczepanski 1960a: 408, 1961: 3; Thompson, W. R. 1943: 91; Tragardh 1925a: 171, 1927a: 208; Vietinghoff 1924: 335; Wiackowski 1957b: 312; Zinovjev 1958: 386; Zinn 1985b. (**hb**) Apfelbeck 1916: 429–439, 1917; Barbey 1901: 23, 77, 1913, 1925: 70; Bargmann 1906; Beffa 1949; Berg 1827; Besceli 1963; Bloch 1776; Borcea 1924; Borodajewsky 1930a; Brandt 1948; Cecconi 1906, 1924; Charvat 1950; Dombrowsky 1887, 1891; Eckstein 1889, 1897, 1926: 579; Eichhoff 1881a: 47, 197; Egger, A. 1974; Escherich 1923b: 463, 483, 601; Feisthamel 1835; Feytaud 1950a; Fuchs 1904a, 1905c: 339, 1906: 46; Gabler 1955; Gornostaev 1916: 312; Györfi 1957; Hadorn 1933; Hartig 1861: 330, 1877: 196; Helmbacher 1924; Hennings 1908c: 218, 1908d; Henschel 1895a: 172; Hess 1898: 348; Hess & Beck 1914: 283, 1927: 339; Hufnagel 1887: 512; Hufnagel & Puzyr 1951: 108, 110; Jaensch 1835: 50; Judeich & Nitsche 1895: 524; Karpinski 1933b: 30; Karpinski & Strawinski 1948: 156; Kholodkovskii 1912: 307; Knotek 1894a: 558; Kozikowsky 1922a: 42; Kraemer 1948: 133, 1950b: 380; Lengerken 1939: 36, 65, 1954: 85; Lindberg 1963: 243; Loos 1913: 406; Lamardonii & Leonardi 1889: 458; Madon 1930: 99; Marcu 1930: 327–336; Mokrzecki 1925: 1–7; Naumann-Etienne 1978a; Numberg 1929: 110; Nuorteva 1956c: 93, 1968, 1970; Nusslin 1898: 281, 1906: 14, 1913: 205, 1927: 321; Ohnesorge 1955: 280; Pfeffer 1941c: 5; Postner 1974: 434; Ratzeburg 1837: 162, 1839:

- 197, 1871c: 81; Reh 1900a: 95; Rhumbler 1927: 321; Rodd 1897: 34; Rozhkov 1970: 143; Rupertsberger 1880: 230; Saalas 1949: 343, 379, 1951: 16; Schimitschek 1930a: 340, 1944: 182, 1955a: 36–37; Schneider-Orelli 1947b: 157; Schwerdtfeger 1944a: 181, 1957a: 186; Sedlaczek 1935a: 163; Seitner 1911: 99; Simmel 1918: 288–291, 1925: 154; Spessivtsev 1913a: 87, 1921b: 221, 1923: 205, 1925d: 103, 1928a: 221, 1929: 678–682; Stark 1926a: 336, 1952: 342; Tragardh 1929a: 313, 1930b: 102, 1930c: 474, 1931: 57, 1939b: 147, 227; Tredl 1908b: 140; Wachtl 1876a: 460, 1901: 381; Weber, H. 1926: 579; Wichmann 1927b: 353; Zinovjev 1958: 386; Zivojinovic 1950: 299–310.
- (ds) Acloque 1896; Ammann & Knabl 1923; Androic 1966: 49; Arnoldi et al. 1955: 712; Badoux 1921; Bakke & Kvamme 1977; Balazy & Michalski 1960; Barth 1896; Ban 1888; Beffa 1949; Belonsov 1916, 1917: 336; Biels 1887; Blanchere & Robert 1889; Blandford 1891a; Borcea 1924: 221–260; Brakman 1966b: 206; Branesik 1871, 1906; Buresh & Lazarov 1956; Calver 1893; Cecconi 1897, 1906; Charvat 1950; Chorbazhiev 1924d; Credler 1868; Crotch 1863; Dejean 1821, 1825; Duftschmidt 1825; Eder 1934; Eggers 1904; Ericson & Sandin 1893; Escalera 1919; Escherich 1923b: 463, 483, 601, 1929, 1930a; Esterberg 1928, 1959; Favre 1890; Florov 1940: 94; Fowler 1882: 64, 1891; Fuchs 1904a, 1905a, 1905c: 339, 1907: 46, 1913; Ganbil 1849: 126; Gauss & Wellenstein 1950; Gemminger & Harold 1872: 2688; Gerhardt 1898a; Geschwind 1918; Gillefors 1966; Goeze 1777: 130; Gornostaev 1917: 308–315; Gredler 1866: 374, 1869: 73, 1875: 115; Grill 1895: 311; Gronzelle 1905; Grune 1979: 117; Gussmann 1919: 62; Gyllenhal 1827: 624; Györfi 1941b; Hagedorn 1910d: 72; Hansen, V. 1939; Heinemann 1908a; Hellen 1947; Hennings 1954: 256, 261; Henschel 1879a, 1895a: 172; Heyden 1879: 140; Heyden, Reitter, & Weise 1883: 181, 1891: 672, 1906: 712; Holdhaus & Denbel 1910: 145; Horion 1935, 1949, 1951, 1956: 123; Hormuzaki 1891: 174; Ihssen 1939: 336; Jazentkovsky 1933: 271; Jazentkovsky 1912: 288; Judeich & Nitsche 1895: 524; Kangas 1971; Karpinski 1925: 216, 1926: 82, 1931: 19, 28, 1932a: 101, 1933a: 290, 1933b: 30; Karpinski & Strawinski 1948: 156; Keler 1925b: 272; Kersten 1933: 76; Kestercanek 1881a: 12; Kinelski 1958: 73; Klefbeck & Sjöberg 1960: 230; Kleine 1912a: 262, 264, 268, 1913a: 35, 1913b: 140, 1934a: 160; Knotek 1892a: 37, 1894a: 558; Koca 1900: 116; Kolbe, W. 1916: 211, 1918: 211; Kovacevic 1957: 67; Kozikowsky 1921: 181, 1922: 42–43; Kraatz 1869: 59; Kraemer 1948: 133; Krivolutskaia 1983; Krol 1877: 34; Kurir 1947c: 14; Lacordaire 1866: 382; Langhoffer 1915c: 158; Leder 1871: 132; Leiler & Prutz 1977; Liegel 1886: 43; Lindberg & Saris 1952: 59; Lokaj 1868: 64; Lomnicki 1886a: 242, 1913b: 148; Loos 1913: 406; Lmardoni & Leonardi 1889: 458; Lundberg 1974: 92, 1979: 31; Lundblad 1950c: 115; Marcu 1926c: 65, 1957b: 213; Matthews & Fowler 1883: 42; Meinert 1887: 70; Mumford 1960: 37; Munster 1928: 259; Novak, P. 1964; Numborg 1927a: 214, 1928b: 88, 112, 1938: 126, 1954: 70, 1960: 158; Nuorteva 1956b: 168, 1971: 67; Nusslin 1898: 281; Orest 1926c: 65; Ortzen 1886: 279; Palm 1946: 122, 1948a: 91; Pfeffer 1924a: 96, 1924b: 471, 1935: 158, 1947c: 4, 1950b: 73, 1976b: 340, 1989a: 63; Pierce, W. D. 1917: 69; Platonoff 1943: 141; Pomerantzev 1907b: 492; Poppins 1900: 108; Postner 1974: 434; Prossen 1913: 83; Rapp 1934: 729; Ratzeburg 1837: 162, 1839: 197; Redtenbacher 1874: 379; Reitter 1894a: 76, 1916: 298, 352; Ruskov 1928c: 61; Saalas 1931: 70; Sahlberg 1900: 106; Schaufuss 1915: 1242; Schaum 1859: 95, 1862: 101; Scheerpeltz & Winkler 1930: 258; Schilsky 1909: 188; Schimitschek 1938b: 115; Schiodte 1873: 103; Seidl 1876: 4; Seidlitz 1872: 394, 1891a: 564, 1891b: 610; Sharp & Fowler 1893: 34; Solia 1893: 217; Sparre-Schneider 1889: 61; Stark 1926a: 336, 1926b: 105, 1926j: 127, 1931a: 26, 1931d: 548, 1952: 342; Stein 1868: 114; Stein & Weise 1877: 164; Stephens 1829a: 145, 1830: 356, 1839: 206; Stierlin 1898: 44; Stierlin & Gautard 1871: 293; Strand 1946: 601; Sturm 1826: 102; Thomson 1865: 368, 1868: 223; Tragardh 1919: 237–248, 1930: 468–480, 1938: 10–14, 1939b: 147, 227; Trappen 1935; Tredl 1907: 14, 1908b: 140; Vappula 1965: 153; Villa & Villa 1833: 26; Wachtl 1876a: 460; Westhoff 1882: 238; Wichmann 1909b: 173, 1927a: 73–74; Yanovskii 1977a; Yanovskii & Dmitrienko 1983; Yanovskii & Tegshzhargal 1985: 410; Zinovjev 1955: 187.
- (tx) Acloque 1896; Balachowsky 1949a: 235; Barbey 1901: 23, 77; Bechstein et al. 1805; Beffa 1949; Bertolini 1872; Blandford 1891a; Branesik 1871; Charvat 1950; Chorbazhiev 1924d; Dejean 1821, 1825; Doebner 1860; Dombrowsky 1887, 1891; Duftschmidt 1825; Eggers 1914: 183–184, 1915b: 13–14, 1940g: 36; Eichhoff 1864: 39, 1868d: 424, 1878b: 183, 1881a: 47, 197, 1883a: 111, 135; Endrodi 1957b; Escherich 1923b: 463, 483, 601; Escherich & Escherich 1897; Fabricius 1801: 387; Fankhauser 1912a; Fanvel 1889; Ferrant 1911; Ferrari 1867a: 32–33, 1867b: 114; Fleischer 1927; Formanek 1907: 37; Fuchs 1913; Gabler 1955; Gebien 1907: 197; Gmelin 1790: 1601; Goeze 1777: 130; Grune 1979: 117; Gyllenhal 1813: 362, 1827: 624; Hagedorn 1910a: 101; Henry 1892: 14; Henschel 1895a: 172; Houlbert 1922a: pl. 1; Jacobson 1895: 521; Jacquelin du Val & Fairmaire 1868: 112; Jansson 1936: 220; Judeich & Nitsche 1895: 524; Karpinski & Strawinski 1948: 156; Knotek 1892a: 37; Koch 1913: 125, 1928: 99, 1932: 119; Kuhn 1913: 1056; Kurenzov 1941: 233; Lacordaire 1866: 382; LeConte 1858: 355; Letzner 1891: 376; Linnaeus 1758: 355, 1767: 143; Lovendal 1898: 131; Lucas

- 1920: 515; Lunardoni & Leonardi 1889: 458; Marshian 1802: 52; Meixner 1937: 1214; Nunberg 1930, 1938: 126, 1954: 70; Nusslin 1911a: 5, 51, 60, 110, 278, 338, 378; Olivier 1795b: 9; Paykull 1800: 155; Pfeffer 1940b: 107, 116, 1941c: 5, 1947c: 4, 1955a: 214, 1976: 340; Pjatnitskii 1931: 171, 173; Portevin 1935: 333; Postner 1974: 434; Quaschik 1953: 35; Ratzeburg 1837: 162, 1839: 197; Redtenbacher 1849a: 359, 851, 1849b: 26, 1874: 379; Reitter 1894a: 76, 1913a: 92, 1916: 298, 352; Rhumbler 1927: 321; Rupertsberger 1880: 230; Saalas 1949: 343, 379; Sahlberg 1836: 150; Schedl 1934f: 1643; Schimitschek 1937: 55, 1955c: 87; Seidlitz 1872: 394, 1891a: 564, 1891b: 610; Seitner 1887: 44; Simmel 1918: 288; Sokanovskii 1929: 670–672, 1954: 19; Spessivtsev 1913a: 87–88, 1922a: 487, 1922b: 221, 1923: 205–206, 1925a: 182, 1925b: 23, 1925d: 103–104, 1926b: 48–50, 1929c: 298, 303, 1931: 61–62; Stark 1927: 227–230, 1952: 342; Stephens 1829a: 145, 1829b: 12, 1830: 356, 1839: 206; Stierlin 1898: 441; Thomson 1865: 368, 1868: 223; Villers 1789: 214. (ms) Eichhoff 1868d: 424; Hartig 1834: 109; Heinemann 1908a; Kozikowsky 1929: 252; Lucas 1920: 515; Reh 1900a: 95; Seitner 1913a: 27.
- fennicus* Eggers 1914: 153. Lectotype ♂; Finland: Karislojo, Turtola; USNM, Washington, designated by Anderson & Anderson 1971: 13. Synonymy: Stark 1952: 342.
References: (bv) Pfeffer 1940b: 116; Tragardh 1930b: 102. (cc) Saalas 1917a: 18, 1928: 652. (hb) Saalas 1913a: 67, 86, 1914: 88, 305; Spessivtsev 1921b: 221; Tragardh 1930b: 102. (ds) Pfeffer 1924b: 471, 1935: 159; Saalas 1913a: 67, 86, 1916: 110–116, 1917a: 18; Schaufuss 1915: 1242. (tx) Anderson, W. H. & Anderson 1971: 13; Balachowsky 1949a: 235; Eggers 1914: 183–184, 1915b: 13–14; Endrodi 1957b: 417; Michalski 1969b: 569; Pfeffer 1940b: 116; Saalas 1913a: 67, 86; Schedl 1934f: 1643; Spessivtsev 1922a: 487, 1922b: 221, 1923: 205–206, 1925d: 103–104, 1926b: 48–50, 1931: 61–62; Tragardh 1930b: 102.
- micrographus sibiricus* Stark 1952: 344. Syntypes, sex?; Siberia, USSR; IZL, Leningrad.
References: (ds) Pfeffer 1976: 340. (tx) Pfeffer 1976: 340; Stark 1952: 344.
- micrographus curtulus* Sokanovskii 1954: 19. Syntypes ?; Poland; not located.
References: (ds) Pfeffer 1976: 340. (tx) Pfeffer 1976: 340; Sokanovskii 1954: 19.
- micrographus longulus* Sokanovskii 1954: 19. Syntypes ?; Poland; not located.
References: (tx) Endrodi 1957b: 417; Kinclaki 1958: 73; Nunberg 1960: 158; Sokanovskii 1954: 19.
- micrograptinus* Wood 1989: 179. Holotype, sex?; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien, automatic.
- Distribution: South America (Brazil).
References: (tx) Wood, S. L. 1989: 179.
- micrographus* Schedl 1972g: 60 (*Breviophthorus*). Holotype, sex?; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien, preoccupied by Linnaeus 1758.
References: (tx) Schedl 1972g: 60, 1979c: 152; Wood, S. L. 1989: 179.
- miniatus* Bright 1981d: 280. Holotype ♀; Cerro Pena Blanca, Honduras; Wood Collection.
Distribution: North America (Guatemala/ Honduras/ Oaxaca in Mexico/ Nicaragua).
Hosts: *Pinus lawsoni*, *P. oocarpa*, *P. pseudostrobus*.
References: (ds) Wood, S. L. 1982b: 1099. (tx) Bright 1981d: 280; Wood, S. L. 1982b: 1099.
- minus* Bright 1976c: 437. Holotype ♀; Hannagan Camp, Greenlee Co., Arizona [USA]; CNCI, Ottawa.
Distribution: North America (Arizona in USA).
Hosts: (?) *Pinus* sp.
References: (ds) Bright 1981d: 197; Wood, S. L. 1982b: 1036. (tx) Bright 1976c: 437, 1981d: 197; McNamara 1984: 757; Wood, S. L. 1982b: 1036.
- minutalis* Wood 1976c: 357. Holotype ♀; Palin, Esquintla, Guatemala; Wood Collection.
Distribution: North America (Guatemala).
Hosts: Shrub.
References: (ds) Bright 1981d: 42; Wood, S. L. 1982b: 1139. (tx) Bright 1981d: 42; McNamara 1984: 757; Wood, S. L. 1976c: 357, 1982b: 1139.
- minutus* Schedl 1963d: 221. Holotype ♂; Brasilien: Matto Grosso, Rio Caraguata; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1963d: 221, 1979c: 157.
- modicus* Blackman 1928b: 94. Holotype ♀; Las Vegas H.S. [Hot Springs], New Mexico [USA]; USNM, Washington.
Figures: Bright 1981d: 344.
Distribution: North America (Chihuahua, Durango in Mexico/ Arizona, California, Nevada, New Mexico, Utah in USA).
Hosts: *Pinus cembroides*, *P. edulis*, *P. monophylla*, *P. sabiniana*.
References: (cc) Bushing & Bright 1965: 203. (hb) Bright & Stark 1973: 112; Chamberlin 1939: 385. (ds) Bright 1964: 170, 1981d: 123; Bright & Stark 1973: 112; Chamberlin 1939: 385; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1035. (tx) Blackman 1928b: 89, 94–95; Bright 1964: 165–170, 1966b: 304, 1981d: 123, 344; Chamberlin 1939: 385; Wood, S. L. 1982b: 1035.
- navis* Blackman 1928b: 95. Holotype ♀; Morgan Hill, California [USA]; USNM, Washington.
Synonymy: Bright 1966b: 304.
References: (hb) Chamberlin 1939: 385. (ds) Chamberlin 1939: 385; Kleine 1934a: 160; Leng & Mutchler 1933: 52. (tx) Blackman

1928b: 95; Bright 1966b: 304; Chamberlin 1939: 385; de Ruette 1970: 110.

molestus Wood 1976c: 362. Holotype ♀; Los Abritos, San Luis Potosi, Mexico; Wood Collection. Distribution: North America (San Luis Potosi in Mexico).

Hosts: *Liquidambar styraciflua*.

References: (ds) Atkinson & Equihua 1988: 98; Bright 1981d: 61, 1986b: 683; Estrada & Atkinson 1988: 212; Wood, S. L. 1982b: 1131. (tx) Bright 1981d: 61; McNamara 1984: 757; Wood, S. L. 1976c: 362, 1982b: 1131.

montezumae Bright 1978: 81. Holotype ♀; 11 km E San Cristobal, Chiapas, Mexico; CNCI, Ottawa. Figures: Bright 1981d: 367.

Distribution: North America (Chiapas in Mexico).

Hosts: *Pinus montezumae*.

References: (hb) Atkinson et al. 1986: 39; Burgos & Saucedo 1983: 119. (ds) Atkinson & Equihua 1985a: 97; Atkinson et al. 1986: 39; Bright 1981d: 272, 1986b: 683; Burgos & Saucedo 1983: 119; Wood, S. L. 1982b: 1079. (tx) Bright 1978: 81, 1981d: 272, 367; McNamara 1984: 757; Wood, S. L. 1982b: 1079.

montivagus Bright 1977: 527. Holotype ♀; Km 77 on Highway 175 about 85 km N Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.

Figures: Bright 1981d: 348.

Distribution: North America (Oaxaca in Mexico).

Hosts: *Pinus ayacahuite*, *P. patula*.

References: (hb) Atkinson et al. 1986: 39. (ds) Atkinson et al. 1986: 39; Bright 1981d: 145; Wood, S. L. 1982b: 1094. (tx) Bright 1977: 527, 1981d: 145, 348; McNamara 1984: 757; Wood, S. L. 1982b: 1094.

mormon Bright 1977: 528. Holotype ♀; Parawan Canyon, Utah [USA]; Wood Collection.

Distribution: North America (Arizona, California, Utah in USA).

Hosts: *Pinus aristata*, *P. contorta*, *P. flexilis*.

References: (ds) Bright 1981d: 236; Wood, S. L. 1982b: 1039. (tx) Bright 1977: 528, 1981d: 236; McNamara 1984: 757; Wood, S. L. 1982b: 1039.

morosovi Spessivtsev 1926b: 48. Syntypes, sex?; Central Russia, Gouv. Kaluga, Orlov; Spessivtsev Collection in IPKE, Eberswalde.

Figures: Tsai & Li 1959: 94, Yin, Huang, & Li 1984: 120.

Distribution: Asia (China/ E USSR), Europe (Czechoslovakia/ Finland/ Poland/ Sweden/ W USSR).

Hosts: *Picea excelsa*, *P. koraiensis*, *P. obovata*.

References: (bv) Grune 1979: 119. (cn) Esterberg 1959; Kontkanen 1932: 62; Pfeffer 1948a: 501, 1950c: 3. (ec) Karpinski 1932a: 103; Pfeffer 1950c: 3, 1960: 345. (hb) Karpinski 1933b: 30, 1955: 79; Karpinski & Strawinski 1948: 156; Postner 1974: 435; Stark 1952: 353. (ds) Esterberg 1959; Grune 1979: 119; Hansen 1939; Hellen

1947; Karpinski 1931: 18, 30, 1932a: 103, 1933b: 30, 1948b: 231, 1955: 79; Karpinski & Strawinski 1948: 156; Kleine 1934a: 160; Klefbeck & Sjöberg 1960: 230; Kontkanen 1932: 62; Lucht 1987: 278; Lundberg 1975: 12, 1977; Numberg 1954: 71; Nuorteva 1971: 71; Pfeffer 1960: 345, 1976: 340, 1989a: 66; Postner 1974: 435; Schedl 1980a: 23, 1981b: 79; Stark 1931a: 26, 1931d: 553, 1952: 353; Strand 1946: 601; Yanovskii & Tegshzhargal 1985: 410; Yin, Huang, & Li 1984: 120; Zinovjev 1955: 187. (tx) Blackman 1928b: 1–184; Grune 1979: 119; Karpinski & Strawinski 1948: 156; Lucht 1987: 278; Numberg 1938: 126, 1954: 71; Pfeffer 1940b: 109, 113, 124, 1955a: 221, 1976: 340; Postner 1974: 435; Schedl 1934f: 1643, 1980a: 23, 1981b: 79; Sokanovskii 1929: 670–672; Spessivtsev 1926b: 48, 1931: 63–64; Stark 1927: 227–230, 1952: 353; Tsai & Li 1959: 94; Yin, Huang, & Li 1984: 120.

morosus Wood 1976c: 362. Holotype ♀; Zamorano, Morazan, Honduras; Wood Collection.

Figures: Bright 1981d: 338.

Distribution: North America (Guatemala/ Honduras/ Chiapas, Veracruz in Mexico).

Hosts: *Eupatorium dalioides*, and a liana.

References: (ds) Bright 1981d: 62; Wood, S. L. 1982b: 1129. (tx) Bright 1981d: 62, 338; McNamara 1984: 757; Wood, S. L. 1976c: 362, 1982b: 1129.

impossae Schedl 1957d: 63. Holotype, sex?; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Berlinia acuminata*, *B. grandiflora*.

References: (bv) Schedl 1960f: 34. (hb) Schedl 1960f: 77, 1962j: 15. (ds) Schedl 1960f: 34, 1962j: 15. (tx) Schedl 1957d: 63, 1962j: 15, 1979c: 160.

mulungensis Schedl 1957d: 60. Holotype ♂?; Congo Belge; Kivu, Chute de la Rutshuru; MRCB, Tervuren.

Distribution: Africa (Kenya/ Zaire).

Hosts: *Caesalpinia decapetala*, and a liana.

References: (hb) Schedl 1962j: 16. (ds) Schedl 1962j: 16. (tx) Schedl 1957d: 60, 1962j: 16, 1979c: 161.

murrayanae Blackman 1922d: 138. Holotype ♀; Grand Lake, Colorado [USA]; USNM, Washington.

Figures: Bright 1981d: 372.

Distribution: North America (Alaska/ Alberta, British Columbia, Manitoba, New Brunswick, Northwest Territories, Ontario in Canada/ Arizona, California, Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming in USA).

Hosts: *Picea glauca*, *P. engelmannii*, *P. pungens*, *Pinus albicaulis*, *P. contorta*, *P. flexilis*, *P. ponderosa*, *P. radiata*.

Notes: (3) Blackman 1928b: 133 (redescribed).

Bright 1981d: 311 (recognized subsp. *aurulentus*; its status is very doubtful).

References: **(cn)** Lindgren 1980a: 68. **(cc)** Matthews 1970; Powell, J. M., Wong, & Melvin 1972: 15; Werner & Holsten 1984. **(hb)** Bright & Stark 1973: 115; Chamberlin 1939: 398, 1958: 160; Lindgren 1980a: 68. **(ds)** Bright 1976d: 183, 1981d: 311; Chamberlin 1925, 1939: 398, 1958: 160; Evans, D. 1978, 1983: 35; Gast et al. 1989: 385; Keen 1929a: 33; Kleine 1934a: 160; Leng & Mutchler 1927: 52; Werner & Holsten 1984; Wood, S. L. 1957c: 401, 1972a: 427, 1982b: 1103. **(tx)** Blackman 1922d: 138, 1928b: 115, 133; Bright 1981d: 311, 372; Chamberlin 1939: 398, 1958: 160; Evans, D. 1983: 35; Keen 1929a: 33; de Ruette 1970: 110; Wood, S. L. 1957c: 401, 1972a: 427, 1977d: 515, 1982b: 1103.

elongatus Swaine 1925b: 194. Holotype ♀: Midday Valley, Merritt, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Blackman 1928b: 133.

References: **(ds)** Kleine 1934a: 159; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 117, 133; Bright 1967b: 675; de Ruette 1970: 109; Swaine 1925b: 194.

gracilis Swaine 1925b: 195. Holotype ♀; Grant Co., Oregon [USA]; CNCI, Ottawa. Synonymy: Wood 1977d: 515.

Notes: (3) Blackman 1928b: 134 (re-described).

References: **(cn)** Lindgren 1980a: 68. **(hb)** Chamberlin 1939: 398, 1958: 160; Lindgren 1980a: 68. **(ds)** Bright 1976d: 182; Chamberlin 1939: 398, 1958; Evans, D. 1983: 35; Evans, D., Lowe, & Hunt 1978; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1948: 54, 1951a: 128, 1972a: 427. **(tx)** Blackman 1928b: 116, 134; Bright 1966b: 301, 1967b: 678; Chamberlin 1939: 398, 1958: 160; Evans, D. 1983: 35; Keen 1929a: 33; de Ruette 1970: 109; Swaine 1925b: 195; Wood, S. L. 1951a: 128, 1972a: 427, 1977d: 515.

cutleri Swaine 1925b: 195. Holotype ♀; Midday Valley, Merritt, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Wood 1977d: 515. Notes: (3) Blackman 1928b: 136 (re-described).

References: **(hb)** Chamberlin 1939: 399, 1958: 161. **(ds)** Bright 1976d: 181; Chamberlin 1939: 399, 1958: 161; Keen 1929a: 34; Kleine 1934a: 159; Leng & Mutchler 1933: 52; Wood, S. L. 1972a: 427. **(tx)** Blackman 1928b: 117, 136; Bright 1967b: 678; Chamberlin 1939: 399, 1958: 161; Keen 1929a: 34; de Ruette 1970: 109; Swaine 1925b: 195; Wood, S. L. 1972a: 427, 1977d: 515.

exilis Swaine 1925b: 196. Holotype ♀; Ochoco National Forest, Oregon [USA]; CNCI, Ottawa. Synonymy: Wood 1977d: 515.

References: **(hb)** Chamberlin 1939: 398. **(ds)**

Chamberlin 1939: 398; Leng & Mutchler 1933: 52; Patterson & Hatch 1945: 153. **(tx)** Blackman 1928b: 116, 134; Chamberlin 1939: 398; de Ruette 1970: 109; Swaine 1925b: 196; Wood, S. L. 1977d: 515.

tenuis Swaine 1925b: 196. Holotype ♀; Merritt, Midday Valley, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Wood 1957c: 401. References: **(hb)** Chamberlin 1939: 398. **(ds)** Chamberlin 1939: 398; Keen 1929a: 33; Kleine 1934a: 161; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 116, 134; Bright 1967b: 679; Chamberlin 1939: 398; Keen 1929a: 33; de Ruette 1970: 111; Swaine 1925b: 196; Wood, S. L. 1957c: 401.

depygus Blackman 1928b: 128. Holotype ♀; Clyde, Colorado [USA]; USNM, Washington. Synonymy: Wood 1977: 515.

References: **(hb)** Chamberlin 1939: 396. **(ds)** Chamberlin 1939: 396; Keen 1929a: 33; Kleine 1934a: 159; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 128; Chamberlin 1939: 396; Keen 1929a: 33; McNamara 1984: 755; Wood, S. L. 1977: 515.

watsoni Schedl 1930: 197. Holotype ♀; Nictor Lake, New Brunswick [Canada]; CNCI, Ottawa. Synonymy: Wood 1977d: 515.

References: **(hb)** Chamberlin 1939: 401. **(ds)** Beanline 1956; Bright 1976d: 188; Chamberlin 1939: 401; Leng & Mutchler 1933: 52. **(tx)** Chamberlin 1939: 401; de Ruette 1970: 112; Schedl 1930: 197–199, 1931b: 166; Wood, S. L. 1964: 65, 1977d: 515.

aurulentus Bright 1966b: 301. Holotype ♀; Shell Ridge at Walnut Creek, Contra Costa Co., California [USA]; CAS, San Francisco. Synonymy: Wood 1977d: 516.

Notes: (3) Bright 1981d: 311 (treated as a subspecies; validity doubtful). Wood 1982b: 1103 (This name is based on females with long frontal hair; this variation occurs sporadically throughout the range of the species).

References: **(hb)** Bright & Stark 1973: 116. **(ds)** Bright 1971d: 311; Bright & Stark 1973: 116. **(tx)** Bright 1966b: 301, 1981d: 311; de Ruette 1970: 108; Wood, S. L. 1977d: 516.

acceptus Bright 1981d: 315. Holotype ♀; Granite Pass, Big Horn Mountains, Wyoming [USA]; CNCI, Ottawa. Synonymy: Wood 1982b: 1103.

References: **(tx)** Bright 1981d: 315; Wood, S. L. 1982b: 1103.

namus Wood 1964: 64. Holotype ♀; Totolapan, Oaxaca, Mexico; Wood Collection.

Distribution: North America (Chiapas, Jalisco, Oaxaca in Mexico).

Hosts: *Bursera* sp.

References: **(cc)** Equihua & Atkinson 1986: 634. **(hb)** Atkinson et al. 1986: 72; Burgos & Saucedo 1983: 123; Equihua & Atkinson 1986: 634. **(ds)**

- Atkinson et al. 1986: 72; Bright 1981d: 78; Burgos & Sancedo 1983: 123; Equihua & Atkinson 1986: 634; Wood, S. L. 1982b: 1125. **(tx)** Bright 1981d: 78; McNamara 1977: 198; Wood, S. L. 1964: 64, 1982b: 1125.
- nebulosus** Wood 1976a: 363. Holotype ♀; Lake Catemaco, Veracruz, Mexico; CNCI, Ottawa. Figures: Bright 1981d: 341. Distribution: North America (Veracruz in Mexico). Hosts: *Bursera* sp. References: **(ds)** Bright 1981d: 100, 1986: 683; Estrada & Atkinson 1988: 212; Wood, S. L. 1982b: 1126. **(tx)** Bright 1981d: 100, 341; McNamara 1984: 757; Wood, S. L. 1976a: 363, 1982b: 1126.
- nemoralis** Wood 1976a: 351. Holotype ♀; Zamorano, Morazan, Honduras; Wood Collection. Distribution: North America (Costa Rica/ Honduras). Hosts: *Aristolochia anguicida*, and a liana. References: **(ds)** Bright 1981d: 55; Wood, S. L. 1982b: 1114. **(tx)** Bright 1981d: 55; McNamara 1984: 757; Wood, S. L. 1976a: 351, 1982b: 1114.
- niger** Schedl 1938a: 187. Holotype, sex?; Porto Alegre [Brasil]; USNM, Washington. Distribution: South America (Brazil). References: **(ds)** Blackwelder 1947: 782. **(tx)** Anderson, W. H. & Anderson 1971: 22; Schedl 1938a: 187–188, 1979c: 166.
- nigricans** Blandford 1904: 236. Lectotype ♂; Quiche Mountains, Guatemala; BMNH, London, designated by Bright 1976b: 184. Figures: Bright 1981d: 356. Distribution: North America (El Salvador/ Guatemala/ Chiapas, Durango, Hidalgo in Mexico). Hosts: *Pinus ayacahuite*, *P. rudis*, *P. strobiformis*. Notes: (3) Bright 1981d: 189 (This is *nitidulus* of Schedl 1956: 18 and Schwerdtfeger 1957: 497). References: **(ec)** Becker, G. 1953, 1955. **(hb)** Becker, G. 1953, 1955. **(ds)** Becker, G. 1953; Blackwelder 1947: 782; Bright 1981d: 189; Ferrer 1942; Hagedorn 1910d: 173; Kleine 1913b: 140, 1914b: 363; Schedl 1963c: 158, 1977e: 42; Wood, S. L. 1982b: 1071. **(tx)** Blandford 1904: 236; Bright 1976b: 184, 1978: 72, 1981d: 189, 356; Hagedorn 1910a: 101; Schedl 1938a: 183, 1940a: 348, 1955g: 18, 20; Wood, S. L. 1982b: 1071.
- chiapensis* Bright 1977: 522. Holotype ♀; 13 km E San Cristobal, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Bright 1978: 72. References: **(tx)** Bright 1977: 522, 1978: 72; McNamara 1984: 754.
- nitellus** Browne 1973a: 291. Holotype, sex?; Zaire: Mwaka, Bulongo; MRCB, Teveren. Distribution: Africa (Zaire). References: **(tx)** Browne 1973a: 291.
- nitidipennis** (Schedl) 1967d: 10 (*Breviophthorus*). Holotype, sex?; Brasilien, Chapeco, 24 07, 52 36, 600 m; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil).
- References: **(ds)** Schedl 1976a: 52. **(tx)** Schedl 1967d: 10–11, 1979c: 169.
- nitidulus** (Mannerheim) 1843: 298. Lectotype ♀; Sitka Island, Alaska; MZU, Helsinki, designated by Bright 1976b: 186. Figures: Bright 1981d: 351, Chamberlin 1958: 155 (galleries). Distribution: North America (Alaska/ Alberta, British Columbia in Canada/ Arizona, California, Idaho, Nevada, Oregon, Utah, Washington, Wyoming in USA). Hosts: *Picea engelmannii*, *P. glauca*, *P. sitchensis*, *Pinus contorta*, *P. muricata*, *P. radiata*. Notes: (3) Blackman 1928b: 108 (redescribed). References: **(bv)** Furniss, M. M., Baker, & Hostetler 1976: 1300; Melsheimer 1853: 87. **(en)** Anonymous 1968g; Chamberlin 1924; Currie 1905: 72; Doane et al. 1936; Essig 1926: 519, 1958: 519; Fall & Cockerell 1907: 217; Gillette 1903: 118; Hopkins 1904a: 17; Lindgren 1980a: 68; Ohmart & Voigt 1982: 345; Schuder 1969: 76; Swaine 1918a: 96, 100. **(ec)** Bushing & Bright 1965: 203; Chamberlin 1918a; Deyrup & Gara 1978: 275; Marsh 1979: 295; Ohmart & Voigt 1982: 340; Schedl 1958d: 187; Werner & Holsten 1984. **(hb)** Bright & Stark 1973: 113; Chamberlin 1939: 388, 1958: 154–155; Doane et al. 1936; Essig 1926: 519, 1958: 519; Hopkins 1904a: 17; Lindgren 1980a: 68; Pierce, W. D. 1906: 290; Schwarz 1894b: 255, 1905: 185; Schwerdtfeger 1957: 496; Swaine 1918a: 96, 100. **(ds)** Anonymous 1968g; Blackwelder 1947: 782; Bright 1976d: 183, 1981d: 167; Bright & Stark 1973: 113; Chamberlin 1917: 355, 1918a: 16, 1925, 1939: 388, 1958: 154–155; Cockerell et al. 1907; Currie 1905; Essig 1926: 519, 1958: 519; Evans, D. 1983: 35; Evans, D., Lowe, & Hunt 1978; Furniss, M. M., Hungerford, & Wicker 1972; Furniss, R. L. & Carolin 1977: 402; Gemminger & Harold 1872: 2688; Hagedorn 1910d: 73; Hamilton 1894a: 35; Henshaw 1885: 147; Keen 1929a: 33, 1938: 32; Kleine 1912a: 218, 1913b: 140, 1934a: 160; Leng 1920: 341; Ohmart & Voigt 1981: 677, 1982: 340; Schedl 1955g: 18–20, 1963c: 158; Schuder 1969: 76; Schwarz 1894: 255, 1900: 537, 1910: 185; Swaine 1909: 135; Van Dyke 1924: 26; Werner & Holsten 1984; Wickham 1896a: 309; Wood, S. L. 1982b: 1095. **(tx)** Blackman 1928b: 91, 108–109; Bright 1976b: 186, 1981d: 167–168, 351; Chamberlin 1939: 388, 1958: 154–155; Eichhoff 1878b: 173; Evans, D. 1983: 35; Fall 1926: 208; Hagedorn 1910a: 101; Hoebeke 1978; Keen 1929a: 33; LeConte 1857: 22, 1868: 157, 1876: 354; Mannerheim 1843: 298 (reprint p. 126), 1852: 359 (reprint p. 152), 1853: 273; Melsheimer 1853: 87; Swaine 1909: 135, 1918a: 96, 100–101; Wood, S. L. 1969c: 115, 1982b: 1095.
- atratus* LeConte 1868: 156. Lectotype ♂; Cabo de los Reyes, California [USA]; MCZ, Can-

bridge, designated by Bright 1967b: 185. Synonymy: LeConte 1876: 354.

References: **(cn)** Chamberlin 1924; Swaine 1918a: 96, 101. **(hb)** Swaine 1918a: 96, 101. **(ds)** Chamberlin 1925; Henshaw 1882: 268; Hopping 1922; Leng 1920: 341; Swaine 1909: 136. **(tx)** Blackman 1928b: 108; Bright 1976b: 185; Eichhoff 1872b: 137, 1878b: 175; LeConte 1868: 156–157, 1876: 354; Schedl 1955g: 21; Swaine 1909: 136, 1918a: 96, 101.

puncticollis LeConte 1874: 71. Lectotype ♀; Calaveras, California [USA]; MCZ, Cambridge, designated by Bright 1976b: 185. Synonymy: Blackman 1928b: 109.

References: **(cn)** Currie 1905: 72; Essig 1926: 519; Fall & Cockerell 1907: 217; Hopkins 1904a: 16; Keen 1929: 53. **(hb)** Essig 1926: 519; Hopkins 1904a: 16; Keen 1929: 53; Pierce, W. D. 1907: 291. **(ds)** Chamberlin 1917: 355, 1925; Cockerell et al. 1907; Currie 1905; Essig 1926: 519, 1958: 519; Hagedorn 1910d: 75; Henshaw 1882: 268, 1885: 147; Keen 1929: 53; Kleine 1913b: 141, 1914b: 387, 1934a: 161; Leng 1920: 341; Swaine 1909: 139. **(tx)** Blackman 1928b: 108–109; Bright 1976b: 185; Hagedorn 1910a: 101; LeConte 1874a: 70–71, 1876: 354; Schedl 1955g: 21; Swaine 1909: 139, 1918a: 101.

nitidus Swaine 1917: 25. Lectotype ♀; Tullochoram, Quebec [Canada]; CNCI, Ottawa, designated by Bright 1967b: 678.

Figures: Bright 1981d: 346.

Distribution: North America (Alaska/ Alberta, British Columbia, New Brunswick, Newfoundland, Northwest Territories, Nova Scotia, Ontario, Quebec, Yukon in Canada/ Colorado, Maine, Montana, Oregon in USA).

Hosts: *Picea engelmannii*, *P. glauca*, *P. mariana*, *Pinus resinosa*, *P. strobus*, rare in other *Pinus* spp.

References: **(cn)** Lindgren 1980a: 69; Ruppel 1967: 78; Swaine 1918a: 94, 98; Swaine & Craighead 1924: 1–27. **(ec)** Werner & Holsten 1954. **(hb)** Chamberlin 1939: 358; Lindgren 1980a: 69; Schwerdtfeger 1957c: 496; Swaine 1918a: 94, 98. **(ds)** Atkinson & Equihua 1988: 98; Beaulne 1956; Bright 1971a: 127, 1976d: 183, 1981d: 129; Chamberlin 1925, 1939: 358; Elias 1982b, 1982d; Kleine 1934a: 160; Leng 1920: 341; Miller, R. F. & Morgan 1982; Morgan, A. V. & Morgan 1980: 1110; Patterson & Hatch 1945: 153; Ruppel 1967: 78; Schwert et al. 1985; Swaine 1919b: 8; Werner & Holsten 1954; Wood, S. L. 1972a: 426, 1982b: 1052. **(tx)** Blackman 1928b: 30, 35; Bright 1967b: 678, 1977: 516, 1981d: 129, 346; Chamberlin 1939: 358; de Ruette 1970: 110; Swaine 1917: 25–26, 1918a: 94, 98, 1919b: 8, 1925b: 192; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1972a: 426, 1982b: 1052.

borealis Swaine 1925b: 195. Holotype ♀; Copper-

mine River, Arctic Canada; CNCI, Ottawa. Synonymy: Bright 1977: 516.

References: **(cn)** Evans, D. 1982; Lindgren 1980a: 68. **(hb)** Chamberlin 1939: 358; Evans, D. 1982; Lindgren 1980a: 68. **(ds)** Bright 1976d: 181; Chamberlin 1939: 358; Evans, D., Lowe, & Hunt 1978; Fall 1926: 208; Kleine 1934a: 158; Leng & Mutchler 1933: 52; Wood, S. L. 1957c: 401. **(tx)** Blackman 1928b: 30, 35; Bright 1967b: 678, 1977: 516; Chamberlin 1939: 358; Evans, D. 1982; Fall 1926: 208; de Ruette 1970: 108; Swaine 1925b: 195–196; Wood, S. L. 1957c: 401.

anceps Blackman 1928b: 31. Holotype ♀; Clyde, Colorado [USA]; USNM, Washington. Synonymy: Bright 1977: 516.

References: **(hb)** Chamberlin 1939: 357. **(ds)** Chamberlin 1939: 357; Keen 1929a: 32; Kleine 1934a: 158; Leng & Mutchler 1933: 10, 52. **(tx)** Blackman 1928b: 29–32; Bright 1977: 516; Chamberlin 1939: 357; Keen 1929a: 32; de Ruette 1970: 107; Wood, S. L. 1964: 59.

varians Schedl 1930: 196. Holotype ♀; Truro, Nova Scotia [Canada]; CNCI, Ottawa. Synonymy: Wood 1957c: 401.

References: **(hb)** Chamberlin 1939: 369. **(ds)** Beaulne 1956; Chamberlin 1939: 369; Leng & Mutchler 1933: 52. **(tx)** Bright 1977: 516; Chamberlin 1939: 369; de Ruette 1970: 112; Schedl 1930: 196, 1931b: 163, 1979c: 264; Wood, S. L. 1957c: 401.

aquilinus Bright 1968c: 604. Holotype ♀; Rampart House, Yukon Territory [Canada]; CNCI, Ottawa. Synonymy: Bright 1977: 516.

References: **(ds)** Bright 1976d: 180. **(tx)** Bright 1968c: 604, 1977: 516; de Ruette 1970: 108.

nocturnus Schedl 1938a: 185. Lectotype ♂; San Geronimo, Verapaz, Guatemala; BMNH, London, designated by Bright 1976b: 186.

Figures: Blackman 1942a: fig. 45, Bright 1981d: 357.

Distribution: North America (Guatemala/ Honduras/ Chiapas, Hidalgo, Veracruz in Mexico).

Hosts: *Pinus lawsonii*, *P. oocarpa*, *P. pseudostrobus*, *P. strobus* var. *chiapensis*, *P. tenuifolia*.

References: **(cn)** Schwerdtfeger 1956b: 48. **(hb)** Atkinson et al. 1986: 39; Schwerdtfeger 1956b: 48, 1957c: 499. **(ds)** Atkinson & Equihua 1988: 98; Atkinson et al. 1986: 39; Blackwelder 1947: 782; Bright 1981d: 192, 1986: 684; Wood, S. L. 1982b: 1087. **(tx)** Bright 1976b: 186, 1977: 517, 1979c: 171, 1981d: 192, 357; Schedl 1938a: 185–186, 1955g: 23, 1979c: 171; Wood, S. L. 1982b: 1087.

hidalgoensis Blackman 1942a: 215. Holotype ♀; Jacala, Hidalgo, Mexico; USNM, Washington. Synonymy: Bright 1977: 516.

References: **(ds)** Blackwelder 1947: 782. **(tx)** Blackman 1942a: 215; Bright 1977: 516–517.

- novateutonicus** Schedl 1964b: 205. Holotype ♂; Nova Tentonia, S. Catarina, Brasilien; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1964b: 205, 1979c: 172.
- novellus** Blackman 1928b: 96. Holotype ♀; Tehachipi, California [USA]; USNM, Washington. Distribution: North America (California in USA). Hosts: *Pinus sabiniana*.
Notes: (3) Bright 1981d: 328 (the holotype is callos, shriveled, and in poor condition; it may be of *infulatus*). Wood 1982b: 1091 (the allotype and female paratype are of *tuberculatus*; type data of this species and of *infulatus* and its synonyms suggest that *novellus* is synonymous with *infulatus*).
References: (hb) Chamberlin 1939: 385. (ds) Chamberlin 1939: 385; Keen 1929a: 33; Kleine 1934a: 160; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1091. (tx) Blackman 1928b: 96; Bright 1971b: 68, 1981d: 328; Chamberlin 1939: 385; Keen 1929a: 33; Wood, S. L. 1982b: 1091.
- nugalis** Wood 1976a: 356. Holotype ♀; Volcan Pacaya, Esquintla, Guatemala, 1300 m; Wood Collection.
Distribution: North America (Guatemala).
Hosts: Liana.
References: (ds) Bright 1981d: 43; Wood, S. L. 1982b: 1138. (tx) Bright 1981d: 43; Wood, S. L. 1976a: 356, 1982b: 1138.
- obtusipennis** Blandford 1904: 240. Lectotype ♂; Balheu, Verapaz, Guatemala; BMNH, London, designated by Bright 1976b: 184.
Figures: Bright 1981d: 336.
Distribution: North America (Guatemala/ Chiapas, Hidalgo, Mexico, Michoacan in Mexico).
Hosts: *Pinus lawsonii*, *P. leiophylla*, *P. strobus* var. *chiapensis*, *P. sp.*
References: (hb) Atkinson et al. 1986: 39; Burgos & Saucedo 1983: 123. (ds) Atkinson & Equihua 1988: 98; Atkinson et al. 1986: 39; Blackwelder 1947: 782; Bright 1981c: 51; Burgos & Saucedo 1983: 123; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 363; Wood, S. L. 1982b: 1128. (tx) Blandford 1904: 240; Bright 1976b: 184, 1981d: 51, 336; Hagedorn 1910a: 101; Schedl 1955g: 18; Wood, S. L. 1982b: 1128.
- obtusus** Schauffuss 1891: 17. Syntypes ♂ ♀; Madagascar; Hamburg Museum, lost, 1 syntype in Eggers Collection, in NHMW, Wien.
Distribution: Madagascar
Notes: (1) Schedl 1979c: 176 (citation of holotype invalid).
References: (ds) Alluaud 1900: 441; Anonymous 1892: 165; Fairmaire 1892b; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 326. (tx) Fairmaire 1892b; Hagedorn 1910a: 101, 1913b: 255; Schauffuss 1891: 17; Schedl 1938a: 183, 1965c: 67, 1977b: 113, 1979c: 176.
- occidentalis** Blackman 1920b: 4. Lectotype ♂; Pitkin, Colorado [USA]; USNM, Washington, designated by Bright 1976b: 183.
Distribution: North America (Colorado, New Mexico, Utah, S Wyoming in USA).
Hosts: *Picea engelmannii*, *P. pungens*, apparently accidental in *Pinus contorta*, *P. flexilis*.
Notes: (3) Blackman 1928b: 110 (redescribed).
References: (hb) Chamberlin 1939: 389. (ds) Bright 1976d: 183, 1981d: 171; Chamberlin 1925, 1939: 389; Keen 1929a: 33; Kleine 1934a: 160; Leng & Mutchler 1927: 52; Snow 1907: 188; Wickham 1898: 312; Wood, S. L. 1972a: 426, 1982b: 1095. (tx) Blackman 1920b: 4–5, 1928b: 91, 110; Bright 1976b: 183, 1981d: 171; Chamberlin 1939: 389; Keen 1929a: 33; McNamara 1984: 757; Wood, S. L. 1972a: 426, 1982b: 1095.
- occlusus** Bright 1976c: 437. Holotype ♀; Yucaran, Paraiso, Honduras; Wood Collection.
Figures: Bright 1981d: 336.
Distribution: North America (Honduras/ Nayarit, Oaxaca in Mexico).
Hosts: *Pinus caribaea*, *P. oocarpa*.
References: (ds) Atkinson & Equihua 1985c: 359; Bright 1981d: 53; Wood, S. L. 1982b: 1128. (tx) Bright 1976c: 437, 1981d: 53, 336; McNamara 1984: 757; Wood, S. L. 1982b: 1128.
- olivierai** Schedl 1972g: 68. Holotype, sex?; Brasilien, Pedra Azul, Minas Gerais; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 68, 1979c: 178.
- opaculus** LeConte 1878b: 623. Holotype ♀; Marquette, Michigan [USA]; MCZ, Cambridge.
Figures: Bright 1981d: 360.
Distribution: North America (Alaska/ all provinces in Canada/ Arizona, California, Colorado, Idaho, Maine, Michigan, Minnesota, Nevada, New Hampshire, New Mexico, New York, Pennsylvania, South Dakota, Utah, West Virginia, Wyoming in USA).
Hosts: *Abies balsamea*, *A. concolor*, *A. lasiocarpa*, *Larix laricina*, *Picea engelmannii*, *P. glauca*, *P. pungens*, *Pinus sylvestris*, *Pseudotsuga menziesii*.
References: (bv) Hosking & Knight 1975. (cn) Blackman 1950; Chamberlin 1924; Doane et al. 1936; James & Linnane 1979; Schuder 1969: 80; Stevens, Lister, & Linnane 1979; Swaine 1918a: 95, 99; Swaine & Craighead 1924: 1–27. (cc) Werner & Holsten 1984. (hb) Baker, W. L. 1972: 255; Blackman 1950; Chamberlin 1939: 366; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Swaine 1918a: 95, 99. (ds) Anonymous 1926c: 519; Ashworth 1977; Blackman 1950; Blatchley & Leng 1916: 633; Bright 1976d: 187, 1981d: 218; Chamberlin 1925, 1939: 366; Deyrup 1981: 8; Deyrup & Atkinson 1987a: 66; Drooz 1985: 363; Elias 1982b, 1982c, 1982d; Hagedorn 1910d: 73; Henshaw 1882: 268, 1885: 147; Hubbard &

- Schwarz 1878a: 665, 1878b: 623; Kleine 1913b: 140, 1934a: 160; Leng 1920: 341; Leonard 1928: 519; Miller, R. F. & Morgan 1952; Morgan, A. V. & Morgan 1950: 1110; Schuder 1969: 80; Swaine 1909: 136; Werner & Holsten 1954; Wood, S. L. 1957c: 401, 1952b: 1037. **(tx)** Blackman 1922b, 1928b: 44, 52; Blatchley & Leng 1916: 633; Bright 1977: 517, 1981d: 218, 360; Chamberlin 1939: 366; Hagedorn 1910a: 101; LeConte 1878b: 623; Schedl 1931b: 165; Swaine 1909: 136, 1918a: 95, 99; Wood, S. L. 1957c: 401, 1952b: 1037.
- abietis* Blackman 1928b: 49. Holotype ♀; Silver Lake, Utah [USA]; USNM, Washington. Synonymy: Bright 1977: 517.
References: **(hb)** Bright & Stark 1973: 107; Chamberlin 1939: 365. **(ds)** Bright & Stark 1973: 107; Chamberlin 1939: 365; Keen 1929a: 32; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 49–50; Bright 1977: 517; Chamberlin 1939: 365; Keen 1929a: 32; Michalski 1969b: 569; de Ruelle 1970: 107.
- albertensis* Blackman 1928b: 50. Holotype ♀; Banff Sp. [Banff Springs], Alberta, Canada; USNM, Washington. Synonymy: Bright 1977: 517.
References: **(cn)** Lindgren 1980a: 67. **(hb)** Chamberlin 1939: 365, 1958: 149; Lindgren 1980a: 67. **(ds)** Bright 1976d: 180; Chamberlin 1939: 365, 1958: 149; Leng & Mutchler 1933: 52; Smith, G. S. 1930; Wood, S. L. 1972a: 425. **(tx)** Blackman 1928b: 44, 50–51; Bright 1977: 517; Chamberlin 1939: 365, 1958: 149; de Ruelle 1970: 107; Wood, S. L. 1972a: 425.
- exiguus* Blackman 1928b: 51. Holotype ♀; Ft. Garland, Colorado [USA]; USNM, Washington. Synonymy: Bright 1977: 517.
References: **(hb)** Chamberlin 1939: 366. **(ds)** Chamberlin 1939: 366; Keen 1929a: 32; Kleine 1934a: 159; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 51; Bright 1966b: 298, 1977: 517; Chamberlin 1939: 366; Keen 1929a: 32; de Ruelle 1970: 109.
- pygmaeus* Schedl 1931b: 165. Holotype ♀; Ste. Anne's, Quebec [Canada]; CNCI, Ottawa. Synonymy: Wood 1957c: 401.
References: **(hb)** Chamberlin 1939: 366. **(ds)** Beaulne 1956; Chamberlin 1939: 366; Leng & Mutchler 1933: 95. **(tx)** Bright 1977: 517; Chamberlin 1939: 366; de Ruelle 1970: 111; Schedl 1931b: 165–166; Wood, S. L. 1957c: 401.
- orarius* Bright 1968c: 607. Holotype ♀; Colwood, Esquimalt District, British Columbia [Canada]; CNCI, Ottawa.
Figures: Ruth 1980: 15.
Distribution: North America (British Columbia in Canada).
Hosts: *Pseudotsuga menziesii*.
- References: **(cn)** Hedlin 1974: 53; Meso 1979: 247; Ruth 1980: 15. **(hb)** Bright 1981d: 255; Hedlin 1974: 53; Hedlin & Ruth 1970: 105–108; Meso 1979: 247; Ruth 1980: 15. **(ds)** Bright 1976d: 184, 1981d: 255; Furniss, R. L. & Carolin 1977: 402; Hedlin & Ruth 1970; Wood, S. L. 1982b: 1061. **(tx)** Bright 1968c: 607–608; 1981d: 255; de Ruelle 1970: 110; Ruth 1980: 15; Wood, S. L. 1982b: 1061.
- ornatus* Blackman 1928b: 102. Holotype ♀; Manitou, Colorado [USA]; USNM, Washington.
Distribution: North America (Colorado, Utah, Wyoming in USA).
Hosts: *Picea pungens*, rare or accidental in *Pinus* spp.
Notes: (3) Bright 1981d: 147 (treated as a synonym of *intextus*).
References: **(hb)** Chamberlin 1939: 387. **(ds)** Chamberlin 1939: 387; Keen 1929a: 33; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Wood, S. L. 1952b: 1090. **(tx)** Blackman 1928b: 102; Chamberlin 1939: 387; Keen 1929a: 33; McNamara 1954: 757; Wood, S. L. 1977d: 516, 1982b: 1090.
- kenti* Blackman 1928b: 141. Holotype ♀; Black Hills, Wyoming [USA]; USNM, Washington. Synonymy: Wood 1977b: 210.
References: **(hb)** Chamberlin 1939: 401. **(ds)** Chamberlin 1939: 401; Keen 1929a: 34; Kleine 1934a: 160; Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 141; Bright 1977: 515; Chamberlin 1939: 401; Keen 1929a: 34; McNamara 1954: 756; Wood, S. L. 1977b: 210.
- linatus* Wood 1964: 65. Holotype ♀; Sanford Canyon, Dixie National Forest, Utah [USA]; Wood Collection. Synonymy: Wood 1977d: 516.
References: **(tx)** McNamara 1977: 198; Wood, S. L. 1964: 65, 1977d: 516.
- ostriacolens* Bright 1986b: 681. Holotype ♀; Mexico: Cuernavaca, Morelos; CNCI, Ottawa.
Distribution: North America (Morelos in Mexico).
Hosts: *Ostrya virginiana*.
References: **(tx)** Bright 1986b: 681.
- pampasae* Schedl 1970e: 92. Holotype, sex?; Pampas (wohl Argentinien); Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
References: **(tx)** Schedl 1970e: 92, 1979c: 183.
- parfentieri* Pjatkitskii 1931: 169. Syntypes, sex?; Kirgisien, USSR: Pjatkitskii Collection.
Distribution: Europe (W USSR).
Hosts: *Picea schrenkiana*.
References: **(hb)** Stark 1952: 345. **(ds)** Kadyrov 1989; Kleine 1934a: 161; Marikovskii 1956: 73; Parfentev 1951: 429–430; Pfeffer 1976: 340; Stark 1952: 345. **(tx)** Pfeffer 1940b: 108, 111, 117, 1976: 340; Pjatkitskii 1931: 169–171; Schedl 1934f: 1643, 1979c: 185; Sokanovskii 1954: 19; Stark 1952: 345.
- parilis* Wood 1976a: 359. Holotype ♀; Buenos Aires, Cortez, Honduras, 2300 m; Wood Collection.

Distribution: North America (Honduras).

Hosts: *Quercus* sp. branch.

References: (ds) Bright 1981d: 103; Wood, S. L. 1982b: 1082. (tx) Bright 1981d: 103; Wood, S. L. 1976a: 359, 1982b: 1082.

paulus Wood 1964: 63. Holotype ♀; 40 km NE Jacala, Hidalgo, Mexico; Wood Collection.

Distribution: North America (Chiapas, Chihuahua, Hidalgo, Michoacan, San Luis Potosi, Tlaxcala, Veracruz in Mexico).

Hosts: *Senecio* sp.

Notes: (3) The host recorded as *Baccharis* sp. was actually *Senecio* (SLW).

References: (hb) Atkinson & Equihua 1985a: 91, 1985c: 359; Atkinson et al. 1986: 40, 72; Burgos & Saucedo 1983: 124. (ds) Atkinson & Equihua 1985a: 91, 1985c: 359; Atkinson et al. 1986: 40, 72; Bright 1981d: 63; Burgos & Saucedo 1983: 124; Wood, S. L. 1982b: 1130. (tx) Bright 1981d: 63; McNamara 1977: 198; Wood, S. L. 1964: 63, 1982b: 1130.

pellitus Schedl 1955g: 23. Lectotype ♀; Quezaltenango, Guatemala; Schedl Collection in NHMW, Wien, designated by Bright 1976b: 186. Figures: Bright 1981d: 360.

Distribution: North America (Guatemala/ Chiapas, Hidalgo, Oaxaca, Puebla, Queretaro in Mexico).

Hosts: *Pinus greggii*, *P. lawsonii*, *P. michoacana*, *P. montezumae*, *P. oocarpa*, *P. pseudostrobus*, *P. rudis*, *P. tenuifolia*, *P. spp.*

References: (hb) Bright 1981d: 216; Schwerdtfeger 1957c: 500. (ds) Bright 1981d: 216; Wood, S. L. 1982b: 1138. (tx) Bright 1976b: 186, 1981d: 216, 360; Schedl 1955g: 18, 23, 1979c: 187; Wood, S. L. 1982b: 1038.

pentaclethrae Schedl 1957d: 62. Holotype ♂²; Congo Belge: Luki, Mayumbe; MRCB, Tervuren. Distribution: Africa (Zaire).

Hosts: *Pentaclethra macrophylla*.

References: (cc) Schedl 1958d: 188, 1962j: 18. (hb) Schedl 1962j: 18. (ds) Schedl 1962j: 18. (tx) Schedl 1957d: 62, 1962j: 18, 1979c: 187–188.

peregrinus Eichhoff 1878b: 193. Holotype, sex²; America meridionalis (Brasilica); Hamburg Museum, lost.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 782; Gonçalves 1935: 414–420; Hagedorn 1910d: 73; Kleine 1913b: 140, 1914b: 336. (tx) Eichhoff 1878a: 389, 1878b: 193; Hagedorn 1910a: 101.

perexiguus Wood 1976a: 355. Holotype ♀; Dominical, Puntarenas, Costa Rica, 3 m; Wood Collection. Distribution: North America (Costa Rica/ Panama).

Hosts: Tree limb.

References: (ds) Bright 1981d: 67; Wood, S. L. 1982b: 1138. (tx) Bright 1981d: 67; McNamara 1984: 757; Wood, S. L. 1976a: 355, 1982b: 1138.

perotei Blackman 1942a: 218. Holotype ♀; Perote, Veracruz, Mexico; USNM, Washington.

Distribution: North America (Mexico, Morelos, Puebla, Veracruz in Mexico).

Hosts: *Pinus leiophylla*, *P. teocote*.

References: (hb) Atkinson et al. 1986: 40; Burgos & Saucedo 1983: 121. (ds) Atkinson & Equihua 1985a: 97; Atkinson et al. 1986: 40; Blackwelder 1947: 782; Bright 1981c: 282; Burgos & Saucedo 1983: 121; Thomas, J. B. 1966; Wood, S. L. 1982b: 1098. (tx) Blackman 1942a: 218; Bright 1981d: 282; McNamara 1984: 757; Wood, S. L. 1982b: 1098.

philippinensis Schedl 1936h: 59. Holotype ♀; Luzon, Manila; Schedl Collection in NHMW, Wien. Distribution: Philippine Islands (Luzon).

References: (ds) Schedl 1966b: 41. (tx) Schedl 1936h: 59, 1938a: 183, 1979c: 192.

pinavorus Bright 1985c: 182. Holotype ♀; Florida: Highlands Co., Lake Placid, 3 miles [5 km] E Archbold Biological Station [USA]; CNCI, Ottawa.

Distribution: Antilles Islands (Dominican Republic in Hispanola), North America (Florida in USA).

Hosts: *Pinus elliottii*, *P. occidentalis*.

References: (cc) Haack et al. 1989. (bv) Haack et al. 1989. (hb) Haack et al. 1989. (ds) Bright 1985c: 176. (tx) Bright 1985c: 176, 182.

pinguis (Blackman) 1928b: 20 (*Myeloborus*). Holotype ♀; Longmont, Colorado [USA]; USNM, Washington.

Distribution: North America (Arizona, Colorado, Nevada, New Mexico, Utah, Wyoming in USA).

Hosts: *Pinus flexilis*, *P. strobiformis*.

Notes: (1) Blackman 1928: 20 (*pinquis*, lapsus calami; name correctly spelled on p. 17, 149, 153, and fig. 60 as *pinguis*).

References: (hb) Chamberlin 1939: 340. (ds) Chamberlin 1939: 340; Keen 1929a: 31; Kleine 1934a: 163; Leng & Mutchler 1933: 53; Wood, S. L. 1972a: 424. (tx) Blackman 1928b: 20; Bright 1976c: 427, 1981d: 210; Chamberlin 1939: 340; Keen 1929a: 31; de Ruelle 1970: 104; Wood, S. L. 1951a: 128, 1972a: 424.

pini Kurenzov 1941a: 176. Lectotype, sex²; Ternei region, Suputinia River, Ussuri, USSR; Kurenzov Collection in ISBN, Novosibirsk, designated by Michalski 1969a: 895.

Figures: Yin, Huang, & Li 1984: 120 (elytral declivity).

Distribution: Asia (China/ E USSR).

Hosts: *Picea koraiensis*, *P. obovata*, *Pinus koraiensis*.

References: (hb) Krivolutsкая 1956: 838, 1960: 78; Kurenzov 1948b: 108, 1950d: 228; Stark 1952: 357. (ds) Krivolutsкая 1956: 838, 1960: 78, 1983; Pfeiffer 1976: 340; Stark 1952: 357; Zinovjev 1955: 187, 190. (tx) Krivolutsкая 1956: 836, 838; Kurenzov 1941a: 176, 233, 1948b: 108; Michalski

1969a: 595, 1969b: 569; Pfeffer 1976: 340; Stark 1952: 357; Yin, Huang, & Li 1954: 120.

pinsapo Pfeffer 1982b: 154. Holotype, sex?, Spain; apparently in Prague Museum.

Distribution: Europe (Spain).

Hosts: *Abies pinsapo*.

References: (tx) Pfeffer 1982b: 154–157.

pityographus (Ratzeburg) 1837: 162 (*Bostrichus*).
Synotypes, sex?: Europe; not located.

Figures: Grune 1979: 114, Joly 1976b: fig. 143, Pfeffer 1989a: pl. 9, Tsai & Li 1959: 94.

Distribution: Asia (Turkey), Europe (Austria/ Belgium/ Bulgaria/ Czechoslovakia/ Denmark/ France/ Germany/ Greece/ Hungary/ Italy/ Poland/ Romania/ Crimea, Ukraine in W USSR/ Yugoslavia).

Hosts: *Picea excelsa*, *P. omorica*, *P. orientalis*, *Abies alba*, *A. borisi*, *A. bornmulleriana*, *A. cephalonica*, *A. pectinata*, *Pinus heldreichii*, *P. peuce*.

References: (ay) Chararas 1977b, (bv) Chararas 1966b, 1977b; Francke et al. 1957; Grune 1979: 115; Hellrigl 1985; Jacobson 1972; Krol & Zabecki 1976; Mori & Pnapoomchareon 1957; Prell 1930c: 631; Tragardh 1930b: 102; Zimr & Landa 1985. (cn) Capecki 1982; Capek et al. 1957; Chararas 1961b: 70, 1961c: 93, 1978; Chorbadzhievo 1929; Gradojevic 1933; Grandi 1951; Hess 1907: 245; Hrubick 1973; Joly 1976; Kailidis & Georgevits 1972; Komarek 1931: 1–256; Konig, E. 1971; Krol 1980a; Krol & Zabecki 1976; Lezhava 1930: 1–15; Lozovoi & Tropin 1965, 1967; Martinek 1953a: 373; Nosek 1951: 106, 1952b: 98, 1959b: 73; Oppermann 1965; Pfeffer 1947c: 204, 1949b: 147, 1950c: 3; Schimitschek 1937c: 55, 1952c: 59; Szujecki 1956: 241–242; Vite 1984b; Wichmann 1927b: 353; Zimr & Landa 1985. (cc) Balazy 1962, 1968; Balazy & Michalski 19641, 1964b; Balazy et al. 1977; Capecki 1982; Chararas 1959d; Cadek 1976; Hellrigl 1985; Heuer & Vite 1984b; Hirschmann & Wisniewski 1982, 1983; Karpinski 1932a: 95, 99, 101; Kielczewski 1976; Kielczewski & Balazy 1966; Kielczewski, Bohdan, & Balazy 1967: 161–163; Kielczewski, Moser, & Wisniewski 1983; Kielczewski & Wisniewski 1978, 1983; Krol 1980a; Krol & Zabecki 1976; Kudela 1981; Lozovoi & Tropin 1965, 1967; Masutti 1959: 268, 299; Michalski & Ratajczak 1989; Nosek 1951: 106, 1952b: 98, 1959b: 87; Numberg 1930: 201; Numberg & Wiackowski 1958: 130; Pfeffer 1932a: 8, 1943b: 182, 1947c: 204, 1949b: 147, 1950c: 3, 1955b: 84, 1957a: 198, 1959: 5, 1960: 346; Ratzeburg 1869a: 79; Rondani 1873: 140; Schimitschek 1941a: 312; Zimr 1985b. (hb) Balachowsky 1963a: 1254; Balazy 1968; Budge 1949; Chararas 1961b: 70, 1961c: 93, 1962c: 363, 1977b; Chorbadzhievo 1929; Grandi 1951; Henschel 1876a: 99, 240; Hess 1907: 245; Joly 1976; Karpinski & Strawinski 1948: 156; Konig, E. 1971; Kudela 1981; Lozovoi & Tropin 1965, 1967; Masutti 1959: 268,

299, 1964; Michalski 1959a: 291; Nordlinger 1856: 27; Nosek 1959b: 87; Numberg 1929a: 111, 1929c: 121, 1947c: 106; Pfeffer 1941b: 1, 1941c: 5, 1989a: 63; Postner 1974: 434; Prell 1930c: 631; Ratzeburg 1837: 136, 1839: 166; Schwerdtfeger 1981: 192; Simmel 1928: 154; Spessivtsev 1923: 205; Stark 1952: 344; Szujecki 1955: 241; Tragardh 1930b: 102; Tschorbadjiev 1929: 169; Wichmann 1927b: 353. (ds) Andersch 1851; Arnoldi et al. 1955: 713; Arru, Covassi, & de Bellis 1966: 32; Benz 1985; Bielz 1851; Borchert 1951; Brakman 1966b: 206; Buresh & Lazarov 1956; Capek et al. 1957; Chapuis & Candeze 1853; Chararas 1962c: 363; Chorbadzhievo 1929; Endrodi 1958b; Escherich 1932b; Forster 1849: 439; Gaubil 1849: 126; Gozis 1875: 80; Gradojevic 1933: 789–790; Gredler 1868: 374; Grune 1979: 115; Hansen, V. 1956; Horion 1951; Jacentkovsky 1939: 76; Jammicky 1960a; Joly 1976; Kailidis & Georgevits 1972; Kaltenbach 1874: 685; Karpinski 1931: 19, 1932a: 95, 99, 101, 1933a: 290, 1948b: 231; Karpinski & Strawinski 1948: 156; Kiefer et al. 1942: 528; Kraatz 1869: 59; Kudela 1946a: 339; Leclercq 1971; Lindemann 1884b: 264; Lucht 1987: 278; Mequignon 1936: 54; Michalski 1957: 165; Negru 1966b: 403; Numberg 1928b: 88, 111, 1938: 126, 1954: 70; Perris 1876a: 255, 1877a: 255; Pfeffer 1931b: 74, 1935: 158, 1947d: 129, 1947e: 2, 1950b: 73, 1960: 346, 1976: 339, 1989a: 63; Pittioni 1943: 175; Pjatnitskii 1930a: 165, 1931: 171–173; Postner 1974: 434; Ratzeburg 1837: 136, 1839: 166; Redtenbacher 1858: 830; Ronbal 1935b: 72, 1936b: 193, 1941: 269; Sainte-Claire & Mequignon 1938: 448; Schaum 1859: 95, 1862: 101; Schedl 1959h: 100, 1961b: 184, 1967c: 71, 1980a: 23, 1981b: 77; Schilsky 1909: 188; Schwerdtfeger 1981: 192; Seidl 1876: 4; Stark 1952: 344; Stein 1868: 114; Stein & Weise 1877: 164; Stolima 1969: 610–627; Strand 1964: 243; Sturm 1843: 230; Szujecki 1955: 241; Tschorbadjiev 1929: 159; Wachtl 1870: 239; West 1935: 184; Wichmann 1927a: 73–74; Zivojinovic 1960: 25. (tx) Bach 1854; Balachowsky 1949a: 235; Chapuis & Candeze 1853; Eggers 1940g: 36; Endrodi 1957a: 307, 1957b; Grune 1979: 114–115; Hansen, V. 1956; Hellrigl 1985; Henschel 1876a: 99, 240; Joly 1976, 1976b: fig. 143; Karpinski 1949: 129; Karpinski & Strawinski 1948: 156; Letzner 1891: 376; Lucht 1987: 278; Negru 1966b: 403; Nordlinger 1848: 242, 1856: 27; Numberg 1929c: 121, 1930, 1938: 126, 1954: 70; Perris 1877a: 255; Pfeffer 1930: 111, 116, 1932b: 22, 1940b: 107, 114, 116, 1941b: 1, 1941c: 5, 1947e: 2, 1976: 339, 1989a: pl. 9; Pjatnitskii 1931: 171; Postner 1974: 434; Ratzeburg 1837: 136, 162, 1839: 166; Redtenbacher 1849: 26, 851, 1858: 830; Schedl 1934f: 1643, 1952f: 87, 1958k: 144, 1959h: 100, 1980a: 23, 1981b: 77; Schimitschek 1937c: 55; Schmidt, G. 1980: 17; Spessivtsev 1923: 205–206, 1926b: 50, 1929a: 678–682,

1929c: 297; Stark 1952: 344; Trappen 1935; Tsai & Li 1959: 94. (**ms**) Chararas 1959d; Escherich 1932b; Michalski 1959a: 291; Mori & Puapoomchareon 1987.

pityographus cribratus Pfeffer 1940: 110. Holotype, sex?, Mt. Chelmon, Greece; Pfeffer Collection in Prague Museum.

Notes: (3) Pfeffer 1976: 335 (cited as a subspecies).

References: (**ds**) Pfeffer 1976: 340, 1989a: 64. (**tx**) Pfeffer 1940: 107, 110, 1976: 340.

pityographus bibractensis Balachowsky 1949a: 236. Holotype ♂; Forêt de St.-Prix (Morvan), France, 800 m; MNHN, Paris. Synonymy: Schedl 1958k: 144.

References: (**ds**) Larroche & Torossian 1971. (**tx**) Balachowsky 1949a: 236; Schedl 1958k: 144.

pityographus maritimus Stark 1952: 344. Holotype, sex?, Caucasus; IZL, Leningrad.

References: (**tx**) Michalski 1969b: 569; Stark 1952: 344.

plaumami (Schedl) 1970e: 102 (*Breviophthorus*). Holotype, sex?, Brasilien, Bocaiva (Pa), 25 08, 49 04, 1000 m; Schedl Collection in NIMW, Wien. Distribution: South America (Brazil).

References: (**tx**) Schedl 1970e: 102, 1979c: 196.

pseudotsugae Swaine 1918a: 99. Holotype ♀; BX Mountain, Vernon District, British Columbia [Canada]; CNCI, Ottawa.

Figures: Bright 1981d: 350.

Distribution: North America (British Columbia in Canada/ Arizona, California, Colorado, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming in USA).

Hosts: *Abies concolor*, *A. grandis*, *A. lasiocarpa*, *A. magnifica*, *Picea engelmannii*, *Pinus lambertiana*, *Pseudotsuga menziesii*, *Tsuga heterophylla*.

Notes: (3) Blackman 1928b: 106 (redescribed).

References: (**cn**) Anonymous 1963k, 1963y; Browne 1968b: 564; Chamberlin 1924; Doane et al. 1936; Hofacker et al. 1989; Ruppel 1967: 78; Schuder 1969: 77; Stark & Borden 1965: 1162–1163; Struble 1937d: 11, 1957: 14; Swaine 1918a: 96, 99. (**ec**) Ashraf & Berryman 1969: 19; Berryman 1968a: 66; Bushing & Bright 1965: 203; Dewey & Clinton 1980; Deyrup & Gara 1978: 275; Struble 1957: 14. (**hb**) Bright & Stark 1973: 114; Browne 1968b: 564; Chamberlin 1939: 387, 1958: 154; Doane et al. 1936; Furniss 1968: 1384–1389; Struble 1937d: 11, 1957: 14; Swaine 1918a: 96, 99. (**ds**) Anonymous 1963k, 1963y; Bedard 1938a; Bright 1976d: 184, 1981d: 162; Bright & Stark 1973: 114; Browne 1968b: 564; Chamberlin 1925, 1939: 387, 1958: 154; Downing 1963: 8; Evans, D. 1983: 35; Furniss, R. L. & Carolin 1977: 402; Cast et al. 1989: 385; Hopping 1922; Keen 1929a: 33; Kleine 1934a: 161; Leng 1920: 341; McComb et al. 1953: 2; Patterson & Hatch 1945:

153; Ruppel 1967: 78; Schuder 1969: 77; Thatcher 1935: 261; Wood, S. L. 1948: 53, 1951a: 128, 1972a: 426, 1982b: 1093. (**tx**) Blackman 1928b: 91, 106–107; Bright 1967b: 678, 1981d: 162, 350; Chamberlin 1939: 387, 1958: 154; Evans, D. 1983: 35; Hoebeke 1978; Keen 1929a: 33, 1938: 32; de Ruelle 1970: 110; Swaine 1918a: 96, 99; Wood, S. L. 1951a: 128, 1972a: 426, 1977d: 516, 1982b: 1093.

thatcheri Bright 1976c: 442. Holotype ♀; Big Sandy Meadow, S28, T5S, R2E, California [USA]; Wood Collection. Synonymy: Wood 1977d: 516.

References: (**tx**) Bright 1976c: 442; McNamara 1984: 758; Wood, S. L. 1977d: 516.

puberulus (LeConte) 1868: 157 (*Cryphalus*). Holotype, sex?, District of Columbia [USA]; MCZ, Cambridge.

Figures: Bright 1981d: 366.

Distribution: North America (New Brunswick, Nova Scotia, Ontario, Quebec in Canada/ District of Columbia, Indiana, Kansas, Kentucky, Maine, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Abies balsamea*, *Picea* spp., *Pinus banksiana*, *P. resinosa*, *P. strobus*, *P. sylvestris* twigs.

Notes: (1) LeConte 1876: 354 (to *Pityophthorus*). (3) Blackman 1928b: 48 (redescribed).

References: (**ay**) Thomas, J. B. 1957: 4, 1967. (**bv**) Kirkendall 1984: 241. (**cn**) Anonymous 1965b; Blackman 1950; Browne 1968b: 564; Chamberlin 1924; Doane et al. 1936; Felt 1906: 752; Hopkins 1899c: 443; Lanier, Abrahamson, & Schoeneck 1983; Lindquist, O. H. & Syme 1981: 88; Packard 1890: 718; Pettit 1928: 1–20; Schuder 1969: 80; Swaine 1918a: 95, 99; Syme & Nystrom 1988: 84. (**cc**) Bushing 1965: 464; Felt 1906: 752; Heqvist 1959: 181; Lanier, Abrahamson, & Schoeneck 1983; Reid 1957b: 7. (**hb**) Baker, W. L. 1972: 255; Blackman 1950; Browne 1968b: 564; Chamberlin 1939: 365; Chittenden 1890; Deyrup & Atkinson 1987a: 66; Deyrup & Kirkendall 1983; Doane et al. 1936; Felt 1906: 752; Hopkins 1899c: 443; Packard 1881: 170–175, 1890: 718, 812; Pierce, W. D. 1907: 291; Schwarz 1885a: 80; Swaine 1918a: 95, 99. (**ds**) Anonymous 1926c: 519, 1965b; Baker, W. L. 1972: 255; Blackman 1919b: 140–142, 1950; Blatchley & Leng 1916: 629; Bright 1971a: 127, 1976d: 187, 1981d: 264; Britton 1920a; Browne 1968b: 564; Chamberlin 1925, 1939: 365; Chittenden 1890; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Dodge 1938; Drooz 1985: 364; Gemminger & Harold 1892: 2688; Hagedorn 1910d: 73; Hamilton 1895a: 378; Hatch 1924; Henshaw 1882: 268, 1885: 147; Hopkins 1893a: 131, 1893b: 209; Hubbard & Schwarz 1878a: 665; Kleine 1913b: 140, 1914b: 391, 1934a:

- 161; Knull 1932: 1196–1203, 1934: 212; Leng 1920: 341; Leonard 1928: 519; Lindquist, O. H. & Syme 1981: 88; Morgan, A. V. & Morgan 1979; Schuder 1969: 80; Schwarz 1888a: 50, 1891: 65; Smith, J. B. 1890: 267, 1900: 362, 1910: 401; Swaine 1909: 137; Syme & Nystrom 1988: 84; Ulke 1902: 56; Wood, S. L. 1982b: 1034. **(tx)** Beal & Massey 1945: 136–137; Blackman 1919b: 140–142, 1922c, 1928b: 44, 48–49; Blatchley & Leng 1916: 629; Bright 1981d: 264, 366; Chamberlin 1939: 365; Dodge 1938: 15; Eichhoff 1878b: 202, 1896: 609–610; Hagedorn 1910a: 101; LeConte 1868: 157, 1876: 354, 1878: 665; Lindquist, O. H. & Syme 1981: 88; Pfeffer 1932b: 23; Swaine 1909: 137, 1918a: 95, 99; Syme & Nystrom 1988: 84; Thomas, J. B. 1957: 4, 1967; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1982b: 1034.
- infans* Eichhoff 1872a: 135. Holotype, sex?, Amer. Bor. (Civitat. unitt.); Hamburg Museum, lost, neotype, ♀, Syracuse, New York [USA], USNM, Washington, designated by Bright 1978: 72.
- Notes: (3) Blackman 1928b: 49 (redescribed). References: **(ds)** Gemminger & Harold 1872: 2688; Leng 1920: 341. **(tx)** Blackman 1928b: 49; Bright 1978: 72; Eichhoff 1872a: 135, 1878b: 187, 1896: 609–610; LeConte 1876: 355; Swaine 1909: 137.
- pubescens* (Marshall) 1802: 58 (*Ips*). Syntypes, sex?, not given, presumably England; not located, presumably in BMNH, London.
- Figures: Pfeffer 1989a: pl. 9.
- Distribution: Africa (Algeria/ Egypt/ Libya/ Morocco/ Tunisia), Asia (Cyprus/ Israel), Europe (Austria/ Belgium/ Corsica/ Czechoslovakia/ Denmark/ England/ France/ Germany/ Greece/ Hungary/ Italy/ Netherlands/ Norway/ Sweden/ Switzerland).
- Hosts: *Abies cephalonica*, *Pinus halepensis*, *P. maritima*, *P. nigra*, *P. pinaster*, *P. pinca*, *P. radiata*, *P. sylvestris*, *P. uncinata*.
- References: **(ay)** Feytaud 1950. **(bv)** Grune 1979: 117; Wichmann, H. E. 1967. **(cn)** Barbey 1925: 280; Browne 1968b: 565; Feytaud 1946, 1950a; Forbes 1910; Grandi 1951; Hanson 1937, 1940a; Joly 1976; Kailidis & Georgevits 1972. **(ec)** Elliot & Morley 1907; Halperin & Holzschuh 1984: 28; Heqvist 1963: 154; Kleine 1908c: 213, 1909a: 47, 78; Kraemer 1950b: 380; Nosek 1959a: 118; Pfeffer 1960: 346; Thompson, W. R. 1943: 91; Wichmann, H. E. 1967. **(hb)** Barbey 1925: 280; Browne 1968b: 565; Chararas 1962c: 363; Dallimore & Munro 1922; Feytaud 1946, 1950a; Gillanders 1908; Grandi 1951; Gyorfi 1957; Halperin & Holzschuh 1984: 28; Joly 1976; Kemner 1919: 175; Kraemer 1950b: 380; Lengerken 1939: 65, 1954: 85; Masutti 1964; Munro 1926: 63; Nosek 1959a: 118; Pfeffer 1941b: 3; Postner 1974: 435; Spessivtsev 1913a: 89, 1921b: 221; Stark 1952: 352; Zocchi 1959: 105. **(ds)** Audras & Schaefer 1957; Bangsholt 1975: 95; Beaulne 1956; Bejer-Petersen & Jorum 1977: 26; Blandford 1891c; Brakman 1966b: 206; Browne 1968b: 565; Butovitsch 1963; Chararas 1962c: 363; Chrystal 1937; Dallimore & Munro 1922: 189–193; Eggers 1918; Endrodi 1958a, 1958b; Escherich 1932b; Everts 1925; Fowler 1891; Georghion 1977: 75; Grune 1979: 117; Hagedorn 1910d: 75; Halperin & Holzschuh 1984: 28; Hanan 1919; Hansen, V. 1939, 1956, 1964: 464; Hellen 1947; Horion 1951; Jansson 1936: 220; Johnson & Halbert 1902: 819; Joly 1976; Kailidis & Georgevits 1972; Karpinski 1949: 125; Kemner 1919: 175; Klefbeck & Sjöberg 1960: 230; Kleine 1912a: 267, 1913a: 35, 1913b: 141, 1934a: 161; Kloft & Hinks 1945: 218; Leclercq 1971; Lucht 1987: 278; Mahler 1987: 232; Munro 1920: 1–35, 1921: 88; Pfeffer 1930b: 120, 1931b: 74, 1935: 158, 1947d: 126–127, 1947e: 12, 1960: 346, 1976: 340, 1984: 277, 1989a: 66; Pittioni 1943: 175; Postner 1974: 435; Reitter 1916: 298; Rouhal 1941: 270; Sainte-Claire & Mequignon 1938: 447; Schaufuss 1915: 1243; Schedl 1964a, 1976c: 71, 1978e: 37, 1981b: 78; Schilsky 1909: 188; Stark 1927b: 90, 1952: 352; Stephens 1829a: 145, 1830: 357, 1839: 207; Tredl 1907: 14; Winter, T. G. 1983: 22; Zocchi 1959: 105. **(tx)** Balachowsky 1949a: 237; Blandford 1891c: 15–17; Duffy 1953; Eggers 1915b: 13–14, 1918: 181; Endrodi 1957b; Fleischer 1927; Gillanders 1908; Grune 1979: 117; Hagedorn 1910a: 101; Hansen, V. 1956, 1964: 464; Joly 1976; Karpinski 1949: 125; Lucht 1987: 278; Marshall 1802: 58; Olivier 1795b: 11; Pfeffer 1933: 3–54, 1940b: 108, 112, 119, 1941b: 3, 1947e: 12, 1955a: 217, 1976: 340, 1989a: pl. 9; Portevin 1935: 333; Postner 1974: 435; Reitter 1913a: 93, 1916: 298; Schedl 1934f: 1643, 1967f: 149, 1980a: 23, 1981b: 78; Sokanovskii 1954: 19; Spessivtsev 1913a: 89, 1922a: 488, 1922b: 221, 1925a: 182, 1926b: 50, 1929c: 298–303, 1931: 63; Stark 1952: 352; Stephens 1829a: 145, 1829b: 12, 1830: 357, 1839: 207. **(ms)** Eggers 1910a, 1910b; Escherich 1932b; Schimitschek 1930b: 407.
- ramulorum* Perris 1856: 191. Syntypes, sex?, Pyrenees, France; not given, presumably MNHN, Paris. Synonymy: Blandford 1891c: 15. References: **(ay)** Nusslin 1911a: 155. **(cn)** Packard 1890: 715; Pfeffer 1933: 43; Wachtl 1883b: 319, 1901: 381; Walsh 1867a: 103. **(cc)** Perris 1852: 504, 1856a: 191, 1862: 189; Pfeffer 1928b: 8, 1933: 43; Wichmann 1959: 413. **(hb)** Barbey 1901: 23, 75; Dombrowsky 1891, 1892; Everts 1903: 758; Henschel 1895a: 173; Nordlinger 1870a: 187; Packard 1890: 715; Perris 1852: 504, 1856a: 191; Peyerimhoff 1915: 61; Rupertsberger 1879: 231, 1880: 230; Wachtl 1901: 381. **(ds)** Acloque 1896; Barthe 1896; Bedel 1888b: 298, 414; Blanchere & Robert 1889; Brancsik 1871, 1906; Bucking

1932; Carpentier & Delaby 1908; Cecconi 1897; Eggers 1904; Everts 1922: 643; Favre 1890; Gozis 1875: 80; Grill 1895: 311; Henschel 1895a: 173; Heyden, Reitter, & Weise 1891: 672, 1906: 711; Kloft & Hinks 1945: 218; Lacordaire 1866: 381; Kunnemann 1921: 58; Mequignon 1936: 15; Micke 1915: 111; Perris 1876a: 255, 1877a: 415; Peyerimhoff 1915: 61, 1933b: 367; Pfeffer 1928b: 8; Ragusa 1924: 116; Reitter 1890b: 212, 1894a: 76; Sainte-Claire 1914: 472; Schilsky 1909: 188; Stein 1868: 114; Stein & Weise 1877: 164; Stierlin 1898: 442; Strauch 1861: 122; Wichmann 1910: 146, 1927: 72. (**tx**) Acloque 1896; Balachowsky 1949a: 237; Barbey 1901: 23, 75; Bedel 1888b: 298, 414; Bertolini 1872; Blandford 1891c: 15–17; Branscik 1871; Carpentier & Delaby 1908; Dombrowsky 1891, 1892; Eggers 1915b: 13–14; Eichhoff 1878b: 178, 1883a: 110, 134; Everts 1903: 758, 1922: 643; Favel 1889; Ferrari 1867a: 32, 1867b: 114; Henschel 1895a: 173; Lacordaire 1866: 381; Kuhnt 1913: 1056; Nordlinger 1870a: 187; Nusslin 1911a: 155; Perris 1856: 196, 1877a: 415; Reitter 1894a: 76, 1901: 101, 1913a: 93; Rupertsberger 1879: 231, 1880: 230; Schedl 1934f: 1643; Spessivtsev 1913: 89; Stierlin 1898: 442; Wichmann 1910a: 146. (**ms**) Eggers 1910b; Wichmann 1959: 413.

pubifrons Bright 1981d: 195. Holotype ♀; Parque Nacional Zoquiapan, Mexico, Mexico; CNCI, Ottawa.

Distribution: North America (Mexico in Mexico).

Hosts: *Pinus hartwegii*.

References: (**hb**) Atkinson et al. 1986: 40. (**ds**) Atkinson & Equihua 1985a: 95; Atkinson et al. 1986: 40; Bright 1981d: 195. (**tx**) Bright 1981d: 195.

pudens (Blackman) 1942a: 199 (*Pityophthoroides*).

Holotype ♀; Cayamas, Cuba; USNM, Washington.

Distribution: Antilles Islands (Cuba/ Dominican Republic in Hispaniola/ Virgin Islands).

References: (**ds**) Blackwelder 1947: 782; Bright 1981c: 156, 1985c: 176. (**tx**) Blackman 1942a: 199; Bright 1985c: 176; Wood, S. L. 1977b: 207.

pudicus Blackman 1942a: 208. Holotype ♂; Guadaluajara, Jalisco, Mexico; USNM, Washington.

Distribution: North America (Jalisco in Mexico).

Hosts: *Sambucus* sp.

References: (**ds**) Blackwelder 1947; Bright 1981d: 80. (**tx**) Blackman 1942a: 208–209; Bright 1981d: 80; Wood, S. L. 1982b: 1025.

pulchellus Eichhoff 1869a: 275. Holotype, sex?;

America septentr.; Hamburg Museum, lost, neotype ♀; Marquette, Michigan [USA]; USNM, Washington, designated by Bright 1978: 73.

Figures: Bright 1981d: 350.

Distribution: North America (Alberta, British Columbia, Manitoba, New Brunswick, Ontario,

Quebec, Saskatchewan, Yukon in Canada/ District of Columbia, Florida, Indiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Jersey, New York, North Carolina, Pennsylvania, Texas, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Pinus banksiana*, *P. palustris*, *P. resinosa*, *P. rigida*, *P. strobus*, *P. virginiana*.

Notes: (**3**) Blackman 1928b: 110 (redescribed). Bright 1981d: 153 (subspecies *tuberculatus* recognized). Wood 1982b: 1091 (treated this taxon as a species distinct from *tuberculatus*, although the hosts hybridize in part of Alberta; intergradation between the beetle species is almost, if not entirely, nonexistent).

References: (**ay**) Thomas, J. B. 1957: 4. (**cn**) Blackman 1950; Doane et al. 1936; Felt 1906: 752; Hopkins 1899c: 442; Lindquist, O. H. & Syme 1981: 88; Martin, J. L. 1964; Swaine 1918a: 97, 102; Syme & Nystrom 1988: 84; Thomas, J. B. 1958: 394. (**ec**) Bushing 1965: 464; Felt 1906: 752; Frederick, Sloan, & Skowron 1976; Heqvist 1959: 181; Reid 1957b: 7; Thomas, J. B. 1958: 394. (**hb**) Baker, W. L. 1972: 255; Beal & Massey 1945: 129; Blackman 1950; Chamberlin 1939: 389; Doane et al. 1936; Felt 1906: 752; Hopkins 1899c: 442; Swaine 1918a: 97, 102. (**ds**) Anonymous 1926c: 519; Baker, W. L. 1972: 255; Beal & Massey 1945: 129; Blackman 1950; Blatchley & Leng 1916: 632; Bright 1976d: 187, 1981d: 152; Chamberlin 1939: 389; Deyrup & Atkinson 1987b: 68; Dodge 1938; Drooz 1985: 364; Evans, D. 1983: 35; Gemminger & Harold 1872: 2689; Hagedorn 1910a: 73; Henshaw 1885: 147; Hopkins 1893c: 132; Kleine 1913b: 140, 1914b: 394, 1934a: 161; Knull 1932: 66; Leng 1920: 341; Leonard 1928: 519; Lindquist, O. H. & Syme 1981: 88; Swaine 1909: 138; Syme & Nystrom 1988: 84; Wood, S. L. 1982b: 1091. (**tx**) Beal & Massey 1945: 129; Blackman 1922c: 125–126, 1928b: 91, 110–111; Blatchley & Leng 1916: 632; Bright 1978: 73, 1981d: 152, 350; Chamberlin 1939: 389; Dodge 1938: 43, 45; Eichhoff 1869a: 275, 1878b: 181, 1896: 609–610; Evans, D. 1983: 35; Hagedorn 1910a: 101; LeConte 1876: 352, 435; Lindquist, O. H. & Syme 1981: 88; Swaine 1909: 138, 1918a: 97, 102; Syme & Nystrom 1988: 84; Thomas, J. B. 1957: 4; Titus, Meikle, & Harrison 1985: 105; Wood, S. L. 1982b: 1091.

hirticeps LeConte 1878b: 623. Lectotype ♂; Marquette, Michigan [USA]; MCZ, Cambridge, designated by Bright 1976b: 185. Synonymy: Eichhoff 1896: 609.

References: (**cc**) Felt 1906: 751–752. (**hb**) Felt 1906: 752; Schwarz 1888a: 80. (**ds**) Chamberlin 1925; Henshaw 1885: 147; Hopkins 1893a: 131, 1893b: 209, 1899: 442; Hubbard & Schwarz 1878a: 665, 1878b: 623; Leng 1920: 341; Schwarz 1886: 41, 1888a: 80; Swaine 1909: 138, 1918a: 102; Ulke 1902: 56. (**tx**) Blackman 1922b: 125–126, 1928b: 110; Bright

- 1976b: 185; Eichlhoﬀ 1896: 609–610; LeConte 1875b: 623; Schwarz 1886: 41, 1888: 80; Swaine 1909: 138.
- pusio* LeConte 1875b: 623. Holotype ♂; Marquette, Michigan [USA]; MCZ, Cambridge. Synonymy: Blackman 1928b: 111. References: (ds) Blatchley & Leng 1916: 632; Hagedorn 1910d: 75; Henshaw 1882: 265, 1885: 147; Hubbard & Schwarz 1878a: 643, 665, 1878b: 623; Kleine 1913b: 141; Leng 1920: 341; Swaine 1909: 140. (tx) Blackman 1922b: 125–126, 1928b: 110; Blatchley & Leng 1916: 632; Hagedorn 1910a: 101; LeConte 1875b: 623; Swaine 1909: 140, 1918a: 102.
- pulicarius* (Zimmermann) 1868: 144 (*Crypturgus*). Lectotype, sex²; Lake Superior [USA]; MCZ, Cambridge. Figures: Bright 1981d: 55, 339. Distribution: Antilles Islands (Cuba), North America (Manitoba, New Brunswick, Nova Scotia, Ontario, Quebec, Saskatchewan in Canada/ Alabama, Arkansas, Connecticut, Delaware, Florida, Georgia, Illinois, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, Texas, Virginia, West Virginia, Wisconsin in USA). Hosts: *Pinus caribaea*, *P. echinata*, *P. palustris*, *P. resinosa*, *P. rigida*, *P. strobus*, *P. sylvestris*, *P. taeda*, *P. virginiana* twigs, uncommon in *Abies balsamea*, *Picea* spp. Notes: (3) Blackman 1928b: 101 (redescribed). References: (bv) Clark 1972. (cn) Anonymous 1962h, 1963r, 1965b; Baker, W. L. 1972: 255; Blackman 1950; Browne 1968b: 565; Clark 1972; Doane et al. 1936; Drooz 1985: 363; Felt 1906: 751, 1933b: 994–995, 1934: 195–200; Hoekstra, Merkel, & Powers 1961; Hopkins 1899c: 442; Kowal 1956a: 9; Lindquist, O. H. & Syme 1981: 88; Mason, G. N. & Jones 1969: 12; Merkel & Kowal 1956: 7; Ollien & Mason 1968: 13; Perry & Wu 1957: 10; Smith, R. H. & Mergen 1959a: 564–565; Swaine 1918a: 95, 99; Syme & Nystrom 1988: 84; Yearian & Warren 1964. (cc) Felt 1906: 751; Hines, J. W. & Heikkinen 1977; Schwarz 1889a: 163; Thomas & Lindquist 1956: 2. (hb) Baker, W. L. 1972: 255; Beal & Massey 1945: 133–134; Blackman 1950; Browne 1968b: 565; Chamberlin 1939: 374; Clark 1972; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Felt 1906: 751; Hopkins 1899c: 442; Ostmark 1968; Swaine 1918a: 95, 99. (ds) Anonymous 1926c: 518, 1963r, 1965b; Beal & Massey 1945: 133–134; Beaulne 1956; Blackman 1950; Blatchley & Leng 1916: 629; Bright 1976d: 187, 1981d: 84, 1985c: 84; Browne 1968b: 565; Chamberlin 1925, 1939: 374; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Dodge 1938; Drooz 1985: 363; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 74; Henshaw 1885: 147; Kirk 1969, 1970; Kleine 1913b: 141, 1934a: 161; Knull 1932: 66, 212; Leng 1920: 341; Leonard 1928: 518; Lindquist, O. H. & Syme 1981: 88; Merkel & Kowal 1956: 7; Ostmark 1968; Proctor 1946: 208; Schuder 1969: 76; Schwarz 1878d: 468, 1889a: 163–164, 1890: 231; Smith, J. B. 1890: 267, 1900: 362, 1910: 401; Swaine 1909: 138; Syme & Nystrom 1988: 84; Ulke 1902: 56; Wood, S. L. 1982b: 1048; Yearian & Warren 1964: 262. (tx) Beal & Massey 1945: 133–134; Blackman 1922b: 101–103, 1928b: 59, 63–64; Blatchley & Leng 1916: 629; Bright 1976b: 187, 1981d: 84–85, 339, 1985c: 176; Britton 1920a: Chamberlin 1939: 374; Dodge 1938: 15, 43–44; Eichlhoﬀ 1878b: 195; Hagedorn 1910a: 101; LeConte 1868: 155, 157, 1876: 353, 1878a: 468; Lindquist, O. H. & Syme 1981: 88; Swaine 1909: 138, 1918a: 95, 99; Wood, S. L. 1977b: 210, 1982b: 1048; Zimmermann 1868: 144. (ms) Hoekstra, Merkel, & Powers 1961; Schuder 1969: 76.
- cubensis* Schedl 1972: 65. Holotype ♂; Cuba, Vinales, Prov. de Pinar del Rio; Schedl Collection in NHMW, Wien. Synonymy: Wood 1977b: 210. References: (tx) Schedl 1972: 65, 1979c: 71; Wood, S. L. 1977b: 210.
- pullus* (Zimmermann) 1868: 143 (*Crypturgus*). Holotype ♀; South Carolina [USA]; MCZ, Cambridge. Distribution: North America (District of Columbia, Florida, Illinois, Maryland, Massachusetts, Michigan, Mississippi, New Jersey, New York, North Carolina, Pennsylvania, South Carolina, Texas, West Virginia in USA). Hosts: *Pinus echinata*, *P. palustris*, *P. strobus*, *P. taeda*, *P. spp.* Notes: (1) LeConte 1876: 352 (to *Pityophthorus*). (3) LeConte 1868: 155 (cited in *Cryphalus*). Blackman 1928b: 118 (redescribed). References: (cn) Blackman 1950; Chamberlin 1924; Doane et al. 1936; Felt 1906: 751; Hopkins 1899c: 357, 442; Swaine 1918a: 98, 103. (ec) Felt 1906: 751. (hb) Beal & Massey 1945: 131; Blackman 1922b: 101, 105–106, 1950; Chamberlin 1939: 394; Doane et al. 1936; Felt 1906: 751; Hopkins 1899c: 357, 442–444; Schwarz 1888a: 80; Swaine 1918a: 98, 103. (ds) Anonymous 1926c: 519; Atkinson et al. 1991: 163; Beal & Massey 1945: 131; Blackman 1922b: 101, 105–106, 1950; Blatchley & Leng 1916: 636; Bright 1981d: 287; Chamberlin 1925, 1939: 394; Deyrup & Atkinson 1987b: 68; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 74; Henshaw 1885: 147; Hopkins 1893a: 129, 1893b: 208; Hubbard & Schwarz 1878: 643; Kirk 1969; Kleine 1913b: 141, 1914b: 391, 1934a: 161; Leng 1920: 341; Leonard 1928: 519; Schwarz 1887: 20, 1888a: 80, 1892: 168; Smith, J. B. 1890: 267, 1900: 362; Swaine

- 1909: 138; Townsend 1889: 235; Ulke 1902: 56; Wood, S. L. 1982b: 1105. **(tx)** Beal & Massey 1945: 131; Blackman 1922b: 101, 105–106, 1928b: 114–118; Blandford 1898c: 5; Blatchley & Leng 1916: 636; Bright 1976b: 187, 1981d: 287; Chamberlin 1939: 394; Eichhoff 1878b: 186; Hagedorn 1910a: 101; LeConte 1868: 155, 157, 1876: 352; Schwarz 1891a: 168; Swaine 1909: 138–139, 1918a: 98, 103; Wood, S. L. 1977d: 516, 1982b: 1105; Zimmernann 1865: 143. **(ms)** Eckstein 1900c.
- cribripennis* Eichhoff 1869a: 274. Holotype, sex?; America septentr.; Hamburg Museum, lost, neotype, sex?; Ripley, Mississippi [USA]; USNM, Washington, designated by Bright 1978: 72. Synonymy: Blackman 1928b: 119. References: **(cc)** Felt 1906: 752. **(hb)** Felt 1906: 751–752. **(ds)** Hopkins 1899: 437–442; Leng 1920: 341; Swaine 1909: 139. **(tx)** Blackman 1928b: 118; Bright 1978: 72; Eichhoff 1869a: 274, 1872b: 137, 1878b: 175, 1896: 609–610; LeConte 1876: 354; Swaine 1909: 139.
- bisulcatus* Eichhoff 1869a: 274. Holotype, sex?; America borealis; Hamburg Museum, lost. Synonymy: LeConte 1876: 352. References: **(bv)** Dixon & Payne 1979b. **(cn)** Hopkins 1899c: 443; Swaine 1918a: 98, 103. **(cc)** Cross & Moser 1971; Dixon & Payne 1979b; Moser, J. C., Cross, & Roton 1971. **(hb)** Blackman 1922b: 108–109; Chamberlin 1939: 400; Hopkins 1899c: 443; Swaine 1918a: 98, 103. **(ds)** Blackman 1922b: 108–109; Chamberlin 1925, 1939: 400; Kirk 1970; Leng 1920: 341; Swaine 1909: 139. **(tx)** Blackman 1922b: 102, 108–109, 1928b: 117, 138; Chamberlin 1939: 400; Eichhoff 1869a: 274, 1878b: 185; LeConte 1876: 352; Swaine 1909: 139, 1918a: 98, 103; Wood, S. L. 1977d: 516.
- punctatus* Eggers 1940a: 130. Holotype ♀; Guadeloupe (Trois Rivières); Eggers Collection, in NHMW, Wien. Distribution: Antilles Islands (Guadeloupe). References: **(cn)** Schuder 1969: 80. **(hb)** Bright 1981d: 287. **(ds)** Bright 1985c: 176; Schuder 1969: 80. **(tx)** Bright 1985c: 176; Eggers 1940a: 130; Schedl 1979c: 204.
- punctifrons* Bright 1966b: 298. Holotype ♀; Frazier Park, Kern Co., California [USA]; CAS, San Francisco. Distribution: North America (California, New Mexico in USA). Hosts: *Pinus edulis*, *P. monophylla*, *P. quadrifolia*. References: **(hb)** Bright & Stark 1973: 106. **(ds)** Bright 1981d: 143; Bright & Stark 1973: 106; Wood, S. L. 1982b: 1035. **(tx)** Bright 1966b: 298, 1981d: 143; de Ruette 1970: 110; Wood, S. L. 1982b: 1035.
- punctiger* (Schedl) 1951m: 108 (*Neodryocoetes*). Holotype, sex?; Mexico, collected by Flohr; Schedl Collection in NHMW, Wien. Distribution: North America (“Mexico”). References: **(ds)** Wood, S. L. 1982b: 1030. **(tx)** Schedl 1951m: 108, 1979c: 205; Wood, S. L. 1982b: 1030.
- pygmaeolus* Schedl 1970e: 93. Holotype, sex?; Brasilien, Caioba, 45 50, 48 40, 10 m; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: **(tx)** Schedl 1970e: 93, 1979c: 207.
- quadriscopinus* Schedl 1966f: 110. Holotype ♂; published as Surinam, Moengo, type in Schedl Collection (♀) labeled Camopi-Oyapock, Guayana (dans nid d’*Atta*); Schedl Collection in NHMW, Wien. Distribution: South America (Suriname). References: **(ds)** Schedl 1966f: 79, 110, 1979c: 208.
- quercinus* Wood 1967d: 40. Holotype ♀; 5 km W El Salto, Durango, Mexico; Wood Collection. Distribution: North America (Durango, Michoacan, Oaxaca in Mexico/ W Texas in USA). Hosts: *Quercus* sp. bole. Notes: (3) Bright 1977: 515 (a synonym of *guatemalensis*). Wood 1982b: 1085 (a distinct species or a very distinct geographical race of *guatemalensis*). References: **(ds)** Wood, S. L. 1982b: 1085. **(tx)** Bright 1977: 515; de Ruette 1970: 111; Wood, S. L. 1967d: 40–41, 1982b: 1085.
- ramiperda* Swaine 1917: 28. Holotype ♀; Ile Perrot, Quebec [Canada]; CNCI, Ottawa. Distribution: North America (Ontario, Quebec in Canada/ Maine, Massachusetts, Michigan, New York, Rhode Island, Vermont, Wisconsin in USA). Hosts: *Pinus resinosa*, *P. strobus*. References: **(cn)** Blackman 1950; Britton 1923: 369; Chamberlin 1924; Pettit 1928: 1–20; Randall 1941: 53, 1943: 53; Sterner & Davidson 1983: 37; Swaine 1918a: 94, 98. **(cc)** Bushing 1965: 462. **(hb)** Baker, W. L. 1972: 259; Blackman 1950; Bright 1981d: 207; Chamberlin 1939: 341; Swaine 1918a: 94, 98. **(ds)** Anonymous 1926c: 518; Baker, W. L. 1972: 259; Beaulne 1956; Blackman 1930; Bright 1976d: 168, 1981d: 207; Chamberlin 1939: 341; Drooz 1985: 364; Kleine 1934a: 161, 163; Leng 1920: 341; Leonard 1928: 518; Proctor 1946: 208; Wood, S. L. 1982b: 1044. **(tx)** Blackman 1928b: 16–17, 22–23; Bright 1967b: 678, 1976c: 426, 1976d: 168, 1981d: 207; Chamberlin 1939: 341; Dodge 1938: 41; Hoebeke 1978; de Ruette 1970: 111; Swaine 1917: 28–29, 1918a: 94, 98, 1925b: 192; Wood, S. L. 1982b: 1044.
- fivasi* Blackman 1928b: 23 (*Myeloborus*). Holotype ♀; Cranberry Lake, New York [USA]; USNM, Washington. Synonymy: Bright 1976c: 426. References: **(cn)** Doane et al. 1936. **(hb)** Baker, W. L. 1972: 259; Blackman 1928b; Bright 1976d: 168; Chamberlin 1939: 341; Doane et

al. 1936. (**ds**) Baker, W. L. 1972: 259; Bright 1976d: 165; Chamberlin 1939: 341; Dodge 1935: 41–42; Leng & Mutchler 1933: 53. (**tx**) Blackman 1928b: 23; Bright 1976c: 426, 1976d: 165; Chamberlin 1939: 341; Dodge 1935: 41–42; Schedl 1938a: 161.

recens Bright 1976c: 435. Holotype ♀; Pine Pass, British Columbia [Canada]; CNCI, Ottawa. Distribution: North America (British Columbia, Yukon in Canada).

Hosts: *Picea glauca*, *P. engelmannii*.

References: (**ds**) Bright 1981d: 142; Wood, S. L. 1982b: 1061. (**tx**) Bright 1976c: 435, 1981d: 142; McNamara 1984: 757; Wood, S. L. 1982b: 1061.

regularis Blackman 1942a: 206. Holotype, sex?; Cayamas, Cuba; USNM, Washington. Distribution: Antilles Islands (Cuba). References: (**ds**) Blackwelder 1947: 782; Bright 1985c: 176. (**tx**) Blackman 1942a: 206; Bright 1985c: 176; Wood, S. L. 1964: 63.

robai (Blackman) 1942a: 200 (*Pityophthoroides*). Holotype ♀; Dept. Santander, Colombia, S. America. 700–1300 m; USNM, Washington. Figures: Blackman 1942a: fig. 28. Distribution: South America (Colombia). Hosts: *Psidium guajava*. Notes: (1) Wood 1977b: 207 (to *Pityophthorus*). References: (**ds**) Blackwelder 1947: 781. (**tx**) Blackman 1942a: 200; Wood, S. L. 1977b: 207.

roppae Schedl 1976a: 69. Holotype ♀; Linhares, E. Santo, Brasil; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (**tx**) Schedl 1976a: 69.

rossicus Eggers 1915b: 13. Holotype ♀; Russia: Tambov; Eggers Collection, in NHMW, Wien. Distribution: Europe (W USSR). References: (**hb**) Stark 1952: 352. (**ds**) Kleine 1934a: 161; Postner 1974: 437; Stark 1952: 352. (**tx**) Eggers 1915b: 13–14; Pfeiffer 1940b: 108, 112, 120, 1979c: 213; Postner 1974: 437; Schedl 1934f: 1643; Spessivtsev 1926: 50; Stark 1952: 352.

rubidus Wood 1978b: 400. Holotype ♀; Madera Canyon, Santa Rita Mts., Arizona [USA]; Wood Collection. Distribution: North America (Durango in Mexico, Arizona in USA). Hosts: *Pinus cooperi*, *P. engelmannii*, *P. leiophylla*, *P. lumholtzi*, *P. ponderosa*. References: (**ds**) Bright 1981d: 301; Wood, S. L. 1982b: 1113. (**tx**) Bright 1981d: 301; Wood, S. L. 1975b: 400, 1982b: 1113.

rudis Blackman 1942a: 212. Holotype ♀; Amecameca, Distrito Federal, Mexico; USNM, Washington. Figures: Blackman 1942a: fig. 46. Distribution: North America (Distrito Federal in Mexico). Hosts: *Abies religiosa*, *Pinus* sp.

References: (**hb**) Atkinson et al. 1986: 40. (**ds**) Atkinson & Equihua 1985a: 95; Atkinson et al. 1986: 40; Blackwelder 1947: 782; Bright 1981d: 223; Wood, S. L. 1982b: 1058. (**tx**) Blackman 1942a: 212; Bright 1981d: 223; McNamara 1984: 757; Wood, S. L. 1982b: 1058.

sachalinensis Krivolutsкая 1956: 835. Holotype, sex?; Sakhalin: Kirovsk region, Tyn river basin, village of Argi-Pag; Institute of Zoology, Academy of Science, Vladivostok. Distribution: Asia (Sakhalin Island in USSR). References: (**hb**) Krivolutsкая 1956: 835. (**ds**) Krivolutsкая 1956: 835; Kurenzov 1967; Pfeiffer 1976: 340. (**tx**) Krivolutsкая 1956: 835, 1958: 163; Pfeiffer 1976: 340.

sambuci Blackman 1942a: 207. Holotype ♂; Jalisco, Mexico; USNM, Washington. Figures: Blackman 1942a: fig. 38, Bright 1981d: 337. Distribution: North America (Jalisco in Mexico). Hosts: *Sambucus* sp. References: (**ds**) Blackwelder 1947: 782; Bright 1981d: 58; Wood, S. L. 1982b: 1132. (**tx**) Blackman 1942a: 207; Bright 1981d: 58, 337; McNamara 1984: 757; Wood, S. L. 1982b: 1132.

sapineus Bright 1981d: 194. Holotype ♀; Atenquique, Jalisco, Mexico; CNCI, Ottawa. Figures: Bright 1981d: 357. Distribution: North America (Jalisco in Mexico). Hosts: *Pinus* sp. References: (**ds**) Atkinson & Equihua 1985c: 360, 1988: 98; Wood, S. L. 1982b: 1056. (**tx**) Bright 1981d: 194, 357; Wood, S. L. 1982b: 1056.

scabridus Schedl 1955g: 24. Lectotype ♀; Quezaltenango, Guatemala; Schedl Collection in NHMW, Wien, designated by Bright 1976b: 186. Figures: Bright 1981d: 365. Distribution: North America (El Salvador/ Guatemala/ Honduras/ Chiapas, Jalisco, Mexico, Michoacan, Morelos, Nayarit, Oaxaca, Veracruz in Mexico). Hosts: *Pinus ayacaluite*, *P. caribaea*, *P. montezumae*, *P. oocarpa*, *P. patula*, *P. pseudostrobus*, *P. rudis*. References: (**hb**) Atkinson et al. 1986: 40; Bright 1981d: 257; Burgos & Sancedo 1983: 117; Schwerdtfeger 1957: 502. (**ds**) Atkinson & Equihua 1985a: 96, 1985c: 360, 1988: 98; Atkinson et al. 1986: 40; Bright 1981d: 257; Burgos & Sancedo 1983: 117; Schedl 1963c: 158, 1977e: 42; Wood, S. L. 1982b: 1050. (**tx**) Bright 1976b: 186, 1981d: 257, 365; Schedl 1955g: 18, 24, 1979c: 220; Wood, S. L. 1982b: 1050.

scalptor Blackman 1928b: 30. Holotype ♀; Julian, California [USA]; USNM, Washington. Distribution: North America (British Columbia in Canada/ California, Idaho, Oregon in USA). Hosts: *Pinus contorta*, *P. coulteri*, *P. jeffreyi*, *P. ponderosa*.

- References: (bv) Moeck, Wood, & Lindahl 1981. (cn) Lindgren 1980: 69. (hb) Bright & Stark 1973: 105; Furniss, M. M. & Johnson 1987: 377; Lindgren 1980: 69. (ds) Bright 1981d: 135; Bright & Stark 1973: 105; Chamberlin 1939: 357; Furniss, M. M. & Johnson 1987: 377; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 152; Wood, S. L. 1982b: 1051. (tx) Blackman 1928b: 30; Bright 1981d: 135; Chamberlin 1939: 357; Hopping 1960a: 867; Keen 1929a: 32; de Ruelle 1970: 111; Wood, S. L. 1982b: 1051.
- scalptus** Bright 1978: 82. Holotype ♀; Aspen Grove, British Columbia [Canada]; CNCI, Ottawa.
Distribution: North America (British Columbia in Canada/ Colorado, Montana in USA).
Hosts: *Pinus ponderosa*.
References: (ds) Bright 1981d: 127; Wood, S. L. 1982b: 1052. (tx) Bright 1978: 82, 1981d: 127; Cast et al. 1989: 385; McNamara 1984: 757; Wood, S. L. 1982b: 1052.
- schrenkianae** Pjatnitskii 1931: 171. Lectotype, sex?; Kirgisien, USSR: Pjatnitskii Collection not located, designated by Michalski 1969a: 895.
Distribution: Asia (E USSR).
Hosts: *Picea schrenkianae*.
References: (ce) Kostin 1964: 112. (hb) Kostin 1960: 133; Stark 1952: 347. (ds) Kleine 1934a: 161; Marikovskii 1956: 73; Parfentev 1951: 429–430; Stark 1952: 347. (tx) Michalski 1969a: 895, 1969b: 569; Pfeffer 1940b: 107, 110, 116; Pjatnitskii 1931: 171; Schedl 1934f: 1643, 1979c: 222; Sokanovskii 1954: 19; Stark 1952: 347.
- schwarzii** Blackman 1928b: 71. Holotype ♀; Las Vegas H.S. [Hot Springs], New Mexico [USA]; USNM, Washington.
Figures: Bright 1981d: 368.
Distribution: North America (Coahuila, Durango, Hidalgo, Nuevo Leon, Queretaro, Tlaxcala in Mexico/ Arizona, New Mexico, W Texas, Utah, Wyoming in USA).
Hosts: *Pinus cembroides*, *P. edulis*, *P. greggii*, *P. leiophylla*, *P. humboltzii*, *P. montezuma*, *P. ponderosa*, *P. teocote*.
References: (hb) Chamberlin 1939: 377. (ds) Atkinson & Equihua 1988: 98; Bright 1981d: 273; Chamberlin 1939: 377; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1080. (tx) Blackman 1928b: 71–72; Bright 1981d: 273, 368; Chamberlin 1939: 377; Keen 1929a: 32; de Ruelle 1970: 111; Wood, S. L. 1982b: 1080.
- schwerdtfegeri** (Schedl) 1955g: 28 (*Conophthorus*). Lectotype ♂; Guatemala: Strasse Quezaltenango-Huehuetenango, Rancho Alegre; Schedl Collection in NHMW, Wien, designated by Bright 1976b: 187.
Figures: Bright 1981d: 340.
Distribution: North America (Guatemala/ Honduras/ Chiapas, Durango, Mexico, Nuevo Leon in Mexico/ Arizona, Nevada, W Texas in USA).
Hosts: *Pinus engelmannii*, *P. oocarpa*, *P. ponderosa*, *P. pseudostrobus*.
References: (cn) Cibrian-Tovar et al. 1986: 47. (hb) Atkinson & Equihua 1985c: 359; Atkinson et al. 1986: 41; Bright 1981d: 88; Burgos & Saucedo 1983: 116; Schwerdtfeger 1957c: 506. (ds) Atkinson & Equihua 1985c: 359, 1985: 98; Atkinson et al. 1986: 41; Bright 1981d: 88; Burgos & Saucedo 1983: 116; Cibrian-Tovar et al. 1986: 47; Rio Mora 1985: 13; Wood, S. L. 1982b: 1047. (tx) Bright 1976b: 187, 1981d: 88, 340; Rio Mora 1985: 13; Schedl 1955g: 18, 28, 1963c: 163, 1979c: 222; Wood, S. L. 1966b: 28, 1982b: 1047.
- islasi** Wood 1962: 80. Holotype ♀; Temascaltepec, Mexico, Mexico: Wood Collection.
Synonymy: Wood 1966b: 28.
References: (tx) Schedl 1979c: 128; Wood, S. L. 1962: 80, 1966b: 28.
- islasi** Schedl 1963c: 163 (*Conophthocrammus*). Holotype ♀; Temascaltepec, Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 28.
References: (tx) Schedl 1963c: 163, 1979c: 128; Wood, S. L. 1966b: 28.
- sctulus** Wood 1976a: 359. Holotype ♀; Volcan Chiriqui [near Cerro Punta], Chiriqui, Panama; Wood Collection.
Distribution: North America (Costa Rica/ Panama).
Hosts: *Quercus* sp. limb.
References: (ds) Bright 1981d: 104; Wood, S. L. 1982b: 1083. (tx) Bright 1981d: 104; McNamara 1984: 757; Wood, S. L. 1976a: 359, 1982b: 1083.
- scriptor** Blackman 1921: 7. Lectotype ♀; Agricultural College [Starkville], Mississippi [USA]; USNM, Washington, designated by Bright 1976b: 184.
Figures: Blackman 1922b: pl. 1, fig. 6.
Distribution: North America (Georgia, Mississippi, North Carolina, Oklahoma, Tennessee, Texas in USA).
Hosts: *Rhus typhina*.
References: (cn) Anonymous 1977f, 1978v; Blackman 1950. (hb) Baker, W. L. 1972: 255; Beal & Massey 1945: 127–128; Blackman 1922b: 106–107, 1950; Chamberlin 1939: 401. (ds) Anonymous 1977f, 1978v; Beal & Massey 1945: 127–128; Blackman 1922b: 106–107, 1950; Bright 1981b: 8; Chamberlin 1939: 401; Deyrup 1981b: 8; Drooz 1985: 363; Kleine 1934a: 161; Leng & Mutchler 1927: 52; Wood, S. L. 1982b: 1118. (tx) Beal & Massey 1945: 127–128; Blackman 1921: 7–8, 1922b: 106–107, 1928b: 114, 142; Bright 1976b: 184, 1981d: 39; Chamberlin 1939: 401; de Ruelle 1970: 111; Wood, S. L. 1982b: 1118.
- segnis** Blackman 1928b: 52. Holotype ♀; Santa Catalina Mts., Arizona [USA]; USNM, Washington.
Figures: Bright 1981d: 358.

Distribution: North America (Durango in Mexico/ Arizona, New Mexico in USA).

Hosts: *Pinus engelmannii*, *P. strobiformis* twigs.

Notes: (3) Bright 1951d: 199 (recognized subspecies *subopacus*). Wood 1952b: 1036 (intergrades with *subopacus*).

References: (hb) Atkinson et al. 1956: 41; Chamberlin 1939: 366. (ds) Atkinson & Equihua 1955a: 95; Atkinson et al. 1956: 40; Bright 1951d: 195; Chamberlin 1939: 366; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Wood, S. L. 1952b: 1036. (tx) Blackman 1928b: 44, 52–53; Bright 1951d: 195–199; Chamberlin 1939: 366; Keen 1929a: 32; McNamara 1954: 757; Wood, S. L. 1952b: 1036.

seiryuensis Murayama 1963c: 393. Holotype, sex?; Seiryu, South Manchuria; Murayama Collection in USNM, Washington.

Distribution: Asia (Manchuria).

Hosts: *Pinus tabulaeformis*.

References: (tx) Murayama 1963c: 393.

separatus Bright 1977: 529. Holotype ♂; Poinsett State Park, South Carolina [USA]; Wood Collection. Distribution: North America (South Carolina in USA).

Hosts: (?) *Pinus* sp. (probably in twigs of living trees).

References: (ds) Bright 1951d: 203; Wood, S. L. 1952b: 1042. (tx) Bright 1977: 529, 1951d: 203; Wood, S. L. 1952b: 1042.

serratus Swaine 1915a: 103. Holotype ♂; Barkhouse Creek, Siskiyou Co., California [USA]; CNCI, Ottawa.

Distribution: North America (S British Columbia in Canada/ California, N Idaho in USA).

Hosts: *Pinus ponderosa*.

References: (cn) Keen 1929: 53; Swaine 1915a: 95, 103. (hb) Bright & Stark 1973: 115; Chamberlin 1939: 395, 1955: 157; Keen 1929: 53; Swaine 1915a: 95, 103. (ds) Bright 1951d: 304; Bright & Stark 1973: 115; Chamberlin 1939: 395, 1955: 157; Hopping 1922; Keen 1929a: 33, 1929c: 53; Kleine 1934a: 161; Leng 1920: 341; Wood, S. L. 1972a: 427, 1952b: 1106. (tx) Blackman 1928b: 115, 122; Bright 1967b: 678, 1951d: 304; Chamberlin 1939: 395, 1955: 157; Keen 1929a: 33; de Ruelle 1970: 111; Swaine 1915a: 95, 103; Wood, S. L. 1972a: 427, 1952b: 1106.

setifer Browne 1965a: 193. Holotype, sex?; Ivory Coast: Adiopodoume; RNH, Leiden.

Distribution: Africa (Ghana/ Ivory Coast).

References: (ds) Schedl 1972e: 252. (tx) Browne 1965a: 193.

setosus Blackman 1928b: 77. Holotype ♀; Monterey, California [USA]; USNM, Washington.

Figures: Bright 1951d: 361.

Distribution: North America (California in USA).

Hosts: *Pinus radiata*, 1 series from *P. muricata*.

References: (cc) Kinn 1971. (hb) Bright & Stark

1973: 109; Chamberlin 1939: 378. (ds) Bright 1951d: 222; Bright & Stark 1973: 109; Chamberlin 1939: 378; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Wood, S. L. 1952b: 1073. (tx) Blackman 1928b: 62, 77–78; Bright 1951d: 222, 361; Chamberlin 1939: 378; Keen 1929a: 32; de Ruelle 1970: 111; Schedl 1931b: 166; Wood, S. L. 1952b: 1073.

sextuberculatus Eggers 1933b: 6. Holotype, sex?; Franz, Guayana (Nonveau Chantier: Bas-Maroni); MNHN, Paris.

Distribution: South America (Cayenne).

References: (ds) Blackwelder 1947: 782. (tx) Eggers 1933b: 6–7.

sichotensis Kurenzov 1941a: 175. Holotype, sex?; Ussuri, USSR; Kurenzov Collection in ISBN, Novosibirsk.

Distribution: Asia (USSR).

Hosts: *Picea ajawensis*.

References: (hb) Krivolutsкая 1956: 838; Kurenzov 1945b: 111; Stark 1952: 358. (ds) Krivolutsкая 1956: 838; Pfeffer 1976: 340; Sokanovskii 1960; Stark 1952: 358. (tx) Krivolutsкая 1956: 838; Kurenzov 1941a: 175, 1945b: 111; Michalski 1969b: 570; Pfeffer 1976: 340; Sokanovskii 1960: 675; Stark 1952: 358.

sierrensis Bright 1971b: 64. Holotype ♀; 1 mile S Onion Valley, Inyo Co., California [USA]; CNCI, Ottawa.

Figures: Bright 1951d: 365.

Distribution: North America (British Columbia in Canada/ California, Colorado, Nevada, Wyoming in USA).

Hosts: *Pinus albicaulis*, *P. aristata*, *P. balfouriana*, *P. flexilis*.

References: (hb) Bright & Stark 1973: 110. (ds) Bright 1951d: 259; Bright & Stark 1973: 110. (tx) Bright 1971b: 64–65, 1951d: 259, 365; McNamara 1977: 198; Wood, S. L. 1952b: 1063.

signatifrons Browne 1975a: 758. Holotype ♂; Tanzania: Mts. Uluguru, Kimboza, 600 m; MRCB, Tervuren.

Distribution: Africa (Tanzania).

References: (tx) Browne 1975a: 758.

similis Eichhoff 1869a: 275. Syntypes, sex?; Venezuela; Hamburg Museum, lost.

Distribution: South America (Venezuela).

References: (ds) Gemminger & Harold 1872: 2689; Kleine 1913b: 141, 1914b: 339. (tx) Eggers 1933b: 3; Eichhoff 1869a: 275, 1878b: 182; Hagedorn 1905a: 412.

sinopae Schedl 1976a: 69. Holotype, sex?; Sinop, M. Grosso, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 69.

sobrinus Wood 1976a: 357. Holotype ♀; 3 km SE Cartago, Cartago, Costa Rica, 1300 m; Wood Collection.

- Distribution: North America (Costa Rica).
 Hosts: Liana.
 References: **(ds)** Bright 1981d: 46; Wood, S. L. 1982b: 1140. **(tx)** Bright 1981d: 46; McNamara 1984: 758; Wood, S. L. 1976a: 357, 1982b: 1140.
- solatus** Wood 1977b: 215. Holotype ♀; 81 km NW Oaxaca, Oaxaca, Mexico; CNCI, Ottawa.
 Distribution: North America (Oaxaca in Mexico).
 Hosts: *Pinus lawsonii*, *P. spp.*
 References: **(ds)** Bright 1981d: 327; Wood, S. L. 1982b: 1108. **(tx)** Bright 1981d: 327; Wood, S. L. 1977b: 215, 1982b: 1108.
- solers** Blackman 1928b: 138. Holotype ♀; Cloudcroft, New Mexico [USA]; USNM, Washington.
 Figures: Bright 1981d: 375.
 Distribution: North America (Chihuahua, Nuevo Leon in Mexico/ Arizona, New Mexico, Utah in USA).
 Hosts: *Abies concolor* twigs, uncommon in *A. lasiocarpa*, *A. magnifica*, *A. religiosa*, *Pseudotsuga menziesii*, *Pinus* sp.
 References: **(hb)** Chamberlin 1939: 401. **(ds)** Bright 1981d: 325; Chamberlin 1939: 401; Keen 1929a: 34; Kleine 1934a: 161; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1108. **(tx)** Blackman 1928b: 117, 138–140; Bright 1981d: 325, 374; Chamberlin 1939: 401; Keen 1929a: 34; de Ruette 1970: 111; Schedl 1930: 197; Wood, S. L. 1982b: 1108.
- solus** Blackman 1928b: 64. Holotype ♂?; Pinal Mountains, Arizona [USA]; USNM, Washington.
 Figures: Blackman 1981d: 348.
 Distribution: North America (Chiapas, Distrito Federal, Durango, Hidalgo, Mexico, Oaxaca, Puebla, Queretaro, Tlaxcala in Mexico/ Arizona, New Mexico in USA).
 Hosts: *Pinus ayacahuite*, *P. edulis*, *P. greggii*, *P. hartwegii*, *P. lawsonii*, *P. leiophylla*, *P. montezumae*, *P. oocarpa*, *P. spp.*
 References: **(hb)** Atkinson et al. 1986: 41; Burgos & Saucedo 1983: 116; Chamberlin 1939: 375. **(ds)** Atkinson & Equihua 1985a: 94, 1985c: 360; Atkinson et al. 1986: 41; Bright 1981d: 140; Burgos & Saucedo 1984: 116; Chamberlin 1939: 375; Leng & Mutchler 1933: 52; Schedl 1960a: 75; Wood, S. L. 1982b: 1148. **(tx)** Blackman 1928b: 64; Bright 1977: 518, 1981d: 140, 348; Chamberlin 1939: 375; Wood, S. L. 1982b: 1048.
- cribratus** Blackman 1942a: 216. Holotype ♀; Mexico, D.F., Mexico; USNM, Washington, preoccupied by Pfeffer 1940. Synonymy: Bright 1977: 518.
 References: **(ds)** Blackwelder 1947: 781. **(tx)** Blackman 1942a: 216; Bright 1977: 518; McNamara 1984: 755; Wood, S. L. 1982b: 1048.
- spadix** Blackman 1942a: 219. Holotype ♀; Rio Frio, Puebla, Mexico; USNM, Washington.
 Distribution: North America (Chihuahua, Durango, Hidalgo, Mexico, Michoacan, Puebla, Oaxaca, Veracruz in Mexico).
 Hosts: *Pinus leiophylla*, *P. pseudostrobus*, *P. spp.*
 References: **(ds)** Atkinson & Equihua 1985a: 97, 1985c: 360; Blackwelder 1947: 782; Bright 1981d: 300; Wood, S. L. 1982b: 1112. **(tx)** Blackman 1942a: 219; Bright 1981d: 300; McNamara 1984: 758; Wood, S. L. 1982b: 1112.
- sparsepilosus** (Schedl) 1935h: 343 (*Gnathophorus*).
 Syntypes ♀; Hamburgfarm on Rio Reventazon, Limon, Costa Rica; Schedl Collection in NHMW, Wien.
 Distribution: North America (Costa Rica).
 Hosts: (?) *Protium* sp. limbs.
 Notes: (1) Wood 1975b: 391 (to *Pityophthorus*).
 References: **(ds)** Blackwelder 1947: 782; Wood, S. L. 1982b: 1135. **(tx)** Bright 1981d: 18; Schedl 1935h: 343, 1979c: 233; Wood, S. L. 1975b: 391, 1982b: 1135.
- speciosus** Wood 1977b: 215. Holotype ♀; 24 km S Valle Nacional, Oaxaca, Mexico, 1300 m; CNCI, Ottawa.
 Distribution: North America (Oaxaca in Mexico).
 References: **(ds)** Bright 1981d: 49; Wood, S. L. 1982b: 1129. **(tx)** Bright 1981d: 49; Wood, S. L. 1977b: 215, 1982b: 1129.
- speculum** Bright 1976c: 440. Holotype ♀; Cerro Potosi, Nuevo Leon, Mexico, 3600 m; CNCI, Ottawa.
 Figures: Bright 1981d: 345.
 Distribution: North America (Hidalgo, Nuevo Leon in Mexico).
 Hosts: *Abies* sp., *Pseudotsuga menziesii*.
 References: **(hb)** Atkinson & Equihua 1985a: 93, 1985c: 350; Bright 1981d: 126. **(ds)** Atkinson & Equihua 1985a: 93, 1985c: 360, 1988: 98; Bright 1976c: 440, 1981d: 126; Wood, S. L. 1982b: 1072. **(tx)** Bright 1976c: 440, 1981d: 126, 345; McNamara 1984: 758; Wood, S. L. 1982b: 1072.
- strictus** Wood 1976a: 354. Holotype ♀; Santa Ana, San Jose, Costa Rica; Wood Collection.
 Distribution: North America (Costa Rica).
 Hosts: *Rhcedia edulis*.
 References: **(ds)** Bright 1981d: 77; Wood, S. L. 1982b: 1124. **(tx)** Bright 1981d: 77; Wood, S. L. 1976a: 354, 1982b: 1124.
- subconcentralis** Schedl 1938a: 183. Lectotype, sex?; Trois Rivières, Guadeloupe, Dufour; USNM, Washington, designated by Anderson & Anderson 1971: 32.
 Distribution: Antilles Islands (Guadeloupe).
 References: **(ds)** Blackwelder 1947: 782; Bright 1985c: 176. **(tx)** Anderson, W. H. & Anderson 1971: 32; Bright 1985c: 176; Eggers 1941a: 108, 1951: 150; Schedl 1938a: 183, 1964m: 317, 1979c: 240.

subcristatus Schedl 1937h: 168. Holotype ♀; Costa Rica. Hamburgfarm, Reventazon, Ebene Limon; Schedl Collection in NHMW, Wien. Figures: Blackman 1942a: 226.

Distribution: North America (Costa Rica/ Nicaragua/ Panama).

Hosts: (?) *Protium* sp. limbs.

Notes: (3) Bright 1977: 513 (to *Gnatholeptus*). Wood 1982b: 1143 (cited in *Gnatholeptus*).

References: (ds) Blackwelder 1947: 782. (tx) Bright 1977: 513, 1981d: 17; Schedl 1937h: 168, 1938a: 183, 1979c: 241; Wood, S. L. 1979b: 135. *zeteki* Blackman 1942a: 226. Holotype ♀; Trinidad River, Panama; USNM, Washington. Synonymy: Bright 1977: 513.

References: (ds) Blackwelder 1947: 782. (tx) Blackman 1942a: 226; Bright 1977: 513; Wood, S. L. 1979b: 135.

subopacus Blackman 1942a: 210. Holotype ♂?; Chalco, Mexico, Mexico; USNM, Washington.

Figures: Blackman 1942a: figs. 41–42.

Distribution: North America (Durango, Mexico, Michoacan, Nuevo Leon, Oaxaca, Puebla, Tlaxcala, Veracruz in Mexico).

Hosts: *Pinus cembroides*, *P. leiophylla*, *P. michoacana*, *P. montezumae*, *Pinus* spp.

Notes: (3) Bright 1981d: 198 (treated as a subspecies of *seguis*). Wood 1982: 1086 (intergrades with *seguis*).

References: (ds) Blackwelder 1947: 782; Bright 1981d: 198; Rio Mora 1985: 11; Thomas, J. B. 1966; Wood, S. L. 1982b: 1086. (tx) Blackman 1942a: 210; Bright 1981d: 198; Rio Mora 1985: 11; Wood, S. L. 1982b: 1086.

elimatus Bright 1976c: 432. Holotype ♀; 85 km NW Oaxaca, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1982b: 1086.

References: (ds) Atkinson & Equihua 1988: 96; Bright 1981d: 196, 1986b: 683; Wood, S. L. 1982b: 1086. (tx) Bright 1976c: 432, 1981d: 196; McNamara 1984: 755; Wood, S. L. 1982b: 1086.

subsimilans Wood 1989: 179. Holotype, sex?; Brasilien, Rio Caraguata, Matto Grosso; Schedl Collection in NHMW, Wien, automatic.

Distribution: South America (Brazil).

References: (tx) Wood, S. L. 1989: 179.

subsimilis Schedl 1966f: 104 (*Breviophthorus*). Holotype, sex?; Brasilien, Rio Caraguata, Matto Grosso; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1955.

References: (cn) Schwerdtfeger 1956b: 48. (hb) Atkinson et al. 1986: 41; Bright 1981d: 301; Schwerdtfeger 1956b: 48, 1957c: 504. (ds) Atkinson et al. 1986: 41; Bright 1981d: 301; Schedl 1966f: 85, 1973d: 155; Wood, S. L. 1982b: 1136. (tx) Bright 1976b: 187, 1981d: 301, 345; Schedl 1955g: 25, 1966f: 104, 1979c:

244; Wood, S. L. 1977b: 210, 1982b: 1136, 1989: 179.

subsimilis Schedl 1955g: 25. Lectotype ♂; G. Ciudad [Guatemala City], Guatemala; Schedl Collection in NHMW, Wien, designated by Bright 1976b: 187.

Figures: Bright 1981d: 371.

Distribution: North America (Guatemala/ Chiapas, Oaxaca, Puebla in Mexico).

Hosts: *Pinus strobus* var. *chiapensis*, *P. tenuifolia*, *P.* sp.

References: (hb) Schwerdtfeger 1957: 504. (tx) Bright 1976b: 187, 1981d: 301, 371; Schedl 1955g: 18, 25, 1979c: 244.

subimpressus Bright 1976c: 441. Holotype ♀; 51 km S Valle Nacional, Oaxaca, Oaxaca; CNCI, Ottawa. Synonymy: Wood 1977b: 210, 1982b: 1136.

Notes: (3) Bright 1981d: 303 (treated as a good species). Wood 1982b: 1136 (this is only one of several minor variations of *subsimilis*).

References: (ds) Bright 1981d: 303. (tx) Bright 1976c: 441, 1981d: 303, 371; McNamara 1984: 758; Wood, S. L. 1977b: 210.

subsulcatus (Schedl) 1963d: 223 (*Breviophthorus*). Holotype, sex?; Brasilien: Parama, Rondon; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1963d: 223, 1979c: 244.

surinamensis Schedl 1961i: 226. Holotype ♀; Surinam, Moengo, Boven, Cottica R.; Cornell University Collection, Ithaca, New York.

Distribution: South America (Suriname).

References: (ds) Schedl 1960a: 78. (tx) Hoebeke 1978; Schedl 1961i: 226, 1964b: 206, 1979c: 247.

suspicious Bright 1972d: 89. Holotype ♀; Blue Mountain Peak, Jamaica, 7400 [feet]; CNCI, Ottawa.

Figures: Bright 1972: 75.

Distribution: Antilles Islands (Jamaica).

References: (ds) Bright 1985c: 176. (tx) Bright 1972: 75, 1972d: 89, 1985c: 176; McNamara 1984: 198.

suturalis Eggers 1932d: 291. Holotype, sex?; Congostaat (Haut-Ucle: Yebo Moto); MRCB, Tervuren. Distribution: Africa (Chana/ Zaïre).

Hosts: *Bussea occidentalis*.

References: (ds) Browne 1965a: 188; Schedl 1962h: 59, 1962j: 20. (tx) Eggers 1932d: 291; Schedl 1938a: 183, 1962j: 20; 1962k: 1080.

tenax Wood 1976a: 354. Holotype ♀; Volcan Pacaya, Esquintla, Guatemala, 1300 m; Wood Collection. Distribution: North America (Guatemala).

Hosts: Tree branch.

References: (ds) Bright 1981d: 74; Wood, S. L. 1982b: 1123. (tx) Bright 1981d: 74; Wood, S. L. 1976a: 354, 1982b: 1123.

- terebrans** Schedl 1970i: 223. Holotype, sex?; Brasilien, Amazonas; Schedl Collection in NHMW, Wien, automatic.
Distribution: South America (Brazil).
References: (tx) Schedl 1970i: 223.
- grandipennis** Schedl 1966f: 111. Holotype, sex?; Brasilien, Amazonas; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1965.
References: (ds) Schedl 1966f: 87. (tx) Schedl 1966f: 111, 1970i: 123, 1977l: 112, 1979c: 112.
- granulicanda** Schedl 1979c: 112. Holotype, sex?; Brasilien, Amazonas; Schedl Collection in NHMW, Wien, automatic.
Notes: (1) Because a synonym was available, this is an unneeded replacement name.
References: (tx) Schedl 1979c: 112.
- thamnis** Bright 1985a: 470. Holotype ♀; Mexico: Pachuca, Edo. Hidalgo; CNCI, Ottawa.
Distribution: North America (Hidalgo in Mexico).
Hosts: *Aluzaania angusta*.
References: (tx) Bright 1985a: 470.
- thomasi** Bright 1976c: 443. Holotype ♀; 17 km SW El Salto, Durango, Mexico, 2600 m; CNCI, Ottawa.
Distribution: North America (Durango in Mexico).
Hosts: *Pinus cooperi*.
References: (ds) Bright 1981d: 122; Wood, S. L. 1982b: 1031. (tx) Bright 1976c: 443, 1981d: 122; McNamara 1984: 758; Wood, S. L. 1982b: 1031.
- timidulus** Wood 1975b: 396. Holotype ♂; Volcan Chiriqui [near Cerro Punta], Chiriqui, Panama; Wood Collection.
Distribution: North America (Panama).
Hosts: Small tree (? *Orcopanax umbigenus*).
References: (ds) Bright 1981d: 96; Wood, S. L. 1982b: 1028. (tx) Bright 1981d: 96; McNamara 1984: 758; Wood, S. L. 1975b: 396, 1982b: 1028.
- timidus** Blandford 1904: 241. Holotype ♀; Mexico (Salle Coll.); BMNH, London.
Distribution: North America ("Mexico").
Notes: (3) Wood 1982b: 1138 (*dimidiatus* Blackman may be a junior synonym).
References: (ds) Blackwelder 1947: 782; Ferrer 1942; Hagedorn 1910d: 75; Kleine 1913b: 141, 1914b: 351; Wood, S. L. 1982b: 1138. (tx) Blandford 1904: 241; Bright 1981d: 328; Hagedorn 1910a: 101; Schedl 1930: 196, 1940a: 348; Wood, S. L. 1982b: 1138.
- togonus** Eggers 1920: 35. Syntypes, sex?; Kete Kratje in Togo; 4 syntypes in MNB, Berlin, 2 in Eggers Collection, in NHMW, Wien.
Distribution: Africa (Ghana/ Togo/ Zaïre).
Hosts: *Carapa grandiflora*, *C. procera*, *Khaya ivorensis*, *Parinari holstii*.
Notes: (1) Schedl 1979: 253 (citation of holotype invalid).
References: (ec) Schedl 1958d: 188. (hb) Schedl 1962j: 21. (ds) Schedl 1962j: 21. (tx) Eggers 1920: 35, 1932d: 291; Schedl 1935a: 183, 1954e: 47, 1962j: 21, 1962k: 1080, 1979c: 253.
- toralis** Wood 1964: 59. Holotype ♀; Beaver Creek, Logan Canyon, Utah [USA]; Wood Collection.
Figures: Bright 1981d: 346.
Distribution: North America (Alberta, British Columbia in Canada/ California, Idaho, Oregon, Utah, Wyoming in USA).
Hosts: *Pinus albicaulis*, *P. flexilis*, *P. monticola*, rare in *P. contorta*.
References: (cn) Lindgren 1980a: 69. (hb) Bright & Stark 1973: 105; Lindgren 1980a: 69. (ds) Bright 1976d: 184, 1981d: 133; Bright & Stark 1973: 105; Wood, S. L. 1982b: 1054. (tx) Bright 1977: 518, 1981d: 133, 346; McNamara 1977: 198; Wood, S. L. 1964: 59, 1982b: 1054.
- confusus** Bright 1966b: 295 (*Myeloborus*). Holotype ♀; Mt. Tallac, Eldorado Co., California [USA]; OSUC, Columbus, preoccupied by Blandford 1904. Synonymy: Bright & Stark 1973: 105.
References: (tx) Bright 1966b: 295, 1977: 518; Bright & Stark 1973: 105.
- collinus** Bright 1968c: 605. Holotype ♀; Terrace Mt., 25 km W Nahm, Osoyoos District, British Columbia [Canada]; CNCI, Ottawa. Synonymy: Bright 1977: 518.
References: (ds) Bright 1976d: 181. (tx) Bright 1968c: 605–606, 1976d: 181, 1977: 518; de Ruelle 1970: 108.
- torridus** Wood 1972d: 76. Holotype ♀; 10 km W High Rolls, Lincoln National Forest, New Mexico, 2000 m [USA]; Wood Collection.
Distribution: North America (New Mexico in USA).
Hosts: *Franseria* sp., probably *deltoides*.
References: (ds) Bright 1981d: 43; Wood, S. L. 1982b: 1117. (tx) Bright 1981d: 43; Wood, S. L. 1972d: 76, 1982b: 1117.
- tragardhi** Spessivtsev 1921a: 219. Syntypes, sex?; Hallnas in Norrland (Nord-Schweden); Spessivtsev Collection, not located.
Distribution: Asia (E USSR), Europe (Finland/ Norway/ Poland/ Sweden/ W USSR).
Hosts: *Picea abies*, *P. excelsa*, *P. obovata*.
References: (bv) Grune 1979: 121; Kangas 1980a: Kinelski 1958: 74. (cn) Esterberg 1959; Juutinen 1960: 8; Kangas 1980a; Kontkanen 1932: 61; Kozikowsky 1929: 253; Wardle 1929: 322. (ec) Kangas 1946b: 23; Karpinski 1932a: 103; Lundberg 1984; Nuorteva 1971; Saalas 1937: 154; Tragardh 1925a: 171. (hb) Karpinski 1933b: 30, 1955: 79; Karpinski & Postner 1974: 435; Stark 1952: 355; Strawinski 1948: 156; Spessivtsev 1921b: 219, 1923: 206, 1925d: 104, 1928a: 221; Stark 1952: 355. (ds) Bakke & Kvamme 1977; Butovitsch & Heqvist 1947; Esterberg 1959; Florov 1949: 97; Grune 1979: 121; Hansen, V. 1939; Hellen 1947; Horion 1951; Karpinski 1931: 18, 28, 1932a: 103, 290, 1933b: 30, 1945b: 231, 1955: 79; Karpinski & Strawinski 1948: 156;

Kinelski 1958: 74; Klefbeck & Sjöberg 1960: 230; Kleine 1934a: 161; Kontkanen 1932: 611; Krivolutskaia 1983; Kurenzov 1967; Lindberg & Saris 1952: 59; Lundberg 1981: 151; Numberg 1938: 126, 1954: 71, 1960b: 158–159; Nuorteva 1971: 67; Pfeiffer 1935: 158, 1976: 340; Postner 1974: 435; Stark 1931a: 26, 1931d: 548, 1952: 355; Yanovskii & Tegshzhargal 1985: 410; Zinovjev 1955: 187. **(tx)** Grune 1979: 121; Jansson 1936: 220; Karpinski & Strawinski 1948: 156; Numberg 1938: 126, 1954: 71; Pfeiffer 1927: 111–112, 1940b: 109, 112, 121, 1955a: 21, 1976: 340; Postner 1974: 435; Schedl 1934f: 1643; Sokanovskii 1929: 670–672, 1960: 677; Spessivtsev 1921a: 219–223, 1922a: 448, 488, 491, 1923: 206, 1925a: 183, 1925b: 27, 1925d: 104, 1926b: 50, 1929c: 298–303, 1931a: 65; Stark 1952: 355. **(ms)** Kozikowsky 1929: 253.

treculiae Schedl 1962k: 1080. Holotype ♂; Congo: Stanleyville; Schedl Collection in NHMW, Wien. Distribution: Africa (Zaire).

Hosts: *Treculia africana*.

References: **(hb)** Schedl 1962j: 22. **(ds)** Schedl 1962j: 22. **(tx)** Schedl 1962j: 22, 1962k: 1080, 1979c: 255.

trepidus Bright 1978: 83. Holotype ♀; Ukiah, California [USA]; USNM, Washington.

Distribution: North America (California in USA).

Hosts: *Pinus radiata*.

References: **(ds)** Bright 1981d: 208; Wood, S. L. 1982b: 1041. **(tx)** Bright 1978: 83, 1981d: 208; McNamee 1984: 758; Wood, S. L. 1982b: 1041.

trunculus Bright 1985a: 470. Holotype ♀; Mexico: Estacion de Biologia, Chamela, Edo. Jalisco; CNCI, Ottawa.

Distribution: North America (Jalisco in Mexico).

References: **(tx)** Bright 1985a: 470.

tuberculatus Eichhoff 1878b: 498. Holotype, sex?; California [USA]; Hamburg Museum, lost, neotype ♀; Black Hills, South Dakota [USA]; USNM, Washington, designated by Bright 1975: 72.

Figures: Kusch 1967: 10.

Distribution: North America (Alaska/ Alberta, British Columbia in Canada/ Baja California Norte, Coahuila, Durango, Nuevo Leon in Mexico/ Arizona, California, Colorado, Idaho, Montana, Nebraska, Nevada, New Mexico, Oregon, South Dakota, W Texas, Utah, Washington, Wyoming in USA).

Hosts: *Pinus aristata*, *P. attenuata*, *P. cembroides*, *P. contorta*, *P. coulteri*, *P. edulis*, *P. flexilis*, *P. jeffreyi*, *P. monophylla*, *P. ponderosa*, *P. sabiniana*, *P. strobiformis*, rare in *Picea engelmannii*, *P. pungens*.

Notes: (1) Schedl 1979c: 258 (invalid neotype designation). (3) Blackman 1928b: 92 (redescribed), p. 94 (aberration named as var. *australis*, nomen nudum, no status). Bright 1981d: 153 (treated as a subspecies of *pulehellus* on basis of host hybrid-

ization; however, there is little or no intergradation between beetle populations).

References: **(cn)** Blackman 1931c; Doane et al. 1936; Felt 1906: 753; Hopkins 1899: 443; Keen 1929: 53; Lindgren 1980a: 69; Ruppel 1967: 78; Schuder 1969: 77; Swaine 1918a: 96, 99. **(cc)** Blackman 1931c; Powell, J. M., Wong, & Melvin 1972: 15. **(hb)** Blackman 1931c; Bright & Stark 1973: 112; Chamberlin 1939: 384, 1958: 153; Doane et al. 1936; Keen 1929: 53; Lindgren 1980a: 69; Swaine 1918a: 96, 99. **(ds)** Atkinson & Equihua 1985c: 360, 1988: 98; Blatchley & Leng 1916: 635; Bright 1976d: 184, 1981d: 152; Bright & Stark 1973: 112; Chamberlin 1925, 1939: 384, 1958: 153; Elias 1982a, 1982b, 1983; Evans, D., Love, & Hunt 1978; Gast et al. 1989: 355; Hopkins 1893a: 132, 1893b: 209; Hopping 1922; Keen 1929: 53, 1929a: 33; Kleine 1913b: 141, 1914b: 391, 1934a: 161; Kusch 1967; Leng 1920: 341; McComb et al. 1953: 2; Patterson & Hatch 1945: 153; Ruppel 1967: 78; Schuder 1969: 77; Smith, G. S. 1930; Still, Tidsbury, & Melvin 1974a: 14; Swaine 1909: 140; Wood, S. L. 1948: 53, 1951a: 128, 1972a: 426, 1982b: 1092. **(tx)** Blackman 1928b: 92–94; Blatchley & Leng 1916: 635; Bright 1971b: 67–68, 1978: 73, 1981d: 152; Chamberlin 1939: 384, 1958: 153; Eichhoff 1878a: 388, 1878b: 498; Hagedorn 1910a: 101; Keen 1929a: 33; Kusch 1967: 10; Swaine 1909: 140, 1918a: 96, 99–102, 1925b: 193; Wood, S. L. 1951a: 128, 1972a: 426, 1982b: 1092.

rugicollis Swaine 1925b: 193. Holotype ♀; Glen Alpine, California [USA]; CNCI, Ottawa. Synonymy: Bright 1971b: 67.

References: **(ds)** Leng & Mutchler 1933: 52. **(tx)** Blackman 1928b: 93–94; Bright 1967b: 678, 1971b: 67; de Ruette 1970: 111; Swaine 1925b: 193–194.

tumidus Blackman 1928b: 58. Holotype ♀; Eureka, California [USA]; USNM, Washington.

Distribution: North America (N California in USA).

References: **(hb)** Bright & Stark 1973: 108; Chamberlin 1939: 369. **(ds)** Bright 1981d: 229; Bright & Stark 1973: 108; Chamberlin 1939: 369; Leng & Mutchler 1933: 52; Wood, S. L. 1982b: 1063. **(tx)** Blackman 1928b: 58; Bright 1981d: 229; Chamberlin 1939: 369; Wood, S. L. 1982b: 1063.

turbiculus Schedl 1938a: 187. Holotype, sex?; Brasilien; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: **(ds)** Blackwelder 1947: 782. **(tx)** Schedl 1938a: 187, 1966f: 86, 1979c: 259.

tutulus Bright 1986a: 643. Holotype ♀; Mexico: Jalapa, Veracruz; CNCI, Ottawa.

Distribution: North America (Veracruz in Mexico).

Hosts: *Rhus radicans*.

References: **(tx)** Bright 1986a: 643.

vegrandis Bright 1986a: 643. Holotype ♀; Mexico: Chetumal, Quintana Roo; CNCI, Ottawa.

- Distribution: North America (Quintana Roo in Mexico).
References: (tx) Bright 1986a: 643.
- venezuelensis** Schedl 1935d: 91. Holotype ♂: Venezuela; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
References: (ds) Blackwelder 1947: 782; Bright 1981d: 233; Wood, S. L. 1982b: 1059. (tx) Bright 1977: 518, 1981d: 233; de Ruelle 1970: 112; Schedl 1935d: 91, 1935a: 183, 1952k: 161, 1979c: 265; Wood, S. L. 1982b: 1059.
- venustus** Blackman 1928b: 75. Holotype ♀; Kaibab National Forest, Arizona [USA]; USNM, Washington.
Figures: Bright 1981d: 362.
Distribution: North America (British Columbia in Canada/ Arizona, California, Colorado, Idaho, Nebraska, New Mexico, South Dakota, Texas, Utah in USA).
Hosts: *Pinus aristata*, *P. contorta*, *P. coulteri*, *P. jeffreyi*, *P. leiophylla*, *P. monticola*, *P. ponderosa*, *P. strobiformis*.
References: (hb) Chamberlin 1939: 377, 1958: 152. (ds) Chamberlin 1939: 377, 1958: 152; Keen 1929a: 32; Kleine 1934a: 161; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 75; Bright 1977: 518, 1981d: 362; Chamberlin 1939: 377, 1958: 152; Keen 1929a: 32.
- artifex** Blackman 1928b: 76. Holotype ♀; Meyers, California [USA]; USNM, Washington. Synonymy: Bright 1977: 518.
References: (hb) Bright & Stark 1973: 110; Chamberlin 1939: 378. (ds) Bright & Stark 1973: 110; Chamberlin 1939: 378; Keen 1929a: 32; Kleine 1934a: 158; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 76; Bright 1977: 518; Chamberlin 1939: 378; Keen 1929a: 32; McNamara 1984: 754.
- vesculus** Wood 1978b: 401. Holotype ♀; Ft. Clayton, Canal Zone, Panama; Wood Collection.
Distribution: North America (Panama).
Hosts: Log phloem.
References: (ds) Bright 1981d: 35; Wood, S. L. 1982b: 1135. (tx) Bright 1981d: 35; Wood, S. L. 1978b: 401, 1982b: 1135.
- vespertinus** Bright 1978: 83. Holotype ♀; 37 km W Durango, Durango, Mexico; Wood Collection.
Distribution: North America (Durango in Mexico).
Hosts: *Pinus* sp.
References: (ds) Atkinson & Equihua 1988: 100; Bright 1981d: 119, 1986: 684; Wood, S. L. 1982b: 1058. (tx) Bright 1978: 83, 1981d: 119; McNamara 1984: 758; Wood, S. L. 1982b: 1058.
- viminalis** Bright 1977: 530. Holotype ♀; Cerro Potosí, Nuevo León, Mexico; CNCI, Ottawa.
Distribution: North America (Nuevo León in Mexico).
Hosts: *Pinus culminicola*.
References: (ds) Bright 1981d: 173; Wood, S. L. 1982b: 1096. (tx) Bright 1977: 530, 1981d: 173; McNamara 1984: 755; Wood, S. L. 1982b: 1096.
- virilis** Blackman 1928b: 143. Holotype ♀; Vermego, New Mexico [USA]; USNM, Washington.
Figures: Bright 1981d: 37, 335.
Distribution: North America (Chihuahua in Mexico/ Arizona, Colorado, S. Idaho, Missouri, New Mexico, Texas, Utah, Wyoming in USA).
Hosts: *Rhus trilobata*.
References: (hb) Chamberlin 1939: 402, 1958: 160. (ds) Bright 1981d: 36; Chamberlin 1939: 402, 1958: 160; Deyrup 1961b: 8; Kleine 1934a: 162; Leng & Mutchler 1933: 52; Wood, S. L. 1946: 55, 1951a: 128, 1960b: 69, 1972a: 427, 1982b: 1119. (tx) Blackman 1928b: 143, 1942a: 223; Bright 1977: 518, 1981d: 36–37, 335; Chamberlin 1939: 402, 1958: 160; de Ruelle 1970: 112; Wood, S. L. 1951a: 128, 1960b: 69, 1972a: 427; 1982b: 1119.
- fortis** Blackman 1928b: 142. Holotype ♀; Montell, Texas [USA]; USNM, Washington. Synonymy: Bright 1977: 518.
References: (hb) Chamberlin 1939: 402. (ds) Chamberlin 1939: 402; Kleine 1934a: 159; Leng & Mutchler 1933: 52. (tx) Blackman 1928b: 142, 1942a: 223; Bright 1977: 518; Chamberlin 1939: 402; de Ruelle 1970: 109.
- virtus** Schedl 1938a: 186. Holotype ♀; Venezuela; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
References: (ds) Blackwelder 1947: 782. (tx) Schedl 1938a: 186, 1979c: 268.
- volvulus** Schedl 1965f: 10. Holotype, sex?; D. Ost Afrika, Usambara Kwai; Schedl Collection in NHMW, Wien.
Distribution: Africa (Kenya/ Tanzania/ Uganda).
Hosts: *Acacia* sp.
References: (ds) Schedl 1962j: 23. (tx) Schedl 1962j: 23, 1965f: 10, 1979c: 268.
- vrydaghi** Nunberg 1973: 13. Holotype ♂; Congo Belge, Parc National Virunga; MRCB, Tervuren.
Distribution: Africa (Zaire).
References: (tx) Nunberg 1973: 13.
- woodi** Bright 1977: 531. Holotype ♀; 11.5 km W Kingston, New Mexico [USA]; CNCI, Ottawa.
Distribution: North America (New Mexico in USA).
Hosts: *Pinus edulis* twigs.
References: (ds) Bright 1981d: 109; Wood, S. L. 1982b: 1039. (tx) Bright 1977: 531, 1981d: 109; McNamara 1984: 758; Wood, S. L. 1982b: 1039.
- zexmenivora** Bright 1985a: 471. Holotype ♀; Mexico: Pachuca, Edo. Hidalgo; CNCI, Ottawa.
Distribution: North America (Hidalgo in Mexico).
Hosts: *Zexmenia* sp.
References: (tx) Bright 1985a: 471.
- zonalis** Bright 1976c: 443. Holotype ♀; Jerome, Arizona [USA]; USNM, Washington.
Distribution: North America (Nuevo León in Mexico/ Arizona in USA).

Hosts: *Pinus cembroides*, *P. ponderosa*.

References: (ds) Bright 1981d: 299; Wood, S. L. 1982b: 1111. (tx) Bright 1976c: 443, 1981d: 299; Wood, S. L. 1982b: 1111.

Corthylini: Corthylini

Corthyli

References: LeConte 1876: 346–347.

Corthylini

References: Wood, S. L. 1978a: 115, 1982b: 73, 1986a: 94, 98–100.

Gnathotrichina

References: Balachowsky 1949a: 241; Schedl 1958k: 144.

Genus *Gnathotrichus* Eichhoff

GNATHOTRICHUS EICHHOFF 1869a: 275. Type-species: *Gnathotrichus corthyloides* Eichhoff = *Tomicus materiarius* Fitch, monobasic.

Gnathotrichoides Blackman 1931b: 267. Type-species: *Cryphalus sulcatus* LeConte, subsequent designation by Wood 1982b: 1155. Synonymy: Wood 1982b: 1155. References: (tx) Blackman 1931b: 267; Wood, S. L. 1982b: 1155.

Ancyloderes Blackman 1938a: 205. Type-species: *Cryphalus pilosus* LeConte, original designation. Synonymy: Wood 1973c: 172. References: (hb) Bright & Stark 1973: 116. (ds) Bright & Stark 1973: 116. (tx) Blackman 1938a: 205–206; Schedl 1950i: 146; Wood, S. L. 1966b: 18, 1973c: 172.

Paraxyleborus Hoffmann 1942: 72. Type-species: *Xyleborus duprezi* Hoffmann = *Tomicus materiarius* Fitch, monobasic. Synonymy: Balachowsky 1949a: 241. References: (tx) Balachowsky 1949a: 241; Hoffmann 1942: 72, 1947: 47.

Prognathotrichus Bright 1972c: 1678. Type-species: *Prognathotrichus primus* Bright, original designation. Synonymy: Wood 1973c: 172. References: (tx) Bright 1972c: 1678; Wood, S. L. 1973c: 172.

Keys: Blackman 1931b: 266, Wood 1982b: 1156. Notes: (1) Blackman 1931b (revision), Wood 1982b: 1155 (revision).

References: (ay) Browne 1961d: 49; Nobuchi 1969a: 67; Schedl 1931a: 1–88. (bv) Chapman, J. A. 1957: 3–4, 1962b: 88; Chapman, J. A. & Kinghorn 1958: 368; Moeck 1970b: 993; Vite 1975b, 1976. (cn) Drake 1921: 201–205; Hagle et al. 1987: 39; Johnson 1958a: 508–511, 1958b: 238; Kinghorn 1961; Snyder 1927: 1–46; Swaine 1913: 87–92. (hb) Bright & Stark 1973: 117; Schedl 1981b: 79; Wood, S. L. 1982b: 1155, 1986a: 98. (ds) Bright & Stark 1973: 117; Patterson & Hatch 1945: 152; Richmond 1953: 86–89; Schedl 1981b: 79; Swaine 1909: 108; Wood, S. L. 1982b: 1155, 1986a: 98. (tx) Arnett 1960: 1045,

1968: 1045; Balachowsky 1949a: 241; Beal & Massey 1945: 60, 120; Blackman 1922b: 98, 1928: 6–16, 1931b: 264–276, 1942a: 178; Blandford 1904: 246; Bright 1976d: 174; Bright & Stark 1973: 117; Chamberlin 1939: 325–332, 1958: 137–140; Dodge 1938: 18; Eichhoff 1869a: 275, 1878b: 405; Hoffmann 1947: 47; LeConte 1876: 350; LeConte & Horn 1883: 517; Schedl 1935h: 342, 1938a: 158–168, 1939h: 45–48, 1951m: 125, 1955g: 4, 1958k: 144, 1981b: 79; Swaine 1918a: 45, 90–91; Wood, S. L. 1961: 48, 1967d: 37–57, 1973c: 172, 1982b: 1155–1166, 1986a: 98–99.

alniphagus Wood 1984: 115. Holotype ♀; Fierro del Toro, Morelos, Mexico; Wood Collection. Distribution: North America (Morelos in Mexico). Hosts: *Alnus firnifolia*.

References: (hb) Atkinson et al. 1986: 47. (ds) Atkinson & Equihua 1985a: 100; Atkinson et al. 1986: 47. (tx) Wood, S. L. 1984: 115.

consentaneus Blandford 1904: 247. Lectotype ♂; Totonicapán, Guatemala: BMNH, London, designated by Wood 1973c: 176.

Distribution: North America (Guatemala/ Chiapas in Mexico).

References: (ds) Blackwelder 1947: 782; Ferrer 1942; Hagedorn 1910d: 96; Kleine 1913b: 158, 1914b: 354. (tx) Blackman 1931b: 266; Blandford 1904: 247; Hagedorn 1910a: 147; Hopkins 1905b: 73; Schedl 1940a: 348, 1955g: 29, 1979c: 64; Wood, S. L. 1973c: 176.

deleoni Blackman 1942a: 227. Holotype ♂; Chalco, Distrito Federal, Mexico; USNM, Washington. Distribution: North America (Distrito Federal, Durango, Mexico, Tlaxcala in Mexico).

Hosts: *Pinus leiophylla*, *P. montezumae*, *P. spp.* Notes: (3) Breeds in shaded-out limbs 7 cm in diameter or larger of living trees.

References: (hb) Atkinson et al. 1986: 47; Burgos & Saucedo 1983: 129; Wood, S. L. 1982b: 1161. (ds) Atkinson & Equihua 1985a: 99, 1988: 90; Atkinson et al. 1986: 47; Blackwelder 1947: 782; Burgos & Saucedo 1983: 99; Dominguez & Carrillo 1976: 141; Wood, S. L. 1982b: 1161. (tx) Blackman 1942a: 227; Wood, S. L. 1967d: 47, 1982b: 1161.

dentatus Wood 1967d: 45. Holotype ♀; 29 km W El Salto, Durango, Mexico; Wood Collection.

Figures: Wood 1967d: 46, 1982b: 1219 (declivity). Distribution: North America (Durango, Michoacan in Mexico).

Hosts: *Quercus* sp.

References: (hb) Atkinson et al. 1986: 47; Burgos & Saucedo 1983: 128. (ds) Atkinson & Equihua 1985a: 98; Atkinson et al. 1986: 47; Burgos & Saucedo 1983: 128; Wood, S. L. 1982b: 1158. (tx) de Ruette 1970: 101; Wood, S. L. 1967d: 45–46, 1982b: 1158.

denticulatus Blackman 1931b: 270. Holotype ♀; Cloudercroft, New Mexico [USA]; USNM, Washington.

Distribution: North America (Chihuahua, Distrito Federal, Durango, Guerrero, Michoacan, Puebla, San Luis Potosi in Mexico/ Arizona, Colorado, SE Montana, New Mexico, South Dakota, W Texas, E Utah in USA).

Hosts: *Pinus ponderosa*, *P. strobiformis*, *P. spp.*, rare in *Abies concolor*, *Picea engelmannii*.

References: (cn) Blackman 1931c: 35–36; Chamberlin 1939: 328–329; Keen 1938: 146; Schuder 1969: 74. (ce) Blackman 1931c: 35–36; Chamberlin 1939: 328–329; Keen 1938: 146; Massey 1961: 358. (hb) Blackman 1931c: 35–36; Chamberlin 1939: 328–329; Keen 1938: 146; Wood, S. L. 1982b: 1163. (ds) Atkinson & Equihua 1985c: 361; Chamberlin 1939: 328–329; Furniss, R. L. & Carolin 1977: 382; Gast et al. 1989: 354; Keen 1938: 146; Leng & Mutchler 1933: 94; Schuder 1969: 74; Snow 1883: 44, 1906: 175, 1907: 185; Wood, S. L. 1982b: 1163. (tx) Blackman 1931b: 270; Chamberlin 1939: 328–329; Hoffmann 1947; Wood, S. L. 1982b: 1163.

imitans Wood 1967d: 48. Holotype ♂; 5 km W El Salto, Durango, Mexico; Wood Collection.

Distribution: North America (Durango, Hidalgo, Mexico in Mexico/ Arizona, New Mexico in USA).

Hosts: *Pinus ponderosa*, *P. sp.*

References: (hb) Wood, S. L. 1982b: 1164. (ds) Furniss, R. L. & Carolin 1977: 382; Wood, S. L. 1982b: 1164. (tx) de Ruelle 1970: 101; Wood, S. L. 1967d: 48–49, 1982b: 1164.

materiaris (Fitch) 1858: 726 (*Tomicus*). Holotype ♀; New York [USA]; USNM, Washington.

Figures: Balachowsky 1949a: 231, 243, Blackman 1922b: pl. 9, fig. 49, Brakman 1966b: 52, Bright 1976d: 204, 213, Dillon & Dillon 1961: 801, Cover et al. 1980: 21, Joly 1976b: 148, Postner 1974: 438.

Distribution: Antilles Islands (Dominican Republic), Europe (introduced: France/ Germany/ Netherlands), North America (New Brunswick, Nova Scotia, Ontario, Quebec in Canada/ Alabama, Arkansas, District of Columbia, Florida, Georgia, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, Oklahoma, Pennsylvania, South Carolina, South Dakota, E Texas, Virginia, Vermont, West Virginia, Wisconsin in USA).

Hosts: *Pinus echinata*, *P. ponderosa*, *P. strobus*, *P. taeda*, *Picea spp.*, *Larix sp.*, *Abies sp.*, *Tsuga sp.*

Notes: (3) Blackman 1931b: 268 (redescribed).

References: (ay) Hopkins 1894g; Kaston 1936: 645; Schedl 1931a: 6–11; Thomas, J. B. 1957: 3, 1967. (bv) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 268; Barr, B. A. 1969: 641; Clemens 1916; Dixon & Payne 1979b; Grune 1979: 123;

Luitjes 1976; Turnbow & Franklin 1980. (cn) Anonymous 1962m; Balch 1945; Beal et al. 1952; Becker, W. D. 1955d, 1959b: 173–174, 1962, 1964c, 1966c; Becker, W. D. & Abbott 1956: 664–666, 1960: 46–47, 1961: 367, 1966; Blackman 1950; Chamberlin 1924, 1939: 327–328; Clemens 1916: 297; Connola et al. 1956: 1–36; Currie 1905: 70; Doane et al. 1936; Doom 1963: 420–423, 1967c; Felt 1901: 495–496, 1926: 247–248, 276, 1930a: 247–248, 276; Felt & Rankin 1932: 408; Fisher, Thompson, & Webb 1953, 1954b: 3–15; Fitch 1858; Garman 1905: 69; Gause 1968: 454–456; Grandi 1951; Herrick 1935: 138; Hopkins 1894: 277, 1899c: 383, 434, 442, 1902c: 14, 1904a: 15; Hubbard 1897b: 30; Jablonskoff 1953: 325; Joly 1976; Packard 1890: 718–720; Pierson 1927: 110; Rodary 1959: 850; Smith, J. B. 1901b: 92; Snyder 1927: 7; Swaine 1918a: 91; Swaine & Craighead 1924: 1–27. (ce) Batra 1963a: 481–487, 1963b: 216; Batra & Michie 1963: 477; Chamberlin 1939: 327–328; Dixon & Payne 1979b; Felt 1906: 371; Hines, J. W. & Heikkinen 1977; Hirschmann & Wisniewski 1982, 1983; Hurlbutt 1967; Luitjes 1976; Morley 1939: 244; Ogaz Ituarte & Cibrián 1980; Rodary 1959: 850; Roeper & French 1981; Rumbold 1931c: 849; Steinhaus 1946: 404; Tomalak, Welch, & Galloway 1989b: 27; Wichmann 1955a: 93, 103; Woodring & Moser 1970. (hb) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 268; Batra 1963b: 221; Beal & Massey 1945: 120–121; Blackman 1919a, 1950; Blandford 1894b; Chamberlin 1939: 327–328; Chittenden 1890, 1899a; Clemens 1916; Doane et al. 1936; Drooz 1985: 372; Felt 1906: 371, 1926: 247–248, 276, 1930a: 247–248, 276; Felt & Rankin 1932: 408; Fitch 1858; Grandi 1951; Herrick 1935: 135; Hopkins 1894g, 1899c: 383, 442, 1904a: 15; Hubbard 1897b: 30; Joly 1968: 508–511, 1976; Luitjes 1976; Morley 1939: 244; Ogaz Ituarte & Cibrián 1980; Ostmark 1968; Packard 1890: 718; Pierce, W. D. 1907: 290; Pierson 1927: 110; Postner 1974: 437; Schwarz 1888a: 80; Skinner 1905: 248; Swaine 1911b: 85, 1918a: 91; Wolcott & Montgomery 1933: 165; Wood, S. L. 1982b: 1160. (ds) Anonymous 1926c: 518, 1962m; Atkinson et al. 1991: 162; Bain 1974: 15; Baker, W. L. 1972: 268; Balachowsky 1949b; Beal & Massey 1945: 120–121; Beaulne 1956; Blackman 1950; Blandford 1894b; Blatchley & Leng 1916: 643; Brakman 1966a: 51, 1966b: 207; Bright 1976d: 176, 1985c: 176; Chamberlin 1925, 1939: 327–328; Chittenden 1890, 1899a: 55–61; Cockerell et al. 1907; Currie 1905; DeLeon 1942a; Deyrup 1981: 8; Dodge 1938; Doom 1967c: 143; Drooz 1985: 372; Fall & Cockerell 1907: 217; Felt 1906: 370, 1924: 276, 1926: 247–248, 276, 1930a: 247–248, 276; Felt & Rankin 1932: 408; Gauss 1968; Gemminger & Harold 1872: 2688; Glick 1939: 37; Grune 1979: 123; Hagedorn 1910d: 96; Hamilton 1895a: 346, 378; Harde 1967; Henshaw 1885: 147;

- Hoffmann 1956: 91; Hopkins 1893a: 128, 1893b: 208, 1902c: 14; Houghton 1905; Howden & Vogt 1951; Hubbard & Schwarz 1878a: 665; Jablonsky 1953: 325; Joly 1968, 1976; Kamp 1979; Kirk 1970; Kleine 1913b: 158, 1914b: 393, 1934a: 171; Leng 1920: 340; Leonard 1928: 518; Licht 1957: 278; MacGillivray, A. & Houghton 1902; Menier 1972; Morgan, Anne, Morgan, & Elias 1985: 1821; Ostmark 1968; Postner 1974: 437; Proctor 1946: 207; Provancher 1877: 566; Rodary 1959: 550; Saint-Albin 1949: 2; Schedl 1966d: 118, 1981b: 79; Schiender 1985; Schiender & Niedersachs 1985; Schwarz 1878d: 468, 1886: 40, 44, 1888a: 50; Smith, J. B. 1910: 401; Swaine 1909: 108, 1911b: 85; Turnbow & Franklin 1980; Ulke 1902: 56; Wichmann 1955a: 93, 103; Wickham 1896a: 309, 1896c: 168; Wolcott & Montgomery 1933: 165; Wood, S. L. 1977a: 70, 1982b: 1160. **(tx)** Balachowsky 1948: 140, 1949a: 242; Beal & Maasey 1945: 120–121; Benoît 1985: 119; Blackman 1922b: 95–96, 1931b: 264–268; Blandford 1894b; Blatchley & Leng 1916: 643; Brakman 1966b: 52; Bright 1976d: 176, 204, 213, 1985c: 176; Chamberlin 1939: 327–328; Clemens 1916; Dillon & Dillon 1961: 801, 811; Dodge 1938: 13; Eichhoff 1869a: 275, 1878b: 406; Fitch 1858: 726; Goyer et al. 1980: 21; Grune 1979: 123; Hagedorn 1910: 147; Hoffmann 1947: 47, 1956: 91; Hopkins 1905b: 73; Hubbard 1897b: 30; Jacques 1951: 353; Joly 1976, 1976b: 148; LeConte 1868: 155, 1876: 350, 1878a: 468; Licht 1957: 278; Postner 1974: 437; Provancher 1877: 566; Schedl 1931a: 6–7, 1981b: 79; Schwarz 1886: 40, 44; Swaine 1909: 108–109, 1918a: 91; Thomas, J. B. 1957: 3, 1967; Titus, Meikle, & Harrison 1985: 71; Wood, S. L. 1982b: 1160; Zimmermann 1868: 143. **(ms)** Eckstein 1900c.
- corthyloides* Eichhoff 1869a: 275. Syntypes, sex²; Carolina [USA]; Hamburg Museum, lost. Synonymy: Eichhoff 1878b: 406. References: **(ay)** Schedl 1931a: 7. **(ds)** Leng 1920: 340; Swaine 1909: 108. **(tx)** Blackman 1931b: 264–268; Eichhoff 1869a: 275, 1878b: 406; Hopkins 1914: 122; LeConte 1876: 350; Schedl 1931a: 7; Swaine 1909: 108, 1918a: 91.
- duprezi* Hoffmann 1936: 43. Holotype ♀; Forêt du Ronvay, Seine-Inferieure, France; MNHN, Paris. Synonymy: Balachowsky 1949a: 242. References: **(ds)** Duprez 1938a; Hoffmann 1936; Sainte-Claire & Mequignon 1938: 449. **(tx)** Balachowsky 1948: 140, 1949a: 242; Hoffmann 1936: 43, 1942: 72, 1947: 47; Wood, S. L. 1982b: 1160.
- nimifrons* Wood 1967d: 47. Holotype ♀; 29 km W El Salto, Durango, Mexico; Wood Collection. Distribution: North America (Durango, Puebla in Mexico/ Arizona in USA). Hosts: *Quercus* sp. References: **(hb)** Wood, S. L. 1982b: 1159. **(ds)** Atkinson & Equihua 1985a: 99; Furniss, R. L. & Carolin 1977: 382; Wood, S. L. 1982b: 1159. **(tx)** Wood, S. L. 1967d: 47, 1982b: 1159.
- nitidifrons* Hopkins 1905b: 72. Holotype ♀; Michoacan, Mexico; USNM, Washington. Distribution: North America (Guatemala/ Durango, Hidalgo, Mexico, Michoacan in Mexico). Hosts: *Pinus cooperi*, *P. leiophylla*, *P. montezumae*. References: **(ay)** Thomas, J. B. 1967. **(hb)** Wood, S. L. 1982b: 1161. **(ds)** Atkinson & Equihua 1985a: 99, 1985c: 361, 1988: 90; Blackwelder 1947: 752; Ferrer 1942; Hagedorn 1910d: 97; Kleine 1913b: 158, 1914b: 354, 1934a: 171; Wood, S. L. 1982b: 1161. **(tx)** Blackman 1931b: 266; Hagedorn 1910a: 147; Hopkins 1905b: 72; Schedl 1940a: 348, 1948f: 268, 1979c: 169; Thomas, J. B. 1967.
- obscurus* Wood 1974a: 55. Holotype ♀; 9 km NE Teziutlan, Puebla, Mexico; Wood Collection. Distribution: North America (Puebla, Tlaxcala in Mexico). Hosts: *Quercus* sp. References: **(hb)** Wood, S. L. 1982b: 1158. **(ds)** Wood, S. L. 1974a: 55, 1982b: 1158. **(tx)** Wood, S. L. 1974a: 55, 1982b: 1158.
- omissus* Wood 1974a: 57. Holotype ♀; Volcan Irazu, Cartago, Costa Rica; Wood Collection. Distribution: North America (Costa Rica/ Guatemala). Hosts: *Orcopanax umbigenus*. References: **(hb)** Wood, S. L. 1982b: 1165. **(ds)** Atkinson & Equihua 1988: 90; Wood, S. L. 1982b: 1165. **(tx)** McNamara 1984: 753; Wood, S. L. 1974a: 56–57, 1982b: 1165.
- perniciosus* Wood 1967d: 47. Holotype ♂; 9 km S Carapan, Michoacan, Mexico; Wood Collection. Distribution: North America (Honduras/ Chiapas, Chihuahua, Distrito Federal, Jalisco, Michoacan, Morelos in Mexico). Hosts: *Pinus chiapensis*, *P. leiophylla*, *P. montezumae*, *P. oocarpa*, *P. pseudostrobus*, *P. spp.* References: **(hb)** Atkinson et al. 1986: 49; Burgos & Saucedo 1984: 129; Wood, S. L. 1982b: 1162. **(ds)** Atkinson & Equihua 1985a: 99, 1985c: 361; Atkinson et al. 1986: 49; Burgos & Saucedo 1984: 129; Furniss, R. L. & Carolin 1977: 382; Schedl 1977e: 42; Wood, S. L. 1982b: 1162. **(tx)** Wood, S. L. 1967d: 47–48, 1982b: 1162.
- pilosus* (LeConte) 1868: 156 (*Cryphalus*). Lectotype, sex²; middle California [USA]; MCZ, Cambridge, designated by Wood 1982b: 1160. Distribution: North America (Durango in Mexico/ Arizona, S California, S Nevada, New Mexico in USA). Hosts: *Quercus* sp. limbs (Keen 1929a: 34); commonly collected at light. Notes: (1) Blackman 1938: 204 (*pilosulus* an invalid spelling). Wood 1973: 172 (to *Gnathotrichus*). (3) Blackman 1938a: 205 (re-described, cited in *Ancyloderes*). References: **(cn)** Swaine 1918a: 94. **(hb)** Bright &

- Stark 1973: 116; Chamberlin 1939: 324–325; Swaine 1918a: 94. (**ds**) Blackwelder 1939; Bright & Stark 1973: 116; Chamberlin 1939: 324–325; Gemminger & Harold 1872: 2688; Hagedorn 1910d: 73; Henshaw 1882: 268, 1885: 147; Keen 1929c: 34; Kleine 1914b: 391; Leng 1920: 340; Snow 1906: 175; Swaine 1909: 136; Wood, S. L. 1982b: 1160. (**tx**) Blackman 1921: 148, 1931a: 225, 1935a: 205–206; Chamberlin 1939: 324–325; Hagedorn 1910a: 101; Keen 1929: 34; LeConte 1868: 154, 156, 1876: 351; Swaine 1909: 136, 1918a: 94; Wood, S. L. 1966b: 18–21, 1982b: 1162. *saltoni* Blackman 1935a: 206. Holotype, sex?, Flagstaff, Arizona [USA]: USNM, Washington. Synonymy: Wood 1966b: 21. References: (**ds**) Blackwelder 1939; Chamberlin 1939: 325. (**tx**) Blackman 1935a: 206; Chamberlin 1939: 325; Wood, S. L. 1966b: 21.
- primus** (Bright) 1972c: 1678 (*Prognathotrichus*). Holotype ♂; Mt. Tzonteluitz, Chiapas, Mexico; CNCI, Ottawa. Figures: Bright 1972c: 1678. Distribution: North America (Chiapas in Mexico). Hosts: Probably *Quercus* sp. References: (**ds**) Wood, S. L. 1982b: 1158. (**tx**) Bright 1972c: 1678; McNamara 1977: 198; Wood, S. L. 1973c: 172, 1982b: 1158.
- retusus** (LeConte) 1868: 155 (*Cryphalus*). Lectotype ♀; California [USA]: MCZ, Cambridge, designated by Wood 1982b: 1163. Figures: Blackman 1931b: 270. Distribution: North America (British Columbia in Canada/ Baja California Norte in Mexico/ California, Idaho, Montana, Nevada, Oregon, NW Utah, Washington in USA). Hosts: *Alnus* sp. (Oregon, Washington only), *Picea engelmannii*, *Pinus contorta*, *P. flexilis*, *P. jeffreyi*, *P. lambertiana*, *P. radiata*, *P. ponderosa*, *Populus trichocarpa* (accidental), *Pseudotsuga menziesii*, *Abies concolor*, *A. magnifica*, *Tsuga heterophylla*. Notes: (3) Blackman 1931b: 271 (*occidentalis* Hopkins, nomen nudum, synonymy). References: (**ay**) Farris 1963: 257; Farris & Funk 1965: 527; Hopkins 1894g; Schedl 1931a: 11; Schneider 1976; Schneider & Rudinsky 1969a, 1969b; Thomas, J. B. 1967. (**bv**) Barr, B. A. 1969: 641; Borden 1984; Borden, Lindgren, & Chong 1980; Borden & McLean 1979; Borden et al. 1980, 1981; Buttrick 1912: 457; Chapman 1963: 675, 1966: 54; Inscoc 1982; Jacobson 1965, 1972; Johnston, B. D. & Slessor 1979; Klassen, Ridgway, & Inscoc 1982; Lindgren 1983; Lindgren et al. 1982, 1983; Liu & McLean 1989; Moeck, Wood, & Lindahl 1981; Nijholt 1980; Ohmart & Voigt 1982: 340; Rudinsky 1966b: 356–361, 1966c: 218–219; Rudinsky & Daterman 1964: 1339; Rudinsky & Schneider 1969a: 1248–1255; Schneider & Rudinsky 1969a: 27–39, 1969b: 995–1003; Shore & McLean 1983: 3, 1985; Zhong & Schowalter 1989. (**cn**) Chamberlin 1924, 1939: 328–329; Essig 1926: 519, 1958: 519; Fall & Cockerell 1907: 217; Felt 1906: 752; Graham 1953; Gray & Borden 1985; Hatch 1938: 194; Hopkins 1902: 14, 1904a: 15; Johnson 1958a: 508; Keen 1929c: 56, 1935: 146; Lindgren 1980a: 63, 1983; Lindgren & Borden 1983; Lindgren et al. 1983; McLean & Borden 1977b; McMullan 1956: 34; Molnar et al. 1969: 117, 1970: 104; Molnar, Ross, & Fiddick 1971: 81; Nijholt 1978a: 1980; Prebble 1944: 50, 1954: 220; Prebble & Graham 1957: 90–112; Ruppel 1967: 47; Shore 1985; Shore & McLean 1985; Silver & Ross 1958: 79, 1960: 98; Stevens, Brewer, & Leatherman 1980: 27; Swaine 1914: 33, 37, 40, 1918a: 90, 91. (**ec**) Borden et al. 1981; Chamberlin 1918a, 1939: 328–329; Chate-lain & Schenk 1983; Deyrup & Gara 1978: 274; Farris 1963; Farris & Funk 1965: 527; Felt 1906: 751; Keen 1935: 146; McLean & Borden 1977; Nakashima 1975: 4; Nijholt 1980, 1981; Ohmart & Voigt 1982: 340; Otvos 1977; Roeper & French 1981; Rudinsky & Schneider 1969a; Schneider 1976; Shore 1985; Steinhaus 1946: 404; Stephen & Dahlsten 1976b: 292. (**hb**) Blandford 1894b; Bright & Stark 1973: 117; Chamberlin 1939: 328–329, 1958; Essig 1926: 519, 1958: 519; Felt 1906: 751; Gast et al. 1989: 385; Hopkins 1894g, 1904a: 15; Johnson 1958a: 508, 1958b: 237; Keen 1929c: 56, 1938: 146; Kimmey 1943: 22; Lindgren 1980a: 63; Nijholt 1978a; Rudinsky & Schneider 1969a; Shore 1985; Stevens, Brewer, & Leatherman 1980: 27; Swaine 1918a: 90–91. (**ds**) Blandford 1894b; Bright 1976d: 176; Bright & Stark 1973: 117; Chamberlin 1917: 325, 1918a: 39, 1925, 1939: 328–329, 1958: 138–139; Chapman 1963: 675; Cockerell et al. 1907; Essig 1926: 519, 1958: 519; Evans, D. 1983: 32, 1985; Furniss, M. M. & Furniss 1972; Furniss, R. L. & Carolin 1977: 381; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 97; Henshaw 1882: 268, 1885: 147; Hopkins 1893a: 128, 1893b: 208, 1894: 277; Hopping 1922; Keen 1929a: 35, 1929c: 56–57, 1935: 146, 1949a: 93; Kleine 1913b: 158, 1914b: 399, 1934a: 171; Lange 1937: 174; Leng 1920: 340; Marchant & Borden 1976; McComb et al. 1953: 4; Miller, S. E. 1983: 99; Murayama 1957a: 37; Nijholt 1980; Ohmart & Voigt 1982: 340; Patterson & Hatch 1945: 152; Rudinsky & Schneider 1969a: 1248–1255; Ruppel 1967: 47; Schedl 1972a: 144; Shore 1985; Smith, C. S. 1930; Swaine 1909: 108, 1914: 33; Wickham 1896a: 309; Wood, S. L. 1951a: 128, 1972a: 428, 1977d: 512, 1982b: 1163. (**tx**) Blackman 1931b: 270–271; Blandford 1894b; Bright 1976d: 176; Chamberlin 1939: 328–329, 1958: 138–139; Eichhoff 1878b: 407, 511; Evans, D. 1983: 32; Hagedorn 1910a: 147; Hopkins 1905b: 73; Johnson 1958b: 237; Keen 1929a: 35; LeConte 1868: 155, 1876: 350; Schedl 1931a: 11, 1935a: 160, 164; Swaine 1909: 108, 1918a: 90–91; Thomas, J. B. 1967; Wood, S. L. 1972a: 428,

- 1952b: 1163. (**ms**) Hatch 1938: 194; Johnston, B. D. & Slessor 1979; Lindgren B. et al. 1982; Prebble 1944: 50.
- alni* Blackman 1931b: 271. Holotype ♀; Hoquiam, Washington [USA]; USNM, Washington. Synonymy: Wood 1977d: 512.
- Notes: (3) The closest relative of *retusus* is *alniplagus* Wood; perhaps the original host of *retusus* was *Alnus* and the adaptation to conifers was secondary, but the original habit persisted in parts of Oregon and Washington (SLW).
- References: (**cn**) Chamberlin 1939: 330; Hatch 1935: 194; Keen 1938: 146; Kinghorn 1957b: 213; Ruppel 1967: 47. (**cc**) Chamberlin 1939: 330; Keen 1938: 146. (**hb**) Chamberlin 1939: 330, 1958: 139; Keen 1938: 146. (**ds**) Chamberlin 1939: 330, 1958: 139; Furniss, R. L. & Carolin 1977: 382; Keen 1938: 146; Leng & Mutchler 1933: 94; Patterson & Hatch 1945: 152; Ruppel 1967: 47. (**tx**) Blackman 1931b: 271; Chamberlin 1939: 330, 1958: 139; Schedl 1979c: 16; Wood, S. L. 1977d: 512.
- sulcatus* (LeConte) 1868: 155 (*Cryphalus*). Holotype ♀; middle California [USA]; MCZ, Cambridge.
- Figures: Atkinson et al. 1986: 48, Blackman 1931b: 273 (antenna), Bright & Stark 1973: 163, Furniss & Carolin 1977: 391, Schwerdtfeger 1960b: 254–257, Shore 1985.
- Distribution: North America (British Columbia in Canada/ El Salvador/ Guatemala/ Honduras/ Distrito Federal, Durango, Hidalgo, Michoacan, Puebla, Tlaxcala, Veracruz in Mexico/ Arizona, California, Colorado, Idaho, Nevada, New Mexico, Oregon, South Dakota, Utah, Washington in USA).
- Hosts: *Abies concolor*, *A. magnifica*, *A. religiosa*, *Pinus leiophylla*, *P. montezumae*, *P. ponderosa*, *P. pseudostrobus*, *P. spp.*, *Pseudotsuga menziesii*, *Tsuga heterophylla*.
- Notes: (3) Blackman 1931b: 274 (re-described).
- References: (**ay**) Farris 1963: 257; Farris & Funk 1965: 527; Schedl 1931a: 13; Schneider 1976; Schneider & Rudinsky 1969a, 1969b; Thomas, J. B. 1967. (**bv**) Barr, B. A. 1969: 641; Borden 1984; Borden, Lindgren, & Chong 1980; Borden & McLean 1979; Borden & Stokkink 1973; Borden et al. 1976, 1980: 108, 1981; Byrne et al. 1974; Cade 1970, 1971; Cade, Hnutford, & Gara 1970: 1014–1015; Chapman 1966: 55; Gagne & Kearby 1978: 1012; Grossman 1988: 9; Inscoc 1982; Inscoc & Beroza 1976: 147; Jacobson 1965, 1972; Johnson 1960b: 4; Johnson, Wright, & Orr 1961: 8; Kinghorn & Chapman 1957: 48; Klassen, Ridgeway, & Inscoc 1982; Lindgren 1983, 1984; Lindgren et al. 1983; Lin & McLean 1989; McLean 1976b, 1980a, 1988; McLean & Borden 1975a, 1975b, 1977a, 1977b, 1979; Moeck 1970b: 993, 1971; Mori 1975c; Nijholt 1980; Oda 1974: 7; Payne & Wood 1981: 492; Plummer et al. 1976; Prebble & Graham 1957: 93; Renwick, Vite, & Billings 1977; Roelofs 1978; Rudinsky 1966b: 356–361, 1966c: 218–219; Rudinsky & Daterman 1964: 1339; Rudinsky & Schneider 1969a: 1248–1255; Schneider & Rudinsky 1969a: 995–1003, 1969b: 27–39; Schuler & Slessor 1977; Shore & McLean 1983; Silverstein 1974; Silverstein & Young 1976: 20; Wood, D. L. 1978, 1979a, 1980b; Wright, R. H. 1974: 134; Zammicio 1981b, 1983; Zhong & Schowalter 1989. (**cn**) Atkins 1976; Boocock 1959d; Boyce 1929; Buttrick 1912: 457; Chamberlin 1915: 40, 1924, 1939: 331–332; Currie 1905: 70; Doane et al. 1936; Essig 1926: 519, 1958: 519; Fiddick & Van Sickle 1980: 20; Fisher, Thompson, & Webb 1953, 1954b: 3–15; Graham 1953, 1963; Graham et al. 1950: 43–44, 48; Gray 1985; Hatch 1935: 194; Hopkins 1902c: 14, 1904a: 15; Hopping & Jenkins 1933: 1–14; Janssen 1955: 10–14; Jardine 1969: 48; Johnson 1958a: 508; Keen 1929c: 56, 1938: 144, 146; Kinghorn 1957b: 213; Lindgren 1980a: 63, 1983, 1984; Lindgren & Borden 1982; Lindgren et al. 1983; Mathers 1935: 14; McBride, C. F. 1950: 46; McBride, C. F. & Kinghorn 1960: 40–52; McLean 1976b, 1980a, 1980b, 1985, 1988; McLean & Borden 1975a, 1975b, 1977a, 1977b, 1979; McMullan 1956: 34–43; Nijholt 1978a, 1980; Ostaff 1974; Payne & Wood 1981: 492; Prebble 1944: 50, 1954: 220; Roelofs 1978; Rudinsky, Terriere, & Allen 1960: 949; Ruppel 1967: 47; Schuler 1969: 74; Schuler & Slessor 1977; Schwerdtfeger 1960b; Shea & Johnson 1962: 5; Shore & McLean 1983; Stevens, Brewer, & Leatherman 1980: 27; Struble 1937d: 11, 1957: 14; Swaine 1914: 37–38, 1918a: 91; Thatcher, T. O. 1961; Wood, D. L. 1979a, 1980b. (**cc**) Becker 1953: 361, 1955; Borden et al. 1981; Cade 1970, 1971; Chamberlin 1918a, 1939: 331–332; Deyrup & Gara 1978: 275; Farris 1963; Farris & Funk 1965: 527; Funk, A. 1970: 1445–1448, 1973; Keen 1938: 144, 146; Kinghorn 1956: 4, 1957a: 3; McLean & Borden 1977a, 1977b; Nakashima 1975: 4; Nijholt 1980; Ostaff 1974; Roeper & French 1981; Rudinsky & Schneider 1969a; Rumbold 1931c: 849; Schneider 1976; Struble 1957: 14; Webb 1945: 71; Wichmann 1955a: 93, 103. (**hb**) Atkinson & Equihua 1985a: 100; Atkinson et al. 1986: 49; Becker, G. 1953, 1955; Bright 1976d: 174; Bright & Stark 1973: 118; Burgos & Saucedo 1983: 130; Cade 1970, 1971; Chamberlin 1939: 331–332, 1958: 139–140; DeLeon 1952: 78–79; Doane & Gilliland 1929: 920–921; Doane et al. 1936; Essig 1926: 519, 1958: 519; Graham 1963; Hopkins 1904a: 15; Johnson 1958a: 508–511, 1958b: 237; Kalshoven 1962: 12; Keen 1929c: 56, 1938: 144, 146; Kimmey & Furniss 1943: 22–23; Lindgren 1980a: 63; McLean 1980a, 1988; McLean & Bennett 1978; McLean & Borden 1977; Nijholt 1978a, 1983; Ostaff 1974;

Pierce, W. D. 1907: 290; Prebble & Graham 1957: 93–112; Rudinsky & Schneider 1969a; Schwerdtfeger 1960b: 254; Stevens, Brewer, & Leatherman 1980: 27; Struble 1937d: 11, 1957: 14; Swaine 1918a: 91; Thatcher, T. O. 1983; Zaniccio 1981b, 1983. (**ds**) Atkinson & Equihua 1985a: 100, 1985c: 361, 1988: 90; Atkinson et al. 1986: 49; Bain 1974: 15; Becker, G. 1953; Blackwelder 1947: 782; Bright 1976d: 174; Bright & Stark 1973: 118; Burgos & Saucedo 1983: 130; Chamberlin 1917: 325, 1918a, 1925, 1939: 331–332, 1958: 139–140; Choo & Woo 1983; Cockerell et al. 1907; Currie 1905: 70; DeLeon 1952; Dominguez & Carrillo 1976: 141; Essig 1926: 519, 1958: 519; Evans, D. 1983; Fall & Cockerell 1907: 217; Furniss, M. M. 1978; Furniss, R. L. & Carolin 1977: 381; Gemminger & Harold 1872: 2689; Hagedorn 1910d: 97; Henshaw 1882: 269; Hopkins 1902c: 14; Hopping 1922; Keen 1929a: 35, 1926c: 56, 1938: 144, 146; Kleine 1913b: 158, 1914b: 354, 391, 1934a: 171; Leng 1920: 340; Marchant & Borden 1976; McLean & Bennett 1978; McLean & Borden 1977; Murayama 1957a: 37; Nijholt 1980; Patterson & Hatch 1945: 152; Rudinsky & Schneider 1969a; Ruppel 1967: 47; Schedl 1940a: 348, 1963c: 158, 1977e: 43; Schuder 1969: 74; Schwerdtfeger 1960b; Swaine 1909: 108, 1914: 38; Thatcher, T. O. 1961; Thomas, J. B. 1966; Wichmann 1955a: 93, 103; Wood, S. L. 1982b: 1164. (**tx**) Atkinson et al. 1986: 48; Blackman 1931b: 274–275; Bright 1976d: 174; Bright & Stark 1973: 163; Chamberlin 1939: 331–332, 1958: 139–140; Eichhoff 1878b: 408, 512; Evans, D. 1983: 32; Furniss, R. L. & Carolin 1977: 391; Hagedorn 1910a: 147; Hopkins 1905a: 73; Hopping & Jenkins 1933; Johnson 1958b: 237; Keen 1929: 35; LeConte 1868: 154–155, 1876: 350; Schedl 1931a: 13, 1940a: 348, 1955g: 18, 29; Schwerdtfeger 1960b: 254–257; Shore 1985; Swaine 1909: 109–110, 1918a: 91; Thomas, J. B. 1967; Wood, S. L. 1977d: 513, 1982b: 1164. (**ms**) Boocock 1959d; Hatch 1938: 194; Kinghorn 1957b: 213; McLean & Bennett 1978; Mori 1975c; Prebble 1944: 50.

aciculatus Blackman 1931b: 272. Holotype ♀; Clondercroft, New Mexico [USA]; USNM, Washington. Synonymy: Wood 1977c: 513. References: (**cn**) Blackman 1931c; Doane et al. 1936; Keen 1938: 146; Schuder 1969: 74. (**cc**) Blackman 1931c; Keen 1938: 146. (**hb**) Baker, W. L. 1972: 268; Blackman 1931c; Chamberlin 1939: 330–331; Doane et al. 1936; Keen 1938: 146. (**ds**) Chamberlin 1939: 330–331; Keen 1938: 146; Leng & Mutchler 1933: 95; Schuder 1969: 74. (**tx**) Blackman 1931b: 272; Chamberlin 1939: 330–331; Schedl 1955g: 29, 1979c: 10; Wood, S. L. 1977d: 513.

Genus *Gnathotrupes* Schedl

GNATHOTRUPES SCHEDL 1951m: 125. Type-species: *Gnathotrupes holivianus* Schedl, monobasic.

Gnathotrypanus Wood 1968b: 9. Type-species: *Gnathotrypanus terabratus* Wood, original designation. Synonymy: Wood 1973c: 172.

References: (**tx**) Wood, S. L. 1968b: 9, 1973c: 172.

Gnathocortus Schedl 1975d: 11. Type-species: *Gnathocortus caliculus* Schedl, original designation. Synonymy: Wood 1984b: 225.

References: (**tx**) Schedl 1975d: 11; Wood, S. L. 1984b: 225.

Gnathomimus Schedl 1975d: 12. Type-species: *Gnathomimus nothofagi* Schedl, original designation. Synonymy: Wood 1984b: 225.

References: (**tx**) Schedl 1975d: 12; Wood, S. L. 1984b: 225.

Gnathoglochinus Schedl 1975d: 16. Type-species: *Gnathoglochinus impressus* Schedl, original designation. Synonymy: Wood 1984b: 225.

References: (**tx**) Schedl 1975d: 16; Wood, S. L. 1984b: 225.

Keys: Wood 1982b: 1167 for Central America.

References: (**hb**) Wood, S. L. 1982b: 1166, 1986a: 99. (**ds**) Browne 1970: 448; Schedl 1951m: 125; Wood, S. L. 1973c: 172, 1982b: 116, 1986a: 99. (**tx**) Browne 1970: 558; Schedl 1951m: 125; Wood, S. L. 1973c: 172, 1982b: 1166–1170, 1986a: 99.

barbifer (Schedl) 1967d: 13 (*Gnathotrichus*). Holotype, sex?, Chiloe, Chile; Schedl Collection in NHMW, Wien.

Distribution: South America (Chile).

Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).

References: (**ds**) Schedl 1972d: 145. (**tx**) Schedl 1967d: 13, 1972d: 145, 1979c: 34; Wood, S. L. 1973c: 172.

bituberculatus (Blandford) 1904: 248 (*Gnathotrichus*). Lectotype ♀; Purulha, Baja Verapaz, Guatemala; BMNH, London, designated by Wood 1982b: 1168.

Distribution: North America (El Salvador/ Guatemala/ Veracruz in Mexico).

Hosts: *Inga* sp., *Quercus* sp.

Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).

References: (**ds**) Blackwelder 1947: 782; Hagedorn 1910d: 96; Kleine 1914b: 368. (**tx**) Blackman 1931b: 266; Blandford 1904: 248; Hagedorn 1910a: 147; Schedl 1955g: 29; Wood, S. L. 1973c: 172, 1982b: 1168–1169.

impressus Schedl 1977e: 44 (*Gnathotrichus*).

Holotype ♀; El Salvador, Metapan, 1600–2200 m; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1975. Synonymy: Wood 1989: 173.

References: (**tx**) Schedl 1977e: 44; Wood, S. L. 1989: 173.

- bolivianus** Schedl 1951m: 126. Holotype ♀; Bolivia; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (tx) Schedl 1951m: 126, 1979c: 43.
- caliculus** (Schedl) 1975d: 12 (*Gnathocortus*). Holotype ♂; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.
Figures: Schedl 1975k: 13.
Distribution: South America (Argentina).
Hosts: *Nothofagus dombeyi*.
Notes: (3) Wood 1984b: 225 (to *Gnathotrupes*).
References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b, (tx) Schedl 1975d: 11, 1975k: 13, 1979c: 50; Wood, S. L. 1984b: 225.
- castaneus** (Schedl) 1972d: 145 (*Gnathotrichus*). Holotype ♂?; Chile; Schedl Collection in NHMW, Wien.
Distribution: South America (Chile).
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: (tx) Schedl 1972d: 145, 1979c: 54; Wood, S. L. 1973c: 172.
- cirratus** Schedl 1975d: 5. Holotype ♀; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
Hosts: *Nothofagus dombeyi*.
References: (tx) Schedl 1975d: 5, 1979c: 58.
- colaphus** Wood 1989: 181. Holotype ♀; La Carbonera Experimental Forest, 50 km W Merida, Merida, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Nectandra* sp., etc.
References: (tx) Wood, S. L. 1989: 181.
- consobrinus** (Eichhoff) 1878b: 409. *America meridionalis* (Chili); Hamburg Museum, lost.
Distribution: South America (Chile).
Notes: (1) Schedl 1972: 150, Wood 1973: 172 (to *Gnathotrupes*).
References: (ds) Blackwelder 1947: 782; Hagedorn 1910d: 96; Kleine 1914b: 332; Schedl 1972d: 151, (tx) Blackman 1931b: 266; Eichhoff 1878b: 409; Hagedorn 1910a: 147; Schedl 1939h: 47, 1951m: 122, 1972d: 151; Wood, S. L. 1973c: 172.
- crecentus** Wood 1974a: 56. Holotype ♀; Volcan, Puntarenas, Costa Rica, 1000 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: Tree limb.
References: (tx) Wood, S. L. 1974a: 56, 1982b: 1169–1170.
- dilutus** Wood 1974a: 56. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Miconia caudata*.
References: (hb) Wood, S. L. 1982b: 1169, (ds) Wood, S. L. 1982b: 1169, (tx) Wood, S. L. 1974a: 56, 1982b: 1169.
- electus** (Wood) 1968b: 10 (*Gnathotrypanus*). Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: Tree limb.
Notes: (1) Wood 1968b: 10, 1982b: 1168 (to *Gnathotrupes*).
References: (ds) Wood, S. L. 1982b: 1168, (tx) Wood, S. L. 1968b: 10, 1973c: 172, 1982b: 1168.
- fimbriatus** (Schedl) 1955e: 259 (*Gnathotrichus*). Lectotype ♀; Chile, P. Arenas, Sud-Chile, Termas de Puyehue; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 152.
Distribution: South America (Chile).
Notes: (1) Wood 1966b: 23 (to *Gnathotrupes*).
References: (ds) Schedl 1980c: 160, (tx) Schedl 1955e: 256, 259, 1972d: 152, 1979c: 97; Wood, S. L. 1966b: 23, 1973c: 172.
- frontalis** (Schedl) 1972d: 146 (*Gnathotrichus*). Holotype ♂; Sudchile, Puerto Puyulmapu; Schedl Collection in NHMW, Wien.
Distribution: South America (Chile).
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: (tx) Schedl 1972d: 146, 1979c: 100; Wood, S. L. 1973c: 172.
- herbertfranzi** (Schedl) 1972d: 147 (*Gnathotrichus*). Holotype ♂; S. Chile, Umg. Malacahuello; Schedl Collection in NHMW, Wien.
Distribution: South America (Chile).
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: (ds) Schedl 1980c: 159, (tx) Schedl 1972d: 147, 1979c: 116; Wood, S. L. 1973c: 172.
- impressus** (Schedl) 1975d: 17 (*Gnathoglochinus*). Holotype ♀; Argentinien, Nahuel Huapi National Park; NHMW, Wien.
Figures: Schedl 1975d: 17.
Distribution: South America (Argentina).
Hosts: *Nothofagus dombeyi*.
Notes: (3) Wood 1984b: 16 (to *Gnathotrupes*).
References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b; Schedl 1980c: 160, (tx) Schedl 1975d: 17, 1979c: 122; Wood, S. L. 1984b: 16.
- longicollis** (Schedl) 1951m: 120 (*Gnathotrichus*). Holotype, sex?; Colombia; Schedl Collection in NHMW, Wien.
Distribution: South America (Colombia).
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: (tx) Schedl 1951m: 120, 1979c: 140; Wood, S. L. 1973c: 172.
- longipennis** (Blanchard) 1851: 429 (*Tomicus*). Syn-types, sex?; Cercanias de Santiago, Chile; not located.
Distribution: South America (Argentina/Chile).
Hosts: *Nothofagus dombeyi*.
Notes: (1) Eichhoff 1878b: 408 (to *Gnathotrichus*), Wood 1973c: 172 (to *Gnathotrupes*).

- Schedl 1972d: 150 (*springer* Eggers, nomen nudum, synonymy).
References: **(ds)** Blackwelder 1947: 782; Bruch 1914a; Elgneta & Jackson 1957: 77; Gemminger & Harold 1872: 2686; Hagedorn 1910d: 96; Kleine 1913b: 161, 1914b: 332; Lacordaire 1866: 383; Schedl 1972d: 148. **(tx)** Blackman 1931b: 266; Blanchard 1851: 429; Eggers 1929e: 49; Eichhoff 1878b: 408; Gay 1852: 429; Hagedorn 1910a: 147; Lacordaire 1866: 383; Lucas 1920: 305; Porter 1932: 106; Schedl 1939h: 47, 1951m: 118, 1955e: 256, 1972d: 148; Wood, S. L. 1973c: 172.
obnixus Schedl 1939h: 47 (*Gnathotrichus*). Syn- types 2 ♀; Puerto Puyuhuapi, Snd-Chile; 1 in IPKE, Eberswalde, 1 in Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 173.
Notes: (1) Schedl 1979c: 175 (citation of holotype invalid). Wood 1973c: 172 (to *Gnathotrupes*).
References: **(ay)** Naumann-Etienne 1978b. **(hb)** Naumann-Etienne 1978b. **(ds)** Blackwelder 1947: 782; Naumann-Etienne 1978b; Schedl 1972d: 149, 1973d: 162, 1975d: 2, 1978c: 293. **(tx)** Schedl 1939h: 47–48, 1972d: 149, 1979c: 174; Wood, S. L. 1973c: 172.
- corthyloides* Schedl 1951d: 20 (*Gnathotrichus*). Lectotype ♀; Chile, Valdivia; Schedl Collection in NHMW, Wien, preoccupied by Eichhoff 1869, designated by Schedl 1979c: 66. Synonymy: Wood 1989: 173.
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: **(tx)** Schedl 1951d: 20, 1955e: 256, 1979c: 66; Wood, S. L. 1973c: 172, 1989: 173.
- sextuberculatus* Schedl 1951m: 118 (*Gnathotrichus*). Holotype, sex?; Chile; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972d: 150.
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: **(ay)** Naumann-Etienne 1978b. **(hb)** Naumann-Etienne 1978b. **(ds)** Hoganson & Ashworth 1982; Naumann-Etienne 1978b; Schedl 1975d: 2. **(tx)** Schedl 1951m: 118, 1955e: 256, 1964m: 312, 1972d: 150, 1979c: 227; Wood, S. L. 1973c: 172.
- quadrituberculatus* Schedl 1951m: 122 (*Gnathotrichus*). Holotype, sex?; Chile; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972d: 150.
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: **(tx)** Schedl 1951d: 17, 1951m: 118, 122, 1955e: 256, 1972d: 150, 1979c: 200; Wood, S. L. 1973c: 172.
- corthyliformis* Schedl 1964m: 312 (*Gnathotrichus*). Lectotype ♀; Chile, Valdivia; Schedl Collection in NHMW, Wien, automatic, designated by Schedl 1979c: 66. Synonymy: Wood 1989: 173.
References: **(ds)** Schedl 1967d: 5, 1972d: 151–152. **(tx)** Schedl 1964m: 312, 1972d: 151–152, 1979c: 66; Wood, S. L. 1973c: 172.
- constrictus* Schedl 1975d: 6. Holotype ♂; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien. Synonymy: Wood 1989: 173.
References: **(ay)** Naumann-Etienne 1978b. **(hb)** Naumann-Etienne 1978b. **(ds)** Naumann-Etienne 1978b. **(tx)** Schedl 1975d: 6, 1979c: 64; Wood, S. L. 1989: 173.
- longiusculus* (Schedl) 1951m: 121 (*Gnathotrichus*). Holotype ♂; Tierra del Fuego, Via Monte; Eggers Collection, in NHMW, Wien.
Distribution: South America (Argentina, Chile).
Hosts: *Nothofagus dombeyi*.
Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).
References: **(ds)** Schedl 1972d: 148. **(tx)** Schedl 1951m: 121, 1972d: 148, 1979c: 142; Wood, S. L. 1973c: 172.
- ciliatus* Schedl 1975d: 4. Holotype ♀; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992a: 89.
References: **(tx)** Schedl 1975d: 4; Wood, S. L. 1992a: 89.
- nanulus* (Schedl) 1972d: 149 (*Gnathotrichus*). Holotype, sex?; Chile; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina/ Chile).
Hosts: *Nothofagus dombeyi*.
References: **(ay)** Naumann-Etienne 1978b. **(hb)** Naumann-Etienne 1978b. **(ds)** Naumann-Etienne 1978b; Schedl 1975d: 2. **(tx)** Schedl 1972d: 149, 1975d: 4, 1979c: 163; Wood, S. L. 1973c: 172.
- nanus* (Eichhoff) 1878b: 410 (*Gnathotrichus*). Holotype ♀; America meridionalis (Chile); Hamburg Museum, lost.
Distribution: South America (Chile).
References: **(ds)** Hagedorn 1910d: 97. **(tx)** Blackman 1931b: 266; Eichhoff 1978b: 410; Hagedorn 1910a: 147; Schedl 1939k: 727; Wood, S. L. 1973c: 172.
- naumanni* (Schedl) 1975d: 15 (*Gnathomimus*). Holotype ♀; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.
Figures: Schedl 1975d: 16.
Distribution: South America (Argentina).
Hosts: *Nothofagus dombeyi*.
Notes: (3) Wood 1954b: 225 (to *Gnathotrupes*).
References: **(ay)** Naumann-Etienne 1978b. **(hb)** Naumann-Etienne 1978b. **(ds)** Naumann-Etienne 1978b. **(tx)** Schedl 1975d: 15–16, 1979c: 164; Wood, S. L. 1984b: 225.
- nectandrae* Wood 1989: 181. Holotype ♂; La Carbonera Experimental Forest, 50 km W Merida, Merida, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Nectandra* sp.
References: **(tx)** Wood, S. L. 1989: 181.

nothofagi (Schedl) 1975d: 13 (*Gnathomimus*).

Holotype ♀: Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

Notes: (3) Wood 1984b: 225 (to *Gnathotrupes*).

References: (tx) Schedl 1975d: 13, 1979c: 171; Wood, S. L. 1984b: 225.

pauciconcarus Schedl 1975d: 7. Holotype ♀;

Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

Notes: (1) This name was originally spelled *pauciconvacus*, an obvious lapsus calami; Schedl 1979c: 186 gave the corrected spelling.

References: (tx) Schedl 1975d: 7, 1979c: 186.

pustulatus Schedl 1975d: 9. Holotype ♀;

Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b; Schedl 1980c: 159, (tx) Schedl 1975d: 9, 1979c: 206.

similis Schedl 1975d: 10. Holotype ♀; Argentinien,

Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b, (tx) Schedl 1975d: 10, 1979c: 229.

solidus Schedl 1975d: 8. Holotype ♂; Argentinien,

Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b, (tx) Schedl 1975d: 8, 1979c: 232.

terebratus (Wood) 1968b: 9 (*Gnathotrypanus*).

Holotype ♂; Volcan, Puntarenas, Costa Rica; Wood Collection.

Figures: Wood 1968b: 5.

Distribution: North America (Costa Rica).

Hosts: Presumably *Pourouma aspera*.

Notes: (1) Wood 1973c: 172, 1982: 1167 (to *Gnathotrupes*).

References: (hb) Wood, S. L. 1982b: 1167, (ds) Wood, S. L. 1982b: 1167, (tx) Wood, S. L. 1968b: 5, 9, 1973c: 172, 1982b: 1167.

rafer (Schedl) 1975d: 3 (*Gnathotrichus*). Holotype,

sex?; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

Notes: (1) Wood 1973c: 172 (to *Gnathotrupes*).

References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b; Schedl 1980c: 159, (tx) Schedl 1975d: 3, 1979c: 263; Wood, S. L. 1973c: 172.

velatus Schedl 1975d: 10. Holotype ♀; Argentinien, Nahuel Huapi National Park; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

Hosts: *Nothofagus dombeyi*.

References: (ay) Naumann-Etienne 1978b, (hb) Naumann-Etienne 1978b, (ds) Naumann-Etienne 1978b; Schedl 1980c: 160, (tx) Schedl 1975d: 10, 1979c: 265.

Genus *Tricolus* Blandford

TRICOLUS BLANDFORD 1905: 286. Type-species: *Tricolus ovicollis* Blandford, subsequent designation by Hopkins 1914: 131.

Pterocyclonoides Schedl 1970e: 101. Type-species: *Pterocyclonoides octodentatus* Schedl, monobasic. Synonymy: Wood 1984b: 228.

References: (tx) Schedl 1970e: 101–102; Wood, S. L. 1984b: 228.

Keys: Wood 1982b: 1170 for Central America.

References: (ay) Nobuchi 1969a: 65, (hb) Wood, S. L. 1982b: 1170, 1986a: 99, (ds) Wood, S. L. 1982b: 1170, 1986a: 99, (tx) Blandford 1905: 286; Bright 1972d: 98; Wood, S. L. 1967c: 119–141, 1982b: 1170–1185, 1986a: 99.

abruptus Schedl 1976a: 83. Holotype, sex?; Carna, Pernambuco, Brasil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 83.

aciculatus Wood 1974a: 62. Holotype ♀; 10 km NE Teziutlan, Puebla, Mexico; Wood Collection.

Distribution: North America (Puebla in Mexico).

Hosts: *Miconia* sp., *Rubus* sp.

References: (hb) Atkinson & Equihua 1985c: 361; Wood, S. L. 1982b: 1183, (ds) Atkinson & Equihua 1985c: 361; Wood, S. L. 1982b: 1183, (tx) McNamara 1984: 759; Wood, S. L. 1974a: 62, 1982b: 1183–1184.

affinis Eggers 1931b: 38. Holotype, sex?; Sao Paulo, Brasil; Prague Museum.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783, (tx) Eggers 1931b: 38; Schedl 1939j: 581, 1979c: 14.

amplus Wood 1974a: 63. Holotype ♀; 10 km NE Teziutlan, Puebla, Mexico; Wood Collection.

Distribution: North America (Puebla in Mexico).

Hosts: Tree limb.

References: (hb) Wood, S. L. 1982b: 1185, (ds) Wood, S. L. 1982b: 1185, (tx) Wood, S. L. 1974a: 63, 1982b: 1185.

ardis Wood 1974a: 58. Holotype ♀; Escasu, San Jose, Costa Rica, 1300 m; Wood Collection.

Figures: Wood 1952b: 1174 (declivity).

Distribution: North America (Costa Rica/ Panama).

Hosts: Tree branches and a seedling.

References: **(hb)** Wood, S. L. 1952b: 1176. **(ds)** Wood, S. L. 1952b: 1176. **(tx)** Wood, S. L. 1974a: 58, 1952b: 1176.

badius Wood 1974a: 60. Holotype ♀; Pandora, Limon, Costa Rica, 50 m; Wood Collection.

Figures: Wood 1952b: 1174 (declivity).

Distribution: North America (Costa Rica/ Panama).

Hosts: *Serjania* sp., a liana, and a tree branch.

References: **(hb)** Wood, S. L. 1952b: 1177. **(ds)** Wood, S. L. 1952b: 1177. **(tx)** Wood, S. L. 1974a: 60, 1952b: 1177.

bicolor Wood 1974a: 63. Holotype ♀; Volcan Poas, Heredia, Costa Rica, 1500 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Tree branch.

References: **(hb)** Burgos & Saucedo 1983: 132; Wood, S. L. 1952b: 1184. **(ds)** Burgos & Saucedo 1983: 132; Wood, S. L. 1952b: 1184. **(tx)** Wood, S. L. 1974a: 63, 1952b: 1184.

bifidus Schedl 1939j: 579. Holotype, sex?; Brasilien; Schedl Collection in NHMW, Wien.

Figures: Pedrosa-Macedo & Schonherr 1955: 53. Distribution: South America (Brazil).

References: **(ds)** Blackwelder 1947: 783; Pedrosa-Macedo & Schonherr 1955: 53; Schedl 1976a: 56. **(tx)** Pedrosa-Macedo & Schonherr 1955: 53; Schedl 1939j: 579, 1979c: 39–40.

capitalis Wood 1974a: 61. Holotype ♀; 13 km SE El Hato del Volcan, Chiriqui, Panama, 1000 m; Wood Collection.

Figures: Wood 1952b: 1174 (declivity).

Distribution: North America (Panama).

References: **(hb)** Wood, S. L. 1952b: 1179. **(ds)** Wood, S. L. 1952b: 1179. **(tx)** Wood, S. L. 1974a: 61, 1952b: 1179.

cecropii Wood 1974a: 58. Holotype ♀; Turrialba, Cartago, Costa Rica, 700 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: *Cecropia peltata* leaf petioles.

Notes: (3) Wood 1982: 1176 (this is a domicile parasite of *Corthylus cecropii* Wood).

References: **(hb)** Wood, S. L. 1952b: 1175–1176; **(ds)** Wood, S. L. 1952b: 1175–1176; **(tx)** Wood, S. L. 1974a: 58, 1952b: 1175–1176.

difodinus Bright 1972a: 1380. Holotype ♂; 21 km N Ocozacoatlán, Chiapas, Mexico; CNCI, Ottawa. Figures: Bright 1972a: 1377, Wood 1952b: 1181 (declivity).

Distribution: North America (Guatemala/ Chiapas in Mexico).

Hosts: Tree branches.

References: **(ec)** Equihua & Atkinson 1986: 634. **(hb)** Atkinson & Equihua 1986a: 422; Equihua & Atkinson 1986: 634; Wood, S. L. 1952b: 1182. **(ds)** Atkinson & Equihua 1985c: 361, 1986a: 422.

1988: 102; Equihua & Atkinson 1986: 634, 1988: 212; Wood, S. L. 1952b: 1182. **(tx)** Bright 1972a: 1377, 1380; McNamara 1977: 199; Wood, S. L. 1952b: 1182.

fenoris Wood 1974a: 60. Holotype ♂?; 15 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Liana.

References: **(hb)** Wood, S. L. 1952b: 1175. **(ds)** Wood, S. L. 1952b: 1175. **(tx)** Wood, S. L. 1974a: 60, 1952b: 1175.

frontalis Wood 1974a: 61. Holotype ♀; 8 km NE Tezintlan, Puebla, Mexico, 1600 m; Wood Collection.

Distribution: North America (Puebla in Mexico).

Hosts: Tree branch.

References: **(ds)** Wood, S. L. 1952b: 1179. **(tx)** Wood, S. L. 1974a: 61, 1952b: 1179.

gracilipennis Schedl 1950i: 170. Lectotype, sex?; Brazil, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 107.

Distribution: South America (Brazil).

References: **(tx)** de Ruelle 1970: 114; Schedl 1950i: 170, 1979c: 107.

gracilis Eggers 1937a: 87. Holotype, sex?; Guadeloupe; USNM, Washington.

Distribution: Antilles Islands (Guadeloupe).

References: **(ds)** Blackwelder 1947: 783; Bright 1955c: 176. **(tx)** Anderson, W. H. & Anderson 1971: 13; Bright 1955c: 176; Eggers 1937a: 87, 1940a: 108; Schedl 1950i: 170, 1954b: 40, 1979c: 108.

inaffectus Wood 1974a: 57. Holotype ♀; Volcan Poas, Heredia, Costa Rica, 2600 m; Wood Collection.

Distribution: North America (Costa Rica).

References: **(ds)** Wood, S. L. 1952b: 1175. **(tx)** Wood, S. L. 1974a: 57, 1952b: 1175.

incisus Schedl 1976a: 84. Holotype, sex?; Caraguatatuba, SP, Res. Flor., 40 m, Brazil; MZUSP, Sao Paulo.

Distribution: South America (Brazil).

References: **(tx)** Schedl 1976a: 84.

inornatus Wood 1974a: 57. Holotype ♀; 15 km SE Cartago, Cartago, Costa Rica, 2500 m; Wood Collection.

Figures: Wood 1952b: 1174 (declivity).

Distribution: North America (Costa Rica).

Hosts: *Siparuna nicaraguaensis*.

References: **(hb)** Atkinson & Equihua 1985c: 361; Wood, S. L. 1952b: 1174. **(ds)** Atkinson & Equihua 1985c: 361; Wood, S. L. 1952b: 1174. **(tx)** Wood, S. L. 1974a: 57, 1952b: 1174.

intrusus Wood 1974a: 58. Holotype ♀; El Laurel, 12 km SW Caracas, Venezuela, 1300 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Cecropia* sp. leaf petioles.

References: (tx) Wood, S. L. 1974a: 58.

minutissimus Schedl 1976a: 54. Holotype, sex?; Linhares, E. Santos [Brazil]; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 54.

naerus Wood 1974a: 61. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.

Figures: Wood 1982b: 1181 (declivity).

Distribution: North America (Costa Rica).

Hosts: *Phoebe mexicana*, *Werklia insignata*, and a liana.

References: (hb) Wood, S. L. 1982b: 1180. (ds) Wood, S. L. 1982b: 1180. (tx) Wood, S. L. 1974a: 61, 1982b: 1180.

nodifer Blandford 1905: 287. Holotype, sex?; Mirandilla, Esquintla, Guatemala; BMNH, London.

Figures: Blandford 1905: pl. 9, Wood 1982b: 1174 (declivity).

Distribution: North America (Costa Rica/ Guatemala/ Veracruz? in Mexico).

Hosts: *Cecropia* sp. leaf petiole, and a tree branch.

References: (hb) Wood, S. L. 1982b: 1178. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1178. (tx) Blandford 1905: 287; Eggers 1937a: 38; Hagedorn 1910a: 149; Wood, S. L. 1966b: 30, 1967c: 139, 1982b: 1178.

triarmatus Schedl 1939j: 578. Holotype, sex?; Colonia, vermutlich Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 30. References: (ds) Blackwelder 1947: 783; Ferrer 1942. (tx) Schedl 1939j: 578, 1940a: 355; Wood, S. L. 1966b: 30.

octodentatus (Schedl) 1970b: 102 (*Pterocyclonoides*). Holotype, sex?; Brasilien, Parana, Rondon; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1970b: 102.

oricollis Blandford 1905: 287. Holotype, sex?; Cerro Zamil, Quezaltenango, Guatemala; BMNH, London.

Figures: Wood 1982b: 1181 (declivity).

Distribution: North America (Costa Rica/ Puebla in Mexico/ Panama).

Hosts: *Miconia* sp., *Phoebe mexicana*, and a liana.

References: (hb) Wood, S. L. 1982b: 1180. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1180. (tx) Blandford 1905: 287; Eggers 1937a: 87; Hagedorn 1910a: 149; Hopkins 1914: 131; Schedl 1939j: 579; Wood, S. L. 1982b: 1180–1183.

parsus Wood 1974a: 59. Holotype ♂?; 15 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.

Figures: Wood 1982b: 1174 (declivity).

Distribution: North America (Costa Rica).

Hosts: Liana.

References: (hb) Wood, S. L. 1982b: 1176. (ds) Wood, S. L. 1982b: 1176. (tx) Wood, S. L. 1974a: 59, 1982b: 1176.

partilis Wood 1974a: 60. Holotype ♂; Volcan, Puntarenas, Costa Rica; Wood Collection.

Figures: Wood 1982b: 1174 (declivity).

Distribution: North America (Costa Rica).

Hosts: Tree branch.

References: (hb) Wood, S. L. 1982b: 1178. (ds) Wood, S. L. 1982b: 1178. (tx) Wood, S. L. 1974a: 60, 1982b: 1178.

peltatus Wood 1974a: 62. Holotype ♀; Cerro Punta near Volcan Chiriqui, Chiriqui, Panama; Wood Collection.

Distribution: North America (Costa Rica/ Panama).

Hosts: Tree branches and a liana.

References: (hb) Wood, S. L. 1982b: 1183. (ds) Wood, S. L. 1982b: 1183. (tx) Wood, S. L. 1974a: 62, 1982b: 1183.

perdilgens Schedl 1950i: 171. Holotype ♀; Jamaica; Schedl Collection in NHMW, Wien.

Figures: Bright 1972d: 59, 75, 95.

Distribution: Antilles Islands (Dominican Republic in Hispanola/ Jamaica).

References: (ds) Bright 1973: 18, 1981c: 156, 1985c: 176. (tx) Bright 1981c: 156, 1985c: 176; Schedl 1950i: 170, 1979c: 188.

ignotus Bright 1972d: 99. Holotype ♀; Hardwar Gap, 4000', St. Andrew Parish, Jamaica; CNCI, Ottawa. Synonymy: Bright 1985c: 176. References: (tx) Bright 1972d: 59, 75, 95, 99, 1985c: 176.

pernanulus Schedl 1939j: 581. Holotype, sex?; Sauda (Brasilien?); Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783; Schedl 1972g: 46. (tx) Schedl 1939j: 581, 1979c: 190.

plamanui Schedl 1954b: 40. Lectotype, sex?; Rondon, Parana, Brazil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 196. Figures: Wood 1982b: 1181 (declivity).

Distribution: North America (Costa Rica/ Panama), South America (Brazil/ Venezuela).

Hosts: Tree branches and lianas.

References: (hb) Wood, S. L. 1982b: 1184. (ds) Wood, S. L. 1982b: 1184. (tx) de Ruelle 1970: 114; Schedl 1954b: 40, 1979c: 196; Wood, S. L. 1982b: 1184.

pumilio Eggers 1937a: 87. Holotype, sex?; Bolivien (Cochabamba); Eggers Collection, in NHMW, Wien.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1937a: 87; Schedl 1939j: 580, 1939k: 726, 1979c: 204.

ruficollis (Fabricius) 1801: 388 (*Bostrichus*). Holotype ♀; published as *America meridionali*, type labeled Essequibo [Guyana]; UZMC, Copenhagen.

Distribution: South America (Guyana).

References: (ds) Dejean 1937; Stein & Weise 1877: 164. (tx) Eichhoff 1868d: 421; Fabricius 1801: 388; Ferrari 1868: 255; Sharp 1879: 102. (ms) Eichhoff 1868d: 421.

rufithorax Wood 1974a: 59. Holotype ♀; Finca Cromaco on Rio Coto Brus, Puntarenas, Costa Rica, 500 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Liana.

References: (hb) Wood, S. L. 1982b: 1177. (ds) Wood, S. L. 1982b: 1177. (tx) Wood, S. L. 1974a: 59, 1982b: 1177.

saunderi Wood 1967c: 139. Holotype ♂; Dominical, Puntarenas, Costa Rica; Wood Collection.

Figures: Wood 1982b: 1174 (declivity).

Distribution: North America (Costa Rica).

Hosts: *Theobroma cacao*, and a liana.

References: (ds) Wood, S. L. 1982b: 1179. (tx) Wood, S. L. 1967c: 139, 1982b: 1179.

scitulus Wood 1974a: 62. Holotype ♀; Tapanti, Cartago, Costa Rica; Wood Collection.

Figures: Wood 1982b: 1181 (declivity).

Distribution: North America (Costa Rica/Panama), South America (Venezuela).

Hosts: *Crotalaria* sp., *Phoebe mexicana*, etc.

References: (hb) Wood, S. L. 1982b: 1182. (ds) Wood, S. L. 1982b: 1182. (tx) Wood, S. L. 1974a: 62, 1982b: 1182.

senex Schedl 1939j: 580. Lectotype ♂; Nova Teutonia, Brazil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 225.

Figures: Schedl 1939j: 580, Terra 1987: 28, Wood 1982b: 1181 (declivity).

Distribution: North America (Costa Rica), South America (Bolivia/ Brazil).

Hosts: *Calandria confusa*, *Siporana nicaraguensis*, etc.

References: (hb) Wood, S. L. 1982b: 1182. (ds) Blackwelder 1947: 783; Nimberg 1971: 58; Schedl 1966f: 91, 1973d: 161, 1978c: 291; Terra 1987: 27; Wood, S. L. 1982b: 1182. (tx) de Ruette 1970: 114; Schedl 1939j: 580, 1979c: 225; Terra 1987: 27–28; Wood, S. L. 1982b: 1181–1183.

simplicis Wood 1974a: 57. Holotype ♀; Palin, Esquinla, Guatemala; Wood Collection.

Distribution: North America (Costa Rica/ Guatemala).

Hosts: *Ficus* sp., and a tree branch.

References: (hb) Wood, S. L. 1982b: 1173. (ds) Wood, S. L. 1982b: 1173. (tx) Wood, S. L. 1974a: 57, 1982b: 1173.

spheniscus Schedl 1939j: 581. Lectotype ♂; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 234. Figures: Pedrosa-Macedo & Schonherr 1985: 54. Distribution: South America (Brazil).

References: (ds) Blackwelder 1947; Pedrosa-Macedo & Schonherr 1985: 54; Schedl 1966f: 91, 1967d: 4, 1970e: 85, 1972g: 46, 1976a: 56. (tx) Pedrosa-Macedo & Schonherr 1985: 54; Schedl 1939j: 581, 1950i: 171, 1979c: 234.

subincisuralis Schedl 1939k: 726. Holotype, sex?; Brasilien, Sta. Catharina; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783; Pedrosa-Macedo & Schonherr 1985: 55; Schedl 1973d: 161, 1976d: 4. (tx) Pedrosa-Macedo & Schonherr 1985: 55; Schedl 1939k: 726, 1979c: 242.

unidentatus Bright 1972d: 98. Holotype ♀; Barbecue Bottom, Trelawny Parish, Jamaica; CNCI, Ottawa.

Figures: Bright 1972: 95.

Distribution: Antilles Islands (Jamaica).

References: (ds) McNamara 1977: 200; Bright 1985c: 176. (tx) Bright 1972d: 95, 98, 1985c: 176.

Genus *Amphicranus* Erichson

AMPHICRANUS ERICHSON 1836: 63. Type-species: *Amphicranus thoracicus* Erichson, monobasic.

Piezorhopalus Guerin-Meneville 1838: 107.

Type-species: *Piezorhopalus nitidulus* Guerin-Meneville = *Amphicranus thoracicus* Erichson, monobasic. Synonymy: Blandford 1905: 290.

References: (tx) Blandford 1905: 290; Guerin-Meneville 1838: 107.

Steganocranus Eichhoff 1878b: 460. Type-species: *Steganocranus dohrni* Eichhoff, monobasic. Synonymy: Blandford 1905: 289.

References: (tx) Blandford 1905: 289; Eichhoff 1878b: 70, 460; Schedl 1939j: 583, 1950i: 176.

Keys: Blandford 1905: 289 and Wood 1982b: 1186 for Central America.

References: (hb) Wood, S. L. 1982b: 1185, 1986a: 99. (ds) Wood, S. L. 1982b: 1185, 1986a: 99. (tx) Blandford 1905: 289; Eggers 1935a: 157; Erichson 1836: 63; Wood, S. L. 1967d: 37–57, 1982b: 1185, 1986a: 99.

acus Wood 1974a: 66. Holotype ♀; El Laurel Experiment Station, 12 km SW Caracas, Venezuela, 1800 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Croton* sp., etc.

References: (tx) Wood, S. L. 1974a: 66.

argutus Wood 1976a: 349. Holotype ♀; Finca La Lola, Limon, Costa Rica; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: *Theobroma cacao*.

References: (ds) Wood, S. L. 1982b: 1193. (tx) Wood, S. L. 1976a: 349, 1982b: 1193.

balteatus Blandford 1905: 291. Holotype, sex?; Cerro Zunil, Quezaltenango, Guatemala, 1300 m; BMNH, London.

Figures: Blandford 1905:pl. 9.

Distribution: North America (Guatemala).

Hosts: "Caldo de Frijol," a tree.

References: (hb) Wood, S. L. 1982b: 1201. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 257, 1914b: 308; Wood, S. L. 1982b: 1201. (tx) Blandford 1905: 291; Hagedorn 1910a: 139; Wood, S. L. 1982b: 1201.

belti Blandford 1905: 292. Holotype, sex?; Chontales, Nicaragua; BMNH, London.

Figures: Blandford 1905:pl. 9.

Distribution: North America (Costa Rica/Nicaragua).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 368. (tx) Blandford 1905: 289, 292; Hagedorn 1910a: 139; Wood, S. L. 1982b: 201.

bipunctatus Eichhoff 1878b: 469. Holotype, sex?; America meridionalis (Nova Granada); Hamburg Museum, lost.

Distribution: South America (Colombia).

References: (ds) Blackwelder 1947: 783; Blandford 1905: 289; Kleine 1913b: 157. (tx) Blandford 1905: 289; Eichhoff 1878b: 469; Hagedorn 1910a: 139.

brasiliensis Eggers 1931b: 37. Holotype, sex?; Brazil: Sao Paulo; Prague Museum.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1931b: 37–38.

brevipennis Blandford 1905: 293. Holotype ♀?; San Isidro, Suchitepequez, Guatemala; BMNH, London.

Figures: Blandford 1905:pl. 9. Wood 1982b: 1191.

Distribution: North America (Costa Rica/Guatemala).

Hosts: Tree branch.

References: (hb) Wood, S. L. 1982b: 1190. (ds) Blackwelder 1947: 783; Bright 1972a: 1385; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 368; Wood, S. L. 1982b: 1190. (tx) Blandford 1905: 293; Hagedorn 1910a: 139; Wood, S. L. 1982b: 1190–1191.

broenei Schedl 1979j: 128. Holotype, sex?; Brasilien, Amapa, Alt. Amapari; Schedl Collection in NHMW, Wien, automatic.

Distribution: South America (Brazil).

References: (tx) Schedl 1979j: 128.

elegantulus Schedl 1978c: 304. Holotype, sex?; Brasilien, Amapa, Alt. Amapari; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1963.

References: (tx) Schedl 1978c: 304; Wood, S. L. 1981: 122.

electus Wood 1981: 122. Holotype, sex?; Brasilien, Amapa, Alta. Amapari; Schedl Collection in NHMW, Wien, automatic.

Notes: (1) This is an unneeded replacement name.

References: (tx) Schedl 1979j: 128; Wood, S. L. 1981: 122.

callosus (Schedl) 1935h: 351 (*Pterocyclon*). Holotype ♀; Coronado [presumably San Isidro de Coronado, San Jose], Costa Rica. 1400–1500 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: (ds) Wood, S. L. 1982b: 1194. (tx) Schedl 1935h: 350–351, 1952a: 450, 1979c: 50; Wood, S. L. 1982b: 1194.

collaris Blandford 1905: 294. Holotype ♀?; Volcan Chiriqui, Panama; BMNH, London.

Figures: Blandford 1905:pl. 9.

Distribution: North America (Costa Rica/Panama).

Hosts: *Phoebe mexicana*, etc.

References: (hb) Wood, S. L. 1982b: 1200. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1200. (tx) Blandford 1905: 294; Hagedorn 1910a: 139; Sampson 1912: 245; Wood, S. L. 1982b: 1200.

cordatus (Bright) 1972a: 1380 (*Tricolus*). Holotype ♂; 184 km S Oaxaca on Highway 131, Oaxaca, Mexico, 2000 m; CNCI, Ottawa.

Figures: Atkinson et al. 1986: 43, Bright 1972a: 1377.

Distribution: North America (Oaxaca in Mexico).

References: (hb) Atkinson et al. 1986: 44; Burgos & Saucedo 1983: 132. (ds) Atkinson et al. 1986: 44; Burgos & Saucedo 1983: 132; Wood, S. L. 1982b: 1203. (tx) Bright 1972a: 1380; Atkinson et al. 1986: 43; Bright 1972a: 1377; McNamara 1977: 199; Wood, S. L. 1982b: 1203.

dohrni (Eichhoff) 1878b: 461 (*Steganocranus*). Syntypes 2, 1 ♀; America meridionali; lost with Hamburg Museum, 1 ♀ in Dohrn Collection at Stettin now in Schedl Collection in NHMW, Wien. Distribution: South America (Brazil/Venezuela). Hosts: Tree bole.

Notes: (1) Schedl 1979c: 83 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 96; Schedl 1966f: 91. (tx) Eichhoff 1878b: 461; Hagedorn 1910a: 139; Hopkins 1914: 129; Schedl 1939j: 583, 1950i: 146, 1951h: 285, 1979c: 83.

caudatus Eggers 1931a: 17. Holotype, sex?; Brazil; MNB, Berlin. Synonymy: Schedl 1950i: 176.

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1931a: 17, 1943a: 384; Schedl 1950i: 146.

eichhoffi Eggers 1931a: 17. Holotype ♂; Bolivia (Cochabamba); Eggers Collection, in NHMW, Wien. Distribution: South America (Bolivia).

- References: **(ds)** Blackwelder 1947: 783. **(tx)** Eggers 1931a: 17, 1943a: 384; Schedl 1950i: 176, 1979c: 87.
- galeatus* (Eggers) 1943a: 384 (*Steganocranus*).
Holotype ♀; Bolivia (Cochabamba); Eggers Collection, in NHMW, Wien. Synonymy: Schedl 1950i: 176.
References: **(tx)** Eggers 1943a: 384; Schedl 1950i: 176, 1979c: 102.
- elegans* Eichhoff 1869a: 276. Holotype, sex?; Mexico; Hamburg Museum, lost.
Figures: Blandford 1905: pl. 9, Wood 1982b: 1191.
Distribution: North America (Costa Rica/ Veracruz in Mexico/ Nicaragua).
Hosts: *Theobroma cacao*.
References: **(ds)** Blackwelder 1947: 783; Blandford 1905: 290; Ferrer 1942; Gemminger & Harold 1872: 2680, 2692; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 353; Wood, S. L. 1982b: 1192. **(tx)** Blandford 1905: 289–290; Eggers 1931a: 18; Eichhoff 1869a: 276, 1878b: 465; Hagedorn 1910a: 139; Schedl 1940a: 359; Wood, S. L. 1982b: 1191–1192.
- elegantulus* Schedl 1963d: 225. Holotype, sex?; Brasilien: Matto Grosso, Rio Caraguata; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: **(tx)** Schedl 1963d: 225, 1979c: 88.
- fastigiatus* Blandford 1905: 296. Holotype ♂; Cerro Zunil, Quezaltenango, Guatemala; BMNH, London.
Figures: Blandford 1905: pl. 9.
Distribution: North America (Costa Rica/ Guatemala).
Hosts: *Conostegia oerstediana*, *Parkinsonia aculeata*.
References: **(hb)** Wood, S. L. 1982b: 1205. **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 368; Schedl 1933c: 164; Wood, S. L. 1982b: 1205. **(tx)** Blandford 1905: 296; Eggers 1931a: 16, 18; Hagedorn 1910a: 139; Schedl 1939j: 583, 1950i: 176; Wood, S. L. 1982b: 1205–1206.
- filliformis* Blandford 1905: 295. Holotype, sex?; Omilteme, Guerrero, Mexico; BMNH, London.
Figures: Blandford 1905: pl. 9.
Distribution: North America (Chiapas, Guerrero in Mexico).
References: **(hb)** Atkinson & Equihua 1985c: 361. **(ds)** Atkinson & Equihua 1985c: 361; Blackwelder 1947: 783; Ferrer 1942; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 353; Wood, S. L. 1982b: 1203. **(tx)** Blandford 1905: 295; Hagedorn 1910a: 139; Schedl 1940a: 359; Wood, S. L. 1982b: 1203.
- fryi* Blandford 1905: 291. Holotype, sex?; Brazil, Rio de Janeiro; BMNH, London.
Distribution: South America (Brazil/ Cayenne/ Ecuador).
References: **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 337. **(tx)** Blandford 1905: 289, 291; Hagedorn 1910a: 139.
- fulgidus* Wood 1976a: 350. Holotype ♀; Finca Los Diamantes [near Dominical], Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Theobroma cacao*.
References: **(ds)** Wood, S. L. 1982b: 1195. **(tx)** Wood, S. L. 1976a: 350, 1982b: 1195.
- gracilis* Eggers 1943a: 383. Holotype, sex?; Bolivia (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia).
References: **(tx)** Anderson, W. H. & Anderson 1971: 13; Eggers 1943a: 383.
- grouvellei* Blandford 1905: 294. Syntypes 2, sex?; tobacco-refuse, probably Brazilian, intercepted at Paris; BMNH, London.
Distribution: South America (Brazil).
References: **(ec)** Wichmann 1955a: 107. **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 337; Wichmann 1955a: 107. **(tx)** Blandford 1905: 290, 294; Hagedorn 1910a: 139.
- hybridus* Blandford 1905: 298. Holotype ♀; Guatemala, Guatemala; BMNH, London.
Figures: Blandford 1905: pl. 9.
Distribution: North America (Guatemala/ Nayarit in Mexico).
References: **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 368; Wood, S. L. 1982b: 1193. **(tx)** Blandford 1905: 298; Eggers 1931a: 19; Hagedorn 1910a: 139; Wood, S. L. 1982b: 1193.
- lesnei* Hagedorn 1903b: 550. Holotype, sex?; Vallee de Cauca, Colombia; MNIN, Paris.
Distribution: South America (Colombia).
References: **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157. **(tx)** Hagedorn 1903b: 550, 1910a: 139.
- macellus* Wood 1974a: 64. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica; 30 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: Tree limb.
References: **(hb)** Wood, S. L. 1982b: 1202. **(ds)** Wood, S. L. 1982b: 1202. **(tx)** Wood, S. L. 1974a: 64, 1982b: 1202.
- melanura* (Blandford) 1904: 272 (*Pterocyclon*). Holotype ♂; Volcan de Chiriqui, Panama; BMNH, London.
Distribution: North America (Costa Rica/ Panama).
Hosts: *Phoebe mexicana*.
References: **(hb)** Wood, S. L. 1982b: 1195. **(ds)** Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1195. **(tx)** Blandford 1904: 272; Hagedorn 1910a: 149; Schedl 1935h: 348; Wood, S. L. 1982b: 1195–1196.
- opacifrons* Schedl 1935h: 350. Holotype ♀; Coro-

nado [presumably San Isidro de Coronado, San Jose], Costa Rica; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 26. References: (tx) Schedl 1935h: 350, 1979c: 179; Wood, S. L. 1966b: 26.

micans Wood 1976a: 350. Holotype ♀; Rio Damitas, Dota Mts., San Jose, Costa Rica; Wood Collection. Distribution: North America (Costa Rica/ Oaxaca, Veracruz in Mexico/ Panama). Hosts: *Protium* sp., *Terminalia* sp.

References: (hb) Wood, S. L. 1982b: 1194. (ds) Atkinson & Equihua 1986a: 422, 1988: 85; Wood, S. L. 1982b: 1194. (tx) Wood, S. L. 1976a: 350, 1982b: 1194.

minor (Eggers) 1935a: 332 (*Anchonocerus*). Holotype ♂; Bolivien (Cochabamba); USNM, Washington.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 20; Eggers 1935a: 332–333.

mirandus Wood 1974a: 63. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica, 30 m; Wood Collection. Distribution: North America (Costa Rica).

Hosts: Tree limb.

References: (hb) Wood, S. L. 1982b: 1197. (ds) Wood, S. L. 1982b: 1197. (tx) Wood, S. L. 1974a: 63, 1982b: 1197–1198.

mucronatus Wood 1974a: 66. Holotype ♀; Volcan de Chiriqui at Cerro Punta, Chiriqui, Panama, 1800 m; Wood Collection.

Distribution: North America (Panama).

Hosts: Tree branch.

References: (ds) Wood, S. L. 1982b: 1205. (tx) Wood, S. L. 1974a: 66, 1982b: 1205.

parilis Wood 1975a: 31. Holotype ♂; 6 km W Tepic, Nayarit, Mexico, 1000 m; Wood Collection.

Distribution: North America (Nayarit in Mexico).

Hosts: Tree branch.

References: (ds) Wood, S. L. 1982b: 1202. (tx) Wood, S. L. 1975a: 31, 1982b: 1202–1203.

plaumanni Schedl 1978c: 305. Holotype, sex?; Brasilien, Nova Tentonia, 300–400 m, 27 11 Br., 52 23 L; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1978c: 305

politus Eichhoff 1869a: 276. Holotype, sex?; Nov. Friburg; IRSNB, Brussels.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783; Blandford 1905: 289; Gemminger & Harold 1872: 2692; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 337; Schedl 1973d: 160. (tx) Blandford 1905: 289; Eichhoff 1869a: 276, 1878b: 467; Hagedorn 1910a: 139.

propugnatus Blandford 1905: 297. Holotype ♀; Cerro Zunil, Quezaltenango, Guatemala; BMNH, London.

Figures: Blandford 1905: pl. 9.

Distribution: North America (Costa Rica/ Guatemala).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 157; Wood, S. L. 1982b: 1203. (tx) Blandford 1905: 297; Hagedorn 1910a: 139; Schedl 1934c: 38; Wood, S. L. 1967d: 55, 1982b: 1203.

armatus Schedl 1934c: 37. Holotype ♀; Turrialba, Costa Rica; Schedl Collection in NHMW, Wien. Synonymy: Wood 1982b: 1203–1204.

References: (ds) Blackwelder 1947: 783. (tx) Schedl 1934c: 37–38, 1979c: 26; Wood, S. L. 1982b: 1203.

quadrinaculatus Schedl 1966f: 127. Holotype, sex?; Chanchamajo, Peruvia; Schedl Collection in NHMW, Wien.

Distribution: South America (Peru).

References: (ds) Schedl 1966f: 92. (tx) Schedl 1966f: 127, 1979c: 208.

rameus Wood 1967d: 55. Holotype ♀; 5 km W El Salto, Durango, Mexico, 2500 m; Wood Collection.

Figures: Wood 1967d: 54, 1982b: 1233 (declivity).

Distribution: North America (Durango in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 1196. (ds) Wood, S. L. 1982b: 1196. (tx) Wood, S. L. 1967d: 54–55, 1982b: 1196, 1233.

rasilis Schedl 1950i: 174. Syntypes ♀; Brazil, Santa Catharina, Nova Tentonia; Schedl Collection in NHMW, Wien, and Plaumann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 210 (citation of holotype invalid).

References: (ds) Pedrosa-Macedo & Schonherr 1985: 56; Schedl 1960f: 92, 1967d: 4, 1970g: 46. (tx) Pedrosa-Macedo & Schonherr 1985: 56; Schedl 1950i: 174, 1979c: 210.

retusus Eichhoff 1869a: 276. Holotype, sex?; Amer. mer. (Cayenne); IRSNB, Brussels.

Distribution: South America (Brazil/ Cayenne/ Ecuador).

References: (ds) Blackwelder 1947: 783; Blandford 1905: 291; Gemminger & Harold 1872: 2692; Hagedorn 1910d: 95; Kleine 1913b: 157, 1914b: 337–338. (tx) Blandford 1904: 289, 291; Eichhoff 1869a: 276, 1878b: 466; Hagedorn 1910a: 139.

schaufussi Blandford 1905: 293. Holotype, sex?; Venezuela; Hamburg Museum, lost.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 158, 1914b: 340. (tx) Blandford 1905: 290, 293; Hagedorn 1910a: 139; Wood, S. L. 1982b: 1199.

- speciosus (Schedl)** 1934c: 38 (*Tricolus*). Holotype, sex?; Hamburgfarm on Rio Reventazon. Limon, Costa Rica; Schedl Collection in NHMW, Wien. Distribution: North America (Costa Rica). Hosts: Tree. References: (ds) Blackwelder 1947: 783; Wood, S. L. 1982b: 1199. (tx) Schedl 1934c: 58, 1979c: 233; Wood, S. L. 1967c: 140, 1982b: 1199.
- spectabilis (Wood)** 1967c: 140 (*Tricolus*). Holotype ♀; Finca La Lola. Limon, Costa Rica; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Theobroma cacao*. References: (hb) Wood, S. L. 1982b: 1199. (ds) Wood, S. L. 1982b: 1199. (tx) Wood, S. L. 1967c: 140, 1982b: 1199–1200.
- spectus Wood** 1980b: 358. Holotype ♂; Pichucalco, Chiapas, Mexico; Wood Collection. Distribution: North America (Chiapas, Veracruz in Mexico). Hosts: *Theobroma cacao*, *Zygia* sp. References: (ds) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 1200. (tx) Wood, S. L. 1980b: 358, 1982b: 1200.
- spinachius (Schedl)** 1939j: 584 (*Staganocranus*). Holotype, sex?; Brasilien; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939j: 584, 1950i: 177, 1979c: 234.
- spinescens Wood** 1974a: 65. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica, 30 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree limb. References: (hb) Wood, S. L. 1982b: 1204. (ds) Wood, S. L. 1982b: 1204. (tx) Wood, S. L. 1974a: 65, 1982b: 1204.
- spinus Wood** 1974a: 65. Holotype ♀; Rincon de Osa, Puntarenas, Costa Rica, 30 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree limb. References: (hb) Wood, S. L. 1982b: 1204. (ds) Wood, S. L. 1982b: 1204. (tx) Wood, S. L. 1974a: 65, 1982b: 1204–1205.
- splendens Wood** 1982a: 223. Holotype ♂; San Rafael, Mexico, Mexico; Wood Collection. Distribution: North America (Mexico in Mexico). Hosts: *Quercus* sp. References: (hb) Atkinson & Equihua 1985a: 101. (ds) Atkinson & Equihua 1985a: 101. (tx) Wood, S. L. 1982a: 223.
- stenodermus (Schedl)** 1963: 166 (*Pterocyclon*). Holotype, sex?; Cordoba [presumably in Veracruz, Mexico]; Schedl Collection in NHMW, Wien. Distribution: North America (Nayarit, Veracruz in Mexico). Hosts: Tree limb. References: (hb) Wood, S. L. 1982b: 1193. (ds) Wood, S. L. 1982b: 1193. (tx) Schedl 1963c: 166, 1979c: 237; Wood, S. L. 1982b: 1193.
- tenuis Blandford** 1905: 295. Holotype ♀; “Mexican” tobacco refuse [probably from Brazil]; BMNH, London. Figures: Blandford 1905: pl. 9. Distribution: South America (Brazil/Venezuela). Notes: All of Blandford’s species named from Mexican tobacco refuse are actually from Brazil. References: (cc) Wichmann 1955a: 107. (hb) Wood, S. L. 1982b: 1197. (ds) Blackwelder 1947: 783; Ferrer 1942; Hagedorn 1910d: 95; Kleine 1913b: 158, 1914b: 354. Wichmann 1955a: 107; Wood, S. L. 1982b: 1197. (tx) Blandford 1905: 295; Hagedorn 1910a: 139; Schedl 1934c: 37, 1940a: 359, 1950i: 174; Wood, S. L. 1982b: 1197, 1202.
- terebella Blandford** 1905: 296. Holotype ♂?; Volcan de Chiriqui, Panama; BMNH, London. Figures: Blandford 1905: pl. 9. Distribution: North America (Panama), South America (Venezuela). Hosts: *Nectandra* sp., Guttiferae (near *Vismia* sp.). References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 158. (tx) Blandford 1905: 296; Hagedorn 1910a: 139.
- theobroma Sampson** 1912: 245. Holotype, sex?; Trinidad; BMNH, London. Distribution: Antilles Islands (Trinidad). Hosts: *Theobroma cacao*. References: (ds) Blackwelder 1947: 783; Kleine 1913b: 158, 1914b: 344. (tx) Sampson 1912: 245.
- thoracicus Erichson** 1836: 64. Holotype, sex?; Brasilien; MNB, Berlin. Distribution: South America (Brazil/ Guyana/ Guyana). References: (ds) Blackwelder 1947: 783; Blandford 1905: 290; Gemminger & Harold 1872: 2692; Hagedorn 1910d: 95; Kleine 1914b: 337–338; Lacordaire 1866: 383; Nunberg 1962: 223; Schedl 1966f: 79, 1967d: 4. (tx) Blandford 1905: 289–290; Castelnau 1840; Erichson 1836: 64; Ferrari 1867a: 48; Hagedorn 1910a: 139; Hopkins 1914: 116; Lacordaire 1866: 383; Lucas 1920: 90. *nitidulus* Guerin-Meneville 1835b: 107 (*Piezorhopalus*). Holotype, sex?; Brazil; not located. Synonymy: Blandford 1904: 290. References: (ds) Gemminger & Harold 1872: 2692; Lacordaire 1866: 383. (tx) Blandford 1904: 290; Eichhoff 1878b: 464; Ferrari 1867: 48; Guerin-Meneville 1835b: 107; Hopkins 1914: 127; Lacordaire 1866: 383.
- tornatilis Wood** 1974a: 64. Holotype ♀; 6 km S San Vito, Puntarenas, Costa Rica; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree. References: (ds) Wood, S. L. 1982b: 1195. (tx) Wood, S. L. 1974a: 64, 1982b: 1195–1199.

torneutes Blandford 1905: 292. Holotype, sex?; Chilpancingo, Guerrero, Mexico; BMNH, London. Figures: Blandford 1905: pl. 9. Distribution: North America (Guerrero at 1300 m in Mexico).

References: **(ds)** Blackwelder 1947: 753; Ferrer 1942; Hagedorn 1910d: 95; Kleine 1914b: 354; Wood, S. L. 1982b: 1201. **(tx)** Blandford 1905: 292; Hagedorn 1910a: 139; Schedl 1940a: 359; Wood, S. L. 1967d: 55, 1982b: 1201.

truncatorus Schedl 1950i: 175. Syntypes ♂; Brazil, Nova Teutonia; Schedl Collection in NHMW, Wien, and Plaumann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 257 (corrects spelling, citation of holotype invalid).

References: **(ds)** Schedl 1967d: 4; Wood, S. L. 1982b: 1201. **(tx)** Schedl 1950i: 175, 1979c: 256; Wood, S. L. 1982b: 1201.

ursus Schedl 1934c: 37. Holotype ♀?; San Jose, Costa Rica, 1000–1500 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: **(ds)** Blackwelder 1947: 753; Wood, S. L. 1982b: 1195. **(tx)** Schedl 1934c: 37, 1979c: 262; Wood, S. L. 1982b: 1198.

vagabundus (Schedl) 1966f: 124 (*Cosmocorynus*). Holotype ♀; Bolivien; Schedl Collection in NHMW, Wien.

Distribution: South America (Bolivia).

References: **(ds)** Schedl 1966f: 50. **(tx)** Schedl 1966f: 80, 124, 1979c: 263.

Genus *Gnatharus* Wood & Yin

GNATHARUS WOOD & YIN 1986: 463. Type-species: *Gnatharus tibetensis* Wood & Yin, original designation.

Notes: (3) This is the only member of the subtribe Corthylina that occurs naturally in an area outside of America.

References: **(hb)** Wood, S. L. 1986a: 99. **(ds)** Wood, S. L. 1986a: 99. **(tx)** Wood, S. L. 1986a: 99; Wood, S. L. & Yin 1986: 463.

tibetensis Wood & Yin 1986: 463. Holotype ♀; Medog, Xizang [Tibet], China; IZAS, Beijing. Distribution: Asia (Xizang [Tibet] in China). References: **(tx)** Wood, S. L. & Yin 1986: 463.

Genus *Monarthrum* Kirsch

MONARTHNUM KIRSCH 1866: 213. Type-species: *Monarthrum chapuisi* Kirsch, monobasic.

Corthylomimus Ferrari 1867a: 48. Type-species: *Bostrichus fasciatus* Say, subsequent designation by Hopkins 1914: 118. Synonymy: Eichhoff 1869c: 300.

References: **(tx)** Eichhoff 1869c: 297–301; Ferrari 1867a: 48; Hagedorn 1910d: 92; Hopkins 1914: 118; LeConte 1876: 153.

Cosmocorynus Ferrari 1867a: 62. Type-species:

Cosmocorynus cristatus Ferrari, monobasic. Synonymy: LeConte 1876: 348.

References: **(tx)** Eggers 1937a: 85; Eichhoff 1869c: 297–301; Ferrari 1867a: 62; Hagedorn 1910d: 92; LeConte 1876: 348; Wood, S. L. 1966b: 19.

Pterocyclon Eichhoff 1869a: 276. Type-species: *Pterocyclon laterale* Eichhoff, subsequent designation by Hopkins 1914: 125, neotype designated for type-species by Wood 1966b: 25. Synonymy: Eichhoff 1869c: 299.

References: **(cn)** Brooks 1917: 1–15; Drake 1921: 201–205; Swabey 1935: 1–35. **(ds)** Swaine 1909: 141. **(tx)** Beal & Massey 1945: 59, 102; Blackman 1922b: 76, 80; Blandford 1898c: 6, 1904: 268–286; Eggers 1935a: 75–77, 153; Eichhoff 1869a: 276, 1869c: 297–301, 1878b: 437; Hagedorn 1910a: 147–149; Schedl 1935h: 343, 1952k: 161, 1959m: 545–557; Swaine 1918a: 45, 86–87; Wood, S. L. 1966b: 19.

Anchonocerus Eichhoff 1878b: 67, 431. Type-species: *Anchonocerus rufipes* Eichhoff = *Pterocyclon ingens* Eichhoff, monobasic. Synonymy: Wood 1977d: 512.

References: **(tx)** Blandford 1905: 250, 266; Eggers 1935a: 329; Eichhoff 1878b: 67, 431; Hagedorn 1910d: 95; Wood, S. L. 1977d: 512.

Phthorinus Eichhoff 1878b: 67, 433. Type-species: *Pterocyclon ingens* Eichhoff, monobasic. Synonymy: Eggers 1935a: 329.

References: **(tx)** Eggers 1935a: 329; Eichhoff 1878b: 67, 433.

Trypocranus Eichhoff 1878b: 67, 435. Type-species: *Trypocranus cincinnatus* Eichhoff, monobasic. Synonymy: Wood 1982b: 1206.

References: Eichhoff 1878b: 67, 435; Eggers 1935a: 76, 329; Schedl 1952k: 161; Wood, S. L. 1982b: 1206.

Eupteroxylon Eggers 1936a: 392. Type-species: *Eupteroxylon comatum* Eggers, monobasic. Synonymy: Wood 1986a: 266.

References: **(tx)** Eggers 1936a: 392–393; Wood, S. L. 1986a: 266.

Keys: Blandford 1904: 269; Wood 1982b: 1207.

Notes: (3) Schedl 1939k: 726–727 (*Pterocyclon perduratum*, nomen nudum).

References: **(ay)** Nobuchi 1969a: 67. **(cn)** Ebeling & Pence 1953: 31; Howard 1897: 84–87; Hubbard 1897: 9–30; Snyder 1927: 1–46. **(hb)** Bright & Stark 1973: 94; Wood, S. L. 1982b: 1206, 1986a: 99. **(ds)** Bright & Stark 1973: 94; Swaine 1909: 142; Wood, S. L. 1982b: 1206, 1986a: 99. **(tx)** Arnett 1960: 1045, 1968: 1045; Bright 1972d: 101, 1976d: 188–189; Bright & Stark 1973: 94; Chamberlin 1939: 279–283, 1958: 128–129; Dodge 1938: 14, 18, 37; Eggers 1935a: 75–78, 1936a: 392; Eichhoff 1869c: 297–301; Ferrari 1867a: 104–115; Kirsch 1866: 213, 1868: 214; LeConte 1876: 347; LeConte & Horn 1883: 517; Swaine 1918a: 86; Wood, S. L. 1964: 19, 1966b:

- 19, 1967d: 37–57, 1977d: 512, 1982b: 1206, 1986a: 99.
- aberrans** Schedl 1977e: 46. Holotype ♂; El Salvador, Metapan, 1600–2200 m; Schedl Collection in NHMW, Wien.
Distribution: North America (El Salvador).
Hosts: *Quercus* sp.
References: (tx) Schedl 1977e: 46.
- abruptum** (Nunberg) 1962: 233 (*Microcorthyus*). Holotype ♂; Estacao Biologica de Boraceia, Salesopolis, Estado de Sao Paulo, Brasilien; DZSA, Sao Paulo.
Distribution: South America (Brazil).
References: (tx) Nunberg 1962: 233.
- adjunctum** (Schedl) 1967d: 15 (*Pterocyclon*). Holotype ♀; Brasilien, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1967d: 15–16, 1979c: 12.
- amphicranoides** (Schedl) 1963d: 225 (*Pterocyclon*). Holotype, sex?; Brasilien: Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1963c: 166, 1963d: 225, 1979c: 18.
- appendicinum** (Schedl) 1967d: 14 (*Pterocyclon*). Holotype ♂; Brasilien, Nova Teutonia, Santa Catarina; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Schedl 1976a: 56. (tx) Schedl 1967d: 14–15.
- bicallosum** (Schedl) 1939j: 577 (*Pterocyclon*). Holotype, sex?; Brasilien; Schedl Collection in NHMW, Wien.
Figures: Schedl 1939j: 578.
Distribution: South America (Brazil).
Hosts: *Cedrela fissilis*.
References: (ds) Blackwelder 1947: 782; Schedl 1966f: 91, 1976a: 56; Schouherr & Pedrosa-Macedo 1981: 53. (tx) Schedl 1939j: 577–578, 1950i: 164–165, 1951h: 285, 1979c: 37.
- bicavum** Wood 1967d: 51. Holotype ♀; Laguna Santa Maria, Nayarit, Mexico, 1000 m; Wood Collection.
Figures: Wood 1967a: 46, 1982b: 1219 (declivity).
Distribution: North America (Nayarit in Mexico).
Hosts: *Quercus* sp.
References: (hb) Wood, S. L. 1982b: 1234. (ds) Wood, S. L. 1982b: 1234. (tx) Wood, S. L. 1967a: 46, 1967d: 51–52, 1982b: 1234.
- bicolor** (Ferrari) 1867a: 56 (*Corthyus*). Holotype ♀; Venezuela [presumably Colonia Tovar]; NHMW, Wien.
Distribution: South America (Venezuela).
Hosts: *Clusia* sp., etc.
References: (ds) Blackwelder 1947: 782; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 92; Kleine 1914b: 340. (tx) Ferrari 1867a: 54, 56; Wood, S. L. 1974d: 283.
- signatus** Ferrari 1867a: 56 (*Corthyus*). Syntypes ♂; Venezuela [presumably Colonia Tovar]; NHMW, Wien. Synonymy: Wood 1974d: 283.
References: (ds) Gemminger & Harold 1872: 2680. (tx) Ferrari 1867a: 54, 56; Wood, S. L. 1974d: 283.
- edentatus** Hagedorn 1903b: 549 (*Plithorius*). Holotype ♂; Colonia Tovar, Venezuela; NHMW, Paris. Synonymy: Wood 1974d: 284.
Notes: (3) Eggers 1935a: 82 (redescribed).
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 95; Kleine 1913b: 158, 1914b: 340. (tx) Eggers 1929e: 51, 1933b: 3, 1935a: 76–83; Hagedorn 1903b: 549, 1910a: 142; Wood, S. L. 1974d: 284.
- bicoloratum** Wood 1974d: 284. Holotype ♂; Mile 10 on Bartica-Potaro Road, British Guiana; BMNH, London, automatic.
Distribution: South America (Guyana).
References: (tx) Wood, S. L. 1974d: 284.
- bicolor** Wood 1968b: 4. Holotype ♂; Mile 10 on Bartica-Potaro Road, British Guiana; BMNH, London, preoccupied by Ferrari 1867.
References: (ds) Hagedorn 1910d: 92; Kleine 1913b: 156. (tx) Hagedorn 1910a: 148; Wood, S. L. 1968b: 4, 1974d: 284.
- bidens** (Blandford) 1904: 277 (*Pterocyclon*). Holotype ♂; Volcan de Chiriqui, Panama; BMNH, London.
Distribution: North America (Costa Rica/ Chiapas in Mexico/ Panama).
Hosts: *Conostegia oerstediana*.
References: (hb) Wood, S. L. 1982b: 1241. (ds) Blackwelder 1947: 782; Bright 1972a: 1385; Hagedorn 1910d: 92; Kleine 1913b: 156, 1914b: 367; Wood, S. L. 1982b: 1241. (tx) Blandford 1904: 277; Hagedorn 1910a: 148; Wood, S. L. 1982b: 1241.
- bidentatum** Wood 1974b: 142. Holotype ♂; 9 km NE Teziutlan, Puebla, Mexico; Wood Collection.
Distribution: North America (Puebla in Mexico).
Hosts: *Alnus* sp., and a tree limb.
References: (hb) Wood, S. L. 1982b: 1240. (ds) Wood, S. L. 1982b: 1240. (tx) Wood, S. L. 1974b: 142, 1982b: 1240.
- bifoveatum** Wood 1974b: 137. Holotype ♀; San Jose, San Jose, Costa Rica; Wood Collection.
Distribution: Antilles Islands (Dominican Republic in Hispanola [?]), North America (Costa Rica), South America (Venezuela).
Hosts: *Spondias purpurea*, and a tree limb.
References: (hb) Wood, S. L. 1982b: 1229. (ds) Bright 1985c: 176; Wood, S. L. 1982b: 1229. (tx) Bright 1985c: 176; Wood, S. L. 1974b: 137, 1982b: 1229.

- bispinum (Blandford)** 1905: 251 (*Pterocyclon*). Holotype ♂; Bugaba, Chiriqui, Panama, 800–1500 ft.; BMNH, London. Figures: Blandford 1905: pl. 9. Distribution: North America (Costa Rica/ Panama). Hosts: Log. References: (ds) Blackwelder 1947: 782; Hagedorn 1910d: 92; Kleine 1913b: 156, 1914b: 342, 367; Wood, S. L. 1982b: 1221. (tx) Blandford 1905: 251; Hagedorn 1910a: 148; Schedl 1952a: 459; Wood, S. L. 1982b: 1221.
- boliviensis Wood** 1989: 178. Holotype ♀; Bolivia, Yungas del Palmar, 2000 m; Schedl Collection in NHMW, Wien, automatic. Distribution: South America (Bolivia). References: (tx) Wood, S. L. 1989: 178.
- bolivianum Schedl** 1970e: 103 (*Cosmocorynus*). Holotype ♀; Bolivia, Yungas del Palmar, 2000 m; Schedl Collection in NHMW, Wien, preoccupied by Eggers 1935. References: (tx) Schedl 1970e: 103; Wood, S. L. 1989: 178.
- brasilensis (Schedl)** 1936i: 107 (*Anchonocerus*). Holotype ♂; Nova Teutonia, S. Catharina, Brasilien; Schedl Collection in NHMW, Wien. Figures: Pedrosa-Macedo & Schonherr 1985: 57. Distribution: South America (Brazil). Notes: (1) Schedl 1959m: 553 (to *Pterocyclon*, =*Monarthrum*). (3) Schedl 1959m: 553 (described female). References: (ds) Blackwelder 1947: 783; Pedrosa-Macedo & Schonherr 1985: 57; Schedl 1966f: 92, 1967d: 4, 1972g: 46, 1976a: 56. (tx) Pedrosa-Macedo & Schonherr 1985: 57; Schedl 1936i: 107, 1951h: 285, 1952k: 161, 1959m: 553, 1979c: 44.
- brittoni (Schedl)** 1970e: 101 (*Pterocyclon*). Holotype ♂; Jamaica: Corn Puss Gap, St. Thomas-Portland Parish line, 2200 ft.; Schedl Collection in NHMW, Wien. Figures: Bright 1972d: 59, 95. Distribution: Antilles Islands (Jamaica). References: (ds) Bright 1972d: 101, 1985c: 176. (tx) Bright 1972d: 59, 95, 101, 1985c: 176; Schedl 1957a: 194, 1970e: 101, 1979c: 45.
- brunneum (Eichhoff)** 1869a: 278 (*Pterocyclon*). Holotype, sex?; Columbia; Hamburg Museum, lost. Distribution: South America (Colombia). References: (ds) Blackwelder 1947: 782; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 92; Kleine 1914b: 342. (tx) Eichhoff 1869a: 278, 1869c: 301, 1878b: 452; Hagedorn 1910a: 145.
- carinatum Wood** 1974b: 139. Holotype ♂; Cerro de la Muerte, San Jose, Costa Rica, 2600 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree hole. References: (hb) Wood, S. L. 1982b: 1237. (ds)
- Wood, S. L. 1982b: 1237. (tx) Wood, S. L. 1974b: 139, 1982b: 1237.
- carinulum Wood** 1974b: 144. Holotype ♂; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Miconia caudata*, *Phoebe mexicana*, and a liana. References: (hb) Wood, S. L. 1982b: 1243. (ds) Wood, S. L. 1982b: 1243. (tx) Wood, S. L. 1974b: 144, 1982b: 1243.
- chapuisi Kirsch** 1866: 213. Holotype ♂; Bogota, Colombia; MNB, Berlin. Distribution: South America (Bolivia/ Colombia). Notes: (3) Eggers 1935a: 79 (redescribed). References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 92; Kleine 1914b: 342. (tx) Eggers 1929e: 51, 1935a: 76–83; Eichhoff 1869a: 277, 1869c: 299, 301, 1878b: 440; Ferrari 1967b: 105; Hagedorn 1910a: 148; Hopkins 1914: 125; Kirsch 1866: 213, 1868: 214; Wood, S. L. 1989: 175.
- bolivianum Eggers** 1935a: 80. Holotype ♀; Bolivia (Cochabamba); USNM, Washington. Synonymy: Wood 1989: 175. References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 7; Eggers 1935a: 80; Schedl 1950i: 176, 1958c: 1, 1979c: 42; Wood, S. L. 1989: 175.
- cinnatum (Eichhoff)** 1878b: 435 (*Trypocranus*). Holotype ♀; America septentrionalis (Bogota); Hamburg Museum, lost. Distribution: South America (Colombia). References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 92; Kleine 1914b: 342. (tx) Eggers 1935a: 77–78; Eichhoff 1878b: 435; Hagedorn 1910a: 148; Hopkins 1914: 131.
- comatum (Eggers)** 1936a: 393 (*Eupteroxylon*). Holotype ♀; Bolivia (Cochabamba); USNM, Washington. Distribution: South America (Bolivia). References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 9; Eggers 1936a: 393–394.
- consimile (Blandford)** 1904: 275 (*Pterocyclon*). Holotype ♀; Volcan de Chiriqui, Panama, 25–4000 ft.; BMNH, London. Distribution: North America (Costa Rica/ Panama). Hosts: *Ochroma* sp. References: (hb) Wood, S. L. 1982b: 1236. (ds) Blackwelder 1947: 782; Hagedorn 1910d: 92; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1236. (tx) Blandford 1904: 275; Hagedorn 1910a: 148; Wood, S. L. 1974d: 284, 1982b: 1236.
- pseudosulcatum Schedl** 1935h: 348 (*Pterocyclon*). Holotype ♀; Vara Blanca, Heredia, Costa Rica; Schedl Collection in NHMW, Wien. Synonymy: Wood 1974: 284. References: (ds) Blackwelder 1947: 783. (tx)

Schedl 1935h: 348, 1979c: 202; Wood, S. L. 1974d: 284.

conversum Wood 1974b: 138. Holotype ♂; 6 km W Tepic, Nayarit, Mexico, 1000 m; Wood Collection. Distribution: North America (Jalisco, Nayarit in Mexico).

Hosts: Tree limb.

References: (hb) Wood, S. L. 1982b: 1236. (ds) Atkinson & Equihua 1985c: 362; Wood, S. L. 1982b: 1236. (tx) Wood, S. L. 1974b: 138, 1982b: 1236.

corculum Wood 1974b: 146. Holotype ♂; Turrialba, Cartago, Costa Rica, 700 m; Wood Collection. Distribution: North America (Costa Rica).

Hosts: *Theobroma cacao*, and a liana.

References: (hb) Wood, S. L. 1982b: 1245. (ds) Wood, S. L. 1982b: 1245. (tx) Wood, S. L. 1974b: 146, 1982b: 1245.

cordatum (Blandford) 1904: 279 (*Pterocyclon*). Holotype ♀; Quiche Mts., Guatemala; BMNH, London.

Distribution: North America (Guatemala/ Panama). Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 1223. (ds) Blackwelder 1947: 782; Hagedorn 1910d: 92; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1223. (tx) Blandford 1904: 279; Hagedorn 1910a: 148; Schedl 1935h: 349, 1952a: 460; Wood, S. L. 1982b: 1223.

corditicum Wood 1974b: 135. Holotype ♂; 30 km E Tulancingo, Hidalgo, Mexico, 2300 m; Wood Collection.

Distribution: North America (Hidalgo, Puebla in Mexico).

Hosts: *Quercus* sp.

References: (hb) Wood, S. L. 1982b: 1222. (ds) Atkinson & Equihua 1988: 92; Wood, S. L. 1982b: 1222. (tx) Wood, S. L. 1974b: 135, 1982b: 1222.

costatum (Eggers) 1937a: 85 (*Brachyspartus*). Syn-types, sex?; Brasilien (Bahia); BMNH, London. Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 784. (tx) Anderson, W. H. & Anderson 1971: 10; Eggers 1937a: 85–86; Schedl 1979c: 67.

cristatum (Ferrari) 1867a: 64 (*Cosmocorynus*). Holotype ♀; Venezuela [presumably Colonia Tovar]; NHMW, Wien.

Distribution: South America (Venezuela).

References: (ds) Gemminger & Harold 1872: 2693. (tx) Eggers 1937a: 84–85; Eichhoff 1868d: 419, 1869c: 301; Ferrari 1867a: 64; Hopkins 1914: 119; Schedl 1979c: 70; Wood, S. L. 1966b: 19. (ms) Eichhoff 1868d: 419.

dentatulum Wood 1989: 178. Holotype ♂; Bolivien (Cochabamba); USNM, Washington, automatic. Distribution: South America (Bolivia).

References: (tx) Wood, S. L. 1989: 178.

dentatum Eggers 1935a: 84. Holotype ♂; Bolivien

(Cochabamba); USNM, Washington, preoccupied by Eggers 1931.

References: (ds) Blackwelder 1947: 783; Schedl 1970e: 91; Wood, S. L. 1982b: 1216. (tx) Anderson, W. H. & Anderson 1971: 11; Eggers 1935a: 84; Schedl 1970e: 97, 1979c: 77; Wood, S. L. 1982b: 1216, 1989: 178.

dentatum (Eggers) 1931a: 19 (*Amphicranus*). Holotype ♂; Mineral del Chico, Hidalgo, Mexico; MNB, Berlin.

Distribution: Antilles Islands (Guadeloupe), North America (Hidalgo in Mexico).

References: (ds) Blackwelder 1947: 783; Schedl 1970e: 91. (tx) Eggers 1931a: 19; Schedl 1940a: 359; Wood, S. L. 1982b: 1216.

denticulatum Wood 1981: 122. Holotype ♂; Guadeloupe (Trois Rivières); Eggers Collection, in NHMW, Wien, automatic.

Distribution: Antilles Islands (Guadeloupe).

References: (ds) Bright 1985c: 176. (tx) Bright 1985c: 176; Wood, S. L. 1981: 122.

dentatum Eggers 1941a: 101 (*Pterocyclon*). Holotype ♂; Guadeloupe (Trois Rivières); Eggers Collection, in NHMW, Wien, preoccupied by Eggers 1931.

Notes: (3) Schedl 1970e: 97 (described female).

References: (tx) Eggers 1941a: 101; Schedl 1940a: 359, 1970e: 97.

dentigerum (LeConte) 1868: 154 (*Cryphalus*). Holotype ♂; middle California [USA]; MCZ, Cambridge.

Figures: Bright & Stark 1973: 163.

Distribution: North America (Baja California Norte in Mexico/ Arizona, California, Oregon, W Texas in USA).

Hosts: *Quercus agrifolia*, *Q.* spp.

Notes: (1) The original (masculine) spelling was *dentiger*.

References: (ay) Francke-Grosman 1956b. (cn) Chamberlin 1939: 281; Doane & Gilliland 1929: 918–920; Doane et al. 1936; Keen 1935: 147; Linsley & MacLeod 1942: 601; Michelbacher & Ortega 1958: 62; Swaine 1918a: 86. (cc) Chamberlin 1939: 281; Francke-Grosman 1956b; Keen 1935: 147; Slaby 1947: 377; Steinhaus 1946: 404; Webb 1945: 65. (hb) Bright & Stark 1973: 95; Chamberlin 1939: 281; Chittenden 1893a: 393; Doane & Gilliland 1929; Doane et al. 1936; Keen 1938: 147; Swaine 1918a: 86; Wood, S. L. 1982b: 1229. (ds) Bright & Stark 1973: 95; Chamberlin 1939: 281, 1955: 128; Furniss, R. L. & Carolin 1977: 398; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 92; Henshaw 1882: 268, 1885: 147; Keen 1929a: 28, 1938: 147; Kleine 1913b: 157, 1914b: 391, 1934a: 171; Leng 1920: 339; Patterson & Hatch 1945: 152; Rivers 1886: 66; Swaine 1909: 142; Wood, S. L. 1959a: 62, 1972a: 425, 1982b: 1229. (tx) Bright & Stark 1973: 163; Chamberlin 1939: 281; Eichhoff 1878b: 449;

- Hagedorn 1910a: 148; Keen 1929a: 28; LeConte 1868: 154, 1876: 348–349; Swaine 1909: 142, 1918a: 86; Wood, S. L. 1959a: 62, 1972a: 428, 1982b: 1229.
- desum** (Wood) 1967d: 52 (*Microcorthyus*). Holotype ♂; 25 km E Morelia, Michoacan, Mexico, 2500 m; Wood Collection.
- Figures: Wood 1967d: 46, 1982b: 1219 (declivity).
Distribution: North America (Chihuahua, Michoacan in Mexico).
Hosts: *Quercus* spp.
References: (hb) Atkinson & Equihua 1985a: 101; Atkinson et al. 1986: 49; Burgos & Saucedo 1983: 136; Wood, S. L. 1982b: 1238. (ds) Atkinson & Equihua 1985a: 101, 1985c: 362; Atkinson et al. 1986: 49; Burgos & Saucedo 1983: 136; Wood, S. L. 1982b: 1238. (tx) Wood, S. L. 1967d: 46, 52, 1982b: 1238.
- difficile** (Blandford) 1904: 276 (*Pterocyclon*). Holotype ♀; Volcan de Chiriqui, Chiriqui, Panama; BMNH, London.
Distribution: North America (Costa Rica/ Panama).
Hosts: Small tree.
References: (ds) Blackwelder 1947; Hagedorn 1910d: 92; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1239. (tx) Blandford 1904: 276; Hagedorn 1910a: 148; Wood, S. L. 1982b: 1239.
- dimidiatum** (Ferrari) 1867a: 57 (*Corthyus*). Syn-types, sex?; Venezuela; NHMW, Wien.
Distribution: North America (Costa Rica/ Panama), South America (Colombia/ Venezuela).
Hosts: *Alexa imperitricia*, *Croton* sp., *Eschweilera corrugata*, etc.
References: (ds) Blackwelder 1947; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 90, 92; Kleine 1913b: 156–157, 1914b: 340, 1934a: 171; Wood, S. L. 1982b: 1227. (tx) Eichhoff 1869c: 301; Ferrari 1867a: 54, 57, 1867b: 105; Hagedorn 1910a: 145, 148; Kirsch 1868: 214; Schedl 1939k: 726–727; Wood, S. L. 1979b: 135, 1982b: 1227.
- moritz** Schedl 1939k: 727 (*Pterocyclon*). Holotype ♂; Venezuela, 1858, Moritz (Schedl 1979c: 159) [Moritz collected primarily near his home in Colonia Tovar]; Schedl Collection in NHMW, Wien. Synonymy: Wood 1979b: 135.
Notes: (3) Schedl 1939k: 727 (cites *perduratum* Schedl, nomen nudum, synonymy).
References: (tx) Schedl 1939k: 727, 1954b: 40, 1979c: 159; Wood, S. L. 1979b: 135.
- distans** (Schedl) 1970e: 97 (*Pterocyclon*). Holotype, sex?; Argentina, Misiones, Dept. Concept. Santa Maria; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina/ Brazil).
References: (ds) Schedl 1970e: 97, 1976a: 56. (tx) Schedl 1970e: 97, 1979c: 83.
- dubiosum** (Schedl) 1976a: 82 (*Pterocyclon*). Holotype, sex?; Carnaru, Pernambuco [Brazil]; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 82.
- duplocordatum** Eggers 1935a: 86. Holotype ♂; Bolivien (Cochabamba); MNHN, Paris.
Distribution: South America (Bolivia).
References: (ds) Blackwelder 1947: 783. (tx) Eggers 1935a: 86.
- durum** (Schedl) 1972e: 74 (*Pterocyclon*). Holotype, sex?; Brasilien, Represa, Rio Grande, Guanabara; Schedl Collection in NHMW, Wien.
Figures: Pedrosa-Macedo & Schonherr 1985: 74.
Distribution: South America (Brazil).
References: (ds) Pedrosa-Macedo & Schonherr 1985: 58. (tx) Pedrosa-Macedo & Schonherr 1985: 58, 74; Schedl 1972e: 74, 1979c: 86.
- egeum** (Blandford) 1904: 280 (*Pterocyclon*). Holotype ♀; San Juan, Verapaz, Guatemala; BMNH, London.
Distribution: North America (Costa Rica/ Guatemala), South America (Brazil/ Colombia).
Hosts: Leguminous trees.
References: (cc) Wichmann 1955a: 107. (hb) Atkinson & Equihua 1985c: 361; Wood, S. L. 1982b: 1226. (ds) Atkinson & Equihua 1985c: 361; Blackwelder 1947: 782; Hagedorn 1910d: 92; Kleine 1913b: 157, 1914b: 367; Schedl 1966f: 92, 1972g: 46, 1973d: 161; Wichmann 1955a: 107; Wood, S. L. 1982b: 1226. (tx) Blandford 1904: 280; Hagedorn 1910a: 148; Schedl 1935b: 347, 1939e: 68, 1939k: 726, 1940c: 208, 1950i: 167; Wood, S. L. 1974d: 284, 1982b: 1226.
- bisetosus** Schedl 1954b: 38 (*Brachyspartus*). Lectotype ♀; Rio Caragnata, Matto Grosso, Brazil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 41. Synonymy: Wood 1974d: 284.
References: (tx) de Ruelle 1970: 98; Schedl 1954b: 38, 1979c: 41; Wood, S. L. 1966b: 18, 24, 1974d: 284, 1979c: 41.
- elegans** (Eichhoff) 1869a: 277 (*Pterocyclon*). Holotype, sex?; Brasilia; Hamburg Museum, lost.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 92; Kleine 1913b: 157, 1914b: 337. (tx) Eggers 1935a: 78; Eichhoff 1869a: 277, 1869c: 301, 1878b: 437; Hagedorn 1910a: 148.
- eumerum** (Schedl) 1952a: 459 (*Pterocyclon*). Syn-types 2 ♂; Argentina, Misiones, Dep. Concep., Sta. Maria; 1 in Schedl Collection in NHMW, Wien, and 1 in Viana Collection.
Distribution: South America (Argentina).
Notes: (1) Schedl 1979c: 92 (citation of holotype invalid).
References: (tx) Schedl 1952a: 459, 1979c: 92.
- excavatus** (Eggers) 1935a: 330 (*Anchonocerus*). Holotype ♂; Bolivien (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia).
References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 13; Eggers 1935a: 330.

exornatum (Schedl) 1939j: 575 (*Pterocyclon*).

Holotype ♂; Colonia (vermutlich Mexico); Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica/ Veracruz in Mexico), South America (Colombia/ Venezuela).

Hosts: *Alexia imperitricia*, *Coffea arabica*, *Ficus* sp.

References: (hb) Wood, S. L. 1982b: 1237. (ds) Blackwelder 1947: 782; Ferrer 1942; Wood, S. L. 1982b: 1237. (tx) Schedl 1939j: 575, 1939k: 727, 1940a: 359, 1950a: 167, 1963c: 167, 1979c: 94; Wood, S. L. 1966b: 25, 1982b: 1237.

gracilicorumm Schedl 1939j: 576 (*Pterocyclon*).

Holotype ♂; Mexico, Jalapa; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 25.

References: (ds) Blackwelder 1947: 782; Ferrer 1942. (tx) Schedl 1939j: 576, 1939k: 727, 1940a: 359, 1979c: 107; Wood, S. L. 1966b: 25.

fasciatum (Say) 1826: 255 (*Bostrichus*). Syntypes, sex?, Sinepuxent Bay, New York [USA]; lost.

Figures: Blackman 1922b: pl. 10, fig. 53, Bright 1976d: 189, Roling & Kearby 1974: 1303, 1305.

Distribution: North America (Arkansas, Connecticut, District of Columbia, Florida, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Massachusetts, Michigan, Mississippi, Missouri, New Jersey, New York, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Texas, Virginia, West Virginia, Wisconsin in USA).

Hosts: *Acer rubrum*, *Carya* spp., *Castanea dentata*, *Liquidambar styraciflua*, *Mimosa* sp., *Nyssa* sp., *Pinus* sp., *Prunus* sp., *Quercus alba*, *Q. digitata*, *Q. rubra*, *Q. spp.*

References: (ay) Batra 1963b: 227; Hopkins 1894g; Lowe, R. E., Giese, & McManus 1967: 451; Roling & Kearby 1974; Schneider 1976. (bv) Atkinson, Foltz, & Connor 1988; Dethier 1947; Haack, R. A., Benjamin, & Haack 1983; Hosking & Knight 1975; Kovach 1986a; Roling & Kearby 1975a, 1975b, 1977; Turnbow & Franklin 1980. (cn) Anonymous 1960q, 1964h, 1973g, 1977f; Beal et al. 1952; Blackman 1950; Brooks 1916: 14, 1917: 1–15; Chamberlin 1939: 281; Christian 1939; Comstock 1880: 274–275; Davis 1895; Doane et al. 1936; Dorsey & Leech 1956: 220–224; Dorsey et al. 1953: 419–420; Felt 1906: 751; Forbes 1894: 23; Gagne & Kearby 1978: 1013; Gahan 1906: 81; Garman 1905: 68–74; Gossard 1911: 206, 1913: 61, 1914: 11; Herrick 1935: 43; Himelick & Curl 1958: 539–541; Hopkins 1899c: 347, 442, 1904: 45; Howard 1897: 85; Hubbard 1897b: 27–28; Kleine 1932a: 304; Kovach 1986a; Kovach & Gorsuch 1985; Lugger 1899a: 308; Packard 1890: 328–520; Quaintance & Siegler 1922: 66, 1931b: 75; Roling & Kearby 1973, 1977; Swaine 1918a: 86. (ec) Batra 1963b: 216; Batra & Michie 1963: 475; Chamberlin 1939: 281; Currie

1905; Felt 1906: 751; Haack, R. A., Benjamin, & Haack 1983; Hagedorn 1907c; Hirschmann 1972a; Jewell 1956: 251; Lowe, R. E., Giese, & McManus 1967: 451; Roling & Kearby 1974, 1977; Schneider 1963: 661, 1976; Slaby 1947: 378; Steinhaus 1946: 404, 1949: 92; Verrall 1943: 135–137, 142–143; Webb 1945: 68; Wertz, Skelly, & Merrill 1971. (hb) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 267; Batra 1963b: 221; Beal & Massey 1945: 102–104; Blackman 1922b, 1950; Chamberlin 1939: 281; Chittenden 1890, 1893a: 391; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Felt 1906: 751; Forbes 1894: 23; Haack, R. A., Benjamin, & Haack 1983; Herrick 1935: 43; Hopkins 1894g, 1899c: 442; Hubbard 1897b: 27; Kleine 1932a: 304; Kovach 1986a; Lugger 1899a: 308; Ostmark 1968; Packard 1890: 328; Pierce, W. D. 1907: 290; Roling & Kearby 1973, 1974a, 1974b; Swaine 1918a: 86. (ds) Anonymous 1926c: 517, 1964h, 1977f; Atkinson et al. 1991: 162; Beal & Massey 1945: 102–104; Beaulne 1956; Blackman 1922b: 81–82, 1950; Blandford 1895b; Blatchley & Leng 1916: 641; Bright 1976d: 191; Britton 1920a; Chamberlin 1925, 1939: 281; Chittenden 1890; Cola 1973; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66; Dodge 1938; Dolphin, Mouzin, & Cleveland 1972: 1600; Droz 1985: 370; Frost & Dietrich 1929; Gagne & Kearby 1978: 1013; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 92; Hamilton 1895a: 346, 378; Henshaw 1885: 147; Hoffmann 1940: 60, 1942: 12; Hopkins 1893a: 128, 1893b: 277; Hubbard & Schwarz 1878a: 665; Kirk 1969, 1970; Kleine 1913b: 157, 1914b: 399, 1932a: 304, 1934a: 171; Lacordaire 1866: 385; Leng 1920: 339; Leonard 1928: 517; Melsheimer 1806: 8, 1853: 87; Ostmark 1968; Roling & Kearby 1974; Schneider & Farner 1969: 413; Smith, J. B. 1900: 361, 1910: 401; Swaine 1909: 143; Turnbow & Franklin 1980; Wood, S. L. 1982b: 1235. (tx) Beal & Massey 1945: 102–104; Blackman 1922b: 81–82; Blandford 1895b; Blatchley & Leng 1916: 641; Bright 1976d: 189, 191; Chamberlin 1939: 281; Dodge 1938: 28; Eggers 1935a: 75; Eichhoff 1868d: 419, 1869a: 277, 1869c: 297–301, 1878b: 442–444; Erichson 1836: 64, 79; Ferrari 1867a: 48–49; Hagedorn 1910a: 148; Hopkins 1914: 118; Hubbard 1897b: 27; Jacques 1951: 352; Lacordaire 1866: 385; LeConte 1868: 153, 1876: 348, 1878a: 468, 665; Lucas 1920: 556; Motschulsky 1856: 10; Roling & Kearby 1974: 1303, 1305; Say 1826: 253–255; Swaine 1909: 143, 1918a: 86; Wood, S. L. 1982b: 1235; Zimmermann 1868: 143. (ms) Eichhoff 1868d: 419; Lucas 1920: 556.

simile Eichhoff 1869a: 277 (*Pterocyclon*). Syntypes, sex?, America bor. [USA]; Hamburg Museum, lost. Synonymy: LeConte 1876: 348. References: (ds) Gemminger & Harold 1872: 2680; Leng 1920: 339; Swaine 1909: 142. (tx)

- Eichhoff 1869a: 277. 1869c: 301; LeConte 1876: 348; Swaine 1909: 142, 1915a: 86.
gracile Eichhoff 1875b: 444 (*Pterocyclon*). Syn-
 types, sex?, Tennessee, Pennsylvania, Carolina
 [USA]; Hamburg Museum, lost. Synonymy:
 Eichhoff 1875b: 444.
 References: (ds) Hagedorn 1910d: 93; Kleine
 1913b: 157, 1914b: 393; Leng 1920: 339;
 Swaine 1909: 142. (tx) Eichhoff 1875b: 444;
 Hagedorn 1910a: 148; Swaine 1909: 142.
- fastigiorum* Wood 1974b: 141. Holotype ♀;
 Tapanti, Cartago, Costa Rica, 1300 m; Wood
 Collection.
 Distribution: North America (Costa Rica).
 Hosts: Liana.
 References: (ds) Wood, S. L. 1982b: 1240. (tx)
 Wood, S. L. 1974b: 141, 1982b: 1240.
- fenestratum* Eggers 1935a: 85. Holotype ♀;
 Bolivien (Cochabamba); MNIN, Paris.
 Distribution: South America (Bolivia).
 References: (ds) Blackwelder 1947: 783. (tx)
 Eggers 1935a: 85, 1979c: 95.
- ferrarii* (Blandford) 1905: 284 (*Pterocyclon*). Holo-
 type ♀; Volcan de Chiriqui, Panama; BMNH,
 London.
 Figures: Blandford 1905: pl. S.
 Distribution: North America (Veracruz in Mexico/
 Panama), South America (Venezuela).
 References: (ds) Blackwelder 1947: 782; Hage-
 dorn 1910d: 92; Kleine 1913b: 157, 1914b: 340,
 367. (tx) Blandford 1905: 284; Hagedorn 1910a:
 148; Wood, S. L. 1982b: 1233.
- fimbriaticorne* (Blandford) 1905: 285 (*Ptero-
 cyclon*). Holotype ♀; Purula, Alta Verapaz, Gua-
 temala; BMNH, London.
 Figures: Blandford 1905: pl. S.
 Distribution: North America (Costa Rica/ Guate-
 mala/ Veracruz in Mexico), South America (Brazil/
 Venezuela).
 Hosts: *Erythrina costaricensis*, *Parkinsonia
 aculeata*, *Spondias purpurea*, *Theobroma cacao*.
 References: (hb) Wood, S. L. 1982b: 1232. (ds)
 Atkinson & Equihua 1985c: 362; Blackwelder
 1947; Bright 1972a: 1385; Hagedorn 1910d: 93;
 Kleine 1913b: 157, 1914b: 367; Wood, S. L.
 1982b: 1232. (tx) Blandford 1905: 285; Hagedorn
 1910a: 148; Wood, S. L. 1974d: 285, 1982b: 1232.
- turbinatum* Schedl 1961i: 230 (*Pterocyclon*).
 Holotype ♂; Cordoba, Veracruz, Mexico; C.A.S.,
 San Francisco. Synonymy: Wood 1974d: 285.
 References: (ds) Schedl 1960a: 75, 1961i: 230,
 1976a: 56; Wood, S. L. 1974d: 285. (tx) Schedl
 1961i: 230, 1979c: 259.
- flohri* (Schedl) 1950i: 168 (*Pterocyclon*). Holotype
 ♀; Mexico; Schedl Collection in NHMW, Wien.
 Distribution: North America ("Mexico").
 References: (ds) Wood, S. L. 1982b: 1228. (tx)
 Schedl 1950i: 168, 1979c: 95; Wood, S. L. 1982b:
 1228.
- fulgens* Schedl 1971f: 153. Holotype ♂; Ecuador:
 Quito; UZMC, Copenhagen.
 Distribution: South America (Ecuador).
 References: (tx) Schedl 1971f: 153, 1979c: 101.
- gibber* (Schedl) 1952a: 460 (*Pterocyclon*). Holotype
 ♀; Argentina, Concepcion; Eggers Collection, in
 NHMW, Wien.
 Distribution: South America (Argentina/ Brazil).
 References: (ds) Schedl 1972g: 46. (tx) Schedl
 1952a: 460, 1979c: 104.
- glabriculum* (Schedl) 1976a: 82 (*Pterocyclon*).
 Holotype, sex?, Brazil: Represa Rio Grande,
 Guanabara; Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (tx) Schedl 1976a: 82.
- glabrifrons* (Blandford) 1904: 278 (*Pterocyclon*).
 Lectotype ♀; Quiche Mts., Guatemala, 7000–
 9000 ft.; BMNH, London.
 Distribution: North America (Guatemala).
 References: (ds) Blackwelder 1947: 782; Hage-
 dorn 1910d: 118; Kleine 1913b: 157, 1914b: 367;
 Wood, S. L. 1982b: 1243. (tx) Blandford 1904:
 278; Hagedorn 1910a: 148; Wood, S. L. 1982b:
 1243.
- gnarum* (Schedl) 1950i: 169 (*Pterocyclon*). Holo-
 type ♀; Mexico; Schedl Collection in NHMW,
 Wien.
 Figures: Bright 1972a: 1384.
 Distribution: North America (Chiapas, Hidalgo,
 Oaxaca in Mexico).
 Hosts: *Quercus* sp.
 References: (hb) Wood, S. L. 1982b: 1220. (ds)
 Wood, S. L. 1982b: 1220. (tx) Schedl 1950i: 169,
 1979c: 106; Wood, S. L. 1974d: 285, 1982b: 1220.
- spinatus* Bright 1972a: 1383 (*Amphicranus*).
 Holotype ♂; 51 km S Valle Nacional, Oaxaca,
 Mexico; CNCI, Ottawa. Synonymy: Wood
 1974d: 285.
 References: (tx) Bright 1972a: 1383–1384;
 McNamara 1977: 194; Wood, S. L. 1974d: 285.
- gracilentum* (Schedl) 1972e: 75 (*Pterocyclon*).
 Holotype, sex?, Brasilien, Jacareacanga, Para;
 Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (tx) Schedl 1972e: 75, 1979c: 107.
- gracilior* (Schedl) 1959m: 553 (*Pterocyclon*). Holo-
 type ♂; Brazil: Nova Teutonia; Schedl Collection
 in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (tx) de Ruette 1970: 113; Schedl
 1959m: 553, 1979c: 107.
- granulatum* Bright 1972a: 1382. Holotype ♂; 24
 km S Valle Nacional, Oaxaca, Mexico, 1300 m;
 CNCI, Ottawa.
 Distribution: North America (Oaxaca, Puebla in
 Mexico).
 Hosts: Tree branch.
 References: (ds) Wood, S. L. 1982b: 1244. (tx)

- Bright 1972a: 1377, 1382; McNamara 1977: 196; Wood, S. L. 1982b: 1244.
- hagedorni** (Schedl) 1939k: 727 (*Pterocyclon*). Holotype, sex?; Campoi Guyanae; MNHN, Paris, automatic. Distribution: South America (Guyana). References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939k: 727.
- dimidiatum** Hagedorn 1903b: 550 (*Pterocyclon*). Holotype, sex?; Campoi Guyanae; MNHN, Paris, preoccupied by Ferrari 1867. References: (tx) Hagedorn 1903b: 550; Schedl 1939k: 727.
- hoegei** (Blandford) 1904: 274 (*Pterocyclon*). Holotype, sex?; Jalapa, Veracruz, Mexico; BMNH, London. Figures: Bright 1972a: 1377. Distribution: North America (Oaxaca, Puebla in Mexico). Hosts: *Alnus* sp., *Miconia* sp. References: (hb) Wood, S. L. 1982b: 1225. (ds) Blackwelder 1947: 783; Ferrer 1942; Hagedorn 1910d: 93; Kleine 1913b: 157, 1914b: 353; Wood, S. L. 1982b: 1225. (tx) Blandford 1904: 274; Hagedorn 1910a: 148; Schedl 1940a: 359; Wood, S. L. 1982b: 1225.
- oaxacaensis** Bright 1972a: 1382. Holotype ♂; 51 km S Valle Nacional, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1982b: 1225. References: (tx) Bright 1972a: 1377, 1382; McNamara 1977: 196; Wood, S. L. 1982b: 1225.
- huachucae** Wood 1959a: 61. Holotype ♂; Miller Canyon, Huachuca Mts., Arizona [USA]; USNM, Washington. Distribution: North America (Arizona in USA). Hosts: *Quercus hypoleucoides*. References: (ds) Furniss, R. L. & Carolin 1977: 398; Wood, S. L. 1959a: 61, 1982b: 1217. (tx) Wood, S. L. 1959a: 61, 1982b: 1217.
- infradentatum** Wood 1974b: 145. Holotype ♂; Rincon de Osa, Puntarenas, Costa Rica, 30 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree limb. References: (ds) Wood, S. L. 1982b: 1244. (tx) Wood, S. L. 1974b: 145, 1982b: 1244.
- ingens** (Eichhoff) 1869a: 278 (*Pterocyclon*). Holotype, sex?; Columbia; IRSNB, Brussels. Distribution: South America (Colombia/ Peru). References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Kleine 1913b: 155, 1914b: 342. (tx) Eggers 1935a: 329–334; Eichhoff 1869a: 278–279, 1869c: 301, 1878b: 433; Hopkins 1914: 127; Lucas 1920: 507. (ms) Lucas 1920: 507.
- rufipes** Eichhoff 1878b: 431 (*Anthonocerus*). Holotype ♀; America meridionalis (Nova Granada); IRSNB, Brussels. Synonymy: Eggers 1935a: 329.
- References: (ds) Hagedorn 1910d: 95; Kleine 1913b: 158, 1914b: 342. (tx) Eggers 1935a: 329; Eichhoff 1878b: 431; Hagedorn 1910a: 141; Hopkins 1914: 116; Lucas 1920: 94. (ms) Lucas 1920: 94.
- assequens** Schedl 1978c: 302 (*Pterocyclon*). Holotype, sex?; Peru, Dep. Cuzco, Marcapata-Thal. Nbl. d. Madre de Dios, 3000 m; Schedl Collection in NIIMW, Wien. Synonymy: Wood 1959: 175. References: (tx) Schedl 1978c: 302; Wood, S. L. 1959: 175.
- insignatum** Wood 1974b: 141. Holotype ♀; Volcan Poas, Heredia, Costa Rica, 2600 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Tree branch, and a shrub. References: (ds) Wood, S. L. 1982b: 1240. (tx) Wood, S. L. 1974b: 141, 1982b: 1240.
- insolitum** (Schedl) 1976a: 83 (*Pterocyclon*). Holotype, sex?; Brazil: Manaus, Amazonas; Schedl Collection in NIIMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1976a: 83.
- intermedium** Schedl 1970e: 105. Holotype ♀; Bolivien, Jungas, de Arepuchio, Chacisacho, ca 1500 m; ZSSM, München. Distribution: South America (Bolivia). References: (tx) Schedl 1970e: 105.
- laevigatum** (Eichhoff) 1869a: 278 (*Pterocyclon*). Holotype, sex?; Brasilia; Hamburg Museum, lost. Distribution: South America (Brazil). References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 93; Kleine 1913b: 157, 1914b: 337; Schedl 1972g: 46, 1976a: 56. (tx) Eichhoff 1869a: 278, 1869c: 301, 1878b: 449; Hagedorn 1910: 145; Schedl 1939j: 574, 1972g: 46.
- laterale** (Eichhoff) 1869a: 278 (*Pterocyclon*). Holotype ♂; Mexico; Hamburg Museum, lost, neotype: Tuxpan, Mexico. BMNH, London, designated by Wood 1966b: 25. Figures: Atkinson et al. 1986: 48. Distribution: North America (Costa Rica/ Guatemala/ Michoacan, Morelos, Puebla, Veracruz in Mexico), South America (Venezuela). Hosts: *Alnus* sp., *Quercus* sp., etc. References: (hb) Atkinson et al. 1986: 50; Burgos & Saucedo 1983: 134; Wood, S. L. 1982b: 1231. (ds) Atkinson et al. 1986: 50; Blackwelder 1947: 783; Blandford 1905: 251; Burgos & Saucedo 1983: 134; Ferrer 1942; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 93; Kleine 1913b: 157, 1914b: 353, 367; Wood, S. L. 1982b: 1231. (tx) Atkinson et al. 1986: 48; Blandford 1905: 251; Eggers 1929c: 51; Eichhoff 1869a: 278, 1869c: 301, 1878b: 439; Hagedorn 1910a: 145; Hopkins 1914: 128; Schedl 1935h: 351, 1936i: 108, 1940a: 359; Wood, S. L. 1966b: 25, 1982b: 1231.
- trifasciatus** Schedl 1950i: 173 (*Cosmocrinus*).

Holotype ♀: Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 25. References: (tx) Schedl 1950i: 173, 1979c: 255; Wood, S. L. 1966b: 25.

latum (Schedl) 1966f: 126 (*Cosmocorynus*). Holotype ♂; Unleserlich, vielleicht Maydes, vermutlich Columbien; Schedl Collection in NHMW, Wien. Distribution: South America (possibly Colombia). References: (ds) Schedl 1966f: 79. (tx) Schedl 1966f: 126, 1979c: 137.

limulum Wood 1974b: 143. Holotype ♂; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: Liana. References: (hb) Wood, S. L. 1982b: 1242. (ds) Wood, S. L. 1982b: 1242. (tx) Wood, S. L. 1974b: 143, 1982b: 242.

lobatum (Ferrari) 1867a: 57 (*Corthylus*). Holotype ♂; Venezuela; NHMW, Wien. Figures: Blandford 1905: pl. 8. Distribution: North America (Costa Rica/ Guatemala), South America (Venezuela). Hosts: *Clusia* sp., *Nectandra* sp., *Spondias purpurea*, etc. References: (hb) Wood, S. L. 1982b: 1230. (ds) Blackwelder 1947: 783; Blandford 1905: 283; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 93; Kleine 1913b: 157, 1914b: 340, 367; Wood, S. L. 1982b: 1230. (tx) Blandford 1905: 283; Ferrari 1867a: 54, 57; Hagedorn 1910a: 148; Wood, S. L. 1982b: 1230.

longus (Schedl) 1950i: 172 (*Anchonocerus*). Holotype ♀; Bolivia, Cochabamba; Eggers Collection, in NHMW, Wien. Distribution: South America (Bolivia). References: (tx) Schedl 1950i: 172, 1979c: 142.

luctuosum (Blandford) 1904: 276 (*Pterocyclon*). Holotype ♂; Cerro Zunil, Guatemala, 4000 ft.; BMNH, London. Distribution: North America (Guatemala/ Puebla in Mexico). Hosts: *Quercus* sp. References: (hb) Wood, S. L. 1982b: 1225. (ds) Atkinson & Equihua 1985c: 361, 1988a: 92; Blackwelder 1947: 783; Hagedorn 1910d: 93; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1225. (tx) Blandford 1904: 276; Hagedorn 1910a: 148; Wood, S. L. 1982b: 1225.

mali (Fitch) 1855: 326 (*Tomicus*). Syntypes, sex?; Middlefield, Massachusetts [USA]; not located. Figures: Blackman 1922b: pl. 10, fig. 54, Dillon & Dillon 1961: 801. Distribution: Antilles Islands (Cuba), North America (British Columbia in Canada, introduced/ Alabama, Arkansas, California, Connecticut, District of Columbia, Florida, Illinois, Indiana, Iowa, Kentucky, Louisiana, Maine,

Maryland, Massachusetts, Michigan, Minnesota, Mississippi, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Tennessee, E Texas, Virginia, Vermont, West Virginia, Wisconsin in USA). Hosts: *Acer rubrum*, *Betula lutea*, *Fagus americana*, *Liquidambar styraciflua*, *Nyassa* sp., *Quercus rubra*, *Tilia* sp.

Notes: (3) Three specimens from California and one from British Columbia have been examined; whether these represent errors in labeling, interceptions, or established populations is not yet known (SIW).

References: (ay) Hopkins 1894g; Schneider 1976. (bv) Atkinson, Foltz, & Connor 1988; Dethier 1947; Meixner 1937: 1218; Roling & Kearby 1975b, 1977; Turnbow & Franklin 1980. (cn) Anonymous 1976c; Beal et al. 1952; Blackman 1950; Brooks 1916: 13, 1917: 1–15; Chamberlin 1939: 282; Christian 1939; Collins & Connola 1955; Connola et al. 1956: 1–36; Craighead 1935: 139; Davis 1895; Doane et al. 1936; Felt 1901: 517–518, 1905: 289–292, 1906: 336, 1924: 277, 1926: 247–248, 277; Felt & Rankin 1932: 145; Fletcher 1888: 15; Garman 1905: 68–70; Gossard 1913: 62, 1914: 12; Herrick 1935: 375; Hopkins 1894b: 295, 1894g: 277, 1898d: 47, 1899c: 347, 442, 1904a: 45, 1905a: 384, 1907b, 1909c: 65, 1910g: 65; Howard 1897: 85; Hubbard 1897b: 26–28; Keen 1938: 147; Kleine 1932a: 304; Leach 1940b: 61; Lintner 1878: 455, 1896: 270; Lugger 1899a: 306, 308; Packard 1890: 93–94; Pierson 1927: 89; Quaintance & Siegler 1922: 66, 1931b: 75; Roling & Kearby 1973, 1977; Saunders 1883: 24; Schneider 1963: 661; Slingerland & Crosby 1914: 198; Swaine 1918a: 86–87; Wheeler 1922: 520. (ec) Batra 1963b: 216; Chamberlin 1939: 282; Currie 1905; Dorsey & Leach 1956: 220–224; Felt 1906: 289; Jewell 1956: 251; Leach 1940b: 61; Roling & Kearby 1977; Schneider 1976; Slaby 1947: 378; Steinhaus 1946: 404; Thompson, W. R. & Simmonds 1964: 30, 1965: 68; Verrall 1941: 552, 1943: 135–137, 142–143; Webb 1945: 68. (hb) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 267; Beal & Massey 1945: 103–104; Blackman 1922b: 80–81, 1950; Chamberlin 1939: 282; Clittenden 1890, 1893a: 392, 1897: 79; Dillon & Dillon 1961: 810; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Drake 1921: 201, 205; Drooz 1985: 370; Felt 1906: 289, 1926: 247–248, 277; Felt & Rankin 1932: 145; Fitch 1856: 5, 8–9; Herrick 1935: 375; Hoffmann 1941; Hopkins 1894g, 1899c: 442, 1904a: 45, 1905a: 384; Hubbard 1897b: 26; Kleine 1932a: 304; Lengerken 1954: 312; Lugger 1899a: 308; Morstatt 1924: 7, 40; Ostmark 1968; Packard 1890: 93; Pierce, W. D. 1907: 290; Pierson 1927: 89; Roling & Kearby 1973; Saunders 1883: 24; Skinner 1905: 248; Swaine 1907: 191, 1918a: 86, 87; Titus & Pratt 1904: 19; Wood, S. L. 1982b: 1234. (ds) Anonymous 1926c: 517, 1976c; Atkinson et al.

- 1991: 162; Beal & Massey 1945: 103–104; Beaulne 1956; Blackman 1922b: 80–81, 1950; Blackwelder 1947: 783; Blandford 1904: 279; Blatchley & Leng 1916: 641; Bright 1976d: 191, 204, 214, 1955c: 176; Britton 1920a; Chamberlin 1925, 1939: 282; Chittenden 1890; Currie 1905; Deyrup 1981b: 9; Deyrup & Atkinson 1987a: 66; Dillon & Dillon 1961: 801; Dodge 1938: 36, 39; Dolphin, Mouzin, & Cleveland 1972: 1600; Drake 1921: 201, 205; Drooz 1985: 370; Felt 1926: 247–248, 277, 1930a: 247–248; Felt & Rankin 1932: 145; Ferrer 1942; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 93; Hamilton 1895a: 378; Henshaw 1885: 147; Hoffmann 1942: 12; Hopkins 1893a: 128, 1893b: 208, 1894: 295; Kaston 1938: 240; Kleine 1913b: 157, 1914b: 378, 384, 1932a: 304, 1934a: 171; Knull 1924: 212; Leng 1920: 339; Leonard 1928: 517; Ostmark 1968; Pechuman 1937: 13; Schedl 1971f: 148; Schwarz 1878: 468, 1886: 44, 1888: 48; Smith, J. B. 1900: 361, 1910: 401; Swaine 1909: 143; Turnbow & Franklin 1950; Wood, S. L. 1982b: 1234. **(tx)** Beal & Massey 1945: 103–104; Benoit 1985: 167; Blackman 1922b: 80–81; Blandford 1904: 270–279; Blatchley & Leng 1916: 641; Bright 1976d: 189, 191, 204, 214, 1955c: 176; Chamberlin 1939: 282; Comstock 1948; Dillon & Dillon 1961: 810; Dodge 1938; Eichhoff 1876a: 378–379, 1878b: 447–449; Fitch 1855: 326; Hagedorn 1910a: 148; Hubbard 1897b: 26; Jacques 1951: 352; LeConte 1868: 154, 1876: 349, 1878a: 465, 468; Meixner 1937: 1218; Schedl 1940a: 359; Schwarz 1886: 44; Swaine 1909: 143, 1918a: 86–87; Titus, Meikle, & Harrison 1985: 89; Wood, S. L. 1982b: 1234; Zimmermann 1868: 143. **(ms)** Swaine 1907: 191.
- longulum* Eichhoff 1869a: 278 (*Pterocyclon*). Syn-types, sex?; Carolina [USA]; Hamburg Museum, lost. Synonymy: LeConte 1876: 349. References: **(ds)** Gemminger & Harold 1872: 2680; Leng 1920: 339; Swaine 1909: 143. **(tx)** Blandford 1904: 279; Eichhoff 1869a: 278, 1869c: 301, 1876a: 378–379, 1878b: 448; LeConte 1876: 349; Swaine 1909: 143, 1918a: 87.
- marcidum* (Schedl) 1970e: 98 (*Pterocyclon*). Holotype, sex?; Bolivien, Yungas del Palmar, 2000 m; Schedl Collection in NHMW, Wien. Distribution: South America (Bolivia). References: **(tx)** Schedl 1970e: 98–99, 1979c: 148.
- minutissimum* (Schedl) 1954b: 40 (*Pterocyclon*). Lectotype ♂; Brasilien: Matto Grosso, Rio Caragnata; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 155. Figures: Bright 1972d: 95. Distribution: South America (Brazil). Notes: (3) Schedl 1959m: 554 (described female). References: **(ds)** Schedl 1972g: 46. **(tx)** Bright 1972d: 95, 102; de Ruette 1970: 113; Schedl 1954b: 40, 1959m: 553–554, 1961i: 230, 1979c: 155. *schedli* Bright 1972d: 102. Lectotype ♂; Brasilien: Matto Grosso, Rio Caragnata; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 155, an unneeded replacement name for *minutissimum* (Schedl 1954). Notes: Bright 1972d: 102 presumed that this species was congeneric with *Microcorthyus minutissimus* Schedl 1952 and introduced a replacement name. The 1952 species is in fact a *Microcorthyus* and is also a junior synonym; therefore, homonymy never existed and the original 1954 name is retained. References: **(tx)** Bright 1972d: 102; Schedl 1972g: 46, 1979c: 155.
- minutum* (Schedl) 1939j: 577 (*Pterocyclon*). Lectotype ♀; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien designated by Schedl 1979c: 157. Distribution: Antilles Islands (Cuba/ Jamaica), North America ("Mexico"), South America (Brazil). References: **(ds)** Blackwelder 1947: 783; Bright 1951c: 155; Numberg 1958a: 480; Schedl 1976a: 56. **(tx)** Numberg 1958a: 480; Schedl 1939j: 577, 1950i: 166, 1952a: 460, 1979c: 157.
- morsum* Wood 1974b: 139. Holotype ♂; Pandora, Limon, Costa Rica; Wood Collection. Distribution: North America (Costa Rica/ Veracruz in Mexico). Hosts: *Quercus* sp., and a tree limb. References: **(hb)** Wood, S. L. 1982b: 1236. **(ds)** Wood, S. L. 1982b: 1236. **(tx)** Wood, S. L. 1974b: 139, 1982b: 1236.
- nevermanni* (Schedl) 1935h: 348 (*Pterocyclon*). Holotype ♂; Kraterlagune, Vulcan Poas, Heredia, Costa Rica; Schedl Collection in NHMW, Wien. Distribution: North America (Costa Rica). Hosts: *Clusia* sp., *Quercus* sp., *Podocarpus oleifolius*. References: **(hb)** Wood, S. L. 1982b: 1223. **(ds)** Blackwelder 1947: 783; Wood, S. L. 1982b: 1223. **(tx)** Schedl 1935h: 348, 1950i: 168, 1979c: 165; Wood, S. L. 1982b: 1223.
- notatum* Wood 1974b: 143. Holotype ♀; Vulcan Poas, Heredia, Costa Rica, 2500 m; Wood Collection. Distribution: North America (Costa Rica/ Panama). Hosts: *Inga* sp., and a leguminous shrub. References: **(hb)** Wood, S. L. 1982b: 1241. **(ds)** Wood, S. L. 1982b: 1241. **(tx)** Wood, S. L. 1974b: 143, 1982b: 1241.
- nudum* (Schedl) 1939j: 574 (*Pterocyclon*). Holotype, sex?; Brasilien, Sta. Catharina; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: **(ds)** Blackwelder 1947: 783; Schedl 1966f: 92, 1976a: 56. **(tx)** Schedl 1939j: 574, 1950i: 164, 1979c: 173.
- obesum* (Schedl) 1970e: 99 (*Pterocyclon*). Holotype, sex?; Peru; Schedl Collection in NHMW, Wien.

Distribution: South America (Peru).

References: (tx) Schedl 1970e: 99, 1979c: 174.

obliquum (Schedl) 1970e: 99 (*Pterocyclon*). Holotype, sex?; Brasilien, Rio Caraguata, Matto Grosso; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Schedl 1970e: 99–100, 1972g: 46. (tx) Schedl 1979c: 174.

obtusum (Eggers) 1935a: 334 (*Anchonocerus*). Holotype ♂; Bolivien (Cochabamba); Eggers Collection, in NHMW, Wien.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1935a: 334; Schedl 1979c: 176.

omissum (Schedl) 1952d: 347 (*Pterocyclon*). Syn-types ♂; Portorico, Mor.; Schedl Collection in NHMW, Wien, automatic.

Distribution: Antilles Islands (Puerto Rico).

References: (ds) Bright 1955c: 176. (tx) Bright 1955c: 176; Schedl 1952d: 347.

opacifrons Schedl 1950i: 167 (*Pterocyclon*). Syn-types ♂; Portorico, Mor.; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1935.

Notes: (1) The replacement name was presented prior to 1960; consequently, even though the senior homonym is now in a different genus, restoration of the 1950 name is not permitted by the Code. Schedl 1979c: 179 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 783. (tx) Schedl 1935h: 350, 1950i: 167, 1952d: 347, 1979c: 179.

parvum (Eggers) 1933b: 22 (*Anchonocerus*). Holotype ♂; Franz. Guayana (Nouveau Chantier); MNHN, Paris.

Distribution: South America (Cayenne/ Guyana).

Hosts: *Peltogyne* sp.

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1933b: 22–23, 1935a: 329–333; Schedl 1935a: 329; Wood, S. L. 1977b: 208.

preclarum Wood 1968b: 6. Holotype ♂; Manaka, British Guiana; BMNH, London. Synonymy: Wood 1977b: 208.

References: (tx) Wood, S. L. 1968a: 5, 1968b: 6, 1977b: 208.

penicillatum (Eichhoff) 1878b: 457 (*Pterocyclon*). Holotype, sex?; America meridionalis (Columbia); Steinheil Munchiensis Collection or Hamburg Museum?

Distribution: South America (Colombia).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 342. (tx) Eggers 1936a: 393; Eichhoff 1878b: 457; Hagedorn 1910a: 149.

pennatum (Schedl) 1963c: 167 (*Pterocyclon*). Holotype ♀; Cordoba, Veracruz, Mexico; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica/ Guate-

mala/ Honduras/ Nayarit, Veracruz in Mexico), South America (Colombia/ Venezuela).

Hosts: *Conostegia oerstediana*, *Miconia pubescens*, "Caldo de Frijol" [a tree], etc.

References: (hb) Atkinson et al. 1986: 50; Wood, S. L. 1982b: 1228. (ds) Atkinson & Equihua 1985c: 361; Atkinson et al. 1986: 50; Wood, S. L. 1982b: 1228. (tx) Schedl 1963c: 167, 1979c: 157; Wood, S. L. 1982b: 1228.

peruanum (Schedl) 1950i: 168 (*Pterocyclon*). Holotype ♂; Peru; Schedl Collection in NHMW, Wien.

Distribution: South America (Peru).

References: (tx) Schedl 1950i: 168, 1979c: 191.

peruvianum Wood 1981: 122. Holotype ♀; Peru, Tambo, Enenas, Com. del Pichis; Schedl Collection in NHMW, Wien, automatic.

Distribution: South America (Peru).

References: (tx) Wood, S. L. 1981: 122.

peruanum Schedl 1978c: 306. Holotype ♀; Peru, Tambo, Enenas, Com. del Pichis; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1950.

References: (tx) Schedl 1978c: 306; Wood, S. L. 1981: 122.

plagiatum (Eichhoff) 1869a: 279 (*Corthylys*). Holotype, sex?; Columbia; Hamburg Museum, lost.

Distribution: South America (Colombia).

References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 94; Kleine 1913b: 157. (tx) Eggers 1936a: 393; Eichhoff 1869a: 279, 1869c: 301, 1878b: 455; Hagedorn 1910a: 149.

plaumanni (Schedl) 1937g: 68 (*Pterocyclon*). Syn-types ♂; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, Strohmeyer Collection, Eberswalde, and Plaumann Collection.

Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 196 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 783; Numborg 1963c: 98; Santoro 1957b: 26; Schedl 1966f: 92. (tx) Schedl 1937g: 68, 1939k: 726, 1950i: 167, 1954b: 38, 1979c: 196.

posticum Wood 1974b: 146. Holotype ♂; Santa Ana, San Jose, Costa Rica, 1300 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Tree limbs, and a liana.

References: (hb) Wood, S. L. 1982b: 1245. (ds) Wood, S. L. 1982b: 1245. (tx) Wood, S. L. 1974b: 146, 1982b: 1245.

praeruptum (Blandford) 1904: 273 (*Pterocyclon*). Holotype ♂; Totonicipan, Guatemala, 85–10,500 ft.; BMNH, London.

Figures: Blandford 1904: pl. 8.

Distribution: North America (Guatemala).

References: (hb) Wood, S. L. 1982b: 1220. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94;

Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1220. (tx) Blandford 1904: 273; Hagedorn 1910a: 149; Schedl 1935h: 348; Wood, S. L. 1982b: 1220.

praeustum (Eggers) 1941a: 100 (*Pterocyclon*). Lectotype ♂; Guadeloupe (Trois Rivières); USNM, Washington, designated by Anderson & Anderson 1971: 26.

Distribution: Antilles Islands (Dominica/ Guadeloupe/ Puerto Rico).

Hosts: *Dacryodes excelsa*, *Inga laurina*.

References: (ds) Bright 1981c: 155, 1985c: 176. (tx) Anderson, W. H. & Anderson 1971: 26; Bright 1985c: 176; Eggers 1941a: 100; Schedl 1950i: 167, 1979c: 198.

proprium Wood 1974b: 136. Holotype ♂; Cerro Punta near Volcan Chiriqui, Panama; Wood Collection.

Distribution: North America (Costa Rica/ Panama). Hosts: *Conostegia oerstediana*, *Inga* sp., and tree limbs.

References: (hb) Wood, S. L. 1982b: 1227. (ds) Wood, S. L. 1982b: 1227. (tx) Wood, S. L. 1974b: 136, 1982b: 1227.

proximum Wood 1974b: 147. Holotype ♂; San Isidro del General, San Jose, Costa Rica, 1000 m; Wood Collection.

Distribution: North America (Costa Rica), South America (Venezuela).

Hosts: *Alexia imperitricia*, *Miconia pubescens*.

References: (ds) Wood, S. L. 1982b: 1245. (tx) Wood, S. L. 1974b: 147, 1982b: 1245.

pseudoscutellare (Schedl) 1935h: 349 (*Pterocyclon*). Holotype ♀; Lagima Volcan Poas, Heredia, Costa Rica; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

Hosts: *Quercus costaricensis*, *Q.* spp.

References: (hb) Wood, S. L. 1982b: 1239. (ds) Blackwelder 1947: 783; Wood, S. L. 1982b: 1239. (tx) Schedl 1935h: 349–350, 1979c: 202; Wood, S. L. 1982b: 1239.

adustum Wood 1974b: 140. Holotype ♂; Cerro de la Muerte, San Jose, Costa Rica; Wood Collection. Synonymy: Wood 1982b: 1239.

References: (tx) Wood, S. L. 1974b: 140, 1982b: 1239.

pumilio (Eichhoff) 1878b: 445 (*Pterocyclon*). Holotype, sex?; America meridionalis (Venezuela); Hamburg Museum, lost.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 340. (tx) Eichhoff 1878b: 445; Hagedorn 1910a: 149; Schedl 1939k: 727.

punctifrons (Blandford) 1904: 278 (*Pterocyclon*). Holotype ♀; Volcan de Chiriqui, Panama; BMNH, London.

Distribution: North America (Costa Rica/ Panama). Hosts: Logs.

References: (hb) Wood, S. L. 1982b: 1239. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1239. (tx) Blandford 1904: 278; Hagedorn 1910a: 149; Wood, S. L. 1982b: 1239.

quadridens (Eichhoff) 1869a: 277 (*Pterocyclon*). Holotype, sex?; Brasilia; Hamburg Museum, lost. Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 337. (tx) Eichhoff 1869a: 277, 1869c: 301, 1878b: 446; Hagedorn 1910a: 149.

dubium Eichhoff 1869a: 277 (*Pterocyclon*). Holotype, sex?; Brasilia; Hamburg Museum, lost. Synonymy: Eichhoff 1878b: 446.

References: (ds) Gemminger & Harold 1872: 2680. (tx) Eichhoff 1869a: 277, 1869c: 301, 1878b: 446.

quadridentatum (Eggers) 1935a: 333 (*Anchonomerus*). Lectotype ♂; Venezuela; USNM, Washington, designated by Anderson & Anderson 1971: 28.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 28; Eggers 1935a: 329, 333–334; Schedl 1979c: 208.

quercicolens Wood 1967d: 49. Holotype ♂; 53 km E Morelia, Michoacan, Mexico; Wood Collection. Figures: Wood 1967a: 46, 1982b: 1219 (declivity). Distribution: North America (Guatemala/ Chihuahua, Durango, Michoacan in Mexico/ Panama). Hosts: *Quercus* spp.

References: (hb) Atkinson et al. 1986: 50; Burgos & Saucedo 1983: 133. (ds) Atkinson et al. 1986: 50; Bright 1972a: 1385; Burgos & Saucedo 1983: 133; Wood, S. L. 1982b: 1219. (tx) de Ruette 1970: 104; Wood, S. L. 1967a: 46, 1967d: 49–50, 1982b: 1219.

quercivorum Schedl 1977e: 46. Holotype ♂; Metapan, El Salvador, 1600–2200 m; Schedl Collection in NHMW, Wien.

Distribution: North America (El Salvador).

Hosts: *Quercus* sp.

References: (tx) Schedl 1977e: 46.

quercum (Wood) 1967d: 53 (*Amphicranus*). Holotype ♂; 5 km W El Salto, Durango, Mexico, 2500 m; Wood Collection.

Figures: Wood 1967d: 54, 1982b: 1233 (declivity). Distribution: North America (Durango in Mexico). Hosts: *Quercus* sp.

References: (tx) Wood, S. L. 1967d: 53–54.

querneum Wood 1967d: 50. Holotype ♂; 53 km E Morelia, Michoacan, Mexico; Wood Collection. Figures: Bright 1972a: 1377, Wood 1967d: 46, 1982b: 1219 (declivity).

Distribution: North America (El Salvador/ Honduras/ Chiapas, Hidalgo, Michoacan in Mexico).

Hosts: *Quercus hondurensis*, *Q.* spp.

References: (hb) Atkinson & Equihua 1985c: 361; Atkinson et al. 1986: 50; Burgos & Saucedo 1983: 133. (ds) Atkinson & Equihua 1985a: 101, 1985c: 361; Atkinson et al. 1986: 50; Burgos & Saucedo 1983: 133; Schedl 1977e: 43; Wood, S. L. 1982b: 1218. (tx) McNamara 1977: 196; Wood, S. L. 1967a: 46, 1967d: 50, 1973c: 179, 1982b: 1218.

bifidus Bright 1972a: 1381. Holotype ♂; 13 km E San Cristobal de las Casas, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 179. References: (tx) Bright 1972a: 1377, 1381; McNamara 1977: 196; Wood, S. L. 1973c: 179.

***robustum* (Schedl)** 1966f: 123 (*Pterocyclon*). Holotype ♀; Las Mercedes, Santa Clara [district in Limon], Costa Rica, 100 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica), South America (Trinidad).

Hosts: *Spondias purpurea*, etc.

References: (hb) Wood, S. L. 1982b: 1224. (ds) Atkinson & Equihua 1986a: 422, 1988: 92; Schedl 1966f: 77; Wood, S. L. 1982b: 1224. (tx) Schedl 1966f: 123, 1979c: 212; Wood, S. L. 1982b: 1224.

***scrobiceps* (Eichhoff)** 1878b: 458 (*Pterocyclon*). Holotype, sex?; America meridionalis (Columbia orientalis); Hamburg Museum, lost.

Distribution: South America (Colombia).

Hosts: *Quercus* sp. (?).

References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 342. (tx) Eggers 1936a: 393; Eichhoff 1878b: 458; Hagedorn 1910a: 149; Schedl 1966f: 76.

***scutellare* (LeConte)** 1857: 59 (*Corthyus*). Holotype ♀; San Jose, California [USA]; MCZ, Cambridge.

Figures: Bright 1976d: 189.

Distribution: North America (British Columbia in Canada/ Baja California Norte in Mexico/ California, Oregon, Washington in USA).

Hosts: *Chrysolepsis* sp., *Lithocarpus densiflorus*, *Quercus agrifolia*, *Q. kelloggii*.

Notes: (1) The original (masculine) spelling was *scutellaris*.

References: (ay) Farris 1965b: 30, 1969: 529; Francke-Grosmann 1956b. (bv) Daterman, Rudinsky, & Nagel 1965. (cn) Chamberlin 1939: 280; Doane et al. 1936; Keen 1938: 147; Ruppel 1967: 64; Swaine 1918a: 86. (ce) Chamberlin 1939: 280; Farris 1965b: 30, 1969: 529; Francke-Grosmann 1956b; Keen 1938: 147; Roeper & French 1981; Slaby 1947: 377; Steinhans 1946: 404; Webb 1945: 65. (hb) Bright 1976d: 190; Bright & Stark 1973: 95; Chamberlin 1939: 280, 1955: 128–129; Chittenden 1893a: 392; Daterman, Rudinsky, & Nagel 1965; DeLeon 1952; Doane & Gilliland 1929: 916–918; Doane et al.

1936; Funk 1965: 929–932; Keen 1938: 147; Roling & Kearby 1974: 1303; Swaine 1918a: 86; Wood, S. L. 1982b: 1216. (ds) Bright 1976d: 190; Bright & Stark 1973: 95; Chamberlin 1939: 280, 1955: 128–129; DeLeon 1952: 79; Evans, D. 1985; Furniss, R. L. & Carolin 1977: 398; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 94; Henshaw 1882: 268, 1885: 147; Keen 1929a: 28, 1938: 147; Kleine 1913b: 157, 1914b: 391, 1934a: 171; Leng 1920: 339; Patterson & Hatch 1945: 152; Ritcher 1955: 16; Rivers 1886: 66; Ruppel 1967: 64; Swaine 1909: 144; Wood, S. L. 1959a: 61, 1972a: 428, 1982b: 1216. (tx) Bright 1976d: 189–190; Bruck 1936a; Chamberlin 1939: 280, 1955: 128–129; Eichhoff 1869a: 277, 1869c: 295, 1878b: 447; Ferrari 1867a: 48–49; Hagedorn 1910a: 149; Keen 1929a: 28; LeConte 1857: 22, 59, 1868: 153, 1876: 348; Schedl 1935h: 347; Swaine 1909: 144, 1918a: 86; Wood, S. L. 1959a: 61, 1966b: 26, 1972a: 428, 1982b: 1216.

cavus LeConte 1868: 153 (*Cryphalus*). Holotype ♂; coast region south of San Francisco, California [USA]; MCZ, Cambridge. Synonymy: LeConte 1876: 348.

References: (ds) Henshaw 1882: 268; Leng 1920: 339; Swaine 1909: 144. (tx) Eichhoff 1878b: 451; LeConte 1868: 153, 1876: 348; Swaine 1909: 144.

***obliquecaudatum* Schedl** 1935h: 351 (*Pterocyclon*). Syntypes ♀; California [USA]; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 26.

References: (ds) Blackwelder 1939. (tx) Schedl 1935h: 351, 1979c: 174; Wood, S. L. 1966b: 26.

***semipallens* (Schedl)** 1954b: 41 (*Pterocyclon*). Lectotype ♂; Brasilien: Matto Grosso, Rio Caragnata; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 224.

Distribution: South America (Brazil).

References: (tx) Schedl 1954b: 41, 1979c: 224.

***sexdentatum* Eggers** 1935a: 83. Lectotype ♀; Bolivien (Cochabamba); USNM, Washington, designated by Anderson & Anderson 1971: 29.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1935a: 83, 329, 331–332; Schedl 1979c: 227.

***sexdenticulum* Wood** 1989: 178. Holotype ♂; Columbien (Aguatal); USNM, Washington, automatic. Distribution: South America (Colombia).

References: (tx) Wood, S. L. 1989: 178.

***sexdentatum* Eggers** 1935a: 331 (*Anchonocerus*). Holotype ♂; Columbien (Aguatal); USNM, Washington, preoccupied by Eggers 1935.

References: (tx) Anderson, W. H. & Anderson 1971: 29; Eggers 1935a: 331–332; Wood, S. L. 1989: 178.

- subductum** (Schedl) 1978c: 303 (*Pterocyclon*).
Holotype, sex?; Peru, Torentoy Canyon (Base Machu-Picchu), 2000–2200 m; Schedl Collection in NHMW, Wien.
Distribution: South America (Peru).
References: (tx) Schedl 1978c: 303.
- subgranulatum** Wood 1974b: 144. Holotype ♂; Cerro Pena Blanca, Honduras, 1900 m; Wood Collection.
Distribution: North America (Honduras).
Hosts: *Persea popenoi*.
References: (hb) Wood, S. L. 1982b: 1243. (ds) Wood, S. L. 1982b: 1243. (tx) Wood, S. L. 1974b: 144, 1982b: 1243.
- subprorum** (Schedl) 1976a: 83 (*Pterocyclon*).
Holotype, sex?; Brazil: Salesopolis, S.P.; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 83.
- subtruncatum** (Schedl) 1978c: 303 (*Pterocyclon*).
Holotype, sex?; Columbién, Cauca, O. Thieme S.; Schedl Collection in NHMW, Wien.
Distribution: South America (Colombia).
References: (tx) Schedl 1978c: 303.
- sulcatum** (Blandford) 1905: 254 (*Pterocyclon*).
Holotype ♂; Volcan de Chiriqui, Panama, 1800 m; BMNH, London.
Distribution: North America (Panama).
Hosts: *Quercus* sp.
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157. (tx) Blandford 1905: 254; Hagedorn 1910a: 149; Schedl 1935h: 348.
- sulcipes** (Schedl) 1978c: 304 (*Pterocyclon*).
Holotype, sex?; Peru, Torentoy Canyon (Base Machu-Picchu), 2000–2200 m; Schedl Collection in NHMW, Wien.
Distribution: South America (Peru).
Notes: (3) This may be a synonym of *lobatum* Ferrari.
References: (tx) Schedl 1978c: 304.
- terminatum** (Blandford) 1904: 280 (*Pterocyclon*).
Holotype ♂; Capetillo, Guatemala; BMNH, London.
Distribution: North America (Guatemala).
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1222. (tx) Blandford 1904: 280; Hagedorn 1910a: 149; Wood, S. L. 1982b: 1222.
- tetradontium** Wood 1974b: 137. Holotype ♂; SE slope of Mt. Colima, Jalisco, Mexico; Wood Collection.
Distribution: North America (Jalisco in Mexico).
References: (hb) Atkinson et al. 1986: 51; Burgos & Saucedo 1983: 134; Wood, S. L. 1982b: 1235. (ds) Atkinson et al. 1986: 51; Burgos & Saucedo 1983: 134; Wood, S. L. 1982b: 1235. (tx) Wood, S. L. 1974b: 137, 1982b: 1235.
- tomicoides** (Blandford) 1904: 273 (*Pterocyclon*).
Lectotype ♂; San Geronimo, Verapaz, Guatemala; BMNH, London, designated by Wood 1982b: 1221.
Figures: Blandford 1904: pl. 8.
Distribution: North America (Guatemala).
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1221. (tx) Blandford 1904: 273; Hagedorn 1910a: 149; Wood, S. L. 1982b: 1221.
- tridentatum** (Schedl) 1971f: 152 (*Cosmocorynus*).
Holotype ♀; Brazil: Lagoa Santa; UZMC, Copenhagen.
Distribution: South America (Brazil).
References: (tx) Schedl 1971f: 152.
- umbrinum** (Blandford) 1904: 275 (*Pterocyclon*).
Holotype ♀; Purula, Verapaz, Guatemala; BMNH, London.
Distribution: North America (Guatemala/ Puebla in Mexico).
Hosts: *Alnus* sp.
References: (hb) Wood, S. L. 1982b: 1225. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1225. (tx) Blandford 1904: 275; Hagedorn 1910a: 149; Wood, S. L. 1982b: 1225.
- unifasciatum** (Schedl) 1970e: 104 (*Cosmocorynus*). Holotype ♀; Columbia, La Estrella H, 1700 m; Frey Collection in NHMBS, Basel.
Distribution: South America (Colombia).
References: (tx) Schedl 1970e: 104.
- vallidum** (Ferrari) 1867a: 55 (*Corthylus*). Lectotype ♂; Mexico; NHMW, Wien, designated by Wood 1974d: 285.
Distribution: North America (Costa Rica/ Veracruz in Mexico/ Panama).
Hosts: *Quercus* spp.
References: (hb) Wood, S. L. 1982b: 1220. (ds) Atkinson & Equihua 1985c: 361; Blackwelder 1947: 783; Blandford 1904: 271; Ferrer 1942; Genminger & Harold 1872: 2680; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 340, 353; Wood, S. L. 1982b: 1220. (tx) Blandford 1904: 271; Eichhoff 1869c: 301; Ferrari 1867a: 54–55, 61; Hagedorn 1910a: 149; Schedl 1940a: 359; Wood, S. L. 1974d: 285, 1982b: 1220.
- mexicanus** Eggers 1931a: 18 (*Amphicraus*).
Holotype ♂; Mexico; MNB, Berlin. Synonymy: Wood 1974d: 285.
References: (ds) Blackwelder 1947: 783. (tx) Eggers 1931a: 18; Schedl 1940a: 359; Wood, S. L. 1974d: 285.
- jalapae** Schedl 1939j: 584 (*Pterocyclon*). Holotype ♂; Jalapa, Veracruz, Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1974d: 285.
References: (ds) Blackwelder 1947: 783; Ferrer 1942. (tx) Schedl 1939j: 584, 1940a: 359, 1979c: 129; Wood, S. L. 1974d: 285.

vernaculum (Schedl) 1952a: 460 (*Pterocyclon*).
Syntypes 2 ♀; Argentina, Dep. Concep., Sta. Maria; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina/ Brazil).
Notes: (1) Schedl 1979c: 266 (citation of holotype invalid).

References: (ds) Schedl 1967d: 4. (tx) de Ruette 1970: 113; Schedl 1950i: 166, 1952a: 460, 1979c: 266.

vicinum (Schedl) 1970e: 100 (*Pterocyclon*). Holotype, sex?; Brasilien, S. Catarina, Luderwaldt; Eggers Collection, in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1970e: 100–101, 1979c: 267.

vittatum (Blandford) 1905: 282 (*Pterocyclon*). Holotype ♀; Volcan de Chiriqui, Panama; BMNH, London.
Figures: Blandford 1904: pl. 8.
Distribution: North America (Costa Rica/ Panama).
Hosts: *Brimellia costaricensis*, *Clusia* sp., *Quercus* sp.
References: (hb) Wood, S. L. 1982b: 1242. (ds) Blackwelder 1947: 783; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 367; Wood, S. L. 1982b: 1242. (tx) Blandford 1905: 282; Hagedorn 1910a: 149; Wood, S. L. 1982b: 1242.

volvulum (Eichhoff) 1869a: 279 (*Pterocyclon*). Holotype, sex?; Columbia; IRSNB, Brussels.
Distribution: South America (Colombia).
References: (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 342. (tx) Eichhoff 1869a: 279, 1869c: 301, 1878b: 454; Hagedorn 1910a: 149.

xalapensis Wood 1987: 548. Holotype ♂; Estacion de Biologia, Chamela, Jalisco, Mexico; Wood Collection.
Distribution: North America (Jalisco in Mexico).
Hosts: *Croton pseudoniveus*.
References: (tx) Wood, S. L. 1987: 548.

Genus *Glochinocerus* Blandford

GLOCHINOCERUS BLANDFORD 1904: 266. Type-species: *Glochinocerus retusipennis* Blandford, subsequent designation by Hopkins 1914: 122.
Keys: Wood 1982b: 1246.

References: (hb) Wood, S. L. 1986a: 99. (ds) Hagedorn 1910d: 92; Wood, S. L. 1986a: 99. (tx) Blandford 1904: 266; Hagedorn 1910a: 142–143; Hopkins 1914: 122; Wood, S. L. 1982b: 1246–1247, 1986a: 99.

gemellus Blandford 1904: 267. Lectotype ♀; Quiche Mountains, El Quiche, Guatemala; BMNH, London, designated by Wood 1982b: 1247.
Figures: Atkinson et al. 1986: 48.
Distribution: North America (Guatemala/ Hidalgo, Mexico, Puebla in Mexico).
Hosts: *Prunus* sp., *Quercus* sp., etc.
References: (hb) Atkinson & Equihua 1985a: 101; Atkinson et al. 1986: 47; Burgos & Saucedo 1983:

136. (ds) Atkinson & Equihua 1985a: 101; Atkinson et al. 1986: 47; Blackwelder 1947: 783; Burgos & Saucedo 1983: 136; Hagedorn 1910d: 92; Kleine 1913b: 156, 1914b: 367; Wood, S. L. 1982b: 1247. (tx) Atkinson et al. 1986: 48; Blandford 1904: 267; Hagedorn 1910a: 143; Wood, S. L. 1982b: 1247.

retusipennis Blandford 1904: 266. Lectotype ♀; Cerro Zumil, Quezaltenango, Guatemala; BMNH, London, designated by Wood 1982b: 1246.
Figures: Blandford 1904: pl. 9.
Distribution: North America (Guatemala).
Hosts: Caldo de Frijol bole, and a tree branch.
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 92; Kleine 1913b: 156, 1914b: 367; Wood, S. L. 1982b: 1246. (tx) Blandford 1904: 266; Hagedorn 1910a: 143; Hopkins 1914: 122; Lucas 1920: 303; Wood, S. L. 1982b: 1246. (ms) Lucas 1920: 303.

Genus *Metacorthylus* Blandford

METACORTHYLUS BLANDFORD 1904: 251, 263. Type-species: *Metacorthylus nigripennis* Blandford, monobasic.

Paracorthylus Wood 1968b: 7. Type-species: *Paracorthylus velutinus* Wood, original designation. Synonymy: Wood 1976a: 347.

References: (tx) Wood, S. L. 1968b: 7, 1976a: 347.

Keys: Wood 1982b: 1248.

References: (hb) Fonseca 1925: 1–8 [these citations erroneous, =*Corthylus* (SLW)]; Wood, S. L. 1982b: 1247, 1986a: 99. (ds) Wood, S. L. 1982b: 1247, 1986a: 99. (tx) Blandford 1904: 251, 263; Wood, S. L. 1976a: 347, 1982b: 1247, 1986a: 99.

concisus (Wood) 1974a: 67 (*Paracorthylus*). Holotype ♀; Moravia, Cartago, Costa Rica, 500 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Log.

Notes: (1) Wood 1976: 347 (to *Metacorthylus*).

References: (hb) Wood, S. L. 1982b: 1249. (ds) Wood, S. L. 1982b: 1249. (tx) Wood, S. L. 1974a: 67, 1976a: 347, 1982b: 1249.

mutilus (Wood) 1974a: 66 (*Paracorthylus*). Holotype ♀; Fort Sherman, Canal Zone, Panama; Wood Collection.

Distribution: North America (Panama).

Notes: (1) Wood 1976a: 347 (to *Metacorthylus*).

References: (ds) Wood, S. L. 1982b: 1248. (tx) Wood, S. L. 1974a: 66, 1976a: 347, 1982b: 1248.

nigripennis Blandford 1904: 263. Holotype ♀; Bugaba, Chiriqui, Panama, 800–1500 ft.; BMNH, London.

Distribution: North America (Panama).

References: (ds) Blackwelder 1947: 782; Hagedorn 1910d: 91; Kleine 1913b: 156, 1914b: 366; Wood, S. L. 1982b: 1251. (tx) Blandford 1904: 263; Fonseca 1925: 1–8; Hagedorn 1910a: 146;

Hopkins 1914: 124; Lucas 1920: 408; Wood, S. L. 1982b: 1251. (ms) Lucas 1920: 408.

velutinus (Wood) 1968b: 7 (*Paracorthylus*). Holotype ♀; Moravia, Cartago Prov., Costa Rica, 500 m; Wood Collection.

Figures: Wood 1968a: 5, 1982b: 1250.

Distribution: North America (Costa Rica), South America (Colombia).

Hosts: Log, and guano seco.

References: (hb) Wood, S. L. 1982b: 1249. (ds) Wood, S. L. 1982b: 1249. (tx) Wood, S. L. 1968a: 5, 1968b: 7, 1976a: 347, 1982b: 1249.

Genus *Microcorthylus* Ferrari

MICROCORTHYLUS FERRARI 1867a: 58. Type-species: *Microcorthylus parvulus* Ferrari, monobasic.

Keys: Wood 1982b: 1252 for Mexico and Central America.

References: (hb) Wood, S. L. 1986a: 99. (ds) Wood, S. L. 1986a: 99. (tx) Eggers 1935a: 75, 153; Ferrari 1867a: 58; Kirsch 1868: 214; Schedl 1939j: 570, 1940a: 359; Wood, S. L. 1967d: 37–57, 1982b: 1251–1259, 1986a: 99.

abruptus Nunberg 1962: 233. Holotype ♀; Estacao Biologica de Boraceia, Salesopolis, Estado de Sao Paulo, Brasilien; DZSA, Sao Paulo.

Figures: Nunberg 1962: 234.

Distribution: South America (Brazil).

Notes: (3) This is probably a *Monarthrum*.

References: (hb) Wood, S. L. 1982b: 1251. (ds) Wood, S. L. 1982b: 1251. (tx) Nunberg 1962: 233–234; Wood, S. L. 1982b: 1251.

bicolor Eggers 1935a: 154. Holotype, sex?; Bolivien, Cochabamba; Eggers Collection, in NHMW, Wien.

Distribution: South America (Bolivia).

References: (ds) Blackwelder 1947: 783. (tx) Eggers 1935a: 153–155; Schedl 1939j: 573, 1979c: 38.

brevis Eggers 1935a: 155. Holotype ♀; Guadeloupe; Eggers Collection, in NHMW, Wien.

Distribution: Antilles Islands (Guadeloupe).

References: (ds) Blackwelder 1947: 783; Bright 1985c: 176. (tx) Bright 1985c: 176; Eggers 1935a: 155–156, 1941a: 108; Schedl 1979c: 47.

castaneus Schedl 1939j: 572. Holotype, sex?; Brasilien, Sta. Catharina; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939j: 572, 1979c: 54.

concisus Wood 1973a: 270. Holotype ♀; Volcan, Puntarenas, Costa Rica, 1000 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: *Croton gossypifolius*, and other tree branches.

References: (hb) Wood, S. L. 1982b: 1259. (ds)

Wood, S. L. 1982b: 1259. (tx) Wood, S. L. 1973a: 270, 1982b: 1259.

contractus Wood 1973a: 274. Holotype ♂; 7 km NW Socopo, Barinas, Venezuela, 200 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Nectandra* sp.

References: (tx) Wood, S. L. 1973a: 274.

curtus Wood 1973a: 274. Holotype ♂; 20 km SW El Vigia, Merida, Venezuela, 50 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: Tree seedling.

References: (tx) Wood, S. L. 1973a: 274.

debilis Wood 1973a: 265. Holotype ♀; Volcan Pacaya, Guatemala, 1300 m; Wood Collection.

Distribution: North America (Guatemala).

Hosts: Tree branch, and a liana.

References: (hb) Wood, S. L. 1982b: 1254. (ds) Atkinson & Equihua 1985c: 362; Wood, S. L. 1982b: 1254. (tx) Wood, S. L. 1973a: 265, 1982b: 1254.

demissus Wood 1973a: 266. Holotype ♀; 9 km NE Teziutlan, Puebla, Mexico; Wood Collection.

Distribution: North America (Costa Rica/ Puebla in Mexico).

Hosts: *Boehmeria ulmifolia*, *Conostegia oerstediana*, *Miconia* sp.

References: (hb) Wood, S. L. 1982b: 1254. (ds) Wood, S. L. 1982b: 1254. (tx) Wood, S. L. 1973a: 266, 1982b: 1254.

dilutus Wood 1973a: 271. Holotype ♂; Piedras Blancas, 10 km E Medellin, Antioquia, Colombia, 2500 m; Wood Collection.

Distribution: South America (Colombia).

Hosts: Guttiferae sp.

References: (tx) Wood, S. L. 1973a: 271.

diversus Wood 1973a: 272. Holotype ♂; La Carbonera Experimental Forest, 50 km NW Merida, Merida, Venezuela, 2500 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Nectandra* sp.

References: (tx) Wood, S. L. 1973a: 272.

glabratus (Ferrari) 1867a: 60 (*Corthylus*). Holotype, sex?; Venezuela [presumably Colonia Tovar]; NHMW, Wien.

Distribution: South America (Brazil/ Venezuela).

Notes: Schedl 1963h: 268 (to *Microcorthylus*).

References: (ds) Bright 1985c: 176; Fleutiaux & Salle 1890: 458; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 93; Kleine 1914b: 340. (tx) Bright 1985c: 176; Ferrari 1867a: 53, 60; Hagedorn 1910a: 148; Schedl 1936i: 108, 1963h: 268.

subopacus Schedl 1939j: 573. Lectotype, sex?; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 243. Synonymy: Schedl 1963h: 268.

- References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939j: 573, 1963l: 268, 1979c: 243.
- grandiclavatus** Eggers 1935a: 156. Holotype ♂; Costa Rica, San Jose, 1300 m; Hamburg Museum, lost, neotype ♂, Escasn, San Jose, Costa Rica. USNM, Washington, designated by Wood 1982b: 1255.
Distribution: North America (Costa Rica/ Panama).
Hosts: Small tree.
References: (hb) Wood, S. L. 1982b: 1255. (ds) Blackwelder 1947: 783; Wood, S. L. 1982b: 1255. (tx) Eggers 1935a: 156; Wood, S. L. 1982b: 1255.
- hostilis** Wood 1973a: 272. Holotype ♀; 30 km N Merida, Merida, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: Tree branch.
References: (tx) Wood, S. L. 1973a: 272.
- inermis** Wood 1973a: 267. Holotype ♀; 15 km SE Cartago, Cartago, Costa Rica, 1500 m; Wood Collection.
Distribution: North America (Costa Rica/ Panama).
Hosts: *Guazuma ulmifolia*, *Siparuna nicaraguensis*.
References: (hb) Wood, S. L. 1982b: 1255. (ds) Wood, S. L. 1982b: 1255. (tx) Wood, S. L. 1973a: 267, 1982b: 1255.
- invalidus** Wood 1973a: 265. Holotype ♀; 6 km W Tepic, Nayarit, Mexico, 1000 m; Wood Collection.
Distribution: North American (Nayarit in Mexico).
Hosts: Tree branch.
References: (hb) Atkinson & Equihua 1955c: 362; Wood, S. L. 1982b: 1255. (ds) Atkinson & Equihua 1955c: 362; Wood, S. L. 1982b: 1255. (tx) Wood, S. L. 1973a: 265, 1982b: 1255.
- lassus** Wood 1973a: 270. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Distribution: North America (Costa Rica/ Panama).
Hosts: Small trees, and a liana.
References: (hb) Wood, S. L. 1982b: 1255. (ds) Wood, S. L. 1982b: 1255. (tx) Wood, S. L. 1973a: 270, 1982b: 1255.
- mexicanus** Eggers 1934a: 81. Holotype, sex?; Mexico; Hamburg Museum, lost.
Distribution: North America ("Mexico").
References: (ds) Blackwelder 1947: 783. (tx) Eggers 1934a: 81–82; Schedl 1940a: 359.
- minimus** Schedl 1950i: 160. Lectotype ♀; Nova Teutonia, Santa Catarina, Brazil; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 154.
Figures: Atkinson & Aquihua 1988: 97, Pedrosa-Macedo & Schonherr 1985: 59
Distribution: Antilles Islands (Cuba/ Jamaica), North America (Costa Rica/ Honduras/ Veracruz in Mexico/ Panama), South America (Brazil).
Hosts: *Acacia pennatula*, *Serjania* sp., and other small branches.
References: (cc) Equihua & Atkinson 1986: 634. (hb) Equihua & Atkinson 1986: 634; Wood, S. L. 1982b: 1256. (ds) Atkinson & Equihua 1988: 92; Bright 1981c: 155, 1985c: 176; Equihua & Atkinson 1986: 634; Pedrosa-Macedo & Schonherr 1985: 59; Schedl 1966f: 91, 1967d: 4, 1976a: 56; Wood, S. L. 1982b: 1256. (tx) Atkinson & Equihua 1988: 97; Bright 1985c: 176; Pedrosa-Macedo & Schonherr 1985: 59; Schedl 1950i: 160, 1979c: 154; Wood, S. L. 1982b: 1256.
- minutissimus** Schedl 1952d: 361. Syntypes, sex?; Jamaica; Schedl Collection in NHMW, Wien.
Synonymy: Wood 1975b: 394.
Notes: (1) Schedl 1979c: 156 (citation of holotype invalid).
References: (ds) Bright 1972d: 102; Schedl 1966f: 91, 1967d: 4, 1971f: 147, 1972g: 46. (tx) Bright 1972d: 102; Schedl 1950i: 160, 1952d: 361, 1979c: 156; Wood, S. L. 1975b: 394.
- obscurus** Eggers 1935a: 155. Holotype, sex?; Bolivia, Cochabamba; USNM, Washington.
Distribution: South America (Bolivia).
References: (ds) Blackwelder 1947: 783. (tx) Anderson, W. H. & Anderson 1971: 22; Eggers 1934a: 81, 1935a: 155; Schedl 1939j: 570–571, 1979c: 175.
- ocularis** Wood 1973a: 266. Holotype ♀; Tapanti, Cartago, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Boehmeria ulmifolia*, *Conostegia oerstediana*, *Siparuna nicaraguensis*.
References: (hb) Wood, S. L. 1982b: 1254. (ds) Wood, S. L. 1982b: 1254. (tx) Wood, S. L. 1973a: 266–267, 1982b: 1254.
- pallidus** Schedl 1939j: 571. Lectotype, sex?; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 182.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 783; Nunberg 1964a: 235; Schedl 1966f: 78. (tx) Nunberg 1956: 235; Schedl 1939j: 570–571, 1979c: 182.
- parvulus** Ferrari 1867a: 53, 58. Holotype ♂; Venezuela [probably Colonia Tovar]; NHMW, Wien.
Distribution: North America (Costa Rica/ Guatemala), South America (Colombia/ Venezuela).
Hosts: *Spondias purpurea*, and other tree branches.
References: (hb) Wood, S. L. 1982b: 1255. (ds) Blackwelder 1947: 783; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 94; Kleine 1913b: 157, 1914b: 340, 366; Wood, S. L. 1982b: 1255. (tx) Eggers 1935a: 153–156; Ferrari 1867a: 53, 58; Hagedorn 1910a: 145; Hopkins 1914: 125; Schedl 1939j: 570; Wood, S. L. 1974d: 283, 1982b: 1255.
- exile** Eichhoff 1878b: 451 (*Pterocyclon*). Holotype ♂; America meridionalis (Nova Granada [probably Colonia Tovar, Colombia]); IRSNB, Brussels. Synonymy: Wood 1974d: 283.
References: (ds) Blackwelder 1947: 783; Hagedorn 1910d: 92; Kleine 1913b: 157. (tx)

Eichhoff 1878b: 444, 451; Hagedorn 1910a: 148; Swaine 1909: 142, 1918a: 88; Wood, S. L. 1974d: 283.

porrectus Schedl 1951b: 292. Lectotype, sex?; Argentinien, Dep. Concepcion, Sta. Maria; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 197.

Distribution: South America (Argentina).

References: (hb) Viana 1964: 126. (ds) Viana 1964: 126. (tx) Schedl 1951b: 292, 1979c: 197.

puerulus Schedl 1939j: 571. Lectotype, sex?; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 203. Figures: Schedl 1939j: 570 (antenna, tibia).

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939j: 570–571, 1950, 1951b: 292, 1979c: 203.

pumilus Wood 1973a: 268. Holotype ♀; 15 km SE Cartago, Costa Rica, 1800 m; Wood Collection.

Distribution: North America (Costa Rica/ Panama).

Hosts: *Conostegia oerstediana*, etc.

References: (hb) Wood, S. L. 1982b: 1257. (ds) Wood, S. L. 1982b: 1257. (tx) Wood, S. L. 1973a: 268, 1982b: 1257.

pusillus Wood 1973a: 269. Holotype ♀; Buenos Aires, Cortez, Honduras, 2300 m; Wood Collection.

Distribution: North America (Guatemala/ Honduras).

Hosts: Tree branch.

References: (hb) Wood, S. L. 1982b: 1257. (ds) Wood, S. L. 1982b: 1257. (tx) Wood, S. L. 1973a: 269, 1982b: 1257.

rufotestaceus Schedl 1939j: 572. Lectotype, sex?; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 215. Distribution: South America (Brasil).

References: (ds) Blackwelder 1947: 783. (tx) Schedl 1939j: 572, 1979c: 215.

suggrandis Schedl 1939j: 574. Lectotype, sex?; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 245. Figures: Pedrosa-Macedo & Schonherr 1955: 60. Distribution: South America (Brazil).

Hosts: *Ocotea puberula*.

References: (ds) Blackwelder 1947: 783; Pedrosa-Macedo & Schonherr 1955: 60; Schedl 1966f: 91; Schonherr & Pedrosa-Macedo 1981: 53. (tx) Numberg 1962: 235; Pedrosa-Macedo & Schonherr 1955: 60; Schedl 1939j: 574, 1979c: 245.

umbratus Wood 1973a: 273. Holotype ♂; La Carbonera Experimental Forest, 50 km NW Merida, Merida, Venezuela, 2500 m; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Nectandra* sp.

References: (tx) Wood, S. L. 1973a: 273.

rescus Wood 1973a: 271. Holotype ♀; Volcan Zumil, Quezaltenango, Guatemala, 1000 m; Wood Collection.

Distribution: North America (Guatemala).

References: (ds) Atkinson & Equihua 1985c: 362; Wood, S. L. 1982b: 1259. (tx) Wood, S. L. 1973a: 271, 1982b: 1259.

vicinus Wood 1977b: 213. Holotype ♀; 51 km S Valle Nacional, Oaxaca, Mexico; CNCI, Ottawa.

Distribution: North America (Chiapas, Oaxaca in Mexico).

Hosts: *Quercus* sp., *Salix* sp.

References: (ds) Wood, S. L. 1982b: 1256. (tx) Wood, S. L. 1977b: 213, 1982b: 1256.

Genus *Corthyrcyclon* Schedl

CORTHYCYCLON SCHEDL 1951m: 128. Type-species: *Corthyrcyclon ustum* Schedl, monobasic.

Keys: Wood 1982b: 1260 for Mexico and Central America.

References: (hb) Wood, S. L. 1986a: 100. (ds) Wood, S. L. 1986a: 100. (tx) Schedl 1951m: 128; Wood, S. L. 1982b: 1259–1264, 1986a: 100.

aztecum (Bright) 1972a: 1374 (*Corthylys*). Holotype ♀; 5.6 km S Suchixtepec, Oaxaca, Mexico; CNCI, Ottawa.

Figures: Bright 1972a: 1370, Wood 1982b: 1262 (antenna).

Distribution: North America (Costa Rica/ Chiapas, Oaxaca, Puebla in Mexico).

Hosts: *Bocconia frutescens*, *Miconia dodecandra*, *M.* sp., *Oreopanax xalapense*, *Quercus* sp., *Siparuna nicaraguensis*, *Vaccinium consanguineum*.

Notes: (1) Wood 1982b: 1263 (to *Corthyrcyclon*).

References: (hb) Wood, S. L. 1982b: 1261. (ds) Wood, S. L. 1982b: 1263. (tx) Bright 1972a: 1370, 1374; McNamara 1977: 195; Wood, S. L. 1982b: 1263–1264.

caliginis Wood 1974b: 148. Holotype ♀; 16 km SE Cartago, Costa Rica, 1800 m; Wood Collection.

Figures: Wood 1982b: 1262.

Distribution: North America (Costa Rica).

Hosts: Liana.

References: (hb) Wood, S. L. 1982b: 1261. (ds) Wood, S. L. 1982b: 1261. (tx) Wood, S. L. 1974b: 148, 1982b: 1261.

ebeninum (Blandford) 1904: 265 (*Brachyspartus*).

Holotype ♀; Volcan de Chiriqui, Chiriqui, Panama, 4–6000 ft.; BMNH, London.

Distribution: North America (Panama).

References: (ds) Blackwelder 1947: 784; Hagedorn 1910d: 91; Kleine 1913b: 156, 1914b: 366; Wood, S. L. 1982b: 1262. (tx) Blandford 1904: 265; Hagedorn 1910a: 140; Wood, S. L. 1982b: 1262.

furcum Wood 1974b: 148. Holotype ♀; Peralta, Cartago, Costa Rica, 500 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: Liana.

References: (hb) Wood, S. L. 1982b: 1261. (ds) Wood, S. L. 1982b: 1261. (tx) Wood, S. L. 1974b: 148, 1982b: 1261.

morulum Wood 1974b: 149. Holotype ♀; 16 km SE Cartago, Costa Rica, 1800 m; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Conostegia oerstediana*.

References: (**hb**) Wood, S. L. 1982b: 1261. (**ds**) Wood, S. L. 1982b: 1261. (**tx**) Wood, S. L. 1974b: 149, 1982b: 1261.

tardum Wood 1974b: 149. Holotype ♀; 16 km SE Cartago, Costa Rica, 1800 m; Wood Collection. Figures: Wood 1982b: 1262 (antenna, tibia).

Distribution: North America (Costa Rica/ Panama). Hosts: *Cecropia* sp. (leaf petioles), *Siparuna nicaraguensis*, *Vaccinium consanguineum*.

References: (**hb**) Wood, S. L. 1982b: 1263. (**ds**) Wood, S. L. 1982b: 1263. (**tx**) Wood, S. L. 1974b: 149, 1982b: 1263.

ustum Schedl 1951m: 128. Holotype ♀; Brasilien; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil).

Notes: (3) Schedl 1959m: 555 (described male). References: (**tx**) Schedl 1951m: 128, 1959m: 555, 1979c: 263.

Genus *Brachyspartus* Ferrari

BRACHYSPARTUS FERRARI 1867a: 65. Type-species: *Brachyspartus moritzi* Ferrari, monobasic.

Thylurcos Schedl 1939j: 567. Type-species: *Brachyspartus moritzi* Ferrari, by subsequent monotypy. Synonymy: Wood 1966b: 18, 1986: 100. References: (**tx**) Schedl 1939j: 567–568; Wood, S. L. 1966b: 18, 1986: 100.

Notes: (1) *Thylurcos* Schedl was based on 2 species; *nevermanni* Schedl was transferred to *Corthylus* (Wood 1982b: 1298) leaving *moritzi* Ferrari as the only original species in the genus; thus, *Thylurcos* became an objective junior synonym of *Brachyspartus*.

References: (**hb**) Wood, S. L. 1986a: 100. (**ds**) Hagedorn 1910d: 91; Wood, S. L. 1986a: 100. (**tx**) Blandford 1904: 264; Eichhoff 1878b: 429; Ferrari 1867a: 65; Hagedorn 1910a: 140; Schedl 1935b: 347, 1939j: 567; Wood, S. L. 1966b: 18, 1986a: 100.

moritzi Ferrari 1867a: 68. Holotype ♀; Venezuela; NHMW, Wien.

Distribution: South America (Venezuela). References: (**ds**) Blackwelder 1947: 784; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 91; Kleine 1913b: 156, 1914b: 340, 342. (**tx**) Eichhoff 1868d: 419, 1878b: 429; Ferrari 1867a: 68; Hagedorn 1910a: 140; Hopkins 1914: 117; Schedl 1939j: 567; Wood, S. L. 1966b: 18, 1992b: 82. (**ms**) Eichhoff 1868d: 419.

obtusus Schedl 1966f: 122 (*Corthylus*). Holotype ♀; Venezuela; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992b: 82.

References: (**tx**) Schedl 1966f: 122, 1979c: 176; Wood, S. L. 1992b: 82.

Genus *Corthylocurus* Wood

CORTHYLOCURUS WOOD 1966b: 18. Type-species: *Brachyspartus barbatus* Blandford, original designation.

Keys: Wood 1982b: 1265 for Mexico and Central America.

References: (**hb**) Wood, S. L. 1986a: 100. (**ds**) Wood, S. L. 1986a: 100. (**tx**) Wood, S. L. 1966b: 18–19, 1973c: 175, 1982b: 1264–1270, 1986a: 100.

aguacatensis (Schedl) 1940a: 357 (*Metacorthylus*). Lectotype ♂; Xochitlan, Morelos, Mexico; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 15.

Figures: Atkinson et al. 1986: 43, Wood 1982b: 1267 (female head, declivity).

Distribution: North America (Michoacan, Morelos, Nayarit in Mexico).

Hosts: *Persca americana*, *Spondias mombin*.

Notes: (1) Wood 1982b: 1266 (designation of lectotype not needed).

References: (**hb**) Atkinson et al. 1986: 44; Burgos & Saucedo 1983: 183; Wood, S. L. 1982b: 1266. (**ds**) Atkinson et al. 1986: 44, 68; Blackwelder 1947: 782; Burgos & Saucedo 1983: 138; Wood, S. L. 1982b: 1266. (**tx**) Atkinson et al. 1986: 43; Schedl 1940a: 357, 1950i: 161, 1979c: 15; Wood, S. L. 1982b: 1266.

barbatus (Blandford) 1904: 265 (*Brachyspartus*). Holotype ♀; Volcan de Chiriqui, Chiriqui, Panama; BMNH, London.

Figures: Blandford 1904: 265, pl. 5, Wood 1982b: 1267 (female head, declivity).

Distribution: North America (Costa Rica/ Oaxaca, Puebla in Mexico/ Panama).

Hosts: *Conostegia oerstediana*, *Miconis* sp., *Spondias purpurea*.

References: (**hb**) Wood, S. L. 1982b: 1269. (**ds**) Atkinson & Equihua 1985c: 362; Blackwelder 1947: 784; Bright 1972a: 1385; Hagedorn 1910d: 91; Kleine 1913b: 156, 1914b: 366; Wood, S. L. 1982b: 1269. (**tx**) Blandford 1904: 265; Hagedorn 1910a: 140; Schedl 1935b: 349, 1939j: 569, 1950i: 162; Wood, S. L. 1966b: 19, 1973c: 175, 1982b: 1269.

brunneus (Nunberg) 1972b: 191 (*Corthylus*). Holotype ♂; Sao Paulo, Itarare (Sampaio); MZUSP, Sao Paulo.

Distribution: South America (Brazil).

References: (**tx**) Nunberg 1972a, 1972b: 191–192.

cinnamatus Bright 1972a: 1379. Holotype ♀; Mexico: 21 km N Ocozautla, Chiapas; CNCI, Ottawa.

Figures: Bright 1972a: 1377, Wood 1982b: 1267 (female head).

Distribution: North America (Costa Rica/ Chiapas in Mexico).

Hosts: Liana.

References: (**hb**) Wood, S. L. 1982b: 1269. (**ds**)

Wood, S. L. 1982b: 1269. (tx) Bright 1972a: 1377, 1379; McNamara 1977: 195; Wood, S. L. 1973c: 175, 1982b: 1269.

costaricensis (Schedl) 1936i: 108 (*Microcorthyus*). Holotype ♀; Turrialba, Cartago, Costa Rica; Schedl Collection in NHMW, Wien.

Figures: Wood 1982b: 1267 (female head, declivity). Distribution: North America (Costa Rica).

Hosts: Lianas.

Notes: (1) Wood 1982b: 1268 (to *Corthylocurus*).

References: (hb) Wood, S. L. 1982b: 1268. (ds) Blackwelder 1947: 783; Wood, S. L. 1982b: 1268. (tx) Schedl 1936i: 108, 1979c: 67; Wood, S. L. 1982b: 1268.

debilis Wood 1974b: 150. Holotype ♀; Beverley, Limon, Costa Rica; Wood Collection.

Figures: Wood, S. L. 1982b: 1267 (female head, declivity).

Distribution: North America (Costa Rica/ Panama). Hosts: *Serjania* sp.

References: (cn) Equihua 1988. (hb) Wood, S. L. 1982b: 1266. (ds) Atkinson & Equihua 1985c: 362, 1988: 86; Estrada & Atkinson 1988: 211; Wood, S. L. 1982b: 1266. (tx) Wood, S. L. 1974b: 150, 1982b: 1266.

mexicanus (Schedl) 1950i: 163 (*Brachyspartus*). Holotype ♂; Comitán, Mexico; Schedl Collection in NHMW, Wien.

Figures: Wood, S. L. 1982b: 1267 (female head, declivity).

Distribution: North America (Costa Rica/ Guatemala/ Chiapas, Oaxaca, Puebla, Veracruz in Mexico/ Panama).

Hosts: *Acacia* sp., *Arbutus* sp., *Inga* sp., *Miconia* sp., *Rhus* sp., *Rubus* sp., and other small branches.

References: (hb) Wood, S. L. 1982b: 1268. (ds) Atkinson & Equihua 1985c: 362; Ferrer 1942; Wood, S. L. 1982b: 1268. (tx) Schedl 1950i: 163, 1979c: 152; Wood, S. L. 1973c: 175, 1974d: 279, 1982b: 1268.

cylindricus Schedl 1963c: 164 (*Corthyus*). Holotype ♀; Jalapa, Veracruz, Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1974d: 279.

References: (tx) Schedl 1963c: 164, 1979c: 73; Wood, S. L. 1974d: 279.

anomalous Bright 1972a: 1378 (*Corthyus*). Holotype ♀; Highway 175, 5.6 km S Suchistepec, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 175.

References: (tx) Bright 1972a: 1378; McNamara 1977: 195; Wood, S. L. 1973c: 175.

signatifrons (Schedl) 1950i: 162 (*Brachyspartus*). Holotype ♀; Tybays or Tybagos [presumably Brazil]; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil?).

References: (tx) Schedl 1950i: 162, 1979c: 228.

tuberculifer (Eggers) 1937a: 86 (*Brachyspartus*). Syntypes 2, sex?; Brasil, Bahia; BMNH, London, 2 Eggers syntypes in NHMW, Wien.

Figures: Terra 1987: 28.

Distribution: South America (Brazil).

Hosts: *Theobroma cacao*.

References: (ds) Blackwelder 1947: 784; Numborg 1976a: 56; Schedl 1976a: 56; Terra 1987: 27. (tx) Anderson, W. H. & Anderson 1971; Eggers 1937a: 86–87; Schedl 1939j: 569, 1979c: 258; Terra 1987: 27–28.

vernaculus (Schedl) 1939j: 569 (*Brachyspartus*). Syntypes ♀; Nova Tentonia, Brazil; Schedl Collection in NHMW, Wien, and Plaumann Collection. Distribution: South America (Brazil).

Notes: (1) Schedl 1979c: 266 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 784; Schedl 1966f: 91, 1967d: 4, 1970e: 85, 1976a: 56. (tx) Schedl 1939j: 569, 1950i: 163, 1979c: 266.

Genus *Corthyus* Erichson

CORTHYLUS ERICHSON 1836: 64. Type-species: *Bostrichus compressicornis* Fabricius, by subsequent monotypy by Ferrari 1867a: 49. Wood 1974c: 202 designated lectotype for type-species.

Morizus Ferrari 1867a: 49. Type-species: *Morizus excisus* Ferrari, monobasic. Synonymy: Eichhoff 1869c: 300.

References: (tx) Blandford 1904: 251; Eichhoff 1869c: 297–301; Ferrari 1867a: 49, 59, 69; Swaine 1909: 90.

Pseudocorthyus Ferrari 1867a: 59. Type-species: *Pseudocorthyus letzneri* Ferrari. Synonymy: Wood 1977d: 511.

References: (tx) Ferrari 1867a: 59; Kirsch 1868: 214; Wood, S. L. 1977d: 511.

Corthylominus Schedl 1972g: 73. Type-species: *Corthylominus gracilis* Schedl, original designation, preoccupied by Ferrari 1867.

References: (tx) Schedl 1972g: 73, 1979c: 107.

Keys: Blandford 1904: 253 for Mexico and Central America; Wood 1982b: 1271 for North and Central America.

References: (ay) Nobuchi 1969a: 68. (hb) Hubbard 1897: 16; Merino & Vasquez 1963: 59–67; Wood, S. L. 1982b: 1270, 1986a: 100. (ds) Hagedorn 1910d: 90–91; Wood, S. L. 1982b: 1270, 1986a: 100. (tx) Arnett 1960: 1045, 1968: 1045; Beal & Massey 1945: 59, 104; Blandford 1904: 251–254; Bright 1972d: 103, 1976d: 191, 1978: 192; Chamberlin 1939: 283–285; Dodge 1938: 18; Eggers 1935a: 76–77, 1940a: 123–141; Eichhoff 1869a: 279, 1869c: 297–301, 1878b: 66, 421; Erichson 1836: 64; Ferrari 1867a: 48, 1867b: 104–115; Fonseca 1927: 585–590; Hagedorn 1910a: 144–145; LeConte 1876: 347; LeConte & Horn 1883: 517; Schedl 1939j: 567, 1940a: 349–350; Wood, S. L. 1966b: 19, 1967b: 37–57, 1967c:

- 119–141, 1974c: 181–202, 1977d: 511, 1982b: 1270–1310, 1986a: 100.
- abbreviatus** Eichhoff 1869a: 279. Holotype ♀; Columbia; IRSNB, Brussels.
Figures: Menschöy & Martin 1974: 6.
Distribution: South America (Colombia/Venezuela).
Hosts: *Ahuus* sp., *Croton* sp., *Ficus* sp., *Vismia* sp.
References: (hb) Menschöy & Martins 1974. (ds) Blackwelder 1947: 784; Blandford 1904: 254; Gemminger & Harold 1872: 2693; Hagedorn 1903: 547, 1910d: 90; Kleine 1913b: 155, 1914b: 342. (tx) Blandford 1904: 254; Eggers 1933b: 2, 21; Eichhoff 1869a: 279, 1869c: 301, 1875b: 425; Hagedorn 1903b: 547, 1910a: 145; Menschöy & Martin 1974: 6.
- abruptedecivis** Schedl 1966f: 121. Holotype ♀; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1966f: 121, 1979c: 8.
- additus** Wood 1974c: 199. Holotype ♀; La Carbonera Experimental Forest, 50 km NW Merida, Merida, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Clusia* sp., *Ficus* sp., *Nectandra* sp., *Rubus* sp., *Vismia* sp.
References: (tx) Wood, S. L. 1974c: 199.
- alienus** Schedl 1966f: 119. Holotype ♀; Argentinien, Tucuman; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
References: (tx) Schedl 1966f: 119, 1979c: 16.
- antennarius** Schedl 1966f: 120. Holotype ♀; Brasilien, Nova Teutonia, Santa Catarina; Schedl Collection in NHMW, Wien.
Figures: Pedrosa-Macedo & Schonherr 1985: 61.
Distribution: South America (Brazil).
References: (ds) Pedrosa-Macedo & Schonherr 1985: 61; Schedl 1976a: 55. (tx) Pedrosa-Macedo & Schonherr 1985: 61; Schedl 1966f: 120, 1979c: 22.
- argentinensis** Schedl 1950i: 157. Holotype ♂; Argentina, Pr. Jujuy, Verzenyi; Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
Notes: (3) Schedl 1976a: 79 (described female).
References: (ds) Schedl 1976a: 53. (tx) Nunberg 1962: 230; Schedl 1950i: 156–157, 1976a: 79, 1979c: 24.
- bifurcus** Schedl 1935h: 345. Holotype ♂; Costa Rica, Lagune Vulkan Poas, 2600 m; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
Hosts: Probably a native bamboo.
References: (ds) Blackwelder 1947: 784; Wood, S. L. 1982b: 1282. (tx) Schedl 1935h: 345, 1979c: 40; Wood, S. L. 1982b: 1282.
- bolivianus** Eggers 1943a: 379. Holotype ♀; Bolivia (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia).
References: (tx) Anderson, W. H. & Anderson 1971: 7; Eggers 1943a: 379–380.
- brunnescens** Wood 1981: 122. Holotype ♀; Volcan Barba, Heredia, Costa Rica; Wood Collection, automatic.
Figures: Wood 1982b: 1291 (female).
Distribution: North America (Costa Rica).
Hosts: Native bamboo.
References: (hb) Wood, S. L. 1982b: 1290. (ds) Wood, S. L. 1982b: 1290. (tx) Wood, S. L. 1981: 122, 1982b: 1290.
- brunneus** Wood 1974c: 185. Holotype ♀; Volcan Barba, Heredia, Costa Rica; Wood Collection, preoccupied by Nunberg 1972.
References: (tx) Wood, S. L. 1974c: 185, 1981: 122.
- calamarius** Wood 1974c: 185. Holotype ♀; Volcan Poas, Heredia, Costa Rica, 2000 m; Wood Collection.
Figures: Wood 1982b: 1280 (female).
Distribution: North America (Costa Rica).
Hosts: Native bamboo.
References: (hb) Wood, S. L. 1982b: 1282. (ds) Wood, S. L. 1982b: 1282. (tx) Wood, S. L. 1974c: 185, 1982b: 1282.
- callidus** Schedl 1973d: 172. Holotype ♀; Brasilien, Guanabara, Rio de Janeiro, Ilha do Governador; MZUSP, Sao Paulo.
Distribution: South America (Brazil).
References: (tx) Schedl 1973d: 172, 1979c: 50.
- calmicoleus** Wood 1974c: 189. Holotype ♀; Volcan Poas, Heredia, Costa Rica, 2500 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: Native bamboo.
References: (tx) Wood, S. L. 1974c: 189, 1982b: 1291.
- canularius** Wood 1974c: 186. Holotype ♀; Cerro de la Muerte, San Jose, Costa Rica, 3200 m; Wood Collection.
Figures: Wood 1982b: 1284 (female).
Distribution: North America (Costa Rica).
Hosts: Native bamboo.
References: (hb) Wood, S. L. 1982b: 1283. (ds) Wood, S. L. 1982b: 1283. (tx) Wood, S. L. 1974c: 186, 1982b: 1283.
- cavifrons** Nunberg 1962: 230. Holotype ♀; Estacao Biologica de Boraceia, Salesopolis, Estado de Sao Paulo, Brasilien; DZSA, Sao Paulo.
Distribution: South America (Brazil).
References: (tx) Nunberg 1962: 230.
- cecropii** Wood 1975a: 31. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Distribution: North America (Costa Rica), South America (Venezuela).
Hosts: *Cecropia peltata* leaf petioles.
References: (hb) Wood, S. L. 1982b: 1278. (ds) Wood, S. L. 1982b: 1278. (tx) Wood, S. L. 1975a: 31, 1982b: 1278.

cirritus Wood 1974c: 200. Holotype ♀; La Carbonera Experimental Forest, 50 km NW Merida, Merida, Venezuela, 1100 m; Wood Collection. Distribution: South America (Venezuela). Hosts: *Nectandra* sp., *Piper* sp., and other small branches.

References: (tx) Wood, S. L. 1974c: 200.

collaris Blandford 1904: 261. Holotype ♂; Cerro Zamil, Guatemala; BMNH, London. Figures: Wood 1982b: 1284 (female). Distribution: North America (Costa Rica/ Guatemala/ Chiapas in Mexico/ Panama).

Hosts: *Acacia* sp., *Crotalaria* sp., *Miconia globuliflora*, *Phoebe mexicana*, *Theobroma cacao*, and other small branches.

Notes: (3) Schedl 1950i: 160 (described female).

References: (ec) Wichmann 1955a: 107. (hb) Wood, S. L. 1982b: 1285. (ds) Blackwelder 1947: 784; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 366; Wichmann 1955a: 107; Wood, S. L. 1982b: 1285. (tx) Blandford 1904: 261; Hagedorn 1910a: 145; Numberg 1962: 230; Schedl 1950i: 160, 1979c: 59; Wood, S. L. 1967c: 138, 1974d: 279, 1982b: 1285.

splendens Wood 1967c: 138. Holotype ♀; 16 km SE Cargago, Cartago, Costa Rica; Wood Collection. Synonymy: Wood 1974d: 279.

References: (tx) Wood, S. L. 1967c: 138, 1974d: 279.

columbianus Hopkins 1895d: 104. Syntypes, sex?; Raleigh Co., West Virginia [USA]; USNM, Washington.

Figures: Baker 1972: 266, Giese 1967: 56, Kabir & Giese 1966a: 886–891, Wood, S. L. 1982b: 1294 (female).

Distribution: North America (District of Columbia, Florida, Georgia, Indiana, Kansas, Maryland, Massachusetts, Missouri, New Jersey, North Carolina, South Carolina, Tennessee, Vermont, Virginia, West Virginia in USA).

Hosts: *Acer rubrum*, *A. saccharinum*, *Castanea dentata*, *Liriodendron tulipifera*, *Platanus occidentales*, *Quercus alba*, *Ulmus* sp.

References: (ay) Giese 1964, 1967; Hopkins 1894g. (bv) Crozier & Giese 1967b; Gagne & Kearby 1979: 300; Kabir 1963, 1964; Kabir & Giese 1966a; Roling & Kearby 1975b, 1977; Turnbow & Franklin 1980. (cn) Anonymous 1960q, 1961h, 1962h, 1963y, 1965b, 1985k: 23; Baker, W. L. 1972: 265; Beal et al. 1952: 59; Bongberg 1962; Burns 1969, 1970: 1–15; Chamberlin 1924, 1939: 284–285; Coulson & Witter 1984: 566; Doane et al. 1936; Drooz 1985: 369; Felt 1906: 702; Hopkins 1894a: 146, 1894b: 300, 313–336, 1895: 104–107, 1896b: 81, 1896c: 76, 1897a: 37, 1897b: 17, 1898a: 106, 1902d: 24, 1909c: 61, 1910d; Hubbard 1897b: 17; Kabir & Giese 1966a; MacAloney & Ewan 1964; Marlatt 1929: 1–39; Mattson & Haack 1987; McCambridge

& Kowal 1957: 7; McManus & Giese 1967; Nord & McManus 1972; Pettinger 1963: 68; Roling & Kearby 1973, 1977; Snyder 1923: 131, 1927: 3; White, W. B. & Giese 1968; Yates 1968. (ec) Baker, W. L. 1972: 265; Chamberlin 1939: 284–285; Crozier & Giese 1967b; Drooz 1985: 369; Felt 1906: 702; Giese 1962, 1966b, 1967b; Hopkins 1897a: 37; Kabir 1963, 1964: 4894–4895; Kabir & Giese 1966a, 1966b; Mattson & Haack 1987; McManus 1969; McManus & Giese 1968; Nord 1973; Oliveria 1975b; Pettinger & Giese 1968; Roling & Kearby 1974: 1308, 1977; Steinhans 1946: 404; White, W. B. & Giese 1968: 1400; Wilson 1959a: 311–317, 1959b: 114. (hb) Anonymous 1985: 23; Baker, W. L. 1972: 265; Beal & Massey 1945: 105–106; Bright 1976d: 193; Burns 1969; Chamberlin 1939: 284–285; Coulson & Witter 1984: 566; Crozier 1968b: 5062; Crozier & Giese 1967a: 55–58, 1967b: 1203–1214; Doane et al. 1936; Drooz 1985: 369; Felt 1906: 702; Giese 1966a, 1966b, 1967: 55; Hoffmann & St. George 1949: 78; Hopkins 1894a, 1894g, 1904b: 325, 327, 1905c: 146; Hubbard 1897b: 17; Kabir 1963, 1964, 1966; Kabir & Giese 1966a: 887; Lucht & Giese 1961; MacAloney & Ewan 1964; McManus 1969: 2479; McManus & Giese 1966, 1967: 410, 1968; Milne 1968b; Milne & Giese 1969: 225–237, 1970: 12–24; Nord 1972, 1973; Nord & Lewis 1970: 155–157; Nord & McManus 1972; Oliveria 1975b; Pettinger & Giese 1963; Roling & Kearby 1973; Wilson 1959: 114; Wood, S. L. 1982b: 1285; Yates 1968. (ds) Anderson, R. L. & Barry 1950: 48; Anonymous 1963y, 1965b; Atkinson et al. 1991: 161; Beal & Massey 1945: 105–106; Blandford 1904: 253; Blatchley & Leng 1916: 643; Bright 1976d: 193; Chamberlin 1939: 284–285; Deyrup 1981b: 8; Drooz 1985: 369; Hagedorn 1910d: 90; Henshaw 1895: 44; Hoffmann 1940: 60, 1942: 12; Hoffmann & St. George 1949: 78; Kabir & Giese 1966a: 887; Kleine 1913b: 155, 1914b: 394, 1934a: 170; Leng 1920: 339; McManus 1969; Nord & McManus 1972; Schuder 1960a: 23; Smith, J. B. 1910: 400; Swaine 1909: 90; Turnbow & Franklin 1980; Wood, S. L. 1982b: 1285. (tx) Baker, W. L. 1972: 266; Beal & Massey 1945: 105–106; Blandford 1904: 253–254; Blatchley & Leng 1916: 643; Bright 1976d: 193; Chamberlin 1939: 284–285; Giese 1967: 56; Hagedorn 1910a: 145; Hopkins 1895d: 104 1915c: pl. 9, fig. 3, pl. 10, fig. 3; Hubbard 1897b: 17–18; Kabir & Giese 1966a: 886–891; Muesebeck 1942: 95, 1950: 130; Schedl 1940a: 354, 1979c: 60; Swaine 1909: 90–91; Wood, S. L. 1982b: 1285. (ms) Anonymous 1985k: 23; Giese 1966a; Nord & Lewis 1970; Tate 1968.

comatus Blandford 1904: 258. Holotype ♀; Cerro Zunil, Guatemala; BMNH, London.

- Figures: Blandford 1904:pl. 8, Wood 1982b: 1292 (female).
 Distribution: North America (Costa Rica/ Guatemala/ Oaxaca, Puebla in Mexico/ Panama).
 Hosts: *Alnus* sp., *Miconia* sp., *Ochroma* sp., *Phoebe mexicana*, *Solanum torvum*, etc.
 References: (hb) Atkinson et al. 1986: 44; Wood, S. L. 1982b: 1292. (ds) Atkinson & Equihua 1985c: 362; Atkinson et al. 1986: 44; Blackwelder 1947: 784; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 366; Wood, S. L. 1982b: 1292. (tx) Blandford 1904: 258; Hagedorn 1910a: 145; Wood, S. L. 1974d: 279, 1982b: 1292.
splendidus Bright 1972a: 1371. Holotype ♀; Highway 175, 5.6 km S Suchixtepec, Oaxaca, Mexico; CNCI, Ottawa. Synonymy: Wood 1974d: 279.
 References: (tx) Bright 1972a: 1370–1371; McNamara 1977: 195; Wood, S. L. 1974d: 279.
- comosus** Wood 1974c: 186. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
 Figures: Wood 1982b: 1284 (female).
 Distribution: North America (Costa Rica).
 Hosts: *Piper* sp.
 References: (hb) Wood, S. L. 1982b: 1283. (ds) Wood, S. L. 1982b: 1283. (tx) Wood, S. L. 1974c: 186, 1982b: 1283–1285.
- compressicornis** (Fabricius) 1801: 388 (*Bostrichus*). Lectotype ♀; published as *America meridionali*, type labeled Essequibo [Guyana]; UZMC, Copenhagen, designated by Wood 1974c: 202.
 Distribution: South America (Guyana/ Suriname).
 Notes: (3) Wood 1974c: 202 (redescribed type). Citations of this species from Colombia and Central America are based on erroneous identifications (SIW).
 References: (ay) Imhoff 1856: 228. (ds) Blackwelder 1947: 784; Blandford 1904: 254; Bright 1972a: 1385; Dejean 1837; Gemminger & Harold 1872: 2693; Hagedorn 1903b: 547, 1910d: 90; Kleine 1913b: 155; Lacordaire 1866: 385; Melsheimer 1853: 87; Wood, S. L. 1961c: 1. (tx) Blandford 1904: 254; Castelnau 1840; Eggers 1929e, 1931b: 39, 1931c: 184, 1934a: 82; Eichhoff 1868: 421, 1869c: 297, 301, 1878b: 422; Fabricius 1801: 388; Ferrari 1867a: 48–49, 61; Hagedorn 1903b: 547, 1910a: 145; Hopkins 1914: 119; Lacordaire 1866: 385; Lucas 1920: 203; Schedl 1933b: 33, 35, 1950i: 153–154, 1979c: 60; Wood, S. L. 1961c: 1, 1967d: 56, 1974c: 202. (ms) Eichhoff 1868d: 421; Lucas 1920: 203.
- concarus** Bright 1972a: 1376. Holotype ♀; 24 km S Valle Nacional, Oaxaca, Mexico; CNCI, Ottawa.
 Distribution: North America (Oaxaca, Puebla in Mexico).
 Hosts: *Alnus* sp., and a tree limb.
 References: (hb) Wood, S. L. 1982b: 1296. (ds) Wood, S. L. 1982b: 1296. (tx) Bright 1972a: 1376; McNamara 1977: 195; Wood, S. L. 1982b: 1296.
- concisus** Wood 1974c: 201. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
 Figures: Blandford 1904:pl. 8, Wood 1982b: 1307, 1309 (female).
 Distribution: North America (Costa Rica/ Chiapas in Mexico/ Panama).
 Hosts: *Conostegia oerstediana*, *Spondias purpurea*, *Theobroma cacao*.
 Notes: (1) Blandford 1904: 255 (this species was cited erroneously as *compressicornis*).
 References: (hb) Wood, S. L. 1982b: 1309. (ds) Atkinson & Equihua 1986a: 422; Kleine 1914b: 366; Wood, S. L. 1982b: 1309. (tx) Wood, S. L. 1974c: 201, 1982b: 1309.
- consimilis** Wood 1974c: 188. Holotype ♀; 9 km NE Teziutlan, Puebla, Mexico, 1600 m; Wood Collection.
 Figures: Wood 1982b: 1291.
 Distribution: North America (Guatemala/ Puebla in Mexico).
 Hosts: *Acacia* sp., and tree branches.
 References: (hb) Wood, S. L. 1982b: 1288. (ds) Wood, S. L. 1982b: 1288. (tx) Wood, S. L. 1974c: 188, 1982b: 1288.
- convexicauda** Eggers 1931b: 40. Syntypes 1 ♂, 1 ♀; Sao Paulo; Prague Museum, 2 Eggers syntypes in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (ds) Blackwelder 1947: 784; Pedrosa-Macedo & Schonherr 1985: 62; Schedl 1973d: 160, 1976a: 56. (tx) Eggers 1931b: 40–41; Pedrosa-Macedo & Schonherr 1985: 62; Schedl 1966f: 76, 1979c: 64.
- bituberculatus** Numberg 1962: 229. Holotype ♂; Barra do Tapirape, Estado de Matto Grosso, Brasilien; DZSA, Sao Paulo. Synonymy: Schedl 1966f: 76.
 References: (tx) Numberg 1962: 228–231; Schedl 1966f: 76.
- convexifrons** Wood 1986c: 270. Holotype ♀; La Mucuy, 20 km W Merida, Merida, Venezuela; Wood Collection.
 Distribution: South America (Venezuela).
 Hosts: *Nectandra* sp., and a tree branch.
 References: (tx) Wood, S. L. 1986c: 270.
- coronatus** Eggers 1933b: 21. Holotype ♀; Venezuela (Colonia Tovar); MNHN, Paris.
 Distribution: South America (Venezuela).
 References: (ds) Blackwelder 1947: 784. (tx) Eggers 1933b: 2, 21–22.
- curiosus** Bright 1972d: 104. Holotype ♀; Hardwar Cap, 4000', St. Andrew Parish, Jamaica; CNCI, Ottawa.
 Figures: Bright 1972d: 59, 75.
 Distribution: Antilles Islands (Jamaica).
 References: (ds) Bright 1985c: 176. (tx) Bright

- 1972d: 59, 75, 104, 1985c: 176; McNamara 1985c: 176.
- dentatus** Eggers 1943a: 382. Holotype ♂?; Bolivia (Cochabamba); Eggers Collection, in NHMW, Wien. Distribution: South America (Bolivia). References: (tx) Eggers 1943a: 382–383; Schedl 1979c: 77.
- detrimentosus** Schedl 1940a: 355. Syntypes, sex?; Sierra de Durango, Mexico; Schedl Collection in NHMW, Wien. Figures: Atkinson et al. 1986: 44. Distribution: North America (Durango, Guerrero, Morelos in Mexico). Hosts: *Arbutus* sp., etc. References: (hb) Atkinson & Equihua 1985a: 102; Atkinson et al. 1986: 44; Burgos & Saucedo 1983: 141; Wood, S. L. 1982b: 1300. (ds) Atkinson & Equihua 1985a: 102, 1985c: 362; Atkinson et al. 1986: 44; Blackwelder 1947: 784; Burgos & Saucedo 1983: 141; Ferrer 1942; Thomas, J. B. 1966; Wood, S. L. 1982b: 1300. (tx) Atkinson et al. 1986: 44; Schedl 1940a: 355, 1979c: 79; Wood, S. L. 1982b: 1300.
- diligens** Wood 1974c: 190. Holotype ♀; Volcan de Agua, Esquinla, Guatemala; Wood Collection. Figures: Wood 1982b: 1292 (female). Distribution: North America (Guatemala). Hosts: *Alnus* sp., *Chusia* sp., *Ficus* sp., etc. References: (hb) Wood, S. L. 1982b: 1293. (ds) Wood, S. L. 1982b: 1293. (tx) Wood, S. L. 1974c: 190, 1982b: 1293.
- discoideus** Blandford 1904: 262. Holotype ♂; Venezuela; BMNH, London. Distribution: South America (Venezuela). References: (ds) Blackwelder 1947: 784; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 340. (tx) Blandford 1904: 262; Hagedorn 1910a: 145; Schedl 1979c: 92.
- donaticus** Wood 1974c: 198. Holotype ♀; Colonia Tovar, Aragua, Venezuela; Wood Collection. Distribution: South America (Venezuela). Hosts: Small trees and branches. References: (tx) Wood, S. L. 1974c: 198.
- dubiosus** (Schedl) 1976a: 81 (*Metacorthylus*). Holotype ♂; Encruzilhada, Bahia, Brazil; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1976a: 81.
- eichhoffi** Schedl 1933b: 34. Holotype ♀; Costa Rica, Iberia, Sta. Clara [district in Limon Province]; Schedl Collection in NHMW, Wien. Distribution: North America (Costa Rica). References: (ds) Blackwelder 1947: 784; Wood, S. L. 1982b: 1310. (tx) Schedl 1933b: 34–35, 1979c: 87; Wood, S. L. 1982b: 1310.
- emarginatus** Eggers 1943a: 380. Lectotype ♂; Bolivia (Cochabamba); USNM, Washington, designated by Anderson & Anderson 1971: 12. Distribution: South America (Bolivia). Notes: (3) Eggers 1943a: 380 (cites *bicolor* Eggers, nomen nudum, synonymy). References: (tx) Anderson, W. H. & Anderson 1971: 12; Eggers 1943a: 380; Schedl 1979c: 90; Wood, S. L. 1966b: 22.
- excisus** (Ferrari) 1867a: 71 (*Morizus*). Holotype, sex?; Venezuela [probably Colonia Tovar]; NHMW, Wien. Distribution: South America (Venezuela). References: (ds) Blackwelder 1947: 784; Blandford 1904: 253; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 90; Kleine 1914b: 340. (tx) Blandford 1904: 253; Eichhoff 1868d: 419, 1869c: 301; Ferrari 1867a: 71, 1867b: 255; Hagedorn 1910a: 145; Hopkins 1914: 125. (ms) Eichhoff 1868d: 419.
- exiguus** Wood 1984: 113. Holotype ♀; Cuatenango, Guerrero, Mexico, 1650 m; Wood Collection. Distribution: North America (Guerrero in Mexico). References: (ds) Kleine 1913b: 155. (tx) Wood, S. L. 1984: 113.
- flagellifer** Blandford 1904: 255. Syntypes, sex?; Toxpan in Mexico, San Geronimo and Pantaleon in Guatemala, and Boquete in Panama; BMNH, London. Figures: Lara & Shenefelt 1965: 174, 176, Wood 1982b: 1304 (female). Distribution: North America (Guatemala/ Chiapas, Morelos, Nayarit, Veracruz in Mexico/ Panama). Hosts: *Acacia* sp., *Persca americana*, and other tree branches. References: (ce) Wichmann 1955a: 107. (hb) Atkinson & Equihua 1985c: 363; Atkinson et al. 1986: 45; Burgos & Saucedo 1983: 143; Lara & Shenefelt 1965: 175; Moreira 1928: 24–25; Wood, S. L. 1982b: 1305. (ds) Atkinson & Equihua 1985c: 363; Atkinson et al. 1986: 45; Blackwelder 1947: 784; Burgos & Saucedo 1983: 143; Ferrer 1942; Guagliumi 1966: 218; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 353, 366; Roba 1935: 338; Wichmann 1955a: 107; Wood, S. L. 1982b: 1305. (tx) Blandford 1904: 255; Eggers 1931b: 41; Hagedorn 1910a: 145; Lara & Shenefelt 1965: 174–176; Schedl 1940a: 350, 1979c: 97; Wood, S. L. 1966b: 22, 1982b: 1305.
- cirrus** Schedl 1940a: 351. Lectotype ♂; Xochitlan, Morelos, Mexico; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 22. References: (ds) Blackwelder 1947: 784; Ferrer 1942. (tx) Schedl 1940a: 351–354, 1979c: 58; Wood, S. L. 1966b: 22.
- nudiusculus** Schedl 1950i: 156. Holotype ♀; Mexico, Comitán; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 22. References: (tx) Schedl 1950i: 156, 1979c: 173; Wood, S. L. 1966b: 22.

fuscus Blandford 1904: 262. Syntypes ♀; "Brazilian" tobacco intercepted at Paris [actual origin was Mexico]; BMNH, London.

Figures: Bustamante & Atkinson 1984: 88. Oranegui & Atkinson 1984: 88.

Distribution: North America (El Salvador/ Guatemala/ Distrito Federal, Mexico in Mexico).

Hosts: *Coffea* sp., *Malus* sp., *Pyrus* sp.

References: (cn) Bustamante & Atkinson 1984; Oranegui & Atkinson 1984. (cc) Wichmann 1955a: 107. (hb) Atkinson & Equihua 1985a: 102. 1985c: 362; Atkinson et al. 1986: 45; Bustamante & Atkinson 1984; Oranegui & Atkinson 1984; Wood, S. L. 1982b: 1283. (ds) Atkinson & Equihua 1985a: 102. 1985c: 362; Atkinson et al. 1986: 45; Blackwelder 1947; Hagedorn 1910d: 90; Kleine 1913b: 155; Wichmann 1955a: 107; Wood, S. L. 1982b: 1289. (tx) Blandford 1904: 262; Bustamante & Atkinson 1984: 88; Hagedorn 1910a: 145; Oranegui & Atkinson 1984: 88; Schedl 1950i: 158; Wood, S. L. 1982b: 1289.

gracilis (Schedl) 1972g: 74 (*Cortylomimus*). Holotype, sex?; Brasilien, Corcovado, Gnanabara; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

Notes: (3) A note written by Schedl in his collection gave the genus as *Cortheloderes*, nomen nudum (SLW).

References: (tx) Schedl 1972g: 74, 1979c: 107.

granulatus Schedl 1935h: 345. Holotype ♀; Costa Rica, Turrialba, 800 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: (ds) Blackwelder 1947: 784; Wood, S. L. 1982b: 1289. (tx) Schedl 1935h: 345–346. 1979c: 110; Wood, S. L. 1982b: 1289.

granulifer Wood 1974c: 181. Holotype ♂; Volcan Poas, Heredia, Costa Rica, 2500 m; Wood Collection.

Distribution: North America (Costa Rica).

Hosts: *Myrica pubescens*, and a tree branch.

References: (tx) Wood, S. L. 1974c: 181.

ingaensis (Schedl) 1939n: 14 (*Metacorthylus*). Holotype ♂; Colombia: El Collegio; BMNH, London.

Distribution: South America (Colombia).

Hosts: *Inga* sp.

References: (ds) Blackwelder 1947: 782. (tx) Schedl 1939n: 14, 1954b: 43, 1979c: 124.

insignis Wood 1974c: 200. Holotype ♀; 24 km E Barbosa, Antioquia, Colombia, 1200 m; Wood Collection.

Distribution: South America (Colombia).

Hosts: *Inga* sp.

References: (tx) Wood, S. L. 1974c: 200.

letzneri Ferrari 1867a: 55, 59. Syntypes, sex?; Venezuela [probably Colonia Tovar]; NHMW, Wien.

Distribution: South America (Colombia/Venezuela).

References: (ds) Blackwelder 1947: 784;

Blandford 1904: 254; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 340. (tx) Blandford 1904: 254; Ferrari 1867a: 55, 59; Hagedorn 1910a: 145; Hopkins 1914: 128; Schedl 1950i: 146; Wood, S. L. 1977b: 208, 1977d: 511.

castaneus Ferrari 1867a: 55, 59. Holotype ♂; Venezuela [probably Colonia Tovar]; NHMW, Wien. Synonymy: Schedl 1950i: 146.

References: (ds) Blackwelder 1947: 784; Blandford 1904: 258; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 340. (tx) Blandford 1904: 258; Ferrari 1867a: 55, 59; Hagedorn 1910a: 145; Schedl 1950i: 146, 158.

strigilatus Eggers 1933b: 20. Holotype ♂; Venezuela (Colonia Tovar); MNHN, Paris. Synonymy: Wood 1977b: 208.

References: (ds) Blackwelder 1947: 784. (tx) Eggers 1933b: 2, 20–21; Wood, S. L. 1977b: 208.

columbianus Schedl 1950i: 158. Holotype ♀; Columbia, Comatan; Schedl Collection in NHMW, Wien, preoccupied by Hopkins 1895. Synonymy: Wood 1977b: 208.

References: (cn) Blackman 1950. (hb) Blackman 1950. (ds) Blackman 1950. (tx) Schedl 1950i: 158, 1952d: 347, 1979c: 60; Wood, S. L. 1977b: 208.

ater Schedl 1952d: 347. Holotype ♀; Columbia, Comatan; Schedl Collection in NHMW, Wien, automatic. Synonymy: Wood 1977b: 208.

References: (tx) Schedl 1952d: 347, 1979c: 60; Wood, S. L. 1977b: 208.

luridus Blandford 1904: 256. Holotype ♀; Volcan de Chiriqui, Chiriqui, Panama; BMNH, London.

Distribution: Antilles Islands (Guadeloupe), North America (Guatemala/ Panama).

References: (ds) Blackwelder 1947: 784; Bright 1985c: 176; Hagedorn 1910d: 90; Kleine 1913b: 155, 1914b: 366; Wood, S. L. 1982b: 1300. (tx) Blandford 1904: 256; Bright 1985c: 176; Hagedorn 1910a: 145; Wood, S. L. 1982b: 1300.

biseriatus Eggers 1943d: 247. Holotype ♀; Guatemala; IPKE, Eberswalde. Synonymy: Wood 1982b: 1300.

References: (tx) Eggers 1943d: 247; Numberg 1962: 229; Wood, S. L. 1982b: 247.

lustratus Wood 1984: 113. Holotype ♀; "Texecal," Maiote Poztlan, Morelos, Mexico; Wood Collection.

Distribution: North America (Morelos in Mexico).

References: (ds) Atkinson et al. 1986: 68; Wood, S. L. 1984: 113.

macrocerus Eichhoff 1869a: 279. Holotype ♀; Colombia; IRSNB, Brussels.

Distribution: South America (Colombia/ Ecuador).

References: (ds) Blackwelder 1947: 784; Blandford 1904: 254; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 90; Kleine 1913b: 155,

- 1914b: 342. (tx) Blandford 1904: 254; Eichhoff 1869a: 279–280, 1878b: 423; Hagedorn 1910a: 145.
- mexicanus** Schedl 1950i: 159. Holotype ♂; Mexico, Comitán; Schedl Collection in NHMW, Wien. Figures: Wood 1982b: 1301 (female). Distribution: North America (Honduras/Chiapas, Puebla, Veracruz in Mexico). Hosts: *Clethra hondurensis*, *Lippia substrigosa*, *Malus* sp., *Orcopanax* sp., *Parathesis serrulata*, *Pinus pseudostrubus*, *Rapanea guyanensis*, *Salix* sp. References: (hb) Atkinson & Equilma 1955a: 103; Atkinson et al. 1956: 46; Burgos & Saucedo 1953: 141; Wood, S. L. 1952b: 1301. (ds) Atkinson & Equilma 1955a: 103, 1955c: 362; Atkinson et al. 1956: 46; Burgos & Saucedo 1953: 141; Wood, S. L. 1952b: 1301. (tx) Atkinson et al. 1956: 46; Nunberg 1962: 229; Schedl 1950i: 156, 159, 1979c: 152; Wood, S. L. 1973c: 175, 1982b: 1301.
- glabrinus** Bright 1972a: 1372. Holotype ♀; 14 km SW Teopisca, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1973c: 175. References: (tx) Bright 1972a: 1370–1372; McNamara 1977: 195; Wood, S. L. 1973c: 175.
- micacirrus** Wood 1984: 114. Holotype ♀; Chilapa, Guerrero, Mexico; Wood Collection. Distribution: North America (Guerrero in Mexico). Hosts: *Ardesia* sp. References: (tx) Wood, S. L. 1984: 114.
- minimus** Wood 1974c: 192. Holotype ♀; La Ceiba, Atlantida, Honduras; Wood Collection. Distribution: North America (Honduras). References: (ds) Wood, S. L. 1982b: 1297. (tx) Wood, S. L. 1974c: 192, 1982b: 1297.
- minutissimus** Schedl 1940a: 353. Lectotype ♀; Mexico: Chiltepec (Oaxaca); Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 156. Distribution: North America (Honduras/ Oaxaca in Mexico). Hosts: *Sivietenia* sp. References: (hb) Wood, S. L. 1982b: 1299. (ds) Blackwelder 1947: 784; Ferrer 1942; Wood, S. L. 1982b: 1299. (tx) Schedl 1940a: 353–354, 1979c: 156; Wood, S. L. 1982b: 1299.
- minutus** Bright 1972a: 1373. Holotype ♀; Lagos de Colores (Lagunas de Montebello National Park), Chiapas, Mexico; CNCI, Ottawa. Figures: Bright 1972a: 1370. Distribution: North America (Chiapas in Mexico). Hosts: *Quercus* sp. References: (ds) Wood, S. L. 1982b: 1303. (tx) Bright 1972a: 1370, 1373; McNamara 1977: 195; Wood, S. L. 1982b: 1303.
- mirabilis** Nunberg 1962: 226. Holotype ♀; Estacao Biologica de Boracéia, Salesopolis, Estado de Sao Paulo, Brasilien; IZW, Warsaw. Figures: Nunberg 1962: 225, 228. Distribution: South America (Brazil).
- References: (tx) Nunberg 1962: 225–226, 228.
- nanus** Wood 1979b: 138. Holotype ♂; 1 km SW Rincon de Osa, Puntarenas, Costa Rica; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Cecropia* sp. leaf petiole. References: (ds) Wood, S. L. 1982b: 1298. (tx) Wood, S. L. 1979b: 138, 1982b: 1298.
- neotardus** Schedl 1950d: 120 (*Corthy cyclon*). Holotype ♂; Caruaru, Pernambuco, Brazil; Schedl Collection in NHMW, Wien, automatic. Distribution: South America (Brazil). References: (tx) Schedl 1950d: 120.
- tardus** Schedl 1976a: 85 (*Corthy cyclon*). Holotype ♂; Caruaru, Pernambuco, Brazil; Schedl Collection in NHMW, Wien, preoccupied by Wood 1974. References: (tx) Schedl 1976a: 85, 1980d: 120; Wood, S. L. 1981: 122.
- tardulus** Wood 1981: 122. Holotype ♂; Caruaru, Pernambuco, Brazil; Schedl Collection in NHMW, Wien, automatic, an unneeded replacement name. References: (tx) Wood, S. L. 1981: 122.
- nevermanni** (Schedl) 1939j: 568 (*Thylurcos*). Holotype ♂; Hamburgfarm on Rio Reventazon, Limón, Costa Rica; Schedl Collection in NHMW, Wien. Distribution: North America (Costa Rica). Notes: (1) Wood 1982b: 1298 (to *Corthy lylus*). References: (ds) Wood, S. L. 1982b: 1298. (tx) Schedl 1939j: 568, 1951m: 127, 1979c: 165; Wood, S. L. 1982b: 1298.
- niger** (Schedl) 1954b: 43 (*Metacorthy lylus*). Lectotype ♀; Brasilien: Sta. Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, designated by Schedl 1979c: 166. Distribution: South America (Brazil). References: (ds) Schedl 1966f: 91, 1970e: 85. (tx) Schedl 1954b: 43, 1979c: 166.
- nolenae** Wood 1974c: 194. Holotype ♀; Huajuapán, 21 km SE Oaxaca, Oaxaca, Mexico; Wood Collection. Distribution: North America (Oaxaca in Mexico). Hosts: *Nolena* sp. fruiting stalk. References: (hb) Wood, S. L. 1982b: 1302. (ds) Wood, S. L. 1982b: 1302. (tx) Wood, S. L. 1974c: 194, 1982b: 1302.
- nudipennis** Schedl 1950i: 155. Syntypes ♀; Brazil, Santa Catarina, Nova Teutonia; Schedl Collection in NHMW, Wien, and Plaumann Collection. Figures: Pedrosa-Macedo & Schonherr 1985: 63. Distribution: South America (Brazil). Notes: (1) Schedl 1979c: 173 (citation of holotype invalid). References: (ds) Pedrosa-Macedo & Schonherr 1985: 63; Schedl 1967d: 4, 1972f: 47. (tx) McNamara 1977: 1195; Pedrosa-Macedo & Schonherr

- 1955: 63; de Ruette 1970: 90; Schedl 1950i: 155, 1961i: 229, 1979c: 172.
- nudus** Schedl 1940a: 354. Syntypes ♀; Mexico: San Antonio Nexapa, Chiapas; Dampf Collection and Schedl Collection in NHMW, Wien.
Distribution: North America (Chiapas, Distrito Federal, Mexico, Michoacan in Mexico).
Notes: (1) Schedl 1979c: 173 (citation of holotype invalid).
References: (hb) Atkinson & Equihua 1955a: 102; Atkinson et al. 1986: 46; Burgos & Saucedo 1983: 140; Wood, S. L. 1982b: 1289. (ds) Atkinson & Equihua 1955a: 102; Atkinson et al. 1986: 46; Blackwelder 1947: 784; Burgos & Saucedo 1983: 140; Ferrer 1942; Wood, S. L. 1982b: 1289. (tx) Schedl 1940a: 354, 1979c: 173; Wood, S. L. 1982b: 1289.
- obliquus** Schedl 1976a: 79. Holotype ♀; Represa Rio Grande, Guanabara [Brazil]; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 79.
- oculatus** Wood 1974c: 184. Holotype ♀; Pandora, Limon, Costa Rica; Wood Collection.
Distribution: North America (Costa Rica).
References: (hb) Wood, S. L. 1982b: 1289. (ds) Wood, S. L. 1982b: 1289. (tx) Wood, S. L. 1974c: 184, 1982b: 1289.
- oliveirai** Schedl 1976a: 80. Holotype ♂; Represa Rio Grande, Guanabara [Brazil]; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 80.
- panamensis** Blandford 1904: 259. Lectotype ♀; Volcan de Chiriqui, Chiriqui, Panama; BMNH, London, designated by Wood 1982b: 1287.
Figures: Blandford 1904: pl. 8, Wood 1982b: 1288 (female).
Distribution: North America (Costa Rica/ Guatemala (?)/ Panama).
Hosts: *Phoebe mexicana*, etc.
References: (hb) Wood, S. L. 1982b: 1287. (ds) Blackwelder 1947: 784; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 366; Schedl 1976a: 56; Wood, S. L. 1982b: 1287. (tx) Blandford 1904: 259; Hagedorn 1910a: 145; Schedl 1933b: 33, 1961i: 230, 1979c: 183; Wood, S. L. 1982b: 1287.
- papulans** Eichhoff 1869a: 280. Holotype, sex?; Brasilia; IRSNB, Brussels.
Figures: Wood 1982b: 1304 (female).
Distribution: Antilles Islands (Eleuthera Islands in Bahamas Islands/ Grenada), North America (Costa Rica/ Honduras/ Michoacan, Nayarit, Oaxaca, Veracruz in Mexico/ Panama/ Florida in USA), South America (Brazil/ Cayenne/ Guyana/ Venezuela).
Hosts: *Persea americana*, *Salix guatemalensis*, *Spondias mombin*, *S. purpurea*, *Theobroma cacao*, and other tree branches.
References: (ds) Blackwelder 1947: 784; Blandford 1904: 254; Gemminger & Harold 1872: 2693; Hagedorn 1903b: 547, 1910d: 91; Kleine 1913b: 155, 1914b: 337; Nimberg 1958a: 480; Pedrosa-Macedo & Schonherr 1985: 64; Schedl 1967d: 4, 1972g: 47, 1973d: 160. (tx) Blandford 1904: 254; Eggers 1931b: 42, 1933b: 2; Eichhoff 1869a: 280, 1869c: 301, 1878b: 427; Hagedorn 1903b: 547, 1910a: 145; Nimberg 1958a: 480; Pedrosa-Macedo & Schonherr 1985: 64.
- spinifer** Schwarz 1891b: 114. Syntypes ♀; Key West, Florida [USA]; USNM, Washington.
Synonymy: Eggers 1933b: 2.
References: (bv) Atkinson, Foltz, & Connor 1988. (ec) Equihua & Atkinson 1986: 633; Wilson 1959b: 118. (hb) Atkinson, Foltz, & Connor 1988; Atkinson et al. 1986: 46, 68; Burgos & Saucedo 1983: 142; Chamberlin 1939: 285; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 633; Kabir & Giese 1966a; Ostmark 1968; Schwarz 1891b: 114; Wood, S. L. 1982b: 1304. (ds) Atkinson & Equihua 1985c: 362, 1986a: 422, 1988: 88; Atkinson et al. 1986: 46, 68, 1991: 162; Blatchley & Leng 1916; Bright 1982b: 164, 1985c: 176; Burgos & Saucedo 1983: 142; Chamberlin 1939: 285; Deyrup & Atkinson 1987a: 66; Equihua & Atkinson 1986: 633; Estrada & Atkinson 1988: 211; Hagedorn 1910d: 91; Henshaw 1895: 44; Kabir & Giese 1966a: 884; Kleine 1913b: 155, 1914b: 384, 1934a: 170; Leng 1920: 339; Ostmark 1968; Schwarz 1891b: 114; Swaine 1909: 91; Wood, S. L. 1977a: 69, 1982b: 1304. (tx) Blatchley & Leng 1916; Bright 1985c: 176; Chamberlin 1939: 285; Eggers 1933b: 2; Hagedorn 1910a: 145; Hopkins 1895d: 104; Kabir & Giese 1966a: 884; Schedl 1940a: 350; Schwarz 1891b: 114; Swaine 1909: 91; Wood, S. L. 1976a: 347, 1977c: 208, 1977d: 512, 1982b: 1304.
- affinis** Fonseca 1925a: 3 (*Metacorthylus*). Syntypes, sex?; Itatiba, Sao Paulo, Brazil; not located, probably in DZSA, Sao Paulo. Synonymy: Eggers 1933b: 2, Wood 1982b: 1304.
References: (cn) Costa Lima 1956. (hb) Costa Lima 1956. (ds) Blackwelder 1947: 784; Costa Lima 1936, 1956; Roba 1935: 338; Schedl 1966f: 91, 1970e: 85, 1973d: 160. (tx) Costa Lima 1956; Fonseca 1925a: 3, 1927: 583–590; Wood, S. L. 1977d: 512, 1982b: 1304.
- affinis** Fonseca 1927: 585. Syntypes, sex?; Itatiba, Sao Paulo, Brazil; not located, probably in DZSA, Sao Paulo. Synonymy: Eggers 1933b: 2, Wood 1982b: 1304.
References: (ds) Kleine 1934a: 171. (tx) Fonseca 1927: 583–590; Wood, S. L. 1977d: 512, 1982b: 1304.
- guayanensis** Eggers 1933b: 22. Syntypes ♂;

- Camopi, Franz, Guayana: MNHN, Paris. Synonymy: Eggers 1933b: 2, Wood 1977b: 208.
References: (ds) Blackwelder 1947: 784. (tx) Eggers 1933b: 2, 22; Wood, S. L. 1977b: 208.
- tommentosus* Schedl 1940a: 350. Syntypes ♀; Mexico: Tuxtepec, Oaxaca; Dampf Collection and Schedl Collection in NHMW, Wien. Synonymy: Wood 1976a: 347.
Notes: (3) Part of the Dampf Collection went to Schedl, part to an Italian collector whose residence was Rome, and part to the Museo Nacional de Historia Natural or Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad México (SLW).
References: (ds) Blackwelder 1947: 784. (tx) Schedl 1940a: 350–351, 1979c: 253; Wood, S. L. 1976a: 347.
- parvulus* Blandford 1904: 261. Holotype ♂; Las Mercedes, Guatemala; BMNH, London.
Distribution: North America (Guatemala).
Notes: (3) *Corthyliis uniseptis* Schedl could be a synonym (SLW).
References: (ds) Blackwelder 1947: 784; Hagedorn 1910d: 91; Kleine 1913b: 155. (tx) Blandford 1904: 261; Ferrari 1867a: 53; Hagedorn 1910a: 145; Schedl 1940a: 353; Wood, S. L. 1982b: 1299.
- peruanus* Schedl 1950i: 155. Holotype ♂; Callanga, Peru; [probably from Eggers Collection] in NHMW, Wien.
Distribution: South America (Peru).
References: (tx) Schedl 1935h: 345–346, 1950i: 154–155, 1979c: 191.
- petilus* Wood 1967d: 56. Holotype ♀; 5 km W El Salto, Durango, Mexico; Wood Collection.
Figures: Wood 1967d: 54, 1982b: 1233.
Distribution: North America (Durango in Mexico/Arizona in USA).
Hosts: *Quercus hypoleucoides*, Q. sp.
References: (tx) Wood, S. L. 1967d: 54, 56–57, 1982b: 1233, 1308.
- pharax* Schedl 1976a: 80. Holotype ♀; Carnaru, Pernambuco, Brasil; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (hb) Wood, S. L. 1982b: 1308. (ds) Pedrosa-Macedo & Schonherr 1985: 65; Wood, S. L. 1982b: 1308. (tx) Pedrosa-Macedo & Schonherr 1985: 65; Schedl 1976a: 80; Wood, S. L. 1982b: 1308.
- pisinnus* Bright 1972d: 103. Holotype ♀; Penlyne Castle, St. Thomas Parish, Jamaica; CNCI, Ottawa.
Distribution: Antilles Islands (Jamaica).
References: (ds) Bright 1985c: 176. (tx) Bright 1972d: 103, 1985c: 176; McNamara 1977: 195.
- praealtus* Schedl 1976a: 80. Holotype ♂; Brasilien, Rio Negro; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 80.
- praeustus* Schedl 1950i: 153. Lectotype ♀; Guatemala; Schedl Collection in NHMW, Wien, designated by Wood 1982b: 1308.
Distribution: North America (Guatemala), South America (Peru).
Notes: (1) Schedl 1979c: 198 (citation of holotype invalid).
References: (ds) Blackwelder 1947: 784; Wood, S. L. 1982b: 1308. (tx) Eggers 1931c: 184; Schedl 1950i: 153, 1979c: 198; Wood, S. L. 1982b: 1308.
- procerus* Bright 1972a: 1376. Holotype ♀; 10 km S Valle Nacional, Oaxaca, Mexico; CNCI, Ottawa.
Figures: Bright 1972a: 1370, 1377, Wood 1982b: 1307 (female).
Distribution: North America (Costa Rica/ Oaxaca in Mexico/ Panama).
Hosts: *Siparuna nicaraguensis*, and a liana.
References: (hb) Wood, S. L. 1982b: 1310. (ds) Wood, S. L. 1982b: 1310. (tx) Bright 1972a: 1370, 1376–1377; McNamara 1977: 195; Wood, S. L. 1982b: 1310.
- ptyocerus* Blandford 1904: 257. Lectotype ♀; Volcan de Chiriquí, Chiriquí, Panama; BMNH, London, designated by Wood 1982b: 1295.
Distribution: North America (Panama).
References: (ds) Blackwelder 1947: 784; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 366; Wood, S. L. 1982b: 1295. (tx) Blandford 1904: 257; Hagedorn 1910a: 145; Wood, S. L. 1982b: 1295.
- pumilus* Wood 1974c: 192. Holotype ♀; Madden Forest, Canal Zone, Panama, 30 m; Wood Collection.
Distribution: North America (Panama).
Hosts: Tree branch.
References: (hb) Wood, S. L. 1982b: 1297. (ds) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 1297. (tx) Wood, S. L. 1974c: 192, 1982b: 1297.
- punctatissimus* (Zimmermann) 1868: 144 (*Crypturgus*). Syntypes, sex?; South Carolina [USA]; not located.
Figures: Wood 1982b: 1288 (female).
Distribution: North America (Ontario in Canada/ Arkansas, Colorado, Connecticut, District of Columbia, Florida, Georgia, Illinois, Kentucky, Massachusetts, Michigan, Minnesota, Mississippi, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, Tennessee, Vermont, Virginia, West Virginia in USA).
Hosts: *Acer saccharinum*, *Berberis* sp., *Carpinus* sp., *Cercocarpus* sp., *Cornus* sp., *Corylus* sp., *Gaylussacia* sp., *Ostrya* sp., *Rhododendron* sp., *Sassafras* sp., *Vaccinium* sp., *Viburnum betulinoides*.
References: (ay) Finnegan 1963, 1967; Hopkins 1894g. (bv) Atkinson, Foltz, & Connor 1988; Meixner 1937: 1218; Rose, A. H. & Lindquist 1982b: 133. (cn) Adrien 1947: 113–117; Baker, W. L. 1972: 265; Beal et al. 1952: 46; Blackman 1950; Boyd 1945: 114–115, 1953: 139–140; Comtois

1988: 182; Daviault 1948: 30; Doane et al. 1936; Drooz 1985: 370; Felt 1905: 50, 65, 1906: 732, 1913: 427, 1915: 36, 1924: 44, 1926: 44, 1933: 45–51; Felt & Rankin 1932: 328, 437; Finnegan 1967; Finnegan, McPhee, & Watson 1959: 2; Fisher, Thompson, & Webb 1953; Hamilton 1933: 62–63; Headlee 1916, 1917: 306–335; Herrick 1935: 179, 322; Howse, Gross, & Rose 1981: 61; Hubbard 1897b: 16–17; Langford & Cory 1939: 50; Lindquist, O. H. & Syme 1981: 32; Martineau 1964b: 45; Packard 1890: 387–390; Pirone 1941: 376; Robert 1947a: 113; Roeper et al. 1987; Rose, A. H. & Lindquist 1982a, 1982b: 133; Sippell, Dance, & Rose 1966: 57; Sippell, MacDonald, & Rose 1960: 56–57, 1961: 56, 1962: 61, 1963: 63, 1964: 62, 1965: 63; Sorachi 1934: 34, 42, 1937: 50, 1941: 50–51; Syme & Nystrom 1988: 33; Twinn 1934: 124; Weigel & Baumhofer 1948: 73–74; White 1933: 631–640; White & Hamilton 1935: 15. (**cc**) Baker, W. L. 1972: 265; Comtois 1988: 182; Felt 1905, 1906: 65, 1915: 36; Finnegan 1963; Roeper 1988; Rose, A. H. & Lindquist 1982b: 133; Rumbold 1931c: 849; Steinhaus 1946: 404; Wilson 1959b: 115. (**hb**) Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 265; Beal & Massey 1945: 104–105; Bellevoye 1898; Blackman 1950; Boyd 1953: 139; Bright 1976d: 192; Chamberlin 1939: 284; Chittenden 1890; Comtois 1988: 182; Deyrup & Atkinson 1987a: 66; Doane et al. 1936; Drooz 1985: 370; Felt 1905, 1906: 65, 1915: 36, 1926: 44; Felt & Rankin 1932: 328, 437; Finnegan 1967; Herrick 1935: 179, 322; Hopkins 1894d: 145, 281, 1894g: 277, 1905c: 145; Hubbard 1897a: 423, 1897b: 16; Kabir & Giese 1966a; Lengerken 1954: 312; Merriam 1883: 84–86; Packard 1890: 387; Pierce, W. D. 1907: 290; Robert 1947a: 113; Roeper et al. 1987; Rose, A. H. & Lindquist 1982a, 1982b: 133; Schwarz 1890: 178, 1891b: 109–115; Smith, E. F. 1896: 318; Sorachi 1937: 50; Weiss 1916: 105; Wenzel 1906: 37; Wood, S. L. 1982b: 1290. (**ds**) Anonymous 1926c: 517, 1964h; Atkinson et al. 1991: 162; Beal & Massey 1945: 105–106; Beaulne 1956; Blackman 1950; Blandford 1904: 253; Blatchley & Leng 1916: 642; Bright 1976d: 192; Chamberlin 1939: 284; Chittenden 1890; Deyrup 1981b: 8; Deyrup & Atkinson 1987a: 66, 1987b: 68; Dodge 1938; Drooz 1985: 370; Felt 1915: 36, 1926: 44; Felt & Rankin 1932: 328, 437; Finnegan 1963: 137, 1967: 50; Fox-Wilson 1938: 2, 318; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 91; Hamilton 1895a: 346, 378; Henshaw 1885: 147; Hopkins 1893a: 127, 1893b: 207; Kabir & Giese 1966a: 884; Kirk 1970; Kleine 1913b: 155, 1914b: 384, 1934a: 170; Knull 1932: 66; Leng 1920: 339; Leonard 1928: 517; Lindquist, O. H. & Syme 1981: 32; Robert 1947a: 113; Roeper et al. 1987; Schwarz 1895b: 109; Smith, J. B. 1900: 361, 1910: 400; Swaine 1909: 91; Syme & Nystrom 1988: 33; Thatcher, T. O. 1951: 80;

Weiss 1915: 473; Wood, S. L. 1982b: 1290. (**tx**) Beal & Massey 1945: 104–105; Benoit 1985: 78; Blandford 1904: 253–254; Blatchley & Leng 1916: 642; Bright 1976d: 192, 204, 214; Chamberlin 1939: 284; Dodge 1938: 13, 38; Eichhoff 1878b: 428, 460; Finnegan 1967; Hagedorn 1910a: 145; Hopkins 1895d: 104; Hubbard 1897b: 16; Jacques 1951: 353; Kabir & Giese 1966a: 884; LeConte 1868: 154, 1876: 347; LeConte & Horn 1883: 517; Lindquist, O. H. & Syme 1981: 32; Meixner 1937: 218; Schedl 1940a: 354, 1950i: 155, 1960a: 77; Schwarz 1891b: 109; Swaine 1909: 91; Syme & Nystrom 1988: 33; Wood, S. L. 1982b: 1290; Zimmerman 1868: 144.

punctatus Eggers 1943a: 382. Holotype ♀; Bolivia (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia).

References: (**cn**) Lavalley & Benoit 1978: 42; Pirone, Dodge, & Rickett 1960: 615; Sippell, Dance, & Rose 1966: 57; Sippell, MacDonald, & Rose 1960: 57. (**tx**) Anderson, W. H. & Anderson 1971: 26; Eggers 1943a: 382.

pusillus Eggers 1943a: 380. Holotype ♂; Bolivia (Cochabamba); USNM, Washington.
Distribution: South America (Bolivia).
References: (**tx**) Anderson, W. H. & Anderson 1971: 27; Eggers 1943a: 380–381.

pygmaeus Wood 1974c: 195. Holotype ♀; Finca La Lola, Limon, Costa Rica; Wood Collection.
Figures: Wood 1982b: 1301 (female).
Distribution: North America (Costa Rica).
Hosts: *Theobroma cacao*, and a tree branch.
References: (**hb**) Wood, S. L. 1982b: 1303. (**ds**) Atkinson & Equihua 1985: 86; Wood, S. L. 1982b: 1303. (**tx**) Wood, S. L. 1974c: 195, 1982b: 1303.

redtenbacheri Ferrari 1867a: 55, 60, 70. Syntypes ♂; Venezuela [probably Colonia Tovar]; NHMW, Wien.

Distribution: South America (Venezuela).
References: (**ds**) Blackwelder 1947: 784; Blandford 1904: 259; Gemminger & Harold 1872: 2680; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 340. (**tx**) Blandford 1904: 259; Ferrari 1867a: 55, 60, 70; Hagedorn 1910a: 145; Schedl 1935h: 345, 1940a: 354, 1950i: 146.

retusifer Wood 1974c: 183. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Figures: Wood 1982b: 1280 (female).
Distribution: North America (Costa Rica).
Hosts: *Phoebe mexicana*, a tree branch, and a liana.
References: (**hb**) Wood, S. L. 1982b: 1281. (**ds**) Wood, S. L. 1982b: 1281. (**tx**) Wood, S. L. 1974c: 183, 1982b: 1281.

retusus Wood 1974c: 182. Holotype ♀; Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Figures: Wood 1982b: 1280 (female).
Distribution: North America (Costa Rica).

- Hosts: *Miconia* sp., *Phoebe mexicana*, *Solanum torvum*.
References: (hb) Wood, S. L. 1952b: 1279. (ds) Wood, S. L. 1952b: 1279. (tx) Wood, S. L. 1974c: 182, 1952b: 1279.
- robustus** Schedl 1936i: 108. Holotype ♀; Brasilien, Nova-Teutonia; Schedl Collection in NHMW, Wien. Figures: Pedrosa-Macedo & Schonherr 1955: 66. Distribution: South America (Brazil).
Hosts: *Inga* sp.
Notes: (3) Schedl 1950d: 120 (described male).
References: (ds) Blackwelder 1947: 754; Pedrosa-Macedo & Schonherr 1955: 66; Schedl 1976a: 56; Schonherr & Pedrosa-Macedo 1951: 54. (tx) Pedrosa-Macedo & Schonherr 1955: 66; Schedl 1936i: 108, 1939j: 564, 1979c: 212, 1950d: 120.
- rubricollis** Blandford 1904: 260. Lectotype ♀; Volcan de Chiriquí, Chiriquí, Panama; BMNH. London, designated by Wood 1952b: 1254. Figures: Blandford 1904: pl. 8, Wood 1952b: 1254 (female).
Distribution: North America (Costa Rica/ Guatemala (?)/ Panama), South America (Colombia).
Hosts: *Miconia caudata*, *Myrica pubescens*, *Werklia insignis*.
References: (hb) Wood, S. L. 1952b: 1256. (ds) Blackwelder 1947: 754; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 366; Schedl 1961c: 1; Wood, S. L. 1961c: 1, 1952b: 1256. (tx) Blandford 1904: 260; Hagedorn 1910a: 145; Schedl 1933b: 33, 1935h: 346, 1950i: 158, 1979c: 214; Wood, S. L. 1961c: 1, 1952b: 1256.
- rufopilosus** Eggers 1931b: 39. Holotype ♀; Sao Paulo [Brazil]; Prague Museum.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 754. (tx) Eggers 1931b: 39–40; Schedl 1954n: 42, 1963d: 224, 1979c: 215.
- sanguineus** Schedl 1935h: 346. Holotype ♀; Costa Rica, [San Isidro de] Coronado; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
Hosts: *Persea americana*.
References: (ds) Blackwelder 1947: 754; Wood, S. L. 1952b: 1257. (tx) Schedl 1935h: 346–347, 1950i: 156, 1979c: 219; Wood, S. L. 1952b: 1257.
- schaufussi** Schedl 1937g: 69. Lectotype ♂; Brasilien, Nova Teutonia; Schedl Collection in NHMW, Wien.
Figures: Pedrosa-Macedo & Schonherr 1955: 66–67, Terra 1957: 29.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 754; Nunberg 1971: 57; Pedrosa-Macedo & Schonherr 1955: 67; Schedl 1966f: 91, 1967d: 4, 1970e: 85, 1972g: 47, 1973d: 160; Terra 1957: 27. (tx) Pedrosa-Macedo & Schonherr 1955: 66–67; Schedl 1937g: 69–70, 1939j: 564, 1950i: 147, 1979c: 221; Terra 1957: 27, 29.
- senticosus** Wood 1956c: 271. Holotype ♀; Jalapa, Veracruz, Mexico; Wood Collection.
Distribution: North America (Veracruz in Mexico).
Hosts: *Psittacanthus schiedeannus*.
References: (tx) Wood, S. L. 1956c: 271.
- sentosus** Wood 1956c: 271. Holotype ♀; Km 32 on Carretera Patzcuaro-Ario de Rosales, Michoacan, Mexico; Wood Collection.
Distribution: North America (Michoacan in Mexico).
Hosts: *Psittacanthus* sp.
References: (tx) Wood, S. L. 1956c: 271.
- sentus** Wood 1974c: 195. Holotype ♀; 2 km SE Cartago, Cartago, Costa Rica, 1300 m; Wood Collection.
Figures: Wood 1952b: 1301 (female).
Distribution: North America (Costa Rica/ Guatemala).
Hosts: *Miconia schlechtendalii*, also in a liana, shrub, and tree.
References: (hb) Wood, S. L. 1952b: 1303. (ds) Wood, S. L. 1952b: 1303. (tx) Wood, S. L. 1974c: 195, 1952b: 1303.
- serratus** Wood 1974c: 197. Tapanti, Cartago, Costa Rica, 1300 m; Wood Collection.
Figures: Wood 1952b: 1307 (female).
Distribution: North America (Costa Rica/ Panama).
Hosts: *Inga* sp., *Miconia globuliflora*, *Phoebe mexicana*, *Siparuna nicaraguensis*, *Werklia insignis*.
References: (hb) Wood, S. L. 1952b: 1306. (ds) Wood, S. L. 1952b: 1306. (tx) Wood, S. L. 1974c: 197, 1952b: 1306.
- serrulatus** Eggers 1934a: 52. Syntypes 1 ♂, 1 ♀; Bolivien (Cochabamba); ♂ MNHN, Paris. ♀ Eggers Collection, in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Blackwelder 1947: 754. (tx) Eggers 1934a: 52; Schedl 1979c: 225.
- simillimus** Schedl 1966f: 118. Holotype ♂; Venezuela; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
References: (tx) Schedl 1966f: 118, 1979c: 230.
- simplex** Wood 1974c: 157. Holotype ♀; Cerro de la Muerte, San Jose, Costa Rica, 3100 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Podocarpus oleifolius*, and other tree branches.
References: (ds) Wood, S. L. 1952b: 1256. (tx) Wood, S. L. 1974c: 157, 1952b: 1256.
- sobrinus** Wood 1974c: 196. Holotype ♀; Turrialba, Cartago, Costa Rica, 700 m; Wood Collection.
Distribution: North America (Costa Rica).
Hosts: *Theobroma cacao*.
References: (hb) Wood, S. L. 1952b: 1305. (ds) Wood, S. L. 1952b: 1305. (tx) Wood, S. L. 1974c: 196, 1952b: 1305.

- spinus Wood** 1974c: 194. Holotype ♂; Fortín de las Flores-Sumidero, Veracruz, Mexico, 900 m; Wood Collection.
Distribution: North America (Veracruz in Mexico).
References: (ds) Wood, S. L. 1982b: 1302. (tx) Wood, S. L. 1974c: 194, 1982b: 1302.
- strigilis Wood** 1974c: 189. Holotype ♀; 16 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.
Distribution: North America (Costa Rica/ Panama).
Hosts: *Cecropia* sp., *Miconia globuliflora*, *Myrica pubescens*, *Ochroma* sp., *Phoebe mexicana*, *Werkliia insignis*.
References: (hb) Wood, S. L. 1982b: 1293. (ds) Wood, S. L. 1982b: 1293. (tx) Wood, S. L. 1974c: 189, 1982b: 1293.
- subasperulus Eggers** 1940a: 141. Lectotype ♀; Guadeloupe; USNM, Washington, designated by Anderson & Anderson 1971: 32.
Distribution: Antilles Islands (Dominica/ Guadeloupe).
References: (tx) Anderson, W. H. & Anderson 1971: 32; Bright 1981c: 153; Eggers 1940a: 141; Schedl 1979c: 240.
- suberratus Wood** 1974c: 197. Holotype ♀; Volcán Poás, Heredia, Costa Rica; Wood Collection.
Figures: Wood 1982b: 1307 (female).
Distribution: North America (Costa Rica).
Hosts: Tree branch, *Vaccinium consanguineum*.
References: (hb) Wood, S. L. 1982b: 1306. (ds) Wood, S. L. 1982b: 1306. (tx) Wood, S. L. 1974c: 197, 1982b: 1306.
- subsulcatus Schedl** 1961i: 230. Holotype ♀; Bolivia, Cochabamba, Prov. Chapara, km 150, Jungas del Palmar; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Schedl 1960a: 77, 1961i: 230.
- suturalis Eggers** 1931b: 41. Lectotype ♂; Brasilien (Espírito Santo); USNM, Washington, designated by Anderson & Anderson 1971: 33.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 784; Pedrosa-Macedo & Schonherr 1985: 68; Schedl 1967d: 4, 1972g: 47, 1973d: 160. (tx) Anderson, W. H. & Anderson 1971: 33; Eggers 1931b: 41–42; Pedrosa-Macedo & Schonherr 1985: 68; Schedl 1950i: 155, 1979c: 247.
- suturifer Schedl** 1963c: 165. Holotype ♀; Córdoba, Veracruz, Mexico; Schedl Collection in NHMW, Wien.
Distribution: North America (Veracruz in Mexico).
References: (ds) Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 1296. (tx) Schedl 1963c: 165, 1979c: 248; Wood, S. L. 1982b: 1296.
- theobromae Nunberg** 1971: 59. Holotype ♂; Ilheus, Bahia, Brasilien; MZUSP, São Paulo.
Figures: Nunberg 1971: 60, Terra 1987: 27, 29.
Distribution: South America (Brazil).
Hosts: *Theobroma cacao*.
References: (ds) Nunberg 1971: 59; Terra 1987: 27. (tx) Nunberg 1971: 59–60; Terra 1987: 27, 29.
- transversus Eichhoff** 1869a: 279. Holotype, sex?; Nov. Granada; IRSNB, Brussels.
Distribution: South America (Colombia).
References: (ds) Blackwelder 1947: 784; Blandford 1904: 254; Gemminger & Harold 1872: 2693; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 342. (tx) Blandford 1904: 254; Eichhoff 1869a: 279, 1869c: 301, 1878b: 426; Hagedorn 1910a: 145.
- trucis Wood** 1974c: 193. Holotype ♀; 16 km SE Cartago, Cartago, Costa Rica, 1800 m; Wood Collection.
Figures: Wood 1982b: 1301 (female).
Distribution: North America (Costa Rica).
Hosts: *Miconia globuliflora*.
References: (hb) Atkinson et al. 1986: 46; Burgos & Sancedo 1983: 140. (ds) Atkinson et al. 1986: 46; Burgos & Sancedo 1983: 140; Wood, S. L. 1982b: 1299. (tx) Wood, S. L. 1974c: 193, 1982b: 1299.
- truncatus Wood** 1985: 272. Holotype ♀; jungle near Leonpampa, Huanuco Department, Peru, 800 m; Wood Collection.
Distribution: South America (Peru).
References: (tx) Wood, S. L. 1985: 272.
- trunculus Wood** 1974c: 191. Holotype ♀; 13 km S El Hato del Volcán, Chiriquí, Panamá; Wood Collection.
Distribution: North America (Costa Rica/ Panama).
Hosts: Small tree, and a liana.
References: (hb) Wood, S. L. 1982b: 1296. (ds) Wood, S. L. 1982b: 1296. (tx) Wood, S. L. 1974c: 191, 1982b: 1296.
- tuberculatus Eggers** 1940a: 140. Lectotype ♂; Guadeloupe; USNM, Washington, designated by Anderson & Anderson 1971: 34.
Distribution: Antilles Islands (Dominica/ Guadeloupe).
References: (ds) Bright 1981c: 153, 1985c: 176. (tx) Anderson, W. H. & Anderson 1971: 34; Bright 1985c: 176; Eggers 1940a: 140, 1941a: 108; Schedl 1979c: 257.
- tulcanus Hagedorn** 1910b: 6. Holotype, sex?; Ecuador - Tulcan; MNHN, Paris.
Distribution: South America (Ecuador).
References: (ds) Blackwelder 1947: 784; Hagedorn 1910d: 91; Kleine 1913b: 155, 1914b: 343. (tx) Hagedorn 1910a: 145, 1910b: 6.
- uniseptis Schedl** 1961i: 229. Holotype ♂; Mexico, Córdoba, Ver.; CAS, San Francisco.
Figures: Bright 1972a: 1370.
Distribution: North America (Chiapas, Veracruz in Mexico).
References: (ds) Atkinson & Equihua 1985c: 362,

- 1986a: 422; Schedl 1960a: 75; Wood, S. L. 1982b: 1299. **(tx)** Schedl 1960a: 75, 1961i: 229, 1979c: 260; Wood, S. L. 1974d: 280, 1982b: 1299.
- reburus* Bright 1972a: 1375. Holotype ♀; Palenque Ruins, Chiapas, Mexico; CNCI, Ottawa. Synonymy: Wood 1974d: 280. References: **(tx)** Bright 1972a: 1370, 1375; McNamara 1977: 195; Wood, S. L. 1974d: 280.
- venustus* Schedl 1951m: 127 (*Thylurcos*). Holotype ♂; Brasilien; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: **(tx)** Schedl 1951m: 127, 1979c: 265.
- villifer* Wood 1974c: 183. Holotype ♀; 15 km SE Cartago, Costa Rica, 1800 m; Wood Collection. Distribution: North America (Costa Rica/ Panama). Hosts: *Siparuna nicaraguensis*, and a liana. References: **(hb)** Wood, S. L. 1982b: 1281. **(ds)** Wood, S. L. 1982b: 1281. **(tx)** Wood, S. L. 1974c: 183, 1982b: 1281.
- villosus* Eggers 1943a: 381. Holotype ♂; Bolivia (Cochabamba); Eggers Collection, in NHMW, Wien. Distribution: South America (Bolivia/ Brazil). Notes: (3) Schedl 1976a: 81 (described female). References: **(ds)** Schedl 1972g: 47, 1973d: 160.
- (tx)** Eggers 1943a: 381; Schedl 1976a: 81, 1979c: 268.
- villus* Bright 1972a: 1371. Holotype ♀; Yerba Buena, 32 km N Bochil, Chiapas, Mexico; CNCI, Ottawa. Figures: Bright 1972a: 1370, Wood 1982b: 1280 (female). Distribution: North America (Costa Rica/ Guatemala/ Chiapas, Puebla in Mexico). Hosts: *Ahuus* sp., *Spondias purpurea*, and other tree branches. References: **(hb)** Wood, S. L. 1982b: 1281. **(ds)** Atkinson & Equihua 1986a: 422; Wood, S. L. 1982b: 1281. **(tx)** Bright 1972a: 1370–1371; McNamara 1977: 195; Wood, S. L. 1982b: 1281.
- zelus* Wood 1974c: 190. Holotype ♀; Cerro de la Muerte, San Jose, Costa Rica; Wood Collection. Figures: Wood 1982b: 1294 (female). Distribution: North America (Costa Rica). Hosts: *Brunellia costaricensis*. References: **(hb)** Wood, S. L. 1982b: 1293. **(ds)** Wood, S. L. 1982b: 1293. **(tx)** McNamara 1977: 195; Wood, S. L. 1974c: 190, 1982b: 1293.

Genera Incertae Sedis

Genus *Bufonius* Eggers

BUFONUS EGGERS 1919: 231. Type-species: *Bufonius obscurus* Eggers, monobasic.

Notes: (1) The holotype and only known specimen of the type-species is lost. The description suggests that is a Hypoborini.

References: (tx) Eggers 1919: 231; Schedl 1961k: 444.

***obscurus* Eggers** 1919: 231. Holotype, sex?; Amani, Ost-Afrika; Hamburg Museum, lost.

Distribution: Africa (Tanzania).

References: (tx) Eggers 1919: 231; Schedl 1961k: 444.

Genus *Carphoborites* Schedl

CARPHOBORITES SCHEDL 1947a: 32. Type-species: *Carphoborites keilbachi* Schedl, present designation.

Notes: (1) This genus is doubtfully distinct from *Carphoborus*. The lapsus calami, *Charphoborites*, was corrected by Schedl on p. 33.

References: (tx) Schedl 1947a: 32.

***keilbachi* Schedl** 1947a: 32. Holotype, sex?; Baltische Bernstein; Geologisch-Palaeontologische Institut Albertus Universitat, Konigsberg.

Distribution: Europe (fossil in Baltic amber).

References: (tx) Schedl 1947a: 32.

***posticus* Schedl** 1947a: 33. Holotype, sex?; Baltische Bernstein; Geologisch-Palaeontologische Institut Albertus Universitat, Konigsberg.

Distribution: Europe (fossil in Baltic amber).

References: (tx) Schedl 1947a: 33.

Genus *Cryphalites* Cockerell

CRYPHALITES COCKERELL 1917: 368. Type-species: *Cryphalites rugosissimus* Cockerell, monobasic.

References: (tx) Cockerell 1917: 368.

***rugosissimus* Cockerell** 1917: 368. Holotype, sex?; Burmese amber; University of Colorado, Boulder [not now present in that collection].

Distribution: Asia (fossil in Burmese amber).

Notes: (1) This specimen was not examined by us, and is unknown to us except for the very brief description. It is tentatively assigned to the Cryphalini.

References: (ds) Keilbach 1985: 255. (tx) Cockerell 1917: 368. (ms) Keilbach 1985: 255.

Genus *Taphramites* Schedl

TAPHRAMITES SCHEDL 1947a: 41. Type-species: *Taphramites gnathotrichus* Schedl, monobasic.

References: (tx) Schedl 1947a: 41–43.

***gnathotrichus* Schedl** 1947a: 42. Holotype, sex?; Baltische Bernstein; Geologisch-Palaeontologische Institut Albertus Universitat, Konigsberg.

Distribution: Europe (fossil in Baltic amber).

Notes: (1) This species probably belongs in or near *Dryocoetes* in the Dryocoetini.

References: (ds) Keilbach 1982: 255. (tx) Schedl 1947a: 42, 1970c: 68–70. (ms) Keilbach 1982: 255.

Genus *Toxophthorus* Wood

TOXOPHTHORUS WOOD 1962: 77. Type-species: *Toxophthorus africanus* Eggers, automatic.

Toxophthorus Eggers 1920: 119. Type-species: *Toxophthorus africanus* Eggers, preoccupied by Schoenherr 1836.

References: (tx) Eggers 1920: 119–120; Schedl 1957d: 14, 1961k: 735; Wood, S. L. 1962: 77.

References: (tx) Wood, S. L. 1962: 77.

***africanus* (Eggers)** 1920: 119 (*Toxophthorus*). Holotype, sex?; Deutsch-Ostafrika; Hamburg Museum, lost.

Distribution: Africa (Tanzania).

Hosts: Copal.

Notes: (1) This species is apparently a Dryocoetini in or near *Tiarophorus* or *Xylocleptes*. Although it was named as a fossil, it is probably a living species.

References: (tx) Eggers 1920: 119; Schedl 1957d: 14, 1961k: 735.

Genus *Xyleborites* Wickham

XYLEBORITES WICKHAM 1913: 26. Type-species: *Xyleborites longipennis* Wickham, monobasic.

References: (tx) Leng 1920: 363; Wickham 1913: 26.

***longipennis* Wickham** 1913: 26. Holotype, sex?; Miocene (Colorado) [USA]; not located.

Distribution: North America (Miocene fossil from Colorado in USA).

Notes: (1) Essential characters were not described. It is uncertain where this species fits in classification or even if it is actually a scolytid.

References: (tx) Leng 1920: 363; Wickham 1913: 26.

Genus *Xylechinites* Hagedorn

XYLECHINITES HAGEDORN 1906: 120. Type-species: *Xylechinites anceps* Hagedorn, monobasic.

References: (tx) Hagedorn 1906: 120; Schedl 1947a: 30–31.

***anceps* Hagedorn** 1906: 120. Holotype, sex?; Baltische Bernstein; Geologisch-Palaeontologische Institut Albertus Universitat, Konigsberg.

Figures: Schedl 1947a: 31 (photograph of type).

Distribution: Europe (fossil in Baltic amber).

Notes: (1) This species appears to be a Tomicini, but it may not be allied to *Xylechinus*. (3) Schedl 1947a: 31 (redescribed).

References: (ds) Keilbach 1982: 255. (tx) Hagedorn 1906: 120; Keler 1928: 28; Schedl 1947a: 31. (ms) Keilbach 1982: 255.

Genus *Cynanchophagus* Axentjev

CYNANCHOPHAGUS AXENTJEV 1987:000. Type-species: *Cynanchophagus cornutus* Axentjev, monobasic.

References: (tx) Axentjev 1987:000.

cornutus Axentjev 1987:000. Holotype ♀;
Tadzhikistan: lowland of river Vakhsh, nature
reserve "Tigrovia balka"; IZL, St. Petersburg/Len-
ingrad.
Distribution: Asia (Tadzhikistan in USSR).

Hosts: *Cynanchum sibiricum*.

Notes: (1) We have not seen the original descrip-
tion or specimens of this species.

References: (tx) Axentjev 1987:000.

Catalog of Platypodidae

Family Platypodidae Erichson

Platypodidae (including the French Platypides)

Notes: In the absence of a comprehensive review of genera for the Platypodidae since the family was named, it was intended that such a study would be completed prior to completion of this catalog. Immovable publication deadlines made that impossible. However, after the catalog was completed but before it was published the review of genera was prepared and will be published in the 1993 *Great Basin Naturalist* (SLW).

References: Balachowsky 1949a: 275; Barbey 1901: 28; Beal & Massey 1945: 62; Bedel 1888b: 385, 404, 421; Blandford 1895b: 89; Broun 1850: 539; Brues & Melander 1932: 445; Chapuis 1865; Costa Lima 1956: 339; Erichson 1847: 138; Handlirsch 1925: 692; Hopkins 1915c: 225; Jacquelin du Val & Fairmaire 1868: 109; Lacordaire 1866: 355; LeConte 1873: 13, 1876: 341; LeConte & Horn 1883: 513; Leng 1920: 337; Lindemann 1876: 151; Lovendal 1889b: 5; Lucas 1920: 50; Nunberg 1954: 13, 92; Reitter 1894: 39, 93, 1913a: 13; Schedl 1939b: 377, 388, 396, 1972f; Strohmeyer 1912c: 1–26; Swaine 1909: 83; Thomson 1865: 377; Tredl 1907: 4.

Platypodinae

References: Blandford 1895b: 89; Strohmeyer 1911: 218, 1912: 4; Swaine 1909: 83.

Platypodini

References: Schedl 1939b: 396.

Platypites

References: Fairmaire 1868: 107.

Platypodae

References: Eichhoff 1881: 305.

Platytarsulidae

References: Schedl 1939b: 381, 387.

Platytarsulinae

References: Schedl 1972f: 83, 269.

Subfamily Coptonotinae Chapuis

Coptonotidae

References: Balachowsky 1949a: 51; Blackman 1944: 77–78; Blackwelder 1947: 788; Chapuis 1869: 11, 1873: 219; Nunberg 1953: 43, 46; Schedl 1939b: 381, 386–387, 1940a: 318, 328, 1962: 3–14.

Coptonotinae

References: Hopkins 1915c: 225, 227; Schedl 1939b: 381, 387.

Chapuisidae

References: Blackman 1944: 77–78; Strohmeyer 1914b: 1–4.

Tribe Coptonotini Chapuis

Coptonotidae

References: Balachowsky 1949a: 51; Blackman 1944: 77–78; Blackwelder 1947: 788; Chapuis 1869: 11, 1873: 219; Nunberg 1953: 43, 46; Schedl 1939b: 381, 386–387, 1940a: 318, 328, 1962d: 314.

Genus *Coptonotus* Chapuis

COPTONOTUS CHAPUIS 1869: 11. Type-species: *Coptonotus cyclopus* Chapuis, monobasic.

References: (ds) Blackwelder 1947: 788; Hagedorn 1910d: 7, (tx) Blackman 1944: 77–78; Blandford 1893d: 427, 441; Chapuis 1869: 11, 1873: 219; Eggers 1935a: 75; Hagedorn 1910a: 34–37; Hopkins 1914: 118, 133; Schaufuss 1905: 89; Schedl 1939b: 386, 1948b: 262, 1962d: 8.

cyclopus Chapuis 1869: 11. Syntypes, sex?; Nouvelle-Grenade; IRSNB, Brussels.

Figures: Equihua & Atkinson 1987: 11, Schedl 1962d: fig. 3.

Distribution: North America (Costa Rica/ Panama), South America (Colombia/ Peru).

Hosts: Probably *Quercus* sp.

References: (ds) Blackwelder 1947: 788; Equihua & Atkinson 1987: 10; Hagedorn 1910d: 7; Kleine 1912b: 162, 1914b: 341, (tx) Blackman 1944: 75; Chapuis 1869: 11, 1873: 219; Equihua & Atkinson 1987: 10–11; Gemminger & Harold 1872: 2678; Hagedorn 1910a: 37; Hopkins 1914: 118, 133; Lucas 1920: 202; Schaufuss 1905: 89; Schedl 1939b: 379, 384, 1962d: 8, 1962q: 485, (ms) Lucas 1920: 202.

striatus Eggers 1933b: 3. Holotype ♀; Franz, Guayana; USNM, Washington.

Distribution: South America (Cayenne/ Peru/ Suriname).

Notes: (1) Schedl 1979c: 237 (holotype designation invalid).

References: (ds) Blackwelder 1947: 788, (tx) Eggers 1933b: 3–4; Schedl 1962d: 8, 1979c: 237.

Genus *Protohyllastes* Wood

PROTOHYLLASTES WOOD 1973d: 83. Type-species: *Protohyllastes annosus* Wood, original designation.

References: (tx) Wood, S. L. 1973d: 83.

annosus Wood 1973d: 84. Holotype ♂; Eungella National Park, Queensland, Australia; Queensland Museum, Brisbane.

Distribution: Australia (Queensland).

References: (tx) Wood, S. L. 1973d: 84.

latirostris (Pascoc) 1873: 197 (*Psepholax*). Holotype ♀; Klawara, N.S. Wales; BMNH, London.

Distribution: Australia (New South Wales).

Notes: (1) Scale color and body shape of these two species are obviously different; however, they

could represent different sexes of the same species. Both holotypes were studied.

References: (tx) Pascoe 1873: 197.

Genus *Scolytotarsus* Schedl

SCOLYTOTARSUS SCHEDL 1937: 404. Type-species: *Scolytotarsus impar* Schedl, monobasic.

References: (tx) Schedl 1937b: 404, 1939b: 387.

impar Schedl 1937b: 405. Holotype, sex?; Wahrscheinlich Kamerun, aus einem Ankauf von Banghaas, Dresden; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon).

References: (tx) Schedl 1936g: 534, 1937b: 405.

maculatus Schedl 1936g: 534. Syntypes, sex?; Queensland: Coen district, Cape York; SAM, Adelaide, and NHMW, Wien.

Distribution: Australia (Cape York in Queensland).

Notes: (3) Voss 196.: 282 (treated in Curculionidae under *Mcypsepholax*).

References: (ds) Brimblecombe 1953: 24; Schedl 1936g: 535. (tx) Schedl 1936g: 534, 1939b: 379–380, 384; Voss 196.: 282.

Tribe Mecopelmini Wood

Mecopelmini

References: Wood, S. L. 1966a: 45.

Genus *Mecopelmus* Blackman

MECOPELMUS BLACKMAN 1944: 78. Type-species: *Mecopelmus zeteki* Blackman, original designation.

References: (tx) Blackman 1944: 78; Schedl 1962d: 12; Wood, S. L. 1966a: 45.

zeteki Blackman 1944: 78. Holotype, sex?: Barro Colorado Island, Panama Canal Zone; USNM, Washington.

Figures: Blackman 1944:figs. 1-5, Equihua & Atkinson 1987: 12, Schedl 1962d:fig. 2.

Distribution: North America (Canal Zone in Panama).

Hosts: *Serjania* sp.

Notes: (3) The species is phloeophagous, monogynous; the larvae make individual mines.

References: (hb) Wood, S. L. 1966a: 45. (ds)

Equihua & Atkinson 1987: 12. (tx) Blackman 1944: 76-80; Equihua & Atkinson 1987: 12; Schedl 1962d: 12; Wood, S. L. 1966a: 45, 1987: 12.

Genus *Protoplatypus* Wood

PROTOPLATYPUS WOOD 1973b: 81. Type-species: *Protoplatypus vetulus* Wood, original designation.

References: (tx) Wood, S. L. 1973b: 81.

vetulus Wood 1973b: 82. Holotype ♂; 8 km NW Bulolo, Morobe, New Guinea; CSIRO, Canberra. Figures: Wood 1973b: 82.

Distribution: New Guinea (near Bulolo in Morobe District).

Hosts: Log (an uncommon species, possibly *Harpullia pedicellaris*).

Notes: (3) The species is phloeophagous, polygynous, and the larvae make individual mines.

References: (hb) Wood, S. L. 1973b: 82. (tx) McNamara 1977: 191; Wood, S. L. 1973b: 82.

Tribe Schedlarini Wood

Schedlarini (automatic replacement)

References: Wood, S. L. 1957b: 103.

Chapnisiidae (automatically replaced by new name when type-genus became a homonym)

References: Brues & Melander 1932: 445; Schedl 1939b: 377; Strohmeier 1914b: 1–6.

Chapnisiides

References: Blandford 1895b: 89, 117.

Chapnisiinae

References: Brues & Melander 1932: 445; Handlirsch 1925: 692; Hopkins 1915c: 225; Schedl 1939b: 377; Strohmeier 1911: 218, 1912c: 3.

Genus *Schedlarius* Wood

SCHEDLARIUS WOOD 1957b: 103. Type-species: *Chapnisia mexicana* Duges, automatic.

Chapnisia Duges 1886: 56. Type-species: *Chapnisia mexicana* Duges, monobasic, preoccupied by Duvivier 1885.

References: (hb) Duges 1885: 58; Wood, S. L. 1957b: 103–104. (tx) Blackman 1944: 76–79; Blandford 1893d: 427, 1895a: 317–318, 1895b: 81, 88, 1896e: 117; Duges 1885: 58, 1888: 141; Hopkins 1914: 118, 134, 1915c: 227; Nunberg 1953: 44, 46; Schedl 1939b: 387, 1962d: 9; Strohmeier 1914b: 1–5, 1914c: 16–17; Wood, S. L. 1957b: 103–104.

References: (tx) Schedl 1962d: 9–11; Wood, S. L. 1957b: 103–104.

mexicanus Duges 1885: 58 (*Chapnisia*). Syntypes, sex?; Mexico; presumably in Duges Collection, Mexico City.

Figures: Blandford 1896e: pl. 5, Burgos & Saucedo 1983: 43, Duges 1885: pl. 5, Equihua & Atkinson 1987: 13, Schedl 1962d: fig. 3.

Distribution: North America (Mexico).

Hosts: *Bursera capallifera*, *B. simarubra*.

Notes: (3) This species is xylophagous, monogynous, and the larvae make long, tortuous, individual tunnels in the xylem; ambrosial fungus is not present although obvious fungal decay of host xylem occurs near larval mines.

References: (hb) Atkinson et al. 1986: 47; Burgos & Saucedo 1983: 42; Duges 1885; Wood, S. L. 1957b: 103. (ds) Atkinson et al. 1986: 47; Blackwelder 1947: 788; Blandford 1896e; Burgos & Saucedo 1983: 42; Equihua 1985: 144; Equihua & Atkinson 1987: 13; Strohmeier 1912c: 3. (tx) Blackman 1944: 77; Blandford 1896e: 117; Burgos & Saucedo 1983: 43; Crowson 1955; Duges 1885: 58, 1888: 141; Equihua & Atkinson 1987: 13; Hopkins 1914: 118, 134; Schedl 1939b: 379–380, 1940a: 328, 1962d: 9; Strohmeier 1911d: 218, 1912c: 3, 1914b: 1–5, 1914c: 16; Wood, S. L. 1957b: 103, 1959b.

Subfamily Tesserocerinae Blandford

Tesserocerini gemmini

References: Blandford 1895b: 115.

Tesserocerini

References: Schedl 1939b: 396, 400.

Tesserocerinae

References: Strohmeier 1914b: 18, 1918: 40, 1920: 44.

Tesserocerariae

References: Strohmeier 1914b: 18–19.

Symmerariae

References: Strohmeier 1914b: 18.

Periommatinae

References: Schedl 1939b: 397, 403, 1972f: 83, 249.

Cenocephalariae

References: Strohmeier 1914b: 18.

Cenocephalini

References: Schedl 1939b: 397.

Platypicerinae

References: Nunberg 1953: 44.

Tribe Diapodini Strohmeier

Diapodariae

References: Strohmeier 1914b: 18.

Diaporinae

References: Schedl 1939b: 397, 404, 1972f: 83, 259.

Genyocerinae

References: Hopkins 1915c: 225.

Genus *Diapus* Chapuis

DIAPUS CHAPUIS 1865: 43, 329. Type-species: *Diapus quadrispinatus* Chapuis, subsequent designation by Hopkins 1914: 121.

Notes: (3) Beeson 1941: 260 (*himalayensis*, nomen nudum, *rostratus*, nomen nudum).

References: (ay) Gardner 1932: 1–9; Nobuchi 1969a: 69. (cn) Beeson 1918: 114–124, 1922: 1–2; Fisher, Thompson, & Webb 1953: 1–21; Garthwaite 1940: 94–106. (hb) Beeson 1910: 222, 1935b: 182–186; Browne 1936b: 1, 1961c: 230; Kalshoven 1960: 31–34, 45; Schedl 1972f: 261. (ds) Beeson 1923: 173, 1941: 335; Munro 1928: 1–29; Schedl 1966b: 102, 1972f: 261, 1977b: 276. (tx) Beeson 1917: 1–29; Blandford 1893d: 426, 1894d: 139; Browne 1955b: 164–182, 1972a: 188; Chapuis 1865: 329, 332; Hopkins 1914: 121, 135, 1915c: 227; Lacordaire 1866: 389, 397–398; Murayama 1925: 216–217, 1934f: 148, 1954: 206; Nobuchi 1973b: 17; Nunberg 1951: 262, 1953: 144; Sampson 1913: 449, 1923: 449; Schedl 1936e: 61, 1937d: 44, 1939b: 391–393, 404, 408, 1957b: 159, 1959a: 516, 1962k: 1052–1053, 1968e: 261–270, 1972f: 259–264, 1977b: 276; Stebbing 1907a: 42, 1914: 613, 628–634; Stroh-

meyer 1911c: 105–107, 1912c: 22, 1914c: 6–19, 46–47, 1918: 1–46, 1942: 286–287; Wood, S. L. 1960a: 3, 6.

aculeatus Blandford 1894d: 139. Syntypes 2 ♂; Higo, Japan; BMNH, London.

Figures: Nobuchi 1973b: pl. 2, figs. 38–39.

Distribution: Asia (Assam, Uttar Pradesh in India/ Japan/Vietnam), Indonesia (Java).

Hosts: *Ahhus nitida*, *Castanea javanica*, *C. tunggurut*, *Quercus dilatata*, *Q. incana*, *Q. semicarpifolia*.

References: (cn) Kleine 1932a: 310, 402. (ec) Beeson 1921b: 22. (hb) Kalshoven 1960c: 46, 1960d: 46; Kleine 1932a: 310, 402; Murayama 1931b: 201. (ds) Beeson 1921b: 22; Blandford 1894c; Kalshoven 1960d: 46; Kleine 1932a: 310, 402; Murayama 1929a: 679, 1931b: 201–203, 1934f: 143, 1936b: 141, 1937c: 582, 1949c: 104, 1950b: 1300, 1953c: 165, 1955: 100–101; Nobuchi 1967: 27, 1973b: 17; Schedl 1962f: 169, 1972f: 263, 1974c: 262, 1975c: 384; Strohmeier 1912c: 22, 1914c: 47. (tx) Blandford 1894d: 139; Murayama 1925b: 234, 1950b: 1300, 1955: 100–101; Nobuchi 1973b: 17, pl. 2; Schedl 1934f: 1647, 1969b: 223, 1972f: 263; Strohmeier 1912c: 22, 1914c: 47.

aculeatus javanus Murayama 1931b: 200. Holotype, sex?; Mt. Gede, 600 m, Java; Murayama Collection in USNM, Washington.

References: (ds) Schedl 1972f: 263. (tx) Murayama 1931b: 200; Schedl 1962f: 263.

amblylaminatus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo: Oomsis L.A., 900 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Canarium* sp., *Castanopsis* sp., *Litsea* sp., *Myristica* sp.

References: (tx) Roberts 1992:(in press).

angustidontus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Gumi L.A., Bulolo, 2000 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Garcinia* spp.

References: (tx) Roberts 1992:(in press).

bilunatus Schedl 1975f: 399. Holotype ♀; New Guinea NE, Mt. Missim, 2400–2800 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Acmena* sp., *Castanopsis* sp., *Elacocarpus* sp., *Litsea* sp., *Macaranga* sp., *Phyllocladus* sp., *Sericolea* sp., *Tenstromia* sp.

References: (tx) Roberts 1992:(in press); Schedl 1975f: 399.

bispinus Schedl 1974d: 471. Holotype ♂; Panganda Logging Area, 2126 m, Watut Valley, Morobe District, New Guinea; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Calophyllum* sp., *Castanopsis acuminatissima*, *Garcinia* sp., *Xanthophyllum* sp.

- Notes: (3) A note in the Schedl Collection says that *latespinis* is a synonym [requires verification].
References: (tx) Schedl 1974d: 471, 1975a: 16.
- borneensis** Browne 1980d: 500. Holotype ♂; Kimanis (Borneo) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Borneo).
References: (tx) Browne 1980d: 500.
- brownei** Schedl 1971c: 398. Holotype ♂; Sarawak, Semengoh; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
References: (ds) Schedl 1972f: 263. (tx) Schedl 1971c: 398–399, 1972f: 263.
- curvidens** Schedl 1965g: 30. Holotype ♂; O. Borneo, Mt. Tibang, 1300 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 263. (tx) Schedl 1965g: 30, 1972f: 263, 1978a: 23.
- elongatus** Schedl 1969b: 225. Holotype ♂; New Guinea: Marafunga, 2800 m, E.H.D.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Cryptocarya* sp., *Dryodaphnae* sp., *Elacocarpus* sp., *Galbulilima* sp., *Phyllocladus* sp., *Planchonella* sp., *Podocarpus* sp., *Spherostenon* sp.
Notes: (3) Schedl 1969b: 225 (in key).
References: (ds) Schedl 1972f: 263. (tx) Roberts 1992:(in press); Schedl 1969b: 225, 1972f: 263, 1978a: 27.
- formosanus** Niisima & Murayama 1925a: 217. Syntypes 4♂, 4♀; Rengeti: Taichu Prov. [Taiwan]; Murayama Collection in USNM, Washington.
Figures: Murayama 1925a: pl. 12, figs. 7–8, pl. 15, figs. 35–47.
Distribution: Asia (Taiwan).
Hosts: *Lithocarpus konishii*, *Psidium guajava*.
References: (hb) Murayama 1931b: 202. (ds) Miwa 1931: 268; Murayama 1925a: 217, 1929a: 679, 1931b: 202, 1937c: 583; Nobuchi 1967: 27; Schedl 1972f: 263. (tx) Murayama 1925a: 217–223, 1925b: 235; Schedl 1972f: 263.
- gestroi** Sampson 1923c: 74. Holotype ♀; Burma: Carin Cheba, 900–1100 m; MCG, Genova.
Distribution: Asia (Burma).
References: (tx) Sampson 1923c: 74–75; Schedl 1939e: 364, 1972f: 263.
- kuperi** Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo: Kuper Range, 2200 m; BPBM, Honolulu.
Distribution: New Guinea.
Hosts: *Nothofagus grandis*, *Syzygium* sp.
References: (tx) Roberts 1992:(in press).
- latespinis** Schedl 1979g: 120. Holotype ♂; Papua, Upper Manki L.A., Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Hosts: *Castanopsis* sp., *Lithocarpus* sp., *Myristica* sp.
Notes: (3) See note under *bispinus* above.
- References: (tx) Roberts 1992:(in press); Schedl 1979g: 120.
- malgassicus** Schedl 1970d: 240. Holotype ♂; Madagascar, Perinet; MNHN, Paris.
Distribution: Madagascar.
Hosts: *Eucalyptus* sp., *Pinus insularis*, *Symphonia* sp.
References: (ds) Schedl 1972f: 263, 1977b: 277. (tx) Schedl 1970d: 240–241, 1972f: 263, 1977b: 277, 1978a: 42.
- minor** Schedl 1954a: 158. Holotype ♂; Java, Mt. Gede, 800 m; Schedl Collection in NHMW, Wien.
Distribution: Asia (Bengal in India), Indonesia (Borneo, Java).
Hosts: *Castanea javanica*, *Castanopsis hystrix*, *Cinnamomum camphora*, *Lindera pulcherrima*, *Quercus lamellosa*.
Notes: (3) Beeson 1961: 260 (*minor*, nomen nudum, synonymy in Browne 1977a: 101).
References: (ds) Beeson 1941 (1961: 260); Bhasin, Roonwal, & Singh 1958; Browne 1977a: 101, 1986c: 662; Kalshoven 1960c: 46; Schedl 1965g: 25, 1971c: 71, 1972f: 263. (tx) Beeson 1941 (1961: 260); Browne 1977a: 101; Schedl 1954a: 145, 158, 1961c: 75, 1969a: 219, 1972f: 263, 1978a: 44.
- minutissimus** Schedl 1969a: 219. Holotype ♂; Sumatra, Singij to Tokyo (Japan), imported; PPST, Tokyo.
Distribution: Asia (Malaya), Indonesia (Borneo, Sumatra).
Hosts: *Calophyllum tetrapterum*.
References: (hb) Beaver & Browne 1978: 617. (ds) Beaver & Browne 1978: 617; Browne 1980c: 484; Nobuchi 1977: 146; Schedl 1971c: 370, 1972c: 263. (tx) Schedl 1962f: 263, 1969a: 219, 1978a: 44.
- molossus** Chapuis 1865: 333. Holotype ♀; Inde orientale, dans la contree d'Almora au Nord-ouest des possessions anglaises; BMNH, London.
Distribution: Asia (Uttar Pradesh in India/ Malaya).
Hosts: *Alnus nepalensis*, *Quercus dilatata*, *Q. lamellosa*, *Q. semicarpifolia*, *Schleichera oleosa*.
References: (ds) Gemminger & Harold 1872: 2702; Lacordaire 1866: 397; Schedl 1962f: 169, 1972f: 263; Strohmeier 1912c: 22, 1914c: 47. (tx) Browne 1977a: 102; Chapuis 1865: 333; Janson 1893: 75; Lacordaire 1866: 397; Schedl 1960c: 63, 1962f: 169, 1972f: 263; Strohmeier 1912c: 22, 1914c: 47.
- impressus* Janson 1893: 74. Syntypes ♂ ♀; Deoband, NW Prov., India; BMNH, London.
Synonymy: Schedl 1972f: 264.
References: (cn) Kleine 1932a: 311, 402; Mathur & Singh 1961a: 29; Pierce, W. D. 1917: 154; Roonwal 1954: 58; Stebbing 1914: 628. (ec) Beeson 1921b; Stebbing 1914: 628. (hb) Beeson 1917: 10; Cotes 1893a; Kleine 1932a: 311, 402; Stebbing 1899: 62, 1906: 414, 1908a: 112, 1914: 328. (ds) Beeson 1916: 23,

- 1916a; 222. 1921b: 21–25, 1941 (1961: 260); Cotes 1893a; Kleine 1932a: 311, 402; Mathur & Singh 1961a: 29; Pierce, W. D. 1917: 154; Roonwal 1954: 58; Strohmeier 1912c: 22, 1914c: 47. **(tx)** Janson 1893: 74; Murayama 1925: 219; Schedl 1972f: 263–264; Stebbing 1908b: 1, 1914: 628; Strohmeier 1912c: 22, 1914c: 47.
- capillatus* Sampson 1913: 449. Holotype ♀; Darjeeling, N. India; BMNH, London.
Notes: (1) Browne 1977a: 102 (treated as a subspecies).
References: **(en)** Kleine 1932a: 310, 402; Pierce, W. D. 1917: 154; Stebbing 1914: 633. **(ec)** Beeson 1921b: 23; Stebbing 1914: 633. **(hb)** Kleine 1932a: 310, 402; Stebbing 1914: 633. **(ds)** Beeson 1921b: 23; Kleine 1932a: 310, 402; Pierce, W. D. 1917: 154; Schedl 1972f: 263; Strohmeier 1914c: 47. **(tx)** Browne 1977a: 102; Sampson 1913: 449–450; Schedl 1939e: 364, 1972f: 263; Stebbing 1914: 633; Strohmeier 1914c: 47.
- nanodontus* Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Oomsis L.A., 800 m; BPBM, Honolulu.
Distribution: New Guinea.
Hosts: *Castanopsis* sp., *Cryptocarya* sp., *Sterculia* sp.
References: **(tx)** Roberts 1992:(in press).
- namus* Schedl 1969b: 225. Holotype ♀; New Guinea: Erave pine forest, S.H.D.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Anisoptera* sp., *Araucaria cunninghamii*, *Calophyllum* sp., *Castanopsis acuminatissima*, *C.* sp., *Cryptocarya* sp., *Elacocarpus* sp., *Haplolobus* sp., *Lithocarpus* sp., *Mangifera* sp., *Myristica* sp., *Phyllocladus* sp., *Sterculia* sp., *Xanthophyllum* sp.
References: **(ds)** Schedl 1972f: 264. **(tx)** Roberts 1992:(in press); Schedl 1969b: 225–226, 1972f: 264, 1978a: 46.
- nebulosus* Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Gumi L.A., 1900 m; BPBM, Honolulu.
Distribution: New Guinea.
Hosts: *Alphitonia* sp., *Caldcluvia* sp., *Calophyllum* sp., *Castanopsis* sp., *Cryptocarya* sp., *Dacrydium* sp., *Elacocarpus* sp., *Garcinia* sp., *Ilex* sp., *Litsa* sp., *Macaranga* sp., *Nothofagus* sp., *Phyllocladus* sp., *Planchonella* sp., *Syzygium* sp., *Xanthomyrtus* sp.
References: **(tx)** Roberts 1992:(in press).
- obtusicornis* Schedl 1965g: 31. Holotype ♀; O. Borneo, Mt. Tibang, 1300 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: **(ds)** Schedl 1972f: 264. **(tx)** Schedl 1965g: 31, 1972f: 264, 1978a: 49.
- oomsis* Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Oomsis L.A., 800 m; BPBM, Honolulu.
Distribution: New Guinea.
- Hosts: *Burkella* sp., *Canarium* sp., *Dysoxylum* sp., *Elacocarpus* sp., *Garcinia* sp., *Myristica* sp.
References: **(tx)** Roberts 1992:(in press).
- oreogenus* Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Gumi L.A., 2,000 m; BPBM, Honolulu.
Distribution: New Guinea.
Hosts: *Galbulilima* sp.
References: **(tx)** Roberts 1992:(in press).
- papuannus* Schedl 1968e: 269. Holotype ♂; Arafinegetu, E. Highlands Dist.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Agathis alba*, *Araucaria cunninghamii*, *Calophyllum* sp., *Castanopsis* sp., *Casuarina* sp., *Endospermum* sp., *Eugenia* sp., *Garcinia* sp., *Glochidion* sp., *Macaranga* sp., *Myristica* sp., *Planchonella* sp., *Podocarpus* sp., *Sterculia* sp., *Syzygium* sp.
Notes: (3) Schedl 1969b: 225 (in key).
References: **(ds)** Schedl 1969b: 226, 1972f: 264, 1979h: 158. **(tx)** Schedl 1968e: 269, 1969b: 225–226, 1972f: 264, 1978a: 51.
- pendleburyi* Schedl 1936d: 18. Lectotype ♂; Malay Peninsula, Selangor: Kuala Lumpur; Schedl Collection in NIIMW, Wien.
Distribution: Asia (Burma/ Assam in India/ Malaya/ Thailand/ Vietnam), Indonesia (Borneo), Philippine Islands.
Hosts: *Dipterocarpus pilosus*, *Dryobalanops aromatica*, *Pentacme suavis*, *Quercus* sp.
Notes: (3) Schedl 1942a: 218 (described female). Beeson 1941 (1961: 260) (*longispinus*, nomen nudum, synonymy in Browne 1977a: 100).
References: **(hb)** Browne 1941. **(ds)** Beeson 1941 (1961: 260); Browne 1949b, 1961c: 233, 1968a: 134, 1977a: 100; Choo, Woo, & Kim 1981: 203; Nobuchi 1977: 145; Ohno, Yoneyama, & Nakazawa 1982b: 11, 1987b: 90; Schedl 1962f: 169, 1966b: 102, 1966g: 34, 1969a: 207, 218, 1972f: 264. **(tx)** Browne 1949b, 1977a: 100; Nobuchi 1985: 330; Schedl 1936d: 18, 1942a: 218, 1942c: 172, 1972f: 264, 1978a: 52; Strohmeier 1942: 285.
- perpygmaeus* Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Oomsis L.A., 850 m; BPBM, Honolulu.
Distribution: New Guinea.
Hosts: *Calophyllum* spp., *Castanopsis* sp., *Cryptocarya* sp., *Lithocarpus* sp., *Neonauclea* sp., *Planchonella* sp., *Pometia* sp., *Xanthophyllum* sp.
References: **(tx)** Roberts 1992:(in press).
- pusillimus* Chapuis 1865: 335. Syntypes 3♂, 1♀ (Schedl 1960: 63); Nouvelle-Guinée, Dorey; BMNH, London.
Distribution: Asia (India, Malaya), Australia, Bismarck Islands (New Britain Island), Indonesia (Borneo), Madagascar, Micronesia (Caroline Islands), New Guinea, Philippine Islands, Samoan Islands.

Hosts: *Afzelia bijuga*, *A. palembanica*, *Agathis alba*, *Albizzia falcata*, *Anisoptera* sp., *Araucaria* sp., *Altingia exelsa*, *Balanocarpus heimii*, *Canarium* sp., *Castanopsis* sp., *Celtis* sp., *Cryptocarya* sp., *Dipterocarpus* sp., *Dracontomelon* sp., *Dryobalanops aromatica*, *Dyera costulata*, *Eugenia argutata*, *Ficus* spp., *Garcinia* sp., *Grevillea robusta*, *Lithocarpus* sp., *Mangifera* sp., *Nothofagus pullei*, *Planchonella sumocensis*, *Pometia* sp., *Pterocymbium* sp., *Shorea bracteolata*, *S. kunstleri*, *S. leprosula*, *S. singkawang*, *Sloanea* sp., *Sterculia* sp., *Syzygium* sp., *Terminalia brassii*, *Xanthophyllum* sp.

Notes: (3) Schedl 1969a: 225 (in key).

References: (ay) Numberg 1951: 262. (bv) Gray, B. 1974b. (cn) Brimblecombe 1956; Brownie 1968b: 233; Hill, D. S. 1983: 679; Wylie & Shanahan 1975. (cc) Browne 1958b; Thompson, W. R. & Simmonds 1964: 15, 1965: 28. (hb) Beaver 1976b: 544; Beaver & Browne 1978: 617; Beeson 1941 (1961: 260–261); Brimblecombe 1956; Browne 1936a, 1941, 1958b, 1961c: 231–232, 1968b: 233; Gray, B. 1974b; Hill, D. S. 1983: 679; Kalshoven 1960c: 46; Wylie & Shanahan 1975. (ds) Beaver 1976b: 544; Beaver & Browne 1975: 305, 1978: 617; Beeson 1941 (1961: 260); Browne 1936a, 1961a: 317, 1961c: 232, 1966: 242, 1968b: 233, 1980a: 370; Chadwick & Nikitin 1968; Choo, Woo, & Kim 1981: 203; Froggatt 1936; Gemminger & Harold 1872: 2702; Hill, D. S. 1983: 679; Kalshoven 1960d: 46; Lacordaire 1866: 397; Nobuchi 1977: 146; Numberg 1961b: 611; Ohno, Yoneyama, & Nakazawa 1982a, 1982b: 5, 1987a: 96, 1987b: 90; Olmo et al. 1988b: 99, 1989: 67; Sampson 1919: 107, 1925: 3, 1926: 121; Schedl 1936d: 8, 22, 1936g: 519, 1936j: 22, 1966b: 103, 1966g: 34, 1968e: 265, 1972f: 264, 1974c: 262, 1975h: 352, 1975i: 347, 1979a: 160; Strohmeier 1912c: 22, 1914c: 47; Wood, S. L. 1960: 6. (tx) Browne 1966: 242; Carne et al. 1980; Chapuis 1865: 335; Lacordaire 1866: 397; Nobuchi 1985: 330; Sampson 1919: 107, 1925: 3; Schedl 1936e: 61, 1939b: 391, 1939e: 337, 1941c: 354–355, 1942c: 165, 1951k: 141–142, 1954a: 158, 1958i: 215, 1960c: 63, 1965g: 31, 1969a: 225, 1972f: 264; Strohmeier 1912c: 22, 1914c: 47; Wood, S. L. 1960a: 6; Wylie & Yule 1977.

grevilliae Lea 1914: 226 (*Crossotarsus*). Holotype ♀; Queensland; Lea Collection at SAM, Adelaide. Synonymy: Schedl 1936g: 519.

References: (cn) Kleine 1932a: 401; Smith, J. H. 1932: 230, 1935: 3. (cc) Smith, J. H. 1935: 3. (hb) Kleine 1932a: 401. (ds) Kleine 1932a: 401; Smith, J. H. 1935: 3. (tx) Browne 1961c: 231; Lea 1914: 226; Schedl 1936g: 519, 1972f: 264; Wood, S. L. 1960a: 6.

***quadriscopinus* Chapuis** 1865: 332. Syntypes 2♂, 1♀ (Schedl 1960c: 62); D l'Inde orientale; IRSNB, Brussels.

Figures: Nakashima 1975: 60–65. Schedl 1962k: 1054, 1972f: 261.

Distribution: Asia (Burma/ Uttar Pradesh in India).

Hosts: *Ahus nepalensis*, *Castanopsis hystrix*, *Cedrus deodara*, *Lindera pulcherrima*, *Machilus odoratissima*, *Prunus nepalensis*, *Quercus incana*, *Q. lamellosa*.

References: (cn) Kleine 1932a: 311, 402. (cc) Beeson 1921b: 23. (hb) Kleine 1932a: 311, 402; Yunn & Hua 1980: 216. (ds) Beeson 1921b, 1941 (1961: 261); Bhasin, Roonwal, & Singh 1955; Gemminger & Harold 1872: 2702; Kleine 1932a: 311, 402; Nobuchi 1977: 144; Schedl 1962f: 169, 1967e: 222, 1968e: 265, 1972f: 264, 1973b: 212; Strohmeier 1912c: 22, 1914c: 47. (tx) Chapuis 1865: 332; Gardiner 1932b; Hopkins 1914: 121; Nakashima 1975: 60–65; Schedl 1936d: 18, 1939g: 2, 1960c: 62, 1962k: 1054, 1972f: 261, 264; Strohmeier 1912c: 22, 1914c: 47.

***quadriscopinus* Schedl** 1969a: 218. Holotype ♂; Philippine Islands, Iligan to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Philippine Islands.

Hosts: Luanan log.

References: (ds) Lacordaire 1866: 397; Schedl 1972f: 264. (tx) Lacordaire 1866: 397; Nobuchi 1985: 330; Schedl 1969a: 218–219, 1972f: 264, 1987a: 61.

***quinquespinatus* Chapuis** 1865: 334. Syntypes: Iles Celebes, Borneo, Morty, Nouvelle-Guinea: 6♂, 1♀ in BMNH, London, 1♂ in IRSNB, Brussels. Figures: Schedl 1972f: 262 (male).

Distribution: Africa (Angola/ Cameroon/ Central African Republic/ Equatorial Guinea/ Fernando Po/ Ghana/ Nigeria/ Sierra Leone/ Zaire/ Zambia), Asia (Bhutan/ Burma/ Andaman Islands, Assam, Uttar Pradesh in India/ Japan/ Malaya/ Sri Lanka/ Taiwan/ Thailand/ Tonkin in Vietnam), Australia, Fiji Islands, Indonesia (Borneo, Buru, Celebes, Java, Sumatra), Madagascar, New Guinea, Philippine Islands, Samoan Islands, Solomon Islands.

Hosts: *Acmencia* sp., *Afzelia palembanica*, *Agathis alba*, *Araucaria humsteadii*, *Artocarpus chaplasha*, *A. elasticus*, *A. scortechini*, *Balanocarpus heimii*, *Canarium euphyllum*, *Castanea argentea*, *Castanopsis tribuloides*, *Dipterocarpus cornutus*, *D. pilosus*, *Dracontomelon* sp., *Dryobalanops aromatica*, *Durio zibethinus*, *Dyera costulata*, *Dysoxylum binectariferum*, *Elacis guineensis*, *Elatiospermum tapos*, *Endospermum malaccense*, *Eucalyptus grandis*, *Eugenia argutata*, *Fagraea gigantea*, *Glochidion* sp., *Hopea* sp., *Horsfieldia* cf. *glabra*, *Lithocarpus* sp., *Mangifera indica*, *Neonaukea* sp., *Pasania sundaica*, *Pentacme suavis*, *Planchonella samoensis*, *Pometia* sp., *Pygeum parviflorum*, *Quercus fagifolia*, *Q. serrata*, *Sarcocephalus cordatus*, *Shorea assamica*, *S. bentongensis*, *S. bracteolata*, *S. faguetiana*, *S. kunstleri*, *S. leprosula*, *S. parvifolia*, *S. robusta*,

Sindora sp., *Sloanea sigun*, *Tectona grandis*, *Terminalia hederica*, *T. bialata*, *T. manni*, *Vatica lanceaefolia*.

References: (ay) Nakashima 1975: 6; Strohmeier 1918: 34; 1920: 42. (bv) Beeson 1917; Loytty-niemi, Beaver, & Loyttyniemi 1985: 28. (cn) Beeson 1915; Browne 1968b: 233; Kalshoven 1924c; Kleine 1932a: 311, 402; Mathur & Singh 1961a: 26, 1961b: 13; Mayne & Donis 1951: 335; Pierce, W. D. 1917: 194; Roberts 1977b: 574; Roonwal 1954: 85; Stebbing 1914: 632. (ce) Beeson 1921b, 1923; Browne 1958b; Mayne & Donis 1951: 335; Nakashima 1975: 6; Stebbing 1914: 632. (hb) Beaver 1976b: 545; Beaver & Browne 1975: 617; Beaver & Loyttyniemi 1985b: 120; Beeson 1916a, 1917: 10, 1923; Browne 1938a, 1941: 65, 1949: 174–189, 1958b: 164–182, 1961c: 230, 1962: 93, 1963a: 264, 1968: 233; Kalshoven 1924c, 1960c: 45; Kleine 1932a: 311, 402; Lepesme 1947: 649; Loyttyniemi, Beaver, & Loyttyniemi 1985: 28; Roberts 1969: 105–106, 1977b: 574; Schedl 1962k: 1053; Stebbing 1914: 632; Thompson, C. H. 1963: 39. (ds) Beaver 1976b: 545; Beaver & Browne 1975: 305; Beaver & Loyttyniemi 1985b: 120; Beeson 1910: 223, 1916a: 223, 1921b: 23, 1923: 165, 1941: 341, 1961: 261; Bhasin, Roonwal, & Singh 1958: 92; Browne 1938a: 24, 29, 1961a: 317, 1961c: 230–231, 1963a: 264, 1966: 242, 1968b: 233, 1980c: 486; Choo & Woo 1983; Ferreira 1965: 1136; Gemminger & Harold 1872: 2702; Kalshoven 1924: 8, 1960d: 45–46; Kleine 1928: 311, 1932a: 311, 402; Lacordaire 1866: 397; Mathur & Singh 1961a: 26, 1961b: 13; Mayne & Donis 1951: 335, 1962: 266; Miller 1932: 19; Miwa 1931: 265; Murayama 1925a: 217, 1929a: 679, 1931: 203, 1934f: 145, 1937c: 583; Nakashima 1975: 3; Nobuchi 1967: 27, 1977: 45, 1980a: Ohno, Yoneyama, & Nakazawa 1982a: 5, 1982b: 11, 1987a: 96, 1987b: 90; Ohno, Yoshioka, et al. 1988b: 99, 1989: 67; Pierce, W. D. 1917: 194; Roberts 1968: 193, 1977b: 574; Roonwal 1954: 85; Sampson 1919: 107, 1926: 120–121; Schedl 1936d: 8, 1936g: 519, 1937b: 407, 1938d: 452, 1959a: 516, 1959c: 168, 1959d: 67, 1959p: 24, 1961c: 70, 1961e: 128, 156, 1962f: 169, 1962i: 73, 1962k: 1053, 1964f: 620, 1965e: 360, 1965g: 25, 1966b: 103, 1966g: 34, 1971c: 367, 1971f: 150, 1971g: 194, 1972f: 264, 1974c: 262, 1975a: 453, 1975c: 384, 1979a: 160; Strohmeier 1911f: 204, 1912c: 22, 1914c: 47; Thompson, C. H. 1963: 39. (tx) Blandford 1894d: 140; Browne 1966: 242; Chapuis 1865: 334; Janson 1893: 75; Lacordaire 1866: 397; Lucas 1920: 235; Murayama 1925a: 217, 1925b: 235; Nobuchi 1985: 330; Sampson 1919: 107; Schedl 1936e: 61, 1937b: 407, 1937d: 44, 1937e: 544, 1939b: 403–404, 1939c: 2, 1939c: 337, 1941c: 354–355, 1941e: 157, 1942a: 172, 1950d: 14, 1951k: 141, 1953g: 245, 1954c: 155, 1954e: 60, 1955b: 286, 1958i: 215, 1960c: 63,

1962k: 1053–1054, 1972f: 264; Stebbing 1914: 632; Strohmeier 1912c: 22, 1914c: 47; Wylie & Yule 1977. (ms) Lucas 1920: 235.

robustus Schedl 1969b: 225. Holotype ♂; New Guinea: Kuni, 11 miles from Mt. Hagen, W.H.D.; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Acmena* sp., *Castanopsis* sp., *Cryptocarya* sp., *Dryodaphnue* sp., *Lithocarpus* sp., *Nothofagus* sp., *Syzygium* sp.

References: (ds) Schedl 1972f: 264. (tx) Schedl 1969b: 225–226, 1972f: 264, 1978a: 63.

spatulifer Browne 1977a: 101. Holotype ♂; Bengal: Darjeeling, Ranbe; BMNH, London.

Distribution: Asia (India).

Hosts: *Beilschmiedia sikkimensis*, *Bucklandia populuca*, *Castanopsis hystrix*, *Cryptomeria japonica*, *Echonicarpus dasycarpus*, *Lindera pulcherrima*, *Litsea elongata*, *Machilus edulis*, *M. spp.*, *Michelia excelsa*, *Nyssa sessiliflora*, *Prunus nepalensis*, *Quercus glauca*, *Q. dilatata*, *Q. lamellosa*, *Symingtonia populnea*, *Symplocos theaeifolia*.

Notes: (3) Beeson 1941 (1961: 260) (*brochus* nomen nudum, *capitalis* nomen nudum, *discolor* nomen nudum, *spatulifer* nomen nudum, synonymy in Browne 1977a: 102).

References: (cn) Mathur & Singh 1960a: 36, 1961a: 82. (ds) Beeson 1941 (1961: 260); Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1960a: 36, 1961a: 82. (tx) Browne 1977a: 101.

spinatus Browne 1980d: 499. Holotype ♂; Semangka (Sumatra) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sumatra).

References: (ds) Ohno, Yoneyama, & Nakazawa 1982b: 11. (tx) Browne 1980d: 499.

spinifer Schedl 1969b: 224. Holotype ♂; New Guinea: Porotop L.M. Station, W.H.D.; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Ficus* sp., *Nothofagus pullei*, *N. sp.*, *Podocarpus* sp., *Syzygium* sp.

References: (ds) Schedl 1972f: 264. (tx) Schedl 1969b: 224–225, 1972f: 264, 1978a: 68.

tonkinensis Schedl 1972m: 201. Holotype ♂; Tonkin: Ngai-Tio, 4800 ft.; BMNH, London.

Distribution: Asia (Tonkin Island in Vietnam).

References: (tx) Schedl 1972m: 201.

truncatus Niisima & Murayama 1934f: 143. Holotype ♂; Arisan (Formosa); Murayama Collection in USNM, Washington.

Distribution: Asia (Taiwan).

References: (ds) Nobuchi 1967: 27, 1977: 147; Schedl 1972f: 264. (tx) Murayama 1934f: 143, 1937c: 583; Nobuchi 1967: pl. 2; Schedl 1972f: 264.

turgidus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Gumi, 2220 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Antidesma* sp., *Castanopsis* sp., *Dryodaphne* sp., *Garcinia* sp., *Glochidion* sp., *Macaranga* sp., *Microcos* sp., *Planchonella* sp., *Spherostenon* sp., *Syzygium* sp., *Terminalia* sp.

References: (tx) Roberts 1992:(in press).

unispineus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mo., Kuper Range, Wau, 2,300 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Caldcluvia* sp., *Castanopsis* sp., *Elaeocarpus* sp., *Litsea* sp., *Nothofagus* sp.

References: (tx) Roberts 1992:(in press).

Genus *Genyocerus* Motschulsky

GENYOCERUS MOTSCHULSKY 1858: 68. Type-species: *Genyocerus albipennis* Motschulsky, monobasic.

Diacavus Schedl 1939: 363. Type-species: *Diapus frontalis* Strohmeier, original designation. Synonymy: Wood 1969c: 118.

References: (tx) Browne 1949d: 73–76, 1960: 201–220, 1961c: 232–233, 1972a: 188; Schedl 1939c: 363–364, 1962f: 163–169, 1972f: 265–266; Wood, S. L. 1969c: 118.

Notes: (1) Wood 1969c: 118 (identity of genus confirmed). (3) Beeson 1941 (1961: 257) (*atkinsoni*, nomen nudum, from Burma).

References: (ay) Nobuchi 1969a: 69. (hb) Schedl 1972f: 265. (ds) Schedl 1959c: 168, 1966b: 101, 1972f: 265. (tx) Browne 1972a: 188; Chapuis 1865: 338; Schedl 1972f: 265; Strohmeier 1942: 286–287; Wood, S. L. 1969c: 118.

abdominalis (Schedl) 1942a: 218 (*Diacavus*). Lectotype ♂; Malaya, Raub; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 9.

Figures: Nakashima 1975: 69 (mycetangia).

Distribution: Asia (introduced in Japan/ Malaya), Indonesia (Borneo), Philippine Islands.

Hosts: *Balanocarpus heimii*, *Parashorea* sp.

References: (ay) Nakashima 1975: 7. (ec) Nakashima 1975: 7. (ds) Browne 1961c: 233; Choo & Woo 1983; Nakashima 1975: 3; Nobuchi 1977: 152, 1980a; Ohno, Yoneyama, & Nakazawa 1982b: 11; Schedl 1959c: 168, 1966g: 34, 1972f: 267. (tx) Browne 1961c: 233; Nakashima 1975: 69; Nobuchi 1985: 330; Schedl 1942a: 218, 1959c: 168, 1972f: 267, 1978a: 9.

albipennis Motschulsky 1858: 68. Holotype ♀; type label reads "Aus Indien or [word illegible]," all known specimens are from Sri Lanka; IZM, Moscow.

Figures: Browne 1970: 582 (as *irregularis*).

Distribution: Asia (Sri Lanka).

Hosts: *Dipterocarpus* sp.

Notes: (1) Wood 1969c: 118 (identity clarified). (3) Beeson 1961: 260 (*zeylanicus*, nomen nudum, synonymy in Browne 1977a: 97).

References: (ds) Gemminger & Harold 1872: 2702; Lacordaire 1866: 398; Schedl 1959a: 515,

1972f: 267. (tx) Beeson 1941 (1961: 260); Blandford 1893d; Browne 1977a: 97; Hopkins 1914: 122; Lacordaire 1866: 398; Motschulsky 1858: 68; Schedl 1959a: 515, 1972f: 277; Strohmeier 1942: 284–287; Wood, S. L. 1969c: 118.

irregularis Browne 1970: 582 (*Diacavus*). Holotype ♂; Ceylon: in gurjun (*Dipterocarpus*) timber from Ceylon, intercepted at Princes Risborough, England; BMNH, London. Synonymy: Wood 1981: 122.

References: (ds) Schedl 1972f: 268. (tx) Browne 1970: 582–583, 1977a: 99; Schedl 1972f: 268; Wood, S. L. 1981: 122.

biporus (Schedl) 1942a: 217 (*Diacavus*). Lectotype ♂; Malaya, Selangor, Buloh F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 16.

Figures: Schedl 1972f: 265.

Distribution: Asia (Burma/ Cambodia/ Malaya/ Vietnam), Indonesia (Borneo).

Hosts: *Cotylelobium flacum*, *Dipterocarpus pilosus*, *Hopea odorata*, *Parashorea stellata*, *Shorea leprosula*, *S. parvifolia*, *Vatica* sp.

Notes: (3) Beeson 1941 (1961: 257) (*biporus* nomen nudum, synonymy in Browne 1977a: 97).

References: (cn) Browne 1949d, 1949e; Mathur & Singh 1960b: 57; Sen-Sarma & Thakur 1956: 37. (hb) Browne 1961c: 233. (ds) Beeson 1941 (1961: 257); Browne 1961a: 318, 1968a: 132–134, 1970: 577, 1977a: 98, 1980c: 484, 1981b: 599; Mathur & Singh 1960d: 57; Nobuchi 1977: 151; Ohno, Yoneyama, & Nakazawa 1982b, 1987a: 96; Schedl 1969a: 206, 1972f: 265, 267. (tx) Browne 1949d: 73, 1977a: 97; Schedl 1942a: 217, 1978a: 16.

compactus (Schedl) 1966g: 40 (*Diacavus*). Holotype ♂; Diapitan, Philippine Islands to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Philippine Islands.

Hosts: Lauan log.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (ds) Browne 1984c: 449; Choo & Woo 1983; Nobuchi 1977: 149; Ohno, Yoneyama, & Nakazawa 1982b: 11; Schedl 1972f: 267. (tx) Nobuchi 1985: 330; Schedl 1966g: 40, 1972f: 267, 1978a: 20; Wood, S. L. 1969c: 118.

compactus dubiosus Schedl 1966g: 41 (*Diacavus*). Lectotype ♂; Aparri, Philippine Islands to Tokyo (Japan), imported; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 20.

Notes: (3) It appears that this is no more than an aberration.

References: (ds) Schedl 1972f: 267. (tx) Schedl 1966g: 41, 1972f: 267, 1978a: 20.

decemspinatus (Schedl) 1942a: 216 (*Diacavus*). Lectotype ♂; Malaya, Batu Jalam; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 23.

Distribution: Asia (Malaya/ Vietnam), Philippine Islands.

Hosts: *Vatica* spp.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (cn) Browne 1949d. (hb) Browne 1949d. (ds) Browne 1961c: 233, 1965a: 190; Nobuchi 1977: 150; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1966g: 34, 1972f: 267. (tx) Browne 1968a: 134; Nobuchi 1985: 330; Schedl 1942a: 216, 1966b: 102, 1966g: 42, 1972f: 267, 1978a: 23; Wood, S. L. 1969c: 118.

diaphanus (Schedl) 1939c: 364 (*Diacavus*). Lectotype ♀; Malaya, Selangor: Sungai Buloh For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 24.

Distribution: Asia (Assam in India/ Malaya/ Vietnam), Indonesia (Sarawak in Borneo).

Hosts: *Anisoptera* sp., *Artocarpus* sp., *Dipterocarpus cornutus*, *Hopca odorata*, *Parashorea lucida*, *Shorea assamica*, *S. luyppochra*, *S. leprosula*, *S. maxwelliana*.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*). (3) Beeson 1941 (1961: 260) (*hopeae* nomen nudum, synonymy in Browne 1977a: 97). Browne 1960: 216–217 and Schedl 1962f: 168 (described male). References: (cn) Mathur & Singh 1961a: 39. (hb) Beaver & Browne 1978: 617; Browne 1941, 1961c: 233. (ds) Beaver & Browne 1978: 617; Beeson 1941 (1961: 257); Browne 1961c: 233, 1977a: 98; Mathur & Singh 1961a: 39; Schedl 1962f: 168, 1972f: 267. (tx) Browne 1960: 216–217, 1962c: 216, 1964: 760, 1977a: 98; Schedl 1939c: 364, 1962f: 168, 1966g: 42, 1972f: 267, 1972p: 159, 1978a: 24; Wood, S. L. 1969c: 118, 1978a: 24.

assamensis Browne 1964: 760 (*Diacavus*). Holotype ♂; India: Assam, Margherata, Lakhimpur Div.; BMNH, London. Synonymy: Schedl 1972f: 267.

Notes: (3) Beeson 1941 (1961: 257) (*assamensis* nomen nudum, synonymy in Browne 1977a: 97).

References: (cn) Sen-Sarma & Thakur 1986: 37. (ds) Beeson 1941 (1961: 257); Browne 1968a: 132–134, 1970: 576, 1977a: 97, 1980d: 490; Mathew 1982, 1987: 185. (tx) Browne 1964: 760, 1977a: 97; Schedl 1972f: 267, 1972p: 159.

dipterocarp Browne 1977b: 98. Holotype ♂; Andaman Islands; BMNH, London.

Figures: Browne 1977b: 99.

Distribution: Andaman Islands.

Hosts: *Canarium euphyllum*, *Dipterocarpus turbinatus*.

Notes: (3) Beeson 1941 (1961: 257) (*dipterocarp* nomen nudum, synonymy in Browne 1977a: 97). References: (cn) Sen-Sarma & Thakur 1986: 37. (ds) Beeson 1941 (1961: 257); Blasin, Roonval, & Singh 1958; Browne 1981b: 598; Ohno, Yoneyama, & Nakazawa 1982b: 11. (tx) Browne 1977b: 98–99.

exilis (Schedl) 1942a: 218 (*Diacavus*). Lectotype ♂; Malaya, Selangor, Bukit Arang; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 118.

Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).

Hosts: *Shorea lacvis*, *Vatica* sp.

Notes: (3) Browne 1962c: 219 (described female).

References: (hb) Browne 1961c: 233. (ds) Browne 1961c: 233, 1980a: 371, 1984c: 449; Ohno, Yoneyama, & Narazawa 1987a: 96; Schedl 1959c: 168, 1972f: 267. (tx) Browne 1962c: 219; Schedl 1942a: 218, 1972f: 267, 1978a: 29.

frontalis (Strohmeyer) 1914f: 5 (*Diapus*). Holotype ♂; Borneo (Sarawak); Stettin Museum, presumably lost.

Distribution: Indonesia (Sarawak in Borneo).

Notes: (1) Schedl 1939c: 363 (to *Diacavus*), Wood 1969c: 118 (to *Genyocerus*).

References: (ds) Schedl 1972f: 267. (tx) Browne 1964: 760; Schedl 1939c: 363–364, 1942a: 216, 1972f: 267; Strohmeyer 1914f: 5–6; Wood, S. L. 1969c: 118.

intermedius (Browne) 1968a: 134 (*Diacavus*). Holotype ♂; Vietnam: near Bien Hoa; BMNH, London.

Distribution: Asia (Malaya/ Vietnam), Philippine Islands.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (ds) Browne 1977a: 99, 1981b: 597; Schedl 1972f: 268. (tx) Browne 1968a: 134, 1977a: 99; Schedl 1972f: 268; Wood, S. L. 1969c: 118.

laticollis (Browne) 1962c: 217 (*Diacavus*). Holotype ♂; Malaya: Selangor, Keping; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Shorea leprosula*.

Notes: (1) Browne 1964: 753 (original spelling of *laticollis* corrected to *laticollis*). Wood 1969c: 118 (to *Genyocerus*).

References: (ds) Schedl 1972f: 268. (tx) Browne 1962c: 217, 1964: 753; Schedl 1972f: 268; Wood, S. L. 1969c: 118.

multiplus (Schedl) 1936h: 67 (*Diapus*). Holotype ♀; Philippine Islands; Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands.

Notes: (1) Wood 1969: 118 (to *Genyocerus*).

References: (ds) Schedl 1966b: 102, 1972f: 268. (tx) Nobuchi 1985: 330; Schedl 1936h: 67, 1939c: 364, 1942a: 216, 1950g: 894, 1972f: 268, 1978a: 45.

papuanus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Wp., Wawoi-Guavi, 150 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Anisoptera* sp.

References: (tx) Roberts 1992:(in press).

philippinensis (Schedl) 1942a: 216 (*Diacavus*). Holotype ♂; Kolambugan, Philippinen; Schedl Collection in NHMW, Wien.

Figures: Nakashima 1975: 67 (mycetangia).

Distribution: Asia (introduced in Japan/ Malaya), Indonesia (Borneo), Philippine Islands.

Hosts: *Dryobalanops oblongifolia*, *Shorea leprosula*, *S. maxwelliana*.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*). (3) Schedl 1966g: 41 (described female).

References: (ay) Nakashima 1975: 7. (ec) Nakashima 1975: 7. (hb) Beaver & Browne 1978: 617. (ds) Beaver & Browne 1978: 618; Nakashima 1975: 3; Nobuchi 1977: 151; Olmo, Yoneyama, & Nakazawa 1987b: 90; Schedl 1966b: 102, 1966g: 34, 1972f: 268. (tx) Nakashima 1975: 67; Nobuchi 1985: 330; Schedl 1942a: 216, 1963j: 483, 1964h: 425, 1966g: 40–41, 1972f: 268, 1978a: 55.

quadridens Browne 1962c: 218 (*Diacavus*). Holotype ♂; Philippines: Mt. Maquiling, Laguna; Forest Products Research Institute, Laguna. Synonymy: Schedl 1964h: 425.

References: (hb) Browne 1961c: 233. (ds) Browne 1961c: 233–234. (tx) Browne 1962c: 218; Schedl 1963j: 483, 1964h: 425.

plumatus (Schedl) 1936h: 66 (*Diapus*). Holotype ♀; Philippines (Luzon?); Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands (Luzon?).

References: (ds) Schedl 1966b: 103, 1972f: 268. (ds) Nobuchi 1985: 330; Schedl 1936h: 66, 1972f: 268, 1978a: 55. (tx) Schedl 1936h: 66.

puer (Schedl) 1969b: 223 (*Diapus*). Holotype ♂; New Guinea: Omsis, Morobe Distr.; CSIRO, Canberra.

Hosts: *Anisoptera polyandra*, *A. sp.*, *Dysoxylum sp.*, *Hopea sp.*

Notes: (1) Roberts 1992:(in press) (subspecies established).

References: (ds) Schedl 1972f: 264. (tx) Roberts 1992:(in press); Schedl 1972f: 264, 1978a: 58.

puer puer:

Distribution: New Guinea.

References: (tx) Roberts 1992:(in press).

trispinatus Schedl 1974d: 471 (*Diacavus*). Holotype ♂; New Guinea Industries Logging Area, Gabensis, Morobe Distr.; CSIRO, Canberra. Synonymy: Roberts 1992:(in press).

References: (ds) Browne 1984c: 449. 1985: 191. (tx) Browne 1985a: 197; Roberts 1992:(in press); Schedl 1974d: 471, 1978a: 75.

puer fergussonus Roberts 1992:(in press). Holotype ♂; Papua New Guinea: Mb. Camadoudou, 50 m; BPBM, Honolulu.

Distribution: New Guinea (Fergusson Island).

References: (tx) Roberts 1992:(in press).

quadrifoveolatus (Schedl) 1970b: 369 (*Diacavus*). Holotype ♂; Sabah, Semporna (Borneo) to Miyako (Japan), imported; PPST, Tokyo.

Figures: Nobuchi 1978a: 80.

Distribution: Indonesia (Borneo).

Hosts: Lanan log.

References: (ds) Browne 1984c: 449; Nobuchi 1977: 149; Schedl 1972f: 268. (tx) Nobuchi 1978a: 80, 1978b: 82; Schedl 1970b: 369–370, 1972f: 268, 1978a: 60.

quadriporus (Schedl) 1942a: 217 (*Diacavus*). Lectotype ♂; Malaya, Selangor, Buloh F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 60.

Distribution: Asia (Malaya), Philippine Islands.

Hosts: *Shorea bracteolata*, *S. parvifolia*.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (hb) Ymus & Hua 1980: 216. (ds) Schedl 1971c: 366, 1972f: 268; Ymus & Hua 1980: 216. (tx) Schedl 1942a: 217, 1972f: 268, 1978a: 60; Wood, S. L. 1969c: 118.

serratus (Schedl) 1936d: 17 (*Diapus*). Lectotype ♂; Malay Peninsula: Pahang, Bilut For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 66.

Distribution: Asia (India/ Malaya), Indonesia (Sarawak in Borneo).

Hosts: *Dipterocarpus borneensis*, *Dryobalanops oblongifolia*, *Hopea ferrea*, *Parashorea lucida*, *Shorea bracteolata*, *S. laevis*, *S. leprosula*, *S. macrocarpa*, *S. maxwelliana*.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (cn) Browne 1949d. (ec) Browne 1958b. (hb) Beaver & Browne 1978: 618; Browne 1961c: 233–235; Browne 1936a, 1941, 1949d: 73, 1958b; Ymus & Hua 1980: 216. (ds) Beaver & Browne 1978: 618; Browne 1936a, 1941, 1949d, 1958b, 1961a: 317, 1961c: 233. (ds) Beaver & Browne 1978: 618; Beeson 1941 (1961: 260); Browne 1936a, 1961a: 317, 1961c: 233; Schedl 1936j: 22, 1959c: 168, 1964c: 305, 1972f: 268. (tx) Schedl 1936d: 17–18, 22, 1939e: 364, 1942a: 216, 1972f: 268, 1978a: 65.

sexporus (Schedl) 1942a: 217 (*Diacavus*). Lectotype ♂; Malaya, Kuala Lumpur; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 66.

Distribution: Asia (Malaya).

Hosts: *Gonystylus sp.*, *Shorea sp.*

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (hb) Browne 1961c: 235. (ds) Browne 1961c: 235; Choo & Woo 1983; Schedl 1971c: 366, 1972f: 268. (tx) Browne 1964: 760, 1970: 552; Schedl 1942a: 217, 1972f: 268, 1978a: 66; Wood, S. L. 1969c: 118.

shoreae Browne 1981b: 605. Holotype ♂; Butuan (Philippines) to Nagoya (Japan), imported; BMNH, London.

Distribution: Philippine Islands.

Hosts: *Shorea sp.*

References: (tx) Browne 1981b: 605.

strohmeyeri Wood 1992b: 78. Syntypes, sex?; Insel Simaloer, westlich Sumatra; IPKE, Eberswalde, automatic.

Distribution: Indonesia (Sumatra).

References: (tx) Wood, S. L. 1992b: 78–79.

albipennis Strohmeier 1942: 284 (*Diapus*). Syntypes, sex?; Insel Simaloer, westlich Sumatra; IPKE, Eberswalde, preoccupied by Motschulsky 1858.

Notes: (1) Schedl 1972f: 267 (to *Diacavus*, =*Genyocerus*).

References: (tx) Schedl 1959a: 515, 1966g: 40–42, 1972f: 267; Strohmeier 1942: 284; Wood, S. L. 1992b: 78–79.

talurae (Stebbing) 1906: 418 (*Diapus*). Holotype ♂; Talamalai Forest, N. Coimbatore Hills, Madras, India; FRI, Dehra Dun.

Figures: Beeson 1917: pls. 1–2, figs. 1–12 (as *furtivus*), Stebbing 1906: pl. 24, fig. 4.

Distribution: Asia (Assam, Bengal, Bihar, Madhya Pradesh, Orissa, Tamil Nadu, Uttar Pradesh in India/ Pakistan/ Thailand/ Vietnam).

Hosts: *Dipterocarpus pilosus*, *Odina wodier*, *Quercus semicarpifolia*, *Shorea robusta*, *S. talura*, *Vateria indica*, *Vatica lanceaefolia*.

Notes: (1) Beeson 1921: 22 (synonymy of *talurae* and *furtivus* cited). Wood 1981 research notes (male syntype found, synonymous with *furtivus*). Schedl 1939e: 364 (to *Diacavus*, =*Genyocerus*).

References: (cn) Pierce, W. D. 1917: 194. (ds) Pierce, W. D. 1917: 194; Schedl 1972f: 264; Strohmeier 1912c: 22. (tx) Beeson 1921: 22; Schedl 1972f: 264; Stebbing 1914: 626; Strohmeier 1912c: 22.

furtivus Sampson 1913: 450 (*Diapus*). Holotype, sex?; Assam; BMNH, London. Synonymy: Beeson 1921: 22, confirmed by Wood in 1981.

Notes: (1) Schedl 1939e: 364 (to *Diacavus*, =*Genyocerus*).

References: (ay) Farris & Funk 1965: 531; Francke-Grosmann 1956b; Numberg 1951: 262. (cn) Beeson 1915: 8–11, 1916: 1–5, 1918: 114–124, 1919: 10–15, 1922: 81–91, 1935:

539–543; Chatterjee 1923: 1–3; Fisher, Thompson, & Webb 1953; Mathur & Singh 1961a: 45; Pierce, W. D. 1917: 194; Sen-Sarma & Thakur 1986: 37; Stebbing 1914: 630; Waterson 1922: 51–94. (ec) Beeson 1921b: 298, 1923: 222; Farris & Runk 1965: 531; Francke-Grosmann 1956b; Stebbing 1914: 630; Thompson, W. R. & Simmonds 1964: 15, 1965: 68. (hb) Beaver & Browne 1975: 305; Beeson 1915b: 298, 1916a: 222, 1923, 1941 (1961: 258); Imms 1925: 510, 1957: 816; Stebbing 1914: 630. (ds) Beaver & Browne 1975: 305; Beeson 1915b, 1916a, 1919b, 1921b, 1923, 1930, 1941 (1961: 258); Mathur & Singh 1961a: 45; Murayama 1934f: 145; Pierce, W. D. 1917: 194; Schedl 1962f: 168, 1972f: 267; Strohmeier 1914c: 47. (tx) Beeson 1915: 298, 1917: 1–29, 1921: 22, 1930; Browne 1968a: 134; Imms 1925: 510, 1957: 816; Sampson 1913: 450–451; Schedl 1939e: 364, 1972f: 267; Stebbing 1914: 630; Strohmeier 1914c: 47.

mirus Sampson 1913: 452 (*Diapus*). Holotype ♀; Assam; BMNH, London. Synonymy: Schedl 1972q: 268.

Notes: (1) Wood 1969c: 118 (to *Genyocerus*).

References: (cn) Pierce, W. D. 1917: 194; Stebbing 1914: 633. (ec) Stebbing 1914: 633. (hb) Stebbing 1914: 633. (ds) Pierce, W. D. 1917: 194; Strohmeier 1914c: 47. (tx) Beeson 1915: 298, 1917: 1–29; Sampson 1913: 452; Schedl 1939e: 364, 1942a: 216, 1972f: 268; Stebbing 1914: 633; Strohmeier 1914c: 47.

tenellus (Schedl) 1966g: 42 (*Diacavus*). Holotype ♂; Diapitan, Philippine Islands to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Philippine Islands.

Hosts: Lauan log.

References: (ds) Nobuchi 1977: 148; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1972f: 268. (tx) Nobuchi 1985: 330; Schedl 1966g: 42, 1972f: 268, 1978a: 72.

Tribe Tesserocerini Blandford

Tesserocerini genuini

References: Blandford 1895b: 115.

Tesserocerini

References: Schedl 1939b: 396, 400.

Tesserocerariae

References: Strohmeier 1914b: 18–19.

Platyarsulinae

References: Schedl 1972f: 83, 269.

Genus *Platyarsulus* Schedl

PLATYTARSULUS SCHEDL 1935: 632. Type-species: *Platyarsulus marshalli* Schedl, monobasic.

References: (hb) Browne 1961c: 235; Schedl 1972f: 273. (ds) Schedl 1972f: 273. (tx) Blackman 1944: 77; Schedl 1935n: 632, 1939b: 387, 1972f: 273.

biconicus Schedl 1954e: 162. Lectotype, sex?; Borneo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 14.

Figures: Schedl 1972f: 272.

Distribution: Indonesia (Borneo).

Hosts: Camphor wood.

References: (ds) Browne 1981b: 599; Schedl 1972a: 274. (tx) Schedl 1954e: 162, 1972f: 272, 274, 1978a: 14.

borneensis Schedl 1942a: 214. Holotype, sex?; Borneo; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Borneo).

References: (ds) Schedl 1972f: 274. (tx) Schedl 1942a: 214, 1972f: 274, 1978a: 17.

declivis Schedl 1942a: 215. Holotype, sex?; Malaya, Temerloh F.R.; Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

Hosts: *Dryobalanops aromatica*.

References: (ds) Browne 1961c: 235, 1983a: 555; Schedl 1964g: 242, 1971c: 366, 1972f: 274. (tx) Schedl 1942a: 215, 1972f: 274, 1978a: 23.

elongatus Schedl 1942a: 215. Lectotype ♂; Malaya, Selangor, Gombak; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 27.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Balanocarpus heimii*, *Dryobalanops aromatica*, *Shorea faguetiana*.

References: (ds) Browne 1961c: 235, 1980c: 483, 1980d: 491, 1984c: 449; Schedl 1959c: 165, 1971c: 371, 1972f: 274. (tx) Nobuchi 1985: 330; Schedl 1942a: 215, 1972f: 274, 1978a: 27.

marshalli Schedl 1935n: 632. Syntypes, sex?; Malay States: Johore North, Kluang, North Borneo: Bettotan near Sandakan; BMNH, London, and Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Dryobalanops aromatica*.

Notes: (1) Schedl 1978a: 43 (restricts location of "types" to those in BMNH, London).

References: (hb) Browne 1941. (ds) Browne

1961c: 235; Schedl 1972f: 274. (tx) Schedl 1935n: 632–633, 1942a: 215, 1972f: 274, 1978a: 43.

mirabilis Menier 1978: 36. Holotype ♂; Santibang?; BMNH, London.

Figures: Menier 1978: 37.

Distribution: Asia (Malaya), Indonesia (Borneo).

References: (tx) Menier 1978: 36.

nanulus Schedl 1942a: 215. Holotype, sex?; Selangor, Bukit, Arang, 1000 Fuss; Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

References: (ds) Browne 1961c: 235; Schedl 1972f: 274. (tx) Schedl 1942a: 215, 1972f: 274, 1978a: 46.

tuberculatus Browne 1983a: 572. Holotype, sex?; Tatau (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: (tx) Browne 1983a: 572.

Genus *Notoplatypus* Lea

NOTOPLATYPUS LEA 1910: 135. Type-species: *Notoplatypus elongatus* Lea, monobasic.

References: (tx) Blackman 1944: 77; Lea 1910: 135; Numberg 1953: 46; Schedl 1936g: 519, 1939b: 387–389, 1972f: 275–277.

elongatus Lea 1910: 136. Syntypes, sex?; N.S. Wales: Ropes Creek, Galston; Lea Collection in SAM, Adelaide.

Figures: Numberg 1953: pl. 9, fig. 7, Schedl 1972f: 275.

Distribution: Australia (New South Wales, Queensland).

Hosts: *Eucalyptus acmenioides*, *E. drepanophylla*.

References: (ds) Schedl 1936g: 519, 1965g: 25, 1971f: 151, 1972a: 144, 1972f: 277, 1975h: 352; Strohmeier 1912c: 19, 1914c: 29. (tx) Hopkins 1914: 125; Lea 1910: 136; Schedl 1936g: 519, 1939b: 379–380, 384, 1941e: 157–158, 1965g: 25, 1972f: 275–277; Strohmeier 1912c: 19, 1914c: 29.

Genus *Tesserocerus* Saunders

TESSEROCERUS SAUNDERS 1836: 155. Type-species: *Platypus (Tesserocerus) insignis* Saunders, monobasic.

Danicerus Spinola 1837: 333 (in Dejean). Type-species: *Danicerus agilis* Spinola = *Tesserocerus insignis* Saunders. Synonymy: Schedl 1972a: 119.

References: (tx) Schedl 1972a: 119; Spinola 1837: 333.

Tesseroplatus Schedl 1935e: 149. Type-species: *Tesseroplatus ursus* Schedl = *Tesserocerus insignis* Saunders, designated by Wood 1992a: 90–91. Synonymy: Schedl 1972: 115.

References: (tx) Schedl 1935e: 149, 1939b: 400, 1972a: 115; Wood, S. L. 1992a: 90–91.

Tesserocephalus Schedl 1936: 103. Type-species:

- Tesserocephalus forficula* Schedl, monobasic.
Synonymy: Schedl 1972a: 116.
References: (tx) Schedl 1936i: 103, 1939b: 400.
References: (ds) Donrojeanni 1965: 9–25; Reichardt 1965b: 162; Schedl 1972f: 116–119. (tx) Chapuis 1865: 7–23; Sampson 1924: 545; Saunders 1836: 155–157; Schedl 1933c: 164, 1935c: 149, 1939b: 383, 400, 1952a: 444, 1960c: 56, 1972f: 116.
- alternantes** Schedl 1977: 47. Holotype ♂; El Salvador, Metapan. 1600–2200 m; Schedl Collection in NHMW, Wien.
Distribution: North America (El Salvador).
Hosts: *Quercus* sp.
References: (tx) Schedl 1977c: 47.
- belti** Sharp 1880: 112. Syntypes, sex?; Chontales [Nicaragua]; BMNH, London.
Distribution: North America (Costa Rica/ Nicaragua).
Hosts: *Cedrus amarga*, *Vismia guianensis*.
References: (cu) Grandi 1951. (hb) Grandi 1951; Paulian 1949a: 989. (ds) Blackwelder 1947: 791; Blandford 1896e; Equihua & Atkinson 1987: 16; Reichardt 1965b: 163; Schedl 1933c: 163, 1972f: 119; Strohmeier 1911f: 204, 1912c: 20. (tx) Blandford 1896e: 114, 117; Equihua & Atkinson 1987: 16; Schedl 1933c: 163, 1940c: 203, 1941e: 157, 1972f: 119; Sharp 1880: 112; Strohmeier 1911f: 204, 1912c: 20.
- bilobus** (Schedl) 1961i: 231 (*Platypus*). Syntypes ♂; Cayenne; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
References: (ds) Schedl 1972f: 119. (tx) Schedl 1961i: 231–232, 1972f: 119, 1978a: 16.
- brasilensis** Evers 1910: 18. Holotype ♂; Curitiba, Parana (Sud-Brasilien); Hamburg Museum, lost.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 791; Schedl 1972f: 119; Strohmeier 1912c: 19, 1914c: 40. (tx) Evers 1910: 18; Schedl 1972f: 119; Strohmeier 1912c: 19, 1914c: 40.
- chapuisi** Schedl 1937d: 43. Lectotype ♂; Brasilien, St. Catharina, Lages Hochland; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 19. Figures: Schedl 1972f: 116.
Distribution: South America (Brazil/ Colombia).
References: (ds) Blackwelder 1947: 791; Schedl 1972f: 119. (tx) Schedl 1937d: 43, 1952a: 444, 1963d: 234, 1972f: 116, 119, 1978a: 19; Wood, S. L. 1966a: 46.
- contractus** Chapuis 1865: 298. Holotype ♂ (Schedl 1960: 57); Bresil, Obydos; BMNH, London.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 119; Strohmeier 1912c: 19. (tx) Chapuis 1865: 298; Schedl 1960c: 57, 1972f: 119; Strohmeier 1912c: 19.
- despectus** Schedl 1971f: 154. Holotype ♂; Brasilien; UZMC, Copenhagen.
Distribution: South America (Brazil).
References: (ds) Schedl 1972f: 119. (tx) Schedl 1971f: 154–155, 1972f: 119.
- devalquei** Chapuis 1865: 300. Syntypes 2♂, 1♀ (Schedl 1960c: 58); Bresil, Ega; BMNH, London.
Distribution: North America (Costa Rica/ El Salvador/ Guatemala/ Nayarit, Veracruz in Mexico/ Nicaragua/ Panama), South America (Argentina/ Bolivia/ Brazil/ Cayenne/ Colombia/ Paraguay/ Peru/ Suriname/ Venezuela).
Hosts: *Caryocar nucifera*, *Hevea brasiliensis*, *Ocotea rodiaei*.
Notes: (3) Dejean 1837: 333 (*appendiculatus* nomen nudum, synonymy in Schedl 1937d: 43).
References: (ds) Blackwelder 1947: 791; Equihua 1985: 144; Equihua & Atkinson 1987: 16; Estrada & Atkinson 1988: 213; Gemminger & Harold 1872: 2700; Nunnberg 1963c: 107; Reichardt 1965b: 163; Schedl 1960a: 80, 1970c: 89, 1970f: 582, 1971f: 151, 1972f: 119, 1972g: 51, 1976a: 59, 1977e: 44, 1978c: 292, 1979e: 59; Strohmeier 1912c: 19, 1914c: 40. (tx) Chapuis 1865: 300; Dejean 1837: 333; Equihua & Atkinson 1987: 16; Reichardt 1965b: 163; Schedl 1936c: 249, 1936e: 61, 1937d: 43, 1938i: 28, 1948d: 38, 1952a: 445, 1952e: 123, 1952h: 74, 1955c: 2, 1960a: 80, 1960c: 58, 1972f: 119; Strohmeier 1912c: 19, 1914c: 40.
- aubei** Chapuis 1865: 301. Syntypes 1♂, 1♀ (Schedl 1960c: 58); Bresil, Ste-Catherine; BMNH, London. Synonymy: Reichardt 1957b: 163, Schedl 1972: 119.
References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Strohmeier 1914c: 40. (tx) Chapuis 1865: 301; Reichardt 1957b: 163, 1965: 163; Schedl 1960c: 58, 1972a: 119; Strohmeier 1914c: 40.
- dejeani** Chapuis 1865: 303. Syntypes; Colombie et du Mexique; 2♂, 1♀ in BMNH, London, 3♂, 2♀ in IRSNB, Brussels, 1♂, 2♀ in MNB, Berlin (Schedl 1960c: 59). Synonymy: Reichardt 1965b: 163, Schedl 1972a: 119.
References: (hb) Kalshoven 1963: 238; Swabey 1935: 7. (ds) Blackwelder 1947: 791; Blandford 1896e: 115; Ferrer 1942; Gemminger & Harold 1872: 2701; Kalshoven 1963: 238; Schedl 1933c: 163, 1963c: 160, 1966f: 77; Strohmeier 1912c: 19, 1914c: 40; Swabey 1935: 7. (tx) Blandford 1896e; Chapuis 1865: 303; Schedl 1937d: 43, 1940a: 328, 1952e: 123, 1952h: 14, 1960c: 59, 1972a: 119; Strohmeier 1912c: 19, 1914c: 40.
- elegans** Chapuis 1865: 295. Syntypes; Colombie; 1♂ BMNH, London, 1♂, 3♀ in IRSNB, Brussels (Schedl 1960c: 57).
Distribution: North America (Costa Rica/ Panama), South America (Colombia/ Peru/ Venezuela).

Hosts: *Enterolobium* sp., *Erythrina costaricensis*, *Lonchocarpus magaritensis*, *Spondias purpurea*.

References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 119; Strohmeier 1912c: 20, 1914c: 40. (tx) Chapuis 1865: 295; Numberg 1939: 241, 1940: 69; Schedl 1937d: 43, 1952h: 74, 1960c: 57, 1972f: 119; Strohmeier 1912c: 20, 1914c: 40.

ericius Blandford 1896e: 115. Syntypes 1♂, ♂ ♀; Mexico (Orizaba), Guatemala (Cerro Zúnil, El Tumbador, Las Mercedes), Panama (Volcan de Chiriqui); BMNH, London.

Figures: Equihua & Atkinson 1987: 15.

Distribution: North America (Belize/ Costa Rica/ Guatemala/ Veracruz in Mexico/ Panama), South America (Brazil).

Hosts: *Bursera sinarubra*, *Vismia guianensis*.

References: (ds) Atkinson & Equihua 1986a: 423; Blackwelder 1947: 791; Equihua 1985: 144; Equihua & Atkinson 1987: 16; Ferrer 1942; Reichardt 1965b: 163; Schedl 1933c: 163, 1966f: 94, 1967d: 5, 1972f: 119; Strohmeier 1912c: 20, 1914c: 40. (tx) Blandford 1896e: 114–115; Equihua & Atkinson 1987: 15, 16; Reichardt 1965b: 163; Schedl 1940a: 327–328, 1941e: 157, 1972f: 119; Strohmeier 1912c: 20, 1914c: 40.

forceps Wood 1966b: 46. Holotype ♂; Pandora, Limon Prov., Costa Rica; Wood Collection.

Figures: Wood 1966b: 47.

Distribution: North America (Costa Rica).

References: (ds) Equihua & Atkinson 1987: 16; Schedl 1972f: 119. (tx) Equihua & Atkinson 1987: 16; Schedl 1972f: 119; Wood, S. L. 1966b: 46–47.

forcipatus Schedl 1972g: 84. Holotype ♂; Brasilien, Jacareacanga, Para; MZUSP, Sao Paulo.

Distribution: South America (Brazil).

References: (ds) Schedl 1976a: 59. (tx) Schedl 1972g: 84, 1978a: 31.

forcifulus (Schedl) 1936i: 103 (*Tesserocephalus*). Holotype ♀; Guyana; Schedl Collection in NHMW, Wien.

Distribution: South America (Guyana).

Notes: (1) Schedl 1972f: 119 (to *Tesserocherus*).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 119. (tx) Reichardt 1965: 163; Schedl 1936i: 103, 1972f: 119.

fronteproductus Schedl 1936i: 102. Holotype ♂; Brasilien; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 119. (tx) Schedl 1936i: 101–103, 1972f: 119, 1978a: 31.

gebieni Evers 1910: 18. Holotype ♀; Lages (Brasilien); Hamburg Museum, lost.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 119; Strohmeier 1912c: 20, 1914c: 40. (tx)

Evers 1910: 18; Schedl 1972f: 119; Strohmeier 1912c: 20, 1914c: 40.

guerini Chapuis 1865: 299. Syntypes; Bresil: 1♂, 2♀ in BMNH, London, 1♂, 1♀ in IRSNB, Brussels (Schedl 1960c: 58).

Distribution: South America (Argentina/ Bolivia/ Brazil/ Cayenne).

Hosts: *Araucaria angustifolia*, *Cedrela fissilis*, *Eschweilera sagotianum*, *Ocotea rodiaei*.

References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1960a: 75, 1966f: 94, 1967d: 5, 1970e: 89, 1971f: 151, 1972f: 119, 1972f: 51, 1976a: 59; Schonherr & Pedrosa-Macedo 1981: 58; Strohmeier 1912c: 20, 1914c: 40. (tx) Chapuis 1865: 299; Schedl 1936c: 249, 1937d: 43, 1939b: 394, 1941e: 157, 1951h: 285, 1960c: 58, 1972f: 119; Strohmeier 1912c: 20, 1914c: 40.

guerini montanus Schedl 1960a: 78. Syntypes, sex?; Corvico, Huarinillas, Bolivia, 1160 m; not given, presumably CAS, San Francisco, and Schedl Collection in NHMW, Wien.

Notes: (1) Schedl 1972f: 119 (not mentioned).

References: (tx) Schedl 1960a: 78.

inermis Guérin-Meneville 1835a: 106. Holotype ♂; Cayenne; MNHN, Paris.

Distribution: South America (Brazil/ Cayenne/ Ecuador/ Guyana/ Peru).

References: (ds) Blackwelder 1947: 791; Chapuis 1865; Gemminger & Harold 1872: 2701; Schedl 1972f: 119, 1976a: 59; Strohmeier 1912c: 20. (tx) Chapuis 1865: 310; Guérin-Meneville 1835a: 106; Lucas 1920: 633; Schedl 1941e: 157, 1960c: 61, 1972f: 119; Strohmeier 1912c: 20. (ms) Lucas 1920: 633.

insignis (Saunders) 1836: 155 (*Platypus*). Holotype ♀; Brasilia circum Monte Video; possibly in BMNH, London.

Distribution: South America (Argentina/ Bolivia/ Brazil/ Paraguay/ Uruguay).

Hosts: *Cedrela fissilis*.

Notes: Dejean 1837: 333 (*Danicerus agilis* nomen nudum, *D. denticornis* nomen nudum, *D. melanocephalus* nomen nudum, synonymy in Schedl 1972f: 119–120).

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (ds) Blackwelder 1947: 791; Blandford 1895b; Chapuis 1865; Gemminger & Harold 1872: 2701; Lacordaire 1866: 392; Santoro 1957b: 25; Schedl 1966f: 94, 1970e: 88, 1971f: 151, 1972f: 119, 1972g: 51, 1976a: 59, 1979e: 59; Schonherr & Pedrosa-Macedo 1981: 58; Strohmeier 1911f: 204, 1912c: 20, 1914c: 40. (tx) Blandford 1895b; Castelnau 1840; Chapuis 1865: 308; Dejean 1837: 333; Guérin-Meneville 1835a; Hopkins 1914: 130; Lacordaire 1866: 392; Lucas 1920: 633; Numberg 1959: 493; Saunders 1836: 155; Schedl 1935e: 150, 1936c: 249, 1936e: 61, 1937d: 43, 1939j: 565, 1941e: 157, 1951h: 285–

- 286, 1952d: 343, 1960c: 60, 1966e: 43, 1972f: 119–120; Sharp 1880: 112; Strohmeier 1911c: 204, 1912c: 20, 1914c: 40. (**ms**) Lucas 1920: 633. *bilamatus* Guérin-Meneville 1838a: 106. Holotype ♀; Bresil: MNHN, Paris. Synonymy: Schedl 1972f: 119. References: (**ds**) Blackwelder 1947: 791. (**tx**) Guérin-Meneville 1838a: 106; Schedl 1972f: 119–120.
- ursus* Schedl 1935e: 149 (*Tesseroplatypus*). Holotype ♀; Parana, Brasilien; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 120. References: (**ds**) Blackwelder 1947: 791. (**tx**) Schedl 1935e: 149, 1972f: 120, 1978a: 78.
- linearis** Chapuis 1865: 302. Syntypes, sex?; Bresil; BMNH, London. Distribution: South America (Brazil). References: (**ds**) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 302; Schedl 1960c: 58, 1972f: 120; Strohmeier 1912c: 20, 1914c: 40.
- morsi** Chapuis 1865: 298. Syntypes 2 ♀; Cayenne: 1 ♀ IRSNB, Brussels, 1 ♀ MNB, Berlin. Distribution: South America (Cayenne). Notes: (3) Schedl 1960: 57 (*mutilatus* Chapuis, nomen nudum, synonymy in Schedl 1972f: 120). References: (**ds**) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 298; Schedl 1937d: 33, 43, 1960c: 57, 1972f: 120; Strohmeier 1912c: 20, 1914c: 40.
- procer** (Erichson) 1847: 135 (*Platypus*). Holotype ♀; not given; MNB, Berlin. Distribution: South America (Bolivia/ Brazil/ Cayenne/ Guyana/ Peru/ Suriname/ Venezuela). Hosts: *Catostemma* sp., *Sinarubra snara*. References: (**ds**) Blackwelder 1947: 791; Chapuis 1865; Gemminger & Harold 1872: 2701; Lacordaire 1866: 392; Schedl 1972f: 120, 1972g: 51, 1976a: 59, 1978c: 293; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 305; Erichson 1847: 135; Lacordaire 1866: 392; Numborg 1939: 241, 1940: 69; Schedl 1937d: 33, 43, 1952d: 343, 1952h: 74, 1960c: 59, 1972f: 120; Strohmeier 1912c: 20, 1914c: 40.
- obtus* Chapuis 1865: 304. Syntypes 1♂, 1♀ (Schedl 1960c: 59); Guyane française, Cayenne et du Bresil, Ega; BMNH, London. Synonymy: Schedl 1960c: 59. References: (**ds**) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1960a: 79; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 304–305; Schedl 1936e: 61, 1941e: 157, 1952d: 343, 1952h: 74, 1958c: 2, 1960a: 79, 1960c: 59, 1972a: 120; Strohmeier 1912c: 20, 1914c: 40.
- productus** Shuckard 1838: 509. Holotype ♀; Brazil?; Shuckard Collection. Distribution: South America (Brazil). References: (**ds**) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Strohmeier 1912c: 20. (**tx**) Schedl 1972a: 120; Shuckard 1838: 509; Strohmeier 1912c: 20.
- quadrident** Schedl 1972g: 54. Holotype ♂; Brasilien, Jacareacanga, Para; MZUSP, Sao Paulo. Distribution: South America (Brazil). Notes: (3) Schedl 1976a: 92 (described female). References: (**tx**) Schedl 1972g: 54, 1976a: 92, 1978a: 60.
- retusus** Guérin-Meneville 1838a: 106. Holotype ♂; Mexique; MNHN, Paris (not found by Schedl 1960c: 56). Distribution: North America (apparently Veracruz in Mexico), South America (Colombia). References: (**ds**) Blackwelder 1947: 791; Chapuis 1865: 294; Equihua & Atkinson 1957: 16; Gemminger & Harold 1872: 2701; Lacordaire 1866: 392; Schedl 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 294; Equihua & Atkinson 1957: 16; Guérin-Meneville 1838a: 106; Lacordaire 1866: 392; Lucas 1920: 633; Schedl 1960c: 56, 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (**ms**) Lucas 1920: 633.
- affinis* Guérin-Meneville 1838: 106. Holotype ♀; Mexique; MNHN, Paris. Synonymy: Schedl 1972f: 120. References: (**tx**) Guérin-Meneville 1838a: 106; Schedl 1960c: 56, 1972f: 120.
- rudis** Chapuis 1865: 297. Holotype ♀ (Schedl 1960c: 47); De la province de Venezuela et Caracas; MNHN, Paris. Distribution: South America (Venezuela). Hosts: *Eschweilera* sp. References: (**ds**) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (**tx**) Chapuis 1865: 297; Schedl 1960c: 47, 1972f: 120; Strohmeier 1912c: 20, 1914c: 40.
- schmutzenhoferi** Schedl 1977e: 48. Holotype ♂; El Salvador, Metapan, 1600–2200 m; Schedl Collection in NHMW, Wien. Distribution: North America (El Salvador). Hosts: *Quercus* sp. References: (**tx**) Schedl 1977e: 48.
- simulatus** Schedl 1936c: 249. Lectotype ♀; Guyane Française; Gourdonville, St.-Laurent du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 66. Distribution: South America (Cayenne). References: (**ds**) Blackwelder 1947: 791; Numborg 1958a: 482, 492; Schedl 1972f: 120. (**tx**) Numborg 1958a: 482, 492; Schedl 1936c: 249, 1960h: 111, 1972f: 120, 1978a: 66.
- spinax** Blandford 1896c: 116. Syntypes 2 ♂; Nicaragua (Chontales), Panama (Volcan de Chiriqui); BMNH, London.

Distribution: North America (Costa Rica/ Nicaragua/ Panama), South America (Brazil/ Colombia/ Guyana).

Hosts: *Caryocar nucifera*, *Dialanthera gordonifolia*, *Slonea multiflora*, *Visnua guianensis*.

References: (ds) Blackwelder 1947: 791; Equihua & Atkinson 1987: 16; Schedl 1972f: 120, 1972g: 52; Strohmeier 1912c: 20, 1914c: 40. (tx) Blandford 1896e: 114, 116; Equihua & Atkinson 1987: 16; Schedl 1933c: 163, 1939j: 565, 1972f: 120, 1972g: 55, 1978a: 68; Strohmeier 1912c: 20, 1914c: 40.

spinolae Chapuis 1865: 307. Syntypes 2; Mexique; 1♂ BMNH, London, 1♂ IRSNB, Brussels (Schedl 1960c: 60).

Distribution: North America (Puebla in Mexico).

Hosts: *Alnus* sp.

References: (ds) Blackwelder 1947: 791; Blandford 1896e: 115; Equihua & Atkinson 1987: 16; Ferrer 1942; Gemminger & Harold 1872: 2701; Schedl 1972f: 120; Strohmeier 1912c: 20, 1914c: 40. (tx) Blandford 1896e: 115; Chapuis 1865: 307; Equihua & Atkinson 1987: 16; Schedl 1940a: 328, 1960c: 60, 1972f: 120; Sharp 1880: 112; Strohmeier 1912c: 20, 1914c: 40.

vegrandis Schedl 1935e: 150 (*Tesseroplattypus*).

Holotype ♀; Mexico; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 120.

References: (ds) Blackwelder 1947: 791; Ferrer 1942. (tx) Schedl 1935e: 150–151, 1940a: 328, 1972f: 120, 1978a: 79.

ustulatus Shuckard 1838: 508. Holotype ♂; Brazil?; Shuckard Collection.

Distribution: South America (Brazil?).

References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2701; Schedl 1972f: 120; Strohmeier 1912c: 20. (tx) Schedl 1972f: 120; Shuckard 1838: 508–509; Strohmeier 1912c: 20.

Genus *Tesserocranulus* Schedl

TESSEROCRANULUS SCHEDL 1933: 164. Type-species: *Tesserocranulus nevermanni* Schedl, original designation.

References: (ds) Schedl 1972f: 121. (tx) Reichardt 1965b: 165; Schedl 1933c: 164, 1939b: 400, 1972f: 121.

nevermanni Schedl 1933c: 164. Holotype ♂; Costa Rica, Hamburgfarn, Reventazon, Ebene Limon; Schedl Collection in NHMW, Wien.

Figures: Reichardt 1965b: 161, Schedl 1972f: 121 (male).

Distribution: North America (Costa Rica), South America (Cayenne).

Hosts: *Eschweileria* sp.

References: (ds) Blackwelder 1947: 791; Equihua & Atkinson 1987: 18; Reichardt 1965b: 165; Schedl 1933c: 164, 1972f: 123, 1976a: 59. (tx) Equihua & Atkinson 1987: 18; Reichardt 1965b:

161, 165; Schedl 1933c: 164, 1935e: 151–152, 1936c: 250, 1940c: 203, 1972f: 121, 123, 1978a: 47.

Genus *Spathidicerus* Chapuis

SPATHIDICERUS CHAPUIS 1865: 42, 311. Type-species: *Spathidicerus thomsoni* Chapuis, subsequent designation by Hopkins 1914: 129.

References: (hb) Schedl 1972f: 256. (ds) Schedl 1966b: 101, 1972f: 256. (tx) Chapuis 1865: 42, 311; Hopkins 1914: 129; Murayama 1925: 199–201; Nunberg 1963d: 357–361; Schedl 1939b: 403, 1941c: 359–360, 1957b: 159, 1960c: 61, 1972f: 256–258.

corporaali Browne 1964: 753. Holotype ♀; Sumatra; BMNH, London.

Distribution: Indonesia (Sumatra).

Notes: (3) Schedl 1972f: 253 (treated in *Periommatius*).

References: (ds) Schedl 1972f: 253. (tx) Browne 1964: 753; Schedl 1972f: 253.

javanus Strohmeier 1913: 165. Syntypes ♂ ♀; Java (Banjoevangi); IPKE, Eberswalde.

Distribution: Indonesia (Java).

References: (ds) Schedl 1972f: 255; Strohmeier 1914c: 43. (tx) Schedl 1972f: 258; Strohmeier 1913: 165, 1914c: 43.

nobilis Chapuis 1865: 315. Syntypes 2♂; Nouvelle-Guinee; BMNH, London.

Distribution: New Guinea.

References: (ds) Browne 1980a: 373; Gemminger & Harold 1872: 2701; Nobuchi 1977: 143; Ohno, Yoshioka, et al. 1988b: 98, 1989: 67; Schedl 1962i: 73, 1972f: 258; Strohmeier 1912c: 21, 1914c: 43. (tx) Chapuis 1865: 315; Schedl 1960c: 61, 1962i: 73, 75, 1972f: 258, 1978a: 47; Strohmeier 1912c: 21, 1914c: 43.

intermedius Schedl 1936g: 519. Syntypes ♂; North-East Papua: Mt. Lamington, 1300–1500 feet; SAM, Adelaide, and Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 258.

References: (ds) Schedl 1936g: 519, 1968f: 537. (tx) Schedl 1936g: 519–520, 1939b: 379, 1941e: 157, 1972f: 255.

papuanus Schedl 1935f: 276. Lectotype ♀; Papua: Kokoda, 1200 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 51.

Distribution: New Guinea.

Notes: (3) Schedl 1972f: 255 (treated in *Periommatius*).

References: (ds) Schedl 1972f: 255. (tx) Schedl 1935f: 276, 1940b: 434, 1942a: 216, 1972f: 255, 1978a: 51.

philippinensis Schedl 1942a: 216. Holotype ♂; Philippines; Schedl Collection in NHMW, Wien. Distribution: Philippine Islands.

Notes: (3) Schedl 1972f: 255 (treated in *Periommatius*).

References: (ds) Schedl 1966b: 101, 1972f: 255.

(**tx**) Nobuchi 1985: 330; Schedl 1942a: 216, 1972f: 255, 1978a: 55.

sumatranus Schedl 1935j: 4. Lectotype ♀; N.O. Sumatra, Tebing-tinggi; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 72. Distribution: Asia (Malaya), Indonesia (Sumatra). Notes: (3) Schedl 1961c: 70, 74 (described male), 1972f: 255 (treated in *Periommatius*). References: (**ds**) Schedl 1972f: 255, 1975a: 453. (**tx**) Schedl 1935j: 4, 1961c: 70, 74, 1972f: 255, 1978a: 72.

thomsoni Chapuis 1865: 314. Holotype ♀; Inde orientale, Chenabor; IRSNB, Brussels (Schedl 1960h: 61). Figures: Schedl 1972f: 256 (male, part of female). Distribution: Indonesia (Celebes, Java, Sumatra), New Guinea. References: (**ay**) Strohmeier 1918: 21. (**bv**) Meixner 1937: 1218. (**cc**) Fuller 1955: 100. (**ds**) Browne 1980a: 371; Genninger & Harold 1872: 2701; Lacordaire 1866: 392; Schedl 1971c: 367, 1972f: 258; Strohmeier 1911f: 204, 1912c: 21, 1914c: 43. (**tx**) Chapuis 1865: 314; Hopkins 1914: 129; Lacordaire 1866: 392; Lucas 1920: 572; Meixner 1937: 1218; Schedl 1937d: 44, 1942c: 165, 1960h: 61, 1972f: 256–258; Strohmeier 1911c: 105, 1912c: 21, 1913: 165, 1914c: 43. (**ms**) Lucas 1920: 572.

Genus *Periommatius* Chapuis

PERIOMMATIUS CHAPUIS 1865: 42, 316. Type-species: *Periommatius longicollis* Chapuis, monobasic.

Asctus Numberg 1958a: 10. Type-species: *Periommatius severini* Strohmeier, designated by Wood 1992a: 90. Synonymy: Schedl 1972f: 250.

References: (**tx**) Numberg 1958a: 10; Schedl 1972f: 250; Wood, S. L. 1992a: 90.

Scitantis Numberg 1966b: 25. Type-species: *Periommatius titschacki* Schedl, original designation. Synonymy: Schedl 1972f: 250.

References: (**tx**) Numberg 1966b: 25; Schedl 1972f: 250.

References: (**hb**) Schedl 1972f: 250. (**ds**) Schedl 1972f: 250. (**tx**) Blackman 1944: 77; Browne 1972a: 179; Chapuis 1865: 42, 316–318; Hopkins 1914: 126, 134, 1915c: 227; Kolbe 1897: 284; Lacordaire 1866: 389, 394; Numberg 1953: 44, 1958a: 1–57, 1966b: 17–39, 1968b: 373–380; Sampson 1924: 132–133; Schedl 1933c: 164, 1933f: 204, 1935c: 154, 1935f: 276, 1936e: 62, 1937d: 44, 1939b: 403–404, 1939g: 169, 1941c: 359–360, 1952i: 7, 1957d: 147, 1960c: 61, 1962k: 1000, 1965f: 3–15, 1969b: 104, 1972f: 250, 1977b: 272; Strohmeier 1911c: 105, 1912a: 17–18, 27–28, 1912c: 21, 1914b: 4–42, 1918: 1–46.

abruptus Numberg 1966b: 26. Holotype ♂; Idanre, Nigeria; Federal Department of Forest Research in Ibadan, Nigeria.

Figures: Numberg 1966b: 39.

Distribution: Africa (Nigeria).

Hosts: *Strombosia pustula*.

References: (**ds**) Numberg 1966b: 20, 26; Roberts 1968: 192; Schedl 1972f: 252. (**tx**) Numberg 1966b: 26, 39; Schedl 1972f: 252.

angustior Schedl 1954e: 87. Lectotype ♂; Suiyani [Gold Coast]; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 11.

Figures: Numberg 1966b: pl. 2, figs. 22, 25.

Distribution: Africa (Ghana/ Nigeria/ Zaire).

Hosts: *Chlorophora excelsa*, *Funtumia elastica*, *Terminalia superba*, *Xylopia* sp.

Notes: (3) Numberg 1958b: 22 (description quoted), Schedl 1972f: 253 (a subspecies of *excisus*).

References: (**ds**) Browne 1983a: 556; Numberg 1968b: 373; Roberts 1960b: 35, 1960e, 1960f; Schedl 1962k: 1029, 1966c: 234, 1967e: 222, 1972f: 253. (**tx**) Numberg 1958b: 22, 1966b: 32; Schedl 1954e: 59, 87, 1962k: 1029, 1972f: 253, 1978a: 11.

simillimus Numberg 1966b: 34. Holotype ♂; Nigerien, Oban; Federal Department of Forest Research in Ibadan, Nigeria. Synonymy: Schedl 1972f: 253.

References: (**ds**) Numberg 1966b: 21; Roberts 1965: 193. (**tx**) Numberg 1966b: 34, 39; Schedl 1972f: 253.

artecircularis Schedl 1966c: 238. Holotype ♂; Congo ex-belge, Luki; Schedl Collection in NHMW, Wien.

Distribution: Africa (Congo/ Zaire).

Hosts: *Xylopia* sp.

References: (**ds**) Schedl 1972f: 252. (**tx**) Schedl 1966c: 238, 1972f: 252, 1978a: 12.

artestrigatus Numberg 1958b: 23. Holotype ♀; Congo Belge: Sankuru; MRCB, Tervuren.

Figures: Numberg 1958b: pl. 1, figs. 1–5.

Distribution: Africa (Uganda/ Zaire).

References: (**ds**) Browne 1972c: 103, 1983a: 556; Schedl 1972: 252. (**tx**) Browne 1972c: 103; Numberg 1958b: 23; Schedl 1972f: 252.

basilewskyi Numberg 1958b: 24. Holotype ♂; Congo Belge: Ruanda, Biumba; MRCB, Tervuren. Distribution: Africa (Zaire).

References: (**ds**) Schedl 1972f: 252. (**tx**) Numberg 1958b: 24; Schedl 1972f: 252.

bicallosus Schedl 1971g: 199. Holotype ♂; Guinea espanol, Mikmeseng; Instituto Espanol de Entom. Distribution: Africa (Equatorial Guinea).

References: (**tx**) Schedl 1971g: 199.

bifrons Schedl 1936b: 134. Holotype, sex?; Kamerun; Schedl Collection in NHMW, Wien.

Figures: Numberg 1966b: pl. 2, figs. 4–11.

Distribution: Africa (Angola/ Cameroon/ Nigeria). Notes: (3) Numberg 1958b: 26 (description quoted), 1966: 27 (described female).

References: (**ds**) Ferreira 1965: 1134; Numberg

1966b: 20; Roberts 1968: 193; Schedl 1959p: 23, 1962k: 1035, 1972f: 252. (tx) Nunberg 1958b: 26, 1966b: 27, 39; Schedl 1936b: 134, 1962k: 1035, 1972f: 252, 1978a: 15

bispinus Strohmeyer 1912c: 19. Syntypes 3 ♂; Kamerun; Strohmeyer Collection.

Figures: Nunberg 1958b: pl. 1, figs. 11–17, pl. 2, figs. 1–4.

Distribution: Africa (Cameroon/ Equatorial Guinea/ Ghana/ Guinea/ Ivory Coast/ Nigeria/ Zaire).

Hosts: *Albizia zygia*, *Celtis soyauxii*, *Xylopia villosa*.

Notes: (3) Nunberg 1968b: 374–375 (named aberration *brevispinatus*, no status).

References: (ay) Strohmeyer 1918: 18, 1920: 22. (ds) Nunberg 1968b: 374; Roberts 1968: 193; Schedl 1962h: 63, 1962k: 1001, 1964f: 620, 1965g: 22, 1966a: 276, 1971e: 3, 1971g: 195, 1972f: 252, 1979b: 417; Strohmeyer 1912c: 2, 1914c: 42; Wood, S. L. 1957e: 1272. (tx) Nunberg 1958b: 27, 1968b: 374–377; Sampson 1924c: 132; Schedl 1937d: 44, 1955d: 270, 1962k: 1001, 1972f: 252; Strohmeyer 1912c: 19–21, 1914c: 42, 1918: 22.

circularis Schedl 1935e: 174. Holotype ♂; Kongo (Haut-Uele; Mauda) (Schedl 1978a: 19); Schedl Collection in NHMW, Wien.

Figures: Nunberg 1966b: pl. 1, figs. 1–2.

Distribution: Africa (Uganda/ Zaire).

Hosts: *Celtis brieii*, *Cyometra haukei*.

References: (ds) Mayne & Donis 1962: 579; Nunberg 1966b: 18; Schedl 1962h: 63, 1962k: 1030, 1972f: 252. (tx) Nunberg 1958b: 32, 1966b: 18; Schedl 1935e: 174, 1941d: 421, 1962k: 1030, 1972f: 252, 1978a: 19.

circularis circulariceps Schedl 1962k: 1030. Holotype ♂; Congo Belge; Yangambi; Schedl Collection in NHMW, Wien.

References: (ds) Schedl 1966c: 234, 1972f: 252. (tx) Schedl 1962k: 1030, 1972f: 252, 1978a: 20.

collaticius Schedl 1941d: 420. Holotype ♂; Ogooue [French West Africa]; Schedl Collection in NHMW, Wien.

Distribution: Africa (Gabon).

References: (ds) Schedl 1972f: 252. (tx) Nunberg 1958b: 12; Schedl 1941d: 420, 1957d: 148, 1962k: 1036, 1972f: 252, 1978a: 20.

congoanus Nunberg 1958b: 33. Holotype ♀; [Belge, Congo] Maniema, bei riv. Kasa-Niemba; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (cn) Mayne & Donis 1951: 332. (ce) Mayne & Donis 1951: 332. (ds) Schedl 1972f: 252. (tx) Nunberg 1958b: 33; Schedl 1972f: 252.

costatus Nunberg 1966b: 28. Holotype ♂; Nigerien, Oban; Federal Department of Forest Research in Ibadan, Nigeria.

Figures: Nunberg 1966b: pl. 2, figs. 12–13.

Distribution: Africa (Nigeria).

References: (ds) Nunberg 1966b: 20; Roberts 1968: 193; Schedl 1972f: 253. (tx) Nunberg 1966b: 39; Schedl 1972f: 253.

defranatus Schedl 1941d: 420. Holotype ♂; Ogooue [French West Africa]; Schedl Collection in NHMW, Wien.

Distribution: Africa (Congo/ Equatorial Guinea/ Gabon/ Zaire).

Notes: (3) Nunberg 1958b: 34 (description quoted). References: (ds) Schedl 1966c: 234, 1971g: 195, 1972f: 253. (tx) Nunberg 1958b: 34; Schedl 1941d: 420, 1962k: 1034, 1972f: 253, 1978a: 24.

elongatus Nunberg 1958b: 35. Holotype ♂; Belgisch-Congo, Elisabethville; MRCB, Tervuren. Figures: Nunberg 1958b: pl. 3, figs. 5–8.

Distribution: Africa (Zaire).

References: (ds) Schedl 1972f: 253. (tx) Nunberg 1958b: 35; Schedl 1972f: 253.

excisus Strohmeyer 1912: 22. Holotype ♂; Kongo; Strohmeyer Collection.

Figures: Mayne & Donis 1962: 279, Nunberg 1958b: pl. 3, figs. 9–12, 1966b: pl. 1, figs. 6–8, 1968b: 377, Roche & Lhoste 1960: fig. c, Schedl 1962k: 1028, 1972f: 250.

Distribution: Africa (Angola/ Cameroon/ Ethiopia/ Equatorial Guinea/ Fernando Po/ Gabon/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Principe Island/ Sao Tome Island/ Sierra Leone/ Spanish Guinea/ Tanzania/ Zaire).

Hosts: Browne 1963a: 247, Schedl 1962k: 1020–1023 (lists many hosts). *Hypodaphnis* sp.

Notes: (3) Nunberg 1958b: 37 (named aberration *sulcatus*, synonymy in Schedl 1972f: 253).

References: (ay) Lhoste & Roche 1960; Roche & Lhoste 1960: 2056. (en) Thompson, G. H. 1960. (ce) Jover 1952: 73–81; Roche & Lhoste 1960: 2056. (hb) Browne 1963a: 247; Jover 1952: 73–81. (ds) Browne 1963a: 247; Cachan 1957: 6, 15; Cola 1971; Ferreira 1965: 1135; Gardner 1957a: 29; Jones, Roberts, & Baker 1959: 13, 23, 45, 53; Mayne & Donis 1960: 98, 1962: 279; Menier 1973a; Nunberg 1966b: 20; Roberts 1968: 193, 1969: 116; Schedl 1962h: 63, 1962k: 1018, 1964f: 620, 1965g: 22, 1966a: 276, 1966c: 234, 1971e: 3, 1971g: 195, 1972e: 285, 1972f: 253, 1972k: 296, 1975h: 352, 1979b: 417; Strohmeyer 1912c: 21, 1914c: 42; Thompson, G. H. 1960, 1963: 55; Webb & Jones 1959: 35–39. (tx) Mayne & Donis 1962: 279; Nunberg 1958b: 36, 1968b: 375, 377; Roche & Lhoste 1960: fig. c; Schedl 1936e: 62, 1937d: 44, 1938d: 452, 1939g: 167, 1941e: 156, 1950c: 205–206, 1950d: 12, 1950e: 213, 1951f: 40, 1953g: 245, 1954d: 875, 1954e: 60, 1955d: 270–271, 1957d: 149, 1959p: 23, 1962k: 1018–1023, 1028–1036, 1972f: 253; Strohmeyer 1912a: 22, 1912c: 21, 1914c: 42, 1920: 1–47.

gracilis Strohmeyer 1912a: 24. Holotype ♀; Kamerun; IPKE, Eberswalde.

Distribution: Africa (Cameroon/ Zaire).

Notes: (3) Numberg 1958b: 37 (description quoted), 1966: 23 (redescribed female).

References: (ay) Strohmeier 1918: 17. (ds) Numberg 1966b: 19; Schedl 1972f: 253; Strohmeier 1911c: 106, 1912c: 21, 1914c: 42. (tx) Numberg 1958b: 37, 1966b: 23; Schedl 1962k: 1036, 1972f: 253; Strohmeier 1911c: 106, 1912a: 24, 1912c: 21, 1914c: 42.

grandis Schedl 1953g: 248. Syntypes, sex?; Kamerun: Umgeb. Kamerunberg, Missellele, Congo Belge: Mayidi, Moto, Stanleyville; MRCB, Tervuren and Schedl Collection in NHMW, Wien. Figures: Numberg 1958b:pl. 3, figs. 19–20.

Distribution: Africa (Cameroon/ Equatorial Guinea/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Uganda/ Zaïre).

Hosts: Schedl 1962k: 1005–1009 (numerous hosts listed).

Notes: (1) Schedl 1978a: 33 (citation of holotype invalid). (3) Numberg 1958b: 37 (description quoted), Schedl 1972f: 254 (treated as a subspecies of *longicollis*).

References: (ds) Browne 1983a: 556; Cachan 1957: 15; Gardner 1957a: 29; Jones, Roberts, & Baker 1959: 13–53; Mayne & Donis 1960: 98, 1962: 280; Numberg 1966b: 19; Roberts 1960b: 36, 39, 1960e, 1960f, 1968: 193; Schedl 1962h: 63, 1962k: 1005, 1964f: 620, 1967e: 222, 1971g: 196, 1972e: 286, 1972f: 620; Thompson, G. H. 1963: 55. (tx) Numberg 1958b: 37; Powell, W. 1987: 28; Sampson 1924: 133; Schedl 1933f: 204, 1939b: 354, 1950d: 12, 1953g: 248, 1954e: 60, 86, 1962h: 63, 1962k: 1005, 1972f: 254, 1978a: 33.

inermis Strohmeier 1912a: 26. Syntypes ♂ ♀; Kinchassa, Kongo; Strohmeier Collection. Figures: Numberg 1958b:pl. 3, figs. 14–18. Distribution: Africa (Zaïre).

Hosts: *Scorodophloeus zenkeri*.

References: (ds) Numberg 1966b: 18; Schedl 1933f: 204, 1962k: 1033, 1972f: 255; Strohmeier 1912c: 21, 1914c: 42. (tx) Numberg 1958b: 12; Sampson 1924c: 132; Schedl 1933f: 204, 1950d: 5, 1962k: 1033, 1972f: 255; Strohmeier 1912a: 18, 26, 28, 1912c: 21, 1914c: 42.

kreczmeri Nunberg 1958b: 38. Holotype ♀; Madagascar Contr., Prov. Moromongo, Ortschaft Perinet; Poln. Inst. Poln. Ak. d. Wiss., Warszawa. Distribution: Madagascar.

Notes: (3) Browne 1972c: 104 (described male). References: (ds) Schedl 1972f: 253. (tx) Browne 1972c: 104–105; Numberg 1958b: 38; Schedl 1972f: 253, 1977b: 273.

latacapus Nunberg 1968b: 378. Holotype ♀; Obudu Cattle Ranch, Nigeria; Federal Department of Forest Research in Ibadan, Nigeria. Figures: Numberg 1968b: 377. Distribution: Africa (Nigeria).

References: (ds) Numberg 1968b: 374; Roberts

1968: 193; Schedl 1972f: 253. (tx) Numberg 1968b: 377–378; Schedl 1972f: 253.

latespinatus Nunberg 1969b: 196. Holotype ♂; Nord-Nigerien, Cyl Nyaki; BMNH, London. Distribution: Africa (Congo/ Nigeria).

References: (ds) Browne 1972c: 100; Schedl 1972f: 254. (tx) Numberg 1969b: 196; Schedl 1972f: 254.

longicollis Chapuis 1865: 318. Syntypes ♂; Cap de Bonne-Esperance; apparently at MNHN, Paris.

Figures: Lhoste & Roche 1960:fig. 1, Numberg 1958b:pl. 4, figs. 6–8, 1966b:pl. 1, figs. 10–14, pl. 2, fig. 3.

Distribution: Africa (Angola/ Cameroon/ Congo/ Equatorial Guinea/ Ethiopia/ Fernando Po/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Madagascar/ Mozambique/ Nigeria/ Ruanda/ Senegal/ South Africa/ Tanzania/ Zaïre/ Zambia/ Zimbabwe).

Hosts: Browne 1963a: 248, Schedl 1962k: 1012–1013, 1017 (numerous hosts listed).

Notes: (3) Numberg 1958b: 40 (description quoted), Schedl 1954e: 59, 70 (*angustus*, nomen nudum, synonymy in Schedl 1957d: 149).

References: (hb) Browne 1963a: 248. (ds) Browne 1963a: 248; Ferreira 1965: 1134; Gemminger & Harold 1872: 2701; Kolbe 1897: 283; Lacordaire 1866: 392; Menier 1973a; Schedl 1962k: 1002, 1965g: 22, 1972f: 254; Strohmeier 1912c: 21, 1914c: 42. (tx) Chapuis 1865: 318; Hopkins 1914: 126, 134; Lacordaire 1866: 392–394; Lucas 1920: 494; Numberg 1958b: 40; Sampson 1924: 132; Schedl 1952j: 7, 1953g: 248, 1954e: 59, 1957d: 149, 1960c: 61–62, 1962k: 1002, 1972f: 254; Strohmeier 1912a: 17–18, 27, 1912c: 21, 1914c: 42. (ms) Lucas 1920: 494.

camerunus Strohmeier 1912a: 21. Syntypes, sex?; Kamerun; IPKE, Eberswalde.

Notes: (1) Schedl 1972f: 254 (a subspecies of *longicollis*). (3) Numberg 1958b: 29 (description quoted, redescribed).

References: (ay) Lhoste & Roche 1960; Roche 1960: 2056–2058; Roche & Lhoste 1960: 385–386; Strohmeier 1918: 18. (cc) Roche & Lhoste 1960: 385–386. (hb) Beaver & Loytyniemi 1985b: 119. (ds) Beaver & Loytyniemi 1985b: 119; Browne 1962: 93, 1963: 248, 1975a: 757, 1975b: 395, 1980c: 495, 1984c: 450; Cachan 1957: 15; Gardner 1957a: 29; Mayne & Donis 1950: 332, 1960: 98, 1962: 278; Numberg 1966b: 18–20, 1968b: 374; Roberts 1968: 193, 1969: 116; Schedl 1933f: 204, 1938d: 452, 1959p: 23, 1962h: 63, 1962k: 1009, 1123, 1964f: 620, 1966c: 236, 1967e: 222, 1971e: 3, 1971g: 196, 1972e: 286, 1972f: 254, 1972k: 296, 1977a: 273, 1982: 279; Strohmeier 1911c: 106, 1912c: 21, 1914c: 42; Thompson, G. H. 1963: 55; Wood, S. L. 1957e: 1272. (tx) Lhoste & Roche 1960:fig. 1; Numberg 1958b: 29; Sampson 1924c: 133; Schedl 1933f: 204, 1935e: 174, 1936b: 135, 1937b: 407,

- 1937d: 44, 1941d: 423, 1941e: 156, 1950d: 5, 12, 19, 1953g: 245, 248, 1954d: 875, 1954e: 60–61, 1955d: 270–271, 1955f: 262, 1957b: 152, 1957d: 148–149, 1959p: 23, 1962h: 63, 1962k: 1009–1018, 1964f: 620, 1964k: 220–221, 1965g: 22, 1966c: 235, 1967e: 222, 1972f: 254, 1977b: 273; Strohmeier 1911c: 106, 1912a: 21–24, 27, 1912c: 21, 1914c: 42, 1918: 22.
- camerunus angustiformis* Schedl 1957d: 149. Lectotype ♂; Congo Belge: Yangambi; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 18.
- Notes: (1) The original description is of an individual variation (aberration) within a series that does not meet the criteria of a species or subspecies. Schedl 1972f: 254 (a variety of *longicollis*).
- References: (ds) Ferreira 1965: 1134; Mayne & Donis 1960: 98, 1962: 279; Roberts 1969: 116; Schedl 1954e: 59, 70, 1959p: 23, 1962h: 63, 1962k: 1014–1016, 1964f: 620, 1966c: 235, 1971g: 195, 1972e: 286, 1972f: 254. (tx) Brownie 1962: 93; Schedl 1957d: 149, 1962k: 1016, 1123, 1972f: 254, 1978a: 18.
- camerunus brevestrigatus* Nunberg 1958b: 32. Holotype ♀; Belgisch Congo, Lulonga-Befale; MRCB, Tervuren. Synonymy: Schedl 1972f: 254.
- References: (ds) Schedl 1972f: 254. (tx) Nunberg 1958b: 11, 32; Schedl 1972f: 254.
- camerunus crebrestrigatus* Nunberg 1972c: 133. Holotype, sex?; Bafulngemba, Cameroon Republic; Roberts Collection. Synonymy: Schedl 1972f: 254.
- Notes: (1) This was originally named as a form, not as a species or subspecies; as such, it has no standing in nomenclature.
- References: (tx) Nunberg 1972c: 133; Schedl 1972f: 254.
- longus* Nunberg 1966b: 25. Holotype ♀; Tshuapa: Bokuma; MRCB, Tervuren.
- Figures: Nunberg 1966b: pl. 1, figs. 15–20.
- Distribution: Africa (Zaire).
- References: (ds) Nunberg 1966b: 20; Schedl 1972f: 254. (tx) Nunberg 1966b: 25, 37; Schedl 1972f: 254.
- luebensis* Nunberg 1958: 13. Holotype ♀?; Belgisch Congo: Luebo; MRCB, Tervuren.
- Figures: Nunberg 1966b: pl. 1, figs. 3–4.
- Distribution: Africa (Zaire).
- References: (ds) Nunberg 1966b: 18; Schedl 1972f: 254. (tx) Nunberg 1958a: 13, 1966b: 21, 37; Schedl 1972f: 254.
- major* Strohmeier 1912a: 20. Syntypes 1♂, 1♀; Kamerun; Strohmeier Collection.
- Figures: Nunberg 1958a: pl. 4, figs. 13–19.
- Distribution: Africa (Cameroon/ Nigeria).
- Notes: (3) Nunberg 1958a: 41 (description quoted, redescribed), Schedl 1972f: 254 (treated as a subspecies of *longicollis*).
- References: (ds) Nunberg 1958a: 41, 1966b: 20; Roberts 1968: 193; Schedl 1933f: 204, 1972f: 254; Strohmeier 1912c: 21, 1914c: 42. (tx) Sampson 1924c: 133; Schedl 1933f: 204, 1939b: 354, 1950d: 12, 1953g: 248, 1954e: 60, 86, 1962k: 1004–1005, 1972f: 254; Strohmeier 1912a: 20–21, 27, 1912c: 21, 1914c: 42.
- minutus* Nunberg 1966b: 29. Holotype ♂; Nigerien, Obar; Federal Department of Forest Research in Ibadan, Nigeria.
- Figures: Nunberg 1966b: pl. 2, figs. 14–16.
- Distribution: Africa (Nigeria).
- References: (ds) Nunberg 1966b: 20; Roberts 1968: 193; Schedl 1972f: 254. (tx) Nunberg 1966b: 29; Schedl 1972f: 254.
- mirabilis* Nunberg 1969b: 200. Syntypes, sex?; Nord-Nigerien, Gyel Nyaki; BMNH, London.
- Distribution: Africa (Cameroon/ Nigeria).
- Notes: (3) Nunberg 1972c: 134 (redescribed).
- References: (ds) Schedl 1972f: 254. (tx) Nunberg 1969b: 200, 1972c: 134; Schedl 1972f: 254.
- mkusii* Strohmeier 1912a: 21. Syntypes ♂ ♀; Deutsch Ostafrika (Mkulusumi-Berg), Nguelo; IPKE, Eberswalde.
- Figures: Nunberg 1958a: pl. 4, figs. 1–8.
- Distribution: Africa (Cameroon/ Kenya/ Tanzania/ Zaire), Madagascar.
- Hosts: *Albizzia zygia*, *Pansinstalia* sp.
- Notes: (3) Nunberg 1958a: 43 (description quoted, redescribed), Schedl 1972f: 254 (treated as a subspecies of *longicollis*).
- References: (hb) Strohmeier 1911a: 182. (ds) Beaver & Loytyniemi 1955b: 120; Nunberg 1958a: 43, 1966b: 19, 1968b: 374; Roberts 1968: 193; Schedl 1933f: 204, 1938d: 452, 1962k: 1004, 1966c: 234, 1972f: 254, 1977b: 275; Strohmeier 1911a: 182, 1912c: 21, 1914c: 42. (tx) Schedl 1933f: 204, 1937d: 44, 1962k: 1004–1005, 1966c: 234, 1972f: 254, 1977b: 275; Strohmeier 1911: 182, 1912a: 21, 27, 1912c: 21, 1914c: 42.
- nigeriensis* Nunberg 1966b: 30. Holotype, sex?; Nigeria, Ore FR; Federal Department of Forest Research, Ibadan, Nigeria.
- Figures: Nunberg 1966b: 39.
- Distribution: Africa (Nigeria).
- Hosts: *Coryphanthe pachyceras*.
- References: (ds) Nunberg 1966b: 21; Schedl 1972f: 254. (tx) Nunberg 1966b: 30, 39; Schedl 1972f: 254.
- nitidicollis* Strohmeier 1912a: 23. Syntypes, sex?; Kamerun, Deutsch Ostafrika; IPKE, Eberswalde.
- Figures: Nunberg 1958a: pl. 4, figs. 1–15.
- Distribution: Africa (Angola/ Cameroon/ Congo/ Ghana/ Ivory Coast/ Nigeria/ Tanzania/ Zaire).
- Hosts: *Chlorophora excelsa*, *Enantia clorantha*, *Oxytigma msoo*.
- Notes: (3) Nunberg 1958a: 44 (description

- quoted, redescribed). Schedl 1972f: 253 (a subspecies of *excisus*).
- References: **(hb)** Strohmeier 1911a: 182. **(ds)** Ferreira 1965: 1135; Nunberg 1952: 22, 1958a: 44, 1965b: 22, 1966b: 19, 1968b: 374; Roberts 1968: 193; Schedl 1933f: 205, 1959p: 23, 1962k: 1024, 1972c: 255, 1972h: 253; Strohmeier 1911a: 182, 1912c: 21, 1914c: 42. **(tx)** Nunberg 1952: 22; Schedl 1933f: 205, 1941e: 157, 1950d: 5, 1953g: 245, 1954d: 575, 1954e: 57, 1962k: 1024, 1972f: 253; Strohmeier 1912a: 23–24, 1912c: 21, 1914c: 42.
- penicillatus** Schedl 1965f: 14. Holotype ♀; Uganda, Mganga; BMNH, London.
Distribution: Africa (Uganda).
References: **(ds)** Schedl 1972f: 255. **(tx)** Schedl 1965f: 14, 1972f: 255, 1978a: 52.
- piceus** Strohmeier 1912a: 24. Holotype ♂; Kamerun; IPKE, Eberswalde.
Figures: Nunberg 1958b: pl. 5, figs. 14–16.
Distribution: Africa (Cameroon/ Congo/ Ivory Coast/ Nigeria/ Sierra Leone/ Zaïre).
References: **(ds)** Browne 1965a: 205, 1972c: 100; Nunberg 1952: 22, 1965b: 22, 1966b: 21; Roberts 1968: 193; Schedl 1972f: 253; Strohmeier 1912c: 21, 1914c: 42. **(tx)** Nunberg 1952: 22, 1958b: 46, 1964: 22; Sampson 1924c: 132; Schedl 1962k: 1026, 1972f: 253; Strohmeier 1912a: 24–27, 1912c: 21, 1914c: 42.
- pseudocamerunus** Nunberg 1958b: 47. Syntypes 2 ♀; Kasenya [Congo], N. Rhodesia; 1 in MRCB, Tervuren, 1 in IZW, Warsaw.
Distribution: Africa (South Africa/ Zaïre/ Zambia/ Zimbabwe).
References: **(ds)** Beaver & Loytyniemi 1985b: 120; Nunberg 1966b: 19; Schedl 1972f: 255. **(tx)** Nunberg 1958b: 47; Schedl 1972f: 255.
- pseudogracilis** Nunberg 1969b: 198. Holotype ♀; Nord-Nigerien, Guel Nyaki; BMNH, London.
Distribution: Africa (Nigeria).
References: **(ds)** Schedl 1972f: 255. **(tx)** Nunberg 1969b: 198; Schedl 1972f: 255.
- pseudomajor** Schedl 1954e: 86. Lectotype ♂; Mpraeso [Gold Coast]; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 58.
Figures: Nunberg 1966b: pl. 2, figs. 18–21.
Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Nigeria).
Hosts: *Ficus exasperata*, *Sterculia oblonga*.
Notes: (3) Nunberg 1958b: 48 (description quoted), 1966b: 33 (described female), Schedl 1972f: 254 (a subspecies of *longicollis*).
References: **(ds)** Nunberg 1966b: 21; Roberts 1968: 193; Schedl 1962k: 1003, 1972e: 286, 1972f: 254; Thompson, G. H. 1963: 55. **(tx)** Browne 1970: 577; Nunberg 1958b: 48, 1966b: 21, 33–34, 39; Schedl 1954e: 60–63, 86, 1962k: 1003, 1972e: 277–294, 1972f: 254, 1978a: 58.
- pseudoseverini** Nunberg 1958b: 15. Holotype ♂; [Congo] Sankuru; Lodja; MRCB, Tervuren.
Figures: Nunberg 1958b: pl. 6, figs. 1–8.
Distribution: Africa (Zaire).
References: **(ds)** Schedl 1972f: 255. **(tx)** Nunberg 1958b: 15; Schedl 1972f: 255.
- quinquespinatus** Nunberg 1958b: 48. Holotype ♂; Congo Belge (Katanga: Lufudizi); MRCB, Tervuren.
Figures: Nunberg 1958b: pl. 6, figs. 9–14.
Distribution: Africa (Zaire).
References: **(ds)** Schedl 1972f: 255. **(tx)** Nunberg 1958b: 48; Schedl 1972f: 255.
- robertsi** Nunberg 1972c: 134. Holotype ♂; Nigeria; Nindam Forest Reserve; Roberts Collection.
Distribution: Africa (Nigeria).
References: **(tx)** Nunberg 1972c: 134.
- severini** Strohmeier 1912a: 25. Syntypes, sex?; Kinchassa, Kongo; IPKE, Eberswalde.
Figures: Nunberg 1958b: pl. 6, figs. 15–22.
Distribution: Africa (Zaire).
Notes: (3) Nunberg 1958b: 18 (description quoted, redescribed), Nunberg 1959b: 127 (reported also as a fossil).
References: **(ds)** Nunberg 1959b: 127, 1966b: 18; Schedl 1933f: 204, 1962k: 1032, 1966c: 235, 1972: 255; Spahr 1981: 79; Strohmeier 1912c: 21, 1914c: 42. **(tx)** Nunberg 1958b: 18, 1959b: 127; Sampson 1924c: 132; Schedl 1933f: 204, 1941d: 422, 1962k: 1032, 1972f: 255; Strohmeier 1912a: 25–28, 1912c: 21, 1914c: 42.
- signatus** Strohmeier 1912a: 25. Syntypes ♂ ♀; Kamerun; IPKE, Eberswalde.
Figures: Nunberg 1958b: pl. 7, figs. 1–5.
Distribution: Africa (Cameroon/ Zaïre).
Notes: (3) Nunberg 1958b: 50 (description quoted, redescribed).
References: **(ds)** Nunberg 1966b: 19; Schedl 1972f: 255; Strohmeier 1912c: 21, 1914c: 42. **(tx)** Nunberg 1958b: 48–50; Schedl 1962k: 1035, 1972f: 255; Strohmeier 1912a: 25, 27, 1912c: 21, 1914c: 42.
- similis** Strohmeier 1912a: 22. Syntypes ♂ ♀; Kamerun; 3♂ in Strohmeier Collection.
Figures: Nunberg 1958b: pl. 7, figs. 6–8, 1966b: pl. 1, figs. 9, 11–13.
Distribution: Africa (Cameroon/ Ghana/ Zaïre).
Hosts: *Bolboceranius bicolor*, *Isoloma brunclii*, *Scorodopliocus zenkeri*.
Notes: (3) Nunberg 1958b: 51 (description quoted), 1966: 24 (described female), Schedl 1972f: 253 (a subspecies of *excisus*).
References: **(ds)** Browne 1980b: 382; Nunberg 1966b: 19; Schedl 1962k: 1029, 1966c: 234, 1972e: 286, 1972f: 253; Strohmeier 1912c: 21, 1914c: 42. **(tx)** Nunberg 1958b: 51, 1966b: 19, 24–25, 37; Schedl 1935e: 174, 1962k: 1029, 1972f: 253; Strohmeier 1912a: 22–27, 1912c: 21, 1914c: 42.

subcircularis Schedl 1941d: 421. Syntypes 2♂, 1♀; Congo Belge: Stanleyville; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Notes: (3) Nunberg 1958b: 20 (description quoted).

References: (ds) Schedl 1962k: 1031, 1972f: 255.

(tx) Nunberg 1958b: 20; Schedl 1941d: 421, 1962k: 1031, 1966c: 239, 1972f: 255, 1978a: 70.

subrobustus Schedl 1957d: 149. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Nunberg 1958b: pl. 7, figs. 17–20.

Distribution: Africa (Ghana/ Nigeria/ Uganda/ Zaire).

Hosts: *Afrostryax lepidophyllus*, *Anonidium mannii*, *Beilschmiedia gilbertii*, *Macaranga* cf. *laucifolia*, *Maesopsis eminii*, *Mitragyne* sp., *Scorodophloeus zenkeri*.

Notes: (3) Schedl 1972f: 253 (treated as a subspecies of *excisus*).

References: (ds) Mayne & Donis 1960: 98, 1962: 280; Schedl 1962k: 1027, 1972e: 286, 1972f: 253.

(tx) Nunberg 1958b: 51; Schedl 1957d: 149, 1962k: 1027, 1972f: 253, 1978a: 71.

substriatus Strohmeier 1912a: 23. Syntypes ♂ ♀; Kamerun; IPKE, Eberswalde.

Figures: Nunberg 1958b: pl. 7, figs. 9–16.

Distribution: Africa (Cameroon/ Fernando Po/ Ghana/ Nigeria/ Togo/ Zaire).

Hosts: *Albizia zygia*, *Drypetes gossweileri*, *Napoleana imperialis*, *Pancovia harnsiana*, *P. laurentii*, *Polyalthia suaveolens*, *Synsepalum subcordatum*, *Ternstroemia ivorensis*, *Xylopia aethiopica*.

Notes: (3) Nunberg 1958b: 52 (description quoted), Schedl 1972f: 253 (a subspecies of *excisus*).

References: (hb) Thompson, G. H. 1963: 55. (ds) Mayne & Donis 1960: 98, 1962: 280; Nunberg 1966b: 21, 1968b: 374; Roberts 1968: 374; Schedl 1933f: 205, 1972e: 286, 1972f: 253; Strohmeier 1912c: 21, 1914c: 42; Thompson, G. H. 1963: 55. (tx) Nunberg 1958b: 52; Schedl 1933f: 205, 1937d: 44, 1954e: 48, 60, 65, 71, 1962k: 1026, 1972f: 253; Strohmeier 1912a: 20, 23, 27, 1912c: 21, 1914c: 42.

suturalis Browne 1972c: 101. Holotype ♂; Central African Republic: Bambari; MRCB, Tervuren.

Figures: Browne 1972c: 102.

Distribution: Africa (Central African Republic).

References: (tx) Browne 1972c: 101–102.

titschaki Schedl 1935e: 154. Lectotype ♂; Deutsch E. Africa: Amani; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 74.

Figures: Nunberg 1958b: pl. 7, fig. 24.

Distribution: Africa (Tanzania).

Notes: (3) Nunberg 1958b: 52 (description quoted), Schedl 1957d: 147 (described female).

References: (ds) Schedl 1962k: 1034, 1972f: 255. (tx) Nunberg 1958b: 52, 1966b: 25; Schedl 1935e:

154, 1937d: 44, 1939b: 390, 403, 1957d: 147–148, 1962k: 1034, 1972f: 255, 1978a: 74.

triquetrus Schedl 1941d: 423. Holotype ♂; Congo Belge: Likimi, Bondia, Monveda; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Notes: (3) Nunberg 1958b: 53 (description quoted), Schedl 1972f: 253 (a subspecies of *excisus*).

References: (ds) Schedl 1972f: 253. (tx) Nunberg 1958b: 53; Schedl 1941d: 423, 1972f: 253, 1978a: 75.

truncatellus Schedl 1957d: 148. Holotype ♂; Congo Belge: Kinchassa; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

References: (ds) Schedl 1972f: 255. (tx) Schedl 1957d: 148, 1962k: 1036, 1972f: 255, 1978a: 75.

truncatiformis Schedl 1941d: 422. Holotype ♂; Congo; Schedl Collection in NHMW, Wien.

Distribution: Africa (Senegal/ Zaire).

Notes: (3) Nunberg 1958b: 20 (description quoted). A note in the Schedl Collection says this is a synonym of *collaticus* [confirmation needed].

References: (ds) Schedl 1972f: 255. (tx) Nunberg 1958b: 20; Schedl 1941d: 422, 1950d: 5, 1962k: 1032, 1972f: 255, 1978a: 75.

planifrons Schedl 1941d: 423. Holotype ♀; Casamance [Senegal]; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 255.

Notes: (3) Nunberg 1958b: 15 (description quoted).

References: (tx) Nunberg 1958b: 15; Schedl 1941d: 422–423, 1972f: 255, 1978a: 55.

truncatipennis Schedl 1966c: 239. Holotype ♂;

Republique du Congo, Dimonika; MNHN, Paris.

Distribution: Africa (Congo/ Zaire).

References: (ds) Schedl 1972f: 255. (tx) Schedl 1966c: 239, 1972f: 255.

truncatus Schedl 1936b: 135. Lectotype ♂; Belgisch Kongo, Likimi; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 76.

Figures: Nunberg 1958b: pl. 7, figs. 21–23.

Distribution: Africa (Equatorial Guinea/ Zaire).

Notes: (3) Nunberg 1958b: 21 (description quoted).

References: (ds) Schedl 1962k: 1031, 1971f: 196,

1972f: 255. (tx) Nunberg 1958b: 21; Schedl 1936b: 135–136, 1950d: 5, 1962k: 1031, 1037,

1972f: 255, 1978a: 76.

Genus *Chaetastus* Nunberg

CHAETASTUS NUNBERG 1953: 44. Type-species: *Symmerus tuberculatus* Chapuis, automatic.

Symmerus Chapuis 1865: 42, 319. Type-species: *Symmerus tuberculatus* Chapuis, monobasic, preoccupied by Walker 1845.

References: (tx) Browne 1972a: 179; Chapuis 1865: 319–321; Hopkins 1914: 136, 1915c: 227; Lacordaire 1866: 389, 395; Nunberg 1953: 44; Sampson 1924: 133; Schedl 1933f:

203, 1936e: 62, 1937d: 44, 1939b: 390–401, 1956d: 145; Strohmeier 1911d: 218, 1912c: 21, 1914: 12–44, 1920: 1–46.

Keys: Browne 1971: 11.

References: (**hb**) Browne 1971c: 8. (**ds**) Browne 1971c: 8. (**tx**) Browne 1971c: 8, 1972a: 179; Numborg 1953: 44; Schedl 1960c: 62, 1962k: 1039.

diversifrons Browne 1971b: 19. Holotype ♂; Uganda: Ruwenzori Forest; BMNH, London.

Figures: Browne 1971b: 13.

Distribution: Africa (Uganda).

Hosts: *Mitragyna stipulosa*, *Olea hochstetteri*, *Pygmean africanum*.

References: (**ds**) Schedl 1972f: 127. (**tx**) Browne 1971b: 13–19; Schedl 1972f: 127.

fradei (Tordo) 1966: 16 (*Symmerus*). Syntypes?; Silva Porto, Angola; not seen.

Figures: Tordo 1966: 16.

Distribution: Africa (Angola).

References: (**tx**) Tordo 1966: 16.

montanus Schedl 1957d: 145. Holotype ♂; Congo Belge; Kivu, Hembe-Bitale; MRCB, Tervuren.

Distribution: Africa (Brundi/ Kenya/ Nigeria/ Ruanda/ Tanzania/ Uganda/ Zaire).

Hosts: Browne 1971: 17 and Schedl 1962k: 1051–1052 (many hosts listed).

Notes: (3) Browne 1971c: 16 (redescribed).

References: (**bv**) Jones 1960c: 231. (**cn**) Curry 1958b: 128–132. (**ds**) Browne 1971c: 16; Gardner 1957a: 30, 1957b: 204; Jones 1960: 3; Mayne & Donis 1962: 288; Roberts 1960: 241, 1962: 1050, 1968: 193; Schedl 1962k: 1050, 1972f: 128. (**tx**) Browne 1971c: 16; Powell, W. 1987: 27; Schedl 1955f: 261, 1955i: 213–214, 1957d: 145–146, 1962h, 1962k: 1050, 1972f: 128, 1978a: 45.

persimilis (Schedl) 1933f: 203 (*Symmerus*). Lectotype ♂; Haut-Uele; Adra, Congo-Kinshasa; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 54.

Figures: Schedl 1962k: 1048 (male).

Distribution: Africa (Cameroon/ Congo/ Fernando Po Island/ Gabon/ Nigeria/ Ruanda/ Sao Tome Island/ Zaire).

Hosts: *Combretodendrum macrophyllum*, *Diospyros crassiflora*, *Garcinia polyantha*, *Pycnanthus angolensis*, *P. kombo*, *Staudtia stipitata*, *Strombosia glaucescens*, *Tridesmostemon classensii*.

References: (**cc**) Schedl 1962k: 1048. (**ds**) Browne 1971c: 15, 1980b: 382; Mayne & Donis 1962: 288; Roberts 1968: 193, 1969: 103; Schedl 1933f: 203, 1962k: 1123, 1972f: 128. (**tx**) Browne 1971c: 15; Schedl 1933f: 203–204, 1936e: 62, 1952i: 60–62, 1955f: 261, 1957d: 146, 1962h: 60–61, 1962k: 1048, 1972f: 128, 1978a: 54.

striatus Browne 1986b: 334. Holotype ♂; Cameroon: Donala to Nagoya (Japan), imported; BMNH, London.

Distribution: Africa (Cameroon).

Hosts: *Paraberlina bifoliolata* log.

References: (**tx**) Browne 1986b: 334.

subalpinus (Schedl) 1957d: 146 (*Symmerus*). Holotype ♂; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Anthocleista nobilis*.

Notes: (3) Browne 1971: 18 (redescribed).

References: (**cc**) Schedl 1962k: 1050. (**ds**) Browne 1971c: 18. (**tx**) Browne 1971c: 18; Schedl 1957d: 145–146, 1962k: 1050, 1972f: 128, 1978a: 69.

tuberculatus (Chapuis) 1865: 321 (*Symmerus*). Holotype ♂ (Schedl 1960: 62); Guinee, Coll. de M. Thomson; IRSNB, Brussels.

Figures: Browne 1971c: 9, Schedl 1962k: 1040 (male).

Distribution: Africa (Angola/ Cameroon/ Central African Republic/ Congo/ Equatorial Guinea/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sao Tome Island/ Sierra Leone/ Tanzania/ Togo/ Uganda/ Zaire/ Zambia).

Hosts: Browne 1971c: 14 and Schedl 1962k: 1043–1046 (many hosts listed).

References: (**bv**) Jones 1960c: 231; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Meixner 1937: 1218. (**cn**) Duffy 1956: 198; Thompson, G. H. 1959b: 420. (**cc**) Roberts 1969a; Schedl 1962k: 1040. (**hb**) Anonymous 1959i: 18; Beaver & Loytyniemi 1985b: 114; Browne 1963a: 248; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Schedl 1962k: 1040. (**ds**) Beaver & Loeytyniemi 1985b: 114; Browne 1963a: 248, 1971c: 12, 1972c: 99, 1980a: 373, 1980c: 485; Cachan 1957: 15; Cola 1971, 1973; Ferreira 1965: 1135; Gardner 1957a: 30, 1957b: 205; Gemminger & Harold 1872: 2702; Jones 1960: 3; Jones, Roberts, & Baker 1959: 13–55; Lacordaire 1866: 395; Mayne & Donis 1960: 100, 1962: 289; Menier 1973a; Roberts 1960b: 35–37, 1960e, 1960f, 1968: 193, 1969: 103–104; Schedl 1933f: 203, 1959p: 23, 1962h: 61, 1962k: 1040, 1964f: 620, 1966c: 235, 1967e: 222, 1971e: 4, 1971g: 194, 1972e: 286, 1972f: 129; Strohmeier 1911f: 204, 1912c: 21; Thompson, G. H. 1959b: 420, 1963: 35; Webb & Jones 1957: 36, 41. (**tx**) Browne 1971c: 12, 19, 1972a: 179; Chapuis 1865: 321–322; Hopkins 1914: 130, 136; Lacordaire 1866: 395; Lucas 1920: 621; Meixner 1937: 1218; Powell, W. 1987: 27; Sampson 1924c: 133; Schedl 1933f: 203–204, 1936e: 62, 1937b: 407, 1937d: 44, 1939b: 391, 1941d: 381, 1941e: 156, 1950d: 5, 1953g: 245, 1954d: 874, 1954e: 48, 60–71, 1955d: 271, 1955f: 261, 1960c: 62, 1962k: 1040, 1972f: 129; Strohmeier 1911f: 203–204, 1911g: 232, 1912c: 21, 1914: 5, 7, 44, 1920: 16. (**ms**) Lucas 1920: 621.

tuberculatus robustus Schedl 1952i: 62 (*Symmerus*). Holotype ♂; Congo Belge, Yangambi; MRCB, Tervuren. Synonymy: Browne 1971c: 12.

References: (**ds**) Mayne & Donis 1960: 100,

1962: 290; Schedl 1962k: 1047, 1972f: 129. (tx) Browne 1971c: 12; Schedl 1952i: 62, 1962k: 1047, 1972f: 129, 1978a: 76.

Genus *Cenocephalus* Chapuis

CENOCEPHALUS CHAPUIS 1865: 325. Type-species: *Cenocephalus thoracicus* Chapuis, monobasic.

Notes: (1) This genus is endemic to tropical America.

References: (tx) Chapuis 1865: 325, 327; Nunberg 1963c: 109; Schedl 1939b: 401, 1960c: 62; Strohmeyer 1911c: 105, 107.

***epistomalis* Wood** 1966a: 47. Holotype ♀; Mile 10 on Bartica-Potaro road, British Guiana; BMNH, London.

Figures: Wood 1966a: 47.

Distribution: South America (Guyana).

References: (ds) Equihua & Atkinson 1987: 18; Schedl 1972f: 130. (tx) Equihua & Atkinson 1987: 18; Schedl 1972f: 130; Wood, S. L. 1966a: 47.

***flexibilis* (Schedl)** 1936c: 249 (*Tesserocranulus*). Holotype ♀; Brazil or Guayana; Schedl Collection in NHMW, Wien.

Distribution: South America (Cayenne/ Suriname).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 123. (tx) Reichardt 1964: 149; Schedl 1936c: 249, 1972f: 123.

° ***hurdi* Schedl** 1962g: 1037. Holotype ♂; Mexican amber, Miocene, Simojovel, U.C.M.P. no. 12976, loc. B-7456 (late Oligocene or early Miocene); University of California Museum of Paleontology, Berkeley.

Figures: Schedl 1962g: 1036.

Distribution: North America (fossil in Mexican amber).

References: (cn) Hosking 1989. (ds) Keilbach 1982: 256; Schedl 1972f: 130; Spahr 1981: 79. (tx) Schedl 1962g: 1036–1037, 1972f: 130. (ms) Keilbach 1982: 256.

***lalolaensis* Nunberg** 1963c: 107. Holotype ♀; Finca La Lola, Limon Prov., Costa Rica; University of Wisconsin Collection, Madison.

Distribution: North America (Costa Rica), South America (Brazil).

Notes: (1) This female remains unidentifiable at the present time.

References: (ds) Equihua & Atkinson 1987: 791; Schedl 1970e: 88. (tx) Nunberg 1963c: 107; Schedl 1972f: 79.

***ornatus* (Schedl)** 1936c: 250 (*Tesserocranulus*). Holotype ♀; Brazil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil/ Cayenne).

References: (ds) Blackwelder 1947: 791. (tx) Schedl 1936c: 250, 1972f: 123, 1978a: 50.

***pulchellus* Schedl** 1935e: 152. Holotype ♀; Guyana; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil/ Guyana/ Suriname).

References: (ds) Blackwelder 1947: 791. (tx) Schedl 1935e: 152, 1972f: 130, 1978a: 59; Wood, S. L. 1966a: 47.

***pulchellus minor* Schedl** 1961i: 233. Holotype ♀; Surinam, Moengo, Boven, Cottica R.; Cornell University Collection, Ithaca, New York.

Notes: (3) Schedl 1970e: 110, 1972g: 83–84 (described male; listed as a species).

References: (ds) Schedl 1960a: 79, 1970e: 88, 1970f: 582, 1972f: 130, 1972g: 51. (tx) Schedl 1961i: 233, 1970e: 110, 1972f: 130, 1972g: 83–84, 1978a: 59.

***pusillus* Schedl** 1936c: 250. Lectotype ♂; French Guyana: Nouveau Chantier, St. Jean du Maroni; Schedl Collection in NHMW, Wien.

Distribution: South America (Cayenne).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 130. (tx) Schedl 1935b: 400, 1936c: 250, 1952d: 365, 1972f: 130, 1978a: 59.

° ***quadrilobus* Schedl** 1962g: 1037. Holotype ♂; Mexican amber, late Oligocene to Miocene, Simojovel, Chiapas, U.C.M.P. no. 12877, loc. B-4117; University of California Museum of Paleontology, Berkeley.

Distribution: North America (fossil in Mexican amber).

References: (ds) Keilbach 1982: 256. (tx) Schedl 1962g: 1035–1037, 1972f: 130. (ms) Keilbach 1982: 256.

° ***rhinoceroide* (Schawaller)** 1981: 10 (*Mitosoma*). Holotype, sex?; Dominikanischem Bernstein (SMNS Inv. Nr. Do. 119–k-1); Staatlichen Museum für Naturkunde, Stuttgart.

Figures: Schawaller 1981: 11 (adult).

Distribution: Antilles Islands (fossil in Dominican Republic amber in Hispanola).

Notes: (3) Three species of *Cenocephalus* have been found in Dominican amber; all three appear to be the three named previously by Schedl from Mexican amber; consequently, the Schawaller type should be re-examined; synonymy is probable [SLW].

References: (ds) Spahr 1981: 79. (tx) Schawaller 1981: 10.

***robustus* Schedl** 1966f: 128. Holotype ♂; Colombian; Schedl Collection in NHMW, Wien.

Distribution: South America (Colombia).

References: (ds) Schedl 1966f: 79, 128, 1972f: 130.

***robustum* Schedl** 1972f: 130. Holotype ♂; Colombian; Schedl Collection in NHMW, Wien.

Notes: (1) This is an unneeded replacement name for *robustus*.

References: (tx) Schedl 1972f: 130.

***rugicollis* Schedl** 1952d: 365. Holotype ♂; Peru; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil/ Peru).

Notes: *Eugenia* sp.

References: (ds) Schedl 1970e: 88, 1972f: 130.

- (**tx**) Reichardt 1964b: 59; Schedl 1952d: 365–366, 1972f: 130, 1978a: 63.
- ° **succinica** Schedl 1962g: 1036. Holotype ♀; Mexican amber, late Oligocene to early Miocene, 12 km SE Sinojovel (U.C.M.P. no. 12718 loc. B-4116, Las Cruces landslide); University of California Museum of Paleontology, Berkeley. Figures: Schedl 1962g: 1036. Distribution: North America (fossil in Mexican amber). References: (**ds**) Keilbach 1982: 256; Spahr 1981: 79. (**tx**) Schedl 1962g: 1036, 1972f: 130. (**ms**) Keilbach 1982: 256.
- thoracicus** Chapuis 1865: 327. Syntypes; Bresil: Rio-Janeiro, Ste-Catherine; BMN11, London. Distribution: South America (Brazil). References: (**ds**) Blackvelder 1947: 791; Gemminger & Harold 1872: 2702; Lacordaire 1866: 397; Santoro 1957b: 26; Schedl 1933c: 163–164, 1967d: 5; Strohmeier 1912c: 22, 1914c: 45. (**tx**) Chapuis 1865: 327–329; Hopkins 1914: 117; Lacordaire 1866: 397; Schedl 1935e: 152, 1937d: 44, 1972f: 130; Strohmeier 1912c: 22, 1914c: 45.
- Genus *Mitosoma* Chapuis
- MITOSOMA** CHAPUIS 1865: 43, 322. Type-species: *Mitosoma crenulata* Chapuis, monobasic.
- Platypicerus** Numberg 1953: 46. Type-species: *Platypicerus hamatus* Numberg, monobasic. Synonymy: Schedl 1972f: 124. References: (**tx**) Numberg 1953: 44, 1963d: 357; Schedl 1957b: 159, 1972f: 124.
- Coccephalonus** Schedl 1965c: 84. Type-species: *Coccephalophonus sulcipennis* Schedl, original designation. Synonymy: Schedl 1972f: 124. References: (**tx**) Schedl 1965c: 84, 1972f: 124. Notes: (3) This genus is endemic to Madagascar. References: (**tx**) Chapuis 1865: 324; Hopkins 1914: 125, 133, 1915c: 227; Lacordaire 1866: 395; Nunberg 1953: 44–46; Schaufuss 1897: 223, 1905: 19; Schedl 1935n: 633, 1936e: 62, 1939b: 401, 1960c: 62, 1961e: 160, 1962k: 1035, 1972f: 124–130, 1977b: 256–267; Strohmeier 1911c: 104, 1911d: 218, 1912c: 22, 1914c: 13–19, 44–46, 1918: 1–46.
- accuratum** Schaufuss 1897: 224. Holotype ♂; Madagascar (Sikora num. 91.); Hamburg Museum, lost. Distribution: Madagascar. Hosts: *Chrysophyllum boivinianum*. Notes: (3) Schedl 1965c: 79 (described female). References: (**ds**) Alluaud 1900: 441; Schedl 1972f: 127, 1977b: 258; Strohmeier 1912c: 22. (**tx**) Schaufuss 1897a: 223–224; Schedl 1936e: 62, 1961e: 162, 1965c: 79, 1972f: 127, 1977b: 257–258, 1978a: 9; Strohmeier 1912c: 22, 1914c: 46, 1918: 16, 21.
- ampliatum** Schedl 1965c: 80. Holotype ♂; Madagascar: Sambava; Schedl Collection in NHMW, Wien. Distribution: Madagascar. References: (**ds**) Schedl 1972f: 127. (**tx**) Schedl 1972f: 127, 1965c: 80, 1978a: 10.
- chapuisi** Strohmeier 1911g: 233. Syntypes ♂ ♀; Madagascar: IRSNB, Brussels, NHR, Stockholm, IPKE, Eberswalde. Distribution: Madagascar. References: (**cc**) Fuller 1958: 100. (**ds**) Schedl 1972f: 127; Strohmeier 1912c: 22, 1914c: 46. (**tx**) Schedl 1972f: 127; Strohmeier 1911c: 104–105, 1911g: 233, 1912c: 22, 1914c: 46.
- crenulatum** Chapuis 1865: 324. Syntypes 2 ♂; Madagascar; MNHN, Paris. Distribution: Madagascar. Hosts: *Canarium boivin*, *Ficus sorocoides* var. *macrophlebia*. Notes: (3) Schedl 1961e: 158 (described female). References: (**ay**) Menier 1976: 348. (**bv**) Menier 1976: 348. (**hb**) Schedl 1977b: 258. (**ds**) Alluaud 1900: 441; Fairmaire 1892b; Gemminger & Harold 1872: 2702; Lacordaire 1866: 396; Schedl 1970d: 234, 1972f: 127, 1977b: 258; Strohmeier 1912c: 22, 1914c: 46. (**tx**) Chapuis 1865: 324; Fairmaire 1892b; Hopkins 1914: 125, 133; Lacordaire 1866: 396; Lucas 1920: 421; Schaufuss 1890: 38, 1897a: 224; Schedl 1960c: 62, 1961e: 158, 1970d: 234, 1972f: 124, 127, 1977b: 258, 1978a: 22; Strohmeier 1912c: 22, 1914c: 46. (**ms**) Lucas 1920: 421.
- dispar** Schedl 1961e: 160. Madagascar, Montagne d'Ambre; IRSM, Madagascar. Distribution: Madagascar. Hosts: *Ficus sorocoides* var. *macrophlebia*. References: (**ds**) Schedl 1972f: 127, 1977b: 260. (**tx**) Schedl 1961e: 160, 1972f: 127, 1977b: 260, 1978a: 25.
- excisum** Schaufuss 1897a: 224. Holotype ♂; Madagascar, Antananarive (Schedl 1977b: 260); Hamburg Museum, lost. Distribution: Madagascar. Hosts: *Aphloia theaeformis*, *Chrysophyllum boivinianum*, *Dracaena* sp., *Elaeocarpus* sp., *Ficus sorocoides* var. *macrophlebia*, *Macphersonia* cf. *madagascarensis*, *Ocotea* sp., *Pinus insularis*, *P. patula*, *Pittosporum* sp., *Terminalia* sp., *Tricocodendron acuminatum*. Notes: (3) Schedl 1961e: 162 (described female). References: (**ds**) Alluaud 1900: 441; Schedl 1970d: 234, 1972f: 128, 1977b: 260; Strohmeier 1912c: 22, 1914c: 46. (**tx**) Schaufuss 1897a: 223–224; Schedl 1961e: 162, 165, 1970d: 234, 1972f: 128, 1977b: 260, 1978a: 29; Strohmeier 1912c: 22, 1914c: 46.
- hamatus** (Numberg) 1953: 49 (*Platypicerus*). Holotype ♀; Central Madagascar, district Moramanga, Perinet; IZW, Warsaw.

Distribution: Madagascar.

Notes: (3) Schedl 1971e: 17 (described female).

References: (ds) Schedl 1972f: 128. (tx) Nunberg 1953: 49–51, 1963d: 357; Schedl 1952b: 159, 1957b: 159, 1971e: 17, 1972f: 128, 1978a: 34.

lobatum Schedl 1961e: 163. Holotype ♂; Madagascar, Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Cedrela* sp., *Dalbergia pterocarpifolia*, *Foetidia chusioidea*, *Garcinia verrucosa*, *Leptolaena multiflora* var. *cuspidata*, *Symphonia* sp.

References: (ds) Schedl 1970d: 235, 1972f: 128, 1977b: 261. (tx) Schedl 1961e: 163, 1970d: 235, 1972f: 128, 1977b: 261, 1978a: 41.

mirabile Nunberg 1967a: 332. Holotype ♀; Madagascar (foret de Fito); MRCB, Tervuren.

Figures: Nunberg 1967a: 339.

Distribution: Madagascar.

References: (tx) Nunberg 1967a: 332, 339.

nigrum Schaufuss 1893: 621. Holotype ♂; Madagascar; Hamburg Museum, lost.

Distribution: Madagascar.

References: (ds) Alluand 1900: 441; Fairmaire 1892b; Schedl 1972f: 128, 1977b: 262; Strohmeier 1912c: 22, 1914c: 46. (tx) Fairmaire 1892b; Schaufuss 1890: 621, 1897a: 223, 1905: 19; Schedl 1972f: 128, 1977b: 262; Strohmeier 1912c: 22, 1914c: 46.

obconiceps Schedl 1970d: 238. Holotype ♂; Madagascar, Perinet; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

Hosts: *Cedrela* sp., *Commiphora* sp., *Dalbergia* sp., *Erythroxylon* sp., *Pinus insularis*, *Ravensara* sp., *Rhodobaena bakeriana*.

References: (ds) Schedl 1970d: 235, 1972f: 128, 1977b: 263. (tx) Schedl 1965c: 83, 1970d: 235, 238, 1972f: 128, 1977b: 263, 1978b: 48.

obliquatum Schedl 1961e: 164. Holotype ♂; Madagascar, Perinet, Betsatsakry, Antaniditra pres Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Cedrela* sp., *Chrysophyllum boivinianum*, *Dalbergia* sp., *Erythroxylon* sp., *Eucalyptus* sp., *Ocotea* sp., *Pinus insularis*, *Rhodobaena bakeriana*, *Symphonia* sp., *Trichila* sp., *Uapaca* sp.

References: (ds) Schedl 1970d: 235, 1972f: 128, 1977b: 264. (tx) Schedl 1961e: 164, 1965c: 80, 1972f: 128, 1977b: 264, 1978a: 48.

obtusum Schedl 1961e: 165. Holotype ♂; Madagascar, Perinet, Betsatsakry pres Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Cussonia vantsilana*, *Dalbergia* sp., *Leptolaena multiflora*, *Symphonia* sp.

References: (ds) Schedl 1970d: 235, 1972f: 128, 1977b: 265. (tx) Schedl 1961e: 165, 1972f: 128, 1977b: 265, 1978a: 49.

octospinosum Schedl 1970d: 239. Holotype ♂; Madagascar, Bemampidy; MNHN, Paris.

Distribution: Madagascar.

Hosts: *Mauloutchia* sp.

References: (ds) Schedl 1972f: 128, 1977b: 265. (tx) Schedl 1970d: 239, 1972f: 128, 1977b: 265, 1978a: 49.

odonticeps Schedl 1961e: 159. Holotype ♂; Madagascar, Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Ravensara perillei*, *R. polyneura*, *Ravenala madagascarensis*, *Syderoxylon* sp., *Trichilia* sp.

References: (ds) Schedl 1972f: 128, 1977b: 266. (tx) Schedl 1961e: 159, 1970d: 235, 1972f: 128, 1977b: 266, 1978a: 49.

paulianum Schedl 1950h: 111. Lectotype ♂; Madagascar, Manjakatampo; Schedl Collection in NHMW, Wien.

Figures: Schedl 1962k: 1038 (male).

Distribution: Madagascar, South Africa.

Hosts: *Chrysophyllum boivinianum*, *Elacocarpus* sp., *Ficus* sp., *Mespilodaphne* sp., *Rhodobaena bakeriana*, *Symphonia* sp., *Tricocodendron acuminatum*.

Notes: (3) Schedl 1961e: 166 (described female).
References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (ds) Schedl 1972f: 128, 1977b: 267. (tx) Schedl 1950h: 111, 1961e: 166, 1962k: 1038, 1972f: 128, 1977b: 267, 1978a: 52.

planum Schaufuss 1897a: 224. Holotype ♂; Madagascar (Sikora num. 91.); Hamburg Museum, lost.

Distribution: Madagascar.

References: (ds) Alluand 1900: 441; Schedl 1972f: 128, 1977b: 268; Strohmeier 1912c: 22, 1914c: 46. (tx) Schaufuss 1897a: 223–224, 1905: 19; Schedl 1972f: 128, 1977b: 268; Strohmeier 1912c: 22, 1914c: 46.

robustum Schaufuss 1890: 623. Holotype ♂; Madagascar; Hamburg Museum, lost.

Distribution: Madagascar.

References: (ds) Alluand 1900: 441; Fairmaire 1892b; Schedl 1972f: 128, 1977b: 268; Strohmeier 1912c: 22, 1914c: 46. (tx) Fairmaire 1892b; Schaufuss 1890: 623, 1897a: 223; Schedl 1972f: 128, 1977b: 268; Strohmeier 1912c: 22, 1914c: 46.

rugosum Schaufuss 1905: 11. Holotype ♂; Madagascar; Hamburg Museum, lost.

Distribution: Madagascar.

References: (ds) Schedl 1972f: 128, 1977b: 268. (tx) Schaufuss 1905: 11; Schedl 1972f: 128, 1977b: 268.

sexspinosum Schedl 1961e: 160. Holotype ♂; Madagascar, Montagne d'Ambre, Plateau 1100 m; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Canarium* sp., *Didymeles* sp., *Ekbergia suavis*, *Elacocarpus* sp., *Eugenia* sp., *Ficus* sp.

References: (ds) Schedl 1972f: 128, 1977b: 269. (tx) Schedl 1961e: 160, 1972f: 128–129, 1977b: 269–270, 1978a: 66.

sulcipennis (Schedl) 1965c: 84 (*Coccephalophonus*). Holotype, sex?: Madagascar, Perinet; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

Notes: (3) Schedl 1979j: 130 (described male).

References: (ds) Schedl 1972f: 129, 1977b: 270. (tx) Schedl 1965c: 84–85, 1971e: 17, 1972f: 129, 1977b: 270, 1978a: 72, 1979j: 130.

suspica Schedl 1970d: 239. Holotype ♂; Madagascar, Sahafanjana (Manambato), Anove; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Mauloutchia* sp.

References: (ds) Schedl 1972f: 129, 1977b: 270. (tx) Schedl 1970d: 239–240, 1972f: 129, 1977b: 270, 1978a: 72.

thoracicum Schedl 1965c: 80. Holotype ♂; Madagascar, Sambava; Schedl Collection in NHMW, Wien.

Figures: Schedl 1972f: 129 (male).

Distribution: Madagascar.

References: (ds) Schedl 1972f: 73. (tx) Schedl 1960c: 62, 1965c: 80, 1972f: 129–130, 1977b: 270, 1978a: 73.

umbonatum Schedl 1972f: 130. Holotype ♂; Madagascar, Sambava; Schedl Collection in NHMW, Wien, automatic.

Notes: (1) This is an unneeded replacement name.

References: (ds) Schedl 1972f: 130. (tx) Schedl 1972f: 130.

truncatipennis Schedl 1961e: 162. Holotype ♀; Madagascar, Perinet, Scierie; IRSM, Madagascar. Distribution: Madagascar.

Hosts: *Eucalyptus robusta*, *Ravensara* sp., *Trichilia* sp.

Notes: (3) Schedl 1965c: 81 (described male).

References: (ds) Schedl 1970d: 235, 1972f: 129, 1977b: 271. (tx) Schedl 1961e: 162, 1965c: 81, 1970d: 235, 1972f: 129, 1977b: 271, 1978a: 75.

truncatipennis tavolae Schedl 1965c: 82. Holotype ♂; Madagascar, Betsatsaky bei Perinet; Schedl Collection in NHMW, Wien.

References: (hb) Schedl 1977b: 271. (ds) Schedl 1972f: 129. (tx) Schedl 1965c: 82, 1972f: 129, 1977b: 271, 1978a: 76.

truncatum Schedl 1965c: 82. Holotype ♂; Madagascar, Kahl Schlag in Betsatsaky bei Perinet; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

References: (ds) Schedl 1972f: 129, 1977b: 272. (tx) Schedl 1965c: 82–83, 1972f: 129, 1977b: 272, 1978a: 76.

vulpinum Schedl 1965c: 83. Holotype ♂; Madagascar, Perinet; IRSM, Madagascar.

Distribution: Madagascar.

Hosts: *Cussonia vantsilana*.

References: (ds) Schedl 1972f: 130, 1977b: 272. (tx) Schedl 1965c: 83, 1972f: 130, 1977b: 272, 1978a: 81.

Subfamily Platypodinae Erichson

See also the citations of Family Platypodidae

Platypodinae

References: Handlirsch 1925: 692; Hopkins 1915c: 227; Lucas 1920: 50; Schedl 1972f: 83, 113; Strohmeier 1912c: 4.

Tribe Platypodini Erichson

Platypodariae

References: Strohmeier 1914b: 18.

Platypodini

References: Lucas 1920: 50; Schedl 1939b: 396.

Crossotarsariae

References: Strohmeier 1914b: 18.

Crossotarsini

References: Schedl 1939b: 397, 401.

Crossotarsinae

References: Schedl 1972f: 83.

Genus *Platypus* Herbst

PLATYPUS HERBST 1793: 128. Type-species: *Bostrichus cylindrus* Fabricius, monobasic.

Cylindra Illiger 1825: 87. Type-species: *Cylindra platypus* Illiger = *Bostrichus cylindrus* Fabricius, monobasic. Synonymy: Chapuis 1865: 97.

References: (tx) Bedel 1888b: 404; Chapuis 1865: 97; Illiger 1825: 87; Swaine 1909: 84; Wood, S. L. 1979a: 1.

Stenoplatypus Strohmeier 1914c: 35. Type-species: *Crossotarsus spinulosus* Strohmeier, subsequent designation by Wood 1992a: 90. Synonymy: Schedl 1939: 398.

References: (tx) Browne 1961c: 216, 1972a: 181; Kalshoven 1960: 31–50; Schedl 1939b: 398, 1972f: 215; Strohmeier 1914c: 35; Wood, S. L. 1979a: 1, 1992a: 90.

Platypinus Schedl 1939b: 397. Type-species: *Platypus curtus* Chapuis, subsequent designation by Wood 1992a: 90. Synonymy: Wood 1979a: 1.

References: (tx) Schedl 1939b: 397; Wood, S. L. 1979a: 1, 1992a: 90.

Austroplatypus Browne 1971a: 49. Type-species: *Platypus incompertus* Schedl, original designation. Synonymy: Wood 1993: (in preparation).

References: (hb) Schedl 1972f: 269–272. (ds) Schedl 1972f: 269–272. (tx) Browne 1971a: 49–50, 1972a: 180; Schedl 1972f: 269.

Keys: Chapuis 1865 (worldwide), Blandford 1894d: 133 and Nobuchi 1973b: 8 (Japan), Blandford 1895b: 91 (Mexico and Central America), for generic groups Schedl 1972f: 169–183, Strohmeier 1914c, Wood ex 1993.

Notes: (3) A major generic revision of Platypodidae was prepared by me after this catalog was complete, but before it was published, in which more than half of the species of *Platypus* were transferred elsewhere. Only a small fraction of those changes are found in this catalog, i. e., *Platyscapulus* and *Treptoplatypus*. The generic revision is expected to be published in 1993 (SLW).

References: (ay) Fuchs 1912: 5–7, 14–33; Gardner 1932: 1–9; Strohmeier 1918: 1–46. (cn) Barbey 1901: 115; Beeson 1916: 1–5, 1918: 114–124; Bennett 1958: 1–35; Blackman 1950: 300, 341–342; Catoni 1921: 21–25; Chamberlin 1918: 34–35, 1958: 33–34; Chellman 1958: 3, 8–9; Curry 1958: 128–132; Dorsey & Leach 1956: 222; Dourojeanni 1965: 9–32; Ferguson, Gibbs, & Thatcher 1960: 27–29; Fisher, Thompson, & Webb 1954: 1–21; Fonseca, Pinto da 1934: 263–289; Froggatt 1926: 359–382; Greene & Ulrich 1931: 227–282; Hagle et al. 1987: 39; Johnson 1958: 508–511; Kelsey 1947: 65–100; Maas 1921: 1–12; Mayne 1917: 1–80; Menzel 1923: 3; Roughley & Welch 1931: 1–27; Sheppard 1925: 50–54; Shuckard 1938: 508–509; Sterrett 1916: 1–88; Swaine 1914: 40, 1916c: 628; Wearne 1923: 412–413; Wolcott 1948: 385; Yazawa, Higochi, & Machii 1957: 1–56. (hb) Beeson 1910: 222–223; Froggatt 1927: 77–79; Hetrick 1967: 297; Johnson 1958: 236; Judeich & Nitsche 1895: 441–442; Kalshoven 1960: 31–34, 1963: 237; Lengerken 1939: 250, 257–258; Munro 1928: 1–29, 1946: 10–43; Neger 1908: 279; Snyder 1927: 13, 40. (ds) Beeson 1938: 295; Blackwelder 1947: 788; Cachan 1957: 5; Crotch 1863: 18; Hetrick 1967: 297; Horn 1933: 173–174; Kolbe 1897: 37, 284; Lara & Shenefelt 1965: 175; Lever 1940: 38; Lovendal 1889: 29, 83; Lundblad 1958: 512; Murayama 1928: 30; Roepke 1911: 10–11; Shuckard 1838: 508–509. (tx) Balachowsky 1949a: 276; Beal & Massey 1945: 62; Bedel 1888b: 385, 404, 421; Beeson 1922: 1–2, 1941: 341–347; Bertolini 1872: 202; Blackman 1922b: 38–39, 1944: 78; Blandford 1893c: 11; 1893d: 442, 1894d: 133, 1895b: 81–95, 114, 193–198; Blatchley & Leng 1916: 581–584; Bright 1972d; Broun 1880: 539–540; Browne 1935b: 182–186, 1936a: 1, 1950b: 649, 1955: 364, 1958b: 164–182, 1960: 213, 1961c: 194–212, 1962e: 652, 655, 1966: 233–257, 1971a: 49–50, 1972a: 184; Chamberlin 1939: 17, 36, 102–109, 1958: 33–34; Chapuis 1865: 22, 97–288; Eichhoff 1864b: 18–22, 42–45, 1881a: 305–307; Endrodi 1957: 420; Erichson 1836: 64; Escherich 1923: 459–460, 633, 638; Gaubil 1849: 125; Guerin-Meneville 1838: 183; Hagedorn 1904: 411–412, 1907: 290–293, 1910a: 1; Herbst 1793: 128; Hopkins 1914: 127, 133, 1915c: 179, 181, 198–199, 202, 227; Hubbard 1897: 14–15; Jacquelin du Val & Fairmaire 1868: 107; Karaman

- 1972: 162; Lacordaire 1866: 355, 389, 391; Latreille 1807: 277; Lea 1910: 136; LeConte 1868: 150, 1876: 343; LeConte & Horn 1883: 513-514; Motschulsky 1858: 68; Murayama 1925: 210-211, 1934: 146; Niisima 1909: 171; Nunberg 1940: 63-69, 1951: 261, 1953: 46; Nusslin 1911a; Ratzeburg 1837: 129-138, 187-188, 1839: 230; Redtenbacher 1845: 95, 151, 167; Reitter 1913a: 111-112, 1916: 306; Sampson 1924a: 545, 1924c: 129; Schedl 1935g: 320, 1935n: 633, 1939b: 383, 388-391, 397, 1940c: 203-208, 1955g: 3, 48, 1957b: 159, 1957d: 118, 1958k: 145, 1959a: 512, 1959n: 545-557, 1962k: 597-599, 1962n: 697-699, 1963j: 483, 486, 1968e: 261-270, 1972f: 169-242, 1977b: 239; Schimitschek 1937: 60; Sedlacek 1902: 2; Shuckard 1840 (1861: 64); Stelbing 1907: 41, 1914: 619-628, 634; Stephens 1839: 205-206; Strohmeier 1911a: 174, 1911b: 26, 1911d: 218, 1912c: 3, 10-19, 1912f: 78, 1914c: 1-22; Swaine 1909: 83-84; Westwood 1836: 35, 1840: 40; Wood, S. L. 1958b: 37-40, 1960a: 3, 6; Zimmermann 1868: 142.
- abietis** Wood 1958b: 39. Holotype ♂; Oak Creek Canyon, Millard Co., Utah [USA]; Wood Collection. Distribution: North America (Arizona, New Mexico, Utah in USA). Hosts: *Abies concolor*. References: (ds) Equihua & Atkinson 1987: 18; Furniss, R. L. & Carolin 1977: 338; Schedl 1972f: 197; Wood, S. L. 1958b: 39, 1979a: 1. (tx) Equihua & Atkinson 1987: 18; Schedl 1972f: 197, 1978b: 9; Wood, S. L. 1958b: 39, 1979a: 1.
- abnormis** Schedl 1975f: 378. Holotype ♀; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien. Distribution: New Guinea. References: (tx) Schedl 1975f: 378.
- abruptifer** Wood 1992a: 90. Holotype ♂; New Guinea: Adi Island to Nagoya (Japan), imported; BMNH, London, automatic. Distribution: New Guinea. Hosts: *Pometia* sp. References: (tx) Wood, S. L. 1992a: 90.
- abruptus** Browne 1986b: 337. Holotype ♂; New Guinea: Adi Island to Nagoya (Japan), imported; BMNH, London, preoccupied by Sampson 1923. References: (tx) Browne 1986b: 337; Wood, S. L. 1992a: 90.
- abruptulus** Browne 1984a: 158. Holotype ♂; Pomio (New Britain) to Nagoya (Japan), imported; BMNH, London. Distribution: New Britain Island. References: (ds) Ohno, Yoshioka, et al. 1989: 65. (tx) Browne 1984a: 158.
- abruptus** Sampson 1923b: 285. Holotype ♂; India: Mandali, Jamnagar, 6000 ft.; BMNH, London. Distribution: Asia (Uttar Pradesh in India). Hosts: *Quercus incana*.
- References: (ds) Schedl 1972f: 212. (tx) Browne 1962e: 643; Murayama 1928a: 286; Sampson 1923b: 285; Schedl 1964c: 301, 1972f: 212.
- acetabuliformis** Roberts 1986: 40. Holotype ♂; New Guinea: Papua, Popondetta; BMNH, London. Distribution: New Guinea. Hosts: *Amoora* sp. Notes: (3) Roberts 1989: 263 (described female). References: (tx) Roberts 1986: 40.
- acetabuliformis comalis** Roberts 1989: 263. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2200 m; NIHMW, Wien. References: (tx) Roberts 1989: 263, 286.
- acuticornifer** Wood 1981: 121. Holotype ♂; Luth. Miss. Sawmill, Porotop, Papua; Schedl Collection in NHMW, Wien, automatic. Distribution: New Guinea. References: (tx) Schedl 1975g: 223; Wood, S. L. 1981: 121 (published in March).
- acuticornis** Schedl 1975g: 223. Holotype ♂; Luth. Miss. Sawmill, Porotop, Papua; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1973. References: (tx) Schedl 1975g: 223.
- longicornis** Browne [1980f] 1981: 215. Holotype ♂; Luth. Miss. Sawmill, Porotop, Papua; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1942. References: (tx) Browne 1980f: 215 (actually published in September-December 1981).
- acuticornis** Schedl 1973f: 77. Holotype ♂; N. slope Mt. Dayman, Maneau Range, M. Bay Dist., 1550 m; AMNH, New York. Distribution: New Guinea. References: (tx) Schedl 1973f: 77, 1978b: 9.
- acutidentatus** Murayama 1956b: 11. Holotype ♂; Palau Island; USNM, Washington. Figures: Murayama 1956b: figs. 1-8, Wood 1960a: 10. Distribution: Micronesia (Palau Island). References: (ds) Schedl 1972f: 208; Wood, S. L. 1960a: 7. (tx) Murayama 1956b: 11; de Ruelle 1970: 68; Schedl 1972f: 208; Wood, S. L. 1960a: 7, 10.
- adnexus** Schedl 1967d: 17. Holotype ♂; Brasilien, Nova Teutonia, Santa Catarina; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (ds) Schedl 1972f: 213. (tx) Schedl 1967d: 17-18, 1972f: 213, 1978b: 9; Wood, S. L. 1966a: 62.
- aduncus** Chapuis 1865: 283. Holotype ♀; Borneo, Sarawak; BMNH, London. (Schedl 1960c: 55). Distribution: Indonesia (Borneo). References: (ds) Sampson 1928b: 392; Schedl 1972f: 208; Strohmeier 1912c: 18, 1914c: 28. (tx) Chapuis 1865: 283; Sampson 1928b: 392; Schedl 1935k: 481, 1941b: 416, 1960c: 55, 1972f: 208; Strohmeier 1912c: 18, 1914c: 28.

advena Schedl 1972i: 54. Holotype ♂; New Guinea (NW), W. of Sentani, 75 m; BPBM, Honolulu.

Distribution: New Guinea.

Hosts: *Alstonia scholaris*, *Pimeleodendron* sp.

Notes: (3) Schedl 1974d: 464 (described female).

References: (ds) Schedl 1972f: 189. (tx) Schedl 1972f: 189, 1972i: 54, 1974d: 465, 1978b: 9.

aequalicinctus Schedl 1948d: 41. Lectotype ♂; Brasil: Sao Paulo: Schedl Collection in NHMW, Wien, designated by Schedl 1978b: 10.

Distribution: South America (Brazil).

References: (ds) Schedl 1972f: 230. (tx) Schedl 1948d: 41, 1972f: 230, 1978b: 10.

aequalis Schedl 1975g: 223. Holotype ♂; Papua, Porotop, Luth. Mission Sawmill; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 223, 1978b: 10.

aequilaterus Roberts 1986: 40. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Xanthophyllum papuanum*.

References: (tx) Roberts 1986: 40.

afzeliae Browne 1972b: 29. Holotype ♂; Thailand: Donglam; BMNH, London.

Distribution: Asia (Thailand).

Hosts: *Afzelia xylocarpa*.

References: (ds) Beaver & Browne 1975: 305. (tx) Browne 1972b: 29.

agathis Browne 1986b: 335. Holotype ♂; Borneo: Bangkuran (Sabah) to Hirao (Japan), imported; BMNH, London.

Distribution: Indonesia (Borneo).

Hosts: *Agathis* sp.

References: (tx) Browne 1986b: 335.

agnatus Schedl 1969a: 217. Holotype ♂; Borneo (Sarawak), Lawas to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Indonesia (Borneo).

Hosts: *Seyara* sp., *Shorea* sp.

References: (ds) Browne 1980c: 484, 1980d: 492, 1984c: 449; Choo, Woo, & Kim 1981: 202; Nobuchi 1977: 141; Ohno et al. 1988b: 97; Schedl 1972f: 208. (tx) Nobuchi 1985: 329; Schedl 1969a: 217, 1972f: 208, 1978b: 10.

algius Schedl 1936j: 33. Lectotype ♂; Malaya Peninsula: Penang Id.; Schedl Collection in NHMW, Wien, designated by Schedl 1978b: 10.

Distribution: Asia (Malaya).

References: (hb) Browne 1941. (ds) Beaver & Browne 1978: 618; Browne 1961c: 206; Schedl 1936j: 33, 1972f: 208. (tx) Schedl 1936j: 33–34, 1941b: 418, 425, 1972f: 208, 1978b: 10.

aliennus Schedl 1963d: 228. Holotype ♂; Surinam: Schedl Collection in NHMW, Wien.

Distribution: South America (Suriname).

Hosts: Bamboo and poesentria logs.

References: (ds) Schedl 1972f: 230, 1972g: 47,

1976a: 57. (tx) Schedl 1963d: 228, 1972f: 230, 1978b: 10.

alternans Chapuis 1865: 156. Holotype ♂ (Schedl 1960c: 24); Venezuela; BMNH, London.

Distribution: North America ("Mexico"), South America (Colombia/ Guyana/ Venezuela).

Notes: (3) Schedl 1935h: 356 (described female).

References: (hb) Cleare 1924: 61–68; Swabey 1935: 7. (ds) Blackwelder 1947: 788; Blandford 1896e: 105; Equihua & Atkinson 1987: 19; Ferrer 1942; Gemminger & Harold 1872: 2697; Schedl 1960a: 79, 1972f: 205; Strohmeier 1912c: 10, 1914c: 24; Swabey 1935: 7. (tx) Blandford 1896e: 105; Chapuis 1865: 156; Equihua & Atkinson 1987: 19; Schedl 1935h: 356, 1937c: 14, 1937d: 38, 1940a: 324, 1960a: 79, 1960c: 24, 1972f: 205, 1978b: 10; Strohmeier 1912c: 10, 1914c: 24.

alternantecostatus (Schedl) 1941c: 364 (*Platyscapus*). Syntypes ♂ ♀; Java, Mt. Gede, 800 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java).

Hosts: *Dysoxylum* sp.

Notes: (1) Schedl 1978a: 10 (citation of holotype invalid).

References: (hb) Kalshoven 1960c: 41, 1960d: 41. (ds) Browne 1984c: 448; Kalshoven 1960d: 41; Schedl 1972f: 186. (tx) Browne 1962e: 653, 1972b: 30; Schedl 1941c: 363–365, 1972f: 186, 1978a: 10.

ambiguus Roberts 1987: 163. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2,200 m; BMNH, London.

Figures: Roberts 1987: 176 (male elytron).

Distribution: New Guinea.

Hosts: *Planchonella* sp.

References: (tx) Roberts 1987: 163, 176.

anaticeps Schedl 1969a: 215. Holotype ♂; Borneo (Kalimantan), Nunukan to Muroran (Japan), imported; PPST, Tokyo.

Distribution: Indonesia (Borneo).

Hosts: Meranti logs.

References: (tx) Schedl 1969a: 215, 1978a: 10.

andrewesi Strohmeier 1910e: 128. Holotype ♂; Nilgiri Hills; H.L. Andrewes Collection.

Distribution: Asia (India/ Malaya).

Notes: (3) Schedl 1942a: 211 (described female).

References: (hb) Browne 1941. (ds) Browne 1961c: 210, 1981a: 125; Mathew 1982, 1987: 188; Schedl 1936j: 21, 1972f: 201; Strohmeier 1912c: 19, 1914c: 28. (tx) Nobuchi 1985: 329; Schedl 1936j: 21, 1939c: 2, 1939e: 337, 1942a: 211, 1972: 201, 1978b: 11; Strohmeier 1910e: 128–129, 1912c: 19, 1914c: 28.

anguis Browne 1984f: 59. Holotype ♂; New Guinea: Morobe District, Bulolo, Snake Creek, 1000 m; BMNH, London.

Distribution: New Guinea.

Hosts: *Aglaia* sp.

References: (tx) Browne 1954f: 59.

angustatulus Wood 1966a: 55. Holotype ♂; 10 km SE Cartago on Pan-American Highway, Cartago Prov., Costa Rica, 5600 feet; Wood Collection.

Figures: Wood 1966a: 52.

Distribution: North America (Costa Rica).

References: (ds) Equihua & Atkinson 1987: 19; Schedl 1972f: 205. (tx) Equihua & Atkinson 1987: 19; Schedl 1972f: 205; Wood, S. L. 1966a: 52, 55.

angustatus Chapuis 1865: 157. Holotype ♂; Mexico; MNHN, Paris (Schedl 1960c: 24).

Distribution: North America (probably Veracruz in Mexico).

References: (cn) Anonymous 1953j: 25. (cc) Anonymous 1953j: 25. (ds) Blackwelder 1947: 788; Blandford 1896e: 104–105; Equihua & Atkinson 1987: 19; Ferrer 1942; Gardiner 1957a; Gemminger & Harold 1872: 2697; Mayne & Donis 1960: 98, 1962: 280; Schedl 1972f: 205; Strohmeier 1912c: 10, 1914c: 24, 36. (tx) Blandford 1896e: 104–105; Chapuis 1865: 157; Equihua & Atkinson 1987: 19; Schedl 1940a: 324, 1950d: 14, 1953g: 244, 1960c: 24, 1972f: 205; Strohmeier 1912c: 10, 1914c: 24, 36.

angusticollis Schedl 1942c: 197. Holotype ♀; New Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (bv) Gray, B. 1974b. (hb) Gray, B. 1974b. (ds) Schedl 1972f: 199. (tx) Schedl 1942c: 197, 1972f: 199, 1978b: 11.

angustior Schedl 1942c: 196. Holotype ♂; New Guinea; Schedl Collection in NHMW, Wien.

Distribution: Asia (India), New Guinea.

References: (ds) Schedl 1969c: 62, 1972f: 200. (tx) Schedl 1942c: 196, 1969b: 233, 1972f: 200.

angustioris Schedl 1948d: 42. Syntypes ♂; Brasil, Sao Paulo; Prague Museum, and Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

Notes: (1) Schedl 1978a: 11 (citation of holotype invalid).

References: (ds) Schedl 1970f: 582, 1972f: 230, 1972g: 47. (tx) Schedl 1948d: 42, 1972f: 230, 1978b: 11.

annexus Wood 1966a: 62. Holotype ♂; Tapanti, Cartago Prov., Costa Rica, 4000 feet; Wood Collection.

Figures: Wood, S. L. 1966a: fig. 20.

Distribution: North America (Costa Rica).

Hosts: *Quercus* sp.

References: (ds) Equihua & Atkinson 1987: 19; Schedl 1972f: 214. (tx) Equihua & Atkinson 1987: 19; Schedl 1972f: 214; Wood, S. L. 1966a: 62, 67.

anomalus Schedl 1975g: 224. Holotype ♀; New Guinea, Porotop, Luth. M. Station; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 224, 1978b: 11.

anoplus Schedl 1975f: 380. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 380.

anthocephali Schedl 1969c: 60. Holotype ♂; India: Bengal, Samsing, 1800 ft., Kalimpong; not in FRI, Dehra Dun.

Distribution: Asia (Bengal in India).

Hosts: *Anthocephalus cadamba*.

Notes: (1) The Schedl "paratypes" apparently are the types; none of the specimens cited by Schedl (1969c: 60) ever reached FRI. (3) Schedl 1969c: 60 (this is *anthocephali* Beeson, nomen nudum).

References: (ds) Schedl 1972f: 208. (tx) Beeson 1941 (1961: 263); Schedl 1969c: 60–61, 1972f: 208, 1978b: 11.

aolai Browne 1986c: 669. Holotype ♂; Solomon Islands: Aola to Nagoya (Japan), imported; BMNH, London.

Distribution: Solomon Islands.

Hosts: *Neonauclea* sp. log.

References: (tx) Browne 1986c: 669.

apertulus Schedl 1975f: 380. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 380.

apertus (Schedl) 1979g: 119 (*Diapus*). Holotype ♂; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

Notes: (1) Roberts 1992:(in press) (to *Platypus*).

References: (tx) Roberts 1992:(in press); Schedl 1979g: 119.

apicalis (White) 1846: 18 (*Crossotarsus*). Holotype ♂; New Zealand, North Island; BMNH, London.

Figures: Milligan 1979: fig. 5.

Distribution: Australia, Chatham Islands, New Zealand.

Hosts: *Cordyline australis*, *Pseudotsuga menziesii*.

References: (bv) Alma & van Boven 1976; Hosking 1977; Milligan 1974, 1982a, 1982b; Neumann & Harris 1974: 136; Ytsma 1986, 1988. (cn) Grehan & Nixon 1978; Hosking 1977; Milligan 1974, 1979, 1982a, 1982b; Zondag 1982. (cc) Hosking 1977; Milligan 1974; Payton 1989; Zervos 1980. (hb) Grehan & Nixon 1978; Laird 1956: 76–84; Milligan 1970a: 61–62, 1979. (ds) Alma & van Boven 1976; Broun 1918: 147; Chapuis 1865: 337; Gemminger & Harold 1872: 2697; Grehan & Nixon 1978; Hosking 1977; Hutton 1904: 219; Laird 1956: 79, 81; Milligan 1979; Nobuchi 1977: 121; Schedl 1972f: 190; Strohmeier 1912c: 5, 1914c: 35; Zondag 1982. (tx) Broun 1880; Browne 1964: 754; Chapuis 1865: 337; Milligan 1979, fig. 5; Schedl 1941e: 154, 1942c: 164, 1960c: 64, 1972f: 190; Strohmeier 1912c: 5, 1914c: 35; White 1846: 18. (ms) Ytsma 1986.

- douci* Chapuis 1865: 237. Syntypes ♂ ♀; de la Nouvelle-Zelande; BMNH, London, MNB, Berlin, IRSNB, Brussels, and MNHN, Paris (Schedl 1960c: 44). Synonymy: Schedl 1941e: 154.
References: (cn) Zondag 1966. (hb) Clark 1932a: 235–243. (ds) Clark 1932a: 235–243; Gemminger & Harold 1872: 2698; Hutton 1904: 219; Strohmeier 1912c: 16, 1914c: 26. (tx) Broun 1880; Browne 1964: 754; Chapuis 1865: 237; Schedl 1935k: 485, 1937d: 40, 1941e: 154, 1960c: 44, 1972f: 190; Strohmeier 1912c: 16, 1914c: 26.
- castaneus* Broun 1880: 542. Holotype ♂; Tairua, New Zealand; Broun Collection. Synonymy: Schedl 1962n: 754.
References: (ds) Hutton 1904: 219; Strohmeier 1912c: 19, 1914c: 28. (tx) Broun 1880: 542; Browne 1964: 754; Schedl 1957d: 118, 1972f: 190; Strohmeier 1912c: 19, 1914c: 28.
- apicaloides* Schedl 1964g: 251. Holotype ♂; Sarawak, Setapok; BMNH, London.
Distribution: Asia (Malaya), Indonesia (Borneo).
Hosts: *Garcinia* sp.
References: (hb) Browne 1961c: 212. (ds) Browne 1961c: 212; Schedl 1972f: 186. (tx) Schedl 1964g: 251, 1972f: 186, 1978b: 12.
- apicatus* Schedl 1975f: 381. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 381.
- apicatus* Schedl 1974d: 466. Holotype ♂; Panganda logging area, Watut Valley, Morobe District, New Guinea, 2126 m; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Castanopsis acuminatissima*.
References: (tx) Schedl 1974d: 466.
- applanatulus* Wood 1992b: 79. Holotype ♂; Manaus, Amazonas; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Wood, S. L. 1992b: 79.
- aplanatus* Schedl 1976a: 85. Holotype ♂; Manaus, Amazonas; Schedl Collection in NHMW, Wien, preoccupied by *applanatus* Wood 1972.
Notes: (1) The names *aplanatus* and *applanatus* are homonyms under the Code even though they differ by one letter.
References: (tx) Schedl 1976a: 85; Wood, S. L. 1992b: 79.
- applanatus* Wood 1972b: 244. Holotype ♂; 30 km E Palmar, Bolivar, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Pouteria egregia*.
References: (tx) Wood, S. L. 1972b: 244.
- araucariae* Schedl 1966e: 44. Holotype ♂; Sudbrasilien; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
Hosts: *Araucaria angustifolia*.
References: (ds) Schedl 1972f: 205, 1976a: 57; Schonherr & Pedrosa-Macedo 1981: 57. (tx) Schedl 1966e: 44, 1972f: 205, 1978b: 12.
- arcuatus* Schedl 1976a: 86. Holotype ♂; V. Vera, Matto Grosso, Brazil, Lon. 55 36. Lat. 12 46; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 86.
- arduus* (Schedl) 1942a: 214 (*Platyscapus*). Holotype ♀; Borneo; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo).
Hosts: *Shorea* sp.
Notes: (3) Schedl 1969a: 218 (described male).
References: (ds) Nobuchi 1977: 123; Schedl 1969a: 206, 1972f: 190. (tx) Browne 1962e: 653; Schedl 1942a: 214, 1969a: 218, 1972a: 190, 1978b: 12.
- areolatus* Chapuis 1865: 179. Holotype ♀; Cuba; Schedl (1960c: 30) was unable to find this type.
Distribution: Antilles Islands (Cuba).
References: (ds) Blackwelder 1947: 788; Equihua & Atkinson 1987: 19; Schedl 1972f: 230; Strohmeier 1912c: 10, 1914c: 24. (tx) Chapuis 1865: 179; Equihua & Atkinson 1987: 19; Schedl 1960c: 30, 1972f: 230; Strohmeier 1912c: 10, 1914c: 24.
- arisaunensis* Murayama 1934f: 135. Holotype ♀; Arisan (Formosa); Murayama Collection in USNM, Washington.
Distribution: Asia (Taiwan).
References: (ds) Murayama 1934f: 135, 1937c: 578; Nobuchi 1967: 26; Schedl 1972f: 194. (tx) Murayama 1934f: 135; Schedl 1972f: 194.
- armaticeps* Schedl 1936d: 15. Lectotype ♂; Malay Peninsula, Perak, Bintang Hijau Res., Chengal; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 12.
Distribution: Asia (India/ Malaya).
Hosts: *Balanocarpus heimii*.
Notes: (3) Beeson 1941 (1961: 263) (*armaticeps*, nomen nudum, not found in FRI by SLW in 1981).
References: (hb) Browne 1941. (ds) Beeson 1941 (1961: 263); Browne 1961c: 202; Schedl 1960f: 206. (tx) Schedl 1936d: 15–16, 1972f: 206, 1978a: 12.
- armatus* Chapuis 1865: 222. Holotype ♂; Colombie; Schedl (1960c: 41) was unable to find this type.
Distribution: North America (Costa Rica), South America (Colombia).
References: (ds) Blackwelder 1947: 788; Blandford 1896e: 109–110; Equihua & Atkinson 1987: 19; Gemminger & Harold 1872: 2697; Schedl 1972f: 226; Strohmeier 1912c: 13, 1914c: 26. (tx) Blandford 1896e: 109–110; Chapuis 1865: 222; Equihua & Atkinson 1987: 19; Lucas 1920: 522;

- Schedl 1939b: 399, 1960c: 41, 1972f: 226; Strohmeyer 1912c: 13, 1914c: 26.
- arrogans** Schedl 1971c: 357. Holotype ♀; Laos: Schedl Collection in NHMW, Wien.
Distribution: Asia (Laos).
References: (ds) Schedl 1972f: 194. (tx) Schedl 1971c: 357–358, 1972f: 194, 1975b: 12.
- artecarinatus** Schedl 1935h: 352. Holotype ♂; Costa Rica, Hamburgfarm, [Rio] Reventazon, Ebene Limon; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
References: (ds) Blackwelder 1947: 788; Equihua & Atkinson 1957: 19; Schedl 1972f: 235. (tx) Equihua & Atkinson 1957: 19; Schedl 1935h: 352–353, 1972f: 235, 1975b: 12.
- artecostatus** (Schedl) 1941c: 363 (*Platyscapus*). Syntypes ♂ ♀; Java, Mnt. Gede, 800 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo, Java).
Notes: (1) Schedl 1975a: 13 (citation of holotype invalid).
References: (hb) Kalshoven 1960c: 41. (ds) Browne 1961c: 213; Kalshoven 1960d: 41; Schedl 1972f: 186. (tx) Browne 1962e: 653; Schedl 1941c: 363–364, 1955h: 498, 1972f: 186, 1975a: 12.
- artesolidus** Schedl 1942a: 201. Lectotype ♂; Malaya, Rotan Tunggal F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1975a: 13.
Distribution: Asia (Malaya), Philippine Islands.
Hosts: *Vatica* sp.
References: (ds) Browne 1961c: 209, 1962c: 201, 1980d: 490, 1983a: 555; Schedl 1966b: 89, 1972f: 197. (tx) Nobuchi 1985: 329; Schedl 1942a: 201, 1972f: 197, 1975a: 13.
- artetrunccatus** Schedl 1979g: 109. Holotype ♂; Papua UMLA [Upper Manki logging area], Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1979g: 109.
- asperatus** Schedl 1976a: 86. Holotype ♂; V. Vera, Matto Grosso, Brazil, Lon. 55° 36', Lat. 12° 46'; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 86.
- aspicis** Roberts & Morimoto 1986: 163. Holotype ♂; New Guinea: Papua, Gumil, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Planchonella* sp.
References: (tx) Roberts & Morimoto 1986: 163.
- assamelae** Browne 1986b: 339. Holotype ♂; Congo: Pointe Noire to Nagoya (Japan), imported; BMNH, London.
Distribution: Africa (Zaire)
Hosts: *Pericopsis elata* log.
- References: (tx) Browne 1986b: 339.
- assimilis** Schedl 1965g: 25. Holotype ♂; O. Borneo, Mt. Tibang, 1300 and 1400 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 207. (tx) Schedl 1965g: 25, 1972f: 207, 1975b: 13.
- associatus** Schedl 1973e: 95. Holotype ♂; Buvusi, West New Britain District; CSIRO, Canberra.
Distribution: New Britain Island.
Hosts: *Eucalyptus deglupta*.
References: (tx) Schedl 1973e: 95, 1975a: 13.
- astutus** Schedl 1965g: 25. Holotype ♂; O. Borneo, Mt. Tibang, 1300 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 200. (tx) Schedl 1965g: 25, 1972f: 200, 1975b: 13.
- atrans** Roberts 1989: 264. Holotype ♂; Papua New Guinea: Mt. Kaindi, 2100–2350 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
References: (tx) Roberts 1989: 264.
- attentus** Schedl 1936c: 234. Holotype ♀; Suriname; Schedl Collection in NHMW, Wien.
Distribution: South America (Guyana/Suriname).
References: (ds) Blackwelder 1947: 788; Schedl 1972f: 235. (tx) Schedl 1936c: 234, 1972f: 235, 1975b: 13.
- auricularis** Chapuis 1865: 125. Syntypes 2 ♂; Ste. Catherine; 1 in MNB, Berlin, 1 in IRSNB, Brussels (Schedl 1960c: 16).
Distribution: South America (Brazil/ Cayenne).
References: (ds) Blackwelder 1947: 788; Gemminger & Harold 1872: 2697; Schedl 1972f: 238; Strohmeyer 1912c: 8, 1914c: 23. (tx) Chapuis 1865: 125; Schedl 1936c: 224, 1937d: 37, 1960c: 16, 1972f: 238; Strohmeyer 1912c: 8, 1914c: 23.
- auritus** Chapuis 1865: 123. Syntypes ♂ ♀; Mexico: Toxpan; BMNH, London.
Distribution: North America (Veracruz in Mexico).
References: (ds) Blackwelder 1947: 788; Blandford 1896c: 97; Equihua & Atkinson 1957: 19; Gemminger & Harold 1872: 2697; Schedl 1972f: 238; Strohmeyer 1912c: 8, 1914c: 23. (tx) Blandford 1896c: 97; Chapuis 1865: 123; Equihua & Atkinson 1957: 19; Schedl 1934e: 210, 1936c: 224, 1940a: 324, 1960c: 16, 1972f: 238; Strohmeyer 1912c: 8, 1914c: 23.
- australis** Chapuis 1865: 240. Holotype ♀; Moreton-Bay, cote orientale de la Nouvelle-Hollande [Australia]; BMNH, London.
Distribution: Australia.
Hosts: *Argyrodendron peralatum*, *Beilschmidia bancroftii*, *Ceratopetalum succirubrum*, *Cryptocarya ablati*, *Doryphora armatica*, *Yanthophyllum ostandrum*.
References: (ay) Ting 1936: 95. (cn) Brimblecombe 1951, 1956; Froggatt 1926b, 1927; Smith,

- J. H. 1932: 232, 1939: 32–35. (**hb**) Brimblecombe 1951, 1956; Froggatt 1926b, 1927. (**ds**) Boissduval 1835; Browne 1981a: 126; Gemminger & Harold 1872: 2697; Nobuchi 1977: 125; Sampson 1925: 1; Schedl 1936g: 515, 1964c: 306, 1966g: 36, 1972a: 144, 1972f: 144, 196, 1975h: 352, 1979a: 159; Strohmeier 1912c: 18, 1914c: 28. (**tx**) Carne et al. 1980; Chapuis 1865: 240; Lea 1904: 105; Sampson 1925: 1; Schedl 1936e: 58–59, 1936g: 515, 1960c: 45, 1972f: 196; Strohmeier 1912c: 18, 1914c: 28.
- crenatus* Chapuis 1865: 287. Holotype ♂; Moreton Bay, Nouvelle-Hollande [Australia]; IRSNB, Brussels (Schedl 1960c: 56). Synonymy: Lea 1904: 105.
- References: (**cn**) Froggatt 1926b, (**hb**) Froggatt 1926b, (**ds**) Gemminger & Harold 1872: 2698; Sampson 1925: 1. (**tx**) Chapuis 1865: 287; Lea 1904: 105; Sampson 1925: 1; Schedl 1960c: 56, 1972f: 196.
- bajulus** Schedl 1936e: 52. Lectotype ♂; Mentawai, Sipora, Sereinu; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 13.
- Distribution: Asia (Malaya), Indonesia (Java, Mentawai, Sumatra).
- Hosts: *Artocarpus elastica*, *Etodia* sp., *Horsfieldia glabra*, *Podocarpus imbricatus*, *Sarcocephalus* sp.
- References: (**hb**) Kalshoven 1960c: 35. (**ds**) Browne 1961c: 198, 1986c: 662; Kalshoven 1960d: 36; Schedl 1972f: 192. (**tx**) Schedl 1936e: 52–53, 1939e: 336, 1941c: 354, 1942a: 172, 1950g: 895–896, 1972f: 192, 1978b: 13.
- bajulus sumatranus* Schedl 1936e: 53. Holotype ♂; Sumatra; Schedl collection in NHMW, Wien (Schedl 1978a: 14).
- Notes: This is apparently an aberration that should have no taxonomic status.
- References: (**ds**) Schedl 1972f: 192. (**tx**) Schedl 1936e: 53, 1972f: 192, 1978a: 14.
- balanocarpus** Schedl 1936d: 15. Lectotype ♂; Malay Peninsula, Pahang; Trannu; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 14.
- Distribution: Asia (Malaya), Indonesia (Borneo).
- Hosts: *Balanocarpus heinuii*, *Shorea ovata*, *S. parvifolia*.
- Notes: (3) Beeson 1941 (1961: 263) (*balanocarpus*, nomen nudum, not now in FRI).
- References: (**hb**) Browne 1936a, 1941, 1961c: 212. (**ds**) Beeson 1961: 263; Browne 1936a, 1961c: 212, 1980b: 381, 1984c: 448; Schedl 1959c: 168, 1971c: 366, 1972f: 185. (**tx**) Schedl 1936d: 15, 1972f: 185, 1978a: 14.
- barbatulus** Schedl 1942a: 209. Holotype ♀; Borneo; Schedl Collection in NHMW, Wien.
- Distribution: Indonesia (Borneo).
- References: (**ds**) Schedl 1972f: 194. (**tx**) Schedl 1942a: 209, 1972f: 194, 1978a: 14.
- barbosai** Schedl 1972g: 75. Holotype ♂; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
- Distribution: South America (Brazil).
- Notes: (3) Schedl 1976a: 87 (described female).
- References: (**tx**) Schedl 1972g: 75, 1976a: 87, 1978a: 14.
- batesi** Chapuis 1865: 139. Holotype ♀; Bresil, Ega.; BMNH, London.
- Distribution: South America (Brazil).
- References: (**ds**) Blackwelder 1947: 788; Gemminger & Harold 1872: 2697; Schedl 1972f: 238; Strohmeier 1912c: 8, 1914c: 23. (**tx**) Chapuis 1865: 139; Schedl 1960c: 19, 1972f: 238; Strohmeier 1912c: 8, 1914c: 23.
- beareri** Browne 1975: 306 in Beaver & Browne. Holotype ♂; Thailand: Chiang Mai, Doi Pui, 1600 m; BMNH, London.
- Distribution: Asia (Thailand).
- Hosts: *Castanopsis* sp., *Quercus* sp.
- References: (**tx**) Beaver & Browne 1975: 306.
- beilschmiediae** Schedl 1957d: 122. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.
- Distribution: Africa (Nigeria/ Zaïre).
- Hosts: *Beilschmiedia gilbertii* var. *glabra*.
- References: (**ds**) Mayne & Donis 1962: 281; Roberts 1968: 281; Schedl 1962k: 639, 1972a: 223. (**tx**) Schedl 1957d: 120, 122, 1962k: 639, 1972f: 223, 1978b: 14.
- bellus** Schedl 1933c: 172. Holotype ♂; Argentina, San Ignacio, Alto Parana; Schedl Collection in NHMW, Wien.
- Distribution: Antilles Islands (Cuba/ Guadeloupe), North America (Mexico), South America (Argentina/ Bolivia/ Brazil/ Colombia/ Paraguay).
- Notes: (3) This is a southern South American species; records from the Antilles Islands and from North America may represent interceptions or errors in identification.
- References: (**ds**) Blackwelder 1947: 788; Santoro 1957b: 25; Schedl 1933c: 172. (**tx**) Schedl 1933c: 172, 1936c: 237, 1936e: 51, 1937d: 39, 1948d: 38, 1952a: 446, 1958f: 35, 1960c: 26, 1972f: 233 (error), 1978b: 14; Wood, S. L. 1973a: 52.
- virgatus* Schedl 1935h: 355. Holotype ♂; Cuba; Schedl Collection in NHMW, Wien. Synonymy: Wood 1973f: 52.
- References: (**ds**) Blackwelder 1947: 791. (**tx**) Schedl 1935h: 355–356, 1960c: 26, 1972f: 233, 1978a: 81; Wood, S. L. 1973f: 52.
- bicarimulatus** Schedl 1942a: 200. Holotype ♂; Malaya, Pahang, Frasers Hills; BMNH, London.
- Distribution: Asia (Malaya).
- References: (**ds**) Browne 1961c: 199; Schedl 1972f: 197. (**tx**) Schedl 1942a: 200, 1972f: 197, 1978a: 14.
- bicaudatus** Schedl 1935h: 359. Holotype ♂; Venezuela; Schedl Collection.

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 788; Schedl 1972f: 213. (tx) Schedl 1935h: 359, 1972f: 213, 1978a: 14.

bicaudatus Browne 1981a: 134. Holotype ♂; Lavas (Borneo) to Tokyo (Japan), imported; BMNH, London.

Distribution: Indonesia (Borneo).

Hosts: *Shorea* sp.

References: (tx) Browne 1981a: 134.

bicolor Montrouzier 1861: 266. Syntypes?; Neu Caledonien; not located.

Distribution: New Caledonia Island.

References: (ds) Beeson 1938b: 295; Schedl 1972f: 242. (tx) Montrouzier 1861: 266; Schedl 1972f: 242.

bicorniger Schedl 1937e: 547. Syntypes ♂; Sarawak, foot of Mt. Dulit, junction of rivers Tinjar and Lejok; BMNH, London.

Distribution: Indonesia (Borneo).

Notes: (1) Schedl 1978a: 15 restricts type to BMNH, London material; the form he designated as the female of this species was later described as the male of *bihamatus* Schedl.

References: (ds) Schedl 1972f: 192. (tx) Schedl 1937e: 547, 1942a: 209, 1972f: 192, 1978a: 15.

bicornis Nunberg 1939: 219. Holotype ♀; Peru; Polischen Zoologischen Staatsmuseums, Warsaw.

Distribution: South America (Peru).

References: (ds) Blackwelder 1947: 788; Schedl 1972f: 238. (tx) Nunberg 1939: 219, 1940: 68, 1959c: 168; Schedl 1960c: 110.

bidens Schedl 1970e: 106. Holotype ♀; Brasilien, Terr. Amapa, Rio Felicio; Departamento de Zoologia, Secretaria Agricultura, Sao Paulo.

Distribution: South America (Brazil).

References: (ds) Schedl 1970e: 86, 1972f: 186.

(tx) Schedl 1970e: 106, 1972f: 186.

bifidus Schedl 1942a: 211. Lectotype ♂; Malaya, Pahang, Buloh F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 15.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: Anonaceae sp., *Canarium* sp., *Dryobalanops oblongifolia*, *Myristica* sp.

References: (hb) Browne 1961c: 212. (ds) Browne 1961a: 313, 1961c: 212; Schedl 1972f: 213. (tx) Schedl 1942a: 211, 1972f: 213, 1978b: 15.

biflexuosus Schedl 1962f: 163. Holotype ♂; Assam, Namluk, Lakhimpur; FRI, Dehra Dun, lost?

Distribution: Asia (Assam in India/ Malaya/ Thailand/ Vietnam).

Hosts: *Dipterocarpus obtusifolius*, *D. pilosus*, *Pinus merkusii*.

Notes: (1) The types which Schedl claimed to return never reached FRI, Dehra Dun, and in all probability are represented by the "paratypes" in the Schedl Collection. (3) Schedl 1962f: 163–164

and 1972f: 197 (cited as *Mesoplatypus beesoni* Sampson, nomen nudum, synonymy).

References: (ds) Beaver & Browne 1975: 306, 1978: 618; Browne 1968a: 132–133, 1970: 576, 1972b: 28; Nobuchi 1977: 127; Schedl 1972f: 197. (tx) Browne 1965a: 206; Schedl 1962f: 163, 1967f: 146–163, 1972f: 197, 1978b: 15.

biformis Chapuis 1865: 255. Holotype ♂; Inde, Dargeling; BMNH, London.

Distribution: Asia (Bengal, Uttar Pradesh in India/ Pakistan).

Hosts: *Pinus roxburghii*.

Notes: (3) Schedl 1971c: 388 (described female).

References: (cn) Beeson 1915a: 317–325; Champion 1922: 168–174, 232–246; Huque 1966; Khan 1975; Kleine 1932a: 310, 401; Pierce, W. D. 1917: 75; Stebbing 1914: 619; Troup 1916: 1–126. (cc) Beeson 1921b: 23; Stebbing 1914: 619. (hb) Beeson 1915a: 317–325, 1916: 222, 1921b: 23; Kleine 1932a: 310, 401; Stebbing 1914: 619. (ds) Beeson 1921b, 1941 (1961: 263); Gemminger & Harold 1872: 2697; Kleine 1932a: 310, 401; Pierce, W. D. 1917: 75; Qadri 1951a: 369, 1951c: 238; Sampson 1921: 25–26; Schedl 1971c: 164, 1972f: 184, 1974a: 88; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 255; Gardiner 1932b; Nakashima 1975: 48–51; Sampson 1921: 25–26; Schedl 1933a: 402, 1960c: 48, 1971c: 388, 1972f: 184, 1978b: 15; Stebbing 1914: 619; Strohmeier 1912c: 16, 1914c: 27.

bifurcus (Schedl) 1938g: 422 (*Crossotarsus*). Syntypes ♂ ♀; Mindanao, Lanao Province, Kolambagan; Division For. Studies and Res., Bureau For., and Schedl Collection in NHMW, Wien.

Figures: Nakashima 1975: 48–51.

Distribution: Asia (Burma/ Malaya/ Vietnam), Australia, Indonesia (Borneo), New Guinea, Philippine Islands (Luzon, Mindanao).

Hosts: *Balanocarpus heimii*, *Cytylelobium* sp., *Dipterocarpus grandiflorus*, *Dryobalanops aromatica*, *Endospermum malaccense*, *Hopsea* sp., *Ocoteles* sp., *Palaquium* sp., *Parashorea lucida*, *P. malaanan*, *Pentaeme* sp., *Shorea bentongensis*, *S. bracteolata*, *S. dasyphylla*, *S. leprosnla*, *S. negrosensis*, *S. parvifolia*, *S. singkawang*.

Notes: (1) Schedl 1978a: 15 (citation of holotype invalid). This was originally treated as a subspecies of *shorcans*. (3) Schedl 1939e: 359 (var. *mitilis*, nomen nudum, an aberration).

References: (ay) Nakashima 1975: 6. (cn) Ishikura 1966. (cc) Nakashima 1975: 6. (hb) Browne 1961c: 210–211. (ds) Browne 1968a: 132–133; Chadwick & Nikitin 1968; Choo, Woo, & Kim 1981: 203; Ishikura 1966; Nakashima 1975: 3; Nobuchi 1977: 117, 1979a: 407; Ohno, Yoneyama, & Nakazawa 1982b: 10, 1987b: 90; Schedl 1936d: 6, 1936j: 21, 1938g: 422, 1966b: 99, 1966g: 33, 1972f: 224. (tx) Nobuchi 1985: 329; Schedl 1936e: 59, 1937e: 544, 1938g: 422, 1939e:

- 1, 1939e: 334, 359, 1942a: 213, 1952b: 366, 1972f: 224, 1978a: 15.
- bigranulatus** Schedl 1980d: 122. Holotype ♂; New Guinea, Wan, Morobe District; Schedl Collection in NHMW, Wien, automatic.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 224, 1980d: 122.
- bituberculatus** Schedl 1975g: 224. Holotype ♂; New Guinea, Wan, Morobe District; Schedl Collection in NHMW, Wien, preoccupied by Nunberg 1967.
References: (tx) Browne 1980f: 215; Schedl 1975g: 224; Wood, S. L. 1981: 121.
- bituberculifer** Wood 1981: 121. Holotype ♂; New Guinea, Wan, Morobe District; Schedl Collection in NHMW, Wien, automatic.
Notes: (1) This is an unneeded replacement name.
References: (tx) Wood, S. L. 1981: 121.
- jeffersi** Browne 1980f: 215. Holotype ♂; New Guinea, Wan, Morobe District; Schedl Collection in NHMW, Wien, automatic.
Notes: (1) This is an unneeded replacement name.
References: (tx) Browne 1980f: 215.
- bihamatus** Schedl 1942a: 209. Syntypes ♂; Sarawak, foot of Mt. Dulit, junction of rivers Tinjar and Lejok; BMNH, London, and Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Sarawak in Borneo).
Notes: (1) The male of this species was originally described as the female of *bicorniger*, an error.
References: (ds) Kolbe 1897: 285; Schedl 1972f: 193. (tx) Schedl 1937e: 547–548, 1942a: 209, 1972f: 193.
- bilobatus** Strohmeier 1911g: 234. Holotype ♂; Peru; Strohmeier Collection.
Distribution: South America (Peru).
References: (ds) Blackwelder 1947: 788; Schedl 1972f: 230; Strohmeier 1912c: 13, 1914c: 25. (tx) Lucas 1920: 522; Schedl 1952h: 72, 1972f: 230; Strohmeier 1911g: 234, 1912c: 13, 1914c: 25.
- bilobipennis** Schedl 1964g: 250. Holotype ♂; Sarawak, Semengoh; BMNH, London.
Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).
Hosts: *Shorea leprosula*, *S. parvifolia*, *S. pauciflora*.
References: (ds) Browne 1961c: 211, 1980c: 483; Schedl 1972f: 207. (tx) Schedl 1964g: 250, 1972f: 207, 1978b: 16.
- binodulus** Chapuis 1865: 130. Holotype ♀; Bresil meridional, Nova Fribourg; IRSNB, Brussels (Schedl 1960c: 17).
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 788; Gemminger & Harold 1872: 2697; Schedl 1972f: 238; Strohmeier 1912c: 8, 1914c: 23. (tx) Chapuis 1865: 130; Schedl 1934e: 211, 1960c: 17, 1972f: 238; Strohmeier 1912c: 8, 1914c: 23.
- biporus** Blandford 1896e: 109. Syntypes 6, ♂ ♀; Mexico, Guatemala: Quiche Mountains, Capetillo, San Geronimo; BMNH, London.
Figures: Blandford 1896e: pl. 4, figs. 19–20.
Distribution: North America (Guatemala/ Hidalgo, Michoacan, Veracruz in Mexico).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 19; Ferrer 1942; Schedl 1972f: 220; Strohmeier 1912c: 13, 1914c: 25. (tx) Blandford 1896e: 109; Equihua & Atkinson 1987: 19; Schedl 1939b: 399, 1940a: 327, 1972f: 220; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1966a: 64.
- bipyramidus** Roberts & Morimoto 1986: 165. Holotype ♂; New Guinea: Papua, Gum, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthomyrtus* sp.
References: (tx) Roberts & Morimoto 1986: 165.
- bisignatus** Schedl 1962f: 164. Holotype ♂; Margherita, Lakhimpur, Assam; FRI, Dehra Dun, lost?
Distribution: Asia (Assam in India/ Malaya).
Hosts: *Shorea assanica*.
Notes: (1) None of the types studied by Schedl, including this one, had reached the FRI by 1981; Schedl's "paratype" is probably the missing type.
References: (ds) Schedl 1972f: 198. (tx) Schedl 1962f: 164, 1972f: 198, 1978b: 16.
- bispinatus** Browne 1984c: 457. Holotype ♂; Tg. Usau (West Irian) to Nagoya (Japan), imported; BMNH, London, automatic.
Distribution: New Guinea.
References: (tx) Browne 1984c: 457.
- trispinatus** Browne 1983a: 561. Holotype ♂; Tg. Usau (West Irian) to Nagoya (Japan), imported, preoccupied by Schedl 1951.
References: (tx) Browne 1983a: 561, 1984c: 457.
- biuncus** Blandford 1896b: 194. Syntypes ♂ ♀; Sumatra; BMNH, London.
Distribution: Asia (Bengal in India/ Malaya), Indonesia (Borneo, Java, Sumatra), Philippine Islands (Luzon).
Hosts: *Actinophora fragrans*, *Albizia falcata*, *Artocarpus elastica*, *Bischofia javanica*, *Cananga odorata*, *Castanospermum* sp., *Dryobalanops aromatica*, *D. biuncus*, *D. oblongifolia*, *Elaeocarpus ganitrus*, *Eusideroxylon zwageri*, *Gluta rengas*, *Heritiera fomes*, *Hevea brasiliensis*, *Millettia delaisensis*, *Pasania* sp., *Persea gratissima*, *Piarua* sp., *Schinus noronhai*, *Semicarpus heterophylla*, *Tectona* sp., *Thea sinensis*, *Xylocarpus granatum*.
References: (en) Ishikura 1966; Ymus & Hua 1980: 216. (hb) Browne 1941 (1961c: 206–207); Kalshoven 1960c: 39, 1960d: 39. (ds) Browne 1961c: 206; Choo, Woo, & Kim 1981: 202;

- Ishikura 1966; Kalshoven 1960d: 39; Murayama 1957a: 37; Nobuchi 1977: 141; Ohno, Yoneyama, & Nakazawa 1982b: 9, 1987b: 89; Schedl 1933c: 176, 1936d: 7, 1962f: 164, 1966b: 89, 1969a: 205, 217, 1972f: 208; Strohmeier 1912c: 18, 1914c: 28; Yunis & Hna 1980: 216. **(tx)** Blandford 1896b: 194; Nobuchi 1985: 329; Schedl 1935j: 1, 1937d: 42, 1941b: 416, 425, 1972f: 208; Strohmeier 1912c: 18, 1914c: 28.
- boettcheri** Schedl 1966b: 89. Holotype ♂; Philippine Is., Mindanao, Kolambugan; MNB, Berlin. Distribution: Philippine Islands (Mindanao). References: **(ds)** Schedl 1966b: 86, 1972f: 198. **(tx)** Schedl 1966b: 89, 1972f: 198, 1978b: 16.
- borneensis** Browne 1983a: 565. Holotype ♂; Lawas (Sarawak) to Nagoya (Japan), imported; BMNH, London. Distribution: Indonesia (Sarawak in Borneo). References: **(tx)** Browne 1983a: 565.
- brasilensis** Nunberg 1959c: 168. Holotype ♂; Brazil; Schedl Collection in NHMW, Wien, automatic. Distribution: South America (Brazil). References: **(ds)** Schedl 1972f: 214, 1976a: 57. **(tx)** Nunberg 1959c: 168; Schedl 1960i: 110, 1972f: 214.
- caudatus** Schedl 1933c: 169. Holotype ♂; Brazil; Schedl Collection in NHMW, Wien, preoccupied by Motschulsky 1863. References: **(ds)** Blackwelder 1947: 789. **(tx)** Nunberg 1959c: 168; Schedl 1933c: 169, 1960i: 110, 1972f: 214.
- brevicaudatus** Nunberg 1939: 232. Holotype ♂; Peru: Lima; Polischen Zoologischen Staatss-museums, Warsaw. Figures: Nunberg 1939: pl. 16, fig. 3, pl. 19, pl. 21, figs. 4, 6, pl. 22, figs. 3, 10. Distribution: South America (Peru). References: **(ds)** Blackwelder 1947: 789; Schedl 1972f: 238. **(tx)** Nunberg 1939: 232–237, 1940: 50–55; Schedl 1952h: 72, 1972f: 238.
- brevicornis** Wood 1966a: 61. Holotype ♂; Villa Mills near Cerro de la Muerte, Costa Rica, 10,000 feet; Wood Collection. Figures: Wood 1966a: 60. Distribution: North America (Costa Rica). Hosts: *Brunellia costaricensis*. References: **(ds)** Equihua & Atkinson 1987: 19; Schedl 1972f: 220. **(tx)** Equihua & Atkinson 1987: 19; Wood, S. L. 1966a: 60–61.
- brevis** Browne 1975: 306 in Beaver & Browne. Holotype ♂; Thailand; Nakhon Ratchasima, Khao Yai; BMNH, London. Distribution: Asia (Thailand). Hosts: Tree. References: **(tx)** Beaver & Browne 1975: 306–307.
- brownei** Schedl 1979j: 128. Holotype ♂; Malaysia: Penang, Telok Bahang; BMNH, London, automatic. Figures: Browne 1977b: 370 (male declivity). Distribution: Asia (Malaya). References: **(tx)** Schedl 1979j: 128.
- pusillus** Browne 1977l: 370. Holotype ♂; Malaysia: Penang, Telok Bahang; BMNH, London, preoccupied by Schedl 1952. References: **(ds)** Beaver & Browne 1978: 620; Mayne & Donis 1962: 286. **(hb)** Beaver & Browne 1978: 620. **(tx)** Browne 1977l: 370; Schedl 1979j: 128.
- brunneus** Chapuis 1865: 228. Holotype ♂; Bresil, Para; BMNH, London (Schedl 1960c: 42). Distribution: South America (Brazil/ Cayenne). References: **(ds)** Blackwelder 1947: 789; Gem-minger & Harold 1872: 2697; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26. **(tx)** Chapuis 1865: 228; Schedl 1936c: 240, 1939j: 565, 1948f: 283, 1960c: 42, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.
- burmanus** (Sampson) 1923c: 72 (*Crossotarsus*). Holotype ♂; Burma; BMNH, London. Distribution: Asia (Burma). References: **(ds)** Beeson 1937: 90–92; Schedl 1972f: 236. **(tx)** Beeson 1937: 90–92; Sampson 1923c: 2; Schedl 1936e: 46, 1939b: 398, 1972f: 236.
- caelestis** Roberts & Morimoto 1986: 165. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London. Distribution: New Guinea. Hosts: *Xanthomyrtus* sp. References: **(tx)** Roberts & Morimoto 1986: 165.
- calamus** Blandford 1894d: 137. Syntypes 16 ♂; Miyanoshta, Oshima, Kiushiu (Iligo, Yuyama, etc.), Japan; BMNH, London. Figures: Nakane et al. 1963: pl. 192, Nakashima 1975: 24, Nobuchi 1973b: pl. 2. Distribution: Asia (Fujian in China/ Japan/ Korea/ Taiwan). Hosts: *Aesculus turbinata*, *Castanopsis cuspidata* var. *sieboldii*, *Cleycla japonica*, *Daphniphyllum glaucescens*, *Fagus crenata*, *Ilex* sp., *Machilus thunbergii*, *Meliosoma myriantha*, *Prunus* sp., *Quercus crispula*, *Q. gilva*, *Q. glauca*, *Q. myrsinacfolia*, *Q. stenophylla*. Notes: (3) Murayama 1925b: 232 (described female). References: **(ay)** Nakashima 1972, 1975: 4, 1978. **(cn)** Anonymous 1980g; Kleine 1932a: 402; Shiraki 1952. **(ce)** Inouye et al. 1955: 120; Nakashima 1972, 1975: 4, 1978; Yasumatsu & Watanabe 1965: 70. **(hb)** Inouye et al. 1955: 120; Kleine 1932a: 402; Murayama 1931b: 196; Nakashima 1972. **(ds)** Anonymous 1980g; Blandford 1894c; Choo 1983: 32; Choo & Woo 1985: 163; Kleine 1932a: 402; Ko 1969: 273; Murayama 1929a: 673, 1931b: 196, 1934f: 136, 1936a: 138, 1937c: 579, 1949c:

- 104, 1950b: 1300, 1953: 25, 1954b: 187; Nakane et al. 1963: 384; Nakashima 1975: 3; Nobuchi 1973b: 11; Schedl 1972f: 209; Shiraki 1952; Stroh-meyer 1912c: 18, 1914c: 28. (tx) Blandford 1894d: 137; Choo 1983: 32; Murayama 1925b: 214, 232, 235, 1936a: 138–139, 1950b: 1300, 1954b: 187; Nakane et al. 1963: 384, pl. 192; Nakashima 1975: 24–27; Nobuchi 1973b: 11, pl. 2; Schedl 1934f: 1647, 1941b: 416, 1972f: 209; Stroh-meyer 1912c: 18, 1914c: 28.
- calamus fukiensis* Schedl 1941a: 43. Holotype ♂; Fukien, Kuatun; Schedl Collection in NHMW, Wien (Schedl 1978a: 17).
References: (ds) Schedl 1972f: 209. (tx) Schedl 1941a: 43, 1972f: 209, 1978a: 17.
- caledoniae* Schedl 1974d: 466. Holotype ♂; Ouenarou, N. Caledoniae; Centre Technique Forestier Tropical, Nogent-Sur-Marne, France. Distribution: New Caledonia Island, New Guinea. Hosts: Kaori.
Notes: (3) Schedl 1975f: 352 (described female). References: (ds) Schedl 1979b: 155. (tx) Schedl 1974d: 466, 1975f: 352, 1978b: 17.
- caligatus* Schedl 1972g: 76. Holotype ♂; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 76, 1978b: 17.
- calignus* Schedl 1936e: 58. Syntypes ♂; Mt. Makiling, Laguna, Luzon, Philippine Islands; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands (Luzon).
Hosts: *Displodiscus paniculatus*.
Notes: (1) Schedl 1978a: 17 (citation of holotype invalid).
References: (ds) Schedl 1966b: 91, 1972f: 227. (tx) Nobuchi 1955: 329; Schedl 1936e: 58, 1972f: 227, 1978a: 17.
- canaliculatus* Schedl 1942a: 199. Syntypes ♂; Malaya, Pahang, Bt. Kajang; BMNH, London.
Distribution: Asia (Malaya).
Hosts: *Dialium* sp.
Notes: (1) Schedl 1978a: 18 (types restricted to BMNH series).
References: (ds) Browne 1961c: 199; Schedl 1972f: 198. (tx) Schedl 1942a: 199, 1972f: 198, 1978a: 18.
- candezei* Chapuis 1865: 257. Syntypes 4 ♂, 1 ♀; Borneo, Malacca; BMNH, London (Schedl 1960c: 49).
Distribution: Asia (Malaya/ Singapore), Indonesia (Borneo, Malacca).
References: (hb) Browne 1941, 1961c: 198–199. (ds) Browne 1961c: 198–199, 1960a: 371; Gem-minger & Harold 1872: 2697; Sampson 1919: 107, 1928b: 394; Schedl 1971f: 150, 1972f: 184; Stroh-meyer 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 257; Sampson 1919: 107, 1928b: 394; Schedl 1939b: 395, 1960c: 49, 1972f: 184; Stroh-meyer 1912c: 16, 1914c: 27.
- capitilanus* Roberts & Morimoto 1986: 166. Holotype ♂; New Guinea: Papua, Cumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Macaranga* sp.
References: (tx) Roberts & Morimoto 1986: 166.
- capito* Browne 1984f: 60. Holotype ♂; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Cinnamomum* sp., *Nothofagus carrii*.
References: (tx) Browne 1984f: 60.
- caravanis* Schedl 1948d: 39. Lectotype ♂; Bolivia, Nor Yungas, Caramavi; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Schedl 1972f: 238. (tx) Schedl 1948d: 39, 1952a: 445, 1972f: 238, 1978a: 18.
- carbonescens* (Beeson) 1937: 100 (*Crossotarsus*). Syntypes ♂ ♀; Queensland: Gargarra; FRI, Dehra Dun.
Distribution: Australia (Queensland).
Hosts: *Beilschmidia bancroftii*.
References: (ds) Browne 1981a: 126; Nobuchi 1977: 131; Schedl 1966g: 36, 1972f: 206, 1975f: 352. (tx) Beeson 1937: 100–101; Schedl 1972f: 206.
- carduus* Schedl 1936c: 238. Lectotype ♂; Guyane Francaise, Noveau Chantier; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 18.
Distribution: South America (Guyenne).
References: (ds) Blackwelder 1947: 759; Schedl 1972f: 219. (tx) Schedl 1936c: 238, 1972f: 219, 1978a: 18.
- carinifer* Schedl 1970e: 106. Holotype ♂; Bolivien, Chapare; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Schedl 1972f: 235. (tx) Schedl 1970e: 106–107, 1972f: 238, 1978b: 18.
- carinifrons* Schedl 1965b: 87. Holotype ♂; Venezu-ela, La Macuy [La Mucuy]; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
Hosts: *Acacia decurrens*.
References: (ds) Schedl 1972f: 218. (tx) Schedl 1965b: 87, 1972f: 218, 1978b: 18.
- caryophyllatus* Schedl 1941c: 357. Syntypes ♂ ♀; Ema (Peninsula Leitimor), Amboina; Kalshoven Collection, and Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Amboine Island).
Hosts: *Eugenia aromatica*. *E. caryophyllata*.
Notes: (1) Schedl 1978a: 18 (citation of holotype invalid).
References: (hb) Kalshoven 1960c: 36. (ds) Kal-shoven 1960c: 36; Schedl 1972f: 188. (tx) Schedl 1941c: 357, 1972f: 188, 1978a: 18.

castigatus (Schedl) 1936e: 47 (*Crossotarsus*).
Lectotype ♂; N. Guinea S.E., Moroka, 1300 m;
Schedl Collection in NHMW, Wien, designated
by Schedl 1975a: 19.

Distribution: New Guinea.

Notes: (1) Schedl 1972f: 237 (to *Platypus*). (3)
Schedl 1942c: 195 (described female).

References: (ds) Schedl 1972f: 237. (tx) Schedl
1936e: 47–48, 1942c: 195, 1972f: 237, 1975a: 19.

cariceps Broun 1880: 541. Holotype ♀; Tairua,
New Zealand; Broun Collection.

Figures: Milligan 1979: fig. 5.

Distribution: New Zealand.

Hosts: Recorded as *Fagus* sp., now [= *Nothofagus* sp.].

References: (bv) Milligan 1974, 1952a; Ytsma 1956,
1958, 1959. (cn) Milligan 1974, 1979, 1952a; Ytsma
1959. (cc) Milligan 1974; Ytsma 1959; Zervos 1980.
(hb) Milligan 1970a: 61–62, 1979; Neumann &
Harris 1974: 135. (ds) Hutton 1904: 219; Milligan
1979; Schedl 1972f: 195; Strohmeyer 1912c: 19,
1914c: 28. (tx) Broun 1880: 541–542; Milligan
1979: fig. 5; Schedl 1939b: 401, 1972f: 195; Stroh-
meyer 1912c: 19, 1914c: 28. (ms) Ytsma 1956.

lobatus Broun 1893: 1253. Syntypes?; New Zea-
land; Broun Collection. Synonymy: Schedl
1939b: 401, 1972f: 195.

References: (ds) Gemminger & Harold 1872:
2699; Hutton 1904: 219; Strohmeyer 1912c:
19, 1914c: 29. (tx) Broun 1893: 1253; Schedl
1939b: 401, 1952a: 443, 1972f: 195; Stroh-
meyer 1912c: 19, 1914c: 29.

carus Strohmeyer 1913: 163. Syntypes ♂ ♀; Nilgiri
Hills; presumably Strohmeyer Collection.

Distribution: Asia (Bhutan/ Burma/ Cambodia/
Assam, Tamil Nadu in India/ Malaya/ Pakistan/
Thailand/ Tonkin Island in Vietnam and Vietnam),
Indonesia (Bali, Borneo, Java, Sumatra, Sumba),
Philippine Islands.

Hosts: *Albizia lucida*, *Amoora rohituka*,
Alphonsea ventricosa, *Bischofia javanica*, *Bombax
malabaricum*, *Castanopsis tribuloides*, *Dalbergia
assamica*, *D. latifolia*, *Diospyros pseudo-evanum*,
Dipterocarpus pilosus, *Dysoxylum* sp., *Elvretia
acuminata*, *Ficus glomerata*, *Gluta travancorica*,
Hevea brasiliensis, *Lagerstroemia parviflora*,
Mallotus albus, *Meliosma simplicifolia*, *Mesua
ferrea*, *Michelia oblonga*, *Nauclea excelsa*, *Nyssa
javanica*, *Pinus merkusii*, *Podocarpus imbricata*,
Pometia pinnata, *Salmalia malabarica*, *Schinus
noronhai*, *S. wallichii*, *Shorea robusta*, *Swietenia
macrophylla*, *Talauma hodgsoni*, *Tectona* sp., *Ter-
minalia catappa*, *Tetraclea nudiflora*, *Vatica
lanceaefolia*, *Vernonia arborea*.

Notes: (3) Schedl 1941b: 422 (review).

References: (bv) Beeson 1917. (cn) Kalshoven
1960c: 48; Mathur & Singh 1960b: 14, 1961a: 45;
Supriana et al. 1978. (hb) Beaver & Browne 1975:
307; Beeson 1916a: 223, 1917; Browne 1961c:
206; Kalshoven 1960c: 36–37. (ds) Beaver &

Browne 1975: 307; Beeson 1916a, 1961: 264;
Browne 1961c: 206, 1980d: 491; Kalshoven 1960c:
37; Mathur 1952, 1957: 185; Mathur & Singh
1960b: 14, 1961a: 45; Numberg 1960b: 161; Num-
berg & Chujo 1961: 358; Ohno, Yoneyama, &
Nakazawa 1952a: 5, 1952b: 10; Schedl 1962b:
157, 1962f: 165, 1964c: 305, 1965a: 430, 1966b:
91, 1971c: 368, 1971f: 150, 1972f: 209, 1975a:
453; Strohmeyer 1914c: 28; Supriana et al. 1978.
(tx) Nobuchi 1955: 329; Schedl 1939c: 1, 1941b:
422, 1942a: 172, 1955d: 273, 1972f: 209; Stroh-
meyer 1913: 162–163, 1914c: 28; Wylie & Yule
1977.

cupulifer Wichmann 1914a: 413. Syntypes 2;
Rotung, India, 1400 ft., and Debrugah,
Assam; not given. Synonymy: Schedl 1941b:
422.

References: (cn) Kleine 1932a: 310, 402;
Roonwal 1954: 47. (cc) Beeson 1921b: 23,
1923. (hb) Beeson 1923; Kleine 1932a: 310,
402. (ds) Beeson 1921b: 23, 1923, 1941 (1961:
264); Bhasin, Roonwal, & Singh 1955; Kleine
1932a: 310, 402; Roonwal 1954: 47. (tx)
Schedl 1941b: 422, 1972f: 209; Wichmann
1914a: 413–414.

celsus Roberts 1979: 84. Holotype ♂; Papua New
Guinea: 8 km south-west of Onim Hut, Mt.
Gihwe, 240 m; BMNH, London.

Distribution: New Guinea.

Hosts: *Nothofagus pullei* (living tree).

References: (tx) Roberts 1979: 84.

chevrolati Chapuis 1865: 251. Syntypes; de la Nou-
velle Guinee et l'île Waigiu; 4 ♂, 2 ♀ BMNH.
London, 1 ♂, 1 ♀ IRSNB, Brussels.

Distribution: New Guinea.

Hosts: *Heritiera trifolia*.

References: (bv) Gray, B. 1974b. (cn) Wylie &
Shanahan 1975. (cc) Roberts 1980. (hb) Gray, B.
1974b; Wylie & Shanahan 1975. (ds) Browne
1980a: 372; Gemminger & Harold 1872: 2698;
Olmo et al. 1988b: 98, 1989: 65; Schedl 1936g:
515, 1962i: 73, 1968e: 264; Strohmeyer 1912c:
18, 1914c: 28. (tx) Chapuis 1865: 151; Schedl
1936e: 56, 1937d: 42, 1940b: 434–435, 1941b:
417, 424, 1941e: 155, 1955b: 282, 1960c: 53,
1972f: 209; Strohmeyer 1912c: 18, 1914c: 28.

chimbui Schedl 1972f: 207. Holotype ♂; New
Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1972f: 207, 1978b: 19.

chiriquensis Wood 1966a: 59. Holotype ♂; near
Cerro Punta, Chiriqui Prov., Panama, 5500 feet;
Wood Collection.

Figures: Schedl 1972f: 237, Wood 1966a: 60.

Distribution: North America (Panama).

Hosts: *Inga* sp., *Ochroma* sp.

References: (ds) Equihua & Atkinson 1957: 21;
Schedl 1972f: 238. (tx) Equihua & Atkinson 1957:

- 21; Schedl 1972f: 237–238; Wood, S. L. 1966a: 59–60.
- cicatricosus Roberts** 1989: 264. Holotype ♂; Papua New Guinea: Gum, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Xanthomyrtus* sp.
References: (tx) Roberts 1989: 264, 286.
- circulariceps Schedl** 1975f: 382. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Hosts: *Castanopsis accuminatissima*.
References: (tx) Schedl 1975f: 382–383.
- circularis Chapuis** 1865: 285. Holotype ♂; Ile de Waigiu; BMNH, London (Schedl 1960c: 55).
Distribution: Indonesia (Sumatra), New Guinea, Philippine Islands.
Notes: (3) Schedl 1936e: 57 (in key, described female).
References: (hb) Browne 1941. (ds) Gemminger & Harold 1872: 2698; Schedl 1972f: 227; Strohmeier 1912c: 18, 1914c: 28. (tx) Chapuis 1865: 285; Lucas 1920: 522; Schedl 1933d: 202–203, 1936e: 56–58, 1939b: 398, 1960c: 55, 1972f: 227, 1978b: 19; Strohmeier 1912c: 18, 1914c: 28.
- circumdentatus Strohmeier** 1910e: 132. Syn-types, sex?; Nilgiri Hills (India orientalis); H.L. Andrews, 1 in Strohmeier Collection.
Distribution: Asia (Karnataka, Tamil Nadu in India).
Hosts: *Machilus macrantha*.
References: (ds) Beeson 1941 (1961: 264); Schedl 1969c: 61, 1972f: 237; Strohmeier 1912c: 19, 1914c: 28. (tx) Schedl 1972f: 237; Strohmeier 1910e: 132, 1912c: 19, 1914c: 28.
- clarkei Browne** 1975b: 395. Holotype ♂; Ethiopia: 15 km E Jimma, Badabuna Forest, 1870 m; MRCB, Tervuren.
Distribution: Africa (Ethiopia).
References: (tx) Browne 1975b: 395.
- clunialis Wood** 1966a: 67. Holotype ♂; near Moravia, Cartago Prov., Costa Rica, 1500 feet; Wood Collection.
Figures: Schedl 1972f: 195, Wood 1966a: 67.
Distribution: North America (Costa Rica).
References: (ds) Equihua & Atkinson 1957: 20; Schedl 1972f: 195. (tx) Equihua & Atkinson 1957: 20; Schedl 1972f: 195; Wood, S. L. 1966a: 67.
- clunivulus Wood** 1966a: 69. Holotype ♂; Ikuribisi, British Guiana; BMNH, London.
Figures: Schedl 1972f: 195, Wood 1966a: 67.
Distribution: South America (Guyana).
Hosts: *Pouteria guianensis*.
References: (ds) Schedl 1972f: 195. (tx) Schedl 1972f: 195; Wood, S. L. 1966a: 67, 69.
- clunivus Wood** 1966a: 68. Holotype ♂; Playon, Puntarenas Prov., Costa Rica, 150 feet; Wood Collection.
Figures: Schedl 1972f: 195, Wood 1966a: 67.
Distribution: North America (Costa Rica).
Hosts: *Cedrela mexicana*.
References: (ds) Equihua & Atkinson 1957: 20; Schedl 1972f: 195. (tx) Equihua & Atkinson 1957: 20; Schedl 1972f: 195; Wood, S. L. 1966a: 67–68.
- coadunatus Schedl** 1975g: 225. Holotype ♂; New Guinea, Wan; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 225, 1978b: 20.
- cognatus Roberts** 1989: 266. Holotype ♂; Papua New Guinea: Upper Manki Logging Area, Bulolo, 1800 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Castanopsis* sp.
References: (tx) Roberts 1989: 266, 286.
- coleopteratus (Schedl)** 1935b: 400 (*Crossotarsus*). Holotype ♂; Philippine Islands; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands.
References: (ds) Schedl 1966b: 91, 1972f: 237. (tx) Nobuchi 1985: 330; Schedl 1935b: 400–401, 1937d: 35, 1952b: 366, 1972f: 237, 1978a: 20.
- collaris Schedl** 1979g: 110. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1979g: 110.
- collatatus Schedl** 1936c: 245. Holotype ♂; Demerara [Guyana]; Schedl Collection in NHMW, Wien.
Distribution: South America (Guyana).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 218. (tx) Schedl 1936c: 245, 1972f: 218, 1978b: 20.
- collinus Browne** 1972c: 105. Holotype ♂; Uganda: Ruwenzori Forest; MRCB, Tervuren.
Distribution: Africa (Uganda).
Hosts: *Dombeya goetzenii*, *Macaranga* sp.
References: (tx) Browne 1972c: 105.
- comoreanus Schedl** 1969d: 15. Holotype ♂; Grande Comore: Niombadjou, 505 m; MNHN, Paris.
Distribution: Comoro Islands in Indian Ocean.
References: (ds) Schedl 1969d: 7, 1972f: 200. (tx) Schedl 1969d: 15–16, 1972f: 200, 1977a: 240.
- complanatus Schedl** 1936c: 245. Syntypes ♂ ♀; Guyane Française: Nonveau Chantier, Les Hattes, Charvin (Bas Maroni), St.-Laurent du Maroni; MNHN, Paris and Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).

Notes: (1) Schedl 1978a: 20 (type restricted to MNHN, Paris material).

References: **(ds)** Blackwelder 1947: 789; Schedl 1972f: 214. **(tx)** Schedl 1936c: 242, 245, 1939b: 399, 1972f: 214, 1975a: 20.

complanus Schedl 1967d: 16. Holotype ♂: Brasilien, Nova Teutonia, Santa Catarina; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: **(ds)** Schedl 1972f: 214. **(tx)** Schedl 1967d: 16–17, 1972f: 214, 1975b: 21.

compositus Say 1824: 324. Holotype ♂: Missouri [USA]; Say Collection, lost.

Figures: Atkinson 1989c, Solomon & Payne 1986: 23. Distribution: Africa (Seychelles Islands), Antilles Islands (Puerto Rico), Madagascar, North America (Costa Rica/ Guatemala/ Mexico/ Alabama, Arkansas, Florida, Illinois, Louisiana, Maryland, Mississippi, Missouri, New York, North Carolina, South Carolina, Tennessee, Texas, Virginia, West Virginia in USA), South America (Argentina/ Brazil/ Cayenne/ Guyana/ Venezuela).

Hosts: *Carya* spp., *Cassia siamea*, *Dacrynodes excelsa*, *Diospyros* sp., *Fraxinus* sp., *Magnolia grandiflora*.

References: **(ay)** Boving & Craighead 1931; Francke-Grosman 1956b: 292; Hopkins 1894g; Khalaf 1980: 339. **(bv)** Atkinson, Foltz, & Connor 1985; Dethier 1947. **(cn)** Anonymous 1895: 72, 1966f, 1966h; Beal & Massey 1945: 65; Beal et al. 1952: 114–115; Blackman 1950: 342–343; Chamberlin 1939: 107–111; Christian 1939: 110; Craighead 1935: 139; Creighton 1945: 706; Doane et al. 1936; Dorsey & Leach 1955: 222; Ebeling 1950: 535, 1959; Felt 1924: 275, 1926: 275, 1930a: 275; Fisher, Thompson, & Webb 1953, 1954b: 3–15; Head 1966a: 4; Herrick 1935: 375; Hopkins 1894: 277, 1896d: 250, 1897b: 151, 1904a: 39, 45, 1905: 381–398, 1907b: 2; Hubbard 1897b: 14; Kowal 1949b: 470–475; Riley 1893b: 17; Rosenfeld 1910: 216; Schluder 1969: 82; Snyder 1927: 10–12; Solomon & Payne 1986. **(cc)** Baker 1963: 222–265; Batra 1963a, 1963b: 216; Chamberlin 1925: 34, 1939: 107–111; Equihua & Atkinson 1986: 624; Felt 1906: 720; Francke-Grosman 1956b: 292; Pierce, W. D. 1908: 350; Verrall 1941: 552, 1943: 125; Webb 1945: 75. **(hb)** Atkinson 1989c; Atkinson, Foltz, & Connor 1985; Baker, W. L. 1972: 273; Beal & Massey 1945: 65; Blackman 1922b: 38–39, 1950; Chamberlin 1939: 107–111; Chittenden 1890; Deyrup & Atkinson 1987a: 64; Doane et al. 1936; Dyakowski 1911; Ebeling 1950; Equihua & Atkinson 1986: 624; Felt 1906: 720, 1926: 275, 1930a: 275; Herrick 1935: 375; Hetrick 1967; Hopkins 1894g, 1904a: 39, 45, 1905a: 384, 1907: 2, 1910: 2; Hubbard 1897a: 426, 1897b: 14; Husson 1955: 350; Ostmark 1968; Pierce, W. D. 1907: 289; Schedl 1977b: 241; Solomon & Payne 1986; White, R. E. 1983: 326. **(ds)**

Anonymous 1926c: 514, 1966f, 1966h, 1979i; Atkinson 1989c; Atkinson & Equihua 1987: 20; Atkinson et al. 1991: 153; Beal & Massey 1945: 65; Blackman 1922b: 38–39, 1950; Blackwelder 1947: 789; Blandford 1895b, 1896c; Blatchley & Leng 1916: 583; Brimley 1938: 245–248; Chamberlin 1925, 1939: 107–111; Chapuis 1865: 163–164; Chittenden 1890; Currie 1905; Deyrup & Atkinson 1987a: 64; Dozier 1918: 374; Drooz 1985: 376; Ebeling 1950, 1959; Equihua 1985: 143; Equihua & Atkinson 1986: 624, 1987: 20; Erichson 1836: 65; Felt 1926: 275, 1930a: 275; Ferrer 1942; Frost 1964: 144; Gemminger & Harold 1872: 2698; Glick 1939: 37, 70; Henshaw 1885: 147; Hoffmann 1940: 60, 1942: 12; Hopkins 1893a: 127, 1893b: 207; Kirk 1969, 1970; Leng 1920: 337; Leonard 1928: 514; Martorell 1945: 468; Melsheimer 1853: 87; Nunberg 1963: 107, 1972b: 189; Ostmark 1968; Riley 1894: 227; Riley & Howard 1891: 418; Rosewell 1920: 148; Schedl 1933c: 171, 1972f: 231, 1977b: 241, 1977d: 281; Schwarz 1878d: 468, 1882: 823; Schwerdtfeger 1960: 255–259; Staines, C. L. 1982; Strohmeier 1911f: 204, 1912c: 10, 1914c: 24; Swaine 1909: 84; Wolcott 1948: 384; Wood, S. L. 1979a: 1. **(tx)** Atkinson 1989c; Beal & Massey 1945: 65; Blackman 1922b: 38–39; Blandford 1895b: 89–94, 1896c: 106; Blatchley & Leng 1916: 583–584; Boving & Craighead 1931; Chamberlin 1939: 107–111; Chapuis 1865: 37, 163–164; Equihua & Atkinson 1987: 20; Erichson 1836: 65; Gardner 1932b: 1; Hubbard 1897b: 14; LeConte 1868: 151, 1876: 344, 1878a: 468; Sampson 1922: 141; Say 1824: 182; Schaufuss 1897: 222; Schedl 1933a: 400, 1937d: 38–39, 1939b: 399, 1940a: 324, 1955g: 48, 1960c: 25, 1961e: 128, 157, 1969d: 9, 12, 1972f: 231, 1977b: 241; Schwarz 1889b: 149; Solomon & Payne 1986: 23; Strohmeier 1911f: 204, 1912c: 10–11, 1914c: 24; Swaine 1909: 84; Wood, S. L. 1979a: 1.

tremiferus Chapuis 1865: 174. Syntypes ♂ ♀: Louisiane et de Texas: 1 ♀ in BMNH, London. 2 ♂, 2 ♀ in IRSNB, Brussels. Synonymy: LeConte 1876: 344.

References: **(cn)** Schwerdtfeger 1960b. **(ds)** Gemminger & Harold 1872: 2700; Leng 1920: 337; Sampson 1928a: 3; Schwerdtfeger 1960b: 258; Strohmeier 1912c: 13, 1914c: 25; Swaine 1909: 84. **(tx)** Chapuis 1865: 174; LeConte 1865: 151, 1876: 344; Sampson 1928a: 3; Schedl 1955g: 48, 1960c: 29, 1972f: 231; Strohmeier 1912c: 13, 1914c: 25; Swaine 1909: 84; Wood, S. L. 1979a: 1.

perforatus Chapuis 1865: 176. Syntypes ♂ ♀: de l'Amerique boreale, de la Louisiane: 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 29). Synonymy: LeConte 1876: 344.

References: **(ds)** Gemminger & Harold 1872: 2699; Leng 1920: 337; Strohmeier 1912c: 11, 1914c: 25; Swaine 1909: 84. **(tx)** Chapuis 1865:

- 176; LeConte 1868: 151, 1876: 344; Schedl 1960c: 29, 1972f: 231; Strohmeier 1912c: 11, 1914c: 25; Swaine 1909: 84; Wood, S. L. 1979a: 1.
- rugosus* Chapuis 1865: 176. Holotype ♂; Amerique boreale, Lonsiane and Texas; IRSNB, Brussels (Schedl 1960c: 29). Synonymy: LeConte 1876: 344.
- References: (ds) Gemminger & Harold 1872: 2700; Leng 1920: 337; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 84. (tx) Chapuis 1865: 176; LeConte 1868: 151, 1876: 344; Schedl 1960c: 29, 1972f: 231; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 84; Wood, S. L. 1979a: 1.
- subcavifrons* Chapuis 1865: 177. Syntypes 2 ♀; de Rio-Janeiro; BMNH, London (Schedl 1960c: 29). Synonymy: Schedl 1972f: 231.
- References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Strohmeier 1912c: 13, 1914c: 25. (tx) Chapuis 1865: 177; Schedl 1960c: 29, 1972f: 231; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1979a: 1.
- rudifrons* Chapuis 1865: 179. Holotype ♀; de Teapa; IRSNB, Brussels (Schedl 1960c: 30). Synonymy: Schedl 1960c: 25, 30.
- References: (ds) Blackwelder 1947: 790; Ferrer 1942; Gemminger & Harold 1872: 2700; Nunberg 1963c: 107; Schedl 1933c: 172; Strohmeier 1912c: 12, 1914c: 25. (tx) Blandford 1896e: 106; Chapuis 1865: 179; Schedl 1933a: 400, 1933e: 172, 1940a: 325, 1960c: 25, 30, 1972f: 231; Strohmeier 1912c: 12, 1914c: 25; Wood, S. L. 1979a: 1.
- dicgensis* Schaufuss 1897a: 222. Holotype ♀; Madagascar; Hamburg Museum, lost. Synonymy: Schedl 1972f: 231.
- References: (cn) Beaver 1988a: 70. (hb) Beaver 1988a: 70. (ds) Alluaud 1900: 441; Beaver 1988a: 70; Schedl 1969d: 9; Strohmeier 1912c: 11, 1914c: 24. (tx) Schaufuss 1897a: 222; Schedl 1961e: 157, 1972f: 231; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 1.
- foraminosus* Schedl 1933a: 400. Holotype ♀; Rio Janeiro, Brazil; BMNH, London. Synonymy: Schedl 1972f: 231.
- References: (ds) Blackwelder 1947: 789. (tx) Schedl 1933a: 400, 1972f: 231, 1978a: 30; Wood, S. L. 1979a: 1.
- concarifrons* Schedl 1975f: 383. Holotype ♀; Karamui, Chimbu Dist., New Guinea; Schedl Collection in NHMW, Wien.
- Distribution: New Guinea.
- References: (tx) Schedl 1975f: 383.
- concentriporus* Roberts 1979: 85. Holotype ♂; Papua New Guinea: 14 km south-west of Onim Hut, Mount Giluwe, 2900 m; BMNH, London.
- Distribution: New Guinea.
- Hosts: *Schizomeria* sp.
- References: (tx) Roberts 1979: 85.
- conciliatus* Schedl 1936c: 225. Holotype ♂; Guyana; Schedl Collection in NHMW, Wien.
- Distribution: South America (Guyana).
- References: (ds) Blackwelder 1947: 789; Schedl 1972f: 238. (tx) Schedl 1936c: 225, 1972f: 238, 1978b: 21.
- concinulus* Blandford 1896e: 111. Syntypes ♂ ♀; Guatemala: Pantaleon, Zapote; BMNH, London.
- Figures: Blandford 1896e: pl. 4, fig. 23.
- Distribution: North America (Guatemala).
- References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 20; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26. (tx) Blandford 1896e: 111–112; Equihua & Atkinson 1987: 20; Schedl 1936c: 244, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.
- conjunctus* Schedl 1957d: 121. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
- Distribution: Africa (Zaire).
- Hosts: *Drypetes* spp.
- Notes: (1) This name is junior to *cupulatus conjunctus* Schedl 1941: 420; however, the senior name appears to be no more than an aberration and, as such, has no status in nomenclature (SLW).
- References: (ds) Mayne & Donis 1962: 281; Schedl 1962k: 640, 1972f: 215. (tx) Schedl 1957d: 120–121, 1962k: 640, 1972f: 215, 1978b: 21.
- connexus* Wood 1966a: 65. Holotype ♂; Volcan Irazu, Cartago Prov., Costa Rica, 7000 feet; Wood Collection.
- Figures: Schedl 1972f: 220; Wood 1966a: 67.
- Distribution: North America (Costa Rica).
- Hosts: *Quercus* sp.
- References: (ds) Equihua & Atkinson 1987: 20; Schedl 1972f: 220. (tx) Equihua & Atkinson 1987: 20; Schedl 1972f: 220; Wood, S. L. 1966a: 65–67.
- consequens* Schedl 1970e: 107. Holotype ♂; Brasilien, Amazonas, Sierra Neblina, N. Rio Canaburi; Zoologischen Sammlungen des Bayerischen Staates, München.
- Distribution: South America (Brazil).
- References: (ds) Schedl 1970e: 86, 1972f: 238. (tx) Schedl 1970e: 107–108, 1972f: 238.
- contaminatus* (Blandford) 1894d: 131 (*Crossotarsus*). Syntypes 4 ♀; Japan: Higo; BMNH, London.
- Figures: Nobuchi 1967: pl. 2, 1973b: pl. 2.
- Distribution: Asia (Fujian in China/ N India/ Japan/ Taiwan).
- Hosts: *Actinodaphne laucifolia*, *Aesculus turbinata*, *Fraxinus commemorialis*, *F. lauginosa*, *F. sieboldi*, *Lindera erythrocarpa*, *Machilus thunbergii*, *Pterocarpa rhoifolia*.
- References: (ds) Blandford 1894c: Murayama 1934f: 137, 1936a: 140, 1937c: 581; Nobuchi 1967: 25, 1973b: 12, 1977: 130; Schedl 1960e:

- 173, 1972f: 200; Strohmeier 1912c: 5, 1914c: 35. (tx) Blandford 1894d: 131–132; Murayama 1936a: 146–147; Nobuchi 1967: pl. 2, 1973b: 12, pl. 2; Schedl 1934f: 1647, 1941a: 43, 1955h: 46, 1972f: 200; Strohmeier 1912c: 5, 1914c: 35.
- contextus** Schedl 1963d: 229. Holotype ♂; Bolivia tropica, Region Chapare, 400 m; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Schedl 1972f: 205. (tx) Schedl 1963d: 229, 1972f: 205, 1978b: 21.
- contractus** Chapuis 1865: 148. Holotype ♂; Colombie; IRSNB, Brussels (Schedl 1960c: 22).
Distribution: North America (Guatemala), South America (Colombia).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 20; Gemminger & Harold 1872: 2698; Schedl 1966f: 77, 1972f: 238; Strohmeier 1912c: 8, 1914c: 23. (tx) Chapuis 1865: 148; Equihua & Atkinson 1987: 20; Schedl 1960c: 22, 1972f: 238; Strohmeier 1912c: 8, 1914c: 23.
- convexicauda** Schedl 1935n: 636. Lectotype ♂; Malaya States: Pahang, Jahit Forest Reserve; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 21.
Distribution: Asia (Malaya), Indonesia (Java).
Hosts: *Pasania sundaica*, *Quercus* sp.
References: (hb) Browne 1935a, 1941. (ds) Beeson 1941 (1961: 264); Browne 1935a, 1961c: 212; Schedl 1972f: 195. (tx) Schedl 1935n: 635–636, 1972f: 195, 1978a: 21.
- convexus** Schedl 1972g: 77. Holotype ♂; Surinam, Moengo; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil/ Suriname).
References: (ds) Schedl 1972f: 187, 1976a: 57. (tx) Schedl 1972f: 187, 1972g: 77, 1978a: 21.
- cordiger** Chapuis 1865: 275. Holotype ♀; de la presqu'île de Malacca (Singapour) et de Borneo (Sarawak); IRSNB, Brussels (Schedl 1960c: 52).
Distribution: Asia (Andaman Islands in India/ Malaya/ Singapore/ Vietnam), Indonesia (Borneo), New Guinea.
Hosts: *Diospyros pyrriocarpa*, *Dipterocarpus indicus*, *D. turbinatus*, *Dysoxylum* sp.
References: (cn) Ishikura 1966. (hb) Browne 1941. (ds) Beeson 1941 (1961: 264); Browne 1961c: 213; Gemminger & Harold 1872: 2698; Ishikura 1966; Schedl 1972f: 186, 1974c: 262; Strohmeier 1912c: 17, 1914c: 28. (tx) Chapuis 1865: 275; Lucas 1920: 522; Schedl 1939b: 399, 1941c: 365, 1960c: 52, 1972f: 186; Strohmeier 1912c: 17, 1914c: 28. (ms) Lucas 1920: 522.
- cordiger bifrons** Chapuis 1865: 276. Syntypes ♀; Borneo (Sarawak); BMNH, London.
References: (ds) Sampson 1928b: 394; Schedl 1972f: 186. (tx) Chapuis 1865: 275–276; Sampson 1928b: 394; Schedl 1972f: 186.
- corniventris** Schedl 1968f: 536. Holotype ♂; New Guinea, River Tor (mouth), 4 km E Hol Maffen; BPBM, Honolulu.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 100. (tx) Schedl 1968f: 535–537, 1972f: 100, 1978a: 21.
- cornutus** Schedl 1942a: 208. Holotype ♂; Borneo; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 190. (tx) Schedl 1942a: 208, 1972f: 190, 1978b: 21.
- coronatus** Schedl 1933c: 170. Holotype ♀; Costa Rica, Vara Blanca, 1700 m; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
Hosts: *Alius acuminata*, *Brannelia costaricensis*.
Notes: (3) Schedl 1939k: 720 (described male).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 20; Schedl 1933c: 170, 1972f: 205. (tx) Equihua & Atkinson 1987: 20; Schedl 1933c: 170, 1939k: 720–722, 1972f: 205, 1978b: 21; Wood, S. L. 1966a: 57, 1966b: 17.
- platyurius** Schedl 1933c: 176. Holotype ♂; Costa Rica, Santiago, 1080 m; Schedl Collection in NHMW, Wien. Synonymy: Wood 1966b: 17.
References: (ds) Blackwelder 1947: 790. (tx) Schedl 1933c: 176, 1972f: 205; Wood, S. L. 1966b: 17.
- costaricensis** Schedl 1936i: 101. Holotype ♂; Costa Rica; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 21; Schedl 1972f: 231. (tx) Equihua & Atkinson 1987: 21; Schedl 1936i: 101–102, 1972f: 231, 1978b: 22.
- costipennis** Schedl 1933a: 398. Holotype ♂; Mexico (Truqui); BMNH, London.
Distribution: North America (apparently Mexico in Mexico).
Notes: (3) Because Truqui collected on the slopes of Volcan Popocatepetl, it is presumed that the type locality was near there; the Truqui notes are at BMNH, London.
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 21; Ferrer 1942; Schedl 1972f: 238. (tx) Equihua & Atkinson 1987: 21; Schedl 1933a: 398–400, 1940a: 324, 1972f: 238.
- crassicornis** Schedl 1975f: 384. Holotype ♂; Mt. Kaindi, M. District, New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Hosts: *Ardisia* sp.
References: (tx) Schedl 1975f: 384.
- inversus** Roberts 1979: 88. Holotype ♂; Papua New Guinea; 8 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London. Synonymy: Roberts 1989: 285.
References: (tx) Roberts 1979: 88, 1989: 285.

crassiusculus Schedl 1935j: 3. Holotype ♂; Sumatra, Medan; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Sumatra).
References: (ds) Schedl 1935j: 3, 1972f: 193. (tx) Schedl 1935j: 3, 1972f: 193, 1978a: 22.

crassus Strohmeier 1913: 162. Holotype ♂; Brit. New Guinea: Aroa Fluss; SMTD, Dresden.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 193; Strohmeier 1914c: 29. (tx) Schedl 1935j: 3, 1936e: 53, 1940b: 435, 1972f: 193; Strohmeier 1913: 162, 1914c: 29.

cribricollis Blandford 1896e: 108. Syntypes 2 ♂; Panama, Bugaba; BMNH, London.
Figures: Blandford 1896e: pl. 4, fig. 18.
Distribution: North America (Panama).
References: (ds) Blackwelder 1947: 789; Equilma & Atkinson 1987: 21; Schedl 1972f: 231; Strohmeier 1912c: 11, 1914c: 24. (tx) Blandford 1896e: 108–109; Equilma & Atkinson 1987: 21; Schedl 1972f: 231; Strohmeier 1912c: 11, 1914c: 24.

cupulatus Schedl 1941b: 420. Syntypes ♂ ♀; Sumatra, Indragiri; Kalshoven Collection and Schedl Collection in NHMW, Wien.
Distribution: Asia (Burma/ Andaman Islands in India/ Malaya/ Vietnam), Indonesia (Borneo, Java, Malacca, Sumatra, Sunda Islands), New Guinea, Philippine Islands.
Notes: (1) Schedl 1978a: 22 (citation of holotype invalid).
References: (cn) Supriana et al. 1978. (hb) Browne 1961c: 204. (ds) Browne 1986b: 333; Olmo, Yoneyama, & Nakazawa 1987b: 89; Schedl 1972f: 209, 1975c: 384, 1975e: 452, 1975j: 294, 1978a: 22, 1979h: 158; Supriana et al. 1978. (tx) Schedl 1941b: 420, 1972f: 209, 1978a: 22.

cupulatus picus Schedl 1941b: 419. Holotype ♂; Malaya: Pahang, Bukit F.R.; Schedl Collection in NHMW, Wien.
References: (hb) Browne 1961c: 203. (ds) Schedl 1941b: 419, 1972f: 209. (tx) Schedl 1972a: 209, 1978a: 22.

cupulatus conjunctus Schedl 1941b: 420. Holotype ♂; Malayische Halbinsel; Schedl Collection in NHMW, Wien.
Notes: (3) This may be no more than an aberration.
References: (hb) Browne 1961c: 203. (ds) Schedl 1972f: 209. (tx) Schedl 1941b: 420–421, 1972f: 209, 1978a: 22.

cupulatus Chapuis 1865: 278. Syntypes 2 ♂, 1 ♀; de l'île de Borneo, Sarawak; BMNH, London (Schedl 1960c: 52).
Distribution: Asia (Bhutan/ Burma/ Hainan Island in China/ Andaman Islands, Tamil Nadu in India/ Malaya/ Sri Lanka/ Thailand/ Tonkin Island in Vietnam), Australia (Victoria), Hawaiian Islands, Indonesia (Malacca, Sarawak in Borneo, Sumatra, Sunda Islands), New Guinea, Philippine Islands (Luzon).

Hosts: *Agathis alba*, *Agave sisalana*, *Albizia falcata*, *Alstonia* sp., *Artocarpus* sp., *Austrobusus nitidus*, *Bombax malabaricum*, *Coclostegia griffithii*, *Croton* sp., *Dipterocarpus pilosus*, *Dolichandrone stipulata*, *Dryobalanops oblongifolia*, *Durio carinatus*, *Dyera costulata*, *Elaeocarpus ganitrus*, *Endospermum malaccense*, *Enterolobium saman*, *Eugenia glomerata*, *E. jambolana*, *Eusideroxylon zwageri*, *Ficus glomerata*, *Garcinia* sp., *Hevea brasiliensis*, *Intsia palembanica*, *Ixonanthes reticulata*, *Lanuca grandis*, *Palaquium* sp., *Parkia speciosa*, *Persea gratissima*, *Polyosma* sp., *Pterocarpus dalbergioides*, *Shorea guiso*, *S. leprosula*, *S. maxicelliana*, *S. ovalis*, *S. robusta*, *S. uliginosa*, *Styrax benzoin*, *Swintonia floribunda*, *Tectona grandis*, *Terminalia belerica*, *T. catappa*, *Vatica stapfiana*, *Xylia dolabriformis*.

References: (ay) Strohmeier 1918: 8. (bv) Beeson 1917; Meixner 1937: 1217. (cn) Beeson 1916: 1–5; Browne 1935b: 182–186, 1935b: 77–86, 1949c; Corbett & Cater 1926b: 261; French 1911; Kalshoven 1960d: 49; Kleine 1932a: 310, 402; Leeftinck 1927: 1–60; Mathur & Singh 1960a: 34, 1960b: 43, 1961a: 45, 1961b: 13; Pierce, W. D. 1917: 218; Stebbing 1914: 626; Yunnus & Hua 1980: 216. (ec) Beeson 1921b; Browne 1958b; Fuller 1958: 100; Speyer 1923: 11–23; Stebbing 1914: 626. (hb) Beaver & Browne 1978: 618; Beeson 1916a: 223, 1917; Browne 1935a, 1941, 1958b, 1961c: 203–204; Kalshoven 1960c: 36; Kleine 1932a: 310, 402; Mercer 1982: 210; Speyer 1923: 23; Stebbing 1914: 626. (ds) Beaver & Browne 1975: 307, 1978: 618; Beeson 1916a: 223, 1921b: 23, 1961: 264; Browne 1935a, 1961c: 203–204, 1980b: 351; Corbett & Cater 1926b: 261; French 1911; Gemminger & Harold 1872: 2698; Kalshoven 1960d: 37; Kleine 1932a: 310, 402; Mathur & Singh 1960a: 34, 1960b: 43, 1961a: 45, 1961b: 13; Mercer 1982: 210; Murayama 1936b: 113; Nobuchi 1977: 135; Olmo, Yoneyama, & Nakazawa 1982b: 10, 1987b: 89; Pierce, W. D. 1917: 218; Sampson 1928b: 391; Schedl 1935k: 479–489, 1936d: 7, 22, 1936g: 515, 1936j: 22, 1937f: 15, 1959c: 168, 1961c: 70, 1962b: 157, 1962f: 165, 1964c: 304, 1965g: 24, 1966b: 92, 1966g: 33, 1969e: 156, 1971c: 367, 1971f: 150, 1972f: 209, 1974c: 262, 1975a: 452; Strohmeier 1911f: 204, 1912c: 18, 1914c: 28; Yunnus & Hua 1980: 216. (tx) Browne 1955: 368; Chapuis 1865: 278; Lea 1910: 135; Lucas 1920: 522; Meixner 1937: 1217; Nobuchi 1955: 329; Sampson 1928b: 391; Schedl 1935n: 635, 1936e: 56, 1937d: 41, 1939b: 394, 1939c: 1, 1941b: 419, 1942a: 172, 1952b: 366, 1953b: 124, 1953c: 291, 1954a: 145, 1958b: 102, 1960c: 53, 1972f: 209; Stebbing 1914: 626; Strohmeier 1911f: 204, 1912c: 18, 1913: 162, 1914c: 28; Wylie & Yule 1977. (ms) Lucas 1920: 522.

cupulatus bhumi Schedl 1979h: 158. Syntypes

♂ ♀; Ostliche Neu Guinea, 4.25–4.30 S, 140 Long., Eipome-Tal, Murggona; Schedl Collection in NHMW, Wien.

References: (tx) Schedl 1979h: 158.

curtatus **Sampson** 1923b: 286. Syntypes ♂ ♀; Panglhora Block, Upper Tondou, India; BMNH, London.

Distribution: Asia (Assam, Bengal in India).

Hosts: *Anoora wallichii*, *Artocarpus lakoocha*, *Beilschmiedia sikkimensis*, *Eugenia jambolana*, *Girtona jentinsiana*, *Machilus odoratissimus*, *Quercus lamellosa*, *Shorea robusta*, *Tetraneles midiflora*, *Vatica lanceaefolia*.

References: (cn) Mathur & Singh 1961a: 45, 1961b: 46; Roonwal 1954: 66, (ds) Beeson 1961: 264; Bhasin, Roonwal, & Singh 1955; Mathur & Singh 1961a: 45, 1961b: 46; Roonwal 1954: 66; Schedl 1962f: 165, (tx) Sampson 1923b: 286; Schedl 1972f: 185.

curtus **Chapuis** 1865: 261. Syntypes ♂; de la presqu'île de Malacca (Singapour), et de l'île de Bornéo (Sarawak); 1 in BMNH, London.

Distribution: Asia (Burma/ Fujian in China/ Assam in India/ Malaya/ Singapore/ Taiwan/ Thailand/ Vietnam), Australia (Queensland), Indonesia (Borneo, Sumatra), Philippine Islands.

Hosts: *Balanocarpus heimii*, *Dipterocarpus baudii*, *D. crinitus*, *Dryobalanops aromatica*, *D. camphora*, *Durio carinatus*, *Endospermum malaccense*, *Hopea odorata*, *Intsia palembanica*, *Koompassia excelsa*, *Parashorea lucida*, *Shorea bentongensis*, *S. faguetiana*, *S. leprosula*, *S. parviflora*, *S. robusta*, *S. pauciflora*, *S. sericeiflora*, *S. uliginosa*, *Swintonia floribunda*.

References: (ay) Nakashima 1975: 5, (bv) Beeson 1917, (cn) Beeson 1915: 8–11, 1916: 1–5; Halperin & Menier 1981; Ishikura 1966; Kleine 1932a: 310, 402; Mathur & Singh 1960b: 5, 1961a: 38; Pierce, W. D. 1917: 194; Sen-Sarma & Thakur 1956: 37; Stebbing 1914: 624; Yunus & Hua 1980: 216, (cc) Beeson 1921b: 24, 1923; Browne 1955b; Nakashima 1975: 5; Stebbing 1914: 624; Thompson, W. R. & Simmonds 1964: 34, 1965: 30, (hb) Beaver & Browne 1978: 619; Beeson 1916a: 222, 1917, 1923; Browne 1935a, 1941, 1955b, 1961c: 199–200; Kalshoven 1960c: 36; Kleine 1932a: 310, 402; Mercer 1982: 210; Stebbing 1914: 624, (ds) Beaver & Browne 1978: 619; Beeson 1916a: 222, 1921b: 24, 1923, 1961: 264; Browne 1935a, 1961a: 312, 1961c: 199–200, 1966: 237, 1965a: 133; Choo, Woo, & Kim 1981: 202; Gemminger & Harold 1872: 2698; Halperin & Menier 1981; Ishikura 1966; Kalshoven 1960d: 36; Kleine 1932a: 402, 310; Mathur & Singh 1960b: 57, 1961a: 38; Mercer 1982: 210; Nakashima 1975: 3; Nobuchi 1967: 26, 1977: 118, 1980a: Ohno, Yoneyama, & Nakazawa 1982b: 10, 1987a: 95, 1987b: 89; Pierce, W. D. 1917: 194; Sampson 1925b: 394; Schedl 1936d: 7, 1953f: 83, 1959c:

168, 1964f: 305, 1964i: 249, 1966b: 92, 1966g: 33, 1971f: 150; Strohmeier 1912c: 17, 1914c: 27; Yunus & Hua 1980: 216, (tx) Browne 1955: 369; Chapuis 1865: 261; Nakashima 1975: 42–47; Nobuchi 1985: 328; Sampson 1925b: 394; Schedl 1935k: 488, 1935n: 635, 1939b: 398, 1941a: 43, 1942a: 172, 1950g: 895–896, 1952b: 365, 1955h: 46, 1960c: 49–50, 1972f: 185, 1978b: 22; Stebbing 1914: 624; Strohmeier 1912c: 17, 1914c: 27.

curtus **artecurtus** Schedl 1960c: 50. Syntypes ♂ ♀; Fukien, Kuatun, 27.40 nBr., oL. 21, 22, 23 [China]; Schedl Collection, and Alexander Konig Museum, Bonn.

Notes: (1) Schedl 1978a: 23 (citation of holotype invalid).

References: (ds) Schedl 1962f: 165, 1964c: 262, 1972f: 185, (tx) Schedl 1960c: 50, 1972f: 185, 1978a: 23.

curvidens **Schedl** 1939k: 719. Holotype ♂; Costa Rica, Hamburgfarm, [Rio] Reventazon, Ebene Limon; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 21; Schedl 1972f: 238, (tx) Equihua & Atkinson 1987: 21; Schedl 1939k: 719, 1972f: 238, 1978a: 23.

cuspidatus **Schedl** 1963d: 229. Holotype ♂; Argentinien: Tucuman, Siambon; Schedl Collection in NHMW, Wien.

Distribution: South America (Argentina).

References: (ds) Schedl 1972f: 205, (tx) Schedl 1963d: 229, 1972f: 205, 1978a: 23.

* **cylindricus** **Burmeister** 1831: 1100. Holotype, sex?; Baltischer Bernstein, Lower Oligocene; not located.

Distribution: Europe (fossil in Lower Oligocene Baltic amber).

References: (hb) Rupertsberger 1879: 231, 1880: 232, (ds) Kugelann 1794: 526, (tx) Burmeister 1831: 1100; Latreille 1807: 277; Kugelann 1794: 526; Olivier 1795b: 4; Rupertsberger 1879: 231, 1880: 232.

cylindrus (**Fabricius**) 1792: 364 (*Bostrichus*). Syntypes 2 ♂; Germania; UZMC, Copenhagen.

Figures: Acatay 1960: 12, Balachowsky 1949a: 23, 274, 277, 1963b: 1290, Bevan 1987: 116, Duffy 1953: 17, Hickin 1963: 257, Lonzil 1961: 108, Pfeffer 1989a: pls. 1, 14, Postner 1974: 484.

Distribution: Africa (Algeria/ Egypt/ Libya/ Morocco/ Tunisia), Asia (Iran/ Turkey), Europe (Austria/ Belgium/ Bulgaria/ Czechoslovakia/ Denmark/ England/ France/ Germany/ Greece/ Hungary/ Italy/ Netherlands/ Norway/ Poland/ Portugal/ Romania/ Sardinia/ Spain/ Sweden/ Switzerland/ Caucasus, Ukraine in W USSR/ Yugoslavia).

Hosts: *Castanea vesca*, *Cerasus avium*, *Fagus orientalis*, *F. sylvatica*, *Fraxinus* spp., *Phytophthora cambivora*, *Prunus* spp., *Quercus* spp.

- References: (ay) Baker, J. M. 1960b, 1965a; Baker, J. M. & Kreger-Van Rij 1964: 433–441; Chapman 1904; Crowson 1938; Dudich 1922: 1–8; Escherich 1923b: 638; Francke-Grosmann 1956b; Fuchs 1912a; Mamaev & Semenova 1961; Menier 1976: 348; Numberg 1928a: 144; Parkin 1943: 64; Roberts 1961c. (bv) Baker, J. B. 1956b, 1960b; Espanol 1967a; Menier 1976: 348. (cn) Aeloque 1914; Androic 1966: 46; Anonymous 1940b: 9, 1966c, 1972c; Barbey 1901: 115; Becker, G. 1950; Bletchley 1962, 1964; Boocock 1959a, 1959d; Breschke 1954: 104; Browne 1952, 1968b: 570; Cermak 1938; Eckstein 1926: 579; Escherich 1916: 272–273, 1923b: 638, 1936; Fagel & Guilleaume 1945; Falck 1916; Fisher 1933, 1937a, 1954a, 1954b; Gabler 1955; Gebien 1857b: 42, 63; Georgescu et al. 1957: 357, 427; Goidanich & Goidanich 1934; Groschke 1953b, 1954b: 107; Hess 1900: 56; Hess & Beck 1927: 348; Jacquist 1951a: 1, 1951b; Joly 1953, 1956: 192, 1960; Jones 1959a; Judeich & Nitsche 1895: 546; Kamp 1954: 6; Kholodkovskii 1912: 287, 323; Kleine 1932a: 310, 401; Kollar 1849b: 3; Konig, E. 1957: 113; Lapeyronie 1948: 274; Lucas 1880: 26; Maksimovic 1959: 3–13; Marcu 1962; Marie 1922: 306–311; Novak, V., Hrozinka, & Stary 1976: 88; Nusslin 1913: 292; Petrovic 1960: 23; Pierce, W. D. 1917: 154; Reissig 1949: 131; Rhumbler 1922: 335, 1927: 352; Roberts 1961c; Schimitschek 1937c: 61, 1944: 227, 1955a: 136, 1955c: 94, 1961a: 154; Schwerdtfeger 1944a: 185, 1957a: 191; Sinreich 1961: 166; Strohmeier 1906b: 329; Thalenhorst 1950: 90; Vrydagh 1947a: 8; Wachtl 1901: 382; Weber, W. 1926: 581; Weber, T. 1965; Wichmann 1927b: 379; Wolff & Krause 1922: 105; Zwolfer 1949: 401. (cc) Apel 1983; Arx & Hennebert 1965; Baker, J. M. 1960b, 1963; Baker, J. M. & Kreger-Van Rij 1964: 433, 441; Favard 1962; Fleischer 1911; Francke-Grosmann 1956b; Gauss & Wellenstein 1950; Hicken 1963; Horn 1933: 173; Kangas 1946b: 41; Kleine 1908c: 222, 1909a: 49, 78; Kostenko 1929; Neger 1909b: 408; Nosek 1959a: 118, 1959b: 85, 87; Novak, P. 1952: 417; Nusslin 1927: 352; Perris 1852: 497, 1856a: 197, 244; Pfeffer 1923a: 332, 1928b: 7; Ratzeburg 1869a: 175; Rondani 1873: 158; Scheerpeltz & Hofler 1948: 281; Schimitschek 1955a: 136; Schwerdtfeger 1944a: 185, 1957a: 191; Sedlaczek 1935a: 157; Slaby 1947: 375; Turnau 1984; Vite 1952a: 99; Wichmann 1955a: 95. (hb) Altum 1881c: 323; Anonymous 1966c, 1972c; Apel 1983; Baeta Neves 1943a; Baker, J. M. 1956a, 1956b, 1960b: 94, 1963: 232–265, 1965a; Balachowsky 1963b: 1290; Barbey 1901: 28, 115; Bargmann 1894d, 1906; Becker, G. 1950; Browne 1968b: 570; Buysson 1910; Cecconi 1905, 1924; Cermak 1938; Chapman 1870a: 103–106, 132–135, 1904; Chittenden 1890; Claudon 1871; Dyakowski 1911; Eckstein 1897, 1915, 1926: 579, 1928; Eichhoff 1881a: 55, 305; Escherich 1923b: 638; Espanol 1964b: 115–116; Falck 1916; Favard 1962; Fisher 1936a; Fleischer 1927; Fuchs 1904a; Furst 1888: 340; Gabler 1955; Gerhard 1908; Girard 1873; Groschke 1953b, 1954b; Henschel 1876a: 217, 219, 242, 1895a: 196; Hergula 1939: 305; Hess 1900: 56; Hess & Beck 1927: 348; Hicken 1963; Husson 1955: 248–356; Jacobi 1906: 148; Jacquiot 1951a: 1–4, 1951b: 283–284; Joly 1953: 3–7; Jones 1959a; Judeich & Nitsche 1895: 546; Karpinski & Strawinski 1948: 157; Karsch 1883: 143; Kholodkovskii 1912: 287, 323; Kleine 1932a: 310, 401; Knotek 1896: 148; Kollar 1849: 3; Konig, E. 1957: 113; Korolev 1959; Lengerken 1954: 325; Lepiney & Mimeur 1932: 46; Liese 1950: 141; Louzil 1961: 27; Lmardoni & Leonardi 1889: 492; Madon 1930: 98; Nordlinger 1855: 186, 1869: 335; Nosek 1959a: 118, 1959b: 85, 87; Novak, V., Hrozinka, & Stary 1976: 88; Nusslin 1898: 283, 1913: 292, 1927: 352; Perris 1852: 497, 1856a: 197, 244; Peverimhoff 1919: 257; Pfeffer 1942a: 2; Postner 1974: 483; Ratzeburg 1837: 168, 188, 1839: 165, 230; Rhumbler 1922: 335, 1927: 352; S. H. 1851: 36; Schimitschek 1944: 227, 1955a: 136; Schwerdtfeger 1944a: 185, 1957a: 191, 1981: 197; Sedlaczek 1935a: 157; Strohmeier 1906b: 329, 1907a: 66; Taschenberg 1880: 247; Vite 1952a: 99; Wachtl 1901: 382; Weber, H. 1926: 581; Wichmann 1927b: 379; Wolff 1927: 76; Wolff & Krause 1922: 105. (ds) Acatay 1960: 12; Aeloque 1896, 1914; Andersch 1851; Androic 1966: 46; Anonymous 1966c, 1972c; Arnoldi et al. 1955: 736; Audras & Schaefer 1957; Balachowsky 1963b: 1290; Barthe 1896; Bau 1888; Bedel 1888b: 404, 421; Bielz 1851, 1887; Blackwelder 1947: 789; Blanchere & Robert 1889; Bletchley 1962; Borchert 1951; Brakman 1966b: 207; Brancsik 1871, 1906; Brown, S. C. S. 1971; Browne 1968b: 570; Buck 1955b: 191; Bucking 1932; Buresh & Lazarov 1956; Buysson 1910; Calver 1854, 1893; Cecconi 1905; Chapuis & Candeze 1853; Chittenden 1890; Chrystal 1937; Debattise 1945; Dejean 1821, 1825, 1837; Dieckmann 1960: 117; Duffy 1945: 175; Eggers 1904, 1912f; Endrodi 1958a: 69, 1958b; Erichson 1836: 65; Ericson & Sandin 1893; Escalera 1919; Escherich 1923b: 638, 1932b: 43; Espanol 1964b; Eyquem 1891; Fedorov 1930; Fisher 1936a, 1937: 1–3; Fowler 1891; Fricken 1889: 357; Fuchs 1904a, 1905a; Gabler 1949b; Gaubil 1849: 125; Gauss & Wellenstein 1950; Gebin 1857a: 392; Gemminger & Harold 1872: 2698; Gozis 1875: 80; Gredler 1866: 375; Grill 1895: 313; Hallett 1923c; Hamilton 1894b: 406; Hansen, V. 1939, 1956; Hellen 1947; Hennig 1954: 265; Henschel 1895a: 196; Heyden 1876; Heyden, Reitter, & Weise 1883: 182, 1891: 674, 1906: 713; Horion 1935, 1951; Hunefeld 1831: 2000; Illiger 1805: 129; Joly 1960; Judeich & Nitsche 1895: 546–547; Kaltenbach 1874: 632; Kangas 1946d: 35; Karpinski 1926: 83, 1931: 20,

- 1932b: 56, 1945a: 173; Karpinski & Strawinski 1948: 157; Keler 1925b: 275; Kersten 1933: 76; Kestercanek 1881a: 12; Klefbeck & Sjöberg 1960: 232; Kleine 1932a: 310, 401; Kloft & Hinks 1945: 218; Koca 1905: 192; Kostenko 1929; Kozikowsky 1921: 181; Kraatz 1869: 60; Kurir 1947c: 22; Lacordaire 1866: 391; Langhoffer 1915c: 158–159, 1929: 1196–1203; Leclercq 1971; Leng 1918: 210; Lentz 1857: 140; Lepiney 1927: 145–174; Lepiney & Mimeur 1932: 46; Liese 1950: 141; Lokaj 1860: 64; Lomnicki 1913b: 149; Lompe 1973; Lucas 1880: 26; Lucht 1957: 280; Luna de Carvalho 1947: 15; Lumardoni & Leonardi 1889: 492; Massee 1947: 126; Matthews & Fowler 1883: 42; Michel 1937: 24–28; Negrin 1966b: 404, 1968a: 457; Novak, P. 1952: 417; Nunberg 1954: 96; Nusslin 1898: 283; Oliveira 1857: 329; Orten 1856: 280; Pachter 1865: 152; Paganetti-Hummiler 1901: 150; Palm 1959: 22, 28, 354, 1966: 18; Parkin 1943: 61–66; Peckaire & Van der Wiel 1946: 76; Perris 1876a: 256, 1877a: 412; Peyerimhoff 1919: 257; Pfeffer 1928b: 7, 1931b: 73, 1936: 90, 1989a: 92; Pierce, W. D. 1917: 154; Postner 1974: 486; Ragusa 1924: 117; Rapp 1934: 738; Ratzeburg 1837: 168, 188, 1839: 165, 230; Redtenbacher 1858: 839; Reissig 1949: 131; Reitter 1869b: 155, 1888b: 280, 1894a: 93, 1916: 306, 352; Romanyk 1959: 425; Roschkamp 1929: 178–180; Roubal 1941: 276; Rye 1866a: 199, 1890: 271; S. H. 1851: 36; Sainte-Claire 1914: 475; Sainte-Claire & Mequignon 1938: 449; Schaschl 1854: 133; Schaufuss 1915: 1201; Schamm 1853: 292, 1859: 96, 1862: 101; Schedl 1961b: 185, 1964a, 1967c: 73, 1971d: 429, 1972f: 189, 1980a: 29, 1981c: 101; Schilsky 1909: 189; Schiodte 1873: 105; Scheerpeltz & Winkler 1930: 253; Scherdtfeger 1981: 197; Seudder 1891: 540; Seidlitz 1872: 389, 1891a: 556, 1891b: 601; Sharp & Fowler 1893: 35; Stark 1927b: 90, 1936e: 142; Stein & Weise 1877: 165; Stephens 1829a: 145, 1830: 355, 1839: 206; Stierlin & Gantard 1871: 294, 1906: 205; Strohmeier 1907e: 170, 1912c: 15, 1914c: 26; Sturm 1843: 231; Swaine 1909: 85; Thomson 1865: 376, 1868: 224; Trappen 1935: 144; Tredl 1907: 19; Vigiani 1943: 102; Villa & Villa 1833: 26; Vite 1953: 41; Vrydagh 1947a: 8; Wanka 1908: 231, 1915: 213; Westhoff 1882: 240; Wichmann 1909b: 172, 1955a: 95; Winter, T. C. 1983: 22; Wolff 1927: 76. (tx) Acatay 1960: 12; Acloupe 1896; Altman 1844; Apel 1983; Bach 1854; Balachowsky 1949a: 277, 1963b: 1290; Barbey 1901: 28, 115; Bechstein et al. 1805: 96; Bedel 1888b: 404, 421; Beeson 1938: 295; Bertolini 1872; Brancsik 1871; Calver 1858; Castelman 1840; Ceballos 1945; Chapuis 1865: 246; Chapuis & Candeze 1853; Chevrolat 1838; Dejean 1821, 1825; Duffy 1953: 17; Eggers 1912f: 29, 1929e: 42; Eichhoff 1864b: 43, 1881a: 55, 108, 305, 1883a: 117, 144; Emden 1942; Endrodi 1957b: 43; Escherich 1923b: 638; Espanol 1964b; Fabricius 1792: 364, 1801: 384; Fauvel 1889; Fleischer 1905; Formanek 1907: 56; Fricken 1889: 357; Fuchs 1912a; Gabler 1949b, 1955; Girard 1873; Hansen 1956; Henry 1892: 15; Henschel 1876a: 217, 219, 242, 1895a: 196; Hickin 1963: 257; Hopkins 1914: 127, 1915c: pl. 9, fig. 18, pl. 12, fig. 18; Houlbert 1922a: 15–16; Iablokoff-Klmzorian 1961: 89; Illiger 1805: 321; Jacobi 1906: 148; Jacquelin du Val & Fairmaire 1868: 108; Judeich & Nitsche 1895: 546; Karpinski & Strawinski 1948: 157; Keler 1928: 1; Korolev 1989; Kuhnt 1913: 1061; Lacordaire 1866: 391; Latreille 1803: 204, 1804: 108; Letzner 1891: 378; Lemis 1886: 183; Lonzil 1961: 108; Lovendal 1889b: 83, 1898; Lucas 1920: 521; Lucht 1987: 280; Lumardoni & Leonardi 1889: 492; Negrin 1966b: 404; Novak, V., Hrozinka, & Stary 1976: 88; Nunberg 1928a: 144, 1953: 52, 1954: 96; Panzer 1795: 285; Perris 1877a: 112; Pfeffer 1932b: 27, 1942a: 2, 1955a: 265, 1989a: pl. 1, 14; Portevin 1935: 339; Postner 1974: 486; Ratzeburg 1837: 168, 188, 1839: 165, 230; Redtenbacher 1849a: 356, 1849b: 26, 1855: 839; Reitter 1894a: 93, 1913a: 112, 1916: 306, 352; Rey 1892b: 31; Rhumbler 1922: 335, 1927: 352; Saunders 1836: 155; Schedl 1934f: 1647, 1937d: 40, 1942a: 200, 1947a: 13, 1952f: 88, 1960c: 47, 1972f: 189, 1980a: 29, 1981c: 101; Schimitschek 1937c: 60–61, 1955c: 94; Seidlitz 1872: 389, 1891a: 556, 1891b: 601; Stephens 1829a: 145, 1829b: 12, 1830: 355, 1839: 206; Stresemann et al. 1889: 357; Strohmeier 1911d: 217, 1912c: 15, 1914c: 26; Swaine 1909: 85; Taschenberg 1880: 247; Thomson 1865: 376, 1868: 224; Westwood 1840: 40. (ms) Boocock 1959a, 1959d; Eggers 1912e; Escherich 1932b; Fisher 1954a, 1954b; Keler 1956: 440; Lucas 1920: 521; Schamm 1854: 148.
- platypus* Duftschmidt 1825: 87 (*Cylindra*). Syntypes ♂; Wien; not located. Synonymy: Chapuis 1865: 246.
References: (tx) Chapuis 1865: 246; Duftschmidt 1825: 87; Schedl 1972f: 189.
- bimaculata* Duftschmidt 1825: 88 (*Cylindra*). Syntypes, sex?; Wien; not located. Synonymy: Chapuis 1865: 246.
References: (ds) Schilsky 1909: 189. (tx) Balachowsky 1949a: 277; Chapuis 1865: 246; Duftschmidt 1825: 88; Espanol 1964: 116; Letzner 1891: 378; Schedl 1972f: 189.
- flavicornis* Kugelann 1872: 496 (*Bostrichus*). Syntypes, sex?; Germany; not located. Synonymy: Strohmeier 1912: 15.
References: (tx) Kugelann 1872: 496; Schedl 1972f: 189.
- cylindriformis* Reitter 1894: 94. Syntypes ♂ ♀; Algier (St. Charles); NHMB, Budapest. Synonymy: Balachowsky 1949a: 277.
Notes: (3) Dejean 1837: 333 (*sulcatus*, nomen nudum, synonymy in Strohmeier 1911: 85–88).
References: (ay) Escherich 1923b: 638; Korolev 1989; Strohmeier 1918: 1. (cn) Acatay 1943a:

- 73, 1949: 73–74; Escherich 1923b: 638; König, E. 1957: 17; Schimitschek 1955a: 136, 1955c: 94; Weber 1926: 581; Wichmann 1927b: 379. (**cc**) Beauverie 1910b; Schimitschek 1955a: 136; Uchiastnova 1985; Vite 1952a: 99. (**hb**) Acatay 1943a: 73; Eckstein 1926: 581; Escherich 1923b: 638; König, E. 1957: 17; Schimitschek 1937: 61, 1955a: 136; Strohmeier 1906b: 329–341, 409–420, 506–511, 1907a: 66, 1910b: 59; Vite 1952a: 99; Weber 1926: 581; Wichmann 1927b: 379. (**ds**) Eckstein 1926: 581; Escherich 1923b: 638; Gabler 1949b; Heyden, Reitter, & Weise 1906: 713; Holdhaus 1912: 454; Lima de Carvalho 1950: 15; Normand 1937: 269; Pfeffer 1947d: 126; Reitter 1894a: 94; Roschkamp 1929: 178–180; Schilsky 1909: 189; Strohmeier 1907c: 170–173, 1911c: 106, 1912c: 251, 1914c: 26; Tredl 1907: 20. (**tx**) Balachowsky 1949a: 277; Eggers 1923a: 214; Escherich 1923b: 638; Espanol 1964: 116; Fleischer 1927; Gabler 1949b; Korolev 1989; Kulmt 1913: 1061; Pfeffer 1989a: pl. 12; Portevin 1935: 339; Reitter 1894a: 93–94, 1913a: 113; Schedl 1934f: 1647, 1972f: 189; Schimitschek 1955c: 94, 1972f: 189; Strohmeier 1910g: 85, 1911c: 106, 1912b: 57, 1914c: 26.
- cymbiformis** Roberts 1979: 86. Holotype ♂; Papua New Guinea: 8 km south-west Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Galbulimima* sp.
References: (**cn**) Schimitschek 1937c: 61, 1938c: 2119, 1939d: 2119, 1944: 227; Schwerdtfeger 1944a: 185, 1957a: 191. (**cc**) Schimitschek 1941a: 317; Schwerdtfeger 1944a: 185, 1957a: 191. (**hb**) Schimitschek 1944: 227; Schwerdtfeger 1944a: 185, 1957a: 191. (**ds**) Strohmeier 1912c: 14. (**tx**) Schimitschek 1937c: 61; Strohmeier 1912c: 14.
- daedalus** Roberts 1986: 41. Holotype ♂; New Guinea: Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (**tx**) Roberts 1986: 41.
- darjeelingensis** Schedl 1969c: 61. Holotype ♂; India: W. Bengal, Lepchajagat, 7000', Rangirum 6000', Darjeeling; FRI, Dehra Dun, not found in 1981 by SLW.
Distribution: Asia (Bengal in India).
Hosts: *Acer campbellii*, *Odina fraxinifolia*, *Symplocos theaeifolia*.
Notes: (1) None of the specimens borrowed by Schedl from FRI, Dehra Dun, was ever returned; the types he cites may be represented by the "paratypes" in his collection. (3) Schedl 1969c: 61 (this is *darjeelingensis* Beeson, nomen nudum), 1971c: 389 (described female).
References: (**cn**) Mathur & Singh 1961a: 83. (**ds**) Beeson 1941: 344, 1961: 265; Mathur & Singh 1961a: 83; Schedl 1971c: 362, 1972f: 189. (**tx**) Schedl 1969c: 61–62, 1971c: 389, 1972f: 189, 1978a: 23.
- darlingtoni** Reichardt 1965: 162. Holotype ♂; Toledo Dist., Brit. Honduras; MCZ, Cambridge.
Distribution: North America (Belize).
References: (**ds**) Schedl 1972f: 239. (**tx**) Reichardt 1965: 162; Schedl 1972f: 239.
- dasycauda** Browne 1980a: 376. Holotype ♂; Keelung (Formosa) to Nagoya (Japan), imported; BMNH, London.
Distribution: Asia (Taiwan).
References: (**tx**) Browne 1980a: 376.
- decens** Sampson 1922b: 147. Holotype ♂; Assam: Nowgong; BMNH, London.
Distribution: Asia (China/ Assam in India/ Malay/ Nepal).
Hosts: *Castanopsis tribuloides*, *Shorea* sp., *Vatica lauceaeifolia*.
References: (**cn**) Mathur & Singh 1961b: 71. (**ds**) Beaver & Browne 1978: 619; Beeson 1961: 265; Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1961b: 71; Schedl 1962f: 165, 1972f: 209, 1973b: 211. (**tx**) Murayama 1956: 13; Sampson 1922b: 147; Schedl 1941b: 416, 1972f: 209.
- deceptor** Wood 1972c: 244. Holotype ♂; La Carbonera Experimental Forest, 50 km W. Merida, Merida, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
Hosts: *Ficus* sp., *Nectandra* sp., *Primus spachrocarpa*.
References: (**tx**) Wood, S. L. 1972c: 244.
- declivis** Sampson 1923c: 72. Lectotype ♂; Labuen, Batjan; BMNH, London, designated by Browne 1970: 580.
Distribution: Indonesia (Batjan Island near Borneo, Java).
References: (**ds**) Beeson 1937: 92; Equihua & Atkinson 1987: 21; Schedl 1972f: 237. (**tx**) Beeson 1937: 90, 92; Browne 1970: 580; Equihua & Atkinson 1987: 21; Sampson 1923c: 72; Schedl 1936c: 46, 1972f: 237.
- declivitatis** Schedl 1941c: 356. Syntypes ♂; Java, Mount Gede, 500 m; Schedl Collection in NHMW, Wien, and RNH, Leiden.
Distribution: Indonesia (Java).
Notes: (1) Schedl 1978a: 23 (citation of holotype invalid, this specimen was lost).
References: (**ds**) Schedl 1972f: 242. (**tx**) Schedl 1941c: 356, 1972f: 242, 1978a: 23.
- decorus** Schedl 1936c: 235. Syntypes ♂; French Guyana, St-Laurent du Maroni; MNHN, Paris, and Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 205. (**tx**) Schedl 1935k: 480, 1936c: 235, 1972f: 205, 1978a: 24.

- deductus** Schedl 1964c: 299. Holotype ♂: Ostkuste Sumatras: Bandar Baroe, 850 m; ZMA, Amsterdam. Distribution: Indonesia (Java, Sumatra). References: (ds) Schedl 1972f: 195. (tx) Schedl 1964c: 299, 1972f: 195, 1978a: 24.
- deflectus** Schedl 1941c: 355. Syntypes ♂ ♀; Java, Mt. Gede, 1000 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien. Distribution: Indonesia (Java), Philippine Islands. References: (hb) Kalshoven 1960c: 40–41. (ds) Kalshoven 1960c: 40–41; Schedl 1972f: 196. (tx) Browne 1966: 235; Nobuchi 1955: 329; Schedl 1941c: 355–356, 1971f: 156, 1972f: 196, 1978a: 24.
- bistriatus** Browne 1966: 235. Holotype ♂; Philippines: Palawan, Pinigisan, Mantalingajan Range, 600 m; UZMC, Copenhagen. Synonymy: Schedl 1971f: 156. References: (tx) Browne 1966: 235; Schedl 1971f: 156, 1972f: 196.
- delicatus** (Schedl) 1936b: 129 (*Crossotarsus*). Syntypes ♂; Belg. Kongo, Kibali-Ituri: Kilo; MRCB, Tervuren, and Schedl Collection in NHMW, Wien. Distribution: Africa (Angola/ Uganda/ Zaïre). Notes: (1) Schedl 1959p: 27 (described female, as *Mesoplatypus*), 1978a: 24 (restricted male type to specimen in MRCB, Tervuren). References: (ds) Ferreira 1965: 1134; Schedl 1959p: 23, 27, 1962k: 996, 1972f: 203. (tx) Schedl 1936b: 129–130, 1962k: 996, 1972f: 203, 1978a: 24.
- dentatus** Dalman 1823: 83. Holotype ♂; Brasilia; Schoenherr Collection. Distribution: North America (Costa Rica), South America (Bolivia/ Brazil/ Colombia). Hosts: *Cecropia peltata*, *Inga vera*, *Jacaranda copatia*. Notes: (3) Schedl 1970e: 108 (described female). References: (ds) Schedl 1970e: 89, 1972f: 239. (tx) Dalman 1823: 83; Schedl 1970e: 108, 1972f: 239, 1978a: 24.
- fortis** Blandford 1896e: 98. Holotype ♂; Colombia; BMNH, London. Synonymy: Schedl 1972f: 238. References: (ds) Blackwelder 1947: 789; Schedl 1966f: 80. (tx) Blandford 1896e: 98; Schedl 1958c: 2, 1972f: 238–239.
- denticollis** Browne 1984f: 61. Holotype ♂; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London. Distribution: New Guinea. Hosts: *Nothofagus* sp. References: (tx) Browne 1984f: 61.
- denticulatus** Browne 1980c: 497. Holotype ♂; Mt. Maunganui (New Zealand) to Kawasaki (Japan), imported; BMNH, London. Distribution: New Zealand. References: (tx) Browne 1980c: 497.
- deperditus** Schedl 1975f: 384. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien. Distribution: New Guinea. References: (tx) Schedl 1975f: 384.
- priscus** Schedl 1975f: 375 (*Crossotarsus*). Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection NHMW, Wien. Synonymy: Roberts 1989: 285. References: (tx) Roberts 1989: 285; Schedl 1975f: 375.
- deplanatus** Wood 1972c: 245. Holotype ♂; La Carbonera Experimental Forest, 50 km W Merida, Merida, Venezuela; Wood Collection. Distribution: South America (Venezuela). Hosts: *Prunus sphaerocarpa*. References: (tx) Wood, S. L. 1972c: 245.
- depressus** Schedl 1979g: 110. Holotype ♀; Papua, Bulolo, Morobe District; Schedl Collection in NHMW, Wien. Distribution: New Guinea. References: (tx) Schedl 1939e: 333, 1979g: 110.
- desultor** Schedl 1939k: 719. Holotype ♂; Costa Rica, San Jose; Schedl Collection in NHMW, Wien. Distribution: North America (Costa Rica). References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 21; Schedl 1972f: 234. (tx) Equihua & Atkinson 1987: 21; Schedl 1939k: 719, 1972f: 239, 1978a: 24.
- detectus** Schedl 1976a: 87. Holotype ♂; Porto Velho, Territ. Rondon, Brazil; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1976a: 87.
- devius** Schedl 1976a: 88. Holotype ♂; Brasil, PA, Santarem; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1976a: 88.
- deyrollei** Chapuis 1865: 127. Syntypes ♂ ♀; Toxpan; 1 ♂, 1 ♀ in BMNH, London, 1 ♂, 1 ♀ in IRSNB, Brussels. Distribution: North America (Costa Rica/ Guatemala/ Veracruz in Mexico/ Panama). Hosts: *Theobroma cacao*. References: (ds) Blackwelder 1947: 789; Blandford 1896e; Equihua & Atkinson 1987: 21; Ferrer 1942; Gemminger & Harold 1872: 2698; Nunberg 1972b: 190; Schedl 1933c: 166, 1972f: 239; Strohmeier 1912c: 8, 1914c: 23. (tx) Blandford 1896e: 97; Chapuis 1865: 127; Equihua & Atkinson 1987: 21; Schedl 1935h: 352, 1937d: 37, 1940a: 324, 1960c: 17, 1972f: 239; Strohmeier 1912c: 8, 1914c: 23.
- dibrachiatus** Roberts 1989: 267. Holotype ♂; Papua New Guinea: Divide Logging Area, Bulolo, 1200 m; MHNW, Wien. Distribution: New Guinea. Hosts: *Terminalia complanata*.

References: (tx) Roberts 1989: 267.

diductus Chapuis 1865: 139. Holotype ♀; Cayenne; IRSNB, Brussels.

Distribution: South America (Cayenne/ Venezuela).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1972f: 239; Strohmeier 1912c: 8, 1914c: 23. (tx) Chapuis 1865: 139; Schedl 1935h: 353–354, 1960c: 20, 1972f: 239; Strohmeier 1912c: 8, 1914c: 23.

diffidens Schedl 1964g: 251. Holotype ♂; Sarawak, Setapok; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: (ds) Schedl 1971c: 370, 1972f: 185. (tx) Schedl 1964g: 251, 1972f: 185, 1978a: 25.

diffinis Schedl 1972i: 55. Holotype ♀; New Guinea (NE), Wan, Morobe Dist., Kunai Creek, 1250 m; BPBM, Honolulu.

Distribution: New Britain Island, New Guinea.

References: (ds) Schedl 1972f: 200. (tx) Schedl 1972f: 200, 1972i: 55, 1978a: 25.

digestus Schedl 1975g: 255. Holotype ♂; Papua, Porotop, Luth. Miss. Sawmill; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 225, 1978b: 25.

dignatus Schedl 1936c: 236. Syntypes ♂; French Guyana, St-Laurent du Maroni; MNHN, Paris, and Schedl Collection in NHMW, Wien.

Figures: Reichardt 1964b: 87.

Distribution: South America (Brazil/ Cayenne/ Guyana/ Suriname).

Hosts: *Eugenia* sp.

Notes: (1) Schedl 1975a: 25 (restricted type to MNHN, Paris material). (3) Schedl 1976a: 88 (described female).

References: (ds) Blackwelder 1947: 789; Schedl 1960d: 79, 1972a: 205. (tx) Reichardt 1964b: 87–88; Schedl 1936c: 236, 1972f: 205, 1976a: 88, 1978a: 25.

dignus Schedl 1964g: 252. Holotype ♂; Sarawak, Setapok; BMNH, London.

Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).

Hosts: *Castanopsis sumatranus*, *Dipterocarpus borneensis*.

Notes: (3) Schedl 1971c: 389 (described female).

References: (hb) Browne 1961c: 218. (ds) Browne 1961a: 315, 1961c: 218; Schedl 1971c: 366, 1972f: 217. (tx) Schedl 1964g: 252, 1971c: 389, 1972f: 217, 1978a: 25.

dimidiatus Chapuis 1865: 153. Syntypes 2 ♂; Cayenne; 1 in MNB, Berlin, 1 in IRSNB, Brussels (Schedl 1960c: 23).

Distribution: South America (Cayenne/ Colombia/ Venezuela).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1972f: 205;

Strohmeier 1911f: 204, 1912c: 10, 1914c: 24. (tx) Chapuis 1865: 153; Lucas 1920: 521; Schedl 1937d: 37, 1939b: 399, 1960c: 22–23, 1972f: 205; Strohmeier 1912c: 10, 1914c: 24. (ms) Lucas 1920: 521.

dipteroearpi Browne 1956b: 338. Holotype ♂; Borneo: Tatan (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

Hosts: *Dipterocarpus* sp.

References: (tx) Browne 1956b: 338.

discollis Chapuis 1865: 254. Syntypes ♂ ♀; Colombia; 1 in MNB, Berlin, 3 in IRSNB, Brussels, 7 ♂, 1 ♀ in BMNH, London (Schedl 1960c: 48). Distribution: North America (Costa Rica/ Guatemala), South America (Bolivia/ Brazil/ Cayenne/ Colombia).

References: (ds) Atkinson & Equihua 1986a: 422; Blackwelder 1947: 789; Blandford 1896e: 112–113; Equihua 1985: 143; Equihua & Atkinson 1987: 22; Gemminger & Harold 1872: 2698; Numberg 1963c: 106; Sampson 1928a: 3; Schedl 1960a: 77, 1970e: 86, 1971f: 150, 1972f: 187, 1972g: 40, 1976a: 57; Strohmeier 1912c: 16, 1914c: 27. (tx) Blandford 1896e: 112–113; Chapuis 1865: 254; Equihua & Atkinson 1987: 22; Sampson 1928a: 3; Schedl 1933a: 402, 1936c: 248, 1937d: 40, 1939b: 399, 1941e: 156, 1960c: 48, 1972f: 187; Strohmeier 1912c: 16, 1914c: 27.

discipennis Schedl 1975g: 226. Holotype ♀; New Guinea, Wan; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 226, 1978a: 25.

discoidalis Schedl 1933a: 402. Holotype ♂; British Guiana, Essequibo R., Moraballi Creek; BMNH, London.

Distribution: South America (Guyana).

References: (ds) Blackwelder 1947: 789; Schedl 1972f: 187. (tx) Schedl 1933a: 402–403, 1972f: 187.

discolor Blandford 1896e: 101. Holotype ♂; Volcan de Chiriqui, Panama; BMNH, London.

Figures: Blandford 1896e: pl. 4, fig. 9.

Distribution: North America (Panama).

References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 22; Schedl 1972f: 214; Strohmeier 1912c: 9, 1914c: 23. (tx) Blandford 1896e: 101; Equihua & Atkinson 1982: 22; Schedl 1936c: 226, 1972f: 214; Strohmeier 1912c: 9, 1914c: 23.

dissimilis Chapuis 1865: 271. Syntypes ♀; de la Nouvelle Grenada [Colombia]; 1 in BMNH, London (Schedl 1960c: 52).

Distribution: South America (Colombia).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1972f: 231; Strohmeier 1912c: 17, 1914c: 27. (tx) Chapuis 1865: 271; Schedl 1935h: 356, 1960c: 52, 1972f: 231; Strohmeier 1912c: 17, 1914c: 27.

- dissipabilis* Schedl 1934e: 212. Syntypes 2 ♂; Bogota und Neu-Granada; Hamburg Museum, lost, and Schedl Collection in NHMW, Wien.
Distribution: Antilles Islands (Guadeloupe), South America (Brazil/ Colombia/ Ecuador/ Venezuela).
Notes: (1) Schedl 1978a: 25 (citation of holotype invalid).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 231, 1972g: 48, 1976a: 57. (tx) Schedl 1934e: 210, 212, 1962p: 210, 1972f: 231, 1978a: 25.
- perplandus* Schedl 1935i: 46. Lectotype ♂; Venezuela, Etat d'Apure; San Fernando de Apure; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 52. Synonymy: Schedl 1962p: 210.
References: (ds) Blackwelder 1947: 790. (tx) Schedl 1935i: 46, 1962p: 210, 1972f: 231, 1978a: 52.
- distinctipes* Schedl 1972i: 55. Holotype ♂; New Guinea (NW), Wau, Kumai Creek, 1270 m; BPBM, Honolulu.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 200. (tx) Schedl 1972f: 200, 1972i: 55, 1978a: 25.
- ponamae* Schedl 1975f: 392. Holotype ♀; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien. Synonymy: Roberts 1989: 285.
Notes: (3) Schedl 1979g: 96 (treated as a synonym of *apicatus*).
References: (tx) Roberts 1989: 285; Schedl 1975f: 392, 1979g: 96.
- distinguendis* Schedl 1934e: 210. Holotype ♂; Costa Rica, Hamburg-Farm, [Rio] Reventazon, Ebene Limon; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 22; Santoro 1957b: 25; Schedl 1972f: 239. (tx) Equihua & Atkinson 1987: 22; Schedl 1934e: 210–211, 1936c: 224, 1972f: 239, 1927g: 26.
- diversipennis* Schedl 1962p: 210. Holotype ♂; Madras, Kattiyur, Wynaad (Perya); FRI, Dehra Dun, not found in 1951 by SIW.
Distribution: Asia (India).
Hosts: *Dipterocarpus indicus*.
Notes: (1) None of the material borrowed by Schedl from FRI was ever returned to FRI; the Schedl "paratype" may be the missing type (SIW).
References: (ds) Schedl 1972f: 204. (tx) Schedl 1962p: 210, 1972f: 204, 1978a: 26.
- dobunabae* Roberts & Morimoto 1986: 167. Holotype ♂; New Guinea; Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Galbulimima belgravcana*.
References: (tx) Roberts & Morimoto 1986: 167.
- dolabratus* Blandford 1896e: 102. Syntypes 5, ♂ ♀; Bugaba, Panama; BMNH, London.
Figures: Blandford 1896e: pl. 4, fig. 12.
Distribution: North America (Panama).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 22; Schedl 1972f: 239; Strohmeyer 1912c: 9, 1914c: 23. (tx) Blandford 1896e: 102; Equihua & Atkinson 1987: 22; Schedl 1940c: 203, 1972f: 239; Strohmeyer 1912c: 9, 1914c: 23.
- dolus* Schedl 1975f: 385. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 385.
- dorsalis* Schedl 1972g: 77. Holotype ♀; Brasilien, Amazonas, Manaos; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1972g: 77.
- dorsatus* Schedl 1975g: 226. Holotype ♀; New Guinea, Sarmi; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 226, 1978b: 26.
- drescheri* Schedl 1941c: 358. Syntypes ♂; Java, Preanger, G. Tangkoeban Prahoë, 4000–5000 Voet; Kalshoven Collection, and Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Java).
Hosts: *Nyssa javanica*.
Notes: (1) Schedl 1978a: 26 (citation of holotype invalid).
References: (hb) Kalshoven 1960e: 40; Schedl 1972f: 207. (tx) Schedl 1941c: 358, 1965g: 28, 1972f: 207, 1978a: 26.
- dryobalanopsis* Browne 1981b: 603. Holotype ♂; Kuala Madai (Sabah, Borneo) to Ganagori (Japan), imported; BMNH, London.
Distribution: Indonesia (Borneo).
Hosts: Kapur log.
References: (ds) Browne 1985b: 290. (tx) Browne 1981b: 603, 1985b: 293.
- dubiosus* Schedl 1933a: 401. Holotype ♂; Brazil, Rio Janeiro; BMNH, London.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 242. (tx) Schedl 1933a: 401, 1972f: 242.
- duplicatus* Schedl 1979g: 111. Holotype ♂; Onenarou, W. Caledonie; Schedl Collection in NHMW, Wien.
Distribution: New Caledonia.
References: (tx) Schedl 1979g: 111.
- duplosignatus* Schedl 1975f: 386. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 386.

- duriovorus** Browne 1955a: 196. Holotype ♂; Tatan (Sarawak) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
Hosts: *Durio* sp.
References: (tx) Browne 1955a: 196.
- durus** Schedl 1936c: 233. Syntypes ♂; Demerara, Brit. Guyana; Hamburg Museum, lost, and Schedl Collection in NHMW, Wien.
Distribution: South America (Guyana).
Notes: (1) Schedl 1975a: 27 (neotype designation apparently invalid).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 239. (tx) Schedl 1936c: 233, 1939b: 399, 1972f: 239, 1978b: 27.
- dyerae** Browne 1956c: 670. Holotype ♂; Borneo: Sandakan (Sabah) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Sabah in Borneo).
Hosts: *Dyera costulata* log.
References: (tx) Browne 1956c: 670.
- echinatus** Roberts & Morimoto 1986: 167. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Cryptocarya* sp.
References: (tx) Roberts & Morimoto 1986: 167.
- efferratus** Schedl 1935h: 357. Holotype ♀; Brazil; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil/ Cayenne/ Guyana).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 231. (tx) Schedl 1935h: 357–358, 1972f: 231, 1978b: 27.
- effetus** Schedl 1935n: 637. Lectotype ♂; Borneo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 27.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 193. (tx) Schedl 1935n: 637, 1965g: 30, 1972f: 193, 1978a: 27.
- egregius** Schedl 1936c: 229. Holotype ♂; probably Guyana; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne/ Guyana?).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 239. (tx) Schedl 1936c: 228–229, 1972f: 239, 1978b: 27.
- elaboratus** Schedl 1975g: 227. Holotype ♂; New Guinea, Wau; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 227, 1978b: 27.
- elongatus** Chapuis 1865: 141. Syntypes 2 ♀; de Caraccas; 1 in BMNH, London, 1 in IRSNB, Brussels (Schedl 1960c: 20).
Distribution: South America (Brazil/ Venezuela).
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 269S; Schedl 1972f: 239; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 141; Schedl 1935g: 313, 1950d: 11, 1952i: 22, 1960c: 20, 1972f: 239, 1978a: 27; Strohmeier 1912c: 9, 1914c: 23.
- emarginatulus** Browne 1981a: 134. Holotype ♂; Dingalan (Philippines) to Tokyo (Japan), imported; BMNH, London.
Distribution: Philippine Islands.
Hosts: *Shorea* sp.
References: (ds) Schedl 1936d: 6; Swaine 1909: 86. (tx) Browne 1981a: 134; Swaine 1909: 86.
- emdeni** Schedl 1935e: 174. Lectotype ♂; D. Neuguinea, Rawlinson Geb. (Schedl 1978a: 28); Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 28.
Distribution: New Guinea.
References: (ds) Schedl 1936d: 15, 1936g: 514, 1972f: 184. (tx) Schedl 1935e: 174–175, 1940b: 434, 1972f: 184, 1978b: 28.
- enixus** Schedl 1936c: 244. Lectotype ♂; Demerara, St.-Jean du Maroni, St.-Laurent du Maroni, Charvein, Bas Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 28.
Distribution: South America (Brazil/ Cayenne/ Guiana).
References: (ds) Atkinson & Equihua 1986a: 422; Blackwelder 1947: 789; Equihua 1985: 143; Equihua & Atkinson 1987: 22; Schedl 1966f: 92, 1972f: 218, 1973d: 164, 1976a: 57. (tx) Equihua & Atkinson 1987: 22; Schedl 1936c: 244, 1972f: 218, 1978a: 28.
- enormis** Schedl 1970a: 130. Holotype ♂; New Guinea, Okasa Pine Forest, Eastern Highlands District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Cryptocarya* sp.
Notes: (3) Schedl 1972i: 57 (described female).
References: (ds) Schedl 1972f: 217. (tx) Schedl 1970a: 130, 1972f: 217, 1972i: 57, 1978a: 28.
- equadorensis** Schedl 1933a: 396. Holotype ♂; Ecuador, Cachahe, low country; BMNH, London.
Distribution: South America (Ecuador/ Peru).
Notes: (3) Browne 1962e: 646 (placed in *Neotrachystus*).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 229. (tx) Browne 1962e: 646; Schedl 1933a: 396, 1939b: 380, 1972f: 229; Wood, S. L. 1966b: 17.
- manipularis** Schedl 1937c: 14. Lectotype ♂; Peru: Cachahe; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 42. Synonymy: Wood 1966b: 17.
References: (ds) Blackwelder 1947: 789. (tx) Schedl 1937c: 14, 1952b: 72, 1972f: 229, 1978a: 42; Wood, S. L. 1966b: 17.
- errans** (Sampson) 1922b: 145. Holotype ♂; Burma: Mohnyin R., Katha; BMNH, London.
Distribution: Asia (Burma/ Assam, Bengal in India).
Hosts: *Acrocarpus fraxinifolius*, *Bombax*

malabricum, *Buchanania latifolia*, *Careya arborea*, *Eugenia jambolana*, *Ficus infectoria*, *Magnolia cathartii*, *Mesua ferrea*, *Phoebe attenuata*, *Tectona grandis*, *Terminalia myriocarpa*, *Wrightia tomentosa*.

Notes: (3) Schedl 1972f: 206 (treated as a subspecies of *indicus*).

References: (cn) Matlur & Singh 1961b: 13, 34, 93. (ds) Beeson 1941 (1961: 265); Bhasin, Roonwal, & Singh 1958; Matlur & Singh 1961b: 13, 34, 93; Schedl 1969c: 68, 1972f: 206. (tx) Sampson 1922b: 145, 1923c: 72; Schedl 1972f: 206.

erraticus Schedl 1972g: 78. Holotype ♂; Brasilien, Jacareacanga, Para; Alvarenga Collection.

Distribution: South America (Brazil).

References: (tx) Schedl 1972g: 78, 1978a: 28.

eugeniae Schedl 1971c: 390. Holotype ♂; Horai, Haldwani, U. P.; Schedl Collection in NHMW, Wien.

Distribution: Asia (Uttar Pradesh in India).

Hosts: *Eugenia jambolana*.

Notes: (3) Beeson 1941: 344, 1961: 265 (*eugeniae*, nomen nudum). The type specimen obviously came from a FRI, Dehra Dun, loan to Schedl dated approximately 1962, that was not returned. References: (ds) Beeson 1941: 344, 1961: 265; Schedl 1972f: 200. (tx) Schedl 1971c: 390–391, 1972f: 200, 1978b: 28.

eugestus Wood 1966a: 64. Holotype ♂; Volcan Irazu, Cartago Prov., Costa Rica, 7000 feet; Wood Collection.

Figures: Schedl 1972f: 213, Wood 1966a: 67.

Distribution: North America (Costa Rica).

Hosts: *Brunellia costaricensis*, *Quercus* sp.

References: (ds) Equihua & Atkinson 1987: 22; Schedl 1972f: 214. (tx) Equihua & Atkinson 1987: 22; Schedl 1972f: 213–214; Wood, S. L. 1966a: 64, 67.

evanidimerius Roberts & Morimoto 1986: 169. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Syzgium* sp.

References: (tx) Roberts & Morimoto 1986: 169.

evanidus Roberts 1986: 42. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Xanthophyllum papuanum*.

References: (tx) Roberts 1986: 42.

eversus Wood 1972b: 251. Holotype ♂; 8 km S Colonia, Valle de Cauca, Colombia; Wood Collection.

Distribution: South America (Colombia).

Hosts: *Lecythis* sp.

References: (tx) Wood, S. L. 1972b: 251.

exaratus Blandford 1896e: 98. Syntypes 5, ♂ ♀; Guatemala: Cerro Zamil, Panajachel, San Geronimo; BMNH, London.

Figures: Blandford 1896e: pl. 4, fig. 5.

Distribution: North America (Guatemala/ Mexico).

References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 22; Schedl 1972f: 239; Strohmeier 1912c: 9, 1914c: 23. (tx) Blandford 1896e: 98; Equihua & Atkinson 1987: 22; Schedl 1939k: 719, 1955b: 30, 1972f: 239; Strohmeier 1912c: 9, 1914c: 23.

excedens Chapuis 1865: 276. Holotype ♂; de la Nouvelle Guinée, Dorey; BMNH, London (Schedl 1960c: 52).

Distribution: Bismarck Islands, Indonesia (Borneo), New Guinea, Philippine Islands (Luzon, Palawan).

References: (bv) Gray, B. 1974b. (cn) Wylie & Shanahan 1977. (hb) Browne 1941; Gray, B. 1974b; Wylie & Shanahan 1975. (ds) Browne 1966: 239, 1980a: 372; Gemminger & Harold 1872: 2698; Olmo, Yoneyama, & Nakazawa 1987b: 89; Olmo et al. 1988b: 98, 1989: 65; Sampson 1928b: 374; Schedl 1938g: 423, 1966b: 92, 1968f: 535, 1972f: 186, 1975i: 347; Strohmeier 1912c: 17, 1914c: 28. (tx) Browne 1970: 579; Chapuis 1865: 276; Nobuchi 1985: 329; Sampson 1928b: 374; Schedl 1935k: 487, 1936h: 65–66, 1936j: 19–35, 1937d: 42, 1941c: 363, 1960c: 52, 1972f: 186, 1978b: 29; Strohmeier 1912c: 17, 1914c: 28; Wylie & Yule 1977.

excellens Schedl 1973e: 95. Holotype ♂; Sattelberg, Papua New Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1973e: 95, 1978a: 29.

excisus Chapuis 1865: 225. Syntypes ♂ ♀; du Mexique, Toxpan; 1 ♂, 3 ♀ in BMNH, London, 1 ♂, 2 ♀ in IRSNB, Brussels.

Distribution: Antilles Islands (Puerto Rico), North America (Costa Rica/ Guatemala/Veracruz in Mexico/ Panama), South America (Argentina/ Brazil/ Cayenne/ Paraguay).

Hosts: *Inga vera*.

References: (cc) Equihua & Atkinson 1986: 624. (hb) Equihua & Atkinson 1986: 624. (ds) Atkinson & Equihua 1986a: 422, 1988: 104; Atkinson et al. 1986: 46; Blackwelder 1947: 789; Blandford 1896e: 110–111; Equihua 1985: 143; Equihua & Atkinson 1986: 624, 1987: 22; Estrada & Atkinson 1988: 212; Ferrer 1942; Gemminger & Harold 1872: 2698; Martorell 1945: 467; Numberg 1972b: 190; Schedl 1966f: 92, 1972f: 218, 1973d: 163, 1976a: 57, 1978c: 292, 1979e: 58; Strohmeier 1912c: 14, 1914c: 26; Wolcott 1936: 316, 1948: 384. (tx) Blandford 1896e: 110–111; Chapuis 1865: 225; Equihua & Atkinson 1987: 22; Schedl 1936c: 239, 1937d: 40, 1940a: 327, 1941e: 156, 1960c: 41–42, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.

eximius Wood 1972b: 248. Holotype ♂; Colonia Tovar, Aragua, Venezuela; Wood Collection.

Figures: Schedl 1972f: 229, Wood 1966a: 52.

Distribution: South America (Venezuela).

References: (tx) Wood, S. L. 1972b: 248.

exitialis Wood 1966a: 51. Holotype ♂; Bartica triangle, British Guiana; BMNH, London.

Figures: Schedl 1972f: 229; Wood 1966a: 52 (male). Distribution: South America (Cayenne/ Guyana).

Hosts: *Licania* sp., *Poweria engleri*.

References: (ds) Schedl 1972f: 229. (tx) Schedl 1972f: 229; Wood, S. L. 1966a: 51–52.

exitiosus Schedl 1936c: 236. Syntypes ♂ ♀; St. Laurent du Maroni, Nouveau Chantier; MNHN, Paris, and Schedl Collection in NHMW, Wien.

Distribution: South America (Cayenne).

Notes: (1) Schedl 1978a: 29 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 789; Schedl 1972f: 229. (tx) Schedl 1936c: 236, 1972f: 229, 1978a: 29; Wood, S. L. 1966a: 51.

exsuperans Beaver 1990a: 283. Holotype ♂; Thailand, Chiang Mai, Doi Pui; BMNH, London.

Figures: Beaver 1990a: 282 (male declivity).

Distribution: Asia (Thailand).

Hosts: *Quercus* sp.

References: (tx) Beaver 1990a: 283.

extensus Schedl 1979g: 111. Holotype ♂; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 111.

falcatus Strohmeier 1908b: 73. Holotype ♂; Himalaya occidentalis (Kulu); Strohmeier Collection.

Distribution: Asia (Uttar Pradesh in India/ Nepal).

Hosts: *Acer campbelli*, *Ahus nepalensis*, *A. nitida*, *Echinocarpus dasygarpus*, *Elaeocarpus lanceaefolius*, *Evodia fraxinifolia*, *Machilus odoratissima*, *Populus* sp., *Quercus lamellosa*.

Notes: (3) Beeson 1961: 265 (*falcator*, nomen nudum, synonymy in Schedl 1969c: 62).

References: (cn) Kleine 1932a: 310, 402; Roonwal 1954: 58. (ec) Beeson 1921b. (hb) Kleine 1932a: 310, 402. (ds) Beeson 1921b: 24, 1941: 344, 1961: 265; Kleine 1932a: 310, 402; Nunberg 1964a: 236; Roonwal 1954: 58; Schedl 1962f: 166, 1969c: 62, 1972f: 209, 1973b: 212; Strohmeier 1912c: 18, 1914c: 28. (tx) Schedl 1941b: 416, 1969c: 62–63, 1972f: 209; Strohmeier 1908b: 73, 1912c: 18, 1914c: 28.

fallax Schedl 1970b: 367. Holotype ♂; Sabah, Sandakan [Borneo]; PPST, Tokyo.

Distribution: Indonesia (Borneo).

Hosts: Seraya.

References: (ds) Browne 1980d: 491, 1981b: 598; Nobuchi 1977: 133; Schedl 1972f: 212. (tx) Nobuchi 1985: 329; Schedl 1970b: 367, 1972f: 212, 1978a: 30.

fenestrallatus Roberts 1989: 268. Holotype ♀; Papua New Guinea: Divide Logging Area, Bulolo, 1200 m; NHMW, Wien.

Distribution: New Guinea.

Hosts: *Terminalia complanata*.

References: (tx) Roberts 1989: 268–269.

ficus Roberts 1970a: 71. Holotype ♂; Nigeria: Ngel Nyaki For. Res., Mambilla Plateau; MRCB, Tervuren.

Distribution: Africa (Congo/ Nigeria).

Hosts: *Ficus eriobotryoides*, *F. macrosperma*.

References: (ds) Browne 1972c: 100. (tx) Roberts 1970a: 71.

flaris Wood 1972: 246. Holotype ♂; Rancho Grande in Pittier National Park, Aragua, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Eschweilera* sp., *Tabebuia* sp.

References: (tx) Wood, S. L. 1972b: 246.

flipennis Schedl 1979g: 111. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 111.

firmus Schedl 1975g: 227. Holotype ♂; New Guinea, Sepik Area; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 227, 1978a: 30.

flavicornis (Fabricius) 1776: 211. Holotype ♂; America boreali [USA]; UZMC, Copenhagen. Figures: Atkinson 1989c, Coyer et al. 1980: 18, White 1983: 324.

Distribution: North America (Mexico (?)/ Alabama, Arkansas, Florida, Georgia, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, E Texas, Virginia, West Virginia in USA).

Hosts: *Pinus* spp.

Notes: (3) Chapuis 1865: 154 (redescribed, =*bidentatus* Dejean 1837: 333, nomen nudum). The occurrence of this species in Mexico could not be confirmed (SLW).

References: (ay) Hopkins 1894g; Stickney 1921: 15. (bv) Atkinson, Foltz, & Connor 1988; Barker, J. A. 1976; Barker, J. A. & Schmid 1976; Coster 1969; Dixon & Payne 1979b; Inscow 1982; Klassen, Ridgway, & Inscow 1982; Madrid, Vite, & Renwick 1972; Renwick, Vite, & Billings 1977. (cn) Anonymous 1966h; Beal et al. 1952: 114–115; Blackman 1950; Chamberlin 1939: 110; Chellman 1958; Doane et al. 1936; Drooz 1985: 375; Fatzinger 1955b; Hertel, Williams, & Merkel 1977; Hopkins 1894: 277; Hubbard 1897b: 15; Lugg 1886: 36; Mead 1966a: 4; Moore, H. B. & McGraw 1976; Williams, L. H. 1988: 73. (cc) Chamberlin 1939: 110; Coster 1969; Dixon & Payne 1979b; Kinn 1984c; Savely 1939: 336. (hb) Atkinson 1989c; Atkinson, Foltz, & Connor 1988; Baker, W. L. 1972: 272; Beal & Massey 1945; Blackman 1922b: 39–40, 1950; Blandford 1896c: 104; Chamberlin 1939: 110; Coster 1969: 1008–

1011; Dalman 1825; Deyrup & Atkinson 1957a: 64; Doane et al. 1936; Drooz 1955: 375; Hetrick 1967; Hopkins 1894g; Hubbard 1897b: 15; Husson 1955: 350; Moore, H. B. & McGraw 1976; Ostmark 1965; White, R. E. 1983. (**ds**) Anonymous 1966h; Atkinson 1989c; Atkinson et al. 1991: 153; Beal & Massey 1945; Blackman 1922b: 39–40; Blackwelder 1947: 759; Blandford 1896e: 104; Blatchley & Leng 1916: 582; Chamberlin 1939: 110; Chapuis 1865; Deyrup & Atkinson 1957a: 64; Drooz 1955: 375; Equihua & Atkinson 1957: 22; Ferrer 1942; Frost 1964: 144; Gemminger & Harold 1872: 2698; Henshaw 1855: 147, 1895: 44; Illiger 1805: 129; Kirk 1969, 1970; Lacordaire 1866: 391; Leng 1920: 337; Melsheimer 1853: 57; Ostmark 1965; Sampson 1928a: 3; Schedl 1933c: 169, 1972f: 223; Schwarz 1878d: 468; Smith, J. B. 1900: 361, 1910: 400; Spahr 1981: 80; Staines, C. L. 1952; Strohmeier 1912c: 10, 1914c: 24; Swaine 1909: 85; Wood, S. L. 1979a: 2. (**tx**) Atkinson 1989c; Beal & Massey 1945: 63–64; Blackman 1922b: 39–40; Blandford 1895: 93–95, 1896e: 104; Blatchley & Leng 1916: 582; Chamberlin 1939: 110; Chapuis 1865: 154–155; Dalman 1825; Dejean 1837: 333; Eggers 1929e: 42; Equihua & Atkinson 1957: 23; Fabricius 1776: 211–212, 1781: 67, 1787: 36, 1792: 364, 1801: 384; Gmelin 1790: 1601; Goyer et al. 1980: 18; Herbst 1793: 118; Hubbard 1897b: 15; Keler 1927d: 232–236; Lacordaire 1866: 391; LeConte 1868: 151, 1876: 343, 1878a: 465; Lucas 1920: 521; Olivier 1795b: 4, 1808: 4; Sampson 1928a: 3; Saunders 1836: 156; Say 1824: 183; Schedl 1935h: 356, 1937c: 14, 1937d: 38, 1940a: 324, 1972f: 223; Strohmeier 1912c: 10, 1914c: 24; Swaine 1909: 85; White, R. E. 1983: 324; Wood, S. L. 1966a: 61, 1979a: 2. (**ms**) Lucas 1920: 521.

flectus Niisima & Murayama 1931: 197. Holotype ♂; Regeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington, automatic. Distribution: Asia (Fujian in China/ Japan/ Taiwan).

Hosts: *Psidium guajava*, *Styrax formosanum*.

Notes: (1) Originally named as a subspecies of *lepidus*. (3) Schedl 1972f: 210 (treated as a subspecies of *lepidus*).

References: (**ds**) Nobuchi 1967: 26; Schedl 1972f: 210. (**tx**) Niisima & Murayama in Murayama 1931: 197; Schedl 1972f: 210.

lepidus formosanus Niisima & Murayama in Murayama 1925a: 214. Holotype ♂; Regeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington, preoccupied by Murayama 1925a: 215, priority by choice of first revisor.

References: (**tx**) Murayama 1925a: 214–215; Niisima & Murayama in Murayama 1931: 197; Schedl 1972f: 210.

murayamensis Schedl 1941a: 43. Holotype ♂;

Regeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington, automatic. Notes: (1) This is an unneeded replacement name.

References: (**ds**) Nobuchi 1967: 26, 1977: 139; Schedl 1966g: 30. (**tx**) Schedl 1941a: 43, 1941b: 416–417, 1972f: 210.

flexiosus Schedl 1936c: 232. Lectotype ♂; Guyane Française; Riviere Lunier; Schedl Collection in NHMW, Wien.

Distribution: South America (Cayenne).

References: (**ds**) Blackwelder 1947: 759; Schedl 1972f: 239. (**tx**) Schedl 1936c: 228, 232, 1972f: 239, 1978b: 30.

forcipatus Schedl 1962k: 1106. Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien. Figures: Schedl 1972f: 203.

Distribution: Africa (Cameroon/ Nigeria).

Hosts: *Authonotha* sp.

Notes: (3) Roberts 1966b: 630 (described female).

References: (**ds**) Roberts 1965: 188, 1969: 116; Schedl 1972f: 202. (**tx**) Roberts 1966b: 630; Schedl 1962k: 1106, 1972f: 202–203, 1975a: 31.

forficula Chapuis 1865: 283. Syntypes ♂; des îles Moluques, Batchian; BMNH, London (Schedl 1960c: 54).

Figures: Nakashima 1975: 55.

Distribution: Asia (Burma/ Cambodia/ Assam in India/ Malay/ Thailand/ Vietnam), Indonesia (Borneo, Java, Moluccas, Sumatra), New Britain Island, New Guinea, New Ireland Island.

Hosts: *Actinophora fragrans*, *Artocarpus elastica*, *Bruguiera parviflora*, *Chrysophyllum roxburghii*, *Koompassia excelsa*, *K. malaccensis*, *Rhizophora mucronata*, *Shorea leprosula*, *Sinodora* sp., *Spondias mangifera*, *Sterculia foetida*.

References: (**ay**) Nakashima 1975: 6. (**cn**) Mathur & Singh 1960a: 84, 1961a: 66, 1961b: 41. (**cc**) Nakashima 1975: 6. (**hb**) Browne 1935a, 1941, 1961c: 206; Kalshoven 1960c: 36. (**ds**) Beaver & Browne 1975: 308; Beeson 1941 (1961: 265); Browne 1935a, 1961c: 206, 1970: 576, 1972b: 28, 1980d: 491; Froggatt 1936; Gemminger & Harold 1872: 2698; Kalshoven 1960d: 38; Mathur & Singh 1960b: 41, 1961a: 66, 1961b: 41; Nakashima 1975: 3; Nobuchi 1977: 133; Ohno et al. 1987: 90, 1988b: 98, 1989: 65; Ohno, Yoneyama, & Nakazawa 1982a, 1982b: 5, 10, 1987: 95; Sampson 1928a; Schedl 1936d: 7, 1936g: 515, 1962f: 166, 1966g: 35, 1969a: 206, 1972f: 209, 1974c: 262, 1975a: 453, 1975e: 452, 1979a: 161; Strohmeier 1911f: 204, 1912c: 18, 1914c: 28. (**tx**) Chapuis 1865: 283; Nakashima 1975: 55; Nobuchi 1985: 329; Sampson 1928a: 3; Schedl 1936d: 7, 1939b: 400, 1939e: 336, 1941b: 416, 1942c: 165, 1953b: 124, 1954a: 145, 1960c: 54, 1972f: 209; Strohmeier 1911f: 204, 1912c: 18, 1914c: 28; Wylie & Yule 1977.

- forficuloides** Schedl 1942a: 212. Syntypes ♂; Malaya, Selangor, Gap; BMNH, London, and Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
Notes: (1) Schedl 1978a: 31 (type restricted to BMNH, London material).
References: (ds) Browne 1961c: 212; Schedl 1972f: 198. (tx) Schedl 1942a: 212, 1972f: 198, 1978b: 31.
- formosanus** Niisima & Murayama in Murayama 1925: 215. Holotype ♂; Rengeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington.
Distribution: Asia (Japan/Taiwan).
Hosts: *Ficus* sp., *Quercus* sp., *Styrax formosanus*.
Notes: (1) Originally named as a subspecies of *lepidus*.
References: (hb) Murayama 1931b: 197. (ds) Miwa 1931: 266; Murayama 1925a: 215, 1929a: 672, 1931b: 197, 1937c: 578. (tx) Murayama 1925a: 215–216, 1925b: 235; Nobuchi 1967: 11–30; Schedl 1941a: 43, 1960h: 111.
- taiwansis** Schedl 1960i: 111. Holotype ♂; Rengeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington automatic.
Notes: (3) Schedl transferred this species from *Platypus* to *Crossotarsus*, thus creating homonymy; then he transferred it back to *Platypus*; this action occurred after 1960, and because of this the original name is restored (SLW).
References: (ds) Nobuchi 1967: 27, 1973b: 16; Schedl 1972f: 189. (tx) Nobuchi 1973b: 16, pl. 2; Schedl 1960i: 111, 1972f: 189.
- fossilatus** Chapuis 1865: 123. Holotype ♀; Bresil, Ega.; BMNH, London (Schedl 1960c: 16).
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1972f: 239; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 123; Schedl 1960c: 16, 1972f: 239; Strohmeier 1912c: 9, 1914c: 23.
- fracticinctus** Schedl 1975f: 386. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 385.
- fracticornis** (Schedl) 1936e: 49 (*Crossotarsus*). Lectotype ♀; N. Guinea S.E., Moroka, 1300 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 31.
Figures: Schedl 1972f: 221.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 221. (tx) Schedl 1936e: 49–50, 1972f: 221, 1978a: 31.
- fractocostis** Schedl 1972i: 57. Holotype ♂; New Guinea (NE), Wan, Morobe Dist., Kunai Creek, 1250 m; BPBM, Honolulu.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 237. (tx) Schedl 1972f: 237, 1972i: 57, 1978a: 31.
- fragosus** Schedl 1936c: 228. Holotype ♂; Guyana; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne/ Guyana).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 239. (tx) Schedl 1936c: 228, 1972f: 239, 1978b: 31.
- franciai** Browne 1962c: 212. Holotype ♂; Philippines: Laguna, Mt. Maquilang; Forest Products Research Institute, Laguna.
Distribution: Philippine Islands (Luzon).
Hosts: *Shorea* sp.
Notes: (1) Browne 1964: 753 (amended the lapsus calami *franciae* to *franciai*). (3) Browne 1985a: 195 (described female).
References: (ds) Browne 1985a: 190; Schedl 1966b: 93, 1972f: 198. (tx) Browne 1962c: 212, 1964: 753, 1970: 577, 1985a: 195; Schedl 1972f: 198.
- fraterculus** Schedl 1941c: 361. Syntypes ♂ ♀; Java, Mount Gede, 800 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya), Indonesia (Borneo, Java), Nicobar Islands.
Hosts: *Castanea argentea*, *Cinnamomum camphora*, *Endriandra rubescens*, *Marrubium muscosa*, *Shorea leprosula*, *Xanthophyllum* sp.
Notes: (1) Schedl 1978a: 31 (citation of holotype invalid).
References: (hb) Kalshoven 1960c: 39. (ds) Browne 1949b, 1961a: 313, 1961c: 211; Kalshoven 1960d: 39; Schedl 1971f: 150, 1972f: 200. (tx) Browne 1949b; Schedl 1941c: 361, 1942a: 172, 1965g: 28, 1972f: 200, 1978a: 31.
- froggatti** Sampson in Froggatt 1926: 259. Lectotype ♂; Sydney, N.S.W., Australia; BMNH, London.
Figures: Brimblecombe 1951: 2 in reprint.
Distribution: Australia (New South Wales, Queensland).
Hosts: *Araucaria cunninghamii*.
Notes: (1) Froggatt (1926b: 259) published this as a new species under Sampson's name after Sampson had designated it as an undescribed species.
References: (cn) Brimblecombe 1951: 69–105, 1956; Browne 1968b: 570; Froggatt 1926b: 259, 1927. (hb) Brimblecombe 1951: 69–105, 1956; Browne 1968b: 570; Froggatt 1926b: 259, 1927. (ds) Browne 1968b: 570. (tx) Browne 1970: 580; Carne et al. 1980; Sampson in Froggatt 1926b: 259; Schedl 1972f: 184.
- pseudoopacus** Schedl 1936g: 517. Lectotype ♂; New South Wales: Dorrig; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1978a: 58.
References: (ds) Schedl 1936g: 517, 1962i: 74, 1972f: 184. (tx) Browne 1970: 580; Schedl 1936g: 517, 1958i: 215, 1972f: 184, 1978a: 58.
- frontebulbifer** Schedl 1979g: 112. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.

References: (tx) Schedl 1979g: 112.

fugax Schedl 1975g: 228. Holotype ♂; New Guinea, War; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 228, 1975b: 31.

fulgens Schedl 1951i: 97. Syntypes 2 ♂; Saigon; Schedl Collection in NHMW, Wien.

Distribution: Asia (Vietnam).

References: (ds) Schedl 1972f: 198, (tx) Nunberg 1960: 624; Schedl 1951i: 97, 1969a: 216, 1972f: 198, 1975a: 32.

fulgidus Browne 1956b: 340. Holotype ♂; New Guinea: Fakfak to Nagoya (Japan), imported: BMNH, London.

Distribution: New Guinea.

Hosts: *Pinelcodendron* log.

References: (tx) Browne 1956b: 340.

fulmeki (Schedl) 1935j: 1 (*Crossotarsus*). Holotype ♂; Sumatra, Medan; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

References: (ds) Schedl 1972f: 201, (tx) Schedl 1935j: 1, 1972f: 201, 1978a: 32.

furcatus Blandford 1895a: 325. Syntypes ♂ ♀; Ceylon, Dikoya; BMNH, London.

Distribution: Asia (Sri Lanka).

Hosts: *Acacia* spp., *Colophyllum walkeri*, *Cupressus macrocarpa*, *Elacocarpus serratus*, *Eucalyptus robusta*, *Gordonia zeylanica*, *Litsaea zeylanica*, *Michelia nilagirica*.

References: (cn) Mathur & Singh 1960a: 40, 1960b: 20; Roonwal 1954: 19, (ds) Beeson 1941 (1961: 265); Mathur & Singh 1960a: 40, 1960b: 20; Roonwal 1954: 19; Schedl 1959a: 512, 1962f: 166, 1972f: 198, (tx) Blandford 1895a: 325; Schedl 1959a: 512, 1972f: 198.

fuscus Chapuis 1865: 134. Syntypes 2 ♂; du Bresil; 1 in MNB, Berlin, 1 in IRSNB, Brussels (Schedl 1960c: 18).

Distribution: South America (Brazil).

Notes: (3) Schedl 1960c: 18 (described female).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1966f: 92, 1967d: 4, 1972f: 239, 1972g: 48, 1976a: 57; Strohmeier 1912c: 9, 1914c: 23, (tx) Chapuis 1865: 134; Schedl 1936c: 230, 1937d: 37, 1951h: 285, 1952d: 363, 1960c: 18, 1972f: 239, 1975b: 32; Strohmeier 1912c: 9, 1914c: 23.

gabensis Schedl 1974d: 467. Holotype ♂; Block 10, New Guinea Industries logging area, Gabensis, Morobe District, CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Terminalia complanata*.

Notes: (1) Roberts 1989: 269 (originally a subspecies of *cordiger*; raised to species rank).

References: (tx) Roberts 1989: 269; Schedl 1974d: 467, 1978b: 21.

gagates Schedl 1976a: 89. Holotype ♂; Rio Caragnata, Matto Grosso, Brazil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (tx) Schedl 1976a: 89.

galbulimimae Roberts & Morimoto 1986: 169. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Galbulimima* sp.

References: (tx) Roberts & Morimoto 1986: 169.

gemellus (Schedl) 1936e: 48 (*Crossotarsus*). Lectotype ♂; N. Guinea, Moroka, 1300 m; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (ds) Schedl 1972f: 221, (tx) Schedl 1936e: 48–50, 1972f: 221, 1978a: 32.

geminatus Chapuis 1865: 239. Syntypes 2♀, ♀; de la Nouvelle-Hollande; 1 in MNB, Berlin, 1 ♀ in MNHN, Paris.

Distribution: Asia (Malaya), Australia, Bougainville Island, Indonesia (Borneo, Celebes, Sumatra, Sumbawa), New Guinea, Philippine Islands (Luzon), Solomon Islands.

Hosts: *Albizzia* sp., *Cassia javanica*, *Polyosma* sp., *Scaphium affine*.

References: (cn) Froggatt 1926b, (hb) Froggatt 1926b, (ds) Beeson 1938b: 295; Browne 1981a: 126; Choo, Woo, & Kim 1981: 202; Gemminger & Harold 1872: 2698; Nobuchi 1977: 124; Ohno, Yoneyama, & Nakazawa 1987a: 95, 1987b: 90; Ohno et al. 1988b: 98, 1989: 65; Schedl 1937d: 40, 1938g: 423, 1965g: 23, 1966b: 93, 1966g: 33, 1969a: 207, 1969e: 156, 1971c: 367, 1972f: 189; Strohmeier 1912c: 16, 1914c: 26, (tx) Chapuis 1865: 239; Nobuchi 1985: 329; Schedl 1960c: 45–46, 1972f: 189; Strohmeier 1912c: 16, 1914c: 26, *turbatus* Chapuis 1865: 242. Syntypes ♂; de l'île

Lucon, Manille; 1 in MNB, Berlin, 1 in IRSNB, Brussels (Schedl 1960c: 46). Synonymy: Schedl 1972f: 190.

References: (hb) Browne 1935a, 1941, (ds) Baer 1886; Beeson 1941 (1961: 267); Browne 1935a; Gemminger & Harold 1872: 2700; Sampson 1928a: 3; Schedl 1936j: 22, 1962i: 75, 1964c: 305; Strohmeier 1912c: 16, 1914c: 27, (tx) Chapuis 1865: 242; Sampson 1928a: 3; Schedl 1937d: 33, 40, 1938g: 423, 1942c: 164, 1952b: 365, 1955b: 282, 1960c: 45–46, 1972f: 190; Strohmeier 1911b: 17, 1912c: 16, 1914c: 217; Wylie & Yule 1977.

gerstaeckeri Chapuis 1865: 240. Syntypes ♂ ♀; des îles Fiji; 1 ♀ BMNH, London, 1 ♂ IRSNB, Brussels.

Distribution: Australia, Fiji Islands, New Guinea.

Hosts: *Swietenia macrophylla*.

References: (bv) Roberts 1977c, (cn) Lever 1938: 14–18, 1940a: 38; Roberts 1977b: 563, 1977c, (hb) Roberts 1977b: 563, 1977c, (ds) Beeson

- 1938b: 295; Gemminger & Harold 1872: 2698; Roberts 1977b: 563; Schedl 1972f: 202, 1979h: 158; Strohmeyer 1912c: 16, 1914c: 26. **(tx)** Chapman 1865: 240; Fairmaire 1881: 468; Schedl 1941e: 154, 1942c: 163–164, 1950f: 35, 41, 1960c: 45, 1972f: 202; Strohmeyer 1912c: 16, 1914c: 26. **(ms)** Roberts 1977c.
- giluwe** Roberts 1979: 87. Holotype ♂; Papua New Guinea: 14 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Ficins* sp., *Syzygium* sp.
References: **(tx)** Roberts 1979: 87.
- glabratus** Roberts 1972: 69. Holotype ♂; West Cameroon: Buea, 4000 feet; BMNH, London.
Distribution: Africa (Cameroon).
Hosts: *Ficus vogeliana*.
References: **(tx)** Roberts 1972: 69.
- globicollis** Schedl 1979g: 112. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1979g: 112.
- glochideus** Roberts & Morimoto 1986: 170. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Planchonella* sp.
References: **(tx)** Roberts & Morimoto 1986: 170.
- godmani** Blandford 1896c: 101. Syntypes ♂ ♀; Guatemala, Sinanja in Vera Paz; BMNH, London.
Figures: Blandford 1896c: pl. 4, fig. 10.
Distribution: North America (Guatemala).
References: **(ds)** Blackwelder 1947: 789; Equihua & Atkinson 1987: 23; Schedl 1972f: 239; Strohmeyer 1912c: 9, 1914c: 23. **(tx)** Blandford 1896c: 101–102; Equihua & Atkinson 1987: 23; Nunberg 1939: 228, 232; Schedl 1972f: 239; Strohmeyer 1912c: 9, 1914c: 23.
- goilalae** Schedl 1975f: 387. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975f: 387.
- gongylodes** Roberts & Morimoto 1986: 170. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Syzygium* sp.
References: **(tx)** Roberts & Morimoto 1986: 170.
- gotoi** Browne 1981b: 604. Holotype ♂; Muara Sabak (Sumatra) to Ishinomaki (Japan), imported; BMNH, London.
Distribution: Indonesia (Sumatra).
References: **(tx)** Browne 1981b: 604.
- gracilicornis** (Schedl) 1936e: 49 (*Crossotarsus*). Syntypes ♂; N. Guinea S.E., Moroka, 1300 m; MCG, Genova, and Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Notes: (1) Schedl 1978a: 33 (citation of holotype invalid). (3) Schedl 1942c: 200 (described female).
References: **(ds)** Schedl 1972f: 221. **(tx)** Schedl 1936e: 49, 1942c: 200, 1972f: 221, 1978a: 33, 1978b: 33.
- gracilior** Browne 1984f: 63. Holotype ♂; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Nothofagus carrii*.
References: **(tx)** Browne 1984f: 63.
- gracilis** Broun 1893: 1254. Syntypes?; New Zealand; Broun Collection.
Figures: Milligan 1979: fig. 5.
Distribution: New Guinea, New Zealand.
References: **(bv)** Milligan 1974, 1982a; Ytsma 1986. **(cn)** Milligan 1974, 1979, 1982a. **(ec)** Milligan 1974; Payton 1989; Zervos 1980. **(hb)** Milligan 1970a, 1979; Neumann & Harris 1974: 135. **(ds)** Broun 1887; Hutton 1904: 219; Milligan 1979; Schedl 1972f: 203; Strohmeyer 1912c: 19, 1914c: 29. **(tx)** Broun 1893: 1254; Milligan 1979: fig. 5; Milligan, Osorne, & Ytsma 1988; Milligan & Ytsma 1988; Schedl 1972f: 203; Strohmeyer 1912c: 19, 1914c: 29; Ytsma 1988. **(ms)** Ytsma 1986.
- inimicus** Broun 1910: 71. Holotype ♂; New Zealand: Raurimu; Broun Collection. Synonymy: Schedl 1972f: 203.
References: **(tx)** Broun 1910: 71; Schedl 1939e: 362, 1941c: 360, 1972f: 203.
- granarius** Schedl 1952d: 363. Holotype ♂; Paraguay, Hohenau; Schedl Collection in NHMW, Wien.
Distribution: South America (Paraguay).
References: **(ds)** Schedl 1972f: 239. **(tx)** Schedl 1952d: 363, 1972f: 239, 1978a: 33.
- grandiporus** Schedl 1961i: 232. Holotype ♀; Bolivia; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: **(ds)** Schedl 1972f: 242. **(tx)** Schedl 1961i: 232, 1972f: 242, 1978a: 33.
- granifer** Schedl 1942a: 200. Holotype ♂; Malaya, Rotan Tunggal F.R.; BMNH, London.
Distribution: Asia (Malaya).
Hosts: *Vatica* sp.
References: **(ds)** Browne 1961c: 199; Schedl 1972f: 213. **(tx)** Schedl 1942a: 200, 1972f: 213.
- granosus** Browne 1986b: 341. Holotype ♂; New Guinea: Fak fak to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: *Penarahan* (Myristicaceae) log.
References: **(tx)** Browne 1986b: 341.

- gramulipennis** Schedl 1975g: 228. Holotype ♂; Australien, New South Wales, Williams R.; Schedl Collection in NHMW, Wien.
Distribution: Australia (New South Wales), New Guinea.
References: (ds) Schedl 1979a: 159. (tx) Schedl 1975g: 228, 1978a: 33.
- gramulosus** Browne 1984c: 455. Holotype ♂; Manokwari (West Irian) to Nagoya (Japan), imported.
Distribution: New Guinea.
Hosts: *Pometia* sp.
References: (tx) Browne 1984c: 455.
- graji** Schedl 1972i: 59. Holotype, sex?; Namarodu, New Ireland Dist.; CSIRO, Canberra.
Distribution: Bougainville Island, New Ireland Island.
References: (ds) Schedl 1972a: 193. (tx) Schedl 1972f: 193, 1972i: 59, 1978a: 34.
- graji immersus** Schedl 1972i: 60. Holotype ♂; Tonolei, Bougainville Dist., New Ireland; CSIRO, Canberra.
References: (ds) Schedl 1972f: 193. (tx) Schedl 1972f: 193, 1972i: 60, 1978a: 34.
- gregalis** Schedl 1936c: 227. Lectotype ♂; Guyane Française, St.-Laurent du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 34.
Distribution: South America (Cayenne/ Guyana).
References: (ds) Blackvelder 1947: 789; Schedl 1934a: 247, 1972f: 239. (tx) Schedl 1934a: 247, 1936c: 227, 1972f: 239, 1978a: 34.
- grumosus** Roberts & Morimoto 1986: 171. New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Agathis alba*.
References: (tx) Roberts & Morimoto 1986: 171.
- guadeloupensis** Schedl 1935i: 45. Lectotype ♂; Guadeloupe; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 34.
Distribution: Antilles Islands (Guadeloupe).
References: (ds) Blackvelder 1947: 789; Schedl 1972f: 214. (tx) Schedl 1935i: 45, 1972f: 214, 1978a: 34.
- gumicus** Roberts 1986: 51. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Galbulimima* sp.
References: (tx) Roberts 1986: 51.
- gumiensis** Roberts 1989: 270. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Garcinia* sp.
References: (tx) Roberts 1989: 270, 286.
- haagi** Chapuis 1865: 173. Syntypes ♂♀; du Mexique: 1 ♂, 1 ♀ in BMNH, London (Schedl 1960c: 28).
Distribution: North America (Veracruz in Mexico/ Panama).
References: (ds) Blackvelder 1947: 789; Blandford 1896c: 107; Equihua & Atkinson 1987: 23; Ferrer 1942; Gemminger & Harold 1872: 2698; Schedl 1972f: 231; Strohmeier 1912c: 11, 1914c: 24. (tx) Blandford 1896c: 101; Chapuis 1865: 173; Equihua & Atkinson 1987: 23; Schedl 1940a: 324, 1952a: 462, 1960c: 28, 1972f: 231; Strohmeier 1912c: 11, 1914c: 24.
- hamaticeps** Schedl 1942a: 205. Lectotype ♂; Malaya, Negri Sembilan, Pasoh F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 34.
Distribution: Asia (Malaya).
References: (ds) Schedl 1972f: 198. (tx) Schedl 1942a: 205, 1972f: 198, 1978a: 34.
- hamaticollis** Schedl 1942a: 204. Lectotype ♂; Malaya, Perak, Trolak F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 34.
Distribution: Asia (India/ Malaya), Indonesia (Borneo).
Hosts: *Amoora spectabilis*, *Garcinia* sp., *Gonystylus bancanus*.
References: (ds) Browne 1961c: 210; Schedl 1969c: 64, 1972f: 186. (tx) Schedl 1942a: 203, 1972f: 186, 1978b: 34.
- garcinia** Browne 1961a: 314. Holotype ♂; Sarawak: Bako National Park; BMNH, London. Synonymy: Schedl 1972f: 186.
References: (ds) Parsons 1963: 352. (tx) Browne 1961a: 314; Schedl 1972f: 186.
- hamatipennis** Schedl 1970b: 364. Holotype ♂; Malaya, Singapore to Nagoya (Japan), imported; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
Hosts: *Segetir*.
References: (ds) Nobuchi 1977: 115; Schedl 1972f: 220. (tx) Schedl 1970b: 364–365, 1972f: 220, 1978b: 34.
- hamatulus** Schedl 1942a: 204. Holotype ♂; Malaya, Bukit, Kajang F.R.; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
Hosts: *Pseudocugenia* sp.
References: (ds) Browne 1961c: 198, 1980d: 492; Schedl 1972f: 206. (tx) Schedl 1942a: 203–204, 1972f: 206, 1978a: 34.
- hamatus** Blandford 1894d: 138. Syntypes 2 ♂; Yuyama and Miyanoshta, Japan; BMNH, London.
Figures: Nobuchi 1973b: pl. 2.
Distribution: Asia (Japan).
Notes: (3) Murayama 1953: 37 (described female).
References: (ds) Blandford 1894c: 579; Murayama 1929a: 673, 1937c: 579, 1954b: 188; Nobuchi

- 1973b: 12; Schedl 1972f: 210; Strohmeier 1912c: 18, 1914c: 28. (**tx**) Blandford 1894d: 138; Murayama 1925b: 235, 1953: 37, 1954b: 188; Nobuchi 1973b: 12, pl. 2; Schedl 1934f: 1647, 1941b: 416, 1972f: 210; Strohmeier 1912c: 18, 1914c: 28.
- hashimotoi** Browne 1986b: 338. Holotype ♂; Fakfak (New Guinea) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: *Nayatoh* (Sapotaceae) log.
References: (**tx**) Browne 1986b: 338.
- hastulifer** Schedl 1959d: 68. Holotype ♂; Crocodile Is., Millingimbi, N. Australia; F. E. Wilson Collection.
Distribution: Australia (Northern Australia).
References: (**ds**) Schedl 1959d: 67, 1972f: 198. (**tx**) Schedl 1959d: 67–68, 1972f: 198, 1978a: 35.
- hebetatus** Roberts 1986: 43. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (**tx**) Roberts 1986: 43.
- heterodoxus** Schedl 1967f: 162. Holotype ♀; Borneo, Xantus; NHMB, Budapest.
Distribution: Indonesia (Borneo).
References: (**ds**) Schedl 1972f: 210. (**tx**) Schedl 1967f: 162–163, 1972f: 210.
- hians** Chapuis 1865: 167. Syntypes ♀; du Bresil; 1 in BMNH, London, 1 apparently lost with Stettin Museum.
Figures: Bright 1972d: 19.
Distribution: Antilles Islands (Bahamas Islands/ Barbados/ Guadeloupe/ Jamaica), North America (Costa Rica/ Panama), South America (Bolivia/ Brazil/ Cayenne/ Guyana/ Paraguay/ Peru/ Suriname/ Venezuela).
Hosts: *Castilloa costaricensis*.
References: (**ds**) Blackwelder 1947: 789; Bright 1972d: 18; Equihua & Atkinson 1987: 23; Gemminger & Harold 1872: 2698; Schedl 1960a: 76, 1963f: 64, 1966f: 92, 1972f: 231, 1972g: 48, 1973a: 368, 1978c: 292; Strohmeier 1912c: 11, 1914c: 24. (**tx**) Bright 1972d: 18–19; Chapuis 1865: 167; Equihua & Atkinson 1987: 23; Keler 1927a: 235; Schedl 1960c: 27, 1972f: 231; Strohmeier 1912c: 11, 1914c: 24.
- apertus** Chapuis 1865: 169. Syntypes ♂; Guadeloupe; 1 in BMNH, London (Schedl 1960c: 27). Synonymy: Schedl 1972f: 231.
References: (**ds**) Blackwelder 1947: 788; Flentiaux & Salle 1890: 455; Gemminger & Harold 1872: 2697; Schedl 1933c: 171; Strohmeier 1912c: 10, 1914c: 24. (**tx**) Chapuis 1865: 169; Keler 1927a: 235; Schedl 1933c: 171, 1934e: 212, 1941c: 155, 1960c: 27, 1972f: 231; Strohmeier 1912c: 10, 1914c: 24.
- perpusillus** Chapuis 1865: 171. Syntypes ♂ ♀; du Bresil. Ega, ♂ Caracas; 1 ♀ MNB, Berlin, 2 ♂ in BMNH, London. Synonymy: Schedl 1972f: 231.
- References: (**ds**) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Numberg 1963c: 107; Strohmeier 1912c: 12, 1914c: 25. (**tx**) Chapuis 1865: 171; Numberg 1939: 241, 1940: 69; Schedl 1936c: 238, 1937d: 39, 1960c: 28, 1972f: 231; Strohmeier 1912c: 12, 1914c: 25, 1960c: 28.
- hinchuachani** Schedl 1939e: 361. Holotype ♀; Hin-chua-Chan, Malaya; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
References: (**ds**) Schedl 1972f: 194. (**tx**) Schedl 1939e: 361, 1972f: 194, 1978a: 35.
- hintzi** Schaufuss 1897b: 103. Syntypes ♂; Gemein [Zaire?]; Hamburg Museum, lost.
Figures: Lhoste & Roche 1960: fig. 2, Mayne & Donis 1962: 282, Roche & Lost 1960: figs. a–b.
Distribution: Africa (Angola/ Cameroon/ Central African Republic/ Congo/ Ethiopia/ Fernando Po Island/ French Guinea/ Gabon/ Ghana/ Ivory Coast/ Kenya/ Malawi/ Mozambique/ Namibia/ Niger/ Nigeria/ Ruanda/ Senegal/ Sierra Leone/ South Africa/ Spanish Guinea/ Sudan/ Tanzania/ Togo/ Uganda/ Zaire/ Zambia/ Zimbabwe).
Hosts: Browne 1963a: 250–251, Schedl 1962k: 609–616 (many hosts listed).
Notes: (3) Numberg 1967b: 325 (named aberration *niger*, no status).
References: (**ay**) Entwistle 1963b; Farris & Funk 1965: 531; Lhoste & Roche 1959: 76–86, 1960; Mayne & Donis 1960: 95; Roche & Lhoste 1960: 2056. (**bv**) Entwistle 1963b; Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**cn**) Alliot & Ivanov 1950: 163–168; Anonymous 1953j: 25; Beaver 1985a: 69; Browne 1961e: 10, 1965b: 571; Cachan 1957: 5–6, 15, 42–53; Halperin & Menier 1951; Mayne & Donis 1951: 337; Roberts 1961a: 41, 1969: 117; Sallenove 1948: 435; Vrydagh 1947a: 5. (**ec**) Anonymous 1953j: 25; Browne 1961e: 10; Farris & Funk 1965: 531; Jover 1952: 73–81; Mayne & Donis 1951: 337; Roche & Lhoste 1960: 2056; Sallenove 1948: 435; Schedl 1958d: 189, 1962k: 600; Smiley & Moser 1984. (**hb**) Beaver 1985a: 69; Beaver & Loytyniemi 1955b: 116; Browne 1963a: 250, 1965b: 571; Entwistle 1963b; Jones, Roberts, & Baker 1959: 13–54; Jover 1952: Loytyniemi, Beaver, & Loytyniemi 1955: 28; Roberts 1961a; Schedl 1958d: 189, 1962k: 600; Thompson, G. H. 1963: 58; Webb & Jones 1957: 25–41. (**ds**) Amaro & Gouveia 1957: 99–100, 131, 141; Beaver 1985a: 69; Beaver & Loytyniemi 1955b: 116; Browne 1961e: 10, 1963a: 250, 1965b: 571, 1972c: 100, 1975a: 757, 1975b: 396; Cola 1971, 1973; Collart 1934: 249; Ferreira 1965: 1129; Gardner 1957a: 29; Halperin & Menier 1951; Hargreaves 1937: 509; Lee 1971: 30; Mayne & Donis 1962: 282; Menier 1973a; Monteiro Guimaraes 1959: 159; Numberg 1952: 22, 1960a: 301, 1965b: 22; Roberts 1960a: 35–37, 1960e,

- 1960f, 1961a: 39–43, 1961b: 53–60, 1968: 188; Schedl 1933f: 192, 1937f: 15, 1938d: 452, 1959p: 22, 1961d: 178, 1962h: 63, 1962k: 600–601, 1964f: 619, 1964i: 249, 1964j: 42, 1965e: 356, 1965g: 21, 1966a: 276, 1966c: 229, 1966g: 37, 1967e: 220, 1968b: 145, 1969d: 4, 1971e: 1, 1971f: 150, 1971g: 196, 1972c: 284, 1972f: 231, 1972k: 296, 1975h: 351, 1977d: 279, 1979b: 416, 1982: 279; Spahr 1981: 50; Strohmeier 1912c: 11, 1914c: 24; Thompson, C. H. 1963: 58; Vrydagh 1947a: 8; Wood, S. L. 1957e: 1272. **(tx)** Eggers 1936b: 33; Lhoste & Roche 1960: fig. 2; Mayne & Donis 1962: 282; Numberg 1952: 22, 1962: 325, 1967b: 325; Powell, W. 1987: 28; Roche & Lhoste 1960: fig. a, b; Schaufuss 1897b: 103; Schedl 1933f: 192, 1936e: 51, 1936f: 102, 1937d: 39, 1937f: 15–17, 1939g: 167, 1941d: 351, 1941e: 155, 1950c: 205, 1950d: 4, 10, 14, 18, 1950e: 212, 1951e: 39, 1951g: 1104, 1952g: 53, 1952j: 5, 1953g: 243, 1954d: 874, 1954e: 48, 58, 1955c: 31, 1955d: 268–271, 1955i: 213, 1962k: 600, 1972f: 231; Strohmeier 1912c: 11, 1912f: 78, 1914c: 24.
- dispar* Schaufuss 1897b: 108. Syntypes ♂ ♀; Gabun; Hamburg Museum, lost. Synonymy: Schedl 1937d: 39.
- References: **(cc)** Hughes & Jackson 1958: 57; Krezal 1959: 611; Vitzthum 1923: 162. **(hb)** Strohmeier 1911a: 182. **(ds)** Schedl 1933f: 192; Strohmeier 1911a: 182, 1911f: 204, 1912c: 11, 1914c: 24. **(tx)** Jeannel 1949: 989; Schaufuss 1897b: 108; Schedl 1933f: 192, 1937d: 39, 1972f: 231; Strohmeier 1912c: 11, 1912f: 78, 1914c: 24.
- penetralis* Sampson 1922: 140. Syntypes ♂ ♀; Portuguese East Africa: Xinavane; BMNH, London. Synonymy: Schedl 1937d: 39.
- References: **(cn)** Slow 1954. **(ds)** Schedl 1933f: 192. **(tx)** Sampson 1922a: 140, 1924c: 129; Schedl 1933f: 192, 1937d: 39, 1972f: 231.
- hippocrepicus** Roberts 1986: 44. Holotype ♂; New Guinea: Papua, Popodetta; BMNH, London. Distribution: New Guinea.
- Hosts: *Amoora* sp.
- References: **(tx)** Roberts 1986: 44.
- hirtellus** Schedl 1936c: 55. Syntypes ♂; Mentawai Island, Sipora Sereimu; MCG, Genova, and Schedl Collection in NHMW, Wien.
- Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo, Mentawai), Samoan Islands.
- Hosts: *Palaquium* sp.
- Notes: (1) Schedl 1978a: 35 (citation of holotype invalid).
- References: **(hb)** Browne 1961c: 200. **(ds)** Beaver 1976b: 545; Browne 1961a: 312, 1961c: 200, 1980a: 372, 1980d: 381, 1986c: 662; Schedl 1964c: 305, 1972f: 185. **(tx)** Browne 1961a: 312, 1970: 575; Schedl 1936c: 55, 1942a: 172, 200, 1964g: 251, 1972f: 185, 1978a: 35.
- hirtus** Schedl 1955b: 310. Holotype ♂; Dutch New Guinea, Maffin Bay; CAS, San Francisco.
- Distribution: New Guinea, Philippine Islands.
- Hosts: *Diospyros* sp.
- Notes: (1) Browne 1966: 235 (raised from subspecies of *solidus* to species rank). (3) Schedl 1972h: 66 (described female).
- References: **(ds)** Browne 1966: 235; Schedl 1972f: 198. **(tx)** Browne 1966: 235; Nobuchi 1985: 329; Schedl 1955b: 282, 310, 1972f: 198, 1972h: 66–67, 1978a: 35, 67.
- histris** (Schedl) 1969a: 217 (*Platyscapus*). Holotype ♂; Borneo (Sarawak), Lawas to Tokyo (Japan), imported; PPST, Tokyo.
- Figures: Nobuchi 1978a: 80.
- Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).
- Hosts: *Shorea* sp.
- References: **(ds)** Browne 1983a: 555; Nobuchi 1977: 122; Schedl 1972f: 190. **(tx)** Nobuchi 1978a: 80, 1978b: 78; Schedl 1969a: 217–218, 1972f: 190.
- holdhausi** Schedl 1935e: 176. Lectotype ♀; Brasilien; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 35.
- Distribution: South America (Brazil).
- References: **(ds)** Blackwelder 1947: 789; Schedl 1972f: 239. **(tx)** Schedl 1935e: 176, 1972f: 239, 1978a: 35.
- horisheusis** Murayama 1928a: 284. Holotype ♂; Formosa; Horisha; Murayama Collection in USNM, Washington.
- Distribution: Asia (Taiwan).
- References: **(ds)** Browne 1972f: 217; Miwa 1931: 266; Murayama 1929a: 674, 1937c: 579; Nobuchi 1967: 26; Schedl 1972f: 217. **(tx)** Murayama 1928a: 284; Schedl 1972f: 217.
- hospes** Schedl 1964g: 253. Holotype ♂; Sarawak, Semengoh; BMNH, London.
- Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).
- Hosts: *Campnosperma* sp.
- References: **(ds)** Schedl 1972f: 227. **(tx)** Schedl 1964g: 253, 1969b: 232, 1972f: 227, 1978a: 35.
- humilis** Chapuis 1865: 229. Holotype ♀; de Caraccas; lost? (Schedl 1960c: 43).
- Distribution: South America (Cayenne/Venezuela).
- Notes: (3) Schedl 1936c: 241 (described male).
- References: **(ds)** Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26. **(tx)** Chapuis 1865: 229; Schedl 1936c: 241, 1939b: 394, 1960c: 43, 1972f: 218, 1978b: 35; Strohmeier 1912c: 14, 1914c: 26.
- hybridus** Schedl 1935b: 395. Holotype ♂; Philippine Islands; Schedl Collection in NHMW, Wien.
- Distribution: Asia (Burma/ Assam in India/ Malaya/ Pakistan/ Thailand/ Vietnam), Indonesia

- (Java), New Guinea, Philippine Islands (Mindanao), Lesser Sunda Islands (Bali).
 Hosts: *Alphonsea ventricosa*, *Diospyros pseudo-ebenum*, *Meilosoma simplicifolia*, *Odina wodier*, *Shorea leprosula*.
 Notes: (3) Browne 1955: 371 (described female), Beeson 1961: 266 (*seni*, nomen nudum, synonymy in Schedl 1969c: 69).
 References: (**hb**) Browne 1961c: 198; Kalshoven 1960c: 36. (**ds**) Beaver & Browne 1975: 309; Browne 1961c: 198, 1981b: 598; Kalshoven 1960: 36; Schedl 1938g: 423, 1962f: 166, 1962i: 73, 1966b: 93, 1969c: 69, 1972f: 192, 1975a: 453. (**tx**) Beeson 1941 (1961: 266); Browne 1955: 371; Nobuchi 1955: 329; Schedl 1935b: 395–396, 1935g: 423, 1941a: 43, 1953c: 291, 1955b: 102, 1972f: 192, 1978a: 36.
- ignotus** Schedl 1936c: 232. Holotype ♂; Suriname; Schedl Collection in NHMW, Wien.
 Distribution: South America (Suriname).
 References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 239. (**tx**) Schedl 1936c: 227, 232, 1972f: 239.
- imberbis** Roberts 1989: 271. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2000 m; NHMW, Wien.
 Distribution: New Guinea.
 Hosts: *Myristica* sp.
 References: (**tx**) Roberts 1989: 271.
- imitatrix** Schedl 1972f: 195–196. Holotype ♂; Brasilien, Jacareacanga, Para; Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (**ds**) Schedl 1972f: 195, 1973d: 163, 1976a: 57. (**tx**) Schedl 1972f: 195–196 (footnote).
- immunis** Schedl 1959m: 556. Holotype ♂; Matto Grosso: Rio Caraguata; Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (**ds**) Schedl 1966f: 93, 1970e: 86, 1972f: 231, 1972g: 48, 1976a: 59. (**tx**) Schedl 1959m: 556, 1972f: 231, 1978a: 36.
- imparidens** Browne 1981b: 604. Holotype ♂; Kimi (New Guinea) to Yatsushiro (Japan), imported; BMNH, London.
 Distribution: New Guinea.
 References: (**tx**) Browne 1981b: 604.
- impolitus** Roberts 1972: 70. Holotype ♂; West Cameroon: Bafut-Ngemba Forest Reserve, 5000 feet; BMNH, London.
 Distribution: Africa (Cameroon).
 Hosts: *Fagara macrophylla*, *Polyscias fulva*.
 References: (**tx**) Roberts 1972: 70.
- imporcatus** Blandford 1896e: 100. Holotype ♂; Panama, Volean de Chiriqui; BMNH, London.
 Distribution: North America (Panama).
 References: (**ds**) Blackwelder 1947: 789; Equihua & Atkinson 1987: 23; Schedl 1972f: 239; Stroh-
 meyer 1912c: 9, 1914c: 23. (**tx**) Blandford 1896e: 100–101; Equihua & Atkinson 1987: 23; Schedl 1933a: 400, 1972f: 239; Strohmeyer 1912c: 9, 1914c: 23.
- impressifrons** Schedl 1942a: 208. Lectotype ♀; Malaya, Buloh F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 36.
 Distribution: Asia (Malaya).
 Hosts: *Shorea leprosula*.
 References: (**ds**) Browne 1961c: 202; Schedl 1972f: 242. (**tx**) Schedl 1942a: 208, 1972f: 242, 1978a: 36.
- impressus** (Strohmeyer) 1912f: 82 (*Crossotarsus*).
 Syntypes ♂; Ostafrika: MNB, Berlin and Strohmeyer Collection.
 Distribution: Africa (Cameroon/ Congo/ Ethiopia/ Fernando Po Island/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sierra Leone/ South Africa/ Tanzania/ Togo/ Uganda/ Zaire).
 Hosts: Schedl 1962k: 655–656 (45 host species listed).
 References: (**bv**) Loyttniemi, Beaver, & Loyttniemi 1985: 28. (**hb**) Beaver & Loyttniemi 1955b: 116; Loyttniemi, Beaver, & Loyttniemi 1985: 28; Roberts 1969: 117; Schedl 1962k: 653–657; Strohmeyer 1911a: 182. (**ds**) Beaver & Loyttniemi 1955b: 116; Mayne & Donis 1951: 336, 1960: 98, 1962: 285; Schedl 1933f: 203, 1962h: 62–63, 1962k: 653, 1107, 1964f: 619, 1965g: 21, 1967e: 220, 1972f: 215, 1979h: 416; Scott 1936: 33; Strohmeyer 1911a: 182, 1912c: 6, 1914c: 36. (**tx**) Eggers 1936b: 33; Schedl 1933f: 202–203, 1937d: 42, 1938d: 452, 1941d: 415, 1950d: 34, 1950g: 896, 1954d: 874, 1954e: 48–73, 1957d: 119–121, 1965: 21, 1970: 228, 1972f: 215; Strohmeyer 1912c: 6, 1912f: 82–85, 1914c: 36.
- mordax** Sampson 1924c: 131. Holotype ♂; Gold Coast, Aburi; BMNH, London. Synonymy: Schedl 1957d: 119.
 References: (**hb**) Thompson, G. H. 1963: 59. (**ds**) Browne 1981a: 127; Mayne & Donis 1960: 98, 1962: 286; Schedl 1933f: 202, 1938d: 452, 1943c: 70; Thompson, G. H. 1963: 59. (**tx**) Sampson 1924c: 131–132; Schedl 1933f: 202, 1954d: 874, 1954e: 48, 58, 1957d: 118–119, 1972f: 215.
- occidentalis** Schedl 1950d: 34. Holotype ♂; Fernando Po; Deutsche Ent. Inst.
 Notes: (3) Schedl 1972f: 215 (treated as a subspecies of *impressus*).
 References: (**ds**) Schedl 1972f: 215. (**tx**) Schedl 1950d: 34, 1962k: 657, 1972f: 215.
- impressus ater** Schedl 1957d: 121. Lectotype ♂; Embu, Kenya; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 36.
 References: (**ds**) Gardner 1957a: 29; Schedl 1962k: 658, 1972f: 215. (**tx**) Schedl 1955i: 213, 1957d: 121, 1962k: 658, 1972f: 215, 1978a: 36.
- malacanthae** Roberts 1966: 630. Holotype ♂;

- Nigeria: Jemma; BMNH, London. Synonymy: Schedl 1972f: 215.
References: (ds) Browne 1972c: 101; Roberts 1966: 630, 1968: 158; Schedl 1972f: 215.
- xylopiæ* Roberts 1966: 633. Holotype ♂; Nigeria: Obudu Cattle Ranch; BMNH, London. Synonymy: Schedl 1970i: 228.
References: (ds) Roberts 1966: 633, 1968: 159, 1969: 120; Schedl 1970i: 228, 1972f: 215.
- inacessus* Schedl** 1936c: 242. Lectotype ♂; Guyane Française, St.-Laurent du Maroni, Bas Maroni, Nouveau Chantier; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 36.
Distribution: North America (San Luis Potosi in Mexico), South America (Brazil/ Cayenne).
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 23; Schedl 1966f: 93, 1970e: 86, 1972f: 218. (tx) Equihua & Atkinson 1987: 23; Schedl 1936c: 242, 1952e: 123, 1972f: 218, 1978a: 36.
- incertus* Schedl** 1970a: 130. Holotype, ♂; New Guinea, Wabag, Western Highlands District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Casuarina oligodermes*.
Notes: (3) Roberts 1989: 272 (described female).
References: (ds) Schedl 1970a: 130–131, 1972f: 198. (tx) Roberts 1989: 272; Schedl 1970a: 130, 1972f: 198, 1978a: 36.
- heteromorphus* Schedl** 1975f: 388. Holotype ♂; Papua, Wabag, Western Highlands District; Schedl Collection in NHMW, Wien. Synonymy: Roberts 1989: 285.
References: (tx) Roberts 1989: 285; Schedl 1975f: 388.
- incognitus* Schedl** 1958h: 499. Lectotype ♀; Sarawak, Setapok; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 37.
Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).
References: (hb) Browne 1961c: 212. (ds) Browne 1961c: 212; Schedl 1972f: 242. (tx) Schedl 1958h: 499, 1972f: 242, 1978a: 37.
- incompertus* Schedl** 1968d: 15. Holotype ♂; Eden, N.S.W., Australia; NMV, Melbourne.
Figures: Browne 1971a: 49, Neumann & Harris 1974: 133, Schedl 1968d: pl. 6.
Distribution: Australia (New South Wales).
Hosts: *Eucalyptus fastigiata*, *E. gigantea*, *E. obliqua*, *E. siberiana*.
Notes: (1) Browne 1971a: 49 (treated in *Austroplatypus*).
References: (bv) Wright, G. M. & Harris 1974. (cn) Harris, Campbell, & Wright 1976; Neumann & Harris 1974. (hb) Harris, Campbell, & Wright 1976; Neumann & Harris 1974; Neumann & Marks 1976; Wright, G. M. & Harris 1974. (tx) Browne 1971a: 49, 1972a: 180; Carne et al. 1980; Neumann & Harris 1974: 133; Schedl 1968d: 15–17, pl. 6, 1972f: 272, 1978a: 37.
- inconstans* Schedl** 1971c: 391. Holotype ♂; Sumatra, Teping Tingi; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Sumatra).
References: (ds) Schedl 1972f: 186. (tx) Schedl 1971c: 391, 1972f: 186, 1978a: 37.
- incostatus* Schedl** 1972h: 68. Holotype, sex?; Dorrego, N.S.W.; BMNH, London.
Distribution: Australia (New South Wales).
Hosts: *Eucalyptus lacopinea*.
References: (tx) Schedl 1972h: 68.
- indicus* Strohmeier** 1910e: 131. Syntypes ♂; Nilgiri Hills (India orientalis); 1 in Andrews Collection, 2 in Strohmeier Collection.
Distribution: Asia (Bangladesh/ Burma/ Assam, Bengal, Karnataka, Tamil Nadu, Uttar Pradesh in India/ Taiwan/ Tonkin Island in Vietnam), Indonesia (Borneo).
Hosts: *Alphonsea ventricosa*, *Atrocarpus integrifolia*, *Bombax* spp., *Dalbergia assamica*, *Dipterocarpus pilosus*, *Ficus* spp., *Meliosma simplicifolia*, *Millettia atropurpurea*, *Poinciana elata*, *Salmalia insignis*, *S. malabarica*, *Schima wallichii*, *Spondias mangifera*, *Talauma hodgsoni*, *Tetrameles nudiflora*.
Notes: (3) Sampson 1922b: 146 (redescribed).
References: (cn) Mathur & Singh 1961a: 10, 1961b: 4; Roonwal 1954: 60. (ce) Beeson 1921b: 24. (ds) Beeson 1921b: 24, 1941 (1961: 265); Browne 1970: 576, 1980b: 350; Mathew 1987: 188; Mathur & Singh 1961a: 10, 1961b: 4; Nobuchi 1977: 132, 1978a: 26; Roonwal 1954: 60; Schedl 1937f: 15, 1962b: 187, 1962f: 166, 1966g: 30, 1969c: 67, 1972f: 206, 1975a: 455, 1975e: 453; Strohmeier 1912c: 19, 1914c: 29. (tx) Beeson 1937: 89–90; Gardner 1932b; Sampson 1922b: 146, 1923c: 2, 1928a: 1–3; Schedl 1942a: 172, 1972f: 206; Strohmeier 1910e: 131, 1912c: 19, 1914c: 29.
- inermis* Sampson** 1923c: 73. Holotype ♂; Burma: Tenasserim, Thagata (Fao); MCG, Genova.
Distribution: Asia (Burma).
References: (ds) Schedl 1972f: 201. (tx) Sampson 1923c: 73–74; Schedl 1972f: 201.
- infuscatus* Browne** 1970: 579. Holotype ♂; Burma: Mohmyin River, Katha; BMNH, London.
Distribution: Asia (Burma).
Hosts: *Wrightia tomentosa*.
References: (ds) Schedl 1972f: 186. (tx) Browne 1970: 579–580; Schedl 1972f: 186.
- insculptus* Schedl** 1967d: 18. Holotype ♂; Brasilien, Nova Tentonia, Santa Catarina; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Schedl 1972f: 219. (tx) Schedl 1967d: 18, 1972f: 219, 1978a: 37.

- insidiosus** Schedl 1935h: 355. Holotype ♂; Petropolis [Brazil]; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 239. (tx) Schedl 1935h: 353–355, 1952d: 364, 1972f: 239, 1978a: 37.
- insignatus** Schedl 1936c: 231. Holotype ♂; Albina, Dutch Guyana; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil/ Suriname).
Notes: (3) Schedl 1976a: 89 (described female).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 239. (tx) Schedl 1935h: 355, 1936c: 230–231, 1952h: 72, 1972f: 239, 1976a: 89, 1978a: 37.
- insularis** Strohmeier 1914f: 4. Syntypes ♂ ♀; Sumatra; Stettin Museum, lost?, and Strohmeier Collection, Eberswalde.
Distribution: Asia (Malaya), Indonesia (Borneo, Java, Malacca, Sumatra).
Hosts: *Albizzia falcata*, *Diospyros* sp., *Elateriospermum tapos*, *Endospermum malaccense*, *Gnelina arborea*, *Hevea brasiliensis*, *Koompassia malaccensis*, *K. malaccensis*, *Pterocarpus indicus*, *Rhizophora mucronata*, *Scaphium affine*, *Shorea bracteolata*, *S. guiso*.
References: (cn) Supriana et al. 1978; Yunus & Hua 1980: 216. (hb) Browne 1961c: 195. (ds) Beeson 1941 (1961: 265); Browne 1961c: 195, 1980a: 371, 1980d: 491; Ohno, Yoneyama, & Nakazawa 1982b: 10, 1987a: 95, 1987b: 90; Schedl 1936d: 7, 1959c: 168, 1961c: 70, 1971c: 367, 1971f: 150, 1972f: 190; Supriana et al. 1978; Yunus & Hua 1980: 216. (tx) Schedl 1937d: 40, 1939e: 336, 1942a: 172, 1954a: 145, 1958b: 102, 1972f: 190; Strohmeier 1914f: 4–5.
- insulindicus** Schedl 1952k: 164. Lectotype ♂; Malaya, Selangor: Sungei Buloh For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 15, automatic.
Distribution: Asia (Burma/ Malaya), Indonesia (Borneo, Java).
Hosts: *Albizzia falcata*, *A. moluccana*, *Artocarpus scortechnii*, *Alstonia* sp., *Dipterocarpus* sp., *Dryobalanops oblongifolia*, *Durio zibethinus*, *Kopsia flavida*, *Parkia speciosa*, *Persca gratissima*, *Pometia* sp., *Shorea leprosula*, *Whitfordiendron pubescens*.
References: (hb) Browne 1961c: 202; Kalshoven 1960c: 36. (tx) Browne 1961c: 202; Kalshoven 1960d: 202; Schedl 1952k: 164, 1960d: 36, 1960i: 110, 1972f: 220.
- bicornis** Schedl 1939e: 360. Lectotype ♂; Selangor: Sungei Buloh For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 15, preoccupied by Numberg 1939.
Notes: (3) Browne 1955: 370 (described female).
References: (cn) Browne 1949c; Hill, D. S. 1983: 515; Mathur & Singh 1961b: 90; Yunus & Hua 1980: 216. (hb) Browne 1938a, 1941; Hill, D. S. 1983: 515. (ds) Beeson 1941 (1961: 263); Browne 1938a; Hill, D. S. 1983: 515; Mathur & Singh 1961b: 90; Yunus & Hua 1980: 216. (tx) Browne 1955: 370; Schedl 1939e: 360–361 (December), 1941c: 354, 1952k: 164, 1972f: 220, 238, 1978a: 15.
- bicornutus** Numberg 1959c: 168. Lectotype ♂; Selangor: Sungei Buloh For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 15, automatic.
Notes: (1) An unneeded replacement name.
References: (tx) Numberg 1959c: 168; Schedl 1959c: 168, 1960h: 110, 1972f: 220, 1978a: 15.
- intermedius** (Schedl) 1937d: 42 (*Stenoplatypus*). Lectotype ♂; Kamerun, Soppo, 800 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 37.
Distribution: Africa (Cameroon/ Ghana/ Sao Tome Island/ Zaire).
Hosts: Schedl 1962k: 650–651 (29 host species listed).
References: (cn) Anonymous 1953j: 25; Mayne & Donis 1951: 337. (cc) Anonymous 1953j: 25; Mayne & Donis 1951: 337. (hb) Jones, Roberts, & Baker 1959: 13; Schedl 1962k: 648–652; Thompson, G. H. 1963: 59; Webb & Jones 1957: 25–26, 37, 41. (ds) Mayne & Donis 1960: 98, 1962: 285; Schedl 1962k: 648–652, 1964f: 619, 1962h: 63, 1972f: 215; Thompson, G. H. 1963: 59. (tx) Schedl 1937d: 42–43, 1950d: 14, 1954e: 58–62, 1957d: 120, 1962k: 648–651, 1972f: 215, 1978a: 37.
- feae** Schedl 1936e: 61 (*Stenoplatypus*). Lectotype ♂; Is. S. Thome, Agua Ize, 400–700 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 30.
Notes: (1) Schedl 1972f: 215 (treated as a subspecies of *intermedius*).
References: (ds) Schedl 1972f: 215. (tx) Schedl 1936e: 61, 1957d: 120, 1962k: 652, 1972f: 215, 1978a: 30.
- interponens** Schedl 1978c: 308. Holotype ♂; Brasilien, Encruzilhada, 980 m, Bahia; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1978c: 308.
- inutilis** (Schedl) 1935k: 479 (*Crossotarsus*). Lectotype ♂; Luzon, Mountain Province, Benguet, Baguio, 1600 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 38.
Distribution: Philippine Islands (Luzon).
References: (ds) Schedl 1966b: 94, 1972f: 200. (tx) Nobuchi 1985: 329; Schedl 1935k: 479–480, 1939b: 398, 1941c: 361, 1955b: 309, 1965g: 29, 1972f: 200, 1978a: 38.
- inviolatus** Schedl 1936c: 230. Lectotype ♂; Guyane Francaise, Nouveau Chantier; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 38.

- Distribution: South America (Brazil/Cayenne).
Notes: (3) Schedl 1972g: 79 (described female).
References: (ds) Blackwelder 1947: 789; Schedl 1970c: 86, 1972f: 239. (tx) Schedl 1936c: 230, 1972f: 239, 1972g: 79, 1978a: 38.
- iriani Browne** 1980e: 377. Holotype ♂; Kimi (West Irian, New Guinea) to Yatsushiro (Japan), imported; BMNH, London.
Distribution: New Guinea.
References: (tx) Browne 1980e: 377.
- irregularis Schedl** 1967d: 18. Holotype ♂; Brasilien, Utinga bei Belem; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Schedl 1972f: 240. (tx) Schedl 1967d: 18–19, 1972f: 240, 1978a: 38.
- irrepertus Schedl** 1936c: 248. Lectotype ♂; Guyane Francaise, St.-Laurent du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 39.
Distribution: South America (Cayenne).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 240. (tx) Schedl 1936c: 247–248, 1972f: 240, 1978a: 39.
- irruptus Schedl** 1935e: 176. Holotype ♂; Albina: Dutch Guyana (Schedl 1978a: 38); Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil/Suriname).
Notes: (3) Schedl 1976a: 89 (described female).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 240. (tx) Schedl 1935e: 176–177, 1936c: 228–229, 1972f: 240, 1976a: 89, 1978b: 39.
- iraguii Roberts & Morimoto** 1986: 171. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Garcinia* sp.
References: (tx) Roberts & Morimoto 1986: 171.
- jamaicensis Bright** 1972d: 17. Holotype ♂; Hardwar Gap, St. Andrew Parish, Jamaica, 4000'.
Figures: Bright 1972d: 19.
Distribution: Antilles Islands (Jamaica).
References: (tx) Bright 1972d: 17–19; McNamara 1977: 191.
- jansonii Chapuis** 1865: 244. Syntypes ♂ ♀; de la Nouvelle Guinee, des iles Moluques et Celebes; BMNH, London.
Figures: Wood 1960a: 12.
Distribution: Admiralty Islands, Asia (Sri Lanka/Taiwan), Australia, Bismarck Islands, Indonesia (Buru, Celebes, Java, Key, Moluccas, Sumatra, Timor), Micronesia (Caroline Islands, Marshall Islands), New Britain, New Guinea, New Hebrides Islands, Philippine Islands (Luzon, Mindanao, Negros), Seychelles Islands, Solomon Islands.
Hosts: *Albizzia* sp., *Alstonia scholaris*, *Althoffia* sp., *Anisoptera* sp., *Autocephalus* sp., *Araucaria cunninghamii*, *A. huastecii*, *Artocarpus elasticus*, *A. incisus*, *Buchanania* sp., *Carallia* sp., *Cocos nucifera*, *Dalbergia latifolia*, *Dysoxylum* sp., *Endospermum medullulosum*, *Evodia* sp., *Ficus* sp., *Garuga* sp., *Hevea brasiliensis*, *Intsia bijuga*, *Laportea* sp., *Litsea domarensis*, *Macaranga* sp., *Mangifera indica*, *M. minor*, *Millettia dehiscens*, *Myristica* sp., *Orctomeles sumatrana*, *Parkia speciosa*, *Planchonella* sp., *Polyalthia* sp., *Pometia pinnata*, *Pterocymbium* sp., *Sloanea forbesii*, *Spondias* sp., *Sterculia campanulata*, *S. foetida*, *Tectonus grandis*, *Terminalia brassii*, *Theobroma cacao*, *Trema orientalis*, *Xanthophyllum* sp.
References: (bv) Gray, B. 1974b. (cn) Froggatt 1926b; Gray, B. & Wylie 1974; Kalshoven 1924c; Kleine 1932a: 402; Roberts 1987; Wylie & Shanahan 1975. (cc) Nakashima 1975: 5. (hb) Froggatt 1926b; Gray, B. 1968: 307, 1974b; Gray, B. & Wylie 1974; Kalshoven 1924c, 1960c: 34; Kleine 1932a: 310, 402; Roberts 1987; Wylie & Shanahan 1975. (ds) Beeson 1938b: 295; Browne 1966: 234; Chadwick & Nikitin 1968; Choo, Woo, & Kim 1981: 202; Froggatt 1936; Gemminger & Harold 1872: 2698; Gray, B. 1968: 307; Kalshoven 1960d: 35; Kleine 1932a: 310, 402; Nakashima 1975: 3; Nobuchi 1977: 125; Nunberg 1961b: 611; Ohno, Yoneyama, & Nakazawa 1987a: 95, 198b7: 90; Ohno et al. 1988b: 98, 1989: 65; Sampson 1926: 120; Schedl 1936g: 514, 1961c: 72, 1962i: 73, 1964c: 304, 1964d: 213, 1965g: 25, 1966b: 94, 1966g: 33, 1968e: 264, 1972f: 190, 1979a: 159; Strohmeier 1912c: 16, 1914c: 27; Wood, S. L. 1960a: 7. (tx) Chapuis 1865: 244; Nobuchi 1985: 329; Sampson 1923c: 3; Schedl 1933e: 105, 1936e: 51–52, 1937d: 40, 1940b: 434–435, 1941c: 354, 1941e: 154, 1942a: 172, 1942c: 164, 1952b: 365, 1955b: 282–283, 1958i: 215, 1960c: 46, 1972f: 190; Strohmeier 1911b: 26, 1912c: 16, 1914c: 27; Wood, S. L. 1960a: 7, 11–12; Wylie & Yule 1977.
- javanus Browne** 1964: 755. Holotype ♂; Java: Malang; BMNH, London.
Distribution: Indonesia (Java).
References: (ds) Schedl 1972f: 210. (tx) Browne 1964: 755; Schedl 1972f: 210.
- jelskii Nunberg** 1939: 226. Holotype ♂; Amable, Peru; IZW, Warsaw.
Distribution: South America (Peru).
References: (ds) Blackwelder 1947: 789; Schedl 1972f: 240. (tx) Nunberg 1939: 226–232; Schedl 1952h: 72, 1972f: 240.
- juvencus Schedl** 1972h: 68. Holotype ♂; Gogol Island Base, Madang District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Evodia* sp., *Pometia* sp.
Notes: (3) Schedl 1974d: 464 (described female).
References: (ds) Schedl 1972f: 224. (tx) Schedl 1972f: 224, 1972h: 68, 1974d: 464, 1978a: 39.
- kalshoveni Schedl** 1941: 360. Syntypes ♀; Java, Preanger, 4000–5000 voet; Kalshoven Collection, and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java).

Notes: (1) Schedl 1978a: 39 (citation of holotype invalid).

References: (ds) Schedl 1972f: 194. (tx) Schedl 1941c: 360–361, 1972f: 194, 1978a: 39.

katoi Browne 1983a: 568. Holotype ♂; Wapoga (West Irian) to Nagoya (Japan), imported; BMNH, London.

Distribution: New Guinea.

Hosts: Matoa log.

References: (tx) Browne 1983a: 568.

keelungensis Browne 1985b: 294. Holotype ♂; Keelung (Formosa) to Nagoya (Japan), imported; BMNH, London.

Distribution: Asia (Taiwan).

Hosts: *Quercus* sp.

References: (tx) Browne 1985b: 294.

kiushuensis Murayama 1936a: 145. Holotype ♀; Ishikochi [Japan]; Murayama Collection in USNM, Washington.

Distribution: Asia (Japan/ Taiwan).

Hosts: *Quercus gilva*, *Q.* spp.

References: (ds) Murayama 1936a: 139–146, 1937c: 577, 1954b: 205; Nobuchi 1973b: 13; Schedl 1972f: 194. (tx) Murayama 1936a: 139–146, 1954b: 205; Nobuchi 1973b: 13; Schedl 1972f: 194.

klapperichi (Schedl) 1941a: 44 (*Stenoplatypus*). Holotype ♂; Fukien, Kiatun, 2300 m, 27.40 n.Br., 117.4 o.L.; Alexander Koenig Museum, Bonn.

Distribution: Asia (Fujian in China).

References: (ds) Schedl 1972f: 217. (tx) Schedl 1941a: 44, 1972f: 217.

kokodaensis (Schedl) 1935f: 274 (*Crossotarsus*). Lectotype ♀; Papua: Kokoda, 1200 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 40.

Distribution: New Guinea.

References: (ds) Schedl 1972f: 200. (tx) Schedl 1935f: 274–275, 1940b: 434–435, 1941e: 152, 1972f: 200, 1978a: 39.

konincki Chapuis 1865: 128. Syntypes ♂ ♀; de la Nouvelle-Grenade, de Bahia; 1 ♂ in MNB, Berlin, 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 17). Distribution: South America (Argentina/ Brazil/ Magdalena Island in Chile/ Colombia/ Guyana/ Venezuela), Maria Madre Island.

Hosts: *Theobroma cacao*.

References: (cn) Idrobo 1958: 75. (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2695; Santoro 1957b: 25; Schedl 1960a: 77–79, 1972f: 240, 1972g: 48, 1976a: 57; Steinhausen 1956: 49; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 128; Schedl 1936c: 224, 1937d: 37, 1955b: 30, 1960c: 17, 1972f: 240; Steinhausen 1956: 49; Strohmeier 1912c: 9, 1914c: 23.

koryoensis (Murayama) 1930b: 24 (*Crossotarsus*). Holotype ♂; Korea; Murayama Collection in USNM, Washington.

Figures: Nobuchi 1967: pl. 2.

Distribution: Asia (Japan/ Korea/ Taiwan/ Ussuri in E USSR).

Hosts: *Acer* sp., *Carpinus laxiflora*, *Quercus acutissima*, *Q. aliena*, *Q. serrata*.

Notes: (3) Arnoldi et al. 1955 (Far East records of *cylindrus* are this species).

References: (ds) Arnoldi et al. 1955: 735–736; Cho 1957; Choo 1983: 32; Choo, Woo, & Nobuchi 1988a: 133; Ko 1969: 273; Murayama 1934f: 142, 1937b: 372, 375, 1937c: 580; Nobuchi 1967: 25; Schedl 1972f: 219. (tx) Choo 1983: 32; Murayama 1930b: 24, 28–31, 35, 1937b: 372, 375; Nobuchi 1967: pl. 2; Schedl 1934f: 1647, 1935k: 484, 1972f: 219.

kusukusensis Murayama 1956b: 13. Holotype ♂; Kusukus, Formosa; Murayama Collection in USNM, Washington.

Distribution: Asia (Taiwan).

Hosts: *Cryptomeria japonica*.

References: (ds) Nobuchi 1967: 26; Schedl 1972f: 210. (tx) Murayama 1956b: 13; Schedl 1972f: 210.

lablabiae Schedl 1979g: 113. Holotype ♂; NE New Guinea, Umboi I., ca 8 km NW Lab Lab, 300 m; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 113.

labrifrons Schedl 1974d: 467. Holotype ♀; Nanwata Banda logging area, Road 14, Bulolo, Morobe District; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Acanthocephalus cadamba*.

References: (tx) Schedl 1974d: 467, 1978a: 40.

laevis Browne 1980a: 376. Holotype ♂; Bialla (New Britain) to Yatsushiro (Japan), imported.

Distribution: New Britain Island.

References: (ds) Ohno et al. 1988b: 98, 1989: 66. (tx) Browne 1980a: 376.

lafertei Chapuis 1865: 144. Syntypes ♂; de la Colombie; 1 in MNB, Berlin, 2 in BMNH, London, 1 in IRSNB, Brussels (Schedl 1960c: 21).

Distribution: North America (Costa Rica/ "Mexico"/ Panama), South America (Brazil/ Colombia/ Guyana/ Venezuela).

References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1957: 23; Gemminger & Harold 1872: 2699; Schedl 1933c: 166, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 144; Equihua & Atkinson 1957: 23; Schedl 1937d: 37, 1940c: 203–204, 1960c: 21, 1971f: 153, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23; Wood, S. L. 1966a: 59.

parysi Chapuis 1865: 145. Syntypes ♂; de la Colombie; 2 in IRSNB, Brussels (Schedl 1960c: 21). Synonymy: Schedl 1960c: 21.

References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865:

145; Schedl 1960c: 21, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.

laminatus Schedl 1964m: 316. Syntypes ♂; Tanandranana, Tamatave [Madagascar]; IRSM, Madagascar, and Schedl Collection in NHMW, Wien.
Distribution: Madagascar.

Notes: (1) Schedl 1978a: 40 (type restricted to IRSM, Madagascar specimens).

References: (ds) Schedl 1972f: 232, 1977b: 244. (tx) Schedl 1961e: 157–158, 1964m: 316, 1972f: 232, 1977b: 244, 1978a: 40.

laosi Schedl 1971c: 391. Holotype ♂; Laos area, Vankly; Schedl Collection in NHMW, Wien.
Distribution: Asia (Laos).

References: (ds) Schedl 1972f: 212. (tx) Schedl 1971c: 391–392, 1972f: 212, 1978a: 40.

lateconcarus Schedl 1942a: 207. Lectotype ♀; Sumatra, Si-Rambe; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 40.
Distribution: Indonesia (Sumatra).

References: (ds) Schedl 1972f: 242. (tx) Schedl 1942a: 207, 1972f: 242, 1978a: 40.

latedeclivis Schedl 1942a: 206. Holotype ♂; Borneo; Schedl Collection in NHMW, Wien, lost.
Distribution: Indonesia (Borneo).

References: (ds) Schedl 1972f: 201. (tx) Browne 1962c: 643; Schedl 1942a: 206, 1972f: 201, 1978a: 40.

latelobatus Browne 1984f: 62. Holotype ♂; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London.
Distribution: New Guinea.

References: (tx) Browne 1984f: 62.

laticollis Chapuis 1865: 250. Syntypes ♂; du Bresil; 2 in BMNH, London (Schedl 1960c: 47).
Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2699; Schedl 1972f: 187; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 250; Schedl 1960c: 47, 1972f: 187; Strohmeier 1912c: 16, 1914c: 27.

latidens Browne 1985b: 296. Holotype ♂; Tg. Mani (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: (tx) Browne 1985b: 296.

latifinis Walker 1859: 261. Holotype ♂; Ceylon; BMNH, London.

Distribution: Asia (Karnataka, Tamil Nadu in India/ Sri Lanka), Indonesia (Borneo).

Hosts: *Bombax malabaricum*, *Diospyros paniculata*, *Doonia zeylanica*, *Eugenia jambolana*, *Ficus asperima*, *Lamaca grandis*, *Mangifera indica*, *Salmaalina malabarica*, *Terminalia bellerica*, *T. paniculata*, *Vateria indica*.

References: (cn) Mathur & Singh 1961a: 13, 1961b: 26; Rau 1936: 35–45; Roonwal 1954: 57. (cc) Mathew 1985. (hb) Speyer 1923: 23. (ds)

Beeson 1961: 265; Blandford 1895a; Gemminger & Harold 1872: 2699; Mathew 1982, 1987: 188; Mathur & Singh 1961a: 13, 1961b: 26; Roonwal 1954: 57; Schedl 1959a: 514, 1962f: 166, 1971a: 278, 1972f: 210; Strohmeier 1912c: 19, 1914c: 29. (tx) Blandford 1895a: 327–328; Motschulsky 1863; Nniberg 1964: 236; Schedl 1937d: 42, 1939b: 398, 1941b: 423–424, 1959a: 514, 1972f: 210; Strohmeier 1912c: 19, 1914c: 29; Walker 1859: 261.

excavatus Chapuis 1865: 280. Holotype ♂; de l'île de Ceylan; BMNH, London (Schedl 1960c: 53). Synonymy: Schedl 1972f: 210.

References: (ds) Gemminger & Harold 1872: 2698; Strohmeier 1912c: 18, 1914c: 28. (tx) Chapuis 1865: 280; Schedl 1937d: 33, 1941b: 417, 1960c: 53, 1972f: 210; Strohmeier 1912c: 18, 1914c: 28.

latreillei Chapuis 1865: 143. Syntypes ♂ ♀; du Mexique; 1 ♂ in MNB, Berlin, 3 ♂, 1 ♀ in BMNH, London, 2 ♂ in IRSNB, Brussels (Schedl 1960c: 21).

Figures: Lara & Shenefelt 1965: 174.

Distribution: North America (Costa Rica/ Guatemala/ Mexico/ Nicaragua), South America (Brazil).

References: (hb) Lara & Shenefelt 1965: 175.

(ds) Blackwelder 1947: 789; Blandford 1896c: 100; Equihua & Atkinson 1987: 24; Ferrer 1942; Gemminger & Harold 1872: 2699; Schedl 1933c: 166, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (tx) Blandford 1896c: 100; Chapuis 1865: 143; Equihua & Atkinson 1987: 24; Lara & Shenefelt 1965: 174; Schedl 1934e: 210, 1940a: 324, 1960c: 21, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.

lawasensis Browne 1970: 578. Holotype ♂; Sarawak: Lawas; BMNH, London.

Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo).

Hosts: *Dryobalanops lanceolata*, *Koordersiodendron pinnatum*.

References: (ds) Schedl 1972f: 211. (tx) Browne 1970: 578–579; Schedl 1972f: 211.

lepidus Chapuis 1865: 282. Syntypes ♂ ♀; des îles Celebes, des îles Moluques, des Philippines; 3 ♂ in BMNH, London, 2 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 54).

Figures: Nobuchi 1967: pl. 2.

Distribution: Africa (Seychelles Islands), Asia (Burma/ India/ Malaya/ Sri Lanka/ Taiwan/ Tonkin Island in Vietnam), Australia, Bismarck Islands, Indonesia (Borneo, Buru, Celebes, Engano, Mentawai, Moluccas, Sumatra), New Guinea, Philippine Islands, Sula, Sunda Islands.

Hosts: *Adinobotrys atropurpureus*, *Albizzia stipulata*, *Anoora rohutaka*, *Artocarpus introcha*, *Camellia sinensis*, *Castanopsis tribuloides*, *Drimycarpus racemosus*, *Elaterspermum tapos*, *Endospermum malaccense*, *Eugenia* spp., *Ficus glomerata*, *Hevea brasiliensis*, *Hibiscus* sp.,

Lannea grandis, *Millettia pendula*, *Parkia speciosa*, *Pithecolobium* sp., *Pometia* sp., *Sarcocephalus cordatus*, *Shorea leprosula*, *Sterculia macrophylla*, *Tectona grandis*, *Vatica lanceaefolia*, *Xanthophyllum glaucum*.

Notes: (3) Schedl 1941b: 424 (review).

References: (cn) Beaver 1958a: 70; Browne 1935b; Corbett 1935: 43–56; Lloyd 1928: 417–426; Mathur & Singh 1960b: 43, 1961b: 71; Yunus & Hua 1980: 217. (hb) Beaver 1958a: 70; Browne 1935a, 1941; Murayama 1931b: 197; Schedl 1977b: 245; Speyer 1923: 23. (ds) Baer 1886: 148; Beaver 1958a: 70; Beeson 1941: 344, 1961: 265; Bhasin, Roonwal, & Singh 1958: 61, 92; Browne 1935a: 1, 1938: 83–84, 1941: 66, 1966: 237, 1980c: 484, 1984c: 449; Choo & Woo 1953; Gemminger & Harold 1872: 2699; Kalshoven 1924b: 357; Leefmans 1927: 6; Lepesme 1947: 649; Mathur & Singh 1960b: 43, 1961b: 71; Mesa 1935: 96; Miwa 1931: 266; Murayama 1925a: 213–214, 1928: 31, 1929a: 673, 1931b: 197, 203, 1934f: 134, 146, 1937c: 579; Nobuchi 1967: 26, 1977: 135; Ohno, Yoneyama, & Nakazawa 1982a: 5, 1987a: 95, 1987b: 90; Ohno, Yoshioka, et al. 1989: 66; Sampson 1914: 382, 1926: 121, 1928a: 3; Schedl 1936d: 7, 1936e: 56, 1936g: 515, 1938g: 423, 1959a: 515, 1966b: 95, 1969d: 9, 12, 1970b: 355, 1972f: 210, 1977b: 245; Strohmeier 1911f: 204, 1912e: 18, 1914c: 28; Yunus & Hua 1980: 217. (tx) Browne 1964: 755; Chapuis 1865: 282; Kalshoven 1960: 38; Murayama 1925a: 213, 1925b: 235; Nobuchi 1967: pl. 2, 1985: 329; Sampson 1914: 382, 1928a: 3; Schedl 1935j: 1, 1937d: 41, 1940b: 434, 1941a: 43, 1941b: 417, 424, 1952b: 366, 1955d: 273, 1959a: 515, 1960c: 54, 1972f: 210, 1977b: 245; Strohmeier 1911b: 17, 27, 1912c: 18, 1914c: 28; Wylie & Yule 1977.

leprosus Browne 1980f: 215. Holotype ♂; Papua, UMLA [Upper Manki logging area], Bulolo, Morobe District; Schedl Collection in NHMW, Wien, automatic.

Distribution: New Guinea.

References: (tx) Browne 1980f: 215.

neotruncatus Schedl 1979g: 114. Holotype ♂; Papua, UMLA, Bulolo, Morobe District; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1972.

References: (tx) Schedl 1978a: 46, 1979g: 114.

lesiniiformis Roberts 1989: 273. Holotype ♂; Papua New Guinea: Divide Logging Area, Bulolo, 1200 m; NHMW, Wien.

Distribution: New Guinea.

Hosts: *Celtis* sp.

References: (tx) Roberts 1989: 273.

levannongi Schedl 1974c: 265. Holotype ♂; Nord-Vietnam, L.Thsi Nuguyen; MHNH, Bucharest.

Distribution: Asia (Vietnam).

References: (tx) Schedl 1974c: 265, 1978a: 41.

lewisi Blandford 1894d: 134. Syntypes 5. ♂ ♀; Miyanoshiita, Kiga, and Yiyama; BMNH, London. Figures: Nakane et al. 1963: pl. 192, Nobuchi 1973b: pl. 2.

Distribution: Asia (Bhutan/ Fujian in China/ Bengal in India/ Japan/ Korea/ Taiwan).

Hosts: *Abies firma*, *Aesculus turbinata*, *Betula grossa*, *B. platyphylla japonicum*, *Castanea crenata*, *Cryptomeria japonica*, *Fagus crenata*, *Kalopanax ricinifolium*, *K. septendobus*, *Quercus acuta*, *Q. acutissima*, *Q. aliena*, *Q. gilva*, *Q. glauca*, *Q. lamellosa*, *Q. mongolica* var. *grosseserrata*, *Q. serrata*.

Notes: (3) Beeson 1941 (1961: 267) *uncacanthurus*, nomen nudum, synonymy in Schedl 1969c: 69.

References: (cn) Kleine 1932a: 402; Shiraki 1952. (cc) Banno, Mikata, & Kodama 1953: 445; Kurenzov 1934a: 58. (hb) Kleine 1932a: 402; Kurenzov 1935a: 44; Murayama 1931b: 197. (ds) Anonymous 1980g; Beeson 1941 (1961: 266); Blandford 1894c; Cho 1957; Choo 1983: 33; Choo & Woo 1985: 163; Kleine 1932a: 402; Ko 1969: 274; Kurenzov 1934a: 58, 1935a: 44, 1936a: 109, 1936b: 350; Miwa 1931: 266; Murayama 1925a: 211, 1929a: 671, 1931b: 197, 1933b: 19, 1934f: 133, 1936a: 139, 1937b: 372, 375, 1937c: 577, 1954b: 188; Nakane et al. 1963: 384; Nobuchi 1967: 26, 1973b: 13; Schedl 1972f: 219, 1975c: 384; Shiraki 1952; Stark 1936e: 142; Strohmeier 1912c: 16, 1914c: 27. (tx) Blandford 1894d: 134; Choo 1983: 33; Murayama 1925a: 211–212, 1925b: 235, 1936a: 139, 146, 1937b: 372, 375, 1954b: 188; Nakane et al. 1963: 384, pl. 192; Nobuchi 1972f: 13, 1973b: pl. 2; Schedl 1934f: 1647, 1969c: 69, 1972f: 219; Strohmeier 1912c: 16, 1914c: 27; Wylie & Yule 1977.

limbatus Chapuis 1865: 253. Syntypes ♂ ♀; Mexique, Toxpan; 1 ♂, 2 ♀ in BMNH, London.

Distribution: North America (Costa Rica/ Guatemala/ Mexico), South America (Brazil).

References: (ds) Blackwelder 1947: 789; Blandford 1896e: 112; Equihua & Atkinson 1987: 24; Ferrer 1942; Gemminger & Harold 1872: 2699; Santoro 1957b: 25; Schedl 1933c: 175, 1972f: 157; Strohmeier 1911f: 204, 1912c: 16, 1914c: 27. (tx) Blandford 1896e: 112; Chapuis 1865: 253; Equihua & Atkinson 1987: 24; Lucas 1920: 522; Schedl 1933a: 402, 1940a: 327, 1960c: 45, 1972f: 157; Strohmeier 1912c: 16, 1914c: 27.

lineaticornis Schedl 1936c: 226. Syntypes ♂; Guyane Française: St.-Laurent du Maroni; MNHN, Paris, and Schedl Collection in NHMW, Wien.

Distribution: South America (Cayenne).

Notes: (1) Schedl 1978a: 41 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 789; Schedl 1972f: 240. (tx) Schedl 1936c: 226, 1972f: 240.

- lineellus** Schedl 1972h: 69. Holotype ♂; Sydney, New South Wales; Schedl Collection in NHMW, Wien.
Distribution: Australia (New South Wales).
References: (ds) Schedl 1972f: 201. (tx) Schedl 1972a: 143–149, 1972f: 201, 1972h: 69, 1978a: 41.
- lingulatus** Roberts & Morimoto 1986: 173. Holotype ♂; Borneo: Niah (Sarawak) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
Hosts: *Lophopetalum* sp. log.
References: (tx) Roberts & Morimoto 1986: 173.
- liraticus** Wood 1966a: 58. Holotype ♂; Rio Danitza, San Jose Prov., Costa Rica, 700 feet; Wood Collection.
Figures: Wood 1966a: 60.
Distribution: North America (Costa Rica).
References: (ds) Equihua & Atkinson 1987: 24; Schedl 1972f: 240. (tx) Equihua & Atkinson 1987: 24; Schedl 1972f: 240; Wood, S. L. 1966a: 58–60.
- liratus** Blandford 1896: 99e. Holotype ♂; Nicaragua, Chontales; BMNH, London.
Distribution: North America (Costa Rica/ El Salvador/ Nicaragua).
Hosts: *Hevea brasiliensis*.
References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 24; Schedl 1972f: 240, 1977e: 44; Strohmeier 1912c: 9, 1914c: 23. (tx) Blandford 1896e: 99–100; Equihua & Atkinson 1987: 24; Schedl 1972f: 240; Strohmeier 1912c: 9, 1914c: 23; Wood, S. L. 1966a: 58.
- longicalcaratus** Roberts 1989: 274. Holotype ♂; Papua New Guinea: Mt. Giluwe, Western Highlands Province, 2900 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Nothofagus pullei*.
References: (tx) Roberts 1989: 274, 286.
- longicollis** Browne 1984c: 455. Holotype ♂; Ambon (Moluccas) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Moluccas).
Hosts: *Canarium* sp., *Shorea* sp.
References: (ds) Ohno, Yoneyama, & Nakazawa 1987a: 95. (tx) Browne 1984c: 455.
- longior** Wood 1966a: 56. Holotype ♂; 16 km SE Cartago, Cartago Prov., Costa Rica; Wood Collection.
Figures: Wood 1966a: 52.
Distribution: Antilles Islands (Puerto Rico), North America (Costa Rica).
References: (ds) Equihua & Atkinson 1987: 24; Schedl 1972f: 205. (tx) Equihua & Atkinson 1987: 24; Schedl 1972f: 205; Wood, S. L. 1966a: 52, 56.
- longipennis** Montrouzier 1861: 265. Holotype ♂; Nouvelle Calédonie; not located.
Distribution: New Caledonia.
References: (ds) Beeson 1938b: 295; Gemminger & Harold 1872: 2699; Lacordaire 1866: 392; Schedl 1972b: 266, 1972f: 196; Strohmeier 1912c: 18, 1914c: 28. (tx) Blandford 1896f: 244; Chapuis 1865: 286; Lacordaire 1866: 392; Montrouzier 1861: 263; Schedl 1939b: 395, 1941e: 156, 1960c: 56, 1972f: 196; Strohmeier 1912c: 18, 1914c: 28.
- longissimus** Roberts 1979: 90. Holotype ♂; Papua New Guinea: 8 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Cloaxylon* sp., *Galbulimima* sp.
References: (tx) Roberts 1979: 90.
- longius** Wood 1966a: 57. Holotype ♂; Volcan Pacaya, Esquintla Prov., Guatemala, 4000 feet; Wood Collection.
Figures: Wood 1966a: 52.
Distribution: North America (Guatemala/ Morelos in Mexico).
Hosts: *Quercus tomentocaulis*.
References: (hb) Atkinson et al. 1986: 12; Burgos & Saucedo 1983: 42. (ds) Atkinson & Equihua 1985a: 103; Atkinson et al. 1986: 12; Burgos & Saucedo 1983: 42; Equihua 1985: 143; Equihua & Atkinson 1987: 42; Schedl 1972f: 205. (tx) Equihua & Atkinson 1987: 24; Schedl 1972: 205; Wood, S. L. 1966a: 52, 57.
- longulus** Chapuis 1865: 158. Syntypes ♂; du Mexique, Toxpan; 2 ♂ in BMNH, London (Schedl 1960c: 24).
Distribution: North America (Costa Rica/ Guatemala/ Chiapas, Veracruz in Mexico), South America (Brazil/ Guyana/ Peru/ Venezuela).
Hosts: *Eschweilera sagotiana*, *Eugenia* sp., *Ocotea rodrigi*, *Tapirira marchandii*.
References: (ds) Blackwelder 1947: 789; Blandford 1896c: 105; Equihua & Atkinson 1987: 24; Ferrer 1942; Gemminger & Harold 1872: 2699; Schedl 1970c: 87, 1972f: 205, 1972g: 50, 1976a: 58; Strohmeier 1912c: 10, 1914c: 24; Wood, S. L. 1966a: 53. (tx) Blandford 1896c: 105; Chapuis 1865: 158; Equihua & Atkinson 1987: 24; Reichardt 1964b: 86; Schedl 1933a: 397, 1937d: 38, 1940a: 324, 1970c: 87, 1972f: 205, 1972g: 50, 1976a: 58; Strohmeier 1912c: 10, 1914c: 24; Wood, S. L. 1966a: 51.
- peregrinus** Schedl 1935h: 356. Holotype ♀; Mexico; Schedl Collection in NHMW, Wien.
Synonymy: Reichardt 1964b: 86.
References: (ds) Blackwelder 1947: 790; Ferrer 1942. (tx) Reichardt 1964b: 86; Schedl 1935h: 356–357, 1940a: 324, 1972f: 205, 1978a: 53.
- lopchuensis** Schedl 1969c: 68. Holotype ♂; India: W. Bengal, Lopchu, 1525 m, Darjeeling; FRI, Dehra Dun, not found in 1981 by SLW.
Distribution: Asia (Bengal in India).
Hosts: *Beilschmiedia sikkimensis*, *Macaranga denticulata*.

Notes: (1) None of the types of Indian species named by Schedl were ever returned to FRI; one non-type is present; this species is not mentioned in Schedl 1978a (SIW). (3) Beeson 1941: 345, 1961: 265 (*lophluensis*, nomen nudum, synonymy in Schedl 1969c: 68).

References: (ds) Beeson 1941: 345, 1961: 265; Bhasin, Roonwal, & Singh 1958; Schedl 1972f: 200. (tx) Schedl 1969c: 68, 1972f: 200.

lophopetali Browne 1985b: 295. Holotype ♂; Niah (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

Hosts: *Lophopetalum* sp.

References: (tx) Browne 1985b: 295.

loriae (Schedl) 1936e: 44 (*Crossotarsus*). Lectotype ♂; N. Guinea, S.E. Moroka, 1300 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 41.

Distribution: New Guinea.

Hosts: *Eudiandra* sp.

Notes: (3) Schedl 1974d: 468 (described female). References: (ds) Schedl 1972f: 200. (tx) Schedl 1936e: 44, 1942a: 207, 1942c: 196, 1972f: 200, 1974d: 468, 1978a: 41.

loricatus (Sampson) 1923c: 71 (*Crossotarsus*). Holotype ♂; Burma: Carin, Ascini Ghacu, 1200–1400 m; MCG, Genova.

Distribution: Asia (Burma/ India/ Malaya/ Vietnam), Indonesia (Borneo, Java, Mentawai, Sumatra), Philippine Islands (Mindoro).

Hosts: *Ficus* sp., *Lucuma malaccensis*, *Pouteria malaccensis*.

Notes: (3) Schedl 1936e: 45 (described female).

References: (hb) Browne 1961c: 200. (ds) Beeson 1941 (1961: 265); Browne 1980b: 381; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1966b: 95, 1971c: 368, 1971f: 150. (tx) Beeson 1937: 91; Nobuchi 1985: 330; Sampson 1923c: 71–72; Schedl 1936e: 45–46, 1937e: 546, 1939e: 335, 1941c: 354, 1942a: 172, 1972f: 237, 1978a: 41.

lucaris Schedl 1979g: 114. Holotype ♂; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 114.

lucasi Chapuis 1865: 243. Holotype ♂; de la Nouvelle Guinée (Dorey); BMNH, London (Schedl 1960c: 46).

Distribution: Asia (Malaya), Indonesia (Java), New Guinea, New Hebrides.

Notes: (3) Schedl 1935f: 275 (described female).

References: (ds) Beeson 1938b: 296; Browne 1961c: 196, 1980d: 492; Gemminger & Harold 1872: 2699; Kalshoven 1924b: 361; Schedl 1936g: 514, 1962i: 73, 1972f: 190; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 243; Sampson 1923c: 3; Schedl 1935f: 275–276, 1939b: 384,

1940b: 434–435, 1960c: 46, 1972f: 190; Strohmeier 1912c: 16, 1914c: 27.

lunatellus Browne 1983a: 569. Holotype ♂; Tg. Usau (West Irian) to Nagoya (Japan), imported. Distribution: New Guinea.

Hosts: Matoa log.

References: (ds) Ohno et al. 1989: 66. (tx) Browne 1983a: 569.

lunatipennis Schedl 1958b: 105. Lectotype ♂; Malaya, Kelantan; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 42.

Distribution: Asia (Malaya).

Hosts: *Sapium baccatum*.

References: (hb) Browne 1961c: 210. (ds) Browne 1961c: 210, 1981a: 126; Schedl 1972f: 201. (tx) Browne 1964: 757; Schedl 1958b: 105, 1972f: 201, 1978a: 42.

lunifer Schedl 1960i: 111. Syntypes ♂; Sumatra, N.-Palawan, Binaluan; MNB, Berlin, and Schedl Collection in NHMW, Wien, automatic.

Distribution: Asia (Burma/ Assam in India/ Malaya), Indonesia (Borneo, Engano, Mentawai, Sumatra), Philippine Islands (Palawan), Si-Oban Island. Hosts: *Azizelia palembanica*, *Albizia* sp., *Coccoceras muticum*, *Shorea beutungeusis*.

Notes: (1) Schedl 1978a: 28 (citation of holotype invalid). (3) Schedl 1961f: 95 (described female). References: (ds) Browne 1966: 237, 1980b: 351, 1980c: 483; Ohno, Yoneyama, & Nakazawa 1982b: 10, 1987b: 90; Schedl 1961f: 95, 1966b: 95, 1971f: 150, 1972f: 201. (tx) Browne 1962e: 654; Schedl 1960i: 111, 1961f: 95, 1972f: 201, 1978a: 28.

emarginatus Schedl 1937d: 34 (*Crossotarsus*).

Syntypes ♂; Sumatra, N.-Palawan, Binaluan; MNB, Berlin, and Schedl Collection in NHMW, Wien, preoccupied by Chapuis 1865.

Notes: (1) Schedl 1978a: 28 (citation of holotype invalid).

References: (cn) Mathur & Singh 1961a: 38. (hb) Browne 1961c: 210. (ds) Beeson 1941 (1961: 253); Browne 1961c: 210, 1983a: 555; Mathur & Singh 1961a: 38; Schedl 1936j: 21. (tx) Schedl 1936e: 43, 1937d: 34, 1939e: 334, 1960c: 35, 1960i: 110–111, 1978a: 28.

luniger Motschulsky 1863: 510. Syntypes 6 ♂ ♀; Montagnes de Nura-Ellia [Ceylon]; IZM, Moscow. Distribution: Africa (Seychelles Islands), Asia (Burma/ Cambodia/ China/ Andaman Islands, Assam, Bengal in India/ Japan/ Malaya/ Sri Lanka/ Taiwan/ Thailand/ Vietnam), Australia (Queensland), Indonesia (Alabal, Borneo, Engano, Java, Mentawai, Moluccas, Sumatra), New Guinea, Philippine Islands (Luzon, Mindoro).

Hosts: *Annoora* sp., *Ehretia acuminata*, *Ficus minnahasse*, *Meliosma simplicifolia*.

References: (hb) Speyer 1923: 23. (ds) Gemminger & Harold 1872: 2699; Schedl 1972f: 242; Strohmeier 1912c: 19, 1914c: 29. (tx) Motschul-

- sky 1863: 510; Schedl 1972f: 242; Strohmeier 1912c: 19, 1914c: 29; Wood, S. L. 1969c: 120.
- caliculus** Chapuis 1865: 280. Syntypes ♂ ♀; Siam; 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 53). Synonymy: Wood 1969c: 120.
Notes: (3) Schedl 1941b: 423 (review).
References: (ay) Nakashima 1975: 5. (cn) Beaver 1988a: 70; Yunus & Hua 1980: 216. (cc) Nakashima 1975: 5; Roberts 1980. (hb) Beaver 1988a: 70. (ds) Beaver 1988a: 70; Beaver & Browne 1975: 307; Blandford 1896a; Browne 1970: 576, 1980d: 491; Gemminger & Harold 1872: 2697; Nakashima 1975: 3; Nobuchi 1977: 137; Ohno, Yoneyama, & Nakazawa 1982a: 4, 1982b: 10, 1987a: 95, 1987b: 89; Schedl 1962b: 187, 1962f: 165, 1962i: 73, 1966b: 90, 1966g: 33, 1971f: 150, 1972f: 209, 1975a: 453, 1979a: 159; Strohmeier 1911f: 204, 1912c: 18, 1914c: 28; Yunus & Hua 1980: 216. (tx) Browne 1964: 755; Chapuis 1865: 280; Murayama 1925b: 234–235, 1928a: 284, 287; Nakashima 1975: 28–33; Nobuchi 1985: 329; Schedl 1937d: 42, 1941b: 417–423, 1952b: 366, 1954a: 145, 1960c: 53, 1972f: 209; Strohmeier 1912c: 18, 1914c: 28; Wood, S. L. 1969c: 120.
- schultzei** Strohmeier 1911b: 26. Syntypes ♂ ♀; Negros, Maao, Philippine Islands; Bureau of Science, Manila. Synonymy: Schedl 1972f: 209.
References: (cn) Ishikura 1966. (ds) Ishikura 1966; Strohmeier 1914c: 28. (tx) Schedl 1936c: 56, 1937d: 42, 1940b: 435, 1941b: 417, 1972f: 209, 1978a: 64; Strohmeier 1911b: 26–28, 1914c: 28.
- luridus** Chapuis 1865: 122. Holotype ♀; du Bresil; Sao Paulo; BMNH, London (Schedl 1960c: 15).
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2699; Schedl 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 122; Lucas 1920: 521; Schedl 1960c: 15, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (ms) Lucas 1920: 521.
- luzonicus** Schedl 1935k: 486. Syntypes ♂ ♀; Luzon, Mountain Province, Benguet, Mount Santo Tomas; W. Schultze Collection, and Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands (Luzon).
Notes: (1) Schedl 1978a: 42 (citation of holotype invalid).
References: (ds) Schedl 1966b: 96, 1972f: 190. (tx) Nobuchi 1985: 329; Schedl 1935k: 486–487, 1939b: 390, 1962b: 193, 1972f: 190, 1978a: 42.
- macer** Browne in Beaver & Browne 1978: 619. Holotype ♂; Malaysia: Penang, Georgetown; BMNH, London.
Figures: Beaver & Browne 1978: 604.
Distribution: Asia (Malaya).
Hosts: Removed from a bird stomach.
References: (tx) Beaver & Browne 1978: 604, 619.
- macroporus** Chapuis 1865: 220. Holotype ♀; de la Colombie; IRSNB, Brussels.
Distribution: South America (Colombia/ Suriname).
Notes: (3) Schedl 1936c: 238 (described male).
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2699; Schedl 1972f: 219; Strohmeier 1912c: 13, 1914c: 25. (tx) Chapuis 1865: 200; Schedl 1936c: 238, 1960c: 40, 1961i: 232, 1972f: 219, 1978a: 42; Strohmeier 1912c: 13, 1914c: 26.
- madagascariensis** Chapuis 1865: 161. Syntypes ♂ ♀; de Madagascar; 1 ♂, 1 ♀ in MNB, Berlin, 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 25).
Distribution: Comoro Islands, Madagascar, Reunion Island, St. Marie Island.
Hosts: *Acridocarpus* sp., *Aphananthe sakalava*, *Canarium* sp., *Cassipourea* sp., *Chrysophyllum boivianum*, *Cinchona succerubra*, *Commiphora* sp., *Dalbergia pterocarpifolia*, *Ekebergia suavis*, *Eugenia* cf. *jambolana*, *Ficus sorocoides*, *Ocotea laevis*, *Olea ambrensis*, *Pachytrophe dimorpha*, *P. obovata*, *Persea americana*, *Pinus* sp., *Plagioscyphus* sp., *Protorhus* sp., *Ravensara pervillei*, *Symphonia* sp., *Vapaca cf. ferruginea*, *Vernonia* sp.
References: (cn) Paulian 1950b: 59. (cc) Schedl 1977b: 247. (hb) Paulian 1950b: 59–60. (ds) Alluaud 1900: 441; Breniere & Dubois 1965: 16, 80; Fairmaire 1892b; Gemminger & Harold 1872: 2699; Heyden 1878: 102; Sampson 1928a: 3; Schedl 1969d: 7, 10, 1970d: 234, 1972f: 233, 1977b: 247; Strohmeier 1912c: 11, 1914c: 25. (tx) Chapuis 1865: 161; Fairmaire 1892b; Lucas 1920: 522; Sampson 1928a: 3; Schaufuss 1897a: 222; Schedl 1937d: 38, 1950h: 105–106, 1951j: 20, 1953d: 70–71, 1960c: 25, 1972f: 233, 1977b: 247; Strohmeier 1912c: 11, 1914c: 25. (ms) Lucas 1920: 522.
- magnus** Browne 1980a: 378. Holotype ♂; Solomon Islands to Taganoura (Japan), imported; BMNH, London.
Distribution: Solomon Islands.
Hosts: *Gonystylus* sp.
References: (ds) Ohno et al. 1988b: 98. (tx) Browne 1980a: 378.
- malaisei** Schedl 1939c: 2. Holotype ♂; NE Burma, Kambaiti, 7000 ft.; NHR, Stockholm.
Distribution: Asia (Burma).
References: (ds) Schedl 1972f: 210. (tx) Murayama 1956b: 13; Schedl 1939c: 2–3, 1972f: 210.
- malignus** Schedl 1936c: 228. Holotype ♂; Dutch Guyana; Schedl Collection in NHMW, Wien.
Distribution: South America (Suriname).
References: (ds) Blackwelder 1947: 789. (tx) Schedl 1936c: 228, 1972f: 240.
- maniensis** Browne 1985a: 197. Holotype ♂; Tg. Mani (Sarawak) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
References: (tx) Browne 1985a: 197.

* *maravignae* **Guerin-Meneville** 1838: 170. Holotype ♂; Sizilianischen (miozanen) Bernstein (Simetit).

Distribution: Europe (Sicilian amber, Miocene).
References: (ds) Handlirsch 1908; Keilbach 1982: 256; Schedl 1960c: 64; Scudder 1891: 569. (tx) Chapuis 1865: 337; Guerin-Meneville 1838: 170; Keler 1928: 3; Schedl 1947a: 14, 1960c: 64. (ms) Keilbach 1982: 256.

marcidus **Blandford** 1896e: 110. Holotype ♂; Panama, Volcan de Chiriqui; BMNH, London.

Figures: Blandford 1896e: pl. 4, fig. 21.
Distribution: North America (Panama).
Hosts: *Quercus* sp.

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 24; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26. (tx) Blandford 1896e: 110; Equihua & Atkinson 1987: 24; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.

margaritaceus **Roberts & Morimoto** 1986: 173. Holotype ♂; New Guinea: Papua, Kaisenik, Wau; BMNH, London.

Distribution: New Guinea.
Hosts: *Nothofagus* sp.
References: (tx) Roberts & Morimoto 1986: 173.

marginatus **Chapuis** 1865: 133. Syntypes ♂; Bresil: Para, Ega; 2 in BMNH, London.

Distribution: South America (Argentina/ Brazil/ Paraguay).
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Schedl 1972f: 240, 1978c: 292; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 133; Schedl 1933a: 400, 1935h: 355, 1936c: 228–232, 1960c: 18, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.

maritimus **Schedl** 1969c: 64. Holotype ♂; Bengal: Sunderbans; FRI, Dehra Dun, not present.

Distribution: Asia (Bengal in India/ Malaya).
Hosts: *Sonneratia apetala*.

Notes: (1) None of the specimens borrowed by Schedl were ever returned to FRI; the Schedl paratypes probably include the type. (3) Beeson 1941: 345, 1961: 265 (*maritimus*, nomen nudum, synonymy in Schedl 1972f: 210).

References: (ds) Beeson 1941: 345, 1961: 265; Choo, Woo, & Kim 1981: 202; Mather & Singh 1961a: 61; Nobuchi 1969a: 206, 1972f: 210. (tx) Schedl 1969c: 64–65, 1972f: 210, 1978a: 43.

matoae **Browne** 1986b: 337. Holotype ♂; New Guinea: Fak fak to Nagoya (Japan), imported; BMNH, London.

Distribution: New Guinea.
Hosts: *Pometia* sp. log.
References: (tx) Browne 1986b: 337.

micrographus **Schedl** 1979g: 114. Holotype ♀; Papua, Upper Manki L.A., Bulolo, Morobe District.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 114.

microlunatus **Browne** 1983a: 571. Holotype ♂; Viru Harbour (Solomon Islands) to Nagoya (Japan), imported; BMNH, London.

Distribution: Solomon Islands.
Hosts: "Basswood" log.
References: (ds) Ohno et al. 1988b: 98. (tx) Browne 1983a: 571.

micrurus **Schedl** 1951i: 96. Syntypes 1 ♂, 1 ♀; Java, Preanger, G. Tangkoeban Praho, 4000–5000 Fuss; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java), New Britain Island.
Hosts: *Dracontomelum maugiferum*.

Notes: (1) Schedl 1978a: 43 (citation of holotype invalid).

References: (ds) Schedl 1972f: 198. (tx) Nunberg 1961: 624; Schedl 1951i: 96, 1962n: 697, 1972f: 198, 1978a: 43.

longicaudatus **Nunberg** 1961b: 623. Holotype ♂; New Britain, Rabaul; BMNH, London. Synonymy: Schedl 1972f: 198.

References: (tx) Nunberg 1961b: 623; Schedl 1972f: 198.

minaciior **Schedl** 1971c: 392. Holotype ♂; Sarawak, Kuching; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).
Hosts: *Vatica* sp.
References: (ds) Schedl 1972f: 213. (tx) Schedl 1971c: 392, 1972f: 213.

minor **Browne** 1980b: 388. Holotype ♂; Vanimo (New Guinea) to Nagoya (Japan), imported; BMNH, London.

Distribution: New Guinea.
References: (ds) Ohno, Yoneyama, & Nakazawa 1987a: 95; Ohno, Yoshioka, et al. 1988b: 98, 1989: 66. (tx) Browne 1980b: 388.

minusculus **Schedl** 1936i: 102. Holotype ♂; Costa Rica; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica), South America (Brazil/ Colombia/ Guyana/ Venezuela).
References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 25; Schedl 1972f: 233, 1976a: 58. (tx) Equihua & Atkinson 1987: 25; Schedl 1936i: 101–102, 1972f: 233, 1978a: 44.

minutissimus **Schedl** 1971e: 13. Holotype ♀; Congo Belge, Yangambi; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).
Hosts: *Macaranga* cf. *spinosa*.
References: (ds) Schedl 1972f: 215. (tx) Schedl 1971e: 13–14, 1972f: 215, 1978b: 44.

mirandus **Schedl** 1969b: 227. Holotype ♂; New Guinea: Karamui, Chimbu District; CSIRO, Canberra.

Distribution: New Guinea
References: (ds) Schedl 1972f: 191. (tx) Schedl 1969b: 227, 1972f: 191, 1978a: 44.

- misoolensis** Browne 1984c: 453. Holotype ♂; Tg. Hoakep (Misool Island, West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: *Maniltoa* sp.
References: (tx) Browne 1984c: 453.
- mixtus** Schedl 1976a: 90. Holotype ♀; Ter. Amapa, Rio Anicohi [Brazil]; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 90.
- mjobergi** Schedl 1965g: 30. Holotype ♂; O. Borneo, Mt. Tibang, 1300 und 1400 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1965g: 24, 1972f: 193. (tx) Schedl 1965g: 24, 30, 1972f: 193, 1978a: 44.
- modestus** Blandford 1894d: 133. Syntypes 4, ♂ ♀; Nikko and Shimidzu Toge [Japan]; BMNH, London. Figures: Nobuchi 1967: pl. 2, 1973b: pl. 2.
Distribution: Asia (Japan/ Taiwan).
Hosts: *Acer palmatum*, *Aesculus turbinata*, *Cercidiphyllum japonicum*, *Fagus crenata*, *Juglans sieboldiana*, *Quercus mongolica* var. *grosseserrata*.
References: (cn) Kleine 1932a: 402. (hb) Kleine 1932a: 402; Murayama 1931b: 195. (ds) Blandford 1894c: 579; Kleine 1932a: 402; Murayama 1929a: 671, 1931b: 195, 1934f: 134, 1936a: 139, 1937c: 578, 1954b: 188; Nobuchi 1967: 26, 1973b: 14; Strohmeier 1912c: 15, 1914c: 27. (tx) Blandford 1894d: 133; Murayama 1925b: 235, 1928a: 283, 1936a: 139, 1954b: 188; Niisima 1910a: 15; Nobuchi 1967: pl. 2, 1973b: 14, pl. 2; Schedl 1934f: 1647, 1972f: 189; Strohmeier 1912c: 16, 1914c: 27.
- modigliani** (Schedl) 1936c: 46 (*Crossotarsus*). Lectotype ♂; Sumatra, Si-Rambe; Schedl Collection in NHMW, Wien, designated by Schedl 1975a: 44.
Figures: Schedl 1972f: 236.
Distribution: Indonesia (Sumatra).
References: (ds) Schedl 1971f: 150, 1972f: 237. (tx) Beeson 1937: 91; Schedl 1936c: 46–47, 1942c: 198, 1972f: 236–237, 1978a: 44.
- morigerus** Schedl 1970a: 131. Holotype ♂; New Guinea, Awande, Eastern Highlands District; CSIRO, Canberra.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 190. (tx) Schedl 1970a: 131–132, 1972f: 190, 1978b: 45.
- morebeensis** Browne 1984f: 65. Holotype ♂; New Guinea: Morobe District, Mount Kaiundi, 2350 m; BMNH, London.
Distribution: New Guinea.
References: (tx) Browne 1984f: 65.
- morosus** Schedl 1975f: 388. Holotype ♂; Kunn, 11 miles from Mt. Hagen, W.H.D., New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 388.
- multisanti** Chapuis 1865: 154. Holotype ♂; Guadeloupe; BMNH, London (Schedl 1960c: 23).
Distribution: Antilles Islands (Guadeloupe).
References: (cn) Cleare 1924: 65–68. (ds) Blackwelder 1947: 790; Fleutiaux & Salle 1890: 458; Gemminger & Harold 1872: 2699; Schedl 1966f: 78, 1970e: 91, 1972f: 205; Strohmeier 1912c: 10, 1914c: 24. (tx) Chapuis 1865: 154; Schedl 1941c: 156, 1960c: 23, 1972f: 205, 1978a: 45; Strohmeier 1912c: 10, 1914c: 24; Wood, S. L. 1966a: 55.
- multiporus** Schedl 1971c: 392. Holotype ♀; Malaya, Pahang; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
References: (ds) Schedl 1972f: 194. (tx) Schedl 1971c: 392–393, 1972f: 194, 1978a: 45.
- muricatus** Roberts 1989: 274. Holotype ♂; Papua New Guinea: Upper Stony Logging Area, Bulolo; NHMW, Wien.
Figures: Roberts 1989: 288.
Distribution: New Guinea.
Hosts: *Syzygium* sp.
References: (tx) Roberts 1989: 274.
- mutatus** Chapuis 1865: 136. Holotype ♀; du Bresil; BMNH, London (Schedl 1960c: 19).
Distribution: South America (Argentina/ Bolivia/ Brazil/ Cayenne/ Paraguay/ Peru/ Uruguay/ Venezuela).
Hosts: *Eucalyptus rostrata*, *Quercus robur*.
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Reichardt 1964b: 86; Schedl 1966f: 93, 1972f: 240, 1972g: 59, 1973d: 163, 1976a: 58; Strohmeier 1910g: 85–88, 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 136; Reichardt 1964b: 86; Schedl 1934e: 211, 1960c: 19, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.
- sulcatus** Chapuis 1865: 137. Syntypes ♂ ♀; du Bresil; 1 ♂ in MNB, Berlin, 3 ♂ in BMNH, London, and 2 ♂, 1 ♀ in IRSNB, Brussels.
Synonymy: Strohmeier 1910g: 85–88, Schedl 1960c: 19.
References: (bv) Santoro 1962c, 1963a. (cn) Bosq 1934: 335; Costa Lima 1956; Ebeling 1950, 1959; Hayward 1941: 91, 1942: 27, 1944: 4; Marelli 1929: 245; Santoro 1957a: 7, 13, 1960a: 965, 1962b: 17–23, 1965b, 1966a, 1967: 70–74; Toscani & De Santis 1960. (cc) Guerrero 1966: 97–103; Santoro 1963a, 1963b. (hb) Costa Lima 1956; Ebeling 1950; Santoro 1957a: 7, 1962b, 1963a, 1965a: 49–58. (ds) Blackwelder 1947: 791; Bosq 1927; Costa Lima 1956; Dejean 1837; Ebeling 1950, 1959; Gemminger & Harold 1872: 2700; Numberg 1958a: 481, 1972b: 190; Ruffinelli & Carbonell 1944: 27; Ruffinelli Rey 1967; Santoro 1957b: 25, 1960a, 1962a; Schedl 1933c: 166, 1960a: 77, 1966f: 81, 94, 1970e: 87, 1972g: 51,

- 1976a: 58, 1978c: 292, 1979e: 58; Strohmeier 1912c: 9, 1914c: 24. **(tx)** Brethes 1909; Chapuis 1965: 137; Costa Lima 1956; Marzelli 1931: 6; Numberg 1958a: 481; Santoro 1965a; Schedl 1935e: 176–177, 1936c: 228, 230, 1936e: 51, 1937d: 37, 1939b: 379, 399, 1939j: 565, 1939m: 170, 1941e: 154, 1948d: 37, 1950i: 148, 1951h: 285, 287, 1952a: 444, 1952e: 123, 1958f: 36, 1960c: 19, 1972f: 240; Strohmeier 1910g: 85–88, 1912c: 9, 1914c: 24.
- plicatus* Brethes 1909: 225. Syntypes ♂ ♀; Buenos Aires, Argentina; Brethes Collection. Synonymy: Bosq 1934: 335.
References: **(cn)** Marelli 1929: 245, 1931: 6; Santoro 1957a: 8. **(hb)** Santoro 1957a: 8. **(ds)** Bruch 1914a; Strohmeier 1912c: 9, 1914c: 23. **(tx)** Bosq 1934: 355; Brethes 1909: 225; Schedl 1939m: 170, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.
- nagakii* Browne 1983a: 567. Holotype ♂; Tanjong Mani (Sarawak) to Toyohashi (Japan), imported; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
Hosts: "Kapur" log.
References: **(tx)** Browne 1983a: 567.
- navarrodeandradei* Marelli 1931: 10. Syntypes?; Brazil; not located.
Distribution: South America (Brazil).
Hosts: *Casuarina* sp., *Eucalyptus* sp.
References: **(cn)** Costa Lima 1956. **(hb)** Costa Lima 1956. **(ds)** Blackwelder 1947: 790; Costa Lima 1936, 1956; Schedl 1972f: 240. **(tx)** Costa Lima 1956; Marelli 1931: 10; Schedl 1972f: 240.
- nebulosus* Roberts 1970: 73. Holotype ♂; Nigeria; Ngel Nyaki Forest Reserve, Mambilla, 1650 m; MRCB, Tervuren.
Distribution: Africa (Nigeria).
Hosts: *Aningeria altissima*.
References: **(ds)** Browne 1972c: 101. **(tx)** Roberts 1970: 73; Schedl 1978b: 46.
- negatus* Schedl 1973e: 96. Holotype ♂; Gogol, Madang District [New Guinea]; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Buchanania* sp., *Ficus* sp., *Myristica* sp., *Spondias* sp.
References: **(tx)** Schedl 1973e: 96, 1978a: 46.
- neglectus* Schedl 1963d: 231. Holotype ♂; Brasilien: Parana, Rondon; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: **(ds)** Schedl 1970f: 87, 1972f: 240. **(tx)** Schedl 1963d: 231, 1972f: 240, 1978a: 46.
- nemorosus* Roberts 1986: 45. Holotype ♂; New Guinea: Papua, Popondetta; BMNH, London.
Distribution: New Guinea.
Hosts: *Myristica* sp.
References: **(tx)** Roberts 1986: 45.
- neopartibilis* Roberts 1989: 276. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo; NHMW, Wien.
Figures: Roberts 1989: 288.
Distribution: New Guinea.
Hosts: *Calbulimima* sp.
References: **(tx)** Roberts 1989: 276, 288.
- neoplicatus* Schedl 1964c: 300. Holotype ♂; Sumatra, Mt. Teleman; RNH, Leiden.
Figures: Schedl 1964c: 301.
Distribution: Indonesia (Sumatra).
References: **(ds)** Schedl 1972f: 191. **(tx)** Schedl 1964c: 300–301, 1972f: 191.
- neosolidus* Schedl 1971e: 14. Holotype ♂; Congo, Camp Putnam, an der Strasse Stanleyville, Irumu; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
Hosts: *Berlinia* sp.
References: **(ds)** Schedl 1972f: 198. **(tx)** Schedl 1971e: 14–15, 1972f: 198, 1978a: 46.
- neotruncatus* Schedl 1972g: 79. Holotype ♂; Brasil, Xingu, Matto Grosso; MZSP, Sao Paulo.
Distribution: South America (Brazil).
References: **(tx)** Schedl 1972g: 79, 1972f: 212.
- niijimai* Murayama 1931b: 197. Syntypes ♂; Nanto [Taiwan]; Murayama Collection in USNM, Washington.
Figures: Nobuchi 1978a: 80.
Distribution: Asia (Taiwan).
Hosts: *Psidium guajava*.
References: **(hb)** Murayama 1931b: 197. **(ds)** Murayama 1931b: 197, 1934f: 134, 1937c: 578; Nobuchi 1967: 26, 1977: 123; Schedl 1972f: 190. **(tx)** Murayama 1931b: 197; Nobuchi 1978a: 80, 1978b: 75; Schedl 1972f: 190.
- nitens* Browne 1980a: 379. Holotype ♂; Ulamona (New Britain) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Britain Island.
References: **(ds)** Ohno, Yoneyama, & Nakazawa 1987a: 95; Ohno et al. 1989: 66. **(tx)** Browne 1980a: 379.
- nitidicollis* Schedl 1935h: 358. Syntypes ♀; Brazil; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: **(ds)** Blackwelder 1947: 790; Schedl 1972f: 240. **(tx)** Schedl 1935h: 358, 1972f: 240, 1978a: 47.
- nitidipennis* Schedl 1948d: 41. Holotype ♂; Guayana Francesa, Riviere Lumier; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
References: **(ds)** Schedl 1972f: 233. **(tx)** Schedl 1948d: 41, 1972f: 233, 1978a: 47.
- nivalis* Sampson 1923c: 73. Holotype ♂; Burma: Tenasserim, Thagata; MCG, Genova.
Distribution: Asia (Burma/ India/ Malaya).
Hosts: *Xanthophyllum* sp.

- References: (**en**) Mathur & Singh 1961b: 94. (**hb**) Browne 1936a, 1941. (**ds**) Beeson 1941 (1961: 266); Browne 1936a, 1961c: 196; Mathur & Singh 1961b: 94; Schedl 1936d: 7, 1972f: 183. (**tx**) Sampson 1923c: 73; Schedl 1972f: 183.
- nocuus** Schedl 1935k: 487. Syntypes ♂ ♀; Luzon, Mountain Province, Mount Santo Tomas; W. Schultze Collection, and Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands (Luzon).
Hosts: *Ficus* sp.
Notes: (1) Schedl 1978: 47 (citation of holotype invalid).
References: (**ds**) Schedl 1966b: 96, 1972f: 186. (**tx**) Nobuchi 1955: 329; Schedl 1935k: 487–488, 1942a: 214, 1972f: 186, 1978a: 47.
- noonadanæ** Browne 1966: 237. Holotype ♂; Philippines: Palawan, Pinigisan, Mantalingajan Range, 600 m; UZMC, Copenhagen.
Distribution: Philippine Islands (Palawan).
References: (**ds**) Schedl 1972f: 210. (**tx**) Browne 1966: 237; Nobuchi 1955: 329; Schedl 1972f: 210.
- norfolkensis** Schedl 1972b: 271. Holotype ♂; Norfolk Island, Middlegate, 120 m, and Bumbora; CSIRO, Canberra.
Distribution: Pacific Ocean Islands (Norfolk Island).
Hosts: *Lagunaria patersonia*.
References: (**tx**) Schedl 1972b: 271, 1978a: 48.
- nothofagus** Roberts 1979: 91. Holotype ♂; Papua New Guinea: 8 km south-west of Onim Hut, Mount Giluwe, 2540 m; BMNH, London.
Figures: Roberts 1979: 93 (male, head of female).
Distribution: New Guinea.
Hosts: *Nothofagus pullei*.
References: (**tx**) Roberts 1979: 91.
- novaeguineensis** Roberts 1989: 277. Holotype ♂; Papua New Guinea: Gumti, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.
Figures: Roberts 1989: 288.
Distribution: New Guinea.
Hosts: *Lithocarpus* sp.
References: Roberts 1989: 277, 288.
- nudatus** Wood 1972b: 243. Holotype ♂; 8 km S Colonia (near Buenaventura), Valle de Cauca, Colombia; Wood Collection.
Distribution: South America (Colombia).
Hosts: *Couma macrocarpa*, *Lecythis* sp., *Sacoglottia procera*.
References: (**tx**) Wood, S. L. 1972b: 243.
- nudiusculus** Roberts 1986: 46. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (**tx**) Roberts 1986: 46.
- nudus** Schedl 1936c: 242. Lectotype ♂; Guyane Française: Camopi, St. Laurent du Maroni, St. Jean du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 48.
Distribution: South America (Cayenne).
References: (**ds**) Blackwelder 1947: 790; Schedl 1972f: 214. (**tx**) Schedl 1936c: 242, 1972f: 214, 1978a: 48.
- obliquecaudatus** Schedl 1936e: 59. Lectotype ♂; New Guinea, Moroka, 1300 m; Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya), New Guinea.
References: (**ds**) Browne 1961c: 210; Schedl 1972f: 211. (**tx**) Numberg 1959: 168; Schedl 1936e: 59–60, 1942a: 210, 1960h: 110, 1972f: 211, 1978b: 48.
- obliquecaudatus** Schedl 1942a: 210. Syntypes ♀; BMNH, London, and Schedl Collection in NHMW, Wien, preoccupied by Schedl 1936.
Notes: (1) Schedl obviously intended to describe the female of this established name, but in so doing inadvertently validated a homonymous synonym.
References: (**tx**) Numberg 1959: 168; Schedl 1936e: 59, 1942a: 210.
- obliquesectus** Schedl 1973f: 77. Holotype ♂; Pengagl, camp east Mt. Wilhelm, Chimbu District, 2770 m; AMNH, New York.
Distribution: New Guinea.
References: (**tx**) Schedl 1973f: 77, 1978a: 48.
- obliquetruncatus** Schedl 1972h: 69. Holotype ♂; Nauwata Banda logging area, Rd 14, Bulolo, M. Dist.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Anthocephalus cadamba*, *Pterygota* sp.
References: (**ds**) Schedl 1972f: 198. (**tx**) Schedl 1972f: 198, 1972h: 69, 1978a: 48.
- obliquus** Wood 1981: 121. Holotype ♂; Neuguinea, Mt. Wilhelm; Schedl Collection in NHMW, Wien, automatic.
Distribution: New Guinea.
References: (**tx**) Wood, S. L. 1981: 121.
- obliquesectus** Schedl 1975g: 229. Holotype ♂; Neuguinea, Mt. Wilhelm; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1973.
References: (**tx**) Schedl 1975g: 229; Wood, S. L. 1981: 121.
- obliteratus** Blandford 1896e: 99. Holotype ♂; Costa Rica, Irazu; BMNH, London.
Distribution: North America (Costa Rica).
References: (**ds**) Blackwelder 1947: 790; Equihua & Atkinson 1987: 25; Schedl 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (**tx**) Blandford 1896e: 99; Brethes 1909; Equihua & Atkinson 1987: 25; Schedl 1940c: 204–205, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.
- obscurus** Schedl 1971c: 393. Holotype ♂; Java, Preanger; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Java), Philippine Islands.
References: (**ds**) Nobuchi 1977: 127; Ohno, Yoneyama, & Nakazawa 1982b: 10; Schedl 1972f: 198. (**tx**) Schedl 1971c: 393–394, 1972f: 198, 1978a: 48.

- obsitus** Schedl 1976a: 90. Holotype ♂; Manaus, Amazonas; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1976a: 90.
- obtectus** (Schedl) 1935n: 641 (*Crossotarsus*). Syn-types ♂; British North Borneo: Mt. Kinabalu, Kamborangah, 7000 feet; BMNH, London, and Schedl Collection in NHMW, Wien. Distribution: Indonesia (Borneo). Notes: (1) Schedl 1978a: 149 (citation of holotype invalid). References: (ds) Schedl 1972f: 200. (tx) Schedl 1935k: 479, 1935n: 641–642, 1936j: 33, 1937e: 544–546, 1972f: 200, 1978a: 49.
- obtusipennis** Schedl 1939c: 2. Holotype ♂; N.E. Burma, Kambaiti, 7000 ft.; NHR, Stockholm. Distribution: Asia (Burma/ Tamil Nadu in India/ Taiwan). Hosts: *Dipterocarpus indicus*. References: (ds) Schedl 1972f: 167, 201, 1975a: 455. (tx) Schedl 1939c: 2, 1962p: 210, 1972f: 201.
- obtusus** Chapuis 1865: 251. Holotype ♂; du Bresil, Bahia; IRSNB, Brussels (Schedl 1960c: 47). Distribution: North America (Costa Rica), South America (Brazil/ Cayenne). References: (ds) Atkinson & Equihua 1986a: 423; Blackwelder 1947: 790; Equihua 1985: 143; Equihua & Atkinson 1987: 25; Gemminger & Harold 1872: 2699; Numberg 1963c: 107; Schedl 1970f: 582, 1972f: 187, 1972g: 49; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 251; Equihua & Atkinson 1987: 25; Murayama 1925: 215–216; Reichardt 1962: 339; Schedl 1933a: 402, 1936c: 248, 1939b: 399, 1939p: 289, 1960c: 47, 1972f: 187; Strohmeier 1912c: 16, 1914c: 27.
- occipitalis** Chapuis 1865: 273. Syntypes ♂ ♀; du Guatemala, Costa Rica; MNB, Berlin (Schedl 1960c: 52). Distribution: North America (Costa Rica/ Guatemala/ Panama). References: (ds) Blackwelder 1947: 790; Blandford 1896c; Equihua & Atkinson 1987: 25; Gemminger & Harold 1872: 2699; Schedl 1966f: 93, 1972f: 184; Strohmeier 1912c: 17, 1914c: 28. (tx) Blandford 1896c; Chapuis 1865: 273; Equihua & Atkinson 1987: 25; Lucas 1920: 522; Schedl 1933c: 175, 1937d: 33, 42, 1939b: 399, 1941e: 156, 1960c: 52, 1972f: 184; Strohmeier 1912c: 17, 1914c: 28; Wood, S. L. 1966a: 54. (ms) Lucas 1920: 522.
- interstitialis** Schedl 1963d: 230. Holotype ♂; Costa Rica: San Jose, 1000–1200 m; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 184. References: (tx) Schedl 1963d: 230, 1972f: 184, 1978a: 35.
- occipitis** Wood 1966a: 54. Holotype ♂; Ikuribisi, British Guiana; BMNH, London. Distribution: South America (Guyana/ Venezuela). Hosts: *Pouteria guianensis*.
- References: (tx) Schedl 1972f: 195; Wood, S. L. 1966a: 54.
- octodentatus** Browne 1955b: 294. Holotype ♂; Keelung (Formosa) to Nagoya (Japan), imported. Distribution: Asia (Taiwan). References: (tx) Browne 1955b: 294.
- octospinosus** Browne 1985a: 193. Kudat (Sabah) to Nagoya (Japan), imported. Figures: Numberg 1960a: 296. Distribution: Indonesia (Borneo). Hosts: *Shorea* sp. References: (tx) Browne 1985a: 193; Numberg 1960a: 296.
- ohnoi** Browne 1981a: 136. Holotype ♂; Singapore to Nagoya (Japan), imported; BMNH, London. Distribution: Asia (Malaya). References: (tx) Browne 1981a: 136.
- olivieri** Chapuis 1865: 132. Holotype ♂; du Bresil, Para, Ega; BMNH, London (Schedl 1960c: 15). Distribution: South America (Brazil). References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Schedl 1972f: 240; Strohmeier 1912c: 9, 1914c: 23. (tx) Chapuis 1865: 132; Schedl 1936c: 230, 1960c: 18, 1972f: 240; Strohmeier 1912c: 9, 1914c: 23.
- omega** Schedl 1970b: 365. Holotype ♂; Malaya, Singapore to Nagoya (Japan), imported; PPST, Tokyo. Figures: Nobuchi 1978a: 80. Distribution: Asia (Malaya). Hosts: *Segetir*. References: (ds) Nobuchi 1977: 134; Schedl 1972f: 210. (tx) Nobuchi 1978a: 80–81; Schedl 1970b: 365–366, 1972f: 210, 1978a: 49.
- omissus** Schedl 1965e: 267. Holotype ♂; Afafiningetu, E. Highlands Dist. [New Guinea]; CSIRO, Canberra. Distribution: New Guinea. Hosts: *Arancaria cunninghamii*. References: (bv) Gray, B. 1974b. (hb) Gray, B. 1974b. (ds) Schedl 1972f: 191. (tx) Schedl 1965e: 267, 1972f: 191, 1978a: 50.
- omnivorus** (Lea) 1904: 104 (*Crossotarsus*). Syn-types ♂ ♀; Tasmania; SAM, Adelaide. Distribution: Australia, Tasmania, New Guinea. Hosts: *Acacia* spp., *Arancaria cunninghamii*, *Eucalyptus* spp., *Flindersia brayleyana*, *F. pubescens*, *Geissois benthamii*. Notes: (1) Schedl 1972f: 206 (to *Platypus*). References: (cn) Browne 1965b: 571; Froggatt 1921: 645–648, 1926b, 1927; Kleine 1932a: 310, 402; Miller 1950: 318; Veitch 1933: 122–127. (cc) Froggatt 1923. (hb) Browne 1965b: 571; Froggatt 1923, 1926b, 1927; Kleine 1932a: 310, 402; Lengerken 1954: 326. (ds) Browne 1965b: 571; Kleine 1932a: 310, 402; Sampson 1925: 2, 1928a: 3; Schedl 1936g: 514, 1962a: 74, 1972f: 206, 1979a: 160, 1980b: 185; Strohmeier 1912c: 19, 1914c: 29. (tx) Beeson 1937: 101; Carne et al.

- 1980; Lea 1904: 104, 1910: 135; Sampson 1925: 2, 1928a: 3, 1928b: 391; Schedl 1939b: 398, 1941c: 153–154, 1942c: 164, 1972f: 206; Strohmeier 1912c: 19, 1914c: 29.
- oneratus** Schedl 1970b: 366. Holotype ♂; Philippine Isl., Dinajawan to Nagoya (Japan), imported; PPST, Tokyo.
Figures: Nobuchi 1978a: 80.
Distribution: Philippine Islands.
Hosts: Lauan log.
References: (ds) Nobuchi 1977: 128; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1972f: 198; (tx) Nobuchi 1978a: 80, 1978b: 80, 1985: 329; Schedl 1970b: 366, 1972f: 198, 1978a: 50.
- opacicauda** Browne 1983a: 565. Holotype ♂; Tg. Usau (West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: Watergum log.
References: (tx) Browne 1983a: 565.
- opacicaudulus** Browne 1983a: 566. Holotype ♂; Tg. Usau (West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: Watergum log.
References: (tx) Browne 1983a: 566.
- opacideclivis** Schedl 1969b: 227. Holotype ♂; New Guinea: Tari Savmill, S. D. H.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Araucaria cunninghamii*, *Macaranga* sp.
References: (ds) Schedl 1972f: 184. (tx) Schedl 1969b: 227–228, 1972f: 184, 1978a: 50.
- opacifrons** Schedl 1936g: 515. Syntypes ♂ ♀; New Guinea and Bougainville; 1 ♀ in SAM, Adelaide, and 1 ♂, 1 ♀ in Schedl Collection in NHMW, Wien.
Distribution: Bougainville Island, New Guinea, Solomon Islands.
Notes: (1) Schedl 1978a: 50 (citation of holotype invalid, male syntype lost).
References: (ds) Schedl 1936g: 515–516, 1972f: 190. (tx) Schedl 1936g: 515, 1942c: 165, 1972f: 190, 1978a: 50.
- opacipennis** Schedl 1975g: 230. Holotype ♂; New Guinea, Wan; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 230, 1978a: 50.
- opacurus** Browne 1980b: 357. Holotype ♂; Jayapura (New Guinea) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
References: (tx) Browne 1980b: 357.
- opimus** Schedl 1975f: 390. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Hosts: *Castanopsis acuminatissima*.
References: (tx) Schedl 1975f: 390.
- orientalis** (Strohmeier) 1912f: 83 (*Steuoplatypus*).
Syntypes ♂ ♀; Ost-Afrika; Strohmeier Collection.
Distribution: Africa (Cameroon/ Congo/ Ethiopia/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Sierra Leone/ Tanzania/ Togo/ Uganda/ Zaïre).
Hosts: Schedl 1962k: 661–667 (many hosts listed).
Notes: (3) Schedl 1972f (etc.) treated this species as *refertus*, although two other names had priority.
References: (hb) Jones, Roberts, & Baker 1959: 13; Strohmeier 1911a: 182; Thompson, C. H. 1963: 59; Webb 1957: 36–41. (ds) Mayne & Donis 1960: 98, 1962: 286; Strohmeier 1911a: 182, 1912c: 6, 1914c: 36; Thompson, C. H. 1963: 59. (tx) Schedl 1954e: 58, 1957d: 118–120, 1962h: 63, 1962k: 663, 1972f: 216; Strohmeier 1911a: 182, 1912c: 6, 1912f: 83, 1914c: 36.
- augustatus** Strohmeier 1912f: 83 (*Crossotarsus*).
Syntypes ♂; Kamerim ud Deutsch Ostafrika am Mkulusumi-Berg; MNB, Berlin, Hamburg Museum, lost, and Strohmeier Collection, preoccupied by Chapuis 1865.
References: (en) Mayne & Donis 1951. (cc) Mayne & Donis 1951. (hb) Gardner 1957: 29; Jones, Roberts, & Baker 1959: 13–54; Webb & Jones 1957: 25–26, 38, 41. (ds) Mayne & Donis 1960: 98, 1962: 280; Strohmeier 1912c: 6. (tx) Schedl 1950d: 4, 11, 14, 1952j: 5, 1953g: 244, 1954e: 48, 1955f: 260, 1957d: 118–122; Strohmeier 1912c: 6, 1912f: 83–85.
- castaneus** Strohmeier 1912f: 84 (*Crossotarsus*).
Syntypes ♂ ♀; West-Usambara (Karasek); Strohmeier Collection. Synonymy: Schedl 1972f: 216.
References: (ds) Strohmeier 1912c: 6, 1914c: 36. (tx) Schedl 1936e: 47–48, 1957d: 118–121, 1972f: 216; Strohmeier 1912c: 6, 1912f: 84–85, 1914c: 36.
- refertus** Schedl 1941d: 415. Holotype ♀; Togo; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1957d: 122.
References: (en) Browne 1968b: 571. (cc) Schedl 1962k: 659. (hb) Browne 1968b: 571; Schedl 1962k: 659. (ds) Browne 1968b: 571; Mayne & Donis 1960: 100, 1962: 286; Roberts 1968: 189; Schedl 1962h: 63, 1962k: 659, 1972e: 284, 1972f: 215. (tx) Numberg 1964: 237; Powell, W. 1987: 28; Schedl 1941d: 415, 1954e: 48, 1957d: 118–122, 1962h: 63, 1962k: 659, 1972f: 215, 1972p: 159, 1978a: 62.
- tetricus** Schedl 1941d: 415. Holotype ♀; Congo; Schedl Collection in NHMW, Wien.
Notes: (1) Schedl 1972f: 216 (treated as a subspecies of *refertus*).
References: (ds) Roberts 1968: 180, 1969: 119; Schedl 1962k: 670. (tx) Schedl 1941d: 415, 1957d: 119, 121, 1962k: 666, 1972f: 216, 1978a: 62.
- pygmaeus** Schedl 1952i: 25. Holotype ♂; Congo Belge; Yaungambi; MRCB, Tervuren.

Notes: (1) Schedl 1972f: 216 (treated as a subspecies of *refertus*).

References: (cn) Thompson, G. H. 1960. (hb) Jones, Roberts, & Baker 1959: 13; Schedl 1962k: 670; Webb & Jones 1957: 37, 41. (ds) Browne 1975b: 396; Mayne & Donis 1960: 100, 1962: 256; Roberts 1968: 189, 1969: 118; Schedl 1962h: 63, 1962k: 670, 1972f: 216; Thompson, G. H. 1960, 1963: 59. (tx) Schedl 1952i: 25–26, 1954e: 58–73, 1957d: 119, 1962h: 63, 1962k: 670, 1972f: 216, 1978a: 60.

refertus montanus Schedl 1957d: 119. Syntypes ♂; Congo Belge: Hembé-Bitale; Schedl Collection in NHMW, Wien.

Notes: This appears to be an aberration with no taxonomic status.

References: (ds) Roberts 1968: 188; Schedl 1962k: 665, 1972f: 216. (tx) Roberts 1969: 118; Schedl 1957d: 119–120, 1962k: 665, 1972f: 216, 1978a: 62.

minutus Roberts 1966: 631. Holotype ♂; Nigeria: Cross River N.F.R.; BMNH, London. Synonymy: Schedl 1972f: 216.

References: (ds) Roberts 1966: 631, 1968: 188, 1969: 118; Schedl 1972f: 216.

refertus vogelii Roberts 1970a: 78. Holotype ♂; Vogel Peak, Nigeria, 1380 m; MRCB, Tervuren. References: (tx) Roberts 1970a: 78.

leonensis Roberts 1970b: 479. Holotype ♂; Sierra Leone: Loma Mountains, 1200 m; BMNH, London. Synonymy: Schedl 1972f: 216.

References: (ds) Browne 1972c: 101. (tx) Roberts 1970b: 479–483; Schedl 1972f: 216, 1972p: 159–160.

refertus lomaensis Roberts 1970b: 483. Holotype ♂; Loma Mountains, 1650 m, Sierra Leone; BMNH, London. Synonymy: Schedl 1972f: 216.

References: (tx) Roberts 1970b: 483; Schedl 1972f: 216.

ornaticeps (Schedl) 1935b: 401 (*Stenoplatypus*). Syntypes ♂ ♀; Mount Maquilang, Laguna, Luzon, Philippine Islands; F. C. Hadden Collection, and Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands.

Notes: (1) Schedl 1978a: 50 (citation of holotype invalid).

References: (ds) Schedl 1966b: 96, 1972f: 217. (tx) Nobuchi 1985: 329; Schedl 1935b: 401–402, 1939c: 4, 1941a: 44, 1964g: 252, 1972f: 217, 1978a: 50.

ornatifrons Schedl 1942: 198. Syntypes ♂; New Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (ds) Schedl 1972f: 203. (tx) Schedl 1942a: 210, 1942c: 198, 1968c: 270, 1972f: 203, 1978a: 50.

ornatus Schedl 1936c: 241. Lectotype ♂; Guyane Française, St.-Jean du Maroni, St. Laurant de Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 50.

Distribution: South America (Cayenne).

References: (ds) Blackwelder 1947: 790; Schedl 1972f: 218. (tx) Schedl 1936c: 241, 1972f: 218, 1978a: 50.

othiodes Schedl 1974d: 469. Holotype ♂; Inakanda logging area, Bulolo, Morobe District; CSIRO, Canberra.

Distribution: New Guinea.

Notes: (3) Schedl 1975f: 391 (described female).

References: (tx) Schedl 1974d: 469, 1975f: 391, 1978a: 50.

otiosus Schedl 1936i: 100. Holotype ♀; Mexico; Schedl Collection in NHMW, Wien.

Distribution: North America (Guatemala/ Durango in Mexico).

Hosts: *Alnus* sp., *Quercus* sp.

References: (hb) Burgos & Sancedo 1983: 44. (ds) Atkinson & Equihua 1985a: 103, 1988: 104; Atkinson et al. 1956: 12–13; Blackwelder 1947: 790; Burgos & Sancedo 1983: 44; Equihua 1985: 143; Equihua & Atkinson 1987: 25; Ferrer 1942; Schedl 1972f: 205; Wood, S. L. 1966a: 53. (tx) Equihua & Atkinson 1987: 25; Schedl 1936i: 100–101, 1940a: 327, 1972f: 205, 1978a: 51; Wood, S. L. 1966a: 53.

ovalicollis Schedl 1936e: 56. Syntypes ♂; Sumatra, Si-Rambe; MCG, Genova, and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

Notes: (1) Schedl 1978a: 51 (citation of holotype invalid).

References: (ds) Schedl 1972f: 196. (tx) Schedl 1936e: 56–57, 1941c: 355, 1972f: 196, 1978a: 51.

ovatus Strohmeier 1914f: 1. Holotype ♂; Birma; Stettin Museum, presumably lost.

Distribution: Asia (Burma/ Malaya), Indonesia (Sarawak in Borneo).

Hosts: *Dipterocarpus* sp., *Eugenia* sp., *Mangifera*, *Swintonia floribunda*, *S. specifera*.

References: (cn) Mathur & Singh 1961a: 81. (hb) Browne 1941, 1961c: 213. (ds) Beaver & Browne 1978: 620; Beeson 1941 (1961: 266); Browne 1961a: 315, 1961c: 213, 1980d: 491; Mathur & Singh 1961a: 81; Schedl 1972f: 191. (tx) Browne 1961a: 315, 1962e: 653; Schedl 1972f: 191; Strohmeier 1914f: 1–4.

oxyurus Dufour 1843: 92. Syntypes²: Des Pyrenees; not located.

Figures: Pfeffer 1989a: pl. 12.

Distribution: Asia (India/ Iran/ Turkey), Europe (Corsica/ France/ Greece/ Italy/ Spain).

Hosts: *Abies cephalonica*, *A. numidica*, *A. pectinata*.

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (cn) Joly 1953: 3–6; Kailidis &

- Georgevits 1972. (**cc**) Kleine 1908c: 222. (**hb**) Eggers 1906; Eichhoff 1881a: 55, 306; Espanol 1964b; Husson 1955: 249; Knotek 1899b: 18, 1901: 576; Pfeffer 1899a: 92; Postner 1974: 486. (**ds**) Acloque 1896; Aube 1950; Barthe 1896; Chapuis 1865; Eggers 1906; Espanol 1985: 115; Gemminger & Harold 1872: 2699; Cogola 1985; Cozis 1815: 80; Herve 1956a; Heyden, Reitter, & Weise 1883: 182, 1891: 674, 1906: 713; Kailidis & Georgevits 1972; Knotek 1899b: 18, 1901: 576; Lacordaire 1866: 391; Lapeyronie 1948: 274; Ortzen 1886: 280; Perris 1876a: 157; Pfeffer 1989a: 92; Postner 1974: 486; Reitter 1894a: 94; Sainte-Claire 1905: 157, 1909: 146, 1914: 475; Sainte-Claire & Mequignon 1938: 449; Schaum 1859: 96, 1862: 101; Schedl 1967c: 73; Siebold 1849: 310; Stein & Weise 1877: 165; Strauch 1861: 122; Strohmeier 1912c: 17, 1914c: 27; Tredl 1907: 20. (**tx**) Acloque 1896; Balachowsky 1949a: 279; Ceballos 1945; Chapuis 1865: 263; Dufour 1843: 263; Eggers 1908c, 1914; Eichhoff 1864b: 43, 1881a: 55, 306, 1883a: 117, 145; Espanol 1964b; Fairmaire 1864: fig. 167; Jacquelin du Val & Fairmaire 1868: 107; Lacordaire 1866: 391; Lucas 1920: 522; Pfeffer 1989a: pl. 12; Portevin 1935: 339; Postner 1974: 486; Reitter 1894a: 94, 1913a: 113; Schedl 1934f: 1647, 1935n: 633, 1960c: 50, 1972f: 198; Strohmeier 1912c: 17, 1914c: 27.
- pahangensis** Schedl 1936j: 34. Lectotype ♂; Malay Peninsula, Pahang; Cameron Highlands, 4800 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 51.
Distribution: Asia (Malaya).
References: (**hb**) Browne 1941, 1961c: 198. (**ds**) Browne 1978b: 51; Nobuchi 1977: 119; Schedl 1936j: 34, 1972f: 193. (**tx**) Schedl 1936j: 34–35, 1972f: 193, 1978a: 51.
- pallidipennis** Blandford 1896e: 111. Holotype ♀; Panama, Bugaba; BMNH, London.
Figures: Blandford 1896e: pl. 4, fig. 22.
Distribution: North America (Panama).
References: (**ds**) Blackwelder 1947: 790; Equihua & Atkinson 1987: 25; Schedl 1972f: 219; Strohmeier 1912c: 14, 1914c: 26. (**tx**) Blandford 1896e: 111; Equihua & Atkinson 1987: 25; Schedl 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.
- pallidus** Chapuis 1865: 284. Syntypes ♀; de la Nouvelle Guinée, Dorey; 2 ♀ in BMNH, London, 1 ♀ in IRSNB, Brussels.
Distribution: Australia, New Guinea, Solomon Islands.
Hosts: *Pterocymbium beccarii*.
Notes: (3) Schedl 1937d: 41 (described male), 1979g: 115 (named an aberration as var. *unimodus*, no status).
References: (**bv**) Gray, B. 1974b. (**cn**) Wylie & Shanahan 1975. (**cc**) Roberts 1980. (**hb**) Gray, B. 1974b; Wylie & Shanahan 1975. (**ds**) Browne 1980c: 483, 1980d: 492; Froggatt 1936; Gemminger & Harold 1872: 2699; Nobuchi 1977: 136; Ohno, Yoneyama, & Nakazawa 1982a: 5, 1982b: 10, 1987a: 95; Ohno et al. 1989: 66; Schedl 1936g: 515, 1962i: 73, 1966b: 96, 1972f: 210, 1979a: 160; Strohmeier 1912c: 18, 1914c: 28. (**tx**) Chapuis 1865: 284; Nobuchi 1985: 329; Schedl 1936e: 56, 1936j: 33–34, 1937d: 41, 1940b: 434, 1941b: 424, 1941e: 155, 1960c: 55, 1972f: 210, 1979g: 115; Strohmeier 1912c: 18, 1914c: 28; Wylie & Yule 1977.
- pallidus sabroni** Schedl 1940b: 433. Holotype ♂; Papua: Kokoda, 1200 ft; Schedl Collection in NHMW, Wien.
References: (**ds**) Schedl 1972f: 210. (**tx**) Schedl 1940b: 433, 1941b: 423, 1946c: 96, 1972f: 210, 1978a: 51.
- panduriformis** Roberts 1989: 278. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, 2200 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Nothofagus* sp.
References: (**tx**) Roberts 1989: 278, 286.
- papuanus** Roberts 1979: 92. Holotype ♂; Papua New Guinea: 12 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Dysoxylum* sp.
References: (**tx**) Roberts 1979: 92.
- papulosus** Roberts 1989: 279. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.
Figures: Roberts 1989: 286.
Distribution: New Guinea.
Hosts: *Myristica* sp.
References: (**tx**) Roberts 1989: 279, 286.
- paradoxus** Schedl 1979g: 115. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (**tx**) Schedl 1979g: 115.
- paralleiventrus** Roberts 1989: 280. Holotype ♂; Papua New Guinea: Divide Logging Area, Bulolo, 1200 m; NHMW, Wien.
Distribution: New Guinea.
Hosts: *Terninalia complanata*.
References: (**tx**) Roberts 1989: 280.
- parallelus** (Fabricius) 1801: 284 (*Bostrichus*). Holotype ♀; America meridionali, type labeled Essequibo [Suriname].
Figures: Atkinson 1989c, Bright 1972d: 19, Equihua & Atkinson 1987: 17, Lara & Schenefelt 1965: 172.
Distribution: Africa (introduced in: Angola/ Cameroon/ Chad/ Congo/ Equatorial Guinea/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Nigeria/ Principe Island/ Sao Tome

Island/ Senegal/ Sierra Leone/ South Africa/ Tanzania/ Togo/ Uganda/ Zaïre), Antilles Islands (Cuba/ Dominican Republic in Hispanola/ Jamaica/ Puerto Rico), Australia (Queensland, introduced), Madagascar (introduced), North America (Belize/ Costa Rica/ El Salvador/ Guatemala/ Honduras/ all states in Mexico/ Nicaragua/ Panama/ S California, S Florida, S Texas in USA), South America (Argentina/ Bolivia/ Brazil/ Cayenne/ Chile/ Colombia/ Ecuador/ Galapagos Islands/ Guyana/ Paraguay/ Peru/ Suriname/ Uruguay/ Venezuela).

Hosts: Schedl 1962k: 624–628, 1979a: 157–164 (many listed).

Notes: (1) Prior to Wood 1973a: 51–52 this name was incorrectly applied to *bellus*. (3) Dejean 1837: 333 (*pracustus*, nomen nudum, *testaceus*, nomen nudum, synonymy in Schedl 1972f: 232). (3) This is the most destructive and the most widely distributed species of Platypodidae in the world (SLW).

References: (cn) Beaver 1988a: 70; Idrobo 1958: 75. (cc) Equihua & Atkinson 1986: 624. (hb) Atkinson 1989c; Beaver 1988a: 70; Burgos & Saucedo 1983: 44; Deyrup & Atkinson 1987a: 64; Equihua & Atkinson 1986: 624. (ds) Allen, A. A. 1976, 1985; Alvarenga 1962: 24; Atkinson 1989c; Atkinson & Equihua 1986a: 423, 1985: 104; Atkinson et al. 1986: 47; Beaver 1988a: 70; Blackwelder 1947: 790; Browne 1975a: 757, 1980d: 491; Burgos & Saucedo 1983: 44; Chapuis 1865: 29, 37, 164; Deyrup & Atkinson 1987a: 64; Drooz 1985: 377; Equihua 1985: 143; Equihua & Atkinson 1986: 624, 1987: 25; Estrada & Atkinson 1988: 212; Gemminger & Harold 1872: 2699; Lacordaire 1866: 391; Leng 1920: 337; Melsheimer 1853: 87; Olmo, Yoneyama, & Nakazawa 1982b: 10; Santoro 1957: 25; Schedl 1933: 193, 1960a: 77, 1965g: 21, 1966f: 81, 1970e: 87, 1972f: 233, 1972g: 50, 1977e: 44, 1978c: 292, 1979a: 160, 1979b: 416, 1979e: 58; Steinhausen 1956: 49; Strohmeier 1912c: 11, 1914c: 25; Swaine 1909: 84; Wood, S. L. 1979a: 2. (tx) Atkinson 1989c; Carne et al. 1980; Chapuis 1865: 29, 37, 164; Equihua & Atkinson 1987: 17, 24; Erichson 1836: 65; Fabricius 1801: 354; Lacordaire 1866: 391; LeConte 1876: 344; Schedl 1933c: 171, 1933e: 193, 1934e: 212, 1936e: 51, 1937d: 39, 1939j: 565, 1939m: 170, 1941e: 155, 1948d: 35, 1952a: 446, 1952h: 73, 1958f: 35, 1960c: 26, 1962k: 624–628, 1972f: 233; Steinhausen 1956: 49; Strohmeier 1912c: 11, 1914c: 25; Swaine 1909: 84; Waterhouse 1890: 554; Wood, S. L. 1973a: 51–52, 1979a: 2; Wylie & Yule 1977.

linearis Stephens 1830: 419. Holotype ♂; Sydenham [England]; BMNH, London. Synonymy: Wood 1979a: 2.

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (cn) Halperin & Menier 1981. (cc) Schedl 1962k: 620. (hb) Cachan

1957: 15; Kalshoven 1963; Lara & Shenefelt 1965: 175; Schedl 1962k: 620. (ds) Anonymous 1965k; Blackwelder 1947: 789; Bright 1972d: 20; Browne 1965a: 205, 1972c: 101; Chapuis 1865: 338; Cola 1971, 1973; Collart 1934: 249; Dejean 1837: 333; Ferreira 1965: 1130; Gemminger & Harold 1872: 2699; Ghesquiere 1966: 41; Halperin & Menier 1981; Kalshoven 1963: 237; Mayne & Donis 1951: 336; Menier 1973a; Mumford 1960: 37, 1961: 31, 43, 1962: 35, 48, 1963: 41, 50, 1964: 30, 1966: 38, 53; Numborg 1971: 59, 1972b: 190; Roberts 1961: 39–43, 1968: 188, 1969: 117–118; Schedl 1941d: 351, 1941e: 155, 1950c: 206, 1950d: 3, 10, 14, 1952j: 5, 1953g: 243, 1954e: 48, 57, 71, 1955d: 270, 1957b: 152, 1957c: 326, 1961e: 157, 1962h: 53, 1962k: 620, 1962m: 64, 1962n: 698, 1963c: 160, 1963f: 64, 1964f: 619, 1964j: 42, 1965e: 357, 1965g: 21, 1966a: 276, 1966c: 230, 1966f: 77, 79, 1966g: 37, 1967e: 220, 1970e: 86, 1970f: 582, 1971e: 2, 1971f: 150, 1971g: 196, 1972e: 284, 1972f: 232, 1972g: 35, 1973a: 368, 1973d: 163, 1974e: 52, 1975h: 351, 1976a: 58, 1976d: 5; Schoneherr & Pedrosa-Macedo 1981: 57; Stephens 1830: 419; Strohmeier 1912e: 19, 1914c: 29; Wolcott 1948: 385; Wood, S. L. 1957e: 1272. (tx) Bright 1972d: 19, 20; Chapuis 1865: 338; Lara & Shenefelt 1965: 175; Sampson 1924: 129; Schedl 1933e: 193, 1936e: 51, 1937b: 407, 1937d: 39, 1941e: 155, 1960c: 64, 1962k: 620, 1972f: 232; Stephens 1830: 419; Strohmeier 1912c: 11, 19, 1914c: 24, 29; Vrydagh 1947: 8; Wood, S. L. 1979a: 2.

pocyi Guerin-Meneville 1838: 183. Syntypes ♂ ♀; de Cuba, de Haiti; 4 ♂, 2 ♀ in BMNH, London, 1 ♂, 1 ♀ IRSNB, Brussels. Synonymy: Schedl 1960c: 64, Wood 1979a: 2.

Notes: (3) Chapuis 1865: 208 (redescribed).

References: (cn) Tsankov et al. 1974: 25. (ds) Blackwelder 1947: 790; Chapuis 1865; Fleutiaux & Salle 1890: 458; Gemminger & Harold 1872: 2699; Lacordaire 1866: 391; Schedl 1933c: 171, 1966f: 80, 1970e: 89; Strohmeier 1912c: 12, 1914c: 25; Wolcott 1936: 316, 1948: 384. (tx) Chapuis 1865: 208; Chevrolat 1838; Guerin-Meneville 1838: 183; Lacordaire 1866: 391; Sagra 1856: 98; Schedl 1933c: 171, 1960c: 37, 1972f: 232; Strohmeier 1912c: 12, 1914c: 25; Wood, S. L. 1979a: 2.

subcostatus Jacquelin-Duval 1857: 238. Synonyms? Cuba; not located. Synonymy: Wood 1979a: 2.

Notes: (3) Chapuis 1865: 210 (redescribed).

References: (ds) Blackwelder 1947: 790; Chapuis 1865: 210; Fleutiaux & Salle 1890: 458; Gemminger & Harold 1872: 2700; Lacordaire 1866: 391; Schedl 1933c: 172; Strohmeier 1912c: 13, 1914c: 25; Wolcott 1936: 317, 1948: 385. (tx) Chapuis 1865: 210; Jacquelin-Duval

- 1857: 238; Lacordaire 1866: 391; Sagra 1856: 99; Schedl 1933c: 172, 1937d: 39, 1941e: 155, 1952a: 444, 1960c: 38, 1972f: 233; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1979a: 2.
- dejeani* Chapuis 1865: 186. Syntypes ♂ ♀; du Bresil, de la Guyanne; 1 ♀ in MNHN, Paris, 1 ♂, 3 ♀ in BMNH, London, and 2 ♂, 3 ♀ in IRSNB, Brussels (Schedl 1960c: 31). Synonymy: Schedl 1972f: 232.
- References: (cn) Costa Lima 1956; Mayne & Donis 1951: 336, (ce) Mayne & Donis 1951: 336, (hb) Costa Lima 1956, (ds) Blackwelder 1947: 789; Blandford 1896e: 107; Costa Lima 1956; Ferrer 1942; Gemminger & Harold 1872: 2698; Hayward 1943: 27; Nunberg 1958a: 482, 1963c: 106; Ruffinelli & Carboneli 1944: 27; Sampson 1928a: 3; Schedl 1933c: 171, 1933f: 192, 1960a: 75-77; Strohmeier 1911f: 204, 1912c: 11, 1914c: 24, (tx) Blandford 1896e: 790; Chapuis 1865: 186; Costa Lima 1956; Nunberg 1958a: 482; Ruffinelli Rey 1967; Sampson 1924c: 129, 1928a: 3; Schedl 1933f: 192-193, 1936e: 51, 1937d: 39, 1939m: 170, 1940a: 324, 1948d: 38, 1950i: 148, 1951m: 72, 74, 1952a: 444, 1952h: 73, 1960c: 31, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 1.
- marseuli* Chapuis 1865: 188. Syntypes ♂ ♀; Bresil, Ste. Catherine; 1 ♀ in BMNH, London, 1 ♂ in IRSNB, Brussels (Schedl 1960c: 32). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 11, 1914c: 25, (tx) Chapuis 1865: 188; Nunberg 1939: 240-241, 1940: 68; Schedl 1941e: 156, 1960c: 32, 1972f: 232; Strohmeier 1912c: 11, 1914c: 25; Wood, S. L. 1979a: 1.
- proximus* Chapuis 1865: 188. Syntypes ♂; du Bresil; 2 ♂ in IRSNB, Brussels (Schedl 1960c: 32). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947; Gemminger & Harold 1872: 2700; Strohmeier 1912c: 12, 1914c: 25, (tx) Chapuis 1865: 188; Schedl 1941e: 156, 1960c: 32, 1972f: 232; Strohmeier 1912c: 12, 1914c: 25; Wood, S. L. 1979a: 2.
- compressus* Chapuis 1865: 191. Syntypes ♂ ♀; Caracas; 2 ♂, 3 ♀ in BMNH, London (Schedl 1960c: 32). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Strohmeier 1912c: 11, 1914c: 24, (tx) Chapuis 1865: 191; Schedl 1936c: 238, 1937d: 39, 1941e: 156, 1952h: 73, 1960c: 32, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 1-2.
- regularis* Chapuis 1865: 192. Holotype ♂; du Bresil; IRSNB, Brussels (Schedl 1960c: 33). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Strohmeier 1912c: 12, 1914c: 25, (tx) Schedl 1936e: 51, 1937d: 39, 1939j: 565, 1940a: 324, 1940c: 203, 1941e: 156, 1948d: 38, 1951h: 287, 1952a: 446, 1952e: 123, 1955b: 274, 1957a: 195, 1958c: 2, 1958f: 35-36, 1972f: 233; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 86; Wood, S. L. 1973a: 51, 1979a: 2.
- reticulatus* Chapuis 1865: 194. Syntypes ♂ ♀; du Mexique; Mexico, Teapa; 1 ♂, 1 ♀ in BMNH, London, 1 ♂, 2 ♀ in IRSNB, Brussels (Schedl 1960c: 33). Synonymy: Wood 1979a: 2.
- References: (ds) Gemminger & Harold 1872: 2700; Leng 1920: 337; Swaine 1909: 86, (tx) Blandford 1896e: 107; Chapuis 1865: 194; Schedl 1960c: 33, 1972f: 233; Swaine 1909: 86; Wood, S. L. 1979a: 2.
- rotundatus* Chapuis 1865: 195. Syntypes ♂ ♀; du Mexique, Izabal; 3 ♂, 2 ♀ in BMNH, London, 2 ♀ in IRSNB, Brussels (Schedl 1960c: 34). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 790; Blandford 1896e: 108; Gemminger & Harold 1872: 2700; Strohmeier 1912c: 12, 1914c: 25, (tx) Blandford 1896e: 108; Chapuis 1865: 195; Schedl 1960c: 34, 1972f: 233; Strohmeier 1912c: 12, 1914c: 25; Wood, S. L. 1979a: 2.
- kraatzii* Chapuis 1865: 196. Syntypes ♂; de la Colombie et du Bresil; 2 ♂ in IRSNB, Brussels (Schedl 1960c: 34). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Strohmeier 1912c: 11, 1914c: 24, (tx) Chapuis 1865: 196; Schedl 1960c: 34, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.
- lebasii* Chapuis 1865: 197. Holotype ♀; de Carthagene [Colombia]; IRSNB, Brussels (Schedl 1960c: 34). Synonymy: Wood 1979a: 2.
- References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 11, 1914c: 24, (tx) Chapuis 1865: 197; Schedl 1960c: 34, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.
- emarginatus* Chapuis 1865: 199. Holotype ♂;

- de Toxpan; BMNH, London (Schedl 1960c: 35). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2698; Leng 1920: 337; Strohmeier 1912c: 11, 1914c: 24. (tx) Blandford 1896c: 107; Chapuis 1865: 199; Schedl 1960c: 35 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Swaine 1909: 86; Wood, S. L. 1979a: 1–2.
- punctulatus* Chapuis 1865: 199. Syntypes ♂; du Texas; 1 ♂ in BMNH, London, 4 ♂ in IRSNB, Brussels (Schedl 1960c: 35). Synonymy: Wood 1979a: 2.
References: (cn) Felt 1913: 275. (ds) Anonymous 1926c: 514; Blackwelder 1947: 790; Ferrer 1942; Gemminger & Harold 1872: 2700; Henshaw 1885: 147; Leng 1920: 337; Leonard 1928: 514; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 85; Tucker 1952: 347; Wolcott 1948: 385. (tx) Chapuis 1865: 199; LeConte 1865: 151, 1876: 345; Schedl 1937d: 39, 1939j: 565, 1940a: 324–325, 1940c: 203, 1941e: 156, 1952e: 123, 1952h: 73, 1955d: 273, 1960c: 35, 1972f: 232; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 85; Wood, S. L. 1979a: 2.
- subaequalis* Chapuis 1865: 200. Holotype ♂; de la Colombie; IRSNB, Brussels (Schedl 1960c: 35). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947; Gemminger & Harold 1872: 2700; Strohmeier 1912c: 13, 1914c: 25. (tx) Chapuis 1865: 200; Schedl 1941e: 156, 1960c: 35, 1972f: 233; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1979a: 2.
- wesmali* Chapuis 1865: 201. Syntypes ♂ ♀; de Buenos-Ayres; 1 ♂, 1 ♀ in IRSNB, Brussel (Schedl 1960c: 35). Synonymy: Wood 1979a: 2.
References: (cn) Costa Lima 1956; Silva, Gayao, & Castro 1959: 12. (hb) Bosq 1934: 335; Costa Lima 1956. (ds) Blackwelder 1947: 791; Bruch 1914a; Costa Lima 1956; Gemminger & Harold 1872: 2700; Sampson 1928a: 3; Schedl 1933c: 172; Strohmeier 1911f: 204, 1912c: 13, 1914c: 25. (tx) Brethes 1909, 1919; Chapuis 1865: 201; Costa Lima 1956; Nummer 1939: 241, 1940: 69; Sampson 1928a: 3; Schedl 1937d: 39, 1938i: 28, 1941e: 156, 1948d: 37, 1951h: 285–286, 1952a: 445, 1952h: 73, 1958f: 36, 1960c: 35, 1972f: 233; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1979a: 2.
- oblongus* Chapuis 1865: 203. Holotype ♀; de Cumana [Venezuela]; IRSNB, Brussels (Schedl 1960c: 35). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 11, 1914c: 25. (tx) Chapuis 1865: 203; Schedl 1960c: 35, 1972f: 232; Strohmeier 1912c: 11, 1914c: 25; Wood, S. L. 1979a: 2.
- difficilis* Chapuis 1865: 204. Bresil, a la Guyane; 2 ♂, 1 ♀ in MNB, Berlin, 2 ♂, 1 ♀ in BMNH, London, 4 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 36). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947: 789; Ferrer 1942; Gemminger & Harold 1872: 2698; Sampson 1928a: 3; Schedl 1933c: 171; Strohmeier 1912c: 11, 1914c: 24; Vital 1951: 177–179. (tx) Chapuis 1865: 204; Sampson 1928a: 3; Schedl 1936c: 238, 1937d: 39, 1940a: 325–326, 1952d: 343, 1952h: 73, 1960c: 36, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.
- praevis* Chapuis 1865: 205. Syntypes ♂ ♀; de la Colombie; 1 ♀ in BMNH, London, 1 ♂, 2 ♀ in IRSNB, Brussels (Schedl 1960c: 36). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Strohmeier 1911f: 204, 1912c: 12, 1914c: 25. (tx) Chapuis 1865: 205; Schedl 1960c: 36, 1972f: 232; Strohmeier 1912c: 12, 1914c: 25; Wood, S. L. 1979a: 2.
- macklini* Chapuis 1865: 207. Syntypes ♂ ♀; de la Colombie et du Bresil; 1 ♀ MNHN, Paris, 4 ♂, 2 ♀ in BMNH, London, 3 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 36–37). Synonymy: Wood 1979a: 2.
References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2699; Schedl 1933c: 171; Strohmeier 1912c: 11, 1914c: 25. (tx) Chapuis 1865: 207; Schedl 1960c: 36–37, 1972f: 232; Strohmeier 1912c: 11, 1914c: 25; Wood, S. L. 1979a: 1.
- erichsoni* Chapuis 1865: 211. Syntypes ♂ ♀; de l'île St. Thomas; 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 38). Synonymy: Wood 1979a: 2.
References: (cn) Lapexronie 1948: 275; Mayne & Donis 1951: 326, 336; Vrydagh 1947a: 8. (cc) Mayne & Donis 1951: 326, 336. (ds) Blackwelder 1947: 789; Collard 1934; Gemminger & Harold 1872: 2698; Hoffmann 1957; Mayne & Donis 1962: 281; Schedl 1933f: 192; Strohmeier 1912c: 11, 1914c: 24; Thompson, G. H. 1963: 58; Vrydagh 1947a: 8; Wood, S. L. 1957c: 1272. (tx) Chapuis 1865: 211; Sampson 1924c: 127; Schedl 1933f: 192–193, 1936e: 51, 1937b: 407, 1937d: 39, 1941d: 351, 1941e: 155, 1950c: 206, 1950d: 3, 10, 14, 1952j: 5, 1953g: 243, 1954e: 48, 57, 1955d: 270, 1957b: 152, 1960c: 12, 35, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.
- laevicollis* Chapuis 1865: 212. Syntypes ♂ ♀; de la Guadeloupe; 1 ♂, 1 ♀ in BMNH, London, 1 ♂ in IRSNB, Brussels (Schedl 1960c: 38). Synonymy: Schedl 1972f: 232, Wood 1979a: 2.
References: (ds) Blackwelder 1947: 789; Fleutiaux & Salle 1890: 458; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 11, 1914c: 24. (tx) Chapuis 1865: 212; Schedl 1941e: 156, 1960c: 39, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.

- congoanus* Duvivier 1891: 363. Syntypes ♂ ♀; Ibenbo, Congo; not located. Synonymy: Wood 1979a: 2.
References: (ds) Schedl 1933f: 192; Strohmeier 1912c: 11, 1914c: 24. (tx) Duvivier 1891: 363; Sampson 1924: 129; Schedl 1933f: 192–193, 1960c: 39, 1972f: 232; Strohmeier 1912c: 11, 1914c: 24; Wood, S. L. 1979a: 2.
- triquetrus* Brethes 1909: 226. Syntypes ♂ ♀; Chaco [Argentina]; Brethes Collection. Synonymy: Wood 1979a: 2.
References: (en) Bosq 1934: 335; Hayward 1942: 27. (ds) Bruch 1914a; Strohmeier 1912c: 13, 1914c: 25. (tx) Brethes 1909: 226; Schedl 1939m: 170, 1972f: 233; Strohmeier 1912c: 13, 1914c: 25; Wood, S. L. 1979a: 2.
- mattai* Brethes 1921: 26. Holotype ♂; Manaos [Brazil]; Brethes Collection. Synonymy: Wood 1979a: 2.
References: (en) Costa Lima 1956. (hb) Costa Lima 1956; Matta 1922: 158, 1929: 495–498. (ds) Blackwelder 1947: 790; Costa Lima 1936, 1956. (tx) Brethes 1921: 26; Costa Lima 1956; Schedl 1972f: 232; Wood, S. L. 1979a: 2.
- paramonovi* Schedl 1972g: 80. Holotype ♂; Guiana; Schedl Collection in NHMW, Wien.
Distribution: South America (Suriname?).
References: (tx) Schedl 1972g: 80, 1978a: 51.
- parapetax* Roberts & Morimoto 1986: 174. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Dysoxylum* sp.
References: (tx) Roberts & Morimoto 1986: 174.
- partibilis* Roberts & Morimoto 1986: 174. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Syzigium* sp.
References: (tx) Roberts & Morimoto 1986: 174.
- particeps* Schedl 1976a: 91. Holotype ♂; Villhena, Rondonia, Lon. 55 36, Lat. 12 46; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (tx) Schedl 1976a: 91.
- partitus* Schedl 1935n: 639. Syntypes ♂; Malay States: Selangor, Ayer Itam; BMNH, London, and Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya), Indonesia (Borneo).
Hosts: *Endospermum malaccense*, Pernarahan, and Sendoksendok logs.
Notes: (1) Schedl 1978a: 51 (type restricted to BMNH, London specimens).
References: (hb) Browne 1941. (ds) Browne 1961c: 198, 1984c: 448; Nobuchi 1977: 118; Schedl 1972f: 193. (tx) Schedl 1935n: 639, 1936c: 52, 1972f: 193, 1978a: 51.
- parvus* Browne 1986b: 340. Holotype ♂; Solomon Islands: Virm Harbor to Nagoya (Japan), imported; BMNH, London.
Distribution: Solomon Islands.
Hosts: *Terminalia* sp.
References: (tx) Browne 1986b: 340.
- pasanaie* Schedl 1958b: 105. Lectotype ♂; Malaya, Selangor; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 51.
Distribution: Asia (Malaya).
Hosts: *Pasania sundaica*.
References: (ds) Browne 1961c: 216; Schedl 1972f: 217. (tx) Schedl 1958b: 105, 1972f: 217, 1978b: 217.
- patens* Roberts 1986: 46. Holotype ♂; New Guinea: Papua, Popondetta; BMNH, London.
Distribution: New Guinea.
Hosts: *Myristica* sp.
References: (tx) Roberts 1986: 46.
- patulus* Chapuis 1865: 190. Holotype ♂; du Bresil; MNHN, Paris (Schedl 1960c: 32).
Distribution: South America (Brazil).
References: (ds) Gemminger & Harold 1872: 2699; Schedl 1972f: 233. (tx) Chapuis 1865: 190; Schedl 1960c: 32, 1972f: 233.
- pectinatus* Roberts 1986: 47. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (tx) Roberts 1986: 47.
- pedellus* Schedl 1975g: 230. Holotype ♂; New Guinea, Wau, 1200 m; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975g: 230, 1978a: 52.
- pedum* Sampson 1928a: 2. Holotype ♂; Sumatra, Sibajak vulkan 1600 m; BMNH, London.
Distribution: Indonesia (Sumatra).
References: (ds) Sampson 1928a: 2; Schedl 1972f: 194. (tx) Sampson 1928a: 2; Schedl 1935g: 320, 1972f: 194, 1978a: 52.
- pennatus* Schedl 1962k: 643. Lectotype ♂; Kenya: S. Kinangop; Schedl Collection in NHMW, Wien.
Distribution: Africa (Kenya/ Nigeria/ Zaire).
Hosts: *Cassipourea* sp., *Erythrina* sp.
References: (ds) Roberts 1968: 188, 1969: 118; Schedl 1972f: 215. (tx) Schedl 1957d: 120, 1962k: 643, 1972f: 215, 1978a: 52.
- perbinodulus* Schedl 1934e: 211. Holotype ♂; Rio de Janeiro, Brazil; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 241. (tx) Schedl 1934e: 211, 1937d: 37, 1972f: 241, 1978a: 52.
- perbrevis* Browne 1986b: 336. Holotype ♂; Borneo: Tanjong Mani (Sarawak) to Nagoya (Japan), imported; BMNH, London.

- Distribution: Indonesia (Sarawak in Borneo).
Hosts: *Rengas* sp. (Anacardiaceae) log.
References: (tx) Browne 1986b: 336.
- percomis** Schedl 1936c: 239. Syntypes ♂; Venezuela, Moritz [presumably Colonia Tovar]; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 218. (tx) Schedl 1936c: 239, 1972f: 218, 1978a: 53.
- perdiligens** Schedl 1936c: 240. Lectotype ♂; Guyane Française, Nouveau Chantier and St.-Laurent du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53.
Distribution: South America (Brazil/ Cayenne).
References: (ds) Blackwelder 1947: 790; Schedl 1966f: 93, 1970e: 87, 1972f: 218, 1972g: 50. (tx) Schedl 1936c: 240, 1972f: 218, 1978a: 53.
- perditus** Schedl 1975f: 391. Holotype ♀; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 391.
- perforans** Schedl 1961i: 232. Holotype ♀; Bolivia, Corvico, Huarinillas; Schedl Collection in NHMW, Wien.
Distribution: South America (Bolivia).
References: (ds) Schedl 1960a: 77, 1972f: 219. (tx) Schedl 1961i: 232, 1972f: 219, 1978a: 53.
- permarginatus** Schedl 1936c: 230. Lectotype ♂; Guyane Française: Nouveau Chantier, St. Jean du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53.
Distribution: South America (Cayenne).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 241. (tx) Schedl 1936c: 225, 230–235, 1972f: 241, 1978a: 53.
- permimicus** Schedl 1936c: 237. Lectotype ♂; Guyane Française: St. Laurent du Maroni and Bas Maroni; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
References: (hb) Kalshoven 1960c: 40, 1960d: 40. (ds) Blackwelder 1947: 790; Kalshoven 1960d: 40; Schedl 1961c: 70, 1972f: 233. (tx) Schedl 1936c: 236–237, 1972f: 233, 1978a: 53.
- permodestus** Schedl 1936c: 235. Lectotype ♀; French Guyana, St. Laurent du Maroni; Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne).
References: (ds) Blackwelder 1947; Schedl 1972f: 241. (tx) Schedl 1936c: 235, 1972f: 241, 1978a: 54.
- pernudus** Schedl 1936c: 243. Syntypes ♂ ♀; Guyana[?]; Schedl Collection in NHMW, Wien.
Distribution: South America (Guyana[?]).
Notes: (1) Schedl 1978a: 54 (citation of holotype invalid).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 214. (tx) Schedl 1936c: 243, 1972f: 214, 1978a: 54.
- perplexus** Bright 1972d: 20. Holotype ♂; Good Hope, Trelawny Parish, Jamaica; CNCI, Ottawa.
Figures: Bright 1972d: 19.
Distribution: Antilles Islands (Jamaica).
References: (tx) Bright 1972d: 19–20; McNamara 1977: 191.
- perrisi** Chapuis 1865: 260. Syntypes ♀; de la presqu'île de Malacca (Singapour) at de l'île de Borneo (Sarawak); 3 ♀ in BMNH, London, 1 ♀ in IRSNB, Brussels.
Distribution: Asia (Malaya/ Singapore), Indonesia (Sarawak in Borneo), Philippine Islands.
Hosts: *Shorea robusta*.
References: (cn) Mathur & Singh 1961a: 45. (hb) Browne 1941; Mercer 1982: 210. (ds) Beaver & Browne 1978: 620; Browne 1961c: 200; Gemminger & Harold 1872: 2699; Mathur & Singh 1961a: 45; Mercer 1982: 210; Sampson 1919: 106, 1928b: 394; Schedl 1962h: 63, 1966b: 97, 1972f: 185; Strohmeier 1912c: 17, 1914c: 27. (tx) Chapuis 1865: 260; Nobuchi 1985: 328; Sampson 1919: 106, 1928b: 394; Schedl 1939b: 398, 1960c: 49, 1972f: 185; Strohmeier 1912c: 17, 1914c: 27.
- pertusus** Chapuis 1865: 170. Holotype ♂; de Caracas; BMNH, London (Schedl 1960c: 27).
Distribution: South America (Brazil/ Venezuela).
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Schedl 1972f: 233, 1976a: 58; Strohmeier 1912c: 11, 1914c: 25. (tx) Chapuis 1865: 170; Keler 1927a: 235; Schedl 1939j: 565, 1948d: 37, 1960c: 27, 1972f: 233, 1978a: 54; Strohmeier 1912c: 11, 1914c: 25.
- peruanus** Nunberg 1939: 238. Holotype ♀; Lima, Peru; Polischen Zoologischen Staatmuseums.
Distribution: South America (Peru).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 241. (tx) Nunberg 1939: 238–241, 1940: 63; Schedl 1952h: 72, 1972f: 241.
- petalinus** Roberts & Morimoto 1986: 175. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Nothofagus* sp.
References: (tx) Roberts & Morimoto 1986: 175.
- petaloideus** Roberts 1979: 93. Holotype ♂; Papua New Guinea: 8 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Nothofagus pullei*.
References: (tx) Roberts 1979: 93.
- petax** Schedl 1974d: 469. Holotype ♂; Latep logging area, Bulolo, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Garuga floribunda*.
Notes: (3) Schedl 1979c: 116 (described female).

References: (tx) Schedl 1974d: 469, 1978a: 54, 1979c: 116.

picinus Schedl 1952i: 22. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 636, 1972f: 222.

Distribution: Africa (Cameroon/ Nigeria/ Zaire).

Hosts: *Anonidium mannii*, *Macaranga* cf. *lanceifolia*, *Polyalthia suaveolens*, *Pycnanthus angolensis*, *Ricinodendron hendelotii*, *Scorodophloeus zenkeri*, *Terminalia superba*.

References: (ds) Mayne & Donis 1960: 98, 1962: 286; Roberts 1968: 188, 1969: 118; Schedl 1962k: 636, 1972f: 223. (tx) Schedl 1952i: 22, 1957d: 120, 1962k: 636, 1108, 1972f: 222–223, 1978a: 55.

piroloti Browne 1972c: 106. Holotype ♂; Congo-Kinshasa: Kivu, Bitale; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Browne 1972c: 106.

pilidens Schedl 1955b: 309. Holotype ♂; New-Guinea, Berlinhafen; originally in Schedl Collection but transferred to NHMB, Budapest (Schedl 1978a: 55).

Distribution: New Guinea.

References: (ds) Schedl 1972f: 200. (tx) Schedl 1955b: 282, 309, 1972f: 200, 1978a: 55.

pini Hopkins 1905b: 71. Holotype ♂; Mexico: Chalco; USNM, Washington.

Figures: Reichardt 1965b: 161.

Distribution: North America (Guatemala/ Honduras/ Durango, Hidalgo, Mexico, Puebla, Tlaxcala in Mexico/ Arizona, New Mexico in USA). Hosts: *Pinus chihuahuana*, *P. leiophylla*, *P. ochoterenai*, *P. ponderosa*, *P. rudis*, *P. pseudostrobus*.

Notes: (3) Reichardt 1965b: 159 (described female).

References: (hb) Atkinson et al. 1986: 12; Burgos & Saucedo 1984: 45. (ds) Atkinson & Equihua 1985a: 103, 1988: 104; Atkinson et al. 1986: 12; Blackwelder 1947: 790; Burgos & Saucedo 1983: 45; Equihua 1985: 144; Equihua & Atkinson 1987: 26; Ferrer 1942; Furniss, R. L. & Carolin 1977: 339; Schedl 1972f: 223, 1977c: 44; Snow 1881: 70, 1907: 188; Strohmeier 1912c: 11, 1914c: 25; Thomas, J. B. 1966; Wickham 1898: 312; Wood, S. L. 1958b: 37, 1979a: 2. (tx) Equihua & Atkinson 1987: 26; Hopkins 1905b: 71; Reichardt 1965b: 159, 161; Schedl 1940a: 324–325, 1972f: 223; Strohmeier 1912c: 11, 1914c: 25; Wood, S. L. 1958b: 37, 1966b: 17, 1979a: 2.

quadridens Schedl 1937d: 38. Lectotype ♂; Mexico, Sierra de Durango und Camelas, Staat Durango; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 60. Synonymy: Wood 1966b: 17.

References: (cn) Schwerdtfeger 1960b: 254–259. (ec) Becker 1953: 363, 1955; Schwerdtfeger 1960b: 254–259. (hb) Becker 1953: 363, 1955. (ds) Becker 1953: 363; Blackwelder 1947: 790; Ferrer 1942; Kirk 1969, 1970;

Schwerdtfeger 1960b: 254–259. (tx) Schedl 1937d: 38, 1940a: 324, 1955g: 48, 1972f: 223, 1978a: 60; Schwerdtfeger 1960b: 254–259; Wood, S. L. 1966b: 17–18, 1979a: 2.

piniperda Schedl 1969c: 65. Holotype ♂; Burma: Kalaw, Shan States; FRI, Dehra Dun, not found in 1981 by SLW.

Distribution: Asia (Burma/ Cambodia/ Thailand).

Hosts: *Pinus insularis*.

Notes: (1) None of the specimens borrowed by Schedl from FRI were ever returned to FRI; the type apparently is represented by “paratypes” in the Schedl Collection. (3) Beeson 1941: 345, 1961: 266 (*piniperda*, nomen nudum, synonymy in Schedl 1972f: 211).

References: (ds) Beaver 1990a: 280; Beeson 1941: 345, 1961: 266; Browne 1980a: 370; Schedl 1972f: 211. (tx) Schedl 1964c: 303, 1969a: 216, 1969c: 65–66, 1972f: 211, 1978a: 55.

piniperda cambodianus Schedl 1969a: 216. Holotype ♂; Cambodia, Phnompenh to Tokyo (Japan), imported; PPST, Tokyo.

References: (ds) Nobuchi 1977: 138; Schedl 1972f: 211. (tx) Schedl 1969a: 216–217, 1972f: 211, 1978a: 55.

planodeclivis Schedl 1942a: 206. Lectotype ♂; Malaya, Pahang, Raub (Chengal); Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 55.

Figures: Nobuchi 1978a: 80.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Balanocarpus heimii*.

References: (ds) Browne 1961c: 229; Nobuchi 1977: 126; Schedl 1972f: 196. (tx) Browne 1962c: 643; Nobuchi 1978a: 80, 1978b: 79; Schedl 1942a: 206, 1972f: 196, 1978a: 55.

platypoides (Browne) 1955: 368 (*Crossotarsus*). Holotype ♂; Malay Peninsula: Kelantan, Sokor; BMNH, London.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Calophyllum* sp., *Castanopsis* spp., *Dipterocarpus borneensis*, *Dryobalanops oblongifolia*, *Pasania sundaica*, *Quercus* sp., *Swietenia macrophylla*.

Notes: (1) Schedl 1972f: 212 (to *Platypus*).

References: (hb) Browne 1961c: 229. (ds) Schedl 1964c: 305, 1972f: 212. (tx) Browne 1955: 368–369, 1961a: 316, 1961c: 229; Schedl 1972f: 212.

plumatus Schedl 1979g: 116. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1979g: 116.

podocarpus Browne 1986c: 669. Holotype ♂; Vietnam: Qui Nhon to Tagonoura (Japan), imported; BMNH, London.

Distribution: Asia (Vietnam).

References: (tx) Browne 1986c: 669.

- politus Chapuis** 1865: 249. Holotype ♀; de l'île de Borneo, Sarawak; BMNH, London (Schedl 1960c: 47).
Distribution: Indonesia (Sarawak in Borneo).
References: (ds) Gemminger & Harold 1872: 2699; Sampson 1928b: 392; Schedl 1972f: 194; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 249; Sampson 1928b: 392; Schedl 1960c: 47, 1972f: 194; Strohmeier 1912c: 16, 1914c: 27.
- porcellinus Schedl** 1975f: 392. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 392.
- porcellus Schedl** 1942a: 209. Holotype ♂; Salomons Inseln; Schedl Collection in NHMW, Wien.
Figures: Nobuchi 1978a: 80.
Distribution: New Guinea, Solomon Islands.
Hosts: *Calophyllum* sp.
Notes: (3) Schedl 1972h: 70 (described female).
References: (ds) Nobuchi 1977: 120; Ohno, Yoshioka, et al. 1988b: 98, 1989: 66; Schedl 1972f: 193. (tx) Nobuchi 1978b: 77, 80; Schedl 1942a: 209, 1972f: 193, 1972b: 70, 1978a: 55.
- poricollis Schedl** 1950i: 147. Holotype ♀; New Guinea; Schedl Collection in NHMW, Wien, automatic.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 199. (tx) Schedl 1950i: 147, 1972f: 199.
- porosus Schedl** 1942c: 193. Holotype ♀; New Guinea; Schedl Collection in NHMW, Wien, preoccupied by Blandford 1896.
References: (ds) Equihua & Atkinson 1987: 27; Schedl 1972f: 205. (tx) Equihua & Atkinson 1987: 27; Schedl 1942c: 193, 1950i: 147, 1972f: 205, 1978b: 56.
- porosulus Browne** 1983a: 564. Holotype ♂; Teminaban (West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
References: (tx) Browne 1983a: 564.
- porosus Blandford** 1896e: 105. Syntypes ♂ ♀; Panama, Volcan de Chiriqui; BMNH, London.
Figures: Blandford 1896e: pl. 4, fig. 16.
Distribution: North America (Panama).
References: (ds) Blackwelder 1947: 790; Strohmeier 1912c: 10, 1914c: 24. (tx) Blandford 1896e: 105–106; Strohmeier 1912c: 10, 1914c: 24.
- porrectus Chapuis** 1865: 150. Holotype ♀; de la Colombie; IRSNB, Brussels (Schedl 1960c: 22).
Distribution: North America (Costa Rica/Panama), South America (Colombia).
References: (ds) Blackwelder 1947: 790; Blandford 1896e: 103–104; Equihua & Atkinson 1987: 27; Gemminger & Harold 1872: 2699; Schedl 1933c: 166, 1972f: 241; Strohmeier 1912c: 9, 1914c: 23. (tx) Blandford 1896e: 103–104; Chapuis 1865: 150; Equihua & Atkinson 1987: 27; Schedl 1960c: 22, 1972f: 241; Strohmeier 1912c: 9, 1914c: 23.
- pouteriae Wood** 1972b: 252. Holotype ♂; 40 km SE Socopo, Barinas, Venezuela; Wood Collection.
Distribution: South America (Colombia/Venezuela).
Hosts: *Pouteria amibacfolia*, *P.* sp.
References: (tx) Wood, S. L. 1972b: 252.
- praealtus Schedl** 1975f: 393. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 393.
- praeceless Schedl** 1969b: 228. Holotype ♂; New Guinea; Porotop L.M. Station, W. H. D.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Nothofagus* sp.
References: (ds) Schedl 1972f: 200. (tx) Schedl 1969b: 228, 1972f: 200, 1978a: 56.
- praepositus Schedl** 1969b: 229. Holotype ♂; New Guinea; Watut Valley, 1200 m, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Araucaria hunsteinii*, *Xanthophyllum* sp.
Notes: (3) Schedl 1972i: 58 (described female).
References: (ds) Schedl 1972f: 200. (tx) Schedl 1969b: 229–230, 1972f: 200, 1972i: 58, 1978a: 56.
- praeteritus Schedl** 1969b: 230. Holotype ♂; New Guinea; Porotop L. M. Station, W. H. D.; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Syzygium* sp.
References: (ds) Schedl 1972f: 193. (tx) Schedl 1969b: 230, 1972f: 193, 1978a: 56.
- praetermissus Schedl** 1979g: 117. Holotype ♂; Papua, Bulolo, Morobe District. Upper Manki L.A.; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1979g: 117.
- prenexus Wood** 1966a: 64. Holotype ♂; 16 km SE Cartago on Pan-American Highway, Cartago Prov., Costa Rica, 5600 feet; Wood Collection.
Figures: Schedl 1972f: 220, Wood 1966a: 67.
Distribution: North America (Costa Rica).
References: (ds) Equihua & Atkinson 1987: 27; Schedl 1972f: 220. (tx) Equihua & Atkinson 1987: 27; Schedl 1972f: 220; Wood, S. L. 1966a: 64, 67.
- principeensis (Schedl)** 1936e: 60 (*Stenoplatypus*).
Lectotype ♂; Is. Principe, Roca Inf. D. Henrique, 100–300 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 56.
Distribution: Africa (Principe Island).
Notes: (3) Schedl 1972f: 215 (treated as a subspecies of *intermedius*), Roberts 1972: 71 (treated as a good species).
References: (ds) Schedl 1972f: 215. (tx) Roberts

1972: 71; Schedl 1936e: 60–61. 1957d: 120, 1962k: 653, 1972f: 215, 1978a: 56.

propinquus Schedl 1959m: 555. Holotype ♂; Brasilien, Nova Tentonia; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (**ds**) Schedl 1967d: 5, 1972f: 219. (**tx**) Schedl 1959m: 555, 1972f: 219, 1978a: 57.

protenus (Schedl) 1939e: 356 (*Crossotarsus*). Lectotype ♂; Malaya, Pahang; Kemasn For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 57.

Distribution: Asia (India, Malaya).

Hosts: *Hopea mengarawan*, *Pometia macrocarpa*, *Xerospermum* sp.

References: (**ds**) Beeson 1961: 266; Browne 1961c: 200; Schedl 1972f: 207. (**tx**) Schedl 1939e: 356–357, 1942a: 172, 1972f: 207, 1978a: 57.

pseudocupulatus Schedl 1935n: 635. Syntypes ♂ ♀; Malay States: Selangor, Rawang, Sungei Buloh, and Kakang; BMNH, London, and Schedl Collection in NHMW, Wien.

Figures: Nakashima 1975: 59, Nobuchi 1978a: 80. Distribution: Asia (Bangladesh/ Burma/ China/ Assam in India/ Malaya/ Thailand/ Tonkin in Vietnam), Australia, Indonesia (Aru Island, Batchian, Borneo, Celebes, Java, Moluccas, Sumatra), New Guinea, Philippine Islands, Sumba Island.

Hosts: *Agave sisalana*, *Alcurites moluccana*, *Alstonia* sp., *Aranga* sp., *Arthrophyllum diversifolium*, *Artocarpus elasticus*, *A. lakoocha*, *Austroboxus nitidus*, *Bouea* sp., *Cassia siamea*, *Castanopsis* sp., *Chrysophyllum roxburghii*, *Dalbergia latifolia*, *Dryobalanops aromatica*, *D. oblongifolia*, *Durio* sp., *Elateryospermum tapos*, *Endospermum malaccense*, *Erythrina* sp., *Eucalyptus deglupta*, *Ficus elastica*, *Garcinia* sp., *Gmelina arborea*, *Gonostylis bancanus*, *Heritiera trifolia*, *Hevea brasiliensis*, *Lithocarpus wallichianus*, *Litsea megacarpa*, *Lophopetalum* sp., *Mangifera* sp., *Millettia dehiscens*, *Parkia speciosa*, *Pithecellobium* sp., *Pometia* sp., *Pouteria malaccensis*, *Rhizophora apiculata*, *Saraca* sp., *Shorea leprosula*, *S. maxwelliana*, *S. parvifolia*, *S. pauciflora*, *Sterculia foetida*, *S. macrophylla*, *Styrax benzoin*, *Terminalia belerica*, *Xanthophyllum* spp., *Xylopia caudata*, *X. cordata*.

Notes: (1) Schedl 1978a: 57 (citation of holotype invalid). (3) Schedl 1941b: 421 (redescription).

References: (**ay**) Nakashima 1975: 6. (**bv**) Gray, B. 1974b. (**cn**) Browne 1949e; Kalshoven 1960d: 48; Supriana et al. 1978. (**cc**) Browne 1958b; Nakashima 1975: 6. (**hb**) Beaver & Browne 1978: 620; Browne 1941, 1958b, 1961c: 205–206; Gray, B. 1974b; Kalshoven 1960c: 36; Mercer 1982: 210. (**ds**) Beaver & Browne 1975: 308, 1978: 620; Beeson 1961: 266; Browne 1961a: 313, 1961c: 205–206, 1972b: 29, 1980d: 492; Choo, Woo, & Kim 1981: 203; Kalshoven 1960d: 48; Mercer

1982: 210; Nakashima 1975: 3; Nobuchi 1977: 137, 1979a: 407; Ohno, Yoneyama, & Nakazawa 1982a: 5, 1982b: 10, 1987b: 90; Ohno, Yoskioka, et al. 1988b: 98; Schedl 1959c: 168, 1961c: 71, 1964c: 304, 1965g: 23, 1965j: 304, 1966c: 97, 1972f: 211, 1974f: 262, 1975g: 294; Supriana et al. 1978. (**tx**) Nakashima 1975: 59; Nobuchi 1985: 329; Schedl 1935n: 635, 1936c: 56, 1939c: 1, 1941b: 421–422, 1942a: 172, 1955i: 215, 1972f: 211, 1978a: 57.

pseudocupulatus sundaensis Schedl 1941b: 421. Holotype ♂; Java, Bandjar; Schedl Collection in NHMW, Wien.

References: (**cn**) Kalshoven 1960d: 48. (**ds**) Browne 1955b: 290; Kalshoven 1960d: 38; Nobuchi 1977: 138; Ohno, Yoneyama, & Nakazawa 1987a: 95; Schedl 1962f: 167, 1969a: 208, 1972f: 211. (**tx**) Nobuchi 1978a: 80, 1978b: 82; Schedl 1941b: 421, 1972f: 211, 1978a: 57.

pseudocupulatus artecanus Schedl 1941b: 422.

Holotype ♂; New Guinea: Papua, Kokoda, 1200 ft.; Schedl Collection in NHMW, Wien.

References: (**ds**) Schedl 1961c: 69, 71, 1965e: 264, 1972f: 211. (**tx**) Schedl 1941b: 422, 1972f: 211, 1978a: 57.

pseudocurtus Schedl 1935n: 635. Lectotype ♀; Malay States: Pahang (Mentakab), Pahang (Tranum), Johore North (Kluang); Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 57.

Distribution: Asia (Malaya/Vietnam).

Hosts: *Dipterocarpus kunstleri*, *Dryobalanops aromatica*, *Shorea faguetiana*, *S. leprosula*, *Intsia palembanica*.

Notes: (3) Browne 1955: 369 (described male).

References: (**hb**) Browne 1941, 1961c: 200. (**ds**) Beeson 1941 (1961: 266); Browne 1961c: 200, 1968a: 132–133, 1980d: 491; Ohno, Yoneyama, & Nakazawa 1982b: 10; Schedl 1972f: 185. (**tx**) Browne 1955: 369; Schedl 1935n: 635–636, 1972f: 185, 1978a: 57.

pseudodignatus Schedl 1963d: 232. Holotype ♂; Brasilien: Sao Paulo, Caraguatatuba, Res. Flor., 40 m; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (**ds**) Schedl 1972f: 241. (**tx**) Schedl 1963d: 232, 1972f: 241, 1978a: 58.

pseudolongulus Schedl 1963d: 233. Holotype ♂; Franz. Guayana; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil/ Cayenne).

References: (**ds**) Schedl 1972f: 205, 1976a: 58. (**tx**) Schedl 1963d: 233, 1972f: 205, 1978a: 58.

pseudoplicatus Schedl 1933c: 168. Holotype ♂; Costa Rica, San Jose, 1000–1200 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: (**ds**) Blackwelder 1947: 790; Equihua

- & Atkinson 1987: 27; Schedl 1933c: 168, 1972f: 241. **(tx)** Equihua & Atkinson 1987: 27; Schedl 1933c: 168, 1972f: 241, 1978b: 58.
- pseudoporus** Schedl 1979g: 117. Holotype ♂; Papua, Upper Manki L.A., Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1979g: 117.
- pseudoselysi** Schedl 1972h: 71. Holotype ♂; New Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(ds)** Schedl 1972f: 224. **(tx)** Schedl 1972f: 224, 1972h: 71, 1978a: 58.
- pseudosindorae** Browne 1983a: 570. Holotype ♂; Wapoga (West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: "Himelio, Matoa" logs.
References: **(ds)** Ohno, Yoneyama, & Nakazawa 1987a: 95. **(tx)** Browne 1983a: 570.
- pseudosolidus** Schedl 1936e: 56. Lectotype ♂; Sumatra, Si-Rambe; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 58.
Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo, Sumatra).
Hosts: *Castanopsis sumatrana*, *Pasania* sp.
Notes: (3) Browne 1955: 370 and Schedl 1971: 394 (described female).
References: **(hb)** Browne 1961c: 209. **(ds)** Schedl 1971c: 370, 1972f: 199. **(tx)** Browne 1955: 370; Nunberg 1961: 624; Schedl 1936e: 56, 1953c: 291, 1971c: 394, 1972f: 199, 1978a: 58.
- pseudoterminatus** Schedl 1936j: 35. Lectotype ♂; Malay Peninsula, Pahang: Rotan Tunggal Forest Reserve; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 58.
Distribution: Asia (Malaya).
References: **(hb)** Browne 1941. **(ds)** Browne 1961c: 213; Schedl 1936j: 35, 1972f: 186. **(tx)** Schedl 1936j: 35, 1941c: 363, 1972f: 186, 1978a: 58.
- ptochoides** Schedl 1970b: 367. Holotype ♂; Philippine Islands, Talomo to Miyako (Japan), imported; PPST, Tokyo.
Distribution: Philippine Islands.
References: **(ds)** Browne 1984c: 499; Nobuchi 1977: 127; Schedl 1972f: 196. **(tx)** Nobuchi 1985: 329; Schedl 1970b: 367–368, 1972f: 196, 1978a: 58.
- puerulus** (Schedl) 1939e: 357 (*Crossotarsus*). Lectotype ♂; Malaya, Perak: Trolak; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 59.
Distribution: Asia (Malaya), Indonesia (Borneo).
Hosts: *Dialium* sp., *Elateriospermum tapos*, *Laurineae* sp., *Lophopetalum* sp., *Nephelium* sp., *Xylopia cordata*.
Notes: (1) Schedl 1972f: 237 (to *Platypus*).
References: **(hb)** Browne 1961c: 200. **(ds)** Beeson 1941 (1961: 254); Browne 1961c: 200, 202, 1981b: 599, 1986c: 662; Schedl 1972f: 237. **(tx)** Schedl 1939e: 357–358, 1941c: 356, 1950g: 895–896, 1972f: 237, 1978a: 59.
- puerulus parvulus** Schedl 1939e: 358 (*Crossotarsus*). Holotype ♂; Kepong, Malaya; Schedl Collection in NHMW, Wien.
References: **(ds)** Schedl 1972f: 237. **(tx)** Schedl 1939e: 358, 1972f: 237, 1978a: 59.
- pulicaris** Chapuis 1865: 165. Holotype ♂; du Bresil, Ste. Catherine; IRSNB, Brussels (Schedl 1960c: 26).
Figures: Bright 1972d: 19.
Hosts: *Cedrela fissula*.
Distribution: Antilles Islands (Guadeloupe/ Jamaica/ Puerto Rico/ Trinidad), South America (Argentina/ Bolivia/ Brazil/ Cayenne/ Colombia/ Paraguay/ Suriname).
References: **(cn)** Ruffinelli & Carbonell 1944: 27. **(ds)** Blackwelder 1947: 790; Bright 1972d: 22; Browne 1981b: 600; Equihua & Atkinson 1987: 27; Gemminger & Harold 1872: 2700; Schedl 1967d: 5, 1972f: 234, 1972g: 40, 1973d: 164, 1976a: 58, 1978c: 292, 1979e: 58; Schonherr & Pedrosa-Macedo 1981: 58; Strohmeyer 1912c: 12, 1914c: 25. **(tx)** Bright 1972d: 19, 22; Chapuis 1865: 165; Equihua & Atkinson 1987: 27; Reichardt 1964b: 88; Schedl 1934e: 212, 1938i: 28, 1939j: 565, 1941e: 155, 1952e: 123, 1959m: 557, 1960c: 26, 1972f: 234; Strohmeyer 1912c: 12, 1914c: 25.
- melanurus** Chapuis 1865: 180. Holotype ♂; de la Colombie; IRSNB, Brussels (Schedl 1960c: 30). Synonymy: Reichardt 1964: 88.
References: **(ds)** Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Nunberg 1958a: 481, 1972b: 190; Strohmeyer 1912c: 11, 1914c: 25. **(tx)** Chapuis 1865: 180; Nunberg 1958a: 481–482; Reichardt 1964: 88; Schedl 1936c: 237, 1948d: 41, 1960c: 30, 1972f: 234; Strohmeyer 1912c: 11, 1914c: 25.
- schaumi** Chapuis 1865: 181. Syntypes ♂ ♀; du Porto-Rico; 1 ♂, 1 ♀ in MNB, Berlin. Synonymy: Reichardt 1964: 88.
References: **(ds)** Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Reichardt 1948: 355; Strohmeyer 1912c: 13, 1914c: 25; Wolcott 1936: 317, 1948: 355. **(tx)** Chapuis 1865: 181; Reichardt 1964: 88; Schedl 1936c: 237, 1937d: 33, 39, 1951h: 287, 1952a: 446, 1972f: 234; Strohmeyer 1912c: 13, 1914c: 25.
- pulvinatus** Roberts & Morimoto 1986: 175. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
References: **(tx)** Roberts & Morimoto 1986: 175.
- quadricornutus** Roberts 1989: 281. Holotype ♂; Papua New Guinea: Upper Stony Logging Area, Bulolo, 1200 m; NHMW, Wien.
References: **(tx)** Roberts 1989: 281.

- pumilus** **Browne** 1955a: 195. Holotype ♂; Kadat (Sabah) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Borneo).
Hosts: *Shorea* sp.
References: **(tx)** Browne 1955a: 195.
- quadricaudatulus** **Schedl** 1934e: 211. Holotype ♂; Caracas [Venezuela]; Schedl Collection in NHMW, Wien.
Distribution: South America (Venezuela).
References: **(ds)** Blackwelder 1947: 790; Schedl 1969c: 66, 1972f: 219. **(tx)** Schedl 1934e: 211, 1972f: 219, 1978a: 60.
- quadricinctus** **Schedl** 1962i: 77. Holotype ♂; New South Wales: Tweed River; Schedl Collection in NHMW, Wien.
Distribution: Australia (New South Wales).
References: **(ds)** Schedl 1962i: 74, 1972f: 201. **(tx)** Schedl 1962i: 77, 1972f: 201, 1978a: 60.
- quadricornis** **Schedl** 1957d: 123. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
Figures: Schedl 1962k: 673, 1972f: 226.
Distribution: Africa (Zaire).
Hosts: *Polyalthia suaveolens*.
References: **(tx)** Mayne & Donis 1960: 100, 1962: 286; Schedl 1957d: 123–124, 1962k: 673, 1972f: 226.
- quadridentatus** (**Olivier**) 1795: 5 (*Scolytus*). Holotype ♂; Amerique septentrionale [USA]; MNHN, Paris, not found, presumed lost.
Figures: Atkinson 1989c.
Distribution: North America (Alabama, Arkansas, Florida, Georgia, S Indiana, Louisiana, Maryland, Mississippi, North Carolina, E Oklahoma, South Carolina, Tennessee, E Texas, Virginia, West Virginia in USA).
Hosts: *Quercus* spp.
Notes: A report of this species from Arizona requires confirmation.
References: **(ay)** Hopkins 1894g: Khalaf 1980: 339. **(cn)** Anonymous 1968g; Blackman 1950; Chamberlin 1939: 110–112; Doane et al. 1936; Hubbard 1897b: 15; Seymour 1968. **(cc)** Chamberlin 1939: 110–112; Felt 1906: 720. **(hb)** Atkinson 1989c; Baker, W. L. 1972: 273; Beal & Massey 1945: 64–65; Beal et al. 1952: 114–115; Blackman 1922b: 39–41, 1950; Chamberlin 1939: 110–112; Doane et al. 1936; Felt 1906: 720; Hetrick 1967; Hopkins 1894g: 277; Hubbard 1897b: 15; Husson 1955: 350. **(ds)** Anonymous 1968g; Atkinson 1989c; Atkinson et al. 1991: 153; Beal & Massey 1945: 64–65; Blackman 1922b: 39–41, 1950; Blatchley & Leng 1916: 583; Chamberlin 1939: 110–112; Drooz 1985: 375; Equihua & Atkinson 1987: 28; Gemminger & Harold 1872: 2700; Henshaw 1885: 147; Hopkins 1893a: 127, 1893b: 207; Schedl 1972f: 223; Schwarz 1878d: 468; Staines, C. L. 1982; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 85; Wood, S. L. 1979a: 2. **(tx)** Atkinson 1989c; Beal & Massey 1945: 64–65; Blackman 1922b: 39–41; Blatchley & Leng 1916: 583; Chamberlin 1939: 110–112; Chapuis 1865: 338; Equihua & Atkinson 1987: 28; Hubbard 1897b: 15; LeConte 1868: 151, 1876: 343–344, 1878a: 468; Olivier 1795b: 5; Schedl 1972f: 223; Strohmeier 1912c: 12, 1914c: 25; Swaine 1909: 85; Wood, S. L. 1972b: 249, 1979a: 2.
- blanchardi** Chapuis 1865: 185. Syntypes ♂; Texas: 2 ♂ in MNB, Berlin, 1 ♂ in MNHN, Paris (Schedl 1960c: 31). Synonymy: LeConte 1876: 344.
References: **(ds)** Blackwelder 1939; Gemminger & Harold 1872: 2697; Leng 1920: 337; Schedl 1933c: 171; Swaine 1909: 86. **(tx)** Chapuis 1865: 185; LeConte 1868: 151, 1876: 344; Schedl 1937d: 39, 1960c: 31, 1972f: 223; Swaine 1909: 86; Wood, S. L. 1979a: 2.
- disciporus** Chapuis 1865: 219. Holotype ♀; de l'Amerique boreale, Tennesse, Coll. de M. Dohrn; presumably Stettin Museum, lost, 1 ♀ not in type series, det. Chapuis in MNB, Berlin. Synonymy: Wood 1958b: 37, 1972b: 249.
References: **(hb)** Chamberlin 1939: 110. **(ds)** Chamberlin 1939: 110; Gemminger & Harold 1872: 2698; Leng 1920: 337; Sampson 1928a: 3; Strohmeier 1912c: 13, 1914c: 26; Swaine 1909: 85; Wood, S. L. 1958b: 37. **(tx)** Chamberlin 1939: 110; Chapuis 1865: 219; LeConte 1868: 151, 1876: 343; Sampson 1928a: 3; Schedl 1937d: 40, 1960c: 31, 1961i: 232, 1972f: 223; Strohmeier 1912c: 13, 1914c: 26; Swaine 1909: 85; Wood, S. L. 1958b: 37, 1972b: 249, 1979a: 1.
- quadrifissilis** **Schedl** 1936e: 58. Lectotype ♂; Mt. Makiling, Laguna, P.I., 400 Fuss; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 60.
Distribution: Asia (Andaman Islands in India/Malaya), Indonesia (Borneo, Mentawai), Philippine Islands.
Hosts: *Bombax insignis*, *Pterocymbium cinctorium*, *Salnalia insignis*, *Scaphium affine*, *Shorea leprosula*, *Sterculia villosa*.
References: **(cn)** Mathur & Singh 1961a: 10. **(hb)** Browne 1941, 1961c: 202. **(ds)** Beeson 1961: 266; Browne 1961a: 313, 1961c: 202, 1980c: 484, 1984c: 449; Mathur & Singh 1961a: 10; Nobuchi 1977: 116; Schedl 1961f: 96, 1966b: 94, 1966g: 33, 1969c: 66, 1972f: 227. **(tx)** Nobuchi 1985: 330; Schedl 1936e: 58, 1939c: 336, 1942a: 172, 1961f: 96, 1972f: 227, 1978a: 60.
- quadrinotatus** **Schedl** 1975g: 231. Holotype ♂; New Guinea, Wan; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
Notes: (3) Roberts 1989: 281 (described female).
References: **(tx)** Schedl 1975g: 231, 1978a: 60.

quadriporus (Beeson) 1937: 96 (*Crossotarsus*).
Syntypes ♂ ♀; Bengal: Darjeeling division.
Debrepani, 6000 feet; FRI, Dehra Dun.
Distribution: Asia (Bhutan/ Bengal in India).
Hosts: *Quercus lamellosa*, *Q. lineata*.
Notes: (1) Schedl 1972f: 219 (to *Platypus*).

References: (ds) Schedl 1972f: 219, 1975c: 384.
(tx) Beeson 1937: 96–97; Schedl 1969c: 66, 1972f: 219, 1978a: 60.

quadrispinatus Chapuis 1865: 221. Syntypes ♂; de la Colombie; 1 ♂ in MNB, Berlin, 1 ♂ in IRSNB, Brussels.

Distribution: South America (Brazil/ Cayenne/ Colombia/ Venezuela).

Hosts: *Cedrela fissilis*.

Notes: (3) Schedl 1936c: 239 (described female).
References: (cn) Morris 1955: 139. (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Sampson 1928a: 3; Schedl 1972f: 219, 1976a: 58; Schonherr & Pedrosa-Macedo 1981: 58; Strohmeier 1912c: 13, 1914c: 26. (tx) Chapuis 1865: 221; Sampson 1928a: 3; Schedl 1934e: 211, 1935i: 44, 1936c: 239, 1937d: 33, 40, 1948f: 283, 1959m: 555, 1960c: 40, 1972f: 219, 1978a: 61; Strohmeier 1912c: 13, 1914c: 26.

quadrispinis Browne 1980d: 496. Holotype ♂; Manokwari (New Guinea) to Nagoya (Japan), imported; BMNH, London.

Figures: *Ficus* sp.

Distribution: New Guinea.

References: (ds) Browne 1984b: 287. (tx) Browne 1980d: 496, 1984c: 292.

quaesitus Schedl 1934e: 210. Holotype ♀; Costa Rica, St. Jose [San Jose]; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 28; Schedl 1972f: 241. (tx) Equihua & Atkinson 1987: 28; Schedl 1934e: 210, 1972f: 241, 1978a: 61.

queenslandi Schedl 1936g: 518. Lectotype ♂; North Queensland; Schedl Collection in NHMW, Wien.

Distribution: Australia (Queensland).

Hosts: *Alphitonia petrici*, *Araucaria cunninghamii*, *Canarium australasicum*, *Juglans* sp.

References: (ds) Schedl 1936g: 518, 1972f: 185, 1980b: 185. (tx) Schedl 1936g: 518–519, 1958i: 214, 1972f: 185, 1978b: 61.

querceus Wood 1972b: 251. Holotype ♂; Volcan de Chiriqui (near Cerro Punta), Chiriqui, Panama; Wood Collection.

Distribution: North America (Costa Rica/ Panama).

Hosts: *Quercus* sp.

References: (ds) Equihua & Atkinson 1987: 28. (tx) Equihua & Atkinson 1987: 28; Wood, S. L. 1972b: 251.

querqi Browne 1980c: 488. Holotype ♂; Keelung (Formosa) to Nagoya (Japan), imported; BMNH, London.

Distribution: Asia (Taiwan).

Hosts: *Quercus* sp.

References: (tx) Browne 1980c: 488.

quercicola Schedl 1971c: 394. Holotype ♂; Malay Penin.: Kelantan Chabong, Jongtrat F.R.; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Quercus* sp.

References: (ds) Schedl 1972f: 185. (tx) Schedl 1971c: 394–395, 1972f: 185.

quercinus Schedl 1942a: 213. Holotype ♂; Malaya, N.S., Pasoh F.R.; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Castanopsis sumatrana*, *Quercus irayi*.

References: (ds) Browne 1961c: 218; Schedl 1972f: 217. (tx) Schedl 1942a: 213, 1972f: 217.

quercivorius Murayama 1925b: 229. Syntypes ♂ ♀; Taiwan: Murayama Collection in USNM, Washington.

Figures: Nobuchi 1973b: pl. 2.

Distribution: Asia (Bengal in India/ Japan/ Taiwan), Indonesia (Java), New Guinea.

Hosts: *Castanopsis cuspidata*, *Cryptomeria japonica*, *Ilex chinensis*, *Lindera erythrocarya*, *Lithocarpus edulis*, *Pasania glabra*, *Prunus* sp., *Quercus acuta*, *Q. acutissima*, *Q. gilva*, *Q. glauca*, *Q. mouglolica* var. *grosseserrata*, *Q. myrsinaefolia*, *Q. salicina*, *Q. serrata*, *Q. sessilifolia*.

Notes: (1) Schedl 1972f: 217 (to *Platypus*).

References: (cn) Anonymous 1980g; Murayama 1954a: 17. (ds) Anonymous 1980g; Murayama 1929a: 678, 1936a: 140, 1937c: 580, 1954a: 17, 1955: 100, 105; Nobuchi 1973b: 14; Schedl 1972f: 217. (tx) Beeson 1937: 94; Murayama 1925b: 229–231, 235, 1928a: 287, 1930b: 28–29, 1936a: 140–141, 1955: 100, 105; Nobuchi 1973b: 14, pl. 2; Schedl 1939c: 4, 1972f: 217, 1972p: 156.

sexfenestratus Beeson 1937: 94 (*Crossotarsus*).

Syntypes ♂ ♀; Bengal: Kalimpong division, Samsingh; FRI, Dehra Dun. Synonymy: Schedl 1972p: 156.

Notes: (1) Schedl 1972f: 217 (to *Platypus*).

References: (ds) Beeson 1941 (1961: 266); Kalshoven 1960d: 41. (tx) Beeson 1937: 94–95; Schedl 1939c: 4, 1941c: 354, 1969c: 66, 1972f: 217, 1972p: 156.

quinquecostatus Chapuis 1865: 149. Syntypes ♂; du Mexique; 2 ♂ in IRSNB, Brussels (Schedl 1960c: 22).

Distribution: North America (El Salvador/ Guatemala/ Mexico).

Hosts: *Hevea brasiliensis*.

References: (ds) Blackwelder 1947: 790; Blandford 1896c: 103; Equihua & Atkinson 1987: 28; Ferrer 1942; Gemminger & Harold 1872: 2700; Schedl 1972f: 241, 1977c: 44; Strohmeier 1912c:

- 9, 1914c: 24. (tx) Blandford 1896e: 103; Chapuis 1865: 149; Equihua & Atkinson 1987: 28; Schedl 1940a: 324, 1960c: 22, 1972f: 241; Strohmeier 1912c: 9, 1914c: 24.
- ramali Schedl** 1940c: 203. Syntypes ♂; Costa Rica, Ramal Parismina, Sta. Clara; Schedl Collection in NHMW, Wien.
Distribution: North America (Costa Rica/ Panama).
References: (ds) Equihua & Atkinson 1987: 28; Schedl 1972f: 241. (tx) Equihua & Atkinson 1987: 28; Schedl 1940c: 203, 1972f: 241, 1978a: 61.
- ramosissimus Roberts & Morimoto** 1986: 176. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Cryptocarya* sp.
References: (tx) Roberts & Morimoto 1986: 176.
- ramulosus Roberts & Morimoto** 1986: 177. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Cryptocarya* sp.
References: (tx) Roberts & Morimoto 1986: 177.
- rasilis Schedl** 1979g: 118. Holotype ♀; Papua, Upper Manki L.A., Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1979g: 118.
- ratzeburgi Chapuis** 1865: 227. Syntypes ♂ ♀; de la Guyanne française, de la Colombie, et du Brésil; 1 ♀ in MNB, Berlin, 2 ♂ in BMNH, London, 1 ♂, 1 ♀ in IRSNB, Brussels.
Distribution: Antilles Islands (Cuba/ Guadeloupe/ Puerto Rico/ Trinidad), North America (Costa Rica/ El Salvador), South America (Argentina/ Bolivia/ Brazil/ Cayenne/ Colombia/ Paraguay/ Peru/ Suriname/ Venezuela).
Hosts: *Erythrina costaricensis*, *Inga* sp., *Lonchocarpus margaritensis*.
References: (hb) Kalshoven 1963; Swabey 1935: 7. (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 28; Gemminger & Harold 1872: 2700; Kalshoven 1963: 238; Martorell 1945: 467–468; Santoro 1957b: 26; Schedl 1933c: 174, 1960a: 78, 1970e: 87, 1972f: 218, 1972g: 50, 1977e: 44; Strohmeier 1912c: 14, 1914c: 26; Swabey 1935: 7; Wolcott 1936: 316, 1948: 385. (tx) Chapuis 1865: 227; Equihua & Atkinson 1987: 28; Schedl 1936e: 55, 1937d: 40, 1939j: 565, 1941e: 156, 1948d: 37, 1952a: 444, 1952e: 123, 1952h: 73, 1957a: 194, 1960a: 78, 1960c: 41–42, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.
- raucis Schedl** 1936c: 224. Syntypes ♂ ♀; Guayana (?); Schedl Collection in NHMW, Wien.
Distribution: South America (Cayenne/ Suriname).
Notes: (1) Schedl 1978a: 62 (citation of holotype invalid).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 241. (tx) Schedl 1936c: 224, 1972f: 241, 1978a: 62.
- refractus Roberts & Morimoto** 1986: 178. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
References: (tx) Roberts & Morimoto 1986: 178.
- reichei Chapuis** 1865: 135. Syntypes ♂ ♀; de la Colombie; 1 ♀ in MNB, Berlin, 1 ♂ in IRSNB, Brussels (Schedl 1960c: 19).
Distribution: North America (Costa Rica/ Nicaragua/ Panama), South America (Brazil/ Colombia).
Hosts: *Ficus* sp., etc.
Notes: (3) Breeds only in healthy, green tissue of living trees.
References: (ds) Blackwelder 1947: 790; Blandford 1896e: 98; Equihua & Atkinson 1987: 28; Gemminger & Harold 1872: 2700; Schedl 1933c: 166, 1972f: 241; Strohmeier 1912c: 9, 1914c: 24. (tx) Blandford 1896e: 98; Chapuis 1865: 135; Equihua & Atkinson 1987: 28; Schedl 1934e: 210, 1935e: 176, 1935h: 354, 1936c: 225, 1960c: 19, 1972f: 241; Strohmeier 1912c: 9, 1914c: 24.
- retusipennis Schedl** 1964c: 301. Holotype ♂; Ostkuste Sumatras, Brastagi, 1300 m; ZMA, Amsterdam.
Distribution: Indonesia (Sumatra).
References: (ds) Schedl 1972f: 212. (tx) Schedl 1964c: 301, 1972f: 212.
- retusulus Schedl** 1975a: 460. Holotype ♂; Vietnam: Tonkin, Kelao; Schedl Collection in NHMW, Wien.
Distribution: Asia (Tonkin Island in Vietnam).
References: (ds) Schedl 1975a: 453. (tx) Schedl 1975a: 453, 460.
- retusus Strohmeier** 1910e: 132. Holotype ♂; Nilgiri Hills (India orientalis); H. L. Andrews Collection.
Distribution: Asia (India).
References: (ds) Schedl 1972f: 212; Strohmeier 1912c: 19, 1914c: 29. (tx) Schedl 1972f: 212; Strohmeier 1910e: 132, 1912c: 19, 1914c: 29.
- rimulosus Schedl** 1950e: 215. Lectotype ♀; Cote d'Ivoire, Adiopodoume, upon Avodire; Schedl Collection in NHMW, Wien.
Distribution: Africa (Ghana/ Ivory Coast).
Hosts: *Turraecanthus africana*.
References: (ce) Jover 1952: 73–81. (hb) Caclian 1957: 6, 15; Jover 1952: 73–81. (ds) Browne 1984b: 287; Schedl 1962h: 63, 1962k: 640, 1972f: 216. (tx) Schedl 1950e: 215, 1954d: 874, 885, 1957d: 120–121, 1962k: 640, 1972f: 216, 1978a: 62.
- roberti Chapuis** 1865: 183. Syntypes ♂; de Madagascar; 1 ♂ in MNB, Berlin, 2 ♂ in MNHN, Paris.
Distribution: Madagascar.
Notes: (3) Browne 1972c: 108 (described female).
References: (ds) Alluand 1900: 441; Fairmaire 1892b; Gemminger & Harold 1872: 2700; Numberg 1964: 237; Schedl 1972f: 234, 1977b: 253;

- Strohmeyer 1912c: 12, 1914c: 25. (tx) Browne 1972c: 108; Chapuis 1865: 183; Fairmaire 1892b; Schanfuß 1897b: 109; Schedl 1936c: 237, 1937d: 33, 39, 1960c: 31, 1961e: 156–157, 1964m: 316, 1967f: 149, 1972f: 234, 1977b: 253, 1978b: 62; Strohmeyer 1912c: 12, 1914c: 25.
- robustus Chapuis** 1865: 146. Holotype ♂; du Venezuela; BMNH, London (Schedl 1960c: 21). Distribution: South America (Venezuela). References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 241; Strohmeyer 1912c: 7, 1914c: 24. (tx) Chapuis 1865: 146; Schedl 1960c: 21, 1972f: 241; Strohmeyer 1912c: 9, 1914c: 24.
- roppai Schedl** 1978c: 308. Holotype ♂; Brasilien, Linhares, E. Santo; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1978c: 308.
- rotundicauda Motschulsky** 1863: 509. Syntypes 2 ♀; des Montagnes de Nura-Ellia, syntypes labeled India Orientalis; IZM, Moscow. Distribution: Asia (Sri Lanka). References: (hb) Speyer 1923: 23. (ds) Gemminger & Harold 1872: 2700; Schedl 1972f: 242; Strohmeyer 1912c: 19, 1914c: 29. (tx) Motschulsky 1863: 509; Schedl 1972f: 242; Strohmeyer 1912c: 19, 1914c: 29.
- rufescens (Strohmeyer)** 1912f: 82 (*Crossotarsus*). Holotype ♂; Kamerun; MNB, Berlin, and Strohmeyer Collection. Figures: Schedl 1962k: 643. Distribution: Africa (Cameroon/ Ivory Coast/ Nigeria/ Sierra Leone/ Zaire/ Zambia). Hosts: *Ochthocosmus africanus*, *Parinarium holstii*, *Terminalia superba*. Notes: (1) Schedl 1972f: 223 (to *Platypus*). References: (ds) Schedl 1933f: 202, 1972f: 223, 1979b: 416; Strohmeyer 1912c: 6, 1914c: 36. (tx) Schedl 1933f: 202, 1935g: 313–314, 1957d: 118, 120–123, 1962k: 641, 1972f: 223; Strohmeyer 1912c: 6, 1912f: 82, 85, 1914c: 36.
- elongatus Schedl** 1935g: 313 (*Stenoplatypus*). Lectotype ♂; Belgian Congo: Sandoa, Lulua; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 27, preoccupied by Chapuis 1865. Notes: (3) Schedl 1972f: 223 (treated as a subspecies of *rufescens*). References: (ds) Schedl 1972f: 223. (tx) Schedl 1933f: 202, 1935g: 313–314, 1950d: 11, 1952i: 23, 1957d: 120, 123, 1959c: 165, 1962k: 642, 1972f: 223, 1978a: 27.
- rufescens parinariae Schedl** 1957d: 122. Holotype ♀; Kivu, Hembe-Bitale, 1800 m; MRCB, Teruren. Notes: (3) Schedl 1972f: 223 (treated as a subspecies of *rufescens*). References: (ds) Roberts 1968: 189, 1969: 119; Schedl 1962k: 642, 1972f: 223. (tx) Schedl 1957d: 120, 122–123, 1962k: 642–643, 1972f: 223, 1972p: 159, 1978a: 63.
- ochthocosmus Roberts** 1970b: 480. Holotype ♂; Sierra Leone: Gola West Forest Reserve; BMNH, London. Synonymy: Schedl 1972f: 223. References: (ds) Beaver & Loytyniemi 1989. (tx) Beaver & Loytyniemi 1989; Roberts 1970b: 480; Schedl 1972f: 223, 1972p: 159.
- rufobrunneus Schedl** 1942a: 207. Holotype ♀; Tenasserim; Schedl Collection in NHMW, Wien. Distribution: Asia (Burma). References: (ds) Schedl 1972f: 242. (tx) Schedl 1942a: 207, 1972f: 242, 1978a: 63.
- rugosifrons Schedl** 1933c: 173. Holotype ♂; Brazil, S. Paulo, Alto da Serra; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil/ Venezuela). References: (ds) Blackwelder 1947: 790; Schedl 1933c: 173, 1972f: 234. (tx) Schedl 1933c: 173, 1972f: 234, 1978a: 63; Wood, S. L. 1992a: 91.
- pretiosus Schedl** 1961i: 233. Holotype ♂; Venezuela, Mt. Duida; CAS, San Francisco. Synonymy: Wood 1992a: 91. References: (ds) Schedl 1960a: 78, 1972f: 234. (tx) Schedl 1961i: 233, 1972a: 234, 1978a: 56; Wood, S. L. 1992a: 91.
- runcinatus Roberts** 1986: 52. Holotype ♂; Papua New Guinea: Mt. Giluwe, 2000 m; BMNH, London. Figures: Roberts 1986: 57. Distribution: New Guinea. Hosts: *Nothofagus pullei*, *Vataca* sp. References: (tx) Roberts 1986: 52, 57.
- sallei Chapuis** 1865: 218. Syntypes 2 ♂; de la Colombie; BMNH, London. Distribution: South America (Colombia/ Venezuela). References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 219; Strohmeyer 1912c: 13, 1914c: 26. (tx) Chapuis 1865: 218; Schedl 1939b: 399, 1948f: 283, 1960c: 40, 1972f: 219; Strohmeyer 1912c: 13, 1914c: 26.
- salvini Blandford** 1896e: 103. Syntypes 2 ♂; Guatemala, Cerro Zunil; BMNH, London. Figures: Blandford 1896e: pl. 4, fig. 14. Distribution: North America (Guatemala). References: (ds) Blackwelder 1947; Equihua & Atkinson 1957: 28; Schedl 1972f: 241; Strohmeyer 1914c: 24. (tx) Blandford 1896e: 103; Equihua & Atkinson 1957: 28; Schedl 1972f: 241; Strohmeyer 1914c: 24.
- sampsoni (Schedl)** 1933f: 202 (*Stenoplatypus*). Lectotype ♂; South Africa, Zululand: Eshowe, and Port St. John: Pondoland; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 63. Distribution: Africa (South Africa).

- Hosts: *Cassipourea gummiflua*, *Fagara capensis*, *Olea laurifolia*, *Virgilia capensis*.
References: (cn) Anonymous 1970c: 12. (cc) Walt & Scott 1971a; Walt, Scott, & van der Klift 1971b: 463. (hb) Browne 1961c: 221. (ds) Anonymous 1970c: 12; Schedl 1933f: 202–203, 1962k: 658, 1969d: 6, 1970h: 179, 1972f: 216, 1975k: 279. (tx) Schedl 1933f: 202–203, 1948c: 665, 1957c: 326, 1957d: 120, 1962k: 658–659, 1972f: 216.
- sandakanensis** Browne 1970: 577. Holotype ♂; Sabah; Sandakan; BMNH, London.
Distribution: Indonesia (Borneo).
References: (ds) Browne 1980c: 484, 1981b: 598; Olmo, Yoneyama, & Nakazawa 1987a: 95; Schedl 1972f: 199. (tx) Browne 1970: 577–578; Schedl 1972f: 199.
- santacruzensis** Mutchler 1925: 233. Holotype ♂; Galapagos Islands; Seymour Bay, Indefatigable Island; AMNH, New York.
Distribution: South America (Galapagos Islands).
References: (ds) Blackwelder 1947: 790; Mutchler 1924: 232; Schedl 1972f: 234, 1974c: 52, 1978c: 293. (tx) Mutchler 1924: 232; Schedl 1972f: 234.
- santiriae** Roberts 1970: 75. Holotype ♂; Nigeria; Ngel Nyaki Forest Reserve, Mambilla Plateau, 1650 m; MRCB, Tervuren.
Distribution: Africa (Nigeria).
Hosts: *Santiria trimera*.
References: (tx) Roberts 1970: 75.
- saravakensis** (Schedl) 1937e: 546 (*Crossotarsus*).
Lectotype ♂; Saravak, foot of Mt. Dulit, junction of rivers Tinjar and Lejok; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 64.
Distribution: Indonesia (Saravak in Borneo).
References: (ds) Schedl 1972f: 224. (tx) Schedl 1937e: 544–547, 1972f: 224, 1978a: 64.
- scalaris** Schedl 1980b: 188. Holotype ♂; Nth [Northern] Queensland, Australia; Schedl Collection in NHMW, Wien.
Distribution: Australia (Queensland).
Hosts: *Flindersia pubescence*.
References: (tx) Schedl 1980b: 188.
- sculptor** Schedl 1975f: 394. Holotype ♀; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 394.
- schenklingi** (Strohmeyer) 1912f: 81 (*Crossotarsus*). Syntypes ♂; Kamerun; MNB, Berlin, and Strohmeyer Collection.
Distribution: Africa (Cameroon/ Congo/ Ghana/ Ivory Coast/ Nigeria/ Sierra Leone/ Tanzania/ Uganda/ Zaïre).
Hosts: Schedl 1962k: 638–639 (15 host species listed).
Notes: (3) Schedl 1939g: 169–170 (described female).
References: (cc) Schedl 1962k: 637. (hb) Browne 1963a: 252; Schedl 1962k: 637. (ds) Browne 1963a: 252; Mayne & Donis 1960: 100, 1962: 287; Roberts 1968: 189, 1969: 119; Schedl 1961m: 85, 1962h: 63, 1962k: 637, 1108, 1964j: 42, 1972f: 223; Strohmeyer 1912c: 6, 1914c: 36. (tx) Schedl 1937d: 42, 1939g: 169–170, 1941d: 414, 1952i: 22, 1957d: 118, 120, 1962k: 637, 1972f: 223, 1978a: 64; Strohmeyer 1912c: 6, 1912f: 81–85, 1914c: 36.
- schmidtii** Chapuis 1865: 150. Holotype ♂; de Cayenne; IRSNB, Brussels (Schedl 1960c: 22).
Distribution: South America (Cayenne).
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 241; Strohmeyer 1912c: 9, 1914c: 24. (tx) Chapuis 1865: 150; Schedl 1936c: 234, 1939b: 399, 1948d: 39–40, 1960c: 22, 1972f: 241; Strohmeyer 1912c: 9, 1914c: 24.
- schultzeanus** Schedl 1966b: 98. Syntypes ♂; Luzon, Mountain Province, Benguet, Mount Santo Tomas; W. Schultze Collection, and Schedl Collection in NHMW, Wien, automatic.
Distribution: Philippine Islands (Luzon).
References: (ds) Schedl 1966b: 98, 1972f: 200. (tx) Nobuchi 1955: 329; Schedl 1966b: 98, 1972f: 200.
- schultzei** Schedl 1935k: 481 (*Crossotarsus*). Syntypes ♂; Mountain Province, Benguet, Mount Santo Tomas; W. Schultze Collection, and Schedl Collection in NHMW, Wien, preoccupied by Strohmeyer 1911.
References: (ds) Murayama 1957a: 37; Strohmeyer 1912c: 18. (tx) Schedl 1935k: 481–482, 1940b: 435, 1966b: 98, 1972f: 200.
- scrobicularis** Roberts 1979: 95. Holotype ♂; Papua New Guinea: 8 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Euodia* sp.
References: (tx) Roberts 1979: 95.
- secretus** Sampson 1921: 25. Syntypes ♂ ♀; Tonkin, Hoabinh; BMNH, London.
Distribution: Asia (Bangladesh/ Bhutan/ Burma/ Assam, Bengal in India/ Laos/ Malaya/ Tonkin Island in Vietnam).
Hosts: *Aesculus pindana*, *Albizia* spp., *Alphonsea ventricosa*, *Anthocephalus cadamba*, *Artocarpus fraxinifolius*, *Cinnamomum* sp., *Dalbergia assamensis*, *Diospyros* spp., *Dipterocarpaceae pilosus*, *Hymenodictyon excelsum*, *Lancea grandis*, *Meliosma simplicifolia*, *Odina woderi*, *Ostodes paniculata*, *Sterculia ornata*, *Terminalia bclerica*.
References: (cn) Kleine 1932a: 310, 402; Mathur & Singh 1960b: 61, 1961a: 70, 1961b: 26; Roonwal & Singh 1954: 33, 40. (cc) Beeson 1921b: 24. (hb) Kleine 1932a: 310, 402. (ds) Beaver & Browne 1975: 308; Beeson 1921b: 24, 1941 (1961: 266); Bhasin, Roonwal, & Singh 1955; Browne 1970: 576, 1972b: 29; Kleine 1932a: 310,

402; Mathur & Singh 1960b: 61, 1961a: 70, 1961b: 26; Roonwal 1954: 33, 40; Sampson 1921: 25; Schedl 1962f: 167, 1971c: 363, 1972a: 191, 1974c: 262, 1975c: 384. **(tx)** Browne 1964: 755, 1972b: 29; Sampson 1921: 25; Schedl 1972f: 191. *cruditus* Schedl 1942a: 212. Lectotype ♂; Tonkin, Region de Hoa-Binh; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 28. Synonymy: Schedl 1972f: 191. References: **(tx)** Schedl 1942a: 212, 1972f: 191, 1978a: 28.

secus Wood 1972b: 248. Holotype ♂; La Carbonera Experimental Forest, 50 km W Merida, Merida, Venezuela; Wood Collection.

Distribution: South America (Venezuela).

Hosts: *Eschweileria* sp., *Nectandra* sp., *Podocarpus raspigliosii*.

References: **(tx)** Wood, S. L. 1972b: 248.

segnis Chapuis 1865: 166. Holotype ♂; de la Nouvelle-Grenade; IRSNB, Brussels (Schedl 1960c: 27). Distribution: Antilles Islands (Guadeloupe), North America (Costa Rica/ Guatemala/ Chiapas, Veracruz in Mexico/ Panama), South America (Bolivia/ Brazil/ Colombia/ Peru/ Venezuela).

References: **(en)** Kleine 1932a: 402. **(cc)** Equihua & Atkinson 1986: 625. **(hb)** Burgos & Saucedo 1983: 46; Equihua & Atkinson 1986: 625; Kleine 1932a: 402. **(ds)** Atkinson & Equihua 1986a: 423, 1988: 104; Atkinson et al. 1986: 47; Blackwelder 1947: 790; Burgos & Saucedo 1983: 46; Equihua 1985: 144; Equihua & Atkinson 1986: 625, 1987: 29; Estrada & Atkinson 1988: 213; Gemminger & Harold 1872: 2700; Kleine 1932a: 402; Numberg 1971: 59; Schedl 1966f: 94, 1972f: 234; Strohmeier 1912c: 13, 1914c: 25; Wille 1943: 315. **(tx)** Chapuis 1865: 166; Equihua & Atkinson 1987: 29; Schedl 1952h: 72, 1960c: 27, 1972f: 234; Strohmeier 1912c: 13, 1914c: 25.

poriferus Chapuis 1865: 168. Syntypes 2 ♂; de Toxpan; BMNH, London (Schedl 1960c: 27). Synonymy: Schedl 1972f: 234.

References: **(ds)** Blackwelder 1947: 790; Blandford 1896c: 106; Ferrer 1942; Schedl 1933c: 172, 1960a: 75, 1970c: 90; Strohmeier 1912c: 12, 1914c: 25. **(tx)** Blandford 1896c: 106; Chapuis 1865: 168; Keler 1927a: 235; Schedl 1940a: 324–325, 1941e: 156, 1960c: 27, 1972f: 234; Strohmeier 1912c: 12, 1914c: 25.

viduus Chapuis 1865: 178. Holotype ♂; Mexico; Haag Collection or lost (Schedl 1960c: 30). Synonymy: Schedl 1972f: 234.

References: **(ds)** Gemminger & Harold 1872: 2700. **(tx)** Chapuis 1865: 178; Schedl 1960c: 30, 1972f: 234.

sellaeformis Roberts & Morimoto 1986: 178. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Cryptocarya* sp.

References: **(tx)** Roberts & Morimoto 1986: 178.

selysi Chapuis 1865: 215. Syntypes 2 ♂, 1 ♀; de la Nouvelle Guinee, Dorey; BMNH, London (Schedl 1960c: 39).

Figures: Gray & Wylie 1974: 81.

Distribution: New Guinea.

Hosts: *Aleurites muluccana*, *Evodia* sp., *Hevea brasiliensis*, *Horsfieldia irya*, *Intsia* sp., *Myristica* sp., *Polyalthia* sp.

References: **(hb)** Gray, B. 1968: 307. **(ds)** Gemminger & Harold 1872: 2700; Gray, B. 1968: 307; Schedl 1962i: 73, 1964c: 305, 1964d, 1968c: 264, 1972f: 224; Smece 1964: 25; Strohmeier 1912c: 13, 1914c: 25. **(tx)** Chapuis 1865: 215; Gray, B. & Wylie 1974: 81; Schedl 1939b: 398, 1940b: 434–435, 1941c: 154, 1960c: 39, 1964d: 213, 1972f: 224; Strohmeier 1912c: 13, 1914c: 25.

semicarinatus Schedl 1942a: 203. Lectotype ♂; Malaya, Pahang, Cameron's Highlands; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 64.

Distribution: Asia (Malaya).

References: **(ds)** Browne 1961c: 198; Schedl 1972f: 206. **(tx)** Schedl 1942a: 203, 1942c: 194, 1972f: 206, 1978a: 64.

semiermis Schedl 1941c: 362. Syntypes ♂ ♀; Java, Mount Gede, 800 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya), Indonesia (Borneo, Java).

Hosts: *Pasania* sp.

Notes: (1) Schedl 1978a: 65 (citation of holotype invalid).

References: **(hb)** Kalshoven 1960c: 41. **(ds)** Browne 1961c: 218; Kalshoven 1960d: 41; Schedl 1971c: 370, 1972f: 217. **(tx)** Schedl 1941c: 362–363, 1942a: 172, 1958h: 498, 1972f: 217.

semigranosus (Sampson) 1925: 2 (*Crossotarsus*). Holotype ♂; Queensland: Malanda; BMNH, London.

Distribution: Australia (Queensland).

Hosts: *Argyrodendron* sp., *Cinnamomum* sp., *Elaeocarpus* sp., *Flindersia pubescens*, *Panax* sp.

Notes: (3) Schedl 1980b: 189 (described female).

References: **(en)** Brimblecombe 1951; Froggatt 1926b, 1927. **(hb)** Brimblecombe 1951; Froggatt 1926b, 1927. **(ds)** Sampson 1925: 2; Schedl 1972f: 185, 1980b: 185. **(tx)** Sampson 1925: 2; Schedl 1936g: 516, 1972f: 185, 1980b: 189.

semiopacus Strohmeier 1913: 163. Holotype ♂?; Brit. Neu-Guinea. Aroa-Fluss; Strohmeier Collection.

Distribution: New Guinea.

Hosts: *Podocarpus* sp.

References: **(ds)** Schedl 1968e: 264, 1968f: 535, 1972f: 185; Strohmeier 1914c: 29. **(tx)** Schedl 1935e: 174, 1936g: 517, 1939b: 398, 1969b: 231, 1972f: 185; Strohmeier 1913: 163, 1914c: 29.

- semisulcatus** Schedl 1969b: 231. Holotype ♂; New Guinea: Kuni, 11 miles from Mt. Hagen; CSIRO, Canberra.
Distribution: New Guinea.
References: (ds) Schedl 1972f: 212. (tx) Schedl 1969b: 231, 1972f: 212, 1978a: 65.
- senexus** Wood 1966a: 66. Holotype ♂; 16 km SE Cartago on Pan-American Highway, Cartago Prov., Costa Rica, 5600 feet; Wood Collection.
Figures: Schedl 1972f: 220, Wood 1966a: 67.
Distribution: North America (Costa Rica).
References: (ds) Equihua & Atkinson 1987: 29; Schedl 1972f: 220. (tx) Equihua & Atkinson 1987: 29; Schedl 1972f: 220; Wood, S. L. 1966a: 66–67.
- sepaloides** Roberts 1959: 281. Holotype ♂; Papua New Guinea: Upper Stony Logging Area, Bulolo, 1200 m; NHMW, Wien.
Figures: Roberts 1959: 286.
Distribution: New Guinea.
Hosts: *Castanopsis* sp.
References: (tx) Roberts 1959: 281, 286.
- sequis** Schedl 1935i: 44. Syntypes ♂; Guadeloupe, env. de Trois-Rivieres; MNHN, Paris, and Schedl Collection in NHMW, Wien.
Distribution: Antilles Islands (Guadeloupe).
Notes: (1) Schedl 1978a: 65 (type restricted to MNHN, Paris specimens).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 219. (tx) Schedl 1935i: 44, 1972f: 219, 1978a: 65.
- sericans** Schedl 1942a: 211. Lectotype ♀; Malaya, Perak, Trolak F.R.; Schedl Collection, designated by Schedl 1978a: 65.
Distribution: Asia (Malaya).
Hosts: *Vatica* sp.
References: (ds) Browne 1961c: 200; Schedl 1972f: 242. (tx) Schedl 1942a: 210–211, 1972f: 242, 1978a: 65.
- setaceus** Chapuis 1865: 234. Syntypes 2 ♂; des îles Philippines; IRSNB, Brussels (Schedl 1960c: 43).
Distribution: Philippine Islands, New Guinea.
Hosts: *Diplodiscus paniculatus*.
Notes: (3) Schedl 1935k: 484 (described female).
References: (cn) Ishikura 1966. (ds) Baer 1886; Browne 1983a: 554; Gemminger & Harold 1872: 2700; Ishikura 1966; Nobuchi 1977: 122; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1966b: 98, 1972f: 191; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 234; Nobuchi 1985: 329; Schedl 1933c: 105, 1935k: 484–486, 1942a: 212, 1960c: 43, 1972f: 191, 1978a: 66; Strohmeier 1911b: 17, 1912c: 16, 1914c: 27.
- severiini** Blandford 1894d: 136. Syntypes (many) ♀; Japan: Nikko, Chiuzenji, and Hakodate; BMNH, London.
Figures: Nakane et al. 1963: pl. 192, Nobuchi 1973b: pl. 2.
Distribution: Asia (Japan/Taiwan).
Hosts: *Acer mono*, *Aesculus turbinata*, *Ahus hirsuta*, *A. japonica*, *Camellia japonica*, *Carpinus japonicus*, *Fagus crenata*, *Fraxinus spachiana*, *F. lanuginosa*, *Prunus* sp., *Quercus gilva*, *Tilia japonica*.
References: (ay) Korolev 1989; Nakashima 1972. (cn) Anonymous 1980g; Ishikura 1966; Kleine 1932a: 402; Uchida et al. 1958: 181. (cc) Banno, Mikata, & Kodama 1983: 445; Inouye et al. 1955: 122; Nakashima 1972. (hb) Inouye et al. 1955: 122; Kleine 1932a: 402; Murayama 1931b: 195; Nakashima 1972. (ds) Anonymous 1980g; Blandford 1894c: Ishikura 1966; Kleine 1932a: 402; Miwa 1931: 266; Murayama 1925a: 212, 1929a: 672, 1931b: 195, 1934f: 133–149, 1937c: 578, 1949c: 104, 1950b: 1299, 1954b: 188; Nakane et al. 1963: 384; Nobuchi 1967: 27, 1973b: 15; Schedl 1934f: 1647, 1972f: 199; Strohmeier 1912c: 16, 1914c: 27. (tx) Blandford 1896e: 136; Korolev 1989; Murayama 1925a: 212, 1925b: 235, 1928a: 283–284, 1950b: 1299, 1954b: 188; Nakane et al. 1963: 384, pl. 2; Nisima 1909: 171, 1910a: 5, 1913a: 5; Nobuchi 1967: 26, 1973b: 15, pl. 2; Schedl 1959d: 68, 1972f: 199; Strohmeier 1912c: 16, 1914c: 27.
- sexcostatus** Chapuis 1865: 126. Holotype ♂; du Bresil; MNHN, Paris (Schedl 1960c: 16).
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 241; Strohmeier 1912c: 9, 1914c: 24. (tx) Chapuis 1865: 126; Schedl 1936c: 225, 1960c: 16, 1972f: 241; Strohmeier 1912c: 9, 1914c: 24.
- sexporus** (Schedl) 1935k: 484 (*Crossotarsus*). Holotype ♀; Philippinen; Schedl Collection in NHMW, Wien.
Distribution: Asia (Vietnam), Indonesia (Java), Philippine Islands.
Hosts: *Castanea argentea*, *C. javanica*, *C. tunggurut*.
Notes: (1) Schedl 1954a: 138 (to *Platypus*).
References: (ds) Schedl 1966b: 98, 1972f: 217, 1974c: 262. (tx) Beeson 1937: 93; Nobuchi 1985: 329; Schedl 1935k: 484, 1939c: 4, 1939e: 335, 1954a: 138, 1972f: 217, 1978a: 66.
- sexforaminatus** Beeson 1937: 92. Syntypes ♂ ♀; Philippine Is.; BMNH, London. Synonymy: Schedl 1939c: 4, 1972f: 335.
References: (tx) Beeson 1937: 92–94; Schedl 1939c: 4, 1939e: 335, 1972f: 217.
- sexualis** Beeson 1937: 95. Syntypes ♂ ♀; United Provinces: Chakrata division, Mandali, 6000 feet; FRI, Dehra Dun.
Distribution: Asia (Uttar Pradesh in India).
Hosts: *Quercus dilatata*.
Notes: (1) Schedl 1972f: 217 (to *Platypus*).
References: (ds) Beeson 1941 (1961: 266); Schedl 1972f: 217. (tx) Beeson 1937: 95–96; Schedl 1939c: 4, 1972f: 217.

- seydeli** (Nunberg) 1967: 326 (*Stenoplatypus*). Holotype ♂; Elisabethville [Congo]; MRCB, Tervuren. Figures: Nunberg 1967b: 339. Distribution: Africa (Zaire). References: (tx) Nunberg 1967b: 326, 338–339.
- shillongensis** Schedl 1969c: 66. Holotype ♂; India: Assam, Shillong, 6000 feet; FRI, Dehra Dun. Distribution: Asia (Assam in India, Malaya). Hosts: *Acer campbelli*, *Livistona jenkinsiana*. Notes: (1) None of the specimens borrowed from FRI by Schedl were ever returned to FRI; the type of this species is probably among the Schedl paratypes (SLW). (3) Beeson 1941: 345, 1961: 266 (*shillongensis*, nomen nudum, synonymy in Schedl 1969c: 66). References: (ds) Beeson 1941: 345, 1961: 266; Schedl 1972f: 189. (tx) Schedl 1969c: 66–67, 1972f: 189, 1978a: 66.
- shoreanus** (Beeson) 1937: 98 (*Crossotarsus*). Syntypes ♂ ♀; Assam: Lakhimpur division, Upper Dihing reserve; FRI, Dehra Dun. Distribution: Asia (Burma/ Cochin in China/ Assam in India/ Malaya/ Sri Lanka/ Vietnam), Indonesia (Borneo), New Guinea, Philippine Islands. Hosts: *Anisoptera cochinchinensis*, *A. polyandra*, *Balanocarpus heimii*, *Dipterocarpus zeylanicus*, *Doonia zeylanicus*, *Hopsea odorata*, *Shorea assamica*, *S. bracteolata*, *S. singkawang*, *Terminalia brassii*. Notes: (1) Schedl 1972f: 224 (to *Platypus*). (3) Schedl 1936d: 7 (*imitator*, nomen nudum, synonymy in Schedl 1972f: 224). References: (cn) Mathur & Singh 1960b: 57, 1961a: 37. (ds) Beeson 1941 (1961: 266); Mathur & Singh 1960b: 57, 1961a: 37; Ohno, Yoneyama, & Nakazawa 1957a: 96; Schedl 1969c: 66, 1972f: 224. (tx) Beeson 1937: 98–99; Schedl 1972f: 224, 1978a: 66; Wylie & Yule 1977.
- doonae** Beeson 1937: 99 (*Crossotarsus*). Syntypes ♂ ♀; Ceylon: S.A.B., Ratnapura, Mawagan-kanda, 500'; FRI, Dehra Dun, not found in 1981 by SLW. Notes: Schedl 1972: 224 (treated as a subspecies). References: (ds) Beeson 1941 (1961: 265); Schedl 1959a: 513. (tx) Beeson 1937: 99–100; Schedl 1959a: 513, 1972f: 224, 1978a: 26.
- bifurcus mutilus** Schedl 1939e: 359 (*Crossotarsus*). Lectotype ♂; Malaya, Selangor: Kepong; Schedl Collection in NHMW, Wien. Notes: (3) Schedl 1972f: 224 (treated as a subspecies), 1942a: 213 (described female). References: (an) Nakashima 1975: 6–8. (ec) Nakashima 1975: 6–8. (ds) Beeson 1941 (1961: 265); Browne 1961a: 314, 1966: 237; Nakashima 1975: 3; Nobuchi 1977: 117; Ohno, Yoneyama, & Nakazawa 1952b: 10; Schedl 1936d: 7, 1959c: 168, 1964i: 249, 1966g: 34, 1972a: 145, 1972f: 224. (tx) Nobuchi 1955: 329; Schedl 1939e: 359, 1942a: 207, 213, 1972f: 224, 1978a: 15; Wylie & Yule 1977.
- shoreanus tersus** Schedl 1972h: 71. Holotype ♂; Wara Sweet logging area, Kui, Morobe District, New Guinea; CSIRO, Canberra. References: (ds) Browne 1983a: 55; Schedl 1972f: 224, 1972h: 71.
- sicarius** Wood 1972b: 250. Holotype ♂; 10 km SE Miri, Barinas, Venezuela; Wood Collection. Distribution: South America (Venezuela). Hosts: *Pouteria anibaefolia*. References: (tx) Wood, S. L. 1972b: 250.
- signatus** Chapuis 1865: 235. Syntypes 3 ♂; de ile de Borneo, Sarawak; BMNH, London. Distribution: Asia (Burma/ India/ Malaya), Indonesia (Sarawak in Borneo, Java, Sumatra). Hosts: *Artocarpus lakoocha*, *Ctenolophon grandifolius*, *Endospermum malaccense*, *Gonystylus bancanus*, *Hevea brasiliensis*, *Kickxia arborea*, *Koompassia malaccensis*, *Lucuma malaccensis*, *Pometia tomentosa*, *Pouteria malaccensis*, *Psychotria viridiflora*, *Schima noronhai*, *Sterculia macrophylla*, *S. urceolata*, *Symplocos* sp., *Xylia dolabriformis*, *X. xylocarpa*. Notes: (3) Schedl 1942a: 208 (described female). References: (cn) Kalshoven 1924: Kleine 1932a: 310, 402; Mathur & Singh 1961a: 70, 1961b: 96; Roonwal 1954: 89; Yunus & Hua 1980: 217. (hb) Browne 1935a, 1941, 1961c: 196; Kalshoven 1960c: 34, 1960d: 34; Kleine 1932a: 310, 402; Mercer 1982: 210. (ds) Beeson 1941 (1961: 266); Browne 1935a, 1961c: 196, 1980c: 484; Gemminger & Harold 1872: 2700; Kalshoven 1924b: 358, 1960d: 34; Kleine 1932a: 310, 402; Mathur & Singh 1961a: 70, 1961b: 96; Mercer 1982: 210; Nobuchi 1977: 119; Roonwal 1954: 89; Sampson 1928b: 392; Schedl 1965g: 24, 1969a: 206, 1971f: 151, 1972f: 194; Strohmeier 1912c: 16, 1914c: 27; Yunus & Hua 1980: 217. (tx) Chapuis 1865: 235; Sampson 1913: 448, 1919: 106, 1928b: 392; Schedl 1937e: 544, 1939b: 398, 1939e: 337, 362–363, 1941c: 354, 1942a: 208, 1960c: 44, 1972f: 194, 1978a: 66; Strohmeier 1912c: 16, 1914c: 27.
- similis** (Nunberg) 1973: 26 (*Stenoplatypus*). Holotype ♂; Massif Ruwenzori, Zalonge, 2100 m, Riv. Nyamwamba, affl. Butahu; MRCB, Tervuren. Distribution: Africa (Zaire). References: (tx) Nunberg 1973: 26.
- simpliciformis** Wood 1966a: 57. Holotype ♂; Tapanti, Cartago Prov., Costa Rica, 4000 feet; Wood Collection. Distribution: North America (Costa Rica). Hosts: *Phoebe mexicana*. References: (ds) Equihua & Atkinson 1987: 29; Schedl 1972f: 206. (tx) Equihua & Atkinson 1987: 29; Schedl 1972f: 206; Wood, S. L. 1966a: 52, 57.
- simulans** Schedl 1941d: 413. Holotype ♂; Syr. [Syrien]; Schedl Collection in NHMW, Wien. Distribution: Asia (Syria/ Turkey).

Hosts: *Alnus* sp., *Liquidambar orientalis*.

References: (ds) Schedl 1961b: 185, 1972f: 191. (tx) Schedl 1941d: 413, 1961b: 186–187, 1972f: 191, 1978a: 66.

siendorae Browne 1980c: 458. Holotype ♂; Singapore to Nagoya (Japan), imported; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Sinodora* sp.

References: (tx) Browne 1980c: 458.

sinensis Schedl 1941a: 43. Holotype ♀; Fukien, Kuatun, 2300 m; Museum Alexander Koenig, Bonn.

Figures: Schedl 1941a: 44.

Distribution: Asia (Fujian in China).

Notes: (1) Schedl 1978a: 67 (Lectotype designation invalid; the designation should have been a lectoallotype because a holotype already existed).

References: (ds) Schedl 1972f: 192. (tx) McNamara 1984: 750; Schedl 1941a: 43, 1953e: 22, 26, 1972f: 192, 1978a: 67.

singalangensis Schedl 1936e: 53. Syntypes ♂; Sumatra, Mte. Singalang; MCG, Genova, and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

Notes: (1) Schedl 1978a: 67 (citation of holotype invalid).

References: (ds) Schedl 1972f: 193. (tx) Schedl 1936e: 53–54, 1972f: 193, 1978a: 67.

singularis Schedl 1975f: 394. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 394.

sinuosus Chapuis 1865: 198. Holotype ♂; de Variñas [now Barinas, Venezuela]; presumably Stettin Museum, lost (Schedl 1960c: 34).

Distribution: South America (Venezuela).

References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 234; Strolmeyer 1912c: 13, 1914c: 25. (tx) Chapuis 1865: 198; Schedl 1960c: 34, 1972f: 234; Strolmeyer 1912c: 13, 1914c: 25.

sirambeensis Schedl 1936e: 59. Syntypes ♂; Sumatra, Si-Rambe; MCG, Genova, and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

Notes: (1) Schedl 1978a: 67 (citation of holotype invalid).

References: (ds) Schedl 1972f: 227. (tx) Schedl 1936e: 59, 1964g: 253, 1972f: 227, 1978a: 67.

sobrinus Schedl 1935h: 353. Syntypes ♂ ♀; Guatemala; Schedl Collection in NHMW, Wien.

Distribution: North America (Guatemala).

Notes: (1) Schedl 1978a: 67 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 29; Schedl 1972f: 241. (tx) Equihua & Atkinson 1987: 29; Schedl 1935h: 353, 1972f: 241, 1978a: 67.

solicitatus Schedl 1975f: 395. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 395.

solidulus Browne 1980: 496. Holotype ♂; Palembang (Sumatra) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sumatra).

References: (ds) Ohno, Yoneyama, & Nakazawa 1982b: 10. (tx) Browne 1980: 496.

solidus Walker 1859: 261. Holotype ♂; Ceylon; BMNH, London.

Figures: Nakashima 1975: 38–41, Nobuchi 1967: pl. 2, 1973b: pl. 2, Wood 1960a: 9.

Distribution: Asia (Burma/ Andaman Islands, Assam, Bengal, Bihar, Karnataka, Orissa, Tamil Nadu in India/ Israel/ Japan/ Korea/ Malaya/ Nepal/ Pakistan/ Singapore/ Sri Lanka/ Taiwan/ Vietnam), Admiralty Islands, Australia (Queensland), Indonesia (Aru, Borneo, Celebes, Damma, Java, Malacca, Sumatra), Micronesia (Caroline Islands, Mariana Islands), New Guinea, Philippine Islands, Solomon Islands.

Hosts: *Acacia catechii*, *Acacia decurrens*, *Acrocarpus fraxinifolia*, *Adina* spp., *Azcleia palembanica*, *Albizia* spp., *Anisoptera* sp., *Anogeissus latifolia*, *Anthocephalus cadamba*, *Artocarpus* spp., *Bombax malabaricum*, *Buchanania* spp., *Butea* spp., *Canarium strictum*, *Carallia suffruticosa*, *Carpinus laxiflora*, *Castanopsis tribuloides*, *Ceratonia siliqua*, *Chloroxylon swietenia*, *Cinnamomum* spp., *Cocos nucifera*, *Cullenia excelsa*, *Dalbergia latifolia*, *D. sissoo*, *Dipterocarpus pilosus*, *Dolichandrone stipulata*, *Drimycarpus racemosus*, *Dryobalanops oblongifolia*, *Duabanga sonneratioides*, *Elateriospermum tapos*, *Eugenia* sp., *Ficus* spp., *Garuga pinnata*, *Gmelina arborea*, *Hevea brasiliensis*, *Holoptelea integrifolia*, *Lannea grandis*, *Litsea polyantha*, *Mallotus albus*, *Mangifera indica*, *Nancelea sessilifolia*, *Odina wodier*, *Palaquium* sp., *Pterocarpus dalbergioides*, *Pterospermum acerifolium*, *Salmalia malabarica*, *Schleichera oleosa*, *S. trijuga*, *Schinus noronhai*, *S. wallichii*, *Semecarpus heterophylla*, *Shorea leprosula*, *S. robusta*, *Sterculia campanulata*, *S. villosa*, *Stereospermum chelonoides*, *Styrax benzoin*, *Suictonia floribunda*, *Syzygium cumini*, *Talauma gioi*, *Tamarindus indica*, *Tectona grandis*, *Terminalia bellerica*, *T. tomentosa*, *Tetrameles nudiflora*.

Notes: (3) Schedl 1937d: 40 (*walkeri*, nomen nudum, synonymy in Schedl 1959a: 513).

- References: **(ay)** Kempers 1932: 69; Nakashima 1975: 5. **(bv)** Beeson 1917; Gray, B. 1974b. **(cn)** Beeson 1915, 1916: 1–5, 1938a; Browne 1938b: 77–86; Chatterjee et al. 1923, 1950; Gray, B. & Wylie 1974; Green 1912a: 2–5, 1916: 608–636; Ishikura 1966; Kalshoven 1924, 1925c: 1, 1960d: 48; Kleine 1932a: 310, 402; Leefmans 1920: 1–90; MacKenzie 1922: 1–14; Mathur & Singh 1960a: 39, 1960b: 6, 1961a: 13, 1961b: 13; Mesa 1931: 18; Ranawera 1959: 78; Roonwal 1954: 18; Roonwal, Chatterjee, & Thapa 1961b: 7; Rutgers 1918: 1–44; Speyer 1923: 11–23; Stebbing 1914: 624; Tammeng-keng & Tantra 1971; Tryon 1917: 49–63; Wylie & Shanahan 1975. **(cc)** Beeson 1921b: 24, 1923; Bhatia 1950; Browne 1955b; Chatterjee & Chatterjee 1951; Halperin & Holzschuh 1984: 29; Mathew 1985; Nakashima 1975: 5; Stebbing 1914: 624. **(hb)** Beeson 1916a: 222, 1917, 1923, 1938a; Browne 1935a, 1941, 1958b, 1961c: 207–209; Chatterjee & Chatterjee 1951; Chatterjee et al. 1950; Gray, B. 1968: 308, 1974b; Gray, B. & Wylie 1974; Halperin & Holzschuh 1984: 29; Kalshoven 1959a: 224–226, 1960c: 39, 1960d: 1; Kleine 1932a: 310, 402; Murayama 1931b: 196; Speyer 1923: 22; Stebbing 1914: 624; Wylie & Shanahan 1975. **(ds)** Beaver & Browne 1975: 308, 1978: 620; Beeson 1916a, 1921b, 1923, 1938a, 1941 (1961: 266); Bhasin, Roonwal, & Singh 1958; Bhatia 1950; Blandford 1895a; Browne 1935a, 1961c: 207, 1966: 235, 1980d: 492, 1984c: 450; Chapuis 1865; Chatterjee & Chatterjee 1951; Chatterjee et al. 1950: 37; Cho 1957; Choo 1983: 33; Choo & Woo 1985: 163; Choo, Woo, & Kim 1981: 203; Gemminger & Harold 1872: 2700; Gray, B. 1968: 308; Green 1912a; Halperin 1976b; Halperin & Holzschuh 1984: 29; Ishikura 1966; Kalshoven 1924b: 358, 1925c: 1, 1960d: 48; Kleine 1932a: 310, 402; Ko 1969: 274; Lacordaire 1866: 392; Mathew 1982, 1987: 188; Mathur & Singh 1960a: 39, 1960b: 6, 1961a: 13, 1961b: 13; Mesa 1935: 96; Miwa 1931: 266; Murayama 1925a: 213, 1929a: 672, 1929b: 2, 1930b: 23, 1931b: 196, 1936b: 113, 1937b: 375, 1937c: 579, 1954b: 205, 1957a: 37; Nakashima 1975: 3; Nobuchi 1967: 27, 1973b: 16, 1977: 129; Numberg 1960b: 161, 1961b: 624; Ohno, Yoneyama, & Nakazawa 1982b: 11, 1987a: 96, 1987b: 90; Ohno, Yoshioka, et al. 1988b: 98, 1989: 67; Roonwal 1954: 18; Roonwal, Chatterjee, & Thapa 1961b: 7; Sampson 1919: 106, 1926: 120; Schedl 1936d: 7, 1936g: 514, 1938g: 423, 1959a: 513, 1961e: 70–71, 1962b: 187, 1962f: 167, 1962i: 75, 1964d: 213, 1965a: 340, 1965g: 23–25, 1966b: 99, 1966g: 34, 1968e: 264, 1971a: 275, 1971c: 368, 1971f: 151, 1972a: 145, 1973b: 212, 1974c: 262, 1975a: 453, 1975e: 453, 1978e: 37; Stroh-meyer 1912c: 17, 1914c: 27; Wood, S. L. 1960a: 7. **(tx)** Chapuis 1865: 267; Choo 1983: 33; Gardiner 1932b; Green 1912a; Kalshoven 1959a: 224; Lacordaire 1866: 392; Lea 1910: 135; Motschul-sky 1863; Murayama 1925a: 213, 1925b: 235, 1930b: 23, 27–28, 31, 1937b: 375, 1950d: 98, 1954b: 205; Nakashima 1975: 38–41; Nobuchi 1967: pl. 2, 1973b: 16, pl. 2, 1985: 329; Sampson 1919: 106; Schedl 1934f: 1647, 1936e: 55–56, 1937d: 40, 1939b: 398, 1939e: 336, 1941e: 156, 1942a: 172, 199, 1942b: 147, 1942c: 164, 1951i: 96, 1953b: 124, 128, 147, 1954a: 145, 1955b: 282–283, 310, 1959a: 513, 1960c: 50, 1969b: 233; Stebbing 1914: 624; Stroh-meyer 1912c: 17, 1912d: 42, 1914c: 27; Walker 1859: 261; Wood, S. L. 1960a: 7–9, 1969c: 120; Wylie & Yule 1977.
- caudatus* Motschulsky 1863: 509. Syntypes 6 ♂; Nura-Ellia, specimens labeled India oriental; IZM, Moscow. Synonymy: Schedl 1959a: 513, Wood 1969c: 120.
- References: **(ds)** Gemminger & Harold 1872: 2698. **(tx)** Motschulsky 1863: 509; Numberg 1959c: 168, 1961b: 624; Schedl 1959a: 513, 1960h: 110, 1972f: 199; Wood, S. L. 1969c: 120.
- cordatus* Motschulsky 1863: 510. Syntypes 4 ♀; Des Montagnes de Nura-Ellia, syntypes labeled India oriental; IZM, Moscow. Synonymy: Wood 1969c: 120.
- References: **(hb)** Speyer 1923: 23. **(ds)** Gemminger & Harold 1872: 2698; Schedl 1972f: 242; Stroh-meyer 1912c: 19, 1914c: 28. **(tx)** Motschulsky 1863: 510; Schedl 1972f: 242; Stroh-meyer 1912c: 19, 1914c: 28; Wood, S. L. 1969c: 120.
- pilifrons* Chapuis 1865: 265. Syntypes 4 ♀; Du continent et de l'Archipel indiens; BMNH, London (Schedl 1960c: 50). Synonymy: Stroh-meyer 1912d: 42.
- References: **(bv)** Beeson 1917. **(cn)** Kalshoven 1925c: 1; Kleine 1932a: 310, 402; Mathur & Singh 1961a: 45. **(hb)** Beeson 1916a: 223, 1917; Kleine 1932a: 310, 402. **(ds)** Beeson 1916a: 223; Gemminger & Harold 1872: 2699; Kalshoven 1925c: 1; Kleine 1932a: 310, 402; Lefroy & Hawlett 1909: 395; Mathur & Singh 1961a: 45; Sampson 1919: 106, 1926: 120; Stroh-meyer 1912c: 17. **(tx)** Beeson 1921: 24; Chapuis 1865: 265; Murayama 1930: 28; Sampson 1919: 106; Schedl 1959a: 513, 1960c: 50, 1972f: 199; Stroh-meyer 1912c: 17, 1912d: 42; Wood, S. L. 1960a: 8.
- solidus exilis* Chapuis 1865: 268. Syntypes, sex?, Iles Celebes et specialement aux environs de Macassar; BMNH, London.
- Notes: (3) Schedl 1972f: 199 (treated as a sub-species). An apparent aberration.
- References: **(ds)** Schedl 1972f: 199. **(tx)** Chapuis 1865: 268; Schedl 1933e: 105, 1972f: 199.
- solidus obtusus* Chapuis 1865: 268. Syntypes, sex?, Ceylan; BMNH, London, preoccupied by Chapuis 1865: 265.

- Notes: (3) Schedl 1972f: 199 (treated as a subspecies). An apparent aberration.
References: (tx) Chapuis 1865: 265; Schedl 1972f: 199.
- solidus rudis* Chapuis 1865: 268. Syntypes, sex?; Iles de Baichian et de Marty dans les Molucques; BMNH, London.
Notes: (3) Schedl 1972f: 199 (treated as a subspecies). It is an apparent aberration.
References: (ds) Schedl 1972f: 199. (tx) Chapuis 1865: 268; Schedl 1972f: 199.
- solomonicus* Browne 1954b: 291. Holotype ♂; Ringi Cove (Solomon Islands) to Nagoya (Japan), imported; BMNH, London.
Distribution: Solomon Islands.
Hosts: *Neonauclaea* sp.
References: (ds) Ohno, Yoshioka, et al. 1958b: 97–98, 1959: 67. (tx) Browne 1954b: 291.
- solutus* Schedl 1933f: 193. Lectotype ♂; Haut-Uele; Moto, Avakubi, Equateur; Boende, Haut-Uele; Abimva, Koteli, Sankuru, Kindu, Likimi, Ituri; Mahagi, Avakubi; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 67. Figures: Schedl 1962k: 600.
Distribution: Africa (Angola/ Cameroon/ Congo/ Equatorial Guinea/ Kenya/ Malawi/ Mozambique/ Nigeria/ Sierra Leone/ Somalia/ South Africa/ Tanzania/ Zaïre/ Zambia), Madagascar.
Hosts: Many listed in Schedl 1962k: 632–634. *Ficus sycomorus*, *Melia azedarach*.
References: (cn) Halperin & Menier 1951; Mayne & Donis 1951: 337; Tordo 1957: 8. (ec) Mayne & Donis 1951: 337; Schedl 1962k: 629. (hb) Schedl 1962k: 629. (ds) Browne 1975a: 757; Ferreira 1965: 1130; Lee 1971: 30; Mayne & Donis 1960: 100, 1962: 287; Menier 1951; Roberts 1968: 188, 1969: 119; Schedl 1934a: 247, 1935d: 452, 1943c: 70, 1959p: 22, 1962k: 629, 1964f: 619, 1965e: 357, 1966c: 230, 1971e: 4, 1971g: 196, 1972f: 234, 1972k: 296, 1975: 279; Spahr 1951: 80. (tx) Powell, W. 1957: 28; Schedl 1933f: 193, 1934a: 247, 1937d: 39, 1939a: 468, 1941d: 381, 1941e: 155, 1948d: 42, 1950d: 4, 11, 14, 1952g: 54, 1952j: 5, 1953g: 243, 1955c: 31, 1955f: 260, 1962k: 600, 629, 1972f: 234, 1978a: 67.
- flavicornis* Dahman 1825: 381, 404–405. Syntypes, sex?; Sierra Leone Africa; not located, preoccupied by Fabricius 1776. Synonymy: Schedl 1972f: 234.
References: (tx) Dahman 1825: 381, 404–405; Schedl 1972f: 234.
- spectabilis* Schedl 1935b: 396. Holotype ♀; Philippine Islands; Schedl Collection in NHMW, Wien.
Distribution: Philippine Islands.
Hosts: *Dipterocarpus grandiflorus*.
References: (ds) Schedl 1938g: 424, 1966b: 100, 1972f: 192. (tx) Nobuchii 1985: 329; Schedl 1935b: 395–397, 1941a: 43, 1941c: 360, 1972f: 192, 1978a: 68.
- spectus* Wood 1972b: 247. Holotype ♂; Rancho Grande in Pittier National Park, Aragua, Venezuela; Wood Collection.
Distribution: South America (Venezuela).
References: (tx) Wood, S. L. 1972b: 247.
- spiculatus* Browne 1972b: 30. Holotype ♂; Malaya; Sungei Buloh; BMNH, London.
Distribution: Asia (Malaya).
References: (ds) Browne 1986c: 663; Ohno, Yoshioka, et al. 1959: 67. (tx) Browne 1972b: 30.
- spiniventris* Schedl 1969b: 232. Holotype ♂; New Guinea; Kumi, 11 miles from Mt. Hagen; CSIRO, Canberra.
Distribution: New Guinea.
Notes: (3) Roberts 1959: 283 (described female).
References: (ds) Schedl 1972f: 227. (tx) Schedl 1969b: 232–233, 1972f: 227, 1978a: 68.
- spinosus* Browne 1981a: 135. Holotype ♂; Way Semangka (Sumatra) to Sakaide (Japan), imported; BMNH, London.
Distribution: Indonesia (Sumatra).
Hosts: *Dipterocarpus* sp.
References: (ds) Mayne & Donis 1960: 98, 1962: 276. (tx) Browne 1981a: 135.
- spinulosus* (Strohmeyer) 1912: 80 (*Stenoplatypus*). Syntypes ♂ ♀; Kamerun, Deutsch-Ostafrika und Französisch-Kongo; MNB, Berlin, Hamburg Museum, lost, Strohmeyer Collection.
Figures: Schedl 1962k: 646, 1972f: 214.
Distribution: Africa (Cameroon/ Congo/ Gabon/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaïre/ Zimbabwe).
Hosts: Schedl 1962k: 646–647 listed 21 host species.
References: (ay) Strohmeyer 1918: 42. (ec) Jover 1952: 73–81; Schedl 1962k: 644. (hb) Browne 1963a: 252; Jones, Roberts, & Baker 1959: 13–14; Jover 1952: 73–81; Schedl 1962k: 644; Webb & Jones 1957: 37, 41. (ds) Browne 1963a: 252; Gardner 1957a: 29; Mayne & Donis 1960: 100, 1962: 288; Numborg 1952: 22, 1965b: 22; Roberts 1968: 189, 1969: 119; Schedl 1961m: 85, 1962h: 63, 1962k: 644, 1967e: 221, 1971e: 4, 1972e: 284, 1972f: 216, 1982: 279; Strohmeyer 1911a: 182, 1912c: 6, 1914c: 36. (tx) Numborg 1952: 22; Powell, W. 1957: 28; Schedl 1937d: 42, 1939b: 398, 1941c: 361, 1950d: 19, 1950e: 212, 1954d: 874, 1957d: 118, 120, 1962k: 644, 646, 1108, 1972f: 216; Strohmeyer 1911a: 182, 1912c: 6, 1912f: 80–82, 85, 1914c: 35–36.
- spimulosus montivagus* Roberts 1970a: 77. Holotype ♂; Nigeria; Ngel Nyaki Forest Reserve, 1650 m, Mambilla Plateau; BMNH, London.
References: (tx) Roberts 1970a: 77.
- squameus* Schedl 1942a: 200. Holotype ♂; Tenasserim; Schedl Collection in NHMW, Wien.
Distribution: Asia (Burma/ Thailand), Indonesia (Borneo, Sumatra).
References: (ds) Beaver & Browne 1975: 309;

- Browne 1972b: 29; Schedl 1964c: 304, 1972f: 183. (tx) Schedl 1942a: 200, 1972f: 183, 1978a: 68.
- squamifer** Schedl 1963d: 233. Holotype ♀; Brasilien: Sao Paulo; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Schedl 1972f: 219. (tx) Schedl 1963d: 233, 1972f: 219, 1978a: 68.
- stenoplicatus** Schedl 1939c: 3. Lectotype ♂; N.E. Burma, Kambaiti, 7000 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 69.
Distribution: Asia (Burma, Fujian in China).
References: (ds) Schedl 1972f: 219. (tx) Schedl 1939c: 3, 1955h: 46, 1972f: 219, 1978a: 69.
- strenuus** Schedl 1969b: 233. Holotype ♂; New Guinea: Togoba [Western Highlands District]; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Anisoptera polyandra*, *Cinnamomum* sp., *Eucalyptus grandis*.
Notes: (3) Schedl 1973i: 96 (described female).
References: (ds) Schedl 1972f: 199. (tx) Schedl 1969b: 233, 1972f: 199, 1973i: 96, 1978a: 69.
- striatodeclivis** Schedl 1975f: 396. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: (tx) Schedl 1975f: 396.
- striatopennis** Schedl 1952a: 463. Syntypes 2 ♂; Argentina, Misiones, Departamento Concepcion Santa Maria; Viana Collection, and Schedl Collection in NHMW, Wien.
Distribution: South America (Argentina).
Notes: (1) Schedl 1978a: 69 (citation of holotype invalid).
References: (ds) Schedl 1972f: 218. (tx) Schedl 1952a: 463, 1972f: 218, 1978a: 69.
- striatopunctatus** Schedl 1942a: 210. Holotype ♀; Borneo; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 201. (tx) Schedl 1942a: 210, 1972f: 201, 1978a: 69.
- striatus** Chapuis 1865: 270. Holotype ♂; du Bresil; BMNH, London (Schedl 1960c: 51).
Distribution: South America (Brazil).
References: (ds) Blackwelder 1947; Gemminger & Harold 1872: 2700; Schedl 1972f: 234; Strohmeier 1912c: 17, 1914c: 27. (tx) Chapuis 1865: 270; Schedl 1960c: 51; Strohmeier 1912c: 17, 1914c: 27.
- strigillatus** Schedl 1941c: 359. Syntypes ♂; Java, Mount Gede, 800 m; Kalshoven Collection, and Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Java).
Hosts: *Castanea javanica*.
Notes: (1) Schedl 1978a: 69 (citation of holotype invalid).
References: (hb) Kalshoven 1960c: 41. (ds) Kalshoven 1960c: 41; Schedl 1972f: 195. (tx) Schedl 1941c: 359–360, 1954a: 145, 158, 1972f: 195, 1978a: 69.
- suavifer** Schedl 1970e: 109. Holotype ♂; Brasilien, Amazonas, Sierra Neblina, M. Rio Canaburi, 1000 m; Zoologischen Sammlungen des Bayrischen Staates, Munchen.
Distribution: South America (Brazil).
References: (ds) Schedl 1970e: 87, 1972f: 241. (tx) Schedl 1970e: 87, 109–110, 1972f: 241.
- subaequalispinosus** Schedl 1936i: 100. Holotype ♂; Bolivien; Schedl Collection in NHMW, Wien.
Figures: Schedl 1972f: 225.
Distribution: South America (Bolivia/ Peru).
References: (ds) Blackwelder 1947: 790; Schedl 1972f: 225. (tx) Schedl 1936i: 100, 1958c: 1–2, 1972f: 225, 1978a: 69.
- subangustior** Schedl 1942a: 207. Lectotype ♀; Malaya, Kuala Lumpur; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 69.
Distribution: Asia (Malaya).
Hosts: *Albizia falcata*, *A. moluccana*.
References: (ds) Browne 1961c: 220; Schedl 1972f: 201. (tx) Schedl 1942a: 207, 1972f: 201, 1978a: 69.
- subcuratus** Browne 1983a: 563. Holotype ♂; Tg. Usan (West Irian) to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: Watergum log.
References: (ds) Browne 1986c: 663; Olmo, Yoneyama, & Nakazawa 1987a: 96. (tx) Browne 1983a: 563.
- subdepressus** (Schedl) 1935b: 397 (*Crossotarsus*). Syntypes ♂ ♀; Mount Maquiling, Laguna Province, and Quezon Park, Tayabas Province, Luzon, Philippine Islands; F. C. Hadden Collection, and Schedl Collection in NHMW, Wien.
Distribution: Asia (Assam in India/ Malaya), Indonesia (Borneo, Sumatra), Philippine Islands (Luzon).
Hosts: *Dipterocarpus pilosus*, *Ehretia acuminata*, *Meliosma simplicifolia*.
Notes: (1) Schedl 1978a: 70 (citation of holotype invalid).
References: (ds) Browne 1981b: 598, 1984c: 151; Nobuchi 1977: 129; Olmo, Yoneyama, & Nakazawa 1987b: 90; Schedl 1934a: 247, 1962f: 167, 1966b: 88, 1966g: 34, 1972f: 202. (tx) Nobuchi 1985: 329; Schedl 1934a: 247, 1935b: 397–398, 1935k: 481, 1937d: 34, 1939b: 394, 395, 1962f: 167, 1969c: 64, 1972f: 202, 1978a: 70.
- sandakanus** Beeson 1937: 64 (*Crossotarsus*). Holotype ♂; Borneo: Sandakan, Baker; BMNH, London. Synonymy: Schedl 1939e: 333.
Notes: (1) Schedl 1972f: 202 (to *Platypus*).
References: (tx) Beeson 1937: 64–65; Schedl 1939e: 333, 1972f: 202.
- latchunatus** Beeson 1937: 66 (*Crossotarsus*).

Holotype ♂; Assam: Sibsagar division, Central Range, Nakachari; FRI, Dehra Dun. Synonymy: Schedl 1972f: 202.

References: (cn) Sen-Sarma & Thakur 1956: 37. (ds) Beeson 1941 (1961: 254); Browne 1961a: 314. (tx) Beeson 1937: 66–67; Browne 1962e: 654; Schedl 1972f: 202.

subgranosus Schedl 1936g: 516. Lectotype ♂; Tasmania: Waratah and Queensland: Dividing Range V.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 70.

Figures: Britton 1970: 621, Neumann & Harris 1974: 134

Distribution: Anstralia (Queensland, Tasmania).

Hosts: *Brachyhiton acerifolium*, *B. populicum*, *Eucalyptus maculata*, *Nothofagus cunninghamii*, *Pterocymbium beccarii*, *Scolopia brounii*.

References: (bv) Elliott, H. J., Madden, & Bashford 1983; Elliott, H. J. et al. 1987. (cn) Brimblecombe 1951; Fisher 1952a; Fisher, Thompson, & Webb 1953; Kile, Elliot, & French 1976; Neumann & Harris 1974; Neumann & Marks 1976. (cc) Batra 1963a; Elliott, H. J. et al. 1987; Kile 1987; Kile & Hall 1988; Neumann & Harris 1974; Slaby 1947: 378; Webb 1945: 57. (hb) Brimblecombe 1951; Elliott & Little 19..: 49, 65; Fisher 1952a; Neumann & Harris 1974; Neumann & Marks 1976. (ds) Hogan 1948: 373; Schedl 1959d: 67–68, 1972f: 185, 1979a: 160. (tx) Britton 1970: 621; Carne et al. 1980; Neumann & Harris 1974: 134; Schedl 1936g: 516–517, 1941e: 155, 1942c: 164, 1958i: 214.

subgranulosa Browne 1955a: 194. Holotype ♂; Lawas (Sarawak) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Sarawak in Borneo).

References: (tx) Browne 1955a: 194.

subitarius Schedl 1936c: 241. Holotype ♂; Brazil, Rio de Janeiro; Schedl Collection in NHMW, Wien.

Figures: Reichardt 1964b: 87.

Distribution: North America (Costa Rica), South America (Argentina/ Brazil).

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 29; Numberg 1963c: 107, 1964a: 237; Schedl 1966f: 94, 1967d: 5, 1970e: 87, 1972f: 218, 1973d: 163, 1976a: 58, 1979e: 58. (tx) Equihua & Atkinson 1987: 29; Reichardt 1964b: 87–89; Schedl 1936c: 241, 1948f: 283, 1972f: 218, 1978a: 70.

perdifficilis Schedl 1936c: 245. Holotype ♂; Rio Grande do Sul [Brazil]; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1948f: 283. References: (ds) Blackwelder 1947: 790. (tx) Schedl 1936c: 240, 245, 1948f: 283, 1972f: 218, 1978a: 53.

subnitens Browne 1983: 571. Holotype ♂; Lever Harbour (Solomon Islands) to Nagoya (Japan), imported; BMNH, London.

Distribution: Solomon Islands.

References: (ds) Ohno, Yoshioka, et al. 1988b: 98. (tx) Browne 1983a: 571.

suboblitaratus Schedl 1940c: 204. Syntypes ♂; Costa Rica, Turrialba, 500 m and San Jose, 1000–1200 m; Neumann Collection and Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

Notes: (1) Schedl 1978a: 71 (citation of holotype invalid).

References: (ds) Equihua & Atkinson 1987: 29; Schedl 1972f: 241. (tx) Equihua & Atkinson 1987: 29; Schedl 1940c: 204, 1972f: 241, 1978a: 71.

subplicatus Schedl 1962b: 193. Holotype ♂; Indochina: Reg. de Hoa-Binh; MNHN, Paris.

Distribution: Asia (Tonkin Island in Vietnam, also Vietnam).

References: (ds) Schedl 1972f: 191. (tx) Schedl 1962b: 193, 1964c: 300, 1972f: 191, 1978a: 71.

subpraeteritus Schedl 1975g: 231. Holotype ♂; Papua, Porotop, Luth. Miss. Sawmill; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975g: 231, 1978a: 71.

subpronus Schedl 1969b: 233. Holotype ♂; New Guinea: Watut Valley 1200 m, Morobe District; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Aglaia* sp.

Notes: (3) Schedl 1975f: 396 (described female).

References: (ds) Schedl 1972f: 201. (tx) Schedl 1969b: 233–234, 1972f: 201, 1975f: 396, 1978a: 71.

subsecretus Browne 1964: 755. Holotype ♂; Burma: Ruby Mines; BMNH, London.

Distribution: Asia (Burma).

References: (ds) Schedl 1972f: 191. (tx) Browne 1964: 755; Schedl 1972f: 191.

subsidarius Schedl 1935j: 2. Holotype ♂; Sumatra, Medan; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Sumatra).

Notes: (1) Schedl 1972f: 202 (to *Platyppus*).

References: (ds) Schedl 1972f: 202. (tx) Schedl 1935j: 2, 1972f: 202, 1978a: 71.

subsimilis Schedl 1936e: 54. Lectotype ♂; Mentawai, Si-Matobe and Sumatra: Siboga; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 71.

Distribution: Asia (Malaya), Indonesia (Mentawai, Sumatra).

Hosts: *Albizzia falcata*, *A. moluccana*.

References: (ds) Beeson 1941 (1961: 267); Browne 1961c: 195; Schedl 1961c: 70, 1972f: 192. (tx) Schedl 1936e: 54, 1939b: 394, 1939e: 336, 1972f: 192, 1978a: 71.

substriatus (Schedl) 1936j: 33 (*Crossotarsus*). Lectotype ♂; Malay Peninsula, Pahang; Cameron Highlands, 4500–4800 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 71.

- Distribution: Asia (Malaya), Indonesia (Sumatra).
 Hosts: *Cinnamomum camphora*.
 Notes: (1) Schedl 1972f: 202 (to *Platypus*).
 References: (hb) Kalshoven 1960c: 41. (ds) Kalshoven 1960c: 41; Schedl 1965g: 23 (as *substrictus*), 1972f: 201–202. (tx) Schedl 1936j: 33, 1972f: 201–202, 1978a: 71.
- subsulcatus** Chapuis 1865: 140. Syntypes 3 ♂; de Cayenne; IRSNB, Brussels (Schedl 1960c: 20).
 Distribution: South America (Brazil/ Cayenne/ Venezuela).
 References: (ds) Blackwelder 1947: 790; Dejean 1837; Gemminger & Harold 1872: 2700; Schedl 1960a: 78, 1972f: 241, 1972g: 51, 1976a: 58. (tx) Chapuis 1865: 140; Schedl 1933c: 167, 1935e: 176, 1936c: 226, 1952a: 444, 1960c: 20, 1972f: 241; Strohmeier 1912c: 9, 1914c: 24.
- subtruncatus** Browne 1980d: 498. Holotype ♂; Wallace Bay (Borneo) to Nagoya (Japan), imported; BMNH, London.
 Distribution: Indonesia (Borneo).
 Hosts: *Nauclea* sp., *Shorea* sp.
 References: (tx) Browne 1980d: 498.
- suffodiens** Sampson 1913: 447. Syntypes ♂ ♀; Lower Burma; BMNH, London.
 Distribution: Asia (Bangladesh/ Burma/ Bengal, Tamil Nadu in India/ Malaya), Indonesia (Borneo, Java), Philippine Islands.
 Hosts: *Adina* spp., *Anthocephalus cadamba*, *Fagraea gigantea*, *Hymenodictyon excelsum*, *Nauclea sessilifolia*, *Pithecolobium saman*, *Samanea saman*, *Vateria indica*.
 References: (cn) Corbett 1935: 43–56; Kleine 1932a: 310, 402; Mathur & Singh 1960b: 61, 1961a: 17, 1961b: 69; Roonwal 1954: 36; Stebbing 1914: 621. (ec) Beeson 1921b: 25; Stebbing 1914: 621. (hb) Browne 1961c: 197–198; Kalshoven 1960d: 35; Kleine 1932a: 310, 402; Stebbing 1914: 621. (ds) Beeson 1921b: 25, 1941 (1961: 267); Browne 1961c: 197–198, 1980a: 372, 1980c: 483, 1981b: 599; Kalshoven 1960d: 35; Kleine 1932a: 310, 402; Mathur & Singh 1960b: 61, 1961a: 17, 1961b: 69; Ohno, Yoneyama, & Nakazawa 1982a: 5, 1987a: 96, 1987b: 90; Roonwal 1954: 36; Sampson 1919: 107, 1925b: 392; Schedl 1962f: 165, 1972f: 193; Strohmeier 1914c: 27. (tx) Sampson 1913: 447–448, 1919: 107, 1925b: 392; Schedl 1939e: 336–360, 1941c: 354, 1942a: 209, 1972f: 193; Stebbing 1914: 621; Strohmeier 1914c: 27.
- sulcaticeps** Schedl 1979g: 118. Holotype ♂; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien.
 Distribution: New Guinea.
 References: (tx) Schedl 1979g: 118.
- sulcinodis** Schedl 1975f: 397. Holotype ♂; Mt. Kaindi, 2250 m. Morobe District; Schedl Collection in NHMW, Wien.
 Distribution: New Guinea.
- Hosts: *Castanopsis accuminatissima*, *Dysoxylum* sp.
 References: (tx) Schedl 1975f: 397.
- sulcipennis** Schedl 1976a: 91. Holotype ♂; Brasilien; Schedl Collection in NHMW, Wien.
 Distribution: South America (Brazil).
 References: (tx) Schedl 1976a: 91.
- suzukii** Browne 1981b: 605. Holotype ♂; Lawas (Sarawak, Borneo) to Tokyo (Japan), imported; BMNH, London.
 Distribution: Indonesia (Sarawak in Borneo).
 Hosts: *Shorea* sp.
 References: (tx) Browne 1981b: 605.
- taiheizanensis** (Murayama) 1932: 485 (*Crossotarsus*). Holotype ♂; Formosa: Taiheizan; Murayama Collection in USNM, Washington.
 Distribution: Asia (Taiwan).
 Hosts: *Castanopsis kawakamii*.
 Notes: (1) Schedl 1935b: 401 (to *Platypus*).
 References: (ds) Murayama 1937c: 580; Nobuchi 1967: 25; Schedl 1972f: 220. (tx) Murayama 1932: 485–486; Schedl 1935b: 401, 1939c: 4, 1941a: 44, 1972f: 220.
- takeharai** Browne 1983a: 563. Holotype ♂; Tg. Usan (West Irian) to Nagoya (Japan), imported; BMNH, London.
 Distribution: New Guinea.
 Hosts: "Matoa" log.
 References: (tx) Browne 1983a: 563.
- tasmanicus** Schedl 1942c: 192. Holotype ♂; Tasmanien; Schedl Collection in NHMW, Wien.
 Distribution: Australia (Tasmania).
 References: (ds) Schedl 1972f: 185. (tx) Schedl 1942c: 192, 1972f: 185, 1978a: 72.
- tayabasi** Schedl 1935k: 481. Holotype ♂; Luzon, Tayabas Province, Quezon Park; Schedl Collection in NHMW, Wien.
 Distribution: Asia (Andaman Islands in India/ Taiwan), Philippine Islands (Luzon, Mindanao).
 Hosts: Matoa log.
 References: (ds) Browne 1980d: 493; Ohno, Yoneyama, & Nakazawa 1987b: 90; Schedl 1966b: 101, 1972f: 202. (tx) Beeson 1937: 65; Nobuchi 1985: 329; Schedl 1935k: 481, 1972f: 202, 1972p: 159, 1978a: 72.
- lunatus** Browne 1964: 757. Holotype ♂; Philippines: Iligan, Mindanao; BMNH, London.
 Synonymy: Schedl 1972f: 202.
 References: (tx) Browne 1964: 757; Schedl 1972f: 202, 1972p: 159.
- lunatulus** Browne 1980d: 497. Holotype ♂; Keelung (Formosa) to Nagoya (Japan), imported; BMNH, London. Synonymy: Wood 1993:(in press).
 References: (ds) Browne 1980d: 497, 1986b: 333. (tx) Wood, S. L. 1993:(in press).

tenellus Schedl 1935k: 455. Syntypes ♂ ♀; Luzon, Mountain Province, Benguet, Mount Santo Tomas; W. Schultze Collection, and Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands (Luzon).

Hosts: *Ficus* sp., *Pinus insularis*.

Notes: (1) Schedl 1978a: 72 (citation of holotype invalid).

References: (ds) Schedl 1966b: 101, 1972f: 191. (tx) Nobuchi 1955: 329; Schedl 1935k: 454–456, 1942c: 198, 1972f: 191, 1978a: 72.

tenuis Murayama 1925b: 233. Holotype ♀; Sapporo, Hokkaido; Murayama Collection in USNM, Washington.

Figures: Nobuchi 1973b:pl. 2.

Distribution: Asia (Japan/ Malaya).

References: (ds) Murayama 1929a: 674, 1937c: 579, 1954b: 205; Schedl 1972f: 211. (tx) Murayama 1925b: 233, 1954b: 205; Nobuchi 1973b:pl. 2; Schedl 1934f: 1647, 1972f: 211.

tenuisculptus Roberts 1979: 95. Holotype ♂; Papua New Guinea: 14 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.

Figures: Roberts 1979: 97.

Distribution: New Guinea.

Hosts: *Ficus* sp.

References: (tx) Roberts 1979: 95–97.

tenuissimus Schedl 1941b: 426. Syntypes ♂ ♀; Java: Mt. Gede, 1000 m; Kalshoven Collection and Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java).

Hosts: *Castanea argentea*, *Eurya japonica*, *Nyssa javanica*.

Notes: (1) Schedl 1978a: 73 (citation of holotype invalid).

References: (hb) Kalshoven 1960c: 39. (ds) Kalshoven 1960d: 39; Schedl 1972f: 211. (tx) Schedl 1941b: 416–426, 1972f: 211, 1978a: 73.

tepidus Schedl 1942a: 210. Lectotype ♀; Lower Burma: Bhamo, Birmamia and Tenasserim, Thagata; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 73.

Distribution: Asia (Burma).

References: (ds) Schedl 1972f: 242. (tx) Browne 1972b: 29; Schedl 1942a: 210, 1972f: 242, 1978a: 73.

terebrans Schedl 1970b: 365. Holotype ♂; Sarawak, Lawas to Tokyo (Japan), imported; PPST, Tokyo.

Distribution: Indonesia (Borneo).

Hosts: Kapur log.

References: (ds) Nobuchi 1977: 120; Schedl 1972f: 194. (tx) Schedl 1970b: 368–369, 1972f: 194, 1978a: 73.

terminatulus Schedl 1971c: 396. Holotype ♂; Malaya, Kelantan; Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya).

References: (ds) Schedl 1972f: 196. (tx) Schedl 1971c: 396, 1972f: 196, 1978a: 73.

ternus Roberts 1986: 48. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Xanthophyllum papuanum*.

References: (tx) Roberts 1986: 48.

tetracerus Beeson 1929: 219. Holotype ♂; Samoa: Upolu, Malololelei, 2000 ft.; BMNH, London.

Distribution: Samoan Islands.

Hosts: *Plauchonella samoensis*.

References: (ds) Beaver 1976b: 545; Beeson 1938b: 296; Schedl 1972f: 211. (tx) Beeson 1929: 219–220; Schedl 1941b: 416, 1951k: 141, 155, 1972f: 211, 1978a: 73.

tiriosensis Reichardt 1965a: 53. Holotype ♂; Brasil, Estado do Para, Tirios (Alto rio Paru d'Oeste); Dep. Zool., Sec. Agric., Sao Paulo.

Distribution: South America (Brazil/ Cayenne/ Guyana/ Suriname).

Hosts: *Catostemma* sp., *Eperua falcata*, *Eschweilera sagottianum*.

References: (ds) Schedl 1972f: 229. (tx) Reichardt 1965a: 53; Schedl 1972f: 229; Wood, S. L. 1992a: 91.

schedli Wood 1966a: 51. Holotype ♂; Manaka, British Guiana; BMNH, London. Synonymy: Wood 1992a: 91.

References: (ds) Schedl 1972f: 229, 1972g: 51. (tx) Schedl 1972f: 229, 1978a: 64; Wood, S. L. 1966a: 51–52, 1992a: 91.

torulosus Roberts & Morimoto 1986: 179. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

References: (tx) Roberts & Morimoto 1986: 179.

tragus Schedl 1939k: 718. Holotype ♂; Brazil: Rio Negro; Schedl Collection in NHMW, Wien.

Figures: Schedl 1939k: 721 (male outline).

Distribution: South America (Brazil).

Hosts: *Hevea* sp.

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 234. (tx) Schedl 1939k: 718, 1972f: 234, 1978a: 74.

transformis Schedl 1936d: 16. Lectotype ♂; Malay Peninsula, Selangor: Kerling Res., Pahang: Temerloh; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 74.

Distribution: Asia (Malaya/ Sri Lanka), Indonesia (Borneo), Philippine Islands.

Hosts: *Balanocarpus heimii*, *Dryobalanops aromatica*, *Shorea* spp. in living trees.

Notes: (3) Schedl 1971c: 396 (described female).

References: (hb) Browne 1936a, 1941, 1961c: 202. (ds) Beeson 1961: 267; Browne 1936a: 120–127, 1961c: 202–203; Nobuchi 1977: 132; Ohno, Yoneyama, & Nakazawa 1987a: 96; Ohno, Yoshioka, et al. 1989: 67; Schedl 1959a: 514,

- 1959c: 168, 1971c: 371, 1972f: 207. (tx) Nobuchi 1985: 329; Schedl 1936d: 16, 1959a: 514, 1971c: 396–397, 1972f: 207, 1978a: 74.
- transindicus** (Beeson) 1937: 89 (*Crossotarsus*). Syntypes ♂ ♀; Tonkin: Hoabinh: FRI, Dehra Dum. Distribution: Asia (Tonkin Island in Vietnam). Notes: (1) Schedl 1972f: 206 (to *Platypus*). (3) Schedl 1972f: 206 (treated as a subspecies of *indicus*). References: (ds) Schedl 1972f: 206. (tx) Beeson 1937: 89; Schedl 1939c: 335, 1972f: 206.
- transversecarinatus** Schedl 1942c: 199. Holotype ♀; Australien, New Queensland; Schedl Collection in NHMW, Wien. Distribution: Australia (Queensland). References: (ds) Schedl 1972f: 242. (tx) Schedl 1942c: 199, 1972f: 242, 1978a: 74.
- transversus** Schedl 1979g: 119. Holotype ♀; Papua, Bulolo, Morobe District, Upper Manki L.A.; Schedl Collection in NHMW, Wien. Distribution: New Guinea. References: (tx) Schedl 1979g: 119.
- triangulatus** Browne 1984c: 454. Holotype ♂; Lever Harbour (Solomon Islands) to Nagoya (Japan), imported; BMNH, London. Distribution: Solomon Islands. Hosts: *Xanthophyllum* sp. References: (ds) Ohno, Yoshioka, et al. 1988b: 98. (tx) Browne 1984c: 454.
- tricuspidatus** Schedl 1954b: 48. Syntypes ♂; Brasilien; Schedl Collection in NHMW, Wien. Distribution: South America (Brazil). References: (tx) Schedl 1954b: 48, 1972f: 234, 1978a: 74.
- trilobipennis** Browne 1983a: 567. Holotype ♂; Sandakan (Sabah) to Nagoya (Japan), imported; BMNH, London. Distribution: Indonesia (Borneo). Hosts: "Erima" log. References: (tx) Browne 1983a: 567.
- triplurus** Browne 1980b: 388. Holotype ♂; Bangkok (Thailand) to Nagoya (Japan), imported; BMNH, London. Distribution: Asia (Thailand). Hosts: *Pterocarpus* sp. References: (tx) Browne 1980b: 388.
- trispinatus** Schedl 1970f: 584. Holotype ♂; Guyane française, Forêt d'Acarouany; Schedl Collection in NHMW, Wien. Distribution: South America (Cayenne). References: (ds) Schedl 1972f: 234. (tx) Schedl 1970f: 584, 1972f: 234, 1978a: 75.
- trispinatus** Schedl 1952a: 462. Syntypes ♂; Prov. Salta, Oran, Buenos Aires, Prov. Corrientes, Brazo Largo; Museo La Plata and Schedl Collection in NHMW, Wien. Distribution: South America (Argentina/ Brazil). Notes: (1) Schedl 1978a: 75 (citation of holotype invalid).
- References: (ds) Schedl 1970e: 87, 1972f: 234, 1979e: 58; Spahr 1981: 80. (tx) Schedl 1951h: 287, 1952a: 462, 1972f: 234, 1978a: 75.
- trispinosus** Chapuis 1865: 184. Holotype ♂; de Rio Janeiro, Bresil: BMNH, London (type not found by Schedl 1960c: 31). Distribution: South America (Argentina/ Brazil/ Venezuela). References: (cn) Santoro 1966a. (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2700; Santoro 1957b: 25; Schedl 1933c: 172, 1965b: 87, 1966f: 94, 1972: 234, 1979e: 59; Strohmeier 1912c: 13, 1914c: 25. (tx) Chapuis 1865: 184; Schedl 1936i: 101, 1951h: 287, 1960c: 31, 1972f: 234, 1978a: 75; Strohmeier 1912c: 13, 1914c: 25.
- quadricuspidatus** Schedl 1948f: 283. Syntypes ♂ ♀; Brasil, Nova Teutonia; Plammann Collection and Schedl Collection in NHMW, Wien. Synonymy: Schedl 1972f: 234. Notes: (1) Schedl 1978a: 60 (citation of holotype invalid). References: (tx) de Ruette 1970: 68; Schedl 1948f: 283, 1972f: 234, 1978a: 60.
- tristis** Roberts 1986: 49. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London. Distribution: New Guinea. Hosts: *Protium* sp. References: (tx) Roberts 1986: 49.
- truncatellus** Browne 1975: 309 in Beaver & Browne. Holotype ♂; Thailand: Nakhon Ratchasima, Khao Yai; BMNH, London. Distribution: Asia (Thailand). Hosts: Tree. References: (tx) Beaver & Browne 1975: 309.
- truncaticauda** Schedl 1968f: 536. Holotype ♂; New Guinea (NW), Bodem, 100 m, 11 km SE of Oerberfaren; apparently in BPBM, Honolulu. Distribution: New Guinea. References: (ds) Schedl 1972f: 212. (tx) Schedl 1968f: 536, 1972f: 212, 1978a: 75.
- truncatigranosus** Schedl 1970a: 132. Holotype ♂; New Guinea, Porotop Lutheran Mission Sawmill, Western Highlands District; CSIRO, Canberra. Distribution: New Guinea. References: (ds) Schedl 1968f: 535, 1972f: 212. (tx) Schedl 1970a: 132, 1972f: 212, 1975f: 398, 1978a: 75.
- truncatipennis** Schedl 1964c: 302. Holotype ♂; New Guinea, Humboldt Bai; Schedl Collection in NHMW, Wien. Distribution: New Guinea. References: (ds) Schedl 1968f: 537, 1972f: 212. (tx) Schedl 1964c: 302, 1972f: 212, 1978a: 76.
- diversiporus** Schedl 1972i: 56. Holotype ♀; Papua, Western Dist., Oriomo Gov. Sta.; BPBM, Honolulu. Synonymy: Roberts 1989: 285.

References: (ds) Schedl 1972f: 217. (tx) Schedl 1972f: 217, 1972i: 56, 1978a: 26.

multifoveolatus Schedl 1975f: 359. Holotype ♀; New Guinea, Madang District, Nr. 1, Ramp, Gogol.; Schedl Collection in NHMW, Wien. Synonymy: Roberts 1989: 285. References: (tx) Roberts 1989: 285; Schedl 1975f: 359.

truncatulus Browne in Beaver & Browne 1978: 621. Holotype ♂; Malaya, Penang, Muka Head; BMNH, London. Distribution: Asia (Malaya). Hosts: *Calophyllum tetrapterum*. References: (tx) Beaver & Browne 1978: 621.

truncaturus Browne 1983a: 562. Holotype ♂; Tatan (Sarawak) to Nagoya (Japan), imported; BMNH, London. Distribution: Indonesia (Sarawak in Borneo). References: (tx) Browne 1983a: 562.

truncatus Chapuis 1865: 269. Syntypes 2 ♂; de l'île de France (Mauritius); 1 in BMNH, London. 1 in IRSNB, Brussels. Distribution: Madagascar, Mauritius Island.

References: (cn) Montia 1932: 9–12. (ds) Alluaud 1900: 441; Gemminger & Harold 1872: 2700; Schedl 1972f: 234; Strohmeier 1912c: 17, 1914c: 27. (tx) Chapuis 1865: 269; Lucas 1920: 522; Schedl 1939b: 399, 1960c: 51, 1972f: 234; Strohmeier 1912c: 17, 1914c: 27. (ms) Lucas 1920: 522.

tuberculatus Chapuis 1865: 131. Holotype ♂; de Cayenne; IRSNB, Brussels (Schedl 1960c: 17). Figures: Mayne & Donis 1962: 259. Distribution: South America (Cayenne). Notes: (3) Schedl 1936c: 226 (described female). References: (ds) Blackwelder 1947: 791; Gemminger & Harold 1872: 2700; Schedl 1970f: 582, 1972f: 241, 1972g: 51; Strohmeier 1912c: 9, 1914c: 24. (tx) Chapuis 1865: 131; Mayne & Donis 1962: 259; Schedl 1933c: 167, 1935e: 176, 1936c: 226, 1960c: 17, 1972f: 241, 1978a: 76; Strohmeier 1912c: 9, 1914c: 24.

tuberculosis Strohmeier 1910e: 128. Holotype ♂; Tasmania; Strohmeier Collection. Distribution: Australia (Tasmania). References: (ds) Schedl 1972f: 199; Strohmeier 1912c: 19, 1914c: 29. (tx) Schedl 1936c: 226, 1972f: 199; Strohmeier 1910e: 128, 1912c: 19, 1914c: 29.

tuberosus (Schedl) 1942a: 214. Holotype ♂; Malaya, Pahang, Cameron's Highlands, 4800 Fuss; Schedl Collection in NHMW, Wien. Distribution: Asia (Malaya). References: (ds) Browne 1961c: 213; Schedl 1972f: 186. (tx) Browne 1962e: 653; Schedl 1942a: 214, 1972f: 186, 1978a: 76.

turcius Schedl 1970b: 366. Holotype ♂; New Guinea, Vanimo to Nagoya (Japan), imported; PPST, Tokyo.

Distribution: New Guinea.

References: (ds) Nobuchi 1977: 131; Schedl 1972f: 201. (tx) Schedl 1970b: 366–367, 1972f: 201, 1978a: 76.

turriiformis Roberts & Morimoto 1986: 180. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.

Distribution: New Guinea.

Hosts: *Xanthophyllum papuanum*.

References: (tx) Roberts & Morimoto 1986: 180.

ugandae Browne 1965a: 205. Holotype ♂; Uganda: Zika; BMNH, London.

Distribution: Africa (Uganda).

References: (ds) Schedl 1972f: 235. (tx) Browne 1965a: 205; Schedl 1972f: 235.

umbonatus Blandford 1896e: 97. Holotype ♂; Guatemala, Cerro Zunil; BMNH, London.

Distribution: North America (Guatemala).

References: (ds) Blackwelder 1947: 791; Equihua & Atkinson 1957: 30; Schedl 1972f: 241; Strohmeier 1912c: 9, 1914c: 24. (tx) Blandford 1896e: 97–98; Equihua & Atkinson 1957: 30; Schedl 1972f: 241; Strohmeier 1912c: 9, 1914c: 24.

umbraticus Schedl 1964c: 303. Holotype ♂; Java: Mt. Ardjoena; ZMA, Amsterdam.

Distribution: Indonesia (Java).

References: (ds) Schedl 1972f: 211. (tx) Schedl 1964c: 303–304, 1972f: 211.

umbrinus Roberts 1989: 283. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.

Figures: Roberts 1989: 288.

Distribution: New Guinea.

Hosts: *Syzygium* sp.

References: (tx) Roberts 1989: 283, 288.

uncatus Browne 1980c: 498. Holotype ♂; Kennedy Bay (Borneo) to Nagoya (Japan), imported; BMNH, London.

Distribution: Indonesia (Borneo).

Hosts: *Nauclea* sp.

References: (tx) Browne 1980c: 489.

uncinatus Blandford 1895a: 327. Syntypes (several) ♂ ♀; Ceylon, Dikoya, Bogawantalawa; BMNH, London.

Distribution: Asia (Burma/ Assam, Bengal, Karnataka, Tamil Nadu in India/ Malaya/ Sri Lanka). Hosts: *Albizia* spp., *Anogeissus latifolia*, *Bassia latifolia*, *Bombax malabaricum*, *Boswellia serrata*, *Bridelia retusa*, *Buchanania latifolia*, *Camellia sinensis*, *Castanopsis tribuloides*, *Cedrela toona*, *Cochlospermum gossypium*, *Diospyros paniculata*, *Dipterocarpus pilosus*, *Doonia zeylanica*, *Eugenia jambolana*, *Ficus glomerata*, *Gmelina arborea*, *Heritiera fomes*, *Salmalia malabarica*, *Shorea* spp., *Terminalia* spp., *Vatica lanceaefolia*.

Notes: (3) Beeson 1941: 265 (*forfex*, nomen nudum, *kanarensis*, nomen nudum, synonymy in Schedl 1960c: 63).

References: (**cn**) Kleine 1932a: 310, 402; Mathur & Singh 1960a: 7, 1960b: 17, 1961a: 13, 1961b: 26; Roonwal 1954: 45; Sen-Sarma & Thakur 1986: 37. (**ee**) Beeson 1921b; Thompson, W. R. 1943: 91. (**hb**) Kleine 1932a: 310, 402; Speyer 1923: 23. (**ds**) Beeson 1919: 1–23, 1921b: 25, 1961: 267; Bhasin, Roonwal, & Singh 1958: 17; Browne 1981b: 599; Kleine 1932a: 310, 402; Mathew 1982, 1987: 188; Mathur & Singh 1960a: 7, 1960b: 17, 1961a: 13, 1961b: 26; Ohno, Yoneyama, & Nakazawa 1982b: 11, 1987b: 90; Roonwal 1954: 45; Schedl 1959a: 515, 1969c: 63, 1971a: 281, 1971c: 362; Waterston 1922: 51–94. (**tx**) Beeson 1941: 265; Blandford 1895a: 327; Schedl 1941b: 416, 1959a: 51, 1960c: 63, 1972f: 211.

rectangulatus Sampson 1913: 448. Syntypes ♂ ♀; Tharrawaddy, Lower Burma; BMNH, London. Synonymy: Schedl 1972f: 221.

References: (**cn**) Kleine 1932a: 310; Mathur & Singh 1961b: 41; Stebbing 1914: 623. (**ee**) Stebbing 1914: 623. (**hb**) Kleine 1932a: 310; Stebbing 1914: 623. (**ds**) Beeson 1921: 24, 1941 (1961: 266); Kleine 1932a: 310; Mathur & Singh 1961b: 41; Schedl 1961c: 167; Strohmeier 1914c: 28. (**tx**) Mathur & Singh 1961b: 41; Sampson 1913: 448–449; Schedl 1941b: 416, 1972f: 211; Stebbing 1914: 623; Strohmeier 1914c: 28.

uniformis Schedl 1970: 234. Holotype ♂ ♀; New Guinea: Porotop L.M. Station, W.H.D.; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Cryptocarya* sp., *Podocarpus* sp., *Syzygium* sp.

References: (**ds**) Schedl 1972f: 185. (**tx**) Schedl 1969b: 234–235, 1972f: 185, 1978a: 77.

ursinus Schedl 1935h: 353. Syntypes ♂ ♀; Brazilien, S. Paulo; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica), South America (Brazil/ Cayenne).

References: (**ds**) Blackwelder 1947: 791; Schedl 1970f: 582, 1972f: 242. (**tx**) Schedl 1935h: 353–354, 1972f: 242, 1978a: 78.

ursus Schedl 1933c: 166. Holotype ♂; Costa Rica, Hamburgfarn, Reventazon, Ebene Limon, 200–300 m; Schedl Collection in NHMW, Wien.

Distribution: North America (Costa Rica).

Notes: (3) Schedl 1934e: 209 (described female).

References: (**ds**) Blackwelder 1947: 791; Equihua & Atkinson 1987: 30; Schedl 1933c: 166, 1972f: 242. (**tx**) Equihua & Atkinson 1987: 30; Schedl 1933c: 166, 1934e: 209, 1972f: 242, 1978a: 78.

ustulatus Chapuis 1865: 224. Syntypes ♂ ♀; du Mexique, Toxpan; 2 ♂, 1 ♀ in BMNH, London, 2 ♂, 1 ♀ in IRSNB, Brussels.

Figures: Bright 1972d: 19.

Distribution: Antilles Islands (Jamaica), North America (Costa Rica/ Guatemala/ Veracruz in

Mexico), South America (Brazil/ Cayenne/ Colombia).

References: (**ds**) Blackwelder 1947: 791; Blandford 1896e: 110; Bright 1972d: 16; Equihua & Atkinson 1987: 30; Gemminger & Harold 1872: 2700; Nunberg 1958a: 482, 1972b: 190; Schedl 1966f: 94, 1967d: 5, 1972f: 218, 1972g: 38, 1976a: 59; Strohmeier 1912c: 14, 1914c: 26; Sturm 1843: 231. (**tx**) Blandford 1896e: 110; Bright 1972d: 16, 19; Chapuis 1865: 224; Equihua & Atkinson 1987: 30; Nunberg 1958a: 482; Schedl 1939b: 399, 1952c: 123, 1960c: 41, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.

petersi Chapuis 1865: 226. Syntypes ♂ ♀; du Bresil; 1 ♂, 1 ♀ in MNB, Berlin, 1 ♂ in IRSNB, Brussels. Synonymy: Schedl 1972f: 219.

References: (**ds**) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Strohmeier 1912c: 14, 1914c: 26. (**tx**) Chapuis 1865: 226; Schedl 1937d: 33, 40, 1952a: 463, 1960c: 42, 1972f: 219; Strohmeier 1912c: 14, 1914c: 26.

obsoletus Chapuis 1865: 228. Syntypes ♂ ♀; de la Colombie; 1 ♂ ex typis in IRSNB, Brussels (Schedl 1960c: 42). Synonymy: Schedl 1972f: 218.

References: (**ds**) Blackwelder 1947: 790; Gemminger & Harold 1872: 2699; Nunberg 1958a: 481; Strohmeier 1912c: 14, 1914c: 26. (**tx**) Chapuis 1865: 228; Nunberg 1958a: 481–482; Schedl 1936c: 242, 1960c: 41–42, 1972f: 218; Strohmeier 1912c: 14, 1914c: 26.

ustus Schedl 1968e: 268. Holotype ♂; Okapa (Okasa), E. Highlands District; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Heritica trifolia*, *Pterocymbium beccarii*.

Notes: (3) Schedl 1969b: 235 (described female).

References: (**ds**) Schedl 1972f: 201, 1979a: 160. (**tx**) Schedl 1968e: 268, 1969b: 235, 1972f: 201, 1978a: 78; Wylie & Yule 1977.

uter Schedl 1968b: 236. Holotype ♂; New Guinea: Karamui, Chimbu District; CSIRO, Canberra.

Distribution: New Guinea.

Notes: (3) Schedl 1975f: 398 (described female).

References: (**ds**) Schedl 1972f: 191. (**tx**) Schedl 1968b: 236, 1972f: 191, 1975f: 398, 1978a: 78.

utibilis (Schedl) 1939e: 360 (*Crossotarsus*). Lectotype ♂; Malaya, Perak: Trolak For. Res. and Pahang: Kemasil For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 78.

Distribution: Asia (Malaya).

Hosts: *Aporosa* (?) sp.

References: (**hb**) Browne 1941. (**ds**) Browne 1961c: 210, 1980c: 483; Schedl 1972f: 202. (**tx**) Schedl 1939e: 360, 1972f: 202, 1978a: 78.

uvarius Roberts 1989: 284. Holotype ♂; Papua New Guinea: Gumi, Watut Logging Area, Bulolo, 2200 m; NHMW, Wien.

Figures: Roberts 1989: 286.

- Distribution: New Guinea.
References: (tx) Roberts 1989: 284, 286.
- vafae Schedl** 1972g: 51. Holotype ♂; Brasilien, Utinga bei Belem, Para; Schedl Collection in NHMW, Wien.
Distribution: South America (Brazil).
References: (ds) Schedl 1976a: 59. (tx) Schedl 1972g: 51, 1978a: 78.
- vaginerris Roberts** 1986: 51. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (tx) Roberts 1986: 51.
- vagus Browne** 1983a: 568. Holotype ♂; Bintulu (Sarawak) to Nagoya (Japan), imported; BMNH, London.
Distribution: Indonesia (Sarawak in Borneo).
Hosts: Pernarahan log.
References: (tx) Browne 1983a: 568.
- valdephlebius Roberts** 1979: 96. Holotype ♂; Papua New Guinea: 12 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.
Figures: Roberts 1979: 93 (male, head of female).
Distribution: New Guinea.
Hosts: *Schizomeria* sp.
References: (tx) Roberts 1979: 93.
- validus Schedl** 1973e: 97. Holotype ♂; New Guinea: Taun logging area, Bulolo, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
References: (tx) Schedl 1973e: 97, 1978a: 78.
- varipennis Schedl** 1968e: 269. Holotype ♂; Afafiningetu, E. Highlands District; CSIRO, Canberra.
Distribution: New Guinea.
Hosts: *Araucaria cunninghamii*.
References: (tx) Schedl 1968e: 269, 1972f: 201, 1978a: 78.
- vegestus Wood** 1966a: 63. Holotype ♂; Pan-American Highway 15 miles NW Cerro de la Muerte, San Jose Prov., Costa Rica, 7500 feet; Wood Collection.
Figures: Schedl 1972f: 213, Wood 1966a: 67.
Distribution: North America (Costa Rica).
Hosts: *Bruellia costaricensis*.
References: (ds) Equihua & Atkinson 1987: 30; Schedl 1972f: 214. (tx) Equihua & Atkinson 1987: 30; Schedl 1972f: 213–214; Wood, S. L. 1966a: 63, 67.
- verelunatus (Beeson)** 1937: 65 (*Crossotarsus*). Holotype ♂; Middle Andaman; FRI, Dehra Dun. Distribution: Asia (Andaman Islands in India/Vietnam).
Hosts: *Artocarpus chaplasha*, *Bassia butyracea*, *Pterocarpus dalbergioides*, *Sterculia* spp., *Terminalia manii*.
References: (cn) Mathur & Singh 1961a: 69, 1961b: 32; Roonwal 1954: 85. (ds) Beeson 1941 (1961: 257); Mathur & Singh 1961a: 69, 1961b: 32; Roonwal 1954: 85; Schedl 1972f: 202, 1974c: 262. (tx) Beeson 1937: 65–66; Schedl 1939e: 360, 1972f: 202, 1978b: 80.
- vernaculus Schedl** 1965g: 29. Holotype ♂; O. Borneo, Mt. Tibang, 1300 und 1400 m; NHR, Stockholm.
Distribution: Indonesia (Borneo).
References: (ds) Schedl 1972f: 201. (tx) Schedl 1965g: 29, 1972f: 201, 1978a: 80.
- vesculus Schedl** 1972i: 58. Holotype ♂; New Guinea (NE), Karimui, 1000 m; BPBM, Honolulu. Distribution: New Guinea.
References: (ds) Schedl 1972f: 191. (tx) Schedl 1972f: 191, 1972i: 58, 1978a: 80.
- vethi Strohmeier** 1913: 161. Syntypes ♂ ♀; Preanger (Java); Strohmeier Collection.
Distribution: Asia (Malaya), Indonesia (Java, Sumatra).
Hosts: *Cinnamomum camphora*, *Dalbergia latifolia*, *Dyera costulata*, *Podocarpus imbricata*, *Schinus cordatus*.
Notes: (3) Schedl 1972f: 193 *suffodiens* var. *major* Sampson, nomen nudum, synonymy).
References: (cn) Browne 1968b: 572; Kalshoven 1960d: 35. (hb) Browne 1961c: 196, 1968b: 572; Kalshoven 1960c: 34. (ds) Browne 1961c: 196, 1968b: 572; Kalshoven 1960d: 35; Strohmeier 1914c: 27. (tx) Browne 1961c: 196, 1968b: 572; Kalshoven 1960d: 63; Schedl 1972f: 193; Strohmeier 1913: 161–162, 1914c: 27.
- suffodiens malayensis** Schedl 1936: 34. Holotype ♂; Malaya (Bukit Cherakash Forest Reserve and Pahang, Memasul Forest Reserve; not given, but added as NHMW, Wien, in Schedl 1978a: 71. Synonymy: Kalshoven 1960: 63.
References: (bv) Bletchly 1961: 15. (cn) Browne & Foenander 1937: 240–254. (ce) Bletchly 1961: 15; Browne 1958b. (hb) Browne 1935a, 1958b. (ds) Beeson 1961: 265; Bletchly 1961: 15; Browne 1935a; Browne & Foenander 1937: 240–254; Schedl 1936j: 34. (tx) Browne 1949b: 910; Kalshoven 1960d: 63; Schedl 1936j: 34, 1939e: 336, 360, 1972f: 193, 1978a: 71.
- spretus Schedl** 1939e: 360. Holotype ♂; Malaya (Bukit Cherakash Forest Reserve and Pahang, Memasul Forest Reserve; Schedl Collection in NHMW, Wien, automatic. Synonymy: Kalshoven 1960d: 63.
References: (hb) Browne 1941. (tx) Browne 1949b: 910; Kalshoven 1960d: 63; Schedl 1939e: 336, 360, 1941c: 354, 1942a: 209.
- vetulus Schedl** 1935n: 638. Lectotype ♂; Malay States: Johore North, Klang; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 80.
Distribution: Asia (Malaya), Indonesia (Borneo, Mentawai, Sumatra).
Hosts: *Dryobalanops aromatica*.

- Notes: (3) Schedl 1936d: 14 (described female).
References: **(hb)** Browne 1941. **(ds)** Beeson 1941 (1961: 267); Browne 1961c: 196, 1981b: 599; Schedl 1965g: 24, 1971c: 370, 1971f: 151, 1972f: 193. **(tx)** Schedl 1935n: 638, 1936d: 14, 1936e: 51, 1939e: 336, 1954c: 155, 1972f: 193, 1978a: 80.
- vetustus** Schedl 1941d: 414. Holotype ♂; Belge Congo: Likimi, Gumba; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: **(ds)** Schedl 1972f: 223. **(tx)** Schedl 1941d: 414, 1957d: 120, 1962k: 637, 1972f: 223, 1978a: 80.
- vexans** Schedl 1972g: 82. Holotype ♂; Brasilien, Cachimbo, Para; MZUSP, Sao Paulo.
Distribution: South America (Brazil).
References: **(tx)** Schedl 1972g: 82, 1978a: 80.
- viticus** Schedl 1974d: 470. Holotype ♀; New Guinea: Taun logging area, Bulolo, Morobe District; CSIRO, Canberra.
Distribution: New Guinea.
References: **(tx)** Schedl 1974d: 470, 1978a: 80.
- vicarius** Schedl 1975g: 232. Holotype ♀; New Guinea, Wan; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975g: 232, 1978a: 81.
- vicinus** Blandford 1896e: 104. Holotype ♂; Mexico; BMNH, London.
Figures: Blandford 1896e: pl. 4, fig. 14.
Distribution: North America (Mexico).
References: **(ds)** Blackwelder 1947: 791; Equihua & Atkinson 1987: 30; Ferrer 1942; Schedl 1972f: 206; Strohmeier 1912c: 10, 1914c: 24. **(tx)** Blandford 1896e: 104; Equihua & Atkinson 1987: 30; Schedl 1940a: 324, 1972f: 206; Strohmeier 1912c: 10, 1914c: 24.
- vitiensis** Roberts 1977b: 571. Holotype ♂; Fiji Islands: Viti Levu, 10 km N Ngaloa Village, 300 m; BMNH, London.
Distribution: Fiji Islands.
Hosts: *Agathis vitiensis*, *Burckella* sp., *Calophyllum vitiense*, *Canarium vitiense*, *Cerbera odollum*, *Cynometra insularis*, *Dacrydium elatum*, *Dysoxylum* sp., *Endospermum macrophyllum*, *Garcinia myrtifolia*, *Gonostylus punctatus*, *Maniltoa* sp., *Myristica castaneifolia*, *Palaquium hornei*, *Parinari insularum*.
References: **(ce)** Roberts 1977b: 571. **(hb)** Roberts 1977b: 571. **(ds)** Roberts 1977b: 571. **(tx)** Roberts 1977b: 571.
- voluticus** Schedl 1936j: 35. Lectotype ♂; Malay Peninsula, Pahang; Trantum, Tersang Forest Reserve, and Perak; Chenderoh; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 29.
Distribution: Asia (Malaya), Bougainville, Indonesia (Java), New Britain, New Guinea, New Ireland.
Hosts: *Diospyros* sp., *Dysoxylum* sp., *Micromcles* sp.
- Notes: (1) Schedl 1941c: 365 (raised to species rank from subspecies of *excedens*). (3) Schedl 1941c: 365 (described female).
References: **(hb)** Browne 1961c: 213; Kalshoven 1960d: 41. **(ds)** Browne 1961c: 213; Kalshoven 1960d: 41; Schedl 1936j: 35, 1968f: 537, 1972f: 186. **(tx)** Browne 1962e: 653; Schedl 1936j: 35, 1941c: 365–366, 1972f: 186, 1978a: 29, 81.
- vonfaberi** Reichardt 1962: 336. Holotype ♂; Brasil, Estado de Matto Grosso, Xingu (lat. 12 00, long. 53 22); DZSA, Sao Paulo.
Distribution: South America (Brazil).
References: **(ds)** Schedl 1972f: 187. **(tx)** Reichardt 1962: 336, 1964: 147; Schedl 1972f: 187.
- uatruui** Roberts 1965: 219. Holotype ♂; Nigeria: Idanre; BMNH, London.
Distribution: Africa (Nigeria/ Sierra Leone).
Hosts: *Xylopia* sp.
References: **(ds)** Roberts 1968: 189, 1969: 120; Schedl 1972f: 228. **(tx)** Roberts 1965: 219; Schedl 1972f: 228.
- uatutensis** Roberts 1986: 53. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: **(tx)** Roberts 1986: 53.
- webberi** Schedl 1939e: 362. Syntypes ♂; Malaya, Selangor: The Gap, 2700 feet; BMNH, London.
Distribution: Asia (Malaya), Indonesia (Sumatra).
Hosts: *Myristica fragrans*.
References: **(hb)** Browne 1941; Kalshoven 1960c: 34. **(ds)** Browne 1961c: 196; Kalshoven 1960d: 34; Schedl 1972f: 194. **(tx)** Schedl 1939e: 362–363, 1941c: 354, 1958b: 102, 1972f: 194, 1978a: 81.
- westwoodi** Chapuis 1865: 236. Syntypes 2 ♂; Ile Borneo, Sarawak; BMNH, London (Schedl 1960c: 44).
Distribution: Asia (Malaya/ Thailand), Indonesia (Borneo, Java, Mentawai, Sumatra), Si-Ohan Island.
Hosts: *Azadirachta bakeri*, *A. palembanica*, *Artocarpus lakoocha*, *Calophyllum* sp., *Cratogeomys arborescens*, *Dyera costulata*, *Endospermum malaccense*, *Eugenia* sp., *Ficus* sp., *Gonostylus bancanus*, *Hevea brasiliensis*, *Koompassia malaccensis*, *Lucuma malaccensis*, *Palaquium* sp., *Pouteria malaccensis*, *Sapium baccatum*, *Scaphium* sp., *Xylopia cordata*.
Notes: (3) Schedl 1936d: 14 (described female).
References: **(cn)** Ishikura 1966; Yunus & Hua 1980: 217. **(cc)** Browne 1958b. **(hb)** Browne 1935a, 1941, 1958b, 1961c: 194; Kalshoven 1960c: 34; Mercer 1952: 210. **(ds)** Beaver & Browne 1975: 309, 1978: 621; Beeson 1941 (1961: 267); Browne 1935a, 1961c: 194–195, 1972b: 29, 1980a: 371; Gemminger & Harold 1872: 2700; Ishikura 1966; Kalshoven 1924b: 358, 1960d: 34; Mercer 1952: 210; Ohno, Yoneyama, & Nakazawa 1952b: 11; Sampson 1919: 106.

- 1925b: 392; Schedl 1936d: 7, 1936j: 22, 1959c: 165, 1971f: 151, 1972f: 192; Strohmeier 1912c: 16, 1914c: 27; Yimms & Hua 1980: 217. **(tx)** Chapuis 1965: 236; Sampson 1913: 448, 1919: 106, 1928b: 392; Schedl 1936d: 14–15, 1937e: 544, 1939e: 336, 1941c: 354, 1972f: 192, 1978a: 51; Strohmeier 1912c: 16, 1914c: 27.
- wilsoni Swaine** 1916a: 98. Holotype ♀; Campbell River, British Colombia [Canada]; CNCI, Ottawa. Figures: Arnett 1960: 1029, Swaine 1916a: pls. 6–7. Distribution: North America (British Columbia in Canada/ California, Idaho, Oregon, Washington in USA). Hosts: *Abies magnifica*, *Pseudotsuga menziesii*, *Tsuga heterophylla*. References: **(ay)** Chapman 1961b: 3; Farris 1969: 529; Farris & Funk 1965: 527–537; Livingston & Berryman 1972: 1799. **(bv)** Prebble & Graham 1957: 91–92; Shore & McLean 1983b. **(cn)** Chamberlin 1917: 356, 1924, 1939: 112; Doane et al. 1936; Fisher, Thompson, & Webb 1953, 1954b: 3–15; Furniss, R. L. 1937: 30; Graham 1939b: Hatch 1938: 192; Johnson 1958a: 508, 1958b: 236–240; Keen 1936: 145; Kowal 1949a: 12–13; Mathers 1935: 14; McMullan 1956: 34; Nijholt 1978a: Prebble 1944: 50; Ruppel 1967: 79; Schuder 1969: 74; Shore 1985; Silver & Ross 1960: 99, 1961: 99; Struble 1937d: 11, 1957: 14; Swaine 1914: 38, 1918a: 38; Thatcher, T. O. 1961. **(cc)** Chamberlin 1918a: 35, 1939: 112; Farris 1969: 529; Farris & Funk 1965: 929–932; Furniss, R. L. & Carolin 1977: 338; Graham 1939b; Keen 1938: 145; Roeper & French 1981; Schedl 1958d: 188; Shore 1985; Struble 1957: 14. **(hb)** Bright & Stark 1973: 10; Chamberlin 1939: 112, 1958: 33–34; DeLeon 1952; Doane et al. 1936; Ferrell 1969: 2238; Furniss, R. L. & Carolin 1977: 338; Graham 1939b; Husson 1955: 350; Johnson 1958a: 508, 1958b: 237; Keen 1938: 145; Kinney & Furniss 1943: 26–27; Nijholt 1978a; Prebble & Graham 1957: 91–92; Shore 1985; Struble 1937d: 11, 1957: 14; Swaine 1918a: 38; Thatcher, T. O. 1961; White, R. E. 1983: 326. **(ds)** Blackwelder & Blackwelder 1948; Chamberlin 1918a: 35, 1925, 1939: 112, 1958: 33–34; Currie 1905; DeLeon 1952: 78; Equihua & Atkinson 1987: 30; Furniss, R. L. & Carolin 1977: 338; Hopping 1922; Keen 1929a: 11, 1938: 145; Leng 1920: 337; Murayama 1957a: 37; Patterson & Hatch 1945: 148; Ruppel 1967: 79; Schedl 1933c: 172, 1972f: 199; Schuder 1969: 74; Shore 1985; Swaine 1914: 38; Thatcher, T. O. 1961; Wood, S. L. 1958b: 39, 1972b: 394. **(tx)** Arnett 1960: 1029, 1968: 1029; Chamberlin 1939: 112, 1958: 33–34; Equihua & Atkinson 1987: 30; Hoebeke 1978; Johnson 1958b: 237; Keen 1929a: 11; de Rente 1970: 68; Schedl 1972f: 199; Swaine 1916a: 97–100, 1918a: 38; Wood, S. L. 1958b: 39, 1972b: 394, 1979a: 2. **(ms)** Hatch 1938: 192; Prebble 1944: 50.
- xanthopus Schedl** 1957d: 124. Holotype ♀; Congo Belge; Yangambi; MRCB, Tervuren. Figures: Schedl 1962k: 673, 1972f: 228. Distribution: Africa (Zaire). Hosts: *Xylopiia* sp. Notes: (3) Schedl 1962k: 1108 (described female). References: **(ds)** Mayne & Donis 1962: 228; Schedl 1962k: 673, 1972f: 228. **(tx)** Schedl 1957d: 124, 1962k: 673, 1108, 1972f: 228, 1978a: 81.
- xylographus Schedl** 1969a: 216. Holotype ♂; Formosa, Kaohsiung to Tokyo (Japan), imported; PPST, Tokyo. Distribution: Asia (Taiwan). Hosts: *Picea* sp., *Pinus* sp. References: **(ds)** Nobuchi 1977: 128; Schedl 1972f: 199. **(tx)** Schedl 1969a: 216, 1972f: 199, 1978a: 82.
- yasiyasi Roberts** 1977b: 560. Holotype ♂; Fiji Islands: Viti Levu, 12 km W Namulolo Village, 200 m; BMNH, London. Figures: Roberts 1977b: 561. Distribution: Fiji Islands. Hosts: *Cleistocalyx* sp., *Syzygium effusum*. References: **(bv)** Roberts 1977b: 560. **(cn)** Essig 1958: 509. **(hb)** Essig 1958: 509; Roberts 1977b: 560. **(ds)** Essig 1958: 509. **(tx)** Roberts 1977b: 560–561.

Genus *Treptoplatypus* Schedl

- TREPTOPLATYPUS SCHEDL** 1939b: 401. Type-species: *Crossotarsus trepanatus* Chapuis, subsequent designation by Wood 1992a: 91. Notes: (2) Browne 1962e: 653 (treated as a synonym of *Platypus*). References: **(tx)** Browne 1962e: 653; Schedl 1939b: 401, 1968e: 261–270, 1972f: 243–245; Wood, S. L. 1992a: 91.
- circulicauda Browne** 1949b: 911. Holotype ♂; Malaya: Kelantan, Bukit Besi; BMNH, London. Distribution: Asia (Malaya), Indonesia (Borneo). Hosts: *Diospyros* sp., *Dipterocarpus* sp. References: **(ds)** Browne 1961c: 209, 1962c: 202, 1981b: 598; Schedl 1972f: 245. **(tx)** Browne 1949b: 911; Schedl 1972f: 245.
- fischeri (Strohmeier)** 1907g: 29 (*Crossotarsus*). Holotype ♀; Sumatra (Regentschaft Palembang); Strohmeier Collection. Distribution: Indonesia (Sumatra). Notes: (1) Schedl 1972f: 245 (to *Treptoplatypus*). References: **(ds)** Schedl 1972f: 245; Strohmeier 1912c: 5, 1914c: 35. **(tx)** Schedl 1972f: 245; Strohmeier 1907g: 29–30, 1912c: 5, 1914c: 35.
- multioporus Schedl** 1968e: 270. Holotype ♀; Okapa (Okasa), E. Highlands District; CSIRO, Canberra. Distribution: New Guinea. Hosts: *Heritiera trifolia*, *Podocarpus* sp.. Notes: (1) Schedl 1978a: 46 (type restricted to CSIRO specimens).

References: **(ds)** Schedl 1972f: 245. **(tx)** Schedl 1968e: 270, 1972f: 245, 1978a: 45.

fastuosus Schedl 1969b: 226 (*Platypus*). Holotype ♂; New Guinea: Maralinga, 2500 m; CSIRO, Canberra. Synonymy: Wood 1992a: 92.

References: **(tx)** Schedl 1969b: 226; Wood, S. L. 1992a: 92.

quadriporus Schedl 1971c: 397. Holotype ♀; China: Fukien: Schedl Collection in NHMW, Wien. Distribution: Asia (Fujian in China).

References: **(ds)** Schedl 1972f: 245. **(tx)** Schedl 1971c: 397, 1972f: 245, 1978a: 61.

subaplanatus (Schedl) 1935n: 634 (*Platypus*). Lectotype ♂; Malay States: Selangor, Ulu Gombak, 17th mile; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 70.

Distribution: Asia (Malaya).

Hosts: Leguminosae sp.

References: **(cn)** Yunus & Hua 1980: 217. **(hb)** Mercer 1982: 210. **(ds)** Browne 1949b: 892–912, 1961c: 209, 1962c: 202; Mercer 1982: 210; Schedl 1972f: 245; Yunus & Hua 1980: 217. **(tx)** Schedl 1935n: 634, 1939b: 401, 1972f: 245, 1978a: 70.

taxicornis (Schedl) 1968f: 537 (*Platypus*). Holotype ♂; New Guinea (NW), Wau, Morobe District, 1050 m; BPBM, Honolulu. Distribution: New Guinea.

Notes: (1) Schedl 1972f: 245 (to *Treptoplatypus*). References: **(ds)** Schedl 1972f: 245. **(tx)** Schedl 1968f: 535–537, 1972f: 245, 1978a: 72.

trepanatus (Chapuis) 1865: 75 (*Crossotarsus*). Holotype ♀; Borneo (Sarawak); BMNH, London (Schedl 1960c: 9).

Figures: Schedl 1972f: 243.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Eugenia* sp., *Gonystylus bancanus*, *Shorea uliginosa*.

References: **(cn)** Supriana et al. 1978. **(hb)** Browne 1961c: 209; Mercer 1982: 210. **(ds)** Browne 1961c: 209; Mercer 1982: 210; Sampson 1928b; Schedl 1972f: 245; Strohmeier 1912c: 5, 1914c: 35; Supriana et al. 1978. **(tx)** Browne 1949b: 911, 1962e: 653; Chapuis 1865: 75–76; Sampson 1928b; Schedl 1935n: 633–634, 1939b: 390, 401, 1939c: 335, 1960c: 9, 1972f: 243, 245; Strohmeier 1907g: 30, 1912c: 5, 1914c: 35; Wood, S. L. 1992b: 91.

Genus *Crossotarsus* Chapuis

CROSSOTARSUS CHAPUIS 1865: 23, 44. Type-species: *Crossotarsus wallacei* Thomson, subsequent designation by Hopkins 1914: 119.

References: **(ay)** Nobuchi 1969a: 69; Strohmeier 1918: 1–46. **(cn)** Beeson 1916: 216–223, 1918: 114–124; Browne 1936: 1; Chatterjee 1917: 1–4; Fisher, Thompson, & Webb 1954: 1–21; Hargreaves 1925: 21–28; Mayne 1917: 1–80; Munro 1928: 1–29; Smith 1932: 229–246; Yano 1919: 453–470. **(hb)** Beeson 1910: 222, 1937: 45–103,

1941: 328–338, 1961: 252–257; Browne 1941: 62–63, 1958b: 164–182, 1960: 313, 1961c: 221–229; Gardner 1932b: 1–6, 9; Kalshoven 1960d: 31–50; Schedl 1972f: 96. **(ds)** Beeson 1938: 295, 1941: 328–338, 1961: 252–257; Kleine 1931: 152, 165; Kolbe 1897: 283; Schedl 1966b: 83, 1972f: 96, 1977b: 234. **(tx)** Beeson 1937: 45–103; Blandford 1894d: 128–129, 1895b: 82, 85, 90; Browne 1950b: 649, 1955: 364, 1962c: 642, 1966: 233–257, 1972a: 187; Chapuis 1865: 22–26, 44–97, figs. 1–29; Choo 1983: 31; Choo, Woo, & Nobuchi 1988b; Hopkins 1914: 119, 136, 1915c: 188–227; Lacordaire 1866: 389–390; Murayama 1925: 201–202, 1928: 29–30, 1934: 147; Nisima 1909: 170; Nobuchi 1973b: 3; Numberg 1953: 46; Sampson 1923: 449; Schedl 1934f: 1647, 1935n: 633, 1936c: 43, 1937d: 33, 1939b: 401, 1959a: 515, 1960c: 5, 1972f: 96–112, 1977b: 234; Stebbing 1907: 19, 1914: 613–619; Strohmeier 1911a: 174, 1911b: 26, 1911c: 104–105, 1911d: 218, 1912c: 4–7, 1912f: 79, 1914c: 5–35.

adursus Beeson 1937: 69. Holotype ♂; Borneo: "E.I.M."; BMNH, London.

Distribution: Indonesia (Borneo).

References: **(ds)** Schedl 1972f: 110. **(tx)** Beeson 1937: 69–70; Schedl 1939c: 334, 1972f: 110.

andamanus Beeson 1937: 82. Syntypes ♂ ♀; Andamans: North Andaman; FRI, Delhra Dum.

Distribution: Asia (Andaman Islands in India).

Hosts: *Diospyros oocarpa*, *Dipterocarpus turbinatus*, *Myristica andamanica*, *Pterocarpus dalbergioides*, *Sagraea elliptica*, *Terminalia* spp.

Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *squamulatus*).

References: **(cn)** Mathur & Singh 1961a: 5, 1961b: 32; Sen-Sarma & Thakur 1986: 37. **(ds)** Mathur & Singh 1961a: 5, 1961b: 32; Schedl 1972f: 107. **(tx)** Beeson 1937: 82; Schedl 1972f: 107, 1978a: 11.

armipennis Lea 1910: 134. Holotype ♀?; N.S. Wales: Macleay Museum, not located.

Distribution: Australia (New South Wales).

References: **(cn)** Froggatt 1927; Smith, J. H. 1935: 3; Tryon 1917. **(cc)** Smith, J. H. 1935: 3. **(hb)** Froggatt 1927. **(ds)** Sampson 1925: 2; Schedl 1936g: 513, 1972a: 110; Smith, J. H. 1935: 3; Strohmeier 1912c: 5, 1914c: 35. **(tx)** Beeson 1937: 72; Browne 1962e: 644; Lea 1910: 134; Sampson 1925: 2; Schedl 1939b: 400, 1939p: 288, 1942c: 165, 1972f: 110; Strohmeier 1912c: 5, 1914f: 7–8, 1914c: 35.

aruensis Beeson 1937: 54. Holotype ♀; Aru Is.: Wokan; FRI, Delhra Dum.

Distribution: Indonesia (Aru Island).

References: **(ds)** Schedl 1972f: 101. **(tx)** Beeson 1937: 54–55; Schedl 1972f: 101.

aureipilus Roberts 1989: 88. Holotype ♂; New Guinea: Papua, Gumi; NHMW, Wien.

Distribution: New Guinea.

Hosts: *Aglaia* sp.

References: (tx) Roberts 1959: 55.

barbatus Chapuis 1865: 66. Syntypes ♂ ♀; Molucques, Ceram, Bouru; 1 ♂, 1 ♀ in BMNH, London, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 5). Distribution: Indonesia (Borneo, Buru, Ceram, Moluccas), New Guinea.

Hosts: *Theobroma cacao*.

References: (ay) Strohmeier 1918: 15. (cc) Roberts 1950. (ds) Browne 1950c: 452, 1951a: 126; Gemminger & Harold 1872: 2696; Ohno, Yoshioka, et al. 1958b: 97; Sampson 1925b: 390; Schedl 1936g: 513; Strohmeier 1912c: 4, 1914c: 33. (tx) Beeson 1937: 56; Chapuis 1865: 66; Sampson 1925b: 390; Schedl 1936d: 12, 1936e: 43, 1937d: 34, 1939b: 402, 1940b: 434–435, 1955b: 282, 1960c: 8, 1972f: 103; Sharp & Muir 1912: 572; Strohmeier 1912c: 4, 1914c: 33; Wylie & Yule 1977.

benecolus Beeson 1937: 50. Holotype ♂; Sumbawa; BMNH, London.

Distribution: Indonesia (Sumbawa).

References: (ds) Schedl 1972f: 101. (tx) Beeson 1937: 50–51; Schedl 1972f: 101.

biconcavus Schedl 1962i: 76. Holotype ♂; Ned. Nieuw Guinea, Sibil Sterrengeb.; ZSSM, Munchen. Distribution: New Guinea, New Ireland.

Hosts: *Theobroma cacao*.

Notes: (3) Schedl 1970: 221 (described female). References: (cn) Roberts 1957; Wylie & Shanahan 1975. (hb) Roberts 1957; Wylie & Shanahan 1975. (ds) Schedl 1962i: 73, 76, 1964d: 212, 1968e: 264, 1969e: 156, 1972f: 101. (tx) Schedl 1962i: 76, 1969b: 221, 1972f: 101, 1975a: 14.

bonrouloiri Chapuis 1865: 55. Syntypes ♂ ♀; Camboje; 1 ♂, 1 ♀ in BMNH, London, 1 ♀ in IRSNB, Brussels (Schedl 1960).

Distribution: Asia (Cambodia/ Assam, Bengal in India/ Malaya/ Thailand), Indonesia (Celebes, Sumatra), Philippine Islands, Sunda Islands.

Hosts: *Buka lindia*, *Citrus* sp., *Cryptocarya wrightiana*, *Dysoxylum binectariferum*, *Engelhardtia spicata*, *Mesua ferrea*, *Sapium eugeniaefolium*, *Shorea robusta*, *Terminalia bellerica*, *Vatica lanceaefolia*, *Xylia xylocarpa*.

References: (bv) Beeson 1917. (cn) Beeson 1916: 1–5; Kleine 1932a: 310, 401; Mathur & Singh 1960a: 35, 1961a: 23, 1961b: 25; Roonwal 1954: 65; Sen-Sarma & Thakur 1956: 37. (cc) Beeson 1921b, 1923. (hb) Beaver & Browne 1975: 302; Beeson 1916a: 222, 1917, 1923; Browne 1961c: 224; Kleine 1932a: 310, 401. (ds) Beaver & Browne 1975: 302; Beeson 1916a, 1921b: 21, 1923, 1961: 253; Bhasin, Roonwal, & Singh 1958; Browne 1961c: 55, 1950c: 484; Gemminger & Harold 1872: 2696; Kleine 1932a: 310, 401; Lacordaire 1866: 389; Mathur & Singh 1960a: 35, 1961a: 23, 1961b: 25; Roonwal 1954: 65; Schedl 1969c: 55, 1972f: 101; Strohmeier 1911f: 203,

1912c: 4, 1914c: 33. (tx) Beeson 1937: 50–53; Chapuis 1865: 55; Lacordaire 1866: 389; Schedl 1937d: 33, 1960c: 6, 1972f: 101; Strohmeier 1912c: 4, 1914c: 33.

brevidens Browne 1975: 302 in Beaver & Browne. Holotype ♂; Thailand: Chiang Mai, Doi Pui, 1450 m; BMNH, London.

Distribution: Asia (Thailand).

Hosts: *Quercus* sp.

References: (tx) References: Beaver & Browne 1975: 302–303.

brownei Schedl 1936d: 12. Lectotype ♂; Malay Peninsula, Pahang; Rotan Tunggal For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 17.

Distribution: Asia (Malaya), Indonesia (Borneo).

Hosts: *Dacryodes rostrata*, *Elateriospermum tapos*, *Gluta elegans*, *Mangifera* sp., *Melanorrhoea curtisii*, *Shorea leprosula*.

Notes: (3) Schedl 1942a: 199 (described female).

References: (cc) Browne 1955b. (hb) Beaver & Browne 1975: 615; Browne 1935a, 1941, 1958b, 1961c: 224; Yunus & Hua 1980: 216. (ds) Beaver & Browne 1975: 615; Beeson 1961: 253; Browne 1935a, 1949b, 1961a: 316, 1961c: 224; Schedl 1972f: 103. (tx) Browne 1949b, 1955: 365; Schedl 1936d: 12–13, 1942a: 199, 1950g: 894, 1972f: 103, 1978a: 17.

desectus Beeson 1937: 58. Holotype ♂; Borneo: Kuching; BMNH, London. Synonymy: Browne 1955: 365.

References: (tx) Beeson 1937: 58; Browne 1955: 365; Schedl 1972f: 103.

caliginosus Roberts 1979: 81. Holotype ♂; Papua New Guinea: 12 km south-west of Onim Hut, Momt Gihwe, 2900m; BMNH, London.

Distribution: New Guinea.

Hosts: *Quintinia* sp.

References: (tx) Roberts 1979: 81.

chalcographus Schedl 1972h: 64. Holotype ♂; New Guinea: Kulolo logging area, Wau, Morobe District; CSIRO, Canberra.

Distribution: Australia, New Guinea.

Hosts: *Calophyllum inophyllum*, *Castanopsis accuminatissima*, *Pasania aspericupula*.

Notes: (1) Five localities are given with no indication of which one is the type locality.

References: (ds) Schedl 1972f: 103, 1980b: 184. (tx) Schedl 1972f: 103, 1972h: 64, 1975a: 19.

cheesmani Schedl 1935f: 275. Lectotype ♂; Papua: Mt. Tafa, 8500 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 19.

Distribution: New Britain Island, New Guinea.

References: (ds) Schedl 1968f: 535, 1972f: 110. (tx) Schedl 1935f: 275, 1935j: 2, 1940b: 434, 1972f: 110, 1978a: 19.

cincinnatus Chapuis 1865: 57. Syntypes 4 ♀; Borneo (Sarawak); BMNH, London.

Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo, Sumatra).

Hosts: *Coccoceras* sp.

Notes: (3) Schedl 1935n: 639 (described male).

References: (**hb**) Browne 1935a, 1941. (**ds**) Beeson 1937, 1941 (1961: 253); Browne 1935a; Gemminger & Harold 1872: 2696; Sampson 1925a: 3, 1925b: 389; Schedl 1936d: 6, 1960c: 6, 1966b: 83, 1972f: 102; Strohmeier 1911f: 203, 1912c: 4, 1914c: 33. (**tx**) Beeson 1937: 54, 56; Browne 1962c: 214; Chapuis 1865: 57; Nobuchi 1985: 328; Sampson 1919: 105, 1925a: 3, 1925b: 389; Schedl 1935n: 639, 1939e: 333, 1952b: 385, 1960c: 6, 1972f: 102; Strohmeier 1912c: 4, 1914c: 33.

penicillatus Chapuis 1865: 64. Holotype ♂; Borneo (Saravak); BMNH, London (Schedl 1960c: 7). Synonymy: Browne 1962c: 214.

References: (**ds**) Gemminger & Harold 1872: 2697; Sampson 1919: 105, 1925b: 389; Strohmeier 1912c: 4, 1914c: 33. (**tx**) Beeson 1937: 54, 56; Browne 1962c: 214; Chapuis 1865: 64; Sampson 1919: 105, 1925b: 389; Schedl 1960c: 7, 1972f: 214; Strohmeier 1912c: 4, 1914c: 33.

cinnamomi Beeson 1937: 97. Syntypes ♂; Bengal: Kalimpong division, Samsingh, 1800 m; FRI, Dehra Dun.

Distribution: Asia (Assam, Bengal, Uttar Pradesh in India).

Hosts: *Cinnamomum cecidodaphne*, *Ilex dipyrrena*.

References: (**cn**) Mathur & Singh 1960b: 63. (**ds**) Beeson 1941 (1961: 253); Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1960b: 63; Schedl 1969c: 55, 1972f: 110. (**tx**) Beeson 1937: 97–98; Schedl 1972f: 110, 1978a: 19.

cliens (Schedl) 1964g: 253 (*Platypus*). Holotype ♂; Sarawak, Merurong Plateau, 2350 ft.; BMNH, London.

Figures: Schedl 1972f: 110.

Distribution: Indonesia (Sarawak in Borneo).

Hosts: *Santiria* sp.

References: (**ds**) Browne 1980a: 372; Schedl 1972f: 111. (**tx**) Browne 1962c: 644; Schedl 1964g: 242, 253, 1972f: 110–111, 1978a: 20.

comatus Chapuis 1865: 59. Holotype ♀; Celebes (Macassar); BMNH, London (Schedl 1960c: 7).

Distribution: Asia (Malaya), Indonesia (Celebes), Philippine Islands.

References: (**cn**) Mesa 1931: 18. (**hb**) Browne 1935a. (**ds**) Beeson 1941 (1961: 253); Browne 1935a; Gemminger & Harold 1872: 2696; Numborg 1964a: 237; Schedl 1966b: 83, 1972f: 102; Strohmeier 1912c: 4, 1914c: 33. (**tx**) Beeson 1937: 55; Chapuis 1865: 59; Nobuchi 1985: 328; Schedl 1960c: 7, 1972f: 102; Strohmeier 1911b: 26, 1912c: 4, 1914c: 33.

concarifrons Schedl 1935b: 399. Syntypes ♀; Philippine Islands; Hamburg Museum, lost, and Schedl Collection in NHMW, Wien.

Distribution: Philippine Islands.

References: (**ds**) Schedl 1966b: 83, 1972f: 112. (**tx**) Nobuchi 1985: 328; Schedl 1935b: 395–399, 1972f: 112.

concinnus Blandford 1894c: 578. Holotype ♀; Japan: Higo; BMNH, London, automatic.

Distribution: Asia (Japan/Taiwan).

References: (**ds**) Blandford 1894c: 578; Murayama 1929a: 675, 1937c: 582, 1954b: 205; Nobuchi 1973b: 5; Schedl 1972f: 100; Strohmeier 1912c: 4, 1914c: 34. (**tx**) Lucas 1920: 209; Murayama 1925b: 235, 1954b: 205; Nobuchi 1973b: 5; Schedl 1934f: 1647, 1939b: 402, 1972f: 100; Strohmeier 1912c: 4, 1914c: 34. (**ms**) Lucas 1920: 209.

chapuisi Blandford 1894d: 129. Holotype ♀; Japan: Higo; BMNH, London, preoccupied by Duvivier 1891.

References: (**tx**) Blandford 1894d: 129, 1894c: 578.

coniferae Stebbing 1906: 411. Syntypes ♂ ♀; North-West Himalayas: Tehri Garhwal forest; FRI, Dehra Dun.

Distribution: Asia (Kashmir, Punjab, Uttar Pradesh in India).

Hosts: *Cedrus deodara*, *Picea morinda*.

References: (**cn**) Kleine 1932a: 310, 401; Roonwal 1954: 12; Stebbing 1914: 613. (**cc**) Beeson 1921b; Stebbing 1914: 613. (**hb**) Beeson 1916: 222; Kleine 1932a: 310, 401; Stebbing 1908a: 112, 1911a: 66, 1914: 613. (**ds**) Beeson 1921b: 21, 1961: 253; Bhasin, Roonwal, & Singh 1955; Kleine 1932a: 310, 401; Roonwal 1954: 12; Schedl 1969c: 56, 1972f: 104; Strohmeier 1912c: 5, 1914c: 35. (**tx**) Schedl 1972f: 104; Stebbing 1906: 411–413, 1910: 66, 1914: 613; Strohmeier 1912c: 5, 1914c: 35.

corrugatus Roberts 1989: 88. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; NHMW, Wien.

Distribution: New Guinea.

Hosts: *Aglaia* sp.

References: (**tx**) Roberts 1989: 88.

coxalis Schedl 1970a: 129. Holotype ♂; New Guinea, Porotop, Lutheran Mission Sawmill, Western Highlands District; CSIRO, Canberra.

Distribution: New Guinea.

Hosts: *Nothofagus* sp., *Podocarpus* sp., *Syzygium* sp.

References: (**bv**) Gray, B. 1974b. (**hb**) Gray, B. 1974b. (**ds**) Schedl 1972f: 103. (**tx**) Schedl 1970a: 129–130, 1972f: 103, 1978a: 22.

dammanus Beeson 1937: 63. Holotype ♂; Damma Is., 92–20, 5944; BMNH, London.

Distribution: Indonesia (Damma, Moluccas).

Hosts: "Canari" log.

Notes: Browne 1981b: 606 (described female).

References: (**ds**) Schedl 1972f: 109. (**tx**) Beeson 1937: 63–64; Browne 1981b: 606; Schedl 1972f: 109.

declivioruatus (Schedl) 1942a: 205 (*Platypus*).
Lectotype ♂; Malaya, Perak, Trolak F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1975a: 23.

Distribution: Asia (Malaya).

Hosts: Rubiaceae sp., *Castanopsis sumatrana*.

Notes: (1) Browne 1962e: 643 (to *Crossotarsus*).

References: (ds) Browne 1961c: 229; Schedl 1972f: 106. (tx) Browne 1962e: 643; Schedl 1942a: 205, 1972f: 106, 1975a: 23.

denticulus Browne 1954f: 57. Holotype ♂; New Guinea: Morobe District, Wau, Kumi Creek, 1500 m; BMNH, London.

Distribution: New Guinea.

References: (tx) Browne 1954f: 57.

denturus Browne 1954f: 55. Holotype ♂; New Guinea: Morobe District, Mount Kaindi, 2350 m; BMNH, London.

Distribution: New Guinea.

Hosts: *Nothofagus* sp.

References: (tx) Browne 1954f: 55.

derosus Schedl 1935j: 2. Holotype ♂; Sumatra, Medan; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Java, Sumatra).

References: (ds) Schedl 1972f: 104. (tx) Beeson 1937: 77; Schedl 1935j: 2, 1938g: 422, 1972f: 104, 1975a: 24.

dolus Schedl 1975f: 373. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 373.

edentatus Beeson 1937: 90. Holotype ♂; Java: Mt. Preanger, 1400 m; Buitenzorg Museum.

Distribution: Indonesia (Java).

Hosts: *Schima noronhai*.

References: (ds) Kalshoven 1960d: 36; Schedl 1972f: 112. (tx) Beeson 1937: 90; Schedl 1972f: 112.

emancipatus Murayama 1934f: 138. Syntypes ♂ ♀; Kiushu; Murayama Collection in USNM, Washington.

Figures: Nobuchi 1973b: pl. 1.

Distribution: Asia (Japan/Taiwan).

Hosts: *Actinodaphne longifolia*, *Castanopsis cuspidata*, *Clochidion hongkongense*, *Lindera erythrocarpa*, *Machilus thunbergii*, *Mallotus japonicus*, *Pasania cuspidata*, *Quercus salicina*.

References: (cn) Kleine 1932a: 401. (hb) Kleine 1932a: 401. (ds) Kleine 1932a: 401; Murayama 1934f: 138, 1936a: 140, 1937c: 582, 1954b: 206, 1955: 100; Nobuchi 1967: 25, 1973b: 5, 1977: 106; Schedl 1972f: 106. (tx) Beeson 1937: 84; Murayama 1928a: 288, 1934f: 138, 1936a: 140, 1954b: 206, 1955: 100–101; Nobuchi 1973b: 5, pl. 1; Schedl 1972f: 106.

emorsus Beeson 1937: 87. Syntypes ♂ ♀; Burma: Katha division, Mohmyin reserve; FRI, Dehra Dun.

Distribution: Asia (Burma), Indonesia (Borneo).

Hosts: *Bauhinia malabarica*, *Careya arborea*, *Sterculia ornata*, *Tectona grandis*, *Xylia dolabriformis*.

Notes: (3) A male type is labeled in the FRI Collection (SIW).

References: (cn) Mathur & Singh 1961a: 70, 1961b: 13. (ds) Beeson 1941 (1961: 253); Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1961a: 70, 1961b: 13; Schedl 1972f: 107. (tx) Beeson 1937: 87–88; Schedl 1939e: 335, 1972f: 107.

externedentatus (Fairmaire) 1849: 78 (*Platypus*).

Holotype ♂; Taïti, M. Vesco; MNHN, Paris.

Figures: Nobuchi 1967: pl. 2, 1973b: pl. 1, Wood 1960a: 7–8.

Distribution: Africa (Mauritius Island/ Seychelles Islands/ South Africa/ Tanzania), Asia (Burma/ Andaman Islands, Assam, Bengal, Karnataka, Tamil Nadu in India/ Japan/ Malaya/ Sri Lanka/ Taiwan/ Thailand/ Tonkin Island in Vietnam), Australia (W Australia), Bismarck Islands, Fiji Islands, Hawaiian Islands, Indonesia (Borneo, Celebes, Java, Sumatra), Madagascar, Mangareva, Micronesia (Caroline Islands, Cook Islands, Kiribati Islands, Palau), Moorea Island, New Zealand, Niue Island, Philippine Islands (Luzon), Raoul Island, Samoan Islands, Society Islands, Tahiti.

Hosts: Anonymous 1970c, Beeson 1937: 73, 1941 (1961: 254–255) (many hosts listed), Browne 1961c (many hosts listed), Schedl 1962: 678–680 (41 host species listed). *Bauhinia* sp., *Buchanania* spp., *Butea monosperma*, *Casuarina cuphyllum*, *Cassia* spp., *Castanopsis cuspidata*, *C. tribuloides*, *Casuarina equisetifolia*, *Cedrela odorata*, *Cinnamomum camphora*, *Cryptomeria japonica*, *Ficus retusa*, *Heritiera fomes*, *Leucaena glauca*, *Machilus thunbergii*, *Odina woderi*, *Quercus* sp., *Shorea* spp., *Terminalia tomentosa*.

Notes: (3) Fairmaire 1850: 51 (redescribed). Wood 1992b: 91 and others have considered *Diapus talurae* Stebbing as a synonym; however, the holotype was found and *talurae* was transferred to *Genyocerus* (SIW).

References: (bv) Gray, B. 1974b; Roberts 1977c. (cn) Anonymous 1967s, 1970c: 11; Beaver 1988a: 69; Ceyer 1960; Pierce, J. R., Wood, & Fujii 1977: 17; Roberts 1977b: 555, 1977c. (ce) Scott & Dutoit 1970; Walt & Scott 1971a, 1971b; Walt, Scott, & van der Klift 1971a: 458, 1971b: 464. (hb) Beaver 1976b: 544, 1988a: 69; Beeson 1929; Gray, B. 1974b; Roberts 1977b: 55, 1977c; Schedl 1962k: 674, 1977b: 235. (ds) Anonymous 1967s, 1970c: 11; Barber 1919: 53–60; Beaver 1976b: 544, 1987b: 65, 1988a: 69, 1990b: 150; Beaver & Browne 1975: 303, 1978: 615; Beaver & Maddison 1990: 1372; Beeson 1915: 297–298, 1929: 218, 1935a: 115, 1935b, 1937: 72–73, 1938b: 295, 1940: 191; Browne 1949: 74, 1966: 241, 1980a: 372, 1980c: 485; Chapuis 1865; Dumbleton 1954: 25, 184; Dupont 1916: 20–22; Gemminger &

Harold 1872: 2696; Lacordaire 1866: 390; Lever 1943a: 82; Miwa 1931: 266; Murayama 1925a: 203, 1929a: 676, 1937c: 582; Nobuchi 1967: 25, 1973b: 5, 1977: 108; Olmo, Yoneyama, & Nakazawa 1982a, 1982b, 1987a: 95, 1987b: 89; Olmo, Yoshioka, et al. 1988b: 97, 1989: 64; Pierce, J. R., Wood, & Fujii 1977: 17; Roberts 1977b: 555; Roonwal 1954: 36; Schedl 1934f: 1647, 1960c: 10, 1961e: 128, 1962b: 187, 1962i: 77, 1962k: 674, 1964c: 305, 1964d: 212, 1965a: 340, 1965g: 23, 1966b: 84, 1969a: 206, 1969c: 56, 1969d: 4, 7, 9, 1970d: 234, 1970h: 179, 1971a: 281, 1972b: 266, 1972f: 111, 1974c: 262, 1975k: 278, 1977b: 235; Strohmer 1911f: 204, 1912c: 5, 1914c: 35; Swezey 1941: 124, 126, 1954: 14, 78, 112, 198; Wood, S. L. 1960a: 7; Zimmerman 1941: 176. **(tx)** Beeson 1929: 218, 1935a: 115, 1935b, 1940; Browne 1962e: 644; Chapuis 1865: 81–82; Fairmaire 1847: 78, 1850: 51, 1866: 81; Lacordaire 1866: 390; Murayama 1925a: 203–204, 1925b: 235; Nobuchi 1967: pl. 2, 1972b: 5, 1973b: pl. 1, 1985: 328; Perkins 1900: 181; Powell, W. 1987: 28; Schedl 1934f: 1647, 1937d: 34, 1939b: 398, 1941e: 153, 1941f: 116, 1948c: 665, 1950f: 35, 41, 1951k: 141, 1952j: 6, 1953b: 124, 1953d: 70–71, 1955b: 286, 1957b: 150–152, 1960c: 10–11, 1962k: 674–675, 1964d: 212, 1972f: 111, 1972p: 156; Sharp (in Perkins) 1900: 173, 182; Sharp & Muir 1912: 572; Stebbing 1914: 628–630; Strohmer 1911f: 204, 1912c: 5, 1914c: 11, 35; Wood, S. L. 1960a: 7–8, 1992b: 91 (error, Note 3 above). **(ms)** Roberts 1977c.

saundersi Chapuis 1865: 80. Syntypes ♂ ♀; Celebes, Borneo; 3 ♂, 2 ♀ in BMNH, London, 1 ♂ IRSNB, Brussels (Schedl 1960c: 10). Synonymy: Schedl 1972f: 111.

References: **(bv)** Beeson 1917; Bletchly 1961: 15. **(cn)** Anonymous 1970c: 11; Beeson 1919: 1–23, 1935a; Browne 1952; Kalshoven 1924c: 310; Lever 1940a: 38, 1943a: 14, 1946: 4; Mathur & Singh 1960a: 45, 1960b: 6, 1960c: 3, 1961a: 5, 1961b: 13; Roonwal 1954: 33; Sen-Sarma & Thakur 1986: 37; Stebbing 1914: 617; Yunus & Hua 1980: 216. **(cc)** Beeson 1921b: 22, 1923; Bhatia 1950; Browne 1958b; Stebbing 1914: 617. **(hb)** Beeson 1915b: 298, 1916a: 222, 1917, 1923, 1933, 1938a; Browne 1935a, 1941, 1958b, 1961c: 218; Kalshoven 1924c, 1960c: 42, 1960d: 1; Kleine 1932a: 310, 401; Speyer 1923: 22; Stebbing 1914: 617. **(ds)** Anonymous 1970c: 11; Beeson 1915b, 1916a: 1–5, 1921b: 22, 1923, 1933: 9, 1935: 539–543, 1938a, 1941 (1961: 254); Bhasin, Roonwal, & Singh 1958; Bhatia 1950; Blandford 1895a, 1898a; Browne 1935a, 1961a: 316, 1961c: 218, 1966: 241, 1980a: 372; Gemminger & Harold 1872: 2697; Kalshoven 1924b: 357; Kleine 1932a: 310, 401; Lever 1940a: 38–42, 1940b: 99, 1941: 80, 1942: 23, 1943a: 14, 1945: 367–377; Mathew 1982, 1987: 188; Mathur & Singh

1960a: 45, 1960b: 6, 1960b: 3, 1961a: 5, 1961b: 13; Olmo, Yoneyama, & Nakazawa 1982b: 9; Roonwal 1954: 33; Sampson 1928a: 2, 1928b: 390; Schedl 1936d: 6, 1936j: 21, 1937f: 15, 1948c: 665, 1950f: 41, 1953b: 124, 1959a: 513, 1971a: 279; Speyer 1923: 11–23; Strohmer 1912c: 6, 1914c: 35. **(tx)** Beeson 1933, 1937: 37, 72; Browne 1962e: 644; Chapuis 1865: 80; Gardner 1932b; Murayama 1925: 203; Sampson 1924b: 89, 1926: 120, 1928a: 2, 1928b: 390; Schedl 1935n: 640, 1936e: 43, 1937d: 34, 1939e: 334, 1941c: 354, 1942a: 172, 206, 1942c: 163–164, 1954a: 145, 1959a: 513, 1960c: 10, 1964g: 253, 1972f: 111; Stebbing 1914: 617; Strohmer 1912c: 6, 1914c: 35, 1914f: 8.

heritiera Stebbing 1906: 420 (*Diapus*). Syntypes ♂ ♀; India, Bengal: Backerganj District, Wazirpur, Sunderbans; FRI, Dehra Dun. Synonymy: Beeson 1915: 298.

References: **(cn)** Stebbing 1914: 628. **(cc)** Stebbing 1914: 628. **(hb)** Stebbing 1914: 628. **(ds)** Strohmer 1912c: 22, 1914c: 47. **(tx)** Stebbing 1914: 628; Strohmer 1912c: 22, 1914: 47.

posticus Broun 1909: 301 (*Platypus*). Holotype ♀; Raoul Island (New Zealand); Broun Collection (damaged). Synonymy: Schedl 1972f: 111.

References: **(tx)** Broun 1909: 301; Schedl 1972f: 111, 1972p: 156.

saundersi usambaricensis Strohmer 1911: 182. Syntypes ♂ ♀; Usambara; IPKE, Eberswalde. Synonymy: Schedl 1972f: 111.

References: **(hb)** Strohmer 1911a: 182. **(tx)** Schedl 1972f: 111; Strohmer 1911a: 182.

saundersi submontanus Beeson 1937: 72. Syntypes ♂ ♀; India, Burma, Ceylon, United Provinces: Dehra Dun division, 2000–2500 ft.; FRI, Dehra Dun. Synonymy: Schedl 1972f: 111.

References: **(cn)** Mathur & Singh 1960b: 6, 1960c: 6, 1961a: 32, 1961b: 26; Sen-Sarma & Thakur 1986: 37. **(ds)** Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1960b: 6, 1960c: 6, 1961a: 32, 1961b: 26. **(tx)** Beeson 1937: 72; Schedl 1972f: 111.

nilgircus Beeson 1937: 73. Syntypes, ♂ ♀; Madras: Nilgiris, Coonoor Valley; FRI, Dehra Dun. Synonymy: Schedl 1972f: 111.

References: **(ds)** Bhasin, Roonwal, & Singh 1958; Mathew 1982, 1987: 188. **(tx)** Beeson 1937: 73; Schedl 1972f: 111.

externespinulosus (Schedl) 1942a: 206 (*Platypus*). Holotype ♂; Malaya: Pahang, Rotan Tinggi F.R.; Schedl Collection, lost, authentic examples in BMNH, London.

Distribution: Asia (Malaya).

Notes: (1) Browne 1961c: 220, 1962e: 644 (to *Crossotarsus*).

References: **(ds)** Browne 1961c: 220; Schedl 1972f:

111. **(tx)** Browne 1962c: 644; Schedl 1942a: 206, 1972f: 111, 1978a: 29.
- fagacearum** **Browne** 1975: 303 in Beaver & Browne. Holotype ♂: Thailand: Chiang Mai, Doi Pui, 1450 m; BMNH, London.
Distribution: Asia (Thailand).
Hosts: *Castanopsis* cf. *acuminatissima*, *Quercus* sp.
References: **(tx)** Beaver & Browne 1975: 303–304.
- fairmairei** **Chapuis** 1865: 79. Syntypes ♂; Inde: 1 ♂ in BMNH, London, 3 ♂ in IRSNB, Brussels (Schedl 1960c: 10).
Distribution: Asia (Punjab, Uttar Pradesh in India/ Japan/ Sri Lanka).
Hosts: *Abies pindrow*, *Cedrus deodara*, *Juglans regia*, *Pinus excelsa*, *Quercus incana*, *Q. semicarpifolia*.
References: **(cn)** Kleine 1932a: 310, 401; Mathur & Singh 1960b: 80; Roonwal 1954: 17, 29; Stebbing 1914: 617. **(cc)** Beeson 1921b: 21; Stebbing 1914: 617. **(hb)** Kleine 1932a: 310, 401; Stebbing 1914: 617. **(ds)** Beeson 1921b: 21, 1937: 73–74, 1961: 253; Blasin, Roonwal, & Singh 1958; Gemminger & Harold 1872: 2697; Kleine 1932a: 310, 401; Mathur & Singh 1960b: 80; Nobuchi 1973b: 6; Roonwal 1954: 17, 29; Schedl 1969c: 56, 1972f: 104; Strohmeier 1911f: 203, 1912c: 5, 1914c: 35. **(tx)** Beeson 1937: 73–74; Chapuis 1865: 79; Nobuchi 1973b: 6; Sampson 1919: 110; Schedl 1960c: 10, 1972f: 104; Stebbing 1914: 617; Strohmeier 1912c: 5, 1912d: 42, 1914c: 35.
- flavomaculatus** **Strohmeier** 1912: 40. Holotype ♂; Hoozan in Formosa, Philippinen; MNB, Berlin. Figures: Nobuchi 1967:pl. 2, 1973b:pl. 1.
Distribution: Asia (China ?/ Japan/ Malaya/ Taiwan), Indonesia (Java), Philippine Islands.
Hosts: *Quercus glauca*.
References: **(hb)** Murayama 1931b: 199. **(ds)** Miwa 1931: 266; Murayama 1925a: 205, 1931b: 199, 1934f: 138, 1937c: 581, 1954b: 206, 1955: 100; Nobuchi 1967: 25, 1973b: 7; Schedl 1966b: 84, 1972f: 104; Strohmeier 1912c: 5, 1914c: 35. **(tx)** Murayama 1925a: 205–206, 1925b: 235, 1954b: 206, 1955: 100–101; Nobuchi 1967:pl. 2, 1973b: 7, pl. 1, 1955: 328; Schedl 1934f: 1647, 1972f: 104; Strohmeier 1912c: 5, 1912d: 40–41, 1914c: 35.
- fluminalis** **Beeson** 1937: 57. Holotype ♀; New Guinea: N.N. Guinea Exped., Albatros Bivak V, Manberano; Buitenzorg Museum.
Distribution: New Guinea.
References: **(ds)** Schedl 1972f: 102. **(tx)** Beeson 1937: 57–58; Schedl 1972f: 102.
- forcipes** **Schedl** 1975f: 374. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975f: 374.
- formosanus** **Strohmeier** 1912: 41. Syntypes ♂ ♀; Kosempo (Formosa); 2 ♂, 1 ♀ in MNB, Berlin, some in Strohmeier Collection.
Distribution: Asia (Taiwan).
References: **(ds)** Murayama 1925a: 206, 1929a: 676, 1937c: 582; Nobuchi 1967: 26; Schedl 1972f: 111; Strohmeier 1912c: 5, 1914c: 35. **(tx)** Murayama 1925a: 206–207, 1925b: 235, 1928a: 287–288; Schedl 1972f: 111; Strohmeier 1912c: 5, 1912d: 40–41, 1914c: 35.
- fractus** **Sampson** 1912: 249. Holotype ♂; Borneo: Kuching, Sarawak; BMNH, London.
Distribution: Asia (Assam in India/ Malaya), Bismarck Islands, Indonesia (Borneo, Java, Sumatra), Philippine Islands (Luzon).
Hosts: *Azelia palembanica*, *Albizia procera*, *Dalbergia latifolia*, *Dryobalanops oblongifolia*, *Eucalyptus* sp., *Grewia* sp., *Palaquium stellatum*, *Quercus* sp., *Styrax benzooin*, *Sucietenia macrophylla*.
Notes: (3) Schedl 1935k: 483 (described female), 1972f: 107 (treated as a subspecies of *squamulatus*).
References: **(cn)** Browne 1933b. **(hb)** Browne 1935a, 1941, 1961c: 227; Kalshoven 1960c: 45. **(ds)** Beeson 1937, 1941 (1961: 254); Browne 1935a, 1961c: 227, 1966: 242, 1980b: 351; Kalshoven 1960c: 45; Ohno, Yoneyama, & Nakazawa 1987b: 89; Schedl 1936d: 6, 1936j: 6, 21, 1938g: 421, 1966b: 87, 1972f: 107; Strohmeier 1914c: 35. **(tx)** Beeson 1937: 79, 81; Nobuchi 1985: 328; Sampson 1912: 249–250, 1919: 106; Schedl 1935k: 483, 1936e: 44–45, 1939c: 334–335, 1941c: 354, 1952b: 366, 1972f: 107, 1978a: 31; Strohmeier 1914c: 35.
- gressitti** **Schedl** 1972i: 53. Holotype ♂; New Guinea, Papua, Western District, Oriomo Gov. Station: BPBM, Honolulu.
Distribution: New Guinea.
References: **(ds)** Schedl 1972f: 104. **(tx)** Schedl 1972f: 104, 1972i: 53, 1978a: 34.
- hastatus** (**Schedl**) 1942c: 194 (*Platytypus*). Holotype ♂; Neu-Guinea; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(ds)** Schedl 1968f: 535, 1972f: 111. **(tx)** Browne 1962c: 643; Schedl 1942c: 194, 1972f: 111, 1978a: 35.
- hebridensis** **Beeson** 1937: 71. Holotype ♂; New Hebrides: Aneityum; BMNH, London.
Distribution: New Hebrides Islands.
References: **(ds)** Beeson 1938b: 295; Schedl 1972f: 111. **(tx)** Beeson 1937: 71–72; Schedl 1972f: 111.
- horni** **Schedl** 1934a: 247. Lectotype ♂; Kerhamanab, Java, Bandoeng; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 35.
Hosts: *Acacia decurrens*, *Cinchona ledgeriana*, *Horsfieldia* cf. *glabra*, *Quercus* sp.
Distribution: Indonesia (Java).
References: **(ds)** Kalshoven 1960d: 40; Schedl 1934a: 247, 1972f: 111. **(tx)** Beeson 1937: 68;

- Browne 1962e: 644; Schedl 1934a: 247, 1939c: 333, 1941c: 354, 1942a: 204, 1963j: 486, 1972f: 111, 1978a: 35.
- biceps* Beeson 1937: 67. Syntypes ♂ ♀; Java: G. Slamet, Batoerraden; Buitenzorg Museum. Synonymy: Schedl 1939c: 333. References: (tx) Beeson 1937: 67–68, 71; Browne 1962e: 644; Schedl 1939c: 333, 1972f: 111.
- inaequidens* Browne 1984c: 456. Holotype ♂; Kaonasugo (Solomon Islands) to Nagoya (Japan), imported; BMNH, London. Distribution: Solomon Islands. Hosts: *Pometia* sp. References: (ds) Ohno, Yoshioka, et al. 1988b: 97. (tx) Browne 1984c: 456.
- indomitus* Chapuis 1865: 84. Holotype ♀; Moluques, de l'île Morty; BMNH, London (Schedl 1960c: 11). Distribution: Asia (Malaya), Australia, Indonesia (Moluccas). References: (ds) Gemminger & Harold 1872: 2697; Schedl 1972f: 111; Strohmeier 1912c: 5, 1914c: 35. (tx) Chapuis 1865: 84; Schedl 1960c: 11, 1972f: 111; Strohmeier 1912c: 5, 1914c: 35.
- cavifrons* Blandford 1896b: 192. Syntypes 2 ♂; Gilolo [Moluccas]; BMNH, London. Synonymy: Schedl 1978a: 111. References: (cn) Smith, J. H. 1935: 3. (ec) Smith, J. H. 1935: 3. (ds) Smith, J. H. 1935: 3; Strohmeier 1912c: 5, 1914c: 35. (tx) Blandford 1896b: 192; Schedl 1960c: 11, 1972f: 111; Strohmeier 1912c: 5, 1914c: 35.
- inermis* Schedl 1937d: 33. Lectotype ♂; West-Sumatra, Padang-Pandjang and Sudost-Borneo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 37. Distribution: Asia (Malaya/ Thailand), Indonesia (Borneo, Mentawai, Sumatra). Hosts: *Azela palembanica*, *Aglaia* sp., *Grewia* sp., *Koompassia excelsa*, *Parkia speciosa*, *Parinarium griffithianum*, *Sterculia macrophylla*. References: (ds) Beaver & Browne 1975: 304; Schedl 1971f: 150, 1972f: 102. (tx) Schedl 1936c: 43, 1937d: 33, 1962p: 209, 1972f: 102, 1978a: 37.
- foederatus* Beeson 1937: 53. Holotype ♂; Federated Malay States; Perak; FRI, Dehra Dun. Synonymy: Schedl 1962p: 209. References: (cn) Browne 1949e. (hb) Browne 1941, 1961c: 223. (ds) Browne 1961c: 223. (tx) Beeson 1937: 53–54; Schedl 1962p: 209, 1972f: 102.
- intermedius* Chapuis 1865: 69. Holotype ♀; des îles Celebes (Macassar); BMNH, London. Distribution: Indonesia (Celebes), New Guinea. References: (ds) Gemminger & Harold 1872: 2697; Schedl 1972a: 103; Strohmeier 1912c: 4, 1914c: 33. (tx) Chapuis 1865: 69; Schedl 1960c: 8, 1972f: 103; Strohmeier 1912c: 4, 1914c: 33.
- javanus* Beeson 1937: 51. Syntypes 1 ♂, 1 ♀; Java: Sockaboeni; FRI, Dehra Dun. Distribution: Indonesia (Java, Sumatra), New Guinea. Hosts: *Albizzia sumatrana*, *Celtis wrightia*, *Dalbergia latifolia*, *Eucalyptus* sp., *Hevea brasiliensis*, *Pometia tomentosa*. References: (hb) Kalshoven 1960c: 44. (ds) Kalshoven 1960c: 44; Schedl 1971c: 367, 1971f: 150, 1972f: 102. (tx) Beeson 1937: 51–52; Schedl 1961c: 74, 1972f: 102.
- keyensis* Strohmeier 1913: 163. Syntypes ♂; Key-Inseln; SMTD, Dresden, and IPKE, Eberswalde. Distribution: Indonesia (Key Island), New Guinea. References: (ds) Schedl 1972f: 102; Strohmeier 1914c: 34. (tx) Schedl 1937d: 34, 1941c: 152, 1972f: 102, 1978a: 39; Strohmeier 1913: 163–164, 1914c: 34.
- kondulensis* Beeson 1937: 86. Syntypes ♂ ♀; Nicobars: Little Nicobar; FRI, Dehra Dun. Distribution: Asia (Nicobar Islands in India). References: (ds) Schedl 1972f: 106. (tx) Beeson 1937: 86–87; Schedl 1939c: 335, 1972f: 106.
- lacordairei* Chapuis 1865: 85. Syntypes 2 ♂; Nouvelle-Guinée; BMNH, London (Schedl 1960c: 11). Distribution: Bismarck Islands, Indonesia (Aru), New Guinea, New Ireland. References: (ds) Browne 1966: 241; Gemminger & Harold 1872: 2697; Nobuchi 1977: 106; Ohno, Yoneyama, & Nakazawa 1987a: 95; Ohno, Yoshioka, et al. 1989: 64; Schedl 1936g: 514, 1972f: 106; Strohmeier 1914c: 35. (tx) Chapuis 1865: 85; Lucas 1920: 209; Schedl 1937d: 34–35, 1960c: 11, 1969b: 222, 1972f: 106; Strohmeier 1914c: 35.
- laevigatus* Chapuis 1865: 70. Syntypes 2 ♀; Îles Molucques (Batchian); BMNH, London. Distribution: Indonesia (Moluccas). References: (ds) Gemminger & Harold 1872: 2697; Schedl 1972f: 103; Strohmeier 1912c: 4, 1914c: 34. (tx) Chapuis 1865: 70; Schedl 1960c: 9, 1972f: 103; Strohmeier 1912c: 4, 1914c: 34.
- laratensis* Beeson 1937: 56. Syntypes ♂ ♀; Larat Is.; 1 ♂, 3 ♀ in BMNH, London. Distribution: Indonesia (Larat Island), Philippine Islands. References: (ds) Nobuchi 1977: 104, 1980a; Ohno, Yoneyama, & Nakazawa 1987b: 89; Schedl 1966b: 84, 1966g: 34, 1972f: 102. (tx) Beeson 1937: 56–57; Nobuchi 1955: 328; Schedl 1972f: 102.
- lecontei* Chapuis 1865: 60. Syntypes ♂ ♀; Île de Luçon; 2 ♂, 1 ♀ in BMNH, London, 2 ♂, 1 ♀ in IRSNB, Brussels, 1 ♀ in MNB, Berlin (Schedl 1960c: 7). Distribution: Asia (Vietnam), Philippine Islands (Luzon). References: (ds) Baer 1886; Browne 1966: 239–240, 1986c: 663; Gemminger & Harold 1872: 2697; Lacordaire 1866: 389; Mesa 1935: 96;

- Nobuchi 1977: 104, 1979a: 407; Olmo, Yoneyama, & Nakazawa 1987b: 89; Schedl 1935g: 421, 1966b: 55, 1971f: 150, 1972f: 102, 1974c: 262; Strohmeier 1912c: 4, 1912e: 251, 1914c: 33. **(tx)** Beeson 1937: 56–57; Chapuis 1865: 60; Hopkins 1914c: 183, 191; Lacordaire 1866: 389; Nobuchi 1985: 328; Schedl 1933e: 105, 1935n: 639, 1937d: 33, 1939b: 392, 1952b: 365, 1960c: 7, 1972f: 102; Strohmeier 1911b: 17, 1912c: 4, 1912e: 251, 1914c: 33.
- longicollis** Browne 1950b: 649. Holotype ♂; Malaya: Selangor, Kepong; BMNH, London. Distribution: Asia (Malaya). Hosts: *Castanopsis sumatrana*, *Lithocarpus wallichianus*, *Pasania sundaica*. References: **(hb)** Beaver & Browne 1978: 615; Browne 1961c: 221. **(ds)** Beaver & Browne 1978: 615; Browne 1961c: 221; Schedl 1972f: 111. **(tx)** Browne 1950b: 649, 1962c: 213, 1962e: 644; Schedl 1972f: 111.
- longicornis** (Schedl) 1942c: 196 (*Platypus*). Holotype ♂; New-Guinea; Schedl Collection in NHMW, Wien. Distribution: New Guinea. Hosts: *Myristica* sp. Notes: (3) Schedl 1972i: 65 (described female). References: **(ds)** Schedl 1972f: 111. **(tx)** Schedl 1942c: 196, 1972f: 111, 1972i: 65, 1978a: 41.
- machili** Beeson 1937: 76. Syntypes ♂ ♀; Bengal: Darjeeling division, Debrepani, 6000 ft.; FRI, Dehra Dun. Distribution: Asia (Bengal in India). Hosts: *Machilus odoratissima*, *Mallotus roxburghianus*, *Symplocos theaeifolia*. References: **(cn)** Mathur & Singh 1961a: 83. **(ds)** Beeson 1941 (1961: 265); Mathur & Singh 1961a: 83; Schedl 1972f: 104. **(tx)** Beeson 1937: 76–77; Schedl 1972f: 104, 1978a: 42.
- majusculus** Sampson 1924b: 89. Syntypes ♂ ♀; New Guinea, Mamberamo River; Brittenzorg Museum? Distribution: New Guinea. References: **(ds)** Schedl 1936g: 514, 1968f: 535, 1972f: 112. **(tx)** Sampson 1924b: 89; Schedl 1937d: 34, 1940b: 434–435, 1941e: 153, 1942a: 213, 1942c: 165, 201, 1972f: 112, 1978a: 42.
- abdominalis** Schedl 1969b: 220. Holotype ♂; New Guinea: Karamui, Chimbu District; CSIRO, Canberra. Synonymy: Schedl 1972f: 112. References: **(tx)** Schedl 1969b: 220–221, 1972f: 112.
- minax** (Walker) 1856: 286 (*Platypus*). Holotype ♂; Ceylon; BMNH, London. Distribution: Asia (Karnataka, Tamil Nadu in India/Sri Lanka). Hosts: *Anacardium occidentale*, *Buchanania lanzan*, *Butea monosperma*, *Calycopteris floribunda*, *Dalbergia latifolia*, *Eugenia* sp., *Gluta travancorica*, *Hevea brasiliensis*, *Lagerstroemia*

lanceolata, *Mangifera indica*, *Pterocarpus marsupium*, *Salmalia malabarica*, *Semicarpus gardneri*, *Spodias mangifera*, *Swietenia macrophylla*, *Tectona grandis*, *Terminalia arjuna*, *T. bellerica*, *T. paniculata*, *T. tomentosa*, *Theobroma cacao*, *Wormia triquetra*, *Xylocarpa*.

Notes: (1) The Walker 1856 article was not seen by us; this species was also named as new in Walker 1859: 260.

References: **(cn)** Holmes 1947: 3, 109; Kleine 1932a: 310, 401; Mathur & Singh 1960b: 14, 1960c: 4, 1961a: 13, 1961b: 13; Roonwal 1954: 67. **(ec)** Beeson 1921b: 22; Chatterjee & Chatterjee 1951. **(hb)** Beeson 1933: 9; Chatterjee & Chatterjee 1951; Kleine 1932a: 310, 401; Speyer 1923: 22. **(ds)** Beeson 1921b, 1933: 9, 1941 (1961: 254); Bhasin, Roonwal, & Singh 1958; Chapuis 1865: 71–73; Chatterjee & Chatterjee 1951; Gemminger & Harold 1872: 2697; Holmes 1947: 109; Kleine 1932a: 310, 401; Lacordaire 1866: 390; Mathur & Singh 1960b: 14, 1960c, 1961a: 13, 1961b: 13; Roonwal 1954: 67; Schedl 1969c: 55, 1971a: 279, 1971f: 150, 1972f: 100, 1975e: 452; Strohmeier 1912c: 4, 1914c: 34. **(tx)** Beeson 1933: 9; Chapuis 1865: 71; Gardner 1932b; Lacordaire 1866: 390; Lucas 1920: 209; Schedl 1937d: 34, 1939b: 402, 1959a: 515, 1960c: 9, 1972f: 100; Strohmeier 1912c: 4, 1914c: 34; Walker 1856: 286, 1859: 260–261. **(ms)** Lucas 1920: 209.

minusculus Chapuis 1865: 68. Syntypes ♂; Nouvelle-Guinee; 1 in BMNH, London, 1 in IRSNB, Brussels (Schedl 1960c: 8).

Distribution: Asia (Malaya), Bismarck Islands, Bougainville Island, Indonesia (Sumatra), New Guinea, New Ireland.

Hosts: *Terminalia complanata*, *Xylocarpus gangeticum*.

References: **(hb)** Browne 1961c: 224. **(ds)** Browne 1961c: 224, 1981b: 599; Gemminger & Harold 1872: 2697; Ohno, Yoshioka, et al. 1988b: 97, 1989: 65; Schedl 1972f: 103; Strohmeier 1912c: 4, 1914c: 34. **(tx)** Browne 1955: 365; Chapuis 1865: 68; Schedl 1960c: 8, 1972f: 103; Strohmeier 1912c: 4, 1914c: 34.

inornatus Chapuis 1865: 65. Syntypes 2 ♀; Nouvelle-Guinee (Dorey); BMNH, London (Schedl 1960c: 8). Synonymy: Browne 1955: 365.

References: **(ds)** Gemminger & Harold 1872: 2697; Strohmeier 1912c: 4, 1914c: 33. **(tx)** Browne 1955: 365; Chapuis 1865: 11; Schedl 1960c: 8, 1972f: 103; Strohmeier 1912c: 4, 1914c: 33.

minusculus grandis Schedl 1975: 374. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District, New Guinea; Schedl Collection in NHMW, Wien.

Notes: (3) This appears to be no more than a size variant.

References: **(tx)** Schedl 1975f: 374.

- novaeguineensis* Roberts 1989: 90. Holotype ♂; New Guinea: Papua, Bulolo; NHMW, Wien.
References: (tx) Roberts 1989: 90.
- mniszewski Chapuis** 1865: 62. Syntypes ♂ ♀; Îles de la Nouvelle Guinée, d'Arou, de Célebes; 4 ♂, 2 ♀ in BMNH, London, 3 ♂ in IRSNB, Brussels (Schedl 1960c: 7).
Distribution: Asia (Malaya), Australia, Bismarck Islands, Indonesia (Aru, Celebes, Key Island), Fergusson Island, New Britain Island, New Guinea, New Ireland, Philippine Islands.
Hosts: *Antidesma* sp., *Cananga odorata*, *Dracontomelon* sp., *Maniltoa* sp., *Pangium* sp., *Polyalthia* sp., *Pometia pinnata*, *Vitex* sp.
References: (cn) Gray, B. & Wylie 1974; Smith, J. H. 1935: 3. (cc) Smith, J. H. 1935: 3. (hb) Gray, B. 1968: 307; Gray, B. & Wylie 1974. (ds) Browne 1966: 239, 1980a: 372, 1980c: 484, 1984c: 449; Gemminger & Harold 1872: 2697; Gray, B. 1968: 307; Schedl 1936g: 513, 1962i: 73, 1966b: 85, 1972f: 102, 1979a: 159; Smith, J. H. 1935: 3; Strohmeier 1912c: 4, 1914c: 33. (tx) Beeson 1937: 57–58; Browne 1964: 758; Chapuis 1865: 62; Lea 1910: 133; Nobuchi 1985: 328; Sampson 1924b: 89; Schedl 1936e: 43, 1937d: 33, 1939b: 384, 394, 1940b: 434–435, 1941e: 53, 1942c: 164, 1952b: 365, 1955b: 282, 1958i: 215, 1960c: 7, 1969b: 221, 1972f: 102; Strohmeier 1912c: 4, 1914c: 33; Wylie & Yule 1977.
- moluccanus Schedl** 1961c: 74. Holotype ♂; Ternate; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Ternate in Moluccas).
References: (ds) Schedl 1961c: 74. 1965g: 23, 1972f: 102. (tx) Schedl 1961c: 74, 1972f: 102, 1978a: 45.
- motui Schedl** 1973f: 76. Holotype ♂; Pengagl, Camp east slopes Mt. Wilhelm, Chimbu District, 2770 m; AMNH, New York.
Distribution: New Guinea.
References: (tx) Schedl 1973f: 76–77.
- mussooriensis Beeson** 1937: 84. Holotype ♂; United Provinces: Mussoorie, Mackinnon Park, 6500 ft.; FRI, Dehra Dun.
Distribution: Asia (Uttar Pradesh in India).
Hosts: *Cedrela serrata*.
Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *terminatus*).
References: (ds) Beeson 1941 (1961: 254); Bhasin, Roonwal, & Singh 1958. (tx) Beeson 1937: 84–85; Schedl 1939e: 355, 1972f: 107.
- nakazawai Browne** 1986b: 342. Holotype ♂; New Guinea: Fakfak to Nagoya (Japan), imported; BMNH, London.
Distribution: New Guinea.
Hosts: *Pometia* sp.
References: (tx) Browne 1986b: 342.
- nepalensis Schedl** 1973b: 212. Holotype ♂; Nepal, Phulchoki b. Kathmandu; Schedl Collection in NHMW, Wien.
Distribution: Asia (Bhutan/ Nepal).
References: (ds) Schedl 1975c: 384. (tx) Schedl 1973b: 212, 1978a: 46.
- niponicus Blandford** 1894d: 130. Syntypes (many) ♂ ♀; Sapporo and Hakodate in Yezo, also Miyashita, Yuyama, etc., and in Kiushiu, Japan; BMNH, London.
Figures: Nakane et al. 1963: pl. 192, Nakashima 1975: 15, 1979: pl. 1, Nobuchi 1973b: pl. 1.
Distribution: Asia (Japan/ Taiwan).
Hosts: *Acer mono*, *A. palmatum*, *Aesculus turbinata*, *Alnus japonica*, *Carpinus laxiflora*, *C. tschonoskii*, *Castanea crenata*, *Clethra barbinervis*, *Euonymus alatus*, *Euptelea polyandra*, *Fagus crenata*, *F. japonica*, *Fraxinus mandshurica* var. *japonica*, *Kalopanax septemlobus*, *Lindera erythrocarpa*, *Machilus thunbergii*, *Magnolia obovata*, *Phellodendron amurense*, *Pterocarya rhoifolia*, *Quercus mongolica* var. *grosseserrata*, *Q. myrsinaefolia*, *Q. salicina*, *Styrax japonica*, *Tilia japonica*, *Tsuga sieboldii*, *Zelkova serrata*.
Notes: (1) The type of *Genyocerus adustipennis* Motschulsky was examined at Moscow in 1968 and, in all probability, is of this species: because no material was available for comparison, synonymy should be confirmed before it is accepted (SLW).
References: (ay) Nakashima 1971, 1975: 4, 1978. (cn) Anonymous 1980g; Kleine 1932a: 401; Uchida et al. 1958: 181; Yazawa, Higochi, & Machii 1957: 1–56. (cc) Banno, Mikata, & Kodama 1983: 445; Inouye et al. 1955: 118; Nakashima 1971, 1975: 4, 1978, 1979; Nakashima, Goto, & Iizuka 1987. (hb) Inouye et al. 1955: 118; Kleine 1932a: 401; Murayama 1931b: 199; Nakashima 1971, 1975: 7. (ds) Anonymous 1980g; Blandford 1894c; Kleine 1932a: 401; Miwa 1931: 266; Murayama 1925a: 207, 1929a: 678, 1931b: 199, 1934f: 141, 1937c: 580, 1949c: 104, 1950b: 1299, 1954b: 189, 1955: 103; Nakane et al. 1963: 384; Nakashima 1975: 3, 1979; Nobuchi 1967: 25, 1973b: 7; Schedl 1969a: 204, 1972f: 105; Strohmeier 1912c: 6, 1914c: 35. (tx) Beeson 1937: 69; Blandford 1894d: 130; Murayama 1925a: 207, 1925b: 235, 1928a: 287, 1950b: 1299, 1954b: 189, 1955: 103; Nakane et al. 1963: 384, pl. 192; Nakashima 1975: 15, 1979: pl. 1; Niisima 1909: 170; Nobuchi 1973b: 7, pl. 1; Schedl 1934f: 1647, 1969b: 222, 1972f: 105; Strohmeier 1912c: 6, 1914c: 35; Wylie & Yule 1977.
- nitens Chapuis** 1865: 77. Holotype ♂; Île de Sula; BMNH, London.
Distribution: Asia (Malaya), Indonesia (Borneo, Sulu).
Hosts: *Araucaria cunninghamii*, *Canarium australasicum*, *Elacocarpus* sp., *Eugenia gustavoioides*, *Euodia* sp., *Pithecellobium* sp.
References: (hb) Browne 1941. (ds) Browne 1961c: 224, 1984c: 448; Choo & Woo 1983; Gemminger

- & Harold 1872: 2697; Nobuchi 1977: 105; Schedl 1936d: 6, 1966g: 35, 1972f: 109; Strohmeier 1912c: 4, 1914c: 34. **(tx)** Chapuis 1865: 77; Lucas 1920: 209; Schedl 1939b: 395, 1954a: 156, 1960c: 10, 1971c: 357, 1972f: 109, 1978a: 47; Strohmeier 1912c: 5, 1914c: 34.
- nitescens Schedl** 1979a: 162. Holotype ♂: Tarzali, Atherton Sub. District, Queensland, Australia; CSIRO, Canberra.
Distribution: Australia (Queensland).
Hosts: *Araucaria cunninghamii*, *Canarium australasicum*, *Elacocarpus* sp., *Eugenia gustavoides*, *Euodia* sp.
Notes: (3) Schedl 1980b: 157 (described female).
References: **(ds)** Schedl 1950b: 155–157. **(tx)** Schedl 1979a: 162.
- nitidior (Schedl)** 1953b: 129 (*Trachygostus*). Syntypes ♀: Hiep-Hoa, East-Vaico River, 40 km W Saigon, Indo China; Barbier Collection and Schedl Collection in NHMW, Wien.
Distribution: Asia (Vietnam).
Notes: (1) Schedl 1978a: 47 (citation of holotype invalid).
References: **(ds)** Schedl 1972f: 112. **(tx)** Browne 1962e: 644; Schedl 1953b: 129–130, 1963j: 486, 1972f: 112, 1978a: 47.
- nitidulus (Schedl)** 1954a: 156 (*Platypus*). Syntypes ♂ ♀: Java, Mt. Gede, 800 m; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo, Java, Sumatra).
Hosts: *Castanea argentea*, *Garcinia* sp., *Nephelium* sp.
Notes: (1) Schedl 1978a: 47 (citation of holotype invalid). Browne 1962e: 643 (to *Crossotarsus*).
References: **(hb)** Kalshoven 1960c: 44. **(ds)** Browne 1961c: 224; Kalshoven 1960d: 44; Schedl 1964c: 304, 1971c: 370, 1972f: 109. **(tx)** Browne 1962e: 643; Schedl 1954a: 145, 156–157, 1972f: 109, 1978a: 47.
- octocostatus Schedl** 1935b: 398. Syntypes ♂ ♀: Mount Maquilang, Laguna Province, Philippine Islands; F. C. Hadden Collection, and Schedl Collection in NHMW, Wien.
Figures: Schedl 1972f: 109.
Distribution: Indonesia (Borneo, Mentawai), Philippine Islands (Luzon).
Notes: (1) Schedl 1978a: 49 (citation of holotype invalid).
References: **(ds)** Browne 1966: 241; Nobuchi 1977: 107, 1979a: 407; Ohno, Yoneyama, & Nakazawa 1987b: 89; Schedl 1938g: 421, 1966b: 56, 1966g: 34, 1972f: 108. **(tx)** Browne 1962e: 644; Nobuchi 1985: 328; Schedl 1935b: 398–399, 1935j: 2, 1936e: 43, 1939b: 398, 1939e: 334, 1963j: 486, 1972f: 108–109, 1978a: 49.
- oligodontus Roberts** 1979: 82. Holotype ♂: Papua New Guinea: 12 km south-west of Onim Int, Momt Giluwe, 2900 m; BMNH, London.
Distribution: New Guinea.
Hosts: *Quintonia* sp.
References: **(tx)** Roberts 1979: 82.
- palatus Beeson** 1937: 55. Holotype ♂; C. Borneo: Bangunay; BMNH, London.
Distribution: Indonesia (Borneo, Ternate in Moluccas), Philippine Islands.
Notes: (3) Browne 1964: 758 (described female).
References: **(ds)** Browne 1966: 239, 1980c: 485, 1986a: 90; Ohno, Yoshioka, et al. 1989: 65; Schedl 1966b: 86, 1972f: 102. **(tx)** Beeson 1937: 55–56; Browne 1964: 758; Nobuchi 1985: 328; Schedl 1972f: 102.
- paucidentatus Roberts** 1989: 90. Holotype ♂; New Guinea: Papua; NHMW, Wien.
Distribution: New Guinea.
Hosts: *Evodia* sp.
References: **(tx)** Roberts 1989: 90.
- pectinatus Browne** 1966: 240. Holotype ♂; Bismarck Island, New Britain: Yalom, Gazelle Peninsula, 1000 m; UZMC, Copenhagen.
Distribution: Bismarck Island, New Britain Island.
References: **(ds)** Schedl 1972f: 102. **(tx)** Browne 1966: 240; Schedl 1972f: 102.
- piceae Stebbing** 1906: 413. Syntypes ♂ ♀; Tehri Garhwal, North-West Himalayas, India; not found in FRI, Dehra Dun.
Distribution: Asia (Uttar Pradesh in India).
Hosts: *Picea* sp.
References: **(ds)** Schedl 1972f: 112; Strohmeier 1912c: 6, 1914c: 35. **(tx)** Schedl 1972f: 112; Stebbing 1906: 413; Strohmeier 1912c: 6, 1914c: 35.
- piceus Chapuis** 1865: 56. Syntypes ♂ ♀; Moluques et d'Arou: 3 ♂, 2 ♀ in BMNH, London, 1 ♂, 1 ♀ in IRSNB, Brussels (Schedl 1960c: 6).
Distribution: Asia (Taiwan), Indonesia (Ami, Moluccas, Sumatra), New Guinea, Philippine Islands.
References: **(ay)** Strohmeier 1918: 13. **(ds)** Gemminger & Harold 1872: 2697; Miwa 1931: 266; Murayama 1925a: 202, 1929a: 674, 1937e: 582; Nobuchi 1967: 25; Schedl 1966b: 86, 1972f: 102; Strohmeier 1912c: 4, 1914c: 33. **(tx)** Beeson 1937: 51–53; Chapuis 1865: 56; Murayama 1925a: 202, 1925b: 235; Nobuchi 1985: 328; Schedl 1937d: 33, 1960c: 6, 1972f: 102; Strohmeier 1912c: 4, 1914c: 33.
- porcatus Roberts** 1988: 91. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; NHMW, Wien.
Distribution: New Guinea.
Hosts: *Litsa* sp.
References: **(tx)** Roberts 1988: 91.
- prociduus Schedl** 1975g: 222. Holotype ♂; New Guinea, Geelvink Bay, 10 m; Schedl Collection in NHMW, Wien.
Distribution: New Guinea.
References: **(tx)** Schedl 1975g: 222, 1978a: 56.

quadricaudatus (Strohmeyer) 1910e: 131 (*Platypus*). Syntypes 3. ♂ ♀; Nilgiri Hills; 1 in H. L. Andrews Collection, 2 in IPKE, Eberswalde.
Distribution: Asia (Tamil Nadu in India).

Hosts: *Canarium strictum*, *Palaquium elliptica*, *Vateria indica*.

Notes: (1) Schedl 1972f: 100 (to *Crossotarsus*).

References: (en) Mathur & Singh 1961b: 69. (ds) Beeson 1941 (1961: 254); Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1961b: 69; Schedl 1972f: 100; Strohmeyer 1912c: 19, 1914c: 34. (tx) Schedl 1978a: 56, 1972f: 100; Strohmeyer 1910e: 131, 1912c: 19, 1913: 165, 1914c: 34.

rasilis Roberts 1979: 83. Holotype ♂; Papua New Guinea: 8 km south-west of Onim Hut, Mount Giluwe, 2450 m; BMNH, London.

Distribution: New Guinea.

Hosts: *Evodia* sp.

References: (tx) Roberts 1979: 83.

rastellus Beeson 1937: 82. Syntypes ♂ ♀; Philippines; BMNH, London.

Distribution: Philippine Islands.

Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *squamulatus*).

References: (ds) Schedl 1966b: 88, 1972f: 107. (tx) Beeson 1937: 82–83; Nobuchi 1985: 328; Schedl 1972f: 107.

rengetensis Nüsiima & Murayama in Murayama 1925a: 208. Syntypes ♂ ♀; Rengeti, Taichu Prov., Formosa; Murayama Collection in USNM, Washington.

Distribution: Asia (Taiwan).

Hosts: *Cinnamomum* spp., *Cryptocarya chinensis*, *Machilus longipaniculata*, *Melia azedarach*, *Pasania cuspidata*, *Taonabo japonica*, *Tetradenia konishii*.

References: (hb) Murayama 1928b: 26. (ds) Miwa 1931: 266; Murayama 1925a: 208, 1928b: 26, 1929a: 678, 1937c: 580; Nobuchi 1967: 25, 1977: 108; Schedl 1972f: 112, 1975a: 455. (tx) Beeson 1937: 97–98; Murayama 1925a: 208–210, 1925b: 235; Schedl 1934f: 1647, 1972f: 112.

saltator Schedl 1935k: 480. Syntypes ♂ ♀; Luzon, Nueva Vizcaya Province, Bayombong; W. Schultze Collection, and Schedl Collection in NHMW, Wien.

Distribution: Asia (Vietnam), Philippine Islands (Luzon).

Notes: (1) Schedl 1978a: 63 (citation of holotype invalid).

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (ds) Browne 1968a: 133, 1980a: 371, 1980c: 484; Nobuchi 1977: 109; Schedl 1966b: 87, 1969a: 205, 1972f: 112. (tx) Nobuchi 1985: 328; Schedl 1935k: 480, 1954a: 157, 1963j: 482, 1964h: 424, 1972f: 112, 1978a: 63.

parvus Browne 1962c: 213 (*Trachyostus*). Holotype ♂; Philippines: Laguna, Mt. Maquiling;

Forest Products Research Institute, Laguna. Synonymy: Schedl 1972f: 112.

Notes: (1) Browne 1962e: 644 (to *Crossotarsus*).

References: (ds) Olmo, Yoshioka, et al. 1988b; Schedl 1966b: 104. (tx) Browne 1962c: 213, 1962e: 644; Nobuchi 1985: 328; Schedl 1963j: 482, 1964h: 424, 1972f: 112.

saltatorinus (Schedl) 1954a: 157 (*Platypus*). Syntypes ♀; Java, Mt. Gede, 800 m; Schedl Collection in NHMW, Wien, and Kalshoven Collection.

Distribution: Asia (Fujian in China), Indonesia (Java).

Notes: (1) Schedl 1978a: 63 (citation of holotype invalid).

References: (ds) Kalshoven 1960d: 40; Schedl 1972f: 105. (tx) Browne 1962e: 644; Schedl 1954a: 145, 157–158, 1972f: 105, 1978a: 63.

schedli Browne 1962c: 215. Holotype ♀; Malaya: Selangor, Kepong; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Azela palembanica*, *Buchanania sessifolia*, *Canarium littorale*, *Castanopsis sumatrana*, *Coccoceras muticum*, *Diplospora* sp., *Dryobalanops oblongifolia*, *Elateriospermum tapos*, *Endospermum malaccense*, *Hymenaea courbaril*, *Mesua ferrea*, *Parkia speciosa*, *Shorea* spp., *Sterculia macrophylla*, *Styrax benzoin*, *Xanthophyllum* sp., *Xerospermum* sp.

Notes: (3) This series had been previously misidentified as *cinnamatus* by Schedl 1935n: 639).

References: (hb) Browne 1961c: 223. (ds) Browne 1961c: 223, 1980b: 381; Olmo, Yoshioka, et al. 1989: 65; Schedl 1972f: 102. (tx) Browne 1962c: 215; Schedl 1935n: 639, 1972f: 102.

scorpius (Schedl) 1942c: 195 (*Platypus*). Holotype ♂; Neu-Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

Notes: (1) Browne 1962e: 643 (to *Crossotarsus*).

References: (ds) Schedl 1972a: 109. (tx) Browne 1962e: 643; Schedl 1942c: 195, 1972f: 109, 1978a: 64.

semicinctus Schedl 1973e: 93. Holotype ♂; Potogalai logging area, Hoskins, West New Britain District; CSIRO, Canberra.

Distribution: New Britain.

Hosts: *Eugenia* sp., *Octomeles sumatrana*.

References: (tx) Schedl 1973e: 93, 1978a: 65.

serratulus (Browne) 1964: 756 (*Platypus*). Holotype ♂; Philippines; BMNH, London.

Distribution: Philippine Islands.

Notes: (1) Schedl 1972f: 103 (to *Crossotarsus*).

References: (ds) Schedl 1972f: 103. (tx) Browne 1964: 756; Schedl 1972f: 103, 1972p: 158.

simplex Murayama 1925b: 231. Syntypes ♂ ♀; Japan: Ayakita (Miyazaki Prov., Kyushu); Murayama Collection in USNM, Washington.

Figures: Nobuchi 1973b: pl. 1.

Distribution: Asia (Japan/ Korea/ Taiwan).

Hosts: *Ilex oldhami*, *Lindera thunbergii*, *Prunus jamasakura*, *Quercus* spp.

References: (cn) Ishikura 1966; Kleine 1932a: 401; Murayama 1954a: 19; Shiraki 1952. (hb) Kleine 1932a: 401. (ds) Cho 1957, 1963; Choo 1983: 31; Ishikura 1966; Kleine 1932a: 401; Ko 1969: 273; Murayama 1929a: 676, 1929b: 2, 1930b: 24, 1931a, 1934f: 135, 1936a: 140–141, 1937b: 375, 1937c: 551, 1949c: 104, 1950b: 1299, 1954a: 19, 1954b: 206, 1955: 100, 105, 1957a: 37; Nobuchi 1967: 25, 1973b: 8, 1977: 105; Schedl 1972f: 105; Shiraki 1952. (tx) Beeson 1937: 77; Choo 1983: 31; Choo & Woo 1985: 163; Murayama 1925b: 231–232, 235, 1928a: 289–290, 1930b: 24, 28, 31, 1931a: 50, 1936a: 140, 1937b: 375, 1950b: 1299, 1954b: 206, 1955: 100, 105; Nobuchi 1973b: 8, pl. 1; Schedl 1934f: 1647, 1972f: 105.

siporanus Schedl 1936e: 44. Lectotype ♂; Mentawai, Sopora Sereinu; Schedl Collection in NHMW, Wien.

Distribution: Asia (Malaya), Indonesia (Borneo, Mentawai, Sumatra), New Guinea.

Hosts: *Aporosa globifera*, *Baccaura parvifolia*, *Schoutenia mastersii*, *Xerospermum* sp.

Notes: (3) Schedl 1962f: 209 (treated in *Platyptus*). References: (hb) Beaver & Browne 1978: 616; Browne 1961c: 228. (ds) Beaver & Browne 1978: 616; Browne 1961c: 228; Schedl 1972f: 107. (tx) Schedl 1936e: 44–45, 1939e: 335, 1962p: 209, 1964d: 213, 1972e: 107, 1978a: 67.

oblongirosus Beeson 1937: 88. Holotype ♂; F.M.S.: Pahang, Raub; FRI, Dehra Dun. Synonymy: Schedl 1962p: 208.

References: (hb) Browne 1941. (tx) Beeson 1937: 88–89; Schedl 1939c: 335, 1962p: 209, 1972f: 107.

siva Beeson 1937: 85. Holotype ♂; Bengal: Kurseong division, Sivakhola, 3500 ft.; FRI, Dehra Dun, not found in 1981 by SLW [20 paratypes were present].

Distribution: Asia (Bengal in India).

Hosts: *Albizzia stipulata*, *Cryptocarya wrightiana*, *Engelhardtia spicata*.

Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *terminatus*).

References: (cn) Mathur & Singh 1960a: 35; Roonwal 1954: 53. (ds) Beeson 1941 (1961: 256); Mathur & Singh 1960a: 35; Roonwal 1954: 53; Schedl 1972f: 107. (tx) Beeson 1937: 85; Schedl 1939c: 365, 1972f: 107, 1978a: 67.

spectrum Schedl 1975f: 376. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

References: (tx) Schedl 1975f: 376.

spinifer (Sampson) 1928b: 393 (*Platyptus*). Syn-types ♂ [♀?]; Borneo: Quop, Sandakan; BMNH, London.

Distribution: Indonesia (Borneo).

Notes: (1) Schedl 1972f: 192 (Sampson's female belongs to a *Platyptus* species). (3) Schedl 1972f: 105 (to *Crossotarsus*).

References: (ds) Sampson 1928b: 393; Schedl 1972f: 105, 192. (tx) Browne 1955: 365; Sampson 1928b: 393; Schedl 1939c: 4, 1972f: 105, 192.

spinipennis Schedl 1972i: 54. Holotype ♂; Tonolei, Bougainville District, New Guinea; CSIRO, Canberra.

Distribution: Bougainville Island.

References: (ds) Schedl 1972f: 103. (tx) Schedl 1972f: 103, 1972i: 54.

squamulatus Chapuis 1865: 87. Syntypes ♀; Java; 1 in BMNH, London. 1 in IRSNB, Brussels (Schedl 1960c: 12).

Figures: Sampson 1912: 249 (male).

Distribution: Asia (Bangladesh/ Burma/ Andaman Islands, Assam, Bengal in India/ Malaya/ Pakistan/ Thailand/ Tonkin Island in Vietnam and Vietnam), Bismarck Island, Indonesia (Borneo, Java, Sumatra), Philippine Islands.

Hosts: *Acacia decurrens*, *Azelia palembanica*, *Albizzia lebbek*, *Amoora rohutuka*, *Anstroboxus nitidus*, *Bauhinia* sp., *Bouea macrophylla*, *Brackenridgea hookeri*, *Bridelia tomentosa*, *Bruguiera parviflora*, *Calophyllum tetrapterum*, *Carya arborca*, *Castanopsis sumatrana*, *Cinnamomum cecidodaphne*, *Cinnamomum iners*, *Coccoceras muticum*, *Cratoxylon arborescens*, *Diospyros* sp., *Durio griffithii*, *Endospermum malaccense*, *Eugenia bernardii*, *E. grandis*, *Ficus* sp., *Fordia* sp., *Grewia* sp., *Heritiera fomes*, *Koompassia excelsa*, *Lithocarpus wallichianus*, *Melanorrhoea* sp., *Ochanostachys amentacea*, *Odina wodier*, *Palaquium* sp., *Parinari griffithianum*, *Parkia speciosa*, *Pasania oocarpa*, *Pithecellobium* sp., *Ponteria malaccensis*, *Schoutenia accrescens*, *Shorea leprosula*, *S. maxwelliana*, *Sloetia elongata*, *Tectona grandis*, *Whitfordiodendron pubescens*.

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (cn) Beeson 1915: 8–11, 1918: 114–124, 1919: 1–23; Hogue 1966; Kalshoven 1960d: 48; Kleine 1932a: 310, 401; Mathur & Singh 1961b: 94; Roonwal 1954: 89. (cc) Beeson 1921b: 22; Browne 1955b. (hb) Beaver & Browne 1978: 616; Beeson 1941 (1961: 256); Browne 1936a, 1941, 1958b. 1961c: 225–228; Kalshoven 1960c: 44, 1960d: 2; Kleine 1932a: 310, 401; Yunus & Hua 1980: 216. (ds) Beaver & Browne 1975: 304, 1978: 616; Beeson 1916: 106, 1921b: 22, 1937: 77–83, 1941 (1961: 256); Bhasin, Roonwal, & Singh 1958; Browne 1936a, 1961a: 316, 1961c: 225, 1966: 241, 1980b: 350; Gemminger & Harold 1872: 2697; Kalshoven 1960d: 48; Kleine

- 1932a: 310, 401; Mathur & Singh 1961b: 94; Murayama 1934f: 140; Ohno, Yoneyama, Nakazawa 1957b: 89; Roonwal 1954: 89; Sampson 1919: 106; Schedl 1936d: 6, 1936j: 22, 1937f: 15, 1966b: 87, 1971f: 150, 1972f: 107, 1974c: 262; Strohmeier 1914c: 35; Yunus & Hua 1950: 216. **(tx)** Beeson 1937: 79–80; Chapuis 1865: 87; Gardner 1932b; Nobuchi 1985: 328; Sampson 1919: 106; Schedl 1935k: 483, 1937d: 35, 1939e: 334, 1941c: 354, 1942a: 172, 1960c: 11, 1972f: 107; Strohmeier 1914c: 35.
- fragrimentus* Sampson 1912: 249. Holotype ♂; Singapore; BMNH, London. Synonymy: Beeson 1937: 79.
- Notes: (3) Schedl 1978a: 107 (treated as a subspecies of *fractus*).
- References: **(cn)** Kalshoven 1924c. **(hb)** Browne 1935a; Kalshoven 1924c. **(ds)** Browne 1935a; Murayama 1934f: 140; Sampson 1928b: 391; Strohmeier 1914c: 35. **(tx)** Beeson 1937: 79; Sampson 1912: 249–250, 1919: 106, 1928b: 391; Schedl 1933e: 105, 1935k: 483, 1972f: 107, 1978a: 107; Strohmeier 1914c: 35.
- fragrimentus minor* Sampson 1928a: 2. Syntypes ♂; Tjiuta Radja, Medan, Sumatra; BMNH, London. Synonymy: Schedl 1972f: 107.
- References: **(ds)** Sampson 1928a: 2–3. **(tx)** Beeson 1937: 81; Sampson 1928a: 2–3; Schedl 1972f: 107.
- squamulatus squamuloides* Beeson 1937: 79. Syntypes ♂; many localities in Bengal, India; FRI, Dehra Dun. Synonymy: Schedl 1972f: 107.
- References: **(cn)** Mathur & Singh 1961b: 3. **(ds)** Bhasin, Roonwal, & Singh 1958; Mathur & Singh 1961b: 3; Schedl 1969c: 58. **(tx)** Beeson 1937: 79–81; Schedl 1972f: 107, 1978a: 68.
- squamulatus sundri* Beeson 1937: 79. Syntypes ♂; Sunderbans Division, Bengal, India; FRI, Dehra Dun. Synonymy: Schedl 1972f: 107.
- References: **(cn)** Chaudry 1966. **(cc)** Chaudry 1966. **(ds)** Chaudry 1966. **(tx)** Beeson 1937: 79–80; Schedl 1972f: 107, 1978a: 68.
- subopacus* Schedl 1969b: 222. Holotype ♂; New Guinea; Watut Valley, 1200 m, Morobe District; CSIRO, Canberra.
- Distribution: New Guinea.
- Hosts: *Aglaia* sp.
- References: **(ds)** Schedl 1972f: 107. **(tx)** Schedl 1969b: 222, 1972f: 107, 1978a: 71.
- subpellucidus* Lea 1910: 134. Syntypes 2 ♂; Queensland: Cairns, Kuranda; Lea Collection.
- Distribution: Australia (Queensland), New Guinea.
- Hosts: *Pterocymbium beccarii*.
- References: **(cn)** Smith, J. H. 1935: 3. **(cc)** Smith, J. H. 1935: 3. **(ds)** Schedl 1936g: 513, 1964c: 306, 1968f: 535, 1972a: 144, 1972f: 103, 1979a: 159; Smith, J. H. 1935: 3; Strohmeier 1912c: 7, 1914c: 35. **(tx)** Lea 1910: 134; Schedl 1942c: 165, 1950g: 894, 1972f: 103; Strohmeier 1912c: 7, 1914c: 35.
- sumbawanus* Beeson 1937: 50. Syntypes 2 ♂, 1 ♀; Sumbawa: B. Aroe Hassa, 2–5,000, Doherty; BMNH, London.
- Distribution: Sunda Islands (Sumbawa).
- References: **(ds)** Schedl 1972a: 102. **(tx)** Beeson 1937: 50–51; Schedl 1972f: 102.
- terminatus* Chapuis 1865: 83. Holotype ♂; Singapore; BMNH, London (Schedl 1960c: 11).
- Distribution: Asia (Andaman Islands, Bengal, Uttar Pradesh in India/ Malaya/ Sri Lanka/ Taiwan), Indonesia (Borneo, Java), New Guinea.
- Hosts: *Aglaia* sp., *Artocarpus elasticus*, *Calophyllum* sp., *Castanea argentea*, *Castanopsis acuminatissima*, *Cinnamomum iners*, *Cupressus macrocarpa*, *Dipterocarpus borneensis*, *D. cornutus*, *Doonia zeylanica*, *Dryobalanops oblongifolia*, *Elaeocarpus petiolatus*, *Eudiodora* sp., *Eugenia* sp., *Gordonia zeylanica*, *Hevea brasiliensis*, *Lagerstroemia hypoleuca*, *Macaranga* spp., *Marumia nuscosa*, *Parinari griffithianum*, *Pasania aspericupula*, *P. sundaica*, *Pithecellobium* sp., *Quercus* sp., *Santiria griffithii*, *Scaphium* sp., *Semecarpus gardneri*, *Shorea leprosula*, *S. parvifolia*, *Swietenia macrophylla*, *Whitfordiodendron pubescens*, *Wormia triquetra*, *Xanthophyllum affine*.
- Notes: (3) Schedl 1936d: 13 (described female).
- References: **(cn)** Roonwal 1954: 37. **(cc)** Browne 1958b. **(hb)** Browne 1936a, 1941, 1958b, 1961c: 228; Kalshoven 1960c: 45; Yunus & Hua 1950: 216. **(ds)** Beeson 1941 (1961: 257); Browne 1936a, 1961a: 317, 1961c: 228–229, 1984b: 287; Gemminger & Harold 1872: 2697; Kalshoven 1960d: 45; Murayama 1929a: 677, 1934f: 141; Nobuchi 1967: 25, 1979a: 407; Ohno, Yoshioka, et al. 1985b: 97; Roonwal 1954: 37; Schedl 1966b: 88, 1969c: 58, 1972f: 107; Strohmeier 1912c: 6, 1914c: 35; Yunus & Hua 1950: 216. **(tx)** Beeson 1937: 84–86, 88; Chapuis 1865: 83; Murayama 1928: 288; Nobuchi 1984: 328; Sampson 1912: 249; Schedl 1936d: 13–14, 1939e: 335, 358, 1941c: 354, 1942a: 205, 1960c: 11, 1972f: 107, 1978a: 73; Strohmeier 1912c: 6, 1914c: 35.
- venustus* Chapuis 1865: 88. Holotype ♀; Ceylan; BMNH, London.
- Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *terminatus*), Blandford 1895a: 324 (described female).
- References: **(cn)** Mathur & Singh 1960a: 40, 1960b: 20, 1961a: 33; Roonwal 1954: 19. **(hb)** Speyer 1923: 22. **(ds)** Beeson 1941 (1961: 257); Blandford 1895a; Gemminger & Harold 1872: 2697; Mathur & Singh 1960a: 40, 1960b: 20, 1961a: 33; Murayama 1934f: 140; Roonwal 1954: 19; Schedl 1959a: 513; Speyer 1923: 11–23; Strohmeier 1914c: 35. **(tx)** Beeson 1937: 84–88; Blandford 1895a: 324; Chapuis 1865: 88; Sampson 1912; Schedl 1937d: 35, 1939b: 398,

- 1939e: 335, 355, 1942a: 205, 1959a: 513, 1960c: 12, 1972f: 107–108; Strohmeier 1914c: 35.
- venustus venustulus* Beeson 1937: 85. Syntypes, sex²; Ceylon: Mawagankanda, Ratnapura, 500 ft.; FRI, Dehra Dun.
- Notes: (3) Schedl 1942a: 205 (described female), 1972f: 107 (treated as a subspecies of *terminatus*).
- References: (cn) Mathur & Singh 1961a: 33, 1961b: 92. (ds) Beeson 1941 (1961: 257); Mathur & Singh 1961a: 33, 1961b: 92; Schedl 1964c: 305. (tx) Beeson 1937: 85–86; Schedl 1942a: 205, 1959a: 514, 1972f: 107, 1978a: 80.
- nicobaricus* Beeson 1937: 86. Syntypes ♂ ♀; Nicobars: Car Nicobar; FRI, Dehra Dun. Synonymy: Wood 1992a: 91.
- References: (cn) Mathur & Singh 1960c: 3; Roonwal 1954: 67. (ds) Mathur & Singh 1960c: 3; Roonwal 1954: 67; Schedl 1972f: 106. (tx) Beeson 1937: 86–87, 1941 (1961: 254); Schedl 1939e: 335, 1972f: 106; Wood, S. L. 1992a: 91.
- terminatus sedulus* Schedl 1972h: 66. Holotype ♂; Kulolo logging area, Wau, M. District, New Guinea; CSIRO, Canberra.
- Notes: (3) Schedl 1972f: 107 (treated as a subspecies of *terminatus*).
- References: (ds) Browne 1983a: 555; Ohno, Yoshioka, et al. 1989: 65; Schedl 1972f: 107. (tx) Schedl 1972f: 107, 1972h: 66, 1978a: 73.
- toralis* Schedl 1975f: 376. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
- Distribution: New Guinea.
- References: (tx) Schedl 1975f: 376.
- trachypennis* (Schedl) 1942a: 213 (*Trachyostus*). Lectotype ♀; Malaya, Perak, Chenderoh; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 74.
- Figures: Schedl 1972f: 104.
- Distribution: Asia (Malaya).
- Hosts: Anouaceae sp.
- Notes: (1) Browne 1962e: 644 (to *Crossotarsus*).
- References: (ds) Browne 1961c: 220; Schedl 1972f: 105. (tx) Browne 1962e: 644; Schedl 1942a: 213–214, 1963j: 486, 1972f: 104–105, 1978a: 74.
- sampsonii* Browne 1955: 366. Holotype ♂; Malay Peninsula: Kelantan, Temangan; BMNH, London. Synonymy: Schedl 1972f: 105.
- References: (hb) Browne 1961c: 221. (ds) Browne 1955: 366, 1961c: 221. (tx) Schedl 1972f: 105, 1972p: 156, 1978a: 63.
- tribunarius* Schedl 1939e: 358. Holotype ♂; Luzon, Philippine Islands; Schedl Collection in NHMW, Wien.
- Distribution: Philippine Islands (Luzon).
- References: (ds) Schedl 1966b: 89, 1972f: 108. (tx) Nobuchi 1985: 328; Schedl 1939e: 358–359, 1972f: 108, 1978a: 74.
- trigonus* Beeson 1937: 83. Holotype ♂; Damma Is.; BMNH, London.
- Distribution: Indonesia (Damma Island).
- References: (ds) Schedl 1972f: 108. (tx) Beeson 1937: 83–84; Schedl 1972f: 108.
- tumidus* Schedl 1975f: 377. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.
- Distribution: New Guinea.
- References: (tx) Schedl 1975f: 377.
- ursus* Schedl 1935b: 640. Lectotype ♂; British North Borneo: Mt. Kinabalu, Kamborangah, 7000 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 75.
- Distribution: Asia (Bengal in India), Indonesia (Borneo).
- Hosts: *Engelhardtia spicata*.
- Notes: (3) Beeson 1937: 60 (*pseudoniponicus* Sampson, nomen nudum, synonymy).
- References: (ds) Schedl 1972f: 112. (tx) Beeson 1937: 69; Browne 1962e: 644, 1970: 582; Schedl 1935b: 640–641, 1939e: 333, 1960i: 111, 1972f: 112, 1978a: 78.
- deceptus* Beeson 1937: 68. Syntypes ♂; Borneo, "E.M. 8"; BMNH, London. Synonymy: Schedl 1939e: 333.
- References: (tx) Beeson 1937: 68; Browne 1962e: 644; Schedl 1939e: 333, 1972f: 112.
- perceptus* Beeson 1937: 70. Syntypes ♂ ♀; Bengal: Kurseong, division, Sivakhola, 3500 ft.; FRI, Dehra Dun. Synonymy: Schedl 1972f: 112.
- References: (tx) Beeson 1937: 70–71, 1941 (1961: 254); Schedl 1939c: 2, 1958b: 105, 1972f: 2, 112.
- ursulus* Schedl 1960g: 111. Lectotype ♂; British North Borneo: Mt. Kinabalu, Kamborangah, 7000 ft.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 78, automatic.
- Synonymy: Browne 1962e: 644.
- Notes: (1) Schedl transferred *ursus* to *Platypus*, renamed it, then transferred it back to *Crossotarsus* in 1960; because of the date on which this was done, the Code mandates that the original name be restored.
- References: (ds) Schedl 1965g: 24. (tx) Browne 1962e: 644, 1970: 582; Schedl 1960g: 111, 1965g: 24, 1972f: 112, 1978a: 78.
- vafer* Schedl 1972h: 66. Holotype ♂; Buin, Bougainville District; CSIRO, Canberra.
- Distribution: Bougainville Island.
- Hosts: *Heritiera trifolia*, *Theobroma cacao*.
- References: (ds) Schedl 1972f: 108. (tx) Schedl 1972f: 108, 1972h: 66, 1978a: 78.
- ventricornis* Schedl 1972h: 67. Holotype ♂; Warangoi, East New Britain District; CSIRO, Canberra.
- Distribution: New Britain.
- Hosts: *Canarium indicum*.

References: **(ds)** Schedl 1972f: 105. **(tx)** Schedl 1972f: 105, 1972h: 67, 1978a: 79.

ventrispinis Schedl 1969b: 222. Holotype ♂; New Guinea, Kun, 11 miles from Mt. Hagen, W.H.D.; CSIRO, Canberra.

Distribution: New Guinea.

References: **(ds)** Schedl 1972f: 105. **(tx)** Schedl 1969b: 222, 1972f: 105, 1978a: 79.

ventrosus Schedl 1973e: 94. Holotype ♂; Upper Manki logging area, Bulolo, Morobe District; CSIRO, Canberra.

Distribution: New Guinea.

Notes: (1) A lapsus calami occurred in the naming of this species. The abstract that introduces the article gives the correct name *ventrosus*, but the heading at the beginning of the description gives the name as *senicinctus*, a name previously used on p. 93 of that article, and an obvious oversight by the printer and by the author.

References: **(tx)** Schedl 1973e: 94, 1978a: 79.

vicinus Schedl 1975f: 378. Holotype, sex?; Karamui, Chimbu D., New Guinea; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

Hosts: Derebe.

References: **(tx)** Schedl 1975f: 378.

wallacei (Thomson) 1858: 343 (*Platypus*). Syntypes ♂; Borneo; BMNH, London (Schedl 1960c: 5).

Figures: Schedl 1972f: 101.

Distribution: Asia (Burma/ India/ Malaya/ Singapore/ Sri Lanka/ Taiwan/ Vietnam), Indonesia (Borneo, Java, Malacca, Sumatra), New Guinea.

Hosts: *Afzelia palembanica*, *Albizia falcata*, *Artocarpus dadah*, *Bouea macrophylla*, *Calophyllum* sp., *Castanopsis* sp., *Croton argyратum*, *Dryobalanops oblongifolia*, *Elaterospermum tapos*, *Endospermum malaccense*, *Eugenia grandis*, *Fagraea gigantea*, *Gironmiera* sp., *Heritiera javanica*, *Hevea brasiliensis*, *Ixonanthes reticulata*, *Koompassia excelsa*, *Litsea megacarpa*, *Ochanostachys amentacea*, *Parashorea lucida*, *Parinarium griffithianum*, *Parkia speciosa*, *Payena* sp., *Pometia pinnata*, *Pterocymbium* sp., *Scaphium affine*, *Schoutenia accrescens*, *Shorea leprosula*, *Sivietenia macrophylla*, *Whitfordiodendron pubescens*.

Notes: (3) Chapuis 1865: 53 (redescribed both sexes).

References: **(ay)** Menier 1976: 348; Strohmeier 1918: 8. **(bv)** Menier 1976: 348. **(cn)** Browne 1935b, 1949c; Corbett & Gater 1926b: 261; Kalshoven 1924b: 357, 1936a: 188; Kleine 1932a: 310, 401; Supriana et al. 1978; Yunus & Hua 1980: 216. **(ce)** Browne 1958b. **(hb)** Beaver & Browne 1978: 616; Browne 1935a, 1941, 1958b, 1961c: 222; Kalshoven 1960c: 43; Kleine 1932a: 310, 401; Mercer 1982: 210. **(ds)** Beaver & Browne 1975: 305, 1978: 616; Beeson 1961: 257; Browne 1935a, 1938b: 77–86, 1961a: 316, 1961c: 222,

1972b: 28, 1980b: 381; Chapuis 1865: 53; Corbett & Gater 1926b: 261; Gemminger & Harold 1872: 2697; Hagen 1890; Kalshoven 1924b: 357; Kleine 1932a: 310, 401; Lacordaire 1866: 389; Mercer 1982: 210; Miwa 1931: 266; Murayama 1925a: 202, 1929a: 675, 1937c: 582; Nobuchi 1967: 26; Sampson 1919: 105, 1928a: 2, 1928b: 390; Schedl 1937f: 15, 1960c: 5, 1964c: 305, 1965g: 24, 1971c: 367, 1971f: 150, 1972f: 102, 1974c: 262, 1975a: 453; Strohmeier 1911f: 203, 1912c: 4, 1914c: 33; Supriana et al. 1978; Yunus & Hua 1980: 216. **(tx)** Beeson 1937: 51, 1938b: 295; Browne 1972b: 28; Chapuis 1865: 53; Lacordaire 1866: 389; Lucas 1920: 209; Murayama 1925a: 202–203, 1925b: 235; Sampson 1919: 105, 1928a: 2, 1928b: 390; Schedl 1936e: 43, 1937d: 33, 1939b: 402, 1939e: 333, 1960c: 5, 1972f: 101–102; Strohmeier 1911f: 203, 1912c: 4, 1914c: 33; Thomson 1858: 343.

uilmoti (Stebbing) 1911b: 36 (*Platypus*). Syntypes ♂ ♀; published as Tons Valley, Jaunsar, labeled Kathian, Jaunsar, NWH; FRI, Dehra Dun.

Distribution: Asia (Kashmir, Uttar Pradesh in India).

Hosts: *Cedrela serrata*, *Cedrus deodara*, *Pinus roxburghii*, *Populus ciliata*, *Quercus incana*.

References: **(cn)** Kleine 1932a: 310; Roonwal 1954: 59. **(cc)** Beeson 1921b: 21. **(hb)** Kleine 1932a: 310; Stebbing 1911a: 6. **(ds)** Beeson 1921b: 21, 1937: 74–76, 1961: 257; Bhasin, Roonwal, & Singh 1958; Kleine 1932a: 310; Roonwal 1954: 59; Schedl 1969c: 59, 1972f: 105; Strohmeier 1914c: 27. **(tx)** Beeson 1937: 74–76; Schedl 1972f: 105; Stebbing 1911b: (reprint p. 36), 1914: 618; Strohmeier 1914c: 27.

Genus *Carchesiopygus* Schedl

CARCHESIOPYGUS SCHEDL 1939b: 402–403. Type-species: *Crossotarsus wollastoni* Chapuis, original designation.

Notes: (1) This genus, as recognized by Schedl 1939b, 1972f, appears to be a conglomeration of unrelated species, most of which belong elsewhere.

References: **(hb)** Browne 1961c: 218; Kalshoven 1960: 31–50. **(ds)** Kendrick & Molnar 1965: 39–43; Schedl 1972f: 93. **(tx)** Browne 1962e: 647; Schedl 1939b: 402–403, 1963j: 484, 1972f: 93.

acanthurus (Beeson) 1937: 61 (*Crossotarsus*). Holotype ♂; Java: Preanger, G. Tongkoeban Prahoe, 4000–5000 ft.; Buitenzorg Museum.

Distribution: Asia (Malaya), Indonesia (Java).

References: **(ds)** Schedl 1972f: 95. **(tx)** Beeson 1937: 61; Browne 1962e: 647; Schedl 1972f: 95.

alternantes Schedl 1971c: 397. Holotype ♂; Borneo; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Borneo).

References: **(ds)** Schedl 1971c: 369, 1972f: 95. **(tx)** Schedl 1971c: 369, 397–398, 1972f: 95, 1978a: 10.

assamensis (Beeson) 1937: 59 (*Crossotarsus*). Syntypes ♂ ♀; Assam: Lakhimpur division, Upper Dihing reserve; FRI, Dehra Dun, not found in 1981 by SIW [1 ♂ paratype in FRI].

Distribution: Asia (Assam in India).

Hosts: *Vatica lanceaefolia*.

Notes: (1) Browne 1962e: 647 (to *Carchesiopygus*).

References: (cn) Mathur & Singh 1961b: 70. (ds) Beeson 1941 (1961: 253); Mathur & Singh 1961b: 70; Schedl 1969c: 59, 1972f: 95. (tx) Beeson 1937: 59–60; Browne 1962e: 647; Schedl 1972f: 95.

dentipennis Schedl 1964i: 246. Holotype ♂; Somersby, New South Wales; Schedl Collection in NHMW, Wien.

Distribution: Australia (New South Wales).

Hosts: *Hakea sericca*.

References: (ds) Schedl 1972f: 95. (tx) Schedl 1964i: 246, 1972f: 95, 1978a: 24.

impariporus (Beeson) 1937: 62 (*Crossotarsus*). Syntypes ♂; Assam: Lakhimpur division, Mokum Reserve, Upper Dihing reserve; FRI, Dehra Dun, not found in 1981 by SIW [paratypes were present]. Distribution: Asia (Assam in India/ Vietnam), Indonesia (Java).

Hosts: *Dipterocarpus pilosus*, *Mallotus albus*, *Mesua ferrea*.

Notes: (1) Browne 1962a: 647 (to *Carchesiopygus*).

References: (ds) Beeson 1941 (1961: 265); Schedl 1965a: 340, 1971c: 362, 1972f: 95, 1974c: 262. (tx) Beeson 1937: 62–63; Browne 1962e: 647; Schedl 1939e: 333, 1962n: 647, 1972f: 95.

lobacanthus Schedl 1969c: 59. Holotype ♂: India: Kausani, Almora; FRI, Dehra Dun.

Distribution: Asia (India).

Hosts: *Quercus incana*.

Notes: (3) Beeson 1961: 254 (*lobacanthus*, nomen nudum, synonymy in Schedl 1972f: 95).

References: (ds) Beeson 1941: 330, 1961: 254; Schedl 1972f: 95. (tx) Browne 1962e: 647; Schedl 1963j: 485, 1969c: 59–60, 1972f: 95.

multidentatus (Strohmeyer) 1913: 163 (*Crossotarsus*). Syntypes ♂ ♀; Preanger, Java; Strohmeyer Collection.

Figures: Schedl 1972f: 93.

Distribution: Asia (Malaya), Indonesia (Java).

Hosts: *Dalbergia latifolia*.

References: (ds) Strohmeyer 1914c: 34. (tx) Schedl 1963j: 483, 1972f: 93, 95; Strohmeyer 1913: 163–165, 1914c: 34.

oculatus (Beeson) 1937: 62 (*Crossotarsus*). Holotype ♀; United Provinces: West Almora, Kumaon [India]; FRI, Dehra Dun.

Distribution: Asia (Uttar Pradesh in India).

References: (ds) Schedl 1972f: 95. (tx) Beeson 1937: 62; Browne 1962e: 647; Schedl 1972f: 95.

psilicanthurus (Beeson) 1937: 60 (*Crossotarsus*). Holotype ♂; Bengal: Kalimpong division. Samsingh, 1800 ft.; FRI, Dehra Dun.

Distribution: Asia (Bengal in India).

Hosts: *Cedrela serrata*.

References: (ds) Beeson 1941 (1961: 254); Schedl 1972f: 95. (tx) Beeson 1937: 60; Browne 1962e: 647; Schedl 1972f: 95.

icollastoni (Chapuis) 1865: 74 (*Crossotarsus*). Syntypes 3 ♂; Borneo (Sarawak); BMNH, London (Schedl 1960c: 7).

Distribution: Asia (Assam in India/ Malaya/ Vietnam), Indonesia (Sarawak in Borneo).

Hosts: *Dyera costulata*, *Vatica lanceaefolia*.

Notes: (3) Schedl identified a Beeson male of this species as *Crossotarsus assamensis* Beeson; synonymy? (SIW).

References: (cn) Mathur & Singh 1961b: 70. (cc) Batra 1971. (hb) Browne 1936a, 1941; Kalshoven 1960c: 42. (ds) Beeson 1937: 58–59, 1941 (1961: 257); Browne 1936a, 1961c: 218; Mathur & Singh 1961b: 70; Sampson 1919: 106, 1925b: 390; Schedl 1936d: 5, 1971c: 363, 1972f: 95, 1974c: 262; Strohmeyer 1912c: 5, 1914c: 34. (tx) Beeson 1937: 58–59; Chapuis 1865: 25, 74–75; Sampson 1919: 106, 1925b: 390; Schedl 1939b: 394, 403, 1960c: 9, 1972f: 95; Strohmeyer 1912c: 5, 1914c: 34.

Genus *Crossotarsinulus* Schedl

CROSSOTARSINULUS SCHEDL 1972: 84–87. Type-species: *Crossotarsus sauteri* Strohmeyer, original designation.

References: (tx) Schedl 1972f: 84–87.

sauteri (Strohmeyer) 1913: 164 (*Crossotarsus*). Syntypes ♂ ♀; Kosempo, Formosa; SMTD, Dresden, and IPKE, Eberswalde.

Distribution: Asia (Taiwan).

References: (ds) Murayama 1925a: 207–208, 1929a: 676; Nobuchi 1967: 25; Schedl 1972f: 87; Strohmeyer 1914c: 35. (tx) Murayama 1925a: 207–208, 1925b: 235; Schedl 1972f: 85–87; Strohmeyer 1913: 164, 1914c: 35.

Genus *Spathicranuloides* Schedl

SPATHICRANULOIDES SCHEDL 1972h: 71. Type-species: *Spathicranuloides moikui* Schedl, original designation.

References: (ds) Schedl 1972f: 135–137. (tx) Schedl 1972f: 135–137, 1972h: 71–72.

Spathicranuloides Schedl 1972f: 135. Type-species: *Spathicranuloides moikui* Schedl, original designation, preoccupied by Schedl 1972h.

References: (tx) Schedl 1972f: 135–137.

moikui Schedl 1972h: 72. Holotype ♂; Tonolei, Bougainville District; CSIRO, Canberra.

Figures: Schedl 1972f: 135.

Distribution: Bougainville Island.

References: (tx) Schedl 1972f: 135, 1972h: 71–72.

Genus *Trachyostus* Schedl

TRACHYOSTUS SCHEDL 1939b: 397, 400. Type-species: *Platypus schaufussi* Strohmeier, subsequent designation by Browne 1962e: 646.

Keys: Browne 1962e: 646 (adults), Roberts 1962a: 29 (larvae).

References: (ay) Nobuchi 1969a: 69. (hb) Browne 1960: 213, 1961c: 218–221; Roberts 1968: 325–352; Schedl 1962k: 698, 1972f: 87, 1977b: 253. (ds) Schedl 1962k: 698, 1966b: 104, 1972f: 87, 1977b: 253. (tx) Browne 1962e: 646, 1972a: 184; Roberts 1962a: 29, 1962b: 241–244; Schedl 1939b: 397, 400, 1939p: 258, 1952i: 7, 1957d: 225, 1962h: 65, 1962k: 698, 1963j: 483, 1964b: 425, 1972f: 87, 1977b: 253.

antongilis Schedl 1961e: 166. Holotype ♂; Madagascar, S. Baie Antongil, Humblot; MNHN, Paris. Figures: Schedl 1972f: 86.

Distribution: Madagascar.

References: (ds) Schedl 1972f: 88, 1977b: 254. (tx) Schedl 1961e: 166, 1972f: 88, 1977b: 254, 1978a: 12.

ater (Strohmeier) 1911g: 230 (*Platypus*). Syntypes 3 ♂; Madagascar (Antongil Bai); IPKE, Eberswalde.

Distribution: Madagascar.

Notes: (1) Schedl 1972f: 88 (to *Trachyostus*). References: (ds) Schedl 1972f: 88; Strohmeier 1912c: 14, 1914c: 26. (tx) Schedl 1955i: 213, 1972f: 88; Strohmeier 1911g: 230–231, 1912c: 14, 1914c: 26.

aterrimus (Schaufuss) 1897b: 107 (*Platypus*). Syntypes ♂; Gabon; IPKE, Eberswalde.

Figures: Mayne & Donis 1962: 291.

Distribution: Africa (Cameroon/ Congo/ Gabon/ Ghana/ Kenya/ Nigeria/ Sierra Leone/ Tanzania/ Uganda/ Zaïre).

Hosts: Schedl 1962k: 700–702 (45 host species listed).

References: (ay) Roberts 1962a: 699. (bv) Roberts 1962a: 699. (cn) Kudler 1978: 16; Roberts 1962a: 699. (cc) Akanbi 1978: 123; Roberts 1969a; Schedl 1962k: 699. (hb) Gardner 1957: 29; Roberts 1961: 8, 1962a: 29, 1967; Schedl 1962k: 699; Strohmeier 1911a: 182. (ds) Collart 1934: 248; Mayne & Donis 1960: 100, 1962: 29; Roberts 1962a: 699, 1968: 189, 1969b: 120; Schedl 1933f: 193–194, 1962k: 699, 1964f: 619, 1966c: 230, 1971g: 196, 1972f: 88; Strohmeier 1911a: 182, 1912c: 14, 1914c: 26; Thompson, G. H. 1963: 69. (tx) Mayne & Donis 1962: 291; Roberts 1962a; Sampson 1924c: 130; Schaufuss 1897b: 107–108; Schedl 1933f: 193–194, 1937d: 40, 1941d: 381, 1941e: 154, 1950d: 5, 1953g: 243, 1954e: 59, 1962k: 699, 1964j: 49, 1972f: 88; Strohmeier 1911a: 182, 1912c: 14, 1913: 161, 1914c: 26.

vastus Strohmeier 1911g: 229 (*Platypus*). Holotype ♂; Kamerun; Strohmeier Collection. Synonymy: Schedl 1972f: 88.

References: (tx) Schedl 1972f: 88; Strohmeier 1911g: 229.

aterrimus minor Roberts 1965: 225. Holotype ♂; river valleys south of Jos Plateau, N. Nigeria; BMNH, London, preoccupied by Schedl 1962.

References: (hb) Roberts 1967. (ds) Roberts 1968: 189; Schedl 1972a: 88. (tx) Roberts 1965: 225; Schedl 1972a: 88.

carinatus Schedl 1962h: 73. Holotype ♂; South Nigeria, Sapele; BMNH, London.

Distribution: Africa (Ghana/ Nigeria).

Hosts: *Ficus asperifolia*.

References: (hb) Roberts 1967. (ds) Roberts 1968: 189, 1969b: 120; Schedl 1962h: 63, 1962k: 710, 1972f: 89. (tx) Schedl 1962h: 73, 1962k: 710, 1972f: 89, 1978a: 18.

decellei Schedl 1962k: 710. Holotype ♂; Congo Belge; Yangambi; Schedl Collection in NHMW, Wien.

Figures: Schedl 1962k: 711 (male).

Distribution: Africa (Zaïre).

Hosts: *Scorodophloeus zeukeri*.

References: (ds) Schedl 1962k: 710, 1972f: 89. (tx) Schedl 1962k: 711, 1972f: 89, 1978a: 23.

fecundus (Sampson) 1924c: 130 (*Platypus*). Holotype ♂; Belgian Congo: Haut Uele and Moto; MRCB, Tervuren.

Distribution: Africa (Uganda/ Zaïre).

Hosts: *Albizia* spp.

Notes: (1) Schedl 1972f: 89 (to *Trachyostus*).

References: (ds) Gardner 1957a: 30; Schedl 1933f: 195, 1962k: 712, 1972f: 89. (tx) Sampson 1924c: 130–131; Schedl 1933f: 195, 1955i: 213, 1962k: 712, 1972f: 89.

fecundus ovalis Schedl 1962k: 713. Holotype ♂; Congo Belge; Yangambi; Schedl Collection in NHMW, Wien.

Notes: (3) This appears to be an aberration.

References: (ds) Mayne & Donis 1960: 100, 1962: 290; Schedl 1972f: 89. (tx) Schedl 1962k: 713, 1972f: 89, 1978a: 30.

fecundus transversus Schedl 1962k: 713. Holotype ♂; Congo Belge; Mayidi; Schedl Collection in NHMW, Wien.

Notes: (3) This appears to be an aberration.

References: (ds) Schedl 1962k: 713, 1972f: 89. (tx) Schedl 1972f: 89, 1978a: 30.

ghanaensis Schedl 1959b: 235. Holotype ♂; Gold Coast, Wiawso; BMNH, London.

Figures: Roberts 1960: pls. 1, 3–4.

Distribution: Africa (Ghana/ Guinea/ Ivory Coast/ Liberia).

Hosts: *Triplochiton scleroxylon*.

References: (ay) Roberts 1961c; Schedl 1962k: 713. (bv) Bletchly 1961: 16; Jones 1960c: 231; Neumann & Harris 1974: 136; Roberts 1960a, 1960b, 1960i, 1962b: 241. (cn) Atuahene 1970; Jones 1960a: 93–94, 1960d: 3; Jones et al. 1959: 10; Kudler 1978: 15; Roberts 1960a, 1960b, 1960i,

- 1966a; Thompson, G. H. 1959b: 420. (**cc**) Browne 1961e: 7–5; Roberts 1960a, 1960b, 1960i: 5, 1961e: 8–34, 1962a: 29–40, 1962b: 241–244. (**hb**) Browne 1961e: 7, 1965b; Duffy 1956: 198–200; Jones et al. 1959: 10; Roberts 1960a: 10, 1960b, 1960i: 5, 1961a, 1962b: 241–244, 1967; Schedl 1972k: 713; Thompson, G. H. 1963: 69. (**ds**) Bletchly 1961: 16; Browne 1960a; Duffy 1956: 198–200; Roberts 1960a: 10, 1960b, 1960i: 5, 1961b: 32, 1966a; Schedl 1962k: 713, 1972e: 286, 1972f: 89; Thompson, G. H. 1959b: 420, 1963: 69. (**tx**) Roberts 1960a, 1960i: pls. 1, 3–4; Schedl 1959b: 235, 1962c: 244, 1962k: 713, 1972f: 89, 1978a: 32.
- interstitialis** Schedl 1964j: 49. Holotype ♂: Ghana: Bibiani: BMNH, London.
Distribution: Africa (Ghana).
References: (**hb**) Roberts 1962a: 33, 1967. (**ds**) Schedl 1972f: 89. (**tx**) Schedl 1962k: 727, 1964j: 49, 1972f: 89, 1978a: 38.
- nobilis** (Schaufuss) 1897a: 221 (*Platytypus*). Syntypes 3 ♂; Madagascar: Diego Suarez; Hamburg Museum, lost.
Distribution: Madagascar.
References: (**ds**) Alluaud 1900: 441; Schedl 1972f: 89, 1977b: 254; Strohmeier 1912c: 16, 1914c: 27. (**tx**) Nunberg 1961b: 625, 1964a: 237; Schaufuss 1897a: 221–222; Schedl 1964j: 49, 1967f: 149, 1972f: 89, 1977b: 254; Strohmeier 1912c: 16, 1914c: 27.
- punctatus** Strohmeier 1911g: 229. Syntypes 2 ♂, 2 ♀; Madagascar: Diego Suarez; Strohmeier Collection. Synonymy: Schedl 1962p: 210.
References: (**tx**) Schedl 1933f: 195, 1952i: 22, 1962p: 210, 1972f: 89; Strohmeier 1911g: 229–230.
- schaufussi** (Strohmeier) 1913: 161 (*Platytypus*). Syntypes ♂ ♀; Kilimanjaro; Strohmeier Collection. Figures: Bright 1972d: 19, Nunberg 1961b: 632, Schedl 1962k: 798, 1972f: 86.
Distribution: Africa (Cameroon/ Congo/ Equatorial Guinea/ Gabon/ Ghana/ Ivory Coast/ Kenya/ Malawi/ Nigeria/ Sierra Leone/ Tanzania/ Togo/ Uganda/ Zaïre), Antilles Islands (introduced: Jamaica).
Hosts: Schedl 1962k: 730–732 (51 host species listed), 1962k: 735–738 (53 host species listed).
References: (**ay**) Menier 1976: 348; Strohmeier 1918: 34. (**bv**) Menier 1976: 348. (**en**) Thompson, G. H. 1960; Webb & Jones 1958: 383. (**cc**) Roberts 1969a. (**hb**) Beaver & Loytyniemi 1955b: 117; Thompson, G. H. 1963: 69; Webb & Jones 1958: 383. (**ds**) Beaver & Loytyniemi 1955b: 117; Bright 1972d: 14; Browne 1961: 8, 1972c: 101; Gardner 1957a: 30; Jones, Roberts, & Baker 1959: 13–14, 23, 36, 55; Lee 1971: 30; Mayne & Donis 1960: 100, 1962: 290; Roberts 1960b: 36, 40, 1960e, 1960f, 1962a: 33, 1968: 189; Schedl 1933f: 195, 1938d: 400, 452, 1972f: 89, 1979b: 417; Strohmeier 1914c: 27; Thompson, G. H. 1960e, 1963: 69. (**tx**) Bright 1972d: 14, 19; Browne 1962e: 646; Nunberg 1961b: 632; Sampson 1924c: 131; Schedl 1933f: 195, 1937d: 40, 1939b: 394, 400–401, 1939p: 288, 1941d: 381, 1941e: 154, 1950e: 212, 1953g: 243, 1954d: 874, 1954e: 48, 59, 1962h: 65, 1962k: 798, 1962n: 698, 1964h: 425, 1972f: 86, 89; Strohmeier 1913: 161, 1914c: 27.
- ugandensis** Nunberg 1961b: 624 (*Platytypus*). Holotype ♀; Uganda, Kampala: BMNH, London. Synonymy: Schedl 1972f: 89.
References: (**tx**) Nunberg 1961b: 624, 632; Schedl 1962h: 65, 1962k: 718, 1972f: 89.
- schaufussi major** Schedl 1962k: 734. Holotype ♂; Congo Belge: Kasai, Makumbi; Schedl Collection in NHMW, Wien.
References: (**ds**) Schedl 1972f: 89. (**tx**) Schedl 1962k: 734, 1972f: 89, 1978a: 64.
- schaufussi minor** Schedl 1962k: 735. Syntypes ♂; Gold Coast: Koforidua; Schedl Collection in NHMW, Wien.
References: (**hb**) Roberts 1967. (**ds**) Mayne & Donis 1960: 100, 1962: 292; Roberts 1968: 189, 1969b: 121; Schedl 1967e: 221, 1972e: 286, 1972f: 89; Thompson, G. H. 1963: 69; Webb & Jones 1957: 37, 41. (**tx**) Schedl 1954e: 59, 1962k: 738, 1972f: 89, 1978a: 64.
- schaufussi medius** Schedl 1962k: 734. Lectotype ♂; Gold Coast, Sunyani; Schedl Collection in NHMW, Wien.
References: (**bv**) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**hb**) Loytyniemi, Beaver, & Loytyniemi 1955: 29; Roberts 1967; Schedl 1962k: 734. (**ds**) Mayne & Donis 1960: 100, 1962: 292; Roberts 1962: 33–35, 1968: 189, 1969b: 121; Schedl 1962h: 63, 1964f: 619, 1971f: 196, 1972e: 286, 1972f: 89; Webb & Jones 1957: 36, 39, 41. (**tx**) Schedl 1952j: 6, 1954e: 59, 1962k: 734, 1972f: 89, 1978a: 64.
- schoutedeni** (Schedl) 1933f: 195 (*Platytypus*). Syntypes ♂; Mongwalu (Kilo), Kwamouth, Sankuru: Komi; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: Schedl 1962k: 740–742 (20 host species listed).
Notes: (1) Schedl 1978a: 64 (type restricted to Tervuren specimens).
References: (**cc**) Schedl 1962k: 740. (**hb**) Schedl 1962k: 740. (**ds**) Mayne & Donis 1960: 100, 1962: 292; Schedl 1933f: 195, 1962k: 740, 1972f: 89. (**tx**) Schedl 1933f: 195, 1962k: 740, 1964j: 49, 1972f: 89, 1978b: 64.
- similis** Nunberg 1967b: 330. Holotype ♀; Elisabethville; MRCB, Tervuren.
Figures: Nunberg 1967: pl. 5, fig. 4.
Distribution: Africa (Zaire).
References: (**tx**) Nunberg 1967b: 330, 339.
- tomentosus** (Strohmeier) 1911g: 231 (*Platytypus*). Holotype ♂; Kamerun; MNB, Berlin.

Figures: Schedl 1962k: 743.

Distribution: Africa (Cameroon/ Ghana/ Nigeria/ Togo/ Zaïre).

Hosts: *Scorodophloeus zenkeri*.

Notes: (3) Schedl 1962h: 73 (described female).

References: (**hb**) Roberts 1967. (**ds**) Roberts 1962a: 33, 1968: 189, 1969b: 121; Schedl 1962h: 63, 1972k: 742, 1972f: 89; Strohmeier 1912c: 19, 1914c: 29. (**tx**) Schedl 1962h: 73, 1962k: 742–743, 1972f: 89, 1978a: 74; Strohmeier 1911g: 231–232, 1912c: 19, 1914c: 29.

Genus *Neotrachyostus* Browne

NEOTRACHYOSTUS BROWNE 1962e: 645. Type-species:

Platypus abbreviatus Chapuis, original designation.

References: (**ds**) Schedl 1972f: 90. (**tx**) Browne 1962e: 645; Schedl 1972f: 90.

abbreviatus (Chapuis) 1865: 28, 35, 106 (*Platypus*). Syntypes ♂ ♀; Bresil; 2 ♂, 1 ♀ in BMNH, London, 2 ♂ in IRSNB, Brussels (Schedl 1960c: 13). Distribution: South America (Brazil).

Notes: (3) Browne 1962e: 645 (to *Neotrachyostus*). References: (**ay**) Menier 1976: 348. (**bv**) Menier 1976: 348. (**ds**) Blackwelder 1947: 788; Schedl 1933c: 166, 1972f: 92, 1976a: 57; Strohmeier 1912c: 8, 1914c: 22. (**tx**) Browne 1962e: 645; Chapuis 1865: 28, 35, 106; Lucas 1920: 521; Schedl 1933c: 166, 1937b: 399, 1939p: 288, 1960c: 13, 1972f: 92; Strohmeier 1912c: 8, 1914c: 22. (**ms**) Lucas 1920: 521.

abdominalis (Schedl) 1936i: 99 (*Platypus*). Holotype ♂; Brasil. Nova Tentonia; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

Notes: (3) Schedl 1972f: 92 (to *Neotrachyostus*, described female).

References: (**ds**) Blackwelder 1947: 788; Schedl 1972g: 51. (**tx**) Schedl 1936i: 99–100, 1972f: 92, 1972g: 51, 1978a: 9.

acutidens (Blandford) 1895e: 96 (*Platypus*). Syntypes 4, ♂ ♀; Guatemala, Capetillo; BMNH, London.

Distribution: North America (Guatemala).

Notes: (1) Browne 1962e: 646 (to *Neotrachyostus*). References: (**ds**) Blackwelder 1947: 788; Equihua & Atkinson 1987: 14; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (**tx**) Blandford 1895e: 93–96; Browne 1962e: 646; Equihua & Atkinson 1987: 14; Schedl 1933a: 938, 1939b: 399, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

concarvus (Chapuis) 1865: 108 (*Platypus*). Holotype ♀; Bresil; BMNH, London (Schedl 1960c: 13).

Distribution: South America (Brazil).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*). References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 22. (**tx**) Browne 1962e: 645; Chapuis 1865: 108–109; Schedl 1960c: 13, 1972f: 92; Strohmeier 1912c: 8, 1914c: 22.

digitalis (Chapuis) 1865: 111 (*Platypus*). Syntypes 4 ♂; Bresil meridional, Nova Fribourg; IRSNB, Brussels (Schedl 1960c: 14).

Distribution: South America (Brazil).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*).

References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 22. (**tx**) Browne 1962e: 645; Chapuis 1865: 111; Reichardt 1964: 85; Schedl 1960c: 14, 1972f: 92; Strohmeier 1912c: 8, 1914c: 22.

distinctus (Chapuis) 1865: 111 (*Platypus*). Holotype ♂; Para.; IRSNB, Brussels (Schedl 1960c: 14). Distribution: South America (Bolivia/ Brazil).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*).

References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 22. (**tx**) Browne 1962e: 645; Chapuis 1865: 111–112; Schedl 1960c: 14, 1972f: 92; Strohmeier 1912c: 8, 1914c: 22.

dohrni (Chapuis) 1865: 115 (*Platypus*). Holotype ♂; Mexico; not found at BMNH, London, or IRSNB, Brussels (Schedl 1960c: 15).

Distribution: North America (Mexico).

Notes: (1) Browne 1962e: 646 (to *Neotrachyostus*). References: (**ds**) Blackwelder 1947: 789; Blandford 1895b: 96; Equihua & Atkinson 1987: 14; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (**tx**) Blandford 1895b: 96; Browne 1962e: 646; Chapuis 1865: 115–116; Equihua & Atkinson 1987: 14; Schedl 1939b: 399, 1939p: 288–289, 1940a: 323, 1960c: 15, 1962e: 646, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

filiformis (Chapuis) 1865: 116 (*Platypus*). Syntypes 2 ♀; Mexico; IRSNB, Brussels (Schedl 1960c: 15). Distribution: North America (Mexico).

References: (**ds**) Blackwelder 1947: 789; Blandford 1896e; Equihua & Atkinson 1987: 14; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (**tx**) Blandford 1896e: 97; Browne 1962e: 646; Chapuis 1865: 116–117; Equihua & Atkinson 1987: 14; Schedl 1940a: 323, 1960c: 15, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

fuscifrons (Chapuis) 1865: 110 (*Platypus*). Holotype ♀; Bresil (Ega); BMNH, London (Schedl 1960c: 14).

Distribution: South America (Brazil).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*). References: (**ds**) Blackwelder 1947: 789; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (**tx**) Browne 1962e: 645; Chapuis 1865: 110–111; Schedl 1960c: 14, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

lobatus (Chapuis) 1865: 109 (*Platypus*). Syntypes ♂; Cayenne; 1 in MNB, Berlin, 2 in IRSNB, Brussels, 1 in BMNH, London (Schedl 1960c: 14). Distribution: South America (Brazil/ Cayenne/ Venezuela).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*).

References: (ds) Blackwelder 1947: 789; Schedl 1960a: 78, 1972f: 92, 1976a: 57; Strohmeier 1912c: 8, 1914c: 23. (tx) Browne 1962e: 645; Chapuis 1865: 109; Schedl 1937d: 37, 1952a: 443, 1960c: 13–14, 1972f: 90, 92; Strohmeier 1912c: 8, 1914c: 23.

obliquus Wood 1966b: 49. Holotype ♂: Volcan, Puntarenas Prov., Costa Rica; Wood Collection. Figures: Wood 1966b: 47 (male). Distribution: North America (Costa Rica). References: (ds) Equihua & Atkinson 1987: 14; Schedl 1972f: 92. (tx) Equihua & Atkinson 1987: 14; Schedl 1972f: 92; Wood, S. L. 1966b: 47, 49–50.

pacificus (Chapuis) 1865: 118 (*Platypus*). Holotype ♀; Cayenne; IRSNB, Brussels (Schedl 1960c: 15). Distribution: South America (Cayenne/ Venezuela).

References: (ds) Blackwelder 1947: 790; Schedl 1933c: 170, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (tx) Browne 1962e: 646; Chapuis 1865: 118–119; Nunberg 1939: 241; Schedl 1960c: 15, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

putzeysi (Chapuis) 1865: 28, 117 (*Platypus*). Syntypes 2 ♀; Mexico, Texpam, Oaxaca; BMNH, London (Schedl 1960c: 15).

Distribution: North America (Guatemala/ Oaxaca, Veracruz in Mexico).

Notes: (1) Browne 1962e: 646 (to *Trachyostus*). References: (ds) Blackwelder 1947: 790; Blandford 1896e: 97; Equihua & Atkinson 1987: 14; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (tx) Blandford 1897e: 97; Browne 1962e: 646; Chapuis 1865: 117–118; Equihua & Atkinson 1987: 14; Schedl 1940a: 323, 1960c: 15, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23; Wood, S. L. 1966.

quadrilobus (Blandford) 1895b: 95 (*Platypus*). Syntypes 4 ♂; Costa Rica, Irazu; BMNH, London. Figures: Schedl 1962g: 1036.

Distribution: North America (Costa Rica).

Hosts: Logs.

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*). (3) This species does not belong to this genus (SLW).

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 14; Schedl 1933c: 166, 1972f: 92; Spahr 1981: 79; Strohmeier 1912c: 8, 1914c: 23. (tx) Blandford 1895b: 93, 95–96; Browne 1962e: 645; Equihua & Atkinson 1987: 14; Nunberg 1939: 241; Schedl 1962g: 1036, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

springi (Chapuis) 1865: 112 (*Platypus*). Syntypes 1 ♂, 1 ♀; Colombie; IRSNB, Brussels (Schedl 1960c: 14).

Distribution: South America (Colombia).

Notes: (1) Browne 1962e: 645 (to *Neotrachyostus*). References: (ds) Blackwelder 1947; Schedl 1972f: 92; Strohmeier 1912c: 8, 1914c: 23. (tx) Browne 1962e: 645; Chapuis 1865: 112–114; Schedl 1960c: 14, 1972f: 92; Strohmeier 1912c: 8, 1914c: 23.

Genus *Platyscapulus* Schedl

PLATYSCAPULUS SCHEDL 1957d: 125. Type-species: *Platypus carinulatus* Chapuis, automatic, subsequent designation by Wood 1992b: 90.

Platyscapus Schedl 1939b: 397, 399. Type-species: *Platypus carinulatus* Chapuis, subsequent designation by Wood 1990b: 90, preoccupied by Huistache 1921.

References: (tx) Browne 1962e: 652; Kalshoven 1960: 31–50; Schedl 1939b: 397, 399, 1939k: 718, 1957d: 125; Wood, S. L. 1979a: 1, 1992b: 90.

Costaroplatus Nunberg 1963c: 109. Type-species: *Costaroplatus shenefelti* Nunberg, original designation. Synonymy: Wood 1979a: 1.

References: (tx) Nunberg 1963c: 109; Schedl 1969f: 555; Wood, S. L. 1979a: 1, 1992b: 90.

Notes: (3) A generic revision of Platypodidae is in progress; the species listed below are transferred into *Platyscapulus* in that study (SLW).

References: (tx) Browne 1961c: 212–213, 1962e: 652; Schedl 1957d: 125; Wood, S. L. 1979a: 1, 1992b: 90.

abditulus (Wood) 1966b: 50 (*Platypus*). Holotype ♂; near Moravia, Cartago Prov., Costa Rica, 1500 feet; Wood Collection.

Figures: Wood 1966b: figs. 5–6.

Distribution: North America (Costa Rica).

References: (ds) Equihua & Atkinson 1982: 18; Schedl 1972f: 235. (tx) Equihua & Atkinson 1987: 18; Schedl 1972f: 235; Wood, S. L. 1966b: 47–50.

abditus (Schedl) 1936c: 246 (*Platypus*). Holotype ♂; Brazil; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 788. (tx) Schedl 1936c: 246, 1972f: 235; Wood, S. L. 1966b: 50.

transitus Schedl 1978b: 309. Holotype ♂; Brasilien, Linhares, E. Santo; Schedl Collection in NHMW, Wien. Synonymy: Wood 1992b: 90.

References: (ds) Schedl 1972f: 235. (tx) Schedl 1936c: 246, 1972f: 235, 1978b: 309; Wood, S. L. 1992b: 90.

carinulatus (Chapuis) 1865: 231 (*Platypus*). Syntypes 1 ♂, 1 ♀; Bresil, Ega; BMNH, London.

Distribution: South America (Brazil/ Cayenne/ Suriname).

Notes: (3) Schedl 1976a: 87 (described female).

References: (ds) Blackwelder 1947: 789; Gemminger & Harold 1872: 2697; Reichardt 1964: 145; Schedl 1960a: 79, 1966f: 92, 1970e: 86, 1972f: 235, 1972g: 47; Strohmeier 1912c: 14, 1914c: 26. (tx) Chapuis 1865: 135; Lucas 1920: 522; Schedl 1960a: 79, 1960c: 43, 1972f: 235, 1976a: 87; Strohmeier 1912c: 14, 1914c: 26. (ms) Lucas 1920: 522.

costellatus (Schedl) 1933c: 175 (*Platypus*). Holotype ♂; Brazil, Ypiranga; Schedl Collection in NHMW, Wien.

Distribution: South America (Brazil).

References: (ds) Blackwelder 1947: 789; Schedl 1933c: 175, 1966f: 92, 1970e: 86, 1972f: 235, 1972g: 47, 1976a: 57. (tx) Reichardt 1962: 333, 335; Schedl 1933c: 175, 1962q: 493, 1972f: 235, 1978a: 22.

luederwaldti Schedl 1933c: 177 (*Platypus*). Holotype ♀; Brazil, Ypiranga; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1962q: 493.

References: (ds) Blackwelder 1947: 789; Schedl 1933c: 177. (tx) Reichardt 1962q: 333; Schedl 1933c: 177, 1962q: 493, 1972f: 235, 1978a: 41.

calix Schedl 1934e: 212 (*Platypus*). Syntypes ♀; Brasilien: Alto da Serra, 700 m; Hamburg Museum, lost, and Schedl Collection in NHMW, Wien. Synonymy: Schedl 1962q: 493. Notes: (1) Schedl 1978a: 17–18 (citation of holotype invalid).

References: (ds) Blackwelder 1947: 789. (tx) Schedl 1934e: 212, 1962q: 493. 1972f: 235, 1978a: 17.

frontalis (Blandford) 1896e: 113 (*Platypus*). Holotype ♀; Bugaba, Panama; BMNH, London.

Figures: Blandford 1896e: pl. 4, fig. 25.

Distribution: North America (Guatemala/ Panama), South America (Brazil).

References: (ds) Blackwelder 1947: 789; Equihua & Atkinson 1987: 23; Schedl 1933c: 177, 1966f: 77, 1972f: 235, 1972g: 48, 1976a: 57; Strohmeier 1912c: 19, 1914c: 29. (tx) Blandford 1896e: 113; Equihua & Atkinson 1987: 23; Schedl 1972f: 235; Strohmeier 1912c: 19, 1914c: 29.

manus (Schedl) 1936c: 247 (*Platypus*). Holotype ♂; Brazil; Schedl Collection in NHMW, Wien.

Figures: Reichardt 1962: 335.

Distribution: South America (Bolivia/ Brazil/ Guyana).

References: (ds) Blackwelder 1947: 790; Schedl 1960a: 79, 1970f: 582, 1972f: 235, 1973a: 369. (tx) Reichardt 1962: 335–336; Schedl 1936c: 247, 1972f: 235, 1978a: 42.

pulchellus (Chapuis) 1865: 230 (*Platypus*). Holotype ♀; de Mexico; IRSNB, Brussels (Schedl 1960c: 43).

Distribution: North America (Belize/ Costa Rica/ Guatemala/ Honduras/ Chiapas, Nayarit, Veracruz in Mexico/ Panama), South America (Brazil/ Guyana/ Suriname/ Venezuela).

Hosts: *Gossypiospermum praecox*.

References: (ce) Equihua & Atkinson 1986: 625. (hb) Equihua & Atkinson 1986: 625. (ds) Atkinson & Equihua 1986a: 423, 1988: 104; Blackwelder 1947: 790; Blandford 1896i: 112; Cola 1973; Equihua 1988: 144; Equihua & Atkinson 1986: 625, 1987: 27; Estrada & Atkinson 1985:

213; Ferrer 1942; Gemminger & Harold 1872: 2700; Nunberg 1963c: 107; Reichardt 1965b: 160; Schedl 1933c: 174, 1960a: 78, 1963c: 160, 1972f: 235; Strohmeier 1912c: 14, 1914c: 26. (tx) Blandford 1895b: 93–95, 1896e: 112; Chapuis 1865: 230; Equihua & Atkinson 1987: 27; Schedl 1936c: 247, 1939b: 399, 1940a: 327, 1956b: 30, 1960c: 43, 1972f: 235; Strohmeier 1912c: 14, 1914c: 26.

pulcher (Chapuis) 1865: 249 (*Platypus*). Holotype ♀; du Bresil; BMNH, London.

Distribution: North America (Mexico), South America (Brazil/ Venezuela).

Notes: (3) Schedl 1933a: 401 (described male).

References: (ds) Blackwelder 1947: 790; Equihua & Atkinson 1987: 27; Ferrer 1942; Gemminger & Harold 1872: 2700; Schedl 1933c: 177, 1972f: 235; Strohmeier 1912c: 16, 1914c: 27. (tx) Chapuis 1865: 249; Equihua & Atkinson 1987: 27; Schedl 1933a: 401, 1936c: 234, 1937d: 40, 1940a: 327, 1941e: 156, 1960c: 47, 1972f: 235; Strohmeier 1912c: 16, 1914c: 27.

pusillimus (Chapuis) 1865: 232 (*Platypus*). Holotype ♂; du Bresil, Bahia; IRSNB, Brussels (Schedl 1960c: 43).

Distribution: South America (Brazil/ Guyana).

References: (ds) Blackwelder 1947: 790; Gemminger & Harold 1872: 2700; Schedl 1972f: 235, 1972g: 50, 1976a: 58; Strohmeier 1912c: 14, 1914c: 26. (tx) Chapuis 1865: 232; Schedl 1959m: 556, 1960c: 43, 1972f: 235, 1978a: 59; Strohmeier 1912c: 14, 1914c: 26.

shenefelti (Nunberg) 1963c: 110 (*Costaroplatus*). Holotype ♀; Costa Rica; University of Wisconsin Collection, Madison.

Figures: Nunberg 1963c: 100

Distribution: North America (Costa Rica).

Notes: (1) Schedl 1969f: 555 (to costellati group in *Platypus*).

References: (ds) Equihua & Atkinson 1987: 29; Schedl 1972f: 235. (tx) Equihua & Atkinson 1987: 29; Nunberg 1963c: 100, 110; Schedl 1969f: 555, 1972f: 235.

subabditus (Schedl) 1935i: 47 (*Platypus*). Lectotype ♀; Guyane française, Saint-Laurent du Maroni; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 69.

Distribution: South America (Cayenne).

Notes: (3) Schedl 1935i: 47 (described female).

References: (ds) Schedl 1972f: 235. (tx) Schedl 1935i: 47, 1972f: 235, 1978a: 69.

turgifrons (Schedl) 1935h: 358 (*Platypus*). Holotype ♀; Equador; Schedl Collection in NHMW, Wien.

Distribution: South America (Ecuador).

References: (ds) Blackwelder 1947: 791; Schedl 1972f: 235. (tx) Schedl 1935h: 358–359, 1972f: 235, 1978a: 76.

umbrosus (Schedl) 1936c: 247 (*Platypus*). Lectotype ♂: Guyane Française: St. Laurent du Maroni, St. Jean du Maroni, Nouveau Chantier; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 77.

Distribution: South America (Cayenne).

References: (ds) Blackvelder 1947: 791; Schedl 1972f: 235, 552. (tx) Schedl 1936c: 247, 1972f: 235, 1978a: 77.

Genus *Baiocis* Browne

BAIOCIS BROWNE 1962e: 651. Type-species: *Crossotarsus peruanulus* Schedl, original designation. References: (ds) Schedl 1972f: 140. (tx) Browne 1962e: 652; Schedl 1972f: 140–142.

anaticeps (Schedl) 1969a: 215 (*Platypus*). Holotype ♂: Borneo (Kalimantan), Nunukan to Muroan (Japan), imported; PPST, Tokyo.

Distribution: Indonesia (Borneo).

Hosts: Meranti log.

References: (ds) Nobuchi 1977: 142; Ohno, Yoneyama, & Nakazawa 1987b: 89; Schedl 1972f: 142. (tx) Nobuchi 1985: 330; Schedl 1969a: 215–216, 1972f: 142.

angusticeps (Schedl) 1942a: 203 (*Platypus*). Syn-types ♀: Malaya, Selangor, Ulu Gombak; BMNH, London.

Distribution: Asia (Malaya).

Hosts: *Shorea faguetiana*.

Notes: (1) Schedl 1978a: 11 (citation of holotype invalid), Browne 1962e: 651 (to *Baiocis*).

References: (ds) Browne 1961c: 211; Schedl 1972f: 197. (tx) Browne 1962e: 651; Schedl 1942a: 203, 1972f: 197, 1978a: 11.

angustiformis (Schedl) 1942a: 202 (*Platypus*). Lectotype ♂: Malaya, Rotan Tunggal F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 11.

Distribution: Asia (Malaya).

Hosts: Anonaceae sp.

Notes: (1) Browne 1962e: 651 (to *Baiocis*).

References: (ds) Browne 1961c: 211; Roberts 1968: 192; Schedl 1972f: 142. (tx) Browne 1962e: 651; Schedl 1942a: 202, 1972f: 142, 1978a: 11.

angustipennis (Schedl) 1942a: 203 (*Platypus*). Lectotype ♂: Malaya, Perak, Trolak F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 11.

Distribution: Asia (Malaya).

Hosts: *Xylopia caudata*.

Notes: (1) Browne 1962e: 651 (to *Baiocis*).

References: (ds) Browne 1961c: 211; Schedl 1971f: 142. (tx) Browne 1962e: 651; Schedl 1942a: 203, 1972f: 142, 1978a: 11.

angustulus (Schedl) 1942a: 202 (*Platypus*). Syn-types ♀: Teibodas, Giava; Schedl Collection in NHMW, Wien.

Distribution: Indonesia (Teibodas, Giava).

References: (ds) Schedl 1972f: 142. (tx) Browne

1962e: 651; Schedl 1942a: 202, 1972f: 142, 1978a: 11.

annularis (Schedl) 1975f: 379. Holotype ♂: Upper Manki logging area, Bulolo, Morobe District; Schedl Collection in NHMW, Wien.

Distribution: New Guinea.

Hosts: *Syzygium* sp.

Notes: (3) Roberts 1989: 262 (described female).

References: (tx) Roberts 1989: 262; Schedl 1975f: 379.

imitatrix Schedl 1973e: 94. Holotype ♂: Upper Manki logging area, Bulolo, Morobe District, New Guinea; CSIRO, Canberra.

Distribution: New Guinea.

References: (tx) Schedl 1973e: 94, 1978a: 36.

incisus (Sampson) 1928: 1 (*Crossotarsus*). Holotype ♂: Sibajak vulkan [Sumatra]; BMNH, London.

Distribution: Indonesia (Sumatra).

Notes: (1) Browne 1962e: 651 (to *Baiocis*).

References: (ds) Schedl 1972f: 142. (tx) Browne 1962e: 651; Sampson 1928: 1; Schedl 1941c: 359, 1972f: 142.

inimicus (Schedl) 1935k: 482 (*Crossotarsus*).

Lectotype ♂: Luzon, Mountain Province, Benguet, Mount Santo Tomas; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 37.

Distribution: New Britain Island, Philippine Islands (Luzon).

Hosts: *Ficus* sp.

Notes: (1) Schedl 1958k: 152 (erroneously placed this species in *Platypus* and gave it an unneeded new name).

References: (ds) Schedl 1972f: 142. (tx) Browne 1962e: 651; Nobuchi 1985: 330; Schedl 1935k: 482, 1939b: 398, 1941c: 258–259, 1958k: 152, 1972f: 142, 1978a: 37.

velatus Schedl 1958k: 152 (*Platypus*). Lectotype ♂: Luzon, Mountain Province, Benguet, Mount Santo Tomas; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 37, automatic.

Notes: (1) This is an unneeded replacement name that resulted from a taxonomic error in generic placement.

References: (ds) Schedl 1965g: 24, 1966b: 101, 1968f: 537. (tx) Browne 1962e: 652; Schedl 1958k: 152, 1972f: 142.

kuntzeni (Schedl) 1937d: 40 (*Platypus*). Lectotype ♂: Deutsch-Neu-Guinea, Hauptlager, Flusslager, Etappenberg, Maifluss; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 40.

Distribution: New Guinea.

Notes: (1) Browne 1983a: 555 (to *Baiocis*).

References: (en) Wylie & Shanahan 1975. (cc) Roberts 1980. (hb) Wylie & Shanahan 1975. (ds) Browne 1983a: 555; Schedl 1936g: 513, 1968e: 264, 1972f: 103. (tx) Browne 1983a: 555; Schedl

- 1936d: 13, 1937d: 40–41, 1940b: 434–435, 1941e: 153, 1955b: 282, 1972f: 103, 1978a: 40.
- umbilosus Roberts & Morimoto** 1986: 162. Holotype ♂; New Guinea: Papua, Gumi, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Garcinia* sp.
References: (tx) Roberts & Morimoto 1986: 162.
- pasohensis (Schedl)** 1939e: 362 (*Platypus*). Syntypes ♂; Malaya, Negri Sembilan: Pasoh For. Res.; BMNH, London and Schedl Collection in NHMW, Wien.
Distribution: Asia (Malaya).
Hosts: *Shorea lacvis*.
Notes: (1) Schedl 1978a: 51 (intended to designate a lectotype, but cited the wrong publication; this invalidated the designation). (3) Schedl 1942a: 203 (described female).
References: (cn) Mathur & Singh 1961a: 39. (ds) Beeson 1941 (1961: 266); Browne 1961c: 211; Mathur & Singh 1961a: 39; Schedl 1972f: 198. (tx) Browne 1962e: 652; Schedl 1939e: 362, 1942a: 203, 1972f: 198, 1978a: 51.
- perangustus (Schedl)** 1942a: 202 (*Platypus*). Lectotype ♂; Malaya, Rotan Tunggal F.R.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53.
Distribution: Asia (Malaya).
References: (ds) Browne 1961c: 212; Schedl 1972f: 200. (tx) Browne 1962e: 652; Schedl 1942a: 202, 1972f: 200, 1978b: 52.
- perinimicus (Schedl)** 1941c: 358 (*Platypus*). Lectotype ♂; Java, Mount Gede, Kalshoven Nr. 20; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53.
Distribution: Indonesia (Java, Sumatra).
Hosts: *Castanea argentea*, *C. javanica*, *Cinnamomum camphora*, *Eurya japonica*.
References: (ds) Kalshoven 1960: 40; Schedl 1972f: 142. (tx) Schedl 1941c: 358–359, 1942a: 203, 1954a: 145, 1961c: 70, 1972f: 142, 1978a: 53.
- pernanulus (Schedl)** 1935k: 482 (*Crossotarsus*). Lectotype ♂; Luzon, Laguna Province, Mount Maquiling, 400 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 54.
Distribution: Asia (India/ Malaya/ Thailand), Australia (Queensland), Indonesia (Borneo, Java), New Guinea, Philippine Islands (Luzon).
Hosts: *Artocarpus lakoocha*, *Castanopsis sumatrana*, *Dryobalanops aromatica*, *Lophopetalum* sp., *Nephelium* sp., *Pometia* sp., *Shorea leprosula*.
Notes: (1) Browne 1962e: 652 (to *Baiocis*).
References: (hb) Browne 1961c: 211. (ds) Beaver & Browne 1975: 302; Beeson 1941 (1961: 266); Browne 1961c: 211, 1962c: 202, 1980a: 371, 1983a: 554, 1984c: 450; Ohno, Yoneyama, & Nakazawa 1987a: 95, 1987b: 89; Schedl 1936g: 514, 1966b: 96, 1969c: 61, 1972f: 143, 1980b: 195. (tx) Browne 1962e: 652; Nobuchi 1955: 330; Schedl 1935k: 482, 1939e: 336, 362, 1942c: 165, 1955i: 214, 1969a: 215, 1972f: 143, 1978a: 54.
- seminitens (Schedl)** 1971c: 395 (*Platypus*). Holotype ♂; India; Schedl Collection in NHMW, Wien.
Distribution: Asia (India).
Notes: (1) Schedl 1978a: 65 (to *Baiocis*).
References: (ds) Schedl 1972f: 143. (tx) Schedl 1971c: 395–396, 1972f: 143, 1978a: 65.
- solomonicus Browne** 1986b: 335. Holotype ♂; Solomon Islands: Lever Harbor to Nagoya (Japan); BMNH, London.
Distribution: Solomon Islands.
Hosts: *Beilschmiedia* sp.
References: (tx) Browne 1986b: 335.
- sublunaris (Schedl)** 1937e: 544 (*Crossotarsus*). Syntypes ♂ ♀; S.O. Borneo; Schedl Collection in NHMW, Wien.
Distribution: Indonesia (Borneo).
Notes: (1) Schedl 1978a: 71 (citation of holotype invalid, to *Baiocis*).
References: (ds) Schedl 1972f: 142–143. (tx) Schedl 1937e: 544, 1972f: 142–143.
- unispinosus Roberts** 1986: 39. Holotype ♂; New Guinea: Papua, Bulolo; BMNH, London.
Distribution: New Guinea.
Hosts: *Xanthophyllum papuanum*.
References: (tx) Roberts 1986: 39.
- variolosus (Schedl)** 1942a: 201 (*Platypus*). Lectotype ♀; Malaya, Perak, Larut Hills, 4500 feet; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 79.
Distribution: Asia (Malaya).
References: (ds) Browne 1961c: 212; Schedl 1971c: 367, 1972f: 242. (tx) Browne 1962e: 652; Schedl 1942a: 201, 1972f: 242, 1978a: 79.

Genus *Cylindropalpus* Strohmeier

- CYLINDROPLATYPUS STROHMEIER 1911a: 174. Type-species: *Cylindropalpus africanus* Strohmeier, monobasic.
References: (hb) Schedl 1958d: 195, 1962k: 959, 1972f: 131, 1977b: 255. (ds) Schedl 1962k: 959, 1972f: 131, 1977b: 255. (tx) Browne 1955: 364, 1962e: 650, 655, 1972a: 183; Hopkins 1914: 120, 132, 1915c: 227; Sampson 1924: 133; Schedl 1935g: 320, 1939b: 397–399, 1941d: 416, 1952i: 7, 1952j: 6, 1957d: 125, 1961e: 157, 1962k: 959, 1972f: 131, 1977b: 255; Strohmeier 1911a: 174, 1912c: 21, 1914c: 12–13, 17–20, 29–30, 1918: 1–46.
- affinis Strohmeier** 1912f: 80. Holotype ♂; West-Usambara; Strohmeier Collection.
Distribution: Africa (Chana/ Tanzania/ Uganda).
Hosts: *Aningeria altissima*, *A. robusta*, *Chrysophyllum metallicum*, *C. natalensis*, *C. perpulchrum*, *Scorodophloeus zenkeri*.
References: (hb) Browne 1963a: 249; Gardner 1957: 30. (ds) Browne 1963a: 249; Mayne & Donis 1960: 98, 1962: 280; Schedl 1962k: 978,

1972f: 133; Strohmeier 1912c: 21, 1914c: 30. (tx) Schedl 1941d: 416, 1950d: 14, 1952j: 6, 1957d: 125, 1962k: 978, 1972f: 133; Strohmeier 1912c: 21, 1912f: 80, 1914c: 30.

auricomans (Schaufuss) 1897b: 106 (*Platyptus*). Holotype ♂; Kamerun; IPKE, Eberswalde.

Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Nigeria/ Sierra Leone/ Togo/ Zaïre/ Zambia).

Hosts: Browne 1963a: 249, Schedl 1962k: 969–971 (many hosts listed).

Notes: (3) Strohmeier 1911a: 174 (described female).

References: (an) Roberts 1961. (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (cn) Cachan 1958; Fougereusse 1957. (cc) Akanbi 1978: 123; Cachan 1957: 1–126, 1958: 393–398; Roberts 1969a; Schedl 1962k: 967; Schimitschek 1964c. (hb) Beaver & Loytyniemi 1985b: 115; Browne 1961b: 23, 1962d: 94, 1963a: 249; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Roberts 1961: 31; Schedl 1962k: 967; Thompson, G. H. 1963: 59. (ds) Beaver & Loytyniemi 1985b: 115; Bletchly 1961: 15; Browne 1963a: 249, 1980c: 486, 1984b: 228; Jones, Roberts, & Baker 1959: 13–14, 44–50, 54; Mayne & Donis 1951: 355, 1960: 98, 1962: 192; Roberts 1968: 192, 1969: 104; Schedl 1933f: 195, 1937b: 407, 1937d: 43, 1938d: 452, 1950d: 14, 1951f: 40, 1961g: 22, 1962h: 61, 1962k: 967, 1120, 1965e: 357, 1965g: 22, 1972f: 133; Strohmeier 1912c: 19, 1914c: 28; Thompson, G. H. 1963: 59. (tx) Hopkins 1914: 120, 132; Sampson 1924: 133; Schaufuss 1897b: 106–107; Schedl 1933f: 195, 1935g: 320, 1938d: 452, 1939b: 398, 1941d: 416, 1951f: 40, 1954d: 874, 1954e: 48, 51, 59, 71–72, 1957b: 152, 1957d: 125, 1962k: 967, 1965g: 22, 1972f: 133; Strohmeier 1911a: 174, 1912c: 19, 21, 1912f: 80, 1914c: 28.

africanus Strohmeier 1911e: 174. Syntypes: Kamerun: 2♂, 1♀ in MNB, Berlin, 1♂, 1♀ in IPKE, Eberswalde. Synonymy: Schedl 1962k: 967, 1972f: 133.

References: (cn) Mayne & Donis 1951: 335. (cc) Mayne & Donis 1951: 335. (ds) Mayne & Donis 1962: 266; Strohmeier 1912c: 21, 1914c: 30. (tx) Hopkins 1914: 120; Sampson 1924c: 133; Schedl 1933f: 195, 1935g: 319–320, 1937b: 407, 1937d: 42–43, 1950d: 14, 1972f: 133; Strohmeier 1911e: 174, 1912c: 21, 1912f: 80, 1914c: 30.

caliginosus Roberts 1966b: 619. Holotype ♂; Nigeria: Gyl Nyaki, Mambilla Plateau, Sardauna Province, over 5000 feet; BMNH, London.

Distribution: Africa (Nigeria).

References: (ds) Roberts 1968: 192, 1969: 104; Schedl 1972f: 134. (tx) Roberts 1966b: 619, 1970a: 65; Schedl 1970i: 221–231, 1972f: 134.

camerunus (Schedl) 1941d: 417 (*Platyscapus*). Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Nigeria/ Sierra Leone).

Hosts: *Afzelia* spp., *Anthonia* spp., *Baphia nitida*, *Barteria nigritiana*, *Chrysophyllum lacourtiannum*, *Pentaclethra celtideana*, *Terminalia superba*, *Triplochiton scleroxylon*.

Notes: (3) Schedl 1954d: 885 (described female).

References: (cc) Akanbi 1978: 123; Jover 1952: 73–81; Roberts 1969a; Schedl 1962k: 977. (hb) Browne 1962d: 94; Jover 1952: 73–81; Schedl 1962k: 973. (ds) Cachan 1957: 15, 43–50; Roberts 1968: 192, 1969b: 104–105; Schedl 1962b: 61, 1962k: 977, 1965e: 357, 1971g: 194, 1972f: 133, 1975h: 352, 1984b: 620; Thompson, G. H. 1963: 60. (tx) Schedl 1941d: 447, 1950e: 212, 1954d: 874, 885–886, 1954e: 59, 65, 71, 1957d: 125, 1962k: 977, 1121, 1972f: 133, 1978a: 18.

cultellus (Schedl) 1953d: 105 (*Platyscapus*). Lectotype ♀; Madagascar, Mt. d'Ambre, 1140 m; Schedl Collection in NHMW, Wien.

Distribution: Madagascar.

Notes: (3) Schedl 1961e: 157 (described male).

References: (ds) Schedl 1972f: 133, 1977b: 256. (tx) Schedl 1953d: 105–106, 1961e: 157, 1972f: 133, 1977b: 256, 1978a: 22.

granulosus (Schedl) 1957d: 126 (*Platyscapulus*). Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Ghana/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sierra Leone/ Tanzania/ Uganda/ Zaïre).

Hosts: *Aningeria robusta*, *Anthonia* spp., *Carapa grandiflora*, *Chrysophyllum* spp., *Combretodendron macrophyllum*, *Drypetes leonensis*, *Khaya anthotheca*, *Microcos pinnatifida*, *Pachystela laurentii*, *Randia acuminata*, *Scorodolobos zenkeri*, *Synsepalum* spp., *Tridesmostemon clausenii*.

References: (cc) Schedl 1958d: 197. (ds) Schedl 1962h: 61, 1962k: 979, 1972f: 134. (tx) Schedl 1957d: 125–127, 1962k: 979, 1972f: 134, 1978a: 33.

granulosus subaffinis Schedl 1957d: 126 (*Platyscapulus*). Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

References: (cc) Schedl 1962k: 979. (hb) Schedl 1962k: 979. (ds) Schedl 1959q: 706, 1962k: 979, 1972f: 134. (tx) Schedl 1952j: 6, 1957d: 126–127, 1959q: 706, 1962h: 61, 1962k: 979–980, 1972f: 134, 1978a: 34.

granulosus fratellus Schedl 1957d: 127 (*Platyscapulus*). Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

References: (cc) Schedl 1962k: 983. (hb) Schedl 1962k: 983. (ds) Mayne & Donis 1960: 100, 1962: 288; Schedl 1958d: 196, 1961m: 84, 1962k: 983–984, 1972f: 134. (tx) Schedl 1957d:

127, 1961m: 84, 1962k: 983–984, 1972f: 134, 1978a: 33.

interpositus (Schedl) 1957d: 127 (*Platyscapulus*). Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Ruanda/ Zaïre).

Hosts: *Chrysophyllum africanum*, *C. lacourtianum*, *C. pruniforme*, *Drypetes leonensis*, *Jubbernardia sereti*, *Pachystela laurentii*, *Strombosiaopsis tetrauda*, *Synsepalum longecuneatum*.

References: (cn) Anonymous 1970c: 12. (cc) Schedl 1962k: 841. (hb) Browne 1963a: 260; Schedl 1962k: 841, 989. (ds) Anonymous 1970c: 12; Browne 1963a: 260; Mayne & Donis 1960: 96, 1962: 271; Schedl 1962k: 841, 989, 1964f: 619, 1964j: 43, 1970h: 179, 1972f: 134, 158, 1975k: 279. (tx) Schedl 1957d: 127–128, 1962k: 841, 989, 1972f: 134, 1978a: 38.

laudatus (Schedl) 1941d: 416 (*Platyscapus*). Syn-types ♂ ♀; ♂ from Niala, Congo, ♀ from Ukaika; Schedl Collection in NHMW, Wien.

Figures: Schedl 1962k: 961, 1972f: 131 (male).

Distribution: Africa (Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Tanzania/ Uganda/ Zaïre).

Hosts: Schedl 1962k: 961–964 (many hosts listed). Notes: (1) Schedl 1978a: 40 (citation of male holotype invalid).

References: (cc) Schedl 1962k: 960, 1972c: 59. (hb) Cardner 1957: 30; Schedl 1962k: 960. (ds) Browne 1984b: 288; Cachan 1957: 15; Mayne & Donis 1960: 98, 1962: 285, 288, 1984b: 288; Roberts 1968: 192, 1969: 105; Schedl 1962k: 960, 1965e: 357, 1972f: 134. (tx) Numberg 1961b: 625; Schedl 1941d: 416, 1950e: 211, 1952j: 6, 1954d: 874, 1955f: 260–261, 1957d: 125, 1962k: 960–961, 1962n: 698, 1972f: 134, 1978a: 40.

maximus Browne 1972c: 109. Holotype ♂; Congo-Kinshasa: Kivu, Bitale; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Browne 1972c: 109.

pernix (Schedl) 1941d: 418 (*Platyscapus*). Holotype ♂; St. Thome [Africa]; Schedl Collection in NHMW, Wien.

Distribution: Africa (Sao Tome Island).

References: (cc) Schedl 1972c: 60. (ds) Schedl 1972f: 134. (tx) Numberg 1967b: 329; Schedl 1941d: 418, 1962k: 989, 1972f: 134, 1978a: 54.

pertinax (Schedl) 1941d: 418 (*Platyscapus*). Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Nigeria/ Sierra Leone/ Zaïre).

Hosts: *Chrysophyllum* spp., *Pterocarpus soyauxii*, *Synsepalum subcordatum*, *Trichilia lanata*, *Tridesmostemon claessensii*.

References: (cc) Schedl 1962k: 988. (hb) Schedl 1962k: 988. (ds) Roberts 1968: 192, 1969: 105; Schedl 1957d: 127, 1962k: 988, 1972f: 134. (tx)

Numberg 1968: 329; Schedl 1941d: 418, 1962k: 988, 1121, 1972f: 134, 1978a: 54.

pumilio (Schedl) 1941d: 419 (*Platyscapus*). Holotype ♂; Congo; Schedl Collection in NHMW, Wien.

Distribution: Africa (Ghana/ Ivory Coast/ Nigeria/ Sierra Leone/ Zaïre).

Hosts: *Anopyxis klaineana*, *Chrysophyllum africanum*, *C. lacourtianum*, *Iringia gabonensis*, *Microdesmis puberula*, *Synsepalum longecuneatum*.

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (cc) Schedl 1962k: 922. (hb) Beaver & Loytyniemi 1985b: 115; Browne 1963a: 250; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Schedl 1962k: 922. (ds) Beaver & Loytyniemi 1985b: 115; Browne 1963a: 250; Mayne & Donis 1962: 286; Roberts 1968: 192, 1969: 105; Schedl 1962h: 61, 1962k: 922, 1972f: 134; Thompson, G. H. 1963: 60. (tx) Schedl 1941d: 419–420, 1952i: 22, 1954e: 59, 71, 1957d: 128, 1961m: 84, 1962h: 61, 1962k: 992, 1121, 1972f: 134, 1978a: 59.

pusillus (Schedl) 1952i: 22 (*Platypus*). Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Chrysophyllum africanum*, *Microdesmis puberula*, *Synsepalum subcordatum*.

References: (cc) Schedl 1962k: 991. (hb) Schedl 1962k: 991. (ds) Schedl 1962k: 991, 1972f: 134. (tx) Schedl 1952i: 22, 1957d: 125, 128, 1962k: 991, 1972f: 134, 1978a: 59.

pusillus Schedl 1957d: 128 (*Platyscapulus*). Holotype ♀; Congo Belge: Yangambi; MRCB, Tervuren, preoccupied by Schedl 1952.

Notes: (1) It is obvious that Schedl intended to describe only the female of an established species, but in so doing he inadvertently validated a new homonymous name.

References: (tx) Schedl 1957d: 128.

sulcatulus Browne 1986c: 670. Holotype ♂; Solomon Islands: Lever Harbor to Nagoya (Japan), imported; BMNH, London.

Distribution: Solomon Islands.

Hosts: *Calophyllum* sp., *Celtis* sp. logs.

References: (tx) Browne 1986c: 670.

tenax (Schedl) 1941d: 419 (*Platyscapus*). Holotype ♀; Fernando-Po; Schedl Collection in NHMW, Wien.

Distribution: Africa (Fernando Po Island).

References: (ds) Schedl 1972f: 134. (tx) Schedl 1941d: 419, 1962k: 989, 1972f: 134, 1978a: 72.

Genus *Triozastus* Schedl

TRIOZASTUS SCHEDL 1939b: 402, 403. Type-species: *Crossotarsus banghaasi* Strohmeier, monobasic.

Keys: Browne 1962e: 635, 1971d: 224.

References: (av) Browne 1972a: 182. (hb) Browne 1971d: 224–225; Schedl 1962k: 680, 1972f: 246. (ds) Browne 1971d: 224–225; Schedl

1952g: 54, 1972f: 246. (**tx**) Browne 1962e: 635, 649, 1971d: 224–225, 1972a: 182; Numberg 1953: 44, 1960: 287–305, 1963a: 567–572; Schedl 1936e: 51, 1939b: 397, 400–403, 1939p: 288–289, 1952i: 7, 1962k: 680–681, 1972f: 246.

banghaasi (Schaufuss) 1905: 87 (*Crossotarsus*). Holotype ♂; Dar-es-Salaam, Afr. Or. [Tanzania]; Hamburg Museum, lost.

Figures: Numberg 1963a: 566, Schedl 1962k: 681. Distribution: Africa (Angola/ Cameroon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Nigeria/ South Africa/ Tanzania/ Uganda/ Zaïre/ Zimbabwe).

Hosts: *Albizzia ferruginea*, *A. zygia*, *Antiaris africana*, *Baphia nitida*, *Blighia welwitschii*, *Bosqueia angolensis*, *Piptadeniastrum africanum*, *Sterculia rhinopetala*, *Strombosia pustulata*, *Trachylobium verrucosum*, *Trichilia heudelotii*, *T. lanata*, *T. rubescens*, *Triplochiton scleroxylon*.

Notes: (1) Schedl 1939b: 402 (to *Triozastus*), Browne 1971d: 228 (established subspecies *banghaasi* and *handeniensis*). (3) Schedl 1935: 314 (described female).

References: (**ay**) Menier 1976: 348. (**bv**) Menier 1976: 348. (**ec**) Akanbi 1978: 123; Schedl 1962k: 682. (**hb**) Browne 1963a: 253; Schedl 1962k: 682; Strohmeier 1911a: 182. (**ds**) Beaver & Loytyniemi 1955b: 117; Browne 1963a: 253, 1971d: 227, 1975a: 757; Menier 1973a; Roberts 1968: 189; Schedl 1962k: 682, 1966a: 276, 1971e: 4, 1972e: 285, 1972f: 248; Strohmeier 1911a: 182, 1911f: 204, 1912c: 6, 1914c: 34. (**tx**) Browne 1971d: 227–228; Numberg 1963a: 566, 568; Powell, W. 1987: 28; Schaufuss 1905: 5, 87; Schedl 1933f: 200, 1935e: 153, 1935g: 314–315, 1939b: 402–403, 1961m: 85, 1962k: 681–682, 1970i: 229, 1972f: 248, 1978a: 14; Strohmeier 1912c: 6, 1914c: 34.

bifurcus Schedl 1941d: 412. Holotype ♂; Mayumbe, Congo; Schedl Collection in NHMW, Wien.

Notes: (3) Schedl 1972f: 248 (treated as a subspecies).

References: (**tx**) Schedl 1941d: 411–413, 1962h: 65, 1962k: 697, 1972f: 248, 1978a: 15.

handeniensis Numberg 1960b: 306. Holotype ♂; Tanganyika Terr., Handeni, 350 m; MRCB, Tervuren.

Notes: (3) Browne 1971d: 228 (treated as a subspecies).

References: (**bv**) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**hb**) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**tx**) Browne 1971d: 228–229; Numberg 1960a: 287–308, 1960b: 153–162, 1963a: 569–571; Schedl 1972f: 248.

brevidens Browne 1971d: 226. Holotype ♂; Ghana: Kintampo; MRCB, Tervuren.

Distribution: Africa (Ghana).

References: (**hb**) Beaver & Browne 1975: 302. (**tx**) Browne 1971d: 226–227.

brownei Beaver in Beaver & Loytyniemi 1955: 118. Holotype ♂; Zambia: Chati, Kitwe; BMNH, London.

Distribution: Africa (Zambia).

References: (**bv**) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**hb**) Beaver & Loytyniemi 1955b: 118; Loytyniemi, Beaver, & Loytyniemi 1955: 28. (**ds**) Beaver & Loytyniemi 1955b: 118.

caliginosus Roberts 1966b: 634. Holotype ♂; Nigeria: Boshi Extension F.R.; BMNH, London.

Distribution: Africa (Nigeria/ an apparent separate geographical race in Ghana, Zaïre).

Hosts: *Bosqueia angolensis*, *Deinbollia* sp., *Pausinystalia macroceras*, *Piptadeniastrum africanum*, *Pittosporum viridiflorum*, *Rauwolfia vomitoria*, *Santiria trimera*, *Schefflera mannii*, *Strombosia* spp.

Notes: (1) Browne 1971d: 233 (established subspecies).

References: (**ds**) Roberts 1968: 189, 1969: 122. (**tx**) Roberts 1966b: 364; Schedl 1970i: 229, 1972f: 248.

caliginosus humilis Browne 1971d: 233. Holotype ♂; Congo-Kinshasa: Yangambi; MRCB, Tervuren.

Notes: (1) Distribution: Ghana and Zaïre.

References: (**tx**) Browne 1971d: 233.

elongatus Schedl 1954e: 83. Lectotype ♂; Mpraeso [Ghana]; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 27.

Figures: Numberg 1963a: 570.

Distribution: Africa (Ghana/ Nigeria).

Hosts: *Albizzia zygia*, *Bussea occidentalis*, *Cleistopholis patens*, *Piptadeniastrum africanum*, *Strombosia pustulata*, *Trichilia prieuriana*.

Notes: (3) Browne 1965a: 207 (described female), Browne 1971d: 229 (a good species), Schedl 1972f: 248 (a subspecies of *banghaasi*).

References: (**ds**) Browne 1971d: 229; Roberts 1968: 189, 1969: 122; Thompson, G. H. 1963: 70. (**tx**) Browne 1965a: 207, 1971d: 229; Numberg 1963a: 569–571; Schedl 1954e: 59, 65, 82–85, 1962h: 64, 1962k: 697, 1972f: 248, 1978a: 27.

marshalli (Sampson) 1924c: 128 (*Crossotarsus*). Holotype ♀; S. Africa, Salisbury; BMNH, London.

Figures: Browne 1971d: 219, Numberg 1963a: 566, 570, Schedl 1972f: 246.

Distribution: Africa (Angola/ Cameroon/ Congo/ Dahomey/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Mozambique/ Nigeria/ Ruanda/ Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaïre/ Zambia/ Zimbabwe).

Hosts: *Afromosia elata*, *Azelia cuanzensis*, *Albizzia gummiifera*, *Alstonia boonei*, *Antiaris welwitschii*, *Bosqueia angolensis*, *Celtis mildbraedii*, *Daniellia oliveri*, *Drypetes gossweileri*, *Ekebergia* sp., *Entandrophragma* sp., *Erythrophloeum guineense*, *Gilbertiodendron dewevrei*,

Grevillea sp., *Guarea cedrata*, *G. thompsonii*, *Macaranga* sp., *Ongokoa gore*, *Parkia biglobosa*, *Polyalthia suaveolens*, *Pygeum africanum*, *Scorodophloeos zenkeri*, *Terminalia superba*, *Trachylobium verrucosum*. Browne 1971d, Schedl 1962k: 687–690 (list many additional hosts).

Notes: (3) Browne 1971d: 230 (a good species), Schedl 1972f: 248 (treated as a subspecies of *banghaasi*).

References: (hb) Beaver & Loytyniemi 1985b: 119. (ds) Beaver & Loytyniemi 1985b: 119; Browne 1971d: 230, 1983a: 556; Mayne & Donis 1960: 100, 1962: 292; Roberts 1968: 189, 1969: 122; Schedl 1933f: 200, 1938d: 452, 1975h: 351; Wood, S. L. 1957e: 1272. (tx) Browne 1971d: 219, 230; Nunberg 1963a: 569–570; Sampson 1924c: 128–129; Schedl 1933f: 200, 1935e: 153, 1935g: 314–315, 1936e: 51, 1937d: 33–36, 1939b: 390, 1941d: 413, 1941e: 153–154, 1950d: 5, 12, 1952g: 69, 1953g: 243, 1955d: 270, 1955f: 261, 1955i: 213, 1962k: 684–685, 1109, 1972f: 248.

pertenuis Schedl 1935g: 315 (*Crossotarsus*). Syn-types ♂; French Equatorial Africa: Lambarene; MNHN, Paris and Schedl Collection in NHMW, Wien. Synonymy: Browne 1971d: 230. Notes: (3) Schedl 1939g: 169 (described female), 1972f: 248 (a subspecies of *banghaasi*).

References: (ds) Mayne & Donis 1960: 100, 1962: 293; Roberts 1968: 189, 1969: 122. (tx) Browne 1971d: 230; Schedl 1935g: 315–316, 1939g: 169, 1954e: 82, 1962k: 688, 1109, 1972f: 248, 1978a: 54.

propatulus Schedl 1935e: 153. Lectotype ♂: Deutsch-Afrika: Amami; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 57. Synonymy: Browne 1971d: 230.

Notes: (1) Schedl 1952g: 54 (to *Triozastus*). (3) Schedl 1937d: 35 (described female).

References: (cn) Curry 1958b: 128–132. (cc) Akanbi 1978: 123; Jover 1952: 73–81; Roberts 1969a: 122. (hb) Gardner 1957: 30; Jones, Roberts, & Baker 1959: 13; Jover 1952: 73–81; Webb & Jones 1957: 36–37, 41. (ds) Cachan 1957: 6, 15; Ferreira 1965: 1130; Mayne & Donis 1960: 100, 1962: 293; Nunberg 1960a: 306; Roberts 1961: 39–43, 1968: 189; Schedl 1934a: 247, 1938d: 451, 1959p: 22, 1962h: 64, 1962k: 691, 1966c: 230, 1967e: 221, 1969a: 203, 1971e: 2, 1971g: 196, 1972e: 285, 1972f: 64, 1975h: 352, 1979b: 417; Thompson, G. H. 1960, 1963: 70; Tordo 1957: 9. (tx) Browne 1971d: 230; Nunberg 1963a: 570–571; Schedl 1934a: 247, 1935e: 153–154, 1935g: 314, 1937b: 407, 1937d: 35–36, 1939g: 169, 1941d: 381, 1950e: 212, 1952g: 54, 1954d: 874, 1954e: 59–71, 1955i: 213, 1962h: 64–65, 1962k: 691, 1972f: 248, 1978a: 57.

dubiosus Schedl 1937d: 36 (*Crossotarsus*). Lectotype ♀; D.-O.-Afrika: Morogara; Schedl Col-

lection in NHMW, Wien, designated by Schedl 1978a: 27. Synonymy: Schedl 1962k: 685.

References: (tx) Nunberg 1963a: 566, 568; Schedl 1935g: 314, 1937d: 35–36, 1939b: 390, 1941e: 154, 1962k: 685, 1972f: 248, 1978a: 27.

pilosulus (Schedl) 1933f: 200 (*Crossotarsus*). Syn-types ♂; Equator: Flandria and Haut-Uele; Moto; MRCB, Tervuren, and Schedl Collection in NHMW, Wien.

Figures: Nunberg 1963a: 570.

Distribution: Africa (Central African Republic/ Nigeria/ Sierra Leone/ Zaire).

Hosts: *Piptadeniastrum africanum*.

Notes: (3) Nunberg 1963a: 569 and Browne 1971d: 225 (a good species), Schedl 1941d: 413, 1972f: 248 (= *marshalli*).

References: (ds) Browne 1971d: 225, 1980c: 486, 1980d: 493; Roberts 1969: 121; Schedl 1933f: 200, 1938d: 452, 1943c: 69; Wood, S. L. 1957e: 1272. (tx) Browne 1971d: 225; Nunberg 1963a: 569–570; Schedl 1933f: 200, 1935e: 153, 1935g: 314, 1937d: 35–36, 1941d: 381–413, 1941e: 153.

Genus *Mesoplatypus* Strohmeyer

MESOPLATYPUS STROHMEYER 1912: 78. Type-species: *Mesoplatypus grandiclavus* Strohmeyer, monobasic. Keys: Browne 1962e: 655.

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1985: 28; Schedl 1972f: 166. (ds) Schedl 1972f: 166. (tx) Browne 1962e: 649, 655, 1965a: 206, 1972a: 184; Hopkins 1914: 124, 133, 1915c: 227; Nunberg 1953: 44, 1965a: 385; Schedl 1939b: 402–403, 1957d: 128, 1962k: 995, 1963j: 483, 1965e: 349–379, 1972f: 166–168; Strohmeyer 1912c: 7, 1914c: 8–37, 1918: 1–46.

appendicinus Schedl 1965e: 374. Holotype ♂; Congo ex belge-Katanga, Bukama; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

References: (ds) Schedl 1972f: 168. (tx) Schedl 1965e: 374, 1972f: 168, 1978a: 12.

bilobatus Nunberg 1969a: 400. Holotype ♂; Bingerville [Ivory Coast]; MRCB, Tervuren.

Distribution: Africa (Ivory Coast).

Notes: (3) Schedl 1972q: 264 (= *nigeriensis*, *venustus*), Beaver & Loytyniemi 1985: 115 (a good species).

References: (ds) Schedl 1972f: 168. (tx) Beaver & Loytyniemi 1985: 115; Nunberg 1969a: 400; Schedl 1972f: 168, 1972q: 264.

calabarius Roberts 1965: 226. Holotype ♂; Nigeria: Obari; BMNH, London.

Distribution: Africa (Nigeria).

Hosts: *Diospyros* sp.

Notes: (3) Schedl 1972: 168 (= *nigeriensis*), Browne 1972: 111 and Beaver & Loytyniemi 1985: 115 (a good species).

References: (ds) Roberts 1968: 192. (tx) Beaver

- & Loytyniemi 1985b: 115; Browne 1972c: 111; Roberts 1965: 226, 1972f: 168.
- curvidens** Browne 1980e: 777. Holotype ♂: Zaire: Park National Albert, secteur nord, Ihunga, affl. rive droite Semliki, 1050 m; MRCB, Tervuren. Distribution: Africa (Zaire). References: (tx) Browne 1980e: 777.
- elegans** Schedl 1962k: 1121. Holotype ♂: Congo: Schedl Collection in NHMW, Wien. Distribution: Africa (Zaire). References: (ds) Schedl 1972f: 168. (tx) Schedl 1962k: 1121, 1972f: 168, 1978a: 27.
- erimaceus** Schedl 1957d: 130. Holotype ♂: Congo Belge: Yangambi; MRCB, Tervuren. Distribution: Africa (Zaire). Hosts: *Tridesmostemon claussensii*. References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (ds) Mayne & Donis 1962: 278; Schedl 1962k: 996, 1966c: 233, 1972f: 168. (tx) Nunberg 1965a: 357; Schedl 1957d: 130, 1962k: 996, 1966c: 238, 1972f: 168, 1978a: 28.
- flavescens** (Strohmeyer) 1912f: 85 (*Crossotarsus*). Syntypes ♂ ♀: Mkululumi-Berg (Deutsch-Ost-Afrika) und West-Usambara; Hamburg Museum, lost, IPKE, Eberswalde. Distribution: Africa (Cameroon/ Equatorial Guinea/ Tanzania). References: (hb) Strohmeyer 1911a: 182. (ds) Schedl 1933f: 202–203, 1972f: 168; Strohmeyer 1911a: 1182, 1912c: 7, 1914c: 34. (tx) Schedl 1933f: 202–203, 1941d: 410, 1962k: 957, 1972f: 168; Strohmeyer 1911a: 182, 1912c: 7, 1912f: 85–86, 1914c: 34.
- grandiclava** Strohmeyer 1912f: 79. Syntypes ♂ ♀: Französisch-Kongo; Strohmeyer Collection. Distribution: Africa (Cameroon/ Equatorial Guinea/ Gabon/ Guinea/ Zaire). References: (ds) Schedl 1962k: 997, 1966c: 233, 1971g: 195, 1972f: 168; Strohmeyer 1912c: 7, 1914c: 37. (tx) Hopkins 1914: 124, 133; Lucas 1920: 406; Nunberg 1965a: 386–387; Schedl 1939b: 394, 403, 1962k: 997, 1972f: 168; Strohmeyer 1912c: 7, 1912f: 78–79, 1914c: 27. (ms) Lucas 1920: 406.
- grandis** Schedl 1950d: 33. Holotype ♀: Congo Belge: Elisabethville; IRSNB, Brussels. Distribution: Africa (Zaire). Notes: (3) Schedl 1965e: 375 (described male). References: (ds) Schedl 1972f: 168. (tx) Schedl 1950d: 33, 1962k: 997, 1965e: 375, 1972f: 168, 1978a: 33.
- imparispinosus** Roberts 1972: 67. Holotype ♂: West Cameroon: Bafut-Ngemba Forest Reserve; BMNH, London. Distribution: Africa (Cameroon). References: (ds) Browne 1986b: 334. (tx) Roberts 1972: 67.
- kitushi** Schedl 1957d: 128. Holotype ♂: Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren. Distribution: Africa (Nigeria/ Zaire). Hosts: *Beilschmiedia lousii*, *Bosquecia angolensis*, *Ficus* sp., *Garcinia polyantha*, *Parinari holstii*, *Syzygium* sp. References: (hb) Schedl 1962k: 998. (ds) Roberts 1968: 192, 1969: 116; Schedl 1962k: 998, 1972f: 168. (tx) Nunberg 1965a: 388; Schedl 1957d: 128, 1962k: 998, 1972f: 168, 1978a: 39.
- parvus** Nunberg 1967b: 333. Holotype ♂: Cote d'Ivoire, Adiopodoume; MRCB, Tervuren. Figures: Nunberg 1967b: 333. Distribution: Africa (Ivory Coast). References: (tx) Nunberg 1967b: 333.
- quadrispinis** Browne 1972c: 110. Holotype ♂: Congo-Kinshasa: Kivu, Bitale; MRCB, Tervuren. Distribution: Africa (Zaire). References: (ds) Ohno, Yoshioka, et al. 1989: 66. (tx) Browne 1972c: 110.
- quinquecinctus** (Schedl) 1941d: 410 (*Doliopygus*). Lectotype ♂: Congo, Kivu: Lugira-Kalehe, 1600 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 61. Figures: Schedl 1962k: 958. Distribution: Africa (Guinea/ Ivory Coast/ Sierra Leone/ Tanzania/ Uganda/ Zaire). Hosts: *Parinari holstii*, *Piptadenia* sp., *Syzygium* sp. Notes: (1) Browne 1970: 580 (to *Mesoplatypus*). References: (ds) Schedl 1962k: 957, 1966c: 232, 1972f: 168, 1979b: 416. (tx) Browne 1970: 580; Nunberg 1966a: 188; Schedl 1941d: 410, 1957d: 144, 1962k: 957–958, 1972f: 168, 1978a: 61.
- quinquecinctus cachani** Schedl 1957d: 145 (*Doliopygus*). Lectotype ♂: Ivory Coast: Adiopodoume; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 61. References: (ds) Schedl 1962k: 958, 1966a: 276, 1967e: 22, 1972f: 168. (tx) Schedl 1957d: 144–145, 1962k: 958, 1972f: 168, 1978a: 61.
- ustus** Schedl 1965e: 376. Holotype ♀: Congo ex belge-Yangambi; Schedl Collection in NHMW, Wien. Distribution: Africa (Zaire). Hosts: *Maba laurentii*. References: (ds) Schedl 1972f: 168. (tx) Schedl 1965e: 374–376, 1972f: 168, 1978a: 78.
- ventricornis** Schedl 1957d: 129. Holotype ♂: Congo Belge: Yangambi; MRCB, Tervuren. Figures: Nunberg 1967a: 338. Distribution: Africa (Ghana/ Nigeria/ Zaire). Hosts: *Bussia occidentalis*, *Celtis soyauxii*, *Dialium corbisieri*. References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Beaver & Loytyniemi 1985b: 116; Browne 1963a: 253; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Schedl 1962k: 998. (ds) Beaver & Loytyniemi 1985b: 116; Browne

- 1963a: 253; Roberts 1968: 192, 1969: 116; Schedl 1962h: 62–63, 1962k: 998, 1972f: 168. **(tx)** Numberg 1965a: 388; Schedl 1957d: 129, 1962k: 998–999, 1972f: 168, 1978a: 79.
- bituberculatus* Numberg 1967a: 325. Holotype ♀; Elisabethville, Congo; MRCB, Tervuren. Synonymy: Beaver & Loytyniemi 1985: 116. Notes: (1) Beaver & Loytyniemi 1985: 116 (to *Mesoplatypus*). References: **(tx)** Beaver & Loytyniemi 1985: 116; Numberg 1967a: 325, 338.
- venustus** Schedl 1962k: 1122. Holotype ♂; Congo; Schedl Collection in NHMW, Wien. Distribution: Africa (Ghana/ Nigeria/ Sierra Leone/ Zaire). Hosts: *Amphimas pterocarpoides*, *Bussea occidentalis*, *Celtis soyauxii*, *Coelocaryon* sp., *Dialium corbisieri*, *Hydodendron gabunensis*, *Monodora myristica*, *Oxystigma mannii*, *Pterocarpus osun*, *Pycnanthus angolensis*. Notes: (3) Schedl 1965c: 376 (described female). References: **(ds)** Schedl 1966c: 233. **(tx)** Schedl 1962k: 112, 1965c: 376, 1972f: 168, 1978a: 80.
- nigeriensis* Roberts 1964: 412. Holotype ♂; Nigeria: Cross River North Forest Reserve. Synonymy: Schedl 1972c: 264. References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(hb)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(ds)** Beaver & Loytyniemi 1985b: 115; Menier 1973a; Roberts 1968: 192, 1969: 115; Schedl 1972f: 168. **(tx)** Browne 1972c: 111; Roberts 1964: 412; Schedl 1972f: 168, 1972c: 264.
- Genus *Doliopygus* Schedl**
- DOLIOPYGUS SCHEDL 1939b: 402–403. Type-species: *Crossotarsus bohemanii* Chapuis, subsequent designation by Schedl 1972f: 143.
- Scutopygus* Numberg 1966a: 187, 189. Type-species: *Crossotarsus rapax* Sampson, subsequent designation by Wood 1992a: 90. References: **(tx)** Numberg 1966a: 187, 189; Wood, S. L. 1992a: 90.
- Pygodolius* Numberg 1966a: 185–189. Type-species: *Crossotarsus vegrandis* Sampson, subsequent designation by Wood 1992a: 90. References: **(tx)** Numberg 1966a: 185–189; Wood, S. L. 1992a: 90.
- Mixopygus* Numberg 1966a: 188. Type-species: *Crossotarsus conradti* Strohmeier, subsequent designation by Wood 1992a: 90. References: **(tx)** Numberg 1966a: 188; Wood, S. L. 1992a: 90.
- Mesopygus* Numberg 1966a: 187–188. Type-species: *Crossotarsus ukerevicensis* Schedl, subsequent designation by Wood 1992a: 90. References: **(tx)** Numberg 1966a: 187–188; Wood, S. L. 1992a: 90.
- Keys: Schedl 1972f: 145 (to species groups only).

- References: **(ay)** Nobuchi 1969a: 69. **(hb)** Curry 1958: 128–132; Schedl 1972f: 143–146. **(ds)** Schedl 1972f: 143–146. **(tx)** Browne 1962c: 648, 1972a: 182; Numberg 1966a: 185; Schedl 1939b: 402–403, 1957d: 1–162, 1962k: 744, 1965c: 52, 1965e: 349–379, 1972f: 143–146.
- abbreviatus** (Strohmeier) 1911g: 224 (*Crossotarsus*). Holotype ♂; Kamerun; Strohmeier Collection. Distribution: Africa (Cameroon/ Nigeria) References: **(cc)** Roberts 1969a. **(ds)** Roberts 1968: 190, 1969b: 106; Schedl 1972f: 157; Strohmeier 1912c: 7, 1914c: 34. **(tx)** Schedl 1962k: 825, 1972f: 157; Strohmeier 1911g: 224–225, 1912c: 7, 1914c: 34.
- abdominales** Schedl 1962k: 1109. Holotype ♂; Congo; Schedl Collection in NHMW, Wien. Distribution: Africa (Zaire). References: **(ds)** Schedl 1972f: 159. **(tx)** Schedl 1941d: 413, 1952g: 56, 1962k: 1109, 1972f: 159, 1978a: 9.
- acutespinatus** Numberg 1966a: 190. Holotype ♂; Congo: Kivu, foret Biambwé, terr. Lubero, 950 m; MRCB, Tervuren. Figures: Numberg 1966a: pl. 2, figs. 1–3. Distribution: Africa (Zaire). References: **(tx)** Numberg 1966a: 190.
- aduncus** Schedl 1961d: 178. Holotype ♂; [Senegal] Parc Natl. du Niokolo-Koba. Badi; MNHN, Paris. Distribution: Africa (Burkina Faso/ Central African Republic/ Ghana/ Nigeria/ Senegal/ Sierra Leone). Hosts: *Daniella oliveri*. References: **(cc)** Roberts 1969a. **(hb)** Browne 1963a: 254. **(ds)** Browne 1963a: 254, 1972c: 99; Roberts 1968: 191, 1969b: 106; Schedl 1962b: 61, 1962k: 885, 1972f: 162. **(tx)** Numberg 1966a: 186; Schedl 1961d: 178, 1962k: 885, 1965g: 21, 1972f: 162, 1978a: 9.
- aequalidentatus** Schedl 1966a: 377. Holotype ♂; Congo ex belge-Elisabethville; Schedl Collection in NHMW, Wien. Distribution: Africa (Zaire/ Zambia). References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(hb)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(ds)** Beaver & Loytyniemi 1985b: 123; Schedl 1972f: 149. **(tx)** Schedl 1965a: 377, 1972f: 149, 1975a: 10.
- semimitratus* Numberg 1966a: 193. Holotype ♀; Congo: Elisabethville; MRCB, Tervuren. Synonymy: Beaver & Loytyniemi 1985: 123. References: **(tx)** Beaver & Loytyniemi 1985: 123; Numberg 1966a: 193.
- alternans** (Strohmeier) 1911g: 224 (*Crossotarsus*). Holotype ♂; Kamerun; MNB, Berlin. Distribution: Africa (Cameroon/ Nigeria). Hosts: *Trichilia* sp. Notes: (3) Roberts 1966b: 620 (described female).

- References: **(ds)** Roberts 1968: 190; Schedl 1962k: 110, 1972f: 157; Strohmeier 1912c: 7, 1914c: 34. **(tx)** Numberg 1966a: 186; Roberts 1966b: 620, 1969b: 106; Sampson 1924: 127; Schedl 1962k: 826, 1110, 1972f: 157; Strohmeier 1911g: 224, 1912c: 7, 1914c: 34.
- angolensis** Schedl 1959p: 26. Holotype, sex?: Dundo, Angola; MRCB, Tervuren.
Distribution: Africa (Angola/ Nigeria/ Zaire).
Hosts: *Dialium* sp.
Notes: (1) Originally named as a subspecies of *exilis*; Browne 1965a: 209 (a good species).
References: **(ds)** Roberts 1968: 190, 1969b: 110; Schedl 1962k: 1113, 1972f: 154. **(tx)** Browne 1965a: 209; Schedl 1959p: 26, 1962k: 779, 1113, 1972f: 154, 1978a: 11.
- arrowi** (Schedl) 1937b: 406 (*Crossotarsus*). Lectotype ♂; N. Rhodesia, N'Change, Belgisch Kongo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 12.
Figures: Schedl 1962k: 927, 1972f: 150.
Distribution: Africa (Zaire/ Zambia/ Zimbabwe).
References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1955: 28. **(hb)** Loytyniemi, Beaver, & Loytyniemi 1955: 28. **(ds)** Beaver & Loytyniemi 1955b: 123; Schedl 1935d: 451, 1962k: 927, 1972f: 149. **(tx)** Numberg 1966a: 187; Schedl 1937b: 406, 1938d: 451, 1957d: 136, 1962k: 927, 1972f: 149–150, 1978a: 12.
- artespinitatus** (Schedl) 1935g: 318 (*Crossotarsus*). Holotype ♂; Belgian Congo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Gabon/ Ghana/ Nigeria/ Uganda/ Zaire).
Hosts: Schedl 1962k: 746 (23 host species listed).
Notes: (3) Schedl 1936b: 131, 1957d: 130 (described female).
References: **(ce)** Schedl 1962k: 745. **(hb)** Schedl 1962k: 745. **(ds)** Mayne & Donis 1960: 96, 1962: 266; Numberg 1964a: 237; Roberts 1961a: 39–43; Schedl 1962h: 61, 1962k: 745, 1967e: 221, 1972f: 153. **(tx)** Numberg 1966a: 186; Powell, W. 1957: 27; Schedl 1935g: 318–319, 1936b: 131, 1936e: 50, 1939g: 168, 1957d: 130, 1962h: 61, 1962k: 745, 1972f: 153, 1978a: 12.
- bituberculatus** Schedl 1936b: 134 (*Crossotarsus*). Holotype ♀; Kongo; Schedl Collection in NHMW, Wien. Synonymy: Schedl 1957d: 131.
References: **(tx)** Numberg 1967b: 338; Schedl 1936b: 134, 1939g: 168, 1957d: 130, 1978a: 16.
- caliginosus** Roberts 1966: 620. Holotype ♂; Nigeria: Boshi Extension F.R.; BMNH, London. Synonymy: Schedl 1972f: 153.
References: **(ds)** Roberts 1968: 189. **(tx)** Roberts 1966: 620; Schedl 1972f: 153.
- ater** Browne 1972c: 117. Holotype ♂; Congo-Kinshasa: Kivu, Bitale; MRCB, Tervuren.
Distribution: Africa (Zaire).
References: **(tx)** Browne 1972c: 117.
- berliniae** Roberts 1970a: 66. Holotype ♂; Nigeria: Mayo Selbe; MRCB, Tervuren.
Distribution: Africa (Nigeria).
Hosts: *Berlinia confusa*.
References: **(tx)** Roberts 1970a: 66.
- bidentatus** (Strohmeier) 1911g: 222 (*Crossotarsus*). Syntypes ♂ ♀; Deutsch-Ost-Afrika; MNB, Berlin, Strohmeier Collection.
Figures: Schedl 1962k: 948–949, 1972f: 164.
Distribution: Africa (Kenya/ Nigeria/ Ruanda/ Tanzania/ Zaire).
Hosts: *Cephalosphaera usambarensis*, *Pachystela laurentii*, *Polyscias kikuyuensis*. Schedl 1962k: 950–951 (15 host species listed).
References: **(hb)** Strohmeier 1911a: 182. **(ds)** Gardner 1957a; Schedl 1962k: 947, 1971e: 4, 1972f: 163; Strohmeier 1911a: 182, 1912c: 7, 1914c: 35. **(tx)** Numberg 1966a: 187; Powell, W. 1957: 27; Schedl 1935g: 319, 1937d: 35–36, 1941d: 411, 1941e: 154, 1957d: 144, 1962h: 64, 1962k: 947–948, 1972f: 163–164; Strohmeier 1911a: 182, 1911g: 222–223, 1912c: 7, 1914c: 11, 35.
- kenyaensis** Schedl 1941d: 411. Syntypes ♂ ♀; Kenya: Mt. Elgon, 2470 m; Schedl Collection in NHMW, Wien.
Notes: (1) Schedl 1978a: 39 (citation of holotype invalid). (3) Schedl 1972f: 163 (treated as a subspecies of *bidentatus*).
References: **(cn)** Browne 1968: 247. **(hb)** Browne 1968: 247. **(ds)** Browne 1968: 247; Gardner 1957a: 28. **(tx)** Numberg 1966a: 187; Powell, W. 1957: 27; Schedl 1941d: 408, 411, 1952j: 6, 1957d: 144, 1962h: 65, 1962k: 955, 1972f: 163, 1978a: 39.
- bidentatus posticalis** Schedl 1957d: 144. Lectotype ♂; Ruanda et du Congo Belge; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 15.
References: **(ce)** Schedl 1962k: 949. **(hb)** Gardner 1957: 28–29; Schedl 1962k: 949. **(ds)** Mayne & Donis 1962: 266; Roberts 1968: 192, 1969: 106; Schedl 1955i: 213, 1962k: 949, 1972f: 163. **(tx)** Numberg 1966a: 189; Schedl 1957d: 144, 1962k: 949, 1972f: 163–164, 1978a: 15.
- leleupi** Numberg 1960b: 305. Holotype ♂; Tanganyika Terr., Bunduki, Uluguru Mts., 1300 m; MRCB, Tervuren. Synonymy: Schedl 1972f: 163.
References: **(tx)** Numberg 1960b: 300, 304, 1966a: 197; Schedl 1962h: 64, 1972f: 163.
- uluguruensis** Numberg 1960b: 306. Holotype, sex?; Tanganyika Terr.: Uluguru Mts., vallee de l'Uhulu-Ndogo, 1500 m; MRCB, Tervuren. Synonymy: Schedl 1972f: 163.
References: **(tx)** Numberg 1960b: 304, 1966a: 197; Schedl 1962h: 65, 1972f: 163.
- bidiscoplanus** Roberts 1970b: 474. Holotype ♂; Sierra Leone: Gola North Forest Reserve; BMNH, London.
Distribution: Africa (Sierra Leone).

- Hosts: *Tarrietia utilis*, *Tieghmella hecklii*.
References: (ds) Schedl 1972f: 149. (tx) Roberts 1970b: 474; Schedl 1972f: 149.
- bijunctispinae** Roberts 1970b: 475. Holotype ♂; Sierra Leone: Gola West Forest Reserve; BMNH, London.
Distribution: Africa (Sierra Leone).
Hosts: *Ochthocosmus africanus*.
References: (ds) Schedl 1972f: 149. (tx) Roberts 1970b: 475; Schedl 1972f: 149.
- bilobatus** (Schedl) 1936b: 127 (*Crossotarsus*). Holotype ♂; Franz. Congo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Gabon/ Nigeria/ Sierra Leone/ Zaïre).
Hosts: *Antrocaryon micraster*, *Dialium* spp., *Distemonanthus benthamianus*, *Klainedoxa gabonensis*, *Lindackeria dentata*, *Macaranga* sp., *Pentaclethra macrophylla*, *Pycauthus angolensis*, *Strombosia scheffleri*.
Notes: (3) Schedl 1950d: 36 (described female).
References: (ec) Schedl 1962k: 748. (hb) Schedl 1962k: 748. (ds) Cola 1973; Mayne & Donis 1962: 267; Schedl 1962k: 748, 1972a: 153. (tx) Nunberg 1966a: 186; Schedl 1936b: 127–128, 1950d: 36, 1957d: 133, 142, 1962k: 748, 1972a: 153, 1978a: 16. *dialiumi* Roberts 1966: 221. Holotype ♂; Nigeria: Alade; BMNH, London. Synonymy: Schedl 1972f: 153.
References: (ds) Roberts 1968: 189, 1969b: 108–109. (tx) Roberts 1965: 221, 1966c: 410; Schedl 1972f: 153.
- bipapillatus** Roberts 1966c: 410. Holotype ♂; Nigeria: Cross River South Forest Reserve; BMNH, London.
Distribution: Africa (Nigeria).
References: (ds) Roberts 1968: 191; Schedl 1972f: 150. (tx) Roberts 1966c: 410; Schedl 1972f: 150.
- bitalei** Schedl 1957d: 135. Holotype ♂; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.
Distribution: Africa (Nigeria/ Ruanda/ Zaïre).
Hosts: *Beilschmiedia lousii*, *Berlinia* sp., *Diospyros* sp., *Lovoa trichiliodes*, *Radlkofera* sp.
References: (ec) Schedl 1958d: 196, 1962k: 928. (hb) Schedl 1958d: 196, 1962k: 928. (ds) Roberts 1968: 191, 1969b: 107; Schedl 1962k: 928, 1972f: 150. (tx) Nunberg 1966a: 187; Schedl 1955f: 261, 1957d: 135, 1962k: 928, 1972f: 150, 1978a: 16. *integerrimus* Roberts 1966b: 625. Holotype ♂; Nigeria: Oban; BMNH, London. Synonymy: Schedl 1972f: 150.
References: (ds) Roberts 1968: 192. (tx) Roberts 1966b: 625; Schedl 1972f: 150.
- bohemani** (Chapuis) 1865: 93 (*Crossotarsus*). Holotype ♂; Cafrerie [Mandibira, D.O.A., given by Schedl 1971: 15]; NHR, Stockholm (Schedl 1960c: 12).
Distribution: Africa (Cameroon/ Chad/ Ethiopia/ Ghana/ Kenya/ Mozambique/ Nigeria/ Senegal/ South Africa/ Sudan/ Tanzania/ Uganda/ Zaïre/ Zambia/ Zimbabwe).
Notes: (3) Schedl 1971c: 15 (described female).
References: (ec) Roberts 1969b: 107. (ds) Browne 1975b: 395; Fahraeus 1871: 661; Schedl 1938d: 451, 1943c: 69, 1962k: 750, 1965e: 358, 1968b: 145, 1971e: 5, 15, 1971f: 150, 1972f: 153, 1975h: 351, 1982: 279; Strohmeier 1911f: 204, 1912c: 6, 1914c: 34. (tx) Browne 1962e: 648; Chapuis 1865: 93; Nunberg 1966a: 186; Schaufuss 1905: 103; Schedl 1936b: 128, 1937d: 35, 1939b: 403, 1941d: 381, 1941e: 154, 1952g: 69, 1952i: 53, 1960c: 12, 1962k: 750, 1971e: 15, 1972f: 153, 1978a: 17; Strohmeier 1911f: 204, 1912c: 6, 1914c: 34.
- bothrocephalus** (Strohmeier) 1907g: 29 (*Crossotarsus*). Holotype ♀; Tanga, Ostafrika; Strohmeier Collection.
Distribution: Africa (Tanzania).
References: (ds) Nunberg 1960a: 301; Schedl 1972f: 112; Strohmeier 1912c: 7, 1914c: 35. (tx) Nunberg 1960: 301, 1966a: 186; Schedl 1962k: 1060, 1123, 1972f: 112; Strohmeier 1907g: 29, 1912c: 7, 1914c: 35.
- brevis** (Strohmeier) 1911g: 228 (*Crossotarsus*). Holotype ♂; Kamerun; Strohmeier Collection.
Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Tanzania).
Hosts: *Azacia africana*, *Albizia zygia*, *Antiaris africana*, *Blighia sapida*, *Bosqueia angolensis*, *Bussea occidentalis*, *Castilloa* sp., *Celtis mildbraedii*, *Cylicodiscus gabunensis*, *Entandrophragma utile*, *Ficus exasperata*, *Guarea cedrata*, *Hexalobus africanus*, *Hymenostegia afzelii*, *Lonchocarpus sericeus*, *Morus mesozygia*, *Panda oleosa*, *Sterculia rhinopetala*, *Strombosia glaucescens*, *Terminalia ivorensis*, *T. superba*, *Trichilia heudelotii*, *Vernonia* sp.
References: (cn) Kleine 1932a: 310, 401. (hb) Aulmann 1912: 58, 1913: 1–126; Browne 1963a: 254; Kleine 1932a: 310, 401; Strohmeier 1911a: 182. (ds) Browne 1963a: 254; Cachan 1957: 15; Kleine 1932a: 310, 401; Mayne & Donis 1962: 267; Schedl 1959p: 22, 1962k: 898, 1972f: 147; Strohmeier 1911a: 182, 1912c: 6, 1914c: 34. (tx) Nunberg 1966a: 187; Schedl 1951p: 40, 1952i: 30, 1954d: 873, 1957d: 142, 1962k: 898, 1972f: 147; Strohmeier 1911a: 182, 1911g: 228–229, 1912c: 6, 1914c: 34.
- bulbifer** Schedl 1962k: 1110. Holotype ♂; Nigeria: Abuja; Schedl Collection in NHMW, Wien.
Figures: Roberts 1964: 402.
Distribution: Africa (Equatorial Guinea/ Nigeria/ Sierra Leone).
Hosts: *Cordia senegalensis*, *Milletia thonningii*.
References: (ec) Roberts 1969a. (ds) Roberts 1968: 191; Schedl 1971g: 194, 1972f: 151. (tx) Schedl 1962k: 1110, 1972f: 151, 1978a: 17. *bulbosus* Roberts 1964: 401. Holotype ♂; Nigeria:

Abuja; BMNH, London. Synonymy: Schedl 1972f: 151.

References: **(ds)** Browne 1984c: 450. **(tx)** Roberts 1964: 401, 1969b: 107; Schedl 1972f: 151.

bulbiventris Schedl 1962k: 1112. Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Ghana).

References: **(ds)** Schedl 1972f: 151. **(tx)** Schedl 1962k: 872, 1972f: 151, 1978a: 17.

umbonatus Browne 1965a: 208. Holotype ♂; Ghana; Kumasí; BMNH, London. Synonymy: Schedl 1972f: 151.

References: **(tx)** Browne 1965a: 208; Schedl 1972f: 151.

caesalpiniae Roberts 1970a: 67. Holotype ♂; Nigeria; Mayo Selbe; MRCB, Tervuren.

Distribution: Africa (Nigeria).

Hosts: *Anthonota macrophylla*, *Berlinia confusa*, *Brachystegia nigerica*.

References: **(tx)** Roberts 1970a: 67.

carapae Schedl 1957d: 132. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Alstonia boonei*, *Carapa procera*.

References: **(ds)** Mayne & Donis 1960: 96, 1962: 267; Schedl 1962k: 751, 1972f: 153. **(tx)** Numberg 1966a: 186; Schedl 1957d: 132, 1962k: 751, 1972f: 153, 1978a: 18.

chapuisi (Duvivier) 1891: 377 (*Crossotarsus*). Holotype ♀; Ibembo, Congo; not given.

Figures: Browne 1972c: 112, Numberg 1960a: 300, 1966a: 197.

Distribution: Africa (Angola/ Bechuanaland/ Cameroon/ Congo/ Equatorial Guinea/ Ghana/ Guinea/ Ivory Coast/ Mozambique/ Nigeria/ Sierra Leone/ South Africa/ Spanish Guinea/ Tanzania/ Togo/ Uganda/ Zaire/ Zambia/ Zimbabwe). Hosts: Browne 1963a: 255, Schedl 1962k: 755–757 (many host species listed).

References: **(ay)** Lhote & Roche 1960; Menier 1976: 348; Strohmeier 1918: 8. **(bv)** Meixner 1937: 1217; Menier 1976: 348. **(cn)** Anonymous 1953j: 25; Browne 1968b: 246; Mayne & Donis 1951: 335. **(cc)** Anonymous 1953j: 25; Fuller 1958: 100; Jover 1952: 73–81; Mayne & Donis 1951: 335; Roberts 1969a. **(hb)** Beaver & Loytyniemi 1955b: 125; Browne 1963a: 255, 1968b: 246; Cachan 1957: 6, 15; Husson 1955: 354; Jones, Roberts, & Baker 1959: 13, 21, 36; Jover 1952: 73–81; Schedl 1962k: 751; Thompson, G. H. 1963: 40; Webb 1957: 26, 36–37, 40. **(ds)** Beaver & Loytyniemi 1955b: 125; Browne 1963a: 255, 1968b: 246, 1975a: 757, 1980a: 373, 1980b: 382; Cola 1973; Ferreira 1965: 1131; Gardner 1957a: 28; Mayne & Donis 1960: 96, 1962: 267; Menier 1973a; Roberts 1960b: 35, 1960e: 35–37, 1960f: 39, 1968: 189, 1969b: 107–108; Schedl 1933f: 196, 1959p: 22, 1962h: 62–64, 1962k: 751, 1964j: 42, 1965c: 358, 1966c: 231, 1967e: 221, 1971g:

194, 1972c: 285, 1975b: 351, 1979b: 416; Strohmeier 1912c: 6, 1914c: 34; Thompson, G. H. 1963: 40. **(tx)** Browne 1972c: 112; Duvivier 1891: 377, 386; Meixner 1937: 1217; Numberg 1966a: 186, 197; Powell, W. 1987: 27; Sampson 1924: 125–126; Schaufuss 1897b: 104–105, 1905: 103; Schedl 1933f: 196, 1936e: 51, 1937b: 407, 1937d: 35, 1939b: 391, 1939g: 167, 1941d: 381, 1941e: 153, 1950c: 205–206, 1950d: 4, 11, 14–15, 19, 1950e: 212, 1951e: 39, 1951f: 40, 1952j: 6, 1953g: 244, 1954d: 873, 1954e: 47, 54, 1955e: 270, 1962k: 751, 1972f: 153; Strohmeier 1911f: 204, 1912c: 6–7, 1914c: 34, 1918: 12–41.

maculatus Schaufuss 1897b: 104 (*Crossotarsus*).

Syntypes ♂ ♀; Sierra Leone; Hamburg Museum, lost. Synonymy: Schedl 1972f: 153.

References: **(ds)** Strohmeier 1911f: 204. **(tx)** Schaufuss 1897b: 104; Schedl 1933f: 196.

chapuisi nitidiventris Schedl 1962k: 757. Lectotype ♂; Belgisch Kongo; Elisabethville; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 19.

References: **(ds)** Schedl 1972f: 153. **(tx)** Schedl 1962k: 757, 1972f: 153, 1978a: 19.

bugiriensis Numberg 1960a: 301. Holotype ♂; Uganda; Bugiri Terr., Longido, Masai District; MRCB, Tervuren. Synonymy: Schedl 1962h: 64.

References: **(tx)** Numberg 1960a: 300–301; Schedl 1962h: 64, 1972f: 153.

citri Schedl 1948c: 667. Lectotype ♂; Newcastle, Natal; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Congo/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sierra Leone/ South Africa/ Tanzania/ Zaire).

Hosts: *Citrus aurantium*. Schedl 1962k: 842–843 (30 host species listed).

References: **(ds)** Schedl 1972f: 157. **(tx)** Browne 1972c: 121; Numberg 1966a: 187; Schedl 1948c: 667, 1962k: 826, 1972f: 157, 1978a: 20.

interpositus Schedl 1952i: 56. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren. Synonymy: Browne 1972c: 121.

References: **(ds)** Cachan 1957: 15; Gardner 1957a: 28; Roberts 1968: 190, 1969b: 111; Schedl 1975k: 279. **(tx)** Browne 1972c: 121; Numberg 1966a: 187; Schedl 1952i: 56–57, 1954d: 873, 1955f: 261, 1962h: 62, 1962k: 841, 1115, 1972f: 158, 1978a: 38.

clarus (Schedl) 1936b: 132 (*Crossotarsus*). Syntypes ♀; Belg. Kongo, Diobo, and Likimi-Mimbo; Collart Collection, and Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Hosts: *Oxytigma oxyphyllum*.

Notes: (1) Schedl 1978a: 20 (citation of holotype invalid).

References: **(ds)** Schedl 1962k: 875, 1972a: 160. **(tx)** Numberg 1966a: 186; Schedl 1936b: 131–132, 1952i: 45, 1962k: 875, 1972f: 160, 1978a: 20.

coelocephalus (Schaufuss) 1905: 103 (reprint p. 5) (*Crossotarsus*). Holotype ♀; Camerunia; Hamburg Museum, lost.

Distribution: Africa (Cameroon/ Gabon/ Ghana/ Kenya/ Nigeria/ Uganda/ Zaïre).

Hosts: Browne 1963a: 256, Schedl 1962k: 759 (many host species listed).

References: (hb) Browne 1962d: 94, 1963a: 256; Schedl 1962k: 757. (ds) Browne 1963a: 256, 1984b: 287; Gardner 1957a: 28; Jones, Roberts, & Baker 1959: 13; Mayne & Donis 1960: 96, 1962: 268; Roberts 1968: 189, 1969b: 108; Schedl 1933f: 196, 1962h: 62, 1962k: 757, 1966c: 231, 1972f: 153; Strohmeyer 1912c: 6, 1914c: 34. (tx) Nunberg 1966a: 186; Schaufuss 1905: 103–104; Schedl 1933f: 196–197, 1937d: 35, 1950d: 11, 1952i: 26, 1955f: 261, 1955i: 213, 1962h: 64, 1962k: 757, 1965g: 21, 1972a: 153; Strohmeyer 1912c: 6–7, 1914c: 34.

tricuspis Schedl 1933f: 196 (*Crossotarsus*).

Lectotype ♂; Sankuru: Lodja, Sandoa, Haut-Uele; Watsa; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 74. Synonymy: Schedl 1972f: 153.

References: (ds) Schedl 1933f: 196. (tx) Schedl 1933f: 196, 1937d: 35, 1950d: 11, 1952i: 26, 1972f: 153, 1978a: 74.

fenestralis Nunberg 1960a: 303. Type not designated; type locality and repository not stated, nomen nudum (?). Synonymy: Schedl 1962h: 64.

References: (tx) Nunberg 1960a: 300, 1960b; Schedl 1962h: 64, 1972f: 153.

congonus Browne 1972c: 115. Holotype ♂; Congo-Kinshasa: Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

References: (tx) Browne 1972c: 115.

conjunctus Schedl 1957d: 140. Holotype ♂; Congo Belge: route Stanleyville-Irumu, Camp Putnam; MRCB, Tervuren.

Figures: Schedl 1962k: 877, 1972f: 161.

Distribution: Africa (Tanzania/ Zaire).

Hosts: *Julbernardia sereti*.

References: (ds) Schedl 1962k: 876, 1972f: 160. (tx) Nunberg 1966a: 186; Schedl 1957d: 140, 1962k: 876–877, 1972f: 160–161, 1978a: 21.

conradti (Strohmeyer) 1911g: 227 (*Crossotarsus*). Syntypes ♂ ♀; Kamerun; MNB, Berlin, Strohmeyer Collection.

Figures: Schedl 1962k: 853, 1972f: 159.

Distribution: Africa (Angola/ Cameroon/ Central African Republic/ Congo/ Equatorial Guinea/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Mozambique/ Nigeria/ Sierra Leone/ Togo/ Zaire).

Hosts: Schedl 1962k: 855–861 (138 host species listed).

References: (ay) Entwistle 1963b; Farris & Funk 1965: 531. (bv) Browne 1962f; Entwistle 1963b. (cn) Anonymous 1953h: 31, 1953j: 25; Browne

1968: 246; Kudler 1978: 16; Thompson, G. H. 1960; Webb & Jones 1958: 383. (cc) Anonymous 1953j: 25; Browne 1961e: 10, 1962d; Farris & Funk 1965: 531; Roberts 1969b: 108; Schedl 1958d: 195–196, 1962k: 852. (hb) Anonymous 1959i: 18; Browne 1961b: 21–22, 24, 1962f, 1963a: 256, 1968: 246; Entwistle 1963b; Roberts 1961: 53–60; Schedl 1958d: 195–196, 1962k: 852; Thompson, G. H. 1963: 40; Webb & Jones 1958: 383. (ds) Browne 1961e: 10, 1963: 256, 1968: 246; Cachan 1957: 5–6, 15, 43–53; Ferreira 1965: 1131; Gardner 1957a: 28; Jones, Roberts, & Baker 1959: 13–52; Jover 1952: 73–81; Kolbe 1897: 283; Mayne & Donis 1960: 96, 1962: 268; Roberts 1960b: 35, 1960e, 1960f, 1968: 191; Schedl 1959p: 22, 1962k: 852, 1965e: 358, 1965g: 21, 1971g: 194, 1972c: 285, 1972f: 159, 1979b: 417; Strohmeyer 1912c: 6, 1914c: 34; Thompson, G. H. 1960, 1963: 40; Tordo 1957: 9; Webb & Jones 1957: 25–40. (tx) Nunberg 1966a: 186; Powell, W. 1957: 27; Schedl 1933f: 201, 1937d: 35, 1938d: 451, 1941e: 153, 1950d: 5, 1950e: 212, 1951e: 39, 1954d: 873, 1954e: 48, 55, 1955d: 270, 1959p: 22, 1962k: 852–853, 1965g: 21, 1972f: 159; Strohmeyer 1911g: 227–228, 1912c: 6–7, 1914c: 34.

vegrandis Sampson 1924c: 126 (*Crossotarsus*).

Holotype ♂; Belgian Congo: Mayumbe; MRCB, Tervuren. Synonymy: Schedl 1972f: 159.

Notes: (1) Schedl 1972f: 159 (to *Doliopygus*). (3) Schedl 1941d: 409 (described female).

References: (cc) Jover 1952. (hb) Jover 1952. (ds) Gardner 1957a; Mayne & Donis 1960: 98, 1962: 278; Schedl 1933f: 201, 1938d: 451. (tx) Nunberg 1966a: 187; Sampson 1924c: 126–127; Schedl 1933f: 201, 1940d: 5, 11, 1941d: 409, 1950d: 11, 1950e: 212, 1951e: 39, 1972f: 159–160, 1978a: 79.

convexus Roberts 1965: 224. Holotype ♂; Nigeria: Bende; BMNH, London.

Distribution: Africa (Nigeria).

Hosts: *Diospyros* sp.

References: (ds) Roberts 1965: 224, 1968: 191; Schedl 1972f: 150. (tx) Roberts 1965: 224; Schedl 1972f: 150.

costatus Schedl 1957d: 136. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Nunberg 1966a: 187.

Distribution: Africa (Nigeria/ Zaire).

Hosts: *Ochthocosmus africanus*.

Notes: (3) Schedl 1962k: 1112 (described female). References: (hb) Schedl 1962k: 930. (ds) Roberts 1968: 191; Schedl 1962k: 930, 1972f: 150. (tx) Nunberg 1966a: 187; Schedl 1957d: 136, 1962k: 930, 1112, 1972f: 150, 1978a: 22.

crinitus (Chapuis) 1865: 90 (*Crossotarsus*). Syntypes ♂ ♀; Senegal et du fleuve Blanc dans la Haute Egypte; MNHN, Paris (Schedl 1960c: 12).

Figures: Nunberg 1966a: 187.

Distribution: Africa (Cameroon/ Congo/ S Egypt/

- Gabon/ Kenya/ Malawi/ Mozambique/ Senegal/ South Africa/ Tanzania/ Zaïre/ Zambia/ Zimbabwe).
Hosts: *Pterocarpus angolensis*.
References: (**cn**) Anonymous 1970c: 11. (**ds**) Anonymous 1970c: 11; Beaver & Loytyniemi 1985b: 130; Browne 1972c: 99, 1984b: 288; Schedl 1938d: 451, 1962h: 62, 1962k: 878, 1969a: 204, 1972f: 161; Spahr 1981: 79; Strohmeier 1911f: 204, 1912c: 6, 1914c: 34; Tordo 1957: 9. (**tx**) Chapuis 1865: 24–25, 90–92; Duvivier 1891: 377; Lucas 1920: 209; Numberg 1966a: 187; Powell, W. 1987: 27; Sampson 1922: 139, 1924c: 126; Schaufuss 1905: 103; Schedl 1936e: 51, 1937b: 406, 1937d: 35, 1939a: 468–469, 1939b: 403, 1941e: 154, 1952i: 26–27, 1953g: 244, 1957b: 152, 1960c: 12, 1962k: 878, 1972f: 161; Strohmeier 1911g: 227–228, 1912c: 6–7, 1914c: 34. *hardenbergi* Sampson 1922a: 137 (*Crossotarsus*).
Holotype ♂; Portuguese East Africa: Xinavane; BMNH, London. Synonymy: Schedl 1951i: 26.
Notes: (3) Schedl 1941d: 408 (described female).
References: (**ds**) Gardner 1957a; Schedl 1933f: 198. (**tx**) Sampson 1922a: 137–139, 1924c: 126; Schedl 1933f: 198, 1937b: 406–407, 1937d: 35, 1941d: 408, 1950d: 12, 1951i: 26–27, 1956a: 35, 1972f: 161, 1978a: 35.
- dentiventris** Browne 1984c: 456. Holotype ♂; Owendo (Gabon) to Nagoya (Japan), imported; BMNH, London.
Distribution: Africa (Gabon).
Hosts: *Gnibourtia* sp.
References: (**tx**) Browne 1984c: 456.
- deruptus** Schedl 1950d: 35. Holotype ♀; Rhodesien (Chirinda); Schedl Collection in NHMW, Wien.
Distribution: Africa (Zimbabwe).
References: (**tx**) Numberg 1966a: 187; Schedl 1950d: 35, 1962k: 826, 1972f: 157, 1978a: 24.
- divaricus** (Schedl) 1936b: 126 (*Crossotarsus*).
Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon/ Zaïre).
Hosts: *Gnarea laurentii*. Schedl 1962k: 866–867 (18 host species listed).
References: (**ds**) Browne 1957b: 334; Schedl 1962k: 871, 1972f: 160. (**tx**) Numberg 1966a: 187; Schedl 1936b: 126–127, 1962k: 871, 1972f: 160, 1978a: 26.
- discrepans** Schedl 1952i: 38. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren. Synonymy: Schedl 1972f: 160.
References: (**cc**) Schedl 1962k: 866. (**hb**) Schedl 1962k: 866. (**ds**) Mayne & Donis 1960: 96, 1962: 268; Schedl 1962k: 866. (**tx**) Numberg 1966a: 186; Schedl 1952i: 38, 1957d: 133, 1962k: 866, 1972f: 160, 1978a: 25.
- divisus** Browne 1972c: 114. Holotype ♂; Congo-Kinshasa: Yangambi; MRCB, Tervuren.
Distribution: Africa (Zaïre).
References: (**tx**) Browne 1972c: 114.
- dolosus** Schedl 1950c: 216. Lectotype ♂; Congo Belge, Vieux-Kilo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 26.
Distribution: Africa (Ivory Coast/ Nigeria/ Zaïre).
Hosts: *Diospyros* sp., *Drypetes leouensis*, *Pachystela laurentii*.
References: (**cc**) Roberts 1969a. (**ds**) Cachan 1957: 15; Roberts 1968: 190, 1969b: 109; Schedl 1962k: 827, 1972f: 157. (**tx**) Numberg 1966a: 187; Schedl 1950c: 216, 1952i: 52, 1962k: 827, 1113, 1972f: 157, 1978a: 26.
- donisi** Schedl 1952i: 49. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
Figures: Schedl 1962k: 761 (male), 1972f: 152 (male).
Distribution: Africa (Zaïre).
Hosts: *Chrysophyllum lacourtianum*, *Pterygopodium oxyphyllum*, *Scorodophloeus zenkeri*. Schedl 1962k: 760–761 (12 host species listed).
References: (**hb**) Schedl 1962k: 760. (**ds**) Mayne & Donis 1960: 96, 1962: 269; Schedl 1962k: 760, 1972f: 153. (**tx**) Numberg 1966a: 186; Schedl 1952i: 49, 1962k: 760–761, 1972f: 152–153, 1978a: 26.
- dubius** (Sampson) 1924c: 126 (*Crossotarsus*).
Holotype ♂; Belgian Congo; MRCB, Tervuren.
Figures: Browne 1961b: pl. 1–2.
Distribution: Africa (Angola/ Bechuanaland/ Cameroon/ Congo/ Equatorial Guinea/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Senegal/ South Africa/ Sudan/ Tanzania/ Uganda/ Zaïre/ Zambia/ Zimbabwe).
Hosts: Schedl 1962k: 764–767 (79 host species listed).
Notes: (1) Originally named as a subspecies of *crinitus*.
References: (**ay**) Entwistle 1963b. (**bv**) Browne 1961b, 1961e: 8; Entwistle 1963b; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Roberts 1960b: 21. (**cn**) Browne 1961b, 1961e: 8, 1968: 246; Jones et al. 1959: 21–52; Kudler 1978: 16; Roberts 1960b: 21, 1966a. (**cc**) Browne 1961b, 1961e: 8, 1962d; Roberts 1960b: 21, 1969a; Schedl 1962k: 761. (**hb**) Beaver & Loytyniemi 1985b: 126; Browne 1961b: 15–27, 1961e: 8, 13, 1963a: 259, 1965b, 1968: 246; Entwistle 1963b; Jones et al. 1959: 21–52; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Roberts 1935e: 35–37, 1961a: 39–43, 1961c: 37, 1962a: 37–38, 1962b: 241, 1969b: 109; Schedl 1962k: 761; Thompson, C. H. 1963: 41; Webb & Jones 1957: 36–40. (**ds**) Beaver & Loytyniemi 1985b: 126; Browne 1961e: 8, 13, 1963a: 259, 1968: 246, 1972c: 99, 1980c: 485, 1984c: 450; Cola 1973; Ferreira 1965: 1132; Lee, R. F. 1971: 29; Mayne & Donis 1960: 96, 1962: 269; Roberts 1960b: 35–39, 1960e, 1960f, 1966a, 1968: 190; Schedl 1938d: 451, 1959p: 23, 1962h: 62,

- 1962k: 761, 1964j: 42, 1965e: 358, 1965g: 21, 1965h: 112, 1966a: 276, 1966c: 231, 1971e: 5, 1971f: 150, 1971g: 194, 1972e: 285, 1972f: 154, 1975b: 351, 1979b: 417; Thompson, G. H. 1963: 41; Wichmann 1955: 97. **(tx)** Browne 1961b: pl. 1, 2; Numberg 1966a: 186; Powell, W. 1987: 27; Sampson 1924c: 126; Schedl 1941d: 408, 1950d: 11, 1952g: 54, 1952i: 27, 1953g: 244, 1954d: 873, 1954e: 55, 1955i: 213, 1962k: 761, 1113, 1972f: 154; Strohmeier 1911g: 226–227.
- paradubius* Roberts 1964: 404. Holotype ♂; Nigeria: Omo Forest Reserve; BMNH, London. Synonymy: Schedl 1972f: 154. References: **(cn)** Browne 1968b: 248; Kudler 1978: 16. **(hb)** Browne 1968b: 248. **(ds)** Browne 1968b: 248; Roberts 1968: 190. **(tx)** Roberts 1964: 404, 1969b: 112–113; Schedl 1972f: 154.
- confusus* Roberts 1966b: 621. Holotype ♂; Nigeria: Oban; BMNH, London. Synonymy: Schedl 1972f: 154. References: **(ds)** Roberts 1968: 189. **(tx)** Roberts 1966b: 621; Schedl 1972f: 154.
- rhizophorae* Roberts 1966b: 629. Holotype ♂; Nigeria: Buguma; BMNH. Synonymy: Schedl 1972f: 154. References: **(ds)** Roberts 1968: 190. **(tx)** Roberts 1966b: 629; Schedl 1972f: 154.
- erichsoni* (Chapuis) 1865: 95 (*Crossotarsus*). Syntypes ♂ ♀; Port Natal; 1 ♂, 2 ♀ in BMNH, London, 1 ♂ in IRSNB, Brussels 1 ♂ in MNB, Berlin, 1 ♂ in MNHN, Paris. Figures: Schedl 1962k: 829, 1972f: 157. Distribution: Africa (Cameroon/ Congo/ Equatorial Guinea/ Gabon/ Ghana/ Guinea/ Kenya/ Mozambique/ Nigeria/ Ruanda/ Sao Tome Island/ Sierra Leone/ South Africa/ Tanzania/ Uganda/ Zaire). Hosts: Schedl 1962k: 830–833 (73 host species listed). References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(cn)** Anonymous 1970c: 12; Browne 1968b: 247; Jones 1959a. **(cc)** Schedl 1962k: 827. **(hb)** Anonymous 1959i: 18; Beaver & Loytyniemi 1985b: 129; Browne 1963a: 259, 1965b: 247; Jones 1959a, 1966: 127; Jones, Roberts, & Baker 1959: 13–21; Loytyniemi, Beaver, & Loytyniemi 1985: 29; Schedl 1962k: 827; Strohmeier 1911a: 182; Thompson, G. H. 1963: 41; Took & Scott 1944: 2–5; Tordo 1957: 9; Webb & Jones 1957: 37, 40. **(ds)** Anonymous 1970c: 12; Beaver & Loytyniemi 1985b: 129; Browne 1963a: 259, 1968b: 247, 1980b: 381; Gardner 1957a: 28; Mayne & Donis 1960: 96, 1962: 269; Roberts 1960b: 35, 1960e, 1960f: 35–39, 1966a, 1968: 190, 1969b: 109; Schedl 1933f: 201, 1935d: 452, 1962k: 827, 1964j: 43, 1965e: 359, 1965g: 21, 1972f: 157; Strohmeier 1911a: 182, 1911f: 204, 1914c: 34; Thompson, G. H. 1963: 41. **(tx)** Chapuis 1865: 24–25, 95–97; Lucas 1920: 209; Numberg 1966a: 187; Powell, W. 1987: 27; Sampson 1924a: 545, 1924c: 127; Schedl 1933f: 201, 1935g: 319–320, 1937c: 14, 1937d: 35–36, 1939b: 403, 1939g: 167, 1941d: 409, 1941e: 154, 1948c: 667, 1950d: 11, 1952g: 55, 1952i: 40, 1952j: 6, 1954e: 55, 1955j: 261, 1957c: 326, 1960c: 12–13, 1962k: 827–829, 1113, 1965g: 21, 1972f: 157; Strohmeier 1911a: 182, 1911f: 204, 1912c: 11, 1914c: 11, 34.
- varius* Sampson 1924a: 545 (*Platypus*). Holotype, sex?; South Africa: Cape Colony, Transkei; BMNH, London. Synonymy: Schedl 1937c: 14, Beaver & Loytyniemi 1985: 129. References: **(tx)** Beaver & Loytyniemi 1985: 129; Sampson 1924a: 545; Schedl 1937c: 14, 1957b: 152, 1972f: 157.
- excavatus* (Sampson) 1924c: 124 (*Crossotarsus*). Holotype ♂; Belgium Congo, Kapiri (Miss. Agric.); MRCB, Tervuren. Figures: Browne 1972c: 112. Distribution: Africa (Cameroon/ Kenya/ Nigeria/ Uganda/ Zaire/ Zambia). Hosts: Schedl 1962k: 775–776 (14 host species listed). References: **(cn)** Hargreaves 1927: 24–27. **(hb)** Roberts 1969b: 109–110. **(ds)** Gardner 1957a: 28. **(tx)** Browne 1972c: 111–112; Numberg 1966a: 186; Powell, W. 1987: 27; Sampson 1924c: 124; Schedl 1933f: 196, 1935g: 318, 1941e: 154, 1955f: 261, 1962h: 62, 1962k: 744, 1972f: 154.
- exilis* (Chapuis) 1865: 92 (*Crossotarsus*). Syntypes ♂ ♀; Fleuve Blanc [White Nile]; MNHN, Paris (Schedl 1960c: 12). Distribution: Africa (Cameroon/ Chad/ Congo/ Equatorial Guinea/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Spanish Guinea/ Sudan/ Tanzania/ Togo/ Uganda/ Zaire/ Zambia/ Zimbabwe). Hosts: Schedl 1962k: 777–778 (23 host species listed). Notes: (3) Schedl 1954e: 57 (*piptadeniae*, nomen nudum, synonymy in Schedl 1972f: 154), Schedl 1957d: 137 (*pygmaeus*, nomen nudum, synonymy in Schedl 1972f: 154). References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1985: 28. **(cn)** Thompson, G. H. 1960. **(ec)** Roberts 1969a: 110; Schedl 1962k: 776. **(hb)** Beaver & Loytyniemi 1985b: 126; Browne 1963a: 260; Jones, Roberts, & Baker 1959: 13–14; Loytyniemi, Beaver, & Loytyniemi 1985: 28; Schedl 1962k: 776; Thompson, G. H. 1963: 41; Webb & Jones 1957: 37–40. **(ds)** Beaver & Loytyniemi 1985b: 126; Browne 1963a: 260; Ferreira 1965: 1132; Gardner 1957a: 28; Mayne & Donis 1960: 96; Roberts 1968: 190; Schedl 1933f: 197, 1959p: 23, 1959q: 706, 1962k: 776, 1964j: 619, 1967e: 221, 1969a: 204, 1971g: 195, 1972f: 154; Strohmeier 1911f: 204, 1912c: 6, 1914c: 34; Thompson, G. H. 1960, 1963: 41. **(tx)** Chapuis

- 1865: 92; Numberg 1966a: 186; Sampson 1924c: 126; Schaufuss 1905: 103; Schedl 1933f: 197, 1936b: 131, 1937d: 35, 1941e: 154, 1953g: 244, 1954e: 48, 56–57, 1955i: 213, 1957d: 131, 1960: 12, 1962k: 776, 1113, 1972f: 154; Strohmeier 1911f: 204, 1912c: 6–7, 1912f: 86, 1914c: 34.
- eximius** Schedl 1941d: 407. Holotype ♂; D.O.A., Nyembe; Schedl Collection in NHMW, Wien.
Distribution: Africa (Ivory Coast/ Tanzania/ Zaire/ Zambia).
Hosts: *Piptadenia africana*.
Notes: (3) Beaver et al. 1955: 130 (described female).
References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (cc) Jover 1952: 73–81. (hb) Jover 1952: 73–81; Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 130; Cachan 1957: 6, 15; Schedl 1962k: 931, 1965g: 21, 1972f: 160. (tx) Numberg 1966a: 187; Schedl 1941d: 406–407, 1951f: 40, 1952g: 55, 1954d: 573, 1962k: 931, 1965g: 21, 1972f: 160, 1978a: 29.
- expletus** Schedl 1941d: 412. Holotype ♀; Kamerun; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon/ Ruanda/ Zaire).
Hosts: Schedl 1962k: 836 (13 host species listed).
Notes: (3) Schedl 1951i: 58 (described female).
References: (cc) Schedl 1962k: 834. (hb) Schedl 1962k: 834. (ds) Mayne & Donis 1960: 96, 1962: 270; Schedl 1962k: 834, 1972f: 158. (tx) Numberg 1966a: 187; Schedl 1941d: 412, 1951i: 58, 1957d: 134, 1962k: 834, 1972f: 158, 1978a: 29.
- falsificus** (Schedl) 1933f: 201 (*Crossotarsus*).
Lectotype ♂; Mayumbe, Soc. Bomeco, and Lucalla, Cie Ind. Bois Congo; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 30.
Distribution: Africa (Nigeria/ Zaire).
Hosts: *Millettia* sp.
References: (cn) Vrydagh 1947a: 8. (cc) Roberts 1969b: 110. (ds) Browne 1980d: 493; Roberts 1965: 190; Schedl 1933f: 202, 1962k: 838, 1972f: 158; Vrydagh 1947a: 8. (tx) Schedl 1933f: 201–202, 1937c: 15, 1950d: 5, 1952i: 56, 1962k: 838, 1972f: 158, 1978a: 31.
- forcipatus** Schedl 1955c: 33. Lectotype ♂; Africa Orient. Ital.: Sagan-Omo, Caschei; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 31.
Distribution: Africa (Mozambique/ Somalia).
References: (ds) Schedl 1972f: 154. (tx) Numberg 1966a: 186; Schedl 1955c: 33, 1962k: 932, 1972f: 154, 1978a: 31.
- fulgens** Schedl 1951i: 51. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.
Distribution: Africa (Nigeria/ Uganda/ Zaire).
Hosts: *Chrysophyllum lacourtianum*, *Cynometra alexandri*, *Dacryodes yangambiensis*.
References: (ds) Gardner 1957a: 28; Mayne & Donis 1962: 270; Roberts 1965: 181, 1969b: 110; Schedl 1962k: 879, 1972f: 161. (tx) Numberg 1966a: 186; Schedl 1952i: 51, 1957d: 140, 1962k: 879, 1972f: 161, 1978a: 32.
- fuscipilosulus** (Schedl) 1936b: 133 (*Crossotarsus*).
Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon).
References: (ds) Schedl 1972f: 150. (tx) Schedl 1936b: 133, 1937c: 14–15, 1962k: 932, 1972f: 150, 1978a: 32.
- galerus** (Schedl) 1936b: 131 (*Crossotarsus*).
Holotype ♀; Franz. Kongo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Congo/ Gabon/ Zaire).
Hosts: *Allogogus brunneus*, *Combretum oblongum*.
References: (cc) Schedl 1962k: 779. (ds) Schedl 1962k: 779, 1972f: 154. (tx) Numberg 1966a: 186; Schedl 1936b: 131, 1962k: 779, 1972f: 154, 1978a: 32.
- ghesquierei** (Schedl) 1933f: 198 (*Crossotarsus*).
Lectotype ♂; Wendji, Sankuru, Haut-Uele, Moto; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 32.
Figures: Schedl 1962k: 887, 1972f: 163.
Distribution: Africa (Cameroon/ Congo/ Ivory Coast/ Kenya/ Malawi/ Nigeria/ Zaire).
Hosts: Schedl 1962k: 887–889 (39 host species listed).
Notes: (3) Schedl 1952i: 57 (described female).
References: (ay) Lhoste & Roche 1960. (cc) Jover 1952: 73–81; Schedl 1962k: 886. (hb) Jover 1952: 73–81; Schedl 1962k: 886. (ds) Browne 1972c: 100; Cachan 1957: 6, 15; Gardner 1957a: 28; Mayne & Donis 1960: 96, 1962: 270; Roberts 1965: 191, 1969b: 110; Schedl 1933f: 198, 200, 1962k: 886, 1966c: 231, 1972f: 162. (tx) Numberg 1966a: 186; Powell, W. 1957: 27; Schedl 1933f: 198, 200, 1937d: 35, 1941d: 381, 1950e: 212, 1952i: 57, 1954d: 873, 1955i: 213, 1956a: 35, 1962k: 886–887, 1114, 1972f: 162–163, 1978a: 32.
- glaber** Schedl 1962k: 1114. Holotype ♂; Congo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: (ds) Schedl 1972f: 162. (tx) Schedl 1962k: 1114, 1972f: 162, 1978a: 32.
- gracilior** Schedl 1954d: 883. Lectotype ♂; Cote d'Ivoire, Lagune Ono; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 33.
Distribution: Africa (Ghana/ Ivory Coast/ Nigeria/ Zaire).
Hosts: *Albizzia zygia*, *Pentaclethra macrophylla*, *Piptadenia africana*, *Theobroma cacao*.
References: (hb) Roberts 1969b: 110; Thompson, C. H. 1963: 41. (ds) Schedl 1962k: 779, 1972f: 154; Thompson, C. H. 1963: 41. (tx) Numberg 1966a: 186; Schedl 1954d: 873, 883, 1954e: 56, 1962k: 779, 1972f: 154, 1978a: 33.
- guineensis** Roberts 1965: 223. Holotype ♂; Nigeria: Oban; BMNH, London.

- Distribution: Africa (Nigeria).
Hosts: *Strombosia pustulata*.
References: (ds) Roberts 1968: 191; Schedl 1972f: 154. (tx) Roberts 1965: 223; Schedl 1972f: 154.
- hartwegi** Schedl 1979d: 452. Holotype ♂; Kamerun, Oku-See; Alexander Koenig Museum, Bonn.
Distribution: Africa (Cameroon).
References: (tx) Schedl 1979d: 452.
- ibex** Schedl 1941d: 411. Holotype ♂; St. Thome Isle; Schedl Collection in NHMW, Wien.
Figures: Schedl 1962k: 956, 1972f: 161.
Distribution: Africa (Sao Tome Island).
References: (ds) Schedl 1972f: 162. (tx) Numberg 1966a: 187; Schedl 1941d: 411, 1962k: 956, 1972f: 161–162, 1978a: 36.
- incilis** Schedl 1957d: 143. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.
Distribution: Africa (Nigeria/ Zaïre).
Hosts: *Cynometra hankci*.
References: (ds) Roberts 1968: 191; Schedl 1962k: 923, 1972f: 148. (tx) Numberg 1966a: 186; Schedl 1957d: 143, 1962k: 923, 1972f: 148, 1978a: 37.
- insitirus** Schedl 1971e: 15. Holotype ♂; Fernando Poo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Fernando Po Island/ Ivory Coast/ Zaïre).
References: (ds) Schedl 1971e: 15, 1972f: 154. (tx) Schedl 1972f: 154, 1978a: 37.
- integratus** Roberts 1964: 409. Holotype ♂; Nigeria; Adauire; BMNH, London.
Distribution: Africa (Nigeria).
Hosts: *Crossandra* sp.
Notes: (3) Schedl 1972f: 151 (treated as a synonym of *ukerweensis*), Beaver & Loytyniemi 1955b: 125 (a good species).
References: (tx) Roberts 1964: 409, 1968: 192; Roberts & Loytyniemi 1955b: 125; Schedl 1972f: 151.
- interjectus** Schedl 1941d: 409. Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon/ Ghana/ Ivory Coast/ Nigeria/ Sierra Leone/ Uganda/ Zaïre).
Hosts: Schedl 1962k: 840 (19 host species listed).
References: (cn) Thompson, G. H. 1960. (ec) Roberts 1969a: 111. (hb) Schedl 1962k: 839; Thompson, G. H. 1963: 41. (ds) Browne 1950b: 381; Cachan 1957: 5, 15; Mayne & Donis 1962: 271; Roberts 1968: 190; Schedl 1961m: 85, 1962h: 62, 1962k: 839, 1964j: 43, 1965e: 355, 1972f: 158; Thompson, G. H. 1960, 1963: 41. (tx) Numberg 1966a: 187; Schedl 1941d: 408–409, 1950e: 215–216, 1954d: 873, 1954e: 56, 1957d: 138, 1961m: 85, 1962h: 62, 1962k: 839, 1115, 1962q: 494, 1972f: 158, 1978a: 37.
- interruptus** (Sampson) 1924c: 124 (*Crossotarsus*).
Holotype ♂; Belgian Congo, Elisabethville (Miss. Agric.); MRCB, Tervuren.
Distribution: Africa (Angola/ Ghana/ Guinea/ Kenya/ Tanzania/ Uganda/ Zaïre/ Zambia/ Zimbabwe).
Hosts: *Acacia* sp., *Antiaris* sp., *Baikava* sp., *Brachystegia* sp.
Notes: (3) Schedl 1936b: 132 (described female).
References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 131; Ferreira 1965: 1132; Gardner 1957a: 28; Schedl 1933f: 198–199, 1959p: 23, 1962k: 890, 1972f: 162, 1975h: 351. (tx) Numberg 1966a: 186; Sampson 1924c: 124–126; Schedl 1933f: 198–199, 1936b: 131–133, 1937b: 406–407, 1937d: 35, 1950d: 12, 1952i: 26–27, 1961d: 178–179, 1962k: 890, 1965g: 21, 1972f: 162, 1978a: 38.
- intritus** (Schedl) 1939g: 167 (*Crossotarsus*). Syn-types ♂ ♀; Insel Ukerewe [Schedl 1979: 38]; Conrad Collection and Schedl Collection in NHMW, Wien.
Distribution: Africa (Ivory Coast/ Tanzania/ Uganda/ Zaïre).
Hosts: *Cynometra alexandrii*, C. sp., *Parinarium curatellifolium*.
Notes: (1) Schedl 1978a: 38 (citation of holotype invalid).
References: (ds) Schedl 1959q: 706, 1962k: 750, 1972f: 154. (tx) Numberg 1966a: 186; Schedl 1939g: 167, 1953g: 244, 1957d: 132, 1962k: 750, 1972f: 154, 1978a: 38.
- jurioni** Schedl 1952i: 37. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.
Figures: Schedl 1962k: 933.
Distribution: Africa (Cameroon/ Nigeria/ Zaïre).
Hosts: *Allophylbus africanus*, *Alstonia boone*, *Blighia welwitschii*, *Carapa procera*, *Celtis brieji*, *Drypetes leonensis*, *Guarea cedrata*, *Microcos pinatifida*, *Microdesmis puberula*, *Pancovia harnsiana*, *P. laurentii*, *Pterygota bequaertii*.
Notes: (3) Schedl 1957: 139 (described female).
References: (ec) Schedl 1962k: 932. (hb) Schedl 1962k: 932. (ds) Mayne & Donis 1960: 96, 1962: 271; Roberts 1968: 192, 1969b: 111; Schedl 1962k: 932, 1972f: 160. (tx) Numberg 1966a: 186; Schedl 1952i: 37, 1953g: 244, 1957d: 139, 1962k: 932–933, 1964j: 48, 1972f: 160, 1978a: 39.
- kakaoensis** Schedl 1941d: 408. Holotype ♂; Kamerun, Victoria Co.; Schedl Collection in NHMW, Wien [data not given in original publication, added in Schedl 1978a: 39].
Distribution: Africa (Cameroon/ Ghana/ Zaïre).
Hosts: Kakao [presumably *Theobroma cacao*].
References: (ds) Schedl 1972e: 255, 1972f: 158. (tx) Numberg 1966a: 187; Schedl 1941d: 408, 1962k: 843, 1972f: 158, 1978a: 39.

karii **Beaver** in Beaver & Loytyniemi 1955b: 127. Holotype ♂; Zambia: Chati, Kitwe; BMNH, London.

Figures: Beaver & Loytyniemi 1955b: 127.

Distribution: Africa (Zambia).

References: **(bv)** Loytyniemi, Beaver, & Loytyniemi 1955: 28. **(hb)** Loytyniemi, Beaver, & Loytyniemi 1955: 28. **(tx)** Beaver & Loytyniemi 1955b: 127.

lateralis **Schedl** 1952i: 35. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 782, 1972f: 152.

Distribution: Africa (Nigeria/ Togo/ Zaire).

Hosts: *Anonidium mannii*, *Guarea laurentii*, *Macaranga* sp., *Mammea africana*, *Ongokea gore*, *Pterygota bequaertii*, *Scorodophloeus zenkeri*. Schedl 1962k: 781–782 (23 host species listed).

References: **(cc)** Schedl 1962k: 781. **(hb)** Schedl 1962k: 781. **(ds)** Mayne & Donis 1960: 96, 1962: 271; Roberts 1968: 190, 1969b: 111; Schedl 1962k: 781, 1972f: 154. **(tx)** Nunberg 1966a: 186; Schedl 1952i: 35, 1957d: 141, 1962k: 781–782, 1972f: 152, 154, 1978a: 40.

lebruni **Schedl** 1952i: 40. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 845.

Distribution: Africa (Nigeria/ Sierra Leone/ Zaire).

Hosts: Schedl 1962k: 844–847 (48 host species listed).

References: **(cc)** Schedl 1962k: 844. **(hb)** Schedl 1962k: 844. **(ds)** Mayne & Donis 1960: 271; Roberts 1968: 190, 1969b: 11; Schedl 1962k: 844, 1972f: 158. **(tx)** Nunberg 1966a: 187; Schedl 1952i: 40–44, 1962k: 844–845, 1972f: 158, 1978a: 40.

lecomtei **Schedl** 1957d: 134. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Fernando Po Island/ Nigeria/ Zaire).

Hosts: *Angylocalyx pynaertii*, *Guarea laurentii*, *Strombosia grandiflora*.

References: **(cc)** Schedl 1962k: 935, 1972c: 60. **(hb)** Schedl 1962k: 935. **(ds)** Roberts 1968: 192, 1969b: 111; Schedl 1962k: 935, 1971g: 195, 1972f: 150. **(tx)** Nunberg 1966a: 187–194; Schedl 1957d: 134, 1962k: 935, 1972f: 150, 1978a: 40.

leferrei **Schedl** 1952i: 53. Holotype ♂; Congo Belge: Kivu, Mulungu; MRCB, Tervuren.

Figures: Mayne & Donis 1962: 272, Schedl 1962k: 785–786, 1972f: 152.

Distribution: Africa (Nigeria/ Zaire).

Hosts: Schedl 1962k: 786–788 (53 host species listed).

References: **(hb)** Schedl 1962k: 785. **(ds)** Mayne & Donis 1960: 96, 1962: 272; Roberts 1968: 190, 1969b: 111–112; Schedl 1962k: 785, 1972f: 154. **(tx)** Mayne & Donis 1962: 272; Nunberg 1966a: 186; Schedl 1952i: 53, 1962k: 785–786, 1972f: 152, 154, 1978a: 40.

lobatus **(Schedl)** 1936b: 128 (*Crossotarsus*). Syn-types ♂♀; Ostafrika: Schedl Collection in NHMW, Wien.

Distribution: Africa (Kenya/ Tanzania/ Zaire/ Zambia).

Hosts: *Acacia lahai*, *A.* spp.

Notes: (1) Schedl 1978a: 41 (citation of holotype invalid).

References: **(ds)** Beaver & Loytyniemi 1955b: 128; Schedl 1938d: 451, 1962k: 789, 1966c: 232, 1971e: 5, 1972f: 155. **(tx)** Nunberg 1966a: 186; Schedl 1936b: 128–129, 1938d: 451, 1939g: 167, 1962k: 789, 1972f: 155, 1978a: 41.

malkini **Schedl** 1956a: 34. Holotype ♂; Nigeria, Ogbomoshos; CAS, San Francisco.

Distribution: Africa (Ghana/ Ivory Coast/ Nigeria/ Sierra Leone).

Hosts: *Cassia* sp., *Sterculia rhinopetala*.

Notes: (3) Schedl 1962k: 1115 (described female).

References: **(hb)** Browne 1963a: 261. **(ds)** Browne 1963a: 261; Roberts 1968: 190, 1969b: 112; Schedl 1962h: 62, 1962k: 789, 1964j: 43, 1972e: 285, 1972f: 155. **(tx)** Nunberg 1966a: 186; Schedl 1956a: 34, 1962h: 72, 1962k: 789, 1115, 1972f: 155, 1978a: 42.

maynei **Schedl** 1952i: 45. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 881.

Distribution: Africa (Zaire).

Hosts: *Chrysophyllum pruniferum*, *Erythrophloeum guineense*, *Pterocarpus soyauxii*, *Scorodophloeus zenkeri*, *Strombosia grandifolia*, *S. scheffleri*.

References: **(ds)** Mayne & Donis 1960: 96, 1963: 272; Schedl 1962k: 880, 1972f: 162. **(tx)** Nunberg 1966a: 186; Schedl 1952i: 45, 1957d: 140, 1962k: 880–881, 1972f: 162, 1978a: 43.

medius **Schedl** 1952i: 28. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 923, 1972f: 147.

Distribution: Africa (Angola/ Cameroon/ Congo/ Gabon/ Ghana/ Kenya/ Nigeria/ Sierra Leone/ Zaire/ Zambia/ Zimbabwe).

Hosts: Schedl 1962k: 913–916 (60 host species listed).

Notes: (1) Originally named as a subspecies of *rapax*. (3) Schedl 1972f: 147 (treated as a subspecies of *brevis*). Browne 1972c: 125 (a good species).

References: **(ds)** Ferreira 1965: 1131; Mayne & Davis 1960: 96–97, 1962: 275; Roberts 1968: 191, 1969b: 107; Schedl 1959p: 22, 1962k: 913–918, 1964f: 619, 1965e: 358, 1967e: 221, 1969a: 205, 1972e: 285; Thompson, C. H. 1963: 43. **(tx)** Browne 1972c: 125; Schedl 1952i: 28–30, 1952j: 7, 1954e: 57–69, 1957d: 142–143, 1962h: 61, 1962k: 913–918, 1110, 1972f: 147, 1978a: 61.

megatoma **Schedl** 1952i: 31. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Cameroon/ Nigeria/ Uganda/ Zaire).

Hosts: Schedl 1962k: 924–925 (31 host species listed).

References: (cc) Schedl 1962k: 923. (ds) Mayne & Donis 1960: 96, 1962: 273; Roberts 1968: 191, 1969b: 187; Schedl 1962k: 923, 1972f: 148. (tx) Nunberg 1966a: 187; Schedl 1952i: 31, 1957d: 143, 1962k: 923, 1972f: 148–149, 1978a: 43.

megatoma vexator Schedl 1957d: 144. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren. Notes: (3) Schedl 1978a: 43 (treated as a subspecies of *megatoma*).

References: (ds) Mayne & Donis 1962: 273; Roberts 1968: 191, 1969b: 115. (tx) Schedl 1957d: 143–144, 1962k: 925, 1119, 1972f: 148, 1978a: 43.

nimicus (Schedl) 1933f: 198 (*Crossotarsus*). Lectotype ♂; Katanga: Mulando; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 43. Figures: Browne 1972c: 119.

Distribution: Africa (Angola/ Cameroon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Malawi/ Mozambique/ Nigeria/ Sierra Leone/ South Africa/ Tanzania/ Zaire/ Zambia/ Zimbabwe).

Hosts: *Albizia* spp., *Brachystegia* spp., *Bussea occidentalis*, *Celtis* spp., *Entandrophragma utile*, *Pterygota macrocarpa*, *Triplochiton scleroxylon*.

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Jones, Roberts, & Baker 1959: 13–53; Loytyniemi, Beaver, & Loytyniemi 1985: 28. (ds) Browne 1972c: 119; Ferreira 1965: 1132; Gardner 1957a: 28; Schedl 1933f: 198, 1938d: 451, 1962k: 891, 1972f: 162; Tordo 1957: 10. (tx) Browne 1972c: 119; Nunberg 1966a: 186, 197; Powell, W. 1987: 27; Schedl 1933f: 198–199, 1936b: 127, 1941e: 154, 1950d: 4, 12, 1952g: 54, 1952i: 48, 1953g: 244, 1957b: 152–153, 1962k: 891, 1965c: 52, 1972f: 162, 1978a: 43, 77.

unispinosus Schedl 1936: 127 (*Crossotarsus*). Holotype ♂; Guinea; Schedl Collection in NHMW, Wien. Synonymy: Browne 1972c: 119, Schedl 1978a: 77.

Notes: (3) Schedl 1954e: 84 (described female).

References: (cc) Roberts 1969b: 115. (hb) Browne 1962: 94, 1963a: 263, 1968: 249; Roberts 1961c: 37; Thompson, G. H. 1963: 43. (ds) Browne 1961: 39–43, 1963a: 263, 1972c: 119, 1968: 249; Cola 1971; Ferreira 1965: 1133; Roberts 1968: 191, 1969b: 115; Schedl 1959p: 23, 1962h: 63, 1962k: 895, 1972e: 285, 1972f: 162, 1975b: 351, 1982: 279; Thompson, G. H. 1963: 43. (tx) Browne 1972c: 119; Nunberg 1959: 167, 1966a: 186; Schedl 1936b: 126–127, 1937b: 406, 1954e: 57, 84, 1955d: 270, 1960h: 111, 1962k: 895, 1119, 1972f: 162, 1978a: 77.

unispinosus Schedl 1937b: 406. Lectotype ♂; N.

Rhodesia, N'Changa; Schedl Collection in NHMW, designated by Schedl 1978a: 77. Synonymy: Schedl 1972f: 162.

Notes: (3) This synonym resulted from inadvertent double naming by Schedl.

References: (tx) Schedl 1937b: 406, 1972f: 162, 1978a: 77.

minimus Schedl 1957d: 131. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Nunberg 1966b: 197.

Distribution: Africa (Nigeria/ Zaire).

Hosts: *Dialium corbisieri*, *Terminalia superba*.

References: (ds) Roberts 1969b: 112; Schedl 1962k: 790, 1972f: 155. (tx) Nunberg 1966b: 186, 197; Schedl 1957d: 131, 1962k: 790, 1972f: 155, 1978a: 43.

minor Schedl 1952i: 28. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Nigeria/ Zaire).

Hosts: Schedl 1962k: 919–920 (25 host species listed).

Notes: (1) Originally named as a subspecies of *rapax*. (3) Schedl 1972f: 147 (treated as a subspecies of *brevis*), Browne 1972c: 125 (a good species).

References: (ds) Browne 1980c: 486; Roberts 1968: 191. (tx) Browne 1972c: 125; Schedl 1952i: 28–30, 1957d: 142–143, 1962k: 919, 1972f: 147, 1972g: 83.

minutissimus (Schedl) 1935g: 316 (*Crossotarsus*). Holotype ♂; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon/ Equatorial Guinea/ Ghana/ Guinea/ Sierra Leone/ Tanzania/ Zaire).

Hosts: *Celtis mildbraedii*, *Guarea thompsonii*, *Occhioecosmus africanus*.

Notes: (3) Roberts 1970b: 478 (described female). References: (ds) Browne 1980b: 382; Schedl 1962h: 62, 1962k: 937, 1971f: 195, 1972f: 150, 1975h: 352. (tx) Nunberg 1966a: 187; Roberts 1970b: 478; Schedl 1935g: 316, 1957d: 136, 1962k: 936–937, 1972f: 150, 1978a: 44.

mitratus Nunberg 1966a: 192. Holotype ♀; Congo: Elisabethville; MRCB, Tervuren.

Figures: Nunberg 1966a: pl. 2, figs. 4–6.

Distribution: Africa (Zaire).

References: (tx) Nunberg 1966a: 192.

modestus Beaver in Beaver & Loytyniemi 1985: 123. Holotype ♂; Zambia: Chati, Kitwe; BMNH, London.

Distribution: Africa (Zambia).

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (tx) Beaver & Loytyniemi 1985: 123.

montanus Roberts 1966b: 626. Holotype ♂; Nigeria: Cyel Nyaki (Mambilla Plateau), Sardauna Prov.; BMNH, London.
Distribution: Africa (Nigeria).
References: (ds) Browne 1980b: 382; Roberts 1966b: 626, 1968: 192; Schedl 1972f: 150. (tx) Schedl 1972f: 150.

multipunctus Roberts 1964: 405. Holotype ♂; Nigeria: Idanre; BMNH, London.
Distribution: Africa (Nigeria).
References: (ds) Roberts 1968: 190; Schedl 1972f: 155. (tx) Roberts 1964: 405; Schedl 1972f: 155.

nairobiensis (Schedl) 1937c: 15 (*Crossotarsus*). Lectotype ♂; British East Africa: Nairobi; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 45.
Distribution: Africa (Ghana/ Kenya/ Tanzania).
Hosts: *Antiaris africana*, *Baphia nitida*, *Cassipourea* sp., *Cordia* sp., *Dombeya* sp., *Ekebergia ruppeliana*, *Eutandophragma cylindricum*, *Eucalyptus* sp., *Mansonia altissima*, *Polyscias* sp., *Turracanthus vignei*, *Warburgia ugandensis*.
References: (cn) Anonymous 1953j: 25; Curry 1958b: 128–132; Thompson, G. H. 1960. (cc) Anonymous 1953j: 25; Jover 1952: 73–81. (hb) Jones, Roberts, & Baker 1959: 13, 21; Jover 1952: 73–81; Thompson, G. H. 1963: 42; Webb & Jones 1957: 25–40. (ds) Gardner 1957a: 29; Schedl 1962k: 847, 1972f: 158; Thompson, G. H. 1960, 1963: 42. (tx) Numberg 1966a: 187; Powell, W. 1987: 27; Schedl 1937c: 15, 1941d: 409, 1950e: 212, 215, 1952j: 6, 1954d: 573, 1954e: 56, 1957d: 138, 1962k: 847, 1972f: 158, 1978a: 45.

nanus (Schedl) 1937d: 37 (*Crossotarsus*). Syntypes ♀; Kamerun; MNB, Berlin, Hamburg Museum, lost, Schedl Collection in NHMW, Wien.
Distribution: Africa (Cameroon/ Tanzania/ Togo).
Notes: (1) Schedl 1978a: 46 (type restricted to MNB, Berlin specimens). (3) Schedl 1972f: 147 (treated as a subspecies of *brevis*).
References: (tx) Numberg 1966a: 187; Schedl 1937d: 37, 1941d: 409, 1952j: 6, 1954e: 48, 1957d: 118, 142, 144, 1962k: 922, 1972f: 147.

neoconjunctus Schedl 1962k: 1115. Holotype ♂; Congo; Schedl Collection in NHMW, Wien.
Distribution: Africa (Zaire).
References: (ds) Schedl 1972f: 162. (tx) Schedl 1962k: 1115, 1972f: 162, 1978a: 46.

nigeriensis Roberts 1971a: 283. Holotype ♂; Nigeria: Boshi Extension Forest Reserve, Obudu Plateau, 5000 feet; BMNH, London.
Distribution: Africa (Nigeria).
Hosts: *Garcinia guetoides*.
References: (tx) Roberts 1971a: 283.

nitens Browne 1970: 581. Holotype ♂; Tanzania: Lushoto, Shume Forest Reserve; BMNH, London.
Distribution: Africa (Tanzania).
Hosts: *Albizia maranguensis*.
References: (ds) Schedl 1972f: 158. (tx) Browne 1970: 581; Schedl 1972f: 158.

nitidulus Schedl 1957d: 141. Holotype ♂; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.
Distribution: Africa (Sierra Leone/ Zaire).
Hosts: *Bussca occidentalis*, *Carapa grandiflora*.
Notes: (3) Schedl 1971e: 15 (described female).
References: (ds) Schedl 1962k: 790, 1971e: 4, 1972f: 155. (tx) Numberg 1966a: 186; Schedl 1957d: 141, 1962k: 790, 1971e: 15, 1972f: 155, 1978a: 47.
leoneusis Roberts 1970b: 477. Holotype ♂; Sierra Leone: Cola West Forest Reserve; BMNH, London. Synonymy: Schedl 1972f: 155.
References: (tx) Roberts 1970b: 477; Schedl 1972f: 155.

notatus Schedl 1962k: 872. Holotype ♂; Gold Coast: MRCB, Tervuren.
Distribution: Africa (Ghana/ Nigeria/ Uganda/ Zaire).
Hosts: *Cynometra alexandri*, *Hymenocardia ulmoides*, *Pentaclethra macrophylla*, *Scorodophloeus zenkeri*, *Triplochiton scleroxylon*.
Notes: (1) Schedl 1978a: 48 (citation of holotype invalid). (3) Browne 1980e: 779 (described female).
References: (hb) Thompson, G. H. 1963: 42. (ds) Gardner 1957a: 29; Roberts 1968: 191; Schedl 1972f: 160; Thompson, G. H. 1963: 42. (tx) Browne 1980e: 779; Numberg 1966a: 188; Schedl 1950d: 35, 1952i: 59, 1954e: 56, 1962k: 872, 1972f: 160, 1978a: 48.

obauensis Roberts 1964: 407. Holotype ♂; Nigeria: Obau; BMNH, London.
Distribution: Africa (Congo/ Nigeria).
References: (ds) Browne 1972c: 100; Roberts 1968: 190; Schedl 1972f: 155. (tx) Roberts 1964: 407; Schedl 1972f: 155.

obuduensis Roberts 1966b: 623. Holotype ♂; Nigeria: Obudu Cattle Ranch; BMNH, London.
Distribution: Africa (Nigeria).
Hosts: *Scottellia coriacea*, *Trichilia* sp.
Notes: (3) Schedl 1972f: 151 (treated as a subspecies of *strombosiopsis*).
References: (ds) Roberts 1966b: 623, 1968: 192; Schedl 1972f: 151. (tx) Roberts 1966b: 623; Schedl 1972f: 151.

occalleseus Schedl 1957d: 134. Holotype ♂; Ruanda: Ihembe; MRCB, Tervuren.
Distribution: Africa (Ruanda/ Zaire).
Hosts: *Drypetes ituriensis*, *Syzygium parvifolium*.
References: (ds) Mayne & Donis 1962: 273; Schedl 1962k: 849, 1972f: 158. (tx) Numberg 1966a: 187; Schedl 1957d: 134, 1962k: 849, 1972f: 158, 1978a: 49.

octodentatus **Browne** 1980e: 778. Holotype ♂; Zaire: Kisangani; MRCB, Tervuren.
Distribution: Africa (Zaire).
References: (tx) Browne 1980e: 778.

omissus **Schedl** 1957: 133. Holotype ♂; Congo Belge: Kivu, Hembe-Bitale; MRCB, Tervuren.
Figures: Schedl 1962k: 874, 1972f: 159.
Distribution: Africa (Zaire).
Hosts: *Radlkofera* sp.
Notes: (3) Schedl 1962k: 1116 (described female).
References: (ds) Schedl 1962k: 873, 1972f: 160. (tx) Numberg 1966a: 186; Schedl 1957d: 133, 1962k: 873–874, 1116, 1972f: 159–160, 1978a: 49.

opifex (**Sampson**) 1922a: 139 (*Crossotarsus*). Holotype ♂; Portuguese East Africa: Xinavane; BMNH, London.
Distribution: Africa (Equatorial Guinea/ Ghana/ Kenya/ Malawi/ Mozambique/ Nigeria/ Senegal/ Sierra Leone/ Sudan/ Tanzania/ Uganda/ Zaire/ Zambia/ Zimbabwe), Madagascar.
Hosts: *Acacia lahai*, *Azela africana*, *Albizia zygia*, *Antiaris africana*, *Bussea occidentalis*, *Hymenostegia afzelii*, *Panda oleosa*, *Piptadeniastrum africanum*, *Triplochiton scleroxylon*.
References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (hb) Browne 1963a: 261; Jones, Roberts, & Baker 1959: 13–53; Loytyniemi, Beaver, & Loytyniemi 1955: 28; Roberts 1969: 112; Thompson, G. H. 1963: 42; Webb & Jones 1957: 37, 40. (ds) Beaver & Loytyniemi 1955b: 128; Browne 1963a: 261, 1972c: 100; Gardner 1957a: 29; Mayne & Donis 1962: 273; Schedl 1933f: 197–198, 1938d: 451, 1962k: 791, 1965c: 52, 1965e: 359, 1968b: 145, 1971f: 150, 1971g: 195, 1972f: 155; Thompson, G. H. 1963: 42. (tx) Numberg 1966a: 186; Powell, W. 1987: 27; Sampson 1922a: 137–139, 1924c: 126; Schedl 1933f: 197–198, 1935g: 319, 1936b: 128, 1939g: 167–168, 1950d: 11, 1952g: 54, 1953g: 244, 246, 1954d: 883, 1954e: 56, 1957d: 131, 1962k: 791, 1117, 1972f: 155.

opulentus **Schedl** 1952i: 59. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
Distribution: Africa (Zaire).
Hosts: *Dacryodes yangambiensis*, *Pterocarpus soyauxii*.
References: (ds) Mayne & Donis 1962: 273; Schedl 1962k: 874, 1972f: 160. (tx) Numberg 1966a: 187; Schedl 1952i: 59, 1962k: 874, 1972f: 160, 1978a: 50.

oraelatus **Roberts** 1964: 403. Holotype ♂; Nigeria: Idanre; BMNH, London.
Distribution: Africa (Nigeria).
Hosts: *Ficus* sp.
References: (ds) Roberts 1968: 191; Schedl 1972f: 148. (tx) Roberts 1964; Schedl 1972f: 148.

perbrevis **Schedl** 1952i: 30. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
Figures: Schedl 1962k: 899.

Distribution: Africa (Congo/ Ghana/ Ivory Coast/ Kenya/ Malawi/ Nigeria/ Sierra Leone/ Uganda/ Zaire).

Hosts: *Azela africana*, *Albizia zygia*, *Blighia sapida*, *Celtis* spp., *Croton* sp., *Cynometra* sp., *Ficus exasperata*, *Khaya* spp., *Lonchocarpus sericeus*, *Mildbradiodendron excelsum*, *Tetrapleura tetraptera*, *Trema guineensis*, *Vernonia conferta*. Schedl 1962k: 901–903 (29 host species listed).

Notes: (3) Schedl 1957d: 142 and 1972f: 146 (treated as a subspecies of *brevis*), Browne 1972c: 147 (a good species).

References: (av) Entwistle 1963b. (bv) Entwistle 1963b. (cn) Thompson, G. H. 1960. (hb) Entwistle 1963b. (ds) Browne 1980a: 373; Cachan 1957: 15; Gardner 1957a: 29; Lee 1971: 29; Mayne & Donis 1960: 96, 1962: 274; Roberts 1968: 191, 1969b: 107; Schedl 1959q: 706, 1975h: 351; Thompson, G. H. 1960. (tx) Browne 1972c: 123; Powell, W. 1987: 27; Schedl 1952i: 30–31, 1954e: 85, 1955i: 213, 1957d: 142–143, 1959q: 706, 1962h: 61, 1962k: 899, 1972f: 146–147, 1978a: 52.

perbrevis solidus Schedl 1954e: 85. Lectotype ♂; Gold Coast: Koforidua; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53. Synonymy: Browne 1972c: 123.

Notes: (3) Schedl 1972f: 148 (treated as a subspecies of *brevis*).

References: (hb) Jones, Roberts, & Baker 1959: 13–51; Thompson, G. H. 1963: 42; Webb & Jones 1957: 35–40. (ds) Roberts 1968: 191, 1969b: 107; Thompson, G. H. 1963: 42. (tx) Browne 1972c: 123; Schedl 1954e: 56, 85, 1957d: 142–143, 1962h: 62, 1962k: 900, 1110, 1972f: 148, 1978a: 53.

perminutissimus (**Schedl**) 1936b: 130 (*Crossotarsus*). Lectotype ♀; Kamerun and Spanish Guinea; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 53.

Distribution: Africa (Cameroon/ Equatorial Guinea/ Ghana/ Nigeria/ Sierra Leone).

Hosts: *Albizia zygia*, *Cassia siamea*, *Parkia* sp., *Piptadenia africana*, *Piptadeniastrum africanum*.
Notes: (1) Schedl 1972f: 155 (to *Doliopygus*). (3) Schedl 1954e: 65 (described female), 1954e: 85 (described male).

References: (hb) Thompson, G. H. 1963: 42. (ds) Roberts 1968: 190, 1969b: 133; Schedl 1962k: 792, 1966a: 276, 1972f: 155; Thompson, G. H. 1963: 42. (tx) Numberg 1966a: 186; Schedl 1936b: 130, 1954e: 57, 65, 85, 1957d: 131, 1962k: 792, 1117, 1972f: 155, 1978a: 53.

praeclarus (**Schedl**) 1933f: 199 (*Crossotarsus*). Lectotype ♂; Katanga: Lubombo, Elisabethville, Dunga; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 56.

Distribution: Africa (Angola/ Cameroon/ Ivory

- Coast/ Mozambique/ Nigeria/ Tanzania/ Zaire/ Zambia/ Zimbabwe).
- Notes: (3) Schedl 1936b: 130 and 1941d: 407 (described female).
- References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 130; Ferreira 1965: 1132; Roberts 1968: 191, 1969b: 113; Schedl 1933f: 199, 1935d: 451, 1959p: 23, 1962h: 62, 1962k: 882, 1965g: 22, 1966c: 232, 1972f: 162, 1979b: 417. (tx) Numberg 1966a: 186; Schedl 1933f: 199, 1934a: 247, 1936b: 130, 1937b: 407, 1937d: 35, 1941d: 407, 1941e: 153, 1952g: 55, 1962h: 62, 1962k: 882, 1117, 1972f: 162, 1978a: 56.
- praemorsus (Dalman)** 1823: 82 (*Platypus*). Holotype ♂; Sierra Leone; Schoenherr Collection. Distribution: Africa (Sierra Leone). References: (ds) Schedl 1972f: 155. (tx) Dalman 1823: 82; Schedl 1972f: 155.
- prolongatus** Schedl 1962k: 1117. Holotype ♂; Senegal; Schedl Collection in NHMW, Wien. Distribution: Africa (Senegal). References: (ds) Schedl 1972f: 155. (tx) Schedl 1962k: 1117, 1972f: 155, 1978a: 57.
- propinquus** Schedl 1954e: 83. Lectotype ♂; Mpraeso; Schedl Collection in NHMW, Wien. Distribution: Africa (Ghana/ Guinea/ Ivory Coast/ Nigeria/ Sierra Leone/ Zaire). Hosts: *Albizia* spp., *Celtis mildbraedii*, *Cola* sp., *Entandrophragma utile*, *Hymenostegia afzchia*, *Mammea africana*, *Pentaclethra macrophylla*, *Piptadenia africana*, *Piptadeniastrum africanum*, *Strombosia glaucescens*. References: (cn) Anonymous 1953j: 25. (cc) Anonymous 1953j: 25. (hb) Browne 1963a: 261; Jones, Roberts, & Baker 1959: 13, 44–53; Thompson, G. H. 1963: 42; Webb & Jones 1957: 25–41. (ds) Browne 1963a: 261; Roberts 1968: 190; Schedl 1962h: 62, 1962k: 793, 1965e: 359, 1966a: 276, 1972f: 155; Thompson, G. H. 1963: 42. (tx) Numberg 1966a: 186; Schedl 1954e: 57, 83, 1962k: 793, 1972f: 155, 1978a: 57.
- proxilus** Schedl 1965e: 378. Holotype ♂; Nig. (vermutlich Nigerien); Schedl Collection in NHMW, Wien. Distribution: Africa (Nigeria). References: (ds) Schedl 1972f: 148. (tx) Schedl 1965e: 378, 1972f: 148, 1978a: 57.
- proximus** Schedl 1957d: 137. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren. Distribution: Africa (Cameroon/ Fernando Po Island/ Ghana/ Nigeria/ Zaire/ Zambia/ Zimbabwe). Hosts: *Bussea occidentalis*, *Guarea laurentii*, *Strombosia grandifolia*, *S. scheffleri*. Notes: (1) Originally named as a subspecies of *ukereveensis*. Beaver & Loytyniemi 1955: 124 (raised to species rank). References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (hb) Beaver & Loytyniemi 1955b: 124; Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 124; Mayne & Donis 1967: 278; Roberts 1968: 192; Schedl 1962h: 62, 1962k: 944, 1964f: 619, 1972e: 285, 1972f: 151. (tx) Beaver & Loytyniemi 1955b: 124; Schedl 1957d: 137, 1962h: 62, 1962k: 944, 1972f: 151, 1978a: 77.
- pseudoserratus** Roberts 1966b: 627. Holotype ♂; Nigeria; Akure; BMNH, London. Distribution: Africa (Nigeria). Hosts: *Cola gigantea*, *Oxytigma manni*, *Pterocarpus mildbraedii*, *Sterculia rhinopetala*. References: (ds) Browne 1980c: 485; Roberts 1968: 190; Schedl 1972f: 155. (tx) Roberts 1966b: 627; Schedl 1972f: 155.
- punctiventris** Schedl 1952i: 52. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren. Distribution: Africa (Cameroon/ Ghana/ Nigeria/ Sierra Leone/ Zaire). Hosts: Schedl 1962k: 850 (11 host species listed). *Alstonia congenensis*, *Klainedoxa gabonensis*, *Scorodophloeus zenkeri*. References: (cc) Schedl 1962k: 849. (hb) Schedl 1962k: 849; Thompson, G. H. 1963: 42. (ds) Mayne & Donis 1960: 96, 1962: 274; Roberts 1968: 190, 1969b: 113; Schedl 1962h: 62, 1962k: 849, 1964f: 619, 1964j: 43, 1965e: 359, 1972f: 158; Thompson, G. H. 1963: 42. (tx) Numberg 1966a: 187; Schedl 1952i: 52, 1954e: 57, 1962k: 849, 1972f: 158, 1978a: 59.
- pygmaeolus** Schedl 1957d: 137. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren. Distribution: Africa (Zaire). Hosts: *Combretum oblongum*, *Jubbernardia secreti*. References: (ds) Schedl 1962k: 937, 1972f: 150. (tx) Schedl 1957d: 137, 1962k: 937, 1966a: 187, 1972f: 150, 1978a: 59.
- rapax (Sampson)** 1924c: 127. Holotype ♂; S. Rhodesia; BMNH, London. Distribution: Africa (Angola/ Cameroon/ Ghana/ Ivory Coast/ Kenya/ Nigeria/ Ruanda/ Sierra Leone/ Uganda/ Zaire/ Zimbabwe). Hosts: Schedl 1962k: 909–911 (56 host species listed). Notes: (3) Schedl 1952i: 27 (described female), 1972f: 148 (treated as a subspecies of *brevis*). References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (cc) Jover 1952: 73–81; Schedl 1962k: 907–912, 1972c: 69. (hb) Beaver & Loytyniemi 1955b: 121; Jover 1952: 73–81; Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 121; Cachan 1957: 6, 15; Cola 1971; Gardner 1957a: 29; Mayne & Donis 1960: 96–97, 1962: 267; Numberg 1966a: 187; Roberts 1968: 191; Schedl 1937d: 37, 1938d: 451; Thompson, G. H. 1960. (tx) Browne 1972c: 123; Mayne & Donis 1960: 96, 1962: 274; Numberg 1966a: 187; Sampson 1924c: 127–128;

- Schedl 1937d: 37, 1941d: 381, 1950d: 5, 1951f: 40, 1952i: 27–31, 1952j: 6–7, 1954d: 873, 1957d: 123, 142–143; 1962k: 907, 1972f: 148, 1978a: 61. *perbrevis brevisculus* Schedl 1952i: 31. Holotype ♂; Congo Belge; Yangambi; Schedl Collection in NHMW, Wien. Synonymy: Browne 1972c: 123.
- Notes: (3) Schedl 1972f: 147 (treated as a subspecies of *brevis*).
- References: (hb) Thompson, G. H. 1963: 42. (ds) Ferreira 1965: 1131; Mayne & Donis 1960: 96, 1962: 274; Thompson, G. H. 1963: 42. (tx) Browne 1972c: 123; Schedl 1952i: 31, 1954e: 56–69, 1957d: 142–143, 1959p: 22, 1962k: 905, 1972f: 147, 1978a: 53.
- brevis interstitialis* Schedl 1957d: 143. Syntype ♂; Congo Belge, Camp Putnam [Schedl 1978a: 17]; Schedl Collection in NHMW, Wien. Synonymy: Browne 1972c: 123.
- Notes: (1) Schedl 1978a: 17 (citation of holotype invalid).
- References: (ec) Schedl 1962k: 912. (hb) Schedl 1962k: 912. (ds) Mayne & Donis 1962: 267; Roberts 1968: 191; Schedl 1962k: 912, 1972f: 147. (tx) Browne 1972c: 123; Schedl 1957d: 143, 1962k: 912, 1972f: 147, 1978a: 17.
- regularis** Schedl 1941d: 406. Holotype ♂; Isle Principe, Rocafn. Henrique; Schedl Collection in NHMW, Wien.
- Distribution: Africa (Principe Island in Gulf of Guinea).
- References: (ds) Schedl 1972f: 155. (tx) Nunberg 1966a: 186; Schedl 1936e: 51, 1941d: 406, 1962k: 794, 1972f: 155, 1978a: 62.
- retusus** Schedl 1957d: 131. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.
- Distribution: Africa (Angola/ Equatorial Guinea/ Ghana/ Nigeria/ Ruanda/ Sierra Leone/ Zaïre).
- Hosts: *Barteria nigritiana*, *Bussea occidentalis*, *Celtis adolphofrederici*, *Chrysophyllum* sp., *Combretodendron africanum*, *C. macrophyllum*, *Macaranga* sp., *Macropsis emini*, *Oxytigma oxyphyllum*, *Paramacrolobium coerulenum*.
- References: (ec) Schedl 1962k: 795. (hb) Browne 1963a: 261; Schedl 1962k: 795. (ds) Browne 1962d: 93, 1963a: 261; Ferreira 1965: 1132; Mayne & Donis 1962: 275; Roberts 1968: 190, 1969b: 113; Schedl 1959p: 23, 1962h: 62, 1962k: 795, 1965e: 359, 1971g: 195, 1972e: 285, 1972f: 155. (tx) Nunberg 1966a: 186; Schedl 1957d: 131, 1962k: 795, 1118, 1972f: 155, 1978a: 62.
- rhodesianus** (Schedl) 1937c: 14 (*Crossotarsus*). Syntypes ♀; Rhodesia: Salisbury; BMNH, London.
- Distribution: Africa (Cameroon/ Fernando Po Island/ Ghana/ Nigeria/ Tanzania/ Uganda/ Zaïre/ Zambia).
- Hosts: Schedl 1962k: 941–942 (20 host species listed).
- Notes: (1) Schedl 1978a: 62 (type restricted to BMNH specimens). (3) Schedl 1972f: 151 (treated as a subspecies of the junior name *ukereveensis*), Beaver et al. 1985: 125 (corrected name priority).
- References: (ds) Gardner 1957: 29. (tx) Schedl 1937c: 14–15, 1957b: 158, 1962k: 945, 1972f: 151, 1978a: 62.
- ukereveensis* Schedl 1939g: 168 (*Crossotarsus*). Syntypes ♂ ♀; Insel Ukerewe [Uganda] [Schedl 1978a: 77]; Conrad Collection, and Schedl Collection in NHMW, Wien. Synonymy: Beaver & Loytyniemi 1955: 124.
- Notes: (1) Schedl 1978a: 77 (citation of holotype invalid).
- References: (ec) Schedl 1962k: 940. (hb) Browne 1963a: 263; Schedl 1962k: 940. (ds) Browne 1963a: 263; Gardner 1957a; Mayne & Donis 1960: 98, 1962: 277; Schedl 1962k: 940, 1971g: 195, 1972f: 151. (tx) Nunberg 1966a: 187; Schedl 1939g: 168–169, 1957b: 152, 158, 1957d: 137, 1962k: 940, 1118, 1972f: 151, 1978a: 77.
- exasperatus** Roberts 1966a: 624. Holotype ♂; Nigeria: Awi; BMNH, London.
- Notes: (1) The status of this name is questionable; Schedl 1972f: 151 treated it as a synonym of *ukereveensis* subspecies *proximus*; Beaver et al. 1985: 125 treated it as a good species.
- References: (ds) Roberts 1968: 191. (tx) Roberts 1966a: 624; Schedl 1972f: 151.
- robustus** Browne 1972c: 113. Holotype ♂; Congo-Kinshasa; Yangambi; MRCB, Tervuren.
- Distribution: Africa (Zaire).
- References: (tx) Browne 1972c: 112–113.
- santiriae** Roberts 1970a: 68. Holotype ♂; Nigeria: Ngel Nyaki, Mambilla Plateau, 1500 m; MRCB, Tervuren.
- Distribution: Africa (Nigeria).
- Hosts: *Santiria trimera*.
- References: (tx) Roberts 1970a: 68.
- schoutedeni** (Sampson) 1924c: 123 (*Crossotarsus*). Holotype ♂; Belgian Congo: Elisabethville (Miss. Agric.); MRCB, Tervuren.
- Distribution: Africa (Angola/ Zaïre/ Zambia).
- Notes: (1) Schedl 1972: 150 (to *Doliopygus*).
- References: (bv) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1955: 28. (ds) Beaver & Loytyniemi 1955b: 124; Ferreira 1965: 1133; Schedl 1959p: 23, 1962k: 938, 1972f: 150. (tx) Nunberg 1966a: 187; Powell, W. 1987: 27; Sampson 1924c: 123–124; Schedl 1937b: 406, 1941d: 407, 1953g: 244, 1957d: 136, 1962k: 938, 1972f: 150.
- semipilosus** (Schedl) 1936b: 133 (*Crossotarsus*). Holotype ♀; Kongo; Schedl Collection in NHMW, Wien.
- Distribution: Africa (Zaire).
- Hosts: *Combretodendron africanum*, *C. macrophyllum*, *Pterocarpus soyauxii*, *Strombosia glaucescens*, *Tridesmostemon claussensii*.

Notes: (1) Schedl 1972f: 155 (to *Doliopygus*). (3) Schedl 1957d: 141 (described male).

References: (ds) Mayne & Donis 1962: 275; Schedl 1962k: 798, 1966c: 232, 1972f: 155. (tx) Numberg 1966a: 186; Schedl 1936b: 133–134, 1957d: 141, 1962k: 798, 1972f: 155, 1978a: 65.

serratulus Schedl 1953g: 246. Lectotype ♂; Congo Belge, Kasenyi, W. Darfur, N. Jebel Murra, Kurra, 5600 ft., and Senegal: Thies; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 65. Distribution: Africa (Ghana/ Senegal/ Sudan/ Zaire).

Notes: (3) Roberts 1971a: 285 (described female). References: (ds) Schedl 1962k: 938, 1968b: 145, 1972f: 155, 1975b: 351. (tx) Numberg 1966a: 186; Powell, W. 1987: 27; Roberts 1971a: 285; Schedl 1953g: 246, 1957d: 132, 1962k: 938, 1972k: 938, 1972f: 155, 1978a: 65.

serratus (Strohmeyer) 1911g: 226 (*Crossotarsus*). Syntypes ♂ ♀; Kamerun and Kilimandjaro; Strohmeyer Collection, and MNB, Berlin.

Distribution: Africa (Angola/ Bechuanaland/ Cameroon/ Congo/ Dahomey/ Equatorial Guinea/ Fernando Po Island/ Gabon/ Ghana/ Guinea/ Ivory Coast/ Kenya/ Nigeria/ Sierra Leone/ Tanzania/ Togo/ Uganda/ Zaire/ Zambia/ Zimbabwe).

Hosts: Schedl 1962k: 802–807 (118 host species listed).

Notes: (1) Schedl 1941d: 351 (to *Doliopygus*).

References: (ay) Entwistle 1963b; Francke-Grosmann 1956b: 275–308; Lhoste & Roche 1960; Strohmeyer 1918: 15, 21. (bv) Entwistle 1963b. (cn) Anonymous 1953j: 25; Browne 1965b: 248; Curry 1958b: 128–132; Mayne & Donis 1951: 335; Vrydagh 1947a: 9. (ce) Anonymous 1953j: 25; Francke-Grosmann 1956b; Jover 1952: 73–81; Mayne & Donis 1951: 335; Roberts 1960: 37–39, 1969b: 113; Schedl 1962k: 798. (hb) Beaver & Loytyniemi 1985b: 128; Browne 1963a: 262, 1968b: 248; Entwistle 1963b; Jover 1952: 73–81; Schedl 1962k: 798; Thompson, G. H. 1963: 43. (ds) Amaro & Gouveia 1957: 100, 130, 141; Beaver & Loytyniemi 1985b: 128; Browne 1962: 94, 1963a: 262, 1968b: 248, 1980b: 382; Cachan 1957: 5, 15, 43–53; Cola 1971; Ferreira 1965: 1133; Gardner 1957a: 29; Jones, Roberts, & Baker 1959: 13, 50; Mayne & Donis 1960: 98, 1962: 275; Menier 1973a; Roberts 1960b: 37, 1960e, 1960f, 1968: 190; Schedl 1933f: 196, 1959p: 23, 1962h: 62, 1962k: 798, 1964f: 619, 1964j: 43, 1965e: 359, 1965g: 22, 1965h: 112, 1966c: 232, 1967e: 221, 1971e: 2, 1971g: 195, 1972e: 285, 1972f: 156, 1975b: 351, 1979b: 417; Strohmeyer 1912c: 7, 1914c: 34; Thompson, G. H. 1963: 43; Vrydagh 1947a: 8; Webb & Jones 1957: 25–41. (tx) Numberg 1966a: 186; Sampson 1924c: 125; Schedl 1933f: 196, 1935g: 318–319, 1936b: 127–132, 1936e: 51, 1937b: 407, 1937d: 35, 1939b:

394, 1941d: 351, 406–407, 1941e: 153, 1950d: 4, 11, 14, 1950e: 212, 1951e: 39, 1951f: 40, 1952g: 54, 1952i: 27, 1953g: 244–245, 1954d: 873, 1954e: 48, 57, 1955d: 270, 1962k: 798, 1118, 1972f: 156; Strohmeyer 1911g: 225–227, 1912c: 7, 1914c: 34.

spatiosus (Schedl) 1935g: 316 (*Crossotarsus*). Lectotype ♂; Belgian Congo: 18 miles SW Elisabethville, Mozambique: Prov. de Gorongosa, Tendos du Sungoae et Riy Vunduzialt, N. Rhodesia: N'Changa, Tanganyika: Nord Rubcho; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 67.

Distribution: Africa (Bechuanaland/ Malawi/ Mozambique/ South Africa/ Tanzania/ Zaire/ Zambia/ Zimbabwe).

Hosts: *Brachystegia* sp.

Notes: (1) Schedl 1972f: 156 (to *Doliopygus*).

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (ds) Beaver & Loytyniemi 1985b: 129; Browne 1980c: 485; Lee 1971: 30; Schedl 1938d: 451, 1962k: 883, 1972f: 156. (tx) Numberg 1966a: 186; Powell, W. 1987: 27; Schedl 1935g: 316–318, 1936b: 126, 1937c: 15, 1941e: 154, 1962k: 883, 1972f: 156, 1978a: 67.

spectabilis Schedl 1952i: 44. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 884.

Distribution: Africa (Zaire).

Hosts: Schedl 1962k: 884–885 (11 host species listed).

References: (ds) Mayne & Donis 1960: 98, 1962: 276; Schedl 1962k: 884, 1972f: 162. (tx) Numberg 1966a: 186; Schedl 1952i: 44, 1962k: 884, 1972f: 162, 1978a: 67.

spinidens (Strohmeyer) 1911g: 225 (*Crossotarsus*). Holotype ♂; Sankuru-Fluss [Zaire]; Schedl Collection in NHMW, Wien.

Distribution: Africa (Zaire).

Notes: (1) Schedl 1972f: 156 (to *Doliopygus*).

References: (ds) Schedl 1972f: 156; Strohmeyer 1912c: 7, 1914c: 34. (tx) Numberg 1966a: 186; Schedl 1962k: 809, 1972f: 156; Strohmeyer 1911g: 225–226, 1912c: 7, 1914c: 34.

staneri Schedl 1952i: 33. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Figures: Schedl 1962k: 810.

Distribution: Africa (Zaire).

Hosts: *Canarium schweinfurthii*, *Entandrophragma* sp., *Guarea thompsonii*, *Tessmannia claessensii*.

References: (ds) Mayne & Donis 1960: 98, 1962: 277; Schedl 1962k: 810, 1972f: 156. (tx) Numberg 1966a: 186; Schedl 1952i: 33, 1962k: 810, 1972f: 156, 1978a: 68.

strombosipsis Schedl 1957d: 137. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Strombosiopsis tetrandra*.

References: (ds) Schedl 1962k: 939, 1967e: 221, 1972f: 151. (tx) Nunberg 1966a: 194; Schedl 1957d: 137, 1962k: 939, 1972f: 151, 1978a: 69.

subditicus (Schedl) 1935g: 319 (*Crossotarsus*). Syntypes ♂; Congo; A. Collart Collection, and Schedl Collection in NHMW, Wien.

Distribution: Africa (Angola/ Cameroon/ Central African Republic/ Ivory Coast/ Kenya/ Nigeria/ Zaïre).

Hosts: Schedl 1962k: 812–814 (28 host species listed).

Notes: (1) Schedl 1978a: 70 (citation of holotype invalid). (3) Schedl 1954d: 884 (described female).

References: (ec) Roberts 1969a; Schedl 1962k: 801. (hb) Gardner 1957c; Schedl 1962k: 811. (ds) Browne 1963a: 205, 1972c: 100; Ferreira 1965: 1133; Gardner 1957a: 29, 1957c: 292; Mayne & Donis 1960: 98, 1962: 277; Menier 1973a; Roberts 1968: 190, 1969b: 114; Schedl 1959p: 23, 1962k: 811, 1964f: 619, 1964j: 43, 1966c: 232, 1967e: 221, 1972f: 156, 1979b: 417. (tx) Nunberg 1966a: 186; Schedl 1935g: 319, 1952i: 38, 1954d: 874, 885, 1954e: 84, 1957d: 131, 1962k: 811, 1972f: 156, 1978a: 70.

subdolosus Schedl 1952i: 52. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Celtis brieji*, *Ficus* sp., *Maba laurentii*, *Piptadenia africana*.

References: (ds) Mayne & Donis 1962: 277; Schedl 1962k: 851, 1972f: 158. (tx) Browne 1970: 581; Nunberg 1966a: 187; Schedl 1952i: 52, 1962k: 851, 1972f: 158, 1978a: 70.

submarginatus Schedl 1962k: 873. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.

Distribution: Africa (Zaire).

Hosts: *Hymenocardia ulmoides*, *Pentaclethra macrophylla*, *Scorodophloeus zenkeri*.

Notes: (1) This was originally named as a subspecies of *notatus*, but was needlessly redescribed as a homonymous new species with a different holotype when it was raised to specific rank (Schedl 1971e: 16).

References: (ds) Schedl 1972f: 160. (tx) Schedl 1962k: 873, 1972f: 160, 1978a: 48.

submarginatus Schedl 1971e: 16. Holotype ♂; Ehemaliger Congo Belge, Yangambi; Schedl Collection in NHMW, Wien, preoccupied by Schedl 1962.

References: (tx) Schedl 1971e: 16.

subnotatus Schedl 1950d: 35. Holotype ♀; Kamerun; Schedl Collection in NHMW, Wien.

Distribution: Africa (Cameroon).

Notes: (3) Schedl 1972f: 148 (treated as a subspecies of *brevis*).

References: (tx) Nunberg 1966a: 187; Schedl 1950d: 35, 1952j: 6, 1957d: 142, 1962k: 907, 1972f: 148, 1978a: 70.

subserratus Browne 1980b: 359. Holotype ♂; Mtwara (Tanzania) to Nagoya (Japan), imported; BMNH, London.

Distribution: Africa (Tanzania).

Hosts: *Pterocarpus angolensis*.

References: (tx) Browne 1980b: 359.

tenuis (Strohmeyer) 1912f: 86 (*Crossotarsus*).

Holotype ♂; Mkulsumi-Berg (Deutsch-Ostafrika); Hamburg Museum, lost, and IPKE, Eberswalde.

Distribution: Africa (Cameroon/ Congo/ Dahomey/ Equatorial Guinea/ Ghana/ Guinea/ Ivory Coast/ Nigeria/ Sierra Leone/ Tanzania/ Uganda/ Zaïre).

Hosts: Schedl 1962k: 819–823 (66 host species listed).

Notes: (1) Schedl 1972f: 156 (to *Doliopygus*).

References: (ay) Menier 1976: 348. (bv) Menier 1976: 348. (ec) Schedl 1962k: 818. (hb) Anonymous 1959i: 18; Browne 1963a: 263; Schedl 1962k: 818; Strohmeyer 1911a: 182. (ds) Browne 1963a: 263; Cola 1973; Jones, Roberts, & Baker 1959: 13–53; Mayne & Donis 1951: 335–336, 1960: 277, 1962: 277; Menier 1973a; Roberts 1960b: 35, 1960e, 1960f: 39, 1968: 190, 1969b: 114; Schedl 1933f: 198, 1962h: 62, 1962k: 818, 1964f: 619, 1966c: 233, 1967e: 222, 1969a: 203, 1971g: 195, 1972f: 156, 1975h: 351; Spahr 1981: 79; Strohmeyer 1911a: 182, 1912c: 7, 1914c: 34. (tx) Nunberg 1966a: 186; Sampson 1924c: 126; Schedl 1933f: 197–198, 1939a: 468–469, 1941d: 381, 1941e: 154, 1950d: 4, 14, 1953g: 245, 1954d: 874, 1955d: 270, 1957d: 130–131, 1962k: 818, 1965c: 52, 1972f: 156, 1978a: 70, 72; Strohmeyer 1911a: 182, 1912e: 7, 1912f: 86, 1914c: 34.

terebrans Schedl 1964j: 48. Holotype ♂; Nigeria: Kano; BMNH, London.

Distribution: Africa (Angola/ Ghana/ Nigeria/ Senegal/ Sierra Leone/ Uganda/ Zaïre).

Hosts: *Acacia dudgeoni*, *Daniellia oliveri*, *Monotes kerstingii*.

References: (ec) Roberts 1969a. (hb) Browne 1963a: 263. (ds) Browne 1963a: 263; Roberts 1968: 192; Schedl 1962k: 939, 1972f: 160. (tx) Nunberg 1966a: 186; Schedl 1962k: 939, 1964j: 48, 1972f: 160, 1978a: 73.

togatus (Schedl) 1937c: 15 (*Crossotarsus*). Lectotype ♀; Rhodesia: Salisbury; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 74.

Distribution: Africa (Zambia/ Zimbabwe).

Notes: (1) Schedl 1972f: 148 (to *Doliopygus*). (3) Schedl 1972f: 148 (treated as a subspecies of *brevis*), Beaver & Loytyniemi 1985b: 122 (described male).

References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (ds) Beaver & Loytyniemi 1985b: 122. (tx) Schedl 1937c: 15, 1950d: 35, 1952j: 6, 1957d: 142, 1962k: 907, 1972f: 48, 1978a: 74.

trichiliae Schedl 1957d: 132. Holotype ♂; Congo Belge; Yangambi; MRCB, Tervuren.

- Distribution: Africa (Zaire).
 Hosts: *Trichilia rubescens*.
 References: (cc) Schedl 1962k: 940. (hb) Schedl 1962k: 940. (ds) Schedl 1962k: 940, 1972f: 156. (tx) Numberg 1966a: 186; Schedl 1957d: 132, 1962k: 940, 1972f: 156, 1978a: 74.
- tridens Roberts** 1970a: 69. Holotype ♂; Nigeria: Idanre Forest Reserve; MRCB, Tervuren.
 Distribution: Africa (Nigeria).
 References: (tx) Roberts 1970a: 69.
- ugandae Schedl** 1956a: 35. Holotype ♂; Uganda: Kampala, Mulange Mahira Forest, 4000 ft.; Schedl Collection in NHMW, Wien.
 Distribution: Africa (Uganda).
 References: (ds) Schedl 1972f: 162. (tx) Numberg 1966a: 186; Schedl 1956a: 32–35, 1962k: 885, 1972f: 162, 1978a: 76.
- uncinatus Schedl** 1953g: 247. Lectotype ♂; Congo Belge: Elisabethville; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 77.
 Distribution: Africa (Angola/ Tanzania/ Zaire/ Zambia).
 References: (bv) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (hb) Loytyniemi, Beaver, & Loytyniemi 1985: 28. (ds) Beaver & Loytyniemi 1985b: 129; Ferreira 1965: 1133; Schedl 1959p: 23, 1962h: 73, 1972f: 156. (tx) Numberg 1966a: 186; Schedl 1953g: 247, 1956a: 34, 1962k: 823, 1972f: 156, 1978a: 77.
- unicornis Schedl** 1952i: 48. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
 Distribution: Africa (Cameroon/ Congo/ Equatorial Guinea/ Ivory Coast/ Nigeria/ Zaire).
 Hosts: Schedl 1962k: 893–895 (45 host species listed).
 References: (cc) Schedl 1962k: 892. (hb) Schedl 1962k: 892. (ds) Browne 1984b: 287; Cola 1971; Mayne & Donis 1960: 98, 1962: 278; Roberts 1968: 191, 1969b: 115; Schedl 1962k: 892, 1964f: 619, 1966c: 233, 1967e: 222, 1972f: 162. (tx) Numberg 1966a: 186; Schedl 1952i: 48, 1954d: 874, 1957d: 130, 1962k: 892–895, 1119, 1972f: 162, 1978a: 77.
- unicus (Schedl)** 1935g: 319 (*Crossotarsus*). Syn-types ♂; Belgian Congo: Kasamvu; A. Collart Collection, and Schedl Collection in NHMW, Wien.
 Distribution: Africa (Kenya/ Nigeria/ Zaire).
 Hosts: *Celtis* sp., *Croton* spp.
 References: (ds) Gardner 1957a: 29; Roberts 1968: 190; Schedl 1962k: 851, 1972f: 158. (tx) Numberg 1966a: 187; Schedl 1935g: 319–320, 1962k: 851, 1972f: 158.
- vilis Schedl** 1962k: 1119. Holotype ♂; Kenya: Schedl Collection in NHMW, Wien.
 Distribution: Africa (Kenya/ Nigeria/ Uganda).
 Notes: (3) Roberts 1972: 72 (described female).
 References: (ds) Browne 1983a: 556, 1984b: 287; Roberts 1968: 192, 1969b: 115; Schedl 1972f: 151. (tx) Roberts 1972: 72; Schedl 1962k: 1119, 1972f: 151, 1978a: 81.

- wittei Schedl** 1952g: 55. Lectotype ♂; Parc Nat. de l'Upemba, Kaswabilenga, 700 m; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 81.
 Distribution: Africa (Zaire).
 References: (ds) Schedl 1972f: 156. (tx) Schedl 1952g: 55, 1962k: 824, 1972f: 156, 1978a: 81.
- woltschei Schedl** 1957d: 138. Holotype ♂; Congo Belge: Yangambi; MRCB, Tervuren.
 Distribution: Africa (Zaire).
 Hosts: *Dialium corbisieri*, *Staudtia stiptata*, *Turraecanthus africana*.
 References: (hb) Schedl 1962k: 945. (ds) Schedl 1962k: 945, 1972f: 151. (tx) Numberg 1966a: 187; Schedl 1957d: 138, 1962k: 945, 1972f: 151, 1978a: 81.
- zonatus Schedl** 1962k: 1120. Holotype ♂; Congo: Kivu; Schedl Collection in NHMW, Wien.
 Distribution: Africa (Ghana/ Zaire).
 References: (ds) Schedl 1972f: 158. (tx) Schedl 1962k: 1120, 1972f: 158, 1978a: 82.

Genus *Dendroplatypus* Browne

- DENDROPLATYPUS BROWNE** 1955: 363. Type-species: *Crossotarsus impar* Schedl, original designation.
 References: (cn) Browne 1968b: 228. (hb) Browne 1961c: 213, 1968b: 228; Schedl 1972f: 138. (ds) Browne 1968b: 228; Schedl 1972f: 138. (tx) Browne 1955: 365, 1962e: 650, 655, 1972a: 180; Schedl 1972f: 138.
- impar (Schedl)** 1936d: 13 (*Crossotarsus*). Lectotype ♂; Malay Peninsula, Johore North: Klang, Lower Perak, Bt. Tunggal For. Res.; Schedl Collection in NHMW, Wien, designated by Schedl 1978a: 36.
 Figures: Schedl 1972f: 138 (male).
 Distribution: Asia (Malaya), Indonesia (Sarawak in Borneo, Sumatra).
 Hosts: *Shorea leprosula*, *S. parvifolia*.
 Notes: (1) Browne 1955: 365 (to *Dendroplatypus*).
 References: (bv) Bletchly 1961: 15; Neumann & Harris 1974: 136. (cn) Beeson 1961 (1941: 254); Browne 1938b, 1952; Browne & Foenander 1937. (hb) Browne 1936b: 13, 1961c: 213–216, 1965b. (ds) Beeson 1941 (1961: 254); Browne 1961c: 213, 1980a: 371; Browne & Foenander 1937; Schedl 1972f: 140. (tx) Browne 1936a: 120–127, 1936b: 3, 1950b: 649, 1955: 364–365, 372, 1972a: 180; Schedl 1936d: 13, 1939b: 391, 1972a: 180, 1972f: 138, 140, 1978a: 36.
- philippinensis (Blandford)** 1896b: 193 (*Platypus*). Holotype ♂; Philippine Is.; BMNH, London.
 Figures: Schedl 1972f: 193 (male).
 Distribution: Philippine Islands (Luzon, Mindanao).
 Notes: (3) Schedl 1935b: 395 (described female).
 References: (ds) Schedl 1966b: 97, 1972f: 204; Strohmeier 1912c: 19, 1914c: 29. (tx) Blandford 1896b: 193; Browne 1962e: 650; Nobuchi 1985: 329; Schedl 1935b: 395, 1939b: 398, 1952b: 366, 1972f: 204, 1978a: 55; Strohmeier 1911b: 17, 1912c: 19, 1914c: 29.

Hosts of Scolytidae and Platypodidae

The following host list includes only those host associations for Scolytidae that were in the computer in January 1992. Almost all associations for Platypodidae are included to August 1992. The user should also consult the hosts listed under beetle species in order to include the maximum number of host associations.

"*L. glauca*"

Hypothenemus parallelus (Hopkins).

"*Mudirira*"

Phrixosoma nigra (Eggers).

"*Nudarira*"

Hylesinopsis niger (Schedl).

Abies alba

Hylurgops glabratus (Zetterstedt), *Hylurgops palliatus* (Gyllenhal), *Pityophthorus cephalonicae* Pfeffer, *Pityophthorus pityographus* (Ratzeburg).

Abies amabilis

Cryphalus ruficollis Hopkins, *Dryocoetes confusus* Swaine, *Pityokteines lasiocarpi* (Swaine), *Pityokteines mystacinus* Wood, *Pseudohylesinus dispar dispar* Blackman, *Pseudohylesinus dispar pullatus* Blackman, *Pseudohylesinus granulatus* (LeConte), *Pseudohylesinus nobilis* Swaine, *Pseudohylesinus sericeus* (Mannerheim).

Abies balsamea

Cryphalus ruficollis Hopkins, *Crypturgus borealis* Swaine, *Pityokteines sparsus* (LeConte), *Pityophthorus angustus* Blackman, *Pityophthorus balsameus* Blackman, *Pityophthorus cariniceps* LeConte, *Pityophthorus opaculus* LeConte, *Pityophthorus puberulus* (LeConte), *Pityophthorus pulicarius* (Zimmermann).

Abies borisi-regis

Pityophthorus pityographus (Ratzeburg).

Abies bornmulleriana

Cryphalus asperatus (Gyllenhal), *Pityophthorus pityographus* (Ratzeburg).

Abies bracteata

Scolytus dentatus Bright.

Abies cephalonica

Cryphalus piceae (Ratzeburg), *Pityophthorus cephalonicae* Pfeffer, *Pityophthorus pityographus* (Ratzeburg), *Pityophthorus pubescens* (Marshall), *Platypus oxyurus* Dufour.

Abies concolor

Crypturgus borealis Swaine, *Dryocoetes confusus* Swaine, *Gnathotrichus denticulatus* Blackman, *Gnathotrichus retusus* (LeConte), *Gnathotrichus sulcatus* (LeConte), *Hylastes gracilis* LeConte, *Pityokteines elegans* Swaine, *Pityophthorus apachae* Bright, *Pityophthorus opaculus* LeConte, *Pityophthorus pseudotsugae* Swaine, *Pityophthorus solers* Blackman, *Platypus abietis* Wood, *Pseudohylesinus dispar dispar* Blackman, *Pseudohylesinus maculosus* Blackman, *Scolytus obelus* Wood, *Scolytus opacus* Blackman, *Scolytus praeceps* LeConte, *Scolytus robustus* Blackman, *Scolytus subscaber* LeConte, *Scolytus ventralis* LeConte.

Abies delavayi

Hylurgops euscultatus Tsai & Huang, *Phloeosinus abietis* Tsai & Yin.

Abies densa

Cryphalus strommeyeri Stebbing.

Abies fabri

Ambrosiodmus rubricollis (Eichhoff), *Cryphalus redikorzei* Berger, *Polygraphus major* Stebbing, *Poly-*

graphus zhungdianensis Tsai & Yin, *Xyleborus emarginatus* Eichhoff, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus germanus* (Blandford).

Abies faxoniana

Cryphalus markangensis Tsai & Li, *Cryphalus pilosus* Tsai & Li, *Cryphalus sinoabietis* Tsai & Li.

Abies firma

Coccotrypes norimasanus (Murayama), *Cryphalus jeholensis* Murayama, *Cryphalus piceae* (Ratzeburg), *Euwallacea validus* (Eichhoff), *Platypus calamus* Blandford, *Platypus lewisi* Blandford, *Polygraphus oblongus* Blandford.

Abies fraseri

Cryphalus ruficollis Hopkins, *Pityophthorus dentifrons* Blackman, *Polygraphus rufipennis* (Kirby).

Abies grandis

Cryphalus pubescens Hopkins, *Cryphalus ruficollis* Hopkins, *Crypturgus borealis* Swaine, *Pityophthorus pseudotsugae* Swaine, *Pseudohylesinus dispar dispar* Blackman, *Pseudohylesinus dispar pullatus* Blackman, *Pseudohylesinus granulatus* (LeConte), *Pseudohylesinus sericeus* (Mannerheim), *Scolytus opacus* Blackman, *Scolytus subscaber* LeConte, *Scolytus ventralis* LeConte.

Abies guatemalensis

Pityophthorus mesembria Bright.

Abies holophylla

Cryphalus kurenzovi Stark, *Cryphalus laricis* Niisima, *Cryphalus piceae* (Ratzeburg), *Cryphalus redikorzei* Berger, *Dryocoetes striatus* Eggers, *Hylurgops interstitialis* (Chapuis), *Pityophthorus abietinus* Wood, *Polygraphus abietis* Kurenzov.

Abies koreana

Ambrosiodmus rubricollis (Eichhoff), *Xyleborinus saxeseni* (Ratzeburg).

Abies lasiocarpa

Cryphalus ruficollis Hopkins, *Crypturgus borealis* Swaine, *Dryocoetes confusus* Swaine, *Dryocoetes sechelti* Swaine, *Hylastes subopacus* Blackman, *Phloeotribus lecontei* Schedl, *Pityokteines lasiocarpi* (Swaine), *Pityokteines minutus* (Swaine), *Pityophthorus absomus* Blackman, *Pityophthorus opaculus* LeConte, *Pityophthorus pseudotsugae* Swaine, *Pityophthorus solers* Blackman, *Pseudohylesinus granulatus* (LeConte), *Pseudohylesinus maculosus* Blackman, *Sciurus pubescens* Swaine, *Scolytus opacus* Blackman.

Abies magnifica

Cryphalus ruficollis Hopkins, *Gnathotrichus retusus* (LeConte), *Gnathotrichus sulcatus* (LeConte), *Pityokteines elegans* Swaine, *Pityophthorus pseudotsugae* Swaine, *Pityophthorus solers* Blackman, *Platypus wilsoni* Swaine, *Pseudohylesinus dispar dispar* Blackman, *Pseudohylesinus granulatus* (LeConte), *Scolytus subscaber* LeConte, *Scolytus ventralis* LeConte.

Abies mariesii

Cryphalus montanus Nobuchi, *Cryphalus piceae* (Ratzeburg), *Polygraphus oblongus* Blandford.

Abies nephrolepis

Cryphalus kurenzovi Stark, *Dryocoetes striatus* Eggers,

Hylastes plumbens Blandford, *Pityogenes scirindensis* Murayama, *Polygraphus jeczocensis* Niisima, *Polygraphus miser* Blandford, *Polygraphus proximus* Blandford.

Abies nobilis

Dolurgus pumilus (Mannerheim).

Abies nordmanniana

Hylurgops palliatus (Gyllenhal).

Abies numidica

Cryphalus numidicus Eichhoff, *Crypturgus numidicus* Ferrari, *Platypus oxyurus* Dufour.

Abies pectinata

Cryphalus asperatus (Gyllenhal), *Cryphalus piccae* (Ratzeburg), *Hylurgops palliatus* (Gyllenhal), *Ips anitinus* (Eichhoff), *Philocotribus spinulosus* (Rey), *Pityogenes irkutensis* Eggers, *Pityokteines curvidens* (Gernar), *Pityokteines spinidens* (Reitter), *Pityokteines vorontzovi* (Jacobson), *Pityophthorus pityographus* (Ratzeburg), *Platypus oxyurus* Dufour.

Abies pindroie

Cryphalus strommeyeri Stebbing, *Dryocoetes quadrisulcatus* Strohmeier, *Pityophthorus deodara* (Stebbing), *Scolytoplatypus siomio* Blandford.

Abies pinsapo

Crypturgus numidicus Ferrari, *Pityophthorus pinsapo* Pfeffer.

Abies procera

Pseudohylesinus dispar pullatus Blackman, *Pseudohylesinus granulatus* (LeConte), *Pseudohylesinus nobilis* Swaine, *Pseudohylesinus sericeus* (Mannerheim), *Pseudohylesinus tsugae* Swaine.

Abies religiosa

Gnathotrichus sulcatus (LeConte), *Pityophthorus abiegus* Wood, *Pityophthorus anthracinus* Bright, *Pityophthorus brighti* Wood, *Pityophthorus rudis* Blackman, *Pityophthorus solers* Blackman, *Pseudohylesinus magnus* Wood, *Pseudohylesinus variegatus* (Blandford), *Scolytus aztecus* Wood, *Scolytus hernosus* Wood, *Scolytus nandus* Wood.

Abies sachalinensis

Cryphalus laticis Niisima, *Cryphalus piccae* (Ratzeburg), *Cryphalus yanaguchii* Inouye & Nobuchi, *Dryocoetes striatus* Eggers, *Polygraphus gracilis* Niisima, *Polygraphus proximus* Blandford.

Abies sibirica

Carphoborus jurinskii Eggers, *Carphoborus rossicus* Semenov, *Carphoborus teplouchovi* Spessivtsev, *Polygraphus subopacus* Thomson.

***Abies* spp.**

Cryphalus latus Eggers, *Cryphalus lepocrinus* Tsai & Li, *Cryphalus markangensis* Tsai & Li, *Cryphalus pilosus* Tsai & Li, *Cryphalus sinobietis* Tsai & Li, *Crypturgus cinereus* (Herbst), *Crypturgus hispidulus* Thomson, *Crypturgus pusillus* (Gyllenhal), *Dendroctonus micans* (Kugelann), *Dendroctonus valens* LeConte, *Dryocoetes affaber* (Mannerheim), *Dryocoetes autographus* (Ratzeburg), *Dryocoetes hectographus* Reitter, *Dryocoetes pini* Niisima, *Dryocoetes rugicollis* Eggers, *Gnathotrichus materiarius* (Fitch), *Hylastes ater* (Paykull), *Hylastes nigrinus* (Mannerheim), *Hylurgops glabratus* (Zetterstedt), *Ips acuminatus* (Gyllenhal), *Ips cembrae* (Heer), *Pityogenes bidentatus* (Herbst), *Pityogenes chalcographus* (Linnaeus), *Pityogenes quadridens* (Hartig), *Pityophthorus jucundus* Blandford, *Pityophthorus micrographus* (Linnaeus), *Pityophthorus speculum* Bright, *Polygraphus horjurenensis* Murayama, *Polygraphus japonicus* Nunberg, *Polygraphus polygraphus* (Linnaeus), *Polygraphus proximus* Blandford,

Scolytoplatypus dainio Blandford, *Scolytoplatypus shogun* Blandford, *Scolytoplatypus tycon* Blandford, *Trypodendron lineatum* (Olivier), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus aquilus* Blandford, *Xylechinus pilosus* (Ratzeburg).

Abies veitchii

Cryphalus sawadai Nobuchi & Takahashi, *Polygraphus fulvipennis* Niisima.

Abies webbiana

Cryphalus strommeyeri Stebbing, *Crypturgus beesonii* Eggers, *Dryocoetes indicus* Stebbing, *Dryocoetes quadrisulcatus* Strohmeier, *Hylastes brunneus* Erichson, *Ips stebbingi* Strohmeier, *Pityophthorus deodara* (Stebbing), *Pityophthorus clatinus* Wood, *Polygraphus aterrinus* Strohmeier, *Polygraphus pini* Stebbing, *Scolytoplatypus raja* Blandford.

Abrus precatorius

Hypothenemus eruditus Westwood.

Abutilon mollissimum

Hypothenemus crudiae (Panzer), *Hypothenemus eruditus* Westwood.

***Abutilon* spp.**

Xylosandrus compactus (Eichhoff).

Acacia auriculaeformis

Hypothenemus dimorphus (Schedl).

Acacia catechu

Euwallacea andamanensis (Blandford), *Hypothenemus acaciae* (Eggers), *Platypus solidus* Walker.

Acacia cavenia

Chramesus spinosus Brethes.

Acacia decurrens

Crossotarsus horni Schedl, *Crossotarsus squamulatus* Chapuis, *Doliopygus excavatus* (Sampson), *Glostatus acaciae* (Schedl), *Hypothenemus eruditus* Westwood, *Platypus carinifrons* Schedl, *Platypus solidus* Walker, *Platypus solutus* Schedl, *Scolytoplatypus nitidus* Eggers, *Scolytoplatypus raja* Blandford, *Xyleborus alluandi* Schanfuss, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky), *Zygophloeus australis* Schedl.

Acacia dudgeoni

Doliopygus terebraus Schedl, *Xyleborinus similans* (Eggers).

Acacia farnesiana

Hypothenemus brunneus (Hopkins), *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff).

Acacia karroo

Xyleborinus acmulus (Wollaston).

Acacia koa

Crossotarsus externudentatus (Fairmaire), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus spinulosus* Blandford.

Acacia lahai

Doliopygus ghesquieri (Schedl), *Doliopygus lefevrei* Schedl, *Doliopygus opifex* (Sampson).

Acacia melanoxylon

Xyleborus affinis Eichhoff.

Acacia mollissima

Chaetastus montanus Schedl, *Scolytoplatypus eichelbaumi* Hagedorn, *Xyleborus atratus* Eichhoff, *Xyleborus clerodendronae* Schedl.

Acacia nigra

Araptus sparsepunctatus (Schedl), *Cnestinus dividuus* Schedl, *Cryptocarenum acaciae* Schedl.

Acacia pennata

Coccotrypes longior (Eggers), *Coccotrypes nubilus* (Blandford), *Eucallacea xanthopus* (Eichhoff), *Rhopalopselion acaciae* (Schedl), *Xyleborus ficus* Eggers, *Xyleborus picus* (Motschulsky).

Acacia pennatula

Cnestinus adusticus Wood, *Cnestinus setulosus* Blandford, *Hylocurus atkinsoni* Wood, *Hypothenemus setosus* (Eichhoff), *Microcorthyus minutus* Schedl, *Pseudothysanoes contrarius* Wood.

Acacia podalyriaefolia

Xyleborinus mimosae (Schedl).

Acacia polyphylla

Xyleborus spinulosus Blandford.

Acacia raddiana

Hypothenemus mateni (Schedl).

Acacia sieberiana

Hypothenemus biseriatus (Eggers), *Rhopalopselion maculatus* (Schedl).

Acacia spp.

Acacis abundans Lea, *Ambrosiodmus aegir* (Eggers), *Ambrosiodmus albizzianus* (Schedl), *Ambrosiodmus camphorae* (Hagedorn), *Ambrosiodmus eichhoffi* (Schreiner), *Ambrosiodmus hagedorni* (Iglesias), *Ambrosiodmus obliquus* (LeConte), *Ambrosiodmus rubricollis* (Eichhoff), *Ambrosiodmus sakoae* (Schedl), *Chramesus acacicolens* Wood, *Chramesus disparilis* Wood, *Chramesus secus* Wood, *Chramesus strigatus* Wood, *Chramesus varius* Wood, *Cladoctonus eggersi* Wichmann, *Cladoctonus natalensis* Eggers, *Cnestus suturalis* (Eggers), *Corthylocurus mexicanus* (Schedl), *Corthylus collaris* Blandford, *Corthylus consimilis* Wood, *Corthylus flagellifer* Blandford, *Cryphalus sylvicola* (Perkins), *Doliopygus interruptus* (Sampson), *Doliopygus lefevrei* Schedl, *Doliopygus subditicus* (Schedl), *Eucallacea fornicatus* (Eichhoff), *Glostus acaciae* (Schedl), *Hadrodemus amorphus* (Eggers), *Hylesinopsis acacicolens* Wood, *Hylesinopsis striatus* (Schedl), *Hylocurus elegans* Eichhoff, *Hylocurus inaequalis* Wood, *Hylocurus scitulus* Wood, *Hypothenemus agnatus* (Eggers), *Hypothenemus camerinus* (Eggers), *Hypothenemus eruditus* Westwood, *Hypothenemus plumieriae* (Nordlinger), *Phloeotribus acaciae* (Lea), *Pityophthorus colvulus* Schedl, *Platypus furcatus* Blandford, *Platypus omnivorus* (Lea), *Platypus solidus* Walker, *Prennobius cavipennis* Eichhoff, *Pseudothysanoes acacicolens* Wood, *Pseudothysanoes aquilus* (Wood), *Scolytotplatypus kitensis* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Thysanoes fimbriicornis* LeConte, *Thysanoes mexicanus* Wood, *Thysanoes texanus* Blackman, *Xyleborinus collarti* (Eggers), *Xyleborinus spinifer* (Eggers), *Xyloctonus striatus* Eggers, *Xylosandrus compactus* (Eichhoff).

Acacia xanthophloea

Cladoctonus amanicus Eggers, *Hypothenemus xanthophloae* (Schedl), *Lamargus xanthophloae* Schedl, *Phloeocurus africanus* (Schedl).

Acalypha neptunica

Xyleborus scobinatus Hagedorn.

Acalypha spp.

Ambrosiodmus albizzianus (Schedl), *Hypothenemus adscitus* (Schedl), *Micracis exilis* Wood, *Xyleborus alluaudi* Schanfuß, *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Acanthocephalus cadamba

Ficicis javanus (Eggers), *Platypus labrifrons* Schedl.

Acanthopanax ricinifolium

Ernoporus acanthopanxi (Niisima).

Acanthopanax spinosum

Phloeosinus pulchellus Blandford.

Acanthopanax spp.

Scolytotplatypus mikado Blandford.

Acanthus pubescens

Xyleborus multispinatus Eggers.

Acer argentum

Taphrocyetus mikuniyamaensis (Murayama).

Acer babinerre

Indocryphalus aceris (Niisima).

Acer caesium

Scolytotplatypus darjeelingi Stebbing, *Scolytotplatypus siomio* Blandford.

Acer campbellii

Eucallacea velatus (Sampson), *Platypus darjeelingensis* Schedl, *Platypus falcatus* Strohmeyer, *Platypus shillonensis* Schedl, *Scolytotplatypus pubescens* Hagedorn.

Acer campestre

Lymanator aceris (Lindemann), *Lymanator coryli* (Perris).

Acer macrophyllum

Procryphalus utahensis Hopkins.

Acer mandshuricum

Indocryphalus aceris (Niisima).

Acer mayrii

Dryocoetes picipennis Eggers.

Acer mono

Crossotarsus niponicus Blandford, *Indocryphalus aceris* (Niisima), *Neopteleobius scutellatus* (Blandford), *Platypus severini* Blandford.

Acer palmatum

Crossotarsus niponicus Blandford, *Platypus modestus* Blandford, *Xyleborus atratus* Eichhoff, *Xyleborus prae-vius* Blandford.

Acer palmatum matsumurae

Crossotarsus niponicus Blandford.

Acer pictum

Dryocoetes aceris Krivolitskaya, *Hyorrhynchus lewisi* Blandford, *Indocryphalus aceris* (Niisima).

Acer platanoides

Lymanator aceris (Lindemann), *Lymanator coryli* (Perris).

Acer pseudoplatanus

Lymanator aceris (Lindemann).

Acer pseudosieboldianum

Indocryphalus aceris (Niisima).

Acer rubrum

Corthylus columbianus Hopkins, *Hypothenemus dissimilis* (Zimmermann), *Hypothenemus interstitialis* (Hopkins), *Thysanoes fimbriicornis* LeConte, *Lymanator decipiens* (LeConte), *Monarthrum fasciatum* (Say), *Monarthrum mali* (Fitch), *Trischidias atomus* (Hopkins), *Xyleborus ferrugineus* (Fabricius), *Xylosandrus zimmermanni* (Hopkins).

Acer rufinerre

Xyleborus aquilus Blandford, *Xyleborus seriatus* Blandford.

Acer saccharinum

Corthylus columbianus Hopkins, *Corthylus punctatissimus* (Zimmermann), *Lymanator decipiens* (LeConte), *Pityophthorus lautus* Eichhoff.

Acer spp.

Amasa amputatus (Blandford), *Ambrosiodmus lewisi* (Blandford), *Ambrosiodmus tachygraphus* (Zimmer-

- mann). *Dryocoetes padi* Stark. *Dryocoetes ussuriensis* Eggers. *Eucallacea calidus* (Eichhoff). *Hylocurus torosus* Wood. *Lymantor decipiens* (LeConte). *Platypus korjocensis* (Murayama). *Scolytoplatypus dainio* Blandford. *Scolytoplatypus mikado* Blandford. *Scolytoplatypus shogun* Blandford. *Scolytoplatypus tycon* Blandford. *Scolytus belokanicus* Stark. *Scolytus koenigi* Schelyrew. *Scolytus laevis* Chapuis. *Taphrotychus lenkoranus* Reitter. *Trypodendron domesticum* (Linnaeus). *Trypodendron signatum* (Fabricius). *Xyleborinus saxeseni* (Ratzeburg). *Xyleborus maiche* (Stark). *Xyleborus obesus* LeConte. *Xyleborus orientalis* Eggers. *Xyleborus sayi* (Hopkins). *Xyleborus pelliculosus* Eichhoff. *Xyleborus pfeili* (Ratzeburg). *Xylosandrus compactus* (Eichhoff). *Xylosandrus germanus* (Blandford). *Xylosandrus mutilatus* (Blandford). *Xyloterinus politus* (Say).
- Acer tataricum**
Lymantor aceris (Lindemann).
- Acer tegmentosum**
Indocryphalus aceris (Niisima).
- Acer turkestanicum**
Scolytus tadzhikistanicus Stark.
- Acer ukurduense**
Indocryphalus aceris (Niisima).
- Achras sapota**
Chaetophloeus insularis (Blackman). *Hypothenemus birmanus* (Eichhoff). *Hypothenemus crudiae* (Panzer). *Hypothenemus hirsutus* (Wood). *Hypothenemus seriatus* (Eichhoff). *Micracisella nanula* (LeConte). *Scolytominus pusillus* (Eggers). *Xyleborus fallax* Eichhoff. *Xyleborus perforans* (Wollaston).
- Acmena spp.**
Diapus bilunatus Schedl. *Diapus quinquespinnatus* Chapuis. *Diapus robustus* Schedl.
- Aconitum stoerkianum**
Thamurgus petzi Reitter.
- Acridocarpus fraxinifolius**
Doliopygus excavatus (Sampson).
- Acridocarpus spp.**
Ambrosiodinus albizzianus (Schedl). *Ambrosiodinus eichhoffi* (Schreiner). *Platypus madagascariensis* Chapuis.
- Acrocarpus fraxinifolius**
Ambrosiodinus eichhoffi (Schreiner). *Doliopygus erichsoni* (Chapuis). *Leptoxyloborus sordicauda* (Motschulsky). *Platypus errans* (Sampson). *Platypus solidus* Walker. *Platypus solutus* Schedl. *Xyleborinus forcipatus* (Schedl). *Xyleborinus sharpi* (Hopkins). *Xyleborus affinis* Eichhoff. *Xyleborus sphenos* Sampson.
- Acrocarpus spp.**
Cyclorhupidion crucifer (Hagedorn). *Doliopygus coclocephalus* (Schaufuss). *Leptoxyloborus concisus* (Blandford). *Prennobius cavipennis* Eichhoff.
- Acrocomia sclerocarpa**
Hypothenemus seriatus (Eichhoff).
- Actinodaphne acuminata**
Xyleborus colvulus (Fabricius).
- Actinodaphne lancifolia**
Platypus calanus Blandford.
- Actinodaphne longifolia**
Phloeosinus kiushuensis Murayama. *Phloeosinus lewisi* Chapuis.
- Actinophora fragrans**
Coptodryas nugax (Schedl). *Platypus biuncus* Blandford. *Platypus forficula* Chapuis. *Xyleborinus andrewesi* (Blandford). *Xylosandrus morigerus* (Blandford).
- Adenanthera microsperma**
Hypothenemus eruditus Westwood.
- Adenanthera pavonina**
Arixyleborus suturalis (Eggers). *Eccoptopterus limbus* Sampson. *Hypothenemus birmanus* (Eichhoff). *Hypothenemus crudiae* (Panzer). *Hypothenemus eruditus* Westwood. *Xyleborus perforans* (Wollaston). *Xylosandrus ater* (Eggers). *Xylosandrus compactus* (Eichhoff). *Xylosandrus maucus* (Blandford). *Xylosandrus morigerus* (Blandford).
- Adenia cisampeloides**
Xylocleptes adeniae (Schedl).
- Adina cordifolia**
Crossotarsus externedentatus (Fairmaire). *Xyleborus perforans* (Wollaston). *Xyleborus similis* Ferrari.
- Adina rubescens**
Xyleborus javanus Eggers. *Xylosandrus crassiusculus* (Motschulsky).
- Adina rubrostipulata**
Platypus impressus (Strohmeyer). *Xyleborus alluaudi* Schauffuss. *Xyleborus principalis* Eichhoff.
- Adina spp.**
Platypus solidus Walker. *Platypus suffodiens* Sampson.
- Adinandra dumosa**
Eucallacea xanthopus (Eichhoff). *Xylosandrus crassiusculus* (Motschulsky).
- Adinobotrys atropurpureus**
Platypus lepidus Chapuis. *Xylosandrus crassiusculus* (Motschulsky).
- Aegle marmelos**
Xyleborinus andrewesi (Blandford). *Xyleborus adusticollis* (Motschulsky). *Xyleborus similis* Ferrari.
- Aerca sanguinolenta**
Xyleborus picus (Motschulsky).
- Aesculus punduana**
Coccotrypes cyperi (Beeson). *Coccotrypes vulgaris* (Eggers). *Platypus secretus* Sampson.
- Aesculus spp.**
Hypothenemus eruditus Westwood. *Hypothenemus interstitialis* (Hopkins). *Scolytoplatypus tycon* Blandford. *Xyleborus dispar* (Fabricius).
- Aesculus turbiuata**
Crossotarsus niponicus Blandford. *Indocryphalus dainichiensis* (Murayama). *Platypus calanus* Blandford. *Platypus lewisi* Blandford. *Platypus modestus* Blandford. *Platypus severini* Blandford. *Xyleborus seriatus* Blandford.
- Afromosia elata**
Chaetastus tuberculatus (Chapuis). *Cyrtogenius chlorophorae* (Schedl). *Doliopygus artemisatus* (Schedl). *Doliopygus conradti* (Schauffuss). *Doliopygus divaricus* (Schedl). *Doliopygus dubius* (Sampson). *Doliopygus lebruni* Schedl. *Doliopygus medius* Schedl. *Doliopygus serratus* (Strohmeyer). *Doliopygus spectabilis* Schedl. *Doliopygus subditicus* (Schedl). *Doliopygus tenuis* (Strohmeyer). *Doliopygus unicornis* Schedl. *Hypothenemus eruditus* Westwood. *Platypus hintzi* Schauffuss. *Platypus solutus* Schedl. *Platypus spinulosus* (Strohmeyer). *Periommatius excisus* Strohmeyer. *Periommatius grandis* Schedl. *Trachyostus aterrimus* (Schauffuss). *Trachyostus schaufussi* (Strohmeyer). *Trachyostus schionedeni* (Schedl). *Triozastus marshalli* (Sampson). *Xyleborus alluaudi* Schauffuss. *Xyleborus ferrugineus* (Fabricius).
- Afrostyax lepidophyllus**
Doliopygus erichsoni (Chapuis). *Doliopygus interjectus*

- Schedl, *Doliopygus rapax* (Sampson), *Eucallacea granosus* (Schedl), *Periommatius subrobustus* Schedl.
- Afzelia africana**
Doliopygus brevis (Strohmeyer), *Doliopygus opifex* (Sampson), *Doliopygus perbrevis* Schedl, *Pityophthorus congonus* Eggers, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer).
- Afzelia bakeri**
Platypus westwoodi Chapuis, *Xyleborus bidentatus* (Motschulsky), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.
- Afzelia bella**
Doliopygus conradti (Strohmeyer), *Doliopygus erichsonii* (Chapuis), *Doliopygus subditicus* (Schedl), *Hypothenemus marshalli* (Eggers), *Hypothenemus morigerus* (Schedl), *Platypus hintzi* Schaufuss, *Polygraphus afzeliae* Schedl, *Polygraphus carphoboroides* Eggers, *Xyleborus alluaudi* Schaufuss, *Xyleborus ustus* Schedl.
- Afzelia bijuga**
Cryphalus scabricollis Eichhoff, *Diapys pusillimus* Chapuis, *Erioporus dispar* (Schedl), *Platypus jansoni* Chapuis, *Xyleborus bidentatus* (Motschulsky).
- Afzelia bipindensis**
Doliopygus chapuisi (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus rapax* (Sampson), *Doliopygus tenuis* (Strohmeyer), *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).
- Afzelia cuanensis**
Triozastus marshalli (Sampson).
- Afzelia palembanica**
Crossotarsus externidentatus (Fairmaire), *Crossotarsus fractus* Sampson, *Crossotarsus incernis* Schedl, *Crossotarsus schedli* Brownie, *Crossotarsus squamulatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Diapys pusillimus* Chapuis, *Diapys quinquispinatus* Chapuis, *Eucallacea destruens* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Hadrodemus globus* (Blandford), *Leptoxyleborus concisus* (Blandford), *Leptoxyleborus puer* (Eggers), *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Platypus linaifer* Schedl, *Platypus pseudocurtus* Schedl, *Platypus solidus* Walker, *Platypus westwoodi* Schedl, *Xyleborus birmanus* Eggers, *Xyleborus emarginatus* Eichhoff, *Xyleborus fallax* Eichhoff, *Xyleborus latus* Eggers, *Xyleborus perforans* (Wollaston), *Xylosandrus morigerus* (Blandford).
- Afzelia spp.**
Cylindropalpus camerunus (Schedl), *Platypus schysi* Chapuis.
- Afzelia xylocarpa**
Platypus afzeliae Brownie.
- Agatha ritiensis**
Cyrtogenius fijianus (Schedl).
- Agathis alba**
Cyrtogenius perakensis (Schedl), *Diapys papuanus* Schedl, *Diapys pusillimus* Chapuis, *Diapys quinque-spinatus* Chapuis, *Hypothenemus eruditus* Westwood, *Platypus cupulatus* Chapuis, *Platypus grumosus* Roberts & Morimoto, *Xyleborus multipunctatus* Brownie.
- Agathis borneensis**
Xyleborus apertus Schedl, *Xyleborus eximius* Schedl.
- Agathis dammara**
Cyrtogenius perakensis (Schedl).
- Agathis palmerstonii**
Xyleborus perforans (Wollaston).
- Agathis spp.**
Platypus agathis Brownie, *Xyleborus agathis* Brownie, *Xyleborus bidentatus* (Motschulsky).
- Agathis ritiensis**
Platypus ritiensis Roberts.
- Agave sisalana**
Platypus cupulatus Chapuis, *Platypus pseudocupulatus* Schedl.
- Agave spp.**
Chramesus annectens (Wood), *Platypus cupulatus* Chapuis, *Platypus pseudocupulatus* Schedl, *Xyleborus perforans* (Wollaston).
- Aglaia spp.**
Crossotarsus auripilus Roberts, *Crossotarsus corrugatus* Roberts, *Crossotarsus incernis* Schedl, *Crossotarsus subopacus* Schedl, *Crossotarsus terminatus* Chapuis, *Platypus anguis* Brownie, *Platypus subpronus* Schedl, *Xyleborus aglaiae* Brownie.
- Ailanthus altissima**
Ambrosiodinus levisi (Blandford), *Hypothenemus eruditus* Westwood, *Xylosandrus discolor* (Blandford).
- Alatonia spatulata**
Acanthotomicus onerosus (Schedl).
- Albizzia adianthifolia**
Ambrosiodinus tropicus (Hagedorn), *Doliopygus chapuisi* (Duvivier), *Eccoptopterus spinosus* (Olivier), *Premnobius cavipennis* Eichhoff, *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus similis* (Eggers), *Xyleborus subtuberculatus* Eggers.
- Albizzia caribaea**
Hypothenemus suspectus Wood.
- Albizzia chinensis**
Eucallacea xanthopus (Eichhoff), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).
- Albizzia coriaria**
Acanthotomicus sexdentatus (Eggers), *Doliopygus chapuisi* (Duvivier), *Platypus parallelus* (Fabricius).
- Albizzia falcata**
Ambrosiodinus wilderi (Beeson), *Crossotarsus wallacei* (Thomson), *Diapys pusillimus* Chapuis, *Eucallacea fornicatus* (Eichhoff), *Platypus biuncus* Blandford, *Platypus cupulatus* Chapuis, *Platypus insularis* Strohmeyer, *Platypus insulindicus* Schedl, *Platypus subangustior* Schedl, *Platypus subsilinus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus perforans* (Wollaston), *Xylosandrus morigerus* (Blandford).
- Albizzia ferruginea**
Doliopygus chapuisi (Duvivier), *Doliopygus exilis* (Chapuis), *Trachyostus aterrimus* (Schaufuss), *Triozastus banghaasi* (Schaufuss), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).
- Albizzia glaberrima**
Chaetastus tuberculatus (Chapuis), *Cryphalus nigericus* Brownie, *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus excavatus* (Sampson), *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus alluaudi* Schaufuss, *Xyleborus picus* (Motschulsky).
- Albizzia grandibracteata**
Doliopygus chapuisi (Duvivier), *Hypothenemus eruditus* Westwood, *Rhopalopselion congonum* (Schedl).

Albizzia giunmifera

Acanthotomicus acuminatus (Schedl), *Acanthotomicus biconicus* (Schedl), *Ambrosiodmus albizzianus* (Schedl), *Ambrosiodmus bostrichoides* (Schedl), *Chaetastus montanus* Schedl, *Chaetastus tuberculatus* (Chapuis), *Coptoborus adjunctus* (Eggers), *Cylindropalpus pumilio* (Schedl), *Doliopygus artespianus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Dryococtoides cristatus* (Fabricius), *Eccoptopterus spinosus* (Olivier), *Hylesinopsis seriatus* (Eggers), *Hypothenemus eruditus* Westwood, *Hypothenemus solitarius* (Schedl), *Perionnatus excisus* Strohmeyer, *Perionnatus grandis* Schedl, *Perionnatus longicollis* Chapuis, *Pityophthorus congonus* Eggers, *Pityophthorus kiruensis* Schedl, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Platypus schenklingi* (Strohmeyer), *Polygraphus coronatus* Eggers, *Premnobius corthyloides* Hagedorn, *Premnobius quadrispinosus* Schedl, *Rhopalopsclion suturalis* (Schedl), *Theoborus ricini* (Eggers), *Tiarophorus elongatus* Schreiner, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus ambasicus* Eggers, *Xyleborus amnecetus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus multispinatus* Eggers, *Xyleborus piceus* (Motschulsky), *Xyleborus sub-tuberculatus* Eggers, *Xyleborus volcuhus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus cylindrotomicus* (Schedl).

Albizzia julibrissin

Xyleborus atratus Eichhoff, *Xyleborus volcuhus* (Fabricius).

Albizzia lebbek

Ambrosiodmus lantanae (Eggers), *Ambrosiodmus sub-nepotulus* (Eggers), *Arixyleborus malayensis* (Schedl), *Chaetastus tuberculatus* (Chapuis), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus squamulatus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Hypothenemus eruditus* Westwood, *Xyleborinus andrevesi* (Blandford), *Xyleborus affinis* Eichhoff, *Xyleborus perforans* (Wollaston), *Xyleborus volcuhus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).

Albizzia lebbekioides

Hypothenemus brunneus (Hopkins), *Hypothenemus eruditus* Westwood.

Albizzia lucida

Platypus curvus Strohmeyer, *Xyleborus perforans* (Wollaston).

Albizzia macrocarpa

Pseudothysanocera murilloi (Blackman).

Albizzia maranguensis

Doliopygus nitens Browne.

Albizzia moluccana

Coptodryas elegans (Sampson), *Eucallacea bicolor* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Platypus insulindicus* Schedl, *Platypus subangustior* Schedl, *Platypus subsimilis* Schedl, *Xyleborus haberkorni* Eggers, *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).

Albizzia odoratissima

Eucallacea fornicatus (Eichhoff), *Xyleborus perforans* (Wollaston).

Albizzia procera

Crossotarsus fractus Sampson, *Xylosandrus morigerus* (Blandford).

Albizzia sekbalaruguma

Cyclorhupidion sulcipenne (Eggers).

Albizzia spp.

Ambrosiodmus cichhoffi (Schreiner), *Ambrosiodmus facetus* (Schedl), *Ambrosiodmus obliquus* (LeConte), *Ambrosiodmus tropicus* (Hagedorn), *Cnestus magnus* Sampson, *Cyclorhupidion crucifer* (Hagedorn), *Cyclorhupidion crucepinne* (Schedl), *Diapus pusillinus* Chapuis, *Diapus quinquespinatus* Chapuis, *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus excavatus* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus propinquus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Eucallacea xanthopus* (Eichhoff), *Hylesinopsis seriatus* (Eggers), *Hypothenemus biseriatus* (Eggers), *Hypothenemus camerinus* (Eggers), *Leptoxyleborus concisus* (Blandford), *Perionnatus excisus* Strohmeyer, *Perionnatus grandis* Schedl, *Platypus biuncus* Blandford, *Platypus curvus* Strohmeyer, *Platypus geminatus* Chapuis, *Platypus hintzi* Schaufuss, *Platypus insularis* Strohmeyer, *Platypus jansoni* Chapuis, *Platypus lunifer* Schedl, *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Platypus secretus* Sampson, *Platypus solidus* Walker, *Platypus uncinatus* Blandford, *Premnobius ambitiosus* (Schaufuss), *Premnobius catipennis* Eichhoff, *Premnobius sex-spinosus* Eggers, *Rhopalopsclion rufus* (Schedl), *Scolytoplattypus armatus* Eggers, *Scolytoplattypus eichelbaumi* Hagedorn, *Trachyostus fecundus* (Sampson), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborinus andrevesi* (Blandford), *Xyleborinus hercae* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus clerodendronae* Schedl, *Xyleborus insignis* Browne, *Xyleborus sphecnus* Sampson, *Xyleborus spinulosus* Blandford, *Xyleborus volcuhus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford), *Xylosandrus mutilatus* (Blandford).

Albizzia stipulata

Coptodryas recidens (Sampson), *Cryphalus scabricollis* Eichhoff, *Platypus lepidus* Chapuis, *Xyleborinus andrevesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).

Albizzia sumatrana

Crossotarsus javanus Beeson, *Eucallacea fornicatus* (Eichhoff).

Albizzia warneckeii

Doliopygus coelocephalus (Schaufuss), *Doliopygus exilis* (Chapuis).

Albizzia welwitschii

Doliopygus chapuisi (Duvivier), *Platypus hintzi* Schaufuss, *Xyleborus volcuhus* (Fabricius).

Albizzia zygia

Chaetastus tuberculatus (Chapuis), *Cyclorhupidion psaltes* (Schedl), *Doliopygus brevis* (Strohmeyer), *Doliopygus chapuisi* (Duvivier), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus excavatus* (Sampson), *Doliopygus exilis* (Chapuis), *Doliopygus*

- gracilior* Schedl, *Doliopygus opifex* (Sampson), *Doliopygus perbrevis* Schedl, *Doliopygus perminutissimus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Eccoptopterus spinosus* (Olivier), *Periommatius bispinus* Strohmeyer, *Periommatius nkusii* Strohmeyer, *Periommatius substriatus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Scolytotplatypus africanus* Eggers, *Trachystus schaufussi* (Strohmeyer), *Triozastus banghaasi* (Schaufuss), *Triozastus elongatus* Schedl, *Xyleborinus sharpeae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus eichhoffianus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus voluculus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).
- Alchornea cordifolia***
Cyclorhipidion agnatieps (Schedl), *Xyleborus ruandae* Schedl.
- Alchornea coriacea***
Ambrosiodmus camphorae (Hagedorn), *Ambrosiodmus inominatus* (Schedl), *Xyleborinus signatipennis* (Schedl), *Xyleborus alluaudi* Schaufuss.
- Alchornea hirtella***
Doliopygus bidentatus (Strohmeyer), *Mimocurcus montanus* (Schedl), *Polygraphus tenuipennis* Schedl, *Premnobius mukuyae* (Schedl), *Xyleborinus similans* (Eggers), *Xyleborus comparabilis* Schedl, *Xyleborus tenellus* Schedl, *Xyleborus truncatulus* Schedl.
- Alchornea* spp.**
Scolytotplatypus kivuensis Schedl, *Xyleborus bezanosano* Schedl.
- Aleurites fordii***
Eucallacea xanthopus (Eichhoff), *Hypothenemus seriatus* (Eichhoff).
- Aleurites moluccana***
Crossotarsus externidentatus (Fairmaire), *Platypus pseudocupulatus* Schedl, *Platypus selysi* Chapuis, *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus rugatus* Blackburn, *Xyleborus whitteni* Beeson.
- Aleurites* spp.**
Eucallacea fornicatus (Eichhoff), *Xylosandrus compactus* (Eichhoff).
- Alexa imperitricia***
Dryocoetoides cristatus (Fabricius), *Monarthrum dimidiatum* (Ferrari), *Monarthrum exornatum* (Schedl), *Monarthrum proximum* Wood, *Taurodemus flavipes* (Fabricius), *Xyleborinus intersetosus* (Blandford), *Xyleborus affinis* Eichhoff, *Xyleborus semipunctatus* Eggers.
- Alibertia edulis***
Premnobius cavipennis Eichhoff.
- Allanblackia floribunda***
Phrixosoma fuscicollis (Schedl), *Polygraphus longipilis* Schedl.
- Allanblackia parvifolia***
Doliopygus serratus (Strohmeyer), *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff.
- Allanblackia* spp.**
Premnobius cavipennis Eichhoff.
- Allanblackia stuhlmannii***
Ctonoxylon flavesceus Hagedorn.
- Allogogus brunneus***
Doliopygus galerus (Schedl).
- Allophyllus africanus***
Doliopygus jurioni Schedl, *Eucallacea granosus* (Schedl), *Premnobius nodulosus* Hagedorn.
- Allophyllus cobbe***
Eucallacea fornicatus (Eichhoff).
- Allophyllus kivuensis***
Eucallacea xanthopus (Eichhoff), *Platypus impressus* (Strohmeyer), *Platypus solutus* Schedl, *Scolytotplatypus armatus* Eggers, *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff.
- Allophyllus macrocarpus***
Xyleborinus spiculatus (Schaufuss), *Xyleborinus spinosus* (Schaufuss), *Xyleborus alluaudi* Schaufuss.
- Alluaudia pourri***
Xyleborus pourriensis Schedl.
- Alluaudia procera***
Xyleborus submolestus Schedl.
- Alniophyllum fortunei***
Xyleborus atratus Eichhoff.
- Alnus acuminata***
Platypus coronatus Schedl, *Scolytodes alni* Wood.
- Alnus crispa***
Trypophloeus striatulus (Mannerheim).
- Alnus firma***
Cryphalus kytoeensis Nobuchi.
- Alnus firmifolia***
Gnathotrichus alniophagus Wood.
- Alnus fruticosa***
Alniophagus alni (Niisima), *Dryocoetes ussuriensis* Eggers, *Ernoporicus longus* (Eggers).
- Alnus glutinosa***
Dryocoetes alni (Georg), *Dryocoetes villosus* (Fabricius), *Taphrorhynchus alni* Pfeffer, *Trypodendron domesticum* (Linnaeus), *Trypophloeus alni* (Lindemann).
- Alnus hepalensis***
Xyleborus lineatus Eggers.
- Alnus hirsuta***
Alniophagus alni (Niisima), *Ambrosiodmus apicalis* (Blandford), *Ambrosiodmus hirsuta* (Eichhoff), *Platypus severini* Blandford, *Trypodendron niponicum* Blandford, *Trypodendron signatum* (Fabricius), *Xyleborus atratus* Eichhoff.
- Alnus incana***
Alniophagus alni (Niisima), *Dryocoetes alni* (Georg), *Trypophloeus alni* (Lindemann).
- Alnus japonica***
Ambrosiodmus apicalis (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Crossotarsus niponicus* Blandford, *Neopteleobius scutulatus* (Blandford), *Platypus severini* Blandford, *Xyleborinus saxenii* (Ratzeburg), *Xyleborus atratus* Eichhoff, *Xyleborus seiryorensis* Murayama.
- Alnus kamtschatica***
Ernoporicus longus (Eggers).
- Alnus maximowiczii***
Ernoporicus longus (Eggers).
- Alnus nepalensis***
Diapus quadrispinatus Chapuis, *Diapus quinquespinnatus* Chapuis, *Platypus falcatus* Strohmeyer, *Scolytotplatypus darjeelingi* Stehling, *Scolytotplatypus pubescens* Hagedorn.
- Alnus nitida***
Cryphalus fulmineus Wood, *Diapus aculeatus* Blandford, *Platypus falcatus* Strohmeyer, *Scolytotplatypus minimus* Hagedorn.
- Alnus rhombifolia***
Alniophagus aspericollis (LeConte).
- Alnus rubra***
Alniophagus aspericollis (LeConte).

Alnus rugosa*Trypophloeus striatulus* (Mannerheim).***Alnus sibirica****Ambrosiodmus apicalis* (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Cryphalus furukawai* Murayama.***Alnus sinuata****Alniphagus hirsutus* Schedl.***Alnus* spp.**

Ambrosiodmus lewisi (Blandford), *Carphothoreus alni* (Bright), *Corthylus abbreviatus* Eichhoff, *Corthylus comatus* Blandford, *Corthylus concavus* Bright, *Corthylus diligens* Wood, *Corthylus villus* Bright, *Cryphalus alni* Krivolutskaya, *Dryocoetes padi* Stark, *Gnathotrichus retusus* (LeConte), *Gymnochilus alni* Wood, *Hyloterus hirtellus* (LeConte), *Monarthrum bidentatum* Wood, *Monarthrum hoegei* (Blandford), *Monarthrum laterale* (Eichhoff), *Monarthrum umbrinum* (Blandford), *Pityophthorus alni* Blackman, *Pityophthorus abnicoleus* Wood, *Platypus otiosus* Schedl, *Platypus simulans* Schedl, *Scolytus platypus daimio* Blandford, *Scolytus platypus mikado* Blandford, *Scolytus laevis* Chapuis, *Tessrococcus spinolae* Chapuis, *Trypodendron gainensis* (Murayama), *Trypophloeus salicis* Hopkins, *Xyleborinus alni* (Nisima), *Xyleborus atratus* Eichhoff, *Xyleborus dispar* (Fabricius), *Xyleborus maiche* (Stark), *Xyleborus pfeili* (Ratzeburg), *Xyleborus seriatus* Blandford, *Xyleborus titubanter* Schedl, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus germanus* (Blandford), *Xyloterminus politus* (Say).

Alnus suaveolens*Trypophloeus rybinskii* Reitter.***Alnus tenuifolia****Alniphagus aspericollis* (LeConte), *Alniphagus hirsutus* Schedl.***Alnus tiliifolia****Scolytus platypus darjeelingi* Stebbing.***Alnus tinctoria glabra****Xyleborus atratus* Eichhoff.***Alnus viridis****Dryocoetes alni* (Georg), *Trypophloeus rybinskii* Reitter.***Aloe vera****Hypothenemus californicus* Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood.***Alphitonia excelsa****Hypothenemus melasomus* (Lea).***Alphitonia petriei****Anasa truncatus* (Erichson), *Platypus queenslandi* Schedl, *Xyleborus similis* Ferrari.***Alphitonia* spp.***Diapus nebulosus* Roberts, *Hyledius nitidicollis* (Motschulsky).***Alphonsea ventricosa****Platypus curvus* Strohmeier, *Platypus hybridus* Schedl, *Platypus indicus* Strohmeier, *Platypus secretus* Sampson.***Alstonia angustifolia****Anasa versicolor* (Sampson).***Alstonia boonei****Ambrosiodmus aegir* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus carapae* Schedl, *Doliopygus jurioni* Schedl, *Platypus hintzi* Schaufuss, *Triozastus marshalli* (Sampson), *Xyleborus consobrinus* Eggers.***Alstonia congensis****Chaetastus tuberculatus* (Chapuis), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus**erichsoni* (Chapuis), *Doliopygus lebruni* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeier), *Doliopygus unicornis* Schedl, *Periommatius grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeier), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xylechorus alluandi* Schaufuss, *Xyleborus colvulus* (Fabricius).***Alstonia scholaris****Crossotarsus externidentatus* (Fairmaire), *Platypus advena* Schedl, *Platypus jansoni* Chapuis.***Alstonia* spp.***Anasa versicolor* (Sampson), *Eccoptopterus spinosus* (Olivier), *Eucallalcea andamanensis* (Blandford), *Periommatius grandis* Schedl, *Platypus cupulatus* Chapuis, *Platypus insulindicus* Schedl, *Platypus pseudocupulatus* Schedl, *Trachyostus aterrimus* (Schaufuss), *Xyleborus minutus* Blandford, *Xylosandrus crassiusculus* (Motschulsky).***Althoffia* spp.***Platypus jansoni* Chapuis, *Platypus solidus* Walker.***Altingia excelsa****Cnestus bicoruis* (Eggers), *Diapus pusillimus* Chapuis, *Xyleborus mucronatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).***Amelanchier oralis****Scolytus amygdali* Guerin-Meneville.***Amelanchier* spp.***Chaetophloeus luterodoxus* (Casey), *Pseudopityophthorus minutissimus* (Zimmermann), *Scolytus rugulosus* (Muller).***Amerimonon brownei****Cnesinus strigicollis* LeConte.***Amherstia nobilis****Hypothenemus eruditus* Westwood.***Amonium* spp.***Xylosandrus morigerus* (Blandford).***Amoora rohataka****Coccotrypes vulgaris* (Eggers), *Crossotarsus squamulatus* Chapuis, *Platypus curvus* Strohmeier, *Platypus lepidus* Chapuis, *Xylosandrus difficilis* (Eggers).***Amoora spectabilis****Coccotrypes longior* (Eggers), *Platypus haematicollis* Schedl.***Amoora* spp.***Platypus acctabuliformis* Roberts, *Platypus hippocrepicus* Roberts, *Platypus luniger* Motschulsky, *Xylosandrus crassiusculus* (Motschulsky).***Amoora wallichii****Coccotrypes cyperi* (Beeson), *Coccotrypes salakensis* (Schedl), *Ambrosiodmus lewisi* (Blandford), *Eucallalcea wallacei* (Blandford), *Leptoxyleborus concisus* (Blandford), *Platypus curtatus* Sampson, *Scolytus platypus pubescens* Hagedorn, *Xyleborinus subgranulatus* (Eggers), *Xyleborus shoreae* (Stebbing).***Amphimas ferrugineus****Acanthotomicus sexdentatus* (Eggers), *Doliopygus chapuisi* (Duvivier), *Doliopygus serratus* (Strohmeier), *Platypus parallelus* (Fabricius), *Xyleborus colvulus* (Fabricius).***Amphimas pterocarpoides****Doliopygus unicornis* Schedl, *Mesoplatypus venustus* Schedl.***Amygdalus communis****Scolytus amygdali* Guerin-Meneville.

Amyris balsamifera

Pseudotsyanoxes securigerus (Blackman).

Anacardium excelsum

Xyleborus volutus (Fabricius).

Anacardium occidentale

Crossotarsus minax (Walker), *Eccoptopterus spinosus* (Olivier), *Hypothenemus macrolabii* (Eggers), *Xyleborinus andreweesi* (Blandford), *Xyleborinus perexiguus* (Schedl), *Xyleborus perforans* (Wollaston), *Xylosandrus manicus* (Blandford).

***Anacardium* spp.**

Xyleborinus andreweesi (Blandford), *Xylosandrus compactus* (Eichhoff).

Anagyris foetida

Liparthrum genistae (Aube).

Audira jamaicensis

Xyleborus ferrugineus (Fabricius).

***Audira* spp.**

Xylosandrus compactus (Eichhoff).

***Andropogon* spp.**

Hypothenemus pubescens Hopkins.

Angophora costata

Xyleborus perforans (Wollaston).

Angylocalyx pynaertii

Acanthotomicus angylocalyx (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus artespianus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus lecontei* Schedl, *Doliopygus medius* Schedl, *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus unicornis* Schedl, *Hypothenemus eruditus* Westwood, *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus pilosellus* (Schedl), *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Auingeria altissima

Cylindropalpus affinis Strohmeyer, *Periommatius grandis* Schedl, *Platypus nebulosus* Roberts.

Auingeria robusta

Chaetastus tuberculatus (Chapuis), *Cylindropalpus affinis* Strohmeyer, *Cylindropalpus granulatus* (Schedl), *Cylindropalpus pumilio* (Schedl), *Doliopygus coelocephalus* (Schauffuss).

***Auingeria* spp.**

Trachyostus aterrimus (Schauffuss).

***Anisopappus* spp.**

Hypothenemus grandis Schedl.

***Anisophyllea* spp.**

Doliopygus bidentatus (Strohmeyer), *Doliopygus crichsoni* (Chapuis), *Platypus solutus* Schedl, *Xyleborus alluaudi* Schauffuss.

Anisoptera cochinchinensis

Platypus shoreanus (Beeson).

Anisoptera curtisii

Webbia pabo Sampson.

Anisoptera polyandra

Cyclorhipidion sulcinoides (Schedl), *Genyocerus puer* (Schedl), *Platypus shoreanus* (Beeson), *Platypus strenuus* Schedl.

***Anisoptera* spp.**

Acanthotomicus emarginatus Browne, *Cyrtogenius anisopterae* (Browne), *Diaprus nanus* Schedl, *Diaprus*

pusillinus Chapuis, *Genyocerus diaphanus* (Schedl), *Genyocerus papuanus* Roberts, *Genyocerus puer* (Schedl), *Platypus jansonii* Chapuis, *Platypus solidus* Walker, *Xyleborus bidentatus* (Motschulsky), *Xyleborus moluccanus* Browne.

Annona cherimolia

Araptus selwaezi (Blackman).

Annona mucosa

Coccotrypes rutschuricus Eggers.

***Annona* spp.**

Euwallacea xanthopus (Eichhoff), *Hypothenemus birnanus* (Eichhoff), *Hypothenemus brunneus* (Hopkins), *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins), *Xyleborus similis* Ferrari, *Xylosandrus compactus* (Eichhoff).

Annona squamosa

Coccotrypes carpophagus (Homung), *Xyleborus perforans* (Wollaston).

Anogeissus acuminata

Ambrosiodinus lantanæ (Eggers), *Coccotrypes nubilis* (Blandford), *Polygraphus anogeissi* Wood, *Sphaerotrypes coimbatorensis* Stebbing, *Xyleborus similis* Ferrari.

Anogeissus latifolia

Platypus solidus Walker, *Platypus uncinatus* Blandford, *Sphaerotrypes coimbatorensis* Stebbing, *Sphaerotrypes globulus* Blandford, *Xyleborus perforans* (Wollaston).

***Anogeissus* spp.**

Ficicis despectus (Walker).

Anonidium manni

Chaetastus tuberculatus (Chapuis), *Coccotrypes ghesquieri* Eggers, *Coccotrypes sparsepilosus* (Eggers), *Coccotrypes subocalis* Eggers, *Cylindropalpus laudatus* (Schedl), *Doliopygus chapuisi* (Divivier), *Doliopygus citri* Schedl, *Doliopygus conradi* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Periommatius subrobustus* Schedl, *Platypus hintzi* Schauffuss, *Platypus impressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus picinus* Schedl, *Platypus schenklingi* (Strohmeyer), *Platypus solutus* Schedl, *Platypus spinulosus* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius).

Anopyxis klaineana

Cylindropalpus pumilio (Schedl), *Xyleborus affinis* Eichhoff.

***Anopyxis* spp.**

Hypothenemus eruditus Westwood.

***Anthocaryon* spp.**

Premnobius cavipennis Eichhoff.

Anthoecephalus cadamba

Dactylipalpus transversus Chapuis, *Euwallacea interjectus* (Blandford), *Platypus anthoecephali* Schedl, *Platypus oblique truncatus* Schedl, *Platypus secretus* Sampson, *Platypus solidus* Walker, *Platypus suffodiens* Sampson.

Anthocephalus spp.

Leptoxyleborus concisus (Blandford), *Platypus jansoni* Chapuis, *Platypus solidus* Walker, *Xyleborinus andrewesi* (Blandford).

Anthocleista nobilis

Chaetastus montanus Schedl, *Chaetastus subalpinus* (Schedl), *Chaetastus tuberculatus* (Chapuis), *Doliopygus chapuisi* (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus punctiventris* Schedl, *Hylesinopsis baphiac* (Schedl), *Periommatatus grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Xyleborinus diversus* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volvulus* (Fabricius).

Anthonota fragrans

Cylindropalpus auricomans (Schaufuss), *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Anthonota macrocarpa

Hypothenemus socialis (Schedl).

Anthonota macrophylla

Coptoborus usagarius (Eggers), *Coptoborus forficatus* (Schedl), *Cylindropalpus auricomans* (Schaufuss), *Cylindropalpus granulatus* (Schedl), *Cylindropalpus laudatus* (Schedl), *Doliopygus caesalpiniae* Roberts, *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Polygraphus kasukunbii* Schedl, *Premnobius longus* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus gracilipennis* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus ananici* Hagedorn, *Xyleborus ferrugineus* (Fabricius), *Xyleborus ficus* Eggers, *Xyleborus subtuberculatus* Eggers, *Xyleborus volvulus* (Fabricius).

Anthonota micraster

Coptoborus usagarius (Eggers), *Premnobius longus* Eggers.

Anthonota spp.

Cylindropalpus camerunus (Schedl), *Platypus forcipatus* Schedl.

Anthurium spp.

Xylosandrus compactus (Eichhoff).

Antiaris africana

Chaetastus tuberculatus (Chapuis), *Doliopygus brevis* (Strohmeyer), *Doliopygus conradi* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus nairobiensis* (Schedl), *Doliopygus opifex* (Sampson), *Doliopygus perbrevis* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus tenuis* (Strohmeyer), *Dryocoetoides cristatus* (Fabricius), *Periommatatus excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus spinulosus* (Strohmeyer), *Polygraphus coronatus* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Triocastus baughasi* (Schaufuss), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasicusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus pseudoambasicus* Schedl, *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Antiaris spp.

Diamerus impar Chapuis, *Doliopygus interruptus* (Sampson), *Premnobius cavipennis* Eichhoff, *Trachyostus aterrimus* (Schaufuss).

Antiaris toxicaria

Chaetastus tuberculatus (Chapuis), *Xyleborus emarginatus* Eichhoff.

Antiaris wehrtschii

Chaetastus tuberculatus (Chapuis), *Doliopygus conradi* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus leferrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus spectabilis* Schedl, *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatatus excisus* Strohmeyer, *Periommatatus grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Triocastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus volvulus* (Fabricius).

Antidesma petiolare

Eucallacca xanthopus (Eichhoff).

Antidesma spp.

Crossotarsus muiszechi Chapuis, *Diapys turgidus* Roberts, *Xylosandrus compactus* (Eichhoff).

Antrocaryon micraster

Diapys quinquespinatus Chapuis, *Doliopygus bilobatus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus leferrei* Schedl, *Hypothenemus eruditus* Westwood, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Premnobius robustulus* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus caudatus* Schedl, *Xyleborus ferrugineus* (Fabricius).

Antrocaryon namanii

Xyleborus affinis Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Aphanamixa rohituka

Eccoptopterus limbus Sampson, *Hadrodemius comans* (Sampson), *Xyleborus perforans* (Wollaston), *Xylosandrus manicus* (Blandford).

Aphananthe aspera

Eucallacca validus (Eichhoff).

Aphananthe sakalava

Platypus madagascariensis Chapuis.

Aphloia theaeformis

Mitosoma excisum Schaufuss.

Apoplanesia paniculata

Chaetophloeus minimus Wood.

Aporosa glohifera

Crossotarsus siva Beeson.

Aporosa spp.

Platypus utilis (Schedl).

Aptandra zenkeri

Doliopygus citri Schedl, *Theoborus ricini* (Eggers), *Xyleborus affinis* Eichhoff.

Apuleia forrea

Spernophthorus apuleiae Costa Lima.

Aralia decaimcana

Xyleborus atratus Eichhoff.

Aralidium pinnatifidum

Arxyleborus leprosulus Schedl, *Eccoptopterus spinosus* (Olivier).

Araucaria angustifolia

Cryptocarenum seriatus Eggers, *Pityophthorus anticus* Schedl, *Platypus araucariae* Schedl, *Tesserocerus guerinii* Chapuis, *Xyleborinus scutosus* (Eichhoff), *Xyleborus*

adelographus Eichhoff, *Xylechinosomus contractus* (Chapuis), *Xylechinosomus minimus* Schedl, *Xylechinosomus pilosus* Wood.

Araucaria araucana

Hylurgonotus antipodius (Eggers).

Araucaria bidwillii

Hylurdretonus piniarius Schedl.

Araucaria brasiliana

Araptus araucariae (Schedl), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xylechinosomus brasiliensis* (Schedl), *Xylechinosomus hirsutus* Schedl.

Araucaria cunninghamii

Coptodryas eucalyptica (Schedl), *Cryphalus araucariae* Schedl, *Cryptoxyleborus gracilior* Browne, *Diapinus nanus* Schedl, *Diapinus papuanus* Schedl, *Eucallacea barbatus* (Hagedorn), *Hyleops glabratus* Schedl, *Hylurdretonus araucariae* Schedl, *Hylurdretonus piniarius* Schedl, *Pachycotes araucariae* Schedl, *Pachycotes australis* Schedl, *Pachycotes clavatus* Schedl, *Pachycotes minor* Wood, *Platypus froggatti* Sampson, *Platypus jansoni* Chapuis, *Platypus opacideclivis* Schedl, *Platypus parallelus* (Fabricius), *Platypus queenslandi* Schedl, *Platypus solidus* Walker, *Platypus varipennis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus gratus* Schedl, *Xyleborus nitellus* Browne, *Xyleborus perforans* (Wollaston), *Xyleborus perplexus* Schedl, *Xyleborus similis* Ferrari, *Xylosandrus pseudosolidus* (Schedl).

Araucaria heterophylla

Pachycotes kuscheli Schedl.

Araucaria humsteini

Chaetophilus impar (Schedl), *Cryphalus brunneus* Browne, *Cryphalus cylindrus* Browne, *Cryphalus diversicolor* Browne, *Diapinus quinquispinatus* Chapuis, *Hylurdretonus corticinus* Wood, *Platypus jansoni* Chapuis, *Platypus praepositus* Schedl.

***Araucaria* spp.**

Ambrosiodinus compressus (Lea), *Diapinus pusillinus* Chapuis, *Hylurgonotus arnaticeps* Schedl, *Hylurgonotus solidus* (Schedl), *Hylurgonotus tuberculatus* (Eggers), *Pachycotes peregrinus* (Chapuis), *Pachycotes villosus* Schedl, *Xylechinosomus valdivianus* (Eggers), *Xylosandrus compactus* (Eichhoff).

Araujia sericifera

Araptus araujiac Brethes.

Arbutus arizonicus

Liparthrum arizonicum Wood.

Arbutus menziesii

Hylodius hirtellus (LeConte).

***Arbutus* spp.**

Cnesinus elegans Blandford, *Cnesinus elegantis* Wood, *Corthylocorus mexicanus* (Schedl), *Corthylus detrimens* Schedl, *Liparthrum arizonicum* Wood, *Micracisella nitidula* Wood, *Micracisella ocellata* Wood, *Pseudothyssanus arbuti* (Wood).

Arbutus xalapensis

Cnesinus setulosus Blandford.

Arceuthobium globosum

Pityophthorus arceuthobii Wood.

Archontophoenix alexandrae

Coccotrypes carpophagus (Hornung).

Archontophoenix cunninghamiana

Coccotrypes congonus Eggers.

Ardisia paniculata

Cnesinus strigicollis LeConte, *Hypothenemus birmanus*

(Eichhoff), *Hypothenemus brunneus* (Hopkins), *Hypothenemus hirsutus* (Wood), *Hypothenemus squamosus* (Hopkins).

***Ardisia* spp.**

Corthylus micacirrus Wood, *Platypus crassicornis* Schedl, *Xylosandrus zimmermanni* (Hopkins).

Areca catechu

Coccotrypes carpophagus (Hornung), *Coccotrypes dactyliperda* (Fabricius), *Hypothenemus eruditus* Westwood, *Xyleborus perforans* (Wollaston).

Arenga saccharifera

Coccotrypes magnus Beeson.

***Arenga* spp.**

Platypus pseudocupulatus Schedl, *Xyleborus affinis* Eichhoff.

Argyrodendron peralatum

Platypus australis Chapuis.

***Argyrodendron* spp.**

Ambrosiodinus compressus (Lea), *Platypus semi-granosus* (Sampson), *Xyleborus similis* Ferrari.

Argyrodendron trifoliolatum

Xyleborus perforans (Wollaston).

Argyrodendron trifoliolatum peralatum

Xyleborus perforans (Wollaston).

Aristolochia anguicida

Pityophthorus uemoralis Wood.

***Armanica* spp.**

Cryphalus malus Nisima.

Artemisia argentea

Liparthrum artemisiae Wollaston.

Arthrophyllum diversifolium

Coptodryas nugas (Schedl), *Cryphalus simplex* (Schedl), *Eccoptopterus limbus* Sampson, *Eccoptopterus spinosus* (Olivier), *Eucallacea fornicatus* (Eichhoff), *Eucallacea xanthopus* (Eichhoff), *Platypus pseudocupulatus* Schedl, *Xyleborus perforans* (Wollaston), *Xylosandrus morigerus* (Blandford).

Arthrosamanea altissima

Doliopygus exilis (Chapuis), *Premnobius nodulosus* Hagedorn, *Xyleborus affinis* Eichhoff, *Xyleborus ambasi-usculus* Eggers.

***Artobotrys* spp.**

Coccotrypes subovalis Eggers.

Artocarpus altilis

Cryphalus trypanoides Beeson.

Artocarpus anisophylla

Diamerus interstitialis (Lea).

Artocarpus chaplasha

Coccotrypes salakensis (Schedl), *Diapinus quinquispinatus* Chapuis, *Leptoxyleborus concisus* (Blandford), *Platypus cereclunatus* (Beeson), *Xyleborus pumilus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Artocarpus dadah

Crossotarsus wallacei (Thomson), *Xyleborinus andreweesi* (Blandford), *Xyleborus haberkorni* Eggers, *Xylosandrus mancus* (Blandford).

Artocarpus diversifolium

Cryphalus artocarpus (Schedl).

Artocarpus elastica

Acanthotomicus artocarpi (Browne), *Coccotrypes cinnamomi* (Eggers), *Coccotrypes marginatus* (Browne), *Crossotarsus terminatus* Chapuis, *Cryphalus artocarpus* (Schedl), *Cryphalus tenuis* Schedl, *Cyclorhpidion polyodon* (Eggers), *Diapinus pusillinus* Chapuis, *Diapinus quinquispinatus* Chapuis, *Eccoptopterus gracilipes* (Eichhoff), *Eucallacea destrucens* (Bland-

ford), *Hypothenemus artocarpi* Browne, *Leptoxyleborus concisus* (Blandford), *Platypus bajulus* Schedl, *Platypus biuncus* Blandford, *Platypus forficula* Chapuis, *Platypus jansoni* Chapuis, *Platypus pseudocupulatus* Schedl, *Xyleborus amphiceranoides* Hagedorn, *Xyleborus apertus* Schedl, *Xyleborus fallax* Eichhoff, *Xyleborus piceus* (Motschulsky), *Xylosandrus difficilis* (Eggers).

Artocarpus fraxinifolius

Coccotrypes vulgaris (Eggers), *Platypus secretus* Sampson.

Artocarpus heterophylla

Hyledius detersus (Chapuis), *Xyleborinus sculptilis* (Schedl).

Artocarpus hirsuta

Xyleborus perforans (Wollaston).

Artocarpus incisus

Cyrtogenius brevior (Eggers), *Platypus jansoni* Chapuis.

Artocarpus integrifolia

Diamerus curvifer (Walker), *Eucallacea interjectus* (Blandford), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Liparthrum artocarpi* Wood, *Platypus indicus* Strohmeier, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus similis* Ferrari, *Xyleborus colvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Artocarpus introcha

Platypus lepidus Chapuis.

Artocarpus kunstleri

Acanthotomicus kepongi (Schedl), *Cryphalus artocarpus* (Schedl), *Cyrtogenius brevior* (Eggers), *Diamerus luteus* Hagedorn, *Eucallacea subparallelus* (Eggers), *Hypothenemus eruditus* Westwood, *Xyleborus amphiceranoides* Hagedorn.

Artocarpus lakoocha

Ambrosiodinus dhingensis (Eggers), *Ambrosiodinus sulcatus* (Eggers), *Baioeis perianthus* (Schedl), *Coccotrypes cyperi* (Beeson), *Cryphalus dorsalis* (Motschulsky), *Cyrtogenius glaber* (Schedl), *Eucallacea icallacci* (Blandford), *Ozopemon obanus* Hagedorn, *Platypus curtatus* Sampson, *Platypus pseudocupulatus* Schedl, *Platypus signatus* Chapuis, *Platypus westwoodi* Chapuis, *Xyleborus perforans* (Wollaston), *Xyleborus pumilus* Eggers, *Xyleborus rufobrunneus* Eggers, *Xyleborus colvulus* (Fabricius).

Artocarpus lanceaefolia

Coccotrypes advena Blandford, *Coccotrypes cinnamomi* (Eggers), *Coptodryas nuxax* (Schedl), *Hypothenemus artocarpi* Browne, *Xyleborus perforans* (Wollaston).

Artocarpus nobilis

Eccoptopterus spinosus (Olivier).

Artocarpus rigida

Coccotrypes advena Blandford, *Coccotrypes cinnamomi* (Eggers), *Cryphalus tenuis* Schedl, *Hypothenemus artocarpi* Browne.

Artocarpus scortechinii

Amasa versicolor (Sampson), *Coccotrypes advena* Blandford, *Coccotrypes cinnamomi* (Eggers), *Coccotrypes marginatus* (Browne), *Coccotrypes petioli* (Browne), *Coptodryas cylindrica* (Eggers), *Cryphalus tenuis* Schedl, *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Eucallacea xanthopus* (Eichhoff), *Hypocryphalus corpulentus* Schedl, *Hypothenemus artocarpi* Browne, *Platypus insulindicus* Schedl, *Xyleborus amphiceranoides* Hagedorn.

Artocarpus spp.

Cladoctonus interruptus (Eggers), *Coccotrypes*

gedeanus (Eggers), *Coccotrypes longior* (Eggers), *Coccotrypes medius* (Eggers), *Cryphalus browni* Wood, *Cryphalus mollis* Schedl, *Cryphalus sylvicola* (Perkins), *Cyrtogenius parvus* Browne, *Eucallacea andamanensis* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Ficicis despectus* (Walker), *Genyocerus diaphanus* (Schedl), *Hadrodennius globus* (Blandford), *Hypocryphalus perminimus* (Schedl), *Hypothenemus artocarpi* Browne, *Hypothenemus eruditus* Westwood, *Leptoxyleborus concisus* (Blandford), *Platypus cupulatus* Chapuis, *Platypus forficula* Chapuis, *Platypus insulindicus* Schedl, *Xyleborinus exiguus* (Walker), *Platypus jansoni* Chapuis, *Platypus solidus* Walker, *Platypus westwoodi* Chapuis, *Xyleborus alluaudi* Schaufuss, *Xyleborus apertus* Schedl, *Xyleborus javanus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus principalis* Eichhoff, *Xyleborus pseudocylindricus* Eggers, *Xylosandrus ater* (Eggers), *Xylosandrus crassiusculus* (Motschulsky).

Artocarpus sticus

Hypothenemus artocarpi Browne.

Arundinaria alpina

Dendrochilus arundinarius Schedl.

Asimina spp.

Hypothenemus eruditus Westwood.

Asimina triloba

Hypothenemus eruditus Westwood, *Trischidias atomus* (Hopkins).

Asparagus spp.

Mimococcus congonus (Schedl), *Xylosandrus compactus* (Eichhoff).

Aspericollis spp.

Hypothenemus eruditus Westwood.

Astragalus spp.

Hypothenemus crudiac (Panzer).

Astrocarym murumura

Coccotrypes surinamensis Schedl.

Astronium graveolens

Stiphlosoma subulatum Wood, *Xyleborus colvulus* (Fabricius).

Aucoumea klaineana

Doliopygus cococephalus (Schaufuss), *Doliopygus conradii* (Strohmeier), *Doliopygus erichsoni* (Chapuis), *Doliopygus serratus* (Strohmeier), *Dryocotoides cristatus* (Fabricius), *Hypothenemus concolor* Hagedorn, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Trachyostus aterrimus* (Schaufuss), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).

Augophora costata

Xyleborus perforans (Wollaston).

Austroboxus nitidus

Crossotarsus squamulatus Chapuis, *Platypus cupulatus* Chapuis, *Platypus pseudocupulatus* Schedl.

Autranella congolensis

Chaetastus tuberculatus (Chapuis), *Coccotrypes cylindricus* (Eggers), *Diapus quinquespinatus* Chapuis, *Doliopygus arteminatus* (Schedl), *Doliopygus cococephalus* (Schaufuss), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeier), *Platypus parallelus* (Fabricius), *Trachyostus schaufussi* (Strohmeier), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).

Acicennia officinalis*Xyleborus ferrugineus* (Fabricius).***Acicennia* spp.***Xyleborus bidentatus* (Motschulsky).***Baccharus* spp.***Micracis sicainei* Blackman, *Pityophthorus paulus* Wood, *Scolytodes constrictus* Wood.***Baccaura parvifolia****Crossotarsus siporani* Schedl.***Baikaea eminii****Xyleborus multipinatus* Eggers, *Xylocleptes baikiae* Schedl.***Baikaea* spp.***Ambrosiodinus albizzianus* (Schedl), *Doliopygus interruptus* (Sampson).***Bailonella toxisperma****Cyclorhipidion agnaticeps* (Schedl).***Balanites wilsoniana****Hypothenemus eruditus* Westwood, *Hypothenemus perhipidus* (Eggers), *Hypothenemus socialis* (Schedl).***Balanocarpus heimii****Arxyleborus mediosectus* (Eggers), *Arxyleborus scabripennis* (Blandford), *Coccotrypes brunneus* (Nunberg), *Coccotrypes gedeanus* (Eggers), *Coccotrypes subcribrosus* (Blandford), *Coptodryas curvidentis* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Cryptoxyleborus dryobalanopsis* Schedl, *Cyclorhipidion agnatum* (Eggers), *Cyclorhipidion pruinatum* (Blandford), *Diapus pusillimus* Chapuis, *Diapus quinquispinatus* Chapuis, *Genyocerus abdominalis* (Schedl), *Hypothenemus cuneolus* (Schedl), *Leptoxyleborus concisus* (Blandford), *Platypus armaticeps* Schedl, *Platypus balanocarpus* Schedl, *Platypus bifurcus* (Schedl), *Platypus curtus* Chapuis, *Platypus planodeclivis* Schedl, *Platypus shoreanus* (Beeson), *Platypus transformis* Schedl, *Platytarsulus elongatus* Schedl, *Schedlia sumatrana* (Hagedorn), *Xyleborus amplexicauda* Hagedorn, *Xyleborus apertus* Schedl, *Xyleborus balanocarpus* Nunberg, *Xyleborus cylindromorphus* Eggers, *Xyleborus dolosus* Blandford, *Xyleborus emarginatus* Eichhoff, *Xyleborus macropterus* Schedl, *Xyleborus persimilis* Eggers, *Xyleborus triangi* Schedl.***Balfourodendron riedelianum****Xyleborus ferrugineus* (Fabricius).***Baloghia lucida****Chaetoptelus tricolor* (Schedl), *Xyleborus perforans* (Wollaston).***Bambusa* spp.***Hypothenemus solitarius* (Schedl), *Scolytplatypus nikado* Blandford.***Bambusa tulda****Coccotrypes barbatus* (Schedl).***Bambusa vulgaris****Chramesus gracilis* Wood, *Chramesus macrocornis* Wood, *Hylocurus nodulus* Wood.***Banksia grandis****Anasa banksiae* (Schedl).***Baphia nitida****Cyclorhipidion crucifer* (Hagedorn), *Cylindropalpus camerunus* (Schedl), *Cylindropalpus laudatus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus interjectus* Schedl, *Doliopygus nairobiensis* (Schedl), *Hypothenemus eruditus* Westwood, *Triozastus banghaasi* (Schaufuss), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers.***Baphia pubescens****Hylesinopsis baphiae* (Schedl), *Hylesinopsis quadratuberculatus* (Schedl), *Hypothenemus biserialis* (Eggers), *Hypothenemus camerunus* (Eggers), *Rhopalopselion grande* Schedl, *Rhopalopselion intermedium* Schedl, *Rhopalopselion nitidum* Schedl, *Rhopalopselion thompsoni* Schedl, *Tiarophorus intermedius* Schedl, *Xyleborus piceus* (Motschulsky).***Baphia soldheidii****Hylesinopsis quadratuberculatus* (Schedl), *Rhopalopselion grande* Schedl, *Rhopalopselion intermedium* Schedl, *Rhopalopselion nitidum* Schedl.***Baphia* spp.***Premnobius cavipennis* Eichhoff, *Rhopalopselion confusum* Eggers, *Xyleborus volutus* (Fabricius).***Barringtonia excelsa****Hypothenemus eruditus* Westwood.***Barringtonia macrostachya****Anasa versicolor* (Sampson).***Barteria fistulosa****Doliopygus lefevrei* Schedl, *Doliopygus tenuis* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Xyleborus affinis* Eichhoff.***Barteria nigriflora****Cylindropalpus camerunus* (Schedl), *Cylindropalpus granulosus* (Schedl), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus dubius* (Sampson), *Doliopygus retusus* Schedl, *Doliopygus tenuis* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Periomminatus excisus* Strohmeyer, *Phrixosoma fuscovillosa* (Schedl), *Platypus hintzi* Schaufuss, *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).***Basjoo* spp.***Hypothenemus eruditus* Westwood.***Bassia butyracea****Platypus verrucatus* (Beeson), *Scolytomimus assamensis* Schedl.***Bassia latifolia****Cryphalus scabricollis* Eichhoff, *Eccoptopterus spinosus* (Olivier), *Platypus uncinatus* Blandford, *Polygraphus anogeissi* Wood, *Scolytomimus assamensis* Schedl, *Scolytomimus minusopis* Wood, *Xyleborus perforans* (Wollaston), *Xyloctonus scolytoideus* Eichhoff.***Bauhinia alba****Hypothenemus californicus* Hopkins, *Hypothenemus columbi* Hopkins, *Hypothenemus javanus* (Eggers).***Bauhinia grandiceps****Hypothenemus brunneus* (Hopkins), *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood.***Bauhinia krugii****Hypothenemus crudiae* (Panzer).***Bauhinia malabarica****Crossotarsus emorsus* Beeson.***Bauhinia purpurea****Hypothenemus eruditus* Westwood, *Xyleborus similis* Ferrari.***Bauhinia* spp.***Crossotarsus externedentatus* (Fairmaire), *Crossotarsus squamulatus* Chapuis, *Hypothenemus eruditus* Westwood, *Xyleborus rugosipennis* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus retusus* (Eichhoff).***Bauhinia tomentosa****Hypothenemus ater* (Eggers), *Hypothenemus bauhiniae* Schedl, *Hypothenemus eruditus* Westwood, *Hypothe-*

- nemus javanus* (Eggers), *Hypothenemus morio* (Eggers), *Hypothenemus mozambicus* Eggers, *Hypothenemus seriatus* (Eichhoff), *Microrhynchus attenuatus* (Eggers), *Scolytomimus assamensis* Schedl, *Xyleborus subasperulus* Eggers, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).
- Bauhinia variegata**
Eucallacea andamanensis (Blandford), *Eucallacea fornicatus* (Eichhoff), *Hypothenemus setosus* (Eichhoff), *Xyleborus perforans* (Wollaston), *Xylosandrus discolor* (Blandford).
- Beilschmiedia bancroftii**
Platypus carbonescens (Beeson).
- Beilschmiedia corbisieri**
Doliopygus lebruni Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus criticus* (Schedl), *Perionnatus excisus* Strohmeyer, *Prennobiobius robustulus* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus ferrugineus* (Fabricius).
- Beilschmiedia gilbertii**
Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Perionnatus excisus* Strohmeyer, *Perionnatus longicollis* Chapuis, *Perionnatus subrobustus* Schedl, *Platypus beilschmiediae* Schedl, *Platypus hintzi* Schaufuss, *Platypus solutus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus consobrinus* Eggers, *Xyleborus volutus* (Fabricius).
- Beilschmiedia louisii**
Ambrosiodinus aegir (Eggers), *Cyclorhynchus agnaticus* (Schedl), *Doliopygus bitalei* Schedl, *Doliopygus rhodesianus* (Schedl), *Mesoplatus kitushi* Schedl, *Platypus orientalis* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus amanicus* Hagedorn, *Xyleborus principalis* Eichhoff.
- Beilschmiedia obtusifolia**
Xyleborus ipidia Schedl.
- Beilschmiedia sikkimensis**
Ambrosiodinus lantanae (Eggers), *Cyclorhynchus hirtum* (Hagedorn), *Diaprus spatulifer* Browne, *Eucallacea icalacci* (Blandford), *Platypus curtatus* Sampson, *Platypus lopchuenis* Schedl.
- Beilschmiedia spp.**
Ambrosiodinus eichhoffi (Schreiner), *Baiocis solominicus* Browne, *Prennobiobius cavipennis* Eichhoff, *Xylosandrus difficilis* (Eggers).
- Belanophora glomerata**
Chaetastus tuberculatus (Chapuis).
- Belotia campbelli**
Trischidias exigua Wood.
- Benzoin spp.**
Phloeosinus pulchellus Blandford, *Scolytotplatypus mikado* Blandford, *Scolytotplatypus tycon* Blandford, *Xylosandrus mutilatus* (Blandford).
- Benzoin thunbergii**
Indocryptophagus pubipennis (Blandford).
- Berberis fremontii**
Thysanococcus berbericolens Wood.
- Berberis nepalensis**
Xyleborus mussooriensis Eggers.
- Berberis spp.**
Corythylus punctatissimus (Zimmermann), *Xylosandrus brevis* (Eichhoff).
- Berchemia scandens**
Thysanococcus berchemiae Blackman.
- Berkheya spekeana**
Rhopalosiphon congonum (Schedl).
- Berlinia acuminata**
Cylindropalpus auricomans (Schaufuss), *Pityophthorus niposae* Schedl, *Platypus parallelus* (Fabricius), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus rothkirchi* Eggers, *Xyleborus volutus* (Fabricius).
- Berlinia auriculata**
Platypus hintzi Schaufuss.
- Berlinia brieji**
Platypus parallelus (Fabricius).
- Berlinia confusa**
Doliopygus berliniae Roberts, *Doliopygus caesalpiniae* Roberts.
- Berlinia grandiflora**
Pityophthorus niposae Schedl.
- Berlinia spp.**
Doliopygus bitalei Schedl, *Doliopygus chapuisi* (Duviol), *Platypus hintzi* Schaufuss, *Platypus neosolidus* Schedl, *Platypus parallelus* (Fabricius).
- Berria amonilla**
Hypothenemus brunneus (Hopkins), *Hypothenemus eruditus* Westwood.
- Bersama abyssinica**
Acanthotomus tanganyikensis (Schedl).
- Bersama spp.**
Chaetastus montanus Schedl, *Scolytotplatypus fasciatus* Hagedorn.
- Bertholletia excelsa**
Hypothenemus obscurus (Fabricius).
- Betonica officinalis**
Thamniurgus kaltenbachii (Bach).
- Betula grossa**
Platypus calamus Blandford, *Platypus lewisi* Blandford.
- Betula japonica**
Xyleborinus alni (Niisima).
- Betula lenta**
Dryocoetes betulae Hopkins, *Trypodendron betulae* Swaine.
- Betula lutea**
Dryocoetes betulae Hopkins, *Monarthrum mali* (Fitch), *Xyleborus ferrugineus* (Fabricius).
- Betula papyrifera**
Dryocoetes betulae Hopkins, *Trypodendron betulae* Swaine.
- Betula platyphylla**
Scolytus betulae Niisima.
- Betula platyphylla japonica**
Platypus lewisi Blandford, *Xyleborinus saxeseni* (Ratzeburg).
- Betula populifera**
Pseudopityophthorus asperulus (LeConte).
- Betula schmidtii**
Xyleborus atratus Eichhoff.
- Betula spp.**
Ambrosiodinus obliquus (LeConte), *Ambrosiodinus tachygraphus* (Zimmermann), *Ernoporicus fagi* (Fabricius), *Hypothenemus erudiae* (Panzer), *Pseudopityophthorus minutissimus* (Zimmermann), *Scolytus daluricus* Chapuis, *Scolytus intricatus* (Ratzeburg), *Scolytus ratzeburgi* Janson, *Taphrorychus betulae* Schedl, *Trypodendron domesticum* (Linnaeus), *Trypodendron signatum*

(Fabricius), *Xyleborus maiche* (Stark), *Xyleborus obesus* LeConte, *Xyleborus sayi* (Hopkins), *Xyleborus seriatus* Blandford, *Xyloterinus politus* (Say).

Bidens pilosa

Hypothenemus californicus Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins).

***Bignonia* spp.**

Hypothenemus crudiae (Panzer).

Biota orientalis

Ambrosiodmus rubricollis (Eichhoff), *Phloeosinus bicolor* (Brulle), *Phloeosinus hopehi* Schedl.

Bischofia javanica

Ambrosiodmus lecontei Hopkins, *Platypus biuncus* Blandford, *Platypus cavius* Strohmeier, *Xylosandrus crassiusculus* (Motschulsky).

Bixa orellana

Eucallaea fornicatus (Eichhoff), *Hypothenemus eruditus* Westwood, *Xyleborus mucronatus* Eggers, *Xylosandrus morigerus* (Blandford).

Blacus fuscipennis

Xyleborinus saxeseni (Ratzeburg).

Blighia sapida

Chaetastus tuberculatus (Chapuis), *Doliopygus brevis* (Strohmeier), *Doliopygus conradti* (Strohmeier), *Doliopygus dubius* (Sampson), *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeier), *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Platypus schencklingi* (Strohmeier), *Triozastus marshalli* (Sampson), *Xyleborinus similans* (Eggers), *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers.

***Blighia* spp.**

Prenubius cavipennis Eichhoff.

Blighia unijugata

Chaetastus tuberculatus (Chapuis).

Blighia welwitschii

Chaetastus tuberculatus (Chapuis), *Doliopygus divaricus* (Schedl), *Doliopygus erichsoni* (Chapuis), *Doliopygus jurioni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus unicornis* Schedl, *Periommatatus excisus* Strohmeier, *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeier), *Triozastus banghaasi* (Schaufuss), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).

Bobea elatior

Xyleborus ferrugineus (Fabricius), *Xyleborus perforans* (Wollaston).

Bocconia frutescens

Corthyoclon aztecum (Bright).

Boehmeria scabra

Hypothenemus californicus Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood.

***Boehmeria* spp.**

Xylosandrus morigerus (Blandford).

Boehmeria ulmifolia

Microcorthyus demissus Wood, *Microcorthyus ocularis* Wood.

***Boieldieuia* spp.**

Hypothenemus eruditus Westwood.

Bolboceranus bicolor

Doliopygus megatoma Schedl, *Periommatatus similis* Strohmeier.

Bombacopsis quinata

Xyleborus affinis Eichhoff.

Bombacopsis quintana

Xyleborus ferrugineus (Fabricius).

Bombax flammeum

Platypus hintzi Schaufuss, *Platypus impressus* (Strohmeier), *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl.

Bombax insigne

Platypus quadrifissilis Schedl, *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Bombax malabaricum

Ambrosiodmus minor (Stebbing), *Cryphalus scabricollis* Eichhoff, *Eucallaea interjectus* (Blandford), *Platypus cavius* Strohmeier, *Platypus cupulatus* Chapuis, *Platypus errans* (Sampson), *Platypus latifinis* Walker, *Platypus solidus* Walker, *Platypus uncinatus* Blandford, *Xyleborinus andrewesi* (Blandford), *Xyleborinus exiguus* (Walker), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus compactus* (Eichhoff).

***Bombax* spp.**

Platypus indicus Strohmeier.

Borassus flabellifer

Coccotrypes carpophagus (Hornung), *Coccotrypes cyperi* (Beeson), *Coccotrypes variabilis* (Beeson), *Xyleborus affinis* Eichhoff.

***Borassus* spp.**

Xyleborus affinis Eichhoff.

Borrichia arborecens

Pityophthorus borrichiae Wood.

Borrichia frutescens

Pityophthorus borrichiae Wood.

Bosqueia angolensis

Chaetastus montanus Schedl, *Chaetastus tuberculatus* (Chapuis), *Ctonoxylon bosqueiae* Schedl, *Diamerus impar* Chapuis, *Diamerus pinelioides* (Schaufuss), *Doliopygus bidentatus* (Strohmeier), *Doliopygus brevis* (Strohmeier), *Doliopygus chapuisi* (Duvivier), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeier), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus serratus* (Strohmeier), *Doliopygus unicornis* Schedl, *Eccoptopterus spinosus* (Olivier), *Hylesinopsis confusus* (Eggers), *Hylesinopsis dubius* Eggers, *Hypothenemus arvecae* (Hornung), *Mesoplatypus kitushi* Schedl, *Periommatatus excisus* Strohmeier, *Periommatatus longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeier), *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Rhopalopselion confusum* Eggers, *Triozastus banghaasi* (Schaufuss), *Triozastus caliginosus* Roberts, *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus principalis* Eichhoff, *Xyleborus scobinatus* Hagedorn, *Xyleborus volutus* (Fabricius).

Bosqueia phoberos

Chaetastus tuberculatus (Chapuis), *Doliopygus chapuisi* (Duvivier).

***Bosqueia* spp.**

Ambrosiodmus albizzianus (Schedl), *Diamerus impar* Chapuis, *Dactylipalpus grouvellei* (Blandford).

Boswellia serrata

Carphoborus boswelliae (Stebbing), *Carphoborus latus*

- Wood, *Platypus uncinatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.
- Bouea macrophylla**
Crossotarsus squamulatus Chapuis, *Crossotarsus wal-lacei* (Thomson).
- Bouea spp.**
Platypus pseudocupulatus Schedl.
- Bourreria harenensis**
Dendrosinus bourreriae Schwarz.
- Brachychiton acerifolium**
Platypus subgranosus Schedl.
- Brachychiton populneum**
Platypus subgranosus Schedl.
- Brachystegia nigerica**
Doliopygus caesalpiniae Roberts, *Platypus hintzi* Schauffuss.
- Brachystegia spp.**
Ambrosiodinus eichhoffi (Schreiner), *Doliopygus interruptus* (Sampson), *Doliopygus minimus* (Schedl), *Doliopygus spatiosus* (Schedl), *Premnobius cavipennis* Eichhoff.
- Brachystegia utilis**
Glostystus scydali (Nunberg), *Polygraphus zambianus* Beaver.
- Brackenridgea hookeri**
Crossotarsus squamulatus Chapuis, *Xylosandrus mancus* (Blandford).
- Bradfordi spp.**
Hypothenemus eruditus Westwood.
- Bramelia costaricensis**
Platypus coronatus Schedl.
- Breynia microphylla**
Eucallacea xanthopus (Eichhoff).
- Bridelia bridelifolia**
Cyclorhipidion agnatices (Schedl), *Xyleborus ruandae* Schedl.
- Bridelia retusa**
Eccoptopterus spinosus (Olivier), *Platypus uncinatus* Blandford.
- Bridelia sclereneuroides**
Hypothenemus tungamcansolus (Schedl), *Platypus hintzi* Schauffuss.
- Bridelia spp.**
Xylosandrus morigerus (Blandford).
- Bridela tomentosa**
Crossotarsus squamulatus Chapuis.
- Bridelia tulasneana**
Xyleborinus quadrispinosus (Eichhoff).
- Brillantaisia leonensis**
Dendrochilus robustus Schedl.
- Brillantaisia spp.**
Xyleborus ambaspennis Schedl, *Xyleborus clerodendronae* Schedl, *Xyleborus truncatulus* Schedl.
- Brosimum alicastrum**
Cladoctonus cubensis (Wood), *Phloeotribus setulosus* Eichhoff, *Pycnarthrum amersum* Wood.
- Brosimum spp.**
Phloeoborus scaber Erichson, *Phloeotribus biguttatus* Blandford, *Phloeotribus pilula* (Erichson), *Phloeotribus setulosus* Eichhoff, *Pycnarthrum brosimi* Wood, *Pycnarthrum subcarinatum* Wood, *Xyleborus affinis* Eichhoff.
- Broussonetia kazinoki**
Cryphalus exiguus Blandford.
- Brounea spp.**
Araptus hymenaeae (Eggers), *Araptus laevigatus* (Eggers), *Chramesus impolitus* Wood, *Dryococtoides cristatus* (Fabricius), *Taurodemus bicornutus* (Wood), *Taurodemus splendidus* (Schauffuss), *Xyleborus volutus* (Fabricius).
- Bruguiera gymorhiza**
Coccotrypes fallax (Eggers), *Cryphalus borneensis* Schedl.
- Bruguiera parviflora**
Crossotarsus squamulatus Chapuis, *Platypus forficula* Chapuis, *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.
- Bumellia costaricensis**
Corthylus zelus Wood, *Monarthrum vittatum* (Blandford), *Platypus brevicornis* Wood, *Platypus eugestus* Wood, *Platypus egestus* Wood.
- Buchanania arborescens**
Xyleborinus andrewesi (Blandford), *Xyleborus metac-nichus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).
- Buchanania lanzan**
Crossotarsus minax (Walker), *Cryphalus scabricollis* Eichhoff, *Xyleborinus andrewesi* (Blandford), *Xyleborus similis* Ferrari, *Xylosandrus gravidus* (Blandford), *Xylosandrus mancus* (Blandford).
- Buchanania latifolia**
Crossotarsus externedentatus (Fairmaire), *Platypus errans* (Sampson), *Platypus uncinatus* Blandford, *Xyleborus perforans* (Wollaston), *Xylosandrus mancus* (Blandford).
- Buchanania sessilifolia**
Crossotarsus schedli Browne, *Ficicis despectus* (Walker), *Xylosandrus crassiusculus* (Motschulsky).
- Buchanania spp.**
Crossotarsus externedentatus (Fairmaire), *Platypus jansoni* Chapuis, *Platypus negatus* Schedl, *Platypus solidus* Walker, *Xyleborinus andrewesi* (Blandford).
- Bucida buceras**
Hypothenemus erudiae (Panzer), *Hypothenemus eruditus* Westwood.
- Bucklandia populnea**
Coccotrypes nubilis (Blandford), *Diapus spatulifer* Browne.
- Buddleja spp.**
Xylosandrus compactus (Eichhoff).
- Budianana sessilifolia**
Hypothenemus javanus (Eggers).
- Buka limlia**
Crossotarsus bonvouloiri Chapuis.
- Bumelia lanuginosa**
Cnesinus strigicollis LeConte.
- Bupleurum fruticosum**
Triotennus grangeri (Peyerimhoff).
- Bupleurum spinosum**
Triotennus grangeri (Peyerimhoff), *Triotennus lepinceyi* Balachowsky.
- Burkella spp.**
Diapus oomsis Roberts, *Platypus vitensis* Roberts.
- Bursaria spinosa**
Cryphalus pilosellus Erichson.
- Bursera capallifera**
Schedlarius mexicanus Duges.
- Bursera instabilis**
Phloeoterus burserae Wood.
- Bursera microphylla**
Cactopinus desertus Bright, *Dendroterus striatus* (LeConte).

Bursera simarubra

Dendroterus eximius Wood, *Dendroterus defectus* Wood, *Dendroterus parilis* Wood, *Dendroterus resolutus* Wood, *Dendroterus sallaei* Blandford, *Dendroterus sodalis* Wood, *Schedlarius mexicanus* Duges, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus spathipennis* Eichhoff, *Xyleborus spinulosus* Blandford, *Xyleborus tumucensis* Hagedorn, *Xyleborus tolvulus* (Fabricius).

Bursera spp.

Cactopinus spinatus Wood, *Dendroterus cognatus* Wood, *Dendroterus decipiens* Wood, *Dendroterus fossifrons* Wood, *Dendroterus luteolus* (Schedl), *Dendroterus mexicanus* Blandford, *Dendroterus modicus* Wood, *Dendroterus sallaei* Blandford, *Hypothenemus crudiae* (Panzer), *Hypothenemus seriatus* (Eichhoff), *Microborus mexicanus* Wood, *Pityophthorus burserae* Wood, *Pityophthorus indigenus* Wood, *Pityophthorus nanus* Wood, *Pityophthorus nebulosus* Wood, *Xyleborus morulus* Blandford.

Bussea occidentalis

Cylindropalpus auricomans (Schaufuss), *Doliopygus brevis* (Strohmeyer), *Doliopygus conradi* (Strohmeyer), *Doliopygus mimicus* (Schedl), *Doliopygus nitidulus* Schedl, *Doliopygus opifex* (Sampson), *Doliopygus proximus* Schedl, *Doliopygus retusus* Schedl, *Hypothenemus concolor* Hagedorn, *Mesoplatypus ventricornis* Schedl, *Mesoplatypus venustus* Schedl, *Periornatus longicollis* Chapuis, *Pityophthorus busseae* Schedl, *Pityophthorus suturalis* Eggers, *Premnobius sexspinosus* Eggers, *Stephanopodius ghanaensis* (Schedl), *Triozastus elongatus* Schedl, *Xyleborinus sharpae* (Hopkins).

Bussea spp.

Scolytoptatypus occidentalis Browne.

Butea frondosa

Coccotrypes longior (Eggers), *Eucallacea andamanensis* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus manicus* (Blandford).

Butea monosperma

Crossotarsus externedentatus (Fairmaire), *Crossotarsus minax* (Walker), *Xyleborus fallax* Eichhoff, *Xyleborus similis* Ferrari, *Xylosandrus morigerus* (Blandford).

Butea spp.

Platypus solidus Walker.

Buxus macrocarpii

Scolytoptatypus obtectus Schedl.

Buxus spp.

Scolytoptatypus fasciatus Hagedorn.

Byronia sandwicensis

Xyleborus obliquus Sharp, *Xyleborus perforans* (Wollaston).

Byrsonima spp.

Xylosandrus compactus (Eichhoff).

Caesalpinia decapetala

Pityophthorus mihungensis Schedl, *Thammurgus granulicollis* Schedl, *Xylocleptes sidanus* (Schedl).

Caesalpinia melanocarpa

Spermiophthorus caesalpiniae Blackman.

Caesalpinia pulcherrima

Hypothenemus eruditus Westwood.

Caesalpinia sappan

Hyorrhynchus lewisi Blandford.

Caesalpinia sopiaria

Cyrtogenius polyphagus (Schedl).

Cajanus cajan

Araptus hymenacae (Eggers), *Cyrtogenius mihungensis* (Schedl), *Hypothenemus brunneus* (Hopkins), *Hypothenemus californicus* Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins), *Hypothenemus seriatus* (Eichhoff).

Calamus caesius

Xylosandrus morigerus (Blandford).

Calamus spp.

Ambrosiodmus brunneipes (Eggers), *Eucallacea limatus* (Schedl), *Hypothenemus eruditus* Westwood, *Leptoxyleborus concisus* (Blandford), *Xyleborus scabricollis* (Schedl), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Caldcluvia spp.

Diapys nebulosus Roberts, *Diapys unispinus* Roberts.

Calliandra confusa

Hypothenemus opacus (Eichhoff), *Hypothenemus seriatus* (Eichhoff), *Micracis grandis* Schedl, *Thysanoes neotropicalis* Wood, *Triculus senex* Schedl, *Xylosandrus zimmermanni* (Hopkins).

Calliandra houstoniana

Micracis festinus Wood.

Callicarpa spp.

Hypothenemus seriatus (Eichhoff), *Xylosandrus compactus* (Eichhoff).

Callitris articulata

Phloeosinus bicolor (Brulle).

Calodendrum capensis

Pityophthorus kenya Schedl.

Calonciton tannifolium

Scolytogenes knabi (Hopkins).

Caloncobia welwitschii

Cryptocarenum pygmaeus Schedl, *Doliopygus donisi* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus rapax* (Sampson), *Hypothenemus ater* (Eggers).

Calophyllum brasiliense antellatum

Araptus pallidus (Blackman).

Calophyllum chapelieri

Xyleborinus cupulatus (Schedl).

Calophyllum cuneifolium

Cryphalus vestitus Blandford.

Calophyllum inophyllum

Crossotarsus chalcographus Schedl, *Crossotarsus externedentatus* (Fairmaire), *Hypothenemus eruditus* Westwood, *Xyleborinus andreweesi* (Blandford), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xylosandrus manicus* (Blandford).

Calophyllum spp.

Crossotarsus squamulatus Chapuis, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Cylindropalpus sulcatus* Browne, *Diapys bispinus* Schedl, *Diapys nanus* Schedl, *Diapys nebulosus* Roberts, *Diapys papuanus* Schedl, *Diapys perpygmaeus* Roberts, *Hypocryphalus laevis* Browne, *Hypocryphalus tutuilaensis* Schedl, *Platypus platypoides* (Browne), *Platypus porcellus* Schedl, *Platypus westcودي* Chapuis.

Calophyllum tetrapterum

Crossotarsus squamulatus Chapuis, *Diapys minutissimus* Schedl, *Platypus truncatus* Browne, *Xylosandrus crassiusculus* (Motschulsky).

Calophyllum citiense

Platypus citiensis Roberts.

Calophyllum walckeri

Arixyleborus seminitens (Blandford).

Calycopteris floribunda*Crossotarsus minax* (Walker).***Calycotome spinosa****Liparthrum genistae* (Anbe), *Phlocotribus rhododactylus* (Marshall).***Calycotome* spp.***Phlocotribus cristatus* (Fairve), *Phlocotribus perfoliatus* (Wollaston).***Camellia japonica****Platypus severini* Blandford, *Xyleborus atratus* Eichhoff, *Xylosandrus brevis* (Eichhoff).***Camellia japonica rusticana****Scolytogenes camelliae* (Nobuchi).***Camellia susangua****Xylosandrus borealis* Nobuchi, *Xylosandrus brevis* (Eichhoff).***Camellia sinensis****Coccotrypes theae* Eggers, *Cryphalus vestitus* Blandford, *Eucallacea fornicatus* (Eichhoff), *Platypus lepidus* Chapuis, *Platypus uncinatus* Blandford, *Scolytomimus assamensis* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford).***Camellia* spp.***Xylosandrus mutilatus* (Blandford).***Camellia thea****Eucallacea fornicatus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).***Camellia theifera****Coccotrypes theae* Eggers, *Xylosandrus morigerus* (Blandford).***Canaoensia maxima****Xyleborus ficus* Eggers.***Camptosperma auriculata****Coccotrypes camptospermae* (Browne), *Coccotrypes cinnamomi* (Eggers), *Coccotrypes gedeanus* (Eggers), *Coccotrypes longior* (Eggers), *Coccotrypes marginatus* (Browne), *Coccotrypes medius* (Eggers), *Cryphalus tenuis* Schedl.***Camptosperma* spp.***Hypothenemus artocarpus* Browne, *Platypus hospes* Schedl, *Platypus jansoni* Chapuis, *Platypus solidus* Walker.***Cananga odorata****Crossotarsus mniszcei* Chapuis, *Platypus biuncus* Blandford.***Canarium australasicum****Platypus queenslandi* Schedl.***Canarium boivin****Mitosoma crenulatum* Chapuis, *Xyleborus alluaudi* Schaufuss.***Canarium commune****Arxyleborus malayensis* (Schedl), *Cryphalus duplosquamosus* Schedl, *Xyleborinus andrewesi* (Blandford), *Xyleborus affinis* Eichhoff.***Canarium euphyllum****Arxyleborus mediosectus* (Eggers), *Arxyleborus medius* (Eggers), *Bothinodroctonus setosus* Wood, *Coccotrypes vulgaris* (Eggers), *Coptodryas perparva* (Sampson), *Crossotarsus externodentatus* (Fairmaire), *Cryphalus scabricollis* Eichhoff, *Diapus quinquespinatus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Genyocerus diptercarpus* Browne, *Xyleborinus andrewesi* (Blandford), *Xyleborinus exiguus* (Walker), *Xyleborus bidentatus*(Motschulsky), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Canarium indicum****Crossotarsus ventricornis* Schedl, *Platypus solidus* Walker.***Canarium littorale****Cnestus aterrimus* (Eggers), *Crossotarsus schedli* Browne, *Cyclorhipidion foersteri* (Hagedorn), *Cyclorhipidion pruinatum* (Blandford), *Hadrodemius globus* (Blandford), *Hypocrypalus malagensis* Schedl, *Xyleborinus perminutissimus* (Schedl).***Canarium patentiuvrium****Cyclorhipidion pruinatum* (Blandford).***Canarium schweinfurthii****Acanthotomicus caclatus* (Schedl), *Ambrosiodmus tropicus* (Hagedorn), *Chaetastus tuberculatus* (Chapuis), *Diapus quinquespinatus* Chapuis, *Doliopygus conradi* (Strohmeyer), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megaloma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus staveri* Schedl, *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius).***Canarium* spp.***Ambrosiodmus oratus* (Eggers), *Ambrosiodmus tropicus* (Hagedorn), *Coccotrypes gedeanus* (Eggers), *Coptoborus capucinulus* (Schedl), *Cyclorhipidion cachani* (Schedl), *Cyclorhipidion foersteri* (Hagedorn), *Cyrtogenius grossepunctatus* (Schedl), *Diapus amblylaminatus* Roberts, *Diapus oomasis* Roberts, *Diapus pusillimus* Chapuis, *Eucallacea xanthopus* (Eichhoff), *Hypocryphalus striatus* Hopkins, *Leptoxyleborus concisus* (Blandford), *Mitosoma sexspinosum* Schedl, *Platypus bifidus* Schedl, *Platypus longicollis* Browne, *Platypus madagascariensis* Chapuis, *Platypus solidus* Walker, *Prennobiobius caripennis* Eichhoff, *Prennobiobius caripennis* Eggers, *Xyleborinus andrewesi* (Blandford), *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborus alluaudi* Schaufuss, *Xyleborus canarii* Browne, *Xyleborus conradi* Hagedorn, *Xyleborus ipidia* Schedl, *Xyleborus kajangensis* Schedl, *Xyleborus multispinatus* Eggers, *Xyleborus posticegranulatus* Schedl, *Xyleborus tunggali* Schedl, *Xylosandrus corthyloides* (Schedl), *Xylosandrus crassiusculus* (Motschulsky).***Canarium strictum****Coccotrypes cardamomi* Schaufuss, *Coccotrypes cyperi* (Beeson), *Crossotarsus quadricaudatus* (Strohmeyer), *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston).***Canarium citiense****Platypus citiensis* Roberts.***Canarium vulgare****Cyrtogenius brevior* (Eggers).***Canavalia maricendbergi****Chramesus pumilus* (Chapuis).***Canavalia* spp.***Chramesus punctatus* Wood, *Hypothenemus brameus* (Hopkins), *Hypothenemus seriatus* (Eichhoff).***Canavalia villosa****Chramesus pumilus* (Chapuis), *Cryptocaremus heveae* (Hagedorn), *Cryptocaremus lepidus* Wood, *Cryptocaremus seriatus* Eggers, *Hypothenemus phumicrae* (Nord-

- linger), *Hypothenemus teretis* Wood, *Stegomerus chiriquensis* Wood, *Stegomerus pygmaeus* Wood.
- Caudiosperma holacobium**
Scolytogenes kuabi (Hopkins).
- Caueia spp.**
Cucinus novateutonicus Schedl.
- Cannabris sativa**
Xylosandrus crassiusculus (Motschulsky).
- Canthium deverei**
Cylindropalpus laudatus (Schedl), *Doliopygus conradti* (Strohmeyer), *Premnobius quadrispinosus* Schedl, *Premnobius robustulus* (Schedl).
- Canthium didymum**
Cyclorhipidion foersteri (Hagedorn).
- Canthium oddoni**
Dryocoetoides cristatus (Fabricius), *Xyleborus piceus* (Motschulsky).
- Canthium spp.**
Ctonoxylon atrum Browne.
- Capparis bifolia**
Hypothenemus californicus Hopkins, *Hypothenemus gossypii* (Hopkins).
- Capparis nobilis**
Hypocryphalus nigrosetosus Schedl.
- Capsicum spp.**
Hypothenemus eruditus Westwood.
- Caragana spp.**
Scolytus schetyrewi Semenov.
- Carallia lucida**
Coccotrypes cyperi (Beeson).
- Carallia suffruticosa**
Platypus solidus Walker.
- Carallia spp.**
Platypus jansoni Chapuis.
- Carapa grandiflora**
Cylindropalpus granulatus (Schedl), *Doliopygus nitidulus* Schedl, *Pityophthorus togonus* Eggers, *Polygraphus dimorphus* Schedl, *Polygraphus granulatus* Eggers, *Polygraphus granulifer* Eggers, *Polygraphus kienensis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ambaspennus* Schedl.
- Carapa guianensis**
Liparthrum carapae Wood.
- Carapa obovata**
Coccotrypes barbatus (Schedl), *Coccotrypes fallax* (Eggers).
- Carapa procera**
Chaetastus tuberculatus (Chapuis), *Cyrtogenius mayumbensis* (Schedl), *Doliopygus carapae* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus expletus* Schedl, *Doliopygus jurionii* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Hypothenemus ater* (Eggers), *Pityophthorus togonus* Eggers, *Polygraphus binotatus* Schedl, *Polygraphus kasukumbii* Schedl, *Premnobius sexspinosus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus piceus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).
- Carapa spp.**
Ambrosiodmus tropicus (Hagedorn), *Scolytoplatypus kienensis* Schedl, *Scolytoplatypus mikado* Blandford, *Xyleborus bidentatus* (Motschulsky).
- Cardwellia sublimis**
Coptodryas eucalyptica (Schedl), *Xyleborus insulindicus* Eggers, *Xyleborus perforans* (Wollaston).
- Careya arborea**
Coccotrypes cyperi (Beeson), *Crossotarsus emorsus* Beeson, *Platypus errans* (Saunders), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus butanali* (Beeson).
- Cargotus urenus**
Coccotrypes dactyliperda (Fabricius).
- Carica papaya**
Crossotarsus externedentatus (Fairmaire), *Hypothenemus crudus* (Panzer), *Hypothenemus eruditus* Westwood, *Xyleborus perforans* (Wollaston), *Xylosandrus crassiusculus* (Motschulsky).
- Carnegiea gigantea**
Cactopinus hubhardi Schwarz.
- Carpinus betulus**
Erioporus fugi (Fabricius), *Erioporus shinanensis* Murayama, *Lymantor coryli* (Perris), *Scolytus azerbaijani* Michalski, *Scolytus jacobsoni* (Spessivtsev), *Scolytus scolytus* (Fabricius), *Scolytus varshavitchi* Michalski, *Taphrorychus villifrons* (Dufour), *Trypodendron domesticum* (Linnaeus).
- Carpinus caroliniana**
Hypothenemus eruditus Westwood.
- Carpinus carpinoideus**
Xyleborus atratus Eichhoff.
- Carpinus cordata**
Cryphalus carpini Berger, *Dryocoetes carpini* Eggers, *Scolytus claviger* Blandford, *Scolytus pubescens* Stark.
- Carpinus japonica**
Platypus severini Blandford, *Scolytus claviger* Blandford.
- Carpinus laxiflora**
Crossotarsus niponicus Blandford, *Cryphalus carpini* Berger, *Cryphalus carpinitorus* Murayama, *Dryocoetes carpinitorus* Choo & Woo, *Platypus koryoensis* (Murayama), *Platypus solidus* Walker, *Scolytus claviger* Blandford, *Taphrorychus moestus* (Blandford), *Xyleborus octidentatus* Murayama, *Xylosandrus germanus* (Blandford), *Xylosandrus mutilatus* (Blandford).
- Carpinus spp.**
Corthylus punctatissimus (Zimmermann), *Pseudopityophthorus minutissimus* (Zimmermann), *Scolytoplatypus tycon* Blandford, *Scolytus carpini* (Ratzeburg), *Scolytus intricatus* (Ratzeburg), *Scolytus pygmaeus* (Fabricius), *Xylosandrus mutilatus* (Blandford).
- Carpinus tschonoskii**
Crossotarsus niponicus Blandford, *Euwallacea validus* (Eichhoff), *Xyleborus seriatus* Blandford.
- Carpinus turczaninowii**
Xyleborus dispar (Fabricius).
- Carpodinus subrepanda**
Ctonoxylon flavescens Hagedorn, *Pseudomicracis bugekeae* (Schedl).
- Cartanopsis cuspidata**
Platypus calamus Blandford.
- Cartanopsis cuspidata sieboldii**
Platypus calamus Blandford.
- Carya alba**
Hypothenemus rotundicollis (Eichhoff).
- Carya arborea**
Crossotarsus squamulatus Chapuis.
- Carya glabra**
Hypothenemus rotundicollis (Eichhoff).
- Carya ovata**
Hypothenemus rotundicollis (Eichhoff).
- Carya spp.**
Ambrosiodmus dextexus (Wood), *Ambrosiodmus lecontei* Hopkins, *Ambrosiodmus obliquus* (LeConte),

Ambrosiodmus rubricollis (Eichhoff), *Ambrosiodmus tachygraphus* (Zimmermann), *Chramesus atkinsoni* Wood, *Chramesus hickoriae* LeConte, *Cnesinus strigicollis* LeConte, *Hylocurus bicornis* (Blackman), *Hylocurus biadatus* Wood, *Hylocurus harnedi* (Blackman), *Hylocurus prolatus* Wood, *Hylocurus rudis* (LeConte), *Hylocurus spadix* Blackman, *Hylocurus torosus* Wood, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus seriatus* (Eichhoff), *Micracisella opacicollis* (LeConte), *Monarthrum fasciatum* (Say), *Platypus compositus* Say, *Pseudopityophthorus minutissimus* (Zimmermann), *Pseudothysanoes dislocatus* (Blackman), *Scolytus quadrispinosus* Say, *Thysanoes fimbriicornis* LeConte, *Thysanoes lobllei* Blackman, *Thysanoes pallens* Wood, *Trischidias atomus* (Hopkins), *Xyleborus celsus* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus sayi* (Hopkins), *Xylosandrus germanus* (Blandford), *Xyloterinus politus* (Say).

Caryota spp.

Xylosandrus crassiusculus (Motschulsky).

Caryota urens

Eucallacea fornicatus (Eichhoff).

Casearia congensis

Platypus internedius (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus spinulosus* (Strohmeyer), *Xyleborus alliaudi* Schauffuss, *Xyleborus picus* (Motschulsky).

Casearia glomerata

Eucallacea silsagarius (Eggers), *Eucallacea velatus* (Sampson), *Eucallacea wallacei* (Blandford).

Casearia spp.

Hylesinus caseariae Wood.

Casimiroa spp.

Xylosandrus compactus (Eichhoff).

Cassia alata

Eucallacea fornicatus (Eichhoff).

Cassia didymobotrya

Xyleborus alliaudi Schauffuss, *Xyleborus picus* (Motschulsky).

Cassia fistula

Crossotarsus externedentatus (Fairmaire), *Dryocetoides cristatus* (Fabricius), *Dryocetoides pseudosolitaris* (Eggers), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus nancus* (Blandford).

Cassia florida

Hypothenemus birmanus (Eichhoff).

Cassia glauca

Hypothenemus seriatus (Eichhoff).

Cassia grandis

Coccotrypes carpophagus (Hommg).

Cassia javanica

Cryphalus capucinus Schedl, *Platypus geminatus* Chapuis.

Cassia laevigata

Ambrosiodmus aegir (Eggers).

Cassia ludheimeriana

Chramesus mimosae Blackman.

Cassia multijuga

Xylosandrus compactus (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford).

Cassia nodosa

Crossotarsus externedentatus (Fairmaire), *Hypothenemus crudiae* (Panzer).

Cassia parkii

Hypothenemus morio (Eggers).

Cassia siamea

Ambrosiodmus bostrichoides (Schedl), *Cyclorhipidion crucifer* (Hagedorn), *Doliopygus chapuisi* (Duvivier), *Doliopygus perminutissimus* (Schedl), *Eucallacea fornicatus* (Eichhoff), *Hypothenemus biseriatus* (Eggers), *Hypothenemus eruditus* Westwood, *Hypothenemus milongensis* (Eggers), *Platypus compositus* Say, *Platypus pseudocupulatus* Schedl, *Prennobiinus corthyloides* Hagedorn, *Prennobiinus longus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus alliaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus picus* (Motschulsky), *Xyleborus volutus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus germanus* (Blandford).

Cassia spp.

Acanthotomicus euphorbiae (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Doliopygus malkini* Schedl, *Hypothenemus glabripennis* (Hopkins), *Hypothenemus opacus* (Eichhoff), *Prennobiinus capucinus* Eichhoff, *Pseudothysanoes concentratus* Wood, *Pseudoxylechinus rugatus* Wood & Huang, *Xyleborus horridus* Eichhoff, *Xylosandrus compactus* (Eichhoff).

Cassia tora

Xylosandrus compactus (Eichhoff).

Cassiniae spp.

Lanurgus spathulatus Schedl, *Lanurgus xylographus* Schedl.

Cassipourea elliotii

Doliopygus ghesquieri (Schedl), *Platypus orientalis* (Strohmeyer).

Cassipourea gummiflua

Platypus sampsoni (Schedl).

Cassipourea spp.

Doliopygus nairobiensis (Schedl), *Platypus madagascariensis* Chapuis, *Platypus pennatus* Schedl, *Scolytotplatypus kiruensis* Schedl, *Xyleborinus aduncus* (Schedl), *Xyleborinus signatipennis* (Schedl), *Xylosandrus hirsutipennis* (Schedl).

Castanea argentea

Arixyleborus castaneae Schedl, *Baiocis perinimicus* (Schedl), *Coptodryas myristicae* (Schedl), *Crossotarsus nitidulus* (Schedl), *Crossotarsus terminatus* Chapuis, *Diapys quinquispinus* Chapuis, *Hadrodemius globus* (Blandford), *Platypus fraternulus* Schedl, *Platypus sexporus* (Schedl), *Platypus tenuissimus* Schedl, *Xyleborus cancellatus* Eggers, *Xyleborus emarginatus* Eichhoff, *Xyleborus fallax* Eichhoff, *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Castanea crenata

Ambrosiodmus apicalis (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Crossotarsus niponicus* Blandford, *Eucallacea validus* (Eichhoff), *Platypus calanus* Blandford, *Xyleborinus saxeus* (Ratzeburg), *Xyleborus ashuensis* Murayama, *Xyleborus gansuensis* Murayama, *Xyleborus laetus* Nisima, *Xyleborus shionomisakiensis* Murayama, *Xyleborus volutus* (Fabricius), *Xylosandrus germanus* (Blandford).

Castanea dentata

Ambrosiodmus obliquus (LeConte), *Ambrosiodmus tachygraphus* (Zimmermann), *Corthylus columbianus* Hopkins, *Hylocurus rudis* (LeConte), *Hylocurus torosus* Wood, *Micracisella opacicollis* (LeConte), *Monarthrum fasciatum* (Say), *Pseudopityophthorus asperulus* (LeConte), *Pseudopityophthorus pubescens* Blackman, *Pseudothysanoes lecontei* Blackman, *Trischidias atomus* (Hopkins).

Castanea javanica

Baiocis perinimicus (Schedl), *Diapus aculeatus* Blandford, *Diapinus minor* Schedl, *Leptoxyleborus semigranulatus* (Schedl), *Platypus sexporus* (Schedl), *Platypus strigillatus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Castanea pubinervis

Neopteleobius scutulatus (Blandford), *Xyleborus rolevus* (Fabricius).

Castanea pumila

Thysanoceros lobdellii Blackman.

Castanea spp.

Amasa schlichii (Stebbing), *Ambrosiodinus rubricollis* (Eichhoff), *Hypothenemus eruditus* Westwood, *Platypus cylindrus* (Fabricius), *Xyleborus ferrugineus* (Fabricius), *Xyleborus pfeili* (Ratzeburg), *Xyleborus sayi* (Hopkins), *Xylosandrus mutilatus* (Blandford), *Xylosterinus politus* (Say).

Castanea tungurru

Arixyleborus minor (Eggers), *Diapus aculeatus* Blandford, *Platypus sexporus* (Schedl), *Xyleborinus andreweesi* (Blandford).

Castanea vesca

Platypus cylindrus (Fabricius), *Taplorhynchus villifrons* (Drufour), *Xyleborus dryographus* (Ratzeburg), *Xyleborus monographus* (Fabricius).

Castanopsis acuminatissima

Arixyleborus sus (Schedl), *Crossotarsus chalcographus* Schedl, *Crossotarsus terminatus* Chapuis, *Diapus bispinus* Schedl, *Diapus nanus* Schedl, *Platypus apicatus* Schedl, *Platypus circulariceps* Schedl, *Platypus opacicauda* Browne, *Platypus sulcinodis* Schedl, *Webbia quadricinctus* Schedl.

Castanopsis cuspidata

Platypus quercivorus Murayama, *Xyleborus bispinus* Nobuchi, *Xyleborus japonicus* Nobuchi, *Xyleborus kaimochii* Nobuchi, *Xyleborus misatocensis* Nobuchi, *Xyleborus wakayamensis* Nobuchi.

Castanopsis cuspidata sieboldii

Ambrosiodinus lewisi (Blandford), *Platypus calamus* Blandford, *Platypus quercivorus* Murayama, *Xyleborus atratus* Eichhoff, *Xyleborus scirytorensis* Murayama.

Castanopsis fargesii

Scolytotlatypus supercilliosus Tsai & Huang.

Castanopsis hystrix

Coccotrypes nubilis (Blandford), *Diapus quadrispinatus* Chapuis, *Diapus quinquespinatus* Chapuis, *Diapus spatulifer* Browne, *Indocryphalus internedius* (Sampson).

Castanopsis indica

Ambrosiodinus lewisi (Blandford), *Eucallacea interjectus* (Blandford).

Castanopsis kawakamii

Platypus taiheizanensis (Murayama).

Castanopsis megacarpa

Arixyleborus leprosulus Schedl.

Castanopsis spp.

Amasa schlichii (Stebbing), *Ambrosiodinus apicalis* (Blandford), *Arixyleborus canaliculatus* (Eggers), *Arixyleborus granifer* (Eichhoff), *Arixyleborus granulifer* (Eggers), *Coccotrypes longior* (Eggers), *Coptoborus bicuspis* (Eggers), *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Cryphalus longior* Browne, *Cyclorhipidion agnatum* (Eggers), *Cyclorhipidion circumcisum* (Sampson), *Cyrtogenius gracilinus* Wood, *Diapus amblylaminatus* Roberts, *Diapus bilunatus* Schedl, *Diapus latespinis* Schedl, *Diapus nanodontus* Roberts, *Diapus nanus* Schedl, *Diapus ne-*

ulosus Roberts, *Diapus papuanus* Schedl, *Diapus perpygmaeus* Roberts, *Diapus pusillinus* Chapuis, *Diapus robustus* Schedl, *Diapus turgidus* Roberts, *Diapus unispinus* Roberts, *Hyledius tuberculosus* (Browne), *Leptoxyleborus depressus* (Eggers), *Leptoxyleborus scabrior* (Schedl), *Platypus cognatus* Roberts, *Platypus decens* Sampson, *Platypus platypoides* (Browne), *Platypus pseudocupulatus* Schedl, *Platypus quercivorus* Murayama, *Platypus sepaloideus* Roberts, *Scolytogenus bicolor* (Browne), *Xyleborus armiger* Schedl, *Xyleborus arnippensis* Schedl, *Xyleborus brevicollis* Browne, *Xyleborus emarginatus* Eichhoff, *Xyleborus pelliculosus* Eichhoff, *Xyleborus perforans* (Wollaston), *Xyleborus seriatus* Blandford, *Xyleborus tunggali* Schedl, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus germanus* (Blandford), *Xylosandrus morigerus* (Blandford).

Castanopsis sumatrana

Amasa schlichii (Stebbing), *Arixyleborus leprosulus* Schedl, *Arixyleborus suturalis* (Eggers), *Baiocis pernanulus* (Schedl), *Cnestus aterrinus* (Eggers), *Coptoborus longicauda* (Browne), *Coptodryas nugax* (Schedl), *Crossotarsus declivioratus* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus longicollis* Browne, *Crossotarsus schedli* Browne, *Crossotarsus squamulatus* Chapuis, *Cyclorhipidion punctatopilosum* (Schedl), *Eccoptopterus limbus* Sampson, *Indocryphalus tropicus* (Browne), *Platypus dignus* Schedl, *Platypus pseudosolidus* Schedl, *Platypus quercinus* Schedl, *Xyleborinus andreweesi* (Blandford), *Xyleborus latus* Eggers.

Castanopsis tribuloides

Crossotarsus externedentatus (Fairmaire), *Diapus quinquespinatus* Chapuis, *Eucallacea bicolor* (Blandford), *Platypus cactus* Strohmeier, *Platypus decens* Sampson, *Platypus lepidus* Chapuis, *Platypus solidus* Walker, *Platypus uncinatus* Blandford, *Xyleborus fallax* Eichhoff, *Xyleborus perforans* (Wollaston).

Castanospermum australe

Xyleborus perforans (Wollaston), *Xyleborus similis* Ferrari.

Castanospermum spp.

Ambrosiodinus compressus (Lea), *Platypus biuncus* Blandford, *Xyleborinus andreweesi* (Blandford).

Castilloa costaricensis

Platypus hians Chapuis.

Castilloa elastica

Xyleborus affinis Eichhoff.

Castilloa spp.

Doliopygus brevis (Strohmeier), *Doliopygus tenuis* (Strohmeier).

Casuarina cunninghamii

Xyleborus perforans (Wollaston).

Casuarina equisetifolia

Crossotarsus externedentatus (Fairmaire), *Xylosandrus crassiusculus* (Motschulsky).

Casuarina oligoderes

Platypus heteromorphus Schedl, *Platypus incertus* Schedl.

Casuarina spp.

Diapus papuanus Schedl, *Hypothenemus melasomus* (Lea), *Platypus navarrodeandradei* Marelli, *Prumnobius sexspinosus* Eggers.

Catha edulis

Strombophorus flavipubens Schedl.

Catostemma spp.

Platypus tiriosensis Reichardt.

Cattleya spp.

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Cayaponia ficifolia

Dendrocraniulus barbatus Schedl., *Dendrocraniulus tayuyaensis* Schedl.

Cayaponia microdonta

Dendrocraniulus consimilis Wood, *Dendrocraniulus costaricensis* Schedl., *Dendrocraniulus diversus* Wood, *Dendrocraniulus guatemalensis* (Hopkins), *Dendrocraniulus naurus* (Blandford), *Dendrocraniulus tardus* Schedl., *Dendrocraniulus vicinus* Wood, *Hypothenemus plumeriae* (Nordlinger).

Ceanothus integerrimus

Pseudothysanocs spinura Wood, *Stenoclyptus sulcatus* (Bruckl).

Ceanothus spp.

Hylocurus hirtellus (LeConte).

Cecropia peltata

Corthyus cecropii Wood, *Platypus dentatus* Dalman, *Scolytodes cecropiavorus* Wood, *Scolytodes cecropii* (Schedl), *Scolytodes piceus* (Blandford), *Tricolus cecropii* Wood, *Xyleborus affinis* Eichhoff.

Cecropia spp.

Araptus costaricensis (Schedl), *Chramesus cecropiae* Wood, *Corthyryclon tardum* Wood, *Corthyus nanus* Wood, *Corthyus strigilis* Wood, *Hypothenemus seriatus* (Eichhoff), *Hypothenemus setosus* (Eichhoff), *Hypothenemus suspectus* Wood, *Hypothenemus teretis* Wood, *Scolytodes acares* Wood, *Scolytodes anceps* Wood, *Scolytodes blandfordi* (Schedl), *Scolytodes cecropicolens* Wood, *Scolytodes chapuisi* Wood, *Scolytodes festus* Wood, *Scolytodes glabrescens* Wood, *Scolytodes jucundus* Wood, *Scolytodes naurus* (Blandford), *Scolytodes parvulus* Wood, *Scolytodes perditus* Wood, *Scolytodes suturalis* Wood, *Theoborus inculcus* (Wood), *Tricolus intrusus* Wood, *Tricolus nodifer* Blandford, *Xyleborinus tribuloides* Wood, *Xyleborus spinulosus* Blandford.

Cedrela fissilis

Ambrosiodmus catharinensis (Eggers), *Monarthrum bicallosum* (Schedl), *Platypus pulicaris* Chapuis, *Platypus quadrispinatus* Chapuis, *Tesserocerus guerini* Chapuis, *Tesserocerus insignis* (Saunders), *Xyleborinus gracilis* (Eichhoff), *Xyleborinus opimulus* (Schedl), *Xyleborinus setosus* (Eichhoff), *Xylosandrus retusus* (Eichhoff).

Cedrela mexicana

Ambrosiodmus decexulus (Wood), *Ambrosiodmus hagedorni* (Iglesias), *Ambrosiodmus lecontei* Hopkins, *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Phlocotribus setulosus* Eichhoff, *Platypus clunis* Wood, *Platypus hintzi* Schauffuss, *Pycnarthrum lucidum* Wood, *Scolytodes cedrelae* Wood, *Xyleborus colvulus* (Fabricius).

Cedrela odorata

Crossotarsus externedentatus (Fairmaire), *Doliopygus chapuisi* (Duvivier).

Cedrela serrata

Carchesiopygus psilanthurus (Beeson), *Crossotarsus musooriensis* Beeson, *Crossotarsus wilmoti* (Stebbing).

Cedrela spp.

Camptocerus auricomus Blandford, *Dendrocraniulus brasiliensis* (Schedl), *Mitosoma lobatum* Schedl, *Mitosoma obliquatum* Schedl, *Mitosoma obconiceps* Schedl, *Scolytodes swieteniae* (Blackman), *Xyleborus affinis*

Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus colvulus* (Fabricius), *Xylosandrus morigerus* (Blandford).

Cedrela toona

Platypus parallelus (Fabricius), *Platypus uncinatus* Blandford, *Xyleborus perforans* (Wollaston), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).

Cedrela toona australis

Xyleborus perforans (Wollaston).

Cedrus amarga

Tesserocerus belti Sharp.

Cedrus atlantica

Hyurgops batnensis (Brisout), *Hyurgops bonvouloiri* (Chapuis).

Cedrus deodara

Crossotarsus coniferac Stebbing, *Crossotarsus fairmairei* Chapuis, *Crossotarsus wilmoti* (Stebbing), *Crypturgus besoni* Eggers, *Crypturgus pusillus* (Gyllenhal), *Diapus quadrispinatus* Chapuis, *Diapus quinquespinatus* Chapuis, *Ips stebbingi* Strohmeyer, *Pityogenes scitus* Blandford, *Pityophthorus cedri* Wood, *Pityophthorus deodara* (Stebbing), *Polygraphus aterrimus* Strohmeyer, *Polygraphus major* Stebbing, *Polygraphus pini* Stebbing, *Scolytoplatypus raja* Blandford, *Scolytoplatypus siomio* Blandford, *Scolytus major* Stebbing.

Cedrus libani

Crypturgus pusillus (Gyllenhal), *Phlocosinus acatayi* Schedl.

Cedrus libanotica

Carphoborus minimus (Fabricius), *Crypturgus numidicus* Ferrari, *Ips robustus* (Knotek), *Phlocosinus acatayi* Schedl.

Cedrus spp.

Hyurgops glabratus (Zetterstedt), *Hyurgops palliatus* (Gyllenhal), *Scolytoplatypus fasciatus* Hagedorn, *Trypodendron lineatum* (Olivier).

Ceiba pentandra

Chaetastus tuberculatus (Chapuis), *Doliopygus serratus* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Platypus hintzi* Schauffuss, *Xylosandrus crassiusculus* (Motschulsky).

Ceiba spp.

Doliopygus dubius (Sampson), *Hypothenemus ceibae* Hopkins, *Scolytodes retifer* Wood.

Ceiba thoningii

Doliopygus conradti (Strohmeyer), *Platypus hintzi* Schauffuss, *Platypus parallelus* (Fabricius), *Xyleborus ferrugineus* (Fabricius), *Xyleborus colvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Celtis adolfi-frederici

Doliopygus retusus Schedl, *Doliopygus serratus* (Strohmeyer), *Platypus intermedius* (Schedl), *Trachyostus aterrimus* (Schauffuss).

Celtis africana

Strombophorus capensis Schedl.

Celtis brieji

Chaetastus tuberculatus (Chapuis), *Chaetastus camerunus* Schauffuss, *Doliopygus artespianatus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus ghesquieri* (Schedl), *Doliopygus jurioni* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus subdolosus* Schedl, *Hypothenemus pubipennis* (Eggers), *Periommatius circularis* Schedl, *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Platy-*

pus hintzi Schaufuss, *Platypus intermedius* (Schedl), *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborinus spinipes* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus subtuberculatus* Eggers.

Celtis brownei

Doliopygus conradti (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).

Celtis durandii

Chaetastus tuberculatus (Chapuis), *Strombophorus celtis* Schedl, *Xyleborus alluaudi* Schaufuss.

Celtis iguanae

Chramesus orinocensis Wood, *Chramesus subopacus* Schaeffer, *Hypothenemus brunneus* (Hopkins), *Micracis festicus* Wood, *Micracis sucinai* Blackman, *Phloeotribus opinus* Wood, *Phloeotribus setulosus* Eichhoff, *Phloeotribus texanus* Schaeffer, *Pseudotlysanoes subpilosus* (Wood).

Celtis kraussiana

Strombophorus celtis Schedl.

Celtis laevigata

Hylocurus parkinsoniae Blackman, *Hypothenemus brunneus* (Hopkins), *Hypothenemus eruditus* Westwood, *Phloeotribus texanus* Schaeffer.

Celtis luzonica

Xylosandrus crassiusculus (Motschulsky), *Xylosandrus ursulus* (Eggers).

Celtis mildbraedii

Ambrosiodmus opacithorax (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus arteminatus* (Schedl), *Doliopygus brevis* (Strohmeyer), *Doliopygus chapuisi* (Duvivier), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus minutissimus* (Schedl), *Doliopygus propinquus* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus subdolosus* Schedl, *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicoloris* Schedl, *Periommatatus excisus* Strohmeyer, *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Platypus solutus* Schedl, *Premmobius ambitiosus* (Schaufuss), *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus consobrinus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Celtis mississippiensis

Hypothenemus eruditus Westwood.

Celtis occidentalis

Chramesus chapuisi LeConte, *Phloeotribus dentifrons* (Blackman), *Phloeotribus frontalis* (Olivier), *Scolytus muticus* Say.

Celtis pallida

Hylocurus langstoni Blackman, *Hypothenemus sparsus* Hopkins.

Celtis sinensis

Cryphalus exiguus Blandford, *Eucallacea validus* (Eichhoff).

Celtis soyauxii

Chaetastus tuberculatus (Chapuis), *Dactylipalpus cameranus* Hagedorn, *Doliopygus bidentatus* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus medius* Schedl, *Doliopygus serratus* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Mesoplatypus ventricornis* Schedl, *Mesoplatypus venustus* Schedl, *Periommatatus bispinus* Strohmeyer, *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Platypus solutus* Schedl, *Premmobius ambitiosus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius).

***Celtis* spp.**

Ambrosiodmus albizzianus (Schedl), *Ambrosiodmus eichloffii* (Schreiner), *Ambrosiodmus obliquus* (LeConte), *Chramesus chapuisi* LeConte, *Chramesus subopacus* Schaeffer, *Coptoborus usagarius* (Eggers), *Cyclorhipidion crucifer* (Hagedorn), *Cyclorhipidion crucipenne* (Schedl), *Cylindropalpus sulcatus* Brovne, *Diamerus impar* Chapuis, *Diapus pusillimus* Chapuis, *Doliopygus coelocephalus* (Schaufuss), *Doliopygus mimicus* (Schedl), *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus unicus* (Schedl), *Hylocurus langstoni* Blackman, *Hylocurus torosus* Wood, *Phloeotribus dentifrons* (Blackman), *Periommatatus grandis* Schedl, *Platypus lesiniformis* Roberts, *Platypus solidus* Walker, *Premmobius cavipennis* Eichhoff, *Premmobius sexspinosus* Eggers, *Pseudotlysanoes lecontei* Blackman, *Scolytogenes onynganus* (Schedl), *Scolytotplatypus kivuensis* Schedl, *Scolytus fagi* Walsh, *Thysanoes finbricornis* LeConte, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus spinipes* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus dispar* (Fabricius), *Xyleborus pseudoambasius* Schedl.

Celtis wrightia

Crossotarsus javanus Beeson.

Celtis zenkerii

Ambrosiodmus camphorae (Hagedorn), *Chaetastus tuberculatus* (Chapuis), *Cyclorhipidion sulcipenne* (Eggers), *Cyclorhipidion teclae* (Schedl), *Doliopygus arteminatus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Periommatatus excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Rhopalopselion conjungens* Schedl, *Strombophorus ericis* (Schaufuss), *Trachyostus aterrimus* (Schaufuss), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus consobrinus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Centrosema plumieri

Hypothenemus eruditus Westwood, *Xylosandrus morigerus* (Blandford).

Cephalosphaera usambariensis

Doliopygus bidentatus (Strohmeyer), *Platypus hintzi* Schaufuss.

Cerasus arium*Platypus cylindrus* (Fabricius).**Ceratonía silíqua***Platypus solidus* Walker, *Taphrorychus ceratoniae* Peyerimhoff.**Ceratonía spp.***Araptus hymenacae* (Eggers).**Ceratopetalum succirubrum***Platypus australis* Chapuis.**Cerbera manghas***Scolytogenes granulatus* (Wood).**Cerbera odollum***Platypus citensis* Roberts.**Cercidiphyllum japonicum***Neopteleobius scutellatus* (Blandford), *Platypus modestus* Blandford, *Xyleborus aquilus* Blandford.**Cercidium microphyllum***Chaetophloeus parkinsoniae* (Blackman).**Cercidium spp.***Hylocurus parkinsoniae* Blackman.**Cercis canadensis***Hypothenemus cruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus rotundicollis* (Eichhoff), *Micracis suturalis* LeConte, *Micracis swainei* Blackman, *Pityophthorus laetus* Eichhoff.**Cercis siliquastrum***Carphoborus perrisi* (Chapuis).**Cercis spp.***Ambrosiodinus rubricollis* (Eichhoff), *Ambrosiodinus tachygraphus* (Zimmermann).**Cercocarpus spp.***Chaetophloeus heterodoxus* (Casey), *Cortylus punctatissimus* (Zimmermann).**Cereus spp.***Cactopinus cactophthorus* Wood, *Cactopinus carinatus* Wood, *Cactopinus granulifer* Wood, *Cactopinus microcornis* Wood, *Cactopinus nasutus* Wood, *Cactopinus niger* Wood.**Ceriops candolleana***Coccotrypes fallax* (Eggers).**Cespedesia macrophylla***Cnemionyx furcescens* Wood, *Xyleborinus dirus* Wood, *Xyleborus affinis* Eichhoff.**Cestrum scandens***Chramesus bicolor* Wood, *Scolytogenes parvatis* (Wood), *Stegomerus pygmaeus* Wood.**Cestrum spp.***Xylosandrus compactus* (Eichhoff).**Chaetacme spp.***Doliopygus excavatus* (Sampson).**Chamaecyparis formosanus***Phloeosinus arisanus* Niisima.**Chamaecyparis nootkatensis***Phloeosinus cupressi* Hopkins, *Phloeosinus keeni* Blackman, *Phloeosinus punctatus* LeConte, *Phloeosinus sequoiae* Hopkins.**Chamaecyparis obtusa***Cryphalus chamaecypariae* Niisima, *Eucallacea validus* (Eichhoff), *Phloeosinus gifuensis* Murayama, *Phloeosinus rudis* Blandford, *Phloeosinus sannohensis* Murayama, *Phloeosinus seriatus* Blandford, *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus aquilus* Blandford, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus germanus* (Blandford).**Chamaecyparis spp.***Phloeosinus lewisi* Chapuis, *Phloeosinus perlatus* Cha-puis, *Phloeosinus thujae* (Perris), *Xyleborus seriatus* Blandford.**Chamaecyparis thyoides***Phloeosinus dentatus* (Say).**Chamaedorea spp.***Coccotrypes dactyliperda* (Fabricius).**Chamerops excelsa***Coccotrypes dactyliperda* (Fabricius).**Chamerops gracilis***Coccotrypes dactyliperda* (Fabricius).**Chamerops humilis***Coccotrypes dactyliperda* (Fabricius).**Chamerops spp.***Coccotrypes dactyliperda* (Fabricius).**Chaetocarpus echinocarpus***Xyleborus parcellus* Wood.**Cheirodendron spp.***Xyleborus hawaiiensis* Perkins, *Xyleborus hiia* Samuelson, *Xyleborus kauaiensis* Perkins, *Xyleborus pcle* Samuelson, *Xyleborus temetiniicus* Beeson.**Cheirodendron trigynum***Xyleborus maitiensis* Perkins, *Xyleborus molokaiensis* Perkins.**Chenopodium ambrosioides***Cryptocarenum seriatus* Eggers.**Chenopodium spp.***Eucallacea xanthopus* (Eichhoff), *Xyleborus principalis* Eichhoff.**Chisocheton glomeratus***Leptoxyleborus concisus* (Blandford).**Chlorocodon ichteii***Mimiocurus congomus* (Schedl).**Chlorophora excelsa***Chaetastus tuberculatus* (Chapuis), *Cyclorhpidion crucipenne* (Schedl), *Cylindropalpus laudatus* (Schedl), *Cyrtogenius chlorophorae* (Schedl), *Doliopygus contradii* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus gluesquierci* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Eucallacea fornicatus* (Eichhoff), *Hypothenemus biseriatus* (Eggers), *Perionnatus angustior* Schedl, *Perionnatus excisus* Strohmeyer, *Perionnatus grandis* Schedl, *Perionnatus nitidicollis* Strohmeyer, *Platypus hintzi* Schauffuss, *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus consobrinus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus subtuberculatus* Eggers, *Xyleborus volucrus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).**Chloroxylon scietenia***Cnestus magnus* Sampson, *Platypus solidus* Walker, *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).**Chomelia spp.***Eucallacea xanthopus* (Eichhoff), *Platypus orientalis* (Strohmeyer).**Chrysalidocarpus spp.***Coccotrypes carpophagus* (Hornung).**Chrysobalanus spp.***Xylosandrus zimmermanni* (Hopkins).

Chrysolepsis spp.

Monarthrum scutellare (LeConte).

Chrysophyllum africanum

Coccotrypes rotundicollis (Eggers), *Cylindropalpus interpositus* (Schedl), *Cylindropalpus pumilio* (Schedl), *Cylindropalpus pusillus* (Schedl), *Doliopygus bidentatus* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus lebruni* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Periommatius grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus spinulosus* (Strohmeyer), *Polygraphus ruandae* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ficus* Eggers.

Chrysophyllum albidum

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Xyloctonus quadricinctus* Schedl.

Chrysophyllum boivinianum

Mitosoma accuratum Schaufuss, *Mitosoma excisum* Schaufuss, *Mitosoma obliquatum* Schedl, *Mitosoma paulianum* Schedl, *Platypus madagascariensis* Chapuis, *Xyleborinus diapiformis* (Schedl), *Xyleborinus forficuloides* (Schedl), *Xyleborinus spiculatus* (Schaufuss), *Xyleborinus spinosus* (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xyleborus antaisaka* Schedl.

Chrysophyllum lacourtianum

Cylindropalpus camerunus (Schedl), *Cylindropalpus interpositus* (Schedl), *Cylindropalpus pumilio* (Schedl), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus fulgens* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus ater* (Eggers), *Periommatius excisus* Strohmeyer, *Platypus intermedius* (Schedl), *Trachyostus aterrinus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson).

Chrysophyllum ludovicianum

Dactylipalpus camerunus Hagedorn, *Hypothenemus biseriatus* (Eggers), *Xyleborus africanus* Eggers.

Chrysophyllum metallicum

Cylindropalpus affinis Strohmeyer.

Chrysophyllum natalensis

Cylindropalpus affinis Strohmeyer.

Chrysophyllum perpulchrum

Cylindropalpus affinis Strohmeyer, *Cylindropalpus pumilio* (Schedl), *Xyleborus ambasiusculus* Eggers.

Chrysophyllum pruniferum

Chaetastus tuberculatus (Chapuis), *Cylindropalpus interpositus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus maynei* Schedl, *Doliopygus negatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius*

longicollis Chapuis, *Platypus hintzi* Schaufuss, *Platypus schenklingsi* (Strohmeyer), *Trachyostus aterrinus* (Schaufuss), *Xyleborus affinis* Eichhoff, *Xyleborus multi-spinatus* Eggers, *Xyleborus colutus* (Fabricius).

Chrysophyllum roxburghii

Coptodryas alpha (Beeson), *Coptodryas chrysophylli* (Eggers), *Coptodryas perparva* (Sampson), *Platypus forficula* Chapuis, *Platypus pseudocupulatus* Schedl.

Chrysophyllum spp.

Cylindropalpus auricomans (Schaufuss), *Cylindropalpus granulatus* (Schedl), *Cylindropalpus pertinax* (Schedl), *Dactylipalpus grouvellei* (Blandford), *Doliopygus retusus* Schedl, *Hypothenemus camerunus* (Eggers), *Premnobius capicennis* Eichhoff, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyloctonus opacus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Chrysophyllum subnudum

Xyleborus alluaudi Schaufuss.

Cicer arietinum

Araptus hymenaeae (Eggers).

Cillaeus contractus

Strombophorus vagans Schedl.

Cinchona calisaya

Eucallacca fornicatus (Eichhoff).

Cinchona ledgeriana

Ambrosiodmus aegir (Eggers), *Crossotarsus horui* Schedl, *Eccoptopterus spinosus* (Olivier), *Eucallacca xanthopus* (Eichhoff), *Hylesinopsis seriatus* (Eggers), *Rhopalosiphon congoum* (Schedl), *Scolytotplatypus hova* Schaufuss, *Xyleborus principalis* Eichhoff, *Xyleborinus quadrispinosus* (Eichhoff).

Cinchona spp.

Arxyleborus tuberculatus (Eggers), *Cnestus aterrinus* (Eggers), *Cnestus bicornis* (Eggers), *Coptoborus usagarius* (Eggers), *Eucallacca xanthopus* (Eichhoff), *Leptoxyleborus concisus* (Blandford), *Scolytotplatypus nitidus* Eggers, *Scolytotplatypus perminis* Schaufuss, *Xyleborinus andrewesi* (Blandford), *Xyleborinus collarti* (Eggers), *Xyleborus emarginatus* Eichhoff, *Xyleborus mucronatus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Cinchona succirubra

Ambrosiodmus camphorae (Hagedorn), *Platypus madagascariensis* Chapuis, *Scolytotplatypus hova* Schaufuss, *Xyleborinus aduncus* (Schedl), *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborus malgasicus* Schedl.

Cinnamomum baileyianum

Xyleborus perforans (Wollaston).

Cinnamomum camphora

Ambrosiodmus lewisi (Blandford), *Arxyleborus camphorae* (Eggers), *Baiocis perinimicus* (Schedl), *Cnestus murayamai* Schedl, *Coptodryas izuensis* (Murayama), *Crossotarsus externedentatus* (Fairmaire), *Diapus minor* Schedl, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Leptoxyleborus semigranulatus* (Schedl), *Phloeosinus camphoratus* Tsai & Yin, *Phloeosinus kumanotoensis* Murayama, *Phloeosinus machilus* (Schedl), *Phloeosinus seriatus* Blandford, *Platypus fraterculus* Schedl, *Platypus substriatus* (Schedl), *Platypus vethi* Strohmeyer, *Scolytotplatypus mikado* Blandford, *Taphrorychus taradakensis* (Murayama), *Xyleborus atratus* Eichhoff, *Xyleborus clerodendronae* Schedl, *Xyleborus kumanotoensis* Murayama, *Xyleborus magnus* Niisima, *Xyleborus perforans* (Wollaston), *Xylosandrus compactus* (Eichhoff), *Xylosandrus*

crassiusculus (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus mutilatus* (Blandford).

Cinnamomum cecicodaphne

Crossotarsus cinnamomi Beeson, *Crossotarsus squamulatus* Chapuis, *Hadrodemius pseudocomans* (Eggers), *Xylosandrus gracilis* (Blandford).

Cinnamomum impressinervium

Indocryptophalus machili Wood.

Cinnamomum iners

Cnestus aterrimus (Eggers), *Crossotarsus squamulatus* Chapuis, *Crossotarsus terminatus* Chapuis, *Xyleborinus exiguus* (Walker), *Xylosandrus compactus* (Eichhoff).

Cinnamomum japonicum

Indocryptophalus pubipennis (Blandford), *Xylosandrus brevis* (Eichhoff).

Cinnamomum mairei

Amasa amputatus (Blandford).

Cinnamomum obtusifolium

Ambrosiodinus lewisi (Blandford), *Sucus niisimai* (Eggers), *Xylosandrus subsimilis* (Eggers).

Cinnamomum randaense

Xyleborus atratus Eichhoff.

***Cinnamomum* spp.**

Amasa amputatus (Blandford), *Ambrosiodinus rubricollis* (Eichhoff), *Crossotarsus reagentensis* Niisima & Murayama, *Crossotarsus squamulatus* Chapuis, *Eucallacea barbatus* (Hagedorn), *Eucallacea fornicatus* (Eichhoff), *Hadrodemius globus* (Blandford), *Platypus capito* Browne, *Platypus secretus* Sampson, *Platypus semigranulosus* (Sampson), *Platypus solidus* Walker, *Platypus strenuus* Schedl, *Xyleborinus andrewesi* (Blandford), *Xyleborus adusticollis* (Motschulsky), *Xyleborus arctocylindrus* Schedl, *Xyleborus justus* Schedl, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus arcuatus* (Sampson), *Xylosandrus ater* (Eggers), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus compactus* (Eichhoff).

Cinnamomum zeylanicum

Coccotrypes cinnamomi (Eggers), *Coccotrypes dactyliperda* (Fabricius), *Coptodryas nudipennis* (Schedl), *Xyleborinus andrewesi* (Blandford), *Xyleborinus subgranulatus* (Eggers), *Xyleborus perforans* (Wollaston), *Xylosandrus compactus* (Eichhoff), *Xylosandrus nudipennis* (Schedl).

Cistanthera leplaei

Cylindropalpus auricomans (Schaufuss), *Doliopygus serratus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl.

Cistanthera papaverifera

Doliopygus conradti (Strohmeyer), *Xyleborus africanus* Eggers.

***Cistanthera* spp.**

Doliopygus conradti (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Poligraphus kasukumbii* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xylosandrus compactus* (Eichhoff).

Citrus aurantifolia

Hypothenemus columbi Hopkins, *Hypothenemus seriatus* (Eichhoff), *Theoborus ricini* (Eggers).

Citrus aurantium

Ambrosiodinus asperatus (Blandford), *Doliopygus citri* Schedl, *Eucallacea fornicatus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Xyleborus affinis* Eichhoff.

Citrus aurantium decumana

Xyleborus perforans (Wollaston).

Citrus decumana

Hypothenemus eruditus Westwood.

Citrus grandis

Xyleborus affinis Eichhoff.

Citrus maxima

Xyleborus perforans (Wollaston).

Citrus medica limonum

Hypothenemus eruditus Westwood.

Citrus sinensis

Cladoctonus interruptus (Eggers), *Xyleborus horridatus* Wood, *Xyleborus perforans* (Wollaston).

***Citrus* spp.**

Ambrosiodinus obliquus (LeConte), *Crossotarsus bonconloui* Chapuis, *Eucallacea fornicatus* (Eichhoff), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus erudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus pubipennis* (Eggers), *Sucus niisimai* (Eggers), *Xyleborus ferrugineus* (Fabricius), *Xyleborus nevai* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus spinulosus* Blandford, *Xylosandrus compactus* (Eichhoff).

Cladrastris shikokiana

Hylesinus costatus Blandford.

Claoxylon polot

Xylosandrus morigerus (Blandford).

Clausena anisata

Eucallacea xanthopus (Eichhoff).

Cleistanthus polystachyus

Phrixosoma fuscicollis (Schedl).

***Cleistanthus* spp.**

Doliopygus conradti (Strohmeyer), *Doliopygus lebruni* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus tenuis* (Strohmeyer), *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer).

***Cleistocalyx* spp.**

Platypus yasyasi Roberts.

Cleistopholis glauca

Coccotrypes subovalis Eggers.

Cleistopholis patens

Acanthotomicus sexdentatus (Eggers), *Chaetastus tuberculatus* (Chapuis), *Doliopygus conradti* (Strohmeyer), *Eccoptopterus spinosus* (Olivier), *Platypus hintzi* Schaufuss, *Triozastus marshalli* (Sampson), *Xylosandrus compactus* (Eichhoff).

***Cleistopholis* spp.**

Premnobius caripennis Eichhoff.

Clematis cirrhosa

Xyloleptes biuncus Reitter.

Clematis orientalis

Taphronurgus exul (Reitter).

Clematis sinensis

Thamnurgus elegans Schedl, *Thamnurgus granulicollis* Schedl, *Xyloleptes sidanus* (Schedl).

***Clematis* spp.**

Chramesus incomptus Wood, *Dacnophthorus clematis* (Wood), *Dacnophthorus rallus* (Wood), *Hypothenemus brunneus* (Hopkins).

Clematis vitalba

Xyloleptes bispinus (Duftschmidt).

Clerodendron colebrookianum

Eucallacea fornicatus (Eichhoff).

Clerodendron infortunatum

Eucallacea fornicatus (Eichhoff), *Hadrodemius metacomans* (Eggers).

***Clerodendron* spp.**

Eucallacea fornicatus (Eichhoff), *Platypus solutus*

- Schedl, *Xyleborus clerodendronae* Schedl, *Xylosandrus compactus* (Eichhoff).
- Clerodendron squamatum**
Hypothenemus erudiae (Panzer).
- Clerodendron villosum**
Cnestus aterrimus (Eggers).
- Clethra barbinervis**
Crossotarsus niponicus Blandford.
- Clethra hondurensis**
Corthylus mexicanus Schedl, *Xyleborus affinis* Eichhoff.
- Cleyera japonica**
Eucallacea validus (Eichhoff), *Platypus calamus* Blandford, *Xyleborus aquilus* Blandford, *Xyleborus atratus* Eichhoff, *Xyleborus seriatus* Blandford, *Xylosandrus germanus* (Blandford).
- Cleyera spp.**
Xyleborus octiesdentatus Murayama.
- Clidemia hirta**
Xylosandrus morigerus (Blandford).
- Clidemia spp.**
Xylosandrus compactus (Eichhoff).
- Clitandra spp.**
Xyleborinus gracilipennis (Schedl).
- Clitandra staudtii**
Ctonoxylon acuminatus Schedl, *Doliopygus expletus* Schedl, *Platypus orientalis* (Strohmeyer), *Polygraphus apicalis* Schedl, *Polygraphus majusculus* Schedl, *Polygraphus ruandae* Schedl.
- Clitoria ternatea**
Hypothenemus eruditus Westwood.
- Cloaxylon spp.**
Platypus longissimus Roberts.
- Clusia flava**
Phrixosoma clusiae Wood.
- Clusia spp.**
Ambrosiodmus ocellatus (Wood), *Corthylus additus* Wood, *Corthylus diligens* Wood, *Microborus aberrans* Wichmann, *Microborus lectus* Wood, *Monarthrum bicolor* (Ferrari), *Monarthrum lobatum* (Ferrari), *Monarthrum nvernanni* (Schedl), *Monarthrum vittatum* (Blandford), *Phloeotribus atavus* Wood, *Phloeotribus collaris* Chapuis, *Scolytodes canaliculus* Wood, *Scolytodes clusiaceolens* Wood, *Scolytodes clusiae* Wood, *Scolytodes clusiavornis* Wood, *Scolytodes comitabilis* Wood, *Scolytodes crinalis* Wood, *Scolytodes fulmineus* Wood, *Scolytodes naevius* Wood, *Scolytodes onnateus* Wood, *Scolytodes praeceps* Wood, *Scolytodes proximus* Wood, *Scolytodes tardus* Wood.
- Clytastoma binatum**
Cnesinus setulosus Blandford.
- Coccoceras muticum**
Crossotarsus schedli Browne, *Crossotarsus squamulatus* Chapuis, *Platypus lunifer* Schedl.
- Coccoceras spp.**
Crossotarsus cinnamatus Chapuis.
- Coccoloba laurifolia**
Cnesinus strigicollis LeConte, *Hypothenemus eruditus* Westwood.
- Coccothrinax alta**
Hypothenemus seriatus (Eichhoff).
- Coccothrinax argentea**
Coccotrypes carpophagus (Hornung).
- Coccothrinax spp.**
Coccotrypes dactyliperda (Fabricius), *Coccotrypes distinctus* (Motschulsky).
- Cochlospermum gossypium**
Platypus uncinatus Blandford.
- Cocos australis**
Coccotrypes circumdatus Fonseca.
- Cocos nucifera**
Platypus jansoni Chapuis, *Platypus solidus* Walker, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus rotulus* (Fabricius), *Xylosandrus morigerus* (Blandford).
- Cocos plumosa**
Coccotrypes dactyliperda (Fabricius).
- Cocos romanzoffiana**
Coccotrypes dactyliperda (Fabricius).
- Cocos spp.**
Coccotrypes dactyliperda (Fabricius).
- Coelocaryon preussii**
Chaestatus tuberculatus (Chapuis), *Platypus parallelus* (Fabricius), *Xyleborus alluandi* Schauffuss, *Xylosandrus crassiusculus* (Motschulsky).
- Coelocaryon spp.**
Mesoplatypus tenuis Schedl, *Premnobius cavipennis* Eichhoff.
- Coelostegia griffithii**
Platypus cupulatus Chapuis.
- Coffea arabica**
Cnesinus adusticus Wood, *Cnesinus coffeae* Schedl, *Cnesinus gracilis* Blandford, *Cnesinus meris* Wood, *Cnesinus robai* Blackman, *Hylocurus elegans* Eichhoff, *Hypothenemus hampei* (Ferrari), *Hypothenemus opacus* (Eichhoff), *Monarthrum exornatum* (Schedl), *Phloeoborus punctatoriosus* Chapuis, *Xyleborus ferox* Blandford, *Xylosandrus compactus* (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford).
- Coffea bukobensis**
Hypothenemus seriatus (Eichhoff), *Xylosandrus compactus* (Eichhoff).
- Coffea canephora**
Xylosandrus compactus (Eichhoff).
- Coffea canephora robusta**
Xylosandrus morigerus (Blandford).
- Coffea excelsa**
Xylosandrus morigerus (Blandford).
- Coffea hybrida**
Dryococtius coffeae (Eggers), *Xylosandrus morigerus* (Blandford).
- Coffea laurentii**
Xylosandrus discolor (Blandford).
- Coffea liberica**
Coccotrypes confusus (Eggers), *Hypothenemus hampei* (Ferrari), *Xyleborus haberkorni* Eggers, *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).
- Coffea quillon**
Xylosandrus compactus (Eichhoff).
- Coffea robusta**
Cnestus bicornis (Eggers), *Cryptocarenum lepidus* Wood, *Dryocoetoides cristatus* (Fabricius), *Hypothenemus eruditus* Westwood, *Hypothenemus hampei* (Ferrari), *Taurodemus perebeae* (Ferrari), *Xylosandrus compactus* (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford).
- Coffea spp.**
Ambrosiodmus aegir (Eggers), *Ambrosiodmus lecontei* Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus grandis* Schedl, *Hypothenemus libericensis*

(Hopkins), *Hypothenemus multidentatus* (Hopkins), *Hypothenemus plumeriacae* (Nordlinger), *Hypothenemus solitarius* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus haberkorni* Eggers, *Xyleborus mucronatus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xyleborus volutus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus retusus* (Eichhoff).

Coffea stenophylla

Xylosandrus compactus (Eichhoff).

Cola acuminata

Doliopygus conradti (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Mioceryphalus attenuatus* (Eggers), *Xylosandrus morigerus* (Blandford).

Cola caricifolia

Rhopalopselion thompsoni Schedl, *Tiarophorus intermedius* Schedl.

Cola chlamydantha

Doliopygus conradti (Strohmeyer), *Doliopygus medius* Schedl, *Hypothenemus eruditus* Westwood, *Perionnatus excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius).

Cola congolana

Doliopygus conradti (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus colae* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus alluaudi* Schaufuss.

Cola diversifolia

Rhopalopselion atrum Eggers.

Cola gigantea

Doliopygus pseudoserratus Roberts, *Doliopygus punctiventris* Schedl, *Platypus hintzi* Schaufuss, *Trachyostus aterrimus* (Schaufuss).

Cola griseiflora

Doliopygus conradti (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus expletus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus colae* (Schedl), *Hypothenemus eruditus* Westwood, *Hypothenemus pubipennis* (Eggers), *Hypothenemus socialis* (Schedl), *Platypus impressus* (Strohmeyer), *Xyleborus alluaudi* Schaufuss.

Cola nitida

Cylindropalpus auricomans (Schaufuss), *Doliopygus serratus* (Strohmeyer), *Mioceryphalus attenuatus* (Eggers), *Perionnatus excisus* Strohmeyer, *Xyleborus barunbuensis* Eggers, *Xylosandrus compactus* (Eichhoff).

***Cola* spp.**

Araptus politus (Blandford), *Doliopygus propinquus* Schedl, *Doliopygus serratus* (Strohmeyer).

***Coleus* spp.**

Eucallacea xanthopus (Eichhoff), *Xyleborinus forficuloides* (Schedl), *Xyleborinus forficulus* (Eggers).

Colophylum walkeri

Platypus furcatus Blandford.

Colutea arborescens

Phloeotribus brevicollis (Kolenati).

Colvillea racemosa

Platypus parallelus (Fabricius).

***Colvillea* spp.**

Eucallacea fornicatus (Eichhoff).

Colydium elongatum

Xyleborinus saxeseni (Ratzeburg).

Colydium lineola

Xyleborinus saxeseni (Ratzeburg).

Combretodendron africanum

Cylindropalpus auricomans (Schaufuss), *Doliopygus lefevrei* Schedl, *Doliopygus retusus* Schedl, *Doliopygus semipilosus* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus scobinatus* Hagedorn.

Combretodendron macrophyllum

Chaetastus persimilis (Schedl), *Chaetastus tuberculatus* (Chapuis), *Coccotrypes cylindricus* (Eggers), *Cylindropalpus granulatus* (Schedl), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus retusus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus semipilosus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicomis* Schedl, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius).

***Combretodendron* spp.**

Ambrosiodinus tropicus (Hagedorn), *Platypus hintzi* Schaufuss, *Prennobia caxipennis* Eichhoff.

Combretum gueinzii

Xyleborus ugandensis Schedl.

Combretum oblongum

Chaetastus tuberculatus (Chapuis), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus erichsoni* (Chapuis), *Doliopygus galerus* (Schedl), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus pygmaeolus* Schedl, *Trachyostus schaufussi* (Strohmeyer).

Combretum paniculatum

Scolytoplatus armatus Eggers.

***Combretum* spp.**

Doliopygus erichsoni (Chapuis), *Scolytus binodus* Wood.

Commidendron robustum

Xyleborinus aemulus (Wollaston).

***Commiphora* spp.**

Mitosoma obconiceps Schedl, *Platypus madagascariensis* Chapuis.

Condalia obtusifolia

Chramesus subopacus Schaeffer, *Hylocurus parkinsoniae* Blackman, *Hypothenemus brunneus* (Hopkins), *Phloeotribus dentifrons* (Blackman), *Phloeotribus texanus* Schaeffer, *Pseudothysanoes heliura* Wood.

***Condalia* spp.**

Chramesus subopacus Schaeffer, *Hypothenemus erectus* LeConte, *Hypothenemus sparsus* Hopkins.

Conocarpus erectus

Cnesinus setulosus Blandford, *Cryptocarenum seriatus* Eggers, *Hypothenemus javanus* (Eggers), *Micraciscella nanula* (LeConte).

Conopharyngia durissima

Phrixosoma garciniae (Schedl), *Phrixosoma uniseriata* (Eggers), *Polygraphus tenuis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky).

Conopharyngia holzkii

Xyloctonus niger Schedl.

Conopharyngia spp.

Chaetastus tuberculatus (Chapuis), *Platypus impressus* (Strohmeyer), *Xyleborinus collarti* (Eggers).

Conostegia oerstediana

Amphicranus fastigiatus Blandford, *Corthyceylon moridum* Wood, *Corthylocurus barbatus* (Blandford), *Corthylus concisus* Wood, *Microcorthylus demissus* Wood, *Microcorthylus ocularis* Wood, *Microcorthylus pumilus* Wood, *Monarthrum bidens* (Blandford), *Monarthrum pennatum* (Schedl), *Monarthrum proprium* Wood, *Taurodemus salvini* (Blandford), *Xyleborus meritus* Wood, *Xyleborus squamulatus* Eichhoff.

Copaifera langsdorfi

Xyleborus parcellus Wood.

Copaifera spp.

Xyleborus spathipennis Eichhoff.

Coprosma spp.

Xylosandrus compactus (Eichhoff).

Cordia caffra

Lamurgus podocarpi Schedl.

Cordia dichotoma

Eucallacea andamanensis (Blandford), *Xylosandrus mancus* (Blandford).

Cordia grandis

Coptodryas alpha (Beeson), *Eucallacea andamanensis* (Blandford).

Cordia millenii

Chaetastus tuberculatus (Chapuis).

Cordia myxa

Xyleborus perforans (Wollaston), *Xylosandrus mancus* (Blandford).

Cordia senegalensis

Doliopygus bulbifer Schedl.

Cordia spp.

Coptoborus pseudotenius (Schedl), *Doliopygus excavatus* (Sampson), *Doliopygus nairobiensis* (Schedl), *Hypothenemus seriatus* (Eichhoff), *Theoborus micarius* (Wood), *Xylosandrus compactus* (Eichhoff).

Cordyline australis

Platypus apicalis (White).

Cordyline terminalis

Xyleborus ferrugineus (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus rugatus* Blackburn.

Cornus controversa

Hyporhynchus lewisi Blandford, *Pseudohyporhynchus wadai* Murayama, *Sucus niisimai* (Eggers).

Cornus florida

Hypothenemus eruditus Westwood, *Xyleborinus saxe-seni* (Ratzeburg), *Xylosandrus germanus* (Blandford).

Cornus macrophylla

Scolytotplatypus minimus Hagedorn, *Scolytotplatypus raja* Blandford.

Cornus spp.

Ambrosiodmus rubricollis (Eichhoff), *Corthylus punctatissimus* (Zimmermann), *Hypothenemus eruditus* Westwood, *Scolytotplatypus daimio* Blandford, *Scolytotplatypus mikado* Blandford, *Xyleborus sayi* (Hopkins), *Xylosandrus mutilatus* (Blandford).

Cornus tartarica

Hypothenemus corni Kurenzov.

Corylus avellana

Dryocoetes alni (Georg), *Lymanator aceris* (Lindemann), *Lymanator coryli* (Perris).

Corylus mandshurica

Cryphalus mandschuricus Eggers, *Xyleborus maiche* (Stark).

Corylus spp.

Corthylus punctatissimus (Zimmermann), *Cryphalus mandschuricus* Eggers, *Scolytus laevis* Chapuis, *Tiarophorus scrutator* (Pandelle).

Cossia arabica

Coccotrypes cyperi (Beeson).

Cotoneaster microphylla

Scolytus abacensis Tsai & Yin, *Scolytus nitidus* Schedl.

Cotoneaster spp.

Scolytus mali (Beckstein), *Scolytus rugulosus* (Muller), *Thammurgus orientalis* Schedl.

Cotylelobium flavum

Genyocerus biporus (Schedl).

Cotynus cogyria

Chaetoptelcus vestitus (Mulsant & Rey).

Couma macrocarpa

Phelloterus anaxeus Wood, *Platypus nudatus* Wood, *Xyleborus adelographus* Eichhoff, *Xyleborus affinis* Eichhoff, *Xyleborus asper* Eggers, *Xyleborus ferrugineus* (Fabricius).

Coicania mexicana

Chaetophlocus heterodoxus (Casey).

Craibia ellioti

Rhopalopschion conjungens Schedl.

Crassocephalum spp.

Eucallacea xanthopus (Eichhoff).

Crataegus spp.

Cnesinus carinatus Wood, *Hypothenemus eruditus* (Panzer), *Hypothenemus eruditus* Westwood, *Scolytus mali* (Beckstein), *Scolytus rugulosus* (Muller), *Xyleborus dispar* (Fabricius).

Craterispermum spp.

Premnobius sexspinosus Eggers.

Cratoxylon arborescens

Crossotarsus squamulatus Chapuis, *Platypus westwoodi* Chapuis.

Cratylia floribunda

Premnobius cavipennis Eichhoff, *Xyleborinus reconditus* (Schedl), *Xyleborus spathipennis* Eichhoff, *Xylosandrus retusus* (Eichhoff).

Crossandra spp.

Doliopygus integratus Roberts.

Crossostylis biflora

Xyleborus ferrugineus (Fabricius).

Crotalaria anagyroides

Xylosandrus morigerus (Blandford).

Crotalaria saltiana

Hypothenemus eruditus Westwood.

Crotalaria spp.

Corthylus collaris Blandford, *Hypothenemus eruditus* Westwood, *Hypothenemus obscurus* (Fabricius), *Xyleborus ferrugineus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Crotalaria striata

Eucallacea fornicatus (Eichhoff).

Crotalaria usaramoensis

Xylosandrus morigerus (Blandford).

Crotalaria spp.

Chramesus marginatus Wood, *Taurodemus perebeae* (Ferrari), *Tricholus scitulus* Wood.

Croton argyратum

Crossotarsus wallacei (Thomson).

Croton gossypifolius

Cnesinus lecontei Blandford, *Microcorthylus concisus* Wood, *Phlocotribus setulosus* Eichhoff.

Croton guianensis

Phloeotribus nebulosus Wood, *Scolytodes resculus* Wood.

Croton haumanianus

Chaetastus tuberculatus (Chapuis), *Doliopygus contradii* (Strohmeyer), *Doliopygus rapax* (Sampson), *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Platypus impressus* (Strohmeyer), *Strombophorus vagans* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus gracilipennis* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky).

Croton macrostachys

Chaetastus montanus Schedl, *Chaetastus tuberculatus* (Chapuis), *Platypus spinulosus* (Strohmeyer).

Croton megalocarpus

Periommatatus grandis Schedl, *Platypus spinulosus* (Strohmeyer).

Croton mubungu

Doliopygus lefevrei Schedl.

Croton pseudoniveus

Chramesus exul Wood, *Hylocurus crotonis* Wood.

***Croton* spp.**

Ambrosiodmus eichhoffi (Schreiner), *Ambrosiodmus obliquus* (LeConte), *Amphicranus acus* Wood, *Chaetastus montanus* Schedl, *Chaetastus tuberculatus* (Chapuis), *Chramesus exul* Wood, *Cucinus equitai* Wood, *Cucinus lecontei* Blandford, *Corthylus abbreviatus* Eichhoff, *Doliopygus citri* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus unicus* (Schedl), *Hypothenemus eruditus* Westwood, *Monarthrum dimidiatum* (Ferrari), *Periommatatus grandis* Schedl, *Phloeotribus demessus* Blandford, *Pityophthorus crotonis* Wood, *Platypus cupulatus* Chapuis, *Platypus impressus* (Strohmeyer), *Platypus spinulosus* (Strohmeyer), *Scolytodes habilis* Wood, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasicusculus* Eggers, *Xylosandrus compactus* (Eichhoff).

Croton sylvaticus

Doliopygus erichsoni (Chapuis).

Cryptocarya ablato

Platypus australis Chapuis.

Cryptocarya chinensis

Crossotarsus rengetensis Niisima & Murayama.

Cryptocarya erythroxylon

Cryphalus asperulus Schedl, *Cryphalus brindleyi* Blandford.

***Cryptocarya* spp.**

Diapys elongatus Schedl, *Diapys nanodontus* Roberts, *Diapys nanus* Schedl, *Diapys nebulosus* Roberts, *Diapys perpygmaeus* Roberts, *Diapys pusillimus* Chapuis, *Diapys robustus* Schedl, *Hylesinus cordipennis* Lea, *Platypus echinatus* Roberts & Morimoto, *Platypus enormis* Schedl, *Platypus ramosissimus* Roberts & Morimoto, *Platypus ramulosus* Roberts & Morimoto, *Platypus sellaeformis* Roberts & Morimoto, *Platypus uniformis* Schedl, *Xyleborinus andrewesi* (Blandford), *Xylosandrus compactus* (Eichhoff).

Cryptocarya wrightiana

Crossotarsus bonvouloiri Chapuis, *Crossotarsus siva* Beeson, *Scolytoplatus cutomoides* Blandford.

***Cryptocery* spp.**

Platypus australis Chapuis.

Cryptolepis buchanani

Scolytogenes darwini Eichhoff.

Cryptomeria japonica

Crossotarsus externidentatus (Fairmaire), *Cryphalus cryptomeriae* Niisima, *Diapys spatulifer* Browne, *Eucallacea validus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Phloeosinus gifucensis* Murayama, *Phloeosinus leucisi* Chapuis, *Phloeosinus rudis* Blandford, *Phloeosinus sannohensis* Murayama, *Platypus kusukusensis* Murayama, *Platypus leucisi* Blandford, *Platypus quercivorus* Murayama, *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus aquilus* Blandford, *Xyleborus seriatus* Blandford, *Xylosandrus mutilatus* (Blandford).

***Cryptomeria* spp.**

Phloeosinus perlatus Chapuis.

Cryptophagus subvittatus

Xyleborinus saxeseni (Ratzeburg).

Ctenolaphon grandifolius

Platypus signatus Chapuis.

***Cucumis* spp.**

Eucallacea fornicatus (Eichhoff).

Cucurbita digitata

Dendrocranulus cucurbitae (LeConte).

Cucurbita foetidissima

Dendrocranulus cucurbitae (LeConte).

Cucurbita pepo

Dendrocranulus knausi (Hopkins).

***Cucurbita* spp.**

Dendrocranulus knausi (Hopkins), *Dendrocranulus limbellus* Wood, *Dendrocranulus modus* Wood.

Cudrania jarauensis

Eucallacea interjectus (Blandford).

Cullenia excelsa

Coccotrypes cardamomi Schaufuss, *Coccotrypes variabilis* (Beeson), *Platypus solidus* Walker.

Cunninghamia lanceolata

Ambrosiodmus rubricollis (Eichhoff), *Eucallacea validus* (Eichhoff), *Phloeosinus bicolor* (Brulle), *Phloeosinus perlatus* Chapuis, *Phloeosinus sinensis* Schedl.

Cunninghamia sinensis

Phloeosinus perlatus Chapuis, *Phloeosinus sinensis* Schedl.

Cupania guatemalensis

Bothrostermus foveatus (Blackman), *Hypothenemus trivialis* Wood, *Xylosandrus zimmermanni* (Hopkins).

Cupressus arizonica

Phloeosinus arizonicus Blackman, *Phloeosinus baumanni* Hopkins, *Phloeosinus cristatus* (LeConte), *Phloeosinus scopulorum neomexicanus* Blackman, *Phloeosinus spinosus* Blackman.

Cupressus benthami

Hypothenemus eruditus Westwood, *Phloeosinus baumanni* Hopkins, *Phloeosinus tacubayae* Hopkins.

Cupressus forbesi

Phloeosinus cristatus (LeConte), *Phloeosinus frontalis* Bruck.

Cupressus guadeloupeusis

Phloeosinus frontalis Bruck.

Cupressus lindleyi

Carphobius pilifer Wood, *Phloeosinus baumanni* Hopkins, *Phloeosinus tacubayae* Hopkins.

Cupressus lusitanica

Carphobius cupressi Wood, *Chaetastus montanus* Schedl, *Hypothenemus eruditus* Westwood, *Phloeosinus*

- baumanni* Hopkins, *Phloeosinus henschi* Reitter, *Phloeosinus tacubayae* Hopkins.
- Cupressus macnabiana**
Phloeosinus hoppingi Swaine.
- Cupressus macrocarpa**
Crossotarsus terminatus Chapuis, *Phloeosinus cristatus* (LeConte), *Phloeosinus cupressi* Hopkins, *Phloeosinus frontalis* Bruck, *Phloeosinus hoppingi* Swaine, *Phloeosinus sequoiae* Hopkins, *Platypus furcatus* Blandford.
- Cupressus sargentii**
Phloeosinus cupressi Hopkins, *Phloeosinus hoppingi* Swaine, *Phloeosinus setosus* Bruck, *Phloeosinus steinerci* Bruck, *Phloeosinus variolatus* Bruck.
- Cupressus sempercirens**
Phloeosinus armatus Reitter.
- Cupressus spp.**
Hypothenemus eruditus Westwood, *Phloeosinus antennatus* Swaine, *Phloeosinus armatus* Reitter, *Phloeosinus bicolor* (Brulle), *Phloeosinus cupressi* Hopkins, *Phloeosinus deleoni* Blackman, *Phloeosinus henschi* Reitter, *Phloeosinus thujae* (Perris).
- Cupressus torulosa**
Phloeosinus jubatus Sampson.
- Cussonia chartacea**
Eucallalcea xanthopus (Eichhoff).
- Cussonia longissima**
Xyleborus ferrugineus (Fabricius), *Xyleborus picus* (Motschulsky).
- Cussonia spp.**
Acanthotomicus octospinosus (Schedl).
- Cussonia rantsilana**
Eucallalcea xanthopus (Eichhoff), *Mitosoma obtusum* Schedl, *Mitosoma vulpinum* Schedl.
- Cyathea spp.**
Eucallalcea oparum (Beeson).
- Cybistax donnell-smithii**
Cnemidophorus splendens (Wood).
- Cydonia oblonga**
Scolytus rugulosus (Muller).
- Cylicodiscus gabunensis**
Doliopygus brevis (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus perbrevis* Schedl, *Platypus hintzi* Schaufuss, *Xyleborinus sharpei* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).
- Cylicodiscus spp.**
Premnobius caripennis Eichhoff.
- Cynodon dactylon**
Hypothenemus eruditus Westwood, *Hypothenemus pubescens* Hopkins.
- Cynometra alexandrii**
Ambrosiodinus aegir (Eggers), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus artespinitus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus fulgens* Schedl, *Doliopygus intritus* (Schedl), *Doliopygus megatoma* Schedl, *Doliopygus notatus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus morio* (Eggers), *Premnobius nodulosus* Hagedorn, *Rhopalosiphon subseriatus* (Schedl), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff.
- Cynometra hankai**
Chaetastus tuberculatus (Chapuis), *Coccotrypes grandis* (Eggers), *Cosmoderes donisi* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus exilis* (Chapuis), *Doliopygus incilis* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus unicomis* Schedl, *Hypothenemus cynometrae* Schedl, *Hypothenemus donisi* (Schedl), *Hypothenemus eruditus* Westwood, *Periommatius circularis* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).
- Cynometra hemitomophylla**
Araptus laevigatus (Eggers), *Coccotrypes cyperi* (Beeson).
- Cynometra insularis**
Platypus citicensis Roberts.
- Cynometra oaxacana**
Pseudochramesus jaliscoensis Wood.
- Cynometra spp.**
Ambrosiodinus tropicus (Hagedorn), *Doliopygus excavatus* (Sampson), *Doliopygus intritus* (Schedl), *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer).
- Cyrtandra spp.**
Xyleborus numfordi Beeson, *Xyleborus temctiniucus* Beeson.
- Cytisus alpinus**
Hylastinus fankhauseri Reitter.
- Cytisus biflorus**
Hylastinus obscurus (Marshall).
- Cytisus hirsutus**
Hylastinus obscurus (Marshall).
- Cytisus laburnum**
Hylastinus fankhauseri Reitter, *Hylastinus obscurus* (Marshall).
- Cytisus nigricans**
Hylastinus obscurus (Marshall).
- Cytisus proliferus**
Liparthrum nigrescens Wollaston.
- Cytisus spp.**
Liparthrum genistae (Aube), *Phloeotribus perfoliatus* (Wollaston), *Phloeotribus rhododactylus* (Marshall).
- Cytisus triflorus**
Hylastinus fankhauseri Reitter, *Phloeotribus pycnimerhoffi* (Eggers).
- Cytisus veldeni**
Phloeotribus hercegovinensis (Seitner).
- Dacrycarpus spp.**
Diapus nebulosus Roberts.
- Dacrydium elatum**
Platypus citicensis Roberts.
- Dacrydium spp.**
Xyleborus partitus Browne.
- Dacryodes deliciosa**
Cylindropalpus auricomans (Schaufuss).
- Dacryodes edulis**
Xyleborus sub tuberculatus Eggers.
- Dacryodes excelsa**
Monarthrum praecustum (Eggers), *Platypus compositus* Say.
- Dacryodes klaineana**
Xyleborus alluaudi Schaufuss.
- Dacryodes pubescens**
Cyclorhipidion callosum (Schedl), *Cyclorhipidion*

nsafukalae (Schedl), *Cyclorhipidion praecursor* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus sub tuberculatus* Eggers.

Dacryodes rostrata

Crossotarsus brownei Schedl, *Cyclorhipidion pruinosum* (Blandford), *Hypocryphalus malayensis* Schedl.

Dacryodes spp.

Ambrosiodinus lecontei Hopkins, *Cyclorhipidion pruinosum* (Blandford).

Dacryodes yangambiensis

Doliopygus fulgens Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus opulentus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Trachyostus aterrimus* (Schaufuss), *Xyleborus affinis* Eichhoff.

Daemonorops spp.

Ambrosiodinus brunneipes (Eggers), *Eucallacea limatus* (Schedl), *Leptoxyleborus concisus* (Blandford).

Dalbergia assamensis

Platypus secretus Sampson.

Dalbergia assamica

Eucallacea sibsagaricus (Eggers), *Platypus cavus* Strohmeyer, *Platypus indicus* Strohmeyer, *Xyleborus piccus* (Motschulsky).

Dalbergia ecastophyllum

Hypothenemus brunneus (Hopkins), *Hypothenemus crudiae* (Panzer).

Dalbergia gastrophyllyum

Hypothenemus birmanus (Eichhoff).

Dalbergia hupeana

Eucallacea validus (Eichhoff).

Dalbergia latifolia

Carchesiopygus multidentatus (Strohmeyer), *Crossotarsus fractus* Sampson, *Crossotarsus javanus* Beeson, *Crossotarsus minax* (Walker), *Ernoporus concentralis* (Eggers), *Platypus jansonii* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus solidus* Walker, *Platypus vethi* Strohmeyer, *Xyleborus haberkorni* Eggers, *Xyleborus perforans* (Wollaston), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus manicus* (Blandford), *Xylosandrus morigerus* (Blandford).

Dalbergia paniculata

Xyleborus perforans (Wollaston).

Dalbergia pterocarpifolia

Mitosoma lobatum Schedl, *Platypus madagascariensis* Chapuis, *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborinus signatipennis* (Schedl), *Xyleborus antaisaka* Schedl, *Xyleborus gibber* Schedl, *Xyleborus sakalava* Schedl.

Dalbergia sissoo

Coccotrypes nubilis (Blandford), *Platypus solidus* Walker, *Xylosandrus crassiusculus* (Motschulsky).

Dalbergia spp.

Arixyleborus minor (Eggers), *Arixyleborus tuberculatus* (Eggers), *Dendrocrandus brasiliensis* (Schedl), *Eucallacea andamanensis* (Blandford), *Mitosoma obconiceps* Schedl, *Mitosoma obliquatum* Schedl, *Mitosoma obtusum* Schedl, *Xyleborus madagascariensis* Schaufuss.

Dalea lasiostachya

Chramesus marginatus Wood.

Dalhousiea spp.

Diamerus impar Chapuis.

Dalium corbisieri

Eucallacea granosus (Schedl).

Daniellia klainei

Cylindropalpus laudatus (Schedl), *Diapus quinque-spinatus* Chapuis, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius).

Daniellia oliveri

Ambrosiodinus eichhoffi (Schreiner), *Doliopygus aduncus* Schedl, *Doliopygus terebrans* Schedl, *Pityophthorus barbifer* Schedl, *Triozastus marshalli* (Sampson).

Daphniphyllum glaucescens

Hypothenemus amakusanus (Murayama), *Platypus calamus* Blandford.

Daphniphyllum macropodium

Xyleborus praecus Blandford, *Xyleborus volutus* (Fabricius).

Daphniphyllum teijsmanni

Platypus calamus Blandford.

Daphnopsis seibertii

Araptus laevigatus (Eggers), *Hypothenemus plumeriae* (Nordlinger).

Dehaasia euneata

Xylosandrus ater (Eggers).

Deimbollia spp.

Chaetastus tuberculatus (Chapuis), *Triozastus caliginosus* Roberts, *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus albaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky).

Delphinium consolida

Thamnurgus delphinii (Rosenhauer).

Delphinium spp.

Thamnurgus delphinii (Rosenhauer).

Dendrobium phalaenopsis

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Dendrobium spp.

Xyleborus perforans (Wollaston), *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Dendrobium urvelli

Xylosandrus morigerus (Blandford).

Dendrobium veratrifolium

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Dendropanax arboreum

Xyleborus affinis Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius), *Xylosandrus laticeps* (Wood).

Derris elliptica

Hypothenemus cruditis Westwood.

Derris microphylla

Xylosandrus morigerus (Blandford).

Derris robusta

Eucallacea fornicatus (Eichhoff), *Hypothenemus cruditis* Westwood.

Derris spp.

Hypothenemus crudiae (Panzer).

Desmodium cephalotes

Eucallacea fornicatus (Eichhoff).

Desmodium cogantii

Chramesus marginatus Wood.

Desmodium flemingia spp.

Xylosandrus morigerus (Blandford).

Desmodium ovalifolium

Xylosandrus compactus (Eichhoff).

Desmoncus spp.

Xyleborus inaequalis Schedl.

Desplatsia deweveri

Doliopygus coelecephalus (Schaufuss), *Doliopygus erichsoni* (Chapuis), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus rapax* (Sampson), *Periommatius grandis* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schontedeni* (Schedl).

Desplatsia subericarpa

Periommatius longicollis Chapuis.

Dialanthera otoba

Xyleborus grossmanni Schedl.

Dialium corbisieri

Ambrosiodinus oratus (Eggers), *Cosmoderes solitarius* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus exilis* (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus minimus* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus woltschei* Schedl, *Hypothenemus socialis* (Schedl), *Mesoplatypus ventricornis* Schedl, *Mesoplatypus venustus* Schedl, *Polygraphus afzelschiae* Schedl, *Strombophorus dialiumi* Schedl, *Strombophorus lukengaee* Schedl, *Triozastus marshalli* (Sampson), *Xyleborinus gracilipennis* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus piceus* (Motschulsky), *Xyleborus voluculus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Dialium excelsum

Cylindropalpus laudatus (Schedl), *Doliopygus coelecephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus lebruni* Schedl, *Doliopygus spectabilis* Schedl, *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatius grandis* Schedl, *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson).

Dialium pachyphyllum

Doliopygus conradti (Strohmeyer), *Doliopygus exilis* (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus tenuis* (Strohmeyer), *Hypothenemus biseriatus* (Eggers), *Platypus hintzi* Schaufuss, *Strombophorus lukengaee* Schedl, *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ficus* Eggers, *Xyleborus multi-spinatus* Eggers, *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Dialium spp.

Acanthotomicus unicus (Schedl), *Doliopygus angolensis* Schedl, *Doliopygus bilobatus* (Schedl), *Platypus canaliculatus* Schedl, *Platypus puerulus* (Schedl), *Strombophorus lukengaee* Schedl, *Xyleborus fallax* Eichhoff.

Dialium tessmanui

Doliopygus medius Schedl, *Periommatius longicollis* Chapuis, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius).

Dialium spp.

Hypothenemus camerinus (Eggers).

Dichaetanthera cordifolia

Pityophthorus dorsalis.

Dichapetalum spp.

Xyleborus truncatulus Schedl.

Dichopetalum thomieri

Hypothenemus aethiops (Schedl).

Dichopsis elliptica

Eucallacea andamanensis (Blandford).

Dichopsis polyantha

Scolytomimus assamensis Schedl.

Dictyosperma album

Coccotrypes dactyliperda (Fabricius).

Didymeles spp.

Mitasoma sexspinosum Schedl.

Dillenia pentagyna

Eucallacea andamanensis (Blandford), *Xyleborinus andrevesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xylosandrus butamali* (Beeson), *Xylosandrus crassiusculus* (Motschulsky).

Dillenia philippinensis

Xyleborus similis Ferrari.

Dillenia retusa

Coccotrypes advena Blandford, *Coccotrypes variabilis* (Beeson).

Dioclea megacarpa

Hypothenemus erudiae (Panzer), *Hypothenemus plumieriae* (Nordlinger), *Stegomernus pygmaeus* Wood, *Theoborus ricini* (Eggers).

Diospyros alboflavescens

Platypus impressus (Strohmeyer), *Platypus intermedius* (Schedl), *Trachyostus schaufussi* (Strohmeyer).

Diospyros crassiflora

Chaetastus persimilis (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus laudatus* (Schedl), *Platypus impressus* (Strohmeyer), *Trachyostus aterrimus* (Schaufuss), *Xyleborus affinis* Eichhoff.

Diospyros eriantha

Xyleborus atratus Eichhoff.

Diospyros kaki

Ambrosiodinus rubricollis (Eichhoff), *Cryphalus exiguus* Blandford, *Hypothenemus eruditus* Westwood, *Xyleborinus attenuatus* (Blandford), *Xyleborus kadoyamaensis* Murayama, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus brevis* (Eichhoff), *Xylosandrus germanus* (Blandford).

Diospyros morrisiana

Xyleborus atratus Eichhoff.

Diospyros oocarpa

Crossotarsus andamanus Beeson, *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston).

Diospyros paniculata

Platypus latifinis Walker, *Platypus uncinatus* Blandford, *Xyleborus perforans* (Wollaston).

Diospyros pseudo-ebenum

Platypus cavius Strohmeyer, *Platypus hybridus* Schedl.

Diospyros pyrrhocarpa

Platypus cordiger Chapuis, *Xyleborus piceus* (Motschulsky).

Diospyros spp.

Ambrosiodinus eichhoffi (Schreiner), *Ambrosiodinus rubricollis* (Eichhoff), *Arxyleborus medius* (Eggers), *Coccotrypes carpophagus* (Homung), *Crossotarsus squamulatus* Chapuis, *Doliopygus bitalei* Schedl, *Doliopygus convexus* Roberts, *Doliopygus dolosus* Schedl, *Mesoplatypus calabaricus* Roberts, *Platypus compositus* Say, *Platypus hirtus* Schedl, *Platypus hybridus* Schedl, *Platypus insularis* Strohmeyer, *Platypus secretus* Sampson, *Platypus volaticus* Schedl,

Treptoplatypus circulicauda Browne, *Xylosandrus compactus* (Eichhoff).

Diospyros suaveolens

Chaetastus tuberculatus (Chapuis).

Dipholis salicifolia

Cnesinus strigicollis LeConte, *Cryptocarenum seriatus* Eggers, *Hypothenemus seriatus* (Eichhoff), *Hypothenemus squamosus* (Hopkins), *Micracisella nanula* (LeConte).

Diphysa robinioidea

Ambrosiodinus lecontei Hopkins, *Hypothenemus erudiae* (Panzer).

Diplodiscus paniculatus

Eucallacea quadraticollis (Eggers), *Platypus setaceus* Chapuis.

Diploglottis australis

Xylosandrus solidus (Eichhoff).

Diplospora spp.

Crossotarsus schedli Browne.

Diprosora viridiflora

Xyleborus atratus Eichhoff.

Dipterocarpus alatus

Cryphalus dipterocarpi Wood.

Dipterocarpus baudii

Arixyleborus suturalis (Eggers), *Coptodryas diversicolor* (Eggers), *Coptodryas nigax* (Schedl), *Coptodryas perparva* (Sampson), *Eccoptopterus limbatus* Sampson, *Eccoptopterus spinosus* (Olivier), *Leptoxyleborus concisus* (Blandford), *Platypus curtus* Chapuis, *Webbia dipterocarpi* Hopkins, *Xyleborus apiculatus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus jacanus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus pseudopilifer* Schedl, *Xyleborus sichus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Dipterocarpus borneensis

Crossotarsus terminatus Chapuis, *Genyocerus serratus* (Schedl), *Platypus dignus* Schedl, *Platypus platypoides* (Browne), *Webbia bakoensis* Browne, *Webbia quattuordecimspinus* Sampson, *Webbia suturalis* Browne.

Dipterocarpus cornutus

Crossotarsus terminatus Chapuis, *Diapus quinquespinatus* Chapuis, *Genyocerus diaphanus* (Schedl), *Xyleborus inprocerus* Sampson, *Xylosandrus mediocris* (Schedl).

Dipterocarpus erinitus

Coccotrypes gedeanus (Eggers), *Platypus curtus* Chapuis, *Xyleborus dolosus* Blandford.

Dipterocarpus curmuk

Xylosandrus mediocris (Schedl).

Dipterocarpus grandiflorus

Arixyleborus rigosipes Hopkins, *Coptodryas confusa* Hopkins, *Cyclorhizipidion dipterocarpi* Hopkins, *Dryocoetius laevis* (Strohmeyer), *Hypocryphalus rotundus* Hopkins, *Platypus bifurcus* (Schedl), *Platypus spectabilis* Schedl, *Webbia dipterocarpi* Hopkins.

Dipterocarpus hasseltii

Diapus quinquespinatus Chapuis.

Dipterocarpus indiens

Platypus cordiger Chapuis, *Platypus diversipennis* Schedl, *Platypus obtusipennis* Schedl, *Xylosandrus butanali* (Beeson).

Dipterocarpus kerrii

Coccotrypes nitidus (Eggers).

Dipterocarpus kuastleri

Cyclorhizipidion agnatum (Eggers), *Platypus pseudo-*

curtus Schedl, *Leptoxyleborus concisus* (Blandford), *Xyleborinus perminutissimus* (Schedl), *Xylosandrus difficilis* (Eggers).

Dipterocarpus obtusifolius

Ambrosiodinus minor (Stebbing), *Platypus biflexuosus* Schedl, *Webbia bifornis* Browne, *Webbia cornutus* Schedl.

Dipterocarpus pilosus

Anasa resceans (Eggers), *Arixyleborus mediosectus* (Eggers), *Carchesiopygus impariporus* (Beeson), *Coccotrypes barbatus* (Schedl), *Coccotrypes monoceros* (Beeson), *Coccotrypes papuanus* (Eggers), *Cryphalus dipterocarpi* Wood, *Diapus pendleburyi* Schedl, *Diapus quinquespinatus* Chapuis, *Genyocerus biporus* Schedl, *Genyocerus furtivus* (Sampson), *Platypus biflexuosus* Schedl, *Platypus curtus* Strohmeyer, *Platypus cupulatus* Chapuis, *Platypus indicus* Strohmeyer, *Platypus secretus* Sampson, *Platypus solidus* Walker, *Platypus subdepressus* (Schedl), *Platypus uncinatus* Blandford, *Xyleborus shoreae* (Stebbing).

Dipterocarpus polosapis

Hypothenemus dipterocarpi Hopkins.

Dipterocarpus serrata

Diapus quinquespinatus Chapuis.

Dipterocarpus spp.

Arixyleborus dipterocarpi Browne, *Arixyleborus medius* (Eggers), *Cnestus suturalis* (Eggers), *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Genyocerus albipennis* Motschulsky, *Leptoxyleborus concisus* (Blandford), *Platypus bifurcus* (Schedl), *Platypus curtus* Chapuis, *Platypus dipterocarpi* Browne, *Platypus insulindicus* Schedl, *Platypus ovatus* Strohmeyer, *Platypus pseudocurtus* Schedl, *Platypus spinosus* Browne, *Schedlia sumatrana* (Hagedorn), *Scolytoplatypus parvus* Sampson, *Treptoplatypus circulicauda* Browne, *Webbia trigintispinus* Sampson, *Xyleborus amphcranoides* Hagedorn, *Xyleborus cavuloides* Browne, *Xyleborus fallax* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Dipterocarpus trimeris

Coccotrypes cyperi (Beeson).

Dipterocarpus turbinatus

Anasa resceans (Eggers), *Arixyleborus mediosectus* (Eggers), *Crossotarsus andamanus* Beeson, *Eccoptopterus spinosus* (Olivier), *Genyocerus dipterocarpi* Browne, *Platypus cordiger* Chapuis, *Sphacrotrypes costatus* Wood, *Xyleborinus andrewesi* (Blandford), *Xyleborus bidentatus* (Motschulsky), *Xyleborus cognatus* Blandford, *Xyleborus najor* (Stebbing), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Dipterocarpus vernicifluus

Coccotrypes incognitus (Schedl).

Dipterocarpus zeylanicus

Arixyleborus granulifer (Eggers), *Arixyleborus mediosectus* (Eggers), *Coccotrypes longior* (Eggers), *Hypothenemus javanus* (Eggers), *Platypus shoreanus* (Beeson), *Webbia ceylonae* Schedl, *Xyleborinus exiguus* (Walker).

Discoglypremma caloneura

Platypus hintzi Schanfuss, *Platypus orientalis* (Strohmeyer), *Prennobilus nodulosus* Hagedorn, *Xyleborus affinis* Eichhoff, *Xyleborus alhuandi* Schanfuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky).

Disholcopsis spp.

Araptus nanulus Wood.

Displodiscus paniculatus*Platypus caliginosus* Schedl.***Dissotis grandiflora****Hypothenemus grandis* Schedl.***Distemonanthus benthamianus****Cylindropalpus auricomans* (Schaufuss), *Doliopygus bilobatus* (Schedl), *Doliopygus dubius* (Sampson), *Platypus hintzi* Schaufuss, *Rhopalopselion thompsoni* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).***Distemonanthus* spp.***Premnobius caricipennis* Eichhoff.***Distylium racemosum****Platypus calanus* Blandford, *Sphaerotrypes pila* Blandford.***Ditoma crenata****Xyleborinus saxseni* (Ratzeburg).***Dolichandrone stipulata****Platypus cupulatus* Chapuis, *Platypus solidus* Walker, *Xyleborinus birmanus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Dolichos lablab****Hypothenemus dolichocolora* Hopkins.***Dombeya goetzenii****Doliopygus lefevrei* Schedl, *Platypus collinus* Browne.***Dombeya mastersii****Doliopygus bidentatus* (Strohmeyer), *Rhopalopselion innaturus* (Schedl).***Dombeya* spp.***Doliopygus nairobiensis* (Schedl), *Eucallacca xanthopus* (Eichhoff), *Platypus orientalis* (Strohmeyer), *Xyleborinus spiculatus* (Schaufuss), *Xyleborinus spinosus* (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xyleborus sakalava* Schedl.***Doonia cordifolia****Sphaerotrypes cristatus* Wood.***Doonia zeylanica****Arixyleborus medius* (Eggers), *Crossotarsus terminatus* Chapuis, *Platypus latifrons* Walker, *Platypus shoreanus* (Beeson), *Platypus ucinatus* Blandford, *Xylosandrus crassiusculus* (Motschulsky).***Doryphora aromatica****Platypus australis* Chapuis.***Doryalis* spp.***Hypothenemus grandis* Schedl.***Dracaena aurea****Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston).***Dracondomerum* spp.***Arixyleborus cariniceps* Schedl.***Dracontomelon* spp.***Cnestus rotundatus* Schedl, *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis.***Dracaena drace****Dactylotrypes longicollis* (Wollaston).***Dracaena* spp.***Mitosoma excision* Schaufuss.***Dracontomelium mangiferum****Platypus micrurus* Schedl.***Dracontomelium* spp.***Crossotarsus mniszewski* Chapuis, *Platypus solidus* Walker.***Drimycarpus ramosus****Platypus lepidus* Chapuis, *Platypus solidus* Walker, *Sphaerotrypes quadrituberculatus* Sampson.***Drimys granatensis****Scolytodes pumilus* Wood.***Dryadodaphne* spp.***Hypocryphalus montanus* Schedl.***Dryobalanops aromatica****Baiocis peruanulus* (Schedl), *Coccotrypes gedeanus* (Eggers), *Cryptoxyleborus dryobalanopsis* Schedl, *Cyclorhipidion prunosum* (Blandford), *Cyclorhipidion sistriophorum* (Hagedorn), *Diapus pendleburyi* Schedl, *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Dryococtius hirsutus* (Schedl), *Hypothenemus cuneolus* (Schedl), *Hypothenemus javanus* (Eggers), *Platypus bifurcus* (Schedl), *Platypus binucus* Blandford, *Platypus curtus* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus pseudocurtus* Schedl, *Platypus transformis* Schedl, *Platypus retulus* Schedl, *Platytarsulus declivis* Schedl, *Platytarsulus elongatus* Schedl, *Platytarsulus marshalli* Schedl, *Webbia quatuordecimspinus* Sampson, *Xyleborus amplexicauda* Hagedorn, *Xyleborus perforans* (Wollaston), *Xyleborus pseudopilifer* Schedl, *Xyleborus similis* Ferrari, *Xyleborus tunggali* Schedl, *Xylosandrus nancus* (Blandford).***Dryobalanops camphora****Platypus curtus* Chapuis.***Dryobalanops lanceolata****Platypus lavasensis* Browne.***Dryobalanops oblongifolia****Amasa schlichii* (Stebbing), *Arixyleborus leprosus* Schedl, *Arixyleborus malayensis* (Schedl), *Arixyleborus minor* (Eggers), *Arixyleborus scabripennis* (Blandford), *Arixyleborus suturalis* (Eggers), *Coptodryas nugax* (Schedl), *Crossotarsus fractus* Sampson, *Crossotarsus schedli* Browne, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Cyclorhipidion agnatum* (Eggers), *Genyocerus philippinensis* (Schedl), *Genyocerus serratus* (Schedl), *Hadrodemus globus* (Blandford), *Platypus bifidus* Schedl, *Platypus binucus* Blandford, *Platypus cupulatus* Chapuis, *Platypus insulindicus* Schedl, *Platypus platypoides* (Browne), *Platypus pseudocupulatus* Schedl, *Platypus solidus* Walker, *Schedlia sumatrana* (Hagedorn), *Scolyto-platypus pusillus* Eggers, *Webbia simplex* (Browne), *Xyleborus approximatus* Schedl, *Xyleborus javanus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus siclus* Schedl, *Xylosandrus ater* (Eggers), *Xylosandrus nancus* (Blandford), *Xylosandrus morigerus* (Blandford).***Dryobalanops* spp.***Scolyto-platypus parvus* Sampson.***Dryodaphnae* spp.***Diapus elongatus* Schedl, *Diapus robustus* Schedl, *Diapus turgidus* Roberts.***Drypetes gossweileri****Ambrosiodinus ovatus* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus coclocephalus* (Schaufuss), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatatus excisus* Strohmeyer, *Periommatatus grandis* Schedl, *Periommatatus longicollis* Chapuis, *Periommatatus*

substriatus Strohmeier. *Platypus hintzi* Schauffuss. *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeier), *Platypus solutus* Schedl, *Trachyostus aterrimus* (Schauffuss), *Trachyostus schaufussi* (Strohmeier), *Trioastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky), *Xyleborus volutus* (Fabricius).

Drypetes ituriensis

Doliopygus lebrunii Schedl, *Doliopygus medius* Schedl, *Doliopygus occallescent* Schedl, *Platypus intermedius* (Schedl), *Platypus solutus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky).

Drypetes leoneis

Coptoborus usagarius (Eggers), *Cylindropalpus granulatus* (Schedl), *Cylindropalpus interpositus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeier), *Doliopygus dolosus* Schedl, *Doliopygus erichsoni* (Chapuis), *Doliopygus jurioni* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Perionnatus longicollis* Chapuis, *Platypus impressus* (Strohmeier), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeier), *Platypus spinulosus* (Strohmeier), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus elongatus* Eggers, *Xyleborus multispinatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Drypetes louisii

Doliopygus erichsoni (Chapuis), *Doliopygus rapax* (Sampson), *Platypus impressus* (Strohmeier), *Platypus intermedius* (Schedl), *Trachyostus aterrimus* (Schauffuss).

Drypetes morocarpa

Doliopygus conradti (Strohmeier), *Doliopygus dubius* (Sampson), *Doliopygus interjectus* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Perionnatus grandis* Schedl, *Perionnatus longicollis* Chapuis, *Platypus impressus* (Strohmeier), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeier), *Xyleborus affinis* Eichhoff.

Drypetes sepiaria

Hylesinus nilgirinus Eggers.

Drypetes spinoso-dentata

Eucallacea xanthopus (Eichhoff), *Platypus orientalis* (Strohmeier).

***Drypetes* spp.**

Dactylipalpus grouvellei (Blandford), *Doliopygus lebrunii* Schedl, *Platypus conjunctus* Schedl, *Platypus intermedius* (Schedl), *Platypus spinulosus* (Strohmeier), *Prennobius cavipennis* Eichhoff, *Xyleborus piccus* (Motschulsky), *Xylosandrus compactus* (Eichhoff).

Drypetes staudtii

Chaetastus tuberculatus (Chapuis).

Drypetes yambataense

Xyleborus multispinatus Eggers.

Duabanga sommeratioides

Eucallacea bicolor (Blandford), *Platypus solidus* Walker.

Dubautia laxa

Xyleborus hawaiiensis Perkins.

***Duboisia* spp.**

Hypothenemus eruditus Westwood.

Duboscia viridiflora

Hypothenemus colae (Schedl), *Hypothenemus eruditus* Westwood, *Hypothenemus socialis* (Schedl).

***Dumoria* spp.**

Prennobius cavipennis Eichhoff.

Durio carinatus

Platypus cupulatus Chapuis, *Platypus curtus* Chapuis.

Durio griffithii

Crossotarsus squamulatus Chapuis.

***Durio* spp.**

Diapus quinquespinatus Chapuis, *Platypus cupulatus* Chapuis, *Platypus duriororus* Browne, *Platypus pseudocupulatus* Schedl, *Xyleborus perforans* (Wollaston).

Durio zibethinus

Diapus quinquespinatus Chapuis, *Platypus insulindicus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus perforans* (Wollaston), *Xyleborus piccus* (Motschulsky), *Xyleborus similis* Ferrari.

Dyera costulata

Carchesiopygus wollastoni (Chapuis), *Diapus pusillinus* Chapuis, *Diapus quinquespinatus* Chapuis, *Hypothenemus javanus* (Eggers), *Platypus cupulatus* Chapuis, *Platypus dyerae* Browne, *Platypus vethi* Strohmeier, *Platypus westwoodi* Chapuis, *Xyleborus apertus* Schedl, *Xyleborus mucronatus* Eggers, *Xyleborus subemarginatus* Eggers.

Dysoxylum amoeroides

Hypothenemus eruditus Westwood, *Xyleborinus andrewesi* (Blandford).

Dysoxylum binectariferum

Crossotarsus boncoulouiri Chapuis, *Diapus quinquespinatus* Chapuis.

Dysoxylum decandrum

Xyleborus perforans (Wollaston), *Xyleborus piccus* (Motschulsky), *Xyleborus volutus* (Fabricius).

Dysoxylum muelleri

Xyleborus perforans (Wollaston).

Dysoxylum pettigrewianum

Xyleborus perforans (Wollaston).

***Dysoxylum* spp.**

Cryphalus vitensis Browne, *Cryphalus waplery* Eichhoff, *Diapus oomsis* Roberts, *Genyocerus puer* (Schedl), *Hadrodemius globus* (Blandford), *Platypus alternantecostatus* (Schedl), *Platypus cavus* Strohmeier, *Platypus cordiger* Chapuis, *Platypus jansoni* Chapuis, *Platypus papuanus* Roberts, *Platypus parapetax* Roberts & Morimoto, *Platypus sulcinodis* Schedl, *Platypus vitensis* Roberts, *Platypus volaticus* Schedl, *Xyleborinus andrewesi* (Blandford).

Echinocarpus dasycarpus

Diapus spatulifer Browne, *Platypus falcatus* Strohmeier, *Xyleborus similis* Ferrari.

Echinocystis macrocarpa

Dendrocraniulus cucurbitae (LeConte).

***Echinopanax* spp.**

Dryocoetes padi Stark.

***Ehlersii* spp.**

Hypothenemus eruditus Westwood.

Ehretia acuminata

Eucallacea sibsagarius (Eggers), *Platypus cavus* Strohmeier, *Platypus luniger* Motschulsky, *Platypus subdepressus* (Schedl), *Xyleborus shoreae* (Stebbing), *Xyleborus similis* Ferrari.

Ekebergia capensis

Hypocryphalus dubiosus Schedl.

Ekebergia ruelpelliana

Chaetastus montanus Schedl, *Doliopygus nairobiensis* (Schedl), *Doliopygus serratus* (Strohmeyer).

Ekebergia spp.

Doliopygus serratus (Strohmeyer), *Platypus madagascariensis* Chapuis, *Scolytoplatypus eichelbaumi* Hagedorn, *Scolytoplatypus kienensis* Schedl, *Triozastus marshalli* (Sampson), *Xyleborinus forficuloides* (Schedl), *Xyleborinus signatipennis* (Schedl).

Ekebergia suavis

Mitosoma sexspinosum Schedl, *Platypus madagascariensis* Chapuis, *Xyleborus alluaudi* Schaufuss.

Elaeagnus pungens

Hypothenemus seriatus (Eichhoff).

Elaeagnus pungens fruitlandi

Hypothenemus eruditus Westwood, *Hypothenemus javanus* (Eggers).

Elaeagnus spp.

Pseudoxylechinus variegatus Wood & Huang, *Scolytus jaroschewskii* Schevyrew, *Scolytus schetyrewi* Semenov, *Trypophloeus klimeschii* Eggers.

Elaeis guineensis

Coccotrypes carpophagus (Hornung), *Coccotrypes congonus* Eggers, *Coccotrypes dactyliperda* (Fabricius), *Coccotrypes nigripes* Eggers, *Diapus quinquespinatus* Chapuis, *Doliopygus conradti* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Hypothenemus perhispidus* (Eggers), *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xylosandrus compactus* (Eichhoff).

Elaeocarpus bifidus

Crossotarsus externedentatus (Fairmaire), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston).

Elaeocarpus decipiens

Sucus niisimai (Eggers).

Elaeocarpus ferrugineus

Anasa versicolor (Sampson), *Leptoxyleborus depressus* (Eggers).

Elaeocarpus ganitrus

Platypus biuncus Blandford, *Platypus cupulatus* Chapuis.

Elaeocarpus grandis

Xyleborus perforans (Wollaston).

Elaeocarpus jackianus

Leptoxyleborus depressus (Eggers).

Elaeocarpus lanceaefolius

Platypus falcatus Strohmeyer.

Elaeocarpus oblongus

Coccotrypes cyperi (Beeson).

Elaeocarpus petiolatus

Anasa versicolor (Sampson), *Ambrosiodmus asperatus* (Blandford), *Arixyleborus suturalis* (Eggers), *Cnestus aterrimus* (Eggers), *Crossotarsus terminatus* Chapuis.

Elaeocarpus sericeus

Ambrosiodmus mahafali (Schedl), *Xylosandrus crassiusculus* (Motschulsky).

Elaeocarpus serratus

Platypus furcatus Blandford.

Elaeocarpus spp.

Anasa versicolor (Sampson), *Ambrosiodmus wilderi* (Beeson), *Coptodryas perparva* (Sampson), *Coptodryas quadricostata* (Schedl), *Cryphalus borneensis* Schedl, *Diapus bilunatus* Schedl, *Diapus elongatus* Schedl, *Diapus nanus* Schedl, *Diapus nebulosus* Roberts, *Diapus oomsis* Roberts, *Diapus unispineus* Roberts,

Mitosoma excisum Schaufuss, *Mitosoma paulinum* Schedl, *Mitosoma sexspinosum* Schedl, *Platypus semi-granosus* (Sampson), *Xyleborinus forficuloides* (Schedl), *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborinus signatipennis* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus antaisaka* Schedl, *Xyleborus insulindicus* Eggers, *Xyleborus madagascariensis* Schaufuss, *Xyleborus minutus* Blandford, *Xyleborus similis* Ferrari, *Xyleborus swezeyi* Beeson, *Xylosandrus crassiusculus* (Motschulsky).

Elaeocarpus tuberculatus

Coccotrypes cardamomi Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).

Elaeodendron spp.

Hypocryphalus dubiosus Schedl.

Elaeophorbium drupifera

Cyrtogenius cribricollis Schedl.

Elateriospermum spp.

Coptodryas nudibrevis (Schedl).

Elateriospermum tapos

Coptodryas perparva (Sampson), *Crossotarsus brownei* Schedl, *Crossotarsus schedli* Browne, *Crossotarsus wallacei* (Thomson), *Cyrtogenius subacuminatus* (Schedl), *Diapus quinquespinatus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Platypus insularis* Strohmeyer, *Platypus lepidus* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus puerulus* (Schedl), *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston).

Ellettaria major

Coccotrypes cardamomi Schaufuss.

Elongatus spp.

Hypothenemus eruditus Westwood.

Embelia schimperi

Xylocleptes adeniae (Schedl).

Enantia chlorantha

Chaetastus tuberculatus (Chapuis), *Periommatius nitidicollis* Strohmeyer.

Enantia polycarpa

Xyleborus ambasiusculus Eggers.

Enantia spp.

Premnobius cavipennis Eichhoff.

Encelia californica

Chaetophilocus maclayi (Bruck).

Encelia farinosa

Chaetophilocus pruinosus (Blackman).

Endiandra palmerstonii

Ambrosiodmus compressus (Lea), *Coptodryas eucalyptica* (Schedl), *Xyleborus insulindicus* Eggers, *Xyleborus perforans* (Wollaston).

Endiandra spp.

Crossotarsus terminatus Chapuis, *Platypus loriae* (Schedl).

Endospermum macrophyllum

Crossotarsus externedentatus (Fairmaire), *Diapus quinquespinatus* Chapuis, *Platypus citiensis* Roberts.

Endospermum malaccense

Coptodryas diversicolor (Eggers), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus schedli* Browne, *Crossotarsus squamulatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Diapus quinquespinatus* Chapuis, *Leptoxyleborus concisus* (Blandford), *Platypus bifurcus* (Schedl), *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Platypus insularis* Strohmeyer, *Platypus lepidus* Chapuis, *Platypus parallelus* (Fabricius), *Platypus pseudocupulatus* Schedl, *Platypus signatus* Chapuis, *Platypus westwoodi* Chapuis, *Xyleborus perforans*

- (Wollaston). *Xyleborus piceus* (Motschulsky), *Xylosandrus morigerus* (Blandford).
- Endospermum medullosum**
Platypus jansoni Chapuis.
- Endospermum spp.**
Diaprus papuanus Schedl, *Xyleborus perforans* (Wollaston).
- Endriandra rubescens**
Platypus fraterculus Schedl.
- Engelhardtia spicata**
Crossotarsus bonvouloiri Chapuis, *Crossotarsus siva* Beeson, *Crossotarsus ursus* Schedl, *Scolytotplatypus parvus* Sampson, *Scolytotplatypus pubescens* Hagedorn, *Scolytotplatypus raja* Blandford.
- Entada gigas**
Araptus laevigatus (Eggers).
- Entadopsis abyssinica**
Doliopygus erichsoni (Chapuis), *Hypothenemus eruditus* Westwood, *Xyleborus affinis* Eichhoff.
- Entandrophragma angolense**
Chaetastus tuberculatus (Chapuis), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus serratus* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus parallelus* (Fabricius), *Polygraphus bicolor* Eggers, *Prennobiobis sexspinosus* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).
- Entandrophragma candollei**
Chaetastus tuberculatus (Chapuis), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Platypus schenklingi* (Strohmeyer), *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius).
- Entandrophragma cylindricum**
Chaetastus tuberculatus (Chapuis), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus nairobiensis* (Schedl), *Platypus hintzi* Schaufuss, *Platypus schenklingi* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus rotulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).
- Entandrophragma excelsum**
Chaetastus montanus Schedl, *Chaetastus tuberculatus* (Chapuis), *Doliopygus bidentatus* (Strohmeyer), *Periommatius grandis* Schedl, *Prennobiobis mukungu* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasicornis* Schedl.
- Entandrophragma spp.**
Ambrosiodinus acir (Eggers), *Ambrosiodinus obliquus* (LeConte), *Chaetastus tuberculatus* (Chapuis), *Doliopygus artespinnatus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus ghesquieri* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus*

serratus (Strohmeyer), *Doliopygus staneri* Schedl, *Doliopygus spectabilis* Schedl, *Doliopygus tenuis* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Polygraphus granulatus* Eggers, *Polygraphus granulifer* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus ferrugineus* (Fabricius).

Entandrophragma utile

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomus* (Schaufuss), *Doliopygus brevis* (Strohmeyer), *Doliopygus chapuisi* (Duvivier), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus nimicus* (Schedl), *Doliopygus perbrevis* Schedl, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).

Enterolobium cyclocarpum

Euwallacea xanthopus (Eichhoff).

Enterolobium saman

Platypus cupulatus Chapuis.

Enterolobium spp.

Euwallacea fornicatus (Eichhoff), *Xyleborus rotulus* (Fabricius).

Eperua falcata

Platypus tiriocensis Reichardt, *Xyleborus parcellus* Wood.

Epicattleya spp.

Xylosandrus compactus (Eichhoff).

Epidendrum spp.

Xylosandrus compactus (Eichhoff).

Epischlia spp.

Hypothenemus bezaziani Peyerimhoff.

Eranthemum spp.

Xylosandrus compactus (Eichhoff).

Eriobotrya japonica

Hypothenemus eruditus Westwood.

Eriocoelem microspermum

Dryocetoctoides cristatus (Fabricius).

Eriocoelem spp.

Doliopygus erichsoni (Chapuis).

Erythrina abyssinica

Euwallacea xanthopus (Eichhoff), *Platypus impressus* (Strohmeyer), *Platypus solutus* Schedl, *Xyleborus alluaudi* Schaufuss, *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff.

Erythrina burtii

Euwallacea xanthopus (Eichhoff).

Erythrina costaricensis

Monarthrum finbriaticorne (Blandford), *Theoborus theobromae* Hopkins, *Xyleborus affinis* Eichhoff, *Xyleborus morulus* Blandford, *Xyleborus posticus* Eichhoff.

Erythrina cristagalli

Araptus hostilis (Blackman).

Erythrina indica

Xyleborus perforans (Wollaston).

Erythrina lithosperma

Euwallacea fornicatus (Eichhoff), *Euwallacea xanthopus* (Eichhoff), *Scolytotplatypus hirsutus* Blackman, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Erythrina orientalis

Euwallacea fornicatus (Eichhoff).

Erythrina peoppigiana*Xyleborus ferrugineus* (Fabricius).***Erythrina* spp.**

Araptus hymenaeae (Eggers), *Doliopygus erichsoni* (Chapuis), *Eucallalca fornicatus* (Eichhoff), *Eucallalca interjectus* (Blandford), *Eucallalca xanthopus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus pennatus* Schedl, *Platypus pseudocupulatus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluandi* Schaufuss, *Xyleborus piccus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).

Erythrina cespertilio*Xyleborus similis* Ferrari.***Erythrocoeca* spp.**

Eucallalca xanthopus (Eichhoff), *Xyleborus alluandi* Schaufuss, *Xyleborus principalis* Eichhoff.

Erythrophleum guineense

Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus artespinatus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus exilis* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus maynei* Schedl, *Doliopygus medius* Schedl, *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicomis* Schedl, *Periommatius excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xyleborus colvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Erythrophleum micranthum

Doliopygus dubius (Sampson), *Doliopygus exilis* (Chapuis), *Doliopygus tenuis* (Strohmeyer).

Erythrophleum* spp.Xyleborus affinis* Eichhoff.***Erythroxydon noragranatense***

Cnestus bicornis (Eggers), *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Erythroxydon sphaeranthrum*Cosmoderes niger* (Schedl).***Erythroxydon* spp.**

Mitosoma obconiceps Schedl, *Mitosoma obliquatum* Schedl.

Eschceilera corrugata

Monarthrum dimidiatum (Ferrari), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).

Eschceilera grata*Xyleborus affinis* Eichhoff.***Eschceilera sagotiana***

Platypus longulus Chapuis, *Platypus tiriosensis* Reichardt, *Xyleborus parvulus* Wood.

***Eschceilera* spp.**

Phelloterus anaxenus Wood, *Phelloterus atroviridis* Wood, *Phelloterus tersus* Wood, *Phloeotribus tetricus* Wood, *Phloeotribus venezuelensis* (Schedl), *Platypus filaris* Wood, *Platypus secus* Wood, *Pseudothysanocis subulatus* (Wood), *Tesseroeranus nevermanni* Schedl, *Xyleborus colvulus* (Fabricius).

Eschceilera subgladulosa*Dryocoetoides severus* Wood.***Eucalyptus acmenoides***

Hypothenemus melasomus (Lea), *Notoplatypus elongatus* Lea.

Eucalyptus bosistoana*Chaetastus tuberculatus* (Chapuis).***Eucalyptus botryoides****Chaetastus montanus* Schedl, *Xyleborus integer* Schedl.***Eucalyptus citridora***

Ambrosiodmus neglectus (Schedl), *Chaetastus montanus* Schedl, *Crossotarsus externedentatus* (Fairmaire), *Doliopygus artespinatus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus excavatus* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus unicomis* Schedl, *Platypus solutus* Schedl, *Xyleborinus forcipatus* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus perforans* (Wollaston), *Xyleborus principalis* Eichhoff, *Xyleborus colvulus* (Fabricius).

Eucalyptus deglupta

Eccoptopterus limbatus Sampson, *Platypus associatus* Schedl, *Platypus pseudocupulatus* Schedl, *Xyleborinus exiguus* (Walker), *Xyleborus gorggae* Schedl, *Xyleborus perforans* (Wollaston).

Eucalyptus drepanophylla

Notoplatypus elongatus Lea, *Xyleborus perforans* (Wollaston).

Eucalyptus excelsa*Dryocoetoides pseudosolitaris* (Eggers).***Eucalyptus fastigata****Austroplatypus incomptus* (Schedl).***Eucalyptus gigantea****Austroplatypus incomptus* (Schedl).***Eucalyptus globulus***

Scolytiplatypus darjeelingi Stebbing, *Xyleborus improbus* Sampson.

Eucalyptus grandis

Diaprius quinquespinatus Chapuis, *Platypus solidus* Walker, *Platypus strenuus* Schedl, *Xyleborus perforans* (Wollaston).

Eucalyptus intermedia*Xyleborus perforans* (Wollaston).***Eucalyptus laeropinea****Platypus incostatus* Schedl.***Eucalyptus maculata***

Crossotarsus externedentatus (Fairmaire), *Platypus subgranosus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus antaisaka* Schedl, *Xyleborus perforans* (Wollaston).

Eucalyptus microcoris*Ambrosiodmus albizzianus* (Schedl).***Eucalyptus obliqua****Austroplatypus incomptus* (Schedl).***Eucalyptus paniculata***

Crossotarsus externedentatus (Fairmaire), *Xyleborus affinis* Eichhoff, *Xyleborus antaisaka* Schedl, *Xyleborus squamulatus* Eichhoff.

Eucalyptus pilularis*Austroplatypus incomptus* (Schedl).***Eucalyptus robusta***

Mitosoma truncatipennis Schedl, *Platypus furcatus* Blandford, *Xyleborus antaisaka* Schedl, *Xyleborus iheringi* Iglesias, *Xyleborus madagascariensis* Schaufuss, *Xyleborus sakalava* Schedl.

Eucalyptus rostrata*Platypus mutatus* Chapuis.

Eucalyptus saligna

Ambrosiodinus eichhoffi (Schreimer), *Chaetastus tuberculatus* (Chapuis), *Doliopygus artespinatus* (Schedl), *Doliopygus erichsoni* (Chapuis), *Platypus solitus* Schedl, *Xyleborus principalis* Eichhoff.

Eucalyptus seeana

Xyleborus perforans (Wollaston).

Eucalyptus siberiana

Austroplatypus incomptus (Schedl).

***Eucalyptus* spp.**

Amasa truncatus (Erichson), *Ambrosiodinus aegir* (Eggers), *Ambrosiodinus albizzianus* (Schedl), *Ambrosiodinus compressus* (Lea), *Ambrosiodinus innominatus* (Schedl), *Ambrosiodinus natalensis* (Schaufuss), *Ambrosiodinus obliquus* (LeConte), *Ambrosiodinus tropicus* (Hagedorn), *Crossotarsus fractus* Sampson, *Crossotarsus javanus* Beeson, *Diapus malgassicus* Schedl, *Doliopygus erichsoni* (Chapuis), *Doliopygus nairobiensis* (Schedl), *Dryocoetoides solitarinus* (Schedl), *Mitosoma obliquatum* Schedl, *Platypus navarrodeandradei* Marelli, *Platypus omnivorus* (Lea), *Prennobilus capipennis* Eichhoff, *Xyleborinus saxescens* (Ratzeburg), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus submolestus* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus solidus* (Eichhoff).

Eucalyptus tereticornis

Xyleborus perforans (Wollaston).

Eucalyptus torrelliana

Coptodryas parva (Lea).

Eucalyptus trachyphloia

Hypothenemus birmanus (Eichhoff).

Euclea nataleensis

Lamurgus podocarpi Schedl.

Eugenia aequa

Dryocoetops eugeniae (Schedl).

Eugenia argutata

Diapus pusillinus Chapuis, *Diapus quinquespinatus* Chapuis.

Eugenia aromatica

Amasa batocerradensis (Schedl), *Platypus caryophyllatus* Blandford.

Eugenia bernardii

Crossotarsus squamulatus Chapuis.

Eugenia buxifolia

Dendrosinus bourreriae Schwarz, *Hypothenemus birmanus* (Eichhoff), *Hypothenemus hirsutus* (Wood), *Hypothenemus seriatus* (Eichhoff).

Eugenia caryophyllata

Platypus caryophyllatus Schedl, *Xylosandrus cylindrotomicus* (Schedl).

Eugenia caryophylla

Xyleborus perforans (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).

Eugenia cumini

Coccotrypes carpophagus (Hornung).

Eugenia formosa

Coccotrypes advena Blandford, *Coccotrypes cyperi* (Beeson), *Coccotrypes papuanus* (Eggers).

Eugenia glomerata

Platypus cupulatus Chapuis.

Eugenia grandis

Crossotarsus squamulatus Chapuis, *Crossotarsus wallacei* (Thomson).

Eugenia jambolana

Amasa eugeniae (Eggers), *Coccotrypes malgasicus* (Schedl), *Coccotrypes nubihus* (Blandford), *Coptodryas elegans* (Sampson), *Eucallacca bicolor* (Blandford), *Platypus cupulatus* Chapuis, *Platypus curtatus* Sampson, *Platypus errans* (Sampson), *Platypus eugeniae* Schedl, *Platypus latifinis* Walker, *Platypus madagascariensis* Chapuis, *Platypus uncinatus* Blandford, *Xyleborinus andreweesi* (Blandford), *Xyleborinus artemisiatus* (Eichhoff), *Xyleborinus forficuloides* (Schedl), *Xyleborinus reconditus* (Schedl), *Xyleborus haberkorui* Eggers, *Xyleborus jambolanaensis* Schedl, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).

Eugenia lineata

Amasa circumcisulus (Schedl).

Eugenia malaccensis

Crossotarsus externedentatus (Fairmaire), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston).

Eugenia polyantha

Xylosandrus morigerus (Blandford).

Eugenia quadribraeteata

Cyclorhipidion agnatum (Eggers).

Eugenia similis

Coccotrypes advena Blandford.

***Eugenia* spp.**

Amasa aspersus (Sampson), *Amasa opalescens* (Schedl), *Arixyleborus malayensis* (Schedl), *Arixyleborus scabripennis* (Blandford), *Arixyleborus suturalis* (Eggers), *Cenocephalus rugicollis* Schedl, *Coccotrypes cyperi* (Beeson), *Coptoborus bicolor* (Browne), *Coptoborus fragilis* (Browne), *Crossotarsus minax* (Walker), *Crossotarsus semicinctus* Schedl, *Crossotarsus terminatus* Chapuis, *Diapus papuanus* Schedl, *Eucallacca fomicatus* (Eichhoff), *Eucallacca laevis* (Eggers), *Eucallacca xanthopus* (Eichhoff), *Lamurgus cribrellus* Schedl, *Mitosoma sexspinosum* Schedl, *Platypus dignatus* Schedl, *Platypus lepidus* Chapuis, *Platypus longulus* Chapuis, *Platypus madagascariensis* Chapuis, *Platypus ovatus* Strohmeier, *Platypus westwoodi* Chapuis, *Scolytotplatypus parvus* Sampson, *Treptotplatypus trepanatus* (Chapuis), *Xyleborus adusticollis* (Motschulsky), *Xyleborus antaisaka* Schedl, *Xyleborus antanala* Schedl, *Xyleborus apertus* Schedl, *Xyleborus javanus* Eggers, *Xylosandrus compactus* (Eichhoff), *Xylosandrus omissus* (Schedl), *Xylosandrus oralis* (Schedl).

Eugenia subglauca

Xyleborinus andreweesi (Blandford).

Euonymus alatus

Crossotarsus niponicus Blandford.

Euonymus atropurpureus

Hypothenemus eruditus Westwood.

***Euonymus* spp.**

Allernoporus euonymi Kurenzov, *Hypothenemus eruditus* Westwood.

Eupatorium daleoides

Pityophthorus hermosus Wood, *Pityophthorus morosus* Wood.

Eupatorium pallescens

Arixyleborus scabripennis (Blandford), *Cnestus nitidipennis* (Schedl), *Cnestus suturalis* (Eggers), *Hadrodemus globus* (Blandford), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Eupatorium spp.

Eucallacea barbatulus (Schedl), *Scolytotplatypus nitidus* Eggers, *Xyleborus ciliatus* Eggers.

Eupatorium tapos

Xyleborus quadrispinosulus Eggers.

Eupatorium tjibeureum

Xylosandrus eupatorii (Eggers).

Euphorbia abyssinica

Triotennus aethiopicus Eggers.

Euphorbia amygdaloides

Thamnuergus varipes Eichhoff.

Euphorbia antiquorum

Cryphalogenes euphorbiae Wood, *Cryphalogenes exiguus* Wood.

Euphorbia balsamifera

Aphanarthrum affine Wollaston, *Aphanarthrum bicinctum* Wollaston, *Aphanarthrum caescens* Wollaston, *Aphanarthrum monodi* Paulian & Villiers, *Liparthrum bicaudatum* Wollaston.

Euphorbia beaumierana

Aphanarthrum mairei Peyerimhoff, *Cisurgus occidentalis* Peyerimhoff, *Colcobothisr alluaudi* (Peyerimhoff), *Triotennus longicollis* Peyerimhoff.

Euphorbia cactiformes

Cisurgus occidentalis Peyerimhoff.

Euphorbia canariensis

Aphanarthrum bicinctum Wollaston, *Aphanarthrum canariensis* Wollaston, *Aphanarthrum pygmaeum* Wollaston, *Cisurgus wollastoni* (Eichhoff), *Colcobothisr luridus* (Wollaston), *Triotennus subretusus* Wollaston.

Euphorbia characias

Thamnuergus characiae Rosenhauer, *Thamnuergus varipes* Eichhoff.

Euphorbia dendroides

Aphanarthrum affine Wollaston, *Liparthrum inarmatum* Wollaston, *Thamnuergus euphorbiae* (Kuster).

Euphorbia dregei

Styracoptinus euphorbiae (Bright).

Euphorbia echinus

Aphanarthrum mairei Peyerimhoff, *Cisurgus occidentalis* Peyerimhoff, *Colcobothisr alluaudi* (Peyerimhoff), *Triotennus longicollis* Peyerimhoff.

Euphorbia gerardiana

Thamnuergus euphorbiae (Kuster), *Thamnuergus varipes* Eichhoff.

Euphorbia handiensis

Aphanarthrum mairei Peyerimhoff, *Colcobothisr alluaudi* (Peyerimhoff).

Euphorbia megalallantica

Thamnuergus mairei Peyerimhoff.

Euphorbia mellifera

Aphanarthrum euphorbiae Wollaston.

Euphorbia nerifolia

Aphanarthrum indicum Wood.

Euphorbia obtusifolia

Aphanarthrum bicinctum Wollaston.

Euphorbia onoclada

Hypothenemus balachowskyi Menier, *Hypothenemus teleforti* (Menier).

Euphorbia piscatoria

Aphanarthrum bicolor Wollaston, *Aphanarthrum piscatorium* Wollaston, *Liparthrum inarmatum* Wollaston.

Euphorbia regis-jubae

Aphanarthrum affine Wollaston, *Aphanarthrum bicin-*

tum Wollaston, *Aphanarthrum bicolor* Wollaston, *Aphanarthrum caescens* Wollaston, *Aphanarthrum jubae* Wollaston, *Liparthrum bicaudatum* Wollaston, *Liparthrum lowei* Wollaston.

Euphorbia resinifera

Aphanarthrum saturatum Peyerimhoff, *Triotennus longicollis* Peyerimhoff.

Euphorbia royleana

Aphanarthrum reticulatum Wood, *Aphanarthrum royalecanum* Wood, *Eucallacea interjectus* (Blandford).

Euphorbia spp.

Aphanarthrum armatum Wollaston, *Aphanarthrum glabrum* Wollaston, *Aphanarthrum piscatorium* Wollaston, *Aphanarthrum reticulatum* Wood, *Aphanarthrum royalecanum* Wood, *Aphanarthrum subglabrum* Israelson, *Aphanarthrum wollastoni* Israelson, *Colcobothisr gerneauxi* Menier, *Cyrtogenius africanus* Wood, *Cyrtogenius scotti* Eggers, *Liparthrum curtum* Wollaston, *Liparthrum lowei* Wollaston, *Triotennus scrofa* (Schedl), *Triotennus villiersi* Schedl.

Euphorbia stenoclada

Hypothenemus euphorbiae (Schedl), *Lamurgus euphorbiae* Schedl, *Lamurgus rugosipes* Schedl.

Euphorbia sudanica

Aphanarthrum duongi Villiers.

Euphorbia teke

Ambrsiodmus albizzianus (Schedl).

Euphorbia tirucalli

Acanthotomicus euphorbiae (Schedl), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus eruditus* Westwood.

Euphorbia tuckeyana

Aphanarthrum hesperidum Wollaston, *Liparthrum loweanum* Wollaston.

Euphorbia wulfenii

Thamnuergus sardus Eggers.

Euphorbia spp.

Xylosandrus compactus (Eichhoff).

Euptelea polyandra

Crossotarsus niponicus Blandford.

Euroschinus falcatus

Coptodryas eucalyptica (Schedl), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Eurya japonica

Baiois perinimicus (Schedl), *Eucallacea velatus* (Sampson), *Platypus tenuissimus* Schedl, *Sucus nissimai* (Eggers), *Xyleborus octiesdentatus* Murayama, *Xyleborus volutus* (Fabricius).

Euryops athanasiae

Thamnuergus euryopsis Schedl.

Eusideroxylon zwageri

Platypus biuncus Blandford, *Platypus cupulatus* Chapuis, *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Euterpe globosa

Coccotrypes robustus Eichhoff.

Euterpe oleracea

Araptus lactigatus (Eggers).

Erodia fraxinifolia

Cyclorhipidion hirtum (Hagedorn), *Platypus falcatus* Strohmeier, *Scolytotplatypus pubescens* Hagedorn, *Xyleborus hagedornianus* Schedl.

Erodia latifolia

Xyleborus piccus (Motschulsky).

Erodia meliaeefolia

Xyleborus atratus Eichhoff.

Erodia rutaecarpu*Cryphalus exiguus* Blandford.***Erodia* spp.***Ambrosiodinus lewisi* (Blandford), *Crossotarsus paucidentatus* Roberts, *Crossotarsus rasilis* Roberts, *Hypothenemus parvulus* Browne, *Platypus bajulus* Schedl, *Platypus jansoni* Chapuis, *Platypus juvenis* Schedl, *Platypus schysi* Chapuis, *Platypus solidus* Walker, *Scolytogenes papuanus* (Schedl).***Excoecaria agallocha****Cryphalus scabricollis* Eichhoff, *Xyleborus bidentatus* (Motschulsky), *Xyleborus cognatus* Blandford.***Excoecaria dallachyana****Scolytogenes atterimus* Wood, *Xyleborus similis* Ferrari.***Exogonium jalapa****Scolytogenes jalapae* (Letzner).***Eysenhardtia polyotucha****Chaetophloeus mexicanus* (Blackman).***Eysenhardtia* spp.***Chaetophloeus mexicanus* (Blackman).***Fagara angolense****Doliopygus chapuisi* (Duvivier), *Doliopygus excavatus* (Sampson), *Doliopygus serratus* (Strohmeyer).***Fagara capensis****Platypus sampsoni* (Schedl), *Xyleborinus acmulus* (Wollaston).***Fagara heitzi****Platypus hintzi* Schauffuss.***Fagara lemairei****Doliopygus expletus* Schedl, *Platypus orientalis* (Strohmeyer), *Xyleborus ferrugineus* (Fabricius).***Fagara macrophylla****Chaetastus tuberculatus* (Chapuis), *Doliopygus conradi* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus interjectus* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatus grandis* Schedl, *Pityophthorus joveri* Schedl, *Platypus hintzi* Schauffuss, *Platypus impolitus* Roberts, *Platypus parallelus* (Fabricius), *Polygraphus tenuis* Schedl, *Prennobius nodulosus* Hagedorn, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus sub tuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).***Fagara* spp.***Euwallacea xanthopus* (Eichhoff), *Leptoxyleborus concisus* (Blandford), *Periommatus longicollis* Chapuis, *Platypus orientalis* (Strohmeyer), *Prennobius capipennis* Eichhoff, *Xyleborus ananicius* Hagedorn, *Xyleborus ferrugineus* (Fabricius), *Xyleborus principalis* Eichhoff.***Fagaropsis angolensis****Chaetastus tuberculatus* (Chapuis).***Fagraea fragrans****Anasa versicolor* (Sampson), *Arixyleborus suturalis* (Eggers), *Xyleborus perforans* (Wollaston), *Xyleborus submarginatus* Eggers.***Fagraea gigantea****Crossotarsus wallacei* (Thomson), *Diapus quinque-spinatus* Chapuis, *Euwallacea fornicatus* (Eichhoff), *Leptoxyleborus concisus* (Blandford), *Platypus suffodiens* Sampson, *Xyleborus emarginatus* Eichhoff, *Xyleborus fallax* Eichhoff.***Fagraea americana****Monarthrum mali* (Fitch).***Fagus crenata****Coccotrypes fagi* (Nobuchi), *Crossotarsus niponicus* Blandford, *Ernoporus shimaneensis* Murayama, *Indocryphalus pubipennis* (Blandford), *Platypus calamus* Blandford, *Platypus lewisi* Blandford, *Platypus modestus* Blandford, *Platypus severini* Blandford, *Taphrorhynchus pilosus* (Murayama), *Xyleborinus attenuatus* (Blandford), *Xyleborinus schaufussi* (Blandford), *Xyleborus atratus* Eichhoff, *Xyleborus seriatus* Blandford, *Xylosandrus mutilatus* (Blandford).***Fagus grandifolia****Cnesinus strigicollis* LeConte, *Dryocoetes betulae* Hopkins, *Hyllocurus rudis* (LeConte), *Hypothenemus interstitialis* (Hopkins), *Hypothenemus rotundicollis* (Eichhoff), *Pseudopityophthorus fagi* Blackman, *Scolytus fagi* Walsh, *Xyleborinus saxescni* (Ratzeburg), *Xyleborus ferrugineus* (Fabricius), *Xyleborus obesus* LeConte.***Fagus japonica****Crossotarsus niponicus* Blandford, *Hyorrhynchus lewisi* Blandford, *Platypus modestus* Blandford.***Fagus multinervis****Euwallacea validus* (Eichhoff), *Xyleborinus saxescni* (Ratzeburg), *Xylosandrus germanus* (Blandford).***Fagus orientalis****Dryocoetes abii* (Georg), *Ernoporus fagi* (Fabricius), *Phloeotribus brevicollis* (Kolenati), *Platypus cylindrus* (Fabricius), *Taphrorhynchus bicolor* (Herbst), *Xyleborus monographus* (Fabricius).***Fagus* spp.***Ambrosiodinus tachygraphus* (Zimmermann), *Hylesinus toranio* (Danthione), *Platypus caticeps* Broun, *Platypus lewisi* Blandford, *Pseudopityophthorus minutissimus* (Zimmermann), *Scolytoplatypus daimio* Blandford, *Scolytoplatypus mikado* Blandford, *Scolytoplatypus shogun* Blandford, *Scolytoplatypus tycon* Blandford, *Scolytus carpini* (Ratzeburg), *Scolytus intricatus* (Ratzeburg), *Scolytus laevis* Chapuis, *Scolytus pygmaeus* (Fabricius), *Trypodendron domesticum* (Linnaeus), *Trypodendron signatum* (Fabricius), *Xyleborus dispar* (Fabricius), *Xyloterinus politus* (Say).***Fagus sylvatica****Ernoporus fagi* (Fabricius), *Ernoporus tiliae* (Panzer), *Platypus cylindrus* (Fabricius), *Taphrorhynchus bicolor* (Herbst), *Taphrorhynchus hirtellus* Eichhoff, *Xyleborus pfeili* (Ratzeburg), *Xyleborus sayi* (Hopkins), *Xylosandrus germanus* (Blandford).***Fanchera* spp.***Euwallacea xanthopus* (Eichhoff).***Faurea saligna****Chaetastus montanus* Schedl, *Cyclorhipidion cruciforme* (Schedl), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Platypus intermedius* (Schedl), *Polygraphus granulifer* Eggers, *Polygraphus kieuensis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss.***Ferdinandia adolphi-federici****Doliopygus coelocephalus* (Schauffuss), *Doliopygus conradi* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus punctiventris* Schedl, *Doliopygus serratus* (Strohmeyer), *Trachyostus atterratus* (Schauffuss), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).***Ferula* spp.***Cisurgus ferulae* Pfeffer.***Ficus ampelos****Xylosandrus morigerus* (Blandford).

Ficus aspera*Ficicis varians* Lea.***Ficus asperifolia****Chaetastus tuberculatus* (Chapuis), *Eccoptypterus spinosus* (Olivier), *Platypus hintzi* Schaufuss, *Platypus spinulosus* (Strohmeyer), *Trachyostus aterrimus* (Schaufuss).***Ficus asperima****Platypus latifinis* Walker, *Xyleborus similis* Ferrari.***Ficus aurea****Cncemonix ficus* (Schwarz), *Cryptocarenum seriatus* Eggers, *Hypothenemus birmanus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Hypothenemus javanus* (Eggers).***Ficus benghalensis****Diamerus variegatus* Schedl.***Ficus benjamina****Platypus curvus* Strohmeyer, *Platypus jansonii* Chapuis, *Xyleborus piceus* (Motschulsky).***Ficus bracteata****Coccotrypes advena* Blandford, *Coccotrypes cinnamomi* (Eggers).***Ficus capensis****Cryphalus kicuensis* Schedl, *Doliopygus bidentatus* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus rhodesianus* (Schedl), *Euwallacea xanthopus* (Eichhoff), *Hylesinopsis confusus* (Eggers), *Hylesinopsis granulatus* (Lepesme), *Hypothenemus eruditus* Westwood, *Hypothenemus lefevrei* (Schedl), *Platypus impressus* (Strohmeyer), *Platypus solutus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus principalis* Eichhoff.***Ficus carica****Ambrosiodmus lewisi* (Blandford), *Aricerus eichhoffi* Blandford, *Cryphalus wapleri* Eichhoff, *Ficicis varians* Lea, *Hadrodemus amorphus* (Eggers), *Hypoborus ficus* Erichson, *Hypothenemus eruditus* Westwood, *Liparthrum curtum* Wollaston.***Ficus chorocarpa****Xyleborinus artemcomans* (Schedl).***Ficus corylifolia****Xyleborus ferrugineus* (Fabricius).***Ficus cotinifolia****Phloeotribus setulosus* Eichhoff.***Ficus elastica****Diamerus fici* Blandford, *Platypus pseudocupulatus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus perforans* (Wollaston).***Ficus eriobotryoides****Chaetastus montanus* Schedl, *Platypus ficus* Roberts.***Ficus exasperata****Chaetastus tuberculatus* (Chapuis), *Diamerus imperfectus* Eggers, *Diapus quinquespinnatus* Chapuis, *Doliopygus brevis* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus medius* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer), *Periommatatus excisus* Strohmeyer, *Periommatatus pseudomajor* Schedl, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus eichhoffianus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus scobinatus* Hagedorn, *Xyleborus colvulus* (Fabricius).***Ficus fistulosa****Scolytoplatypus nitidus* Eggers, *Xyleborus perforans* (Wollaston).***Ficus glabella****Diamerus minor* Eggers.***Ficus glabrata****Hylocurus disparilis* Wood.***Ficus glomerata****Coccotrypes cyperi* (Beeson), *Coccotrypes variabilis* (Beeson), *Cryphalus dorsalis* (Motschulsky), *Platypus curvus* Strohmeyer, *Platypus cupulatus* Chapuis, *Platypus lepidus* Chapuis, *Platypus uncinatus* Blandford, *Xyleborus perforans* (Wollaston).***Ficus hauli****Xylosandrus ursulus* (Eggers).***Ficus hispida****Euwallacea malloti* (Eggers).***Ficus infectoria****Platypus errans* (Sampson), *Xyleborus perforans* (Wollaston), *Xyleborus pumilus* Eggers.***Ficus laurifolia****Cladoctonus major* (Eggers).***Ficus macrophylla****Acanthotomicus australis* (Schedl), *Aricerus eichhoffi* Blandford.***Ficus macrosperma****Platypus ficus* Roberts.***Ficus mallatocarpa****Periommatatus grandis* Schedl.***Ficus melleri****Cryphalus pallidus* Eichhoff.***Ficus minahassae****Cryphalus densepilosus* Schedl, *Platypus luniger* Motschulsky, *Xyleborus affinis* Eichhoff.***Ficus mucoso****Hypothenemus eruditus* Westwood, *Xyleborus ficus* Eggers.***Ficus natalensis****Eccoptypterus spinosus* (Olivier), *Xyleborus madagascariensis* Schaufuss.***Ficus pilosa****Hypothenemus eruditus* Westwood.***Ficus polita****Hylesinopsis ficus* (Schedl), *Hylesinopsis joveri* (Schedl), *Hypothenemus pubipennis* (Eggers).***Ficus preussii****Doliopygus conradti* (Strohmeyer), *Platypus hintzi* Schaufuss.***Ficus religiosa****Coccotrypes vulgaris* (Eggers), *Diamerus variegatus* Schedl, *Dactylipalpus transversus* Chapuis, *Erniodadius corpidentus* (Sampson), *Xyleborinus artemistriatus* (Eichhoff), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Ficus roxburghii****Coccotrypes longior* (Eggers).***Ficus rubiginosa****Aricerus eichhoffi* Blandford.***Ficus rumphii****Cryphalus dorsalis* (Motschulsky), *Diamerus variegatus* Schedl.***Ficus seretii****Ambrosiodmus tropicus* (Hagedorn), *Xyleborus fallax* Eichhoff.***Ficus soroceoides****Ambrosiodmus triton* (Schaufuss), *Euwallacea xantho-*

pus (Eichhoff), *Mitosoma crenulatum* Chapuis, *Mitosoma dispar* Schedl, *Mitosoma excisum* Schaufuss, *Platypus madagascariensis* Chapuis, *Xyleborinus spiculatus* (Schaufuss), *Xyleborinus spinosus* (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xyleborus sakalava* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus hirsutipennis* (Schedl).

Ficus spp.

Acanthotomicus fici Browne, *Acanthotomicus kepongi* (Schedl), *Acanthotomicus perexiguus* (Blandford), *Ambrosiodmus declivispinatus* (Schedl), *Ambrosiodmus eichhoffi* (Schreiner), *Ambrosiodmus obliquus* (LeConte), *Araptus aztecus* (Wood), *Araptus consobrinus* Wood, *Araptus facetus* Wood, *Araptus obsoletus* (Blandford), *Araptus speciosus* Wood, *Aricerus chapuisi* Blandford, *Aricerus eichhoffi* Blandford, *Arixyleborus scabripennis* (Blandford), *Baioeis inimicus* (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cnemomyx evidens* Wood, *Cnesinus setulosus* Blandford, *Cnestus nitidus* (Schedl), *Coccotrypes gedeanus* (Eggers), *Coccotrypes medius* (Eggers), *Coccotrypes petioli* (Browne), *Corthylus abbreviatus* Eichhoff, *Corthylus additus* Wood, *Corthylus diligens* Wood, *Crossotarsus squamulatus* Chapuis, *Cryphalus exiguus* Blandford, *Cryphalus scabricollis* Eichhoff, *Cryphalus sylvicola* (Perkins), *Cryphalus wapleri* Eichhoff, *Cyclorhupidion crucifer* (Hagedorn), *Diamerus ater* Hagedorn, *Diamerus curcifer* (Walker), *Diamerus fici* Blandford, *Diamerus impar* Chapuis, *Diamerus pulverulentus* Gerstaecker, *Diamerus variegatus* Schedl, *Diapys pusillimus* Chapuis, *Diapys spinifer* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus medius* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus spectabilis* Schedl, *Euwallacea fornicatus* (Eichhoff), *Euwallacea interjectus* (Blandford), *Euwallacea quadratocollis* (Eggers), *Ficicis despectus* (Walker), *Ficicis javanus* (Eggers), *Ficicis maculipennis* (Schedl), *Ficicis philippinensis* (Eggers), *Glostatus acaciae* (Schedl), *Gymnochilus consocius* (Blandford), *Gymnochilus minor* (Blandford), *Gymnochilus reitteri* Eichhoff, *Gymnochilus zonatus* Eichhoff, *Hylesinopsis fasciatus* (Hagedorn), *Hylesinopsis hirsutus* (Schedl), *Hylesinopsis sandiarabiae* (Schedl), *Hypoecryphalus peruvianus* (Schedl), *Hypothenemus brunneus* (Hopkins), *Hypothenemus columbi* Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus erectus* LeConte, *Hypothenemus eruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus mahus* (Schedl), *Hypothenemus seriatus* (Eichhoff), *Hypothenemus trivialis* Wood, *Indocryphalus pubipennis* (Blandford), *Leptoxyleborus concisus* (Blandford), *Liparthrum americanum* Wood, *Liparthrum mexicanum* Wood, *Mesoplatypus kitushi* Schedl, *Mitosoma paulianum* Schedl, *Mitosoma sexspinosum* Schedl, *Monarthrum exornatum* (Schedl), *Periommatatus excisus* Strohmeyer, *Periommatatus grandis* Schedl, *Phloeoborus scaber* Erichson, *Phloeotribus fici* Wood, *Phloeotribus geminus* Wood, *Phloeotribus maurus* Wood, *Phloeotribus opimus* Wood, *Phloeotribus setulosus* Eichhoff, *Platypus deceptor* Wood, *Platypus formosanus* Nisima & Murayama, *Platypus gilveii* Roberts, *Platypus hintzi* Schaufuss, *Platypus indicus* Strohmeyer, *Platypus jansonii* Chapuis, *Platypus loricatus* (Sampson), *Platypus negatus* Schedl, *Platypus nocuus* Schedl, *Platypus orientalis* (Strohmeyer), *Platypus pseudocupulatus* Schedl, *Platypus quadrispinis* Browne, *Platypus reichii* Chapuis, *Platypus solidus*

Walker, *Platypus tenellus* Schedl, *Platypus westwoodi* Chapuis, *Polygraphus kivuensis* Schedl, *Prennobiobius mukunguiae* (Schedl), *Prennobiobius sexspinosus* Eggers, *Pycnarthrum hispidum* (Ferrari), *Pycnarthrum reticulatum* Schedl, *Pycnarthrum setulosum* Waterhouse, *Scolytodes amoenus* Wood, *Scolytodes cubensis* (Schedl), *Scolytodes erinophilus* Wood, *Scolytodes exiguus* Wood, *Scolytodes facetus* Wood, *Scolytodes ficicola* Wood, *Scolytodes ficivorus* Wood, *Scolytodes genialis* Wood, *Scolytodes lepidus* Wood, *Scolytodes micidius* Wood, *Scolytodes pseudopiceus* Wood, *Scolytodes reticulatus* (Wood), *Scolytodes rugicollis* (Schedl), *Scolytodes schwarzi* (Hopkins), *Scolytodes tenuis* (Wood), *Scolytodes venustus* Wood, *Scolytoplatus kivuensis* Schedl, *Scolytoplatus pusillus* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Tricholus simplicis* Wood, *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus amanicus* Hagedorn, *Xyleborus ambasi-pennis* Schedl, *Xyleborus ambasiusculus* Eggers, *Xyleborus barumbiensis* Eggers, *Xyleborus comparabilis* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus multi-spinatus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus posticus* Eichhoff, *Xyleborus similis* Ferrari, *Xyleborus sparsipilosus* Eggers, *Xyleborus tenellus* Schedl, *Xyleborus truncatulus* Schedl, *Xyleborus rotundus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus curtulus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Ficus stenocarpa

Ficicis varians Lea, *Xyleborus perforans* (Wollaston).

Ficus thunningii

Cryphalus bellus Schedl, *Cryphalus pallidus* Eichhoff.

Ficus variegata

Xyleborinus andrewesi (Blandford).

Ficus vogelii

Periommatatus grandis Schedl, *Periommatatus longicollis* Chapuis, *Platypus glabratus* Roberts, *Platypus hintzi* Schaufuss.

Ficus wightiana

Cryphalus ficivorus Murayama, *Cryphalus furukawai* Murayama.

Ficus xylophylla

Coccotrypes cinnamomi (Eggers).

Ficus zenkeri

Doliopygus chapuisi (DuRoi), *Doliopygus serratus* (Strohmeyer).

Fissicalyx fendleri

Xyleborus ferrugineus (Fabricius).

Fissistigma elegans

Acacis malayanus Browne, *Cnestus aterrimus* (Eggers), *Coptodryas curcidentis* (Schedl), *Euwallacea fornicatus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Flacourtia spp.

Xylosandrus compactus (Eichhoff).

Flemingia strobilifera

Xylosandrus morigerus (Blandford).

Flindersia acuminata

Xyleborus insulindicus Eggers.

Flindersia bourjotiana

Xyleborus perforans (Wollaston).

Flindersia brayleyana

Coptodryas eucalyptica (Schedl), *Xyleborus perforans* (Wollaston).

Flindersia laevis*Xyleborus perforans* (Wollaston).***Flindersia pubescens****Peridryocetes queenslandi* Schedl, *Platypus scalaris* Schedl, *Platypus semigranosus* (Sampson).***Flindersia schotiana****Xyleborus emarginatus* Eichhoff.***Flindersia* spp.***Ambrosiodinus compressus* (Lea), *Hylesinus nilgirinus* Eggers, *Xylosandrus compactus* (Eichhoff).***Flourensia cernua****Chaetophloeus sulcatus* (Wood).***Foetida elusoides****Ambrosiodinus inominatus* (Schedl), *Mitosoma lobatum* Schedl, *Xyleborinus mitosomipennis* (Schedl).***Fordia* spp.***Crossotarsus squamulatus* Chapuis, *Hadrodemius comans* (Sampson).***Franzeria* spp.***Pityophthorus franzeriae* Wood, *Pityophthorus torridus* Wood.***Fraxinus americana****Hylesinus aculeatus* Say, *Hylesinus californicus* (Swaine), *Hylesinus criddlei* (Swaine), *Hylesinus fasciatus* LeConte, *Hylesinus pruinosis* Eichhoff.***Fraxinus anomala****Hylesinus mexicanus* (Wood).***Fraxinus chinensis****Ambrosiodinus rubricollis* (Eichhoff).***Fraxinus commemoralis****Platypus contaminatus* (Blandford).***Fraxinus excelsa****Hylesinus crenatus* (Fabricius).***Fraxinus excelsior****Hylesinus eos* Spessivtsev, *Hylesinus orni* Fuchs, *Hylesinus toranio* (Danthione), *Phloeotribus caucasicus* Reitter, *Pteleobius kraatzi* (Eichhoff), *Scolytus kirschi* Skalitzky, *Scolytus scolytus* (Fabricius).***Fraxinus excelsior hookeri****Hylesinus macmahoni* (Stebbing).***Fraxinus floribunda****Hylesinus macmahoni* (Stebbing).***Fraxinus japonica****Platypus calamus* Blandford.***Fraxinus lanuginosa****Platypus severini* Blandford.***Fraxinus lanuginosa serrata****Hylesinus cingulatus* Blandford, *Hylesinus laticollis* Blandford.***Fraxinus latifolia****Hylesinus californicus* (Swaine), *Hylesinus oregonus* (Blackman).***Fraxinus longicauspis****Hylesinus cingulatus* Blandford.***Fraxinus mandshurica****Erioporicus spessivtzevi* Berger, *Hylesinus chodkovskiyi* Berger, *Hylesinus costatus* Blandford, *Hylesinus crenatus* (Fabricius), *Hylesinus eos* Spessivtsev, *Hylesinus lubarskii* Stark, *Hylesinus nobilis* Blandford, *Hylesinus praxidini* Stark, *Hylesinus shabliovskiyi* Kurenzov, *Procrpyphalus fraxini* (Berger), *Scolytus trispinosus* Strohmeier, *Xyleborus dispar* (Fabricius), *Xyleborus maiche* (Stark).***Fraxinus mandshurica japonica****Crossotarsus niponicus* Blandford, *Hylesinus cingulatus* Blandford, *Hylesinus laticollis* Blandford.***Fraxinus ornus****Hylesinus orni* Fuchs, *Hylesinus toranio* (Danthione), *Phloeotribus muricatus* (Eggers).***Fraxinus oxyphylla****Hylesinus crenatus* (Fabricius), *Phloeotribus fraxini* (Eggers).***Fraxinus pennsylvanica****Hylesinus californicus* (Swaine), *Hylesinus criddlei* (Swaine).***Fraxinus rhynchophylla****Hylesinus cingulatus* Blandford, *Hylesinus laticollis* Blandford.***Fraxinus sieboldiana****Hylesinus costatus* Blandford, *Platypus contaminatus* (Blandford).***Fraxinus sogdiana****Hylesinus pritenkii* Sokanovskii.***Fraxinus spaethiana****Hylesinus laticollis* Blandford, *Platypus severini* Blandford.***Fraxinus* spp.***Hylesinus aculeatus* Say, *Hylesinus botscharnikovi* Stark, *Hylesinus crenatus* (Fabricius), *Hylesinus guatemalensis* (Wood), *Hylesinus mandshuricus* Eggers, *Hylesinus tupolevi* Stark, *Hylesinus varius* (Fabricius), *Hylesinus wachli* Reitter, *Phloeotribus mayeti* (Guillebeau), *Platypus compositus* Say, *Platypus cylindrus* (Fabricius), *Scolytoplatypus daimio* Blandford, *Scolytoplatypus tycon* Blandford, *Scolytus japonicus* Chapuis, *Trypodendron domesticum* (Linnaeus), *Trypodendron signatum* (Fabricius), *Xyleborus dispar* (Fabricius), *Xyleborus sayi* (Hopkins), *Xylosandrus compactus* (Eichhoff), *Xylosandrus germanus* (Blandford), *Xylosterinus politus* (Say).***Fraxinus uhdei****Hylesinus aztecus* Wood.***Fraxinus velutina****Hylesinus californicus* (Swaine).***Fremontia californica****Pseudothysanotes hopkinsi* Blackman.***Freyinetia arborea****Coccotrypes dactyliperda* (Fabricius), *Xyleborus ignobilis* Perkins.***Freyinetia hombronii****Xylosandrus morigerus* (Blandford).***Freyinetia* spp.***Xyleborus agamus* Perkins.***Fuchsia* spp.***Xylosandrus morigerus* (Blandford).***Funtumia elastica****Ambrosiodinus acgir* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus conradti* (Strohmeier), *Platypus orientalis* (Strohmeier), *Xyleborinus similans* (Eggers).***Funtumia latifolia****Doliopygus erichsoni* (Chapuis), *Periommatius grandis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus piceus* (Motschulsky).***Galactia spiciformis****Hypothenemus brunneus* (Hopkins), *Hypothenemus californicus* Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Hypothenemus squamosus* (Hopkins).***Galbulinima belgraveana****Platypus dobmanniae* Roberts & Morimoto.

Galbulimima spp.

Diapus elongatus Schedl, *Diapus oregonus* Roberts, *Platypus cymbiformis* Roberts, *Platypus galbulimimae* Roberts & Morimoto, *Platypus gunicus* Roberts, *Platypus longissimus* Roberts, *Platypus noepartibilis* Roberts.

Galiniera coffeoides

Xyleborinus collarti (Eggers).

Ganua spp.

Cryphalus nitens Browne.

Garcinia gnetoides

Doliopygus nigeriensis Roberts.

Garcinia merguensis

Coccotrypes gedeanus (Eggers), *Coccotrypes tunggali* (Schedl), *Xyleborus bidentatus* (Motschulsky).

Garcinia myrtifolia

Platypus citicensis Roberts.

Garcinia polyantha

Chaetastus persimilis (Schedl), *Dactylipalpus parricida* Eggers, *Doliopygus artespianus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus serratus* (Strohmeyer), *Mesoplatypus kitushi* Schedl, *Periommatius longicollis* Chapuis, *Phrixosoma major* (Eggers), *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Polygraphus longipilis* Schedl, *Polygraphus sulcatus* Schedl, *Premnobius hystrix* (Schedl), *Premnobius nodulosus* Hagedorn, *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xyleborus scobinatus* Hagedorn, *Xylosandrus crassiusculus* (Motschulsky).

Garcinia punctata

Chaetastus tuberculatus (Chapuis), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Hypothenemus eruditus* Westwood, *Periommatius longicollis* Chapuis, *Phrixosoma garciniae* (Schedl), *Phrixosoma uniseriata* (Eggers), *Platypus intermedius* (Schedl), *Polygraphus subsulcatus* Schedl, *Premnobius mukunyeae* (Schedl), *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus pilosellus* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus katangensis* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Garcinia smeathmannii

Chaetastus montanus Schedl.

Garcinia spp.

Amasa schlichii (Stebbing), *Ambrosiodmus tropicus* (Hagedorn), *Baiocis nubilosus* Roberts & Morimoto, *Coptodryas nugax* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus nitidulus* (Schedl), *Cryphalus ater* Browne, *Diamerus impar* Chapuis, *Diamerus luteus* Hagedorn, *Diapus angustidentus* Roberts, *Diapus hispinus* Schedl, *Diapus nubilosus* Roberts, *Diapus oomsis* Roberts, *Diapus papuanus* Schedl, *Diapus pusillinus* Chapuis, *Diapus turgidus* Roberts, *Hypocryphalus sandakanensis* (Schedl), *Peridryocetes nitens* (Schedl), *Phrixosoma psaltes* (Hagedorn), *Platypus apicaloides* Schedl, *Platypus cupulatus* Chapuis, *Platypus gumienis* Roberts, *Platypus hamaticollis* Schedl, *Platypus itagaii* Roberts & Morimoto, *Platypus pseudocupulatus* Schedl, *Polygraphus tenuis* Schedl, *Premnobius caripennis* Eichhoff, *Scolytogenes gracilis* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus brevipennis* Schedl, *Xyleborus colvulus* (Fabricius).

Garcinia verrucosa

Ambrosiodmus camphorae (Hagedorn), *Mitosoma*

lobatum Schedl, *Polygraphus creber* Schedl, *Xyleborinus cupulatus* (Schedl), *Xyleborinus forficuloides* (Schedl).

Gardenia spp.

Xyleborus nitens Browne.

Garuga floribunda

Platypus petax Schedl.

Garuga pinnata

Crossotarsus externedentatus (Fairmaire), *Cryphalus scabricollis* Eichhoff, *Eucallacca andamanensis* (Blandford), *Eucallacca interjectus* (Blandford), *Platypus solidus* Walker, *Xyleborinus andrevesi* (Blandford), *Xyleborus perforans* (Wollaston).

Garuga spp.

Platypus jansonii Chapuis, *Xyleborinus andrevesi* (Blandford).

Geumarouna panamensis

Xyleborus affinis Eichhoff.

Gaylussacia spp.

Corthylus punctatissimus (Zimmermann), *Xylosandrus crassiusculus* (Motschulsky).

Geijera salicifolia

Platypus solidus Walker.

Geniosporum paludosum

Eucallacca xanthopus (Eichhoff), *Hypothenemus grandis* Schedl, *Thamurgus granulicollis* Schedl, *Xylocleptes sidanus* (Schedl).

Genista horrida

Liparthrum genistae (Aube).

Genista numidica

Liparthrum genistae (Aube), *Phloeotribus peyerimhoffi* (Eggers).

Genista spp.

Phloeotribus cristatus (Fauvel), *Phloeotribus rhododactylus* (Marshall).

Geranium spp.

Hypothenemus eruditus Westwood.

Germari spp.

Hypothenemus eruditus Westwood.

Gilbertiodendron dewevrei

Coccotrypes grandis (Eggers), *Doliopygus lebruni* Schedl, *Doliopygus tenuis* (Strohmeyer), *Hypothenemus biserialis* (Eggers), *Hypothenemus macrolobii* (Eggers), *Hypothenemus socialis* (Schedl), *Trachyostus aterrimus* (Schaufuss), *Triozastus marshalli* (Sampson), *Xyleborus ambasiusculus* Eggers.

Gilbertiodendron spp.

Hypothenemus camerunus (Eggers).

Giricidia sepium

Hypothenemus brunneus (Hopkins).

Gironniera spp.

Crossotarsus wallacei (Thomson).

Girstoua jentiusiana

Platypus curtatus Sampson.

Gleditschia japonica

Neopteleobius scutellatus (Blandford).

Gleditschia spp.

Dryocoetoides pseudosolitaris (Eggers), *Xyleborus dispar* (Fabricius), *Xyleborus spinulosus* Blandford.

Gleditschia triacanthos

Hylocurus langstoni Blackman, *Thysanoes fimbriicornis* LeConte.

Giricidia maculata

Xylosandrus crassiusculus (Motschulsky).

Gliricidia sepium

Eucallacea fornicatus (Eichhoff), *Xyleborus perforans* (Wollaston).

Glochidion spp.

Diaprus papuanus Schedl, *Diaprus quinquespinatus* Chapuis, *Diaprus turgidus* Roberts, *Eucallacea xanthopus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Gluta elegans

Crossotarsus browni Schedl, *Cryphalus substriatus* Schedl.

Gluta renga

Platypus biuncus Blandford.

Gluta spp.

Diaprus quinquespinatus Chapuis.

Gluta tourtour

Xylosandrus crassiusculus (Motschulsky).

Gluta travancorica

Coccotrypes cyperi (Beeson), *Crossotarsus minas* (Walker), *Eccoptopterus spinosus* (Olivier), *Pityophthorus glutae* Wood, *Platypus carus* Strohmeier, *Xylosandrus crassiusculus* (Motschulsky).

Glycine max

Hypothenemus crudiae (Panzer).

Gmelina arborea

Anasa versicolor (Eichhoff), *Ambrosiodinus lantanae* (Eggers), *Coccotrypes salakensis* (Schedl), *Coptodryas mus* (Eggers), *Eccoptopterus sagittarius* Schedl, *Eucallacea andamanensis* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Eucallacea malloiti* (Eggers), *Eucallacea velatus* (Sampson), *Platypus insularis* Strohmeier, *Platypus pseudocupulatus* Schedl, *Platypus solidus* Walker, *Platypus uncinatus* Blandford, *Premnobius ambitiosus* (Schauffuss), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus pseudopilifer* Schedl, *Xylosandrus difficilis* (Eggers), *Xylosandrus metagermanus* (Schedl).

Gomphia serrata

Xylosandrus manicus (Blandford).

Gonioma camassi

Xyleborinus aenulus (Wollaston).

Gonystylus bancanus

Platypus hamaticollis Schedl, *Platypus pseudocupulatus* Schedl, *Platypus signatus* Chapuis, *Platypus westwoodi* Chapuis, *Treptoplatypus trepanatus* (Chapuis).

Gonystylus confusus

Coccotrypes medius (Eggers).

Gonystylus punctatus

Platypus titiensis Roberts.

Gonystylus spp.

Genyocerus sexporus (Schedl), *Platypus magnus* Browne, *Platypus pseudocupulatus* Schedl, *Platypus westwoodi* Chapuis, *Xylosandrus ursulus* (Eggers).

Gordonia zeylanica

Crossotarsus terminatus Chapuis, *Platypus furcatus* Blandford.

Gossampinus heptaphylla

Xyleborus piceus (Motschulsky).

Gossweilerodendron balsamiferum

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schauffuss), *Cylindropalpus laudatus* (Schedl), *Doliopygus conradti* (Strohmeier), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus expletus* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl,

Doliopygus medius Schedl, *Doliopygus megatoma* Schedl, *Doliopygus perbreris* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeier), *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeier), *Doliopygus unicornis* Schedl, *Perionunatus grandis* Schedl, *Platypus parallelus* (Fabricius), *Platypus schenklingi* (Strohmeier), *Strombophorus ericis* (Schauffuss), *Trachyostus aterrimus* (Schauffuss), *Trachyostus schaufussi* (Strohmeier), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus eichhoffianus* Schedl, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Gossweilerodendron spp.

Ambrosiodinus eichhoffi (Schreiner), *Ambrosiodinus obliquus* (LeConte), *Ambrosiodinus tropicus* (Hagedorn), *Diamerus impar* Chapuis.

Gossypiospermum praecox

Platypus pulchellus Chapuis.

Gossypium hirsutum

Ambrosiodinus eichhoffi (Schreiner), *Ambrosiodinus tropicus* (Hagedorn), *Hypothenemus carbonarius* (Eggers), *Hypothenemus eruditus* Westwood, *Hypothenemus javanus* (Eggers), *Premnobius ambitiosus* (Schauffuss), *Premnobius circumspinosus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus sub-tuberculatus* Eggers, *Xyleborus volutus* (Fabricius).

Gossypium spp.

Hypothenemus eruditus Westwood, *Hypothenemus gossypii* (Hopkins), *Pseudotlysanoes excavatus* (Wood), *Pseudotlysanoes obesus* (Wood), *Xylosandrus compactus* (Eichhoff).

Gossypium tomentosum

Hypothenemus eruditus Westwood.

Gouania glandulosa

Eucallacea xanthopus (Eichhoff), *Xyleborinus spiculatus* (Schauffuss), *Xyleborinus spinosus* (Schauffuss), *Xyleborus alluaudi* Schauffuss, *Xyleborus madagascariensis* Schauffuss.

Gouldia spp.

Xylosandrus compactus (Eichhoff).

Graptophyllum spp.

Xylosandrus compactus (Eichhoff).

Grevillea pierrard

Ambrosiodinus albizzianus (Schedl).

Grevillea robusta

Ambrosiodinus albizzianus (Schedl), *Cryphalus asperidus* Schedl, *Cryphalus puberulus* Schedl, *Cryphalus subcompactus* Lea, *Diaprus pusillinus* Chapuis, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus excavatus* (Sampson), *Eucallacea fornicatus* (Eichhoff), *Platypus solutus* Schedl, *Xyleborus perforans* (Wollaston), *Xyleborus principalis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).

Grevillea spp.

Ambrosiodinus aegir (Eggers), *Ambrosiodinus obliquus* (LeConte), *Chaetastus montanus* Schedl, *Doliopygus citri* Schedl, *Doliopygus erichsoni* (Chapuis), *Scolytotplatypus fasciatus* Hagedorn, *Triozastus marshalli* (Sampson), *Xylosandrus brevis* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).

Grewia asiatica

Hypothenemus brunneus (Hopkins), *Hypothenemus crudiae* (Panzer).

Grewia laerigata

Xylosandrus morigerus (Blandford).

Grewia latifolia

Hypothenemus cruditus Westwood, *Xylosandrus ater* (Eggers).

Grewia mildbraedii

Chaetastus tuberculatus (Chapuis), *Doliopygus expletus* Schedl, *Euwallacea xanthopus* (Eichhoff), *Perionimatus grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus solutus* Schedl, *Platypus spinulosus* (Strohmeyer), *Polygraphus dimorphus* Schedl, *Triozastus marshalli* (Sampson), *Xyleborus principalis* Eichhoff.

Grewia paniculata

Cnestus aterrimus (Eggers), *Xylosandrus manicus* (Blandford).

***Grewia* spp.**

Arxyleborus suturalis (Eggers), *Crossotarsus fractus* Sampson, *Crossotarsus inermis* Schedl, *Crossotarsus squamulatus* Chapuis, *Eccoptopterus limbus* Sampson, *Scolytoplatypus parvus* Sampson, *Xyleborus perforans* (Wollaston), *Xylosandrus manicus* (Blandford).

Grewia tiliaefolia

Coccotrypes nubilus (Blandford), *Euwallacea andamanensis* (Blandford), *Xyleborus perforans* (Wollaston).

Grewia tomentosa

Xyleborus javanus Eggers.

Grewia vestita

Coccotrypes nubilus (Blandford), *Xyleborus perforans* (Wollaston).

Guarea quara

Hypothenemus seriatus (Eichhoff).

***Guapira* spp.**

Dendrosinus mexicanus Wood.

Guarea cedrata

Chaetastus tuberculatus (Chapuis), *Cryphalus bellus* Schedl, *Cylindropalpus auricomans* (Schaufuss), *Cylindropalpus laudatus* (Schedl), *Doliopygus artespianus* (Schedl), *Doliopygus brevis* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus juronii* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Euwallacea granosus* (Schedl), *Hypothenemus cruditus* Westwood, *Perionimatus excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Platypus solutus* Schedl, *Platypus spinulosus* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus althaudi* Schaufuss, *Xyleborus ambasicusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus sub-tuberculatus* Eggers, *Xyleborus colvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Guarea laurentii

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Cylindropalpus laudatus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus dicaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus lecontei* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus proximus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus spectabilis* Schedl, *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Euwallacea xanthopus* (Eichhoff), *Microrhynchus congonus* Eggers, *Perionimatus grandis* Schedl, *Platypus hintzi* Schaufuss,

Platypus impressus (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus schencklingi* (Strohmeyer), *Platypus solutus* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus althaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus colvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

***Guarea* spp.**

Premnobius cavipennis Eichhoff.

Guarea thompsonii

Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus coeloccephalus* (Schaufuss), *Doliopygus dicaricus* (Schedl), *Doliopygus erichsoni* (Chapuis), *Doliopygus lebrunii* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minutissimus* (Schedl), *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus stauri* Schedl, *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Platypus hintzi* Schaufuss, *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus althaudi* Schaufuss, *Xyleborus ambasicusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus colvulus* (Fabricius).

Guarea trichiloides

Hypothenemus trinitatis (Hopkins), *Pycnarthrum fulgidum* Wood.

***Gnatteria* spp.**

Coptoborus tolimanus (Eggers), *Hypothenemus bolivianus* (Eggers), *Taurodemus splendidus* (Schaufuss).

Gnazuma ulmifolia

Cnesinus punctatus Blandford, *Coptoborus catulus* (Blandford), *Microcortylus inermis* Wood.

***Gnattarda* spp.**

Xyleborus biconicus Eggers.

Guerina abelana

Pseudothyasanoes guerinae (Schedl).

Guibourtia arnoldiana

Enucladus guiboutiae (Schedl), *Microrhynchus guiboutiae* Schedl, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius).

***Guibourtia* spp.**

Doliopygus dentiventris Browne.

Guizotia schultzei

Euwallacea xanthopus (Eichhoff).

Gulbertiodendron deverei

Hypothenemus perluspidus (Eggers).

***Guttiferae* spp.**

Xyleborus vismieae Wood.

***Gymnarion* spp.**

Xyleborus affinis Eichhoff.

Gymnosporia luteola

Traglostus exornatus Schedl.

***Gymnosporia* spp.**

Platypus orientalis (Strohmeyer), *Scolytoplatypus fasciatus* Hagedorn, *Scolytoplatypus kienensis* Schedl.

Gynotroches ouillarisi

Xylosandrus morigerus (Blandford).

***Ilaesia* spp.**

Xylosandrus compactus (Eichhoff).

Ilaesia abyssinica

Cyclorhipidion agnaticeps (Schedl), *Euwallacea xanthopus* (Eichhoff), *Hypothenemus longipennis* (Eggers),

Platypus impressus (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Xyleborus ruidae* Schedl.

Agania spp.

Acanthotomicus curvidens (Schedl), *Ambrosiodinus aegir* (Eggers), *Platypus orientalis* (Strohmeyer).

Hakea sericea

Carchesiopygus dentipennis Schedl, *Hypocryphalus moorei* Schedl.

Hamamelis japonica

Indocryphalus pubipennis (Blandford), *Xyleborinus attenuatus* (Blandford).

Hamamelis spp.

Hypothenemus eruditus Westwood, *Pityophthorus lautus* Eichhoff, *Pseudopityophthorus minutissimus* (Zimmermann), *Xylosandrus brevis* (Eichhoff).

Hamamelis virginiana

Hypothenemus eruditus Westwood, *Lymantor decipiens* (LeConte), *Micracisella opacicollis* (LeConte).

Hannoa klaineana

Ambrosiodinus opacithorax (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatous longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus internedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Haplolobus spp.

Diapus nanus Schedl.

Hardwickia pinnata

Coccotrypes cardamomi Schaufuss.

Haronga paucilata

Acanthotomicus harongae (Schedl), *Acanthotomicus kivuensis* (Schedl), *Doliopygus conradti* (Strohmeyer), *Phrixosoma nigra* (Eggers), *Platypus orientalis* (Strohmeyer), *Xyleborinus collarti* (Eggers), *Xylosandrus crassiusculus* (Motschulsky).

Harpullia pedicellaris

Protoplatypus vetulus Wood.

Harpullia spp.

Ambrosiodinus incertus (Schedl), *Xyleborus venustulus* Schedl.

Harungana madagascariensis

Hypothenemus plumieriae (Nordlinger), *Phrixosoma fuscovillosa* (Schedl), *Pseudomicracis harunganae* (Schedl), *Xylosandrus hirsutipennis* (Schedl).

Heathi spp.

Hypothenemus eruditus Westwood.

Hechtia podantha

Chramesus annectens (Wood).

Hedera colchica

Kissophagus novaki Reitter.

Hedera helix

Kissophagus lederae (Schmidt), *Kissophagus novaki* Reitter, *Kissophagus nuesslini* Reitter, *Scolytogenes indicus* Wood, *Scolytotplatypus siomio* Blandford.

Hedera spp.

Kissophagus erinacellus Wichmann.

Helianthus spp.

Hypothenemus eruditus Westwood.

Helichrysum pandurctum

Cyrtogenius mulungensis (Schedl).

Heritiera fomes

Coccotrypes litoralis (Beeson), *Coccotrypes salakensis* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus squamulatus* Chapuis, *Eucallacea bicolor* (Blandford), *Platypus biuncus* Blandford, *Platypus uncinatus* Blandford, *Xyleborinus artestriatus* (Eichhoff), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Heritiera javanica

Crossotarsus wallacei (Thomson), *Eccoptopterus limbus* Sampson, *Xyleborus tunggali* Schedl.

Heritiera littoralis

Xyleborus similis Ferrari.

Heritiera spp.

Scolytotplatypus parvus Sampson.

Heritiera trifolia

Crossotarsus vafer Schedl, *Platypus chevrolati* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus ustus* Schedl, *Treptotplatypus multiporus* Schedl, *Xyleborus cyclops* Schedl.

Hevea brasiliensis

Ambrosiodinus obliquecaudata Motschulsky, *Chaetastus tuberculatus* (Chapuis), *Coptoborus pseudotenuis* (Schedl), *Coptoborus vespatoriinus* (Schedl), *Coptodryas corporaali* (Eggers), *Crossotarsus minax* (Walker), *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Doliopygus conradti* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Eccoptopterus spinosus* (Olivier), *Eucallacea andamanensis* (Blandford), *Eucallacea comptus* (Sampson), *Eucallacea fornicatus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus biserialis* (Eggers), *Hypothenemus eruditus* Westwood, *Hypothenemus javanus* (Eggers), *Periommatous longicollis* Chapuis, *Platypus cupulatus* Chapuis, *Platypus hintzi* Schaufuss, *Platypus insularis* Strohmeyer, *Platypus jansoni* Chapuis, *Platypus lepidus* Chapuis, *Platypus liratus* Blandford, *Platypus parallelus* (Fabricius), *Platypus pseudocupulatus* Schedl, *Platypus quinquecostatus* Chapuis, *Platypus schysi* Chapuis, *Platypus signatus* Chapuis, *Platypus solidus* Walker, *Platypus westwoodi* Chapuis, *Premnobius quadrispinosus* Schedl, *Premnobius robustulus* (Schedl), *Tesserocerus dewalquei* Chapuis, *Theoborus ricini* (Eggers), *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus exiguus* (Walker), *Xyleborinus heveae* (Schedl), *Xyleborinus polyalthiae* (Schedl), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambastusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus ficus* Eggers, *Xyleborus horridus* Eichhoff, *Xyleborus imbellis* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff, *Xyleborus setulosus* Eggers, *Xyleborus similis* Ferrari, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus morigerus* (Blandford).

Hevea spp.

Ambrosiodinus albizzianus (Schedl), *Coptodryas nuxax* (Schedl), *Crossotarsus javanus* Beeson, *Hypothenemus canerinus* (Eggers), *Hypothenemus eruditus* Westwood, *Platypus biuncus* Blandford, *Platypus cavus* Strohmeyer, *Platypus tragus* Schedl, *Premnobius cavipennis* Eichhoff, *Xyleborus similis* Ferrari, *Xylosandrus discolor* (Blandford).

Hexalobus africanus*Doliopygus brevis* (Strohmeyer).***Hexalobus crispiflorus****Platypus parallelus* (Fabricius).***Hibiscus macrophyllus****Erniocladus corpulentus* (Sampson), *Eucallacea xanthopus* (Eichhoff), *Margadillius minor* Schedl, *Xylosandrus manicus* (Blandford).***Hibiscus rosa-sinensis****Hypothenemus eruditus* Westwood.***Hibiscus spp.****Margadillius parvulus* Eggers, *Platypus lepidus* Chapuis, *Platypus pseudocupulatus* Schedl, *Xylosandrus compactus* (Eichhoff).***Hibiscus syriacus****Ernioporus caucasicus* (Lindemann), *Ernoporus tiliae* (Panzer).***Hibiscus tiliaceus****Cryphalus major* Stebbing, *Cryphalus neglectus* Schedl, *Hypocryphalus corpulentus* Schedl, *Hypothenemus diptercarpi* Hopkins, *Ptilopodius marquesanus* Beeson, *Ptilopodius ramosus* Beeson, *Xyleborus ferrugineus* (Fabricius).***Hildegardia burteri****Chaetastus tuberculatus* (Chapuis).***Hippocratea spp.****Hypothenemus eruditus* Westwood.***Hippomane mancinella****Cnemionyx opacus* Wood, *Cnemionyx splendens* (Wood).***Hirtella glandulosa****Dryococtoides pseudosolitarinus* (Eggers).***Hirtella spp.****Dryococtoides rusticus* Wood.***Hirtella triunda****Taurodenus bicornutus* (Wood).***Histiogaster carpio****Xyleborinus saxeseni* (Ratzeburg).***Holigarna arnottiana****Eucallacea andamanensis* (Blandford), *Xyleborus perforans* (Wollaston), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus terminatus* (Eggers).***Holoptelea grandis****Doliopygus medius* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer), *Xyleborus alluaudi* Schaufuss.***Holoptelea integrifolia****Platypus solidus* Walker.***Homalanthus spp.****Leptoxyleborus concisus* (Blandford), *Xyleborus ferrugineus* (Fabricius).***Homalium foetidum****Xyleborus emarginatus* Eichhoff.***Homalium letestui****Chaetastus tuberculatus* (Chapuis).***Homalium planiflorum****Xyleborus antaisaka* Schedl.***Homalium spp.****Coccotrypes rotundicollis* (Eggers), *Doliopygus conradi* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus intermedius* (Schedl).***Hoepa beccariana****Webbia cornutus* Schedl, *Webbia duodecimsinatus* Schedl, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus manicus* (Blandford).***Hoepa ferrea****Arixyleborus scabripennis* (Blandford), *Coptodryas perparta* (Sampson), *Cryptoxyleborus nanus* Browne, *Eucallacea linatus* (Schedl), *Genyocerus serratus* (Schedl), *Hadrodemius comans* (Sampson), *Leptoxyleborus concisus* (Blandford), *Webbia bituberculatus* Browne, *Webbia cornutus* Schedl, *Webbia duodecimsinatus* Schedl, *Xyleborus minutus* Blandford, *Xyleborus pseudopilifer* Schedl, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus manicus* (Blandford).***Hoepa mengarawan****Platypus protenus* (Schedl).***Hoepa odorata****Arixyleborus medius* (Eggers), *Eccoptopterus limbis* Sampson, *Genyocerus biporus* (Schedl), *Genyocerus diaphanus* (Schedl), *Hadrodemius comans* (Sampson), *Platypus curtus* Chapuis, *Platypus shoreanus* (Beeson), *Webbia cornutus* Schedl, *Webbia trigintispinatus* Sampson, *Xylosandrus crassiusculus* (Motschulsky).***Hoepa parviflora****Ambrosiodmus lewisi* (Blandford), *Sphaerotrypes subtectus* Browne, *Xylosandrus compactus* (Eichhoff).***Hoepa spp.****Cnestus suturalis* (Eggers), *Diapus quinquespinatus* Chapuis, *Genyocerus puer* (Schedl), *Platypus bifureus* (Schedl), *Schedlia sumatrana* (Hagedorn), *Sphaerotrypes philippinensis* Strohmeyer, *Taphrodasus divisis* (Browne), *Webbia diptercarpi* Hopkins, *Webbia quatuordecimsinatus* Sampson, *Webbia suturalis* Browne, *Xyleborinus perminutissimus* (Schedl), *Xyleborus anplexicauda* Hagedorn.***Hoepa wightiana****Eucallacea andamanensis* (Blandford), *Sphaerotrypes subtectus* Browne, *Xyleborus perforans* (Wollaston), *Xylosandrus crassiusculus* (Motschulsky).***Horsfieldia glabra****Crossotarsus horni* Schedl, *Diapus quinquespinatus* Chapuis, *Platypus bajulus* Schedl.***Horsfieldia irya****Platypus schysi* Chapuis.***Hoslundia opposita****Eucallacea xanthopus* (Eichhoff).***Hoslundia spp.****Ambrosiodmus obliquus* (LeConte).***Hovea spp.****Ambrosiodmus rubricollis* (Eichhoff).***Hovenia dulcis****Hypothenemus brunneus* (Hopkins).***Hovea belmoreana****Coccotrypes dactyliperda* (Fabricius).***Hullettia dumosa****Xylosandrus manicus* (Blandford).***Humiristrum exselsum****Dryococtoides insculptus* Wood.***Hura polyandra****Cnemionyx equilmui* Wood.***Hydnocarpus spp.****Xyleborus perforans* (Wollaston).***Hydnocarpus wightiana****Xyleborus perforans* (Wollaston).***Hydodendron gabunense****Chaetastus tuberculatus* (Chapuis), *Hypothenemus biseriatus* (Eggers), *Mesoplatypus venustus* Schedl, *Premnobius quadrispinosus* Schedl, *Xyleborus ferrugineus* (Fabricius).

Hymenaea courbaril

Araptus hymenaeae (Eggers), *Crossotarsus schedli* Brownie, *Hypothenemus obscurus* (Fabricius).

Hymenocardia capensis

Laurigus bicolor Schedl.

Hymenocardia uluoides

Doliopygus conradti (Strohmeyer), *Doliopygus notatus* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus submarginatus* Schedl, *Periommatius grandis* Schedl, *Xyleborus annectens* Schedl.

Hymenodictyon excelsum

Coptodryas recidens (Sampson), *Eucallacca interjectus* (Blandford), *Platypus secretus* Sampson, *Platypus suffodiens* Sampson, *Xyleborus pumilus* Eggers.

Hymenostegia afzelii

Doliopygus brevis (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus exilis* (Chapuis), *Doliopygus opifex* (Sampson), *Doliopygus perbrevis* Schedl, *Doliopygus propinquus* Schedl, *Dryocoetoides cristatus* (Fabricius), *Periommatius longicollis* Chapuis, *Stephanopodius ghanaensis* (Schedl), *Trachyostus aterrimus* (Schaufuss), *Xyleborus alluaudi* Schaufuss.

***Hymenostegia* spp.**

Prennobilis caripennis Eichhoff.

Hypericum lanceolatum

Cyclorhpidion agnaticops (Schedl), *Pseudomicracis bugekae* (Schedl), *Xyleborus amanicus* Hagedorn, *Xyleborus integer* Schedl, *Xyleborus principalis* Eichhoff.

Hyphaene guineensis

Coccotrypes carpophagus (Hornung).

***Hyphaene* spp.**

Coccotrypes dactyliperda (Fabricius).

***Hyphorbe* spp.**

Coccotrypes dactyliperda (Fabricius).

***Hypodaphnis* spp.**

Periommatius excisus Strohmeyer.

***Hyracraea* spp.**

Hypothenemus seriatus (Eichhoff).

***Ibuga* spp.**

Xyleborus amanicus Hagedorn.

Ichthyomethia communis

Ambrosiodinus lecontei Hopkins, *Hypothenemus brunneus* (Hopkins), *Hypothenemus columbi* Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus squamosus* (Hopkins).

Icica altissima

Acanthotomicus bidentis Wood, *Acanthotomicus ocularis* (Wood), *Xyleborinus celatus* Wood, *Xyleborus affinis* Eichhoff.

Ilex anomala

Xyleborinus saxseni (Ratzeburg), *Xyleborus ferrugineus* (Fabricius), *Xyleborus molokaisensis* Perkins, *Xyleborus obliquus* Sharp.

Ilex chinensis

Platypus calamus Blandford, *Platypus quercivorus* Murayama.

Ilex dipyrrena

Crossotarsus cinnamomi Beeson, *Scolytoplatus euto-moides* Blandford.

Ilex macropoda

Coptodryas kirishimanus (Murayama).

Ilex mitis

Chaetastus montanus Schedl.

Ilex oldhami

Crossotarsus simplex Murayama, *Xyleborus osunensis* Murayama.

***Ilex* spp.**

Ambrosiodinus apicalis (Blandford), *Ambrosiodinus rubricollis* (Eichhoff), *Cryphalus sylvicola* (Perkins), *Diapus nebulosus* Roberts, *Eucallacca calidus* (Eichhoff), *Platypus calamus* Blandford, *Xyleborus dispar* (Fabricius), *Xylosandrus compactus* (Eichhoff).

Illicium religiosum

Platypus calamus Blandford, *Polygraphus magnus* Murayama, *Xyleborus octiesdentatus* Murayama, *Xyleborus ohtensis* Nobuchi.

Indigofera arrecta

Hypothenemus grandis Schedl, *Xylosandrus compactus* (Eichhoff).

Indigofera galegoides

Hypothenemus eruditus Westwood.

***Indigofera* spp.**

Xylosandrus compactus (Eichhoff).

Indigofera suffruticosa

Xylosandrus compactus (Eichhoff).

Inga alba

Xyleborus volutus (Fabricius).

Inga laurina

Monarthrum praecustum (Eggers), *Xyleborus affinis* Eichhoff.

Inga marginata

Xyleborus ferrugineus (Fabricius).

Inga paterno

Chramesus ritiosus Wood, *Chramesus xylophagus* Wood, *Hylocurus inaequalis* Wood, *Scolytus laetus* Wood, *Scolytus propinquus* Blandford.

***Inga* spp.**

Chramesus vastus Wood, *Cnesinus elegans* Blandford, *Corthyllocinus mexicanus* (Schedl), *Corthylus ingaensis* (Schedl), *Corthylus insignis* Wood, *Corthylus serratus* Wood, *Corthylus robustus* Schedl, *Dryocoetoides capucinus* (Eichhoff), *Dryocoetoides monachus* (Blandford), *Dryocoetoides pileatus* Wood, *Dryocoetoides verrucosus* Wood, *Gnathotrupes bituberculatus* (Blandford), *Hylocurus elegans* Eichhoff, *Hylocurus simplex* Blandford, *Hylocurus verrucosus* Wood, *Hypothenemus erudiae* (Panzer), *Hypothenemus erectus* LeConte, *Hypothenemus opacus* (Eichhoff), *Hypothenemus solocis* Wood, *Hypothenemus suspectus* Wood, *Hypothenemus teretis* Wood, *Micracis lepidus* Wood, *Micracis suainei* Blackman, *Monarthrum notatum* Wood, *Monarthrum proprium* Wood, *Phloeoborus punctatorugosus* Chapuis, *Phloeotribus ingae* Wood, *Phloeotribus squamiger* Wood, *Platypus chiriquensis* Wood, *Scolytodes ingatoris* Wood, *Taurodennus bicornutus* (Wood), *Taurodennus varius* (Wood), *Theoborus villosulus* (Blandford), *Xyleborinus celatus* Wood, *Xyleborus ferrugineus* (Fabricius), *Xyleborus squamulatus* Eichhoff, *Xylosandrus compactus* (Eichhoff), *Xylosandrus laticeps* (Wood).

Inga vera

Chramesus rotundatus (Chapuis), *Dryocoetoides pseudosolitaris* (Eggers), *Platypus dentatus* Dalman, *Platypus excisus* Chapuis, *Xyleborus affinis* Eichhoff.

Inocarpus edulis

Crossotarsus externedentatus (Fairmaire), *Xyleborus volutus* (Fabricius).

***Inocarpus* spp.**

Crossotarsus externedentatus (Fairmaire), *Xylosandrus compactus* (Eichhoff).

Ipomoea cathartica

Hypothenemus californicus Hopkins, *Hypothenemus*

- cruditus* Westwood, *Hypothenemus hirsutus* (Wood), *Hypothenemus seriatus* (Eichhoff).
- Ipomoea littoralis***
Hypothenemus californicus Hopkins.
- Ipomoea pres-caprae***
Cryptocarenum seriatus Eggers, *Scolytogenes knabi* (Hopkins).
- Ipomoea* spp.**
Scolytogenes hirtus (Wood), *Scolytogenes rusticus* (Wood), *Scolytogenes truncis* (Wood).
- Ipomoea turpethum***
Scolytogenes aspericollis (Eichhoff).
- Irauma xanthophyllum***
Xyleborus xanthophyllus Schedl.
- Irringia gabouensis***
Chaetastus tuberculatus (Chapuis), *Cylindropalpus pumilio* (Schedl), *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff.
- Irringia grandifolia***
Ambrosiodmus aegir (Eggers), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus lefevrei* Schedl, *Doliopygus serratus* (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).
- Isoloma bruneelii***
Doliopygus crichsoni (Chapuis), *Doliopygus tenuis* (Strohmeyer), *Eucallacea xanthopus* (Eichhoff), *Periommatus similis* Strohmeyer, *Xyleborinus polyalthiae* (Schedl), *Xyleborus affinis* Eichhoff.
- Isoloma* spp.**
Prennobiobus cavipennis Eichhoff.
- Isonandra polyantha***
Eucallacea bicolor (Blandford), *Scolytomimus assamensis* Schedl.
- Isonandra* spp.**
Xyleborinus andrewesi (Blandford).
- Ira imbricata***
Hypothenemus californicus Hopkins, *Hypothenemus gossypii* (Hopkins).
- Ixonanthes reticulata***
Crossotarsus wallacei (Thomson), *Platypus cupulatus* Chapuis.
- Ixora parviflora***
Eucallacea andamanensis (Blandford).
- Jacaranda copaia***
Platypus dentatus Dahman, *Xyleborus semipunctatus* Eggers, *Xyleborus volutus* (Fabricius).
- Jacaranda* spp.**
Prennobiobus cavipennis Eichhoff, *Xylosandrus laticeps* (Wood).
- Jacksouia* spp.**
Xyleborus perforans (Wollaston).
- Jacobiua* spp.**
Xylosandrus compactus (Eichhoff).
- Jambosa malaccensis***
Crossotarsus externedentatus (Fairmaire).
- Jasminium abyssinicum***
Dendrochilus jasminiae Schedl.
- Jasminium* spp.**
Xylosandrus compactus (Eichhoff).
- Jatropha cinerea***
Liparthrum albosetum Bright.
- Jatropha curcas***
Xyleborus perforans (Wollaston).
- Jatropha kebella***
Xyleborus perforans (Wollaston).
- Jatropha spathulata***
Dendroterus texanus Wood.
- Jatropha* spp.**
Hypothenemus macrolobii (Eggers).
- Juglans californica***
Pityophthorus juglandis Blackman.
- Juglans cinerea***
Xyleborus ferrugineus (Fabricius).
- Juglans major***
Pityophthorus juglandis Blackman.
- Juglans mandshurica***
Cryphalus mandshuricus Eggers, *Hylesinus eos* Spessivtsev, *Xyleborus aequalis* (Reitter).
- Juglans mandshurica sieboldiana***
Platypus modestus Blandford.
- Juglans nigra***
Ambrosiodmus rubricollis (Eichhoff), *Hylesinus crenatus* (Fabricius), *Hypothenemus bezaziani* Peyerimhoff, *Hypothenemus crudiac* (Panzer), *Hypothenemus cruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Pityophthorus juglandis* Blackman, *Pityophthorus laetus* Eichhoff, *Pseudothysanoes lecontei* Blackman.
- Juglans regia***
Ambrosiodmus rubricollis (Eichhoff), *Coccotrypes nubilus* (Blandford), *Crossotarsus fairmairei* Chapuis, *Dryocotes himalayensis* Strohmeyer, *Scolytus nitidus* Schedl, *Scolytus scolytus* (Fabricius), *Sphaerotrypes juglans* Tsai & Yin, *Trypodendron domesticum* (Linnaeus), *Xyleborinus artetriatus* (Eichhoff), *Xyleborinus subgranulatus* (Eggers), *Xylosandrus discolor* (Blandford).
- Juglans sieboldiana***
Cryphalus juglans Nüßina, *Hylesinus laticollis* Blandford.
- Juglans* spp.**
Ambrosiodmus apicalis (Blandford), *Ambrosiodmus tachygraphus* (Zimmermann), *Hylesinus cingulatus* Blandford, *Hylesinus varius* (Fabricius), *Hypothenemus cruditus* Westwood, *Eucallacea validus* (Eichhoff), *Platypus quercuslandi* Schedl, *Scolytoplatus tyeon* Blandford, *Sphaerotrypes juglans* Krivolutskaia & Kupyanskaya, *Xyleborus ferrugineus* (Fabricius), *Xyleborus sayi* (Hopkins), *Xylosandrus germanus* (Blandford).
- Julbernardia brieji***
Cylindropalpus auricomans (Schaufuss), *Doliopygus divaricus* (Schedl).
- Julbernardia paniculata***
Glostatus multipinosus (Schedl), *Polygraphus zambianus* Beaver.
- Julbernardia seretii***
Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Cylindropalpus interpositus* (Schedl), *Doliopygus conjunctus* Schedl, *Doliopygus ghesquierei* (Schedl), *Doliopygus lefevrei* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus pygmaeolus* Schedl, *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicoloris* Schedl, *Trachyostus aterrimus* (Schaufuss), *Triozastus marslalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).
- Juniperus chinensis***
Phloeosinus hopchi Schedl, *Phloeosinus rudis* Blandford, *Phloeosinus shensi* Tsai & Yin.

Juniperus communis*Phloeosinus turkestanicus* Semenov.***Juniperus deppeana****Carphobius arizonicus* Blackman, *Phloeosinus delconii* Blackman, *Phloeosinus furnissi* Blackman, *Phloeosinus hoferi* Blackman, *Phloeosinus scopulorum neomexicanus* Blackman, *Phloeosinus serratus* (LeConte), *Phloeosinus spinosus* Blackman.***Juniperus excelsa****Phloeosinus acatayi* Schedl.***Juniperus flaccida****Phloeosinus delconii* Blackman.***Juniperus monosperma****Phloeosinus furnissi* Blackman, *Phloeosinus scopulorum neomexicanus* Blackman, *Phloeosinus serratus* (LeConte).***Juniperus occidentalis****Phloeosinus hoppingi* Swaine, *Phloeosinus punctatus* LeConte, *Phloeosinus serratus* (LeConte), *Phloeosinus swaini* Bruck.***Juniperus osteosperma****Phloeosinus furnissi* Blackman, *Phloeosinus hoferi* Blackman, *Phloeosinus scopulorum neomexicanus* Blackman, *Phloeosinus serratus* (LeConte).***Juniperus procera****Glostatus acaciae* (Schedl).***Juniperus pseudosabina****Phloeosinus turkestanicus* Semenov.***Juniperus scopulorum****Phloeosinus hoferi* Blackman, *Phloeosinus hoppingi* Swaine, *Phloeosinus keeni* Blackman, *Phloeosinus scopulorum scopulorum* Swaine, *Phloeosinus serratus* (LeConte).***Juniperus* spp.***Phloeosinus bicolor* (Brulle), *Phloeosinus cristatus* (LeConte), *Phloeosinus delconii* Blackman, *Phloeosinus henschii* Reitter, *Phloeosinus palearis* Wood, *Phloeosinus perlatus* Chapuis, *Phloeosinus serratus* (LeConte), *Phloeosinus thujae* (Perris).***Juniperus virginiana****Phloeosinus dentatus* (Say).***Kairiballi* spp.***Xyleborus parcellus* Wood.***Kalmia latifolia****Xyleborus sayi* (Hopkins).***Kalopanax ricinifolium****Emporicus senecioides* (Kurenzov), *Platypus lewisi* Blandford, *Xyleborus orientalis* Eggers.***Kalopanax septenlobum****Crossotarsus niponicus* Blandford, *Platypus lewisi* Blandford, *Xyleborus seriatus* Blandford.***Kaya floribunda****Xylosandrus crassiusculus* (Motschulsky).***Khaya anthotheca****Cylindropalpus granulosus* (Schedl), *Doliopygus perbrevis* Schedl, *Rhopalopselion immaturus* (Schedl), *Xyleborus consobrinus* Eggers.***Khaya grandifolia****Chaetastus tuberculatus* (Chapuis), *Polygraphus coronatus* Eggers, *Xyleborus affinis* Eichhoff.***Khaya ivorensis****Chaetastus tuberculatus* (Chapuis), *Diapus quinque-spinatus* Chapuis, *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus*(Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus scobinatus* Hagedorn, *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).***Khaya klainea****Platypus hintzi* Schaufuss.***Khaya nyasica****Eucallacea khayae* (Schedl).***Khaya senegalensis****Pityophthorus fallax* (Hagedorn), *Xylosandrus manicus* (Blandford).***Khaya* spp.***Ambrosiodinus eichhoffi* (Schreiner), *Chaetastus tuberculatus* (Chapuis), *Cyclorhpidion crucipenne* (Schedl), *Doliopygus artespiniatus* (Schedl), *Doliopygus perbrevis* Schedl, *Doliopygus unicornis* Schedl, *Pityophthorus hintzi* Schedl, *Platypus hintzi* Schaufuss, *Polygraphus bicolor* Eggers, *Polygraphus granulosus* Eggers, *Polygraphus granulifer* Eggers, *Premnobius cavipennis* Eichhoff, *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus principalis* Eichhoff, *Xyleborus volutus* (Fabricius).***Kickxia arborea****Platypus signatus* Chapuis.***Kigelia lanceolata****Platypus solutus* Schedl, *Rhopalopselion congoum* (Schedl).***Klaineana* spp.***Ambrosiodinus eichhoffi* (Schreiner).***Klainedoxa gabonensis****Acanthotomicus sexdentatus* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Coccotrypes rotundicollis* (Eggers), *Doliopygus artespiniatus* (Schedl), *Doliopygus bilobatus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus ghesquierei* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus spectabilis* Schedl, *Doliopygus unicornis* Schedl, *Hypothenemus eruditus* Westwood, *Micriophthalus klainedoxae* Schedl, *Periommatius excisus* Strohmeyer, *Periommatius longicollis* Chapuis, *Pityophthorus africanus* Wood, *Pityophthorus africanus* Eggers, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Trachytus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus caudatus* Schedl, *Xyleborus clerodendronae* Schedl, *Xyleborus nitidulus* Eggers, *Xyleborus volutus* (Fabricius).***Klainedoxa gabonensis oblongifolia****Xylosandrus cylindrotomicus* (Schedl).***Klainedoxa* spp.***Premnobius cavipennis* Eichhoff.***Kleinhovia hospita****Hadrodemius globus* (Blandford).***Knightia excelsa****Xyleborinus saxeni* (Ratzeburg).***Koompassia excelsa****Arixyleborus granulifer* (Eggers), *Coptodryas diversicolor* (Eggers), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus inermis* Schedl, *Crossotarsus squamulatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Eccoptopterus limbus* Sampson, *Platypus curtus* Chapuis, *Platypus forficula* Chapuis, *Xyleborus emarginatus* Eichhoff, *Xyleborus fallax* Eichhoff.

Koompassia malaccensis

Ozopemon rugatus (Blandford). *Platypus forficula* Chapuis. *Platypus insularis* Strohmeier. *Platypus signatus* Chapuis. *Platypus westwoodi* Chapuis. *Xyleborus apertus* Schedl. *Xyleborus fallax* Eichhoff. *Xyleborus perforans* (Wollaston). *Xyleborus picus* (Motschulsky). *Xyleborus volutus* (Fabricius).

Koompassia spp.

Arixyleborus granifer (Eichhoff).

Koordersiodendron pinnatum

Platypus laevis Brownie.

Kopsia flavida

Platypus insulindicus Schedl. *Xyleborinus andrewesi* (Blandford). *Xyleborus adusticollis* (Motschulsky).

Kosteletzkyia adoensis

Hypothenemus grandis Schedl.

Kotschya africana

Thammurgus granulicollis Schedl.

Kududarula spp.

Ambrosiodmus obliquecaudata Motschulsky.

Kydia calycina

Ernocladius corpulentus (Sampson). *Eucallacea interjectus* (Blandford). *Xyleborus fallax* Eichhoff. *Xyleborus perforans* (Wollaston). *Xyleborus similis* Ferrari.

Lactuca spp.

Hypothenemus grandis Schedl.

Lagenaria vulgaris

Cosmoderes monilicollis Eichhoff.

Lagerstroemia flos-reginae

Xylosandrus crassiusculus (Motschulsky).

Lagerstroemia hypoleuca

Crossotarsus terminatus Chapuis. *Xyleborus cognatus* Blandford.

Lagerstroemia lanceolata

Crossotarsus minax (Walker). *Sphaerotrypes globulus* Blandford.

Lagerstroemia parviflora

Crossotarsus externedentatus (Fairmaire). *Platypus cavius* Strohmeier. *Sphaerotrypes globulus* Blandford. *Xyleborus perforans* (Wollaston).

Lagerstroemia speciosa

Xylosandrus crassiusculus (Motschulsky).

Lagunaria leucantha

Hypothenemus eruditus Westwood.

Lagunaria patersonia

Platypus norfolkensis Schedl.

Landeria pulcherrima

Coccotrypes nubilus (Blandford).

Landolphia owariensis

Platypus intermedius (Schedl). *Platypus solutus* Schedl.

Landolphia robusta

Ctonoxylon flavescens Hagedorn.

Landolphia spp.

Scolytogenes sodalis (Schedl).

Launea grandis

Coccotrypes nubilus (Blandford). *Crossotarsus externedentatus* (Fairmaire). *Platypus cupulatus* Chapuis. *Platypus latifinis* Walker. *Platypus lepidus* Chapuis. *Platypus secretus* Sampson. *Platypus solidus* Walker. *Xyleborinus andrewesi* (Blandford). *Xyleborinus artetriatus* (Eichhoff). *Xyleborus shorae* (Stebbing). *Xylosandrus crassiusculus* (Motschulsky).

Launea wehrschii

Doliopygus serratus (Strohmeier). *Periommatius grandis* Schedl. *Platypus hintzi* Schaufuss. *Platypus parallelus*

(Fabricius). *Xyleborus affinis* Eichhoff. *Xyleborus ferrugineus* (Fabricius). *Xyleborus picus* (Motschulsky).

Launea zenkeri

Platypus hintzi Schaufuss.

Lausium domesticum

Coptodryas nudipennis (Schedl). *Cyrtogenius brevior* (Eggers). *Xyleborinus andrewesi* (Blandford). *Xyleborus adusticollis* (Motschulsky). *Xylosandrus nudipennis* (Schedl).

Lausium spp.

Coptodryas elegans (Sampson).

Lantana salvifolia

Eucallacea xanthopus (Eichhoff). *Xyleborus principalis* Eichhoff.

Lantana spp.

Ambrosiodmus lantanae (Eggers). *Eucallacea formicatus* (Eichhoff). *Xylosandrus compactus* (Eichhoff). *Xylosandrus terminatus* (Eggers).

Laportea spp.

Platypus jansoni Chapuis. *Platypus solidus* Walker.

Larix americana

Hylurgops rugipennis pinifex Fitch.

Larix dahurica

Carphoborus cholodkovskyi Spessivtsev. *Cryphalus latus* Eggers. *Hylurgops spessivtzevi* Eggers. *Polygraphus proximus* Blandford.

Larix dahurica koreana

Dryocoetes infuscatus Murayama. *Polygraphus horyurensis* Murayama.

Larix europaea

Crypturgus hispidulus Thomson. *Hylastes cunicularius* Erichson. *Orthotomicus laricis* (Fabricius). *Pityogenes irkutensis* Eggers. *Pityokteines curvidens* (Germar). *Pityokteines spinidens* (Reitter). *Pityophthorus buyssoni* Reitter. *Pityophthorus glabratus* Eichhoff.

Larix gmelini

Cryphalus latus Eggers. *Cryphalus premayaensis* Murayama. *Polygraphus sachalinensis* Eggers.

Larix griffithiana

Trypodendron dorjitenzingi Schmutzenhofer.

Larix laricina

Carphoborus dunni Swaine. *Dendroctonus simplex* LeConte. *Dendroctonus valens* LeConte. *Pityophthorus consimilis* LeConte. *Pityophthorus opaculus* LeConte. *Polygraphus rufipennis* (Kirby). *Scolytus piccae* (Swaine).

Larix leptolepis

Cryphalus laricis Nisima. *Dryocoetes karamatsu* Sawamoto. *Polygraphus kisoensis* Nisima. *Xyleborus seriatus* Blandford.

Larix lyallii

Pityophthorus alpinensis G. Hopping. *Scolytus laricis* Blackman.

Larix occidentalis

Dendroctonus pseudotsugae Hopkins. *Scolytus laricis* Blackman. *Xylechinus montanus* Blackman.

Larix olgensis coreana

Trypodendron gaimaensis (Murayama).

Larix potanini

Polygraphus chungdianensis Tsai & Yin.

Larix sibirica

Carphoborus teplouchovi Spessivtsev. *Pityophthorus lichtensteini* (Ratzeburg). *Pityophthorus micrographus* (Linnaeus). *Polygraphus polygraphus* (Linnaeus). *Polygraphus seriatus* Reitter. *Scolytus moravitzii* Semenov.

Larix spp.

Cryphalus asperatus (Gyllenhal), *Cryphalus latus* Eggers, *Dendroctonus micans* (Kugelann), *Dryocoetes affaber* (Mannerheim), *Dryocoetes autographus* (Ratzeburg), *Dryocoetes baicalicus* Reitter, *Dryocoetes pini* Nüßima, *Dryocoetes rigicollis* Eggers, *Gnathotrichus materiarius* (Fitch), *Hylastes nigrinus* (Mannerheim), *Hylurgops glabratus* (Zetterstedt), *Hylurgops palliatus* (Gyllenhal), *Ips acuminatus* (Gyllenhal), *Ips cembrae* (Heer), *Ips typographus* (Linnaeus), *Orthotomicus caelatus* (Eichhoff), *Orthotomicus suturalis* (Gyllenhal), *Pityogenes chalcographus* (Linnaeus), *Scolytus moravici* Semenov, *Xyleborinus saxenii* (Ratzeburg), *Xylechinus pilosus* (Ratzeburg).

Larrea tridentata

Chaetophilocus fasciatus (Blackman).

Lasiodiscus spp.

Ambrosiodmus albizzianus (Schedl).

Lauraceae spp.

Xyleborus piccus (Motschulsky).

Laurineae spp.

Platypus puerulus (Schedl).

Laurus alba

Thamurgus kaltenbachii (Bach).

Laurus camphora

Ambrosiodmus camphorae (Hagedorn).

Laurus canariensis

Liparthrum nigrescens Wollaston.

Laurus nobilis

Hypothenemus hystrix (Eggers), *Liparthrum colchicum* Semenov, *Liparthrum laurivorus* Schedl.

Lebrunia sushaie

Chaetastus montanus Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus scobinatus* Hagedorn.

Lecaniodiscus cupanioides

Doliopygus erichsoni (Chapuis), *Doliopygus medius* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Trachyostus schoutedeni* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Lecaniodiscus spp.

Ambrosiodmus eichhoffi (Schreiner), *Prennobia cavipennis* Eichhoff.

Lecythis costaricensis

Phloeoborus punctatorugosus Chapuis.

Lecythis spp.

Phelloterus anaxeus Wood, *Platypus eversus* Wood, *Platypus nudatus* Wood, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).

Leea crispa

Xylosandrus crassiusculus (Motschulsky).

Leea sambucina

Xylosandrus crassiusculus (Motschulsky).

Leea spp.

Xyleborinus andrewesi (Blandford).

Leontotis africana

Eucallacea xanthopus (Eichhoff).

Leptoderris laurentii

Doliopygus conradti (Strohmeyer), *Xyleborus alluaudi* Schaufuss.

Leptolaena multiflora

Ambrosiodmus artemis (Schedl), *Mitosoma lobatum* Schedl, *Mitosoma obtusum* Schedl, *Polygraphus amoenus* (Schaufuss), *Xyleborinus forficuloides* (Schedl).

Leptonychia bantangensis

Xyleborus bantabensis Eggers.

Leptospermum ericoides

Anasa truncatus (Erichson).

Leucaenia glauca

Crossotarsus externedentatus (Fairmaire), *Hypothenemus brunneus* (Hopkins), *Hypothenemus eruditus* Westwood, *Hypothenemus javanus* (Eggers), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Leucaenia leucocephala

Eccoptopterus spinosus (Olivier).

Leucaenia pulverulenta

Chramesus mimosae Blackman, *Hylocurus atkinsoni* Wood.

Leucaenia spp.

Ambrosiodmus wilderi (Beeson), *Eucallacea fornicatus* (Eichhoff), *Xyleborus sweczyi* Beeson, *Xylosandrus compactus* (Eichhoff).

Leucas spp.

Mimicurus congonus (Schedl).

Leucosceptrum canum

Eucallacea velatus (Sampson), *Eucallacea wallacei* (Blandford).

Lezjavai spp.

Hypothenemus eruditus Westwood.

Libocedrus decurrens

Phloeosinus antennatus Swaine, *Phloeosinus fulgens* Swaine, *Phloeosinus hoppingi* Swaine, *Phloeosinus punctatus* LeConte, *Phloeosinus sequoiae* Hopkins, *Phloeosinus vandykei* Swaine.

Licania spp.

Phelloterus anaxeus Wood, *Platypus exitialis* Wood, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).

Ligustrum spp.

Ambrosiodmus rubricollis (Eichhoff), *Laururgus barbatus* Eggers.

Limaciopsis loangensis

Doliopygus artespianus (Schedl).

Lindackeria dentata

Doliopygus bilobatus (Schedl), *Doliopygus conradti* (Strohmeyer).

Lindera benzoin

Xyleborus sayi (Hopkins).

Lindera erythrocarpa

Crossotarsus niponicus Blandford, *Platypus quercivorus* Murayama, *Xylosandrus germanus* (Blandford), *Xylosandrus mutilatus* (Blandford).

Lindera latifolia

Coptodryas recidens (Sampson), *Eucallacea bicolor* (Blandford), *Xyleborus glabratus* Eichhoff.

Lindera pulcherrima

Diapys quadrispinatus Chapuis, *Diapys quinquespinatus* Chapuis, *Diapys spatulifer* Brownie, *Scolytogenes cylonicus* (Schedl).

Lindera spp.

Ambrosiodmus obliquus (LeConte), *Scolytoplatus tycon* Blandford, *Xylosandrus brevis* (Eichhoff), *Xylosandrus mutilatus* (Blandford).

Lindera thunbergii

Crossotarsus simplex Murayama.

Lippia spp.

Ambrosiodmus aegir (Eggers).

Lippia substrigosa*Corthylus mexicanus* Schedl.***Liquidambar orientalis****Platypus simulans* Schedl.***Liquidambar* spp.***Xylosandrus compactus* (Eichhoff).***Liquidambar styraciflua***

Cnecinus strigicollis LeConte, *Dryocoetes betulae* Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus seriatus* (Eichhoff), *Monarthrum fasciation* (Say), *Monarthrum mali* (Fitch), *Pityophthorus liquidambaris* Blackman, *Pityophthorus molestus* Wood, *Xyleborus ferrugineus* (Fabricius).

***Liriodendron* spp.**

Ambrosiodinus tachygraphus (Zimmermann), *Xyleborus dispar* (Fabricius), *Xyleborus obesus* LeConte.

Liriodendron tulipifera

Corthylus columbianus Hopkins, *Trischidias atomus* (Hopkins), *Xyleborus sayi* (Hopkins), *Xylosandrus germanus* (Blandford).

Liriosma singularis

Prennobia cuspideus Eichhoff, *Taurodenus perebeae* (Ferrari), *Xyleborinus reconditus* (Schedl), *Xyleborus rugosipennis* Schedl.

Lithocarpus amygdalifolia*Xyleborus atratus* Eichhoff.***Lithocarpus densiflorus****Monarthrum scutellare* (LeConte).***Lithocarpus edulis****Platypus quercitorius* Murayama.***Lithocarpus konishii****Diapus formosanus* Niisima & Murayama.***Lithocarpus* spp.**

Diapus latespinis Schedl, *Diapus nanus* Schedl, *Diapus perpygmaeus* Roberts, *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Diapus robustus* Schedl, *Platypus noraguineensis* Roberts, *Xyleborus arniger* Schedl, *Xyleborus arnippennis* Schedl.

Lithocarpus stipitata*Xyleborus atratus* Eichhoff.***Lithocarpus ternaticupula****Xyleborus atratus* Eichhoff.***Lithocarpus wallichianus***

Anasa schliehi (Stebbing), *Crassotarsus longicollis* Browne, *Crassotarsus squamulatus* Chapuis, *Cyclorhipidion punctatopilosum* (Schedl), *Platypus pseudocupulatus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Litsea anara

Phloeocranus bruchoides Schedl, *Xylosandrus pygmaeus* (Eggers).

Litsea cassiaefolia*Xylosandrus compactus* (Eichhoff).***Litsea dolichocarpa****Xyleborus atratus* Eichhoff.***Litsea domarensis***

Hyledius nitidicollis (Motschulsky), *Platypus jansoni* Chapuis, *Sucus niisimai* (Eggers).

Litsea elongata

Diapus spatulifer Browne, *Scolytotplatypus pubescens* Hagedorn, *Scolytotplatypus raja* Blandford, *Xyleborus glabratus* Eichhoff.

Litsea megacarpa

Crassotarsus wallacci (Thomson), *Hadrodemus globus* (Blandford), *Platypus pseudocupulatus* Schedl, *Xylosandrus manicus* (Blandford).

Litsea polyantha

Coccotrypes myristicae (Roepke), *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston).

***Litsea* spp.**

Crassotarsus porcatus Roberts, *Diapus amblylaminatus* Roberts, *Diapus bilunatus* Schedl, *Diapus nebulosus* Roberts, *Diapus unispinus* Roberts, *Phloeocranus bruchoides* Schedl, *Scolytotplatypus darjeelingi* Stebbing, *Xyleborus perforans* (Wollaston).

Litsea umbrosa*Xyleborus eggersi* Beeson.***Litsea zeylanica***

Ambrosiodinus asperatus (Blandford), *Platypus furcatus* Blandford.

Liristona cochinchinensis

Coccotrypes barbatus (Schedl), *Coccotrypes nitidus* (Eggers).

Liristona jenkinsiana*Platypus shillongensis* Schedl.***Liristona* spp.**

Coccotrypes carpophagus (Hornung), *Coccotrypes dactylipeda* (Fabricius), *Xyleborus perforans* (Wollaston).

Lobelia bequaertii*Thamniurgus lobeliae* Eggers.***Lobelia gibberoa***

Cyrtogenus novolae (Schedl), *Thamniurgus africanus* Eggers, *Thamniurgus zukwaleae* Scott.

***Lobelia* spp.**

Thamniurgus africanus Eggers, *Thamniurgus punctatissimus* Eggers, *Xyleborus piceus* (Motschulsky).

Lobelia telekii*Thamniurgus lobeliae* Eggers.***Lobelia wollastonii****Thamniurgus lobeliae* Eggers.***Lodoicea callipyge****Xyleborus perforans* (Wollaston).***Lodoicea sechellarum***

Xyleborus ferrugineus (Fabricius), *Xyleborus perforans* (Wollaston).

Lomatia hirsuta*Xylechinus spathifer* Schedl.***Lonchocarpus capassa****Liparthrum australis* Schedl.***Lonchocarpus castilloi****Chramesus crenatus* Wood, *Cusinus strigicollis* LeConte.***Lonchocarpus constrictus****Chramesus vitiosus* Wood.***Lonchocarpus guatemalensis****Chramesus vitiosus* Wood.***Lonchocarpus margaritensis***

Scolytus dimidiatus Chapuis, *Xyleborus ferrugineus* (Fabricius).

Lonchocarpus sericeus

Doliopygus brevis (Strohmeyer), *Doliopygus perbrevis* Schedl, *Platypus hintzi* Schanfuss, *Tiarophorus intermedius* Schedl.

***Lonchocarpus* spp.**

Chramesus periosus Wood, *Chramesus securus* Wood, *Hylesinopsis rhodesianus* (Eggers), *Pseudothysanoc squameus* Wood, *Rhopalopschion lonchocarpae* (Schedl), *Scolytus dimidiatus* Chapuis, *Scolytus propinquus* Blandford, *Strombophorus interstitialis* Browne.

Lonicera caprifolium*Xylosandrus morigerus* (Blandford).

Lonicera spp.

Hypothenemus eruditus Westwood, *Xylosandrus discolor* (Blandford).

Lophira alata

Cylindropalpus auricomans (Schaufuss), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Premnobius corthyloides* Hagedorn, *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).

Lophira procera

Platypus parallelus (Fabricius), *Xyleborus affinis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Lophira spp.

Premnobius cavipennis Eichhoff.

Lophopetalum spp.

Ambrosiodinus sarawakensis (Eggers), *Baiocis pernanulus* (Schedl), *Platypus lingulatus* Roberts & Morimoto, *Platypus lophopetali* Browne, *Platypus pseudocupulatus* Schedl, *Platypus puerulus* (Schedl).

Loranthus spp.

Cyrtogenius polyphagus (Schedl), *Thamniurgus granulicollis* Schedl, *Xylocleptes sidanus* (Schedl), *Xylosandrus abruptulus* (Schedl).

Loroea broeiuii

Chaetastus tuberculatus (Chapuis), *Doliopygus artemis* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus excavatus* (Sampson), *Xyleborus alluaudi* Schaufuss.

Loroea klaineana

Xyleborus affinis Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).

Loroea spp.

Doliopygus excavatus (Sampson), *Periommatius excisus* Strohmeier, *Periommatius longicollis* Chapuis.

Loroea trichilioides

Chaetastus tuberculatus (Chapuis), *Doliopygus artemis* (Schedl), *Doliopygus bitalei* Schedl, *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeier), *Doliopygus dubius* (Sampson), *Hypothenemus eruditus* Westwood, *Periommatius grandis* Schedl, *Platypus parallelus* (Fabricius), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus amanicus* Hagedorn, *Xyleborus principalis* Eichhoff, *Xyleborus sphenos* Sampson, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Lucuma malaccensis

Platypus loriceatus (Sampson), *Platypus signatus* Chapuis, *Platypus westwoodi* Chapuis.

Luffa acutangula

Dendrocraniulus cucurbitae (LeConte), *Dendrocraniulus sobrinus* Wood.

Luffa aegyptiaca

Cosmoderes monilicollis Eichhoff.

Luffa spp.

Xylosandrus crassiusculus (Motschulsky).

Lupinus spp.

Chramesus editus (Bright).

Lysiloma bahamensis

Hypothenemus brunneus (Hopkins), *Hypothenemus hirsutus* (Wood), *Hypothenemus squamosus* (Hopkins).

Maba abyssinica

Xyleborus elongatus Eggers, *Xyleborus ugandensis* Schedl.

Maba laurentii

Doliopygus citri Schedl, *Doliopygus conradti* (Strohmeier), *Doliopygus subdolosus* Schedl, *Doliopygus tenuis* (Strohmeier), *Mesoplatus ustus* Schedl, *Platy-*

pus hintzi Schaufuss, *Xyleborus multispinatus* Eggers, *Xyleborus scobinatus* Hagedorn.

Maba sanduicensis

Crossotarsus externidentatus (Fairmaire).

Maba spp.

Glostatus acaciae (Schedl), *Scolytoptatypus fasciatus* Hagedorn, *Trachyostus aterrimus* (Schaufuss).

Macadamia indica

Coccotrypes cyperi (Beeson).

Macadamia spp.

Eucallacea fornicatus (Eichhoff), *Xylosandrus compactus* (Eichhoff).

Macaranga aleuritoides

Hypothenemus quadridentatus Browne.

Macaranga andamanica

Diamerus curvifer (Walker).

Macaranga harteri

Periommatius grandis Schedl, *Periommatius longicollis* Chapuis, *Platypus orientalis* (Strohmeier).

Macaranga denticulata

Coccotrypes cyperi (Beeson), *Coccotrypes longior* (Eggers), *Coccotrypes nubilus* (Blandford), *Diamerus nigrescens* Eggers, *Eucallacea interjectus* (Blandford), *Eucallacea celatus* (Sampson), *Eucallacea wallacei* (Blandford), *Platypus lopchuensis* Schedl, *Scolytoptatypus raja* Blandford.

Macaranga gigantea

Coccotrypes cinnamomi (Eggers), *Coccotrypes medius* (Eggers).

Macaranga kilimandjarica

Crossotarsus externidentatus (Fairmaire).

Macaranga laucifolia

Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus donisi* Schedl, *Doliopygus erichsoni* (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus minor* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Hypothenemus eruditus* Westwood, *Periommatius excisus* Strohmeier, *Periommatius subrobustus* Schedl, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeier), *Platypus picinus* Schedl, *Platypus schenklengi* (Strohmeier), *Platypus spinulosus* (Strohmeier), *Trachyostus schaufussi* (Strohmeier), *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius).

Macaranga monandra

Xylosandrus crassiusculus (Motschulsky).

Macaranga neomilbraediana

Eucallacea xanthopus (Eichhoff).

Macaranga schweinfurthii

Chaetastus tuberculatus (Chapuis).

Macaranga spinosa

Platypus minutissimus Schedl, *Xyleborinus sharpae* (Hopkins).

Macaranga spp.

Amasa versicolor (Sampson), *Chaetastus montanus* Schedl, *Crossotarsus terminatus* Chapuis, *Cyclorhipidion bicinctum* (Schedl), *Diapus bilunatus* Schedl, *Diapus papuanus* Schedl, *Diapus nebulosus* Roberts, *Diapus turgidus* Roberts, *Doliopygus bidentatus* (Strohmeier), *Doliopygus bilobatus* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus retusus* Schedl, *Platypus capitilunatus* Roberts & Morimoto, *Platypus collinus* Browne, *Platypus jansonii* Chapuis, *Platypus opacideclivis* Schedl, *Platypus picinus* Schedl, *Platypus solidus* Walker, *Prem-*

nobius cavipennis Eichhoff, *Prennubius sexspinosus* Eggers, *Scolytoplatypus mikado* Blandford, *Scolytoplatypus occidentalis* Browne, *Scolytoplatypus truncatus* Browne, *Triozastus marshalli* (Sampson), *Xyleborinus gracilipennis* (Schedl), *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborinus similans* (Eggers), *Xyleborus truncatulus* Schedl, *Xylosandrus difficilis* (Eggers), *Xylosandrus morigerus* (Blandford).

Machilus edulis

Diapus spatulifer Browne.

Machilus japonica

Platypus calannus Blandford, *Scolytus aratus* Blandford, *Xyleborus machili* Niisima.

Machilus longipaniculata

Crossotarsus rengetensis Niisima & Murayama, *Xyleborus atratus* Eichhoff.

Machilus macrautha

Coccotrypes cinnamomi (Eggers), *Coccotrypes variabilis* (Beeson), *Platypus circumdentatus* Strohmeyer.

Machilus odoratissima

Crossotarsus externedentatus (Fairmaire), *Crossotarsus machili* Beeson, *Diapus quadrispinatus* Chapuis, *Diapus quinquespinatus* Chapuis, *Eucallacea velatus* (Sampson), *Indocryphalus machili* Wood, *Phloeosinus machilus* (Schedl), *Phloeosinus squamulatus* Chapuis, *Platypus curtatus* Sampson, *Platypus falcatus* Strohmeyer, *Scolytoplatypus minimus* Hagedorn, *Scolytoplatypus raja* Blandford, *Xyleborus improbus* Sampson, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).

Machilus spp.

Amasa amputatus (Blandford), *Ambrosiodmus apicalis* (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Diapus spatulifer* Browne, *Eucallacea interjectus* (Blandford), *Eucallacea validus* (Eichhoff), *Hadrodemus amorphus* (Eggers), *Hadrodemus artemcomans* (Schedl), *Scolytoplatypus mikado* Blandford, *Scolytoplatypus tycon* Blandford, *Xyleborus aquilus* Blandford, *Xyleborus kumamotoensis* Murayama, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus germanus* (Blandford).

Machilus thunbergii

Cnestus murayamai Schedl, *Crossotarsus niponicus* Blandford, *Dryocactiops coffeae* (Eggers), *Indocryphalus pubipennis* (Blandford), *Phloeosinus pulchellus* Blandford, *Platypus calannus* Blandford, *Platypus contaminatus* (Blandford), *Sucus niisimai* (Eggers), *Taphrocyetus taradakensis* (Murayama), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus atratus* Eichhoff, *Xyleborus colvulus* (Fabricius), *Xylosandrus brevis* (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus mutilatus* (Blandford).

Maclura pomifera

Hypothenemus seriatus (Eichhoff), *Liparthrum squamosum* (Blackman).

Macphersonia madagascariensis

Mitosoma excisum Schaufuss, *Xyleborinus spiculatus* (Schaufuss), *Xyleborus alluaudi* Schaufuss.

Macrolobium heudelotii

Ambrosiodmus tropicus (Hagedorn), *Cyclorhipidion cachani* (Schedl), *Cylindropalpus auricomans* (Schaufuss), *Doliopygus serratus* (Strohmeyer), *Xyleborinus similans* (Eggers), *Xyleborus ambasiusculus* Eggers, *Xyleborus subtuberculatus* Eggers.

Macrolobium macrophyllum

Cylindropalpus auricomans (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers,

Xyleborus rothkirchi Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Macrolobium spp.

Ambrosiodmus tropicus (Hagedorn), *Coptoborus usagarius* (Eggers), *Hypothenemus macrolobii* (Eggers), *Prennubius cavipennis* Eichhoff, *Prennubius quadrispinosus* Schedl, *Prennubius sexspinosus* Eggers, *Xyleborinus sharpae* (Hopkins), *Xyleborus ambasiusculus* Eggers, *Xyleborus conradti* Hagedorn, *Xyleborus multispinatus* Eggers, *Xyleborus subtuberculatus* Eggers, *Xylosandrus corthyloides* (Schedl), *Xylosandrus crassiusculus* (Motschulsky).

Madhuca spp.

Amasa striatotruncatus (Schedl), *Crossotarsus externedentatus* (Fairmaire).

Madrilus edulis

Xyleborus lineatus Eggers.

Maesa lanceolata

Eucallacea xanthopus (Eichhoff).

Maesa rufescens

Chaetastus montanus Schedl, *Doliopygus artespiniatus* (Schedl), *Doliopygus bidentatus* (Strohmeyer), *Eucallacea xanthopus* (Eichhoff), *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Theoborus ricini* (Eggers), *Xyleborus alluaudi* Schaufuss, *Xyleborus principalis* Eichhoff.

Maesa spp.

Chaetastus montanus Schedl, *Eucallacea malloti* (Eggers), *Scolytoplatypus kicuensis* Schedl, *Xyleborus ruandae* Schedl.

Maesa tenera

Xylosandrus brevis (Eichhoff).

Maesobotrya spp.

Scolytoplatypus kicuensis Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus alluaudi* Schaufuss.

Maesopsis eminii

Ambrosiodmus albizzianus (Schedl), *Coptoborus usagarius* (Eggers), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus retusus* Schedl, *Doliopygus serratus* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Periommatius subrobustus* Schedl, *Platypus hintzi* Schaufuss, *Scolytoplatypus africanus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus colvulus* (Fabricius).

Maesopsis spp.

Prennubius cavipennis Eichhoff.

Magnolia cathartii

Platypus errans (Sampson).

Magnolia foetida

Hypothenemus eruditus Westwood.

Magnolia glauca

Cnesinus strigicollis LeConte, *Hypothenemus eruditus* Westwood.

Magnolia grandiflora

Platypus compositus Say.

Magnolia obovata

Crossotarsus niponicus Blandford, *Eucallacea validus* (Eichhoff), *Xyleborinus attenuatus* (Blandford).

Magnolia spp.

Hypothenemus erudiae (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Scolytoplatypus shoguen* Blandford.

Magnolia virginiana

Hypothenemus eruditus Westwood.

Malacantha spp.

Chaetastus montanus Schedl, *Doliopygus bidentatus* (Strohmeyer).

Malaisia scandens

Cryphalus waplery Eichhoff.

Mallotus albus

Carchesiopygus impariporus (Beeson), *Hadrodemius pseudocomans* (Eggers), *Platypus solidus* Walker, *Webbia trigintispinatus* Sampson.

Mallotus japonicus

Eucallacca validus (Eichhoff), *Platypus calamus* Blandford, *Xyleborus aquilus* Blandford, *Xyleborus seriatus* Blandford, *Xyleborus volutus* (Fabricius).

Mallotus philippinensis

Coccotrypes nubilus (Blandford), *Eucallacca bicolor* (Blandford), *Eucallacca malloti* (Eggers), *Xyleborinus artestriatus* (Eichhoff), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Mallotus ricinoides

Cryphalus malloti Schedl, *Scolytplatypus pusillus* Eggers.

Mallotus roxburghianus

Crossotarsus machili Beeson.

Mallotus spp.

Xyleborinus andrewesi (Blandford).

Malus formosana

Xyleborus atratus Eichhoff.

Malus punila

Ambrosiodmus apicalis (Blandford), *Scolytus aratus* Blandford, *Xylosandrus crassiusculus* (Motschulsky).

Malus sieboldii

Xyleborus montanus Niisima.

Malus spp.

Corthylus mexicanus Schedl, *Hypothenemus eruditus* Westwood, *Scolytus laevis* Chapuis, *Scolytus mali* (Beckstein), *Scolytus pomi* Yin & Huang, *Scolytus rugulosus* (Müller), *Scolytus semenovi* (Spessivtsev), *Scolytus tadzhikistanicus* Stark, *Tiarophorus scrutator* (Pandelle), *Xyleborinus saxensis* (Ratzeburg).

Malus sylvestris

Xylosandrus pseudosolidus (Schedl).

Malvastrum spp.

Hypothenemus californicus Hopkins, *Xylosandrus compactus* (Eichhoff).

Mammea africana

Doliopygus coeloccephalus (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus expletus* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus propinquus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Perionnatus excisus* Strohmeyer, *Perionnatus longicollis* Chapuis, *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).

Mammea americana

Coccotrypes cyperi (Beeson).

Mammea spp.

Ambrosiodmus eichhoffi (Schreiner).

Mangifera caesia

Xyleborinus andrewesi (Blandford).

Mangifera foetida

Xyleborus emarginatus Eichhoff.

Mangifera indica

Ambrosiodmus diversipennis (Schedl), *Ambrosiodmus*

fraterculus (Schaufuss), *Ambrosiodmus hagedorni* (Iglesias), *Coccotrypes confusus* (Eggers), *Coccotrypes cyperi* (Beeson), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus minax* (Walker), *Cryptocarenum seriatus* Eggers, *Cyclorhupidion mangoense* (Schedl), *Diamerus mangiferae* Brownie, *Diamerus nigristosus* Eggers, *Diapus quinquespinatus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Eucallacca interjectus* (Blandford), *Eucallacca xanthopus* (Eichhoff), *Ficicis despectus* (Walker), *Hypocryphalus mangiferae* (Stebbing), *Hypothenemus birnanus* (Eichhoff), *Hypothenemus californicus* Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins), *Hypothenemus javanus* (Eggers), *Hypothenemus mangiferae* Schedl, *Hypothenemus morio* (Eggers), *Hypothenemus seriatus* (Eichhoff), *Hypothenemus setosus* (Eichhoff), *Platypus jansoni* Chapuis, *Platypus latifinis* Walker, *Platypus solidus* Walker, *Premnobius ambitiosus* (Schaufuss), *Xyleborinus andrewesi* (Blandford), *Xyleborinus artestriatus* (Eichhoff), *Xyleborinus forficuloides* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus biconicus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus haberkorni* Eggers, *Xyleborus metacuneolus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xyleborus spinulosus* Blandford, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus manicus* (Blandford), *Xylosandrus pseudosolidus* (Schedl).

Mangifera minor

Platypus jansoni Chapuis.

Mangifera odorata

Hypocryphalus mangiferae (Stebbing), *Xyleborus metacuneolus* Eggers, *Xylosandrus compactus* (Eichhoff).

Mangifera spp.

Crossotarsus browni Schedl, *Diapus nanus* Schedl, *Diapus pusillimus* Chapuis, *Eucallacca fornicatus* (Eichhoff), *Platypus ovatus* Strohmeyer, *Platypus pseudocupulatus* Schedl, *Premnobius catipennis* Eichhoff.

Manihot glaziovii

Xyleborus affinis Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky).

Manihot utilissima

Hypothenemus bahianiae Schedl, *Platypus parallelus* (Fabricius), *Xyleborus ferrugineus* (Fabricius).

Manikara hexandra

Scolytomimus rectus Wood.

Manikara kauki

Coccotrypes carpophagus (Hornung).

Maniltoa spp.

Crossotarsus mniszecchi Chapuis, *Platypus misoolensis* Brownie, *Platypus ritiensis* Roberts.

Mansoniea altissima

Doliopygus nairobiensis (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky).

Manuglieta insignis

Dactylipalpus transversus Chapuis.

Marcgravia spp.

Premnobius catipennis Eichhoff, *Xyleborinus gracilis* (Eichhoff), *Xyleborus parcellus* Wood, *Xyleborus rugosipennis* Schedl.

Marumia muscosa

Crossotarsus terminatus Chapuis, *Platypus fraterculus* Schedl, *Xylosandrus morigerus* (Blandford).

Mauloutchia spp.

Mitosoma octospinosum Schedl, *Mitosoma suspicax* Schedl, *Sauroptilius sauropterus* (Schedl).

Mauria glauca

Pityophthorus debilis Wood.

Maytenus acuminatus

Lanurgus gracilis Schedl, *Lanurgus spatulatus* Schedl.

Maytenis spp.

Scolytodendron fasciatus Hagedorn.

Medicago sativa

Hylastinus obscurus (Marsham).

Melaleuca leucadendron

Xyleborus perforans (Wollaston).

Melaleuca spp.

Xylosandrus compactus (Eichhoff).

Melanorrhoea curtisii

Crossotarsus brownei Schedl, *Xyleborus haberkornii* Eggers.

Melanorrhoea spp.

Arixyleborus scabripennis (Blandford), *Crossotarsus squamulatus* Chapuis, *Hadrodemius comans* (Sampson), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus nancus* (Blandford).

Melanorrhoea usitata

Polygraphus thitsi Wood, *Polygraphus vexator* (Browne).

Melanthera brownii

Cyrtogenius polyphagus (Schedl), *Xylocleptes sidanus* (Schedl).

Melastoma malabathricum

Eucallalcea fornicatus (Eichhoff).

Melastoma spp.

Xyleborinus saxeseni (Ratzeburg), *Xylosandrus compactus* (Eichhoff).

Melia azedarach

Crossotarsus rengetensis Niisima & Murayama, *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Melia spp.

Cnestus suturalis (Eggers), *Xylosandrus compactus* (Eichhoff).

Melia azedarach japonica

Ambrosiodmus rubricollis (Eichhoff).

Melicocca bijugata

Xyleborus ferrugineus (Fabricius), *Xyleborus colvulus* (Fabricius), *Xylosandrus laticeps* (Wood).

Melicoccus spp.

Xylosandrus compactus (Eichhoff).

Melicope neurococca

Ficicis varians Lea.

Melilotus spp.

Hylastinus obscurus (Marsham).

Meliosma cuneifolia

Xylosandrus brevis (Eichhoff).

Meliosma myriantha

Platypus calamus Blandford.

Meliosma simplicifolia

Platypus catus Strohmeyer, *Platypus indicus* Strohmeyer, *Platypus luniger* Motschulsky, *Platypus secretus* Sampson.

Melochia odorata

Xyleborus immersus Schedl.

Melochia spp.

Xylosandrus compactus (Eichhoff).

Melochia umbellata

Xylosandrus crassiusculus (Motschulsky).

Melodorum fulgens

Eruoporus melodori (Hopkins).

Memocylon spp.

Coptodryas nuxax (Schedl).

Menispermum spp.

Araptus trepidus Wood.

Meriania spp.

Scolytodes serenus Wood.

Mesogordonia spp.

Xyleborinus quadrispinosus (Eichhoff), *Xyleborus submolestus* Schedl.

Mespilodaphne spp.

Cyclorhpidion scalptor (Schedl), *Mitosoma paulianum* Schedl.

Mespilus germanica

Scolytus amygdali Guerin-Meneville.

Mespilus spp.

Scolytus rugulosus (Müller).

Mesua ferrea

Carchesiopygus impariporus (Beeson), *Coccotrypes monoceros* (Beeson), *Coccotrypes papuanus* (Eggers), *Crossotarsus bonvouloiri* Chapuis, *Crossotarsus schedli* Browne, *Cryphalus scabriscollis* Eichhoff, *Dactylipalpus transversus* Chapuis, *Hadrodemius metacomans* (Eggers), *Platypus catus* Strohmeyer, *Platypus errans* (Sampson), *Xyleborinus andrewesi* (Blandford), *Xyleborus fallax* Eichhoff, *Xyleborus mucronatulus* Eggers, *Xylosandrus mesuae* (Eggers).

Mesua myriocarpa

Ambrosiodmus lewisi (Blandford).

Metopium toxiferum

Cryptocarenum seriatus Eggers, *Hypothenemus hirsutus* (Wood), *Pityophthorus concentricus* Eichhoff.

Metrosideros collina polymorpha

Xyleborinus saxeseni (Ratzeburg), *Xyleborus similinus* Perkins.

Metrosideros excelsa

Anasa truncatus (Erichson).

Michelia champaca

Ambrosiodmus lantanae (Eggers), *Coptodryas mus* (Eggers), *Cyclorhpidion hirtum* (Hagedorn).

Michelia excelsa

Diapus spatulifer Browne.

Michelia nilagirica

Platypus furcatus Blandford.

Michelia oblonga

Platypus catus Strohmeyer.

Miconia caudata

Corthylus rubricollis Blandford, *Monarthrum carinulum* Wood, *Theoborus pristis* (Wood).

Miconia dodecandra

Corthyocyclon aztecum (Bright).

Miconia globuliflora

Corthylus collaris Blandford, *Corthylus serratus* Wood, *Corthylus strigilis* Wood, *Corthylus tricus* Wood, *Scolytodes pumilus* Wood, *Xyleborus affinis* Eichhoff.

Miconia pubescens

Monarthrum pennatum (Schedl), *Monarthrum proximum* Wood.

Miconia schlechtendalii

Corthylus sentus Wood, *Micracis lignicolus* Wood.

Miconia spp.

Corthyocyclon aztecum (Bright), *Corthylocorus barbatus* (Blandford), *Corthylocorus mexicanus* (Schedl), *Corthylus comatus* Blandford, *Corthylus retusus* Wood, *Dryocoetoides capucinus* (Eichhoff), *Hypothenemus*

erectus LeConte, *Hypothenemus interstitialis* (Hopkins), *Micracis swainci* Blackman, *Microcorthyus demissus* Wood, *Monarthrum hoegei* (Blandford), *Scolytodes immanis* Wood, *Taurodemus perebeae* (Ferrari), *Tricolus aciculatus* Wood, *Tricolus ovicollis* Blandford, *Xyleborus meritus* Wood.

Microcos coriacea

Doliopygus conradti (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Premnobius quadripinosus* Schedl, *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Microcos pinnatifida

Cylindropalpus granulatus (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus jurioni* Schedl, *Doliopygus interjectus* Schedl, *Doliopygus medius* Schedl, *Doliopygus subditicus* (Schedl), *Hypothenemus eruditus* Westwood, *Hypothenemus macrolobii* (Eggers), *Premnobius nodulosus* Hagedorn, *Xyleborinus gracilipennis* (Schedl), *Xyleborus tenellus* Schedl, *Xyleborus xylotripes* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Microcos spp.

Diaprus turgidus Roberts.

Microdesmis puberula

Cylindropalpus pumilio (Schedl), *Cylindropalpus pusillus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus jurioni* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff.

Microglossa volubilis

Cyrtogenius movoliae (Schedl), *Hypothenemus atterrimulus* Wood.

Micromeles spp.

Dryocoetes padi Stark, *Platypus volaticus* Schedl.

Mikania scandens

Dendrochilus mikaniae Schedl.

Mildbraediendendron excelsum

Chaetastus tuberculatus (Chapuis), *Doliopygus perbrevis* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alhuaudi* Schaufuss.

Millettia atropurpurea

Platypus indicus Strohmeyer.

Millettia dehiscens

Platypus biuncus Blandford, *Platypus jansoni* Chapuis, *Platypus pseudocupulatus* Schedl.

Millettia drastica

Doliopygus conradti (Strohmeyer), *Hypothenemus eruditus* Westwood.

Millettia duchesnei

Doliopygus serratus (Strohmeyer), *Strombophorus crenatus* Hagedorn, *Strombophorus millettiae* Schedl, *Strombophorus vagans* Schedl.

Millettia ferruginea

Cyrtogenius millettiae (Schedl), *Cyrtogenius naihungensis* (Schedl), *Euwallacea xanthopus* (Eichhoff), *Hypothenemus lefevrei* (Schedl), *Platypus impressus* (Strohmeyer), *Xyleborinus collarti* (Eggers), *Xyleborus alhuaudi* Schaufuss, *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff.

Millettia hylobia

Rhopalopselion congonum (Schedl).

Millettia laurentii

Hypothenemus eruditus Westwood.

Millettia megasperma

Scolytogenes tricolor (Lea).

Millettia pendula

Platypus lepidus Chapuis.

Millettia spp.

Acanthotomicus biconicus (Schedl), *Doliopygus falsificus* (Schedl), *Tiarophorus capucinus* Schedl, *Xyleborus multispinatus* Eggers.

Millettia sutherlandii

Doliopygus erichsoni (Chapuis).

Millettia thomningii

Doliopygus bulbifer Schedl.

Mimosa ercvendbergii

Chramesus unicornis Wood.

Mimosa spp.

Chaetophloeus andinus Wood, *Chramesus mimosae* Blackman, *Hylocurus parkinsoniae* Blackman, *Hypothenemus brunneus* (Hopkins), *Hypothenemus dolosus* Wood, *Hypothenemus erectus* LeConte, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus opacus* (Eichhoff), *Hypothenemus rotundicollis* (Eichhoff), *Micracisella nanula* (LeConte), *Micracisella opacithorax* (Schedl), *Monarthrum fasciatum* (Say), *Pseudothyrsanoes acaciae* (Blackman), *Theoborus coartatus* (Sampson), *Thysanoes inornatus* Wood, *Thysanoes texanus* Blackman.

Mimusops africana

Xyleborus volvulus (Fabricius).

Mimusops caffra

Xyloctonus scolytoides Eichhoff.

Mimusops djave

Platypus parallelus (Fabricius).

Mimusops elengi

Scolytomimus mimusopsis Wood.

Mimusops heckelii

Chaetastus tuberculatus (Chapuis), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus volvulus* (Fabricius).

Mimusops littoralis

Scolytomimus pusillus (Eggers), *Xyleborus bidentatus* (Motschulsky), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Mimusops spp.

Xyleborus ferrugineus (Fabricius).

Mitragyna spp.

Ambrosiodinus eichhoffi (Schreiner), *Chaetastus tuberculatus* (Chapuis), *Doliopygus dubius* (Sampson), *Periommatius excisus* Strohmeyer, *Periommatius subrobustus* Schedl, *Pityophthorus hintzi* Schedl, *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).

Mitragyna stipulosa

Chaetastus diversifrons Browne, *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff, *Xyleborus piceus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).

Mitrastemon spp.

Coccotrypes barbatus (Schedl).

Moerua spp.

Ambrosiodinus signiceps (Schedl).

Monodora myristica

Chaetastus tuberculatus (Chapuis), *Hypothenemus pubipennis* (Eggers), *Mesoplatypus venustus* Schedl, *Xyleborus alhuaudi* Schaufuss.

Monotes kerstingii*Doliopygus terebrans* Schedl.**Morinda citrifolia***Xyleborus perforans* (Wollaston).**Morinda lucida***Chaetastus tuberculatus* (Chapuis), *Doliopygus conradti* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus lefevrei* Schedl., *Xyleborus affinis* Eichhoff, *Xyleborus subtuberculatus* Eggers.**Moringa oleifera***Hypothenemus eruditus* Westwood.**Moringa pterygosperma***Scolytogenes indicus* Wood.**Morisonia americana***Cucinus setulosus* Blandford.**Morus acidosa***Xyleborus atratus* Eichhoff.**Morus alba***Ambrosiodmus rubricollis* (Eichhoff), *Chramesus setosus* Wood, *Cryphalus exiguus* Blandford, *Ficicis varians* Lea, *Phloeotribus frontalis* (Olivier), *Thysanocis fimbriicornis* LeConte, *Trypodendron domesticum* (Linnaeus), *Xyleborus atratus* Eichhoff.**Morus bombycis***Cryphalus exiguus* Blandford, *Xyleborus atratus* Eichhoff.**Morus lactea***Chaetastus tuberculatus* (Chapuis), *Hylesinopsis dubius* Eggers, *Xyleborus piceus* (Motschulsky).**Morus mesozygia***Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Hylesinopsis dubius* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus coliculus* (Fabricius).**Morus mexozygia***Doliopygus brevis* (Strohmeyer), *Doliopygus perbrevis* Schedl., *Platypus hintzi* Schaufuss, *Xyleborus alluaudi* Schaufuss.**Morus rubra***Hylocurus langstoni* Blackman, *Hypothenemus columbi* Hopkins, *Hypothenemus crudiæ* (Panzer), *Hypothenemus interstitialis* (Hopkins), *Phloeotribus frontalis* (Olivier).**Morus spp.***Ambrosiodmus rubricollis* (Eichhoff), *Doliopygus rapax* (Sampson), *Hypothenemus eruditus* Westwood, *Premnobius catipennis* Eichhoff, *Trachyostus aterrimus* (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xyleborus atratus* Eichhoff, *Xyleborus praeceus* Blandford, *Xylosandrus germanus* (Blandford).**Motandra guineensis***Acanthotomicus biconicus* (Schedl).**Mucuna andreaana***Araptus politus* (Blandford).**Mucuna fauceitti***Araptus politus* (Blandford).**Mucuna spp.***Araptus mucinae* (Blackman).**Muelenbeckia tamnifolia***Stegomerus montanus* Wood.**Muntingia calabura***Xylosandrus compactus* (Eichhoff).**Muonnia spp.***Hypothenemus seriatus* (Eichhoff).**Murraya koenigii***Crossotarsus externedentatus* (Fairmaire), *Xylosandrus crassiusculus* (Motschulsky).**Murraya paniculata***Xylebomis perforans* (Wollaston).**Murraya spp.***Xylosandrus compactus* (Eichhoff).**Musa basjoo***Hypothenemus eruditus* Westwood.**Musanga cecropioides***Cryphalus bellus* Schedl, *Doliopygus chapuisi* (Duvi-vier), *Doliopygus conradti* (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Eucallacea pandae* (Schedl), *Hypothenemus biseriatus* (Eggers), *Hypothenemus eruditus* Westwood, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Polygraphus musangae* (Schedl), *Premnobius quadrispinosus* Schedl, *Serratus similis* (Eggers), *Xyleborinus gracilipennis* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus caudatus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).**Musanga spp.***Ambrosiodmus eichhoffi* (Schreiner).**Myrianthus arboreus***Doliopygus donisi* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus unicornis* Schedl, *Periommatius excisus* Strohmeyer, *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).**Myrianthus holstii***Xyleborus alluaudi* Schaufuss, *Xyleborus clerodendronae* Schedl.**Myrianthus spp.***Ambrosiodmus tropicus* (Hagedorn), *Premnobius catipennis* Eichhoff, *Xyleborus ambastusculus* Eggers.**Myrica faya***Xyleborinus savaseni* (Ratzeburg).**Myrica pubescens***Araptus nigrellus* Wood, *Corthylus granulifer* Wood, *Corthylus rubricollis* Blandford, *Corthylus strigilis* Wood, *Xyleborus meritus* Wood.**Myrica salicifolia***Doliopygus erichsoni* (Chapuis), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus lefevrei* (Schedl), *Xyleborus principalis* Eichhoff.**Myrica spp.***Hylocurus hintellus* (LeConte), *Xylosandrus germanus* (Blandford).**Myristica andamanica***Crossotarsus andamanus* Beeson, *Eucallacea bicolor* (Blandford), *Xyleborinus andrewesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.**Myristica castaneaeefolia***Crossotarsus externedentatus* (Fairmaire), *Platypus citiensis* Roberts, *Xyleborus perforans* (Wollaston).**Myristica cinnamomea***Coccotrypes variabilis* (Beeson).**Myristica dactyloides***Arxyleborus granulifer* (Eggers), *Coccotrypes longior* (Eggers), *Coccotrypes salakensis* (Schedl), *Coccotrypes vulgaris* (Eggers), *Cryphalus nigricans* Schedl, *Hyledius ceylonicus* (Schedl), *Hyledius concinnulus* (Walker), *Hyledius regalis* (Wood).**Myristica fragrans***Anasa batoerradensis* (Schedl), *Arxyleborus malayensis* (Schedl), *Coccotrypes myristicae* (Roepke),

Coccotrypes priesneri Schedl, *Coptodryas myristicac* (Schedl), *Coptodryas nugax* (Schedl), *Hypocryphalus sumatranus* (Schedl), *Hypothenemus obscurus* (Fabricius), *Hyledius nitidicollis* (Motschulsky), *Platypus webberi* Schedl, *Xyleborus fulculus* Schedl, *Xyleborus setulosus* Eggers.

Myristica furfuracae

Coriaceophilus xyloctonoides Schedl, *Hypocryphalus granulatus* Schedl, *Hyledius cribratus* (Blandford).

Myristica indica

Xyleborinus andrewesi (Blandford).

Myristica longifolia

Hadrodemius pseudoconans (Eggers), *Hyledius jiri* (Wood).

Myristica philippinensis

Dactyliphalus transversi Chapuis, *Hyledius cribratus* (Blandford), *Hyledius nitidicollis* (Motschulsky).

***Myristica* spp.**

Coccotrypes barbatus (Schedl), *Coccotrypes tapatapaoanus* (Schedl), *Crossotarsus longicornis* (Schedl), *Diamerus gravis* Brownie, *Diapus amblylaminatus* Roberts, *Diapus latespinis* Schedl, *Diapus nebulosus* Roberts, *Diapus oosis* Roberts, *Diapus papuanus* Schedl, *Hypocryphalus basilirtus* (Beeson), *Hypothenemus eruditus* Westwood, *Hyledius imitans* (Eggers), *Platypus bifidus* Schedl, *Platypus imberbis* Roberts, *Platypus negatus* Schedl, *Platypus nemorosus* Roberts, *Platypus papulosus* Roberts, *Platypus patens* Roberts, *Platypus schysi* Chapuis, *Xylosandrus crassiusculus* (Motschulsky).

***Myrmidone* spp.**

Hypothenemus eruditus Westwood.

Myroxylon toluiferum

Xyleborus affinis Eichhoff.

***Myrsine* spp.**

Xyleborinus saxescii (Ratzeburg), *Xylosandrus compactus* (Eichhoff).

***Myrtaceae* spp.**

Platypus madagascariensis Chapuis.

Napoleona imperialis

Doliopygus conradti (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Periommatius substriatus* Strohmeyer, *Platypus hintzi* Schaufuss, *Premnobius perspinidens* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus volculus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

***Napoleona* spp.**

Ambrosiodmus eichloffii (Schreiner), *Premnobius cavipennis* Eichhoff.

Nauclea diderichii

Chaetastus tuberculatus (Chapuis), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus piceus* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky).

Nauclea excelsa

Platypus cavius Strohmeyer.

Nauclea orientalis

Coptodryas eucalyptica (Schedl).

Nauclea sessilifolia

Platypus solidus Walker, *Platypus suffodiens* Sampson.

***Nauclea* spp.**

Platypus solidus Walker, *Platypus subtruncatus* Brownie, *Platypus uncatus* Brownie, *Premnobius cavipennis* Eichhoff.

Nectandra angustifolia

Xylosandrus compactus (Eichhoff).

***Nectandra* spp.**

Amphicramus terebella Blandford, *Araptus rufopalliat* Eichhoff, *Cnesinus bicostatus* Schedl, *Cnesinus intermedius* Schedl, *Cnesinus teretis* Wood, *Corthylus additus* Wood, *Corthylus cirritus* Wood, *Corthylus convexifrons* Wood, *Cryptocarenum brevicollis* Eggers, *Gnathotrupes colaphus* Wood, *Gnathotrupes nectandrae* Wood, *Hylocurus flagellatus* Wood, *Hypothenemus opacus* (Eichhoff), *Microcorthylus contractus* Wood, *Microcorthylus diversus* Wood, *Microcorthylus umbratus* Wood, *Monarthrum lobatum* (Ferrari), *Platypus deceptor* Wood, *Platypus secus* Wood, *Scolytodes decorus* Wood, *Sternobothrus bicaudatus* (Blandford), *Sternobothrus sculpturatus* (Blandford), *Taurodemus bicornutus* (Wood), *Xylosandrus retusus* (Eichhoff).

Neesia altissima

Xyleborus cancellatus Eggers.

Nemosoma elongatum

Xyleborinus saxescii (Ratzeburg).

Neobuxbaumia mezealensis

Cactopinus burjosi Wood.

Neodopsis baroni

Polygraphus amoenus (Schaufuss), *Xyleborinus forficuloides* (Schedl), *Xyleborus sakalava* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

***Neonauclea* spp.**

Diapus perygmaeus Roberts, *Diapus quinquispinatus* Chapuis, *Platypus aolai* Brownie, *Platypus solomonicus* Brownie.

Neotina boursii

Xyleborinus forficuloides (Schedl).

Neowashingtonia robusta

Coccotrypes carpophagus (Hornung).

Nephelium lappaceum

Xylosandrus manicus (Blandford).

Nephelium lichi

Coccotrypes advena Blandford, *Coccotrypes confusus* (Eggers), *Coccotrypes papuanus* (Eggers), *Euwallacea forficatus* (Eichhoff), *Hypothenemus birmanus* (Eichhoff).

Nephelium longana

Anasa beesoni (Eggers).

***Nephelium* spp.**

Anasa striatotruncatus (Schedl), *Arixyleborus suturalis* (Eggers), *Baiois perianthus* (Schedl), *Coccotrypes rhizophorae* (Hopkins), *Coptodryas diversicolor* (Eggers), *Crossotarsus nitidulus* (Schedl), *Platypus puerulus* (Schedl), *Xyleborinus pomelianus* (Schedl).

Nerium oleander

Araptus dubiosus (Schedl), *Araptus oleanderi* (Schedl).

***Nerium* spp.**

Microborus lautus Wood.

Nesogordonia papaverifera

Xyleborus affinis Eichhoff.

***Nesogordonia* spp.**

Polygraphus kasukumbii Schedl.

Newtonia buchananii

Platypus intermedius (Schedl).

Nicotiana tabacum*Xyleborus spinulosus* Blandford.***Nipa* spp.***Leptoxyleborus concisus* (Blandford).***Nolena* spp.***Chramesus annexens* (Wood), *Corthylus nolenae* Wood.***Nothofagus carrii****Platypus capito* Browne, *Platypus gracilior* Browne.***Nothofagus cunuaughamii****Platypus subgranosus* Schedl.***Nothofagus dombeyi****Gnathotrupes caliculus* (Schedl), *Gnathotrupes cirratus* Schedl, *Gnathotrupes impressus* (Schedl), *Gnathotrupes longipennis* (Blanchard), *Gnathotrupes longinervis* (Schedl), *Gnathotrupes nanulus* (Schedl), *Gnathotrupes naumanni* (Schedl), *Gnathotrupes nothofagi* (Schedl), *Gnathotrupes pauciconvexus* Schedl, *Gnathotrupes pauciconvexus* Schedl, *Gnathotrupes pustulatus* Schedl, *Gnathotrupes similis* Schedl, *Gnathotrupes solidus* Schedl, *Gnathotrupes vaser* (Schedl), *Gnathotrupes velatus* Schedl, *Phloeotribus nahueliac* (Schedl).***Nothofagus grandis****Diapus kuperi* Roberts.***Nothofagus obliqua****Sinophloeus porteri* Brethes.***Nothofagus pullei****Diapus pusillinus* Chapuis, *Diapus spinifer* Schedl, *Platypus celsus* Roberts, *Platypus inversus* Roberts, *Platypus longicalcaratus* Roberts, *Platypus nothofagus* Roberts, *Platypus petaloideus* Roberts, *Platypus runcinatus* Roberts.***Nothofagus pumilio****Xylechinus maculatus* Schedl.***Nothofagus* spp.***Crossotarsus coxalis* Schedl, *Crossotarsus dentatus* Browne, *Cryphalus nothofagi* Browne, *Diapus nebulosus* Roberts, *Diapus robustus* Schedl, *Diapus spinifer* Schedl, *Diapus unispinus* Roberts, *Hypothenemus incernis* Browne, *Platypus denticollis* Browne, *Platypus margaritaceus* Roberts & Morimoto, *Platypus panduriformis* Roberts, *Platypus petalinus* Roberts & Morimoto, *Platypus praececlens* Schedl, *Scolytogenes papuanus* (Schedl).***Nothopanax* spp.***Eucallacea fornicatus* (Eichhoff), *Hypothenemus eruditus* Westwood.***Nuxia conferta****Cyclorhipidion guincense* (Eggers).***Nyssa aquatica****Hypothenemus eruditus* Westwood, *Xylosandrus germanus* (Blandford).***Nyssa javanica****Platypus cavus* Strohmeier, *Platypus drescheri* Schedl, *Platypus tenuissimus* Schedl.***Nyssa sessiliflora****Diapus spatulifer* Browne, *Eucallacea bicolor* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Nyssa* spp.***Monarthrum fasciatum* (Say), *Monarthrum mali* (Fitch), *Xyleborus ferrugineus* (Fabricius), *Xyleborus sayi* (Hopkins).***Nyssa sylvatica****Hypothenemus eruditus* Westwood.***Obscurus* spp.***Hypothenemus eruditus* Westwood.***Ochanostachys amentacea****Crossotarsus squamulatus* Chapuis, *Crossotarsus wallacci* (Thomson), *Xyleborus perforans* (Wollaston), *Xyleborus colvulus* (Fabricius).***Ochua arborea****Lamirgus capensis* Schedl.***Ochrocarpos* spp.***Hypothenemus nuticus* (Schedl), *Polygraphus aequalis* Schedl, *Polygraphus acquatus* Schedl, *Polygraphus amoenus* (Schaufuss).***Ochroma lagopus****Eucallacea fornicatus* (Eichhoff), *Xylosandrus morigerus* (Blandford).***Ochroma* spp.***Ambrosiodmus hagedorni* (Iglesias), *Corthylus comatus* Blandford, *Corthylus strigilis* Wood, *Hypothenemus javanus* (Eggers), *Hypothenemus opacus* (Eichhoff), *Hypothenemus seriatus* (Eichhoff), *Hypothenemus trivialis* Wood, *Monarthrum consimile* (Blandford), *Phloeotribus vesculus* Wood, *Platypus chiriquensis* Wood, *Sampsonius dampfi* Schedl, *Taurodemus perebeae* (Ferrari), *Theoborus theobromae* Hopkins, *Xyleborinus bicornatulus* (Wood), *Xyleborus caribicus* Eggers, *Xyleborus ferox* Blandford, *Xyleborus ferrugineus* (Fabricius), *Xyleborus spinulosus* Blandford.***Ochroma velutina****Scolytodes ochromae* Wood.***Ochthocosmus africanus****Doliopygus artespinatus* (Schedl), *Doliopygus bijunctus* Roberts, *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeier), *Doliopygus costatus* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minutissimus* (Schedl), *Doliopygus serratus* (Strohmeier), *Doliopygus spectabilis* Schedl, *Doliopygus tenuis* (Strohmeier), *Platypus hintzi* Schaufuss, *Platypus intermedius* (Schedl), *Platypus rufescens* (Strohmeier), *Platypus schenklingi* (Strohmeier), *Trachyostus schoutedeni* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus multispinatus* Eggers, *Xyleborus rothkirchi* Eggers, *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).***Ocotea bullata****Xyleborinus acmulus* (Wollaston).***Ocotea catesbyana****Hypothenemus birmanus* (Eichhoff), *Hypothenemus brunneus* (Hopkins), *Hypothenemus dissimilis* (Zimmermann), *Hypothenemus interstitialis* (Hopkins), *Hypothenemus javanus* (Eggers), *Hypothenemus seriatus* (Eichhoff), *Xylosandrus zimmermanni* (Hopkins).***Ocotea guianensis****Taurodemus splendidus* (Schaufuss).***Ocotea laevis****Xyleborus sartor* Schedl.***Ocotea laevis****Platypus madagascariensis* Chapuis.***Ocotea puberula****Microcorthylus suggrandis* Schedl.***Ocotea racemosa****Xyleborus sakalata* Schedl.***Ocotea rodiaei****Platypus longulus* Chapuis.

Ocotea spp.

Ambrosiodinus innominatus (Schedl), *Chaetastus montanus* Schedl, *Mitosoma excisum* Schanfuss, *Mitosoma obliquatum* Schedl, *Pagiocerus frontalis* (Fabricius), *Premnobius cavipennis* Eichhoff.

Ocotea usambarensis

Chaetastus montanus Schedl, *Chortastus orientalis* Schedl, *Doliopygus bidentatus* (Strohmeyer).

Octomeles sumatrana

Crossotarsus semirictus Schedl, *Platypus jansoni* Chapuis, *Platypus solidus* Walker, *Xyleborus emarginatus* Eichhoff.

Odina fraxinifolia

Platypus darjeelingensis Schedl.

Odina wodier

Bothinodroctonus indicus Wood, *Coccotrypes nubilus* (Blandford), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus squamulatus* Chapuis, *Cryphalus scabricollis* Eichhoff, *Eucallacea xanthopus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Genyocerus furtivus* (Sampson), *Platypus secretus* Sampson, *Platypus solidus* Walker, *Polygraphus anogeissi* Wood, *Xyleborinus andrewesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Olea africana

Lanurgus oleae Schedl, *Lanurgus xylographus* Schedl.

Olea ambrensis

Platypus madagascariensis Chapuis, *Xyleborinus aduncus* (Schedl), *Xyleborinus quadrispinis* (Schedl), *Xyleborinus spiculatus* (Schaufuss).

Olea capensis

Lanurgus oleaeformis Schedl, *Lanurgus spathulatus* Schedl, *Lanurgus xylographus* Schedl, *Xyleborinus acmulus* (Wollaston).

Olea cuspidata

Hylesinus cingulatus Blandford, *Hylesinus macmahoni* (Stebbing).

Olea europaea

Carphoborus perrisi (Chapuis), *Chaetoptelius vestitus* (Mulsant & Rey), *Coccotrypes dactyliperda* (Fabricius), *Hylesinus californicus* (Swaine), *Hylesinus torauio* (Danthione), *Phlocotribus brevicollis* (Kolenati), *Phlocotribus maroccanus* (Guillebeau), *Phlocotribus scarabaeoides* (Bernard), *Scolytus pygmaeus* (Fabricius), *Xylosandrus compactus* (Eichhoff).

Olea hochstetteri

Chaetastus diversifrons Browne, *Chaetastus montanus* Schedl, *Ctonoxylon griseum* Schedl, *Ctonoxylon hamatum* Schedl, *Cyclorhipidion agnaticeps* (Schedl), *Lanurgus oleae* Schedl.

Olea laurifolia

Lanurgus oleaeformis Schedl, *Platypus sampsoni* (Schedl).

Olea spp.

Ambrosiodinus cichhoffi (Schreiner), *Hylesinus varius* (Fabricius), *Tiarophorus gardneri* Schedl.

Olea welwitschii

Chaetastus montanus Schedl, *Ctonoxylon hamatum* Schedl, *Ctonoxylon spathifer* Schedl.

Olmea recta

Chramesus tibialis Wood.

Olmediella spp.

Xylosandrus compactus (Eichhoff).

Oncostemum spp.

Xyleborinus signatipennis (Schedl).

Ongokea gore

Chaetastus tuberculatus (Chapuis), *Doliopygus arteminatus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus lateralis* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Eucallacea xanthopus* (Eichhoff), *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schanfuss, *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Trachyostus schoutedeni* (Strohmeyer), *Xyleborus alhuandi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Ononis natrix

Hylastinus obscurus (Marsham).

Opuntia spp.

Aphanarthrum indicum Wood.

Orbiguya oleifera

Coccotrypes cyperi (Beeson).

Oreodoxa spp.

Coccotrypes dactyliperda (Fabricius).

Oreopanax capitatum

Bothrostermus foveatus (Blackman), *Cnesinus denotatus* Wood, *Cnesinus porcatus* Blandford.

Oreopanax nubigenus

Gnathotrichus omissus Wood, *Pityophthorus timidulus* Wood, *Scolytodes irazuensis* Wood, *Scolytodes punctiferus* Wood, *Xylechinus avarus* Wood.

Oreopanax spp.

Cnesinus porcatus Blandford, *Corthylus mexicanus* Schedl, *Xylechinus marmoratus* Blandford, *Xylechinus tessellatus* Blandford.

Oreopanax xalapense

Corthycyclon aztecum (Bright), *Scolytodes irazuensis* Wood, *Scolytodes venustus* Wood, *Xylechinus fuliginosus* Blandford, *Xylechinus marmoratus* Blandford, *Xylechinus mexicanus* Wood.

Oricia suaveolens

Xyleborus alhuandi Schaufuss.

Origanum vulgare

Thammurgus kaltenbachii (Bach).

Ormosia formosana

Xyleborus atratus Eichhoff.

Ormosia hosiei

Xylosandrus mutilatus (Blandford).

Osbeckia aspera

Sueus niisimai (Eggers), *Xylosandrus mesuae* (Eggers).

Osmanthus fragrans

Xylosandrus mutilatus (Blandford).

Osmanthus spp.

Xylosandrus compactus (Eichhoff).

Ostodes paniculata

Platypus secretus Sampson.

Ostrya virginica

Pseudopityophthorus asperulus (LeConte), *Pseudothy-sanoes lecontei* Blackman.

Ostrya carpinifolia

Corthylus punctatissimus (Zimmermann), *Scolytus carpinii* (Ratzeburg).

Ostyra spp.

Scolytus intricatus (Ratzeburg), *Tiarophorus scrutator* (Pandelle).

Ougeinia dalbergioides

Hypothenemus eruditus Westwood, *Xylechinus ougeniae* Wood, *Xylosandrus crassiusculus* (Motschulsky).

Oxyanthus speciosus

Chaetastus montanus Schedl, *Xyleborus alluaudi* Schauffuss.

Oxystigma manni

Doliopygus pseudoserratus Roberts, *Mesoplatypus cernatus* Schedl.

Oxystigma msoo

Periommatius nitidicollis Strohmeyer.

Oxystigma oxyphyllum

Chaetastus tuberculatus (Chapuis), *Coccotrypes rotundicollis* (Eggers), *Cylindropalpus laudatus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus citri* Schedl, *Doliopygus clarus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus retusus* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schauffuss, *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Platypus schenklingi* (Strohmeyer), *Platypus solutus* Schedl, *Premnobius quadrispinosus* Schedl, *Strombophorus erichsoni* (Schauffuss), *Trachyostus aterrimus* (Schauffuss), *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus annectens* Schedl, *Xyleborus colvulus* (Fabricius).

Pachylobus spp.

Premnobius cavipennis Eichhoff.

Pachycereus pectenaboriginum

Cactopinus mexicanus Wood.

Pachycereus spp.

Cactopinus atkinsoni Wood.

Pachyelasma spp.

Coccotrypes rotundicollis (Eggers), *Premnobius cavipennis* Eichhoff.

Pachyelasma tessmanii

Doliopygus conradti (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Platypus hintzi* Schauffuss, *Premnobius longus* Eggers.

Pachylobus pubescens=Dacryodes buetneri

Acanthotomicus pachylobius (Schedl).

Pachylobus deliciosus

Coccotrypes confusus (Eggers), *Ctonoxylon spathifer* Schedl, *Cyclorhipidion cachani* (Schedl), *Cylindropalpus auricomans* (Schauffuss), *Euwallacea pandae* (Schedl), *Platypus schenklingi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Pachylobus spp.

Xyleborus alluaudi Schauffuss, *Xyleborus piceus* (Motschulsky).

Pachypodanthium staudtii

Xyleborus colvulus (Fabricius).

Pachystela laurentii

Cylindropalpus granulatus (Schedl), *Cylindropalpus interpositus* (Schedl), *Doliopygus bidentatus* (Strohmeyer), *Doliopygus dolosus* Schedl, *Premnobius mikuniae* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus ambaspicatus* Schedl, *Xyleborus comparabilis* Schedl.

Pachystela spp.

Ambrosiodinus aegir (Eggers).

Pachytrope dimepate

Euwallacea xanthopus (Eichhoff), *Platypus madagascariensis* Chapuis, *Xyleborinus quadrispinus* (Schedl), *Xyleborus alluaudi* Schauffuss, *Xylosandrus crassiusculus* (Motschulsky).

Pachytrope obovata

Platypus madagascariensis Chapuis.

Padus asiatica

Cryphalus scopiger Berger.

Padus maaki

Dryocoetes padi Stark.

Padus pubescens

Cryphalus scopiger Berger.

Pajanelia longifolia

Euwallacea fornicatus (Eichhoff).

Palaquium elliptica

Crossotarsus quadricaudatus (Strohmeyer), *Xyleborinus andreweesi* (Blandford).

Palaquium gutta

Arixyleborus scabripennis (Blandford), *Hyledius crib-ratus* (Blandford), *Xyleborus adusticollis* (Motschulsky), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus manicus* (Blandford).

Palaquium hornei

Platypus citiensis Roberts.

Palaquium maingayi

Arixyleborus leprosus Schedl, *Arixyleborus minor* (Eggers), *Arixyleborus suturalis* (Eggers), *Crossotarsus externedentatus* (Fairmaire), *Liparthrum palaquius* (Schedl), *Scolytomimus pusillus* (Eggers).

Palaquium pseudorostratum

Scolytomimus assamensis Schedl.

Palaquium spp.

Crossotarsus squamulatus Chapuis, *Platypus bifurcus* (Schedl), *Platypus cupulatus* Chapuis, *Platypus hirtellus* Schedl, *Platypus solidus* Walker, *Platypus westwoodi* Chapuis, *Webbia cylindricus* Schedl, *Webbia sara-wakensis* Schedl, *Xyleborus apertus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus major* (Stebbing).

Palaquium stellatum

Anasa foreicollis (Browne), *Anasa versicolor* (Sampson), *Arixyleborus grandis* (Schedl), *Coptoborus bicolor* (Browne), *Coptoborus fragilis* (Browne), *Crossotarsus fractus* Sampson, *Hypocryphalus cylindrus* (Browne), *Scolytomimus pusillus* (Eggers), *Webbia costulatus* Schedl, *Xyleborinus andreweesi* (Blandford), *Xylosandrus ater* (Eggers).

Palisandra rouge

Ambrosiodinus artemgranulatus (Schedl).

Palmetia pinnata

Scolytomimus philippinensis (Eggers).

Panax spp.

Platypus semigranosus (Sampson), *Xyleborinus spiculatus* (Schaufuss), *Xyleborus sakalava* Schedl.

Pancovia harnisiana

Chaetastus tuberculatus (Chapuis), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus jurionii* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Hypothenemus eruditus* Westwood, *Hypothenemus solitarius* (Schedl), *Periommatatus excisus* Strohmeyer, *Periommatatus grandis* Schedl, *Periommatatus substriatus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky), *Xyleborus volculus* (Fabricius).

Pancovia laurentii

Doliopygus conradti (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus jurionii* Schedl, *Doliopygus medius* Schedl, *Doliopygus tenuis* (Strohmeyer), *Eucallacea pandae* (Schedl), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus concolor* Hagedorn, *Periommatatus substriatus* Strohmeyer, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piccus* (Motschulsky), *Xyleborus rothkirchi* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Panda oleosa

Cylindropalpus laudatus (Schedl), *Doliopygus brevis* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus expletus* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus opifex* (Sampson), *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Eucallacea pandae* (Schedl), *Periommatatus excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus schenklingi* (Strohmeyer), *Premnobius mukunyei* (Schedl), *Premnobius nodulosus* Hagedorn, *Scolytotplatypus neglectus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus ficus* Eggers, *Xyleborus subtuberculatus* Eggers, *Xyleborus volculus* (Fabricius).

Panda spp.

Premnobius cavipennis Eichhoff, *Scolytotplatypus kieuensis* Schedl.

Pandanus spp.

Ptilopodius dubiosus Wood.

Pangium spp.

Crossotarsus mnischei Chapuis.

Pausinystalia spp.

Periommatatus mkusii Strohmeyer.

Parabaena sagittata

Eucallacea andamanensis (Blandford).

Parabenzoin praecox

Phloeosinus pulchellus Blandford, *Sucus niisimai* (Eggers), *Xylosandrus brevis* (Eichhoff), *Xylosandrus mutilatus* (Blandford).

Parabenzoin spp.

Scolytotplatypus tycon Blandford.

Paraberlina bifoliolata

Chaetastus striatus Browne.

Paramacrolobium coeruleum

Doliopygus ghesquieri (Schedl), *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus retusus* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus volculus* (Fabricius).

Parartocarpus spp.

Xyleborus amphicranoides Hagedorn, *Xyleborus apertus* Schedl.

Parashorea lucida

Crossotarsus wallacei (Thomson), *Genyocerus diaphanus* (Schedl), *Genyocerus serratus* (Schedl), *Platypus bifurcus* (Schedl), *Platypus curtus* Chapuis.

Parashorea malaenan

Platypus bifurcus (Schedl).

Parashorea plicata

Sphaerotypus quadrituberculatus Sampson.

Parashorea spp.

Genyocerus abdominalis (Schedl), *Platypus bifurcus* (Schedl).

Parashorea stellata

Genyocerus biporus (Schedl).

Parashorea tomentella

Platypus sandakanensis Browne.

Parathesis serrulata

Corthylus mexicanus Schedl.

Parinarium curatellifolium

Doliopygus chapuisi (Duvivier), *Doliopygus intritus* (Schedl).

Parinarium excelsum

Dryocoetoides indolatus Wood, *Taurodemus flavipes* (Fabricius), *Taurodemus splendidus* (Schaufuss).

Parinarium glabra

Platypus intermedius (Schedl), *Platypus spinulosus* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus principalis* Eichhoff.

Parinarium griffithianum

Crossotarsus incrimis Schedl, *Crossotarsus squamulatus* Chapuis, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Xyleborus latus* Eggers.

Parinarium holstii

Doliopygus conradti (Strohmeyer), *Doliopygus donisi* Schedl, *Doliopygus erichsoni* (Chapuis), *Doliopygus tenuis* (Strohmeyer), *Mesoplatypus kitushi* Schedl, *Mesoplatypus quinquecinctus* (Schedl), *Pityophthorus togonus* Eggers, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Platypus rufescens* (Strohmeyer), *Platypus schenklingi* (Strohmeyer), *Xyleborus ambasiipennis* Schedl, *Xyleborus parinarie* Schedl.

Parinarium insularum

Platypus citiensis Roberts.

Parinarium kerstingi

Doliopygus interjectus Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).

Parinarium mobota

Diapus quinquespinatus Chapuis.

***Parinarium* spp.**

Ambrosiodinus sarawakensis (Eggers), *Hypocryphalus striatus* Hopkins, *Leptoxyleborus concisus* (Blandford), *Ozopemon parinarii* Hopkins, *Prennobiobius quadrispinosus* Schedl, *Prennobiobius robustulus* (Schedl).

Parinarium tenuifolium

Doliopygus leferrei Schedl, *Doliopygus rapax* (Sampson).

Parishia insignis

Xyleborus cognatus Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Paritricia excelsa

Taurodemus varians (Fabricius).

Parkia bicolor

Chactastus tuberculatus (Chapuis), *Doliopygus erichsoni* (Chapuis), *Doliopygus expletus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Parkia biglobosa

Triozastus marshalli (Sampson).

Parkia filicoidea

Chactastus tuberculatus (Chapuis), *Doliopygus chapuisi* (Duvivier).

Parkia oliveri

Doliopygus tenuis (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Xyleborus volutus* (Fabricius).

Parkia speciosa

Arixyleborus tuberculatus (Eggers), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus inermis* Schedl, *Crossotarsus schedli* Browne, *Crossotarsus squamulatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Platypus cupulatus* Chapuis, *Platypus insulindicus* Schedl, *Platypus jansonii* Chapuis, *Platypus lepidus* Chapuis, *Platypus pseudocupulatus* Schedl, *Xyleborinus andrewesi* (Blandford), *Xyleborinus exiguus* (Walker), *Xyleborus affinis* Eichhoff, *Xyleborus haberkornii* Eggers, *Xyleborus perforans* (Wollaston).

***Parkia* spp.**

Doliopygus perminutissimus (Schedl), *Scolytoplatypus occidentalis* Browne.

Parkinsonia aculeata

Amphicranus fastigiatus Blandford, *Monarthrum fimbriaticorne* (Blandford).

Parmentiera edulis

Hypothenemus eruditus Westwood, *Hypothenemus gossypii* (Hopkins).

Paropsia schiebeiiana

Doliopygus conradti (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Dryococtoides cristatus* (Fabricius), *Eucallacea pandae* (Schedl), *Hypothenemus biseriatus* (Eggers), *Hypothenemus criticus* (Schedl), *Hypothenemus eruditus* Westwood, *Platypus hintzi* Schaufuss, *Prennobiobius longus* Eggers, *Prennobiobius marginatus* Eggers, *Trachyostus schaufussi* (Strohmeyer).

Parrottia jacquemontiana

Scolytoplatypus sionioi Blandford.

Parthenium argentatum

Pityophthorus mexicanus Blackman.

Parthenocissus quinquefolia

Hypothenemus squamosus (Hopkins).

***Partocarpus* spp.**

Eucallacea arteleacris (Schedl).

Pasania uspericupula

Crossotarsus chalcographus Schedl, *Crossotarsus terminatus* Chapuis.

Pasania cuspidata

Crossotarsus emancipatus Murayama, *Crossotarsus rengetensis* Niisima & Murayama, *Xyleborus atratus* Eichhoff, *Xyleborus exesus* Blandford, *Xyleborus nagaensis* Murayama.

Pasania glabra

Platypus quercivorus Murayama.

Pasania oocarpa

Crossotarsus squamulatus Chapuis.

***Pasania* spp.**

Amasa schlichii (Stebbing), *Amasa versicolor* (Sampson), *Crossotarsus squamulatus* Chapuis, *Cyclorhipidion circumcissum* (Sampson), *Diapus quinquespinatus* Chapuis, *Leptoxyleborus concisus* (Blandford), *Platypus biuncus* Blandford, *Platypus pseudosolidus* Schedl, *Platypus semiernis* Schedl, *Webbia piscecauda* Browne.

Pasania sundaca

Arixyleborus scabripennis (Blandford), *Coptodryas diversicolor* (Eggers), *Crossotarsus longicollis* Browne, *Crossotarsus terminatus* Chapuis, *Cyclorhipidion punctatopilosum* (Schedl), *Diapus quinquespinatus* Chapuis, *Leptoxyleborus concisus* (Blandford), *Platypus convexicauda* Schedl, *Platypus pasanae* Schedl, *Platypus platypoides* (Browne), *Xyleborus ciliatoformis* Schedl, *Xyleborus latus* Eggers.

***Pascoria* spp.**

Hypothenemus ater (Eggers).

Paspalum vaginatum

Hypothenemus californicus Hopkins, *Hypothenemus eruditus* Westwood, *Hypothenemus pubescens* Hopkins.

Passiflora latifolia

Hypothenemus erudiae (Panzer), *Hypothenemus eruditus* Westwood.

Passiflora mollissima

Chramesus hispinus Wood.

Passiflora multiflora

Hypothenemus eruditus Westwood.

***Passiflora* spp.**

Xylosandrus compactus (Eichhoff).

***Passing* spp.**

Xyleborus separandus Schedl.

Paullinia pinnata

Hypothenemus grandis Schedl.

Pausinystalia macroceras

Triozastus caliginosus Roberts.

Pavonia rutshuruensis

Doliopygus erichsoni (Chapuis).

***Payena* spp.**

Crossotarsus wallacei (Thomson).

Pedialanthus macrocarpus

Araupis attenuatus Wood.

Pegauum parmala

Thammurgus brylinskiji Reitter.

Pelargonium hortorum

Amasa amputatus (Blandford).

***Pelea* spp.**

Xyleborus oahuensis Perkins, *Xylosandrus compactus* (Eichhoff).

Pellacalyx saccardianus

Xyleborus similis Ferrari.

***Peltogyne* spp.**

Monarthrum parvum (Eggers).

Peltophorum pterocarpum*Xyleborinus gracilipennis* (Schedl).**Penarahan spp.***Platypus granosus* Brownie.**Pentaclethra eetveldeana***Cylindropalpus camerunus* (Schedl), *Hylesinopsis horridus* (Eggers), *Hypothenemus eruditus* Westwood, *Periommatatus longicollis* Chapuis, *Pityophthorus congonus* Eggers.**Pentaclethra macrophylla***Coptoborus usagaricus* (Eggers), *Diapys quinquespinatus* Chapuis, *Doliopygus bilobatus* (Schedl), *Doliopygus conradti* (Strohmeyer), *Doliopygus donisi* Schedl, *Doliopygus exilis* (Chapuis), *Doliopygus gracilior* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus notatus* Schedl, *Doliopygus propinquus* Schedl, *Dryococtoides cristatus* (Fabricius), *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus submarginatus* Schedl, *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus ater* (Eggers), *Hypothenemus biseriatus* (Eggers), *Hypothenemus concolor* Hagedorn, *Hypothenemus eruditus* Westwood, *Pityophthorus pentaclethrae* Schedl, *Platypus parallelus* (Fabricius), *Platypus spinulosus* (Strohmeyer), *Premnobius mukunyeae* (Schedl), *Premnobius nodulosus* Hagedorn, *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xyleborus ficus* Eggers, *Xyleborus multispinatus* Eggers, *Xyleborus pentaclethrae* Schedl, *Xyleborus principalis* Eichhoff, *Xyleborus rothkirchi* Eggers, *Xyleborus scobinatus* Hagedorn, *Xyleborus subgranosus* Schedl, *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius).**Pentaclethra spp.***Ambrosiodmus tropicus* (Hagedorn).**Pentacme contorta***Sphaerotrypes quadrituberculatus* Sampson.**Pentacme maris***Cryptoxyleborus subnaccus* Schedl.**Pentacme spp.***Platypus bifurcus* (Schedl).**Pentacme suavis***Coptodryas perparva* (Sampson), *Cryptoxyleborus turbineus* (Sampson), *Diapys pendleburyi* Schedl, *Diapys quinquespinatus* Chapuis, *Sphaerotrypes pentacme* Wood, *Sphaerotrypes quadrituberculatus* Sampson.**Pentadesma butyracea***Coccotrypes rotundicollis* (Eggers).**Pentadesma lebrunii***Chaetastus montanus* Schedl, *Coptoborus usagaricus* (Eggers), *Doliopygus erichsoni* (Chapuis), *Platypus solutus* Schedl, *Polygraphus longipilis* Schedl, *Polygraphus tenuipennis* Schedl, *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus ambasi-pennis* Schedl, *Xyleborus principalis* Eichhoff, *Xyleborus splenos* Sampson, *Xyleborus truncatulus* Schedl.**Pentaspadon spp.***Xyleborus apertus* Schedl.**Peptadenia pericillei***Xyleborinus signatipennis* (Schedl).**Peraphyllum ramosissimum***Chaetophloeus heterodoxus* (Casey).**Perebea integrifolia***Taurodemus perebeae* (Ferrari).**Pericopsis elata***Platypus assamela*.**Periploca graeca***Liparthrum arnoldi* Semenov.**Perrottetia sandwicensis***Xyleborinus saxeseni* (Ratzeburg).**Perrottetia spp.***Xyleborus ferrugineus* (Fabricius), *Xylosandrus compactus* (Eichhoff).**Persea americana***Ambrosiodmus aegir* (Eggers), *Araptus mendicus* Wood, *Araptus placetulus* Wood, *Araptus schwarzi* (Blackman), *Cnesinus carinatus* Wood, *Cnesinus costulatus* Blandford, *Cnesinus electinus* Wood, *Cnesinus gracilis* Blandford, *Cnesinus mycelitis* Wood, *Cnesinus prominulus* Wood, *Cnesinus pullus* Blandford, *Cnesinus punctatus* Blandford, *Coccotrypes cyperi* (Beeson), *Corthylocorus aguacatensis* (Schedl), *Corthylus flagellifer* Blandford, *Corthylus papulans* Eichhoff, *Corthylus sanguineus* Schedl, *Crossotarsus externedentatus* (Fairmaire), *Cryptocarenum diadematus* Eggers, *Doliopygus erichsoni* (Chapuis), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins), *Hypothenemus seriatus* (Eichhoff), *Pagiocerus frontalis* (Fabricius), *Phloeocleptus obscurus* Wood, *Platypus madagascariensis* Chapuis, *Platypus solutus* Schedl, *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus aplanatidecicus* Schedl, *Xyleborus perforans* (Wollaston).**Persea borbonia***Ambrosiodmus lecontei* Hopkins, *Cryptocarenum seriatus* Eggers, *Hypothenemus birmanus* (Eichhoff), *Hypothenemus interstitialis* (Hopkins), *Hypothenemus javanus* (Eggers), *Hypothenemus seriatus* (Eichhoff).**Persea grattissima***Coccotrypes dactyliperda* (Fabricius), *Crossotarsus externedentatus* (Fairmaire), *Eucallacea fornicatus* (Eichhoff), *Pityophthorus kiruensis* Schedl, *Platypus biuncus* Blandford, *Platypus cupulatus* Chapuis, *Platypus insulindicus* Schedl, *Xyleborus affinis* Eichhoff, *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).**Persea indica***Hypothenemus longipennis* (Eggers), *Xylosandrus compactus* (Eichhoff).**Persea pittieri***Araptus schwarzi* (Blackman).**Persea popenoi***Monarthrum subgranulatum* Wood.**Persea spp.***Ambrosiodmus obliquus* (LeConte), *Chramesus atkinsoni* Wood, *Eucallacea fornicatus* (Eichhoff), *Pagiocerus frontalis* (Fabricius), *Phloeocleptus ardis* Wood, *Phloeocleptus atkinsoni* Wood, *Phloeocleptus cristatus* Wood, *Phloeocleptus spicatus* Wood, *Pseudothysanoceros perseae* Wood, *Xylosandrus compactus* (Eichhoff).**Perymanium grande***Pityophthorus hermosus* Wood.**Petraea volubilis***Eucallacea fornicatus* (Eichhoff).**Phalocarpus septentrionalis***Hypothenemus birmanus* (Eichhoff), *Hypothenemus eruditus* Westwood.**Phaseolus lunatus***Hypothenemus eruditus* Westwood.

Phaseolus spp.

Chramesus puntilus (Chapuis).

Phellodendron amurense

Crossotarsus niponicus Blandford, *Eucallacea validus* (Eichhoff), *Xyleborus aequalis* (Reitter), *Xyleborus maiche* (Stark), *Xylechinus bergeri* Spessivtzev.

Phellodendron sachalinensis

Eidophelus imitans Eichhoff.

Phellodendron spp.

Scolytoplatypus tycon Blandford.

Phialodiscus hancoensis

Pityophthorus joveri Schedl.

Phialodiscus plurijugatus

Platypus orientalis (Strohmeyer).

Philibertella clausa

Hypothenemus eruditus Westwood, *Hypothenemus seriatus* (Eichhoff).

Phoebe attenuata

Platypus errans (Sampson).

Phoebe hainesiana

Coccotrypes vulgaris (Eggers).

Phoebe lanceolata

Ambrosiodinus lewisi (Blandford), *Phloeosinus phoebe* Wood, *Xyleborus rufobrunneus* Eggers, *Xyleborus shoreae* (Stebbing), *Xylosandrus crassiusculus* (Motschulsky).

Phoebe mexicana

Amphicranus collaris Blandford, *Amphicranus melanura* (Blandford), *Corthylus collaris* Blandford, *Corthylus comatus* Blandford, *Corthylus panamensis* Blandford, *Corthylus retusifera* Wood, *Corthylus retusus* Wood, *Corthylus serratus* Wood, *Corthylus strigilis* Wood, *Monarthrum carinatum* Wood, *Platypus simpliciformis* Wood, *Scolytodes phoebeae* Wood, *Scolytodes piceus* (Blandford), *Tricolus naevus* Wood, *Tricolus ovicollis* Blandford, *Tricolus scitulus* Wood, *Xyleborus meritus* Wood, *Xyleborus prolatus* Wood.

Phoenix canariensis

Coccotrypes dactyliperda (Fabricius), *Dactylotrypes longicollis* (Wollaston).

Phoenix dactylifera

Coccotrypes carpophagus (Homung), *Coccotrypes dactyliperda* (Fabricius), *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston).

Phoenix paludosa

Leptoxyleborus concisus (Blandford).

Phoenix reclinata

Coccotrypes dactyliperda (Fabricius), *Platypus impressus* (Strohmeyer).

Phoenix rupicola

Coccotrypes dactyliperda (Fabricius).

Phoenix spp.

Coccotrypes dactyliperda (Fabricius), *Coccotrypes distinctus* (Motschulsky).

Phoenix sylvestris

Coccotrypes dactyliperda (Fabricius).

Phoradendron bolleanum

Pseudothysanoes funebris Wood.

Phoradendron flavescens

Pseudothysanoes leechii Wood, *Pseudothysanoes phoradendri* Blackman.

Phoradendron longifolium

Pseudothysanoes funereus Wood.

Phoradendron robinsonii

Pseudothysanoes viscicornis Wood.

Phoradendron robustissimum

Chaetophloeus phoradendri Wood, *Pseudothysanoes furvus* Wood, *Pseudothysanoes viscicornis* Wood.

Phoradendron spp.

Chaetophloeus confinis Wood, *Chaetophloeus phoradendri* Wood, *Chaetophloeus struthanthii* Wood, *Chramesus priscus* Wood, *Hypothenemus solocis* Wood, *Hypothenemus squamosus* (Hopkins), *Pseudothysanoes amassius* Wood, *Pseudothysanoes bullatus* Wood, *Pseudothysanoes fimbriatus* Wood, *Pseudothysanoes furvatus* Wood, *Pseudothysanoes furvescens* Wood, *Pseudothysanoes graniticus* Wood, *Pseudothysanoes peniculus* Wood, *Pseudothysanoes tenellus* Wood, *Pseudothysanoes verticilis* Wood, *Xylosandrus curtulus* (Eichhoff).

Photinia japonica

Eucallacea formicatus (Eichhoff).

Phragmites communis

Hypothenemus eruditus Westwood.

Phyllanthus discoideus

Doliopygus dubius (Sampson), *Doliopygus lefevrei* Schedl, *Doliopygus serratus* (Strohmeyer).

Phyllanthus emblica

Xyleborinus artistriatus (Eichhoff).

Phyllanthus flexuosus

Indocryphalus pubipennis (Blandford).

Phyllanthus guineensis

Eucallacea xanthopus (Eichhoff).

Phyllanthus spp.

Hypothenemus grandis Schedl.

Phyllocladus hypophylla

Phloeosinus phyllocladus Bright.

Phyllocladus spp.

Diapus bilunatus Schedl, *Diapus elongatus* Schedl, *Diapus nanus* Schedl, *Diapus nebulosus* Roberts.

Phyllostachys spp.

Scolytoplatypus mikado Blandford.

Phytelephas macrocarpa

Coccotrypes cyperi (Beeson), *Coccotrypes dactyliperda* (Fabricius).

Phytolacca spp.

Eucallacea xanthopus (Eichhoff).

Phytophthora cambivora

Platypus cylindrus (Fabricius), *Xyleborus dryographus* (Ratzeburg).

Piarna spp.

Platypus biuncus Blandford.

Picea abies

Pityophthorus micrographus (Linnaeus), *Pityophthorus tragardhi* Spessivtzev.

Picea ajanensis

Crypturgus tuberosus Niisima, *Hylastes brunneus* Erichson, *Hylastes plumbicus* Blandford, *Orthotomicus starki* Spessivtzev, *Pityogenes foveolatus* Eggers, *Pityogenes sciridensis* Murayama, *Pityophthorus jucundus* Blandford, *Pityophthorus sichotensis* Kurenzov, *Tomiscus puellus* (Reitter).

Picea asperata

Cryphalus myialopiceus Tsai & Li, *Cryphalus saltuarius* Weise, *Hylurgops eusulcatus* Tsai & Huang, *Polygraphus angustus* Tsai & Yin, *Polygraphus squameus* Yin & Huang, *Polygraphus szecmaensis* Tsai & Yin, *Polygraphus zhongdianensis* Tsai & Yin, *Scolytus sinopiceus* Tsai, *Xyleborinus saxenii* (Ratzeburg).

Picea chihuahuana

Pityophthorus bassetti Blackman, *Pseudothysanoes conferae* (Wood).

Picea engelmannii

Carphoborus carri Swaine, *Carphoborus intermedius* Wood, *Carphoborus piceae* Wood, *Carphoborus sansoni* Swaine, *Cryphalus ruficollis* Hopkins, *Crypturgus borealis* Swaine, *Dolurgus pumilus* (Mannerheim), *Dryocoetes caryi* Hopkins, *Dryocoetes confusus* Swaine, *Gnathotrichus denticulatus* Blackman, *Gnathotrichus retusus* (LeConte), *Hylastes macer* LeConte, *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mannerheim), *Hylurgops subcostulatus subcostulatus* (Mannerheim), *Ips borealis swainei* Hopping, R., *Ips borealis lanieri* Wood, *Ips hunteri* Swaine, *Ips perturbatus* (Eichhoff), *Ips pilifrons pilifrons*, *Ips pilifrons utahensis* Wood, *Ips pilifrons sulcifrons* Wood, *Ips tridens engelmanni* Swaine, *Phlocotribus lecontei* Schedl, *Pityokteines lasiocarpi* (Swaine), *Pityophthorus aquilus* Blackman, *Pityophthorus bassetti* Blackman, *Pityophthorus carinulatus* Swaine, *Pityophthorus murrayanae* Blackman, *Pityophthorus nitidulus* (Mannerheim), *Pityophthorus nitidus* Swaine, *Pityophthorus occidentalis* Blackman, *Pityophthorus opaculus* LeConte, *Pityophthorus pseudotsugae* Swaine, *Pityophthorus recens* Bright, *Pityophthorus tuberculatus* Eichhoff, *Polygraphus convexifrons* Wood, *Polygraphus hoppingi* Swaine, *Polygraphus rufipennis* (Kirby), *Scierus annectens* LeConte, *Scierus pubescens* Swaine, *Scolytus opaculus* Blackman, *Trypodendron rufitarsis* (Kirby), *Xylechinus montanus* Blackman.

Picea excelsa

Carphoborus rossicus Semenov, *Carphoborus teplouchovi* Spessivtsev, *Cryphalus asperatus* (Gyllenhal), *Cryphalus piceae* (Ratzeburg), *Cryphalus piceus* Eggers, *Cryphalus saltuarius* Weise, *Dendroctonus micans* (Kugelann), *Hylastes angustatus* (Herbst), *Hylastes brunneus* Erichson, *Hylastes cunicularius* Erichson, *Hylastes plumbicus* Blandford, *Hylurgops glabratus* (Zetterstedt), *Hylurgops inouyei* Nobuchi, *Hylurgops palliatus* (Gyllenhal), *Ips amittinus* (Eichhoff), *Ips duplicatus* (Sahlberg), *Orthotomicus starki* Spessivtsev, *Phlocotribus spinulosus* (Rey), *Pityogenes saalasi* Eggers, *Pityophthorus araki* (Sawamoto), *Pityophthorus excelsus* (Ratzeburg), *Pityophthorus micrographus* (Linnaeus), *Pityophthorus morosovi* Spessivtsev, *Pityophthorus pityographus* (Ratzeburg), *Pityophthorus tragardhi* Spessivtsev, *Trypodendron laeae* Eggers.

Picea glauca

Carphoborus andersoni Swaine, *Carphoborus carri* Swaine, *Carphoborus sansoni* Swaine, *Cryphalus ruficollis* Hopkins, *Crypturgus borealis* Swaine, *Dendroctonus punctatus* LeConte, *Dryocoetes caryi* Hopkins, *Dryocoetes granicollis* (LeConte), *Ips borealis borealis* Swaine, *Ips borealis thomasi* Hopping, G., *Ips borealis lanieri* Wood, *Ips perturbatus* (Eichhoff), *Ips tridens tridens*, *Ips tridens engelmanni* Swaine, *Phlocosinus pini* Swaine, *Phlocotribus piceae* Swaine, *Pityophthorus balsameus* Blackman, *Pityophthorus bassetti* Blackman, *Pityophthorus biovalis* Blackman, *Pityophthorus cariniceps* LeConte, *Pityophthorus cascoensis* Blackman, *Pityophthorus concaeus* Blackman, *Pityophthorus consimilis* LeConte, *Pityophthorus dentifrons* Blackman, *Pityophthorus intextus* Swaine, *Pityophthorus murrayanae* Blackman, *Pityophthorus nitidulus* (Mannerheim), *Pityophthorus nitidus* Swaine,

Pityophthorus opaculus LeConte, *Pityophthorus recens* Bright, *Polygraphus convexifrons* Wood, *Polygraphus rufipennis* (Kirby), *Scierus annectens* LeConte, *Xylechinus americanus* Blackman.

Picea glehnii

Cryphalus piceae (Ratzeburg), *Cryphalus piceus* Eggers, *Crypturgus tuberosus* Niisima, *Hylurgops inouyei* Nobuchi, *Polygraphus gracilis* Niisima, *Xyleborus septentrionalis* Niisima.

Picea jezoensis

Cryphalus jezoensis Inouye & Nobuchi, *Cryphalus laticis* Niisima, *Cryphalus niponensis* Inouye & Nobuchi, *Cryphalus piceae* (Ratzeburg), *Cryphalus piceus* Eggers, *Cryphalus sichotensis* Kurenzov, *Crypturgus tuberosus* Niisima, *Dryocoetes nijimai* Nobuchi, *Hylastes cunicularius* Erichson, *Hylastes plumbicus* Blandford, *Hylurgops glabratus* (Zetterstedt), *Hylurgops inouyei* Nobuchi, *Hylurgops interstitialis* (Chapuis), *Hylurgops longipilus* (Reitter), *Hylurgops spessivtsevi* Eggers, *Hylurgops interstitialis* (Chapuis), *Hylurgops palliatus* (Gyllenhal), *Ips duplicatus* (Sahlberg), *Pityogenes forcolatus* Eggers, *Pityogenes scirindensis* Murayama, *Polygraphus gracilis* Niisima, *Polygraphus proximus* Blandford, *Polygraphus sachalinensis* Eggers, *Polygraphus squamulatus* Niisima, *Tomiscus puellus* (Reitter), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus septentrionalis* Niisima.

Picea koraiensis

Carphoborus taiwensis Murayama, *Carphoborus teplouchovi* Spessivtsev, *Crypturgus tuberosus* Niisima, *Hylurgops glabratus* (Zetterstedt), *Pityophthorus morosovi* Spessivtsev, *Pityophthorus pini* Kurenzov, *Polygraphus proximus* Blandford, *Polygraphus sachalinensis* Eggers.

Picea likiangensis

Polygraphus junanensis Sokanovskii.

Picea mariana

Crypturgus alutaceus Schwarz, *Pityophthorus biovalis* Blackman, *Pityophthorus briscoei* Blackman, *Pityophthorus nitidus* Swaine.

Picea microsperma

Hylurgops palliatus (Gyllenhal).

Picea morinda

Crossotarsus conferae Stebbing, *Cryphalus major* Stebbing, *Dryocoetes indicus* Stebbing, *Ips stebbingi* Strohmeier, *Pityogenes spessivtsevi* Lebedev, *Polygraphus major* Stebbing, *Polygraphus pini* Stebbing, *Scolytoplatypus raja* Blandford, *Scolytoplatypus siomio* Blandford.

Picea morrissonicola

Polygraphus taiwanensis Schedl.

Picea obovata

Carphoborus teplouchovi Spessivtsev, *Cryphalus piceus* Eggers, *Cryphalus saltuarius* Weise, *Hylastes brunneus* Erichson, *Hylastes cunicularius* Erichson, *Hylastes plumbicus* Blandford, *Hylurgops glabratus* (Zetterstedt), *Hylurgops spessivtsevi* Eggers, *Ips duplicatus* (Sahlberg), *Orthotomicus starki* Spessivtsev, *Pityogenes rudnei* Sokanovskii, *Pityogenes saalasi* Eggers, *Pityogenes scirindensis* Murayama, *Pityophthorus flavimaculatus* Murayama, *Pityophthorus lapponicus* Stark, *Pityophthorus lichtensteini* (Ratzeburg), *Pityophthorus micrographus* (Linnaeus), *Pityophthorus morosovi* Spessivtsev, *Pityophthorus pini* Kurenzov, *Pityophthorus tragardhi* Spessivtsev, *Polygraphus sachalinensis* Eggers.

Picea omorika

Pityophthorus micrographus (Linnaeus), *Pityophthorus pityographus* (Ratzeburg).

Picea orientalis

Cryphalus saltuarius Weise, *Pityophthorus pityographus* (Ratzeburg).

Picea pungens

Carphoborus brevisetosus Wood, *Carphoborus sansoni* Swaine, *Ips hunteri* Swaine, *Ips pilifrons sulcifrons* Wood, *Phloeotribus lecontei* Schedl, *Pityokteines ornatus* (Swaine), *Pityophthorus bassetti* Blackman, *Pityophthorus carinulatus* Swaine, *Pityophthorus murrayanae* Blackman, *Pityophthorus occidentalis* Blackman, *Pityophthorus opaculus* LeConte, *Pityophthorus ornatus* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Polygraphus rufipennis* (Kirby).

Picea rubens

Carphoborus carri Swaine, *Carphoborus dumii* Swaine, *Cryphalus rubentis* Hopkins, *Cryphalus ruficollis* Hopkins, *Crypturgus borealis* Pjatuinskii, *Dendroctonus punctatus* LeConte, *Dryocoetes caryi* Hopkins, *Pityophthorus angustus* Blackman, *Pityophthorus bioculis* Blackman, *Pityophthorus briscoli* Blackman, *Pityophthorus dentifrons* Blackman, *Polygraphus rufipennis* (Kirby), *Xylechinus americanus* Blackman.

Picea schrenkiana

Ips hauseri Reitter, *Pityogenes spessivtzevi* Lebedev, *Pityophthorus kirgisicus* Pjatuinskii, *Pityophthorus parfentievi* Pjatuinskii, *Pityophthorus schrenkianae* Pjatuinskii.

Picea sitchensis

Cryphalus pubescens Hopkins, *Dendroctonus punctatus* LeConte, *Dolurgus pumilus* (Mannerheim), *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mannerheim), *Ips concinnus* (Mannerheim), *Ips plastographus maritimus* Lanier, *Ips tridens tridens*, *Ips tridens engelmanni* Swaine, *Pityophthorus nitidulus* (Mannerheim), *Pseudohylesinus sitchensis* Swaine.

Picea spinulosa

Trypodendron dorjitenzingi Schmutzenhofer.

***Picea* spp.**

Carphoborus sansoni Swaine, *Cryphalus latus* Eggers, *Crypturgus cinereus* (Herbst), *Crypturgus hispidulus* Thomson, *Crypturgus pusillus* (Gyllenhal), *Crossotarsus piccae* Stebbing, *Dendroctonus rufipennis* Kirby, *Dendroctonus valens* LeConte, *Dryocoetes affaber* (Mannerheim), *Dryocoetes autographus* (Ratzeburg), *Dryocoetes formosanus* Nobuchi, *Dryocoetes hectographus* Reitter, *Dryocoetes pinii* Niisima, *Dryocoetes ruficollis* Eggers, *Gnathotrichus materiarius* (Fitch), *Hylastes angusticollis* Eggers, *Hylastes ater* (Paykull), *Hylastes nigrinus* (Mannerheim), *Hylastes opacus* Erichson, *Hylurgops imitator* (Reitter), *Hylurgops porosus* (LeConte), *Hylurgops rugipennis pinifex* Fitch, *Hylurgops subcostulatus subcostulatus* (Mannerheim), *Hypothenemus interstitialis* (Hopkins), *Ips acuminatus* (Gyllenhal), *Ips orientalis* Wood & Yin, *Ips sexdentatus* (Boemer), *Ips typographus* (Linnaeus), *Orthotomicus caclatus* (Eichhoff), *Orthotomicus laricus* (Fabricius), *Orthotomicus proximus* (Eichhoff), *Orthotomicus suturalis* (Gyllenhal), *Pityogenes bidentatus* (Herbst), *Pityogenes chalcographus* (Linnaeus), *Pityogenes quadridens* (Hartig), *Pityokteines curvidens* (Germar), *Pityophthorus balsameus* Blackman, *Pityophthorus carinatus carinatus* Bright, *Pityophthorus consimilis* LeConte, *Pityophthorus lautus* Eichhoff,

Pityophthorus puberulus (LeConte), *Pityophthorus pulicarius* (Zimmermann), *Platypus xylographus* Schedl, *Polygraphus jezoensis* Niisima, *Polygraphus kisoensis* Niisima, *Polygraphus poligraphus* (Linnaeus), *Polygraphus rufipennis* (Kirby), *Polygraphus subopacus* Thomson, *Scolytoplatypus tycon* Blandford, *Scolytus piccae* (Swaine), *Tomicus minor* (Hartig), *Tomicus piniperda* (Linnaeus), *Trypodendron lineatum* (Olivier), *Xyleborinus saxeseni* (Ratzeburg), *Xylechinus montanus* Blackman, *Xylechinus pilosus* (Ratzeburg), *Xyloterinus politus* (Say).

***Pimeleodendron* spp.**

Platypus advena Schedl, *Platypus fulgidus* Browne.

Pinus albicaulis

Dendroctonus ponderosae Hopkins, *Ips latidens* (LeConte), *Ips montanus* (Eichhoff), *Pityogenes fossifrons* (LeConte), *Pityophthorus absonus* Blackman, *Pityophthorus aquilus* Blackman, *Pityophthorus confertus* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus murrayanae* Blackman, *Pityophthorus toralis* Wood.

Pinus aristata

Carphoborus declivis Wood, *Conophthorus ponderosae* Hopkins, *Pityophthorus absonus* Blackman, *Pityophthorus aquilus* Blackman, *Pityophthorus boycei* Swaine, *Pityophthorus crassus* Blackman, *Pityophthorus jeffreyi* Blackman, *Pityophthorus mormoni* Bright, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman.

Pinus armandii

Cryphalus chidingensis Tsai & Li, *Cryphalus lipingensis* Tsai & Li, *Cryphalus pseudochinlingensis* Tsai & Li, *Dendroctonus armandi* Tsai & Li, *Hylastes techangensis* Tsai & Huang, *Hylurgops junnanicus* Sokanovskii, *Hylurgops longipilus* (Reitter), *Hylurgops major* Eggers, *Polygraphus junnanicus* Sokanovskii, *Polygraphus sinensis* Eggers, *Polygraphus squameus* Yin & Huang, *Polygraphus verrucifrons* Tsai & Yin, *Polygraphus zhungdianensis* Tsai & Yin, *Tomicus pilifer* (Spessivtsev), *Xyleborinus saxeseni* (Ratzeburg).

Pinus attenuata

Hylurgops porosus (LeConte), *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mannerheim), *Ips mexicanus* (Hopkins), *Ips paraconfusus* Lanier, *Pityokteines ornatus* (Swaine), *Pityophthorus californicus* Bright, *Pityophthorus carneli* Swaine, *Pityophthorus tuberculatus* Eichhoff.

Pinus ayacahuite

Dendroctonus adjunctus Blandford, *Dendroctonus approximatus* Dietz, *Dendroctonus mexicanus* Hopkins, *Hylurgops planirostris* (Chapuis), *Pityoborus secundus* Blackman, *Pityophthorus abstrusus* Bright, *Pityophthorus annexens* LeConte, *Pityophthorus brevis* Blackman, *Pityophthorus crassus* Blackman, *Pityophthorus cristatus* Wood, *Pityophthorus cuspidatus* Blackman, *Pityophthorus delicatus* Wood, *Pityophthorus discretus* Wood, *Pityophthorus festus* Wood, *Pityophthorus glabratus* (Schedl), *Pityophthorus impexus* Bright, *Pityophthorus montivagus* Bright, *Pityophthorus nigricans* Blandford, *Pityophthorus scabridus* Schedl, *Pityophthorus solus* Blackman.

Pinus balfouriana

Dendroctonus ponderosae Hopkins, *Hylurgops rugipennis pinifex* Fitch, *Ips montanus* (Eichhoff), *Pityogenes fossifrons* (LeConte), *Pityophthorus absonus* Blackman.

Pinus banksiana

Conophthorus resinosae Hopkins, *Cryphalus fulvus* Niisima, *Dendroctonus murrayanae* Hopkins, *Ips perroti* Swaine, *Ips pilifrons thatcheri* Wood, *Ips pini* (Say), *Phloeosinus pini* Swaine, *Pityogenes plagiatus* (LeConte), *Pityophthorus balsameus* Blackman, *Pityophthorus briscoli* Blackman, *Pityophthorus cariniceps* LeConte, *Pityophthorus cavatus* Bright, *Pityophthorus concavus* Blackman, *Pityophthorus consimilis* LeConte, *Pityophthorus dentifrons* Blackman, *Pityophthorus puberulus* (LeConte), *Pityophthorus pulchellus* Eichhoff, *Trypodendron rufitarsis* (Kirby), *Trypodendron scabricollis* (LeConte).

Pinus brutia

Pityogenes pennidens (Reitter).

Pinus canariensis

Crypturgus concolor (Wollaston), *Hylastes lowei* Paiva, *Ips nobilis* (Wollaston).

Pinus caribaea

Pityoborus hondurensis Wood, *Pityophthorus annectens* LeConte, *Pityophthorus ocellus* Bright, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus scabridus* Schedl, *Xyleborinus sharpae* (Hopkins).

Pinus cembra

Hylurgops glabratus (Zetterstedt), *Pityogenes conjunctus* Reitter, *Pityophthorus lichtensteini* (Ratzeburg).

Pinus cembroides

Carphoborus convexifrons Wood, *Carphoborus pini-colens* Wood, *Conophthorus edulis* Hopkins, *Ips hoppingi* Lanier, *Pityophthorus arcanus* Bright, *Pityophthorus barberi* Blackman, *Pityophthorus confertus* Swaine, *Pityophthorus crassus* Blackman, *Pityophthorus deletus* LeConte, *Pityophthorus keeni* (Blackman), *Pityophthorus lecontei* Bright, *Pityophthorus leiophyllae* Blackman, *Pityophthorus modicus* Blackman, *Pityophthorus schuicarzi* Blackman, *Pityophthorus subopacus* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus zonalis* Bright, *Pityotrichus barbatus* (Blackman).

Pinus chiapensis

Gnathotrichus perniciosus Wood.

Pinus chihuahuana

Platypus pini Hopkins.

Pinus clausa

Carphoborus bicornis Wood.

Pinus contorta

Carphoborus declivis Wood, *Carphoborus intermedius* Wood, *Carphoborus ponderosae* Swaine, *Conophthorus ponderosae* Hopkins, *Crypturgus borealis* Swaine, *Dendroctonus murrayanae* Hopkins, *Dendroctonus ponderosae* Hopkins, *Dolurgus pumilus* (Mannerheim), *Gnathotrichus retusus* (LeConte), *Hylastes longicollis* Swaine, *Hylurgops porosus* (LeConte), *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mannerheim), *Hylurgops rugipennis pinifex* Fitch, *Hylurgops subcostatus subcostulatus* (Mannerheim), *Ips emarginatus* (LeConte), *Ips latidens* (LeConte), *Ips pini* (Say), *Ips plastographus maritimus* Lanier, *Phloeosinus antennatus* Swaine, *Pityogenes knecheli* Swaine, *Pityokteines ornatus* (Swaine), *Pityophthorus absomus* Blackman, *Pityophthorus aquilus* Blackman, *Pityophthorus boycei* Swaine, *Pityophthorus carinatus monticolae* Bright, *Pityophthorus carinulatus* Swaine, *Pityophthorus confertus* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus crassus* Blackman, *Pityophthorus digestus* (LeConte), *Pityophthorus fuscus* Black-

man, *Pityophthorus indigus* Wood, *Pityophthorus mormon* Bright, *Pityophthorus murrayanae* Blackman, *Pityophthorus nitidulus* (Mannerheim), *Pityophthorus occidentalis* Blackman, *Pityophthorus scalptor* Blackman, *Pityophthorus toralis* Wood, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman, *Pseudohylesinus pini* Wood, *Sciurus annectens* LeConte, *Trypodendron rufitarsis* (Kirby), *Xyleborus intrusus* Blandford.

Pinus cooperi

Gnathotrichus nitidifrons Hopkins, *Ips mexicanus* (Hopkins), *Pityophthorus crassus* Blackman, *Pityophthorus rubidus* Wood, *Pityophthorus thomasi* Bright.

Pinus coulteri

Dendroctonus brevicornis LeConte, *Dendroctonus ponderosae* Hopkins, *Ips paraconfusus* Lanier, *Pityophthorus carmeli* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus electus* Blackman, *Pityophthorus scalptor* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman, *Xyleborus intrusus* Blandford.

Pinus culminicola

Pityophthorus crassus Blackman, *Pityophthorus culminicola* Bright, *Pityophthorus megas* Bright, *Pityophthorus riminalis* Bright.

Pinus deusiflora

Ambrosiodmus rubricollis (Eichhoff), *Cryphalus fulvus* Niisima, *Cryphalus jeholensis* Murayama, *Cryphalus laricis* Niisima, *Cryphalus piceae* (Ratzeburg), *Cryphalus oblongus* Niisima, *Dryocoetes uniseriatus* Eggers, *Eucallacca validus* (Eichhoff), *Hylastes parallelus* Chapuis, *Hylastes plumbus* Blandford, *Hylurgops interstitialis* (Chapuis), *Orthotomicus multidentatus* (Murayama), *Orthotomicus tosaensis* (Murayama), *Pityophthorus jucundus* Blandford, *Polygraphus kisoensis* Niisima, *Xyleborus aquilus* Blandford, *Xyleborus atratus* Eichhoff, *Xyleborus festinus* Eichhoff, *Xyleborus septentrionalis* Niisima.

Pinus discolor

Conophthorus edulis Hopkins.

Pinus douglasiana

Conophthorus ponderosae Hopkins.

Pinus durangensis

Conophthorus ponderosae Hopkins, *Dendroctonus rhizophagus* Thomas & Bright, *Hylastes gracilis* LeConte, *Hylurgops incomptus* (Blandford), *Ips mexicanus* (Hopkins), *Pityophthorus crassus* Blackman, *Pityophthorus festus* Wood.

Pinus echinata

Carphoborus bifurcus Eichhoff, *Conophthorus echinatae* Wood, *Crypturgus alutaceus* Schwarz, *Crypturgus borealis* Swaine, *Dendroctonus frontalis* Zimmermann, *Dendroctonus terebrans* (Olivier), *Gnathotrichus materiarius* (Fitch), *Pityogenes meridianus* Blackman, *Pityogenes plagiatus* (LeConte), *Pityophthorus confusus* Blandford, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus pullus* (Zimmermann), *Trypodendron scabricollis* (LeConte), *Xyleborus pubescens* Zimmermann.

Pinus edulis

Carphoborus convexifrons Wood, *Carphoborus pini-colens* Wood, *Conophthorus edulis* Hopkins, *Dendroctonus ponderosae* Hopkins, *Hylastes fulgidus* Blackman, *Hylastes gracilis* LeConte, *Ips confusus* (LeConte), *Ips latidens* (LeConte), *Pityokteines ornatus* (Swaine), *Pityophthorus annectens* LeConte, *Pityophthorus barberi* Blackman, *Pityophthorus blandus* Black-

man, *Pityophthorus brevis* Blackman, *Pityophthorus confertus* Swaine, *Pityophthorus crassus* Blackman, *Pityophthorus deletus* LeConte, *Pityophthorus keeni* (Blackman), *Pityophthorus lecontei* Bright, *Pityophthorus modicus* Blackman, *Pityophthorus punctifrons* Bright, *Pityophthorus schwarzi* Blackman, *Pityophthorus solus* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus woodi* Bright, *Pityotrichus barbatus* (Blackman).

Pinus eliottii

Ambrosiodmus catharinensis (Eggers), *Ambrosiodmus hagedornii* (Iglesias), *Carphoborus bifurcus* Eichhoff, *Dendroctonus terebrans* (Olivier), *Hylastes exilis* Chapuis, *Pityophthorus annectens* LeConte, *Pityophthorus anticus* Schedl, *Pityophthorus concentricus* Eichhoff, *Pityophthorus confusus* Blandford, *Pityophthorus pinavornus* Bright, *Theoborus villosulus* (Blandford), *Xyleborinus gracilis* (Eichhoff), *Xyleborinus linearicollis* (Schedl), *Xyleborinus schoenherrii* (Schedl), *Xyleborus adelographus* Eichhoff.

Pinus engelmannii

Carphoborus pinicoleus Wood, *Conophthorus apacheae* Hopkins, *Dendroctonus approximatus* Dietz, *Dendroctonus frontalis* Zimmermann, *Dendroctonus rhizophagus* Thomas & Bright, *Hylurgops incomptus* (Blandford), *Pityophthorus annectens* LeConte, *Pityophthorus crassus* Blackman, *Pityophthorus cristatus* Wood, *Pityophthorus intentus* Bright, *Pityophthorus rubidus* Wood, *Pityophthorus schuerdtfergeri* (Schedl), *Pityophthorus seguis* Blackman.

Pinus excelsa

Crossotarsus fairmairei Chapuis, *Cryphalus major* Stebbing, *Cryphalus strommeyeri* Stebbing, *Dryocoetes indicus* Stebbing, *Hylastes brunneus* Erichson, *Ips stebbingi* Strohmeier, *Pityogenes scitus* Blandford, *Pityogenes spessittsevi* Lebedev, *Pityophthorus deodara* (Stebbing), *Polygraphus pini* Stebbing, *Scolytus major* Stebbing.

Pinus flexilis

Carphoborus pinicoleus Wood, *Conophthorus ponderosae* Hopkins, *Dendroctonus ponderosae* Hopkins, *Gnathotrichus retusus* (LeConte), *Hylurgops porosus* (LeConte), *Hylurgops rugipennis pinifex* Fitch, *Ips latidens* (LeConte), *Ips woodi* Thatcher, *Pityogenes fossifrons* (LeConte), *Pityophthorus absonus* Blackman, *Pityophthorus acutus* Blackman, *Pityophthorus amplus* (Blackman), *Pityophthorus aquilus* Blackman, *Pityophthorus boycei* Swaine, *Pityophthorus confertus* Swaine, *Pityophthorus crassus* Blackman, *Pityophthorus deletus* LeConte, *Pityophthorus grandis* Blackman, *Pityophthorus mormon* Bright, *Pityophthorus murrayanae* Blackman, *Pityophthorus occidentalis* Blackman, *Pityophthorus pinguis* (Blackman), *Pityophthorus toralis* Wood, *Pityophthorus tuberculatus* Eichhoff, *Pityotrichus hesperius* Bright.

Pinus gerardiana

Carphoborus zhohi (Stebbing), *Ips stebbingi* Strohmeier, *Pityogenes scitus* Blandford, *Pityogenes spessittsevi* Lebedev, *Pityophthorus chilgoza* Wood, *Polygraphus major* Stebbing, *Polygraphus trenchii* Stebbing.

Pinus glabra

Dendroctonus frontalis Zimmermann.

Pinus greggii

Carphoborus costatus Wichmann, *Pityophthorus crassus* Blackman, *Pityophthorus pellitus* Schedl, *Pityophthorus schwarzi* Blackman, *Pityophthorus solus* Blackman.

Pinus griffithii

Carphoborus costatus Wichmann, *Polygraphus aterrinus* Strohmeier, *Polygraphus major* Stebbing, *Polygraphus setosus* Schedl, *Polygraphus squameus* Yin & Huang, *Polygraphus chungdianensis* Tsai & Yin, *Trypodendron dorjitenzingi* Schmutzenhofer.

Pinus halepensis

Carphoborus bonnairi Brisout, *Carphoborus marani* Pfeffer, *Carphoborus pini* Eichhoff, *Crypturgus mediterraneus* Eichhoff, *Crypturgus numidicus* Ferrari, *Hylastes linearis* Erichson, *Hylurgus micklitzii* Wachtl, *Ips erosus* (Wollaston), *Orthotomicus proximus* (Eichhoff), *Pityogenes calcaratus* (Eichhoff), *Pityogenes porifrons* Eggers, *Pityophthorus pubescens* (Marshall).

Pinus hartwegii

Dendroctonus adjunctus Blandford, *Dendroctonus approximatus* Dietz, *Ips mexicanus* (Hopkins), *Pityogenes mexicanus* Wood, *Pityophthorus blandulus* Schedl, *Pityophthorus cortezi* Bright, *Pityophthorus crassus* Blackman, *Pityophthorus furuissi* Bright, *Pityophthorus pubifrons* Bright, *Pityophthorus solus* Blackman, *Pseudotsyanoxenus pini* Wood.

Pinus heldreichii

Pityophthorus balcanicus Pfeffer, *Pityophthorus henscheli* Seitner, *Pityophthorus pityographus* (Ratzeburg).

Pinus hondurensis

Xyleborus affinis Eichhoff.

Pinus insularis

Ambrosiodmus camphorae (Hagedorn), *Ambrosiodmus inominatus* (Schedl), *Ambrosiodmus triton* (Schaufuss), *Coccotrypes recticollis* (Schedl), *Cryphalus malloti* Schedl, *Cryphalus sylvicola* (Perkins), *Cyrtogenius luteus* (Blandford), *Diaprus vulgassicus* Schedl, *Hylastes techangensis* Tsai & Huang, *Mitosoma excisum* Schaufuss, *Mitosoma onconiceps* Schedl, *Mitosoma obliquatum* Schedl, *Platypus piniperla* Schedl, *Platypus tenellus* Schedl, *Polygraphus junnanicus* Sokanovskii, *Polygraphus szemaecensis* Tsai & Yin, *Scolytos platypus pusillus* Eggers, *Tomicus brevipilosus* (Eggers), *Xyleborinus spinifer* (Eggers), *Xyleborus antaisaka* Schedl, *Xyleborus madagascariensis* Schaufuss, *Xyleborus pinicola* Eggers, *Xyleborus sakalava* Schedl, *Xyleborus submolestus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Pinus jeffreyi

Cactopinus pini Blackman, *Carphoborus simplex* LeConte, *Conophthorus ponderosae* Hopkins, *Dendroctonus jeffreyi* Hopkins, *Dendroctonus ponderosae* Hopkins, *Gnathotrichus retusus* (LeConte), *Hylurgops porosus* (LeConte), *Hylurgops reticulatus* Wood, *Hylurgops subcostatus subcostulatus* (Mannerheim), *Ips emarginatus* (LeConte), *Ips latidens* (LeConte), *Ips paraconfusus* Lanier, *Ips pilifrons thatcheri* Wood, *Ips pini* (Say), *Pityogenes carinulatus* (LeConte), *Pityokteines ornatus* (Swaine), *Pityophthorus confertus* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus digestus* (LeConte), *Pityophthorus electus* Blackman, *Pityophthorus jeffreyi* Blackman, *Pityophthorus scalptor* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman, *Trypodendron rufitarsis* (Kirby), *Xyleborus ferrugineus* (Fabricius), *Xyleborus intrusus* Blandford.

Pinus koraiensis

Cryphalus fulvus Niisima, *Cryphalus laricis* Niisima, *Cryphalus piccus* Eggers, *Dryocoetes baicalicus* Reitter, *Dryocoetes infuscatus* Murayama, *Hylastes brunneus* Erichson, *Hylastes cunicularius* Erichson, *Hylastes*

plumbeus Blandford, *Hylurgops interstitialis* (Chapuis), *Hylurgops longipilus* (Reitter), *Hylurgops spessirtsevi* Eggers, *Hylurgops transbaicalicus* Eggers, *Pityogenes foreolatus* Eggers, *Pityogenes irkutensis* Eggers, *Pityophthorus lapponicus* Stark, *Pityophthorus pini* Kirenzov, *Polygraphus sachalinensis* Eggers, *Scolytus moravici* Semenov, *Tomicus brevipilosus* (Eggers), *Tomicus pilifer* (Spessirtsev), *Tomicus puellus* (Reitter), *Xyleborinus saxesenii* (Ratzeburg).

Pinus lambertiana

Cactopinus koebelei Blackman, *Carphoborus pinicolens* Wood, *Conophthorus ponderosae* Hopkins, *Cryphalus pubescens* Hopkins, *Dendroctonus ponderosae* Hopkins, *Gnathotrichus retusus* (LeConte), *Hylurgops rugipennis rugipennis* (Mammerheim), *Ips latidens* (LeConte), *Ips paraconfusus* Lanier, *Pityophthorus confinis* LeConte, *Pityophthorus dolus* Wood, *Pityophthorus pseudotsugae* Swaine.

Pinus laevis

Dendroctonus mexicanus Hopkins, *Pityophthorus annectens* LeConte, *Pityophthorus cuspidatus* Blackman, *Pityophthorus durus* Blackman, *Pityophthorus lepidus* Bright, *Pityophthorus miniatus* Bright, *Pityophthorus nocturnus* Schedl, *Pityophthorus obtusipennis* Blandford, *Pityophthorus pellitus* Schedl, *Pityophthorus solatus* Wood, *Pityophthorus solus* Blackman.

Pinus leiophylla

Carphoborus convexifrons Wood, *Carphoborus pinicolens* Wood, *Conophthorus conicolens* Wood, *Conophthorus ponderosae* Hopkins, *Dendroctonus adjunctus* Blandford, *Dendroctonus approximatus* Dietz, *Dendroctonus mexicanus* Hopkins, *Dendroctonus parallelocollis* Chapuis, *Gnathotrichus delongi* Blackman, *Gnathotrichus nitidifrons* Hopkins, *Gnathotrichus perniciosus* Wood, *Gnathotrichus sulcatus* (LeConte), *Hylurgops incomptus* (Blandford), *Hylurgops planirostris* (Chapuis), *Hylurgops subcostulatus alternans* (Chapuis), *Ips mexicanus* (Hopkins), *Pityoborus secundus* Blackman, *Pityophthorus aciculatus* Bright, *Pityophthorus acutus* Blackman, *Pityophthorus amnectens* LeConte, *Pityophthorus barberi* Blackman, *Pityophthorus clarus* Blackman, *Pityophthorus crassus* Blackman, *Pityophthorus cristatus* Wood, *Pityophthorus delicatus* Wood, *Pityophthorus durus* Blackman, *Pityophthorus festus* Wood, *Pityophthorus grandis* Blackman, *Pityophthorus intentus* Bright, *Pityophthorus leiophyllae* Blackman, *Pityophthorus obtusipennis* Blandford, *Pityophthorus perotei* Blackman, *Pityophthorus rubidus* Wood, *Pityophthorus schwarzi* Blackman, *Pityophthorus solus* Blackman, *Pityophthorus spadix* Blackman, *Pityophthorus subopacus* Blackman, *Pityophthorus venustus* Blackman, *Platypus pini* Hopkins, *Xyleborus intrusus* Blandford.

Pinus leucodermis

Crypturgus numidicus Ferrari, *Ips robustus* (Knotek), *Pityogenes bistridentatus* (Eichhoff).

Pinus luchuensis

Orthotomicus kunijoshii Nobuchi, *Xyleborus atratus* Eichhoff.

Pinus lungholtzii

Carphoborus mexicanus Bright, *Pityophthorus crassus* Blackman, *Pityophthorus cristatus* Wood, *Pityophthorus cuspidatus* Blackman, *Pityophthorus rubidus* Wood, *Pityophthorus schwarzi* Blackman.

Pinus massoniana

Cryphalus massonians Tsai & Li, *Cyrtogenius luteus*

(Blandford), *Eucallacca interjectus* (Blandford), *Eucallacca validus* (Eichhoff), *Hylurgops longipilus* (Reitter), *Hylurgops major* Eggers, *Xyleborus aquilus* Blandford, *Xyleborus pinicola* Eggers.

Pinus maximowiczii

Cryphalus furukawai Murayama, *Xyleborus muticus* Blandford.

Pinus merkusii

Platypus biflexuosus Schedl, *Platypus cavus* Strohmeyer, *Polygraphus sumatranus* Schedl, *Xyleborus pfeili* (Ratzeburg).

Pinus mexicana

Xyleborus intrusus Blandford.

Pinus michoacana

Conophthorus michoacanae Wood, *Pityophthorus confusus* Blandford, *Pityophthorus durus* Blackman, *Pityophthorus pellitus* Schedl, *Pityophthorus subopacus* Blackman.

Pinus monophylla

Cactopinus koebelei Blackman, *Cactopinus pini* Blackman, *Carphoborus frontalis* Wood, *Carphoborus pinicolens* Wood, *Conophthorus monophyllae* Hopkins, *Dendroctonus ponderosae* Hopkins, *Hylastes gracilis* LeConte, *Ips confusus* (LeConte), *Pityophthorus arcanus* Bright, *Pityophthorus blandus* Blackman, *Pityophthorus confinis* LeConte, *Pityophthorus deletus* LeConte, *Pityophthorus grandis* Blackman, *Pityophthorus keenii* (Blackman), *Pityophthorus modicus* Blackman, *Pityophthorus punctifrons* Bright, *Pityophthorus tuberculatus* Eichhoff.

Pinus montana

Cryphalus piceus Eggers, *Hylurgops glabratus* (Zetterstedt), *Pityogenes conjunctus* Reitter.

Pinus montezumae

Conophthorus ponderosae Hopkins, *Dendroctonus adjunctus* Blandford, *Dendroctonus approximatus* Dietz, *Dendroctonus mexicanus* Hopkins, *Dendroctonus parallelocollis* Chapuis, *Gnathotrichus delongi* Blackman, *Gnathotrichus nitidifrons* Hopkins, *Gnathotrichus perniciosus* Wood, *Gnathotrichus sulcatus* (LeConte), *Hylastes gracilis* LeConte, *Hylastes retifer* Wood, *Ips bonansea* (Hopkins), *Ips mexicanus* (Hopkins), *Pityophthorus arcuobii* Wood, *Pityophthorus crassus* Blackman, *Pityophthorus dispar* Bright, *Pityophthorus micans* Bright, *Pityophthorus montezumae* Bright, *Pityophthorus pellitus* Schedl, *Pityophthorus scabridus* Schedl, *Pityophthorus schwarzi* Blackman, *Pityophthorus solus* Blackman, *Pityophthorus subopacus* Blackman, *Pseudothysanoes pini* Wood.

Pinus monticola

Conophthorus ponderosae Hopkins, *Dendroctonus ponderosae* Hopkins, *Hylastes longicollis* Swaine, *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mammerheim), *Ips emarginatus* (LeConte), *Ips latidens* (LeConte), *Ips montanus* (Eichhoff), *Pityogenes fossifrons* (LeConte), *Pityophthorus absorus* Blackman, *Pityophthorus boycei* Swaine, *Pityophthorus confertus* Swaine, *Pityophthorus venustus* Blackman, *Pityophthorus toralis* Wood, *Scolytus monticolae* (Swaine), *Trypodendron rufitarsis* (Kirby).

Pinus mugo

Hylurgops glabratus (Zetterstedt), *Pityophthorus glabratus* Eichhoff, *Pityophthorus henscheli* Seigner, *Pityophthorus knoteki* Reitter, *Pityophthorus pubescens* (Marshall).

Pinus mugo rotundata

Pityophthorus glabratus Eichhoff.

Pinus muricata

Dolburgus pumilus (Mannerheim), *Hylurgops reticulatus* Wood, *Hylurgops rugipennis rugipennis* (Mannerheim), *Ips mexicanus* (Hopkins), *Ips plas-tographus maritimus* Lanier, *Pityophthorus carneli* Swaine, *Pityophthorus nitidulus* (Mannerheim), *Pityophthorus setosus* Blackman, *Pseudohylesinus pini* Wood.

Pinus niger

Pityogenes trepanatus (Nordlinger).

Pinus nigra

Cryphalus jeholensis Murayama, *Orthotomicus pini-rora* Schedl, *Pityogenes bistridentatus* (Eichhoff), *Pityogenes conjunctus* Reitter, *Pityophthorus balcanicus* Pfeffer, *Pityophthorus buyssoni* Reitter, *Pityophthorus carniolicus* Wichmann, *Pityophthorus glabratus* Eichhoff, *Pityophthorus lichtensteini* (Ratzeburg), *Pityophthorus pubescens* (Marshall).

Pinus nigra mauretanicus

Pityophthorus mauretanicus Peyerimhoff.

Pinus nigricans

Ips mannsfeldi (Wachtl).

Pinus obovata

Carphoborus rossicus Semenov.

Pinus occidentalis

Ambrosiodinus lecontei Hopkins, *Hypothenemus eruditus* Westwood, *Pityophthorus antillicus* Bright, *Pityophthorus pinavorus* Bright.

Pinus ochotereuui

Ambrosiodinus funebris (Schedl), *Pityophthorus attenuatus* Blackman, *Pityophthorus blandulus* Schedl, *Pityophthorus confusus* Blandford, *Pityophthorus dispar* Bright, *Pityophthorus glabratus* (Schedl), *Platypus pini* Hopkins.

Pinus oocarpa

Araptus sobrinus Wood, *Dendroctonus frontalis* Zimmermann, *Dendroctonus mexicanus* Hopkins, *Dendroctonus parallelocollis* Chapuis, *Gnathotrichus perniciosus* Wood, *Pityoborus frontalis* Wood, *Pityoborus hondurensis* Wood, *Pityoborus secundus* Blackman, *Pityophthorus blandulus* Schedl, *Pityophthorus cacuminatus* Blandford, *Pityophthorus confusus* Blandford, *Pityophthorus crassus* Blackman, *Pityophthorus delicatus* Wood, *Pityophthorus festus* Wood, *Pityophthorus euterpes* Bright, *Pityophthorus miniatus* Bright, *Pityophthorus nocturnus* Schedl, *Pityophthorus ocellus* Bright, *Pityophthorus pellitus* Schedl, *Pityophthorus scabridus* Schedl, *Pityophthorus schweirdtfegeri* (Schedl), *Pityophthorus solus* Blackman, *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).

Pinus palustris

Crypturgus alutaceus Schwarz, *Dendroctonus frontalis* Zimmermann, *Dendroctonus terebrans* (Olivier), *Hylastes exilis* Chapuis, *Pityophthorus annectens* LeConte, *Pityophthorus confusus* Blandford, *Pityophthorus pulchellus* Eichhoff, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus pullus* (Zimmermann), *Xyleborus ferrugineus* (Fabricius), *Xyleborus pubescens* Zimmermann.

Pinus parviflora

Eucallacea validus (Eichhoff), *Hylurgops interstitialis* (Chapuis), *Tomicus brevipilosus* (Eggers).

Pinus patula

Hylurgops longipennis (Blandford), *Hylurgops planirostris* (Chapuis), *Mitosoma excisum* Schauffuss, *Pityophthorus ciliatus* Blackman, *Pityophthorus crassus*

Blackman, *Pityophthorus montivagus* Bright, *Pityophthorus scabridus* Schedl, *Xyleborus spinulosus* Blandford.

Pinus peuce

Pityophthorus knoteki Reitter, *Pityophthorus lichtensteini* (Ratzeburg), *Pityophthorus pityographus* (Ratzeburg).

Pinus pinaster

Carphoborus bonnairei Brisout, *Crypturgus mediterraneus* Eichhoff, *Crypturgus numidicus* Ferrari, *Hylastes angustatus* (Herbst), *Hylastes linearis* Erichson, *Ips crosus* (Wollaston), *Pityogenes calcaratus* (Eichhoff), *Pityogenes porifrons* Eggers, *Pityophthorus pubescens* (Marshall).

Pinus pinea

Pityogenes pennidens (Reitter), *Pityophthorus pubescens* (Marshall).

Pinus pithusa

Crypturgus numidicus Ferrari.

Pinus ponderosa

Carphoborus costatus Wichmann, *Carphoborus perplexus* Wood, *Carphoborus pinicolens* Wood, *Carphoborus ponderosae* Swaine, *Carphoborus simplex* LeConte, *Conophthorus ponderosae* Hopkins, *Crypturgus borealis* Swaine, *Dendroctonus adjunctus* Blandford, *Dendroctonus approximatus* Dietz, *Dendroctonus brevicornis* LeConte, *Dendroctonus frontalis* Zimmermann, *Dendroctonus jeffreyi* Hopkins, *Dendroctonus mexicanus* Hopkins, *Dendroctonus ponderosae* Hopkins, *Gnathotrichus denticulatus* Blackman, *Gnathotrichus initalis* Wood, *Gnathotrichus materiarius* (Fitch), *Gnathotrichus retusus* (LeConte), *Gnathotrichus sulcatus* (LeConte), *Hylastes fulgidus* Blackman, *Hylastes gracilis* LeConte, *Hylastes longicollis* Swaine, *Hylastes macer* LeConte, *Hylastes nigricans* (Mannerheim), *Hylurgops porosus* (LeConte), *Hylurgops reticulatus* Wood, *Hylurgops subcostatus subcostulatus* (Mannerheim), *Hylurgops subcostulatus alternans* (Chapuis), *Ips bonansea* (Hopkins), *Ips emarginatus* (LeConte), *Ips knausi* Swaine, *Ips latidens* (LeConte), *Ips lecontei* Swaine, *Ips paracensus* Lanier, *Ips pilifrons thacheri* Wood, *Ips pini* (Say), *Pityoborus secundus* Blackman, *Pityogenes carinulatus* (LeConte), *Pityojekines ornatus* (Swaine), *Pityophthorus acutus* Blackman, *Pityophthorus amplus* (Blackman), *Pityophthorus annectens* LeConte, *Pityophthorus aquilus* Blackman, *Pityophthorus arcanus* Bright, *Pityophthorus barberi* Blackman, *Pityophthorus blandus* Blackman, *Pityophthorus boycei* Swaine, *Pityophthorus brevis* Blackman, *Pityophthorus clarus* Blackman, *Pityophthorus comosus* Blackman, *Pityophthorus confertus* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus crassus* Blackman, *Pityophthorus cuspidatus* Blackman, *Pityophthorus deletus* LeConte, *Pityophthorus digestus* (LeConte), *Pityophthorus dolus* Wood, *Pityophthorus electus* Blackman, *Pityophthorus grandis* Blackman, *Pityophthorus immanis* Blackman, *Pityophthorus insulatus* Blackman, *Pityophthorus ingens* Blackman, *Pityophthorus intentus* Bright, *Pityophthorus jeffreyi* Blackman, *Pityophthorus lecontei* Bright, *Pityophthorus leechi* Wood, *Pityophthorus levis* Wood, *Pityophthorus murrayanae* Blackman, *Pityophthorus rubidus* Wood, *Pityophthorus scalptor* Blackman, *Pityophthorus sculptus* Bright, *Pityophthorus schwarzi* Blackman, *Pityophthorus schweirdtfegeri* (Schedl), *Pityophthorus serratus* Swaine, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman, *Pityophthorus zonalis* Bright, *Pityotrichus barbatus* (Blackman), *Platypus*

pini Hopkins, *Trypodendron rufitarsis* (Kirby), *Xyleborus intrusus* Blandford.

Pinus pringlei

Pityophthorus durus Blackman, *Pityophthorus glabratus* (Schedl).

Pinus pseudostrobus

Corthylus mexicanus Schedl, *Dendroctonus adjunctus* Blandford, *Dendroctonus mexicanus* Hopkins, *Dendroctonus vitei* Wood, *Gnathotrichus perniciosus* Wood, *Gnathotrichus sulcatus* (LeConte), *Hylurgops incomptus* (Blandford), *Hylurgops planirostris* (Chapuis), *Ips bonansea* (Hopkins), *Pityoborus hirtellus* Wood, *Pityoborus secundus* Blackman, *Pityophthorus aciculatus* Bright, *Pityophthorus delicatus* Wood, *Pityophthorus diglyphus* Blandford, *Pityophthorus glabratus* (Schedl), *Pityophthorus minutus* Bright, *Pityophthorus nocturnus* Schedl, *Pityophthorus pellitus* Schedl, *Pityophthorus scabridus* Schedl, *Pityophthorus schuerdtfegeri* (Schedl), *Pityophthorus spadix* Blackman, *Platypus pini* Hopkins.

Pinus pumila

Crypturgus tuberosus Niisima, *Hylurgops spessitzevi* Eggers, *Pityogenes foveolatus* Eggers, *Pityogenes japonicus* Nobuchi, *Polygraphus kisoensis* Niisima, *Polygraphus meakanensis* Niisima.

Pinus quadrifolia

Pityophthorus punctifrons Bright.

Pinus radiata

Carphoborus radiatae Swaine, *Conophthorus radiatae* Hopkins, *Dolurgus pumilus* (Mannerheim), *Gnathotrichus retusus* (LeConte), *Hylurgops porosus* (LeConte), *Hylurgops reticulatus* Wood, *Hylurgops rugipennis* *rugipennis* (Mannerheim), *Ips mexicanus* (Hopkins), *Ips plastographus maritimus* Lanier, *Pityophthorus californicus* Bright, *Pityophthorus carmeli* Swaine, *Pityophthorus murrayanae* Blackman, *Pityophthorus nitidulus* (Mannerheim), *Pityophthorus pubescens* (Marshall), *Pityophthorus setosus* Blackman, *Pityophthorus trepidus* Bright.

Pinus resinosa

Conophthorus resinosae Hopkins, *Hylurgops rugipennis* *pinifex* Fitch, *Ips perroti* Swaine, *Pityogenes plagiatus* (LeConte), *Pityophthorus amectens* LeConte, *Pityophthorus balsameus* Blackman, *Pityophthorus cariniceps* LeConte, *Pityophthorus cavatus* Bright, *Pityophthorus concavus* Blackman, *Pityophthorus consimilis* LeConte, *Pityophthorus intextus* Swaine, *Pityophthorus nitidus* Swaine, *Pityophthorus puberulus* (LeConte), *Pityophthorus pulchellus* Eichhoff, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus ramiperda* Swaine, *Trypodendron scabricollis* (LeConte).

Pinus rigida

Cryphalus fulvus Niisima, *Dendroctonus frontalis* Zimmermann, *Dendroctonus terebrans* (Olivier), *Pityophthorus consimilis* LeConte, *Pityophthorus pulchellus* Eichhoff, *Pityophthorus pulicarius* (Zimmermann), *Xyleborinus saxeseni* (Ratzeburg).

Pinus roxburghii

Carphoborus costatus Wichmann, *Coccotrypes nubilis* (Blandford), *Crossotarsus wilnoti* (Stebbing), *Cryphalus major* Stebbing, *Hylurgus indicus* Wood, *Ips longifolia* (Stebbing), *Liparthrum longifolia* (Stebbing), *Pityogenes scitus* Blandford, *Pityogenes spessitzevi* Lebedev, *Pityophthorus deodara* (Stebbing), *Platypus bififormis* Chapuis, *Polygraphus aterrimus* Strohmeier, *Polygraphus difficilis* Wood, *Polygraphus longifolia*

Stebbing, *Polygraphus setosus* Schedl, *Platypus bififormis* Chapuis.

Pinus rudis

Dendroctonus adjunctus Blandford, *Dendroctonus approximatus* Dietz, *Dendroctonus mexicanus* Hopkins, *Pityophthorus blandulus* Schedl, *Pityophthorus glabratus* (Schedl), *Pityophthorus nigricans* Blandford, *Pityophthorus pellitus* Schedl, *Pityophthorus scabridus* Schedl, *Platypus pini* Hopkins, *Xyleborus intrusus* Blandford.

Pinus sabiniana

Carphoborus simplex LeConte, *Ips spinifer* (Eichhoff), *Pityophthorus confinis* LeConte, *Pityophthorus modicus* Blackman, *Pityophthorus novellus* Blackman, *Pityophthorus tuberculatus* Eichhoff.

Pinus serotina

Dendroctonus terebrans (Olivier).

Pinus siberica

Pityophthorus lapponicus Stark.

Pinus spp.

Ambrosiodmus apicalis (Blandford), *Ambrosiodmus fraterculus* (Schaufuss), *Ambrosiodmus rusticus* (Wood), *Carphoborus bifurcus* Eichhoff, *Carphoborus minimus* (Fabricius), *Carphoborus pinicolens* Wood, *Coccotrypes cinnamomi* (Eggers), *Coccotrypes medius* (Eggers), *Conophthorus coniperda* (Schwarz), *Conophthorus mexicanus* Wood, *Conophthorus resinosae* Hopkins, *Crypturgus cinereus* (Herbst), *Crypturgus hispidulus* Thomson, *Crypturgus pusillus* (Gyllenhal), *Dendroctonus brevicornis* LeConte, *Dendroctonus micans* (Kugelann), *Dendroctonus parallelocollis* Chapuis, *Dendroctonus valens* LeConte, *Dryocoetes affaber* (Mannerheim), *Dryocoetes autographus* (Ratzeburg), *Dryocoetes hectographus* Reitter, *Dryocoetes pini* Niisima, *Dryocoetes rugicollis* Eggers, *Gnathotrichus deleari* Blackman, *Gnathotrichus denticulatus* Blackman, *Gnathotrichus imitans* Wood, *Gnathotrichus perniciosus* Wood, *Gnathotrichus sulcatus* (LeConte), *Hylastes asperatus* Wood, *Hylastes ater* (Paykull), *Hylastes attenuatus* Erichson, *Hylastes exilis* Chapuis, *Hylastes flohri* (Eggers), *Hylastes fulgidus* Blackman, *Hylastes gracilis* LeConte, *Hylastes linearis* Erichson, *Hylastes macer* LeConte, *Hylastes mexicanus* Wood, *Hylastes niger* Wood, *Hylastes nigrinus* (Mannerheim), *Hylastes opacus* Erichson, *Hylastes parallelus* Chapuis, *Hylastes porculus* Erichson, *Hylastes salebrosus* Eichhoff, *Hylastes tenuis* Eichhoff, *Hylocurus beckeri* Heqvist, *Hylocurus rivalis* Wood, *Hylurgops glabratus* (Zetterstedt), *Hylurgops imitator* (Reitter), *Hylurgops incomptus* (Blandford), *Hylurgops interstitialis* (Chapuis), *Hylurgops longipennis* (Blandford), *Hylurgops palliatus* (Gyllenhal), *Hylurgops planirostris* (Chapuis), *Hylurgops porosus* (LeConte), *Hylurgops subcostatus* *subcostulatus* (Mannerheim), *Hylurgops subcostulatus alternans* (Chapuis), *Hylurgus ligniperda* (Fabricius), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Ips acuminatus* (Gyllenhal), *Ips amitinus* (Eichhoff), *Ips avulsus* (Eichhoff), *Ips bonansea* (Hopkins), *Ips calligraphus* (Germar), *Ips cembrae* (Heer), *Ips confusus* (LeConte), *Ips emarginatus* (LeConte), *Ips grandicollis* (Eichhoff), *Ips integer* (Eichhoff), *Ips sexdentatus* (Boerner), *Ips typographus* (Linnaeus), *Monarthrum fasciatum* (Say), *Orthotomicus caelatus* (Eichhoff), *Orthotomicus laricis* (Fabricius), *Orthotomicus longicollis* (Gyllenhal), *Orthotomicus proximus* (Eichhoff), *Orthotomicus suturalis* (Gyllenhal), *Pityoborus comatus* (Zimmermann), *Pityoborus fron-*

talis Wood, *Pityoborus hirtellus* Wood, *Pityoborus rubentis* Wood, *Pityoborus securinus* Blackman, *Pityoborus relictus* Wood, *Pityogenes bidentatus* (Herbst), *Pityogenes carinulatus* (LeConte), *Pityogenes chalcographus* (Linnaeus), *Pityogenes irkutensis* Eggers, *Pityogenes quadridens* (Hartig), *Pityokteines curvidens* (Germar), *Pityophthorus abstrusus* Bright, *Pityophthorus annectens* LeConte, *Pityophthorus aztecus* Bright, *Pityophthorus cacuminatus* Blandford, *Pityophthorus chalcocensis* Hopkins, *Pityophthorus ciliatus* Blackman, *Pityophthorus consimilis* LeConte, *Pityophthorus cristatus* Wood, *Pityophthorus declivisetosus* Bright, *Pityophthorus deleoni* (Blackman), *Pityophthorus delicatus* Wood, *Pityophthorus discretus* Wood, *Pityophthorus durus* Blackman, *Pityophthorus excelsus* (Ratzeburg), *Pityophthorus festus* Wood, *Pityophthorus impexus* Bright, *Pityophthorus ineditus* Bright, *Pityophthorus ingens* Blackman, *Pityophthorus jeffreyi* Blackman, *Pityophthorus laticeps* Bright, *Pityophthorus leiophyllae* Blackman, *Pityophthorus lepidus* Bright, *Pityophthorus litos* Bright, *Pityophthorus minus* Bright, *Pityophthorus nitidus* Swaine, *Pityophthorus obtusipennis* Blandford, *Pityophthorus ornatus* Blackman, *Pityophthorus pellitus* Schedl, *Pityophthorus pullus* (Zimmermann), *Pityophthorus rudis* Blackman, *Pityophthorus sapineus* Bright, *Pityophthorus separatus* Bright, *Pityophthorus solatus* Wood, *Pityophthorus solers* Blackman, *Pityophthorus solus* Blackman, *Pityophthorus spadix* Blackman, *Pityophthorus subopacus* Blackman, *Pityophthorus subsimilis* Schedl, *Pityophthorus vespertinus* Bright, *Platypus flavicornis* (Fabricius), *Platypus madagascariensis* Chapuis, *Platypus pini* Hopkins, *Platypus xylographus* Schedl, *Polygraphus horjuncensis* Murayama, *Polygraphus poligraphus* (Linnaeus), *Polygraphus proximus* Blandford, *Polygraphus subopacus* Thomson, *Polygraphus szemacensis* Tsai & Yin, *Polygraphus verrucifrons* Tsai & Yin, *Scolytoplatypus tycon* Blandford, *Tomicus destruens* (Vollaston), *Tomicus minor* (Hartig), *Tomicus piniperda* (Linnaeus), *Trischichia atomus* (Hopkins), *Xyleborinus saxoseni* (Ratzeburg), *Xyleborus atratus* Eichhoff, *Xyleborus dispar* (Fabricius), *Xyleborus curygraphus* (Ratzeburg), *Xyleborus pinivorus* Brownie, *Xyleborus pubescens* Zimmermann, *Xyleborus seriatus* Blandford, *Xylosandrus germanus* (Blandford), *Xyloterinus politus* (Say).

Pinus strobiformis

Conophthorus ponderosae Hopkins, *Dendroctonus ponderosae* Hopkins, *Dendroctonus terebrans* (Olivier), *Gnathotrichus denticulatus* Blackman, *Ips woodi* Thatcher, *Pityogenes fossifrons* (LeConte), *Pityophthorus acutus* Blackman, *Pityophthorus amplus* (Blackman), *Pityophthorus brevicornatus* Bright, *Pityophthorus brevis* Blackman, *Pityophthorus citrus* Bright, *Pityophthorus confertus* Swaine, *Pityophthorus confinis* LeConte, *Pityophthorus crassus* Blackman, *Pityophthorus cuspidatus* Blackman, *Pityophthorus deletus* LeConte, *Pityophthorus durus* Blackman, *Pityophthorus immanis* Blackman, *Pityophthorus infundatus* Blackman, *Pityophthorus intentus* Bright, *Pityophthorus nigricans* Blandford, *Pityophthorus pinguis* (Blackman), *Pityophthorus seignis* Blackman, *Pityophthorus tuberculatus* Eichhoff, *Pityophthorus venustus* Blackman, *Pityotrachelus hesperis* Bright.

Pinus strobus

Conophthorus coniperda (Schwarz), *Dendroctonus frontalis* Zimmermann, *Dendroctonus murrayanae*

Hopkins, *Dendroctonus terebrans* (Olivier), *Gnathotrichus materiarius* (Fitch), *Hylurgops inouyei* Nobuchi, *Hylurgops rugipennis* pilifex Fitch, *Ips latidens* (LeConte), *Ips pilifrons* thatcheri Wood, *Ips pini* (Say), *Pityogenes hopkinsi* Swaine, *Pityophthorus angustus* Blackman, *Pityophthorus balsameus* Blackman, *Pityophthorus bioalis* Blackman, *Pityophthorus briseoi* Blackman, *Pityophthorus carinatus* carinatus Bright, *Pityophthorus cariniceps* LeConte, *Pityophthorus consimilis* LeConte, *Pityophthorus dentifrons* Blackman, *Pityophthorus laudus* Eichhoff, *Pityophthorus nitidus* Swaine, *Pityophthorus puberulus* (LeConte), *Pityophthorus pulchellus* Eichhoff, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus pullus* (Zimmermann), *Pityophthorus ramipeda* Swaine, *Polygraphus rufipennis* (Kirby).

Pinus strobus chiapensis

Pityophthorus nocturnus Schedl, *Pityophthorus obtusipennis* Blandford, *Pityophthorus subsimilis* Schedl.

Pinus sylvestris

Carphoborus cholodkovskiy Spessivtsev, *Carphoborus jurinskii* Eggers, *Cryphalus laricis* Nüssima, *Hylastes angustatus* (Herbst), *Hylastes bruniceus* Erichson, *Hylastes cunicularius* Erichson, *Hylurgops glabratus* (Zetterstedt), *Hylurgops inouyei* Nobuchi, *Ips duplicatus* (Sahlberg), *Ips mannsfeldi* (Wachtl), *Ips robustus* (Knotek), *Pityogenes calcaratus* (Eichhoff), *Pityogenes conjunctus* Reitter, *Pityogenes irkutensis* Eggers, *Pityogenes trepanatus* (Nordlinger), *Pityophthorus buyssoni* Reitter, *Pityophthorus carniolicus* Wichmann, *Pityophthorus glabratus* Eichhoff, *Pityophthorus lichtensteini* (Ratzeburg), *Pityophthorus opaculus* LeConte, *Pityophthorus puberulus* (LeConte), *Pityophthorus pubescens* (Marshall), *Pityophthorus pulicarius* (Zimmermann), *Polygraphus seriatus* Reitter, *Xyleborus klinganensis* Murayama.

Pinus tabulaeformis

Cryphalus jeholensis Murayama, *Cryphalus pseudochinlingensis* Tsai & Li, *Cryphalus pseudotabulaeformis* Tsai & Li, *Cryphalus tabulaeformis* Tsai & Li, *Hylastes techangensis* Tsai & Huang, *Hylurgops junmanicus* Sokanovskii, *Hylurgops longipilus* (Reitter), *Hylurgops major* Eggers, *Pityophthorus scirpus* Murayama, *Polygraphus junmanicus* Sokanovskii, *Polygraphus sinensis* Eggers, *Polygraphus squameus* Yin & Huang, *Polygraphus szemacensis* Tsai & Yin, *Polygraphus verrucifrons* Tsai & Yin, *Tomicus pilifer* (Spessivtsev), *Xyleborus emarginatus* Eichhoff.

Pinus tabulaeformis mukdensis

Cryphalus fulvus Nüssima.

Pinus taeda

Carphoborus bifurcus Eichhoff, *Crypturgus alutaceus* Schwarz, *Dendroctonus frontalis* Zimmermann, *Dendroctonus terebrans* (Olivier), *Gnathotrichus materiarius* (Fitch), *Hylastes exilis* Chapuis, *Hypothenemus crudiae* (Pauper), *Pityogenes meridianus* Blackman, *Pityophthorus annectens* LeConte, *Pityophthorus confusus* Blandford, *Pityophthorus consimilis* LeConte, *Pityophthorus pulicarius* (Zimmermann), *Pityophthorus pullus* (Zimmermann), *Trypodendron scabricollis* (LeConte), *Xyleborus ferrugineus* (Fabricius).

Pinus taicaiensis

Encallacea validus (Eichhoff).

Pinus tenuifolia

Dendroctonus adjunctus Blandford, *Dendroctonus mexicanus* Hopkins, *Dendroctonus parallelocollis* Chapuis, *Hyllocurus clarki* Wood, *Pityophthorus delicatus*

Wood, *Pityophthorus festus* Wood, *Pityophthorus nocturnus* Schedl, *Pityophthorus pellitus* Schedl, *Pityophthorus subsimilis* Schedl, *Xyleborus affinis* Eichhoff.

Pinus teocote

Conophthorus teocotum Wood, *Dendroctonus approximatus* Dietz, *Dendroctonus mexicanus* Hopkins, *Dendroctonus vitci* Wood, *Pityophthorus crassus* Blackman, *Pityophthorus leiophyllae* Blackman, *Pityophthorus perotei* Blackman, *Pityophthorus schuarzi* Blackman.

Pinus thunbergiana

Cryphalus fulvus Nisima, *Cryphalus furukawai* Murayama, *Cryphalus jicholensis* Murayama, *Cryphalus laricis* Nisima, *Eucallacca validus* (Eichhoff), *Orthotomicus nankinensis* Kirenzov & Kononov, *Orthotomicus tosaensis* (Murayama), *Pityophthorus jucundus* Blandford.

Pinus torreyana

Pityophthorus carmeli Swaine.

Pinus tuberculata

Dolargus pumilus (Mannerheim).

Pinus virginiana

Carphoborus bicornis Wood, *Carphoborus bifurcus* Eichhoff, *Dendroctonus frontalis* Zimmermann, *Hylurgops rugipennis pinifex* Fitch, *Pityogenes plagiatus* (LeConte), *Pityophthorus cariniceps* LeConte, *Pityophthorus consimilis* LeConte, *Pityophthorus pulchellus* Eichhoff, *Pityophthorus pulicarius* (Zimmermann).

Pinus wachaoensis

Conophthorus ponderosae Hopkins, *Pityophthorus digestus* (LeConte).

Pinus yunnanensis

Cryphalus szechuanensis Tsai & Li, *Hylastes techangensis* Tsai & Huang, *Hylurgops major* Eggers, *Ips chinensis* Kurenzov & Kononov, *Polygraphus junnanicus* Sokanovskii, *Polygraphus szenaeensis* Tsai & Yin, *Polygraphus verrucifrons* Tsai & Yin, *Xyleborus emarginatus* Eichhoff.

Piper guahameuse

Cryphalus sylvicola (Perkins).

Piper spp.

Cuestus suturalis (Eggers), *Corthylus cirritus* Wood, *Corthylus comosus* Wood, *Hadrodemius globus* (Blandford), *Xyleborus haberkornii* Eggers.

Piptadenia africana

Chaetastus tuberculatus (Chapuis), *Doliopygus chapuisi* (Duxivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus exilis* (Chapuis), *Doliopygus eximius* Schedl, *Doliopygus expletus* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus gracilior* Schedl, *Doliopygus lateralis* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus perminutissimus* (Schedl), *Doliopygus propinquus* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus spectabilis* Schedl, *Doliopygus subdolosus* Schedl, *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Hypothenemus ater* (Eggers), *Hypothenemus biserialis* (Eggers), *Periommatius longicollis* Chapuis, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Piptadenia buehananii

Cryptocorus spinipennis Schedl, *Doliopygus exilis* (Chapuis).

Piptadenia percillei

Xyleborus alluaudi Schaufuss, *Xyleborus malgasicus* Schedl.

Piptadenia spp.

Mesoplatus quinquenectus (Schedl), *Pseudochramesus semibrunneus* (Eggers).

Piptadeniastrium africanum

Chaetastus montanus Schedl, *Doliopygus opifex* (Sampson), *Doliopygus perminutissimus* (Schedl), *Doliopygus propinquus* Schedl, *Periommatius grandis* Schedl, *Triozastus banghaasi* (Schaufuss), *Triozastus caliginosus* Roberts, *Triozastus elongatus* Schedl, *Triozastus pilosulus* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus barumbucensis* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Piptadeniastrium spp.

Prennobiobius cavipennis Eichhoff.

Pipturus albidus

Xyleborus ferrugineus (Fabricius), *Xyleborus hawaiiensis* Perkins, *Xyleborus perforans* (Wollaston).

Pipturus spp.

Xyleborinus saxeseni (Ratzeburg), *Xylosandrus compactus* (Eichhoff), *Xylosandrus morigerus* (Blandford).

Piscidia piscipala

Cnemidophorus vagabundus (Wood).

Pisonia spp.

Xyleborus ferrugineus (Fabricius).

Pisonia umbellifera

Xyleborus ferrugineus (Fabricius).

Pistacia atlantica

Chaetoptelius vestitus (Mulsant & Rey).

Pistacia integerrima

Chaetoptelius vestitus (Mulsant & Rey).

Pistacia lentiscus

Chaetoptelius vestitus (Mulsant & Rey), *Phloeotribus scarabacoides* (Bernard).

Pistacia spp.

Carphoborus perrisi (Chapuis), *Phloeotribus mayeti* (Guillebeau).

Pistacia terebenthus

Chaetoptelius vestitus (Mulsant & Rey).

Pistacia vera

Chaetoptelius vestitus (Mulsant & Rey).

Pithecolobium bubalinum

Cyclorhipidion foersteri (Hagedorn).

Pithecolobium dulce

Xyleborus perforans (Wollaston).

Pithecolobium guadeloupense

Cryptocarus seriatus Eggers, *Hypothenemus hirsutus* (Wood), *Hypothenemus seriatus* (Eichhoff), *Micracisella nanula* (LeConte).

Pithecolobium lobatum

Xyleborinus andrewesi (Blandford), *Xyleborus setulosus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Pithecolobium pinnatum

Xyleborus asper Eggers, *Xyleborus ferrugineus* (Fabricius).

Pithecolobium saman

Ambrosiodinus pithecolobius (Schedl), *Platypus suffodiens* Sampson.

Pithecolobium splendens

Crossotarsus externodentatus (Fairmaire).

***Pithecolobium* spp.**

Crossotarsus squamulatus Chapuis, *Crossotarsus terminatus* Chapuis, *Cyclorhipidion pruinatum* (Blandford), *Platypus lepidus* Chapuis, *Platypus pseudocupulatus* Schedl, *Xylosandrus compactus* (Eichhoff).

Pithecellobium unguis-cati

Hypothenemus hirsutus (Wood), *Hypothenemus squamosus* (Hopkins).

Pittosporum resiniferum

Cryphalus resiniferi Schedl.

Pittosporum spathicalyx

Euwallacea xanthopus (Eichhoff).

***Pittosporum* spp.**

Ambrosiodinus compressus (Lea), *Cyrtogenius dimorphus* (Schedl), *Euwallacea xanthopus* (Eichhoff), *Hypothenemus seriatus* (Eichhoff), *Mitosoma excisum* Schaufuss, *Xyleborinus quadrispinus* (Schedl), *Xyleborinus spiculatus* (Schaufuss), *Xyleborinus spinosus* (Schaufuss), *Xyleborus alluaudi* Schaufuss, *Xylosandrus compactus* (Eichhoff).

Pittosporum tobira

Cryphalus birosinensis (Murayama).

Pittosporum viridiflorum

Chaetastus montanus Schedl, *Triozastus caliginosus* Roberts.

***Plagioscyphus* spp.**

Crossotarsus nitens Chapuis, *Platypus madagascariensis* Chapuis, *Prennobilus catipennis* Eichhoff.

Planchonella laurifolia

Platypus semigranulosus (Sampson).

Planchonella pohlmaniana

Xyleborus perforans (Wollaston), *Xyleborus volutus* (Fabricius).

Planchonella samoensis

Cryphalus samoensis Beeson, *Diapus pusillinus* Chapuis, *Diapus quinquespinatus* Chapuis, *Platypus tetracerus* Beeson, *Xyleborus longicollis* Browne.

Planchonella sandwicensis

Xyleborus lanaiensis Perkins.

***Planchonella* spp.**

Diapus elongatus Schedl, *Diapus nebulosus* Roberts, *Diapus papuanus* Schedl, *Diapus perpygmaeus* Roberts, *Diapus quinquespinatus* Chapuis, *Diapus turgidus* Roberts, *Platypus ambiguus* Roberts, *Platypus aspicius* Roberts & Morimoto, *Platypus glouchideus* Roberts & Morimoto, *Platypus jansoni* Chapuis.

Planchonella andamanica

Xyleborus cognatus Blandford, *Xyleborus similis* Ferrari.

***Planchonella* spp.**

Xyleborus similis Ferrari.

Planera aquatica

Hylocurus floridensis Atkinson.

Platanus acerifolia

Hypothenemus eruditus Westwood.

Platanus occidentalis

Corthylius columbianus Hopkins.

***Platanus* spp.**

Xyleborus dispar (Fabricius).

Plathymenia reticulata

Xyleborus rugosipennis Schedl.

***Platycarya* spp.**

Xylosandrus mutilatus (Blandford).

Platygladus orientalis

Phloeosinus thujae (Perris).

Plectocomia elongata

Xyleborus inaequalis Schedl.

***Electronia* spp.**

Xyleborus cancellatus Eggers.

Pleiocarpa micrantha

Scolytogenes pleiocarpae (Schedl).

Pleiocarpa tubicina

Hypothenemus perhispidus (Eggers), *Platypus intermedius* (Schedl), *Scolytogenes pleiocarpae* (Schedl), *Xyleborus alluaudi* Schaufuss, *Xyleborus principalis* Eichhoff.

***Pleiogynum* spp.**

Ambrosiodinus lecontei Hopkins.

Pleomele aurea

Xyleborus rugatus Blackburn.

Plerandra grayi

Scolytogenes fijianus (Schedl).

***Plochodion* spp.**

Xyleborinus longus (Eggers).

Plumeria rubra

Scolytodes plumeriae Wood, *Scolytodes plumericolens* Wood, *Scolytodes tenuatus* Wood.

***Plumeria* spp.**

Scolytopsis laticollis Wood.

Podocarpus falcata

Lanurgus spathulatus Schedl, *Traglostus brevisetosus* Schedl, *Xyleborinus acmulus* (Wollaston).

Podocarpus imbricata

Platypus bajulus Schedl, *Platypus cavius* Strohmeier, *Platypus vethi* Strohmeier, *Scolytoplatypus nitidus* Eggers.

Podocarpus indicus

Coccotrypes cinnamomi (Eggers), *Coccotrypes medius* (Eggers).

Podocarpus latifolia

Ambrosiodinus eichhoffi (Schreiner), *Hypocryphalus caplandicus* Schedl, *Lanurgus oleaeformis* Schedl.

Podocarpus nageia

Phloeosinus osunensis Murayama.

Podocarpus oleifolia

Corthylius simplex Wood, *Monarthrum nevermanni* (Schedl), *Phlocotribus atavus* Wood.

Podocarpus raspiglosii

Araptus impensus (Wood), *Platypus secus* Wood.

***Podocarpus* spp.**

Acanthotomicus curvidens (Schedl), *Coccotrypes leverii* Browne, *Crossotarsus coxalis* Schedl, *Diapus elongatus* Schedl, *Diapus papuanus* Schedl, *Diapus spinifer* Schedl, *Doliopygus bidentatus* (Strohmeier), *Lanurgus podocarpi* Schedl, *Phloeosinus kinabaluensis* Bright, *Phloeosinus podocarpi* Browne, *Platypus fastuosus* Schedl, *Platypus orientalis* (Strohmeier), *Platypus semiopacus* Strohmeier, *Platypus uniformis* Schedl, *Scolytodes gemmaeus* Wood, *Scolytoplatypus kienensis* Schedl, *Treptoplatypus multiporus* Schedl.

Poinciana elata

Ambrosiodinus lewisi (Blandford), *Euwallacea interjectus* (Blandford), *Platypus indicus* Strohmeier, *Xyleborinus andreweesi* (Blandford).

Poinciana regia

Coccotrypes confusus (Eggers), *Euwallacea xanthopus* (Eichhoff), *Hypothenemus erudiae* (Pauzer), *Hypothenemus liberiensis* (Hopkins), *Hypothenemus melasonus* (Lea), *Micracis burgosi* Wood, *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston).

Poinsettia heterophylla

Hypothenemus brunneus (Hopkins), *Hypothenemus gossypii* (Hopkins).

Polyalthia simiarum

Coccotrypes carpophagus (Hornung).

***Polyalthia* spp.**

Crossotarsus mniszechi Chapuis, *Platypus jansoni* Chapuis, *Platypus selysi* Chapuis.

Polyalthia suaveolens

Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus divaricus* (Schedl), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus expletus* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Hypothenemus macrolobii* (Eggers), *Hypothenemus socialis* (Schedl), *Perionnatus excisus* Strohmeyer, *Perionnatus grandis* Schedl, *Perionnatus longicollis* Chapuis, *Perionnatus substriatus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus picinus* Schedl, *Platypus quadricornis* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Xyleborinus polyalthiae* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xyleborus scobinatus* Hagedorn, *Xyleborus volutus* (Fabricius).

Polygonum tomentosum

Hypothenemus grandis Schedl.

***Polyosma* spp.**

Platypus cupulatus Chapuis, *Platypus geminatus* Chapuis.

Polypodium lycopodioides

Hypothenemus eruditus Westwood, *Hypothenemus perhispidus* (Eggers).

***Polyporus* spp.**

Ambrosiodmus natalensis (Schaufuss).

Polyscias fulva

Chaetastus montanus Schedl, *Chaetastus tuberculatus* (Chapuis), *Platypus inpolitus* Roberts, *Xyleborus alluaudi* Schaufuss.

Polyscias kikuyuensis

Chaetastus montanus Schedl, *Chaetastus tuberculatus* (Chapuis), *Doliopygus bidentatus* (Strohmeyer), *Doliopygus citri* Schedl, *Platypus orientalis* (Strohmeyer), *Tiarophorus gardneri* Schedl.

***Polyscias* spp.**

Chaetastus montanus Schedl, *Doliopygus citri* Schedl, *Doliopygus nairobiensis* (Schedl), *Doliopygus rapax* (Sampson).

Pometia macrocarpa

Platypus protenus (Schedl).

Pometia pinnata

Crossotarsus mniszechi Chapuis, *Crossotarsus wallacei* (Thomson), *Platypus cavius* Strohmeyer, *Platypus jansoni* Chapuis, *Xyleborinus andreweesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus perplexus* Schedl, *Xylosandrus ater* (Eggers), *Xylosandrus manicus* (Blandford).

***Pometia* spp.**

Baioeis pernamulus (Schedl), *Crossotarsus inaequidens* Browne, *Crossotarsus nakazawai* Browne, *Diapus perpygmaeus* Roberts, *Diapus pusillimus* Chapuis, *Diapus*

quincuspinnatus Chapuis, *Platypus abruptifer* Wood, *Platypus granulatus* Browne, *Platypus insulindicus* Schedl, *Platypus jurencus* Schedl, *Platypus lepidus* Chapuis, *Platypus matoae* Browne, *Platypus pseudocupulatus* Schedl, *Xyleborinus andreweesi* (Blandford), *Xyleborinus pomelianus* (Schedl).

Pometia tomentosa

Crossotarsus javanus Beeson, *Platypus signatus* Chapuis.

Pongamia glabra

Phlocoditica curtus (Eggers).

***Popowia* spp.**

Ctonoxylon kituensis Schedl.

Populus acuminata

Trypophloeus populi Hopkins.

Populus alba

Scolytus kirschi Skalitzky, *Trypophloeus granulatus* (Ratzeburg).

Populus angustifolia

Trypophloeus populi Hopkins.

Populus ciliata

Crossotarsus wilmot (Stebbing).

Populus deltoides

Eucallacea validus (Eichhoff), *Hypothenemus seriatus* (Eichhoff), *Trypodendron retusum* (LeConte).

Populus diversifolia

Trypophloeus klimeschi Eggers.

Populus glandulosa

Eucallacea validus (Eichhoff).

Populus grandidentata

Trypodendron retusum (LeConte).

Populus nigra

Scolytus scolytus (Fabricius).

Populus pruinosa

Trypophloeus klimeschi Eggers.

Populus pyramidalis

Trypophloeus binodulus (Ratzeburg), *Trypophloeus granulatus* (Ratzeburg).

Populus sieboldii

Xyleborus volutus (Fabricius).

***Populus* spp.**

Ambrosiodmus apicalis (Blandford), *Ambrosiodmus lewisi* (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Eucallacea fornicatus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Hyllocurus hirtellus* (LeConte), *Micracis swaini* Blackman, *Platypus falcatus* Strohmeyer, *Scolytoplatus tycon* Blandford, *Trypodendron signatum* (Fabricius), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus emarginatus* Eichhoff, *Xyleborus pfeili* (Ratzeburg), *Xylosandrus crassiusculus* (Motschulsky).

Populus suaveolens

Trypophloeus klimeschi Eggers.

Populus tremula

Trypophloeus binodulus (Ratzeburg), *Trypophloeus granulatus* (Ratzeburg), *Trypophloeus kurenzovi* Nunberg, *Trypophloeus palmi* Hansen, *Trypophloeus tremulae* Stark, *Xyleborus pfeili* (Ratzeburg).

Populus tremuloides

Procrisphalus mucronatus (LeConte), *Trypodendron retusum* (LeConte), *Trypophloeus populi* Hopkins, *Trypophloeus thatcheri* (Wood), *Xyleborus obesus* LeConte.

Populus trichocarpa

Gnathotrichus retusus (LeConte), *Trypophloeus populi* Hopkins, *Trypophloeus thatcheri* (Wood).

Potakowskia tacaco

Dendrocranulus declivis Schedl. *Dendrocranulus vicinus* Wood.

Poupartia caffra

Ambrosiodinus sakoae (Schedl).

Pourouma aspera

Gnathotripes terbratus (Wood).

Pouteria anibaefolia

Platypus pouteriae Wood. *Platypus sicarius* Wood. *Xyleborus affinis* Eichhoff. *Xyleborus ferrugineus* (Fabricius).

Pouteria campechana

Chaetophloeus pouteriae Wood.

Pouteria egregia

Platypus applanatus Wood. *Xyleborus affinis* Eichhoff. *Xyleborus rotulus* (Fabricius).

Pouteria guianensis

Platypus chuniculus Wood. *Platypus occipitis* Wood. *Xyleborus rotulus* (Fabricius).

Pouteria malaccensis

Crossotarsus squamulatus Chapuis. *Platypus loricatus* (Saunders). *Platypus pseudocupulatus* Schedl. *Platypus signatus* Chapuis. *Platypus westwoodi* Chapuis.

Pouteria spp.

Dryocoetoides rusticus Wood. *Dryocoetoides severus* (Schedl). *Hypothenemus bolivianus* (Eggers). *Platypus pouteriae* Wood. *Premnobius cavipennis* Eichhoff. *Sampsonius expulsus* Wood. *Taurodenus ebenus* (Wood). *Taurodenus splendidus* (Schaufuss). *Xyleborinus reconditus* (Schedl). *Xyleborus commixtus* Blandford. *Xyleborus parcellus* Wood.

Poiceria exitialis

Platypus exitialis Wood.

Pritchardia pacifica

Coccotrypes dactyliperda (Fabricius).

Pritchardia thurstoni

Coccotrypes carpophagus (Hornig).

Pronia copaifer

Coccotrypes cyperi (Beeson).

Prosopis chilensis

Hypothenemus rugifer (Schedl).

Prosopis glandulosa

Pseudothysanoes turibowi Wood.

Prosopis juliflora

Thysanoes texanus Blackman.

Prosopis nudiflora

Xylosandrus compactus (Eichhoff).

Prosopsis pallida

Xyleborinus saxeseni (Ratzeburg).

Prosopis spp.

Chaetophloeus fasciatus (Blackman). *Hypothenemus erectus* LeConte. *Hypothenemus interstitialis* (Hopkins). *Hypothenemus rotundicollis* (Eichhoff). *Micracisella nanula* (LeConte). *Micracisella opacithorax* (Schedl). *Xylosandrus compactus* (Eichhoff).

Protea spp.

Thamniurgus grandicollis Schedl.

Protium copal

Gnatholeptus semiernis (Nimberg).

Protium macgregorii

Platypus solidus Walker.

Protium nerrosium

Taurodenus ebenus (Wood). *Taurodenus varulus* (Wood). *Xyleborinus celatus* Wood.

Protium spp.

Amphicranus micans Wood. *Camptocerus acneipennis*

(Fabricius). *Camptocerus auricomus* Blandford. *Camptocerus niger* (Fabricius). *Cuemonyx exiguus* (Blandford). *Cuemonyx proticorus* Wood. *Coptoborus usagarius* (Eggers). *Cryphalus nitidipennis* Browne. *Cryptocarenum lepidus* Wood. *Cryptocarenum pilosus* Eggers. *Gnatholeptus panamensis* Blackman. *Gnatholeptus shannoni* (Blackman). *Hypothenemus bolivianus* (Eggers). *Hypothenemus opacus* (Eichhoff). *Microborus limatus* Wood. *Pericryptophalus sobrinus* Wood. *Pityophthorus sparsepilosus* (Schedl). *Pityophthorus suberibratus* Schedl. *Platypus tristis* Roberts. *Premnobius cavipennis* Eichhoff. *Taurodenus bicornutus* (Wood). *Taurodenus flatipes* (Fabricius). *Xyleborus biuncus* Browne. *Xyleborus ferrugineus* (Fabricius). *Xyleborus protii* Browne.

Protium tenuifolium

Camptocerus rectus Wood. *Xylosandrus laticeps* (Wood).

Protorkus spp.

Platypus madagascariensis Chapuis.

Prunus americana

Phloeotribus liminaris (Harris).

Prunus angustifolia

Phloeotribus liminaris (Harris).

Prunus apetala

Xyleborus lactus Niisima.

Prunus armeniaca

Cryphalus malus Niisima. *Scolytoplatypus minimus* Hagedorn. *Scolytus nitidus* Schedl. *Xyleborinus saxeseni* (Ratzeburg).

Prunus avium

Polygraphus grandiclava Thomson.

Prunus cerasa

Polygraphus grandiclava Thomson.

Prunus communis

Phloeotribus pruni Wood.

Prunus davidiana

Ambrosiodinus rubricollis (Eichhoff).

Prunus domestica

Hypothenemus birmanus (Eichhoff). *Phloeotribus porteri* Bruch. *Scolytoplatypus fasciatus* Hagedorn.

Prunus donarium spontanea

Xyleborus atratus Eichhoff.

Prunus grayana

Xylechinus gummeus (Murayama).

Prunus jamasakura

Coccotrypes uaidaijincensis (Murayama). *Crossotarsus simplex* Murayama.

Prunus maximowiczii

Dryocoetes cerasi Eggers.

Prunus nune

Cryphalus malus Niisima.

Prunus nepalensis

Diapus quadrispinatus Chapuis. *Diapus quinque-spinatus* Chapuis. *Diapus spatulifer* Browne. *Scolytoplatypus darjeelingi* Stebbing. *Scolytoplatypus pubescens* Hagedorn. *Scolytoplatypus raja* Blandford.

Prunus padus

Cryphalus scopiger Berger. *Lymantor accris* (Lindemann). *Lymantor coryli* (Perris). *Polygraphus grandiclava* Thomson. *Scolytoplatypus siomio* Blandford. *Xylechinus padus* Wood.

Prunus parviflora

Tomicus multisetosus (Murayama).

Prunus persica

Cryphalus malus Niisima. *Hypothenemus crudiae*

- (Panzer). *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Phloeotribus destructor* Wood, *Phloeotribus discrepans* Blandford, *Phloeotribus liminaris* (Harris), *Phloeotribus pruni* Wood, *Scolytus pomi* Yin & Huang, *Xylosandrus crassiusculus* (Motschulsky).
- Prunus pseudo-cerasus***
Cryphalus nalis Nisima.
- Prunus salasii***
Ambrosiodinus natalensis (Schaufuss), *Eucallacea xanthopus* (Eichhoff), *Hypothenemus grandis* Schedl.
- Prunus salicina***
Cryphalus nalis Nisima.
- Prunus sargentii***
Cryphalus nalis Nisima.
- Prunus serotina***
Hypothenemus eruditus Westwood, *Liparthrum pruni* Wood, *Phloeotribus destructor* Wood, *Phloeotribus liminaris* (Harris), *Phloeotribus perniciosus* Wood, *Phloeotribus pruni* Wood, *Xyleborinus saxeseni* (Ratzeburg).
- Prunus serrulata spontanea***
Eucallacea validus (Eichhoff).
- Prunus sphaerocarpa***
Acorthylus pruni (Wood), *Platypus calamus* Blandford, *Platypus deceptor* Wood, *Platypus deplanatus* Wood.
- Prunus* spp.**
Ambrosiodinus apicalis (Blandford), *Ambrosiodinus lewisi* (Blandford), *Ambrosiodinus rubricollis* (Eichhoff), *Chaetophilocus heterodoxus* (Casey), *Coccotrypes naidaijiniensis* (Murayama), *Glochinocerus gemellus* Blandford, *Hypothenemus eruditus* Westwood, *Monarthrum fasciatum* (Say), *Phloeotribus destructor* Wood, *Phloeotribus pruni* Wood, *Platypus calamus* Blandford, *Platypus cylindrus* (Fabricius), *Platypus quereivorus* Murayama, *Platypus severini* Blandford, *Polygraphus parvulus* Murayama, *Polygraphus ssiori* Nisima, *Scolytplatypus mikado* Blandford, *Scolytus amygdali* Guérin-Meneville, *Scolytus aratus* Blandford, *Scolytus ensifer* Eichhoff, *Scolytus gretschkii* Sokanovskii, *Scolytus japonicus* Chapuis, *Scolytus kirschi* Skalitzy, *Scolytus mali* (Beckstein), *Scolytus pygmaeus* (Fabricius), *Scolytus rugulosus* (Müller), *Scolytus scheygrewi* Semenov, *Trypodendron domesticum* (Linnaeus), *Xyleborinus attenuatus* (Blandford), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus praeceus* Blandford, *Xyleborus seriatus* Blandford, *Xylosandrus germanus* (Blandford), *Xylosandrus pseudosolidus* (Schedl).
- Prunus ssiori***
Platypus calamus Blandford, *Polygraphus shariensis* Nisima.
- Prunus tomentosa***
Scolytus pomi Yin & Huang.
- Prunus triloba***
Cryphalus pruni Eggers.
- Prunus virginiana***
Phloeotribus pruni Wood.
- Prunus vulgaris***
Phloeotribus pruni Wood.
- Prunus yedoensis***
Xyleborus atratus Eichhoff.
- Pseudoeugenia* spp.**
Platypus hamatulus Schedl.
- Pseudomorus brunoniana***
Ficicis varians Lea.
- Pseudoolmedia laevigata***
Dryocoetoides severus Wood, *Taurodenus splendidus* (Schaufuss), *Taurodenus varians* (Fabricius), *Xyleborinus reconditus* (Schedl).
- Pseudoolmedia* spp.**
Phloeotribus minor Wood.
- Pseudospondias microcarpa***
Chaetastus tuberculatus (Chapuis), *Platypus hintzi* Schaufuss, *Scolytplatypus africanus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).
- Pseudospondias* spp.**
Trachyostus aterrimus (Schaufuss).
- Pseudotsuga douglasii***
Cryphalus asperatus (Gyllenhal).
- Pseudotsuga flahuati***
Hylastes asperatus Wood, *Hylastes gracilis* LeConte.
- Pseudotsuga macrocarpa***
Carphoborus blaisdelli Swaine, *Dendroctonus pseudotsugae* Hopkins, *Pseudohylesinus nebulosus serratus* Bruck, *Scolytus oregoni* Blackman.
- Pseudotsuga menziesii***
Carphoborus pseudotsugae Wood, *Carphoborus vandykei* Bruck, *Cryphalus pubescens* Hopkins, *Cryphalus ruficollis* Hopkins, *Dendroctonus pseudotsugae* Hopkins, *Dryocoetes autographus* (Ratzeburg), *Dryocoetes cristatus* Inouye & Nobuchi, *Gnathotrichus retusus* (LeConte), *Gnathotrichus sulcatus* (LeConte), *Hylastes ater* (Paykull), *Hylastes nigrinus* (Mannerheim), *Hylastes ruber* Swaine, *Hylurgops reticulatus* Wood, *Hylurgops rugipennis* (Mannerheim), *Hylurgops subcostulatus* (Mannerheim), *Phloeosinus autumnatus* Swaine, *Phloeotribus lecontei* Schedl, *Pityokteines elegans* Swaine, *Pityokteines lasiocarpi* (Swaine), *Pityophthorus arcuatus* Bright, *Pityophthorus intentus* Bright, *Pityophthorus malleatus* Bright, *Pityophthorus opaculus* LeConte, *Pityophthorus orarius* Bright, *Pityophthorus pseudotsugae* Swaine, *Pityophthorus solers* Blackman, *Pityophthorus speculum* Bright, *Platypus apicalis* (White), *Platypus wilsoni* Swaine, *Pseudohylesinus nebulosus nebulosus* (LeConte), *Pseudohylesinus sericeus* (Mannerheim), *Pseudothysanoes conferae* (Wood), *Scierus annexens* LeConte, *Scolytus hermosus* Wood, *Scolytus monticola* (Swaine), *Scolytus oregoni* Blackman, *Scolytus reflexus* Blackman, *Scolytus inispinosus* LeConte, *Scolytus virgatus* Bright, *Xyleborus intrusus* Blandford.
- Pseudotsuga* spp.**
Trypodendron lineatum (Olivier).
- Psidium guajava***
Diaprus formosanus Nisima & Murayama, *Eucallacea fornicatus* (Eichhoff), *Eucallacea xanthopus* (Eichhoff), *Phloeoborus punctatorugosus* Chapuis, *Pityophthorus robai* (Blackman), *Platypus flectus* Nisima & Murayama, *Platypus nijimai* Murayama, *Scolytogenes samoanus* (Browne), *Taurodenus perebae* (Ferrari), *Xyleborus affinis* Eichhoff, *Xyleborus subtuberculatus* Eggers.
- Psittacanthus schiedeanus***
Chaetophilocus struthanthi Wood, *Cnesinus uogueriae* Atkinson, *Corthylus senticosus* Wood.
- Psittacanthus* spp.**
Corthylus senticosus Wood, *Hypothenemus indigenus* Wood, *Pseudothysanoes plumalis* Wood, *Pseudothysanoes tenellus* Wood.
- Psychotia* spp.**
Xyleborinus quadrispinosus (Eichhoff), *Xyleborus alluaudi* Schaufuss, *Xylosandrus hirsutipennis* (Schedl).

Psychotia viridiflora*Platypus signatus* Chapuis.***Ptelea trifoliata****Phlocotribus scabricollis* (Hopkins).***Pteleopsis hylo dendron****Chaetastus tuberculatus* (Chapuis), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Dryocetoides cristatus* (Fabricius), *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl.***Pteridium aquilinum****Coccotrypes cardamomi* Schaufuss, *Coccotrypes pterydophytæ* (Schedl).***Pternandra coerulea****Anasa versicolor* (Sampson).***Pterocarpus angoleusis****Doliopygus conradti* (Strohmeyer), *Doliopygus crinitus* (Chapuis), *Doliopygus subserratus* Browne.***Pterocarpus dalbergioides****Crossotarsus andamanus* Beeson, *Platypus cupulatus* Chapuis, *Platypus solidus* Walker, *Platypus verchunatus* (Beeson), *Xyleborinus exiguus* (Walker), *Xyleborus bidentatus* (Motschulsky), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus pfeili* (Ratzeburg), *Xyleborus similis* Ferrari.***Pterocarpus indicus****Crossotarsus externedentatus* (Fairmaire), *Eucallacea limatus* (Schedl), *Hypocryphalus perminutus* (Schedl), *Platypus insularis* Strohmeyer, *Xyleborus granulipes* Schedl, *Xyleborus similis* Ferrari.***Pterocarpus marsupium****Ambrosiodinus diligensis* (Eggers), *Crossotarsus minax* (Walker), *Erioporus concentricus* (Eggers), *Eucallacea andamanensis* (Blandford), *Eucallacea interjectus* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Pterocarpus mildbraedii****Chaetastus tuberculatus* (Chapuis), *Doliopygus pseudoserratus* Roberts, *Trachyostus aterrimus* (Schaufuss).***Pterocarpus osun****Mesoplatypus tenuis* Schedl.***Pterocarpus soyauxii****Ambrosiodinus oratus* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus pertinax* (Schedl), *Doliopygus coelocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus maynei* Schedl, *Doliopygus opulentus* Schedl, *Doliopygus semipilosus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Hypothenemus eruditus* Westwood, *Prennobiobius longus* Eggers, *Prennobiobius robustulus* (Schedl), *Prennobiobius scapinosus* Eggers, *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus candatus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius).***Pterocarpus* spp.***Ambrosiodinus natalensis* (Schaufuss), *Erioporus armatus* (Browne), *Erioporus tuberculatus* (Browne), *Leptoxyleborus concisus* (Blandford), *Platypus triplurus* Browne.***Pterocarpus stenoptera****Sphaerotrypes juglausi* Tsai & Yin.***Pterocarpus tinctorius****Cylindropalpus laudatus* (Schedl), *Doliopygus per-**brevis* Schedl, *Hypothenemus pubipennis* (Eggers), *Hypothenemus solitarius* (Schedl), *Platypus parallelus* (Fabricius), *Xyleborus affinis* Eichhoff, *Xyleborus ficus* Eggers.***Pterocarya rhoifolia****Crossotarsus niponicus* Blandford, *Scolytogenes fujisanus* (Nobuchi).***Pterocymbium beccarii****Crossotarsus subpellucidus* Lea, *Eucallacea barbatus* (Hagedorn), *Eucallacea wallacei* (Blandford), *Platypus pallidus* Chapuis, *Platypus parallelus* (Fabricius), *Platypus solidus* Walker, *Platypus subgranosus* Schedl, *Platypus ustus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus similis* Ferrari.***Pterocymbium cinctiorum****Platypus quadrifissilis* Schedl.***Pterocymbium* spp.***Crossotarsus wallacei* (Thomson), *Diapus pusillinus* Chapuis, *Hyledius cribratus* (Blandford), *Platypus jansoni* Chapuis.***Pteromalus bimaculatus****Xyleborinus saxeseni* (Ratzeburg).***Pterospermum acerifolium****Platypus solidus* Walker, *Xyleborus perforans* (Wollaston), *Xylosandrus discolor* (Blandford).***Pterospermum heyneanum****Coccotrypes longior* (Eggers).***Pterospermum* spp.***Xyleborinus andrewesi* (Blandford).***Pterygopodium oxyphyllum****Doliopygus donisi* Schedl.***Pterygota bequaertii****Chaetastus tuberculatus* (Chapuis), *Doliopygus jurioni* Schedl, *Doliopygus lateralis* Schedl, *Doliopygus lateralis* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditicus* (Schedl), *Doliopygus unicornis* Schedl, *Platypus hintzi* Schaufuss, *Platypus intermedius* (Schedl), *Strombophorus vagans* Schedl, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).***Pterygota macrocarpa****Cylindropalpus auricomans* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus medius* Schedl, *Doliopygus mimicus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Platypus hintzi* Schaufuss, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).***Pterygota* spp.***Platypus oblique truncatus* Schedl, *Prennobiobius caripennis* spp.***Ptychosperma* spp.***Coccotrypes dactyliperda* (Fabricius).***Pueraria thumbergiana****Ambrosiodinus levisi* (Blandford).***Punica granatum****Eucallacea xanthopus* (Eichhoff), *Platypus solutus* Schedl, *Xyleborus piccus* (Motschulsky).***Punica* spp.***Xyleborus dispar* (Fabricius).***Purshia mexicana****Chaetophloeus heterodoxus* (Casey).***Pycanthus angoleusis****Chaetastus persimilis* (Schedl), *Chaetastus tuberculatus* (Chapuis), *Doliopygus bilobatus* (Schedl), *Doliopygus*

conradti (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus interjectus* Schedl, *Doliopygus lefevrei* Schedl, *Eucallacea gramosus* (Schedl), *Mesoplatypus venustus* Schedl, *Periommatus excisus* Strohmeyer, *Periommatus grandis* Schedl, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus orientalis* (Strohmeyer), *Platypus parallelus* (Fabricius), *Platypus picinus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus nitidulus* Eggers, *Xyleborus picus* (Motschulsky), *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Pycanthus kombo

Chaetastus persimilis (Schedl), *Chaetastus tuberculatus* (Chapuis), *Doliopygus conradti* (Strohmeyer), *Doliopygus serratus* (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xyleborus volutus* (Fabricius).

Pycanthus marchalianus

Xyleborus ferrugineus (Fabricius).

***Pycanthus* spp.**

Ambrosiodinus eichhoffi (Schreiner).

Pycnostachys coerulea

Eucallacea xanthopus (Eichhoff), *Rhopalopselion congonum* (Schedl).

Pygeum africanum

Chaetastus diversifrons Browne, *Chaetastus montanus* Schedl, *Cylindropalpus laudatus* (Schedl), *Hylesinopsis fasciatus* (Hagedorn), *Sphaerotrypes pygeumii* Schedl, *Triozastus marshalli* (Sampson).

Pygeum parviflorum

Diapus quinquespinatus Chapuis.

Pyrus betulaeolia

Sphaerotrypes pyri Tsai & Yin.

Pyrus communis

Cnesinus strigicollis LeConte.

Pyrus lanata

Dryocoetes himalayensis Strohmeyer, *Scolytoplatus sionio* Blandford.

Pyrus malus

Chaetophloeus heterodoxus (Casey), *Cryphalus malus* Nüßima, *Hypothenemus africanus* (Hopkins), *Hypothenemus crudus* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus malus* (Schedl), *Lymantrix coryli* (Perris), *Scolytus nitidus* Schedl.

***Pyrus* spp.**

Hypothenemus eruditus Westwood, *Scolytoplatus daimio* Blandford, *Scolytus japonicus* Chapuis, *Scolytus mali* (Beckstein), *Scolytus rugulosus* (Müller), *Xylosandrus germanus* (Blandford), *Xyleborus dispar* (Fabricius).

Pyrus ussuriensis

Scolytus jacobsoni (Spessivtsev).

Qualea ingens

Dryocoetoides flavus (Fabricius), *Taurodemus perebeae* (Ferrari), *Xyleborinus reconditus* (Schedl), *Xyleborus spinulosus* Blandford.

Qualea wittrockii

Hypothenemus plumieriae (Nordlinger), *Sampsonius dampfi* Schedl.

Quercus acuta

Ambrosiodinus rubricollis (Eichhoff), *Coptodryas kirishimanus* (Murayama), *Platypus calamus* Blandford.

Platypus lewisi Blandford, *Platypus quercivorus* Murayama, *Xyleborus atratus* Eichhoff, *Xyleborus miyazakensis* Murayama, *Xyleborus nameranus* Murayama, *Xyleborus pelliculosus* Eichhoff, *Xyleborus tsukubanus* Murayama.

Quercus acutissima

Platypus koryoensis (Murayama), *Platypus lewisi* Blandford, *Platypus quercivorus* Murayama, *Scolytogenes imitans* Eichhoff, *Sphaerotrypes pila* Blandford.

Quercus afares

Taphrorychus villifrons (Dufour).

Quercus agrifolia

Gnathotrichus pilosus (LeConte), *Monarthrum dentigerum* (LeConte), *Monarthrum scutellare* (LeConte), *Pseudopityophthorus agrifoliae* Blackman, *Pseudopityophthorus pubipennis* (LeConte).

Quercus alba

Corthylus columbianus Hopkins, *Micracisella opacicollis* (LeConte), *Monarthrum fasciatum* (Say), *Pseudopityophthorus pubescens* Blackman, *Xyleborus ferrugineus* (Fabricius).

Quercus aliena

Platypus koryoensis (Murayama), *Platypus lewisi* Blandford.

Quercus borealis

Pseudopityophthorus pubescens Blackman, *Xyleborus ferrugineus* (Fabricius).

Quercus cerris

Erioporicus fagi (Fabricius), *Taphrorychus hirtellus* Eichhoff.

Quercus coccinea

Xyleborus ferrugineus (Fabricius).

Quercus costaricensis

Monarthrum pseudoscutellare (Schedl).

Quercus crispula

Platypus calamus Blandford.

Quercus dentata

Taphrorychus hewetti (Stebbing).

Quercus digitata

Monarthrum fasciatum (Say).

Quercus dilatata

Diapus aculeatus Blandford, *Diapus molossus* Chapuis, *Diapus spatulifer* Browne, *Platypus sexualis* Beeson, *Sphaerotrypes querci* Stebbing.

Quercus emoryi

Pseudopityophthorus opacicollis Blackman, *Pseudothythanoes sedulus* Blackman.

Quercus fagifolia

Diapus quinquespinatus Chapuis.

Quercus falcata

Micracisella opacicollis (LeConte), *Xyleborus ferrugineus* (Fabricius).

Quercus gambellii

Pseudothythanoes sedulus Blackman.

Quercus garrayana

Pseudopityophthorus pubipennis (LeConte).

Quercus gilliana

Scolytus querci Yin & Huang.

Quercus gilva

Amasa calamoides (Murayama), *Diapus spatulifer* Browne, *Platypus calamus* Blandford, *Platypus kiushuensis* Murayama, *Platypus lewisi* Blandford, *Platypus quercivorus* Murayama, *Platypus severini* Blandford.

Quercus glauca

Coptodryas kirishimana (Murayama), *Crossotarsus flavomaculatus* Strohmeyer, *Eucallacea velatus* (Samp-

son), *Platypus calamus* Blandford, *Platypus quercivorus* Murayama.

Quercus griseus

Pseudopityophthorus ratapaii Blackman.

Quercus grosseserrata

Eucallacea validus (Eichhoff).

Quercus hondai

Platypus calamus Blandford.

Quercus hondurensis

Monarthrum quercuum Wood, *Pseudopityophthorus granulifer* Wood, *Pseudopityophthorus pruinosis* (Eichhoff).

Quercus humboldtii

Cnesinus deperditus Wood, *Cnesinus fulgidus* Wood, *Phloeotribus remorsus* Wood, *Pseudopityophthorus colombianus* Wood.

Quercus hypoleucoides

Corthylihus petilus Wood, *Monarthrum huachucae* Wood, *Pseudopityophthorus granulatus* Blackman, *Pseudopityophthorus pruinosis* (Eichhoff), *Pseudopityophthorus yacapaii* Blackman.

Quercus ilex

Taphrorychus minor Eggers.

Quercus incana

Carchesiopygus lobacanthus Schedl, *Crossotarsus fairmairei* Chapuis, *Crossotarsus uilmoti* (Stebbing), *Diapus aculeatus* Blandford, *Diapus quadrispinatus* Chapuis, *Diapus quinquespinatus* Chapuis, *Platypus abruptus* Sampson, *Polygraphus querci* Wood, *Scolytroplatypus pubescens* Hagedorn, *Sphaerotrypes querci* Stebbing.

Quercus jordanae

Xyleborus rufus Schedl.

Quercus kelloggii

Monarthrum scutellare (LeConte), *Pseudopityophthorus agrifoliae* Blackman, *Pseudopityophthorus pubipennis* (LeConte).

Quercus lamellosa

Ambrosiodmus apicalis (Blandford), *Arixyleborus moestus* (Eggers), *Diapus minor* Schedl, *Diapus molossus* Chapuis, *Diapus quadrispinatus* Chapuis, *Diapus quinquespinatus* Chapuis, *Diapus spatulifer* Browne, *Indoceryphalus intermedius* (Sampson), *Platypus curtatus* Sampson, *Platypus falcatus* Strohmeier, *Platypus lewisi* Blandford, *Platypus quadriporus* (Beeson), *Pseudoxylechinus indicus* Wood, *Scolyto-platypus darjeelingi* Stebbing, *Xyleborus improbus* Sampson, *Xyleborus inarmatus* Eggers, *Xylechinus darjeelingensis* Schedl.

Quercus lanuginosa

Taphrorychus heuetti (Stebbing).

Quercus lineata

Platypus quadriporus (Beeson).

Quercus marylandica

Pseudopityophthorus pruinosis (Eichhoff), *Trischidias atomus* (Hopkins).

Quercus mirbeckii

Taphrorychus villifrons (Dufour).

Quercus mongolica

Platypus quercivorus Murayama, *Xyleborus atratus* Eichhoff, *Xyleborus lactus* Niisima, *Xyleborus punctulatus* Kurenzov, *Xyleborus starki* Numberg.

Quercus mongolica grosseserrata

Crossotarsus niponicus Blandford, *Platypus calamus* Blandford, *Platypus lewisi* Blandford, *Platypus modestus* Blandford, *Platypus quercivorus* Murayama, *Xyleborus aquilus* Blandford.

Quercus myrsinaefolia

Acanthotomicus spinosus Blandford, *Ambrosiodmus rubricollis* (Eichhoff), *Crossotarsus niponicus* Blandford, *Platypus calamus* Blandford, *Platypus quercivorus* Murayama, *Sphaerotrypes pila* Blandford, *Xyleborus japonicus* Nobuchi, *Xyleborus naneranus* Murayama, *Xylosandrus compactus* (Eichhoff).

Quercus nigra

Pseudopityophthorus asperulus (LeConte), *Pseudopityophthorus pruinosis* (Eichhoff).

Quercus pedunculata

Taphrorychus villifrons (Dufour).

Quercus phellos

Xyleborus ferrugineus (Fabricius).

Quercus phillyraeoides

Xyleborus misatoensis Nobuchi, *Xyleborus wakayamensis* Nobuchi.

Quercus pubescens

Taphrorychus villifrons (Dufour).

Quercus robur

Platypus mutatus Chapuis, *Taphrorychus villifrons* (Dufour).

Quercus rubra

Monarthrum fasciatum (Say), *Monarthrum mali* (Fitch), *Pseudopityophthorus asperulus* (LeConte), *Pseudopityophthorus pubescens* Blackman.

Quercus salicina

Crossotarsus niponicus Blandford, *Platypus calamus* Blandford, *Platypus quercivorus* Murayama.

Quercus sapotaeifolia

Micracis lignicolus Wood, *Pseudopityophthorus granulifer* Wood.

Quercus semicarpifolia

Crossotarsus fairmairei Chapuis, *Diapus aculeatus* Blandford, *Diapus molossus* Chapuis, *Genyocerus furticus* (Sampson), *Scolytus querci* Yin & Huang, *Sphaerotrypes querci* Stebbing, *Taphrorychus heuetti* (Stebbing).

Quercus serrata

Diapus quinquespinatus Chapuis, *Platypus calamus* Blandford, *Platypus koryoensis* (Murayama), *Platypus lewisi* Blandford, *Platypus quercivorus* Murayama, *Xyleborus atratus* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Quercus sessilifolia

Platypus calamus Blandford, *Platypus quercivorus* Murayama.

Quercus sopataefolia

Pseudopityophthorus pruinosis (Eichhoff).

Quercus spicata

Coccotrypes papuanus (Eggers).

***Quercus* spp.**

Anasa schlichii (Stebbing), *Ambrosiodmus apicalis* (Blandford), *Ambrosiodmus devexus* (Schedl), *Ambrosiodmus ferus* (Wood), *Ambrosiodmus funebris* (Schedl), *Ambrosiodmus lewisi* (Blandford), *Ambrosiodmus rubricollis* (Eichhoff), *Amphicranus rameus* Wood, *Amphicranus splendens* Wood, *Chramesus dentatus* Schaeffer, *Cnesinus elegantis* Wood, *Cnesinus gibbulus* Wood, *Cnesinus porcatus* Blandford, *Cnesinus strigicollis* LeConte, *Corthylicyclon aztecum* (Bright), *Corthylihus minutus* Bright, *Corthylihus petilus* Wood, *Crossotarsus fractus* Sampson, *Crossotarsus horni* Schedl, *Crossotarsus oligodontus* Roberts, *Crossotarsus simplex* Murayama, *Crossotarsus terminatus* Chapuis, *Cryphalus felis* Wood, *Cyclorhipidion circumcissum*

(Sampson), *Cyclorhpidion subobtusum* (Schedl), *Diaprus pendleburgi* Schedl, *Dryocoetius coffeae* (Eggers), *Eucallacea xanthopus* (Eichhoff), *Glochinoscerus gemellus* Blandford, *Gnathotrichus dentatus* Wood, *Gnathotrichus nimifrons* Wood, *Gnathotrichus obscurus* Wood, *Gnathotrichus pilosus* (LeConte), *Gnathotrichus primus* (Bright), *Gnathotrupes bituberculatus* (Blandford), *Hadrodemius amorphus* (Eggers), *Hylesinus crenatus* (Fabricius), *Hylesinus varius* (Fabricius), *Hylocurus femineus* Wood, *Hylocurus hirtellus* (LeConte), *Hylocurus incompitus* Wood, *Hylocurus longipennis* Wood, *Hylocurus parkinsoniae* Blackman, *Hypothenemus birmanus* (Eichhoff), *Hypothenemus columbi* Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus dissimilis* (Zimmermann), *Hypothenemus eruditus* Westwood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus rotundicollis* (Eichhoff), *Hypothenemus scriatus* (Eichhoff), *Leptoxyleborus concisus* (Blandford), *Lymantor coryli* (Perris), *Micracis amplius* Wood, *Micracis evanescens* Wood, *Micracis incertus* Wood, *Micracis lignator* Blackman, *Micracisella adnata* Wood, *Micracisella knulli* (Blackman), *Micracisella mimetica* Wood, *Micracisella nanula* (LeConte), *Micracisella scitula* Wood, *Microcorthyus vicinus* Wood, *Monarthrum aberrans* Schedl, *Monarthrum bicavum* Wood, *Monarthrum cordatum* (Blandford), *Monarthrum cordicatum* Wood, *Monarthrum dentigerum* (LeConte), *Monarthrum desum* (Wood), *Monarthrum fasciatum* (Say), *Monarthrum gnarum* (Schedl), *Monarthrum laterale* (Eichhoff), *Monarthrum luctuosum* (Blandford), *Monarthrum morsum* Wood, *Monarthrum netemanni* (Schedl), *Monarthrum pseudoscutellare* (Schedl), *Monarthrum quercicolens* Wood, *Monarthrum quercivorus* Schedl, *Monarthrum quercus* (Wood), *Monarthrum quercuum* Wood, *Monarthrum scrobiceps* (Eichhoff), *Monarthrum sulcatum* (Blandford), *Monarthrum validum* (Ferrari), *Monarthrum vittatum* (Blandford), *Phloeosinus pulchellus* Blandford, *Phloeotribus quercinus* Wood, *Pityophthorus conspectus* Wood, *Pityophthorus guatemalensis* Blandford, *Pityophthorus lautus* Eichhoff, *Pityophthorus medialis* Wood, *Pityophthorus melanurus* Wood, *Pityophthorus parilis* Wood, *Pityophthorus quercinus* Wood, *Pityophthorus scitulus* Wood, *Platypus annexus* Wood, *Platypus cugestus* Wood, *Platypus formosanus* Niisima & Murayama, *Platypus connexus* Wood, *Platypus convexicauda* Schedl, *Platypus cylindrus* (Fabricius), *Platypus jansoni* Chapuis, *Platypus kiusuensis* Murayama, *Platypus nardicus* Blandford, *Platypus otiosus* Schedl, *Platypus platypoides* (Browne), *Platypus quadridentatus* (Olivier), *Platypus quercus* Wood, *Platypus querci* Browne, *Platypus quercicola* Schedl, *Platypus quercivorus* Murayama, *Pseudopityophthorus agrifoliae* Blackman, *Pseudopityophthorus asperulus* (LeConte), *Pseudopityophthorus comosus* Bright, *Pseudopityophthorus declivis* Wood, *Pseudopityophthorus denticulus* Wood, *Pseudopityophthorus festinus* Wood, *Pseudopityophthorus granulatus* Blackman, *Pseudopityophthorus hondurensis* Wood, *Pseudopityophthorus limbatus* Eggers, *Pseudopityophthorus minutissimus* (Zimmermann), *Pseudopityophthorus opacicollis* Blackman, *Pseudopityophthorus peregrinus* Wood & Yin, *Pseudopityophthorus pruinosis* (Eichhoff), *Pseudopityophthorus singularis* Wood, *Pseudopityophthorus squamosus* Bright, *Pseudopityophthorus tenuis* Wood, *Pseudopityophthorus virilis* Wood, *Pseudopityophthorus yavapaii* Blackman, *Pseudo-*

thysanoes brunneus Wood, *Pseudothysanoes costatus* Wood, *Pseudothysanoes lecontei* Blackman, *Pseudothysanoes quercicolens* Wood, *Pseudothysanoes quercinus* (Wood), *Pseudothysanoes quercus* Wood, *Pseudothysanoes securus* Wood, *Pseudothysanoes scitulus* Blackman, *Schedlia vulpinus* (Schedl), *Scolytodes anabilis* Wood, *Scolytodes radiatus* Wood, *Scolytodes daimio* Blandford, *Scolytodes mikado* Blandford, *Scolytodes sionio* Blandford, *Scolytodes tycon* Blandford, *Scolytus intricatus* (Ratzeburg), *Scolytus laevis* Chapuis, *Scolytus pygmaeus* (Fabricius), *Scolytus scolytus* (Fabricius), *Scolytus sulcifrons* Rey, *Sphaerotrupes yunnanensis* Tsai & Yin, *Tessrocercus alternans* Schedl, *Tessrocercus schmutzenhoferi* Schedl, *Thysanoes adonis* Wood, *Thysanoes berchemiae* Blackman, *Thysanoes fimbricornis* LeConte, *Thysanoes lobdelli* Blackman, *Thysanoes pallens* Wood, *Thysanoes subsulcatus* Wood, *Thysanoes xylophagus* Blackman, *Tiarophorus scrutator* (Pandelle), *Trypodendron domesticum* (Linnaeus), *Webbia piscecauda* Browne, *Xyleborinus aenulus* (Wollaston), *Xyleborinus attenuatus* (Blandford), *Xyleborinus exiguus* (Walker), *Xyleborinus gracilicornis* (Schedl), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus adusticollis* (Motschulsky), *Xyleborus affinis* Eichhoff, *Xyleborus armipennis* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus eurgraphus* (Ratzeburg), *Xyleborus excusus* Blandford, *Xyleborus kumamotoensis* Murayama, *Xyleborus monographus* (Fabricius), *Xyleborus obesus* LeConte, *Xyleborus olmoi* Browne, *Xyleborus okinosuensis* Murayama, *Xyleborus onoharacensis* Murayama, *Xyleborus perforans* (Wollaston), *Xyleborus pfeili* (Ratzeburg), *Xyleborus quercicola* Eggers, *Xyleborus scirporensis* Murayama, *Xyleborus scriatus* Blandford, *Xyleborus xylographus* (Say), *Xylosandrus brevis* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus germanus* (Blandford), *Xylosandrus manicus* (Blandford), *Xyloterinus politus* (Say).

Quercus stenophylla

Ambrosiodinus lewisi (Blandford), *Platypus calamus* Blandford, *Xyleborinus attenuatus* (Blandford), *Xyleborinus saxeseni* (Ratzeburg), *Xyleborus kojimai* Murayama, *Xyleborus pelliculosus* Eichhoff, *Xyleborus volulus* (Fabricius).

Quercus texana

Hypothenemus eruditus Westwood.

Quercus tomentocaulis

Platypus longius Wood.

Quercus tomentosicupula

Xyleborus atratus Eichhoff.

Quercus velutina

Hylocurus rudis (LeConte), *Pseudopityophthorus asperulus* (LeConte).

Quercus virginiana

Araptus nannus Wood.

Quercus williamsii

Micracis lignicolus Wood.

Quercus wislizenii

Pseudopityophthorus agrifoliae Blackman.

Quercus wrayi

Cyclorhpidion punctatopilosum (Schedl), *Platypus quercinus* Schedl.

***Quintinia* spp.**

Crossotarsus caliginosus Roberts, *Margadillius magnus* Browne.

Quisqualis indica

Hypothenemus californicus Hopkins, *Hypothenemus crudiae* (Panzer), *Hypothenemus cruditus* Westwood.

***Radikofera* spp.**

Doliopygus bitalei Schedl, *Doliopygus omissus* Schedl, *Eucallacea xanthopus* (Eichhoff), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Xyleborus althaudi* Schaufuss, *Xyleborus amanicus* Hagedorn, *Xyleborus principalis* Eichhoff.

Randia acuminata

Cylindropalpus granulatus (Schedl).

Randia anisophylla

Anasa versicolor (Sampson), *Ambrosiodinus rubricollis* (Eichhoff), *Leptoxyleborus concisus* (Blandford).

Randia congolana

Doliopygus serratus (Strohmeyer), *Hypothenemus perhispidus* (Eggers), *Micryphalus nitidus* (Schedl), *Xyleborus althaudi* Schaufuss, *Xylosandrus crassiusculus* (Motschulsky).

***Randia* spp.**

Pseudothysanocs truncatus Wood, *Pseudothysanocs resculus* Wood, *Xyleborinus andrewesi* (Blandford).

Randia uliginosa

Eucallacea andamanensis (Blandford).

Rapanea guyanensis

Corthylus mexicanus Schedl.

Rapanea melanophloeos

Xyleborinus acmulus (Wollaston).

Rapanea rhododendroides

Chaetastus montanus Schedl.

Raucolfia caffra

Eucallacea xanthopus (Eichhoff), *Premnobius minor* Eggers.

Raucolfia inebrians

Rhopalopschion immaturus (Schedl).

Raucolfia vomitoria

Triozastus caliginosus Roberts.

Rarensara perrillei

Platypus madagascariensis Chapuis, *Mitosoma odonticeps* Schedl.

Rarensara polyneura

Mitosoma odonticeps Schedl.

***Rarensara* spp.**

Eucallacea xanthopus (Eichhoff), *Hypothenemus abruptus* (Schedl), *Mitosoma obconiceps* Schedl, *Mitosoma truncatipennis* Schedl, *Xyleborinus mitosomipennis* (Schedl), *Xyleborus madagascariensis* Schaufuss, *Xyleborus sakalava* Schedl, *Xylosandrus mancus* (Blandford).

Ravenala madagascariensis

Mitosoma odonticeps Schedl.

***Rengas* spp.**

Platypus perbrevis Browne.

Retama bovei

Phlocotribus pycrinhoffi (Eggers).

Retama sphaerocarpa

Liparthrum genistae (Aube), *Phlocotribus rhodactylus* (Marshall).

***Retama* spp.**

Phlocotribus cristatus (Fauvel), *Phlocotribus maroccanus* (Guillebeau).

Reynoldsia tahitensis

Xyleborus munifordii Beeson.

Reynosia septentrionalis

Hypothenemus hirsutus (Wood).

Rhacoma crossopetalum

Cryptocarenum scriatus Eggers.

Rhamnus betulaeifolia

Chramesus setosus Wood.

Rhamnus capreaefolia

Chramesus chapuisi LeConte.

Rhamnus cathartica

Lymanator coryli (Perris).

Rhamnus frangula

Lymanator coryli (Perris).

Rhamnus lanceolata

Hypothenemus dissimilis (Zimmermann), *Hypothenemus rotundicollis* (Eichhoff).

***Rhamnus* spp.**

Hylocurus hirtellus (LeConte), *Hypothenemus scriatus* (Eichhoff), *Hypothenemus sparsus* Hopkins.

Rheedia edulis

Camptocerus auricomus Blandford, *Micracis carinulus* Wood, *Microborus boops* Blandford, *Phrixosoma minor* Wood, *Phrixosoma rude* Blandford, *Pityophthorus galeritus* Wood, *Pityophthorus strictus* Wood.

Rheedia madruno

Phrixosoma crebra Wood, *Phrixosoma frustrata* Wood, *Phrixosoma obesa* Blackman, *Phrixosoma viriosa* Wood, *Taurodemus bicornutus* (Wood), *Xyleborus schildi* Schedl.

***Rheedia* spp.**

Hypothenemus javanus (Eggers).

Rhizophora apiculata

Crossotarsus externedentatus (Fairmaire), *Platypus pseudocupulatus* Schedl, *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston).

Rhizophora mangle

Coccotrypes rhizophorae (Hopkins), *Hypothenemus birmanus* (Eichhoff), *Hypothenemus crudiae* (Panzer), *Hypothenemus cruditus* Westwood, *Hypothenemus javanus* (Eggers), *Micracisella nanula* (LeConte), *Thysanocs pallens* Wood, *Trischidias atomus* (Hopkins), *Trischidias minutissima* Wood, *Xyleborus productus* Hagedorn.

Rhizophora mucronata

Coccotrypes litoralis (Beeson), *Coccotrypes rhizophorae* (Hopkins), *Ernaporus incrimis* (Schedl), *Platypus forciflua* Chapuis, *Platypus insularis* Strohmeyer, *Xyleborus bidentatus* (Motschulsky), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus sinilis* Ferrari.

***Rhizophora* spp.**

Coccotrypes fallax (Eggers), *Cryphalus negrosensis* Browne, *Premnobius cavipennis* Eichhoff, *Xylosandrus compactus* (Eichhoff).

Rhodannia trinervia

Anasa versicolor (Sampson), *Arixyleborus scabripennis* (Blandford), *Eccoptypterus limbus* Sampson, *Hadrodemus comans* (Sampson), *Xyleborus approximatus* Schedl, *Xyleborus minutus* Blandford.

Rhodobaena bakeriana

Mitosoma obconiceps Schedl, *Mitosoma obliquatum* Schedl, *Mitosoma paulianum* Schedl.

***Rhododendron* spp.**

Corthylus punctatissimus (Zimmermann), *Hypothenemus interstitialis* (Hopkins).

Rhoicissus erythrodes

Eucallacea xanthopus (Eichhoff), *Xyleborus principalis* Eichhoff.

Rhus ambigua*Xyleborus seriatus* Blandford.***Rhus aromatica****Hypothenemus distinctus* Wood, *Pityophthorus hylocuroides* Wood.***Rhus chinensis****Xylosandrus discolor* (Blandford), *Xylosandrus germanus* (Blandford).***Rhus diversilobata****Cactopinus rhois* Blackman.***Rhus glabra****Hypothenemus seriatus* (Eichhoff).***Rhus integrifolia****Chaetophloeus hystrix* (LeConte), *Chaetophloeus penicillatus* (Bruck).***Rhus lancea****Hypothenemus malus* (Schedl), *Lauregus rhusi* Schedl.***Rhus laurina****Chaetophloeus hystrix* (LeConte), *Stenochyptus sulcatus* (Bruck).***Rhus microphylla****Chaetophloeus penicillatus* (Bruck).***Rhus mucronata****Xyleborinus aemulus* (Wollaston).***Rhus orata****Chaetophloeus hystrix* (LeConte), *Chaetophloeus penicillatus* (Bruck).***Rhus radicans****Pityophthorus lautus* Eichhoff.***Rhus* spp.***Acanthotomicus bidentatus* (Schedl), *Ambrosiodmus rubricollis* (Eichhoff), *Ambrosiodmus tachygraphus* (Zimmermann), *Chaetophloeus lasius* Wood, *Chaetophloeus sulcatus* (Wood), *Corthylocurus mexicanus* (Schedl), *Glostatus acaciae* (Schedl), *Hylesinopsis confusus* (Eggers), *Hylocurus hirtellus* (LeConte), *Hypothenemus interstitialis* (Hopkins), *Micracisella nigrella* Wood, *Pityophthorus debilis* Wood, *Platypus solutus* Schedl, *Polygraphus natalensis* Eggers, *Thysanoes inornatus* Wood, *Xyleborinus saxseni* (Ratzeburg).***Rhus succedanea****Sueus niisimai* (Eggers).***Rhus sylvestris****Ambrosiodmus lewisi* (Blandford), *Lymantria kabei* Murayama.***Rhus taitensis****Platypus solidus* Walker, *Xyleborus viaticus* Schedl.***Rhus toxicodendron****Pityophthorus crinalis* Blackman, *Trischidias atomus* (Hopkins).***Rhus toxicodendron radicans****Cryphalus rhusi* Niisima.***Rhus trichocarpa****Indocryphalus pubipennis* (Blandford).***Rhus trilobata****Cactopinus rhois* Blackman, *Chaetophloeus penicillatus* (Bruck), *Pityophthorus viridis* Blackman.***Rhus typhina****Lymantria decipiens* (LeConte), *Pityophthorus lautus* Eichhoff, *Pityophthorus scriptor* Blackman.***Rhus vernix****Pityophthorus crinalis* Blackman.***Rhus villosa****Platypus solutus* Schedl.***Ribes* spp.***Chaetophloeus heterodoxus* (Casey).***Ricinodendron africanum****Chaetastus tuberculatus* (Chapuis), *Doliopygus lefevrei* Schedl, *Platypus hintzi* Schauffuss, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volvulus* (Fabricius).***Ricinodendron heudelotii****Ambrosiodmus ocatius* (Eggers), *Chaetastus tuberculatus* (Chapuis), *Coccotrypes rotundicollis* (Eggers), *Doliopygus conradti* (Strohmeyer), *Doliopygus interjectus* Schedl, *Doliopygus medius* Schedl, *Doliopygus serratus* (Strohmeyer), *Euwallacea pandae* (Schedl), *Hypothenemus eruditus* Westwood, *Periommatius excisus* Strohmeyer, *Platypus hintzi* Schauffuss, *Platypus impressus* (Strohmeyer), *Platypus picinus* Schedl, *Platypus solutus* Schedl, *Prennobiobius nodulosus* Hagedorn, *Trachyostus aterrimus* (Schauffuss), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus piceus* (Motschulsky), *Xyleborus sub-tuberculatus* Eggers, *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).***Ricinus communis****Acanthotomicus tanganyikaensis* (Schedl), *Euwallacea fornicatus* (Eichhoff), *Euwallacea xanthopus* (Eichhoff), *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Pityophthorus kivuensis* Schedl, *Theoborus ricini* (Eggers), *Xyleborus principalis* Eichhoff, *Xyleborus volvulus* (Fabricius).***Ricinus* spp.***Euwallacea fornicatus* (Eichhoff).***Rinorea* spp.***Dactylipalpus groutelleyi* (Blandford).***Rinorea icelchitschii****Doliopygus subditivus* (Schedl), *Xyleborus piceus* (Motschulsky).***Ritchiea apprealiana****Euwallacea xanthopus* (Eichhoff), *Xyleborus ambasi-pennis* Schedl, *Xyleborus principalis* Eichhoff.***Robinia neomexicana****Chramesus asperatus* Schaeffer.***Robinia pseudacacia****Euwallacea fornicatus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Trischidias atomus* (Hopkins), *Trypodendron domesticum* (Linnaeus).***Robinia* spp.***Chramesus chapuisi* LeConte, *Hylesinus varius* (Fabricius), *Hylocurus ruber* Wood, *Hypothenemus eruditus* Westwood.***Rollinia orethopetala****Coccotrypes carpophagus* (Hornung).***Rollinia* spp.***Xylosandrus compactus* (Eichhoff).***Rosa* spp.***Cnesinus elegans* Blandford, *Xyleborus dispar* (Fabricius).***Roupala complicata****Araptus vinnulus* Wood.***Roupala* spp.***Araptus furvescens* Wood, *Araptus furvus* Wood.***Rubus* spp.***Cnesinus elegans* Blandford, *Cnesinus fulgens* Wood, *Cnesinus lucaris* Wood, *Corthylocurus mexicanus* (Schedl), *Corthyus additus* Wood, *Cryphalus balanopselaphus* Eggers, *Hypothenemus erectus* LeConte,

- Micracisella nitidula* Wood, *Tricholus aciculatus* Wood, *Xylosandrus compactus* (Eichhoff).
- Rudaea amazonica**
Xyleborus spinulosus Blandford.
- Rumex spp.**
Cyrtogenius nubungensis (Schedl).
- Sabal bermudana**
Coccotrypes dactyliperda (Fabricius).
- Sabal palmetto**
Coccotrypes carpophagus (Hornung).
- Sabia parviflora**
Hypothenemus scriatus (Eichhoff).
- Sabina chinensis**
Phloeosinus hopeii Schedl, *Phloeosinus perlatus* Chapuis, *Phloeosinus shensi* Tsai & Yin.
- Saccharum officinarum**
Xyleborus affinis Eichhoff, *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).
- Saccharum spontaneum**
Hypothenemus bauhianae Schedl, *Hypothenemus eruditus* Westwood, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus perforans* (Wollaston), *Xyleborus volutus* (Fabricius).
- Sacogluthia procera**
Phelloterus anaxeus Wood, *Platypus nudatus* Wood, *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius).
- Sacogluthia spp.**
Phelloterus atrocis Wood.
- Sagraea elliptica**
Crossotarsus andamanus Beeson.
- Sagraea laurina**
Xylosandrus crassiusculus (Motschulsky).
- Sahaje jahara**
Sphacrotypes bengalensis Wood.
- Salacia erecta**
Mimiocurus kikusa (Schedl), *Mimiocurus rugicollis* (Schedl).
- Salix babylonica**
Hypothenemus californicus Hopkins.
- Salix caprea**
Trypophloeus binodulus (Ratzeburg).
- Salix fragilis**
Trypophloeus binodulus (Ratzeburg).
- Salix guatemalensis**
Corthylus papulans Eichhoff, *Micracis grandis* Schedl.
- Salix nigra**
Trischidius atomus (Hopkins).
- Salix sachalinensis**
Trypophloeus dejeri Stark.
- Salix scouleriana**
Procrpyphalus utahensis Hopkins, *Trypophloeus striatulus* (Mannerheim).
- Salix silesiaca**
Trypophloeus rybinskii Reitter.
- Salix spp.**
Ambrosiodinus levisi (Blandford), *Corthylus mexicanus* Schedl, *Cryphalus exiguus* Blandford, *Hylocurus hirtellus* (LeConte), *Hylocurus microcoris* Wood, *Hypothenemus columbi* Hopkins, *Hypothenemus scriatus* (Eichhoff), *Lymanator alaskanus* Wood, *Lymanator decipiens* (LeConte), *Micracis carinulatus* Wood, *Micracis detentus* Wood, *Micracis scainui* Blackman, *Micracis tribulatus* Wood, *Micracisella knulli* (Blackman), *Microcorthylus vicinus* Wood, *Procrpyphalus utahensis* Hopkins, *Scolytus scheryzei* Semenov, *Scolytus scolytus* (Fabricius), *Taphrorychus machnoeskii* (Sokanovskii), *Trypophloeus binodulus* (Ratzeburg), *Trypophloeus uiger* Stark, *Trypophloeus salicis* Hopkins.
- Salix tetrasperma**
Scolytoplatypus minimus Hagedorn, *Xyleborus haberkorni* Eggers.
- Salnalia insignis**
Eucallacea andamanensis (Blandford), *Eucallacea interjectus* (Blandford), *Platypus indicus* Strohmeier, *Platypus quadrifissilis* Schedl, *Xyleborus cognatus* Blandford.
- Salnalia malabarica**
Ambrosiodinus minor (Stebbing), *Crossotarsus minax* (Walker), *Eucallacea andamanensis* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Platypus cavius* Strohmeier, *Platypus indicus* Strohmeier, *Platypus latifinis* Walker, *Platypus solidus* Walker, *Platypus uncinatus* Blandford.
- Samanea saman**
Doliopygus chapuisi (Duvivier), *Hypothenemus eruditus* Westwood, *Platypus suffodiens* Sampson, *Xyleborus perforans* (Wollaston), *Xyleborinus andrewesi* (Blandford), *Xyleborinus exiguus* (Walker), *Xyleborus spinulosus* Blandford.
- Samanea spp.**
Eucallacea fornicatus (Eichhoff), *Xylosandrus compactus* (Eichhoff).
- Sambucus canadensis**
Hypothenemus eruditus Westwood, *Xylosandrus compactus* (Eichhoff).
- Sambucus chinensis**
Xylosandrus crassiusculus (Motschulsky).
- Sambucus javanica**
Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).
- Sambucus spp.**
Pityophthorus coronarius Blackman, *Pityophthorus exquisitus* (Blackman), *Pityophthorus pudicus* Blackman, *Pityophthorus sambuci* Blackman, *Xylosandrus compactus* (Eichhoff).
- Sandoricum spp.**
Xylosandrus crassiusculus (Motschulsky).
- Santalum album**
Xyleborus similis Ferrari.
- Santalum spp.**
Ficicis despectus (Walker), *Xylosandrus compactus* (Eichhoff).
- Santiria griffithii**
Crossotarsus terminatus Chapuis, *Cyclorhipidion foersteri* (Hagedorn), *Cyclorhipidion pruinatum* (Blandford).
- Santiria spp.**
Crossotarsus cliens (Schedl).
- Santiria trimera**
Chactastus montanus Schedl, *Doliopygus santiriae* Roberts, *Platypus santiriae* Roberts, *Triozastus caliginosus* Roberts.
- Sapindus oahuensis**
Xyleborus lanaiensis Perkins.
- Sapium baccatum**
Platypus linatipennis Schedl, *Platypus westwoodii* Chapuis.
- Sapium biloculare**
Cnemomyx confinis (Wood).

Sapium ellipticum*Xyleborus scobinatus* Hagedorn.***Sapium eugeniaefolium****Crossotarsus bonvouloiri* Chapuis, *Eucallacea andamanensis* (Blandford), *Eucallacea sibsagaricus* (Eggers), *Eucallacea tristis* (Eggers), *Eucallacea wallacei* (Blandford), *Xyleborinus subgranulatus* (Eggers).***Sapium guineense****Xyleborus voltrulus* (Fabricius).***Sapium mannianum****Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus serratus* (Strohmeyer), *Pityophthorus kivuensis* Schedl, *Platypus solutus* Schedl, *Xyleborus affinis* Eichhoff, *Xyleborus principalis* Eichhoff.***Sapium* spp.***Chaetastus tuberculatus* (Chapuis).***Sapium thelocarpum****Scolytodes pumilus* Wood.***Sapota gonocarpa****Dendrocaninus brasiliensis* (Schedl).***Saraca indica****Cryptocarenum diadematus* Eggers.***Saraca* spp.***Eucallacea andamanensis* (Blandford), *Leptoxyleborus concisus* (Blandford), *Platypus pseudocupulatus* Schedl.***Sarcocephalus cordatus****Diapys quinquespinnatus* Chapuis, *Eucallacea interjectus* (Blandford), *Platypus lepidus* Chapuis, *Xyleborus emarginatus* Eichhoff, *Xyleborus fischeri* Hagedorn, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.***Sarcocephalus* spp.***Platypus bajulus* Schedl, *Premnobius caripennis* Eichhoff, *Premnobius scaxpinosus* Eggers, *Xyleborus alhuandi* Schaufuss, *Xyleborus nitidulus* Eggers.***Sarcocephalus trillesii****Cylindropalpus laudatus* (Schedl), *Doliopygus conradi* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Hypothenemus eruditus* Westwood, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus voltrulus* (Fabricius).***Sarcostemma clausum****Araptus dentifrons* Wood.***Sarothamnus scoparius****Phloeotribus rhododactylus* (Marsham).***Sassafras albidum****Xyleborus sayi* (Hopkins).***Sassafras* spp.***Corthylus punctatissimus* (Zimmermann).***Saurauia pentapetala****Hadrodemius comans* (Sampson).***Scaevola frutescens****Scolytogenes nubilus* (Wood).***Scaphium affine****Crossotarsus wallacei* (Thomson), *Eucallacea fornicatus* (Eichhoff), *Platypus geminatus* Chapuis, *Platypus insularis* Strohmeyer, *Platypus quadrifissilis* Schedl.***Scaphium* spp.***Anasa striatotruncatus* (Schedl), *Coptodryas nuxax* (Schedl), *Crossotarsus terminatus* Chapuis, *Cyclorhi-**pidion agnatum* (Eggers), *Leptoxyleborus concisus* (Blandford), *Platypus westwoodi* Chapuis.***Schefflera aromatica****Cryphalus tenuis* Schedl.***Schefflera manni****Triozastus caliginosus* Roberts.***Schima cordatus****Platypus rethi* Strohmeyer.***Schima noronhae****Crossotarsus edentatus* Beeson, *Platypus biuncus* Blandford, *Platypus catus* Strohmeyer, *Platypus signatus* Chapuis, *Platypus solidus* Walker.***Schima* spp.***Ambrosiodmus apicalis* (Blandford), *Xyleborus seriatus* Blandford.***Schima superba****Ambrosiodmus rubricollis* (Eichhoff), *Xyleborus pfeili* (Ratzeburg), *Xylosandrus discolor* (Blandford).***Schima wallichii****Platypus catus* Strohmeyer, *Platypus indicus* Strohmeyer, *Platypus solidus* Walker, *Xyleborus schoreae* (Stebbing).***Schinus* spp.***Eucallacea fornicatus* (Eichhoff), *Xylosandrus compactus* (Eichhoff).***Schizolobium excelsum****Chaetastus tuberculatus* (Chapuis).***Schizolobium* spp.***Xyleborus affinis* Eichhoff.***Schizomeria* spp.***Platypus concentriporus* Roberts, *Platypus valdephlebius* Roberts.***Schleichera oleosa****Diapys molossus* Chapuis, *Eucallacea andamanensis* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Platypus solidus* Walker, *Xylosandrus morigerus* (Blandford).***Schleichera trijuga****Hypothenemus crudiae* (Panzer), *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston).***Schotia latifolia****Xyleborinus acmulus* (Wollaston).***Schoutenia accrescens****Anasa versicolor* (Sampson), *Crossotarsus squannulatus* Chapuis, *Crossotarsus wallacei* (Thomson).***Schoutenia mastersii****Crossotarsus siporamus* Schedl.***Schoutenia orata****Hypothenemus eruditus* Westwood.***Sclerocarya biffra****Ambrosiodmus eichhoffi* (Schreiner).***Sclerocarya birrea****Xyleborinus sclerocaryae* (Schedl).***Sclerocarya caffra****Xyleborinus sclerocaryae* (Schedl).***Sclerocarya* spp.***Ambrosiodmus eichhoffi* (Schreiner).***Sclerosperma mannii****Coccotrypes congenus* Eggers, *Coccotrypes nigripes* Eggers.***Scolopia brownii****Platypus subgranosus* Schedl.***Scorodocarpus borneensis****Cnestus aterrimus* (Eggers).***Scorodophloeus zenkeri****Chaetastus tuberculatus* (Chapuis), *Cylindropalpus affinis* Strohmeyer, *Cylindropalpus auricomans* (Schaufuss).

fuss), *Cylindropalpus granulosus* (Schedl), *Cylindropalpus laudatus* (Schedl), *Dactylipalpus canerinus* Hagedorn, *Doliopygus chapuisi* (Duvivier), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus donisi* Schedl, *Doliopygus ditarsicus* (Schedl), *Doliopygus donisi* Schedl, *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus lateralis* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus maynei* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus notatus* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus submarginatus* Schedl, *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Hypothenemus cruditus* Westwood, *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius inermis* Strohmeyer, *Periommatius longicollis* Chapuis, *Periommatius similis* Strohmeyer, *Periommatius subrobustus* Schedl, *Platypus hintzi* Schaufuss, *Platypus inpressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Platypus picinus* Schedl, *Trachyostus aterrinus* (Schaufuss), *Trachyostus decellei* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Trachyostus tomentosus* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborinus polyalthiae* (Schedl), *Xyleborinus similans* (Eggers), *Xyleborinus spinipes* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus katangensis* Eggers, *Xyleborus picus* (Motschulsky), *Xyleborus ustus* Schedl, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Scottellia coriacea
Doliopygus obduensis Roberts.

Scutia connersonii
Hypothenemus scutiae (Schedl).

Scutia myrtina
Hypothenemus marshalli (Eggers), *Hypothenemus morigerus* (Schedl).

Seaforthia spp.
Coccotrypes dactyliperda (Fabricius).

Sechium edule
Cnesinus degener Wood, *Dendrocraniulus declivis* Schedl, *Dendrocraniulus gracilis* Wood, *Dendrocraniulus limus* Wood, *Dendrocraniulus mexicanus* Wood, *Dendrocraniulus tardulus* Wood.

Semecarpus anacardium
Crossotarsus externedentatus (Fairmaire), *Xyleborinus artetriatus* (Eichhoff).

Semecarpus curtisii
Coccotrypes cinnamomi (Eggers).

Semecarpus gardneri
Amasa versicolor (Sampson), *Crossotarsus minax* (Walker), *Crossotarsus terminatus* Chapuis.

Semecarpus heterophylla
Platypus biuncus Blandford, *Platypus solidus* Walker.

Semecarpus merrillana
Xyleborus similis Ferrari.

Semecarpus spp.
Diapus quinquespinatus Chapuis.

Senecio erechthitoides
Xyleborinus forficuloides (Schedl).

Senecio erioneuron
Miniocurus ruenzoricensis (Schedl).

Senecio spp.

Pityophthorus paulus Wood, *Thammurgus senecionis* Schedl, *Thammurgus wittvi* Eggers.

Sequoia gigantea

Phlocosinus punctatus LeConte.

Sequoia sempervirens

Cryphalus pubescens Hopkins, *Phlocosinus cupressi* Hopkins, *Phlocosinus punctatus* LeConte, *Phlocosinus sequoiae* Hopkins.

Sequoia spp.

Phlocosinus bicolor (Brulle), *Phlocosinus thujae* (Perris).

Sericolea spp.

Diapus bilunatus Schedl.

Serjania mexicana

Cnesinus setulosus Blandford, *Stegomerus vulgaris* Wood.

Serjania racemosa

Hypothenemus crudiae (Panzer), *Micracisella serjaniae* Wood.

Serjania spp.

Bothrostermus forcatus (Blackman), *Bothrostermus hirsutus* Wood, *Bothrostermus truncatus* Eichhoff, *Cnesinus degener* Wood, *Cnesinus elegans* Blandford, *Cnesinus gracilis* Blandford, *Cnesinus retifer* Wood, *Cnesinus squamosus* Wood, *Corthylocorus debilis* Wood, *Cryptocareus dialematus* Eggers, *Cryptocareus lepidus* Wood, *Eupagiocerus ater* Eggers, *Hypothenemus brunneus* (Hopkins), *Hypothenemus columbi* Hopkins, *Hypothenemus erectus* LeConte, *Hypothenemus indigenus* Wood, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus plumieriae* (Nordlinger), *Hypothenemus seriatus* (Eichhoff), *Hypothenemus solocis* Wood, *Hypothenemus squamosus* (Hopkins), *Hypothenemus tereis* Wood, *Hypothenemus trivialis* Wood, *Mecopeplus zetecki* Blackman, *Micracisella divaricata* Wood, *Micracisella similis* Wood, *Micracisella squamatula* Wood, *Micracisella striata* Wood, *Micracisella vescula* Wood, *Microcorthyus minimus* Schedl, *Scolytogenes knabi* (Hopkins), *Stegomerus mexicanus* Wood, *Stegomerus vulgaris* Wood, *Tricolus badius* Wood, *Xylosandrus curtulus* (Eichhoff).

Serjania triquetra

Phloeotribus opimus Wood, *Stegomerus vulgaris* Wood.

Sesbania spp.

Euwallacea xanthopus (Eichhoff), *Xyleborus alluaudi* Schaufuss, *Xyleborus principalis* Eichhoff.

Sejara spp.

Platypus agnatus Schedl.

Shiia sieboldii

Xyleborus pelliculosus Eichhoff, *Xyleborus volutus* (Fabricius).

Shiia spp.

Leptoxyleborus concisus (Blandford).

Shorea acuminata

Cnestus aterrinus (Eggers), *Dendroplatypus impar* (Schedl), *Dryococtops nitidus* (Schedl), *Xyleborus bidentatus* (Motschulsky), *Xyleborus volutus* (Fabricius).

Shorea ussamica

Diapus quinquespinatus Chapuis, *Euwallacea andamanensis* (Blandford), *Euwallacea interjectus* (Blandford), *Genyocerus diaphanus* (Schedl), *Platypus bisignatus* Schedl, *Platypus shoreanus* (Beeson), *Sphacrotypes siculikeus* Stebbing, *Xyleborus perforans* (Wollaston).

Shorea balanocarpoides

Arixyleborus minor (Eggers), *Arixyleborus suturalis* (Eggers), *Webbia quattuordecimspinitus* Sampson, *Xyleborus ciliatiformis* Schedl, *Xyleborus latus* Eggers.

Shorea bentoungensis

Diapus quinquespinatus Chapuis, *Platypus bifurcus* (Schedl), *Platypus curtus* Chapuis, *Platypus lunifer* Schedl.

Shorea bracteolata

Cnestus bicornis (Eggers), *Cyclorhipidion agnatum* (Eggers), *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Genyocerus quadriporus* (Schedl), *Genyocerus serratus* (Schedl), *Hadrodemius comans* (Sampson), *Platypus bifurcus* (Schedl), *Platypus insularis* Strohmeier, *Platypus shoreanus* (Beeson), *Webbia bicornis* (Schedl), *Xyleborus perforans* (Wollaston), *Xylosandrus ater* (Eggers), *Xylosandrus gravidus* (Blandford), *Xylosandrus mancus* (Blandford).

Shorea curtisii

Amasa versicolor (Sampson), *Coptodryas confusa* (Hopkins), *Coptodryas nugax* (Schedl), *Coptodryas perparva* (Sampson), *Cryptoxyleborus subnervus* Schedl, *Dendroplatypus impar* (Schedl), *Webbia bituberculatus* Browne, *Webbia duodecimspinitus* Schedl, *Xyleborus minutus* Blandford, *Xylosandrus crassiusculus* (Motschulsky).

Shorea dasphylla

Dendroplatypus impar (Schedl), *Platypus bifurcus* (Schedl), *Xylosandrus mediocris* (Schedl).

Shorea exima

Taphrodasus percorthylus (Schedl).

Shorea faquetiana

Baiocis angusticeps (Schedl), *Diapus quinquespinatus* Chapuis, *Platypus curtus* Chapuis, *Platypus pseudocurtus* Schedl, *Platytarsulus elongatus* Schedl.

Shorea guiso

Platypus cupulatus Chapuis, *Platypus insularis* Strohmeier, *Xylosandrus crassiusculus* (Motschulsky).

Shorea gysbertsiana

Cryphalus rugosus (Schedl), *Cryptoxyleborus opacicaudulus* Schedl, *Ptilopodius squamosus* Schedl, *Xyleborinus micrographus* (Schedl).

Shorea hypochra

Genyocerus diaphanus (Schedl).

Shorea kunstleri

Diapus pusillimus Chapuis, *Diapus quinquespinatus* Chapuis.

Shorea laevis

Baiocis pasohensis (Schedl), *Genyocerus exilis* (Schedl), *Genyocerus serratus* (Schedl), *Xyleborus triangi* Schedl.

Shorea lepidota

Dendroplatypus impar (Schedl).

Shorea leprosula

Amasa versicolor (Sampson), *Amasa striatotruncatus* (Schedl), *Arixyleborus granulifer* (Eggers), *Arixyleborus leprosolus* Schedl, *Arixyleborus minor* (Eggers), *Arixyleborus scabripennis* (Blandford), *Arixyleborus suturalis* (Eggers), *Baiocis pernambus* (Schedl), *Cnestus bicornis* (Eggers), *Coccotrypes subcubrosus* (Blandford), *Coptodryas abbreviata* (Schedl), *Coptodryas corporaali* (Eggers), *Coptodryas curvidentis* (Schedl), *Coptodryas nugax* (Schedl), *Coptodryas undulata* (Sampson), *Crossotarsus brownei* Schedl, *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus squamulatus* Chapuis, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Cryptoxyleborus confusus* Browne, *Cryptoxyleborus opacicaudulus* Schedl,

Cyclorhipidion agnatum (Eggers), *Cyclorhipidion arnaticeps* (Schedl), *Cyclorhipidion foersteri* (Hagedorn), *Dendroplatypus impar* (Schedl), *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Eccoptopterus gracilipes* (Eichhoff), *Eccoptopterus limbatus* Sampson, *Eccoptopterus spinosus* (Olivier), *Eucallacea xanthopus* (Eichhoff), *Ficicis javanus* (Eggers), *Genyocerus biporus* (Schedl), *Genyocerus diaphanus* (Schedl), *Genyocerus laticollis* (Browne), *Genyocerus philippinensis* (Schedl), *Genyocerus serratus* (Schedl), *Hadrodemius comans* (Sampson), *Hypothenemus cuniculus* (Schedl), *Hypothenemus eruditus* Westwood, *Leptoxyleborus concisus* (Blandford), *Platypus bifurcus* (Schedl), *Platypus bilopennis* Schedl, *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Platypus forcicula* Chapuis, *Platypus fraterculus* Schedl, *Platypus hybridus* Schedl, *Platypus impressifrons* Schedl, *Platypus insulindicus* Schedl, *Platypus lepidus* Chapuis, *Platypus pseudocurtus* Schedl, *Platypus quadrifissilis* Schedl, *Platypus solidus* Walker, *Schedia sumatranus* (Hagedorn), *Sphaerotrypes minutus* Browne, *Taphrodasus percorthylus* (Schedl), *Webbia bifornis* Browne, *Webbia cornutus* Schedl, *Webbia cylindricus* Schedl, *Xyleborinus andrewesi* (Blandford), *Xyleborus apertus* Schedl, *Xyleborus approximatus* Schedl, *Xyleborus emarginatus* Eichhoff, *Xyleborus javanus* Eggers, *Xyleborus kajanensis* Schedl, *Xyleborus latus* Eggers, *Xyleborus major* (Stebbing), *Xyleborus perforans* (Wollaston), *Xyleborus semistriatus* Schedl, *Xylosandrus ater* (Eggers), *Xylosandrus difficilis* (Eggers), *Xylosandrus mancus* (Blandford), *Xylosandrus morigerus* (Blandford).

Shorea macroptera

Amasa versicolor (Sampson), *Arixyleborus suturalis* (Eggers), *Coptodryas nugax* (Schedl), *Coptodryas perparva* (Sampson), *Cryptoxyleborus opacicaudulus* Schedl, *Cyclorhipidion agnatum* (Eggers), *Dendroplatypus impar* (Schedl), *Genyocerus serratus* (Schedl), *Hypothenemus eruditus* Westwood, *Ptilopodius shoreae* Schedl, *Webbia cornutus* Schedl, *Webbia cylindricus* Schedl, *Webbia micrographus* (Schedl), *Xyleborus approximatus* Schedl, *Xyleborus macropterus* Schedl, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus difficilis* (Eggers), *Xylosandrus mancus* (Blandford).

Shorea materialis

Xyleborus perforans (Wollaston).

Shorea maxcelliana

Coptodryas perparva (Sampson), *Crossotarsus squamulatus* Chapuis, *Genyocerus diaphanus* (Schedl), *Genyocerus philippinensis* (Schedl), *Genyocerus serratus* (Schedl), *Hadrodemius comans* (Sampson), *Platypus cupulatus* Chapuis, *Platypus pseudocupulatus* Schedl, *Webbia bituberculatus* Browne, *Webbia duodecimspinitus* Schedl, *Webbia suturalis* Browne, *Xyleborus cylindromorphus* Eggers, *Xyleborus haberkorni* Eggers, *Xyleborus punilus* Eggers, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus mancus* (Blandford).

Shorea multiflora

Coptodryas confusa Hopkins.

Shorea negrosensis

Platypus bifurcus (Schedl).

Shorea ovalis

Amasa versicolor (Sampson), *Arixyleborus granulifer* (Eggers), *Arixyleborus scabripennis* (Blandford), *Cnestus bicornis* (Eggers), *Crossotarsus externedentatus* (Fairmaire), *Dendroplatypus impar* (Schedl),

Leptoxyleborus concisus (Blandford), *Platypus cupulatus* Chapuis, *Xyleborinus andrevesi* (Blandford), *Xyleborus macropterus* Schedl, *Xylosandrus difficilis* (Eggers).

Shorea ovata

Coptodryas perparva (Sampson), *Cryptoxyleborus opacicaudulus* Schedl, *Platypus balanocarpus* Schedl, *Xyleborus haberkorni* Eggers, *Xylosandrus maucus* (Blandford).

Shorea pauciflora

Ptilopodius squamosus Schedl.

Shorea parrifolia

Anasa cervicolar (Sampson), *Arixyleborus scabripennis* (Blandford), *Coptodryas confusa* Hopkins, *Coptodryas quadricostata* (Schedl), *Crossotarsus terminatus* Chapuis, *Cryptoxyleborus nanus* Browne, *Cyclorhipidion sexspiniatum* (Schedl), *Dendroplatypus impar* (Schedl), *Diapus quinquespinatus* Chapuis, *Genyocerus biporus* (Schedl), *Genyocerus quadriporus* (Schedl), *Platypus balanocarpus* Schedl, *Platypus bifurcus* (Schedl), *Platypus bilobipennis* Schedl, *Platypus curtus* Chapuis, *Platypus pseudocupulatus* Schedl, *Ptilopodius squamosus* Schedl, *Sphaerotrypes rufopalliatius* Schedl, *Webbia obtusispinosus* Schedl, *Xyleborus javanus* Eggers, *Xyleborus persimilis* Eggers.

Shorea pauciflora

Dendroplatypus impar (Schedl), *Platypus bilobipennis* Schedl, *Platypus curtus* Chapuis, *Platypus pseudocupulatus* Schedl.

Shorea platycarpa

Cryptoxyleborus subnervus Schedl, *Dendroplatypus impar* (Schedl).

Shorea robusta

Anasa schlichii (Stebbing), *Ambrosiodmus minor* (Stebbing), *Coccotrypes carpophagus* (Hornung), *Coccotrypes mibilis* (Blandford), *Coptodryas elegans* (Sampson), *Coptodryas perparva* (Sampson), *Coptodryas recidens* (Sampson), *Coptodryas undulata* (Sampson), *Crossotarsus boureouloiri* Chapuis, *Crossotarsus externedentatus* (Fairmaire), *Cryptoxyleborus turbinus* (Sampson), *Diapus quinquespinatus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Eucallacea andamanensis* (Blandford), *Eucallacea aplanatus* (Wichmann), *Eucallacea bicolor* (Blandford), *Eucallacea interjectus* (Blandford), *Genyocerus furtivus* (Sampson), *Platypus curvus* Strohmeier, *Platypus cupulatus* Chapuis, *Platypus curtatus* Sampson, *Platypus curtus* Chapuis, *Platypus perrisi* Chapuis, *Platypus solidus* Walker, *Sphaerotrypes globulus* Blandford, *Sphaerotrypes sitcalikensis* Stebbing, *Webbia pabo* Sampson, *Xyleborinus andrevesi* (Blandford), *Xyleborinus artistriatus* (Eichhoff), *Xyleborus glabratus* Eichhoff, *Xyleborus haberkorni* Eggers, *Xyleborus major* (Stebbing), *Xyleborus mucronatulus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus mesuae* (Eggers).

Shorea rugosa

Cyclorhipidion kelantanum (Schedl), *Xyleborus perforans* (Wollaston), *Xyleborus persimilis* Eggers.

Shorea scabrida

Cyclorhipidion tuberculosissimum (Eggers).

Shorea scutulata

Ambrosiodmus lantanac (Eggers), *Xyleborus lateccari-natus* Schedl.

Shorea sericea

Platypus curtus Chapuis, *Xyleborus macropterus* Schedl.

Shorea sideroxylon

Cryphalus substriatus Schedl.

Shorea singkawang

Dendroplatypus impar (Schedl), *Diapus pusillimus* Chapuis, *Platypus bifurcus* (Schedl), *Platypus shoreanus* (Beeson).

Shorea spp.

Anasa schlichii (Stebbing), *Ambrosiodmus restrictus* (Schedl), *Arixyleborus medius* (Eggers), *Arixyleborus okadae* Browne, *Arixyleborus rugosipes* Hopkins, *Arixyleborus scabripennis* (Blandford), *Cnestus suturalis* (Eggers), *Coccotrypes gedeanus* (Eggers), *Coptodryas alpha* (Beeson), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus schelli* Browne, *Crossotarsus squamulatus* Chapuis, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Cryptoxyleborus naevus* Schedl, *Cryptoxyleborus nanus* Browne, *Cryptoxyleborus shoreae* Browne, *Cryptoxyleborus subnervus* Schedl, *Cyclorhipidion agnatum* (Eggers), *Diapus pusillimus* Chapuis, *Diapus quinquespinatus* Chapuis, *Eccoptopterus sagittarius* Schedl, *Eucallacea fomicatus* (Eichhoff), *Genyocerus serratus* (Schedl), *Genyocerus sexporus* (Schedl), *Genyocerus shoreae* Browne, *Hadrodemus comans* (Sampson), *Leptoxyleborus concisus* (Blandford), *Leptoxyleborus puer* (Eggers), *Leptoxyleborus scabrior* (Schedl), *Platypus agnatus* Schedl, *Platypus arduus* (Schedl), *Platypus bicandatus* Browne, *Platypus bifurcus* (Schedl), *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Platypus decens* Sampson, *Platypus emarginatus* Browne, *Platypus franciae* Browne, *Platypus histrix* (Schedl), *Platypus longicollis* Browne, *Platypus octospinosus* Browne, *Platypus pseudocupulatus* Schedl, *Platypus pumilus* Browne, *Platypus subtruncatus* Browne, *Platypus szechui* Browne, *Platypus transformis* Schedl, *Platypus uncinatus* Blandford, *Scolytplatypus parvus* Sampson, *Sphaerotrypes incernis* Browne, *Webbia bakoensis* Browne, *Webbia cylindricus* Schedl, *Webbia dasyurus* Browne, *Webbia dipterocarpi* Hopkins, *Webbia multidentatus* Browne, *Webbia platypoides* Eggers, *Webbia quattuordecimspinosus* Sampson, *Webbia simplex* (Browne), *Xyleborus abscissus* Browne, *Xyleborus amphicranoides* Hagedorn, *Xyleborus dolosus* Blandford, *Xyleborus excusus* Schedl, *Xyleborus fallax* Eichhoff, *Xyleborus granurus* Browne, *Xyleborus amplexicauda* Hagedorn, *Xyleborus decumanus* Schedl, *Xyleborus javanus* Eggers, *Xyleborus latus* Eggers, *Xyleborus minutus* Blandford, *Xyleborus nigrescens* Browne, *Xyleborus obiensis* Browne, *Xyleborus perforans* (Wollaston), *Xyleborus pseudopilifer* Schedl, *Xyleborus pseudocylindricus* Eggers, *Xyleborus shoreae* (Stebbing), *Xyleborus sichus* Schedl, *Xyleborus trispinatus* Browne, *Xyleborus tuberculosus* Browne, *Xyleborus tunggalii* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus difficilis* (Eggers), *Xylosandrus maucus* (Blandford).

Shorea sumatraua

Cnestus bicornoides (Schedl), *Cnestus bicornis* (Eggers), *Xyleborinus penninutissimus* (Schedl), *Xyleborus javanus* Eggers, *Xylosandrus ater* (Eggers), *Xylosandrus compactus* (Eichhoff), *Xylosandrus maucus* (Blandford).

Shorea tahura

Diaprus tahurae Stebbing, *Genyocerus furtivus* (Sampson), *Xyleborinus andrewesi* (Blandford).

Shorea teysmanniana

Dendroplatypus inpar (Schedl).

Shorea uliginosa

Arixyleborus scabripennis (Blandford), *Cyclorhipidion agnatum* (Eggers), *Dendroplatypus inpar* (Schedl), *Leptoxyleborus depressus* (Eggers), *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Tetroplatypus trepanatus* (Chapuis), *Xyleborus apertus* Schedl, *Xyleborus perforans* (Wollaston).

Sicydium taminifolium

Dendrocranus consimilis Wood.

Sida rhombifolia

Hypothenemus crudiae (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus gossypii* (Hopkins), *Hypothenemus seriatus* (Eichhoff), *Thamnurgus granulicollis* Schedl, *Xyloctetes sidanus* (Schedl).

Sideroxylon inerme

Xyloctonus maculatus Schedl.

Sideroxylon longepetiolatum

Scolytomimus andamanensis Wood.

Sideroxylon macranthum

Eucallacea bicolor (Blandford), *Xyleborus piceus* (Motschulsky), *Xyleborus similis* Ferrari.

Sideroxylon sandwicensis

Crossotarsus externodentatus (Fairmaire).

Sideroxylon spp.

Chaetastus montanus Schedl.

Simaruba amara

Xyleborus ferrugineus (Fabricius).

Sindora siamensis

Acacis trahax (Sampson).

Sindora spp.

Diaprus quinquespinatus Chapuis, *Platypus forficula* Chapuis, *Platypus sindorae* Browne.

Siparuna guianensis

Pericoryphalus sobrinus Wood.

Siparuna nicaraguensis

Corthyceylon aztecum (Bright), *Corthyceylon tardum* Wood, *Corthylyus procerus* Bright, *Corthylyus serratus* Wood, *Corthylyus villiger* Wood, *Microcorthylyus inermis* Wood, *Microcorthylyus ocellaris* Wood, *Tricholus inornatus* Wood, *Tricholus senex* Schedl.

Sloanea forbesii

Platypus jansoni Chapuis, *Platypus solidus* Walker.

Sloanea multiflora

Xyleborinus interstosus (Blandford), *Xyleborus affinis* Eichhoff, *Xyleborus geayi* Hagedorn.

Sloanea spp.

Diaprus pusillimus Chapuis, *Xyleborus parcellus* Wood.

Sloetia elongata

Crossotarsus externodentatus (Fairmaire), *Cryphalus substriatus* Schedl, *Hypocryphalus perminimus* (Schedl), *Leptoxyleborus concisus* (Blandford).

Smilax aspera

Chaetoptelius vestitus (Mulsant & Rey).

Smilax china

Cryphalus jeholensis Murayama, *Xylosandrus brevis* (Eichhoff).

Smilax spp.

Chramesus exilis Wood, *Cucsinus strigicollis* LeConte, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood.

Smithia thymodora

Eucallacea xanthopus (Eichhoff).

Solanum aculeastrum

Coptoborus usagarius (Eggers), *Cyrtogenius mulungensis* (Schedl), *Eucallacea xanthopus* (Eichhoff), *Xyleborinus collarti* (Eggers), *Xyleborus piceus* (Motschulsky), *Xyleborus principalis* Eichhoff.

Solanum flamigni

Eucallacea xanthopus (Eichhoff).

Solanum spp.

Eucallacea xanthopus (Eichhoff), *Xylosandrus compactus* (Eichhoff).

Solanum torcum

Corthylyus comatus Blandford, *Corthylyus retusus* Wood.

Soumeratia apelata

Platypus maritimus Schedl, *Xyleborus bidentatus* (Motschulsky).

Sophora japonica

Ambrosiodmus rubricollis (Eichhoff), *Hypothenemus eruditus* Westwood, *Xylosandrus discolor* (Blandford).

Sorbus ahnifolia

Xyleborus dispar (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Sorbus aucuparia

Pteleobius krautzi (Eichhoff).

Sorbus commixta

Cryphalus mahus Niisima, *Polygraphus nigriclytris* Niisima, *Scolytus ellipticus* Murayama.

Sorbus spp.

Scolytus mali (Bechstein), *Scolytus rugulosus* (Muller), *Trypodendron domesticum* (Linnaeus).

Sorghum saccharatum

Hypothenemus eruditus Westwood.

Sorindeia lemairei

Doliopygus conradi (Strohmeyer), *Polygraphus tropicus* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus solutus* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Sorocea illicifolia

Prennobiobius catipennis Eichhoff.

Spartium junceum

Liparthrum genistae (Anbe), *Phloeotribus cristatus* (Favet), *Phloeotribus pseudocristatus* (Pfeffer), *Phloeotribus rhododactylus* (Marsham), *Phloeotribus sharpi* (Guillebeau).

Spartium scoparium

Hylastinus obscurus (Marsham).

Spartocystus nubigenus

Liparthrum curtum Wollaston.

Spathodea campanulata

Doliopygus erichsoni (Chapuis), *Micocryphalus attenuatus* (Eggers).

Sphaeracula fasciculata

Pseudothysanocles bartoni Bruck.

Spherostenon spp.

Diaprus elongatus Schedl, *Diaprus turgidus* Roberts.

Spirostachys africana

Styracoptinus murex (Blandford).

Spirostachys spp.

Scolytogenes spirostachius (Schedl).

Spondianthus preusii

Chaetastus tuberculatus (Chapuis), *Diaprus quinquespinatus* Chapuis, *Doliopygus dubius* (Sampson), *Hypothenemus morio* (Eggers), *Xyleborus ferrugineus* (Fabricius).

Spondias dulcis

Xyleborus ferrugineus (Fabricius).

Spondias mangifera

Coccotrypes longior (Eggers), *Crossotarsus minax* (Walker), *Diaprus quinquespinatus* Chapuis, *Eucallacea interjectus* (Blandford), *Platypus forficula* Chapuis, *Platypus indicus* Strohmeyer, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Spondias mombini

Acanthotomicus analogus (Wood), *Acanthotomicus fortis* (Wood), *Acanthotomicus ipsiformis* Wood, *Acanthotomicus minimus* (Schedl), *Corthylocurus aguacatenis* (Schedl), *Corthythus papulans* Eichhoff, *Doliopygus serratus* (Strohmeyer), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Styphlosoma granulatum* Blandford, *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus morulus* Blandford, *Xyleborus piceus* (Motschulsky), *Xylosandrus laticeps* (Wood).

Spondias purpurea

Corthylocurus barbatus (Blandford), *Corthythus concisus* Wood, *Corthythus papulans* Eichhoff, *Microcorthythus parvulus* Ferrari, *Monarthrum bifoveatum* Wood, *Monarthrum fimbriatocorne* (Blandford), *Monarthrum lobatum* (Ferrari), *Monarthrum robustum* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus ferrugineus* (Fabricius), *Xyleborus posticus* Eichhoff.

Spondias spp.

Crossotarsus externedentatus (Fairmaire), *Eucallacea fornicatus* (Eichhoff), *Platypus jansoni* Chapuis, *Platypus negatus* Schedl, *Xylosandrus compactus* (Eichhoff).

Stachys sylvatica

Thammurgus kaltenbachi (Bach).

Stachytarpheta spp.

Xylosandrus compactus (Eichhoff).

Staphylea trifolia

Phlocotribus scabricollis (Hopkins).

Staudtia gaboneusis

Chortastus minimus Hagedorn, *Coccotrypes rotundicollis* (Eggers).

Staudtia stipitata

Chaetastus persimilis (Schedl), *Chaetastus tuberculatus* (Chapuis), *Chortastus camerunus* Schaufuss, *Chortastus medius* Eggers, *Chortastus minimus* Hagedorn, *Cylindropalpus laudatus* (Schedl), *Doliopygus artespinitus* (Schedl), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus lebruni* Schedl, *Doliopygus lefevrei* Schedl, *Doliopygus megaloma* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicornis* Schedl, *Doliopygus woltschei* Schedl, *Hypothenemus cruditus* Westwood, *Periommatius excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Platypus solutus* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Xyleborinus gracilipennis* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus nitidulus* Eggers, *Xyleborus rothkireli* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Stephegyne diversifolia

Ptilopodius stephegymis Hopkins, *Xyleborus bidentatus* (Motschulsky).

Sterculia alata

Ambrosiodmus funereus (Lea), *Eucallacea interjectus* (Blandford), *Xyleborinus andrewesi* (Blandford), *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky).

Sterculia bequaertii

Chaetastus tuberculatus (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus expletus* Schedl, *Doliopygus ghesquieri* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus piceus* (Motschulsky).

Sterculia campanulata

Eucallacea interjectus (Blandford), *Platypus jansoni* Chapuis, *Platypus solidus* Walker, *Xyleborus cognatus* Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus pfcili* (Ratzeburg), *Xyleborus punilus* Eggers, *Xyleborus similis* Ferrari.

Sterculia colorata

Eucallacea sibsagaricus (Eggers), *Eucallacea tristis* (Eggers), *Eucallacea wallacei* (Blandford), *Xylosandrus crassiusculus* (Motschulsky).

Sterculia dawei

Chaetastus tuberculatus (Chapuis).

Sterculia foetida

Platypus forficula Chapuis, *Platypus jansoni* Chapuis, *Platypus pseudocupulatus* Schedl.

Sterculia macrophylla

Crossotarsus inermis Schedl, *Crossotarsus schedli* Browne, *Eucallacea xanthopus* (Eichhoff), *Ficicis despectus* (Walker), *Hypothenemus javanus* (Eggers), *Platypus lepidus* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus signatus* Chapuis, *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky).

Sterculia oblonga

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Diaprus quinquespinatus* Chapuis, *Doliopygus expletus* Schedl, *Doliopygus rapax* (Sampson), *Periommatius pseudomajor* Schedl, *Platypus orientalis* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Sterculia ornata

Crossotarsus emorsus Beeson, *Eucallacea interjectus* (Blandford), *Platypus secretus* Sampson, *Xyleborus perforans* (Wollaston).

Sterculia pruriens

Hypothenemus teretis Wood.

Sterculia quinqueloba

Doliopygus serratus (Strohmeyer), *Doliopygus unicornis* Schedl, *Periommatius excisus* Strohmeyer, *Polygraphus bicolor* Eggers, *Xyleborus affinis* Eichhoff, *Xyleborus ambasiusculus* Eggers, *Xyleborus principalis* Eichhoff, *Xyleborus volutus* (Fabricius).

Sterculia rhinopetala

Cylindropalpus auricomans (Schaufuss), *Doliopygus brevis* (Strohmeyer), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus malkini* Schedl, *Doliopygus pseudoserratus* Roberts, *Doliopygus tenuis* (Strohmeyer), *Periommatius excisus* Strohmeyer, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Trachyostus aterrimus* (Schaufuss), *Triozastus banghaasi* (Schaufuss), *Xyleborinus similans*

(Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus barinbiensis* Eggers.

Sterculia spp.

Ambrosiodmus tropicus (Hagedorn), *Arixyleborus medius* (Eggers), *Diaprus nanodontus* Roberts, *Diaprus nanus* Schedl, *Diaprus papuanus* Schedl, *Diaprus pusillimus* Chapuis, *Dryocoetoides cristatus* (Fabricius), *Eucallacea fornicatus* (Eichhoff), *Hypothecinus eruditus* Westwood, *Platypus jansonii* Chapuis, *Platypus verehnutus* (Beeson), *Premnobius catipennis* Eichhoff, *Scolytotplatypus occidentalis* Browne, *Xyleborus piceus* (Motschulsky).

Sterculia tragacantha

Doliopygus conradti (Strohmeyer), *Xyleborus alluaudi* Schaufuss, *Xyleborus piceus* (Motschulsky).

Sterculia urceolata

Platypus signatus Chapuis.

Sterculia villosa

Eucallacea interjectus (Blandford), *Platypus quadrifissilis* Schedl, *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston), *Xyleborus punilus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Stereospermum chelenoides

Platypus solidus Walker.

Stereospermum neuranthum

Leptoxyleborus sordicauda (Motschulsky).

Stereospermum spp.

Leptoxyleborus sordicauda (Motschulsky).

Stereospermum suaveolens

Scolytogenes indicus Wood.

Storckia spp.

Cryphalus striatus Browne.

Straussia spp.

Xyleborus ferrugineus (Fabricius), *Xyleborus perforans* (Wollaston).

Streblus elongatus

Hadrodemus comans (Sampson).

Strombosia glaucescens

Chaetastus persimilis (Schedl), *Chaetastus tuberculatus* (Chapuis), *Cylindropalpus laudatus* (Schedl), *Doliopygus brevis* (Strohmeyer), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lateralis* Schedl, *Doliopygus lebrunii* Schedl, *Doliopygus medius* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus propinquus* Schedl, *Doliopygus semipilosus* (Schedl), *Periommatous excisus* Strohmeyer, *Periommatous grandis* Schedl, *Periommatous longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus parallelus* (Fabricius), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Strombosia grandifolia

Ambrosiodmus oratus (Eggers), *Chaetastus tuberculatus* (Chapuis), *Doliopygus lecomtei* Schedl, *Doliopygus maynei* Schedl, *Doliopygus proximus* Schedl, *Xylocleptes sidanus* (Schedl).

Strombosia javanica

Anasa versicolor (Sampson), *Coptodryas nuxag* (Schedl).

Strombosia pustulata

Chaetastus tuberculatus (Chapuis), *Ctonoxylon spathi-*

fer Schedl, *Doliopygus guineensis* Roberts, *Doliopygus interjectus* Schedl, *Periommatous abruptus* Numberg, *Triozastus banghaasi* (Schaufuss), *Triozastus elongatus* Schedl, *Xyleborus ambasiusculus* Eggers.

Strombosia scheffleri

Dactylipalpus grouvellei (Blandford), *Doliopygus bicentatus* (Strohmeyer), *Doliopygus bilobatus* (Schedl), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus lebrunii* Schedl, *Doliopygus leferrei* Schedl, *Doliopygus maynei* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus proximus* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus unicornis* Schedl, *Platypus impressus* (Strohmeyer), *Platypus spinulosus* (Strohmeyer), *Thamurgus granulicollis* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Xyleborus ferrugineus* (Fabricius).

Strombosia spp.

Ambrosiodmus obliquus (LeConte), *Premnobius nodulosus* Hagedorn, *Tiarophorus decellei* Browne, *Triozastus caliginosus* Roberts.

Strombosia tetrandra

Cylindropalpus interpositus (Schedl), *Doliopygus citri* Schedl.

Strombosia spp.

Ambrosiodmus tropicus (Hagedorn).

Strombosia tetrandra

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Dendrochilus strombosiosis* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebrunii* Schedl, *Doliopygus leferrei* Schedl, *Doliopygus megatoma* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus strombosiosis* Schedl, *Doliopygus spectabilis* Schedl, *Eucallacea xanthopus* (Eichhoff), *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus intermedius* (Schedl), *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus piceus* (Motschulsky), *Xyleborus strombosiosis* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Struthanthus deppeanus

Chaetophloeus struthanthi Wood.

Struthanthus spp.

Cnesinus elegans Blandford, *Cryptocarenum spatulatus* Wood.

Struthanthus venetus

Chaetophloeus struthanthi Wood, *Micracisella monadisi* Wood, *Pseudothyranos vallatus* Wood.

Strychnos decussata

Scolytogenes grobleri (Schedl).

Strychnos nux-tomica

Eucallacea andamanensis (Blandford).

Strychnos spp.

Premnobius catipennis Eichhoff.

Stuartia monodelpha

Platypus calamus Blandford, *Xyleborus atratus* Eichhoff.

Styrax benzoin

Coccotrypes variabilis (Beeson), *Coptodryas recidens* (Sampson), *Crossotarsus fractus* Sampson, *Crossotarsus schedli* Browne, *Crossotarsus wallacei* (Thomson),

Hypothenemus glabripennis (Hopkins), *Hypothenemus styx* (Schedl), *Platypus cupulatus* Chapuis, *Platypus pseudocupulatus* Schedl, *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus manicus* (Blandford).

Styrax formosanus

Platypus flectus Nisima & Murayama, *Platypus formosanus* Nisima & Murayama.

Styrax japonicus

Crossotarsus niponicus Blandford, *Platypus calanus* Blandford, *Trypodendron pulchellus* (Murayama), *Xyleborus aquilus* Blandford, *Xyleborus atratus* Eichhoff, *Xylosandrus germanus* (Blandford), *Xylosandrus mutilatus* (Blandford).

Styrax obassia

Xylosandrus borealis Nobuchi, *Xylosandrus brevis* (Eichhoff).

Sussonia spp.

Lanurgus podocarpi Schedl.

Scietenia macrophylla

Anasa versicolor (Sampson), *Arixyleborus suturalis* (Eggers), *Cnестus aterrimus* (Eggers), *Cnестus bicornoides* (Schedl), *Coccotrypes cyperi* (Beeson), *Coptodryas nuxax* (Schedl), *Crossotarsus externidentatus* (Fairmaire), *Crossotarsus fractus* Sampson, *Crossotarsus minax* (Walker), *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Eccoptopterus limbatus* Sampson, *Eccoptopterus sagittarius* Schedl, *Eccoptopterus spinosus* (Olivier), *Hadrodemius comans* (Sampson), *Hadrodemius globus* (Blandford), *Hypothenemus birmanus* (Eichhoff), *Hypothenemus eruditus* Westwood, *Hypothenemus glabripennis* (Hopkins), *Platypus carus* Strohmeier, *Platypus gerstaeckeri* Chapuis, *Platypus platypoides* (Browne), *Xyleborus approximatus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus haberkornii* Eggers, *Xyleborus jatanus* Eggers, *Xyleborus quadrispinosulus* Eggers, *Xyleborus volcubus* (Fabricius), *Xylosandrus ater* (Eggers), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus gravidus* (Blandford), *Xylosandrus manicus* (Blandford), *Xylosandrus morigerus* (Blandford), *Xylosandrus mutilatus* (Blandford), *Xylosandrus terminatus* (Eggers).

Scietenia mahagoni

Arixyleborus malayensis (Schedl), *Cnестus suturalis* (Eggers), *Eucallacea quadratocollis* (Eggers), *Scolytotplatypus cutomoides* Blandford, *Xyleborus affinis* Eichhoff, *Xyleborus haberkornii* Eggers, *Xyleborus metacuculus* Eggers, *Xylosandrus compactus* (Eichhoff), *Xylosandrus discolor* (Blandford), *Xylosandrus gravidus* (Blandford), *Xylosandrus manicus* (Blandford), *Xylosandrus morigerus* (Blandford).

Scietenia spp.

Cnестus magnus Sampson, *Coccotrypes variabilis* (Beeson), *Corthylus minutissimus* Schedl, *Hypothenemus dolosus* Wood, *Scolytodes guyanaensis* (Schedl), *Scolytodes spadix* (Blackman), *Scolytodes scieteniae* (Blackman), *Theoborus ricini* (Eggers), *Xylosandrus compactus* (Eichhoff).

Scintonia floribunda

Coccotrypes cyperi (Beeson), *Coccotrypes variabilis* (Beeson), *Coccotrypes vulgaris* (Eggers), *Coptodryas perparca* (Sampson), *Platypus cupulatus* Chapuis, *Platypus curtus* Chapuis, *Platypus ovatus* Strohmeier, *Platypus solidus* Walker, *Xyleborus fallax* Eichhoff.

Scintonia specifera

Eccoptopterus limbatus Sampson, *Platypus ovatus* Strohmeier, *Xylosandrus crassiusculus* (Motschulsky).

Syagrus ueddeleum

Coccotrypes dactyliperda (Fabricius).

Syderoxylon spp.

Mitosoma odonticeps Schedl.

Symingtonia populnea

Diapus spatulifer Browne, *Eucallacea andamanensis* (Blandford).

Symphonia clusoides

Polygraphus amoenus (Schaufuss).

Symphonia globulifera

Dactylipalpus grouvellei (Blandford), *Eucallacea xanthopus* (Eichhoff), *Phrixosoma minor* Wood, *Phrixosoma obesa* Blackman, *Polygraphus majusculus* Schedl.

Symphonia spp.

Diapus malgassicus Schedl, *Mitosoma lobatum* Schedl, *Mitosoma obliquatum* Schedl, *Mitosoma obtusum* Schedl, *Mitosoma paulianum* Schedl, *Platypus madagascariensis* Chapuis, *Xyleborus sakalava* Schedl.

Symphonia urophylla

Polygraphus creber Schedl.

Symplocos loha

Ambrosiodinus asperatus (Blandford), *Xylosandrus arcuatus* (Sampson).

Symplocos myrtacea

Platypus calanus Blandford.

Symplocos prunifolia

Xyleborus volcubus (Fabricius).

Symplocos spp.

Ambrosiodinus apicalis (Blandford), *Diamerus curvifer* (Walker), *Leptoxyleborus punctatissimus* (Eichhoff), *Platypus signatus* Chapuis, *Xyleborus cancellatus* Eggers, *Xyleborus emarginatus* Eichhoff, *Xyleborus perforans* (Wollaston), *Xyleborus piceus* (Motschulsky).

Symplocos theaeolia

Crossotarsus machili Beeson, *Cyclorhipidion hirtum* (Hagedorn), *Diapus spatulifer* Browne, *Eucallacea andamanensis* (Blandford), *Eucallacea wallacei* (Blandford), *Indocryphalus intermedius* (Sampson), *Platypus darjeelingensis* Schedl, *Scolytotplatypus darjeelingi* Stebbing, *Scolytotplatypus raja* Blandford, *Xyleborus eggersi* Beeson, *Xyleborus graveleyi* Wichmann, *Xyleborus hagedornianus* Schedl, *Xyleborus lineatus* Eggers.

Synsarpia spp.

Xylosandrus compactus (Eichhoff).

Synsepalum dulcificum

Doliopygus medius Schedl, *Perionomatus longicollis* Chapuis, *Xyleborus affinis* Eichhoff.

Synsepalum longecuneatum

Cylindropalpus interpositus (Schedl), *Cylindropalpus pumilio* (Schedl), *Doliopygus minor* Schedl, *Hypothenemus concolor* Hagedorn, *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss.

Synsepalum spp.

Ambrosiodinus tropicus (Hagedorn), *Cylindropalpus granulatus* (Schedl), *Premnobius caripennis* Eichhoff.

Synsepalum subcordatum

Chaetastus tuberculatus (Chapuis), *Cylindropalpus pertinax* (Schedl), *Cylindropalpus pusillus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus leferrei* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeier), *Doliopygus subditus* (Schedl), *Doliopygus tenuis*

Tectona spp.

Hypothenemus cruditus Westwood, *Platypus bimucus* Blandford, *Platypus carus* Strohmeier, *Platypus jansoni* Chapuis.

Tenstromia spp.

Diapus bilunatus Schedl.

Tephrosia candida

Eucallacea fornicatus (Eichhoff), *Xylosandrus discolor* (Blandford).

Tephrosia maxima

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Tephrosia spp.

Hypothenemus cruditus Westwood, *Xyleborus principalis* Eichhoff, *Xylosandrus morigerus* (Blandford).

Tephrosia vogelii

Cyrtogenius milletiae (Schedl), *Eucallacea fornicatus* (Eichhoff), *Eucallacea xanthopus* (Eichhoff), *Xyleborus alluaudi* Schaufuss, *Xyleborus piccus* (Motschulsky), *Xyleborus principalis* Eichhoff, *Xylosandrus morigerus* (Blandford).

Terminalia amazonia

Xyleborus affinis Eichhoff, *Xyleborus volvulus* (Fabricius).

Terminalia arjuna

Crossotarsus minax (Walker), *Sphaerotrypes globulus* Blandford.

Terminalia bellerica

Crossotarsus bonvouloiri Chapuis, *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus minax* (Walker), *Diapus quinquespinatus* Chapuis, *Eccoptyterus spinosus* (Olivier), *Eucallacea andamanensis* (Blandford), *Eucallacea bicolor* (Blandford), *Eucallacea interjectus* (Blandford), *Platypus cupulatus* Chapuis, *Platypus latifinis* Walker, *Platypus pseudocupulatus* Schedl, *Platypus secretus* Sampson, *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Terminalia bialata

Arixyleborus medius (Eggers), *Coccotrypes salakensis* (Schedl), *Coptodryas redicens* (Sampson), *Diapus quinquespinatus* Chapuis, *Eucallacea bicolor* (Blandford), *Leptoxyleborus concisus* (Blandford), *Xyleborinus andrewesi* (Blandford), *Xyleborinus exiguis* (Walker), *Xyleborus bidentatus* (Motschulsky), *Xyleborus perforans* (Wollaston), *Xyleborus pfeili* (Ratzeburg), *Xyleborus similis* Ferrari.

Terminalia brassii

Diapus pusillimus Chapuis, *Platypus jansoni* Chapuis, *Platypus shoreanus* (Beeson).

Terminalia catappa

Eucallacea fornicatus (Eichhoff), *Platypus carus* Strohmeier, *Platypus cupulatus* Chapuis.

Terminalia chebula

Leptoxyleborus sordicauda (Motschulsky), *Xyleborus perforans* (Wollaston).

Terminalia citrina

Xyleborus perforans (Wollaston).

Terminalia complanata

Crossotarsus minusculus Chapuis, *Platypus dibrachiatatus* Roberts, *Platypus fenestrallatus* Roberts, *Platypus gabensis* Schedl, *Platypus paralleliventris* Roberts.

Terminalia conferta

Xyleborus piccus (Motschulsky).

Terminalia edulis

Coptoborus terminaliae Hopkins, *Cyclorhupidion terminaliae* Hopkins, *Xyleborus affinis* Eichhoff.

Terminalia guianensis

Scolytopsis orinocanus Wood.

Terminalia icorensis

Acanthotomicus subinimunitus (Schedl), *Chaetastus tuberculatus* (Chapuis), *Doliopygus brevis* (Strohmeier), *Doliopygus chapuisi* (Duvivier), *Doliopygus dubius* (Sampson), *Doliopygus serratus* (Strohmeier), *Periommatatus excisus* Strohmeier, *Periommatatus substriatus* Strohmeier, *Platypus hintzi* Schaufuss, *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus conradti* Hagedorn, *Xyleborus eichhoffianus* Schedl, *Xyleborus piccus* (Motschulsky), *Xyleborus rothkirchi* Eggers, *Xyleborus subtuberculatus* Eggers, *Xyleborus volvulus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Terminalia mani

Diapus quinquespinatus Chapuis, *Platypus verchinatus* (Beeson), *Xyleborus perforans* (Wollaston).

Terminalia mantaly

Ambrosiodmus mahafali (Schedl).

Terminalia myriocarpa

Ambrosiodmus lewisi (Blandford), *Ambrosiodmus minor* (Stebbing), *Ambrosiodmus rubricollis* (Eichhoff), *Cnestus turalis* (Eggers), *Coccotrypes cyperi* (Beeson), *Coccotrypes papuanus* (Eggers), *Coccotrypes vulgaris* (Eggers), *Eucallacea interjectus* (Blandford), *Eucallacea velatus* (Sampson), *Hadrodemus metacomans* (Eggers), *Hadrodemus pseudocomans* (Eggers), *Platypus errans* (Sampson), *Xyleborinus subgranulatus* (Eggers), *Xyleborus haberkorni* Eggers, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus subsimilis* (Eggers).

Terminalia paniculata

Crossotarsus minax (Walker), *Platypus latifinis* Walker, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.

Terminalia procera

Eucallacea bicolor (Blandford), *Leptoxyleborus concisus* (Blandford), *Xyleborinus andrewesi* (Blandford), *Xyleborus perforans* (Wollaston), *Xylosandrus discolor* (Blandford).

Terminalia sepicana

Platypus solidus Walker.

Terminalia spp.

Ambrosiodmus hagedorni (Iglesias), *Ambrosiodmus lecontei* Hopkins, *Ambrosiodmus tropicus* (Hagedorn), *Amphicranus micans* Wood, *Crossotarsus andamanus* Beeson, *Cryphalus terminaliae* Browne, *Cyrtogenius prepartus* Browne, *Diapus turgidus* Roberts, *Eidophilus borneensis* Browne, *Eucallacea interjectus* (Blandford), *Margadillius fulvus* Browne, *Margadillius terminaliae* Browne, *Mitrosoma excisum* Schaufuss, *Platypus madagascariensis* Chapuis, *Platypus parvus* Browne, *Platypus uncinatus* Blandford, *Premnobius cavipennis* Eichhoff, *Taurodemus sharpi* (Blandford), *Theoborus ricini* (Eggers), *Xyleborinus gracilis* (Eichhoff), *Xyleborus cognatus* Blandford, *Xyleborus glabratus* Browne, *Xyleborus pumilus* Eggers, *Xyleborus shoreae* (Stebbing), *Xyleborus spinulosus* Blandford, *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford).

Terminalia superba

Chaetastus tuberculatus (Chapuis), *Cylindropalpus*

auricomans (Schaufuss), *Cylindropalpus camerunus* (Schedl), *Diapys quinquispinatus* Chapuis, *Doliopygus brevis* (Strohmeyer), *Doliopygus chapuisi* (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus divaricus* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus medius* Schedl, *Doliopygus minimus* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus punctiventris* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Dryocoetoides cristatus* (Fabricius), *Eccoptypterus spinosus* (Olivier), *Hypothenemus concolor* Hagedorn, *Periommatius angustior* Schedl, *Periommatius excisus* Strohmeyer, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus parallelus* (Fabricius), *Platypus picinus* Schedl, *Premnobius quadrispinosus* Schedl, *Platypus rufescens* (Strohmeyer), *Platypus solutus* Schedl, *Platypus spinulosus* (Strohmeyer), *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufussi* (Strohmeyer), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus scolinatus* Hagedorn, *Xyleborus subasperulus* Eggers, *Xyleborus subtuberculatus* Eggers, *Xyleborinus sharpae* (Hopkins), *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Terminalia tomentosa

Coccotrypes longior (Eggers), *Coccotrypes mibilis* (Blandford), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus minax* (Walker), *Diamerus nigrisctosus* Eggers, *Euwallacea andamanensis* (Blandford), *Sphaerotrypes globulus* Blandford, *Platypus forficula* Chapuis, *Platypus solidus* Walker, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus butanali* (Beeson), *Xylosandrus crassiusculus* (Motschulsky).

Ternstroemia gymnanthera

Xyleborus atratus Eichhoff.

Ternstroemia japonica

Platypus calamus Blandford.

Tessmannia africana

Doliopygus chapuisi (Duvivier), *Doliopygus conradti* (Strohmeyer), *Doliopygus ghesquieri* (Schedl), *Doliopygus lebruni* Schedl, *Doliopygus spectabilis* Schedl, *Xylosandrus crassiusculus* (Motschulsky).

Tessmannia anomala

Doliopygus conradti (Strohmeyer), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).

Tessmannia claessensii

Doliopygus stanceri Schedl.

Tessmannia spp.

Premnobius caripennis Eichhoff.

Tetradenia konishii

Crossotarsus rengetensis Niisima & Murayama.

Tetrameles nudiflora

Euwallacea andamanensis (Blandford), *Euwallacea interjectus* (Blandford), *Platypus cavius* Strohmeyer, *Platypus curtatus* Sampson, *Platypus indicus* Strohmeyer, *Platypus solidus* Walker, *Xyleborus similis* Ferrari.

Tetraplasandra spp.

Xyleborus ferrugineus (Fabricius), *Xyleborus hawaiiensis* Perkins.

Tetrapleura tetraptera

Chaetastus tuberculatus (Chapuis), *Cylindropalpus auricomans* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus erichsoni* (Chapuis), *Doliopygus exilis* (Chapuis), *Doliopygus perbrevis* Schedl, *Doliopygus rapax* (Sampson), *Doliopygus serratus* (Strohmeyer), *Doliopygus subditivus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Doliopygus unicomis* Schedl, *Hypothenemus biseriatus* (Eggers), *Platypus hintzi* Schaufuss, *Platypus parallelus* (Fabricius), *Theoborus ricini* (Eggers), *Trachyostus schoutedeni* (Schedl), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Tetrorchidium didymotemon

Platypus spinulosus (Strohmeyer).

Teucrium pseudoscorodonia

Thamnurgus delphinii (Rosenhaner).

Teucrium scorodonia

Thamnurgus kaltenbachii (Bach).

Thalia geniculata

Xyleborus ferrugineus (Fabricius), *Xyleborus ficus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).

Thea japonica

Sphaerotrypes pila Blandford.

Thea sinensis

Platypus biuncus Blandford, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus morigerus* (Blandford).

Theobroma cacao

Ambrosiodmus hagedorni (Iglesias), *Amphicranus argutus* Wood, *Amphicranus elegans* Eichhoff, *Amphicranus fulgidus* Wood, *Amphicranus spectabilis* (Wood), *Amphicranus spectus* Wood, *Amphicranus theobroma* Sampson, *Araptus columbianus* (Schedl), *Coccotrypes carpophagus* (Hornung), *Coccotrypes cyperi* (Beeson), *Coptoborus pseudotenuis* (Schedl), *Coptoborus tolimanus* (Eggers), *Coptoborus vespatorius* (Schedl), *Coptodryas nugax* (Schedl), *Corthyllocorus tuberculifer* (Eggers), *Corthylus collaris* Blandford, *Corthylus concisus* Wood, *Corthylus papulans* Eichhoff, *Corthylus pygmaeus* Wood, *Corthylus sobrinus* Wood, *Corthylus theobromae* Nunberg, *Crossotarsus minax* (Walker), *Crossotarsus cafer* Schedl, *Cryphalus theobromae* Nunberg, *Cyclorhipidion popondettae* (Browne), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus gracilior* Schedl, *Doliopygus kakaensis* Schedl, *Doliopygus lefevrei* Schedl, *Dryocoetoides cristatus* (Fabricius), *Dryocoetoides flavus* (Fabricius), *Euwallacea destruens* (Blandford), *Euwallacea fornicatus* (Eichhoff), *Euwallacea interjectus* (Blandford), *Hypothenemus ater* (Eggers), *Hypothenemus atratus* (Schedl), *Hypothenemus concolor* Hagedorn, *Hypothenemus criticus* (Schedl), *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus grandis* Schedl, *Hypothenemus intricatus* (Schedl), *Hypothenemus liberiensis* (Hopkins), *Hypothenemus morio* (Eggers), *Hypothenemus obscurus* (Fabricius), *Hypothenemus perhispidus* (Eggers), *Hypothenemus pupipennis* (Eggers), *Hypothenemus seriatus* (Eichhoff), *Hypothenemus setosus* (Eichhoff), *Hypothenemus solitarius* (Schedl), *Hypothenemus suspectus* Wood, *Hypothenemus teretis* Wood, *Microrhipalus attenuatus* (Eggers), *Monarthrum corculum* Wood, *Monarthrum fimbriaticorne* (Blandford), *Platypus deyrrollei* Chapuis, *Platy-*

- pus hintzi* Schaufuss, *Platypus jansoni* Chapuis, *Platypus parallelus* (Fabricius), *Pseudothysanoes atomus* Wood, *Sampsonius dampfi* Schedl, *Scolytogenes samonius* (Browne), *Taurodenus ebenus* (Wood), *Taurodenus pandulus* (Wood), *Taurodenus percheae* (Ferrari), *Taurodenus sharpi sharpi* (Blandford), *Taurodenus splendens* (Schaufuss), *Taurodenus varians* (Fabricius), *Theoborus coartatus* (Sampson), *Theoborus ricini* (Eggers), *Theoborus theobromae* Hopkins, *Theoborus villosulus* (Blandford), *Tricolus saundersi* Wood, *Xyleborinus andreevsi* (Blandford), *Xyleborinus bicornatulus* (Wood), *Xyleborinus gracilis* (Eichhoff), *Xyleborinus intersetosus* (Blandford), *Xyleborinus protinus* (Wood), *Xyleborinus reconditus* (Schedl), *Xyleborinus subulcatus* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus asper* Eggers, *Xyleborus barinbuensis* Eggers, *Xyleborus caribicus* Eggers, *Xyleborus commixtus* Blandford, *Xyleborus ferox* Blandford, *Xyleborus ferrugineus* (Fabricius), *Xyleborus ficus* Eggers, *Xyleborus haberkorni* Eggers, *Xyleborus horridatus* Wood, *Xyleborus lacunatus* Wood, *Xyleborus macer* Blandford, *Xyleborus multispinatus* Eggers, *Xyleborus parallelocollis* Eggers, *Xyleborus parallelus* Eggers, *Xyleborus perforans* (Wollaston), *Xyleborus posticus* Eichhoff, *Xyleborus quadrispinosulus* Eggers, *Xyleborus squamulatus* Eichhoff, *Xyleborus subasperulus* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky), *Xylosandrus discolor* (Blandford), *Xylosandrus laticeps* (Wood), *Xylosandrus manicus* (Blandford), *Xylosandrus morigerus* (Blandford), *Xylosandrus ursinus* (Hagedorn).
- Theobroma silvestre**
Hypocryphalus obscurus Hopkins.
- Theobroma spp.**
Xyleborus similis Ferrari, *Xylosandrus compactus* (Eichhoff).
- Thespesia populnea**
Erucoladus corpulentus (Sampson), *Eucallacca andamanensis* (Blandford), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff).
- Theretia nereifolia**
Hypothenemus eruditus Westwood.
- Theretia orata**
Liparthrum theretiae Wood.
- Theretia spp.**
Liparthrum cracentis Wood, *Pityophthorus costabilis* Wood, *Pityophthorus costatulus* Wood.
- Thrinax argentea**
Coccotrypes carpophagus (Hornung).
- Thujopsis dolabrata**
Phloeosinus gifuensis Murayama, *Phloeosinus lewisi* Chapuis, *Phloeosinus rudis* Blandford, *Phloeosinus sanniohensis* Murayama, *Xyleborinus saxseni* (Ratzeburg).
- Thujopsis spp.**
Phloeosinus perlatus Chapuis.
- Thunbergia grandiflora**
Emcallacca velatus (Sampson).
- Thuja japonica**
Phloeosinus rudis Blandford.
- Thuja occidentalis**
Hypothenemus eruditus Westwood, *Phloeosinus canadensis* Swaine, *Phloeosinus dentatus* (Say), *Phloeosinus punctatus* LeConte.
- Thuja orientalis**
Phloeosinus hopchi Schedl.
- Thuja plicata**
Phloeosinus cupressi Hopkins, *Phloeosinus punctatus* LeConte, *Phloeosinus sequoiae* Hopkins.
- Thuja spp.**
Phloeosinus bicolor (Brulle), *Phloeosinus cristatus* (LeConte), *Phloeosinus henschi* Reitter, *Phloeosinus thujae* (Perris).
- Thuja standishii**
Xyleborus seriatus Blandford.
- Tieghmella hecklii**
Doliopygus bidiscoplanus Roberts.
- Tilia americana**
Pseudothysanoes rigidus (LeConte).
- Tilia amurensis**
Ernoporus tiliae (Panzer), *Eucallacca validus* (Eichhoff), *Scolytus koltzei* Reitter, *Xyleborinus alni* (Niisima).
- Tilia cienserrata**
Pseudothysanoes multispinatus Wood.
- Tilia cordata**
Ernoporus tiliae (Panzer), *Pseudoxylechinus tiliae* (Niisima).
- Tilia cordata japonica**
Crossotarsus niponicus Blandford, *Platypus severini* Blandford, *Hypothenemus sapporocensis* (Niisima), *Xyleborus seriatus* Blandford.
- Tilia intermedia**
Ernoporus caucasicus (Lindemann), *Ernoporus tiliae* (Panzer), *Hylastinus tiliae* Semenov.
- Tilia mandshurica**
Scolytus koltzei Reitter.
- Tilia parvifolia**
Ernoporus caucasicus (Lindemann), *Ernoporus tiliae* (Panzer), *Hylastinus tiliae* Semenov.
- Tilia rubra**
Ernoporus caucasicus (Lindemann).
- Tilia spp.**
Hylesinus crenatus (Fabricius), *Monarthrum nuli* (Fitch), *Scolytus laevis* Chapuis, *Trypodendron signatum* (Fabricius), *Xyleborinus saxseni* (Ratzeburg), *Xyleborus sayi* (Hopkins).
- Tilia tomentosa**
Ernoporus caucasicus (Lindemann).
- Tiliacora acuminata**
Miniocurus beasoni Wood.
- Tococa formicaria**
Xyleborus parcellus Wood.
- Toddalia asiatica**
Eucallacca xanthopus (Eichhoff), *Xyleborus principalis* Eichhoff.
- Toonia calantas**
Xyleborus haddeni Schedl.
- Toonia sinensis**
Leptoxyleborus concisus (Blandford), *Xyleborus haberkorni* Eggers, *Xyleborus piccus* (Motschulsky).
- Toonia spp.**
Xylosandrus compactus (Eichhoff).
- Toonia sureni**
Xylosandrus manicus (Blandford).
- Torrubia longifolia**
Dendrosinus bourreriae Schwarz, *Hypothenemus squamosus* (Hopkins).
- Toulcia pulcinata**
Xyleborus affinis Eichhoff, *Xyleborus asper* Eggers.
- Toromita guianensis**
Hypothenemus opacus (Eichhoff).

Toxicodendron spp.

Hypothenemus erectus LeConte, *Hypothenemus indicus* Wood, *Pityophthorus cornutus* Wood, *Pityophthorus detentus* Wood.

Toxicodendron trichocarpum

Eidophelus imitans Eichhoff.

Toxoxylon pomiferum

Cnestus strigicollis LeConte.

Trachelospermum asiaticum

Scolytogenes scolytomimoides (Nobuchi).

Trachylobium narrucosum

Hypothenemus seriatus (Eichhoff).

Trachylobium verrucosum

Triozastus banghaasi (Schaufuss), *Triozastus marshalli* (Sampson).

Treculia africana

Chaetastus tuberculatus (Chapuis), *Doliopygus coclocephalus* (Schaufuss), *Doliopygus serratus* (Strohmeyer), *Hylesinopsis dubius* Eggers, *Doliopygus erichsoni* (Chapuis), *Pityophthorus treculiae* Schedl, *Trachyostus schaufussi* (Strohmeyer), *Trachyostus schoutedeni* (Schedl), *Xyleborus alluaudi* Schaufuss.

Trema floridana

Hypothenemus birmanus (Eichhoff), *Hypothenemus brunneus* (Hopkins), *Hypothenemus eruditus* Westwood, *Hypothenemus javanus* (Eggers), *Hypothenemus seriatus* (Eichhoff).

Trema guineense

Ambrosiodmus aegir (Eggers), *Cyclorhipidion crucipenne* (Schedl), *Doliopygus perbrevis* Schedl, *Hypothenemus biseriatus* (Eggers), *Hypothenemus pubipennis* (Eggers), *Platypus hintzi* Schaufuss, *Platypus spinulosus* (Strohmeyer), *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus consobrinus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus multispinatus* Eggers.

Trema orientalis

Ambrosiodmus artemgranulatus (Schedl), *Ambrosiodmus mahafali* (Schedl), *Doliopygus erichsoni* (Chapuis), *Eucallacea xanthopus* (Eichhoff), *Pityophthorus kivuensis* Schedl, *Platypus jansoni* Chapuis, *Xyleborinus quadrispinus* (Schedl), *Xyleborinus spiculatus* (Schaufuss), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus antaisaka* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus principalis* Eichhoff, *Xyleborus subtuberculatus* Eggers, *Xylosandrus morigerus* (Blandford).

Tricalysia viridiflora

Xyleborus kunamotoensis Murayama.

Trichilia arborea

Hypothenemus seriatus (Eichhoff), *Micracis swainiei* Blackman.

Trichilia gilgiana

Coccotrypes cylindricus (Eggers), *Doliopygus dubius* (Sampson), *Doliopygus serratus* (Strohmeyer), *Platypus parallelus* (Fabricius), *Triozastus marshalli* (Sampson).

Trichilia heudelotii

Cylindropalpus auricomans (Schaufuss), *Doliopygus brevis* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus erichsoni* (Chapuis), *Doliopygus interjectus* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus serratus* (Strohmeyer), *Periommatus grandis* Schedl, *Periommatus longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus orientalis* (Strohmeyer), *Trachyostus schaufussi* (Strohmeyer), *Triozastus banghaasi* (Schaufuss), *Triozastus marshalli* (Sampson),

Xyleborinus collarti (Eggers), *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Trichilia lanata

Cylindropalpus pertinax (Schedl), *Doliopygus coclocephalus* (Schaufuss), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus interjectus* Schedl, *Platypus hintzi* Schaufuss, *Premnobius quadrispinosus* Schedl, *Scolytoplatypus eichelbanni* Hagedorn, *Triozastus banghaasi* (Schaufuss), *Xyleborus affinis* Eichhoff.

Trichilia prieureana

Doliopygus conradti (Strohmeyer), *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus subditivus* (Schedl), *Hypothenemus eruditus* Westwood, *Triozastus elongatus* Schedl, *Xyleborinus sharpae* (Hopkins), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Trichilia propinqua

Cnemidophorus testitus (Eggers), *Xyleborus affinis* Eichhoff.

Trichilia rubescens

Chaetastus tuberculatus (Chapuis), *Doliopygus citri* Schedl, *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus lefevrei* Schedl, *Doliopygus serratus* (Strohmeyer), *Doliopygus trichiliae* Schedl, *Triozastus banghaasi* (Schaufuss), *Triozastus marshalli* (Sampson), *Xyleborinus sharpae* (Hopkins).

Trichilia spp.

Dactylipalpus camerunus Hagedorn, *Doliopygus alternans* (Strohmeyer), *Doliopygus conradti* (Strohmeyer), *Doliopygus interjectus* Schedl, *Doliopygus obduens* Roberts, *Doliopygus serratus* (Strohmeyer), *Mitosoma obliquatum* Schedl, *Mitosoma odonticeps* Schedl, *Mitosoma truncatepennis* Schedl, *Platypus parallelus* (Fabricius), *Premnobius cavipennis* Eichhoff, *Scolytoplatypus occidentalis* Brownie, *Xyleborinus diversus* (Schedl), *Xyleborinus gracilipennis* (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus clerodendronae* Schedl, *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius).

Trichilia velutischi

Trachyostus schaufussi (Strohmeyer).

Trichoscypha arborea

Cyclorhipidion daosi (Schedl).

Trichoscypha spp.

Doliopygus chapuisi (Duvivier), *Doliopygus serratus* (Strohmeyer), *Platypus parallelus* (Fabricius).

Tricocodendron acuminatum

Mitosoma excisum Schaufuss, *Mitosoma paulianum* Schedl, *Xyleborinus spiculatus* (Schaufuss).

Tridesmostemon claussensii

Chaetastus persimilis (Schedl), *Cylindropalpus granulatus* (Schedl), *Cylindropalpus pertinax* (Schedl), *Doliopygus dubius* (Sampson), *Doliopygus semipilosus* (Schedl), *Doliopygus tenuis* (Strohmeyer), *Hypothenemus criticus* (Schedl), *Hypothenemus eruditus* Westwood, *Mesoplatypus crinaceus* Schedl, *Platypus impressus* (Strohmeyer), *Premnobius quadrispinosus* Schedl, *Triozastus marshalli* (Sampson), *Xyleborus affinis* Eichhoff, *Xyleborus volutus* (Fabricius).

Tridesmostemon spp.

Premnobius cavipennis Eichhoff.

Trifolium pratense

Hylastinus obscurus (Marshall).

Triopteris jamaicensis

Hypothecinus eruditus Westwood.

Triplochiton scleroxylon

Ambrosiodmus neglectus (Schedl), *Chaetastus tuberculatus* (Chapuis), *Ctenoxylon flavescens* Hagedorn, *Cylindropalpus auricomans* (Schaufuss), *Cylindropalpus canerianus* (Schedl), *Cylindropalpus pumilio* (Schedl), *Cyrtogenius cribripennis* Schedl, *Diamerus imperfectus* Eggers, *Doliopygus chapuisi* (Duviérier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus interjectus* Schedl, *Doliopygus medius* Schedl, *Doliopygus notatus* Schedl, *Doliopygus opifex* (Sampson), *Doliopygus serratus* (Strohmeyer), *Dryococtoides cristatus* (Fabricius), *Hypothecinus eruditus* Westwood, *Hypothecinus pubipennis* (Eggers), *Leptoxyleborus scabrior* (Schedl), *Periommatius excisus* Strohmeyer, *Periommatius grandis* Schedl, *Periommatius longicollis* Chapuis, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus parallelus* (Fabricius), *Polygraphus coronatus* Eggers, *Scolytogenes braderi* (Browne), *Trachyostus ghanaensis* Schedl, *Triozastus banghaasi* (Schaufuss), *Xyleborinus diversus* (Schedl), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiusculus* Eggers, *Xyleborus eichhoffianus* Schedl, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xyleborus pseudoambasius* Schedl, *Xyleborus subtuberculatus* Eggers, *Xyleborus volutus* (Fabricius), *Xylosandrus crassiusculus* (Motschulsky).

Triplochiton spp.

Prennobiis catipennis Eichhoff.

Tristania spp.

Anasa opalescens (Schedl), *Xylosandrus compactus* (Eichhoff).

Tristania ichiteana

Xyleborus haberkornii Eggers, *Xylosandrus mancus* (Blandford).

Triumfetta spp.

Ambrosiodmus ovatus (Eggers), *Hypothecinus grandis* Schedl.

Trochis racemosa

Xyleborus naeae Blandford.

Tsuga chinensis

Sphaerotrypes magnus Tsai & Yin, *Sphaerotrypes tsugae* Tsai & Yin.

Tsuga diversifolia

Dryocoetes brevipilosus Murayama.

Tsuga heterophylla

Dendroctonus pseudotsugae Hopkins, *Dryocoetes autographus* (Ratzeburg), *Gnathotrichus retusus* (LeConte), *Gnathotrichus sulcatus* (LeConte), *Hylobius reticulatus* Wood, *Phloeosinus punctatus* LeConte, *Pityophthorus pseudotsugae* Swaine, *Platypus wilsoni* Swaine, *Pseudohylesinus granulatus* (LeConte), *Pseudohylesinus nebulosus* (LeConte), *Pseudohylesinus sericeus* (Mannerheim), *Pseudohylesinus tsugae* Swaine.

Tsuga mertensiana

Pseudohylesinus dispar dispar Blackman, *Pseudohylesinus tsugae* Swaine.

Tsuga sieboldii

Crossotarsus niponicus Blandford, *Euwallacea validus* (Eichhoff).

Tsuga spp.

Cryphalus taiwanus Schedl, *Gnathotrichus materiarius* (Fitch), *Hylastes nigrinus* (Mannerheim), *Scolytus*

tsugae (Swaine), *Xyleborus dispar* (Fabricius), *Xyleborus scriatus* Blandford, *Xyloterinus politus* (Say).

Tsuga yunnanensis

Xyleborinus saxeni (Ratzeburg).

Tulasneana spp.

Euwallacea xanthopus (Eichhoff).

Turpinia latifolia

Euwallacea destrucis (Blandford).

Turpinia nepalensis

Coccotrypes nubilis (Blandford), *Xyleborus eggersi* Beeson.

Turpinia pomifera

Anasa nobilis (Eggers), *Hadrodemus globus* (Blandford), *Xyleborus cancellatus* Eggers, *Xyleborus fallax* Eichhoff, *Xyleborus haberkornii* Eggers.

Turpinia spp.

Euwallacea andamanensis (Blandford).

Turraeanthus africana

Chaetastus tuberculatus (Chapuis), *Coptoborus forficatus* (Schedl), *Cyclorhipidion crucifer* (Hagedorn), *Doliopygus chapuisi* (Duviérier), *Doliopygus conradti* (Strohmeyer), *Doliopygus dubius* (Sampson), *Doliopygus ghesquieri* (Schedl), *Doliopygus interjectus* Schedl, *Doliopygus medius* Schedl, *Doliopygus rhodesianus* (Schedl), *Doliopygus serratus* (Strohmeyer), *Doliopygus wolschei* Schedl, *Dryococtoides cristatus* (Fabricius), *Hypothecinus eruditus* Westwood, *Periommatius excisus* Strohmeyer, *Platypus hintzi* Schaufuss, *Platypus impressus* (Strohmeyer), *Platypus rimulosus* Schedl, *Platypus spinulosus* (Strohmeyer), *Polygraphus granulifer* Eggers, *Prennobiis mukunyeae* (Schedl), *Prennobiis nodulosus* Hagedorn, *Prennobiis quadrispinosus* Schedl, *Xyleborinus collarti* (Eggers), *Xyleborinus sharpae* (Hopkins), *Xyleborinus similans* (Eggers), *Xyleborus affinis* Eichhoff, *Xyleborus africanus* Eggers, *Xyleborus alluaudi* Schaufuss, *Xyleborus ambasiipennis* Schedl, *Xyleborus ambasiusculus* Eggers, *Xyleborus ferrugineus* (Fabricius), *Xyleborus picus* (Motschulsky), *Xyleborus tenellus* Schedl, *Xyleborus turraeanthus* Schedl, *Xylosandrus compactus* (Eichhoff), *Xylosandrus crassiusculus* (Motschulsky).

Turraeanthus spp.

Ambrosiodmus eichhoffi (Schreiner), *Prennobiis catipennis* Eichhoff.

Turraeanthus vignei

Doliopygus nairobiensis (Schedl), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schaufuss, *Xyleborus picus* (Motschulsky).

Uapaca spp.

Mitosoma obliquatum Schedl, *Platypus madagascariensis* Chapuis, *Xyleborinus dentellus* (Schedl).

Ulex europaeus

Hylastinus obscurus (Marsham), *Phloeotribus rhododactylus* (Marsham).

Ulmus americana

Hylocurus rudis (LeConte).

Ulmus campestris

Hylesinus elatus Niisima, *Hyorrhynchus lewisi* Blandford, *Pteleobius kraatzii* (Eichhoff), *Pteleobius vittatus* (Fabricius), *Scolytus chikisanii* Niisima, *Scolytus frontalis* Blandford.

Ulmus carpinifolia

Scolytus chikisanii Niisima, *Scolytus ecksteini* Butovitsch, *Scolytus jaroschewskii* Scheyvrew, *Scolytus centrosus* Scheyvrew, *Scolytus zaitzevi* Butovitsch.

Ulmus effusa

Pteleobius kraatzii (Eichhoff), *Pteleobius vittatus* (Fabricius).

Ulmus japonica

Hylesinus elatus Niisima, *Neopteleobius scutulatus* (Blandford), *Scolytus butovitschii* Stark, *Scolytus chikisanii* Niisima, *Scolytus jacobsoni* (Spessivtsev), *Scolytus pubescens* Stark, *Scolytus semenovi* (Spessivtsev), *Tripodendron signatum* (Fabricius), *Xyleborus atratus* Eichhoff.

Ulmus laciniata

Scolytus chikisanii Niisima, *Scolytus jacobsoni* (Spessivtsev), *Scolytus semenovi* (Spessivtsev).

Ulmus laevis

Scolytus kozikowskii Michalski.

Ulmus lanceifolia

Scolytus chelogaster Schedl, *Scolytus squamosus* Yin & Huang.

Ulmus montana

Enoporicus caucasicus (Lindemann), *Pteleobius kraatzii* (Eichhoff), *Pteleobius vittatus* (Fabricius).

Ulmus propinqua

Scolytus butovitschii Stark, *Scolytus jacobsoni* (Spessivtsev), *Scolytus pubescens* Stark, *Scolytus semenovi* (Spessivtsev), *Scolytus ventrosus* Scheyrev.

Ulmus pumila

Eucallacea validus (Eichhoff), *Hypothenemus modestus* (Murayama), *Pseudothysanoes mongolica* (Sokanovskii), *Scolytus butovitschii* Stark, *Scolytus parviclaviger* Yin & Huang, *Sphaerotrypes ulmi* Tsai & Yin.

Ulmus rubra

Trischidiast atomus (Hopkins).

***Ulmus* spp.**

Ambrosiodinus apicalis (Blandford), *Corthylus columbianus* Hopkins, *Corthylus punctatissimus* (Zimmermann), *Hylocurus langstoni* Blackman, *Hylurgopinus rufipes* (Eichhoff), *Hypothenemus eruditus* Westwood, *Pteleobius kraatzii* (Eichhoff), *Scolytus aratus* Blandford, *Scolytus ensifer* Eichhoff, *Scolytus japonicus* Chapuis, *Scolytus kirschi* Skalitzky, *Scolytus laevis* Chapuis, *Scolytus mali* (Beckstein), *Scolytus multistriatus* (Marsham), *Scolytus pygmaeus* (Fabricius), *Scolytus scheyrevici* Semenov, *Scolytus scolytus* (Fabricius), *Scolytus sulcifrons* Rey, *Scolytus trispinosus* Strohmeier, *Taphrorychus vilifrons* (Dufour), *Thysanoes berchemiae* Blackman, *Xyleborus atratus* Eichhoff, *Xyleborus curygraphus* (Ratzeburg), *Xyleborus maiche* (Stark), *Xyleborus pfeili* (Ratzeburg), *Xylosandrus germanus* (Blandford), *Xyloterinus politus* (Say).

Ulmus wallichiana

Scolytus kashmirensis Schedl.

***Umbellularia* spp.**

Hylocurus hirtellus (LeConte).

Uniola paniculata

Hypothenemus californicus Hopkins.

Urena lobata

Hypothenemus eruditus Westwood.

***Urena* spp.**

Hypothenemus eruditus Westwood, *Hypothenemus seriatus* (Eichhoff).

***Urea* spp.**

Xyleborinus saxeseni (Ratzeburg).

Urophyllum lyallii

Xyleborus albaudi Schaufuss, *Xylosandrus hirsutipennis* (Schedl).

***Urostigma* spp.**

Cryphalus walkeri (Blandford).

Usambara occidentalis

Peruophorus brevicollis (Strohmeier).

Vaccinium consanguineum

Corthycyclon aztecum (Bright), *Corthycyclon tardum* Wood, *Corthylus subserratus* Wood.

***Vaccinium* spp.**

Corthylus punctatissimus (Zimmermann).

Vachellia farnesiana

Hypothenemus brunneus (Hopkins).

Valeriana scandens

Cnesinus adustus Schedl.

Valleris haynii

Scolytogenes indicus Wood.

Vanda coerulea

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

***Vanda* spp.**

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Vanda teres

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Vanda tricolor

Xylosandrus compactus (Eichhoff), *Xylosandrus morigerus* (Blandford).

Vateria copalifera

Sphaerotrypes cristatus Wood, *Xylosandrus manicus* (Blandford).

Vateria indica

Coccotrypes advena Blandford, *Coccotrypes cardamomi* Schaufuss, *Coccotrypes cyperi* (Beeson), *Coccotrypes variabilis* (Beeson), *Coccotrypes vateriae* (Beeson), *Crossotarsus quadricaudatus* (Strohmeier), *Eucallacea andamanensis* (Blandford), *Genyocerus furtivus* (Sampson), *Platypus latifinis* Walker, *Platypus suffodiens* Sampson, *Xylosandrus butanali* (Beeson), *Xylosandrus crassiusculus* (Motschulsky).

Vatica heteroptera

Xylosandrus crassiusculus (Motschulsky).

Vatica lanceaefolia

Ambrosiodinus lewisi (Blandford), *Arixyleborus malayensis* (Schedl), *Carchesiopygus assamensis* (Beeson), *Carchesiopygus uollastoni* (Chapuis), *Coptodryas alpha* (Beeson), *Crossotarsus bontouloiri* Chapuis, *Diapus quinquespinatus* Chapuis, *Eucallacea tristis* (Eggers), *Genyocerus furtivus* (Sampson), *Platypus cactus* Strohmeier, *Platypus curtatus* Sampson, *Platypus decens* Sampson, *Platypus lepidus* Chapuis, *Platypus uncinatus* Blandford, *Sphaerotrypes vaticanus* Beeson, *Xyleborinus subgranulatus* (Eggers), *Xyleborus shoreae* (Stebbing), *Xyleborus similis* Ferrari, *Xylosandrus gravidus* (Blandford).

***Vatica* spp.**

Acacis borneensis Brownie, *Cnestus suturalis* (Eggers), *Coptodryas bella* (Sampson), *Coptodryas confusa* Hopkins, *Coptodryas obtusicollis* (Schedl), *Coptodryas perparva* (Sampson), *Cyclorhipidion punctillicolle* (Schedl), *Cyrtogenius vaticae* (Nunberg), *Genyocerus biporus* (Schedl), *Genyocerus decemspinatus* (Schedl), *Genyocerus exilis* (Schedl), *Genyocerus serratus* (Schedl), *Leptoxyleborus scabrior* (Schedl), *Ozopemon obanus* Hagedorn, *Platypus artemisoides* Schedl, *Platypus granifer* Schedl, *Platypus minacior* Schedl, *Platypus sericans* Schedl.

- Sphaerotrypes biserialis* Schedl, *Sphaerotrypes minutus* Browne, *Webbia bakoensis* Browne, *Webbia seriata* (Browne), *Webbia trepanicauda* (Eggers), *Xyleborus macropterus* Schedl, *Xyleborus muasi* Browne, *Xyleborus pseudopilifer* Schedl.
- Vatica stapfiana**
Platypus cupulatus Chapuis.
- Varzea spp.**
Platypus runcinatus Roberts.
- Vepris stolzii**
Doliopygus lefevrei Schedl, *Periommatus excisus* Stroh-meyer.
- Vepris undulata**
Xyleborinus aemulus (Wollaston).
- Verbena spp.**
Hypothenemus californicus Hopkins.
- Verbesina agricultorum**
Cnesinus amictens Wood, *Hypothenemus erectus* LeConte.
- Verbesina laciniata**
Hypothenemus eruditus Westwood.
- Vernonia amygdalina**
Eucallacea xanthopus (Eichhoff), *Xyleborus alluaudi* Schauffuss, *Xyleborus piccus* (Motschulsky).
- Vernonia appendiculata**
Eucallacea xanthopus (Eichhoff), *Pityoplathorus madagascariensis* Schedl.
- Vernonia arborea**
Platypus carus Stroymeyer, *Xyleborus affinis* Eichhoff, *Xylosandrus crassiusculus* (Motschulsky).
- Vernonia auriculifera**
Thamnurgus africanus Eggers.
- Vernonia conferta**
Doliopygus erichsoni (Chapuis), *Doliopygus lateralis* Schedl, *Doliopygus medius* Schedl, *Doliopygus minor* Schedl, *Doliopygus perbrevis* Schedl, *Doliopygus rhodesianus* (Schedl), *Eucallacea pandae* (Schedl), *Platypus hintzi* Schauffuss, *Platypus orientalis* (Stroh-meyer), *Trachyostus schaufussi* (Strohmeyer), *Xyleborus affinis* Eichhoff, *Xyleborus alluaudi* Schauffuss, *Xyleborus ferrugineus* (Fabricius), *Xyleborus solutus* Schedl, *Xyleborus subtuberculatus* Eggers, *Xylosandrus crassiusculus* (Motschulsky).
- Vernonia lasiopus**
Ambrosiodmus aegir (Eggers).
- Vernonia spp.**
Doliopygus brevis (Strohmeyer), *Doliopygus conradi* (Strohmeyer), *Doliopygus perbrevis* Schedl, *Platypus madagascariensis* Chapuis, *Xyleborinus quadrispinosus* (Eichhoff), *Xyleborus bezanozano* Schedl, *Xyleborus principalis* Eichhoff, *Xylosandrus hirsutipennis* (Schedl).
- Veronica arboreseens**
Xyleborus adusticollis (Motschulsky).
- Vernonia spp.**
Ambrosiodmus tropicus (Hagedorn), *Hypothenemus grandis* Schedl.
- Viburnum betuloides**
Corthylus punctatissimus (Zimmermann).
- Viburnum dilatatum**
Cryphalus riburni Stark.
- Viburnum spp.**
Xylosandrus brevis (Eichhoff).
- Vigna sinensis**
Hypothenemus eruditus Westwood.
- Villebrunnea spp.**
Xylosandrus morigerus (Blandford).
- Vinea spp.**
Xylosandrus compactus (Eichhoff).
- Virecta multiflora**
Hypothenemus grandis Schedl, *Thamnurgus grandicollis* Schedl.
- Virgilia capensis**
Platypus sampsoni (Schedl), *Xyleborinus aemulus* (Wollaston).
- Virgilia oroboides**
Hypocryphalus caplandicus Schedl.
- Virola spp.**
Cnemonyx insignis Wood, *Cnemonyx maculicornis* (Blandford), *Scolytodes declivistriatus* Schedl.
- Virola surinamensis**
Xyleborus grossmanni Schedl.
- Virola icarburgii**
Xyleborus asper Eggers, *Xyleborus sparsipilosus* Eggers.
- Viscum album**
Liparthrum bartschti Muhl.
- Vismia cayennensis**
Cnesinus reticulatus Wood.
- Vismia guianensis**
Cnemonyx panamensis (Blandford), *Hypothenemus trivialis* Wood, *Scolytodes cedrelae* Wood, *Tesseroecerus belti* Sharp, *Xyleborinus interstosus* (Blandford), *Xyleborus vismiae* Wood.
- Vismia spp.**
Amphicranus terebella Blandford, *Cnemonyx minusculus* (Blandford), *Cnemonyx vismiacolens* Wood, *Cnesinus lucaris* Wood, *Corthylus abbreviatus* Eichhoff, *Corthylus additus* Wood, *Hypothenemus erectus* LeConte, *Hypothenemus interstitialis* (Hopkins), *Hypothenemus squamosus* (Hopkins), *Sampsonius dampfi* Schedl, *Taurodenus godmani* (Blandford), *Taurodenus sanguinicollis* (Blandford), *Xyleborus ferrugineus* (Fabricius), *Xyleborus volutus* (Fabricius).
- Vitex altissima**
Eucallacea andamanensis (Blandford), *Xylosandrus crassiusculus* (Motschulsky).
- Vitex congoensis**
Doliopygus lefevrei Schedl, *Doliopygus rapax* (Sampson).
- Vitex euneata**
Pernophorus abhorrens Eggers.
- Vitex hildebrandii**
Doliopygus chapuisi (Duvivier).
- Vitex negundo**
Ambrosiodmus rubricollis (Eichhoff).
- Vitex peduncularis**
Eccoptypterus spinosus (Olivier), *Eucallacea andamanensis* (Blandford), *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari.
- Vitex pubescens**
Cnestus aterrimus (Eggers), *Hypothenemus eruditus* Westwood, *Xyleborus approximatus* Schedl, *Xyleborus haberkorni* Eggers, *Xylosandrus ater* (Eggers), *Xylosandrus manicus* (Blandford), *Xylosandrus pygmaeus* (Eggers).
- Vitex spp.**
Ambrosiodmus triton (Schauffuss), *Crossotarsus muszelschii* Chapuis, *Xyleborus perforans* (Wollaston), *Xyleborus similis* Ferrari, *Xylosandrus compactus* (Eichhoff).
- Vitex wehertschii**
Doliopygus medius Schedl.

Vitis discolor

Xyleborus piccus (Motschulsky).

Vitis spp.

Ambrosiodmus apicalis (Blandford), *Cnesinus elegans* Blandford, *Cryphalus felis* Wood, *Cryptocarenum heveae* (Hagedorn), *Hypothenemus birmanus* (Eichhoff), *Hypothenemus dissimilis* (Zimmermann), *Hypothenemus interstitialis* (Hopkins), *Hypothenemus jarmanus* (Eggers), *Hypothenemus ritis* Browne, *Micracisella nanula* (LeConte), *Premnobius catipennis* Eichhoff, *Xylosandrus compactus* (Eichhoff), *Xylosandrus germanus* (Blandford), *Xylosandrus morigerus* (Blandford).

Vitis vinifera

Hypothenemus eruditus Westwood, *Xylosandrus discolor* (Blandford).

Walsura robusta

Xyleborus perforans (Wollaston).

Warburgia ugandensis

Dolopygus nairobiensis (Schedl).

Washingtonia filifera

Coccotrypes carpophagus (Hornung), *Coccotrypes dactyliperda* (Fabricius).

Washingtonia sonorae

Coccotrypes dactyliperda (Fabricius).

Webbia spp.

Hypothenemus eruditus Westwood.

Weigela hortensis

Xylosandrus brevis (Eichhoff).

Weinmannia spp.

Xyleborinus cupulatus (Schedl), *Xyleborinus forficuloides* (Schedl).

Wendlandia bicuspidata

Liparthrum brincki (Schedl).

Wendlandia tinctoria

Scolytplatypus minimus Hagedorn.

Werklia insignis

Corthylus rubricollis Blandford, *Corthylus serratus* Wood, *Corthylus strigilis* Wood, *Tricolus naevus* Wood.

Whitfordiodendron pubescens

Anasa versicolor (Sampson), *Coptodryas nugax* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus schedli* Browne, *Crossotarsus squamulatus* Chapuis, *Crossotarsus terminatus* Chapuis, *Crossotarsus wallacei* (Thomson), *Eucallacea fornicatus* (Eichhoff), *Eucallacea linatus* (Schedl), *Eucallacea xanthopus* (Eichhoff), *Platypus insulindicus* Schedl.

Widdringtonia juniperoides

Lanurgus widdringtoniae Schedl.

Wikstroemia spp.

Xylosandrus compactus (Eichhoff).

Wistaria floribunda

Hypothenemus kraunhiiae (Murayama), *Scolytogenes badius* (Nobuchi), *Xyleborus kraunhiiae* Niisima.

Wistaria sinensis

Hypothenemus eruditus Westwood.

Wistaria spp.

Chramesus wisteriae Wood, *Hypothenemus erudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff).

Wormia ferruginea

Scolytogenes crenatus (Sampson).

Wormia triquetra

Crossotarsus minax (Walker), *Crossotarsus terminatus* Chapuis.

Wrightia tinctoria

Eucallacea andamanensis (Blandford), *Xylosandrus crassiusculus* (Motschulsky).

Wrightia tomentosa

Platypus errans (Sampson), *Platypus infuscatus* Browne.

Xanthomyrtus spp.

Diaprus nebulosus Roberts, *Platypus bipyramidus* Roberts & Morimoto, *Platypus caelestis* Roberts & Morimoto, *Platypus cicatricosus* Roberts.

Xanthophyllum affine

Crossotarsus terminatus Chapuis, *Xylosandrus crassiusculus* (Motschulsky).

Xanthophyllum curtisii

Eucallacea linatus (Schedl).

Xanthophyllum glaucum

Platypus lepidus Chapuis.

Xanthophyllum ostrandum

Platypus australis Chapuis.

Xanthophyllum papuanum

Baioeis unispinosus Roberts, *Platypus acquilaterus* Roberts, *Platypus daedalus* Roberts, *Platypus evanidus* Roberts, *Platypus hebetatus* Roberts, *Platypus nudiusculus* Roberts, *Platypus pectinatus* Roberts, *Platypus ternus* Roberts, *Platypus turiformis* Roberts & Morimoto, *Platypus vaginerris* Roberts, *Platypus watutensis* Roberts.

Xanthophyllum spp.

Ambrosiodmus sarawakensis (Eggers), *Coptodryas nugax* (Schedl), *Crossotarsus externedentatus* (Fairmaire), *Crossotarsus schedli* Browne, *Cyrtogenius glaber* (Schedl), *Diaprus bispinus* Schedl, *Diaprus nanus* Schedl, *Diaprus perpygmaeus* Roberts, *Diaprus pusillimus* Chapuis, *Eccoptopterus spinosus* (Olivier), *Eucallacea andamanensis* (Blandford), *Eucallacea linatus* (Schedl), *Leptoxyleborus concisus* (Blandford), *Leptoxyleborus punctatissimus* (Eichhoff), *Hyledius xanthophylli* (Browne), *Platypus fraterculus* Schedl, *Platypus jansoni* Chapuis, *Platypus nivalis* Sampson, *Platypus praepositus* Schedl, *Platypus pseudocupulatus* Schedl, *Platypus triangulatus* Browne, *Xyleborus adusticollis* (Motschulsky), *Xyleborus fallax* Eichhoff, *Xyleborus javanus* Eggers, *Xylosandrus assequens* Schedl, *Xylosandrus difficilis* (Eggers).

Xanthostemon pubescens

Xyleborus perforans (Wollaston).

Xeiopia spp.

Cryptocarenum lepidus Wood.

Xerospermum intermedium

Anasa beesonii (Eggers).

Xerospermum spp.

Crossotarsus schedli Browne, *Crossotarsus siporanus* Schedl, *Cyclorhizidion sisyrinchiorum* (Hagedorn), *Platypus protenus* (Schedl), *Xyleborinus pomelianus* (Schedl).

Xylia dolabriformis

Coccotrypes cyperi (Beeson), *Crossotarsus emorsus* Beeson, *Eucallacea velatus* (Sampson), *Platypus cupulatus* Chapuis, *Platypus signatus* Chapuis, *Xyleborus birmanus* Eggers, *Xyleborus fallax* Eichhoff.

Xylia xylocarpa

Crossotarsus bonvouloiri Chapuis, *Crossotarsus minax* (Walker), *Eucallacea andamanensis* (Blandford), *Eucallacea fornicatus* (Eichhoff), *Eucallacea interjectus* (Blandford), *Eucallacea velatus* (Sampson), *Leptoxyleborus concisus* (Blandford), *Leptoxyleborus*

- sordicauda* (Motschulsky), *Platypus signatus* Chapuis, *Xyleborus fallax* Eichhoff.
- Xylocarpus gangeticus**
Crossotarsus minusculus Chapuis, *Xyleborus bidentatus* (Motschulsky).
- Xylocarpus graucatum**
Platypus biuncus Blandford, *Xyleborus bidentatus* (Motschulsky).
- Xylopia aethiopica**
Doliopygus lefevrei Schedl, *Hypothenemus eruditus* Westwood, *Periommatius substriatus* Strohmeier, *Platypus hintzi* Schaufuss, *Platypus schenklingi* (Strohmeier), *Xyleborus annectens* Schedl.
- Xylopia caudata**
Baiois angustipennis (Schedl), *Platypus pseudocupulatus* Schedl.
- Xylopia cordata**
Platypus pseudocupulatus Schedl, *Platypus puerulus* (Schedl), *Platypus westwoodi* Chapuis.
- Xylopia hypolampa**
Platypus parallelus (Fabricius).
- Xylopia sericea**
Dryocoetoides flavus (Fabricius).
- Xylopia spp.**
Doliopygus conradti (Strohmeier), *Periommatius angustior* Schedl, *Periommatius artecircularis* Schedl, *Platypus hintzi* Schaufuss, *Platypus watutensis* Roberts, *Platypus xanthopus* Schedl, *Premnobius cavipennis* Eichhoff.
- Xylopia rullotii**
Coccotrypes suboratis Eggers.
- Xylopia villosa**
Cylindropalpus laudatus (Schedl), *Periommatius bispinus* Strohmeier, *Periommatius grandis* Schedl, *Platypus hintzi* Schaufuss, *Trachyostus aterrimus* (Schaufuss), *Trachyostus schaufuss* (Strohmeier), *Xyleborus ferrugineus* (Fabricius).
- Xylosma hawaiiense**
Xyleborinus saxeseni (Ratzeburg), *Xyleborus scabratus* Schedl.
- Xylosma monospora**
Cryphalus nyalubombeae Schedl, *Doliopygus erichsoni* (Chapuis), *Lanurgus olcaeformis* Schedl, *Xyleborinus acmulus* (Wollaston).
- Xylosma suaveolens**
Xyleborus perforans (Wollaston).
- Yucca spp.**
Cactopinus depressus Bright, *Hypothenemus crudiae* (Panzer), *Hypothenemus eruditus* Westwood, *Hypothenemus seriatus* (Eichhoff), *Pseudothysanoes frondicolens* Wood, *Pseudothysanoes yuccae* (Wood), *Pseudothysanoes yuccavorus* Wood.
- Zanthoxylum flutum**
Ambrosiodinus lecontei Hopkins.
- Zanthoxylum rhetsum**
Euallacea andamanensis (Blandford).
- Zea mays**
Hypothenemus californicus Hopkins, *Pagiocerus frontalis* (Fabricius).
- Zelkora carpinifolia**
Scolytus ensifer Eichhoff.
- Zelkora formosana**
Xyleborus taiwanensis Browne.
- Zelkora serrata**
Coccotrypes naidaijinensis (Murayama), *Crossotarsus niponicus* Blandford, *Euallacea validus* (Eichhoff), *Indocryphalus pubipennis* (Blandford), *Neopteleobius scutulatus* (Blandford).
- Zelkora spp.**
Scolytplatypus tycon Blandford.
- Zexmenia spp.**
Pityophthorus diligens.
- Zhauna golungensis**
Xyleborus alluaudi Schaufuss.
- Zhauna spp.**
Xyleborus alluaudi Schaufuss.
- Ziziphus jujuba**
Amasa amputatus (Blandford), *Xylosandrus germanus* (Blandford).
- Zygia spp.**
Amphicranus spectus Wood.

Bibliography Supplement

The following bibliography is a supplement to Wood & Bright (1987: 1–685) and lists only that literature missed by the original volume or published since 1987. While the original intent was to cite the names of all species of Scolytidae and Platypodidae mentioned in this literature, copies of many items could not be found (these are marked by an asterisk [*]) and several others were unintentionally overlooked by indexers for reasons unknown to us.

A

- ABGRALL, JEAN FRANÇOIS. 1986. Utilisation de phéromones artificielles contre *Ips typographus* [Utilization of artificial pheromones against *Ips typographus*]. English and Russian summaries. Eppo Bulletin 16(4): 633–637. (bv cn).
- ABGRALL, JEAN FRANÇOIS, AND DANIEL P. SCHVESTER. 1987. Observations sur le piégeage de *Ips typographus* L. après chablis [Observations on the trapping of *Ips typographus* L.]. English summary. Revue Forestière Française 39: 359–377. (cn).
- ABRAHAO, J., AND O. WEGMÜLLER. 1974. Criação do insecto *Hypocryphalus mangiferae* em culturas puras de *Ceratocystis frimbriata*. Revista Sociedade Brasileira de Fitopatologia 3: 31–32. (ec).
- ADLUNG, KARL GUNTHER, P. SCHICKE, AND J. O'SVATH. 1986. Analysis of an investigation on the control of the spruce bark beetle (*Ips typographus* L.) by means of pheromones: 1. Planning and evaluation of the individual trial [In German, English summary]. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 93: 462–475. (bv cn).
- AIHERN, F. J. 1985. The effects of bark beetle stress on the foliar spectral reflectance of lodgepole pine. International Journal of Remote Sensing 9: 1451–1465. (cn).
- AISAGBONIH, C. I. 1988. Pest incidence in marketed date palm fruits in Dutse, Kano State, Nigeria. Date Palm Journal 6: 287–295. (cn).
- AITKEN-SOUC, P. 1955. Quelques maladies et pestes cancrantes du caféier. Feuille d'Extension 59. 5 p. (cn).
- AKERS, R. PATRICK. 1959. Counterturns initiated by decrease in rate of increase of concentration. Possible mechanism of chemotaxis by walking female *Ips paraconfusus* bark beetles. Journal of Chemical Ecology 15: 183–205. (bv).
- AKERS, R. PATRICK, AND DAVID LEE WOOD. 1989a. Olfactory orientation responses by walking female *Ips paraconfusus* bark beetles I. Chemotaxis assay. Journal of Chemical Ecology 15: 3–24. (bv).
- . 1989b. Olfactory orientation responses by walking female *Ips paraconfusus* bark beetles II. In an anemotaxis assay. Journal of Chemical Ecology 15: 1147–1159. (bv).
- AKSENT'EV, S. I., AND L. N. UCHASTNOVA. 1986. Problem of the symbiosis of xylophagous beetles and ambrosia fungi [In Russian, English summary]. Nauchnye Doklady Vyshehei Shkoly, Biologicheskie nauki, Moscow 5: 5–22. (ay ec).
- ALBERT, A. M., AND HERMANN BOGENSCHULTZ. 1957. Die Bedeutung nicht aqualer Arthropoden-Verteilung bei Untersuchungen zur Belastbarkeit von Ökosystemen [The importance of unequal arthropod distribution in investigations of the tolerance of ecosystems]. English summary. Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie, Mitteilungen 5: 77–81. (hb).
- ALBUQUERQUE LEAO, MOACIR DE. 1941. O expurgo pelas ondas-curtas. Sociedade Brasileira de Agronomia, Rio de Janeiro, Revista 4: 319–325. (cn).
- *ALDEN. 1930. [*Scolytus rugulosus*]. Georgia State Board of Entomology, Atlanta, Bulletin 71: 31. ().
- *ALDERMAN, GIBBINGS, AND RUMSEY. 1913. [*Phloeosinus liminaris*, *Scolytus rugulosus*]. West Virginia Circular 7: 30, 35. ().
- *ALEKSEEV, A. V. 1957. K nachozhdeniu korojeda roda *Thamnurgus* Eichl. v Kurskoj oblasti. Uchenye zapiski Orechovo-Zujevskij pedagogiceskij institut 5: 159–163. ().
- ALEXANDER, K. N. A. 1988. Some records of *Xyloterus signatus* (L.) (Coleoptera: Scolytidae). Entomologist's Gazette 39: 326. (ds).
- ALLARD, C. B., AND D. MOORE. 1989. *Heterorhabditis* sp. nematodes as control agents for coffee berry borer *Hypothenemus hampei* Scolytidae. Journal of Invertebrate Pathology 54: 45–48. (ec cn).
- ALLEN, ANTHONY A. 1952. Coleoptera at Khole Park, Sevenoaks, Kent. Entomologists Record 64: 224–225. (ds).
- . 1955a. *Platypus parallelus* (F.) Col., Scolytidae again captured at light in S.E. London. Entomologist's Monthly Magazine 121: 141. (ds).
- . 1955b. The earliest British capture of *Scolytus laevis* Chapuis (Col., Scolytidae). Entomologist's Monthly Magazine 121: 195. (ds).
- . 1989. [Note to D. Nash]. Entomologist's Record and Journal of Variation 101: 152. (ds).
- ALLSOPE, P. G. 1957. *Hypothenemus californicus* Hopkins (Coleoptera: Scolytidae) recorded from Australia. Australian Entomological Society, Journal 26: 322. (ds).
- AMMAN, GENE DOYLE. 1955a. A test of lodgepole pine hazard rating methods for mountain pine beetle infestation in southeastern Idaho. Pages 186–200 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 240 p. (ec hb).
- . 1955b. The beetle: behavior, biology, and life cycle. Pages 2–7 in M. D. McGregor and D. M. Cole, Integrating management strategies for the mountain pine beetle with multiple-resource management of lodgepole pine. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, General Technical Report INT-174. (bv hb).
- . 1985a. Lodgepole pine selection by mountain pine beetle in relation to growth and vigor following thinning. Pages 197–213 in T. L. Payne and H. Saarenmaa,

- Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn).
- _____. 1958b. Mountain pine beetle (*Dendroctonus ponderosae* Hopkins) infestations in relation to growth and vigor of lodgepole pine (*Pinus contorta* Douglas) following thinning. International Congress of Entomology, Proceedings 15: 414. (cn).
- _____. 1959a. Proceedings: Symposium on the management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (bv cn ec hb).
- _____. 1959b. Why partial cutting in lodgepole pine stands reduces losses to mountain pine beetle. Pages 45–59 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- AMMAN, GENE DOYLE, AND JOHN A. ANHOLD. 1959. Preliminary evaluation of hazard and risk rating variables for mountain pine beetle infestations in lodgepole pine stands. Pages 22–27 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- AMMAN, GENE DOYLE, EUGENE D. LESSARD, LYNN A. RASMUSSEN, AND CURTIS G. O'NEIL. 1955. Lodgepole pine vigor, regeneration, and infestation by mountain pine beetle following partial cutting on the Shoshone National Forest, Wyoming. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, Research Paper INT-396. 8 p. (cn).
- AMMAN, GENE DOYLE, AND MARK D. MCGREGOR. 1955. Assessing stand hazard and risk hazard rating and predicting tree loss in managed stands. Pages 29–30 in M. D. McGregor and D. M. Cole, Integrating management strategies for the mountain pine beetle with multiple-resource management of lodgepole pine. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, General Technical Report INT-174. (cn).
- AMMAN, GENE DOYLE, MARK D. MCGREGOR, AND ROBERT E. DOLPH, JR. 1955. Mountain pine beetle. United States Department of Agriculture, Forest Service, Forest Insect and Disease Leaflet 2. 11 p. (cn ec hb).
- AMMAN, GENE DOYLE, MARK D. MCGREGOR, RICHARD FRANKLIN SCHMITZ, AND R. D. OAKES. 1955. Susceptibility of lodgepole pine to infestation by mountain pine beetles following partial cutting of stands. Canadian Journal of Forest Research 15: 655–695. (cn).
- AMMAN, GENE DOYLE, AND JUDITH E. PASEK. 1956. Mountain pine beetle in ponderosa pine: effects of phloem thickness and egg gallery density. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, Research Paper INT-367. 7 p. (ec hb).
- AMMAN, GENE DOYLE, AND LASZLO SAFRANYIK. 1955. Insects of lodgepole pine: impacts and control. Pages 107–124 in D. M. Baumgartner, R. G. Krebill, J. T. Arnott, and G. F. Weetman, Symposium proceedings, Washington State University, Cooperative Extension Service. (cn).
- AMMAN, GENE DOYLE, AND RICHARD FRANKLIN SCHMITZ. 1955. Mountain pine beetle-lodgepole pine interactions and strategies for reducing tree losses. Ambio 17: 62–65. (cn).
- AMMAN, GENE DOYLE, RALPH W. THIER, MARK D. MCGREGOR, AND RICHARD FRANKLIN SCHMITZ. 1959. Efficacy of verbenone in reducing lodgepole pine infestation by mountain pine beetles in Idaho. Canadian Journal of Forest Research 19: 60–64. (cn).
- ANDERBRANT, OLLE. 1955. Dispersal of reemerged spruce bark beetles, *Ips typographus* (Coleoptera, Scolytidae): a mark-recapture experiment. Journal of Applied Entomology 99: 21–25. (bv hb).
- _____. 1956. A model for the temperature and density dependent reemergence of the bark beetle *Ips typographus*. Entomologia Experimentalis et Applicata 40: 51–55. (ec hb).
- _____. 1955a. Reproductive strategies in *Ips typographus*. International Congress of Entomology, Proceedings 15: 409. (hb).
- _____. 1955b. Survival of parent and brood adult bark beetles, *Ips typographus*, in relation to size, lipid content and reemergence or emergence day. Physiological Entomology 13: 121–129. (ay hb).
- _____. 1955c. Reproduction and competition in the spruce bark beetle *Ips typographus*. Dissertation (published), Department of Ecology, Lund University, Sweden. 106 p. (ec hb).
- _____. 1959. Reemergence and second brood in the bark beetle *Ips typographus*. Holarctic Ecology 12: 494–500. (hb).
- ANDERBRANT, OLLE, AND J. LOFQVIST. 1955. Relation between first and second brood production in the bark beetle *Ips typographus*. Oikos 53: 357–365. (hb).
- ANDERBRANT, OLLE, AND FREDRIK SCHLYTER. 1957a. Differences in morphology and sexual size dimorphism between the Dutch elm disease vectors *Scolytus laevis* and *Scolytus scolytus* (Col., Scol.). Zeitschrift für Angewandte Entomologie 103: 375–386. (ay).
- _____. 1957b. Ecology of the Dutch elm disease vectors *Scolytus laevis* and *Scolytus scolytus* (Coleoptera: Scolytidae) in southern Sweden. Journal of Applied Ecology 24: 539–550. (bv ec).
- _____. 1959. Causes and effects of individual quality in bark beetles. Holarctic Ecology 12: 455–493.
- ANDERBRANT, OLLE, FREDRIK SCHLYTER, AND G. BIRGERSSON. 1955. Interspecific competition affecting parents and offspring in the bark beetle *Ips typographus*. Oikos 45: 89–98. (bv ec hb).
- ANDERBRANT, OLLE, FREDRIK SCHLYTER, AND J. LOFQVIST. 1955a. Dynamics of tree attack in the bark beetle *Ips typographus* under semi-epidemic conditions. International Congress of Entomology, Proceedings 15: 412. (bv hb).
- _____. 1955b. Dynamics of tree attack in the bark beetle *Ips typographus* under semi-epidemic conditions. Pages 35–51 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv hb).
- ANDERSON, W. C., R. W. GULDIN, AND J. M. VASIEVICH. 1955. Risk assessments of investments in loblolly pine plantations threatened by bark beetles. Pages 325–334 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest

- Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56, 383 p. (cn).
- _____. 1987. Assessing the risk of insect attack in plantation investments. *Journal of Forestry* 85: 46-47. (cn).
- ANDREEV, E. A. 1955. Fauna and ecology of *Proctolaelaps* Acarosejidae mites found in galleries made by bark beetles in Moscow Oblast Russian SFSR, USSR [In Russian]. *Biologicheski nauki*, Moscow 10: 34-37. (ec).
- *ANDREWS, A. W. 1923. The Coleoptera of the Shiras Expedition to Whitefish Point, Chippewa County, Michigan. *Papers of the Michigan Academy of Science* 1: 293-390. ().
- ANDREWS, MARK W., GEORGE L. BARNES, AND KEN PINKSTON. 195.. Dutch elm disease and its control. Oklahoma State University Cooperative Extension Service, Pest Management Series, Extension Facts No. 7602. 4 p. (cn).
- ANDRYSZAK, N. A., B. STAFFAN LINDGREN, and P. L. PAYNE. 1987. Electrophysiological responses by *Trypodendron lineatum* (Ol.) (Coleoptera, Scolytidae) from western Canada and Germany to lineatin and isomers of alpha-pinene. *Journal of Applied Entomology* 103: 23-25. (ay).
- ANGUS, R. B. 1964. Some northern Scottish Coleoptera. *Entomologist's Monthly Magazine* 100: 172-182. (ds).
- _____. 1965a. Further Coleoptera from Cumberland, Westmorland and northern part of Lancashire. *Entomologist's Monthly Magazine* 101: 4-8. (ds).
- _____. 1965b. Further notes on northern Scottish Coleoptera. *Entomologist's Monthly Magazine* 101: 9-12. (ds).
- _____. 1965c. Some new records of Coleoptera from Pembrokeshire. *Entomologist's Monthly Magazine* 101: 12-13. (ds).
- ANHOLD, JOHN A. 1986a. Biological evaluation of mountain pine beetle infestation: Grey's River Road, Bridger-Teton National Forest. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-S6-9. (cn).
- _____. 1986b. Biological evaluation of Douglas-fir beetle: Davis Hill, Buffalo Road, Bridger-Teton National Forest, 1985. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report S6-12. 5 p. (cn).
- _____. 1986c. Mountain pine beetle pheromone trapping, Roosevelt Ranger District, Ashley National Forest, 1985. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-S6-11. 4 p. (cn).
- _____. 1986d. MPB campground spray evaluation: Wasatch-Cache and Ashley National Forests. United States Department of Agriculture, Forest Service, Biological Evaluation Report S6-14. 7 p. (cn).
- _____. 1986e. Spruce beetle survey in Big Cottonwood Canyon, Salt Lake Ranger District, Wasatch-Cache National Forest, 1985. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-S6-16. 7 p. (cn).
- _____. 1987. Biological evaluation of spruce beetle populations, Big Cottonwood Canyon, Salt Lake Ranger District, Wasatch-Cache National Forest 1987. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-S6-17. 4 p. (cn).
- _____. 1988. Evaluation of preventive treatment for mountain pine beetle in high-value stands of lodgepole pine. United States Department of Agriculture, Forest Service, Intermountain Region (Ogden, Utah), Forest Pest Management Report R4-S8-3. 5 p. (cn).
- ANHOLD, JOHN A., AND MICHAEL J. JENKINS. 1987. Potential mountain pine beetle (Coleoptera: Scolytidae) attack of lodgepole pines as described by stand density index. *Environmental Entomology* 16: 735-742. (cn).
- ANONYMOUS. 1896c. *Bostrichus dispar* bei Munster in Spalierbäumen. Münster Westfälischen Provinzial-Vereins für Wissen-Schaft und Kunst, Jahresbericht, Zool.-Botan. Sekt. 1896: 25. (cn).
- _____. 1907. United States Department of Agriculture, yearbook, 1906. Washington, D.C. 720 p. (cn).
- *_____. 1920b. Nursery and orchard insect pests [*Scolytus rugulosus*]. Missouri Agricultural Experiment Station, Columbia, Bulletin 176. 35 p. ().
- _____. 1930j. Ulmensterben. Ulmensplintkafer und Schützimpfung. *Entomologische Zeitschrift* 44: 231. (cn).
- *_____. 1944c. [*Aricercus fici* (=eichloffii)]. *Agricultural Gazette, New South Wales [Australia]* 55(1): 21-22. ().
- *_____. 1951h. [*Scolytus rugulosus*]. Washington State Agriculture Extension Bulletin 419 (revised): 30. ().
- _____. 1956p. Southwest bug damage tops \$12 million. *Timberman* 58: 110. (cn).
- *_____. 1964z. A special edition of pine stem boring insects [In Japanese]. *Shinrin Boeki Nyusu [Forest Protection News]* 13(15). ().
- _____. 1967w. Parasitic wasp. *Chemical and Engineering News*, April 24: 14. (ec).
- _____. 1970w. Forest insect conditions in the United States, 1969. United States Department of Agriculture, Forest Service, Washington, D.C. 39 p. (cn).
- _____. 1971v. The population dynamics of the southern pine beetle in east Texas: the epidemic 1958-1965. *Southern Forest Research Institute, Progress Report July/August 1971: 13-15. (hb).*
- _____. 1971z. Spruce beetle attacks trees baited with frontaline. *Southern Forest Research Institute, Progress Report July/August 1971: 16. (bv).*
- _____. 1971aa. Control alternatives using frontalure: sticky traps. *Southern Forest Research Institute, Progress Report Sept./Oct. 1971: 17-19. (bv cn).*
- _____. 1971ab. Frontalure-cacodylic acid tested for control of southern pine beetle on the eastern shore of Virginia. *Southern Forest Research Institute, Progress Report Sept./Oct. 1971: 20. (cn).*
- _____. 1971ac. Grandipso: attractant for the southern five-spined engraver [*Ips grandicollis*] isolated and identified. *Southern Forest Research Institute, Progress Report Jan./Feb. 1971: 2-3. (bv).*
- _____. 1971ad. Winter control of the southern pine beetle with frontalure. *Southern Forest Research Institute, Progress Report Jan./Feb. 1971: 4. (cn).*
- _____. 1971ae. The population dynamics of the southern pine beetle in east Texas: seasonal population trends. *Southern Forest Research Institute, Progress Report Nov./Dec. 1971: 22-24. (hb).*
- _____. 1971af. A new compound that inhibits southern pine beetle response to attractants. *Southern Forest Research Institute, Progress Report May/June 1971: 9-10. (bv).*
- _____. 1971ag. Douglure: a powerful tool in manipulating the Douglas-fir beetle. *Southern Forest Research*

- Institute, Progress Report May/June 1971: 11–12. (cn).
- _____. 1971ah. Forest insect and disease conditions in the United States, 1970. United States Department of Agriculture, Forest Service, Washington, D.C. 44 p. (cn).
- _____. 1972u. The attractant of the six-spined engraver beetle, *Ips calligraphus*, isolated and identified. Southern Forest Research Institute, Progress Report Mar./Apr. 1972: 6. (bv).
- _____. 1972v. The cybernetics of pheromone production in the engraver beetles. Southern Forest Research Institute, Progress Report Mar./April 1972: 7–8. (bv).
- _____. 1972w. Trans-verbenol: a pheromone common to many bark beetle species? Southern Forest Research Institute, Progress Report Mar./April 1972: 8. (bv).
- _____. 1972x. The Cadillac beetle, *Platypus flavicornis*. Southern Forest Research Institute, Progress Report May/June 1972: 10–11. (cn).
- _____. 1972y. Dendroctol, a new weapon against the Douglas-fir beetle. Southern Forest Research Institute, Progress Report May/June 1972: 11–12. (cn).
- _____. 1972z. Forest insect and disease conditions in the United States, 1972. United States Department of Agriculture, Forest Service, Washington, D.C. O-506-654. 72 p. (cn).
- _____. 1973r. Geographical distribution and physiological differences among *Dendroctonus frontalis* populations. Southern Forest Research Institute, Progress Report Mar./April 1973: 6–8. (ay ds).
- _____. 1975v. Biological evaluation: Groundhog Park and Albert Park, Del Norte Ranger District, Rio Grande National Forest. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-75-9. 1 p. (cn).
- _____. 1975w. Spruce evaluation: Dolores Ranger District, San Juan National Forest. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-75-11. 2 p. (cn).
- _____. 1975x. Spruce beetle biological evaluation in the "Greenhorn Sale Area." Pike and San Isabel National Forests. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-75-13. 1 p. (cn).
- _____. 1975y. Biological evaluation of Douglas-fir beetle near Douglas Pass. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-75-15. 1 p. (cn).
- _____. 1975z. Biological evaluation of spruce beetle near the Workman and Fernwood Sales, Rio Grande National Forests. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-75-18. 2 p. (cn).
- _____. 1975aa. Biological evaluation: mountain pine beetle infestation, Cassia Division, Twin Falls Ranger District, Sawtooth National Forest. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah. 7 p. (cn).
- _____. 1975ab. Barkbilleskader i skogbruket. Norges Offentlige Utredninger. Universitetsforlaget, Oslo. NOU 1975: 20. 33 p. (cn).
- _____. 1976q. Biological evaluation: mountain pine beetle, Cassia Division, Twin Falls Ranger District, Sawtooth National Forest. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah. 3 p. (cn).
- _____. 1976ad. Insect and disease conditions of pinyon pine and Utah juniper in Mesa Verde National Park, Colorado. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-78-4. (cn).
- _____. 1976ae. Mexican observe work on southern pine beetle. United States Department of Agriculture, Southern Pine Beetle Research and Applications Program, Pineville, Louisiana, News Release. (ms).
- _____. 1979ab. Forest insect and disease conditions in Alaska, 1977. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska, Series R10-31. 11 p. (cn).
- _____. 1979ac. Forest insect and disease conditions in Alaska, 1978. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska, Report 62. 35 p. (cn).
- _____. 1980x. Forest insect and disease conditions in Alaska, 1979. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska, Report 115. 17 p. (cn).
- _____. 1980y. Kisatchie National Forest initiates southern pine beetle hazard rating. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, Southern Pine Beetle News, February 17: 1. (cn).
- _____. 1980z. Cooperative magazine warns of southern pine beetle crisis. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, Southern Pine Beetle News, August, 23: 4. (cn ms).
- _____. 1980aa. Computer will help determine profit in SPB-killed timber. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, Southern Pine Beetle News, August, 23: 3. (cn).
- _____. 1981w. Forest insect and disease conditions in Alaska, 1980. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska, Report 146. 17 p. (cn).
- _____. 1982m. New report details toxicant studies performed under ESPBRAP. Pest Management News 32: 1–2. (cn).
- _____. 1982n. South Carolina southern pine beetle outbreak collapses. Pest Management News 32: 1. (cn).
- _____. 1983h. The Forest Pest Control Section: twenty years of progress in forest protection. Texas Forest News 62(fall): 12–15. (cn).
- _____. 1983i. Forest insect and disease conditions in Alaska, 1981–82. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska, Report 173. 24 p. (cn).
- _____. 1983j. IPM of southern pine forests. Pest Management News 40: 1–8. (cn).
- _____. 1983k. Twenty years of progress in forest protection. Texas Forest News 62: 12–15. (cn).
- _____. 1984h. Texas forest pest report 1982–1983. Texas Forest Service Publication 136. 31 p. (cn).
- _____. 1984i. Forest insect and disease conditions in Alaska in 1983. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Report R-10. 25 p. (cn).
- _____. 1984j. Forest protection. Pages 72–97 in Wattle

- Research Institute, Pietermaritzburg, South Africa, Annual Report. (cn ec hb).
- _____. 1954k. Annual report 1953. Maine Forest Service, Entomological Division, Augusta, Maine. (cn).
- _____. 1955a. Amendments to the movement of spruce wood order. Page 32 in Sixty-fifth annual report and accounts of the Forestry Commission for the year ended 31 March 1955, together with the Comptroller and Auditor-General's report of the accounts. London, United Kingdom. (cn).
- _____. 1955b. Breeding *Rhizophagus grandis*. Page 35 in Sixty-fifth annual report and accounts of the Forestry Commission for the year ended 31 March 1955, together with the Comptroller and Auditor-General's report on the accounts. London, United Kingdom. (ec hb).
- _____. 1955c. *Dendroctonus micans*. Pages 36–37 in Sixty-fifth annual report and accounts of the Forestry Commission for the year ended 31 March 1955 together with the Comptroller and Auditor-General's report on the accounts. London, United Kingdom. (cn ec).
- _____. 1955d. Pheromone traps for exotic bark beetles. Page 37 in Sixty-fifth annual report and accounts of the Forestry Commission for the year ended 31 March 1955, together with the Comptroller and Auditor-General's report on the accounts. London, United Kingdom. (cn).
- _____. 1955e. Cocoa. Parliamentary Paper, Parliament of Fiji, No. 62. (cn).
- _____. 1955f. Department of Entomology. Pages 114–141 in Biennial report of the Waite Agricultural Institute, South Australia, 1954–1955, Glen Osmond, South Australia. (cn ec).
- _____. 1955g. Entomology/Nematology. Pages 91–123 in Thirty-eight annual detailed technical report 1954–1955, Chikmagalar, India, Central Coffee Research Institute. (cn).
- _____. 1955h. Forest insect and disease conditions in Alaska. Region 10, 1954 Report. Journal of Applied Entomology 149: 1–26. (cn).
- _____. 1955i. Forest pest conditions in California, 1954. California Forest Pest Control Action Council, California Department of Forestry, Sacramento, California. 26 p. (cn).
- _____. 1955j. Forest protection. Pages 69–79 in Institute for Commercial Forest Research, University of Natal, Pietermaritzburg, South Africa, Annual Report 1955. (cn ec hb).
- _____. 1955k. Insects and diseases of trees in the South. United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia, General Report RS–GR–5. 98 p. (cn hb ms).
- _____. 1955m. *Dendroctonus micans*. Pages 18–19 in Sixty-fifth annual report and accounts of the Forestry Commission for the year ended 31 March 1955, together with the Comptroller and Auditor-General report on the accounts. London, United Kingdom. (cn).
- _____. 1955n. *Ips grandicollis* in Queensland pine plantations. Queensland Department of Forestry, Information Sheet 17. 2 p. (cn).
- _____. 1955p. The bark beetles: killers of pine and spruce. PesTopics No. 1, British Columbia Ministry of Forests, Pest Management, Protection Branch. 2 p. (cn ms).
- _____. 1956a. Outbreak of *Dendroctonus micans*, the great spruce beetle. Plant Health Information Sheet 10, Forestry Commission, United Kingdom. 1 p. (cn).
- _____. 1956b. Forest pest conditions in California, 1955. California Forest Pest Control Action Council, California Department of Forestry, Sacramento, California. 24 p. (cn).
- _____. 1956c. Entomology. Pages 41–44 in Report on forest research, Forestry Commission, London, United Kingdom. (cn).
- _____. 1956d. Outbreak of *Dendroctonus micans*, the great spruce bark beetle. Progress report. Plant Health Information Sheet 9, Forestry Commission, United Kingdom. 4 p. (cn).
- _____. 1956e. Semiochemical baits integrated into pine management program. IPM Practitioner 8: 8. (cn).
- _____. 1956f. Mountain pine beetle control program 1950–1956, a success story. Alberta Forestry, Lands and Wildlife, Forest Service Publication 1/143. 12 p. (cn ms).
- _____. 1957a. *Dendroctonus micans*, the great spruce beetle. General control strategy. Plant Health Information Sheet 11, Forestry Commission, United Kingdom. (cn).
- _____. 1957b. Entomology/Nematology. Pages 90–212 in Coffee Board Research Department, India, Thirty-ninth Annual Report 1955–1956. (cn hb).
- _____. 1957c. Entomology. Pages 49–54 in Report on forest research, Forestry Commission, London, United Kingdom. (cn).
- _____. 1957d. Forest pest conditions report for the Northeastern Area, 1956. United States Department of Agriculture, Forest Service, Northeastern Area, Broomall, Pennsylvania, Forest Pest Management Report NA–FR–34. 35 p. (cn).
- _____. 1957e. Forest pest conditions in California, 1956. California Forest Pest Council, California Department of Forestry, Sacramento, California. 28 p. (cn).
- _____. 1957f. Rx for SPB: Cut! Cut! Cut! The American Tree Farmer 6: 13. (cn ms).
- _____. 1957g. Final environmental impact statement for the suppression of the southern pine beetle, Southern Region. Volume 1: Chapters I–VI; Volume 2: Chapter VII; Volume 3: Appendices. United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia, Management Bulletin RS–MB2. (cn).
- _____. 1957h. Regional Research Station. Kalpetta, Kerala. Plant Protection. Pages 197–199 in Coffee Board Research Department, India. (ec).
- _____. 1957i. Mountain pine beetle trapping with the Lindgren funnel trap. Phero Tech Technical Bulletin. 2 p. (cn).
- _____. 1957j. Le grand bostryche, le stenographe *Ips sexdentatus* [The pine engraver beetle, *Ips sexdentatus*]. Informations-Forêt 1. 5 p. (cn hb).
- _____. 1957k. Bark beetles: a threat to our forests. British Columbia Ministry of Forests and Lands. [Unpaginated]. 20 p. (cn ms).
- _____. 1957m. Forest and shade tree insect and disease conditions for Maine: a summary of the 1956 situation. Insect and Disease Management Division, summary report 1. Maine Forest Service, Augusta, Maine. 22 p. (cn).
- _____. 1957n. Mountain pine beetle: a menace to pine forests. PesTopics No. 24. British Columbia Ministry of Forests, Pest Management, Protection Branch. 4 p. (cn).
- _____. 1958a. *Dendroctonus micans*, the great spruce bark

- beetle. General control strategy. Plant Health Information Sheet 12, Forestry Commission, United Kingdom. 2 p. (cn).
- _____. 1955b. Forest pest conditions in California, 1957. California Forest Pest Council, California Department of Forestry, Sacramento, California. 25 p. (cn).
- _____. 1955c. Biological evaluation of a mountain pine beetle infestation on the Sawtooth National Recreation Area, Idaho, 1957. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report 55-04. 8 p. (cn).
- _____. 1955d. Forest insect and disease conditions in Alaska, 1955. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, Forest Pest Management Report R10-55-C. 16 p. (cn).
- _____. 1955e. Montana forest pest conditions and program highlights 1955. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana. Forest Pest Management Report 59-2. 21 p. (cn).
- _____. 1955f. TSF [Texas Forest Service] grid hazard. Southern pine beetle fact sheet number 43. United States Department of Agriculture, Forest Service, Protection Report RS-PR9. 2 p. (cn).
- _____. 1955g. Entomology. Pages 44-48 in Report on forest research, Forestry Commission, London. United Kingdom. (cn).
- _____. 1955h. *Dendroctonus micans*. Pages 36-37 in Sixty-seventh annual report and accounts 1956-1957. Forestry Commission, Edinburgh. United Kingdom. (cn).
- _____. 1955i. Pheromone traps for exotic bark beetles. Page 37 in Sixty-seventh annual report and accounts 1956-1957. Forestry Commission, Edinburgh, United Kingdom. (cn).
- _____. 1955j. Insect control. Sixty-seventh annual report and accounts 1956-1957. Forestry Commission, Edinburgh, United Kingdom. (cn).
- _____. 1955k. Ecology and control of *Dendroctonus* bark beetles. Report on the International Symposium, Borzhomy, Georgia, USSR, 22-27 September 1957. (cn ec).
- _____. 1955m. Southern pine beetle demonstration areas. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, SPB Update. 4 p. (cn).
- _____. 1955n. Southern pine beetle Fact Sheet number 43: TFS [Texas Forest Service] grid hazard. United States Department of Agriculture, Forest Service, Forest Pest Management, Protection Report RS-PR9. 2 p. (cn).
- _____. 1955p. Texas court case. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, SPB Update, August. (ms).
- _____. 1955q. Southern pine beetle outlook for 1955. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana, SPB Update, August. (cn).
- _____. 1959a. *Dendroctonus micans*, the great spruce bark beetle. General control strategy. Plant Health Information Sheet 13, Forestry Commission, United Kingdom. 2 p. (cn).
- _____. 1959b. Forest insect and disease conditions in Ontario, summer 1959. Government of Canada, Forestry Service, Ontario Region, Survey Bulletin. 27 p. (cn).
- _____. 1959c. *Hypothenemus hampei* Ferrari. CAB International Institute of Entomology, Distribution Maps of Pests 170 (revised). 3 p. (ds).
- _____. 1955a. Identification and control of three Front Range insects. Colorado State Forest Service, Colorado State University, Fort Collins, Circular 121-0155. 2 p. [Undated, issued about 1956]. (cn).
- _____. 1955b. Dutch elm disease in Colorado. Colorado State Forest Service and Colorado State University Extension Service, Circular. 6 p. [Undated, issued about 1956]. (cn).
- *ANTHON, EDWARD W. 1949b. [*Scolytus rugulosus*]. Washington State Horticulture Association, Proceedings 45: 205-206. ().
- *ANTHON, EDWARD W., ET AL. 1950. [*Scolytus rugulosus*]. Washington State Agriculture Extension Bulletin 49: 23-24. ().
- ANTROBIUS, W. L., J. E. TAYLOR AND S. KOHLER. 1959. Montana forest pest conditions and program highlights. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, and Montana Department of State Lands and Forests, Division Report 90-2. 8 p. (cn).
- APPLEJOHN, M. J. 1965. Status of insects in the Swastika district. Pages E35-42 in Annual district reports of the Forest Insect and Disease Survey, Ontario, 1964. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- ARNETT, ROSS HAROLD, JR. 1968. The beetles of the United States: a manual for identification. American Entomological Institute, Ann Arbor, Michigan. 1112 p. (reprinted 1971). (tx).
- ARNO, STEPHEN F., AND R. J. HOFF. 1959. Silvics of whitebark pine (*Pinus albicaulis*). United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah. Research Paper INT-253. 11 p. (cn).
- ARNOLD, DON C., AND STEVEN R. JIRACEK. 1952. Beetles of Oklahoma pines, Oklahoma State University Agricultural Experiment Station, MP-113. 16 p. (cn).
- ASCENCIO CERDA, VICTOR E., AND B. E. SERRATO B. 1954a. Evaluación de cinco insecticidas organofosforados para el combate de *Dendroctonus mexicanus* Hopks. en el área de explotación forestal de Atenquique, Jal. [Evaluation of five organophosphorus insecticides against *Dendroctonus mexicanus* in the forest of Atenquique, Jalisco]. Ciencia Forestal 9: 42-64. (cn).
- _____. 1954b. Evaluación de Gusathion-M20 (Azinfos Metil) y lindano contra el *Dendroctonus mexicanus* Hopks. en el área de explotación forestal de Atenquique, Jal. [Evaluation of gusathion M-20 (Azinphos methyl) and lindane for controlling *Dendroctonus mexicanus* in the forest of Atenquique, Jalisco]. Ciencia Forestal 9: 3-22. (cn).
- ASHIRU, MISILU O., AND ROBERT IMRE GARA. 1951. Towards an improved fat extraction technique in the Douglas-fir beetle, *Dendroctonus pseudotsugae* Hopk. West African Science Association, Journal 26: 29-39. (ay ms).
- ASHWORTH, ALLAN CHARLES, AND A. M. CYANCARA. 1953. Paleogeology of the southern part of the Lake Agassiz basin. Pages 135-156 in J. T. Teller and Lee Clayton, Glacial Lake Agassiz. Geological Society of Canada, Special Paper 26. (ds).
- ASHWORTH, ALLAN CHARLES, AND J. W. HOGANSON. 1957. Coleoptera bioassociations along an elevational gradient in the Lake Region of southern Chile, and comments on the postglacial development of the

- fauna. Entomological Society of America, Annals 50: 865-895. (ds).
- °ATKINSON, GEORGE FRANCIS. 1889. [*Scolytus rugulosus*]. South Carolina Agricultural Experiment Station, n. s., Bulletin 4: 79-80. ().
- ATKINSON, THOMAS HARRIS. 1989a. New species and notes on Mexican Hylesininae (Coleoptera: Scolytidae). Insecta Mundi 3: 57-64. (ds tx).
- . 1989b. New synonymy, new species, and notes on Scolytidae (Coleoptera) from southeastern United States. Coleopterists Bulletin 43: 325-337. (ds tx).
- . 1989c. The species of *Platypus* of Florida (Coleoptera: Platypodidae). Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular 321. 4 p. (ds hb tx).
- ATKINSON, THOMAS HARRIS, AND ARMANDO EQUIHUA MARTINEZ. 1985a. Lista comentada de lost coleopteros Scolytidae y Platypodidae del Valle de Mexico. Folia Entomologica Mexicana 65: 63-108. (ds hb).
- . 1985b. Notes on biology and distribution of Mexican and Central American Scolytidae (Coleoptera). I. Hylesininae, Scolytinae except Cryphalini and Corthylini. Coleopterists Bulletin 39: 227-235. (ds hb).
- . 1985c. Notes on biology and distribution of Mexican and Central American Scolytidae (Coleoptera). II. Scolytinae: Cryphalini and Corthylini. Coleopterists Bulletin 39: 355-363. (ds hb).
- . 1986a. Biology of bark and ambrosia beetles (Coleoptera: Scolytidae and Platypodidae) of a tropical rain forest in southeastern Mexico with an annotated checklist of species. Entomological Society of America, Annals 79: 414-423. (ds hb).
- . 1986b. Biology of the Scolytidae and Platypodidae (Coleoptera) in a tropical deciduous forest at Chame-la, Jalisco, Mexico. Florida Entomologist 69: 303-310. (ds hb).
- . 1985. Notas sobre la biologia de Scolytidae y Platypodidae (Coleoptera) de Mexico y Centro-america. Folia Entomologica Mexicana 76: 53-105. (ds hb).
- ATKINSON, THOMAS HARRIS, JOHN L. FOLTZ, AND M. D. CONNOR. 1988. Flight patterns of phloem-and-wood-boring Coleoptera (Scolytidae, Platypodidae, Curculionidae, Buprestidae, Cerambycidae) in a north Florida slash pine plantation. Environmental Entomology 17: 259-265. (bv hb).
- ATKINSON, THOMAS HARRIS, JOHN L. FOLTZ, AND ROBERT CLEVELAND WILKINSON. 1988. *Xyleborus crassiusculus* (Motschulsky): an Asian ambrosia beetle recently introduced into Florida (Coleoptera: Scolytidae). Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Entomology Circular 310. (cn).
- ATKINSON, THOMAS HARRIS, EDGAR MARTINEZ-FERNANDEZ, EDUARDO SAUCEDO-CEPEDES, AND ARMANDO BURCOS-SOLORIO. 1986. Scolytidae y Platypodidae (Coleoptera) asociados a selva baja y comunidades derivadas en el estado de Morelos, Mexico. Folia Entomologica Mexicana 69: 41-82. (ds hb).
- ATKINSON, THOMAS HARRIS, R. J. RABAGLIA, AND DONALD EDWARD BRIGHT, JR. 1990. Newly detected exotic species of *Xyleborus* (Coleoptera: Scolytidae) with a revised key to species in eastern North America. Canadian Entomologist 122: 93-104. (ds tx).
- ATKINSON, THOMAS HARRIS, R. J. RABAGLIA, S. B. PECK, AND J. L. FOLTZ. 1991. New records of Scolytidae and Platypodidae (Coleoptera) from the United States and the Bahamas. Coleopterists Bulletin 45: 152-164. (ds hb).
- ATKINSON, THOMAS HARRIS, EDUARDO SAUCEDO-CEPEDES, EDGAR MARTINEZ-FERNANDEZ, AND ARMANDO BURCOS-SOLORIO. 1986. Coleopteros Scolytidae y Platypodidae asociados con las comunidades vegetales de clima templado y frio en el estado de Morelos, Mexico. Acta Zoologica Mexicana, nueva serie, 17: 1-58. (ds hb).
- °ATUAHENE, S. K. N. 1970. The economic effect of insect pests on the timber industry in Ghana. Forest Product Research Institute, Ghana, Technical Newsletter 4: 4-8. ().
- AULLO, MANUEL. 1918. [*Hylastes ater*]. Boletin de la R. Sociedad Espanola de Entomologia, Saragossa 1: 163-171. (ds).
- AUSTARA, ØYSTEIN, ERIKKI ANNILA, BRODER BEJER-PETERSEN, AND BENGT EHNSTROM. 1983. Insect pests in the forest of the Nordic countries 1977-1981. Fauna Norvegica Series B 31(1): 5-15. (cn ds).
- AUSTARA, ØYSTEIN, ALF BAKKE, AND F. MIDTGAARD. 1986. Response in *Ips typographus* to logging waste odors and synthetic pheromones. Journal of Applied Entomology 10: 194-195. (bv).
- AUSTARA, ØYSTEIN AND F. MIDTGAARD. 1986. On the longevity of *Ips typographus* L. adults. Journal of Agricultural Entomology 102: 106-111. (hb).
- AXENT'JEV, S. I. 1957. New genus and species of shot-hole borers from Tadzhikistan (Coleoptera, Scolytidae). Pages 150-151 in A. G. Kireitschuk, Systematics and geographical distribution of Coleoptera [In Russian]. USSR Academy of Science, Proceedings of the Zoological Institute, Leningrad. 161 p. (tx).
- AYER, W. A., L. M. BROWNE, M. C. FENG, H. ORSZANSKA, AND M. SAEEDI-GHOMI. 1986. The chemistry of the blue stain fungi: Part I. Some metabolites of *Ceratocystis* species associated with mountain pine beetle infected lodgepole pine. Canadian Journal of Chemistry 64: 904-909. (ec).

B

- BAADER, E. J. 1989. Comparative studies on semiochemicals in *Pityogenes* spp. (Coleoptera: Scolytidae) and their potential usage in pest management systems [In German, English summary]. Journal of Applied Entomology 107: 1-31. (bv cn).
- BACCETTI, BACCIO, ANNA G. BURRINI, ANNA MAGNANO, AND LUIGI MAGNANO. 1985. Spermatozoi e sistematica dei Curculionioidei. Pages 155-161 in Atti XIV Congresso Nazionale Italiano di Entomologia, Palermo. 918 p. (ay).
- BAEKSTROM, P., Ü. JACOBSSON, T. NORIN, AND C. R. UNELIUS. 1988. Synthesis and characterization of all four isomers of methyl 2,4-decadienoate for an investigation of the pheromone components of *Pityogenes chalcographus*. Tetrahedron 44: 2541-2545. (bv ms).
- BAERT, L. L. A., AND J. P. MAELFAIT. 1977. Contribution to the knowledge of the arachno- and entomofauna of different wood habitats. Part II. Influence of the micro-relief upon epedalic Coleoptera. Societe Royale Belge d'Entomologie, Bulletin et Annales 113: 101-110. (ec).
- BAISIER, M., AND J. C. GREGOIRE. 1985. Factors influencing oviposition in *Rhizophagus grandis* (Coleoptera: Rhizophagidae), specific predator of the bark beetle *Dendroctonus micans* (Coleoptera: Scolytidae).

- Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent 53: 1159–1167. (ec).
- BAISIER, M., J. C. GREGOIRE, K. DELINTE, AND O. BONNARD. 1955. The role of spruce monoterpene derivatives as oviposition stimuli for *Rhizophagus grandis*, a predator of the bark beetle *Dendroctonus micans*. Pages 359–365 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan (eds.), Mechanisms of woody plant defenses against insects: search for pattern. Springer, Berlin. xiv + 416 p. (ec).
- BAKER, BRUCE HOWARD, BRUCE B. HOSTETLER, AND THOMAS A. LAURENT. 1975. Forest insect and disease conditions in Alaska, 1974. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska. 13 p. (cn).
- BAKER, BRUCE HOWARD, AND THOMAS A. LAURENT. 1974. Note on dead and dying larch in Alaska. Pages 6–7 in J. F. Chancellor and H. B. Toko, Forest insect and disease conditions in the United States, 1973. United States Department of Agriculture, Forest Service, Washington, D.C. 55 p. (cn).
- BAKER, G. T., AND T. EVAN NEEKEER. 1986. Sensory receptors on the ovipositor of *Thanasimus dubius* (F.) (Coleoptera: Cleridae). Societe Entomologique de France. Annales 22: 49–52 (ec).
- BAKER, PETER S. 1959. A sampling plan for a control project against the coffee berry borer (*Hypothenemus hampei*) in Mexico. Tropical Pest Management 35: 169–172. (cn).
- BAKER, PETER S., J. F. BARRERA, AND J. E. VALENZUELA. 1959. The distribution of the coffee berry borer *Hypothenemus hampei* in southern Mexico: a survey for a biocontrol project. Tropical Pest Management 35: 163–165. (cn).
- BAKKE, ALF. 1955a. Deploying pheromone-baited traps for monitoring *Ips typographus* populations. Journal of Applied Entomology 99: 33–39. (cn).
- . 1955b. Methods and effects of suppressing bark beetle populations. Pages 169–174 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO Conference proceedings, Canada Department of the Environment, Forestry Service, and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 239 p. (cn).
- . 1957. Repression of *Ips typographus* infestation in stored logs by semiochemicals. Scandinavian Journal of Forest Research 2: 179–185. (bv cn).
- . 1958a. Potential use of semiochemicals for integrated control of bark beetles in Europe. Pages 257–261 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- . 1958b. Potential use of semiochemicals for integrated control of bark beetles in Europe. International Congress of Entomology, Proceedings 18: 415. (cn).
- . 1959. The recent *Ips typographus* outbreak in Norway: experiences from a control program. Holarctic Ecology 12: 151–159. (cn).
- BAKKE, ALF AND R. LIE. 1959. Mass trapping. Pages 67–87 in A. R. Jutsum and R. F. S. Gordon, Insect pheromones in plant protection. John Wiley and Sons, New York. (bv).
- BAKKE, ALF, TORFINN SÆTHIER, AND OYSTEIN AUSTARA. 1959. Response by *Ips typographus* to pheromone dispensers stored for 5 years. Scandinavian Journal of Forest Research 4: 393–394. (bv).
- BAKKE, ALF, TORFINN SÆTHIER, AND TORSTEIN KVAMME. 1953. Mass trapping of the spruce bark beetle *Ips typographus*: pheromone and trap technology. Meddelelser fra Norsk Institutt for Skogforskning 38: 35. (cn).
- BAKKE, ALF AND JEAN PIERRE VITE. 1957. Methylbutynol no efficient replacement for the natural pheromone component in *Ips typographus*. Naturwissenschaften 74: 292–293. (bv).
- BAKKE, JAN M., J. KRANE, AND T. SKJETNE. 1959. Confirmations of bark beetle pheromones: the importance of the medium. Acta Chemica Scandinavica 43: 777–782. (bv ms).
- BALAKRISHNAN, M. M., P. K. VINODKUMAR, AND C. B. PRAKASAN. 1959. Impact of the predator *Eupelmus* sp. (Hymenoptera: Eupelmidae) on the incidence of *Xylosandrus compactus*. Journal of Coffee Research 19: 88–90. (ec).
- BALAZI, STANISLAW, JACEK MICHALSKI, AND E. RATAJCZAK. 1957. Materiały do znajomości wrogów naturalnych *Ips acuminatus* Gyll. (Coleoptera: Scolytidae) [Contribution to the knowledge of the natural enemies of *Ips acuminatus* Gyll.]. Polskie Pismo Entomologiczne 57: 735–745. (ec).
- *BALDWIN, C. H. 1916. [*Scolytus rugulosus*]. Page 131 in Eighth annual report of the Indiana State Entomologist. ().
- BALICK, M. J., D. C. FURTH, AND G. COOPER-DIVER. 1978. Biochemical and evolutionary aspects of arthropod predation on ferns. Oecologia (Berlin) 35: 55–59. (bv ec).
- BAITENSWEILER, WERNER. 1955. "Waldsterben": forest pests and air pollution. Journal of Applied Entomology 99: 77–85. (ec).
- BANERJEE, BARUNDEB. 1953. Arthropoda accumulation on tea in young and old habitats. Ecological Entomology 8: 117–123. (ec).
- BANGSHOLT, F. 1975. Fjerde tillæg til "Fortegnelse over Danmarks biller" (Coleoptera). Entomologiske Meddelelser 43: 65–96. (ds).
- *BARBER, H. S. 1919. [*Pagiocerus rimosus* (=frontalis)]. Entomological Society of Washington, Proceedings. 21: 59. ().
- BARBOSA, PEDRO, AND MICHAEL R. WAGNER. 1959. Introduction to forest and shade tree insects. Academic Press, San Diego, California. xi + 639 p. (cn ec hb).
- BARGER, JACK H., AND WILLIAM N. CANNON, JR. 1957. Response of smaller European elm bark beetles to pruning wounds on American elm. Journal of Arboriculture 13: 102–104. (bv).
- BARR, BARBARA A., D. L. HANSON, AND C. S. KOEHLER. 1978. Red turpentine beetle: a pest of pines. Division of Agricultural Sciences, University of California, Berkeley. Leaflet 21055. 4 p. (cn hb).
- BARRICOLI, J. A. F., LELAND CHANDLER, AND N. F. LOPES. 1955. Rosposta fisiologica do feijoeiro (*Phaseolus vulgaris* L.) ao forate a suas consequencias entomologicas. III. Efeito sobre as pragas e producao da cultura, no plantio da "seca" [Physiological response of *Phaseolus vulgaris* to phorate and the entomological consequences] [English summary]. Revista Ceres 35: 370–387. (cn).
- BARRY, PATRICK J. 1955. Biological evaluation of southern pine beetle infestations on the Tyger and Enoree Ranger Districts, Sumter National Forest, South Carolina. United States Department of Agriculture,

- Forest Service, Southern Region, Atlanta, Georgia, Forest Pest Management Report 85-1-22. (cn).
- *BARTIL, AGNAR. 1902. Grantoken, Dens Aarsager og Forbyggels. Det Kongelige Selskab for Norge vel, Kristiania (=Oslo). 6S p. ().
- BARTINDALE, G. C., AND G. R. BARTINDALE. 1948. Coleoptera of the Macclesfield district. *Entomologist's Monthly Magazine* 84: 130-135. (ds).
- BARTOS, DALE L. 1985a. Microclimate as a factor in lodgepole pine (*Pinus contorta* Douglas) stands and tree selection by mountain pine beetles (*Dendroctonus ponderosae* Hopkins). Pages 215-230 in T. L. Payne and H. Saarenmaa (eds.), Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv ec).
- _____. 1985b. Microclimate as a factor in lodgepole pine (*Pinus contorta* Douglas) stand and tree selection by mountain pine beetles (*Dendroctonus ponderosae* Hopkins). International Congress of Entomology, Proceedings 15: 414. (bv ec).
- BARTOS, DALE L., AND GENE DOYLE AMMAN. 1989. Microclimate: an alternative to tree vigor as a basis for mountain pine beetle infestations. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, Research Paper INT-400. 10 p. (ec).
- BARTOS, DALE L., AND K. B. DOWNING. 1985. Development of an expert system for endemic populations of mountain pine beetle in lodgepole pine forests. Pages 232-233 in Proceedings: International symposium on advanced technology in natural resources management. Resource Technology SS, Fort Collins, Colorado, 20-23 June 1985. American Society for Photogrammetry and Remote Sensing, Falls Church, Virginia. (cn).
- _____. 1989. Evolution of a research prototype expert system for endemic populations of mountain pine beetle in lodgepole pine forests. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, Research Note INT-390. 8 p. (cn).
- BASHAM, JACK TUCKER. 1959. Studies in forest pathology: XX, investigations of the pathological deterioration in killed balsam fir. *Canadian Journal of Botany* 37: 292-326. (cn).
- _____. 1956. Biological factors influencing stem deterioration rates and salvage planning in balsam fir killed after defoliation by spruce budworm. *Canadian Journal of Forest Research* 16: 1217-1229. (cn).
- BATRA, LEKH RAJ. 1985. Ambrosia beetles and their associated fungi: research trends and techniques. *Indiana Academy of Science, Proceedings, Plant Science* 94: 137-145. (ec).
- BATRA, LEKH RAJ, S. W. T. BATRA, AND TOSHIO NAKASHIMA. 1986. Some techniques to study ambrosia beetles and their associated fungi. *Hokkaido Musashi Women's Junior College, Memoirs* 18: 73-94. (ec ms).
- BATTISTI, A. 1985. *Dendroctonus micans* (Kugelann) in Italia (Coleoptera: Scolytidae). *Frustula Entomologica* 7-8: 631-637. (ay cn ec ds).
- BATTISTI, A., R. MENARDI, AND G. SALA. 1986. Sulla presenza del coleottero scolidite *Dendroctonus micans* Kugelann in boschi di abete rosso del Veneto [On the presence of the scolytid beetle *Dendroctonus micans* Kugelann in Norway spruce woods in the Venice district] [English summary]. *Italia Forestale e Montana* 41: 197-203. (cn ec hb).
- BAYLIS, N. T. 1986. Observations of damage of a secondary nature following a wild fire at the Otterford State Forest. *South African Forestry Journal* 137: 36-37. (cn).
- *BEACH. 1899. [*Scolytus rugulosus*]. New York State Agricultural Experiment Station, Bulletin 170: 315-445. ().
- BEAVER, ROGER A. 1954. The biology of the ambrosia beetle, *Sucus niissimai* (Eggers) Col.: Scolytidae, in Fiji. *Entomologist's Monthly Magazine* 120: 99-102. (hb).
- _____. 1956. The taxonomy, mycangia and biology of *Hypothenemus curtipes* (Schedl), the first known cryptaline ambrosia beetle (Coleoptera: Scolytidae). *Entomologica Scandinavica* 17: 131-135. (ec hb ds tx).
- _____. 1957a. Biological studies on bark beetles of the Seychelles (Col., Scolytidae). *Journal of Applied Entomology* 104: 11-23. (cn ds hb).
- _____. 1957b. The bark and ambrosia beetles (Coleoptera: Scolytidae and Platypodidae) of Tonga. *New Zealand Entomologist* 9: 64-70. (cn hb ds tx).
- _____. 1958a. Biological studies on ambrosia beetles of the Seychelles (Col., Scolytidae and Platypodidae). *Journal of Applied Entomology* 105: 62-73. (bv cn hb).
- _____. 1958b. Patterns of bark and ambrosia beetles in tropical forests. International Congress of Entomology, Proceedings 15: 410. (hb).
- _____. 1959a. Bark and ambrosia beetles (Coleoptera: Scolytidae) newly recorded from Fiji, and their potential economic importance. *South Pacific Journal of Natural Science* 9: 1-7. (ds).
- _____. 1959b. Insect-fungus relationships in the bark and ambrosia beetles. Pages 121-143 in N. Wilding, N. M. Collins, P. M. Hammond, and J. F. Webber (eds.), *Insect-fungus interactions*. Academic Press, London. xvi + 344 p. (ec).
- _____. 1990a. New records and new species of bark and ambrosia beetles from Thailand (Coleoptera: Scolytidae and Platypodidae). *Deutsche Entomologische Zeitschrift, Neue Folge*, 37: 279-284. (ds tx).
- _____. 1990b. The bark and ambrosia beetles of Kiribati, South Pacific (Col., Scolytidae and Platypodidae). *Entomologist's Monthly Magazine* 126: 149-151. (ds tx).
- _____. 1991. New synonymy and taxonomic changes in Pacific Scolytidae (Coleoptera). *Naturhistorisches Museum Wien, Annalen* 92: 57-97. (tx).
- BEAVER, ROGER A., AND K. LOYTTYNIEMI. 1955a. Bark and ambrosia beetles (Coleoptera: Scolytidae) of Zambia. *Revue de Zoologie Africaine* 99: 63-85. (hb ds tx).
- _____. 1955b. The platypodid ambrosia beetles of Zambia (Coleoptera: Platypodidae). *Revue de Zoologie Africaine* 99: 113-134. (hb ds tx).
- _____. 1959. Further observations on the bark and ambrosia beetles of Zambia (Coleoptera: Scolytidae and Platypodidae). *Revue Zoologie Africaine* 103: 255-290. (ds tx).
- BEAVER, ROGER A., AND P. A. MADDISON. 1990. The bark and ambrosia beetles of the Cook Islands and Niue (Coleoptera: Scolytidae and Platypodidae). *Journal of Natural History* 24: 1365-1375. (ds tx).
- *BECK, GUNTHER. 1856. Coleoptera, Scolytidae. Pages 626-627 in M. A. Becker (ed.), *Fauna von Hertenstein in Niederosterreich. II. Teil H. Halbband*. ().
- *BECKER. 1913. [*Scolytus rugulosus*]. *Arkansas Horticultural Society, Proceedings* 1912-1913: 29. ().
- BECKLEY, PAUL R. 1959. Case history: application of risk assessment, Flathead National Forest. Pages 34-36

- in G. D. Amman, Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah. General Technical Report INT-262. 119 p. (cn).
- BEDARD, WILLIAM DELLES, JR., K. Q. LINDAHL, JR., PAUL E. TILDON, AND DAVID LEE WOOD 1985. Behavior of the western pine beetle during host colonization. *Journal of Chemical Ecology* 11: 1249-1262. (bv).
- BEESON, CYRIL FREDERICK CHERRINGTON. 1934. The role of insects in dying off of sal. *Indian Forester* 1934: 539-543. (bh).
- *BEGEMANN, H. 1927c. [*Xyleborus compactus*]. *Archiv f. v. d. Koffeculture* 1: 273. ().
- BEGLEY, MICHAEL J., AND J. F. GROVE. 1955. Metabolic products of *Phomopsis oblonga*. Part 1: 3 5A 6,7,8,9,9A 9B octahydro-7 9B-dimethylnaphtho-1 2-C-furan-1-311-one oblongide. *Journal of the Chemical Society Perkin, Transactions* 1: 861-864. (bv ms).
- *BEJER-PETERSEN, BRODER. 1959. An der ventes staerkere angreb af *Hylesinus micans*? Borstlig Budstikke 19(22): 87-88. ().
- . 1955a. *Dendroctonus micans* in Denmark. Pages 2-19 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of the European Communities and the University Libre de Bruxelles, Bruxelles, Belgium. Proceedings of seminar, 3-4 October 1954. 141 p. (ec).
- . 1955b. Sitkagran og "micans." [English summary]. *Dansk Skooforenings Tidsskrift* 73: 37-42. (ec).
- *BEKMAN. 1929. [*Pagiodercus zcav (=frontalis)*]. *Izv. prikl. Ent.* 4(1): 151-166. ().
- BELANGER, ROGER P. 1989. Silviculture, a remedy for southern pine beetle problems. Society of American Foresters National Convention, Proceedings 1985: 187-191. (cn).
- BELANGER, ROGER P., J. F. GODBEE, T. MILLER, AND R. S. WEBB. 1983. Salvage cutting in southern pine forests in South Carolina, Georgia, and Alabama. United States Department of Agriculture, Forest Service, SE-24: 352-356. (cn).
- BELANGER, ROGER P., ROY L. HEDDEN, AND MICHAEL R. LENNARTZ. 1988. Potential impact of the southern pine beetle on red-cockaded woodpecker colonies in the Georgia Piedmont. *Southern Journal of Applied Forestry* 12: 194-199. (ec).
- BELANGER, ROGER P., ROY L. HEDDEN, AND FRANK H. TAITER. 1986. Managing Piedmont forests to reduce losses from the littleleaf disease-southern pine beetle complex. United States Department of Agriculture, Forest Service, Cooperative State Research Service, Integrated Pest Management Handbook. Agriculture Handbook 649. 19 p. (cn ec).
- BENNET, E. M., AND TERRI A. TATTAR. 1985. Bluestain fungi and insect vector interaction in Japanese black and Scots pine mortality. *Arboricultural Journal* 12: 237-247. (ec).
- BENOIT, PAUL. 1985. Nomenclatura insectorum canadensium. Noms d'insectes au Canada. Insect names in Canada. Government of Canada, Canadian Forestry Service, Laurentian Forest Research Centre, Sainte-Foy, Quebec. 299 p. (tx).
- . 1986. Nomenclatura insectorum canadensium. Noms d'insectes au Canada. Insect names in Canada. Supplementum. Government of Canada, Forestry Service, Laurentian Forest Research Centre, Sainte-Foy, Quebec. 114 p. (tx).
- BENOIT, PAUL AND R. BLAIS. 1984. Pertes de bois causes par le dendroctone du maleze. *Phytoprotection* 65: 89. (bv cn).
- BENOIT, PAUL, GASTON LAFLAMME, G. BONNEAU, AND R. PICHIER. 1982. Insectes et maladies des arbres Quebec, 1981. *Forest Conservation* 45(10): 1-19. (cn).
- . 1983. Insectes et maladies des arbres Quebec, 1982. *Forest Conservation* 49(10): 1-23. (cn).
- *BENTZ, BARBARA J. 1984. Phenetic and phylogenetic relationships among *Dendroctonus* (Coleoptera: Scolytidae) bark beetles. Unpublished thesis, University of Idaho, Moscow, Idaho. 109 p. ().
- BENTZ, BARBARA J., C. K. LISTER, JOHN M. SCHMID, S. A. MATA, L. A. RASMUSSEN, AND D. HANEMAN. 1989. Does verbenone reduce mountain pine beetle attacks in susceptible stands of ponderosa pine? United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado, Research Note RM-495. 4 p. (cn).
- BENZ, GEORG. 1955a. *Cryphalus abietis* (Ratz.) and *Ips typographus* (L.) new for Turkey, and a note on the tree killing capacity of *Pityophthorus pityographus* (Ratz.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 58: 275. (cn ds).
- . 1955b. *Dendroctonus micans* in Turkey: the situation today. Pages 43-47 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of the European Communities and the Universite Libre de Bruxelles, Belgium. Proceedings of seminar, 3-4 October 1954. 141 p. (ec).
- BENZ, GEORG, P. BOVEY, AND P. JUXOD. 1986. On the specific attraction of the males of the six-toothed spruce bark beetle, *Pityogenes chalcographus* (L.) to a mixture of synthetic pheromones of the eight-toothed spruce bark beetle, *Ips typographus* (L.) (Coleoptera, Scolytidae). *Experimentia* (Basel) 42: 325-326. (bv).
- BERGDAIL, DALE R. 1982. Occurrence of the pinewood nematode in eastern larch. Pages 47-55 in J. E. Appleby and R. B. Malek, Proceedings of the 1982 national pinewilt workshop, Rosemont, Illinois. (bv cn).
- BERGDAIL, DALE R., D. L. K. SMELTZER, AND S. S. HALIK. 1984. Components of a conifer wilt disease complex in the northeastern states [abstract]. Proceedings of the joint U.S.-Japan pine wilt disease seminar, Honolulu, Hawaii. (cn).
- . 1986. Components of a conifer wilt disease complex in the northeastern United States. Pages 152-157 in V. Dropkin, The resistance mechanisms of pines against pine wilt disease, 7-11 May 1984, Honolulu, Hawaii. Japan Society for the Promotion of Science. (ec).
- BERISFORD, CHARLES WAYNE, AND U. EUGENE BRADY. 1986. Field evaluation of fenitrothion for post-attack control of the southern pine beetle. *Journal of Entomological Science* 21: 139-141. (cn).
- BERISFORD, CHARLES WAYNE, U. EUGENE BRADY, CARL WARREN FATZINGER, AND B. H. EBEL. 1986. Evaluation of a repellent for prevention of attacks by three species of southern pine bark beetles (Coleoptera: Scolytidae). *Journal of Entomological Science* 21: 316-318. (cn).
- BERISFORD, CHARLES WAYNE, AND DONALD L. DAHLSTEN

1989. Biological control of *Ips grandicollis* (Eichhoff) (Coleoptera: Scolytidae) in Australia: a preliminary evaluation. Pages 81–93 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ec).
- BERRISFORD, CHARLES WAYNE, G. L. DEBARR, AND THOMAS LEE PAYNE. 1955. Utilization of pheromones in forest pest management in insects and diseases of southern forests. Pages 92–96 in R. A. Gover and J. P. Jones, Insects and diseases in southern forests. 34th annual symposium proceedings. Louisiana Agricultural Experiment Station, Baton Rouge, Louisiana. (bv cn).
- BERRISFORD, CHARLES WAYNE, R. F. MIZELL III, L. H. KUDON, AND K. D. WARE. 1955. Line intersect sampling techniques for estimating *Ips typographus* in bagging residue. Pages 13–25 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. General Technical Report SO-56. 353 p. (cn ms).
- BERRISFORD, CHARLES WAYNE, AND THOMAS LEE PAYNE. 1955a. Regional variation: a potential factor in the integration of behavioral chemicals into southern pine beetle management. Pages 275–282 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn).
- _____. 1955b. Regional variation: a potential factor in the integration of behavioral chemicals into southern pine beetle management. International Congress of Entomology, Proceedings 15: 415. (bv cn).
- BERRIOS, C., JUAN M. MENENDEZ, AND M. RODRIGUEZ. 1957. Presencia del genero *Ips* (Coleoptera, Scolytidae) sobre nuevas especies de pinos en el norte de la provincia de Matanzas [Presence of *Ips* on new species of pines in the north of Matanzas Province] [English summary]. Revista Forestal Baracoa 17: 113–115. (cn ds).
- *BERRY, P., AND M. VAQUERO. 1957. Lista de insectos clasificados de El Salvador. Boletín Técnico 21: 9–50. ().
- BERRYMAN, ALAN ANDREW. 1967. Entomological considerations in logging managed forests. Pages 73–110 in R. W. Bruce, Logging managed forests. Washington State University Press, Pullman. (cn ec).
- *_____. 1954. Threshold theory and its application to pest population management. Pages 9–57 in Pest and pathogen control: strategic, tactical and policy models. International Series on Applied Systems Analysis. 13. ().
- _____. 1957. The theory and classification of outbreaks. Pages 3–30 in P. Barbosa and J. C. Schultz, Insect outbreaks. Academic Press, San Diego, California. xiv + 578 p. (cn ec).
- _____. 1955a. Dynamics of forest insect populations: patterns, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec hb).
- _____. 1955b. Pests and the stability of forest ecosystems. Northwestern Environmental Journal 4: 351–355. (cn ec).
- _____. 1959. Adaptive pathways in scolytid-fungus associations. Pages 145–159 in N. Wilding, N. M. Collins, P. M. Hammond, and J. F. Webber, Insect-fungus interactions. Academic Press, London. xvi + 344 p. (cc).
- BERRYMAN, ALAN ANDREW, B. DENNIS, KENNETH FRANCIS RAFFA, AND NILS CHR. STENSETH. 1955. Evolution of optimal group attack, with particular reference to bark beetles (Coleoptera: Scolytidae). Ecology 66: 895–903. (bv).
- BERRYMAN, ALAN ANDREW, AND GEORGE T. FERRELL. 1955. The fir engraver beetle in western states. Pages 555–577 in A. A. Berryman, Dynamics of forest insect populations: patterns, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec).
- BERRYMAN, ALAN ANDREW, KENNETH FRANCIS RAFFA, J. A. MILLSTEIN, AND NILS CHR. STENSETH. 1959. Interaction dynamics of bark beetle aggregation and conifer defense rates. Oikos 56: 256–263. (cc tx).
- BERRYMAN, ALAN ANDREW, AND RONALD WILLIAM STARK. 1955a. Assessing the risk of stand destruction by pests. Pages 163–169 in D. M. Baumgartner, R. G. Krebil, J. T. Armott, and G. F. Weetman, Lodgepole pine: the species and its management. Cooperative Extension Service, Washington State University, Pullman. (cn ec).
- _____. 1955b. Assessing the risk of forest insect outbreaks. Journal of Applied Entomology 99: 199–205. (cn).
- BERRYMAN, ALAN ANDREW, AND NILS CHR. STENSETH. 1959. A theoretical basis for understanding and managing biological populations with particular reference to the spruce bark beetle. Holarctic Ecology 12: 357–394. (cn).
- BERRYMAN, ALAN ANDREW, NILS CHR. STENSETH, AND A. S. ISAEV. 1957. Natural regulation of herbivorous forest insect populations. Oecologia (Berlin) 71: 174–184. (cn hb).
- BETREM, JOHAN GEORGE. 1936. Insectenbestrijding. Bergcultures, Batavia 10: 196–201. (cn).
- *BEUTENMULLER, W. 1951. Bibliographical catalogue of the described transformations of North American Coleoptera. New York Microscopical Society 7: 1–52. ().
- BEVAN, DERMOT. 1957. Forest insects: a guide to insects feeding on trees in Britain. Forestry Commission, Handbook 1. (cn hb tx).
- BHAT, P. K. 1957. A review of entomological research in coffee with reference to major insect pests. Journal of Coffee Research 17: 77–79. (cn).
- BHAT, S. S., AND K. SREEDHARAN. 1955. Association of *Ambrosiella xylebori* Brader, with the shot-hole beetle *Xyleborus compactus* Eichhoff, a pest of Robusta coffee. Journal of Coffee Research 15: 54–57. (ec).
- *BICKHARDT HEINRICH. 1911a. Book review: Prof. J. Rouhal, Die Coleopterenfauna von Litauen. Entomologische Blätter 7: 71. ().
- *_____. 1911b. Book review: Dr. Max Wolff, Die Borkenkäfer, ihre Schaden um ihre Bekämpfung. Entomologische Blätter 7: 70–71. ().
- *_____. 1911c. Verzeichnis der Spezialisten für Coleopteren. Entomologische Blätter 7: 25–44. ().
- BIGGER, M. 1955. The insect pests of forest plantation trees in Solomon Islands [Scolytidae, p. 115–132]. Solomon Islands Forest Record No. 4. Overseas Development Natural Resources Institute, Kent, United Kingdom. 190 p. (ec).
- BILLINGS, RONALD FORREST. 1953a. Southern pine beetle in Honduras: new approach to an old problem. Texas Forest News 62: 2–5. (cn hb).
- _____. 1953b. SPB [southern pine beetle]: twenty years of

- progress in forest protection. *Texas Forest News* 62: 12–15. (cn).
- _____. 1954d. SPB [southern pine beetle] control in wilderness areas of Texas. *Texas Forest News* 63: 12–13. (cn).
- _____. 1955a. Implementing forest pest management: the forester's role. Pages 111–119 in R. A. Goyer and J. P. Jones, *Insects and diseases of southern forests*. 34th Annual Forestry Symposium, Louisiana State University, Baton Rouge, Louisiana. 135 p. (cn).
- _____. 1955b. Southern pine bark beetle and associated insects: effects of rapidly-released host volatiles on response to aggregation pheromones. *Journal of Applied Entomology* 99: 453–491. (bv ec).
- _____. 1956a. Coping with forest insect pests in southern wilderness areas, with emphasis on the southern pine beetle. Pages 120–125 in D. L. Kulhavy and R. N. Conner (eds.), *Wilderness and natural areas in the eastern United States: a management challenge*. Symposium proceedings, Stephen F. Austin University, Nacogdoches, Texas, 13–15 May 1955. 416 p. (cn).
- _____. 1956b. Forest pests and control programs in Mexico. *Texas Forest News* 65(summer): 14–16. (cn).
- _____. 1956c. Southern pine beetle demonstration project: final synopsis. *Texas Forest News* 65(spring): 8–9. (cn).
- _____. 1957. Southern pine beetle inhibitor: a promising new approach to direct control. *Texas Forest News* 66(2): 7–9. (cn).
- _____. 1955a. Forecasting southern pine beetle infestation trends with pheromone traps. Pages 295–306 in T. L. Payne and H. Saaremaa (eds.), *Integrated control of scolytid beetles*. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn).
- _____. 1955b. Forecasting southern pine beetle infestation trends with pheromone traps. *International Congress of Entomology. Proceedings* 18: 415. (bv cn ec).
- _____. 1955c. Forestry in the Dominican Republic revisited. *Texas Forest News* 67(3): 16–18. (cn ds).
- _____. 1955d. Tornado-damaged pines are easy targets for pine bark beetle attack. *Texas Forestry* 29(1): 2. (cn).
- _____. 1959. SPB [southern pine beetle] update. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, Pineville, Louisiana. 4 p. (cn).
- BILLINGS, RONALD FORREST, AND CHARLES MELROSE BRYANT 1953a. New approach developed to forecast SPB [southern pine beetle] outbreaks. *Texas Forestry* 24(5): 1. (cn).
- BILLINGS, RONALD FORREST, CHARLES MELROSE BRYANT, HERBERT A. PASE III, K. A. WILSON, AND C. WALKER 1955. Southern pine beetle technology transfer in Texas. Pages 55–63 in G. D. Hertel, S. J. Branham, and K. M. Swain, Sr., *Technology transfer in integrated forested forest pest management in the South*. United States Department of Agriculture, Forest Service, General Technical Report SE-34. 77 p. (cn hb).
- BILLINGS, RONALD FORREST, CHARLES MELROSE BRYANT, AND K. H. WILSON. 1955. Development, implementation, and validation of a large area hazard- and risk-rating system for southern pine beetle. Pages 226–232 in S. J. Branham and R. C. Thatcher, *Integrated pest management research symposium: the proceedings*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- BILLINGS, RONALD FORREST, AND RICHARD A. GOYER. 1957. New approaches to control southern pine beetle. *Forest Farmer* 99: 453–491. (cn).
- BILLINGS, RONALD FORREST, EDWARD H. HOLSTEN, AND ANDRUS EGLITIS 1972. Insects associated with *Pinus radiata* D. Don in Chile. *Turrialba* 22: 105–109. (ec ds).
- BILLINGS, RONALD FORREST, AND FORREST E. VARNER. 1986. Why control southern pine beetle infestations in wilderness areas?: the Four Notch and Huntsville State Park experiences. Pages 129–134 in D. L. Kulhavy and R. N. Conner, *Wilderness and natural areas in the eastern United States: a management challenge*. Symposium proceedings, 13–15 May 1955. Stephen F. Austin University, Nacogdoches, Texas. 416 p. (cn).
- BILLINGS, RONALD FORREST, AND JAMES D. WARD. 1954. How to conduct a southern pine beetle aerial detection survey. *Texas Forest Service Circular* 267. 19 p. (cn).
- BING, J. W. 1955. Seasonal abundance and distribution of insects associated with *Ips grandicollis* in Nacogdoches County, Texas. Unpublished thesis, Stephen F. Austin University, Nacogdoches, Texas. 111 p. (cn ec).
- BIRGERSSON, GÖRAN. 1955. The semiochemical system of *Ips typographus*: quantitative variation in production and release, and behavioural activity. [Unpublished?] dissertation, University of Göteborg, Sweden. (bv).
- _____. 1959. Host tree resistance influencing pheromone production of *Ips typographus* (Coleoptera: Scolytidae). *Holarctic Ecology* 12: 451–456. (bv ec).
- BIRGERSSON, GÖRAN, AND GUNNAR BERGSTROM 1959. Volatiles released from individual spruce bark beetle entrance holes: quantitative variations during the first week of attack. *Journal of Chemical Ecology* 15: 2465–2484. (bv).
- BIRGERSSON, GÖRAN, AND A. LEUFVEN. 1955. The influence of host response to *Ips typographus* and fungal attack on production of semiochemicals. *Insect Biochemistry* 15: 761–770. (bv cn).
- BIRGERSSON, GÖRAN, F. SCHLYTER, G. BERGSTROM, AND J. LOFQVIST. 1955. Individual variations in aggregation pheromone content of the bark beetle, *Ips typographus*. *Journal of Chemical Ecology* 14: 1737–1760. (bv).
- BISTROM, OLOF, AND R. VÄISÄNEN. 1955. Ancient-forest invertebrates of the Pyhä-Häkki National Park in central Finland. *Acta Zoologica Fennica* 185: 1–69. (ds).
- BLACKMAN, MAULSBY WILLETT. 1940c. [Book review]. *Entomological Society of Washington, Proceedings* 42: 46–47. (ms).
- BLACKWELL, MEREDITH J., ROBERT BRIDGES, JOHN CONRAD MOSER, AND THIELMA J. PERRY. 1956. Hypophoretic dispersal of a *Ptychodiphora* anamorph. *Science* 232: 993–995. (ec).
- BLACKWELL, MEREDITH J., JOHN CONRAD MOSER, AND J. WISNIEWSKI. 1959. Ascospores of *Ptychodiphora* on mites associated with beetles in trees and wood. *Mycological Record* 92(3): p. ? (ec).
- BLACKWELL, MEREDITH J., THIELMA J. PERRY, J. ROBERT BRIDGES, AND JOHN CONRAD MOSER. 1956. A new species of *Ptychodiphora* and its thaxteriella anamorph. *Mycologia* 78: 605–612. (ec).
- BLAIR, K. G. 1940. A note on *Xyleborus sampsoni* Donisth.

- (Col., Scolytidae). *Entomologist's Monthly Magazine* 76: 40. (tx).
- BLAKE, E. A., MICHAEL R. WAGNER, AND THOMAS W. KOEBER. 1956. Insects destructive to ponderosa [pine]. Pages 238–242 in R. C. Shearer (ed.), *Conifer tree seed in the inland Mountain West: symposium proceedings, 5–6 August 1955, Missoula, Montana*. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station (Ogden, Utah), General Technical Report. (cn).
- _____. 1959. Relative effect of seed and cone insects on ponderosapine in northern Arizona. *Journal of Economic Entomology* 52: 1691–1694. (cn).
- BLANCHÉ, C. A., JOHN DEAVOURS HODGES, AND T. EVAN NEBEKER. 1955. Changes in bark beetle susceptibility indicators in a lightning-struck loblolly pine. *Canadian Journal of Forest Research* 15: 397–399. (cn).
- BLANCHÉ, C. A., JOHN DEAVOURS HODGES, T. EVAN NEBEKER, AND D. M. MOEHRING. 1953. Southern pine beetle: the host dimension. *Mississippi Agriculture and Forestry Experiment Station. Bulletin* 917. 29 p. (ec).
- BLANCHÉ, C. A., T. EVAN NEBEKER, JOHN DEAVOURS HODGES, B. L. KARR, AND J. J. SCHMITT. 1955. Effect of thinning damage on bark beetle susceptibility indicators in loblolly pine. Pages 471–479 in E. Shoulders, *Third Biennial Southern Silvicultural Research Conference: proceedings, 7–8 November 1954, Atlanta, Georgia*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station (Research Triangle Park, North Carolina), General Technical Report SO-54. (cn).
- *BLEASDELL, G. G. 1937. The Rhynchophora of Iowa. *Iowa State College Journal of Science* 11: 404–445. ().
- BLINOVA, S. L. 1971. Investigations of crop and forest insect pest nematodes in the USSR. *International Congress of Entomology, Proceedings* 13: 55. (ec).
- BOCHER, JENS. 1955. The Coleoptera of Greenland. *Meddelelser om Grønland, Bioscience* 26: 1–100. (ds).
- BOEIM, H. 1958. Bark beetle slot traps. Page 437 in *Communications of the Federal Biological Institute for Agriculture and Forestry, Berlin-Dahlem, No. 245. Forty-sixth German Plant Protection Convention*. Paul Parey, Berlin. xvii + 524 p. (cn).
- *BOGDANOVA, D. A. 1954. O roli nasekomykh-kislofagov v lesnykh soobshchestvakh Sibiri. Pages 5–10 in *Nasekomye v ekosistemakh lesnoj zony Sibiri, Izdatelstvo universiteta*. ().
- _____. 1956. Trunk pests of pine in a fire-damaged forest of the upper Obregion [In Russian, English summary]. *Biologicheskikh Nauk* 3: 71–76. (cn).
- _____. 1957. Pine xylophages in the area of industrial pollution [In Russian]. *Ekologiya* 1957: 87–89. (ec).
- _____. 1955. Complexes of xylophagous insects in pine cultures contaminated by root fungus [In Russian, English summary]. *Izvestiya Sibirskogo Otdeleniya Akademii Nauk SSSR* 3: 25–28. (ec).
- BOLLENBACHER, B., AND K. E. GIBSON. 1956. Mountain pine beetle: a land manager's perspective. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 6–15. 5 p. (cn).
- BOMBOSCH, SIEGFRIED. 1957. Eine neue Möglichkeit zum einsatz von Borkenkäferpheromonen? [A new possibility for the use of bark beetle pheromones?] [English summary]. *Journal of Applied Entomology* 103: 360–363. (bv cn).
- _____. 1958a. A new possibility to use bark beetle pheromones? *International Congress of Entomology, Proceedings* 18: 415. (cn).
- _____. 1958b. Some considerations on the use of bark beetle pheromones. Pages 263–265 in T. L. Payne and H. Saarenmaa, *Integrated control of scolytid bark beetles*. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv).
- BOMBOSCH, SIEGFRIED, I. ENGLER, H. GOSSENAUER, AND B. IFFMANN. 1955. Über die Rolle von Pheroprax bei der Desiedlung der Fichte durch den Buchdrucker (*Ips typographus* L.) [In German, English summary]. *Journal of Applied Entomology* 100: 455–463. (bv).
- BOMBOSCH, SIEGFRIED, AND M. JOHANN. 1955. Notes on the host selection of *Ips typographus*. Pages 43–51 in L. Safranik, *The role of the host in the population dynamics of forest insects*. IUFRO conference proceedings, Canada Department of the Environment, Canadian Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 240 p. (bv).
- BORDEN, JOHN HARVEY. 1955. Aggregation pheromones. Pages 257–255 in G. A. Kerkut and L. I. Gilbert, *Comprehensive insect physiology, biochemistry, and pharmacology*. Pergamon Press, Oxford. (bv).
- _____. 1955. The striped ambrosia beetle. Pages 579–596 in A. A. Berryman, *Dynamics of forest insect populations: patterns, causes, implications*. Plenum Press, New York. xx + 603 p. (bv cn ec hb).
- _____. 1959. Semiochemicals and bark beetle populations: exploitation of natural phenomena by pest management strategists. *Holarctic Ecology* 12: 501–510. (bv).
- BORDEN, JOHN HARVEY, L. J. CHONG, AND D. J. BERGVINSON. 1958. Assessment of two pine oil treatments to protect stands of lodgepole pine from attack by the mountain pine beetle. *Entomological Society of British Columbia, Journal* 55: 28–33. (cn).
- BORDEN, JOHN HARVEY, L. J. CHONG, AND T. E. LACEY. 1956. Pre-logging baiting with semiochemicals for the mountain pine beetle, *Dendroctonus ponderosae*, in high hazard stands of lodgepole pine. *Forestry Chronicle* 62: 20–23. (cn).
- BORDEN, JOHN HARVEY, L. J. CHONG, AND B. S. LINDGREN. 1990. Redundancy in the semiochemical message required to induce attack on lodgepole pines by the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). *Canadian Entomologist* 122: 769–777. (bv).
- BORDEN, JOHN HARVEY, D. W. A. HUNT, D. R. MILLER, AND K. N. SLESSOR. 1956. Orientation in forest Coleoptera: an uncertain outcome of responses by individual beetles to variable stimuli. Pages 97–109 in T. L. Payne, M. C. Birch, and C. E. J. Kennedy, *Mechanisms in insect olfaction*. Clarendon Press, Oxford. 349 p. (bv).
- BORDEN, JOHN HARVEY, AND T. E. LACEY. 1955. Semiochemical-based manipulation of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins: a component of lodgepole pine silviculture in the Merritt timber supply area of British Columbia. *Journal of Applied Entomology* 99: 139–145. (bv cn).
- BORDEN, JOHN HARVEY, AND B. STAFFAN LINDGREN. 1958a. The role of semiochemicals in IPM of the mountain pine beetle. Pages 247–255 in T. L. Payne and H. Saarenmaa, *Integrated control of scolytid bark bee-*

- tles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn).
- _____. 1955b. The role of semiochemicals in integrated pest management of the mountain pine beetle (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 15: 414. (bv cn).
- BORDEN, JOHN HARVEY, A. M. PIERCE, H. D. PIERCE, JR., J. L. CHONG, A. J. STOCK, AND A. C. OEHLISCHLAGER. 1957. Semiochemicals produced by western balsam bark beetle, *Dryocoetes confusus* Swaine (Coleoptera: Scolytidae). Journal of Chemical Ecology 13: 823-836. (bv).
- BORDEN, JOHN HARVEY, LEE C. RYKER, L. J. CHONG, H. D. PIERCE, JR., B. D. JOHNSTON, AND A. C. OEHLISCHLAGER. 1957. Response of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae), to five semiochemicals in British Columbia lodgepole pine forests. Canadian Journal of Forest Research 17: 118-128. (bv).
- BORGES, P. A., AND A. R. M. SERRANO. 1959. New records of the coleopterous fauna (Insecta, Coleoptera) from the Azores. Boletim do Museu Municipal do Funchal 41, Art. 209. 24 p. (ds).
- BORSODI, D., T. GAL AND S. KISS. 1955. A betűzsol (Ips typographus L.) elleni vedekezes aggregacios feromonnal. Novevényvellem 21: 219. (bv).
- *BOSQ, JUAN M. 1934. [*Platypus plicatus*]. Boletín del Ministerio de Agricultura, Argentina 36(4): 335. ().
- BOUCEK, ZDENEK. 1980. A revision of the genus *Monacon* Waterston (Hymenoptera: Chalcidoidea: Perilampinae), parasites of ambrosia beetles (Coleoptera: Platypodidae). Bulletin of Entomological Research 70: 73-96. (ec).
- BOUCHOT L., F. LIEUTIER, AND D. DEBOUZIE. 1955a. Colonization dynamics of *Tomicus piniperda* L. and *Ips sexdentatus* Boern. (Coleoptera: Scolytidae) on Scots pine. International Congress of Entomology, Proceedings 18: 439. (bv).
- _____. 1955b. Spatial and temporal distribution of attacks by *Tomicus piniperda* L. and *Ips sexdentatus* Boern. (Coleoptera, Scolytidae) on *Pinus sylvestris*. Journal of Applied Entomology 106: 356-371. (bv).
- *BOURNE, A. I., AND WALTHAM WHITCOMB. 1937. Department of Entomology. Pages 46-57 in Annual report, 1936. Massachusetts Agricultural Experiment Station, Bulletin 339. ().
- BOUSEFIELD, WAYNE E., R. EDER, AND DAILE BENNETT. 1955. Users' guide and documentation for insect and disease survey. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 85-19. 19 p. (cn).
- BOUSEFIELD, WAYNE E., S. K. HAGLE, AND S. KOHLER. 1955. Montana forest pest conditions and program highlights, 1957. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Timber, Cooperative Forestry and Pest Management Report 85-2. 15 p. (cn).
- BOUTZ, GARY E., J. W. BREWER, AND J. N. BISHOP. 1985. Capture patterns of *Scolytus multistriatus* (Marsh.) (Col., Scolytidae) attracted to a pheromone-baited trap. Journal of Applied Entomology 99: 366-370. (bv cn).
- *BOWDITCH, F. C. 1896. List of Mt. Washington Coleoptera. Supplement to Psyche, II. 11 p. ().
- BOWERS, W. W., AND JOHN HARVEY BORDEN. 1990. Evidence for a male-produced aggregation pheromone in the four-eyed spruce bark beetle, *Polygraphus rufipennis* (Kirby) (Col., Scolytidae). Journal of Applied Entomology 111: 292-299. (bv).
- BOWERS, W. W., JOHN HARVEY BORDEN, AND G. GRIES. 1985. Aggregation pheromone of the four-eyed spruce bark beetle, *Polygraphus rufipennis* Kirby (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 18: 430.
- *BOYD, W. M. 1945. [*Scolytus rugulosus*]. New Jersey Department of Agriculture, Circular 355: 131-135. ().
- BRAND, J. M., AND F. SCOTT. 1985. Enzymatic stereospecificity and bark beetle pheromones. Fort Hare Papers 5(1): 52-54. 70. (bv ms).
- BRANHAM, SUSAN J., AND ROBERT CLIFFORD THATCHER. 1985. Integrated pest management symposium: proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 353 p. (cn ec).
- BRANHAM, SUSAN J., ROBERT CLIFFORD THATCHER, GARLAND N. MASON, AND GERRARD D. HERTEL. 1985. Integrated pest management in the South: highlights of a 5-year program. United States Department of Agriculture, Forest Service, Agriculture Information Bulletin 491. 19 p. (cn ms).
- *BRAUN, RUDOLF. 1949. Erfahrungen aus der Borkenkäfer bekämpfung im Schwarzkieferngebiet. Österreich Forst und Holzwirtschaft 3: 66-65. ().
- BRAUNER, K. 1955. Der Kupferstecher: Ein winziger, aber gefährlicher Forstschädling. Allgemeine Forstzeit-schrift 12: 255-259. (cn).
- BREWER, S. D., R. A. BECK, AND RICHARD A. ROEPER. 1955. Observations of the gallery habits of *Trypodendron retusum* (Coleoptera: Scolytidae) infesting aspen in central Michigan. Great Lakes Entomologist 21: 5-8. (lb).
- BREWER, WAYNE. 1978. Dutch elm disease: elm bark beetles and insects that weaken elms. Colorado State University Extension Service, Service in Action No. 5.506. 2 p. (cn).
- BRIDGES, JOHN ROBERT. 1985. Relationship of symbiotic fungi to southern pine beetle population trends. Pages 127-135 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 353 p. (ec).
- _____. 1987a. Effects of terpenoid compounds on growth of symbiotic fungi associated with southern pine beetle. Phytopathology 77: 83-85. (ec).
- _____. 1987b. Prevalence of *Unikaryon minutum* (Microsporidia: Nosematidae) infection in outbreak populations of the southern pine beetle (Coleoptera: Scolytidae). Journal of Invertebrate Pathology 49: 334-335. (ec).
- BRIDGES, JOHN ROBERT, AND JOHN CONRAD MOSER. 1986. Relationship of phoretic mites (Acari: Tarsenemidae) to the bluestaining fungus, *Ceratocystis minor*, in trees infested by southern pine beetle (Coleoptera: Scolytidae). Environmental Entomology 15: 951-953. (ec).
- BRIDGES, JOHN ROBERT, W. A. NETTLETON, AND M. D. CONNER. 1985. Southern pine beetle (Coleoptera: Scolytidae) infestations without the bluestain fungus, *Ceratocystis minor*. Journal of Economic Entomology 78: 325-327. (ec).
- BRIDGES, JOHN ROBERT, AND THIELMA J. PERRY. 1987. *Ceratocystiopsis ranaculosus* sp. n. associated with

- the southern pine beetle. *Mycologia* 79: 630–633. (cc).
- BRIDGES, JOHN ROBERT, W. T. THOENEY, AND A. E. TIARKS. 1988a. Techniques for studying bark beetle dispersal. Pages 307–319 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (hb ms).
- _____. 1988b. Techniques for studying bark beetle dispersal. International Congress of Entomology, Proceedings 15: 416. (hb ms).
- BRIGHT, DONALD EDWARD, JR. 1985a. New species and records of North American *Pityophthorus* (Coleoptera: Scolytidae), Part IV: the scriptor group. Great Basin Naturalist 45: 467–475. (tx).
- _____. 1985b. New species and records of North American *Pityophthorus* (Coleoptera: Scolytidae), Part V: the juglandis group. Great Basin Naturalist 45: 476–482. (tx).
- _____. 1985c. Studies on West Indian Scolytidae (Coleoptera), 3: checklist of Scolytidae of the West Indies with descriptions of new species and taxonomic notes. *Entomologischen Arbeiten aus dem Museum G. Frey* 33–34: 169–188. (tx).
- _____. 1986a. New species and new records of North American *Pityophthorus* (Coleoptera: Scolytidae), Part VI: the lantus group. Great Basin Naturalist 46: 641–645. (tx).
- _____. 1986b. New species and new records of North American *Pityophthorus* (Coleoptera: Scolytidae), Part VII. Great Basin Naturalist 46: 679–684. (tx).
- _____. 1987a. A review of the Scolytidae (Coleoptera) of the Azores with description of a new species of *Phloeosinus*. *Bocagiana* 107. 5 p. (ds tx).
- _____. 1987b. Notes on the occurrence of *Xyleborinus gracilis* (Eichhoff) in the United States (Coleoptera: Scolytidae). *Coleopterists Bulletin* 41: 335. (ds).
- _____. 1988a. Use of the aedeagus in the systematics of the Scolytidae (Coleoptera), with phylogenetic implications. International Congress of Entomology, Proceedings 15: 409. (ay tx).
- _____. 1988b. *Xylosandrus germanus* (Blandford), an ambrosia beetle new to Canada. Pages 93–94 in M. J. Sarazin, The Canadian agriculture insect pest review 66. 117 p. (ds).
- _____. 1989. Two new species of *Phloeosinus* Chapuis from Mount Kinabalu, Borneo, with taxonomic notes (Coleoptera: Scolytidae). *Coleopterists Bulletin* 43: 79–82. (tx).
- _____. 1990. A new species of *Liparthrum* from Borneo with notes on its generic placement (Coleoptera: Scolytidae). *Coleopterists Bulletin* 44: 485–488. (tx).
- _____. 1991a. Platypodidae (Curculionoidea). Page 616 in F. W. Stehr, Immature insects, Volume 2. Kendall Hunt, Dubuque, Iowa. 975 p. (hb tx).
- _____. 1991b. Scolytidae (Curculionoidea). Pages 613–616 in F. W. Stehr, Immature insects, Volume 2. Kendall Hunt, Dubuque, Iowa. 975 p. (hb tx).
- _____. 1991c. Studies in Xyleborini 2: review of the genus *Sampsonius* Eggers (Coleoptera: Scolytidae). *Studies on Neotropical Fauna and Environment* 26: 11–25. (ds tx).
- BRIGHT, DONALD EDWARD, JR. AND ROBERT E. SKIDMORE. 1991. Two new records of Scolytidae (Coleoptera) from Canada. *Coleopterists Bulletin* 45: 368. (ds).
- BROWN, K. W. 1961. Entomological studies from a high tower in Mpanga Forest, Uganda. XI. Observations on Coleoptera. Royal Entomological Society of London, Transactions 113: 353–355. (hb).
- BROWN, M. W., T. EVAN NEBECKER, AND C. R. HONEA. 1987. Thinning increases loblolly pine vigor and resistance to bark beetles. *Southern Journal of Applied Forestry* 11: 28–31. (cn).
- BROWN, WILLIAMSON JAMES. 1930. Coleoptera of the north slope of the Gulf of the St. Lawrence. *Canadian Entomologist* 62: 239–246. (ds tx).
- BROWNE, FRANCIS GEORGE. 1986a. Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, XIII. *Kontyo* 54: 89–99. (ds tx).
- _____. 1986b. Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, XIII. *Kontyo* 54: 333–343. (ds tx).
- _____. 1986c. Bark beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, XV. *Kontyo* 54: 661–671. (ds tx).
- BRUN, L. O., V. GAUDICHON, AND C. MARCILLAUD. 1989. Provisional method for detecting endosulfan resistance in coffee berry borer *Hypothenemus hampei* (Coleoptera, Scolytidae). *FAO Plant Protection Bulletin* 37: 125–129. (ay cn).
- BRUN, L. O., C. MARCILLAUD, V. GAUDICHON, AND D. M. SUCKLING. 1989. Endosulfan resistance in *Hypothenemus hampei* (Coleoptera: Scolytidae) in New Caledonia. *Journal of Economic Entomology* 82: 1311–1316. (cn).
- BRUNSON. 1929. [Note on *Scolytus rugulosus*]. *Southern Cultivator* 87: 11. (S. (cn)).
- BRYANT, C. H. 1985. Hazard rating systems and southern pine beetle prevention in Texas. Pages 480–484 in E. Shoulders (ed.), Third Biennial Southern Silvicultural Research Conference: proceedings, 7–8 November 1984, Atlanta, Georgia. United States Department of Agriculture, Forest Service, Southeastern Forest Experiment Station (Research Triangle Park, North Carolina), General Technical Report SO-54. (cn).
- BRYANT, C. H., H. A. PASE III, AND RONALD FORREST BILLINGS. 1982. Microcomputer aids southern pine beetle management in Texas. Pages 121–124 in Microcomputers: a new tool for foresters. Conference proceedings, 18–20 May 1982, Purdue University. Society of American Foresters, Publication 82-05. (cn).
- BUCK, F. D. 1952. Some coleopterous records from Hampstead Heath. *Entomological Gazette* 3: 101–103. (ds).
- _____. 1955b. A provisional list of the Coleoptera of Epping Forest. *Entomologist's Monthly Magazine* 91: 174–192. (ds).
- BUTH, P. N. 1989. Six new Danish roundworms (Nematoda) found in the gut of tree- and fungus-dwelling beetles. *Flora and Fauna* 95: 61–64. (cc).
- *BUISMANN, CHRISTINE J. 1930. The Dutch elm disease [abstract]. *Phytopathology* 20: 111–112. ().
- *_____. 1932d. Das grosse Ulmensterben. *Deutsche Forst. Zeit.* 47: 141–142. ().
- *_____. 1932e. De olmen of iepenziekte. *Antwerpen: Drukkerij K. Dirix van Riet, Muizenstrat* 19: 1–15. ().
- *_____. 1932f. Verslag van de phytopathologische onderzoekingen over de iepenziekte verricht in het Laboratorium "Willie Commeln Scholten."

- gedurende 1931. Tijdschrift over Plantenziekten 35: 17–36. ().
- °. 1933c. De iepenziekte. Tijdschrift der Nederlandsche Heideemaatschappij 45: 219–237. ().
- °. 1933d. Proeven over waterverduimping pij bladeren van verschillende iepensoorten. Tijdschrift over Plantenziekten 39: 35–41. ().
- °. 1934b. De iepenziekte in America. Nederlandsche Boschbouw Tijdschrift 7: 439–440. ().
- °. 1934c. The area of distribution of the *Ceratostomella* (*Graphium*) elm disease. Mededeelingen van het phytopathologisch Laboratorium "Willie Commelin Scholten," Baarn 13: 35–46. ().
- °. 1935a. Het iepenziekte probleem in Italie. Nederlandsche Boschbouw-Tijdschrift 8: 179–180. ().
- °. 1935b. Over de iepenziekte buiten Nederland. Natuur en techniek, 7 July. ().
- °. 1935c. Verslag over de onderzoeken betreffende de iepenziekte, verricht in het Phytopathologisch Laboratorium "Willie Commelin Scholten" de Baarn, gedurende 1934. Tijdschrift over Plantenziekten 41(5): 104–120. ().
- °. 1936b. Stand van de iepenziekte in Amerika volgens de proceedeances van de Ile Shade Tree Conference (August 1935) 9: 77–80. ().
- °. 1936c. Verlag van de onderzoeken over de iepenziekte, verriicht in hat Phytopathologisch Laboratorium Willie Commelin Scholten de Baarn, dedurende 1935. Tijdschrift over Plantenziekten 42: 21–44. ().
- ° BUISMANN, CHRISTINE J., JOHANNA WESTERDIJK, AND S. G. A. DOORENBOS. 1932. Wat kunnen de nederlandse boomkweekers doen in verband met de iepenziekte? Tijdschrift over Plantenziekten 35: 37–40. ().
- BURKE, HORACE R., AND DONALD MORGAN ANDERSON. 1976. Systematics of larvae and pupae of American Curculionoidea: status report, historical review and bibliography. Southern Entomologist 1: 56–73. (tx ms).
- BURKHART, H. E., H. L. HANEY, JR., J. D. NEWBERRY, W. A. LEUSCHNER, C. L. MORRIS, AND D. D. REED. 1986. Evaluation of thinning for reduction of losses from southern pine beetle attack in loblolly pine stands. Southern Journal of Applied Forestry 10: 105–108. (cn).
- BURKS, BERNARD DEWITT. 1973. North American species of *Calosota* Curtis (Hymenoptera: Enpeluidae). Washington Academy of Science, Journal 63: 26–31. (ec).
- BURRELL, A. G., L. MAGNANO, A. R. MAGNANO, C. SCALA, AND B. BACCETTI. 1986. Spermatozoa and phylogeny in Curculionoidea superfamily. Pages 69–73 in M. Cresti and R. Dallai (eds.), Biology of reproduction and cell motility in plants and animals. (ay tx).
- . 1988. Spermatozoa and phylogeny of Curculionoidea (Coleoptera). International Journal of Insect Morphology and Embryology 17: 1–50. (ay tx).
- BUSH, P. B., J. W. TAYLOR, C. K. MCMAHON, AND D. G. NEARY. 1987. Residues of lindane and chlorpyrifos in firewood and woodsmoke. Journal of Entomological Science 22: 131–139. (cn).
- BYCILAWSKA, S. 1983. Występowanie wazniejszych kambio-1 Ksylofagow sosny w drzewostanach uszkodzonych przez Huragany. Sylwan 127: 45–52. (ec).
- BYERS, GEORGE WILLIAM, AND JAY B. KARREN. 1968. Catalogue of the types in the Snow Entomological Museum, Part VI (Coleoptera). University of Kansas Science Bulletin 45: 1–20. (tx).
- BYERS, JOHN ALLEN. 1957. Interactions of pheromone component odor plumes of western pine beetle. Journal of Chemical Ecology 13: 2143–2153. (bv).
- . 1958a. Novel diffusion-dilution method for release of semiochemicals: testing pheromone component ratios on western pine beetle. Journal of Chemical Ecology 14: 199–212. (bv ms).
- . 1958b. Upwind flight orientation to pheromone in western pine beetle tested with rotating windvane traps. Journal of Chemical Ecology 14: 189–195. (bv).
- . 1958c. Host tree susceptibility, suitability, and unsuitability recognized by some European bark beetles. International Congress of Entomology, Proceedings 18: 429. (bv).
- . 1959a. Behavioral mechanisms involved in reducing competition in bark beetles. Holarctic Ecology 12: 446–476. (bv ec).
- . 1959b. Chemical ecology and bark beetles. Experimentia (Basel) 45: 271–283. (bv ec).
- BYERS, JOHN ALLEN, O. ANDERBRANT, AND J. LOFQVIST. 1959. Effective attraction radius: a method for comparing species attractants and determining densities of flying insects. Journal of Chemical Ecology 15: 749–766. (bv ec).
- BYERS, JOHN ALLEN, GORAN BIRGERSSON, J. LOFQVIST, AND GUNNAR BERGSTROM. 1955. Synergistic pheromones and monoterpenes enable aggregation and host recognition by a bark beetle. Naturwissenschaften 75: 153–155. (bv).
- BYERS, JOHN ALLEN, H. E. HOGBERG, C. R. UNELIUS, GORAN BIRGERSSON, AND J. LOFQVIST. 1959. Structure-activity studies on aggregation pheromone components of *Pityogenes chalcographus* (Coleoptera: Scolytidae): all stereoisomers of chalcogran and methyl-2 4-decadienoate. Journal of Chemical Ecology 15: 655–696. (bv).
- BYERS, JOHN ALLEN, B. S. LAMME, AND J. LOFQVIST. 1959. Host tree unsuitability recognized by pine shoot beetles in flight. Experimentia (Basel) 45: 489–492. (bv).
- BYERS, JOHN ALLEN, B. S. LAMME, J. LOFQVIST, F. SCHLTER, AND GUNNAR BERGSTROM. 1955. Olfactory recognition of host-tree susceptibility by pine shoot beetles. Naturwissenschaften 72: 324–326. (bv).
- BYERS, JOHN ALLEN, AND J. LOFQVIST. 1959. Flight initiation and survival in the bark beetle *Ips typographus* (Coleoptera: Scolytidae) during the spring dispersal. Holarctic Ecology 12(4): 432–440. (hb).
- BYKOV, A. A. 1957. Characteristics of the reproduction of bark beetles in pine plantations [In Russian]. Zashchita Rastenii, Moscow 3: 35. (hb).
- BYLER, JAMES W., WAYNE E. BOUSFIELD, AND S. KNIOLER. 1986. Montana forest pest conditions and program highlights, 1985. United States Department of Agriculture, Forest Service, Northern Region (Missoula, Montana), Cooperative Forestry and Pest Management Report 86–2, 19 p. (cn).

C

- CAHILL, DONN B. 1967. Bark beetle evaluation, Bighorn National Forest. United States Department of Agriculture, Forest Service, Washington, D.C. [1967?]. 3 p. (cn).
- CALL, RICHARD J. 1959. Case history: Northern Region Forest Service Eastside Zone-Mountain pine beetle considerations. Pages 41–42 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest

- Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 199 p. (cn).
- CALTRELL, R. M. 1969. Annual district report: Grande Prairie-Peace River District, 1968. Pages 55–63 in Annual reports of the Forest Insect and Disease Survey: Alberta-Northwest Territories-Yukon Region, 1968. Canada Department of Fisheries and Forestry, Canadian Forest Service, Forest Research Laboratory (Calgary, Alberta), Information Report A-X-22. (cn).
- _____. 1970. Annual district report: Grande Prairie-Peace River District, 1969. Pages 36–42 in Annual reports of the Forest Insect and Disease Survey: Alberta-Northwest Territories-Yukon Region, 1969. Canada Department of Fisheries and Forestry, Canadian Forestry Service, Forest Research Laboratory (Calgary, Alberta), Information Report A-X-30. (cn).
- CAMBI, G. 1915. Il "punternoio" e il "fleotripide" dell'olivo (*Phloeotribus scarabaeoides* e *Phloeotribus oleae*) nel chianti. *Revista di Agricoltura* 35: 601. (ec).
- CAMERON, DAWN. 1989. Biological evaluation of spruce beetle populations, Curtis Creek Area, Ogden RD, Wasatch-Cache NF, United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-S9-3. 3 p. (cn).
- CAMERON, M., AND A. C. GATTO. 1907. A list of the Coleoptera of the Maltese Islands. *Entomological Society of London, Transactions* 55: 383–403. (ds).
- CAMERON, R. SCOTT. 1987a. Southern pine bark beetles in the urban environment. *Journal of Arboriculture* 13: 145–151. (bv cn).
- _____. 1987b. Southern pine bark beetles in the urban environment. Texas Forest Service, Circular 274. 7 p. (cn).
- CAMERON, R. SCOTT, AND RONALD FORREST BILLINGS. 1988. Southern pine beetle: factors associated with spot occurrence and spread in young plantations. *Southern Journal of Applied Forestry* 12: 205–214. (ec).
- CANE, JAMES H., MOLLY WILDORF STOCK, DAVID LEE WOOD, AND S. J. GAST. 1990. Phylogenetic relationships of *Ips* bark beetles (Coleoptera: Scolytidae): electrophoretic and morphometric analysis of the *grandicollis* group. *Biochemical Systematics and Ecology* 18: 359–365. (ay tx).
- CANE, JAMES H., AND DAVID LEE WOOD. 1988. Phylogenetic divergence of semiochemical communications of *Ips* pine bark beetles (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 18: 409. (bv tx).
- CANNON, WILLIAM N., JR. 1955. Gallery construction and oviposition of *Scolytus multistriatus* (Coleoptera: Scolytidae) in relation to temperature and adult density. *Environmental Entomology* 15: 641–643. (bv ec bb).
- CANNON, WILLIAM N., JR., JACK H. BARGER, AND L. L. GROTH. 1955. Seasonal detection of visible Dutch elm disease symptoms. *Journal of Arboriculture* 11: 233–235. (cn).
- CANNON, WILLIAM N., JR., JACK H. BARGER, AND DAVID P. WORLEY. 1950. Dutch elm disease control: performance and cost. United States Department of Agriculture, Forest Service, Northeastern Forest Experiment Station, Research Paper NE-457. 8 p. (cn).
- _____. 1952. Dutch elm disease control: economics of girdling diseased elms to improve sanitation performance. *Journal of Arboriculture* 8: 129–135. (ec).
- CAPEK, MIROSLAV. 1957. Lumcikovité (Hymenoptera: Braconidae) ako parazitoídy podkorných a drevo-kazných škodcov duba - vektorov tracheomykoz [The braconids as parasitoids of oak bark-mining and wood-boring insects—carriers of tracheomycoses] [English summary]. *Biologia (Bratislava)* 41: 535–542. (ec).
- *CAPPELLI, M., AND L. MASUTTI. 1965. Note preliminari sul deperimento dell'abete rosso nella stazione di Foresta (Merano). Gruppo giornalistico dell'edagricole Bologna, XVI, 5: 29–49. ().
- CARLE, PIERRE R., AND DANIEL SCHWESTER. 1975. Perspectives d'avenir du pin maritime en provenance. *Revue Forestière Française* 27: 339–350. (cn ec).
- _____. 1983. Pyrethroids and integrated pest control against xylophagous forest insects. Pages 113–118 of volume 2 in International conference on integrated plant protection: proceedings, 4–9 July 1983, Budapest. (cn).
- CARPENTER, STEVEN E. 1985. *Ambrodiscus*, a new genus of inoperculate discomycetes from ambrosia beetle galleries. *Mycologia* 80: 320–323. (ec).
- CARPENTER, STEVEN E., MARK E. HARMON, ELAINE R. INGHAM, RICK G. KELSEY, JOHN D. LATTIN, AND TIMOTHY D. SCHOWALTER. 1988. Early patterns of heterotroph activity in conifer logs. *Royal Society of Edinburgh* 94S: 33–43. (ec).
- *CEIANU, I. 1966. Observatii asupra formarii entomofaunei in culturile forestiere noi din zona de Cimpie. Pages 267–287 in Ministerul Invatamintului, Cercetari de Ecologie Animala. ().
- CEREZKE, HERBERT F. 1989. Mountain pine beetle aggregation semiochemical use in Alberta and Saskatchewan, 1983–1987. Pages 108–113 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (bv cn).
- CEREZKE, HERBERT F., AND F. J. EMOND. 1989. Forest insect and disease conditions in Alberta, Saskatchewan, Manitoba, and the Northwest Territories in 1987. Government of Canada, Forestry Service, Northern Forestry Centre, Edmonton, Alberta, Information Report NOR-X-300. 25 p. (cn).
- CHAMPION, GEORGE CHARLES. 1917. *Pediacus depressus* Herbst, a species frequenting pines in the working district [*Hylastes ater*, *Hylastes palliatus*, *Myelophilus piniperda*, *Tomicus laricis*]. *Entomologists' Monthly Magazine* 53: 173–174. (ec).
- CHIAN, H. T., U. RAZANI, AND F. E. PUTZ. 1984. A preliminary study on planting of *Rhizophora* species in an avicennia forest at the Matang mangroves. Pages 340–345 in S. Soemodihardjo et al., *Ekosistem Mangrove: seminar proceedings*. Jakarta, Indonesia: Lembaga Ilmu Pengetahuan Indonesia. (cn).
- *CHANDLER, STEWART C. 1913. [*Scolytus rugulosus*]. Missouri State Board of Agriculture, Special Bulletin 57. 53 p. ().
- *_____. 1950. [*Phloeotribus liminaris*, *Scolytus rugulosus*]. Illinois Natural History Survey, Circular 43: 18–19. ().
- CHANDRA, AVINASHI, AND S. I. AHMAD. 1955. Review of current status of taxonomy of the family Scolytidae in India. *Indian Forester* 111: 622–629. (tx).
- CHAPIN, JOAN B., AND A. D. OLIVER. 1986. New records for

- Xylosandrus* and *Xyleborus* species (Coleoptera: Scolytidae). Entomological Society of Washington, Proceedings 55: 650-653. (ds).
- *CHARRAS, CONSTANTIN. 1950. Etude ecophysologique des Scolytidae parasites des conifères. I. Les Scolytidae spécifique de *Cedrus atlantica* [Physiological study of scolytid pests of conifers. I. Host-specific Scolytidae of *Cedrus atlantica*]. Recherche Forestieres au Maroc, Annales 20: 205-267. ().
- _____. 1956. Selection de la plante-hôte par les Scolytidae et mecanismes d'installation des insectes (attraction primaire et attraction secondaire) [Selection of the host-plant by Scolytidae and installation mechanisms (primary and secondary attraction)]. Societe Entomologique de France 91: 137-162. (bv).
- CHAUDHRY, GHULAM ULLAH, M. ISMAIL CHAUDHRY, AND NASRULLAH KHAN MALIK. 1970. Survey of insect fauna of Pakistan. Vol. 2. Biological Science Research Division, Forest Entomology Branch, Pakistan Forest Institute, Peshawar. 205 p. (cn ds).
- CHEN, N. M., AND JOHN HARVEY BORDEN. 1959. Adverse effect of fenoxycarb on reproduction by the California five-spined ips, *Ips paraconfusus* Lanier (Coleoptera: Scolytidae). Canadian Entomologist 121: 1059-1065. (ec hb).
- CHEN, N. M., JOHN HARVEY BORDEN, AND H. D. PIERCE, JR. 1955. Effect of juvenile hormone analog fenoxycarb on pheromone production by *Ips paraconfusus* (Coleoptera: Scolytidae). Journal of Chemical Ecology 14: 1057-1095. (bv).
- CHENG, J. D. 1959. Streamflow changes after clear-cut logging of a pine beetle infested watershed in southern British Columbia, Canada. Water Resources Research 25: 449-456. (cn ec).
- CHENICLET, C., C. BERNARD-DAGAN, AND G. PAULY. 1955. Terpene biosynthesis under pathological conditions. Pages 117-130 in W. J. Mattson, N. Levieux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (cn).
- CHENIER, J. V. R., AND B. J. R. PHILOGENE. 1959a. Evaluation of three trap designs for the capture of conifer-feeding beetles and other forest Coleoptera. Canadian Entomologist 121: 159-167. (cn ms).
- _____. 1959b. Field responses of certain forest Coleoptera to conifer monoterpenes and ethanol. Journal of Chemical Ecology 15: 1729-1745. (bv).
- *CHITTENDEN, FRANK HURLBUT. 1905. [*Scolytus rugulosus*]. United States Department of Agriculture, Yearbook 1905: 346-347, fig. 85. ().
- CHONG, J. MICHAEL, AND EDUARDO K. MAR. 1959. Enantioselective syntheses of *endo*- and *exo*-brevicomins via alpha-alkoxytannanes. Tetrahedron 45: 7709-7716. (bv ms).
- CHOO, HO YUL, H. K. KAYA, AND P. SHEA. 1959. *Parasitylenchus orthotomici* new species (Tylenchida: Allantonematidae) from *Orthotomicus angulatus* (Coleoptera: Scolytidae) with note on parasitism. Korean Journal of Applied Entomology 25: 1-3. (ec).
- CHOO, HO YUL, H. K. KAYA, P. SHEA, AND E. M. NOFFSINGER. 1957. Ecological study of nematode parasitism in *Ips* beetles from California and Idaho. Journal of Nematology 19: 495-502. (ec).
- CHOO, HO YUL, AND K. S. WOO. 1955. A list of Korean bark and ambrosia beetles, and their host plants. Korean Journal of Plant Protection 24: 163-167. (ds).
- _____. 1959a. Four new species of Scolytidae (Coleoptera) from Korea. Korean Journal of Applied Entomology 25: 57-60. (tx).
- _____. 1959b. Supplementary notes on the bark and ambrosia beetles of Korea. Korean Journal of Applied Entomology 25: 4-9. (ds).
- CHOO, HO YUL, K. S. WOO, AND AKIRA NOBUCHI. 1955a. Classification of Korean bark and ambrosia beetles by their galleries [In Korean, English summary]. Korean Journal of Applied Entomology 27: 131-137. (hb ds).
- _____. 1955b. Key to the subfamilies, tribes and genera of Korean Platypodidae and Scolytidae (Coleoptera) [In Korean, English summary]. Korean Journal of Applied Entomology 27: 135-143. (tx).
- CHOO, HO YUL, K. S. WOO, AND C. D. PARK. 1955. Scolytid beetles, including two unrecorded species attracted to an attractant. Korean Journal of Applied Entomology 27: 190-193. (bv ds).
- *CHORBADZHIVO, P. 1926c. [*Scolytus rugulosus*]. Rapp. Ann. Sta. Agron. Etat Sofia. Sofia 1926: 175-241. ().
- CHOW, YIEN S., Y. CHEN, R. S. TSAI, AND Y. W. MENG. 1955. Cross attractions of the pine beetle pheromone in Taiwan. Institute of Zoology, Academia Sinica, Bulletin 27(1): 67-72. (bv).
- CHRISTIANSEN, ERIK. 1954. Bartraernes forsvær mot barkbilleangrep [In Norwegian]. Naturen 1954(2): 43-48. (cn ec).
- _____. 1955a. *Ceratocystis polonica* inoculated in Norway spruce: blue-staining in relation to inoculum density, sesinosis and tree growth. European Journal of Forest Pathology 15: 160-167. (cn ec).
- _____. 1955b. *Ips/Ceratocystis*-infection of Norway spruce: what is a deadly dosage? Journal of Applied Entomology 99: 6-11. (bv cn ec).
- _____. 1955. Influence of Norway spruce trees against the Eurasian spruce bark beetle, *Ips typographus*, and its symbiotic fungi. International Congress of Entomology. Proceedings 15: 439. (ec).
- _____. 1959. Bark beetles and air pollution. Meddelelser fra det Norsk Institutt for Skogforskning 42: 101-107. (ec).
- CHRISTIANSEN, ERIK, AND ALF BAKKE. 1955a. The spruce bark beetle of Eurasia. Pages 479-503 in A. A. Berryman, Dynamics of forest insect populations: patterns, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec).
- _____. 1955b. Granbarkbiller. Hva nyere forskning har lært oss [The spruce bark beetle *Ips typographus*: a review of our present knowledge] [In Norwegian, English summary]. Norsk Institutt for Skogforskning, Rapport 4. 21 p. (bv cn ec).
- CHRISTIANSEN, ERIK, AND ANDERS ERICSSON. 1956. Starch reserves in *Picea abies* in relation to defense reaction against a bark beetle transmitted blue-stain fungus, *Ceratocystis polonica*. Canadian Journal of Forest Research 16: 78-83. (ec).
- CHRISTIANSEN, ERIK, RICHARD H. WARING, AND ALAN ANDREW BERRYMAN. 1957. Resistance of conifers to bark beetle attack: searching for general relationships. Forest Ecology and Management 22: 95-106. (cn ec).
- CHU, YAU-L AND MEI-LIN HSIAO. 1951. A tentative bibliography of Formosan Coleoptera (1967-1980) [Scolytidae-Platypodidae, p. 86-89]. Pathologist and Entomologist, N. T. U. [Taiwan]. (ds).
- CHURCHER, J. JOSEPH. 1955. Detection of spruce beetle (*Dendroctonus rufipennis*) infestation using aerial photographs. Forestry Abstracts 46: 635. (cn).

- CIBRIAN TOVAR, DAVID B. II, EBEL, HARRY O. YATES, III, AND J. JULIO MENDEZ MONTIEL. 1956. Insectes de conos y semillas de los coníferas de México [Cone and seed insects of Mexican conifers]. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station (Research Triangle Park, North Carolina), General Technical Report. 110 p. (cn ds hb).
- CIESLA, WILLIAM M. 1988. Pine bark beetles: a new pest management challenge for Chilean foresters. *Journal of Forestry* 86: 27–31. (cn ds).
- CIESLA, WILLIAM M., MARK D. MCGREGOR, AND H. E. MEYER. 1971. Status of bark beetle infestations near Blue Mountain, Missoula District, Lolo National Forest, Montana. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Report 71–6. 4 p. (cn).
- CIESLA, WILLIAM M., AND OSVALDO RAMIREZ GREZ. 1988. *Hylurgus ligniperla* (Fabricius) (Coleoptera: Scolytidae) [In Spanish]. Protección Fitosanitaria Forestal, Corporación Nacional Forestal, Folleto de Divulgación Año 5, Santiago, Chile, No. 15. 7 p. (cn hb).
- CLARKE, L. J., AND G. C. CAREW. 1984b. Forecast of forest insect and disease conditions in Newfoundland and Labrador in 1984. Woody Points. Government of Canada Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland 13(4): 1–4. (cn).
- _____. 1985. Forest insect and disease conditions in Newfoundland and Labrador in 1984. Government of Canada. Canadian Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland, Information Report N-X-229. 31 p. (cn).
- _____. 1986. Forest insect and disease conditions in Newfoundland and Labrador in 1985. Government of Canada. Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland, Information Report N-X-21. 33 p. (cn).
- _____. 1987. Forest insect and disease conditions in Newfoundland and Labrador in 1986. Government of Canada. Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland, Information Report N-X-259. 36 p. (cn).
- _____. 1988. Forest insect and disease conditions in Newfoundland and Labrador in 1987. Government of Canada. Canadian Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland, Information Report N-X-265. 37 p. (cn).
- CLAYDON, N., J. F. GROVE, AND M. POPE. 1985. Elm bark beetle boring and feeding deterrents from *Phomopsis oblonga*. *Phytochemistry* 24: 937–943. (cn ec).
- COBB, FIELDS W., JR. 1988. *Leptographium wageneri*, cause of black-stain root disease: a review of its discovery, occurrence and biology with emphasis on pinyon and ponderosa pine. Pages 41–62 in T. C. Harrington and F. W. Cobb, Jr. (eds.), *Leptographium* root diseases on conifers. American Phytopathological Society Press, St. Paul, Minnesota. (ec).
- COBOS SUAREZ, J. M., AND E. MARTIN BERNAL. 1987. Metodos para la determinación del periodo de vuelo de los coleopteros escolitidos. *Información Técnica Económica Agraria* [Spain] 75: 65–72. (hb).
- COHEN, T., AND J. MATZ. 1957. Method of synthesizing brevicomin and using same in beetle control. U.S. Patent–4695584. 22 September 1957. Official Gazetteer, U.S. Patent Trademark Office 1084(4): 2045. (cn ms).
- COLE, DENNIS M. 1989. Preventive strategies for lodgepole pine/mountain pine beetle problems: opportunities with immature stands. Pages 64–69 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- *COLE, DENNIS M., AND MARK D. MCGREGOR. 1988. Stand culture/bark beetle relationships of immature tree stands in the inland Mountain West. Pages 278–285 in *Future forests of the Mountain West: a stand culture symposium*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report 243 p. ().
- COLE, WALTER ECKLE, GENE DOYLE AMMAN, AND CHESTER E. JENSEN. 1955. Mountain pine beetle dynamics in lodgepole pine forests. Part III. Sampling and modeling of mountain pine beetle populations. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-188. 46 p. (cn).
- COLE, WALTER ECKLE, AND MARK D. MCGREGOR. 1985. Reducing or preventing mountain pine beetle outbreaks in lodgepole pine stands by selective cutting. Pages 175–186 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service, and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1983. 239 p. (cn).
- COLLING, GENE F. 1955. Stalking a forest killer. United States Department of Agriculture, Forest Service, Intermountain Research Station and Northern Region, Ogden, Utah. Video tape, color, three-quarter or half inch. 19 minutes. (cn).
- *COLLINGS. 1915. [*Hylastes ater*]. *Journal of the Board of Agriculture*, London 22(8): 789–791. ().
- COLLINS, DONALD LEWIS. 1938. Feeding habits of *Scolytus multistriatus* Marsham with reference to Dutch elm disease. *Journal of Economic Entomology* 31: 196–200. (hb).
- *COLPI, C., AND L. MASUTTI. 1984. Reperti sull'entomofauna epigea di popolamenti di *Abies viridis* (Chaix) D. C. nel parco naturale di Paneveggio-Pale di S. Martino (Dolomiti Trentine). *Studi Trentini di Scienze Naturali, Acta Biologica*. Trento 61: 197–237. ().
- COOK, STEPHEN P., AND FRED PAUL HAIN. 1985a. Comparison of loblolly and shortleaf pine bolts as hosts of the southern pine beetle, *Dendroctonus frontalis* Zimmermann (Coleoptera: Scolytidae). *Environmental Entomology* 14: 332–335. (bv hb).
- _____. 1985b. Qualitative examination of the hypersensitive response of loblolly pine, *Pinus taeda* L., inoculated with two fungal associates of the southern pine beetle, *Dendroctonus frontalis* Zimmermann (Coleoptera: Scolytidae). *Environmental Entomology* 14: 396–400. (cn ec).
- _____. 1986. Defensive mechanisms of loblolly pine and shortleaf pine against attack by southern pine beetle, *Dendroctonus frontalis* Zimmermann, and its fungal associate, *Ceratocystis minor* (Hedgecock) Hunt. *Journal of Chemical Ecology* 12: 1397–1406. (ec).
- _____. 1987a. Four parameters of the wound response of loblolly and shortleaf pines to inoculation with the blue-staining fungus associated with the southern pine beetle. *Canadian Journal of Botany* 65: 2403–2409. (ec).

- _____. 1957b. Susceptibility of trees to southern pine beetle, *Dendroctonus frontalis* (Coleoptera: Scolytidae). *Environmental Entomology* 16: 9–14. (ec).
- _____. 1958a. Resistance mechanisms of loblolly and shortleaf pines to southern pine beetle attack. Pages 295–304 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (ec).
- _____. 1958b. Toxicity of host monoterpenes to *Dendroctonus frontalis* and *Ips typographus* (Coleoptera: Scolytidae). *Journal of Entomological Science* 23: 287–292. (cn).
- _____. 1958c. Wound responses of loblolly and shortleaf pines attacked or reattacked by *Dendroctonus frontalis* Zimmermann (Coleoptera: Scolytidae) or its fungal associate, *Ceratocystis minor* (Hedgecock) Hunt. *Canadian Journal of Forest Research* 18: 33–37. (cn ec).
- _____. 1959. Within-tree population parameters of *Dendroctonus frontalis* (Coleoptera: Scolytidae) in nylon mesh-enclosed *Pinus chinata*. *Environmental Entomology* 18: 1025–1031. (hb).
- COOK, STEPHEN P., FRED PAUL HAIN, AND P. B. NAPPEN. 1986. Seasonality of the hypersensitive response by loblolly and shortleaf pines to inoculation with a fungal associate of the southern pine beetle (Coleoptera: Scolytidae). *Journal of Entomological Science* 21: 283–285. (ec).
- COPE, R., D. BRUSS, T. SCHWARTZ, B. P. MUNDY, AND K. LIPKOWITZ. 1987. Studies on bicyclic ketals: a tandem oxymercuration-solvolmercuration approach to natural products. American Chemical Society, Abstracts of Papers. 193. (ms).
- CORREA LOPEZ, ALFONSO DE JESUS. 1989. Genitales de las especies Mexicanas del genero *Ips* DeGeer (Coleoptera: Scolytidae) como una herramienta en la identificacion. Published thesis, Colegio de Postgraduados, Institucion de Enseñanza e Investigacion en Ciencias Agricolas, Centro de Entomologia y Acarologia, Chapingo, Mexico. 65 p. (ay tx).
- *CORY, EARNEST NEAL. 1923. [*Scolytus rugulosus*]. Maryland State Agricultural Society, Report 6: 160 (1921–1922). ().
- COSTA, TANIA C. S. VILLACORTA, AND A. VILLACORTA. 1989. Modelo acumulativo para *Hypothenemus hampei* (Ferrari, 1867) (Coleoptera: Scolytidae) com base em suas exigencias termicas [A summation model for *Hypothenemus hampei* based on its thermal unit requirements]. *Anais de Sociedade Entomologica do Brasil* 18(supplement): 91–99. (ec hb).
- COSTA LIMA, ANGELO MOREIRA DA. 1962. Insectos do Brasil. Vol. 12: Hymenopteros, Part 2. Escola Nacional de Agronomia, Serie didatica No. 14, Servico Grafico. B. G. E., Rio de Janeiro. 393 p., 141 figs. (ec).
- *COTTA. 1919. [*Hylastes ater*]. L'Italia Agricola, Piacenza 5(1–3): 10–16, 70–80. ().
- COUDROY, J. P. 1981a. Caracteristiques comparees des systemes de ponts des trois especes de *Pityokteines* Fuchs (Coleopteres, Scolytides) en Vallee d'Ossau (Pyrenees Atlantiques): *P. curvidens* Germar, *P. spinidens* Reitter et *P. vorontzovi* Jacobson. Documents d'Ecologie Pyreneenne 2: 45–57. (hb).
- _____. 1981b. Contribution a l'etude ecologique et biocenotique des insectes xylophages du sapin (*Abies alba*) en Vallee d'Ossau, Pyrenees Atlantiques. Documents d'Ecologie Pyreneenne 2: 75–78. ().
- _____. 1982. La biocenose des insectes xylophages du pin a crachets (*Pinus uncinata* Ram) dans la reserve du Neouvielle (H. P.) et le massif du Pic d'Anie (P. A.). *Acta Biologica Montana* 2/3: 77–86. ().
- *_____. 1984. Enchaînement biocenotique des insectes xylophages du sapin (*Abies alba* L.) en Vallee d'Ossau: descriptions des deux premiers groupes d'especes. Documents d'Ecologie Pyreneenne 3/4: 97–100. ().
- COULSON, ROBERT N., GENE DOYLE AMMAN, DONALD L. DAHLSTEN, C. J. DEMARS, JR. AND F. M. STEPHEN. 1985. Forest-bark beetle interactions: bark beetle population dynamics. Pages 61–80 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. 256 p. (cn ec).
- COULSON, ROBERT N., R. O. FLAMM, PAUL E. PULLEY, THOMAS LEE PAYNE, E. J. RYKIEL, AND T. L. WAGNER. 1986. Response of the southern pine bark beetle guild (Coleoptera: Scolytidae) to host disturbance. *Environmental Entomology* 15: 850–858. (bv).
- COULSON, ROBERT N., R. O. FLAMM, T. L. WAGNER, E. J. RYKIEL, P. S. H. SHARPE, THOMAS LEE PAYNE, AND S. K. LIN. 1985. Population dynamics of initiation and growth of southern pine beetle infestations. Pages 136–153 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report (SO-56. 383 p. (cn hb).
- COULSON, ROBERT N., L. A. GRAHAM, AND C. N. LOVELADY. 1985a. Intelligent geographic information system for predicting the distribution, abundance, and location of southern pine beetle infestations in forest landscapes. Pages 283–294 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- _____. 1985b. Intelligent geographic information system for predicting bark beetle infestations. *International Congress of Entomology, Proceedings* 18: 415. (cn).
- COULSON, ROBERT N., M. C. SAUNDERS, D. K. LOH, F. L. OLIVERIA, D. DRUMMOND, P. J. BARRY, AND K. M. SWAN. 1989. Knowledge system environment for integrated pest management in forest landscapes: the southern pine beetle (Coleoptera: Scolytidae). *Entomological Society of America, Bulletin* 35(2): 26–32. (cn).
- COULSON, ROBERT N., M. C. SAUNDERS, K. D. LOH, E. J. RYKIEL, THOMAS LEE PAYNE, PAUL E. PULLEY, AND L. C. HU. 1985. A decision support system for the southern pine beetle. Pages 35–46 in R. A. Goyer and J. P. Jones, Insects and diseases of southern forests. 34th annual forestry symposium, Division of Continuing Education, Louisiana State University, Baton Rouge, Louisiana. (cn).
- COULSON, ROBERT N., M. C. SAUNDERS, THOMAS LEE PAYNE, R. O. FLAMM, T. L. WAGNER, P. B. HENNIER, AND E. J. RYKIEL. 1985. A conceptual model of the role of lightning in the epidemiology of the southern pine beetle. Pages 136–146 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Canadian Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada. September 1983. 240 p. (ec).
- COULSON, ROBERT N., AND J. A. WITTER. 1984. Forest ento-

- mology: ecology and management. John Wiley & Sons, New York. x + 669 p. (cn hb).
- COULSON, ROBERT N., et al. 1989. An overview of the TAMBEETLE model of *Dendroctonus frontalis* population dynamics. *Holarctic Ecology* 12: 445–450. (hb cn).
- COVASSI, MARCO. 1983. Sulla presenza dello *Scolytus kirschi* Skal. an *Ulmus carpiniifolia* Gled in Toscana (Coleoptera: Scolytidae). Pages 135–136 in Atti XII Congresso Nazionale Italiano de Entomologia, Roma, 5–9 Novembre 1980. Volume II. Accademia Nazionale Italiana de Entomologia, Roma. (ds).
- *COVASSI, MARCO, AND L. MASUTTI. 1978. Aperçu sur les principaux problèmes posés par les insectes forestiers en Italie. Estratto da Annali dell Istituto sperimentale per la Zoologia Agraria 6(1978–1979): 27–32. ().
- *_____. 1980. Generalità sull entomofauna degli olmi, con particolare riguardo ai coleotteri scolidi vettori della grafiosi. Estratto da Informatore Fitopatologico 1(30): 19–26. ().
- CRANSHAW, WHITNEY S., AND DAVID LEATHERMAN. 198.. Ips beetles: characteristics and control. Colorado State University Cooperative Extension Service, Service in Action [Circular] No. 5.55S. (cn).
- CRESSAP, VERNON L. M. 1977. Mountain pine beetle: Salida and San Carlos Ranger Districts, Pike and San Isabel National Forests, Colorado. United States Department of Agriculture, Forest Service, Rocky Mountain Region (Lakewood, Colorado), Biological Evaluation R2–77–7. 3 p. (cn).
- CROOKSTON, NICHOLAS LEE II, AND RONALD WILLIAM STARK. 1985. Forest-bark beetle interactions: stand dynamics and prognoses. Pages 81–103 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. 256 p. (cn ec).
- CROSBY, DAVID, AND DONALD J. CURTIS. 1970. Forest insect and disease conditions in Alaska during 1969. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska. 15 p. (cn).
- CROSS, EARLE ALBRIGHT, AND JOHN CONRAD MOSER. 1975. A new dimorphic species of *Pyemotes* and a key to previously described forms (Acarina: Tarsone-moidea). *Entomological Society of America, Annals* 68: 723–732. (ec).
- CROSS, EARLE ALBRIGHT, JOHN CONRAD MOSER, AND G. RACK. 1981. Some new forms of *Pyemotes* (Acari: Pyemotidae) from forest insects, with remarks on polymorphism. *International Journal of Acarology* 7: 179–196. (ec).
- CROTCH, GEORGE ROBERT. 1863. A catalog of British Coleoptera [Scolytidae, p. 18]. Cambridge University Press, [printed by] C. J. Clay. (ds).
- CROWSON, ROY ALBERT. 1957. A second Scottish locality for *Scolytus intricatus* Ratzeburg (Col., Scolytidae). *Entomologist's Monthly Magazine* 123: 126. (ds).
- CROWSON, ROY ALBERT, AND F. A. HUNTER. 1964. Some Coleoptera associated with old trees in Grimsthorpe Park, Lincs. *Entomologist's Monthly Magazine* 100: 198–200. (ds).
- and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn).
- DALE, JOHN W., AND G. DE NITTO. 1986. Tree mortality associated with a high water table. Oak Knoll Ranger District, Klamath National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report S6–14. (ec).
- DALE, JOHN W., AND J. KLIEJUNAS. 1986. Mountain pine beetle activity in three stands of lodgepole pine near Truckee, California. United States Department of Agriculture, Forest Service, Pacific Southwest Region (San Francisco, California), Forest Pest Management Report S6–9. (cn).
- *DAMIANO. 1963. [*Hylesinus oleiperda* (=toranio)]. *Rivista de Agricoltura Subtropica e Tropicale*, Firenze 57: 45S. ().
- DANKS, HUGH V., AND R. G. FOOTITT. 1989. Insects of the boreal zone of Canada. *Canadian Entomologist* 121: 625–690. (ds).
- *DANTHIONE. 1888. *Byrrhus toranio*. Page 270 in [J. Et. ?] Bernard, *Memoir pour servir a l'Histoire Naturelle de l'olivier*, sec. II: Des insectes qui vivent sur l'olivier. *Memoir pour servir a l'Histoire Naturelle de la Provence* 6: 271–272. ().
- *DAS, A. K., M. K. DEVROY, AND B. MITRA. 1988. Insect borers of mangroves in the Bay Islands. *Andaman Science Association, Journal* 4: 32–38. ().
- DAVID, C. T., D. A. TILES, AND DAVID LEE WOOD. 1979. Factors associated with tree failure of giant sequoia: entomological aspects. Page 239 in *Proceedings: First conference on scientific research in national parks*. (cn ec).
- DAVIDSON, RALPH HOWARD, AND W. F. LYON. 1987. Insect pests of farm, garden, and orchard. Edition S. J. Riley & Sons, New York. xiii + 640 p. (cn hb).
- DAVIDSON, W. F. 1961. Notes on Cumberland and Westmorland Coleoptera. *Entomologist's Monthly Magazine* 97: 15–21. (ds).
- *DAVIS. 1937. [*Scolytus rugulosus*]. *Indiana Academy of Science, Proceedings* 46: 230–239 (1936). ().
- *_____. 1947. [*Scolytus rugulosus*]. *Indiana Academy of Science, Proceedings* 56: 150. ().
- DAVIS, MERRILL S., AND W. B. WHITE. 1959. A tool for assessing the impact of mountain pine beetle and related management strategies. Pages 37–40 in G. D. Amman, *Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- *DEAN, GEORGE ADAMS. 1918. [*Scolytus rugulosus*]. *Kansas State Horticultural Society, Transactions* 34: 197. ().
- *_____. 1922. [*Scolytus rugulosus*]. *Kansas State Horticultural Society, Biennial Report* 36: 181. ().
- *DEAN, GEORGE ADAMS, AND LEONARD MARION PEARS. 1913. [*Phloeotribus liminaris*]. *Kansas Agric. Educ.* 1: 2, 76. ().
- DEANGELIS, J. D., JOHN DEAVOURS HODGES, AND T. EVAN NEBECKER. 1986. Phenolic metabolites of *Ceratocystis minor* from laboratory cultures and their effects on transpiration in loblolly pine seedlings. *Canadian Journal of Botany* 64: 151–155. (ec).
- DEANGELIS, J. D., T. EVAN NEBECKER, AND JOHN DEAVOURS

- HODGES. 1956. Influence of tree age and growth rate on the radial resin duct system in loblolly pine. *Canadian Journal of Botany* 64: 1046-1049. (bv).
- DEARBORN, R. G., AND D. A. STARK. 1956. Forest and shade tree insect and disease conditions for Maine: summary issue for the 1955 season. Maine Department of Conservation, Forest Service, Entomology Laboratory Report (Augusta, Maine) 85-18. 16 p. (cn).
- DEBARR, G. L., AND L. R. BARBER. 1958. Use of prescribed burning to control the white pine cone beetle, *Conophthorus coniperda* (Schwarz) (Coleoptera: Scolytidae) in an eastern white pine seed orchard. International Congress of Entomology, Proceedings 15: 439. (cn).
- DECAZY, B., AND H. OCHOA. 1958. Eficacia de la aplicacion de un insecticida a bajo volumen sobre la broca del cafe *Hypothenemus hampei* Ferr. [Effectiveness of low-volume insecticide application on the coffee berry borer] [English summary]. *Boletin de Promecafe* 40: 6-10. (cn).
- DEDEK, W., AND J. PAPE. 1988. Integrated pest control in forest management: combined use of pheromones and insecticides for attracting and killing the bark beetle *Ips typographus* L. Studies with 32p-labelled methamidophos in the ascending sap of spruce. *Forest Ecology and Management* 26: 47-61. (cn).
- DEDEK, W., J. PAPE, F. GRIMMER, AND H. J. KORNER. 1988. Integrated pest control in forest management: combined use of pheromones and insecticides for attracting and killing the bark beetle *Ips typographus*. II. Effects of methamidophos treatment following bark penetration into the ascending sap of pheromone-baited spruce. *Forest Ecology and Management* 26: 63-76. (cn).
- DEJONG, MART C. M., AND P. GRIPMA. 1956. Competition between larvae of *Ips typographus*. *Entomologia Experimentalis et Applicata* 41: 121-133. (ec hb).
- DEJONG, MART C. M., AND HANNU SAARENMAA. 1955. A mechanistic simulation model for the movement and competition of bark beetle larvae (Coleoptera, Scolytidae). *Ecological Modelling* 27: 109-135. (ec).
- DEJONG, MART C. M., AND M. W. SABELIS. 1988. How bark beetles avoid interference with squatters: an ess for colonization by *Ips typographus*. *Oikos* 51: 88-96. (ec).
- *DELGUERCIO, GIACOMO. 1913. Nuova contribuzione alla conoscenza dei nemici dell'olivo. V. Osservazioni preliminari intorno ai costumi dell'*Llesino* dell'olivo (*Hylesinus oleiperda*). *Redia* 9: 70-74. ().
- * _____. 1939. [*Hylesinus oleiperda* (=toranio)]. *Redia Firenze* [Florence] 51 [page or volume number? An obvious error in this citation exists; vol. 25 was published in 1939]. ().
- DELL, T. R., JACQUELINE L. ROBERTSON, AND MICHAEL I. HAWERTY. 1953. Estimate of cumulative change of state with the Weibull function. *Entomological Society of America, Bulletin* 29(4): 38-40. (ec).
- DELOBEL, ALEX. 1992. Les cossettes de manioc, un important reservoir d'insectes des denrees stockees en Afrique central. *Revue de Zoologie Africaine* 106: 17-25. (ec ds).
- DEMAKOV, YU. P. 1958. Accuracy requirements in counting pine bark beetles [In Russian, English summary]. *Lesnoe Khozyaistvo* 3: 35-38. (hb).
- DEMARS, CLARENCE JOHN, JR., DONALD L. DAHLSTEN, N. X. SHARPBACK, AND D. L. ROWNEY. 1956. Tree utilization and density of attacking and emerging populations of the western pine beetle (Coleoptera: Scolytidae) and its natural enemies, Bass Lake, California 1970-1971. *Canadian Entomologist* 118: 881-900. (bv ec hb).
- DEMARS, CLARENCE JOHN, JR., GEORGE T. FERRELL, AND W. J. OSTROSINA. 1958. Host-insect/disease interactions in drought-stressed white fir stands at Lake Tahoe, California. Pages 135-146 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv ec).
- DENITTO, GREGG. 1984. Preliminary evaluation of the pest situation at the Zephyr Cove Special Use Area, Lake Tahoe Basin Management Unit. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 54-36. (cn).
- DENITTO, GREGG, AND J. W. DALE. 1982a. An evaluation of insect and disease conditions at recreation areas of the Little Grass Valley Reservoir, La Porte Ranger District, Plumas National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-31. (cn).
- _____. 1982b. An evaluation of the Illinois Sale Area, La Porte Ranger District, Plumas National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-26. (cn).
- DENITTO, GREGG, AND J. PIERCE. 1981. An evaluation of existing and potential forest disease and insect problems in the Catavee and Kinnikinnick Campgrounds, Pineridge Ranger District, Sierra National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 81-15. (cn).
- _____. 1983. An evaluation of conifer mortality on the San Bernardino National Forest between May 1981 and May 1982. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-35. (cn).
- _____. 1955. Biological evaluation of pests at Los Prietos Ranger Station and oak dieback. Santa Barbara Rd., Los Padres National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 55-33. (cn).
- DENITTO, GREGG, AND D. SCHULTZ. 1982a. Evaluation of the level and causes of tree mortality on the Dome Timber Sale, Cannell Meadow Ranger District, Sequoia National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-47. (cn).
- _____. 1982b. Evaluation of the Ridge 2 Sale, Tule River District, Sequoia National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-41. (cn).
- DENITTO, GREGG, AND J. WENZ. 1983. Biological evaluation of the Crane Valley, Denver Church, and Little Denver Church Campgrounds, Bass Lake Ranger District, Sierra National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-15. (cn).

- DESLARZES, K., C. LE NEDIC, AND D. CHERIX. 1957. Efficiency and selectivity of bark beetle *Ips typographus* traps in terms of their color and kind of forest area. *Société Vaudoise des Sciences Naturelles, Bulletin* 78: 349–362. (cn).
- DEVADAS, V. C., KANDASAMI, N., MURALEEDHARAN, K. V. S. N., RAO, K. G., UDAYBIANU, AND R. VARATHARAJAN. 1989. Mid-cycle chemical control of *Xyleborus formicatus* Eichhoff (Scolytidae: Coleoptera), the shot-hole borer of tea. *Journal of Plantation Crops* 16(supplement): 179–183. (cn).
- DEVDAIRANI, T. G. 1973. Two new species of nematodes from the beetle *Taphrorychus bicolor* Herbst from the eastern Georgian SSR [In Georgian, Russian and English summaries]. *Sobshch. Akad. Nauk Gruz. SSR* 70: 213–216. (ec).
- DEYRUP, MARK A. 1981. Deadwood decomposers. *Natural History* (3): 84–91. (ec ms).
- . 1957. *Trischlidia exigua* Wood, new to the United States, with notes on the biology of the genus (Coleoptera: Scolytidae). *Coleopterists Bulletin* 41: 339–343. (hb ds).
- . 1955. Scolytid diversity: key to biological riddles (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 18: 410. (hb).
- DEYRUP, MARK A., AND THOMAS HARRIS ATKINSON. 1987a. Comparative biology of temperate and subtropical bark and ambrosia beetles (Coleoptera: Scolytidae, Platypodidae) in Indiana and Florida. *Great Lakes Entomologist* 20: 59–66. (bv hb).
- . 1987b. New distribution records of Scolytidae from Indiana and Florida. *Great Lakes Entomologist* 20: 67–68. (ds).
- °DICKASON, ELVIS ARNIE, AND RAY THOMAS EVERLY. 1955. [*Hylastinus obscurus*]. *Oregon State Extension Bulletin* 749: 13–14. ().
- DICKENS, JOSEPH CLIFTON, AND THOMAS LEE PAYNE. 1985. Chemical messengers and insect behavior. Pages 201–230 in N. B. Mandova, *Handbook of natural pesticides: methods*. Volume 1: Theory, practice and detection. CRC Press, Boca Raton, Florida. (bv ec).
- DICKENS, JOSEPH CLIFTON, THOMAS LEE PAYNE, LEE C. RYKER, AND JULIUS ALEXANDER RUDINSKY. 1985. Multiple acceptors for pheromonal enantiomers on single olfactory cells in the Douglas-fir beetle, *Dendroctonus pseudotsugae* Hopk. (Coleoptera: Scolytidae). *Journal of Chemical Ecology* 11: 1359–1370. (av bv).
- DIMITRI, L. 1955. Einsatz biotechnischer Verfahren zur Populationssenkung der Borkenkäfer. *Allgemeine Forstzeitschrift* 12: 254–256. (cn).
- DIMITRI, L., E. KONIG, E. NIEMEYER, AND O. VAUFEL. 1986. Der Dreifallenstern: eine möglichkeit zur steigerung der effektivität von Borkenkäferfallen [The "Three-trap star": a possibility of increasing the effectiveness of bark beetle traps]. *Forst- und Holzwirt* 41: 171–173. (cn).
- DIX, MARY ELLEN, AND DAVID LEATHERMAN. 1988. Insect management in windbreaks. *Agriculture, Ecosystems and Environment* 22/23: 513–537. (cn).
- DIX, MARY ELLEN, JUDITH E. PASEK, MARK O. HARRELL, AND FREDERICK P. BANENDALE. 1986. Common insect pests of trees in the Great Plains [Scolytidae, p. 15, 25–26]. *Great Plains Agricultural Council Publication No. 119*, and Nebraska Cooperative Extension Service EC 86–154S. 44 p. (cn hb).
- °DIXON. 1913. [*Scolytus rugulosus*]. *Texas Department of Agriculture Bulletin* 32: 136. ().
- DIXON, E. A., N. IBRAHIM, H. WIESER, H. F. CEREZKE, AND A. WHITEHEAD. 1988a. Interaction of bark beetles with pheromone mimics. *American Chemical Society (Abstracts)* 3(1). (bv).
- . 1988b. On the complementarity of electroantennogram signals and tree baiting and trapping catches for a series of bicyclic ketal pheromone analogs. *American Chemical Society (Abstracts)* 3(1). (bv).
- DIXON, E. A., N. IBRAHIM, H. WIESER, D. R. CRUMP, F. X. WEBSTER, AND G. D. PRESTWICH. 1988. Isolation and identification of the metabolic products from mountain pine beetle exposed to exo-brevicomin, frontalin and analogs of these. *American Chemical Society (Abstracts)* 3(1). (bv).
- DIXON, WAYNE N. 1983. Clonal specificity of *Ips, Pityophthorus* spp. (Coleoptera: Scolytidae) in a slash pine seed orchard. *Florida Entomologist* 66: 514–517. (bv).
- °DOMINIK, J. 1965. Z doswiadczen nad moziwoscia zerowania owadów w drewnie cisá (*Taxus baccata* L.) [Experiments on the possibility of insects feeding in the wood of yew]. *Sylvan* 6: 55–60. ().
- DONISTHORPE, HORACE ST JOHN KELLY HERBERT. 1931. An annotated list of the additions to the British coleopterous fauna, made since the publication of the supplementary volume (VI) of Fowler's *Coleoptera of the British Islands*. *Entomologist's Monthly Magazine* 67: 119–126, 164–175. (ds).
- . 1932. A second capture of *Aulonium trisulcatum* Geoff. in Windsor forest. *Entomologist's Monthly Magazine* 68: 14. (ec).
- ° ———. 1939. A preliminary list of the Coleoptera of Windsor Forest. *Nathaniel Lloyd & Co., Ltd., Blackfriars*. 124 p. ().
- DOVER, S. 1984. The main insects of pistacio in Israel [In Hebrew]. Pages 11–39 in *Proceedings: Third symposium of pistacio growers in Israel*. Sede Boger, 6 August 1984. Ben-Gurion University of the Negev, The Blaustein International Center for Desert Studies, Israel. (cn).
- DOWDING, P. 1984. The evolution of insect-fungus relationships in the primary invasion of forest timber. In A. D. M. Rayner and D. W. H. Walton. *Invertebrate-microbial interactions*. *British Mycological Symposium* 6. Cambridge University Press. (ec).
- DOZIER, H. L. 1918. An annotated list of Gainesville, Florida, Coleoptera. *Entomological News* 29: 370–374. (ds).
- DRACZYNSKA, B., C. CAGARA, A. SIEWINSKI, A. RYMKIEWICZ, A. ZABZA, AND A. LEUYEN. 1955. Biotransformation of pinenes. XVII. Transformation of alpha- and beta-pinenes by means of *Armillariella mellea* (honey fungus), a parasite of woodlands. *Journal of Basin Microbiology* 25: 487–492. (ms).
- DRANE, A. B. 1985. A second Northants. Locality for *Emoporus caucasicus* Lindemann (Col., Scolytidae) and notes on some other beetles. *Entomologist's Monthly Magazine* 121: 107. (ds).
- DROOZ, ARNOLD T. 1955. Insects of eastern forests. United States Department of Agriculture, Forest Service, *Miscellaneous Publication* 1426. 608 p. (cn ec hb).
- DUFFY, EVELYN ARTHUR JOSEPH. 1945. The coleopterous fauna of the Hants-Surrey border. *Entomologist's Monthly Magazine* 51: 169–179. (ds).
- DU, J. W. 1987. Chemical communication in the bark beetle [In Chinese. English summary]. *Chinese Journal of Biological Control* 3(2): 91–95. (bv).
- DU MERLE, P. 1988. Studies of forest insect pests in southern

- France. Revue Forestiere Francaise (Nancy) 40: 297-301. (cn).
- DUBBEL, VOLKER. 1985. Untersuchungen zum Einsatz von Lockstoff: Fallen gegen den Gestreiften Nutzholzborkenkäfer *Trypodendron lineatum* Oliv. Published dissertation, Forstl. Fak. Univ. Freiburg. (bv).
- DUBBEL, VOLKER, K. KERCK, M. SOHRT AND S. MANGOLD. 1985. Influence of trap color on the efficiency of bark beetle pheromone traps. Journal of Applied Entomology 99: 59-64. (cn).
- DUBBEL, VOLKER, O. VAUPEL AND L. DEMITRI. 1985. Untersuchungen zur Wirksamkeit und Ökologischen Vertraglichkeit von Borkenkäfer fallen. Holz-Zentralblatt 111: 357-359. (cn).
- DUELL, P., M. STUDER AND W. NAF. 1986. Der Borkenkäferflug ausserhalb des Waldes [The flight of bark beetles outside forest areas] [English summary]. Journal of Applied Entomology 102: 139-145. (bv cn lb).
- DUNCAN, B. 1987. An illustrated guide to the identification and distribution of the species of *Dendroctonus* Erichson (Coleoptera: Scolytidae) in British Columbia. Entomological Society of British Columbia, Journal 54: 101-112. (ds tv).
- DUNN, J. P., T. W. KIMMERER AND G. L. NORDIN. 1986. Attraction of the twolined chestnut borer, *Agrilus bilineatus* (Weber) (Coleoptera: Buprestidae) and associated borers to volatiles of stressed white oak. Canadian Entomologist 118: 503-509. (bv cn).
- DWINELL, L. D. 1985. Distribution of pine wood nematode in southern pine beetle-killed loblolly pines colonized by pine sawyers. Phytopathology 78: 1555. (ec).
- DYER, E. D. A., AND P. M. HALL. 1983. Effect of high density frontal baiting on attack distribution of *Dendroctonus rufipennis* in spruce plots. Entomological Society of British Columbia, Journal 80: 14-19. (bv).
- E**
- *EASTHAM. 1915. [*Scolytus rugulosus*]. Entomological Society of British Columbia, Victoria, Proceedings 7: 18-21. ().
- ECKSTEIN, KARL. 1939d. Das Bohrmehl des Waldgartners, *Myelophilus piniperda* L. Arbeiten der physiologische und angewandte entomologie aus Berlin-Dahlem 6: 32-41. (hb).
- EDMONDS, R. L., AND A. EGLITIS. 1989. The role of the Douglas-fir beetle and wood borers in the decomposition of and nutrient release from Douglas-fir logs. Canadian Journal of Forest Research 19: 853-859. (cn).
- *EDSON, LEWIS J. 1967. An annotated and illustrated key to the species of the genus *Scolytus* (Coleoptera, Scolytidae) attacking coniferous trees of the Nearctic region. Unpublished thesis, Humboldt State College, California. ().
- EDWARDS, B. M. 1982. Operational control of the Douglas-fir beetle. Potlatch Corporation, Wood Products, Western Division, Forestry Research Report RR-82-2. 7 p. (cn).
- EGGER, ALFRED. 1974b. Zur Biologie von *Hylastinus fankhouseri* Reitt. Centralblatt für das Gesamte Forstwesen 91: 65-74. (hb).
- _____. 1987. Pheroprax-Versuche gegen den Buchdrucker (*Ips typographus* L.) 1978-1981 [Pheroprax trials against the spruce bark beetle *Ips typographus* 1978-1981] [English summary]. Centralblatt für das Gesamte Forstwesen 104: 157-190. (cn).
- *EGGERS, HANS. 1925. Kulturschädliche Borkenkäfer des indischen Archipels. Entomologischen Berichten 121-144: 84-88. ().
- EGLITIS, ANDRIS. 1985. Permanent plots for monitoring population trends of the spruce beetle in Glacier Bay National Park. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Report R181 (Technical Report R10-85-1). (cn).
- _____. 1986. Spruce beetle in Glacier Bay National Park, 1985 update. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Biological Evaluation R10-86-1. 17 p. (cn).
- _____. 1987. Spruce beetle in Glacier Bay National Park, 1986 update. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Forest Pest Management Report R10-87-4. 13 p. (cn).
- _____. 1985a. Effects of high water on vegetation during glacial closure of Russell Fiord in 1987. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Forest Pest Management Report R10-85-4. 10 p. (cn).
- _____. 1985b. Spruce mortality in Yakutat Forelands, 1987. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Biological Evaluation R10-85-2. 9 p. (cn).
- EGLITIS, ANDRIS, AND PAUL E. HENNON. 1985. Effects of high water on vegetation during glacial closure of Russell Fiord in 1987. United States Department of Agriculture, Forest Service, Alaska Region (Anchorage, Alaska), Biological Evaluation R10-85-4. 10 p. (cn).
- EIINSTROM, BENGT. 1985. Insektsskador i svenska skogar sedan 1970 [Insect damage in Swedish forests since 1970] [English summary]. Sveriges Skogvårdsförbunds Tidskrift 2: 11-18. (cn).
- EIBER, T. 1989. Popular literature on *Ips pini* in Minnesota. Minnesota Department of Natural Resources. (cn ms).
- EIDMANN, HUBERTUS H. 1985a. Silviculture and insect problems. Journal of Applied Entomology 99: 105-112. (cn).
- _____. 1985b. Skogsinsekter-problem och forskning [Forest insect problems and research in Sweden] [English summary]. Sveriges Skogvårdsförbunds Tidskrift 2: 3-8. (cn).
- _____. 1985c. Control and monitoring the spruce bark beetle *Ips typographus* in Sweden using pheromones. Pages 205-221 in Biological and biotechnical control of forest pests: symposium proceedings, Tabor. (cn).
- _____. 1985d. Niceni a signalizace kurovce *Ips typographus* ve svedsku s ponioci feromonu. Pages 132-138 in Biological and biotechnical control of forest pests: symposium proceedings, Tabor. (cn).
- _____. 1987a. Der Befall von Scolytiden und Cerambyciden an Stammabschnitten fremdländischer Koniferen in Schweden [Infestation of logs of exotic conifers in Sweden by scolytids and cerambycids] [English summary]. Journal of Applied Entomology 103: 278-283. (bv cn).
- _____. 1987b. Attractants and deterrents of natural origin for the suppression of bark beetle attacks. Pages 56-62 in Proceedings: European seminar on wood production and harvesting. Bologna. Vol. 3. (cn).

- EIDMANN, HUBERTUS H., P. BAECKSTROM, S. HARDING, T. NORIN, J. VRKOC, AND J. WESLIEN. 1957. Methylbutanol effectively replaces methylbutenol, a pheromone component of *Ips typographus* (L.) (Coleoptera: Scolytidae). *Journal of Chemical Ecology* 13: 1555–1560. (bv).
- EIDMANN, HUBERTUS H., AND GÖRAN BIRGERSSON. 1958. Semiochemicals in the East Himalaya spruce bark beetle. *Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz* 61: 147–148. (bv).
- *EIDMANN, HUBERTUS H., AND T. INGESTAD. 1963. Ernährungsstand, Zuwachs und Insektenbefall in einer Kiefernkultur. *Studia forestalia suecica*, Stockholm 12: 1–22. ().
- EIDMANN, HUBERTUS H., AND A. KLINGSTROM. 1976. Skadegörelse i skogen. Svampar-Insekter-Ryggradsdjur. Lts forlag: Centraltryckeriet AB, Borås. 288 p. (cn hb).
- EIDMANN, HUBERTUS H., J. WESLIEN, S. HARDING, P. BAECKSTROM, T. NORIN, AND J. VRKOC. 1956. A compound replacing a natural pheromone component of the spruce bark beetle. *Naturwissenschaften* 73: 629–630. (bv).
- EIDMANN, HUBERTUS H., N. WIERSMA. 1959. Development and reproduction in the East Himalaya spruce bark beetle. *Journal of Applied Entomology* 107: 411–417. (hb).
- EISENHOUR, D. R. 1959. Untersuchungen zur Entwicklung der ökologischen Stabilität vor Eichenbeständen im nordöstlichen Harzvorland [Investigation towards improving the ecological stability of oak stands in the northeastern foothills of the Harz Mountains] [In German, English summary]. *Beiträge für die forstwirtschaft* 23: 55–62. (cc).
- *ELDRIDGE, R. H. 1953. Introduced pine bark beetles in New South Wales. *Forest Protection Series*, Forestry Commission of New South Wales 5: 1–3. ().
- ELFYNG, B., AND BO LANGSTROM. 1954. Kronskadegörelse och tillväxtreaktion i ett margborreskadat bestånd [Crown damage and growth response in a pine stand attacked by *Tomicus piniperda*]. *Sveriges Skogsvårdsförbunds Tidskrift*, nr. 1. (cn).
- *ELGSTRAND. 1921. [*Hyllastes ater*]. Skogen, Stockholm 1921: 225–229. ().
- ELGUETA D., MARIO, AND D. JACKSON S. 1957. Nombre actual de las especies de Curculionidae (Insecta: Coleoptera) tratadas en la obra de Gay. *Revista Chilena de Entomología* 15: 71–78. (tx).
- ELIAS, SCOTT A. 1982d. Holocene insect fossils from two sites at Ennadai Lake, Keewatin, Northwest Territories, Canada. *Quaternary Research* 17: 371–390. (ds).
- . 1955. Paleoenvironmental interpretations of Holocene insect fossil assemblages from four high-altitude sites in the Front Range, Colorado, USA. *Revue de Paleobiologie* 5: 31–45. (ds).
- ELIAS, SCOTT A., S. K. SHORT, AND P. V. CLARK. 1956. Paleoenvironmental interpretations of the Late Holocene, Rocky Mountain National Park, Colorado, USA. *Revue de Paleobiologie* 5: 127–142. (ds).
- ELLIOTT, H. J., R. BASHFORD, S. CANDY, AND G. A. KILE. 1955. The role of the ambrosia beetle, *Platypus subgranosus* Schedl, as an indicator of infection of myrtle, *Nothofagus cunninghamii* (Hook.) Gerst., by the pathogenic fungus, *Chalara australis* Walker and Kile, in Tasmanian rainforest. *International Congress of Entomology, Proceedings* 18: 430. (cc).
- ELLIOTT, H. J., G. A. KILE, S. G. CANDY, AND D. A. RATKOWSKY. 1957. The incidence and spatial pattern of *Nothofagus cunninghamii* (Hook.) Gerst. attacked by *Platypus subgranosus* Schedl in Tasmania's cool temperate rainforest. *Australian Journal of Ecology* 12: 125–138. (bv ec).
- ELLIOTT, H. J., G. A. KILE, AND J. L. MADDEN. 1952a. Ambrosia beetle attack on myrtle. Pages 67–73 in *Symposium: Tasmanian rainforests, recent research results*, November 1952. Organized by: Forest Ecology Research Fund. (cn ec).
- ELLIOTT, H. J., AND D. W. DELITTLE. 19... Insect pests of trees and timber in Tasmania. *Forestry Commission, Tasmania*. (cn hb).
- ELMAGRABY, SALAH E., H. A. DEVINE, S. D. DILLERY, AND J. E. DE STEIGUER. 1957. Optimal partitioning of a seasonal distribution. Pages 476–484 in P. E. Dress and R. C. Field, *Systems analysis in forest resources: symposium proceedings*, 1955, Society of American Foresters. Georgia Center for Continuing Education, Athens, Georgia. 555 p. (cn hb).
- EMOND, F. J. 1969. Annual district report: West-Central District, 1968. Pages 33–34 in *Annual District Reports of the Forest Insect and Disease Survey*, Alberta-Northwest Territories-Yukon Region, 1968. Canada Department of Fisheries and Forestry, Canadian Forestry Service, Forest Research Laboratory, Calgary, Alberta. Information Report A-X-22. (cn).
- ENDRODI SEBO. 1956. The families of Anthribidae, Nemomychidae, Attelabidae, Curculionidae and Scolytidae of the Rhynchophora in the Kiskunsag National Park (Coleoptera). Pages 197–215 in S. Mahunka (ed.), *Natural history of the national parks of Hungary*, vol. 4. The fauna of Kiskunsag National Park, vol. 1. Akademiai Kiado, Budapest, Hungary. 490 p. (ds).
- EQUIHUA MARTINEZ, ARMANDO. 1955. Nuevos registros de localidades y huéspedes de Coleopteros Platypodidae Americanos. *Folia Entomologica Mexicana* 66: 143–144. (ds).
- . 1955. Coleopteros Scolytidae atraídos a trampas NTP-50 en el área norte de la Reserva de la Biosfera "sain Ka'an", Quintana Roo, Mexico. *Folia Entomologica Mexicana* 74: 179–180. (cn).
- EQUIHUA MARTINEZ, ARMANDO, AND THOMAS HARRIS ATKINSON. 1956. Annotated checklist of bark and ambrosia beetles (Coleoptera: Scolytidae and Platypodidae) associated with a tropical deciduous forest at Chamela, Jalisco, Mexico. *Florida Entomologist* 69: 619–635. (ds ec hb).
- . 1957. *Catálogo de Platypodidae (Coleoptera) de Norte y Centroamérica* [Catalogue of the Platypodidae (Coleoptera) of North and Central America] [English summary]. *Folia Entomologica Mexicana* 75: 5–31. (ds tx).
- ERICSSON, A., C. HELLQVIST, BO LANGSTROM, S. LARSSON, AND O. TENOW. 1955. Effects on growth of simulated and induced shoot pruning by *Tomicus piniperda* as related to carbohydrate and nitrogen dynamics in Scots pine. *Journal of Applied Ecology* 22: 105–124. (bv).
- ESTRADA VALENCIA, ANTONIO, AND THOMAS HARRIS ATKINSON. 1955. Scolytidae y Platypodidae (Coleoptera) de Escarcega, Campeche, Mexico. *Biogeografía, biología, importancia económica y una lista comentada de especies* [English summary]. *Anales Instituto de Biología UNAM* 55. Series Zoológica 1: 199–220. (cn ds).
- EVANS, DAVID. 1950. Important insects: British Columbia

- coastal forests. Canada Department of Agriculture, Forest Insect Investigations, Division of Entomology, Forest Insect Survey, Annual Report 1949: 107. (cn ds).
- _____. 1955. Annotated checklist of insects associated with Garry oak in British Columbia. Government of Canada, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-262. 36 p. (ds).
- EVANS, G. O. 1955. A revision of the British Aceosejiniae (Acarina: Mesostigmata). Zoological Society of London, Proceedings 131: 177-229. (ec).
- EVANS, HUGH F. 1955. Great spruce bark beetle, *Dendroctonus micans*: an exotic pest new to Britain. Antenna 9: 117-121. (cn).
- _____. 1957. Biological control of *Dendroctonus micans* in the USSR. Entopath News 89: 5-7. (ec).
- EVANS, HUGH F., AND C. J. KING. 1959. Biological control of *Dendroctonus micans* (Coleoptera: Scolytidae): British experience of rearing and release of *Rhizophagus grandis* (Coleoptera: Rhizophagidae). Pages 109-125 in D. L. Kullhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ec).
- EVANS, HUGH F., C. J. KING, AND D. WAINHOUSE. 1955. *Dendroctonus micans* in the United Kingdom. The results of two years experience in survey and control. Pages 20-34 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium, Proceedings of Seminar, 3-4 October 1954. 141 p. (cn).
- EVEARAERTS, C., J. C. GREGOIRE, AND J. MERLIN. 1955. The toxicity of Norway spruce monoterpenes to two bark beetle species and their associates. Pages 335-344 in W. J. Mattson, J. Levieux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (cn).
- F**
- *FAGEL, G., AND H. GUILLEAUME. 1935. [*Hylesinus oleiperda* (=toranio)]. Societe Entomologique de Belgique, Bulletin et Annales 75: 335. ().
- *FALL, HENRY CLINTON. 1901. List of the Coleoptera of southern California, with notes on habits and distribution and descriptions of new species. Occasional Papers of the California Academy of Sciences 5: 1-252. ().
- FANG, S. Y., C. S. WANG, AND D. F. CHI. 1984. Trapping experiment with *Ips* pheromone in Funling Natural Conservation Area in Heilongjiang Province. Northeastern Forestry Institute (China), Journal 12: 43-47. (cn).
- FANKHAUSER, F., JR. 1894. Oberförster Wilhelm Eichhoff. Schweizerische Zeitschrift für das Forstwesen 1894: 67-69. (ms).
- FARGO, WALTER SCOTT, TERRENCE L. WAGNER, AND ROBERT N. COULSON. 1955. Factors influencing the growth of multiple-tree infestations of *Dendroctonus frontalis*. Researches on Population Ecology 27: 25-35. (cn).
- FATZINGER, CARL WARREN. 1955a. Attraction of the black turpentine beetle (Coleoptera: Scolytidae) and other forest Coleoptera to turpentine-baited traps. Environmental Entomology 14: 765-775. (cn hb).
- _____. 1955b. Turpentine-baited traps capture black turpentine beetles and other forest Coleoptera but do not prevent attacks on pines. Pages 26-33 in S. J. Branham and R. C. Thatcher, Integrated pest management research: symposium proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station (New Orleans, Louisiana), General Technical Report SO-56. 353 p. (cn).
- FATZINGER, CARL WARREN, B. D. SIEGFRIED, ROBERT CLEVELAND WILKINSON, AND J. L. NATION. 1957. Trans-verbenol, turpentine, and ethanol as trap baits for the black turpentine beetle, *Dendroctonus terebrans*, and other forest Coleoptera in north Florida. Journal of Entomological Science 22: 201-209. (bv cn).
- FAULDS, W. 1955. Improved techniques for the laboratory rearing of *Thanastmus formicarius*. New Zealand Journal of Forest Science 15: 157-190. (cn).
- _____. 1959. *Hylastes ater* (Paykull), black pine bark beetle and *Hylurgus ligniperda* (Fabricius), golden haired bark beetle (Coleoptera: Scolytidae). Pages 271-275 in P. J. Cameron, R. L. Hill, J. Bain, and W. P. Thomas, A review of biological control of invertebrate pests and weeds in New Zealand 1874-1957. Technical Communication (Commonwealth Institute of Biological Control), Wallingford, Oxon, United Kingdom. (cn ec).
- FELDMAN, R. M., T. L. WAGNER, G. L. CURRY, ROBERT N. COULSON, AND PETER J. H. SHARPE. 1955. A methodology for biophysical modeling using TAMBEETLE as an example. Pages 195-201 in S. J. Branham and R. C. Thatcher, Integrated pest management research: symposium proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 353 p. (cn ms).
- *FELT, EPHRAIM PORTER. 1906. [*Hylurgops glabratus* (=rugipennis pinifex)]. New York State Museum, Memoir 5(2): 649, 665-666, fig. 194. ().
- *FERNALD, HENRY TORSEY. 1921. Applied entomology, an introductory textbook of insects in their relations to man. Edition 1. McGraw-Hill, New York. [1926?]. ().
- FERREIRA, GODOFRICO W. S. 1955. Desenvolvimento de *Tomicus pinor* (L.) em arvores derrubados pelo vento e em toros em estaleiro em pinhal bravo (Coleoptera: Scolytidae). Sociedade Portuguesa de Entomologia 1955(suplemento 1): 285-290. (cn).
- FERREIRA, J. M. S., AND J. P. MORIN. 1955. Ocorrência de *Xyleborus ferrugineus* (Fabricius, 1801) e *Xyleborus affinis* (Eichhoff, 1867) (Coleoptera: Scolytidae) sobre coqueiro no estado de Sergipe. Anais da Sociedade Entomologica do Brasil 14: 327-330. (cn hb ds).
- FERREIRA, MARIA CORTINA. 1963. Catalogo dos Coleopteros de Mocambique [Scolytidae, p. 664-674; Platypodidae, p. 675-679]. Revista de Entomologia de Mocambique 6: 533-1005. (ds).
- _____. 1955a. Impacte dos predadores da classe insecta nos escolidideos do pinheiro bravo. Sociedade Portuguesa de Entomologia, Boletim 4(suplemento 1): 107-114. (ec).
- _____. 1955b. Sobre a presença de *Tomicus minor* (Hartig) em Portugal continental (Coleoptera: Scolytidae). Sociedade Portuguesa de Entomologia, Boletim 4(suplemento 1): 281-283. (ds).
- _____. 1956. Investigação a hilesina do pinheiro [*Tomicus*

- piniperda*]. Gazeta das Aldeias 1986(Dezembro): 10-11. (cn hb).
- _____. 1987. Pragas relacionadas com o fogo em resinosas: o gorgulho do pinheiro. Floresta 8(Novembro-Dezembro): 22-23. (cn hb).
- _____. 1988a. Pragas do pinheiro bravo. Tecnologias Agrarias 1(1): 12-13. (cn hb).
- _____. 1988b. Impacto dos incendios nas pragas subcorticais em pinhal. Tecnologias Agrarias 1: 13-14. (cn hb).
- _____. 1989. Contribution to the bioecological study of *Phloeosinus aubei* Perris (Coleoptera, Scolytidae) [In French]. International Congress of Coleopterology, Abstracts 1989: 46. (ay lw ec).
- FERREIRA, MARIA CORINTA AND GODOFRICO W. S. FERREIRA. 1985a. Impacto do bostrio e da hilesina em pinheiro bravo, em Portugal Continental (Scolytidae: Coleoptera). Sociedade Portuguesa de Entomologia, Boletim 4(suplemento 1): 91-106 (cn).
- _____. 1985b. Contribuicao para o estudo da hibernacao dos escolitideos do pinheiro bravo em Portugal Continental (Coleoptera). Sociedade Portuguesa de Entomologia, Boletim 4(suplemento 1): 115-122. (cn hb).
- _____. 1985c. Identificacao e bioecologia dos escolitideos associados com o pinheiro bravo em Portugal Continental. Estacao Florestal Nacional, Curso de pragas florestais integrado na Proteccao da Floresta contra pragas. 15 p. (cn).
- _____. 1986a. Importancia da causa das arvores nas biocenoses das essencias florestais. Boletim Agricola, ser. 2, 38: 5-6. (cn).
- _____. 1986b. Pragas do pinheiro bravo em Portugal: escolitideos [Pests of maritime pine in Portugal: scolytids] [English summary]. Boletim Agricola 36(Janeiro-Marco): 1-4. (cn hb).
- _____. 1986c. Ravageurs rapportes avec le feu dans les peuplements de *Pinus pinaster*. Reuniao luso-francesa em Marvao, Secco de Incendios florestais. 5 p. (cn).
- _____. 1986d. Pragas relacionadas com incendios em pinhal bravo. Congresso Florestal Nacional [Portugal], Comunicacoes 1: 169-172. (cn).
- _____. 1986e. Scolytids associated with fires in *Pinus pinaster* forests. OEPP/EEPO Bulletin 16: 507. (cn).
- _____. 1986f. Pragas do pinheiro bravo em Portugal: escolitideos. Buletin Agricola, ser. 2, 36: 1-4. (cn).
- _____. 1987a. Ataques de insectos relacionados com praticas silvcolas: o genero *Hylastes* Erichson. Boletim Agricola, ser. 2, 43: 5-6. (cn).
- _____. 1987b. Escolitideos vectores de fungos em pinhal bravo. Boletim do Instituto dos Productos Florestais, Resinosos 56: 35-36. (cn).
- _____. 1987c. Impacto do fogo nos escolitideos associados com o pinheiro bravo (*Pinus pinaster* Ait.). Floresta 4-5: 12-14. (cn).
- _____. 1988a. Pragas do pinheiro manso. Pages 1-24. pl. I-XI in Encontro sobre pinheiro manso. Alaracer do Sal, 25 e 26 Nov. 1988. Sociedade Portuguesa de Ciencias Florestais. (cn).
- _____. 1988b. Impacto do fogo nos escolitideos associados com o pinheiro bravo. Pages 1-7 in Simposio sobre: a floresta e o ordenamento do espaco de montanha. Vila Real, Maio de 1988. Sociedade Portuguesa de Ciencias Florestais. (cn).
- _____. 1988c. Ataques de insectos relacionados com praticas de silvicultura. Floresta 2: 20-21. (cn).
- _____. 1988d. Desequilibrios causados pelos incendios florestais nas populacoes de insectos. Symposium: Jornadas cientificas sobre incendios florestais, 23 a 25 de Novembro de 1988. Universidade de Coimbra, Coimbra, Portugal. 7 p. (cn).
- _____. 1989a. Investagacao: pragas do sobreiro, *Platypus cylindrus* F. (Coleoptera, Platypodidae). Gazeta das Aldeias 2956(Marco 1989): 29-30. (cn).
- _____. 1989b. *Platypus cylindrus* F. (Coleoptera, Platypodidae), praga de *Quercus suber* L. [In Spanish, English summary]. Boletim de Sanidad Vegetal, Plagas 15: 301-306. (ay cn ds).
- _____. 1989c. Succession of bark beetles and wood-borers (Coleoptera) infesting *Pinus pinaster* in Portugal. International Congress of Coleopterology, Abstracts 1989: 47. (ec hb).
- _____. 1990a. Pages 30-31, 49-75 in Pragas das resinosas: Guia de Campo. Ministerio da Agricultura, Pescas e Alimentacao, Direccao-Geral de Planeamento & Agricultura, Lisboa. Serie Divulgacao, No. 3. 105 p. (cn hb ds).
- _____. 1990b. Pages 45-53, 63-65 in Pragas dos viveiros florestais, das plantacoes e da regeneracao natural: Guia de Campo. Ministerio da Agricultura, Pescas e Alimentacao, Direccao-Geral de Planeamento & Agricultura, Lisboa. Serie Divulgacao, No. 4. 132 p. (cn hb ds).
- _____. 1991. *Scolytus* spp. Pages 100-113 in Pragas das Folhosas: Guia de Campo. Ministerio da Agricultura, Pescas e Alimentacao, Direccao-Geral de Planeamento & Agricultura. Lisboa. Serie Divulgacao, No. 5. 192 p. (hb ds tx).
- FERRELL, GEORGE THOMAS. 1986a. Fir engraver. United States Department of Agriculture, Forest Service, Insect and Disease Leaflet 13. 8 p. (cn hb ds).
- _____. 1986b. Using indicator plants to assess susceptibility of California red fir and white fir to the fir engraver beetle. United States Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station (Berkeley, California), Research Note PSW-385. 5 p. (cn).
- _____. 1988. Wound-induced oleoresins of *Abies concolor*: is it part of host resistance to the fir engraver *Scolytus ventralis*? Pages 305-312 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (cn).
- FERRELL, GEORGE THOMAS, C. J. DE MARS, JR., C. B. WILLIAMS, JR. AND W. J. OTROSINA. 1988. Fir engraver outbreaks and tree mortality related to forest site and stand characteristics. Northwest Environmental Journal 4: 334-335. (bv cn).
- FERRELL, GEORGE THOMAS, W. J. OTROSINA AND C. J. DE MARS, JR. 1988. White fir characteristics associated with wound response to *Trichosporium symbioticum* fungal symbiont of the fir engraver beetle. Phytopathology 78: 1533. (ec).
- FIEDLER, CARL E. 1989. Utilization opportunities for reducing mountain pine beetle damage in lodgepole pine. Pages 70-74 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- FILIP, G. M., E. CHRISTIANSEN AND C. A. PARKS. 1989. Secondary resin production increases with vigor of *Abies grandis* inoculated with *Trichosporium symbioticum* in northeastern Oregon. United States Department

- of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon, Research Note PNW-459, 10 p. (ec).
- FLAMM, RICHARD O., S. P. COOK, TERENCE L. WAGNER, PAUL E. PULLEY, AND ROBERT N. COULSON. 1957. Reemergence and emergence of *Ips avulsus* and *Ips calligraphus* (Coleoptera: Scolytidae). *Environmental Entomology* 16: 569-576. (ec hb).
- FLAMM, RICHARD O., AND ROBERT N. COULSON. 1958. Traumatized hosts: their influence on the population dynamics of the southern pine bark beetle guild. Pages 345-355 in W. J. Mattson, J. Levieux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (hb).
- FLAMM, RICHARD O., ROBERT N. COULSON, P. BECKLEY, PAUL E. PULLEY, AND TERENCE L. WAGNER. 1959. Maintenance of a phloem-inhabiting guild. *Environmental Entomology* 18: 351-357. (ec).
- FLAMM, RICHARD O., ROBERT N. COULSON, AND THOMAS LEE PAYNE. 1955. The southern pine beetle. Pages 531-553 in A. A. Berryman, Dynamics of forest insect populations: pattern, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec).
- FLAMM, RICHARD O., TERENCE L. WAGNER, S. P. COOK, PAUL E. PULLEY, ROBERT N. COULSON, AND T. M. MCARDLE. 1957. Host colonization by cohabiting *Dendroctonus frontalis*, *Ips avulsus*, and *I. calligraphus* (Coleoptera: Scolytidae). *Environmental Entomology* 16: 390-399. (ec).
- FLECHTMANN, CARLOS A. H. 1955. On the biology of *Ameroseius dendroctagis* (Acari, Mesostigmata, Ameroseiidae). *Revista Brasileira de Zoologia* 2: 397-399. (ec).
- FLECHTMANN, CARLOS H. W., AND C. A. H. FLECHTMANN. 1955. A new species of *Ameroseius* (Acari: Mesostigmata, Ameroseiidae) from Brasil. *Revista Brasileira de Zoologia* 2: 393-396. (ec).
- FLORES L., JAIME, AND DONALD EDWARD BRIGIT, JR. 1957. A new species of *Conophthorus* from Mexico: descriptions and biological notes (Coleoptera: Scolytidae). *Coleopterists Bulletin* 41: 151-154. (hb tx).
- *FOLSOM, JUSTUS WATSON. 1909b. [Hylastinus obscurus]. Pages 92-96 in Illinois State Entomologist, Report 25. ().
- FOLTZ, JOHN L., J. A. CORNEIL, AND R. M. REICH. 1955. Procedures for sampling six-spined *Ips* populations in slash pine. Pages 6-12 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 353 p. (cn ms).
- FORBES, R. S., G. R. UNDERWOOD, D. C. EIDT, F. G. MUMING, AND W. J. CARROLL. 1955. Pages 9-21 in Atlantic provinces, forest insect survey. Canada Department of Forestry, Forest Entomology and Pathology Branch, Forest Insect and Disease Survey, Annual Report 1955. (cn).
- *FORD, L. B. 1956a. Attack dynamics of the spruce beetle, *Dendroctonus rufipennis* (Kirby), in south-central Alaska. Unpublished dissertation, University of Washington, Seattle. 171 p. ().
- _____. 1956b. Attack dynamics of the spruce bark beetle, *Dendroctonus rufipennis* (Kirby), in south-central Alaska. Dissertation Abstracts International (Science & Engineering) 47: 1350. (bv).
- FORNAL, A. B., JASKULSKA, AND A. KOLK. 1956. Evaluation of the effectiveness of catching the lineate bark beetle *Trypodendron lineatum* Ol. in traps with synthetic aggregation pheromone [In Polish]. *Sylvan* 130: 41-52. (bv cn).
- FORSSE, ERIK. 1957. Flight duration of *Ips typographus* L.: insensitivity to nematode infection. *Journal of Applied Entomology* 104: 326-328. (ec hb).
- _____. 1959. Migration in bark beetles with special reference to the spruce bark beetle *Ips typographus*. Published dissertation, Sveriges Lantbruksuniversitet, Uppsala. (hb).
- FORSSE, ERIK, AND CH. SOLBRECK. 1955. Migration in the bark beetle *Ips typographus* L.: duration, timing and height of flight. *Journal of Applied Entomology* 100: 47-57. (hb).
- FORSTER, H. W. 1949. *Ips sexdentatus* Boerner (Col., Scolytidae) in Surrey. *Entomologist's Monthly Magazine* 55: 153. (ds).
- FOSTER, H. R. 1961. Status of insects in the Lake Erie District. Pages 106-123 in Annual district report of the forest biology rangers of Ontario, 1960. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- _____. 1962. Status of insects in the Lake Erie District. Pages 113-134 in Annual district report of the Forest Insect and Disease Survey Ontario, 1961. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- _____. 1963. Status of insects and tree diseases in the northern forest region. Pages E1-S in Annual district reports of the Forest Insect and Disease Survey Ontario, 1962. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- _____. 1964. Status of insects and tree diseases in the northern forest region. Pages E1-9 in Annual district reports of the Forest Insect and Disease Survey Ontario, 1963. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- *FOX, J. W. 1959. The role of vectored fungi in the biology of *Ips* species. Unpublished dissertation, University of California, Berkeley. 107 p. ().
- FOX, J. W., AND DAVID LEE WOOD. 1955. Some roles of associated fungi in the life history of the bark beetle, *Ips paraconfusus* (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 18: 439. (ec hb).
- FOX, J. W., DAVID LEE WOOD, AND JAMES H. CANE. 1955. Incipient premating barriers to reproductive isolation in two sibling *Ips* species (Coleoptera: Scolytidae) from southern California: miscegenation persists. International Congress of Entomology, Proceedings 18: 429. (hb).
- FRANCKE, WITTKO. 1955. Identification and synthesis of new pheromones. *Agriculture, Ecosystems and Environment* 21: 21-30. (bv ms).
- FRANCKE, WITTKO, J. BARTELS, H. SCHMUTZENHOFER, U. KOHNLE, AND JEAN PIERRE VITE. 1955. The odor-bouquet of *Ips schmutzenhoferi* Holschuh (Col.: Scol.). *Zeitschrift für Naturforschung, C (Biosciences)* 43: 955-960. (bv).
- FRANCKE, WITTKO, M. L. PAN, J. BARTELS, W. A. KONIG, JEAN PIERRE VITE, S. KRAWIELITZKI, AND U. KOHNLE. 1956. The odour bouquet of three pine engraver beetles (*Ips* spp.). *Journal of Applied Entomology* 101: 453-461. (bv).
- FRANCKE, WITTKO, M. L. PAN, W. A. KOENIG, K. MORI, P.

- PAUPOOMCHAREON, H., HEUER, AND JEAN PIERRE VITE. 1957. Identification of "pityol" and "grandisol" as pheromone components of the bark beetle, *Pityophthorus pityographus*. *Naturwissenschaften* 74: 343-345. (bv).
- FRANSEN, JACOBUS JOHAN. 1933. De kleine iepenspintkever *Scolytus (Eccoptogaster) multistriatus* Mrsh. als verbreider der iepenziekte. Comité Inzake Bestuidering en Bestrijding van de Iepenziekte, Mededeeling 13: 35-43. (cn).
- *FRANZ, H. 1974. Die Nordost-Alpen im Spiegel ihrer Landtierwelt, Band IV [Scolytidae, p. 660-680]. Universitätsverlag Wagner, Innsbruck-München. 707 p. ().
- FREDERICKS, S. E., AND M. J. JENKINS. 1955. Douglas-fir beetle (*Dendroctonus pseudotsugae* Hopkins, Coleoptera: Scolytidae) brood production on Douglas-fir defoliated by western spruce budworm (*Choristoneura occidentalis* Freeman, Lepidoptera: Tortricidae) in Logan Canyon, Utah. *Great Basin Naturalist* 45: 348-351. (lb).
- FRENCH, JOHN RICHARD JOSEPH, PETER J. ROBINSON, AND GEORGE MINKO. 1952. Gamma irradiating elm billets reduces their attractiveness to the smaller elm bark beetle, *Scolytus multistriatus* (Marshall). *Zeitschrift für Angewandte Entomologie* 94: 175-179. (cn ec).
- FRENCH, JOHN RICHARD JOSEPH, PETER J. ROBINSON, GEORGE MINKO, AND PETER J. PAHL. 1954. Response of the European elm bark beetle, *Scolytus multistriatus*, to host bacterial isolates. *Journal of Chemical Ecology* 10: 1133-1149. (ec).
- FROST, C. A. 1912. Collecting Coleoptera in a Maine sawmill yard. *Canadian Entomologist* 44: 304-308. (ds).
- FRY, J. M. 1959. Natural enemy databank 1957. A catalog of natural enemies of arthropods derived from records in the CIBC Natural Enemy Databank. CAB International Institute of Biological Control, CAB International, Wallingford, Oxon. 155 p. (ec).
- FU, HUI-EN, BIAO WU, AND XUE-CHENG NING. 1954. A preliminary experiment on attraction with pheromone of bark beetles. *Kunchong Zhishi [Insect Knowledge]* 21: 213-215. (bv).
- FU, W. J., Y. T. QI, H. G. QIU, B. J. SHEN, AND X. D. WU. 1955. Studies on the aggregation pheromone of *Ips subelongatus*. I. Characteristics of aggregation behaviour and the extraction of active materials [In Chinese, English summary]. *Contributions from Shanghai Institute of Entomology* 8: 61-66. (bv).
- FUCHS, ANTON GILBERT. 1906. Nachtrag zur ersten Veröffentlichung über die Borkenkäfer Karntens. *Naturwissenschaftliche Zeitschrift für Land- und Forstwirtschaft* 4: 291-301. (lb).
- *_____. 1931. Die Genera: 1. *Rhabditolaimus* Fuchs, 2. *Neodiplogaster* Cobb, 3. *Tylenchodon* Fuchs. *Centralblatt für das Gesamte Forstwesen* 57: 177-194. ().
- FUCHS, M. G., AND JOHN HARVEY BORDEN. 1955. Pre-emergence insecticide applications for control of the mountain pine beetle *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Entomological Society of British Columbia, Journal* 52: 25-28. (cn).
- FUHRER, ERWIN. 1985. Air pollution and the incidence of forest insect problems. *Journal of Applied Entomology* 99: 371-377. (ec).
- FULLER, LLOYD R., AND D. W. JOHNSON. 1955. Evaluation of a survey technique to detect mortality caused by root diseases and bark beetles on the San Juan National Forest, 1952. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Biological Evaluation R2-S5-1. 8 p. (cn).
- *FURNISS, MALCOLM MACFARLANE, AND R. CAMPOS BOLANOS. 1954. Comparative adult anatomy of sympatric populations of *Dendroctonus rhizophagus* Thomas & Bright and *Dendroctonus valens* LeConte in Chihuahua, Mexico. Pages 97-105 in *Proceedings 2nd simposio nacional sobre parasitologia forestal*, 17-20 Febrero 1952, Cuernavaca, Mexico. ().
- FURNISS, MALCOLM MACFARLANE, AND J. B. JOHNSON. 1957. List of Idaho Scolytidae (Coleoptera) and notes on new records. *Great Basin Naturalist* 47: 375-382. (lb ds).
- _____. 1959. Description of the gallery and larva of *Dendroctonus punicatus* LeConte (Coleoptera: Scolytidae). *Canadian Entomologist* 121: 757-762. (lb tx).
- FURNISS, MALCOLM MACFARLANE, H. SOLHEIM, AND E. CHRISTIANSEN. 1955. Mechanism of transmission of the pathogenic blue-stain fungus, *Ophiostoma polonicum*, by *Ips typographus* in Norway spruce. *International Congress of Entomology, Proceedings* 15: 430. (ec).
- FURNISS, MALCOLM MACFARLANE, J. Y. WOO, MARK A. DEYRUP, AND THOMAS HARRIS ATKINSON. 1957. Prothoracic mycangium on pine-infesting *Pityophorus* spp. (Coleoptera: Scolytidae). *Entomological Society of America, Annals* 50: 692-696. (ay ec).
- FURUTA, KIMITO. 1959. A comparison of endemic and epidemic populations of the spruce beetle (*Ips typographus japonicus* Niisima) (Coleoptera: Scolytidae) in wind damaged forest in Hokkaido. *Journal of Applied Entomology* 107: 259-295. (bv ec).
- FURUTA, KIMITO, HIDEKI MORI, YUKIO TERASAKI, AND IKUO TAKAHASHI. 1956. An experiment for the control of *Ips typographus japonicus* Niisima (Coleoptera: Scolytidae) using pheromone traps [In Japanese]. *Japanese Forestry Society* 68(2): 75-77. (lv cn).
- FURUTA, KIMITO, IKUO TAKAHASHI, SYOICHI ANDO, AND MAKOTO INOUE. 1955. Reproduction and mass trapping of *Ips typographus japonicus* Niisima (Coleoptera: Scolytidae) in wind damaged forest in Hokkaido. *Bulletin of the Tokyo University Forests* 74: 39-65. (cn).

G

- GABEEV, V. N., AND E. V. GNAT. 1956. Silvicultural and physiological characteristics of trees damaged by *Dendroctonus* [In Russian, English summary]. *Izvestiya Sibirskogo Otdeleniya Akademii Nauk SSSR* 3: 24-25. (cn).
- *GADEK, K. 1950. Die Entomofauna der Weisstanne (*Abies alba* Mill.) in den polnischen Mittelgebirge (Gory Swietokrzyskie). *Acta Musei Reginalhradecensis* 5. A Supplementum: 35-45. ().
- *_____. 1957. Analiza szkodliwych czynników kształtujących stan zagrożenia drzewostanów Babiogorskiego Parku Narodowego [The analysis of harmful factors influencing the degree of danger of tree stands in the Babia Gora National Park]. *Ochrona przyrody* 45: 210-235. ().
- GAEDIKE, R. 1959. Book review: S. L. Wood and D. E. Bright, Jr., 1957. A catalog of Scolytidae and Platypodidae (Coleoptera), Part 1: Bibliography. *Coleopterists Bulletin* 43: 120. (ds tx ms).

- GAL, T., S. KISS, O. BORSODI AND A. ELEKES. 1955. Integrált vedekezes a betűzész (Ips typographus L.) ellen [Integrated protection against the typographer (Ips typographus L.)]. Növényvédelem 21: 487-490. (cn ec).
- GALVAO, JOSE MARTINS DE MIRA. 1956. Novo processo de luta contra o caruncho da oliveira. Direccao Geral dos Servicos Agricolas, Brigada Tecnica da XIV Regiao-Beja, Folha de Divulgacao No. 48, Serie III, No. 10, 14 p. (cn).
- GAMBILIEL, HERVE A., REN G. GATES, MARGARET K. CAFFEY, MOQUIN AND TIMOTHY D. PAINE. 1955. Variation in the chemistry of loblolly pine in relation to the infection by the blue-stain fungus. Pages 177-185 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56, 383 p. (cn ec).
- GANGLBAUER, LUDWIG. 1882. Beitrage zur Kenntnis der Coleopteren-Fauna des Erzherzogthumes Osterreich. Wiener Entomologische Zeitung 1: 118-120. (ds).
- GARA, ROBERT IRE. 1955. Interactions between fires, fungi, mountain pine beetles, and lodgepole pine in south-central Oregon. Northwest Environmental Journal 4: 355-358. (ec).
- GARA, ROBERT IRE, J. K. AGEE, W. R. LITCKE, AND D. R. GEISLER. 1986. Fire wounds and beetle scars. Journal of Forestry 84: 47-50. (cn).
- GARA, ROBERT IRE, W. R. LITCKE, J. K. AGEE, D. R. GEISLER, J. D. STUART AND C. H. DRIVER. 1955. Influence of fires, fungi and mountain pine beetle on development of a lodgepole pine forest in south-central Oregon. Pages 153-162 in D. M. Baumgartner et al., Lodgepole pine: the species and its management. Symposium proceedings. Cooperative Extension, Washington State University, Pullman, Washington. (cn).
- GARRWAY, ERIC. 1986. The biology of Ips calligraphus and Ips grandicollis (Coleoptera: Scolytidae) in Jamaica. Canadian Entomologist 118: 113-121. (hb ds).
- *GASPAR, C., AND C. VERSTRAETEN. 1972. Recherches sur l'ecosysteme foret. Serie C: Le chenais a Galeobdolon et a Ovalis de Mesnil-Eglise (Ferage). Contribution 26. Biocenose des coleopteres. Bulletin Societe Royale Scientifique Liege 41: 227-249. ().
- GAST, SANDRA J., MALCOLM MAC FARLANE FURNESS, JAMES B. JOHNSON, AND MICHAEL A. IVIE. 1959. List of Montana Scolytidae (Coleoptera) and notes on new records. Great Basin Naturalist 49: 351-356. (ds).
- GAST, SANDRA J., MOLLY WILFORD STOCK AND MALCOLM MAC FARLANE FURNESS. 1955. A demonstration of host attraction in Ips pini (Say) (Coleoptera: Scolytidae) in Idaho. International Congress of Entomology, Proceedings 18: 429. (bv).
- *GATES, D. E., AND L. L. PETERS. 1962. Insects in Kansas. Kansas State University Extension Service B-94. ().
- GAUTHIEAU, E. J. 1970. Annual district report: MacKenzie District, 1969. Pages 43-52 in Annual district reports of the Forest Insect and Disease Survey: Alberta-Northwest Territories-Yukon Region, 1969. Canada Department of Fisheries and Forestry, Canadian Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-30. (cn).
- GWYALIS, V. M., AND B. YU YAKATIS. 1987. Application of reduced doses of attractant for catching bark beetles (Ips typographus L.) [In Russian, English summary]. Khimorestseptsiya Nasekomym 9: 115-117. (bv cn).
- GEHRKEN, UNN. 1955. Physiology of diapause in the adult bark beetle, Ips acuminatus Gyll., studied in relation to cold hardiness. Journal of Insect Physiology 31: 909-916. (ay ec hb).
- _____. 1959. Supercooling and thermal hysteresis in the adult bark beetle Ips acuminatus Gyll. Journal of Insect Physiology 35: 347-352. (ay hb).
- GEISTHARDT, M. 1986. Ergänzende Meldungen zur kaferfauna der kapverdische Inseln (Coleoptera) [Scolytidae, p. 78-79]. Courier Forschungs-Institut Senckenberg 91: 69-80. (ds).
- _____. 1955. Tabellarische übersicht zur Verbreitung der Coleoptern aus dem Capverdischen Inseln [Scolytidae, p. 199-200]. Courier Forschungs-Institut Senckenberg 105: 193-210. (ds).
- GEMMA, J. N., S. S. WASTL AND G. C. HARTMAN. 1985. Antagonism of entomogenous fungal extract to the Dutch elm disease fungus, Ceratocystis ulmi. New York Entomological Society, Journal 93: 1109-1112. (ec).
- GENTRY, J. W. 1955. Beetle pest conditions. Coleopterists Bulletin 9: 25. (ds).
- GIBB, K. S., AND J. M. FISHER. 1986. Co-ordination of the life-cycle of Contortylenculus grandicollis (Nematoda: Allantonematidae) with that of its host Ips grandicollis (Scolytidae). Nematologica 32: 222-233.
- GIBBS, JOHN N., H. M. HEYBROEK, AND FRANCIS W. HOLMES. 1972. Aggressive strain of Ceratocystis ulmi in Britain. Nature [London] 236(5342): 121-122. (ec).
- GIBSON, KENNETH E. 1955a. Permanent mountain pine beetle population trend plots: an update 1955. United States Department of Agriculture, Forest Service, Northern Region (Missoula, Montana), Forest Pest Management Report 55-14, 8 p. (hb).
- _____. 1955b. Will trap trees reduce spruce beetle populations? Pages 42-43 in Thirty-sixth annual Western Forest Insect Work Conference: proceedings, 4-7 March 1955. Boulder Colorado. Canada Department of Environment, Canadian Forestry Service, Northern Forest Research Centre, Edmonton, Alberta, 54 p. (cn).
- _____. 1955a. Partial cutting of lodgepole pine stands to reduce tree losses to mountain pine beetle. Northwest Environmental Journal 4: 337-338. (cn).
- _____. 1955b. Permanent mountain pine beetle population trend plots. Northwest Environmental Journal 4: 336-337. (cn hb).
- _____. 1959a. Mountain pine beetle status—western United States. Pages 4-5 in G. D. Amman, Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262, 119 p. (cn).
- _____. 1959b. Partial cutting (sanitation thinning) to reduce mountain pine beetle-caused mortality. Pages 45-47 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262, 119 p. (cn).
- GIBSON, KENNETH E., AND D. D. BENNETT. 1955. Carbaryl prevents attacks on lodgepole pine by the mountain pine beetle. Journal of Forestry 83: 109-112. (cn).
- GIBSON, KENNETH E., W. E. BOUSFIELD, SANDRA J. GAST AND

- R. D. OAKES. 1955. Status and predicted trend of mountain pine beetle infestations in upper Yak River and Sullivan Creek drainages. Three Rivers and Rexford Ranger Districts. Kootenai National Forest, 1955. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana. Timber, Cooperative Forestry and Pest Management Report 75 p. (cn).
- GIBSON, KENNETH E., AND D. J. DOOLING. 1955. Status of insects and diseases on USA eastside national forest lands and adjoining state and private lands, 1954. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana. Forest Pest Management Report 55-16. 6 p. (cn).
- GIBSON, KENNETH E., S. K. HAGLE, AND S. KOHLER. 1987. Montana forest pest conditions and program highlights, 1986. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana. Cooperative Forestry and Pest Management Report 87-2. 29 p. (cn).
- GIBSON, KENNETH E., R. L. JAMES, ROBERT LADD LIVINGSTON, ET AL. 1984. Idaho forest pest conditions and program summary, 1983. United States Department of Agriculture, Forest Service, Northern and Intermountain Regions and Idaho Department of Lands, Coeur d'Alene, Idaho, Report 84-I. 40 p. (cn).
- GIBSON, KENNETH E., MARK D. MCGREGOR, AND GENE DOYLE ANMAN. 1985. Demonstration of the effectiveness of basal area cutting to reduce tree killing by the mountain pine beetle in ponderosa pine on Crow and Northern Cheyenne Indian Reservations, Montana 1984. Establishment Report. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 85-8. 9 p. (cn).
- GIBSON, KENNETH E., MARK D. MCGREGOR, AND R. D. OAKES. 1985. Mountain pine beetle infestation in ponderosa pine on Crow/ Northern Cheyenne Indian Reservations, Montana, 1984. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 85-9. 16 p. (cn).
- GIBSON, KENNETH E., AND R. D. OAKES. 1987. Mountain pine beetle status report, Northern Region, 1986. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 87-7. 57 p. (cn).
- GIL SANCHEZ, LUIS A. 1982. Escolítidos parasitos de coníferas de la provincia de Madrid. Ser. Forestal de Medio Ambiente y contra incendios. Dip. provincial de Madrid. [Internal document] 57 p. (cn).
- _____. 1984. Los Hylesininae Ibericos parasitos de coníferas (Col., Scolytidae). An. INIA, Ser. Forestal 5: 167-199. (ec ds).
- GIL SANCHEZ, LUIS A., AND JUAN A. PJARES ALONSO. 1986. Los escolítidos de las coníferas en la península ibérica. Monografías INIA N-53. Madrid. 194 p. (cn).
- GIL SANCHEZ, LUIS A., JUAN A. PJARES ALONSO, AND MANUEL G. DE VIEDMA. 1985. Estudios acerca de la atracción primaria en Scolytidae (Coleoptera) parasitos de coníferas. Boletín de la Estación Central de Ecología 14(27): 107-125. (bv cn).
- GILES, D. R. 1986a. Harvesting and processing beetle-killed pine. Pages 15-17 in R. W. Nielson, Harvesting and processing of beetle-killed timber. Proceedings of a symposium. Special publication, Forintek Canada Corp. SP-26. 53 p. (cn).
- _____. 1986b. Salvage and storage of beetle-killed timber. Pages 10-14 in R. W. Nielson, Harvesting and processing of beetle-killed timber. Proceedings of a symposium. Special publication, Forintek Canada Corp. SP-26. 53 p. (cn).
- GILLERFORS, COSTA. 1955. Second contribution to the coleopterous fauna of the Azores. Bocagiana 118. 6 p. (ds).
- * GILLETTE, CLARENCE PRESTON, AND G. LIST. 1920. [*Scolytus rugulosus*]. Page 21 in Colorado State Entomologist, Circular 28. 64 p. ().
- GIRLING, M. A., AND J. GREIG. 1955. A first fossil record of *Scolytus scolytus*, elm bark beetle. Its occurrence in elm decline deposits from London. England and the implications for Neolithic elm disease. Journal of the Archaeological Society 12: 347-352. (ds).
- * GLADITSCH, SIEGFRIED. 1976. Weitere kalerstfunde für südwest-Deutschland mit je einem erstfund für Mitteleuropa und Deutschland. 9. Beitrag zur Faunistik der südwestdeutschen Coleopteren. Beitr. naturk. Forsch. SudwDtl. 35: 149-167. ().
- GLOWACKA, B., M. WAJLAND, AND W. MILCZYŃSKI. 1955. Możliwości przyspieszenia terminu chemicznych zabiegów zabezpieczających nie korowane drewno sosnowe przed cetyncem wickszym [Possibilities of earlier control treatment for the protection of unbarked Scots pine timber against the large pine-shoot beetle]. Sylwan 132: 62-69. (cn).
- GNANAHARAN, R., V. V. SUDHENDRAKUMAR, AND K. S. NAIR. 1955. Protection of cashew wood in storage against wood borers. Material und Organismen 20: 65-74. (cn).
- GOBBI, GIOVANNI. 1959. Coleotterofauna xilofaga del pino loricato (*Pinus leucodermis*) sul Massiccio del Pollino (Coleoptera) [The beetle fauna of *Pinus leucodermis* from the massif of Mount Pollino] [English summary]. Bollettino Associazione Romana di Entomologia 43: 55-62 (1958). (ds).
- * GOGOLA, E. 1982. Roczna doba tazyby jedli z aspektu prevencie proti jedlovym podkornikom (Scolytidae) [Season of silver fir cutting from the aspect of the prevention against silver fir bark beetle]. Pages 25-31 in International Scientific Conference, 7-9 September 1982, VSLD Zvolen. ().
- _____. 1985a. *Platypus oxyurus*, new record (Coleoptera, Platypodidae): a new faunal species in Central Europe and its occurrence in Slovakia. Biologia (Bratislava) 40: 149-152. (ds).
- * _____. 1985b. Vyvoj kornika jedloveho (*Cryphalus piccae* Ratz., Col., Scol.) v laboratornych a prirodnych podmienkach [Development of *Cryphalus piccae* under laboratory and natural conditions]. Acta Facultatis Forestalis, Zvolen 27: 105-116. ().
- * _____. 1985c. Zintenzivnit ochranu jedle proti podkornikom. Les 41: 17-20. ().
- * _____. 1986. Jadrohlod jedlovy (*Platypus oxyurus* Duf.), nový drevokazný škodca jedle na Slovensku [Platypus oxyurus, a new ligniperdous pest of fir in Slovakia]. Lesnícky časopis, Bratislava 32: 17-24. ().
- GOHEEN, D. J., FIELDS WHITE COBB JR., DAVID LEE WOOD, AND D. L. ROWNEY. 1985. Visitation frequencies of some insect species on *Ceratocystis wagneri* infected and apparently healthy ponderosa pines. Canadian Entomologist 117: 1535-1543. (bv ec).
- GOLDHAMMER, D. S. 1985. Reproduction in the southern pine beetle, *Dendroctonus frontalis* (Coleoptera: Scolytidae): the symbiotic fungi component. Kansas Entomological Society, Journal 61: 361. (ec bv).
- GOLDHAMMER, D. S., F. M. STEPHEN, AND TIMOTHY D. PAINE

1959. Average radial growth rate and chlamydospore production of *Ceratocystis minor*, *Ceratocystis minor* var. *barrasii* and SJB 122 in culture. Canadian Journal of Botany 67: 3495–3505. (cc).
- GONZALES, M. D. 1958. Ciclo de vida y hábitos de la broca del fruto del café *Hypothenemus hampei* Ferr. 1956. en El Salvador [Life cycle and habits of the coffee berry borer in 1956 in El Salvador] [English summary]. Boletín de Promecafe 40: 11. (hb).
- COPE, B., AND S. K. ROY. 1956. Shade tree pest survey in the Dooars. Two and a Bud 33(1–2): 26–27. (cn ds).
- GORJACEVA, V. L., AND G. I. ANDREEVA. 1961. Himicheskaya bor'ba s koroedami tipografom dvojnikom na mestah ih zimovka [Chemical control of *Ips typographus* and *I. duplicatus* in their overwintering grounds]. Lesn. Z., Arhangel 4: 162–164. (cn).
- GORSUCH, C. S., AND J. KOVACH. 1956. Ambrosia beetles: problem or symptom? Pages 91–93 in E. I. Zenn, Stone fruit tree decline workshop: proceedings. United States Department of Agriculture, Agriculture Research Service. (cn).
- *GOSSARD, H. A. 1905. [*Scolytus rugulosus*]. Page 51 in Ohio Agricultural Experiment Station, Bulletin 195. ().
- *GOSSARD, H. A., AND WALTON. 1922. [*Scolytus rugulosus*]. Ohio Agricultural Experiment Station, Wooster, Bulletin 357: 83–126. ().
- GOULD, F. 1953. Book review: Review of bark beetles in North American conifers. Science 220: 708–709. (ins).
- GOULDING, HEATHER A., D. J. HALL, K. F. RAFFA, AND A. J. MARTIN. 1958. Wisconsin woodlands: identifying and managing pine pests in Wisconsin. Cooperative Extension Service, University of Wisconsin. Madison. (cn ms).
- GOYER, RICHARD ALBRICHT, G. L. LENHARD, T. EVAN NEBEKER, AND J. J. SCHMITT. 1955. Distinguishing immatures of insect associates of southern pine bark beetles. Pages 1–19 in Integrated pest management handbook. United States Department of Agriculture, Forest Service, Cooperative State Research Service, Washington, D.C. Agriculture Handbook 641. (cc).
- GOZIS, MAURICE PERROT DES. 1856. Recherche de l'espece typique de quelques anciens genres. Rectifications synonymiques et notes diverses [cited in Weise, Deutsche Entomologische Zeitschrift 31: 350–352 (1857)]. Montlucon, Impr. Herbin. 36 p. (cn).
- GRAHAM, DAVID A., AND G. MILLER. 1959. Canada/ U.S. mountain pine beetle/ lodgepole pine program 1951–1959. Pages 1–3 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- *GRASER, K. 1965. Ein Fund von *Xyloterus* (= *Trypodendron*) *domesticus* L. (Col.). Entomologische Berichten 1965: 38–39. ().
- GRASSE, P. P. 1949. Traite de Zoologie [Famille des Ipidae Reitter 1906, pages 955–958; Famille des Chapuisidae Blandford 1895, page 955; Famille des Platypodidae Chapuis 1865, p. 955]. IX: 771–1077. (tx).
- GRAY, D. R., AND JOHN HARVEY BORDEN. 1955. Ambrosia beetle attack on logs before and after processing through a dryland sorting area. Forestry Chronicle 61: 299–302. (cn).
- _____. 1959. Containment and concentration of mountain pine beetle (Coleoptera: Scolytidae) infestations with semiochemicals validation by sampling of baited and surrounding zones. Journal of Economic Entomology 52: 1399–1405. (bv cn).
- *GREEN, EDWARD ERNEST. 1916. [*Phloeotribus puncticollis* (= *setulosus*)]. Pages 605–636 in Third International Congress of Tropical Agriculture: proceedings, London. ().
- GREGOIRE, JEAN CLAUDE. 1955a. *Dendroctonus micans* in Belgium: the situation today. Pages 45–62 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium, Proceedings of seminar, 3–4 October 1954. 141 p. (cn).
- _____. 1955b. Host colonization strategies in *Dendroctonus*: larval gregariousness or mass attack by adults? Pages 147–154 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings. Canada Department of the Environment, Canadian Forest Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 240 p. (bv hb).
- _____. 1955. The greater European spruce beetle. Pages 455–478 in A. A. Berryman, Dynamics of forest insect populations: patterns, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec hb).
- GREGOIRE, JEAN CLAUDE, M. BAISIER, J. MERLIN, AND Y. NACCACHE. 1959. Interactions between *Rhizophagus grandis* (Coleoptera: Rhizophagidae) and *Dendroctonus micans* (Coleoptera: Scolytidae) in the field and the laboratory: their application for the biological control of *D. micans* in France. Pages 95–105 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ec).
- GREGOIRE, JEAN CLAUDE, AND J. MERLIN. 1955. *Dendroctonus micans*: the evolution of a brood system. Pages 80–86 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium, Proceedings of seminar, 3–4 October 1954. 141 p. (bv hb).
- GREGOIRE, JEAN CLAUDE, J. MERLIN, R. JAFFUEL, PH. DENIS, P. LAFONT, AND DANIEL SCHIVESTER. 1956. Elevage a petite et moyenne echelle du predateur *Rhizophagus grandis* Gyll. en vue de la lutte biologique contre *Dendroctonus micans* Kug. [Small-scale and medium-scale rearing of the pest *Rhizophagus grandis* Gyll. for biological control of *Dendroctonus micans* Kug.]. Revue Forestiere Francaise (Nancy) 35: 457–464. (cn).
- GREGOIRE, JEAN CLAUDE, J. MERLIN, J. M. PASTEELS, R. JAFFUEL, G. VOULAND, AND DANIEL SCHIVESTER. 1955a. Biocontrol of *Dendroctonus micans* by *Rhizophagus grandis* Gyll. (Col., Rhizophagidae) in the Massif Central (France): a first appraisal of the mass-rearing and release methods. Journal of Applied Entomology 99: 182–190. (cn ec).
- _____. 1955b. Mass-rearings and releases of *Rhizophagus grandis* in Lozere. Pages 122–125 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and the University Libre de Bruxelles,

- Bruxelles, Proceedings of seminar, 3-4 October 1984. 141 p. (ec).
- GREGOIRE, JEAN CLAUDE, AND J. M. PASTEELS. 1984. Biological control of bark beetles: (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium, Proceedings of seminar, 3-4 October 1984. 141 p. (bv ec).
- GRIES, GERHARD. 1955. Zur Frage der Dispersion des Buchdruckers (*Ips typographus* L.). Journal of Applied Entomology 99: 12-20. (ay bv hb).
- . 1956. Zur Bedeutung des Reifungsfrasses für die Dispersion des Kupferstechers, *Pityogenes chalcographus* L. (Coleoptera: Scolytidae) [On the importance of maturation feeding for the dispersal of the engraver beetle, *Pityogenes chalcographus*] [English summary]. Journal of Applied Entomology 73: 267-279. (ec hb).
- GRIES, GERHARD, A. LEUFVENS, J. P. LAFONTAINE, H. D. PIERCE, JR., JOHN HARVEY BORDEN, D. VANDERWEL, AND A. C. OEHELENSCHLAGER. 1990. New metabolites of alpha-pinene produced by the mountain pine beetle, *Dendroctonus ponderosae* (Coleoptera: Scolytidae). Insect Biochemistry 20: 365-371. (ay bv).
- GRIES, GERHARD, D. R. MILLER, M. J. SMIRLE, AND J. H. BORDEN. 1988. New semiochemicals in the pine engraver, *Ips pini* (Say) (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 18: 429. (bv).
- GRIES, GERHARD, R. NOLTE, AND W. SANDERS. 1989. Computer simulated host selection in *Ips typographus*. Entomologia Experimentalis et Applicata 53: 211-217. (cn ec).
- GRIES, GERARD, H. D. PIERCE, JR., B. S. LINDGREN, AND JOHN HARVEY BORDEN. New techniques for capturing and analyzing semiochemicals for scolytid beetles (Coleoptera: Scolytidae). Journal of Economic Entomology 81: 1715-1720. (bv ms).
- GRIJPM, PETER. 1955. *Dendroctonus micans* (Kug.) in the Netherlands: the situation today. Pages 35-42 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and the Université Libre de Bruxelles, Belgium, Proceedings of seminar, 3-4 October 1984. 141 p. (cn).
- GRIJPM, PETER, AND W. SCHIURING. 1984a. De ontwikkeling van de bastkevers, *Ips typographus*, *Ips cembrae* en *Pityogenes chalcographus* in niet-marktwaardig dunning shout. Nederlands Bosbouw tijdschrift 56: 159-164. (cn hb).
- . 1984b. Vergelijking van twee insecticide-lokstof combinaties op vangstammen voor bestrijding van de letterzetter (*Ips typographus*) [Comparison of two insecticide-pheromone combinations on trap trees for the control of the spruce bark beetle]. Nederlands Bosbouw tijdschrift 56: 295-302. (bv cn).
- GRILLO, H., AND M. ALVAREZ. 1984. Resultados parciales en la determinación de la entomofauna de la empresa de cítricos "Victoria de Giron" [Partial results of the survey of the entomofauna at the citrus enterprise "Victoria de Giron"] [English summary]. Centro Agrícola 11: 123-124. (cn).
- GRISDALE, D. G. 1962. Status of insects and tree diseases in the northern forest region. Pages 304-316 in Annual district reports of the Forest Insect and Disease Survey Ontario, 1962. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- GRITSCH, J. W. 1957. Control of bark beetles using baits. Pflanzenarzt 40: 3-4. (cn).
- *GROHMANN. 1913. [*Hylastes cunicularius*]. Tharandter Forstliches Jahrbuch, Berlin 64(4): 325-361. ().
- GROOT, PETER DE. 1986. Cone and twig beetles (Coleoptera: Scolytidae) of the genus *Conophthorus*: an annotated bibliography. Canada Department of the Environment, Forestry Service, Forest Pest Management Institute, Information Report FPM-X-76. 36 p. (cn ms).
- GROOTAERT, P., G. HAGHEBAERT, AND M. POLLET. 1957. Some *Madetera* (Diptera: Dolichopodidae) associated with Scolytidae (Coleoptera) from elms. Societe Royale Belge d'Entomologie, Bulletin & Annales 123: 350-352. (ec).
- GROSSMAN, JOEL. 1985. ESA '87 IPM highlights, Part 6. The IPM Practitioner 10(11-12): 7-10. (bv).
- GROVE, JOHN F. 1955. Metabolic products of *Phomopsis oblonga*. Part 2, Phomopsolide A and B, tiglic esters of two 6-substituted 5,6-Dihydro-5-hydroxypyran-2-ones. Journal of the Chemical Society, Transactions 4: S65-S70. (ms).
- GRUENWALD, M. 1986. Ecological segregation of bark beetles (Coleoptera, Scolytidae) of spruce. Journal of Applied Entomology 101: 176-187. (ec).
- GUANGJING, M. 1989. Air pollution and forest decline in China. Pages 51-54 in J. B. Bucher and I. Bucher-Wallin, Air pollution and forest decline. Fourteenth International Meeting for Specialists in Air Pollution Effects on Forest Ecosystems, Interlaken, Switzerland. 192 p. (ec).
- GUSTELEVA, L. A. 1982. The interaction of wood-decomposing insects with microorganisms [In Russian]. Pages 56-67 in A. S. Isaev, Konsortivnye Svyazi Dereva I Dendrofilnykh Nasekomykh. Novosibirsk, USSR [Canada Department of Environment, Translation OOENV TR-2274. 1983]. (cn ec).

H

- HAACK, ROBERT A. 1955. Voltinism and diurnal emergence-flight patterns of *Ips calligraphus* (Coleoptera: Scolytidae) in Florida. Florida Entomologist 68(4): 658-677. (ec hb).
- HAACK, ROBERT A., DANIEL M. BENJAMIN, AND KEVIN D. HAACK. 1983. Buprestidae, Cerambycidae, and Scolytidae associated with successive stages of *Agrilus bilineatus* (Coleoptera: Buprestidae) infestation of oaks in Wisconsin. Great Lakes Entomologist 16(2): 47-55. (ec).
- HAACK, ROBERT A., RONALD FORREST BILLINGS, AND ANDREAS M. RICHTER. 1989. Life history parameters of bark beetles (Coleoptera: Scolytidae) attacking West Indian pine in the Dominican Republic. Florida Entomologist 72: 591-603. (bv ec hb).
- HAACK, ROBERT A., JOHN L. FOLTZ, AND ROBERT CLEVELAND WILKINSON. 1984. Longevity and fecundity of *Ips calligraphus* (Coleoptera: Scolytidae) in relation to slash pine phloem thickness. Entomological Society of America, Annals 77: 657-662. (hb).
- . 1985. Effects of temperature and slash pine phloem thickness on *Ips calligraphus* life processes. Pages 102-113 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. General Technical Report SO-56. 353 p. (ec).

- HAACK, ROBERT A., AND W. J. MATTSO. 1959. They nibbled while the forests burned. *Natural History* (January): 56-57. (ms).
- HAACK, ROBERT A., AND FRANK SLANSKY, JR. 1957. Nutritional ecology of wood-feeding Coleoptera, Lepidoptera, and Hymenoptera. Pages 449-486 in F. Slansky, Jr. and J. G. Rodriguez. *Nutritional ecology of insects, mites, and spiders*. John Wiley, New York. (av ec).
- HAACK, ROBERT A., AND ROBERT CLEVELAND WILKINSON. 1957. Phoresy by *Dendrochernes pseudoscorpions* on Cerambycidae (Coleoptera) and Aulacidae (Hymenoptera) in Florida. *American Midland Naturalist* 117: 369-373. (ec).
- HAACK, ROBERT A., ROBERT CLEVELAND WILKINSON, JOHN L. FOLTZ. 1957. Plasticity in life-history traits of the bark beetle *Ips calligraphus* as influenced by phloem thickness. *Oecologia* (Berlin) 72: 32-38. (hb).
- HAACK, ROBERT A., ROBERT CLEVELAND WILKINSON, JOHN L. FOLTZ, AND JEFFRY A. CORNELL. 1957. Spatial attack pattern, reproduction, and brood development of *Ips calligraphus* (Coleoptera: Scolytidae) in relation to slash pine phloem thickness: a field study. *Environmental Entomology* 16: 425-436. (bv hb).
- HAANSTAD, J. D., AND D. M. MORRIS. 1955. Microbial symbiotes of the ambrosia beetle *Xyloterinus politus*. *Microbial Ecology* 11: 267-276. (ec).
- HABERMANN, M., AND SCHOFF. 1957. Untersuchungen zu Laborzucht, Jungkaferschupf und Adultfrass von *Scolytus intricatus* (Ratz.) (Col., Scolytidae) [Studies on laboratory rearing, emergence of young adults and feeding by adults of *Scolytus intricatus*] [English summary]. *Journal of Applied Entomology* 104: 519-525. (bv hb).
- _____. 1955. Freilanduntersuchungen zu Flugaktivität, Adultfrass und Bruterfolg von *Scolytus intricatus* (Ratz.) (Col., Scolytidae) [Field studies on the flight activity, adult feeding and brood success of *Scolytus intricatus*] [English summary]. *Journal of Applied Entomology* 106: 252-261. (hb).
- HAESSELBARTH, ERASMUS. 1955. Determination list of entomophagous insects 10. International Union for Biological Sciences, International Organization for Biological Control of Noxious Animals and Plants, West Palaearctic Regional Section. 61 p. (ec).
- HAGLE, SUSAN K., S. TUNNOCK, K. E. GIBSON, AND C. J. GILLIGAN. 1957. Field guide to diseases and insect pests of Idaho and Montana forests. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, State and Private Forestry. 123 p. (cn hb).
- HAIGHT, R. G., J. D. BRODIE, AND W. G. DAHMS. 1955. A dynamic programming algorithm for optimization of lodgepole pine management. *Forest Science* 31: 321-330. (cn).
- HAIN, FRED P., AND A. BEN ALYA. 1955. Interactions of the southern pine beetle with competitor species and meteorological factors. Pages 114-126 in S. J. Branham and R. C. Thatcher. *Symposium proceedings: Integrated pest management research*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (ec).
- HAIN, FRED P., AND S. P. COOK. 1955a. Host resistance and its utility for controlling southern pine beetle in North Carolina. Pages 77-83 in T. L. Payne and H. Saarenmaa. *Integrated control of scolytid bark beetles*. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- _____. 1955b. Host resistance and its utility for controlling southern pine beetle in North Carolina. *International Congress of Entomology, Proceedings* 15: 413. (cn).
- HAIN, FRED P., S. P. COOK, P. A. MATSON, AND K. G. WILSON. 1955. Factors contributing to southern pine beetle host resistance. Pages 154-160 in S. J. Branham and R. C. Thatcher. *Symposium proceedings: Integrated pest management research*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (ec).
- HAJEK, ANN E., AND DONALD L. DAHLSTEN. 1955a. Factors influencing the mortality of *Scolytus multistriatus* (Coleoptera: Scolytidae) in elm branches in California. *Canadian Entomologist* 117: 819-828. (bv ec).
- _____. 1955b. Insect and mite associates of *Scolytus multistriatus* (Coleoptera: Scolytidae) in California. *Canadian Entomologist* 117: 409-421. (ec).
- HALL, P. M. 1956. Mountain pine beetle symposium: proceedings, 16-18 April 1955, Smithers, British Columbia. British Columbia Ministry of Forests, Pest Management Report 7. 155 p. (cn ec hb).
- _____. 1959. The use of mountain pine beetle aggregation semiochemicals in British Columbia. Pages 101-107 in G. D. Amman. *Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (bv cn).
- HALL, RALPH CORBIN. 1952. Uses of remote sensing in forest pest damage appraisal. *Proceedings of a seminar held 5 May 1951, Edmonton, Alberta*. Government of Canada, Canadian Forestry Service, Northern Forestry Centre, Edmonton, Alberta, Information Report NOR-X-235. viii + 60 p. (cn).
- HALPERIN, J. 1954. The elm bark beetle [*Scolytus multistriatus*] and its control [In Hebrew]. *Can Vanof* 40: 120-124. (cn).
- _____. 1955. The main insect pests of pistacio in Israel [In Hebrew]. Pages 11-39 in S. Dover. *Third symposium of pistacio growers in Israel: proceedings, 8 June 1954*. Sede Boquer. Ben-Gurion University of the Negev, The Blaustein International Center for Desert Studies, Sede Boquer. (cn ec).
- _____. 1956a. The elm bark beetle *Scolytus kirschi* and its control in Israel. *EPPO Bulletin* 16: 593-595 (cn ec ds).
- _____. 1956b. Braconidae (Hymenoptera) associated with forest and ornamental trees and shrubs in Israel. *Phytoparasitica* 14: 119-135. (ec).
- _____. 1957. Practical application of research results on insect pests of ornamental and forest trees in Israel [Abstract of lecture: Third Israeli Convention of Agricultural Entomology, Bet Dagan, 9 April 1957]. *Phytoparasitica* 15: 151. (cn ec).
- HAMEL, DENNIS R. 1959. Registration status of bark beetle semiochemicals. Pages 118-119 in G. D. Amman. *Management of lodgepole pine to minimize the losses to the mountain pine beetle: symposium proceedings*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).

- HAMILTON, JOHN. 1891. Notes on Coleoptera, No. 7. Canadian Entomologist 23: 60–65. (hb ds).
- HAMILTON, W. DOUGLAS. 1967. Extraordinary sex ratios. Science 156: 477–488. (lb).
- _____. 1979. Wingless and fighting males in fig wasps and other insects. Pages 167–220 in M. S. Blum and N. A. Blum, Sexual selection and reproductive competition in insects. Academic Press, New York. 463 p. (hb).
- HAMILTON, W. DOUGLAS, PAWEŁ SVHIRA, AND CARLTON S. KOEHLER. 1988. Pine tree deaths by bark beetles. University of California, Cooperative Extension, Division of Agriculture and Natural Resources Leaflet 7069. 2 p. (cn).
- HANISCH, Meg A., H. DANIEL BROWN, AND EDWARD A. BROWN. 1983. Dutch elm disease management guide. United States Department of Agriculture, Forest Service, and Extension Service. Bulletin 1. 23 p. (cn).
- HANSEN, E. M., D. J. GOHEEN, P. F. HESSBURG, J. J. WITCOSKY, AND T. D. SCHWALTER. 1988. Biology and management of black-stain root disease in Douglas-fir. Pages 63–80 in T. C. Harrington and F. W. Cobb, Jr., Leptographium root diseases in conifers. American Phytopathological Society, St. Paul, Minnesota. vii + 149 p. (ec).
- HANSEN, T. E. 1986. Hibernation physiology of *Polygraphus poligraphus* larvae [In Russian, English summary]. Metsanduslikud murimused, Estonian SSR 21: 103–113. (ay lb).
- HARD, JOHN S. 1985. Spruce beetles attack slowly growing spruce. Forest Science 31: S39–S50. (cn).
- _____. 1987. Vulnerability of white spruce with slowly expanding lower boles on dry cold sites to early seasonal attack by spruce beetles in south-central Alaska. Canadian Journal of Forest Research 17: 428–435. (bv ec).
- _____. 1989. Sequence of trees attacked by spruce beetles in a mature even-aged spruce stand in south-central Alaska. Northwest Science 63: 5–12. (bv ec).
- HARD, JOHN S., AND EDWARD H. HOLSTEN. 1985. Managing white and Lutz spruce stands in south-central Alaska for increased resistance to spruce beetle. United States Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon, General Technical Report PNW-188. ii + 21 p. (cn ec).
- * HARDE, K. W. 1981. Der Kosmos-Kaferfuhrer. Kosmos Verlag, Stuttgart. ().
- * _____. 1984. A field guide in colour to beetles. Edited and with additional introductory material by P. M. Hammond. Octopus Books, London. 334 p. ().
- HARDING, SUSANNE, E. B. LAPIS, AND BRODER BEJER-PETERSEN. 1986. Observations on the activity and development of *Pityogenes chalcographus* L. (Col., Scolytidae) in stands of Norway spruce in Denmark. Journal of Applied Entomology 102: 237–244. (ec).
- HARDING, SUSANNE, AND H. P. RAVN. 1985. Seasonal activity of *Ips typographus* L. (Col., Scolytidae) in Denmark. Journal of Applied Entomology 99: 123–131. (hb).
- * HARRINGTON, THOMAS CHARLES. 1980. Release of airborne basidiospores from the pouch fungus *Cryptoporus volvatus*. Mycologia 72: 926–936. ().
- HARRINGTON, THOMAS CHARLES, FIELDS WHITE COBB, JR., AND J. LOWNSBERRY. 1985. Activity of *Hylastes nigrinus* a vector of *Verticicladiella wagneri*, in thinned stands of Douglas-fir. Canadian Journal of Forest Research 15: 519–523. (ec lb).
- HARRINGTON, WILLIAM HAGUE. 1894. An entomological trip to Copper Cliff, Ont. Canadian Entomologist 26: 9–16. (ds).
- HARRIS, JOHN W. E. 1960b. The balsam woolly aphid, *Adelges piceae* (Ratz.), in British Columbia. Canada Department of Agriculture, Research Branch, Bi-monthly Progress Report 16(2): 3–4. (cn).
- _____. 1971. Aerial photography (35 mm): aid to forest pest surveys. Canada Department of Fisheries and Environment, Canadian Forestry Service, Bi-monthly Research Notes 27: 20. (cn ms).
- _____. 1972. High-level photography for insect damage surveillance in British Columbia. First Canadian Symposium on Remote Sensing, Ottawa 1971: 145–148. (cn).
- _____. 1974a. ERTS imagery and small-scale photography in modern forest management. British Columbia Lands Service, Survey and Mapping Branch, Victoria. Users Handbook B.C. ERTS-1: 26–27. (cn ms).
- _____. 1974b. Small scale imagery in forest pest surveys in British Columbia. Canadian Surveyor 28: 155–161. (cn ms).
- HARRIS, JOHN W. E., A. F. DAWSON, AND R. G. BROWN. 1979. Detection des chablis, foyers potentiels d'infestation des scolytides, au moyen de techniques simples de photographie aeriennne. Canada Department of Fisheries and Forestry, Service des forets, Revue bimensuelle de recherches 35: 9–10. (cn ms).
- HARRIS, S. A. 1955. Beetles blanket woodlands. Louisiana Conservation 37: 28–29. (cn ms).
- HART, DENNIS. 1984. Economic analysis of treating a root disease-bark beetle complex, the cause of tree mortality on McCloud Flats, Shasta-Trinity National Forests. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84–10. (cn).
- HASTINGS, F. L., U. E. BRADY, AND A. S. JONES. 1989. Lindane and fenitrothion reduce soil and litter mesofauna on Piedmont and Appalachian sites. Environmental Entomology 18: 245–250. (cn).
- HATCH, MELVILLE HARRISON. 1939. List of the Coleoptera taken by Mr. George P. Engelhardt in Alaska in 1938. Brooklyn Entomological Society, Bulletin 34: 45–50. (ds).
- _____. 1941. A second supplement to the indices to the keys to and local lists of Nearctic Coleoptera. New York Entomological Society, Journal 49: 21–42. (tx ms).
- HAVERTY, MICHAEL I., AND PATRICK J. SHEA. 1985. Protection of blister rust-resistant western white pine cones from insect damage with permethrin and fenvalerate. Pages 246–250 in Conifer tree seed in the inland Mountain West: symposium proceedings, 5–6 August 1985, Missoula, Montana. Society of American Foresters 55–10. (cn).
- HAVERTY, MICHAEL I., PATRICK J. SHEA, AND RICHARD W. HALL. 1985. Effective residual life of carbaryl for protecting ponderosa pine from attack by the western pine beetle (Coleoptera: Scolytidae). Journal of Economic Entomology 78: 197–199. (cn).
- HAVERTY, MICHAEL I., PATRICK J. SHEA, AND LAWRENCE E. STIPE. 1988. Protection of disease resistant western white pine seed from insect damage. Western Journal of Applied Forestry 3: 18–20. (cn).
- HAWBOLDT, L. S. 1948. Forest insects of the season. Pages 35–81 in Nova Scotia Department of Lands and Forests, Report. Forest Entomology. (cn).
- HAWKSWORTH, FRANK G. 1985. Insect-dwarf mistletoe asso-

- ciations. Pages 49–50 in Thirty-sixth annual Western Forest Insect Work Conference: proceedings, Boulder, Colorado, 4–7 March 1955. Canada Department of Environment, Canadian Forestry Service, Northern Forest Research Centre, Edmonton, Alberta. 54 p. (bv cn).
- *HAY, AND WOOTE. 1955. [*Scolytus rugulosus*]. United States Department of Agriculture, Forest Service, Central States Forest Experiment Station, Technical Paper 145: 6–9, 11–12. ().
- HAYES, A. J., AND D. R. PARIHAR. 1957. Some observations on the breeding biology of *Scolytus scolytus* (Coleoptera: Scolytidae), a vector of Dutch elm disease in Scotland. *Arboricultural Journal* 11: 333–344. (hb).
- HE, L. F., Y. P. LI AND W. J. FU. 1958. Method for sexing live beetles of *Ips subelongatus* (Coleoptera: Scolytidae). Contributions from Shanghai Institute of Entomology 8: 73–76. (ay).
- *HEADLEE, THOMAS J. 1915. [*Scolytus rugulosus*]. Pages 631–695 in New Jersey Agricultural Experiment Station, New Brunswick, Report for 1914. ().
- HEDDEN, ROY L. 1953. Evaluation of loblolly pine thinning regimes for reduction of losses from southern pine beetle attack. United States Department of Agriculture, Forest Service, Southeastern Forest Experiment Station (Research Triangle Park, North Carolina), General Technical Report SE-24: 371–375. (cn).
- _____. 1955. Simulation of southern pine beetle-associated timber loss using CLEMBEETLE. Pages 288–291 in S. J. Barnham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- HEDDEN, ROY L., AND ROGER P. BELANGER. 1955. Predicting susceptibility to southern pine beetle attack in Coastal Plain, Piedmont and Southern Appalachians. Pages 232–238 in S. J. Barnham and R. C. Thatcher, Integrated pest management research: symposium proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (ec).
- HEDDEN, ROY L., AND P. L. LORIO, JR. 1955. Rating stand susceptibility to southern pine beetle attack on national forests in the Gulf Coastal Plain. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, Research Paper SO-221. 5 p. (cn).
- HEDDEN, ROY L., AND T. EVAN NEEBEK. 1954. Integrated forest pest management in pine stands (0–5 years): insects. Pages 39–53 in S. J. Barnham and G. D. Hertel, Integrated forest pest management: symposium proceedings, Athens, Georgia. [Publisher?]. (cn).
- HELIOVAARA, KARI AND ARJA LILJA. 1959. Sinistajasienet ja kaarnakuoriaiset Pohjois-Euroopassa [Blue stain fungi and the bark beetles in northern Europe] [In Finnish, English summary]. *Luonnon Tutkija* 93: 136–143. (ec).
- HELIOVAARA, KARI AND K. PUUKKO. 1956. Note on *Ips acuminatus* (Coleoptera) on the south coast of Finland. *Notulae Entomologicae* 66: 179. (ds).
- HELLAND, INGE S., O. ANDERBRANT AND J. M. HOFF. 1959. Modelling bark beetle flight: a review. *Holarctic Ecology* 12: 427–431. (cn hb).
- HELLQVIST, C. 1954. Produktion av storre margborre i tallstubb [Production of *Tomicus piniperda* in pine stump]. *Sveriges Skogsvardsforbunds Tidskrift*, nr. 1. (cn hb).
- HELLRIGL, K. 1955. Über Borkenkäfer (Col., Scolytidae) in Zweigen der Zirbe (*Pinus cembra*) in Südtirol [Observations on bark beetles in branches of *Pinus cembra* in south Tyrol]. *Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz* 58: 108–110. (ec).
- HELLRIGL, K., AND W. SCHWENKE. 1955. Begleitinsekten in Buchdrucker-Pheromonfallen in Südtirol [Accompanying insects in *Ips typographus* pheromone traps in south Tyrol]. *Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz* 58: 47–50. (ec).
- HENSHAW, SAMUEL. 1855. List of Coleoptera of America, north of Mexico. American Entomological Society, George B. Cresson, Philadelphia. 4 + 161 p. (ds).
- HEQVIST, KARL JOHAN. 1969. Notes on Cerocephalini with descriptions of new genera and species (Hymenoptera: Chalcidoidea, Pteromalidae). *Entomological Society of Washington, Proceedings* 71: 449–467. (ec).
- HERBAUT, C., K. LACHICER-SILLOU, J. Y. LE GALLO, AND M. EL-ILDRISI-EL AAMMARI. 1957. Etude du cycle biologique de *Phloeotribus scarabaeoides* Bernard (Coleoptera, Scolytidae) dans la region de Marrakech [A study of the biological cycle of *Phloeotribus scarabaeoides* in the Marrakesh region] [English summary]. *Bulletin de l'Institut Scientifique, Université Mohammed* 11: 187–192. (hb).
- *HERBERT, FRANK B. 1919. [*Phloeosinus* spp.]. *Journal of Economic Entomology* 12: 337. ().
- _____. 1920b. [*Phloeosinus cristatus*]. United States Department of Agriculture, Bulletin 535. 22 p. ().
- HERGER, P. 1955a. Zur Insektenfauna der Umgebung des Brisen-Haldgrates, 1200–2400 m, Kanton Nidwalden. VII, Coleoptera 3: Scolytidae und Curculionidae. *Entomologische Berichte, Luzern* 13: 93–95. (ds).
- _____. 1955b. Zur Insektenfauna der Umgebung der Vogelwarte Sempach, Kanton Luzern. XV, Coleoptera 3: Cerambycidae, Chrysomelidae, Scolytidae und Nachträge. *Entomologische Berichte Luzern* 13: 77–80. (ds).
- _____. 1955c. Zur Insektenfauna von Pilatus-Kulm, 2060 m, Kanton Nidwalden. VIII, Coleoptera 3: Scolytidae und Curculionidae. *Entomologische Berichte Luzern* 13: 91–92. (ds).
- HERITAGE, STEWART, S. COLLINS, AND H. F. EVANS. 1959. A survey of damage by *Hylobius abietis* and *Hylastes* spp. in Britain. Pages 36–42 in R. I. Alfaro and S. C. Glover, Insects affecting reforestation: biology and damage. Government of Canada, Department of Forestry, Pacific Forestry Centre, Victoria, British Columbia. 256 p. (cn).
- HERTEL, GERARD D., SUSAN J. BRANHAM AND KENNETH M. SWAIN, SR. 1955. Technology transfer in integrated forest pest management in the South. United States Department of Agriculture, Forest Service, Southeastern Forest Experiment Station (Research Triangle Park, North Carolina), General Technical Report SE-34. 77 p. (cn).
- HESJEDAL, K., AND T. EDLAND. 1958. Angrep av lauvtrebarkbille i frukthagen [Attack of the deciduous tree bark beetle in fruit orchards] [In Norwegian, English summary]. *Gartneryrket* 78: 115–117. (cn ec hb).

- HEWITT, C. G. 1915. Report of the Dominion Entomologist for the year ending March 31, 1915. Canada Department of Agriculture, Ottawa, Ontario. (cn).
- *HEYDEN, LUCAS FRIEDRICH JULIUS DOMINICUS VON. 1884. Beitrag zur Coleopterenfauna der Insel Askold und anderer Theile des Amurgebietes. Deutsche Entomologische Zeitschrift 28: 273-300. (.)
- HICKS, ROY R., JR., JACK E. COSTER, AND GARLAND N. MASON. 1957. Forest insect hazard rating. Journal of Forestry 55: 20-25. (cn).
- HIGHLEY, L., AND T. A. TATTAR. 1955. *Leptographus terabrantis* and black turpentine beetles associated with blue stain and mortality of black and Scots pines on Cape Cod, Massachusetts. Plant Disease 69: 525-530. (ec).
- . 1957. Patterns of bluestain discoloration and associated organisms in Japanese black and Scots pines on Cape Cod, Massachusetts. Arboricultural Journal 11: 105-113. (ec).
- HILBURN, D. J., AND R. D. GORDON. 1959. Coleoptera of Bermuda. Florida Entomologist 72: 673-692. (ds).
- HILL, DENNIS S. 1957. Agricultural insect pests of temperate regions and their control. Cambridge University Press, Cambridge. 659 p. (cn ds hb).
- *HINKLEY, ALDEN DEXTER. 1963. Trophic records of some insects, mites, and ticks in Fiji. Department of Agriculture, Fiji, Bulletin 45. 116 p. (.)
- *HINTON, HOWARD EVEREST. 1951. Biology of insect eggs. Pergamon Press. 1125 p. (.)
- HINTZE-PODUEAL, C., AND A. DRUSCHKE. 1955. Untersuchungen zur Besiedlungsdichte und Parasitierung des kleinen bunten eschenbastkafers *Leperisinus varius* (F.) [Investigations on the population density and parasitic infestation of *Leperisinus varius*] [English summary]. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 61: 241-245. (ec hb).
- HIRSCHMANN, WERNER, AND W. RUHM. 1953. Ein "Haustier" des Buchdruckers? Acarologie 30: 22-23. (ec).
- HO, Y. C., H. K. KAYA, AND P. SHEA. 1959. *Parasitylenchus orthotomici* sp. n. (Tylenchia: Allantonematidae) from *Orthotomicus angulatus* (Coleoptera: Scolytidae), with notes on parasitism. Korean Journal of Applied Entomology 28: 1-3. (ec).
- *HO, Y. C., AND S. W. KUN. 1959. Supplementary notes on the bark and ambrosia beetles of Korea. Korean Journal of Applied Entomology 28: 4-9. (.)
- HOBBS, A. J. 1955. Effects of air-photo scale on early detection of mountain pine beetle infestations. Forestry Abstracts 46: 636-637. (cn).
- HOBSON, K. R., I. KUBO, AND DAVID LEE WOOD. 1955. Host colonization behavior of the red turpentine beetle *Dendroctonus valens*. International Congress of Entomology, Proceedings 18: 439. (bv).
- *HOCHMUT, R. 1952. La proteccion contra las plagas y enfermedades forestales en Cuba. Silvicultura Tropica et Subtropica 9: 95-111. (.)
- *HOCHMUT, R., AND V. NOVAK. 1958. Moznosti aktivizace nekterych kalamitnich hynzich skudcu imisnim zatizenim lesnich porostu. Lesnictvi 34(LX1), (1): 93-96. (.)
- HODGES, GARY V. 1955. The High Country Project. Pages 26-28 in Thirty-sixth annual Western Forest Insect Work Conference: proceedings, 4-7 March 1955, Boulder, Colorado. Canada Department of Environment, Canadian Forestry Service, Northern Forest Research Centre, Edmonton, Alberta. 54 p. (cn).
- HODGES, JOHN DEAVOURS, T. EVAN NEBEKER, C. A. BLANCHE, R. HONEA, T. H. FISHER, AND T. P. SCHULTZ. 1959. Southern pine beetle-microorganisms-host interactions: influence of compounds produced by *Ceratocystis minor*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-74: 567-572. (ec).
- HODGES, JOHN DEAVOURS, T. EVAN NEBEKER, J. D. DEANGELIS, AND C. A. BLANCHE. 1955. Host/beetle interactions: influence of associated microorganisms, tree disturbance, and host vigor. Pages 161-165 in S. J. Branham and R. C. Thatcher. Integrated pest management research: symposium proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station (New Orleans, Louisiana), General Technical Report SO-56. 383 p. (ec).
- HODGES, JOHN DEAVOURS, T. EVAN NEBEKER, J. D. DEANGELIS, B. L. KARR, AND C. A. BLANCHE. 1955. Host resistance and mortality: a hypothesis based on the southern pine beetle-microorganism-host interactions. Entomological Society of America 31: 31-35. (cn ec).
- HODGKINSON, ROBERT SIDNEY. 1956. Use of trap trees for spruce beetle management in British Columbia, 1979-1984: a review with recommendations. British Columbia Ministry of Forests, Pest Management Report 5. 39 p. (cn).
- HOEBEKE, E. RICHARD. 1959. *Pityogenes bidentatus* (Herbst), a European bark beetle new to North America (Coleoptera: Scolytidae). New York Entomological Society, Journal 97: 305-308. (ds).
- *HOERNER. 1929. [*Phloeotribus liminaris*]. Maryland Agricultural Experiment Station, Bulletin 310: 449-465. (.)
- HOFACKER, THOMAS H., ROBERT C. LOOMIS, AND R. T. GILSTRAP. 1955. Forest insect and disease conditions in the United States, 1957. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C. 102 p. (cn).
- HOFACKER, THOMAS H., ROBERT C. LOOMIS, AND SUSAN M. TUCKER. 1957. Forest insect and disease conditions in the United States, 1956. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C. 94 p. (cn).
- HOFACKER, THOMAS H., ROBERT C. LOOMIS, AND A. J. WORRAL. 1959. Forest insect and disease conditions in the United States, 1955. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C. 102 p. (cn).
- HOFFARD, WILLIAM H. 1955. Southern pine beetle. A would-be manager of southern forests. Pages 32-37 in R. C. Loomis, and Thomas H. Hofacker, What else is growing in our forests? Insect and disease conditions in the United States 1979-1983. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C., General Technical Report WO-46. 93 p. (cn).
- HOFFMAN, J., A. KEYSOR, A. KNAPP, AND J. WEATHERBY. 1957. Forest insect and disease conditions, Intermountain Region. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report. 22 p. (cn).
- . 1958. Forest insect and disease conditions, Intermountain Region, 1957. United States Department of Agriculture, Forest Service, Forest Pest Management, Intermountain Region, Ogden, Utah. 20 p. (cn).
- HOFFMANN, ADOLPHE. 1952a. Repertoire analytique des

- especes animales nuisibles aux cultures en France (Metropole et Departements d'outre-mer) avant present d'interessantes particularites en 1950. *Annales des Epiphyties* 1: 103-109. (cn ds).
- HOGBERG, H. E., E. HEDENSTROM, R. ISAKSSON, AND A. B. WASSGREN. 1957. Preparation of the four stereoisomers of chalcogran, pheromone components of *Pityogenes chalcographus* and of both enantiomers of gamma-caprolactone, pheromone component of *Trogoderma granarium*. *Acta Chemica Scandinavica, B (Organic Chemistry and Biochemistry)* 41: 694-697. (bv ms).
- *HOGUE, C. L., AND S. E. MILLER. 1980. Entomofauna of Cocos Island, Costa Rica (including Arachnida and other terrestrial Arthropoda): first progress report on studies. 37 p. ().
- HOLMES, FRANCIS WILLIAM. 1956. Recorded Dutch elm disease distribution in North America as of 1955. *Plant Disease Reporter* 40(4): 351-353. (cn).
- _____. 1973. Preliminary research report on chemical treatments for Dutch elm disease. University of Massachusetts, Shade Tree Laboratories. Photocopied report dated 10 August 1973. 2 p. (cn).
- _____. 1974. Benlate and Dutch elm disease. *Trees* 1974(April-June): 28-29. (cn).
- _____. 1976a. Benomyl phosphate and Dutch elm disease, H. University of Massachusetts, Shade Tree Laboratories. Report issued with a reprint copy of *Phytopathology* 62: 939-940 [1972]. 4 p. (cn).
- _____. 1976b. Benomyl phosphate and the Dutch elm disease, III. Pages 17-19 in 31st annual conference of Dutch elm disease; proceedings, Waltham, Massachusetts. (cn).
- _____. 1976c. Benomyl phosphate versus "Benlate" benomyl in elms already afflicted with Dutch elm disease. *American Phytopathological Society, Proceedings* 3: 266. (cn).
- _____. 1976d. The American elm fights back. *Horticulture Magazine* 1976(January): 72-75. (cn).
- _____. 1980. Bark beetles, *Ceratocystis ulmi*, and Dutch elm disease. Pages 133-147 in *Vectors of plant pathogens*. Academic Press. (cn).
- _____. 1980. Dutch elm disease. *Northeastern States Extension Plant Pathologists* [Undated circular issued about 1980]. 1 p. (cn).
- HOLSTEN, EDWARD H. 1981. Spruce beetle: Chugach National Forest and adjacent lands. United States Department of Agriculture, Forest Service, State and Private Forestry, Alaska Region, Anchorage, Alaska. Biological Evaluation R10-81-1. 15 p. (cn).
- _____. 1990. Spruce beetle activity in Alaska: 1920-1989. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, Forest Pest Management Report R10-90-18. 28 p. (cn).
- HOLSTEN, EDWARD H., A. EGLITIS, AND THOMAS A. LAURENT. 1981. Alaska Region. Pages 65-71 in H. D. Brown and P. W. Orr, *Forest insect and disease conditions in the United States, 1979*. United States Department of Agriculture, Forest Service, Washington, D.C., General Technical Report WO-20. 91 p. (cn).
- HOLSTEN, EDWARD H., P. E. HENNON, AND RICHARD ALLEN WERNER. 1985. Insects and diseases of Alaskan forests. United States Department of Agriculture, Forest Service, Alaska Region, Report 151. Juneau, Alaska. 217 p. (cn).
- HOLSTEN, EDWARD H., AND RICHARD ALLEN WERNER. 1985a. Evaluation of a controlled release formulation of methylcyclohexenone (MCH) in preventing spruce beetle attacks in Alaska. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska. State and Private Forestry, Report R10-9. 9 p. (cn).
- _____. 1985b. Evaluation of a controlled release formulation of methylcyclohexenone (MCH) in preventing spruce beetle attacks in Alaska. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska. State and Private Forestry, Report R10-12. 9 p. (cn).
- _____. 1987. Engraver beetles in Alaska forests. United States Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, Oregon. undated circular. GPO 795-322. 6 p. (cn).
- HOLSTEN, EDWARD H., K. P. ZOGAS, R. L. WOLFE, AND RICHARD ALLEN WERNER. 1979. Resurrection Creek spruce beetle infestation: a three year interim report. United States Department of Agriculture, Forest Service, Alaska Region, Anchorage, Alaska, State and Private Forestry, Report. (cn).
- HOLTHUIS, L. B. 1976. Comment on the proposed use of the plenary powers to conserve the generic name *Phloeotribus* Latreille, 1804. *Bulletin of Zoological Nomenclature* 32: 208-208. (tx).
- *HOLZEL, EMIL. 1936. II. Nachtrag zum Verzeichnis der bisher in Karnten beobachteten Kafer. *Carinthia* 11, 126: 47-56. ().
- HOLZSCHUH, CAROLUS. 1977. Bemerkenswerte Kaferfunde in Osterreich II. *Koleopterologische Rundschau* 53: 27-69. (ds).
- *_____. 1983. Bemerkenswerte Kaferfunde in Osterreich III [Important beetle finds in Austria]. *Mitteilungen der Forestlichen Bundesversuchsanstalt-Wien, Osterreichischer Agrarverlag*, 1141 Wien, FBVA, 148 (Scolytidae): 77: 81. ().
- _____. 1985. Eine neue Art der Gattung *Ips* aus Bhutan (Coleoptera, Scolytidae) [A new species of the genus *Ips* from Bhutan] [English summary]. *Entomologica Basiliensia* 12: 451-455. (tx).
- HONEA, C. R., T. EVAN NEBEKER, AND J. D. DEANGELIS. 1987. Selection of southern pine beetle hazard rating system for Mississippi. Pages 544-549 in E. P. Jones, Jr., 4th biennial Southern Silviculture Research Conference: proceedings. United States Department of Agriculture, Forest Service, General Technical Report SE-42. (cn).
- HONEA, C. R., T. EVAN NEBEKER, AND T. J. STRAKA. 1987. Economic evaluation of seven southern pine beetle hazard rating systems. *Journal of Environmental Management* 24: 259-266. (cn).
- HOPKINS, ANDREW DELMAR. 1906d. Insect enemies of forest reproduction. United States Department of Agriculture, Yearbook 1905: 249-256. (cn).
- _____. 1921. Contributions toward a monograph of the scolytid beetles. Contents and Index. United States Department of Agriculture, Technical Bulletin series 17: 233-247. (cn ms).
- *HORN, GEORGE HENRY. 1872. Coleoptera. Pages 352-392 in *Preliminary Report of the United States Geological Survey of Wyoming*. ().
- _____. 1894. The Coleoptera of Baja California. *California Academy of Science, Proceedings*, second series, 4: 302-449. (ds).
- HORNTVEDT, RICHARD. 1985. Resistance of *Picea abies* to *Ips typographus* tree response to monthly inoculations with *Ophiostoma polonicum*, a beetle transmitted blue-stain fungus. *Scandinavian Journal of Forest Research* 3: 107-114. (ec).

- HOSKING, GORDON P. 1954. Forestry pest problems of the South Pacific: the price of progress. *New Zealand Journal of Forestry* 29: 185-192. (cn).
- . 1959. Beech forest health implications for management. *New Zealand Journal of Forest Science* 19: 290-293. (cn).
- HOSKING, GORDON P., AND D. J. KERSHAW. 1955. Red beech death in the Manua Valley, South Island, New Zealand. *New Zealand Journal of Botany* 23: 201-211. (cn).
- HOSTETLER, BRUCE B., P. A. RUSIL AND T. A. LAURENT. 1976. Forest insect and disease conditions in Alaska, 1975. United States Department of Agriculture, Forest Service, Alaska Region (Juneau, Alaska). 12 p. (cn).
- HOUGHTON, C. O. 1905. A list of Coleoptera taken on the summit of Mt. Seward, N. Y. *Entomological News* 16: 50-51. (ds).
- HOULE, C., G. C. HARTMANN AND S. S. WASTI. 1957. Ineffectivity of eight species of entomogenous fungi to the larvae of the elm bark beetle, *Scolytus multistriatus* (Marshall). *New York Entomological Society, Journal* 95: 14-18. (ec).
- *HOWARD, LELAND OSSIAN. 1898. [*Hylastinus trifolii* (=obscurus)]. United States Department of Agriculture, Yearbook for 1897. ().
- HUBER, MR. AND E. T. WILLIAMS. 1953. Biological evaluation of pine bark beetles on the Andrew Pickens Ranger District, Sumter National Forest. United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia. Forest Pest Management Report 83-1-6. (cn).
- HUIHTA, VEIKKO, R. HYVONEN, P. KAASALAINEN, A. KOSKENNIEMI, J. MUONA, I. MAKELA, M. SULANDER, AND P. VILKAMAA. 1986. Soil fauna of Finnish coniferous forests. *Annales Zoologici Fennici* 23: 345-360. (ds).
- HUNT, D. W. A. 1986. Absence of fatty acid germination inhibitors for conidia of *Beauveria bassiana* on the integument of the bark beetle *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Canadian Entomologist* 118: 837-838. (ec).
- HUNT, D. W. A., AND JOHN HARVEY BORDEN. 1988. Response of mountain pine beetle, *Dendroctonus ponderosae* Hopkins, and pine engraver, *Ips pini* (Say), to ipsdienol in southwestern British Columbia. *Journal of Chemical Ecology* 14: 277-293. (bv).
- . 1989. Terpene alcohol pheromone production by *Dendroctonus ponderosae* and *Ips paraconfusus* (Coleoptera: Scolytidae) in the absence of readily culturable microorganisms. *Journal of Chemical Ecology* 15: 1433-1464. (bv).
- HUNT, D. W. A., JOHN HARVEY BORDEN, B. S. LINDGREN, AND G. GRIES. 1989. The role of autoxidation of alpha-pinene in the production of pheromones of *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *Canadian Journal of Forest Research* 19: 1275-1282. (bv).
- HUNT, D. W. A., JOHN HARVEY BORDEN, H. D. PIERCE, JR., K. N. SLESSOR, G. G. S. KING, AND E. K. CZYZEWSKA. 1986. Sex-specific production of ipsdienol and myrcenol by *Dendroctonus ponderosae* (Coleoptera: Scolytidae) exposed to myrcene vapours. *Journal of Chemical Ecology* 12: 1579-1586. (bv).
- HUNT, D. W. A., AND MICHAEL J. SMIRLE. 1988. Partial inhibition of pheromone production in *Dendroctonus ponderosae* (Coleoptera: Scolytidae) by polysubstrate monooxygenase inhibitors. *Journal of Chemical Ecology* 14: 529-536. (bv).
- HUNT, F. A. 1955. Wildlife issues in national forests. Pages 101-125 in W. J. Chandler, Audubon Wildlife Report 1955/1959. Academic Press, San Diego, California. xviii + 517 p. (ms).
- HUNT, R. S., AND D. J. MORRISON. 1956. Black-stain root disease on lodgepole pine in British Columbia. *Canadian Journal of Forest Research* 16: 996-999. (ec).
- HUNT, R. S., AND G. O. POINAR. 1971. Culture of a *Parasitorhabditis* sp. (Rhabditida: Protorhabditinae) on a fungus. *Nematologia* 17: 321-322. (ec).
- HUNTER, PRESTON EUGENE, R. M. T. ROSARIO, AND JOHN CONRAD MOSER. 1989. Two new species of *Ereynetes* (Acari: Prostigmata, Ereynetidae) associated with bark beetles. *Journal of Entomological Science* 24: 16-20. (ec).
- HURFORD, A. W. 1952b. Save the elms program increasingly effective. *Connecticut Woodlands* 17: 5. (cn ms).
- HURST, G. D., AND D. S. MATTESON. 1986. Chiral quaternary centers via boronic esters. *American Chemical Society (Abstracts)* 192. (bv ms).

I

- IACOBAEUS, H., AND A. LINDELOW. 1955. Insekter i obarkade contorta stammar. *Sveriges Skogsvardsforbunds Tidskrift* 55-2: 43-45. (cn).
- *IENISTEA, M. AL. 1933. Contributiuni noi la fauna coleopterelor Romaniei. *Buletinul Societatii Studentilor in Stinte Naturale* 4: 50-62. ().
- *ISAAC. 1934. [*Scolytus rugulosus*]. Pages 161-166 in Institute of Agricultural Research, Pusa, Science Report, Delhi, for 1932-1933. ().
- ISHIKUBO, SHIGERU. 1962. Physiological and ecological studies on the pine bark beetles. *Bulletin of the Faculty of Education, University of Kagoshima, Japan* 14: 26-81. (ay ec).
- ISRAELSON, GUNNAR. 1985. Notes on the coleopterous fauna of the Azores, with description of new species of *Atheta* Thomson (Coleoptera). *Boletim do Museu Municipal do Funchal [Madeira]* 37(165): 5-19. (ds).
- . 1990a. Further notes on the coleopterous fauna of the Azores, with speculations on its origin. *Bocagiana, Museum Municipal do Funchal [Madeira]* 135: 1-8. (ds).
- . 1990b. Key to the Macaronesian Hypoborini, with description of two new species (Coleoptera, Scolytidae). *Bocagiana, Museum Municipal do Funchal [Madeira]* 137: 1-11. (tx).
- ISRAELSON, GUNNAR, ANTONIO MACHADO, PEDRO OROMI, AND THURE PALM. 1982. Novedades para la fauna Coleopterologica de las Islas Canarias [Scolytidae, p. 133]. *Vieraea [Tenerife]* 11: 109-134. (ds).
- ISTRATE, GEORGE I. 1980. Date biometrice si raportul numeric al sexelor la *Dendroctonus micans* Kug. (Coleoptera, Scolytidae) in moliditurile din nordul Romaniei [In Romanian, French summary]. *Muzeul de Stiintele Naturii Bacau, Studii si Comunicari* 10-12: 59-65 (1977-1979). (hb).
- IVES, W. G. II., AND H. R. WONG. 1955. Tree and shrub insects of the prairie provinces. Government of Canada, Forestry Service, Northern Forestry Centre, Edmonton, Alberta. Information Report NOR-X-292. 327 p. (cn hb).

J

- JACKSON, G. G. 1962. Status of insects in the Kenora District. Pages 444-457 in *Annual District Reports of the Forest Insect and Disease Survey Ontario, 1961*. Canada Department of Agriculture, Forest Biology Division, Sault Ste Marie, Ontario. (cn).

- JACOBSON, MARTIN, AND STANLEY A. HALL. 1967. Attractants produced by one sex to lure both sexes (assembling scents). Pages 49-72 in *Insect sex attractants*. Interscience Publishers, John Wiley & Sons, Inc., New York, London, Sydney. (bv).
- JACTEL, H., AND F. LIEUTIER. 1957. Effects of attack density on fecundity of the Scots pine beetle *Ips sexdentatus* Born. (Col.: Scolytidae). *Journal of Applied Entomology* 104: 190-204. (bv).
- JAGER, D. 1955. Borkenkäferüberwachung mit Hubschrauber. *Allgemeine Forstzeitschrift* 5: 91. (cn).
- JAKAITIS, B. J., AND V. GAVELIS. 1957. Question of perfecting barrier traps for attraction of the bark beetle *Ips typographus* L. [In Russian]. *Khemoretsepsiya Nasekomykh* 9: 118-122. (cn).
- JAMES, ROBERT L., S. TUNAOCK, ROBERT LADD LIVINGSTON, J. W. SCHWANDT, D. P. BECKMAN, AND K. A. KNAPP. 1955. Idaho forest pest conditions and program summary, 1954. United States Department of Agriculture, Forest Service, Northern and Intermountain Regions and Idaho Department of Lands, Coeur d'Alene, Idaho, Report 55-1. 35 p. (cn).
- *JANCARIK, V., MILO KNIZEK, J. LISKA, J. PAREZ, V. PICHOVA, AND M. SVECOVA. 1991. Zhodnoceni vyskytu lesnich skodlivych cinitelu a jejich ocekavany stav v roce 1991 [Evaluation of occurrence of forest harmful factors and prognosis of their occurrence in 1991]. *Lesnicka prace* 5. ().
- *JANCARIK, V., M. SROT, J. PAREZ, P. ZAHRADNIK, AND MILO KNIZEK. 1990. Analýza stavu škodlivých činitelů lesních dřevin a prognóza jejich výskytu v roce 1990 [Analysis of forest harmful factors and prognosis of their occurrence in 1990]. *Lesnická práce* 5. 9 p. ().
- JANIN, J. L., AND F. LIEUTIER. 1955. Existence de fécondations précoces dans le cycle biologique de *Tomicus piniperda* L. (Coleoptera, Scolytidae) en forêt d'Orléans [The existence of early mating in the life cycle of *Tomicus piniperda* in the forest of Orléans] [English summary]. *Agronomie S(2)*: 169-172. (hb).
- JANSONS, V. 1963. Status of insects in the Geraldton District. Pages F26-33 in *Annual district reports of the Forest Insect and Disease Survey Ontario, 1962*. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- _____. 1964. Status of insects in the Geraldton District. Pages F15-24 in *Annual District Reports of the Forest Insect and Disease Survey Ontario, 1962*. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- JENNINGS, DANIEL T., AND HERBERT ALLEN PASE III. 1956. Spiders preying on *Dendroctonus frontalis* (Coleoptera: Scolytidae). *Entomological News* 97: 227-229. (ec).
- JOHANN, M. 1956a. Zur Anlockung des Buchdruckers (*Ips typographus* L.). I. Die Lockwirkung natürlichen Brutmaterials [Attraction of the engraver beetle (*Ips typographus*)]. I. The attractant effect of natural breeding material [English summary]. *Journal of Applied Entomology* 101: 332-342. (bv).
- _____. 1956b. Zur Anlockung des Buchdruckers (*Ips typographus* L.). II. Möglichkeiten der Integration Natürliches Brutmaterial in die Bekämpfung mit Pheromon [Attraction of the engraver beetle (*Ips typographus*)]. II. Possibilities of the integration of natural breeding materials into control by pheromones [English summary]. *Journal of Applied Entomology* 102: 115-123. (cn).
- *JOHANNSEN. 1913. [*Hylobogopinus rufipes*]. Maine Agricultural Experiment Station. 15 p. ().
- JOHNSON, RICHARD LEE. 1954. Bark beetles of interior Douglas-fir/grand fir forest type. Pages 103-107 in D. M. Baumgartner and R. Mitchell, *Silvicultural management strategies for pests of the interior Douglas-fir and grand fir forest types: proceedings of a symposium*. Cooperative Extension, Washington State University, Pullman, Washington. (cn hb).
- *JOHNSON, CHARLES WILLISON. 1900. [*Scolytus rugulosus*]. Page 393 in *New York Agricultural Experiment Station, Bulletin* 195. ().
- JOHNSON, DAVID W., AND C. G. O'NEIL. 1986. Forest pest conditions in the Rocky Mountain Region for 1985. United States Department of Agriculture, Forest Service, Timber, Forest Pest, and Cooperative Forestry Management (Denver, Colorado). 48 p. (cn).
- JOHNSON, M. A., AND R. CROTEAU. 1957. Biochemistry of conifer resistance to bark beetles and their fungal symbionts. Pages 76-92 in G. Fuller and W. D. Nes, *Ecology and metabolism of plant lipids*. American Chemical Society Symposium Series 325. vii + 374 p. (ec).
- JOHNSON, WARREN T., AND HOWARD H. LYON. 1955. *Insects that feed on trees and shrubs*. Edition 2. Comstock Publishing Associates, Cornell University Press, Ithaca, New York and London. 556 p. (cn).
- JOLY, R. 1976b. Les insectes ennemis des pins. Volume 2. Ecole nationale du Génie Rural, des eaux et des Forêts: Centre de Nancy. Unpaginated. (hb).
- JONG, M. C. M. DE, AND P. GRIJPSMA. 1956. Competition between larvae of *Ips typographus*. *Entomologia Experimentalis et Applicata* 41: 121-133. (ec).
- JONG, M. C. M. DE, AND M. W. SABELIS. 1959. How bark beetles avoid interference with squatters: a correction. *Oikos* 54: 125. (hb).
- JOY, NORMAN H. 1909. Three new British Coleoptera. *Entomologist's Monthly Magazine* 45: 268-269. (ds).
- *JULIASOVA, G., AND P. HRUBIK. 1975. Príčina usychania borievok (*Juniperus* sp.) na Slovensku [Causes of drying up of junipers in Slovakia]. *Lesníctví* 21(XLVIII), 11: 1019-1029. ().

K

- KAARIK, AINO. 1975. Skydda skogen och virket! Insekter och blånad. *Skogen* 62: 727. (ms).
- KADYROV, A. KH. 1955. K faune korozedov (Coleoptera, Scolytidae) drevnykh porod jugozapachovo Tadzhikistana [On the fauna of scolytids (Coleoptera, Scolytidae) of tree species of south-western Tadzhikistan] [In Russian, English summary]. *Entomologicheskoe Obozrenie* 67: 42-47. Translation in *Ent. Revue* 68 (1): 47-61. (ds).
- *_____. 1959a. Bark beetles (Coleoptera, Scolytidae) in central Asia USSR. *Izv. Akad. Nauk Tadzh. SSR Otd. Biol. Nauk* 2: 23-27. ().
- _____. 1959b. On the bark beetles (Coleoptera, Scolytidae) of southwestern Tajikistan [In Russian, English summary]. *Entomologicheskoe Obozrenie* 68(1): 12-18. Translation in *Ent. Revue*. (ds).
- KALIDIS, D. S. 1955. Dry periods and secondary dying and bark beetle epidemics in forests of Greece [In Greek, English summary]. Publication, Ergasterio Ulonikes, Tmesa Dasologias Kai Fusikou Periballontos, Aristoteleio Panepistemio Thessalonikes 2. 16 p. (cn ds).
- KALIDIS, D. S., AND S. MARKALAS. 1955. Durreperioden in

- Zusammenhang mit sekundärem Absterben und Massenvermehrungen rindenbittender Insekten in den Wäldern Griechenlands [Dry periods and secondary dying and bark beetle epidemics in forests of Greece [In German. English summary]. *Anzeiger für Schadlingskunde Pflanzen- und Umweltschutz* 61: 25–30. (cn ds).
- *KAKULIYA, G. A. 1963a. A new species of nematode, *Tylaphelenchus paramonovi* Kakulia n. sp. [In Russian]. *Doklady Akademii Nauk Gruzinskoi SSR* 32: 649–654. ().
- *_____. 1963b. A new species of nematode of the four-striped engraver, *Cryptophelenchus quadridens* n. sp. [In Russian]. *Doklady Akademii Nauk Gruzinskoi SSR* 31: 433–435. ().
- _____. 1967. A new species of nematode, *Contortylechus proximus* sp. n. from *Orthotomicus proximus* Eichhoff [In Russian]. *Doklady Akademii Nauk Gruzinskoi SSR* 48: 459–460. (ec).
- KAKULIYA, G. A., AND S. L. LAZAREVSKAYA. 1965. *Ectaphelenchus piniperdae* sp. n., a new nematode of the larger pine shoot beetle [In Russian]. *Trudy Gel'mintol. Lab. Akademii Nauk SSR* 15: 54–56. (ec).
- KALISZEWSKI, MAREK J., EARLE ALBRIGHT CROSS, JOHN CONRAD MOSER, AND T. W. PHILLIPS. 1985. Systematic position, biology, and ecology of *Scolytomotes hockeji* n. sp., n. gen., n. tribe (Acari: Dolichoeyboidea): a new parasitoid of bark beetles (Scolytidae). *International Congress of Entomology, Proceedings* 15: 435. (ec).
- *KALSHOVEN, LOUIS GEORGE EDMUND. 1926c. Primaire aantasting van houtige gewassen door *Xyleborus* soorten. *Algemeen Landbouweekblad* 11: 172–178. ().
- KAMP, HANS J. 1979. Erwähnenswerte Scolytiden-Funde aus Deutschland. *Entomologische Blätter für Biologie und Systematik der Käfer* 74: 183. (ds).
- KAPLAN, M., AND T. MOKRZYCKI. 1985. A contribution to the knowledge of the occurrence of the pine shoot beetles *Tomicus minor* and *T. piniperda* in Scots pine thickets in Niedzwiedy Forest District [In Polish, English and Russian summaries]. *Sylvan* 132(6): 35–39. (cn).
- KARASEVICZ, D., AND W. MERRILL. 1956. Volume of stain and decay not related to ambrosia beetle attack in oaks killed following gypsy moth defoliation. *Annual meeting of the American Phytopathological Society (Northeastern Division)*, 6–8 November 1955, 76: 654. (ec).
- KARG, W. 1955. Eine neue Raubmilbenart der Gattung *Proctolaelaps* Berlese 1923 (Acarina, Parasitiformes) am Grossen Oberbaumsplintkafer (*Scolytus [Eccoptogaster] mali* Bechst.). [A new predatory mite species of the genus *Proctolaelaps* Berlese 1923 on the great fruit bark beetle [English summary]. *Archiv für Phytopathologie und Pflanzenschutz* 24: 515–517. (ec).
- KARPINSKI, C., JR., ROY L. HEDDEN, ROGER P. BELANGER, AND T. S. PRICE. 1954. Guidelines for managing pine bark beetles in Georgia. *Georgia Forestry Commission*. 61 p. (cn).
- *KARPINSKI, J. J. 1937. Korniki zebrane w Parku Narodowym na Czarnohorze w lecie 1937 r. [Ipides recoltés dans le Parc National de Czarnohora durant l'été 1937]. [Source?]. 1 p. ().
- KATTIAR, K. H., AND V. SHARMA. 1957. Influence of bark thickness on the host preference of hymenopterous parasites of bark beetles. *Indian Society of Desert Technology, Transactions* 12: 121–122. (ec).
- KATOVICH, S. A., AND R. J. LAVIGNE. 1986. The applicability of available hazard rating systems for mountain pine beetle in lodgepole pine stands of southwestern Wyoming. *Canadian Journal of Forest Research* 16(2): 222–225. (cn).
- KEILBACH, ROLF. 1952. Bibliographie und Liste der Arten tierischer Einschlüsse in fossilen Harzen sowie ihrer Aufbewahrungsorte. *Deutsche Entomologische Zeitschrift, new series*, 29: 129–256. (ds ms).
- *KELER, STEFAN. 1926. [*Ips crosus*]. *International Congress of Entomology, Proceedings* 3: 169–170. ().
- KELLEHER, JAMES S. 1956. The Canadian Agriculture Insect Pest Review, Volume 62, 1954. *Canada Department of Agriculture, Research Branch, Ottawa, Ontario*. 107 p. (cn ds).
- KELLEY, WAYNE. 1956. Managing southern forests to reduce southern pine beetle impacts: long and short-term strategies and research needs. *United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia*. 37 p. (cn).
- KELLEY, WAYNE, ROGER P. BELANGER, MICHAEL D. CONNER, TOMMY R. DELL, J. E. DE STEIGUER, ROSS KESTER, ROBERT N. KITCHENS, PETER L. LORIO, JR., AND KENNETH M. SWAIN. 1956a. Long-term strategies and research needs for managing southern forests to reduce southern pine beetle impacts. *United States Department of Agriculture, Forest Service, Atlanta, Georgia*. 37 p.
- _____. 1956b. Short-term strategies for managing southern forests to reduce southern pine beetle impacts. *United States Department of Agriculture, Forest Service, Atlanta, Georgia*. 32 p. (cn).
- KENNY, P. T. M., S. Y. TAMURA, A. FREDENHAGEN, Y. NAYA, K. NAKANISHI, K. NISHIJAMA, M. SUGIURA, H. KITA, AND H. KOMURA. 1959. Symbiotic micro-organisms of insects: a potential new source for biologically active substances. *Pesticide Science* 27: 117–131. (cn ec).
- KESKİNALEMDAR, E., S. ALKAN, AND Y. AKSU. 1957. Artvin ilinde *Ips typographus* (L.) (Coleoptera, Scolytidae) 'un biyolojisi ve mücadelsi üzerinde calismalar [Studies on the biology and control of *Ips typographus* in Artvin Province] [In Turkish, English summary]. *Türkiye 1. Entomoloji Kongresi Bildirileri*, 13–16 Ekim 1957, Ege Üniversitesi, Bornova, İzmir. Bornova/İzmir, Turkey; Ege Üniversitesi Atatürk Kültür Merkezi 1957: 737–742. (cn hb).
- KEYSOR, ANN M. 1956a. Evaluation of pest conditions, Kaler and Dodds Hollows, Vernal Ranger District, Ashley National Forest. *United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report* 56–2. 4 p. (cn).
- _____. 1956b. Evaluation of pest conditions, North Fork Provo River and Broadhead Meadows, Kamas Ranger District, Wasatch National Forest. *United States Department of Agriculture, Forest Service, Intermountain Region (Ogden, Utah), Forest Pest Management Report* R4–56–7. 4 p. (cn).
- _____. 1956c. Evaluations of pest conditions, Old Lilly Lake Burn, Evanston Ranger District, Ashley National Forest. *United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report* 56–3. 3 p. (cn).
- KFIR, R. 1956. Releases of natural enemies against the pine bark beetle *Orthotomicus crosus* (Wollaston) in

- South Africa. Entomological Society of Southern Africa, Journal 49: 391–392. (cn ec).
- KHALAF, K. T. 1980. Micromorphology of beetle elytra, using simple replicas. Florida Entomologist 63: 307–340. (av).
- KIELCZEWSKI, BOHDAN, AND S. SENICZAK. 1966. The problem of predation of bark beetle eggs by mites. Ekologia Polska B, 12: 161–163 [Translation, United States Department of Commerce, National Technical Information Service, Springfield, Virginia]. (ec).
- KIESEL, KLAUS. 1955. Bibliographie der neueren Borkenkäferliteratur unter besonderer Berücksichtigung der Aggregationspheromone. Freiburger Waldschutzabhandlungen, Vol. 7. 279 p. (bv ms).
- KILE, G. A. 1957. Lack of a direct role for *Platypus subgranosus* (Coleoptera) in the dissemination of *Chalara australis*, the cause of a vascular disease of *Nothofagus cunninghamii*. XIV International Botanical Congress (Abstracts) 15: 405. (ec).
- KILE, G. A., AND M. F. HALL. 1955. Assessment of *Platypus subgranosus* as a vector of *Chalara australis*, causal agent of a vascular disease of *Nothofagus cunninghamii*. New Zealand Journal of Forest Science 15: 166–186. (ec).
- KILE, G. A., AND J. WALKER. 1957. *Chalara australis* sp. nov. (Hyphomycetes): a vascular pathogen of *Nothofagus cunninghamii* (Fagaceae) in Australia and its relationship to other *Chalara* species. Australian Journal of Botany 35: 1–32. (ec).
- KING, C. J., AND H. F. EVANS. 1955. The rearing of *Rhizophagus grandis* and its release against *Dendroctonus micans* in the United Kingdom. Pages 57–97 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium. Proceedings of seminar, 3–4 October 1954. 141 p. (ec ms).
- KINN, DONALD N. 1956a. Incidence of the pinewood nematode in a southern pine beetle infestation in central Louisiana. Journal of Entomological Science 21: 114–117. (ec).
- . 1956b. Studies of the flight capabilities of *Dendroctonus frontalis* and *Ips calligraphus*: preliminary findings using tethered beetles. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, Research Note SO-324. 3 p. (bv hb).
- KINN, DONALD N., AND C. DEISTER. 1957. Pinewood nematode: decontamination of North American wood chips for pulp and paper research in Europe. Tappi Journal 70(5): 131–132. (ec).
- KINN, DONALD N., AND L. M. ROTON. 1959. Rearing the sixspined ips (Coleoptera: Scolytidae) free of mite associates. Entomological Society of America, Annals 52: 60–63 (hb ms).
- KINUURA, H., AND K. KANAMITSU. 1955. Scolytidae and their host trees in central Japan. International Congress of Entomology, Proceedings 15: 440. (ds).
- KINZER, H. GRANT, AND J. M. REEVES. 1955. Chemical treatments for brood control and suppression of *Dendroctonus adjunctus* attacks on ponderosa pine. Southwestern Entomologist 10: 244–252. (cn).
- KIRCHHOFF, J. F., AND E. FUIRER. 1955. Häufigkeit und Verbreitung von *Malamocha scolyti* Purrini bei *Dryocoetes autographus* in einigen Gebieten Nord- und Nordwest-deutschlands [Abundance and distribution of *Malamocha scolyti* Purrini in *Dryocoetes autographus* in some areas of north and north-west Germany] [English summary]. Forstwissenschaftliches Centralblatt 104: 373–380. (ec).
- KIRK, VERNON MILES, AND E. U. BLUBAUGH, JR. 1975. A list of the beetles of South Dakota. South Dakota State University Agricultural Experiment Station, Technical Bulletin 42. 139 p. (ds).
- KIRKENDALL, LAWRENCE RICHARD. 1958. Evolution of scolytid gallery systems. International Congress of Entomology, Proceedings 15: 409. (hb tx).
- . 1959. Within-harem competition among *Ips* females, an overlooked component of density-dependent larval mortality. Holarctic Ecology 12: 477–487. (ec).
- KIRKENDALL, LAWRENCE RICHARD, AND N. C. STENSETH. 1959. Population dynamics of bark beetles with special reference to *Ips typographus*: contributions of applied bark beetle studies to basic research in ecology and population biology. Holarctic Ecology 12: 526–527. (hb).
- KIRSTEN, J. F. 1958. Sable Forest Entomology Unit. Plant Protection News, South Africa No. 11(3). (ms).
- *KLAGES, H. G. 1901. Supplement to Dr. John Hamilton's list of the Coleoptera of southwestern Pennsylvania. Annals of the Carnegie Museum 1: 265–294. ().
- *KLAUSNITZER, B., AND G. FÖRSTER. 1976. Zur Kenntnis der Parasiten und Episiten des Bruchdruckers *Ips typographus* L. (Col., Scol.). Entomologische Berichten 1976: 11–43. ().
- KLEIN, E. 1955. Lebendbefall durch den Nutzholzborkenkäfer. Allgemeine Forstzeitschrift 12: 264. (cn).
- KLIEJUNAS, JOHN. 1955. A biological evaluation of black stain root disease and associated pests in Timber Mountain Area, Doublehead Ranger District, Modoc National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 85–22. (cn).
- KLIEJUNAS, JOHN, AND J. PIERCE. 1956. A biological evaluation of proposed recreation sites at Sourgrass and Middle Fork, Stanislaus National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 56–5. (cn).
- KLIMETZEK, D., E. J. BAADER, AND W. HILBIG. 1951b. Die Eignung von Lockstoff-Fallen zur Überwachung der Ulmensplintkäfer. Mitteilungen der Deutschen Gesellschaft für Angewandte Entomologie 2: 311–313. (bv cn).
- KLIMETZEK, D., J. BARTELS, AND W. FRANCKE. 1959. The pheromone system of the elm bark beetle *Ptilcobius vittatus* F. (Coleoptera: Scolytidae). Journal of Applied Entomology 107: 518–523. (bv).
- KLIMETZEK, D., J. KOHLER, S. KROHN, AND W. FRANCKE. 1959. Das Pheromon-System des Waldrehen-Borkenkäfers *Xylocleptes bispinus* Duft. (Col., Scolytidae) [The pheromone system of the Clematis bark beetle] [English summary]. Journal of Applied Entomology 107: 304–309. (bv).
- KLIMETZEK, D., J. KOHLER, JEAN PIERRE VITE, AND U. KOHNLE. 1956. Dosage response to ethanol mediates host selection by "secondary" bark beetles. Naturwissenschaften 73: 270–272. (bv).
- KLIMETZEK, D., AND JEAN PIERRE VITE. 1956. Die Wirkung insektenbuertiger Duftstoffe auf das Aggregationsverhalten des Mediterranen Kiefernborckenkäfers *Orthotomicus erosus* [The role of insect produced attractants on the aggregation behaviour of the Mediterranean pine engraver beetle

- Orthotomicus erosus* [English summary]. Journal of Applied Entomology 101: 239–243. (bv).
- . 1989. Die tierischen Schädlinge der Fichte. In H. Schmidt-Vogt, Die Fichte, Vol. 2, Part 2. Parey, Hamburg, Berlin. (cn).
- *KLIMSCH, E. 1899. Die Kaferwelt der Umgebung Klagenfurts, besonders jene der Satnitz. Carinthia II, 89: 136–152. ().
- KLIPSTEIN, E. L. 1985. Rassendifferenzierung beim Kupferstecher. Allgemeine Forstzeitschrift 12: 263. (ay).
- . 1986a. Cytologie und Spermapolyploidie bei *Pityogenes chalcographus* L. (Col., Scolytidae) [Cytology and spermapolyploidy in *Pityogenes chalcographus*] [English summary]. Journal of Applied Entomology 102: 285–295. (ay).
- . 1986b. Intraspezifische differenzierung in der Epidemiologie und Ökologie von *Pityogenes chalcographus* L. Bastardierung [Intraspecific differentiation in epidemiology and ecology of *Pityogenes chalcographus* with regard to hybridization] [English summary]. Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz 59: 131–135. (ay hb).
- *KLOET, G. S., AND W. D. HINCKS. 1945. A check list of British insects. Kloet & Hincks, Stockport. 477 p. ().
- KNAPP, ANDREW K. 1985. Forest insect and disease conditions: Intermountain Region 1984. United States Department of Agriculture, Forest Service, State and Private Forestry, Intermountain Region, Ogden, Utah. 23 p. (cn).
- KNAPP, ANDREW K., J. C. WEATHERBY, J. T. HOFFMAN, V. KALVE, AND L. LA MADELINE. 1989. Forest insect and disease conditions: Intermountain Region 1988. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah. 31 p. (cn).
- . 1990. Forest insect and disease conditions: Intermountain Region. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah. 38 p. (cn).
- KNAUER, K. H. 1989. An administrative perspective on North American bark beetles and biological control opportunities. Pages 231–236 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ms).
- KNEIFF, FRITZ. 1939. Schlechtes Wachstum der Sitkafichte. Mitteilungen der Deutschen Dendrologischen Gesellschaft 52: 170. (cn hb).
- KNIZEK, MILOS. 1983. Faunistic records from Czechoslovakia: Coleoptera, Scolytidae, *Pityogenes trepanatus* (Nord.). Acta Entomologica Bohemoslovaca 80: 399. (ds).
- . 1988. Faunistic records from Czechoslovakia: Coleoptera, Scolytidae, *Xyleborus alni* Nisima. Acta Entomologica Bohemoslovaca 85: 396. (ds).
- . 1989. K životnímu jubileu Profesora Antonína Pfeffer. Lesnická Práce 1989: 45–46. (ms).
- . 1991. Book review: Antonin Pfeffer, Kurovcoviti (Scolytidae) a Jadrólodoviti (Platypodidae). Acta Entomologica Bohemoslovaca 88: 376. (tx ms).
- KNOWLTON, GEORGE FRANKLIN. 1931. Notes on Utah Coleoptera. Florida Entomologist 15: 10. (ds).
- KNOWLTON, GEORGE FRANKLIN, AND G. P. TAYLOR. 1952. Beetles—Coleoptera: records and notes, largely from Utah. Utah Agricultural Experiment Station, Mimeograph Series 359: 1–32. (ds).
- KO, J. H., AND K. MORIMOTO. 1985. Loss of tree vigor and the role of boring insects in red pine stands heavily infested by the pine needle gall midge in Korea. Esakia 23: 151–155. (cn).
- KOBAYASHI, FUJIO. 1978. Pine bark beetle problem in Japan, referring to the discovery of the pine wood nematode, *Bursaphelenchus lignicola* (Mamiya & Kiyohara). Anzeiger für Schädlingkunde, Pflanzenschutz, Umweltschutz 51: 76–79. (cn).
- KOEHLER, CARLTON SMITH, R. H. HUNT, D. FROEHLICH, AND J. GEIGER. 1988. Protecting trees when building on forested land. University of California, Division of Agriculture and Natural Resources, Cooperative Extension Leaflet 2134S. 11 p. (cn ms).
- KOEHLER, CARLTON SMITH, J. J. MCKELVEY, JR., W. L. ROELOFS, II., H. SHOREY, ROBERT MILTON SILVERSTEIN, AND DAVID LEE WOOD. 1977. Advancing toward operational behavior-modifying chemicals. Pages 395–400 in H. H. Shorey and J. J. McKelvey, Jr., Chemical control of insect behavior: theory and application. John Wiley & Sons, New York. (bv cn).
- KOEHLER, U. 1986. Problems of infesting standing trees by *Blastophagus piniperda* [In German, English summary]. Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz 59: 145–147. (cn).
- KOEBER, THOMAS. 1985. Plantation insect problems. Page 51 in Thirty-sixth annual Western Forest Insect Work Conference: proceedings, 4–7 March 1985, Boulder, Colorado. Canada Department of Environment, Canadian Forestry Service, Northern Forest Research Centre, Edmonton, Alberta. 54 p. (cn ec).
- KOHLER, STEVE, O. J. DOOLING, AND W. E. BOUSFIELD. 1985. Montana forest pest conditions and program highlights, 1984. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Cooperative Forestry and Pest Management Report 85–2. 23 p. (cn).
- KOHNLE, U. 1984. Beziehungen zwischen Austrocknung von Fangbäumen und ihrer Attraktivität für Fichtenborkenkäfer. Allgemeine Forst- und Jagdzeitung 155: 270–274. (bv cn).
- * ———. 1985a. Untersuchungen über die Pheromonsysteme sekundärer Borkenkäfer (Col., Scolytidae). Unpublished dissertation, University of Freiburg, Bertoldstr., Germany. ().
- . 1985b. Untersuchungen über die Pheromonsysteme sekundärer Borkenkäfer (Col., Scolytidae). Journal of Applied Entomology 100: 197–215. (bv hb).
- KOHNLE, U., W. FRANCKE, AND ALF BAKKE. 1985. *Polygraphus poligraphus* (L.): response to enantiomers of beetle specific terpene alcohols and a bicyclic ketol. Journal of Applied Entomology 100: 5–8. (bv).
- KOHNLE, U., S. KOPP, AND W. FRANCKE. 1986. Inhibition of the attractant pheromone response in *Ips acuminatus* (Gyll.) by *Ips sexdentatus* (Boerner) (Coleoptera, Scolytidae). Journal of Applied Entomology 101: 316–319. (bv).
- KOHNLE, U., M. MUSSONG, V. DUBBEL, AND W. FRANCKE. 1987. Acetophenone in the aggregation of the beech bark beetle *Taphrorychus bicolor* (Col., Scolytidae). Journal of Applied Entomology 103: 249–252. (bv).
- KOHNLE, U., HEINRICH SCHMUTZENHOFFER, J. BARELS, AND W. FRANCKE. 1988. Oxygenated terpenes in the chemical communication system of the bark beetle, *Ips schmutzenhoferi* (Col., Scolytidae), a species

- recently described for the southeastern Himalaya. *Journal of Applied Entomology* 106: 46-51. (bv).
- KOHNLE, U., JEAN PIERRE VITE, C. ERBACHER, J. BARTELS, AND W. FRANCKE. 1988. Aggregation response of European engraver beetles of the genus *Ips* mediated by terpenoid pheromones. *Entomologia Experimentalis et Applicata* 49: 43-54. (bv).
- KOLBE, W. 1957a. Anmerkungen zur Arthropodenfauna im Staatswald Burgholz unter besonderer Berücksichtigung der Borkenkäfer (Scolytidae) [Remarks on the arthropod fauna of the Burgholz State Forest with particular reference to bark beetles] [English summary]. *Decheniana* 104: 73-75. (cn ds).
- . 1957b. Der Einfluss von NA-PCP auf die Arthropodenfauna in Staatswald Burgholz in Solingen unter besonderer Berücksichtigung der Coleopteren [The influence of NA-PCP on the arthropod fauna in the Burgholz State Forest in Solingen with particular reference to the Coleoptera] [English summary]. *Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie, Mitteilungen* 5: 82-86. (cn ec).
- *KOLOMIETS, N. G. 1981. Interaction between the European spruce bark beetle (*Dendroctonus micans* Kugel., Col., Scol.) and Scots pine (*Pinus sylvestris* L.) in western Siberia [abstract]. The role of insect-plant relationships in the population dynamics of forest pests. UFIRO/MAB meeting, 24-28 August 1981, Irkutsk, USSR. 65: 28. ().
- *KOLUBAJIV, SERGEJ. 1962. Výsledky chovu entomofagu (cizopasníku a dravců) lmyzích škůdců, hlavně lesních, získané v období 1934-1958. *Rozpravy Československé akademie věd* 72(6): 1-73. ().
- KONDO, EDWARD S., AND B. H. MOODY. 1957. Forest insect and disease conditions in Canada, 1956. Government of Canada, Forestry Service, Forest Insect and Disease Survey, Ottawa, Ontario. 128 p. (cn).
- KONDO, EDWARD S., AND R. G. TAYLOR. 1954. Forest insect and disease conditions in Canada, 1953. Canada Department of Environment, Forestry Service, Forest Insect and Disease Survey, Ottawa, Ontario. 73 p. (cn).
- . 1955. Forest insect and disease conditions in Canada 1954. Government of Canada, Forestry Service, Forest Insect and Disease Survey, Ottawa, Ontario. 76 p. (cn).
- . 1956. Forest insect and disease conditions in Canada, 1955. Government of Canada, Forestry Service, Forest Insect and Disease Survey, Ottawa, Ontario. 107 p. (cn).
- KONGKATHIP, N., B. KONGKATHIP AND R. SOOKKHO. 1987. A short efficient synthesis of (plus or minus) frontalin and its analogue. *Journal of the Science Society of Thailand* 13: 239-242. (bv ms).
- KONIG, H. ERWIN. 1984. Aus der Forschungsarbeit der Abt. Waldschutz. *Mitteilungen der Forstlichen Versuchs- und Forschungsanstalt Baden-Württemberg* 108: 133-150. (cn).
- . 1988. Mass-trapping of *Trypodendron lineatum* Ol. (Col., Scolytidae): effect of infestation of cut timber temporarily left in the forest. *Journal of Applied Entomology* 106: 262-265. (bv cn).
- KONIG, H. ERWIN, AND WOLFRAM BERWIG. 1986. Kalk statt insektizide gegen gestreiften nitzholzborckenkafer? *Allgemeine Forstzeitung* 41: 326-327. (cn).
- . 1987. Kalkspritzung als vorsorge gegen nitzholzborckenkaferbefall nicht bewahrt [Lime spraying not successful in protecting timber against bark beetle attack] [English summary]. *Allgemeine Forstzeitung* 41: 354-355. (cn).
- KONOPKA, JOSEF E., SOBOCKY, AND M. CAPEK. 1987. Poskodzovanie Lesov SSR a jeho prognóza do roku 2000 [Damage to forests of the Slovak SR and forecasts up to year 2000] [In Slovak, English summary]. *Lesnícky Časopis* 33: 49-61. (cn).
- KOPPONEN, M., AND M. NUORTeva. 1973. Über subfossile Waldinsekten aus dem Moor Pilonisno in Südfinnland. *Acta Entomologica Fennica* 29: 1-84. (ds).
- KORENCHENKO, E. A. 1987a. *Cryptaphlenchus diversispicularis* n. sp. (Thylenchida, Aphelenchoididae), a new nematode of the bark beetle, *Ips subelongatus* (Coleoptera, Ipsidae) [In Russian, English summary]. *Parazitologiya* 21: 73-75. (ec).
- . 1987b. *Parasitylenchus klimenkorum* n. sp. (Nematoda, Allantonematidae), a parasite of *Orthotomicus laricis* (Coleoptera, Ipsidae) [In Russian, English summary]. *Parazitologiya* 21: 567-576. (ec).
- KOROLEV, S. G. 1989. The morphology of the larvae of *Platypus* Herbst beetles (Coleoptera: Platypodidae) in connection with the characteristics of their ecology [In Russian]. *Entomologicheskoe Obozrenie* 68: 353-360. Translation in *Ent. Revue*, (ay tx).
- *KOVACEVIC, ZELJKO. 1928. [*Scolytus rugulosus*]. Verhandlungen der Deutschen Gesellschaft für Angewandte Entomology 7: 33-41. ().
- *KOVACH, JOSEPH. 1956a. Life cycle, seasonal distribution and tree responses to scolytid beetles in South Carolina peach orchards. Unpublished dissertation, Clemson University, Clemson, South Carolina. 88 p. ().
- . 1956b. Life cycle, seasonal distribution and tree responses to scolytid beetles in South Carolina peach orchards. *Dissertation Abstracts International* 47(5-B): 1845. (bv cn hb).
- KOVACH, JOSEPH, AND CLYDE S. GORSUCH. 1955. Survey of ambrosia beetle species infesting South Carolina peach orchards and a taxonomic key for the most common species. *Journal of Agricultural Entomology* 2: 235-247. (cn hb).
- KOVACIC, DAVID ALLEN, M. I. DYER, AND A. T. CRINGAN. 1985. Understory biomass in ponderosa pine following mountain pine beetle infestation. *Forest Ecology and Management* 13: 53-65. (cn).
- KOWAL, ROMUALD JOSEPH, AND BERNARD H. EBEL. 1972. Insects attacking forest trees in the South. *Forest Farmer, Manual Edition* 31(7): 24-30. (cn).
- KOZLOV, V. I. 1966. On the biology of elongata bark beetle (*Ips elongatus* Motsch.) in the larch forests of the lower Angara region [In Russian]. Pages 74-76 in A. I. Cherepanov, The fauna and ecology of Arthropoda from Siberia. Academy of Sciences of the USSR, Siberian Branch, Novosibirsk. (hb).
- KRISTIN, ANTON. 1986. Some results in the research of pheromone preparations on *Ips typographus* and notes to their selective effect [In Slovak]. *Sprawy Slovenskej zoologickej spoločnosti pri SAV, Bratislava* 12: 198-201. (bv cn).
- KRISTIN, ANTON, AND FERDINAND BENDA. 1986. Trapping of non-target Coleoptera species in pheromone traps to *Ips typographus* [In Slovak]. *Prace Slov. entomol. spol. SAV, Bratislava* 6: 118-127. (bv ec).
- KRISTIN, ANTON, STEFAN VARKONDA, LUDOVIT KURUC, AND FERDINAND BENDA. 1985. The new findings in development of new pheromone preparations against the spruce bark beetle *Ips typographus* L. and notes to their selective effect. Pages 242-259 in Conference

- on biological and biotechnological control of forest pests in Tabor: proceedings. Forestry and Game Management Research Institute Jiloviste, Strmady. (bv cn).
- KRIVOSHEINA, N. P., AND B. M. MAMAEV 1986. Regional complexes of trunk insects of fir [In Russian, English summary]. *Izvestiya Sibirskogo Otdeleniya Akademii Nauk SSSR, Biologicheskiy Khim. Nauk* (2): 97–103. (ds).
- KRIVOSHEINA, N. P., AND T. B. TOKGAEV 1985. The formation of complexes of trunk insects on irrigated lands in the foothills of the Kopet-Dag [In Russian, English summary]. *Izvestiya Akademii Nauk Turkmenskoy SSR, Biologicheskikh Nauk* 5: 34–39. (cn).
- *KROL, ALFRED. 1982. Szkodniki wtorne sosny wejmutki (*Pinus strobus* L.) na wybranej powierzchni lesnego zakladu doswiadczalnego w Krynicy [Secondary pests of *Pinus strobus* L. on a sample area of the Krynica forest experimental station]. *Lesnictwo, Z.* 14, 172: 91–99. (i).
- KROL, ALFRED, AND ALF. BAKKE 1985. Effect of distance between the pheromone traps on the effectiveness of the spruce bark beetle *Ips typographus* aggregation. *Acta Agrar Silvestria Ser Silvestris* 24: 21–30. (bv).
- . 1985. Comparison of trap trees and pheromone loaded pipe traps in attracting *Ips typographus* L. (Col., Scolytidae). *Canadian Entomologist* 56: 437–445. (bv cn).
- KROL, ALFRED, DANIEL KUBISZ, JERZY R. STARZYK, AND MACIEJ WITRYLAK 1987. Badania nad przwabianiem drwalnika paskowanego, *Trypodendron lineatum* Ol., Col., Scolytidae, do rurowych pułapek feromonowych w lesnym zakładzie doświadczalnym w Krynicy, białe sadyki [Studies on the attraction of ambrosia beetle, *Trypodendron lineatum* to pheromone pipe-traps in the Forest Experiment Station in Krynica] [In Polish, English summary]. *Zeszyty Naukowe Akademii Rolniczej Im. Hugona Kollataja w Krakowie, Sesja Naukowa* 17: 151–172. (bv cn).
- KROMBEIN, KARL VON VORSE, PAUL D. HURD, JR., D. R. SMITH, AND BARNARD DEWITT BURKS 1979. Catalog of Hymenoptera in America North of Mexico. Smithsonian Institution, Washington D.C. 2735 p. (ec).
- KUBO, ISAO, S. KOMATSU, T. IWAGAWA, AND DAVID LEE WOOD 1986. Analytical and preparative separation of bark beetle pheromones by high performance liquid chromatography. *Journal of Chromatography* 363: 309–314. (bv ms).
- KUDON, LOUIS HARRY, AND CHARLES WAYNE BERISFORD 1985. Host preference behavior of insect parasites in the presence of southern pine beetle and one or more *Ips* species. Pages 74–86 in S. J. Branham and R. C. Thatcher, Integrated pest management research: symposium proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (ec).
- KUDRIHALT, U., M. PANIKE, AND K. WUST 1988. Wood destroying beetles: an updated literature survey. Bundesanstalt für Materialforschung und -Prüfung (BAM). Dokumentation Biologische Materialprüfung Series. 314 p. (ms).
- KULHAVY, DAVID LUMIR 1986. Forest protection. Pages 114–145 in D. L. Kulhavy and R. N. Conner, Wilderness and natural areas in the eastern United States: a management challenge. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. (cn).
- KULHAVY, DAVID LUMIR, R. A. GOYER, J. W. BING, AND M. A. RILEY 1989. *Ips* spp. natural enemy relationships in the Gulf Coast states. Pages 157–167 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (ec).
- KULHAVY, DAVID LUMIR, AND MITCHELL C. MILLER 1989. Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, School of Forestry, Nacogdoches, Texas. 255 p. (cn ec).
- KULHAVY, DAVID LUMIR, J. H. MITCHELL, AND R. N. CONNER 1985a. The southern pine beetle and the red-cockaded woodpecker: potential for interaction. International Congress of Entomology, Proceedings 15: 416. (cn ec).
- . 1985b. The southern pine beetle and the red-cockaded woodpecker: potential for interaction. Pages 337–343 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn ec).
- KURENZOV, ALEKSEI IVANOVICH 1963b. Entomological fauna of Kamchatka and the Bering arc problem in zoogeography. Pages 113–119 in J. L. Gressitt, Pacific basin biogeography: a symposium. Pacific Science Congress, Honolulu, Hawaii, 1961. Bishop Museum Press 10. 563 p. (ds tx).
- KVAMME, TORSTEIN 1955. New records of Norwegian Coleoptera II: species new to the fauna and notes on some little known species. *Fauna Norvegica* 32: 49–51. (ec ds).
- . 1958. *Trypodendron piceum* Strand (Col., Scolytidae): flight period and response to synthetic pheromones. *Fauna Norvegica* 35: 65–70. (hb).
- KVAMME, TORSTEIN, AND RUNE AXELSON 1953. Sphecidae (Hym., Aculeata) recorded from bark beetle pipe traps and some faunal notes from south Norway. *Fauna Norvegica, ser. B.* 30: 57–59. (ec).

L

- LACHANCE, DENIS, P. BENOIT, B. BONNEAU, AND GASTON LAFLAMME 1981. Insectes et maladies des arbres Quebec, 1980. *Forest Conservation* 47(9): 9. (cn).
- LACHANCE, DENIS, P. BENOIT, GASTON LAFLAMME, B. BONNEAU, AND R. PICHER 1984. Insectes et maladies des arbres Quebec, 1983. *Forest Conservation* 50(10): 7. (cn).
- LACHANCE, M. W., AND D. G. PFEIFFER 1959. Seasonal occurrences of wood boring beetles in Virginia commercial vineyards. *Hortscience* 24: 223. (cn ec).
- LANDIN, BENGT OLOF 1953. Insekter i farg. Almqvist and Wiksell, Stockholm. (hb ms).
- LANGWALD, J. 1989. Ein Beitrag zur Habitatselektion des Ameisenbuntkäfers, *Thanosinus formicarius* L. (Col., Cleridae) [Studies on the habitat selection of the bark beetle predator *Thanosinus formicarius*] [English summary]. *Anzeiger für Schädlingsskunde Pflanzen- und Umweltschutz* 62: 55–90. (ec).
- LANGOR, DAVID WILLIAM 1984. Flight ability of re-emerging eastern larch beetles, *Dendroctonus simplex*

- LeConte (Coleoptera: Scolytidae), following egg gallery formation and oviposition in tamarack, *Larix laricina* (Du Roi) K. Koch, in Newfoundland [Abstract]. Entomological Society of Alberta, 32nd annual meeting: proceedings. Mount Royal College, Calgary Alberta. 52 p. (bv hb).
- _____. 1955. Ecology of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae), in Newfoundland. Unpublished thesis, University of Newfoundland, St. Johns, Newfoundland. xvi + 197 p. (bv ec hb).
- _____. 1957. Flight muscle changes in the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae). Coleopterists Bulletin 41: 351–357. (ay).
- _____. 1958. Host effects on the population genetics and dynamics of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 18: 431. (ay hb).
- _____. 1959. Host effects on the phenology, development, and mortality of field populations of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). Canadian Entomologist 12: 149–157. (bv ec hb).
- _____. 1991. Arthropods and nematodes co-occurring with eastern larch beetle, *Dendroctonus simplex* (Col., Scolytidae) in Newfoundland. Entomophaga 36: 303–313. (ec).
- LANGOR, DAVID WILLIAM, AND ARTHUR G. RASKE. 1957a. Emergence, host attack, and overwintering behaviour of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae), in Newfoundland. Canadian Entomologist 119: 975–983 (bv hb).
- _____. 1957b. Reproduction and development of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae), in Newfoundland. Canadian Entomologist 119: 985–992. (hb).
- _____. 1958a. Annotated bibliography of the eastern larch beetle, *Dendroctonus simplex* LeConte (Coleoptera: Scolytidae). Government of Canada, Canadian Forestry Service, Newfoundland Forestry Centre, St. Johns, Newfoundland, Information Report N-X-266. 38 p. (ms).
- _____. 1958b. Mortality factors and life tables of the eastern larch beetle, *Dendroctonus simplex* (Coleoptera: Scolytidae) in Newfoundland. Environmental Entomologist 17: 959–963. (ec).
- _____. 1959a. A history of the eastern larch beetle *Dendroctonus simplex* (Coleoptera: Scolytidae) in North America. Great Lakes Entomologist 22: 139–154. (cn).
- _____. 1959b. The eastern larch beetle, another threat to our forests (Coleoptera: Scolytidae). Forestry Chronicle 1959(August): 276–279. (cn).
- LANGOR, DAVID WILLIAM, AND JOHN R. SPENSE. 1991. Host effect on allzyme and morphological variation of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). Canadian Entomologist 123: 395–410. (ay ec).
- LANGSTROM, BO E. 1956. Attack density and brood production of *Tomicus piniperda* in thinned *Pinus sylvestris* stems as related to felling date and latitude in Sweden. Scandinavian Journal of Forest Research 1: 351–357. (bv hb).
- LANGSTROM, BO E., AND C. HELLQVIST. 1955. *Pinus contorta* as a potential host for *Tomicus piniperda* L. and *Tomicus minor* (Hart.) (Col., Scolytidae) in Sweden. Journal of Applied Entomology 99: 174–181. (bv ec hb).
- _____. 1958a. Scots pine resistance against *Tomicus piniperda* as related to tree vitality and attack density. Pages 121–133 in T. L. Payne and H. Saarenmaa. Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn ec).
- _____. 1958b. Scots pine resistance against *Tomicus piniperda* as related to tree vitality and attack density. International Congress of Entomology, Proceedings 18: 413. (cn ec).
- LANIER, GERALD NORMAN. 1951. Larch beetle (*Dendroctonus simplex*). Page 3 in Forest insect and disease problems for New York state, 1950–51. Committee on insects and diseases, New York section, Syracuse, New York. (cn).
- _____. 1957a. Discovery of type material of some of Eichhoff's North American species of *Ips* and *Orthotomicus* (Coleoptera: Scolytidae). Coleopterists Bulletin 41: 107–110. (tx).
- _____. 1957b. The validity of *Ips cribricollis* (Eichh.) (Coleoptera: Scolytidae) as distinct from *I. grandicollis* (Eichh.) and occurrence of both species in Central America. Canadian Entomologist 119: 179–187. (ay tx).
- _____. 1959. Trap trees for control of Dutch elm disease. Journal of Arboriculture 15: 105–111. (ec).
- LANIER, GERALD NORMAN, J. P. HENDRICH, AND J. E. FLORES. 1958a. Biosystematics of the *Dendroctonus frontalis* (Coleoptera: Scolytidae) complex. Entomological Society of America, Annals 81: 403–418. (ay ds tx).
- _____. 1958b. Biosystematics of the *Dendroctonus frontalis* complex. International Congress of Entomology, Proceedings 18: 409. (ay tx).
- LANIER, GERALD NORMAN, AND A. H. JONES. 1955. Trap trees for elm beetles: augmentation with pheromone baits and chlorpyrifos. Journal of Chemical Ecology 11: 11–20. (bv cn).
- LANIER, GERALD NORMAN, AND LAWRENCE RICHARD KIRKENDALL. 1956. Karyology of pseudogamous *Ips* bark beetles. Hereditas 105: 87–96. (ay).
- LANIER, GERALD NORMAN, D. C. SCHUBERT, AND P. D. MANION. 1958. Dutch elm disease and elm yellows in central New York. Out of the frying pan into the fire. Plant Disease 72: 159–194. (ec).
- LANIER, GERALD NORMAN, S. A. TEALE, AND J. A. PAJARES. 1991. Biosystematics of the genus *Ips* (Coleoptera: Scolytidae) in North America: review of *Ips calligraphus* group. Canadian Entomologist 123: 1103–1124. (ay ds tx).
- LANNE, B. S., P. IVARSSON, P. JOHNSON, G. BERGSTROM, AND A. B. WASSGREN. 1959. Biosynthesis of 2 methyl-3-butene-2-ol, a pheromone component of *Ips typographus* (Coleoptera: Scolytidae). Insect Biochemistry 19: 163–168. (bv ms).
- LANNE, B. S., F. SCHLYTER, J. A. BYERS, J. LOFQVIST, A. LEUFVEN, G. BERGSTROM, J. N. C. VAN DER PERS, R. UNELIUS, P. BAECKSTROM, AND T. 1957. Differences in attraction to semiochemicals present in sympatric pine shoot beetles, *Tomicus minor* and *T. piniperda*. Journal of Chemical Ecology 13: 1045–1067. (bv).
- LANZ, W., AND J. BAESKOW. 1956. Bekämpfung des Kuperstechers durch Hackschneitzung von Fichtenschwamholz [Control of *Pityogenes chalcographus* by chipping of small-diameter Norway spruce wood] [English summary]. Forst- und Holzwirt 41: 178–182. (cn).

- LAPIS, E. B. 1955. Geographical and altitudinal distribution of the six-spined engraver beetle (*Ips calligraphus* Germar, Coleoptera: Scolytidae) in the Philippines. *Sylvatrop* 10: 211-217. (ec hb).
- LAPIS, E. B., AND H. O. SAN VALENTIN 1982. Protection of *Rhizophora mucronata* Lam. seedlings against *Pocillips fallax* Eggers (Coleoptera: Scolytidae) by cultural methods. *Sylvatrop* 7: 171-176. (cn).
- LARA, R. R. 1966. El combate directo de *Dendroctonus frontalis* Zimm. por derribo, descortezamiento y quema de la corteza de las arboles infestados. *Bosques* 3: 5-11. (cn hb).
- LAUT, JOHN G. 1955. Operational use of mountain pine beetle pheromones. Page 15 in *Thirty-sixth annual Western Forest Insect Work Conference, Proceedings*, 4-7 March 1955, Boulder, Colorado. Canada Department of Environment, Forestry Service, Northern Forest Research Centre, Edmonton, Alberta. 54 p. (bv cn).
- LAUT, JOHN G., LAWRENCE B. HELLBURG, DAVID A. LEATHERMAN, MICHAEL E. SCHOMAKER, AND KATE NISHIJIMA. 1978. Colorado forest insect and disease conditions, 1977. Colorado State Forest Service, Insect and Disease Division, Colorado State University, Fort Collins. 74 p. (cn).
- LAUT, JOHN G., MICHAEL E. SCHOMAKER, TERESA M. STIEGER, AND JOHN METZLER. 1979. Dutch elm disease: a bibliography (revised). Colorado State Forest Service, Colorado State University, Fort Collins. 165 p. (unpaginated). (ms).
- LAUT, JOHN G., AND THERESA M. STIEGER. 1980. Dutch elm disease: a bibliography, addendum 1979. Colorado State Forest Service, Colorado State University, Fort Collins. 26 p. (unpaginated). (ms).
- _____. 1981. Dutch elm disease: a bibliography, addendum 1980. Colorado State Forest Service, Colorado State University, Fort Collins. 30 p. (unpaginated). (ms).
- LAYTON, C. R. 1966. Annual district report: Peace River District, 1965. Pages 87-95 in *Annual district reports of the Forest Insect and Disease Survey Alberta*, 1965. Canada Department of Forestry, Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-1. (cn).
- _____. 1967. Annual district report: Lac La Biche District, 1966. Pages 62-72 in *Annual district reports of the Forest Insect and Disease Survey Alberta-Northwest Territories-Yukon region*, 1966. Canada Department of Forestry, Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-6. (cn).
- _____. 1969. Annual district report: Northwest District, 1968. Pages 45-54 in *Annual district reports of the Forest Insect and Disease Survey, Alberta-Northwest Territories-Yukon region*, 1968. Canada Department of Fisheries and Forestry, Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-22. (cn).
- _____. 1970. Annual district report: Northwest District, 1969. Pages 30-35 in *Annual district reports of the Forest Insect and Disease Survey Alberta-Northwest Territories-Yukon region*, 1969. Canada Department of Fisheries and Forestry, Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-30. (cn).
- LEATHERMAN, DAVID A. 1984. The mountain pine beetle. Colorado State Forest Service, Colorado State University, Fort Collins, [Circular, undated, ca 1984], No. 4. 2 p. (cn).
- LEATHERMAN, DAVID A., AND WHITNEY S. CRENSHAW. 1986. Firewood and house log insects in Colorado. Colorado State University Cooperative Extension Service, Service in Action, [Circular] No. 5.563. 2 p. (cn).
- LEATHERMAN, DAVID A., HEIDI E. MCNULTY, MICHAEL E. SCHOMAKER, AND DENNIS L. LYNCH. 1986. Aspen: a guide to common problems in Colorado. Colorado State Forest Service, Colorado State University, Fort Collins. 35 p. (cn).
- *LEBARON, WILLIAM. 1874. Outlines in entomology. Part first. Including the order Coleoptera. Fourth annual report on the Noxious and Beneficial Insects of the State of Illinois. ().
- LECUONA, R. E., P. M. FERNANDES, S. B. AIXES, AND E. BLEICHER. 1986. Patogenicidade de *Metarhizium anisopliae* (Metsch.) Sorok., a broca-do-café, *Hypothenemus hampei* (Ferrari, 1867) (Coleoptera: Scolytidae) [Pathogenicity of *Metarhizium anisopliae* to the coffee berry borer, *Hypothenemus hampei*] [In Portuguese, English summary]. *Anais da Sociedade Entomologica do Brasil* 15(supl.): 21-27. (ec).
- LEE, K. S. 1989. Geographic information systems (GIS) use in forest pest management. A simulated study on mountain pine beetle infestations [In Korean]. Korean Forestry Society, Journal 75: 168-176. (cn).
- LEE, Y. JIM, E. T. OSWALD, AND J. W. E. HARRIS. 1974. A preliminary evaluation of ERTS imagery for forest land management in British Columbia. Pages 87-101 in *Second Canadian symposium on remote sensing*, 29 April-1 May, University of Guelph, Guelph, Ontario. (cn ms).
- LEECH, HUGH BORDIN. 1973. [Book review]. *Pan-Pacific Entomologist* 49: 282-283. (ms).
- *LEEFMANS, SALOMON. 1928b. Takkenboek als bessenboek. *Bergcultures* 2: 924-925. ().
- LEGOWSKI, D. 1987. Obserwacje zmian opadu cętny na terenie nadleśnictwa Niedzwiedzi w latach 1977-1983 [Observations on twig fall in Niedzwiedzi Forest District in 1977-83] [In Russian, English summary]. *Sylvan* 131(9): 39-47. (cn).
- LELUAN, MONIQUE, G. LELUAN, AND CONSTANTIN CHARARAS. 1987. Caracteristiques physiologiques de l'arbrehote et installation de differents insectes secondaires (Scolytidae et Cerambycidae) [Physiological characteristics of host trees and establishment of various secondary insects (Scolytidae and Cerambycidae)] [English summary]. *Comptes Rendus de l'Academie des Sciences, III (Sciences de la Vie)* 305: 423-426. (bv).
- LEMPERIERE, G., AND D. BAILEY. 1986. Limousin: observation sur les manifestations du dendroctone de l'Epicea [Comments on the outbreaks of spruce bark beetle in Limousin]. *Forêts de France* 290: 18-22. (bv cn hb).
- LEMPERIERE, G., ET AL. 1989. Infestation des peuplements de *Picea* par *Dendroctonus micans* Kug. (Coleoptera: Scolytidae) en Limousin (France) et essais de lutte biologique avec le prédateur *Rhizophagus grandis* Gyll. (Coleoptera: Rhizophagidae) [Infestation of *Picea* population by *Dendroctonus micans* in Limousin, France and attempts at biological control with the predator *Rhizophagus grandis*] [English subtitle]. *Le Naturalist Canadien* 115: 235-243. (cn hb).
- *LENG, CHARLES WILLIAM, AND WILLIAM THOMPSON DAVIS. 1924. List of the Coleoptera of Staten Island, New

- York. Staten Island Institute of Arts and Sciences, Proceedings 2: 1-52. ().
- LENGERKEN, HANNS GERHARD VON. 1954. Die Brutfürsorge und Brutplageinstinkte der Käfer. Zweite Auflage. Akademische Verlagsgesellschaft Geest & Portig K.-G., Leipzig. (hb).
- LEON, J., AND NIRY BARRIO. 1987. Influencia de las fases lunares en la atracción de arboles trampa para escolitidos del género *Ips* en pinares de Baracoa [Influence of the phases of the moon in the attraction of trap trees for scolytids of the genus *Ips* in pine forests of Baracoa] [English summary]. *Revista Forestal Baracoa* 17: 45-56. (bv).
- *LEONTOVIC, R., J. PATOCKA, AND J. GREK. 1987. Vyskyt a význam hromadného hynutia dubov vo svete a na Slovensku [Occurrence and importance of mass decay of oak in the world and in Slovakia]. *Vedecke práce Vyskumneho ustavu lesneho hospodarstva vo Zvolene* 1987: 15-32. ().
- *LESNE, PIERRE. 1917. [*Scolytus rugulosus*]. *Journal de l'Agriculture Pratique*, Paris 30: 222-224. ().
- LESSARD, EUGENE D. 1985a. High country integrated pest management project: post suppression evaluation 1955. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Timber, Forest Pest and Cooperative Forestry Management, Biological Evaluation R2-55-4. 17 p. + appendix. (cn).
- _____. 1955b. Root diseases and bark beetles. Page 16 in Thirty-sixth annual Western Forest Insect Work Conference, Proceedings, 4-7 March 1955, Boulder, Colorado. Canada Department of Environment, Canadian Forestry Service, Northern Forest Research Centre, Edmonton, Alberta. 54 p. (cn ec).
- _____. 1959. Mountain pine beetle population manipulation strategies: past suppression practices. Pages 51-52 in G. D. Amman, Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- LESSARD, EUGENE D., MFG. HANSEN, AND MARY LU EILERS. 1956. Annotated bibliography: mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae). United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Timber, Forest Pest and Cooperative Forestry Management, Technical Report R2-35. 158 p. (cn ms).
- LESSARD, EUGENE D., D. W. JOHNSON, T. E. HINDS, AND W. H. HOPKINS. 1955. Association of *Amillaria* root disease with mountain pine beetle infestations on the Black Hills National Forest, South Dakota. United States Department of Agriculture, Forest Service, Forest Pest Management, Methods Application Group, Fort Collins, Colorado, Report 55-4. 6 p. + appendix. (cn).
- LEUFVEN, ANDERS, G. BERGSTROM, AND E. FALSEN. 1955. Oxygenated monoterpenes produced by yeasts isolated from *Ips typographus* (Coleoptera: Scolytidae) and grown in phloem medium. *Journal of Chemical Ecology* 14: 353-362. (ec).
- LEUFVEN, ANDERS, AND G. BIRGERSSON. 1957. Quantitative variation of different monoterpenes around galleries of *Ips typographus* (Coleoptera: Scolytidae) attacking Norway spruce. *Canadian Journal of Botany* 65: 1035-1044. (bv cn).
- LEUFVEN, ANDERS, AND L. NEHLS. 1956. Quantification of different yeasts associated with the bark beetle, *Ips typographus*, during its attack on a spruce tree. *Microbial Ecology* 12: 237-243. (ec).
- LEUNIS, JOHANNES. 1856. Synopsis der Thierkinde, ein handbuch für höhere Lehranstalten. Zweiter Band [Scolytidae, p. 178-183]. Hahn'sche Buchhandlung, Hannover. (cn hb).
- LEUSCHNER, WILLIAM A., AND P. BERCK. 1955a. Decision analysis. Pages 177-189 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn).
- _____. 1955b. Impacts on forest uses and values. Pages 105-120 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn).
- *LEVER, R. J. A. W. 1935. Entomological notes, 3. Timber-boring beetles. *Fiji Agriculture Journal* 9: 15, 18. ().
- *_____. 1942. Entomological notes, 1. A shot-hole borer attacking rubber trees. *Fiji Agriculture Journal* 13: 24. ().
- LEVIEUX, JEAN, H. JACTEL, AND FRANCOIS LIEUTIER. 1985. Etude préliminaire de la variabilité de la pression de séve de clones de Pin sylvestre dans le centre de la France [Preliminary study of the sap pressure variability between several clones of Scots pine in the center of France] [English summary]. *Annales des Sciences Forestières* 45: 341-355. (cn).
- LEVIEUX, JEAN, FRANCOIS LIEUTIER, AND A. DELPLANQUE. 1955a. Les scolytes ravageurs de l'épicéa [Scolytid pests of Norway spruce] [English summary]. *Revue Forestière Française (Nancy)* 37: 347-355. (cn ec hb).
- _____. 1955b. Les scolytes ravageurs du pin sylvestre [Scolytid pests of Scots pine] [English summary]. *Revue Forestière Française (Nancy)* 37: 431-440. (cn hb).
- LEVIEUX, JEAN, FRANCOIS LIEUTIER, JOHN CONRAD MOSER, AND T. J. PERRY. 1959. Transportation of phytopathogenic fungi by the bark beetle *Ips sexdentatus* Boerner and associated mites. *Journal of Applied Entomology* 108: 1-11. (ec).
- LEWIS, K. J., AND S. A. ALEXANDER. 1956. Insects associated with the transmission of *Verticilladiella procera*. *Canadian Journal of Forest Research* 16(6): 1330-1333. (ec).
- LEXON, BERT R. 1939. An experiment in the use of sodium arsenite in the thinning ponderosa pine. *Journal of Forestry* 37: 259-262. (cn).
- LI KUAN SHENG. 1959. Scolytidae species on *Pinus armandii* and the preliminary biology observation of *Dendroctonus* sp. [In Chinese]. Report of preliminary studies on forest insect pests. 234 p. (hb).
- LI KUAN SHENG, SHI QUAN CHANG, AND AO HENG YI. 1954. Shaanxi forest tree diseases and insects, pictorial monograph, Volume 2 [In Chinese]. Shaanxi Province Peoples Publication House, Xian. 344 p. (hb ds).
- LI KUAN SHENG, DANG XINDE, AND SHI QUAN CHANG. 1977. Shaanxi forest tree diseases and insects, pictorial monograph, Volume 1 [In Chinese]. Shaanxi Province Peoples Publication House, Xian. 218 p. (hb ds).
- LI KUAN SHENG, AND J. ZHOU. 1950. Forest insects in China [In Chinese]. Chinese Forestry Press. 218 p. (ec hb).
- LIBBEY, L. M., LEE C. RYKER, AND K. L. YANDELL. 1955. Laboratory and field studies of volatiles released by *Dendroctonus ponderosae* Hopkins (Coleoptera:

- Scolytidae). *Journal of Applied Entomology* 100: 381–392. (bv).
- *LICHTENSTEIN, JEAN L., AND FRANÇOIS PICARD. 1920. [*Scolytus rugulosus*]. *Société Entomologique de France, Paris, Bulletin* 1920: 54–55. ().
- LIEBHOLD, ANDREW M., P. BERCK, N. A. WILLIAMS, AND DAVID LEE WOOD. 1955. Estimating and valuing western pine beetle impacts. *Forest Science* 32: 325–338. (cn).
- LIEUTIER, FRANÇOIS. 1955. Influence de nematodes parasites sur l'hémolymphe des adultes d'*Ips sexdentatus* Boern. (Coleoptera: Scolytidae) [Influence of parasitic nematodes on the hemolymph of *Ips sexdentatus* Boern.] [English summary]. *Ann. Parasitol. Hom. Comp.* 60(2): 195–204. (ay ec).
- LIEUTIER, FRANÇOIS, AND ALAN ANDREW BERRYMAN. 1955a. Elicitation of defensive reactions in conifers. Pages 313–320 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (ec).
- . 1955b. Preliminary histological investigations of the defense reactions of three pines to *Ceratocystis clavigera* and two chemical elicitors. *Canadian Journal of Forest Research* 15: 1243–1247. (ec).
- LIEUTIER, FRANÇOIS, C. CHENICLET, AND J. GARCIA. 1959. Comparison of the defense reactions of *Pinus pinaster* and *Pinus sylvestris* to attacks by two bark beetles (Coleoptera: Scolytidae) and their associated fungi. *Environmental Entomology* 18: 228–234. (ec).
- LIEUTIER, FRANÇOIS, T. FAURE, AND J. GARCIA. 1955. Attacks by scolytids and dieback of the Scots pine in the Provence Cote d'Azur region France. *Revue Forestière Française* (Nancy) 40: 224–232. (cn ec).
- LIEUTIER, FRANÇOIS, AND G. T. FERRELL. 1955a. Relationships between indexes of tree vigour and the induced defense reaction of Scots pine to a fungus associated with *Ips sexdentatus* (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 15: 430. (ec).
- . 1955b. Relationships between indexes of tree vigour and the induced defense reaction of Scots pine to a fungus associated with *Ips sexdentatus* Boern. (Coleoptera: Scolytidae). Pages 163–178 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (ec hb).
- *LIEUTIER, FRANÇOIS, R. HAM, M. C. HAM, AND J. GARCIA. 1956. Une méthode de marquage individuel des coleopteres scolytides pour les études de laboratoire [A method of marking individual scolytid beetles for laboratory studies] [English summary]. *Agronomie* 6: 773–776. ().
- LIEUTIER, FRANÇOIS, M. JASTRABSKY, AND P. BONNAFE. 1955a. Modifications of the hemolymph and ovarian protein pattern induced by parasitic nematodes in *Ips sexdentatus* (Coleoptera: Scolytidae) [In French, English summary]. *Ann. Parasitol. Hom. Comp.* 60(6): 715–726. (ay ec).
- . 1955b. Variations of the proteinograms of the hemolymph, ovaries and fat body during the adult life of *Ips sexdentatus* (Coleoptera: Scolytidae) [In French, English summary]. *Bull. Soc. Zool. France* 109(3): 279–300. (ay ec).
- LIEUTIER, FRANÇOIS, AND JEAN LEVIEUX. 1955. Les relations coniferes-scolytides: importance et perspectives de recherches [The extent of bark beetle/conifer interactions and proposed research] [English summary]. *Annales des Sciences Forestières* 42: 359–370. (cn).
- LIEUTIER, FRANÇOIS, AND A. YART. 1959. Temperature preference of the fungi associated with *Ips sexdentatus* Boern. and *Tomicus piniperda* L. (Coleoptera: Scolytidae) [In French]. *Annales des Sciences Forestières* 46: 411–415. (ec).
- LIEUTIER, FRANÇOIS, A. YART, J. GARCIA, M. C. HAM, M. MORELET, AND JEAN LEVIEUX. 1959. Champignons phytopathogènes associés à deux coleoptères Scolytidae du pin sylvestre (*Pinus sylvestris* L.) et étude préliminaire de leur agressivité envers l'hôte [English summary]. *Annales des Sciences Forestières* 46: 210–216. (ec).
- LIEUTIER, FRANÇOIS, A. YART, J. GARCIA, B. POUPINEL, AND JEAN LEVIEUX. 1958. Do fungi influence the establishment of bark beetles in Scots pine? Pages 321–334 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (ec).
- LIII, M. P., AND F. M. STEPHEN. 1953. Use of computer simulation models to predict expected tree mortality and monetary loss from SPB spots: a research update. United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia, Forest Pest Management Report SPB fact sheet 26. 2 p. (cn).
- LINDELOW, AKE, AND R. AXELSSON. 1955. Svarta bastborrarforbisedda skadegorare i skogsplanteringar. *Skogen* 2–85: 43–44. (ec).
- LINDELOW, AKE, AND H. IACOBÆUS. 1955. Insekter i obarkade contortastammar [Insects found in unbarked stems of lodgepole pine] [In Swedish, English summary]. *Sveriges Skogvårdsförbunds Tidskrift* 2: 43–45. (cn).
- LINDELOW, AKE, AND J. WESLIEN. 1956. Sex-specific emergence of *Ips typographus* L. (Coleoptera: Scolytidae) and flight behavior in response to pheromone sources following hibernation. *Canadian Entomologist* 118: 59–67. (bv hb).
- LINDGREN, B. STAFFAN. 1956. *Trypodendron lineatum* (Coleoptera: Scolytidae) breeding in big leaf maple, *Acer macrophyllum*. *Entomological Society of British Columbia, Journal* 83: 44. (ds).
- . 1955. Application of semiochemicals for management of *Dendroctonus* bark beetles. *Northwest Environmental Journal* 4: 327–325. (cn).
- . 1990. Ambrosia beetles: semiochemical technology offers potential management tool. *Journal of Forestry* 1990(February): 8–11. (cn ms).
- LINDGREN, B. STAFFAN, AND JOHN HARVEY BORDEN. 1959. Semiochemicals of the mountain pine beetle (*Dendroctonus ponderosae* Hopkins). Pages 83–85 in G. D. Amman, Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (bv cn).
- LINDGREN, B. STAFFAN, JOHN HARVEY BORDEN, C. H. CUSHION, L. J. CHONG, AND C. J. HIGGINS. 1959. Reduction of mountain pine beetle (Coleoptera: Scolytidae) attacks by verbenone in lodgepole pine stands in British Columbia. *Canadian Journal of Forest Research* 19: 65–68. (cn).
- LINDGREN, B. STAFFAN, MARK D. MCGREGOR, ROBERT D. OAKES, AND HUBERT E. MEYER. 1958. Effects of

- MCH and baited Lindgren traps on Douglas-fir beetle attacks on felled trees. *Journal of Applied Entomology* 105: 289–294. (bv cn).
- _____. 1989. Suppression of spruce beetle attacks by MHIC released from bubble caps. *Western Journal of Applied Forestry* 4(2): 49–52. (bv cn).
- LINDSTROM, M., T. NORIN, G. BIRGERSSON, AND F. SCHYTER. 1989. Variation of enantiomeric composition of alpha pinene in Norway spruce and its influence on the production of verbenol isomers by *Ips typographus* in the field. *Journal of Chemical Ecology* 15: 541–548. (bv).
- LINNANE, JAMES P. 1985. Biological evaluation—spruce beetle-population trend and tree losses. Fort Apache Indian Reservation, Arizona. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico, Forest Pest Management Report R3–55–5. 31 p. (cn lb).
- *LINSLEY 1944. [*Pagiocerus* spp.]. *Hilgardia* 16(4): 202. ().
- LINTON, D. A., AND LASZLO SAFRANYIK. 1988. The spruce beetle, *Dendroctonus rufipennis* (Kirby): an annotated bibliography 1857–1987. Government of Canada, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-298. 39 p. (ms).
- LINTON, D. A., LASZLO SAFRANYIK, AND R. BETTS. 1987. Field techniques for rearing and marking mountain pine beetle for use in dispersal studies. Entomological Society of British Columbia. *Journal* 84: 53–57. (cn lb).
- *LISKA, J., MILOS KNIZEK, AND P. KAPITOLA. 1989. Vazne ohrozeni blatkovych prostu na raselinisti Zofinka v jiznich Cechach [Heavy damage of *Pinus rotundata* on the Zofinka peat-hog in southern Bohemia]. *Ziva* 6: 247–248. ().
- *LISKA, J., P. ZAHRADNIK, V. PICHIOVA, AND MILOS KNEZEK. 1991. Zhodnoceni ochrany lesa proti hmyzim skudum v CSFR ve srovnani s ostatnimi staty [Evaluation of forest protection against insect pests in Czechoslovakia compared to other countries]. Pages 1–8 in *Sbornik konference, Ekologicka ochrana lesu-Prachatic, 4–5 September 1991, Spolana*. ().
- LISTER, C. KENDALL, AND J. G. LAUT. 1957. MPB pheromone trials in Colorado, 1955. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Timber, Forest Pest and Cooperative Forestry Management, Technical Report R2–40. 8 p. (bv cn).
- LIU, Y. B., AND J. A. McLEAN. 1989. Field evaluation of responses of *Gnathotrichus sulcatus* and *Gnathotrichus retusus* (Coleoptera: Scolytidae) to semiochemicals. *Journal of Economic Entomology* 82: 1687–1690. (bv).
- LIU, Z. F., B. Y. ZHAO, S. F. ZHAO, M. L. SHENG, Q. H. ZHANG, Z. Q. CUI, D. CHU, Y. J. SUN, S. B. XU, AND X. D. ZHANG. 1989. A report on monitoring stem borers in burned areas in Daxinganling Mountain [In Chinese]. *Forest Science and Technology* 2: 12–15. (cn).
- LIVESEY, F. 1966. Status of insects in the Tweed District. Pages A22–31 in *Annual district reports of the Forest Insect and Disease Survey Ontario, 1965*. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- LIVINGSTON, ROBERT LADD. 1987a. Evaluation of Douglas-fir beetle infestations, Pend Oreille Lake Supervisory Area, October 1987. Idaho Department of Lands (Coeur d'Alene, Idaho), Report 87–15. 4 p. (cn).
- _____. 1987b. Evaluation of pine engraver, western pine beetle, and western spruce budworm on the southwest Idaho Supervisory Area, March 1987. Idaho Department of Lands, Coeur d'Alene, Idaho, Report 87–4. 5 p. (cn).
- _____. 1989. Detection and survey methods. Page 21 in G. D. Amman, Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- *LIVINGSTON, ROBERT LADD, AND J. W. SCHWANDT. 1987. Evaluation of bark beetle and root rot activity in the Mad Dog Timber Sale, Priest Lake Supervisory Area, May 1987. Idaho Department of Lands, Coeur d'Alene, Report 87–6. 5 p. ().
- LIVINGSTON, ROBERT LADD, J. W. SCHWANDT, D. P. BECKMAN, S. GUST, ET AL. 1988. Idaho forest pest conditions and program summary, 1987. United States Department of Agriculture, Forest Service, Northern and Intermountain Regions and Idaho Department of Lands (Coeur d'Alene, Idaho), Report 88–1. 28 p. (cn).
- LODGE, H. P. 1945. Catalogue of the beetles of Alabama. Geological Survey of Alabama. Monograph 11. 172 p. (ds).
- LOGAN, J. A., AND GENE DOYLE AMMAN. 1986. A distribution model for egg development in mountain pine beetle. *Canadian Entomologist* 118: 361–372. (ec).
- LOOMIS, ROBERT C., AND T. H. HOFACKER. 1981. Forest insect and disease conditions in the United States, 1980. United States Department of Agriculture, Forest Service, Report, Washington, D.C. 36 p. (cn).
- LOOMIS, ROBERT C., T. H. HOFACKER, AND S. M. TUCKER. 1985. Forest insect and disease conditions in the United States 1984. United States Department of Agriculture, Forest Service, Washington, D.C., vi + 90 p. (cn).
- _____. 1986. Forest insect and disease conditions in the United States 1985. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C. 95 p. (cn).
- LORIO, PETER LEONCE, JR. 1986. Growth-differentiation balance: a basis for understanding southern pine beetle-tree interactions. *Annales Entomologici Fennici* 14: 259–273. (cn).
- _____. 1988. Growth differentiation-balance relationships in pines affect their resistance to bark beetles (Coleoptera: Scolytidae). Pages 73–92 in W. J. Mattson, J. Leveux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York, xiv + 416 p. (ec).
- LORIO, PETER LEONCE, JR., AND SUSAN J. BRANHAM. 1988. Where'd the beetle go? *Forests and People* 36(3): 18–20, 36–37. (cn ms).
- *LORIO, PETER LEONCE, JR. AND J. D. HODGES. 1985. Theories of interactions among bark beetles, associated microorganisms, and host trees. Pages 485–492 in United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-54. ().
- LORIO, PETER LEONCE, JR., AND ROBERT A. SOMMERS. 1985. Potential use of soil maps to estimate southern pine beetle risk. Pages 239–245 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, South-

- ern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- _____. 1956. Evidence of competition for photosynthates between growth processes and oleoresin synthesis in *Pinus taeda* L. Pages 301–306 in R. J. Luxmoore, J. J. Landsberg, and M. R. Kaufmann, Coupling of carbon, water and nutrient interactions in woody plant soil systems. *Tree Physiology* 2(1–3). (cn).
- LOSEKRUIG, R. C. 1955. Befall von Buchenaltholzer durch den Laubnutzholzborkenkäfer [Attack on old beech trees by *Trypodendron domesticum*] [English summary]. *Allgemeine Forstzeitschrift* 34: 942–943. (bv cn).
- * LOYTTYNEMI, KARL. 1950a. A preliminary annotated list of pine insects in Zambia. Division of Forest Research, Zambia. Research Note 26. 13 p. ().
- * _____. 1950b. A preliminary study of insects attacking freshly cut logs of some exotic plantation tree species in Zambia. Division of Forest Research, Zambia. Research Note 27. 17 p. ().
- * _____. 1950c. A preliminary annotated list of Eucalyptus insects in Zambia. Division of Forest Research, Zambia. Research Note 28. 19 p. ().
- LOYTTYNEMI, KARL, ROGER A. BEWER, AND R. LOYTTYNEMI. 1955. Annual flight patterns of timber insects on Miombo woodland in Zambia. *Platypodidae* (Coleoptera). *Annales Entomologici Fennici* 51: 27–29. (bv hb).
- LOYTTYNEMI, KARL K., HELIOVAARA, AND S. REPO. 1955a. No evidence of a population pheromone in *Tomicus piniperda* (Coleoptera, Scolytidae): a field experiment. *Annales Entomologici Fennici* 54: 93–95. (bv).
- _____. 1955b. Pine turpentine strongly attracts *Hylastes brunneus* (Coleoptera, Scolytidae). *Annales Entomologici Fennici* 54: 145–148. (bv).
- LUCIET, WILHELM H. 1957. Die Käfer Mitteleuropas. Katalog. Goecke and Evers, Krefeld. 342 p. (ds tx).
- LUIK, A. 1956. Temperature as the principal exogenous regulator of winter dormancy in xylophages [In Russian, English summary]. *Metsanduslikud uurimused, Estonian SSR* 21: 98–102. (ec hb).
- LUKASHIIEVA, N. V. 1956. Successional changes in the complex of xylophilous Diptera on pine [In Russian]. *Trudy Vsesoyuznogo Entomologicheskogo Obschestva* 65: 89–92. (ec).
- LUNDBERG, STIG. 1974. Skallbaggar funna i Messaureområdet, Lule lappmark (Coleoptera) [Beetles found in the Messaure area, Lule Lapmark (Coleoptera)] [In Swedish, English summary]. *Entomologisk Tidskrift* 95: 85–94. (ds).
- _____. 1979. Fangst av skallbaggar med hjälp av fönsterfallor [Catching beetles with windowtraps (Coleoptera)] [In Swedish, English summary]. *Entomologisk Tidskrift* 100: 29–32. (ds).
- _____. 1980. För Norrbotten nya skallbaggar under tioårsperioden 1969–75 [Beetles new to the province of Norrbotten during the years 1969–75] [In Swedish, English summary]. *Entomologisk Tidskrift* 101: 147–150. (ds).
- _____. 1981. Gotska Sandons skallbaggsfauna: nytillskott och intressanta arter [The beetle fauna of Gotska Sandon: new findings and interesting species] [In Swedish, English summary]. *Entomologisk Tidskrift* 102: 147–154. (ds).
- _____. 1985a. Catalogus Coleopterorum Sueciae 1986: rattelser och tillägg. *Entomologisk Tidskrift* 109: 81–85. (ds tx).
- _____. 1955b. Nagra intressanta skallbaggsfynd 1 till Norrbotten importerat barrvirke [Some interesting records of beetles in coniferous timber imported into Norrbotten] [In Swedish, English summary]. *Entomologisk Tidskrift* 109: 49–50. (ds).
- LUNDHOLM, BENGT. 1959. Pest control and science foundations. *Holarctic Ecology* 12: 354–356. (ms).
- LUO, LU-XI. 1974. A preliminary report on the life cycle and control methods of *Sphacrotypes coimbatorensis* Stebbing. *Kunchong Zhishi* 1: 32–33. (cn).
- _____. 1984. Study on the damage of the camphor *Xyleborus* (*X. interjectus* Blandford) and its control. *Kunchong Zhishi* 21: 166–167. (cn).
- LUTYK, P. 1954. Żerzy cecynka wiekszego (*Tomicus Myelophilus piniperda* L.) w pedach świerka i jodły [Feeding of *Tomicus piniperda* and shoots of Norway spruce and silver fir]. *Sylvan* 128(3): 65–68. (bv ec).
- _____. 1955. Występowanie i liczebność generacji siostrzanej cecynka wiekszego (*Tomicus piniperda* L.) w Polsce [Occurrence and size of a sister generation of the pine shoot beetle *Tomicus piniperda* in Poland] [In Polish, English and Russian summaries]. *Sylvan* 132(10): 53–60. (hb).
- LYONS, L. A. 1957b. Insects affecting seed production in red pine II: *Diorctetria disculsa* Heinrich, *D. abietella* (D. & S.), and *D. cambiiicola* Dyar (Lepidoptera: Phycitidae). *Canadian Entomologist* 89: 70–79. (cn ec).

M

- MACELSKI, M. 1954. Biotechnicki insekticidi s posebnim osvrtom na feromone [Biotechnical insecticides with particular emphasis on pheromones]. *Poljoprivredna Znanstvena Smotra* 67: 609–620. (cn).
- MACGILLIVRAY, A., AND C. O. HOUGHTON. 1902. A list of insects taken in the Adirondack Mountains. New York, *Entomological News* 13: 247–253. (ds).
- MACLAUCHLAN, L. E., JOHN HARVEY BORDEN, M. R. CACKETTE, AND J. M. D'ACURIA. 1957. A rapid multisample technique for detection of trace elements in trees by energy-dispersive x-ray fluorescence spectroscopy. *Canadian Journal of Forest Research* 17: 1124–1130. (ms).
- MACLAUCHLAN, L. E., JOHN HARVEY BORDEN, AND J. M. D'ACURIA. 1955. Distribution of arsenic in lodgepole pines treated with MSMA. *Western Journal of Applied Forestry* 3: 37–40. (cn).
- MACLAUCHLAN, L. E., JOHN HARVEY BORDEN, J. M. D'ACURIA, AND L. A. WHEELER. 1955. Distribution of arsenic in MSMA-treated lodgepole pines infested by the mountain pine beetle, *Dendroctonus ponderosae* (Coleoptera: Scolytidae), and its relationship to beetle mortality. *Journal of Economic Entomology* 81: 274–280. (cn).
- MADDEN, JOHN LEO H. D. PIERCE, JR., JOHN HARVEY BORDEN, AND A. BUTTERFIELD. 1955. Sites of production and occurrence of volatiles in Douglas-fir beetle, *Dendroctonus pseudotsugae* Hopkins. *Journal of Chemical Ecology* 14: 1305–1317. (bv).
- MAGAS, LASZLO P. 1977b. Eastern larch beetle: another Maritime forest enemy. Canada Department of Environment, Canadian Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Pith to Peridern 11(7): 28. (cn).
- _____. 1979. Forest pest conditions in the Maritimes in 1978 with an outlook for 1979. Pages 7–9 in *Canada*

- Department of Environment, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-98. 34 p. (cn).
- _____. 1982b. Maritimes region. Pages 17–28 in Annual report of the Forest Insect and Disease Survey, 1978. Canada Department of Environment, Forestry Service, Ottawa, Ontario. (cn).
- _____. 1984b. Important forest pests of larch in the Maritimes. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Technical Note 98. 4 p. (cn).
- _____. 1985. Forest pest conditions in the Maritimes in 1984. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-154. 49 p. (cn).
- _____. 1986a. A synopsis of forest pest conditions in the Maritimes in 1986. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Technical Note 172. 4 p. (cn).
- _____. 1986b. Forest pest conditions in the Maritimes in 1985. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-159. 55 p. (cn).
- _____. 1987. Forest pest conditions in the Maritimes in 1986. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-161. 67 p. (cn).
- _____. 1988. Forest pest conditions in the Maritimes in 1987. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-166. 109 p. (cn).
- _____. 1989. Forest pest conditions in the Maritimes in 1988. Forestry Canada, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-174. 76 p. (cn).
- MAHLER, VIGGO. 1987. Sjette tillæg til Fortegnelse over Danmarks biller (Coleoptera) [Sixth supplement to the list of Danish Coleoptera] [In Danish, English summary]. Entomologiske Meddelelser 54: 181–235. (ds).
- MAITI, P. K., AND NIVEDITA SAHA. 1986. Contributions to the knowledge of the bark and timber beetles (Scolytidae: Coleoptera) of the Andaman and Nicobar Islands. Records of the Zoological Survey of India, Miscellaneous Publication. Occasional Paper 86. 152 p. (ds tx).
- _____. 1987. Indian species of the genus *Arixyleborus* Hopkins (Scolytidae: Coleoptera). Bulletin of the Zoological Survey of India 8: 1–12. (ds tx).
- _____. 1988. Descriptions of three new species of the genus *Hyorrhynchus* Blandford (Coleoptera, Scolytidae). Proceedings of the Zoological Society (Calcutta, India) 41: 41–51. (tx).
- MAJZLAN, OTO, I. RYCHLIK, M. HOLECOVA, AND T. KOZISEK. 1987. Beetles of the Abrod State Nature Reserve [In Czech, English summary]. Ochrana Přírody 8: 177–201. (ds).
- MAKSYMOWIC, MIŁOS. 1986a. Laboratorijska ispitivanja parazita potkornjaka bresta [Laboratory investigations on the parasites of elm bark beetles] [In Serbo-Croatian, English summary]. Zastita Bilja 37: 5–20. (ec).
- *MAKSYMOWIC, J. K. 1977. Trockenheit 1976 und Borkenkäfersituation im Frühjahr 1977. Schweizerischen Zeitschrift für Forstwesen 128: 154–160. ().
- _____. 1987. Erstmaliger Massenbefall des Schwarzen Nutzholzborkenkäfer *Xylosandrus germanus* Blandf., in der Schweiz. [First mass attack of *Xylosandrus germanus* in Switzerland] [French, Italian, English summaries]. Schweizerische Zeitschrift für Forstwesen 138(3): 215–227. (bv cn ec).
- MAKOV, KONSTANTIN. 1905. Die schädlichsten Insekten und Pflanzenkrankheiten, welche an den Kulturpflanzen in Bulgarien während des Jahres 1903 geschädigt haben. Pflanzenkrankheiten 15: 50–53. (cn).
- *MAMAIEV. 1929. [*Hylastes ater*]. Izv. sibirsk. kraev. Stantz. Rast. 3(6): 134–142. ().
- MAMAIEV, YU. B. 1986. Foci of xylophages [In Russian]. Zashchita Rastenii Moscow 9: 31. (bv).
- MANKINS, J. V. 1980. El barrenador de la corteza de las coníferas de Centroamérica, género *Ips* [The bark beetles of Central American conifers, genus *Ips*] [English summary]. Nota Técnica, Escuela Nacional de Ciencias Forestales, Honduras No. 2. 4 p., 10 figs. (cn hb).
- MANKOVSKA, B. 1987. The content of Pb and Cd in the developmental stages of important oak pests (Coleoptera). Biologia 42: 565–569. (ec).
- MANOJLOVIC, B. 1986b. Prilog proucavanju napada potkornjaka bresta sa posebnim osvrtom na znacaj malog brestovog potkornjaka *Scolytus multistriatus* Marsh. (Coleoptera, Scolytidae) [Contribution of the study of attack of elm by bark beetles with particular reference to the smaller elm bark beetle *Scolytus multistriatus*] [In Serbo-Croatian, English summary]. Zastita Bilja 37: 297–310. (bv hb).
- _____. 1986c. Prilog proucavanju razvika malog brestovog potkornjaka *Scolytus multistriatus* Marsh. (Coleoptera, Scolytidae) [Contribution to study of the development of *Scolytus multistriatus*]. Zastita Bilja 37: 251–270. (ec hb).
- MANSINGH, AJAI AND L. F. RHODES. 1985. Residual toxicity of various insecticidal formulations to the coffee berry borer, *Hypothenemus hampei* Ferrari (Scolytidae: Coleoptera). Insect Science and Its Application 6: 209–212. (cn).
- MANVILLE, J. F., L. H. McMULLEN, AND K. J. REIMER. 1988. Impact and role of monosodium methanearsonate on attack and progeny production by the Douglas-fir beetle (Coleoptera: Scolytidae) in lethal trap trees. Journal of Economic Entomology 8: 1691–1697. (cn hb).
- MARKOV, V. A. 1985. The great spruce bark beetle in the forests of the Ryazan region [In Russian]. Lesnoe Khozaystvo 9: 59–60. (cn).
- MARTIN, E., AND JOSE M. COBOS. 1986. Graves ataques de perforadores en los abetares de Anso (Huesca) [Serious attacks by borers in the fir plantations of Anso (Huesca)] [English summary]. Boletín de Sanidad Vegetal, Plagas [Spain] 12: 297–298. (cn).
- MARTINAT, P. J. 1987. The role of climatic variation and weather in forest insect outbreaks. Pages 241–268 in P. Barbosa and J. C. Schultz, Insect outbreaks. Academic Press, San Diego, California. xiv + 578 p. (ec).
- MARTINEAU, RENE. 1985. Insectes nuisibles des forêts de l'est du Canada. Marcel Broquet, LaPrairie, Quebec. 283 p. (cn hb).
- MASON, CARLAND N., PETER L. LORIO, JR., ROGER P. BELANGER, AND W. L. NETTLETON. 1985. Rating the susceptibility of stands to southern pine beetle attack.

- United States Department of Agriculture, Forest Service, Cooperative State Research Service, Washington, D.C., Agriculture Handbook 645. 31 p. (cn).
- MASUTTI, LUIGI. 1968b. Coleotteri Scolitidi dei pini d'altitudine in alcuni biotopi delle alpi e sul Massiccio del M. Pollino. *Archivio Botanico e Biografico Italiano* 44: 213-223. (cc ds).
- * _____. 1977. Vicissitudini di insetti e di piante nella storia recente delle biocenosi terrestri [Insect and plant vicissitudes in the recent history of terrestrial biocenoses]. Gruppo geomorfologico ed agrario, Il montanaro d'Italia-Monti e boschi 27(3): 37-44. (.)
- MATHEW, G. 1955. Some coleopteran predators associated with timber beetles in Kerala (India). *Entomon* 10(2): 179-181. (cc).
- _____. 1956. Insects associated with forest plantations of *Gmelina arborea* Roxb. in Kerala, India. *Indian Journal of Forestry* 9: 308-312. (ds).
- _____. 1957. Insect borers of commercially important stored timber in the state of Kerala, India. *Journal of Stored Products Research* 23: 185-190. (ds).
- MATSON, PAMELA A., F. P. HAIN, AND W. MAWBY. 1957. Indices of tree susceptibility to bark beetles vary with silviculture treatment in a loblolly pine plantation. *Forest Ecology and Management* 22: 107-118. (cn).
- MATTESON, DONALD S., K. M. SADIU, AND M. L. PETERSON. 1956. Chirally selective synthesis via pinanediol boronic esters: insect pheromones, diols, and an amino alcohol. *Journal of the American Chemical Society* 108: 810-819. (ms).
- MATTSON, WILLIAM JOHN. 1956. Competition for food between two principal cone insects of red pine, *Pinus resinosa*. *Environmental Entomology* 15: 88-92. (cc).
- MATTSON, WILLIAM JOHN, AND ROBERT A. HAACK. 1957. The role of drought stress in provoking outbreaks of phytophagous insects. Pages 365-407 in P. Barbosa and J. C. Schultz. *Insect outbreaks*. Academic Press, San Diego, California. xiv + 578 p. (cn cc).
- MATUASHVILI, S. I., G. A. TSILOSANI, V. PALWANDISHVILI, AND T. SH. IMNADZE. 1974. Effect of some crystalliferous entomopathogenic microorganisms on the European spruce beetle (*Dendroctonus micans* Kugel.) [In Russian, Armenian]. *Biol. Zh. Armenia* 1974: 60-64. (cc).
- MAWBY, WILLIAM DAVID, AND FRED P. HAIN. 1955a. Evaluation of procedures for estimating within-spot populations of *Dendroctonus frontalis* (Coleoptera: Scolytidae) during a collapsing epidemic. *Canadian Entomologist* 117: 1083-1091. (cn).
- _____. 1955b. The large-scale prediction of southern pine beetle populations. Pages 53-55 in S. J. Branham and R. C. Thatcher. *Integrated pest management research symposium: the proceedings*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- MAWBY, WILLIAM DAVID, FRED P. HAIN, AND C. A. DOGGETT. 1959. Endemic and epidemic populations of southern pine beetle: implications of the two-phase model for forest management. *Forest Science* 35: 1075-1087. (hb cn).
- MAY, CURTIS, AND W. L. BAKER. 1952. Insects and spread of forest-tree diseases. Pages 677-682 in A. Stefferud, Insects. United States Department of Agriculture, Washington, D.C. Yearbook 1952. xviii + 780 p. (cn ms).
- MAZUR, SŁAWOMIR. 1955. Umagi o występowaniu niektórych chrząszczy podkorowych w zerowiskach cetynic wieszego (*Tomicus piniperdis* L.) [Remarks on the occurrence of some subcortical beetles in the breeding sites of *Tomicus piniperdis*] [In Polish, English summary]. *Sylvan* 129(8): 35-42. (cc).
- MAZZONE, HORACE M., AND JOHN WILLIAM PEACOCK. 1955. Prospects for control of Dutch elm disease: biological considerations. *Journal of Arboriculture* 11: 285-292. (cn).
- MBONDJI, P. M. 1959. Etude epidemiologique d'*Hypothenemus hampei* (Coleoptera: Scolytidae), ravageur des baies du cafeeier, dans deux regions du Cameroun [Epidemiological study of *Hypothenemus hampei* a pest of coffee berries in 2 regions of Cameroun] [English subtitle]. *Le Naturaliste Canadien* 115: 245-249. (cc cn).
- MCCAIN, ARTHUR H., C. S. KOEHLER, AND S. A. TJOSVOLD. 1957. Pitch canker threatens California pines. *California Agriculture* 41(11-12): 22-23. (cc).
- MCCLURE, H. E. 1943. Further notes on aero-plankton of Kentucky. *Entomological News* 54: 37-45. (hb).
- MCDONALD, L., B. MANLY, J. LOCKWOOD, AND J. LOGAN. 1959. Estimation and analysis of insect populations. *Lecture Notes in Statistics* 55. 492 p. (cn hb).
- MCGREGOR, MARK D. 1955. Mountain pine beetle. The conflict between people and the beetle. Pages 16-23 in R. C. Loomis, S. Tucker, and T. H. Hofacker. *What else is growing in our forests? Insect and disease conditions in the United States 1979-1983*. United States Department of Agriculture, Forest Service, Forest Pest Management, Washington, D.C., General Technical Report WO-46. 93 p. (cn hb).
- _____. 1955. The place of partial cutting lodgepole pine in integrated management of mountain pine beetle. *International Congress of Entomology, Proceedings* 18: 414. (cn).
- MCGREGOR, MARK D., GENE DOYLE ANIMAN, RICHARD FREDERICK SCHMITZ, AND ROBERT D. OAKES. 1957. Partial cutting lodgepole pine stands to reduce losses to the mountain pine beetle. *Canadian Journal of Forest Research* 17: 1234-1239. (cn).
- MCGREGOR, MARK D., W. E. BOUSFIELD, R. L. JAMES, AND R. G. EDER. 1956. Biological evaluation of the Bridger-Derby, Deer Creek and Iron Mountain Management Units, Big Timber Ranger District, Gallatin National Forest, Montana. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 85-15. 13 p. (cn).
- MCGREGOR, MARK D., AND DENNIS M. COLE. Integrating management strategies for the mountain pine beetle with multiple-resource management of lodgepole pine forests. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, General Technical Report INT-174. 68 p. (cn cc).
- MCGREGOR, MARK D., KENNETH E. GIBSON, SCOTT TUNNOCK, LAWRENCE E. STIPE, H. E. MEYER, AND ROBERT D. OAKES. 1955. Status of mountain pine beetle infestations, Northern Region 1954. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Forest Pest Management Report 85-25. 57 p. (cn).
- MCGREGOR, MARK D., AND MITCHEL C. MILLER. 1959. Cross-attraction surveys for insect enemies of southern pine beetle. Pages 201-212 in D. L. Kullbavy and M. C. Miller. *Potential for biological control of *Dendroctonus* and *Ips* bark beetles*. Center for Applied Studies, School of Forestry, Stephen F.

- Austin State University, Nacogdoches, Texas. 255 p. (bv cc).
- MCGREGOR, MARK D., ROBERT D. OAKES, AND H. E. MEYER. 1955. Evaluation of Douglas-fir mortality from Douglas-fir beetle from 1952 through 1954 following 3 methyl-2-cyclohexan-1-one application. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana. Forest Pest Management Report 55-7, 19 p. (cn).
- MCGREGOR, MARK D., B. STEELE, P. SHEA, AND W. BOUSFIELD. 1959. Baiting and cutting to manage mountain pine beetle infestations. Pages 89-100 in G. D. Amman. Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- MCGUFFIN, W. C., AND ROBERT WILLIAM REID. 1951. Alberta and Rocky Mountain National Parks. Canada Department of Agriculture, Science Service, Division of Forest Biology, Forest Insect and Disease Survey, Annual Report 1950: 96-105 (cn).
- *MCLANE, LEONARD S. 1936. [*Hylesinus fraxini* (=varius)]. Journal of Economic Entomology 29: 768. ().
- *_____. 1937. Entomology in relation to industry. Journal of Economic Entomology 30: 6. ().
- MCLEAN, JOHN ALEXANDER. 1955. Ambrosia beetles: a multi-million dollar degrade problem of saw logs in coastal British Columbia. Forestry Chronicle 61: 295-298. (cn).
- _____. 1955. Ambrosia beetles in the forests of the Northwest. Northwest Environmental Journal 4: 324-327. (bv cn hb).
- MCLEAN, JOHN ALEXANDER, ALF BAKKE, AND HANS NIEMAYER. 1957. An evaluation of three traps and two lures for the ambrosia beetle *Trypodendron lineatum* (Oliv.) (Coleoptera: Scolytidae) in Canada, Norway and West Germany. Canadian Entomologist 119: 273-280. (cn).
- MCLEAN, JOHN ALEXANDER, AND EVELINE STORKINK. 1955a. Challenges in implementing ambrosia beetle pest management programs in British Columbia. Pages 179-187 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic Institute and State University, Blacksburg, Virginia. 355 p. (cn).
- _____. 1955b. Challenges in implementing ambrosia beetle pest management problems in British Columbia. International Congress of Entomology, Proceedings 15: 413. (cn).
- MCLEAN, JOHN ALEXANDER, AND J. TUYTEL. 1955. Marking forest insects: evaluation of two systems for the systematic introduction of rubidium into Douglas-fir trees. Canadian Journal of Forest Research 15: 19-23. (ms).
- McMULLEN, LESLIE HOWARD, AND LASZLO SAFRANYIK. 1955. Some effects of pine oil on mountain pine beetle (Coleoptera: Scolytidae) at different population levels. Entomological Society of British Columbia, Journal 52: 29-30. (bv cn).
- McMULLEN, LESLIE HOWARD, LASZLO SAFRANYIK, AND D. A. LINTON. 1956. Suppression of mountain pine beetle infestations in lodgepole pine forests. Government of Canada, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-276. 20 p. (cn).
- McMULLEN, LESLIE HOWARD, LASZLO SAFRANYIK, D. A. LINTON, AND R. BETTS. 1958. Survival of self-marked mountain pine beetles emerged from logs dusted with fluorescent powder. Entomological Society of British Columbia, Journal 55: 25-28. (hb ms).
- MEDUNA, J. 1956. Fannistic records from Czechoslovakia: Hymenoptera, Pteromalidae. Acta Entomologica Bohemoslovaca 53: 232. (cc).
- MEISNER, J., M. WEISSENBERG, D. BLUMBERG, AND K. R. S. ASCHER. 1955. Date palm fruit stone extracts as phagostimulants for the adult date stone beetle, *Coccotrypes dactyliperds* F. (Coleoptera: Scolytidae). Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 92: 305-309. (ay).
- MEIVILLE, R. V. 1979. *Xyleborus* Eichhoff, 1864 (Col., Scol.) conserved under the plenary powers, Opinion 1146. Bulletin of Zoological Nomenclature 36: 151-153. (tx).
- MENDEL, ZVI. 1955. Predation of *Orthotomicus erosus* (Col., Scolytidae) by the Syrian woodpecker *Picoides syriacus*, Aves, Picidae. Journal of Applied Entomology 100: 355-360. (cc).
- _____. 1956a. Comparison between two methods for the determination of the hymenopterous parasitoid complex of bark beetles (Scolytidae). Journal of Applied Entomology 101: 275-282. (cc ms).
- _____. 1956b. Discovery of the southern pine beetle, *Dendroctonus frontalis*, in Israel. Phytoparasitica 14: 319-320. (ds).
- _____. 1956c. Hymenopterous parasitoids of bark beetles (Scolytidae) in Israel: host relation, host plant, abundance and seasonal history. Entomophaga 31: 113-125. (cc).
- _____. 1956d. Hymenopterous parasitoids of bark beetles (Scolytidae) in Israel: relationships between host and parasitoid size, and sex ratio. Entomophaga 31: 127-137. (cc).
- _____. 1957. Major pests of man-made forests in Israel: origin, biology, damage and control. Phytoparasitica 15: 131-137. (cn hb).
- _____. 1958a. Attraction of *Orthotomicus erosus* and *Pityogenes calcaratus* to a synthetic aggregation pheromone of *Ips typographus*. Phytoparasitica 16: 109-117. (bv).
- _____. 1958b. Effects of food, temperature, and breeding conditions on the life span of adults of three cohabiting bark beetles (Scolytidae) parasitoids (Hymenoptera). Environmental Entomology 17: 293-298. (cc).
- _____. 1958c. The relation of bast scale and bark beetle outbreaks to management of pine plantations in Israel. Pages 329-336 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn cc).
- _____. 1958d. The relation of bast scale and bark beetle outbreaks to management of pine plantations in Israel. International Congress of Entomology, Proceedings 15: 416. (cn cc).
- MENDEL, ZVI, Y. GOLAN, Z. MADAR, AND Z. SOLET. 1953. Insect pests and diseases of cypress in Israel [In Hebrew, English summary]. La-Yaaran 33: 37-41. (cn).
- MENDEL, ZVI, AND E. GUREVITZ. 1955. Hymenopterous parasitoids of the almond bark beetles in Israel. Alon Hanotea 39: 1057-1060. (cc).
- MENDEL, ZVI, Z. MADAR, AND Y. GOLAN. 1953. Chemical control of pine bark beetles (preliminary trials). La-Yaaran 33: 41-43. (cn).
- _____. 1955. Comparison of the seasonal occurrence and

- behavior of seven pine bark beetles (Coleoptera: Scolytidae) in Israel. *Phytoparasitica* 13: 21–32. (bv hb).
- MENDEL, ZVI, H. PODDLER, AND H. LIVNE. 1989. Establishment sequence and seasonal development of *Anthonomus ruficornis* Olivier (Coleoptera: Curculionidae), a predator of bark beetles in pine plantations in Israel. *Acta Oecologica, Oecologia Applicata* 10: 103–114. (ec).
- MERCER, CHRISTOPHER W. L. 1952. Studies on the forest insects of Sarawak. A terminal report, Vols. I and II. Forest Department of Sarawak, Kuching, Sarawak. vii + 290 p. (cn ds).
- MERCER, CHRISTOPHER W. L., D. LEWIS, AND HWEL ROBERTS. 1986. A trial in Labu TRP [timber rights purchase] to test the effectiveness of three insecticides against ambrosia beetle attack in logs. *Klinkii* 3: 25–39. (cn).
- *MERCET. 1926. [*Pityogenes quadridens*]. *Revista de Fitopatologia, Madrid* 1926: 40–47. ().
- MERLIN, J., JEAN CLAUDE GREGOIRE, M. BAISIER, AND J. M. PASTEELS. 1955. Some new data on the biology of *Rhizophagus grandis* (Coleoptera: Rhizophagidae). Pages 107–121 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Universite Libre de Bruxelles, Belgium, Proceedings of seminar, 3–4 October 1954. 141 p. (ec).
- MERLIN, J., C. PARMENTIER, AND JEAN CLAUDE GREGOIRE. 1956. The feeding habits of *Rhizophagus dispar* (Col., Rhizophagidae), an associate of bark beetles. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* 51: 915–923. (ec).
- MERRILL, LAURA D., DAVID LEE WOOD, AND JAMES H. CANE. 1985. Hybridization barriers in Group IX *Ips*: *I. hoppingi* Lanier and *I. confusus* (LeConte) (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 18: 430. (hb).
- MICHAELS, P. J., D. E. SAPPINGTON, AND P. J. STENGER. 1985. Objective prediction of climate-related changes in the distribution of southern pine beetle. Pages 41–52 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (ec).
- _____. 1986. SPBCMP, a program to assess the likelihood of major changes in the distribution of the southern pine beetle. *Southern Journal of Applied Forestry* 10: 158–161. (ec ms).
- MICHAELS, P. J., P. J. STENGER, AND D. E. SAPPINGTON. 1986. Modeling changes in the epidemic range of southern pine beetle using temperature and objective moisture indicators. *Theor. Appl. Clim.* 37: 39–50. (ec ms).
- MICHALSKI JACEK. 1985. On the occurrence and significance of nematodes (Nematoda) in the galleries of *Tomicus piniperda* L. (Coleoptera, Scolytidae). *Bull. Soc. Amis Sci. Lett. Poznan, ser. B, Sci. Biol.* 26: 129–137. (ec).
- MICHALSKI JACEK, AND E. RATAJCZAK. 1989. *Korniki* (Coleoptera: Scolytidae) wraz z towarzyszącą im fauną w Gorach Świętokrzyskich [Bark beetles along with their accompanying fauna in the Świętokrzyskie Mountains, Poland] [In Polish, English summary]. *Fragmenta Faunistica* 32: 279–315. (ec).
- *MIKKOLA, M., G. D. PIERCE, K. LÖYTTYNIEMI, AND A. C. MUBITA. 1979. Late stock-scion incompatibility in *Pinus merkusii* de Vries in Zambia. *Commonwealth Forestry Review* 55: 207–209. ().
- MILES, BRUCE R. 1987. Tragedy of the Four Notch. *American Forests* 93(3–4): 26–29, 76. (cn).
- MILLAR, J. G., C. H. ZHAO, GERALD NORMAN LANIER, D. P. O'CALLAHAN, M. GRIGGS, J. R. WEST, AND ROBERT MILTON SILVERSTEIN. 1986. Components of moribund American elm trees as attractants to elm bark beetles, *Hylurgopinus rufipes* and *Scolytus multistriatus*. *Journal of Chemical Ecology* 12: 583–605. (bv).
- *MILLER, D., AND A. F. CLARK. 1935b. [*Hylastes ater*]. *Bulletin of Entomological Research* 26: 149–154. ().
- MILLER, DANIEL R., AND JOHN HARVEY BORDEN. 1955. Life history and biology of *Ips latidens* (LeConte) (Coleoptera: Scolytidae). *Canadian Entomologist* 117: 559–571. (cn ec hb).
- _____. 1990. Beta-phellandrene: kairomone for pine engraver, *Ips pini* (Say) (Coleoptera: Scolytidae). *Journal of Chemical Ecology* 16: 2519–2531. (bv).
- MILLER, DANIEL R., JOHN HARVEY BORDEN, AND K. N. SLESSOR. 1989. Inter- and intrapopulation variation of the pheromone, ipsdienol produced by male pine engravers, *Ips pini* (Say) (Coleoptera: Scolytidae). *Journal of Chemical Ecology* 15: 233–247. (bv).
- MILLER, DANIEL R., J. L. MADDEN, AND JOHN HARVEY BORDEN. 1986. Primary attraction of *Ips latidens* (LeConte) and *Hylastes gracilis* LeConte (Coleoptera: Scolytidae) to high-girdled lodgepole pine, *Pinus contorta* var. *latifolia* Engelm. *Canadian Entomologist* 118: 85–88. (bv).
- MILLER, L. KEITH, AND RICHARD ALLEN WERNER. 1987. Cold-hardiness of adult and larval spruce beetles *Dendroctonus rufipennis* (Kirby) in interior Alaska. *Canadian Journal of Zoology* 65: 2927–2930. (hb).
- MILLER, MITCHELL C. 1985. The effect of *Monochamus titillator* (F.) (Col., Cerambycidae) foraging on the emergence of *Ips calligraphus* (Germ.) (Col., Scolytidae) insect associates. *Journal of Applied Entomology* 100: 189–197. (ec).
- _____. 1986a. Survival of within-tree *Ips calligraphus* (Col.: Scolytidae): effect of insect associates. *Entomophaga* 31: 39–45. (ec).
- _____. 1986b. Within-tree effects of bark beetle insect associates on the emergence of *Ips calligraphus* (Coleoptera: Scolytidae). *Environmental Entomology* 15: 1104–1108. (ec).
- MILLER, MITCHELL C., JOHN CONRAD MOSER, AND DONALD N. KINN. 1985. More information on bark beetle alternatives. *IPM Practitioner* 7(11–12): 14. (ec).
- MILLER, MITCHELL C., JOHN CONRAD MOSER, MARK D. MCGREGOR, JEAN CLAUDE GREGOIRE, M. BAISIER, DONALD L. DAHLSTEN, AND RICHARD ALLEN WERNER. 1987. Potential for biological control of native North American *Dendroctonus* beetles (Coleoptera: Scolytidae). *Entomological Society of America, Annals* 80: 417–428. (cn ec).
- MILLER, MITCHELL C., ET AL. 1989. Responses of insect associates of allied species to *Dendroctonus* and *Ips* bark beetles. In D. L. Kulhavy and M. C. Miller, Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (bv ec).
- MILLER, NORMAN CECIL EGERTON. 1941. Insects associated with cocoa *Theobromus cacao* in Malaya. *Bulletin of Entomological Research* 32: 1–15. (cn).
- MILLER, RANDALL F., W. D. FITZGERALD, AND D. N. BUIHAY

1957. Fossil Coleoptera from the postglacial spruce-pine transition period near Minesing Swamp, Ontario, Canada. *Canadian Journal of Earth Sciences* 24: 2099–2103. (ds).
- MILLER, RANDALL F., ALAN V. MORGAN, AND S. R. HICOCK. 1955. Pre-vashon fossil Coleoptera of Fraser age from the Fraser Lowland British Columbia. *Canadian Journal of Earth Sciences* 22: 498–505. (ds).
- MILLER, RANDALL H., AND ALAN ANDREW BERRYMAN. 1956. Carbohydrate allocation and mountain pine beetle attack in girdled lodgepole pines. *Canadian Journal of Forest Research* 16: 1036–1040. (bv).
- MILLER, RANDALL H., ALAN ANDREW BERRYMAN, AND C. A. RYAN. 1956. Biotic elicitors of defense reactions in lodgepole pine. *Phytochemistry* 25: 611–612. (cn).
- MILLER, RANDALL H., H. STUART WHITNEY, AND ALAN ANDREW BERRYMAN. 1956. Effects of induced translocation stress and bark beetle attack *Dendroctonus ponderosae* on heat pulse velocity and the dynamic wound response of lodgepole pine *Pinus contorta* var. *latifolia*. *Canadian Journal of Botany* 64: 2669–2674. (bv cn).
- MILLER-WEEKS, MARGARET. 1955. Forest pest conditions report for the Northeastern Area, 1954. United States Department of Agriculture, Forest Service, Northeastern Area (Broomall, Pennsylvania). Forest Pest Management Report NA-FR-32. 22 p. (cn).
- MILLIGAN, ROBERT H., G. O. OSBORNE, AND G. YTMA. 1958. Evidence for an aggregation pheromone in *Platypus gracilis* Broun (Col., Platypodidae). *Journal of Applied Entomology* 106: 20–24. (bv).
- MILLIGAN, ROBERT H., AND G. YTMA. 1958. Pheromone dissemination by male *Platypus apicalis* White and *P. gracilis* Broun (Col., Platypodidae). *Journal of Applied Entomology* 106: 113–115. (bv).
- MILLS, J. R. 1955. Host location: an important consideration in the use of exotic natural enemies for the biocontrol of native scolytids. *International Congress of Entomology, Proceedings* 18: 416. (cn ec).
- MILLS, NICHOLAS J. 1955. Some observation on the role of predation in the natural regulation of *Ips typographus* populations. *Journal of Applied Entomology* 99: 209–215. (ec).
- _____. 1956. A preliminary analysis of the dynamics of within tree populations of *Ips typographus* (L.) (Coleoptera: Scolytidae). *Journal of Applied Entomology* 102: 402–416. (ec lb).
- MILLS, NICHOLAS J., AND K. KRUGER. 1958. Host location: an important factor in the use of exotic natural enemies for the biocontrol of native scolytids. Pages 321–325 in T. L. Payne and H. Saaremaa. *Integrated control of scolytid bark beetles*. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn ec).
- MILLS, NICHOLAS J., AND J. SCHLUP. 1959. The natural enemies of *Ips typographus* in central Europe: impact and potential use in biological control. Pages 131–146 in D. L. Kuhlavy and M. C. Miller. *Center for Applied Studies, School of Forestry*, Stephen F. Austin State University, Nacogdoches, Texas. 255. (cn ec).
- *MIRZOIAN, S. A. 1977. Dendrofilnye nasekomye lesov i parkov Armenii. Nauchno-issledovatel'skij institut zascity rastenij MSCH Armjanskij SSR, Izdatel'stvo Ajastan, Jerevan 1977: 27–43, 166–181, 346–357 (etc.?). ().
- MITCHELL, RUSSEL G. 1959. Mixed host strategies for mountain pine beetle control in Oregon. Pages 60–63 in G. D. Amman. *Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah. General Technical Report INT-262. 119 p. (cn).
- MITCHELL, RUSSEL G., R. E. MARTIN, AND J. STUART. 1953. Catfaces on lodgepole pine: fire scars or strip kills by the mountain pine beetle? *Journal of Forestry* 51: 598–601. (cn).
- MIZELL, RUSSELL F., T. EVAN NEBEKER, AND JAMES L. FRAZIER. 1981. A new electro-optical activity monitor for determining locomotor activity of *Thanasimus dubius* (F.) in the laboratory. *Georgia Entomological Society, Journal* 16: 479–484. (ec).
- MOCETTINI, PHILIP J., JR. AND D. B. CAHILL. 1959. Evaluation of spruce beetle population trends on the Payette National Forest, Idaho. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report 89–2. 12 p. (cn).
- *MOECK, HENRY ADOLF. 1967. Electron microscopic studies of antennal sensilla in the ambrosia beetle *Trypodendron lineatum* (Olivier) (Scolytidae). Unpublished thesis, University of British Columbia. 91 p. (av).
- *_____. 1975. Host selection behavior of bark beetles (Coleoptera: Scolytidae) attacking *Pinus ponderosa*, with special emphasis on the western pine beetle, *Dendroctonus brevicornis* LeC. Unpublished dissertation, University of California, Berkeley. 62 p. ().
- _____. 1955. A bucket emergence trap for corticolous insects. *Canadian Entomologist* 120: 855–865. (cn ms).
- MOECK, HENRY ADOLF, LASZLO SAFRANYIK, CLARENCE S. SIMMONS, AND CAROL M. LAWKO. 1955. *Ips tridens* (Coleoptera: Scolytidae) attracted by ipsdienol plus cis-verbenol. *Canadian Entomologist* 117: 955–960. (bv).
- MOHR, P., AND C. TAMM. 1957. Synthesis of endo-1,3 dimethyl-2,9-dioxabicyclo-3,3,1-nonane. *Tetrahedron Letters* 28: 395–396. (bv ms).
- MOLLER, W. J., AND J. E. DEVAY. 1968. Insect transmission of *Ceratocystis fimbriata* in deciduous fruit orchards. *Phytopathology* 58: 1499–1505. (ec).
- *MONCAYO, E. R. 1958. Plagas del cacao en los departamentos de Santander y Antioquia, Colombia. Pages 261–269 in Septima Conferencia Interamericana de Cacao, Palmeira, Colombia. ().
- *MONTGOMERY, B. E., AND J. M. AMOS. 1941. Contributions to a list of the Coleoptera of the Clark County State Forest. *Indiana Academy of Science, Proceedings* 50: 251–255. ().
- *MONTROUZIER, P. 1861. Essai sur la faune entomologique de la Nouvelle Calédonie et des Iles des Pin. Art. Lifu, etc. *Société Entomologique de France, Annales*, ser. 3, 8: 229–308, ser. 4, 1: 265–306. ().
- MOODY, B. H. 1955. Forest insect and disease conditions in Canada 1957. Government of Canada, Forestry Service, Forest Insect and Disease Survey, Ottawa, Ontario. 92 p. (cn).
- MOODY, B. H., AND H. F. CEREZKE. 1955. Forest insect and disease conditions in Alberta, Saskatchewan, Manitoba and the Northwest Territories in 1954 and predictions for 1955. Government of Canada, Canadian Forestry Service, Northern Forestry Centre (Edmonton, Alberta), Information Report NOR-X-269. 21 p. (cn).
- MOODY, B. H., P. SINGH, AND L. J. CLARKE. 1952. Newfoundland

- land Region. Pages 7–16 in Annual report of the Forest Insect and Disease Survey, 1978. Canada Department of Environment, Canadian Forestry Service, Ottawa, Ontario. (cn).
- MORAL, L. G. 1957. Aantastingen door insecten en mijten op bomen en struiken in 1956. Nederlands Bosbouw tijdschrift 59: 373–381. (cn).
- _____. 1958. Aantastingen van insecten en mijten in bossen en natuurgebieden in 1957. Nederlands Bosbouw tijdschrift 60: 352–360. (cn).
- MORGAN, ALAN V. 1958a. *Polygraphus convexifrons* Wood (Coleoptera: Scolytidae): a range extension to Ungava Bay, Quebec, Canada. Coleopterists Bulletin 42: 69–72. (ds).
- _____. 1958b. Zoogeography and migration of scolytid species during the last 125,000 years in North America. International Congress of Entomology, Proceedings 18: 411. (ds).
- *MORGAN, ANNE 1973. Late Pleistocene environmental changes indicated by fossil insect faunas of the English midlands. Boreas 2: 173–212. ().
- MORGAN, ANNE, ALAN V. MORGAN, AND S. A. ELLIS. 1955. Holocene insects and paleoecology of the Au Sable River, Michigan. Ecology 66: 1817–1828. (ds).
- MORGAN, F. D. 1959. Forty years of *Sirex noctilio* and *Ips grandicollis* in Australia. New Zealand Journal of Forest Science 19: 198–209. (cn).
- MORI, KENJI AND Y. IGARASHI 1955. Pheromone synthesis CIV. Synthesis of the enantiomers of alpha phellandren-5-ol (p mentha-1 5-dien-5-ol): a monoterpene from bark beetles. Liebigs Annalen der Chemie (1): 93–96. (bv ms).
- MORI, KENJI, M. KUHARA, AND M. SUZUKI 1959. Stereoselective synthesis of (1R,3R,5S)-1,3-Dimethyl-2,9-dioxabicyclo[3.3.1]nonane. Chem. Pharm. Bull. [Tokyo] 37: 1078–1079. (bv ms).
- MORI, KENJI, AND J. J. OGOCIE 1955. An enzyme-mediated synthesis of both enantiomers of seudenol and 1-methyl-2-cyclohexen-1-ol, the aggregation pheromones of *Dendroctonus pseudotsugae*. Liebigs Annalen der Chemie 9: 903–905. (bv ms).
- MORI, KENJI, AND P. PUAPOOMCHARON. 1957. Conversion of the enantiomers of sulcatol 6-methyl-5-hepten-2-ol to the enantiomers of pityol trans-2-1 hydroxy-1-methylethyl-5-methyltetrahydrofuran, a male-specific attractant of the bark beetle *Pityophthorus pityographus*. Liebigs Annalen der Chemie 3: 271–272. (bv ms).
- _____. 1955. Pheromone synthesis CVI. Synthesis of all of the four stereoisomers of tetrahydro-2 2 6-trimethyl-2h-pyran-3-ol: a volatile compound from the elm bark beetle *Pteleobius vittatus*. Liebigs Annalen der Chemie (2): 175–177. (bv ms).
- _____. 1959. Pheromone synthesis CXVII. Conversion of the enantiomers of sulcatol to the enantiomers of CIS pityol, a volatile compound from the elm bark beetle *Pteleobius vittatus*. Liebigs Annalen der Chemie 12: 1261–1262. (bv ms).
- MORI, KENJI AND Y. B. SEU 1955. A new synthesis of levo-alpha multistriatin, the pheromone of the smaller European elm bark beetle [*Scolytus multistriatus*]. Tetrahedron 44: 1035–1035. (bv ms).
- MORIMOTO, KATSURA 1955. Some observations on the classification of the subfamilies of Curculionidae. International Congress of Entomology, Proceedings 18: 40. (ts).
- MORITZ, B., AND E. FUHRER 1955. Rindeninhaltsstoffe der Fichte (*Picea abies*): Analysen an Borkenkäfer-fangbaumen [Phloem contents of *Picea abies*: analyses on bark beetle trap trees] [English summary]. Journal of Applied Entomology 105: 502–509. (ec).
- MORON, M. A., J. F. CAMAL, AND O. CANUL 1956. Analisis de la entomofauna necrofila de la zona norte de la reserva "Sian Ka'an," Quintana Roo, Mexico [Analysis of the necrophilous entomofauna of the northern area of the biosphere reserve "Sian Ka'an," Quintana Roo, Mexico] [English summary]. Folia Entomologica Mexicana 69: 83–95. (cn ms).
- MORRISON, DUNCAN J., AND RICHARD S. HUNT 1955. Black-stain root diseases on lodgepole pine in British Columbia, Canada. Phytopathology 75: 1295. (ec).
- MOSER, JOHN CONRAD 1975. Biosystematics of the straw itch mite with special reference to nomenclature and dermatology. Royal Entomological Society of London, Transactions 127(2): 185–191. (ec).
- _____. 1985. Use of sporothecae by phoretic *Tarsonemus* mites to transport ascospores of coniferous bluestain fungi. Br. [British?] Mycological Society, Transactions 84: 750–753. (cn ec).
- _____. 1959a. An exotic predator for the biological control of the black turpentine beetle. Entomological Society of America, Symposium. (ec).
- _____. 1959b. Inoculative release of an exotic predator for the biological control of the black turpentine beetle. Pages 189–200 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ec).
- MOSER, JOHN CONRAD, AND SUSAN J. BRANHAM 1985. Bugs that eat bugs: biological control research offers hope for southern pine bark beetle management. Forest Farmer 47(3): 17–20. (cn).
- MOSER, JOHN CONRAD, AND JOHN ROBERT BRIDGES 1956. *Tarsonemus* (Acarina: Tarsonemidae) mites phoretic on the southern pine beetle (Coleoptera: Scolytidae): attachment sites and numbers of bluestain (Ascomycetes: Ophiostomataceae) ascospores carried. Entomological Society of Washington, Proceedings 55: 297–299. (ec).
- MOSER, JOHN CONRAD, HUBERTUS H. EIDMANN, AND J. R. REGNANDER 1959. The mites associated with *Ips typographus* in Sweden. Annales Entomologici Fennici 55: 23–27. (ec).
- MOSER, JOHN CONRAD, THELMA J. PERRY, AND HALVOR SOLHEIM 1959. Ascospores hypophoretic on mites associated with *Ips typographus*. Mycological Research 93: 513–517. (ec).
- MOSER, JOHN CONRAD, ROBERT A. SOMMERS, PETER L. LORIO, JR., JOHN ROBERT BRIDGES, AND JEFFREY J. WITCOSKY 1957. Southern pine beetles attack felled green timber. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. Research Note SO-342. 7 p. (bv).
- MOSER, JOHN CONRAD, AND WILLIAM A. THOMPSON 1956. Temperature thresholds related to flight of *Dendroctonus frontalis* Zimm. (Col., Scolytidae). Agronomie 6: 905–910. (ec).
- MUESEBECK, CARL, FREDERICK WILLIAM, KARL VON VORSE KROMBEIN, AND H. K. THOMAS 1951. Hymenoptera of America north of Mexico, synoptic catalog, United States Department of Agriculture, Monograph 2. 1420 p. (ec).
- MOULSON, D. C. 1955. The development of a climbing tree delimeter/debarker. Forest Engineering Research

- Institute of Canada, Technical Note TN-51. 17 p. (cn ms).
- MOUSSION, G. 1955. Que faire contre le depérissement des thuyas cause par les insectes xylophages? [What can be done against thuya wilt caused by wood-eating insects?]. *Phytoma* 402: 61-64. (cn).
- MOZOLEVSKAYA, EKATERINA G. 1955. The density of a population of bark beetles and its informative significance [In Russian, English summary]. *Zoologicheskii Zhurnal* 64: 1334-1341. (cn hb).
- MOZOLEVSKAYA, EKATERINA G., N. V. KRYLOVA, N. K. BELOVA, AND I. N. OSIPOV. 1957. The ecology of bark beetle transmitters of Dutch elm disease. *Zashchita Rastenii Moscow* 7: 37-40. (cc hb).
- MURACEK, ZDENEK. 1955. Elektrodynamická aplikace chemických přípravků vochrane lesu [Electrodynamic application of chemical preparations in forest protection] [In Czech, English, Russian summaries]. *Lesnictví [Prague]* 31(7): 629-640. (cn ms).
- MUKAI, YUKIO, T. SUGIMURA, H. WATANABE, AND K. WAKAMORI. 1957. Extraction of areas infested by pine bark beetles using Landsat MSS data. *Photogrammetric Engineering and Remote Sensing* 53: 77-81. (cn).
- *MÜLLER-THURGAU, H., A. OSTERWALDER, AND OTTO SCHNEIDER-ORELLI. 1917. [*Scolytus rugulosus*]. *Landwirtschafth. Jahrbuch der Schweiz* 1917: 416-426. ().
- MULOCK, PETER, AND ERIK CHRISTIANSEN. 1986. The threshold of successful attack by *Ips typographus* on *Picea abies*: a field experiment. *Forest Ecology and Management* 14: 125-132. (bv cn).
- MULSANT, MARTIEL ETIENNE, AND CLAUDIUS REY. 1852. Description d'une espèce nouvelle de Coleoptère du genre *Bostrichus*. *Academie des sciences, belles-lettres et arts de Lyon, Classe de sciences, serie 2*, 2: 205-206. (tx).
- MUNOZ, HERNANDEZ, R. 1955. Muestreo en fincas para determinar la poblacion de broca (*Hypothenemus hampei* Ferr.) y metodologia para calcular el nivel de dano economico [Field sampling to determine populations of *Hypothenemus hampei* and a method for calculating economic damage levels] [English summary]. *Boletin de Promecafe* 35: 4-14. (cn ms).
- MURALEEDHARAN, N. 1953. Entomology. Annual Report, Tea Scientific Department, United Planter's Association of Southern India 56: 86-106. (cn hb).
- _____. 1954. Entomology. Annual Report, Tea Scientific Department, United Planter's Association of Southern India 57: 72-91. (cn hb).
- _____. 1956. The shot-hole borer of tea: its biology, ecology and control. United Planter's Association of Southern India. Handbook of Tea Culture, series 19. 4 p., 3 figs. (cn hb).
- _____. 1957. Entomological research in tea in southern India. *Journal of Coffee Research* 17: 80-83. (cn).
- MURALEEDHARAN, N., AND B. RADHAKRISHNAN. 1959. Recent studies on tea pest management in south India. United Planter's Association of Southern India. Bulletin 43: 16-29. (cn).
- MURALEEDHARAN, N., R. SELVASUNDARAM, AND B. RADHAKRISHNAN. 1955. Natural enemies of certain tea pests occurring in southern India. *Insect Science and its Application* 9: 647-654. (cc).
- *MURAYAMA, JOZO J. 1947. [*Hyurgops palliatus*]. *Kontyu* 16(1): 27. ().
- MURTHA, P. A., AND R. COZENS. 1955. Color infra-red photo interpretation and ground surveys to evaluate spruce beetle attack. *Canadian Journal of Remote Sensing* 11: 177-187. (cn).
- MURTHA, P. A., AND R. J. WIART. 1957. PC-based digital image analysis for mountain pine beetle green attack: preliminary results. *Canadian Journal of Remote Sensing* 13: 92-95. (cn).
- _____. 1959. Cluster analysis of pine crown foliage patterns aid identification of mountain pine beetle current-attack. *Photogrammetric Engineering and Remote Sensing* 55: 83-86. (cn).
- MUSSCHIE, G., B. GARBOUS, V. BAUTERS, AND A. PANIS. 1957. Etude preliminaire du peuplement d'Hymenopteres de l'olivier dans le nord-ouest de la Tunisie [Preliminary study of the population of Hymenoptera on olive trees in the north-west of Tunisia] [English summary]. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* 52: 303-309. (cn).
- MUSTAPARTA, HANNA, AND T. J. ALMAAS. 1959. How does the olfactory system in insects discriminate between odors of own and related species? *Fauna Norvegica* 36: 81-86. (bv).
- MUSTAPARTA, HANNA, B. A. TOMMERAS, AND GERALD NORMAN LANIER. 1955. Pheromone receptor cell specificity in interpopulational hybrids of *Ips pini* (Coleoptera: Scolytidae). *Journal of Chemical Ecology* 11: 999-1000. (ay bv).

N

- *NAKANE, TAKEHIRO. 1963. A list of Coleoptera from the Shiretoko Peninsula, Hokkaido, Japan (Insecta). *Sci. Rep. Kyoto Pref. Univ. (Nat. Sci. & Liv. Sci.)*, ser. A, 3(5): 237-245. ().
- NAKASHIMA, TOSHIHO. 1950. Sexual division of labor in reproductive behavior, especially the transmission of mutualistic fungi by ambrosia beetles [abstract]. *International Congress of Entomology, Proceedings* 16(9S-1.6): 285. (bv).
- _____. 1959. Observations on the ambrosia fungus, *Ambrosiella* sp., growing in the gallery of *Scolytoplatypus shogun* (Coleoptera: Scolytidae) and on the concurrent damage of wood tissue. *Faculty of Agriculture, Hokkaido University, Journal* 64: 99-105. (cc).
- NAKASHIMA, TOSHIO, CHIE GOTO, AND TOSHIHIKO IIZUKA. 1957. The primary and auxiliary ambrosia fungi isolated from the ambrosia beetles, *Scolytoplatypus shogun* Blandford (Coleoptera: Scolytidae) and *Crossotarsus niponicus* Blandford (Coleoptera: Platypodidae). *Faculty of Agriculture, Hokkaido University, Journal* 63: 185-208. (cc).
- NAKASHIMA, TOSHIO, T. OTOMO, Y. KATO, AND TOSHIHIKO IIZUKA. 1958. Scanning electron microscope observations on the ambrosia fungi growing in the galleries where the larvae of ambrosia beetles are growing. *International Congress of Entomology, Proceedings* 18: 410. (cc).
- NAKATA, T., S. NAKAO, S. TAKAO, T. TANAKA, AND T. OISHI. 1955. Stereoselective synthesis of 1 R 3 R 5 S-1,3 dimethyl-2,9-dioxabicyclo-3.3.1-nonane. *Tetrahedron Letters* 26: 73-74. (ms).
- NAKAYAMA, K., AND P. S. TERRA. 1956. Atratividade de substancias e de ramos de cacauero sobre *Xylosandrus morigerus* (Blandford, 1894) (Coleoptera, Scolytidae) [Attraction of *Xylosandrus morigerus* to different substances and branches of cacao plants]. *Revista Theobroma* 16: 155-160. (bv).

- NASH, DAVID R. 1959. Notes on British *Orthotomicus* (Col.: Scolytidae) including *suturalis* Gyllenhal new to Wiltshire. *Entomologist's Record and Journal of Variation* 10: 151-152. (ds).
- *NAVARRO DE ANDRADE, EDMUNDO 1917. [*Hylesinus oleiperda* (=toranio)]. *Boletino de Agricultura Tecnica y Economica*, Madrid 7(78): 554-563. ().
- NEBEKER, T. EVAN 1985. Influence of forestry practices on bark beetle populations: a perspective. Pages 52-59 in *Thirty-fourth annual Forestry Symposium*, Division of Continuing Education, Louisiana State University, Baton Rouge. (hb).
- NEBEKER, T. EVAN, AND ROY L. HEDDEN 1984. Integrated forest pest management in pine stands (6+ yrs.): insects. Pages 116-125 in S. J. Branham and G. D. Hertel, *Symposium proceedings: Integrated pest management*. Athens, Georgia. (cn).
- NEBEKER, T. EVAN, AND JOHN DEAVOURS HODGES 1985. Thinning and harvesting practices to minimize site and stand disturbance and susceptibility to bark beetle and disease attacks. Pages 263-271 in S. J. Branham and R. C. Thatcher, *Symposium proceedings: Integrated pest management research*. United States Department of Agriculture, Forest Service, General Technical Report SO-56. (cn).
- NEBEKER, T. EVAN, JOHN DEAVOURS HODGES, ROBERT K. KARR, AND DAVID M. MOEHRRING 1985. Thinning practices in southern pines with pest management recommendations. United States Department of Agriculture, Forest Service, Technical Bulletin 1703: 3-36. (cn).
- NEBEKER, T. EVAN, JOHN DEAVOURS HODGES, C. R. HONEA, AND C. A. BLANCHIE 1988. Preformed defensive system in loblolly pine: variability and impact on management practices. Pages 147-162 in T. L. Payne and H. Saarenmaa, *Integrated control of scolytid bark beetles*. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv ec).
- NEBEKER, T. EVAN, D. R. HOUSTON, AND JOHN DEAVOURS HODGES 1986. Forest pests: influence of forest management practices on pest population dynamics and forest productivity. Pages 101-112 in T. C. Hennessey, P. M. Dougherty, S. V. Kossuth, and J. D. Johnson, *Stress physiology and forest productivity*. Martinus-Nijhoff Publishers. (cn).
- NEBEKER, T. EVAN, RUSSELL F. MIZELL III, AND NORMAN J. BEDWELL 1984. Management of bark beetle populations. Impact of manipulating predator cues and other control tactics. Pages 23-33 in W. Y. Garner and J. Harvey, *Chemical and biological control agents in forestry*. American Chemical Society Symposium Series 238, Seattle, Washington, 20-25 March 1983. (cn).
- NEBEKER, T. EVAN, G. C. PURSER, AND RUSSELL F. MIZELL III 1980. Collection and maintenance of *Thanosinus dubius* (F.) for biological and behavioral studies. *Georgia Entomological Society, Journal* 15: 406-412. (ec).
- NECHILEBA, ALOIS 1928c. Response to K. E. Schedl comments on: Verkummern und Verderben von Brutten forstschadlicher Insekten. *Anzeiger fur Schadlingsskunde* 4(8): 111.
- NEE, LUDOVIC 1989. Quantification de l'intensite des attaques d'*Ips typographus* L. sur *Picea abies* Karst. [Quantification of the intensity of the attacks of *Ips typographus* on the Norway spruce]. *Journal of Applied Entomology* 107: 296-303. (cn).
- NEE, LUDOVIC, AND R. JANSSENS 1986. Influence de divers types de pieges et de facteurs environnementaux sur les captures d'ipides forestiers [The influence of different types of traps and environmental factors on the captures of forest ips]. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* 51(3a): 907-913. (cn ec).
- NEGRU, STEPHAN 1970. Noi plante-gazda ale Scolitidelor (Col., Scolytidae) din Romania [New plant hosts for Romanian Scolytidae]. *Comm. Zool.* 1970: 151-160. (ds).
- * ———. 1971. Scolitidi (Coleoptera, Scolytidae) della del futura bacino d'accumulazione 'Portile de Filer.' *Travaux de Museum d'Histoire Naturelle Grigore Antipa, Bucharest* 11: 175-189. ().
- NEIVA, ARTHUR A., AND ROSARIO AVERNA-SACCA 1927. Os entomofagos cryptogamicos na broca do cafeeiro (*Stephanoderes hampei* Ferr.) encontrados em S. Paulo. [Source journal not copied by loaning library] 1927[or later]: 10-24, 195-213. (cn ec).
- *NELSON 1931. [*Hylastes ater*]. Department of Scientific and Industrial Research, New Zealand. *Annual Report* 1930: 1. ().
- NETTLETON, W. A., M. D. CONNOR, AND G. W. RYAN 1985. Evaluation and application of the TAMBEETLE and Arkansas southern pine beetle spot growth models in the Gulf Coastal Plain. Pages 202-205 in S. J. Branham and R. C. Thatcher, *Symposium proceedings: Integrated pest management research*. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans. Louisiana, General Technical Report SO-56. 353 p. (cn).
- NEUMANN, FREDERICK G. 1987. Introduced bark beetles on exotic trees in Australia with special reference to infestations of *Ips grandicollis* in pine plantations. *Australian Forestry* 50: 166-178. (cn ec hb).
- NEUMANN, FREDERICK G., AND GEORGE MINKO 1985a. Aspects of the biology, survey and control of the introduced smaller European elm bark beetle, *Scolytus multistriatus*, a potential carrier of Dutch elm disease in Victoria. [Australia] Department of Conservation, Forests and Lands, State Forests and Lands Service, Research Branch Report 250. 21 p. (hb ds).
- . 1985b. Studies on the introduced smaller European elm bark beetle, *Scolytus multistriatus*, a potential vector of Dutch elm disease in Victoria. *Australian Forestry* 48: 116-126. (hb).
- NEUMANN, FREDERICK G., AND J. MOREY 1984. Studies on the introduced bark beetle *Ips grandicollis* (Eichhoff) in Victorian Radiata pine populations. *Australian Forest Research* 14: 283-300. (hb).
- NEVES, N., F. MONIZ, N. D. AZEVEDO, M. C. FERREIRA, AND G. W. S. FERREIRA 1986. Present phytosanitary situations of Portuguese forests. European and Mediterranean Plant Protection Organization, *Bulletin* 16: 505-508. (cn).
- NEWELL, R. 1977. Forest insect and disease highlights. Canada Department of Fisheries and Environment, Canadian Forestry Service, Maritimes Forestry Research Centre, Fredricton, New Brunswick, *Pith to Periderm* 11(2): 11. (cn).
- . 1978. Forest insect and disease highlights. Canada Department of Fisheries and Environment, Canadian Forestry Service, Maritimes Forestry Research Centre, Fredricton, New Brunswick, *Pith to Periderm* 12(1): 3. (cn).
- *NICHOLLS 1923. [*Acacis abundans*]. Tasmania Depart-

- ment of Agriculture and Stock, Report 1922-1923: 15-17. ().
- NIELSON, R. W. 1986. Harvesting and processing of beetle-killed timber. Proceedings: seminar sponsored by Forintek Canada Corporation and Cof, Northern Interior Lumber Sector, 10 May 1985, Prince George, British Columbia. Special publication. Forintek Canada Corp. SP-26. 53 p. (cn).
- NIELSON, R. W., AND J. F. G. MACKAY. 1986. Sorting of dry and green lodgepole pine before kiln drying. Pages 31-34 in R. W. Nielson, Harvesting and processing of beetle-killed timber. Proceedings: seminar sponsored by Forintek Canada Corporation and Cof, Northern Interior Lumber Sector, 10 May 1985, Prince George, British Columbia. Special Publication. Forintek Canada Corp. SP-26. 53 p. (cn).
- *NIEMEYER, HANS. 1979. Die Entwicklung der Borkenkäferpopulationen nach 1972 und ihre Bekämpfung. Pages 5-37 in W. Altenkirch et al., Aus dem Walde, Dokumentation der Sturmkatastrophe vom 13 November 1972. Part V. Mitteilungen aus der Niedersächsischen Landesforstverwaltung 31. 190 p. ().
- _____. 1985a. Field response of *Ips typographus* L. (Col., Scolytidae) to different trap structures and white versus black flight barriers. Journal of Applied Entomology 99: 44-51. (cn).
- _____. 1985b. Freilandbeobachtungen zum Anflugverhalten und zur visuellen Orientierung des Buchdrückers (*Ips typographus* L.) an Pheromonfallen [Field studies on flight behaviour and visual orientation of *Ips typographus* in relation to pheromone traps] [English summary]. Forst- und Holzwirt 40: 85-92. (bv).
- _____. 1985c. Test und Effektivität von Borkenbefallen [Testing and effectiveness of bark beetle traps] [English summary]. Forst- und Holzwirt 40: 32-40. (cn).
- NIGAM, PRAKASH CHANDRA. 1975. Chemical insecticides. Pages 5-24 in M. L. Prebble, Aerial control of forest insects in Canada. Canada Department of the Environment, Canadian Forestry Service, Ottawa, Catalog No. Fo-23/19/1975. (cn).
- NIIJOU, P., LUDOVIC NEE AND L. WATERKEYN. 1989. Variabilité inter- et intra-individuelle de quelques caractéristiques anatomiques de l'écorce de l'épicéa commun (*Picea abies* [L.] Karsten) en Ardenne belge. Annales des Sciences Forestières 46: 85-95. (cn).
- NILSSON, A. C. 1987. Nyetablering av en grandbillefauna i det nordligste Norge som følge av innplantning av gran [Colonization by a spruce beetle fauna in northernmost Norway following planting of spruce] [In Norwegian, English summary]. Entomologiske Meddelelser 55: 169-170. (hb).
- NOBUCHI, AKIRA. 1979c. Treatment of *Xyleborus* species attacks with boric acid on pressure treated timber [In Japanese]. [Source journal?] No. 15: 11-13. (cn).
- *_____. 1979d. Bark and ambrosia beetles collected by Dr. Kintaro Baba (Col., Scol. and Platyp.). Studies on Scolytidae XVI. [Source?]: 115-121. ().
- _____. 1980d. Ambrosia beetles injurious to bed logs of Shiitake mushroom (I). Forest Pests 29(5, No. 335): 51-57. (cn ds).
- _____. 1980e. Ambrosia beetles injurious to bed logs of Shiitake mushroom (II). Forest Pests 29(6, No. 339): 109-115. (cn ds).
- _____. 1980f. Bark beetles in imported timber I [In Japanese]. [Source journal?] No. 66. 75 p. (cn).
- _____. 1980g. Bark beetles in imported timber II [In Japanese]. [Source journal?] No. 67. (cn).
- _____. 1980h. Studies on Scolytidae, XIX: Formosan Platypodidae and Scolytidae collected by Dr. Yan-I Chin (Coleoptera). Japan Coleopterological Society, Osaka 34: 93-97. (ds).
- _____. 1982. Bostrichidae and Scolytidae (Coleoptera) from Minami-Iwojima Island. Pages 357-359 in Nature Conservation Bureau, Environment Agency of Japan, December. (ds).
- _____. 1985a. The Platypodidae (Coleoptera) of the Philippines. Japan Agriculture Research Quarterly 18: 327-330. (ds tx).
- _____. 1985b. Check-list of Coleoptera of Japan: Platypodidae. Coleopterists Association of Japan 29: 1-3. (ds tx).
- _____. 1985c. Check-list of Coleoptera of Japan: Scolytidae. Coleopterists Association of Japan 30: 1-32. (ds tx).
- NOBUCHI, AKIRA AND MASASHI KAMIOCHI. 1985. Bark and ambrosia beetle fauna (Coleoptera: Platypodidae and Scolytidae) in Wakayama Prefecture [Japan]. Kinokuni 27: 1-11. (ds).
- NOGUERA-MARTINEZ, FILIPE A., AND THOMAS HARRIS ATKINSON. 1990. Biogeography and biology of bark and ambrosia beetles (Coleoptera: Scolytidae and Platypodidae) of a mesic montane forest in Mexico, with an annotated checklist of species. Entomological Society of America, Annals 53: 453-466. (hb ds).
- NOHIRA, T., AND S. OGAWA. 1986. Scolytidae (Coleoptera) collected by chemical traps for Matsukimushi [In Japanese]. Japanese Forestry Society, Journal 65: 249-250. (ds).
- NORD, JOHN C. 1971. Soft maple + CTB [Columbian timber beetle] = soft maple (WHND) [worm holes no defect] = dollars? Southern Timberman 1971 (December 15): 140-142. (cn ms).
- NORD, JOHN C., ALICE S. JONES AND FELTON L. HASTINGS. 1985. Field tests of insecticides for control of black turpentine beetle. Pages 272-280 in S. J. Branham and R. C. Thatcher, Integrated pest management research symposium: the proceedings. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- NORDANSTIG, G. 1976. Granbarkborrens ekonomiska betydelse. Pages 13-14 in Skogsskydd [In Swedish]. Sveriges Skogsvårdsförbund. (cn).
- *NORMAND, H. 1937. [*Hylesinus fraxini* (=varius)]. Societe d'Histoire Naturelle de Afrique Nord 25: 265. ().
- NORRIS, DALE MEIXIN. 1987. Electrochemistry of phytochemicals as repellents or antifeedants to insects. Pages 147-152 in V. Labeyrie, G. Fabres, and D. Lachaise, Insect-plants. Sixth International Symposium on Insect-Plant Relationships, Proceedings. Series Entomologica 41. 549 p. (ay cn).
- NOSEK, JOSEF. 1952b. K biocenologii fauny krovcu v Beskydech [Contribution a la biocenologie de la faune d'ipides des les Beskydes]. Praha, Československa Akademie vfed. 1(15): 95-100. (ec ds).
- NOTMAN, II. 1920. Coleoptera collected at Windsor, Broome Co., New York, 26 May to 5 June 1918, with notes and descriptions. New York Entomological Society, Journal 28: 178-194. (ds).
- *NOVAK, VLADIMIR. 1965. Ochrana neodkorneného suroveho dreva proti hmyzu a boj proti krovcom. Metodiky pro zavadení výsledku výzkumu do praxe. Ústav

- vedecké a technické informací. Ústředí zemědělského a potravinářského výzkumu 1: 1–27, 1 pl. ().
- °. 1971. Moderní prostředky ochrany lesů boji se škodlivými biotickými činiteli. Pages 225–234 in *Sborník konference VULHM Lesy a lesní hospodářství v rozvoji současné společnosti*. Praha 28–29 September 1971. ().
- NOVAK, VLADIMÍR, AND P. ZAHRADNÍK. 1988. The results of the use of Linoprap pheromone preparation to reduce the population of lineate bark beetle *Xyloterus lineatus* Ol. [In Czech]. *Lesnictví* (Prague) 34: 499–512. (cn).
- O**
- °OCETE, R., E. OCETE R., AND A. PEREZ. 1986. Contribución al estudio de la perforación de galerías del "barrenillo del olivo" (*Phloeotribus scarabaeoides* Bern.) (Coleoptera, Scolytidae) en Dos Hermanas (Seville) [Contribution to the study of the perforation of galleries by *Phloeotribus scarabaeoides* in Dos Hermanas (Seville)] [English summary]. *Boletim da Sociedade Portuguesa de Entomologia* supplement 1: 153–160. ().
- OCHOA M., H., AND B. DECAZY. 1987. Efecto biológico de aplicación de plaguicida a bajo volumen sobre broca del fruto del café *Hypothenemus hampei* Ferr. (Estudio preliminar) [Biological effect of low volume application of pesticide on *Hypothenemus hampei*, preliminary studies] [English summary]. *Revista Cafetalera* 27S: 21–28. (cn).
- OCHOA M., H., B. DECAZY, AND R. LOTODE. 1988. Estudio de un método de muestreo para determinar el índice de compate de la broca del fruto del café *Hypothenemus hampei* [A study of a sampling method to determine the control index for the coffee berry borer] [English summary]. *Revista Cafetalera* 287: 16–25. (cn).
- OCHOA M., H., ET AL. 1989. *Hypothenemus hampei* Ferr., en función de diferentes porcentajes de infestación [Determination of crop loss due to *Hypothenemus hampei* as a function of different percentages of infestation] [English subtitle]. *Revista Cafetalera* 303: 23–27. (cn).
- O'CONNER, BARRY M., AND JOHN CONRAD MOSER. 1985. Phylogenetic relationships of the Algophagidae (Acari: Astigmata), with descriptions of a new subfamily, genus, and species. *Entomological Society of America, Annals* 78: 783–789. (ec).
- O'CONNOR, J. P., T. G. WINTER, AND J. A. GOOD. 1991. A review of Irish Scolytidae (Insecta: Coleoptera). *Irish Naturalists Journal* 23: 389–425. (ds).
- O'DONNELL, B. P., THOMAS LEE PAYNE, AND K. D. WALSH. 1986. Effect of pine oil on landing and attack by the southern pine beetle (Coleoptera: Scolytidae). *Journal of Entomological Science* 21: 319–321. (bv cn).
- OEHLISCHLAGER, A. C., AND B. D. JOHNSTON. 1987. Synthesis of the enantiomers of endobrevicomin. *Journal of Organic Chemistry* 52: 940–944. (bv ms).
- OEHLISCHLAGER, A. C., C. G. S. KING, H. D. PIERCE, JR., A. M. PIERCE, K. N. SLESSOR, J. G. MILLAR, AND JOHN HARVEY BORDEN. 1987. Chirality of Macrolidase pheromones of grain beetles in the genera *Oryzaephilus* and *Cryptolestes* and its implications for species specificity. *Journal of Chemical Ecology* 13(6): 1543–1554. (ec ms).
- OEHLISCHLAGER, A. C., A. M. PIERCE, H. D. PIERCE, JR., AND JOHN HARVEY BORDEN. Chemical communication in cucujid grain beetles. *Journal of Chemical Ecology* 14(11): 2071–2095. (bv ec).
- OEHLISCHLAGER, A. C., AND D. VANDERSEL. 1987. Biosynthesis of pheromones in Coleoptera. *American Chemical Society (Abstract)* 194: Agro 36. (bv ms).
- OHMART, CLIFFORD P. 1982. Insects associated with *Pinus radiata* throughout the world: an annotated bibliography. [Australia] Commonwealth Scientific and Industrial Research Organization, Division of Forest Research, Canberra, Division Report 9. 79 p. (ds ms).
- °. 1989. Why are there so few tree-killing bark beetles associated with angiosperms? *Oikos* 54: 242–245. (tx ms).
- OHNESORGE, BERNHART. 1953. Der Einfluss von Geruchs- und Geschmacksstoffen auf die Wahl der Frasspflanzen beim grossen braunen Rüsselkafer *Hylobius abietis* L. *Beiträge zur Entomologie* 3: 437–467. (ec).
- OHINO, SHIZUO. 1989. Studies on Scolytidae and Platypodidae (Coleoptera) found on imported logs at Japanese ports. I: keys to the subfamilies, tribes and genera. *Research Bulletin of the Plant Protection Service Japan* 25: 7–22. (tx).
- OHINO, SHIZUO, K. YONEYAMA, AND H. NAKAZAWA. 1982a. The Scolytidae and Platypodidae (Coleoptera) from Celebes, found in logs at Nagoya port. *Research Reports, Nagoya Plant Protection Station* [Journal title in Japanese] 25: 3–6. (ds).
- °. 1982b. The Scolytidae and Platypodidae (Coleoptera) from Sumatra and Mentawai Islands, found in logs at Nagoya port. *Research Reports, Nagoya Plant Protection Station* [Journal title in Japanese] 25: 7–12. (ds).
- °. 1987a. The Scolytidae and Platypodidae (Coleoptera) from Philippines, found in logs at Nagoya port. *Research Bulletin of the Plant Protection Service Japan* 23: 87–91. (ds).
- °. 1987b. The Scolytidae and Platypodidae (Coleoptera) from Moluccas Islands, found in logs at Nagoya port. *Research Bulletin of the Plant Protection Service Japan* 23: 93–97. (ds).
- OHINO, SHIZUO, K. YOSHIOKA, N. UCHIDA, K. YONEYAMA, AND K. TSUKAMOTO. 1989. The Scolytidae and Platypodidae (Coleoptera) from Bismark Archipelago found in logs at Nagoya port Japan. *Research Bulletin of the Plant Protection Service Japan* 25: 59–69. (ds).
- OHINO, SHIZUO, K. YOSHIOKA, K. YONEYAMA, AND H. NAKAZAWA. 1988a. The Scolytidae and Platypodidae (Coleoptera) from Solomon Islands, found in logs at Nagoya port. I. *Research Bulletin of the Plant Protection Service Japan* 24: 91–95. (ds).
- °. 1988b. The Scolytidae and Platypodidae (Coleoptera) from Solomon Islands, found in logs at Nagoya port. II. *Research Bulletin of the Plant Protection Service Japan* 24: 97–99. (ds).
- °OKOŁOW, CZESŁAW. 19... Badania entomologiczne w Białowieżskim i Kampinoskim parku narodowym (stan, potrzeby) [Entomological investigations in national parks: Białowieża and Kampinos (present state, needs)]. *Entomologia i gospodarka narodowa* 19...: 19–36. ().
- °. 1968. Cykl rozwojowy jesiennika czarnego, *Hylesinus crenatus* Fabr. (Col., Scol.) [Life cycle of the large ash bark beetle, *Hylesinus crenatus*]. *Polskie pismo entomologiczne, Wrocław* 38: 627–630. ().
- °. 1980. *Polygraphus punctifrons* Thomson, eine interessante, boreale Borkenkaferart. Ein Beitrag zur

- Biologie und Ökologie. Acta Musei Reginae-hradecensis S. A Supplementum 1980: 75-76. ().
- OKUDA, HIROSHI AND S. SUZUKI. 1955. Efficacy and persistence of fenitrothion for prevention of the larch ips, *Ips cembrae* (Heer) (Coleoptera: Scolytidae) [In Japanese, English summary]. Japanese Journal of Applied Entomology and Zoology 29(4): 326-329. (cn).
- O'NEIL, CURTIS G. 1956. Mountain pine beetle in state, county, and private lands on Casper/Muddy Mountain, Wyoming. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, State and Private Forestry, Biological Evaluation R2-55-9. 5 p. (cn).
- . 1955. Mountain pine beetle on Laramie Peak, Medicine Bow National Forests, Wyoming. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Biological Evaluation R2-55-3. 19 p. (cn).
- O'NEIL, CURTIS G., AND M. SHARON. 1955. Forest pest conditions in the Rocky Mountain Region 1957. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Timber, Forest Pest and Cooperative Forestry Management, Technical Report. 33 p. (cn).
- OPANASENKO, F. I., AND A. P. KONONENKO. 1966. Specific composition and ecological peculiarities of trunk pests inhabiting the fir in north-eastern Alta [In Russian]. Pages 83-86 in A. I. Cherepanov, The fauna and ecology of Arthropoda from Siberia. USSR Academy of Sciences, Siberian Branch, Novosibirsk. (hb).
- OPPERMANN, T. A. 1955. Rinden- und Holzbrutende Insekten an Immissionsgeschädigten Fichten und Kiefern [Bark and wood insect pests of pollution-damaged spruce and pine] [English summary]. Holz-Zentralblatt 111(14): 213-217. (cn).
- ORMEROD, ELEANOR ANNE. 1890. [*Xyleborus dispar*, p. 330-334]. Manual of injurious insects. ().
- OTROSINA, W. J., AND G. T. FERRELL. 1959. Effects of water stress on white fir wound response to *Trichosporium symbioticum* fungal symbiont of the fir engraver beetle. Phytopathology 79: 1164. (ec).
- OTSUKA, T., S. J. SEYBOLD, DAVID LEE WOOD, AND I. KUBO. 1955. Optical resolution of ipsdienol and ipsenol, aggregation pheromones of bark beetles *Ips* spp. American Chemical Society (Abstract) 196:Orgn 386. (bv ms).
- OTT, H. J. 1959. Ökologische Untersuchungen zur Bedeutung von Windwurfslächen für die Insektenfauna, Teil I. Waldhygiene 17: 193-247. (cn ec).
- OTTO, H. J. 1955. Sykiculture according to site conditions as a method of forest protection. Journal of Applied Entomology 99: 190-195. (cn ec).
- OTVOS, IMRE S. 1977. Observations on the food of three forest-dwelling lizards in California. Herpetological Review 8: 6-7. (ec).
- OTVOS, IMRE S., RONALD WILLIAM STARK. 1955. Arthropod food of some forest-inhabiting birds. Canadian Entomologist 117: 971-990. (ec).
- OVERHULSER, D. L. 1956. Southwest Oregon forest insect pests. Pages 29-34 in O. T. Helgeson, Forest pest management in southwest Oregon. Forest Research Laboratory, Oregon State University, Corvallis, Oregon. (cn).
- OWEN, DONALD R. 1956. Annual Southwestern Region pest conditions report, 1955. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico, Forest Pest Management Report R3-56-4. 17 p. (cn).
- OWEN, DONALD R., K. Q. LINDAHL, JR., DAVID LEE WOOD, AND J. PARMETER. 1957. Pathogenicity of fungi isolated from *Dendroctonus valens*, *D. brevicornis*, and *D. ponderosae* to ponderosa pine seedlings. Phytopathology 77: 631-636. (ec).

P

- PAGE, MARION, MICHAEL I. HAWERTY, AND CHARLES E. RICIMOND. 1955. Residual activity of carbaryl protected lodgepole pine against mountain pine beetle, Dillon, Colorado, 1952 and 1953. United States Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station, Berkeley, California, Research Note PSW-375. 4 p. (cn).
- PAGE, P. C. B., C. M. RAYNER, AND I. O. SUTHERLAND. 1956. Stereospecific synthesis of exo- and endo-1,3-dimethyl-2,9-dioxabicyclo-3,3,1-nonane. Tetrahedron Letters 27(30): 3535-3535. (bv ms).
- PAINE, TIMOTHY D., C. A. BLANCHE, T. EVAN NEBEKER, FREDERICK M. STEPHEN. 1955. Composition of loblolly pine resin defenses: comparison of induced lesion and sapwood resin. Canadian Journal of Forest Research 17: 1202-1206. (cn ec).
- PAINE, TIMOTHY D., R. G. CATES, AND FREDERICK M. STEPHEN. 1958. Moisture stress, tree suitability, and southern pine beetle population dynamics. International Congress of Entomology, Proceedings 18: 413. (ec hb).
- PAINE, TIMOTHY D., AND FREDERICK M. STEPHEN. 1957a. Fungi associated with the southern pine beetle: avoidance of induced defense response in loblolly pine. Oecologia [Berlin] 74: 377-379. (ec).
- . 1957b. Influence of tree stress and site quality on the induced defense system of loblolly pine. Canadian Journal of Forest Research 17: 569-571. (ec).
- . 1957c. Response of loblolly pine to different inoculum doses of *Ceratocystis minor*, a blue-stain fungus associated with *Dendroctonus frontalis*. Canadian Journal of Botany 65: 2093-2095. (ec).
- . 1957d. The relationship of tree and crown class to the induced-plant defenses of loblolly pine. Canadian Journal of Botany 65: 2090-2092. (ec).
- . 1958. Induced defenses of loblolly pine, *Pinus taeda*: potential impact on *Dendroctonus frontalis* within-tree mortality. Entomologia Experimentalis et Applicata 46: 39-46. (ec).
- PAINE, TIMOTHY D., FREDERICK M. STEPHEN, AND R. G. CATES. 1955. Induced defenses against *Dendroctonus frontalis* and associated fungi: variation in loblolly pine resistance. Pages 169-176 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- . 1955a. Moisture stress, tree suitability, and southern pine beetle population dynamics. Pages 85-103 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (ec hb).
- . 1955b. Phenology of an induced response in loblolly pine following inoculation of fungi associated with the southern pine beetle. Canadian Journal of Forest Research 18: 1556-1562. (cn ec).

- PAINÉ, TIMOTHY D., FREDERICK M. STEPHEN, AND G. N. MASON. 1955. A risk model integrating stand hazard and southern pine beetle population level. Pages 201–212 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 240 p. (cn).
- PAINÉ, TIMOTHY D., FREDERICK M. STEPHEN, G. W. WALLIS, AND J. F. YOUNG. 1955. Seasonal variation in host tree defense to the southern pine beetle. *Arkansas Farm Research* 34: 5. (cn).
- PAIVA, MARIA ROSA AND K. KIESEL. 1955a. Field response of *Trypodendron* spp. (Col., Scolytidae) to different concentrations of lineatin and alpha-pinene. *Journal of Applied Entomology* 99: 442–448. (bv).
- . 1955b. Lineatin biosynthesis and interspecific communication in *Trypodendron* spp. (Col., Scolytidae). *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 4(4–6): 402–405. (bv).
- PAIVA, MARIA ROSA, M. F. PESSOA, AND JEAN PIERRE VITE. 1955. Reduction in the pheromone attractant response of *Orthotomicus erosus* (Woll.) and *Ips sexdentatus* Boern. (Col., Scolytidae). *Journal of Applied Entomology* 106: 198–200. (bv).
- PAIVA, MARIA ROSA AND JEAN PIERRE VITE. 1954. As feromonas de agregacao no controlo de Scolytidae: algumas implicacoes ecologicas [The aggregation pheromones in the control of Scolytidae: some ecological implications] [English summary]. *Boletim da Sociedade Portuguesa de Entomologia* 11, 20: 225–235. (bv cn).
- *PAJARES ALONSO, JUAN A. 1957. Contribucion al conocimiento de los escolitidos vectores de la grafiosis en la peninsula ibérica. Unpublished dissertation, Universidad Politecnica de Madrid, Spain. 242 p. ().
- PAJARES ALONSO, JUAN A., AND M. J. AREVALO. 1957. Protección de los olmos contra los insectos vectores de la grafiosis [Protection of elms against vectors of Dutch elm disease] [English summary]. *Boletim de Sanidad Vegetal, Plagas* 13(3): 311–325. (cn).
- PAJARES ALONSO, JUAN A., AND LUIS GIL. 1955. La grafiosis de los olmos. Ministerio de Agricultura, Pesca y Alimentación, Publicaciones de Extensión Agraria, Hojas Divulgadoras 19. 24 p. (cn).
- *PALM, THURE WILHELM. 1965. [*Hylesinus oleiperda* (= *toranio*)]. *Opuscula Entomologica* 30: 167. ().
- . 1956. Skallbagsstudier pa Stenshuvid. *Entomologisk Tidskrift* 107: 53–55. (ds).
- PARIHAR, D. R., AND A. J. HAYES. 1957. Biology of *Scolytus laevis* Chapuis (Coleoptera: Scolytidae), a vector of Dutch elm disease in Scotland. *Scottish Forestry* 41: 155–196. (hb).
- PARK, J. D., AND B. H. BYUN. 1955. Trapping the overwintering pine bark beetle, *Tomicus piniperda* L. (Coleoptera: Scolytidae), by turpentine [In Japanese, English captions and tables]. *Research Reports of the Forestry Research Institute (Seoul)* 36: 126–129. (cn).
- PARMETER, JOHN R., JR., GAREY W. SLAUGHTER, M. M. CHEN, DAVID LEE WOOD, AND H. A. STUBBS. 1959. Single and mixed inoculations of ponderosa pine with fungal associates of *Dendroctonus* sp. *Phytopathology* 79: 768–772. (ec).
- PATOCKA, JAN. 1955. Biology and forestry importance of *Scolytus intricatus* Ratz. (Coleoptera: Scolytidae) [In Slovak]. *Zpravy lesnického výskumu, Zvolen* 3: 17–21. (cn hb).
- PATOCKA, JAN, AND J. NOVOTNY. 1957. Úcast hmyzu na hromadnom hynuti dubov na Slovensku [Share of insects in the mass decay of oak in Slovakia]. *Vedecké práce Výskumného ústavu lesného hospodárstva vo Zvolene* 36: 59–90. (cn ec).
- *PAWŁOWSKI, JERZY. 1972. The beetles (Coleoptera) of Babia Góra Mountain [Polish western Carpathians]. *Acta Zoologica Cracoviensia* 12: 19–251. ().
- *PAX, FERDINAND. 1921. [*Hylastes angustatus*]. *Jahresheft des Vereins für Schlesische Insektenkunde zu Breslau, Breslau* 1921: 44–56. ().
- PAYNE, THOMAS LEE. 1970. Sex communication between insects. Pages 57–64 in Texas Agricultural Experiment Station Conference, Proceedings. (bv hb).
- . 1980a. Behavioral chemicals in bark beetle manipulation [abstract]. Joint Congress, European Chemoreception Research Organization and International Symposium on Olfaction and Taste, Netherlands. (bv).
- . 1980b. Influence of inhibitors on southern pine beetle behavior [abstract]. Biology and behavior of bark and ambrosia beetles: symposium proceedings. International Congress of Entomology. Proceedings 16: 284. (bv).
- . 1980c. Pheromonal control of *Dendroctonus frontalis* and its application to pest management [abstract]. Symposium proceedings: Pheromonal control of insect behavior and its application to pest management. International Congress of Entomology. Proceedings 16: 222. (bv cn).
- . 1986a. Olfactory basis of insect enemies of allied species. Symposium proceedings: Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Entomological Society of America. (bv cn).
- . 1986b. Olfaction and vision in host finding by a bark beetle. Pages 111–116 in T. L. Payne, M. C. Birch, and C. E. J. Kennedy, Mechanisms in insect olfaction. Clarendon Press, Oxford. xvi + 364 p. (bv).
- . 1955. Pheromone plus cut and leave for southern pine beetle suppression. International Congress of Entomology. Proceedings 15: 415. (bv cn).
- PAYNE, THOMAS LEE, N. A. ANDRYSZAK, H. WIESER, E. A. DIXON, N. IBRAHIM, AND J. COERS. 1955. Antennal olfactory and behavioral response of southern pine beetle, *Dendroctonus frontalis*, to analogs of its aggregation pheromone frontalin. *Journal of Chemical Ecology* 14: 1217–1226. (bv).
- PAYNE, THOMAS LEE, AND RONALD FORREST BILLINGS. 1955. Effect of pheromone plus tree cutting on suppression of southern pine beetle infestations. Pages 267–273 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn).
- . 1959. Evaluation of (s)-verbenone applications for suppressing southern pine beetle (Coleoptera: Scolytidae) infestations. *Journal of Economic Entomology* 52: 1702–1705. (cn).
- PAYNE, THOMAS LEE, RONALD FORREST BILLINGS, J. D. DELORME, N. A. ANDRYSZAK, J. BARTELS, W. FRANCKE, AND JEAN PIERRE VITE. 1957. Kairomonal-pheromonal system in the black turpentine beetle, *Dendroctonus terebrans* (Ol.). *Journal of Applied Entomology* 103: 15–22. (bv).
- PAYNE, THOMAS LEE, MARTIN C. BIRCH, AND C. E. J. KENNEDY

1956. Mechanisms in insect olfaction. Clarendon Press, Oxford. xvi + 364 p. (bv).
- PAYNE, THOMAS LEE, JACK E. COSTER, AND J. V. RICHIERSON. 1984. Insect predator-prey coevolution via enantiomeric specificity in a kairomone-pheromone system. *Journal of Chemical Ecology* 10: 487-492. (bv ec).
- PAYNE, THOMAS LEE, AND ROBERT N. COULSON. 1985. Role of visual and olfactory stimuli in host selection and aggregation behavior by *Dendroctonus frontalis*. Pages 73-82 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Canadian Forestry Service, and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1983. 240 p. (bv).
- PAYNE, THOMAS LEE, L. H. KUDON, CHARLES W. BERISFORD, P. P. O'DONNELL, AND KIERON D. WALSH. 1955. Effects of frontalure in suppressing southern pine beetle spot growth under endemic and epidemic population levels. Pages 251-257 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (bv cn).
- PAYNE, THOMAS LEE, L. H. KUDON, KIERON D. WALSH, AND CHARLES W. BERISFORD. 1955. Influence of infestation density on suppression of *D. frontalis* infestations with attractant. *Journal of Applied Entomology* 99: 39-43. (bv cn).
- PAYNE, THOMAS LEE, AND J. V. RICHIERSON. 1955. Pheromone-mediated competitive replacement between two bark beetle populations: influence of infestation suppression. *Journal of Applied Entomology* 99: 131-135. (bv ec).
- PAYNE, THOMAS LEE, AND H. SAARENMAA. 1988. Integrated control of scolytid bark beetles. IUFRO Working Party and the XVII International Congress of Entomology Symposium, Vancouver, British Columbia. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- PAYTON, I. J. 1989. Fungal (*Sporothrix*) induced mortality of Kamahia (*Weinmannia racemosa*) after attack by pin-hole borer (*Platypus* spp.). *New Zealand Journal of Botany* 27: 359-365. (ec).
- PEACOCK, JOHN WILLIAM, ROY A. CUTHBERT, AND GERALD NORMAN LANIER. 1981. Deployment of traps in a barrier strategy to reduce populations of the European elm bark beetle, and the incidence of Dutch elm disease. Pages 155-174 in E. R. Mitchell, Management of insect pests with semiochemicals. Concepts and practice. Plenum Press, New York. 514 p. (bv cn).
- PECK, OSWALD. 1963. A catalogue of the Nearctic Chalcidoidea (Insecta: Hymenoptera). Canadian Entomologist, Supplement 30. 1092 p. (ec).
- PECK, STEWART B., JARMILA KUKALOVA-PECK, AND C. BORDON A. 1989. Beetles (Coleoptera) of an oil-bird cave: Cueva del Guacharo, Venezuela. *Coleopterists Bulletin* 43: 151-156. (hb).
- PEDROSA-MACEDO, JOSE HENRIQUE. 1955. Insect pests and their control in pine plantations in Brazil. Pages 149-161 in Protection of forests in the tropics: noxious insects to pine and eucalyptus plantations in the tropics. Universidade Federal do Parana, Curitiba, Brazil. (cn ds).
- PEDROSA-MACEDO, JOSE HENRIQUE, AND J. SCHONHERR. 1985. Manual dos Scolytidae nos reflorestamentos Brasileiros [Manual of the Scolytidae in reforested areas in Brazil] [English summary]. Universidade Federal do Parana, Curitiba, Parana, Brazil. vi + 69 + 11 p. (ds tx).
- PERFECTO, I. 1985. Variation in attack rates among subpopulations of *Coccotrypes carpophagus* utilizing *Eutrepe globosa* seeds at three locations in Puerto Rico. *Tropical Ecology* 29: 114-120. (bv cn).
- PERICART, J., AND J. HALPERIN. 1989. The Anthocoridae of Israel (Heteroptera). *Phytoparasitica* 17: 91-99. (ec).
- PERROUD, B. P. 1864. *Bostrichus Boieldieu*. Page 185 in B. P. Perroud and X. Montrouzier, Essai sur la faune entomologique de Kanala (Nouvelle-Caledonie) et description de quelques especes nouvelles on peu connue. Societe Linneenne de Lyon. Annales, ser. 2, 11: 46-257. (tx).
- PERSON, H. L. 1940. The clerid *Enoclerus lecontei* (Woll.) as a factor in the control of the western pine beetle. *Journal of Forestry* 38: 390-396. (ec).
- PERUSQUA ORTIZ, JUSTINA. 1982. Insectos asociados a los descortezadores *Dendroctonus* spp. de los pinos [English summary]. Secretaria de Agricultura y Recursos Hidraulicos, Subsecretaria Forestal y de la Fauna, Instituto Nacional de Investigaciones Forestales, Mexico, D.F., Boletin Technico 83. 33 p. (cn ec).
- PETRENKO, E. S. 1966. The ipidofauna of felling areas in the pine forest of Krasnojarsk Angara region [In Russian]. Pages 77-80 in A. I. Cherepanov, The fauna and ecology of Arthropoda from Siberia. USSR Academy of Sciences, Siberian Branch. Novosibirsk. (hb).
- PETTY, T. M., AND C. C. SHAW. 1986. Isolation of *Fomitopsis pinicola* from in-flight bark beetles (Coleoptera, Scolytidae). *Canadian Journal of Botany* 64: 1507-1509. (ec).
- *PETTIT. 1899. [*Scolytus rugulosus*]. Michigan Agricultural Experiment Station, Bulletin 125(175): 363-365. ().
- _____. 1904. [*Scolytus rugulosus*]. Michigan Agricultural Experiment Station, Bulletin 24: 34, 53-55, 59, fig. 53. ().
- PETTY, J. 1967. Annual district report: Grande Prairie, Slave Lake district, 1966. Pages 73-82 in District reports of the Forest Insect and Disease Survey, Alberta-Northwest Territories-Yukon region, 1966. Canadian Department of Forestry and Rural Development, Forestry Research Laboratory, Calgary, Alberta, Information Report A-X-6. (cn).
- PFEFFER, ANTONIN. 1947e. Urcovaci tabulky nejdnelejších hmyzích škůdců, jejichž na našich dřevinách. Zvláštní otisk z Male encyklopedie lesniců, díl I., oddíl V. Nakladem vlastním. Tiskem knihtiskárny Bohumil Novotný, Písek. 28 p. (ds tx).
- *_____. 1948. [*Pityogenes quadridens*]. Entomologické Listy, Folia Entomologica, Brno 9(1-4): 116. ().
- *_____. 1971. Hmyz jako bioindikátor [Insekten als Bioindikatoren]. Bioindicators of landscape deterioration, Praha 1971: 83-85. ().
- *_____. 1976. Insekten als Indikatoren von Veränderungen in der Bestandzusammensetzung der Südböhmischen Moore (Einleitung, Problematik, Südböhmische Späke, Ökologische Grundlagen, Methodik, Zusammenfassung der Ergebnisse Schlussfolgerung). Quaestiones Geobiologicae 16: 74-98. ().
- *_____. 1977. Famistic records from Czechoslovakia. Col., Scol., *Trypophloeus spiculatus*, *T. granulatus*, *T.*

- ryhinski*. Acta Entomologica Bohemoslovaca 74: 207. ().
- _____. 1975. Wirkungen von Luftverunreinigungen auf die freilebende Tierwelt. Schweizerischen Zeitschrift für Forstwesen 129: 362–367. (ec).
- _____. 1980. Insektenfauna der Hochgebirgswälder Mitteleuropas. Acta Musei Reginahradensis S. A. Supplementum 1980: 77–80. (ec).
- _____. 1984. Synopsis der gattung *Hylastinus* Bedel (Col., Scolytidae). Acta Entomologica Jugoslavica 20: 9–13. (ds tx).
- _____. 1985a. Sexualdimorphismus der Arten der Gattung *Carphoborus* Eichhoff und taxonomische Bemerkungen zu den einzelnen Arten (Coleoptera: Scolytidae) [Sexual dimorphism of species of the genus *Carphoborus* Eichhoff and taxonomic notes on the individual species] [English summary]. Acta Entomologica Bohemoslovaca 82: 468–475. (ay ds tx).
- _____. 1985b. Zur taxonomischen Stellung der Gattung *Saliciphilus* Sokanovsky (Coleoptera, Scolytidae) [The taxonomic position of the genus *Saliciphilus*] [English summary]. Acta Entomologica Bohemoslovaca 82: 138–142. (tx).
- _____. 1987. *Taphrocetes* gen. n. und die Heterogenität der Gattung *Taphrorychus* Eichhoff (Coleoptera, Scolytidae) [*Taphrocetes* gen. n. and heterogeneity of the genus *Taphrorychus*] [English summary]. Acta Entomologica Bohemoslovaca 84: 22–26. (tx).
- _____. 1989a. Kurovcoviti Scolytidae a jadrohlodoviti Platypodidae. Československa Akademie Veg, Praha. 137 p. (ds tx).
- _____. 1989b. Taxonomischer Status einiger Arten der Gattung *Xyloterus* Erichson (Coleoptera, Scolytidae) [The taxonomic position of some species of the genus *Xyloterus*] [English summary]. Acta Entomologica Bohemoslovaca 86: 129–136. (ds tx).
- _____. 1991. The taxonomic position of genus *Taphromurgus* (Coleoptera, Scolytidae). Acta Entomologica Bohemoslovaca 88: 211–214. (tx).
- _____. 1992. Nomenclatural note on *Crypturgus mediterraneus* (Coleoptera, Scolytidae). Acta Entomologica Bohemoslovaca 89: 69–70. (tx).
- PFEFFER, ANTONIN, AND MILOS KNIZEK. 1989. Problematika kurovcu introdukovanych do Evropy. Lesnické Prace 1989: 311–312. (cn ds).
- PFEFFER, ANTONIN, AND VACLAV ZUMR. 1983. Communities of Coleoptera on the silver fir (*Abies alba*). Acta Entomologica Bohemoslovaca 80: 401–412. (ec).
- PFEIFFER, DOUGLAS G. 1985. Ambrosia beetles in Virginia vineyards. Virginia Journal of Science 36: 91. (cn ec).
- PHILLIPS, T. W. 1988. Chemical signals and male strategies in Scolytidae. International Congress of Entomology. Proceedings 18: 430. (bv).
- PHILLIPS, T. W., THOMAS HARRIS ATKINSON, AND JOHN L. FOLTZ. 1989. Pheromone-based aggregation in *Orthotomicus caelatus* Eichhoff (Coleoptera: Scolytidae). Canadian Entomologist 121: 933–940. (bv).
- PHILLIPS, T. W., J. L. NATION, ROBERT CLEVELAND WILKINSON, AND JOHN L. FOLTZ. 1989. Secondary attraction and field activity of beetle-produced volatiles in *Dendroctonus terebrans*. Journal of Chemical Ecology 15: 1513–1534. (bv).
- PHILLIPS, T. W., A. J. WILKENING, THOMAS HARRIS ATKINSON, J. L. NATION, ROBERT CLEVELAND WILKINSON, AND JOHN L. FOLTZ. 1988. Synergism of turpentine and ethanol as attractants for certain pine-infesting beetles (Coleoptera). Environmental Entomology 17: 456–462. (bv).
- °PHILLIPSEN, W. J., AND M. E. ASCERNO. 1980. Native elm bark beetle control. Minnesota Tree Line Journal 27. 2 p. ().
- PHILLIPSEN, W. J., M. E. ASCERNO, AND V. R. LANDWEHR. 1986. Colonization, emergence, and survival of *Hylurgopinus rufipes* and *Scolytus multistriatus* (Coleoptera: Scolytidae) in insecticide-treated elm wood. Journal of Economic Entomology 79: 1349–1350. (cn).
- °PIC, MAURICE. 1915. Notes diverses, descriptions et diagnostics. L'Echange. Revue Linneenne 31(362): 1 [etc.]. ().
- PIERCE, A. M., AND H. D. PIERCE, JR. 1955. Beetle attractant: U. S. Patent 4,560,551. Official Gazetteer. United States Patent Trademark Office 106(4): 1682. (cn ms).
- PIERCE, H. D., JR., J. E. CONN, A. C. OEHLISCHLAGER, AND JOHN HARVEY BORDEN. 1987. Monoterpene metabolism in female mountain pine beetles, *Dendroctonus ponderosae* Hopkins, attacking ponderosa pine. Journal of Chemical Ecology 13: 1455–1480. (ay bv).
- PIERCE, JOHN, AND J. KLIEJUNAS. 1981. Biological evaluation of conifer mortality on the proposed Smith Salvage Sale, Klamath National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 51–25. (cn).
- PIGNAL, M. C., CONSTANTIN CHARARAS, AND M. BOURGEAY-CAUSSE. 1987. Isolement et etude de levures du tractus digestif d'*Ips sexdentatus*, Coleoptere parasite des coniferes [Isolation and study of yeasts from the digestive tract of *Ips sexdentatus*, a beetle pest of conifers] [English summary]. Comptes Rendus de l'Academie des Sciences, III (Sciences de la Vie) 304(17): 449–452. (ay).
- _____. 1988. Yeasts from *Ips sexdentatus* (Scolytidae): enzymatic activity and vitamin excretion. Mycopathologia 103: 43–48. (ec).
- PILECKIS, S. 1986. Idomioji entomologija. Vilnius Mokslas. 240 p. (ms).
- PILNY, JERRY J., AND ALAN V. MORGAN. 1987. Paleontology and paleoecology of a possible Sangamonian site near Innerkip, Ontario. Quaternary Research 28: 157–174. (ds).
- PILNY, JERRY J., ALAN V. MORGAN, AND ANNE MORGAN. 1987. Paleoclimatic implications of a Late Wisconsinan insect assemblage from Rostock, southwestern Ontario. Canadian Journal of Earth Sciences 24: 617–630. (ds).
- PINKSTON, KEN, RAY D. EIKENBARY, AND DON ARNOLD. 198.. Shade tree borers. Oklahoma State University Cooperative Extension Service, Extension Facts No. 7315. 6 p. (cn).
- PIOU, D., AND FRANCOIS LIEUTIER. 1989. Symptomatology observations and possible roles of *Ophiostoma minus* Hedge. (Ascomycetes: Ophiostomatales) and *Tomicus piniperda* L. (Coleoptera: Scolytidae) in the dieback of Scots pine in the Orleans forest France [In French]. Annales des Sciences Forestieres 46: 39–54. (ec).
- PLANK, M. E., T. A. SNELGROVE, AND T. D. FAHEY. 1986. Volume and value recovery from live and dead lodgepole pine. Pages 27–30 in R. W. Nielson, Symposium proceedings: Harvesting and processing of beetle-killed timber. Special publication, Forintek Canada Corp. SP-26. 53 p. (cn).
- PLAZA, ESPERANZA. 1983. Los representantes espanoles de las tribus Crypturgini y Pityophthorini (Col., Scolytidae). Eos 59: 223–241. (ds tx).

- POHL-APPEL, G., AND K. RENNER. 1957. Coleopterologische analyse des inhaltes von Borkenkäfer-Pheromoufallen im raum Bielefeld [Coleopterological analysis of the content of bark beetle pheromone traps in the Bielefeld area] [In German, English summary]. *Decheniana* 140: 79-86. (cc).
- POPP, M. P., ROBERT CLEVELAND WILKINSON, E. J. JOKELA, R. B. HARDING, AND T. W. PHILLIPS. 1989. Effects of slash pine phloem nutrition on reproductive performance of *Ips calligraphus* (Coleoptera: Scolytidae). *Environmental Entomology* 18: 795-799. (hb).
- POWELL, JOHN M. 1964. The mountain pine beetle, *Dendroctonus ponderosae* Hopk. (d) bioclimatic studies. Page 117 in Canada Department of Forestry, Forest Entomology and Pathology Branch, Annual Report for 1964. (cc).
- _____. 1971. The arthropod fauna collected from the command blister rust, *Cronartium comandrae*, on lodgepole pine in Alberta. *Canadian Entomologist* 103: 908-918. (cc).
- POWELL, W. 1980. A revised checklist of forest Coleoptera in the reference collection at the Forestry Research Institute of Malawi. *Forestry Research Institute of Malawi*, Forestry Research Record 59. 32 p. (tx).
- PRICE, TERRY S. 1957. Pine bark beetles in urban areas. Georgia Forestry Commission, Macon, Georgia. (cn).
- _____. 1958a. A guide to common forest pests in Georgia. Georgia Forestry Commission 3-1-235, November 1958. 40 p. (cn tx).
- _____. 1958b. Pine bark beetles in urban areas. Georgia Forestry Commission, undated circular [ca 1958]. 5 p. (cn).
- PRICE, TERRY S., AND COLEMAN DOGGETT. 1952. A history of southern pine beetle outbreaks in the southeastern United States: revised. Georgia Forestry Commission 1952(April): 1-35. (cn).
- PRICE, TERRY S., AND KERRY G. THOMAS. 1952. Little bug—big trouble. *TOPS* 1952(December): 4-5, 20, 22. (cn).
- *PRIHODA, A. 1952. Z bionomie kurovce *Lymantria coryli* Perr. Ochrana prirody. Praha 4: 126. ().
- * _____. 1964. Kurovec *Limantia aceris* Lindemann a houba *Diaporthe fibrosa* (Pers.) Fuckel v Cechach [Le scolytid *Lymantria aceris* et le champignon *Diaporthe fibrosa* en Boheme]. *Sbornik Vysoke školy zemědělské v Praze* 1964: 373-377. ().
- * _____. 1982. Hubové nákazy jaseňov po napadnutí hřokazom jaseňovým *Hylesinus fraxini* (Panzer) [Fungal infections of ash trees after attack by the common ash bark beetle *Hylesinus fraxini*]. *Lesnický časopis*, Bratislava 28: 19-27. ().
- PRIOR, C. 1956. Sudden death of cocoa in Papua New Guinea associated with *Phytophthora palmivora* cankers invaded by bark beetles. *Annals of Applied Biology* 109: 535-543. (cc).
- PRONOS, JOHN, AND J. W. DALE. 1951. Biological evaluation at Bull Meadow Timber Sale on the Goosenest Ranger District, Klamath National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 81-4. (cn).
- PRONOS, JOHN, J. W. DALE, AND J. E. BORRECCO. 1951. Evaluation of pests on the Mt. Pinos Ranger District, Los Padres National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 81-17. (cn).
- PRONOS, JOHN, AND D. SCHULTZ. 1984a. An evaluation of pest conditions in Rancheria Campground, Pineridge Ranger District, Sierra National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-05. (cn).
- _____. 1984b. An evaluation of pests in Buck Meadow Campground, Kings River Ranger District, Sierra National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-04. (cn).
- *PORTMAN, R. W., AND MANIS. 1954. [*Ilyastinus obscurus*, *Scolytus rugulosus*]. Idaho Agricultural Extension Bulletin 216: 16, 51-52. ().
- PROVANCHER, L. 1878. Additions and corrections a la faune Coleopterologique de la province de Quebec, part 2. C. Darveau, Quebec. 18 p. (ds).
- *PRUEFFER. 1948. [*Crypturgus cinereus*]. *Torinis. Stud. Soc. Torinensis, Sec. E (Zool.)* 112: 23. ().
- PULLEY, PAUL E. 1979. Accuracy and precision of the topological mapping procedure for estimating within-tree populations of bark beetles. *Research in Population Ecology* 21: 43-52. (cn ms).
- PULLEY, PAUL E., R. O. FLAMM, AND ROBERT N. COULSON. 1955. Estimating *Dendroctonus frontalis* (Coleoptera: Scolytidae) daily infestation dynamics. *Researches on Population Ecology* 30: 193-214. (cn ec hb).
- PURRINI, KURTISH. 1955. On disease agents of insect pests of wild palms and forests in Tanzania. *Journal of Applied Entomology* 99: 237-240. (cc).
- PURRINI, KURTISH, AND J. WEISER. 1955. Ultrastructural study of the microsporidian *Chytridiopsis typographi* (Chytridiopsida: Microspora) infecting the bark beetle *Ips typographus* (Scolytidae: Coleoptera) with new data on spore dimorphism. *Journal of Invertebrate Pathology* 45: 66-74. (cc).

Q

- QIU, DEXUN, AND HUO SHAO-TANG. 1958. The preliminary observation on the habit of *Ips typographus* (Linnaeus) [In Chinese]. *Kunhong Zhishi* 4: 264-268. (hb).
- QIU, H. G., W. J. FU, Y. T. QI, L. F. HE, AND X. D. LING. 1958. Studies on the aggregation pheromone of *Ips subelongatus*, II: the relationship between aggregation behaviour and its host plant tree [In Chinese, English summary]. *Contributions from Shanghai Institute of Entomology* 5: 67-72. (bv).
- *QUAINTANCE, ALTUS LACY. 1899. [*Scolytus rugulosus*]. United States Department of Agriculture, Division of Entomology: n. s., Bulletin 20: 56-59. ().
- * _____. 1901. [*Scolytus rugulosus*]. Maryland Agricultural Experiment Station, Report 1901: 103, fig. 20. ().
- * _____. 1905. [*Scolytus rugulosus*]. United States Department of Agriculture, Yearbook 1905: 346. ().
- *QUAINTANCE, ALTUS LACY, AND EDOUARD HORACE SIEGLER. 1915a. [*Scolytus rugulosus*]. United States Department of Agriculture, Farmers Bulletin 908: 89. ().
- * _____. 1915b. [*Scolytus rugulosus*]. United States Department of Agriculture, Farmers Bulletin 1270: 66. ().

R

- RABAGLIA, ROBERT JOSEPH. 1954. Aspects of the chemical ecology of twig-crotch feeding by *Scolytus multi-striatus* (Coleoptera: Scolytidae). Unpublished dis-

- sertation, State University of New York, Syracuse. 102 p. (ec hb).
- RABASSE, J. M. 1979. Les insectes ravageurs des cypres en France. Pages 217–222 in V. Grasso and P. Raddi, Seminario: Il cypressso malattie e difesa. Comunità Economica Europea, Direzione Generale dell'Agricoltura, Agimied Sottogruppo Fitopatologie Mediterranea. (cn).
- RADHAKRISHNAN, S., P. K. RAMAIAH, AND P. K. BHAT. 1957. Methodology to estimate yield loss in coffee due to insect pests. *Journal of Coffee Research* 17: 90–93. (cn ms).
- RAFFA, KENNETH FRANCIS. 1955a. Host orientation behavior of *Dendroctonus ponderosae*: integration of token stimuli host and defenses. Pages 369–390 in W. J. Mattson, J. Levieux, and C. Bernard-Dagan, Mechanisms of woody plant defenses against insects: search for pattern. Springer, New York. xiv + 416 p. (bv).
- _____. 1955b. The mountain pine beetle in western North America. Pages 505–530 in A. A. Berryman, Dynamics of forest insect populations: patterns, causes, implications. Plenum Press, New York. xx + 603 p. (bv cn ec).
- _____. 1959. Genetic engineering of trees to enhance resistance to insects. *Bioscience* 39: 524–534. (cn).
- RAFFA, KENNETH FRANCIS, AND ALAN ANDREW BERRYMAN. 1956. A mechanistic computer model of mountain pine beetle populations interacting with lodgepole pine stands and its implications for forest managers. *Forest Science* 32: 789–805. (cn hb).
- _____. 1957. Interacting selective pressures in conifer-bark beetle systems: a basis for reciprocal adaptations? *American Naturalist* 129(2): 234–263. (bv ec).
- RAFFA, KENNETH FRANCIS, ALAN ANDREW BERRYMAN, J. SIMASKO, W. TEAL, AND B. L. WONG. 1955. Effects of grand fir monoterpenes on the fir engraver *Scolytus ventralis* (Coleoptera: Scolytidae), and its symbiotic fungus. *Environmental Entomology* 14: 552–556. (ec cn).
- RAFFA, KENNETH FRANCIS, AND K. D. KLEFZIG. 1959. Chiral escape of bark beetles from predators responding to a bark beetle pheromone. *Oecologia [Berlin]* 80: 566–569. (bv).
- RAFFA, KENNETH FRANCIS, AND EUGENE B. SMALLEY. 1955a. Host resistance to invasion by lower stem and root infesting insects of pine: response to controlled inoculations with the fungal associate *Leptographium terebrantis*. *Canadian Journal of Forest Research* 18(6): 675–681. (ec).
- _____. 1955b. Response of red and jack pines to inoculation with microbial associates of the pine engraver, *Ips pini* (Coleoptera: Scolytidae). *Canadian Journal of Forest Research* 18: 551–556. (ec).
- _____. 1955c. Seasonal and long-term responses of host trees to microbial associates of the pine engraver *Ips pini*. *Canadian Journal of Forest Research* 18: 1624–1634. (cn ec).
- RAIMO, BERNARD J. 1955. A ponderosa pine complex: *Choristoneura lambertiana*, *Neophasia menapia* (F. & F.), *Zelleria hainbachii* (Busck), *Pityophthorus aquilus* Blackman on the San Juan National Forest and adjacent lands. United States Department of Agriculture, Forest Service, Rocky Mountain Region, Lakewood, Colorado, Timber, Forest Pest and Cooperative Forestry Management, Biological Evaluation R2–55–5. 10 p. (cn).
- RAISANEN, HANNU, L. LAINE, I. KERO, AND T. KALEVA. 1956. Alustavia tutkimustuloksia hyonteis- ja sienituhoista pystykarstiissa maimikoissa [Preliminary study on insect and fungal damage in pruned Scots pine stands] [In Finnish, English summary]. *Folia Forestalia* 663. 18 p. (cn hb).
- *RAKKE. 1960. [*Hylurgops palliatus*]. Meddelelser fra det Norske Skogforsoksvesen, Vollebakk. 1960: 315. ().
- RAMESH, P. K. 1957. Observations on crop loss in robusta coffee due to mealybug and shot-hole borer. *Journal of Coffee Research* 17: 94–95. (cn).
- RAMISCH, H. 1956. Zur Wirtsfindung von *Trypodendron domesticum* und *Trypodendron lineatum* (Coleoptera: Scolytidae) [Host finding in *Trypodendron domesticum* and *Trypodendron lineatum*] [English summary]. *Journal of Applied Entomology* 73: 159–197. (bv hb).
- RAMISCH, H., AND S. BOMBOSCH. 1956. Ein Beitrag zur Brutbaumerkennung von *Trypodendron lineatum* (Ol.) und *Trypodendron domesticum* (L.) [A contribution to breeding-site recognition by *Trypodendron lineatum* and *Trypodendron domesticum*] [English summary]. *Journal of Applied Entomology* 102: 422–431. (bv hb).
- RANASINGHE, M. A. S. K., AND R. L. WICKREMASINGHE. 1955. Use of acetates to induce host resistance to *Xyleborus fornicatus* (Coleoptera: Scolytidae) attacking tea. Pages 251–256 in Proceedings: British Crop Protection Conference: pests and diseases. British Crop Protection Council, London. (cn ec).
- *RANDUSKA, P. 1951. Aktivizacia podkornikov v porastach poskodenyh poziarom v SPR Kysel, CHKO Slovensky raj. Autoreferat., Vysoka skola lesnicka a drevarska Zvolen, Lesnicka fakulta. 23 p. ().
- RANE, K. K., AND T. A. TATTAR. 1957. Pathogenicity of blue-stain fungi associated with *Dendroctonus terebrans*. *Plant Disease* 71: 579–583. (ec).
- *RAPP, OTTO. 1910. Zur Coleopterologischen Sammelreise nach Karnten. *Entomologische Blätter* 6: 75–81. ().
- RAFFAFORT, NANCY G., AND DAVID LEE WOOD. 1955a. Host relationships in the Douglas-fir twig mining beetle, *Pityophthorus orarius* Bright (Coleoptera: Scolytidae), in a northern California seed orchard. In G. E. Miller, Proceedings of the Third IUFO Seed and Cone Insects Working Conference, 26–29 June 1955. Victoria, British Columbia (cn hb).
- _____. 1955b. *Pityophthorus orarius* Bright (Coleoptera: Scolytidae) in a Douglas-fir seed orchard: effect of clonal source and cone crop on rate of attack. *International Congress of Entomology, Proceedings* 18: 431. (bv).
- RASKE, ARTHUR G., J. D. ROWE, AND H. O. SCHOOLEY. 1975. Preliminary report on an outbreak of the eastern larch beetle in Newfoundland. Government of Canada, Forestry Service, Newfoundland Forest Research Centre, St. John's, Newfoundland. File report: (unpublished) study 1–2. 22 p.
- RASMUSSEN, LYNX A. 1957. Mountain pine beetle selection of dwarf mistletoe and comandra blister rust infected lodgepole pine. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, Research Note INT-367. 3 p. (bv).
- RAUCH, P. A., B. EWING, DAVID LEE WOOD, AND WILLIAM DELLES BEDARD, JR. 1905. The information system for research and project management. Pages 420–428 in C. B. Huffaker, New technology in pest control. John Wiley & Sons, New York. (cn).
- RAUF, A., D. M. BENJAMIN, AND R. A. CECICH. 1955. Insects

- affecting seed production of jack pine and life tables of conelet and cone mortality in Wisconsin. *Forest Science* 31(2): 271–281. (cn).
- RAWN, HANS PETER. 1955. Expansion of the populations of *Ips typographus* (L.) (Coleoptera, Scolytidae) and their local dispersal following gale disaster in Denmark. *Journal of Applied Entomology* 99: 26–33. (bv cn ec).
- REDFERN, DEREK B. 1959. The roles of the bark beetle *Ips cembrae*, the woodwasp *Urocerus gigas* and associated fungi in dieback and death of larches. Pages 195–204 in N. Wilding, N. M. Collins, P. M. Hammond, and J. F. Bebbler. Insect-fungus interactions. Academic Press, London. xiv + 344 p. (ec).
- REDFERN, DEREK B., JOHN T. STOKLEY, II, STEELE, AND DAVID W. WINTER. 1957. Dieback and death of larch caused by *Ceratocystis laricicola* sp. nov. following attack by *Ips cembrae*. *Plant Pathology* 36: 467–480. (ec).
- REDLICH, HARTMUT, W. BURNS, W. FRANCKE, V. SCHURIG, THOMAS LEE PAYNE, JEAN PIERRE VITE. 1957. Chiral building units from carbohydrates. 13. Identification of the absolute configuration of endo-brevicommin from *Dendroctonus frontalis* and synthesis of both enantiomers from D-Ribose. *Tetrahedron* 43: 2029–2034. (ms).
- REED, A. N. F., AND J. W. HANOVER. 1955. Chemical factors in bark beetle preference among Douglas-fir and western larch. American Chemical Society (abstracts) No. 190. (bv).
- REED, A. N. F., J. W. HANOVER, AND MALCOLM MACFARLANE. 1956. Douglas-fir and western larch: chemical and physical properties in relation to Douglas-fir bark beetle attack. *Tree Physiology* 1(3): 277–287. (bv).
- REED, W. J., AND D. ERRICO. 1957. Techniques for assessing the effects of pest hazards on longrun timber supply. *Canadian Journal of Forest Research* 17: 1455–1465. (cn ms).
- REID, J. C., AND A. MANSINGH. 1955. Economic losses due to *Hypothenemus hampei* Ferr. during processing of coffee berries in Jamaica. *Tropical Pest Management* 31: 55–59. (cn).
- *REN, Z., AND X. DANG. 1959. The preliminary report on investigation and control of bark beetles attacking *Pinus armandii* [In Chinese]. Report of preliminary studies on forest insect pests. 234 p. ().
- *REN, Z., AND J. ZHOU. 1962. The outbreak reason and role of scolytids on *Pinus armandii* [In Chinese, Russian summary]. Shaanxi Province Forestry Society, Proceedings of annual forestry scientific meeting 1962: 1–19. ().
- *REN, Z., ET AL. 1965. The annual summary on integrated control of scolytids on *Pinus armandii*: a corpus of forestry scientific study references in Shaanxi Province [In Chinese]. Peoples Press of Shaanxi 1965: 157–164. ().
- RENWICK, JOHN ALAN ALEXANDER. 1958. Comparative mechanisms of host selection by insects attacking pine trees and crucifers. Pages 303–316 in K. C. Spencer. Chemical mediation of coevolution. Academic Press, San Diego. (bv).
- RIHODES, LLEWELLYN F., AND A. MANSINGH. 1956. Distribution of the coffee berry borer *Hypothenemus hampei* Ferr. in Jamaica and an assessment of the chemical control programme (1979–1952). *Insect Science and Its Application* 7: 505–510. (cn).
- RICHMOND, CHARLES E. 1955. Effectiveness of two pine oils for protecting lodgepole pine from attack by mountain pine beetle (Coleoptera: Scolytidae). *Canadian Entomologist* 117: 1445–1446. (cn).
- *RIEDL, HELMUT WOLFGANG. 1973. Aspects of the feeding behavior of *Scolytus multistriatus*, the primary vector of Dutch elm disease, and a critical evaluation of present chemical control measures. Unpublished dissertation, Michigan State University, East Lansing. 195 p. ().
- *RILEY, CHARLES VALENTINE, AND LELAND OSSIAN HOWARD. 1891. [*Scolytus rugulosus*]. *Insect Life* 3: 295. ().
- RILEY, M. A., AND RICHARD ALBRIGHT COYER. 1956. Impact of beneficial insects on *Ips* spp. (Coleoptera: Scolytidae) bark beetles in felled loblolly and slash pines in Louisiana. *Environmental Entomology* 15: 1220–1224. (ec).
- . 1958. Seasonal abundance of beneficial insects and *Ips* spp. engraver beetles (Coleoptera: Scolytidae) in felled loblolly and slash pines in Louisiana. *Journal of Entomological Science* 23: 357–365. (ec).
- *RINGS, ROY W. 1956. [*Phloeotribus liminaris*, *Scolytus rugulosus*]. Ohio State University, Bulletin 765: 44–45. ().
- RIO MORA, ADOLFO A. DEL. 1954. Insectos que afectan la producción de semilla en las especies de pinos de la Meseta Tarasca, Michoacan [Insects that affect seed production of the pine species of La Meseta Tarasca, Michoacan]. Secretaria de Agricultura y Recursos Hidraulicos, Instituto Nacional de Investigaciones Forestales, Proyecto de Protección Forestal del Centro de Investigaciones Forestales de Occidente, INIF, SF-SARH, Mexico, D. F. Boletín Técnico 97. 38 p. (ec hb).
- . 1955. Clave para la determinación de algunas especies de *Pityophthorus* (Coleoptera: Scolytidae) en pinos de la Meseta Tarasca, Michoacan [Identification key for some species of *Pityophthorus* on pines on La Meseta Tarasca] [English summary]. Secretaria de Agricultura y Recursos Hidraulicos, Instituto Nacional de Investigaciones Forestales, Mexico, Boletín Técnico 125. 18 p. (ds tx).
- RISBERG, BIRGER. 1955. Uppskattning av granbarkborreskad i Vaernland 1971–1982 genom korrigering och komplettering av tidigare flyginventeringsresultat [Estimation of *Ips typographus* damage in Vaernland 1971–1982 by correcting and supplementing earlier results]. Sveriges Skogvardsforbunds Tidskrift 2: 21–32. (cn).
- RITZENGRUBER, O., AND E. FUJIRER. 1956. Isoenzymanalyse verschiedener Populationen von *Pityogenes chalcographus* L. (Coleoptera, Scolytidae): I. Methoden-anpassung, Enzympolymorphismus [Isoenzyme analysis of various populations of *Pityogenes chalcographus*: I. Adaptation of methods, enzyme polymorphism] [English summary]. *Journal of Applied Entomology* 101: 187–194. (ay ms).
- ROBERTS, J. H., F. P. HAIN, AND L. B. NUNALLY. 1957. Genetic structure of southern pine beetle populations. *Forest Science* 33: 52–69. (ay).
- ROBERTS, HYWELL. 1956. New Platypodidae (Coleoptera) from the rain forests of Papua New Guinea. *Esakia* 24: 37–58. (tx).
- . 1957. Forest insect pests of Papua New Guinea: 2, pin-hole borers (shot hole borers) attacks on logs, lumber and living trees. *Harvest* 12: 91–96. (cn hb).

- . 1959. New *Crossotarsus* from Papua New Guinea (Platypodidae: Coleoptera). *Koleopterologische Rundschau* 59: 87–94. (tx).
- . 1992. Diapodini of Papua New Guinea (Platypodidae). In press [a Bishop Museum journal]. (tx).
- ROBERTS, HYWELL, AND K. MORIMOTO. 1957. New Platypodidae (Coleoptera) from the mountain forests of Papua New Guinea. *Esakia* 25: 161–181. (tx).
- ROBERTSON, F. DALE. 1957. Record of decision, USDA Forest Service, Suppression of the southern pine beetle: final environmental impact statement: deciding official, F. Dale Robertson. United States Department of Agriculture, Forest Service, Southern Region, Atlanta, Georgia. (cn).
- ROBERTSON, J. L. 1953. [Book review]. *Entomological Society of America, Bulletin* 29(2): 71. (ms).
- ROBINSON, M. N., JEAN CLAUDE GREGOIRE, AND L. DE VAS. 1955. A method of sexing live *Dendroctonus micans*. Pages 63–67 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Universite Libre de Bruxelles, Belgium, Proceedings of seminar, 3–4 October 1954. 141 p. (ay ms).
- RODOVALHO, BENTO DE TOLEDO. 1925b. Defendendo o cafe o repasse. *Rural, Revista da Sociedade Rural Brasileira* 6(59): 188–192. (cn).
- RODRIGUES MACHADO BENASSI, V. L., AND E. BERTI F. 1959. (Coleoptera, Scolytidae) in Espirito Santo State. Nota sobre a ocorrencia de *Cephalonomia* sp. (Hymenoptera, Bethyidae) parasitando a broca-do-cafe, *Hypothenemus hampei* (Ferrari, 1867) (Coleoptera, Scolytidae) no estado do Espirito Santo [Note on the occurrence of *Cephalonomia* sp. parasitizing the coffee berry borer, *Hypothenemus hampei*] [English subtitle]. *Revista de Agricultura (Piracicaba)* 64: 105–106. (ec).
- ROEDIGER, K. J. 1955. Ubervachung des Kupferstechers mit Chatoprax [Monitoring the spruce wood engraver with chalcoprax] [English summary]. *Gesunde Pflanzen* 40: 192–193. (cn hb).
- ROEFER, RICHARD A. 1955. Interaction between the ambrosia beetle *Corthylihus punctatissimus* and its mutualistic fungi. *International Congress of Entomology, Proceedings* 18: 410. (ec).
- ROEFER, RICHARD A., B. J. PALIK, D. V. ZESTOS, P. G. HESCH, AND C. D. LARSEN. 1957. Observations of the habits of *Corthylihus punctatissimus* (Coleoptera: Scolytidae) infesting maple saplings in central Michigan. *Great Lakes Entomologist* 20: 173–176. (cn hb).
- ROEFER, RICHARD A., D. V. ZESTOS, B. J. PALIK, AND LAWRENCE RICHARD KIRKENDALL. 1957. Distribution and host plants of *Corthylihus punctatissimus* (Coleoptera: Scolytidae) in the lower peninsula of Michigan. *Great Lakes Entomologist* 20: 69–70. (ds).
- ROGERS, TERRANCE J., AND P. F. HESSBURG. 1955. Annual southwestern region pest conditions report, 1954. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico, Forest Pest Management Report R3–55–9. 15 p. (cn).
- ROGERS, TERRANCE J., AND H. M. MAFFEI. 1957. Annual southwestern region pest conditions report 1956. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico, Forest Pest Management Report R3–57–4. 14 p. (cn).
- . 1958. Annual southwest region pest conditions report, 1957. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico, Forest Pest Management Report R3–58–2. 15 p. (cn).
- ROMERO CASADO, J. 1955. *Gossyparia ulmi* Geoffroy (Homoptera: Eriococcidae) una causa mas de debilitamiento de los olmos. Estudio morfologico y bionomico [*Gossyparia ulmi*, one more cause of weakening the elms: studies on morphology and bionomics]. *Boletin del Servicio de Defensa Contra Plagas e Inspeccion Fitopatologica* 11(1): 45–55. (cn).
- ROMIE, W. H., D. H. KNIGHT, AND J. B. YAVITT. 1956. Mountain pine beetle outbreaks in the Rocky Mountains: regulators of primary productivity. *American Naturalist* 127: 484–494. (ec).
- *ROSENFELD. 1910. [*Scolytus rugulosus*]. *Journal of Economic Entomology* 3: 215. (j).
- ROTON, LARY M. 1957. Promising treatment for southern pine beetles. *American Papermaker* (October): 30–32. (cn).
- . 1959. Bark beetle treatment: U. S. Patent 4,524,665. Official Gazetteer, United States Patent Trademark Office 110(4): 2602. (cn ms).
- *ROUBAL, JAN. 1940. Coleoptera v pude karpatskeho smiseného lesa. Quae Coleoptera in hino saltuum carpathicorum vivunt? *Vestnik C. Zool. Spolec. Prave* 8: 97–130. (j).
- ROY, D. N., J. R. PURDY, P. A. PERUMAL, AND J. K. GRACE. 1955. Effect of application equipment on the distribution of chlorpyrifos applied for Dutch elm disease vector control. *Entomological Society of Ontario, Proceedings* 119: 63–65. (cn).
- RUHM, WALTER. 1956. Zwei im Xylem der Araukarie, *Araucaria araucana* (Mol.) Koch brutende Xylomyetophage und koprophage Borkenkaferarten (Col.: Scolytidae) [English summary]. *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg* 8(128): 265–280. (bv ec hb).
- RUMANN, J. 1956. Kalkschutz gegen Nutzholzborkenkafer [Use of lime for protection against *Xyloterus lineatus*]. *Holz-Zentralblatt* 112(47): 661–662. (cn).
- RUSH, PETER A. 1956. Forest pest conditions report for the Northeastern Area, 1955. United States Department of Agriculture, Forest Service, Northeastern Area (Broomall, Pennsylvania), Forest Pest Management Report NA-FR-33. 35 p. (cn).
- RUSH, PETER A., T. A. LAURENT, L. C. YARGER, AND R. K. LAWRENCE. 1977. Forest insect and disease conditions in Alaska, 1976. United States Department of Agriculture, Forest Service, Alaska Region, Juneau, Alaska. 12 p. (cn).
- *RUTHERFORD, A. 1914d. Some insect pests of 1913 [*Xyleborus compactus*]. *Ceylon Department of Agriculture, Bulletin* 15: 2. 7. (j).
- RUTLEDGE, D. N., CONSTANTIN CHARARAS, AND C. J. DUCAUZE. 1955. Investigating the hypothesis of primary attraction in conifer-specific Scolytidae using a chemometrical approach. Pages 391–409 in W. J. Mattson, J. Leveux, and C. Bernard-Dagen, Mechanisms of woody plant defenses against insects: search for patterns. Springer, New York. xiv + 416 p. (bv).
- RYKER, LEE C. 1955. Acoustic studies of *Dendroctonus* bark beetles. *Florida Entomologist* 71: 447–461. (bv).
- RYKIEL, E. J., JR., ROBERT N. COUSLON, P. J. H. SHARPE, T. F. H. ALLEN, AND R. O. FLAMM. 1955. Disturbance propagation by bark beetles as an episodic landscape phenomenon. *Landscape Ecology* 1: 129–140. (cn).

S

- SAARENMAA, HANNU. 1953. Modeling the spatial pattern and interspecific competition in *Tomicus piniperda* (Coleoptera, Scolytidae). Communications Instituti Forestalis Fenniae 118, 40 p. (en ec).
- _____. 1955a. The role of temperature in the population dynamics of *Tomicus piniperda* (L.) (Col., Scolytidae) in northern conditions. Journal of Applied Entomology 99: 224-236. (ec hb).
- _____. 1955b. Within-tree population dynamics models for integrated management of *Tomicus piniperda* (Coleoptera, Scolytidae). Communications Instituti Forestalis Fenniae 128, 56 p. (en ec hb).
- _____. 1957. Tuhohyonteisten ja sinistymän esiintyminen myrskyn kauttamissa puissa Lapissa 1953-56 [Insect attack and blue stain in windthrown trees in Lapland in 1953-56] [In Finnish, English summary]. Folia Forestalia [Helsinki] 696, 15 p. (bv ec).
- _____. 1955a. Attack and reproduction of bark beetles in windthrown pine in northern Finland. Pages 69-75 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, 355 p. (bv hb).
- _____. 1955b. Attack and reproduction of bark beetles in wind-blown pine in northern Finland. International Congress of Entomology, Proceedings 18: 412. (bv).
- _____. 1959. A model for the timing of swarming of *Tomicus piniperda* (Coleoptera: Scolytidae). Holarctic Ecology 12: 441-444. (hb).
- SAFRANYIK, LASZLO. 1955a. Infestation incidence and mortality in white spruce stands by *Dendroctonus rufipennis* Kirby (Coleoptera, Scolytidae) in central British Columbia. Journal of Applied Entomology 99: 86-93. (en).
- _____. 1955b. The role of the host in the population dynamics of forest insects. IUFRO conference proceedings. Government of Canada, Forestry Service and United States Department of Agriculture, Forest Service, 240 p. (bv ec hb).
- _____. 1955c. The relationship between density of emerged *Dendroctonus ponderosae* (Coleoptera: Scolytidae) and density of exit holes in lodgepole pine. Canadian Entomologist 117: 267-275. (hb).
- _____. 1955a. Estimating attack and brood totals and densities of the mountain pine beetle in individual lodgepole pine trees. Canadian Entomologist 120: 323-331. (en ms).
- _____. 1955b. The population biology of the spruce beetle in western Canada and implications for management. Pages 3-23 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia, 355 p. (en hb).
- _____. 1955c. The role of population biology in development of management guidelines to reduce losses from the spruce beetle *Dendroctonus rufipennis* in western Canada. International Congress of Entomology, Proceedings 18: 412. (en).
- _____. 1959b. Mountain pine beetle: biology overview. Pages 9-13 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262, 119 p. (bv ec hb).
- SAFRANYIK, LASZLO, AND D. A. LINTON. 1955. Influence of competition on size, brood production and sex ratio in spruce beetles (Coleoptera: Scolytidae). Entomological Society of British Columbia, Journal 52: 52-56. (ec hb).
- _____. 1957a. Line intersect sampling for the density and bark area of logging residue susceptible to the spruce beetle, *Dendroctonus rufipennis* (Kirby). Government of Canada, Canadian Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-295, 10 p. (en).
- _____. 1957b. Patterns of landing of spruce beetles, *Dendroctonus rufipennis* (Coleoptera: Scolytidae), on baited lethal trap trees. Entomological Society of British Columbia, Journal 54: 21-32. (ec hb).
- _____. 1955. Distribution of attacks on spruce stumps by the spruce beetle, *Dendroctonus rufipennis* (Kirby) (Coleoptera: Scolytidae), and effects on the length of egg galleries. Canadian Entomologist 120: 85-94. (bv hb).
- SAFRANYIK, LASZLO, R. SILVERSIDES, L. H. McMULLEN, AND D. A. LINTON. 1959. An empirical approach to modeling the local dispersal of the mountain pine beetle (*Dendroctonus ponderosae* Hopk.) (Col., Scolytidae) in relation to source of attraction, wind direction and speed. Journal of Applied Entomology 105: 495-511. (bv hb).
- SAFRANYIK, LASZLO, AND H. S. WHITNEY. 1955. Development and survival of axenically reared mountain pine beetles, *Dendroctonus ponderosae* (Coleoptera: Scolytidae), at constant temperatures. Canadian Entomologist 117: 185-192. (ec hb).
- SAHA, NIVEDITA, AND P. K. MAITI. 1954. On a collection of scolytid beetles (Scolytidae: Coleoptera) from Sikkim, India. Record of the Zoological Survey of India 51(3-4): 1-5. (ds).
- _____. 1957. Description of hitherto unknown males of three species of scolytid beetles (Scolytidae: Coleoptera) from India. Bulletin of the Zoological Survey of India 5(1-3): 71-76. (ds tx).
- SAIHOTA, T. S., AND S. H. FARRIS. 1950. Inhibition of flight muscle degeneration by precocene II in the spruce bark beetle, *Dendroctonus rufipennis* (Kirby) (Coleoptera: Scolytidae). Canadian Journal of Zoology 55: 378-381. (ay).
- SAIHOTA, T. S., AND F. G. PEET. 1955a. Changes in the quality of scolytid populations: their determination and relation to scolytid outbreaks. Pages 25-33 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia, 355 p. (hb).
- _____. 1955c. Computer-assisted measurement and analysis of chromatin distribution for determining quality differences among bark beetle (Scolytidae) populations. Pages 171-180 in In paths from a viewpoint: the Wellington festschrift on insect ecology. Memoir of the Entomological Society of Canada 146. (en ec).
- SAIHOTA, T. S., F. G. PEET, AND I. IBARAKI. 1957. Manipulations of egg-gallery length to vary brood density in spruce beetle *Dendroctonus rufipennis* (Coleoptera: Scolytidae): effects on brood survival and quality. Entomological Society of British Columbia 54: 59-63. (hb).
- SAJAN, R. J. 1951. Results of forest insect and disease surveys in the eastern region of Ontario, 1950. Page 12 in Canada Department of Environment, Forestry Service, Great Lakes Forest Research Centre, Sault Ste. Marie, Ontario. (en).
- SALAZAR, M. A. 1952. Historia del *Dendroctonus frontalis* Zimm. en Honduras (Coleoptera: Scolytidae). Pages

- 222–232 in *Actas de las IV Jornadas de Reforestación, Corporación Hondureña de Desarrollo Forestal, Tegucigalpa, Honduras*. (cn).
- SALDARRIAGA VELEZ, A. 1985. El *Pagiocerus frontalis* (F.), plaga del maíz almacenado: biología hábitos y notas ecológicas [*Pagiocerus frontalis*, a pest of stored maize: biology, behavior and ecological notes] [English summary]. *Revista Colombiana de Entomología* 10(3–4): 9–14. (ay bv lb).
- SALOM, SCOTT M., AND JOHN ALEXANDER McLEAN. 1988. Semiochemicals for capturing the ambrosia beetle, *Trypodendron lineatum*, in multiple-funnel traps in British Columbia. *Entomological Society of British Columbia, Journal* 85: 34–39. (bv cn).
- . 1989. Influence of wind on the spring flight of *Trypodendron lineatum* (Olivier) (Coleoptera: Scolytidae) in a second-growth coniferous forest. *Canadian Entomologist* 121: 109–119. (ec lb).
- SAMSON, P. R. 1984. The biology of *Roptrococcus xylophagorum* (Hym., Torymidae), with a note on its taxonomic status. *Entomophaga* 29: 287–298. (ec).
- SAMSON, P. R., AND J. SMIBERT. 1986. Preliminary studies on the efficacy and establishment of *Roptrococcus xylophagorum* (Hym.: Torymidae), a parasitoid of *Ips grandicollis* (Col.: Scolytidae), in Australia. *Entomophaga* 31: 173–182. (ec).
- SANCHEZ, LUIS A. GIL, AND JUAN A. PJARES ALONSO. 1986. Los escolitidos de las coníferas en la península ibérica. Instituto Nacional de Investigaciones Agrarias, Madrid, Spain. 194 p. (cn ds).
- SANDERS, W. 1987a. Ein beitrage zur habitatwahl der nutzholzborkenkäfer *Trypodendron lineatum* und *T. domesticum* [A contribution to the habitat selection of the ambrosia beetles *Trypodendron lineatum* and *T. domesticum*] [English summary]. *Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz* 60: 65–67. (lb).
- . 1987b. Untersuchungen über die aktivitätsdichte des buckdrückers *Ips typographus* in laubwäldern und in offener landschaft. [Studies on the activity of the engraver beetle *Ips typographus* in deciduous forests and in the open field] [English summary]. *Journal of Applied Entomology* 103: 240–249. (lb).
- *SANDERSON. 1912. [*Scolytus rugulosus*]. *Insect pests*, edition 1: 544. ().
- *———. 1921. [*Phloeotribus liminaris*, *Scolytus rugulosus*]. *Insect pests*, edition 2: 477, 582. ().
- *———. 1931. [*Scolytus rugulosus*]. *Insect pests*, edition 3: 300–302. ().
- SANTIS, L. DE, AND A. E. C. DE SUREDA. 1986. Una subfamilia de Chalcidoideos nueva para la Republica Argentina (Insecta, Hymenoptera) [A subfamily of Chalcidoidea new for Argentina] [English summary]. *Revista de Investigación, Centro de Investigaciones para la Regulación de Poblaciones de Organismos Nocivos, Argentina* 4(1–4): 63–66. (ec).
- SARTWELL, CHARLES, JR. 1970. *Ips pini* attack density in ponderosa pine slash as related to felling date in eastern Washington. United States Department of Agriculture, Forest Service, Research Note PNW-131, Portland, Oregon. 8 p. (cn ec).
- . 1988. Thinning *Pinus ponderosa* for silvicultural control of *Dendroctonus ponderosae* (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 18: 441. (cn).
- SASAKAWA MITSUHIRO, AND Y. KAWAGUCHI. 1987. Initial attack of the minute pine bark beetle, *Cryphalus fulvus* Niisima (Coleoptera: Scolytidae). *Scientific Reports of the Kyoto Prefectural University, Agriculture* 39: 12–19. (bv).
- SASAKAWA, MITSUHIRO, AND Y. YOSHIYASU. 1983. Stridulatory organs of the Japanese pine bark beetles (Coleoptera: Scolytidae). *Kontyu* 51: 493–501. (ay).
- SAUNDERS, M. C., P. K. LOH, ROBERT N. COULSON, E. J. RYKIEL, THOMAS LEE PAYNE, P. E. PULLEY, AND L. C. HU. 1985. Development and implementation of the southern pine beetle decision support system. Pages 335–364 in S. J. Bramham and R. C. Thatchner, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- *SAUVARD, D. 1988. Capacités de multiplication de *Tomicus piniperda* L. (Coleoptera: Scolytidae) selon la densité d'attaque et diverse caractéristiques de l'arbre [Multiplication capacity of *Tomicus piniperda* according to the density of attack and various characteristics of the tree] [English summary]. Unpublished dissertation, University of Orleans, Orleans, France. 88 p. ().
- . 1989. Capacités de multiplication de *Tomicus piniperda* L. (Col., Scolytidae): 1. effets de la densité d'attaque [Reproductive capacity of *Tomicus piniperda*: 1. effects of attack density] [English summary]. *Journal of Applied Entomology* 105: 164–181. (ec lb).
- SAUVARD, D., FRANÇOIS LIEUTIER, AND JEAN LÉVÉQUE. 1987. Répartition spatiale et dispersion de *Tomicus piniperda* L. (Coleoptera: Scolytidae) en forêt d'Orleans, France [English summary]. *Annales des Sciences Forestières* 44: 417–434. (lb).
- . 1988. L'Hylesine du pin (*Tomicus piniperda* L.) en forêt d'Orleans: répartition-dégâts-lutte [*Tomicus piniperda* in the forest of Orleans, France: distribution, damage, control]. *Revue Forestière Française* 40: 13–19. (cn).
- SCHAUFF, M. E. 1988. The species of *Entedon* in America north of Mexico (Hymenoptera: Eulophidae). *New York Entomological Society, Journal* 96: 30–62. (ec).
- *SCHEDL, KARL EDUARD. 1949. Nachtrag zur Gesamtliteratur der Borkenkäfer (Ipidae und Platypodidae) von Kleine 1939. *Zentralblatt für das Gesamtgebiet der Entomologie* 3: 63–107. ().
- *SCHEERFELTZ, OTTO, AND ALBERT WINKLER. 1930. [*Hylurgops synonymus*]. *Tierwelt Mitteilungen* 1930: 26, 256. ().
- SCHMITSCHKE, ERVIN. 1939d. Über Forstschadlingssauftreten in der Türkei. *International Congress of Entomology, Proceedings* 7: 2104–2131. (cn).
- SCHILUP, J. 1987. Untersuchungen zur populationsdynamik von *Ips typographus* L. und zur Biologie seines häufigsten Prädatoren *Thanasimus formicarius* L. Unpublished dissertation, University of Bern, Switzerland. (ec lb).
- SCHILATER, FREDRIK. 1985. Attack density regulation in *Ips typographus*: natural mechanisms and manipulation for management. *International Congress of Entomology, Proceedings* 18: 412. (bv cn).
- SCHILATER, FREDRIK, AND O. ANDERBRANT. 1987. Dutch elm disease (*Ceratocystis ulmi*) and elm bark beetles (*Scolytus* spp.) in Malmö 1985: distribution, phenology and practical measures in an integrated control program. *Naxtskyddsnotiser* 51: 2–10. (cn lb ds).
- . 1988. Pheromone production and response governing attack dynamics in *Ips typographus*. *Internation-*

- tional Congress of Entomology, Proceedings 18: 409. (bv).
- . 1959. Mass attack on trees by *Ips typographus* induced by sex-specific pheromone: a model of attack dynamics. *Holarctic Ecology* 12: 415–426. (bv).
- SCHLYTER, FREDRIK, O. ANDERBRANT, G. LINDQUIST, AND A. JANSSON. 1957. Ahmsjuka (*Ceratocystis ulmi*) och ahmsplintborrar (*Scolytus* spp.) i Malmö 1955: förekomst, fenologi och praktiska åtgärder inom ett integrerat kontrollprogram [Dutch elm disease (*Ceratocystis ulmi*) and elm bark beetles (*Scolytus* spp.) in Malmö 1955: distribution, phenology and practical measures in an integrated control program] [In Swedish, English summary]. *Växtskyddsnotiser* 51: 2–10. (ec).
- SCHLYTER, FREDRIK, AND GÖRAN BIRGERSSON. 1959. Individual variation in bark beetle and moth pheromones: a comparison and an evolutionary background. *Holarctic Ecology* 12(4): 457–465. (bv).
- SCHLYTER, FREDRIK, GÖRAN BIRGERSSON, J. A. BYERS, J. LOFQVIST, AND G. BERGSTROM. 1957. Field response of spruce bark beetle, *Ips typographus*, to aggregation pheromone candidates. *Journal of Chemical Ecology* 13: 701–716. (bv).
- SCHLYTER, FREDRIK, GÖRAN BIRGERSSON, AND A. LEUFVEN. 1959. Inhibition of attraction to aggregation pheromone by verbenone and ipsenol: density regulation mechanisms in bark beetle *Ips typographus*. *Journal of Chemical Ecology* 15: 2263–2277. (bv).
- SCHLYTER, FREDRIK, J. A. BYERS, AND J. LOFQVIST. 1957. Attraction to pheromone sources of different quantity, quality, and spacing: density-regulation mechanisms in bark beetle *Ips typographus*. *Journal of Chemical Ecology* 13: 1503–1523. (bv).
- SCHLYTER, FREDRIK, J. A. BYERS, J. LOFQVIST, A. LEUFVEN, AND GÖRAN BIRGERSSON. 1955. Reduction of attack density of the bark beetles *Ips typographus* and *Tomicus piniperda* on host bark by verbenone inhibition of attraction to pheromone and host kairomone. Pages 53–68 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv).
- SCHLYTER, FREDRIK, AND J. LOFQVIST. 1956. Response of walking spruce bark beetles *Ips typographus* to pheromone produced in different attack phases. *Entomologia Experimentalis et Applicata* 41: 219–230. (bv).
- SCHLYTER, FREDRIK, J. LOFQVIST, AND J. A. BYERS. 1957. Behavioural sequence in the attraction of the bark beetle *Ips typographus* to pheromone sources. *Physiological Entomology* 12: 185–196. (bv).
- SCHMID, JOHN MICHAEL. 1957. Partial cutting in MPB [mountain pine beetle]-susceptible pine stands: will it work and for how long? United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado, General Technical Report RM-149: 243–245. (cn).
- SCHMID, JOHN MICHAEL, S. A. MATA, AND R. D. AVERILL. 1959. Containment of small infestations of the mountain pine beetle in ponderosa pine. United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado, Research Note RM-493. 4 p. (cn).
- SCHMID, JOHN MICHAEL, S. A. MATA, AND WILLIAM FRANCIS MCCAMBRIDGE. 1955. Natural falling of beetle-killed ponderosa pine. United States Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colorado, Research Note RM-454. 3 p. (cn).
- SCHMID, JOHN MICHAEL, J. C. MITCHELL, AND S. A. MATA. 1956. Ponderosa pine conelet and cone mortality in central Arizona. *Great Basin Naturalist* 46: 445–448. (cn).
- SCHMITT, JEFFREY J., T. EVAN NEBEKER, C. A. BLANCHIE, AND JOHN DEAVOURS HODGES. 1955. Physical properties and monoterpene composition of xylem oleoresin along the bole of *Pinus taeda* in relation to southern pine beetle attack distribution. *Canadian Journal of Botany* 66: 156–160. (bv).
- SCHMITZ, RICHARD FRANKLIN. 1955a. Understanding scolytid problems in lodgepole pine forests: the need for an integrated approach. Pages 231–245 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (bv cn ec hb).
- . 1955b. Understanding scolytid problems in lodgepole pine forests: the need for an integrated approach. *International Congress of Entomology, Proceedings* 18: 414. (bv cn ec hb).
- . 1959. Efficacy of verbenone for preventing infestation of high-value lodgepole pine stands by the mountain pine beetle. Pages 75–80 in G. D. Amman, *Proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle: symposium proceedings*. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- SCHMITZ, RICHARD FRANKLIN, MARK D. MCGREGOR, GENE DOYLE AMMAN, AND ROBERT D. OAKES. 1959. Effect of partial cutting treatments of lodgepole pine stands on the abundance and behavior of flying mountain pine beetles. *Canadian Journal of Forest Research* 19: 566–574. (cn ec hb).
- *SCHMUTZENHOFER, HEINRICH. 1955a. Insektenspuren an Berindetem Nadelholz [Signs of insect damage in unbarked softwood]. Österreichische Agrarverlag, Wien, Austria. 166 p. (i).
- . 1955b. The use of pheromones for insect pest control methods, part II: aggregation pheromones. Pages 209–213 in *Protection of forests in the tropics: noxious insects to pine and eucalypt plantations in the tropics*. Protecáo de Universidade Federal do Paraná, Curitiba [Brazil]. (bv cn).
- . 1955a. Mass outbreaks of *Ips* bark beetles in Bhutan and the revision of the genus *Ips* DeGeer for the Himalayan region. Pages 345–355 in T. L. Payne and H. Saaremaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (ay cn hb tx).
- . 1955b. Mass outbreaks of *Ips* bark beetles in Bhutan and the revision of the genus *Ips* for the Himalayan region. *International Congress of Entomology, Proceedings* 18: 416. (ay cn hb tx).
- . 1955c. Zum Nachweis der Gattung *Trypodendron* im Himalaja (Coleoptera, Scolytidae). *Entomologica Basiliensia* 12: 487–490. (tx).
- SCHNEIDER, ISOLDE A. 1965c. Frau Professor Dr. Helene Francke-Grosman zum 65. Geburtstag. *Journal of Applied Entomology* 55: 301–303. (ms).
- . 1955a. A new locality in northwestern Germany for *Cnathotrichus materiarius* (Coleoptera: Scolytidae) introduced from North America. *Neue Entomol. Nachr.* 17: 39–42. (ds).

- _____. 1955b. *Gnathotrichus materiarius* Fitch (Col., Scolytidae) in Pheromonfallen von *Ips cembrae* (Heer) (Col., Scolytidae), ein neuer Fundort für NW-Deutschland. Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz 58: 50–51. (ds).
- _____. 1957. Verbreitung, Pilzübertragung und Brutsystem des Ambrosiakäfers *Xyleborus affinis* im Vergleich mit *X. mascarensis* (Coleoptera: Scolytidae) [Distribution, fungus transfer and brood system of the ambrosia beetle *Xyleborus affinis* in comparison with *X. mascarensis* (Coleoptera: Scolytidae)] [English summary]. Entomologia Generalis 12: 267–275. (ec hb ds).
- *SCHOENHERR, J. 1955. Systematic review on Neotropical bark beetles. Contribuição a taxonomia e ecologia dos escolitídeos do Brasil. In Proteção de florestas nos trópicos: insectos nocivos ao plantio de *Pinus* e *Eucalyptus* nos trópicos. Universidade Federal do Paraná, Curitiba [Brazil]. ().
- *SCHOOL, 1922. [*Scolytus rugulosus*]. Texas Department of Agriculture, Bulletin 72: 76. ().
- SCHOMAKER, MICHAEL L. 1951. Colorado forest pest index. Colorado State Forest Service, Insect and Disease Division, Colorado State University, Fort Collins. 57 p. (cn).
- SCHOPF, A. 1955. Zum Einfluss der Photoperiode auf die Entwicklung und Kalteresistenz des Buchdruckers, *Ips typographus* L. (Col., Scolytidae). Anzeiger für Schädlingkunde Pflanzen- und Umweltschutz 58: 73–75. (bv hb).
- _____. 1959. Die Wirkung der Photoperiode auf die Induktion der Imaginaldiapause von *Ips typographus* (L.) (Col., Scolytidae) [The effect of photoperiod on the induction of the imaginal diapause of *Ips typographus* (L.)] [English summary]. Journal of Applied Entomology 107: 275–288. (ec hb).
- SCHOPF, R. 1957. Zur Frage der Bedeutung von Tieren als Vektoren Pflanzenpathogener Mikroorganismen in Wirtschaftswäldern [On the importance of animals as vectors of plant pathogenic microorganisms in forest ecosystems] [English, French summaries]. Allgemeine Forst- und Jagdzeitung 155: 7–11. (ec).
- *SCHÖYEN, THOB. HIORTH. 1914. [*Hylastes ater*]. Tidskrift for Skogbruk, Kristiania 1914: 448–449. ().
- SCHREIBER, LAWRENCE R., EDWARD E. CONWAY, AND JOHN W. PEACOCK. 1956. Aggressiveness, competitiveness and stability of tolerance of benzimidazole-tolerant strains of *Ceratocystis ulmi*. Plant Disease 70: 154–158. (ec).
- SCHREIBER, LAWRENCE R., AND JOHN W. PEACOCK. 1975. Dutch elm disease and its control. United States Department of Agriculture, Forest Service, Agriculture Information Bulletin 193. 14 p. (revised 1979). (cn).
- *_____. 1955. Competitiveness of benzimidazole-tolerant and sensitive strain of *Ceratocystis ulmi* [abstract]. Phytopathology 75: 1308. (cn).
- SCHROEDER, LEIF MARTIN. 1957. Attraction of the bark beetle *Tomicus piniperda* to Scots pine trees in relation to tree vigor and attack density. Entomologia Experimentalis et Applicata 44: 53–58. (bv cn).
- _____. 1958a. Attraction of the bark beetle *Tomicus piniperda* and some other bark- and wood-living beetles to the host volatiles of alpha-pinene and ethanol. Entomologia Experimentalis et Applicata 46: 203–210. (bv).
- _____. 1958b. Host finding and colonization in the bark beetle *Tomicus piniperda* (L.) (Coleoptera: Scolytidae). International Congress of Entomology, Proceedings 18: 441. (bv).
- *_____. 1958c. Host recognition in *Tomicus piniperda* (Coleoptera: Scolytidae) and other bark beetles attacking Scots pine. Published dissertation, Swedish University of Agricultural Science, Sweden. Printed at SLU, Uppsala, ISBN 91–576–3363–0. 29 p. ().
- SCHROEDER, LEIF MARTIN, AND HUBERTUS H. EIDMANN. 1956. The effects of pure and blended atmospheric gases on the survival of three bark beetle species. Journal of Applied Entomology 101: 353–359. (ec).
- _____. 1957. Gallery initiation by *Tomicus piniperda* (Coleoptera: Scolytidae) on Scots pine trees baited with host volatiles. Journal of Chemical Ecology 13: 1591–1599. (bv).
- SCHROEDER, LEIF MARTIN, AND A. LINDELOW. 1959. Attraction of scolytids and associated beetles by different absolute amounts and proportions of alpha pinene and ethanol. Journal of Chemical Ecology 15: 807–818. (bv).
- SCHROEDER, LEIF MARTIN, AND B. RISBERG. 1959. Establishment of a new brood in *Tomicus piniperda* L. (Coleoptera, Scolytidae) after a second hibernation. Journal of Applied Entomology 108: 27–34. (hb).
- SCHUBERT, C. 1955. Die Aktivitätsdynamik der peroxidase (EC 1.11.1.7) aus dem Kupferstecher (*Pityogenes chalcographus*) (Coleoptera, Scolytidae) und dem Bast der Europäischen Fichte (*Picea abies* [L.] Karsten) [Changes in peroxidase activity in the bark beetle *Pityogenes chalcographus* and the phloem of European spruce]. Zoologischer Anzeiger 221: 355–367. (ay).
- SCHULP, J. 1957. Untersuchungen zur Populationsdynamik von *Ips typographus* L. und zur Biologie seines häufigsten Praedators *Thanasimus formicarius* L. Unpublished dissertation, University of Bern, Switzerland. (hb ec).
- SCHULTZ, DAVID E., AND J. ALLISON. 1951. A biological evaluation of existing and potential forest insect and disease problems in Connel Camp, Lobo-oso Group Camp, and snag-thinning study plots, San Bernardino National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 51–15. (cn).
- _____. 1952. Tree mortality on the Cleveland National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 52–12. (cn).
- SCHULTZ, DAVID E., AND WILLIAM DELLES BEDARD. 1957. California five-spined Ips. United States Department of Agriculture, Forest Service, Insect and Disease Leaflet 102. 8 p. (bv cn hb).
- SCHULTZ, DAVID E., G. DENITTO, AND J. ALLISON. 1951. Biological evaluation of the Upper Boulder Sale, Greenville Ranger District, Plumas National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 51–23. (cn).
- SCHULTZ, DAVID E., AND J. KLIEJUNAS. 1951. Evaluation of true fir twig mortality on the Goosenest Ranger District, Klamath National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 51–34. (cn).
- _____. 1953. Evaluation of Medicine Lake Campgrounds,

- Doublehead Ranger District, Modoc National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-34. (cn).
- . 1954. Evaluation of mortality in Sugar Pine Reservoir Campground, Foresthill Ranger District, Tahoe National Forest, United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-06. (cn).
- SCHULTZ, DAVID E., AND J. PRONOS. 1982. Evaluation of Jeffrey pine mortality at Mineral King, Sequoia Kings Canyon National Parks, United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-34. (cn).
- SCHULTZ, DAVID E., AND B. ROETTERING. 1984. A biological evaluation of lodgepole pine mortality in Parks Creek Drainage, Mt. Shasta Ranger District, Shasta-Trinity National Forest, United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-26. (cn).
- *SCHULTZE, W. 1915. A catalogue of Philippine Coleoptera. Department of the Interior, Bureau of Science, Manila 7: 1-198. ().
- SCHURIG, V., U. LEYFER, AND U. KOHNLE. 1955. Enantiomer composition and absolute configuration of terpen-4-ol from the bark beetle *Polygraphus poligraphus*. *Naturwissenschaften* 72(4): 211 p. (bv ms).
- SCHURING, W. 1951. Aantastingen door insecten en mijten op bomen en struiken in 1950. *Nederlands Bosbouw Tijdschrift* 53: 173-183. (cn).
- . 1952. Aantastingen door insecten en mijten op bomen en struiken in 1951. *Nederlands Bosbouw-tijdschrift* 54: 124-133. (cn).
- . 1953. Aantastingen door insecten en mijten op bomen en struiken in 1952. *Nederlands Bosbouw-tijdschrift* 55: 124-133 [?]. (cn).
- . 1954. Aantastingen door insecten en mijten op bomen en struiken in 1953. *Nederlands Bosbouw-tijdschrift* 56: 156-195. (cn).
- . 1955. Aantastingen door insecten en mijten op bomen en struiken in 1954. *Nederlands Bosbouw-tijdschrift* 57: 367-374. (cn).
- . 1956. Aantastingen door insecten en mijten op bomen en struiken in 1955. *Nederlands Bosbouw-tijdschrift* 58: 363-369. (cn).
- SCHIVESTER, DANIEL. 1951b. Methodes de lutte contre les scolytides des arbres fruitiers. *Revue de Zoologie Agricole et Appliquee* 1951(2e trimestre, nos. 4-6): 3-7. (cn).
- . 1953. Sur le cycle evolutif de *Xyleborus saxeseni* Ratz. (Coleoptera, Scolytidae). *Revue de Zoologie Agricole et Appliquee* 1953 (4e trimestre, nos. 10-12): 1-2 (unpaginated). (cn).
- . 1970. Desarrollos recientes en el conocimiento de los insectos devastadores del pino maritimo en zona Mediterranea Francesa. *Boletin del Servicio de Plagas Forestales* 13(26): 173-180. (cn).
- . 1955. Les insectes et la foret francaise. *Revue Forestiere Francaise* (Nancy) 37(special issue): 45-64. (cn).
- . 1956. Problemes entomologiques forestiers en zone mediterrannee francaise. *European and Mediter-ranean Plant Protection Organization, Bulletin* 16: 603-612. (cn).
- SCHWANDT, JOHN W., ROBERT LADD LIVINGSTON, D. P. BECKMAN R. L. JAMES S. TUNNOCK, AND K. A. KNAPP. 1986. Idaho forest pest conditions and program summary. 1985. United States Department of Agriculture, Forest Service, Northern and Intermountain Regions and Idaho Department of Lands, Coeur d'Alene, Idaho, Report 86-1. 25 p. (cn).
- SCHWARZ, MARIE BEATRICE. 1924. *Botrytis stephanoderis* n. sp. Bally and *Botrytis bassiana* Bals. *Bulletin du Jardin botanique de Buitenzorg* 3(6): 68-69. (cc).
- SCHWENKE, WOLFGANG. 1955. Beziehungen zwischen tierischen Schadlingen und Baumkrankungen [Relationships between animal pests and tree diseases]. *Forestwissenschaftliches Centralblatt* 104: 220-225. (cn).
- SCHWERTFEGER, FRITZ. 1951. Die Waldkrankheiten. Edition 4 [Scolytidae, p. 152-197]. Paul Parey, Berlin. 486 p., 242 figs. (bv ds hb).
- SCHWERT, DONALD P., T. W. ANDERSON, ANNE MORGAN, ALAN V. MORGAN, AND P. F. KARROWS. 1955. Changes in late Quaternary vegetation and insect communities in southwestern Ontario. *Quaternary Research* 23: 205-226. (ds).
- SCHWERT, DONALD P., AND ALAN V. MORGAN. 1908. Paleo-environmental implications of a late glacial insect assemblage from northwestern New York. *Quaternary Research* 13: 93-110. (ds).
- SEDLACZEK, WALTER. 1904. Review of: Keller, Untersuchungen über der Hohenverbreitung forstschadlicher Tiere in der Schweiz. *Zentralblatt für das Forstwesen* 30: 357-360. (ms).
- SEKENDIZ, O. A. 1957. *Pityokteines curvidens* (Germ.) (Coleoptera, Scolytidae) in dogu karadeniz orman-larinda, dogu goknari (*Abies nordmanniana* Stev. Spach) uzer unde ki zararları ve biyolojisi [On the damage and biology of *Pityokteines curvidens* on Nordmann fir *Abies nordmanniana* in the eastern Black Sea forest region] [In Turkish, English summary]. *Türkiye 1. Entomoloji Kongresi Bildirileri*, 13-16 Ekim 1957. Ege Üniversitesi, Nornova, Izmir. Bornova/Izmir, Turkey. Ege Üniversitesi Atatürk Kultur Merkezi 1957: 209-218. (cn ec hb).
- *SELANDER, J. 1956. Seasonal life history and economic importance of the major timber beetles of the Zambian teak forest. Pages 257-301 in G. D. Pierce, The Zambezi teak forests. 1st International Conference on the Teak Forests of Southern Africa, proceedings. Livingstone, Zambia, 18-24 March 1954. Forest Department, Ndola, Zambia. ().
- SELLENSCHLO, U. 1956. Beifänge in Borkenkafer-Pheromonfallen [Non-target captures in bark beetle pheromone traps] [English summary]. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 6: 371-372. (cc).
- *SELM, E. 1957. The Hylesiniinae of Turkey. *Istanbul Üniversitesi Orman Fakultesi Dergisi*, seri A, 37(1): 67-88. ().
- SEN-SARMA, P. K., AND M. L. THAKUR. 1956. Insect pests of Dipterocarpaceae and their management in India. *Indian Journal of Forestry* 9: 29-50. (cn).
- SEREZ, M. 1957. Verwendung des Aggregationspheromonen-praparatis "Ipslure" gegen den Mediteranen Kiefernborckenkafer, *Ips (Orthotomicus) crosus* (Woll.) (Col., Scolytidae) [Use of the aggregation-pheromone preparation "Ipslure" against the Mediterranean pine bark beetle *Ips crosus*]. *Anzei-*

- ger für Schadlingskunde Pflanzen- und Umweltschutz 60: 4-95. (cn).
- SEREZ, M., AND J. SCHONHERR. 1985. Bekämpfung von *Ips sexdentatus* (Boern.) (Col., Scolytidae) mit synthetischem Lockstoff Ipslure [Mass trapping of *Ips sexdentatus* with the synthetic pheromone Ipslure]. Journal of Applied Entomology 100: 24-26. (bv).
- *SERRANO, A. R. M. 1982. Contribuição para o conhecimento do povoamento, distribuição e origem dos coleópteros do Arquipélago dos Açores (Insecta, Coleoptera). Boletim Museu Municipal Funchal 34(147): 67-104. ().
- SEU, Y. B., AND KENJI MORI. 1986. A new synthesis of (-)-exobrevicomin. Agricultural and Biological Chemistry 50: 2923-2924. (bv ms).
- SEYBOLD, STEVEN J., J. E. MILSTEAD, T. OHTSUKA, I. KUBO, AND DAVID L. WOOD. 1988. The isolation and identification of the aggregation pheromone of *Ips lecontei* Swaine. International Congress of Entomology, Proceedings 1S: 441. (bv).
- SHAMOUN, S. F., AND M. P. LEVI. 1955. A chemical and microscopic study of decayed earlywood and latewood of loblolly pine killed by the southern pine beetle (*Dendroctonus frontalis*). Wood and Fiber Science 17: 22-28. (cn ms).
- SHARON, MIKE, AND C. G. O'NEIL. 1985. Forest pest conditions in the Rocky Mountain Region for 1984. United States Forest Service, Rocky Mountain Region, Timber, Forest Pest and Cooperative Forestry Management, Denver, Colorado, 36 p. (cn).
- SHARP, DAVID. 1870. [Title?]. Entomologist's Monthly Magazine 6: 256. (tx).
- SHARPE, PETER J. H., R. J. NEWTON, AND R. D. SPENCE. 1986. Forest pests: the role of phloem osmotic adjustment in the defensive response of conifers to bark beetle attack. Pages 113-132 in T. C. Hemmessey et al., Forestry sciences: stress physiology and forest productivity. Martinus Nijhoff, Dordrecht, Netherlands. viii + 239 p. (cn).
- SHARPE, PETER J. H., AND HSIN-JI WU. 1955. A preliminary model of host susceptibility to bark beetle attack. Pages 108-127 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September, 1983. 240 p. (cn).
- SHARPE, PETER J. H., HSIN-JI WU, R. G. GATES, AND J. D. GOESCHL. 1985. Energetics of pine defense systems to bark beetle attack. Pages 206-224 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- SHATILOV, O. A. 1985. The biology of little known species of bark beetles (Coleoptera, Ipsidae) of the Altai [In Russian, English summary]. Pages 54-57 in A. I. Cherepanov, Sistematika i biologiya chlenistonogikh i gel'mintov. (hb).
- SHAVLIASHVILI, I. A., AND D. G. ZHARKOV. 1985. Effects of ecological factors on the interaction between populations of *Dendroctonus micans* and *Ips typographus* (Coleoptera: Scolytidae). Pages 227-232 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Canadian Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1983. 240 p. (ec).
- SHAW, CHARLES C. III, ANDRIS EGLITIS, THOMAS H. LAURENT, AND PAUL E. HENNON. 1955. Decline and mortality of *Chamaecyparis nootkatensis* in southern Alaska, a problem of long duration but unknown cause. Plant Disease 69: 13-17. (cn).
- SHAW, M. R. 1959. *Cosmophorus cembrae* Ruschka, new to Britain (Hymenoptera: Braconidae: Euphorinae). Entomologist's Gazette 40: 241-243. (ec).
- SHAW, S. R. 1988. Euphorine phylogeny: the evolution of diversity in host-utilization by parasitoid wasps (Hymenoptera: Brachonidae). Ecological Entomology 13: 323-335. (ec).
- SHCHERBAK, G. I. 1985. Contribution to the study of the Dendrolaelapinae of North America (Multi-dendrolaelaps, Rhodacaridae, Gamasina) [In Russian, English summary]. Vestnik Zoologii 5: 25-36. (ec).
- SHEA, KEITH R. 1985. Integrated forest pest management in the South. Pages 3-4 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- SHEA, PATRICK J. 1989. Use of chemicals to protect trees from mountain pine beetle attack. Pages 114-117 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- SHEA, PATRICK J., MICHAEL I. HAWERTY, AND GARY E. DATERMAN. 1986. Impact of insects and methodology for monitoring insects in western white pine seed orchards. Pages 147-153 in Second conference of the cone and seed insects: proceedings. 3-5 September 1986, Briancon. International Union of Forestry Research Organizations. (cn ms).
- SHEA, PATRICK J., AND MARK D. MCGREGOR. 1987. A new formulation and reduced rates of carbaryl for protecting lodgepole pine from mountain pine beetle attack. Western Journal of Applied Forestry 2: 114-116. (cn).
- *SHERMAN. 1903. [*Scolytus rugulosus*]. North Carolina Agricultural Experiment Station, Bulletin 186: 5-6. 10. 21. ().
- SHERMAN, JOHN D., JR. 1910. A list of Labrador Coleoptera. New York Entomological Society, Journal 15: 173-197. (ds).
- SHORE, TERRY L. 1955. Ambrosia beetles. Canada Department of Environment, Canadian Forestry Service, Pacific Forest Research Centre, Victoria, British Columbia. Pest Leaflet FPL 72. 4 p. (cn ec hb).
- SHORE, TERRY L., PAULA A. BOUDEWYN, ELEANOR R. GARDNER, AND ALAN J. THOMSON. 1989. A preliminary evaluation of hazard rating systems for the mountain pine beetle in lodgepole pine stands in British Columbia. Pages 28-33 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- SHORE, TERRY L., AND JOHN ALEXANDER McLEAN. 1955. A

- survey for the ambrosia beetles *Trypodendron lineatum* and *Gnathotrichus retusus* (Coleoptera: Scolytidae) in a sawmill using pheromone-baited traps. Canadian Entomologist 117: 49-55. (bv cn).
- . 1988. The use of mark-recapture to evaluate a pheromone-based mass trapping program for ambrosia beetles in a sawmill. Canadian Journal of Forest Research 18: 1113-1117. (cn).
- SHORE, TERRY L., JOHN ALEXANDER MCLEAN, AND J. C. ZANUNCIO. 1987. Reproduction and survival of the ambrosia beetle *Trypodendron lineatum* (Oliv.) (Coleoptera: Scolytidae) in Douglas-fir and western hemlock logs. Canadian Entomologist 119: 131-139. (ec lb).
- SHRIMPTON, D. MALCOLM, AND A. J. THOMPSON. 1955. Relationship between phloem thickness and lodgepole pine growth characteristics. Canadian Journal of Forest Research 15: 1004-1005. (ec).
- SIEBER, ROBERT, AND G. BENZ. 1985. The diapause of the birch engraver, *Scolytus ratzeburgi* Janson (Col., Scolytidae), its termination by chilling, and manipulation with ecdysterone. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 58: 193-195. (ec lb).
- SIEBOLD, CARL THEODOR ERNST VON. 1849. Über Leon Dufour's Beitrag zur Käfer-Fauna der Pyrenäen. Stettiner Entomologische Zeitung 10: 306-311. (ds).
- SIEGFRIED, BLAIR D., CARL WARREN FATZINGER, ROBERT CLEVELAND WILKINSON, JR., AND JAMES L. NATION. 1986. In-flight responses of the black turpentine beetle (Coleoptera: Scolytidae) to individual monoterpenes, turpentine, and paraquat-treated slash pines. Environmental Entomology 15: 710-714. (bv).
- *SIERPINSKY, Z. 1980. Tannenschadlinge im Święty Krzyż-Gebirge in Polen. Acta Musei Reginaebradecensis S. A. Supplementum 1980: 254-256. ().
- SHIVERSTEIN, ROBERT MILTON. 1985. Chiral semiochemicals. Pages 121-140 in T. E. Acree and D. M. Soderlund, Semiochemistry: flavors and pheromones. Proceedings of the American Chemical Society Symposium, August 1983. Washington, D.C. (bv ms).
- *SIMANDL, JIRI, AND ZDENEK KLETECKA. 1987. Community of xylophagous beetles (Coleoptera) on *Sarothamnus scoparius* in Czechoslovakia. Acta Entomologica Bohemoslovaca 84: 321-329. ().
- *SIMIONESCU, ADAM. 1987. Protecția rasinoaselor împotriva dăunătorilor de tulpină [Protection of resinous forests against stem insect pests] [In Romanian, English summary]. București, Editura Ceres. 397 p. ().
- *SINADSKII, JU. V. 1958. Pests of the eastern clematis in the woods of the Amu-Darya floodlands. Bulletin M.o.-va ist. prirody, otd. biologii 63: 143-144. ().
- SINGH, PRITAM, B. H. MOODY, AND L. J. CLARKE. 1982. Newfoundland Region. Pages 11-15 in Annual Report of the Forest Insect and Disease Survey, 1979. Canada Department of Environment, Forestry Service, Ottawa, Ontario. (cn).
- *SINREICH, A. 1967. Fannistische Untersuchungen (Arthropoden und Mollusken) an einem Edelkastanienstandort am südlichen Rand der Thermalpen. Mitt. forstl. Bundesversuchsanstalt Wien 76: 33-57. ().
- SJODIN, K., L. M. SCHROEDER, H. H. EIDMANN, T. NORIN, AND S. WOLD. 1959. Attack rates of scolytids and composition of volatile wood constituents in healthy and mechanically weakened pine trees. Scandinavian Journal of Forest Research 4: 379-392. (bv).
- SKATTEBOL, L., AND Y. STENSTROM. 1955. Synthesis of racemic lineatin: an aggregation pheromone component of *Trypodendron lineatum*. Acta Chem. Scand., Ser. B, Org. Chem. Biochem. 39: 291-304. (bv ms).
- SKELLY, JOHN M., ET AL. 1988. Diagnosing injury to eastern forest trees. A manual for identifying damage caused by air pollution, pathogens, insects, and abiotic stresses. United States Department of Agriculture, Forest Service, Atlanta, Georgia, and Pennsylvania State University, University Park. 122 p. (cn lb ms).
- SKIDMORE, P., AND C. JOHNSON. 1965. A preliminary list of the Coleoptera of Merioneth, North Wales. Entomologist Gazette 20: 139-225. (ds).
- SLANKIS, A. I. 1967b. A new species of nematode, *Aphelenchoides macromicrus* sp. n. (Tylenchida), from the bark beetle, *Ips typographus* L. [In Russian]. Mater. nauk. confer. bse. obshch. Gel'mintol. Moscow 5: 279-282. (ec).
- . 1974. Four new species of Sphaerulariidae from Ipidae (Col.) [In Russian, English summary]. Parazitologiya 8: 57-62. (ec).
- SLANSKY, F., JR., AND ROBERT A. HAACK. 1986. Age-specific flight behavior in relation to body weight and lipid content of *Ips calligraphus* reared in slash pine bolts with thick or thin inner bark (phloem). Entomologia Experimentalis et Applicata 40: 197-207. (ay bv).
- SLESSOR, KEITH N., G. G. S. KING, D. R. MILLER, M. L. WINSTON, AND T. L. CUTFORTH. 1985. Determination of chirality of alcohol or latent alcohol semiochemicals in individual insects. Journal of Chemical Ecology 11: 1659-1667. (ms).
- SLOSSEN, ANNIE TRUMBULL. 1895. Additional list of insects taken in alpine region of Mt. Washington. Entomological News 6: 4-7. (ds).
- . 1906. Additional list of insects taken in alpine region of Mt. Washington. Entomological News 17: 323-326. (ds).
- SMILEY, ROBERT L., AND JOHN CONRAD MOSER. 1984. Notes and a key to separate normal and heteromorphic males of *Pycnomerus giganticus* Cross, Moser, & Rack and *P. dimorphus* Cross & Moser (Acrid: Pyrometidae). International Journal of Acarology 10: 11-14. (ec).
- . 1985. A new species, key to females, and distribution records for *Heterotarsonemus* (Acrid: Tarsonemidae). International Journal of Acarology 11: 247-253. (ec).
- *SMITH, I. 1921. [*Xyleborus perforans*]. Virgin Islands Agricultural Experiment Station, St. Croix. Bulletin 2. 23 p. ().
- SMITH, ANTHONY H., AND JOHN O. JOHNSON. 1955. Forest conditions in the state of New Mexico. Pages 11-14 in T. J. Rogers and P. F. Hessburg, Annual Southwestern Region Pest Conditions Report, 1954. United States Department of Agriculture, Forest Service, Southwestern Region, Albuquerque, New Mexico. Forest Pest Management Report R-3-85-9. 18 p. (cn).
- SMITH, D. P. 1987. Final environmental impact statement for the suppression of the southern pine beetle, Southern Region. United States Department of Agriculture, Forest Service, Southern Region, Management Bulletin R-S-MIB2. (cn).
- SMITH, I. M. 1955. Pests and disease problems in European forests. FAO Plant Protection Bulletin 33: 159-164. (cn).
- SMITH, MICHAEL T., R. BUSCH, THOMAS LEE PAYNE, AND J. C. DICKENS. 1988. Antennal olfactory responsiveness of three sympatric *Ips* species (*Ips avulsus* [Eichhoff], *Ips calligraphus* [Germar], *Ips grandicollis*

- [Eichhoff]], to intra- and interspecific behavioral chemicals. *Journal of Chemical Ecology* 14: 1289-1304. (ay hv).
- SMITH, RICHARD HARRISON. 1977. Book review: The insects and arachnids of Canada, Part 2: the bark beetles of Canada and Alaska, Coleoptera: Scolytidae. Entomological Society of America, Bulletin 23: 148. (ds tx ms).
- _____. 1986. Trapping western pine beetles with baited toxic trees. United States Department of Agriculture, Forest Service, Pacific Southwest Forest Experiment Station, Berkeley, California, Research Note PSW-382. 9 p. (cn).
- SMITH, RICHARD S., JR., AND J. W. DALE. 1981. Biological evaluation of white fir mortality in Horton and Dotta canyons, Plumas National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 81-21. (cn).
- SMITH, RICHARD S., JR., J. PIERCE, D. SCHULTZ, AND J. PRONOS. 1984. Biological evaluation of North Grove and Squaw Hollow campgrounds in Calaveras Big Tree State Park. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-07. (cn).
- SMITH, RICHARD S., JR., AND B. ROETTERING. 1982. A biological evaluation of three years of pest-caused tree mortality on the San Bernardino National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 82-4. (cn).
- *SNAPP. 1928. [*Scolytus rugulosus*]. United States Department of Agriculture, Farmers Bulletin 1557: 27-28. [Revised 1936]. ().
- *_____. 1941. [*Scolytus rugulosus*]. United States Department of Agriculture, Farmers Bulletin 1861: 17-19. [Revised 1954]. ().
- SNOW, G. A., W. H. HOFFARD, C. E. CORDELL, AND A. G. KAIS. 1989. Pest management in longleaf pine stands. Pages 128-134 in Symposium proceedings: Management of longleaf pine. 4-6 April 1989, Long Beach, Mississippi. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-75. (cn).
- *SOLBRECK, C. 1985. Insect migration strategies and population dynamics. Pages 641-662 in M. A. Rankin, D. Checkley, J. Cullen, C. Kitting, and P. Thomas, Migration mechanisms and adaptive significance. Contributions to Marine Science 68. ().
- SOLJIEIM, HALVOR. 1986. Species of Ophiostomataceae isolated from *Picea abies* infested by the bark beetle *Ips typographus*. *Nordic Journal of Botany* 6: 199-207. (ec).
- SOLOMON, JAMES D. 1985. Using hardwood lumber with insect, fungus, and bird defects as character-marked woods. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, Research Paper SO-226. 11 p. (cn).
- SOLOMON, JAMES D., FRANCIS I. MCCracken, ROBERT L. ANDERSON, ROBERT LEWIS, JR., FORREST L. OLIVERIA, THEODORE H. FILER, AND PATRICK J. BARRY. 1957. Oak pests: a guide to major insects, diseases, air pollution and chemical injury. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, Protection Report RS-PR7. 70 p. (cn).
- SOLOMON, JAMES D., AND J. A. PAYNE. 1986. A guide to the insect borers, primers, and girdlers of pecan and hickory. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station (New Orleans, Louisiana), General Technical Report SO-64. 31 p. (cn lib).
- *SOWER, LONNE L., AND GARY EDWARD DATERMAN. 1986. Use of behavioral chemicals in western U. S. forests. In I. Chaudry and C. Thies, Proceedings: 13th international symposium on controlled release of bioactive materials. 3-6 August 1986, Norfolk, Virginia. The Controlled Release Society, Lincolnshire, Illinois. ().
- SPEERS, CHARLES F. 1956. Radioisotopes in forest insect studies. Association of Southern Agricultural Workers, Proceedings 53: 130. (cn).
- SPRECHER VON BERNEGG, ANDREAS. 1960. Tropische und subtropische Weltwirtschaftspflanzen ihre Geschichte, kultur und volkswirtschaftliche Bedeutung, Teil III, Genusspflanzen, Band 2. Ferdinand Enke, Stuttgart. (cn).
- STAACK, J. 1985. Vom Fangbaum zur fälle: Geschichtliche Entwicklung der Borkenkäferbekämpfung [From trap tree to (pheromone) trap: the historical development of bark beetle control]. *Forst- und Holzwirt* 40: 27-31. (cn).
- STADNITSKII, G. V. 1986. Xylophagous insects in forest ecological systems [In Russian]. *Zashchita Rastenii* Moscow 12: 18-20. (hv).
- STAINES, C. L., JR. 1982. Distributional records of Platypodidae (Coleoptera) in Maryland. *Entomological Society of Washington, Proceedings* 84: 858-859. (ds).
- STAMBULL, M. 1988. The problem of coffee insect pests in Tanzania. *Taro Newsletter* 3(2): 10-14. (cn).
- STARK, RONALD WILLIAM, AND ALAN ANDREW BERRYMAN. 1971. Techniques for sampling western pine beetle populations and estimating the efficiency of natural control factors. Pages 191-206 in G. P. Patil, E. C. Pielou, and E. E. Waters, Statistical ecology: vol. 2. Pennsylvania State University Press, State College. (cn ms).
- STARK, RONALD WILLIAM, AND WILLIAM E. WATERS. 1985. Concept and structure of a forest pest management system. Pages 49-60 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn).
- STARK, RONALD WILLIAM, WILLIAM E. WATERS, AND DAVID LEE WOOD. 1985. Summary. Pages 191-201 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine bark beetle ecosystems. John Wiley & Sons, New York. (cn).
- *STARNES. 1899. [*Scolytus rugulosus*]. Georgia Agricultural Experiment Station Bulletin 42: 225-229 (1898). ().
- *_____. 1904. [*Scolytus rugulosus*]. Georgia Agricultural Experiment Station, Bulletin 67: 253-254. ().
- *STARZAK, JERZY R. 1974. Charakterystyka szkodliwej entomofauny lesnej rezerwatu Lipówka w Puszczy Niepolomickiej [Characteristics of the noxious forest entomofauna of the Reserve Lipówka in the Niepolomice Forest near Krakow]. *Acta Zoologica Cracoviensia, Krakow* 19: 237-252. ().
- _____. 1976. Obwczyn-*Stenostola dubia* (Laich) (Col., Cerambycidae)- w polsce, jego morfologia i biologia [Scolytidae, p. 206]. *Folia Forestalia Polonica, ser. A*, 22: 197-208. (ec).

- . 1977. Morphology, biology and life history of *Carilia* (= *Neogaurotes* Pod.) *virginica* (L.) (Col., Cerambycidae) [cites *Ips typographus*, *Pityophthorus pityographus*, *Polygraphus poligraphus*, *Dryocoetes autographus*]. Zeitschrift für Angewandte Entomologie 53: 269–251. (ec).
- . 1955. Badania nad bionomią i znaczeniem gospodarczym rzemlika plamistego, *Saperda scalaris* (L.) (Col., Cerambycidae) w Puszczy Niepolomickiej koło Krakowa [Scolytidae, p. 452]. Polskie Pismo Entomologiczne 58: 465–487. (ec).
- STARZYK, JERZY R., ALFRED KRÓL, DANIEL KUBISK, MACIEJ WITRYLAK, AND MALGORZATA OSSOWSKA. 1987. Wykorzystanie substancji feromonowych do zwalczania gatunków z rodzaju drwaliki (*Trypodendron* Steph.) w lasach górskich. Sylwan 131: 71–81. (ec).
- STARZYK, JERZY R., AND JOANNA KULACZEK. 1955. Studies on the infestation of boles and branches of *Abies alba* Mill. with cambio- and xylophagous insects at the Forest Experimental Station in Krynica (Beskid Sadecki Mts.). Pages 153–164 in IVth symposium on the protection of forest ecosystems. Warsaw Agricultural University. (ec).
- STARZYK, JERZY R., AND M. LESSAER. 1978. Studies on the distribution, morphology and biology of *Tetrops starki* Chev. (Col., Cerambycidae) [Scolytidae, p. 45]. Zeitschrift für Angewandte Entomologie 86: 35–46. (ec).
- STARZYK, JERZY R., AND MARIAN J. LUSZCZAK. 1952. Owady floemokambio- i ksylofagiczne spalowanych drzew iglastych i liściastych na wybranych powierzchniach w lesnym zakładzie doświadczalnym w krynicy. Acta Agraria et Silvestria 21: 99–119. (ec).
- STARZYK, JERZY R., AND ADAM SEK. 1953. Zgrupowania i zespoły owadów w pniakach świerkowych na wybranych powierzchniach w lesnym zakładzie doświadczalnym w krynicy (beskid sadecki). Acta Agraria et Silvestria 22: 71–86. (ec).
- STARZYK, JERZY R., AND MARIA SIEMEK. 1957. Zgrupowania i zespoły owadów kambio- i ksylofagicznych na strzałach i galeziach świerków II/III klasy wieku w lesnym zakładzie doświadczalnym w krynicy. Zeszyty Naukowe Akademii Rolniczej Im. H. Kołłataja w Krakowie, No. 213: 103–119. (ec).
- STARZYK, JERZY R., AND KRYSZYNA STARZYK. 1951. Owady kambiofagiczne, kambio-ksylofagiczne i ksylofagiczne w drzewostanach Puszczy Niepolomickiej. Studia Ośrodka Dokumentacji Fizjograficznej 9: 255–290. (ec).
- STARZYK, JERZY R., AND MAREK STYCZYŃSKI. 1954. Owady kambio- i ksylofagiczne w tyczkowinach i drógowinach jodecwych lesnego zakładu doświadczalnego w krynicy/beskid sadecki/. Zeszyty Naukowe Akademii Rolniczej Im. H. Kołłataja w Krakowie, No. 154: 103–112. (ec).
- STARZYK, JERZY R., AND ZBIGNIEW WITKOWSKI. 1951. Changes of the parameters describing the cambio- and xylophagous insect communities during the secondary succession of the oak-hornbeam association of the Niepolomice Forest near Krakow. Zeitschrift für Angewandte Entomologie 91: 525–533. (ec).
- . 1953. Z miary zgrupowań owadów kambio- i ksylofagicznych towarzyszące sukcesji wtórnej lasu w gradach Puszczy Niepolomickiej [In Polish, English summary]. Zakład Ochrony Przyrody i Zasobów Naturalnych Polskiej Akademii Nauk, ser. A., 27: 101–115. (ec).
- STARZYK, JERZY R., MACIEJ WITRYLAK, DANIEL KUBISK, TADEUSZ KAZMIERZAK, MARIAN LUSZCZAK, PRZEMISŁAW SAWIŃKO, AND MALGORZATA OSSOWSKA. 1957. Effect of severity of thinnings on the mortality of trees and occurrence of cambio- and xylophagous insects in the mountain forest stands composed of *Abies alba* Mill. and *Picea abies* (L.) Karst. Pages 179–189. IV symposium on the protection of forest ecosystems, Warsaw Agricultural University. (ec).
- STARZYK, JERZY R., AND GRZEGORZ WOJCIEK. 1955. Badania nad owadami kambio- i ksylofagicznymi jodli i świerka w różnych fazach rozwojowych drzewostanu w lesnym zakładzie doświadczalnym krynicy [In Polish, English summary]. Zeszyty Naukowe Akademii Rolniczej Im. H. Kołłataja w Krakowie, No. 197: 119–132. (cn).
- STAUBLI, A., AND H. HON. 1955. Les ennemis de nos vergers. Une nouvelle serie de 5 planches en couleur [The enemies of our orchards: a new series of 5 color plates]. Revue Suisse de Viticulture, d'Arboriculture et d'Horticulture 20: 347–354, 359–366. (cn).
- STEIGUER, J. E. DE, AND R. L. HEDDEN. 1955. Effects of aerial detection schedules on the age of southern pine beetle infestations. Forest Science 34: 229–235. (cn).
- STEIGUER, J. E. DE, R. L. HEDDEN, AND J. M. PYE. 1957. Optimal level of expenditure to control the southern pine beetle. United States Department of Agriculture, Forest Service, Southeastern Forest Experiment Station (Triangle Park, North Carolina), Research Paper SE-263. 30 p. (cn).
- STEJSKAL, J. V. 1920. Další příspěvek ku výzkumu broukové fauny Radhoště. Klub Přírodovědecký v Prostějově, Vestník 1919: 19–21. (ds).
- STENSETH, NILS CHRISTIAN. 1959a. A model for the conquest of a tree by bark beetles. Holarctic Ecology 12: 405–414. (cn lb).
- . 1959b. A simple population model for bark beetles providing general guidelines for the application of aggregation and anti-aggregation pheromones. Holarctic Ecology 12: 395–407. (bv cn).
- STENSETH, NILS CHRISTIAN, AND LAWRENCE RICHARD KIRKENDALL. 1959. Population dynamics of bark beetles with special reference to *Ips typographus*. Holarctic Ecology 12(4): 352–353. (lb).
- STEPHEN, FREDERICK M., AND M. P. LIII. 1955. A *Dendroctonus frontalis* infestation growth model: organization, refinement, and utilization. Pages 156–194 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research, United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 353 p. (cn).
- STEPHEN, FREDERICK M., M. P. LIII, TIMOTHY D. PAINE, AND G. W. WALLIS. 1955a. Influence of tree resistance on southern pine beetle population dynamics. International Congress of Entomology, Proceedings 15: 413. (cn lb).
- . 1955b. Using acute stress to modify tree resistance: impact on within-tree southern pine beetle populations. Pages 105–119 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles, Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn ec).
- STEPHEN, FREDERICK M., M. P. LIII, AND G. W. WALLIS. 1955. Southern pine beetle IPM. Arkansas Farm Research 37: 5. (cn).
- STEPHEN, FREDERICK M., AND TIMOTHY D. PAINE. 1955a. Conceptual model of infestation probability based

- on bark beetle abundance and host tree susceptibility: an addendum. *Environmental Entomology* 14:iv. (ms).
- _____. 1985b. Seasonal patterns of host tree resistance to fungal associates of the southern pine beetle. *Journal of Applied Entomology* 99: 113-122. (cn ec).
- STEPHEN, FREDERICK M., TIMOTHY D. PAINE, AND M. P. LILL. 1985. Mechanisms of tree resistance to bark beetles: synthesis and applications. Pages 69-82 in 34th Annual Forestry Symposium, Division of Continuing Education, Louisiana State University, Baton Rouge, Louisiana. (ec).
- STIPE, LAWRENCE E., R. L. JAMES, ROBERT LADD LIVINGSTON, ETAL. 1987. Idaho forest pest conditions and program summary, 1986. United States Department of Agriculture, Forest Service, Northern and Intermountain Regions and Idaho Department of Lands, Coeur d'Alene, Idaho, Report 87-1. 32 p. (cn).
- STOCK, A. J., JOHN HARVEY BORDEN, T. L. PRATT, II, D. PIERCE, JR. AND B. D. JOHNSTON. 1990. Entobrevicomin: an antiaggregation pheromone for the western balsam bark beetle, *Dryocoetes confusus* Swaine (Coleoptera: Scolytidae). *Canadian Entomologist* 122: 935-940. (bv hb).
- STOCK, A. J., AND R. A. GORLEY. 1989. Observations on a trial of broadcast burning to control an infestation of the mountain pine beetle *Dendroctonus ponderosae*. *Canadian Entomologist* 121: 521-523. (cn).
- STOCK, MOLLY WILFORD, JEAN CLAUDE GREGOIRE, AND MALCOLM MACFARLANE FURNISS. 1987. Electrophoretic comparison of European *Dendroctonus micans* and ten North American *Dendroctonus* species (Coleoptera: Scolytidae). *Pan-Pacific Entomologist* 63: 353-357. (ay).
- STONE, C. 1985. A new species of *Proctolaelaps* (Acari: Ascidae) from New South Wales, Australia. *Acarologia* 29: 319-327. (ec).
- STONE, C., AND J. A. SIMPSON. 1987. Influence of *Ips grandicollis* on the incidence and spread of bluestain fungi in *Pinus elliotii* billets in north-eastern New South Wales. *Australian Forestry* 50: 86-94. (ec).
- STRESSMANN, ERWIN, H. HANNEMANN, B. KLAUSNITZER, AND K. SENGLAUB. 1989. Exkursionsfauna für die Gebiete der DDR und der BRD. Band 2/1 Wirbellose. Teil 1. Insekten, 8 Auflage [Scolytidae and Platypodidae, p. 350-357]. Volk und Wissen Volkseigener, Berlin. 504 p. (hb ds tv).
- *STRICKLAND. 1916. [*Scolytus rugulosus*]. New York Department of Agriculture, Bulletin 79: 1099. ().
- *STRITTMATTER, M. 1985. Versuche zur Isolierung des Populationslockstoffes von *Cryphalus piceae* (Ratz.). Unpublished thesis, University of Freiburg. ().
- STRJCEK, J. 1988. Faunistic records from Czechoslovakia. Hymenoptera. *Acta Entomologica Bohemoslovaca* 55: 475-477. (ec).
- STROBEL, GARY A., AND F. SUGAWARA. 1986. The pathogenicity of *Ceratocystis monia* to lodgepole pine. *Canadian Journal of Botany* 64: 113-116. (ec).
- *STROHMAYER, HEINRICH. 1911i. [*Platypus cylindrus*]. Montevideo. *Anales del Museo Nacional*, ser. 2, 1(3): 85-88. ().
- STRONGMAN, D. B. 1987. A method for rearing *Dendroctonus ponderosae* Hopk. (Coleoptera: Scolytidae) from eggs to pupae on host tissue with or without a fungal complement. *Canadian Entomologist* 119: 207-208. (ec hb).
- STURGEON, KAREEN BARBARA, AND JEFFERY B. MITTON. 1986a. Allozyme and morphological differentiation of mountain pine beetles *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae) associated with host tree. *Evolution* 40(2): 290-302. (ay tv).
- _____. 1986b. Biochemical diversity of ponderosa pine and predation by bark beetles (Coleoptera: Scolytidae). *Journal of Economic Entomology* 79: 1064-1068. (cn).
- STURGEON, KAREEN BARBARA, AND J. L. ROBERTSON. 1985. Mitochondrial polysubstrate monooxygenase activity in western and mountain pine beetles (Coleoptera: Scolytidae). *Entomological Society of America, Annals* 75: 1-4. (ay).
- SUKARTANA, P. 1988. The possibility of rearing ambrosia beetles on artificial diets. *Malaysian Forester* 48: 347-352. (hb).
- *SUKHOVOLSKII, V. G. 1981. Tree-xylophage interaction and dielectric properties of tree tissues [abstract]. The role of insect-plant relationship in the population dynamics of forest pests, IUFRO/MAB Meeting, 24-25 August 1981, Irkutsk, USSR 57: 47. ().
- _____. 1982. A change in the dielectric properties of the tissues of a tree affected by insects [In Russian]. Pages 41-56 in A. S. Isaev, Konsortivnye svyazi dereva i dendrofilnykh nasekomykh. Novosibirsk, USSR, Nauka, Siberian Section. Translation, Canada Department of Environment, OOENV TR-2255 (1983). 28 p. (ec).
- *SUKOVATOVA, L. M., L. S. MILOVIDOVA, AND K. S. TRUBACHEVA. 1987. Detecting the entomopathogenic fungus *Cordyceps militaris* Fr. Lk. in the southern Ob River area in Tomsk Oblast Russian SFSR USSR [In Russian]. *Mikol Fitopatol* 21: 528-529. ().
- SULISTYOWATI, E. 1986. Masalah hama bubuk buah kopi, *Hypothenemus hampei* Ferr. (Coleoptera, Scolytidae) dan usaha pengendaliannya [The problem of the coffee berry borer, *Hypothenemus hampei*, and its control] [In Hindi, English summary]. *Pelita Perkebunan* 2: 10-18. (cn hb).
- SUMER, S. 1987. The distribution of cypress (*Cupressus* L.) in Turkey and the current status of its pests and diseases, especially cypress canker disease. *Istanbul Üniversitesi Orman Fakültesi Dergisi*, ser. A, 37: 46-66. (cn).
- *SUMMERS, S. V. 1874. Catalogue of the Coleoptera from the region of Lake Pontchartrain, La. Buffalo Society of Natural Science, Bulletin 2: 78-99. ().
- SUSUT, J. P. 1967. Annual district report: Peace River District, 1966. Pages 83-91 in Annual district reports of the Forest Insect and Disease Survey for Alberta-Northwest Territories-Yukon region, 1966. Canada Department of Forestry and Rural Development, Canadian Forestry Service, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-6. (cn).
- _____. 1969. Annual district report: Yukon Territory, 1968. Pages 79-86 in Annual district reports of the Forest Insect and Disease Survey, Alberta-Northwest Territories-Yukon region, 1968. Canada Department of Fisheries and Forestry, Forest Research Laboratory, Calgary, Alberta, Information Report A-X-22. (cn).
- SVTHRA, PAVEL. 1987. Suitability of elm firewood to bark beetle attack stored under polyethylene sheeting. *Journal of Arboriculture* 13: 164-166. (bv hb).
- _____. 1988. To dig or not to dig the red turpentine beetle from infested pines. *Grower Points* 25(5): 2-3. (cn ms).
- _____. 1989. How to protect drought stressed pine from red

turpentine beetle attacks, without lindane. *Grower Points* 25(5): 3-4. (cn).

*SWAINE, JAMES MALCOLM. 1917b. [*Ips radiatae*]. *Canadian Entomologist* 49: 356. ().

———. 1919c. The Entomological Branch: the balsam injury in Quebec and its control. *Agricultural Gazette of Canada* 6: 227-233. (cn ec).

*SWARUP, G., AND RAJAN. 1955. Mechanism of insect control by nematodes. Pages 49-59, 254-258 in R. K. Bhan-tor, S. Verma, and S. I. Farooqi. Non-insect pests and predators. Impact of non-insect pests and predators on food production and environment: symposium proceedings. New Delhi, India. ().

SWEDENBORG, P. D., R. L. JONES, M. E. ASCERNO, AND V. R. LANDWEHR. 1955. *Hylurgopinus rufipes* (Eichhoff) (Coleoptera: Scolytidae): attraction to broodwood, host colonization behavior, and seasonal activity in central Minnesota. *Canadian Entomologist* 120: 1041-1050. (bv hb).

SWEDENBORG, P. D., R. L. JONES, AND LEE C. RYKER. 1959. Stridulation and associated behavior of the native elm bark beetle *Hylurgopinus rufipes* Eichhoff (Coleoptera: Scolytidae). *Canadian Entomologist* 121: 245-252. (bv).

SWINGLE, ROGER ULASSES. 1950. Report on the Dutch elm disease. *Arborists' News* 15(10): 113-114. (cn).

SYED, AKBAR, AND KENNETH GRAHAM. 1957. Response of the mountain pine beetles, *Dendroctonus ponderosae* Hopkins, to ethanol in a laboratory olfactometer. *Canadian Entomologist* 119: 459-490. (bv).

SYME, PAUL D., AND K. L. NYSTROM. 1955. Insects and mites associated with Ontario forests: classification, common names, main hosts, and importance. Canada Department of Forestry, Forestry Service, Great Lakes Forestry Centre, Sault Ste. Marie, Ontario, Information Report 0-X-392. 131 p. (cn ds tx).

SYMONS, THOMAS BADDELEY. 1904. Entomological notes for the year in Maryland. United States Department of Agriculture, Division of Entomology, Bulletin 46. (cn).

SZONTAGI, P. 1955. Fitofag rovarok szerepe a kocsanytalan tolgypusztulasaban [The role of phytophagous insects in the decay of sessile oak] [English title]. *Novényvedelem* 21(5): 219. (cn).

T

TABASHNIK, BRUCE E., W. J. MATTHEWSON, AND J. R. MILLER. 1955. Host acceptance behavior of the red pine cone beetle (*Conophthorus resinosae*). *Entomologia Experimentalis et Applicata* 37: 3-7. (bv).

TAHA, HAMDI A., AND FREDERICK M. STEPHEN. 1954. Modeling with imperfect data: a case study simulating a biological system. *Simulation* 42: 109-115. (cn).

TAIT, SUSAN M., CHARLES G. SHAW III, AND ANDRIS EGLITIS. 1955. Occurrence of insect and disease pests on young-growth Sitka spruce and western hemlock in southeastern Alaska. Pacific Northwest Forest and Range Experiment Station, Portland, Oregon, Research Note PNW-433. 16 p. (cn).

TAKAHASHI, K. 1959. Flight activity of insects sampled with a truck trap. IV: flight activity of Curculionidae (Coleoptera). *Japanese Journal of Entomology* 57: 402-405. (hb).

*TALBERT, THOMAS JESSE. 1946. [*Phloeotribus liminaris*]. *Gen. Hort.* 1946: 190. ().

TANDON, P. L., AND A. VERGHESE. 1957. New insect pests of

certain fruit crops. *Indian Journal of Horticulture* 44(1-2): 121-122. (cn).

*TANG, Z. T., Z. W. JIN, Z. J. ZHOU, C. H. WANG, AND S. Z. LI. 1955. Some tropical hardwood destroying insects and their control methods [In Chinese, English summary]. *Journal of Nanjing Institute of Forestry* 1: 35-43. ().

TEALE, S. A., AND GERALD NORMAN LANIER. 1955. Seasonality of pheromone specificity in *Ips pini* (Coleoptera: Scolytidae). *International Congress of Entomology, Proceedings* 18: 429. (bv).

TEILLON, H. BRENTON, BARBARA S. BURNS, AND RONALD S. KELLEY. 1951. Forest insect and disease conditions in Vermont, calendar year 1951. *Forestry, Parks, and Recreation, Montpelier, Vermont*. (cn).

———. 1952. Forest insect and disease conditions in Vermont, calendar year 1952. *Forestry, Parks, and Recreation, Montpelier, Vermont*. (cn).

———. 1953. Forest insect and disease conditions in Vermont, calendar year 1953. *Forestry, Parks, and Recreation, Montpelier, Vermont*. (cn).

———. 1956. Forest insect and disease conditions in Vermont, calendar year 1956. *Department of Forests, Parks, and Recreation, Montpelier, Vermont*. 40 p. (cn).

TEILLON, H. BRENTON, RONALD S. KELLEY, AND E. E. KEENAN. 1979. Forest insect and disease conditions in Vermont, calendar year 1979. *Forestry, Parks, and Recreation, Montpelier, Vermont*. (cn).

TEILLON, H. BRENTON, RONALD S. KELLEY, AND B. D. SCHULTZ. 1950. Forest insect and disease conditions in Vermont, calendar year 1950. *Forestry, Parks, and Recreation, Montpelier, Vermont*. (cn).

TENKOCOVA, I., AND J. MITUCHI. 1957. Nematodes new for the fauna of the Czechoslovak Socialist Republic with the affinity to scolytids (Coleoptera: Scolytidae). *Helminthologia* 24: 281-291. (ec).

TERASAKI, YUKIO, NABIAKI YOSHIDA, KENJI FUKUYAMA, AND KIMITO FURUTA. 1957. Response of *Larix leptolepis* to inoculated *Ips cembrae* [In Japanese]. *Tokyo Daigaku Nogakubu Enshurin Hokoku* [Bulletin of the Tokyo University Forests] 77: 19-30. (cn).

TERRA, PAULO S. 1957. Guia para identificacao de escolitideos (Coleoptera, Scolytidae) associados ao cacaneiro (*Theobroma cacao* L.) no sul da Bahia [Guide for the identification of the ambrosia beetles (Coleoptera, Scolytidae) associated with the cacao tree in the south of Bahia]. *Revista Theobroma* 17: 17-30. (cs tx).

*THALENHORST, W. 1979. Ablauf der Borkenkäfer-Kalamität in Naturwaldreservaten (NWR). Pages 35-47 in *Aus dem Walde*, 1979, Dokumentation der Sturmkatastrophe vom 13. November 1972. Part V. Mitteilungen aus der Niedersächsischen Landesforstverwaltung 31. 190 p. ().

THAPA, R. S., AND PRATAP SINGH. 1956. Large-scale mortality of deodar tree by the bark borer, *Scolytus major* Stebbing (Scolytidae, Coleoptera) in Kulu Forest Division, Himachal Pradesh. *Indian Forester* 112: 392-398. (cn ec hb).

THATCHER, ROBERT CLIFFORD, AND M. P. CONNOR. 1955. Identification and biology of southern pine bark beetles. United States Department of Agriculture, Forest Service, Agricultural Handbook 634. 14 p. (bv hb).

THATCHER, ROBERT CLIFFORD, GARLAND N. MASON, AND GERARD D. HERTEL. 1955. Integrated pest management in the South: an overview. Pages 364-375 in S.

- J. Granham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- THIRUGNANASUNTHARAN, K. 1957. The use of insecticides in tea. *Tea Bulletin* 7: 19-26. (cn).
- THIRUGNANASUNTHARAN, K., AND I. C. R. JAYACHANDRAN. 1959. Recent observations on resistance and susceptibility of tea clones to the shot-hole borer beetle, *Xyleborus fornicatus* Eichl. (Coleoptera: Scolytidae). *Sri Lanka Journal of Tea Science* 58: 50-55. (cn).
- THOMAS, P. R. 1956. Infestation by pine and spruce bark beetles in British Columbia and its effect on kraft and mechanical pulping. Pages 35-47 in R. W. Nielson, Harvesting and processing of beetle-killed timber. Proceedings of a symposium. Special publication, Forintek Canada Corporation SP-26. 53 p. (cn).
- THOMAS, TERRI L., AND J. K. AGEI. 1956. Prescribed fire effects on mixed conifer forest structure at Crater Lake Oregon. *Canadian Journal of Forest Research* 16: 1052-1057. (bv).
- THOMPSON, WILLIAM A. 1955. SPB-Microbeetles: a simulation system for evaluating economic performance of management alternatives for stands attacked by southern pine beetles. Pages 292-299 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383. (cn).
- THOMPSON, WILLIAM A., AND JOHN CONRAD MOSER. 1956. Temperature thresholds related to flight of *Dendroctonus frontalis* Zimm. (Col.: Scolytidae). *Agronomie* 6: 905-910. (ec hb).
- THOMSON, A. J., LASZLO SAFRANYIK, D. MALCOLM SHIMPTON, AND H. S. WHITNEY 1955. A theory of mountain pine beetle population response to weather-induced changes in host resistance. Pages 128-135 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September 1953. 240 p. (ec hb).
- THOMSON, M. J. 1964. 1964. Status of insects in the Fort Frances district. Pages G36-42 in Annual district reports of the Forest Insect and Disease Survey Ontario, 1963. Canada Department of Agriculture, Forest Biology Division, Sault Ste Marie, Ontario. (cn).
- TILDEN, PAUL E. 1955. Remedial treatment of lodgepole pine infested with mountain pine beetle: efficacy of three insecticides. United States Department of Agriculture, Forest Service, Pacific Southwest Forest Experiment Station, Berkeley, California, Research Note PSW-374. 4 p. (cn).
- TILDEN, PAUL E., WILLIAM DELLES BEDARD, JR. 1955. Field response of *Dendroctonus brevicomis* to exo-brevicomin, frontalin and myrcene released at two proportions and three levels. *Journal of Chemical Ecology* 11: 757-766. (bv).
- _____. 1955. Effect of verbenone on response of *Dendroctonus brevicomis* to exo-brevicomin, frontalin, and myrcene. *Journal of Chemical Ecology* 14: 113-122. (bv).
- TILDEN, PAUL E., WILLIAM DELLES BEDARD, JR., DAVID LEE WOOD, AND LLOYD E. BROWNE. 1957. Interruption of response of *Dendroctonus brevicomis* to attractive pheromone by release of pheromone at several rates and spacings. *Journal of Chemical Ecology* 13: 85-97. (bv).
- TITUS, F. A. 1957. Pests that changed the forests of Funday National Park. Government of Canada, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-165. 33 p. (cn).
- TITUS, F. A., O. A. MEKLE, AND K. J. HARRISON. 1955. Scientific and common names of insects and mites of interest in the Maritime Provinces. Canada Department of Environment, Forestry Service, Maritimes Forest Research Centre, Fredricton, New Brunswick, Information Report M-X-155. 130 p. (tx).
- TKACZ, BORYS M. 1956. Evaluation of pest conditions: Indian Creek Recreation Lands, Carson City District, Bureau of Land Management. United States Department of Agriculture, Forest Service, Intermountain Region, Ogden, Utah, Forest Pest Management Report R4-56-S. 6 p. (cn).
- TKACZ, BORYS, AND RICHARD FREDERICK SCHMITZ. 1956. Association of an endemic mountain pine beetle population with lodgepole infected by *Armillaria* root disease in Utah. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, Research Note INT-353. 7 p. (ec).
- TOMALAK, MAREK, H. E. WELCH, AND T. D. GALLOWAY. 1955. Interaction of parasitic nematode *Parasitaphelenchus oldhami* (Nematoda: Aphelenchoididae) and a bacterium in Dutch elm disease vector *Hylurgopinus rufipes* (Coleoptera: Scolytidae). *Journal of Invertebrate Pathology* 52: 301-305. (ec).
- _____. 1959a. Effect of crowding on *Sulphurectylenchus* spp. (Nematoda: Allantonematidae) in the hemocoel of their bark beetle hosts (Coleoptera: Scolytidae). *Canadian Entomologist* 121: 521-522. (ec).
- _____. 1959b. Nematode parasites of bark beetles (Scolytidae) in southern Manitoba, with descriptions of three new species of *Sulphurectylenchus* Ruhm (Nematoda: Allantonematidae). *Canadian Journal of Zoology* 67: 2497-2505. (ec).
- _____. 1959c. Pathology of *Aphelenchoides ptyokteini* (Nematoda: Aphelenchidae) in the malpighian tubules of *Pityokteines sparsus* (Coleoptera: Scolytidae). *Journal of Invertebrate Pathology* 53: 140-141. (ec).
- _____. 1959d. Parasitism of *Parasitorhabditis obtusa* and *Parasitorhabditis autographi* (Nematoda: Rhabditidae) in the digestive tract of their bark beetle (Coleoptera: Scolytidae) hosts. *Journal of Invertebrate Pathology* 53: 57-63. (ec).
- TOMMERAS, B. A. 1955. Specialization of the olfactory receptor cells in the bark beetle *Ips typographus* and its predator *Thanosinus fornicarius* to bark beetle pheromones and host tree volatiles. *Journal of Comparative Physiology, A (Sensory, Neural, and Behavioral Physiology)*, 157(3). (ay).
- * _____. 1957. Olfaction in bark beetle communities: inter-specific interactions in regulation of colonization density, predator-prey relationship and host attrac-

- tion. Unpublished dissertation, University of Trondheim, Norway, 79 p. ().
- . 1988. The clerid beetle, *Thanasimus formicarius*, is attracted to the pheromone of the ambrosia beetle, *Trypodendron lineatum*. *Experimentia* [Basel] 44: 536–537. (ec).
- . 1989. Host selection by odorous compounds from host and non-host trees in bark beetles. *Fauna Norvegica* 36: 75–80. (bv ms).
- TOMMERAS, B. A., S. MADSEN, AND H. MUSTAPARTA. 1989. Concentration dependent influence of ex-brevicomin on the pheromone attraction of the bark beetle *Ips typographus* L. (Coleoptera, Scolytidae). *Journal of Applied Entomology* 105: 24–26. (bv).
- TOMMERAS, B. A., AND H. MUSTAPARTA. 1987. Chemoreception of host volatiles in the bark beetle *Ips typographus*. *Journal of Comparative Physiology, A* (Sensory, Neural, and Behavioral Physiology) 161: 705–710. (bv).
- . 1989. Single cell responses of pheromones, host and non-host volatiles in the ambrosia beetle *Trypodendron lineatum*. *Entomologia Experimentalis et Applicata* 52: 141–148. (bv).
- TOMMERAS, B. A., H. MUSTAPARTA, O. GEDARAAS, AND T. ANTHONSEN. 1987. Single cell responses combined with gas chromatographical separation of host volatiles in bark beetles [abstract]. *Chemical Senses* 12: 193–194. (ay bv).
- TOMMERAS, B. A., H. MUSTAPARTA, AND JEAN CLAUDE GREGOIRE. 1985a. Electrophysiological recordings from olfactory receptor cells in *Dendroctonus micans* and *Rhizophagus grandis*. Pages 95–106 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles: *Dendroctonus micans*. Commission of European Communities and Universite Libre de Bruxelles, Belgium, Proceedings of seminar, 3–4 October 1984. 141 p. (ay).
- . 1985b. Electrophysiological recordings from olfactory receptor cells in *Dendroctonus micans* and *Rhizophagus grandis*. Pages 98–106 in R. Gilles, Animals and environmental stress. Pergamon Press. (ay bv).
- TOTANI, K., AND T. SUGIMOTO. 1957. Studies on the relationship of the interspecies of the genus *Xyleborus* Eichhoff (Coleoptera, Scolytidae) by the proventriculus [In Japanese, English summary]. *Research Bulletin of the Plant Protection Service, Japan* 23: 39–43. (ay).
- TOTH, JOZSEF. 1985. Kartetele es eletmodja magyarorszagon [Damage and habits of *Ips sexdentatus* Boer. in Hungary] [English title]. *Novenyvedelem* 21: 97–100. (cn hb).
- *TOWNSEND, CHARLES HENRY TYLER. 1928. The contribution to a knowledge of the coleopterous fauna of the lower Rio Grande Valley in Texas and Tamaulipas, with biological notes and special reference to geographical distribution. *Texas Academy of Science, Transactions* 5: 51–101. ().
- TRAUBOTH, V. 1985. Der Einsatz von Fensterfallen zur Ermittlung des Flugrhythmus vom Buchdrucker *Ips typographus* im Jahr 1983 [Use of window traps to determine the night activity pattern of *Ips typographus* in 1983]. *Beiträge für die Forestwirtschaft* 19: 42–44. (cn hb).
- TREAT, A. E. 1971. [Book review]. *New York Entomological Society, Journal* 79: 235. (ms).
- TREILAN, I. R., G. L. KAD, AND S. GUPTA. 1986. A new synthesis of racemic frontalin, the pheromone of *Dendroctonus* bark beetles. *Indian Journal of Chemistry* 25B: 1243–1244. (bv cn).
- TRIESELMANN, R. A. 1963. Status of insects in the Gogama district. Pages D41–55 in *Annual district reports of the Forest Insect and Disease Survey Ontario, 1962*. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- TRIGGIANI, O. 1984. *Tomicus (Blastophagus) piniperda* (Coleoptera, Scolytidae, Hylesininae): biologia, danni e controllo nel litorale Ionico [Tomicus piniperda: biology, damage and control on the Ionic Coast]. *Entomologica* 19: 5–21. (cn ec hb).
- TRINNELL, J. R. 1962. Status of insects in the Gogama district. Pages 266–283 in *Annual district reports of the Forest Insect and Disease Survey Ontario, 1961*. Canada Department of Agriculture, Forest Biology Division, Sault Ste. Marie, Ontario. (cn).
- TRIPLEHORN, CHARLES A. 1990. Review of the genus *Corticus* (Coleoptera: Tenebrionidae) of America north of Mexico. *Entomological Society of America, Annals* 83(3): 277–656. (ec).
- TRIPLEHORN, CHARLES A., AND JOHN CONRAD MOSER. 1970. Two new species of *Corticus* from Mexico and Honduras (Coleoptera: Tenebrionidae). *Coleopterists Bulletin* 24: 47–50. (ec).
- TRIPP, H. A., AND J. K. ROBINS. 1968a. Annual report of the Forest Insect and Disease Survey: Alberta-Northwest Territories-Yukon region, 1967. Canada Department of Forestry, Forest Research Laboratory, Calgary, Alberta, Information Report. (cn).
- TROFIMOV, V. N., AND V. A. LIPATIN. 1986. Variability of stem insect communities [In Russian]. *Lesovedenie* 1986(1): 51–57. (hb).
- *TROOP. 1894. [*Scolytus rugulosus*]. *Indiana Agricultural Experiment Station, Bulletin* 53: 126–130. ().
- TROSTLE, GALEN E. 1986. Distinguishing mated and unmated mountain pine beetles in alcohol-preserved specimens. United States Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station, Ogden, Utah, Research Note INT-356. 3 p. (ay).
- TSUNEDA, A., S. MURAKAMI, K. NISHIMURA, AND M. MIYAJI. 1986. Pleomorphism and conidiogenesis in *Rhinoctadiella atrovirens* isolated from beetle galleries. *Canadian Journal of Botany* 64: 1112–1119. (ec).
- TUBBS, P. K. 1986. Opinion 1408. *Hypocryphalus mangiferae* (Stebbing, 1914) given nomenclatural precedence over *Cryphalus inops* Eichhoff, 1872, and *Hypothenemus griseus* Blackburn, 1885 (Insecta, Coleoptera). *Bulletin of Zoological Nomenclature* 43: 245–246. (tx).
- *TULLGREN. 1916. [*Hylastes opacus*, *Pityogenes quadridens*, *Scolytus rugulosus*]. *Medd. från Centralanstalten för Jorsbruksforsk.* No. 152, *Entomologiska Avelningen* No. 27: 104. ().
- TUMLINSON, J. H. 1985. Beetles: pheromonal chemists par excellence. Pages 367–380 in P. A. Hedlin, Bioregulators for pest control. American Chemical Society, Washington, D.C. (bv).
- TUNNOCK, SCOTT, O. J., DOOLING, AND S. KOHLER. 1984. Montana forest pest conditions and program highlights, 1983. United States Department of Agriculture, Forest Service, Northern Region, Missoula, Montana, Cooperative Forestry and Pest Management Report 84–2. 31 p. (cn).
- TUNSET, K., A. C. NILSEN, AND J. ANDERSON. 1988. A new trap design for primary attraction of bark beetles and

- bark weevils (Col., Scolytidae and Curculionidae). *Journal of Applied Entomology* 106: 266-269. (cn).
- TURNBOW, ROBERT HAROLD, JR., ROBERT N. COULSON, L. HU, AND RONALD FORREST BILLINGS. 1982. Procedural guide for using the interactive version of the TAMBEETLE model of southern pine beetle population and spot dynamics. Texas Agricultural Experiment Station MP-1518. 24 p. (cn ms).
- *TWIGG. 1911. [*Phloeotribus liminaris*]. Kansas Horticultural Society 1911: 319-320. (.)
- ## U
- UCHASTNOVA, L. N. 1955. The complex of fungi associated with the galleries of *Platypus cylindriciformis* (Coleoptera: Platypodidae) and *Xyleborus monographus* (Coleoptera: Scolytidae). *Biologicheski nauki*, Moscow 2: 47-50. (ec).
- _____. 1958. Some data on the mycetangia of bark and ambrosia beetles. *Biologicheski nauki* 5: 38-41. (ay ec).
- ULRICH, ANTON. 1954b. Ein Grossversuch der Fichtenborkenkäferbekämpfung mit Hexa-Nebel. Pages 491-496 in Wellenstein. Forstschutzstelle Südwest, Ringingen. 496 p. (cn).
- *ULTEE, A. J. 1930. Verslag over 1929 [*Xyleborus compactus* biology]. Mededeelingen van het Proefstation Malang 75: 28. (.)
- *_____. 1931a. Verslag over 1930 [*Xyleborus compactus* biology]. Mededeelingen van het Proefstation Malang 80: 17. (.)
- *_____. 1931b. [*Xyleborus compactus*]. *Bergcultures* 5: 915. (.)
- *_____. 1931c. [*Xyleborus compactus*]. *Bergcultures* 5: 1090-1100. (.)
- *URBAN, J. 1986. Škodlivý výskyt lykohuba zrnitého na jasaněch oslabených nedostatkem vody. *Lesnická práce*, Praha 1986: 354-361. (.)
- ## V
- *VAJDA, Z. 1974. Nauka o Zastiti summa. Škol. knjiga, Zagreb [Yugoslavia]. (.)
- *_____. 1974. Uzročnici susenja slavonskih suma. Pages 221-235 in Jug. Ak. zn. i um.: Zb. o stotoj obl. sum. Jugoist. Slavonije, Zagreb [Yugoslavia]. (.)
- *VALENTA, V. T. 1981. Interrelations of pine and spruce xylophages with the food objects [abstract]. The role of insect-plant relationship in the population dynamics of forest pests, IUFRO/MAB meeting, 24-28 August 1981, Irkutsk, USSR 90: 11. (.)
- *VALTEROVA, I., B. KALINOVA, MILOŠ KNIZEK, A. SVATOS, AND J. VRKOC. 1990. Main components of the aggregation pheromone of the Cuban bark beetle *Ips interstitialis* [abstract]. Conference on Insect Chemical Ecology, 12-18 August 1990, Tabor, Czechoslovakia 1990: 42. (.)
- VANDENBURG, JAMES H. 1989. Silvicultural strategies to minimize mountain pine beetle losses: an overview. Pages 43-44 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- *VAN DER GOOT, P. 1934. [*Xyleborus compactus*]. Mededeelingen van het Institute voor Plantenziekten 83: 39-69. (.)
- *_____. 1935. [*Xyleborus compactus*]. Mededeelingen van het Institute voor Plantenziekten 84: 36-51, 66-67, 71-84. (.)
- VANDERWEL, DESIREE, AND A. C. OEHLISCHLAGER. 1987. Biosynthesis of pheromones and endocrine regulation of pheromone production in Coleoptera. Pages 175-215 in G. D. Prestwich and G. L. Blomquist. *Pheromone biochemistry*. Academic Press, Orlando, Florida. 565 p. (bv ms).
- VANDERWEL, DESIREE, A. C. OEHLISCHLAGER, S. M. SINGH, S. RAMASWAMY, T. CUTFORTH, H. D. PIERCE, JR., AND JOHN HARVEY BORDEN. 1985. Biosynthesis of pheromones by selected species of grain and bark beetles. Chemical Congress of North America [abstracts] 3(1): Agro 1. (bv ms).
- *VAN DINE, D. L. 1913a. [*Xyleborus* spp.]. Puerto Rico Sugar Producer's Association Experiment Station, Rio Piedras, Puerto Rico, Bulletin 5: 25-46. (.)
- *_____. 1913b. [*Xyleborus* spp.]. *Journal of Economic Entomology* 6: 251-257. (.)
- *VAN DYKE, EDWIN COOPER. 1953. The Coleoptera of the Galapagos Islands. Occasional Papers of the California Academy of Science 22: 1-181. (.)
- VAN SICKLE, G. ALLEN. 1989. Status of mountain pine beetle in western Canada. Pages 6-8 in G. D. Amman, Symposium proceedings: Management of lodgepole pine to minimize losses to the mountain pine beetle. United States Department of Agriculture, Forest Service, Intermountain Research Station, Ogden, Utah, General Technical Report INT-262. 119 p. (cn).
- VASECHKO, G. I. 1985. The problems that need to be solved for improvement of bark beetle control. *Journal of Applied Entomology* 106: 1-12. (cn).
- VASIEVICI, J. M., AND WILLIAM A. THOMPSON. 1985. I-T-E-M-s: an integrated method to project southern pine stand development. Pages 300-327 in S. J. Bramham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana, General Technical Report SO-56. 383 p. (cn).
- VASSILAIA-ALEXOPOULOU, P., P. A. MOURIKIS, AND C. T. BUCHELOS. 1986. *Coccotrypes dactyliperda* Fabr. a new species in the Greek fauna. *Annales de l'Institute Phytopathologique Benaki* 15: 87-89. (ay ds).
- VAUPEL, O., L. DIMITRI, E. KÖNIG, AND W. BERWIG. 1986. Zur optimierung des Falleneinsatzes bei Buchdrucker und Gestreiftem Nutzholzborkenkäfer [Optimization of the use of traps against *Ips typographus* and *Trypodendron lineatum*]. *Allgemeine Forstzeitschrift* 23: 572-574. (cn).
- VAZQUEZ, L. L. 1988. *Ambrosiodmus lecontei* Hopkins (Coleoptera: Scolytidae) como perforador del tallo del café [*Ambrosiodmus lecontei*, a coffee stem borer] [English summary]. *Revista de Protección Vegetal* 3: 271-272. (cn ds).
- VAZQUEZ, L. L., AND S. MONTEAGUDO. 1988. *Xylosandrus compactus* (Coleoptera, Scolytidae): nuevo insecto danino para el café en Cuba [*Xylosandrus compactus*: new coffee pest in Cuba] [English summary]. *Revista de Protección Vegetal* 3: 67-73. (cn ds).
- VEITCH, R. 1923. Minor pests of sugarcane in Fiji. Agricultural Report, Colonial Sugar Refining Company, Limited, Sydney, No. 7. 30 p. (cn).
- VERDCOURT, B. 1952. Further records of Bedfordshire Coleoptera. *Entomologist's Monthly Magazine* 58: 73-80. (ds).

- VIERECK, H. L. 1925. New genera and species of Ichneumonidae in the Canadian National Collection. *Canadian Entomologist* 57: 71-78. (ec).
- VILLA CASTELLO, J. 1955. Enemigos naturales y organismos asociados al descortezador de pinos *Dendroctonus adjunctus* Blandford en el Navado de Colima [Natural enemies and organisms associated with the pine bark beetle *Dendroctonus adjunctus* in Nevado de Colima, Mexico]. Instituto Nacional de Investigaciones, Boletín Técnico 121. 22 p. (ec).
- VILLACORTA, A. 1955. Dieta meridica para criacao de sucessivys geracoes de *Hypothenemus hampei* (Ferrari, 1867) (Coleoptera: Scolytidae). *Anais da Sociedade Entomologica do Brasil* 14: 315-319. (hb).
- VILLAIN, L. 1955. Concentraciones letales de insecticidas sobre la broca *Hypothenemus hampei* Ferr. 1867 [Lethal concentrations of insecticides on the coffee berry borer *Hypothenemus hampei*] [English summary]. *Boletín de Promecafe* 40: 11. (cn).
- VINODKUMAR, P. K., M. M. BALAKRISHNAN, AND T. S. GOVINDARAJAN. 1956. *Eupelmus* sp. (Hymenoptera: Eupelmidae), a new predator of *Xylosandrus compactus*. *Journal of Coffee Research* 16: 35-37. (ec).
- VITE, JEAN PIERRE. 1979b. Borkenkäfer-lockstoff: mit fälle oder fangbaum gegen den buchdrucker? *Forstliche Mitteilungen* 32: 115. (bv).
- _____. 1979c. Mitt lockstoff-fallen gegen Nutzholzborkenkäfer. *Holz-Zentralblatt* 105: 1507-1509. (bv).
- _____. 1957. Das Borkenkäferproblem aus der sicht der Pheromoneinsatzes [The bark beetle problem from the standpoint of pheromone application] [English summary]. *Forstarchiv* 58: 239-243. (bv cn).
- _____. 1958a. Current issues in the pest management of scolytid bark beetles. Pages 1-2 in T. L. Payne and H. Saarenmaa, Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- _____. 1958b. Current issues in the pest management of scolytid bark beetles. *International Congress of Entomology, Proceedings* 18: 412. (cn).
- _____. 1959a. Compositions for combating bark beetles. U.S. Patent 4539383, 13 June 1959. Official Gazetteer, U.S. Patent Trademark Office 1103(2): 1262. (cn).
- _____. 1959b. The European struggle to control *Ips typographus*: past, present, and future. *Holarctic Ecology* 12: 520-525. (cn).
- VITE, JEAN PIERRE, RONALD FORREST BILLINGS, C. W. WARE, AND K. MORI. 1955. Southern pine beetle: enhancement or inhibition of aggregation response mediated by enantiomers of *endo*-brevicomin. *Naturwissenschaften* 72: 99-100. (bv).
- VITE, JEAN PIERRE, ROBERT IMRE GARA, AND H. D. VON SCHELLER. 1955. Southern pine beetle: enhancement or inhibition of aggregation response mediated by entiomers of *endo*-brevicomin. *Naturwissenschaften* 72: 99. (bv).
- VITE, JEAN PIERRE, L. B. GERKEN, AND GERALD NORMAN LANIER. 1976. Ulmensplintkafer: Anlockversuche mit synthetische pheromone im Overhaintal. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 83: 166-171. (bv).
- VITE, JEAN PIERRE, AND P. SAUERWEIN. 1979. Zum einsatz von lockstoff-fallen gegen borkenkäfer. *Gesunde Pflanzen* 31: 217-223. (bv).
- VOGEL, W. R. 1956. Zur Schwermetallbelastung der Borkenkäfer [Heavy metal contamination of bark beetles] [English summary]. *Entomologia Experimentalis et Applicata* 42: 259-269. (ay).
- _____. 1958. Die Belastung von Arthropoden mit Blei und Cadmium in unterschiedlich schadstoffexponierten Waldgebieten [The lead and cadmium burdens of arthropods in forest areas with different levels of exposure to pollutants] [English summary]. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 61: 205-216. (ay).
- VOGLER, DETLEV, AND J. DALE. 1952. Biological evaluation of tree mortality at Fort Bidwell Indian Reservation. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report S2-23. (cn).
- VOGLER, DETLEV, AND D. SCHULZ. 1955. A biological evaluation of pest conditions at Sierra Summit Ski area, Pineridge Ranger District, Sierra National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report S5-21. (cn).
- VOLTZ, H. A. 1958. Monoterpenes governing host selection in the bark beetles *Hylurgops palliatus* and *Tomicus piniperda*. *Entomologia Experimentalis et Applicata* 47: 31-35. (bv).
- VOOLMA, K. K. 1956. Entomophages of *Dendroctonus micans* in Estonia [In Russian, English summary]. *Metsanduslikud nurimused, Estonian SSR* 21: 89-97. (ec).
- *VORMAND, H. 1939. Contribution au catalogue des Coléoptères de la Tunisie. *Bull. Soc. Hist. Natur. Afr. Nord* 29: 481-515. ().
- VORONTSOV, A. I. 1955. Ways of development of forest entomology [In Russian]. *Zashchita Rastenii Moscow* 7: 23-24. (cn).
- VORONTSOV, A. I., ET AL. 1957. Protection of urban plantings. *Zashchita Rastenii Moscow* 7: 30-40. (cn).
- VOULAND, G., M. GIRAUD, AND DANIEL SCHVESTER. 1955. La période tenerale et l'envol chez *Dendroctonus micans* Kng. (Coleoptera, Scolytidae). Pages 68-79 in J. C. Gregoire and J. M. Pasteels, Biological control of bark beetles (*Dendroctonus micans*). Commission of European Communities and Université Libre de Bruxelles, Belgium, Proceedings of seminar, 3-4 October 1954. 141 p. (hb).
- VULINEC, KEVINA, AND R. A. DAVIS. 1954. Coleoptera types in the Charles Dury collection of the Cincinnati Museum of Natural History [Ohio, USA]. *Coleopterists Bulletin* 38: 232-239. (tx).
- VYPLEL, G. 1955. Untersuchungen zur Fortpflanzungs-Physiologie des Kupferstechers. *Allgemeine Forstzeitschrift* 12: 262-263. (ay).

W

- WAGNER, ERNST. 1954c. Entstehung und Verlauf der Borkenkäferplage 1945-1951 in Rheinland-Pfalz. Pages 165-177 in *Forstschutzstelle Sudwest, Ringingen*. 496 p. (cn).
- _____. 1954d. Beiträge zur Erforschung für die forstliche Praxis wichtiger Zusammenhänge der Borkenkäferkatastrophe 1945-1949 im Bad. Forstbezirk St. Blasien. Pages 191-224 in *Forstschutzstelle Sndwest, Ringingen*. 496 p. (cn ec).
- WAGNER, MICHAEL R., STEPHEN K. N. ATUAHENE, AND JOSEPH R. COBBINAH. 1958. Fauna of West Africa: forest insects of Ghana. [Date and publisher not indicated]

- on reprint, possibly 1987 or 1988. This might be a preliminary draft of: Forest Entomology in West Tropical Africa: Forest Insects of Ghana, 1991, Series Entomologica 47, 210 p.] Chapters 4-6: 93-225. (cn ds).
- WAGNER, TERENCE L., W. S. FARGO, R. O. FLAMM, ROBERT N. COULSON, AND P. E. PULLEY 1987. Development and mortality of *Ips calligraphus* (Coleoptera: Scolytidae) at constant temperatures. Environmental Entomology 16: 484-496. (cc hb).
- WAGNER, TERENCE L., R. O. FLAMM, AND ROBERT N. COULSON 1985. Strategies for cohabitation among the southern pine bark beetle species: comparisons of life-process biologies. Pages 87-101 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. General Technical Report SO-56. 383 p. (bv ec hb).
- _____. 1986. A temperature-dependent model of reemergence of *Ips calligraphus* (Coleoptera: Scolytidae). Canadian Entomologist 118: 901-911. (ec).
- WAGNER, TERENCE L., R. O. FLAMM, P. B. HENNIER, W. WU, AND ROBERT N. COULSON 1985. A temperature-dependent model of reemergence of *Ips avulsus* (Coleoptera: Scolytidae). Environmental Entomology 17: 192-198. (ec).
- WAGNER, TERENCE L., R. O. FLAMM, H. I. WU, W. S. FARGO, AND ROBERT N. COULSON 1987. Temperature-dependent model of life cycle development of *Ips calligraphus* (Coleoptera: Scolytidae). Environmental Entomology 16: 497-502. (ec hb).
- WAGNER, TERENCE L., P. B. HENNIER, R. O. FLAMM, AND ROBERT N. COULSON 1985. Development and mortality of *Ips avulsus* (Coleoptera: Scolytidae) at constant temperatures. Environmental Entomology 17: 181-191. (ec hb).
- WALL, R. E. 1988. Deterioration of severely defoliated balsam fir in relation to stand age, spacing and foliar protection. Canadian Journal of Forest Research 18: 490-497. (ec).
- WALSER, D. C. 1986. Processing dead timber into veneer and plywood. Pages 20-26 in R. W. Nielson, Harvesting and processing of beetle-killed timber. Proceedings of a symposium. Special publication, Forintek Canada Corp. SP-26. 53 p. (cn).
- WANG, CHANG-SHAN. 1981. A study of the Korean bark beetle (*Blastophagus pilifer* Spess.) [In Chinese]. Kunchong Zhishi 18: 165-167. (hb).
- *WANKA, THEODOR V. 1920. [*Hylastinus obscurus*]. Entomologische Blätter 16: 202-213. ().
- *WARBURTON. 1917. [*Scolytus rugulosus*]. Royal Agricultural Society of England, London, Journal 78: 209-219. ().
- WARD, J. D., C. W. DULL, G. W. RYAN, AND M. C. REMON. 1985. Estimating southern pine beetle caused timber losses over extensive areas. Pages 34-40 in S. J. Branham and R. C. Thatcher, Symposium proceedings: Integrated pest management research. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. General Technical Report SO-56. 383 p. (cn).
- WARREN, G. R. 1957. Sapwood decay systems of spruce budworm killed balsam fir in Newfoundland, Canada. Canadian Journal of Plant Pathology 9: 286-287. (ec).
- *WATERHOUSE, CHARLES OWEN 1874. Notice of a small collection of Coleoptera from Jamaica, with descriptions of new species from the West Indies. Entomological Society of London, Transactions 26: 303-311. ().
- *_____. 1854. On the coleopterous insects collected by Mr. H. O. Forbes in the Timor-Lont Islands. Zoological Society of London, Proceedings 1854: 213-218. ().
- WATERS, WILLIAM E. 1985a. Monitoring bark beetle populations and beetle-caused damage. Pages 141-175 in W. E. Waters, R. W. Stark, and D. L. Wood. Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn).
- _____. 1985b. The pine-bark beetle ecosystem: a pest management challenge. Pages 1-48 in W. E. Waters, R. W. Stark, and D. L. Wood. Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn ec hb).
- _____. 1985c. Workshop II: population and impact assessment. Discussion group C. Quantifying impacts on the other forest resources. Pages 255-261 in C. Sanders, R. W. Stark, E. J. Mullins, and J. Murphy, Recent advances in spruce budworm research. Minister of Supply and Services, Ottawa, Ontario. (cn).
- _____. 1986. A comprehensive approach to monitoring in management systems for forest pests. Forest Ecology and Management 15: 3-21. (cn).
- WATERS, WILLIAM E., AND ELLIS B. COWLING 1976. Integrated forest pest management: a silvicultural necessity. Pages 149-177 in J. L. Apple and R. F. Smith. Integrated pest management. Plenum Publishing Corporation, New York. (cn).
- WATERS, WILLIAM E., AND ROBERT W. GRAEBNER 1986. Sampling aerial photographs for estimation of pest-caused tree mortality. Western Journal of Applied Forestry 1: 84-89. (cn).
- WATERS, WILLIAM E., AND RONALD WILLIAM STARK 1980. Forest pest management: concept and reality. Annual Review of Entomology 25: 479-509. (cn).
- WATERS, WILLIAM E., RONALD WILLIAM STARK, AND DAVID LEE WOOD 1985. Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. 304 p. (cn).
- WAUTHIOZ, V. 1952. La lutte contre l'Hylobe est entree dans sa phase decisive. Societe Royale Forestiere de Belgique, Bulletin 59: 1-14. (ec).
- WAWRZYNSKI, ROBERT P., ANDREW D. KAHN, WHITNEY S. CRANSHAW, AND DAVID A. LEATHERMAN 1986. Insect pests of Colorado trees. Colorado State University Cooperative Extension Service, Bulletin 506a. 60 p. (cn).
- WEBBER, JOAN F. 1988. Impact of D-factor mycoviruses on infection by the elm pathogen *Ophiostoma ulmi*. Phytopathology 78(12, part 1): 1593. (ec).
- WEBBER, JOAN F., C. M. BRASIER, AND A. G. MITCHELL 1988. The role of the saprophytic phase in Dutch elm disease. Pages 298-313 in G. F. Pegg, and P. G. Ayres, Fungal infection of plants. Cambridge University Press, Cambridge, England. xiii + 428 p. (ec).
- WEBBER, JOAN F., AND J. N. GIBBS 1989. Insect dissemination of fungal pathogens of trees. Pages 161-193 in N. Wilding, N. M. Collins, P. M. Hammond, and J. F. Webber, Insect-fungus interactions. Academic Press, London. xvi + 344 p. (ec).
- WEBER, BARBARA CATHERINE 1984. The short- and long-term effects of insect attacks on trees. Pages 17-20 in M. M. Harris and A. M. Spearing, Research in forest productivity, use, and pest control. United States Department of Agriculture, Forest Service,

- Northeastern Station, General Technical Report NE-90. (cn).
- WEBER, BARBARA CATHERINE, AND JOHN EDWIN McPIERSON. 1985. Relation between attack by *Xylosandrus germanus* (Coleoptera: Scolytidae) and disease symptoms in black walnut. Canadian Entomologist 117: 1275-1277. (bv ec).
- _____. 1991. Seasonal flight patterns of Scolytidae (Coleoptera) in black walnut plantations in North Carolina and Illinois. Coleopterists Bulletin 45: 45-56. (hb ds).
- WEBER, H. C. 1942. Division of Forestry. Pages 64-97 in Statistical report for biennium ending June 30, 1942. Minnesota Department of Conservation. (cn).
- *WEBSTER, FRANCIS MARION. 1893a. [*Hylastinus obscurus*]. Indiana Academy of Science, Proceedings 1893: 84. ().
- *_____. 1893b. [*Hylastinus obscurus*]. Ohio Agricultural Experiment Station, Bulletin 51: 120. ().
- *_____. 1893c. [*Phloeotribus lininaris*]. Canadian Entomologist 25: 111. ().
- *_____. 1894. [*Hylastinus obscurus*]. Ohio Agricultural Experiment Station. Annual Report 1894: 31. ().
- *_____. 1895. [*Hylastinus obscurus*, *Scolytus rugulosus*]. Ohio Agricultural Experiment Station, Bulletin 68: 23-25, 31-33, figs. 3-6. ().
- *_____. 1899. [*Scolytus rugulosus*]. Ohio Farmer 1899(25 May): 449. ().
- WEGENSTEINER, R., E. LORBEER, AND E. FUEHRER. 1889. Additional attraction of *Pityogenes chalcographus* L. (Coleoptera, Scolytidae) to chalcopras-baited traps with spruce branches. Journal of Applied Entomology 108: 250-259. (bv).
- WEI, HOU-JIAN. 1959. Methods for observing habits of bark beetles inside their host tree [In Chinese]. Kunchong Zhishi 5: 237-238. (hb ms).
- _____. 1960. Methods for observing habits of *Blastophagus piniperda* Linnaeus inside the bark of trees [In Chinese]. Kunchong Zhishi 6: 117-118, 131. (hb ms).
- WEIR, H. J., AND W. D. BIGGS. 1974. Forest insect and disease conditions in the southwest survey region, 1973. Canada Department of Fisheries and Forestry, Forestry Service, Great Lakes Forestry Research Centre, Sault Ste. Marie, Ontario, Information Report O-X-194. 26 p. (cn).
- WEIR, H. J., M. J. THOMSON, D. C. CONSTABLE AND C. G. JONES. 1986. A review of important forest insect and disease problems in the Kirkland district of Ontario, 1950-1980. Canada Department of Environment, Forestry Service, Great Lakes Forest Research Centre, Sault Ste. Marie, Ontario, Miscellaneous Report 34. 133 p. (cn).
- *WEISS, HARRY B. 1915. [*Scolytus rugulosus*]. Canadian Entomologist 47: 165-166. ().
- WELCH, D. S., G. W. HERRICK, AND R. W. CURTIS. 1934. The Dutch elm disease. Cornell University Extension Service, Bulletin 290. 19 p. (cn).
- WELCH, R. COLIN. 1968b. The Coleoptera of Monks Wood National Nature Reserve, Huntingdonshire, first supplement, 1965. Entomologist's Gazette 19: 9-20. (ds).
- _____. 1968c. Some Coleoptera from Loch Lomond National Nature Reserve. Entomologist's Monthly Magazine 104: 119-122. (ds).
- _____. 1970. The Coleoptera of Monks Wood National Nature Reserve, Huntingdonshire, second supplement, 1966-1968. Entomologist's Gazette 21: 133-141. (ds).
- _____. 1980. The Coleoptera of Monks Wood National Nature Reserve, Cambridgeshire (Vice County), fifth supplement, 1974-1979. Entomologist's Gazette 31: 263-273. (ds).
- *_____. 1983. Coleoptera in the Inner Hebrides. Royal Society of Edinburgh, Proceedings 83B: 505-529. ().
- _____. 1987. *Xyleborus saxeseni* (Ratz.) (Col., Scolytidae) attracted to blue ear in forest of Dean, Gloucestershire. Entomologist's Monthly Magazine 123: 161. (bv).
- WENZ, JOHN. 1984. Posttreatment evaluation: Jeffrey pine beetle suppression project, South Shore Recreation Complex, Lake Tahoe Basin Management Unit. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 84-02. (cn).
- _____. 1985. Evaluation of bark beetle conditions in the Glenbrook Burn, Lake Tahoe Basin Management Unit. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 85-19. (cn).
- WENZ, JOHN, AND G. DENITTO. 1983. Evaluation of pest conditions in the Kiva/Taylor Creek Visitor Center Recreation Areas on the Lake Tahoe Basin Management Unit. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-06. (cn).
- WENZ, JOHN, AND J. PRONOS. 1983. An evaluation of insect and disease conditions in mixed conifer stands underburned for fuels reduction on the Quincy Ranger District, Plumas National Forest, United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-02. (cn).
- WENZ, JOHN, AND R. S. SMITH, JR. 1983. An evaluation of the Jackass Insect Salvage Sale Area, Truckee Ranger District, Tahoe National Forest. United States Department of Agriculture, Forest Service, Pacific Southwest Region, San Francisco, California, Forest Pest Management Report 83-04. (cn).
- WERNER, RICHARD ALLEN. 1986a. The eastern larch beetle in Alaska. United States Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station, Portland, Oregon. Research Paper PNW-357. ii + 13 p. (bv hb).
- _____. 1986b. Association of plants and phytophagous insects in taiga forest ecosystems. Pages 205-212 in K. Van Cleve, F. S. Chapin III, P. W. Flanagan, et al., Forest ecosystems in the Alaskan taiga: a synthesis of structure and function. Springer, New York. (cn ec).
- _____. 1988a. Recommendations for suppression of an *Ips perturbatus* outbreak in interior Alaska using integrated control. Pages 189-195 in T. L. Payne and H. Saarenmaa. Integrated control of scolytid bark beetles. Virginia Polytechnic and State University, Blacksburg, Virginia. 355 p. (cn).
- _____. 1988b. Suppression of an *Ips perturbatus* outbreak in interior Alaska using integrated control. International Congress of Entomology, Proceedings 18: 414. (cn).
- WERNER, RICHARD ALLEN, JOHN S. HARD, AND EDWARD H. HOLSTEN. 1985. The development of management strategies to reduce the impact of the spruce beetle in south-central Alaska. Northwest Environmental Journal 4: 319-358. (cn).

- WERNER, RICHARD ALLEN, FELTON L. HASTINGS, EDWARD H. HOLSTEN, AND ALICE S. JONES. 1986. Carbaryl and lindane protect white spruce from attack by spruce beetles (Coleoptera: Scolytidae) for three growing seasons. *Journal of Economic Entomology* 79: 1121–1124. (cn).
- WERNER, RICHARD ALLEN, AND EDWARD H. HOLSTEN. 1985a. Effect of phloem temperature on development of spruce beetles in Alaska. Pages 155–163 in L. Safranyik, The role of the host in the population dynamics of forest insects. IUFRO conference proceedings, Canada Department of the Environment, Forestry Service, and United States Department of Agriculture, Forest Service, Banff, Alberta, Canada, September, 1983. 240 p. (cc lib).
- . 1985b. Factors influencing generation times of spruce beetles in Alaska. *Canadian Journal of Forest Research* 15: 438–443. (cc lib).
- WERNER, RICHARD ALLEN, EDWARD H. HOLSTEN, AND FELTON L. HASTINGS. 1986. Evaluation of pine oil for protecting white spruce from spruce beetle (Coleoptera: Scolytidae) attack. *Entomological Society of British Columbia Journal* 83: 3–5. (cn).
- WESLIEN, JAN. 1989. The role of trapping in population management of the spruce bark beetle *Ips typographus* L. Dissertation, Swedish University of Agricultural Sciences, Uppsala. 98 p. (cn).
- WESLIEN, JAN, ERIKKI ANNILA, ALF BAKKE, BRODER BEJER PETERSEN, HUBERTUS H. EIDMANN, K. NARVESTAD, A. NIKULA, AND H. P. RAVN. 1989. Estimating risks for spruce bark beetle [*Ips typographus* (L.)] damage using pheromone-baited traps and trees. *Scandinavian Journal of Forest Research* 4: 87–98. (cn).
- WESLIEN, JAN, AND H. BYLUND. 1988. The number and sex of spruce bark beetles *Ips typographus* L. caught in pheromone traps as related to flight season trap type and pheromone release. *Journal of Applied Entomology* 106: 488–493. (bv cn lib).
- WESLIEN, JAN, AND A. LINDELOW. 1989. Trapping a local population of spruce bark beetles *Ips typographus* (L.): population size and origin of trapped beetles. *Holarctic Ecology* 12: 511–514. (cn lib).
- WESTCOTT, C. 1970. The gardener's bugbook. Doubleday & Co., Inc., Garden City, New York. 625 p. (cn).
- WETTSTEIN, OTTO. 1959. Die Lebensgemeinschaft an einem Birkenstamm. *Entomologisches Nachrichtenblatt Österreichischer und Schweizer Entomologen* 11: 42–59. (cc).
- WHITEHEAD, ARMAND TOYN. 1986. Electroantennogram responses by mountain pine beetles, *Dendroctonus ponderosae* Hopkins, exposed to selected semiochemicals. *Journal of Chemical Ecology* 12: 1603–1621. (ay bv).
- WHITEHEAD, ARMAND TOYN, D. T. SCOTT, RICHARD FRANKLIN SCHMITZ, AND KENJI MORI. 1989. Electroantennogram responses by mountain pine beetles, *Dendroctonus ponderosae* Hopkins, exposed to selected chiral semiochemicals. *Journal of Chemical Ecology* 15: 2089–2100. (ay bv).
- *WHITEHEAD, W. E. 1932. The morphology of the head-cap-sule of some coleopterous larvae. *Canadian Journal of Research* 6: 227–252. ().
- WHITESSELL, J. K., AND C. M. BUCHANAN. 1986. Synthesis of (-)- and (+)-frontalin. *Journal of Organic Chemistry* 51(26): 5443–5445. (bv cn).
- WHITNEY, HARVEY S., R. J. BANDONI, AND F. OBERWINKLER. 1987. *Entomocorticium dendroctoni* gen. et sp. nov. (Basidiomycotina), a possible nutritional symbiote of the mountain pine beetle in lodgepole pine in British Columbia. *Canadian Journal of Botany* 65: 95–102. (cc).
- WHITNEY, HARVEY S., AND M. M. Z. KHARADY. 1984. Some results on low-level microwave treatment of the mountain pine beetle and the darkling beetle. *IEEE Transactions Microwave Theory Technology* 32: 798–803. (cc).
- WHITTLE, K., AND DONALD MORGAN ANDERSON. 1987. Pests not known to occur in the United States or of limited distribution. European spruce beetle. United States Department of Agriculture, Animal and Plant Health Inspection Service (Hyattsville, Maryland). 12 p. (cn ds tx).
- *WICHMANN, HEINRICH E. 1916b. [*Kissophagus orinacellus*, *Scolytus pygmaeus*]. *Zeitschrift für Pflanzenkrankheiten* 26: 432–433. ().
- *WICKHAM, HENRY FREDERICK. 1893. Report on an entomological reconnaissance of southern Alaska and adjacent portions of British Columbia. Laboratory of Natural History, State University of Iowa, Bulletin 2(3): 202–233. ().
- *———. 1909. A list of the Coleoptera of Iowa. Laboratory of Natural History, State University of Iowa, Bulletin 6(1): 1–40. ().
- *WICKMAN, BOYD E. 1990. The battle against bark beetles in Crater Lake National Park, 1925–1934. United States Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, Oregon. General Technical Report GTR-259. 40 p. ().
- *WILANSKY. 1920. [*Phloeotribus piccae* (=lecontei), *Hylestinus oleiperda* (=californicus)]. California State Department of Agriculture, Monthly Bulletin 9: 339–340. ().
- WILKEN, C. 1864. Zur Fauna des Oberharzes. *Berliner Entomologische Zeitschrift* 8: 369–373. (ds).
- WILKINSON, D. S. 1935. On some braconids (Hym.). *Stylops: a Journal of Taxonomic Entomology* 4: 71–72. (cc).
- WILKINSON, ROBERT CLEVELAND, AND ROBERT A. HAACK. 1989. Within-tree distribution of pine bark beetles (Coleoptera: Scolytidae) in Honduras. *Ceiba* 28: 115–133. (cc ds).
- *WILLE, JOHANNES E. 1943. [*Pagiocerus rimosus* (=frontalis)]. *Estac. Expt. Agric. de Molina*, Lima, Peru. 1943: 364–366. ().
- WILLIAMS, J. O. 1988. Occurrence of *Xyleborus ferrugineus* (Fabr.) (Coleoptera: Scolytidae) on yam tubers in Nigeria. *Insect Science and its Application* 9: 41–42. (cn).
- WILLIAMS, LONNIE H. 1985. Wood-inhabiting insects and their control: producer and user viewpoints. Pages 67–76 in Wood protection techniques and the use of treated wood in construction. United States Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, Louisiana. Proceedings 4725S. (cn).
- WILLIAMS, NANCY E., J. A. WESTGATE, D. D. WILLIAMS, ANNE MORGAN, AND ALAN V. MORGAN. 1981. Invertebrate fossils (Insecta: Trichoptera, Diptera, Coleoptera) from the Pleistocene Scarborough Formation at Toronto, Ontario, and their paleoenvironmental significance. *Quaternary Research* 16: 146–166. (ds).
- WILLIAMS, T. M., C. A. GERSHAM, R. L. HEDDEN, AND E. R. BLOOD. 1989. Impact of southern pine beetle infestation on nutrient retention mechanisms in loblolly pine. United States Department of Agriculture, Forest Service, Southern Forest Experiment Sta-

- tion, New Orleans, Louisiana. General Technical Report SO-74: 465-472. (cn).
- *WILLSON 1937. [*Scolytus rugulosus*]. Canadian Journal of Research, Section C, 15: 331. ().
- WILSON, HARLEY FROST. 1915. Orchard insect pests and methods of control. Pacific Horticultural Correspondence School, Portland, Oregon. Meade Press, Orenco, Oregon. (cn lib).
- WILSON, LOUIS F. 1962. Insect damage to field-piled pulpwood in northern Minnesota. Journal of Economic Entomology 55: 510-516. (cn).
- WINGFIELD, M. J., L. A. STRAUSS, AND G. D. TRIBE. 1985. Fungi associated with three pine bark beetles in South Africa. Phytopathology 75: 133S. (ec).
- WINGFIELD, M. J., P. S. VAN WYK, AND P. W. J. VAN WYK. 1989. Conidial development in the anamorph of *Ophiostoma cucullatum*. Mycological Research 93: 91-95. (ec).
- WINKLER, ALBERT. 1932. Catalogus Coleopterorum regionis palaearctica [Scolytidae by K. E. Schedl, p. 1633-1647]. Wien, Austria. 13 parts in 4 volumes. (ds).
- WINTER, T. G. 1955. Is *Ips typographus* (Linnaeus) (Coleoptera: Scolytidae) a British insect? Entomologist's Gazette 36: 153-160. (ds).
- WINTER, T. G., AND D. A. BURDEKIN. 1987. The poem and the pest. Quarterly Journal of Forestry 81: 234-238. (ms).
- WISNIEWSKI, JERZY. 1950. Zwei neue mit *Proctolaelaps xyloteri* Sams. (Mesostigmata, Blattisociidae) verwandte Proctolaelaps-Arten aus Polen. Acarologia 21: 3-8. (ec).
- WISNIEWSKI, JERZY, AND JACEK MICHALSKI. 1983. Gangsystematik der parasitiformes, Teil 432. Studium einer neuen *Dendrolaelaps*-Art aus Sibirien. Acarologie 30: 101-103. (ec).
- WITANACHIT, JAYANTHI P. 1986. Response of *Ips grandicollis* (Eichhoff) adults (Coleoptera: Scolytidae) to sites in bark initiated by others of the same species. Australian Entomological Society, Journal 25: 41-46. (bv).
- WITCOSKY, J. J. 1959. Root beetles, stand disturbance, and management of black-stain root disease in plantations of Douglas-fir. Pages 55-70 in R. I. Alfaro and S. G. Clover, Insects affecting reforestation: biology and damage. Government of Canada, Canadian Forestry Service, Pacific and Yukon Region, Victoria, British Columbia. (cn ec).
- WITCOSKY, J. J., AND EVERETTE HANSEN. 1985. Root-colonizing insects recovered from Douglas-fir in various stages of decline due to black-stain root disease. Phytopathology 75: 399-402. (ec).
- WITCOSKY, J. J., TIMOTHY D. SCHOWALTER, AND EVERETTE M. HANSEN. 1986a. *Hylastes nigrinus* (Coleoptera: Scolytidae), *Pissodes fasciatus*, and *Stereomus carinatus* (Coleoptera: Curculionidae) as vectors of black-stain root disease of Douglas-fir. Environmental Entomology 15: 1090-1095. (ec).
- _____. 1986b. The influence of time of precommercial thinning on the colonization of Douglas-fir by three species of root-colonizing insects. Canadian Journal of Forest Research 16: 745-749. (bv).
- _____. 1987. Host-derived attractants for the beetles *Hylastes nigrinus* (Coleoptera: Scolytidae) and *Stereomus carinatus* (Coleoptera: Curculionidae). Environmental Entomology 16: 1310-1313. (bv).
- *WOLCOTT, GEORGE N. 1933. [*Scolytus rugulosus* parasites]. Economic Entomology of West Indies 1933: 201-202. ().
- _____. 1945. The insects of Puerto Rico. Journal of Agriculture of the University of Puerto Rico 32: 1-975. (ds).
- WOLFE, MAX. 1927. Unsere Käfer. Ullstein, Berlin. (ds).
- WOLLASTON, THOMAS VERNON. 1860b. On additions to the Madeiran Coleoptera [Scolytidae, p. 110]. Annals and Magazine of Natural History 3(6): 100-111. (ds tx).
- _____. 1861. Certain coleopterous insects from the island of Ascension [Scolytidae, p. 303-304]. Annals and Magazine of Natural History 3(7): 299-304. (ds tx).
- _____. 1871. On additions to the Atlantic Coleoptera [Scolytidae, p. 262-264]. Entomological Society of London, Transactions 19: 203-314. (ds tx).
- _____. 1877. Coleoptera Sanctae-Helenae [Scolytidae, p. 79-81]. Van Voorst, London. xxv + 256 p. (ds tx).
- WOOD, COLIN S., AND G. A. VAN SICKLE. 1987. Forest insect and disease conditions: British Columbia & Yukon 1986. Government of Canada, Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-257. 35 p. (cn).
- _____. 1989. Forest insect and disease conditions: British Columbia & Yukon 1988. Government of Canada, Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-306. 33 p. (cn).
- WOOD, COLIN S., G. A. VAN SICKLE, AND L. M. HUMBLE. 1987. Forest insect and disease conditions: British Columbia & Yukon 1987. Government of Canada, Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-296. 40 p. (cn).
- WOOD, COLIN S., G. A. VAN SICKLE, AND T. L. SHORE. 1985. Forest insect and disease conditions: British Columbia & Yukon 1984. Government of Canada, Forestry Service, Pacific Forestry Centre, Victoria, British Columbia, Information Report BC-X-259. 32 p. (cn).
- WOOD, DAVID LEE. 1978. Status of behavior-modifying chemicals (IMC) in forest insect management. Pages 66-73 in Proceedings of the 29th annual Western Forest Insect Work Conference, Durango, Colorado, 7-8 March 1978. (bv cn).
- _____. 1979a. Attraction—Coleoptera: Disruption—Coleoptera. Pages 21-32, 91-96 in W. L. Roelofs, Establishing efficacy of sex attractants and disruptants for insect control. Entomological Society of America, Miscellaneous Publication. (bv cn).
- _____. 1979b. The use of microbial agents and unconventional chemicals in forest pest management. Pages 25-33 in R. D. Gail, The Pinchot Institute for Conservation Studies/ Proceedings for Integrated Pest Management Colloquium. United States Department of Agriculture, Forest Service, Technical Report WO-14. (bv cn ec).
- _____. 1981. Obituary: Julius A. Rudinsky, 1917-1980. Entomological Society of America, Bulletin 27: 292-293. (ms).
- _____. 1984a. Book review: M. C. Birch and K. F. Haynes, Insect pheromones (Studies in Biology No. 147, Edward Arnold, Limited, London and Baltimore, 1982. 58 p.). Journal of Chemical Ecology 10: 189-191. (bv ms).
- _____. 1984b. Book review: J. B. Mitton and K. B. Sturgeon, Bark beetles in North American conifers: a system for the study of evolutionary biology (University of Texas Press, Austin. 527 p.). Quarterly Review of Biology 59: 197-198. (ms).
- _____. 1988. Forest health management for the future: insect problems of coniferous forests. Society of

- American Foresters, Proceedings 1958: 114–120. (cn ms).
- WOOD, DAVID LEE, R. PATRICK AKERS, DONALD R. OWEN, AND JOHN R. PARMETER, JR. 1986. The behavior of bark beetles colonizing ponderosa pine. Pages 91–103 in B. E. Juniper and T. R. E. Southwood. Insects and the plant surface. Edward Arnold, Limited, London. (bv ec).
- WOOD, DAVID LEE, JOHN H. PARMETER, JR., AND DONALD L. DAHLSTEN. 1988. Interrelationships among *Dendroctonus* and *Ips* bark beetles, their vectored fungi and host physiology. International Congress of Entomology, Proceedings 18: 410. (ec).
- WOOD, DAVID LEE, AND P. A. RAUCH. 1980. Approach to research and forest management for the western pine beetle: principal research results. Pages 429–442 in C. B. Huffaker, New technology of pest control. John Wiley & Sons, New York. (cn).
- WOOD, DAVID LEE, RONALD WILLIAM STARK, WILLIAM E. WATERS, WILLIAM DELLES BEDARD, JR., AND FIELDS WHITE COBB, JR. 1985. Treatment tactics and strategies. Pages 121–139 in W. E. Waters, R. W. Stark, and D. L. Wood, Integrated pest management in pine-bark beetle ecosystems. John Wiley & Sons, New York. xvii + 256 p. (cn).
- WOOD, DAVID LEE, ROBERT MILTON SILVERSTEIN, AND MINORU NAKIJIMA. 1970. Control of insect behavior by natural products. Academic Press, New York. (cn).
- WOOD, STEPHEN LANE. 1961e. An alternative proposal to the suggested validation of *Myelophilus* Eichhoff, 1878 (Insecta: Coleoptera). Bulletin of Zoological Nomenclature 18: 319–321. (tx).
- _____. 1972i. Family Platypodidae. Page 394 in M. H. Hatch. The beetles of the Pacific Northwest, Part 5. University of Washington, Publications in Biology No. 16. 662 p. (ds tx).
- _____. 1984e. New synonymy and new species of American bark beetles (Coleoptera: Scolytidae), part X. Great Basin Naturalist 44: 113–119. (tx).
- _____. 1985. New synonymy and new species of bark beetles (Coleoptera: Scolytidae). Great Basin Naturalist 45: 266–275. (tx).
- _____. 1986a. A reclassification of the genera of Scolytidae (Coleoptera). Great Basin Naturalist Memoirs 10. 126 p. (tx).
- _____. 1986b. New *Pseudoxylechinus* (Coleoptera: Scolytidae) from India. Great Basin Naturalist 43: 468. (tx).
- _____. 1986c. New synonymy and new species of American bark beetles (Coleoptera: Scolytidae), part XI. Great Basin Naturalist 46: 265–273. (tx).
- _____. 1987. Six new Scolytidae (Coleoptera) from Mexico. Great Basin Naturalist 47: 547–550. (tx).
- _____. 1988a. Nomenclatural changes and new species of Scolytidae (Coleoptera). Great Basin Naturalist 48: 31–38. (tx).
- _____. 1988b. Nomenclatural changes and new species of Scolytidae (Coleoptera), part II. Great Basin Naturalist 48: 188–195. (tx).
- _____. 1988c. Nomenclatural changes and new species of Scolytidae (Coleoptera), part III. Great Basin Naturalist 48: 196–201. (tx).
- _____. 1988d. Systematic position of the Scolytidae and Platypodidae (Coleoptera). International Congress of Entomology, Proceedings 18: 40. (tx).
- _____. 1988e. Recent advances in knowledge of the distribution and classification of the Scolytidae (Coleoptera). International Congress of Entomology, Proceedings 18: 410. (tx).
- _____. 1989. Nomenclatural changes and new species of Scolytidae (Coleoptera), part IV. Great Basin Naturalist 49: 167–185. (tx).
- _____. 1992a. Nomenclatural changes and new species of Platypodidae and Scolytidae (Coleoptera), part II. Great Basin Naturalist 52: 78–88. (tx).
- _____. 1992b. Nomenclatural changes in Scolytidae and Platypodidae (Coleoptera). Great Basin Naturalist 52: 89–92. (tx).
- _____. ex1993. Revision of the genera of Platypodidae (Coleoptera). Great Basin Naturalist 53: (in review).
- WOOD, STEPHEN LANE, AND DONALD EDWARD BRIGHT, JR. 1987. A catalog of Scolytidae and Platypodidae (Coleoptera), part 1: bibliography. Great Basin Naturalist Memoirs 11. 685 p. (ms).
- WOOD, STEPHEN LANE, AND FU-SHENG HUANG. 1986. New genus of Scolytidae (Coleoptera) from Asia. Great Basin Naturalist 46: 465–467. (tx).
- WOOD, STEPHEN LANE, GEORGE C. STEVENS, AND HUMBERTO J. LEZAMA. 1991a. Los Scolytidae de Costa Rica: clave de generos y de la subfamilia Hylesinae (Coleoptera). Revista de Biología Tropical 39: 125–148. (ds tx).
- _____. 1991b. Scolytidae (Coleoptera) de Costa Rica II. Clave para la subfamilia Scolytinae, tribus: Scolytini, Ctenophorini, Micracini, Ipini, Dryocoetini, Xyleborini y Cryphalini. Revista de Biología Tropical 39: 279–306. (ds tx).
- WOOD, STEPHEN LANE, AND HUI-FEN YIN. 1986. Relict occurrence of three "American" Scolytidae (Coleoptera) in Asia. Great Basin Naturalist 46: 461–464. (tx).
- WOODSON, GEORGE. 1955. Utilization of beetle-killed southern pine. United States Department of Agriculture, Forest Service, Forest Pest Management (Washington, D.C.), General Technical Report WO-47. 27 p. (cn).
- *WORNLE, A. 1950. Die kafer von Nordtirol. Pages 376–383 in Universitätsverlag Wagner, Innsbruck-München. ().
- WORRALL, J. J., AND T. C. HARRINGTON. 1988. Etiology of canopy gaps in spruce-fir forests at Crawford Notch, New Hampshire. Canadian Journal of Forest Research 18: 1463–1469. (cn).
- WITTEWAAL, J. 1964. Volksleesboek over schadelijke en nuttige insekten. Hoitsema, Groningen. (cn hb).
- WULF, A. 1955. Zur Umweltverträglichkeit von Borkenkäferbekämpfungsmitteln [Environmental compatibility of bark beetle control agents]. Allgemeine Forstzeitschrift 12: 265–266. (cn).

X

- *XAMBEU, VINCENT. 1908. Faune Entomologique des Pyrenées-orientales, supplement aux Hemipteres-Homopteres du Catalogue de la Faune des Environs de Ria [Scolytidae, p. 96–97]. L'Echange. ().

Y

- YADAV, J. S., V. VIDYASAGAR, AND P. S. REDDY. 1986. Enantiospecific synthesis of dextro exo-brevicomin from D-xylose. Carbohydrate Research 156: 236–240. (bv ms).
- *YACDYEV, A. 1987. Pests of ornamental plants in towns of the Turkmen SSR USSR [In Russian]. [Source?] 1: 47–50. ().
- *YAKAITIS, B. YU. 1985. Nine species of Coleoptera new to the Lithuanian SSR, found in 1979–1984 [In Rus-

- sian, English summary]. Pages 14–18 in V. Ionaitis and V. Petrauskas. ().
- YAKAITIS, B. YU. AND V. M. GAVYALIS. 1985. The stationing of barrier traps with attractive substances for the regulation of numbers of the engraver beetle [In Russian, English summary]. *Acta Entomologica Lituania* 5: 53–57. (cn).
- _____. 1987. On the question of the improvement of barrier traps for the attraction of bark beetles (*Ips typographus* L.) [In Russian, English summary]. *Khemoretseptsiya Nasekomykh* 9: 118–122. (cn).
- YAMAOKA, Y., R. H. SWANSON, AND Y. HIRATSUKA. 1987. Inoculation experiments of lodgepole pine with four blue-stain fungi associated with mountain pine beetle monitored by a heat pulse velocity VPH instrument. *Canadian Journal of Plant Pathology* 9(3): 287. (ec).
- YANG, J., AND L. JI. 1989b. Amount of Masson pine pests in acid rain area. *Journal of Ecology* 5: 54–55. (cn ec).
- YANG, ZHONGQI. 1987. A preliminary survey of parasitic wasps of *Dendroctonus armandi* Tsai & Li (Coleoptera, Scolytidae) in Quinling Mountains with description of three new species and a new Chinese record (Hymenoptera: Pteromalidae). *Entomotaxonomia* 9(3): 175–184. (ec).
- _____. 1989a. A new species of *Ropalophorus* Curtis (Hymenoptera: Braconidae) parasitizing *Polygraphus poligraphus* adult (Coleoptera: Scolytidae) with a key to the world species of the genus [In Chinese, English summary]. *Entomotaxonomia* 11(1–2): 91–96. (ec).
- _____. 1989b. One new species and other pteromalids parasitizing bark beetles in Shaanxi, China (Hymenoptera, Chalcidoidea, Pteromalidae). *Entomotaxonomia* 11(1–2): 101–103. (ec).
- _____. 1989c. *Dendroctonus armandi* Tsai & Li (Coleoptera: Scolytidae) in China: its natural enemies and their potential as biological control agents. Pages 147–156 in D. L. Kulhavy and M. C. Miller, Potential for biological control of *Dendroctonus* and *Ips* bark beetles. Center for Applied Studies, School of Forestry, Stephen F. Austin State University, Nacogdoches, Texas. 255 p. (cn ec hb).
- *YANOVSKII, V. M. 1980. K faune lesnykh nasekomykh Mongolii [On the fauna of forest insects of Mongolia]. *Nasekomye Mongolii*, Nauka, Leningrad 7: 17–27. ().
- *_____. 1982. Vozdejstvie entomofagov na dinamiku cislennosti korojedov. Ctenia pamjati Nikolaja Alexandrovici Cholodkovskovo, Nauka, Leningrad 1982: 25–53. ().
- _____. 1986. Natural enemies of bark beetles [In Russian, English summary]. *Zashchita Rastenii* Moscow 1986: 26–29. (ec).
- _____. 1989. Forest Coleoptera of the west part of the zone of Kansk-Achinsk'fiel energetic complex (Krasnoyarsk Territory) [In Russian]. *Entomologicheskoe Obozrenie* 68: 57–67. (ds).
- YANOVSKII, V. M., AND D. TEGSIZHARGAL. 1985. Bark beetles (Coleoptera, Scolytidae) of the Mongolian People's Republic [In Russian]. Pages 404–417 in *Insects of Mongolia*. Number 9. Academy of Sciences of the USSR, Zoological Institute. 576 p. (ds).
- YIN, HUI-FEN. 1983a. A brief account on the study of symbiotic relationship of *Xyleborus* spp. with fungi [In Chinese]. *Forest Pest and Disease* 1983(2): 26–30. (ec).
- _____. 1983b. Some quarantine problems of exotic scolytids in the imported wood [In Chinese]. *Forest Pest and Disease* 1983(1): 36–38, 42. (cn).
- YIN, HUI-FEN, HUANG FU-SHENG, AND LI ZHAO-LIN. 1984. Coleoptera: Scolytidae [In Chinese]. *Economic Insect Fauna of China*, Fasc. 29. Science Press, Beijing. 205 p. (hb ds tx).
- YOSHIDA, N., Y. AKITA, C. KOIZUMI, AND K. FUKUYAMA. 1986. Field test of pheromone trap for *Ips typographus* in the windthrown forest caused by a typhoon. *Annual Report, Hokkaido Branch, Forestry and Forest Products Research Institute (Sapporo)* 1985: 54–55. (bv cn).
- YOSHIKAWA, K. 1987a. A study of the subcortical insect community in pine trees, II: vertical distribution. *Applied Entomology and Zoology* 22: 195–206. (ec).
- _____. 1987b. A study of the subcortical insect community in pine trees, III: species correlation. *Applied Entomology and Zoology* 22: 207–215. (ec).
- YOSHIKAWA, K., H. TAKEDA, K. SONE, AND E. I. SHIBATA. 1986. A study of the subcortical insect community in pine trees, I: oviposition and emergence periods of each species. *Applied Entomology and Zoology* 21: 255–265. (bv ec hb).
- YOUNG, CAMERON. 1985. Coming of age in the Flathead: how the British Columbia Forest Service contended with the mountain pine beetle infestation of southeastern British Columbia 1976–1986. *British Columbia Ministry of Forests, Pest Management Report No.* 10. 31 p. (cn).
- YTSMÄ, G. 1986. Inducing attack by male *Platypus* (Col., Platypodidae) on wood billets in the laboratory. *Journal of Applied Entomology* 102: 210–212. (bv ms).
- _____. 1988a. Pheromone traps for pinhole borer management. What's new in forest research, No. 165. *Forest Research Institute, Minister of Forestry, New Zealand*. 4 p. (bv cn).
- _____. 1988b. Stridulation in *Platypus apicalis*, *P. caviceps*, and *P. gracilis* (Col., Platypodidae). *Journal of Applied Entomology* 105: 256–261. (bv hb).
- _____. 1989. Colonization of southern beech by *Platypus caviceps* (Coleoptera: Platypodidae). *Journal of Chemical Ecology* 15: 1171–1176. (bv cn ec).
- YU, C. M., AND Q. H. ZHANG. 1988. Biological activity and excretion law of aggregation pheromone of larch bark beetle [In Chinese, English summary]. *Journal of Northeast Forestry University, China* 16: 1–7. (bv).
- *YUNAR, K. H. 1986. Species composition of Hymenopteran parasites of bark beetles on conifers in Estonia [In Russian, English summary]. *Metsanduslikus Uurimused, Estonian SSR* 21: 79–85. ().

Z

- ZACH, PETER. 1991. Anwendung von Photoelektroden beim Studium der kambio- und xylophagen Coleopteren. *Anzeiter für Schädlingkunde, Pflanzenschutz, Umweltschutz* 64: 34–37. (ec ms).
- *ZAHRADNIK, P., AND MILOŠ KNIZEK. 1990. Bronci ve feromonových lapacích na kůrovce (Scolytidae) [Kafer (Col.), die in pheromonephallen für Borkenkäfer (Scolytidae) getroffen]. *Sborník referátů, III. colloquium coleopterologicum 1990, Sumava, Zelená Ruda* 1990: 33–36. ().
- *ZAHRADNIK, P., MILOŠ KNIZEK, P. KAPITOLA, A. RODZIEWICZ, AND A. KOLK. 1990. Porovnání účinnosti používaných typu feromonových odpárníků k lákání lykožroutů smrkového (*Ips typographus*) [Comparison of effects of pheromones which are used in catching of

- the bark beetle *Ips typographus*]. Zpravy lesnického výzkumu 4: 23–27. ().
- * _____. 1991. Porovnání odchytových vlastností nových typů feromonových lapací na lýkožravce smrkového (*Ips typographus* L.) [Comparison of catching characteristics of a new type of pheromone traps for *Ips typographus*]. Zpravy lesnického výzkumu 1: 7–15. ().
- ZAKLAMA, S. F., A. M. AWADALLAH, AND F. F. ABDALLAH. 1979. Chemical control of the shot-hole borer, *Scolytus amygdali* Gner, on almond trees. Entomological Society of Egypt, Bulletin 11: 157–192. (cn).
- *ZESCH, F., AND O. REINECKE. 1881. List of the Coleoptera observed and collected in the vicinity of Buffalo. Buffalo Society of Natural Science, Bulletin 4: 2–15. ().
- ZHAD, J. N., AND B. CAO. 1987. Bionomics and control of *Phloeosinus perlatus* Chapuis [In Chinese, English summary]. Insect Knowledge 24(4): 227–230. (cn hb).
- ZHANG, LU-HONG, YU CHENG-MING, AND TAN GUI-ZHONG. 1955. A discussion on the control of some bark beetle species infesting forests in Dailing, northeast. Kunchong Zhishi 4(1): 27–30. (cn).
- ZHAO, KU-JIE, SHENG FENG-JU, AND HUJI MAIMATI. 1954. A preliminary study on *Scolytus rugulosus* Ratzeburg. Kunchong Zhishi 21(1): 27–31. (cn).
- ZHONG, H., AND TIMOTHY D. SCHOWALTER. 1959. Conifer bole utilization by wood-boring beetles in western Oregon. Canadian Journal of Forest Research 19: 943–947. (ec).
- *ZHOU, JIA-XI. 1963. The recent investigation of control of Scolytidae species on *Pinus armandii* by forest management measure [In Chinese]. A corpus of forestry scientific study references in Shaanxi Province. Peoples Press of Shaanxi 1963: 1–9. ().
- _____. 1953. Explanation on scolytids' damage frequently found in northwest of China [In Chinese]. Kunchong Zhishi 20(3): 134–136. (cn).
- ZIEGLER, K. 1955. Unerwünschte Beifänge in Weissen Borkenkäferfallen [Undesired trapping of non-target species in white bark beetle traps]. Allgemeine Forstzeitung 12: 256–257. (cn).
- *ZIVOJINOVIC, SVETISLAV. 1950b. La faune des insectes du Domaine Forestier de Majdanpek. Academie Serbe des Sciences, Monographies. T. CLX. Institut d'Ecologie et de Biogeographie 2: 1–262. ().
- ZOLUBUS, P. I. 1957. On the flight of spruce bark beetles to a source of chemical attractants [In Russian, English summary]. Khimoretseptsiya Nasekomykh 9: 75–83. (bv).
- *ZORRILLA, M. A. 19... Especies del Genero *Ips* (Col., Scol.); descortezadores de pinos en Cuba. [Source? (manuscript 57 p.)]. ().
- ZOTOVA, S. L. 1957. Trapping the bark beetles *Ips typographus* and *Ips duplicatus* with pheromone traps [In Russian]. Lesnoe Khozyaistvo 12: 57–58. (bv).
- *ZOUFAL, V. 1910. Ein Ausfluss auf Mostarsko-blato am 6 September 1909. Entomologische Blätter 6: 53–56. ().
- *ZUMR, VACLAV. 1950. Prostorové rozmišteni v rozsahu kmene, rojení a letová aktivita hlavních druhů kůrovce na smrku ztepilém (*Picea excelsa* L.). Československá akademie. 25 p. ().
- * _____. 1954. Prostorové rozmišteni kůrovce (Coleoptera, Scolytidae) na smrku ztepilém (*Picea excelsa* Link.) a jejich indifference podle lesních vegetačních stupňů [Spatial distribution of bark beetles in Norway spruce and their indifference]. Lesnictví [Prague] 30: 509–522. ().
- _____. 1955a. Atraktivita přípravku linoprax pro dřevokaze carkovanceho. *Trypodendron lineatum* (Ol.) (Coleoptera, Scolytidae) [Attractiveness of Linoprax pheromone for the lineate bark beetle *Trypodendron lineatum*]. Lesnictví [Prague] 31: 97–105. (bv).
- _____. 1955b. Společenstvo brouků (Coleoptera) v chodbach kůrovce (Scolytidae) na smrku ztepilém (*Picea excelsa* Link.) v jižních Čechách [Communities of beetles in the feeding sites of bark beetles on Norway spruce in South Bohemia]. Sborník Jihočeského muzea v Českých Budejovicích, Přírodní vědy 25: 45–48. (ec).
- * _____. 1955c. Biologie a ekologie lýkožravce smrkového (*Ips typographus*) a ochrana proti němu. Studie CSAV, Academia, Praha, vol. 17. 124 p. ().
- * _____. 1956a. Podkorní společenstvo brouků (Coleoptera) na smrku ztepilém *Picea excelsa* Link. [The community of bark-inhabiting beetles (Coleoptera) on Norway spruce] [In Czech, with English, Russian, German summaries]. Lesnictví [Prague] 32: 67–79. ().
- _____. 1956b. Reakce přirozených hmyzích nepřátel lýkožravce smrkového, *Ips typographus* (L.) (Coleoptera, Scolytidae), na feromon pheroprax [Reaction of insect natural enemies of the engraver beetle *Ips typographus* to the pheromone pheroprax] [In Czech, with English, Russian, German summaries]. Lesnictví [Prague] 32: 431–444. (ec).
- _____. 1957a. K ekologii méné známého lýkohuba haluzkového *Phthorophloeus spinulosus* Rey 1853 (Coleoptera, Scolytidae) na smrku ztepilém (*Picea excelsa* Link.) [On the ecology of the little known bark beetle *Phthorophloeus spinulosus* on Norway spruce] [In Russian, English summary]. Sborník Jihočeského Muzea v Českých Budejovicích, Přírodní Vědy 27: 17–25. (hb).
- _____. 1957b. Omezení gradací lýkožravce smrkového *Ips typographus* (L.) (Coleoptera, Scolytidae), pomocí feromonových lapací [Reduction of outbreaks of the engraver beetle *Ips typographus* using pheromone traps] [In Czech, English summary]. Lesnictví [Prague] 33: 49–64. (cn).
- _____. 1955. Účinnost agregací feromonu Chalcoprax proti lýkožravci lesklemu, *Pityogenes chalcographus* (L.), (Col., Scol.) [The effectiveness of the aggregation pheromone Chalcoprax in the control of spruce wood engraver *Pityogenes chalcographus* L. (Coleoptera, Scolytidae)]. Lesnictví [Prague] 34: 459–495. (bv cn).
- _____. 1959. Attractiveness of the terpene alpha pinene to the large pine shoot beetle *Blastophagus piniperda* L. (Coleoptera, Scolytidae). Journal of Applied Entomology 107: 141–144. (bv).
- ZUMB, VACLAV AND M. LANDA. 1955. Mit Pheromon-fallen Gegen Borkenkäfer in den Fichtenbeständen von Grätzen [Use of pheromone traps for bark beetle control in spruce stands in the Novohradské hory mountains]. Allgemeine Forstzeitung 96: 204–205. (bv cn).
- ZUMR, VACLAV, V. NEMEC, AND P. STARY. 1955. Seasonal changes in the nutrient content in bodies of *Ips typographus* L. (Col., Scolytidae). Journal of Applied Entomology 100: 464–465. (av).
- *ZUMR, VACLAV AND J. SIMANDL. 1955. Vyskyt dřevníka *Xyleborus cryptographus* Ratzeburg 1839 (Col., Scol.) na Blatenské [Occurrence of *Xyleborus cryptographus* near Blatná. Sbor. Jihočes. Muz. v Čes. Budejovicích, Přír. vědy 25: 32. ().

*ZVIEREZOMB-ZUBOVSKY AND ROSTOFF 1918. [*Hylesinus oleiperda* (=toranio), *Phloeotribus caucasicus*, *Pteleobius kraatzi*, *Scolytus rugulosus*]. Don Province Bureau for Control of Agricultural Pests. Report for 1917. 36 p. ().

ZWOLINSKI, J. B., AND C. J. GELDENHUYS. 1958. Insect damage to timbers from the indigenous forests: preliminary results. South African Forestry Journal 146: 7-11. (cn).

Addendum

Nine and one-half years have passed since this catalog was started. During those years significant world events have occurred, some of which have impacted the catalog. Several nations have ceased to exist, while other new ones have come into existence. The consequent changes in names or in new spellings of old names have been of concern. While we have attempted to incorporate many of those changes into the catalog, others have not been included. For example, the recent fragmentation of the former Soviet Union (USSR) into independent republics was not included. We have, however, included most, if not all, of the name changes among African nations. In 1957 it was announced by the news media that French Guiana was to become independent and would be named "Cayenne." On the strength of that announcement, we used the name Cayenne throughout the catalog. Unfortunately, the official change in name has not yet occurred.

The basic text of the catalog, including most notes, but excluding "References," was prepared by SLW. Notes prepared by DEB bear his initials. About half of the host records for Platypodidae and about one-third of those for Xyleborini came from DEB. While most of the literature citations listed under References, published from 1758 to 1959, came from SLW, his files also cover literature to 1972 or 1973 in reasonable detail and are included, as were other incidental references to 1991. Approximately two-thirds of the citations listed under References, published from 1960 to 1991, came from DEB except in *Dendroctonus*, *Ips*, Xyleborini, and Platypodidae, where the percentage of his contributions was substantially higher.

The funds needed for publication of Part 2 of the catalog, both volumes 2A and 2B, came from four sources. Approximately 57 percent came from funds conserved from grants of the U.S. sponsors (USDA and NSF), 34 percent from Canada, 2 percent from Brigham Young University, and 7 percent from S. L. Wood. The Canadian portion of publication support for Part 2 of this catalog was funded by Forestry Canada and the British Columbia Ministry of Forests under the Canada-British Columbia Partnership Agreement on Forest Resource Development: FRDA II (1991-1995). Dr. R. Asselin, Director, CLBRR, Agriculture Canada; Dr. L. Carlson, Deputy Director, Forestry Canada; and Mr. A. Shortreid, Forestry Technology Transfer Officer, Forestry Canada, arranged for and expedited the Canadian funds used for publication.

We regret the spelling error on p. 7 in the Introduction of the name of Derrick Bells.

Family Platypodidae Index

This index includes all Latin names used in this catalog for Platypodidae. Family-group names (family, subfamily, tribe) and names applied below the rank of subspecies (including aberrations, variations, nomen nudums) are given in regular type. Valid names of genera and species are in bold type. The names of synonyms of genera and species, subgeneric names, and subspecific names are given in *italics*. The names of fossil species are preceded by an asterisk (*). Fossil species are given at the beginning of the index and are also listed alphabetically with other names.

° <i>cylindricus</i> Burmeister, <i>Platypus</i>	1130	<i>aequalidentatus</i> Schedl, <i>Doliopygus</i>	1223
° <i>hurdi</i> Schedl, <i>Cenocephalus</i>	1109	<i>aequalis</i> Schedl, <i>Platypus</i>	1115
° <i>maravignae</i> Guérin-Meneville, <i>Platypus</i>	1157	<i>aequilaterus</i> Roberts, <i>Platypus</i>	1115
° <i>quadrilobus</i> Schedl, <i>Cenocephalus</i>	1109	<i>affinis</i> Guérin-Meneville, <i>Tesserocerus</i> (= <i>retusus</i>)	1100
° <i>rhinoceroide</i> (Schawaller), (<i>Mitosoma</i>), <i>Cenocephalus</i>	1109	<i>affinis</i> Strohmeier, <i>Cylindropalpus</i>	1217
° <i>succinicaptus</i> Schedl, <i>Cenocephalus</i>	1110	<i>africanus</i> Strohmeier, <i>Cylindropalpus</i> (= <i>auricomans</i>)	1218
<i>abbreviatus</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1213	<i>afzeliae</i> Browne, <i>Platypus</i>	1115
<i>abbreviatus</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1223	<i>agathis</i> Browne, <i>Platypus</i>	1115
<i>abditulus</i> (Wood), (<i>Platypus</i>), <i>Platyscapulus</i>	1214	<i>agilis</i> Dejean, (<i>Damicerus</i>), <i>Tesserocerus</i> (= <i>insignis</i>)	1099
<i>abditus</i> (Schedl), (<i>Platypus</i>), (<i>Platyscapulus</i>)	1214	<i>agnatus</i> Schedl, <i>Platypus</i>	1115
<i>abdominales</i> Schedl, <i>Doliopygus</i>	1223	<i>albipennis</i> Motschulsky, <i>Genyocerus</i>	1093
<i>abdominalis</i> (Schedl), (<i>Diacavus</i>), <i>Genyocerus</i>	1093	<i>albipennis</i> Strohmeier, (<i>Diapus</i>), <i>Genyocerus</i> (= <i>strohmeyeri</i>)	1096
<i>abdominalis</i> (Schedl), (<i>Platypus</i>), <i>Neotrachyostus</i>	1213	<i>algius</i> Schedl, <i>Platypus</i>	1115
<i>abdominalis</i> Schedl, <i>Crossotarsus</i> (= <i>majusculus</i>)	1202	<i>alienus</i> Schedl, <i>Platypus</i>	1115
<i>abietis</i> Wood, <i>Platypus</i>	1114	<i>alternans</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1223
<i>abnormis</i> Schedl, <i>Platypus</i>	1114	<i>alternans</i> Chapuis, <i>Platypus</i>	1115
<i>abruptifer</i> Wood, <i>Platypus</i>	1114	<i>alternantecostatus</i> (Schedl), (<i>Platyscapus</i>), <i>Platypus</i>	1115
<i>abruptulus</i> Browne, <i>Platypus</i>	1114	<i>alternantes</i> Schedl, <i>Carchesiopygus</i>	1209
<i>abruptus</i> Browne, <i>Platypus</i> (= <i>abruptifer</i>)	1114	<i>alternantes</i> Schedl, <i>Tesserocerus</i>	1098
<i>abruptus</i> Nunberg, <i>Periommatas</i>	1102	<i>ambiguus</i> Roberts, <i>Platypus</i>	1115
<i>abruptus</i> Sampson, <i>Platypus</i>	1114	<i>amblylaminatus</i> Roberts, <i>Diapus</i>	1088
<i>acanthurus</i> (Beeson), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1209	<i>ampliatum</i> Schedl, <i>Mitosoma</i>	1110
<i>accuratum</i> Schaufuss, <i>Mitosoma</i>	1110	<i>anaticeps</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1216
<i>acetabuliformis</i> Roberts, <i>Platypus</i>	1114	<i>anaticeps</i> Schedl, <i>Platypus</i>	1115
<i>aculeatus</i> Blandford, <i>Diapus</i>	1088	<i>andamanus</i> Beeson, <i>Crossotarsus</i>	1195
<i>aculeatus javanus</i> Murayama, <i>Diapus</i> (= <i>aculeatus</i>)	1088	<i>andrewesi</i> Strohmeier, <i>Platypus</i>	1115
<i>acutespinatus</i> Nunberg, <i>Doliopygus</i>	1223	<i>angolensis</i> Schedl, <i>Doliopygus</i>	1224
<i>acuticornifer</i> Wood, <i>Platypus</i>	1114	<i>anguis</i> Browne, <i>Platypus</i>	1115
<i>acuticornis</i> Schedl, <i>Platypus</i>	1114	<i>angustatulus</i> Wood, <i>Platypus</i>	1116
<i>acuticornis</i> Schedl, <i>Platypus</i> (= <i>acuticornifer</i>)	1114	<i>angustatus</i> Chapuis, <i>Platypus</i>	1116
<i>acutidens</i> (Blandford), (<i>Platypus</i>), <i>Neotrachyostus</i>	1213	<i>angustatus</i> Strohmeier, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>orientalis</i>)	1162
<i>acutidentatus</i> Murayama, <i>Platypus</i>	1114	<i>angusticeps</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1216
<i>adnexus</i> Schedl, <i>Platypus</i>	1114	<i>angusticollis</i> Schedl, <i>Platypus</i>	1116
<i>aduncus</i> Chapuis, <i>Platypus</i>	1114	<i>angustidontis</i> Roberts, <i>Diapus</i>	1055
<i>aduncus</i> Schedl, <i>Doliopygus</i>	1223	<i>angustiformis</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1216
<i>adursus</i> Beeson, <i>Crossotarsus</i>	1195	<i>angustiformis</i> Schedl, <i>Periommatas</i> (= <i>longicollis</i>)	1105
<i>adustipennis</i> Motschulsky, <i>Genyocerus</i>	1203	<i>angustior</i> Schedl, <i>Periommatas</i>	1102
<i>advena</i> Schedl, <i>Platypus</i>	1115	<i>angustior</i> Schedl, <i>Platypus</i>	1116
<i>aequalicinctus</i> Schedl, <i>Platypus</i>	1115	<i>angustioris</i> Schedl, <i>Platypus</i>	1116
		<i>angustipennis</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1216
		<i>angustulus</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1216

<i>angustus</i> Schedl, <i>Periomnatus</i> (= <i>longicollis</i>)	1104	<i>astutus</i> Schedl, <i>Platypus</i>	1118
<i>annexus</i> Wood, <i>Platypus</i>	1116	<i>ater</i> (Strohmeyer), (<i>Platypus</i>), <i>Trachyostus</i>	1211
<i>annosus</i> Wood, <i>Protolylastes</i>	1084	<i>ater</i> Browne, <i>Doliopygus</i>	1224
<i>annularis</i> Schedl, (<i>Platypus</i>), <i>Baiocis</i>	1216	<i>ater</i> Schedl, <i>Platypus</i> (= <i>impressus</i>)	1147
<i>anomalus</i> Schedl, <i>Platypus</i>	1116	<i>aterrimus</i> (Schaufuss), (<i>Platypus</i>), <i>Trachyostus</i>	1211
<i>anoplus</i> Schedl, <i>Platypus</i>	1116	<i>aterrimus minor</i> Roberts, <i>Trachyostus</i> (= <i>aterrimus</i>)	1211
<i>anthocephali</i> Beeson, <i>Platypus</i> (= <i>anthocephali</i>)	1116	<i>atkinsoni</i> Beeson, (<i>Diacatus</i>), <i>Genyocerus</i> (=sp.?)	1093
<i>anthocephali</i> Schedl, <i>Platypus</i>	1116	<i>atrans</i> Roberts, <i>Platypus</i>	1118
<i>antongilis</i> Schedl, <i>Trachyostus</i>	1211	<i>attentus</i> Schedl, <i>Platypus</i>	1118
<i>aolai</i> Browne, <i>Platypus</i>	1116	<i>anbei</i> Chapuis, <i>Tesserocherus</i> (= <i>dewalquei</i>)	1098
<i>apertulus</i> Schedl, <i>Platypus</i>	1116	<i>aureipilus</i> Roberts, <i>Crossotarsus</i>	1195
<i>apertus</i> Chapuis, <i>Platypus</i> (= <i>hians</i>)	1145	<i>auricomans</i> (Schaufuss), (<i>Platypus</i>), <i>Cylindropalpus</i>	1218
<i>apertus</i> Schedl, (<i>Diapus</i>), <i>Platypus</i>	1116	<i>auricularis</i> Chapuis, <i>Platypus</i>	1118
<i>apicalis</i> (White), (<i>Crossotarsus</i>), <i>Platypus</i>	1116	<i>auritus</i> Chapuis, <i>Platypus</i>	1118
<i>apicaloides</i> Schedl, <i>Platypus</i>	1117	<i>australis</i> Chapuis, <i>Platypus</i>	1118
<i>apicatus</i> Schedl, <i>Platypus</i>	1117	<i>Austroplatypus</i> Browne (= <i>Platypus</i>)	1113
<i>aplanatus</i> Schedl, <i>Platypus</i> (= <i>applanatulus</i>)	1117	<i>Baiocis</i> Browne	1216
<i>appendicinus</i> Schedl, <i>Mesoplatypus</i>	1221	<i>bajulus</i> Schedl, <i>Platypus</i>	1119
<i>appendiculatus</i> Dejean, <i>Tesserocherus</i> (= <i>dewalquei</i>)	1098	<i>bajulus sumatranus</i> Schedl, <i>Platypus</i> (= <i>bajulus</i>)	1119
<i>applanatulus</i> Wood, <i>Platypus</i>	1117	<i>balanocarpus</i> Beeson, <i>Platypus</i> (= <i>balanocarpus</i> Schedl)	1119
<i>applanatus</i> Wood, <i>Platypus</i>	1117	<i>balanocarpus</i> Schedl, <i>Platypus</i>	1119
<i>araucariae</i> Schedl, <i>Platypus</i>	1117	<i>banghaasi</i> (Schaufuss), (<i>Crossotarsus</i>), <i>Triozastus</i>	1220
<i>arcuatus</i> Schedl, <i>Platypus</i>	1117	<i>banghaasi banghaasi</i> (Schaufuss), <i>Triozastus</i> (= <i>banghaasi</i>)	1220
<i>arduus</i> (Schedl), (<i>Platyscapus</i>), <i>Platypus</i>	1117	<i>barbatulus</i> Schedl, <i>Platypus</i>	1119
<i>areolatus</i> Chapuis, <i>Platypus</i>	1117	<i>barbatus</i> Chapuis, <i>Crossotarsus</i>	1196
<i>arisannensis</i> Murayama, <i>Platypus</i>	1117	<i>barbosai</i> Schedl, <i>Platypus</i>	1119
<i>armaticeps</i> Beeson, <i>Platypus</i> (= <i>armaticeps</i> Schedl)	1117	<i>basilewskyi</i> Nunberg, <i>Periomnatus</i>	1102
<i>armaticeps</i> Schedl, <i>Platypus</i>	1117	<i>batesi</i> Chapuis, <i>Platypus</i>	1119
<i>armatus</i> Chapuis, <i>Platypus</i>	1117	<i>beaveri</i> Browne, <i>Platypus</i>	1119
<i>armipemis</i> Lea, <i>Crossotarsus</i>	1195	<i>beesoni</i> Sampson, (<i>Mesoplatypus</i>), <i>Platypus</i> (= <i>biflexuosus</i>)	1120
<i>arrogans</i> Schedl, <i>Platypus</i>	1118	<i>beilschmiediae</i> Schedl, <i>Platypus</i>	1119
<i>arrowi</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1224	<i>bellus</i> Schedl, <i>Platypus</i>	1119
<i>artecarinatus</i> Schedl, <i>Platypus</i>	1118	<i>belti</i> Sharp, <i>Tesserocherus</i>	1098
<i>artecatus</i> Schedl, <i>Platypus</i> (= <i>pseudocupulatus</i>)	1172	<i>benevolus</i> Beeson, <i>Crossotarsus</i>	1196
<i>artecircularis</i> Schedl, <i>Periomnatus</i>	1102	<i>berliniae</i> Roberts, <i>Doliopygus</i>	1224
<i>artecostatus</i> (Schedl), (<i>Platyscapus</i>), <i>Platypus</i>	1118	<i>bicallosus</i> Schedl, <i>Periomnatus</i>	1102
<i>artecurtus</i> Schedl, <i>Platypus</i> (= <i>curtus</i>)	1130	<i>bicarinulatus</i> Schedl, <i>Platypus</i>	1119
<i>artesolidus</i> Schedl, <i>Platypus</i>	1118	<i>bicaudatus</i> Schedl, <i>Platypus</i>	1119
<i>artespinitus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1224	<i>bicaudatus</i> Browne, <i>Platypus</i>	1120
<i>artestrigatus</i> Nunberg, <i>Periomnatus</i>	1102	<i>biceps</i> Beeson, <i>Crossotarsus</i> (= <i>horni</i>)	1201
<i>artetrunatus</i> Schedl, <i>Platypus</i>	1118	<i>bicolor</i> Montrouzier, <i>Platypus</i>	1120
<i>aruensis</i> Beeson, <i>Crossotarsus</i>	1195	<i>biconcavus</i> Schedl, <i>Crossotarsus</i>	1196
<i>Asetus</i> Nunberg, (= <i>Periomnatus</i>)	1102	<i>biconicus</i> Schedl, <i>Platytarsulus</i>	1097
<i>asperatus</i> Schedl, <i>Platypus</i>	1118	<i>bicorniger</i> Schedl, <i>Platypus</i>	1120
<i>aspicis</i> Roberts & Morimoto, <i>Platypus</i>	1118	<i>bicomis</i> Beeson, <i>Platypus</i> (= <i>insulindicus</i>)	1149
<i>assamelae</i> Browne, <i>Platypus</i>	1118	<i>bicornis</i> Schedl, <i>Platypus</i> (= <i>insulindicus</i>)	1149
<i>assamensis</i> (Beeson), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1210	<i>bicornutus</i> Nunberg, <i>Platypus</i> (= <i>insulindicus</i>)	1149
<i>assamensis</i> Beeson, <i>Genyocerus</i> (= <i>diaphanus</i>)	1094	<i>bideus</i> Schedl, <i>Platypus</i>	1120
<i>assamensis</i> Browne, (<i>Diacatus</i>), <i>Genyocerus</i> (= <i>diaphanus</i>)	1094	<i>bidentatus</i> Dejean, <i>Platypus</i> (= <i>flavicornis</i>)	1139
<i>assimilis</i> Schedl, <i>Platypus</i>	1118		
<i>associatus</i> Schedl, <i>Platypus</i>	1118		

<i>bidentatus</i> (Strohmeyer), (Crossotarsus), Doliopygus	1224	<i>brasilensis</i> Evers, Tesserocerus	1095
<i>bidentatus posticalis</i> Schedl, Doliopygus (= <i>bidentatus</i>)	1224	<i>brasilensis</i> Nunberg, Platypus	1122
<i>bidiscoplanus</i> Roberts, Doliopygus	1224	<i>brevestrigatus</i> Nunberg, Periommatatus (= <i>longicollis</i>)	1105
<i>bifidus</i> Schedl, Platypus	1120	<i>brevicaudatus</i> Nunberg, Platypus	1122
<i>biflexuosus</i> Schedl, Platypus	1120	<i>brevicornis</i> Wood, Platypus	1122
<i>biformis</i> Chapuis, Platypus	1120	<i>brevidens</i> Browne, Crossotarsus	1196
<i>bifrons</i> Chapuis, Platypus (= <i>cordiger</i>)	1128	<i>brevidens</i> Browne, Triozastus	1220
<i>bifrons</i> Schedl, Periommatatus	1102	<i>brevis</i> (Strohmeyer), (Crossotarsus), Doliopygus	1225
<i>bifurcus</i> (Schedl), (Crossotarsus), Platypus	1120	<i>brevis</i> Browne, Platypus	1122
<i>bifurcus</i> Schedl, Triozastus (= <i>banghaasi</i>)	1220	<i>brevis interstitialis</i> Schedl, Doliopygus (= <i>rapax</i>)	1237
<i>bifurcus mutilus</i> Schedl, (Crossotarsus), Platypus (= <i>shoreanus</i>)	1151	<i>brevispinatus</i> Nunberg, Periommatatus (= <i>bispinus</i>)	1103
<i>bigranulatus</i> Schedl, Platypus	1121	<i>breviusculus</i> Schedl, Doliopygus (= <i>rapax</i>)	1237
<i>bilamatus</i> Guerin-Meneville, Tesserocerus (= <i>insignis</i>)	1100	<i>brochus</i> Beeson, Diapus (= <i>spatulifer</i>)	1092
<i>bilamatus</i> Schedl, Platypus	1121	<i>brownei</i> Beaver, Triozastus	1220
<i>bijunctispinae</i> Roberts, Doliopygus	1225	<i>brownei</i> Schedl, Crossotarsus	1196
<i>bilobatus</i> (Schedl), (Crossotarsus), Doliopygus	1225	<i>brownei</i> Schedl, Diapus	1059
<i>bilobatus</i> Nunberg, Mesoplatypus	1221	<i>brownei</i> Schedl, Platypus	1122
<i>bilobatus</i> Strohmeyer, Platypus	1121	<i>brunneus</i> Chapuis, Platypus	1122
<i>bilobipennis</i> Schedl, Platypus	1121	<i>bugiriensis</i> Nunberg, Doliopygus (= <i>chapuisi</i>)	1226
<i>bilobus</i> (Schedl), (Platypus), Tesserocerus	1095	<i>bulbifer</i> Schedl, Doliopygus	1225
<i>bilunatus</i> Schedl, Diapus	1058	<i>bulbiventris</i> Schedl, Doliopygus	1226
<i>bimaculata</i> Duftschmidt, (Cylindra), Platypus (= <i>cyindra</i>)	1132	<i>bulbosus</i> Roberts, Doliopygus (= <i>bulbifer</i>)	1225
<i>binodulus</i> Chapuis, Platypus	1121	<i>burmannus</i> (Sampson), (Crossotarsus), Platypus	1122
<i>bipapillatus</i> Roberts, Doliopygus	1225	<i>cachani</i> Schedl, Mesoplatypus (= <i>quinquicinctus</i>)	1222
<i>biporus</i> (Schedl), (Diacatus), Genyocerus	1093	<i>caelestis</i> Roberts & Morimoto, Platypus	1122
<i>biporus</i> Beeson, Genyocerus (= <i>biporus</i>)	1093	<i>caesalpiniae</i> Roberts, Doliopygus	1226
<i>biporus</i> Blandford, Platypus	1121	<i>calabarius</i> Roberts, Mesoplatypus	1221
<i>bipyramidus</i> Roberts & Morimoto, Platypus	1121	<i>calamus</i> Blandford, Platypus	1122
<i>bisignatus</i> Schedl, Platypus	1121	<i>calanus fukiensis</i> Schedl, Platypus (= <i>calamus</i>)	1123
<i>bispinatus</i> Browne, Platypus	1121	<i>caledoniae</i> Schedl, Platypus	1123
<i>bispinus</i> Schedl, Diapus	1058	<i>caliculus</i> Chapuis, Platypus (= <i>luniger</i>)	1156
<i>bispinus</i> Strohmeyer, Periommatatus	1103	<i>caligatus</i> Schedl, Platypus	1123
<i>bistriatus</i> Browne, Platypus (= <i>deflectus</i>)	1134	<i>caliginosus</i> Roberts, Crossotarsus	1196
<i>bitalei</i> Schedl, Doliopygus	1225	<i>caliginosus</i> Roberts, Cylindropalpus	1215
<i>bituberculatus</i> Nunberg, Mesoplatypus (= <i>ventricornis</i>)	1223	<i>caliginosus</i> Roberts, Doliopygus (= <i>artespinitus</i>)	1224
<i>bituberculatus</i> Schedl, (Crossotarsus), Doliopygus (= <i>artespinitus</i>)	1224	<i>caliginosus</i> Roberts, Triozastus	1220
<i>bituberculatus</i> Schedl, Platypus (= <i>bigranulatus</i>)	1121	<i>caliginosus humilis</i> Browne, Triozastus (= <i>caliginosus</i>)	1220
<i>bituberculifer</i> Wood, Platypus (= <i>bigranulatus</i>)	1121	<i>caliginosus</i> Schedl, Platypus	1123
<i>biuncus</i> Blandford, Platypus	1121	<i>calix</i> Schedl, (Platypus), Platyscapulus (= <i>costellatus</i>)	1215
<i>blanchardi</i> Chapuis, Platypus (= <i>quadridentatus</i>)	1174	<i>cambodianus</i> Schedl, Platypus (= <i>piniperda</i>)	1170
<i>blumi</i> Schedl, Platypus (= <i>cupulatus</i>)	1129	<i>camerunus</i> (Schedl), (Platyscapus), Cylindropalpus	1215
<i>boettcheri</i> Schedl, Platypus	1122	<i>camerunus</i> Strohmeyer, Periommatatus (= <i>longicollis</i>)	1104
<i>bohemani</i> (Chapuis), (Crossotarsus), Doliopygus	1225	<i>camerunus angustiformis</i> Schedl, Periommatatus (= <i>longicollis</i>)	1105
<i>bonvouloiri</i> Chapuis, Crossotarsus	1196	<i>camerunus brevestrigatus</i> Nunberg, Periommatatus (= <i>longicollis</i>)	1105
<i>borneensis</i> Browne, Diapus	1059	<i>camerunus crebrestrigatus</i> Nunberg, Periommatatus (= <i>longicollis</i>)	1105
<i>borneensis</i> Browne, Platypus	1122	<i>canaliculatus</i> Schedl, Platypus	1123
<i>borneensis</i> Schedl, Platytarsulus	1097	<i>candezei</i> Chapuis, Platypus	1123
<i>bothrocephalus</i> (Strohmeyer), (Crossotarsus), Doliopygus	1225	<i>capillatus</i> Sampson, Diapus (= <i>molossus</i>)	1090
		<i>capitalis</i> Beeson, Diapus (= <i>spatulifer</i>)	1092

<i>capitilanus</i> Roberts & Morimoto, <i>Platypus</i> . . .	1123	<i>clunalis</i> Wood, <i>Platypus</i> . . .	1125
<i>capito</i> Browne, <i>Platypus</i> . . .	1123	<i>cluniculus</i> Wood, <i>Platypus</i> . . .	1125
<i>carupae</i> Schedl, <i>Doliopygus</i> . . .	1226	<i>clunis</i> Wood, <i>Platypus</i> . . .	1125
<i>caranensis</i> Schedl, <i>Platypus</i> . . .	1123	<i>coadunatus</i> Schedl, <i>Platypus</i> . . .	1125
<i>carbonescens</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1123	<i>Coccephalonus</i> Schedl (= <i>Mitosoma</i>) . . .	1110
<i>Carchesiopygus</i> Schedl . . .	1209	<i>coelocephalus</i> (Schanfuss), <i>Doliopygus</i> . . .	1227
<i>carduus</i> Schedl, <i>Platypus</i> . . .	1123	<i>cognatus</i> Roberts, <i>Platypus</i> . . .	1125
<i>carinatus</i> Schedl, <i>Trachyostus</i> . . .	1211	<i>coleopteratus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1125
<i>carinifer</i> Schedl, <i>Platypus</i> . . .	1123	<i>collaris</i> Schedl, <i>Platypus</i> . . .	1125
<i>carinifrons</i> Schedl, <i>Platypus</i> . . .	1123	<i>collatatus</i> Schedl, <i>Platypus</i> . . .	1125
<i>carinulatus</i> (Chapuis), (<i>Platypus</i>), <i>Platyscapulus</i> . . .	1214	<i>collaticius</i> Schedl, <i>Periommatus</i> . . .	1103
<i>caryophyllatus</i> Schedl, <i>Platypus</i> . . .	1123	<i>collinus</i> Browne, <i>Platypus</i> . . .	1125
<i>castaneus</i> Broun, <i>Platypus</i> (= <i>apicalis</i>) . . .	1117	<i>comalis</i> Roberts, <i>Platypus</i> (= <i>acetabuliformis</i>) . . .	1114
<i>castaneus</i> Strohmeier, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>orientalis</i>) . . .	1162	<i>comatus</i> Chapuis, <i>Crossotarsus</i> . . .	1197
<i>castigatus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1124	<i>comoreanus</i> Schedl, <i>Platypus</i> . . .	1125
<i>caudatus</i> Motschulsky, <i>Platypus</i> (= <i>solidus</i>) . . .	1183	<i>compactus</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i> . . .	1093
<i>caudatus</i> Schedl, <i>Platypus</i> (= <i>brasilensis</i>) . . .	1122	<i>compactus</i> dubiosus Schedl, (<i>Diacarus</i>), <i>Genyocerus</i> (= <i>compactus</i>) . . .	1093
<i>caviceps</i> Broun, <i>Platypus</i> . . .	1124	<i>complanatus</i> Schedl, <i>Platypus</i> . . .	1125
<i>cavifrons</i> Blandford, <i>Crossotarsus</i> (= <i>indomitus</i>) . . .	1201	<i>complanus</i> Schedl, <i>Platypus</i> . . .	1126
<i>cavus</i> Strohmeier, <i>Platypus</i> . . .	1124	<i>compositus</i> Say, <i>Platypus</i> . . .	1126
<i>celsus</i> Roberts, <i>Platypus</i> . . .	1124	<i>compressus</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>) . . .	1166
<i>Cenocephalariae</i> Strohmeier (= <i>Tesserocerini</i>) . . .	1085	<i>concarifrons</i> Schedl, <i>Crossotarsus</i> . . .	1197
<i>Cenocephalini</i> Strohmeier (= <i>Tesserocerini</i>) . . .	1085	<i>concarifrons</i> Schedl, <i>Platypus</i> . . .	1127
<i>Cenocephalus</i> Chapuis . . .	1109	<i>concarus</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i> . . .	1213
<i>Chaetastus</i> Nunberg . . .	1107	<i>concentriporus</i> Roberts, <i>Platypus</i> . . .	1127
<i>chalcographus</i> Schedl, <i>Crossotarsus</i> . . .	1196	<i>conciliatus</i> Schedl, <i>Platypus</i> . . .	1127
<i>chapuisi</i> (Duvivier), (<i>Crossotarsus</i>), <i>Doliopygus</i> . . .	1226	<i>concinulus</i> Blandford, <i>Platypus</i> . . .	1127
<i>chapuisi</i> Blandford, <i>Crossotarsus</i> (= <i>concinus</i>) . . .	1197	<i>concinus</i> Blandford, <i>Crossotarsus</i> . . .	1197
<i>chapuisi</i> Schedl, <i>Tesserocerus</i> . . .	1095	<i>confusus</i> Roberts, <i>Doliopygus</i> (= <i>dubius</i>) ¹ . . .	1229
<i>chapuisi</i> Strohmeier, <i>Mitosoma</i> . . .	1110	<i>congoanus</i> Duvivier, <i>Platypus</i> (= <i>parallelus</i>) . . .	1168
<i>chapuisi nitidiventris</i> Schedl, <i>Doliopygus</i> (= <i>chapuisi</i>) . . .	1226	<i>congoanus</i> Nunberg, <i>Periommatus</i> . . .	1103
<i>Chapuisia</i> Duges (= <i>Schedlarini</i>) . . .	1087	<i>congonus</i> Browne, <i>Doliopygus</i> . . .	1227
<i>Chapuisidae</i> Blandford (= <i>Schedlarini</i>) . . .	1084, 1087	<i>coniferae</i> Stebbing, <i>Crossotarsus</i> . . .	1197
<i>Chapuisidae</i> Blandford (= <i>Schedlarini</i>) . . .	1087	<i>conjunctus</i> Schedl, <i>Doliopygus</i> . . .	1227
<i>Chapuisinae</i> Blandford (= <i>Schedlarini</i>) . . .	1087	<i>conjunctus</i> Schedl, <i>Platypus</i> . . .	1127
<i>cheesmani</i> Schedl, <i>Crossotarsus</i> . . .	1196	<i>conjunctus</i> Schedl, <i>Platypus</i> (= <i>cupulatus</i>) . . .	1129
<i>cheerolati</i> Chapuis, <i>Platypus</i> . . .	1124	<i>connexus</i> Wood, <i>Platypus</i> . . .	1127
<i>chimbui</i> Schedl, <i>Platypus</i> . . .	1124	<i>conradti</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i> . . .	1227
<i>chiriquensis</i> Wood, <i>Platypus</i> . . .	1124	<i>consequens</i> Schedl, <i>Platypus</i> . . .	1127
<i>cicatricosus</i> Roberts, <i>Platypus</i> . . .	1125	<i>contaminatus</i> (Blandford), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1127
<i>cincinnatus</i> Chapuis, <i>Crossotarsus</i> . . .	1196	<i>contextus</i> Schedl, <i>Platypus</i> . . .	1125
<i>cinnamomi</i> Beeson, <i>Crossotarsus</i> . . .	1197	<i>contractus</i> Chapuis, <i>Platypus</i> . . .	1128
<i>circulariceps</i> Schedl, <i>Periommatus</i> (= <i>circularis</i>) . . .	1103	<i>contractus</i> Chapuis, <i>Tesserocerus</i> . . .	1095
<i>circulariceps</i> Schedl, <i>Platypus</i> . . .	1125	<i>conrexicauda</i> Schedl, <i>Platypus</i> . . .	1125
<i>circularis</i> Chapuis, <i>Platypus</i> . . .	1125	<i>convexus</i> Roberts, <i>Doliopygus</i> . . .	1227
<i>circularis</i> Schedl, <i>Periommatus</i> . . .	1103	<i>convexus</i> Schedl, <i>Platypus</i> . . .	1125
<i>circularis circulariceps</i> Schedl, <i>Periommatus</i> (= <i>circularis</i>) . . .	1103	<i>Coptonotidae</i> Chapuis (= <i>Coptonotinae</i>) . . .	1084
<i>circulicauda</i> Browne, <i>Treptoplatypus</i> . . .	1194	<i>Coptonotinae</i> Chapuis . . .	1084
<i>circumdentatus</i> Strohmeier, <i>Platypus</i> . . .	1125	<i>Coptonotini</i> Chapuis . . .	1084
<i>citri</i> Schedl, <i>Doliopygus</i> . . .	1226	<i>Coptonotus</i> Chapuis . . .	1084
<i>clarkei</i> Browne, <i>Platypus</i> . . .	1125	<i>cordatus</i> Motschulsky, <i>Platypus</i> (= <i>solidus</i>) . . .	1183
<i>clarus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i> . . .	1226	<i>cordiger</i> Chapuis, <i>Platypus</i> . . .	1128
<i>cliens</i> (Schedl), (<i>Platypus</i>), <i>Crossotarsus</i> . . .	1197	<i>cordiger bifrons</i> Chapuis, <i>Platypus</i> (= <i>cordiger</i>) . . .	1128

<i>cordiger gabensis</i> Schedl, <i>Platypus</i> (=gabensis) . . .	1142	<i>dasycauda</i> Browne, <i>Platypus</i> . . .	1133
<i>corniventris</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1128	<i>decellei</i> Schedl, <i>Trachyostus</i> . . .	1211
<i>cornutus</i> Schedl, <i>Platypus</i> . . .	1128	<i>decemspinatus</i> (Schedl), (<i>Diacavus</i>), <i>Genyocerus</i> . . .	1093
<i>coronatus</i> Schedl, <i>Platypus</i> . . .	1128	<i>decens</i> Sampson, <i>Platypus</i> . . .	1133
<i>corporaali</i> Browne, <i>Spathidicerus</i> . . .	1101	<i>deceptor</i> Wood, <i>Platypus</i> . . .	1133
<i>corrugatus</i> Roberts, <i>Crossotarsus</i> . . .	1197	<i>deceptus</i> Beeson, <i>Crossotarsus</i> (=ursus) . . .	1208
<i>costaricensis</i> Schedl, <i>Platypus</i> . . .	1128	<i>declivornatus</i> (Schedl), (<i>Platypus</i>), <i>Crossotarsus</i> . . .	1198
<i>Costaroplatus</i> Numberg, (<i>Platyscapulus</i>) . . .	1214	<i>declivis</i> Sampson, <i>Platypus</i> . . .	1133
<i>costatus</i> Nunberg, <i>Periommatas</i> . . .	1103	<i>declivis</i> Schedl, <i>Platytarsulus</i> . . .	1097
<i>costatus</i> Schedl, <i>Doliopygus</i> . . .	1227	<i>declivitatis</i> Schedl, <i>Platypus</i> . . .	1133
<i>costellatus</i> (Schedl), (<i>Platypus</i>), <i>Platyscapulus</i> . . .	1215	<i>decorus</i> Schedl, <i>Platypus</i> . . .	1133
<i>costipennis</i> Schedl, <i>Platypus</i> . . .	1128	<i>deductus</i> Schedl, <i>Platypus</i> . . .	1134
<i>coxalis</i> Schedl, <i>Crossotarsus</i> . . .	1197	<i>deflectus</i> Schedl, <i>Platypus</i> . . .	1134
<i>crassicornis</i> Schedl, <i>Platypus</i> . . .	1128	<i>defranatus</i> Schedl, <i>Periommatas</i> . . .	1103
<i>crassiusculus</i> Schedl, <i>Platypus</i> . . .	1129	<i>dejeani</i> Chapuis, <i>Platypus</i> (=parallelus) . . .	1166
<i>crassus</i> Strohmeier, <i>Platypus</i> . . .	1129	<i>dejeani</i> Chapuis, <i>Tesserocherus</i> (=deicalquei) . . .	1098
<i>crebrestrigatus</i> Numberg, <i>Periommatas</i> (=longicollis) . . .	1105	<i>delicatus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> . . .	1134
<i>crenatus</i> Chapuis, <i>Platypus</i> (=australis) . . .	1119	Dendroplatypus Browne . . .	1240
<i>crenulatum</i> Chapuis, <i>Mitosoma</i> . . .	1110	<i>dentatus</i> Dalman, <i>Platypus</i> . . .	1134
<i>cribricollis</i> Blandford, <i>Platypus</i> . . .	1129	<i>denticollis</i> Browne, <i>Platypus</i> . . .	1134
<i>erinitus</i> (Chapuis), (<i>Crossotarsus</i>), <i>Doliopygus</i> . . .	1227	<i>denticornis</i> Dejean, (<i>Danicerus</i>), <i>Tesserocherus</i> (=insignis) . . .	1099
<i>Crossotarsariae</i> Strohmeier (=Crossotarsini) . . .	1113	<i>denticulatus</i> Browne, <i>Platypus</i> . . .	1134
<i>Crossotarsinae</i> Strohmeier (=Crossotarsini) . . .	1113	<i>denticulus</i> Browne, <i>Crossotarsus</i> . . .	1198
<i>Crossotarsini</i> Strohmeier (=Crossotarsini) . . .	1113	<i>dentipennis</i> Schedl, <i>Carchesiopygus</i> . . .	1210
Crossotarsinulus Schedl . . .	1210	<i>denticentris</i> Browne, <i>Doliopygus</i> . . .	1228
Crossotarsus Chapuis . . .	1195	<i>denturus</i> Browne, <i>Crossotarsus</i> . . .	1198
<i>cultellus</i> (Schedl), (<i>Platyscapus</i>), <i>Cylindropalpus</i> . . .	1218	<i>deperditus</i> Schedl, <i>Platypus</i> . . .	1134
<i>cupulatus</i> Schedl, <i>Platypus</i> . . .	1129	<i>deplanatus</i> Wood, <i>Platypus</i> . . .	1134
<i>cupulatus conjunctus</i> Schedl, <i>Platypus</i> (=cupulatus) . . .	1129	<i>depressus</i> Schedl, <i>Platypus</i> . . .	1134
<i>cupulatus</i> Chapuis, <i>Platypus</i> . . .	1129	<i>derosus</i> Schedl, <i>Crossotarsus</i> . . .	1198
<i>cupulatus bhuni</i> Schedl, <i>Platypus</i> (=cupulatus) . . .	1129	<i>deruptus</i> Schedl, <i>Doliopygus</i> . . .	1228
<i>cupulatus picus</i> Schedl, <i>Platypus</i> (=cupulatus) . . .	1129	<i>desectus</i> Beeson, <i>Crossotarsus</i> (=broicuei) . . .	1196
<i>cupidifer</i> Wichmann, <i>Platypus</i> (=cavus) . . .	1124	<i>despectus</i> Schedl, <i>Tesserocherus</i> . . .	1098
<i>curtatus</i> Sampson, <i>Platypus</i> . . .	1130	<i>desultor</i> Schedl, <i>Platypus</i> . . .	1134
<i>curtus</i> Chapuis, <i>Platypus</i> . . .	1130	<i>detectus</i> Schedl, <i>Platypus</i> . . .	1134
<i>curtus artemcurtus</i> Schedl, <i>Platypus</i> (=curtus) . . .	1130	<i>devius</i> Schedl, <i>Platypus</i> . . .	1134
<i>curvidens</i> Browne, <i>Mesoplatypus</i> . . .	1222	<i>deicalquei</i> Chapuis, <i>Tesserocherus</i> . . .	1098
<i>curvidens</i> Schedl, <i>Diapus</i> . . .	1089	<i>deyrollei</i> Chapuis, <i>Platypus</i> . . .	1134
<i>curvidens</i> Schedl, <i>Platypus</i> . . .	1130	<i>Diacavus</i> Schedl (=Genyocerus) . . .	1093
<i>cuspidatus</i> Schedl, <i>Platypus</i> . . .	1130	<i>dialiumi</i> Roberts, <i>Doliopygus</i> (=bilobatus) . . .	1225
<i>cyclopus</i> Chapuis, <i>Coptonotus</i> . . .	1084	<i>diaphanus</i> (Schedl), (<i>Diacavus</i>), <i>Genyocerus</i> . . .	1094
<i>Cylindra</i> Illiger (=Platypus) . . .	1113	<i>Diapodariae</i> Strohmeier (=Diapodini) . . .	1088
* <i>cylindricus</i> Burmeister, <i>Platypus</i> . . .	1130	<i>Diapodinae</i> Strohmeier (=Diapodini) . . .	1088
<i>cylindriformis</i> Reitter, <i>Platypus</i> (=cylindrus) . . .	1132	<i>Diapodini</i> Strohmeier . . .	1088
Cylindropalpus Strohmeier . . .	1217	Diapus Chapuis . . .	1088
<i>cylindrus</i> (Fabricius), (<i>Bostrichus</i>), <i>Platypus</i> . . .	1130	<i>dibrachiatus</i> Roberts, <i>Platypus</i> . . .	1134
<i>cymbiformis</i> Roberts, <i>Platypus</i> . . .	1133	<i>diductus</i> Chapuis, <i>Platypus</i> . . .	1135
<i>daedalus</i> Roberts, <i>Platypus</i> . . .	1133	<i>diegensis</i> Schanfuß, <i>Platypus</i> (=compositus) . . .	1127
<i>Danicerus</i> Spinola (=Tesserocherus) . . .	1097	<i>difficilis</i> Chapuis, <i>Platypus</i> (=parallelus) . . .	1167
<i>dammanus</i> Beeson, <i>Crossotarsus</i> . . .	1197	<i>diffidens</i> Schedl, <i>Platypus</i> . . .	1135
<i>darjeelingensis</i> Beeson, <i>Platypus</i> (=darjeelingensis) . . .	1133	<i>diffinis</i> Schedl, <i>Platypus</i> . . .	1135
<i>darjeelingensis</i> Schedl, <i>Platypus</i> . . .	1133	<i>digestus</i> Schedl, <i>Platypus</i> . . .	1135
<i>darlingtoni</i> Reichardt, <i>Platypus</i> . . .	1133	<i>digitalis</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i> . . .	1213
		<i>dignatus</i> Schedl, <i>Platypus</i> . . .	1135
		<i>dignus</i> Schedl, <i>Platypus</i> . . .	1135

<i>dimidiatus</i> Chapuis, <i>Platypus</i>	1135	<i>elegans</i> Schedl, <i>Mesoplatypus</i>	1222
<i>dipteroearpi</i> Beeson, <i>Genyocerus</i> (= <i>dipteroearpi</i>)	1094	<i>elongatus</i> Chapuis, <i>Platypus</i>	1137
<i>dipteroearpi</i> Browne, <i>Genyocerus</i>	1094	<i>elongatus</i> Lea, <i>Notoplatypus</i>	1097
<i>dipteroearpi</i> Browne, <i>Platypus</i>	1135	<i>elongatus</i> Nunberg, <i>Periommatatus</i>	1103
<i>discicollis</i> Chapuis, <i>Platypus</i>	1135	<i>elongatus</i> Schedl, (<i>Stenoplatypus</i>), <i>Platypus</i> (= <i>rufescens</i>)	1177
<i>discipennis</i> Schedl, <i>Platypus</i>	1135	<i>elongatus</i> Schedl, <i>Diapus</i>	1089
<i>disciporus</i> Chapuis, <i>Platypus</i> (= <i>quadridentatus</i>)	1174	<i>elongatus</i> Schedl, <i>Platyarsulus</i>	1097
<i>discoidalis</i> Schedl, <i>Platypus</i>	1135	<i>elongatus</i> Schedl, <i>Triozastus</i>	1220
<i>discolor</i> Beeson, <i>Diapus</i> (= <i>spatulifer</i>)	1092	<i>emancipatus</i> Murayama, <i>Crossotarsus</i>	1198
<i>discolor</i> Blandford, <i>Platypus</i>	1135	<i>emarginatulus</i> Browne, <i>Platypus</i>	1137
<i>discrepan</i> Schedl, <i>Doliopygus</i> (= <i>divaricus</i>)	1228	<i>emarginatus</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1166
<i>dispar</i> Schaufuss, <i>Platypus</i> (= <i>hintzi</i>)	1146	<i>emarginatus</i> Schedl, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>lunifer</i>)	1155
<i>dispar</i> Schedl, <i>Mitosoma</i>	1110	<i>emdeni</i> Schedl, <i>Platypus</i>	1137
<i>dissimilis</i> Chapuis, <i>Platypus</i>	1135	<i>emorsus</i> Beeson, <i>Crossotarsus</i>	1198
<i>dissipabilis</i> Schedl, <i>Platypus</i>	1136	<i>enixus</i> Schedl, <i>Platypus</i>	1137
<i>distinctipes</i> Schedl, <i>Platypus</i>	1136	<i>enormis</i> Schedl, <i>Platypus</i>	1137
<i>distinctus</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1213	<i>epistomalis</i> Wood, <i>Cenocephalus</i>	1109
<i>distinguendis</i> Schedl, <i>Platypus</i>	1136	<i>equadorensis</i> Schedl, <i>Platypus</i>	1137
<i>divaricus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1228	<i>erichsoni</i> (Chapuis), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1229
<i>diversifrons</i> Browne, <i>Chaetastus</i>	1108	<i>erichsoni</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167
<i>diversipennis</i> Schedl, <i>Platypus</i>	1136	<i>ericius</i> Blandford, <i>Tesseroerus</i>	1099
<i>diversiporus</i> Schedl, <i>Platypus</i> (= <i>truncatipennis</i>)	1189	<i>erinaceus</i> Schedl, <i>Mesoplatypus</i>	1222
<i>divisus</i> Browne, <i>Doliopygus</i>	1228	<i>errans</i> (Sampson), (<i>Crossotarsus</i>), <i>Platypus</i>	1137
<i>dobunabe</i> Roberts & Morimoto, <i>Platypus</i>	1136	<i>erraticus</i> Schedl, <i>Platypus</i>	1138
<i>dohrni</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1213	<i>eruditus</i> Schedl, <i>Platypus</i> (= <i>secretus</i>)	1179
<i>dolabratus</i> Blandford, <i>Platypus</i>	1136	<i>eugeniae</i> Beeson, <i>Platypus</i> (= <i>eugeniae</i>)	1138
<i>Doliopygus</i> Schedl	1223	<i>eugeniae</i> Schedl, <i>Platypus</i>	1138
<i>dolosus</i> Schedl, <i>Doliopygus</i>	1228	<i>eugustus</i> Wood, <i>Platypus</i>	1138
<i>dolus</i> Schedl, <i>Crossotarsus</i>	1198	<i>evanidinericius</i> Roberts & Morimoto, <i>Platypus</i>	1138
<i>dolus</i> Schedl, <i>Platypus</i>	1136	<i>evanidus</i> Roberts, <i>Platypus</i>	1138
<i>donisi</i> Schedl, <i>Doliopygus</i>	1228	<i>eversus</i> Wood, <i>Platypus</i>	1138
<i>doonae</i> Beeson, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>shoreanus</i>)	1181	<i>exuratus</i> Blandford, <i>Platypus</i>	1138
<i>dorsalis</i> Schedl, <i>Platypus</i>	1136	<i>exasperatus</i> Roberts, <i>Doliopygus</i> (= <i>rhodesianus</i>)	1237
<i>dorsatus</i> Schedl, <i>Platypus</i>	1136	<i>excavatus</i> (Sampson), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1229
<i>donci</i> Chapuis, <i>Platypus</i> (= <i>apicalis</i>)	1117	<i>excavatus</i> Chapuis, <i>Platypus</i> (= <i>latifinis</i>)	1152
<i>dreseheri</i> Schedl, <i>Platypus</i>	1136	<i>excedens</i> Chapuis, <i>Platypus</i>	1138
<i>dryobalanopsis</i> Browne, <i>Platypus</i>	1136	<i>excellens</i> Schedl, <i>Platypus</i>	1138
<i>dubiosus</i> Schedl, (<i>Crossotarsus</i>), <i>Triozastus</i> (= <i>marshalli</i>)	1221	<i>excisum</i> Schaufuss, <i>Mitosoma</i>	1110
<i>dubiosus</i> Schedl, (<i>Diacarus</i>), <i>Genyocerus</i> (= <i>compactus</i>)	1093	<i>excisus</i> Chapuis, <i>Platypus</i>	1138
<i>dubiosus</i> Schedl, <i>Platypus</i>	1136	<i>excisus</i> Strohmeier, <i>Periommatatus</i>	1103
<i>dubius</i> (Sampson), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1228	<i>exilis</i> (Chapuis), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1229
<i>duplicatus</i> Schedl, <i>Platypus</i>	1136	<i>exilis</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i>	1094
<i>duplosignatus</i> Schedl, <i>Platypus</i>	1136	<i>exilis</i> Chapuis, <i>Platypus</i> (= <i>solidus</i>)	1183
<i>duriororus</i> Browne, <i>Platypus</i>	1137	<i>eximius</i> Schedl, <i>Doliopygus</i>	1230
<i>durus</i> Schedl, <i>Platypus</i>	1137	<i>eximius</i> Wood, <i>Platypus</i>	1138
<i>dyerae</i> Browne, <i>Platypus</i>	1137	<i>exitialis</i> Wood, <i>Platypus</i>	1139
<i>echinatus</i> Roberts, <i>Platypus</i>	1137	<i>exitiosus</i> Schedl, <i>Platypus</i>	1139
<i>edentatus</i> Beeson, <i>Crossotarsus</i>	1198	<i>expletus</i> Schedl, <i>Doliopygus</i>	1230
<i>efferatus</i> Schedl, <i>Platypus</i>	1137	<i>exsuperans</i> Beaver, <i>Platypus</i>	1139
<i>effetus</i> Schedl, <i>Platypus</i>	1137	<i>extensus</i> Schedl, <i>Platypus</i>	1139
<i>egregius</i> Schedl, <i>Platypus</i>	1137	<i>externedentatus</i> (Fairmaire), (<i>Platypus</i>), <i>Crossotarsus</i>	1198
<i>elaboratus</i> Schedl, <i>Platypus</i>	1137	<i>externespinulosus</i> (Schedl), (<i>Platypus</i>), <i>Crossotarsus</i>	1199
<i>elegans</i> Chapuis, <i>Tesseroerus</i>	1098	<i>fugacerum</i> Browne, <i>Crossotarsus</i>	1200

- fairmairei* Chapuis, *Crossotarsus* 1200
- falcator* Beeson, *Platypus* (= *falcatus*) 1139
- falcatus* Strohmeier, *Platypus* 1139
- fallax* Schedl, *Platypus* 1139
- falsificus* (Schedl), (*Crossotarsus*), *Doliopygus* 1230
- fastuosus* Schedl, (*Platypus*), *Treptoplatypus*
(= *multioporus*) 1195
- feae* Schedl, *Platypus* (= *intermedius*) 1149
- fecundus* (Sampson), (*Platypus*), *Trachyostus* 1211
- fecundus ovalis* Schedl, *Trachyostus* (= *fecundus*) 1211
- fecundus transversus* Schedl, *Trachyostus*
(= *fecundus*) 1211
- fenestralis* Nunberg, *Doliopygus*
(= *coleocephalus*) 1227
- fenestrallatus* Roberts, *Platypus* 1139
- fergussonus* Roberts, *Genyocerus* (= *puer*) 1095
- figus* Roberts, *Platypus* 1139
- filaris* Wood, *Platypus* 1139
- filiformis* (Chapuis), (*Platypus*), *Neotrachyostus* 1213
- filipennis* Schedl, *Platypus* 1139
- firinus* Schedl, *Platypus* 1139
- fischeri* (Strohmeier), (*Crossotarsus*),
Treptoplatypus 1194
- flatesceus* (Strohmeier), (*Crossotarsus*),
Mesoplatypus 1222
- flavicornis* (Fabricius), *Platypus* 1139
- flavicornis* Dalman, *Platypus* (= *solutus*) 1184
- flavicornis* Kugelann, (*Bostrichus*), *Platypus*
(= *cylindrus*) 1132
- flavomaculatus* Strohmeier, *Crossotarsus* 1200
- flectus* Niisima & Murayama, *Platypus* 1140
- flexibilis* (Schedl), (= *Tesserocranulus*),
Cenocephalus 1109
- flexiosus* Schedl, *Platypus* 1140
- fluminalis* Beeson, *Crossotarsus* 1200
- foderatus* Beeson, *Crossotarsus* (= *inernis*) 1201
- foraminosus* Schedl, *Platypus* (= *compositus*) 1127
- forcipes* Wood, *Tesserocerus* 1099
- forcipatus* Schedl, *Doliopygus* 1230
- forcipatus* Schedl, *Platypus* 1140
- forcipatus* Schedl, *Tesserocerus* 1099
- forcipes* Schedl, *Crossotarsus* 1200
- forfex* Beeson, *Platypus* (= *uncinatus*) 1190
- forficula* Chapuis, *Platypus* 1140
- forficuloides* Schedl, *Platypus* 1141
- forficulus* (Schedl), (*Tesseroccephalus*),
Tesserocerus 1099
- formosanus* Niisima & Murayama, *Diapus* 1089
- formosanus* Niisima & Murayama, *Platypus* 1141
- formosanus* Niisima & Murayama, *Platypus*
(= *flectus*) 1140
- formosanus* Strohmeier, *Crossotarsus* 1200
- fortis* Blandford, *Platypus* (= *dentatus*) 1134
- fossulatus* Chapuis, *Platypus* 1141
- fractinectus* Schedl, *Platypus* 1141
- fracticornis* (Schedl), (*Crossotarsus*), *Platypus* 1141
- fractocostis* Schedl, *Platypus* 1141
- fractus* Sampson, *Crossotarsus* 1200
- fradei* (Tordo), (*Symmeris*), *Chactastus* 1108
- fragmentus* Sampson, *Crossotarsus*
(= *squamulatus*) 1207
- fragmentus minor* Sampson, *Crossotarsus*
(= *squamulatus*) 1207
- fragosus* Schedl, *Platypus* 1141
- franciae* Browne, *Platypus* (= *franciai*) 1141
- franciai* Browne, *Platypus* 1141
- fratellus* Schedl, *Cylindropalpus* (= *gramulosus*) 1218
- fraterculus* Schedl, *Platypus* 1141
- froggatti* Sampson, *Platypus* 1141
- frontalis* (Strohmeier), (*Diapus*), *Genyocerus* 1094
- frontalis* (Blandford), (*Platypus*), *Platyscapulus* 1215
- frontebulbifer* Schedl, *Platypus* 1141
- fronteproductus* Schedl, *Tesserocerus* 1099
- fugax* Schedl, *Platypus* 1142
- fukiensis* Schedl, *Platypus* (= *calamus*) 1123
- fulgens* Schedl, *Doliopygus* 1230
- fulgens* Schedl, *Platypus* 1142
- fulgidus* Browne, *Platypus* 1142
- fulmeki* (Schedl), (*Crossotarsus*), *Platypus* 1142
- furcatus* Blandford, *Platypus* 1142
- furtirus* (Sampson), (*Diapus*), *Genyocerus* 1096
- fuscifrons* (Chapuis), (*Platypus*), *Neotrachyostus* 1213
- fuscipilosulus* (Schedl), (*Crossotarsus*),
Doliopygus 1230
- fuscus* Chapuis, *Platypus* 1142
- gabensis* Schedl, *Platypus* 1142
- gagates* Schedl, *Platypus* 1142
- galbulimimae* Roberts & Morimoto, *Platypus* 1142
- galerus* (Schedl), (*Crossotarsus*), *Doliopygus* 1230
- garciniae* Browne, *Platypus* (= *hamaticollis*) 1144
- gebieni* Evers, *Tesserocerus* 1099
- gemellus* (Schedl), (*Crossotarsus*), *Platypus* 1142
- geminatus* Chapuis, *Platypus* 1142
- Genyocerinae* Hopkins (= *Diapodini*) 1088
- Genyocerus* Motschulsky 1093
- gerstaeckeri* Chapuis, *Platypus* 1142
- gestroi* Sampson, *Diapus* 1089
- ghanaensis* Schedl, *Trachyostus* 1211
- ghesquierei* (Schedl), (*Crossotarsus*), *Doliopygus* 1230
- gilncei* Roberts, *Platypus* 1143
- glaber* Schedl, *Doliopygus* 1230
- glabratus* Roberts, *Platypus* 1143
- globicollis* Schedl, *Platypus* 1143
- glochideus* Roberts & Morimoto, *Platypus* 1143
- godmani* Blandford, *Platypus* 1143
- goilalae* Schedl, *Platypus* 1143
- gongyloides* Roberts, *Platypus* 1143
- gotoi* Browne, *Platypus* 1143
- gracilicornis* (Schedl), (*Crossotarsus*), *Platypus* 1143
- gracilior* Browne, *Platypus* 1143
- gracilior* Schedl, *Doliopygus* 1230
- gracilis* Broun, *Platypus* 1143
- gracilis* Strohmeier, *Periommatatus* 1103

<i>granarius</i> Schedl, <i>Platypus</i>	1143	<i>hirtus</i> Schedl, <i>Platypus</i>	1146
<i>grandiclava</i> Strohmeier, <i>Mesoplatypus</i>	1222	<i>histrix</i> (Schedl), (<i>Platyscapus</i>), <i>Platypus</i>	1146
<i>grandiporus</i> Schedl, <i>Platypus</i>	1143	<i>holdhausi</i> Schedl, <i>Platypus</i>	1146
<i>grandis</i> Schedl, <i>Crossotarsus</i> (= <i>minusculus</i>)	1202	<i>hopeae</i> Beeson, <i>Genyocerus</i> (= <i>diaphanus</i>)	1094
<i>grandis</i> Schedl, <i>Mesoplatypus</i>	1222	<i>horishensis</i> Murayama, <i>Platypus</i>	1146
<i>grandis</i> Schedl, <i>Periommatius</i>	1104	<i>horni</i> Schedl, <i>Crossotarsus</i>	1200
<i>granifer</i> Schedl, <i>Platypus</i>	1143	<i>hospes</i> Schedl, <i>Platypus</i>	1146
<i>granosus</i> Browne, <i>Platypus</i>	1143	<i>humilis</i> Browne, <i>Triozastus</i> (= <i>caliginosus</i>)	1220
<i>granulipennis</i> Schedl, <i>Platypus</i>	1144	<i>humilis</i> Chapuis, <i>Platypus</i>	1146
<i>granulosus</i> (Schedl), (<i>Platyscapulus</i>), <i>Cylindropalpus</i>	1218	* <i>hurdi</i> Schedl, <i>Cenocephalus</i>	1109
<i>granulosus</i> Browne, <i>Platypus</i>	1144	<i>hybridus</i> Schedl, <i>Platypus</i>	1146
<i>granulosus</i> fratellus Schedl, (<i>Platyscapulus</i>), <i>Cylindropalpus</i> (= <i>granulosus</i>)	1218	<i>ibex</i> Schedl, <i>Doliopygus</i>	1231
<i>granulosus</i> subaffinis Schedl, (<i>Platyscapulus</i>), <i>Cylindropalpus</i> (= <i>granulosus</i>)	1218	<i>ignotus</i> Schedl, <i>Platypus</i>	1147
<i>grayi</i> Schedl, <i>Platypus</i>	1144	<i>imberbis</i> Roberts, <i>Platypus</i>	1147
<i>grayi</i> immersus Schedl, <i>Platypus</i> (= <i>grayi</i>)	1144	<i>imitator</i> Schedl, <i>Platypus</i> (= <i>shoreanus</i>)	1181
<i>gregalis</i> Schedl, <i>Platypus</i>	1144	<i>imitatrix</i> Schedl, <i>Platypus</i>	1147
<i>gressitti</i> Schedl, <i>Crossotarsus</i>	1200	<i>imitatrix</i> Schedl, <i>Baiocis</i>	1216
<i>grevilleae</i> Lea, (<i>Crossotarsus</i>), <i>Diapus</i> (= <i>pusillimus</i>)	1091	<i>immersus</i> Schedl, <i>Platypus</i> (= <i>grayi</i>)	1144
<i>grumosus</i> Roberts & Morimoto, <i>Platypus</i>	1144	<i>immunis</i> Schedl, <i>Platypus</i>	1147
<i>guadeloupensis</i> Schedl, <i>Platypus</i>	1144	<i>impar</i> (Schedl), (<i>Crossotarsus</i>), <i>Dendroplatypus</i>	1240
<i>guerini</i> Chapuis, <i>Tesserocerus</i>	1099	<i>impar</i> Schedl, <i>Scolytotarsus</i>	1085
<i>guerini</i> montanus Schedl, <i>Tesserocerus</i> (= <i>pusillimus</i>)	1099	<i>imparidens</i> Browne, <i>Platypus</i>	1147
<i>guineensis</i> Roberts, <i>Doliopygus</i>	1230	<i>impariporus</i> (Beeson), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1210
<i>gumicus</i> Roberts, <i>Platypus</i>	1144	<i>imparispinosus</i> Roberts, <i>Mesoplatypus</i>	1222
<i>gumiensis</i> Roberts, <i>Platypus</i>	1144	<i>impolitus</i> Roberts, <i>Platypus</i>	1147
<i>haagi</i> Chapuis, <i>Platypus</i>	1144	<i>imporcatus</i> Blandford, <i>Platypus</i>	1147
<i>hamaticeps</i> Schedl, <i>Platypus</i>	1144	<i>impressifrons</i> Schedl, <i>Platypus</i>	1147
<i>hamaticollis</i> Schedl, <i>Platypus</i>	1144	<i>impressus</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Platypus</i>	1147
<i>hamatipennis</i> Schedl, <i>Platypus</i>	1144	<i>impressus</i> Janson, <i>Diapus</i> (= <i>molossus</i>)	1089
<i>hamatulus</i> Schedl, <i>Platypus</i>	1144	<i>impressus</i> ater Schedl, <i>Platypus</i> (= <i>impressus</i>)	1147
<i>hamatus</i> (Nunberg), <i>Mitosoma</i>	1110	<i>inaccessus</i> Schedl, <i>Platypus</i>	1148
<i>hamatus</i> Blandford, <i>Platypus</i>	1144	<i>inaequidens</i> Browne, <i>Crossotarsus</i>	1201
<i>handeniensis</i> Nunberg, <i>Triozastus</i> (= <i>banghaasi</i>)	1220	<i>incertus</i> Schedl, <i>Platypus</i>	1148
<i>hardenbergi</i> Sampson, (<i>Crossotarsus</i>), <i>Doliopygus</i> (= <i>crinitus</i>)	1228	<i>incilis</i> Schedl, <i>Doliopygus</i>	1231
<i>hartwegi</i> Schedl, <i>Doliopygus</i>	1231	<i>incisus</i> (Sampson), (<i>Crossotarsus</i>), <i>Baiocis</i>	1216
<i>hashimotoi</i> Browne, <i>Platypus</i>	1145	<i>incognitus</i> Schedl, <i>Platypus</i>	1148
<i>hastatus</i> (Schedl), (<i>Platypus</i>), <i>Crossotarsus</i>	1200	<i>incompertus</i> Schedl, <i>Platypus</i>	1148
<i>hastulifer</i> Schedl, <i>Platypus</i>	1145	<i>inconstans</i> Schedl, <i>Platypus</i>	1148
<i>haticollis</i> Browne, <i>Genyocerus</i> (= <i>laticollis</i>)	1094	<i>incostatus</i> Schedl, <i>Platypus</i>	1148
<i>hebetatus</i> Roberts, <i>Platypus</i>	1145	<i>indicus</i> Strohmeier, <i>Platypus</i>	1148
<i>hebridensis</i> Beeson, <i>Crossotarsus</i>	1200	<i>indomitus</i> Chapuis, <i>Crossotarsus</i>	1201
<i>heritierae</i> Stebbing, (<i>Diapus</i>), <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199	<i>inermis</i> Guérin-Meneville, <i>Tesserocerus</i>	1099
<i>heterodoxus</i> Schedl, <i>Platypus</i>	1145	<i>inermis</i> Sampson, <i>Platypus</i>	1148
<i>heteromorphus</i> Schedl, <i>Platypus</i> (= <i>incertus</i>)	1148	<i>inermis</i> Schedl, <i>Crossotarsus</i>	1201
<i>hians</i> Chapuis, <i>Platypus</i>	1145	<i>inermis</i> Strohmeier, <i>Periommatius</i>	1104
<i>himalayensis</i> Beeson, <i>Diapus</i> (= sp.?)	1088	<i>infuscatus</i> Browne, <i>Platypus</i>	1148
<i>hinchuachani</i> Schedl, <i>Platypus</i>	1145	<i>inimicus</i> (Schedl), (<i>Crossotarsus</i>), <i>Baiocis</i>	1216
<i>hintzi</i> Schauffuss, <i>Platypus</i>	1145	<i>inimicus</i> Bromm, <i>Platypus</i> (= <i>gracilis</i>)	1143
<i>hippocrepicus</i> Roberts, <i>Platypus</i>	1146	<i>inornatus</i> Chapuis, <i>Crossotarsus</i> (= <i>minusculus</i>)	1202
<i>hirtellus</i> Schedl, <i>Platypus</i>	1146	<i>insculptus</i> Schedl, <i>Platypus</i>	1148
		<i>insidiosus</i> Schedl, <i>Platypus</i>	1149
		<i>insignatus</i> Schedl, <i>Platypus</i>	1149
		<i>insignis</i> (Saunders), (<i>Platypus</i>), <i>Tesserocerus</i>	1099
		<i>insiticus</i> Schedl, <i>Doliopygus</i>	1231
		<i>insularis</i> Strohmeier, <i>Platypus</i>	1149

<i>insulindicus</i> Schedl, <i>Platypus</i>	1149	<i>kuperi</i> Roberts, <i>Diapus</i>	1059
<i>integerrimus</i> Roberts, <i>Doliopygus</i> (= <i>bitalei</i>)	1225	<i>kusukusensis</i> Murayama, <i>Platypus</i>	1151
<i>integratus</i> Roberts, <i>Doliopygus</i>	1231	<i>lablabiae</i> Schedl, <i>Platypus</i>	1151
<i>interjectus</i> Schedl, <i>Doliopygus</i>	1231	<i>labrifrons</i> Schedl, <i>Platypus</i>	1151
<i>intermedius</i> (Browne), (<i>Diacarus</i>), <i>Genyocerus</i>	1094	<i>lacordairei</i> Chapuis, <i>Crossotarsus</i>	1201
<i>intermedius</i> (Schedl), <i>Platypus</i>	1149	<i>laevicollis</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167
<i>intermedius</i> Chapuis, <i>Crossotarsus</i>	1201	<i>laevigatus</i> Chapuis, <i>Crossotarsus</i>	1201
<i>intermedius</i> Schedl, <i>Spathidicerus</i> (= <i>nobilis</i>)	1101	<i>laeris</i> Browne, <i>Platypus</i>	1151
<i>interponens</i> Schedl, <i>Platypus</i>	1149	<i>lafertei</i> Chapuis, <i>Platypus</i>	1151
<i>interpositus</i> (Schedl), (<i>Platyscapulus</i>), <i>Cylindropalpus</i>	1219	<i>lalolaensis</i> Nunberg, <i>Cenocephalus</i>	1109
<i>interpositus</i> Schedl, <i>Doliopygus</i> (= <i>citri</i>)	1226	<i>laminatus</i> Schedl, <i>Platypus</i>	1152
<i>interruptus</i> (Sampson), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1231	<i>laosi</i> Schedl, <i>Platypus</i>	1152
<i>interstitialis</i> Schedl, <i>Doliopygus</i> (= <i>rapax</i>)	1237	<i>laratensis</i> Beeson, <i>Crossotarsus</i>	1201
<i>interstitialis</i> Schedl, <i>Platypus</i> (= <i>occipitalis</i>)	1161	<i>lateconcarus</i> Schedl, <i>Platypus</i>	1152
<i>interstitialis</i> Schedl, <i>Trachyostus</i>	1212	<i>latedecliris</i> Schedl, <i>Platypus</i>	1152
<i>intritus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1231	<i>lutelobatus</i> Browne, <i>Platypus</i>	1152
<i>inutilis</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1149	<i>latelunatus</i> Beeson, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>subdepressus</i>)	1155
<i>inversus</i> Roberts, <i>Platypus</i> (= <i>crassicornis</i>)	1128	<i>lateralis</i> Schedl, <i>Doliopygus</i>	1232
<i>involutus</i> Schedl, <i>Platypus</i>	1149	<i>latescapus</i> Nunberg, <i>Periommatius</i>	1104
<i>iriani</i> Browne, <i>Platypus</i>	1150	<i>latespinatus</i> Nunberg, <i>Periommatius</i>	1104
<i>irregularis</i> Browne, (<i>Diacarus</i>), <i>Genyocerus</i> (= <i>albipennis</i>)	1093	<i>latespinis</i> Schedl, <i>Diapus</i>	1059
<i>irregularis</i> Schedl, <i>Platypus</i>	1150	<i>laticollis</i> (Browne), (<i>Diacarus</i>), <i>Genyocerus</i>	1094
<i>irrepertus</i> Schedl, <i>Platypus</i>	1150	<i>laticollis</i> Chapuis, <i>Platypus</i>	1152
<i>irruptus</i> Schedl, <i>Platypus</i>	1150	<i>latidens</i> Browne, <i>Platypus</i>	1152
<i>ivagaii</i> Roberts & Morimoto, <i>Platypus</i>	1150	<i>latifinis</i> Walker, <i>Platypus</i>	1152
<i>jamaicensis</i> Bright, <i>Platypus</i>	1150	<i>latreillei</i> Chapuis, <i>Platypus</i>	1152
<i>jansoni</i> Chapuis, <i>Platypus</i>	1150	<i>latrirostris</i> (Pascoe), (<i>Psepholax</i>), <i>Protolylastes</i>	1054
<i>javanus</i> Beeson, <i>Crossotarsus</i>	1201	<i>laudatus</i> (Schedl), (<i>Platyscapus</i>), <i>Cylindropalpus</i>	1219
<i>javanus</i> Browne, <i>Platypus</i>	1150	<i>lawasensis</i> Browne, <i>Platypus</i>	1152
<i>javanus</i> Murayama, <i>Diapus</i> (= <i>aculeatus</i> ?)	1088	<i>lebasii</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1166
<i>javanus</i> Strohmeier, <i>Spathidicerus</i>	1101	<i>lebrunii</i> Schedl, <i>Doliopygus</i>	1232
<i>jeffersi</i> Browne, <i>Platypus</i> (= <i>bigranulatus</i>)	1121	<i>lecomtei</i> Schedl, <i>Doliopygus</i>	1232
<i>jelskii</i> Nunberg, <i>Platypus</i>	1150	<i>lecontei</i> Chapuis, <i>Crossotarsus</i>	1201
<i>jurioni</i> Schedl, <i>Doliopygus</i>	1231	<i>leferrei</i> Schedl, <i>Doliopygus</i>	1232
<i>juvencus</i> Schedl, <i>Platypus</i>	1150	<i>leleupi</i> Nunberg, <i>Doliopygus</i> (= <i>bidentatus</i>)	1224
<i>kakaoensis</i> Schedl, <i>Doliopygus</i>	1231	<i>leonensis</i> Roberts, <i>Doliopygus</i> (= <i>nitidulus</i>)	1234
<i>kalshoveni</i> Schedl, <i>Platypus</i>	1150	<i>leonensis</i> Roberts, <i>Platypus</i> (= <i>orientalis</i>)	1163
<i>kanarensis</i> Beeson, <i>Platypus</i> (= <i>uncinatus</i>)	1190	<i>lepidus</i> Chapuis, <i>Platypus</i>	1152
<i>karii</i> Beaver, <i>Doliopygus</i>	1231	<i>lepidus formosanus</i> Nüßima & Murayama, <i>Platypus</i> (= <i>flectus</i>)	1140
<i>katoi</i> Browne, <i>Platypus</i>	1151	<i>leprosus</i> Browne, <i>Platypus</i>	1153
<i>keelungensis</i> Browne, <i>Platypus</i>	1151	<i>lesiniformis</i> Roberts, <i>Platypus</i>	1153
<i>kenyaensis</i> Schedl, <i>Doliopygus</i> (= <i>bidentatus</i>)	1224	<i>levannongi</i> Schedl, <i>Platypus</i>	1153
<i>keyensis</i> Strohmeier, <i>Crossotarsus</i>	1201	<i>lewisi</i> Blandford, <i>Platypus</i>	1153
<i>kitushi</i> Schedl, <i>Mesoplatypus</i>	1222	<i>limbatus</i> Chapuis, <i>Platypus</i>	1153
<i>kiushuensis</i> Murayama, <i>Platypus</i>	1151	<i>linearis</i> Chapuis, <i>Tesserocerus</i>	1100
<i>klapperichi</i> (Schedl), (<i>Stenoplatypus</i>), <i>Platypus</i>	1151	<i>linearis</i> Stephens, <i>Platypus</i> (= <i>parallelus</i>)	1165
<i>kokodaensis</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1151	<i>lineaticornis</i> Schedl, <i>Platypus</i>	1153
<i>kondulensis</i> Beeson, <i>Crossotarsus</i>	1201	<i>lineellus</i> Schedl, <i>Platypus</i>	1154
<i>konincki</i> Chapuis, <i>Platypus</i>	1151	<i>lingulatus</i> Roberts & Morimoto, <i>Platypus</i>	1154
<i>koryoensis</i> (Murayama), (<i>Crossotarsus</i>), <i>Platypus</i>	1151	<i>liraticus</i> Wood, <i>Platypus</i>	1154
<i>kraatzi</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1166	<i>liratus</i> Blandford, <i>Platypus</i>	1154
<i>kreczmeri</i> Nunberg, <i>Periommatius</i>	1104	<i>lobacanthus</i> Beeson, <i>Carchesiopygus</i> (= <i>lobacanthus</i> Schedl)	1210
<i>kuntzeni</i> (Schedl), (<i>Platypus</i>), <i>Baociis</i>	1216	<i>lobacanthus</i> Schedl, <i>Carchesiopygus</i>	1210

<i>lobatum</i> Schedl. <i>Mitosoma</i>	1111	<i>malkini</i> Schedl. <i>Doliopygus</i>	1232
<i>lobatus</i> (Chapuis), (Platypus), <i>Neotrachyostus</i>	1213	<i>maniensis</i> Browne, <i>Platypus</i>	1156
<i>lobatus</i> (Schedl), (Crossotarsus), <i>Doliopygus</i>	1232	<i>manipularis</i> Schedl. <i>Platypus</i> (= <i>equadorensis</i>)	1137
<i>lobatus</i> Broun, <i>Platypus</i> (= <i>caviceps</i>)	1124	<i>manus</i> (Schedl), (Platypus), <i>Platyscapulus</i>	1215
<i>lonacensis</i> Roberts, <i>Platypus</i> (= <i>orientalis</i>)	1163	° <i>maravignae</i> Guérin-Meneville, <i>Platypus</i>	1157
<i>longicaudatus</i> Nunberg, <i>Platypus</i> (= <i>micrurus</i>)	1157	<i>marcidus</i> Blandford, <i>Platypus</i>	1157
<i>longicalcaratus</i> Roberts, <i>Platypus</i>	1154	<i>margaritaceus</i> Roberts & Morimoto, <i>Platypus</i>	1157
<i>longicollis</i> Browne, <i>Crossotarsus</i>	1202	<i>marginatus</i> Chapuis, <i>Platypus</i>	1157
<i>longicollis</i> Browne, <i>Platypus</i>	1154	<i>maritimus</i> Beeson, <i>Platypus</i> (= <i>maritimus</i> Schedl)	1157
<i>longicollis</i> Chapuis, <i>Periommatas</i>	1104	<i>maritimus</i> Schedl, <i>Platypus</i>	1157
<i>longicornis</i> (Schedl), (Platypus), <i>Crossotarsus</i>	1202	<i>marseuli</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1166
<i>longicornis</i> Browne, <i>Platypus</i> (= <i>acuticornifer</i>)	1114	<i>marshalli</i> (Sampson), (Crossotarsus), <i>Trizastus</i>	1220
<i>longior</i> Wood, <i>Platypus</i>	1154	<i>marshalli</i> Schedl. <i>Platytarsulus</i>	1097
<i>longipennis</i> Montrouzier, <i>Platypus</i>	1154	<i>matoae</i> Browne, <i>Platypus</i>	1157
<i>longispinus</i> Beeson, <i>Diapus</i> (= <i>pendleburyi</i>)	1090	<i>mattai</i> Brethes, <i>Platypus</i> (= <i>parallelus</i>)	1168
<i>longissimus</i> Roberts, <i>Platypus</i>	1154	<i>maximus</i> Browne, <i>Cylindropalpus</i>	1219
<i>longius</i> Wood, <i>Platypus</i>	1154	<i>maynei</i> Schedl. <i>Doliopygus</i>	1232
<i>longulus</i> Chapuis, <i>Platypus</i>	1154	Mecopelmini Wood	1086
<i>longus</i> Nunberg, <i>Periommatas</i>	1105	Mecopelmus Blackman	1086
<i>lopchuensis</i> Beeson, <i>Platypus</i> (= <i>lopchuensis</i> Schedl)	1155	<i>medius</i> Schedl. <i>Doliopygus</i>	1232
<i>lopchuensis</i> Schedl, <i>Platypus</i>	1154	<i>medius</i> Schedl, <i>Trachyostus</i> (= <i>schaufussi</i>)	1212
<i>lophopetali</i> Browne, <i>Platypus</i>	1155	<i>megatoma</i> Schedl. <i>Doliopygus</i>	1232
<i>loriae</i> (Schedl), (Crossotarsus), <i>Platypus</i>	1155	<i>megatoma</i> vexator Schedl, <i>Doliopygus</i> (= <i>megatoma</i>)	1233
<i>loricatus</i> (Sampson), (Crossotarsus), <i>Platypus</i>	1155	<i>melanocephalus</i> Dejean, (<i>Damicerus</i>), <i>Tesseroerus</i> (= <i>insignis</i>)	1099
<i>lucaris</i> Schedl, <i>Platypus</i>	1155	<i>melanurus</i> Chapuis, <i>Platypus</i> (= <i>pulicaris</i>)	1173
<i>lucasi</i> Chapuis, <i>Platypus</i>	1155	Mesoplatypus Strohmeier	1221
<i>luebensis</i> Nunberg, <i>Periommatas</i>	1105	<i>Mesopygus</i> Nunberg (= <i>Doliopygus</i>)	1223
<i>luederwaldti</i> Schedl, (Platypus), <i>Platyscapulus</i> (= <i>costellatus</i>)	1215	<i>mexicanus</i> Duges, (<i>Chapuisia</i>), <i>Schedlarius</i>	1087
<i>lunatellus</i> Browne, <i>Platypus</i>	1155	<i>micrographus</i> Schedl, <i>Platypus</i>	1157
<i>lunatipennis</i> Schedl, <i>Platypus</i>	1155	<i>microlunatus</i> Browne, <i>Platypus</i>	1157
<i>lunatulus</i> Browne, <i>Platypus</i> (= <i>tayabasi</i>)	1157	<i>micrurus</i> Schedl, <i>Platypus</i>	1157
<i>lunatus</i> Browne, <i>Platypus</i> (= <i>tayabasi</i>)	1157	<i>mimicus</i> (Schedl), (Crossotarsus), <i>Doliopygus</i>	1233
<i>lunifer</i> Schedl, <i>Platypus</i>	1155	<i>minacior</i> Schedl, <i>Platypus</i>	1157
<i>luniger</i> Motschulsky, <i>Platypus</i>	1155	<i>minax</i> (Walker), (Platypus), <i>Crossotarsus</i>	1202
<i>luridus</i> Chapuis, <i>Platypus</i>	1156	<i>minimus</i> Schedl, <i>Doliopygus</i>	1233
<i>luzonicus</i> Schedl, <i>Platypus</i>	1156	<i>minor</i> Beeson, <i>Diapus</i> (= <i>minor</i> Schedl)	1089
<i>macer</i> Browne, <i>Platypus</i>	1156	<i>minor</i> Browne, <i>Platypus</i>	1157
<i>machili</i> Beeson, <i>Crossotarsus</i>	1202	<i>minor</i> Roberts, <i>Trachyostus</i> (= <i>aterrimus</i>)	1211
<i>macroporus</i> Chapuis, <i>Platypus</i>	1156	<i>minor</i> Sampson, <i>Crossotarsus</i> (= <i>squamulatus</i>)	1207
<i>maculatus</i> Schaufuss, (Crossotarsus), <i>Doliopygus</i> (= <i>chapuisi</i>)	1226	<i>minor</i> Schedl, <i>Cenocrphalus</i> (= <i>pulehellus</i>)	1109
<i>maculatus</i> Schedl, <i>Scolhyotarsus</i>	1085	<i>minor</i> Schedl, <i>Diapus</i>	1089
<i>madagascariensis</i> Chapuis, <i>Platypus</i>	1156	<i>minor</i> Schedl, <i>Doliopygus</i>	1233
<i>maeklini</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167	<i>minor</i> Schedl, <i>Trachyostus</i> (= <i>sclaufussi</i>)	1212
<i>magnus</i> Browne, <i>Platypus</i>	1156	<i>minusculus</i> Chapuis, <i>Crossotarsus</i>	1202
<i>major</i> Sampson, <i>Platypus</i> (= <i>vethi</i>)	1192	<i>minusculus</i> Schedl, <i>Platypus</i>	1157
<i>major</i> Strohmeier, <i>Periommatas</i>	1105	<i>minusculus grandis</i> Schedl, <i>Crossotarsus</i> (= <i>minusculus</i>)	1202
<i>major</i> Schedl, <i>Trachyostus</i> (= <i>schaufussi</i>)	1212	<i>minusculus novaguineensis</i> Roberts, <i>Crossotarsus</i> (= <i>minusculus</i>)	1203
<i>majusculus</i> Sampson, <i>Crossotarsus</i>	1202	<i>minutissimus</i> (Schedl), (Crossotarsus), <i>Doliopygus</i>	1233
<i>malacanthae</i> Roberts, <i>Platypus</i> (= <i>impressus</i>)	1147	<i>minutissimus</i> Schedl, <i>Diapus</i>	1089
<i>malaisei</i> Schedl, <i>Platypus</i>	1156	<i>minutissimus</i> Schedl, <i>Platypus</i>	1157
<i>malayensis</i> Schedl, <i>Platypus</i> (= <i>vethi</i>)	1192	<i>minutus</i> Nunberg, <i>Periommatas</i>	1105
<i>malgassicus</i> Schedl, <i>Diapus</i>	1089	<i>minutus</i> Roberts, <i>Platypus</i> (= <i>orientalis</i>)	1163
<i>malignus</i> Schedl, <i>Platypus</i>	1156		

<i>mirabile</i> Numberg, <i>Mitosoma</i>	1111	<i>nebulosus</i> Roberts, <i>Platypus</i>	1159
<i>mirabilis</i> Menier, <i>Platytrarsulus</i>	1097	<i>negatus</i> Schedl, <i>Platypus</i>	1159
<i>mirabilis</i> Nunberg, <i>Periommatatus</i>	1105	<i>neglectus</i> Schedl, <i>Platypus</i>	1159
<i>mirandus</i> Schedl, <i>Platypus</i>	1157	<i>nenorosus</i> Roberts, <i>Platypus</i>	1159
<i>mirus</i> Sampson, (<i>Diapus</i>), <i>Genyocerus</i> (= <i>furtivus</i>)	1096	<i>neoconjunctus</i> Schedl, <i>Doliopygus</i>	1234
<i>misoolensis</i> Browne, <i>Platypus</i>	1155	<i>neopartibilis</i> Roberts, <i>Platypus</i>	1159
<i>Mitosoma</i> Chapuis	1110	<i>neoplicatus</i> Schedl, <i>Platypus</i>	1159
<i>mitratus</i> Nunberg, <i>Doliopygus</i>	1233	<i>neosolidus</i> Schedl, <i>Platypus</i>	1159
<i>Mixopygus</i> Nunberg (= <i>Doliopygus</i>)	1223	<i>Neotrachyostus</i> Browne	1213
<i>mixtus</i> Schedl, <i>Platypus</i>	1155	<i>neotruncatus</i> Schedl, <i>Platypus</i>	1159
<i>njobergi</i> Schedl, <i>Platypus</i>	1155	<i>neotruncatus</i> Schedl, <i>Platypus</i> (= <i>leprosus</i>)	1153
<i>nkusii</i> Strohmeier, <i>Periommatatus</i>	1105	<i>nepalensis</i> Schedl, <i>Crossotarsus</i>	1203
<i>nniszechi</i> Chapuis, <i>Crossotarsus</i>	1203	<i>nevermanni</i> Schedl, <i>Tesserocranulus</i>	1101
<i>modestus</i> Beaver, <i>Doliopygus</i>	1233	<i>nicobaricus</i> Beeson, <i>Crossotarsus</i> (= <i>terminatus</i>)	1205
<i>modestus</i> Blandford, <i>Platypus</i>	1155	<i>niger</i> Nunberg, <i>Platypus</i> (= <i>hiintzi</i>)	1145
<i>modigliani</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1155	<i>nigeriensis</i> Nunberg, <i>Periommatatus</i>	1105
<i>moikui</i> Schedl, <i>Spathicranuloides</i>	1210	<i>nigeriensis</i> Roberts, <i>Doliopygus</i>	1234
<i>molossus</i> Chapuis, <i>Diapus</i>	1059	<i>nigeriensis</i> Roberts, <i>Mesoplatypus</i> (= <i>venustus</i>)	1223
<i>moluccanus</i> Schedl, <i>Crossotarsus</i>	1203	<i>nigrum</i> Schaufuss, <i>Mitosoma</i>	1111
<i>montanus</i> Roberts, <i>Doliopygus</i>	1234	<i>nijimai</i> Murayama, <i>Platypus</i>	1159
<i>montanus</i> Schedl, <i>Chaetastus</i>	1105	<i>nulgiricus</i> Beeson, <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199
<i>montanus</i> Schedl, <i>Platypus</i> (= <i>orientalis</i>)	1163	<i>niponicus</i> Blandford, <i>Crossotarsus</i>	1203
<i>montanus</i> Schedl, <i>Tesserocerus</i> (= <i>guerini</i>)	1099	<i>nitens</i> Browne, <i>Doliopygus</i>	1234
<i>montivagus</i> Roberts, <i>Platypus</i> (= <i>spinulosus</i>)	1154	<i>nitens</i> Browne, <i>Platypus</i>	1159
<i>mordax</i> Sampson, <i>Platypus</i> (= <i>impressus</i>)	1147	<i>nitens</i> Chapuis, <i>Crossotarsus</i>	1203
<i>morigerus</i> Schedl, <i>Platypus</i>	1155	<i>nitescens</i> Schedl, <i>Crossotarsus</i>	1204
<i>morebeensis</i> Browne, <i>Platypus</i>	1158	<i>nitidicollis</i> Schedl, <i>Platypus</i>	1159
<i>morosus</i> Schedl, <i>Platypus</i>	1155	<i>nitidicollis</i> Strohmeier, <i>Periommatatus</i>	1105
<i>morsi</i> Chapuis, <i>Tesserocerus</i>	1100	<i>nitidior</i> (Schedl), (<i>Trachyostus</i>), <i>Crossotarsus</i>	1204
<i>motui</i> Schedl, <i>Crossotarsus</i>	1203	<i>nitidipennis</i> Schedl, <i>Platypus</i>	1159
<i>multisanti</i> Chapuis, <i>Platypus</i>	1155	<i>nitidiventris</i> Schedl, <i>Doliopygus</i> (= <i>chapuisi</i>)	1226
<i>multidentatus</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1210	<i>nitidulus</i> (Schedl), (<i>Platypus</i>), <i>Crossotarsus</i>	1204
<i>multifoccolatus</i> Schedl, <i>Platypus</i> (= <i>truncatipennis</i>)	1190	<i>nitidulus</i> Schedl, <i>Doliopygus</i>	1234
<i>multiaporus</i> (Schedl), (<i>Diapus</i>), <i>Genyocerus</i>	1094	<i>nivalis</i> Sampson, <i>Platypus</i>	1159
<i>multiaporus</i> Schedl, <i>Platypus</i>	1155	<i>nobilis</i> (Schaufuss), (<i>Platypus</i>), <i>Trachyostus</i>	1212
<i>multiaporus</i> Schedl, <i>Treptoplatypus</i>	1194	<i>nobilis</i> Chapuis, <i>Spathidicerus</i>	1101
<i>multiapunctus</i> Roberts, <i>Doliopygus</i>	1234	<i>nocuus</i> Schedl, <i>Platypus</i>	1160
<i>murayamaensis</i> Schedl, <i>Platypus</i> (= <i>flexus</i>)	1140	<i>noouadanae</i> Browne, <i>Platypus</i>	1160
<i>muricatus</i> Roberts, <i>Platypus</i>	1155	<i>norfolkensis</i> Schedl, <i>Platypus</i>	1160
<i>mussooriensis</i> Beeson, <i>Crossotarsus</i>	1203	<i>notatus</i> Schedl, <i>Doliopygus</i>	1234
<i>mutatus</i> Chapuis, <i>Platypus</i>	1155	<i>nothofagus</i> Roberts, <i>Platypus</i>	1160
<i>mutilatus</i> Chapuis, <i>Tesserocerus</i> (= <i>morsi</i>)	1100	<i>Notoplatypus</i> Lea	1097
<i>mutilus</i> Schedl, <i>Platypus</i> (= <i>bifurcus</i>)	1120	<i>norueguineensis</i> Roberts, <i>Platypus</i>	1160
<i>mutilus</i> Schedl, <i>Platypus</i> (= <i>shoreanus</i>)	1151	<i>noraguineensis</i> Roberts, <i>Crossotarsus</i> (= <i>minusculus</i>)	1203
<i>uagakii</i> Browne, <i>Platypus</i>	1159	<i>nubilosus</i> Roberts & Morimoto, <i>Baiocis</i>	1217
<i>uairobiensis</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1234	<i>nudatus</i> Wood, <i>Platypus</i>	1160
<i>nakazawai</i> Browne, <i>Crossotarsus</i>	1203	<i>nudiusculus</i> Roberts, <i>Platypus</i>	1160
<i>nanodontus</i> Roberts, <i>Diapus</i>	1090	<i>nudus</i> Schedl, <i>Platypus</i>	1160
<i>nanulus</i> Schedl, <i>Platytrarsulus</i>	1097	<i>obanensis</i> Roberts, <i>Doliopygus</i>	1234
<i>nanus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1234	<i>obconiceps</i> Schedl, <i>Mitosoma</i>	1111
<i>nanus</i> Schedl, <i>Diapus</i>	1090	<i>obliquatum</i> Schedl, <i>Mitosoma</i>	1111
<i>navarrodeandradei</i> Marcelli, <i>Platypus</i>	1159	<i>obliquecaudatus</i> Schedl 1942, <i>Platypus</i> (= <i>obliquecaudatus</i> Schedl 1936)	1160
<i>nebulosus</i> Roberts, <i>Diapus</i>	1090	<i>obliquecaudatus</i> Schedl 1936, <i>Platypus</i>	1160

<i>obliquesectus</i> Schedl 1975, <i>Platypus</i> (= <i>obliquus</i>)	1160	<i>ornatus</i> (Schedl), (<i>Tesseroecranulus</i>), <i>Cenocephalus</i>	1109
<i>obliquesectus</i> Schedl 1973, <i>Platypus</i>	1160	<i>ornatus</i> Schedl, <i>Platypus</i>	1163
<i>obliquetruncatus</i> Schedl, <i>Platypus</i>	1160	<i>othiodes</i> Schedl, <i>Platypus</i>	1163
<i>obliquus</i> Wood, <i>Neotrachyostus</i>	1214	<i>otiosus</i> Schedl, <i>Platypus</i>	1163
<i>obliquus</i> Wood, <i>Platypus</i>	1160	<i>oralicollis</i> Schedl, <i>Platypus</i>	1163
<i>obliteratus</i> Blandford, <i>Platypus</i>	1160	<i>ovalis</i> Schedl, <i>Trachyostus</i> (= <i>fecundus</i>)	1211
<i>oblongirostris</i> Beeson, <i>Crossotarsus</i> (= <i>siporanus</i>)	1206	<i>ovatus</i> Strohmeier, <i>Platypus</i>	1163
<i>oblongus</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167	<i>oxyurus</i> Dufour, <i>Platypus</i>	1163
<i>obscurus</i> Schedl, <i>Platypus</i>	1160	<i>pucificus</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1214
<i>obsitus</i> Schedl, <i>Platypus</i>	1161	<i>pahangensis</i> Schedl, <i>Platypus</i>	1164
<i>obsoletus</i> Chapuis, <i>Platypus</i> (= <i>ustulatus</i>)	1191	<i>palatus</i> Beeson, <i>Crossotarsus</i>	1204
<i>obtectus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1161	<i>pallidipennis</i> Blandford, <i>Platypus</i>	1164
<i>obtusicornis</i> Schedl, <i>Diapus</i>	1090	<i>pallidus</i> Chapuis, <i>Platypus</i>	1164
<i>obtusipennis</i> Schedl, <i>Platypus</i>	1161	<i>pallidus sabroni</i> Schedl, <i>Platypus</i> (= <i>pallidus</i>)	1164
<i>obtusum</i> Schedl, <i>Mitosoma</i>	1111	<i>panduriformis</i> Roberts, <i>Platypus</i>	1164
<i>obtusus</i> Chapuis, <i>Platypus</i>	1161	<i>papuanus</i> Roberts, <i>Genyocerus</i>	1094
<i>obtusus</i> Chapuis, <i>Platypus</i> (= <i>solidus</i>)	1183	<i>papuanus</i> Roberts, <i>Platypus</i>	1164
<i>obtusus</i> Chapuis, <i>Tesseroecerus</i> (= <i>procer</i>)	1100	<i>papuanus</i> Schedl, <i>Diapus</i>	1090
<i>obuduensis</i> Roberts, <i>Doliopygus</i>	1234	<i>papuanus</i> Schedl, <i>Spathidicrus</i>	1101
<i>occalleseus</i> Schedl, <i>Doliopygus</i>	1234	<i>papulosus</i> Roberts, <i>Platypus</i>	1164
<i>occidentalis</i> Schedl, <i>Platypus</i> (= <i>impressus</i>)	1147	<i>paradoxus</i> Schedl, <i>Platypus</i>	1164
<i>occipitalis</i> Chapuis, <i>Platypus</i>	1161	<i>paradubius</i> Roberts, <i>Doliopygus</i> (= <i>dubius</i>)	1229
<i>occipitis</i> Wood, <i>Platypus</i>	1161	<i>parallelicentrus</i> Roberts, <i>Platypus</i>	1164
<i>ochthocosmus</i> Roberts, <i>Platypus</i> (= <i>rufescens</i>)	1177	<i>parallelus</i> (Fabricius), (<i>Bostrichus</i>), <i>Platypus</i>	1164
<i>octostatus</i> Schedl, <i>Crossotarsus</i>	1204	<i>paramonori</i> Schedl, <i>Platypus</i>	1168
<i>octodentatus</i> Browne, <i>Doliopygus</i>	1235	<i>parapetax</i> Roberts & Morimoto, <i>Platypus</i>	1168
<i>octodentatus</i> Browne, <i>Platypus</i>	1161	<i>parinaiae</i> Schedl, <i>Platypus</i> (= <i>rufescens</i>)	1177
<i>octospinosum</i> Schedl, <i>Mitosoma</i>	1111	<i>partibilis</i> Roberts, <i>Platypus</i>	1168
<i>octospinosus</i> Browne, <i>Platypus</i>	1161	<i>particeps</i> Schedl, <i>Platypus</i>	1168
<i>oculatus</i> (Beeson), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1210	<i>partitus</i> Schedl, <i>Platypus</i>	1168
<i>odonticeps</i> Schedl, <i>Mitosoma</i>	1111	<i>parculus</i> Schedl, <i>Platypus</i> (= <i>puerulus</i>)	1173
<i>ohnoi</i> Browne, <i>Platypus</i>	1161	<i>parvus</i> Browne, (<i>Trachyostus</i>), <i>Crossotarsus</i> (= <i>saltator</i>)	1205
<i>oligodontus</i> Roberts, <i>Crossotarsus</i>	1204	<i>parvus</i> Browne, <i>Platypus</i>	1168
<i>olivieri</i> Chapuis, <i>Platypus</i>	1161	<i>parvus</i> Nunberg, <i>Mesoplatypus</i>	1222
<i>omega</i> Schedl, <i>Platypus</i>	1161	<i>parysi</i> Chapuis, <i>Platypus</i> (= <i>lafertei</i>)	1151
<i>omissus</i> Schedl, <i>Doliopygus</i>	1235	<i>pasanaie</i> Schedl, <i>Platypus</i>	1168
<i>omissus</i> Schedl, <i>Platypus</i>	1161	<i>pasohensis</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1217
<i>omnirorus</i> (Lea), (<i>Crossotarsus</i>), <i>Platypus</i>	1161	<i>patens</i> Roberts, <i>Platypus</i>	1168
<i>oneratus</i> Schedl, <i>Platypus</i>	1162	<i>patulus</i> Chapuis, <i>Platypus</i>	1168
<i>oomsis</i> Roberts, <i>Diapus</i>	1090	<i>paucidentatus</i> Roberts, <i>Crossotarsus</i>	1204
<i>opacicauda</i> Browne, <i>Platypus</i>	1162	<i>paulianum</i> Schedl, <i>Mitosoma</i>	1111
<i>opacicaudulus</i> Browne, <i>Platypus</i>	1162	<i>pectinatus</i> Browne, <i>Crossotarsus</i>	1204
<i>opacideclivis</i> Schedl, <i>Platypus</i>	1162	<i>pectinatus</i> Roberts, <i>Platypus</i>	1168
<i>opacifrons</i> Schedl, <i>Platypus</i>	1162	<i>pedellus</i> Schedl, <i>Platypus</i>	1168
<i>opacipennis</i> Schedl, <i>Platypus</i>	1162	<i>pedum</i> Sampson, <i>Platypus</i>	1168
<i>opacurus</i> Browne, <i>Platypus</i>	1162	<i>pendleburgi</i> Schedl, <i>Diapus</i>	1090
<i>opifex</i> (Sampson), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1235	<i>penctralis</i> Sampson, <i>Platypus</i> (= <i>hintzi</i>)	1146
<i>opimus</i> Schedl, <i>Platypus</i>	1162	<i>penicillatus</i> Chapuis, <i>Crossotarsus</i> (= <i>cineinnatus</i>)	1197
<i>opulentus</i> Schedl, <i>Doliopygus</i>	1235	<i>penicillatus</i> Schedl, <i>Periommatas</i>	1106
<i>oraelatus</i> Roberts, <i>Doliopygus</i>	1235	<i>pennatus</i> Schedl, <i>Platypus</i>	1168
<i>oreogenus</i> Roberts, <i>Diapus</i>	1090	<i>perangustus</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1217
<i>orientalis</i> (Strohmeier), (<i>Stenoplatypus</i>), <i>Platypus</i>	1162	<i>perbinodulus</i> Schedl, <i>Platypus</i>	1168
<i>ornaticeps</i> (Schedl), (<i>Stenoplatypus</i>), <i>Platypus</i>	1163	<i>perbrevis</i> Browne, <i>Platypus</i>	1168
<i>ornatifrons</i> Schedl, <i>Platypus</i>	1163	<i>perbrevis</i> Schedl, <i>Doliopygus</i>	1235

- perbrevis brevisculus* Schedl, *Doliopygus* 1237
 (= *rapax*)
- perbrevis solidus* Schedl, *Doliopygus* 1235
 (= *perbrevis*)
- perceptus* Beeson, *Crossotarsus* (= *ursus*) 1208
- percomis* Schedl, *Platypus* 1169
- perdifficilis* Schedl, *Platypus* (= *subitarius*) 1186
- perdiligens* Schedl, *Platypus* 1169
- perditus* Schedl, *Platypus* 1169
- peregrinus* Schedl, *Platypus* (= *longulus*) 1154
- perforans* Schedl, *Platypus* 1169
- perfossus* Chapuis, *Platypus* (= *compositus*) 1126
- perinimicus* (Schedl), (*Platypus*), *Baiocis* 1217
- Periommatinae* Schedl (= *Tesserocerini*) 1088
- Periommatus* Chapuis 1102
- permarginatus* Schedl, *Platypus* 1169
- permimicus* Schedl, *Platypus* 1169
- perminutissimus* (Schedl), (*Crossotarsus*),
Doliopygus 1235
- permodestus* Schedl, *Platypus* 1169
- pernanulus* (Schedl), (*Crossotarsus*), *Baiocis* 1217
- pernix* (Schedl), (*Platyscapus*), *Cylindropalpus* 1219
- pernudus* Schedl, *Platypus* 1169
- perplandus* Schedl, *Platypus* (= *dissipabilis*) 1136
- perplexus* Bright, *Platypus* 1169
- perpusillus* Chapuis, *Platypus* (= *hians*) 1145
- perpygmaeus* Roberts, *Diapus* 1090
- perrisi* Chapuis, *Platypus* 1169
- persimilis* (Schedl), (*Symmerus*), *Chaetastus* 1108
- pertenuis* Schedl, (*Crossotarsus*), *Triozastus*
 (= *marshalli*) 1221
- pertinax* (Schedl), (*Platyscapus*), *Cylindropalpus* 1219
- pertusus* Chapuis, *Platypus* 1169
- peruanus* Nunberg, *Platypus* 1169
- petalinus* Roberts & Morimoto, *Platypus* 1169
- petaloideus* Roberts, *Platypus* 1169
- petax* Schedl, *Platypus* 1169
- petersi* Chapuis, *Platypus* (= *ustulatus*) 1191
- philippinensis* (Blandford), (*Platypus*),
Dendroplatypus 1240
- philippinensis* (Schedl), (*Diacavus*), *Genyocerus* 1095
- philippineusis* Schedl, *Spathidicerus* 1101
- pieceae* Stebbing, *Crossotarsus* 1204
- piceus* Chapuis, *Crossotarsus* 1204
- piceus* Schedl, *Platypus* (= *cupulatus*) 1129
- piceus* Strohmer, *Periommatus* 1106
- picius* Schedl, *Platypus* 1170
- pierloti* Browne, *Platypus* 1170
- pilidens* Schedl, *Platypus* 1170
- pilifrons* Chapuis, *Platypus* (= *solidus*) 1183
- pilosulus* (Schedl), (*Crossotarsus*), *Triozastus* 1221
- pini* Hopkins, *Platypus* 1170
- piniperda* Beeson, *Platypus* (= *piniperda* Schedl) 1170
- piniperda* Schedl, *Platypus* 1170
- piniperda cambodianus* Schedl, *Platypus*
 (= *piniperda*) 1170
- piptadeniae* Schedl, *Doliopygus* (= *exilis*) 1229
- planifrons* Schedl, *Periommatus* (= *truncatiformis*) 1107
- planodecliris* Schedl, *Platypus* 1170
- planum* Schaufuss, *Mitosoma* 1111
- Platypicerinae* Nunberg (= *Tesserocerini*) 1088
- Platypicerus* Nunberg (= *Mitosoma*) 1110
- Platypides* Erichson (= *Platypodidae*) 1084
- Platypites* Erichson (= *Platypodidae*) 1084
- Platypinus* Schedl (= *Platypus*) 1113
- Platypodae* Erichson (= *Platypodidae*) 1084
- Platypodariae* Erichson (= *Platypodini*) 1113
- Platypodidae* Erichson 1084
- Platypodinae* Erichson 1084, 1113
- Platypodini* Erichson 1084, 1113
- platypoides* (Browne), (*Crossotarsus*), *Platypus* 1170
- platypus* Duftschmidt, (*Cylindra*), *Platypus*
 (= *cylindrus*) 1132
- Platypus* Herbst 1113
- Platyscapulus* Schedl 1214
- Platyscapus* Schedl (= *Platypus*) 1214
- Platyarsulidae* Schedl, part (= *Platypodini*) 1084
- Platyarsulinae* Schedl, part (= *Platypodini*) 1084, 1097
- Platyarsulus* Schedl 1097
- platyurius* Schedl, *Platypus* (= *coronatus*) 1128
- plicatus* Brethes, *Platypus* (= *mutatus*) 1159
- plumatus* (Schedl), (*Diapus*), *Genyocerus* 1095
- plumatus* Schedl, *Platypus* 1170
- podocarpi* Browne, *Platypus* 1170
- poeyi* Guerin-Meneville, *Platypus* (= *parallelus*) 1165
- politus* Chapuis, *Platypus* 1171
- ponamae* Schedl, *Platypus* (= *distinctipes*) 1136
- porcatus* Roberts, *Crossotarsus* 1204
- porcellinus* Schedl, *Platypus* 1171
- porcellus* Schedl, *Platypus* 1171
- poricollis* Schedl, *Platypus* 1171
- poriferus* Chapuis, *Platypus* (= *segnis*) 1179
- porosulus* Browne, *Platypus* 1171
- porosus* Blandford, *Platypus* 1171
- porosus* Schedl, *Platypus* (= *poricollis*) 1171
- porrectus* Chapuis, *Platypus* 1171
- postialis* Schedl, *Doliopygus* (= *bidentatus*) 1224
- posticus* Broun, (*Platypus*), *Crossotarsus*
 (= *externedentatus*) 1199
- pouteriae* Wood, *Platypus* 1171
- praealtus* Schedl, *Platypus* 1171
- praececlens* Schedl, *Platypus* 1171
- praeclarus* (Schedl), (*Crossotarsus*), *Doliopygus* 1235
- praemorsus* (Dalman), (*Platypus*), *Doliopygus* 1236
- praepositus* Schedl, *Platypus* 1171
- praeteritus* Schedl, *Platypus* 1171
- praetermissus* Schedl, *Platypus* 1171
- praenstus* Dejean, *Platypus* (= *parallelus*) 1165
- praevius* Chapuis, *Platypus* (= *parallelus*) 1167
- prenexus* Wood, *Platypus* 1171
- pretiosus* Schedl, *Platypus* (= *rugosifrons*) 1177
- principeensis* (Schedl), (*Stenoplatypus*), *Platypus* 1171

<i>priscus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>deperditus</i>)	1134	<i>pusillinus</i> Chapuis, <i>Diapus</i>	1090
<i>procer</i> (Erichson), (<i>Platypus</i>), <i>Tesseroerus</i>	1100	<i>pusillimus</i> (Chapuis), (<i>Platypus</i>), <i>Platyscapulus</i>	1215
<i>prociidus</i> Schedl, <i>Crossotarsus</i>	1204	<i>pusillus</i> (Schedl) 1952, (<i>Platypus</i>), <i>Cylindropalpus</i>	1219
<i>productus</i> Shuckard, <i>Tesseroerus</i>	1100	<i>pusillus</i> Browne, <i>Platypus</i> (= <i>broenei</i>)	1122
<i>prolongatus</i> Schedl, <i>Doliopygus</i>	1236	<i>pusillus</i> Schedl 1957, (<i>Platyscapulus</i>), <i>Cylindropalpus</i> (= <i>pusillus</i> 1952)	1219
<i>propatulus</i> Schedl, <i>Triozastus</i> (= <i>marshalli</i>)	1221	<i>pusillus</i> Schedl, <i>Cenocephalus</i>	1109
<i>propinquus</i> Schedl, <i>Doliopygus</i>	1236	<i>putzeysi</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1214
<i>propinquus</i> Schedl, <i>Platypus</i>	1172	<i>pygmaeolus</i> Schedl, <i>Doliopygus</i>	1236
<i>protenus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1172	<i>pygmaeus</i> Schedl, <i>Doliopygus</i> (= <i>exilis</i>)	1229
<i>Protohyastes</i> Wood	1054	<i>pygmaeus</i> Schedl, <i>Platypus</i> (= <i>orientalis</i>)	1162
<i>Protoplatypus</i> Wood	1056	<i>Pygodolus</i> Nunberg (= <i>Doliopygus</i>)	1223
<i>proxilus</i> Schedl, <i>Doliopygus</i>	1236	<i>quadricaudatus</i> Schedl, <i>Platypus</i>	1174
<i>proximus</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1166	<i>quadricaudatus</i> (Strohmeyer), (<i>Platypus</i>), <i>Crossotarsus</i>	1204
<i>proximus</i> Schedl, <i>Doliopygus</i>	1236	<i>quadricinctus</i> Schedl, <i>Platypus</i>	1174
<i>pseudocamerunus</i> Nunberg, <i>Periommatas</i>	1106	<i>quadricornis</i> Schedl, <i>Platypus</i>	1174
<i>pseudocupulatus</i> Schedl, <i>Platypus</i>	1172	<i>quadricornutus</i> Roberts, <i>Platypus</i> (= <i>pulvinatus</i>)	1173
<i>pseudocupulatus arceus</i> Schedl, <i>Platypus</i> (= <i>pseudocupulatus</i>)	1172	<i>quadricuspidatus</i> Schedl, <i>Platypus</i> (= <i>trispinosus</i>)	1189
<i>pseudocupulatus sundensis</i> Schedl, <i>Platypus</i> (= <i>pseudocupulatus</i>)	1172	<i>quadridens</i> Browne, (<i>Diacarus</i>), <i>Genyocerus</i> (= <i>philippinensis</i>)	1095
<i>pseudocurtus</i> Schedl, <i>Platypus</i>	1172	<i>quadridens</i> Schedl, <i>Platypus</i> (= <i>pini</i>)	1170
<i>pseudodignatus</i> Schedl, <i>Platypus</i>	1172	<i>quadridens</i> Schedl, <i>Tesseroerus</i>	1100
<i>pseudogracilis</i> Nunberg, <i>Periommatas</i>	1106	<i>quadridentatus</i> (Olivier), (<i>Scolytus</i>), <i>Platypus</i>	1174
<i>pseudolongulus</i> Schedl, <i>Platypus</i>	1172	<i>quadrifissilis</i> Schedl, <i>Platypus</i>	1174
<i>pseudomajor</i> Schedl, <i>Periommatas</i>	1106	<i>quadriforeolatus</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i>	1095
<i>pseudoniponicus</i> Sampson, <i>Crossotarsus</i> (= <i>ursus</i>)	1208	<i>quadrilobus</i> (Blandford), (<i>Platypus</i>), <i>Neotrachyostus</i>	1214
<i>pseudoopacus</i> Schedl, <i>Platypus</i> (= <i>froggatti</i>)	1141	* <i>quadrilobus</i> Schedl, <i>Cenocephalus</i>	1109
<i>pseudoplicatus</i> Schedl, <i>Platypus</i>	1172	<i>quadrinotatus</i> Schedl, <i>Platypus</i>	1174
<i>pseudoporus</i> Schedl, <i>Platypus</i>	1173	<i>quadriporus</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i>	1175
<i>pseudoselysi</i> Schedl, <i>Platypus</i>	1173	<i>quadriporus</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i>	1095
<i>pseudoserratus</i> Roberts, <i>Doliopygus</i>	1236	<i>quadriporus</i> Schedl, <i>Treptoplatypus</i>	1195
<i>pseudoseverini</i> Nunberg, <i>Periommatas</i>	1106	<i>quadrispinatus</i> Chapuis, <i>Diapus</i>	1091
<i>pseudosindora</i> Browne, <i>Platypus</i>	1173	<i>quadrispinatus</i> Chapuis, <i>Platypus</i>	1175
<i>pseudosolidus</i> Schedl, <i>Platypus</i>	1173	<i>quadrispinis</i> Browne, <i>Mesoplatypus</i>	1222
<i>pseudoternatus</i> Schedl, <i>Platypus</i>	1173	<i>quadrispinis</i> Browne, <i>Platypus</i>	1175
<i>psilicanthurus</i> (Beeson), (<i>Crossotarsus</i>), <i>Car-</i> <i>chesiopygus</i>	1210	<i>quadrispinosus</i> Schedl, <i>Diapus</i>	1091
<i>ptchoides</i> Schedl, <i>Platypus</i>	1173	<i>quaesitus</i> Schedl, <i>Platypus</i>	1175
<i>puer</i> (Schedl), (<i>Diapus</i>), <i>Genyocerus</i>	1095	<i>queenslandi</i> Schedl, <i>Platypus</i>	1175
<i>puerfergussonus</i> Roberts, <i>Genyocerus</i> (= <i>puer</i>)	1095	<i>querceus</i> Wood, <i>Platypus</i>	1175
<i>puer puer</i> (Schedl), (<i>Diapus</i>), <i>Genyocerus</i> (= <i>puer</i>)	1095	<i>querci</i> Browne, <i>Platypus</i>	1175
<i>puerulus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1173	<i>quercicola</i> Schedl, <i>Platypus</i>	1175
<i>puerulus parvulus</i> Schedl, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>puerulus</i>)	1173	<i>quercinus</i> Schedl, <i>Platypus</i>	1175
<i>pulchellus</i> (Chapuis), (<i>Platypus</i>), <i>Platyscapulus</i>	1215	<i>quercivorus</i> Murayama, <i>Platypus</i>	1175
<i>pulchellus</i> Schedl, <i>Cenocephalus</i>	1109	<i>quinquecinctus</i> (Schedl), (<i>Doliopygus</i>), <i>Mesoplatypus</i>	1222
<i>pulchellus minor</i> Schedl, <i>Cenocephalus</i> (= <i>pulchellus</i>)	1109	<i>quinquecinctus cachani</i> Schedl, (<i>Doliopygus</i>), <i>Mesoplatypus</i> (= <i>quinquecinctus</i>)	1222
<i>pulcher</i> (Chapuis), (<i>Platypus</i>), <i>Platyscapulus</i>	1215	<i>quinquecostatus</i> Chapuis, <i>Platypus</i>	1175
<i>pulicaris</i> Chapuis, <i>Platypus</i>	1173	<i>quinquespinatus</i> Chapuis, <i>Diapus</i>	1091
<i>pulvinatus</i> Roberts & Morimoto, <i>Platypus</i>	1173	<i>quinquespinatus</i> Nunberg, <i>Periommatas</i>	1106
<i>pumilio</i> (Schedl), (<i>Platyscapus</i>), <i>Cylindropalpus</i>	1219	<i>ramali</i> Schedl, <i>Platypus</i>	1176
<i>pumilus</i> Browne, <i>Platypus</i>	1174	<i>ramosissimus</i> Roberts & Morimoto, <i>Platypus</i>	1176
<i>punctatus</i> Strohmeyer, <i>Trachyostus</i> (= <i>nobilis</i>)	1212	<i>ramulosus</i> Roberts & Morimoto, <i>Platypus</i>	1176
<i>punctiventris</i> Schedl, <i>Doliopygus</i>	1236		
<i>punctulatus</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167		

<i>rapax</i> (Sampson). <i>Doliopygus</i>	1236	<i>saltatorinus</i> (Schedl). (<i>Platypus</i>), <i>Crossotarsus</i>	1205
<i>rasilis</i> Roberts. <i>Crossotarsus</i>	1205	<i>salvini</i> Blandford. <i>Platypus</i>	1177
<i>rasilis</i> Schedl. <i>Platypus</i>	1176	<i>sampsoni</i> Browne. <i>Crossotarsus</i> (= <i>trachypennis</i>)	1205
<i>rastellus</i> Beeson. <i>Crossotarsus</i>	1205	<i>sampsoni</i> (Schedl). (<i>Stenoplatypus</i>), <i>Platypus</i>	1177
<i>ratzeburgi</i> Chapuis. <i>Platypus</i>	1176	<i>sandakauensis</i> Browne. <i>Platypus</i>	1175
<i>raucus</i> Schedl. <i>Platypus</i>	1176	<i>sandakanus</i> Beeson. (<i>Crossotarsus</i>). <i>Platypus</i> (= <i>subdepressus</i>)	1185
<i>rectangulatus</i> Sampson. <i>Platypus</i> (= <i>uncinatus</i>)	1191	<i>sautacruzensis</i> Mutchler. <i>Platypus</i>	1175
<i>refertus</i> Schedl. <i>Platypus</i> (= <i>orientalis</i>)	1162	<i>santiriae</i> Roberts. <i>Doliopygus</i>	1237
<i>refertus lomaensis</i> Roberts. <i>Platypus</i> (= <i>orientalis</i>)	1163	<i>santiriae</i> Roberts. <i>Platypus</i>	1175
<i>refertus montanus</i> Schedl. <i>Platypus</i> (= <i>orientalis</i>)	1163	<i>sarawakensis</i> (Schedl). <i>Platypus</i>	1175
<i>refertus togelii</i> Roberts. <i>Platypus</i> (= <i>orientalis</i>)	1163	<i>saundersi</i> Chapuis. <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199
<i>refractus</i> Roberts & Morimoto. <i>Platypus</i>	1176	<i>saundersi submontanus</i> Beeson. <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199
<i>regularis</i> Chapuis. <i>Platypus</i> (= <i>parallelus</i>)	1166	<i>saundersi usambariensis</i> Strohmeier. <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199
<i>regularis</i> Schedl. <i>Doliopygus</i>	1237	<i>sauteri</i> (Strohmeier). (<i>Crossotarsus</i>). <i>Crossotarsinus</i>	1210
<i>reichei</i> Chapuis. <i>Platypus</i>	1176	<i>scalaris</i> Schedl. <i>Platypus</i>	1175
<i>rengetensis</i> Niisima & Murayama. <i>Crossotarsus</i>	1205	<i>scalptor</i> Schedl. <i>Platypus</i>	1175
<i>reticulatus</i> Chapuis. <i>Platypus</i> (= <i>parallelus</i>)	1166	<i>schaufussi</i> (Strohmeier). (<i>Platypus</i>). <i>Trachyostus</i>	1212
<i>retusipennis</i> Schedl. <i>Platypus</i>	1176	<i>schaufussi major</i> Schedl. <i>Trachyostus</i> (= <i>schaufussi</i>)	1212
<i>retusulus</i> Schedl. <i>Platypus</i>	1176	<i>schaufussi medius</i> Schedl. <i>Trachyostus</i> (= <i>schaufussi</i>)	1212
<i>retusus</i> Guerin-Meneville. <i>Tesserocerus</i>	1100	<i>schaufussi minor</i> Schedl. <i>Trachyostus</i> (= <i>schaufussi</i>)	1212
<i>retusus</i> Schedl. <i>Doliopygus</i>	1237	<i>schaumi</i> Chapuis. <i>Platypus</i> (= <i>pulcaris</i>)	1173
<i>retusus</i> Strohmeier. <i>Platypus</i>	1176	Schedlarini Wood	1087
° <i>rhinoceroide</i> (Schawaller). (<i>Mitosoma</i>). <i>Cenocephalus</i>	1109	Schedlarius Wood	1087
<i>rhizophorae</i> Roberts. <i>Doliopygus</i> (= <i>dubius</i>)	1229	schedli Browne. <i>Crossotarsus</i>	1205
<i>rhodesianus</i> (Schedl). (<i>Crossotarsus</i>). <i>Doliopygus</i>	1237	schedli Wood. <i>Platypus</i> (= <i>tiroiensis</i>)	1155
<i>rimulosus</i> Schedl. <i>Platypus</i>	1176	schenklingi (Strohmeier). (<i>Crossotarsus</i>). <i>Platypus</i>	1175
<i>roberti</i> Chapuis. <i>Platypus</i>	1176	schmidt i Chapuis. <i>Platypus</i>	1175
<i>robertsi</i> Nunberg. <i>Periomnatus</i>	1106	schmutzenhoferi Schedl. <i>Tesserocerus</i>	1100
<i>robustum</i> Schedl. <i>Cenocephalus</i> (= <i>robustus</i>)	1109	schoutedeni (Sampson). (<i>Crossotarsus</i>). <i>Doliopygus</i>	1237
<i>robustum</i> Schaufuss. <i>Mitosoma</i>	1111	schoutedeni (Schedl). (<i>Platypus</i>). <i>Trachyostus</i>	1212
<i>robustus</i> Browne. <i>Doliopygus</i>	1237	schultzeanus Schedl. <i>Platypus</i>	1175
<i>robustus</i> Chapuis. <i>Platypus</i>	1177	schultzei Schedl. (<i>Crossotarsus</i>). <i>Platypus</i> (= <i>schultzeanus</i>)	1175
<i>robustus</i> Schedl. <i>Cenocephalus</i>	1109	schultzei Strohmeier. <i>Platypus</i> (= <i>luniger</i>)	1156
<i>robustus</i> Schedl. <i>Diapus</i>	1092	Scolytotarsus Schedl	1085
<i>robustus</i> Schedl. (<i>Symmerus</i>). <i>Chaetastus</i> (= <i>tuberculatus</i>)	1105	scorpius (Schedl). (<i>Platypus</i>). <i>Crossotarsus</i>	1205
<i>roppai</i> Schedl. <i>Platypus</i>	1177	scrobicularis Roberts. <i>Platypus</i>	1175
<i>rostratus</i> Beeson. <i>Diapus</i> (= sp.?)	1085	<i>Scutopygus</i> Nunberg (= <i>Doliopygus</i>)	1223
<i>rotundatus</i> Chapuis. <i>Platypus</i> (= <i>parallelus</i>)	1166	secretus Sampson. <i>Platypus</i>	1175
<i>rotundicauda</i> Motschulsky. <i>Platypus</i>	1177	secus Wood. <i>Platypus</i>	1179
<i>rudifrons</i> Chapuis. <i>Platypus</i> (= <i>compositus</i>)	1127	sedulus Schedl. <i>Crossotarsus</i> (= <i>terminatus</i>)	1205
<i>rudis</i> Chapuis. <i>Platypus</i> (= <i>solidus</i>)	1184	segnis Chapuis. <i>Platypus</i>	1179
<i>rudis</i> Chapuis. <i>Tesserocerus</i>	1100	scyllaeformis Roberts & Morimoto. <i>Platypus</i>	1179
<i>rufescens</i> (Strohmeier). (<i>Crossotarsus</i>). <i>Platypus</i>	1177	selysi Chapuis. <i>Platypus</i>	1179
<i>rufescens parinariae</i> Schedl. <i>Platypus</i> (= <i>rufescens</i>)	1177	semicarinatus Schedl. <i>Platypus</i>	1179
<i>rufobrunneus</i> Schedl. <i>Platypus</i>	1177	semicinctus Schedl. <i>Crossotarsus</i>	1205
<i>rugicollis</i> Schedl. <i>Cenocephalus</i>	1109	semiermis Schedl. <i>Platypus</i>	1179
<i>rugosifrons</i> Schedl. <i>Platypus</i>	1177		
<i>rugosum</i> Schaufuss. <i>Mitosoma</i>	1111		
<i>rugosus</i> Chapuis. <i>Platypus</i> (= <i>compositus</i>)	1127		
<i>rugulosus</i> Chapuis. <i>Platypus</i> (= <i>parallelus</i>)	1166		
<i>runcinatus</i> Roberts. <i>Platypus</i>	1177		
<i>sabroni</i> Schedl. <i>Platypus</i> (= <i>pallidus</i>)	1164		
<i>sallei</i> Chapuis. <i>Platypus</i>	1177		
<i>saltator</i> Schedl. <i>Crossotarsus</i>	1205		

<i>semigranulosus</i> (Sampson), (<i>Crossotarsus</i>), <i>Platypus</i>	1179	<i>sira</i> Beeson, <i>Crossotarsus</i>	1206
<i>seminitratus</i> Nunberg, <i>Doliopygus</i> (= <i>aequalidentatus</i>)	1223	<i>sobrinus</i> Schedl, <i>Platypus</i>	1182
<i>seminitens</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1217	<i>solicitatus</i> Schedl, <i>Platypus</i>	1182
<i>semiotopus</i> Strohmeier, <i>Platypus</i>	1179	<i>solidulus</i> Browne, <i>Platypus</i>	1182
<i>semipilosus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1237	<i>solidus</i> Schedl, <i>Doliopygus</i> (= <i>perbrevis</i>)	1235
<i>semisulcatus</i> Schedl, <i>Platypus</i>	1180	<i>solidus</i> Walker, <i>Platypus</i>	1182
<i>senex</i> Wood, <i>Platypus</i>	1180	<i>solidus exilis</i> Chapuis, <i>Platypus</i> (= <i>solidus</i>)	1183
<i>seni</i> Beeson, <i>Platypus</i> (= <i>hybridus</i>)	1147	<i>solidus obtusus</i> Chapuis, <i>Platypus</i> (= <i>solidus</i>)	1183
<i>sepaloideus</i> Roberts, <i>Platypus</i>	1180	<i>solidus rudis</i> Chapuis, <i>Platypus</i> (= <i>solidus</i>)	1184
<i>sequius</i> Schedl, <i>Platypus</i>	1180	<i>solomonicus</i> Browne, <i>Baiocis</i>	1217
<i>sericans</i> Schedl, <i>Platypus</i>	1180	<i>solomonicus</i> Browne, <i>Platypus</i>	1184
<i>serratulus</i> (Browne), (<i>Platypus</i>), <i>Crossotarsus</i>	1205	<i>solutus</i> Schedl, <i>Platypus</i>	1184
<i>serratulus</i> Schedl, <i>Doliopygus</i>	1238	<i>Spathicranuloides</i> Schedl	1210
<i>serratus</i> (Schedl), (<i>Diapus</i>), <i>Genyocerus</i>	1095	<i>Spathidicerus</i> Chapuis	1101
<i>serratus</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1238	<i>spathulifer</i> Beeson, <i>Diapus</i> (= <i>spatulifer</i>)	1092
<i>setaceus</i> Chapuis, <i>Platypus</i>	1180	<i>spatiosus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1238
<i>Setantus</i> Nunberg, <i>Periommatatus</i>	1102	<i>spatulifer</i> Browne, <i>Diapus</i>	1092
<i>severini</i> Blandford, <i>Platypus</i>	1180	<i>spectabilis</i> Schedl, <i>Doliopygus</i>	1238
<i>severini</i> Strohmeier, <i>Periommatatus</i>	1106	<i>spectabilis</i> Schedl, <i>Platypus</i>	1184
<i>sexcostatus</i> Chapuis, <i>Platypus</i>	1180	<i>spectrum</i> Schedl, <i>Crossotarsus</i>	1206
<i>sexfenestratus</i> Beeson, (<i>Crossotarsus</i>), <i>Platypus</i> (= <i>quercivorus</i>)	1175	<i>spectus</i> Wood, <i>Platypus</i>	1184
<i>sexforaminatus</i> Beeson, <i>Platypus</i> (= <i>sexporus</i>)	1180	<i>spiculatus</i> Browne, <i>Platypus</i>	1184
<i>sexporus</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i>	1095	<i>spinatus</i> Browne, <i>Diapus</i>	1092
<i>sexporus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i>	1180	<i>spinax</i> Blandford, <i>Tesserocerus</i>	1100
<i>sexspinosum</i> Schedl, <i>Mitosoma</i>	1111	<i>spinidens</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1238
<i>sexualis</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i>	1180	<i>spinifer</i> (Sampson), (<i>Platypus</i>), <i>Crossotarsus</i>	1206
<i>seydeli</i> (Nunberg), (<i>Stenoplatypus</i>), <i>Platypus</i>	1181	<i>spinifer</i> Schedl, <i>Diapus</i>	1092
<i>shenefelti</i> (Nunberg), (<i>Crossotarsus</i>), <i>Platyscapulus</i>	1215	<i>spinipennis</i> Schedl, <i>Crossotarsus</i>	1206
<i>shillongensis</i> Beeson, <i>Platypus</i> (= <i>shillongensis</i> Schedl)	1181	<i>spiniventris</i> Schedl, <i>Platypus</i>	1184
<i>shillongensis</i> Schedl, <i>Platypus</i>	1181	<i>spinolae</i> Chapuis, <i>Tesserocerus</i>	1101
<i>shoreae</i> Browne, <i>Genyocerus</i>	1095	<i>spinosus</i> Browne, <i>Platypus</i>	1184
<i>shoreanus</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i>	1181	<i>spinulosus</i> (Strohmeier), (<i>Stenoplatypus</i>), <i>Platypus</i>	1184
<i>shoreanus tersus</i> Schedl, <i>Platypus</i> (= <i>shoreanus</i>)	1181	<i>spinulosus montivagus</i> Roberts, <i>Platypus</i> (= <i>spinulosus</i>)	1184
<i>sicarius</i> Wood, <i>Platypus</i>	1181	<i>spretus</i> Schedl, <i>Platypus</i> (= <i>rethi</i>)	1192
<i>signatus</i> Chapuis, <i>Platypus</i>	1181	<i>springi</i> (Chapuis), (<i>Platypus</i>), <i>Neotrachyostus</i>	1214
<i>signatus</i> Strohmeier, <i>Periommatatus</i>	1106	<i>squameus</i> Schedl, <i>Platypus</i>	1184
<i>similis</i> (Nunberg), (<i>Stenoplatypus</i>), <i>Platypus</i>	1181	<i>squamifer</i> Schedl, <i>Platypus</i>	1185
<i>similis</i> Nunberg, <i>Trachyostus</i>	1212	<i>squamulatus</i> Chapuis, <i>Crossotarsus</i>	1206
<i>similis</i> Strohmeier, <i>Periommatatus</i>	1106	<i>squamulatus squamuloides</i> Beeson, <i>Crossotarsus</i> (= <i>squamulatus</i>)	1207
<i>simillimus</i> Nunberg, <i>Periommatatus</i> (= <i>angustior</i>)	1102	<i>squamulatus sundri</i> Beeson, <i>Crossotarsus</i> (= <i>squamulatus</i>)	1207
<i>simplex</i> Murayama, <i>Crossotarsus</i>	1205	<i>squamuloides</i> Beeson, <i>Crossotarsus</i> (= <i>squamulatus</i>)	1207
<i>simpliciformis</i> Wood, <i>Platypus</i>	1181	<i>staneri</i> Schedl, <i>Doliopygus</i>	1238
<i>simulans</i> Schedl, <i>Platypus</i>	1181	<i>Stenoplatypus</i> Strohmeier (= <i>Platypus</i>)	1113
<i>simulatus</i> Schedl, <i>Tesserocerus</i>	1100	<i>stenoplicatus</i> Schedl, <i>Platypus</i>	1185
<i>sindorae</i> Browne, <i>Platypus</i>	1182	<i>streuius</i> Schedl, <i>Platypus</i>	1185
<i>sinensis</i> Schedl, <i>Platypus</i>	1182	<i>striatodeclivis</i> Schedl, <i>Platypus</i>	1185
<i>singalangensis</i> Schedl, <i>Platypus</i>	1182	<i>striatopennis</i> Schedl, <i>Platypus</i>	1185
<i>singularis</i> Schedl, <i>Platypus</i>	1182	<i>striatopunctatus</i> Schedl, <i>Platypus</i>	1185
<i>sinuosus</i> Chapuis, <i>Platypus</i>	1182	<i>striatus</i> Browne, <i>Chaetastus</i>	1108
<i>siporanus</i> Schedl, <i>Crossotarsus</i>	1206	<i>striatus</i> Chapuis, <i>Platypus</i>	1185
<i>sirambeensis</i> Schedl, <i>Platypus</i>	1182	<i>striatus</i> Eggers, <i>Coptonotus</i>	1084

<i>strigillatus</i> Schedl. <i>Platypus</i>	1185	<i>sulcipemmis</i> Schedl. <i>Platypus</i>	1187
<i>strohmeieri</i> Wood, <i>Genyocerus</i>	1096	<i>sumatranus</i> Schedl. <i>Platypus</i> (= <i>bajulus</i>)	1119
<i>strombosiosis</i> Schedl, <i>Doliopygus</i>	1238	<i>sumatranus</i> Schedl, <i>Spathidicerus</i>	1102
<i>suavifer</i> Schedl. <i>Platypus</i>	1185	<i>sumbaicanus</i> Beeson, <i>Crossotarsus</i>	1207
<i>subabditus</i> (Schedl), (<i>Platypus</i>), <i>Platyscapulus</i> . .	1215	<i>sundaensis</i> Schedl, <i>Platypus</i> (= <i>pseudocupulatus</i>) . .	1172
<i>subaequalis</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>) . . .	1167	<i>sundri</i> Beeson, <i>Crossotarsus</i> (= <i>squamulatus</i>) . . .	1207
<i>subaequalispinosus</i> Schedl, <i>Platypus</i>	1185	<i>suspica</i> Schedl, <i>Mitosoma</i>	1112
<i>subaffinis</i> Schedl, <i>Cylindropalpus</i> (= <i>granulosus</i>) .	1218	<i>suturalis</i> Browne, <i>Periommatius</i>	1107
<i>subalpinus</i> (Schedl), (<i>Symmerus</i>), <i>Chactastus</i> . .	1108	<i>suzukii</i> Browne, <i>Platypus</i>	1187
<i>subangustior</i> Schedl, <i>Platypus</i>	1185	<i>Symmerariae</i> Strohmeier (= <i>Tesserocerini</i>)	1088
<i>subaplanatus</i> (Schedl), (<i>Platypus</i>), <i>Treptoplatypus</i>	1195	<i>Symmerus</i> Chapuis (= <i>Chactastus</i>)	1107
<i>subbeavifrons</i> Chapuis, <i>Platypus</i> (= <i>compositus</i>) .	1127	<i>taiheizanensis</i> (Murayama), (<i>Crossotarsus</i>), <i>Platypus</i>	1187
<i>subcircularis</i> Schedl, <i>Periommatius</i>	1107	<i>taicancensis</i> Schedl, <i>Platypus</i> (= <i>formosanus</i>)	1141
<i>subcostatus</i> Jacquelin-Duval, <i>Platypus</i> (= <i>parallelus</i>)	1165	<i>takeharai</i> Browne, <i>Platypus</i>	1187
<i>subcuratus</i> Browne, <i>Platypus</i>	1165	<i>talurae</i> (Stebbing), (<i>Diapus</i>), <i>Genyocerus</i>	1096
<i>subdepressus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> .	1185	<i>tasmanicus</i> Schedl, <i>Platypus</i>	1187
<i>subditicus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i> .	1239	<i>tavolae</i> Schedl, <i>Mitosoma</i> (= <i>truncatipennis</i>)	1112
<i>subdolosus</i> Schedl, <i>Doliopygus</i>	1239	<i>taxicornis</i> (Schedl), (<i>Platypus</i>), <i>Treptoplatypus</i> . .	1195
<i>subgranosus</i> Schedl, <i>Platypus</i>	1186	<i>tayabasi</i> Schedl, <i>Platypus</i>	1187
<i>subgranulosa</i> Browne, <i>Platypus</i>	1186	<i>tenax</i> (Schedl), (<i>Platyscapus</i>), <i>Cylindropalpus</i> . . .	1219
<i>subitarius</i> Schedl, <i>Platypus</i>	1186	<i>tenellus</i> (Schedl), (<i>Diacarus</i>), <i>Genyocerus</i>	1096
<i>sublunaris</i> (Schedl), (<i>Crossotarsus</i>), <i>Baiocis</i> . . .	1217	<i>tenellus</i> Schedl, <i>Platypus</i>	1188
<i>submarginatus</i> Schedl 1962, <i>Doliopygus</i>	1239	<i>tenuis</i> (Strohmeier), (<i>Crossotarsus</i>), <i>Doliopygus</i> .	1239
<i>submarginatus</i> Schedl 1971, <i>Doliopygus</i> (= <i>submarginatus</i> 1962)	1239	<i>tenuis</i> Murayama, <i>Platypus</i>	1188
<i>submontanus</i> Beeson, <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199	<i>tenuisculptus</i> Roberts, <i>Platypus</i>	1188
<i>subnitens</i> Browne, <i>Platypus</i>	1186	<i>tenuissimus</i> Schedl, <i>Platypus</i>	1188
<i>subnotatus</i> Schedl, <i>Doliopygus</i>	1239	<i>tepidus</i> Schedl, <i>Platypus</i>	1188
<i>suboblitaratus</i> Schedl, <i>Platypus</i>	1186	<i>terebrans</i> Schedl, <i>Doliopygus</i>	1239
<i>subopacus</i> Schedl, <i>Crossotarsus</i>	1207	<i>terebrans</i> Schedl, <i>Platypus</i>	1188
<i>subpellucidus</i> Lea, <i>Crossotarsus</i>	1207	<i>terminatulus</i> Schedl, <i>Platypus</i>	1188
<i>subpublicus</i> Schedl, <i>Platypus</i>	1186	<i>terminatus</i> Chapuis, <i>Crossotarsus</i>	1207
<i>subpraeteritus</i> Schedl, <i>Platypus</i>	1186	<i>terminatus sedulus</i> Schedl, <i>Crossotarsus</i> (= <i>terminatus</i>)	1208
<i>subpronus</i> Schedl, <i>Platypus</i>	1186	<i>ternus</i> Roberts, <i>Platypus</i>	1188
<i>subrobustus</i> Schedl, <i>Periommatius</i>	1107	<i>tersus</i> Schedl, <i>Platypus</i> (= <i>shoreanus</i>)	1181
<i>subsecretus</i> Browne, <i>Platypus</i>	1186	<i>Tesseroccephalus</i> Schedl (= <i>Tesserocerus</i>)	1097
<i>subserratus</i> Browne, <i>Doliopygus</i>	1239	<i>Tesserocerariae</i> Blandford (= <i>Tesserocerini</i>)	1088, 1097
<i>subsidiarius</i> Schedl, <i>Platypus</i>	1186	<i>Tesserocerinae</i> Blandford	1088
<i>subsimilis</i> Schedl, <i>Platypus</i>	1186	<i>Tesserocerini</i> Blandford	1088, 1097
<i>substriatus</i> (Schedl), (<i>Crossotarsus</i>), <i>Platypus</i> . .	1186	<i>Tesserocerus</i> Saunders	1097
<i>substriatus</i> Strohmeier, <i>Periommatius</i>	1107	<i>Tesserocranulus</i> Schedl	1101
<i>subsulcatus</i> Chapuis, <i>Platypus</i>	1187	<i>Tesseroplatypus</i> Schedl (= <i>Tesserocerus</i>)	1097
<i>subtruncatus</i> Browne, <i>Platypus</i>	1187	<i>testaceus</i> Dejean, <i>Platypus</i> (= <i>parallelus</i>)	1165
° <i>succiniceps</i> Schedl, <i>Cenocephalus</i>	1110	<i>tetracerus</i> Beeson, <i>Platypus</i>	1188
<i>suffodiens</i> Sampson, <i>Platypus</i>	1187	<i>tetricus</i> Schedl, <i>Platypus</i> (= <i>orientalis</i>)	1162
<i>suffodiens malayensis</i> Schedl, <i>Platypus</i> (= <i>vethi</i>) .	1192	<i>thomsoni</i> Chapuis, <i>Spathidicerus</i>	1102
<i>sulcaticeps</i> Schedl, <i>Platypus</i>	1187	<i>thoracicum</i> Schedl, <i>Mitosoma</i>	1111
<i>sulcatulus</i> Browne, <i>Cylindropalpus</i>	1219	<i>thoracicus</i> Chapuis, <i>Cenocephalus</i>	1110
<i>sulcatus</i> Chapuis, <i>Platypus</i> (= <i>mutatus</i>)	1185	<i>tiriosensis</i> Reichert, <i>Platypus</i>	1188
<i>sulcatus</i> Dejean, <i>Platypus</i> (= <i>cylindrus</i>)	1132	<i>titschaki</i> Schedl, <i>Periommatius</i>	1107
<i>sulcatus</i> Nunberg, <i>Periommatius</i> (= <i>excisus</i>)	1103	<i>togatus</i> (Schedl), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1239
<i>sulcinodis</i> Schedl, <i>Platypus</i>	1187	<i>tomentosus</i> (Strohmeier), (<i>Platypus</i>), <i>Trachyostus</i>	1212
<i>sulcipemmis</i> (Schedl), (<i>Coccecephalophonus</i>), <i>Mitosoma</i>	1112	<i>tonkinensis</i> Schedl, <i>Diapus</i>	1092
		<i>toralis</i> Schedl, <i>Crossotarsus</i>	1208

<i>tortulosus</i> Roberts & Morimoto, <i>Platypus</i>	1188	<i>tuberculatus robustus</i> Schedl. (<i>Synnumerus</i>), <i>Chaetastus</i> (= <i>tuberculatus</i>)	1108
<i>Trachyostus</i> Schedl.	1211	<i>tuberculosis</i> Strohmeier, <i>Platypus</i>	1190
<i>trachypennis</i> (Schedl.), (<i>Trachyostus</i>), <i>Crossotarsus</i>	1208	<i>tuberosus</i> (Schedl.), <i>Platypus</i>	1190
<i>tragus</i> Schedl., <i>Platypus</i>	1188	<i>tumidus</i> Schedl., <i>Crossotarsus</i>	1208
<i>transformis</i> Schedl., <i>Platypus</i>	1188	<i>turbatus</i> Chapuis, <i>Platypus</i> (= <i>geminatus</i>)	1142
<i>transiudicus</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i>	1189	<i>turcicus</i> Schedl., <i>Platypus</i>	1190
<i>transitus</i> Schedl., (<i>Platypus</i>), <i>Platyscapulus</i> (= <i>abditus</i>)	1214	<i>turgidus</i> Roberts, <i>Diapus</i>	1092
<i>transversecurinatus</i> Schedl., <i>Platypus</i>	1189	<i>turgifrons</i> (Schedl.), (<i>Platypus</i>), <i>Platyscapulus</i>	1215
<i>transversus</i> Schedl., <i>Platypus</i>	1189	<i>turriiformis</i> Roberts & Morimoto, <i>Platypus</i>	1190
<i>transversus</i> Schedl., <i>Trachyostus</i> (= <i>secundus</i>)	1211	<i>ugandae</i> Browne, <i>Platypus</i>	1190
<i>tremiferus</i> Chapuis, <i>Platypus</i> (= <i>compositus</i>)	1126	<i>ugandae</i> Schedl., <i>Doliopygus</i>	1240
<i>trepanatus</i> (Chapuis), (<i>Crossotarsus</i>), <i>Treptoplatypus</i>	1195	<i>ugandensis</i> Nunberg, (<i>Platypus</i>), <i>Trachyostus</i> (= <i>schaufussi</i>)	1212
<i>Treptoplatypus</i> Schedl.	1194	<i>ukereccensis</i> Schedl., (<i>Crossotarsus</i>), <i>Doliopygus</i> (= <i>rhodesianus</i>)	1237
<i>triangulatus</i> Browne, <i>Platypus</i>	1189	<i>uluguruensis</i> Nunberg, <i>Doliopygus</i> (= <i>bidentatus</i>)	1224
<i>tribuarius</i> Schedl., <i>Crossotarsus</i>	1208	<i>umbonatum</i> Schedl., <i>Mitosoma</i> (= <i>thoracicum</i>)	1112
<i>trichiliae</i> Schedl., <i>Doliopygus</i>	1239	<i>umbonatus</i> Blandford, <i>Platypus</i>	1190
<i>tricuspidatus</i> Schedl., <i>Platypus</i>	1189	<i>umbonatus</i> Browne, <i>Doliopygus</i> (= <i>bulbiventris</i>)	1226
<i>tricuspis</i> Schedl., (<i>Crossotarsus</i>), <i>Doliopygus</i> (= <i>coelocephalus</i>)	1227	<i>umbraticus</i> Schedl., <i>Platypus</i>	1190
<i>trideus</i> Roberts, <i>Doliopygus</i>	1240	<i>umbrinus</i> Roberts, <i>Platypus</i>	1190
<i>trigonus</i> Beeson, <i>Crossotarsus</i>	1208	<i>umbrosus</i> (Schedl.), (<i>Platypus</i>), <i>Platyscapulus</i>	1216
<i>trilobipennis</i> Browne, <i>Platypus</i>	1189	<i>uncacanthurus</i> Beeson, <i>Platypus</i> (= <i>lewisi</i>)	1153
<i>Triozaustus</i> Schedl.	1219	<i>uncatus</i> Browne, <i>Platypus</i>	1190
<i>triplurus</i> Browne, <i>Platypus</i>	1189	<i>uncinatus</i> Blandford, <i>Platypus</i>	1190
<i>triquetrus</i> Brethes, <i>Platypus</i> (= <i>parallelus</i>)	1168	<i>uncinatus</i> Schedl., <i>Doliopygus</i>	1240
<i>triquetrus</i> Schedl., <i>Periommatas</i>	1107	<i>unicornis</i> Schedl., <i>Doliopygus</i>	1240
<i>trispinatus</i> Schedl., <i>Platypus</i>	1189	<i>unicus</i> (Schedl.), (<i>Crossotarsus</i>), <i>Doliopygus</i>	1240
<i>trispinatus</i> Schedl., (<i>Diacarus</i>), <i>Genyocerus</i> (= <i>puer puer</i>)	1095	<i>uniformis</i> Schedl., <i>Platypus</i>	1191
<i>trispinatus</i> Browne, <i>Platypus</i> (= <i>bispinatus</i>)	1121	<i>unimodus</i> Schedl., <i>Platypus</i> (= <i>pallidus</i>)	1164
<i>trispinatus</i> Schedl., <i>Platypus</i>	1189	<i>unispineus</i> Roberts, <i>Diapus</i>	1093
<i>trispinosus</i> Chapuis, <i>Platypus</i>	1189	<i>unispinosus</i> Roberts, <i>Baiois</i>	1217
<i>tristis</i> Roberts, <i>Platypus</i>	1189	<i>unispinosus</i> Schedl. 1936, (<i>Crossotarsus</i>), <i>Doliopygus</i> (= <i>mimicus</i>)	1233
<i>truncatellus</i> Browne, <i>Platypus</i>	1189	<i>unispinosus</i> Schedl. 1937, <i>Doliopygus</i> (= <i>mimicus</i>)	1233
<i>truncatellus</i> Schedl., <i>Periommatas</i>	1107	<i>ursinus</i> Schedl., <i>Platypus</i>	1191
<i>truncatocauda</i> Schedl., <i>Platypus</i>	1189	<i>ursulus</i> Schedl., <i>Crossotarsus</i> (= <i>ursus</i>)	1208
<i>truncatiformis</i> Schedl., <i>Periommatas</i>	1107	<i>ursus</i> Schedl., (<i>Tesseroplatypus</i>), <i>Tesserocerus</i> (= <i>insignis</i>)	1100
<i>truncatigranulosus</i> Schedl., <i>Platypus</i>	1189	<i>ursus</i> Schedl., <i>Crossotarsus</i>	1208
<i>truncatipennis</i> Schedl., <i>Mitosoma</i>	1112	<i>ursus</i> Schedl., <i>Platypus</i>	1191
<i>truncatipennis</i> Schedl., <i>Periommatas</i>	1107	<i>usambariensis</i> Strohmeier, <i>Crossotarsus</i> (= <i>externedentatus</i>)	1199
<i>truncatipennis</i> Schedl., <i>Platypus</i>	1189	<i>ustulatus</i> Chapuis, <i>Platypus</i>	1191
<i>truncatipennis tavolae</i> Schedl., <i>Mitosoma</i> (= <i>truncatipennis</i>)	1112	<i>ustulatus</i> Shuckard, <i>Tesserocerus</i>	1101
<i>truncatulus</i> Browne, <i>Platypus</i>	1190	<i>ustus</i> Schedl., <i>Mesoplatypus</i>	1222
<i>truncatum</i> Schedl., <i>Mitosoma</i>	1112	<i>ustus</i> Schedl., <i>Platypus</i>	1191
<i>truncaturus</i> Browne, <i>Platypus</i>	1190	<i>uter</i> Schedl., <i>Platypus</i>	1191
<i>truncatus</i> Chapuis, <i>Platypus</i>	1190	<i>utibilis</i> (Schedl.), (<i>Crossotarsus</i>), <i>Platypus</i>	1191
<i>truncatus</i> Niisima & Murayama, <i>Diapus</i>	1092	<i>urarius</i> Roberts, <i>Platypus</i>	1191
<i>truncatus</i> Schedl., <i>Periommatas</i>	1107	<i>vafer</i> Schedl., <i>Crossotarsus</i>	1208
<i>tuberculatus</i> (Chapuis), (<i>Synnumerus</i>), <i>Chaetastus</i>	1108	<i>vafer</i> Schedl., <i>Platypus</i>	1192
<i>tuberculatus</i> Browne, <i>Platytarsulus</i>	1097	<i>vaginensis</i> Roberts, <i>Platypus</i>	1192
<i>tuberculatus</i> Chapuis, <i>Platypus</i>	1190	<i>vagus</i> Browne, <i>Platypus</i>	1192
		<i>valdephlebius</i> Roberts, <i>Platypus</i>	1192
		<i>validus</i> Schedl., <i>Platypus</i>	1192

<i>variolosus</i> (Schedl), (<i>Platypus</i>), <i>Baiocis</i>	1217	<i>vicinus</i> Blandford, <i>Platypus</i>	1193
<i>varipennis</i> Schedl, <i>Platypus</i>	1192	<i>vicinus</i> Schedl, <i>Crossotarsus</i>	1209
<i>varius</i> Sampson, (<i>Platypus</i>), <i>Doliopygus</i> (= <i>erichsoni</i>)	1229	<i>viduus</i> Chapuis, <i>Platypus</i> (= <i>seguis</i>)	1179
<i>vastus</i> Strohmeyer, (<i>Platypus</i>), <i>Trachyostus</i> (= <i>aterrimus</i>)	1211	<i>vilis</i> Schedl, <i>Doliopygus</i>	1240
<i>vagestus</i> Wood, <i>Platypus</i>	1192	<i>virgatus</i> Schedl, <i>Platypus</i> (= <i>bellus</i>)	1119
<i>vegrandis</i> Sampson, (<i>Crossotarsus</i>), <i>Doliopygus</i> (= <i>conradti</i>)	1227	<i>vitiensis</i> Roberts, <i>Platypus</i>	1193
<i>vegrandis</i> Schedl, (<i>Tesseroplatypus</i>), <i>Tesserocerus</i> (= <i>spinolae</i>)	1101	<i>vogelii</i> Roberts, <i>Platypus</i> (= <i>orientalis</i>)	1163
<i>velatus</i> Schedl, (<i>Platypus</i>), <i>Baiocis</i> (= <i>inimicus</i>) . . .	1216	<i>rolaticus</i> Schedl, <i>Platypus</i>	1193
<i>ventricornis</i> Schedl, <i>Crossotarsus</i>	1208	<i>ronfaberi</i> Reichardt, <i>Platypus</i>	1193
<i>ventricornis</i> Schedl, <i>Mesoplatypus</i>	1222	<i>vulpinum</i> Schedl, <i>Mitosoma</i>	1112
<i>ventrispinis</i> Schedl, <i>Crossotarsus</i>	1208	walkeri Schedl, <i>Platypus</i> (= <i>solidus</i>)	1183
<i>ventrosus</i> Schedl, <i>Crossotarsus</i>	1209	<i>wallacei</i> (Thomson), (<i>Platypus</i>), <i>Crossotarsus</i> . . .	1209
<i>venustulus</i> Beeson, <i>Crossotarsus</i> (= <i>terminatus</i>) . .	1207	<i>watbrui</i> Roberts, <i>Platypus</i>	1193
<i>venustus</i> Chapuis, <i>Crossotarsus</i> (= <i>terminatus</i>) . .	1207	<i>watutensis</i> Roberts, <i>Platypus</i>	1193
<i>venustus</i> Schedl, <i>Mesoplatypus</i>	1223	<i>webberi</i> Schedl, <i>Platypus</i>	1193
<i>venustus venustulus</i> Beeson, <i>Crossotarsus</i> (= <i>terminatus</i>)	1207	<i>wesmaeli</i> Chapuis, <i>Platypus</i> (= <i>parallelus</i>)	1167
<i>verehunatus</i> (Beeson), (<i>Crossotarsus</i>), <i>Platypus</i> . .	1192	<i>westwoodi</i> Chapuis, <i>Platypus</i>	1193
<i>vernaculus</i> Schedl, <i>Platypus</i>	1192	<i>wilmoiti</i> (Stebbing), (<i>Platypus</i>), <i>Crossotarsus</i> . . .	1209
<i>resculus</i> Schedl, <i>Platypus</i>	1192	<i>wilsoni</i> Swaine, <i>Platypus</i>	1194
<i>rethi</i> Strohmeyer, <i>Platypus</i>	1192	<i>wittei</i> Schedl, <i>Doliopygus</i>	1240
<i>retulus</i> Schedl, <i>Platypus</i>	1192	<i>wollastoni</i> (Chapuis), (<i>Crossotarsus</i>), <i>Carchesiopygus</i>	1210
<i>retulus</i> Wood, <i>Protoplatypus</i>	1086	<i>woltschei</i> Schedl, <i>Doliopygus</i>	1240
<i>retustus</i> Schedl, <i>Platypus</i>	1193	<i>xanthopus</i> Schedl, <i>Platypus</i>	1194
<i>vexans</i> Schedl, <i>Platypus</i>	1193	<i>xylographus</i> Schedl, <i>Platypus</i>	1194
<i>vexator</i> Schedl, <i>Doliopygus</i> (= <i>megatoma</i>)	1233	<i>xylopiac</i> Roberts, <i>Platypus</i> (= <i>impressus</i>)	1145
<i>viaticus</i> Schedl, <i>Platypus</i>	1193	<i>yasiyasi</i> Roberts, <i>Platypus</i>	1194
<i>vicarius</i> Schedl, <i>Platypus</i>	1193	<i>zeteki</i> Blackman, <i>Mecopelmus</i>	1086
		<i>zeylanicus</i> Beeson, <i>Genyocerus</i> (= <i>albipennis</i>)	1093
		<i>zonatus</i> Schedl, <i>Doliopygus</i>	1240

Family Scolytidae Index

This index includes all Latin names used in this catalog for Scolytidae. Family-group names (family, subfamily, tribe) and names applied below the rank of subspecies (including aberrations, variations, nomen nudums) are given in regular type. Valid names of genera and species are in bold type. The names of synonyms of genera and species, subgeneric names, and subspecific names are given in italics. The names are listed in alphabetical order by the computer; however, the user must realize that the computer reads parentheses () as a letter of the alphabet preceding the letter "a." The names of fossil species are preceded by an asterisk (*). Fossil species are given at the beginning of the index and are also listed alphabetically with other names.

- * *americanus* Wickham, *Hylastes* 43
- * *anceps* Hagedorn, *Xylechinites* 1082
- * *arcessitus* (Scudder), (*Phlocosinites*),
 Phlocosinus 240
- * *assimilis* (Schedl), (*Phlocosinites*), *Phlocosinus* . . . 241
- * *aterites* Schedl, *Hylastes* 49
- * *brunni* (Hagedorn), (*Phlocosinites*), *Phlocosinus* . . 243
- * *carbonarius* Scudder, *Dryocoetes* 580
- * *corpulentus* Schedl, *Hylurgops* 30
- * *diluvialis* Wickham, *Dryocoetes* 582
- * *dromiscens* Scudder, *Hylesinus* 79
- * *dubius* (Hagedorn), (*Myelophilites*), *Hylurgops* . . 31
- * *electricus* (Germar), (*Hylesinites*), *Hylurgops* . . . 31
- * *emmi* (Hagedorn), (*Cryphalus*), *Hypothenemus* . . . 918
- * *excavatus* (Hagedorn), (*Xyleborus*), *Xyleborinus* . . 807
- * *extractus* Scudder, *Hylesinus* 80
- * *facilis* Heer, *Hylesinus* 80
- * *gnathotrichus* Schedl, *Taphramites* 1082
- * *granulatus* (Schedl), (*Hylescerites*), *Hylurgops* . . . 33
- * *hydropicus* (Wickham), (*Apidocephalus*),
 Hylesinus 80
- * *immaturus* Schedl, *Taphrorychus* 559
- * *impressum* Scudder, *Trypodendron* 635
- * *keilbachi* Schedl, *Carphoborites* 1082
- * *lineatus* Foster, *Hylesinus* 81
- * *longipennis* Wickham, *Xyleborites* 1082
- * *piger* Wickham, *Hylurgops* 38
- * *pilosellus* Schedl, *Hylurgops* 39
- * *posticus* Schedl, *Carphoborites* 1082
- * *priscus* Eggers, *Xyleborus* 765
- * *pseudocrucifer* (Schedl), (*Xyleborus*),
 Cyclorhipidion 702
- * *regimontanus* (Hagedorn), (*Phlocosinites*),
 Phlocosinus 252
- * *rehi* (Hagedorn), (*Phlocosinites*), *Phlocosinus* . . . 252
- * *robustus* (Schedl), (*Phlocosinites*), *Phlocosinus* . . . 252
- * *rugosissimus* Cockerell, *Cryphalites* 1082
- * *schellwieni* (Hagedorn), (*Hylastites*), *Hylurgops* . . 41
- * *sexspinosus* (Schedl), (*Phlocosinites*),
 Phlocosinus 255
- * *squalidens* (Scudder), (*Hylastes*), *Phlocosinus* . . . 255
- * *tiburtinus* Claus, *Scolytus* (= *intricatus*) 334
- * *tuberculatus* Schedl, *Hylurgops* (= *tuberculifer*) . . . 43
- * *tuberculifer* (Schedl), (*Phlocosinites*),
 Phlocosinus 258
- * *tuberculifer* Wood, *Hylurgops* 43
- * *wolffi* (Schedl), (*Phlocosinites*), *Phlocosinus* 259
- * *worthini* Scudder, (*Polygraphus*), not Scolytidae . . 284
- 12-spinatus* Schedl, *Webbia* (= *duodecimspinatus*) . . . 831
- 14-spinatus* Sampson, *Webbia*
 (= *quattuordecimspinatus*) 832
- 26-spinatus* Sampson, *Webbia* (= *trigintispinatus*) . . . 833
- 30-spinatus* Sampson, *Webbia* (= *trigintispinatus*) . . . 833
- abachidsei* Stark, *Carphoborus* (= *perrisi*) 308
- abaensis* Tsai & Yin, *Scolytus* 322
- abberrans* Schedl, *Xyleborus* 705
- abbreviata* (Schedl), (*Xyleborus*), *Coptodryas* 822
- abbreviatipennis* Schedl, *Xyleborus* 705
- abbreviatus* (Schedl), (*Cryptocleptes*),
 Pseudothysanoes 411
- abbreviatus* Eggers, *Crypturgus* (= *numidicus*) 625
- abbreviatus* Eichhoff, *Corthylus* 1070
- abbreviatus* Schedl, *Cryphalus* 873
- abbreviatus* Schedl, *Pityophthorus* 978
- abbreviatus* Schedl, *Pseudochramesus* 261
- abdominalis* Hopkins, *Hypothenemus* (= *columbi*) . . . 913
- abeillei* Guillebeau, (*Phlocophthorus*),
 Phlocotribus (= *perfoliatus*) 226
- aberrans* Browne, *Hypothenemus* 905
- aberrans* Schedl, *Chramesus* 263
- aberrans* Schedl, *Monarthrum* 1051
- aberrans* Wichmann, *Microborus* 383
- aberrans* Wood, *Hyllocurus* 422
- aberrans* Wood, *Spermophthorus* 971
- abhorrens* Eggers, *Pernophthorus* 202
- abhorrens* Wichmann, (*Eccoptogaster*), *Scolytus*
 (= *multistriatus*) 353
- abiegnus* Wood, *Pityophthorus* 978
- abietinus* Fabricius, (*Bostrichus*), *Tomicus*
 (= *piniperda*) 136
- abietinus* Kono & Tamanuki, *Dryocoetes*
 (= *striatus*) 585
- abietinus* Wood, *Pityophthorus* 978
- abietiperda* Bechstein, (*Bostrichus*), *Hylurgops*
 (= *palliatus*) 38
- abictis* Blackman, *Pityophthorus* (= *opaculus*) 1018
- abictis* Blackman, *Scolytus* (= *opacus*) 355
- abictis* Hopkins, *Dryocoetes* (= *confusus*) 581
- abictis* Kurenzov, *Pityophthorus* (= *abietinus*) 978
- abictis* Kurenzov, *Polygraphus* 284

- abietis* Ratzeburg, (*Bostrichus*), *Cryphalus*
(=*asperatus*) 874
- abietis* Tsai & Yin, *Phloeosinus* 240
- abitusus* Bright, *Pityophthorus* 978
- abnormalis* Bright, *Pityophthorus* 978
- abnormis* Eichhoff, (*Xyleborus*), *Eccoptopterus*
(=*spinus*) 821
- abruptedeclevis* Schedl, *Corthylus* 1070
- abruptoides* (Schedl), (*Xyleborus*), *Xylosandrus* 787
- abruptulus* (Schedl), (*Pocillips*), *Coccotrypes* 591
- abruptulus* (Schedl), (*Xyleborus*), *Xylosandrus* 787
- abruptum* (Nunberg), (*Microcorthylus*),
Monarthrum 1051, 1065
- abruptus* (Schedl), (*Pocillips*), *Coccotrypes* 591
- abruptus* (Schedl), (*Stephanoderes*),
Hypothenemus 905
- abruptus* Eggers, (*Xyleborus*), *Amasa* (= *resectus*) 684
- abruptus* Nunberg, *Microcorthylus* 1051, 1065
- abruptus* Nunberg, *Xylocleptes* 547
- abruptus* Sampson, (*Xyleborus*), *Xylosandrus*
(=*mancus*) 797
- abruptus* Schedl, *Arixyleborus* 665
- abruptus* Schedl, *Tricolus* 1042
- abscissus* Browne, *Xyleborus* 705
- absonus* (Schedl), (*Cryphalomorphus*),
Scolytogenes 858
- absonus* Blackman, *Pityophthorus* 978
- abstrusus* Bright, *Pityophthorus* 979
- abundans* Lea, *Acacasis* 193
- abyssinicus* (Schedl), (*Hoplitantus*), *Xylocleptes* 547
- Acacasis* Lea 193
- acaciae* (Blackman), (*Cryptocleptes*),
Pseudothyasanoes 412
- acaciae* (Eggers), (*Cryphalus*), *Hypothenemus* 906
- acaciae* (Lea), (*Phloeophthorus*), *Phloeotribus* 218
- acaciae* (Schedl), (*Apoglostatus*), *Glostatus* 835
- acaciae* (Schedl), (*Haplogenus*), *Rhopalopselion* 96
- acaciae* Schedl, *Cryptocarenum* 902
- acacicolens* Wood, *Chramesus* 263
- acacicolens* Wood, *Hylesinopsis* 92
- acacicolens* Wood, *Pseudothyasanoes* 412
- acanthodes* Schedl, *Xyleborus* (= *multispinatus*) 754
- acanthopanxi* (Niisima), (*Cryphalus*), *Erioporus* 854
- Acanthophorus* Strohmeyer (= *Pernophorus*) 202
- Acanthotomicus* Blandford 478
- acanthurus* (Lea), (*Tomicus*), *Xyleborus* 705
- Acanthurus* Eichhoff (= *Diamerus*) 194
- acanthus* Schedl, *Xyleborus* (= *multispinatus*) 754
- acanthus acanthodes* Schedl, *Xyleborus*
(=*multispinatus*) 754
- acanthus acuticornis* Schedl, *Xyleborus*
(=*multispinatus*) 754
- acanthus minus* Schedl, *Xyleborus*
(=*multispinatus*) 754
- acares* (Wood), (*Cryptocleptes*),
Pseudothyasanoes 412
- acares* Wood, *Scolytodes* 358
- acatayi* Schedl, *Phloeosinus* 240
- acceptus* Bright, *Pityophthorus* (= *murrayanae*) 1014
- accinctus* Wood, *Araptus* 953
- accomodatus* Schedl, *Xyleborus* (= *adelographus*) 706
- acerini* Blackman, *Pityophthorus* (= *lautus*) 1006
- aceris* (Lindemann), (*Dryocoetes*), *Lymantor* 554
- aceris* (Niisima), (*Xyloterus*), *Indocryphalus* 648
- aceris* Hopkins, *Procryphalus* (= *utahensis*) 848
- aceris* Knotek, *Scolytus* (= *koenigi*) 338
- aceris* Krivolutskaia, *Dryocoetes* 570
- aceris* Kurenzov, *Xyleborus* (= *orientalis*) 757
- achillei* Reitter, *Hylastinus* 64
- aciculatus* Blackman, *Gnathotrichus* (= *sulcatus*) 1039
- aciculatus* Bright, *Pityophthorus* 979
- aciculatus* Bruck, *Phloeosinus* (= *serratus*) 255
- aciculatus* Schedl, *Coccotrypes* 591
- aciculatus* Schedl, *Hypocryphalus* 868
- aciculatus* Wood, *Tricolus* 1042
- acilis* Wood, *Xyleborus* (= *improvidus*) 744
- Acorthylus* Brethes 852
- Acrantus* Broun (= *Chactoptelius*) 103
- Acryphalus* Tsai & Li (= *Cryphalus*) 872
- aculeatus* LeConte, *Micracis* (= *suturalis*) 433
- aculeatus* Say, *Hylesinus* 74
- acuminatum* Schedl, *Ctonoxylon* 836
- acuminatus* (Gyllenhal), (*Bostrichus*), *Ips* 485
- acuminatus* (Schedl), (*Eidophelus*), *Cyrtogenius* 562
- acuminatus* (Schedl), (*Minips*), *Acanthotomicus* 478
- acuminatus* (Schedl), (*Neopityophthorus*),
Pityophthorus 979
- acuminatus* Bright, *Pseudopityophthorus*
(=*singularis*) 976
- acuminatus* Browne, *Pityophthorus* (= *hintzi*) 1002
- acuminatus* Schedl, (*Pocillips*), *Coccotrypes*
(=*salakensis*) 611
- acuminatus* Schedl, *Cnemidophorus* 313
- acuminatus* Schedl, *Cnecinus* 205
- acuminatus* Schedl, *Hylocurus* (= *dimorphus*) 424
- acuminatus* Schedl, *Miniocurus* 949
- acuminatus* Schedl, *Scolytoplatypus*
(=*eichelbaumi*) 402
- acuminatus* Schedl, *Xyleborus* (= *grossmanni*) 741
- acuminatus* Wood, *Scolytodes* (= *cecropiavorus*) 390
- acus* Wood, *Amphicranus* 1045
- acuteclavatus* (Hagedorn), (*Chramesus*),
Pseudochramesus 262
- acutedentatus* Schedl, *Hylocurus* 422
- acuticornis* Schedl, *Xyleborus* (= *multispinatus*) 754
- acutipennis* Eichhoff, *Micracis* 431
- acutus* (Schedl), (*Xyleborus*), *Webbia* 829
- acutus* Blackman, *Pityophthorus* 979
- acutus* Wood, *Dendrocraudus* 550
- adamsoni* Beeson, *Xyleborus* 705
- addendus* (Schedl), (*Xyleborus*), *Ambrosiodmus* 670
- additus* Wood, *Corthylus* 1070
- adelographus* Eichhoff, *Xyleborus* 706
- adeniae* (Schedl), (*Dryocoetes*), *Xylocleptes* 547
- adherescens* Schedl, *Xylosandrus* 787

<i>Adiacretus</i> Hagedorn (= <i>Hypothenemus</i>)	905	<i>affinis</i> Blandford. <i>Dryocoetes</i>	571
<i>adjunctum</i> (Schedl). (<i>Pterocyclon</i>), <i>Monarthrum</i>	1051	<i>affinis</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>multistriatus</i>)	353
<i>adjunctus</i> (Eggers). (<i>Xyleborus</i>), <i>Coptoborus</i>	662	<i>affinis</i> Eggers. <i>Bothrostermus</i>	215
<i>adjunctus</i> Blandford. <i>Deudroctonus</i>	137	<i>affinis</i> Eggers. <i>Tricholus</i>	1042
<i>adjunctus corruptus</i> Schedl. (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>adjunctus</i>)	662	<i>affinis</i> Eichhoff. <i>Xyleborus</i>	706
<i>adnata</i> Wood. <i>Micracisella</i>	429	<i>affinis</i> Fonseca. (<i>Metacorthylus</i>), <i>Corthylus</i> (= <i>papulans</i>)	1076
<i>adonis</i> Wood. <i>Thysanoes</i>	419	<i>affinis</i> Fonseca. <i>Corthylus</i> (= <i>papulans</i>)	1076
<i>adossuarius</i> Schedl. <i>Xyleborus</i>	706	<i>affinis</i> Schedl. <i>Cyrtogenius</i> (= <i>subgranosus</i>)	569
<i>Adryocoetes</i> Eggers (= <i>Coptodryas</i>)	822	<i>affinis</i> Stephens. (<i>Tomicus</i>), <i>Polygraphus</i> (= <i>poligraphus</i>)	292
<i>adscitus</i> (Schedl). (<i>Stephanoderes</i>), <i>Hypothenemus</i>	906	<i>affinis</i> Strohmeier. <i>Cladoctonus</i>	235
<i>adspersus</i> Schedl. <i>Xyleborus</i> (= <i>ruficanus</i>)	783	<i>affinis</i> fuscobrunneus Eichhoff. <i>Xyleborus</i> (= <i>affinis</i>)	705
<i>adunbratus</i> Blandford. <i>Xyleborus</i> (= <i>pfeili</i>)	763	<i>affinis</i> mascarensis Eichhoff. <i>Xyleborus</i> (= <i>affinis</i>)	705
<i>aduncus</i> Schedl. (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>aduncus</i>)	804	<i>affinis</i> parvus Eichhoff. <i>Xyleborus</i> (= <i>affinis</i>)	705
<i>aduncus</i> (Schedl). (<i>Xyleborus</i>), <i>Xyleborinus</i>	804	<i>afiamalus</i> Schedl. <i>Hypocryphalus</i>	565
<i>aduncus profundus</i> Schedl. (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>aduncus</i>)	804	<i>africanulus</i> Wood. <i>Pityophthorus</i>	979
<i>adusticollis</i> (Motschulsky). (<i>Tomicus</i>), <i>Xyleborus</i>	706	<i>africanus</i> (Browne). (<i>Orthotomicus</i>), <i>Acanthotomicus</i>	478
<i>adusticus</i> Wood. <i>Cnesinus</i>	205	<i>africanus</i> (Eggers). (<i>Pseudoploceotribus</i>), <i>Hylesinopsis</i>	92
<i>adustum</i> Wood. <i>Monarthrum</i> (= <i>pseudoscutellare</i>)	1061	<i>africanus</i> (Eggers). (<i>Toxophorus</i>), <i>Toxophthorus</i>	1082
<i>adustus</i> (Eggers). (<i>Hylocurosoma</i>), <i>Scolytodes</i>	385	<i>africanus</i> (Hopkins). (<i>Stephanoderes</i>), <i>Hypothenemus</i>	906
<i>adustus</i> (Eggers). (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	670	<i>africanus</i> (Numberg). (<i>Carphoborus</i>), <i>Cardroctonus</i>	283
<i>adustus</i> Eggers. <i>Cryptocarenum</i> (= <i>seriatus</i>)	904	<i>africanus</i> (Schedl). (<i>Hylocurus</i>), <i>Phloeocorus</i>	429
<i>adustus</i> Schedl. <i>Cnesinus</i>	205	<i>africanus</i> (Schreiner). (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	563
<i>advena</i> Blandford. <i>Coccotrypes</i>	592	<i>africanus</i> Browne. <i>Xylechinus</i>	115
<i>advena</i> Schedl. <i>Chramesus</i>	263	<i>africanus</i> Eggers. <i>Hypocryphalus</i> (= <i>mangiferae</i>)	869
<i>advena</i> Schedl. <i>Cnesinus</i>	205	<i>africanus</i> Eggers. (<i>Metahylastes</i>), <i>Cyrtogenius</i> (= <i>africus</i>)	563
<i>aegir</i> (Eggers). (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	670	<i>africanus</i> Eggers. <i>Dactylipalpus</i>	101
<i>aegyptiacus</i> Pic. <i>Scolytus</i> (= <i>amygdali</i>)	323	<i>africanus</i> Eggers. <i>Pityophthorus</i>	979
<i>aemulus</i> (Wollaston). (<i>Tomicus</i>), <i>Xyleborinus</i>	804	<i>africanus</i> Eggers. <i>Scolytotypus</i>	401
<i>aeneipennis</i> (Fabricius). (<i>Hylesinus</i>), <i>Camptocerus</i>	315	<i>africanus</i> Eggers. <i>Thamurgus</i>	542
<i>aeonii</i> Lindberg. <i>Aphanarthrum</i>	616	<i>africanus</i> Eggers. <i>Xyleborus</i>	709
<i>aequaliclavatus</i> Schedl. <i>Hypothenemus</i> (= <i>fuscolis</i>)	926	<i>africanus</i> Schedl. (<i>Aluiphagus</i>), <i>Hylesinopsis</i> (= <i>acacicolens</i>)	92
<i>aequalis</i> (Reitter). (<i>Anisandrus</i>), <i>Xyleborus</i>	706	<i>africanus</i> Schedl. (<i>Hylesinus</i>), <i>Hylesinopsis</i> (= <i>secutus</i>)	95
<i>aequalis</i> (Schedl). (<i>Ozodendron</i>), <i>Cyrtogenius</i>	562	<i>africanus</i> Schedl. (<i>Necodryocoetes</i>), <i>Pityophthorus</i> (= <i>africanulus</i>)	979
<i>aequalis</i> Schedl. <i>Polygraphus</i>	284	<i>africanus</i> Schedl. (<i>Stephanorhopalus</i>), <i>Eruoporicus</i>	549
<i>aequalis solidus</i> Schedl. <i>Polygraphus</i> (= <i>aequalis</i>)	284	<i>africanus</i> picinus Schedl. <i>Xyleborus</i> (= <i>africanus</i>)	709
<i>aequatorensis</i> Eggers. <i>Xyleborus</i>	706	<i>africus</i> Wood. <i>Cyrtogenius</i>	563
<i>aequatus</i> Schedl. <i>Polygraphus</i>	284	<i>Afrochramesus</i> Schedl. (= <i>Chortastus</i>)	311
<i>aequipunctatus</i> Eggers. <i>Scolytodes</i>	385	<i>Afromicracis</i> Schedl. (= <i>Microcryphalus</i>)	410
<i>aequipunctatus</i> Nüssima. <i>Scolytus</i> (= <i>aratus</i>)	324	<i>Afrotryptus</i> Bright (= <i>Styracoptinus</i>)	251
<i>aesculi</i> Ferrari. (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxeseni</i>)	814	<i>afzeliae</i> Schedl. <i>Polygraphus</i>	254
<i>aesculinus</i> Bright. <i>Pseudopityophthorus</i> (= <i>opacicollis</i>)	974	<i>agamus</i> Perkins. <i>Xyleborus</i>	709
<i>aethiopicus</i> (Eggers). (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	563	<i>agathis</i> (Browne). (<i>Ozodendron</i>), <i>Cyrtogenius</i>	563
<i>aethiopicus</i> Eggers. <i>Triotemus</i>	540	<i>agathis</i> Browne. <i>Xyleborus</i>	709
<i>aethiops</i> (Schedl). (<i>Lepiceroidea</i>), <i>Hypothenemus</i>	906	<i>aglaiae</i> Browne. <i>Xyleborus</i>	709
<i>aethiops</i> Schedl. <i>Xyloctonus</i>	840	<i>agnaticeps</i> (Schedl). (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	697
<i>afer</i> (Schedl). (<i>Cryphalophilus</i>), <i>Scolytogenes</i>	855		
<i>affaber</i> (Mannerheim). (<i>Bostrichus</i>), <i>Dryocoetes</i>	570		
<i>affine</i> Wollaston. <i>Aphanarthrum</i>	616		
<i>affinis</i> (Schedl). (<i>Pelicerus</i>), <i>Cyrtogenius</i>	563		

- agnatum* (Eggers), (*Xyleborus*), *Cyclorhipidion* . . . 697
agnatus (Eggers), (*Stephanoderes*), *Hypothenemus* . . . 906
agnatus Blackman, *Pityophthorus* (= *confertus*) . . . 989
agnatus Blandford, *Scolytus* (= *dahuricus*) . . . 328
agnatus Eggers, *Chortastus* (= *minimus*) . . . 311
agnatus Schedl, *Miocryphalus* . . . 410
agraphus Schedl, *Xyleborus* . . . 709
agrifoliae Blackman, *Pseudopityophthorus* . . . 972
aguacatensis (Schedl), (*Metacortylus*),
Cortyllocurus . . . 1068
aitutakii Beaver & Maddison, *Ptilopodius* . . . 899
aizawai Kono, *Pityogenes* (= *seirindensis*) . . . 458
alaskanus Blackman, *Phloeosinus* (= *pinii*) . . . 250
alaskanus Wood, *Lymantria* . . . 555
albanicus Apfelbeck, *Pityogenes* (= *bistridentatus*) . . . 444
albertensis Blackman, *Pityophthorus* (= *opaculus*) . . . 1018
albidum Wichmann, (*Liparthrum*) (= *genistae*) . . . 277
albipilis Reitter, (*Stephanoderes*), *Hypothenemus*
 (= *leprieuri*) . . . 933
albizzianus (Schedl), (*Xyleborus*), *Ambrosiodmus* . . . 670
albosetum Bright, *Liparthrum* . . . 275
algidus Schedl, *Xyleborus* . . . 709
alienus (Schedl), (*Xyleborus*), *Xyleborinus* . . . 804
alienus Eichhoff, *Hylocurus* . . . 422
alienus Eichhoff, *Pityophthorus* . . . 979
alienus Schedl, (*Cryphalomorphus*), *Scolytogenes*
 (= *knabi*) . . . 563
alienus Schedl, *Cortylus* . . . 1070
alienus Schedl, *Glochiphorus* . . . 271
alienus Wood, *Cnesinus* . . . 205
Allarthrum Hagedorn (= *Cryphalus*) . . . 872
allecta (Schedl), (*Xyleborus*), *Schedlia* . . . 822
Allernoporus Kurenzov . . . 852
alluadi (Lepesme), (*Kissophagus*), *Hylesinopsis* . . . 93
alluadi (Peyerimhoff), (*Aphanarthrum*),
Coleobothrus . . . 615
alluadi Schaufuss, *Xyleborus* . . . 709
alni (Bright), (*Chaetophloeus*), *Carphotoreus* . . . 237
alni (Georg), (*Bostrichus*), *Dryocoetes* . . . 571
alni (Lindemann), (*Cryphalus*), *Trypophloeus* . . . 842
alni (Niisima), (*Hylastes*), *Alniphagus* . . . 73
alni (Niisima), (*Xyleborus*), *Xyleborinus* . . . 804
alni Blackman, *Gnathotrichus* (= *retusus*) . . . 1038
alni Blackman, *Pityophthorus* . . . 979
alni Krivolutskaya, *Cryphalus* . . . 873
alni Mulsant & Rey, (*Bostrichus*), *Xyleborus*
 (= *pfeili*) . . . 763
alni Pfeffer, *Taphronychus* . . . 557
alni Wood, *Gymnochilus* . . . 386
alni Wood, *Scolytodes* . . . 385
alnicolens Wood, *Pityophthorus* . . . 979
Alniphagini (= *Hylesinini*) . . . 64
alniphagus Nobuchi, *Emoporicus* . . . 849
Alniphagus Swaine . . . 73
alniphagus Wood, *Gnathotrichus* . . . 1034
alpha (Beeson), (*Xyleborus*), *Coptodryas* . . . 823
alpinensis G. Hopping, *Pityophthorus* . . . 979
alpinus Eggers, *Pityogenes* (= *calcaratus*) . . . 444
alpinus Schedl, *Xyleborus* (= *clerodendronae*) . . . 719
alsapanicus (Schedl), (*Xyleborus*), *Ambrosiodmus* . . . 670
alter (Eggers), (*Xyleborus*), *Dryocoetoides* . . . 656
alter Eggers, (*Stephanoderes*), *Hypothenemus*
 (= *hirmanus*) . . . 910
alternans (Schedl), (*Cryphalomorphus*),
Scolytogenes . . . 555
alternans (Schedl), (*Ips*), *Acanthotomicus* . . . 475
alternans Browne, *Hypothenemus* (= *criticus*) . . . 914
alternans Chapuis, *Hydurgops* (= *subcostulatus*) . . . 42
alternans Eggers, *Dryocoetes* (= *autographus*) . . . 575
alternans Eichhoff, *Xyleborus* (= *volculus*) . . . 782
alternans Schedl, *Hylesinus* (= *macmahoni*) . . . 81
alternantes Schedl, *Cyrtogenius* . . . 563
alternatus (Eggers), (*Problechilus*), *Gymnochilus* . . . 386
alternatus (Eggers), (*Stephanoderes*),
Hypothenemus . . . 906
alternatus (Eggers), (*Thannurgides*),
Coccotrypes . . . 593
alternatus Eggers, *Hylocurus* . . . 422
alternus Wood, *Hylocurus* . . . 422
altilis Schedl, *Xyleborus* . . . 710
alutaceus (Schaufuss), (*Stephanoderes*),
Ctonoxylon . . . 836
alutaceus Schwarz, *Crypturgus* . . . 620
alvarengai Bright, *Sampsonius* . . . 655
alvarengai Schedl, *Pityophthorus* . . . 990
amabilis Chamberlin, *Cryphalus* (= *ruficollis*) . . . 892
amabilis Wood, *Scolytodes* . . . 355
amakusanus (Murayama), (*Stephanoderes*),
Hypothenemus . . . 906
amanicum Hagedorn, *Ctonoxylon* . . . 836
amanicus (Eggers), (*Cryphalus*), *Scolytogenes* . . . 555
amanicus Eggers, *Cladoctonus* . . . 238
amanicus Hagedorn, *Xyleborus* . . . 710
ananicus ominosus Schedl, *Xyleborus*
 (= *amanicus*) . . . 710
amarantum (Schedl), (*Xyleborus*), *Cyclorhipidion* . . . 697
Amasa Lea . . . 651
amassius Wood, *Pseudothysanoses* . . . 412
amatus Schedl, *Acanthotomicus* . . . 478
amazonicus (Eggers), (*Loganius*), *Cnemomyx* . . . 313
amazonicus (Eggers), (*Neodryocoetes*), *Araptus* . . . 953
amazonicus Eggers, (*Stephanoderes*), *Hypothenemus*
 (= *obscurus*) . . . 937
amazonicus Schedl, (*Ceratolepis*), *Cnemomyx*
 (= *amazonicus*) . . . 313
amazonicus Schedl, *Scolytus* . . . 322
ambasinotatus Schedl, *Xyleborus* . . . 710
ambasipennis Schedl, *Xyleborus* . . . 710
ambasius Hagedorn, *Xyleborus* (= *conradti*) . . . 720
ambasiusculus Eggers, *Xyleborus* . . . 711
ambiguus Blandford, *Hylastes* . . . 43
ambitosius (Schaufuss), (*Xylocleptes*),
Premnobius . . . 651
ambitus Wood, *Microborus* . . . 383
Ambrosiodmus Hopkins . . . 670

<i>americanum</i> Wood, <i>Liparthrum</i>	275	<i>* anceps</i> Hagedorn, <i>Xylechinites</i>	1082
<i>americanus</i> Blackman, <i>Xylechinus</i>	115	<i>anceps</i> Wood, <i>Scolytodes</i>	388
<i>americanus</i> Dejean, <i>Phloeotribus</i> (= <i>scarabaeoides</i>)	230	<i>Anchonocerus</i> Eichhoff (= <i>Monarthrum</i>)	1050
<i>americanus</i> Hopkins, <i>Dryocoetes</i> (= <i>autographus</i>)	578	<i>Ancyloderes</i> Blackman (= <i>Gnathotrichus</i>)	1034
<i>* americanus</i> Wickham, <i>Hylastes</i>	43	<i>andamanensis</i> (Blandford), (<i>Xyleborus</i>), <i>Euwallacea</i>	685
<i>amersum</i> Wood, <i>Pycnarthrum</i>	384	<i>andamanensis</i> Beeson, (<i>Xyloctonus</i>), <i>Scolytomimus</i>	838
<i>amiculus</i> Wood, <i>Pityophthorus</i>	980	<i>andamanensis</i> Wood, <i>Scolytomimus</i>	838
<i>amiskwienensis</i> Hopping, <i>Ips</i> (= <i>tridens engelmanni</i>)	528	<i>andersoni</i> Swaine, <i>Carphoborus</i>	302
<i>amitinus</i> (Eichhoff), (<i>Tomicus</i>), <i>Ips</i>	489	<i>andersoni</i> Wood, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943
<i>amitinus montanus</i> Fuchs, <i>Ips</i> (= <i>amitinus</i>)	491	<i>andinus</i> Wood, <i>Chaetophloeus</i>	271
<i>amoenus</i> (Schaufuss), (<i>Hylurgus</i>), <i>Polygraphus</i>	284	<i>andresi</i> Eggers, <i>Phloeosinus</i> (= <i>armatus</i>)	241
<i>amoenus</i> Blandford, <i>Pityophthorus</i>	980	<i>andrewesi</i> (Blandford), (<i>Xyleborus</i>), <i>Xyleborinus</i>	804
<i>amoenus</i> Schedl, <i>Xyleborus</i> (= <i>asper</i>)	713	<i>andriani</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698
<i>amoenus</i> Wood, <i>Scolytodes</i>	388	<i>anepotulus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	670
<i>amorphus</i> (Eggers), (<i>Xyleborus</i>), <i>Hadrodemus</i>	818	<i>angeri</i> Pfeffer, <i>Pityophthorus</i> (= <i>buyssoni</i>)	985
<i>amphicauda</i> Browne, <i>Xyleborus</i>	711	<i>angolanus</i> Wood, <i>Hylesinopsis</i>	93
<i>Amphicranidae</i> (= <i>Corthylini</i> , <i>Corthylina</i>)	949	<i>angolensis</i> (Schedl), (<i>Afromicracis</i>), <i>Micriophalus</i>	410
<i>amphicranoides</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1051	<i>angolensis</i> (Schedl), (<i>Aridianerus</i>), <i>Hylesinopsis</i> (= <i>angolanus</i>)	93
<i>amphicranoides</i> Hagedorn, <i>Xyleborus</i>	711	<i>angolensis</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	475
<i>amphicranoides</i> Schedl, <i>Prennobius</i>	651	<i>angolensis</i> Schedl, <i>Hylesinopsis</i>	93
<i>amphicranoides latecavatus</i> Eggers, <i>Xyleborus</i> (= <i>amphicranoides</i>)	711	<i>angolensis</i> Schedl, <i>Xyleborus</i>	711
<i>amphicranoides parvior</i> Browne, <i>Xyleborus</i> (= <i>amphicranoides</i>)	711	<i>angulatus</i> (Eichhoff), (<i>Tomicus</i>), <i>Orthotomicus</i>	467
<i>amphicranulus</i> Eggers, <i>Xyleborus</i> (= <i>fallax</i>)	735	<i>angulatus</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	475
<i>Amphicranus</i> Erichson	1045	<i>angulatus</i> Schedl, (<i>Xyleboricus</i>), <i>Arixyleborus</i> (= <i>mediosectus</i>)	668
<i>amplexicauda</i> Hagedorn, <i>Xyleborus</i>	711	<i>angustatus</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>perminutissimus</i>)	809
<i>ampliatus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>angustatus</i> (Herbst), (<i>Bostrichus</i>), <i>Hylastes</i>	44
<i>ampliatus</i> Schedl, <i>Cucinus</i>	205	<i>angustatus</i> Browne, <i>Scolytus</i>	324
<i>amplipennis</i> Eichhoff, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739	<i>angustatus</i> Eichhoff, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxseni</i>)	814
<i>amplipennis</i> Wood, <i>Micracis</i>	431	<i>angusticollis</i> Eggers, <i>Hylastes</i>	45
<i>amplipennis</i> Hopkins, <i>Hypothenemus</i> (= <i>columbi</i>)	913	<i>angusticollis</i> Schedl, <i>Polygraphus</i>	285
<i>amplipennis</i> Schedl, <i>Cucinus</i>	205	<i>angustior</i> (Eggers), (<i>Cryphalus</i>), <i>Hypocryphalus</i>	868
<i>amplus</i> (Blackman), (<i>Myeloborus</i>), <i>Pityophthorus</i>	980	<i>angustior</i> Eggers, <i>Camptocerus</i>	318
<i>amplus</i> Wood, <i>Phloeotribus</i>	218	<i>angustior</i> Eggers, <i>Xyleborus</i>	711
<i>amplus</i> Wood, <i>Tricolus</i>	1042	<i>angustus</i> Blackman, <i>Pityophthorus</i>	980
<i>amputatus</i> (Blandford), (<i>Xyleborus</i>), <i>Amasa</i>	682	<i>angustus</i> Tsai & Yin, <i>Polygraphus</i>	285
<i>anaurensis</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>ratzeburgi</i>)	363	<i>angylocalyx</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	478
<i>amygdali</i> Cucrin-Meneville, <i>Scolytus</i>	322	<i>Anisandrus</i> Ferrari (= <i>Xyleborus</i>)	704
<i>amygdali rufipennis</i> Brancsik, <i>Scolytus</i> (= <i>amygdali</i>)	323	<i>anisandrus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>hagedorni</i>)	674
<i>Anacretus</i> Duges (= <i>Xyleborus</i>)	704	<i>anisopterae</i> (Browne), (<i>Ozodendron</i>), <i>Cyrtogenius</i>	563
<i>analis</i> Schedl, <i>Aricerus</i>	217	<i>anisopterae</i> Browne, <i>Xyleborus</i>	711
<i>analis</i> Schedl, <i>Xyleborus</i>	711	<i>ankius</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	859
<i>analogus</i> (Wood), (<i>Mimips</i>), <i>Acanthotomicus</i>	478	<i>ankius</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	805
<i>analogus</i> LeConte, (<i>Hylurgus</i>), <i>Tomicus</i> (= <i>piniperda</i>)	136	<i>annectens</i> (Wood), (<i>Prochramesus</i>), <i>Chramesus</i>	263
<i>analogus</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>intersectus</i>)	808	<i>annectens</i> LeConte, <i>Pityophthorus</i>	980
<i>anatolicus</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>amygdali</i>)	323	<i>annectens</i> LeConte, <i>Sciurus</i>	29
<i>anaxeus</i> Wood, <i>Phelloterus</i>	971	<i>annectens</i> Schedl, <i>Xyleborus</i>	711
<i>Anaxyleborus</i> Wood (= <i>Amasa</i>)	681	<i>annectens</i> Wood, <i>Cucinus</i>	205
<i>anceps</i> Blackman, <i>Pityophthorus</i> (= <i>nitidus</i>)	1016	<i>annexus</i> Schedl, <i>Xyleborus</i>	712

- annulicollis* Enderlein, *Coleobothrus* (= *luridus*) . . . 615
Anodius Motschulsky (= *Dryocoetes*) . . . 569
anogeissi Wood, *Polygraphus* . . . 285
anomalus (Schedl), (*Xyleborus*), *Amasa* . . . 682
anomalus Bright, (*Corthylus*), *Corthylocurus*
 (= *mexicanus*) . . . 1069
anona Hopkins, (*Xylocleptes*), *Dendrocraniulus*
 (= *carbonarius*) . . . 550
anona Hopkins, *Coccotrypes* (= *carpophagus*) . . . 596
anoplum (Schedl), (*Xyleborus*), *Cyclorhipidion* . . . 698
antaisaka Schedl, *Xyleborus* . . . 712
antanala Schedl, *Xyleborus* . . . 712
antennarius Schedl, *Ernoporus* . . . 854
antennarius Schedl, *Corthylus* . . . 1070
antennatus Schedl, *Scolytus* . . . 324
antennatus Swaine, *Phloeosinus* . . . 240
anthracinus Bright, *Pityophthorus* . . . 980
anticus (Schedl), (*Erioschidius*), *Cosmoderes* . . . 901
anticus Schedl, *Pityophthorus* . . . 981
antillicus Bright, *Pityophthorus* . . . 981
antipodus (Eggers), (*Blastophagus*), *Hylurgonotus* . . . 187
antipodus Schedl, *Hylesinus* . . . 75
antoinei Peyerimhoff, *Trietennus* . . . 541
aomoriensis Nobuchi, *Scolytus* . . . 324
apachae Bright, *Pityophthorus* . . . 981
apache Lanier, *Ips* . . . 491
apachecae Hopkins, *Conophthorus* . . . 963
apatoides Eichhoff, (*Dryocoetes*), *Hypothenemus*
 (= *tristis*) . . . 946
apertus Schedl, *Xyleborus* . . . 712
apfelbecki Eggers, *Crypturgus* (= *cinereus*) . . . 623
Aphanarthrum Wollaston . . . 616
Aphanocleptus Wood (= *Pseudothysanoes*) . . . 411
aphodioides Villa, (*Bostrichus*), *Crypturgus*
 (= *pusillus*) . . . 628
apicalis (Blandford), (*Xyleborus*), *Ambrosiodinus* . . . 670
apicalis (Schedl), (*Cryphalomorphus*),
Scolytogenes . . . 859
apicalis (Schedl), (*Ips*), *Acanthotomicus* . . . 478
apicalis Endrodi, *Trypodendron* (= *domesticum*) . . . 633
apicalis Schedl, *Polygraphus* . . . 285
apicalis Wood, *Hypothenemus* . . . 906
apicatus Beeson, *Coccotrypes* (= *longior*) . . . 605
apicatus Schedl, (*Poeecilips*), *Coccotrypes* (= *longior*) . . . 605
apicenotatus Schedl, *Pityophthorus* . . . 981
apicenotatus Schedl, *Xyleborus* (= *eximius*) . . . 735
apicipenne (Schedl), (*Xyleborus*), *Cyclorhipidion* . . . 698
apicipennis Schedl, *Pityophthorus* . . . 981
apiculatus Schedl, *Pityophthorus* . . . 981
apiculatus Schedl, *Xyleborus* . . . 712
Apidocephalus Wickham (= *Hylesinus*) . . . 74
aplanatideclivis Schedl, *Xyleborus* . . . 712
aplanatus (Wichmann), (*Xyleborus*), *Euvallacea* . . . 686
aplanatus Schedl, *Ozopemon* . . . 589
aplanatus Schedl, *Pityophthorus* (= *digestus*) . . . 995
Apoglostatus Schedl (= *Glostatus*) . . . 835
Apoxyleborus Wood (= *Xylosandrus*) . . . 787
appendicinum (Schedl), (*Pterocyclon*),
Monarthrum . . . 1051
approximatus (Schedl), (*Cryphalomorphus*),
Scolytogenes . . . 859
approximatus Dietz, *Dendroctonus* . . . 138
approximatus Hopkins, (*Stephanoderes*),
Hypothenemus (= *interstitialis*) . . . 931
approximatus Hopkins, *Cryphalus* (= *ruficollis*) . . . 891
approximatus Schedl, *Xyleborus* . . . 712
apuleiae Costa Lima, *Spermophthorus* . . . 971
aquilonius Bright, *Pityophthorus* (= *nitidus*) . . . 1016
aquilonius (Nobuchi), (*Taenioglyptes*), *Cryphalus* . . . 873
aquilus (Wood), (*Cryptulocleptus*),
Pseudothysanoes . . . 412
aquilus Blackman, *Pityophthorus* . . . 981
aquilus Blandford, *Xyleborus* . . . 712
aquilus Wood, *Camptocerus* (= *opacicollis*) . . . 320
aquilus Wood, *Chramesus* . . . 263
arabiae Schedl, *Hylesinopsis* . . . 93
arakii (Sawamoto), (*Cladoborus*), *Pityophthorus* . . . 951
aralidii Numberg, *Arixyleborus* (= *leprosulus*) . . . 667
Araptidae Eichhoff (= *Corthylini*, *Pityophthorina*) . . . 949
Araptus Eichhoff . . . 952
aratus Blandford, *Scolytus* . . . 324
aratus intermedius Kurenzov, *Scolytus* (= *aratus*) . . . 324
araucariae (Schedl), (*Conophthoracranulus*),
Araptus . . . 953
araucariae Schedl, (*Xylogopinus*), *Hylurdretonus*
 (= *corticinus*) . . . 156
araucariae Schedl, *Cryphalus* . . . 873
araucariae Schedl, *Hylurdretonus* . . . 156
araucariae Schedl, *Pachycotes* . . . 157
araucariae Schedl, *Xylechinusomus* (= *brasilienis*) . . . 105
araujiae Brethes, (*Xyleborus*), *Araptus* . . . 953
arbuti (Wood), (*Cryptulocleptus*),
Pseudothysanoes . . . 412
arbuti Hopkins, (*Xyleborus*), *Xyleborinus*
 (= *saxeseni*) . . . 815
arcaneus Bright, *Pityophthorus* . . . 981
*** arcessitus** (Scudder), (*Phloeosinutes*),
Phloeosinus . . . 240
arceuthobii Wood, *Pityophthorus* . . . 981
Archaeoscolytus Butovitsch (= *Scolytus*) . . . 321
Archeophalus Schedl (= *Hypothenemus*) . . . 905
arcticollis Blandford, (*Xyleborus*), *Cyclorhipidion*
 (= *pruinusum*) . . . 701
arcturus Samuelson, *Xyleborus* . . . 712
ardis Wood, *Phloeocleptus* . . . 421
ardis Wood, *Tricolus* . . . 1042
ardius Schedl, (*Xyleborus*), *Ambrosiodinus*
 (= *desectus*) . . . 673
areccae (Hornung), (*Bostrichus*), *Hypothenemus* . . . 906
areccae Eggers, *Coccotrypes* (= *dactyliperda*) . . . 600
argentinae Blackman, *Phloeotribus* . . . 218
argentinensis (Eggers), (*Prionosceles*), *Scolytodes* . . . 355
argentinensis (Schedl), (*Plathophloeus*),
Phloeotribus . . . 218
argentinensis Eggers, *Pityophthorus* . . . 982

<i>argentinensis</i> Eggers, <i>Scolytopsis</i> (= <i>puncticollis</i>)	321	<i>artelaevis</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	686
<i>argentinensis</i> Nunberg, <i>Neocryphus</i>	552	<i>artelineatus</i> (Beeson), (<i>Xyleborus</i>), <i>Xyleborinus</i>	805
<i>argentinensis</i> Schedl, <i>Chramesus</i>	263	<i>artelongus</i> Schedl, (<i>Xyleborus</i>), <i>Premobius</i> (= <i>longus</i>)	653
<i>argentinensis</i> Schedl, <i>Corthyus</i>	1070	<i>artemarginatus</i> Schedl, (<i>Xyleborus</i>), <i>Leptoxyleborus</i> (= <i>semigranulatus</i>)	660
<i>argentinensis</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	924	<i>artemisiae</i> Wollaston, <i>Liparthrum</i>	275
<i>argentinensis</i> Schedl, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739	<i>Artepityophthorus</i> Schedl (= <i>Cyrtogenius</i>)	562
<i>argentiniae</i> (Schedl), (<i>Bostrichus</i>), <i>Pseudothysanoes</i>	412	<i>artepunctatus</i> Eggers, <i>Dryocoetes</i> (= <i>autographus</i>)	575
<i>argentiinae</i> (Schedl), (<i>Breviophthorus</i>), <i>Araptus</i>	953	<i>artepinulosus</i> Schedl, <i>Xyleborus</i> (= <i>spinulosus</i>)	775
<i>argentiinae</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	982	<i>artestriatus</i> (Browne), (<i>Tacnioglyptes</i>), <i>Cryphalus</i>	873
<i>argentiinae</i> Schedl, <i>Dendrodicticus</i>	437	<i>artestriatus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xyleborinus</i>	805
<i>argutus</i> (Wood), (<i>Eidophelus</i>), <i>Hemicryphalus</i>	867	<i>artestriatus</i> Schedl, <i>Scolytus</i> (= <i>submarginatus</i>)	377
<i>argutus</i> Wood, <i>Amphicranus</i>	1045	<i>artestrigosus</i> Schedl, <i>Bothrostermus</i>	215
<i>Aricerus</i> Blandford	217	<i>artenuis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptoborus</i>	662
<i>Aridiamerus</i> Schedl (= <i>Hylesinopsis</i>)	92	<i>artifex</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698
<i>aries</i> (Schedl), (<i>Artepityophthorus</i>), <i>Cyrtogenius</i>	563	<i>artifex</i> Blackman, <i>Pityophthorus</i> (= <i>venustus</i>)	1033
<i>aries</i> Schedl, <i>Xyleborus</i>	713	<i>artocarpi</i> (Browne), (<i>Ips</i>), <i>Acanthotomicus</i>	478
<i>arisanus</i> Eggers, <i>Xylechinus</i>	115	<i>artocarpi</i> Browne, <i>Hypothenemus</i>	908
<i>arisanus</i> Nüsima, <i>Phlocosinus</i>	240	<i>artocarpus</i> (Schedl), (<i>Ericryphalus</i>), <i>Cryphalus</i>	873
<i>aristatae</i> Bright, <i>Pityophthorus</i> (= <i>aquilus</i>)	981	<i>artocarpus</i> Beeson, <i>Cryphalus</i> (= <i>artocarpus</i> Schedl)	874
<i>Arixyleborus</i> Hopkins	665	<i>artocarpus</i> Beeson, <i>Liparthrum</i> (= <i>artocarpus</i>)	275
<i>arizonicum</i> Wood, <i>Liparthrum</i>	275	<i>artocarpus</i> Schedl, <i>Cryphalus</i> (= <i>brownei</i>)	877
<i>arizonicus</i> Blackman, <i>Carphobius</i>	311	<i>artocarpus</i> Wood, <i>Liparthrum</i>	275
<i>arizonicus</i> Blackman, <i>Phlocosinus</i>	240	<i>artus</i> (Wood), (<i>Gnathophthorus</i>), <i>Dacnophthorus</i>	969
<i>arizonicus</i> Hopkins, <i>Dendroctonus</i> (= <i>frontalis</i>)	152	<i>arundinarius</i> Schedl, <i>Dendrochilus</i>	438
<i>armandi</i> Tsai & Li, <i>Dendroctonus</i>	138	<i>arundinis</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	908
<i>armaticeps</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698	<i>asaroriensis</i> Beeson, <i>Hypothenemus</i> (= <i>eruditus</i>)	924
<i>armaticeps</i> Schedl, <i>Hylurgonotus</i>	187	<i>ascendens</i> Schedl, <i>Pityophthorus</i>	982
<i>armatus</i> Wollaston, <i>Aphanarthrum</i>	616	<i>ascitus</i> Wood, <i>Hypothenemus</i>	908
<i>armatus</i> (Browne), (<i>Euptilius</i>), <i>Erioporus</i>	554	<i>ashanti</i> Schedl, <i>Pityophthorus</i>	982
<i>armatus</i> (Schaufuss), (<i>Xyleborus</i>), <i>Xyleborinus</i>	805	<i>ashantius</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	478
<i>armatus</i> Blandford, <i>Phlocotribus</i>	218	<i>ashuensis</i> (Murayama), (<i>Xyloterus</i>), <i>Trypodendron</i>	632
<i>armatus</i> Comolli, <i>Scolytus</i> (= <i>pygmaeus</i>)	359	<i>ashuensis</i> Murayama, <i>Xyleborus</i>	713
<i>armatus</i> Eggers, <i>Pernophorus</i>	202	<i>asiaticus</i> Pjatnitskii, <i>Phlocotribus</i> (= <i>mayeti</i>)	224
<i>armatus</i> Eggers, <i>Scolytoplatypus</i>	401	<i>asininae</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922
<i>armatus</i> Reitter, <i>Phlocosinus</i>	240	<i>asper</i> (Broun), (<i>Tomicus</i>), <i>Hypocryphalus</i>	868
<i>armatus</i> Schedl, (<i>Phlocosinopsis</i>), <i>Hyledius</i> (= <i>spinifer</i>)	261	<i>asper</i> (Schedl), (<i>Pocilips</i>), <i>Coccotrypes</i>	593
<i>armatus</i> Schedl, <i>Amphicranus</i> (= <i>propugnatus</i>)	1048	<i>asper</i> Eggers, <i>Xyleborus</i>	713
<i>armatus</i> Schedl, <i>Cryphalus</i>	873	<i>asper</i> Erichson, <i>Phloeoborus</i>	98
<i>armeniacus</i> Reitter, <i>Thamnurgus</i>	542	<i>asper</i> Sampson, <i>Hyledius</i> (= <i>nitidicollis</i>)	260
<i>armifer</i> (Schedl), (<i>Xyleborus</i>), <i>Webbia</i>	829	<i>asper</i> Swaine, <i>Hylastes</i> (= <i>gracilis</i>)	55
<i>armiger</i> Schedl, <i>Xyleborus</i>	712	<i>asperatus</i> (Blandford), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	671
<i>armillatus</i> Schedl, <i>Xyleborus</i>	712	<i>asperatus</i> (Gyllenhal), (<i>Bostrichus</i>), <i>Cryphalus</i>	874
<i>armipennis</i> Schedl, <i>Xyleborus</i>	712	<i>asperatus</i> Blandford, <i>Phlocotribus</i> (= <i>setulosus</i>)	232
<i>arnoldi</i> Semenov, <i>Liparthrum</i>	275	<i>asperatus</i> Brethes, <i>Acorthylus</i>	853
<i>arquatus</i> (Sampson), (<i>Xyleborus</i>), <i>Xylosandrus</i>	787	<i>asperatus</i> Schaeffer, <i>Chramesus</i>	263
<i>artecomanus</i> (Schedl), (<i>Xyleborus</i>), <i>Hadrodemius</i>	818	<i>asperatus</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943
<i>artecineolus</i> Schedl, (<i>Xyleborus</i>), <i>Theoborus</i> (= <i>coartatus</i>)	661	<i>asperatus</i> Wood, <i>Hylastes</i>	45
<i>artecylindrus</i> Schedl, <i>Xyleborus</i>	712	<i>aspericauda</i> Eggers, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>gracilis</i>)	808
<i>artegranulatus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	671		
<i>artegrapha</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	823		
<i>artehybridus</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	697		

- aspericollis* (Beeson), (*Thamnurgides*),
Coccotrypes 593
- aspericollis* (Eichhoff), (*Lepicerus*), *Scolytogenes* . . . 859
- aspericollis* (LeConte), (*Hylesinus*), *Alphitagus* . . . 73
- aspericollis* Eggers, *Pseudodiamerus* 194
- aspericollis* Schedl, *Chramesus* 263
- aspericollis* Strohmeier, *Phlocoborus* 98
- asperipennis* Wollaston, (*Cryphalus*), *Hypothenemus*
 (= *eruditus*) 920
- asperipennis* Eggers, (*Xyleborus*), *Eucallacea*
 (= *velatus*) 695
- asperipunctatus* Eggers, *Xyleborus* 713
- asperrinus* Schedl, *Xyleborus* 713
- aspersus* (Sampson), (*Xyleborus*), *Amasa* 682
- asperulus* (Eggers), (*Xyleborus*), *Dryococtoides* . . . 656
- asperulus* (LeConte), (*Cryphalus*),
Pseudopityophthorus 972
- asperulus* Eggers, *Phloeotribus* 219
- asperulus* Eichhoff, (*Stephanoderes*), *Hypothenemus*
 (= *obscurus*) 937
- asperulus* LeConte, (*Micracis*), *Micracisella*
 (= *opacicollis*) 431
- asperulus* Schedl, *Chramesus* 263
- asperulus* Schedl, *Cryphalus* 876
- assamensis* Eggers, (*Xyleborus*), *Eucallacea*
 (= *velatus*) 695
- assamensis* Schedl, *Scolytominus* 839
- assamensis* Stebbing, (*Tomicus*), *Xyleborus*
 (= *shoreae*) 772
- assamensis* Stebbing, *Sphaerotrypes*
 (= *sicalikensis*) 201
- assequens* Schedl, (*Pterocyclon*), *Monarthrum*
 (= *ingens*) 1057
- assequens* Schedl, *Xylosandrus* 787
- assiduus* (Schedl), (*Xyleborus*), *Coptoborus* 662
- ° *assimilis* (Schedl), (*Phlocosinites*), *Phlocosinus* . . . 241
- assimilis* Boheman, *Scolytus* (= *rugulosus*) 369
- assimilis* Eggers, *Xyleborus* 713
- assimilis* Schedl, *Glostatus* 835
- assitus* Wood, *Pityophthorus* 982
- associatus* Schedl, *Xyleborus* 713
- astutus* Schedl, *Xyleborus* 713
- atakorae* (Schedl), (*Rhopalopschion*), *Hylesinopsis* . . 93
- atava* (Schedl), (*Xyleborus*), *Coptodryas* 823
- atavus* Wood, *Cnesinus* 205
- atavus* Wood, *Phloeotribus* 219
- ater* (Eggers), (*Negritus*), *Scolytogenes* 559
- ater* (Eggers), (*Prionosceles*), *Scolytodes* 388
- ater* (Eggers), (*Stephanoderes*), *Hypothenemus* . . . 908
- ater* (Eggers), (*Xyleborus*), *Xylosandrus* 787
- ater* (Nunberg), (*Trypographus*), *Hylesinopsis* 93
- ater* (Paykull), (*Bostrichus*), *Hylastes* 45
- ater* Browne, *Cryphalus* 876
- ater* Eggers, (*Hyllocurosoma*), *Scolytodes*
 (= *aterrimus*) 388
- ater* Eggers, (*Thamnurgides*), *Coccotrypes*
 (= *barbatus*) 593
- ater* Eggers, *Carphoborus* (= *minimus*) 306
- ater* Eggers, *Dendrosinus* 236
- ater* Eggers, *Dryococtes* 572
- ater* Eggers, *Eupagiocerus* 214
- ater* Eggers, *Ozopemon* 589
- ater* Fabricius, (*Bostrichus*), *Pityogenes*
 (= *bidentatus*) 441
- ater* Hagedorn, *Diamerus* 194
- ater* Nobuchi, *Eruoporicus* 849
- ater* Nunberg, *Cryphalominus* 835
- ater* Schedl, (*Cryphalophilus*), *Scolytogenes*
 (= *uncatus*) 866
- ater* Schedl, *Cnesinus* 205
- ater* Schedl, *Corthylus* (= *letzneri*) 1074
- ater* Schedl, *Xylocleptes* 547
- ° *aterites* Schedl, *Hylastes* 49
- aterrimus* Wood, *Hypothenemus* 908
- aterrimus* (Eggers), (*Xyleborus*), *Cnestus* 801
- aterrimus* (Schedl), (*Stephanoderes*),
Hypothenemus 908
- aterrimus* Eggers, *Camptocerus* 318
- aterrimus* Eggers, *Hylastes* (= *brunneus*) 51
- aterrimus* Eggers, *Pityophthorus* 982
- aterrimus* Schedl, (*Lepicroides*), *Hypothenemus*
 (= *aterrimus*) 908
- aterrimus* Schedl, (*Pocillips*), *Coccotrypes*
 (= *nitidus*) 607
- aterrimus* Strohmeier, *Polygraphus* 285
- aterrimus* Wood, *Scolytodes* 388
- atkinsoni* Bright, *Pityophthorus* 982
- atkinsoni* Wood, *Cactopinus* 435
- atkinsoni* Wood, *Chramesus* 264
- atkinsoni* Wood, *Hyllocurus* 422
- atkinsoni* Wood, *Phloeocleptes* 421
- atlanticus* Bright, *Chaetophloeus* 271
- atlanticus* Schedl, *Phloeotribus* (= *setulosus*) 233
- atoma* (Hopkins), (*Hypothenemus*), *Trischidiidae* . . . 947
- atomarius* Chapuis, *Hylesinus* 75
- atomus* (Wood), (*Eidophelus*), *Hemicryphalus* 867
- atomus* LeConte, *Crypturgus* (= *pusillus*) 628
- atomus* Wood, *Pityophthorus* 982
- atomus* Wood, *Pseudothysanocis* 412
- atratus* LeConte, *Pityophthorus* (= *nitidulus*) 1015
- atratus* (Blandford), (*Loganius*), *Cnemidox* 313
- atratus* (Blandford), (*Prionosceles*), *Scolytodes* . . . 388
- atratus* (Schedl), (*Stephanoderes*), *Hypothenemus* . . 909
- atratus* Browne, (*Mimips*), *Acanthotomicus*
 (= *euphorbiae*) 450
- atratus* Chapuis, *Scolytus* 324
- atratus* Eichhoff, *Xyleborus* 713
- atratus atratus* (Blandford), (*Prionosceles*),
Scolytodes (= *atratus*) 388
- atratus panamensis* (Wood), (*Prionosceles*),
Scolytodes (= *atratus*) 389
- atrocis* Wood, *Cladotomus* 235
- atrocis* Wood, *Cnesinus* 205
- atrocis* Wood, *Phelloterus* 971
- atrodechloris* Wood, *Cnesinus* (= *adustus*) 205

<i>atrotibialis</i> (Eggers), (Prionosecles), Scolytodes	389	<i>babai</i> Murayama, Cryptalaus	876
<i>atrum</i> Browne, Ctonoxylon	836	<i>baculum</i> Beeson, Xyleborus	714
<i>atrum</i> Eggers, Rhopalopselion	96	<i>badius</i> (Nobuchi), (Cryptalomorphus), Scolytogenes	859
<i>attenuatus</i> (Blandford), (Xyleborus), Xyleborinus	805	<i>badius</i> Eichhoff, Xyleborus (=volvulus)	782
<i>attenuatus</i> (Eggers), (Stephanoderes), Miocryptalaus	410	<i>badius</i> Schedl, Chramesus	264
<i>attenuatus</i> Blackman, Pityophthorus	982	<i>badius</i> Wood, Tricolus	1043
<i>attenuatus</i> Erichson, Hylastes	49	<i>baguenai</i> (Schedl), (Afrochramesus), Chortastus	311
<i>attenuatus</i> Motschulsky, (Phloeotrogus), Leptoxyleborus (=sordicauda)	660	<i>baicalicus</i> Eggers, Pityogenes (=conjunctus)	451
<i>attenuatus</i> Nobuchi, (Pseudocosmoderes), Cosmoderes (=monilicollis)	901	<i>baikalicus</i> Reitter, Dryocoetes	579
<i>attenuatus</i> Wood, Araptus	953	<i>baikiae</i> Schedl, Xylocleptes	547
<i>atticus</i> Eggers, Crypturgus (=cinereus)	623	<i>bakeri</i> (Eggers), (Stephanoderes), Hypocryptalaus	868
<i>attritus</i> Peyerimhoff, Carphoborus (=pinii)	308	<i>bakeri</i> (Sampson), (Hylesinus), Ficicis	90
<i>aubei</i> Perris, (Hylesinus), Phloeosinus (=bicolor)	242	<i>bakeri</i> Hopkins, Coccotrypes (=carpophagus)	596
<i>auctor</i> Blackman, Pityophthorus (=leiophyllae)	1007	<i>bakoensis</i> Browne, Webbia	829
<i>augustae</i> Eggers, Ozopemon	589	<i>balachowskyi</i> Menier, Hypothenemus	909
<i>aulmanni</i> Hagedorn, (Cryptalaus), Hypothenemus (=seriatus)	941	<i>balanocarpi</i> Nunberg, Xyleborus (=persimilis)	762
<i>auratum</i> Hagedorn, Ctonoxylon	837	<i>balanopselaphus</i> Eggers, Cryptalaus	576
<i>auratus</i> Eggers, Eccoptopterus (=limbus)	820	<i>balbalanus</i> Eggers, Xyleborus	714
<i>auratus</i> Eggers, Xyleborus (=squamosus)	821	<i>balcanicus</i> Eggers, Scolytus (=carpini)	327
<i>auricomus</i> Blandford, Camptocerus	319	<i>balcanicus</i> Pfeiffer, Pityophthorus	982
<i>aurilegulus</i> Schaufuss, Xyleborus	714	<i>balgensis</i> Murayama, Carphoborus (=minimus)	307
<i>aurilentus</i> Bright, Pityophthorus (=murrayanae)	1014	<i>baloghi</i> (Schedl), (Stephanoderes), Hypothenemus	909
<i>australis</i> (Schedl), (Dryocoetes), Dryocoetiops	587	<i>baloghi</i> Schedl, Scolytomimus	839
<i>australis</i> (Schedl), (Ips), Acanthotomicus	479	<i>balsameus</i> Blackman, Pityophthorus	982
<i>australis</i> (Schedl), (Lepicerinus), Scolytogenes	859	<i>balsameus</i> Hopkins, Cryptalaus (=ruficollis)	891
<i>australis</i> Blackman, Pityophthorus (=tuberculatus)	1032	<i>balsameus</i> LeConte, (Tomicus), Pityokteines (=sparsus)	464
<i>australis</i> Schedl, Liparthrum	275	<i>balteatus</i> Blandford, Amphieramus	1046
<i>australis</i> Schedl, Pachycotes	187	<i>baluchistani</i> Schedl, Scolytus (=rugulosus)	370
<i>australis</i> Schedl, Phloeosinus	241	<i>bambesanus</i> Eggers, (Poclipis), Coccotrypes (=sannio)	612
<i>australis</i> Schedl, Phloeotribus (=pilula)	227	<i>bambesanus</i> Eggers, (Stephanoderes), Hypothenemus (=areccae)	908
<i>australis</i> Schedl, Xyleborus	714	<i>bambesanus</i> Eggers, Xyleborus	714
<i>australis</i> Schedl, Xylechinus	115	<i>bambusae</i> Beeson, (Thammurgides), Coccotrypes (=barbatus)	593
<i>australis</i> Schedl, Zygothlocus	271	<i>bambusae</i> Browne, Hypothenemus	909
<i>austriacus</i> (Guillebeau), (Phloeophthorus), Phloeotribus (=rhododactylus)	230	<i>bambusae</i> Browne, Ptilopodius	899
<i>austriacus</i> Wachtl, (Tomicus), Pityogenes (=trepanatus)	459	<i>bananensis</i> Eggers, (Stephanoderes), Hypothenemus (=javanus)	932
<i>autographus</i> (Ratzeburg), (Bostrichus), Dryocoetes	572	<i>bangensis</i> (Eggers), (Cryptalomorphus), Scolytogenes	859
<i>autographus</i> sachalinensis Sokanovskii, Dryocoetes (=autographus)	578	<i>bangensis</i> Eggers, Sphaerotrypes	198
<i>avarus</i> Wood, Xylechinus	115	<i>banjocwangi</i> Schedl, (Xyleborus), Xylosandrus (=mutilatus)	799
<i>avulsus</i> (Eichhoff), (Tomicus), Ips	491	<i>banksiae</i> (Schedl), (Xyleborus), Amasa	682
<i>azerbaidzhanicus</i> Michalski, Scolytus	324	<i>banksianae</i> McPherson, Conophthorus (=resinosae)	967
<i>aztecum</i> (Bright), (Corthylus), Corthycyclon	1067	<i>banosus</i> (Eggers), (Hoplites), Cladoctonus	238
<i>aztecus</i> (Wood), (Sphenoceros), Araptus	953	<i>banosus</i> (Hagedorn), (Hexacolus), Scolytodes	389
<i>aztecus</i> Bright, Pityophthorus	982	<i>baphiae</i> (Schedl), (Glochicopterus), Hylesinopsis	93
<i>aztecus</i> Wood, Dendroctonus (=parallelocollis)	159	<i>barbatogranosus</i> Schedl, Xyleborus (=setulosus)	771
<i>aztecus</i> Wood, Hylesinus	75	<i>barbatoides</i> Schedl, Xyleborus	714
<i>aztecus</i> Wood, Scolytus	324	<i>barbatomorphus</i> (Schedl), (Xyleborus), Euwallacea	686
<i>abadjanidis</i> Eggers, Liparthrum (=arnoldi)	275	<i>barbatulus</i> (Schedl), (Xyleborus), Euwallacea	686
<i>babai</i> (Murayama), (Thammurgus), Coccotrypes	593	<i>barbatus</i> (Blackman), (Pityophilus), Pityotrichus	970

- barbatus* (Blandford), (*Brachyspartus*),
Corthylocurus 1065
- barbatus* (Eggers), (*Mimulus*), *Cnemomyx* 314
- barbatus* (Hagedorn), (*Xyleborus*), *Eucallacea* 686
- barbatus* (Schedl), (*Thammurgides*), *Coccotrypes* 593
- barbatus* Eggers, *Chramesus* (= *dentatus*) 265
- barbatus* Eggers, *Lamurgus* 408
- barbatus* Hagedorn, *Sphaerotrypes* 198
- barbatus* Schedl, (*Ceratolepis*), *Cnemomyx*
 (= *errans*) 314
- barbatus* Schedl, *Dendrocranulus* 550
- barbatus* Schedl, *Scolytus* 325
- barberi* Blackman, *Pityophthorus* 983
- barberi* Blackman, *Pseudothysanoes* (= *sedulus*) 417
- barberi* Hopkins, *Dendroctonus* (= *brevicomis*) 143
- barbeyi* Strohmer, *Crypturgus* (= *numidicus*) 625
- barbieri* Schedl, *Cryptoxyleborus* 828
- barbifer* (Schedl), (*Gnathotrichus*),
Gnathotrupes 1039
- barbifer* Schedl, *Pityophthorus* 983
- barinensis* Wood, *Scolytus* 325
- bartoni* Bruck, *Pseudothysanoes* 412
- bartschti* Muhl, *Liparthrum* 275
- barumbuensis* Eggers, *Xyleborus* 714
- barumbucensis mendosus* Schedl, *Xyleborus*
 (= *barumbuensis*) 714
- basalis* Schedl, *Xyleborus* 714
- basihirtus* (Beeson), (*Cryphalus*), *Hypocyphalus* 868
- basilaris* (Wood), (*Cryphalomorphus*),
Scolytogenes 859
- basjoo* Niisima, (*Cryphalus*), *Hypothenemus*
 (= *eruditus*) 921
- bassetti* Blackman, *Pityophthorus* 983
- bassiae* Beeson, *Polygraphus* (= *anogeissi*) 285
- bassiae* Schedl, *Scolytominus* (= *assamensis*) 839
- bassiatorius* Hopkins, *Coccotrypes* (= *dactyliperda*) 601
- basutoae* Schedl, *Polygraphus* 285
- batensis* (Brisout), (*Hylastes*), *Hylurgops* 30
- batoensis* Eggers, *Diamerus* 195
- batoensis* Eggers, *Xyleborus* 714
- batoerradensis* (Schedl), (*Pseudoxyleborus*),
Amasa 682
- bauhaniae* Schedl, *Hypothenemus* (= *areccae*) 908
- bauhaniae* (Schedl), (*Stylotenus*), *Hypothenemus* 909
- baumanni* Hopkins, *Phloeosinus* 241
- beameri* Wood, *Hypothenemus* (= *gossypii*) 926
- beaveri* (Browne), (*Xyleborus*), *Xyleborinus* 805
- beaveri* Wood, *Cnesinus* 206
- beckeri* Bright, *Xyleborus* 714
- beckeri* Heqvist, *Hyllocurus* 422
- beckeri* Thatcher, *Dendroctonus* (= *valens*) 186
- beesoni* (Eggers), (*Pseudoxyleborus*), *Amasa* 682
- beesoni* Eggers, *Crypturgus* 620
- beesoni* Wood, *Mimicurus* 949
- bella* (Sampson), (*Xyleborus*), *Coptodyras* 823
- bellus* (Schedl), (*Cryptocleptes*), *Pseudothysanoes* 412
- bellus* Blackman, *Pityophthorus* (= *confusus*) 990
- bellus* Schedl, *Cryphalus* 876
- belokanicus* Stark, *Scolytus* 325
- belti* Blandford, *Amphicranus* 1046
- belti* Blandford, *Phloeoborus* 98
- bengalensis* Stebbing, (*Dryocoetes*), *Xylosandrus*
 (= *crassiusculus*) 792
- bengalensis* Wood, *Sphaerotrypes* 198
- benguensis* (Schedl), (*Xyleborus*), *Eucallacea* 687
- benguetus* Blackman, *Scolytoplatypus*
 (= *papuanus*) 405
- berbericolens* Wood, *Thysanoes* 419
- berchemiae* Blackman, *Thysanoes* 419
- berezinae* Stark, *Trypophloeus* (= *binodulus*) 844
- bergeri* Spessivtsev, *Xylechinus* 115
- berchemiae* Blackman, *Thysanoes* (= *berchemiae*) 419
- betsileo* Schedl, *Xyleborus* 714
- betulae* Hopkins, *Dryocoetes* 579
- betulae* Niisima, *Scolytus* 325
- betulae* Schedl, *Taphrorychus* 557
- betulae* Swaine, *Trypodendron* 632
- bezanozano* Schedl, *Xyleborus* 714
- bezaziani* Peyerimhoff, *Hypothenemus* 909
- bialowiczensis* Karpinski, *Pityogenes* (= *irkutensis*) 453
- bibractensis* Balachowsky, *Pityophthorus*
 (= *pityographus*) 1021
- bicallosum* (Schedl), (*Pterocyclon*), *Monarthrum* 1051
- bicallosus* Eggers, *Scolytus* (= *malii*) 344
- bicariniatus* (Nobuchi), (*Taenioglyptes*),
Cryphalus 876
- bicaudatum* Wollaston, *Liparthrum* 276
- bicaudatus* (Blandford), (*Bothrostermus*),
Sternobothrus 216
- bicaudatus* (Eggers), (*Ips*), *Acanthotomicus* 479
- bicavum* Wood, *Monarthrum* 1051
- bicinctulus* Schedl, *Xyleborus* (= *biconicus*) 715
- bicinctum* (Schedl), (*Xyleborus*), *Cyclorhipidion* 698
- bicinctum* Wollaston, *Aphanarthrum* 616
- bicinctum obsitum* Wollaston, *Aphanarthrum*
 (= *bicinctum*) 617
- bicinctum vestitum* Wollaston, *Aphanarthrum*
 (= *bicinctum*) 617
- bicinctus* Schedl, *Bothriodactonus* 310
- bicinctus* Schedl, *Cnesinus* 206
- bicinctus* Schedl, *Hypothenemus* 909
- bicinctus* Schedl, *Scolytus* 325
- bicinctus* Schedl, *Xyleborus* (= *biconicus*) 715
- bicolor* (Blandford), (*Xyleborus*), *Eucallacea* 687
- bicolor* (Browne), (*Cryphalomorphus*),
Cryphalus 876
- bicolor* (Browne), (*Streptocranus*), *Coptoborus* 662
- bicolor* (Brulle), (*Hylesinus*), *Phloeosinus* 241
- bicolor* (Eggers), (*Hexacolus*), *Scolytodes* 389
- bicolor* (Eggers), (*Lepersinus*), *Hylesinus* 76
- bicolor* (Ferrari), (*Corthylius*), *Monarthrum* 1051
- bicolor* (Herbst), (*Bostrichus*), *Taphrorychus* 557
- bicolor* Dejean, (*Bostrichus*), *Pityogenes*
 (= *chalcographus*) 446

<i>bicolor</i> Eggers, (<i>Problechilus</i>), <i>Gymnochilus</i> (= <i>reitteri</i>)	357	<i>bidentatus</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	479
<i>bicolor</i> Eggers, <i>Cuesinus</i>	206	<i>bidentatus</i> Browne, <i>Hypocryphalus</i>	865
<i>bicolor</i> Eggers, <i>Cortylus</i> (= <i>emarginatus</i>)	1073	<i>bidentatus</i> Schedl, <i>Hylocurus</i>	423
<i>bicolor</i> Eggers, <i>Hypothenemus</i> (= <i>eruditus</i>)	923	<i>bidentatus</i> <i>carniolica</i> Fuchs, <i>Pityogenes</i> (= <i>bidentatus</i>)	442
<i>bicolor</i> Eggers, <i>Microborus</i>	353	<i>bidentis</i> Wood, <i>Acanthotomicus</i>	479
<i>bicolor</i> Eggers, <i>Microcortylus</i>	1065	<i>bifidus</i> Bright, <i>Monarthrum</i> (= <i>querneum</i>)	1062
<i>bicolor</i> Eggers, <i>Polygraphus</i>	285	<i>bifidus</i> Schedl, <i>Tricolus</i>	1043
<i>bicolor</i> Eggers, <i>Scolytodes</i> (= <i>eggersi</i>)	391	<i>biformis</i> Browne, <i>Webbia</i>	830
<i>bicolor</i> Eggers, <i>Scolytus</i>	325	<i>bifoveatum</i> Wood, <i>Monarthrum</i>	1051
<i>bicolor</i> Eggers, <i>Sphaerotrypes</i>	198	<i>bifureus</i> Eichhoff, <i>Carphoborus</i>	302
<i>bicolor</i> Philippi, <i>Hylesinus</i>	76	<i>bifureus</i> Schedl, <i>Cortylus</i>	1070
<i>bicolor</i> Sampson, <i>Coccotrypes</i>	593	<i>biguttatus</i> Blandford, <i>Phloeotribus</i>	219
<i>bicolor</i> Schedl, <i>Dryocoetius</i>	557	<i>bimaculatus</i> (Eggers), (<i>Hapalogenius</i>), <i>Rhopalopselion</i>	96
<i>bicolor</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	924	<i>bimaculatus</i> (Eggers), (<i>Xyleborus</i>), <i>Cnestus</i>	802
<i>bicolor</i> Schedl, <i>Lanurgus</i>	408	<i>bimaculatus</i> (Schedl), (<i>Leperisinus</i>), <i>Chaetoptelius</i>	103
<i>bicolor</i> Sokanovskii, <i>Polygraphus</i> (= <i>poligraphus</i>)	295	<i>bimarginatus</i> Eggers, <i>Xyloctonus</i>	840
<i>bicolor</i> Strohmeier, <i>Cyrtogenius</i>	563	<i>binodatus</i> Wood, <i>Hylocurus</i>	423
<i>bicolor</i> Wollaston, <i>Aphanarthrum</i>	617	<i>binodosus</i> Nunberg, <i>Prennobius</i> (= <i>corthyloides</i>)	653
<i>bicolor</i> Wood, <i>Chiraneus</i>	264	<i>binodulus</i> (Ratzeburg), (<i>Bostrichus</i>), <i>Trypophlocus</i>	843
<i>bicolor</i> Wood, <i>Monarthrum</i> (= <i>bicoloratum</i>)	1051	<i>binodus</i> Reitter, <i>Kissophagus</i> (= <i>novaki</i>)	69
<i>bicolor</i> Wood, <i>Scolytominus</i>	839	<i>binodus</i> Wood, <i>Scolytus</i>	325
<i>bicolor</i> Wood, <i>Tricolus</i>	1042	<i>binotatus</i> Schedl, <i>Polygraphus</i>	285
<i>bicolor</i> <i>siculus</i> Eggers, <i>Taphrorychus</i> (= <i>bicolor</i>)	557	<i>biographus</i> LeConte, <i>Xyleborus</i> (= <i>celsus</i>)	718
<i>bicolor</i> <i>unimodus</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>bicolor</i>)	687	<i>biorbis</i> Blackman, (<i>Micracis</i>), <i>Hylocurus</i> (= <i>rudis</i>)	427
<i>bicoloratum</i> Wood, <i>Monarthrum</i>	1051	<i>biovalis</i> Blackman, <i>Pityophthorus</i>	983
<i>biconcarus</i> Blackman, <i>Hylocurus</i>	423	<i>bipunctatus</i> Eichhoff, <i>Amphicranus</i>	1046
<i>biconicus</i> (Schedl), (<i>Mycloborus</i>), <i>Acanthotomicus</i>	479	<i>birmanus</i> (Eggers), (<i>Phloeosinus</i>), <i>Hyledius</i>	259
<i>biconicus</i> Eggers, <i>Xyleborus</i>	714	<i>birmanus</i> (Eichhoff), (<i>Triarnocerus</i>), <i>Hypothenemus</i>	909
<i>bicornatulus</i> (Wood), (<i>Xyleborus</i>), <i>Xyleborinus</i>	806	<i>birmanus</i> Eggers, (<i>Cryphalus</i>), <i>Coriacephilus</i> (= <i>coriaceus</i>)	854
<i>bicornioides</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	802	<i>birmanus</i> Eggers, <i>Coccotrypes</i>	593
<i>bicornis</i> (Eggers), (<i>Xyleborus</i>), <i>Cnestus</i>	802	<i>birmanus</i> Eggers, <i>Hyporhynchus</i>	191
<i>bicornis</i> (Schedl), (<i>Xyleborus</i>), <i>Webbia</i>	830	<i>birmanus</i> Eggers, <i>Xyleborus</i>	716
<i>bicornis</i> Browne, (<i>Hyloperus</i>), <i>Cryptocurus</i> (= <i>spinipennis</i>)	72	<i>birosimensis</i> (Murayama), (<i>Enocryphalus</i>), <i>Cryphalus</i>	877
<i>bicornis</i> Wood, <i>Acacis</i>	193	<i>biseriatus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	910
<i>bicornis</i> Wood, <i>Carphoborus</i>	302	<i>biseriatus</i> Browne, <i>Dactylipalpus</i>	101
<i>bicornus</i> (Blackman), (<i>Micracis</i>), <i>Hylocurus</i>	423	<i>biseriatus</i> Eggers, <i>Chortastus</i> (= <i>minimus</i>)	311
<i>bicornus</i> Wood, <i>Cuesinus</i>	206	<i>biseriatus</i> Eggers, <i>Cortylus</i> (= <i>luridus</i>)	1074
<i>bicornutus</i> (Wood), (<i>Xyleborus</i>), <i>Taurodenus</i>	785	<i>biseriatus</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>curtulus</i>)	793
<i>bicostatus</i> (Sampson), (<i>Xyleborus</i>), <i>Anasa</i>	682	<i>biseriatus</i> Schedl, <i>Sphaerotrypes</i>	198
<i>bicostatus</i> Schedl, <i>Cuesinus</i>	206	<i>biseriatus</i> Schedl, <i>Xyloctonus</i>	840
<i>bicristatus</i> Chapuis, <i>Carphoborus</i> (= <i>bifureus</i>)	303	<i>bisetosus</i> Schedl, (<i>Brachyspartus</i>), <i>Monarthrum</i> (= <i>egeum</i>)	1054
<i>bicuspis</i> (Eggers), (<i>Xyleborus</i>), <i>Coptoborus</i>	662	<i>bismarcensis</i> Browne, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>libra</i>)	825
<i>bicuspis</i> Reitter, <i>Xylocleptes</i>	547	<i>bispinatus</i> Eichhoff, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739
<i>bidens</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1051	<i>bispinatus</i> Schedl, <i>Cuesinus</i>	206
<i>bidens</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	479	<i>bispinosulus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	671
<i>bidens</i> Browne, <i>Hypothenemus</i>	909	<i>bispinosus</i> (Eggers), (<i>Ips</i>), <i>Acanthotomicus</i>	479
<i>bidens</i> Fabricius, (<i>Bostrichus</i>), <i>Pityogenes</i> (= <i>bidentatus</i>)	441		
<i>bidens</i> Wood, (<i>Minips</i>), <i>Acanthotomicus</i> (= <i>bidentis</i>)	479		
<i>bidentatum</i> Wood, <i>Monarthrum</i>	1051		
<i>bidentatus</i> (Herbst), (<i>Bostrichus</i>), <i>Pityogenes</i>	439		
<i>bidentatus</i> (Motschulsky), (<i>Phloeotrogus</i>), <i>Xyleborus</i>	715		

- bispinosus* Beaver, *Glostatus* 535
bispinulus Eggers, *Trypophloeus* (= *granulatus*) 545
bispinum (Blandford), (*Pterocyclon*),
Monarthrum 1052
bispinus (Duftschmid), (*Bostrichus*), *Xylecloptes* 547
bispinus (Schedl), (*Xyleborus*), *Coptoborus* 662
bispinus Guyon, (*Bostrichus*), *Pityogenes*
 (= *bidentatus*) 442
bispinus Nobuchi, *Xyleborus* 716
bispinus Wood, *Chramesus* 264
bistridentatus (Eichhoff), (*Tomicus*), *Pityogenes* 442
bisulcatus Eichhoff, *Pityophthorus* (= *pullus*) 1025
bisuleatus Schedl, *Cnesinus* 206
bituberculatum (Eggers), (*Xyleborus*),
Cyclorhipidion 698
bituberculatum Hagedorn, *Rhopalopsclion* 96
bituberculatum Wollaston, *Liparthrum* 276
bituberculatus (Blandford), (*Gnathotrichus*),
Gnathotrupes 1039
bituberculatus Browne, *Webbia* 530
bituberculatus Eggers, (*Stephanoderes*),
Hypothenemus (= *brunneus*) 911
bituberculatus Eggers, *Prennobius* (= *caripennis*) 652
bituberculatus Hagedorn, *Xylecloptes* 549
bituberculatus Numberg, *Corthylus*
 (= *conexicauda*) 1072
bituberculatus Puzyr, *Scolytus* 325
bituberculatus Schedl, *Cnesinus* 206
biuncus Browne, *Xyleborus* 716
biuncus Reitter, *Xylecloptes* 549
biuncus Schedl, *Xyleborus* (= *ferrugineus*) 736
bivittata Kirby, (*Apatte*), *Trypodendron* (= *lineatum*) 642
blackmani (Schedl), (*Conophthocramulus*),
Pityophthorus 953
blackmani Bright, *Pityophthorus* (= *brighti*) 955
blackmani Nunberg, (*Renocis*), *Chaetophloeus*
 (= *mexicanus*) 273
blackmani Schedl, *Cnesinus* 206
blackmani Schedl, *Phloeosinus* (= *sequoiae*) 254
blackwelderi Blackman, *Phloeosinus* (= *eupressi*) 245
blaisdelli Swaine, *Carphoborus* 303
blandfordi (Schedl), (*Hexacolus*), *Scolytodes* 359
blandfordi Sampson, *Hyorrhynchus* 191
blandfordi Schaufuss, *Sphaerotrypes* 195
blandfordi Stebbing, (*Tomicus*), *Ips* (= *stebbingi*) 527
blanditus Wood, *Araptus* 953
blandulus Schedl, *Pityophthorus* 953
blandus Blackman, *Pityophthorus* 953
blandus Schedl, *Xyleborus* 716
Blastophagus Eichhoff (= *Tomicus*) 122
bobiriac Schedl, *Xyleborus* (= *pseudoambasius*) 766
bodei Eggers, *Phloeoborus* 98
bodoanus Reitter, *Xyleborus* 716
boeni Schedl, *Xyleborus* (= *autanala*) 712
boettcheri Eggers, *Sphaerotrypes* 195
boettcheri Schedl, *Cryphalus* 577
boieldicui Perrond, (*Bostrichus*), *Hypothenemus*
 (= *eruditus*) 920
boleti Marsham, (*Ips*), *Hylastes* (= *ater*) 46
boliviae (Blackman), (*Campitocerus*), *Cnemonyx* 314
boliviae (Schedl), (*Hexacolus*), *Scolytodes* 359
boliviae (Schedl), (*Stephanopodius*), *Styphlosoma* 950
boliviae Blackman, *Phloeotribus* (= *schoenbachii*) 232
boliviae Wood, (*Hoplitophthorus*), *Cladoctonus*
 (= *corumbensis*) 235
bolivianum Eggers, *Monarthrum* (= *chapuisi*) 1052
bolivianum Schedl, (*Cosmocorynus*), *Monarthrum*
 (= *boliviensis*) 1052
bolirianus (Eggers), (*Isophthorus*),
Acanthotomicus 479
bolirianus (Eggers), (*Stephanoderes*),
Hypothenemus 910
bolirianus (Schedl), (*Neodryocoetes*), *Araptus* 953
bolivianus Eggers, (*Prionosceles*), *Scolytodes*
 (= *boliriensis*) 359
bolirianus Eggers, *Corthylus* 1070
bolivianus Eggers, *Cryptocarenum* (= *seriatus*) 904
bolivianus Eggers, *Phloeotribus* (= *setulosus*) 233
bolirianus Eggers, *Pityophthorus* 954
bolirianus Eggers, *Scolytodes* 359
bolirianus Eggers, *Xyleborus* 716
bolirianus Schedl, *Chramesus* 264
bolirianus Schedl, *Gnathotrupes* 1040
bolirianus Schedl, *Scolytus* 325
boliriensis Wood, *Monarthrum* 1052
boliriensis Wood, *Scolytodes* 359
bombycinus Browne, *Scolytoplatypus*
 (= *eutomoides*) 403
bonanseai (Hopkins), (*Tomicus*), *Ips* 492
bonnairei Brisout, *Carphoborus* 303
bonnairei Reitter, (*Dendrosinus*), *Chramesus*
 (= *rotundatus*) 265
bonrouloiri (Chapuis), (*Hylastes*), *Hylurgops* 30
boops Blandford, *Microborus* 353
borassi Beeson, *Coccotrypes* (= *dactyliperda*) 602
borealis Hopkins, *Dendroctonus* (= *rufipennis*) 180
borealis Karpinski, *Carphoborus*
 (= *cholodkorskyi*) 304
borealis Nobuchi, *Xylosandrus* 757
borealis Swaine, *Crypturgus* 620
borealis Swaine, *Ips* 493
borealis Swaine, *Pityophthorus* (= *nitidus*) 1016
borealis Swaine, *Trypodendron* (= *lineatum*) 643
borealis borealis Swaine, *Ips* (= *borealis*) 493
borealis lanieri Wood, *Ips* (= *borealis*) 494
borealis swainei R. Hopping, *Ips* (= *borealis*) 493
borealis thomasi G. Hopping, *Ips* (= *borealis*) 493
borneensis (Eggers), (*Dendrugus*), *Coccotrypes* 593
borneensis (Schedl), (*Phloeosinus*), *Hyledius* 259
borneensis Browne, *Acacacis* 193
borneensis Browne, *Eidophelus* 567
borneensis Browne, *Xyleborus* 716
borneensis Eggers, *Ozopemon* 559

<i>borneensis</i> Eggers, <i>Xyleborus</i> (= <i>amplexicauda</i>)	711	<i>brasiliensis</i> Schedl, (<i>Ceratolepis</i>), <i>Cnemidophorus</i>	314
<i>borneensis</i> Schedl, <i>Acanthotomicus</i>	479	(= <i>errans</i>)	
<i>borneensis</i> Schedl, <i>Arixyleborus</i> (= <i>granifer</i>)	666	<i>brasiliensis</i> Schedl, (<i>Hexacolus</i>), <i>Scolytodes</i>	359
<i>borneensis</i> Schedl, <i>Cryphalus</i>	577	(= <i>brasilianus</i>)	
<i>borneensis</i> Schedl, <i>Cyrtogenius</i>	563	<i>brasiliensis</i> Schedl, <i>Cryphalus</i>	577
<i>Boroxylon</i> Hopkins (= <i>Xyleborus</i>)	704	<i>brasiliensis</i> Schedl, <i>Dryocoetes</i> (= <i>autographus</i>)	575
<i>borrichiae</i> Wood, <i>Pityophthorus</i>	984	<i>brasiliensis</i> Schedl, <i>Phrixosoma</i>	189
<i>bosqueiae</i> Schedl, <i>Ctonoxylon</i>	837	<i>brasiliensis</i> Schedl, <i>Pityophthorus</i> (= <i>dimorphus</i>)	996
<i>bosqui</i> (Schedl), (<i>Phacrylus</i>), <i>Acorthylus</i>	853	<i>brasiliensis</i> Schedl, <i>Pseudochramesus</i>	262
Bostrichidae, part (= <i>Scolytidae</i>)	28	<i>bravoii</i> Bright, <i>Pityophthorus</i>	954
<i>Bostrichips</i> Schedl (= <i>Pseudothysanops</i>)	411	<i>brasiliensis</i> (Blackman), (<i>Renocis</i>), <i>Chaetophloeus</i>	271
<i>bostrichoides</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	671	<i>brevi</i> Eggers, <i>Liparthinus</i> (= <i>mori</i>)	278
<i>Bostrichus</i> , part (= <i>Scolytidae</i>)	28	<i>brevicauda</i> Wichmann, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>thoracicus</i>)	378
<i>bosicelliae</i> (Stebbing), (<i>Cryphalus</i>), <i>Carphoborus</i>	304	<i>brevicollis</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	911
<i>Bothinodroctonus</i> Schedl	310	<i>brevicollis</i> (Kolenati), (<i>Hylesinus</i>), <i>Phlocotribus</i>	219
<i>Bothrosterni</i> Blandford (= <i>Bothrosternini</i>)	205	<i>brevicollis</i> (Strohmeyer), (<i>Acanthophorus</i>), <i>Peruophorus</i>	202
<i>Bothrosterninae</i> Blandford (= <i>Bothrosternini</i>)	205	<i>brevicollis</i> Browne, <i>Xyleborus</i>	716
<i>Bothrosternoides</i> Schedl	194	<i>brevicollis</i> Eggers, <i>Cryptocarenum</i>	902
<i>Bothrosternus</i> Eichhoff	215	<i>brevicollis</i> Eggers, <i>Dimerus</i>	195
<i>Bothryperus</i> Hagedorn (= <i>Phrixosoma</i>)	159	<i>brevicomatus</i> Bright, <i>Pityophthorus</i>	954
<i>botschmarki</i> Stark, (<i>Hylesinus</i>)	76	<i>brevicomis</i> LeConte, <i>Dendroctonus</i>	138
<i>bouyeriae</i> Schwarz, <i>Dendrosinus</i>	236	<i>brevicornis</i> Kirby, (<i>Apate</i>), <i>Polygraphus</i> (= <i>rufipennis</i>)	298
<i>boycei</i> Swaine, <i>Pityophthorus</i>	984	<i>brevidentatus</i> Eggers, <i>Xyleborus</i> (= <i>bidentatus</i>)	715
<i>Brachydendrulus</i> Schedl (= <i>Araptus</i>)	952	<i>Breviophthorus</i> Schedl (= <i>Pityophthorus</i>)	977
<i>Brachyspartus</i> Ferrari	1065	<i>brevior</i> (Eggers), (<i>Pelicerus</i>), <i>Cyrtogenius</i>	563
<i>braderi</i> (Browne), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	859	<i>brevior</i> (Eggers), [<i>brevius</i>], (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698
<i>bradfordi</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>brevior</i> Eggers, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>vulgaris</i>)	614
<i>brahma</i> Blandford, <i>Scolytoplatypus</i> (= <i>eutomoides</i>)	403	<i>brevior</i> Schedl, <i>Hypocryphalus</i>	868
<i>brasiliannus</i> Wood, <i>Scolytodes</i>	359	<i>brevipennis</i> Blandford, <i>Amphicranus</i>	1046
<i>brasiliensis</i> (Eggers), (<i>Prionosceles</i>), <i>Scolytodes</i>	359	<i>brevipennis</i> Kurenzov, <i>Scolytus</i> (= <i>aratus</i>)	324
<i>brasiliensis</i> (Schedl), (<i>Anchonomerus</i>), <i>Monarthrus</i>	1052	<i>brevipennis</i> Reitter, <i>Crypturgus</i> (= <i>numidicus</i>)	625
<i>brasiliensis</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	984	<i>brevipennis</i> Schedl, (<i>Xyleborus</i>), <i>Anasa</i> (= <i>schlichti</i>)	684
<i>brasiliensis</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	479	<i>brevipilosus</i> (Eggers), (<i>Blastophagus</i>), <i>Tomicus</i>	123
<i>brasiliensis</i> (Schedl), (<i>Micryphalus</i>), <i>Cryptocarenum</i>	902	<i>brevipilosus</i> (Schedl), (<i>Hylesinus</i>), <i>Ficicis</i>	90
<i>brasiliensis</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	953	<i>brevipilosus</i> Beeson, (<i>Thammurgides</i>), <i>Coccotrypes</i> (= <i>nubilus</i>)	608
<i>brasiliensis</i> (Schedl), (<i>Phthorophloeus</i>), <i>Phlocotribus</i>	219	<i>brevipilosus</i> Eggers, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>myristicae</i>)	825
<i>brasiliensis</i> (Schedl), (<i>Pseudohylesinus</i>), <i>Xylechinonotus</i>	105	<i>brevipilosus</i> Eggers, <i>Coccotrypes</i> (= <i>surinamensis</i>)	613
<i>brasiliensis</i> (Schedl), (<i>Stephanopodius</i>), <i>Styphlosoma</i>	950	<i>brevipilosus</i> Murayama, <i>Dryocoetes</i>	550
<i>brasiliensis</i> (Schedl), (<i>Xylocleptes</i>), <i>Dendrocranus</i>	550	<i>brevipilosus</i> Murayama, <i>Taphrorychus</i> (= <i>pilosus</i>)	560
<i>brasiliensis</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>obliquus</i>)	678	<i>brevipilosus</i> Schedl, <i>Cryphalus</i>	577
<i>brasiliensis</i> Eggers, <i>Amphicranus</i>	1046	<i>breris</i> (Eggers), (<i>Problechilus</i>), <i>Gymnochilus</i>	356
<i>brasiliensis</i> Eggers, <i>Scolytopsis</i>	320	<i>breris</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	911
<i>brasiliensis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915	<i>breris</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xylosandrus</i>	787
<i>brasiliensis</i> Nunberg, <i>Chramesus</i>	264	<i>breris</i> Blackman, <i>Pityophthorus</i>	984
<i>brasiliensis</i> Nunberg, <i>Hylocurus</i>	423	<i>brevi</i> Browne, (<i>Ptilopodius</i>), <i>Cryphalus</i> (= <i>tetricus</i>)	897
<i>brasiliensis</i> Nunberg, <i>Premnobius</i> (= <i>ambitosus</i>)	651	<i>brevi</i> Eggers, <i>Bothrosternus</i>	215
		<i>brevi</i> Eggers, <i>Coccotrypes</i> (= <i>carpophagus</i>)	594

- brevis* Eggers, *Microcorthyus* 1065
brevis Eggers, *Ozopemon* 589
brevis Panzer, (*Bostrichus*), *Xyleborus* (= *dispar*) 729
brevisetosus Bright, *Chranesius* (= *opacicollis*) 267
brevisetosus Bright, *Cladoctonus* 238
brevisetosus Browne, *Sphaerotrypes* 198
brevisetosus Eggers, *Pityophthorus* 984
brevisetosus Schedl, *Cnemonyx* 314
brevisetosus Schedl, *Cryphalus* (= *scabricollis*) 894
brevisetosus Schedl, *Traglostus* 410
brevisetosus Wood, *Carphoborus* 304
brevis (Eggers) [brevior], (*Xyleborus*),
Cyclorhipidion 698
breviusculus Chapuis, *Phloeoborus*
(= *punctatorugosus*) 100
breviusculus Schedl, *Xyleborus* (= *minutus*) 751
brighti Schedl, *Xyleborus* (= *hoerdenae*) 743
brighti Wood, *Cnesinus* 206
brighti Wood, *Pityophthorus* 984
brimblecombei (Schedl), (*Cryphalomorphus*),
Scolytogenes 859
brimblecombei Schedl, *Cryphalus* 877
brincki (Schedl), (*Dacryophthorus*), *Liparthrum* 276
brincki (Schedl), (*Metahylesinus*), *Hylesinopsis* 93
briscoei Blackman, *Pityophthorus* 985
brittaini Swaine, (*Pseudocryphalus*), *Chaetophloeus*
(= *heterodoxus*) 272
brittoni (Schedl), (*Pterocydon*), *Monarthrum* 1052
brosimi Wood, *Pycnarthrum* 384
brownei Beaver, *Hypothenemus* 911
brownei Bright, *Schedlia* 822
brownei Schedl, (*Xyleborus*), *Webbia* (= *seriata*) 832
brownei Schedl, *Amphicranus* 1046
brownei Schedl, *Ozopemon* 589
brownei Schedl, *Xylocleptes* 549
brownei Wood, *Cryphalus* 877
Brownia Nunberg (= *Ambrosiodmus*) 670
bruchii (Hagedorn), (*Hexacosus*), *Scolytodes* 389
bruchii Schedl, (*Scolytus*), *Scolytopsis*
(= *puncticollis*) 321
bruchoides Schedl, *Phloeocranus* 236
brucki Bright, *Pityophthorus* (= *deletus*) 994
brunigi Browne, *Scolytomimus* (= *pusillus*) 839
brunneicollis Hopkins, (*Stephanoderes*),
Hypothenemus (= *erectus*) 918
brunneipennis Hopkins, *Hypothenemus* (= *columbi*) 913
brunneipes (Eggers), (*Xyleborus*), *Ambrosiodmus* 671
brunneipes Nunberg, (*Stephanoderes*),
Hypothenemus (= *camerunus*) 912
brunnescens Wood, *Corthyus* 1070
brunneum (Eichhoff), (*Pterocydon*),
Monarthrum 1052
brunneus (Browne), (*Phloeosinus*), *Hyledius* 259
brunneus (Hopkins), (*Stephanoderes*),
Hypothenemus 911
brunneus (Nunberg), (*Corthyus*),
Corthylocurus 1065
brunneus (Nunberg), (*Hylonius*), *Xylocleptes* 549
brunneus (Nunberg), (*Pocilips*), *Coccotrypes* 594
brunneus Blackman, (*Renocius*), *Chaetophloeus*
(= *heterodoxus*) 272
brunneus Browne, *Cryphalus* 877
brunneus Browne, *Xyleborus* 716
brunneus Eggers, *Polygraphus* 285
brunneus Erichson, *Hylastes* 50
brunneus Heymons, *Sphaerotrypes* 198
brunneus Nunberg, *Chortastus* (= *medius*) 311
brunneus Nunberg, *Coccotrypes* (= *brunnipes*) 594
brunneus Schedl, *Hylurgonotus* (= *tuberculatus*) 187
brunneus Wood, *Corthyus* (= *brunnescens*) 1070
brunneus Wood, *Pseudothyssanoes* 412
° brunni (Hagedorn), (*Phloeosinites*), *Phloeosinus* 243
brunnipes Wood, *Coccotrypes* 594
bryanti Browne, *Sphaerotrypes* 198
bryanti Sampson, *Xyleborus* 716
brylinskyi Reitter, *Thammurgus* 542
bucco Schaufuss, *Xyleborus* (= *similis*) 773
bucharicus Michalski, *Scolytus* (= *rugulosus*) 364
buckhorni Blackman, *Phloeosinus* (= *punctatus*) 252
buculus Schedl, *Sampsonius* 655
Bufonius Eggers 1082
bugekeae (Schedl), (*Hyllocurus*), *Pseudomicracis* 429
bullatus Wood, *Pseudothyssanoes* 412
bulmerineqi Kolenati, (*Bostrichus*), *Taphrorychus*
(= *rillifrons*) 561
buloloensis Browne, *Cryphalus* 877
burgdorfi Hopkins, *Xyleborus* (= *spathipennis*) 774
burgosi Wood, *Micracis* 432
burjosi Wood, *Cactopinus* 435
burkei Blackman, *Pityophthorus* (= *confertus*) 990
burmanicus Beeson, *Polygraphus* (= *junnanicus*) 289
burmanicus Beeson, (*Xyleborus*), *Eutwallacea*
(= *andamanensis*) 686
burserae Wood, *Phloeoterus* 952
burserae Wood, *Pityophthorus* 985
burnensis (Eggers), (*Cryphalomorphus*),
Scolytogenes 859
buseki (Hopkins), (*Xyleborus*), *Xyleborinus* 806
buseki Blackman, (*Neodryocotes*), *Araptus*
(= *eruditus*) 956
buseki Hopkins, (*Stephanoderes*), *Hypothenemus*
(= *obscurus*) 937
busseae Schedl, *Pityophthorus* 985
butamali (Beeson), (*Xyleborus*), *Xylosandrus* 788
butovitschi Eggers, *Scolytus* (= *butovitschi* Stark) 325
butovitschi Stark, *Scolytus* 325
buxtoni Beeson, *Xyleborus* 716
buyssoni Reitter, *Pityophthorus* 985
cachani (Schedl), (*Xyleborus*), *Cyclorhipidion* 698
cachoeirinhae Schedl, *Xyleborus* 716
cacographus LeConte, (*Tomicus*), *Ips*
(= *grandicollis*) 508
cactophthorus Wood, *Cactopinus* 435
Cactopinae Chamberlini (= *Cactopinini*) 435
Cactopinini Chamberlin 435

<i>Cactopinus</i> Bright (= <i>Cactopinus</i>)	435	<i>camerunus</i> Eggers, (<i>Ozophagus</i>), <i>Polygraphus</i>	
Cactopinus Schwarz	435	(= primus)	296
<i>cacuminatus</i> Blandford, <i>Pityophthorus</i>	985	<i>camerunus</i> Eggers, (<i>Pseudocrypturgus</i>), <i>Microborus</i>	
cacuminatus Eggers, <i>Xyleborus</i>	716	(= boops)	383
<i>caclator</i> Blackman, <i>Pityophthorus</i> (= aquilus)	981	camerunus Eggers, <i>Pityophthorus</i>	986
<i>caclator</i> Browne, (<i>Xyleborus</i>), <i>Cryptoxyleborus</i>		camerunus Hagedorn, <i>Dactylipalpus</i>	101
(= nanus)	828	camerunus Hagedorn, <i>Strombophorus</i>	202
caelatus (Eichhoff), (<i>Tomicus</i>), <i>Orthotomicus</i>	467	<i>camerunus</i> Hagedorn, <i>Xyleborus</i> (= alluaudi)	710
caelatus (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	479	camerunus Schaufuss, <i>Chortastus</i>	311
<i>caelatus</i> Blanchard, (<i>Phloeotripes</i>), <i>Phloeoborus</i>		<i>camerunus</i> rugosus Eggers, <i>Xyleborus</i> (= alluaudi)	710
(= scaber)	100	<i>canopinus</i> Hagedorn, (<i>Xyleborus</i>), <i>Taurodenus</i>	
<i>caclebs</i> Blandford, <i>Taurodenus</i> (= godmani)	785	(= splendidus)	786
caesalpiniae Blackman, <i>Spermophthorus</i>	971	camphorae (Eggers), (<i>Webbia</i>), <i>Arixyleborus</i>	665
caesius Hagedorn, <i>Diamerus</i>	195	camphorae (Hagedorn), (<i>Xyleborus</i>),	
calamarius Wood, <i>Corthylus</i>	1070	<i>Ambrosiodinus</i>	672
calamoides (Murayama), (<i>Xyleborus</i>), <i>Anasa</i>	682	camphoratus Tsai & Yin, <i>Phloeosinus</i>	243
<i>calapanus</i> Eggers, (<i>Thamnurgides</i>), <i>Coccotrypes</i>		camptospermae (Browne), (<i>Poclipis</i>),	
(= myristicae)	606	<i>Coccotrypes</i>	594
<i>calcaratus</i> Dejean, <i>Pityogenes</i>		<i>Camptoceridae</i> Chapuis (= <i>Scolytini</i>)	313
(= calcaratus Eichhoff)	444	<i>Camptocerinae</i> Chapuis (= <i>Scolytini</i>)	313
calcaratus (Eichhoff), (<i>Tomicus</i>), <i>Pityogenes</i>	444	<i>Camptocerini</i> Chapuis (= <i>Scolytini</i>)	313
caliculus (Schedl), (<i>Gnathocortus</i>),		Camptocerus Latreille	318
<i>Gnathotripes</i>	1040	<i>canadensis</i> Blackman, <i>Hylastes</i> (= porculus)	61
californicus (Swaine), (<i>Leperisinus</i>), <i>Hylesinus</i>	76	<i>canadensis</i> Chamberlin, <i>Cryphalus</i> (= ruficollis)	892
californicus Bright, <i>Pityophthorus</i>	985	<i>canadensis</i> Swaine, <i>Phloeosinus</i>	243
<i>californicus</i> Essig, (<i>Leperisinus</i>), <i>Hylesinus</i>		<i>canadensis</i> Swaine, <i>Pityophthorus</i> (= cariniceps)	986
(= californicus Swaine)	76	<i>canadensis</i> Swaine, <i>Xyleborus</i> (= xylographus)	784
<i>californicus</i> Hopkins, (<i>Xylocleptes</i>), <i>Dendrocranius</i>		canaliculatus (Eggers), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	665
(= cucurbitae)	550	canaliculus Wood, <i>Scolytodes</i>	389
<i>californicus</i> Hopkins, <i>Dendroctonus</i> (= rufipennis)	173	canalis Wood, <i>Scolytodes</i>	389
californicus Hopkins, <i>Hypothenemus</i>	912	canariense Wollaston, <i>Aphanarthrum</i>	617
<i>californicus</i> LeConte, <i>Scolytus</i> (= scolytus)	376	<i>canariense</i> longipes Israelson, <i>Aphanarthrum</i>	
<i>californicus</i> Schedl, (<i>Plesiophthorus</i>), <i>Dendroterus</i>		(= canariense)	617
(= striatus)	952	<i>canariensis</i> Eggers, <i>Coccotrypes</i> (= carpophagus)	596
californicus Wood, <i>Xyleborus</i>	716	canarii Browne, <i>Xyleborus</i>	716
caliginis Wood, <i>Corthyccyclon</i>	1067	canarivorus Browne, <i>Xyleborus</i>	716
callidus Schedl, <i>Corthylus</i>	1070	cancellatus (Chapuis), (<i>Bothrostermus</i>),	
calligraphus (Germar), (<i>Tomicus</i>), <i>Ips</i>	494	<i>Sternobothrus</i>	216
callosum (Schedl), <i>Cyclorhipidion</i>	698	cancellatus Blandford, <i>Hyllocurus</i>	423
callosus (Schedl), (<i>Pterocyclon</i>), <i>Amphicranus</i>	1046	cancellatus Eggers, <i>Xyleborus</i>	717
calmicolens Wood, <i>Corthylus</i>	1070	<i>cancellatus</i> pronunciatus Eggers, <i>Xyleborus</i>	
calvus Schedl, <i>Xylechinus</i> (= imperialis)	116	(= cancellatus)	717
camela (Eggers), (<i>Xyleborus</i>), <i>Coptodryas</i>	823	candidus (Nobuchi), (<i>Cryphalomorphus</i>),	
camelliae (Nobuchi), (<i>Cryphalomorphus</i>),		<i>Scolytogenes</i>	860
<i>Scolytogenes</i>	860	canescens Wollaston, <i>Aphanarthrum</i>	617
camerunum Hagedorn, <i>Ctonoxylon</i>	837	<i>canescens</i> polyspiniger Israelson, <i>Aphanarthrum</i>	
<i>camerunum</i> hirsutum Hagedorn, <i>Ctonoxylon</i>		(= canescens)	617
(= flavescens)	837	<i>canescens</i> simplex Wollaston, <i>Aphanarthrum</i>	
camerunus (Eggers), (<i>Minips</i>), <i>Acanthotomicus</i>	479	(= canescens)	617
camerunus (Eggers), (<i>Stephanoderes</i>),		cannularius Wood, <i>Corthylus</i>	1070
<i>Hypothenemus</i>	912	canus (Browne), (<i>Phloeosinus</i>), <i>Hyledius</i>	259
camerunus (Hagedorn), (<i>Araptus</i>),		<i>canus</i> Blackman, <i>Chramesus</i> (= subopacus)	269
<i>Pseudomicracis</i>	429	canus Niisima, <i>Xyleborus</i>	717
camerunus (Hagedorn), (<i>Hypaspistes</i>),		capensis (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698
<i>Tiarophorus</i>	539	capensis (Schedl), (<i>Dacryophthorus</i>), <i>Xylechinus</i>	115
camerunus (Schedl), (<i>Dendrocranius</i>),		capensis Schedl, <i>Ctonoxylon</i>	837
<i>Xylocleptes</i>	549	capensis Schedl, <i>Lamurgus</i>	408
		capensis Schedl, <i>Strombophorus</i>	202

- capensis* Schedl. *Thammurgus* 542
capitalis Beeson, *Hypothenemus* (= *ureccue*) 907
capitalis Wood, *Tricolus* 1043
capito Schaufuss, *Xyleborus* (= *similis*) 773
caplandicus Numberg, (*Pseudohylocurus*), *Lamurgus*
 (= *podocarpi*) 409
caplandicus Schedl. *Hypocryphalus* 865
capronatus Perris, (*Dryocoetes*), *Taphrorychus*
 (= *villifrons*) 561
capucinicollis Schedl. *Cryphalus* 577
capucinoides Eggers, (*Xyleborus*), *Dryocoetoides*
 (= *capucinus*) 657
capucinoides Eggers, *Cryphalus* 877
capucinomorphus Schedl. *Cryphalus* 577
capucinulus (Schedl.), (*Streptocranus*),
Coptoborus 663
capucinus (Eichhoff), (*Xyleborus*), *Dryocoetoides* 656
capucinus (Schedl.), (*Cryphalops*), *Emoporius* 549
capucinus Schedl. *Cryphalus* 577
capucinus Schedl. *Tiarophorus* 539
caracicolai Hopkins, *Dryocoetoides* 657
caraibica Schedl. *Phixosoma* 159
caraibicus Eggers, *Cryptocarenum* (= *heveae*) 903
caraibicus Eggers, *Pagiocerus* (= *frontalis*) 214
caraibicus Eggers, *Xyleborus* 717
caraibicus Schedl. (*Cryphalomorphus*), *Scolytogenes*
 (= *knabi*) 863
caraibicus Schedl. (*Pocilips*), *Coccotrypes*
 (= *cyperii*) 599
carapae Wood, *Liparthrum* 276
carbonarius (Eggers), (*Stephanoderes*),
Hypothenemus 912
carbonarius (Ferrari), (*Xylocleptes*),
Dendrocramulus 550
carbonarius Eggers, *Xyleborus* (= *conradti*) 720
carbonarius Fitch, *Hylastes* (= *porculus*) 60
carbonarius Schedl. *Cnesinus* 206
 ° *carbonarius* Scudder, *Dryocoetes* 550
cardamomi Schaufuss, *Coccotrypes* 594
Cardoctonus Schedl. 253
caribacus Blackman, (*Neodryocoetes*), *Araptus*
 (= *hymenaeae*) 957
carinatum Wood, *Monarthrum* 1052
carinatum Wood, *Pycnarthrum* 354
carinatus Bright, *Pityophthorus* 956
carinatus Browne, *Margadillius* 595
carinatus Chapuis, *Scolytus* 325
carinatus Eggers, *Sphaerotrypes* 199
carinatus Eggers, *Sternobothrus* 216
carinatus Wood, *Cactopinus* 435
carinatus Wood, *Cnesinus* 206
carinatus carinatus Bright, *Pityophthorus*
 (= *carinatus*) 956
carinatus monticolae Bright, *Pityophthorus*
 (= *carinatus*) 956
carinensis (Eggers), (*Dendrugus*), *Coccotrypes* 594
carinensis Eggers, *Xyleborus* 717
cariniceps LeConte, *Pityophthorus* 956
cariniceps Schedl. *Arxyleborus* 666
carinifer Schedl. *Glostatus* 535
carinifrons Atkinson, *Hylocurus* 423
carinifrons (Blandford), (*Pityophthorus*),
Araptus 953
carinipennis Eichhoff, (*Xyleborus*), *Ambrosiodmus*
 (= *obliquecaudata*) 677
carinulatus (LeConte), (*Cryphalus*), *Pityogenes* 445
carinulatus Eggers, *Xyleborus* 717
carinulatus Swaine, *Pityophthorus* 956
carinulatus Wood, *Micracis* 432
carinulum Wood, *Monarthrum* 1052
carinulus Wood, *Micracis* 432
caritus (Wood), (*Cryptulocleptus*),
Pseudothysanops 412
carmeli Swaine, *Pityophthorus* 957
carniolica Fuchs, *Pityogenes* (= *bidentatus*) 442
carniolicus Wichmann, *Pityophthorus* 957
Carphobius Blackman 311
Carphoboridae Nussli (= *Polygraphini*) 253
Carphoborites Schedl. 1052
carphoboroides (Numberg), (*Trogloeditica*),
Acacasis 193
carphoboroides Eggers, *Polygraphus* 255
Carphoborus Eichhoff 302
Carphodicticini Wood 437
Carphodicticus Wood 437
Carphotoreus Wood 237
carpini (Ratzeburg), (*Eccoptogaster*), *Scolytus* 325
carpini Berger, *Cryphalus* 577
carpini Eggers, *Dryocoetes* 550
carpini Eggers, *Sphaerotrypes* 199
carpini Stark, *Dryocoetes* (= *carpini* Eggers) 550
carpinivorus Choo & Woo, *Dryocoetes* 550
carpinivorus Murayama, *Cryphalus* 575
carpophagus (Hornung), (*Bostrichus*),
Coccotrypes 594
Carpophilocus Schedl. (= *Cyrtogenius*) 562
Carposinus Hopkins (= *Cyrtogenius*) 562
carri Swaine, *Carphoborus* 304
carya Riley, *Scolytus* (= *quadrispinosus*) 361
caryae Walsh, *Scolytus* (= *quadrispinosus*) 361
caryi Hopkins, *Dryocoetes* 550
cascoensis Blackman, *Pityophthorus* 957
caseariae Wood, *Hylesinus* 76
cassavacensis Schedl. *Hypothenemus* (= *seriatus*) 943
cassiae Eichhoff, (*Stephanoderes*), *Hypothenemus*
 (= *obscurus*) 937
castaneae Schedl. *Arxyleborus* 666
castaneus (Schedl.), (*Gnathotrichus*),
Gnathotrupes 1040
castaneus Browne, (*Dryocoetes*), *Cyrtogenius*
 (= *perakensis*) 565
castaneus Browne, *Xylocleptes* 549
castaneus Ferrari, *Corthylus* (= *letzneri*) 1074
castaneus Ratzeburg, (*Eccoptogaster*), *Scolytus*
 (= *muli*) 344

<i>castaneus</i> Schedl. <i>Microcorthyus</i>	1065	<i>cedri cedri</i> Brisout, <i>Phlocosinus</i> (= <i>cedri</i>)	244
<i>castaneus</i> Wood. (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>cedri libani</i> Balachowsky, <i>Phlocosinus</i> (= <i>cedri</i>)	244
<i>catenatus</i> (Schedl), (<i>Macraciops</i>), <i>Lanurgus</i>	408	<i>cedri maura</i> Balachowsky, <i>Phlocosinus</i> (= <i>cedri</i>)	244
<i>Catenophorus</i> Nunberg	237	<i>cedris</i> Eichhoff, <i>Crypturgus</i> (= <i>pusillus</i>)	628
<i>catharinensis</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	672	<i>ceibae</i> Hopkins, <i>Hypothenemus</i>	912
<i>catulus</i> (Blandford), (<i>Xyleborus</i>), <i>Coptoborus</i>	663	<i>celatus</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	988
<i>catulus</i> Blackman, <i>Pityophthorus</i> (= <i>boycei</i>)	984	<i>celatus</i> Wood, <i>Xyleborinus</i>	806
<i>caucasicus</i> (Lindemann), (<i>Ernoporus</i>), <i>Eruoporus</i>	849	<i>celoides</i> Hagedorn, <i>Xyleborus</i>	717
<i>caucasicus</i> Butovitch, <i>Scolytus</i> (= <i>rugulosus</i>)	370	<i>celsus</i> Eichhoff, <i>Xyleborus</i>	717
<i>caucasicus</i> Reitter, <i>Phloeotribus</i>	219	<i>celtis</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	480
<i>caucasicus</i> Reitter, <i>Thannurgus</i>	542	<i>celtis</i> Schedl, <i>Strombophorus</i>	202
<i>caudatulus</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	479	<i>cembrae</i> (Heer), (<i>Bostrichus</i>), <i>Ips</i>	497
<i>caudatum</i> Schedl, <i>Ctonoxylon</i>	837	<i>cembrae</i> Seitner, <i>Polygraphus</i> (= <i>grandiclava</i>)	287
<i>caudatus</i> (Browne), (<i>Mimips</i>), <i>Acanthotomicus</i>	479	<i>cembroides</i> Wood, <i>Conophthorus</i> (= <i>edulis</i>)	964
<i>caudatus</i> Browne, (<i>Hyloperus</i>), <i>Cryptocurus</i> (= <i>spinipennis</i>)	72	<i>cephalonica</i> Pfeffer, <i>Pityophthorus</i>	988
<i>caudatus</i> Eggers, <i>Amphicranus</i> (= <i>dohrui</i>)	1046	<i>ceramensis</i> (Schedl), (<i>Xyleborus</i>), <i>Leptoxyleborus</i>	659
<i>caudatus</i> Eggers, <i>Scolytus</i>	327	<i>cerasi</i> Eggers, <i>Dryocoetes</i>	580
<i>caudatus</i> Schedl, <i>Xyleborus</i>	717	<i>cerasi</i> Eggers, <i>Xyleborus</i> (= <i>dispar</i>)	730
<i>caudatus</i> Wood, <i>Phloeocleptus</i>	421	<i>cerasi</i> Stark, <i>Dryocoetes</i> (= <i>cerasi</i> Eggers)	580
<i>caratus</i> Bright, <i>Pityophthorus</i>	987	<i>Ceratolepis</i> Chapuis (= <i>Cnemomyx</i>)	313
<i>caratus</i> Browne, <i>Xyleborus</i>	717	<i>ceratoniae</i> Peyerimhoff, <i>Taphrorychus</i>	559
<i>catenarius</i> Zimmermann, (<i>Hylurgus</i>), <i>Hylastes</i> (= <i>porculus</i>)	60	<i>ceylonae</i> Schedl, <i>Webbia</i>	830
<i>catifrons</i> Mannerheim, (<i>Bostrichus</i>), <i>Trypodendron</i> (= <i>lineatum</i>)	643	<i>ceylonicus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	860
<i>catifrons</i> Nunberg, (<i>Mimips</i>), <i>Acanthotomicus</i> (= <i>biconicus</i>)	479	<i>ceylonicus</i> (Schedl), (<i>Phlocosinus</i>), <i>Hyledius</i>	259
<i>carifrons</i> Nunberg, <i>Cortulys</i>	1070	<i>ceylonicus</i> (Schedl), (<i>Ptilopodius</i>), <i>Ernoporus</i>	850
<i>capipennis</i> (Schedl), (<i>Styracopterus</i>), <i>Styracoptinus</i>	281	<i>ceylonicus</i> Schedl, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>cacipennis</i> Eichhoff, <i>Premnobius</i>	651	<i>Chaetophloeus</i> LeConte	271
<i>cacipennis spinosus</i> Hagedorn, <i>Premnobius</i> (= <i>ambitosus</i>)	651	<i>Chaetophorus</i> Fuchs (= <i>Chaetoptelius</i>)	103
<i>caruloides</i> Browne, <i>Xyleborus</i>	717	<i>Chaetoptelius</i> Fuchs	103
<i>carulus</i> Browne, <i>Xyleborus</i>	717	<i>chagnoni</i> Swaine, <i>Ips</i> (= <i>grandicollis</i>)	508
<i>cacus</i> LeConte, (<i>Cryphalus</i>), <i>Monarthrum</i> (= <i>scutellare</i>)	1062	<i>chalconensis</i> Hopkins, <i>Pityophthorus</i>	988
<i>ceanothi</i> Blackman, <i>Stenoclyptus</i> (= <i>sulcatus</i>)	419	<i>chalcographus</i> (Linnaeus), (<i>Dermestes</i>), <i>Pityogenes</i>	445
<i>cecropiae</i> Wood, <i>Chramesus</i>	264	<i>chalconensis</i> Hopkins, <i>Pityophthorus</i> (= <i>chalconensis</i>)	988
<i>cecropiavorus</i> Wood, <i>Scolytodes</i>	390	<i>Chalcohyus</i> Blackman (= <i>Pseudothysanoes</i>)	411
<i>cecropiavorus acuminatus</i> Wood, <i>Scolytodes</i> (= <i>cecropiavorus</i>)	390	<i>chamaecipariae</i> Nisima, <i>Cryphalus</i>	878
<i>cecropiavorus cecropiavorus</i> Wood, <i>Scolytodes</i> (= <i>cecropiavorus</i>)	390	<i>chamberlini</i> Blackman, <i>Phlocosinus</i> (= <i>punctatus</i>)	252
<i>cecropiavorus punctifer</i> Wood, <i>Scolytodes</i> (= <i>cecropiavorus</i>)	390	<i>chamberlini</i> Swaine, <i>Ips</i> (= <i>concinuus</i>)	500
<i>cecropicoleus</i> Wood, <i>Scolytodes</i>	390	<i>championi</i> (Blandford), (<i>Eulytocerus</i>), <i>Phloeotribus</i>	219
<i>cecropii</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	390	<i>chapini</i> (Blackman), (<i>Renocis</i>), <i>Chaetophloeus</i>	271
<i>cecropii</i> Wood, <i>Corthyus</i>	1070	<i>chapuisi</i> Blandford, (<i>Phloeophthorus</i>), <i>Phloeotribus</i> (= <i>spinulosus</i>)	234
<i>cedrelae</i> Wood, <i>Scolytodes</i>	390	<i>chapuisi</i> Blandford, <i>Ariceus</i>	217
<i>cedri</i> Beeson, <i>Pityophthorus</i> (= <i>cedri</i> Wood)	987	<i>chapuisi</i> Kirsch, <i>Monarthrum</i>	1052
<i>cedri</i> Brisout, <i>Phlocosinus</i>	244	<i>chapuisi</i> LeConte, <i>Chramesus</i>	264
<i>cedri</i> Wood, <i>Pityophthorus</i>	987	<i>chapuisi</i> Wood, <i>Scolytodes</i>	390
		<i>chapuisii</i> Eichhoff, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>dissimilis</i>)	917
		<i>characiae</i> Rosenhauer, <i>Thannurgus</i>	542
		<i>charpentierae</i> Schedl, <i>Camptocerus</i>	319
		<i>chelogaster</i> Schedl, <i>Scolytus</i>	327
		<i>cherenus</i> Eggers, <i>Dianerus</i>	195
		<i>chiapensis</i> Bright, <i>Pityophthorus</i> (= <i>nigricans</i>)	1015

- chientzuangensis* Tsai & Li, *Cryphalus*
(=*tabulaeformis*) 596
- chikisani** Niisima, *Scolytus* 327
- chilensis** (Schedl), (*Conophthoracranulus*), *Araptus* 954
- chilgoza* Beeson, *Pityophthorus* (= **chilgoza** Wood) 958
- chilgoza** Wood, *Pityophthorus* 958
- chiliensis** (Nunberg), (*Siquamosinus*), *Xylechinus* 115
- chiliensis* Eggers, *Phlocotribus* (= **willi**) 235
- Chilodendron* Schedl (= *Hylesinopsis*) 92
- Chiloxylon** Schedl 587
- chimbui** (Schedl), (*Pocilips*), *Coccotrypes* 597
- chimbui** (Schedl), (*Xyleborus*), *Coptodryas* 823
- chimbui* Schedl, (*Xyleborus*), *Coptodryas* (= **rosseli**) 827
- chimbui* Schedl, (*Xyleborus*), *Coptodryas*
(=**chimbui**) 823
- chinensis** Kurenzov & Kononov, *Ips* 500
- chinlingensis** Tsai & Li, *Cryphalus* 578
- chir* Beeson, *Hylurgus* (= **indicus**) 119
- chir* Beeson, *Pityophthorus* (= **deodara**) 994
- chiricahua* Blackman, *Phloeosinus* (= **cristatus**) 245
- chirindaensis** (Schedl), (*Dryocoetes*), *Cyrtogenius* 564
- chiriquensis** (Blandford), *Acanthotomicus* 450
- chiriquensis* Eggers, *Pagiocerus* (= **frontalis**) 214
- chiriquensis** Wood, *Stegomerus* 852
- chlorophorae** (Schedl), (*Dryocoetes*), *Cyrtogenius* 564
- chloropus* Duftschmid, *Hylastes* (= **ater**) 49
- chloroticus* Dejean, *Ips* (= **calligraphus**) 494
- cholodkovskiy** Berger, *Hylesinus* 76
- cholodkovskiy** Spessivtsev, *Carphoborus* 304
- Chondronoderes* Schedl (= *Hypothenemus*) 905
- Chortastus** Schaufuss 311
- Chramesus** LeConte 262
- chrysophylli** (Eggers), (*Xyleborus*), *Coptodryas* 823
- chujoi** Schedl, *Xyleborus* 718
- cicatricosus** (Blandford), *Dactylipalpus* 101
- cicatricosus** (Schedl), (*Lepicerinus*), *Scolytogenes* 860
- ciliaris* Rey, *Scolytus* (= **scolytus**) 371
- ciliatipennis** Schedl, *Microcaphalus* 410
- ciliatipes** Blandford, *Cryphalus* 578
- ciliatoformis** Schedl, *Xyleborus* 718
- ciliatum* Eggers, *Liparthrum* (= **curtum**) 276
- ciliatus** (Hagedorn), (*Pocilips*), *Acanthotomicus* 450
- ciliatus** Blackman, *Pityophthorus* 958
- ciliatus** Eggers, *Xyleborus* 718
- ciliatus* Schedl, *Gnathotrupes* (= **longiusculus**) 1041
- cinchonae* Schedl, (*Hapalogenius*), *Hylesinopsis*
(=**seriatus**) 95
- cinchonae* Veen, *Xyleborus* (= **emarginatus**) 733
- cincinmatum** (Eichhoff), (*Trypocranus*),
Monarthrum 1052
- cincinmatum** (Blandford), (*Pityophthorus*),
Pseudopityophthorus 972
- cincinmatum** Bright, *Corthylocorus* 1065
- cinctipennis* Schedl, (*Xyleborus*), *Xyleborinus*
(=**saxeseni**) 816
- cinctipes** Schedl, *Xyleborus* 718
- cinctus* Chapuis, *Camptocerus* (= **suturalis**) 320
- cinerascens** Fairmaire, *Diamerus* 195
- cinereotestaceus** (Motschulsky), (*Hypoborus*),
Cryphalus 575
- cinereus** (Herbst), (*Bostrichus*), *Crypturgus* 621
- cinereus* Herbst, (*Bostrichus*), *Eruoporus* (= **tiliae**) 557
- cinereus* Swaine, (*Leperisinus*), *Hylesinus*
(=**aculeatus**) 75
- cingulatus** Blandford, *Hylesinus* 77
- cinnamomi** (Eggers), (*Thannurgides*),
Coccotrypes 597
- cinnamomi* Tsai & Yin, *Phloeosinus* (= **machilus**) 249
- circulicauda* Browne, (*Xyleborus*), *Amasa*
(=**cylindriciformis**) 652
- circumcinctus** Schedl, *Prennobius* 652
- circumcisulus** (Schedl), (*Xyleborus*), *Amasa* 652
- circumciscum** (Sampson), (*Xyleborus*),
Cyclorhipidion 695
- circumciscus** Schedl, *Webbia* 830
- circumdatus** Fonseca, *Coccotrypes* 597
- circumdentatus* Schedl, (*Xyleborus*), *Prennobius*
(=**circumspinitus**) 652
- circumspinitus** Eggers, *Prennobius* 652
- circumspinosus** Schedl, *Xyleborus* 718
- cirratus* Nunberg, (*Loganius*), *Cnemonyx*
(=**cliffornis**) 314
- cirratus** Schedl, *Gnathotrupes* 1040
- cirritus** Wood, *Corthylus* 1070
- cirrus* Schedl, *Corthylus* (= **flagellifer**) 1073
- Cissophagus* (= **Kissophagus**) 67
- Cisurgus** Reitter 630
- citri* Beeson, (*Xyleborus*), *Ambrosiodmus*
(=**asperatus**) 671
- citri* Ebeling, *Hypothenemus* (= **eruditus**) 924
- citrus* Blackman, *Pityophthorus* (= **annectens**) 950
- Cladoborus* Sawamoto (= *Pityophthorus*) 977
- Cladoctonus** Strohmeier 237
- Cladoctoporus* Schedl (= *Triotenus*) 540
- clarki** Wood, *Hyllocorus* 423
- clarus** Blackman, *Pityophthorus* 955
- clarus* Wood, *Eupagiocerus* (= **dentipes**) 214
- clavatus** Schedl, *Pachycotes* 157
- claviger** Blandford, *Scolytus* 325
- clavus* Wollaston, *Hylastes* (= **linearis**) 56
- clematis** (Wood), (*Gnathophthorus*),
Dacnophthorus 969
- clerodendronae** Schedl, *Xyleborus* 718
- clerodendronae* alpinus Schedl, *Xyleborus*
(=**clerodendronae**) 719
- clerodendronae* morosus Schedl, *Xyleborus*
(=**clerodendronae**) 719
- clivus** Bright, *Pityophthorus* 958
- cloudcrofti* Swaine, *Ips* (= **grandicollis**) 509
- clunicus* Hopkins, *Conophthorus* (= **coniperda**) 964
- clusiacolens** Wood, *Scolytodes* 390
- clusiae** Wood, *Phrixosoma* 159
- clusiae** Wood, *Scolytodes* 390
- clusiavorus** Wood, *Scolytodes* 390
- Cnemonyx** Eichhoff 313

Cnesinus LeConte	205	Comesiella DelGuercio (=Phloeotribus)	218
Cnestus Sampson	801	comitabilis Wood, Scolytodes	390
<i>coadunatus</i> Marsham, (<i>Ips</i>), <i>Pteleobius</i> (= <i>rittatus</i>)	72	<i>commixtus</i> Blackman, (<i>Renocis</i>), <i>Chaetophloeus</i> (= <i>heterodoxus</i>)	272
coartatus (Sampson) , (<i>Xyleborus</i>), <i>Theoborus</i>	661	commixtus Blandford, Xyleborus	719
coccotrypanoides (Schedl) , (<i>Lepicerinus</i>), <i>Scolytogenes</i>	860	communis (Schaufuss) , (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	860
Coccotrypes Eichhoff	591	<i>communis</i> Schaufuss, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	921
<i>coccotrypoides</i> Eggers, (<i>Xyleborus</i>), <i>Theoborus</i> (= <i>villosulus</i>)	662	comosus Blackman, Pityophthorus	988
<i>coficola</i> Campos Novaes, (<i>Xyleborus</i>), <i>Hypothenemus</i> (= <i>hampei</i>)	930	comosus Bright, Hypothenemus	913
coffae (Eggers) , (<i>Dryocoetes</i>), <i>Dryocoetiops</i>	557	comosus Bright, Pseudopityophthorus	972
<i>coffae</i> Hagedorn, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>hampei</i>)	930	comosus Wood, Corthylus	1072
coffae Schedl, Cnesinus	206	compactus (Eichhoff) , (<i>Xyleborus</i>), <i>Xylosandrus</i>	788
<i>coffae</i> Wurth, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>morigerus</i>)	795	compactus Lea, Cryphalus	875
coffeiceus (Schedl) , (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	672	comparabilis Schedl, Xyleborus	719
<i>coffeivorus</i> Weele, (<i>Xyleborus</i>), <i>Hypothenemus</i> (= <i>hampei</i>)	930	compressicornis (Fabricius) , (<i>Bostrichus</i>), <i>Corthylus</i>	1072
<i>cognatus</i> Blackman, <i>Cnesinus</i> (= <i>setulosus</i>)	211	compressus (Lea) , (<i>Xylopertha</i>), <i>Ambrosiodmus</i>	672
<i>cognatus</i> Blackman, <i>Pityophthorus</i> (= <i>cariniceps</i>)	986	comptus (Sampson) , (<i>Xyleborus</i>), <i>Eucallacea</i>	687
cognatus Blandford, Xyleborus	719	<i>comptus</i> Blackman, <i>Pityophthorus</i> (= <i>confertus</i>)	990
<i>cognatus</i> Wood, <i>Dendroterus</i>	950	concarifrons (Schedl) , (<i>Xylocleptes</i>), <i>Thamnurgus</i>	542
coimbatorensis Stebbing, Sphaerotrypes	199	concarus Blackman, Pityophthorus	988
colae (Schedl) , (<i>Stephanoderes</i>), <i>Hypothenemus</i>	912	concarus Bright, Corthylus	1072
colaphus Wood, Gnathotrypes	1040	centralis (Schedl) , (<i>Cryphalophilus</i>), <i>Scolytogenes</i>	860
colchicum Semenov, Liparthrum	276	centralis (Schedl) , (<i>Neodryocoetes</i>), <i>Araptus</i>	954
Coleobothrus Enderlein	615	centralis Eggers, Emiopus	855
collaris (Schedl) , (<i>Pocilips</i>), <i>Coccotrypes</i>	597	centralis Eichhoff, Pityophthorus	989
collaris Blandford, Amphicranus	1046	<i>centralis</i> Hopkins, <i>Trypophloeus</i> (= <i>salicis</i>)	847
collaris Blandford, Corthylus	1070	<i>centralis</i> Schedl, (<i>Ctenyophthorus</i>), <i>Araptus</i> (= <i>confluens</i>)	954
collaris Chapuis, Phloeotribus	219	centralis Schedl, Margadillius	895
<i>collaris</i> Eggers, (<i>Xyleborus</i>), <i>Eccoptopterus</i> (= <i>gracilipes</i>)	820	centralis Wood, Pseudothysanoes	413
collaris Schedl, Pityophthorus	988	concentus Wood, Xyleborus	719
collarti (Eggers) , (<i>Xyleborinus</i>), <i>Xyleborinus</i>	806	concinulus (Walker) , (<i>Hylurgus</i>), <i>Hyledius</i>	259
<i>collinus</i> Bright, <i>Pityophthorus</i> (= <i>toralis</i>)	1031	concinus (Mannerheim) , (<i>Bostrichus</i>), <i>Ips</i>	500
collis Niisima, Xyleborus	719	<i>concinus</i> Beeson, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>elegans</i>)	824
colombianus (Blackman) , (<i>Cryptocleptes</i>), <i>Pseudothysanoes</i>	412	concinus Wood, Pityophthorus	989
colombianus Wood, Cnesinus	206	concisus (Blandford) , (<i>Xyleborus</i>), <i>Leptoxyleborus</i>	659
colombianus Wood, Pseudopityophthorus	972	concisus (Wood) , (<i>Paracorthylus</i>), <i>Metacorthylus</i>	1064
<i>coloradensis</i> Wood, <i>Cryphalus</i> (= <i>ruficollis</i>)	892	concisus Wood, Corthylus	1072
colossus (Blandford) , (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	672	concisus Wood, Microcorthylus	1065
columbi Hopkins, Hypothenemus	912	concitatus (Schedl) , (<i>Xyleborus</i>), <i>Anasa</i>	682
columbianus (Schedl) , (<i>Hexacolus</i>), <i>Scolytodes</i>	390	concolor (Wollaston) , (<i>Aphanarthrum</i>), <i>Crypturgus</i>	623
columbianus (Schedl) , (<i>Neopityophthorus</i>), <i>Araptus</i>	954	concolor Hagedorn, Hypothenemus	913
columbianus Hopkins, Corthylus	1071	conditus Schedl, Xyleborus	719
<i>columbianus</i> Schedl, <i>Corthylus</i> (= <i>letzneri</i>)	1074	conditus Wood, Araptus	954
columbianus Schedl, Dendrocranius	550	conditus Wood, Dendrocranius	550
comans (Sampson) , (<i>Xyleborus</i>), <i>Hadrodemius</i>	819	confertus (Schedl) , (<i>Pocilips</i>), <i>Coccotrypes</i>	597
comatum (Eggers) , (<i>Eupteroxylon</i>), <i>Monarthrum</i>	1052	confertus Swaine, Pityophthorus	989
comatus (Zimmermann) , (<i>Crypturgus</i>), <i>Pityophthorus</i>	965	confinis (Blandford) , (<i>Pityophthorus</i>), <i>Araptus</i>	954
comatus Blandford, Corthylus	1071	confinis (Eggers) , (<i>Webbia</i>), <i>Arxyleborus</i>	666
		confinis (Wood) , (<i>Loganius</i>), <i>Cnemionyx</i>	314

- confinis* Eggers, (*Xyleborus*), *Eucallacca*
(=*icallacei*) 695
- confinis* LeConte, *Pityophthorus* 990
- confinis* Wood, *Chaetophloeus* 271
- confinis* Wood, *Dendrocraniulus* 551
- confinis* Wood, *Dendroterus* (= *mexicanus*) 951
- confluens* (Schedl), (*Ctenyophthorus*), *Araptus* 954
- confluens* Schedl, *Xyleborus* 719
- conformis* Dejean, *Ips* (= *calligraphus*) 494
- conformis* Koll, *Xyleborus* (= *ferrugineus*) 736
- confractus* Bright, *Pityophthorus* 990
- confragosus* (Sampson), (*Cryptalomorphus*),
Scolytogenes 860
- confusa* Hopkins, *Coptodryas* 823
- Confusoscolytus* Tsai & Huang (= *Scolytus*) 322
- confusum* Eggers, *Rhopalopselion* 96
- confusus* (Eggers), (*Hylcurosoma*), *Scolytodes* 390
- confusus* (Eggers), (*Kissophagus*), *Hylesinopsis* 93
- confusus* (Eggers), (*Poecilips*), *Coccotrypes* 597
- confusus* (Eggers), (*Stephanoderes*),
Hypothenemus 913
- confusus* (LeConte), (*Tomicus*), *Ips* 500
- confusus* Blandford, *Pityophthorus* 990
- confusus* Bright, *Pityophthorus* (= *toralis*) 1031
- confusus* Browne, *Cryptoxyleborus* 828
- confusus* Eggers, (*Eccoptogaster*), *Scolytus*
(= *japonicus*) 335
- confusus* Eggers, (*Mimips*), *Acanthotomicus*
(= *biconicus*) 479
- confusus* Eggers, *Cryphalus* 878
- confusus* Eggers, *Polygraphus* 286
- confusus* Eichhoff, *Xyleborus* (= *ferrugineus*) 738
- confusus* Hopkins, *Margadillius* 899
- confusus* Swaine, *Dryocoetes* 580
- congonum* (Schedl), (*Hapalogenius*), *Rhopalopselion* 96
- congonus* (Eggers), (*Dryocoetes*), *Cyrtogenius* 564
- congonus* (Eggers), (*Mimips*), *Acanthotomicus* 480
- congonus* (Schedl), (*Brachydendrilus*),
Mimicurus 949
- congonus* Eggers, (*Poecilips*), *Coccotrypes*
(= *rotundicollis*) 610
- congonus* Eggers, *Coccotrypes* 597
- congonus* Eggers, *Microcryphalus* 410
- congonus* Eggers, *Pityophthorus* 991
- congonus* Eggers, *Polygraphus* 286
- congonus* Eggers, *Sphaerotrypes* 199
- congonus* Hagedorn, (*Stephanoderes*),
Hypothenemus (= *setosus*) 944
- congonus* Hagedorn, (*Xyleborus*), *Ambrosiodmus*
(= *eichhoffi*) 673
- congonus* Hagedorn, *Xylocleptes* 549
- congonus* Nunberg, *Catenophorus* 237
- congonus* Schedl, *Caridoctonus* 283
- congonus* Schedl, *Scolytoplatypus* 401
- congruens* (Schedl), (*Mimips*), *Acanthotomicus* 480
- congruens* Schedl, *Xyleborus* 719
- conicolens* Wood, *Conophthorus* 963
- conidens* Browne, *Scolytus* 328
- conidens* Eggers, *Xyleborus* 720
- conifer* (Hagedorn), (*Xyleborus*), *Sampsonius* 655
- coniferae* (Wood), (*Aphanocleptus*),
Pseudothysanoes 413
- coniferae* Stebbing, (*Pityophthorus*), *Pityogenes*
(= *scitus*) 457
- coniperda* (Schwarz), (*Pityophthorus*),
Conophthorus 963
- conjunctus* (Schedl), (*Mimips*), *Acanthotomicus* 480
- conjunctus* Reitter, *Pityogenes* 451
- conjungens* Schedl, *Rhopalopselion* 96
- Conophthocraniulus* Schedl (= *Pityophthorus*) 977
- Conophthorus* Hopkins 963
- conradti* Hagedorn, *Xyleborus* 720
- conradti* Schedl, *Ctonoxylon* 537
- conscriptus* Bright, *Pityophthorus* 991
- consentaneus* Blandford, *Gnathotrichus* 1034
- consimile* (Blandford), (*Pterocyclon*),
Monarthrum 1052
- consimilis* (Eggers), (*Xyleborus*), *Ambrosiodmus* 672
- consimilis* LeConte, *Pityophthorus* 991
- consimilis* Wood, *Corthylus* 1072
- consimilis* Wood, *Dendrocraniulus* 551
- consobrinus* (Eichhoff), *Gnathotrypes* 1040
- consobrinus* Blandford, *Cosmoderes* 901
- consobrinus* Eggers, *Xyleborus* 720
- consobrinus* Wood, *Araptus* 954
- consocius* (Blandford), (*Problechilus*),
Gymnochilus 386
- conspiciens* Schedl, (*Xyleborus*), *Coccotrypes*
(= *cyperi*) 599
- conspicuous* (Schedl), (*Xyleborus*), *Ambrosiodmus* 672
- conspicuous* Wood, *Pityophthorus* 991
- constrictus* Schedl, (*Cryphalus*), *Hypocryphalus*
(= *perminimus*) 871
- constrictus* Schedl, *Gnathotrypes* (= *longipennis*) 1041
- constrictus* Schedl, *Hypocryphalus* (= *perminimus*) 871
- constrictus* Wood, *Scolytodes* 391
- consularis* (Schedl), (*Xyleborus*), *Amasa* 682
- continentalis* Eggers, *Xyleborus* 720
- contortae* Hopkins, *Conophthorus* (= *ponderosae*) 965
- contortus* Schedl, *Phloeotribus* 219
- contractus* (Chapuis), (*Hylastes*), *Xylechinus* 105
- contractus* Chapuis, *Phloeotribus* 220
- contractus* Schedl, *Cladoctonus* 238
- contractus* Wood, *Microcorthylus* 1065
- contractus* Wood, *Scolytodes* 391
- contrarius* Wood, *Pseudothysanoes* 413
- controversae* Murayama, (*Sphaerotrypes*), *Sueus*
(= *niisimai*) 191
- conversum* Wood, *Monarthrum* 1053
- convexa* Bright, *Schedlia* 822
- convexicauda* Eggers, *Corthylus* 1072
- convexicauda* Eggers, *Xyleborus* 720
- convexifrons* Hopkins, *Dendroctonus* (= *adjunctus*) 137
- convexifrons* Wood, *Carphoborus* 304

<i>convexifrons</i> Wood, <i>Corthylus</i>	1072	<i>corpulentus</i> Schedl, <i>Hypocryphalus</i>	565
<i>convexifrons</i> Wood, <i>Polygraphus</i>	286	<i>corpulentus</i> sundri Schedl, (<i>Margadillius</i>), <i>Enicoladus</i> (= <i>corpulentus</i>)	557
<i>convexus</i> (Schedl), (<i>Afronicracis</i>), <i>Micocryphalus</i>	410	<i>corrugatum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698
<i>convexus</i> Bright, <i>Pseudopityophthorus</i> (= <i>pruinus</i>)	975	<i>corrugatus</i> (Schedl), (<i>Stephanorhopalus</i>), <i>Scolytogenes</i>	860
<i>convexus</i> Schedl, <i>Scolytus</i>	328	<i>corrugatus</i> Swaine, <i>Crypturgus</i> (= <i>borealis</i>)	620
<i>cooki</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>hampei</i>)	930	<i>corruptus</i> (Schedl), (<i>Xyleborus</i>), <i>Prennobiis</i>	653
<i>Coptoborus</i> Hopkins	662	<i>corruptus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>adjunctus</i>)	662
<i>Coptodryas</i> Hopkins	822	<i>corruptus</i> Wood, <i>Pityophthorus</i>	992
<i>Coptodryas</i> Schedl (= <i>Cnemonyx</i>)	313	<i>corsicum</i> Eichhoff, <i>Liparthrum</i> (= <i>mori</i>)	279
<i>Coptogaster</i> Illiger (= <i>Scolytus</i>)	321	<i>corsicus</i> Eggers, (<i>Myelophilus</i>), <i>Tomicus</i> (= <i>minor</i>)	127
<i>Coptosomus</i> Schedl (= <i>Cnemonyx</i>)	313	<i>corsicus</i> Eggers, <i>Crypturgus</i> (= <i>cinereus</i>)	623
<i>coracinus</i> Wood, <i>Cnesinus</i>	206	<i>corsicus</i> Eggers, <i>Trypophloeus</i> (= <i>rybinskii</i>)	546
<i>coracinus</i> Wood, <i>Pseudothysanocs</i>	413	<i>corsicus</i> Guillebeau, (<i>Phloeophthorus</i>), <i>Phlocotribus</i> (= <i>pubifrons</i>)	227
<i>corculum</i> Wood, <i>Monarthrum</i>	1053	<i>corsicus</i> Guillebeau, <i>Phlocotribus</i> (= <i>cristatus</i>)	220
<i>cordatum</i> (Blandford), (<i>Pterocydon</i>), <i>Monarthrum</i>	1053	<i>cortezi</i> Bright, <i>Pityophthorus</i>	992
<i>cordatus</i> (Bright), (<i>Tricolus</i>), <i>Amphicranus</i>	1046	<i>Cortheloderes</i> Schedl (= <i>Corthylus</i> , = <i>gracilis</i>)	1074
<i>cordatus</i> Hagedorn, <i>Strombophorus</i>	202	<i>Corthyacydon</i> Schedl	1067
<i>cordatus</i> Hagedorn, <i>Xyleborus</i> (= <i>emarginatus</i>)	733	<i>Corthyli</i> LeConte (= <i>Corthylini</i>)	949, 1034
<i>cordicticum</i> Wood, <i>Monarthrum</i>	1053	<i>Corthyidae</i> LeConte (= <i>Corthylini</i>)	949
<i>cordipennis</i> Lea, <i>Hylesinus</i>	77	<i>corthyiformis</i> Schedl, (<i>Gnathotrichus</i>), <i>Gnathotrupes</i> (= <i>longipennis</i>)	1041
<i>Coriacephilus</i> Schedl	553	<i>Corthylina</i> LeConte, part (= <i>Corthylini</i>)	1034
<i>coriaceus</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Coriacephilus</i>	854	<i>Corthylinae</i> LeConte (= <i>Corthylini</i>)	949
<i>coriaceus</i> Schedl, (<i>Erioschidias</i>), <i>Cosmoderes</i> (= <i>monilicollis</i>)	901	<i>Corthylini</i> LeConte	949
<i>corni</i> Kurenzov, <i>Hypothenemus</i>	913	<i>Corthylocurus</i> Wood	1065
<i>corniculatulus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>respatorius</i>)	665	<i>corthyloides</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	790
<i>corniculatus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>respatorius</i>)	665	<i>corthyloides</i> Eichhoff, <i>Gnathotrichus</i> (= <i>materiarius</i>)	1036
<i>corniger</i> Wood, <i>Chramesus</i>	264	<i>corthyloides</i> Hagedorn, <i>Prennobiis</i>	653
<i>cornicornus</i> Murayama, <i>Xyleborus</i>	720	<i>corthyloides</i> Schedl, (<i>Gnathotrichus</i>), <i>Gnathotrupes</i> (= <i>longipennis</i>)	1041
<i>cornutum</i> Eggers, <i>Ctenoxylon</i>	837	<i>Corthylomimus</i> Ferrari (= <i>Monarthrum</i>)	1050
<i>cornutus</i> Axentjev, <i>Cynanchophagus</i>	1083	<i>Corthylomimus</i> Schedl (= <i>Corthylus</i>)	1069
<i>cornutus</i> Schaufuss, <i>Xyleborus</i>	720	<i>Corthylus</i> Erichson	1069
<i>cornutus</i> Schedl, <i>Webbia</i>	830	<i>corticalis</i> Eichhoff, <i>Pityophthorus</i>	992
<i>cornutus</i> Wood, <i>Cnesinus</i>	207	<i>corticinus</i> Wood, <i>Hylurdretonus</i>	186
<i>coronarius</i> Blackman, <i>Pityophthorus</i>	992	<i>corticiperda</i> Erichson, <i>Hylastes</i> (= <i>linearis</i>)	56
<i>coronatus</i> (Chapuis), (<i>Phloeosinus</i>), <i>Chaetophloeus</i>	271	<i>corticus</i> Beeson, (<i>Thammurgides</i>), <i>Coccotrypes</i> (= <i>nubilus</i>)	608
<i>coronatus</i> Eggers, <i>Corthylus</i>	1072	<i>corumbensis</i> (Eggers), (<i>Hoplites</i>), <i>Cladoctonus</i>	238
<i>coronatus</i> Eggers, <i>Polygraphus</i>	286	<i>corumbensis</i> Eggers, <i>Chramesus</i>	264
<i>coronatus</i> Eggers, <i>Taphroxyelus</i>	559	<i>coryli</i> (Perris), (<i>Tomicus</i>), <i>Lymantor</i>	555
<i>coronatus</i> Eichhoff, <i>Xyleborus</i> (= <i>spathipennis</i>)	774	<i>coryli</i> Stark, <i>Cryphalus</i>	878, 884
<i>coronatus</i> Wood, <i>Cryptocarenum</i> (= <i>brevicollis</i>)	902	<i>Cosmocorynus</i> Ferrari (= <i>Monarthrum</i>)	1050
<i>corporaali</i> (Eggers), (<i>Xyleborus</i>), <i>Coptodryas</i>	823	<i>Cosmoderes</i> Eichhoff	900
<i>corpulentus</i> (Sampson), (<i>Cryphalus</i>), <i>Enicoladus</i>	557	<i>cosmoderoides</i> Murayama, <i>Hypothenemus</i>	913
<i>corpulentus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	860	<i>costabilis</i> Wood, <i>Pityophthorus</i>	992
<i>corpulentus</i> (Schedl), (<i>Erioschidias</i>), <i>Cosmoderes</i>	901	<i>costabilis</i> Wood, <i>Scolytodes</i>	391
<i>corpulentus</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	954	<i>costalimai</i> Blackman, <i>Pityophthorus</i>	992
<i>corpulentus</i> Browne, <i>Coccotrypes</i>	598	<i>costalimai</i> Numberg, (<i>Enicophloeus</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	926
<i>corpulentus</i> Eggers, <i>Xyleborus</i>	720	<i>costalimai</i> Schedl, <i>Dendrocramulus</i>	551
° <i>corpulentus</i> Schedl, <i>Hylurgops</i>	30	<i>costaricensis</i> (Schedl), (<i>Microcorthylus</i>), <i>Corthylocurus</i>	1069

- costaricensis* (Schedl), (*Neodryocoetes*), *Araptus* . . . 954
costaricensis Blandford, *Xyleborus* . . . 720
costaricensis Nunberg, *Sampsonius* (= *dampfi*) . . . 655
costaricensis Schedl, *Dendrocraniulus* . . . 551
costaricensis Wood, *Micracis* (= *grandis*) . . . 432
costata (Schedl), (*Bothryperus*), *Phrixosoma* . . . 189
costatomorphus Schedl, *Xyleborus* . . . 721
costatulus Wood, *Pityophthorus* . . . 992
costatum (Eggers), (*Brachyspartus*),
Monarthrum . . . 1053
costatus (Chapuis), (*Bothrosternus*),
Sternobothrus . . . 216
costatus (Eichhoff), (*Stephanoderes*),
Hypothenemus . . . 913
costatus Blandford, *Hylesinus* . . . 77
costatus Chapuis, *Camptocerus* . . . 319
costatus Wichmann, *Carphoborus* . . . 305
costatus Wood, *Pityophthorus* . . . 992
costatus Wood, *Pseudothyanoes* . . . 413
costatus Wood, *Sphaerotrypes* . . . 199
costellatus Chapuis, *Scolytus* . . . 328
costiceps Broun, *Dendrotrypes* . . . 106
costifera Bright, *Pityophthorus* . . . 992
costipennis (Schedl), (*Xyleborus*), *Coptodryas* . . . 823
costipennis Schedl, *Scolytomimus* . . . 539
costulatus Schedl, *Webbia* . . . 830
costulatus Blackman, *Pseudochramesus* . . . 262
costulatus Blandford, *Cnesinus* . . . 207
costulatus Eggers, *Xyleborus* . . . 721
covinus Reitter, *Xyleborus* (= *pfeili*) . . . 762
cracens (Wood), (*Gnathophthorus*),
Dacnophthorus . . . 969
cracensis Bright, *Pityophthorus* . . . 992
cracensis Wood, *Liparthrum* . . . 276
Craniodicticus Blandford . . . 437
crassinus Wood, *Pseudothyanoes* . . . 413
crassitarsus Schedl, (*Xyleborus*), *Coptodryas*
 (= *recidens*) . . . 827
crassiusculus (Motschulsky), (*Phloeotrogus*),
Xylosandrus . . . 790
crassiventris (Schedl), (*Pocilips*), *Coccotrypes* . . . 598
crassulus Wood, *Araptus* . . . 954
crassus (Eggers), (*Dryocoetes*), *Peridryocoetes* . . . 561
crassus (Schedl), (*Thamnophtorus*), *Araptus* . . . 954
crassus Beeson, (*Hylesinus*), *Ficicis* (= *despectus*) . . . 91
crassus Blackman, *Pityophthorus* . . . 992
crassus Hagedorn, (*Xyleborus*), *Ambrosiodmus*
 (= *minor*) . . . 676
crassus Wood, *Araptus* (= *crassulus*) . . . 954
crassus Wood, *Scolytodes* . . . 391
craterigerus (Nunberg), (*Orthotomicus*),
Acanthotomicus . . . 480
crebellus Reitter, *Crypturgus* (= *pusillus*) . . . 629
creber (Schedl), (*Cryphalomorphus*),
Scolytogenes . . . 8560
creber (Schedl), (*Pocilips*), *Coccotrypes* . . . 598
creber Schedl, *Cnemonyx* . . . 314
creber Schedl, *Polygraphus* . . . 286
crebra Wood, *Phrixosoma* . . . 189
crecentus Wood, *Gnathotrupes* . . . 1040
crenatulus Duftschmidt, (*Hylesinus*), *Hylastinus*
 (= *obscurus*) . . . 67
crenatum Hagedorn, *Ctonoxylon* . . . 537
crenatus (Fabricius), (*Bostrichus*), *Hylesinus* . . . 77
crenatus (Guillebeau), (*Elzearius*), *Phlocotribus* . . . 220
crenatus (Sampson), (*Cryphalomorphus*),
Scolytogenes . . . 860
crenatus Eggers, (*Xyleborus*), *Dryocoetoides*
 (= *cristatus*) . . . 657
crenatus Hagedorn, *Strombophorus* . . . 203
crenatus Olivier, (*Scolytus*), *Hylastinus* (= *obscurus*) . . . 66
crenatus Wood, *Chramesus* . . . 264
crenulatus Eggers, *Xyleborus* . . . 721
cressatyi Bruck, *Carphoborus* (= *blaisdelli*) . . . 303
cribratus (Blandford), (*Phloeosinus*), *Hyledius* . . . 260
cribratus Blackman, *Pityophthorus* (= *solus*) . . . 1029
cribratus Browne, *Xylocleptes* . . . 549
cribratus Eggers, *Xyleborus* . . . 721
cribratus Pfeffer, *Pityophthorus* (= *pityographus*) . . . 1021
cribrellus Schedl, *Lanurgus* . . . 408
cribricollis (Schedl), (*Neodryocoetes*), *Araptus* . . . 955
cribricollis Eichhoff, (*Tomicus*), *Ips*
 (= *grandicollis*) . . . 508
cribricollis Eichhoff, *Pagiocerus* . . . 213
cribricollis Schedl, *Cyrtogenius* . . . 564
cribripennis (Schedl), (*Neodryocoetes*), *Araptus* . . . 955
cribripennis Eggers, *Taphrorychus* (= *ceratoniae*) . . . 559
cribripennis Eggers, *Xyleborus* . . . 721
cribripennis Eichhoff, *Pityophthorus* (= *pullus*) . . . 1025
cribripennis Schedl, *Coriacephilus* . . . 854
cribripennis Schedl, *Cyrtogenius* . . . 564
criddlei (Swaine), (*Leperisimus*), *Hylesinus* . . . 79
criddlei Swaine, (*Pseudocryphalus*), *Chactophloeus*
 (= *heterodoxus*) . . . 272
crinalis Blackman, *Pityophthorus* . . . 993
crinalis Wood, *Scolytodes* . . . 391
crinitulus Wood, *Xyleborus* . . . 721
crinitus Blackman, *Hyllocurus* (= *hirtellus*) . . . 425
crinitus Schedl, *Xyleborus* . . . 721
crinitus Wood, *Scolytodes* . . . 391
cristatuloides Schedl, (*Xyleborus*), *Ambrosiodmus*
 (= *asperatus*) . . . 671
cristatulus (Schedl), (*Xyleborus*), *Ambrosiodmus* . . . 672
cristatum (Ferrari), (*Cosmocorynus*),
Monarthrum . . . 1053
cristatus (Fabricius), (*Bostrichus*), *Dryocoetoides* . . . 657
cristatus (Fauvel), (*Phloeophthorus*), *Phlocotribus* . . . 220
cristatus (LeConte), (*Hylesinus*), *Phloeosinus* . . . 244
cristatus (Schedl), (*Phacrylus*), *Neocryphus* . . . 552
cristatus Chapuis, *Phlocoborus* . . . 98
cristatus Hagedorn, (*Xyleborus*), *Ambrosiodmus*
 (= *apicalis*) . . . 671
cristatus Inouye & Nobuchi, *Dryocoetes* . . . 552

<i>cristatus</i> Mannerheim, (<i>Hylastes</i>), <i>Hylurgops</i> (= <i>subcostulatus</i>)	42	Crypturgi LeConte (=Crypturgini)	615
<i>cristatus</i> Schedl, <i>Chranesius</i> (= <i>spinosus</i>)	269	Crypturgidae LeConte (=Crypturgini)	615
<i>cristatus</i> Wood, <i>Carphodicticus</i>	437	Crypturginae LeConte (=Crypturgini)	615
<i>cristatus</i> Wood, <i>Phlococleptus</i>	421	Crypturgini LeConte	615
<i>cristatus</i> Wood, <i>Pityophthorus</i>	992	Crypturgina LeConte (=Crypturgini)	615
<i>cristatus</i> Wood, <i>Scolytus</i>	328	Crypturgus Erichson	620
<i>cristatus</i> Wood, <i>Sphaerotrypes</i>	199	<i>esikii</i> Endrodi, <i>Thamnurgus</i>	542
<i>criticus</i> (Schedl), (<i>Cryphalus</i>), <i>Hypothenemus</i>	913	Ctenophoridae Chapuis (=Ctenophorini)	383
<i>criticus</i> Eichhoff, <i>Hylastes</i> (= <i>tenuis</i>)	63	Ctenophorini Chapuis	383
<i>criticus</i> Schedl, <i>Xyleborus</i> (= <i>perforans</i>)	762	Ctenophorus Chapuis (= <i>Scolytodes</i>)	387
<i>crotonis</i> Wood, <i>Hyllocurus</i>	423	Ctenyophthorus Schedl (= <i>Pityophthorus</i>)	977
<i>crotonis</i> Wood, <i>Pityophthorus</i>	992	Ctonocryphus Schedl (= <i>Glostatus</i>)	835
<i>cruciatus</i> Schedl, <i>Xyleborus</i>	721	Ctonoxylon Hagedorn	836
<i>crucifer</i> (Hagedorn), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	698	<i>cubanus</i> Eggers, (<i>Thamnurgides</i>), <i>Coccotrypes</i> (= <i>advena</i>)	592
<i>crucifer</i> <i>cruciferinum</i> Schedl, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>crucifer</i>)	699	<i>cubensis</i> (Blackman), (<i>Neodryocotes</i>), <i>Araptus</i>	955
<i>cruciferinum</i> Schedl, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>crucifer</i>)	699	<i>cubensis</i> (Schedl), (<i>Hexacohus</i>), <i>Scolytodes</i>	391
<i>cruciforme</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	699	<i>cubensis</i> (Wood), (<i>Hoplitophthorus</i>), <i>Cladoctonus</i>	238
<i>crucipenne</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	699	<i>cubensis</i> Blackman, <i>Cucinus</i>	207
<i>crudiae</i> (Panzer), (<i>Bostrichus</i>), <i>Hypothenemus</i>	914	<i>cubensis</i> Blackman, <i>Micracis</i>	432
<i>cruralis</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	802	<i>cubeus</i> Bright, <i>Chaetophloeus</i>	272
<i>cruralis</i> Eggers, <i>Cnestus</i> (= <i>cruralis</i> Schedl)	802	<i>cubensis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>erectus</i>)	918
Cryphali Lindemann (=Cryphalini)	842	<i>cubensis</i> Schedl, <i>Pityophthorus</i> (= <i>pulicarius</i>)	1024
Cryphalidae Lindemann (=Cryphalini)	842	<i>cubensis</i> Wood, <i>Scolytopsis</i> (= <i>puncticollis</i>)	321
Cryphalina Lindemann (=Cryphalini)	842	<i>cucullatus</i> Blandford, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>brevis</i>)	788
Cryphalinae Lindemann (=Cryphalini)	842	<i>cucurbitae</i> (LeConte), (<i>Xylocleptes</i>), <i>Dendrocraniulus</i>	551
Cryphalini Lindemann	842	<i>cucurbitae</i> Schedl, <i>Dendrocraniulus</i> (= <i>schedli</i>)	553
Cryphalites Cockerell	1082	<i>culcitatus</i> (Blandford), (<i>Epomadius</i>), <i>Scolytodes</i>	391
Cryphalogenes Wood	857	<i>culminicola</i> Bright, <i>Pityophthorus</i>	993
Cryphaloideae Lindemann (=Cryphalini)	842	<i>Cumatotomicus</i> Ferrari (= <i>Ips</i>)	484
cryphuloides (Eichhoff), (<i>Triamocerus</i>), <i>Hypothenemus</i>	916	<i>cuneatus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Coptoborus</i>	663
Cryphuloides Formenek (= <i>Coccotrypes</i>)	591	<i>cuneidentis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	806
<i>cryphaloides</i> Schedl, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>confusa</i>)	823	<i>cuneiformis</i> Schedl, <i>Xyleborus</i>	722
Cryphalominetes Browne (= <i>Stephanopodius</i>)	853	<i>cuneipennis</i> Schedl, <i>Xyleborus</i>	722
Cryphalominus Browne (= <i>Stephanopodius</i>)	853	<i>cuneolosus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	806
Cryphalominus Eggers	835	<i>cuneolus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	916
Cryphalomorphus Schanfuss (= <i>Scolytogenes</i>)	858	<i>cuneolus</i> Eggers, <i>Xyleborus</i>	722
<i>cryphalomorphus</i> Schedl, <i>Hypothenemus</i> (= <i>brunneus</i>)	911	<i>cunicularius</i> Erichson, <i>Hylastes</i>	51
Cryphalophilus Schedl (= <i>Scolytodes</i>)	858	<i>cupressi</i> Hopkins, <i>Phlocosinus</i>	245
Cryphalops Reitter (= <i>Ernoporus</i>)	854	<i>cupressi</i> Wood, <i>Carphobius</i>	311
Cryphalus Erichson	872	<i>cupulatus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	806
Cryphyophthorus Schedl	274	<i>curiosus</i> Bright, <i>Corythus</i>	1072
<i>Cryptarthrum</i> Blandford (= <i>Cryphalus</i>)	872	<i>curtidentis</i> Schedl, <i>Xyleborus</i>	722
Cryptocareus Eggers	902	<i>curtipennis</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	916
<i>Cryptocleptes</i> Blackman, <i>Pseudothysanoes</i>	411	<i>curtuloides</i> Eggers, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>curtulus</i>)	793
Cryptocorus Schedl	72	<i>curtulus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xylosandrus</i>	793
cryptographus (Ratzeburg), (<i>Bostrichus</i>), <i>Xyleborus</i>	721	<i>curtulus</i> Sokanovskii, <i>Pityophthorus</i> (= <i>micrographus</i>)	1012
<i>cryptolepis</i> Schedl, <i>Scolytogenes</i> (= <i>durivini</i>)	861	<i>curtum</i> Wollaston, <i>Liparthrum</i>	276
<i>cryptomeriae</i> Nüßima, <i>Cryphalus</i>	878	<i>curtus</i> (Eggers), (<i>Kissoplagus</i>), <i>Phlocoditica</i>	236
Cryptoxyleborus Schedl	828	<i>curtus</i> (Schedl), (<i>Ozodendron</i>), <i>Cyrtogenius</i>	564
<i>Cryptuloceptus</i> Wood (= <i>Pseudothysanoes</i>)	411		

- curtus* Bright, *Pseudopityophthorus* (= *declivis*) 972
- curtus* Eggers, (*Thamnuroides*), *Coccotrypes*
(= *myristicae*) 606
- curtus* Eggers, *Xyleborus* (= *spathipennis*) 774
- curtus* Wood, *Microcorthyus* 1065
- curvatus* Browne, *Xyleborus* 722
- curridens* (Germar), (*Tomicus*), *Pityokteines* 460
- curridens* (Schedl), (*Brachydendrus*),
Acanthotomicus 480
- curridens heterodon* Wachtl, (*Tomicus*), *Pityokteines*
(= *spinidens*) 466
- curridentis* (Schedl), (*Xyleborus*), *Coptodryas* 824
- currifer* (Walker), (*Hylesinus*), *Dianerus* 195
- curripenne* (Schedl), (*Xyleborus*), *Cyclorhipidion* 699
- curviventralis* Niisima, *Scolytus* (= *chikisanii*) 327
- cuspidatus* Blackman, *Pityophthorus* 993
- cuspidatus* Eggers, *Hylocurus* 423
- cuspidis* Wood, *Araptus* (= *tenellus*) 962
- cuspidis* Wood, *Pseudothysanoes* 413
- cuspidus* (Schedl), (*Xyleborus*), *Taphrolasus* 827
- cutleri* Swaine, *Pityophthorus* (= *murrayanae*) 1014
- cyclopus* (Schedl), (*Pelicerus*), *Cyrtogenius* 564
- cyclopus* Schedl, *Xyleborus* 723
- Cyclorhipidion* Hagedorn 697
- cylihraceus* (Schedl), (*Stephanoderes*),
Hypothenemus 916
- cylicridica* (Eggers), (*Xyleborus*), *Coptodryas* 824
- cylicricollis* Eggers, *Crypturgus* (= *pusillus*) 629
- cylicricus* (Eggers), (*Poecilips*), *Coccotrypes* 598
- cylicricus* (Eggers), (*Xylocleptes*), *Thamnuroides* 542
- cylicricus* (Schedl), (*Cryphalomorphus*),
Scolytogenes 860
- cylicricus* (Schedl), (*Erioschidias*), *Cosmoderes* 901
- cylicricus* Browne, *Cryphalus* 878
- cylicricus* Eggers, *Ozopemon* (= *obanus*) 590
- cylicricus* Hopkins, (*Stephanoderes*),
Hypothenemus (= *plumeriae*) 938
- cylicricus* Schedl, (*Corthylius*), *Corthylocurus*
(= *mexicanus*) 1069
- cylicricus* Schedl, *Chramesus* 264
- cylicricus* Schedl, *Coccotrypes* (= *robustus*) 610
- cylicricus* Schedl, *Hypothenemus* (= *eruditus*) 924
- cylicricus* Schedl, *Phloeotribus* 220
- cylicricus* Schedl, *Scolytodes* 391
- cylicricus* Schedl, *Webbia* 830
- cylicriformis* (Schedl), (*Xyleborus*), *Amasa* 682
- cylicripennis* Schedl, (*Xyleborus*), *Coptodryas*
(= *myristicae*) 825
- cylicripennis* Schedl, *Hypocryphalus* 868
- cylicripennis* Schedl, *Hypothenemus* (= *eruditus*) 925
- cylicromorphus* Eggers, *Xyleborus* 723
- cylicrotomicus* (Schedl), (*Pseudoxyleborus*),
Xylosandrus 793
- Cylicrotomicus* Eggers (= *Scolytogenes*) 858
- cylicrus* (Browne), (*Dacryphalus*),
Hypocryphalus 868
- cylicrus* Browne, *Cryphalus* 878
- cylicrus* Schedl, *Xyleborus* 723
- Cynanchophagus* Axentjev 1082
- cyuometrae* Schedl, *Hypothenemus* 916
- cyperi* (Beeson), (*Thamnuroides*), *Coccotrypes* 598
- Cyrtogenius* Strohmeyer 562
- Cyrtotomicus* Ferrari (= *Ips*) 484
- cytisi* Eggers, *Liparthrum* (= *genistae*) 277
- Dacnophthorus* Wood 969
- Dacryophthorus* Schedl (= *Liparthrum*) 274
- Dacryostactus* Schaufuss 281
- Dacryphalus* Hopkins (= *Hypocryphalus*) 868
- Dactylipalpi* (= *Hylesinini*) 64
- Dactylipalpus* Chapuis 101
- dactyliperda* (Fabricius), (*Bostrichus*),
Coccotrypes 599
- Dactylopselaphus* Gemminger & Harold
(= *Dactylipalpus*) 101
- Dactylotrypes* Eggers 554
- dahuricus* Chapuis, *Scolytus* 328
- daimio* Blandford, *Scolytotylatus* 402
- dainichiensis* (Murayama), (*Xyloterus*),
Indocryphalus 648
- dalbergiae* Eggers, (*Xyleborus*), *Euwallacea*
(= *sibsagaricus*) 693
- dalmatinum* Eggers, *Liparthrum* (= *mori*) 279
- dalmatinus* Eggers, *Phloeotribus* (= *scarabaeoides*) 230
- dampfi* Schedl, *Sampsonius* 655
- danicus* Eggers, *Crypturgus* (= *pusillus*) 629
- daosi* (Schedl), (*Xyleborus*), *Cyclorhipidion* 699
- darjeelingensis* Schedl, *Xylechinus* 115
- darjeelingi* Stebbing, *Scolytotylatus* 402
- darwinensis* Schedl, (*Stephanoderes*), *Hypothenemus*
(= *seriatus*) 943
- darwini* Eichhoff, *Scolytogenes* 860
- darwini* Schedl, *Scolytomimus* (= *dilutus*) 839
- darwini* Schedl, *Xyleborus* 723
- dasyurus* (Browne), (*Xyleborus*), *Amasa* 683
- dasyurus* Browne, *Webbia* 830
- dearmatus* Eggers, *Xyleborus* (= *subcostulatus*) 776
- debilis* Chapuis, (*Hylesinus*), *Ptelcobius* (= *cittatus*) 70
- debilis* Wood, *Corthylocurus* 1069
- debilis* Wood, *Microcorthyus* 1065
- debilis* Wood, *Pityophthorus* 992
- decellei* (Nunberg), (*Trypographus*), *Hylesinopsis* 93
- decellei* Browne, *Tiarophorus* 539
- deceptus* Schedl, *Arixyleborus* 666
- decipiens* (LeConte), (*Xylocleptes*), *Lymantor* 556
- decipiens* Browne, (*Poecilips*), *Coccotrypes*
(= *papuanus*) 608
- decipiens* Eggers, (*Xyleborus*), *Cyclorhipidion*
(= *pruinotum*) 701
- decipiens* Wood, *Dendroterus* 951
- decliridepressus* Schedl, *Glostatus* 835
- decligramulatus* Schedl, (*Xyleborus*), *Xylosandrus*
(= *crassiusculus*) 792
- declivis* (Schedl), (*Carposinus*), *Cyrtogenius* 564
- declivis* Eggers, *Prennobius* 653
- declivis* Eichhoff, *Xyleborus* 723

<i>declivis</i> Reitter, <i>Thamurgus</i> (= <i>kaltenbuchi</i>)	544	<i>Dendrodicticus</i> Schedl	437
<i>declivis</i> Sampson, <i>Coccotrypes</i>	602	<i>Dendrographus</i> Schedl	561
<i>declivis</i> Schedl, <i>Dendrocranulus</i>	551	<i>Dendrosinus</i> Chapuis	236
<i>declivis</i> Wood, <i>Carphoborus</i>	305	<i>Dendroterus</i> Blandford	950
<i>declivis</i> Wood, <i>Pseudopityophthorus</i>	972	<i>Dendrotrupes</i> Broun	106
<i>declivisetosus</i> Bright, <i>Pityophthorus</i>	993	<i>Dendrotrypanum</i> Schedl (= <i>Indocryphalus</i>)	648
<i>declivispinatus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	672	<i>Dendrurgus</i> Eggers (= <i>Coccotrypes</i>)	591
<i>declivistriatus</i> Schedl, <i>Scolytodes</i>	391	<i>denotatus</i> Wood, <i>Cnesinus</i>	207
<i>decor</i> Boieldieu, (<i>Tomicus</i>), <i>Xyleborinus</i> (= <i>saxeseni</i>)	514	<i>densatus</i> Schedl, <i>Xyleborus</i>	723
<i>decorulus</i> Wood, <i>Araptus</i>	955	<i>densepilosus</i> Schedl, <i>Cryphalus</i>	878
<i>decorus</i> (Bright), (<i>Neodryocotes</i>), <i>Araptus</i>	955	<i>densepilosus</i> Schedl, <i>Hypocryphalus</i>	868
<i>decorus</i> Schedl, (<i>Xyleborus</i>), <i>Webbia</i> (= <i>micrographus</i>)	531	<i>densepunctatus</i> Schedl, <i>Xyleceptes</i>	549
<i>decorus</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>micrographus</i>)	509	<i>denseseriatus</i> Eggers, <i>Xyleborus</i>	723
<i>decorus</i> Wood, <i>Araptus</i> (= <i>decorulus</i>)	955	<i>densicornis</i> Schedl, <i>Xyleborus</i>	723
<i>decorus</i> Wood, <i>Scolytodes</i>	391	<i>dentatulum</i> Wood, <i>Monarthrum</i>	1053
<i>decretus</i> Eichhoff, (<i>Tomicus</i>), <i>Orthotomicus</i> (= <i>caelatus</i>)	465	<i>dentatulus</i> Browne, <i>Xyleborus</i>	723
<i>decumans</i> Schedl, <i>Xyleborus</i>	723	<i>dentatum</i> (Eggers), (<i>Amphicranus</i>), <i>Monarthrum</i>	1053
<i>decumanus</i> Erichson, (<i>Hylesinus</i>), <i>Hyhurgops</i> (= <i>glabratus</i>)	33	<i>dentatum</i> Eggers, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>denticulatum</i>)	1053
<i>defectus</i> Wood, <i>Dendroterus</i>	951	<i>dentatum</i> Eggers, <i>Monarthrum</i> (= <i>dentatulum</i>)	1053
<i>defensus</i> Blandford, <i>Xyleborus</i>	723	<i>dentatum</i> Numberg, <i>Rhopalopselion</i>	96
<i>definitus</i> Wood, <i>Bothrosterus</i>	215	<i>dentatus</i> (Say), (<i>Hyhurgus</i>), <i>Phloeosinus</i>	246
<i>deformatus</i> Browne, <i>Xyleborus</i>	723	<i>dentatus</i> Blandford, <i>Xyleborus</i>	724
<i>degener</i> Lindberg, <i>Liparthrum</i> (= <i>nigrescens</i>)	279	<i>dentatus</i> Bright, <i>Scolytus</i>	329
<i>degener</i> Wood, <i>Cnesinus</i>	207	<i>dentatus</i> Eggers, <i>Corthylus</i>	1073
<i>degener</i> Wood, <i>Pityophthorus</i>	993	<i>dentatus</i> Eggers, <i>Webbia</i>	530
<i>dejevi</i> Eggers, <i>Trypophloeus</i> (= <i>dejevi</i> Stark)	544	<i>dentatus</i> Schaeffer, <i>Chramesus</i>	264
<i>dejevi</i> Stark, <i>Trypophloeus</i>	544	<i>dentatus</i> Schedl, <i>Acanthotomicus</i>	480
<i>deleoni</i> (Blackman), (<i>Myeloborus</i>), <i>Pityophthorus</i>	993	<i>dentatus</i> Sturm, <i>Ips</i> (= <i>pini</i>)	517
<i>deleoni</i> Blackman, <i>Gnathotrichus</i>	1034	<i>dentatus</i> Wood, <i>Gnathotrichus</i>	1034
<i>deleoni</i> Blackman, <i>Phloeosinus</i>	245	<i>dentellus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	506
<i>deleoni</i> Bright, <i>Pityophthorus</i> (= <i>californicus</i>)	986	<i>dentibarbis</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>forficuloides</i>)	807
<i>deletus</i> LeConte, <i>Pityophthorus</i>	993	<i>denticollis</i> Schedl, <i>Hylocurus</i>	423
<i>delicatum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	699	<i>denticulatum</i> Wood, <i>Monarthrum</i>	1053
<i>delicatus</i> (Schedl), (<i>Paraglostatus</i>), <i>Glostatus</i>	535	<i>denticulatus</i> (Wichmann), (<i>Trigonogenius</i>), <i>Pityophthorus</i>	994
<i>delicatus</i> Schedl, <i>Hypothenemus</i>	916	<i>denticulatus</i> Blackman, <i>Gnathotrichus</i>	1035
<i>delicatus</i> Wood, <i>Araptus</i>	955	<i>denticulatus</i> Browne, <i>Webbia</i>	530
<i>delicatus</i> Wood, <i>Pityophthorus</i>	994	<i>denticulatus</i> Eggers, (<i>Pityophthorus</i>), <i>Araptus</i> (= <i>eggersianus</i>)	956
<i>delphinii</i> (Rosenhauer), (<i>Bostrichus</i>), <i>Thamurgus</i>	542	<i>denticulatus</i> Sturm, <i>Orthotomicus</i> (= <i>laricis</i>)	469
<i>demaioni</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>kirschii</i>)	337	<i>denticulatus</i> Wood, <i>Chramesus</i>	265
<i>demissus</i> Blandford, <i>Phloeotribus</i>	220	<i>denticulosus</i> Sokanovskii, <i>Hylesinus</i> (= <i>tupolevi</i>)	85
<i>demissus</i> Blackman, <i>Pityophthorus</i> (= <i>absonus</i>)	978	<i>denticulus</i> (Eggers), (<i>Ips</i>), <i>Acanthotomicus</i>	480
<i>demissus</i> Wood, <i>Chramesus</i>	264	<i>denticulus</i> Motschulsky, (<i>Anodius</i>), <i>Xyleborus</i> (= <i>perforans</i>)	761
<i>demissus</i> Wood, <i>Microcorthylus</i>	1065	<i>denticulus</i> Wood, <i>Pseudopityophthorus</i>	973
<i>demissus</i> Wood, <i>Xyleborus</i>	723	<i>dentifrons</i> (Blackman), (<i>Phthorophloeus</i>), <i>Phloeotribus</i>	221
<i>Dendriops</i> Schedl (= <i>Cosmoderes</i>)	900	<i>dentifrons</i> Blackman, <i>Pityophthorus</i>	994
<i>Dendrochilus</i> Schedl	435	<i>dentifrons</i> Wood, <i>Araptus</i>	955
<i>Dendrocranulus</i> Schedl	550	<i>dentiger</i> LeConte, <i>Monarthrum</i> (= <i>dentigerum</i>)	1053
<i>Dendroctonides</i> Nusslin (= <i>Tomicini</i>)	103	<i>dentigerum</i> (LeConte), (<i>Cryphalus</i>), <i>Monarthrum</i>	1053
<i>Dendroctonus</i> Erichson	136	<i>dentigerum</i> Schedl, <i>Ctonoxylon</i>	837
		<i>dentipennis</i> Browne, <i>Xyleborus</i>	724

- dentipes* Blandford, *Eupagiocerus* 214
dentipes Schedl, *Chramesus* 265
deodara (Stebbing), (*Cryphalus*), *Pityophthorus* 994
deodara Stebbing, *Scolytus* (=major) 340
deperditus Wood, *Cnecinus* 207
deplanatulus Schedl, *Xyleborus* 724
deplanatus Eggers, *Chramesus* 265
deplanatus Eggers, *Xyleborus* 724
deprecator (Schedl), (*Pachynoderes*),
Hypothenemus 916
deprecator Schaufuss, *Pityophthorus* 995
depressurus Browne, *Xyleborus* 724
depressus (Browne), (*Mimidendrulus*),
Cyrtogenius 564
depressus (Eggers), (*Thamnurgides*), *Coccotrypes* 602
depressus (Eggers), (*Xyleborus*), *Leptoxyleborus* 660
depressus Bright, *Cactopinus* 435
depressus Eichhoff, (*Stephanoderes*), *Hypothenemus*
 (=setosus) 944
depygis Blackman, *Pityophthorus*
 (=murrayanae) 1014
derelictus Hagedorn, *Xyleborus* 724
Dermestes (=Scolytidae) 25
Deropria Enderlein 620
derupteterminatus (Schedl), (*Xyleborus*),
Xylosandrus 793
deruptulus (Schedl), (*Xyleborus*), *Xylosandrus* 794
desectus (Eggers), (*Xyleborus*), *Ambrosiodmus* 672
desertus Bright, *Cactopinus* 435
desertus Schedl, *Xyleborus* 724
despectus (Walker), (*Hylesinus*), *Ficicis* 90
despectus Schedl, *Phlocotribus* 221
despectus Schedl, *Xyleborus* 724
destrictum (Schedl), (*Xyleborus*), *Cyclorhipidion* 699
destructor Eggers, *Sinophloeus* (=porteri) 105
destructor Olivier, *Scolytus* (=scolytus) 375
destructor Wood, *Phlocotribus* 221
destruens (Blandford), (*Xyleborus*), *Eucallacea* 687
destruens (Wollaston), (*Hydurgus*), *Tomicus* 123
desultorius Bright, *Pityophthorus* 995
desum (Wood), (*Microcorthyus*), *Monarthrum* 1054
detectus Schedl, *Cyrtogenius* 564
detectus Schedl, *Pityophthorus* 995
detectus Schedl, *Xyleborus* 724
detentus Wood, *Micracis* 432
detentus Wood, *Pityophthorus* 995
detersus (Chapuis), (*Phloeosinus*), *Hyledius* 260
detractus Wood, *Sampsonius* 655
detrimentosus Schedl, *Corthyus* 1073
detritus Eggers, *Xyleborus* 724
derexulus (Wood), (*Xyleborus*), *Ambrosiodmus* 673
derexus (Schedl), (*Xyleborus*), *Ambrosiodmus* 673
derexus Wood, (*Xyleborus*), *Ambrosiodmus*
 (=derexulus) 673
derius (Schedl), (*Cryphalophilus*), *Scolytogenes* 861
derius Schedl, (*Neodryocoetes*), *Arauptus*
 (=pallidus) 960
derius Schedl, *Xyleborus* 724
dexter (Sampson), (*Cryphalus*), *Hypothenemus* 916
deyrollei (Blandford), (*Pityophthorus*), *Arauptus* 955
diadematus Eggers, *Cryptocarenum* 903
dialiumi Schedl, *Strombophorus* 203
Diamerides Browne (=Phlococranus) 236
Diamerinae Hagedorn (=Diamerini) 193
Diamerini Hagedorn 193
Diamerus Erichson 194
diapiformis (Schedl), (*Xyleborus*), *Xyleborinus* 806
dichrous Eichhoff, *Xyleborus* 724
dietzi Hopkins, *Dendroctonus* (=rufipennis) 173
differens Hopkins, (*Stephanoderes*), *Hypothenemus*
 (=crudiae) 915
difficile (Blandford), (*Pterocyclon*), *Monarthrum* 1054
difficilis (Eggers), (*Xyleborus*), *Xylosandrus* 794
difficilis (Schedl), (*Micracis*), *Pseudomicracis* 429
difficilis Beeson, (*Urdugraphus*), *Polygraphus*
 (=difficilis) 254, 286
difficilis Wood, *Polygraphus* 256
difformis (Schedl), (*Loganius*), *Cnemonyx* 314
difodinus Bright, *Tricolus* 1043
digestus (LeConte), (*Cryphalus*), *Pityophthorus* 995
diglyphus Blandford, *Pityophthorus* 996
diglyptus Schedl, *Xyleborus* 724
dihingense Schedl, (*Xyleborus*), *Cyclorhipidion*
 (=dihingicum) 699
dihingensis (Eggers), (*Xyleborus*), *Ambrosiodmus* 673
dihingensis Eggers, *Xyleborus* (=rufobrunneus) 769
dihingicum Wood, *Cyclorhipidion* 699
dilatatiformis (Schedl), (*Xyleborus*), *Eucallacea* 655
dilatatus Schedl, *Xyleborus* (=similis) 774
dilatatus Eichhoff, *Xyleborus* (=similis) 773
diligens Schedl, *Xyleborus* (=sphenos) 775
diligens Wood, *Corthyus* 1073
diligens Wood, *Pityophthorus* 996
dilutus Blandford, *Scolytomimus* 839
dilutus Eichhoff, *Cryphalus* (=scabricollis) 894
dilutus Wood, *Gnathotrypes* 1040
dilutus Wood, *Hylocurus* 423
dilutus Wood, *Microcorthyus* 1065
 ° *diluvialis* Wickham, *Dryocoetes* 552
dimidiatum (Ferrari), (*Corthyus*), *Monarthrum* 1054
dimidiatum Hagedorn, (*Pterocyclon*), *Monarthrum*
 (=hagedorni) 1057
dimidiatus Blackman, *Pityophthorus* 996
dimidiatus Chapuis, *Scolytus* 329
dimidiatus Eggers, *Xyleborus* 724
diminutivus Bright, *Pityophthorus* 996
dimorphus (Schedl), (*Cryptocleptes*),
Pseudothysanoides 413
dimorphus (Schedl), (*Dryocoetes*), *Cyrtogenius* 564
dimorphus (Schedl), (*Metahylesinus*), *Hylesinopsis* 93
dimorphus (Schedl), (*Micracis*), *Hylocurus* 424
dimorphus (Schedl), (*Stephanoderes*),
Hypothenemus 916
dimorphus Schedl, *Cryphalus* (=sylvicola) 896

<i>dimorphus</i> Schedl. <i>Pityophthorus</i>	996	<i>distinctus</i> Wood, <i>Hypothenemus</i>	917
<i>dimorphus</i> Schedl. <i>Polygraphus</i>	286	<i>distinguendus</i> Eggers, <i>Xyleborus</i>	730
<i>dinoderoides</i> (Blandford), (<i>Dryocoetes</i>), <i>Taphrorychus</i>	559	<i>divaricata</i> Wood, <i>Micracisella</i>	429
<i>dipteroearpi</i> (Hopkins), <i>Cyclorhipidion</i>	699	<i>diversepilosus</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	794
<i>dipteroearpi</i> Beeson, <i>Cryphalus</i> (= <i>dipteroearpi</i>)	575	<i>diversicauda</i> Browne, <i>Webbia</i>	531
<i>dipteroearpi</i> Beeson, (<i>Thamniurgides</i>), <i>Coccotrypes</i> (= <i>barbatus</i>)	593	<i>diversicolor</i> (Eggers), (<i>Xyleborus</i>), <i>Coptodryas</i>	824
<i>dipteroearpi</i> Browne, <i>Arixyleborus</i>	666	<i>diversicolor</i> Browne, <i>Cryphalus</i>	579
<i>dipteroearpi</i> Hopkins, (<i>Ozopemon</i>), <i>Dryocoetiops</i> (= <i>laevis</i>)	555	<i>diversicolor</i> Eggers, (<i>Ozopemon</i>), <i>Dryocoetiops</i> (= <i>laevis</i>)	585
<i>dipteroearpi</i> Hopkins, <i>Hypothenemus</i>	916	<i>diversipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	673
<i>dipteroearpi</i> Hopkins, <i>Webbia</i>	530	<i>diversus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	806
<i>dipteroearpi</i> Wood, <i>Cryphalus</i>	575	<i>diversus</i> Bright, <i>Pityophthorus</i>	996
<i>dirus</i> Wood, <i>Xyleborinus</i>	806	<i>diversus</i> Wood, <i>Dendrocraniulus</i>	551
<i>discedens</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	391	<i>diversus</i> Wood, <i>Microcorthyus</i>	1065
<i>discedens</i> Eggers, <i>Scolytodes</i> (= <i>discriminatus</i>)	391	<i>dividuus</i> Schedl, <i>Cucinus</i>	207
<i>discedens</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>erectus</i>)	915	<i>divisus</i> (Browne), (<i>Webbia</i>), <i>Taphrodasus</i>	525
<i>discifer</i> Eichhoff, <i>Hylocurus</i>	424	<i>djuguensis</i> Eggers, <i>Pityophthorus</i>	996
<i>discoideus</i> Blandford, <i>Corthyus</i>	1073	<i>docta</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	824
<i>discolor</i> (Blandford), (<i>Xyleborus</i>), <i>Xylosandrus</i>	794	<i>dohrni</i> (Eichhoff), (<i>Steganoecranus</i>), <i>Amphicranus</i>	1046
<i>discrepans</i> Blandford, <i>Phloeotribus</i>	221	<i>dohrni</i> Wollaston, (<i>Tomicus</i>), <i>Xyleborinus</i> (= <i>saxeseni</i>)	514
<i>discrepans</i> Schedl, <i>Hypocryphalus</i>	869	<i>doliaris</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	683
<i>discrepans</i> Schedl, <i>Xyleborus</i>	724	<i>dolichocola</i> Hopkins, <i>Hypothenemus</i>	917
<i>discretus</i> Eggers, <i>Xyleborus</i>	725	<i>dolosus</i> Blandford, <i>Xyleborus</i>	730
<i>discretus</i> Eichhoff, <i>Cryphalus</i> (= <i>scabricollis</i>)	593	<i>dolosus</i> Wood, <i>Hypothenemus</i>	917
<i>discretus</i> Wood, <i>Cucinus</i>	207	<i>Dolurgocleptes</i> Schedl	540
<i>discretus</i> Wood, <i>Pityophthorus</i>	996	<i>Dolurgus</i> Eichhoff	615
<i>discriminatus</i> Wood, <i>Scolytodes</i>	391	<i>dolus</i> Schedl, <i>Hylesinus</i>	79
<i>dislocatus</i> (Blackman), (<i>Cryptocleptes</i>), <i>Pseudohysanocs</i>	413	<i>dolus</i> Wood, <i>Pityophthorus</i>	996
<i>dispar</i> (Eggers), (<i>Stephanoderes</i>), <i>Stephanopodius</i>	553	<i>domesticum</i> (Linnaeus), (<i>Dermestes</i>), <i>Trypodendron</i>	633
<i>dispar</i> (Fabricius), (<i>Apatе</i>), <i>Xyleborus</i>	725	<i>donaticus</i> Wood, <i>Corthyus</i>	1073
<i>dispar</i> (Schedl), (<i>Cryphalops</i>), <i>Enioporus</i>	555	<i>donisi</i> (Schedl), (<i>Eriocryphalus</i>), <i>Hypothenemus</i>	917
<i>dispar</i> (Schedl), (<i>Micracidendron</i>), <i>Saurotosis</i>	419	<i>donisi</i> (Schedl), (<i>Erioschidias</i>), <i>Cosmoderes</i>	901
<i>dispar</i> Blackman, <i>Pseudohylesinus</i>	109	<i>donisthorpei</i> Formanek, (<i>Cryphaloides</i>), <i>Coccotrypes</i> (= <i>carpophagus</i>)	595
<i>dispar</i> Bright, <i>Pityophthorus</i>	996	<i>donisthorpei</i> Schedl, <i>Xyleborus</i> (= <i>dryographus</i>)	732
<i>dispar</i> <i>dispar</i> Blackman, <i>Pseudohylesinus</i> (= <i>dispar</i>)	109	<i>dorjitenzingi</i> Schmutzenhofer, <i>Trypodendron</i>	635
<i>dispar</i> pullatus Blackman, <i>Pseudohylesinus</i> (= <i>dispar</i>)	109	<i>dorsalis</i> (Motschulsky), (<i>Hypoborus</i>), <i>Cryphalus</i>	579
<i>dispar</i> rugulosus Eggers, <i>Xyleborus</i> (= <i>dispar</i>)	730	<i>dorsalis</i> Schedl, <i>Pityophthorus</i>	997
<i>disparilis</i> Wood, <i>Chramesus</i>	265	<i>dorsalis</i> Schedl, <i>Xyleborus</i>	730
<i>disparilis</i> Wood, <i>Hylocurus</i>	424	<i>dorsosignatus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	917
<i>dissidens</i> Wood, <i>Hylocurus</i>	424	<i>dorsosulcatus</i> Beeson, <i>Xyleborus</i>	730
<i>dissidens</i> Wood, <i>Xyleborus</i> (= <i>titubanter</i>)	779	<i>dossuarius</i> Eggers, <i>Xyleborus</i>	730
<i>dissimilis</i> (Eggers), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	666	<i>draconis</i> Enderlein, <i>Dactyloctrypes</i> (= <i>longicollis</i>)	554
<i>dissimilis</i> (Nobuchi), (<i>Taeniohyptes</i>), <i>Cryphalus</i>	578	<i>drakei</i> Blackman, <i>Pseudohysanocs</i> (= <i>rigidus</i>)	416
<i>dissimilis</i> (Zimmermann), (<i>Crypturgus</i>), <i>Hypothenemus</i>	916	<i>drescheri</i> Eggers, <i>Eccoptyterus</i>	520
<i>dissimilis</i> Hagedorn, <i>Diamerus</i> (= <i>curvifer</i>)	195	<i>drescheri</i> Eggers, <i>Hyorrhynchus</i> (= <i>lewisi</i>)	192
<i>dissimilis</i> Schedl, <i>Scolytodes</i>	391	* <i>dromiscens</i> Scudder, <i>Hylesinus</i>	79
<i>dissimilis</i> Wood, <i>Hylocurus</i>	424	<i>dryobalanopsis</i> Schedl, <i>Cryptoxyleborus</i>	525
<i>dissimulatus</i> Wood, <i>Xyleborus</i>	730	<i>Dryocoetes</i> Eichhoff	569
<i>dissolutus</i> Wood, <i>Pityophthorus</i>	996	<i>Dryocoetina</i> Lindemann (= <i>Dryocoetini</i>)	539
<i>distans</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1054	<i>Dryocoetinae</i> Lindemann (= <i>Dryocoetini</i>)	539
<i>distinctus</i> (Motschulsky), (<i>Anodius</i>), <i>Coccotrypes</i>	602	<i>Dryocoetini</i> Lindemann	539

- Dryocoetinus* Balachowsky (= *Dryocoetes*) 569
- Dryocoetiops* Schedl** 587
- Dryocoetoideae* Lindemann (= *Dryocoetini*) 539
- Dryocoetoides* Hopkins** 656
- dryocoetoides* Numberg, *Cryptogenius*
(= *cribricollis*) 564
- dryographus* (Ratzeburg), (Bostrichus),**
Xyleborus 730
- dryographus* Schedl, *Cnesinus* (= *diriduus*) 207
- Dryotomicus* Wood (= *Phloeotribus*) 218
- Dryotomus* Chapuis (= *Phloeotribus*) 217
- dubiosus* (Schedl), (Pterocyclon), Monarthrum** 1054
- dubiosus* (Schedl), (Metacorthylus), Corthylus** 1073
- dubiosus* (Schedl), (Prionosceles), Scolytodes** 391
- dubiosus* (Schedl), (Thammophthorus), Araptus** 955
- dubiosus* Perkins, Xyleborus** 732
- dubiosus* Schedl, Cryptophalus** 579
- dubiosus* Schedl, Hypocryphalus** 569
- dubiosus* Schedl, Hypothenemus (= *eruditus*)** 924
- dubiosus* Schedl, Phloeosinus** 247
- dubiosus* Wood, Ptilopodius** 599
- dubium* Eichhoff, (Pterocyclon), *Monarthrum*
(= *quadrideus*) 1061
- dubius* (Eggers), (Dryocoetes), Cryptogenius** 564
- * *dubius* (Hagedorn), (Myelophilites), Hyurgops** 31
- dubius* (Schedl), (Neodyocoetes), Araptus** 955
- dubius* (Schedl), (Stylotenus), Hypothenemus** 917
- dubius* Blandford, Phloeosinus (= *pulchellus*)** 251
- dubius* Eggers, (Pocillips), Coccotrypes (= *sannio*)** 611
- dubius* Eggers, (Xyleborus), Euwallacea**
(= *xanthopus*) 696
- dubius* Eggers, Hylesinopsis** 93
- dubius* Eichhoff, Crypturgus (= *mediterraneus*)** 625
- dubius* Eichhoff, Phloeotribus (= *setulosus*)** 232
- dubius* Schedl, Hylocurus** 424
- dubius* Schedl, Miocryphalus** 410
- dubius* Swaine, Ips (= *tridens engelmanni*)** 525
- dulcispei* Beeson, Sphaerotrypes (= *subtectus*)** 201
- dumineus* Beeson, Cryptophalus (= *fulmineus*)** 580
- dunni* Swaine, Carphoborus** 305
- duodecimspinus* Schedl, Webbia** 831
- duodecimspinus* Schedl, Xyleborus** 732
- duongi* Villiers, Aphanarthrum** 618
- duplex* Browne, Xyleborus (= *deplanatus*)** 724
- duplicatus* (Sahlberg), (Bostrichus), Ips** 502
- duplicatus* (Schedl), (Mimips), Acanthotomicus** 450
- duplicatus* Eggers, (Xyleborus), Eccoptopterus**
(= *limbus*) 520
- duplicatus* Schedl, (Xyleborus), Euwallacea**
(= *quadraticollis*) 693
- duploarmatus* Browne, Xyleborus** 732
- duplocordatum* Eggers, Monarthrum** 1054
- duplopilosus* (Browne), (Pocillips), Coccotrypes** 602
- duplosquamosus* Schedl, Cryptophalus** 579
- duplosquamosus* Schedl, Pseudochramesus** 262
- duponti* Montrouzier, Xyleborus (= *perforans*)** 761
- duprezi* Hoffmann, Gnathotrichus**
(= *materiarius*) 1036
- durangoensis* Wood, Pseudopityophthorus** 973
- durum* (Schedl), (Pterocyclon), Monarthrum** 1054
- durus* Blackman, Pityophthorus** 997
- durus* Schedl, (Protopityophthorus), Dendrographus**
(= *pygmaeus*) 561
- calacensis* Eggers, (Archephalus), Hypothenemus**
(= *eruditus*) 925
- ebenium* (Blandford), (Brachyspartus),**
Corthyocyclon 1067
- ebenus* (Wood), (Xyleborus), Tawrodenus** 755
- ebianensis* Huang & Yin, Hyorrhynchus** 191
- ebriosus* Niisina, (Xyleborus), Xylosandrus**
(= *crassiusculus*) 792
- Eccoptogaster* Gyllenhal (= *Scolytus*)** 321
- Eccoptogasteridae* Reitter (= *Scolytidae*)** 25
- Eccoptogasteridae* Reitter (= *Scolytinae*)** 313
- Eccoptogasteridae* Reitter (= *Scolytini*)** 313
- Eccoptogasterinae* Reitter (= *Scolytinae*)** 313
- Eccoptogasterini* Reitter (= *Scolytinae*)** 313
- Eccoptogasterini* Reitter (= *Scolytini*)** 313
- eccoptopterus* (Schedl), (Xyleborus),**
Eccoptopterus 520
- Eccoptopterus* Motschulsky** 519
- echinatae* Wood, Conophthorus** 964
- ecksteini* Butovitsch, Scolytus** 329
- edentatus* Hagedorn, (Phthorus), Monarthrum**
(= *bicolor*) 1051
- editus* (Bright), Chramesus** 265
- edulis* Hopkins, Conophthorus** 964
- effeminatus* Wood, Hylocurus** 424
- egenum* (Blandford), (Pterocyclon), Monarthrum** 1054
- egenus* Blandford, Hylocurus** 424
- eggersi* (Schedl), (Brachydendrulus), Araptus** 955
- eggersi* (Schedl), (Cryptophalopus),**
Scolytogenes 561
- eggersi* (Schedl), (Hexacolus), Scolytodes** 391
- eggersi* Beeson, Xyleborus** 732
- eggersi* Hagedorn, Coccotrypes (= *dactyliperda*)** 601
- eggersi* Schedl, (Pocillips), Coccotrypes (= *cyperi*)** 599
- eggersi* Schedl, Cryptophyphthorus** 274
- eggersi* Schedl, Cryptoxyleborus** 525
- eggersi* Schedl, Diamerus** 195
- eggersi* Schedl, Phloeotribus (= *neglectus*)** 225
- eggersi* Schedl, Phloeotribus (= *demessus*)** 220
- eggersi* Schedl, Pityophthorus** 997
- eggersi* Stark, Erioporus (= *tiliae*)** 557
- eggersi* Wichmann, Cladoctonus**
eggersi Wood, (Renocis), *Chaetophloeus*
(= *mexicanus*) 273
- Eggersia* Lebedev (= *Pityogenes*)** 439
- eggersianus* (Schedl), (Pityophthorus), Araptus** 955
- eggersianus* Schedl, (Xyleborus), Cryptoxyleborus**
(= *eggersi*) 525
- ehlersi* Eichhoff, (Stephanoderes), Hypothenemus**
(= *eruditus*) 921

<i>ehlersi rottroui</i> Peyערimhoff, <i>Hypothenemus</i> (= <i>eruditus</i>)	923	<i>elongatulus</i> Schedl, <i>Pityophthorus</i>	997
<i>eichelbaumi</i> Hagedorn, <i>Scolytoplatypus</i>	402	<i>elongatulus</i> Wood, <i>Cyrtogenius</i>	564
<i>eichhoffi</i> (Schreiner), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	673	<i>elongatum</i> (Eggers), (<i>Aphanarthrum</i>), <i>Deropria</i>	620
<i>eichhoffi</i> Blandford, <i>Aricerus</i>	217	<i>elongatum</i> Eggers, <i>Liparthrum</i> (= <i>mori</i>)	278
<i>eichhoffi</i> Eggers, <i>Amphicranus</i>	1046	<i>elongatus</i> (Eggers), (<i>Dendurgus</i>), <i>Coccotrypes</i>	602
<i>eichhoffi</i> Eggers, <i>Hylastinus</i> (= <i>obscurus</i>)	65	<i>elongatus</i> (Eggers), (<i>Erioschidias</i>), <i>Cosmoderes</i>	901
<i>eichhoffi</i> Ferrari, <i>Dryocoetes</i>	582	<i>elongatus</i> (Eggers), (<i>Pelicerus</i>), <i>Cyrtogenius</i>	565
<i>eichhoffi</i> Hopkins, <i>Dryocoetes</i> (<i>betulae</i>)	579	<i>elongatus</i> (Schedl), (<i>Hylesinus</i>), <i>Ficicis</i>	91
<i>eichhoffi</i> Reitter, <i>Scolytus</i>	329	<i>elongatus</i> (Schedl), (<i>Landolphianus</i>), <i>Laururgus</i>	408
<i>eichhoffi</i> Schiauffuss, <i>Xyleborus</i> (= <i>eichhoffianus</i>)	732	<i>elongatus</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	450
<i>eichhoffi</i> Schedl, <i>Corthylus</i>	1073	<i>elongatus</i> (Schedl), (<i>Thamnophthorus</i>), <i>Araptus</i>	956
<i>eichhoffianus</i> Schedl, <i>Xyleborus</i>	732	<i>elongatus</i> Chapuis, <i>Phloeoborus</i> (= <i>rudis</i>)	100
<i>Eidophelinae</i> Murayama (= <i>Cryphalini</i>)	842	<i>elongatus</i> Eggers, <i>Hylastinus</i> (= <i>obscurus</i>)	65
<i>Eidophelus</i> Eichhoff	867	<i>elongatus</i> Eggers, (<i>Hylocurosoma</i>), <i>Scolytodes</i> (= <i>elongatissimus</i>)	392
<i>Ekkoptogaster</i> Herbst (= <i>Scolytus</i>)	321	<i>elongatus</i> Eggers, <i>Hyorthynclius</i>	191
<i>elaboratus</i> Schedl, <i>Cryphalus</i>	879	<i>elongatus</i> Eggers, <i>Strombophorus</i>	203
<i>elaeocarpus</i> Beeson, <i>Coccotrypes</i> (= <i>dactyliperda</i>)	602	<i>elongatus</i> Eggers, <i>Xyleborus</i>	733
<i>elaphas</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	917	<i>elongatus</i> Herbst, (<i>Bostrichus</i>), <i>Hylurgus</i> (= <i>ligniperda</i>)	121
<i>elatinus</i> Wood, <i>Pityophthorus</i>	997	<i>elongatus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	923
<i>elatus</i> Niisima, <i>Hylesinus</i>	79	<i>elongatus</i> Lovendal, (<i>Tomicus</i>), <i>Pityogenes</i> (= <i>trepanatus</i>)	459
<i>eldaricus</i> Stark, <i>Chactoptelius</i> (= <i>vestitus</i>)	104	<i>elongatus</i> (Nobuchi), (<i>Ericryphalus</i>), <i>Cryphalus</i>	579
<i>electinus</i> Wood, <i>Cnesinus</i>	207	<i>elongatus</i> Schedl, (<i>Eidophelus</i>), <i>Cyrtogenius</i> (= <i>elongatulus</i>)	564
* <i>electrinus</i> (Germar), (<i>Hylesinus</i>), <i>Hylurgops</i>	31	<i>elongatus</i> Schedl, (<i>Ozodendron</i>), <i>Cyrtogenius</i> (= <i>elongatissimus</i>)	564
<i>electus</i> (Wood), (<i>Gnathotrypanus</i>), <i>Gnathotrypes</i>	1040	<i>elongatus</i> Schedl, <i>Cladoctonus</i>	238
<i>electus</i> Blackman, <i>Pityophthorus</i>	997	<i>elongatus</i> Schedl, <i>Cryphalus</i>	879
<i>electus</i> Wood, <i>Amphicranus</i> (= <i>broicnei</i>)	1046	<i>elongatus</i> Schedl, <i>Micocryphalus</i>	411
<i>electus</i> Wood, <i>Cnesinus</i>	207	<i>elongatus</i> Schedl, <i>Scolytodes</i> (= <i>trispinosus</i>)	400
<i>elegans</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1054	<i>elongatus</i> Schedl, <i>Scolytus</i>	329
<i>elegans</i> (Sampson), (<i>Xyleborus</i>), <i>Coptodryas</i>	824	<i>elongatus</i> Schedl, <i>Thamnurgus</i>	543
<i>elegans</i> Blandford, <i>Cnesinus</i>	207	<i>elongatus</i> Schreiner, <i>Tiarophorus</i>	539
<i>elegans</i> Eichhoff, <i>Hylocurus</i>	424	<i>elongatus</i> Swaine, <i>Pityophthorus</i> (= <i>murayanae</i>)	1014
<i>elegans</i> Krivolitskaya, <i>Eidophelus</i> (= <i>imitans</i>)	867	<i>elsae</i> Eggers, <i>Pseudomicracis</i>	429
<i>elegans</i> Schedl, <i>Cosmoderes</i>	901	<i>Elzearius</i> Guillebeau (= <i>Phlocotribus</i>)	218
<i>elegans</i> Schedl, <i>Phlocoditica</i>	236	<i>emarginatus</i> (LeConte), (<i>Tomicus</i>), <i>Ips</i>	503
<i>elegans</i> Schedl, <i>Pityophthorus</i>	997	<i>emarginatus</i> (Numberg), (<i>Trypographus</i>), <i>Hylesinopsis</i>	94
<i>elegans</i> Schedl, <i>Thamnurgus</i>	543	<i>emarginatus</i> Browne, <i>Acanthotomicus</i>	450
<i>elegans</i> Swaine, <i>Pityokteines</i>	462	<i>emarginatus</i> Eggers, <i>Corthylus</i>	1073
<i>elegans</i> Thomson, <i>Hylesinus</i>	80	<i>emarginatus</i> Eggers, <i>Xyloctonus</i> (= <i>scolytoides</i>)	841
<i>elegans</i> Wichmann, (<i>Pseudothamnurgus</i>), <i>Tiarophorus</i> (= <i>normandi</i>)	540	<i>emarginatus</i> Eichhoff, <i>Xyleborus</i>	733
<i>elegantis</i> Wood, <i>Cnesinus</i>	208	<i>emarginatus</i> Hopkins, <i>Coptoborus</i> (= <i>respatorius</i>)	665
<i>elegantulus</i> Schedl, <i>Amphicranus</i>	1047	<i>emarginatus</i> Schedl, <i>Hypothenemus</i> (= <i>obscurus</i>)	937
<i>elegantulus</i> Schedl, <i>Amphicranus</i> (= <i>broicnei</i>)	1046	<i>emarginatus</i> Wichmann, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>scheryweici</i>)	371
<i>elevatus</i> Eggers, <i>Xyleborus</i>	732	<i>emarginatus</i> semicircularis Schedl, <i>Xyleborus</i> (= <i>emarginatus</i>)	733
<i>climatus</i> Bright, <i>Pityophthorus</i> (= <i>subopacus</i>)	1030	* <i>emmi</i> (Hagedorn), (<i>Cryphalus</i>), <i>Hypothenemus</i>	918
<i>ellipticus</i> Chapuis, <i>Phloeoborus</i>	99	<i>emmi</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>emmi</i> Hagedorn)	918
<i>ellipticus</i> Eggers, (<i>Hexacolus</i>), <i>Scolytodes</i> (= <i>maurus</i>)	395	<i>endroedyi</i> (Schedl), (<i>Hemihylesinus</i>), <i>Hylesinopsis</i>	94
<i>ellipticus</i> Murayama, <i>Scolytus</i>	329	<i>engadinensis</i> Fuchs, <i>Ips</i> (= <i>cembrae</i>)	497
<i>elongatissimus</i> Wood, <i>Cyrtogenius</i>	564	<i>engelmanni</i> Hopkins, <i>Dendroctonus</i> (= <i>rufipennis</i>)	178
<i>elongatissimus</i> Wood, <i>Scolytodes</i>	391		
<i>elongatulus</i> (Schedl), (<i>Poeicilips</i>), <i>Coccotrypes</i>	602		
<i>elongatulus</i> Schedl, <i>Dendrochilus</i>	438		

- engelmanni* Swaine, *Ips* (= *tridens*) 528
engelmanni Wood, *Carphoborus* (= *sansonii*) 309
enixus Blackman, *Phloeosinus* (= *dentatus*) 246
ensifer Eichhoff, *Scolytus* 329
Eocryphalus Kurenzov (= *Erioporicus*) 848
eos Spessivtsev, *Hylesinus* 80
epicaris Wood, *Thysanocs* 420
episternatus Schedl, (*Scolytus*), *Scolytopsis*
 (= *puncticollis*) 321
epistomalis Schedl, (*Pityophthorus*), *Gnatholeptus*
 (= *punamensis*) 970
Epomalius Blandford (= *Scolytodes*) 358
Epsips Beeson (= *Hypothenemus*) 905
equidens Wood, *Hylocurus* 424
equihuai Bright, *Pityophthorus* 997
equihuai Wood, *Cnemonyx* 314
equihuai Wood, *Cnesinus* 208
erectus LeConte, *Hypothenemus* 918
ericius (Schaufuss), (*Diamerus*), *Hylesinopsis* 94
ericius (Schaufuss), (*Diamerus*), *Strombophorus* 203
Erieryphalus Hopkins (= *Cryphalus*) 872
erinacea (Eggers), (*Xyleborus*), *Coptodryas* 824
erinacellus Wichmann, *Kissophagus* 68
erinaceus Schedl, *Chramesus* 265
erinaceus Schedl, *Phloeotribus* 221
Erineophilus Hopkins (= *Scolytodes*) 358
erineophilus Wood, *Scolytodes* 392
Erincosinus Blackman (= *Liparthrum*) 274
Erioschidias Schedl (= *Cosmoderes*) 900
Ernocladius Wood 857
Eriocryphalus Murayama (= *Cryphalus*) 872
Eriophloeus Nunberg (= *Hypothenemus*) 905
Eriopocerus Balachowsky (= *Erioporicus*) 848
Erioporicus Berger 848
Erioporides Hopkins (= *Scolytogenes*) 858
Erioporinae Nusslin (= *Cryphalini*) 842
Erioporus Thomson 854
erosus (Wollaston), (*Tomicus*), *Ips* 504
erosus Schedl, *Phloeotribus* 221
erosus melanurus Reitter, *Ips* (= *robustus*) 522
errans (Blandford), (*Ceratolepis*), *Cnemonyx* 314
errans Blandford, *Hylocurus* 424
erraticum Schedl, *Cryphalus* 879
erraticus Schedl, *Pityophthorus* 997
eruditus (Schedl), (*Neopityophthorus*), *Araptus* 956
eruditus Westwood, *Hypothenemus* 919
crythrinae Eggers, (*Stephanoderes*), *Hypothenemus*
 (= *eruditus*) 924
erythrinae Hopkins, *Margadillius* 899
esau Gredler, *Hylesinus* (= *toranio*) 85
espinosai Brethes, *Pityophthorus* 997
Estenoborus Reitter (= *Carphoborus*) 302
esuriens Blandford, *Scolytus* 330
Ethadopselaphus Blandford (= *Dactylipalpus*) 100
eucalyptica (Schedl), (*Xyleborus*), *Coptodryas* 824
eugeniae (Eggers), (*Xyleborus*), *Amasa* 683
eugeniae (Schedl), (*Dryocoetes*), *Dryocoetips* 587
Eulepiops Schedl (= *Cyrtogenius*) 562
Eulytocerus Blandford (= *Phloeotribus*) 218
eumerum (Schedl), (*Pterocyclon*), *Monarthrum* 1054
evonymi Kurenzov, *Alleroporius* 852
Eupagiocerus Blandford 214
eupatorii (Eggers), (*Xyleborus*), *Xylosandrus* 794
euphorbiae (Bright), (*Afrotrypetus*),
Styracoptinus 281
euphorbiae (Kuster), (*Bostrichus*), *Thamurgus* 543
euphorbiae (Schedl), (*Mimips*), *Acanthotomicus* 480
euphorbiae (Schedl), (*Stephanoderes*),
Hypothenemus 925
euphorbiae Schedl, *Lamurgus* 408
euphorbiae Wollaston, *Aphanarthrum* 618
euphorbiae Wood, *Cnemonyx* 315
euphorbiae Wood, *Cryphalogenes* 857
eupolyphagus Beeson, *Hypothenemus* (= *areccae*) 907
Eupteroxylon Eggers (= *Monarthrum*) 1050
Euptilius Schedl (= *Erioporus*) 854
Eurydactylus Hagedorn (= *Eccoptopterus*) 819
eurygraphus (Ratzeburg), (*Bostrichus*),
Xyleborus 733
euryopsis Schedl, *Thamurgus* 543
eurypterus Schedl, *Chramesus* 265
eusulcatus Tsai & Huang, *Hyलगорps* 31
euterpes Bright, *Pityophthorus* 997
eutomoides Blandford, *Scolytolatypus* 402
Euvallacea Hopkins 685
eranesens Wood, *Micracis* 432
evidens Wood, *Cnemonyx* 315
evonymi Hopkins, (*Stephanoderes*), *Hypothenemus*
 (= *eruditus*) 923
exactus (Schedl), (*Xyleborus*), *Amasa* 683
exaratus Blandford, *Xyleborus* 734
exasperatus Schedl, *Coccotrypes* (= *carpophagus*) 597
excaratus (Eggers), (*Anchonocerus*),
Monarthrum 1054
 * *excaratus* (Hagedorn), (*Xyleborus*),
Xyleborinus 806
excaratus (Schedl), (*Mimips*), *Acanthotomicus* 480
excaratus (Wood), (*Cryptulocleptus*),
Pseudothysanocs 413
excaratus Schedl, *Coccotrypes* 602
excarus Schedl, *Xyleborus* 734
excellens (Schedl), (*Cryphalomorphus*),
Scolytogenes 861
excellens Schedl, *Pityophthorus* 997
excisus (Ferrari), (*Morizus*), *Corthylus* 1073
exesus Blandford, *Xyleborus* 734
exesus Say, (*Bostrichus*), *Ips* (= *calligraphus*) 496
exigialis Wood, *Araptus* 956
exigua Wood, *Trischidias* 947
exiguus (Blandford), (*Loganius*), *Cnemonyx* 315
exiguus (Browne), (*Euptilius*), *Erioporus* 855
exiguus (Walker), (*Bostrichus*), *Xyleborinus* 806
exiguus Blackman, *Pityophthorus* (= *opaculus*) 1018

<i>exiguus</i> Blandford, <i>Cryphalus</i>	579	<i>fallax</i> Eggers, (<i>Minips</i>), <i>Acanthotomicus</i> (= <i>quadriloberculatus</i>)	453
<i>exiguus</i> Wood, <i>Cortylus</i>	1073	<i>fallax</i> Eggers, <i>Ips</i> (= <i>cembrae</i>)	499
<i>exiguus</i> Wood, <i>Cryphalogenes</i>	557	<i>fallax</i> Eichhoff, <i>Xyleborus</i>	735
<i>exiguus</i> Wood, <i>Scolytodes</i>	392	<i>fallax</i> Hagedorn, (<i>Meringopalpus</i>), <i>Gymnochilus</i> (= <i>zonatus</i>)	357
<i>exile</i> Eichhoff, (<i>Pterocyclon</i>), <i>Microcortylus</i> (= <i>parvulus</i>)	1066	<i>fallax</i> Wichmann, <i>Hylurgops</i> (= <i>batuensis</i>)	30
<i>exilis</i> (Wood), (<i>Loganius</i>), <i>Cnemonyx</i>	315	<i>fallaxoides</i> Schedl, <i>Xyleborus</i>	735
<i>exilis</i> Chapuis, <i>Hylastes</i>	53	<i>falsus</i> Schedl, <i>Xyleborus</i>	735
<i>exilis</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>pseudotenuis</i>)	664	<i>familiaris</i> (Schedl), (<i>Xyleborus</i>), <i>Prennobiis</i>	653
<i>exilis</i> Swaine, <i>Pityophthorus</i> (= <i>murrayanae</i>)	1014	<i>fankhauseri</i> Reitter, <i>Hylastinus</i>	64
<i>exilis</i> Wood, <i>Chramesus</i>	265	<i>farinosus</i> Blandford, <i>Hypothenemus</i> (= <i>birmanus</i>)	910
<i>exilis</i> Wood, <i>Micracis</i>	432	<i>fasciatum</i> (Say), (<i>Bostrichus</i>), <i>Monarthrum</i>	1055
<i>eximius</i> (Schedl), (<i>Erichsidias</i>), <i>Ernoporicus</i>	850	<i>fasciatus</i> (Blackman), (<i>Renocis</i>), <i>Chaetophloeus</i>	272
<i>eximius</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	480	<i>fasciatus</i> (Hagedorn), (<i>Kissophagus</i>), <i>Hylesinopsis</i>	94
<i>eximius</i> Schedl, <i>Hypothenemus</i>	925	<i>fasciatus</i> Fabricius, (<i>Hylesinus</i>), <i>Camptocerus</i> (= <i>suturalis</i>)	320
<i>eximius</i> Schedl, <i>Pityophthorus</i>	997	<i>fasciatus</i> Hagedorn, <i>Scolytoplatypus</i>	403
<i>eximius</i> Schedl, <i>Xyleborus</i>	734	<i>fasciatus</i> LeConte, <i>Hylesinus</i>	80
<i>eximius</i> Wood, <i>Dendroterus</i>	951	<i>fasciatus</i> Reitter, <i>Scolytus</i> (= <i>kirschi</i>)	337
<i>eximius</i> Wood, <i>Scolytodes</i>	392	<i>fastigatus</i> Schedl, <i>Xyleborus</i>	735
<i>exornatum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1055	<i>fastigatus</i> Blandford, <i>Amphicranus</i>	1047
<i>exornatus</i> Schedl, <i>Traglostus</i>	410	<i>fastigiorum</i> Wood, <i>Monarthrum</i>	1056
<i>expers</i> (Blandford), (<i>Hypothenemus</i>), <i>Scolytogenes</i>	861	<i>fastigius</i> (Wood), (<i>Loganius</i>), <i>Cnemonyx</i>	315
<i>explicitus</i> Wood, <i>Pityophthorus</i>	995	<i>fauxeli</i> Reitter, <i>Scolytus</i> (= <i>rugulosus</i>)	369
<i>expressus</i> Schedl, <i>Sphaerotripes</i>	199	<i>fefferi</i> Keler, (<i>Ips</i>), <i>Orthotomicus</i> (= <i>proximus</i>)	475
<i>expulsus</i> Wood, <i>Sampsonius</i>	656	<i>felis</i> Beeson, <i>Cryphalus</i> (= <i>felis</i> Wood)	550
<i>exquisitus</i> (Blackman), (<i>Neodryocotes</i>), <i>Pityophthorus</i>	995	<i>felis</i> Wood, <i>Cryphalus</i>	550
<i>exsulpta</i> (Eggers), (<i>Xyleborus</i>), <i>Coptodryas</i>	824	<i>felix</i> (Schedl), (<i>Xyleborus</i>), <i>Prennobiis</i>	653
<i>exsculptus</i> (Ratzeburg), <i>Pityophthorus</i>	995	<i>femineus</i> Wood, <i>Hylocurus</i>	424
<i>exsectus</i> Perkins, <i>Xyleborus</i>	735	<i>femoratus</i> Eggers, <i>Xyleborus</i> (= <i>spathipennis</i>)	774
<i>exsectus</i> Schedl, <i>Pityophthorus</i>	999	<i>fenestratum</i> Eggers, <i>Monarthrum</i>	1056
<i>extensa</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	824	<i>fennicus</i> Eggers, <i>Pityophthorus</i> (= <i>micrographus</i>)	1012
<i>externedentatus</i> Schedl, <i>Hypothenemus</i>	926	<i>fenoris</i> Wood, <i>Trichus</i>	1043
* <i>extractus</i> Scudder, <i>Hylesinus</i>	80	<i>ferinus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	794
<i>exul</i> (Reitter), (<i>Thamnurgus</i>), <i>Taphronurgus</i>	546	<i>ferox</i> Blandford, <i>Xyleborus</i>	735
<i>exul</i> Wood, <i>Chramesus</i>	265	<i>ferrarii</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056
<i>exultus</i> (Wood), (<i>Xyleborus</i>), <i>Coptoborus</i>	663	<i>ferreirai</i> (Schedl), (<i>Styracopterus</i>), <i>Styracoptinus</i>	251
<i>fabricii</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>apicalis</i>)	671	<i>ferrugineus</i> (Fabricius), (<i>Bostrichus</i>), <i>Xyleborus</i>	735
<i>facetus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	673	<i>ferrugineus</i> Boheman, (<i>Bostrichus</i>), <i>Xyleborus</i> (= <i>similis</i>)	773
<i>facetus</i> Wood, <i>Araptus</i>	956	<i>ferrugineus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>multidentatus</i>)	936
<i>facetus</i> Wood, <i>Scolytodes</i>	392	<i>ferrugineus</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	923
<i>facialis</i> Schedl, <i>Scolytus</i>	331	<i>ferulae</i> Pfeffer, <i>Cisurgus</i>	630
* <i>facilis</i> Heer, <i>Hylesinus</i>	80	<i>ferus</i> (Wood), (<i>Ambrosiosmus</i>), <i>Ambrosiodmus</i>	673
<i>fagi</i> (Fabricius), (<i>Apaté</i>), <i>Ernoporicus</i>	850	<i>festivum</i> Schedl, <i>Ctenoxylon</i>	837
<i>fagi</i> (Nobuchi), (<i>Pocillips</i>), <i>Coccotripes</i>	603	<i>festivus</i> Eichhoff, <i>Xyleborus</i>	739
<i>fagi</i> Blackman, <i>Pseudopityophthorus</i>	973	<i>festivus</i> Schedl, <i>Cyrtogenius</i>	565
<i>fagi</i> Walsh, <i>Scolytus</i>	330	<i>festivus</i> Wood, <i>Micracis</i>	432
<i>falcarius</i> Schedl, <i>Xyleborus</i>	735	<i>festivus</i> Wood, <i>Pseudopityophthorus</i>	973
<i>fallaciosus</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	450	<i>festus</i> Wood, <i>Pityophthorus</i>	999
<i>fallaciosus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>usagarius</i>)	665	<i>festus</i> Wood, <i>Scolytodes</i>	392
<i>fallax</i> (Eggers), (<i>Pocillips</i>), <i>Coccotripes</i>	603	<i>fici</i> Blandford, <i>Diameris</i>	195
<i>fallax</i> (Hagedorn), (<i>Trigonogenius</i>), <i>Pityophthorus</i>	999	<i>fici</i> Beeson, (<i>Eriocryphalus</i>), <i>Cryphalus</i> (= <i>scabricollis</i>)	893
<i>fallax</i> Costa Lima, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915	<i>fici</i> Browne, <i>Acanthotomicus</i>	481

- fici* Lea, *Aricerus* (= *eichhoffi*) 217
fici Browne, *Hypocryphalus* 869
fici Wood, *Phlocotribus* 221
fici Wood, *Pycnarthrum* (= *reticulatum*) 355
Ficicis Lea 90
ficicolens Wood, *Scolytodes* 392
Ficiphagus Murayama (= *Ficicis*) 90
ficivorus Murayama, *Cryphalus* 550
ficivorus Wood, *Scolytodes* 392
ficus (Schedl), (*Kissophagus*), (*Hylesinopsis*) 94
ficus (Schwarz), (*Loganius*), (*Cnemomyx*) 315
ficus Beeson, *Cryphalus* (= *scabricollis*) 593
ficus Eggers, *Xyleborus* 740
ficus Erichson, *Hypoborus* 250
ficus Hopkins, (*Stephanoderes*), (*Hypothenemus*)
 (= *seriatus*) 942
fiebrigi Hopkins, (*Stephanoderes*), (*Hypothenemus*)
 (= *seriatus*) 942
figuratus Schedl, *Xyleborus* 740
fijianus (Schedl), (*Lepicerinus*), (*Scolytogenes*) 561
fijianus (Schedl), (*Ozopemon*), (*Cyrtogenius*) 565
fijianus (Schedl), (*Pociclops*), (*Coccotrypes*) 603
fijianus (Schedl), (*Xyleborus*), (*Xylosandrus*) 794
fijianus Schedl, (*Stephanopodius*), (*Hypothenemus*)
 (= *dorsosignatus*) 917
filiformis (Schedl), (*Xyleborus*), (*Eucallacea*) 685
filliformis Blandford, *Amphicranus* 1047
filum (Reitter), (*Crypturgus*), (*Cisurgus*) 630
filum Schedl, *Dendrochilus* 435
fimbriaticorne (Blandford), (*Pterocyclon*),
Monarthrum 1056
fimbriatus (Schedl), (*Gnathotrichus*),
Gnathotrupes 1040
fimbriatus Wood, *Pseudothysanocs* 413
fimbricornis LeConte, *Thysanocs* 420
fiorii Eggers, (*Hylastinus*), (*Pagiocerus*) (= *frontalis*) 213
fischeri Hagedorn, *Xyleborus* 740
fiskei Blackman, *Scolytus* (= *unispinosus*) 350
fitchi Hopkins, *Xyleborus* (= *intrusus*) 745
fixasi Blackman, (*Myeloborus*), (*Pityophthorus*)
 (= *ramiperda*) 1025
flabellifer Sampson, *Cryphalus* (= *scabricollis*) 593
flagellatus Wood, *Hylocurus* 425
flagellifer Blandford, *Corthylus* 1073
flaglerensis Blackman, *Hylocurus* 425
flavescens Hagedorn, *Ctonoxylon* 537
flavescens Hopkins, (*Stephanoderes*), (*Hypothenemus*)
 (= *interstitialis*) 931
flavescens usambaricum Eggers, *Ctonoxylon*
 (= *flavescens*) 537
flavicollis Enderlein, *Coleobotrus* (= *luridus*) 616
flavicollis Hopkins, (*Stephanoderes*), (*Hypothenemus*)
 (= *eruditus*) 923
flavicornis (Blandford), (*Dryocoetes*),
Dryocoetops 557
flavicornis (Chapuis), (*Loganius*), (*Cnemomyx*) 315
flavicornis Chevrolat, *Scolytus* (= *multistriatus*) 353
flavicornis Lindberg, *Hylastes* (= *linearis*) 56
flavimaculatus Murayama, *Pityophthorus* 999
flavipennis Krausse, *Tomicus* (= *minor*) 123
flavipennis Schedl, *Xyleborus* 740
flavipes (Fabricius), (*Bostrichus*), (*Taurodemus*) 755
flavipes Hopkins, (*Hypothenemus*) 923
flavipes Panzer, *Hylurgus* (= *ligniperda*) 121
flavipubeus Schedl, *Strombophorus* 203
flavopannus Huang & Yin, *Hyorrhynchus* 192
flavopilosus Schedl, *Cnesinus* (= *setulosus*) 211
flavopilosus Schedl, *Xyleborus* 740
flavosquamosus Hopkins, (*Hypothenemus*)
 (= *eruditus*) 922
flavus (Fabricius), (*Hylesinus*), (*Dryocoetoides*) 657
flavus Hopkins, (*Hypothenemus*) 926
flavus Krausse, *Tomicus* (= *minor*) 123
flavus Stephens, (*Tomicus*), (*Xyleborus*)
 (= *dryographus*) 732
fletcheri Hopkins, (*Dendroctonus*) (= *rufipennis*) 173
flectiauxi Blandford, *Xyleborus* (= *subcostatus*) 776
flexilis Hopkins, (*Conophthorus*) (= *ponderosae*) 966
flexicostatus Schedl, (*Xyleborus*), (*Coptodryas*)
 (= *elegans*) 524
floccosus Hagedorn, *Dactylipalpus* 102
flohri (Eggers), (*Hylurgops*), (*Hylastes*) 54
flohri (Schedl), (*Pterocyclon*), (*Monarthrum*) 1056
flohri Schedl, *Xyleborus* (= *horridus*) 743
floridensis Atkinson, *Hylocurus* 425
floridensis Blackman, (*Tachyderes*), (*Cryptocareus*)
 (= *seriatus*) 904
floridensis Hopkins, (*Ernoporides*), (*Scolytogenes*)
 (= *knabi*) 563
floridensis Hopkins, (*Stephanoderes*), (*Hypothenemus*)
 (= *seriatus*) 942
floridensis Hopkins, (*Xylocleptes*), (*Dendrocramulus*)
 (= *carbonarius*) 550
floridensis Hopkins, (*Xyleborus*), (*Xyleborinus*)
 (= *saxseni*) 515
floridensis Schedl, (*Coccotrypes*) (= *distinctus*) 602
foederatus Schedl, *Xyleborus* 740
foersteri (Hagedorn), (*Xyleborus*), (*Cyclorhpidion*) 699
foratus Wood, *Cnesinus* 205
foratus Wood, (*Pityophthorus*) (= *comosus*) 955
forcipatus (Schedl), (*Xyleborus*), (*Xyleborinus*) 507
forcicatus (Schedl), (*Xyleborus*), (*Coptoborus*) 663
forciculoides (Schedl), (*Xyleborus*), (*Xyleborinus*) 507
forciculoides dentibaris Schedl, (*Xyleborus*),
Xyleborinus (= *forciculoides*) 507
forciculoides pinguis Schedl, (*Xyleborus*), (*Xyleborinus*)
 (= *forciculoides*) 507
forciculus (Eggers), (*Xyleborus*), (*Xyleborinus*) 507
formosae Wood, *Xyleborus* 740
formosanus (Schedl), (*Orosiotes*), (*Cyrtogenius*) 565
formosanus (Schedl), (*Xylechinus*),
Phlocosinopsioides 239
formosanus Browne, (*Ptilopodius*) 599
formosanus Browne, (*Xyleborus*) (= *formosae*) 740
formosanus Eggers, (*Scolytus*) (= *frontalis*) 331

<i>formosanus</i> Eggers, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>maucus</i>)	797	<i>frontalis</i> Schedl. <i>Lanurgus</i>	405
<i>formosanus</i> Nobuchii, <i>Dryocoetes</i>	582	<i>frontalis</i> Kurenzox, <i>Polygraphus</i> (= <i>sachalinensis</i>)	295
<i>formosanus</i> Schedl, <i>Cryphalus</i>	580	<i>frontalis</i> Wood, <i>Araptus</i>	956
<i>formosanus</i> Schedl, <i>Hypocryphalus</i>	569	<i>frontalis</i> Wood, <i>Cnesinus</i>	205
<i>formosus</i> Bright, <i>Pityophthorus</i>	999	<i>frontalis</i> Wood, <i>Carpoborus</i>	305
<i>fornicator</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacca</i> (= <i>fornicatus</i>)	690	<i>frontalis</i> Wood, <i>Pityoborus</i>	965
<i>fornicatus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Eucallacca</i>	688	<i>frontalis</i> Wood, <i>Tricolus</i>	1043
<i>fortis</i> (Wood), (<i>Mimips</i>), <i>Acanthotomicus</i>	481	<i>frontalis</i> Zimmermann, <i>Dendroctonus</i>	143
<i>fortis</i> Blackman, <i>Pityophthorus</i> (= <i>virilis</i>)	1033	<i>frontis</i> Wood, <i>Araptus</i>	956
<i>fossifrons</i> (LeConte), (<i>Pityophthorus</i>), <i>Pityogenes</i>	451	<i>frontoglabratus</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	392
<i>fossifrons</i> Wood, <i>Araptus</i>	956	<i>frugalis</i> Wood, <i>Araptus</i>	956
<i>fossifrons</i> Wood, <i>Dendroctonus</i>	950	<i>frustrata</i> Wood, <i>Phrixosoma</i>	159
<i>fonqueti</i> (Schedl), <i>Cyclorhpidion</i>	699	<i>fryi</i> Blandford, <i>Amphicranus</i>	1047
<i>foreatus</i> (Blackman), (<i>Cnesinus</i>), <i>Bothrostermus</i>	215	<i>fuchsi</i> Reitter, <i>Scolytus</i> (= <i>scolytus</i>)	376
<i>foreicollis</i> (Browne), (<i>Xyleborus</i>), <i>Anasa</i>	683	<i>fugax</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	561
<i>foreifrons</i> (Schedl), (<i>Thamnophthorus</i>), <i>Araptus</i>	956	<i>fugax</i> Schedl, <i>Cryphalus</i>	580
<i>foreolatus</i> Eggers, <i>Pityogenes</i>	451	<i>fujisanus</i> (Nobuchi), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	561
<i>fragilis</i> (Browne), (<i>Streptocranus</i>), <i>Coptoborus</i>	663	<i>fukiensis</i> Eggers, <i>Tomicus</i> (= <i>brevipilosus</i>)	123
<i>fragosus</i> Schedl, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>nugax</i>)	526	<i>fukiensis</i> Eggers, <i>Xyleborus</i>	740
<i>frankei</i> Wichmann, <i>Scolytus</i> (= <i>schetyrewi</i>)	371	<i>fulgens</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	561
<i>franseriae</i> Wood, <i>Pityophthorus</i>	999	<i>fulgens</i> (Schedl), (<i>Pocilips</i>), <i>Coccotrypes</i>	603
<i>fraseri</i> Hopkins, <i>Cryphalus</i> (= <i>ruficollis</i>)	891	<i>fulgens</i> (Schedl), (<i>Xyleborus</i>), <i>Anasa</i>	683
<i>fraterculus</i> (Schaufuss), <i>Ambrosiodmus</i>	674	<i>fulgens</i> Schedl, <i>Monarthrum</i>	1056
<i>fraternus</i> Blandford, (<i>Xyleborus</i>), <i>Eucallacca</i> (= <i>xanthopus</i>)	696	<i>fulgens</i> Schedl, <i>Pityophthorus</i>	999
<i>fraxini</i> (Berger), (<i>Ernoporus</i>), <i>Procryphalus</i>	545	<i>fulgens</i> Swaine, <i>Phloeosinus</i>	247
<i>fraxini</i> (Eggers), (<i>Phloeophthorus</i>), <i>Phloeotribus</i>	221	<i>fulgens</i> Wood, <i>Cnesinus</i>	205
<i>fraxini</i> Panzer, (<i>Bostrichus</i>), <i>Hylesinus</i> (= <i>varius</i>)	86	<i>fulgidum</i> Wood, <i>Pycnarthrum</i>	354
<i>fraxini</i> Stark, <i>Scolytus</i> (= <i>japonicus</i>)	335	<i>fulgidus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	561
<i>fraxinoides</i> Beeson, <i>Hylesinus</i> (= <i>macmahoni</i>)	51	<i>fulgidus</i> Blackman, <i>Hylastes</i>	54
<i>fraxinoides</i> Schedl, (<i>Lepersinus</i>), <i>Hylesinus</i> (= <i>macmahoni</i>)	51	<i>fulgidus</i> Wood, <i>Amphicranus</i>	1047
<i>freiurgi</i> Schedl, <i>Xylechinus</i>	115	<i>fulgidus</i> Wood, <i>Cnesinus</i>	205
<i>frenatus</i> (Schedl), (<i>Thamnophthorus</i>), <i>Araptus</i>	956	<i>fulgidus</i> Wood, <i>Dendrocraniulus</i>	551
<i>freyi</i> Schedl, (<i>Problochelus</i>), <i>Gymnochilus</i> (= <i>zonatus</i>)	357	<i>fuliginosus</i> (Eggers), (<i>Xyleborus</i>), <i>Arixyleborus</i>	666
<i>freyi</i> Schedl, <i>Phlocoborus</i>	99	<i>fuliginosus</i> Blandford, <i>Cryphalus</i>	550
<i>frigidus</i> (Schedl), (<i>Orosiotes</i>), <i>Cyrtogenius</i>	565	<i>fuliginosus</i> Blandford, <i>Xylechinus</i>	115
<i>frigidus</i> Blackburn, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxensis</i>)	514	<i>fulmineus</i> Wood, <i>Cryphalus</i>	550
<i>froggatti</i> Nimberg, <i>Hypocryphalus</i> (= <i>perminimus</i>)	571	<i>fulmineus</i> Wood, <i>Scolytodes</i>	392
<i>frondicolens</i> Wood, <i>Pseudothysanoces</i>	413	<i>fulripennis</i> (Nobuchi), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	561
<i>frontale</i> Eggers, <i>Liparthrum</i> (= <i>curtum</i>)	276	<i>fulripennis</i> Niisima, <i>Polygraphus</i>	256
<i>frontalis</i> (Fabricius), (<i>Bostrichus</i>), <i>Pagiocerus</i>	213	<i>fulvulus</i> Schedl, <i>Xyleborus</i>	740
<i>frontalis</i> (Olivier), (<i>Scolytus</i>), <i>Phloeotribus</i>	221	<i>fulvus</i> (Murayama), (<i>Xyleborus</i>), <i>Eucallacca</i>	690
<i>frontalis</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	999	<i>fulvus</i> Browne, <i>Margadillius</i>	599
<i>frontalis</i> (Schedl), (<i>Gnathotrichus</i>), <i>Gnathotrupes</i>	1040	<i>fulvus</i> Niisima, <i>Cryphalus</i>	580
<i>frontalis</i> (Wood), (<i>Erioschidias</i>), <i>Cosmoderes</i>	901	<i>fulvus</i> Schedl, <i>Xyleborus</i> (= <i>fulvulus</i>)	740
<i>frontalis</i> Blandford, <i>Scolytus</i>	331	<i>funiatus</i> DeGeer, <i>Tomicus</i> (= <i>piniperda</i>)	125
<i>frontalis</i> Bruck, <i>Phloeosinus</i>	247	<i>funebri</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	674
<i>frontalis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>brunneus</i>)	911	<i>funebri</i> Wood, <i>Pseudothysanoces</i>	413
<i>frontalis</i> Schedl, (<i>Gnathocranus</i>), <i>Araptus</i> (= <i>frontis</i>)	956	<i>funereus</i> (Lea), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	674
		<i>funereus</i> Wood, <i>Pseudothysanoces</i>	414
		<i>funerium</i> Wood, <i>Pycnarthrum</i>	354
		<i>funestus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	674
		<i>fungicola</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>arecae</i>)	907

- furcatus* Marsham, (*Ips*), *Pteleobius* (= *vittatus*) 72
furnissi Blackman, *Phloeosinus* 247
furnissi Blackman, *Pseudohylesinus* (= *nobilis*) 112
furnissi Bright, *Pityophthorus* 999
furukawai Murayama, *Cryphalus* 880
furcatus Wood, *Pseudothysanoes* 414
furrescens Wood, *Araptus* 957
furrescens Wood, *Cnemonyx* 315
furrescens Wood, *Pseudothysanoes* 414
furrum Wood, *Corthycyclon* 1067
furrus (Sampson), (*Dryocoetes*), *Coccotrypes* 603
furrus Wood, *Araptus* 957
furrus Wood, *Phloeotribus* 222
furrus Wood, *Pseudothysanoes* 414
fuscatus Eichhoff, *Xyleborus* (= *ferrugineus*) 737
fuscescens Stephens, (*Hylurgus*), *Hylastinus*
 (= *obscurus*) 67
fuscicollis (Eichhoff), (*Stephanoderes*),
Hypothenemus 926
fuscicollis Hagedorn, *Ozopemon* 589
fuscipennis (Chapuis), (*Phloeotribus*), *Hylesinopsis* . 94
fuscipennis Krausse, *Tomicus* (= *minor*) 123
fuscipilosus (Eggers), (*Xyleborus*), *Amasa* 683
fusciseriatus Eggers, *Xyleborus* (= *spimulosus*) 775
fuscobrunneus Eichhoff, *Xyleborus* (= *affinis*) 708
fuscovillosa (Schedl), (*Bothryperus*), *Phrixosoma* . 189
fuscum Hagedorn, *Ctonoxylon* (= *camerunum*) 837
fuscus Blackman, (*Renocis*), *Chaetophloeus*
 (= *heterodoxus*) 272
fuscus Blackman, *Pityophthorus* 999
fuscus Blandford, *Corthylus* 1074
fuscus Duftschmidt, (*Hylesinus*), *Hylurgops*
 (= *palliatius*) 38
fuscus Eggers, (*Xyleborus*), *Hadrodemius* (= *globus*) . 819
fuscus Marsham, (*Ips*), *Taphrorychus* (= *bicolor*) . . 558
fushimensis Murayama, (*Hylurgops*), *Hylastes*
 (= *plumbeus*) 60
fuyugei Schedl, *Xyleborus* 740
gaimaensis (Murayama), (*Xyloterus*),
Trypodendron 635
galeatus (Blandford), (*Xyleborus*), *Euwallacea* . . 690
galeatus (Eggers), (*Steganoecranus*), *Amphicranus*
 (= *eichhoffi*) 1047
galeritus Eichhoff, *Cnemonyx* 315
galeritus Wood, *Pityophthorus* 999
gallicus Eggers, *Kissophagus* (= *hederae*) 68
galoanus (Browne), (*Xyleborus*), *Euwallacea* . . . 690
gambetti Blackman, *Pseudothysanoes* (= *sedulus*) . . 417
ganshoensis Murayama, *Xyleborus* 740
garambaensis Nunberg, *Cryphalus* 881
garciae Schedl, (*Stephanoderes*), *Hypothenemus*
 (= *stigmaeus*) 945
garciniae (Schedl), (*Bothryperus*), *Phrixosoma* . . 189
gardneri Schedl, *Tiarophorus* 539
garrulus Schedl, *Cnesinus* 208
gaujoni Fairmaire, *Phloeoborus* 99
gaunersdorferi Reitter, *Crypturgus* (= *pusillus*) . . . 628
geayi Hagedorn, *Xyleborus* 740
gedeanus (Eggers), (*Thannurgides*), *Coccotrypes* . 603
gedeanus (Schedl), (*Xyleboricus*), *Arixyleborus* . . 666
gemellus Blandford, *Glochinocerus* 1064
geminatus (Hagedorn), (*Xyleborus*), *Amasa* 683
geminatus Zetterstedt, (*Bostrichus*), *Ips*
 (= *acuminatus*) 489
geminus Wood, *Phloeotribus* 222
genealis Wood, *Araptus* 957
 Genera Incertae Sedis 1082
genialis Wood, *Scolytodes* (= *gennaeus*) 392
genialis Wood, *Scolytodes* 392
genistae (Aube), (*Hypoborus*), *Liparthrum* 277
gennaeus Wood, *Scolytodes* 392
gentilis (Schedl), (*Xyleborus*), *Coptoborus* 663
gentilis Schedl, (*Pityophthorus*), *Gnatholeptus*
 (= *shannoni*) 970
geoffroi Goeze, (*Dermestes*), *Scolytus* (= *scolytus*) . 374
georgi Knotek, *Liparthrum* (= *genistae*) 277
georgiae Hopkins, (*Stephanoderes*), *Hypothenemus*
 (= *seriatus*) 941
georgiae Hopkins, *Trischidius* 947
gergeri Eggers, (*Hylastes*), *Hylurgops* (= *batnensis*) . 30
germanus (Blandford), (*Xyleborus*), *Xylosandrus* . 795
germanus Bright, *Pityophthorus* 1000
germari Eichhoff, (*Stephanoderes*), *Hypothenemus*
 (= *eruditus*) 921
germeauxi Menier, *Colcophorus* 616
geschwindi Seitner, (*Phloeophthorus*), *Phloeotribus*
 (= *cristatus*) 220
gezei Lepesme, *Xyleborus* 741
ghanaensis (Schedl), (*Cryphalomorphus*),
Scolytogenes 861
ghanaensis (Schedl), (*Hypocryphalus*),
Stephanopodius 853
ghanaensis Schedl, *Hypothenemus* (= *fuscicollis*) . . 926
ghanaensis Schedl, *Pityophthorus* 1000
ghesquierei Eggers, *Coccotrypes* 603
gibber (Schedl), (*Pterocyclon*), *Monarthrum* 1056
gibber Blackman, *Chramesus* (= *asperatus*) 263
gibber Schedl, *Xyleborus* 741
gibbosus Wood, *Cnesinus* 208
gibbulus Wood, *Cnesinus* 208
gibbus (Chapuis), (*Nemophilus*), *Cnesinus* 208
gibbus Fabricius, (*Hylesinus*), *Camptocerus*
 (= *aeneipennis*) 318
gifensis Murayama, *Phloeosinus* (= *gifuensis*) . . . 247
gifuensis Murayama, *Phloeosinus* 247
giganteus (Schedl), (*Micracis*), *Hylocurus* 425
giganteus Schedl, *Cryphalus* 881
giganteus Schedl, *Ozopemon* 589
giganteus Schedl, *Stephanopodius* 553
gigas Schedl, *Cryphalus* 881
gillerforsi Bright, *Phloeosinus* 247
gilvipes Blandford, (*Xyleborus*), *Ambrosiodmus*
 (= *obliquus*) 677
glabellus Schedl, (*Stephanoderes*), *Hypothenemus*
 (= *hampei*) 930

<i>glaber</i> (Eggers), (<i>Loganius</i>), <i>Cnemomyx</i>	316	<i>Glostatus</i> Schedl	835
<i>glaber</i> (Eichhoff), (<i>Hexacolus</i>), <i>Scolytodes</i>	392	<i>glutae</i> Wood, <i>Pityophthorus</i>	1001
<i>glaber</i> (Schedl), (<i>Eulepiops</i>), <i>Cyrtogenius</i>	565	<i>Glyptoderes</i> Eichhoff, <i>Trypophloeus</i>	842
<i>glaber</i> (Schedl), (<i>Problechilus</i>), <i>Gymnochilus</i>	386	<i>gnarum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056
<i>glaber</i> Eggers, (<i>Xyleborus</i>), <i>Amasa</i> (= <i>schlich</i>)	684	Gnatharus Wood & Yin	1050
<i>glaber</i> Eggers, <i>Scolytodes</i> (= <i>glaberrimus</i>)	393	<i>Gnathoborus</i> Schedl (= <i>Araptus</i>)	953
<i>glaber</i> Eggers, <i>Scolytoplatypus</i>	403	<i>Gnathocortus</i> Schedl (= <i>Gnathotrupes</i>)	1039
<i>glaber</i> Wood, (<i>Prionosecles</i>), <i>Scolytodes</i> (= <i>glabrescens</i>)	392	<i>Gnathocranus</i> Schedl (= <i>Araptus</i>)	953
<i>glaberrimus</i> Schedl, (<i>Xyleborus</i>), <i>Cnestus</i> (= <i>aterrimus</i>)	802	<i>Gnathoglochinus</i> Schedl (= <i>Gnathotrupes</i>)	1039
<i>glaberrimus</i> Wood, <i>Scolytodes</i>	392	Gnatholeptus Blackman	970
<i>glabinus</i> Bright, <i>Corthylus</i> (= <i>mexicanus</i>)	1075	<i>Gnathomimus</i> Schedl (= <i>Gnathotrupes</i>)	1039
<i>glabratellus</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	925	<i>Gnathophorus</i> Schedl (= <i>Pityophthorus</i>)	977
<i>glabratulus</i> (Schedl), (<i>Ctenyophthorus</i>), <i>Pityophthorus</i>	1000	<i>Gnathophthorus</i> Wood (= <i>Pityophthorus</i>)	977
<i>glabratus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	926	<i>Gnathotrichina</i> Balachowsky, part (= <i>Corthylina</i>)	1034
<i>glabratus</i> Browne, <i>Xyleborus</i>	741	<i>Gnathotrichoides</i> Blackman (= <i>Gnathotrichus</i>)	1034
<i>glabratus</i> (Ferrari), (<i>Corthylus</i>), <i>Microcorthylus</i>	1065	Gnathotrichus Eichhoff	1034
<i>glabratus</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	393	° <i>gnathotrichus</i> Schedl, <i>Taphramites</i>	1052
<i>glabratus</i> (Schedl), (<i>Loganius</i>), <i>Cnemomyx</i>	316	Gnathotrupes Schedl	1039
<i>glabratus</i> (Zetterstedt), (<i>Hylurgus</i>), <i>Hylurgops</i>	31	<i>Gnathotrypatus</i> Wood (= <i>Gnathotrupes</i>)	1039
<i>glabratus</i> Eichhoff, <i>Pityophthorus</i>	1000	<i>godmani</i> (Blandford), (<i>Xyleborus</i>), <i>Taurodennus</i>	785
<i>glabratus</i> Eichhoff, <i>Xyleborus</i>	741	<i>golbachii</i> Schedl, <i>Scolytus</i>	331
<i>glabratus</i> Schedl, <i>Hylcops</i>	237	<i>goliathoides</i> Murayama, (<i>Phlocosinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	91
<i>glabratus</i> Schedl, <i>Hypocryphalus</i>	869	<i>goloanus</i> (Browne), (<i>Xyleborus</i>), <i>Eucallacca</i>	690
<i>glabratus</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	925	<i>golovjankoi</i> Pjatnitzky, <i>Orthotomicus</i>	469
<i>glabrellus</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	393	<i>goniomma</i> Enderlein, <i>Aphanarthrum</i> (= <i>mairei</i>)	619
<i>glabrescens</i> Wood, <i>Scolytodes</i>	392	<i>gorggae</i> Schedl, <i>Xyleborus</i>	741
<i>glabricollis</i> Schedl, <i>Araptus</i> (= <i>confinis</i>)	954	<i>gorontalosus</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	524
<i>glabriculum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056	<i>gossypii</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	926
<i>glabrifrons</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056	<i>gracile</i> Eichhoff, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>fasciatum</i>)	1056
<i>glabripennis</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	926	<i>gracilens</i> Wood, <i>Araptus</i>	957
<i>glabripennis</i> Schedl, (<i>Xyleborus</i>), <i>Cnestus</i> (= <i>aterrimus</i>)	802	<i>gracilens</i> Wood, <i>Cnemomyx</i>	316
<i>glabrum</i> Wollaston, <i>Aphanarthrum</i>	618	<i>gracilentum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056
<i>glabrum nudum</i> Israelson, <i>Aphanarthrum</i> (= <i>glabrum</i>)	618	<i>gracilentus</i> (Schedl), (<i>Neodryococtes</i>), <i>Araptus</i>	957
<i>glandis</i> Beeson, (<i>Thammurgides</i>), <i>Coccotrypes</i> (= <i>papuanus</i>)	608	<i>gracilicornis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	807
<i>glaucus</i> (Sampson), (<i>Xyleborus</i>), <i>Amasa</i>	683	<i>gracilicornum</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>exornatum</i>)	1055
<i>globosus</i> (Eggers), (<i>Troglogitica</i>), <i>Acacis</i>	193	<i>gracilior</i> (Schedl), (<i>Apoglostatus</i>), <i>Glostatus</i>	835
<i>globosus</i> (Eichhoff), (<i>Hylesinus</i>), <i>Dendrosinus</i>	237	<i>gracilior</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1056
<i>globosus</i> Hagedorn, (<i>Hapalogenius</i>), <i>Hylesinopsis</i> (= <i>fuscipennis</i>)	94	<i>gracilior</i> Browne, <i>Cryptoxyleborus</i>	828
<i>globosus</i> Hagedorn, <i>Chramesus</i>	265	<i>gracilipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	807
<i>globosus</i> Strohmeier, <i>Glochiphorus</i>	271	<i>gracilipennis</i> Schedl, <i>Tricolus</i>	1043
<i>globulus</i> Blandford, <i>Sphaerotrypes</i>	199	<i>gracilipes</i> (Eichhoff), (<i>Platydictylus</i>), <i>Eccoptopterus</i>	520
<i>globulus</i> Stebbing, (<i>Chramesus</i>), <i>Sphaerotrypes</i> (= <i>querci</i>)	201	<i>gracilis</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xyleborinus</i>	808
<i>globus</i> (Blandford), (<i>Xyleborus</i>), <i>Hadrodennus</i>	519	<i>gracilis</i> (Schedl), (<i>Corthylomimus</i> , <i>Cortheloderes</i>), <i>Corthylus</i>	1074
<i>Glochicopterus</i> Schedl (= <i>Hylesinopsis</i>)	92	<i>gracilis</i> (Schedl), (<i>Lepicerinus</i>), <i>Scolytogenes</i>	861
Glochinocerus Blandford	1064	<i>gracilis</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	481
<i>Glochiphorus</i> Strohmeier	271	<i>gracilis</i> (Schedl), (<i>Neodryococtes</i>), <i>Araptus</i>	957
		<i>gracilis</i> (Schedl), (<i>Ozodendron</i>), <i>Cyrtogenius</i>	565
		<i>gracilis</i> (Schedl), (<i>Phacrylus</i>), <i>Acorthylus</i>	553
		<i>gracilis</i> Blackman, <i>Pseudopityophthorus</i> (= <i>asperulus</i>)	972
		<i>gracilis</i> Blandford, <i>Cnesinus</i>	208

- gracilis* Browne, *Cyrtogenius* (= *gracillinus*) 565
- gracilis* Browne, *Eidophelus* 867
- gracilis* Browne, *Webbia* 831
- gracilis* Eggers, (*Stephanoderes*), *Hypothencmus*
(= *eruditus*) 924
- gracilis* Eggers, *Amphicramus* 1047
- gracilis* Eggers, *Tricolus* 1043
- gracilis* Eichhoff, *Pycnarthrum* (= *hispidum*) 385
- gracilis* LeConte, *Hylastes* 54
- gracilis* Niisima, *Polygraphus* 286
- gracilis* Schedl, *Lamurgus* 408
- gracilis* Schedl, *Scolytodes* 393
- gracilis* Swaine, *Pityophthorus* (= *murrayanae*) 1014
- gracilis* Wood, *Chramesus* 265
- gracilis* Wood, *Dendrocraniulus* 551
- gracillinus* Wood, *Cyrtogenius* 565
- granatus* Schedl, *Xyleborus* 741
- grande* Schedl, *Rhopalopselion* 96
- grandiclava* Thomson, *Polygraphus* 286
- grandiclavatus* Eggers, *Microcorthyus* 1066
- grandicollis* (Eichhoff), (*Tomicus*), *Ips* 506
- grandicollis* Swaine, *Hylurgops* (= *incomptus*) 34
- grandidentatus* Schedl, *Scolytotplatypus*
(= *eichelbaumi*) 402
- grandis* (Beeson), (*Pelicerus*), *Cyrtogenius* 565
- grandis* (Eggers), (*Pocilips*), *Coccotrypes* 604
- grandis* (Erichson), (*Phlocotrupes*), *Phloeoborus* 99
- grandis* (Schedl), (*Hexacolus*), *Scolytodes* 393
- grandis* (Schedl), (*Neodryocoetes*), *Araptus* 957
- grandis* (Schedl), (*Xyleboricus*), *Arixyleborus* 666
- grandis* Blackman, *Pityophthorus* 1001
- grandis* Browne, *Acanthotomicus* 481
- grandis* Chamberlin, *Cryphalus* (= *ruficollis*) 892
- grandis* Eichhoff, *Xyleborus* 741
- grandis* Kurebuzov, *Scolytus* (= *trispinosus*) 378
- grandis* Nunberg, *Polygraphus* 287
- grandis* Schedl, (*Cryphalomorphus*), *Hypothencmus*
(= *ingens*) 931
- grandis* Schedl, *Cnesinus* 209
- grandis* Schedl, *Cryphalominus* 835
- grandis* Schedl, *Hypothencmus* 926
- grandis* Schedl, *Micracis* 432
- grandis* Schedl, *Scolytodes* (= *eximius*) 392
- grandis* Schedl, *Sphaerotrypes* 199
- grandis* Schedl, *Trypophloeus* 844
- grandis* Swaine, *Pseudohylesinus* (= *sericeus*) 112
- grangeri* (Peyerimhoff), (*Lymantor*), *Triotenius* 541
- graniceps* (Eichhoff), (*Dryocoetes*), *Coccotrypes* 604
- graniceps* Eichhoff, *Coccotrypes* (= *graniceps*) 604
- granicollis* (LeConte), (*Xyleborus*), *Dryocoetes* 582
- granicollis* Eichhoff, (*Phlocophthorus*), *Phlocotribus*
(= *frontalis*) 222
- granifer* (Eichhoff), (*Xyleborus*), *Arixyleborus* 666
- granifer* borneensis Schedl, *Arixyleborus*
(= *granifer*) 666
- granifer* Browne, *Diamerus* 195
- graniger* Eichhoff, (*Dendroctonus*), *Phloeosinus*
(= *dentatus*) 246
- graniger* Schedl, *Xyleborus* 741
- granistriatus* (Schedl), (*Orosiotes*), *Cyrtogenius* 565
- granistriatus* Eggers, (*Xyleborus*), *Arixyleborus*
(= *imitator*) 667
- graniticus* Wood, *Pseudothysanops* 414
- granosus* (Schedl), (*Xyleborus*), *Eucwallacea* 690
- granosus* Chapuis, *Hylastes* (= *porculus*) 61
- granosus* Eichhoff, *Phloeoborus* 99
- granosus* Schedl, *Phlocosinus* 247
- granularis* Schedl, *Xyleborus* (= *rolulus*) 783
- granulatum* Blandford, *Styphlosoma* 950
- granulatum* Bright, *Monarthrum* 1056
- granulatum* Eggers, *Trypodendron* (= *lineatum*) 643
- granulatus* (Eggers), (*Pagiocerus*), *Chramesus* 265
- granulatus* (Ferrari), (*Xylocleptes*),
Acanthotomicus 481
- granulatus* (LeConte), (*Hylastes*),
Pseudohylesinus 110
- granulatus* (Lepesme), (*Kissophagus*),
Hylesinopsis 94
- granulatus* (Ratzeburg), (*Bostrichus*),
Trypophloeus 844
- ° *granulatus* (Schedl), (*Hylescrites*), *Hylurgops* 33
- granulatus* (Wood), (*Cryphalomorphus*),
Scolytogenes 862
- granulatus* Blackman, *Pseudopityophthorus* 973
- granulatus* Bruck, *Phloeosinus* (= *frontalis*) 247
- granulatus* Eggers, *Diamerus* 195
- granulatus* Eggers, *Hylastinus* (= *obscurus*) 65
- granulatus* Eggers, *Hyorrhynchus* (= *leicisi*) 192
- granulatus* Eggers, *Phloeoborus* 99
- granulatus* Eggers, *Polygraphus* 287
- granulatus* Niisima, *Polygraphus* (= *japonicus*) 288
- granulatus* Schedl, (*Neodryocoetes*), *Araptus*
(= *tenellus*) 962
- granulatus* Schedl, *Corthyus* 1074
- granulatus* Schedl, *Hypocryphalus* 869
- granulatus* Schedl, *Ozopemon* 589
- granulatus* Swaine, *Pityophthorus* (= *consimilis*) 991
- granulatus* Wood, *Cactopinus* 435
- granulicauda* (Eggers), (*Xyleborus*),
Dryocoetoides 657
- granulicauda* Schedl, *Arixyleborus* 666
- granulicauda* Schedl, (*Pocilips*),
Coccotrypes (= sp.?) 591
- granulicauda* Schedl, *Pityophthorus* (= *terebrans*) 1031
- granulicauda* Schedl, *Polygraphus* 288
- granulicollis* (Schedl), (*Neodiamerus*), *Acacacis* 193
- granulicollis* Schedl, (*Dendriops*), *Cosmoderes*
(= *monilicollis*) 901
- granulicollis* Schedl, *Thammurgus* 543
- granulifer* (Beeson), (*Pelicerus*), *Cyrtogenius* 565
- granulifer* (Eggers), (*Xyleborus*), *Arixyleborus* 666
- granulifer* Eggers, *Polygraphus* 288
- granulifer* Eggers, *Strombophorus* 203

<i>granulifer</i> Motschulsky, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	90	<i>groutellei</i> (Blandford), (<i>Ethadopselaphus</i>), <i>Dactylipalpus</i>	102
<i>granulifer</i> Reitter, <i>Scolytus</i> (= <i>jaroschewskii</i>)	336	<i>groutellei</i> Blandford, <i>Amphicranus</i>	1047
<i>granulifer</i> Wood, <i>Cactopinus</i>	435	<i>guadeloupanus</i> Wood, <i>Araptus</i>	957
<i>granulifer</i> Wood, <i>Corthylus</i>	1074	<i>guadeloupensis</i> Eggers, <i>Cuesinus</i>	209
<i>granulifer</i> Wood, <i>Pseudopityophthorus</i>	973	<i>guadeloupensis</i> Numberg, (<i>Pityophthorus</i>), <i>Araptus</i> (= <i>eggersianus</i>)	956
<i>granulifer</i> Wood, <i>Thysanocs</i>	420	<i>guadeloupensis</i> Schedl, (<i>Brachydendrus</i>), <i>Araptus</i> (= <i>guadeloupanus</i>)	957
<i>granulipennis</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	957	<i>guadeloupensis</i> Schedl, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>laevigatus</i>)	958
<i>granulipennis</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>andamanensis</i>)	686	<i>guadeloupensis</i> Schedl, <i>Hypothenemus</i> (= <i>plumeriae</i>)	939
<i>granulipennis</i> Schedl, <i>Chramesus</i>	266	<i>guanaguatensis</i> Duges, <i>Xyleborus</i> (= <i>rolulus</i>)	782
<i>granulipennis</i> Schedl, <i>Phloeosinus</i>	247	<i>guatemalensis</i> (Hopkins), (<i>Xylocleptes</i>), <i>Dendrocraniulus</i>	551
<i>granulipennis</i> Schedl, <i>Pityophthorus</i> (= <i>terebrans</i>)	1031	<i>guatemalensis</i> (Wood), (<i>Leperisinus</i>), <i>Hylesinus</i>	80
<i>granulipennis</i> Schedl, <i>Pityophthorus</i>	1001	<i>guatemalensis</i> Blandford, <i>Pityophthorus</i>	1001
<i>granulipennis</i> Schedl, <i>Xylocleptes</i>	549	<i>guatemalensis</i> Hopkins, (<i>Ambrosiodmus</i>), <i>Ambrosiodmus</i> (= <i>hagedorni</i>)	674
<i>granulipes</i> Schedl, <i>Xyleborus</i>	741	<i>guatemalensis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915
<i>granulosus</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	1001	<i>guatemalensis</i> Hopkins, <i>Dryocoetoides</i> (= <i>capucinus</i>)	657
<i>granulosus</i> Schedl, <i>Xyleborus</i> (= <i>partitus</i>)	758	<i>guayanensis</i> Eggers, <i>Corthylus</i> (= <i>papulans</i>)	1076
<i>grauurus</i> Browne, <i>Xyleborus</i>	741	<i>guayanensis</i> Eggers, <i>Phloeoborus</i>	99
<i>graphus</i> Duftschmidt, <i>Hylastes</i> (= <i>angustatus</i>)	44	<i>guayanensis</i> Eggers, <i>Xyleborus</i> (= <i>tumucensis</i>)	779
<i>gratiosus</i> Schedl, <i>Xyleborus</i>	741	<i>gueriniae</i> (Schedl), (<i>Bostrichips</i>), <i>Pseudothysanocs</i>	414
<i>gratus</i> Schedl, <i>Xyleborus</i>	741	<i>guianae</i> Blackman, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>hymenaeae</i>)	958
<i>gravelyi</i> Wichmann, <i>Xyleborus</i>	741	<i>guiboutiae</i> (Schedl), (<i>Microcryphalus</i>), <i>Ernoeladius</i>	857
<i>gravidus</i> (Blandford), (<i>Dryocoetes</i>), <i>Ozopemon</i>	589	<i>gildi</i> Blackman, <i>Ips</i> (= <i>latidens</i>)	511
<i>gravidus</i> (Blandford), (<i>Xyleborus</i>), <i>Xylosandrus</i>	796	<i>guillebeui</i> Reitter, (<i>Phloeophthorus</i>), <i>Phloeotribus</i> (= <i>pubifrons</i>)	228
<i>grayi</i> Schedl, <i>Cryphalus</i> (= <i>tetricus</i>)	897	<i>guineensis</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhupidion</i>	699
<i>grenadensis</i> Hopkins, <i>Xyleborus</i> (= <i>rolulus</i>)	783	<i>guineensis</i> Hagedorn, (<i>Hyphaeni</i>), <i>Coccotrypes</i> (= <i>sp.</i> ?)	591
<i>gretschkini</i> Sokanovskii, <i>Scolytus</i>	331	<i>gummensis</i> (Murayama), (<i>Pruniphagus</i>), <i>Xylechinus</i>	116
<i>Gretschkinia</i> Sokanovskii (= <i>Pseudothysanocs</i>)	411	<i>gundlachi</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>lecontei</i>)	675
<i>griseopubescens</i> Schedl, <i>Diamerus</i>	195	<i>gunneri</i> Schedl, <i>Pityophthorus</i>	1001
<i>griseum</i> Schedl, <i>Ctonoxylon</i>	837	<i>guttifer</i> (Schedl), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	667
<i>griseus</i> Blackburn, (<i>Hypothenemus</i>), <i>Hypocryphalus</i> (= <i>mangiferae</i>)	870	<i>guyanaensis</i> (Schedl), <i>Scolytodes</i>	393
<i>griseus</i> Eggers, <i>Polygraphus</i> (= <i>poligraphus</i>)	295	Gymnochilus Eichhoff	386
<i>griseus</i> Marsham, (<i>Ips</i>), <i>Hylesinus</i> (= <i>varius</i>)	86	<i>haagi</i> Eichhoff, (<i>Dendroctonus</i>), <i>Phloeosinus</i> (= <i>dentatus</i>)	246
<i>griseus</i> Swaine, <i>Pseudohylesinus</i> (= <i>nebulosus nebulosus</i>)	110	<i>haberkorni</i> Eggers, <i>Xyleborus</i>	741
<i>grisseopuberulus</i> Schedl, <i>Coccotrypes</i> (= <i>carpophagus</i>)	597	<i>habilis</i> Wood, <i>Scolytodes</i>	393
<i>grobleri</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	862	<i>haddeni</i> Schedl, <i>Xyleborus</i>	742
<i>grobleri</i> Schedl, <i>Microcryphalus</i>	411	Hadrodemius Wood	818
<i>grossepunctatus</i> (Schedl), (<i>Policerus</i>), <i>Cyrtogenius</i>	565	<i>haemorrhoidalis</i> Marsham, (<i>Ips</i>), <i>Hylesinus</i> (= <i>varius</i>)	89
<i>grossepunctatus</i> Browne, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>puncticollis</i>)	865	<i>haemorrhous</i> Schmidberger, <i>Scolytus</i> (= <i>rugulosus</i>)	369
<i>grossepunctatus</i> Eggers, <i>Ozopemon</i>	589	<i>haesitus</i> Schedl, <i>Xyleborus</i>	742
<i>grossepunctatus</i> Schedl, <i>Thannurgus</i>	543	<i>hagedorni</i> (Iglesias), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	674
<i>grossmanni</i> Schedl, <i>Xyleborus</i>	741	<i>hagedorni</i> (Schedl), (<i>Prionosceles</i>), <i>Scolytodes</i>	393
<i>grossopunctatus</i> Schedl, (<i>Xyleborus</i>), <i>Coccotrypes</i> (= <i>gedeanus</i>)	603	<i>hagedorni</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1057
<i>grossopunctatus</i> Schedl, <i>Dendrocraniulus</i> (= <i>macilentus</i>)	552		
<i>grossus</i> Chapuis, <i>Phloeoborus</i>	99		
<i>grothi</i> Hagedorn, (<i>Cryphalus</i>), <i>Trypophloeus</i> (= <i>binodulus</i>)	844		

<i>hagedornii</i> Eggers, (Coccotrypes), <i>Cryphalus</i> (= <i>walkeri</i>)	895	<i>herbertfranzi</i> (Schedl), (<i>Gnathotrichus</i>), <i>Gnathotrypes</i>	1040
<i>hagedornii</i> Eggers, <i>Sphacrotypes</i>	199	<i>hercegovinensis</i> (Seitner), (<i>Phloeophthorus</i>), <i>Phloeotribus</i>	222
<i>hagedornii</i> Stebbing, <i>Xyleborus</i> (= <i>hagedornianus</i>)	742	<i>hercegovinensis</i> Eggers, <i>Phloeosinus</i> (= <i>bicolor</i>)	243
<i>hagedornianus</i> Schedl, <i>Xyleborus</i>	742	<i>hermosus</i> Wood, <i>Pityophthorus</i>	1002
<i>Hagedornius</i> Lucas (= <i>Pityophthorus</i>)	976	<i>hermosus</i> Wood, <i>Scolytus</i>	331
<i>halli</i> Schedl, <i>Xyleborus</i>	742	<i>herrarai</i> Hopkins, <i>Pityophthorus</i> (= <i>chalcoensis</i>)	955
<i>Halystus</i> Schedl	253	<i>hesperidum</i> Wollaston, <i>Aphanarthrum</i>	615
<i>hamamelidis</i> Blackman, <i>Pityophthorus</i> (= <i>latus</i>)	1006	<i>hesperius</i> Bright, <i>Pityophthorus</i> (= <i>concarus</i>)	959
<i>hamamelidis</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>hesperius</i> Bright, <i>Pityotrichus</i>	970
<i>hamatum</i> Schedl, <i>Ctenoxylon</i>	835	<i>Heteroborips</i> Reitter (= <i>Xyleborus</i>)	704
<i>hamatus</i> Hagedorn, <i>Scolytoplatypus</i> (= <i>eutomoides</i>)	403	<i>heterodon</i> Wachtl, (<i>Tomicus</i>), <i>Pityokteines</i> (= <i>spinidens</i>)	466
<i>hamatus</i> LeConte, (<i>Xyleborus</i>), <i>Pityogenes</i> (= <i>carinulatus</i>)	445	<i>heterodoxus</i> (Casey), (<i>Renocis</i>), <i>Chaetophloeus</i>	272
<i>hampei</i> (Ferrari), (<i>Cryphalus</i>), <i>Hypothenemus</i>	927	<i>heterolepis</i> Costa Lima, <i>Hypothenemus</i> (= <i>arecae</i>)	907
<i>Hapalogenius</i> Hagedorn (= <i>Rhopalopselion</i>)	96	<i>heveae</i> (Hagedorn), (<i>Stephanoderes</i>), <i>Cryptocarenum</i>	903
<i>Hapalophloeus</i> Schedl (= <i>Hylesinopsis</i>)	92	<i>heveae</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	505
<i>harnedi</i> (Blackman), (<i>Micracis</i>), <i>Hylocurus</i>	425	<i>heveti</i> (Stebbing), (<i>Dryocoetes</i>), <i>Taphrorhynchus</i>	559
<i>harongae</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	451	Hexacolidae Chapuis (= <i>Ctenophorini</i>)	353
<i>harringtoni</i> (Blackman), (<i>Tachyderes</i>), <i>Cryptocarenum</i>	903	Hexacolidae Chapuis (= <i>Ctenophorini</i>)	353
<i>harringtoni</i> Blackman, <i>Phloeotribus</i>	222	Hexacolinae Chapuis (= <i>Ctenophorini</i>)	353
<i>harringtoni</i> Blackman, <i>Pseudochramesus</i>	262	Hexacolini Chapuis	353
<i>harunganae</i> (Schedl), (<i>Micracis</i>), <i>Pseudomicracis</i>	429	<i>Hexacolinus</i> Schedl (= <i>Scolytodes</i>)	355
<i>hashimotoi</i> Browne, <i>Xyleborus</i>	742	<i>Hexacolus</i> Eichhoff (= <i>Scolytodes</i>)	357
<i>hastatum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700	<i>heydeni</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>acuminatus</i>)	459
<i>hatanakai</i> Browne, <i>Xyleborus</i>	742	<i>hicoriae</i> LeConte, <i>Chramesus</i>	266
<i>hatanakai</i> Browne, <i>Webbia</i>	831	<i>hidalgoensis</i> Blackman, <i>Pityophthorus</i> (= <i>nocturnus</i>)	1016
<i>hattori</i> Kono, <i>Cryphalus</i> (= <i>piceae</i>)	555	<i>hiika</i> Samuelson, <i>Xyleborus</i>	742
<i>hauseri</i> Reitter, <i>Ips</i>	509	<i>himalabietis</i> Beeson, <i>Hylastes</i> (= <i>brunneus</i>)	50
<i>hawaiiensis</i> Perkins, <i>Xyleborus</i>	742	<i>himalayensis</i> Beeson, (<i>Thamurgides</i>), <i>Coccotrypes</i> (= <i>nubilus</i>)	605
<i>hawaiiensis</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943	<i>himalayensis</i> Eggers, <i>Ambrosiodinus</i> (= <i>minor</i>)	676
<i>heathi</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	923	<i>himalayensis</i> Stebbing, <i>Hylastes</i> (= <i>Curculionidae</i>)	43
<i>hebes</i> Schedl, <i>Phloeotribus</i>	222	<i>himalayensis</i> Stebbing, (<i>Cryphalus</i>), <i>Pityophthorus</i> (= <i>deodara</i>)	995
<i>hectographus</i> Reitter, <i>Dryocoetes</i>	552	<i>himalayensis</i> Stebbing, <i>Polygraphus</i> (= <i>longifolia</i>)	259
<i>hederae</i> (Schmidt), (<i>Hylesinus</i>), <i>Kissophagus</i>	65	<i>himalayensis</i> Stebbing, <i>Scolytoplatypus</i> (= <i>raja</i>)	406
<i>helpferi</i> Eggers, <i>Sphacrotypes</i>	199	<i>himalayensis</i> Strohmeyer, <i>Dryocoetes</i>	553
<i>helpferi</i> Villa, (<i>Hylurgus</i>), <i>Hylurgops</i> (= <i>palliatus</i>)	35	<i>hintzi</i> Schedl, <i>Pityophthorus</i>	1002
<i>heliura</i> Wood, <i>Pseudothysanoes</i>	414	<i>hirsutipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	796
<i>helopioides</i> Schedl, <i>Cryphalus</i>	551	<i>hirsutulus</i> Schedl, <i>Arixyleborus</i>	667
<i>helveticus</i> Guillebeau, (<i>Phloeophthorus</i>), <i>Phloeotribus</i> (= <i>cristatus</i>)	220	<i>hirsutum</i> Hagedorn, <i>Ctenoxylon</i> (= <i>flavescens</i>)	537
<i>hemibitalis</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>usagarius</i>)	665	<i>hirsutus</i> (Nobuchi), (<i>Taenioglyptes</i>), <i>Cryphalus</i>	551
<i>Hemicyphalus</i> Schedl	866	<i>hirsutus</i> (Schedl), (<i>Dryocoetes</i>), <i>Dryocoetiops</i>	557
<i>Hemihylesinus</i> Schedl (= <i>Hylesinopsis</i>)	92	<i>hirsutus</i> (Schedl), (<i>Trypographus</i>), <i>Hylesinopsis</i>	94
<i>hendrickxi</i> Schedl, (<i>Streptocranus</i>), <i>Coptoborus</i> (= <i>usagarius</i>)	664	<i>hirsutus</i> (Wood), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	930
<i>henscheli</i> Knotek, <i>Hylesinus</i> (= <i>varius</i>)	59	<i>hirsutus</i> Blackman, <i>Scolytoplatypus</i>	403
<i>henscheli</i> Reitter, <i>Carphoborus</i> (= <i>bonnairei</i>)	303	<i>hirsutus</i> Bright, <i>Pseudopityophthorus</i> (= <i>tenuis</i>)	976
<i>henscheli</i> Seitner, <i>Pityophthorus</i>	1001	<i>hirsutus</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>concinuus</i>)	500
<i>henschi</i> Reitter, <i>Phloeosinus</i>	247	<i>hirsutus</i> Lea, (<i>Xylopertha</i>), <i>Xyleborus</i> (= <i>perforans</i>)	761
<i>henshawi</i> Hopkins, (<i>Ericryphalus</i>), <i>Cryphalus</i> (= <i>sylvicola</i>)	596	<i>hirsutus</i> Schedl, <i>Abiophagus</i>	73
<i>herbellae</i> Strohmeyer, <i>Pityogenes</i> (= <i>bistridentatus</i>)	444	<i>hirsutus</i> Schedl, <i>Dendrosinus</i> (= <i>ater</i>)	236

<i>hirsutus</i> Schedl. <i>Glostatus</i>	855	<i>hopeae</i> Browne, <i>Xyleborus</i>	742
<i>hirsutus</i> Schedl. <i>Xylechinosis</i>	105	<i>hopehi</i> Schedl. <i>Phloeosinus</i>	248
<i>hirsutus</i> Wood, <i>Bothrostermus</i>	215	<i>hopkinsi</i> Beeson, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739
<i>hirsutus</i> Wood, <i>Scolytodes</i>	393	<i>hopkinsi</i> Blackman, <i>Pityophthorus</i> (= <i>digestus</i>)	995
<i>hirtellum</i> Schedl. <i>Ctonoxylon</i>	838	<i>hopkinsi</i> Blackman, <i>Pseudothysanoes</i>	414
<i>hirtellus</i> (LeConte), (<i>Micracis</i>), <i>Hyloterus</i>	425	<i>hopkinsi</i> Browne, <i>Hypothenemus</i> (= <i>multidentatus</i>)	936
<i>hirtellus</i> Eichhoff, <i>Taphrorychus</i>	559	<i>hopkinsi</i> Swaine, <i>Pityogenes</i>	452
<i>hirtellus</i> Schedl. <i>Cyrtogenius</i>	565	<i>Hoplites</i> Eggers (= <i>Cladoctonus</i>)	238
<i>hirtellus</i> Schedl. <i>Phloeotribus</i>	222	<i>Hoplitantus</i> Wood (= <i>Cladoctonus</i>)	238
<i>hirtellus</i> Schedl. (<i>Xyleborus</i>), <i>Theoborus</i> (= <i>theobromae</i>)	661	<i>Hoplitophthorus</i> Wood (= <i>Cladoctonus</i>)	238
<i>hirtellus</i> Wood, <i>Pityoborus</i>	968	<i>hoppingi</i> Lanier, <i>Ips</i>	509
<i>hirticeps</i> LeConte, <i>Pityophthorus</i> (= <i>pulehellus</i>)	1023	<i>hoppingi</i> Swaine, <i>Phloeosinus</i>	248
<i>hirtipennis</i> (Eggers), (<i>Xyleborus</i>), <i>Arixyleborus</i>	667	<i>hoppingi</i> Swaine, <i>Polygraphus</i>	288
<i>hirtipennis</i> Schedl. <i>Campocerus</i> (= <i>suturalis</i>)	320	<i>hornus</i> (Schedl), (<i>Eidophelus</i>), <i>Cyrtogenius</i>	565
<i>hirtipennis</i> Schedl. <i>Hypothenemus</i>	930	<i>horridatus</i> Wood, <i>Xyleborus</i>	742
<i>hirtipes</i> Schedl. <i>Xyleborus</i>	742	<i>horridicus</i> Wood, <i>Xyleborus</i>	743
<i>hirtum</i> (Hagedorn), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700	<i>horridulus</i> Browne, <i>Xyleborus</i>	743
<i>hirtuosus</i> Beeson, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>hirtum</i>)	700	<i>horridus</i> (Eggers), (<i>Hylesinus</i>), <i>Hylesinopsis</i>	94
<i>hirtus</i> (Eggers), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	565	<i>horridus</i> Eichhoff, <i>Cryphalus</i>	881
<i>hirtus</i> (Wood), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	862	<i>horridus</i> Eichhoff, <i>Xyleborus</i>	743
<i>hirtus</i> Wood, <i>Phloeotribus</i>	222	<i>horridus</i> Graham, <i>Cryphalus</i> (= <i>theobromae</i>)	897
<i>hispaniolum</i> Bright, <i>Liparthrum</i>	277	<i>horyurensis</i> Murayama, <i>Polygraphus</i>	288
<i>hispaniolus</i> Bright, <i>Pityophthorus</i>	1002	<i>hostilis</i> (Blackman), (<i>Neodryocoetes</i>), <i>Araptus</i>	957
<i>hispidulus</i> Eggers, <i>Phloeotribus</i>	222	<i>hostilis</i> Wood, <i>Microcthyus</i>	1066
<i>hispidulus</i> LeConte, (<i>Cryphalus</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	914	<i>hova</i> Schaufuss, <i>Scolytotplatypus</i>	403
<i>hispidulus</i> Thomson, <i>Crypturgus</i>	624	<i>hova</i> Schedl. <i>Xyleborus</i>	743
<i>hispidum</i> (Ferrari), (<i>Hypoborus</i>), <i>Pycnarthrum</i>	384	<i>howardi</i> Hopkins, <i>Xyleborus</i> (= <i>iutrusus</i>)	745
<i>hispidus</i> (Eggers), (<i>Metalhylesinus</i>), <i>Hylesinopsis</i>	94	<i>howdenae</i> Bright, <i>Xyleborus</i>	743
<i>hispidus</i> (Klug), (<i>Hylesinus</i>), <i>Diamerus</i>	196	<i>howdeni</i> Bright, <i>Chaetophloeus</i>	272
<i>hispidus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>arecae</i>)	907	<i>huachucae</i> Blackman, <i>Pseudothysanoes</i>	414
<i>hispidus</i> Eggers, <i>Cuesinus</i>	209	<i>huachucae</i> Wood, <i>Monarthrum</i>	1057
<i>hispidus</i> Eggers, <i>Kissophagus</i> (= <i>hederae</i>)	68	<i>huangi</i> (Browne), (<i>Xyleborus</i>), <i>Coptodryas</i>	824
<i>hispidus</i> Eggers, <i>Pseudopityophthorus</i>	973	<i>huapiae</i> (Schedl), (<i>Phthorophloeus</i>), <i>Phloeotribus</i>	222
<i>hispidus</i> Eggers, <i>Strombophorus</i>	203	<i>hubbardi</i> Blackman, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>politus</i>)	961
<i>hitcaoca</i> Beeson, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915	<i>hubbardi</i> Blackman, <i>Pityophthorus</i> (= <i>infulatus</i>)	1003
<i>hobohmi</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	861	<i>hubbardi</i> Hopkins, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>hoegei</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1057	<i>hubbardi</i> Hopkins, <i>Xyleborus</i> (= <i>colvulus</i>)	783
<i>hoferi</i> Blackman, (<i>Lepersinus</i>), <i>Hylesinus</i> (= <i>californicus</i>)	76	<i>hubbardi</i> Schwarz, (<i>Bothrostermus</i>), <i>Pagiocerus</i> (= <i>frontalis</i>)	213
<i>hoferi</i> Blackman, <i>Phloeosinus</i>	248	<i>hubbardi</i> Schwarz, <i>Cactopinus</i>	435
<i>holdhausi</i> Wichmann, <i>Trypophloeus</i> (= <i>alni</i>)	842	<i>hudsonicus</i> LeConte, <i>Ips</i> (= <i>perturbatus</i>)	516
<i>Holonthogaster</i> Gemminger & Harold (= <i>Hyledius</i>)	259	<i>huelmetanus</i> Schedl, <i>Dendrocranius</i> (= <i>maurus</i>)	552
<i>holtzi</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>aegir</i>)	670	<i>humilis</i> (Blanchard), (<i>Hylastes</i>), <i>Xylechinosis</i>	105
<i>holtzi</i> Strohmeier, <i>Thamurgus</i> (= <i>delphinii</i>)	543	<i>humilis</i> Blackman, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>hymenaeae</i>)	958
<i>Honarius</i> Broun (= <i>Chaetoptelius</i>)	103	<i>hunauensis</i> Browne, <i>Xyleborus</i>	743
<i>Homocoryphalus</i> Lindemann (= <i>Hypothenemus</i>)	904	<i>hunteri</i> Swaine, <i>Ips</i>	509
<i>hondurensis</i> Wood, <i>Micracisella</i>	430	<i>hyaspistis</i> Schedl, <i>Tiarophorus</i>	539
<i>hondurensis</i> Wood, <i>Pityoborus</i>	968	<i>hybridus</i> Blandford, <i>Amphicranus</i>	1047
<i>hondurensis</i> Wood, <i>Pseudopityophthorus</i>	973	<i>hybridus</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	696
<i>hoodi</i> Blackman, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>hymenaeae</i>)	958	° <i>hydropicus</i> (Wickham), (<i>Apidocephalus</i>), <i>Hylesinus</i>	80
		<i>Hylastes</i> Erichson	43
		<i>Hylastes</i> LeConte (= <i>Hylastini</i>)	29

<i>Hylastides</i> LeConte (=Hylastini)	29	<i>Hypoborina</i> Nusslin (=Hypoborini)	271
<i>Hylastini</i> LeConte	29	<i>Hypoborinae</i> Nusslin (=Hypoborini)	271
<i>Hylastinoides</i> LeConte (=Hylesinini)	64	<i>Hypoborini</i> Nusslin	271
<i>Hylastinoides</i> Spessivitzev (=Alniaphagus)	73	<i>Hypoborus</i> Erichson	279
<i>Hylastinus</i> Bedel	64	<i>Hypocryphalus</i> Hopkins	868
<i>Hylastites</i> Hagedorn (=Hylurgops)	30	<i>Hypopityophthorus</i> Bright (=Pityophthorus)	977
<i>Hyledius</i> Sampson	259	<i>Hypothenemus</i> Westwood	904
<i>Hyleops</i> Schedl	237	<i>Hypothenoides</i> Hopkins (=Scolytogenes)	858
<i>Hylescierites</i> Schedl (=Hylurgops)	30	<i>hystricoides</i> Browne, Xyleborus	743
<i>Hylesinae</i> Erichson (=Hylesininae)	28	<i>hystrix</i> (Eggers), (Adiaeretus), Hypothenemus	931
<i>Hylesinen</i> Erichson (=Hylesininae)	28	<i>hystrix</i> (LeConte), (Hylesinus), Chaetophloeus	273
<i>Hylesini</i> Erichson (=Hylesinae, Hylesini)	28, 64	<i>hystrix</i> (Schedl), (Xyleborus), Prcmuobius	653
<i>Hylesinidae</i> Erichson (=Hylesininae)	28	<i>hystrix</i> Abeille de Perrin, Cisurgus (=wollastini)	631
<i>Hylesinoides</i> Erichson (=Hylesininae)	28	<i>hystrix</i> Wood, Phloeotribus	222
<i>hylesiniformis</i> (Schedl), (Ptilopodius),		iconographus Dejean, (Bostrichus), Ips	
<i>Cryphlyophthorus</i>	274	(=acuminatus)	455
<i>Hylesinina</i> Erichson (=Hylesininae)	28	<i>idahoensis</i> Hopkins, Procryphalus (=mucronatus)	848
<i>Hylesininae</i> Erichson	28	<i>idoneus</i> Blackman, Pityophthorus (=digestus)	995
<i>Hylesinini</i> Erichson	28, 64	<i>ignobilis</i> Perkins, Xyleborus	743
<i>Hylesinites</i> Germar (=Hylurgops)	29	<i>ignotus</i> (Schedl), (Micracis), Pseudomicracis	429
<i>hylesinopsis</i> (Schedl), (Cryphalomorphus),		<i>ignotus</i> Bright, Tricolus (=perdilignis)	1044
<i>Scolytogenes</i>	862	<i>ignotus</i> Schedl, Pityophthorus	1002
<i>Hylesinopsis</i> Eggers	92	<i>iheringi</i> Iglesias, Xyleborus	743
<i>Hylesinosoma</i> Lea (=Aricerus)	217	<i>ikelaensis</i> Nunberg, Pityophthorus	1002
<i>Hylesinus</i> Erichson (=Hylesininae)	28	<i>illepidus</i> Schedl, (Xyleborus), Ambrosiodmus	
<i>Hylesinus</i> Fabricius	74	(=obliquus)	678
<i>Hyllocuri</i> Eichhoff (=Micracini)	408	<i>illustrius</i> (Schedl), (Xyleborus), Euwallacea	690
<i>Hyllocuridae</i> Eichhoff (=Micracini)	408	<i>imbellis</i> Blandford, Xyleborus	743
<i>Hyllocurinae</i> Eichhoff (=Micracini)	408	<i>imbricorus</i> Eichhoff, Phloeoborus (=asper)	95
<i>hyllocuroides</i> Wood, Pityophthorus	1002	<i>imitans</i> (Eggers), (Phloeosinus), Hyledius	260
<i>Hyllocurosoma</i> Eggers (=Scolytodes)	388	<i>imitans</i> (Eggers), (Prionosceles), Scolytodes	393
<i>Hyllocurus</i> Eichhoff	422	<i>imitans</i> (Eggers), (Trigonogenius), Pityophthorus	1002
<i>Hylonius</i> Nunberg (=Xylocleptes)	547	<i>imitans</i> Eggers, (Pocillips), Coccotrypes	
<i>Hyloperus</i> Browne (=Cryptocurus)	72	(=ghesquieri)	603
<i>Hylurdretonus</i> Schedl	186	<i>imitans</i> Eggers, Dactylipalpus	102
<i>Hylurgidae</i> LeConte (=Tomicini)	103	<i>imitans</i> Eggers, (Xyleborus), Euwallacea (=piceus)	693
<i>Hylurgides</i> LeConte (=Tomicini)	103	<i>imitans</i> Eggers, Microborus (=aberrans)	383
<i>Hylurgini</i> LeConte (=Tomicini)	103	<i>imitans</i> Eggers, Sphaecotrypes (=pila)	200
<i>Hylurgina</i> LeConte (=Tomicini)	103	<i>imitans</i> Eichhoff, Eidophelus	867
<i>hylurgoides</i> (Schedl), (Coptodryas), Cnemonyx	316	<i>imitans</i> Schedl, Hypocryphalus	869
<i>hylurgoides</i> Schedl, Chramesus	266	<i>imitans</i> Wood, Gnathotrichus	1035
<i>Hylurgonotus</i> Schedl	186	<i>imitator</i> (Eggers), (Webbia), Arixyleborus	667
<i>Hylurgopina</i> Balachowsky (=Hylastini)	29	<i>imitator</i> (Reitter), (Hylastes), Hylurgops	33
<i>Hylurgopinus</i> Swaine	106	<i>imitator</i> Schedl, (Xyleborus), Dryocoetoides	
<i>Hylurgops</i> LeConte	29	(=asperulus)	656
<i>hylurgulus</i> (Eggers), (Phloeosinus), Hyledius	260	<i>imitator</i> Sokanovskii, Alniaphagus (=alni)	73
<i>Hylurgulus</i> Eggers (=Hyledius)	257	<i>imitatrix</i> (Schedl), (Erioschidias), Cosmoderes	901
<i>hylurgulus</i> Schedl, Phloeotribus	222	<i>imitatrix</i> (Schedl), (Neodryocoetes), Araptus	958
<i>Hylurgus</i> Latreille	119	<i>immanis</i> Blackman, Pityophthorus	1002
<i>hymenaeae</i> (Eggers), (Neodryocoetes), Araptus	957	<i>immanis</i> Wood, Scolytodes	393
<i>Hyorrhynchinae</i> Hopkins (=Hyorrhynchini)	191	<i>immaturus</i> (Schedl), (Hapalogenius),	
<i>Hyorrhynchini</i> Hopkins	191	<i>Rhopalopselion</i>	96
<i>Hyorrhynchus</i> Blandford	191	<i>immaturus</i> Blackburn, Xyleborus (=perforans)	761
<i>Hypaspistes</i> Hagedorn (=Tiarophorus)	539	* <i>immaturus</i> Schedl, Taphrorychus	559
<i>Hypertentus</i> Hagedorn (=Araptus)	952	<i>immersus</i> Schedl, Xyleborus	743
<i>Hypphaene</i> Hagedorn (=Coccotrypes)	591	<i>immitatrix</i> Schedl, Xyleborus	743
		<i>immitus</i> Bright, Pityoborus (=secundus)	969

<i>immune</i> Eggers, <i>Rhopalopseleon</i>	97	<i>incomptus</i> (Wood), (<i>Eidophelus</i>), <i>Hemicryphalus</i>	867
<i>immunis</i> Nunberg, <i>Xylocleptes</i> (= <i>brounei</i>)	549	<i>incomptus</i> Wood, <i>Chramesus</i>	266
<i>immunitis</i> (Browne), (<i>Minips</i>), <i>Acanthotomicus</i>	481	<i>incomptus</i> Wood, <i>Hylocurus</i>	425
<i>impar</i> (Schedl), (<i>Hylesinus</i>), <i>Chaetoptelius</i>	103	<i>inconstans</i> Schedl, <i>Xyleborus</i>	744
<i>impar</i> Chapuis, <i>Diamerus</i>	196	<i>inconueniens</i> Schedl, <i>Xyleborus</i> (= <i>sparsipilosus</i>)	774
<i>impar</i> Eggers, <i>Xyleborus</i>	743	<i>incultus</i> (Wood), (<i>Xyleborus</i>), <i>Theoborus</i>	661
<i>impar</i> Schedl, <i>Cyrtogenius</i>	566	<i>incurtus</i> Eggers, (<i>Xyleborus</i>), <i>Leptoxyleborus</i> (= <i>concisus</i>)	659
<i>impar</i> Schedl, <i>Hylocurus</i>	425	<i>indefessus</i> Bright, <i>Pityophthorus</i>	1003
<i>impar nanus</i> Hagedorn, <i>Diamerus</i> (= <i>impar</i>)	196	<i>indicum</i> Wood, <i>Aphanarthrum</i>	618
<i>impensus</i> (Wood), (<i>Thamnophthorus</i>), <i>Araptus</i>	958	<i>indicus</i> Beeson, <i>Scolytogenes</i>	862
<i>imperfectus</i> Eggers, <i>Diamerus</i>	196	<i>indicus</i> Eggers, (<i>Thamnurgides</i>), <i>Coccotrypes</i> (= <i>cyperi</i>)	598
<i>imperialis</i> (Schedl), (<i>Pseudochramesus</i>), <i>Nylechinus</i>	116	<i>indicus</i> Eichhoff, <i>Cryphalus</i> (= <i>dorsalis</i>)	879
<i>imperialis</i> Eichhoff, <i>Hylesinus</i> (= <i>aculeatus</i>)	75	<i>indicus</i> Eichhoff (= <i>Xyleborus</i>), <i>Eucallacea</i> (= <i>piceus</i>)	692
<i>impexus</i> Bright, <i>Pityophthorus</i>	1002	<i>indicus</i> Schedl, <i>Xylocleptes</i>	549
<i>impexus</i> Schedl, <i>Xyleborus</i>	743	<i>indicus</i> Stebbing, <i>Cryphalus</i> (= <i>strohmeyeri</i>)	595
<i>impolitus</i> Wood, <i>Chramesus</i>	266	<i>indicus</i> Stebbing, <i>Dryocoetes</i>	583
<i>imporcatus</i> Wood, <i>Chramesus</i>	266	<i>indicus</i> Wood, <i>Bothinodroctonus</i>	310
<i>impressifrons</i> Hopkins, (<i>Hypothencenus</i>), <i>Trischidias</i> (= <i>atomus</i>)	947	<i>indicus</i> Wood, <i>Hylurgus</i>	119
* <i>impressum</i> Scudder, <i>Trypodendron</i>	635	<i>indicus</i> Wood, <i>Pseudoxylechinus</i>	114
<i>impressus</i> (Schedl), (<i>Gnathoglochinus</i>), <i>Gnathotrupes</i>	1040	<i>indicus</i> Wood, <i>Scolytogenes</i>	862
<i>impressus</i> (Wood), (<i>Loganius</i>), <i>Chemonyx</i>	316	<i>indicus subcoriaceus</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>piceus</i>)	693
<i>impressus</i> Eggers, <i>Coccotrypes</i>	604	<i>indigenus</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700
<i>impressus</i> Eichhoff, <i>Xyleborus</i> (= <i>ferrugineus</i>)	738	<i>indigenus</i> Wood, <i>Hypothencenus</i>	931
<i>impressus</i> Rey, <i>Phloeosinus</i> (= <i>thujae</i>)	258	<i>indigenus</i> Wood, <i>Pityophthorus</i>	1003
<i>impressus</i> Schedl, (<i>Gnathotrichus</i>), <i>Gnathotrupes</i> (= <i>bituberculatus</i>)	1039	<i>indigenus</i> Wood, <i>Pityophthorus</i> (= <i>indigus</i>)	1003
<i>impressus</i> Wood, <i>Scolytodes</i>	393	<i>indigenus</i> Wollaston, (<i>Hylesinus</i>), <i>Chaetoptelius</i> (= <i>vestitus</i>)	104
<i>improbis</i> Sampson, <i>Xyleborus</i>	744	<i>indigus</i> Wood, <i>Pityophthorus</i>	1003
<i>improcerus</i> Sampson, <i>Xyleborus</i>	744	<i>indocorus</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>wallacei</i>)	695
<i>improvidus</i> Schedl, <i>Xyleborus</i>	744	<i>Indocryphalus</i> Eggers	648
<i>inaequalis</i> Schedl, <i>Xyleborus</i>	744	<i>indolatus</i> Wood, <i>Dryocoetoides</i>	658
<i>inaequalis</i> Wood, <i>Hylocurus</i>	425	<i>indonesianus</i> Browne, <i>Xyleborus</i>	744
<i>inaequidens</i> Schedl, <i>Pityophthorus</i>	1003	<i>industrius</i> Sampson, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>caripennis</i>)	652
<i>inaequipunctatus</i> Butovitsch, <i>Scolytus</i> (= <i>pygmaeus</i>)	358	<i>ineditus</i> Bright, <i>Pityophthorus</i>	1003
<i>inaffectatus</i> (Schedl), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658	<i>inermis</i> (Schedl), (<i>Stephanorhopalus</i>), <i>Eruoporus</i>	855
<i>inaffectus</i> Wood, <i>Tricolus</i>	1043	<i>inermis</i> Browne, <i>Cyrtogenius</i>	566
<i>inarmatum</i> Wollaston, <i>Liparthrum</i>	277	<i>inermis</i> Browne, <i>Hypothencenus</i>	931
<i>inarmatus</i> Eggers, <i>Xyleborus</i>	744	<i>inermis</i> Browne, <i>Sphaerotrypes</i>	199
<i>inceptis</i> Wood, <i>Pityophthorus</i> (= <i>exquisitus</i>)	998	<i>inermis</i> Eggers, <i>Diamerus</i>	196
<i>incertus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	674	<i>inermis</i> Eichhoff, <i>Xyleborus</i> (= <i>xylographus</i>)	784
<i>incertus</i> Schedl, <i>Xyleborus</i> (= <i>rugosipennis</i>)	769	<i>inermis</i> Nunberg, <i>Phloeosinus</i> (= <i>henschi</i>)	248
<i>incertus</i> Wood, <i>Micracis</i>	432	<i>inermis</i> Wood, <i>Microcorythylus</i>	1066
<i>incisus</i> Schedl, <i>Tricolus</i>	1043	<i>infans</i> Eichhoff, <i>Pityophthorus</i> (= <i>puberulus</i>)	1022
<i>inclinans</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	481	<i>infans</i> Hagedorn, (<i>Xyleborus</i>), <i>Coccotrypes</i> (= <i>subcribrosus</i>)	612
<i>incognitus</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	604	<i>inferior</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	675
<i>incognitus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothencenus</i>	931	<i>infidelis</i> Wood, <i>Camptocerus</i>	319
<i>incognitus</i> Eggers, <i>Scolytus</i>	331	<i>infimus</i> Schedl, <i>Cryphalus</i>	881
<i>incommodus</i> (Blandford), (<i>Pityophthorus</i>), <i>Araptus</i>	958	<i>infimus</i> Schedl, <i>Pityophthorus</i>	1003
<i>incompositus</i> (Blandford), (<i>Pityophthorus</i>), <i>Araptus</i>	958	<i>infradentatum</i> Wood, <i>Monarthrum</i>	1057
<i>incomptus</i> (Blandford), (<i>Hylastes</i>), <i>Hylurgops</i>	33	<i>infucatus</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>duplicatus</i>)	503

- infulatus* Blackman, *Pityophthorus* 1003
infuscatus Murayama, *Dryocoetes* 583
ingae (Blackman), (*Prionosceles*), *Scolytodes* 393
ingae Wood, *Phloeotribus* 223
ingaeensis (Schedl), (*Metacorthylus*), *Corthylus* 1074
ingatorus Wood, *Scolytodes* 393
ingens (Eichhoff), (*Pterocyclon*), *Monarthrum* 1057
ingens (Schedl), (*Stephanoderes*), *Hypothenemus* 931
ingens Blackman, *Pityophthorus* 1003
ingens Wood, *Chramesus* 266
inhabilis Bright, *Pityophthorus* 1003
inimicus Wood, *Micracis* 432
iniquus Blackman, *Pityophthorus* (= *boycei*) 954
inominatus (Schedl), (*Xyleborus*), *Ambrosiodinus* 675
inoblitus (Schedl), (*Loganius*), *Camptocerus* 319
inoblitus (Schedl), (*Xyleborus*), *Ambrosiodinus* 675
inopinatus (Schedl), (*Dryocoetes*), *Dryocoetiops* 587
inopinatus (Schedl), (*Xyleborus*), *Ambrosiodinus* 675
inops Eichhoff, (*Cryphalus*), *Hypocryphalus*
 (= *mangiferae*) 570
inops Wood, *Pityophthorus* 1003
inornatum Wood, *Pyenarthrum* 385
inornatus Wood, *Thysanoes* 420
inornatus Wood, *Tricolus* 1043
inouyei Nobuchi, *Hylurgops* 34
inquietus Blackman, *Pityophthorus* (= *deletus*) 994
insculptus Wood, *Dryocoetoides* 655
insidiosus (Schedl), (*Loganius*), *Cnemonyx* 316
insignatum Wood, *Monarthrum* 1057
insignis Browne, *Hypothenemus* 931
insignis Browne, *Xyleborus* 744
insignis Eichhoff, (*Xyleborus*), *Taurodemus*
 (= *varius*) 756
insignis Wood, *Cnemonyx* 316
insignis Wood, *Corthylus* 1074
insinuatus Wood, *Araptus* (= *deyrollei*) 955
insitivus Schedl, *Xyleborus* 744
insolitum (Schedl), (*Pterocyclon*), *Monarthrum* 1057
insolitus Bright, (*Xyleborus*), *Xyleborinus*
 (= *andrevesi*) 805
insuetus Bright, *Pityophthorus* 1004
insulare Blair, *Pyenarthrum* 385
insularis (Blackman), (*Cryptocleptes*),
 Pseudothyanoes 414
insularis (Blackman), (*Renocis*), *Chaetophloeus* 273
insularis (Eggers), (*Ips*), *Acanthotomicus* 451
insularis (Eggers), (*Problechilus*), *Gymnochilus* 356
insularis (Nobuchi), (*Cryphalomorphus*),
 Scolytogenes 562
insularis (Schedl), (*Hexacohus*), *Scolytodes* 393
insularis Beeson, *Scolytominus* (= *pusillus*) 840
insularis Eggers, (*Dryocoetes*), *Coccotrypes*
 (= *cyperi*) 599
insularis Eggers, (*Neodryocoetes*), *Araptus*
 (= *hymenaeae*) 957
insularis Eggers, (*Neopityophthorus*), *Araptus*
 (= *laevigatus*) 955
insularis Eggers, *Cnesinus* 209
insularis Eggers, *Coccotrypes* (= *cyperi*) 599
insularis Eggers, (*Thamurgides*), *Coccotrypes*
 (= *nitidus*) 607
insularis Eggers, *Phloeotribus* 223
insularis Eggers, *Sphaerocryptes* 199
insularis Perkins, *Hypothenemus* (= *eruditus*) 921
insularis Schedl, (*Scolytocleptes*), *Scolytominus*
 (= *pusillus*) 839
insularis Schedl, (*Xylochilus*), *Dendroterus*
 (= *sallaci*) 952
insularis Sharp, *Xyleborus* (= *ferrugineus*) 739
insularum Beeson, (*Hylesinus*), *Ficicis* (= *despectus*) 91
insularum Krivolutsкая, *Hypothenemus*
 (= *krivolutskae*) 933
insulindicus Eggers, *Coccotrypes* (= *myristicae*) 606
insulindicus Eggers, *Xyleborus* 744
integer (Eichhoff), (*Tomicus*), *Ips* 509
integer Eichhoff, *Coccotrypes* (= *carpophagus*) 595
integer Schedl, *Xyleborus* 744
intentus Bright, *Pityophthorus* 1004
interjectus (Blandford), (*Xyleborus*), *Eucallacea* 690
interjectus Wood, *Araptus* (= *foreifrons*) 956
intermedium Schedl, *Ctonoxylon* 835
intermedium Schedl, *Monarthrum* 1057
intermedium Schedl, *Rhopalopselion* 97
intermedius (Eggers), (*Xyleborus*), *Coptodryas* 525
intermedius (Sampson), (*Xyloterus*),
 Indocryphalus 645
intermedius Eggers, (*Pocillips*), *Coccotrypes*
 (= *rotundicollis*) 610
intermedius Eggers, *Phloeoborus* 99
intermedius Eggers, *Strombophorus* 203
intermedius Ferrari, *Cryphalus* 551
intermedius Kurenzov, *Scolytus* (= *aratus*) 324
intermedius Schedl, *Cnesinus* 209
intermedius Schedl, *Dendrochilus* 438
intermedius Schedl, *Hylocurus* 426
intermedius Schedl, *Tiarophorus* 539
intermedius Sokanovskii, *Scolytus* (= *rugulosus*) 370
intermedius Wood, *Carphoborus* 305
interponens Schedl, *Hypocryphalus* 569
interponens Schedl, *Xyleborus* 744
interpositus Schedl, *Hylocurus* 426
interpunctatus (Eggers), (*Prionosceles*),
 Scolytodes 393
interpunctatus Blandford, *Xyleborus* 744
interpunctatus Schedl, *Thamurgus* 543
interpunctus Eichhoff, (*Tomicus*), *Ips*
 (= *perturbatus*) 516
interpunctus Hopkins, (*Stephanoderes*),
 Hypothenemus (= *interstitialis*) 931
interruptus (Eggers), *Cladoctonus* 239
interruptus Eggers, (*Xyleborus*), *Eucallacea*
 (= *xanthopus*) 696
interruptus Mannerheim, (*Bostrichus*), *Ips*
 (= *tridens tridens*) 527
interruptus Schedl, *Hylocurus* 426

<i>intersestosus</i> (Blandford), (<i>Xyleborus</i>), <i>Xyleborinus</i>	808	<i>izuensis</i> (Murayama), (<i>Xyleborus</i>), <i>Coptodryas</i>	825
<i>intersestosus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	924	<i>izucensis</i> Nobuchi, <i>Phloeosinus</i> (= <i>pulchellus</i>)	251
<i>interstitialis</i> (Chapuis), (<i>Hylastes</i>), <i>Hylurgops</i>	34	<i>jacobsoni</i> (Eggers), (<i>Dendrugus</i>), <i>Coccotrypes</i>	604
<i>interstitialis</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	931	<i>jacobsoni</i> (Spessivtsev), (<i>Eccoptogaster</i>), <i>Scolytus</i>	334
<i>interstitialis</i> (Lea), (<i>Hylesinus</i>), <i>Diamerus</i>	196	<i>jaintianus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	796
<i>interstitialis</i> Browne, <i>Strombophorus</i>	203	<i>jalapae</i> (Letzner), (<i>Bostrichus</i>), <i>Scolytogenes</i>	862
<i>interstitialis</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>calligraphus</i>)	496	<i>jalapae</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>validum</i>)	1063
<i>interstitialis</i> Eichhoff, <i>Xyleborus</i> (= <i>voluulus</i>)	782	<i>jaliscoensis</i> Wood, <i>Pseudochramcesus</i>	262
<i>intextus</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>andamanensis</i>)	686	<i>jamaicensis</i> Bright, <i>Xyleborus</i> (= <i>scaber</i>)	770
<i>intextus</i> Swaine, <i>Pityophthorus</i>	1004	<i>jambolanaensis</i> Schedl, <i>Xyleborus</i>	745
<i>intonsus</i> Wood, <i>Pityoborus</i> (= <i>secundus</i>)	969	<i>jandiacus</i> Enderlein, <i>Colcobotrus</i> (= <i>alluaudi</i>)	615
<i>intricatus</i> (Ratzeburg), (<i>Eccoptogaster</i>), <i>Scolytus</i>	331	<i>japonicus</i> (Eggers), (<i>Pocilips</i>), <i>Coccotrypes</i>	604
<i>intricatus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	932	<i>japonicus</i> (Niisima), (<i>Cryphalus</i>), <i>Hypothenemus</i>	932
<i>intricatus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>catulus</i>)	663	<i>japonicus</i> Chapuis, <i>Scolytus</i>	335
<i>intrusus</i> Blandford, <i>Xyleborus</i>	745	<i>japonicus</i> Eggers, <i>Crypturgus</i> (= <i>cinereus</i>)	621
<i>intrusus</i> Wood, <i>Tricolus</i>	1043	<i>japonicus</i> Niisima, <i>Ips</i> (= <i>typographus</i>)	535
<i>inurbanus</i> (Broun), (<i>Apate</i>), <i>Mesoscolytus</i>	818	<i>japonicus</i> Nobuchi, <i>Ernoporus</i>	555
<i>invalidus</i> Wood, <i>Microcorthyus</i>	1066	<i>japonicus</i> Nobuchi, <i>Pityogenes</i>	453
<i>injoensis</i> Bright, <i>Pityophthorus</i> (= <i>absonus</i>)	979	<i>japonicus</i> Nobuchi, <i>Xyleborus</i>	746
<i>Ipidae</i> Bedel (= <i>Scolytidae</i>)	28	<i>japonicus</i> Nunberg, <i>Polygraphus</i>	288
<i>Ipidae</i> Bedel (= <i>Ipini</i>)	438	<i>jaroschewskii</i> Schevyrew, <i>Scolytus</i>	335
<i>Ipina</i> Bedel (= <i>Ipini</i>)	438	<i>jasminiae</i> Schedl, <i>Dendrochilus</i>	438
<i>Ipinae</i> Bedel (= <i>Ipini</i>)	438	<i>jasperi</i> Swaine, <i>Pityokteines</i> (= <i>minutus</i>)	463
<i>Ipini</i> Bedel	438	<i>javanus</i> (Eggers), (<i>Hylesinus</i>), <i>Ficicis</i>	91
<i>ipidia</i> Schedl, <i>Xyleborus</i>	745	<i>javanus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	932
<i>Ips</i> DeGeer	484	<i>javanus</i> Chapuis, <i>Scolytus</i> (= <i>multistriatus</i>)	353
<i>ipsiformus</i> Wood, <i>Acanthotomicus</i>	481	<i>javanus</i> Eggers, (<i>Dryocoetes</i>), <i>Dryocoetips</i> (= <i>coffeae</i>)	587
<i>iranicus</i> Eggers, <i>Scolytus</i> (= <i>eichhoffi</i>)	329	<i>javanus</i> Eggers, <i>Scolytoplatypus</i>	403
<i>irazuensis</i> Wood, <i>Scolytodes</i>	393	<i>javanus</i> Eggers, <i>Xyleborus</i>	746
<i>iriani</i> Browne, <i>Arxyleborus</i>	667	<i>javanus</i> Schedl, <i>Cryphalus</i> (= <i>walkeri</i>)	898
<i>irkutensis</i> Eggers, <i>Pityogenes</i>	452	<i>javanus</i> Schedl, <i>Philopodius</i>	599
<i>irrasus</i> Blandford, <i>Xylechinus</i>	116	<i>jeannili</i> Lepesme, <i>Thamniurgus</i> (= <i>senecionis</i>)	545
<i>irregularis</i> Eggers, <i>Phloeoborus</i>	99	<i>jeffreyi</i> Blackman, <i>Pityophthorus</i>	1004
<i>irregularis</i> Eggers, <i>Pityophthorus</i>	1004	<i>jeffreyi</i> Hopkins, <i>Dendroctonus</i>	152
<i>irregularis</i> Eggers, <i>Xyleborus</i>	745	<i>jeholensis</i> Murayama, <i>Cryphalus</i>	882
<i>irresolutus</i> Walker, <i>Hylesinus</i>	80	<i>jemeniae</i> Schedl, <i>Thamniurgus</i>	544
<i>irretitus</i> Browne, <i>Xylocleptes</i>	549	<i>jezoensis</i> Inoye & Nobuchi, <i>Cryphalus</i>	882
<i>irritans</i> Schedl, <i>Pityophthorus</i> (= <i>indigus</i>)	1003	<i>jezoensis</i> Niisima, <i>Polygraphus</i>	288
<i>ishidai</i> Niisima, <i>Xyleborus</i>	745	<i>jiri</i> (Wood), (<i>Olonthogaster</i>), <i>Hyledius</i>	260
<i>islasi</i> Schedl, <i>Pityophthorus</i> (= <i>schwerdtfegeri</i>)	1027	<i>johansenii</i> Swaine, <i>Dendroctonus</i> (= <i>punctatus</i>)	173
<i>islasi</i> Wood, <i>Pityophthorus</i> (= <i>schwerdtfegeri</i>)	1027	<i>jolireti</i> Nunberg, <i>Thamniurgus</i>	544
<i>isolatus</i> Bright, <i>Bothrosternus</i>	215	<i>jongaensis</i> Schedl, <i>Xyleborus</i>	746
<i>Isophthorus</i> Schedl (= <i>Acanthotomicus</i>)	478	<i>joreri</i> (Schedl), (<i>Trypographus</i>), <i>Hylecinopsis</i>	94
<i>italus</i> Eggers, <i>Dryocoetes</i>	583	<i>joveri</i> Schedl, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>quadrispinosus</i>)	654
<i>itatiayacensis</i> Schedl, <i>Xyleborus</i> (= <i>mutabilis</i>)	754	<i>joreri</i> Schedl, <i>Pityophthorus</i>	1004
<i>ituriensis</i> Eggers, <i>Pityophthorus</i> (= <i>djuguensis</i>)	996	<i>jubae</i> Wollaston, <i>Aphanarthrum</i>	618
<i>ituriensis</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>plumeriae</i>)	939	<i>jubae tuberculatus</i> Wollaston, <i>Aphanarthrum</i> (= <i>jubae</i>)	618
<i>ivoriensis</i> Nunberg, <i>Premnobius</i> (= <i>corthyloides</i>)	653	<i>jubatus</i> Sampson, <i>Phloeosinus</i>	249
<i>ivoriensis</i> Nunberg, <i>Serrastus</i> (= <i>similis</i>)	283	<i>jucundus</i> (Chapuis), (<i>Ceratolepis</i>), <i>Cnemomyx</i>	316
		<i>jucundus</i> Blandford, <i>Pityophthorus</i>	1004
		<i>jucundus</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>cylindrotomicus</i>)	793

- juvundus* Wood, *Scolytodes* 394
judeichi Kirsch, (*Bostrichus*), *Ips* (= *duplicatus*) 503
judenkoi (Schedl), (*Xyleborus*), *Coptodryas* 525
juglandis Beeson, *Scolytus* (= *nitidus*) 354
juglandis Blackman, *Hypothenemus* (= *eruditus*) 923
juglandis Blackman, *Pityophthorus* 1005
juglansi Niisima, *Cryphalus* 552
juglansi Tsai & Yin, *Sphaerotrypes* 200
juglansis Krivolutsкая & Kupianskaya, *Sphaerotrypes* 200
Jugoeryphalus Tsai & Li (= *Cryphalus*) 572
jujuya Blackman, *Phloeotribus* 223
juniperi Doebner, (*Dendroctonus*), *Phloeosinus* (= *thujae*) 255
juniperi Swaine, *Phloeosinus* (= *serratus*) 254
jumanicus Sokanovskii, *Hylurgops* 34
jumanicus Sokanovskii, *Ips* (= *sexdentatus*) 526
jumanicus Sokanovskii, *Polygraphus* 289
jurinskii Eggers, *Carphoborus* 305
justus Schedl, *Xyleborus* 746
kabei Murayama, *Lymantria* 556
kadoyamaensis Murayama, *Xyleborus* 746
kagoshimensis (Nobuchi), (*Taeniothytes*), *Cryphalus* 552
kaimochii (Nobuchi), (*Nipponopolygraphus*), *Polygraphus* 289
kaimochii Nobuchi, *Xyleborus* 746
kajangensis Schedl, *Xyleborus* 746
kalambanganus Schedl, *Hypocryphalus* 569
kalimpongensis Maiti & Saha, *Hyorrhynchus* 192
kalopanacis Kurenzov, *Xyleborus* (= *orientalis*) 757
kalshoveni (Schedl), (*Notoxyleborus*), *Xyleborus* 746
kalshoveni Eggers, *Scolytomimus* (= *pusillus*) 539
kalshoveni Schedl, (*Ptilopodius*), *Cosmoderes* (= *sericeus*) 902
kalshoveni Schedl, (*Stephanoderes*), *Hypothenemus* (= *jaranus*) 933
kaltenbachii (Bach), (*Bostrichus*), *Thamnuargus* 544
kanawhae (Hopkins), (*Erioporus*), *Erioporicus* 551
kaniksu Blackman, *Phloeosinus* (= *punctatus*) 252
karamani Reitter, *Cisurgus* (= *maurus*) 630
karamatsu Sawamoto, *Dryocoetes* 554
kashmirensis Schedl, *Scolytus* 336
kasukumbii Schedl, *Polygraphus* 259
kaszabi Schedl, *Strombophorus* 203
katangensis (Eggers), (*Mimips*), *Acanthotomicus* 451
katangensis Eggers, *Xyleborus* 746
katoi Brownei, *Xyleborus* 746
kauaiensis Perkins, *Xyleborus* 746
keeni (Blackman), (*Myeloborus*), *Pityophthorus* 1005
keeni Blackman, *Phloeosinus* 249
keeni Blackman, *Pseudohylesinus* (= *tsugae*) 114
keeni Hopkins, (*Dendroctonus*) (= *rufipennis*) 173
** keilbachi* Schedl, *Carphoborites* 1052
kelantanensis (Browne), (*Ips*), *Acanthotomicus* 451
Kelantaninus Nunberg (= *Cyclorhipidion*) 697
kelantanum (Schedl), (*Xyleborus*), *Cyclorhipidion* 700
kenti Blackman, *Pityophthorus* (= *ornatus*) 1015
kenyae Lepesme, *Thamnuargus* (= *wittei*) 546
kenyae Schedl, *Glostatus* 535
kenyae Schedl, *Pityophthorus* 1005
kenyaensis (Schedl), (*Afromicracis*), *Miocryphalus* 411
kepongi (Schedl), (*Dryocoetes*), *Dryocoetiops* 555
kepongi (Schedl), (*Ips*), *Acanthotomicus* 451
kersianus Browne, *Xyleborus* 747
kesiyae Browne, *Cryphalus* 552
khasianus Murayama, (*Blastophagus*), *Tomicus* (= *brevipilosus*) 123
khayae (Schedl), (*Xyleborus*), *Eucallacea* 691
khinganensis Murayama, *Xyleborus* 747
kikusae (Schedl), (*Mimiphthorus*), *Mimicurus* 949
kikusae occidentalis Schedl, (*Mimiphthorus*), *Mimicurus* (= *kikusae*) 949
kinabaluensis Bright, *Phloeosinus* 249
kinangopensis Lepesme, *Thamnuargus* (= *senecionis*) 546
kirgisicus Pjatnitskii, *Pityophthorus* 1005
kirishimanus (Murayama), (*Xyleborus*), *Coptodryas* 525
kirschi Skalitzky, *Scolytus* 336
kisoensis Niisima, *Polygraphus* 259
Kissophagus Chapuis 67
kiushuensis Murayama, *Phloeosinus* 249
kivuensis (Nunberg), (*Troglogitica*), *Acacis* 193
kivuensis (Schedl), (*Mimips*), *Acanthotomicus* 451
kivuensis Eggers, (*Xyleborus*), *Eucallacea* (= *xanthopus*) 696
kivuensis Nunberg, (*Tiarophorus*), *Cyrtogenius* (= *bicolor*) 563
kivuensis Schedl, *Cryphalus* 552
kivuensis Schedl, *Ctonoxylon* 535
kivuensis Schedl, *Pityophthorus* 1005
kivuensis Schedl, *Polygraphus* 259
kivuensis Schedl, *Scolytoplatypus* 404
klainedoxae Schedl, *Miocryphalus* 411
klapperichi (Schedl), (*Pocillips*), *Coccotrypes* 604
klapperichi (Schedl), (*Xyleborus*), *Cnestus* 502
klapperichi Bright, *Ambrosiodinus* 675
kleinei Eggers, *Pycnarthrum* 355
klimeschi Eggers, *Trypophloeus* 545
knabi (Hopkins), (*Erioporidae*), *Scolytogenes* 562
knausi (Hopkins), (*Xylocleptes*), *Dendrocraulus* 552
knausi Swaine, *Hylurgops* (= *planirostris*) 39
knausi Swaine, *Ips* 510
knechteli Swaine, *Pityogenes* 453
knochei Eggers, *Polygraphus* 259
knoteki Reitter, *Pityophthorus* 1005
knulli (Blackman), (*Micracis*), *Micracisella* 430
koebelei Blackman, *Cactopinus* 436
kochelei Hopkins, *Hypothenemus* (= *eruditus*) 922
kochelei Lea, *Ficicis* (= *despectus*) 91
koenigi Schevyrew, *Scolytus* 337
kojimai Murayama, *Xyleborus* 747
kolbei (Hagedorn), (*Allarthrum*), *Cryphalus* 552

<i>kolbei</i> Schaufuss, <i>Dacryostactus</i>	252	<i>laevigatus</i> (Eggers), (<i>Pityophthorus</i>), <i>Araptus</i>	955
<i>koltzei</i> Reitter, <i>Scolytus</i>	335	<i>laevigatus</i> Chapuis, (<i>Ctenophorus</i>), <i>Scolytodes</i> (= <i>chapuisi</i>)	390
<i>kononovi</i> Kurenzov, <i>Scolytus</i> (= <i>semenovi</i>)	376	<i>laevigatus</i> Ferrari, <i>Scolytodes</i>	394
<i>kororensis</i> (Wood), (<i>Xyleborus</i>), <i>Eucallacea</i>	691	<i>laevigatus</i> Schedl, (<i>Hexacolus</i>), <i>Scolytodes</i> (= <i>laevigatulus</i>)	394
<i>kostini</i> Sokanovskii, <i>Scolytus</i> (= <i>jarochevskii</i>)	336	<i>laevipennis</i> Schedl, <i>Xyleborinus</i> (= <i>mitosomipennis</i>)	809
<i>kozikowskii</i> Michalski, <i>Scolytus</i>	338	<i>laevis</i> (Browne), (<i>Ozodendron</i>), <i>Cyrtogenius</i>	566
<i>kraatzii</i> (Eichhoff), (<i>Hylesinus</i>), <i>Pteleobius</i>	69	<i>laevis</i> (Eggers), (<i>Prionosecles</i>), <i>Scolytodes</i>	394
<i>kraatzii</i> Eichhoff, <i>Xyleborus</i> (= <i>perforans</i>)	761	<i>laevis</i> (Eggers), (<i>Xyleborus</i>), <i>Eucallacea</i>	691
<i>kraatzii philippinensis</i> Eichhoff, <i>Xyleborus</i> (= <i>perforans</i>)	761	<i>laevis</i> (Schedl), (<i>Neopityophthorus</i>), <i>Araptus</i>	955
<i>kraunthiae</i> (Murayama), (<i>Cryphalus</i>), <i>Hypothenemus</i>	933	<i>laevis</i> (Strohmeyer), (<i>Ozopemon</i>), <i>Dryocoetops</i>	585
<i>kraunthiae</i> Niisima, <i>Xyleborus</i>	747	<i>laevis</i> Browne, <i>Cryphalus</i>	883
<i>krausseii</i> Wichmann-Rosenhof, <i>Hypothenemus</i> (= <i>leprieuri</i>)	933	<i>laevis</i> Browne, <i>Hypocryphalus</i>	869
<i>krinaeus</i> Eggers, <i>Phloeosinus</i> (= <i>henschii</i>)	248	<i>laevis</i> Chapuis, <i>Scolytus</i>	339
<i>krivolutskayae</i> Wood, <i>Hypothenemus</i>	933	<i>laevis</i> Eggers, (<i>Hylocurosoma</i>), <i>Scolytodes</i> (= <i>laevicarpus</i>)	394
<i>krivolutskianus</i> Krivolutskaya, <i>Polygraphus</i> (= <i>poligraphus</i>)	295	<i>laevis</i> Eggers, <i>Strombophorus</i>	203
<i>kroaticus</i> Fuchs, <i>Hylastinus</i> (= <i>obscurus</i>)	67	<i>laeviusculus</i> Blandford, <i>Xyleborus</i> (= <i>bidentatus</i>)	715
<i>kuchingensis</i> (Browne), (<i>Xyleborus</i>), <i>Webbia</i>	831	<i>lambertianae</i> Hopkins, <i>Conophthorus</i> (= <i>ponderosae</i>)	966
<i>kucniemannii</i> Reitter, <i>Hypothenemus</i> (= <i>obscurus</i>)	937	<i>lamottei</i> Chapuis, <i>Pycnarthrum</i> (= <i>hispidum</i>)	385
<i>kumamotoensis</i> Murayama, <i>Phloeosinus</i>	249	<i>lamuensis</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	933
<i>kumamotoensis</i> Murayama, <i>Xyleborus</i>	747	<i>lanaiensis</i> Perkins, <i>Xyleborus</i>	747
<i>kumatoensis</i> (Niisima), (<i>Orosiotes</i>), <i>Cyrtogenius</i>	566	<i>landolphiae</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	863
<i>kunala</i> Strohmeyer, <i>Scolytoplattypus</i> (= <i>sionio</i>)	407	<i>Landolphianus</i> Schedl (= <i>Lamurgus</i>)	408
<i>kuniyoshii</i> Nobuchi, (<i>Ips</i>), <i>Orthotomicus</i>	469	<i>langstoni</i> Blackman, <i>Hylocurus</i>	426
<i>kurenzovi</i> Nunberg, <i>Trypophloeus</i>	845	<i>lauguidus</i> Eichhoff, <i>Pityophthorus</i>	1006
<i>kurenzovi</i> Schedl, <i>Trypophloeus</i> (= <i>kurenzovi</i>)	846	<i>lanieri</i> Wood, <i>Ips</i> (= <i>borealis</i>)	494
<i>kurenzovi</i> Stark, <i>Cryphalus</i>	882	<i>lantanae</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	675
<i>kurilensis</i> Krivolutskaya, <i>Cryphalus</i>	882	<i>Lauurgus</i> Eggers	408
<i>kuscheli</i> Schedl, <i>Coccotrypes</i>	604	<i>laponicus</i> Stark, <i>Pityophthorus</i>	1006
<i>kuscheli</i> Schedl, <i>Pachycotes</i>	187	<i>largipennis</i> Toledo Piza Junior, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915
<i>kuscheli</i> Schedl, <i>Pityophthorus</i>	1005	<i>laricis</i> (Fabricius), (<i>Bostrichus</i>), <i>Orthotomicus</i>	469
<i>kushkensis</i> Sokanovskii, <i>Carphoborus</i> (= <i>perrisi</i>)	308	<i>laricis</i> Blackman, <i>Scolytus</i>	340
<i>kyotoensis</i> Nobuchi, <i>Cryphalus</i>	882	<i>laricis</i> Niisima, <i>Cryphalus</i>	883
<i>labiatu</i> Schedl, <i>Scolytus</i> (= <i>thoracicus</i>)	378	<i>lasiocarpus</i> (Swaine), (<i>Orthotomicus</i>), <i>Pityokteines</i>	463
<i>laboulbenei</i> Decaux, <i>Coccotrypes</i> (= <i>dactyliperda</i>)	601	<i>lasius</i> Wood, <i>Chaetophloeus</i>	273
<i>laciniatus</i> Hagedorn, <i>Xyleborus</i>	747	<i>lassus</i> Wood, <i>Microcorthyus</i>	1066
<i>lacordairei</i> (Chapuis), (<i>Bothrosternus</i>), <i>Sternobothrus</i>	216	<i>latecarinatus</i> Schedl, <i>Xyleborus</i>	747
<i>lacunatus</i> Wood, <i>Xyleborus</i>	747	<i>latecavatus</i> Eggers, <i>Xyleborus</i> (= <i>amphicranoides</i>)	711
<i>laetus</i> Niisima, <i>Xyleborus</i>	747	<i>latecompressus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	675
<i>laetus</i> Schedl, <i>Cnesinus</i>	209	<i>latecornis</i> Schedl, <i>Xyleborus</i>	747
<i>laetus</i> Wood, <i>Pityophthorus</i>	1005	<i>latedeclivis</i> Schedl, (<i>Ips</i>), <i>Acanthotomicus</i> (= <i>sumatranus</i>)	484
<i>laetus</i> Wood, <i>Scolytus</i>	339	<i>laterale</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1057
<i>laeve</i> Eggers, <i>Trypodendron</i>	635	<i>lateralis</i> Swaine, <i>Pityophthorus</i> (= <i>concentralis</i>)	989
<i>laevicollis</i> (Eggers), (<i>Problechilus</i>), <i>Gymnochilus</i>	386	<i>latetruncatus</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	683
<i>laevicollis</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	604	<i>laticaudatus</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	683
<i>laevicollis</i> Schedl, <i>Cnesinus</i> (= <i>dividuus</i>)	207	<i>laticeps</i> (Wood), (<i>Xyleborus</i>), <i>Xylosandrus</i>	796
<i>laevicarpus</i> Wood, <i>Scolytodes</i>	394	<i>laticeps</i> Bright, <i>Pityophthorus</i>	1006
<i>laevifrons</i> Michalski, <i>Scolytus</i> (= <i>schevyrevi</i>)	370	<i>laticollis</i> (Browne), (<i>Pocillips</i>), <i>Coccotrypes</i>	604
<i>laevigatulus</i> Wood, <i>Scolytodes</i>	394		
<i>laevigatum</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1057		
<i>laevigatus</i> (Blandford), (<i>Hypothenemus</i>), <i>Cryptocarenum</i>	903		

<i>laticollis</i> Blandford, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>artestriatus</i>)	805	<i>Leiparthrum</i> Wollaston (= <i>Liparthrum</i>)	275
<i>laticollis</i> Blandford, <i>Hylesinus</i>	80	<i>lenis</i> Blackman, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>schedli</i>)	961
<i>laticollis</i> Browne, <i>Cryphalus</i>	883	<i>lenis</i> Wood, <i>Pityophthorus</i>	1007
<i>laticollis</i> Browne, <i>Hypocryphalus</i>	869	<i>lenis</i> Wood, <i>Taurodennus</i> (= <i>sharpi</i>)	786
<i>laticollis</i> Eggers, <i>Polygraphus</i> (= <i>proximus</i>)	296	<i>lenkoranus</i> Eggers, <i>Scolytus</i> (= <i>intricatus</i>)	334
<i>laticollis</i> Swaine, <i>Ips</i> (= <i>pini</i>)	521	<i>lenkoranus</i> Reitter, <i>Taphrorychus</i>	559
<i>laticollis</i> Wollaston, <i>Aphanarthrum</i> (= <i>pygmaeum</i>)	619	<i>leonhardi</i> Eggers, <i>Dryocoetes</i> (= <i>alutii</i>)	572
<i>laticollis</i> Wood, <i>Scolytopsis</i>	320	<i>leonii</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>sulcifrons</i>)	378
<i>latidens</i> (LeConte), (<i>Tomicus</i>), <i>Ips</i>	510	<i>Lepersinus</i> Reitter (= <i>Hylesinus</i>)	74
<i>laticor</i> Eggers, (<i>Pocellips</i>), <i>Coccotrypes</i> (= <i>rotundicollis</i>)	610	<i>Lepicerinus</i> Hinton (= <i>Scolytogenes</i>)	858
<i>laticor</i> Eggers, <i>Ficicis</i> (= <i>despectus</i>)	90	<i>Lepiceroides</i> Schedl (= <i>Hypothenemus</i>)	905
<i>laticor</i> Eggers, <i>Premnobius</i> (= <i>caripennis</i>)	652	<i>Lepicerus</i> Eichhoff (= <i>Scolytogenes</i>)	858
<i>latipennis</i> Schedl, <i>Xyleborus</i>	747	<i>lepidus</i> (Wichmann), (<i>Pityogenes</i>), <i>Acanthotomicus</i>	482
<i>latipilis</i> Schedl, <i>Camptocerus</i>	319	<i>lepidus</i> Bright, <i>Pityophthorus</i>	1007
<i>latisulcatus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	675	<i>lepidus</i> Bright, <i>Xyleborus</i>	748
<i>lativentris</i> Schedl, <i>Xyleborus</i>	747	<i>lepidus</i> Wood, <i>Araptus</i>	958
<i>latum</i> (Schedl), (<i>Cosmocorymus</i>), <i>Monarthrum</i>	1058	<i>lepidus</i> Wood, <i>Cryptocaremus</i>	903
<i>latus</i> Beeson, <i>Carphoborus</i> (= <i>latus</i> Wood)	305	<i>lepidus</i> Wood, <i>Micracis</i>	432
<i>latus</i> Eggers, <i>Cryphalus</i>	883	<i>lepidus</i> Wood, <i>Scolytodes</i>	394
<i>latus</i> Eggers, (<i>Phloeosinus</i>), <i>Hyledius</i> (= <i>nitidicollis</i>)	260	<i>lepineyi</i> Balachowsky, <i>Triotennus</i>	540
<i>latus</i> Eggers, <i>Ozopemon</i>	589	<i>Lepisomus</i> Kirby (= <i>Polygraphus</i>)	283
<i>latus</i> Eggers, <i>Polygraphus</i> (= <i>bicolor</i>)	285	<i>lepocrinus</i> Tsai & Li, <i>Cryphalus</i>	883
<i>latus</i> Eggers, <i>Strombophorus</i>	203	<i>leprieuri</i> (Perris), (<i>Dryocoetes</i>), <i>Hypothenemus</i>	933
<i>latus</i> Eggers, <i>Xyleborus</i>	747	<i>leprosulus</i> (Browne), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	863
<i>latus</i> Eggers, <i>Xyloctonus</i>	840	<i>leprosulus</i> Browne, <i>Cryphalus</i>	883
<i>latus</i> Lepesme, <i>Thammurgus</i> (= <i>lobeliae</i>)	544	<i>leprosulus</i> Schedl, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>undulata</i>)	827
<i>latus</i> Wichmann, (<i>Phloeophthorus</i>), <i>Phlocotribus</i> (= <i>cristatus</i>)	220	<i>leprosulus</i> Schedl, <i>Arixyleborus</i>	667
<i>latus</i> Wood, <i>Carphoborus</i>	305	<i>leprosus</i> Browne, <i>Glostatus</i>	536
<i>laudatus</i> Wood, <i>Araptus</i>	958	<i>Leptoxyleborus</i> Wood	659
<i>laurivorus</i> Schedl, <i>Liparthrum</i>	277	<i>leptus</i> (Bright), (<i>Neodryocoetes</i>), <i>Araptus</i>	958
<i>lautus</i> Eichhoff, <i>Pityophthorus</i>	1006	<i>lesnei</i> (Eggers), (<i>Hapalogenius</i>), <i>Rhopalopselion</i>	97
<i>lautus</i> Wood, <i>Microborus</i>	383	<i>lesnei</i> Hagedorn, <i>Amphicranus</i>	1047
<i>leui</i> (Schedl), (<i>Xylechinus</i>), <i>Phloeosinopsioides</i>	239	<i>Letznerella</i> Reitter (= <i>Scolytogenes</i>)	858
<i>lebronneci</i> Beeson, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915	<i>letzneri</i> Ferrari, <i>Corthylus</i>	1074
<i>lecontei</i> Blackman, <i>Pseudothysanoes</i>	414	<i>leverensis</i> Browne, <i>Xyleborus</i>	748
<i>lecontei</i> Blandford, <i>Cnesinus</i>	209	<i>leveri</i> Browne, <i>Coccotrypes</i>	605
<i>lecontei</i> Bright, <i>Pityophthorus</i>	1006	<i>levis</i> Blackman, (<i>Hexacolus</i>), <i>Scolytodes</i> (= <i>chapuisi</i>)	390
<i>lecontei</i> Chapuis, (<i>Rhopalopleurus</i>), <i>Chramesus</i> (= <i>hicoriae</i>)	266	<i>levis</i> Wood, <i>Phlocotribus</i>	223
<i>lecontei</i> Hopkins, <i>Ambrosiodmus</i>	675	<i>levis</i> Wood, <i>Pityophthorus</i>	1007
<i>lecontei</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiae</i>)	915	<i>lewckianus</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>lewisi</i>)	676
<i>lecontei</i> Schedl, <i>Phlocotribus</i>	223	<i>lewisi</i> (Blandford), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	675
<i>lecontei</i> Swaine, <i>Hylurgops</i> (= <i>porosus</i>)	39	<i>lewisi</i> Blandford, <i>Hyporhynchus</i>	192
<i>lecontei</i> Swaine, <i>Ips</i>	511	<i>lewisi</i> Chapuis, <i>Phloeosinus</i>	249
<i>lecontei</i> Swaine, <i>Pityogenes</i> (= <i>plagiatus</i>)	455	<i>lezjavai</i> Pjatnitsky, <i>Hypothenemus</i> (= <i>eruditus</i>)	924
<i>lectus</i> Wood, <i>Microborus</i>	384	<i>libani</i> Balachowsky, <i>Phloeosinus</i> (= <i>cedri</i>)	244
<i>lederi</i> Reitter, (<i>Cryphalus</i>), <i>Ernoporus</i> (= <i>tiliae</i>)	857	<i>liberiensis</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	933
<i>leechi</i> Wood, <i>Pityophthorus</i>	1007	<i>liberiensis</i> Hopkins, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>leechi</i> Wood, <i>Pseudothysanoes</i>	414	<i>libidus</i> Wood, <i>Scolytodes</i>	394
<i>lefevrei</i> (Browne), (<i>Mimips</i>), <i>Acanthotomicus</i>	481	<i>libra</i> (Eggers), (<i>Xyleborus</i>), <i>Coptodryas</i>	825
<i>lefevrei</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	933		
<i>leiophyllae</i> Blackman, <i>Pityophthorus</i>	1007		

<i>librocedri</i> Swaine, <i>Xyleborinus</i> (= <i>saxesei</i>)	815	<i>liquidambarae</i> Wood, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943
<i>lichtensteini</i> (Ratzeburg), (<i>Bostrichus</i>), <i>Pityophthorus</i>	1007	<i>liquidambaris</i> Hopkins, <i>Dryocoetes</i> (= <i>betulae</i>)	579
<i>lifuanus</i> Fauvel, <i>Hylastes</i>	55	<i>liquidambaris</i> Blackman, <i>Pityophthorus</i>	1009
<i>lignator</i> Blackman, <i>Micracis</i>	432	<i>liratus</i> (Wood), (<i>Loganius</i>), <i>Cnemomyx</i>	316
<i>lignicolus</i> Wood, <i>Micracis</i>	433	<i>Lissoclastus</i> Schauffuss (= <i>Diamerus</i>)	194
<i>ligniperda</i> (Fabricius), (<i>Bostrichus</i>), <i>Hylurgus</i>	119	<i>litoralis</i> (Beeson), (<i>Thamnurgides</i>), <i>Coccotrypes</i>	605
<i>lignivorus</i> Browne, <i>Diamerus</i>	196	<i>litos</i> Bright, <i>Pityophthorus</i>	1009
<i>lignographus</i> Schedl, <i>Xyleborus</i>	748	<i>litseae</i> Browne, (<i>Diamerides</i>), <i>Phloeocranus</i> (= <i>bruchoides</i>)	236
<i>likiangensis</i> Tsai & Huang, <i>Hylurgops</i> (= <i>longipilus</i>)	35	<i>littoralis</i> Perkins, <i>Xyleborus</i>	748
<i>likiangensis</i> Tsai & Yin, <i>Polygraphus</i> (= <i>rudis</i>)	297	<i>lobatum</i> (Ferrari), (<i>Corthylus</i>), <i>Monarthrum</i>	1058
<i>lina</i> Eggers, <i>Dendrosinus</i> (= <i>bourreriae</i>)	237	<i>lobatus</i> Eggers, <i>Sternobothrus</i>	216
<i>limatus</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	691	<i>lobdelli</i> Blackman, <i>Thysanoes</i>	420
<i>limatus</i> Wood, <i>Dendrocraniulus</i>	552	<i>lobeliae</i> Eggers, <i>Thamnurgus</i>	544
<i>limatus</i> Wood, <i>Microborus</i>	384	<i>loebli</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	676
<i>limatus</i> Wood, <i>Pityophthorus</i> (= <i>ornatus</i>)	1018	<i>loebli</i> Schedl, (<i>Poeclips</i>), <i>Dryocoetops</i> (= <i>laevis</i>)	588
<i>limax</i> (Schedl), (<i>Sphenoceros</i>), <i>Araptus</i>	959	<i>loevendali</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>laevis</i>)	340
<i>limbatus</i> (Blandford), (<i>Dryocoetes</i>), <i>Dendrocraniulus</i>	552	<i>Loganius</i> Chapuis (= <i>Cnemomyx</i>)	313
<i>limbatus</i> (Eggers), (<i>Hylocurosoma</i>), <i>Scolytodes</i>	394	<i>lomatiae</i> Schedl, (<i>Pteleobius</i>), <i>Xylechinus</i> (= <i>spathifer</i>)	115
<i>limbatus</i> Eggers, <i>Pseudopityophthorus</i>	973	<i>lonchocarpeae</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopselion</i>	97
<i>limbatus</i> Eggers, <i>Sphaerotrypes</i>	200	<i>longchirtus</i> Nunberg, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	697
<i>limbatus</i> Fabricius, <i>Trypodendron</i> (= <i>domesticum</i>)	635	<i>longicauda</i> (Browne), (<i>Streptocranus</i>), <i>Coptoborus</i>	663
<i>limbellus</i> Wood, <i>Dendrocraniulus</i>	552	<i>longicollis</i> (Blandford), (<i>Loganius</i>), <i>Cnemomyx</i>	316
<i>limbus</i> Sampson, <i>Eccoptopterus</i>	820	<i>longicollis</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	394
<i>liminaris</i> (Harris), (<i>Tomicus</i>), <i>Phloeotribus</i>	223	<i>longicollis</i> (Eggers), (<i>Thamnurgides</i>), <i>Coccotrypes</i>	605
<i>limitaris</i> Wood, <i>Dendrocraniulus</i>	552	<i>longicollis</i> (Gyllenhal), (<i>Bostrichus</i>), <i>Orthotomicus</i>	472
<i>limulum</i> Wood, <i>Monarthrum</i>	1058	<i>longicollis</i> (Schedl), (<i>Gnathotrichus</i>), <i>Gnathotrupes</i>	1040
<i>linus</i> Wood, <i>Dendrocraniulus</i>	552	<i>longicollis</i> (Wollaston), (<i>Xyloterus</i>), <i>Dactylotrypes</i>	554
<i>linderae</i> Hopkins, (<i>Ambrosiodmus</i>), <i>Ambrosiodmus</i> (= <i>obliquus</i>)	677	<i>longicollis</i> Browne, <i>Xyleborus</i>	748
<i>linearicollis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	508	<i>longicollis</i> Eggers, <i>Cnecinus</i>	209
<i>linearis</i> (Schedl), (<i>Thamnophthorus</i>), <i>Araptus</i>	959	<i>longicollis</i> Eggers, <i>Dryocoetes</i> (= <i>autographus</i>)	578
<i>linearis</i> Eggers, (<i>Poeclips</i>), <i>Coccotrypes</i> (= <i>longior</i>)	605	<i>longicollis</i> Peyerimhoff, <i>Triotemus</i>	541
<i>linearis</i> Eggers, <i>Dendrocraniulus</i> (sp.?)	550	<i>longicollis</i> Schedl, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>araujaiae</i>)	953
<i>linearis</i> Erichson, <i>Hylastes</i>	55	<i>longicollis</i> Swaine, <i>Hylastes</i>	56
<i>linearis</i> Schedl, <i>Xyleborus</i> (= <i>dryographus</i>)	732	<i>longideclivis</i> Wood, <i>Xyleborus</i> (= <i>deplanatus</i>)	724
<i>lineatiformis</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>longidens</i> Eggers, <i>Xyleborus</i>	748
<i>lineatopunctatus</i> (Eggers), (<i>Xyleborus</i>), <i>Cyrtogenius</i>	566	<i>longidens</i> Swaine, <i>Ips</i> (= <i>latidens</i>)	511
<i>lineatum</i> (Olivier), (<i>Bostrichus</i>), <i>Trypodendron</i>	636	<i>longifolia</i> (Stebbing), (<i>Cryphalus</i>), <i>Liparthrum</i>	278
<i>lineatus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	934	<i>longifolia</i> (Stebbing), (<i>Tomicus</i>), <i>Ips</i>	511
<i>lineatus</i> Eggers, <i>Phloeotribus</i>	224	<i>longifolia</i> Stebbing, <i>Polygraphus</i>	289
<i>lineatus</i> Eggers, <i>Xyleborus</i>	748	<i>longifoliae</i> Stebbing, (<i>Hylastes</i>) (= <i>Curculionidae</i>)	43
* <i>lineatus</i> Foster, <i>Hylesinus</i>	81	<i>longior</i> (Eggers), (<i>Poeclips</i>), <i>Coccotrypes</i>	605
<i>lineatus</i> Kurenzov, <i>Scolytus</i> (= <i>ratzeburgi</i>)	364	<i>longior</i> Browne, <i>Cryphalus</i>	883
<i>lineellus</i> Endrodi, <i>Trypodendron</i> (= <i>lineatum</i>)	636	<i>longipennis</i> (Blanchard), (<i>Tomicus</i>), <i>Gnathotrupes</i>	1040
<i>lineigera</i> Guillebeau, (<i>Phloeophthorus</i>), <i>Phloeotribus</i> (= <i>pubifrons</i>)	227	<i>longipennis</i> (Blandford), (<i>Hylastes</i>), <i>Hylurgops</i>	34
<i>lineigera</i> Guillebeau, <i>Phloeotribus</i> (= <i>eristatus</i>)	220	<i>longipennis</i> (Browne), (<i>Taenioglyptes</i>), <i>Cryphalus</i>	883
<i>linnaeivorii</i> Schedl, <i>Xylocleptes</i>	549		
<i>Liparthrum</i> Wollaston	274		
<i>lipingensis</i> Tsai & Li, <i>Cryphalus</i>	883		
<i>lipperti</i> Henschel, (<i>Tomicus</i>), <i>Pityogenes</i> (= <i>calcaratus</i>)	444		

<i>longipennis</i> (Eggers), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	863	<i>luctuosus</i> (Eggers), (<i>Xyleborus</i>), <i>Euacallacea</i>	691
<i>longipennis</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	934	<i>luedericaldi</i> Eggers, <i>Pagiocerus</i>	214
<i>longipennis</i> Browne, <i>Cyrtogenius</i>	566	<i>lugubris</i> (Eggers), (<i>Xyleborus</i>), <i>Euacallacea</i>	692
<i>longipennis</i> Eggers, <i>Scolytodes</i>	394	<i>lukengeae</i> Schedl, <i>Strombophorus</i>	203
<i>longipennis</i> Eggers, <i>Xyleborus</i>	748	<i>lunulatus</i> Eggers, <i>Phloeoborus</i>	99
° <i>longipennis</i> Wickham, <i>Xyleborites</i>	1052	<i>luridus</i> (Wollaston), (<i>Aphanarthrum</i>), <i>Coleobothrus</i>	616
<i>longipennis</i> Wood, <i>Hylocurus</i>	426	<i>luridus</i> Blandford, <i>Corthylus</i>	1074
<i>longipes</i> Israelson, <i>Aphanarthrum</i> (= <i>canariense</i>)	617	<i>luridus annulicollis</i> Enderlein, <i>Coleobothrus</i> (= <i>luridus</i>)	616
<i>longipilis</i> Browne, <i>Hypocryphalus</i>	569	<i>luridus flavicollis</i> Enderlein, <i>Coleobothrus</i> (= <i>luridus</i>)	616
<i>longipilis</i> Schedl, <i>Hypothenemus</i>	934	<i>lustratus</i> Wood, <i>Corthylus</i>	1074
<i>longipilis</i> Schedl, <i>Polygraphus</i>	289	<i>luteolus</i> (Schedl), (<i>Plesiophthorus</i>), <i>Dendrocterus</i>	951
<i>longipilis</i> Schedl, <i>Tragostus</i>	410	<i>luteus</i> (Blandford), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	566
<i>longipilosus</i> Schedl, <i>Scolytus</i> (= <i>submarginatus</i>)	377	<i>luteus</i> Hagedorn, <i>Diamerus</i>	196
<i>longipilum</i> Eggers, <i>Ctenoxylon</i>	838	<i>luteus</i> Schedl, <i>Xyleborus</i>	748
<i>longipilus</i> (Reitter), (<i>Hylastes</i>), <i>Hylurgops</i>	34	<i>luzonicus</i> (Schedl), (<i>Dryocoetes</i>), <i>Coccotrypes</i>	605
<i>longipilus</i> Eggers, <i>Phloeotribus</i>	224	<i>luzonicus</i> Eggers, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>morigerus</i>)	799
<i>longipilus</i> Eggers, <i>Xyleborus</i>	748	<i>luzonicus</i> Eggers, <i>Scolytotrypes</i> (= <i>papuanus</i>)	405
<i>longipilus</i> Schedl, <i>Cryphalus</i>	883	<i>Lymantria</i> Lovendal	554
<i>longipilus</i> Schedl, <i>Pityophthorus</i>	1009	<i>macellus</i> Wood, <i>Amphicranus</i>	1047
<i>longipilus</i> Schedl, <i>Thamnurgus</i>	544	<i>macer</i> (Bright), (<i>Neodryocoetes</i>), <i>Araptus</i>	959
<i>longisetosus</i> (Nobuchi), (<i>Taenioglyptes</i>), <i>Cryphalus</i>	883	<i>macer</i> Blandford, <i>Xyleborus</i>	748
<i>longispinis</i> (Browne), (<i>Xyleborus</i>), <i>Coptoborus</i>	663	<i>macer</i> LeConte, <i>Hylastes</i>	56
<i>longius</i> (Eggers), (<i>Xyleborus longior</i>), <i>Cyclorhipidion</i>	700	<i>macgregori</i> Blackman, <i>Scolytotrypes</i> (= <i>papuanus</i>)	405
<i>longiusculus</i> (Schedl), (<i>Gnathotrichus</i>), <i>Gnathotrichus</i>	1041	<i>machili</i> Niisima, <i>Xyleborus</i>	749
<i>longula</i> Nunberg, <i>Micracis</i>	433	<i>machili</i> Wood, <i>Indocryphalus</i>	648
<i>longulum</i> Eichhoff, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>mali</i>)	1059	<i>machilus</i> (Schedl), (<i>Hylesinus</i>), <i>Phloeosinus</i>	249
<i>longulus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	808	<i>machnorskii</i> (Sokanovskii), (<i>Hypothenemus</i>), <i>Taphrotychus</i>	559
<i>longulus</i> Kolenati, <i>Hylurgus</i> (= <i>ligniperda</i>)	121	<i>macilentus</i> (Blandford), (<i>Dryocoetes</i>), <i>Dendrocranius</i>	552
<i>longulus</i> Sokanovskii, <i>Pityophthorus</i> (= <i>micrographus</i>)	1012	<i>maclayi</i> (Bruck), (<i>Pseudocryphalus</i>), <i>Chaetophloeus</i>	273
<i>longus</i> (Eggers), (<i>Enoporus</i>), <i>Enoporiens</i>	851	<i>macmahoni</i> (Stebbing), (<i>Sphaerotrypes</i>), <i>Hylesinus</i>	81
<i>longus</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	808	<i>macrocerus</i> Eichhoff, <i>Corthylus</i>	1074
<i>longus</i> (Schedl), (<i>Anthonocerus</i>), <i>Monarthrum</i>	1058	<i>macrocornis</i> Wood, <i>Chramesus</i>	267
<i>longus</i> Eggers, <i>Premnobius</i>	653	<i>Macrocryphalus</i> Nobuchi (= <i>Hypothenemus</i>)	905
<i>longus</i> LeConte, <i>Hylastes</i> (= <i>gracilis</i>)	54	<i>macrographus</i> Eichhoff, <i>Pityophthorus</i> (= <i>exsculptus</i>)	998
<i>longus</i> Nunberg, <i>Micryphalus</i>	411	<i>macrolobii</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	934
<i>lopehuensis</i> Beeson, <i>Euacallacea</i> (= <i>interjectus</i>)	690	<i>macropterus</i> Schedl, <i>Xyleborus</i>	749
<i>loranthus</i> Schedl, <i>Margadillius</i>	899	<i>maculatum</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopschion</i>	97
<i>loricatus</i> (Schedl), (<i>Xyleborus</i>), <i>Euacallacea</i>	690	<i>maculatus</i> Beeson, <i>Scolytomimus</i>	839
<i>loiceanum</i> Wollaston, <i>Liparthrum</i>	278	<i>maculatus</i> Browne, <i>Cnestus</i>	802
<i>loicei</i> Paiva, <i>Hylastes</i>	56	<i>maculatus</i> Browne, <i>Hypocryphalus</i> (= <i>sandakanensis</i>)	572
<i>loicei</i> Wollaston, <i>Liparthrum</i>	278	<i>maculatus</i> Nunberg, <i>Scolytotrypes</i> (= <i>fasciatus</i>)	403
<i>lubarskii</i> Stark, <i>Hylesinus</i>	51	<i>maculatus</i> Schedl, <i>Xyleclinus</i>	116
<i>lubricus</i> Schedl, <i>Xyleborus</i>	748	<i>maculatus</i> Schedl, <i>Xylocotinus</i>	840
<i>lucaris</i> Wood, <i>Cnestus</i>	209	<i>maculicollis</i> Sharp, <i>Hypothenemus</i> (= <i>birmanus</i>)	910
<i>lucasi</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942		
<i>lucidum</i> Wood, <i>Pycnarthrum</i>	385		
<i>lucidum</i> Wood, <i>Bothrosternus</i>	215		
<i>luctuosum</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1058		

<i>maculicornis</i> (Blandford), (<i>Ceratolepis</i>), <i>Cnemidophorus</i>	316	<i>major</i> Stebbing, <i>Polygraphus</i>	289
<i>maculipennis</i> (Schedl), (<i>Hylesinus</i>), (<i>Ficicis</i>)	91	<i>major</i> Stebbing, <i>Scolytus</i>	340
<i>maculosus</i> Blackman, <i>Pseudohylesinus</i>	110	<i>major</i> Strohmeier, <i>Cyrtogenius</i>	566
<i>maculosus</i> Kirby, (<i>Ips</i>), (<i>Pteleobius</i> (= <i>vittatus</i>))	70	<i>major</i> Strohmeier, <i>Ozopemon</i>	589
<i>madagascariensis</i> (Schedl), (<i>Euptilius</i>), <i>Cosmoderes</i>	901	<i>majus</i> (Eggers), (<i>Trypodendron</i>), (<i>Indocryphalus</i>)	648
<i>madagascariensis</i> (Schedl), (<i>Micracis</i>), <i>Pseudomicracis</i>	429	<i>majus</i> Wood, <i>Scolytodes</i>	394
<i>madagascariensis</i> Schaufuss, <i>Xyleborus</i>	749	<i>majusculus</i> Schedl, <i>Polygraphus</i>	290
<i>madagascariensis</i> Schedl, (<i>Eriocryphalus</i>), <i>Hypothenemus</i> (= <i>douisi</i>)	917	<i>majusculus</i> Schedl, <i>Xyleborus</i>	749
<i>madagascariensis</i> Schedl, <i>Cyrtogenius</i>	566	<i>malaccensis</i> (Schedl), (<i>Dryocoetes</i>), (<i>Dryocoetops</i>)	588
<i>madagascariensis</i> Schedl, <i>Hypothenemus</i>	934	<i>malaisi</i> Eggers, (<i>Indocryphalus</i> (= <i>intermedius</i>))	648
<i>madagascariensis</i> Schedl, <i>Pityodendron</i>	970	<i>malayanus</i> Browne, <i>Acacacis</i>	193
<i>madagascariensis</i> Schedl, <i>Pityophthorus</i>	1009	<i>malayensis</i> (Schedl), (<i>Dryocoetes</i>), (<i>Cyrtogenius</i>)	566
<i>madagascariensis</i> Schedl, <i>Ptilopodius</i>	900	<i>malayensis</i> (Schedl), (<i>Lepiceriodes</i>), <i>Hypothenemus</i>	934
<i>madagascariensis</i> Schedl, <i>Scolytoplatypus</i> (= <i>peruivus</i>)	406	<i>malayensis</i> (Schedl), (<i>Xyleboricus</i>), (<i>Arxyleborus</i>)	667
<i>magna</i> Blackman, <i>Plurixosoma</i>	189	<i>malayensis</i> Browne, <i>Xyleborus</i>	749
<i>magnificus</i> Wood, <i>Xyleborus</i>	749	<i>malayensis</i> Schedl, <i>Bothrosteronoides</i>	194
<i>magnispinatus</i> Beaver, <i>Xyleborus</i>	749	<i>malayensis</i> Schedl, (<i>Phloeosinus</i>), (<i>Hyledius</i>) (= <i>cribratus</i>)	260
<i>magnus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	934	<i>malayensis</i> Schedl, <i>Hypocryphalus</i>	869
<i>magnus</i> Beeson, <i>Coccotrypes</i>	605	<i>malgasicus</i> (Schedl), (<i>Pocilips</i>), (<i>Coccotrypes</i>)	605
<i>magnus</i> Browne, <i>Arxyleborus</i>	667	<i>malgasicus</i> Schedl, <i>Xyleborus</i>	749
<i>magnus</i> Browne, <i>Margadillius</i>	899	<i>malgasicus</i> Schedl, (<i>Dohrgeocleptes</i>)	540
<i>magnus</i> Murayama, <i>Polygraphus</i> (= <i>proximus</i>)	296	<i>mali</i> (Beckstein), (<i>Bostrichus</i>), (<i>Scolytus</i>)	341
<i>magnus</i> Niisima, <i>Xyleborus</i>	749	<i>mali</i> (Fitch), (<i>Tomicus</i>), (<i>Monarthrum</i>)	1058
<i>magnus</i> Sampson, <i>Cnestus</i>	802	<i>mali</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922
<i>magnus</i> Tsai & Yin, <i>Sphaerotrypes</i>	200	<i>malleatus</i> Bright, <i>Pityophthorus</i>	1009
<i>magnus</i> Wood, <i>Pseudohylesinus</i>	110	<i>malloti</i> (Eggers), (<i>Xyleborus</i>), (<i>Eucallacea</i>)	692
<i>mahafali</i> (Schedl), (<i>Xyleborus</i>), (<i>Ambrosiodinus</i>)	676	<i>malloti</i> Schedl, <i>Cryphalus</i>	884
<i>maiche</i> (Stark), (<i>Anisandrus</i>), (<i>Xyleborus</i>)	749	<i>malayi</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	934
<i>maiche</i> Eggers, <i>Xyleborus</i> (= <i>maiche</i> Stark)	749	<i>malus</i> (Schedl), (<i>Stephanoderes</i>), (<i>Hypothenemus</i>)	934
<i>maillieri</i> Eggers, <i>Scolytus</i> (= <i>amygdali</i>)	323	<i>malus</i> Niisima, <i>Cryphalus</i>	884
<i>mainensis</i> Blackman, <i>Cryphalus</i> (= <i>ruficollis</i>)	892	<i>mamilliae</i> Wood, (<i>Phloeographus</i>), (<i>Halystus</i>) (= <i>namibiae</i>)	283
<i>mairei</i> Peyerimhoff, <i>Aphanarthrum</i>	619	<i>mamibillae</i> (Browne), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	676
<i>mairei</i> Peyerimhoff, <i>Thamnuergus</i>	544	<i>mamillatus</i> Chapuis, <i>Phloeoborus</i>	99
<i>major</i> (Eggers), (<i>Bothryperus</i>), (<i>Plurixosoma</i>)	189	<i>manicus</i> (Blandford), (<i>Xyleborus</i>), (<i>Xylosandrus</i>)	796
<i>major</i> (Eggers), (<i>Hoplites</i>), (<i>Cladoctonus</i>)	239	<i>manicus</i> Wood, <i>Pseudothysanoes</i>	414
<i>major</i> (Eggers), (<i>Loganinus</i>), (<i>Camptocerus</i>)	319	<i>manicus</i> formosanus Eggers, (<i>Xyleborus</i>), (<i>Xylosandrus</i>) (= <i>manicus</i>)	797
<i>major</i> (Eggers), (<i>Prionosceles</i>), (<i>Scolytodes</i>)	394	<i>mandibulare</i> Wollaston, <i>Liparthrum</i>	278
<i>major</i> (Schedl), (<i>Mimips</i>), (<i>Acanthotomicus</i>)	482	<i>mandibularis</i> Blackman, <i>Gnatholeptus</i> (= <i>shannoni</i>)	970
<i>major</i> (Stebbing), (<i>Phloeosinus</i>), (<i>Xyleborus</i>)	749	<i>mandibularis</i> Bright, <i>Chaetophloeus</i>	273
<i>major</i> Browne, <i>Cryptoxyleborus</i>	828	<i>mandibularis</i> Schedl, <i>Pityophthorus</i>	1009
<i>major</i> Browne, <i>Hypothenemus</i>	934	<i>mandibularis</i> Schedl, <i>Thamnuergus</i>	545
<i>major</i> Eggers, (<i>Blastophagus</i>), (<i>Tomicus</i>) (= <i>piniperda</i>)	136	<i>mandibularis</i> Wood, <i>Pseudothysanoes</i>	415
<i>major</i> Eggers, (<i>Nigritus</i>), (<i>Scolytogeus</i> (= <i>darwini</i>))	861	<i>mandli</i> Eggers, (<i>Eccoptogaster</i>), (<i>Scolytus</i>) (= <i>japonicus</i>)	335
<i>major</i> Eggers, <i>Hylurgops</i>	35	<i>mandschuricus</i> Eggers, <i>Cryphalus</i>	884
<i>major</i> Eggers, <i>Scolytodes</i> (= <i>majus</i>)	394	<i>mandschuricus</i> Schedl, <i>Scolytus</i> (= <i>chikisanii</i>)	327
<i>major</i> Schedl, <i>Dendrocraniulus</i>	552	<i>mandshuricus</i> Eggers, <i>Hylesinus</i>	81
<i>major</i> Schedl, <i>Pagiocerus</i>	214	<i>mandarctanus</i> Beeson, <i>Hypothenemus</i> (= <i>dipteroearpi</i>)	916
<i>major</i> Schedl, <i>Strombophorus</i>	203	<i>mangiferae</i> (Stebbing), (<i>Cryphalus</i>), <i>Hypocryphalus</i>	869
<i>major</i> Stebbing, <i>Cryphalus</i>	884		
<i>major</i> Stebbing, <i>Phloeotribus</i>	224		

- mangiferae* Browne, *Diamerus* 196
mangiferae Eggers, *Hypocryphalus*
 (= *mangiferae* Stebbing) 570
manghissiensis Lezhaya, *Scolytus* (= *rugulosus*) 370
mangoense (Schedl), (*Xyleborus*), *Cyclorhpidion* 700
mangororus Schedl, *Hypothenemus* 934
maniensis Browne, *Xyleborus* 750
manni Blackman, *Phloeotribus* (= *pilula*) 227
manni Blackman, *Pseudochranes* 262
mannsfeldi (Wachtl), (*Tomicus*), *Ips* 512
marani Pfeffer, *Carphoborus* 305
marcidum (Schedl), (*Pterocyclon*), *Monarthrum* 1059
marcidus (Schedl), (*Xyleborus*), *Xyleborinus* 808
margadilaonis Hopkins, *Margadillius* 899
Margadillius Hopkins 898
marginatulus Schedl, (*Xyleborus*), *Premnobius*
 (= *marginatus*) 653
marginatus (Browne), (*Poecilips*), *Coccotrypes* 606
marginatus (Hagedorn), (*Xestips*), *Xyloceptes* 549
marginatus Chapuis, *Scolytus* 344
marginatus Duftschmidt, (*Hylesinus*), *Hylurgops*
 (= *palliat*) 38
marginatus Eggers, *Arixyleborus* (= *malayensis*) 667
marginatus Eggers, (*Xyleborus*), *Leptoxyleborus*
 (= *concisus*) 659
marginatus Eggers, *Phloeotribus* 224
marginatus Eggers, *Premnobius* 653
marginatus Wood, *Chramesus* 267
marginatus Wood, *Scolytodes* 394
marginicollis Eggers, *Cnesinus* 209
marginicollis Schedl, *Xyleborus* 750
maritimus Lanier, *Ips* (= *plastographus*) 521
maritimus Reitter, *Hylurgus* (= *ligniperda*) 119
maritimus Stark, *Pityophthorus* (= *pityographus*) 1021
markangensis Tsai & Li, *Cryphalus* 884
marmoratus Blandford, *Xylechinus* 116
marmoratus Strohmeier, *Dactylipalpus*
 (= *cicatricosus*) 101
maroantsetrae Schedl, *Pityophthorus* 1009
maroccanus (Guillebeau), (*Phloeophthorus*),
Phloeotribus 224
maronicus (Eggers), (*Xyleborus*), *Dryocoetoides* 658
marovayji Schedl, *Hypothenemus* (= *seriatus*) 943
marquesanus Beeson, *Ptilopodius* 900
marshalli (Eggers), (*Stephanoderes*),
Hypothenemus 934
marshami Rye, (*Tomicus*), *Dryocoetes* (= *alni*) 572
martiniquensis Eggers, *Hypothenemus* (= *areccae*) 908
marylandicae Hopkins, (*Hypothenemus*), *Trischidias*
 (= *atomus*) 947
mascareniformis Eggers, *Xyleborus* 750
mascarensis Eichhoff, *Xyleborus* (= *affinis*) 708
mascarensis Hagedorn, (*Xyleborus*), *Xylosandrus*
 (= *crassiusculus*) 792
masoni Beeson, (*Thanurgidae*), *Coccotrypes*
 (= *myristicae*) 607
massonianus Tsai & Li, *Cryphalus* 884
matangi Sampson, *Diamerus* 197
materiaris (Fitch), (*Tomicus*), *Guathotrichus* 1035
mateui (Schedl), (*Stephanoderes*), *Hypothenemus* 934
mauiensis Perkins, *Xyleborus* 750
mauiensis Schedl, *Hypothenemus* (= *eruditus*) 925
maulei Roubal, *Crypturgus* (= *pusillus*) 629
maura Balachowsky, *Phlososinus* (= *cedri*) 244
mauretanicus Peyerimhoff, *Pityophthorus* 1009
mauritanus (Schedl), (*Cryphalommorphus*),
Scolytogenes 863
mauritanus Browne, (*Poecilips*), *Coccotrypes*
 (= *nubilus*) 608
mauritanus Menier, *Xyloctonus* 840
maurus (Blandford), (*Dryocoetes*),
Dendrocranulus 552
maurus (Blandford), (*Prionosceles*), *Scolytodes* 394
maurus Eggers, *Cisurgus* 630
maurus Wood, *Phloeotribus* 224
mayeti (Guillebeau), (*Phloeophthorus*),
Phloeotribus 224
mayeti Guillebeau, *Phloeotribus* (= *eristatus*) 220
mayumbensis (Schedl), (*Dryocoetes*),
Cyrtogenius 566
mayumbensis Eggers, *Hypothenemus* (= *areccae*) 906
meakanensis Nisima, *Polygraphus* 290
meccadanus Reitter, *Taphrorychus* (= *hirtellus*) 559
medialis Wood, *Araptus* 959
medialis Wood, *Pityophthorus* 1009
medialis Wood, *Scolytodes* 394
mediocris (Schedl), (*Xyleborus*), *Xylosandrus* 797
mediosectus (Eggers), (*Xyleboricus*),
Arixyleborus 667
mediterraneus (Eggers), (*Dryocoetes*),
Tiarophorus 540
mediterraneus Eggers, (*Eccoptogaster*), *Scolytus*
 (= *rugulosus*) 369
mediterraneus Eichhoff, *Crypturgus* 624
medius (Eggers), (*Mimips*), *Acanthotomicus* 482
medius (Eggers), (*Poecilips*), *Coccotrypes* 606
medius (Eggers), (*Prionosceles*), *Scolytodes* 395
medius (Eggers), (*Webbia*), *Arixyleborus* 668
medius Eggers, *Chortastus* 311
medius Eggers, *Scolytodes* (= *medialis*) 395
medius Wood, *Hyloecarus* 426
megas Bright, *Pityophthorus* 1009
mekoi Schedl, *Cryphalus* (= *wapleri*) 898
melaenus Eichhoff, *Dryocoetes* 584
melanarius (Schedl), (*Cosmoderes*),
Hypothenemus 935
melanarius Schedl, (*Xyleborus*), *Ambrosiodmus*
 (= *obliquus*) 678
melanranis Beeson, *Xyleborus* (= *lineatus*) 748
melanocephalus (Blandford), (*Hexacohus*),
Scolytodes 395
melanocephalus Fabricius, (*Bostrichus*), *Hylesinus*
 (= *rarius*) 86
melanotus Beeson, *Polygraphus* (= *aterrimus*) 285
melanura (Blandford), (*Pterocyclon*),
Amphicranus 1047

<i>melanurus</i> Reitter, <i>Ips</i> (= robustus)	522	<i>mexicanus</i> Eggers, (<i>Neodryocoetes</i>), <i>Araptus</i>	
<i>melanurus</i> Wood, <i>Pityophthorus</i>	1009	(= politus)	961
<i>melas</i> (Eggers), (<i>Xyleborus</i>), <i>Coptoborus</i>	663	<i>mexicanus</i> Eggers, (<i>Renocis</i>), <i>Chactophloeus</i>	
<i>melasomus</i> (Lea), (<i>Cryphalus</i>), <i>Hypothenemus</i>	935	(= mexicanus Blackman)	273
<i>melli</i> Schedl, (<i>Xyleborus</i>), <i>Hadrodemus</i>		<i>mexicanus</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	
(= amorphus)	515	(= obliquus)	675
<i>melodori</i> (Hopkins), (<i>Stephanorhopalus</i>),		<i>mexicanus</i> Eggers, <i>Microcorthyus</i>	1066
<i>Eruoporus</i>	555	<i>mexicanus</i> Hopkins, <i>Dendroctonus</i>	153
<i>mendicus</i> (Wood), (<i>Cryptocleptus</i>),		<i>mexicanus</i> Lacordaire, <i>Phloeotribus</i> (=sp.?)	218
<i>Pseudothyssanus</i>	415	<i>mexicanus</i> Nunberg, <i>Cucinus</i> (= blackmani)	206
<i>mendicus</i> Wood, <i>Araptus</i>	959	<i>mexicanus</i> Schedl, (<i>Breviophthorus</i>), <i>Araptus</i>	
<i>mendosus</i> Schedl, <i>Xyleborus</i> (= barumbuensis)	714	(= dentifrons)	955
<i>mendosus</i> Wood, <i>Pityophthorus</i>	1009	<i>mexicanus</i> Schedl, (<i>Ctenyophthorus</i>), <i>Araptus</i>	
<i>menoni</i> Browne, <i>Scolytinus</i> (= pusillus)	839	(= tenellus)	962
<i>merionale</i> Endrodi, <i>Trypodendron</i> (= lineatum)	636	<i>mexicanus</i> Schedl, <i>Chramesus</i> (= pumilus)	268
<i>meridensis</i> Wood, <i>Liparthrum</i>	278	<i>mexicanus</i> Schedl, <i>Corthyus</i>	1075
<i>meridensis</i> Wood, <i>Xyleborus</i> (= mutabilis)	754	<i>mexicanus</i> Schedl, <i>Micracis</i> (= swainei)	434
<i>meridianus</i> Blackman, <i>Micracis</i> (= suturalis)	433	<i>mexicanus</i> Wood, <i>Cactopinus</i>	436
<i>meridianus</i> Blackman, <i>Pityogenes</i>	454	<i>mexicanus</i> Wood, <i>Conophthorus</i>	964
<i>meridionale</i> Eggers, <i>Trypodendron</i> (= lineatum)	643	<i>mexicanus</i> Wood, <i>Dendroctonus</i>	552
<i>meridionalis</i> (Nobuchi), (<i>Taenioglyptes</i>),		<i>mexicanus</i> Wood, <i>Dendrosinus</i>	237
<i>Cryphalus</i>	554	<i>mexicanus</i> Wood, <i>Hylastes</i>	57
<i>Meringopalpus</i> Hagedorn (= <i>Gymnochilus</i>)	356	<i>mexicanus</i> Wood, <i>Microborus</i>	384
<i>merinjaki</i> Sampson, <i>Diamerus</i>	197	<i>mexicanus</i> Wood, <i>Pityogenes</i>	454
<i>meris</i> Wood, <i>Cucinus</i>	209	<i>mexicanus</i> Wood, <i>Stegomerus</i>	552
<i>meritus</i> Wood, <i>Xyleborus</i>	750	<i>mexicanus</i> Wood, <i>Thysanocis</i>	420
<i>meruensis</i> Nunberg, <i>Xyloctepes</i> (= marginatus)	549	<i>mexicanus</i> Wood, <i>Xylechinus</i>	116
<i>mesembria</i> Bright, <i>Pityophthorus</i>	1009	<i>micaceus</i> Wood, <i>Araptus</i>	959
<i>mesoleiulus</i> Schedl, <i>Xyleborus</i>	750	<i>micaceus</i> Wood, <i>Hyllocurus</i>	426
<i>mesoleius</i> Schedl, <i>Hypothenemus</i> (= eruditus)	925	<i>micacirrus</i> Wood, <i>Corthyus</i>	1075
<i>Mesoscolytus</i> Broun	515	<i>micans</i> (Eggers), (<i>Stephanoderes</i>), <i>Scolytogenes</i>	563
<i>mesuae</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	797	<i>micans</i> (Kugclann), (<i>Bostrichus</i>), <i>Dendroctonus</i>	154
<i>metacomans</i> (Eggers), (<i>Xyleborus</i>), <i>Hadrodemus</i>	519	<i>micans</i> Bright, <i>Pityophthorus</i>	1010
<i>Metacorthyus</i> Blandford	1064	<i>micans</i> Wood, <i>Amphicranus</i>	1045
<i>metacrucifer</i> Browne, (<i>Xyleborus</i>), <i>Cyclorhipidion</i>		<i>micans</i> Wood, <i>Pseudopityophthorus</i> (= limbatus)	973
(= crucipenne)	699	<i>micarius</i> (Wood), (<i>Xyleborus</i>), <i>Theoborus</i>	661
<i>metacuneolus</i> Eggers, <i>Xyleborus</i>	750	<i>michoacanae</i> Wood, <i>Conophthorus</i>	964
<i>metagermanus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	797	<i>micidus</i> Wood, <i>Scolytodes</i>	395
<i>Metahylastes</i> Eggers (= <i>Cyrtogenius</i>)	562	<i>micklitzi</i> Wachtl, <i>Hylurgus</i>	122
<i>Metahylesinus</i> Eggers (= <i>Hylesinopsis</i>)	92	<i>Micracidae</i> LeConte (= <i>Micracini</i>)	405
<i>metanepotulus</i> (Eggers), (<i>Xyleborus</i>), <i>Eucallacea</i>	692	<i>Micracidendron</i> Schedl (= <i>Mimicoccus</i>)	949
<i>methneri</i> Eggers, <i>Ctonoxylon</i>	535	<i>Micracides</i> LeConte (= <i>Micracini</i>)	405
<i>menseli</i> Reitter, <i>Xyleborus</i>	750	<i>Micracinae</i> LeConte (= <i>Micracini</i>)	405
<i>mexicanum</i> Wood, <i>Liparthrum</i>	275	<i>Micracini</i> LeConte	405
<i>mexicanus</i> (Blackman), (<i>Renocis</i>), <i>Chactophloeus</i>	273	<i>Micraciops</i> Schedl (= <i>Lanurgus</i>)	405
<i>mexicanus</i> (Hopkins), (<i>Tomicus</i>), <i>Ips</i>	512	<i>Micracis</i> LeConte	431
<i>mexicanus</i> (Schedl), (<i>Brachyspartus</i>),		<i>Micracisella</i> Blackman	429
<i>Corthylocurus</i>	1069	<i>Micracisoides</i> Blackman (= <i>Hyllocurus</i>)	422
<i>mexicanus</i> (Wood), (<i>Leperisinus</i>), <i>Hylesinus</i>	81	<i>Microborus</i> Blandford	383
<i>mexicanus</i> Blackman, <i>Pityophthorus</i>	1010	<i>microcornis</i> Wood, <i>Cactopinus</i>	436
<i>mexicanus</i> Blackman, <i>Pseudohylesinus</i>		<i>microcornis</i> Wood, <i>Hyllocurus</i>	426
(= variegatus)	114	<i>Microcorthyus</i> Ferrari	1065
<i>mexicanus</i> Blandford, <i>Dendroctonus</i>	951	<i>micrographus</i> (Linnaeus), (<i>Dermestes</i>),	
<i>mexicanus</i> Bright, <i>Carphoborus</i>	305	<i>Pityophthorus</i>	1010
<i>mexicanus</i> Bright, <i>Sampsonius</i>	656	<i>micrographus</i> (Schedl), (<i>Cryptographus</i>),	
<i>mexicanus</i> Eggers, (<i>Amphicranus</i>), <i>Monarthrum</i>		<i>Webbia</i>	831
(= validum)	1063	<i>micrographus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	509

- micrographus* Schedl, (*Breviophthorus*),
Pityophthorus (= *micrograptinus*) 1012
- micrographus curtulus* Sokanovskii, *Pityophthorus*
 (= *micrographus*) 1010
- micrographus longulus* Sokanovskii, *Pityophthorus*
 (= *micrographus*) 1010
- micrographus sibiricus* Stark, *Pityophthorus*
 (= *micrographus*) 1010
- micrograptinus* Wood, *Pityophthorus* 1012
- Microperus* Wood (= *Coptodryas*) 822
- micropilosus* Wood, *Araptus* 959
- microporosus* Wood, *Chramesus* 267
- mikado* Blandford, *Scolytoplatypus* 404
- mikaniae* Schedl, *Dendrochilus* 438
- mikuniyamensis* (Murayama), (*Pseudopocellips*),
Taphrorychus 560
- miles* (LeConte), (*Cryphalus*), *Hypothenemus* 935
- militaris* Eggers, *Polygraphus* 290
- milletiae* (Schedl), (*Dryocoetes*), *Cyrtogenius* 566
- milletiae* Schedl, *Strombophorus* 204
- mimetica* Wood, *Micracisella* 430
- mimicus* (Schedl), (*Mimips*), *Acanthotomicus* 482
- mimicus* Schedl, *Cryphalus* (= *scabricollis*) 593
- mimicus* Schedl, (*Cryphalus*), *Hypocryphalus*
 (= *mangiferae*) 570
- Mimidendrulus* Schedl (= *Cyrtogenius*) 562
- Mimiocurus* Schedl 949
- Mimiophthorus* Schedl (= *Mimiocurus*) 949
- Mimips* Eggers (= *Acanthotomicus*) 478
- mimosae* (Schedl), (*Xyleborus*), *Xyleborinus* 809
- mimosae* Blackman, *Chramesus* 267
- minus* Schedl, *Xyleborus* (= *multispinatus*) 754
- minusopis* Wood, *Scolytominus* 839
- minax* Schedl, *Cnesinus* 209
- mindanaensis* Eggers, *Xyleborus* 750
- mindoroensis* Schedl, *Cyrtogenius* (= *brevior*) 563
- mindoroensis* Schedl, *Hypocryphalus* 570
- miniatus* Bright, *Pityophthorus* 1012
- minimus* (Fabricius), (*Hylesinus*), *Carphoborus* 305
- minimus* (Schedl), (*Pocellips*), *Coccotrypes* 606
- minimus* Eggers, *Cryphalus* 884
- minimus* Hagedorn, *Chortastus* 311
- minimus* Hagedorn, *Scolytoplatypus* 404
- minimus* Schedl, *Microcorthyus* 1066
- minimus* Schedl, *Xyleborus* 750
- minimus* Schedl, *Xylechinus* 105
- minimus* Stebbing, (*Polygraphus*), *Crypturgus*
 (= *pusillus*) 629
- minimus* Wood, *Chaetophloeus* 273
- minimus* Wood, *Corthyus* 1075
- minitropis* Wood, *Cnesinus* 209
- minor* (Blackman), (*Cryptocleptes*),
Pseudothysanoes 415
- minor* (Blandford), (*Problechilus*), *Gymnochilus* 357
- minor* (Eggers), (*Adiaeretus*), *Hypothenemus* 935
- minor* (Eggers), (*Anchonocerus*), *Amphicranus* 1048
- minor* (Eggers), (*Negritus*), *Scolytogenes* 863
- minor* (Eggers), (*Pelicerus*), *Cyrtogenius* 566
- minor* (Eggers), (*Prionosceles*), *Scolytodes* 395
- minor* (Eggers), (*Xyleboricus*), *Arixyleborus* 668
- minor* (Hartig), (*Dendroctonus*), *Tomicus* 123
- minor* (Schedl), (*Landolphianus*), *Lanurgus* 408
- minor* (Stebbing), (*Phlocosinus*), *Ambrosiodmus* 676
- minor* Browne, *Webbia* (= *biformis*) 530
- minor* Eggers, (*Cryphalomorphus*), *Scolytogenes*
 (= *eggersi*) 861
- minor* Eggers, (*Dendrugus*), *Coccotrypes*
 (= *advena*) 592
- minor* Eggers, *Chramesus* 267
- minor* Eggers, *Craniodicticus* 437
- minor* Eggers, *Diamerus* 197
- minor* Eggers, *Dryocoetes* (= *villosus*) 557
- minor* Eggers, *Prennobiis* 653
- minor* Eggers, *Strombophorus* 204
- minor* Eggers, *Taphrorychus* 560
- minor* Eggers, *Xylechinus* 116
- minor* Lindemann, *Polygraphus* (= *subopacus*) 301
- minor* Schedl, *Acacis* 193
- minor* Schedl, *Cnemonyx* 316
- minor* Schedl, *Hypocryphalus* 870
- minor* Schedl, *Margadillius* 899
- minor* Stebbing, *Polygraphus* (= *pini*) 291
- minor* Stebbing, *Scolytus* (= *major*) 340
- minor* Swaine, *Xyleborus* (= *sayi*) 770
- minor* Wood, *Cnesinus* 209
- minor* Wood, *Hylocurus* (= *elegans*) 424
- minor* Wood, *Pachycotes* 187
- minor* Wood, *Phloeotribus* 225
- minor* Wood, *Phrixosoma* 159
- minor robustus* Schedl, *Cyrtogenius* (= *minor*) 566
- minulus* (Wood), (*Cryptocleptes*),
Pseudothysanoes 415
- minulus* Eggers, *Cnemonyx* (= *barbatus*) 314
- Minulus* Eggers (= *Cnemonyx*) 313
- minulus* Wood, *Chramesus* 267
- minus* Bright, *Pityophthorus* 1012
- minuscule* (Blandford), (*Loganius*), *Cnemonyx* 317
- minuscule* Eggers, (*Xyleborus*), *Coptodryas*
 (= *recidens*) 827
- minuscule* Schedl, *Cnesinus* 209
- minuscule* Schedl, *Polygraphus* (= *occidentalis*) 291
- minutalis* Wood, *Pityophthorus* 1012
- minutissima* Wood, *Trischidias* 947
- minutissimum* (Schedl), (*Pterocyclon*),
Monarthrum 1059
- minutissimus* (Schedl), (*Brachydendrulus*),
Araptus 959
- minutissimus* (Schedl), (*Dryocoetes*),
Peridryocoetes 561
- minutissimus* (Schedl), (*Pocellips*), *Coccotrypes* 606
- minutissimus* (Zimmermann), (*Crypturgus*),
Pseudopityophthorus 973
- minutissimus* Eggers, (*Xyleborus*), *Coptodryas*
 (= *recidens*) 827

<i>minutissimus</i> Schedl, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (=knabi)	863	<i>mjoebergi</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	606
<i>minutissimus</i> Schedl, (<i>Hexacolimus</i>), <i>Scolytodes</i> (= <i>minutissimus</i>)	395	<i>mkulumusius</i> (Eggers), (<i>Stephanoderes</i>), <i>Stephanopodius</i>	853
<i>minutissimus</i> Schedl, (<i>Hexacolus</i>), <i>Scolytodes</i> (= <i>minutus</i>)	395	<i>mkulanusius</i> Hagedorn, <i>Xyleborus</i> (=amanicus)	710
<i>minutissimus</i> Schedl, <i>Corthylus</i>	1075	<i>modestus</i> (Murayama), (<i>Cryphalus</i>), <i>Hypothenemus</i>	935
<i>minutissimus</i> Schedl, <i>Eidophelus</i>	867	<i>modestus</i> Murayama, <i>Hyliurgops</i> (=spessirtseri)	41
<i>minutissimus</i> Schedl, <i>Hypothenemus</i> (=pubescens)	939	<i>modicus</i> Blackman, <i>Pityophthorus</i>	1012
<i>minutissimus</i> Schedl, <i>Lanurgus</i>	409	<i>modicus</i> Wood, <i>Dendroterus</i>	951
<i>minutissimus</i> Schedl, <i>Microcorthylus</i> (=minimus)	1066	<i>modus</i> Wood, <i>Dendrocramulus</i>	553
<i>minutissimus</i> Schedl, <i>Polygraphus</i>	290	<i>moestus</i> (Blandford), (<i>Dryocoetes</i>), <i>Taphrorhynchus</i>	560
<i>minutissimus</i> Schedl, <i>Ptilopodius</i>	900	<i>moestus</i> (Eggers), (<i>Xyleborus</i>), <i>Arixyleborus</i>	668
<i>minutissimus</i> Schedl, <i>Scolytodes</i>	395	<i>molhogani</i> Sampson, <i>Scolytodes</i> (=guyanaensis)	393
<i>minutissimus</i> Schedl, <i>Strombophorus</i>	204	<i>molestulus</i> (Wood), (<i>Xyleborus</i>), <i>Theoborus</i>	661
<i>minutissimus</i> Schedl, <i>Tricolus</i>	1044	<i>molestus</i> Wood, <i>Pityophthorus</i>	1013
<i>minutulus</i> Schedl, <i>Hypothenemus</i> (=eruditus)	925	<i>mollis</i> Blackman, <i>Pityophthorus</i> (=infulatus)	1003
<i>minutus</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1059	<i>mollis</i> Schedl, <i>Cryphalus</i>	885
<i>minutus</i> (Swaine), (<i>Dryocoetes</i>), <i>Pityokteines</i>	463	<i>molokaiensis</i> Perkins, <i>Xyleborus</i>	751
<i>minutus</i> Blackman, <i>Hylastes</i> (=tenuis)	63	<i>moluccanus</i> Browne, <i>Xyleborus</i>	751
<i>minutus</i> Blandford, <i>Eidophelus</i>	867	<i>monacensis</i> Fuchs, <i>Pityogenes</i> (=irkutensis)	453
<i>minutus</i> Blandford, <i>Phloeosinus</i> (=lewisi)	249	<i>monacensis</i> bialowiezensis Karpinski, <i>Pityogenes</i> (= <i>irkutensis</i>)	453
<i>minutus</i> Blandford, <i>Xyleborus</i>	750	<i>monachus</i> (Blandford), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658
<i>minutus</i> Bright, <i>Corthylus</i>	1075	<i>monadis</i> Wood, <i>Micracisella</i>	430
<i>minutus</i> Browne, <i>Hypocryphalus</i>	571	<i>Monarthrum</i> Kirsch	1050
<i>minutus</i> Browne, <i>Sphaerotrypes</i>	200	<i>Monebius</i> Hopkins (=Pycnarthrum)	384
<i>minutus</i> Butovitschi, <i>Scolytus</i> (=pygmaeus)	358	<i>mongolica</i> (Sokanovskii), (<i>Gretschkinia</i>), <i>Pseudothysanoes</i>	415
<i>minutus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942	<i>monilicollis</i> Eichhoff, <i>Cosmoderes</i>	901
<i>minutus</i> Hopkins, <i>Margadillius</i>	899	<i>monoceros</i> (Beeson), (<i>Thamnurgides</i>), <i>Coccotrypes</i>	606
<i>minutus</i> Panzer, (<i>Bostrichus</i>), <i>Hylesinus</i> (=varius)	86	<i>monodi</i> Paulian & Villiers, <i>Aphanarthrum</i>	619
<i>minutus</i> Schedl, (<i>Cryphalus</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	925	<i>monographus</i> (Fabricius), (<i>Bostrichus</i>), <i>Xyleborus</i>	751
<i>minutus</i> Schedl, <i>Coccotrypes</i>	606	<i>monophyllae</i> Blackman, <i>Pityophthorus</i> (=deletus)	994
<i>minutus</i> Schedl, <i>Pityophthorus</i>	1012	<i>monophyllae</i> Hopkins, <i>Conophthorus</i>	965
<i>minutus</i> Swaine, <i>Phloeosinus</i> (=swainei)	256	<i>montana</i> Kurenzov, <i>Scolytus</i> (=jacobsoni)	334
<i>minutus</i> Wood, <i>Scolytodes</i>	395	<i>montanum</i> Eggers, <i>Ctonoxylon</i>	838
<i>Miocryphalus</i> Schedl	410	<i>montanus</i> (Bright), (<i>Neodryocoetes</i>), <i>Araptus</i>	959
<i>mirabilis</i> (Schedl), (<i>Streptocramus</i>), <i>Coptoborus</i>	663	<i>montanus</i> (Eichhoff), (<i>Tomicus</i>), <i>Ips</i>	513
<i>mirabilis</i> Nunberg, <i>Corthylus</i>	1075	<i>montanus</i> (Schedl), (<i>Micracidendron</i>), <i>Mimicurus</i>	949
<i>mirandus</i> Wood, <i>Amphicramus</i>	1048	<i>montanus</i> Blackman, <i>Xylechinus</i>	116
<i>mirandus</i> Wood, <i>Stegomerus</i>	852	<i>montanus</i> Bright, <i>Pseudopityophthorus</i> (= <i>hondurensis</i>)	973
<i>mirus</i> (Wood), (<i>Cryptulocleptes</i>), <i>Pseudothysanoes</i>	415	<i>montanus</i> Eggers, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>vulgaris</i>)	614
<i>misatoensis</i> Nobuchi, <i>Xyleborus</i>	751	<i>montanus</i> Fuchs, <i>Ips</i> (=amitinus)	491
<i>miser</i> Blandford, <i>Polygraphus</i>	290	<i>montanus</i> Niisima, <i>Xyleborus</i>	753
<i>mississippiensis</i> Blackman, (<i>Phthorophloeus</i>), <i>Phloeotribus</i> (=liminaris)	224	<i>montanus</i> Nobuchi, <i>Cryphalus</i>	885
<i>mitosomiformis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809	<i>montanus</i> Schedl, <i>Hypocryphalus</i>	871
<i>mitosomipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809	<i>montanus</i> Schedl, <i>Mimicurus</i> (=monticulus)	949
<i>mitosomus</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>spinus</i>)	817	<i>montanus</i> Schedl, <i>Polygraphus</i>	290
<i>mixtecus</i> Bright, <i>Phloeotribus</i> (=armatus)	219	<i>montanus</i> Schedl, <i>Xyleborus</i> (=tenellus)	778
<i>mixtus</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	683	<i>montanus</i> Wood, <i>Stegomerus</i>	852
<i>miyalopiceus</i> Tsai & Li, <i>Cryphalus</i>	884	<i>montezumae</i> Bright, <i>Pityophthorus</i>	1013
<i>miyazakiensis</i> Murayama, <i>Xyleborus</i>	751	<i>monticolae</i> (Swaine), (<i>Eccoptogaster</i>), <i>Scolytus</i>	345

- monticolae* Bright, *Pityophthorus* (= *carinatus*) . . . 956
monticolae Hopkins, *Conophthorus* (= *ponderosae*) . . . 966
monticolae Hopkins, *Dendroctonus* (= *ponderosae*) . . . 165
monticolus Schedl, (*Xyleborus*), *Coptoborus*
 (= *usagarius*) . . . 664
monticulus Wood, *Mimiocurus* . . . 949
montivagus Bright, *Pityophthorus* . . . 1013
moorei Schedl, *Hypocryphalus* . . . 871
moravitzii Semenov, *Scolytus* . . . 345
moreirai Eggers, *Coccotrypes* (= *dactyliperda*) . . . 601
mori (Aube), (*Hypoborus*), *Liparthrum* . . . 278
morigerus (Blandford), (*Xyleborus*), *Xylosandrus* . . . 797
morigerus (Schedl), (*Stephanoderes*),
Hypothenemus . . . 935
morigerus Wood, *Araptus* . . . 959
morinda Stebbing, *Cryphalus* (= *major*) . . . 884
morio (Eggers), (*Stephanoderes*), *Hypothenemus* . . . 935
morio (Eggers), (*Xyleborus*), *Arxyleborus* . . . 668
morio (Schedl), (*Loganius*), *Camptocerus* . . . 319
moriperda Hopkins, (*Phloeophthorus*), *Phloeotribus*
 (= *frontalis*) . . . 222
moritzi Ferrari, *Brachyspartus* . . . 1068
moritzi Schedl, (*Pterocyclon*), *Monarthrum*
 (= *dimidiatum*) . . . 1054
Morizus Ferrari (= *Corthylus*) . . . 1069
mormon Bright, *Pityophthorus* . . . 1013
morokensis (Eggers), (*Dendrugus*), *Coccotrypes* . . . 606
morosovi Spessivtsev, *Pityophthorus* . . . 1013
morosus Schedl, *Hypothenemus* . . . 935
morosus Schedl, *Xyleborus* (= *clerodendronae*) . . . 719
morosus Wood, *Pityophthorus* . . . 1013
mostatti Hagedorn, (*Xyleborus*), *Xylosandrus*
 (= *compactus*) . . . 789
morsum Wood, *Monarthrum* . . . 1059
morulum Wood, *Corthycyclon* . . . 1068
morulus (Schedl), (*Hexacolus*), *Scolytodes* . . . 395
morulus Blandford, *Xyleborus* . . . 753
moschatae Schaufuss, (*Stephanoderes*),
Hypothenemus (= *obscurus*) . . . 937
moseri Eggers, *Sphaerotrypes* . . . 200
moroliae (Schedl), (*Minidendrus*), *Cyrtogenius* . . . 566
moroliae Schedl, *Strombophorus* . . . 204
mozambiquensis Eggers, *Hypothenemus* . . . 935
mpangae Browne, *Xyleborus* . . . 753
mpossae Schedl, *Pityophthorus* . . . 1013
muasi Browne, *Xyleborus* . . . 753
mucronatoides Schedl, *Xyleborus* . . . 753
mucronatulus Eggers, *Xyleborus* . . . 753
mucronatus (LeConte), (*Cryphalus*),
Procryphalus . . . 848
mucronatus (Wood), (*Cryptoleptus*),
Pseudothysanocs . . . 415
mucronatus Blandford, *Craniodicticus* . . . 437
mucronatus Eggers, *Webbia* . . . 831
mucronatus Eggers, *Xyleborus* . . . 753
mucronatus Wood, *Amphicraus* . . . 1048
mucronifer Wollaston, (*Cryphalus*), *Hypothenemus*
 (= *crudiae*) . . . 914
mueunae (Blackman), (*Neodryocoetes*), *Araptus* . . . 959
mukunye (Schedl), (*Xyleborus*), *Premnobius* . . . 654
mulongensis (Eggers), (*Stephanoderes*),
Hypothenemus . . . 935
multidentatulus (Schedl), (*Stephanoderes*),
Hypothenemus . . . 935
multidentatus (Hopkins), (*Stephanoderes*),
Hypothenemus . . . 936
multidentatus (Murayama), (*Ips*), *Orthotomicus* . . . 473
multidentatus Browne, *Webbia* . . . 831
multidentatus Schedl, (*Stephanoderes*),
Hypothenemus (= *multidentatulus*) . . . 935
multigranosum (Schedl), (*Xyleborus*),
Cyclorhipidion . . . 700
multipunctatus (Schedl), (*Stephanoderes*),
Hypothenemus . . . 936
multipunctatus Browne, *Xyleborus* . . . 753
multipunctatus Browne, *Xyleborus* . . . 753
multiseriatus Schedl, *Pseudochramesus*
 (= *harringtoni*) . . . 262
multisetosus (Murayama), (*Blastophagus*), *Tomicus*
 (= *brevipilosus*) . . . 123
multispinatus Eggers, *Xyleborus* . . . 753
multispinatus Wood, *Pseudothysanocs* . . . 415
multispinosus (Hagedorn), (*Platydictylus*),
Eccoptopterus . . . 820
multispinosus (Schedl), (*Ctonocryphus*),
Glostatus . . . 836
multistriatus (Marshall), (*Ips*), *Scolytus* . . . 345
multistriatus (Wood), (*Hexacolus*), *Scolytodes* . . . 395
multistriatus therondi Hoffmann, *Scolytus*
 (= *multistriatus*) . . . 353
mulungensis (Schedl), (*Dryocoetes*), *Cyrtogenius* . . . 567
mulungensis Schedl, *Pityophthorus* . . . 1013
mumfordi Beeson, *Xyleborus* . . . 754
mundulus (Broun), (*Homarus*), *Chaetoptelius* . . . 103
mundus Blackman, *Pityophthorus* (= *briscoei*) . . . 985
mundus Wood, *Dendroctonus* (= *luteolus*) . . . 951
mundus Wood, *Scolytus* . . . 354
murayamai Browne, *Cnestus*
 (= *murayamai* Schedl) . . . 802
murayamai Schedl, *Cnestus* . . . 802
murex (Blandford), (*Styracopterus*),
Styracoptinus . . . 251
muricatus (Eggers), (*Phloeophthorus*),
Phloeotribus . . . 225
muricus Eichhoff, (*Xyleborus*), *Xyleborinus*
 (= *exiguus*) . . . 807
murilloi (Blackman), (*Cryptoleptus*),
Pseudothysanocs . . . 415
murrayanae Blackman, *Pityophthorus* . . . 1013
murrayanae Hopkins, *Dendroctonus* . . . 157
murudensis Browne, *Xyleborus* . . . 754
mus (Eggers), (*Xyleborus*), *Coptodryas* . . . 825
mus (Schedl), (*Cryphalomorphus*), *Scolytogenes* . . . 863
musangae (Schedl), (*Chortastus*), *Polygraphus* . . . 290
mussooriensis Eggers, *Xyleborus* . . . 754

<i>mustus</i> Schedl. <i>Xyleborus</i>	754	<i>nanus</i> Browne. <i>Hylurdectonus</i>	186
<i>mutabilis</i> Schedl. <i>Scolytotlatypus</i>	404	<i>nanus</i> Eggers. <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>mutabilis</i> Schedl. <i>Xyleborus</i>	754	<i>nanus</i> Hagedorn. <i>Diamerus</i> (= <i>impar</i>)	196
<i>muticus</i> (Schedl). (<i>Stephanoderes</i>), <i>Hypothenemus</i>	936	<i>nanus</i> Hagedorn. <i>Hypothenemus</i> (= <i>crudiae</i>)	914
<i>muticus</i> Blandford. <i>Xyleborus</i>	754	<i>nanus</i> Schedl. <i>Polygraphus</i>	290
<i>muticus</i> Hagedorn. <i>Scolytotlatypus</i> (= <i>dainio</i>)	402	<i>nanus</i> Schedl. <i>Scolytotlatypus</i>	404
<i>muticus</i> Say. <i>Scolytus</i>	354	<i>nanus</i> Wood. <i>Corthylus</i>	1075
<i>mutilatus</i> (Blandford). (<i>Xyleborus</i>). <i>Xylosandrus</i>	799	<i>nanus</i> Wood. <i>Phloeotribus</i>	225
<i>mutilus</i> (Wood). (<i>Paracorthylus</i>). <i>Metacorthylus</i>	1064	<i>nanus</i> Wood. <i>Pityophthorus</i>	1014
<i>myelitis</i> Wood. <i>Cucinus</i>	209	<i>nardus</i> (Schedl). (<i>Hexacolus</i>). <i>Scolytodes</i>	395
<i>Myeloborus</i> Blackman (= <i>Pityophthorus</i>)	976	<i>nasutus</i> Wood. <i>Cactopinus</i>	436
<i>Myelophilites</i> Hagedorn (= <i>Hylurgops</i>)	30	<i>natalensis</i> (Eggers). (<i>Stephanoderes</i>), <i>Micryphalus</i>	411
<i>Myelophilus</i> Eichhoff (= <i>Tomicus</i>)	122	<i>natalensis</i> (Schauffuss). (<i>Xyleborus</i>). <i>Ambrosiodmus</i>	676
<i>myllus</i> Browne. <i>Xyleborus</i>	754	<i>natalensis</i> (Schedl). (<i>Archcephalus</i>). <i>Hypothenemus</i>	936
<i>myristicae</i> (Roepke). (<i>Thamnurgides</i>). <i>Coccotrypes</i>	606	<i>natalensis</i> Eggers. <i>Cladoctonus</i>	239
<i>myristicae</i> (Schedl). (<i>Xyleborus</i>). <i>Coptodryas</i>	825	<i>natalensis</i> Eggers. <i>Polygraphus</i>	290
<i>myristicae</i> Hopkins. <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>natalis</i> Blackman. <i>Pityophthorus</i> (= <i>lautus</i>)	1006
<i>myrmedon</i> Eichhoff. (<i>Stephanoderes</i>). <i>Hypothenemus</i> (= <i>eruditus</i>)	921	<i>naumanni</i> (Schedl). (<i>Gnathominus</i>). <i>Gnathotrupes</i>	1041
<i>mystacinus</i> Wood. <i>Pityokteines</i>	463	<i>narus</i> Blackman. <i>Pityophthorus</i> (= <i>modicus</i>)	1012
<i>naerius</i> Wood. <i>Scolytodes</i>	395	<i>neardus</i> Schedl. <i>Xyleborus</i> (= <i>sayi</i>)	770
<i>naerus</i> Schedl. <i>Cryptoxyleborus</i>	828	<i>nebulosus</i> (LeConte). (<i>Hylesinus</i>). <i>Pseudohylesinus</i>	110
<i>naerus</i> Wood. <i>Triculus</i>	1044	<i>nebulosus</i> Motschulsky. (<i>Hypoborus</i>). <i>Cryphalus</i> (= <i>dorsalis</i>)	879
<i>naganiensis</i> Schedl. <i>Hyledius</i> (= <i>cribratus</i>)	260	<i>nebulosus nebulosus</i> LeConte. <i>Pseudohylesinus</i> (= <i>nebulosus</i>)	111
<i>naganus</i> Beeson. <i>Diamerus</i> (= <i>fici</i>)	195	<i>nebulosus serratus</i> Bruck. <i>Pseudohylesinus</i> (= <i>nebulosus</i>)	111
<i>nagaoensis</i> Murayama. <i>Xyleborus</i>	754	<i>nebulosus</i> Wood. <i>Cucinus</i>	210
<i>nahueliae</i> (Schedl). (<i>Phthorophloeus</i>). <i>Phloeotribus</i>	225	<i>nebulosus</i> Wood. <i>Phloeotribus</i>	225
<i>naidaijinensis</i> (Murayama). (<i>Dryocoetes</i>). <i>Coccotrypes</i>	607	<i>nebulosus</i> Wood. <i>Pityophthorus</i>	1015
<i>nakaneii</i> Nobuchi. <i>Scolytus</i>	354	<i>neocopinus</i> (Bright). (<i>Trypophellos</i>). <i>Liparthrum</i>	279
<i>nakazawai</i> Browne. <i>Xyleborus</i>	754	<i>nectandrae</i> Wood. <i>Gnathotrupes</i>	1041
<i>nameranus</i> Murayama. <i>Xyleborus</i>	755	<i>neganensis</i> Schedl. (<i>Phloeosinus</i>). <i>Hyledius</i> (= <i>cribratus</i>)	241, 260
<i>namibiae</i> (Schedl). (<i>Xyleborus</i>). <i>Xyleborinus</i>	809	<i>neglectum</i> Schedl. <i>Aphanarthrum</i>	619
<i>namibiae</i> Schedl. <i>Halystus</i>	283	<i>neglectus</i> (Schedl). (<i>Xyleborus</i>). <i>Ambrosiodmus</i>	677
<i>namosianus</i> Browne. <i>Hypothenemus</i>	936	<i>neglectus</i> Beeson. (<i>Ericryphalus</i>). <i>Cryphalus</i> (= <i>neglectus</i> Schedl)	885
<i>nandarivatus</i> Schedl. (<i>Xyleborus</i>). <i>Eucallacea</i> (= <i>destruens</i>)	688	<i>neglectus</i> Schedl. <i>Chramesus</i>	267
<i>nanellus</i> Wood. <i>Hypothenemus</i>	936	<i>neglectus</i> Schedl. <i>Cryphalus</i>	885
<i>nanellus</i> Wood. <i>Scolytodes</i>	395	<i>neglectus</i> Schedl. <i>Phloeotribus</i>	225
<i>nankinensis</i> Kurenzov & Kononov. <i>Orthotomicus</i>	473	<i>neglectus</i> Schedl. <i>Scolytotlatypus</i>	404
<i>namula</i> (LeConte). (<i>Micracis</i>). <i>Micracisella</i>	430	<i>Negritus</i> Eggers (= <i>Scolytogenes</i>)	858
<i>nanulus</i> (Schedl). (<i>Gnathotrichus</i>). <i>Gnathotrupes</i>	1041	<i>negrosensis</i> Browne. <i>Cryphalus</i>	885
<i>nanulus</i> (Wood). (<i>Cryphalomorphus</i>). <i>Scolytogenes</i>	863	<i>neirai</i> Eggers. <i>Xyleborus</i>	755
<i>nanulus</i> Schedl. (<i>Stephanoderes</i>). <i>Hypothenemus</i> (= <i>eruditus</i>)	925	<i>Nemobius</i> Chapuis (= <i>Pycnarthrum</i>)	384
<i>nanulus</i> Schedl. <i>Hylesinus</i>	81	<i>Nemopagocerus</i> Schedl (= <i>Enpagocerus</i>)	214
<i>nanulus</i> Wood. <i>Araptus</i>	959	<i>Nemophilus</i> Chapuis (= <i>Cucinus</i>)	205
<i>nanulus</i> Wood. <i>Phloeocleptus</i>	421	<i>nemoralis</i> Wood. <i>Pityophthorus</i>	1015
<i>nanus</i> (Blandford). (<i>Xyleborus</i>). <i>Xylosandrus</i>	799	<i>neoadjunctus</i> (Schedl). (<i>Xyleborus</i>). <i>Coptoborus</i>	663
<i>nanus</i> (Eichhoff). (<i>Gnathotrichus</i>). <i>Gnathotrupes</i>	1041	<i>neocaripennis</i> (Schedl). (<i>Xyleborus</i>). <i>Cyclorhipidion</i>	700
<i>nanus</i> Browne. <i>Cryptoxyleborus</i>	828	<i>neocrucifer</i> (Schedl). (<i>Xyleborus</i>). <i>Cyclorhipidion</i>	700

<i>Neocryphalus</i> Eggers (= <i>Scolytogenes</i>)	555	<i>niger</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943
Neocyphus Numberg	552	<i>niger</i> Marsham, (<i>Hps</i>), <i>Hylastes</i> (= <i>ater</i>)	46
neocylindricus Schedl, <i>Xyleborus</i>	755	<i>niger</i> Numberg, <i>Xyleborus</i> (= <i>squamulatus</i>)	776
<i>Neodiamerus</i> Schedl (= <i>Acacacis</i>)	193	niger Sampson, <i>Xyleborus</i>	755
<i>Neodryocoetes</i> Eggers (= <i>Araptus</i>)	952	<i>niger</i> Schedl, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>advena</i>)	593
neofacialis Schedl, <i>Scolytus</i>	354	niger Schedl, <i>Cryphalus</i>	585
<i>Neoglostatus</i> Schedl (= <i>Pseudothyanoes</i>)	411	niger Schedl, <i>Dactylipalpus</i>	102
<i>neogracilis</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>gracilis</i>)	805	niger Schedl, <i>Pityophthorus</i>	1015
<i>neohybridus</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	697	niger Schedl, <i>Xyloctonus</i>	840
<i>Neohylesinus</i> Eggers (= <i>Phrixosoma</i>)	189	<i>niger</i> Sokanovskii, <i>Pityogenes</i> (= <i>saalasi</i>)	457
<i>Neohyorrhynchus</i> Schedl (= <i>Sucus</i>)	191	niger Stark, <i>Trypophloeus</i>	846
<i>neomexicanus</i> Blackman, <i>Phloeosinus</i> (= <i>scopulorum</i>)	253	<i>niger</i> Stebbing, <i>Polygraphus</i> (= <i>aterrimus</i>)	285
<i>Neomips</i> Schedl (= <i>Pityophthorus</i>)	977	<i>niger</i> Wood, (<i>Loganius</i>), <i>Cnemonyx</i> (= <i>nigrellus</i>)	317
<i>Neophloeotribus</i> Eggers (= <i>Phloeotribus</i>)	215	niger Wood, <i>Cactopinus</i>	436
<i>Neopityophthorus</i> Schedl (= <i>Araptus</i>)	952	niger Wood, <i>Cnesinus</i>	210
Neopteleobius Nobuchi	72	niger Wood, <i>Hylastes</i>	57
neosecabridus Schedl, <i>Xyleborus</i>	755	<i>nigericus</i> Browne, <i>Cryphalus</i>	555
neosphenos (Schedl), (<i>Xyleborus</i>), <i>Coptoborus</i>	663	<i>nigericus</i> Browne, <i>Xyleborus</i> (= <i>erinitus</i>)	721
neotardus Schedl, <i>Corthylus</i>	1075	nigra (Eggers), (<i>Bothryperus</i>), <i>Phrixosoma</i>	190
<i>Neotomicus</i> Fuchs (= <i>Orthotomicus</i>)	467	nigra Wood, <i>Micracisella</i>	430
neotropicalis Wood, <i>Thysanoes</i>	420	nigrella Wood, <i>Micracisella</i>	430
<i>neotropicus</i> Schedl, <i>Phloeosinus</i> (= <i>serratus</i>)	255	nigrellus Wood, <i>Araptus</i>	959
neotruncatus (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	653	nigrellus Wood, <i>Cnemonyx</i>	316
<i>Neoxyloctonus</i> Eggers, <i>Scolytominus</i>	535	nigrescens Browne, <i>Xyleborus</i>	755
<i>nephelii</i> Eggers, (<i>Thamnurgides</i>), <i>Coccotrypes</i> (= <i>rhizophorae</i>)	610	<i>nigrescens</i> Eggers, <i>Diamerus</i> (= <i>ritsemiae</i>)	195
nepocranus (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	677	nigrescens Schedl, <i>Strombophorus</i>	204
<i>nepos</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>funereus</i>)	674	nigreseens Wollaston, <i>Liparthrum</i>	279
<i>nepos robustus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>funereus</i>)	674	nigricans Blandford, <i>Pityophthorus</i>	1015
<i>nepotulomorphus</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>asperatus</i>)	671	<i>nigricans</i> Eggers, <i>Eucallacea</i> (= <i>fornicatus</i>)	655
<i>nepotulus</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>asperatus</i>)	671	<i>nigricans</i> Kurenzov, <i>Polygraphus</i> (= <i>proximus</i>)	296
neptunus Schaufuss, <i>Xyleborus</i>	755	<i>nigricans</i> Schedl, <i>Cnemonyx</i> (= <i>niger</i>)	317
<i>nesianus</i> Beeson, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739	nigricans Schedl, <i>Cryphalus</i>	555
nevermanni (Schedl), (<i>Pterocyclus</i>), <i>Monarthrum</i>	1059	<i>nigricaps</i> Kirby, (<i>Apate</i>), <i>Polygraphus</i> (= <i>rufipennis</i>)	295
nevermanni (Schedl), (<i>Thylurcos</i>), <i>Corthylus</i>	1075	<i>nigricollis</i> Eggers, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>advena</i>)	592
<i>nevermanni</i> Schedl, <i>Eupagiocerus</i> (= <i>ater</i>)	214	<i>nigricollis</i> Hagedorn, (<i>Xyleborus</i>), <i>Dryocoetoides</i> (= <i>granulicauda</i>)	657
nevermanni Schedl, <i>Scolytus</i>	354	<i>nigricollis</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922
<i>nevermanni</i> Schedl, <i>Xyleborus</i> (= <i>costaricensis</i>)	721	nigrielytris Niisima, <i>Polygraphus</i>	290
<i>nibarani</i> Beeson, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	nigrina (Schedl), (<i>Ernoporus</i>), <i>Trischidias</i>	947
nigellatus (Schedl), (<i>Lepicerinus</i>), <i>Scolytogenes</i>	864	nigrinus (Mannerheim), (<i>Hylurgus</i>), <i>Hylastes</i>	57
niger (Bright), (<i>Neodryocoetes</i>), <i>Araptus</i>	959	nigrinus Schedl, <i>Microcryphalus</i>	411
niger (Eggers), (<i>Ceratolepis</i>), <i>Cnemonyx</i>	317	nigripennis (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	677
niger (Fabrieius), (<i>Hylesinus</i>), <i>Camptocerus</i>	319	nigripennis Blandford, <i>Metacorthylus</i>	1064
niger (Schedl), (<i>Erioschidias</i>), <i>Cosmoderes</i>	901	<i>nigripennis</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	923
niger (Schedl), (<i>Hapalogenius</i>), <i>Hylesinopsis</i>	94	<i>nigripennis</i> Mader, <i>Tomicus</i> (= <i>minor</i>)	123
niger (Schedl), (<i>Metacorthylus</i>), <i>Corthylus</i>	1075	nigripes Eggers, <i>Coccotrypes</i>	607
niger Eggers, <i>Coccotrypes</i>	607	<i>nigripes</i> Reitter, <i>Phloeosinus</i> (= <i>bicolor</i>)	242
<i>niger</i> Geoffroy, <i>Scolytus</i> (= <i>scolytus</i>)	375	nigrisetosus Eggers, <i>Diamerus</i>	197
		<i>nigritus</i> Gyllenhal, (<i>Bostrichus</i>), <i>Orthotomicus</i> (= <i>suturalis</i>)	477
		nigrirestis (Schedl), (<i>Ctenocryphus</i>), <i>Glostatus</i>	836
		<i>nigroaffinis</i> Beeson, <i>Xyleborus</i> (= <i>adamsoni</i>)	705
		nigronitens (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	607

<i>nigropiceus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	936	<i>nitidulus</i> Guerin-Meneville, (<i>Piezorhopalus</i>), <i>Amphicranus</i> (= <i>thoracicus</i>)	1049
<i>nigropilosus</i> Eggers, <i>Xyleborus</i>	755	<i>nitidulus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943
<i>nigrosetosus</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacca</i>	692	<i>nitidulus</i> Schedl, <i>Thammurgus</i>	545
<i>nigrosetosus</i> Hagedorn, <i>Xylechinus</i>	116	<i>nitidum</i> Schedl, <i>Rhopalopselion</i>	97
<i>nigrosetosus</i> Schedl, <i>Hypocryphalus</i>	871	<i>nitidus</i> (Eggers), (<i>Dendrugus</i>), <i>Coccotrypes</i>	607
<i>nigrum</i> Sokanovskii, <i>Trypodendron</i> (= <i>lineatum</i>)	643	<i>nitidus</i> (Eggers), (<i>Prionosceles</i>), <i>Scolytodes</i>	396
<i>niijimai</i> Nobuchi, <i>Dryocoetes</i>	584	<i>nitidus</i> (Hagedorn), (<i>Lepicrus</i>), <i>Cyrtogenius</i>	567
<i>niisimai</i> (Eggers), (<i>Hyortlynchus</i>), <i>Sucus</i>	191	<i>nitidus</i> (Schedl), (<i>Dryocoetes</i>), <i>Dryocoetiops</i>	588
<i>niligrinus</i> Eggers, <i>Hylesinus</i>	52	<i>nitidus</i> (Schedl), (<i>Eidophelus</i>), <i>Micryphalus</i>	411
<i>nimifrons</i> Wood, <i>Cnathotrichus</i>	1036	<i>nitidus</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	803
<i>niponensis</i> Inouye & Nobuchi, <i>Cryphalus</i>	855	<i>nitidus</i> Blackman, <i>Cnesinus</i> (= <i>blackmani</i>)	206
<i>niponicum</i> Blandford, <i>Trypodendron</i>	644	<i>nitidus</i> Browne, <i>Thammurgus</i>	545
<i>niponicus</i> Murayama, <i>Hylurgops</i> (= <i>interstitialis</i>)	34	<i>nitidus</i> Eggers, (<i>Adryocoetes</i>), <i>Coptodryas</i> (= <i>pulla</i>)	826
<i>Nipponopolygraphus</i> Nobuchi (= <i>Polygraphus</i>)	284	<i>nitidus</i> Eggers, <i>Chramesus</i> (= <i>opacicollis</i>)	267
<i>nitellus</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	395	<i>nitidus</i> Eggers, <i>Cnesinus</i>	210
<i>nitellus</i> Browne, <i>Pityophthorus</i>	1015	<i>nitidus</i> Eggers, <i>Ips</i>	514
<i>nitellus</i> Browne, <i>Xyleborus</i>	755	<i>nitidus</i> Eggers, <i>Pagiocerus</i> (= <i>frontalis</i>)	214
<i>nitellus</i> Schedl, <i>Thammurgus</i>	545	<i>nitidus</i> Eggers, <i>Pityogenes</i> (= <i>seirindensis</i>)	458
<i>nitens</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	396	<i>nitidus</i> Eggers, <i>Scolytotplatypus</i>	405
<i>nitens</i> (Schedl), (<i>Ozodendron</i>), <i>Peridryocoetes</i>	561	<i>nitidus</i> Schedl, (<i>Ptilopodius</i>), <i>Eidophelus</i> (= <i>imitans</i>)	867
<i>nitens</i> Browne, <i>Cnestus</i> (= <i>aterrimus</i>)	802	<i>nitidus</i> Schedl, <i>Scolytus</i>	354
<i>nitens</i> Browne, <i>Cryphalus</i>	855	<i>nitidus</i> Swaine, <i>Hylastes</i> (= <i>gracilis</i>)	55
<i>nitens</i> Browne, <i>Xyleborus</i>	755	<i>nitidus</i> Swaine, <i>Phlocosinus</i> (= <i>eupressi</i>)	245
<i>nitens</i> Wood, <i>Cnemionyx</i> (= <i>atratus</i>)	314	<i>nitidus</i> Swaine, <i>Pityophthorus</i>	1016
<i>nitidicollis</i> (Eggers), (<i>Plathorophloeus</i>), <i>Phloeotribus</i>	225	<i>nitidus</i> Swaine, <i>Trypophloeus</i> (= <i>striatulus</i>)	847
<i>nitidicollis</i> (Motschulsky), (<i>Olonthogaster</i>), <i>Hyleidus</i>	260	<i>nitidus orientalis</i> Eggers, <i>Cyrtogenius</i> (= <i>nitidus</i>)	567
<i>nitidicollis</i> (Reitter), (<i>Dryocoetes</i>), <i>Tiarophorus</i>	540	<i>nobilis</i> (Eggers), (<i>Pseudoxyleborus</i>), <i>Amasa</i>	683
<i>nitidicollis</i> Chapuis, <i>Phloeoborus</i> (= <i>punctatorugosus</i>)	100	<i>nobilis</i> (Wollaston), (<i>Tomicus</i>), <i>Ips</i>	514
<i>nitidicollis</i> Eggers, <i>Scolytotplatypus</i>	405	<i>nobilis</i> Blandford, <i>Hylesinus</i>	52
<i>nitidicollis</i> Hagedorn, (<i>Xyleborus</i>), <i>Dryocoetoides</i> (= <i>granulicauda</i>)	657	<i>nobilis</i> Swaine, <i>Pseudohylesinus</i>	111
<i>nitidicollis</i> Schedl, <i>Hypocryphalus</i>	871	<i>nobuchii</i> Schedl, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>punctatulus</i>)	865
<i>nitidifrons</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>multidentatus</i>)	936	<i>nobuchii</i> Choo & Woo, <i>Polygraphus</i>	291
<i>nitidifrons</i> Hopkins, <i>Cnathotrichus</i>	1036	<i>nocturnus</i> Schedl, <i>Pityophthorus</i>	1016
<i>nitidiloides</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	803	<i>nodatus</i> Wood, <i>Scolytus</i>	354
<i>nitidior</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	683	<i>nodicornis</i> Wichmann, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>nodulus</i>)	355
<i>nitidipennis</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	1015	<i>nodifer</i> Blandford, <i>Tricolus</i>	1044
<i>nitidipennis</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	960	<i>nodifer</i> Reitter, <i>Scolytus</i> (= <i>multistriatus</i>)	353
<i>nitidipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	803	<i>nodosum</i> Eggers, <i>Ctonoxylon</i>	838
<i>nitidipennis</i> Browne, <i>Cryphalus</i>	855	<i>nodulosus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	677
<i>nitidipennis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942	<i>nodulosus</i> Hagedorn, <i>Premnobius</i>	654
<i>nitidipennis</i> Roubal, <i>Xyleborus</i> (= <i>monographus</i>)	751	<i>nodulus</i> (Wichmann), (<i>Eccoptogaster</i>), <i>Scolytus</i>	355
<i>nitidipennis</i> Schedl, (<i>Pociclipis</i>), <i>Coccotrypes</i> (= <i>longior</i>)	605	<i>nodulus</i> Wood, <i>Hylocurus</i>	426
<i>nitidissimus</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	396	<i>noguerae</i> Atkinson, <i>Cnesinus</i>	210
<i>nitidula</i> Wood, <i>Micracisella</i>	430	<i>uolena</i> Wood, <i>Corthylus</i>	1075
<i>nitidulus</i> (Mannerheim), <i>Pityophthorus</i>	1015	<i>Nomebius</i> Navas (= <i>Pycnarthrum</i>)	384
<i>nitidulus</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	960	<i>nonseptis</i> Schedl, (<i>Phlocosinus</i>), <i>Cladoctonus</i> (= <i>banosus</i>)	238
<i>nitidulus</i> Chapuis, <i>Scolytus</i> (= <i>mali</i>)	344	<i>norfolkensis</i> Schedl, <i>Xyleborus</i>	755
<i>nitidulus</i> Eggers, <i>Xyleborus</i>	755	<i>norimasanus</i> (Murayama), (<i>Dryocoetes</i>), <i>Coccotrypes</i>	607
		<i>normandi</i> (Eggers), (<i>Thammurgus</i>), <i>Tiarophorus</i>	540
		<i>noosi</i> Schedl, <i>Xyleborus</i>	755

<i>notatum</i> Wood, <i>Monarthrum</i>	1059	<i>numidicus</i> Eichhoff, <i>Cryphalus</i>	885
<i>notatus</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	396	<i>numidicus</i> Ferrari, <i>Crypturgus</i>	625
<i>notatus</i> Eggers, (<i>Pocilips</i>), <i>Coccotrypes</i> (= <i>vulgaris</i>)	614	<i>numidicus abbreviatus</i> Eggers, <i>Crypturgus</i> (= <i>numidicus</i>)	625
<i>notatus</i> Eggers, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739	<i>nunbergi</i> Michalski, <i>Scolytus</i>	355
<i>nothofagi</i> (Schedl), (<i>Gnathominus</i>), <i>Gnathotrupes</i>	1042	<i>nuperus</i> Bright, <i>Xyleborus</i>	755
<i>nothofagi</i> Browne, <i>Cryphalus</i>	885	<i>nutans</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700
<i>Notoxyleborus</i> Schedl (= <i>Xyleborus</i>)	704	<i>nutans</i> Schedl, <i>Xyleborus</i> (= <i>rugatus</i>)	769
<i>noumeanus</i> (Browne), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	567	<i>nyalubombeae</i> Schedl, <i>Cryphalus</i>	886
<i>noragrandensis</i> Eggers, <i>Xyleborus</i>	755	<i>nyssae</i> Hopkins, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739
<i>noragineanus</i> Schedl, <i>Xyleborus</i> (= <i>similis</i>)	773	<i>oahuensis</i> Perkins, <i>Xyleborus</i>	756
<i>noraki</i> Reitter, <i>Kissophagus</i>	69	<i>oahuensis</i> Samuelson, <i>Xyleborus</i> (= <i>scabratus</i>)	770
<i>norateutonicus</i> (Schedl), (<i>Gnathocranus</i>), <i>Araptus</i>	960	<i>oahuensis</i> Schedl, <i>Hypothenemus</i> (= <i>arecae</i>)	908
<i>norateutonicus</i> (Schedl), (<i>Phthorophloeus</i>), <i>Phloeotribus</i>	225	<i>oaxacaensis</i> Bright, <i>Monarthrum</i> (= <i>hoegei</i>)	1057
<i>norateutonicus</i> (Schedl), (<i>Ptilopodius</i>), <i>Hypothenemus</i>	936	<i>obanus</i> Hagedorn, <i>Ozopenon</i>	589
<i>norateutonicus</i> Schedl, (<i>Problechilus</i>), <i>Gymnochilus</i> (= <i>concisus</i>)	356	<i>obelus</i> Wood, <i>Scolytus</i>	355
<i>norateutonicus</i> Schedl, <i>Cucinus</i>	210	<i>obesa</i> Blackman, <i>Phrixosoma</i>	190
<i>norateutonicus</i> Schedl, <i>Pityophthorus</i>	1017	<i>obesum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1059
<i>norateutonicus</i> Schedl, <i>Scolytus</i>	355	<i>obesus</i> (Hopkins), (<i>Dacryphalus</i>), <i>Hypocryphalus</i>	871
<i>norateutonicus</i> Schedl, <i>Xyleborus</i> (= <i>posticus</i>)	765	<i>obesus</i> (Schedl), (<i>Micraclops</i>), <i>Lanurgus</i>	409
<i>novellus</i> Blackman, <i>Pityophthorus</i>	1017	<i>obesus</i> (Wood), (<i>Cryptulocleptus</i>), <i>Pseudothysanocs</i>	415
<i>norus</i> (Eggers), (<i>Xyleborus</i>), <i>Leptoxyleborus</i>	660	<i>obesus</i> Browne, <i>Hyporhynchus</i>	192
<i>notus</i> Bright, <i>Xyleborus</i> (= <i>howdenae</i>)	743	<i>obesus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>jaranus</i>)	932
<i>noxius</i> Ratzeburg, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>pygmaeus</i>)	359	<i>obesus</i> Kirsch 1875, <i>Phloeotribus</i> (= <i>pilula</i>)	227
<i>noxius</i> Sampson, (<i>Xyleborus</i>), <i>Euwallacea</i> (= <i>andamanensis</i>)	656	<i>obesus</i> LeConte, <i>Xyleborus</i>	756
<i>nsafukalae</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700	<i>obesus</i> Mannerheim, (<i>Hylurgus</i>), <i>Dendroctonus</i> (= <i>rufipennis</i>)	176
<i>nubilus</i> (Blandford), (<i>Dryocoetes</i>), <i>Coccotrypes</i>	607	<i>obesus</i> Swaine, <i>Pseudohylesinus</i> (= <i>tsugae</i>)	113
<i>nubilus</i> (Wood), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	864	<i>obesus</i> Wood, <i>Araptus</i>	960
<i>nubilus</i> Blackman, (<i>Ceratolepis</i>), <i>Cnemonyx</i> (= <i>ficus</i>)	315	<i>obesus</i> Wood, <i>Scolytodes</i>	396
<i>nubilus</i> Blandford, <i>Phloeotribus</i>	225	<i>obesus</i> minor Swaine, <i>Xyleborus</i> (= <i>sayi</i>)	770
<i>nubilus</i> Samuelson, <i>Xyleborus</i>	755	<i>obiensis</i> Browne, <i>Xyleborus</i>	756
<i>nuciferus</i> Schedl, (<i>Pocilips</i>), <i>Coccotrypes</i> (= <i>advena</i>)	593	<i>oblicatus</i> Schedl, <i>Xyleborus</i> (= <i>obliquus</i>)	757
<i>nudibrevis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	825	<i>obliquecaudata</i> Motschulsky (<i>Phloeotrogus</i>), <i>Ambrosiodmus</i>	677
<i>nudifrons</i> Motschulsky, (<i>Phlocosinus</i>), <i>Hyledius</i> (= <i>concinulus</i>)	259	<i>obliquecaudatum</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>scutellare</i>)	1062
<i>nudipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	825	<i>obliquesectum</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700
<i>nudipennis</i> Schedl, <i>Corthylus</i>	1075	<i>obliquum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1060
<i>nudiusculus</i> Schedl, <i>Corthylus</i> (= <i>flagellifer</i>)	1073	<i>obliquus</i> (LeConte) (<i>Pityophthorus</i>), <i>Ambrosiodmus</i>	677
<i>nudum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1059	<i>obliquus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>interstitialis</i>)	932
<i>nudum</i> Israelson, <i>Aphanarthrum</i> (= <i>glabrum</i>)	618	<i>obliquus</i> Chapuis, <i>Phloeotribus</i> (= <i>pilula</i>)	227
<i>nudus</i> (Nunberg), (<i>Xyleborus</i>), <i>Cnestus</i>	803	<i>obliquus</i> Schedl, <i>Corthylus</i>	1076
<i>nudus</i> (Schedl), (<i>Thamnophtorus</i>), <i>Araptus</i>	960	<i>obliquus</i> Schedl, <i>Cryphalus</i> (= <i>sylvicola</i>)	896
<i>nudus</i> Nunberg, <i>Strombophorus</i> (= <i>crenatus</i>)	203	<i>obliquus</i> Sharp, <i>Xyleborus</i>	756
<i>nudus</i> Schedl, <i>Corthylus</i>	1076	<i>oblitus</i> Perris, (<i>Tonicus</i>), <i>Orthotomicus</i> (= <i>longicollis</i>)	473
<i>nudus</i> Swaine, <i>Pityophthorus</i> (= <i>consimilis</i>)	991	<i>oblongus</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	396
<i>nuesslini</i> Reitter, <i>Kissophagus</i>	69	<i>oblongus</i> (Eggers), (<i>Pseudophloeotribus</i>), <i>Hylesinopsis</i>	95
<i>nugalis</i> Wood, <i>Pityophthorus</i>	1017	<i>oblongus</i> (Nobuchii), (<i>Macrocryphalus</i>), <i>Hypothenemus</i>	936
<i>nugax</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	826	<i>oblongus</i> Blandford, <i>Polygraphus</i>	291
<i>numidicus</i> Brisout, <i>Scolytus</i>	355		

<i>oblongus</i> Eggers, (<i>Poecilips</i>), <i>Coccotrypes</i> (= <i>longior</i>)	605	<i>occidentalis</i> Blackman, <i>Pityophthorus</i>	1017
<i>oblongus</i> Niisima, <i>Cryptobius</i>	886	<i>occidentalis</i> Browne, <i>Scolytolplatypus</i>	405
<i>oblongus</i> Schedl, <i>Xyleborus</i>	757	<i>occidentalis</i> Eggers, <i>Camptocerus</i>	319
<i>obnixus</i> Schedl, (<i>Gnathotrichus</i>), <i>Gnathotripes</i> (= <i>longipennis</i>)	1041	<i>occidentalis</i> Hopkins, <i>Gnathotrichus</i> (= <i>retusus</i>)	1037
<i>obscura</i> Schedl 1962, <i>Pseudodiamerus</i> (= <i>obscurus</i>)	194	<i>occidentalis</i> Peyerimhoff, <i>Cisurgus</i>	630
<i>obscura</i> Schedl 1963, <i>Pseudodiamerus</i> (= <i>obscurus</i>)	194	<i>occidentalis</i> Schedl, (<i>Miniophthorus</i>), <i>Minicoccus</i> (= <i>kikusuae</i>)	949
<i>obscuriceps</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	925	<i>occidentalis</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>areccae</i>)	908
<i>obscurus</i> (Eggers), (<i>Neodryocoetes</i>), <i>Araptus</i>	960	<i>occidentalis</i> Schedl, <i>Polygraphus</i>	291
<i>obscurus</i> (Fabricius), (<i>Hylesinus</i>), <i>Hypothenemus</i>	936	<i>occidentalis</i> Schedl, <i>Strombophorus</i>	204
<i>obscurus</i> (Marshall), (<i>Ips</i>), <i>Hylastinus</i>	64	<i>occidentalis minusculus</i> Schedl, <i>Polygraphus</i> (= <i>occidentalis</i>)	291
<i>obscurus</i> (Wood), (<i>Hexacolus</i>), <i>Scolytodes</i>	396	<i>occlusus</i> Bright, <i>Pityophthorus</i>	1017
<i>obscurus</i> Chapuis, <i>Hylastes</i> (= <i>plumbeus</i>)	60	<i>ocellata</i> Wood, <i>Micracisella</i>	430
<i>obscurus</i> Eggers, <i>Bifonus</i>	1052	<i>ocellatus</i> (Wood), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	678
<i>obscurus</i> Eggers, <i>Microcorthyus</i>	1066	<i>ochromae</i> Wood, <i>Scolytodes</i>	396
<i>obscurus</i> Eggers, <i>Pernophorus</i>	202	<i>octiesdentatus</i> Murayama, <i>Xyleborus</i>	757
<i>obscurus</i> Eggers, <i>Pseudodiamerus</i>	194	<i>octodecimspinus</i> Sampson, <i>Webbia</i> (= <i>diptercarpi</i>)	830
<i>obscurus</i> Eggers, <i>Xylechinus</i>	116	<i>octodentatus</i> (Schedl), (<i>Pterocyclonoides</i>), <i>Tricolus</i>	1044
<i>obscurus</i> Eichhoff, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>setosus</i>)	944	<i>octodentatus</i> Browne, <i>Acanthotomicus</i>	482
<i>obscurus</i> Ferrari, (<i>Cryptobius</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	920	<i>octodentatus</i> Paykull, (<i>Bostrichus</i>), <i>Ips</i> (= <i>typographus</i>)	538
<i>obscurus</i> Hopkins, <i>Hypocryptobius</i>	871	<i>octospinosus</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809
<i>obscurus</i> Rey, <i>Coccotrypes</i> (= <i>dactyliperda</i>)	600	<i>octospinosus</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	482
<i>obscurus</i> Schedl, (<i>Xyleborus</i>), <i>Webbia</i> (= <i>similis</i>)	832	<i>octospinosus</i> Nunberg, <i>Scolytolplatypus</i> (= <i>eichelbaumi</i>)	402
<i>obscurus</i> Wood, <i>Gnathotrichus</i>	1036	<i>ocularis</i> (Schedl), (<i>Cryptalomorphus</i>), <i>Scolytogenes</i>	864
<i>obscurus</i> Wood, <i>Phloeocleptus</i>	421	<i>ocularis</i> (Wood), (<i>Minips</i>), <i>Acanthotomicus</i>	482
<i>obsitum</i> Wollaston, <i>Aphanarthrum</i> (= <i>bicinctum</i>)	617	<i>ocularis</i> Blandford, <i>Cnesinus</i>	210
<i>obsolete</i> (Blandford), (<i>Pityophthorus</i>), <i>Araptus</i>	960	<i>ocularis</i> Wood, <i>Microcorthyus</i>	1066
<i>obstipus</i> Schedl, <i>Xyleborus</i>	757	<i>oculatus</i> Wood, <i>Corthyus</i>	1076
<i>obtectus</i> Schedl, <i>Scolytolplatypus</i>	405	<i>olausi</i> Hagedorn, <i>Xyleborus</i> (= <i>princeps</i>)	765
<i>obtrusus</i> Sampson, <i>Xyleborus</i> (= <i>tonuceus</i>)	779	<i>olmoi</i> Browne, <i>Xyleborus</i>	757
<i>obtusatus</i> Schedl, <i>Xyleborus</i>	757	<i>oltoensis</i> Nobuchi, <i>Xyleborus</i>	757
<i>obtusicollis</i> (Schedl), (<i>Poecilips</i>), <i>Coccotrypes</i>	608	<i>okadai</i> Browne, <i>Araxyleborus</i>	668
<i>obtusicollis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	826	<i>okinosenensis</i> Murayama, <i>Xyleborus</i>	757
<i>obtusicornis</i> Schedl, <i>Sampsonius</i>	656	<i>okounensis</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>crassiusculus</i>)	792
<i>obtusipennis</i> Blandford, <i>Pityophthorus</i>	1017	<i>oleae</i> Fabricius, (<i>Bostrichus</i>), <i>Phloeotribus</i> (= <i>scarabaeoides</i>)	231
<i>obtusipennis</i> Eggers, <i>Xyleborus</i> (= <i>ferrugineus</i>)	736	<i>oleae</i> Schedl, <i>Lanurgus</i>	409
<i>obtusipennis</i> Schedl, <i>Hylacurus</i>	426	<i>oleaeformis</i> Schedl, <i>Lanurgus</i>	409
<i>obtusipennis</i> Schedl, <i>Webbia</i>	831	<i>oleanderi</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	960
<i>obtusispinosus</i> Schedl, <i>Webbia</i>	831	<i>oleiperda</i> Fabricius, (<i>Bostrichus</i>), <i>Hylesinus</i> (= <i>toranio</i>)	83
<i>obtusitruuncatus</i> (Schedl), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658	<i>oleiphilus</i> DelGuercio, <i>Phloeotribus</i> (= <i>scarabaeoides</i>)	232
<i>obtusum</i> (Eggers), (<i>Anchouocerus</i>), <i>Monarthrum</i>	1060	<i>olireirai</i> Schedl, <i>Corthyus</i>	1076
<i>obtusum</i> Eggers, <i>Trypodendron</i> (= <i>signatum</i>)	648	<i>olivierai</i> Schedl, <i>Pityophthorus</i>	1017
<i>obtusum</i> Eggers, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>circumcisum</i>)	698	<i>Olonthogaster</i> Motschulsky (= <i>Hyledius</i>)	259
<i>obtusum</i> Eggers, <i>Diamerus</i>	197	<i>ominosus</i> Schedl, <i>Xyleborus</i> (= <i>amanicus</i>)	710
<i>obtusum</i> Eggers, <i>Pityogenes</i> (= <i>bidentatus</i>)	442	<i>omissum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1060
<i>obtusum</i> Schaufuss, <i>Pityophthorus</i>	1017	<i>omissus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	799
<i>obtusum</i> Schedl, (<i>Corthyus</i>), <i>Brachyspartus</i> (= <i>moritzi</i>)	1068		
<i>occidentalis</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopschion</i>	97		
<i>occidentalis</i> Bedel, <i>Phloeotribus</i> (= <i>scarabaeoides</i>)	230		

<i>omissus</i> Eichhoff, (<i>Tomicus</i>), <i>Orthotomicus</i> (= <i>proximus</i>)	475	<i>opinatatum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	700
<i>omissus</i> Schedl, <i>Coccotrypes</i>	605	<i>opinus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809
<i>omissus</i> Wood, <i>Gnathotrichus</i>	1036	<i>opinus</i> (Wood), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	678
<i>ommateus</i> Wood, <i>Scolytodes</i>	396	<i>opinus</i> Blackman, <i>Pityophthorus</i> (= <i>carinulatus</i>)	986
<i>oneratus</i> Schedl, <i>Xyleborus</i>	757	<i>opinus</i> Wood, (<i>Acrantus</i>), <i>Chaetoptelius</i>	103
<i>onerosus</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	452	<i>opinus</i> Wood, <i>Phloeotribus</i>	225
<i>onerosus</i> Schedl, <i>Xyleborus</i> (= <i>nascareuiformis</i>)	750	<i>opinus</i> Wood, <i>Scolytodes</i>	396
<i>onoharaeensis</i> Murayama, <i>Xyleborus</i>	757	<i>optatus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	678
<i>Onthotomicus</i> Ferrari (= <i>Orthotomicus</i>)	467	<i>opulentus</i> Schedl, <i>Xyleborus</i>	757
<i>onynganus</i> (Schedl), (<i>Letznerella</i>), <i>Scolytogenes</i>	864	<i>oralis</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	799
<i>opacicauda</i> Eggers, (<i>Xyleborus</i>), <i>Amasa</i> (= <i>resectus</i>)	684	<i>orarius</i> Bright, <i>Pityophthorus</i>	1015
<i>opacicaudulus</i> Schedl, <i>Cryptoxyleborus</i>	828	<i>orbatus</i> Blandford, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>germanus</i>)	796
<i>opacicollis</i> (Eggers), (<i>Loganius</i>), <i>Camptocerus</i>	319	<i>orbicaudatus</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	684
<i>opacicollis</i> (Eggers), (<i>Prionosceles</i>), <i>Scolytodes</i>	396	<i>orbicularis</i> Schedl, <i>Webbia</i>	831
<i>opacicollis</i> (LeConte), (<i>Micracis</i>), <i>Micracisella</i>	430	<i>orbiculatus</i> (Eggers), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	668
<i>opacicollis</i> Blackman, <i>Pseudopityophthorus</i>	974	<i>orbiculatus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800
<i>opacicollis</i> Eggers, <i>Chramesus</i>	267	<i>orbis</i> Browne, (<i>Xyleborus</i>), <i>Dryococtops</i> (= <i>laevis</i>)	558
<i>opacicollis</i> Eggers, <i>Phloeotribus</i>	225	<i>oregoui</i> Blackman, <i>Scolytus</i>	355
<i>opacicollis</i> Eggers, <i>Polygraphus</i> (= <i>pinus</i>)	296	<i>oregonis</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>pinii</i>)	520
<i>opacicollis</i> Eggers, <i>Scolytoplatypus</i>	405	<i>oregonus</i> (Blackman), (<i>Leperisinus</i>), <i>Hylesinus</i>	82
<i>opacicollis nitidus</i> Eggers, <i>Chramesus</i> (= <i>opacicollis</i>)	267	<i>orientale</i> Schedl, <i>Rhopalopschion</i>	97
<i>opacifrons</i> Beeson, (<i>Thamniurgides</i>), <i>Coccotrypes</i> (= <i>salakeensis</i>)	611	<i>orientalis</i> (Eggers), (<i>Metahylesinus</i>), <i>Hylesinopsis</i>	95
<i>opacifrons</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>pubescens</i>)	939	<i>orientalis</i> (Eggers), (<i>Ozophagus</i>), <i>Polygraphus</i>	291
<i>opacifrons</i> Reitter, <i>Pityogenes</i> (= <i>calcaratus</i>)	445	<i>orientalis</i> (Schedl), (<i>Mimiopterus</i>), <i>Mimiocturus</i>	949
<i>opacifrons</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>omissum</i>)	1060	<i>orientalis</i> Eggers, <i>Camptocerus</i>	320
<i>opacifrons</i> Schedl, <i>Amphicranus</i> (= <i>melanura</i>)	1047	<i>orientalis</i> Eggers, <i>Cryphalus</i>	886
<i>opacipeunus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>interstitialis</i>)	932	<i>orientalis</i> Eggers, (<i>Pelicerus</i>), <i>Cyrtogenius</i> (= <i>nitidus</i>)	567
<i>opacithorax</i> (Schedl), (<i>Micracis</i>), <i>Micracisella</i>	431	<i>orientalis</i> Eggers, <i>Dactylipalpus</i>	102
<i>opacithorax</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	678	<i>orientalis</i> Eggers, <i>Prennobiobius</i>	654
<i>opacithorax</i> Schedl, <i>Phloeoborus</i> (= <i>scaber</i>)	101	<i>orientalis</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>multistriatus</i>)	353
<i>opaculus</i> LeConte, (<i>Hylesinus</i>), <i>Hylurgopinus</i> (= <i>rufipes</i>)	109	<i>orientalis</i> Eggers, <i>Xyleborus</i>	757
<i>opaculus</i> LeConte, <i>Pityophthorus</i>	1017	<i>orientalis</i> Krivolutskaia, (<i>Blastophagus</i>), <i>Tomicus</i> (= <i>puellus</i>)	136
<i>opaculus</i> Schedl, <i>Sternobothrus</i>	216	<i>orientalis</i> Kurenzov, <i>Dryocoetes</i> (= <i>infuscatus</i>)	583
<i>opacus</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	937	<i>orientalis</i> Schedl, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>braderi</i>)	859
<i>opacus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	864	<i>orientalis</i> Schedl, <i>Aphanarthrum</i>	619
<i>opacus</i> Blackman, <i>Scolytus</i>	355	<i>orientalis</i> Schedl, <i>Cardiocratus</i>	283
<i>opacus</i> Broun, (<i>Acrantus</i>), <i>Chaetoptelius</i> (= <i>mnudulus</i>)	103	<i>orientalis</i> Schedl, <i>Chortastus</i>	312
<i>opacus</i> Eggers, <i>Diamerus</i>	197	<i>orientalis</i> Wood & Yin, <i>Ips</i>	514
<i>opacus</i> Erichson, <i>Hylastes</i>	57	<i>orientalis aceris</i> Kurenzov, <i>Xyleborus</i> (= <i>orientalis</i>)	757
<i>opacus</i> Schedl, <i>Hypocryphalus</i> (= <i>mangiferae</i>)	870	<i>orientalis kalopanaxiae</i> Kurenzov, <i>Xyleborus</i> (= <i>orientalis</i>)	757
<i>opacus</i> Schedl, <i>Pseudochramesus</i>	262	<i>orientalis pilosiusculus</i> Kurenzov, <i>Dryocoetes</i> (= <i>infuscatus</i>)	554
<i>opacus</i> Schedl, <i>Xyloctonus</i>	840	<i>orinocanus</i> Wood, <i>Scolytopsis</i>	320
<i>opacus</i> Wood, <i>Cnemionyx</i>	317	<i>orinocensis</i> Wood, <i>Chramesus</i>	267
<i>opacus</i> Wood, <i>Scolytodes</i>	396	<i>ornatus</i> (Swaine), (<i>Orthotomicus</i>), <i>Pityokteines</i>	463
<i>opalescens</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	683	<i>ornatus</i> Blackman, <i>Pityophthorus</i>	1015
<i>oparunus</i> (Beeson), (<i>Xyleborus</i>), <i>Eucallacea</i>	692	<i>orni</i> Fuchs, <i>Hylesinus</i>	82
<i>operosus</i> Schedl, <i>Xyleborus</i>	757	<i>Orosiotes</i> Niisima (= <i>Cyrtogenius</i>)	562
<i>opienensis</i> Tsai & Li, <i>Cryphalus</i> (= <i>sinoabietis</i>)	895	<i>Orthaspistes</i> Hagedorn (= <i>Tiarophorus</i>)	539

<i>Orthotomicus</i> Ferrari	467	<i>pallidus</i> Eichhoff, <i>Cryphalus</i>	886
<i>Orthotomides</i> Wood (= <i>Pityokteines</i>)	459	<i>pallidus</i> Escalera, <i>Tomicus</i> (= <i>pini</i> <i>perda</i>)	128
<i>ostriacolens</i> Bright, <i>Pityophthorus</i>	1018	<i>pallidus</i> Hopkins, <i>Hypothenemus</i> (= <i>plumeriae</i>)	938
<i>osumiensis</i> Murayama, <i>Phlocosinus</i>	249	<i>pallidus</i> Schedl, <i>Microcorthyus</i>	1066
<i>osumiensis</i> Murayama, <i>Xyleborus</i>	758	<i>pallipes</i> Sturm, <i>Ips</i> (= <i>pini</i>)	517
<i>ougeniae</i> Wood, <i>Xylechinus</i>	116	<i>palmarum</i> Eggers, <i>Coccotrypes</i>	608
<i>oralicollis</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>wallacei</i>)	695	<i>palmeri</i> Hopkins, <i>Coptoborus</i>	664
<i>oralicollis</i> Schedl, <i>Hypocryphalus</i>	871	<i>palmi</i> Hansen, <i>Trypophloeus</i>	846
<i>oralis</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	396	<i>palmicola</i> Hornung, (<i>Bostrichus</i>), <i>Coccotrypes</i> (= <i>dactyliperda</i>)	601
<i>ovalis</i> Eggers, (<i>Phloeophthorus</i>), <i>Phlocotribus</i> (= <i>cristatus</i>)	220	<i>pampasue</i> Schedl, <i>Pityophthorus</i>	1018
<i>oralis</i> Schedl, <i>Chramesus</i>	267	<i>panamensis</i> (Blandford), (<i>Loganius</i>), <i>Cnemomyx</i>	317
<i>oratus</i> (Eggers), (<i>Dryotonus</i>), <i>Phlocotribus</i>	225	<i>panamensis</i> Blackman, <i>Chramesus</i> (= <i>pumilus</i>)	268
<i>oratus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	678	<i>panamensis</i> Blackman, <i>Cucinus</i> (= <i>setulosus</i>)	211
<i>oratus</i> Chapuis, <i>Phlocoborus</i> (= <i>asper</i>)	98	<i>panamensis</i> Blackman, <i>Gnatholeptus</i>	970
<i>oratus</i> Wood, <i>Micraxis</i>	433	<i>panamensis</i> Blandford, <i>Corthyus</i>	1076
<i>oricollis</i> Blandford, <i>Tricolus</i>	1044	<i>panamensis</i> Wood, <i>Scolytodes</i> (= <i>atratus</i>)	389
<i>oxyurus</i> Schedl, <i>Cryptoxyleborus</i>	829	<i>pandae</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	692
<i>Ozodendron</i> Schedl 1957 (= <i>Cyrtogenius</i>)	562	<i>pandulus</i> (Wood), (<i>Xyleborus</i>), <i>Taurodemus</i>	785
<i>Ozodendron</i> Schedl 1964 (= <i>Cyrtogenius</i>)	562	<i>pannuceus</i> Wood, <i>Scolytodes</i>	396
<i>Ozopemon</i> Hagedorn	588	<i>papatrae</i> Schedl, <i>Xyleborus</i>	758
<i>Ozophagus</i> Eggers (= <i>Polygraphus</i>)	284	<i>papuae</i> Wood, <i>Cyrtogenius</i>	567
<i>pabo</i> Sampson, <i>Webbia</i>	831	<i>papuanus</i> (Eggers), (<i>Dendriurgus</i>), <i>Coccotrypes</i>	608
<i>Pachycotes</i> Sharp	187	<i>papuanus</i> (Eggers), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	567
<i>pachylobius</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	482	<i>papuanus</i> (Schedl) 1974, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	864
<i>Pachynoderes</i> Schedl (= <i>Hypothenemus</i>)	905	<i>papuanus</i> (Schedl), (<i>Margadillius</i>), <i>Erioporus</i>	855
<i>pacificus</i> (Beeson), (<i>Hylesinus</i>), <i>Ficicis</i>	91	<i>papuanus</i> (Schedl), (<i>Phlocosinus</i>), <i>Hyledius</i>	261
<i>pacificus</i> (Schedl), (<i>Lepicerinus</i>), <i>Scolytogenes</i>	864	<i>papuanus</i> Blandford, <i>Xyleborus</i>	758
<i>pacificus</i> Beeson, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>papuanus</i> Browne, (<i>Euptilius</i>), <i>Erioporus</i> (= <i>antennarius</i>)	854
<i>pacificus</i> Bright, <i>Phlocotribus</i>	225	<i>papuanus</i> Eggers, (<i>Pelicerus</i>), <i>Cyrtogenius</i> (= <i>papuae</i>)	567
<i>pacificus</i> Nunberg, <i>Xyleborus</i> (= <i>vulcanus</i>)	783	<i>papuanus</i> Eggers, <i>Hylesinus</i> (= <i>cordipennis</i>)	77
<i>pacificus</i> Schedl, <i>Ptilopodius</i>	900	<i>papuanus</i> Eggers, <i>Ozopemon</i>	590
<i>pacificus</i> Wood, <i>Phlocosinus</i>	249	<i>papuanus</i> Eggers, <i>Scolytplatypus</i>	405
<i>padi</i> Krivolutskaya, <i>Cryphalus</i> (= <i>scopiger</i>)	894	<i>papuanus</i> Schedl 1979, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>paradoxus</i>)	864
<i>padi</i> Stark, <i>Dryocoetes</i>	584	<i>papuanus</i> Schedl, (<i>Xylechinus</i>), <i>Phlocosinopsioides</i> (= <i>triseriatus</i>)	239
<i>padus</i> Beeson, (<i>Alniaphagus</i>), <i>Xylechinus</i> (= <i>padus</i>)	117	<i>papuanus</i> Schedl, (<i>Xylocryptus</i>), <i>Scolytogenes</i> (= <i>papuensis</i>)	864
<i>padus</i> Wood, <i>Xylechinus</i>	117	<i>papuanus</i> Schedl, <i>Cosmoderes</i>	902
<i>paganus</i> Eichhoff, <i>Cryphalus</i>	886	<i>papuanus</i> Schedl, <i>Cryphalus</i> (= <i>tetricus</i>)	897
<i>Pagiocerus</i> Eichhoff	213	<i>papuanus</i> Schedl, (<i>Eidophelus</i>), <i>Cyrtogenius</i> (= <i>papuensis</i>)	567
<i>palaquius</i> (Schedl), (<i>Phloeochilus</i>), <i>Liparthrum</i>	279	<i>papuanus</i> Schedl, <i>Scolytus</i> (= <i>multistriatus</i>)	353
<i>palatus</i> Wood, <i>Xyleborus</i>	758	<i>papuensis</i> Wood, <i>Cyrtogenius</i>	567
<i>palauensis</i> (Wood), (<i>Phlocotrypetus</i>), <i>Liparthrum</i>	279	<i>papuensis</i> Wood, <i>Scolytogenes</i>	864
<i>palauanus</i> Eggers, <i>Sphacotrypes</i>	200	<i>papulaus</i> Eichhoff, <i>Corthyus</i>	1076
<i>palauanus</i> Schedl, <i>Cryphalus</i>	886	<i>paraconfusus</i> Lanier, <i>Ips</i>	514
<i>palearis</i> Wood, <i>Phlocosinus</i>	250	<i>paraconrexa</i> Bright, Schedl	822
<i>paleatus</i> Blandford, <i>Cucinus</i>	210	<i>Paracorthyus</i> Wood (= <i>Metacorthyus</i>)	1064
<i>palembangensis</i> Schedl, <i>Xyleborus</i> (= <i>quadrispinosulus</i>)	768	<i>paradoxus</i> (Schedl), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658
<i>pallens</i> Wood, <i>Thysanoes</i>	420	<i>paradoxus</i> Schedl, <i>Hypothenemus</i>	938
<i>palliatus</i> (Gyllenhal), (<i>Hylesinus</i>), <i>Hyhurgops</i>	35	<i>paradoxus</i> Wood, <i>Scolytogenes</i>	864
<i>pallidipennis</i> (Eggers), (<i>Xyleborus</i>), <i>Cnestus</i>	803	<i>Paraglostatus</i> Schedl (= <i>Glostatus</i>)	835
<i>pallidipennis</i> Murayama, (<i>Tosaxyleborus</i>), <i>Cnestus</i> (= <i>murayamai</i>)	802		
<i>pallidum</i> (Chapuis), (<i>Nemobius</i>), <i>Pycnarthrum</i>	385		
<i>pallidus</i> (Blackman), (<i>Neodryocoetes</i>), <i>Araptus</i>	960		

- paraguayensis* Eggers, *Dendrosinus* 237
- paraguayensis* Hopkins, (*Stephanoderes*),
 Hypothenemus (= *cruidiae*) 915
- paraguayensis* Schedl, (*Xyleborus*), *Xyleborinus*
 (= *saxeseni*) 815
- paraguayensis* Schedl, *Cnesinus* 210
- parallelocollis* Chapuis, *Dendroctonus* 158
- parallelocollis* Eggers, *Xyleborus* 758
- parallelocollis* Eichhoff, *Crypturgus* (= *pusillus*) . . . 628
- parallelus* (*Hopkins*), (*Stephanoderes*),
 Hypothenemus 938
- parallelus* (Schedl), (*Hexacolus*), *Scolytodes* 397
- parallelus* Chapuis, *Hylastes* 59
- parallelus* Eggers, *Xyleborus* 758
- parallelus* Schedl, *Dendrocraniulus*
 (= *guatemalensis*) 552
- paranae* (Schedl), (*Brachydendrulus*), *Araptus* 960
- Parasphaerotrypes* Murayama (= *Sucus*) 191
- Paraxyleborus* Hoffmann (= *Gnathotrichus*) 1034
- parcellus* Wood, *Xyleborus* 758
- parcius* (Schedl), (*Ips*), *Acanthotomicus* 482
- parcius* Schedl, *Hypothenemus* (= *eruditus*) 925
- parcus* Wood, *Chramesus* 267
- pardous* Eggers, *Xyleborus* 758
- parfentievi* Pjatnitskii, *Pityophthorus* 1018
- parilis* Schedl, *Hypothenemus* (= *eruditus*) 925
- parilis* Wood, *Amphicranus* 1048
- parilis* Wood, *Dendroterus* 951
- parilis* Wood, *Pityophthorus* 1018
- parinarie* Schedl, *Xyleborus* 758
- parinari* (*Browne*), (*Ozodendron*), *Cyrtogenius* . . . 567
- parinari* Hopkins, *Ozopemon* 590
- parkinsoniae* (*Blackman*), (*Renocis*),
 Chaetophlocus 273
- parkinsoniae* Blackman, *Hylocurus* 426
- parricida* Eggers, *Dactylipalpus* 102
- parsus* Wood, *Tricolus* 1044
- partilis* Wood, *Tricolus* 1044
- partitus* Browne, *Xyleborus* 758
- parva* (*Lea*), (*Xylopertha*), *Coptodryas* 826
- parva* Blackman, *Phrixosoma* 190
- parvatis* (*Wood*), (*Cryphalomorphus*),
 Scolytogenes 864
- parviclariger* Yin & Huang, *Scolytus* 356
- parvicornis* Wood, *Cnesinus* 210
- parvidentatus* Schedl, *Scolytplatypus*
 (= *eichelbaumi*) 402
- parvior* Browne, *Xyleborus* (= *amphicranoides*) . . . 711
- parvipunctatus* Eggers, *Xyleborus* 758
- parvispinosus* Schedl, *Xyleborus*
 (= *quadrspinosulus*) 768
- parvispinosus* Palembangensis Schedl, *Xyleborus*
 (= *quadrspinosulus*) 768
- parvulus* Blandford, *Corthylus* 1077
- parvulus* Browne, *Hypothenemus* 938
- parvulus* Eggers, *Margadillius* 899
- parvulus* Eichhoff, *Xyleborus* (= *similis*) 773
- parvulus* Ferrari, *Microcorthylus* 1066
- parvulus* Murayama, *Polygraphus* 291
- parvulus* Nisima, *Cryphalus* 886
- parvulus* Wood, *Scolytodes* 397
- parvum* (*Eggers*), (*Anchonoceus*), *Monarthrum* . . . 1060
- parvus* (*Eggers*), (*Hylocurosoma*), *Scolytodes* 397
- parvus* (*Hopkins*), (*Hypothenoides*), *Scolytogenes* . . 864
- parvus* (*Nunberg*), (*Loganius*), *Cucmonyx* 317
- parvus* Beeson, (*Thannurgides*), *Coccotrypes*
 (= *nubilus*) 607
- parvus* Blackman, (*Tachyderes*), *Cryptocarenum*
 (= *herae*) 903
- parvus* Blackman, *Hylastes* (= *tenuis*) 63
- parvus* Browne, *Cryphalus* 886
- parvus* Browne, *Cyrtogenius* 567
- parvus* Eggers, *Diamerus* 197
- parvus* Eggers, *Hylurgops* (= *palliatu*) 38
- parvus* Eichhoff, *Xyleborus* (= *affinis*) 708
- parvus* Hopkins, *Hypothenemus* (= *eruditus*) 922
- parvus* Sampson, *Coccotrypes* 609
- parvus* Sampson, *Scolytplatypus* 405
- parvus* Wood, *Phloeocleptus* 422
- passerini* Keremidiev, *Phloeosinus* (= *sp.?*) 239
- patchi* Blackman, *Pityophthorus* (= *balsameus*) . . . 983
- paucegranulatus* Eggers, *Scolytplatypus* 405
- pauciconcavus* Schedl, *Gnathotrupes* 1042
- paucus* Wood, *Ambrosiodinus* 678
- pauliani* (*Lepesme*), (*Kissophagus*), *Hylesinopsis* . . 95
- paulus* Wood, *Pityophthorus* 1019
- pauper* Endrodi, *Trypodendron* (= *lineatum*) 636
- paykulli* Duftschmidt, (*Hylesinus*), *Hylurgops*
 (= *glabratus*) 33
- peccanis* Hopkins, (*Stephanoderes*), *Hypothenemus*
 (= *seriatus*) 943
- peccanus* Hopkins, (*Xyleborus*), *Xyleborinus*
 (= *saxeseni*) 815
- pectinicornis* Balachowsky, *Phloeotribus*
 (= *cristatus*) 220
- pedella* (Schedl), (*Xyleborus*), *Coptodryas* 826
- pegani* Eggers, *Thannurgus* 545
- peguensis* Eggers, (*Xyleborus*), *Leptoxyleborus*
 (= *concisus*) 659
- pele* Samuelson, *Xyleborus* 758
- pelicerinus* Schedl, (*Hexacolus*), *Scolytodes*
 (= *notatus*) 396
- Pelicerus* Eggers (= *Cyrtogenius*) 562
- peliciformis* (Schedl), (*Xyleborus*), *Peridryocoetes* . 562
- pelicipennis* (Schedl), (*Hexacolus*), *Scolytodes* . . . 397
- peliohi* Beaver, *Dryocoetops* 588
- pelliculosum* Hagedorn, *Cyclorhipidion* 701
- pelliculosus* Eichhoff, *Xyleborus* 758
- pellitus* (Schedl), (*Erioschidias*), *Cosmoderes* 902
- pellitus* Schedl, (*Hylesinus*), *Ficicis* (= *pacificus*) . . 91
- pellitus* Schedl, *Ambrosiodinus* (= *lewisi*) 676
- pellitus* Schedl, *Pityophthorus* 1019
- peltatus* Wood, *Tricolus* 1044
- penangensis* (Schedl), (*Phloeosinus*), *Hyledius* . . . 261

<i>penangensis</i> Brownie, (<i>Streptocranus</i>), <i>Coptoborus</i> (= <i>capucinulus</i>)	663	<i>perfacilis</i> Eggers, <i>Ozopemon</i>	590
<i>penicillatum</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1060	<i>perfoliatus</i> (Wollaston), (<i>Phloeophthorus</i>), <i>Phlocotribus</i>	225
<i>penicillatus</i> (Bruck), (<i>Renocis</i>), <i>Chaetophloeus</i>	273	<i>perforans</i> (Wollaston), (<i>Tomicus</i>), <i>Xyleborus</i>	759
<i>penicillatus</i> (Hagedorn), (<i>Xyleborus</i>), <i>Taphrodasus</i>	825	<i>perforatus</i> Beeson, <i>Pityogenes</i> (= <i>spessirtseri</i>)	455
<i>penicillatus</i> Reitter, <i>Scolytus</i> (= <i>intricatus</i>)	334	<i>perhispidus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	938
<i>penicillus</i> Schedl, <i>Scolytus</i> (= <i>propinguus</i>)	357	<i>Peridryocoetes</i> Schedl (= <i>Peridryocoetes</i> Wood)	561
<i>peniculus</i> Wood, <i>Chramesus</i>	267	<i>Peridryocoetes</i> Wood	561
<i>peniculus</i> Wood, <i>Pseudothysanoes</i>	415	<i>Periocryphalus</i> Wood	947
<i>pennatum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1060	<i>periosus</i> Wood, <i>Chramesus</i>	267
<i>pennatus</i> (Schedl), (<i>Micracis</i>), <i>Pseudomicracis</i>	429	<i>peritus</i> Blandford, <i>Hypothenemus</i> (= <i>birmanus</i>)	910
<i>pennatus</i> Schedl, <i>Microcryphalus</i>	411	<i>perkinsi</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910
<i>pennatus</i> Schedl, <i>Sampsonius</i>	656	<i>perlaetus</i> Schedl, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>pelliculosum</i>)	701
<i>pennidens</i> (Reitter), (<i>Tomicus</i>), <i>Pityogenes</i>	454	<i>perlaetus</i> Schedl, <i>Polygraphus</i>	291
<i>pentaclethrae</i> Schedl, <i>Pityophthorus</i>	1019	<i>perlatus</i> Chapuis, <i>Phlocosinus</i>	250
<i>pentaclethrae</i> Schedl, <i>Xyleborus</i>	759	<i>perlongus</i> Eggers, <i>Xyleborus</i>	762
<i>pentacme</i> Wood, <i>Sphaerotrypes</i>	200	<i>permagnus</i> (Eggers), (<i>Prionosceles</i>), <i>Scolytodes</i>	397
<i>perakensis</i> (Schedl), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	567	<i>permarginatus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	678
<i>perakensis</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>wallacei</i>)	695	<i>perminimus</i> (Schedl), (<i>Cryphalus</i>), <i>Hypocryphalus</i>	871
<i>peramplioides</i> Schedl, <i>Xyleborus</i>	759	<i>perminor</i> Schedl, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>minor</i>)	654
<i>peramplus</i> Schedl, <i>Xyleborus</i>	759	<i>perminutissimus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809
<i>perappositus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	935	<i>permirus</i> Schauffuss, <i>Scolytoplatypus</i>	405
<i>perbrevis</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>fornicatus</i>)	690	<i>pernanulus</i> Schedl, <i>Trichus</i>	1044
<i>percorthyloides</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>corthyloides</i>)	790	<i>pernicosus</i> Wood, <i>Gnathotrichus</i>	1036
<i>percorthylus</i> (Schedl), (<i>Xyleborus</i>), <i>Taphrodasus</i>	825	<i>pernicosus</i> Wood, <i>Phlocotribus</i>	226
<i>percristatus</i> Eggers, <i>Xyleborus</i>	759	<i>pernitidus</i> Eggers, (<i>Poecilips</i>), <i>Coccotrypes</i> (= <i>adrena</i>)	592
<i>percuneolus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809	<i>pernitidus</i> Schedl, <i>Xyleborus</i> (= <i>minutus</i>)	751
<i>perdeclivis</i> Schedl, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>declivis</i>)	653	<i>pernodulus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>nodulosus</i>)	677
<i>perdiligens</i> Schedl, <i>Trichus</i>	1044	<i>Peruophorus</i> Strohmeyer	202
<i>perdiligens</i> Schedl, <i>Xyleborus</i>	759	<i>pernotus</i> Schedl, <i>Xyleborus</i>	762
<i>perdiligens diligens</i> Schedl, <i>Xyleborus</i> (= <i>spheos</i>)	775	<i>perorientalis</i> Schedl, <i>Xyleborus</i> (= <i>orientalis</i>)	757
<i>perditor</i> Blandford, <i>Coccotrypes</i>	609	<i>perotei</i> Blackman, <i>Pityophthorus</i>	1019
<i>perditum</i> Wood, <i>Pycnarthrum</i>	355	<i>perparva</i> (Sampson), (<i>Xyleborus</i>), <i>Coptodryas</i>	826
<i>perditus</i> Wood, <i>Scolytodes</i>	397	<i>perpilosellum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701
<i>perdlx</i> Schedl, <i>Xyleborus</i> (= <i>javanus</i>)	746	<i>perplexus</i> Schedl, <i>Glostatus</i>	836
<i>perduratum</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>dimidiatum</i>)	1050, 1054	<i>perplexus</i> Schedl, <i>Scolytodes</i>	397
<i>perebeae</i> (Ferrari), (<i>Amphicranus</i>), <i>Taurodenus</i>	755	<i>perplexus</i> Schedl, <i>Xyleborus</i>	762
<i>peregrinus</i> (Chapuis), (<i>Hylastes</i>), <i>Pachycotes</i>	187	<i>perplexus</i> Wood, <i>Carphoborus</i>	307
<i>peregrinus</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	452	<i>perplexus</i> Wood, <i>Cuesinus</i>	210
<i>peregrinus</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>carpini</i>)	327	<i>perpunctatus</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	938
<i>peregrinus</i> Eggers, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxeseni</i>)	815	<i>perpunctatus</i> Schedl, <i>Xyleborus</i>	762
<i>peregrinus</i> Eichhoff, <i>Pityophthorus</i>	1019	<i>perpusillus</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809
<i>peregrinus</i> Schedl, <i>Cyrtogenius</i>	565	<i>perpusillus</i> Wood, <i>Scolytodes</i>	397
<i>peregrinus</i> Wood & Yin, <i>Pseudopityophthorus</i>	974	<i>perquadrispinosus</i> Schedl, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>quadrispinosus</i>)	654
<i>perexiguus</i> (Blandford), (<i>Tomicus</i>), <i>Acanthotomicus</i>	452	<i>perrisi</i> (Chapuis), (<i>Hylesinus</i>), <i>Carphoborus</i>	307
<i>perexiguus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809		
<i>perexiguus</i> Wood, <i>Pityophthorus</i>	1019		

- perrisi* Guillebeau, (*Phloeophthorus*), *Phlocotribus*
(=*spinulosus*) 234
- perroti* Swaine, *Ips* 515
- perseae* Wood, *Pseudothysanoes* 415
- persicae* Hopkins, (*Thanaurgides*), *Coccotrypes*
(=*adrena*) 592
- persimilis* Eggers, *Hylesinus* (= *nigrinus*) 52
- persimilis* Eggers, *Xyleborus* 762
- perspectus* (Schedl), (*Plesiophthorus*),
Dendroterus 951
- persphenos* Schedl, (*Xyleborus*), *Xyleborinus*
(=*andreweesi*) 805
- perspinidens* (Schedl), (*Xyleborus*), *Premnobius* 654
- perspiner* Schedl, (*Xyleborus*), *Premnobius*
(=*spinifer*) 655
- pertinax* (Schedl), (*Hylesinus*), *Ficicis* 91
- pertortuosus* (Schedl), (*Xyleborus*),
Ambrosiodinus 678
- pertuberculatus* Eggers, (*Xyleborus*), *Coptodryas*
(=*mugax*) 526
- pertuberculatus* Eggers, *Phloeosinus* 250
- perturbatus* (Eichhoff), (*Tomicus*), *Ips* 515
- pertusus* (Wood), (*Gnathophthorus*),
Dacnophthorus 969
- peruanum* (Schedl), (*Pterocyclon*), *Monarthrum* 1060
- peruanum* Schedl, *Monarthrum* (= *peruvianum*) 1060
- peruanus* Eggers, *Scolytopsis* 320
- peruanus* Schedl, *Chramesus* 267
- peruanus* Schedl, *Corthylyus* 1077
- peruensis* Schedl, *Scolytus* 356
- peruensis* Schedl, *Phloeotribus* 226
- peruviana* Eggers, *Phrixosoma* 190
- peruvianum* Wood, *Monarthrum* 1060
- peruvianus* Schedl, *Xyleborus* 762
- perversus* Hagedorn, (*Xyleborus*), *Taurodemus*
(=*varians*) 786
- petilus* Wood, *Corthylyus* 1077
- petioli* (Browne), (*Pocilips*), *Coccotrypes* 609
- petioli* Beaver, *Dryocoetops* 555
- petzi* Reitter, *Thanaurgus* 545
- pexus* Schedl, *Cryphalus* 886
- peyerimhoffi* (Eggers), (*Phloeophthorus*),
Phlocotribus 226
- peyerimhoffi* Pfeffer, *Liparthrum* (= *genistae*) 277
- pfelti* (Ratzeburg), (*Bostrichus*), *Xyleborus* 762
- Phacrylus* Schedl (= *Acorthylyus*) 552
- pharax* Schedl, *Corthylyus* 1077
- phascoli* Eggers, (*Stephanoderes*), *Hypothenemus*
(=*eruditus*) 919
- Phellodendrophagus* Krivolutskaya (= *Eidophelus*) 867
- Phelloterus* Wood 971
- philippinensis* (Eggers), (*Hylesinus*), *Ficicis* 91
- philippinensis* (Eggers), (*Neoxylotomus*),
Scolytomimus 539
- philippinensis* (Eggers), (*Pelicerus*), *Cyrtogenius* 565
- philippinensis* (Schedl), (*Erioschidias*),
Cosmoderes 902
- philippinensis* Eggers, (*Dendrugus*), *Coccotrypes*
(=*adrena*) 592
- philippinensis* Eggers, (*Ips*), *Acanthotomicus*
(=*perexiguus*) 452
- philippinensis* Eichhoff, *Xyleborus* (= *perforans*) 761
- philippinensis* Hopkins, (*Stephanoderes*),
Hypothenemus (= *javanus*) 932
- philippinensis* Schedl, (*Phloeosinus*), *Hyledius*
(=*nitidicollis*) 261
- philippiensis* Schedl, *Coccotrypes* (= *adrena*) 592
- philippinensis* Schedl, *Pityophthorus* 1019
- philippinensis* Schedl, *Webbia* 532
- philippinensis* Strohmeier, *Sphaerotrypes* 200
- Phloeobori* Blandford (= *Hylesinini*) 64
- Phloeoborini* Blandford (= *Hylesinini*) 64
- Phloeoborus* LeConte 97
- Phlocochilus* Schedl (= *Liparthrum*) 274
- Phloeocleptus* Wood 421
- Phloeocranus* Schedl 236
- Phloeocurus* Wood 429
- Phloeoditica* Schedl 236
- Phlocographus* Wood (= *Halystus*) 253
- Phloeophthorus* Wollaston (= *Phlocotribus*) 217
- Phloeosina* Nusslin (= *Phloeosinini*) 236
- Phloeosinides* Nusslin (= *Phloeosinini*) 236
- Phloeosinina* Nusslin (= *Phloeosinini*) 236
- Phloeosinini* Nusslin 236
- Phloeosinites* Hagedorn (= *Phloeosinus*) 239
- phloeosinites* Schedl, *Chramesus* 267
- phloeosinoides* Browne, *Phlocoditica* 236
- Phloeosinopsioides* Schedl 239
- Phloeosinopsis* Schedl 1964 (= *Phloeosinopsioides*) 239
- Phloeosinopsis* Schedl 1936 (= *Hyledius*) 259
- Phloeosinus* Chapuis 239
- Phloeoterus* Wood 952
- Phloeotribi* Chapuis (= *Phloeotribini*) 217
- Phloeotribidae* Chapuis (= *Phloeotribini*) 217
- Phloeotribina* Chapuis (= *Phloeotribini*) 217
- Phloeotribinae* Chapuis (= *Phloeotribini*) 217
- Phloeotribini* Chapuis 217
- phloeotriboides* Schedl, *Chramesus* 265
- Phloeotribus* Latreille 217
- Phloeotrogus* Motschulsky (= *Ambrosiodinus*) 670
- Phloeotrupes* Erichson (= *Phlocoborus*) 97
- Phloeotrupidae* Chapuis, part (= *Hylesinini*) 64
- Phloeotrupides* Chapuis, part (= *Hylesinini*) 64
- Phloeotrupinae* Chapuis, part (= *Hylesinini*) 64
- Phloeotrupini* Chapuis, part (= *Hylesinini*) 64
- Phlocotrypsetus* Wood (= *Liparthrum*) 274
- Phloiortribus* Latreille (= *Phlocotribus*) 215
- phoebe* Wood, *Phloeosinus* 250
- phoebeae* Wood, *Scolytodes* 397
- phoenicola* Beeson, *Coccotrypes* (= *carpophagus*) 596
- phoradendri* Blackman, *Pseudothysanoes* 415
- phoradendri* Wood, *Chaetophloeus* 274
- photophilus* Wood, *Micracis* (= *sucainei*) 434
- Phrixosoma* Blandford 159

<i>Phrixosomini</i> Wood	189	<i>pilipeme</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701
<i>Phthorius</i> Eichhoff (= <i>Monarthrum</i>)	1050	<i>pilipunctatus</i> Browne, <i>Xyleborus</i>	763
<i>Plithorophloeoides</i> Nusslin (= <i>Phloeotribini</i>)	217	<i>pilosellus</i> (Browne), (<i>Minips</i>), <i>Acanthotomicus</i>	482
<i>Plithorophloeus</i> Rey (= <i>Phloeotribus</i>)	217	<i>pilosellus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	809
<i>phyllocladus</i> Bright, (<i>Phloeosinus</i>)	250	<i>pilosellus</i> Erichson, <i>Cryphalus</i>	889
<i>piceae</i> (Ratzeburg), (<i>Bostrichus</i>), <i>Cryphalus</i>	886	* <i>pilosellus</i> Schedl, <i>Hylurgops</i>	39
<i>piceae</i> (Swaine), (<i>Eccoptogaster</i>), <i>Scolytus</i>	356	<i>pilosiusculus</i> Kurenzov, <i>Dryocoetes</i> (= <i>infuscatus</i>)	583
<i>piceae</i> Blackman, <i>Pityophthorus</i> (= <i>lautus</i>)	1006	<i>pilosulum</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701
<i>piceae</i> Hopkins, <i>Dryocoetes</i> (= <i>affaber</i>)	571	<i>pilosulus</i> (Eggers), (<i>Problechilus</i>), <i>Gymnochilus</i>	387
<i>piceae</i> Pfeffer, <i>Hylastes</i> (= <i>substriatus</i>)	62	<i>pilosulus</i> Browne, <i>Cryphalus</i>	889
<i>piceae</i> Swaine, <i>Phloeosinus</i> (= <i>pini</i>)	250	<i>pilosulus</i> LeConte, (<i>Cryphalus</i>), <i>Gnathotrichus</i> (= <i>pilosus</i>)	1036
<i>piceae</i> Swaine, <i>Phloeotribus</i>	226	<i>pilosulus</i> Schedl, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>piceae</i> Wood, <i>Carphoborus</i>	308	<i>pilosus</i> (Eggers), (<i>Ips</i>), <i>Acanthotomicus</i>	483
<i>piceaperda</i> Hopkins, <i>Dendroctonus</i> (= <i>rufipennis</i>)	177	<i>pilosus</i> (Eggers), (<i>Xyleborus</i>), <i>Arixyleborus</i>	668
<i>piceum</i> Strand, <i>Trypodendron</i> (= <i>laeve</i>)	636	<i>pilosus</i> (LeConte), (<i>Cryphalus</i>), <i>Gnathotrichus</i>	1036
<i>piceus</i> (Blandford), (<i>Hexacolus</i>), <i>Scolytodes</i>	397	<i>pilosus</i> (Murayama), (<i>Pseudopoeccilips</i>), <i>Taphrozylus</i>	560
<i>piceus</i> (Motschulsky), (<i>Anodius</i>), <i>Euwallacea</i>	692	<i>pilosus</i> (Ratzeburg), (<i>Hylesinus</i>), <i>Xylechinus</i>	117
<i>piceus</i> Blackman, <i>Scolytoplatypus</i> (= <i>javanus</i>)	404	<i>pilosus</i> Blandford, <i>Dryocoetes</i>	584
<i>piceus</i> Bright, <i>Pityophthorus</i> (= <i>deletus</i>)	994	<i>pilosus</i> Eggers, (<i>Hyorrhynchus</i>), <i>Sucus</i> (= <i>niisimai</i>)	191
<i>piceus</i> Eggers, <i>Cryphalus</i>	889	<i>pilosus</i> Eggers, <i>Cuesinus</i>	210
<i>piceus</i> Marsham, (<i>Ips</i>), <i>Hylurgops</i> (= <i>palliatius</i>)	38	<i>pilosus</i> Eggers, <i>Cryptocarenum</i>	903
<i>piceus</i> Stephens, (<i>Hylurgus</i>), <i>Hylastinus</i> (= <i>obscurus</i>)	67	<i>pilosus</i> Eggers, <i>Dendrocraniulus</i>	553
<i>picecauda</i> Schedl, <i>Webbia</i>	832	<i>pilosus</i> Eggers, <i>Hylastinus</i> (= <i>obscurus</i>)	67
<i>picicolor</i> Stephens, <i>Scolytus</i> (= <i>intricatus</i>)	334	<i>pilosus</i> Hopkins, <i>Hypothenemus</i>	938
<i>picinus</i> Schedl, <i>Xyleborus</i> (= <i>africanus</i>)	709	<i>pilosus</i> Sasaki, <i>Cryphalus</i> (= <i>exiguus</i>)	880
<i>picipennis</i> Eggers, <i>Dryocoetes</i>	584	<i>pilosus</i> Schedl, <i>Hylcurus</i>	426
<i>picipennis</i> Eggers, <i>Phloeotribus</i>	226	<i>pilosus</i> Tsai & Li, <i>Cryphalus</i>	889
<i>picipennis</i> Stephens, <i>Hylesinus</i> (= <i>rarius</i>)	89	<i>pilosus</i> Wood, <i>Xylechinusomus</i>	105
<i>Piezorhopalus</i> Guerin-Meneville (= <i>Amphicranus</i>)	1045	<i>pilosus</i> Yin & Huang, <i>Scolytus</i>	356
* <i>piger</i> Wickham, <i>Hylurgops</i>	38	<i>pilula</i> (Erichson), (<i>Hylesinus</i>), <i>Phloeotribus</i>	226
<i>pila</i> Blandford, <i>Sphaerotrypes</i>	200	<i>pilula</i> Hagedorn, <i>Xylosandrus</i> (= <i>mutillatus</i>)	799
<i>pilatus</i> Wood, <i>Cuesinus</i>	210	<i>pinelioides</i> (Schaufuss), (<i>Lissoclastus</i>), <i>Diamerus</i>	197
<i>pileatulus</i> Schedl, <i>Xyleborus</i>	763	<i>pinastri</i> Bechstein, (<i>Bostrichus</i>), <i>Ips</i> (= <i>sexdentatus</i>)	525
<i>pileatus</i> Wood, <i>Dryocoetoides</i>	688	<i>pinavorus</i> Bright, <i>Pityophthorus</i>	1019
<i>pilicornis</i> Wood, <i>Trietennus</i>	541	<i>pindrow</i> Beeson, <i>Pityophthorus</i> (= <i>deodara</i>)	994
<i>pilidens</i> Reitter, <i>Pityogenes</i> (= <i>bistridentatus</i>)	444	<i>Pinetoscolytus</i> Butovitch (= <i>Scolytus</i>)	322
<i>pilidens albanicus</i> Apfelbeck, <i>Pityogenes</i> (= <i>bistridentatus</i>)	444	<i>pinguis</i> (Blackman), (<i>Mycloborus</i>), <i>Pityophthorus</i>	1019
<i>pilifer</i> (Eggers), (<i>Hylesinus</i>), <i>Strombophorus</i>	204	<i>pinguis</i> Browne, <i>Xyleborus</i>	764
<i>pilifer</i> (Eggers), (<i>Problechilus</i>), <i>Gymnochilus</i>	387	<i>pinguis</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>forficuloides</i>)	807
<i>pilifer</i> (Spessivtsev), (<i>Myelophilus</i>), <i>Tomicus</i>	128	<i>pinguis</i> Wood, <i>Dendrocraniulus</i>	553
<i>pilifer</i> Eggers, <i>Cryphalus</i>	889	<i>pini</i> (Say), (<i>Bostrichus</i>), <i>Ips</i>	517
<i>pilifer</i> Eggers, <i>Xyleborus</i>	763	<i>pini</i> Blackman, <i>Cactopinus</i>	436
<i>pilifer</i> Schedl, <i>Hypocryphalus</i>	871	<i>pini</i> Eggers, <i>Cryphalus</i> (= <i>fulvus</i>)	880
<i>pilifer</i> Schedl, <i>Pityophthorus</i> (= <i>cascoensis</i>)	987	<i>pini</i> Eggers, <i>Scolytus</i> (= <i>morawitzi</i>)	345
<i>pilifer</i> Wood, <i>Carphobius</i>	311	<i>pini</i> Eichhoff, <i>Carphoborus</i>	308
<i>pilifer</i> Wood, <i>Scolytodes</i>	397	<i>pini</i> Eichhoff, <i>Xyleborus</i> (= <i>pubescens</i>)	766
<i>pilifrons</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	397	<i>pini</i> Hopkins, (<i>Carposinus</i>), <i>Cyrtogenius</i> (= <i>luteus</i>)	566
<i>pilifrons</i> Browne, (<i>Poeccilips</i>), <i>Coccotrypes</i> (= <i>cyperi</i>)	599	<i>pini</i> Hopkins, (<i>Piperius</i>), <i>Cryphalus</i> (= <i>sylicola</i>)	896
<i>pilifrons</i> Swaine, <i>Ips</i>	516	<i>pini</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942
<i>pilifrons pilifrons</i> Swaine, <i>Ips</i> (= <i>pilifrons</i>)	517	<i>pini</i> Kurenzov, <i>Pityophthorus</i>	1019
<i>pilifrons sulcifrons</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517	<i>pini</i> Niisima, <i>Dryocoetes</i>	584
<i>pilifrons thatcheri</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517	<i>pini</i> Stebbing, <i>Polygraphus</i>	291
<i>pilifrons utahensis</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517		
<i>piliger</i> Schedl, <i>Hypocryphalus</i>	871		

- pini* Swaine, *Phloeosinus* 250
- pini* Wood, *Pseudohylesinus* 112
- pini* Wood, *Pseudothysanoes* 416
- pinarius* Schedl, *Hylurdectonus* 186
- pinicola* Hopkins, *Dendroctonus* (= *frontalis*) 143
- pinicola* Bedel, *Hylastes* (= *ater*) 49
- pinicola* Eggers, *Xyleborus* 764
- pinicolens* Wood, *Carphoborus* 308
- pinifex* Fitch, (*Hylastes*), *Hylurgops* (= *rugipennis*) 40
- piniperda* (Linnaeus), (*Dermestes*), *Tomicus* 128
- pinivora* Schedl, *Orthotomicus* 473
- pinivorus* Browne, *Xylebomis* 764
- pinnatus* Eggers, *Scolytus* 356
- pinsapo* Pfeffer, *Pityophthorus* 1020
- Piperius* Hopkins (= *Cryphalus*) 872
- piscatorium* Wollaston, *Aphanarthrum* 619
- piscecauda* Browne, *Webbia* 832
- piscinus* Bright, *Corthylius* 1077
- pistaciae* Wichmann, *Phloeotribus* (= *mayeti*) 224
- pistor* Schedl, (*Stephanoderes*), *Hypothenemus* (= *javanus*) 933
- pithecolobius* (Schedl), (*Xyleborus*),
 Ambrosiodmus 679
- Pityoborus* Blackman 968
- Pityoceragenes* Balachowsky (= *Pityogenes*) 439
- Pityodendron* Schedl 971
- pityogenes* (Schedl), (*Xyleborus*), *Cyclorhipidion* 701
- Pityogenes* Bedel 438
- Pityogenina* Balachowsky, part (= *Ipini*) 438
- pityographus* (Ratzeburg), (*Bostrichus*),
 Pityophthorus 1020
- pityographus* *bibractensis* Balachowsky,
 Pityophthorus (= *pityographus*) 1021
- pityographus* *cribratus* Pfeffer, *Pityophthorus*
 (= *pityographus*) 1021
- pityographus* *maritimus* Stark, *Pityophthorus*
 (= *pityographus*) 1021
- Pityokteines* Fuchs 459
- Pityophilus* Blackman (= *Pityotrichus*) 969
- Pityophthoridae* Eichhoff, part (= *Corthylini*) 949
- Pityophthoridae* Eichhoff (= *Pityophthorina*,
 Corthylini) 949
- Pityophthorinae* Eichhoff (= *Pityophthorina*,
 Corthylini) 949
- Pityophthorina* Eichhoff, part (= *Corthylini*) 949
- pityophthorinus* (Schedl), (*Lepicerinus*),
 Scolytogenes 864
- Pityophthoroides* Blackman (= *Pityophthorus*) 977
- Pityophthorus* Eichhoff 976
- Pityotrichus* Wood 969
- placatus* Wood, *Araptus* (= *tenuis*) 962
- placetulus* Wood, *Araptus* 960
- plagiatum* (Eichhoff), (*Corthylius*), *Monarthrum* 1060
- plagiatus* (LeConte), (*Xyleborus*), *Pityogenes* 454
- plagiatus* Wood, *Phloeocleptus* 422
- planicolle* (Schedl), (*Chilodendron*), *Xylechinus* 118
- planicollis* Schedl, *Cryphalus* 889
- planicollis* Zimmermann, *Xyleborus* 764
- planipennis* Schedl, *Xyleborus* 764
- planirostris* (Chapuis), (*Hylastes*), *Hylurgops* 39
- planodeclivis* Browne, *Xyleborus* (= *ipidia*) 745
- planotruncatum* (Schedl), (*Xyleborus*),
 Cyclorhipidion 701
- plastographus* (LeConte), (*Tomicus*), *Ips* 521
- plastographus* *maritimus* Lanier, *Ips*
 (= *plastographus*) 521
- plastographus* *plastographus* (LeConte),
 (*Tomicus*), *Ips* (= *plastographus*) 521
- Platydactylus* Eichhoff (= *Eccoptopterus*) 819
- platypoides* Eggers, *Webbia* 831
- platystylus* Wichmann, *Scolytus* (= *claviger*) 328
- platyurus* Browne, *Xylebomis* 764
- plaumanni* (Schedl), (*Brachydendruhus*), *Araptus* 920
- plaumanni* (Schedl), (*Breviophthorus*),
 Pityophthorus 1021
- plaumanni* (Schedl), (*Cryptocleptes*),
 Pseudothysanoes 416
- plaumanni* (Schedl), (*Pterocyclon*), *Monarthrum* 1060
- plaumanni* Schedl, *Amphicranus* 1048
- plaumanni* Schedl, *Cresinus* 210
- plaumanni* Schedl, *Trichus* 1044
- pleiades* Samuelson, *Xyleborus* 764
- pleiocarpae* (Schedl), (*Cryphalomorphus*),
 Scolytogenes 864
- Plesiophthorus* Schedl (= *Dendroterus*) 950
- plicatus* Wood, *Scolytodes* (= *ragicollis*) 398
- plumalis* Wood, *Pseudothysanoes* 416
- plumbeus* Blandford, *Hylastes* 59
- plumeriae* (Nordlinger), (*Bostrichus*),
 Hypothenemus 938
- plumeriae* Wood, *Scolytodes* 397
- plumericolens* Wood, *Scolytodes* 397
- pluridentatus* Schedl, *Eccoptopterus*
 (= *multispinosus*) 820
- podocarpi* Browne, *Phloeosinus* 251
- podocarpi* Schedl, *Lanurgus* (= *podocarpi*) 409
- podocarpi* Schedl, *Lanurgus* 409
- Pocilips* Schauffuss (= *Coccotrypes*) 591
- poligraphus* (Linnaeus), (*Dermestes*),
 Polygraphus 291
- politus* (Blandford), (*Pityophthorus*), *Araptus* 960
- politus* (Browne), (*Pocilips*), *Coccotrypes* 609
- politus* (Say), (*Bostrichus*), *Xyloterinus* 649
- politus* Eichhoff, *Amphicranus* 1048
- politus* Hagedorn, *Xyleborus* 764
- polonicus* Karpinski, *Dryocortes* (= *autographus*) 573
- polonicus* Karpinski, *Pityophthorus*
 (= *cephalonicae*) 988
- polyalthiae* (Schedl), (*Xyleborus*), *Xyleborinus* 810
- Polygraphi* Chapuis (= *Polygraphini*) 283
- Polygraphidae* Chapuis (= *Polygraphini*) 283
- Polygraphina* Chapuis (= *Polygraphini*) 283
- Polygraphini* Chapuis 283
- Polygraphus* Erichson 283
- polygraphus*, *Polygraphus* (= *poligraphus*) 292

<i>polynesiæ</i> Schedl. <i>Hypocryphalus</i>	871	<i>pourriensis</i> Schedl. <i>Xyleborus</i>	765
<i>polyodon</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701	<i>pouteriæ</i> Wood. <i>Chactophloeus</i>	274
<i>polyphagus</i> (Schedl), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	568	<i>præaltus</i> Bright, <i>Pityophthorus</i> (= <i>deletus</i>)	994
<i>polyphagus</i> Costa Lima, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>crudiæ</i>)	915	<i>præaltus</i> Schedl. <i>Corthylus</i>	1077
<i>polyphagus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>areccæ</i>)	907	<i>præcellens</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	939
<i>polyspringer</i> Israelson, <i>Aphanarthrum</i> (= <i>causceus</i>)	617	<i>præceps</i> LeConte, <i>Scolytus</i>	357
<i>pomacearum</i> Butovitsch, <i>Scolytus</i> (= <i>luevis</i>)	339	<i>præceps</i> Wood, <i>Scolytodes</i>	397
<i>pomelianus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	810	<i>præcursor</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701
<i>pomi</i> Yin & Huang, <i>Scolytus</i>	357	<i>præda</i> (Browne), (<i>Ptilopodius</i>), <i>Scolytogenes</i>	865
<i>porioris</i> Bedel, <i>Scolytus</i> (= <i>rugulosus</i>)	364	<i>præfrictus</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>pini</i>)	520
<i>ponderosæ</i> Blackman, <i>Pityophthorus</i> (= <i>digestus</i>)	995	<i>præmorsus</i> Eichhoff, (<i>Tomicus</i>), <i>Ips</i> (= <i>calligraphus</i>)	496
<i>ponderosæ</i> Hopkins, <i>Conophthorus</i>	964	<i>prænotatus</i> Gredler, (<i>Phloeophthorus</i>), <i>Phloeosinus</i> (= <i>bicolor</i>)	243
<i>ponderosæ</i> Hopkins, <i>Dendroctonus</i>	159	<i>præruptum</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1060
<i>ponderosæ</i> Swaine, <i>Carphoborus</i>	305	<i>præruptus</i> (Eggers), (<i>Pseudothamnurgus</i>), <i>Tiarophorus</i>	540
<i>ponderosæ</i> Swaine, <i>Ips</i> (= <i>calligraphus</i>)	497	<i>præstans</i> Wood, <i>Xyleborus</i>	765
<i>ponderosæ</i> Swaine, <i>Trypodendron</i> (= <i>rufitarsus</i>)	645	<i>præusta</i> (Eggers), (<i>Xyleborus</i>), <i>Schedlia</i>	822
<i>poundanus</i> Eggers, <i>Perniphorus</i>	202	<i>præustum</i> (Eggers), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1061
<i>poundanus</i> Schedl, <i>Glostatus</i>	836	<i>præustus</i> Schedl. <i>Corthylus</i>	1077
<i>popondettæ</i> (Browne), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701	<i>præurius</i> Blandford, <i>Xyleborus</i>	765
<i>populi</i> Blackman, <i>Micracis</i> (= <i>sicainei</i>)	434	<i>prævidui</i> Stark, <i>Hylesinus</i>	83
<i>populi</i> Hopkins, <i>Procryphalus</i> (= <i>mucronatus</i>)	848	<i>præclarum</i> Wood, <i>Monarthrum</i> (= <i>parvum</i>)	1060
<i>populi</i> Hopkins, <i>Trypophloeus</i>	846	<i>premayaensis</i> Murayama, <i>Cryphalus</i>	889
<i>populi</i> Kurenzov, <i>Trypophloeus</i> (= <i>kurenzovi</i>)	846	<i>Prennobilis</i> Eichhoff	651
<i>populi</i> Swaine, <i>Xyleborus</i> (= <i>obesus</i>)	756	<i>Prennophilus</i> Browne (= <i>Prennobilis</i>)	651
<i>porcatus</i> Blandford, <i>Cuesinus</i>	210	<i>preparvus</i> Browne, <i>Cyrtogenius</i>	568
<i>porcatus</i> Chapuis, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	90	<i>priesneri</i> Schedl. <i>Coccotrypes</i>	609
<i>porculus</i> Erichson, <i>Hylastes</i>	60	<i>primitus</i> (Schedl), (<i>Taphroterus</i>), <i>Taphrorychus</i>	560
<i>poricollis</i> (Blandford), (<i>Pityophthorus</i>), <i>Araptus</i>	961	<i>primus</i> (Bright), (<i>Prognathotrichus</i>), <i>Gnathotrichus</i>	1037
<i>porifrons</i> Eggers, <i>Pityogenes</i>	455	<i>primus</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopselion</i>	97
<i>porographus</i> Dejean, (<i>Bostrichus</i>), <i>Ips</i> (= <i>acuminatus</i>)	485	<i>primus</i> Wichmann, <i>Polygraphus</i>	296
<i>porosus</i> (LeConte), (<i>Hylastes</i>), <i>Hylurgops</i>	39	<i>princeps</i> Blandford, <i>Xyleborus</i>	765
<i>porosus</i> Wood, <i>Cryptocarenum</i> (= <i>heveæ</i>)	903	<i>principalis</i> Eichhoff, <i>Xyleborus</i>	765
<i>porrectus</i> Schedl, <i>Microcorthylus</i>	1067	<i>Prionosecles</i> Blandford (= <i>Scolytodes</i>)	357
<i>porteri</i> Brethes, <i>Sinophloeus</i>	105	* <i>priscus</i> Eggers, <i>Xyleborus</i>	765
<i>porteri</i> Brethes, <i>Xylechinus</i>	115	<i>priscus</i> Wood, <i>Chramesus</i>	268
<i>porteri</i> Bruch, <i>Phloeotribus</i>	227	<i>pristis</i> (Wood), (<i>Xyleborus</i>), <i>Theoborus</i>	661
<i>portoricensis</i> Schedl, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>pallidus</i>)	960	<i>privatus</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>bicolor</i>)	687
<i>possyeti</i> Stark, <i>Scolytus</i> (= <i>dakuricus</i>)	329	<i>Problechilidae</i> Eichhoff (= <i>Ctenophorini</i>)	383
<i>posticegraulatus</i> Schedl, <i>Xyleborus</i>	764	<i>Problechilus</i> Eichhoff (= <i>Gymnochilus</i>)	386
<i>posticipilosus</i> Schedl, <i>Xyleborus</i>	764	<i>procer</i> Eichhoff, <i>Xyleborus</i>	765
<i>posticepunctatus</i> Eggers, <i>Thamnurgus</i>	545	<i>procerior</i> Schedl, (<i>Xyleborus</i>), <i>Cyclorhipidion</i> (= <i>terminaliæ</i>)	703
<i>posticespinatum</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	701	<i>procerrimus</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	693
<i>posticestriatus</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800	<i>procerrissimus</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	693
<i>posticoides</i> Schedl, <i>Xyleborus</i>	764	<i>procerus</i> (Erichson), (<i>Phloeotrupes</i>), <i>Phloeoborus</i>	99
<i>posticum</i> Wood, <i>Monarthrum</i>	1060	<i>procerus</i> Bright, <i>Corthylus</i>	1077
<i>posticus</i> Eichhoff, <i>Xyleborus</i>	764	<i>procerus</i> Schedl, <i>Cryphalus</i>	889
* <i>posticus</i> Schedl, <i>Carphoborites</i>	1082	<i>Prochramesus</i> Wood (= <i>Chramesus</i>)	262
<i>potens</i> Schedl, <i>Polygraphus</i>	295	<i>prociduus</i> (Wood), (<i>Loganius</i>), <i>Cnemonyx</i>	317
<i>potens</i> Schedl, <i>Xyleborus</i> (= <i>approximatus</i>)	712	<i>Procryphalus</i> Hopkins	847

- productus* Hagedorn, *Scolytus* (=marginatus) 344
- productus* Hagedorn, *Xyleborus* 765
- profanus* Schedl, *Phloeotribus* 227
- profundus* Schedl, (*Xyleborus*), *Xyleborinus*
(=adamicus) 804
- Progenius* Blandford (=Xyleborus) 704
- Prognathotrichus* Bright (=Gnathotrichus) 1034
- prolatus* Wood, *Hyllocurus* 426
- prolatus* Wood, *Xyleborus* 765
- prolixus* Schedl, *Xyleborus* 766
- prominulus* Wood, *Cnestinus* 210
- pronunciatus* Eggers, *Xyleborus* (=cancellatus) 717
- propinquus* Blandford, *Scolytus* 357
- propinquus* Eichhoff, *Xyleborus* (=pubescens) 767
- propinquus* Nunberg, *Diamerus* 197
- proprium* Wood, *Monarthrum* 1061
- propugnatus* Blandford, *Amphicranus* 1048
- prosorovi* Kurenzov & Kononov, *Dendroctonus*
(=armandi) 135
- prosper* Schedl, (*Stephanoderes*), *Hypothenemus*
(=javanus) 933
- prostratus* Peyerimhoff, *Phloeosinus* (=thujae) 255
- protensus* (Eggers), (*Xyleborus*), *Cnestus* 803
- protii* Browne, *Xyleborus* 766
- protinus* (Wood), (*Xyleborus*), *Xyleborinus* 810
- proticorus* Wood, *Cnemidophorus* 317
- Protopityophthorus* Schedl (=Dendrographus) 561
- Prowebbia* Browne (=Webbia) 829
- proximus* (Niisima), (*Xyloterus*), *Trypodendron* 644
- proximum* Wood, *Monarthrum* 1061
- proximus* (Eggers), (*Cryptalus*), *Coriaceophilus* 854
- proximus* (Eichhoff), (*Tomicus*), *Orthotomicus* 473
- proximus* Blandford, *Polygraphus* 296
- proximus* Chapuis, *Scolytus* 357
- proximus* Eggers, *Xyleborus* (=affinis) 709
- proximus* Wood, *Scolytodes* 398
- pruinosulus* Browne, *Xyleborus* 766
- pruinotum* (Blandford), (*Xyleborus*),
Cyclorhipidion 701
- pruinotus* (Blackman), (*Renocis*), *Chaetophloeus* 274
- pruinotus* (Eichhoff), (*Pityophthorus*),
Pseudopityophthorus 974
- pruinotus* Eichhoff, *Hylesinus* 83
- pruni* (Wood), (*Phacrylus*), *Acorthylus* 853
- pruni* Eggers, *Cryptalus* 859
- pruni* Hopkins, *Hypothenemus* (=eruditus) 922
- pruni* Ratzeburg, (*Eccoptogaster*), *Scolytus* (=mali) 342
- pruni* Wood, *Liparthrum* 279
- pruni* Wood, *Phloeotribus* 227
- Pruniphagus* Murayama (=Xylechinus) 114
- prutenskyi* Sokanovskii, *Hylesinus* 83
- psaltes* (Hagedorn), (*Bothryperus*), *Phrixosoma* 190
- psaltes* (Schedl), (*Xyleborus*), *Cyclorhipidion* 702
- Pseudoacacis* Schedl (=Acacis) 193
- pseudoacuminatus* (Schedl), (*Hexacolus*),
Scolytodes 398
- pseudoambasius* Schedl, *Xyleborus* 766
- pseudoangustatus* Schedl, (*Xyleborus*), *Xyleborinus*
(=saxeseni) 815
- pseudobarbatus* Schedl, (*Xyleborus*), *Eucallacea*
(=destruens) 688
- pseudobicolor* Eggers, (*Hexacolus*), *Scolytodes*
(=notatus) 396
- pseudobrasiliensis* Eggers, (*Xyleborus*), *Ambrosiodmus*
(=obliquus) 678
- pseudobrunneus* Schedl, *Polygraphus* 296
- pseudocaudatus* Eggers, *Scolytus* (=caudatus) 327
- pseudochinlingensis* Tsai & Li, *Cryptalus* 859
- Pseudochramesus* Blackman 261
- pseudocitri* (Schedl), (*Xyleborus*), *Ambrosiodmus* 679
- pseudococcytripes* Eggers, (*Xyleborus*),
Theoborus (=theobromae) 661
- pseudocolossus* (Schedl), (*Xyleborus*),
Ambrosiodmus 679
- pseudocomans* (Eggers), (*Xyleborus*),
Hadrodemus 819
- Pseudocorthylus* Ferrari (=Corthylus) 1069
- Pseudocosmoderes* Nobuchi (=Cosmoderes) 900
- pseudocostellatus* Schedl, *Scolytus* 357
- pseudocristatus* (Pfeffer), (*Phloeophthorus*),
Phloeotribus 227
- * *pseudocrucifer* (Schedl), (*Xyleborus*),
Cyclorhipidion 702
- Pseudocryptalus* Ferrari (=Cryptalus) 872
- Pseudocryptalus* Swaine (=Chaetophloeus) 271
- Pseudocrypturgus* Eggers (=Microborus) 383
- pseudocylindricus* Eggers, *Xyleborus* 766
- Pseudodiamerus* Eggers 194
- pseudoforsteri* (Schedl), (*Xyleborus*),
Cyclorhipidion 702
- pseudogracilis* Schedl, (*Xyleborus*), *Xyleborinus*
(=saxeseni) 815
- Pseudohylesinus* Eggers (=Hylesinopsis) 92
- Pseudohylesinus* Swaine 109
- Pseudohyllocurus* Nunberg (=Lanurgus) 408
- Pseudohyorrhynchus* Murayama 192
- pseudohystrix* (Schedl), (*Xyleborus*), *Premnobius* 654
- pseudompar* Schedl, *Hyllocurus* 427
- pseudomajor* Schedl, *Xyleborus* 766
- Pseudomicracis* Blackman (=Micracisella) 429
- Pseudomicracis* Eggers 429
- pseudomovellae* Nunberg, *Strombophorus*
(=vittatus) 204
- Pseudophloeotribus* Eggers (=Hylesinopsis) 92
- pseudopiceus* Wood, *Scolytodes* 398
- pseudopilifer* Schedl, *Xyleborus* 766
- pseudopityogenes* (Eggers), (*Xyleborus*),
Xyleborinus 810
- Pseudopityophthorus* Swaine 971
- Pseudopoclipus* Murayama (=Taphrorhynchus) 557
- Pseudopolygraphus* Seither (=Polygraphus) 254
- pseudoprocer* Schedl, *Xyleborus* (=declivis) 723
- pseudopunctulus* (Schedl), (*Xyleborus*), *Cnestus* 803
- pseudorudis* Schedl, *Xyleborus* 766
- pseudoscurbicollis* Atkinson, *Phloeotribus* 227

<i>pseudoscutellare</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1061	<i>pubescens</i> Wood, <i>Cryptocarenum</i>	903
<i>pseudosolidus</i> (Schedl), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800	<i>pubescens</i> Wood, <i>Scolytodes</i>	398
<i>pseudosolitaris</i> (Eggers), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658	<i>pubescens</i> Zimmermann, <i>Xyleborus</i>	766
<i>pseudosolitaris</i> <i>schizolobius</i> Schedl, (<i>Xyleborus</i>), <i>Dryocoetoides</i> (= <i>pseudosolitaris</i>)	658	<i>pubifer</i> (Schedl), (<i>Xyleborus</i>), <i>Hadrodemius</i>	819
<i>pseudosulcatum</i> Schedl, (<i>Pterocyclon</i>), <i>Monarthrum</i> (= <i>consimile</i>)	1052	<i>pubifer</i> Schedl, <i>Xyloctonus</i>	840
<i>pseudosuturalis</i> Schedl, <i>Cnestus</i>	803	<i>pubifrons</i> (Guillebeau), (<i>Phloeophthorus</i>), <i>Phlocotribus</i>	227
<i>pseudotabulaeformis</i> Tsai & Li, <i>Cryphalus</i>	889	<i>pubifrons</i> Guillebeau, <i>Phlocotribus</i> (= <i>cristatus</i>)	220
<i>pseudotenuis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptoborus</i>	664	<i>pubifrons</i> Bright, <i>Pityophthorus</i>	1023
<i>pseudothammurgus</i> Eggers (= <i>Tiarophorus</i>)	539	<i>pubipennis</i> (Blandford), (<i>Trypodendron</i>), <i>Indocryphalus</i>	648
<i>Pseudothyasnoes</i> Blackman	411	<i>pubipennis</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	939
<i>pseudotsugae</i> Chamberlin, <i>Phloeosinus</i> (= <i>autematus</i>)	240	<i>pubipennis</i> (LeConte), (<i>Bostrichus</i>), <i>Pseudopityophthorus</i>	975
<i>pseudotsugae</i> Hopkins, <i>Dendroctonus</i>	168	<i>pubipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	826
<i>pseudotsugae</i> Swaine, <i>Dryocoetes</i> (= <i>autographus</i>)	578	<i>pubens</i> (Blackman), (<i>Pityophthoroides</i>), <i>Pityophthorus</i>	1023
<i>pseudotsugae</i> Swaine, <i>Pityophthorus</i>	1021	<i>pubicus</i> Blackman, <i>Pityophthorus</i>	1023
<i>pseudotsugae</i> Wood, <i>Carphoborus</i>	309	<i>puellus</i> (Reitter), (<i>Myelophilus</i>), <i>Tomicus</i>	136
<i>pseudotalidus</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>interjectus</i>)	691	<i>puellus</i> <i>orientalis</i> Krivolutsкая, (<i>Blastophagus</i>), <i>Tomicus</i> (= <i>puellus</i>)	136
<i>Pseudowebbia</i> Browne (= <i>Webbia</i>)	829	<i>puer</i> (Eggers), (<i>Xyleborus</i>), <i>Leptoxyleborus</i>	660
<i>Pseudoxyleborus</i> Eggers (= <i>Amasa</i>)	681	<i>puer</i> (Schedl), (<i>Hexacolus</i>), <i>Scolytodes</i>	398
<i>Pseudoxylechinus</i> Wood & Huang	114	<i>puerarae</i> Choo & Woo, <i>Scolytogenes</i>	865
<i>psidii</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>puerulus</i> Schedl, <i>Microcorthyus</i>	1067
<i>Pteleobius</i> Bedel	69	<i>pulchellus</i> (Murayama), (<i>Xyloterus</i>), <i>Trypodendron</i>	644
<i>Pterocyclon</i> Eichhoff (= <i>Monarthrum</i>)	1050	<i>pulchellus</i> (Nobuchi), (<i>Taenioglyptes</i>), <i>Cryphalus</i>	890
<i>Pterocyclonoides</i> Schedl (= <i>Tricolus</i>)	1042	<i>pulchellus</i> Blandford, <i>Phloeosinus</i>	251
<i>pteryodphytae</i> (Schedl), (<i>Pocilips</i>), <i>Coccotrypes</i>	609	<i>pulchellus</i> Eichhoff, <i>Pityophthorus</i>	1023
<i>pterygosperma</i> Schedl, <i>Hypothenemus</i> (= <i>eruditus</i>)	921	<i>pulchripes</i> Schedl, <i>Xyleborus</i> (= <i>perlongus</i>)	762
<i>Ptilopodius</i> Hopkins	899	<i>pulcerrimus</i> Schedl, <i>Xyleborus</i> (= <i>perlongus</i>)	762
<i>ptyocerus</i> Blandford, <i>Corthyus</i>	1077	<i>pulicarius</i> (Zimmermann), (<i>Crypturgus</i>), <i>Pityophthorus</i>	1024
<i>puberulus</i> (Blandford), (<i>Xyleborus</i>), <i>Arixyleborus</i>	668	<i>pulla</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	826
<i>puberulus</i> (Chapuis), (<i>Dryotomus</i>), <i>Phlocotribus</i>	227	<i>pullatus</i> Blackman, <i>Pseudohylesinus</i> (= <i>dispar</i>)	109
<i>puberulus</i> (LeConte), (<i>Cryphalus</i>), <i>Pityophthorus</i>	1021	<i>pullus</i> (Zimmermann), (<i>Crypturgus</i>), <i>Pityophthorus</i>	1024
<i>puberulus</i> LeConte, <i>Phlocotribus</i> (= <i>lecontei</i>)	223	<i>pullus</i> Blandford, <i>Cnesinus</i>	211
<i>puberulus</i> Schedl, <i>Cryphalus</i>	890	<i>pullus</i> Wood, <i>Periocryphalus</i>	948
<i>pubescens</i> (Marshall), (<i>Ips</i>), <i>Pityophthorus</i>	1022	<i>pulvereus</i> Blackman, <i>Pseudopityophthorus</i> (= <i>pruinosis</i>)	975
<i>pubescens</i> (Schedl), (<i>Neodryocoetes</i>), <i>Araptus</i>	961	<i>pulverulentus</i> Eichhoff, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	941
<i>pubescens</i> Blackman, <i>Pseudopityophthorus</i>	975	<i>pulverulentus</i> Gerstaecker, <i>Diamerus</i>	197
<i>pubescens</i> Eggers, <i>Scolytus</i> (= <i>pubescens</i> Stark)	358	<i>pumilio</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1061
<i>pubescens</i> Fabricius, (<i>Anthrabus</i>), <i>Hylesinus</i> (= <i>rarius</i>)	86	<i>pumilio</i> Eggers, <i>Tricolus</i>	1044
<i>pubescens</i> Fabricius, (<i>Bostrichus</i>), <i>Polygraphus</i> (= <i>poligraphus</i>)	294	<i>pumilio</i> Eichhoff, <i>Dryocoetes</i>	584
<i>pubescens</i> Hagedorn, <i>Scolytotplatypus</i>	406	<i>pumilionides</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865
<i>pubescens</i> Hopkins, <i>Cryphalus</i>	890	<i>pumilus</i> (Chapuis), (<i>Rhopalopleurus</i>), <i>Chramesus</i>	268
<i>pubescens</i> Hopkins, <i>Hypothenemus</i>	939	<i>pumilus</i> (Mannerheim), (<i>Hylastes</i>), <i>Dohurgus</i>	615
<i>pubescens</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>schedli</i>)	940	<i>pumilus</i> (Wood), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865
<i>pubescens</i> Schedl, <i>Coccotrypes</i>	609	<i>pumilus</i> (Wood), (<i>Cryptulocleptus</i>), <i>Pseudothyasnoes</i>	416
<i>pubescens</i> Schedl, <i>Traglostus</i>	410	<i>pumilus</i> Eggers, (<i>Loganius</i>), <i>Cnemidophorus</i> (= <i>exiguus</i>)	315
<i>pubescens</i> Stark, <i>Scolytus</i>	357		
<i>pubescens</i> Swaine, <i>Dryocoetes</i> (= <i>affaber</i>)	571		
<i>pubescens</i> Swaine, <i>Sciurus</i>	29		

- pumilus* Eggers, *Cnesinus* 211
- pumilus* Eggers, *Xyleborus* 767
- pumilus* Wood, *Corthylus* 1077
- pumilus* Wood, *Dendrocraniulus* 553
- pumilus* Wood, *Microcorthylus* 1067
- pumilus* Wood, *Phloeosinusopsioides* 239
- pumilus* Wood, *Scolytodes* 395
- punctatissimus* (Eichhoff), (*Xyleborus*),
 Leptoxyleborus 660
- punctatissimus* (Schedl), (*Neodryocoetes*),
 Araptus 961
- punctatissimus* (Zimmermann), (*Crypturgus*),
 Corthylus 1077
- punctatissimus* Eggers, *Thamniurgus* 545
- punctatopilosum* (Schedl), (*Xyleborus*),
 Cyclorhipidion 702
- punctatorugosus* (Schedl), (*Micracis*), *Hylocurus* 427
- punctatorugosus* Chapuis, *Phlocoborus* 99
- punctatostratus* Schedl, *Cryphalus* 890
- punctatulus* Nobuchi, *Scolytogenes* 865
- punctatum* (Eggers), (*Xyleborus*),
 (*Cyclorhipidion*) 702
- punctatus* (Eggers), (*Thamniurgides*), *Coccotrypes* 609
- punctatus* (Eggers), (*Kissophagus*), *Hylesinopsis* 95
- punctatus* (Eggers), (*Hylocurosoma*), *Scolytodes* 395
- punctatus* (Schedl), (*Cryphalomorphus*),
 Scolytogenes 865
- punctatus* (Schedl), (*Phlocotrupes*), *Phlocoborus* 100
- punctatus* Blandford, *Cnesinus* 211
- punctatus* Browne, *Cryphalus* 890
- punctatus* Browne, *Xylocleptes* (= *brownei*) 549
- punctatus* Eggers, (*Pociclipis*), *Coccotrypes*
 (= *salakensis*) 611
- punctatus* Eggers, (*Stephanoderes*), *Hypothenemus*
 (= *hampei*) 930
- punctatus* Eggers, *Corthylus* 1078
- punctatus* Eggers, *Pagiocerus* 214
- punctatus* Eggers, *Pityophthorus* 1025
- punctatus* Hopkins, (*Xylocleptes*), *Dendrocraniulus*
 (= *eucurbitae*) 551
- punctatus* LeConte, *Dendroctonus* 173
- punctatus* LeConte, *Phloeosinus* 251
- punctatus* Muller, *Scolytus* (= *scolytus*) 374
- punctatus* Nobuchi, (*Cryphalomorphus*),
 Scolytogenes (= *punctatulus*) 865
- punctatus* Ratzeburg, (*Eccoptogaster*), *Scolytus*
 (= *rugulosus*) 369
- punctatus* Wood, *Chramesus* 268
- punctatus* Wood, *Phloeocleptus* 422
- puncticollis* (Schedl), (*Lepicrinus*), *Scolytogenes* 865
- puncticollis* Blandford, *Dendrosinus* 237
- puncticollis* Blandford, *Scolytopsis* 321
- puncticollis* Chapuis, *Phloeotribus* (= *setulosus*) 232
- puncticollis* Eggers, *Diamerus* 197
- puncticollis* Hopkins, (*Stephanoderes*),
 Hypothenemus (= *erectus*) 918
- puncticollis* LeConte, *Pityophthorus* (= *nitidulus*) 1015
- punctifer* Wood, *Scolytodes* (= *cecropiavorus*) 390
- punctifer* Wood, *Scolytodes* (= *punctiferus*) 395
- punctiferus* Wood, *Scolytodes* 395
- punctifrons* (Blandford), (*Pterocyclon*),
 Monarthrum 1061
- punctifrons* Bright, *Pityophthorus* 1025
- punctifrons* Hopkins, *Hypothenemus* (= *eruditus*) 923
- punctifrons* Schedl, *Cryptocarenum* 903
- punctifrons* Thomson, *Polygraphus* (= *poligraphus*) 295
- punctiger* (Schedl), (*Neodryocoetes*),
 Pityophthorus 1025
- punctilicolle* (Schedl), (*Xyleborus*),
 Cyclorhipidion 702
- punctipennis* (Schedl), (*Xyleborus*), *Coptodryas* 826
- punctipennis* Eggers, (*Thamniurgides*), *Coccotrypes*
 (= *rhizophorae*) 610
- punctipennis* Eggers, *Xyloctonus* 840
- punctipennis* Hopkins, *Hypothenemus* (= *eruditus*) 923
- punctipennis* Hopkins, *Trypophloeus* (= *striatulus*) 847
- punctipennis* LeConte, (*Xyleborus*), *Orthotomicus*
 (= *caelatus*) 469
- punctipennis* Schedl, *Cryphalus* 890
- punctipennis* Schedl, *Microcryphalus* 411
- punctulatus* Eggers, *Coccotrypes* (= *carpophagus*) 596
- punctulatus* Eggers, *Cryphalus* (= *kurenzovi*) 852
- punctulatus* Kurenzov, *Xyleborus* 767
- punctulatus* Schedl, (*Xyleborus*), *Cnestus* (= *nudus*) 803
- pupillatus* Eggers, *Hylastes* (= *brunneus*) 50
- pupillatus* Eggers, *Phlocoborus* (= *bodei*) 98
- pusillimus* Schedl, *Cryphalus* 890
- pusillimus* Wood, *Scolytodes* 395
- pusillus* (Eggers), (*Hylocurosoma*), *Scolytodes* 395
- pusillus* (Eggers), (*Neoxyloctonus*), *Scolytomimus* 839
- pusillus* (Eggers), (*Webbia*), *Arxyleborus* 668
- pusillus* (Gerstaecker), (*Hylesinus*), *Rhopalopselion* 97
- pusillus* (Gyllenhal), (*Bostriehus*), *Crypturgus* 625
- pusillus* (Schedl), (*Xyleborus*), *Xylosandrus* 800
- pusillus* Blackman, *Hylastes* (= *tenuis*) 63
- pusillus* Eggers, *Corthylus* 1078
- pusillus* Eggers, *Dryocoetes* 554
- pusillus* Eggers, *Hypothenemus* (= *eruditus*) 923
- pusillus* Eggers, *Scolytoplatus* 406
- pusillus* Eggers, *Strombophorus* 204
- pusillus* Harris, (*Tomicus*), *Pseudopityophthorus*
 (= *minutissimus*) 974
- pusillus* Schedl, *Cnesinus* 211
- pusillus* Schedl, *Cryphalus* 890
- pusillus* Wollaston, (*Aphanarthrum*), *Cisurgus*
 (= *wollastoni*) 630
- pusillus* Wood, *Microcorthylus* 1067
- pusillus* Wood, *Pityophthorus* (= *attenuatus*) 982
- pusio* Eggers, *Xyleborus* 767
- pusio* LeConte, *Pityophthorus* (= *pulchellus*) 1024
- pustulatus* Schedl, *Gnathotrupes* 1042
- putonii* Eichhoff, (*Hylesinus*), *Pteleobius* (= *kraatzii*) 70
- putputensis* Browne, *Xyleborus* 767
- Pycnarthrum* Eichhoff 384
- pygeumi* Schedl, *Sphaerotrypes* 200

<i>pygmaeolus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865	<i>quadrioculata</i> (Eggers), (<i>Neohylesinus</i>), <i>Phrixosoma</i>	190
<i>pygmaeolus</i> Schedl, <i>Pityophthorus</i>	1025	<i>quadrioculatus</i> (Hagedorn), (<i>Spongotarsus</i>), <i>Polygraphus</i>	296
<i>Pygmaeoscolytus</i> Butovitsch (= <i>Scolytus</i>)	322	<i>quadrioculatus</i> Browne, <i>Scolytomimus</i> (= <i>assamensis</i>)	839
<i>pygmaeum</i> Eggers, <i>Ctenoxylon</i>	838	<i>quadriporus</i> Beaver, <i>Cryptoxyleborus</i>	829
<i>pygmaeum</i> Wollaston, <i>Aphanarthrum</i>	619	<i>quadriscignatus</i> Schedl, <i>Xyleborus</i>	768
<i>pygmaeum laticollis</i> Wollaston, <i>Aphanarthrum</i> (= <i>pygmaeum</i>)	619	<i>quadriscopinatus</i> Schedl, <i>Pityophthorus</i>	1025
<i>pygmaeus</i> (Eggers), (<i>Pelicerus</i>), <i>Dendrographus</i>	561	<i>quadriscopinis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	810
<i>pygmaeus</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800	<i>quadriscopinosulus</i> Eggers, <i>Xyleborus</i>	768
<i>pygmaeus</i> (Fabricius), (<i>Bostrichus</i>), <i>Scolytus</i>	358	<i>quadriscopinosus</i> (Eggers), (<i>Xylocleptes</i>), <i>Acanthotomicus</i>	483
<i>pygmaeus</i> Eggers, <i>Polygraphus</i> (= <i>coronatus</i>)	286	<i>quadriscopinosus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xyleborinus</i>	810
<i>pygmaeus</i> Eichhoff, <i>Coccotrypes</i> (= <i>carpophagus</i>)	595	<i>quadriscopinosus</i> Blackman, <i>Hylocurus</i>	427
<i>pygmaeus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	923	<i>quadriscopinosus</i> Eggers, <i>Sampsonius</i>	656
<i>pygmaeus</i> Schedl, <i>Cryptocarenum</i>	903	<i>quadriscopinosus</i> Say, <i>Scolytus</i>	359
<i>pygmaeus</i> Schedl, <i>Lanurgus</i>	409	<i>quadriscopinosus</i> Schedl, <i>Prennobiis</i>	654
<i>pygmaeus</i> Schedl, <i>Micracis</i> (= <i>sicainei</i>)	434	<i>quadrisculatus</i> Strohmeier, <i>Dryocoetes</i>	584
<i>pygmaeus</i> Schedl, <i>Pityophthorus</i> (= <i>opaculus</i>)	1018	<i>quadrituberculatus</i> (Schedl), (<i>Isophthorus</i>), <i>Acanthotomicus</i>	483
<i>pygmaeus</i> Wood, <i>Corthylus</i>	1078	<i>quadrituberculatus</i> (Schedl), (<i>Metahylesinus</i>), <i>Hylesinopsis</i>	95
<i>pygmaeus</i> Wood, <i>Stegomerus</i>	852	<i>quadrituberculatus</i> Sampson, <i>Sphaerotrypes</i>	200
<i>pyri</i> Peck, (<i>Scolytus</i>), <i>Xyleborus</i> (= <i>dispar</i>)	729	<i>quadrituberculatus</i> Schedl, (<i>Gnathotrichus</i>), <i>Gnathotripes</i> (= <i>longipennis</i>)	1041
<i>pyri</i> Ratzeburg, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>mali</i>)	343	<i>quadrituberculatus</i> Schedl, <i>Margadillius</i>	899
<i>pyri</i> Tsai & Yin, <i>Sphaerotrypes</i>	200	<i>quaesitus</i> Schedl, <i>Cnesinus</i>	211
<i>quadraticolle</i> Eichhoff, <i>Pycnarthrum</i> (= <i>hispidum</i>)	385	<i>quasimodo</i> Browne, <i>Xyleborus</i>	768
<i>quadraticollis</i> (Eggers), (<i>Xyleborus</i>), <i>Eucallacea</i>	693	<i>quattuordecimcostatus</i> Schedl, <i>Webbia</i> (= <i>quattuordecimspinatus</i>)	832
<i>quadraticollis</i> Chapuis, <i>Dactylipalpus</i> (= <i>transversus</i>)	102	<i>quattuordecimspinatus</i> Sampson, <i>Webbia</i>	832
<i>quadratus</i> Blandford, <i>Xyleborus</i>	767	<i>quattuordecimspinatus</i> Schedl, <i>Webbia</i> (= <i>quattuordecimspinatus</i> Sampson)	832
<i>quadrinictus</i> Schedl, <i>Webbia</i>	832	<i>queenslandi</i> (Schedl), (<i>Erioschidias</i>), <i>Cosmoderes</i>	902
<i>quadrinictus</i> Schedl, <i>Xyloctonus</i>	841	<i>queenslandi</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	609
<i>quadriscostata</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	826	<i>queenslandi</i> Schedl, <i>Peridryocoetes</i>	562
<i>quadricuspe</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	702	<i>quercki</i> Stebbing, <i>Sphaerotrypes</i>	201
<i>quadridens</i> (Browne), (<i>Hypothenemus</i>), <i>Scolytogenes</i>	865	<i>quercki</i> Wood, <i>Polygraphus</i> (= <i>kaimochii</i>)	289
<i>quadridens</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1061	<i>quercicola</i> Eggers, <i>Xyleborus</i>	768
<i>quadridens</i> (Hartig), (<i>Bostrichus</i>), <i>Pityogenes</i>	455	<i>quercicoleus</i> Wood, <i>Monarthrum</i>	1061
<i>quadridens</i> (Schedl), (<i>Cryphalops</i>), <i>Erioporus</i>	855	<i>quercicolens</i> Wood, <i>Pseudothysanocis</i>	416
<i>quadridens</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	483	<i>quercinus</i> (Wood), (<i>Cryptocleptes</i>), <i>Pseudothysanocis</i>	416
<i>quadridens</i> Blackman, <i>Camptocerus</i>	320	<i>quercinus</i> Wood, <i>Phloeotribus</i>	228
<i>quadridens</i> Eggers, <i>Prennobiis</i>	654	<i>quercinus</i> Wood, <i>Pityophthorus</i>	1025
<i>quadridens</i> Eggers, <i>Xyleborus</i> (= <i>bidentatus</i>)	715	<i>querciperda</i> Schwarz, (<i>Pityophthorus</i>), <i>Pseudopityophthorus</i> (= <i>pruinosis</i>)	975
<i>quadrideus</i> Pjatnitskii, <i>Orthotomicus</i> (= <i>golorjankoi</i>)	469	<i>quercivorum</i> Schedl, <i>Monarthrum</i>	1061
<i>quadridens</i> Schedl, <i>Xyloctonus</i>	840	<i>quercum</i> (Wood), (<i>Amphicranus</i>), <i>Monarthrum</i>	1061
<i>quadridens</i> Wood, <i>Chramesus</i>	268	<i>quercus</i> Eichhoff, (<i>Xyloterus</i>), <i>Trypodendron</i> (= <i>signatum</i>)	647
<i>quadridens</i> Wood, <i>Scolytomimus</i>	840	<i>quercus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>rotundicollis</i>)	940
<i>quadridentatum</i> (Eggers), (<i>Anchonocerus</i>), <i>Monarthrum</i>	1061	<i>quercus</i> Hopkins, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxseni</i>)	815
<i>quadridentatus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>interstitialis</i>)	932		
<i>quadridentatus</i> Schedl, (<i>Xyleborus</i>), <i>Prennobiis</i> (= <i>quadridens</i>)	654		
<i>quadridentatus</i> Sturm, (<i>Bostrichus</i>), <i>Ips</i> (= <i>acuminatus</i>)	485		
<i>quadrimaculatus</i> Schedl, <i>Amphicranus</i>	1048		

<i>quercus</i> Kurenzov, <i>Xyleborus</i> (=starki)	776	<i>regius</i> Hagedorn, <i>Ozopemon</i>	590
<i>querneum</i> Wood, <i>Monarthrum</i>	1061	<i>regularis</i> (Schedl), (<i>Pocilips</i>), <i>Coccotrypes</i>	609
<i>querneus</i> Wood, <i>Pseudothysanoes</i>	416	<i>regularis</i> Blackman, <i>Pityophthorus</i>	1026
<i>quinquedincatum</i> Adams, (<i>Bostrichus</i>), <i>Trypodendron</i> (=signatum)	647	<i>° rehi</i> (Hagedorn), (<i>Phloeosinus</i>), <i>Phloeosinus</i>	252
<i>radiatae</i> Hopkins, <i>Conophthorus</i>	966	<i>reinoseri</i> Schedl, <i>Pycnarthrum</i> (=hispidum)	385
<i>radiatae</i> Hopkins, (<i>Tomicus</i>), <i>Ips</i> (=mexicanus)	513	<i>reitteri</i> Eichhoff, <i>Gymnochilus</i>	357
<i>radiatae</i> Swaine, <i>Carphoborus</i>	309	<i>remorsus</i> Wood, <i>Phlocotribus</i>	228
<i>radiatus</i> Wood, <i>Scolytodes</i>	398	<i>Renocis</i> Casey (=Chaetophloeus)	271
<i>radulosus</i> Blandford, <i>Phloeoborus</i> (=cristatus)	95	<i>repositus</i> Schedl, <i>Xyleborus</i>	768
<i>ragusae</i> Reitter, <i>Cisurgus</i>	630	<i>resecans</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	684
<i>raja</i> Blandford, <i>Scolytplatypus</i>	406	<i>resectus</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	684
<i>rallus</i> (Wood), (<i>Gnathophthorus</i>), <i>Dacnophthorus</i>	969	<i>resiniferae</i> Peyerimhoff, <i>Cisurgus</i>	630
<i>rameus</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacca</i> (=bicolor)	657	<i>resiniferi</i> Schedl, <i>Cryphalus</i>	890
<i>rameus</i> Wood, <i>Amphicranus</i>	1048	<i>resinosae</i> Hopkins, <i>Conophthorus</i>	967
<i>ramicola</i> Reitter, <i>Dryocoetes</i>	595	<i>° resinosus</i> Schedl, <i>Xyleborus</i>	768
<i>ramipera</i> Swaine, <i>Pityophthorus</i>	1025	<i>resolutus</i> Wood, <i>Dendroterus</i>	951
<i>ramosus</i> Beeson, <i>Ptilopodius</i>	900	<i>restrictus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	679
<i>ramosus</i> Bright, <i>Pityoborus</i> (=secundus)	969	<i>retamae</i> Perris, (<i>Hylesinus</i>), <i>Phlocotribus</i> (=rhododactylus)	229
<i>ramulorum</i> Perris, <i>Pityophthorus</i> (=pubescens)	1022	<i>reticulatum</i> Schedl, <i>Pycnarthrum</i>	355
<i>ramulorum</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (=cylindrotomicus)	793	<i>reticulatum</i> Wood, <i>Aphanarthrum</i>	619
<i>ranasinghei</i> Wood, <i>Sphaerotrypes</i>	201	<i>reticulatus</i> (Wood), (<i>Hexacolus</i>), <i>Scolytodes</i>	398
<i>rapandus</i> Schedl, <i>Xyleborus</i>	768	<i>reticulatus</i> Bright, <i>Sampsonius</i>	656
<i>rapanus</i> Beeson, <i>Xyleborus</i> (=adamsoni)	705	<i>reticulatus</i> Chapuis, <i>Cnesinus</i>	211
<i>rasilis</i> Schedl, <i>Amphicranus</i>	1048	<i>reticulatus</i> Wood, <i>Hylurgops</i>	40
<i>ratamae</i> Blackman, <i>Thysanoes</i> (=texanus)	421	<i>reticulus</i> Wood, <i>Cnesinus</i>	211
<i>ratzeburgi</i> Ferrari, (<i>Cryphalus</i>), <i>Ernoporius</i> (=tiliae)	557	<i>retifer</i> Wood, <i>Cnesinus</i>	211
<i>ratzeburgi</i> Janson, <i>Scolytus</i>	361	<i>retifer</i> Wood, <i>Hylastes</i>	61
<i>ratzeburgi</i> Kolenati, (<i>Bostrichus</i>), <i>Xyleborus</i> (=dispar)	729	<i>retifer</i> Wood, <i>Scolytodes</i>	398
<i>raucus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	679	<i>reticentriculus</i> Tsai & Yin, <i>Polygraphus</i> (=rudis)	297
<i>reburrus</i> Bright, <i>Corthylus</i> (=uniseptis)	1081	<i>retrusus</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (=saxeseni)	815
<i>recarus</i> Wood, <i>Cnemonyx</i>	317	<i>retusicollis</i> Zimmermann, <i>Xyleborus</i> (=ferrugineus)	738
<i>recarus</i> Wood, <i>Pseudothysanoes</i>	416	<i>retusifer</i> Wood, <i>Corthylus</i>	1078
<i>recens</i> Bright, <i>Pityophthorus</i>	1026	<i>retusiformis</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (=ater)	787
<i>recidens</i> (Sampson), (<i>Xyleborus</i>), <i>Coptodryas</i>	827	<i>retusipennis</i> Blandford, <i>Glochinnocerus</i>	1064
<i>reconditus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	810	<i>retusipennis</i> Blandford, <i>Hyllocurus</i>	427
<i>rectangulus</i> Ferrari, (<i>Tomicus</i>), <i>Ips</i> (=erosus)	505	<i>retusum</i> (LeConte), (<i>Xyloterus</i>), <i>Trypodendron</i>	644
<i>recticollis</i> (Schedl), (<i>Pocilips</i>), <i>Coccotrypes</i>	609	<i>retusus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800
<i>rectus</i> LeConte, (<i>Tomicus</i>), <i>Ips</i> (=pini)	521	<i>retusus</i> (LeConte), (<i>Cryphalus</i>), <i>Gnathotrichus</i>	1037
<i>rectus</i> Schedl, <i>Hyllocurus</i> (=brasiliensis)	423	<i>retusus</i> Eichhoff, <i>Amphicranus</i>	1048
<i>rectus</i> Wood, <i>Camptocerus</i>	320	<i>retusus</i> Olivier, (<i>Scolytus</i>), <i>Xylocleptes</i> (=bispinus)	548
<i>rectus</i> Wood, <i>Scolytominus</i>	840	<i>retusus</i> Wood, <i>Corthylus</i>	1078
<i>recurtus</i> Browne, (<i>Streptocranus</i>), <i>Coptoborus</i> (=bicuspis)	662	<i>reunionis</i> Schedl, <i>Xyleborus</i>	768
<i>redikorzeri</i> Berger, <i>Cryphalus</i>	890	<i>revocabile</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	702
<i>reditus</i> Wood, <i>Dendrocranulus</i>	553	<i>rhizophagus</i> Thomas & Bright, <i>Dendroctonus</i>	173
<i>redtenbacheri</i> Ferrari, <i>Corthylus</i>	1078	<i>rhizophorae</i> (Hopkins), (<i>Spermatoplex</i>), <i>Coccotrypes</i>	609
<i>refertus</i> Wood, <i>Araptus</i>	961	<i>rhizophorae</i> Eggers, (<i>Dendrugus</i>), <i>Coccotrypes</i> (=rhizophorae)	609
<i>reflexus</i> Blackman, <i>Scolytus</i>	364	<i>rhodesianus</i> (Eggers), (<i>Pseudophlocotribus</i>), <i>Hylesinopsis</i>	95
<i>reflexus</i> Browne, <i>Hypoerythralus</i>	571	<i>rhodesianus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	679
<i>regalis</i> (Wood), (<i>Olonthogaster</i>), <i>Hyleidius</i>	261	<i>rhododactylus</i> (Marshall), (<i>Ips</i>), <i>Phlocotribus</i>	228
<i>° regimontanus</i> (Hagedorn), (<i>Phloeosinites</i>), <i>Phloeosinus</i>	252		
<i>regius</i> (Schedl), (<i>Lepersinus</i>), <i>Hylesinus</i>	83		

<i>rhododactylus</i> Ratzeburg, (<i>Hylesinus</i>), <i>Phlocotribus</i> (= <i>spinulosus</i>)	234	<i>robustus</i> Schedl. <i>Corthylus</i>	1079
<i>rhois</i> Blackman, <i>Cactopinus</i>	436	<i>robustus</i> Schedl. <i>Dendrochilus</i>	438
<i>rhois</i> Blackman, <i>Stenoclyptus</i> (= <i>sulcatus</i>)	419	<i>robustus</i> Schedl. <i>Hylocurus</i>	426
<i>rhois</i> Swaine, <i>Pityophthorus</i> (= <i>lautus</i>)	1006	<i>robustus</i> Schedl. <i>Micracis</i> (= <i>sucainei</i>)	434
<i>rhois acerini</i> Blackman, <i>Pityophthorus</i> (= <i>latus</i>)	1006	<i>robustus</i> Schedl. (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>funereus</i>)	674
<i>rhois hamamelidis</i> Blackman, <i>Pityophthorus</i> (= <i>lautus</i>)	1006	<i>rodgeri</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>bicolor</i>)	687
<i>rhois sucainei</i> Blackman, <i>Pityophthorus</i> (= <i>lautus</i>)	1006	<i>rodgeri priratus</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>bicolor</i>)	687
<i>Rhopalocryphus</i> Nunberg (= <i>Glostatus</i>)	835	<i>roeri</i> Schedl. <i>Xylechinus</i>	118
<i>Rhopalopleurus</i> Chapuis (= <i>Chramesus</i>)	262	<i>rollinae</i> Hopkins, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>Rhopalopselion</i> Hagedorn	96	<i>roppae</i> Schedl. <i>Pityophthorus</i>	1026
<i>rhusi</i> Nüsima, <i>Cryphalus</i>	890	<i>rosseli</i> (Schedl), (<i>Taphroterus</i>), <i>Miniocurus</i>	950
<i>rhusi</i> Schedl, <i>Lanurgus</i>	409	<i>rosseli</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	827
<i>ribbentropi</i> Stebbing, (<i>Tomicus</i>), <i>Ips</i> (= <i>stebbingi</i>)	526	<i>rossicus</i> Eggers, <i>Pityophthorus</i>	1026
<i>ricini</i> (Eggers), (<i>Xyleborus</i>), <i>Theoborus</i>	661	<i>rossicus</i> Semenov, <i>Carphoborus</i>	309
<i>riehli</i> Eichhoff, <i>Xyleborus</i> (= <i>bidentatus</i>)	715	<i>rostratus</i> Schedl, <i>Cnestus</i>	803
<i>rigidus</i> (LeConte), (<i>Cryphalus</i>), <i>Pseudothysanoes</i>	416	<i>rothkirchi</i> Eggers, <i>Xyleborus</i>	768
<i>rileyi</i> Hopkins, <i>Xyleborus</i> (= <i>volculus</i>)	783	<i>rotrovi</i> Peyerimhoff, <i>Hypothenemus</i> (= <i>eruditus</i>)	923
<i>rimosus</i> Eichhoff, <i>Pagiocerus</i> (= <i>frontalis</i>)	213	<i>rotundatus</i> (Chapuis), (<i>Rhopalopleurus</i>), <i>Chramesus</i>	268
<i>rimskii</i> Kurenzov, <i>Scolytus</i> (= <i>jacobsoni</i>)	334	<i>rotundatus</i> Schedl, <i>Cnestus</i>	803
<i>rimulosus</i> Schedl, <i>Xyleborus</i>	768	<i>rotundicollis</i> (Eggers), (<i>Pocillips</i>), <i>Coccotrypes</i>	610
<i>ritchei</i> Sampson, (<i>Hypothenemus</i>), <i>Scolytogenes</i> (= <i>knabi</i>)	863	<i>rotundicollis</i> (Eichhoff), (<i>Stephanodere</i>), <i>Hypothenemus</i>	939
<i>ritsemae</i> (Eichhoff), (<i>Acanthurus</i>), <i>Diamerus</i>	197	<i>rotundicollis</i> Browne, <i>Xyleborus</i>	768
<i>rivalis</i> Wood, <i>Hylocurus</i>	427	<i>rotundicollis</i> Eggers, <i>Dryocoetes</i>	585
<i>robai</i> (Blackman), (<i>Pityophthoroides</i>), <i>Pityophthorus</i>	1026	<i>rotundicollis</i> Reitter, <i>Hylastes</i> (= <i>cunicularius</i>)	53
<i>robai</i> Blackman, <i>Cnesinus</i>	211	<i>rotundus</i> Hopkins, <i>Hypocryphalus</i>	871
<i>robertsi</i> (Browne), (<i>Minips</i>), <i>Acanthotomicus</i>	483	<i>roubali</i> Pfeffer, <i>Taphroterchus</i> (= <i>cillifrons</i>)	560
<i>robertsi</i> Browne, <i>Xyleborus</i> (= <i>barumbuensis</i>)	714	<i>royaleanum</i> Wood, <i>Aphanarthrum</i>	619
<i>robiniae</i> Hopkins, (<i>Hypothenemus</i>), <i>Trischlidias</i> (= <i>atomus</i>)	947	<i>ruandae</i> (Schedl), (<i>Erioschidias</i>), <i>Cosmoderes</i>	902
<i>robustipennis</i> Schedl, <i>Xyleborus</i>	768	<i>ruandae</i> Schedl, <i>Polygraphus</i>	296
<i>robustulus</i> (Schedl), (<i>Xyleborus</i>), <i>Premnobius</i>	654	<i>ruandae</i> Schedl, <i>Xyleborus</i>	768
<i>robustulus</i> Wood, <i>Coccotrypes</i>	610	<i>rubentis</i> Hopkins, <i>Cryphalus</i>	891
<i>robustum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1062	<i>rubentis</i> Wood, <i>Pityoborus</i>	968
<i>robustus</i> (Eggers), (<i>Hylesinus</i>), <i>Ficicis</i>	92	<i>ruber</i> (Eichhoff), (<i>Xyleborus</i>), <i>Taurodemus</i>	785
<i>robustus</i> (Knotek), (<i>Tomicus</i>), <i>Ips</i>	522	<i>ruber</i> Swaine, <i>Hylastes</i>	61
<i>robustus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865	<i>ruber</i> Wood, <i>Hylocurus</i>	427
<i>robustus</i> (Schedl), (<i>Mimidendrus</i>), <i>Xylocleptes</i>	549	<i>rubescens</i> Krausse, <i>Tomicus</i> (= <i>piniperda</i>)	128
<i>robustus</i> (Schedl), (<i>Phacrylus</i>), <i>Acorthylus</i>	853	<i>rubicundulus</i> Swaine, <i>Phlocosinus</i> (= <i>punctatus</i>)	251
* <i>robustus</i> (Schedl), (<i>Phlocosinus</i>), <i>Phlocosinus</i>	252	<i>rubidus</i> Beeson, (<i>Thannurgides</i>), <i>Coccotrypes</i> (= <i>papuanus</i>)	608
<i>robustus</i> (Schedl), (<i>Thamnophthorus</i>), <i>Araptus</i>	961	<i>rubidus</i> Wood, <i>Pityophthorus</i>	1026
<i>robustus</i> Blackman, <i>Hypothenemus</i> (= <i>seriatus</i>)	943	<i>rubricollis</i> (Eichhoff), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	679
<i>robustus</i> Blackman, <i>Scolytus</i>	364	<i>rubricollis</i> Blandford, <i>Corthylus</i>	1079
<i>robustus</i> Eggers, <i>Thannurgus</i>	545	<i>rubripennis</i> Reitter, <i>Tomicus</i> (= <i>piniperda</i>)	128
<i>robustus</i> Eichhoff, <i>Coccotrypes</i>	610	<i>rubripes</i> (Eggers), (<i>Pityophthorus</i>), <i>Dendrocranulus</i>	553
<i>robustus</i> Eichhoff, <i>Cryphalus</i>	891	<i>rubripes</i> Eggers, <i>Polygraphus</i> (= <i>jezoensis</i>)	288
<i>robustus</i> Pfeffer, (<i>Pityophthorus</i>) (= <i>lichtensteini</i>)	1007	<i>rude</i> Blandford, <i>Plurixosoma</i>	190
<i>robustus</i> Schedl, <i>Cladoctonus</i> (= sp.?)	238	<i>rudis</i> (LeConte), (<i>Micracis</i>), <i>Hylocurus</i>	427
<i>robustus</i> Schedl, <i>Coccotrypes</i> (= <i>robustulus</i>)	610	<i>rudis</i> Blackman, <i>Pityophthorus</i>	1026
<i>robustus</i> Schedl, (<i>Pelicerus</i>), <i>Cyrtogenius</i> (= <i>minor</i>)	566	<i>rudis</i> Blandford, <i>Phlocosinus</i>	252
<i>robustus</i> Schedl, (<i>Trogloditica</i>), <i>Hylesinus</i> (= <i>nilgrinus</i>)	82		
<i>robustus</i> Schedl, <i>Chramesus</i>	268		

- rudis* Eggers, (*Xyleborus*), *Euwallacea*
(=*xanthopus*) 696
- rudis* Eggers, *Polygraphus*** 297
- rudis* Eichhoff, *Phloeotribus* (=*setulosus*) 232
- rudis* Erichson, *Phloeoborus*** 100
- rudis* Wood, *Dendrocraniulus*** 553
- rudis likiangensis* Tsai & Yin, *Polygraphus* (=*rudis*) 297
- rudis retiventriculus* Tsai & Yin, *Polygraphus*
(=*rudis*) 297
- rudueri* Sokanovskii, *Pityogenes*** 457
- rufescens* Hopkins, (*Stephanoderes*), *Hypothenemus*
(=*obscurus*) 937
- rufescens* Stephens, (*Hylurgus*), *Hylurgops*
(=*palliatius*) 35
- ruficauda* Eggers, *Scolytotplatypus* (=*parvus*) 405
- ruficeps* Perkins, *Hypothenemus*** 940
- ruficollis* (Fabricius), (*Bostrichus*), *Tricolus*** 1045
- ruficollis* Hopkins, *Cryphalus*** 891
- ruficollis* Wood, *Stenoclyptus*** 419
- ruficollis coloradensis* Wood, *Cryphalus*
(=*ruficollis*) 892
- rufipennis* (Kirby), (*Apate*), *Polygraphus*** 297
- rufipennis* Brancsik, *Scolytus* (=*amygdali*) 323
- rufipennis* Kirby, *Dendroctonus*** 173
- rufipes* (Eichhoff), (*Hylastes*), *Hylurgopinus*** 106
- rufipes* Eggers, *Xyleborus*** 765
- rufipes* Eichhoff, (*Anthonocerus*), *Monarthrum*
(=*ingens*) 1057
- rufitarsis* (Kirby), (*Apate*), *Trypodendron*** 645
- rufithorax* Eichhoff, (*Xyleborus*), *Dryocoetoides*
(=*capucinus*) 657
- rufithorax nigricollis* Hagedorn, (*Xyleborus*),
Dryocoetoides (=*granulicauda*) 657
- rufithorax* Wood, *Tricolus*** 1045
- rufobrunneus* Eggers, *Xyleborus*** 769
- rufobrunneus dihingensis* Eggers, *Xyleborus*
(=*rufobrunneus*) 769
- rufoniger* (Schedl), (*Xyleborus*), *Euwallacea*** 693
- rufonitidum* (Schedl), (*Xyleborus*), *Cyclorhipidion*** 702
- rufonitidus* Schedl, *Sternobothrus*** 216
- rufopalliatius* Eichhoff, *Araptus*** 961
- rufopalliatius* Hopkins, *Hypothenemus* (=*columbi*)** 913
- rufopalliatius* Schedl, *Sphaerotrypes*** 201
- rufopiceus* Eggers, *Xyleborus* (=*ferrugineus*)** 739
- rufopilosus* Eggers, *Corthylus*** 1079
- rufopilosus* Schedl, *Cryphalus*** 892
- rufostriatus* Sokanovskii, *Scolytus* (=*jaroschewskii*) 336
- rufotestaceus* Schedl, *Microcorthylus*** 1067
- rufulus* Schedl, *Chiloxylon*** 557
- rufus* (Eggers), (*Spongotarsus*), *Polygraphus*** 295
- rufus* (Schedl), (*Hapalogenius*), *Rhopalopselion*** 97
- rufus* Marsham, (*Ips*), *Hylurgops* (=*palliatius*) 35
- rufus* Schedl, *Xyleborus*** 769
- rugatus* (Blandford), (*Dryocoetes*), *Ozopemon*** 590
- rugatus* Blackburn, *Xyleborus*** 769
- rugatus* Blandford, *Phloeoborus* (=*asper*) 98
- rugatus* Wood & Huang, *Pseudoxylechinus*** 114
- rugicollis* (Blandford), (*Xyleborus*),
*Ambrosiodinus*** 650
- rugicollis* (Eggers), (*Orosiotes*), *Cyrtogenius*** 565
- rugicollis* (Eggers), (*Thamnurgides*), *Coccotrypes*** 611
- rugicollis* (Schedl), (*Hexacolus*), *Scolytodes*** 395
- rugicollis* (Schedl), (*Mimiodendrus*), *Mimiocurus*** 950
- rugicollis* Browne, (*Mimidendrus*), *Cyrtogenius*
(=*ruginosus*) 565
- rugicollis* Eggers, *Dryocoetes*** 555
- rugicollis* Schedl, *Hyledius* (=sp.?) 259
- rugicollis* Swaine, *Pityophthorus* (=*tuberculatus*) 1032
- rugifer* (Schedl), (*Stephanoderes*), *Hypothenemus*** 940
- rugifer* Schedl, (*Pachynoderes*), *Hypothenemus*
(=*ruginosus*) 940
- ruginosus* Wood, *Cyrtogenius*** 565
- ruginosus* Wood, *Hypothenemus*** 940
- rugipennis* (Mannerheim), (*Hylastes*), *Hylurgops*** 40
- rugipennis* (Schedl), (*Carpophloeus*), *Cyrtogenius*** 565
- rugipennis* Eggers, *Phloeoborus* (=*rudis*) 100
- rugipennis* Schedl, (*Xyleborus*), *Xyleborinus*
(=*artestriatus*) 805
- rugipennis pinifex* (Fitch), (*Hylastes*), *Hylurgops*
(=*rugipennis*) 40
- rugipennis rugipennis* (Mannerheim), *Hylurgops*
(=*rugipennis*) 40
- rugosipennis* Schedl, *Xyleborus*** 769
- rugosipennis incertus* Schedl, *Xyleborus* 769
- rugosipes* Hopkins, *Arixyleborus*** 665
- rugosipes* Schedl, *Lanurgus*** 409
- * *rugosissimus* Cockerell, *Cryphalites*** 1052
- rugosus* (Schedl), (*Eriocryphalus*), *Cryphalus*** 892
- rugosus* Eggers, *Xyleborus* (=*alluaudi*) 710
- rugosus* Schedl, *Ptilopodius*** 900
- rugosus* Swaine, *Phloeosinus* (=*serratus*) 254
- ruguloides* Sokanovskii, *Scolytus* (=*kirschi*) 336
- Ruguloscolytus* Butovitsch (=*Scolytus*) 321
- rugulosus* (Eggers), (*Loganinus*), *Cucumonyx*** 317
- rugulosus* (Muller), (*Bostrichus*), *Scolytus*** 364
- rugulosus* (Schedl), (*Mimidendrus*), *Xylocleptes*** 550
- rugulosus* Eggers, *Dryocoetes* (=*ussuriensis*) 556
- rugulosus* Eggers, (*Poclipils*), *Coccotrypes*
(=*rotundicollis*) 610
- rugulosus* Eggers, *Phloeotribus*** 230
- rugulosus* Eggers, *Scolytodes*** 399
- rugulosus* Eggers, *Xyleborus* (=*dispar*) 730
- rugulosus baluchistani* Schedl, *Scolytus*
(=*rugulosus*) 370
- rugulosus intermedius* Sokanovskii, *Scolytus*
(=*rugulosus*) 370
- rugulosus samarkandicus* Butovitsch, *Scolytus*
(=*rugulosus*) 370
- rugulosus sanctahuciae* Hoffmann, *Scolytus*
(=*rugulosus*) 370
- rugulosus similis* Butovitsch, *Scolytus*
(=*rugulosus*) 370
- rumseyi* Hopkins, *Hypothenemus* (=*eruditus*) 922
- russulus* (Schedl), (*Xyleborus*), *Euwallacea*** 693
- russus* Swaine, *Phloeosinus* (=*randykei*)** 255

<i>rusti</i> Blackman, <i>Phloeosinus</i> (= <i>punctatus</i>)	252	<i>sampsoni</i> Nunberg, <i>Webbia</i> (= <i>quattuordecimspinus</i>)	832
<i>rusticus</i> (Wood), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865	<i>sampsoni</i> Stebbing, <i>Pityophthorus</i> (= <i>deodara</i>)	995
<i>rusticus</i> (Wood), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	650	Sampsonius Eggers	655
<i>rusticus</i> Wood, <i>Dryococtoides</i>	655	<i>sanctaluciae</i> Hoffmann, <i>Scolytus</i> (= <i>rugulosus</i>)	370
<i>rutschurueusis</i> Eggers, <i>Coccotrypes</i>	611	<i>sandakanensis</i> (Schedl), (<i>Cryphalus</i>), <i>Hypocryphalus</i>	872
<i>ruenzoriensis</i> (Schedl), (<i>Mimiophthorus</i>), <i>Mimiocurus</i>	950	<i>sandragotoensis</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	680
rybinskii Reitter, <i>Trypophloeus</i>	846	<i>sanguineus</i> Schedl, <i>Corthylus</i>	1079
<i>rybinskii salicis</i> Stark, <i>Trypophloeus</i> (= <i>rybinskii</i>)	847	<i>sanguinicollis</i> (Blandford), (<i>Xyleborus</i>), <i>Taurodenus</i>	786
<i>saalasi</i> Eggers, <i>Pityogenes</i>	457	<i>sannio</i> (Schaufuss), (<i>Pocillips</i>), <i>Coccotrypes</i>	611
<i>saalasi niger</i> Sokanovskii, <i>Pityogenes</i> (= <i>saalasi</i>)	457	<i>sannohensis</i> Murayama, <i>Phloeosinus</i>	253
<i>sabahensis</i> Bright, <i>Liparthrum</i>	279	<i>sansoni</i> Swaine, <i>Carphoborus</i>	309
<i>sabinianae</i> Hopping, (<i>Orthotomicus</i>), <i>Ips</i> (= <i>spinifer</i>)	526	<i>sapineus</i> Bright, <i>Pityophthorus</i>	1026
<i>sacchari</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>sapporoensis</i> (Niisima), (<i>Cryphalus</i>), <i>Hypothenemus</i>	940
<i>sacchari</i> Hopkins, <i>Xyleborus</i> (= <i>affinis</i>)	709	<i>sapporoensis</i> Eggers, <i>Polygraphus</i> (= <i>poligraphus</i>)	292, 295
<i>sachalinensis</i> Eggers, <i>Polygraphus</i>	295	<i>sarawakensis</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	650
<i>sachalinensis</i> Krivolutskaia, <i>Pityophthorus</i>	1026	<i>sarawakensis</i> (Schedl), (<i>Dryococtes</i>), <i>Cyrtogenius</i>	568
<i>sachalinensis</i> Michalski, <i>Scolytus</i> (= <i>esuriens</i>)	330	<i>sarawakensis</i> Schedl, <i>Cryphalus</i>	893
<i>sachalinensis</i> Sokanovskii, <i>Dryococtes</i> (= <i>autographus</i>)	575	<i>sarawakensis</i> Schedl, <i>Webbia</i>	832
<i>sachalinensis frontalis</i> Kurenzov, <i>Polygraphus</i> (= <i>sachalinensis</i>)	299	<i>sardus</i> Eggers, <i>Thamnurgus</i>	545
<i>sachtleni</i> Schedl, <i>Xylechinus</i>	105	<i>sardus</i> Strohmeier, <i>Dryococtes</i> (= <i>villosus</i>)	557
<i>saginata</i> Mannerheim, <i>Polygraphus</i> (= <i>rufipeunis</i>)	295	sartor Schedl, <i>Xyleborus</i>	769
sagittarius Schedl, <i>Eccoptopterus</i>	520	<i>sassaensis</i> (Eggers), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	940
<i>sahlbergi</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>ratzeburgi</i>)	363	satoi Schedl, <i>Xyleborus</i>	769
sakalava Schedl, <i>Xyleborus</i>	769	<i>saturatum</i> Peyerimhoff, <i>Aphanarthrum</i>	619
sakoe (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	650	<i>saudiarabiae</i> (Schedl), (<i>Chilodendron</i>), <i>Hylesinopsis</i>	95
salakeensis (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	611	saundersi Wood, <i>Tricolus</i>	1045
salebrosus Eichhoff, <i>Hylastes</i>	61	<i>sauropteroideus</i> Schedl, (<i>Xyleborus</i>), <i>Sauroptilius</i> (= <i>sauropterus</i>)	971
<i>Saliciphilus</i> Sokanovskii (= <i>Taphrorychus</i>)	556	<i>sauropterus</i> (Schedl), (<i>Xyleborus</i>), <i>Sauroptilius</i>	971
<i>salicis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942	Sauroptilius Browne	971
<i>salicis</i> Hopkins, <i>Procryphalus</i> (= <i>utahensis</i>)	848	Saurotocis Wood	419
<i>salicis</i> Hopkins, <i>Trypophloeus</i>	847	<i>savadae</i> Nobuchi & Takahashi, <i>Cryphalus</i>	893
<i>salicis</i> Stark, <i>Trypophloeus</i> (= <i>rybinskii</i>)	847	<i>saxseni</i> (Ratzeburg), (<i>Bostrichus</i>), <i>Xyleborinus</i>	810
sallaei Blandford, <i>Dendroterus</i>	952	sayi (Hopkins), (<i>Anisandrus</i>), <i>Xyleborus</i>	769
<i>saltoni</i> Blackman, <i>Gnathotrichus</i> (= <i>pilosus</i>)	1037	scaber Erichson, <i>Phloeoborus</i>	100
saltuarius Weise, <i>Cryphalus</i>	892	<i>scaber</i> Marsham, (<i>Ips</i>), <i>Hylesinus</i> (= <i>toranio</i>)	85
saltrini (Blandford), (<i>Xyleborus</i>), <i>Taurodemus</i>	785	scaber Schedl, <i>Xyleborus</i>	770
<i>samarkandicus</i> Butovitch, <i>Scolytus</i> (= <i>rugulosus</i>)	370	<i>scaber</i> Swaine, <i>Hylastes</i> (= <i>porculus</i>)	61
sambesianus Eggers, <i>Hypothenemus</i>	940	scabiosus Blandford, <i>Xylechinus</i>	118
sambuci Blackman, <i>Pityophthorus</i>	1026	scabratus Blandford, <i>Phloeotribus</i>	230
samoanus (Browne), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	865	scabratus Schedl, <i>Xyleborus</i>	770
samoanus (Eggers), (<i>Dryococtes</i>), <i>Cyrtogenius</i>	568	scabricollis (Hopkins), (<i>Phloeophthorus</i>), <i>Phloeotribus</i>	230
samoanus Schedl, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	91	scabricollis (LeConte), (<i>Xyloterus</i>), <i>Trypodendron</i>	645
samoanus Schedl, <i>Eidophelus</i>	867	scabricollis (Schedl), (<i>Ozopenon</i>), <i>Xyleborus</i>	770
samoensis Beeson, <i>Cryphalus</i>	893	scabricollis Browne, <i>Cyrtogenius</i>	568
<i>samoensis</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>piceus</i>)	693	scabricollis Eichhoff, <i>Cryphalus</i>	893
<i>sampsoni</i> Donisthorpe, <i>Xyleborus</i> (= <i>dryographus</i>)	732		
<i>sampsoni</i> Eggers, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>mutilatus</i>)	799		

- scabridus* Schedl, (*Xyleborus*), *Ambrosiodinus*
(=*aegir*) 670
- scabridus* Schedl, *Pityophthorus* 1026
- scabrifrons* St., (*Hylesinus*), *Hylastes*
(=*cunicularius*) 51
- scabrior* (Schedl), (*Xyleborus*), *Leptoxyleborus* 660
- scabrior* Schedl, (*Xyleborus*), *Ambrosiodinus*
(=*neglectus*) 677
- scabripennis* (Blandford), (*Xyleborus*),
Arixyleborus 669
- scabripennis* Schedl, *Cryphalus* 594
- scabripennis* Zimmermann, *Hylastes* (= *salebrosus*) . . 62
- scalaris* (Schedl), (*Xyleborus*), *Ambrosiodinus* 680
- scaliger* Hagedorn, (*Loganius*), *Cnemonyx*
(=*flavicornis*) 315
- sculptor* (Schedl), (*Xyleborus*), *Cyclorhipidion* 702
- sculptor* Blackman, *Pityophthorus* 1026
- sculptor* Schedl, *Polygraphus* 299
- scalptus* Bright, *Pityophthorus* 1027
- scandinarius* Lekander, *Hylastes* 62
- scapulare* (Schedl), (*Xyleborus*), *Cyclorhipidion* . . . 702
- scarabaeoides* (Bernard), (*Scolytus*), *Phloeotribus* . . 230
- schabliovskii* Stark, *Lymantria* (= *aceris*) 554
- schaufussi* (Blandford), (*Xyleborus*), *Xyleborinus* . . 516
- schaufussi* Blandford, *Amphicranus* 1045
- schaufussi* Schedl, *Corthylus* 1079
- schedli* (Blackman), (*Neodryocoetes*), *Araptus* 961
- schedli* Bright, *Monarthrum* (= *minutissimum*) 1059
- schedli* Browne, *Hypothenemus* 940
- schedli* Eggers, *Xyleborus* (= *ferrugineus*) 739
- schedli* Wood, *Cnemonyx* (= *errans*) 314
- schedli* Wood, *Dendrocranius* 553
- Schedlia* Browne 822
- * *schellucien* (Hagedorn), (*Hylastites*), *Hyhurgops* . . 41
- schenklingi* Hagedorn, *Chortastus* 312
- schervyei* Semenov, *Scolytus* 370
- schildi* Schedl, *Xyleborus* 770
- schimitscheki* Eggers, *Taphrorychus* 560
- schizolobius* Schedl, (*Xyleborus*), *Dryocoetoides*
(=*pseudosolitarius*) 655
- schliehi* (Stebbing), (*Xyleborus*), *Amasa* 654
- schmutzenhoferi* Holtzschuh, *Ips* (= *stebbingi*) 527
- schoenbachi* Kirsch, *Phloeotribus* 232
- schoenherri* (Schedl), (*Xyleborus*), *Xyleborinus* . . . 516
- schoenherri* Schedl, *Cucinus* 211
- schoutedeni* Eggers, *Xyleborus* 770
- schreineri* Eggers, (*Xyleborus*), *Xyleborinus*
(=*sharpae*) 516
- schreineri* Eichhoff, (*Cryphalus*), *Ernioporus*
(=*caucasicus*) 549
- schrenkianae* Pjatnitskii, *Pityophthorus* 1026
- schultzei* (Schedl), (*Dryocoetes*), *Dryocoetops* 555
- schultzei* Schedl, (*Xyleborus*), *Eucallacea*
(=*fornicatus*) 690
- schumensis* Eggers, *Phloeosinus* (= *bicolor*) 243
- schwarz* (Blackman), (*Thamniophthorus*), *Araptus* . . 961
- schwarz* (Hopkins), (*Erineophilus*), *Scolytodes* 399
- schwarz* Blackman, *Hylocurus* 428
- schwarz* Blackman, *Pityophthorus* 1027
- schwarz* Hopkins, (*Cosmoderes*), *Hypothenemus*
(=*eruditus*) 922
- schwarz* Hopkins, *Xyleborus* (= *rolulus*) 753
- schuerdtfegeri* (Schedl), (*Conophthorus*),
Pityophthorus 1027
- Scierus* LeConte 29
- scitula* Wood, *Micracisella* 431
- scitulus* Wood, *Hylocurus* 428
- scitulus* Wood, *Pityophthorus* 1027
- scitulus* Wood, *Tricolus* 1045
- scitus* Blandford, *Pityogenes* 457
- sclerocaryae* (Schedl), (*Xyleborus*), *Xyleborinus* . . . 516
- scobinatus* Hagedorn, *Xyleborus* 770
- scobinosus* Eichhoff, *Hylastes* (= *salebrosus*) 61
- scobipennis* Chapuis, (*Hylesinus*), *Ficicis*
(=*despectus*) 90
- scolytus* Sulzer, *Scolytus* (= *scolytus*) 374
- Scolytidae Latreille 28
- Scolytidae verae (= *Scolytini*) 313
- Scolytidarii Latreille (= *Scolytidae*) 28
- Scolytides Latreille (= *Scolytini*) 313
- Scolytinae Latreille 313
- Scolytini Latreille 313
- Scolytocheilus* Reitter (= *Scolytus*) 321
- scolytocheilus* Schedl (= *Scolytomimus*) 835
- Scolytodes* Ferrari 357
- Scolytogenes* Eichhoff 855
- scolytoides* Eichhoff, *Xyloctonus* 841
- scolytomimoides* (Nobuchi), (*Cryphalomorphus*),
Scolytogenes 865
- Scolytomimus* Blandford 835
- Scolytoplatypus* Blandford (= *Scolytoplatypodini*) . . . 401
- Scolytoplatypidae* Blandford (= *Scolytoplatypodini*) . . 401
- Scolytoplatypini* Blandford (= *Scolytoplatypodini*) . . 401
- Scolytoplatypodidae* Blandford
(= *Scolytoplatypodini*) 401
- Scolytoplatypodinae* Blandford
(= *Scolytoplatypodini*) 401
- Scolytoplatypodini* Blandford (= *Scolytoplatypodini*) . . 401
- Scolytoplatypus* Schaufuss 401
- Scolytopsis* Blandford 320
- scolytus* (Fabricius), (*Bostriculus*), *Scolytus* 371
- Scolytus* Geoffroy 28, 321
- scopiger* Berger, *Cryphalus* 594
- scopulorum* Hopkins, *Xyleborus* (= *intrusus*) 745
- scopulorum* Hopkins, *Conophthorus*
(=*ponderosae*) 965
- scopulorum* Swaine, *Phloeosinus* 253
- scopulorum* *neomexicanus* Blackman, *Phloeosinus*
(=*scopulorum*) 253
- scopulorum* *scopulorum* Swaine, *Phloeosinus*
(=*scopulorum*) 253
- scorpium* (Schedl), (*Xyleborus*), *Cyclorhipidion* . . . 702
- scotti* Eggers, *Cyrtogenius* 565
- scriba* Gozis, *Cryphalus* (= *saltuarius*) 593

<i>scriptor</i> Blackman, <i>Pityophthorus</i>	1027	<i>semipallens</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1062
<i>scrobiceps</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1062	<i>semipilosus</i> Eggers, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>collarti</i>)	806
<i>scrofa</i> (Schedl), (<i>Cladoctoporus</i>), <i>Trietennus</i>	541	<i>semipolitus</i> Schedl, <i>Xyleborus</i>	771
<i>scrutator</i> (Pandelle), (<i>Thammurgus</i>), <i>Tiarophorus</i>	540	<i>semipunctatus</i> Eggers, <i>Xyleborus</i>	771
<i>sculptilis</i> (Schedl), (<i>Pocetilis</i>), <i>Coccotrypes</i>	611	<i>semipunctatus</i> Wood, <i>Scolytodes</i>	399
<i>sculptilis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	816	<i>semirostris</i> Hopping, <i>Ips</i> (= <i>tridens engelmanni</i>)	528
<i>sculpturatus</i> (Blandford), (<i>Bothrosternus</i>), <i>Sternobothrus</i>	216	<i>semirudis</i> Blandford, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	696
<i>sculpturatus</i> Eichhoff, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>rotundicollis</i>)	940	<i>semirufus</i> Reitter, <i>Thammurgus</i>	545
<i>scutellare</i> (LeConte), (<i>Corthylus</i>), <i>Monarthrum</i>	1062	<i>semirufus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>obliquecaudata</i>)	677
<i>scutiae</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	940	<i>semistriatus</i> Schedl, <i>Xyleborus</i>	771
<i>scutulatus</i> (Blandford), (<i>Hylesinus</i>), <i>Neopteleobius</i>	72	<i>semitruncatus</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>cyliandrotonicus</i>)	793
<i>sechelti</i> Swaine, <i>Dryocoetes</i>	585	<i>senchalensis</i> Beeson, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>andamauensis</i>)	686
<i>sechii</i> Nunberg, <i>Dendrocraniulus</i>	553	<i>senecionis</i> Schedl, <i>Thammurgus</i>	545
<i>secundus</i> Blackman, <i>Pityoborus</i>	968	<i>senegambiensis</i> (Schedl), (<i>Hapalogenius</i>), <i>Hylesinopsis</i>	95
<i>securigerus</i> (Blackman), (<i>Chalcophylus</i>), <i>Pseudothyanoes</i>	417	<i>senex</i> Schedl, <i>Tricolus</i>	1045
<i>securus</i> (Schedl), (<i>Erieryphalus</i>), <i>Cryphalus</i>	894	<i>senex</i> Wichmann, <i>Pityophthorus</i> (= <i>buyssoni</i>)	985
<i>securus</i> Wood, <i>Chramesus</i>	268	<i>sensarmai</i> Maiti & Saha, <i>Hyorrhynchus</i>	192
<i>securus</i> Wood, <i>Dendrocraniulus</i>	553	<i>senticosus</i> Wood, <i>Corthylus</i>	1079
<i>securus</i> Wood, <i>Pseudothyanoes</i>	417	<i>sentosus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Xyleborinus</i>	816
<i>secus</i> Wood, <i>Chramesus</i>	268	<i>sentosus</i> Wood, <i>Corthylus</i>	1079
<i>secus</i> Wood, <i>Hylocurus</i>	428	<i>sentus</i> Wood, (<i>Hoplitophthorus</i>), <i>Cladoctonus</i> (= <i>interruptus</i>)	239
<i>secutus</i> Wood, <i>Hylesinopsis</i>	95	<i>sentus</i> Wood, <i>Corthylus</i>	1079
<i>sedlaceki</i> (Schedl), (<i>Ozodendron</i>), <i>Cyrtogennius</i>	568	<i>sentus</i> Wood, <i>Micracis</i>	433
<i>sedulus</i> Blackman, <i>Pseudothyanoes</i>	417	<i>seoulensis</i> Choo & Woo, <i>Hypothenemus</i>	940
<i>segnis</i> Blackman, <i>Pityophthorus</i>	1027	<i>separandus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866
<i>seirindensis</i> Murayama, <i>Pityogenes</i>	458	<i>separandus</i> Schedl, <i>Xyleborus</i>	771
<i>seiryorensis</i> Murayama, <i>Xyleborus</i>	770	<i>separatus</i> Bright, <i>Pityophthorus</i>	1028
<i>seiryuensis</i> Murayama, <i>Pityophthorus</i>	1028	<i>sepicola</i> Lovendal, <i>Lymantor</i> (= <i>coryli</i>)	556
<i>sejugatus</i> Schedl, (<i>Xyleborus</i>), <i>Leptoxyleborus</i> (= <i>depressus</i>)	660	<i>septentrionalis</i> Eggers, <i>Hylastes</i> (= <i>plumbeus</i>)	60
<i>sejugatus</i> Schedl, <i>Cryphalus</i>	894	<i>septentrionalis</i> Nisima, <i>Xyleborus</i>	771
<i>semenori</i> (Kurenzov), (<i>Eocryphalus</i>), <i>Eruoporus</i>	851	<i>septentrionis</i> Mannerheim, (<i>Bostrichus</i>), <i>Dryocoetes</i> (= <i>autographus</i>)	577
<i>semenori</i> (Spessivtsev), (<i>Eccoptogaster</i>), <i>Scolytus</i>	376	<i>sequoiae</i> Hopkins, <i>Phloeosinus</i>	253
<i>semibrunneus</i> (Eggers), (<i>Chramesus</i>), <i>Pseudochramesus</i>	262	<i>sercinus</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>xanthopus</i>)	696
<i>semicarinatus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	680	<i>serenus</i> Wood, <i>Scolytodes</i>	399
<i>semicastaneus</i> Mannerheim, (<i>Bostrichus</i>), <i>Dryocoetes</i> (= <i>autographus</i>)	578	<i>seriata</i> (Browne), (<i>Pseudonebbia</i>), <i>Webbia</i>	832
<i>semicircularis</i> Schedl, <i>Xyleborus</i> (= <i>emarginatus</i>)	733	<i>seriatus</i> (Eggers), (<i>Pseudophloeotribus</i>), <i>Hylesinopsis</i>	95
<i>semicostatus</i> (Schedl), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	658	<i>seriatus</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	940
<i>semiernis</i> (Nunberg), (<i>Pityophthorus</i>), <i>Gnatholeptus</i>	970	<i>seriatus</i> Blandford, <i>Phloeosinus</i>	254
<i>semiernis</i> Schedl, <i>Xyleborus</i>	771	<i>seriatus</i> Blandford, <i>Xyleborus</i>	771
<i>semigranulosus</i> Blandford, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>crassiusculus</i>)	792	<i>seriatus</i> Eggers, <i>Camptocerus</i>	320
<i>semigranulatus</i> (Schedl), (<i>Dryocoetes</i>), <i>Dryocoetiops</i>	588	<i>seriatus</i> Eggers, <i>Cryptocarenum</i>	904
<i>semigranulatus</i> (Schedl), (<i>Xyleborus</i>), <i>Leptoxyleborus</i>	660	<i>seriatus</i> LeConte, (<i>Pityophthorus</i>), <i>Pityoborus</i> (= <i>comatus</i>)	968
<i>seminitens</i> (Blandford), (<i>Xyleborus</i>), <i>Arizyleborus</i>	669	<i>seriatus</i> Reitter, <i>Polygraphus</i>	299
<i>sempiopacus</i> Eichhoff, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>crassiusculus</i>)	791		

- sericeus* (Mannerheim), (*Hylurgus*),
Pseudohylesinus 112
- sericeus* (Schedl), (*Erioscladius*), *Cosmoderes* 902
- sericeus* Chapuis, *Phloeoborus* (= *scaber*) 100
- sericeus* Marsham, *Ptelcobius* (= *rittatus*) 72
- sericeus* Motschulsky, (*Hylesinus*), *Cryphalus*
 (= *dorsalis*) 579
- serjaniae* Wood, *Micracisella* 431
- Serrastus* Nunberg 253
- serratus* (LeConte), (*Hylesinus*), *Phlocosinus* 254
- serratus* Bruck, *Pseudohylesinus* (= *nebulosus*) 111
- serratus* Eggers, *Phloeotribus* 232
- serratus* Fabricius, (*Bostrichus*), *Taurodemus*
 (= *varians*) 756
- serratus* Panzer, (*Bostrichus*), *Erioporus* (= *fagi*) 551
- serratus* Swaine, *Pityophthorus* 1028
- serratus* Swaine, *Xyleborus* (= *obesus*) 756
- serratus* Wood, *Corthylus* 1079
- serratus* Wood, *Eupagiocerus* (= *ater*) 214
- serrifer* (Hagedorn), (*Chortastus*), *Serrastus* 253
- serrifer* Wichmann, *Phlocosinus* (= *thujae*) 258
- serrulatus* Eggers, *Corthylus* 1079
- setifer* (Schedl), (*Mimips*), *Mimiocurus* 950
- setifer* (Wood), (*Cryphalomorphus*), *Scolytogenes* 866
- setifer* Browne, *Pityophthorus* 1028
- setifer* Eggers, *Ctenoxylon* 838
- setifer* Schedl, *Pseudochramesus* 262
- setiger* Schedl, *Chramesus* 269
- setistriatus* (Lea), (*Cryphalus*), *Cosmoderes* 902
- setosus* (Blandford), (*Hexacolus*), *Scolytodes* 399
- setosus* (Eggers), (*Kissophagus*), *Phloeoditica* 236
- setosus* (Eichhoff), (*Hypoborus*), *Hypothenemus* 943
- setosus* Beeson, *Polygraphus* (= *setosus*) 299
- setosus* Beeson, (*Thammurgides*), *Coccotrypes*
 (= *advena*) 592
- setosus* Blackman, *Pityophthorus* 1028
- setosus* Bruck, *Phlocosinus* 255
- setosus* Eggers, *Cnesinus* 211
- setosus* Schedl, *Polygraphus* 299
- setosus* Schedl, *Scolytoplatypus* (= *papuanus*) 405
- setosus* Wood, *Bothinodroctonus* 310
- setosus* Wood, *Cactopinus* 436
- setosus* Wood, *Chramesus* 269
- setulosum* Waterhouse, *Pycnarthrum* 356
- setulosus* (Eggers), (*Loganius*), *Cnemonyx* 317
- setulosus* Blandford, *Cnesinus* 211
- setulosus* Eggers, *Microborus* (= *aberrans*) 353
- setulosus* Eggers, *Xyleborus* 771
- setulosus* Eichhoff, *Phloeotribus* 232
- setulosus* Schedl, *Strombophorus* 204
- seulensis* Murayama, *Scolytus* (= *schevyrewi*) 371
- severus* Bright, *Pityoborus* (= *frontalis*) 968
- severus* Wood, *Dryocoetoides* 655
- sexdentatum* Eggers, (*Anchonocerus*), *Monarthrum*
 (= *sexdenticulum*) 1062
- sexdentatum* Eggers, *Monarthrum* 1062
- sexdentatus* (Boerner), (*Dermestes*), *Ips* 522
- sexdentatus* (Eggers), (*Mimips*), *Acanthotomicus* 453
- sexdentatus* (Eggers), (*Xyleborus*), *Ambrosiodinus* 680
- sexdentatus* Eggers, *Sampsonius* 656
- sexdentatus* Olivier, (*Scolytus*), *Pityogenes*
 (= *chalcographus*) 450
- sexdentatus* junianicus Sokanovskii, *Ips*
 (= *sexdentatus*) 526
- sexdenticulum* Wood, *Monarthrum* 1062
- sexnotatus* (Schedl), (*Xyleborus*), *Premnobius* 655
- sexspinatum* (Schedl), (*Xyleborus*),
Cyclorhipidius 702
- sexspinosus* (Schedl), (*Ips*), *Acanthotomicus* 453
- ° *sexspinosus* (Schedl), (*Phlocosinites*),
Phlocosinus 255
- sexspinosus* Eggers, *Premnobius* 655
- sexspinosus* Motschulsky, *Eccoptopterus*
 (= *spinosus*) 821
- sexspinosus pluridentatus* Schedl, *Eccoptopterus*
 (= *multispinosus*) 520
- sextuberculatus* Eggers, *Pityophthorus* 1028
- sextuberculatus* Schedl, (*Gnathotrichus*),
Gnathotrupes (= *longipennis*) 1041
- sextuberculatus* Schedl, *Xyleborus* 771
- seydeli* (Nunberg), (*Rhopalocryphus*), *Glostatus* 536
- seydeli* (Schedl), (*Mimips*), *Acanthotomicus* 483
- shablionskyi* Kurenzov, *Hylesinus* 83
- shanhaiensis* Yin & Huang, *Scolytus* 376
- shamoni* (Blackman), (*Pityophthorus*),
Gnatholeptus 970
- shanonum* Beeson, (*Thammurgides*), *Coccotrypes*
 (= *rhizophorae*) 610
- shariensis* Niisima, *Polygraphus* 299
- sharpae* (Hopkins), (*Xyleborus*), *Xyleborinus* 816
- sharpi* (Blandford), (*Xyleborus*), *Taurodemus* 756
- sharpi* (Guillebeau), (*Phloeophthorus*),
Phloeotribus 233
- sharpi* Guillebeau, *Phloeotribus* (= *cristatus*) 220
- sharpi lenis* (Wood), *Taurodemus* (= *sharpi*) 756
- sharpi sharpi* Blandford, *Taurodemus* (= *sharpi*) 756
- shensi* Tsai & Yin, *Phlocosinus* 255
- shepardi* Blackman, *Pityophthorus* (= *intextus*) 1004
- shikisanii*, *Scolytus* (= *chikisanii*) 327
- shimanensis* Murayama, *Erioporus* 555
- shinanoensis* Yono, *Ips* (= *cembrae*) 499
- shionomisakiensis* Murayama, *Xyleborus* 771
- shiva Maiti & Saha*, *Hyorrhynchus* 192
- shiva Maiti & Saha*, *Xyleborus* 772
- shogun* Blandford, *Scolytoplatypus* 406
- shoreae* (Stebbing), (*Tomicus*), *Xyleborus* 772
- shoreae* Browne, *Cryptoxyleborus* 529
- shoreae* Schedl, *Ptilopodius* 900
- shoshone* Hopkins, *Dendroctonus* (= *murayanae*) 158
- shotoensis* Murayama, *Phlocosinus* (= *rudis*) 252
- sibiricus* Eggers, (*Eccoptogaster*), *Scolytus*
 (= *ratzeburgi*) 363
- sibiricus* Nunberg, *Pityophthorus* (= *abietinus*) 975
- sibericus* Stark, *Pityophthorus* (= *micrographus*) 1012
- sibsagaricus* (Eggers), (*Xyleborus*), *Euwallacea* 693

<i>sichotensis</i> Kurenzov, <i>Cryphalus</i>	594	<i>similis</i> Eggers, <i>Phlocosinus</i>	255
<i>sichotensis</i> Kurenzov, <i>Pityophthorus</i>	1028	<i>similis</i> Eggers, <i>Xyleborus</i> (= <i>rufobrunneus</i>)	769
<i>sichus</i> Schedl, <i>Xyleborus</i>	772	<i>similis</i> Eichhoff, <i>Pityophthorus</i>	1028
<i>sicula</i> DelGuercio, (<i>Comesiella</i>), <i>Phlocotribus</i> (= <i>pubifrons</i>)	228	<i>similis</i> Ferrari, <i>Xyleborus</i>	772
<i>siculus</i> Eggers, <i>Scolytus</i> (= <i>koenigi</i>)	338	<i>similis</i> Hagedorn, <i>Dactylipalpus</i> (= <i>groucellei</i>)	102
<i>siculus</i> Eggers, <i>Taphrorhynchus</i> (= <i>bicolor</i>)	558	<i>similis</i> Hopkins, <i>Hypothenemus</i> (= <i>sparsus</i>)	945
<i>siculus</i> Eggers, <i>Thamniurgus</i>	546	<i>similis</i> LeConte, <i>Dendroctonus</i> (= <i>rufipennis</i>)	177
<i>siculus</i> Ferrari, <i>Hypoborus</i> (= <i>ficus</i>)	281	<i>similis</i> Nunberg, <i>Cladotomus</i>	239
<i>sidanus</i> (Schedl), (<i>Dryocoetes</i>), <i>Xylocleptes</i>	550	<i>similis</i> Schedl, <i>Gnathotripes</i>	1042
<i>sidneyanus</i> (Nordlinger), (<i>Bostrichus</i>), <i>Cryphalus</i>	894	<i>similis</i> Wood, <i>Micracisella</i>	431
<i>sierraeonensis</i> Eggers, (<i>Poecilips</i>), <i>Coccotrypes</i> (= <i>confusus</i>)	597	<i>simillimus</i> Perkins, <i>Xyleborus</i>	774
<i>sierrensis</i> Bright, <i>Pityophthorus</i>	1028	<i>simillimus</i> Schedl, <i>Corthylus</i>	1079
<i>signis</i> Hagedorn, <i>Eucallacca</i> (= <i>xanthopus</i>)	696	<i>simmeli</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>intricatus</i>)	334
<i>signatifrons</i> (Schedl), (<i>Brachyspartus</i>), <i>Corthylurus</i>	1069	<i>simoni</i> (Reitter), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	944
<i>signatifrons</i> Browne, <i>Pityophthorus</i>	1028	<i>simplex</i> (Browne), (<i>Cryptoxyleborus</i>), <i>Webbia</i>	833
<i>signatipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	816	<i>simplex</i> (Schedl), (<i>Eriocryphalus</i>), <i>Cryphalus</i>	894
<i>signatipennis</i> Schedl, <i>Chramesus</i>	269	<i>simplex</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	483
<i>signatum</i> (Fabricius), (<i>Apate</i>), <i>Trypodendron</i>	646	<i>simplex</i> Blandford, <i>Hylocurus</i>	428
<i>signatus</i> Ferrari, (<i>Corthylus</i>), <i>Monarthrum</i> (= <i>bicolor</i>)	1051	<i>simplex</i> LeConte, <i>Carphoborus</i>	309
<i>signatus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>funereus</i>)	674	<i>simplex</i> LeConte, <i>Dendroctonus</i>	180
<i>signatus</i> Strohmeier, <i>Phlocoborus</i>	101	<i>simplex</i> Rey, <i>Hylastes</i> (= <i>opacus</i>)	59
<i>signiceps</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	680	<i>simplex</i> Schedl, <i>Hylocurus</i> (= <i>brasilensis</i>)	423
<i>signifer</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	680	<i>simplex</i> Wollaston, <i>Aphanarthrum</i> (= <i>canescens</i>)	617
<i>silraiae</i> Schedl, <i>Cyrtogenius</i>	568	<i>simplex</i> Wood, <i>Corthylus</i>	1079
<i>silvanus</i> Schedl, <i>Cryphalus</i>	894	<i>simplex</i> Wood, <i>Phlocotribus</i>	233
<i>silvaticus</i> Bright, <i>Scolytus</i> (= <i>hermosus</i>)	331	<i>simplex</i> Wood, <i>Pseudotlysanocis</i>	417
<i>silvestris</i> Beeson, <i>Coccotrypes</i> (= <i>longior</i>)	605	<i>simplicidens</i> Wood, <i>Phlocotribus</i>	233
<i>silvestris</i> Beeson, <i>Xyleborus</i> (= <i>volulus</i>)	783	<i>simplicis</i> Wood, <i>Chramesus</i>	269
<i>similans</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	816	<i>simplicis</i> Wood, <i>Tricolus</i>	1045
<i>similaris</i> Schedl, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>onynganus</i>)	864	<i>simulatus</i> Bright, <i>Xyleborus</i>	774
<i>similaris</i> Schedl, <i>Xyleborus</i> (= <i>malgasicus</i>)	750	<i>sindorae</i> Browne, <i>Acanthotomicus</i>	483
<i>simile</i> Eggers, <i>Pycnarthrum</i> (= <i>kleinei</i>)	355	<i>sinense</i> (Eggers), (<i>Trypodendron</i>), <i>Indocryphalus</i>	649
<i>simile</i> Eichhoff, (<i>Pterocydon</i>), <i>Monarthrum</i> (= <i>fasciatum</i>)	1055	<i>sinensis</i> Eggers, <i>Polygraphus</i>	299
<i>similis</i> (Eggers), (<i>Chortastus</i>), <i>Serrastus</i>	283	<i>sinensis</i> Eggers, <i>Scolytus</i> (= <i>scheryrewi</i>)	370
<i>similis</i> (Eggers), (<i>Dendriurgus</i>), <i>Coccotrypes</i>	612	<i>sinensis</i> Eggers, <i>Xyleborus</i>	774
<i>similis</i> (Eggers), (<i>Loganius</i>), <i>Cnemidomys</i>	317	<i>sinensis</i> Schedl, <i>Phlocosinus</i>	255
<i>similis</i> (Eggers), (<i>Prionosceles</i>), <i>Scolytodes</i>	399	<i>sinensis</i> Tsai & Huang, <i>Scolytoplatypus</i> (= <i>mikado</i>)	404
<i>similis</i> (Eggers), (<i>Stephanoderes</i>), <i>Cryphalus</i>	894	<i>sinensis</i> Wood & Huang, <i>Pseudoxylechinus</i>	114
<i>similis</i> (Eggers), (<i>Xyleborinus</i>), <i>Webbia</i>	832	<i>singalangicus</i> Eggers, <i>Ozopemon</i> (= <i>theklae</i>)	590
<i>similis</i> Blackman, <i>Cnesinus</i> (= <i>costulatus</i>)	207	<i>singularis</i> Bright, <i>Pityophthorus</i> (= <i>blandus</i>)	983
<i>similis</i> Blackman, <i>Pseudohylesinus</i> (= <i>sugae</i>)	114	<i>singularis</i> Wood, <i>Hylocurus</i>	428
<i>similis</i> Butovitsch, <i>Scolytus</i> (= <i>rugulosus</i>)	370	<i>singularis</i> Wood, <i>Pseudopityophthorus</i>	976
<i>similis</i> Eggers, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	91	<i>sinoabietis</i> Tsai & Li, <i>Cryphalus</i>	895
<i>similis</i> Eggers, (<i>Nigrites</i>), <i>Scolytogenes</i> (= <i>darwini</i>)	861	<i>sinoabietis opienensis</i> Tsai & Li, <i>Cryphalus</i> (= <i>sinoabietis</i>)	895
<i>similis</i> Eggers, (<i>Trigonogenius</i>), <i>Pityophthorus</i> (= <i>eggersi</i>)	997	<i>sinopae</i> Schedl, <i>Pityophthorus</i>	1028
<i>similis</i> Eggers, <i>Dryocoetes</i> (= <i>alni</i>)	572	<i>Sinophloeus</i> Brethes	105
<i>similis</i> Eggers, <i>Ozopemon</i>	590	<i>sinopiceus</i> Tsai, <i>Scolytus</i>	376
<i>similis</i> Eggers, <i>Phlocoborus</i> (= <i>signatus</i>)	101	<i>sinuosa</i> (Schedl), (<i>Bothryperus</i>), <i>Phrixosoma</i>	190
		<i>siobanus</i> Eggers, (<i>Xyleborus</i>), <i>Eucallacca</i> (= <i>andamanensis</i>)	686
		<i>sionio</i> Blandford, <i>Scolytoplatypus</i>	406
		<i>sionxensis</i> Bright, <i>Pityophthorus</i> (= <i>boycei</i>)	984
		<i>sipolisii</i> Fairmaire, <i>Phlocoborus</i> (= <i>procerus</i>)	99
		<i>siporanus</i> (Eggers), (<i>Dryocoetes</i>), <i>Cyrtogenius</i>	568

- siporanus* Hagedorn, (*Xyleborus*), *Eucallacca*
(=*wallacci*) 695
- siranbeanus* (Eggers), (*Xyleborus*), *Amasa* 654
- siranbeanus* Eggers, *Ozopemon* (= *theklae*) 590
- sisyrnophorum* (Hagedorn), (*Xyleborus*),
Cyclorhipidion 702
- sitchensis* Swaine, *Pseudohylesinus* 113
- sircalikensis* Stebbing, *Sphaerotrypes* 201
- smithi* Schedl, *Pityophthorus* (= *fuscus*) 999
- sobrinus* (Schedl), (*Stephanoderes*),
Hypothenemus 944
- sobrinus* Blackman, *Scolytus* (= *unispinosus*) 350
- sobrinus* Eichhoff, (*Xyleborus*), *Xyleborinus*
(=*saxeseni*) 514
- sobrinus* Wood, *Araptus* 961
- sobrinus* Wood, *Corthylus* 1079
- sobrinus* Wood, *Dendrocaulus* 553
- sobrinus* Wood, *Pericoryphalus* 945
- sobrinus* Wood, *Pityophthorus* 1025
- socialis* (Schedl), (*Stephanoderes*), *Hypothenemus* 944
- societatis* Beeson, *Xyleborus* (= *affinis*) 709
- socius* Blackman, *Pityophthorus* (= *deletus*) 994
- sodalis* (Schedl), (*Cryphalomorphus*), *Scolytogenes* 566
- sodalis* Blandford, *Phloeotribus* (= *setulosus*) 232
- sodalis* Wood, *Dendroterus* 952
- solatus* Wood, *Pityophthorus* 1029
- solers* Blackman, *Pityophthorus* 1029
- solicitatus* Wood, *Chramesus* 269
- solidus* (Eichhoff), (*Xyleborus*), *Xylosandrus* 500
- solidus* (Schedl), (*Blastophagus*), *Hyurgonotus* 157
- solidus* Schedl, *Gnathotrupes* 1042
- solidus* Schedl, *Polygraphus* (= *aequalis*) 254
- solitariceps* Schedl, (*Xyleborus*), *Theoborus*
(=*ricini*) 661
- solitariformis* (Schedl), (*Xyleborus*),
Dryocoetoides 655
- solitarius* (Schedl), (*Xyleborus*), *Dryocoetoides* 655
- solitaripennis* Schedl, (*Xyleborus*), *Dryocoetoides*
(=*paradoxus*) 655
- solitarius* (Schedl), (*Erioschidias*), *Cosmoderes* 901
- solitarius* (Schedl), (*Stephanoderes*),
Hypothenemus 944
- solitarius* Hagedorn, (*Xyleborus*), *Dryocoetoides*
(=*cristatus*) 657
- solocis* Wood, *Hypothenemus* 945
- solomonicus* (Browne), (*Phloeosinus*), *Hyledius* 261
- solomonicus* (Browne), (*Poecilips*), *Coccotrypes* 612
- solomonicus* (Schedl), (*Xyleborus*), *Eucallacca* 693
- soltani* Hopkins, (*Stephanoderes*), *Hypothenemus*
(=*seriatus*) 942
- soltani* Hopkins, *Xyleborus* (= *ferrugineus*) 739
- solus* Blackman, *Pityophthorus* 1029
- solutus* Schedl, *Xyleborus* 774
- sordicauda* (Motschulsky), (*Phloeotrogus*),
Leptoxyleborus 660
- sordicaudulus* Eggers, (*Xyleborus*), *Leptoxyleborus*
(=*concisus*) 659
- sordicaudulus* peguensis Eggers, (*Xyleborus*),
Leptoxyleborus (= *concisus*) 659
- sordidus* (Blandford), (*Trippodendron*),
Indocoryphalus 649
- sordisus* (Nobuchi), *Taenioglyptes*, *Cryphalus* 595
- soussouensis* Eggers, (*Stephanoderes*),
Hypothenemus (= *mallyi*) 934
- spadix* (Blackman), (*Prionosecles*), *Scolytodes* 399
- spadix* Blackman, *Hylocurus* 425
- spadix* Blackman, *Pityophthorus* 1029
- sparsedentatus* (Schedl), (*Stephanoderes*),
Hypothenemus 945
- sparsopilosus* (Eggers), (*Poecilips*), *Coccotrypes* 612
- sparsopilosus* (Schedl), (*Gnathophorus*),
Pityophthorus 1029
- sparsopilosus* Schedl, *Cryphalus* 595
- sparsopunctatus* (Schedl), (*Neodryocoetes*),
Araptus 961
- sparserrugosus* (Schedl), (*Poecilips*), *Coccotrypes* 612
- sparsipilosus* Eggers, *Xyleborus* 774
- sparsipunctatus* (Eggers), (*Cyrtogenius*),
Xylocleptes 550
- sparsus* (LeConte), (*Xyleborus*), *Pityokteines* 464
- sparsus* Hopkins, *Hypothenemus* 945
- spartii* Nordlinger, (*Hylesinus*), *Phloeotribus*
(=*rhododactylus*) 229
- spathifer* Schedl, *Ctonoxylon* 535
- spathifer* Schedl, *Xylechinus* 115
- spathipennis* Eichhoff, *Xyleborus* 774
- spathipennis* ohansi Hagedorn, *Xyleborus*
(=*princeps*) 765
- spatulatus* Schedl, *Hypocoryphalus* 572
- spatulatus* Schedl, *Lamurgus* 409
- spatulatus* Schedl, *Stronbophorus* 204
- spatulatus* Blandford, (*Xyleborus*), *Leptoxyleborus*
(=*punctatissimus*) 660
- spatulatus* Schedl, *Tragostus* 410
- spatulatus* Wood, *Cryptocarenum* 904
- speciosus* (Schedl), (*Tricolus*), *Amphicranus* 1049
- speciosus* (Schedl), (*Xyleborus*), *Xyleborinus* 517
- speciosus* Wood, *Araptus* 962
- speciosus* Wood, *Pityophthorus* 1029
- spectabilis* Schedl, *Acacacis* (= *sp.*?) 193
- spectabilis* (Wood), (*Tricolus*), *Amphicranus* 1049
- spectus* Wood, *Amphicranus* 1049
- speculum* Bright, *Pityophthorus* 1029
- Spermatoplex* Hopkins (= *Coccotrypes*) 591
- Spermophthorus* Costa Lima 971
- spessirtseri* Eggers, *Hyurgops* 41
- spessirtseri* Lebedev, *Pityogenes* 455
- spessirtzeri* Berger, *Erioporicus* 551
- spesswzeffi* Kurenzov, *Hyurgops* (= *spessirtseri*) 41
- Sphaerosinus* Eggers (= *Phrixosoma*) 159
- Sphaerotrypes* Blandford 195
- Sphaerotrypini* Murayama (= *Diamerini*) 193
- sphaerotrypoides* Murayama, *Sucus* (= *niisimai*) 191
- spheniscus* Schedl, *Tricolus* 1045
- Sphenoceros* Schedl (= *Araptus*) 952

<i>spheuos</i> Sampson, <i>Xyleborus</i>	774	<i>spinus</i> DeGeer, (<i>Ips</i>), <i>Pityogenes</i> (= <i>chalcographus</i>)	450
<i>spiculatus</i> Browne, <i>Xyleborus</i>	775	<i>spinus</i> Eggers, <i>Strombophorus</i> (= <i>pusillus</i>)	204
<i>spicatus</i> (Wood), (<i>Cryptulocleptus</i>), <i>Pseudothysanoes</i>	417	<i>spinus</i> Hagedorn, (<i>Adiactetus</i>), <i>Hypothenemus</i> (= <i>elaphus</i>)	918
<i>spicatus</i> Browne, <i>Xyleborus</i>	775	<i>spinus</i> Hagedorn, <i>Premnobius</i> (= <i>ambitosus</i>)	651
<i>spicatus</i> Wood, <i>Phlocleptus</i>	422	spinus Wood, <i>Amphicranus</i>	1049
<i>spiculatus</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	spinus Wood, <i>Corthylus</i>	1080
<i>spiculatus</i> (Schaufuss), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	<i>Spinuloscolytus</i> Butovitch (= <i>Scolytus</i>)	321
<i>spiculatus</i> Eggers, <i>Trypophloeus</i> (= <i>binodulus</i>)	844	spinulosus (Rey), (<i>Phlocophthorus</i>), <i>Phlocotribus</i>	233
spinachius (Schedl), (<i>Steganocranus</i>), <i>Amphicranus</i>	1049	spinulosus Blandford, <i>Xyleborus</i>	775
<i>spinachius</i> Schedl, (<i>Xyleborus</i>), <i>Webbia</i> (= <i>armifer</i>)	829	spinura Wood, <i>Pseudothysanoes</i>	417
<i>spinata</i> (Schedl), (<i>Microrhynchus</i>), <i>Trischidia</i>	947	spirostachius (Schedl), (<i>Ptilopodius</i>), <i>Scolytogenes</i>	866
<i>spinatifer</i> Wood, <i>Pseudothysanoes</i>	417	splendens (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866
<i>spinatus</i> (Schedl), (<i>Bostrichips</i>), <i>Pseudothysanoes</i>	417	splendens (Wood), (<i>Loganius</i>), <i>Cnemomyx</i>	318
<i>spinatus</i> Bright, (<i>Amphicranus</i>), <i>Monarthrum</i> (= <i>gnarum</i>)	1056	<i>splendens</i> Blackman, <i>Phlocosinus</i> (= <i>fulgens</i>)	247
<i>spinatus</i> Eggers, <i>Xyleborus</i>	775	<i>splendens</i> Sokanovskii, <i>Taphrorychus</i> (= <i>rillifrons</i>)	560
<i>spinatus</i> Wood, <i>Cactopinus</i>	436	splendens Wood, <i>Amphicranus</i>	1049
<i>spinatus</i> Wood, <i>Pseudothysanoes</i> (= <i>spinatifer</i>)	417	<i>splendens</i> Wood, <i>Corthylus</i> (= <i>collaris</i>)	1070
<i>spinescens</i> Wood, <i>Amphicranus</i>	1049	splendidulus (Schedl), (<i>Neodryococtes</i>), <i>Araptus</i>	962
<i>spinibarbe</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	splendidus (Schaufuss), (<i>Xyleborus</i>), <i>Taurodennus</i>	756
<i>spinicarinatus</i> Beaver, <i>Glostatus</i>	836	<i>splendidus</i> Bright, <i>Corthylus</i> (= <i>comatus</i>)	1072
spinicollis (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	945	<i>Spongocerini</i> (= <i>Scolytolatypodini</i>)	401
<i>spinicornis</i> Schedl, <i>Xyleborus</i>	775	<i>Spongocerus</i> Blandford (= <i>Scolytolatypus</i>)	401
<i>spinideus</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	<i>Spongotarsus</i> Hagedorn (= <i>Polygraphus</i>)	283
<i>spinideus</i> (Reitter), (<i>Ips</i>), <i>Pityokteinus</i>	465	squalidens (Scudder), (<i>Hylastes</i>), <i>Phlocosinus</i>	255
<i>spinideus</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	483	<i>Squamasinus</i> Nunberg (= <i>Xylechinus</i>)	114
<i>spinideus</i> Schedl, <i>Scolytus</i>	377	squamatilis (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866
<i>spinifer</i> (Browne), (<i>Phlocosinus</i>), <i>Hyleidius</i>	261	squamatilis (Schedl), (<i>Xyleborus</i>), <i>Webbia</i>	833
<i>spinifer</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	squamatula Wood, <i>Micracisella</i>	431
<i>spinifer</i> (Eichhoff), (<i>Tomicus</i>), <i>Ips</i>	526	squamatus Wood, <i>Phlocotribus</i>	234
<i>spinifer</i> Eggers, <i>Ctenoxylon</i>	838	squameus Schedl, <i>Cryphalus</i>	895
<i>spinifer</i> Eggers, <i>Premnobius</i>	655	squameus Wood, <i>Pseudothysanoes</i>	417
<i>spinifer</i> Schwarz, <i>Corthylus</i> (= <i>papulaus</i>)	1076	squameus Yin & Huang, <i>Polygraphus</i>	299
<i>spinifex</i> Blandford, <i>Hyloterus</i>	428	squamifer (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	612
<i>spiniger</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	squamifer Wood, <i>Cnemomyx</i>	318
<i>spinipennis</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	squamiger Schedl, <i>Xylechinus</i>	119
<i>spinipennis</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	612	squamiger Wood, <i>Phlocotribus</i>	234
<i>spinipennis</i> Eggers, <i>Phlocotribus</i> (= <i>setulosus</i>)	232	squamipennis Schedl, <i>Peridryococtes</i>	562
<i>spinipennis</i> Eichhoff, (<i>Acanthotomicus</i>), <i>Diamerus</i> (= <i>curcifer</i>)	195	squamiger Chapuis, <i>Camptocerus</i> (= <i>niger</i>)	319
<i>spinipennis</i> Schedl, <i>Cryptocurus</i>	72	squamosus (Blackman), (<i>Erinosinus</i>), <i>Liparthrum</i>	279
<i>spinipennis</i> Schedl, (<i>Eidophelus</i>), <i>Xyleborinus</i> (= <i>spiniposticus</i>)	817	squamosus (Eggers), (<i>Pseudophlocotribus</i>), <i>Hylesinopsis</i>	95
<i>spinipennis</i> Schedl, <i>Webbia</i>	833	squamosus (Eggers), (<i>Troglogitica</i>), <i>Acacis</i>	193
<i>spinipes</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	squamosus (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	945
<i>spiniposticus</i> Wood, <i>Xyleborinus</i>	817	squamosus (Schedl), (<i>Blastophagus</i>), <i>Polygraphus</i>	299
<i>spinulosus</i> Schedl, <i>Xyleborus</i> (= <i>spinulosus</i>)	775	squamosus (Schedl), (<i>Lepicerinus</i>), <i>Scolytogenes</i>	866
<i>spinus</i> (Olivier), (<i>Scolytus</i>), <i>Eccoptopterus</i>	820	squamosus (Schedl), (<i>Neoglostatus</i>), <i>Pseudothysanoes</i>	417
<i>spinus</i> (Schaufuss), (<i>Xyleborus</i>), <i>Xyleborinus</i>	817	squamosus (Schedl), (<i>Phlocosinus</i>), <i>Xylechinus</i>	119
<i>spinus</i> (Schedl), (<i>Lepiceroides</i>), <i>Hypothenemus</i>	945	<i>squamosus</i> Blackman, <i>Phlocosinus</i> (= <i>sequoiae</i>)	253
<i>spinus</i> Blackman, <i>Phlocosinus</i>	255	squamosus Bright, <i>Pseudopityophthorus</i>	976
<i>spinus</i> Blandford, <i>Acanthotomicus</i>	483		
<i>spinus</i> Brethes, <i>Chramesus</i>	269		

- squamosus* Murayama, *Hylurgops* (= *spessirtseri*) . . . 41
squamosus Nunberg, *Stephanopodius* . . . 853
squamosus Schedl, *Glostatus* . . . 836
squamosus Schedl, *Ptilopodius* . . . 900
squamosus Wood, *Cnesinus* . . . 212
squamosus Yin & Huang, *Scolytus* . . . 377
squamulatus (Beaver), (*Apoxyleborus*),
Xylosandrus . . . 800
squamulatus Chapuis, *Phloeosinus* . . . 255
squamulatus Eichhoff, *Xyleborus* . . . 775
squamulatus Niisima, *Polygraphus* . . . 299
squamulatus Redtenbacher, (*Hylesinus*),
Carphoborus (= *minimus*) . . . 307
squamulatus niger Nunberg, *Xyleborus*
 (= *squamulatus*) . . . 776
squamulosus (Eggers), (*Cylindrotomicus*),
Scolytogenes . . . 866
squamulosus (Eggers), (*Xyleborus*), *Eccoptyterus* . . . 821
squamulosus Eggers, (*Ernoporus*), *Acorthylus*
 (= *bosqui*) . . . 853
squamulosus Strohmeier, *Cryphalus* . . . 895
squamulosus auratus Eggers, (*Xyleborus*),
Eccoptyterus (= *squamulosus*) . . . 821
squamulosus duplicatus Eggers, (*Xyleborus*),
Eccoptyterus (= *limbus*) . . . 820
ssiori Niisima, *Polygraphus* . . . 299
stachi Nunberg, *Hylocurus* (= *robustus*) . . . 427
starhoni Reitter, *Dryocoetes* (= *villosus*) . . . 587
starki Eggers, (*Blastophagus*), *Tomicus* (= *puellus*) . . . 136
starki Eggers, (*Hylurgops*), *Hylastes*
 (= *cunicularius*) . . . 53
starki Eggers, *Ernoporus* (= *tiliae*) . . . 857
starki Kurenzov, *Scolytus* (= *japonicus*) . . . 335
starki Nunberg, *Xyleborus* . . . 776
starki Spessivtzev, *Orthotomicus* . . . 475
stebbingi Strohmeier, *Ips* . . . 526
Steganocranus Eichhoff (= *Amphicranus*) . . . 1045
Stegomerus Wood . . . 852
Stenoclyptus Blackman . . . 419
stenodermus (Schedl), (*Pterocyclon*),
Amphicranus . . . 1049
stenographus (Schedl), (*Xyleborus*),
Cryptoxyleborus . . . 829
stenographus Duftschmidt, (*Bostrichus*), *Ips* (= *sex-*
dentatus) . . . 525
stenographus Schedl, *Xyloctonus* (= *aethiops*) . . . 840
Stephanoderes Eichhoff (= *Hypothenemus*) . . . 904
Stephanopodius Schedl . . . 853
Stephanorhopalus Hopkins (= *Ernoporus*) . . . 854
stephegynis Hopkins, *Ptilopodius* . . . 900
stephegynis Hopkins, *Xyleborus* (= *bidentatus*) . . . 715
sterculiae Hopkins, (*Stephanoderes*), *Hypothenemus*
 (= *birmanus*) . . . 910
Sternobothrus Eggers . . . 216
stierlini Eggers, (*Cryphalus*), *Scolytogenes*
 (= *aspericollis*) . . . 859
stigmatosus (Schedl), (*Stephanoderes*),
Hypothenemus . . . 945
stoeckleini Schedl, *Phloeosinus* (= *henschi*) . . . 248
streblicola Hopkins, *Euwallacea* . . . 693
Streptocranus Schedl (= *Coptoborus*) . . . 662
striata (Eggers), (*Sphaerosinus*), *Phrixosoma* . . . 190
striata Wood, *Micracisella* . . . 431
striatopunctatus Lea, (*Cryphalus*), *Hypothenemus*
 (= *eruditus*) . . . 921
striatotruncatus (Schedl), (*Xyleborus*), *Amasa* . . . 654
striatulus (Mannerheim), (*Cryphalus*),
Trypophloeus . . . 847
striatulus (Schedl), (*Hyorrhynchus*), *Sucus* . . . 191
striatulus Browne, *Cryphalus* . . . 895
striatulus Browne, (*Cryphalomorphus*),
Hypothenemus (= *brounei*) . . . 911
striatulus Browne, *Xyleborus* . . . 776
striatulus Hagedorn, *Camptocerus* (= *niger*) . . . 319
striatulus Schedl, *Hypothenemus* (= *seriatus*) . . . 943
striatulus Wood, *Coccotrypes* (= *tahitensis*) . . . 613
striatulus Wood, *Scolytodes* . . . 399
striatus (Eggers), (*Hexacolus*), *Scolytodes* . . . 399
striatus (Eggers), (*Phthorophloeus*), *Phloeotribus* . . . 234
striatus (LeConte), (*Cryphalus*), *Dendroterus* . . . 952
striatus (Schedl), (*Metahylesinus*), *Hylesinopsis* . . . 95
striatus Browne, *Cryphalus* . . . 895
striatus Eggers, (*Thaumurgides*), *Coccotrypes*
 (= *tahitensis*) . . . 613
striatus Eggers, *Bothrostermus* . . . 215
striatus Eggers, *Chramesus* . . . 269
striatus Eggers, *Coccotrypes* . . . 612
striatus Eggers, *Cryphalominus* . . . 835
striatus Eggers, *Diamerus* . . . 198
striatus Eggers, *Dryocoetes* . . . 555
striatus Eggers, (*Problechilus*), *Gymnochilus*
 (= *reitteri*) . . . 387
striatus Eggers, *Hylesinus* (= *laticollis*) . . . 81
striatus Eggers, *Phloeotribus* (= *neglectus*) . . . 225
striatus Eggers, *Pseudodiamerus* . . . 194
striatus Eggers, (*Hylocurosoma*), *Scolytodes*
 (= *striatulus*) . . . 399
striatus Eggers, *Trietennus* . . . 541
striatus Eggers, *Xyloctonus* . . . 841
striatus Hopkins, *Hypocryphalus* . . . 872
striatus Nobuchi, *Taphrorychus* . . . 560
striatus Wood, *Chramesus* (= *strigatus*) . . . 269
strictus Wood, *Pityophthorus* . . . 1029
strigatus Wood, *Chramesus* . . . 269
strigicollis LeConte, *Cnesinus* . . . 212
strigicollis Schedl, *Coccotrypes* . . . 612
strigilatus Eggers, *Corthylus* (= *letzneri*) . . . 1074
strigilatus Reitter, *Scolytus* (= *mali*) . . . 341
strigilatus Wichmann, *Cryphalus* . . . 895
strigilis Wood, *Chramesus* . . . 269
strigilis Wood, *Corthylus* . . . 1080
strigillatus Chapuis, (*Nemophilus*), *Cnesinus*
 (= *strigicollis*) . . . 212
strigipennis Schedl, *Cryphalus* . . . 895
strigipennis Schedl, *Scolytus* . . . 377

<i>strohmeyeri</i> Eggers, <i>Scolytoplatus</i> (= <i>fasciatus</i>)	403	<i>subcostulatus subcostulatus</i> (Mannerheim).	
<i>strohmeyeri</i> Schedl. (<i>Xyleborus</i>), <i>Ambrosiodmus</i>		<i>Hylurgops</i> (= <i>subcostulatus</i>)	42
(= <i>rubricollis</i>)	679	<i>subcrenulatus</i> Eggers, <i>Xyleborus</i>	776
<i>strohmeyeri</i> Stebbing, <i>Cryphalus</i>	895	<i>subcristatus</i> Schedl. <i>Pityophthorus</i>	1030
<i>strombiformis</i> (Schedl.) (<i>Xyleborus</i>), <i>Eucallacca</i>	693	<i>subcristosus</i> (Blandford) (<i>Xyleborus</i>),	
<i>Strombophorini</i> Schedl (= <i>Diamerini</i>)	193	<i>Coccotrypes</i>	612
<i>Strombophorus</i> Hagedorn	202	<i>subcristosus</i> (Eggers), (<i>Hexacolus</i>), <i>Scolytodes</i>	399
<i>strombosiopsis</i> Schedl. <i>Dendrochilus</i>	435	<i>subcristosus</i> Eggers, <i>Crypturgus</i> (= <i>cinereus</i>)	623
<i>strombosiopsis</i> Schedl. <i>Xyleborus</i>	776	<i>subcylindricus</i> (Schedl.) (<i>Pocilips</i>), <i>Coccotrypes</i>	613
<i>Strophionocerus</i> Sampson, <i>Scolytoplatus</i>	401	<i>subcylindricus</i> Eggers, (<i>Stephanoderes</i>),	
<i>struthanthi</i> Wood, <i>Chaetophloeus</i>	274	<i>Hypothenemus</i> (= <i>eruditus</i>)	924
<i>stinuosus</i> Schedl. <i>Xylosandrus</i> (= <i>curtulus</i>)	793	<i>subcylindricus</i> Schedl. (<i>Cryphalus</i>), <i>Hypocryphalus</i>	
<i>Stylotenus</i> Schedl (= <i>Hypothenemus</i>)	905	(= <i>mangiferae</i>)	870
<i>Styphlosoma</i> Blandford	950	<i>subdentatus</i> Browne, <i>Xyleborus</i>	776
<i>Styracopterus</i> Blandford (= <i>Styracoptinus</i>)	281	<i>subdentatus</i> Browne, <i>Xyleborus</i>	776
<i>Styracoptinus</i> Wood	281	<i>subdepressus</i> Eggers, (<i>Dryococtes</i>), <i>Coccotrypes</i>	
<i>styrax</i> (Schedl.) (<i>Stephanoderes</i>), <i>Hypothenemus</i>	945	(= <i>cyperi</i>)	599
<i>suavi</i> (Schedl.) (<i>Pocilips</i>), <i>Coccotrypes</i>	612	<i>subdepressus</i> Eggers, <i>Cryphalus</i> (= <i>piceae</i>)	889
<i>suavi</i> Schedl. <i>Xyleborus</i>	776	<i>subdepressus</i> Rex, (<i>Xyleborus</i>), <i>Xyleborinus</i>	
<i>subacuminatus</i> (Eggers), (<i>Pocilips</i>), <i>Coccotrypes</i>	612	(= <i>saxseni</i>)	814
<i>subacuminatus</i> (Schedl.) (<i>Pelicerus</i>), <i>Cyrtogenius</i>	568	<i>subdepressus</i> Schedl. <i>Coccotrypes</i>	612
<i>subacuminatus</i> (Schedl.) (<i>Stephanoderes</i>),		<i>subdepressus</i> Schedl. <i>Lamargus</i>	409
<i>Hypothenemus</i>	945	<i>subdolosus</i> Schedl. <i>Xyleborus</i>	777
<i>subadjunctus</i> Schedl. (<i>Xyleborus</i>), <i>Coptoborus</i>		<i>subductum</i> (Schedl.) (<i>Pterocyclon</i>), <i>Monarthrum</i>	1063
(= <i>usagaricus</i>)	664	<i>subductus</i> Schedl. <i>Xyleborus</i>	777
<i>subaffinis</i> Eggers, <i>Xyleborus</i> (= <i>affinis</i>)	709	<i>subelongatus</i> Hopkins, <i>Hypothenemus</i>	
<i>subagnatum</i> Schedl. <i>Cyclorhipidion</i>		(= <i>pubescens</i>)	939
(= <i>subagnatum</i>)	703	<i>subelongatus</i> Motschulsky, (<i>Tomicus</i>), <i>Ips</i>	
<i>subagnatum</i> Wood, <i>Cyclorhipidion</i>	703	(= <i>cembrae</i>)	499
<i>subagnatus</i> Eggers, <i>Cyclorhipidion</i> (= <i>subagnatum</i>)	703	<i>submarginatus</i> Eggers, <i>Xyleborus</i>	777
<i>subagnatus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i>		<i>subglabratus</i> Schedl. <i>Hypothenemus</i> (= <i>arecae</i>)	908
(= <i>javanus</i>)	933	<i>subglabrum</i> Isackson, <i>Aphanarthrum</i>	620
<i>subalpinus</i> Eggers, <i>Hylastes</i> (= <i>cunicularius</i>)	51	<i>subgranosus</i> (Schedl.) (<i>Eidophelus</i>), <i>Cyrtogenius</i>	569
<i>subaplanatus</i> Schedl. (<i>Pocilips</i>), <i>Coccotrypes</i>		<i>subgranosus</i> Schedl. <i>Xyleborus</i>	777
(= <i>cyperi</i>)	599	<i>subgranosus affinis</i> Schedl. <i>Cyrtogenius</i>	
<i>subasperulus</i> Eggers, <i>Corthylus</i>	1080	(= <i>subgranosus</i>)	569
<i>subasperulus</i> Eggers, <i>Xyleborus</i>	776	<i>subgranulatum</i> Wood, <i>Monarthrum</i>	1063
<i>subcarinatum</i> Wood, <i>Pycnarthrum</i>	356	<i>subgranulatus</i> (Eggers), (<i>Xyleborus</i>),	
<i>subcarinatus</i> (Schedl.) (<i>Phloeosinus</i>), <i>Hyledius</i>	261	<i>Xyleborinus</i>	817
<i>subcarinatus</i> Eggers, <i>Xyleborus</i>	776	<i>subgranulatus</i> Schedl. <i>Cryphalus</i>	895
<i>subcompactus</i> Lea, <i>Cryphalus</i>	895	<i>subgranulatus</i> Schedl. <i>Hylocurus</i>	428
<i>subconcentricus</i> Hopkins, (<i>Stephanoderes</i>),		<i>subimmunitus</i> (Schedl.) (<i>Minips</i>),	
<i>Hypothenemus</i> (= <i>eruditus</i>)	923	<i>Acanthotomicus</i>	484
<i>subconcentricus</i> Hopkins, <i>Cryphalus</i> (= <i>pubescens</i>)	890	<i>subimpressus</i> Bright, <i>Pityophthorus</i> (= <i>subsimilis</i>)	1030
<i>subconcentricus</i> Schedl. <i>Pityophthorus</i>	1029	<i>subimpressus</i> Eggers, (<i>Dryococtes</i>), <i>Coccotrypes</i>	
<i>subconfusus</i> Eggers, <i>Scolytus</i> (= <i>japonicus</i>)	335	(= <i>cyperi</i>)	599
<i>subcoriaceus</i> Eggers, <i>Eucallacca</i> (= <i>piceus</i>)	693	<i>subincisuralis</i> Schedl. <i>Tricolus</i>	1045
<i>subcostatus</i> Eggers, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	91	<i>subitus</i> Schedl. <i>Xyleborus</i> (= <i>ferrugineus</i>)	739
<i>subcostatus</i> Eggers, <i>Phloeotribus</i>	234	<i>sublaevis</i> (Eggers), (<i>Webbia</i>), <i>Arixyleborus</i>	669
<i>subcostatus</i> Eggers, <i>Xylocotonus</i>	841	<i>sublaevis</i> Eggers, <i>Coccotrypes</i> (= <i>sannio</i>)	611
<i>subcostatus</i> Eichhoff, <i>Xyleborus</i>	776	<i>sublinearis</i> Eggers, <i>Xyleborus</i>	777
<i>subcostatus dearnatus</i> Eggers, <i>Xyleborus</i>		<i>sublongus</i> Eggers, <i>Coccotrypes</i> (= <i>carpophagus</i>)	594
(= <i>subcostatus</i>)	776	<i>sublongus</i> Eggers, <i>Xyleborus</i>	777
<i>subcostulatus</i> (Mannerheim), (<i>Hylastes</i>),		<i>submarginatus</i> Blandford, <i>Xyleborus</i> (= <i>similis</i>)	773
<i>Hylurgops</i>	42	<i>submarginatus</i> Schedl. <i>Scolytus</i>	377
<i>subcostulatus alternans</i> Chapuis, (<i>Hylastes</i>),		<i>submolestus</i> Schedl. <i>Xyleborus</i>	777
<i>Hylurgops</i> (= <i>subcostulatus</i>)	42	<i>submontanus</i> Schedl. <i>Xyleborus</i> (= <i>tenellus</i>)	778
		<i>submuricatus</i> Eichhoff, <i>Cryphalus</i>	895

<i>subnaevis</i> Schedl, <i>Cryptoxyleborus</i>	529	<i>subsulcatus</i> Eggers, <i>Diamerus</i> (= <i>interstitialis</i>)	196
<i>subnepotulus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	650	<i>subsulcatus</i> Schedl, <i>Coccotrypes</i>	613
<i>subnitida</i> Blackman, <i>Micracisella</i>	431	<i>subsulcatus</i> Schedl, <i>Corthylus</i>	1080
<i>subnitidus</i> Schedl, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>advena</i>)	593	<i>subsulcatus</i> Schedl, <i>Polygraphus</i>	301
<i>subobtusum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	<i>subsulcatus</i> Wood, <i>Thysanoes</i>	421
<i>subopacicollis</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	943	<i>subtectus</i> Browne, <i>Sphaerotrypes</i>	201
<i>subopacus</i> Blackman, <i>Hylastes</i>	62	<i>subtilis</i> Schedl, <i>Xyleborus</i>	777
<i>subopacus</i> Blackman, <i>Pityophthorus</i>	1030	<i>subtriatus</i> Schedl, (<i>Cryphalomorphus</i>), <i>Scolytogenes</i> (= <i>knabi</i>)	863
<i>subopacus</i> Eggers, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>despectus</i>)	91	<i>subtruncatum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1063
<i>subopacus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>obscurus</i>)	936	<i>subtruncatus</i> Schedl, <i>Xyleborus</i>	777
<i>subopacus</i> Schaeffer, <i>Chramesus</i>	269	<i>subtuberculatus</i> Eggers, (<i>Pocillips</i>), <i>Coccotrypes</i> (= <i>rotundicollis</i>)	610
<i>subopacus</i> Schedl, <i>Bothrostermus</i>	215	<i>subtuberculatus</i> Eggers, <i>Chramesus</i>	269
<i>subopacus</i> Schedl, <i>Microcorthylus</i> (= <i>glabratus</i>)	1065	<i>subtuberculatus</i> Eggers, <i>Xyleborus</i>	777
<i>subopacus</i> Thomson, <i>Polygraphus</i>	300	<i>subtuberculatus</i> Schedl, <i>Cryphalus</i>	596
<i>subopacus minor</i> Lindemann, <i>Polygraphus</i> (= <i>subopacus</i>)	301	<i>subuculae</i> (Browne), (<i>Proncebbia</i>), <i>Webbia</i>	533
<i>suboralis</i> Eggers, <i>Coccotrypes</i>	613	<i>subulatum</i> Wood, <i>Styphlosoma</i>	950
<i>suboratus</i> Blandford, <i>Phloeotribus</i>	234	<i>subulatus</i> (Wood), (<i>Cryptoleptus</i>), <i>Pseudothysanoes</i>	415
<i>subparallelus</i> (Eggers), (<i>Xyleborus</i>), <i>Euallacea</i>	694	<i>subvestitus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>arecae</i>)	907
<i>subparallelus</i> Eggers, (<i>Hexacolus</i>), <i>Scolytodes</i> (= <i>notatus</i>)	396	<i>subvestitus</i> Schedl, <i>Cryphalus</i>	596
<i>subpilosus</i> (Wood), (<i>Cryptocleptes</i>), <i>Pseudothysanoes</i>	417	<i>subvulgaris</i> (Browne), (<i>Pocillips</i>), <i>Coccotrypes</i>	613
<i>subplanatus</i> Eggers, <i>Xyleborus</i>	777	<i>succicus</i> Eggers, <i>Dryocoetes</i>	555
<i>subpronum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1063	<i>Sueinae</i> Murayama (= <i>Hyorrhynchini</i>)	191
<i>subpruinosis</i> Browne, <i>Xyleborus</i>	777	<i>Sueus</i> Murayama	191
<i>subretusus</i> Wollaston, <i>Triotennus</i>	541	<i>suggrandis</i> Schedl, <i>Microcorthylus</i>	1067
<i>subscaber</i> LeConte, <i>Scolytus</i>	377	<i>sulcaticeps</i> Schedl, <i>Xyleborus</i>	777
<i>subseriatus</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopsclion</i>	97	<i>sulcatulus</i> Eggers, <i>Xyleborus</i>	775
<i>subseriatus</i> Eggers, <i>Coccotrypes</i> (= <i>carpophagus</i>)	594	<i>sulcatum</i> (Blandford), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1063
<i>subserriatus</i> Wood, <i>Corthylus</i>	1050	<i>sulcatus</i> (Bruck), (<i>Pseudothysanoes</i>), <i>Stenoclyptus</i>	419
<i>subsimilans</i> Wood, <i>Pityophthorus</i>	1030	<i>sulcatus</i> (Eggers), (<i>Metalylesinus</i>), <i>Hylesinopsis</i>	95
<i>subsimiliformis</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800	<i>sulcatus</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	650
<i>subsimilis</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	800	<i>sulcatus</i> (LeConte), (<i>Cryphalus</i>), <i>Gnathotrichus</i>	1035
<i>subsimilis</i> Schedl, (<i>Breviophthorus</i>), <i>Pityophthorus</i> (= <i>subsimilans</i>)	1030	<i>sulcatus</i> (Nunberg), (<i>Neodryocoetes</i>), <i>Araptus</i>	962
<i>subsimilis</i> Schedl, <i>Arixyleborus</i> (= <i>canaliculatus</i>)	665	<i>sulcatus</i> (Schedl), (<i>Eidophelus</i>), <i>Cyrtogenius</i>	569
<i>subsimilis</i> Schedl, <i>Pityophthorus</i>	1030	<i>sulcatus</i> (Wood), (<i>Renocis</i>), <i>Chaetophloeus</i>	274
<i>subspinosis</i> Eggers, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>saxeseni</i>)	515	<i>sulcatus</i> Bright, <i>Pityophthorus</i> (= <i>immanis</i>)	1002
<i>substriatus</i> Schedl, (<i>Eulytocerus</i>), <i>Phloeotribus</i> (= <i>championi</i>)	219	<i>sulcatus</i> Bright, <i>Sampsonius</i>	656
<i>substriatus</i> Schedl, <i>Cryphalus</i>	896	<i>sulcatus</i> Browne, <i>Sphaerotrypes</i>	201
<i>substriatus</i> Strohmeier, <i>Hylastes</i>	62	<i>sulcatus</i> Eggers, <i>Chortastus</i> (= <i>minimus</i>)	312
<i>substrigatus</i> Blackman, <i>Cnesinus</i> (= <i>gracilis</i>)	205	<i>sulcatus</i> Eggers, <i>Cucinus</i>	212
<i>subsulcatus</i> (Eggers), (<i>Xyleborus</i>), <i>Xyleborinus</i>	515	<i>sulcatus</i> Eggers, <i>Hyburgops</i>	42
<i>subsulcatus</i> (Schedl), (<i>Breviophthorus</i>), <i>Pityophthorus</i>	1030	<i>sulcatus</i> LeConte, <i>Scolytus</i> (= <i>mali</i>)	344
<i>subsulcatus</i> (Schedl), (<i>Eidophelus</i>), <i>Cyrtogenius</i>	569	<i>sulcatus</i> Marsham, (<i>Ips</i>), <i>Hylesinus</i> (= <i>crenatus</i>)	77
<i>subsulcatus</i> Browne, <i>Lamurgus</i>	409	<i>sulcatus</i> Schedl, (<i>Breviophthorus</i>), <i>Araptus</i> (= <i>cribricollis</i>)	955
		<i>sulcatus</i> Schedl, <i>Polygraphus</i>	301
		<i>sulcatus</i> Schedl, <i>Xylechinus</i>	119
		<i>sulcicauda</i> Schedl, <i>Xyleborus</i>	775
		<i>sulcifrons</i> Chapuis, <i>Phloeotribus</i>	234
		<i>sulcifrons</i> Eichhoff, <i>Phloeoborus</i>	101
		<i>sulcifrons</i> Rey, <i>Scolytus</i>	377
		<i>sulcifrons</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517

<i>sulcinodis</i> (Schedl), (<i>Hylesinus</i>), <i>Ficicis</i>	92	<i>suturalis</i> Wood, <i>Scolytodes</i>	399
<i>sulcinoides</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	<i>suturifer</i> Schedl, <i>Corthylus</i>	1080
<i>sulciperne</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1063	<i>swainei</i> Blackman, <i>Micracis</i>	433
<i>sulciperne</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	<i>swainei</i> Blackman, <i>Pityophthorus</i> (= <i>lautus</i>)	1006
<i>sulciperne</i> Schedl, (<i>Ozopenon</i>), <i>Coccotrypes</i> (= <i>gedeanus</i>)	603	<i>swainei</i> Bruck, <i>Carphoborus</i> (= <i>simplex</i>)	310
<i>sumatрана</i> (Hagedorn), (<i>Xyleborus</i>), <i>Schedlia</i>	822	<i>swainei</i> Bruck, <i>Phloeosinus</i>	256
<i>sumatranus</i> (Blandford), (<i>Dryocoetes</i>), <i>Ozopenon</i>	590	<i>swainei</i> Drake, (<i>Anisandrus</i>), <i>Xyleborus</i> (= <i>dispar</i>)	730
<i>sumatranus</i> (Eggers), (<i>Phloeosinus</i>), <i>Hyledius</i>	261	<i>swainei</i> Eggers, <i>Hylastes</i> (= <i>porculus</i>)	61
<i>sumatranus</i> (Schedl), (<i>Chiloxylon</i>), <i>Scolytogenes</i>	866	<i>swainei</i> R. Hopping, <i>Ips</i> (= <i>borealis</i>)	493
<i>sumatranus</i> (Schedl), (<i>Dacryphalus</i>), <i>Hypocryphalus</i>	872	<i>swazezi</i> Beeson, <i>Xyleborus</i>	778
<i>sumatranus</i> Eggers, <i>Coccotrypes</i> (= <i>papuanus</i>)	608	<i>swazezi</i> Schedl, <i>Cryphalus</i> (= <i>sylicicola</i>)	896
<i>sumatranus</i> Eggers, (<i>Hylesinus</i>), <i>Ficicis</i> (= <i>bakeri</i>)	90	<i>swieteniae</i> (Blackman), (<i>Hexacolus</i>), <i>Scolytodes</i>	399
<i>sumatranus</i> Eggers, (<i>Hylurgulus</i>), <i>Hyledius</i> (= <i>hylurgulus</i>)	260	<i>sylicicola</i> (Perkins), (<i>Hypothenemus</i>), <i>Cryphalus</i>	896
<i>sumatranus</i> Schedl, <i>Eidophelus</i>	867	<i>sylicicola</i> obliquus Schedl, <i>Cryphalus</i> (= <i>sylicicola</i>)	896
<i>sumatranus</i> Schedl, (<i>Erinosinus</i>), <i>Cryptophlyphthorus</i> (= <i>eggersi</i>)	274	<i>symphoniae</i> Browne, <i>Polygraphus</i> (= <i>majusculus</i>)	290
<i>sumatranus</i> Schedl, <i>Polygraphus</i>	301	<i>syrtuschecki</i> (Wichmann), (<i>Dendrosinus</i>), <i>Acacis</i>	194
<i>sumatranus</i> Strohmeier, <i>Acanthotomicus</i>	484	<i>syzygii</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	818
<i>sundaensis</i> Eggers, (<i>Dendrugus</i>), <i>Coccotrypes</i> (= <i>myristicae</i>)	606	<i>syzygii</i> Numberg, <i>Xyleborus</i> (= <i>tenellus</i>)	778
<i>sundaensis</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>fuscicollis</i>)	926	<i>szechuanensis</i> Tsai & Li, <i>Cryphalus</i>	896
<i>sundaensis</i> Eggers, <i>Xyleborus</i>	778	<i>szechuanensis</i> <i>teichangensis</i> Tsai & Li, <i>Cryphalus</i> (= <i>szechuanensis</i>)	896
<i>sundaensis</i> Schedl, <i>Cryphalus</i>	896	<i>szemaoensis</i> Tsai & Yin, <i>Polygraphus</i>	301
<i>sundri</i> Schedl, (<i>Margadillius</i>), <i>Ernoeladius</i> (= <i>corpulentus</i>)	557	<i>szentivanyi</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>colossus</i>)	672
<i>superbulus</i> (Schedl), (<i>Xyleborus</i>), <i>Coptoborus</i>	664	<i>taboensis</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>rubicollis</i>)	679
<i>superbus</i> (Schedl), (<i>Xyleborus</i>), <i>Coptoborus</i>	664	<i>tabogae</i> (Blackman), (<i>Neodryocoetes</i>), <i>Araptus</i>	962
<i>superbus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>superbulus</i>)	664	<i>tabulaeformis</i> Tsai & Li, <i>Cryphalus</i>	896
<i>supercilliosus</i> Tsai & Huang, <i>Scolytoplatypus</i>	407	<i>tabulaeformis</i> <i>chienlungensis</i> Tsai & Li, <i>Cryphalus</i> (= <i>tabulaeformis</i>)	896
<i>surinamensis</i> Schedl, <i>Coccotrypes</i>	613	<i>Tachyderes</i> Blackman (= <i>Cryptocarcinus</i>)	902
<i>surinamensis</i> Schedl, <i>Pityophthorus</i>	1030	<i>tachygraphus</i> (Zimmermann), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	680
<i>sus</i> (Schedl), (<i>Xyleborus</i>), <i>Arixyleborus</i>	669	<i>tachygraphus</i> Sahlberg, (<i>Bostrichus</i>), <i>Xyleborus</i> (= <i>dispar</i>)	729
<i>suspectus</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	484	<i>tacubayae</i> Hopkins, <i>Phloeosinus</i>	256
<i>suspectus</i> Bright, <i>Hylastes</i>	62	<i>tadzhikistanicus</i> Stark, <i>Scolytus</i>	378
<i>suspectus</i> Wood, <i>Hypothenemus</i>	946	<i>taedae</i> Hopkins, <i>Conophthorus</i> (= <i>coniperda</i>)	964
<i>suspicious</i> Bright, <i>Pityophthorus</i>	1030	<i>Taeniocerus</i> Blandford (= <i>Scolytoplatypus</i>)	401
<i>sutirifer</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	484	<i>Taenioceri</i> Blandford (= <i>Scolytoplatypodini</i>)	401
<i>suturale</i> Eggers, <i>Trypodendron</i> (= <i>signatum</i>)	648	<i>Taenioglyptes</i> Bedel (= <i>Cryphalus</i>)	572
<i>suturalis</i> (Eggers), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	669	<i>tahitensis</i> (Beeson), (<i>Thamniurgides</i>), <i>Coccotrypes</i>	613
<i>suturalis</i> (Eggers), (<i>Xyleborus</i>), <i>Cnestus</i>	803	<i>taichuensis</i> Schedl, <i>Xyleborus</i>	778
<i>suturalis</i> (Fabricius), (<i>Hylesinus</i>), <i>Camptocerus</i>	320	<i>taihokuensis</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>suturalis</i> (Gyllenhal), (<i>Bostrichus</i>), <i>Orthotomicus</i>	476	<i>taireiensis</i> Murayama, <i>Carphoborus</i>	310
<i>suturalis</i> (Schedl), (<i>Hapalogenius</i>), <i>Rhopalopselion</i>	97	<i>taitonus</i> Eggers, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>mutilatus</i>)	799
<i>suturalis</i> Browne, <i>Cyrtogenius</i>	569	<i>taivanensis</i> Browne, <i>Xyleborus</i>	778
<i>suturalis</i> Browne, <i>Webbia</i>	833	<i>taivanensis</i> Schedl, <i>Polygraphus</i>	301
<i>suturalis</i> Eggers, <i>Bothrostermus</i>	215	<i>taivanus</i> Schedl, <i>Cryphalus</i>	896
<i>suturalis</i> Eggers, <i>Corthylus</i>	1080	<i>takahashii</i> (Browne), (<i>Phloeosinus</i>), <i>Hyledius</i>	261
<i>suturalis</i> Eggers, <i>Phloeotribus</i>	234	<i>takahashii</i> Nobuchi, <i>Ernoporicus</i>	852
<i>suturalis</i> Eggers, <i>Pityophthorus</i>	1030	<i>takeharai</i> Browne, <i>Xyleborus</i>	778
<i>suturalis</i> LeConte, <i>Micracis</i>	433	<i>takinoyensis</i> Murayama, <i>Xyleborus</i>	778
<i>suturalis</i> Redtenbacher, <i>Hylesinus</i> (= <i>toranio</i>)	85	<i>talauricus</i> (Eggers), (<i>Xyleboricus</i>), <i>Arixyleborus</i>	669

<i>tatunai</i> (Browne), (<i>Xyleborus</i>), <i>Eucallacea</i>	694	<i>tecleae</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703
<i>talyphensis</i> Michalski, <i>Scolytus</i> (= <i>ensifer</i>)	330	<i>tectonae</i> Nunberg, <i>Xyleborus</i> (= <i>hagedornianus</i>)	742
<i>tamarindi</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942	<i>tectonae</i> Stehling, (<i>Cryphalus</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	921
<i>tanganikaensis</i> Schedl, <i>Xyleborus</i> (= <i>amanicus</i>)	710	<i>tectus</i> Beeson, <i>Sphaerotrypes</i> (= <i>querci</i>)	201
<i>tanganus</i> Eggers, <i>Coccotrypes</i> (= <i>ductyliperdu</i>)	602	<i>tectus</i> Eggers, <i>Campococcus</i> (= <i>orientalis</i>)	320
<i>tanganus</i> Hagedorn, <i>Xyleborus</i> (= <i>ferrugineus</i>)	739	<i>tectus</i> Schedl, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>declivispinatus</i>)	672
<i>tanganus</i> Schaufuss, <i>Sphaerotrypes</i>	201	<i>tegalensis</i> Eggers, (<i>Xyleborus</i>), <i>Ambrosiodmus</i> (= <i>levisi</i>)	676
<i>tanganikaensis</i> (Schedl), (<i>Ips</i>), <i>Acanthotomicus</i>	484	<i>teichangensis</i> Tsai & Li, <i>Cryphalus</i> (= <i>szechuanensis</i>)	896
<i>tanganikaensis</i> Schedl, <i>Hypoborus</i>	251	<i>temetiucius</i> Beeson, <i>Xyleborus</i>	778
<i>tanibe</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703	<i>teminabani</i> Browne, <i>Xyleborus</i>	778
<i>tantalus</i> Schedl, <i>Xyleborus</i> (= <i>obliquus</i>)	756	<i>tenax</i> Wood, <i>Pityophthorus</i>	1030
<i>tantillus</i> Lea, (<i>Cryphalus</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	921	<i>tenebrosus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	681
<i>tanzanicus</i> Browne, <i>Polygraphus</i> (= <i>granulatus</i>)	287	<i>tenebrosus</i> Sahlberg, <i>Hylurgops</i> (= <i>glabratus</i>)	33
<i>tapatapaoanus</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	613	<i>tenella</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	827
<i>tapatapaoensis</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>fornicatus</i>)	690	<i>tenellus</i> (Schedl), (<i>Neodryocotes</i>), <i>Araptus</i>	962
<i>Taphramites</i> Schedl	1082	<i>tenellus</i> Schedl, <i>Xyleborus</i>	778
<i>Taphroborus</i> Nunberg (= <i>Cyrtogenius</i>)	562	<i>tenellus</i> Wood, <i>Pseudothysanoes</i>	418
<i>Taphrocotes</i> Pfeffer (= <i>Taphrorychus</i>)	557	<i>tenerrimus</i> Sahlberg, (<i>Hylesinus</i>), <i>Crypturgus</i> (= <i>cinereus</i>)	623
<i>Taphrodasus</i> Wood	827	<i>teninabani</i> Browne, <i>Xyleborus</i>	778
<i>Taphroungus</i> Reitter	546	<i>tenuigratum</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703
<i>Taphropterus</i> Schedl (= <i>Hemicryphalus</i>)	866	<i>tenipennis</i> Browne, <i>Xyleborus</i> (= <i>sulcicauda</i>)	778
<i>Taphrorychina</i> Reitter (= <i>Dryocoetini</i>)	539	<i>tenipennis</i> Schedl, <i>Polygraphus</i>	301
<i>Taphrorychini</i> Reitter (= <i>Dryocoetini</i>)	539	<i>tenius</i> (Blackman), (<i>Neodryocotes</i>), <i>Araptus</i>	962
<i>Taphrorychinina</i> Reitter (= <i>Dryocoetini</i>)	539	<i>tenius</i> (Schedl), (<i>Carposinus</i>), <i>Cyrtogenius</i>	569
<i>Taphrorychus</i> Eichhoff	556	<i>tenius</i> (Schedl), (<i>Minips</i>), <i>Acanthotomicus</i>	484
<i>Taphrostenosis</i> Schedl (= <i>Cyrtogenius</i>)	562	<i>tenius</i> (Wood), (<i>Hexacolus</i>), <i>Scolytodes</i>	399
<i>Taphroterus</i> Schedl (= <i>Taphrorychus</i>)	557	<i>tenius</i> Blandford, <i>Amphicranus</i>	1049
<i>taprobanus</i> (Blandford), (<i>Dryocotes</i>), <i>Coccotrypes</i>	613	<i>tenius</i> Browne, <i>Scolytomimus</i>	840
<i>taradakeensis</i> (Murayama), (<i>Pseudopocillips</i>), <i>Taphrorychus</i>	560	<i>tenius</i> Eichhoff, <i>Hylastes</i>	62
<i>tarawai</i> Beaver, <i>Ptilopodius</i>	900	<i>tennis</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922
<i>tardulus</i> Wood, <i>Corthylus</i> (= <i>neotardus</i>)	1075	<i>tenus</i> Schedl, (<i>Xyleborus</i>), <i>Coptoborus</i> (= <i>pseudotenius</i>)	664
<i>tardulus</i> Wood, <i>Dendrocraniulus</i>	553	<i>tenus</i> Schedl, <i>Cryphalus</i>	897
<i>tardum</i> Wood, <i>Corthyccyon</i>	1068	<i>tenus</i> Schedl, <i>Polygraphus</i>	301
<i>tardus</i> Schedl, <i>Corthylus</i> (= <i>neotardus</i>)	1075	<i>tenus</i> Swaine, <i>Pityophthorus</i> (= <i>murrayanae</i>)	1014
<i>tardus</i> Schedl, <i>Dendrocraniulus</i>	553	<i>tenus</i> Wood, <i>Pseudopityophthorus</i>	976
<i>tardus</i> Wood, <i>Scolytodes</i>	399	<i>teocotum</i> Wood, <i>Conophthorus</i>	968
<i>tarsalis</i> Forster, (<i>Hylesinus</i>), <i>Phloeotribus</i> (<i>rhododactylus</i>)	229	<i>teplouchovi</i> Spessivtsev, <i>Carphoborus</i>	310
<i>tarsalis</i> Schedl, <i>Eccoptopterus</i>	822	<i>terebella</i> Blandford, <i>Amphicranus</i>	1049
<i>tamayii</i> Eggers, (<i>Xylechinus</i>), <i>Xylechinomimus</i> (= <i>contractus</i>)	105	<i>terebrans</i> (Olivier), (<i>Scolytus</i>), <i>Dendroctonus</i>	181
<i>tauricus</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>jaroschewskii</i>)	336	<i>terebrans</i> Schedl, <i>Pityophthorus</i>	1031
<i>Taurodemus</i> Wood	785	<i>terebratus</i> (Wood), (<i>Gnathotrypanus</i>), <i>Gnathotrypes</i>	1042
<i>taxicola</i> Lezhava, <i>Scolytus</i> (= <i>rugulosus</i>)	370	<i>teres</i> (Blackman), (<i>Neodryocotes</i>), <i>Araptus</i>	962
<i>taxicornis</i> Schedl, <i>Xyleborus</i>	778	<i>teres</i> Blandford, <i>Cnesinus</i>	212
<i>taxodii</i> Blackman, <i>Phloeosinus</i>	256	<i>tereticollis</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	685
<i>taxodii</i> <i>taxodii</i> Blackman, <i>Phloeosinus</i> (= <i>taxodii</i>)	256	<i>teretis</i> Wood, <i>Cnesinus</i>	212
<i>taxodii</i> <i>taxodicolens</i> Wood, <i>Phloeosinus</i> (= <i>taxodii</i>)	256	<i>teretis</i> Wood, <i>Hypothenemus</i>	946
<i>taxodicolens</i> Wood, <i>Phloeosinus</i> (= <i>taxodii</i>)	256	<i>terminaliae</i> (Hopkins), (<i>Terminalinus</i>), <i>Cyclorhipidion</i>	703
<i>tayuyaensis</i> Schedl, <i>Dendrocraniulus</i>	553	<i>terminaliae</i> Browne, <i>Cryphalus</i>	897
<i>techangensis</i> Tsai & Huang, <i>Hylastes</i>	62		

<i>terminuliae</i> Browne, <i>Margadillius</i>	599	<i>thitsi</i> Wood, <i>Polygraphus</i>	301
<i>terminaliae</i> Hopkins, <i>Coptoborus</i>	664	<i>thomasi</i> Bright, <i>Pityophthorus</i>	1031
<i>Terminalius</i> Hopkins (= <i>Cyclorhipidion</i>)	697	<i>thomasi</i> G. Hopping, <i>Ips</i> (= <i>borealis</i>)	493
<i>terminalis</i> Flores & Bright, <i>Conophthorus</i>	965	<i>thomasi</i> Wood, <i>Pseudothysanoes</i>	418
<i>terminatum</i> (Blandford), (<i>Pterocyclus</i>), <i>Monarthrum</i>	1063	<i>thompsoni</i> Schedl, <i>Rhopalopselion</i>	97
<i>terminatus</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	501	<i>thomsoni</i> Ferrari, (<i>Ernoporus</i>), <i>Ernoporicus</i> (= <i>fagi</i>)	551
<i>ternatus</i> Eggers, (<i>Dendrugus</i>), <i>Coccotrypes</i> (= <i>advena</i>)	592	<i>thoracicus</i> Chapuis, <i>Scolytus</i>	378
<i>tersus</i> Wood, <i>Phelloterus</i>	971	<i>thoracicus</i> Erichson, <i>Amphicranus</i>	1049
<i>tertius</i> Blackman, <i>Pityoborus</i> (= <i>secundus</i>)	969	<i>thoracicus</i> Hopkins, <i>Hypothenemus</i> (= <i>californicus</i>)	912
<i>tessellatus</i> Blandford, <i>Xylechinus</i>	119	<i>thoracicus</i> Lea, <i>Anasa</i> (= <i>truncatus</i>)	655
<i>testaceus</i> Fabricius, (<i>Bostrichus</i>), <i>Tomicus</i> (= <i>piniperda</i>)	135	<i>thoracicus</i> Panzer, (<i>Bostrichus</i>), <i>Xyleborus</i> (= <i>dispar</i>)	729
<i>testaceus</i> Walker, (<i>Bostrichus</i>), <i>Xyleborus</i> (= <i>perforans</i>)	760	<i>thrinacis</i> Hopkins, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>testudo</i> (Eggers), (<i>Xyleborus</i>), <i>Xylosandrus</i>	501	<i>thujae</i> (Perris), (<i>Hylesinus</i>), <i>Phloeosinus</i>	256
<i>testudo</i> Eggers, <i>Strombophorus</i>	204	<i>Thylacrus</i> Schedl (= <i>Brachyspartus</i>)	1065
<i>teteforti</i> (Menier), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946	<i>Thysanoes</i> LeConte	419
<i>tetracanthus</i> Beeson, <i>Coptodryas</i> (= <i>quadriricostata</i>)	527	<i>Tiarophorus</i> Schreiner	539
<i>tetradontium</i> Wood, <i>Monarthrum</i>	1063	<i>tibetensis</i> Wood & Huang, <i>Pseudoxylechinus</i>	114
<i>tetricus</i> (Schedl), (<i>Erieryphalus</i>), <i>Cryphalus</i>	597	<i>tibetensis</i> Wood & Yin, <i>Gnatharus</i>	1050
<i>tetricus</i> Wood, <i>Phloeotribus</i>	235	<i>tibialis</i> Wood, <i>Chramesus</i>	269
<i>texanus</i> Blackman, <i>Phloeosinus</i> (= <i>scopulorum</i>)	253	* <i>tiburtinus</i> Claus, <i>Scolytus</i> (= <i>intricatus</i>)	334
<i>texanus</i> Blackman, <i>Thysanoes</i>	421	<i>tigrensis</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>eruditus</i>)	925
<i>texanus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	942	<i>tiliae</i> (Niisima), (<i>Kissophagus</i>), <i>Pseudoxylechinus</i>	114
<i>texanus</i> Schaeffer, <i>Phloeotribus</i>	235	<i>tiliae</i> (Panzer), (<i>Apate</i>), <i>Ernoporus</i>	555
<i>texanus</i> Wood, <i>Dendroterus</i>	952	<i>tiliae</i> Semenov, <i>Hylastinus</i>	67
<i>thailundicus</i> (Schedl), (<i>Euptilus</i>), <i>Ernoporus</i>	555	<i>timidulus</i> Wood, <i>Pityophthorus</i>	1031
<i>Thamnophthorus</i> Schedl (= <i>Araptus</i>)	952	<i>timidus</i> Blandford, <i>Pityophthorus</i>	1031
<i>Thamnurgides</i> Hopkins (= <i>Coccotrypes</i>)	591	<i>timidus</i> Schedl, <i>Xyleborus</i>	779
<i>Thamnurgina</i> Nusslin (= <i>Dryocoetini</i>)	539	<i>tinianensis</i> Wood, <i>Liparthrum</i>	279
<i>Thamnurginae</i> Nusslin (= <i>Dryocoetini</i>)	539	<i>timitus</i> Schedl, <i>Xyleborus</i>	779
<i>Thamnurgus</i> Eichhoff	541	<i>titubanter</i> Schedl, <i>Xyleborus</i>	779
<i>thamnus</i> Bright, <i>Pityophthorus</i>	1031	<i>toba</i> Wichmann, <i>Scolytopsis</i> (= <i>puncticollis</i>)	321
<i>thatcheri</i> (Wood), (<i>Cryphalus</i>), <i>Trypophloeus</i>	547	<i>todo</i> Kono, <i>Xyleborus</i>	779
<i>thatcheri</i> Bright, <i>Pityophthorus</i> (= <i>pseudotsugae</i>)	1021	<i>togonus</i> (Eggers), (<i>Pseudohylesinus</i>), <i>Hylesinopsis</i>	95
<i>thatcheri</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517	<i>togonus</i> Eggers, <i>Pityophthorus</i>	1031
<i>Thaumasimus</i> Reitter (= <i>Chramesus</i>)	262	<i>tolimanus</i> (Eggers), (<i>Xyleborus</i>), <i>Coptoborus</i>	664
<i>theae</i> Eggers, (<i>Xyleborus</i>), <i>Coptodryas</i> (= <i>myristicae</i>)	825	<i>tolimanus</i> (Schedl), (<i>Prionosceles</i>), <i>Scolytodes</i>	400
<i>theae</i> Eggers, <i>Coccotrypes</i>	613	<i>tomentosus</i> (Eggers), (<i>Xyleborus</i>), <i>Hadrodemius</i>	519
<i>theklae</i> Hagedorn, <i>Ozopemon</i>	590	<i>tomentosus</i> Eichhoff, (<i>Pityophthorus</i>), <i>Pseudopityophthorus</i> (= <i>pruinosis</i>)	975
<i>Theoborus</i> Hopkins	660	<i>tomentosus</i> Schedl, <i>Corthylus</i> (= <i>papulans</i>)	1077
<i>theobroma</i> Sampson, <i>Amphicranus</i>	1049	<i>Tomici</i> (= <i>Ipini</i>)	435
<i>theobromae</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>libericensis</i>)	933	<i>Tomicidae</i> Thomson (= <i>Scolytidae</i>)	28, 103
<i>theobromae</i> Hopkins, <i>Theoborus</i>	661	<i>Tomicidae</i> (= <i>Ipini</i>)	435
<i>theobromae</i> Nunberg, <i>Corthylus</i>	1050	<i>Tomicides</i> Thomson (= <i>Tomicini</i>)	103
<i>theobromae</i> Nunberg, <i>Cryphalus</i>	597	<i>Tomicides</i> (= <i>Ipini</i>)	435
<i>theocallus</i> Bright, <i>Cnesinus</i>	212	<i>Tomicinae</i> Thomson (= <i>Tomicini</i>)	103
<i>therondi</i> Hoffman, <i>Scolytus</i> (= <i>multistriatus</i>)	353	<i>Tomicinae</i> (= <i>Ipini</i>)	435
<i>theretiae</i> Wood, <i>Liparthrum</i>	279	<i>Tomicini</i> Thomson	103
<i>thitsi</i> Beeson, (<i>Spongotarsus</i>), <i>Polygraphus</i> (= <i>thitsi</i> Wood)	301	<i>Tomicini</i> (= <i>Ipini</i>)	435
		<i>tomicoides</i> (Blandford), (<i>Pterocyclus</i>), <i>Monarthrum</i>	1063
		<i>tomicoides</i> (Eggers), (<i>Xyleborus</i>), <i>Ambrosiodmus</i>	681
		<i>tomicoides</i> (Schedl), (<i>Micracidendron</i>), <i>Saurotaxis</i>	419

<i>Tomicus</i> Latreille	122	<i>trepidus</i> Wood, <i>Araptus</i>	962
<i>tongaensis</i> Schedl, <i>Hypocryphalus</i>	872	<i>tresmariae</i> (Schedl), (<i>Hylocurus</i>), <i>Phloeocleptus</i>	422
<i>tongatapai</i> (Schedl), (<i>Mimiothorus</i>), <i>Mimiothorus</i>	950	<i>tricolor</i> Beeson, <i>Coccotrypes</i> (= <i>carpophagus</i>)	596
<i>tonkinensis</i> (Schedl), (<i>Dryocoetes</i>), <i>Dryocoetops</i>	585	<i>triangi</i> Schedl, <i>Xyleborus</i>	779
<i>tonkinensis</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	694	<i>triangularis</i> (Schedl), (<i>Xyleborus</i>), <i>Cnestus</i>	503
<i>tonsus</i> (Hagedorn), (<i>Dryocoetes</i>), <i>Xyleborus</i>	779	<i>triangularis</i> Schedl, <i>Hypocryphalus</i>	872
<i>tonsus</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866	<i>triangularis</i> Wood, <i>Cnesinus</i>	212
<i>tonsus</i> Blackman, <i>Pityophthorus</i> (= <i>intexus</i>)	1004	<i>triarmatus</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>scolytus</i>)	376
<i>tonsus</i> Eggers, <i>Hypothenemus</i> (= <i>seriatus</i>)	943	<i>triarmatus</i> Schedl, <i>Trichus</i> (= <i>nodifer</i>)	1044
<i>toracalis</i> Endrodi, <i>Trypodendron</i> (= <i>domesticum</i>)	633	<i>Triarmocerus</i> Eichhoff (= <i>Hypothenemus</i>)	905
<i>toralis</i> Wood, <i>Pityophthorus</i>	1031	<i>tribulatus</i> Wood, <i>Micracis</i>	434
<i>toranio</i> (Danthione), (<i>Byrrhus</i>), <i>Hylesinus</i>	83	<i>tribulatus</i> Wood, <i>Xyleborus</i>	779
<i>tornatilis</i> Wood, <i>Amphicranus</i>	1049	<i>tribuloides</i> Wood, <i>Xyleborinus</i>	515
<i>torneutes</i> Blandford, <i>Amphicranus</i>	1050	<i>tribulosus</i> Wood, <i>Xyleborinus</i>	515
<i>torosus</i> Wood, <i>Hylocurus</i>	428	<i>tricolor</i> (Lea), (<i>Cryphalus</i>), <i>Scolytogenes</i>	866
<i>torquatus</i> Eichhoff, <i>Xyleborus</i> (= <i>rolulus</i>)	781	<i>tricolor</i> (Schedl), (<i>Acrantus</i>), <i>Chaetoptelius</i> (= <i>versicolor</i>)	104
<i>torreyanae</i> Swaine, <i>Pityophthorus</i> (= <i>carmeli</i>)	987	<i>tricolor</i> (Schedl), (<i>Leperisinus</i>), <i>Chaetoptelius</i>	103
<i>torridus</i> Wood, <i>Pityophthorus</i>	1031	<i>Tricolus</i> Blandford	1042
<i>tortuosus</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	651	<i>tridens</i> (Mannerheim), (<i>Bostrichus</i>), <i>Ips</i>	527
<i>torulus</i> Wood, <i>Scolytus</i>	375	<i>tridens</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	454
<i>torus</i> Wood, <i>Micracis</i>	434	<i>tridens</i> engelmanni Swaine, <i>Ips</i> (= <i>tridens</i>)	525
<i>tosaensis</i> (Murayama), (<i>Ips</i>), <i>Orthotomicus</i>	478	<i>tridens tridens</i> , <i>Ips</i> (= <i>tridens</i>)	527
<i>Tasaxyleborus</i> Murayama (= <i>Cnestus</i>)	501	<i>tridentatum</i> (Schedl), (<i>Cosmocorynus</i>), <i>Monarthron</i>	1063
<i>toxicodendri</i> Hopkins, (<i>Hypothenemus</i>), <i>Trischidias</i> (= <i>atoma</i>)	947	<i>tridentatus</i> (Schedl), (<i>Mimips</i>), <i>Acanthotomicus</i>	454
<i>Toxophorus</i> Eggers (= <i>Toxophthorus</i>)	1052	<i>tridentatus</i> Eggers, <i>Ips</i> (= <i>robustus</i>)	522
<i>Toxophthorus</i> Wood	1052	<i>tridentatus</i> Hopkins, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>sparsus</i>)	945
<i>tragardi</i> Spessivtsev, <i>Pityophthorus</i>	1031	<i>trifasciatus</i> Schedl, (<i>Cosmocorynus</i>), <i>Monarthron</i> (= <i>laterale</i>)	1057
<i>Traglostus</i> Schedl	410	<i>trifolii</i> Muller, (<i>Dermestes</i>), <i>Hylastinus</i> (= <i>obscurus</i>)	66
<i>trahax</i> (Sampson), (<i>Troglostica</i>), <i>Acacacis</i>	194	<i>trigintispinatus</i> Sampson, <i>Webbia</i>	833
<i>transatlanticus</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>plumieriae</i>)	939	<i>Trigonogenius</i> Hagedorn (= <i>Pityophthorus</i>)	976
<i>transbaicalicus</i> Eggers, <i>Hylurgops</i>	42	<i>trimaeculatus</i> Schedl, (<i>Problechilus</i>), <i>Gymnochilus</i> (= <i>consocius</i>)	356
<i>transcaspicus</i> Eggers, (<i>Eccoptogaster</i>), <i>Scolytus</i> (= <i>scheryzei</i>)	371	<i>trinidadiansis</i> Schedl, <i>Xyleborus</i> (= <i>caraibicus</i>)	717
<i>transcaspicus</i> Semenov, <i>Phloeosinus</i> (= <i>bicolor</i>)	243	<i>trinitatis</i> (Hopkins), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>transitus</i> Schedl, <i>Cnesinus</i>	212	<i>trionatus</i> Eichhoff, <i>Scolytus</i> (= <i>multistriatus</i>)	345
<i>transitus</i> Schedl, <i>Xyleborus</i>	779	<i>Triotennus</i> Wollaston	540
<i>transversalis</i> Blandford, <i>Dendrosinus</i>	237	<i>Trischidias</i> Hopkins	946
<i>transversalis</i> Eggers, <i>Scolytus</i>	375	<i>triseriatus</i> Schedl, (<i>Phloeosinopsis</i>), <i>Phloeosinopsioides</i>	239
<i>transversarius</i> Schedl, <i>Phloeosinus</i>	255	<i>trispinatus</i> Browne, <i>Xyleborus</i>	779
<i>transversum</i> Blandford, <i>Pycnarthrum</i> (= <i>hispidum</i>)	355	<i>trispinatus</i> Schedl, <i>Hylocurus</i>	425
<i>transversus</i> Chapuis, <i>Dactylipalpus</i>	102	<i>trispinosus</i> Eggers, <i>Scolytodes</i>	400
<i>transversus</i> Chapuis, <i>Phloeotribus</i>	235	<i>trispinosus</i> Strohmeier, <i>Scolytus</i>	375
<i>transversus</i> Eichhoff, <i>Corthylus</i>	1050	<i>tristichus</i> Ferrari, (<i>Dryocoetes</i>), <i>Tapirorychus</i> (= <i>bicolor</i>)	557
<i>trapezicollis</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	694	<i>tristichus</i> Wood, <i>Xyleborus</i>	779
<i>treeuliae</i> Schedl, <i>Pityophthorus</i>	1032	<i>tristis</i> (Eggers), (<i>Xyleborus</i>), <i>Eucallacea</i>	694
<i>treddi</i> (Reitter), (<i>Cryphalus</i>), <i>Hypothenemus</i>	946	<i>tristis</i> (Eichhoff), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>treddi</i> Reitter, <i>Trypophloeus</i> (= <i>granulatus</i>)	545	<i>tristis</i> Blandford, <i>Hylesinus</i> (= <i>costatus</i>)	77
<i>tremulae</i> Stark, <i>Trypophloeus</i>	547	<i>tritici</i> Hopkins, <i>Hypothenemus</i> (= <i>californicus</i>)	912
<i>trenchi</i> Stebbing, <i>Polygraphus</i>	301	<i>triton</i> (Schaufuss), <i>Ambrosiodinus</i>	651
<i>trepanatus</i> (Nordlinger), (<i>Bostrichus</i>), <i>Pityogenes</i>	455		
<i>trepanatus</i> Wichmann, (<i>Pteleobius</i>), <i>Neopteleobius</i> (= <i>scutulatus</i>)	72		
<i>trepanicauda</i> (Eggers), (<i>Xyleborus</i>), <i>Webbia</i>	833		
<i>trepidus</i> Bright, <i>Pityophthorus</i>	1032		

<i>trivialis</i> Wood, <i>Hypothenemus</i>	946	<i>tuberculatus</i> (Chapuis), (<i>Rhopalopleurus</i>), <i>Chramesus</i>	270
<i>Trogoditica</i> Sampson (= <i>Acaecis</i>)	193	<i>tuberculatus</i> (Eggers), (<i>Dryotomus</i>), <i>Phlocotribus</i>	235
<i>trolaki</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	681	<i>tuberculatus</i> (Eggers), (<i>Hylurgus</i>), <i>Hylurgonotus</i>	187
<i>tropicalis</i> Wood, <i>Pseudopityophthorus</i> (= <i>pruinus</i>)	975	<i>tuberculatus</i> (Eggers), (<i>Ips</i>), <i>Acanthotomicus</i>	484
<i>tropicus</i> (Browne), (<i>Trypodendron</i>), <i>Indocryphalus</i>	649	<i>tuberculatus</i> (Eggers), (<i>Xyleboricus</i>), <i>Arxyleborus</i>	669
<i>tropicus</i> (Hagedorn), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	681	<i>tuberculatus</i> (Schedl), (<i>Eidophelus</i>), <i>Cyrtogenius</i>	569
<i>tropicus</i> (Schedl), (<i>Orthotomicus</i>), <i>Acanthotomicus</i>	484	<i>tuberculatus</i> Bright, (<i>Neodryocoetes</i>), <i>Araptus</i> (= <i>macer</i>)	959
<i>tropicus</i> Eggers, <i>Polygraphus</i>	301	<i>tuberculatus</i> Bright, <i>Carphoborus</i> (= <i>pimicolens</i>)	308
<i>tropicus</i> Eichhoff, <i>Coccotrypes</i> (= <i>dactyliperda</i>)	601	<i>tuberculatus</i> Eggers, <i>Corthylus</i>	1050
<i>trucis</i> (Wood), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866	<i>tuberculatus</i> Eggers, <i>Hylurgops</i>	43
<i>trucis</i> Wood, <i>Corthylus</i>	1080	<i>tuberculatus</i> Eggers, <i>Phlocotribus</i> (= <i>demessus</i>)	220
<i>truncatellus</i> (Schedl), (<i>Xyleborus</i>), <i>Dryocoetoides</i>	655	<i>tuberculatus</i> Eggers, <i>Sternobothrus</i>	216
<i>truncatellus</i> Schedl, (<i>Xyleborus</i>), <i>Xylosandrus</i> (= <i>cylindrotomicus</i>)	793	<i>tuberculatus</i> Eichhoff, <i>Pityophthorus</i>	1032
<i>truncaticaula</i> Browne, <i>Xyleborus</i>	779	<i>tuberculatus</i> Hagedorn, <i>Diamerus</i> (= <i>pimelioides</i>)	197
<i>truncatiferus</i> (Schedl), (<i>Xyleborus</i>), <i>Amasa</i>	685	<i>tuberculatus</i> Motschulsky, (<i>Anodius</i>), <i>Xyleborus</i> (= <i>perforans</i>)	761
<i>truncatiformis</i> (Eggers), (<i>Xyleborus</i>), <i>Amasa</i>	685	<i>tuberculatus</i> Schedl, (<i>Minips</i>), <i>Acanthotomicus</i> (= <i>tuberculifer</i>)	484
<i>truncatipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Xyleborinus</i>	818	<i>tuberculatus</i> Schedl, <i>Cladotomus</i>	239
<i>truncatorus</i> Schedl, <i>Amphicranus</i>	1050	<i>tuberculatus</i> Schedl, <i>Cryphalus</i>	897
<i>truncatulus</i> Schedl, <i>Xyleborus</i>	779	* <i>tuberculatus</i> Schedl, <i>Hylurgops</i> (= <i>tuberculifer</i>)	43
<i>truncatus</i> (Erichson), (<i>Tomicus</i>), <i>Amasa</i>	685	<i>tuberculatus</i> Strohmeier, <i>Ozopemou</i>	590
<i>truncatus</i> Bright, <i>Pseudopityophthorus</i> (= <i>declivis</i>)	972	<i>tuberculatus</i> Wollaston, <i>Aphanarthrum</i> (= <i>jubae</i>)	619
<i>truncatus</i> Browne, <i>Scolytotatus</i>	407	<i>tuberculatus</i> Wood, <i>Thysanoes</i>	421
<i>truncatus</i> Eichhoff, <i>Bothrostermus</i>	215	<i>tuberculifer</i> (Eggers), (<i>Brachyspartus</i>), <i>Corthylocurus</i>	1069
<i>truncatus</i> Sharp, <i>Xyleborus</i> (= <i>vulcanus</i>)	783	<i>tuberculifer</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703
<i>truncatus</i> Stebbing, (<i>Acanthotomicus</i>), <i>Amasa</i> (= <i>schlichii</i>)	684	* <i>tuberculifer</i> (Schedl), (<i>Phlocosinus</i>), <i>Phlocosinus</i>	258
<i>truncatus</i> Wood, <i>Corthylus</i>	1080	<i>tuberculifer</i> Schedl, <i>Cyrtogenius</i>	569
<i>truncatus</i> Wood, <i>Micracis</i> (= <i>lignator</i>)	433	<i>tuberculifer</i> Wood, <i>Acanthotomicus</i>	484
<i>truncatus</i> Wood, <i>Pseudothysanoes</i>	418	* <i>tuberculifer</i> Wood, <i>Hylurgops</i>	43
<i>trunculus</i> Bright, <i>Pityophthorus</i>	1032	<i>tuberculosissimus</i> (Eggers), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	703
<i>trunculus</i> Wood, <i>Corthylus</i>	1080	<i>tuberculosus</i> (Browne), (<i>Phlocosinus</i>), <i>Hyledius</i>	261
<i>trux</i> (Schedl), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	681	<i>tuberculosus</i> Browne, <i>Xyleborus</i>	779
<i>trux</i> Schedl, <i>Arxyleborus</i>	669	<i>tuberculosus</i> Hagedorn, <i>Hypothenemus</i> (= <i>eruditus</i>)	921
<i>trypanaeoides</i> Wollaston, (<i>Tomicus</i>), <i>Xyleborus</i> (= <i>ferrugineus</i>)	737	<i>tuberculosus</i> Herbst, (<i>Bostrichus</i>), <i>Xyleborus</i> (= <i>monographus</i>)	753
<i>trypanoides</i> Beeson, <i>Cryphalus</i>	897	<i>tuberopectus</i> Huang & Yin, <i>Hyorhynchus</i>	192
<i>Trypanophellos</i> Bright (= <i>Liparthrum</i>)	274	<i>tuberosus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>trypanus</i> Sampson, <i>Cryphalus</i>	897	<i>tuberosus</i> Niisima, <i>Crypturgus</i>	629
<i>Trypocranus</i> Eichhoff (= <i>Monarthrum</i>)	1050	<i>Tabuloscolytus</i> Bntovitsch (= <i>Scolytus</i>)	321
<i>Trypodendrina</i> Tredl (= <i>Xyloterini</i>)	632	<i>tulcanus</i> Hagedorn, <i>Corthylus</i>	1080
<i>Trypodendrinae</i> Tredl (= <i>Xyloterini</i>)	632	<i>tumidulus</i> Blandford, <i>Chramesus</i> (= <i>pumilus</i>)	268
<i>Trypodendron</i> Stephens	632	<i>tumidulus</i> Wood, <i>Pseudothysanoes</i>	418
<i>Trypographus</i> Schedl (= <i>Hylesinopsis</i>)	92	<i>tumidus</i> (Schedl), (<i>Xyleborus</i>), <i>Euwallacea</i>	694
<i>Trypophloeinae</i> Nusslin (= <i>Cryphadini</i>)	842	<i>tumidus</i> Blackman, <i>Pityophthorus</i>	1032
<i>Trypophloeus</i> Fairmaire	842	<i>tumencensis</i> Hagedorn, <i>Xyleborus</i>	779
<i>tsugae</i> (Swaine), (<i>Eccoptogaster</i>), <i>Scolytus</i>	378	<i>tungamransolus</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>tsugae</i> Swaine, <i>Pseudohylesinus</i>	113	<i>tnuggali</i> (Schedl), (<i>Pocillips</i>), <i>Coccotrypes</i>	613
<i>tsugae</i> Swaine, <i>Xyleborinus</i> (= <i>saxseni</i>)	815		
<i>tsugae</i> Tsai & Yin, <i>Sphaerotrypes</i>	201		
<i>tsukubanus</i> Murayama, <i>Xyleborus</i>	779		
<i>tuberculatum</i> Wollaston, <i>Aphanarthrum</i> (= <i>jubae</i>)	618		
<i>tuberculatus</i> (Browne), (<i>Euptilius</i>), <i>Ernoporus</i>	857		

- tunggali* Schedl, *Xyleborus* 780
- tupolevi* Eggers, (*Leperisinus*), *Hylesinus*
(=*tupolevi*) 85
- tupolevi* Stark, *Hylesinus* 85
- tupolevi* denticulosus Sokanovskii, *Hylesinus*
(=*tupolevi*) 85
- turbiculus* Schedl, *Pityophthorus* 1032
- turbinatum* Schedl, (*Pterocyclon*), *Monarthrum*
(=*fimbriaticorne*) 1056
- turbinatus* Maiti & Saba, *Webbia* 833
- turbineus* (Sampson), (*Xyleborus*),
Cryptoxyleborus 829
- turgidus* (Schedl), (*Xyleborus*), *Ambrosiodmus* 681
- turkestanicus* Semenov, *Phloeosinus* 255
- turnbowi* Wood, *Pseudothysanoes* 418
- turraeanthus* Schedl, *Xyleborus* 780
- tutuilaensis* Schedl, *Hypocyphalus* 872
- tutuilaensis* Beeson, (*Thamniurgides*), *Coccotrypes*
(=*advena*) 592
- tutulus* Bright, *Pityophthorus* 1032
- tycon* Blandford, *Scolytplatypus* 407
- typographus* (Linnaeus), (*Dermestes*), *Ips* 525
- typographus* japonicus Niisima, *Ips*
(=*typographus*) 535
- uapouensis* (Beeson), (*Ericryphalus*), *Cryphalus* 897
- uchimappensis* Murayama, *Polygraphus*
(=*japonicus*) 288
- ugandae* Schedl, *Polygraphus* 301
- ugandaensis* Schedl, *Xyleborus* 780
- ugandensis* Nunberg, *Thamniurgus* (= *senecionis*) 546
- ulmi* Redtenbacher, *Scolytus* (= *multistriatus*) 353
- ulmi* Tsai & Yin, *Sphaerotrypes* 202
- umbraticus* (Schedl), (*Neodryocoetes*), *Araptus* 962
- umbratulus* (Schedl), (*Xyleborus*), *Anasa* 685
- umbratum* (Eggers), (*Xyleborus*), *Cyclorhpidion* 703
- umbratus* (Schedl), (*Mimiophthorus*), *Mimicoccus* 950
- umbratus* Schedl, (*Conophthoeranus*),
Dendroterus (= *mexicanus*) 951
- umbratus* Wood, *Microcorthyus* 1067
- umbrinum* (Blandford), (*Pterocyclon*),
Monarthrum 1063
- uncatus* (Schedl), (*Cryphalophilus*), *Scolytogenes* 866
- uncatus* Schedl, *Xyleborus* 780
- uncinatus* (Eichhoff), (*Xylocleptes*),
Dendrocranus 553
- uncinatus* Wood, (*Mimips*), *Acanthotomicus*
(=*granulatus*) 481
- unctus* Wood, *Dactylipalpus* (= *niger*) 102
- uncus* (Schedl), (*Ips*), *Acanthotomicus* 484
- undatus* (Schedl), (*Xyleborus*), *Xyleborinus* 818
- undulata* (Sampson), (*Xyleborus*), *Coptodryas* 827
- unicolor* Eichhoff, (*Xyloterus*), *Xyloterinus*
(=*politus*) 650
- unicolor* Hopkins, (*Stephanoderes*), *Hypothenemus*
(=*eruditus*) 923
- unicornis* Nobuchi, *Hyorrhynchus* 192
- unicornis* Wood, *Chramesus* 270
- unicornis* Wood, *Micracis* 434
- unicus* (Schedl), (*Eidophelus*), *Cyrtogenius* 569
- unidentatus* Bright, *Tricholus* 1045
- unidentatus* Fabricius, (*Bostrichus*), *Taurodemus*
(=*varians*) 786
- unifasciatum* (Schedl), (*Cosmocorymus*),
Monarthrum 1063
- uniformis* (Schedl), (*Mimips*), *Acanthotomicus* 484
- uniformis* Endrodi, *Hylesinus* (= *varius*) 86
- uniformis* Schedl, *Xylechinus* 119
- uniformis* Wood & Huang, *Pseudoxylechinus* 114
- unimodus* (Schedl), (*Cryptocleptes*),
Pseudothysanoes 415
- unimodus* Beeson, (*Xyleborus*), *Eucallacea*
(=*bicolor*) 687
- unipunctatus* (Blandford), (*Hexacolus*),
Scolytodes 400
- uniseptis* Schedl, *Corthyus* 1080
- uniseriata* (Eggers), (*Bothryperus*), *Phrixosoma* 190
- uniseriatum* Schedl, *Ctonoxylon* 835
- uniseriatum* Schedl, *Pycnarthrum* 386
- uniseriatus* (Schedl), (*Mimiophthorus*),
Mimicoccus 950
- uniseriatus* Eggers, (*Stephanoderes*), *Hypothenemus*
(=*crudiæ*) 915
- uniseriatus* Eggers, (*Thamniurgides*), *Coccotrypes*
(=*duplopilosus*) 602
- uniseriatus* Eggers, (*Xyleborus*), *Anasa* (= *schlichii*) 684
- uniseriatus* Eggers, *Coccotrypes* 613
- uniseriatus* Eggers, *Dryocoetes* 586
- uniseriatus* Eggers, *Ozopemon* 590
- uniseriatus* Eggers, *Phlocotribus* 235
- uniseriatus* Schedl, *Camptocerus* (= *major*) 319
- unispinosus* LeConte, *Scolytus* 379
- unispinosus* Scheyvrew, *Scolytus* (= *jaroschewskii*) 336
- upoluensis* (Schedl), (*Xyleborus*), *Ambrosiodmus* 681
- upoluensis* Schedl, *Cryphalus* 897
- Urdugraphus* Beeson (= *Polygraphus*) 284
- urichi* Sampson, (*Xyleborus*), *Dryocoetoides*
(=*cristatus*) 657
- ursa* (Eggers), (*Xyleborus*), *Xylosandrus* 801
- ursinus* (Hagedorn), (*Xyleborus*), *Xylosandrus* 801
- ursulus* (Eggers), (*Xyleborus*), *Xylosandrus* 801
- ursus* Eggers, (*Xyleborus*), *Hadrodemius* (= *globus*) 819
- ursus* Schedl, *Amphicranus* 1050
- ursus* Sokanovskii, *Dryocoetes* (= *baikalicus*) 579
- ursus* fuscus Eggers, (*Xyleborus*), *Hadrodemius*
(=*globus*) 819
- usagaricus* (Eggers), (*Neocryphalus*), *Scolytogenes* 866
- usagaricus* (Eggers), (*Xyleborus*), *Coptoborus* 664
- usagaricus* hembibitalei Schedl, (*Xyleborus*),
Coptoborus (= *usagaricus*) 665
- usagaricus* subadjunctus Schedl, (*Xyleborus*),
Coptoborus (= *usagaricus*) 664
- usambaricum* Eggers, *Ctonoxylon* (= *flavescens*) 837
- usambaricus* Schedl, *Stephanopodius* 853
- usambaricus* Schedl, *Xylocleptes* 550
- usitata* (Schedl), (*Xyleborus*), *Schedlia* 852

<i>ussuriensis</i> Berger & Kholodkovskii, <i>Scolytotplatypus</i> (= <i>tycon</i>)	407	<i>varians</i> Lea, <i>Ficicis</i>	92
<i>ussuriensis</i> Eggers, <i>Cryphalus</i> (= <i>kurenzori</i>)	552	<i>varians</i> Schedl, <i>Pityophthorus</i> (= <i>nitidus</i>)	1016
<i>ussuriensis</i> Eggers, <i>Dryocoetes</i>	586	<i>varicus</i> Schedl, <i>Arixyleborus</i> (= <i>sus</i>)	669
<i>ussuriensis</i> Kurenzov, <i>Scolytus</i> (= <i>japonicus</i>)	335	<i>variegatus</i> (Blandford), (<i>Hylastes</i>), <i>Pseudohylesinus</i>	114
<i>ussuriensis</i> Reitter, <i>Ips</i>	538	<i>variegatus</i> (Chapuis), (<i>Phlocosinus</i>), <i>Xylechinus</i>	119
<i>ussuriensis</i> Schedl, <i>Polygraphus</i> (= <i>proximus</i>)	296	<i>variegatus</i> (Eggers), (<i>Pseudophloeotribus</i>), <i>Hylesinopsis</i>	96
<i>ussuriensis</i> <i>rigulosus</i> Eggers, <i>Dryocoetes</i> (= <i>ussuriensis</i>)	586	<i>variegatus</i> Beeson, <i>Diamerus</i> (= <i>variegatus</i>)	198
<i>usticus</i> Wood, <i>Xyleborus</i> (= <i>discretus</i>)	725	<i>variegatus</i> Eggers, <i>Chramesus</i>	270
<i>ustulatus</i> Hagedorn, <i>Xyleborus</i> (= <i>alluaudi</i>)	709	<i>variegatus</i> Eggers, <i>Sphaerotrypes</i>	202
<i>ustum</i> Schedl, <i>Corthyceylon</i>	1065	<i>variegatus</i> Eggers, <i>Strombophorus</i> (= <i>ericus</i>)	203
<i>ustus</i> Schedl, <i>Xyleborus</i>	780	<i>variegatus</i> Schedl, <i>Diamerus</i>	198
<i>usurpatus</i> Wood, <i>Sampsonius</i>	656	<i>variegatus</i> Wood & Huang, <i>Pseudoxylechinus</i>	114
<i>utahensis</i> Hopkins, <i>Procrlyphalus</i>	848	<i>variipennis</i> (Schedl), (<i>Xyleborus</i>), <i>Cyrtogenius</i>	569
<i>utahensis</i> Swaine, <i>Phlocosinus</i> (= <i>serratus</i>)	254	<i>variolatus</i> Bruck, <i>Phlocosinus</i>	259
<i>utahensis</i> Wood, <i>Ips</i> (= <i>pilifrons</i>)	517	<i>variolosus</i> Perris, <i>Hylastes</i> (= <i>linearis</i>)	56
<i>uter</i> Eggers, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>variolosus</i> Schedl, <i>Cryphalus</i>	897
<i>uter</i> Schedl, <i>Scolytotplatypus</i>	407	<i>varipes</i> Eichhoff, <i>Thammurgus</i>	546
<i>uyltenboogaarti</i> Eggers, <i>Dactylotrypes</i> (= <i>longicollis</i>)	554	<i>varius</i> (Fabricius), (<i>Bostriehus</i>), <i>Hylesinus</i>	85
<i>vachelliae</i> Blackman, <i>Thysanoes</i> (= <i>texanus</i>)	421	<i>varius</i> (Schedl), (<i>Cryphalomorphus</i>), <i>Scolytogenes</i>	866
<i>vafer</i> (Schedl), (<i>Gnathotrichus</i>), <i>Gnathotripes</i>	1042	<i>varius</i> Schedl, (<i>Problechilus</i>), <i>Gymnochilus</i> (= <i>minor</i>)	387
<i>vafer</i> Blandford, <i>Hypothenemus</i> (= <i>arecae</i>)	907	<i>varius</i> Wood, <i>Chramesus</i>	270
<i>vafr</i> (Schedl), (<i>Xyleborus</i>), <i>Coptodryas</i>	827	<i>varius</i> Wood, <i>Scolytodes</i>	400
<i>vagabundus</i> (Schedl), (<i>Cosmocorynus</i>), <i>Amphicranus</i>	1050	<i>varshalovitchi</i> Michalski, <i>Scolytus</i>	380
<i>vagabundus</i> (Wood), (<i>Loganius</i>), <i>Cnemidox</i>	318	<i>varulus</i> (Wood), (<i>Xyleborus</i>), <i>Tanrodemus</i>	787
<i>vagabundus</i> Blandford, <i>Hylaeus</i>	428	<i>vaspatorius</i> Hagedorn, <i>Ambrosiodemus</i> (= <i>desectus</i>)	672
<i>vagabundus</i> Schedl, <i>Xyleborus</i> (= <i>rolulus</i>)	783	<i>vastans</i> Chapuis, <i>Hylastes</i> (= <i>gracilis</i>)	54
<i>vagans</i> (Schedl), (<i>Xyleborus</i>), <i>Cyclorhipidion</i>	704	<i>vastus</i> Wood, <i>Chramesus</i>	270
<i>vagens</i> Eggers, (<i>Phlocosinus</i>), <i>Hyledius</i> (= <i>nitidicollis</i>)	260	<i>vastus</i> Wood, <i>Eupagiocerus</i>	214
<i>vagens</i> Schedl, <i>Strombophorus</i>	204	<i>vateriae</i> (Beeson), (<i>Thammurgides</i>), <i>Coccotrypes</i>	614
<i>valdicianus</i> (Eggers), (<i>Xylechinus</i>), <i>Xylechinonotus</i>	105	<i>vaticae</i> (Nunberg), (<i>Taphroborus</i>), <i>Cyrtogenius</i>	569
<i>valens</i> LeConte, <i>Dendroctonus</i>	183	<i>vegrandis</i> Bright, <i>Pityophthorus</i>	1032
<i>valens</i> Sampson, <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>velatus</i> (Sampson), (<i>Xyleborus</i>), <i>Euwallacea</i>	694
<i>validicornis</i> Schedl, (<i>Xyleborus</i>), <i>Xyleborinus</i> (= <i>artelineatus</i>)	805	<i>velatus</i> Schedl, <i>Gnathotripes</i>	1042
<i>validus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Euwallacea</i>	694	<i>velutinus</i> (Wood), (<i>Paracorthyus</i>), <i>Metacorthyus</i>	1065
<i>validus</i> Blandford, <i>Hypothenemus</i> (= <i>erectus</i>)	918	<i>velutinus</i> Wood, <i>Dryocoetoides</i>	659
<i>validus</i> <i>valens</i> Sampson, <i>Hypothenemus</i> (= <i>birmanus</i>)	910	<i>velutinus</i> Wood, <i>Pityoborus</i>	969
<i>vallatus</i> Wood, <i>Pseudothysanoes</i>	418	<i>venezuelensis</i> (Schedl), (<i>Phlocophthorus</i>), <i>Phlocotribus</i>	235
<i>vallidum</i> (Ferrari), (<i>Corthyus</i>), <i>Monarthrum</i>	1063	<i>venezuelensis</i> Schedl, <i>Pityophthorus</i>	1033
<i>vancouveri</i> Swaine, <i>Ips</i> (= <i>montanus</i>)	513	<i>ventralis</i> LeConte, <i>Scolytus</i>	380
<i>vandykei</i> Bruck, <i>Carphoborus</i>	310	<i>ventralis</i> Sharp, <i>Pachycotes</i> (= <i>peregrinus</i>)	187
<i>vandykei</i> Swaine, <i>Phlocosinus</i>	258	<i>ventrosus</i> Scheyvrew, <i>Scolytus</i>	382
<i>vaurynae</i> Browne, <i>Xyleborus</i>	780	<i>venturina</i> Hopkins, (<i>Xyloleptes</i>), <i>Dendrocranius</i> (= <i>cucurbitae</i>)	551
<i>variabilis</i> (Beeson), (<i>Thammurgides</i>), <i>Coccotrypes</i>	614	<i>venustulus</i> Schedl, <i>Xyleborus</i>	780
<i>variabilis</i> (Browne), (<i>Mimips</i>), <i>Acanthotomicus</i>	484	<i>venustulus</i> Wood, <i>Scolytodes</i>	400
<i>variabilis</i> Schedl, <i>Xyleborus</i> (= <i>caraibicus</i>)	717	<i>venustus</i> Blackman, <i>Pityophthorus</i>	1033
<i>variabilis</i> Sokanovskii, <i>Scolytus</i> (= <i>scolytus</i>)	371	<i>venustus</i> Schedl, (<i>Thyureos</i>), <i>Corthyus</i>	1081
<i>variabilis</i> Wood, <i>Chramesus</i>	270	<i>venustus</i> Schedl, <i>Ptilopodius</i>	900
<i>varians</i> (Fabricius), (<i>Bostriehus</i>), <i>Tanrodemus</i>	786	<i>venustus</i> Wood, <i>Scolytodes</i>	400
		<i>verax</i> Schedl, (<i>Xyleborus</i>), <i>Amasa</i> (= <i>schlichii</i>)	684
		<i>verdicus</i> Wood, <i>Pseudothysanoes</i>	418
		<i>vernaculum</i> (Schedl), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1064

- vernaculus* (Schedl), (*Brachyspartus*),
Corthylocurus 1069
- vernaculus* Schedl, *Xyleborus* 750
- verrucifrons* Tsai & Yin, *Polygraphus* 301
- verrucosus* Wood, *Dryocoetoides* 659
- verrucosus* Wood, *Hylocurus* 428
- versicolor* (Sampson), (*Xyleborus*), *Amasa* 685
- versicolor* Wood, *Chaetoptelinus* 104
- verticillus* Wood, *Pseudothyrsanoes* 418
- rescula* Wood, *Micracisella* 431
- resculus* Wood, *Araptus* 962
- resculus* Wood, *Hypothenemus* 946
- resculus* Wood, *Phloeotribus* 235
- resculus* Wood, *Pityophthorus* 1033
- resculus* Wood, *Pseudothyrsanoes* 418
- resculus* Wood, *Scolytodes* 400
- rescus* Wood, *Microcorthyus* 1067
- respatorius* (Schedl), (*Xyleborus*), *Coptoborus* 665
- respatorius* Hagedorn, *Ambrosiodinus* (= *desectus*) 672
- respertinus* Bright, *Pityophthorus* 1033
- restigator* (Schedl), (*Xyleborus*), *Cryptoxyleborus* 829
- restitum* Wollaston, *Aphanarthrum* (= *bicinctum*) 617
- restitus* (Eggers), (*Loganius*), *Cnemomyx* 318
- restitus* (Eggers), (*Problechilus*), *Scolytodes* 400
- restitus* (Mulsant & Rey), (*Hylesinus*),
Chaetoptelinus 104
- restitus* Blandford, *Cryphalus* 897
- restitus* Brown, *Dendrotrypes* (= *costiceps*) 106
- restitus* Eggers, *Cnesinus* 212
- restitus* Eggers, *Phloeotribus* 235
- restitus* Schedl, *Xyleborus* (= *adusticollis*) 706
- rexans* (Schedl), (*Xyleborus*), *Dryocoetoides* 659
- rexator* (Browne), (*Spongotarsus*), *Polygraphus* 302
- rexator* Reitter, *Scolytus* (= *koltzei*) 338
- rexator* Schedl, *Cnesinus* 212
- rianai* Schedl, (*Conophthoracranulus*),
Spermophthorus (= *apuleiae*) 971
- rianai* Schedl, *Cnemomyx* (= *floricornis*) 315
- rianai* Schedl, *Hylocurus* 428
- rianai* Schedl, *Hypothenemus* (= *eruditus*) 925
- rianai* Schedl, *Pseudochramesus* 262
- riaticus* Schedl, *Xyleborus* 780
- riburni* Eggers, *Cryphalus* (= *riburni* Stark) 898
- riburni* Stark, *Cryphalus* 897
- vicarius* Beeson, (*Thamnurgides*), *Corcotypes*
 (= *advena*) 593
- vicarius* Eichhoff, *Xyleborus* (= *pfeili*) 763
- vicinalis* Wood, *Dendrocrannulus* 554
- vicinum* (Schedl), (*Pterocydon*), *Monarthrum* 1064
- vicinus* (Eggers), (*Prionosecles*), *Scolytodes* 400
- vicinus* Comolli, (*Hylesinus*), *Kissophagus*
 (= *hederae*) 68
- vicinus* Eichhoff, *Xyleborus* 780
- vicinus* LeConte, (*Xyleborus*), *Orthotomicus*
 (= *caelatus*) 469
- vicinus* Wood, *Dendrocrannulus* 554
- vicinus* Wood, *Microcorthyus* 1067
- victoris* Mulsant & Rey, (*Bostrichus*), *Dryocoetes*
 (= *autographus*) 578
- ridius* Eichhoff, *Xyleborus* 750
- vietnamensis* Schedl, *Polygraphus* 302
- rigilans* (Schedl), (*Xyleborus*), *Cyclorhipidion* 704
- rilis* (Blandford), (*Phloeosinus*), *Hyledius* 261
- rilliersi* (Lepesme), (*Thamnurgus*), *Dendrochilus* 435
- rilliersi* Schedl, *Trietennus* 541
- rillifer* Wood, *Corthyus* 1081
- rillifrons* (Dufour), (*Bostrichus*), *Taphrorychus* 560
- rillifrons* Wood, *Hylocurus* 429
- rillosulus* (Blandford), (*Xyleborus*), *Theoborus* 661
- rillosulus* Lacordaire, *Phloeotribus* (= sp.?) 215
- rillosus* (Fabricius), (*Bostrichus*), *Dryocoetes* 586
- rillosus* Eggers, *Corthyus* 1080
- rillosus* Herbst, *Dryocoetes* (= *autographus*) 576
- rillosus* Ratzeburg, (*Bostrichus*), *Xyleborus*
 (= *cryptographus*) 722
- rillosus* Schedl, (*Xyleborus*), *Theoborus*
 (= *rillosulus*) 662
- rillosus* Schedl, *Pachycotes* 185
- rillosus minor* Eggers, *Dryocoetes* (= *rillosus*) 586
- rillosus starhoni* Reitter, *Dryocoetes* (= *rillosus*) 586
- rilus* Bright, *Corthyus* 1081
- riminalis* Bright, *Pityophthorus* 1033
- vinealis* Bright, (*Neodryocoetes*), *Araptus*
 (= *tabogae*) 962
- vinealis* Wood, *Chramesus* 270
- vinealis* Wood, *Dendrocrannulus* 554
- rimulus* Wood, *Araptus* 963
- vinogradovi* (Semenov), (*Phloeophthorus*),
Phloeotribus (= *rhododactylus*) 230
- virentis* Hopkins, (*Stephanoderes*), *Hypothenemus*
 (= *seriatus*) 942
- virgatus* Bright, *Scolytus* 352
- virginianae* Hopkins, *Conophthorus* (= *resenosae*) 967
- virilis* Blackman, *Pityophthorus* 1033
- virilis* Wood, *Pseudopityophthorus* 976
- viriosa* Wood, *Phrixosoma* 190
- virtus* Schedl, *Pityophthorus* 1033
- viruensis* Browne, *Xyleborus* 780
- viscicolens* Wood, *Pseudothyrsanoes* 418
- viscivorus* Wood, *Pseudothyrsanoes* 418
- vismiacolens* Wood, *Cnemomyx* 318
- vismiae* Eggers, (*Loganius*), *Cnemomyx*
 (= *minusculus*) 317
- vismiae* Wood, *Xyleborus* 780
- Vitaderes* Beeson (= *Cosmoderes*) 901
- ritei* Wood, *Dendroctonus* 156
- ritiensis* Browne, *Cryphalus* 897
- ritiosus* Schedl, *Xyleborus* (= *adelographus*) 706
- ritiosus* Wood, *Chramesus* 270
- ritis* Browne, *Hypothenemus* 946
- ritatum* (Blandford), (*Pterocydon*), *Monarthrum* 1064
- ritatus* (Fabricius), (*Bostrichus*), *Pteleobius* 70
- ritatus* Eggers, *Strombophorus* 204
- ritatus* Schedl, *Xylechinus* 119

<i>rittifrons</i> Blandford, <i>Dendrosinus</i>	237	<i>woodi</i> Schedl. (<i>Xyleborus</i>), <i>Ambrosiodinus</i> (= <i>devexus</i>)	673
<i>rittiger</i> Eichhoff, <i>Trypodendron</i> (= <i>lineatum</i>)	643	<i>woodi</i> Thatcher, <i>Ips</i>	538
<i>ritulus</i> Wood, <i>Micracis</i>	434	° <i>worthini</i> Scudder, (<i>Polygraphus</i>), not <i>Scolytidae</i>	284
<i>roarotrae</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	695	<i>xalapae</i> Atkinson, <i>Chramesus</i>	270
<i>rolastos</i> (Schedl), (<i>Thamniophthorus</i>), <i>Araptus</i>	963	<i>xalapae</i> Wood, <i>Pseudopityophthorus</i>	976
<i>rolcanus</i> Wood, <i>Scolytodes</i>	400	<i>xalapensis</i> Wood, <i>Monarthrum</i>	1064
<i>rolustus</i> Eggers, <i>Dendrocranius</i> (=sp?)	550	<i>xanthophloae</i> (Schedl), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946
<i>rolulum</i> (Eichhoff), (<i>Pterocyclon</i>), <i>Monarthrum</i>	1064	<i>xanthophloae</i> Schedl, <i>Lanurgus</i>	409
<i>rolulus</i> (Fabricius), (<i>Bostrichus</i>), <i>Xyleborus</i>	750	<i>xanthophylli</i> (Browne), (<i>Phloeosinus</i>), <i>Hyledius</i>	261
<i>rolulus</i> Schedl, <i>Pityophthorus</i>	1033	<i>xanthophyllus</i> Schedl, <i>Xyleborus</i>	783
<i>rorontzowi</i> (Jacobson), (<i>Tomicus</i>), <i>Pityokteines</i>	466	<i>xanthopus</i> (Eichhoff), (<i>Xyleborus</i>), <i>Eucallacea</i>	695
<i>ryrdaghi</i> (Nunberg), (<i>Apoglostatus</i>), <i>Glostatus</i>	836	<i>xanthopus</i> Dejean, (<i>Bostrichus</i>), <i>Eucallacea</i> (= <i>xanthopus</i> Eichhoff)	696
<i>ryrdaghi</i> Nunberg, <i>Pityophthorus</i>	1033	xaverti Reitter, <i>Polygraphus</i> (= <i>subopacus</i>)	300
<i>rulcanus</i> Perkins, <i>Xyleborus</i>	783	<i>Xelyborus</i> Schedl (= <i>Webbia</i>)	829
<i>rulgaris</i> (Eggers), (<i>Dendrognus</i>), <i>Coccotrypes</i>	614	<i>Xenophthorus</i> Wood & Yin (= <i>Pseudopityophthorus</i>)	971
<i>rulgaris</i> Schauffuss, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>seriatus</i>)	941	<i>Xestips</i> Hagedorn (= <i>Xylocleptes</i>)	547
<i>rulgaris</i> Wood, <i>Stegomerus</i>	852	<i>Xylebori</i> LeConte (= <i>Xyleborini</i>)	651
<i>rupina</i> (Schedl), (<i>Xyleborus</i>), <i>Schedlia</i>	822	<i>Xyleboricus</i> Eggers (= <i>Arixyleborus</i>)	665
<i>rsachtli</i> Reitter, <i>Hylesinus</i>	89	<i>Xyleboridae</i> LeConte (= <i>Xyleborini</i>)	651
<i>rsadai</i> Murayama, <i>Pseudohyorrhynchus</i>	192	<i>Xyleborinae</i> LeConte (= <i>Xyleborini</i>)	651
<i>wakayamensis</i> Nobuchi, <i>Xyleborus</i>	353	<i>Xyleborina</i> LeConte (= <i>Xyleborini</i>)	651
<i>walkeri</i> (Blandford), (<i>Cryptarthrum</i>), <i>Cryphalus</i>	898	<i>Xyleborini</i> Leconte	651
<i>wallacei</i> (Blandford), (<i>Hylesinus</i>), <i>Ficicis</i>	92	<i>Xyleborinus</i> Reitter	803
<i>wallacei</i> (Blandford), (<i>Xyleborus</i>), <i>Eucallacea</i>	695	<i>Xyleborips</i> Reitter (= <i>Xyleborus</i>)	704
<i>wallacei</i> indocorus Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>wallacei</i>)	695	<i>Xyleborites</i> Wickham	1082
<i>wapleri</i> Eichhoff, <i>Cryphalus</i>	898	<i>Xyleborus</i> Eichhoff	704
<i>waringi</i> Curtis, (<i>Bostrichus</i>), <i>Trypodendron</i> (= <i>signatum</i>)	647	<i>Xylechinides</i> Nusslin (= <i>Tomicini</i>)	103
<i>watsoni</i> Schedl, <i>Pityophthorus</i> (= <i>murrayanae</i>)	1014	<i>Xylechinites</i> Hagedorn	1082
<i>webbi</i> Blackman, <i>Hylastes</i> (= <i>porculus</i>)	61	<i>Xylechinops</i> Browne (= <i>Xylechinus</i>)	114
<i>webbi</i> Hopkins, <i>Hypothenemus</i> (= <i>eruditus</i>)	922	<i>Xylechinus</i> Chapuis	114
<i>webbi</i> Hopkins, <i>Xyleborus</i> (= <i>subcostatus</i>)	776	<i>Xylochilus</i> Schedl (= <i>Dendroterus</i>)	950
Webbia Hopkins	829	<i>Xylocleptes</i> Ferrari	547
<i>Webbinae</i> Hopkins (= <i>Xyleborini</i>)	651	<i>xylocranellus</i> Schedl, (<i>Xyleborus</i>), <i>Premnobius</i> (= <i>cavipennis</i>)	652
<i>whitfordi</i> Schedl, (<i>Xyleborus</i>), <i>Eucallacea</i> (= <i>fornicatus</i>)	690	<i>Xylocryptus</i> Schedl (= <i>Scolytogenus</i>)	858
<i>whitteni</i> Beeson, <i>Xyleborus</i> (= <i>perforans</i>)	761	<i>Xyloctonidae</i> Eichhoff (= <i>Xyloctonini</i>)	835
<i>wickhami</i> Blackman, <i>Scolytus</i> (= <i>reflexus</i>)	364	<i>Xyloctoninae</i> Eichhoff (= <i>Xyloctonini</i>)	835
<i>wickhami</i> Hopkins, <i>Dendroctonus</i> (= <i>rufipennis</i>)	173	<i>Xyloctonini</i> Eichhoff	835
<i>widdringtoniae</i> Schedl, <i>Lanurgus</i>	409	<i>xyloctonoides</i> Schedl, <i>Coriacephilus</i>	854
<i>wieslanderi</i> Swaine, <i>Ips</i> (= <i>mexicanus</i>)	512	<i>xyloctonus</i> (Schedl), (<i>Ctonocryptus</i>), <i>Glostatus</i>	836
<i>wilderi</i> (Beeson), (<i>Xyleborus</i>), <i>Ambrosiodinus</i>	681	<i>Xyloctonus</i> Eichhoff	840
<i>willei</i> Schedl, <i>Phloeotribus</i>	235	<i>Xylogopinus</i> Schedl (= <i>Hylurdretonus</i>)	186
<i>winkleri</i> (Reitter), (<i>Stephanoderes</i>), <i>Hypothenemus</i>	946	<i>xylographus</i> (Say), (<i>Bostrichus</i>), <i>Xyleborus</i>	783
<i>wisteriae</i> Wood, <i>Chramesus</i>	270	<i>xylographus</i> Sahlberg, (<i>Bostrichus</i>), <i>Pityogenes</i> (= <i>chalcographus</i>)	451
<i>wittei</i> Eggers, <i>Thamurgus</i>	546	<i>xylographus</i> Schedl, <i>Lanurgus</i>	409
° <i>wolffi</i> (Schedl), (<i>Phloeosinities</i>), <i>Phloeosinus</i>	259	<i>xylophagus</i> Blackman, <i>Thysanoes</i>	421
<i>wollastoni</i> (Eichhoff), (<i>Crypturgus</i>), <i>Cisurgus</i>	630	<i>xylophagus</i> Wood, <i>Chramesus</i>	270
<i>wollastoni</i> Israelson, <i>Aphanarthrum</i>	620	<i>Xylosandrus</i> Reitter	787
<i>woodi</i> Bright, <i>Phloeosinus</i> (= <i>hoppingi</i>)	248	<i>Xyloteri</i> Lindemann (= <i>Xyloterini</i>)	632
<i>woodi</i> Bright, <i>Pityophthorus</i>	1033	<i>Xyloteridae</i> Lindemann (= <i>Xyloterini</i>)	632
<i>woodi</i> Browne, <i>Scolytominus</i>	840	<i>Xyloterinae</i> Lindemann (= <i>Xyloterini</i>)	632
		<i>Xyloterini</i> Lindemann	632

<i>Xyloterina</i> Lindemann (=Xyloterini)	632	<i>zapotecus</i> Bright, <i>Cnesinus</i> (= <i>elegantis</i>)	208
<i>Xyloterinus</i> Swaine	649	<i>zeae</i> Eggers, <i>Pagiocerus</i> (= <i>frontalis</i>)	214
<i>Xyloterioidea</i> Lindemann (=Xyloterini)	632	<i>zeae</i> Schedl, (<i>Stephanoderes</i>), <i>Hypothenemus</i> (= <i>californicus</i>)	912
<i>xyloteroides</i> Eggers, <i>Xyleborus</i>	785	<i>zealandicus</i> Wood, <i>Dendrotrupes</i>	106
<i>Xyloterus</i> Erichson, <i>Trypodendron</i>	632	<i>zelus</i> Wood, <i>Corthylus</i>	1081
<i>xylotrupes</i> (Eichhoff), (<i>Pityophthorus</i>), <i>Araptus</i>	963	<i>zeteki</i> Blackman, <i>Pityophthorus</i> (= <i>subcribratus</i>)	1030
<i>xylotrupes</i> Schedl, <i>Xyleborus</i>	785	<i>zexmenicora</i> Bright, <i>Pityophthorus</i>	1033
<i>yablonianus</i> Murayama, <i>Scolytus</i>	352	<i>zeylanicus</i> Wood, <i>Acacasis</i>	194
<i>yakushimanus</i> (Murayama), (<i>Xyleborus</i>), <i>Arixyleborus</i>	669	<i>zhobi</i> (Stebbing), (<i>Phlocosinus</i>), <i>Carphoborus</i>	310
<i>yamaguchii</i> Inouye & Nobuchi, <i>Cryphalus</i>	595	<i>zhungdianensis</i> Tsai & Yin, <i>Polygraphus</i>	302
<i>yasamatsui</i> Nobuchi, <i>Pseudohylesinus</i> (= <i>sericeus</i>)	113	<i>zicsii</i> (Schedl), (<i>Xyleborus</i>), <i>Eucallacea</i>	697
<i>yavapaii</i> Blackman, <i>Pseudopityophthorus</i>	976	<i>zimmermani</i> Schedl, <i>Cryphalus</i>	595
<i>yohoensis</i> Swaine, <i>Ips</i> (= <i>tridens engelmanni</i>)	525	<i>zimmermanni</i> Schedl, <i>Ptilopodius</i> (= <i>murquesanus</i>)	900
<i>yuccae</i> (Wood), (<i>Cryptocleptes</i>), <i>Pseudothysanoes</i>	415	<i>zimmermanni</i> (Hopkins), (<i>Anisandrus</i>), <i>Xylosandrus</i>	501
<i>yuccarorus</i> Wood, <i>Pseudothysanoes</i>	419	<i>zimmermanni</i> Wickham, (<i>Phloeotribus</i>) <i>Curculionidae</i>	215
<i>yukonis</i> Fall, <i>Hylastes</i> (= <i>nigrinus</i>)	57	<i>zonalis</i> Bright, <i>Pityophthorus</i>	1033
<i>yunnanensis</i> Tsai & Yin, <i>Sphaerotrypes</i>	202	<i>zonatus</i> Eichhoff, <i>Gymnochilus</i>	357
<i>zuchvatkini</i> (Krivolutskaia), (<i>Ecoryphalus</i>), <i>Ernoporicus</i>	552	<i>zukevalae</i> Scott, <i>Thanurgus</i>	546
<i>zaitzevi</i> Butovitsch, <i>Scolytus</i>	352	<i>Zygophloeus</i> Schedl	271
<i>zambianus</i> Beaver, <i>Polygraphus</i>	302		



3 2044 072 231 335

