


2/2

QK21
.G7
G8





JOHN TRADESCANT THE ELDER

EARLY
BRITISH BOTANISTS
AND THEIR GARDENS

BASED ON UNPUBLISHED WRITINGS OF
GOODYER, TRADESCANT, AND OTHERS

BY

R. T. GUNTHER, M.A., F.L.S.

LIBRARIAN AND RESEARCH FELLOW OF MAGDALEN COLLEGE

LIBRARY
NEW YORK
BOTANICAL
GARDENS

With Nine Plates and Twenty-one other Illustrations

OXFORD
PRINTED BY FREDERICK HALL FOR THE AUTHOR
AT THE UNIVERSITY PRESS

1922

G7
G8

LIBRARY
NEW YORK
BOTANICAL
GARDEN

PREFACE

THE following accounts of some Botanists of the Elizabethan and Jacobean age have gathered around the literary remains of one who, but twelve years ago, was introduced to us as 'A forgotten Botanist of the seventeenth Century'. By a strange hazard we can now come closer to John Goodyer through his own writings than to any of the contemporaries whose names have been writ larger on the roll of the history of botany: and through him, other botanists of distinction have been made known, who otherwise would have remained in almost total oblivion; for as a modern authority has recently discovered, 'Every writer of the period owed help from Goodyer in one way or another'.¹

The Goodyer papers serve to illustrate missing chapters in the histories of Botany and Horticulture in that most interesting period of British Science, the hundred years which preceded the foundation of the Royal Society. Authors of standard histories of British Botany, largely based on German authority, have been apt to skim rapidly over this period, in which several of our countrymen were in some respects well abreast of Linnaeus. And these manuscripts with all the annotated books, which Goodyer bequeathed to Magdalen College in 1664, are probably the completest and most useful collection, for a study of English Botany that was not merely pre-Linnean, but was pre-Morisonian and pre-Raian as well.

In his scientific attitude of mind Goodyer was superior to several of the first members of the Royal Society.

¹ White, *Bristol Flora*, p. 57.

MAR 11 1932

He had no use for the superstitions of Ashmole or Aubrey, nor would he, like Sir Kenelm Digby, have fed his wife on capons fattened with the flesh of vipers in order to preserve her beauty. Nor would he, like the credulous Sir R. Moray, have seen tiny geese, perfectly shaped, in little shells adhering to trees among the western islands of Scotland.

His notes begin in 1616, show the period of his greatest activity to have been in 1621, and become fewer after 1633. The material came into my hands in the form of thousands of scraps of paper in disorder and in various handwritings. These had first to be sorted and bound; and then, although Goodyer could, and did generally write a remarkably clear hand, his jotted notes are scribbles, and exceedingly difficult to read. In some cases weeks elapsed before the meaning of the more difficult passages dawned upon me, and even with expert help, there are still unread words in our text. Our readers will kindly remember that many notes, that we have printed as indications of occupations and interests, were solely intended for the eye of the writer.

Further biographical details have been gleaned from visits to various parts of the country, from ledgers relating to College estates, from the parish registers of several Hampshire towns, from the account books of the Weston Charity at Petersfield, from wills at Somerset House.

As the work progressed new facts relating to Goodyer's botanical contemporaries emerged, which were scarcely less interesting than those relating to himself. Except in the papers which we now describe, there is no other surviving record of their work for Botany or Horticulture. This is due in some measure to the disturbances of the Civil War, partly too to the death of such workers as

How, Dale, and Johnson at an early age, and before the complete publication of their work, and partly to the absence at that early period of any School of English Botany or of any botanical journal.

Thanks to Goodyer we are now able to print much new matter relating to the plant-records of Sir John Salusbury, William Mount, Richard Shanne, Walter Stonehouse, William How, Dr. John Dale, and others, to publish many 'first evidences' of the plants of Kent, Hampshire, and other counties, and to list the garden plants grown by John Coys, John Parkinson, the elder Tradescant, and Morison; and in some instances from their original writings. To many, our lists of pre-Linnean plant-names may appear uninteresting, but we believe that such publication is a necessary preliminary to the preparation of any comprehensive monograph on the subject of the introduction of plants into English Gardens, whence a few, e.g. the Italian Ivy-leaved Toadflax, have run wild all over the country.

If in this compilation I have disentangled a few of the knots in that ancient skein of names and dates, I rest satisfied. I know that the fabric is left with plenty of 'ends' for other workers, and, like the Irishman's net, is full of holes.

It remains for me to acknowledge my obligations to my College, not only for having given me the opportunity of finding and arranging the Goodyerian manuscripts, but also for having made a most substantial contribution towards the heavy cost of the printing. St. John's and Jesus Colleges have likewise assisted with grants in aid of the publication of the plant records of How and Salusbury, distinguished members of their respective Societies, and the Delegates of the Clarendon Press have assisted financially at a very difficult time and by the loan

of their block of the Tradescant portrait. Mr. J. Murray has lent blocks of Lobel and Parkinson. Miss Lacell permitted me to look over Goodyer's house in Petersfield and Mr. C. Branfill Russell pointed out the vestiges of Coys's house and garden. The design of a partridge with a *good ear* of wheat in its bill, which is impressed on the binding of this volume, is the crest of Mr. Edward Goodyear, who has kindly lent the stamps with which all the books bequeathed to Magdalen College by his kinsman have been marked.

I have derived much advantage from the printed works of my predecessors, Canon Vaughan the 'discoverer', Miss Wotton the 'pioneer', and Mr. Druce the 'producer' of the forgotten Hampshire botanist of the Seventeenth Century. On certain doubtful points I have had the advantage of the experience of Dr. Church, and in the reading of difficult passages, of Messrs. Salter, Driver, Craster, and Gambier-Parry.

My grateful acknowledgements are due to my friends Sir David Prain, Professor Keeble, the officers of the Botanical Department of the British Museum, Dr. Rendel and Mr. James Britten; Dr. Stapf of the Kew Herbarium, and especially Dr. Daydon Jackson, the biographer of Gerard and Turner. To all I must express my thanks for much valuable assistance. I have also to thank my wife for sacrificing much time in the labour of revising both manuscript and proofs.

R. T. GUNTHER.

MAGDALEN COLLEGE.

February 1922.

CONTENTS

	PAGE
I	
LIFE OF JOHN GOODYER	I

II	
DESCRIPTIONS OF PLANTS BY GOODYER	100

III	
THE GOODYER'S BOTANICAL LIBRARY	197

IV	
NOTES ON CONTEMPORARY BOTANISTS	233

V	
LISTS OF PLANTS GROWN IN ENGLISH GARDENS	303

VI	
LISTS OF EXOTIC PLANTS	358

VII	
GOODYER'S MISCELLANEOUS PAPERS	372

INDEXES

GOODYER'S LIST OF PLANTS (INDEX I)	387
II. INDEX OF PLANTS	402
III. INDEX OF PERSONS, PLACES, AND THINGS	409

LIST OF ILLUSTRATIONS

	PAGE
John Tradescant	<i>Frontispiece</i>
Map of South-east Hampshire	4
Mapledurham House	7
The Garden at Stubbers	<i>facing</i> 16
Mill-mountain	22
Jerusalem Artichoke	23
Sheet Mill	<i>facing</i> 28
Goodyer's four Elms	39-42
Goodyer's signature	55
Plan of Petersfield	64
Goodyer's House in the Spain	<i>facing</i> 64
Goodyer's House in the Spain	66
Interlinear Translation of Dioscorides	<i>facing</i> 84
Drawings by Goodyer	99
Potamogeton	124-125
Description of Yew	169
The Heath at Petersfield	<i>facing</i> 188
Notes in Ray's <i>Catalogus</i>	223
Lobel	246
{ Old Testimonial to Lobel	<i>facing</i> 248
{ Sonnet dedicated to Lobel	" 249
Signatures to Lobel's second Testimonial	250
John Parkinson	266
{ " Letter	<i>facing</i> 276
{ First Draft of a British Flora by How	<i>facing</i> 276
Yucca	313
Tradescant's Title-Page	334

THE LIFE OF JOHN GOODYER

JOHN GOODYER was born at Alton in Hampshire in 1592.¹ Possibly he first saw the light in a house belonging to Magdalen College, Oxford, the College which both through its landed property in the Petersfield district, through Goodyer's relationship to its tenants and bailiffs, and through the scientific members of its foundation, was so fully to win his confidence and affection, that in his last will and testament he bequeathed to it his most cherished possession, his botanical library and manuscripts. These materials are now available for the reconstruction of his life and work.

His father, Reginald Goodyer, appears several times in the College books.² There is a copy of the terms on which the then President, William Langton, and the Fellows of the College leased to 'Reginald Goodyear', a yeoman of the parish of St. Laurence in Alton, the farm called Beeches Place and the wood called Priors Reade now or late in the tenure of Henry Mervyn, Esq., at an annual rent of £4 3s. 4d. together with two quarters of 'good, sweete, & marchantable wheate, and furthermore 3 quarters of good sweete malte to be delivered within the said College', or their value in money according to the prices of the Oxford market.

¹ I have no knowledge of an entry in any Baptismal Register. The year has probably been calculated from his Marriage Licence which states his age as forty in 1632.

² The entry in Ledger K is dated 27 July 1614, and on f. 188 there is a further entry made in 1619, when the name is spelt Reginald Goodier. According to information from Miss Wotton, Reginald Goodyer had previously paid taxes direct from Alton in 1600, and was described in a Star Chamber Case as a yeoman of Alton in 1605. Then (*Star Chamber Proc. James I*, 204-13) he gave evidence for Sir Richard Pawlett, showing intimate acquaintance with the manor of Herriard, where his father was living in 1572, and with the hamlet of Southropp. It is possible that he was a sub-tenant of Beeches Place under Henry Mervyn, before he held it under a direct lease from the College.

Concerning Reginald's family history we have no knowledge. His wife Ann, who predeceased him, bore him two sons and two daughters: Lewis, b. 1579, d. 1655, Rose, Ann, and John, b. 1592. Lewis had at least eight children and left numerous descendants; Rose married William Yalden; Ann married Richard Pratt and had three sons; and John, the subject of this memoir, had an only daughter Elizabeth.

The Yaldens were still more closely connected with Magdalen College. In 1593 William Yalden, perhaps the father of John's brother-in-law, leased from Magdalen farm-lands in Sheet, near Petersfield, known as Brooke-land, Skindre, Shirk leyes, and Pulyns. As early as 1587 (30 Elizabeth) with 'Dorothie his wyffe' he rented Sheet mills, and in 1596 acting as the College bailiff, collected the College rents, £171 5s. 3d. from Selborne and £18 9s. 6d. from Petersfelde. In 1597, he was appointed steward for the holding of courts and leets within the borough of Petersfield, by Th. Hanbury of Buriton, who had recently purchased the property. The lease for the Sheet mills mentions 'water mills, a wheat mill and a malt mill, together with all the waters, watercourses, ponds, fishinge, banckes, baies, and fludgates thereunto belonging, with free libertie to digg turfe in the great moore'. Even the legal document is redolent of the natural amenities, for which the rent was £5 6s. 8d. and half a crown in 1618. William Yalden, described as of the diocese of Chichester, was the College Clerk of the Account from 1616 to 1643; and among the other College tenants were Crusophilus Yalden at Roplie Farm (1618) and Henrie Yalden in the Spaine in Petersfield.

John Goodyer's nephew and heir, the Rev. Edmund Yalden, son of William Yalden of Sheet, *gen.*, became a Demy and Fellow of Magdalen, 1630 to 1642, when he resigned his fellowship on being presented to the Rectory of Compton in Surrey. Sheet was a botanical locality often mentioned by Goodyer.

The Goodyer country is perhaps better known to, though less visited by the English reader, than any other inland area in England. Its natural features and attractions have been made widely known by the premier work on nature study in our language, the classic *Natural History of Selborne*, by Gilbert White.

This country in which John Goodyer lived extends among the chalk hills at the junction of the North and South Downs, on and around one of the principal watersheds of south Britain. Born in the valley of the Wey, whose waters flow into the Thames, he passed his young manhood at Droxford on the Meon, which runs straight down to the Solent six miles west of Portsmouth; while at Petersfield he lived by the sources of the Rother, whose waters, mingling with those of the Arun, enter the English Channel at Arundel. A cyclist could visit all his homes in an afternoon.

As Gilbert White said a century and a half later, in this district 'so diversified with such a variety of hill and dale, aspects and soils, it is no wonder that great choice of plants should be found. Chalks, clays, sands, sheep-walks and downs, bogs, heaths, woodlands, and champaign fields cannot but furnish an ample *Flora*. The deep rocky lanes abound with *filices*, and the pastures and moist woods with *fungi*. If in any branch of botany we may seem to be wanting, it must be in the large aquatic plants, which are not to be expected on a spot far removed from rivers, and lying up amidst the hill country at the spring heads'. And yet perhaps it was just this upland character of his native country that caused Goodyer to pay especial attention to water plants whenever he came across them, an attention of which the reward was several most remarkable discoveries. He added at least a dozen aquatics to the British flora.

Nor are the literary associations of this favoured spot confined to the name of Gilbert White. Droxford is remembered as the village where Izaak Walton passed

the last years of his life, Steventon was the cradle and Chawton the inspiration of the genius of Jane Austen, for if *Pride and Prejudice* and *Sense and Sensibility* were composed at the former, *Emma*, *Mansfield Park*, and *Persuasion* were produced at the latter; while Buriton was the home of the childhood of Edward Gibbon, who always



SKETCH-MAP OF SOUTH-EAST HAMPSHIRE.

cherished fond recollections of its natural beauties. 'The aspect of the adjacent grounds was various and cheerful: the Downs commanded the prospect of the sea, and the long hanging woods in sight of the house could not perhaps have been improved by art or expense.' Such is the appreciation of the place in the *Autobiography* of the great historian.

The localities most frequently mentioned in the Goodyer papers are Alton on the high road from London to Winchester and about half-way between Farnham and Alresford, three miles east of which is Ropley. To the north-east and south-west of Alton are Holybourne and Chawton respectively. Bramshott and Liphook lie to the east, Selborne to the west of Woolmer Forest.

The Petersfield localities lie in a ring of about two miles radius from the town: Steep due north; Sheet to north-east on the London-Portsmouth road, Durford Abbey due east, Buriton due south, overhung by Butser Hill on the west. The Forest of Bere is about eight miles south of Petersfield; while in the south-west quarter lie Hambledon, seven miles, and Droxford and Soberton about eight miles distant. Petersfield is about twenty miles from Southampton and about sixteen miles from Winchester: Idsworth is about six miles south. Bursledon Ferry over the Hamble is about four miles from Southampton.

In Goodyer's day, according to a survey of the manor of East Meon taken on 31 July 1647, 'The "bacon" (beacon-fire) on Butser Hill was usually supplied out of the coppices of Hyden woods both with timber and fuel. Stroud Common was overgrown with bushes which the tenants claim a right unto for making and mending their fences, but the great wood belonging to the lord was of late destroyed except some very little and young oaks all at present not worth above 30s.' In the defining of the boundaries of the manor, several large trees are mentioned which must have been well-known landmarks to Goodyer. They included the yew-tree at Wheatham Green, 'a great oak standing in the midst of Chescombe, and so abutting upon the manor of Berriton and Mapledurham upon the south-east', and a great ash standing on the side of Butser Hill.¹

We have no information as to Goodyer's schooling. We know that he was not educated either at Winchester

¹ *Vict. County Hist. Hants*, iii, p. 67.

or at Oxford: it may be that he went to the local Grammar School at Alton.

His translations of Theophrastus and Dioscorides show that, like many men of science, he appreciated the classics, and that he was a considerable Greek scholar for his time. His library included Latin, Greek, French, Italian, Dutch, and German books: his notes show that he was well able to find his way about them, even though he was not very familiar with 'Duch and the Teutonick'.


As regards his walk in life we have evidence from many scraps of paper, often torn, upon which he scribbled botanical notes, drafted letters, kept accounts, wrote prescriptions, &c. The original writings upon these papers generally relate to legal proceedings, village affairs, the collection of tithe and taxes, while two or three documents definitely associate Goodyer with Sir Thomas Bilson, Knight, of West Mapledurham, as his steward or agent.

Goodyer's service with the Bilsons is of some importance, because all who have hitherto written of him have followed Johnson in describing him as John Goodyer of Mapledurham, thereby suggesting that he was a landed proprietor in that place. Moreover, the name by confusion with the well-known Mapledurham on the Thames, has led to the concealment of John's identity, as, for instance, when the editor of the *Flora of Hampshire* remarks in 1883 that as Maple Durham is in Oxfordshire, Goodyer's Maple Durham was a possible misprint for Maple Durwell, near Basingstoke!

A clear account of the exact relations of the Hampshire Mapledurhams is given in the *Victoria County History of Hampshire*. They are situate in the parish of Buriton in the hundred of Finchdean. From an original manor of Mapledurham, dating from before the conquest, were divided a chief manor of Mapledurham, formerly held by the Gibbons, and now by the Bonham-Carter family, and the manor of West Mapledurham which Bishop Bilson purchased: it is now held by a member of the Legge family. Then there is Weston, a tithing in the parish of

Buriton, which is believed to have been roughly co-extensive with the manor of West Mapledurham, and likewise belongs to Mr. Legge. The old manor-house of West Mapledurham was pulled down in 1829, and there is no tradition of any other old house in the district. The present Farm House of Weston was built in 1776.¹

Sir Thomas Bilson's father, Thomas, Bishop of Winchester 1597-1616, purchased the manor of West Mapledurham in



MAPLEDURHAM HOUSE.

1605 from the widow and sons of the recusant Henry Shelley. On the death of the Bishop in 1616, the manor descended to the eldest son Thomas 'aged twenty-four and more' and on the latter's death in 1649 to the second son Leonard.

The manor-house was a house with a history. During the occupation of the Shelleys it had been a centre where Papists foregathered during the latter part of the sixteenth century. There in 1586 the recusant Edward Jones used daily to 'consociate withal and heard mass every day'. There were 'priest's holes' which must have been a great joy to the Bilson boys: 'there is a hollow place in the parlour by the livery cupboard where two men may well

¹ Information from Capt. P. Seward.

lie together, which has many times deceived the searchers'; and elsewhere 'under a little table is a vault, with a grate of iron for a light into the garden, as it were the window of a cellar, and against the grate groweth rosemarye'. It is said that sometimes as many as six or seven priests were in hiding at the same time.

Goodyer may have entered the service of Sir Thomas in 1616 or 1617; he was certainly working for him for the next seven or eight years, and he may actually have been dwelling in Mapledurham House at the time that he was corresponding with Johnson in 1632-3. Thus he would have been correctly designated as 'of Mapledurham', though not as a landed proprietor there.

Sir Thomas Bilson of Mapledurham, Knight, had married at Wickham, 6 August 1612, Susanna the youngest daughter of Sir William Uvedale of Wickham, Kt. (a surname which occurs among Goodyer's notes), having issue Thomas Bilson of Buriton, born 1614, who married Edith Betisworth of Roegatt in 1640,¹ and Leonard, baptized 5 December 1616, who was named by Goodyer as one of the executors of his will, and whose monument may be seen at the west end of Buriton Church.

How long Goodyer remained with Sir Thomas we do not know. We deem it certain that he had periodically to visit the neighbouring towns and outlying farms, and even to ride up to London on his master's business. But botany was his hobby, and he probably endeavoured to combine so far as possible business with pleasure and visits to his friends' gardens with sittings in courts.

It would be easy to interweave writings of contemporary local interest into the life of our hero. The lists of the villagers assessed for tithes and taxes, the picturesque roll-call of Armour-bearers and Spearmen and 'Peionors of Beritun' (= Pioneers of Buriton), were probably all part of his daily life, but we feel bound to keep within limits, and have relegated most of the contemporary documents

¹ *Bishop of London Marr. Lic.* For other children, see p. 96 note.

to the Appendix. At the same time these documents probably illustrate his varied avocations more definitely than anything else that has come down to us, and his biography would be incomplete without a mention of them.

Of special interest are :

1. A Deposition of Arthur Hyde of Weston in the parish of Buriton concerning Weston Farm in the possession of Sir Thomas Bilson, Knight (p. 375).

2. A Petition from Fra. Waller 'most humbly intreating yo^r good worship S^r Thomas Bilson and to you and yo^r man Ma^r. Good-yeer greeting . . .' (p. 375). Written before 19 July 1621.

3. Notes of acres held by S^r Tho. (Bilson) and ten other persons in an unmentioned parish (p. 379).

4. Receipt for £250 received by W. Inkferbie and Richard Bell from Sir Tho. Bilson, Knight, at his Mansion House called West Mapledurham in the county of Southampton (p. 374). Dated 15 November, but unfortunately the year, possibly 1620 or 1622, is not mentioned.

All these and other papers are endorsed with notes in Goodyer's hand ; and even more convincing evidence as to his occupation is supplied by the draft of a letter, which we provisionally assign to the year 1618.

Sep.	3-4		
	5-6	hervest	at my being w th you I spak of
Oct.	11-12	moving cocks	Canary Wine w ^{ch} my masters mother
	12-14	lay by Coppels	M ^{ris} Bilson was used to . . . for to
	15-16	mow gard. & cutt hedges	you, you told Mr. Hall had not made
No.	14	cover hartichokes	payment for all that had fetch in her

name, now he protesteth he hath. You then faithfully promised me if my master S^r Thomas Bilson did send for any, you would send him of your best, which you affirmed to be as good as any was in England. For my lord Bishoppe Bilson was wont to comend your tast. Nowe my master entreates you to send him by this bearer Robt Palmer who you comend for his honesty, 4 or 5 gallons of such Canary Sack as you promised me to send in a sweet vessel. And the next weeke after we know what quantum you have sent you shall receive your money by the same bearer.

And so I rest

Your loving freind,

JOHN GOODIER.

[MS. f. 6.]

It is likely that Goodyer found a distraction from thoughts on the grievous troubles of the times in his botanical studies. During his early years the bitter strife between Papists and Protestants reached an extreme intensity, as first one and then the other of the conflicting parties gained the ascendancy, but in all his multifarious writings there is no note as to the side to which he belonged. It may be assumed that he shared the general tendency of scientific men to take a 'philosophic' view of life, showing some disregard of the petty, transient events which chiefly absorb the attention of mean minds. He appears to have been a man who felt most at peace when his thoughts were reposing on the larger and more enduring aspects of the moral and material world. Yet all round him was turmoil. In his own county he would have known many who could have told lurid tales of the heavy blows of the 'Hammer of Heretics'. As a young man he might have seen one of the greatest Englishmen, the immortal Raleigh, undergo trial, imprisonment, and execution, and the Pilgrim Fathers driven abroad to seek that most elemental of all liberties, the liberty of worship: they sailed from Southampton in 1620. In the prime of life the Civil Wars robbed him both of his best years for scientific work, and of his great friend, Dr. T. Johnson, killed at Basing House, to the great loss of natural science.

There is always a danger in reading into fragmentary documents more than was really meant, and yet there is a great temptation to a Magdalen man to recall an incident relating to a Magdalen College choir boy who became a Bishop of Winchester and Visitor of the College. The story gives one a vivid idea of the troubled state of Hampshire in the boyhood of John Goodyer, and greatly enhances the human interest attaching to his papers about 'pioneers'¹ and about Vachell² and Uvedale.³ Thomas Cooper, translated from Lincoln to Winchester in 1584, was sorely troubled by the number of Romish recusants

¹ p. 380.

² p. 376.

³ p. 161.

in Hampshire, and made it a matter of conscience in 1586 to petition 'for certaine Orders to repress the bouldness and waiewardnes of the recusants in the Countie of Southampton', and also that 'an hundred or two of obstinate recusants, lustie men well hable to labour, maie by some convenient Commission be taken up and sent into Flaunders as Pioners and labourers, whereby the Country shall be disburdened of a companie of dangerous persons, and the residue y^t remaine be put in some feare y^t theie maie not so safe revolte as now they doe'.

The council turned a favourable ear to the bishop's appeal, and wrote to the sheriff and certain of the justices authorizing the suggested sudden searches and ordering them to follow the bishop's directions.¹ That is why the occurrence among the Goodyer papers of a list of pioneers of Buriton carries with it ominous suggestions.

Bishop Cooper died in 1594, when John Goodyer was two years old, and was succeeded by Thomas Bilson, with whose son, Sir Thomas Bilson, Goodyer was most closely associated. Recusancy was still being punished by imprisonment in Winchester gaol, but the prisoners benefited by the general sympathy of the public and were frequently released. Indeed, Bishop Bilson found that the manor of Woodcot, Hants, given for the safe keeping of the gaol, had actually been inherited by a recusant, one Anthony Uvedale, lately deceased, and had passed to his seven-year old grandson, Anthony Brewning, whose parents were recusants. The penalty for the wealthier recusants, that was enforced about 1590, was the seizure of two-thirds of their land; and among those whose names appear in the Recusant Rolls at the Record Office are Anthony Uvedale (the hereditary keeper of Winchester gaol) of Woodcote, near Alresford, and Stephen Vachell of Heath House, Buriton, both of whom are mentioned in the Goodyer papers.

A few years later an evil system prevailed of farming

¹ *Acts of Privy Council*, 1586-7, p. 125; *Vict. County Hist. Hants*, ii, p. 82.

out the recusancy fines for a fixed sum, and the payments by Henry Shelley of Petersfield, among others, were allotted to certain servants of James I.¹

Goodyer was in the habit of jotting down lists of plants and odd notes on the backs and covers of letters, on petitions, or on any stray scrap of paper that came handy. In a way this is fortunate, for often the date of the document helps us to date the notes which he put upon it. Intermingled are the figures of calculations, shopping lists, series of days of the month with the Sundays ticked, names or plants and books, notes for excursions, medical prescriptions, names of litigants, taxpayers, and the like.

The sums done on the papers show that his arithmetical practice did not include the 12 times table, but that he first multiplied by 2 and then by 10, and added the results as in long multiplication.

It is through these papers that we infer Goodyer to have had a training corresponding to that obtained in a solicitor's office at the present day.

The oldest document in the collection, a fragment, bearing the name Edward Cole, is dated when Goodyer was only sixteen years of age, but it would probably have come into his hands at a later date as part of an account with which Sir Thomas Bilson was concerned. Edward Cole may have been the Mayor of Winchester who in the year before the Armada contributed £50 to the war fund, and who is still remembered in Winchester as the founder of Christ's Hospital. On the back of this paper Goodyer jotted down a botanical note dated 1624.

Then there is an Order from one of H.M. Justices of the Peace residing at New Alresford, but holding court at Bishops Waltham, to John Rowland of Ropley, requiring him to take his 'Corporall oth for the dewe Execution and performance of the office of Tithingman-shippe for the Tithinge of Roplie'. Dated 23 September 1614. This document is written in a clerkly hand: whether John Goodyer's or

¹ *Dom. State Papers, James I*, xlix, pp. 54-80.

not does not much matter. And there are several similar papers. The deduction we would make is that the training which he received at this period left its mark on all his botanical work. He acquired an exceptionally neat handwriting, and learnt the value of methodical habits and the importance of dating every note and recorded circumstance relating to his observations. His descriptions of plants are most conscientiously dated. Many of his books are clearly marked with the day and year of purchase, the cost of binding, and the cost of carriage from London, even though this came, as in one case, to £0 os. od.

Sometimes the exact hour of the day at which a piece of work was begun or finished is recorded with the precision that might have been expected of his astrologer contemporaries. But there is no evidence that Goodyer himself believed in horoscopes: his writings are severely scientific records of actual observation.

1616

From his practice of dating his books, we know for certain that Goodyer was contemplating the scientific study of botany in 1616. During the winter of this year he added some very important works to his library. In each case the date and price are neatly written inside the cover of the book.

		<i>s. d.</i>	
13	Novemb. 1616	2 6	Clusius, <i>Curae posteriores</i> , 1611.
12	Decemb. 1616	16 0	{ Clusius, <i>Rariorum plantarum</i> , 1601. Pona, <i>Monte Baldo</i> , fol. 1601.
15	Feb. 1616	15 0	
28	Feb. 1616	3 6	Bauhin, <i>Phytopinax</i> , 1596.
10	Marcii 1616	4 0	2nd part
12	Mar. 1616	4 0	1st part
17	Mar. 1616	1 6	the bindinge them together

} 9s. 6d. Lobel,
Plantarum, 1605.

The conclusion is irresistible. Such books were not to be bought in Hampshire: Goodyer must have spent the winter in London, and have begun to make a serious study of the literature of botany. He was twenty-four years of age; and if, as we surmise, he were already in the service of

Sir Thomas Bilson, whose father, the bishop, had recently been buried in Westminster Abbey, he might well have gone up to London on his master's business. A paper that throws much light on this visit to London is one on which he wrote out a list, dated the 24th and 25th of March, of the more remarkable plants in the garden of Mr. Coys of Stubbers, a place a few miles east of London in Essex. On the same sheet of paper are a list of drugs and two London addresses, 'de Laune in ye black friers', and 'Mr. Cole y^t married Mr. Lobel's daughter in Lyne Street'. This last is a most interesting note, because it gives the correct spelling of a name which from the time of Pulteney onwards has generally appeared in literature as 'Coel'. James Cole was a London merchant, mentioned by Johnson as exceedingly well experienced in the knowledge of simples.

The fact of Goodyer's visiting James Cole at this time would have a special significance. The eminent botanist Mathias de L'Obel, Cole's distinguished father-in-law, who had been living with him, had just died, leaving his botanical writings to his son-in-law. Lobel, as he wrote his name in this country, was the youngest of the triumvirate of great Flemish botanists, Dodoens, Clusius, L'Obel. He had brought to this country the learning of his master Rondelet of Montpellier and the botanical illustrations of that prince of printers, patron of botanists, Christof Plantin of Antwerp; he had made a special study of English plants and during his last years had been engaged on a new botanical work, the *Illustrationes Stirpium*. To Goodyer the name of Lobel, like that of Gerard, was probably a household word: he was known as the only botanist in Britain on whose scientific accuracy a student could rely. There is nothing more probable than that Goodyer, learning of his death on 3rd March, would lose no time in visiting Cole, and so get to hear of the manuscripts, which later, after the death of their editor How, did eventually come into his charge and were bequeathed

by him to Magdalen College. In addition to the copy of the *Plantarum Historia* mentioned above, and purchased within a week of the death of Lobel, Goodyer owned five other editions by the same author: one of these, the edition of 1576, was a presentation copy from Lobel to Dr. Martin Ramerus or Rhamneirus, a Licentiate of the Royal College of Physicians. A copy of the *Icones Stirpium* with notes by William Mount, and with some figures coloured, may have been among Goodyer's first botany books as a boy. We know that he could draw and that he painted in water-colour.

Goodyer's miscellaneous papers show that even while he was occupied with the business of estate management, or of the Courts, or in dealings with tithing officers, his mind would always be turning to the plants which he had seen in the gardens of his friends or had found on his excursions.

The earlier notes are scrappy, but of great historical importance, for they place our knowledge of early English gardening on a far surer basis than heretofore. The notes become progressively more methodical until in 1621 their number and finish show that descriptive botany had become the principal object of the author's life. Whether Goodyer ever had any thoughts of publishing these descriptions under his own name we do not know, but of many he made both rough and fair copies: some of the latter he handed over to his friend Dr. Thomas Johnson for inclusion in the new edition of Gerard's *Herbal* to be mentioned in 1633.

His first gardening friends included Parkinson, Coys, and Franqueville, who have often been mentioned in histories of early gardening, but now for the first time have we anything in the nature of lists of the plants actually growing in their gardens. Last year when editing the long list of plants grown by Walter Stonehouse in 1640 at Darfield in Yorkshire, it was pointed out that only three earlier garden lists were then known, those of the Holborn

garden of Gerard (1596), of George Gibbes' garden at Bath (1634), and of the Lambeth garden of the elder Tradescant (1634). We are now able to add several other lists of intermediate date which comprised the more novel plants then being grown by horticulturists.

The earliest of Goodyer's garden notes go back to the year 1616, when he first gathered seeds of *Astragalus lusitanicus* 'in the garden of my good friend Mr. John Parkinson an Apothecarie of London Anno 1616' (f. 107), and about the same time he noticed in the same garden in Long Acre, *Pisum arvense*, which he called 'P. maculatum Boelii' and 'Ervilia silvestris Dodonaei' (*Lathyrus Ochrus*). The two former were no doubt two of the many new plants introduced into English gardens by Guillaume Boel from Portugal and Spain in 1608. Parkinson has put it on record that Boel 'gathered there about two hundred sorts of seeds . . . of all which seeds I had my part, and by sowing them saw the faces of a great many excellent plants but many of them came not to maturitie with me, and most of the other whereof I gathered ripe seed one yeare, by unkindly yeares that fell afterwards, have perished likewise'.¹

Goodyer grew all three plants from seed in his own garden at Droxford and described them in 1621.

The next garden of which we have his notes was that of Wm. Coys² at Stubbers, North Okington, in Essex. It was already old-established, and well known to Gerard in 1597 for its exotic plants, and was visited by Lobel during the early years of the seventeenth century. The latter botanist was moreover greatly impressed by Coys' methods of brewing beer and ales.

We are glad to think that this garden is not only still in existence but is in the possession of the family which succeeded the Coyses at the end of the seventeenth century. Through the curtesy of the present owner, Mr. Champion Branfill Russell, I was privileged to make a pilgrimage

¹ Parkinson, *Theatrum*, p. 1108.

² See p. 312.




The Lime Avenue



Common Elm 'Witch' Elm

Elms in Spring



Plan

THE GARDEN AT STUBBERS

to Stubbers this spring, and there on the day of the eclipse of the sun to celebrate the tercentenary of Goodyer's visit. The gardens were much altered by Repton at the end of the eighteenth century, and the house has been practically rebuilt, but sufficient remains to indicate the general arrangement of the old sites.

The flower garden of William Coys almost certainly coincided with what is now known as the Mulberry garden at the south-east corner of the house. The middle line of an old enclosed garden is indicated by a fine avenue of small but ancient limes, whose great age is hardly apparent at first sight. The trees were pollarded early, and their trunks being bark-bound have ceased to grow for many years. A very old-fashioned pink rose, a plant of which has been sent to Shakespeare's garden at Stratford-on-Avon, may also date from Coys' time. And to my great joy I found a more certain link with him in the pretty little Ivy-leaved Toadflax that is still growing on the older walls around this classic spot, where it was first grown as a garden plant in England and before it had become the common wall-weed that it now is. Another plant that may be directly descended from Coys' garden is the Yellow Fig-wort (*Scrophularia vernalis* L.), which comes up sporadically in the borders.

But of even greater botanical interest are the elms on the west side of the garden. At the time of my visit, one differed in the most conspicuous manner from its neighbours by being later and quite leafless while they were almost in full leaf. Had Goodyer seen the elms between Romford and Stubbers under the same circumstances, he could hardly have failed to remark on the matter to Coys, who would then have informed him of the local name 'Witch Elm', and would have expounded the special qualities of its wood, which was more 'desired for naves of carts' than common elm.

The garden of William Coys has always been famous in the annals of horticulture because it was there in 1604

that the *Yucca* first flowered in England. Now, thanks to Goodyer, we are able to print a list of 126 names of plants that were growing at Stubbers in March 1616-17, two lists including plants given by Coys to Goodyer in 1621 or 1622, and a longer list in which 324 plants are marked with the letter C (= Coys).¹ This last list was made by Goodyer between 1618 and 1625.

The earlier of these lists of Coys' garden is the oldest known MS. list of an English garden, in which the plants are properly distinguished by their scientific names. I have no doubt but that the names, which are in Goodyer's handwriting, were supplied by Coys, who may have determined many of them himself. In the history of English horticulture it therefore comes between the printed Catalogue of the Trees, Fruits, and Plants grown by John Gerard in his garden at Holborn² and the lists of George Gibbes and John Tradescant, both of which are printed below. It is entitled

Mr. Coys his garden 24 and 25 of March 1616, 1617.

Then follow ten names of plants distinguished with *g* (= Gerard), and then twenty-four others which may or may not have formed part of the same collection. The latter include the American novelties :

Battata Virginiana.
 Prunus Virginianus. Pishimon.
 Cerastus Virginianus.
 Vitis Virginiana.
 Anonymos Sumas Virginianum.
 Solanum Virginianum.

In after years Goodyer received many plants and seeds from Mr. Coys, which he grew in his Hampshire garden at Droxford, and described in detail when they flowered.

¹ See p. 387.

² Gerard's printed *Catalogus arborum*, &c., of 1596 is only represented by a single copy, which has been admirably edited by Dr. B. D. Jackson in 1876.

1617

It was inevitable at the outset that the medicinal interest of herbs should be much to the fore. Botanical lore was practically confined to physicians, apothecaries, and herbalists; and perhaps this strong human interest supplied the motive power for the beginnings of the science of botany. All English botany books of early days were practically treatises on *Materia Medica*, and even Parkinson found a medicinal use for nearly every plant.

Goodyer's development took the same course. The notes made during his early period show that a medicinal interest in herbs was certainly present to his mind, though it may not have provided the principal incentive to his studies. In 1617 he abstracted the names of forty-six exotic medicinal plants and products from Lobel's writings, and added the following list of drugs :

Terra sigillata	Tartar
Gum Elemni	Ladanum
Aurmi vulgar	Castor
Virga aurea	Paeina ppi
Mirrh	Bac mirti
Bolus armenic	Acacia
Caious	Spodium
Borax	Bdellium
Litharg. auri	

And on 28 April 1617 he added :

2. Corall dissolved & swetned with Juyce of Granates
 - Syrup of Rheubarbe made of ye extracte
 - The Extracte of Rhubarbe
 - Pearle dissolved
 - Liquor solutionis Scamonii 1610 23 Marcij
 - Balsamum
 - Scamonium corrected 1610 23 Marcij
 - Hirae simpl. ʒiiij pro quat. viribus
 - Manna 2 oz.
 - Santali Citrini incisi ʒij
 - Olibanum
 - Epithimum ʒij
 - Oximel simpl.
 - Dia Codion simp.
 - Dia moron.
- de Laune in y^e black friers.

[MS. f. 24

The study of drugs was evidently going strong about this time, for in this very year, 1617, the Society of Apothecaries of London, the *Societas Pharmaceutica Londinensis*, severed its connexion with the Grocers' Company, and received a new charter from James I. Goodyer appears to have obtained his list of drugs from one of the early masters of the Society, Mr. de Laune,¹ an apothecary in Blackfriars, whose name and address he noted on the back of the paper, and who is known as having sent from Burgundy to Gerard plants of *Gentiana maior*, the Great Felwoort or Bald money, now known as *Gentiana lutea* L.

The two Scammony preparations are dated 1610, when he would have been eighteen years of age. And in the first list the figure 16 before five drugs may show that he became possessed of them in that year. The five are *Hermodactyli non venenati offic.* Lo. 146; *Rha Ponticum antiquorum* Lo. 289; *Aloe* Lo. 374; *Glans unguentaria* Lo. o. 101; *Araba lacca*.

Towards the close of his life he again revived his interest in medical matters. He bought medical books, he wrote out prescriptions, and he copied notes from medical

¹ By finding the will of William Delaune (P. C. C. Wood 23) at Somerset House I have obtained evidence that the de Laune visited by Goodyer is likely to have been the son, Paul de Laune. WM. DELAUNE, Preacher of the word of God and Physician, willed that he should be buried near his late wife. To Gedeon his eldest son he left his lands in Blackfryars near Ludgate which he had purchased of Sir Wm. More; to poor of French church in London £4; to poor of Blackfriars and of Church of Norwich and Deepe 20s.; to kindred beyond the sea £6 (to be administered by s. Nathaniel); to s.-in-law Mary Desloges, widow of Cornelis Tance, £3; to s. Paul the newe house, greate Court & house of office, and the Apothecary stuffe and furniture.

The relations mentioned include sons, Gideon, Peter, Nathaniel, and Paul; grandsons Abraham s. of Gideon, and Henry s. and h. to Isaack Delaune dec.; das. Sara, Ester, and Elizabeth; sons-in-law Chambleyn and Nathaniel Mary. The will is dated 20 November 1610; proved 12 March 1610. Both William and Paul were on the Roll of the College of Physicians. Gidcon de Laune made money as Apothecary to Mary, the Queen Mother, and died worth £80,000. Shakespeare's friend, Sir W. Davenant, was his great acquaintance (Aubrey, *Lives*). Paul is believed to have died in Jamaica, c. 1654: he was a relative of Dr. John Argent see p. 250.

authors into the margins of his herbals; all of which goes to show that he was turning his knowledge to practical use by attending to the medicinal needs of his neighbours.

But John Goodyer, though mindful of the medicinal and economic value of plants, clearly saw that the science of botany could not be advanced without detailed morphological descriptions from living plants. His ideals were those of his great predecessors Fuchs and Brunfels: but whereas they put their observations into their drawings, Goodyer put his into descriptions.

One of his local discoveries of 1617 was the Woolly Thistle, 'Corona fratrum quorundam' (*Carduus eriophorus* L.), which he found 'wilde neare London highwaie on the East part of Haliborne in Hampshire'.

Another was the Purging or Cathartic Flax, or Mill-mountaine, a plant of known medicinal virtue which used to grow plentifully in localities where 'Carduus acaulis septentrionalium L'obellii' was also found, namely, 'in the unmanured inclosures of Hampshire on chalky downes and on Purfleet hils in Essex, and in many other places'. He owed his information concerning the properties of this plant to one of his apothecary acquaintances.

'I came to know this herbe by the name of Mil-mountaine, and his vertue by this meanes. On the second of October 1617 going by M^r. Colsons shop an Apothecary of Winchester in Hampshire, I saw this herbe lying on his stall, which I had seene growing long before: I desired of him to know the name of it, he told me that it was called Mill-mountain, and he also told me that beeing at Doctour Lake his house at Saint Crosse a mile from Winchester, seeing a man of his have this hearbe in his hand, he desired the name; hee told him as before, and also the use of it, which is this.

Take a handfull of Mill-mountaine, the whole plant, leaves, seedes, flowers and all, bruise it and put it in a small tunne or pipkin of a pinte filled with white Wine, and set in on the embers to infuse all night, and drinke that wine in the morning fasting, and hee said it would give eight or tenne stooles. This Doctour Lake

was afterward made Bishop of Bath and Wells, who alwaies used this hearbe for his purge, after the said manner, as his man affirmed.¹

But the outstanding event of this year was the introduction of the Jerusalem Artichoke to English gardens




MILL-MOUNTAIN.

and cookery, undoubtedly the result of a visit to the London garden of John Franqueville. Of this garden we are also able to give a list, but as Goodyer was there at

¹ *Ger. emac.* 560. For Goodyer's description of the plant, see p. 112. The use of Honewort, as Corn Parsley was locally called in Hampshire, steeped in beer to cure swellings of the cheek, was also duly recorded by him, and will be again referred to under 1625.

the end of the year, when it would be looking its worst, the list is short (p. 326).



THE JERUSALEM ARTICHOKE.

We here quote his description of the circumstances of his success with the new 'wonderfull increasinge' vegetable, leaving the account of the plant for a later chapter.

'Where this plant groweth naturallie I knowe not. In Anno 1617 I received two small rootes thereof from Master *Franquevill* of London, no bigger then hens eggs, the one I planted, the other I gave to a friend, mine brought me a peck of rootes, wherewith I stored Hampsheire.'—MS. 11, f. 117; *Ger. emac.* 754.

The friend is almost certainly William Coys, for whom he afterwards wrote out a very long list of plants (17 pages), with notes and references to the works of Lobel and Gerard, and to their occurrence in the gardens of Coys, Parkinson, and Franqueville. In this list after the entry relating to the Artichoke '*Heliotropium indicum vel virginianum*', he added the note, 'you had lately planted it when I was at your house 25 Martii 1617'.

Goodyer's quaint description of *The Vertues* of the Artichoke have often been quoted: but it is a story that does not lose in the repetition.

'Theis rootes are dressed divers waies; some boile them in water, and after stewe them with sack and butter, addinge a little Ginger: others bake them in pies, puttinge Marrow, Dates, Ginger, Reasons of the Sunne, Sack, &c. Others some other way, as they are led by their skill in Cookerie. But in my iudgement, which way soever they be drest and eaten they stirre and cause a filthie loathsome stinking winde within the bodie, thereby causing the belly to bee pained and tormented, and are a meat more fit for swine, than men: yet some say they have usually eaten them, and have found no such windie qualitie in them.'—MS. 11, f. 117; *Ger. emac.* 754.

But before making any attempt to account for the tastes of past generations, it is necessary to know precisely both how they prepared their food and how they arranged their dietary. The transition from the age of beer to that of tea and coffee must have profoundly modified the national palate. And this is well illustrated by a very striking instance of an acquired taste, quoted by Goodyer's contemporary Parkinson. Chocolate, he described as a drink 'well pleasing and accepted with the greatest among the Indians, who account nothing of more esteeme; but to the Christians it seemeth a wash fitter for hogs, yet by use even accepted by them also in the want of better'.

The cultivation of the other, the Globe Artichoke, appears to have spread through the southern counties of England some half a century earlier, and the vegetable that figured in legal documents in Goodyer's lifetime is more likely to have been the Globe than the Jerusalem Artichoke. For in one of the Bodleian charters relating to Kent there is a stipulation that the tenant of a garden in Wateringbury is not to have 'the benefit of the Sparrow Grass beds and Hartichoaks in the Garden'. An Hartichoke Garden, 44 perches in extent, was one of the features of Henrietta Maria's garden at Wimbledon, and at the time of the survey of 1649 contained plants and roots to the value of £1 10s.¹ Her gardener probably grew both sorts.

Every one who has hitherto written about Goodyer appears to have tacitly assumed that he spent his early manhood at Mapledurham, and that there his first experiments in horticulture were made. This, however, was not the case, for although he was no doubt often at Mapledurham, his home and, what interests us more particularly at present, his garden were at Droxford. It was Droxford, therefore, that must have the honour of having been the first village in England to produce the new Artichokes in quantity. How completely all recollection of this important horticultural event has disappeared, is proved by the fact that even the modern 'discoverer' of John Goodyer, Canon Vaughan, did not know that his botanical predecessor had ever dwelt in Droxford. Yet Canon Vaughan, while Rector of that parish, when engaged in researches into the forgotten details of Izaak Walton's life there, passed the history of Droxford through a hair sieve. But then Hampshire is a forgetful county. Do not the lives of Walton, of Goodyer, of Gibbon, and above all, of Jane Austen show it?

The papers that have been reposing for the last two and a half centuries in the manuscript room of Magdalen are evidence that Goodyer lived and gardened at Droxford,

¹ *Archaeologia*, vol. x.

but unfortunately their information goes no further. There is no hint as to where his house may have stood, or where the first Artichokes increased so amazingly. The village, writes Canon Vaughan, is but little changed since the time when Izaak Walton spent the last days of his life resting in 'the cool shade of the honeysuckle hedge' and watching the moorhens 'on the gliding stream'. The old mill is still standing, on the bridge of which 'the aged angler must often have lingered, as he watched the rush of water making pleasant music beneath his feet' and thought of his fishings in swift streams 'full of great stores of trout'. Nor would it have been very different in Goodyer's early manhood. Then, as now, the church was flanked on either side by the Manor House and the Rectory, with their gardens, orchards, and pastures sloping down to the clear running Meon. The underground tunnel from the Manor House below the Rectory garden to the house beyond, may or may not have existed: at present it is only a reality in local tradition. Of greater permanence are those details of uncultured nature, of the birds and flowers, which the Canon has described so well: his sketch of the wild life of 'Longmead' bordering the Meon is for all time: that is Droxford as Goodyer knew it. He would have recognized the spot where the green Alkanet (*Anchusa sempervirens*) puts forth its rich blue flowers and he would have revelled in the rare beauty of the Yellow Meadow Rue in its season, but it is doubtful whether Coral Root would have escaped him, had it been growing near Church Mead, where Canon Vaughan tells us it grew within recent years. Certainly there is no evidence that Goodyer knew it in Droxford before he found it at Mayfield. Botanists tell us that Coral Root is only found in this one locality in the whole of Hampshire, and we may not unreasonably deduce that Goodyer will have followed his usual practice of lifting roots found away from home and planting them in his garden. The Hampshire colony may be descended from a garden-escape. Is it a clue to the site of his garden?

In after years Goodyer cultivated many other new fruits and vegetables, including the Virginian water-melon or Pompion, 'no bigger nor larger than a great apple', but none are now so well known as the Artichoke, his first horticultural triumph, and one with which his name will always be connected. Indeed, it may be that this early success inspired those assiduous searchings for new and rare plants that characterized the whole of his active career.

1618

Several papers dated 1618 show Goodyer's connexion with the Bilson household, probably, as we have said, in the capacity of steward or agent. In the corner of a roughly scribbled letter, printed on page 9, is a fragment of a diary, with a reference to the fact that his 'hartichokes' required protection in cold weather.

At the same time his interest in field botany was steadily increasing. He made several journeys on horseback in his own and neighbouring counties and up to London, perhaps on his master's business, but certainly to his own profit and advancement as a botanist.

On one of these visits he would almost certainly have again visited the well-known garden in Long Acre belonging to John Parkinson who, like Coys, had received many rare plants collected by William Boel in Spain in 1607, an acquisition all the more desirable since many of his own fine plants had perished during the 'most violent frosty winter' that preceded. In 1618 Parkinson was gloating over his various horticultural triumphs, including the Great Double Yellow Spanish Daffodil which he claimed to have been the first to grow.¹

On the 9th of April Goodyer noted 'Cowslipps 2-in-a-hose' and 'Primrose 2-in-a-hose' at Sheet, probably in the garden of William Yalden, who almost certainly lived in a new house by the mill on the upper waters of the Rother. The

¹ Parkinson, *Paradise*, p. 103.

house is still standing, and even though more than three hundred years old, is a comfortable home. A stone in the wall records the year 1606, when it was built, and the year 1867, when it was repaired. The timber-framed walls of the upper story are partly filled in with wattle and dab, and partly with a warm red brick. The old kitchen chimney is so wide that it can only be swept by climbing inside; and there is, it is said, but one man now living who is willing to strip to his trousers and bring down the soot. This was the home of Rose Yalden, Goodyer's sister: plants in the garden at Sheet are oft-times mentioned in his papers, the last being some Sweet Potatoes in 1637.

To plants found near his own home at Droxford, he frequently refers. In a wood by Strugnells in the Thetcher, near Droxford, he found *Lunaria minor* on 21st May, and again on 1 June. These are the first notices of Moonwort in Hampshire. He records plants collected (and incidentally his own movements) at Chawton¹ (10 June), Sheet² (7 July), Chawton³ (18 July), Tichfield⁴ (20 July), Hounslow Heath⁵ (30 July). In August he was travelling in Wiltshire, visiting Warminster⁶ (21 August), and Venny Sutton,⁷ now Sutton-Veny (27 August). And nearer home he visited Winchester, Southampton, and Emsworth.

At Winchester he noted that 'Willowe' was the popular name for 'Chamaenerion Gesneri', which is therefore the first evidence of the Willow herb (*Epilobium angustifolium* L.) in Hampshire. It is a plant which has since been

¹ 'Herba Paris, some with five and six leaves.' The first notice for Hants of *Paris quadrifolia*: Gilbert White noted it again in 1778.

² 'Scabiosa minima hirsuta.' Probably the first record of *Jasione montana* L.


³ 'Jacea flore albo.' Perhaps a white variety of the Greater Knapweed (*Centaurea Scabiosa* L.).

⁴ 'Eryngium marinum.' First evidence of the Sea Holly (*Eryngium maritimum* L.) in Hants. Eryngo roots candied were a favourite delicacy in the days of James I, but they were then obtained from the East coast. Sir John Salusbury, c. 1580, knew their medicinal value.

⁵ 'Plantago aquatica stellata.' First record of *Damasonium Alisma* Mill., the rarest of our Water Plantains.

⁶ 'Colchicum flo. albo et purpureo.'

⁷ 'Ebulus', 'called Scots Blood'. *Sambucus Ebulus* L. or Danewort.



SHEET MILL

reported as abundant in the valley from Beech Farm, where he lived as a boy, to Meadstead. He may, therefore, have known the plant near home, but may not have been familiar with its Winchester name. The Willow herb remained unrecorded in Hampshire until Gilbert White mentioned it again in his Garden Kalendar for September 1765.

A more striking discovery was that of 'Lactuca sylvestris vera ingrato odore'. 'This wilde stinkinge lettice I found wild on the walls and dry bancks of earth at Southampton. Anno 1618.' This is the first county record of *Lactuca virosa* L. He sent seeds to Parkinson, in whose garden Johnson saw it growing about 1630. A detailed description was written on 13 September 1621.

By a mill at Emsworth he found 'Anagallis aquatica tertia', the Brookweed or Water Pimpernel (*Samolus Valerandi* L.), thus making his first contribution to the paludal flora of his county. It was not recorded in Hampshire again for more than two hundred years.

During the last two weeks in August he was travelling in south Wiltshire and passed through Salisbury. The finds on this trip are mentioned in a draft letter dated 7 November and written in London. The name of the person to whom it was written does not appear, but from the fact that the names of the plants mentioned are included in a long list which Goodyer sent to his friend Coys, I strongly suspect that it was addressed to him.

S^r after my service remembered . . . I wrote unto you a letter from Droxford y^e 8 of Sept. last, and therein enclosed certaine stalks & seeds of an herbe which I found in Wiltshere somethinge like to *Lysimachia*, [I found it in Wiltshere (*erased*)] also I sent you some of the leaves & stalks of *Plantago aquatica stellata* which I found in Hounslowe Heath. I therein wrote unto you that I found *Colchicum fl. albo & purpureo* in Wiltshere, in flower in August last, I have of this roots of them at Droxford, if you want I can furnish you with some of them. I doubt you received not that letter, for that our foot-post died by the waie going towards London, since whose death we wanted a foot-post for

a year, but now we have another who did lye at y^e King's armes in Shoe Lane, the place where the old foot-post lay, but nowe he is removed to another place which yet I doe not know. I have left at M^{rs} Capells (for Mr. Capell¹ they tell me is dedd since my last being with you) a small plant of our Wich Hasell which I take to be *Ulmus latifolia lobelii*: it came from seed this last year. I have seen y^e leaves of them this somer twice as brode as I saw any of y^e leaves of our Common Elm. This desiring that God do bless you with health and happiness to your great Comfort, I comitt you to Gods protection and rest

to you my bounden

[Unsigned]

From my Lodging at the Red Lyon
in Fleet Street, London, the 7 of November 1618.

Yf at any tyme you wright to me you may direct your letters to be left at Mr. Tho. Johnsons [a Tayler (*erased*)] at the signe of the Raynedecare without Temple Barre neare St. Clements Church, where our foot-post shall endeavor to have them. [MS. f. 2]

This letter is of great interest because it is the earliest evidence for the Willow herb, 'something like to Lysimachia' (he found it on 27 August), and for the Autumn Crocus in Wiltshire. It settles the question of the priority of the discovery of the Water Plantain, *Damasonium Alisma*, as a British plant, in favour of our author rather than of Johnson. It also dates his first recognition of the Wych Elm (*Ulmus montana* Stokes) as a distinct species of English timber tree. Apparently it was growing at Droxford and was known to Goodyer by its south-country name of Witch Hazel. Now this name has been generally superseded by the northern name of Wych Elm. I have no doubt that by this observation at an early period of his career, his eye became trained and apt to distinguish the two other elms which he was the first to describe. His

¹ At a later period CAPEL was quite one of the best-known names in connexion with English gardens. Sir Henry Capel began horticulture at Kew, his brother Lord Capel made one of the most beautiful gardens of the seventeenth century at Cassiobury, and Mary Somerset (*née* Capel) had a botanic garden at Badmington, but there is of course no evidence that Goodyer's friend was 'born in the Purple'.

descriptions of the four species are reprinted on page 38, but the first draft of his description of the Broad-leaved Elm may appropriately appear here. It is written on the back of a document dated 1614.

The broad-leaved Elme or Witch Hasell groweth to be a very greate tree, and very high, especially when he groweth in woods amongst other trees, the tymber in y^e yonge trees is very white, his branches or boughes are grosse & bigge and doe spread themselves broader [and hang more downwards than those of the first]. The leaves are rougher and much broader and longer than those of y^e Comon Elme, stript or cutt about y^e edges, of a dark greene colour, nearer resembling y^e leaves of y^e Hasell, which is y^e cause of y^e name.

The one side of y^e leaves are allwaies longer and doe come nearer to y^e boughes whereon they were then the other, like unto y^e leaves of all y^e rest of y^e kind of elmes, as Lobell's figure of the Common Elm well expresseth; the leaves of the older trees are very much narrower than those of y^e younge ones, also on y^e leaves of this Elme doe growe blisters or bladders in which at y^e springe are little creatures about y^e bignes of Bed-fleas, which in y^e somer (as I take it) turn into flies growinge wings like those of the Comon Elme. The barke on y^e out side is blacker than that of y^e Comon Elme & is very tough. So that when y^e sappe is uppe, it will strippe & peelee from the lower end of y^e boughes to y^e toppe, without breakinge, whereof are made cords & ropes.

The roots spread far abroad in y^e earth & are very tough, which makes the tree with much difficultie to be grubbed uppe.

Whereon doe growe y^e flowers nearly compacted together & like unto y^e pointells or chives growinge in midle of flowers, which afterwards turn into flatt seed, more long than broade, not much unlike to Arach seed; and doe for the most parte fall away before or shortly after, the leaves springe forth, and some hang on a greate parte of the somer. They flower at the latter end of febr or beg of March.

[MS. f. 4; *Ger. emac.* p. 1481 (much altered)]

The following lines are written on the cover of a letter addressed to 'Nicholas Everender at his brother's house in Sedlescombe in Sussex. This to M^r Samuell Shute at Mr. Thomas Caltherste at ye albus Lion in Dea(?)side'. On the other side is written the draft of the letter already

quoted, dated by Goodyer from his lodging at the Red Lyon in Fleet Street, on the 7 of November 1618.

‘Even such is tyme, that takes in trust
 Our youth, our ioyes and all we have
 And paies us but with age and dust
 Who in the darke and silent grave
 When we have wanderd all our waies
 Shutts uppe the storie of our daies
 And from which earth and grave and dust
 The Lord shall raise me uppe I trust.’ [MS. f. 2

Being, regrettably, unfamiliar with the poem I consulted my friend Sir Walter Raleigh, our Professor of English Literature, in the confident expectation that he would identify it for me. The strange coincidence could hardly have been anticipated that the lines should be known as the ‘Last Poem of Sir Walter Raleigh’ of Elizabethan fame. There is a tradition that they were written in the Tower, shortly before the execution, but whether this be so or not, it is surely remarkable that Goodyer should have had a copy on the back of a letter written within a few days of the execution on 28 October 1618.

As a twelve-year old boy Goodyer must have heard of the infamous trial at Winchester, when the people first pelted Raleigh with tobacco pipes and later were won over by the eloquence of his defence and the dauntlessness of his bearing. He would assuredly have taken a sympathetic interest in his great contemporary not only for his heroic career, but as a pioneer familiar with the splendours of the tropical vegetation of the New World. Goodyer’s Artichoke culture was perhaps modelled on that of the Potato. Whether he was a smoker, we do not know, but about 1618 he paid 4*d.* for ‘tabaco’, apparently at Alresford. It was only a few years before, that James I had stigmatized smoking as ‘a custom loathsome to the eye, hateful to the nose, harmful to the brain, dangerous to the lungs, and in the black stinking fume thereof nearest resembling the horrible Stygian smoke of the pit that is bottomless’.

In 1619 Goodyer took to growing two kinds of tobacco at Droxford.

There are several lists of garden plants among the Goodyer papers, but unfortunately they have no localities or dates attached to them, with the exception of a short list of plants, without date, and not in Goodyer's hand (see p. 347). It is headed *To have from Mr. Gibbes*. Of the plants enumerated only six are noted by Johnson as occurring in the garden of Mr. George Gibbes of Bath in 1634.

Another list in his own autograph may refer to plants growing at Droxford (p. 327).

1619

Of the year 1619 there is but little to record except the discovery and description of two plants, one of which was new to the British flora. He also wrote the description of the *Linum catharticum*, part of which has already been quoted.

On the 3rd of August he found the Small Woodruff or Squinancy-wort (*Asperula cynanchica* L.), and thus he independently discovered one of the denizens of the chalk hills which had been previously, though probably unknown to him, noticed by Lobel in the Isle of Wight: 'cretaceis gaudet montosis prope Drayton e regione Vectis Insulae'. Goodyer's observation of 1619 was printed by his friend Johnson in 1633, and Lobel's earlier note was printed by his admirer, Dr. How, in 1655. The Procumbent Marshwort (*Apium nodiflorum*) is recorded on the authority of Goodyer¹ by How: we are now able to print the original description (p. 114).

A further success was the addition of a new Hedge-Parsley to our flora (*Caucalis nodosa* Scop.²). Later on Goodyer discovered the Field Hedge-Parsley 'amongst wheat plentifully near Petersfield'.³

His well-known description of the Dewberry (*Rubus caesius* L.), distinguishing it from other Brambles under

¹ How, *Phytologia*.

² Druce, *Goodyer*, p. 1.

³ Merret, *Pinax*, 24.

the name of 'Rubus repens fructu caesio', was written in September of this year, but the original MS., from which Johnson's account was printed, is not now extant. It was almost certainly taken from a Hampshire plant, but since Goodyer omitted to mention the locality, he is not quoted as a first recorder in Townsend's local *Flora*, in which seventy species of brambles are now distinguished: truly a thorny subject! Hampshire, perhaps with one exception, is stated to be richest in brambles of all the counties in England.

In his garden he raised some specimens of the forbidden plant, tobacco. 'In anno 1619 I received the seedes hereof from Mr. Anthony Uvedale¹ who that yere intended to plante greate store thereof, and was hindered of his purpose by a proclamation sett forth by Authoritie.'

1620

The next two years were perhaps the most eventful in his career as a botanist. His excursions were rewarded by the discovery of several other plants new to the British flora, and many new plants imported from the south of Europe thrived so well at Droxford, that he was able to give the first English description of them.

On the 28th of April we find him making observations on the Oak, Walnut, and Chestnut, being especially in-

¹ Anthony Uvedale, the Governor of Winchester Gaol, has already been mentioned. The Uvedales were a Hampshire family who lived at Wickham, and intermarried with the May family, living at Mayfield in Sussex.

The name of Uvedale is well known in the annals of early horticulture. Robert Uvedale (1642-1722) became master of the grammar school at Enfield, Middlesex, about the year of Goodyer's death. He was 'a great lover of plants, and, having an extraordinary art in managing them, is become master of the greatest and choicest collection of exotic greens that is perhaps anywhere in this land. His greens take up six or seven houses or roomsteads. His orange trees and largest myrtles fill up his biggest house, and . . . those more nice and curious plants that need closer keeping are in warmer rooms, and some of them stoved when he thinks fit' (Gibson, *Account of several Gardens near London*, 1691, *Archaeologia*, 1794, xii. 188). He was thus one of the earliest possessors of hothouses in England. One of his former pupils is said to have brought him from Mount Lebanon a cedar which was recently flourishing at Enfield.

The copy of Turner's *Herball*, 1568, in the Oxford Physic Garden, which belonged to *W. Clowes* in 1571, bears the signature of 'Rob. Udall.'

terested in their flowers, buds, and galls. He added to these notes and wrote them out afresh on 9 May 1622: they are printed under that date in Johnson's *Gerard*.

Business at this time brought him into touch with the lands and villagers of Soberton, a hamlet about a mile and a half south of Droxford. It was there that he gathered the 'Oenanthe Angustifolia'. This has been identified by Druce as the first record of the Marsh Parsley Oenanthe in Britain, but Townsend (*Flora*) considers that it was the Sulphur-wort Dropwort (*Oenanthe silaifolia* Bieb.) that was found.

'This 19 of May 1620 I found this wild in East Hoo in ye parish of Subberton about 7 miles from Petersfield in Hampshire in a hedgerowe about a flightshott from ye then dwelling-house of Mr. William Browne on ye south part of ye said house and ye 18 of June 1620 I saw it there in flower.'¹

Merrett² gives the locality as East How. William Browne of Hoe had purchased the manor of Soberton in 1619.³

A few days later an excursion to the New Forest yielded another new plant, the Narrow-leaved Lungwort (*Pulmonaria angustifolia*), whose fine blue flowers attracted his attention on 25 May 'in a Wood by Holbury House'. It is a rarity in Britain; Hampshire and the Isle of Wight being the only stations where it grows wild. The Forest children call it 'Joseph and Mary'.⁴

In the summer he described the following plants:

20 June	Antirrhinum minus.	<i>Linaria minor</i> Desf.
2 July	Pastinaca aquatica minor.	<i>Sium erectum</i> Huds.
5 „	Rapunculum silvestre Tragi.	<i>Phyteuma orbiculare</i> L.
5 „	Eufrasia altera Dod.	<i>Bartsia Odontites</i> L.
5 „	Anthyllis montana.	<i>Thesium humifusum</i> DC.
8 „	Carduus acaulis septentrionalium L'Ob.	<i>Carduus acaulis</i> L.
22 „	Melampyrum luteum.	<i>Melampyrum sylvaticum</i> L.
26 „	Herba Doria altera.	<i>Senecio sarracenicus</i> L.
10 Aug.	Panicum.	<i>Echinochloa Crus-galli</i> L.
12 „	Acinos.	<i>Ocimum Basilicum</i> L.
18 „	(Aster conyzoides).	<i>Jasonia tuberosa</i> DC.
	Sium siifoliis.	<i>Carum segetum</i> , Benth.

¹ Goodyer quoted by How. MS. note to *Phytologia*, p. 81.

² *Pinax*, p. 84.

³ *Vict. County Hist. Hants*, iii. 263.

⁴ J. Vaughan.

He found both the Water Parsnip (*Sium erectum* Huds.) and the Bastard Toadflax (*Thesium humifusum* DC.), near Droxford, the former growing in the River Meon, the latter 'wild on the side of a chalkie hill in an inclosure on the right hand by the way, as you goe from Droxford to Poppie Hill'. Both these references are the earliest we have to these plants in Britain, as also is his excellent description of the Honewort or Corn Parsley (*Carum segetum*), which he again wrote out for the new *Herbal* in 1632.

Further detailed descriptions were written of plants as they flowered in his own garden or in those of his friends, for instance *Acinos odoratissimum*: 'I first found growing in the garden of Mr. William Yalden in Sheete near Petersfield in Hampshire, Anno 1620, amongst sweet Marjerom, and which by chance they bought with the seedes thereof. It is to be considered whether the seedes of sweete Marjerom degenerate and send forth this herbe or not, 11th October 1621, John Goodyer' (*Ger. emac.* 677).

But the outstanding event of the year was a visit to the garden of his 'worthy friend and excellent Herbarist of happy memorie Mr. William Coys of Stubbers in the parish of North Okington in Essex'. Goodyer came away with the seeds of many rare and little known plants, including several that had been collected by Boelius in Spain and had been brought over by him to England. Among other novelties which were thus introduced into the Droxford garden in 1620 were:

Acarna flore rubro.	<i>Carlina lanata</i> L.
Aracus maior Baeticus Boelii.	<i>Vicia sativa</i> β <i>linearis</i> Lange.?
Cattaria tuberosa radice Baetica Boelii.	<i>Nepeta tuberosa</i> L.
Ervilia silvestris Dodonaei.	<i>Lathyrus Ochrus</i> L.
Faba veterum serratis foliis Boelii.	<i>Vicia Faba</i> L. var.
Geranium Baeticum Boelii.	<i>Erodium gruinum</i> Willd.?
Gramen cristatum Baeticum Boelii.	<i>Cynosurus echinatus</i> L.
Gramen tremulum maximum.	<i>Briza maxima</i> L.
Herba Doria altera 26 July 1620.	<i>Senecio sarracenicus</i> L.
Hieracium stellatum Boelii.	<i>Rhagadiolus edulis</i> Gaertn.
„ medio nigrum fl. maior B.	<i>Tolpis barbata</i> Gaertn.

Hieracium medio nigrum fl. minore B.	<i>Tolpis umbellata</i> β minor Lange.
„ lanosum.	<i>Hieracium andryatoides</i> Vill.
Horminum silvestre tertium Clusii.	<i>Salvia verticillata</i> L.
Jacea capitulis hirsutis Boelii.	?
Lagopus trifolius maior Baeticus.	<i>Trifolium ligusticum</i> Balb.?
Legumen pallidum Vlissiponense.	<i>Vicia lutea</i> β laevigata Boiss.
Malva flore amplo Baetica aestiva.	<i>Malva moschata</i> L.
Petum indicum folio pene obtuso.	<i>Nicotiana Tabacum</i> v. <i>brasiliensis</i> .
Scabiosa flore rubro. Scabiosa sexta Indica Clusii.	<i>Scabiosa atropurpurea</i> L.
Silibum minus fl. nutante Boelii.	<i>Notobasis syriaca</i> Cass.
Valeriana mexicana.	<i>Valeriana cornucopiae</i> L.

Most of the plants raised from Coys' seed did so well with him that he was able to describe them as they flowered in the following summer.

It has been suggested¹ that on his way home Goodyer passed Rickmansworth, and there 'in the ponds about Moore Park' found 'Sium alterum Olusatii facie'. There is, however, no ground for this supposition; Goodyer himself wrote that he visited Moore Park in 1625.

We owe our first description of a new kind of English Elm to this same journey into Essex.

When we consider that many of our contemporaries are so deficient in powers of observation as to be unable to tell one tree from another, it is remarkable that Goodyer should have been so quick in distinguishing the various species of Elm. Two of these had already been recognized by Gerard in 1597, and were redescribed by Goodyer for the second edition in 1633: to these he now added a third, on his way to visit Mr. Coys:

'I observed it growing very plentifully as I rode betweene Rumford and . . . Stubbers in the year 1625, intermixed with the first kinde (the Common Elm), but easily to be discerned apart, and is in those parts usually called Witch Elme.'

The name of this Elm has caused considerable confusion in the minds of those who have not attentively read the words 'in those parts'. The Elm is certainly not the Wych Elm ordinarily so called, which Goodyer knew as the Witch Hasell or Broadest-leaved Elm; but, as

¹ Druce, *Goodyer*, p. 2.

Mr. Russell told me on my recent visit to Stubbers, it is the tree which is still locally known in that part of Essex as the 'Witch Elm', a name which is thus proved to have survived locally since Elizabethan times. Goodyer named it 'Vlmus folio glabro' or Smooth-leaved Elm, a name that has been adopted by Miller in the form of *Ulmus glabra*, the name by which it is still widely known, although Elwes prefers the name *U. nitens* Moench. for it.

Systematists of the eighteenth and nineteenth centuries have brought the species of Elms into such a state of confusion that it is a relief once more to get back to the simpler system of Goodyer, whose accounts of the four species of Elms were doubtless redrafted in 1632, but the original observations were much earlier. Johnson prefaces the descriptions with a short introduction :

'Our Author [Gerard] only described two Elms, and those not so accurately but that I thinke I shall give the Reader content, in exchanging them for better received from Mr. *Goodyer* ; which are these.'—T. Johnson.

Ulmus campestris Sm.

Vlmus vulgatissima folio lato scabro. The common Elme.

This Elme is a very great high tree, the barke of the young trees, and boughes of the Elder, which are usually lopped or shred, is smooth and very tough, and wil strip or pil from the wood a great length without breaking : the bark of the body of the old trees as the trees grow in bignesse, teares or rents, which makes it very rough. The innermost wood of the tree is of reddish yellow or brownish colour, and curled, and after it is drie, very tough, hard to cleave or rent, whereof naves of Carts are most commonly made : the wood next the barke, which is called the sap, is white. Before the leaves come forth the flowers appeare, about the end of March, which grow on the twigs or branches, closely compacted or thrust together, and are like to the chives growing in the middle of most flowers, of a reddish colour : after which come flat seed, more long than broad, not much unlike the garden Arach seed in forme and bignesse, and doe for the most part fall away before or shortly after the leaves spring forth, and some hang on a great part of the Sommer : the leaves grow on the twigges, of a darke greene colour ; the middle size whereof are two inches broad, and three inches

long, some are longer and broader, some narrower and shorter, rough or harsh in handling on both sides, nickt or indented about the edges, and many times crumpled, having a nerve in the middle, and many smaller nerves growing from him: the leafe on one side of the nerve is alwaies longer than on the other. On these leaves oftentimes grow blisters or small bladders, in which at the spring



THE COMMON ELM.¹

are little wormes, about the bignesse of Bed-fleas. This Elme is common in all parts of England, where I have travelled.

Ulmus minor Miller.

Ulmus minor folio angusto scabro. The Narrow leaved Elme.

This tree is like the other, but much lesser and lower, the leaves are usually about two inches and a halfe long, and an inch or an

¹ This figure originally appeared in *Mathiolus*, 144; was copied in *Lobel, Obs.* 607. 1; reproduced in *Ger. emac.* 1480. 1, and reappeared in *Parkinson*, 1404. 1. 6. It shows blisters on the leaves.

inch and a quarter broad, nickt or indented about the edges, and hath one side longer than the other, as the first hath, and are also harsh or rough on both sides, the barke or rinde will also strip as the first doth: hitherto I have not observed either the flowers or seed, or blisters on the leaves, nor have I had any sight of the timber, or heard of any use thereof. This kinde I have seene growing but once, and that in the hedges by the high way as I rode.



THE NARROW-LEAVED ELM.¹


betweene Christ Church and Limmington in the New Forrest in Hampshire, about the middle of September 1624, from whence I brought some small plants of it, not a foot in length, which now, 1633, are risen up ten or twelve foot high, and grow with me by the first kinde, but are easily to be discerned apart, by any that will looke on both.

¹ *Dodoens*, 837; reproduced in *Ger. emac.* 1480. 2; copied in *Parkinson*, 1404. 4. The leaves have been attacked by insects.

Wych Elm. *Ulmus montana* Stokes.

Ulmus folio latissimo scabro. Witch Hasell, or the broadest leaved Elme.

This groweth to be a very great tree, and also very high, especially when he groweth in woods amongst other trees: the barke on the outside is blacker than that of the first, and is also very rough, so that when there is plenty of sap it will strip or




THE WYCH ELM.¹

peele from the wood of the boughes from the one end to the other, a dozen foot in length or more, without breaking, whereof are often made cords or ropes: the timber hereof is in colour neere like the first; it is nothing so firme or strong for naves of Carts as the first is, but will more easily cleave; this timber is also covered with a white sappe next the barke: the branches or young

¹ *Ger. emac.* 1481. 3; copied in *Parkinson*, 1404. 2. (Stokes' edit. of *Withering*, 1787.)

boughes are grosser and bigger, and do spread themselves broader and hang more downwards than those of the first; the flowers are nothing but chives, very like those of the first kind: the seed is also like, but something bigger: the leaves are much broader and longer than any of the kindes of Elme, usually three or foure inches broad, and five or six inches long, also rough or harsh in handling on both sides, snipt or indented about the edges, neere



THE SMOOTH-LEAVED ELM.¹

resembling the leaves of the Hasell: the one side of the leaves are also most commonly longer than the other, also on the leaves of this Elme are sometimes blisters or bladders like those on the first kinde. This prospereth and naturally groweth in any soile moist or dry, on high hills, and in low vallies in good plenty in most places in Hampshire, wher it is commonly called Witch Hasell.

¹ *Gerard*, 1297. 2; copied in *Ger. emac.* 1481. 4; recopied in *Parkinson*, 1403. 3. Stokes considered this figure as 'rather a variety of *U. campestris*'.

Old men affirme, that when long boughes were in great use, there were very many made of the wood of this tree, for which purpose it is mentioned in the statutes of England by the name of Witch Hasell, as 8. *El.* 10. This hath little affinitie with *Carpinus*, which in Essex is called Witch Hasell.

Smooth Leaved Elm. *Ulmus glabra* Miller.

Ulmus folio glabro. Witch Elme, or smooth leaven Elme.

This kinde is in bignesse and height like the first, the boughes grow as those of the Witch Hasell doe, that is hanged more downwards than those of the common Elme, the barke is blacker than that of the first kinde, it will also peelee from the boughes: the flowers are like the first, and so are the seeds: the leaves in forme are like those of the first kinde, but are smooth in handling on both sides. My worthy friend and excellent Herbarist of happy memorie Mr. *William Coys* of Stubbers in the parish of Northokington in Essex told me, that the wood of this kinde was more desired for naves of Carts than the wood of the first. I observed it growing very plentifully as I rode between Rumford and the said Stubbers, in the yeere 1620 intermixed with the first kinde, but easily to be discerned apart, and is in those parts usually called Witch Elme.—*Ger. emac.* 1479-82.

A copy of a Memorandum, dated 15 November, in Goodyer's handwriting is printed on p. 374.

Part of the winter was spent in working through Gerard's *Herbal* and in extracting lists of plants, one of which (MS. f. 83) was dated 16 Januarii 1620.

1621

Goodyer signalized his twenty-ninth year by turning out more descriptions of new or rare plants in the three months of July, August, and September than in all the rest of his life. There must have been some special reason for this great output. It is reasonable to think that he found an incentive in the large number of novelties he had seen in the gardens of Coys and Parkinson. There was also the added interest of possession, of the triumph of the horticulturist: seeds given to him by Coys in the preceding year were coming up well. What would they become?

What new benefits to mankind might they not be made to yield? Many of them growing at Droxford under his own eye were either not described or were imperfectly described in Gerard's *Herbal*; there was room for a new edition, or at any rate for a Supplement. The new plants might not be hardy enough to survive the next winter, and Boel might never be able to procure the same seed again. Whatever the reasons may have been, Goodyer's enthusiasm certainly reached a red-heat this summer.

It was the enthusiasm that every true naturalist feels when he is conscious of being the first to obtain an insight into unknown processes and phenomena. What plans for publication he may have had we know not, nor do we know what hindrances arose, but there followed a hiatus and these notes of Goodyer's were laid aside for twelve years, and then a selection of them were freely placed at the disposal of Dr. Johnson for the new edition of Gerard.

The descriptions conform rigidly to the style in vogue at the period. The habit of the plant, its stalks, branches, leaves, flowers, seed, root, duration are all considered in order, and in many cases notes are added on the season, hardiness, and locality. The technical words such as 'footstalk' in the sense of petiole, 'knee' = node, 'bosome' = axil (of a leaf), 'pointell' = pistil, 'cheives' = stamens, and umbell in the modern sense are in the language of Gerard. Occasionally the descriptions are brightened by refreshing comparisons. The flower of *Monotropa* 'resembleth ye suck-bottle which children use to suck their drinke out of'; 'seed no bigger than a flea'; 'clammie as Bird-lime'; 'woollie like Spiders webbs'; a colour is 'orange tawnie velvett'; a herb 'heateth and burneth the mouth'; the distribution of the pollen of the Yew is thus described, 'if you shall beate on them stones into this tree about the end of February, there will proceed and fly from the flowers an abundance of dustie smoke'.

Moreover, at this period of his scientific life he might have written as Ray did in June 1667 that he had been

at work 'in gathering up into a catalogue all such plants as I had found at any time growing wild in England, not in order to the present publishing of them, but for my own use, possibly one day they may see the light'.

It must be remembered that at this period there was no English botany book worthy of the name in existence. The standard works used in this country were the *Herbal* of Dr. William Turner, printed in 1568 at Cologne, and illustrated, *not* with drawings of British plants, but with reduced copies of some 400 figures drawn from continental plants and previously used at Basel to illustrate the 1545 edition of Fuchs' great work. That Fuchs' original drawings were of considerable beauty, and that nothing to equal them could have been engraved in England, is admitted; but as illustrations to an English *Flora* they were misleading.

In 1574 a further impetus was given to the study of English botany by the publication of Henry Lyte's edition of Dodoens' *Herbal*, which, as the preface shows, has considerable claims to originality rather than to being a mere translation, but for this again the small, inferior copies of Fuchs' figures were requisitioned.

The next and best known of English *Herbals*, that of Gerard, appeared in 1597, but again it was a compilation from foreign works, so imperfectly 'accomodated unto our English nation' that Lobel was requested by the printer to correct the blunders, and still Goodyer was able to discover many others.

Consequently, like many other botanists, Goodyer must have been eternally plagued with the attempt to make English plants fit the descriptions and figures of continental writers—a labour that has been aptly compared with the endeavour to fit square pegs in round holes. He was, however, sufficiently gifted to perceive the futility of the attempt, and to recognize at an early period the need for accurate descriptions made from the living, flowering plant beside him.

Among foreign authors he would have relied largely on Matthiolum, Bauhin, and on Lobel. One of the most important works that appeared during his working life was the *Pinax* of Kaspar Bauhin, the result of forty years of toil, published in 1623, and acquired by Goodyer in the same year. This work carried still further Lobel's idea that natural affinity must be the foundation of a truly natural system : it went far to clear up the confusion in the nomenclature of the day, which had resulted from the fact that different names had often been given to the same plant by different authors. The *Pinax*, as Reynolds Green has well put it, 'not only got rid of much confusion by setting forth the different synonyms in use, but it introduced greater terseness of description, and helped to restrict the inordinate length of names'. Green did not think that Bauhin's work had exerted any great influence in England :¹ but we have noted a profound influence upon Goodyer's work, who prepared and, had times been favourable, would assuredly have published a *Pinax* of the British flora on the lines of Bauhin, whose works he had read, marked, and inwardly digested from cover to cover.

Goodyer's excursions in search of plants were rewarded by the finding on 2 June of two species of Pondweeds new to the British aquatic flora. Both were growing quite near home, the one actually in Droxford, the other just over the Sussex border at Durford on the upper waters of the Rother. A paper, MS. f. 6, suggests that either Goodyer, or more probably Sir T. Bilson, was interested in land that once belonged to the old Priory at Durford.

On 29 June, about a mile from his native town of Alton, he found a plant which he described as 'Nidus avis flore et caule violaceo purpureo colore'. This he 'found wilde in the border of a field called Marborne, neere Habridge in Haliborne . . . being the land of one

¹ On this point Dr. Church notes that in Oxford 'Bauhin's *Pinax* prompted *Sherard's Pinax* of Dillenius, and hence was responsible in England for the Sherardian Professor of Botany'.

William Balden.¹ In this place also groweth wild the thistle called *Corona fratrum*'. The identity of the species has been the subject of speculation. The most recent review of the position is that of Mr. Druce, who follows Sir J. E. Smith in considering that the plant was probably the Purple Broomrape (*Orobanche purpurea*) growing parasitically upon the roots of the common Yarrow which is now abundant in a locality, which Mr. Druce believes that he has correctly identified with the 'field called Marborne'. Townsend, on the other hand, suggests that the plant was the rare Violet Helleborine, *Epipactis violacea* Dur., which he found near Alton; a view with which Dr. Stapf agrees. In his annotated copy of Tabernaemontanus' *Icones*, Goodyer has added a note and a reference to the picture of *Orobanche* 1 to the effect that it is *Nidus Avis* Lob. o. 356 or '86 (b. 2)', which is evidently a reference to *Orchis abortiva violacea* in Bauhin's *Pinax*.²

Although closely tied to his work and his garden during the early part of the summer, Goodyer appears to have paid short visits to the coast, at Haylinge on 20 August, where he found Sea Heath (*Frankenia laevis* L.), and *Diotis maritima*, which was formerly much more widely distributed than it is now; to Southsea on the 30th; and to Bursledon Ferry on 3 September. Here he saw the Sea Heath in flower and described it a second time. At Bellmere Pond he found a white-flowered Eyebright in flower on 24 August, and at Southsea Castle he saw Climbing Fumitory in flower on 30 August.

Altogether he described at least ninety species of plants during the summer, and when October brought him a respite from his labours it was but natural that he should

¹ Parish Records might possibly show whether 'Balden' is an old error for Yalden, the name of Goodyer's kinsfolk.

² Yet another view is printed by the editors of the catalogue of the *Morisonian Herbarium*. The entry is 'Orobanche radice compacta major flore violaceo, nobis. No specimen. In agro Hantoniensi prope Alton oppidum invenit D. Goodyer. This is usually referred to *Limodorum abortivum* Sw., but Goodyer's plant (of which no specimen is known to exist) was probably *Helleborine purpurata* Druce'. Vines and Druce, p. 173.

wish to compare notes with his friend Coys. His description of 'Jacea capitulis hirsutis Boelii' certainly implies that it was taken on 10 October 1621 in Coys' own garden. And a letter from his friend, Laurence Davis, about gold weights addressed to him at Droxford on 9 November, implies that he was home again then.

To my very loving freind Mr. John Goodier at
Droxford give this.

L. S.

Mr. Goodyer I have sent you by Maye yo^r Gould waights wth those other that you desired extraordinary, only the Elz. Angell serves both for the Edward & Elzabeth waight, theye cost 6^s. And yo^r peice I could gett but 8^s 10^d for; the Remynder I have returnd by Maye. And I am gladd to have this occasion to expresse my desire to bee esteemed.

Yo^r ever loving freind

Laurence Davis.

I praye remind mee kindly to yo^r fellows

Patience, Mr. Parker, and Henry Henly.

[MS. f. 9

This letter was endorsed by Goodyer 'Rec. 9 November 1621' and annotated 'rec. 3^s againe'.

6

His last work this winter was to distinguish between 'the Yew bearing Acorns and berryes' and 'the Yew with only flowers'.

A number of the papers in which Sir Thomas Bilson's affairs are mentioned belong to 1621 and 1622. Shopping notes occur on MS. f. 46. 'Morgan is to doe the fan of a pinke color'. 'The fans handell is to be doble gilt'. Among the things to be purchased are '20 yards of galome lace', buttons, fustian, cambic, oyle skins, 50 needles, and other such feminine gear, p. 382. The addresses 'Little Minories' and 'Barbican' on the same paper indicate that business had again taken him to London.

1622

The principal journey of 1622 was to Oxford. It is not difficult to find good reasons for the visit. The relations that existed between Magdalen College and Goodyer's

own county, with William Yalden the College Clerk of the Account, his brother-in-law, and in addition his scientific interests may rather make us wonder that he should not have visited Oxford more frequently. It was but the year before, that Magdalen had granted to the University five acres of land, then in process of being laid out as a Botanic Garden, and Goodyer may have contributed some of the first plants. Among the then fellows of the College was Walter Stonehouse whose garden-lists subsequently came into Goodyer's possession, and among the new Demies was Sampson Johnson, the friend of Thomas Johnson and an early authority on the purgative action of various kinds of rhubarb.¹

In May Goodyer completed his notes on the flowers and galls of the oak, and on the cachryes² of the walnut, chestnut, alder, and birch. Oak-galls had long had a peculiar interest, because a gardener by 'looking whether there be in them eyther Flyes, Wormes, or Spiders' could presage battle, dearth, and scarcity or plague.³

On 2 July 'in the high waie neare Abington leadinge towards Oxford' he saw the fine Woolly Thistle (*Cnicus eriophorus*), which he had already found by a Hampshire roadside in 1617 (f. 107). On the 5th while exploring 'by the Rivers side and in the water diches about Oxford' he saw the Great Water Parsnip (*Sium latifolium* L.) before the flowers were fully formed, and continuing his walk on the west part of Gloster Hall, now Worcester College, he there noted for the first time the Wood Club Rush (*Scirpus sylvaticus* L.) (f. 7 v.).

¹ Thomas Johnson wrote 'My friend Mr. Sampson Johnson, Fellow of Magdalen College in Oxford, assures me, that the Physitions of Vienna in Austria use scarce any other [medicine] at this day than the Rubarb of the Antients which grows in Hungary not far from thence: and they prefer it before the dried Rubarb brought out of Persia and the East Indies, because it hath not so strong a binding facultie as it, neither doth it heate so much; onely it must be used in somewhat a larger quantitie' (*Ger. emac.* 396). S. Johnson had no doubt opportunity of testing this statement during his year's leave of absence from College, beginning on 14 March 1630-1.

² See p. 174.

³ Thomas Hill, *The profitable Arte of Gardening*, 1574.

In the autumn he paid his customary visit to Coys,¹ who on 29 September was able to show him 'Scabiosa flore rubro' in flower (cf. 8 October 1621).

During the winter months he was evidently occupied with a literary labour that will be referred to in the following year when it was finished.

In February a stray note on the Curled Parsley (*Apium crispum*) growing at Idsworth suggests an excursion down the steep wooded hills beyond the South Downs in the midst of very beautiful country, then part of the Forest of Bere. And in March he received seeds of twenty-two garden plants from Coys, including two kinds of 'aples of love' or tomatoes, p. 325.

1623

His botanical descriptions now cease for a while. We have only one note, on a species of *Acorus* dated July 1623. It may be that he was unable to pursue his hobby with the same activity as heretofore: perhaps he had to leave Droxford—for we find no mention of his garden there after this date. But the advancement of botanical learning still remained the prevailing occupation of his leisure, only from henceforth we find it taking another form, a desire to make the plant lore of antiquity available for English readers, and to this he devoted the winter months.

By this time he had collected a considerable number of botanical works both of his contemporaries and of the great Masters of the past. A beautifully bound copy in Italian binding of the Aldine folio edition of Theophrastus (1497) was among them. The margins of this book he inscribed with the numbers of the chapters, and translated the whole into English. The ten books *De Plantis* fill 238 pages of his neat small script, and the six books *De Causis Plantarum* take 256 more. We do not know when

¹ This is proved by a marginal note on p. 130 of his copy of Parkinson's *Theatrum*. There he renames Parkinson's No. 8, *Aster Virgineus*, as 'Helianthemum radice repente Virginianum. Mr. William Coys of Stubbers in Essex was wont to call it so. 26 Sept. 1622 I sawe it in his garden. John Goodyer'.

he began the translation, but he noted the dates on which he completed the chapters *De Causis Plantarum*.

Chap.	2	finis	25 Febr. 1622
	„	3	„ 8 Marcii 1622
	„	4	„ 18 Marcii 1622
	„	5	„ 28 Jan. 1623

At the end is written

‘Finis 6^{to} die Februarij 1623.

This booke was sent from bindinge the six and twentieth day of September 1661.

The bindinge—the cleane paper—the claspes 4^s.’

This, so far as we know, was the only English translation of the *De Plantis* until 1919, when Sir Arthur Hort published his translation; but Goodyer’s still remains the only version of the *De Causis* in the English language.

1624

We find him resuming his botanical excursions in 1624. He made a second trip into Wiltshire, and noted the occurrence of Sanfoin (*Onobrychis sativa* Lam.) at Langford and Stapleford on 24 July. A carefully prepared itinerary for a tour in the New Forest in September of the same year shows his route :

From hyve to Lymington	Fr. hampto to Redbridg	4
from Lymington to Christchurch	fro Redbridge to Ringw	16
from Christchurch to Poole 8 myle	fro Ringwood to Langham	6
from Poole to Wareham 7 myles	fr Langham to Poole	4
to Wimborne 6		
to Brainford 10		
to lichett 3		
from Wareham to Wooll & Briden 4 myle	Ringewood	2 4
frō wooll to Waymouth 9 myle	Poole	1 2
to Dorchester 9 myle	Wooll	2 4
to Woodbury hill 4 myle	horse hire	1 0
to Winfruite 2 to Osmunton 4	passag	0 4
to bean 3	Waymouth	3 0
fro Waymouth to Woobury 14	Christchurch	3 0
to Wimborne 8 fro	Lymington pass	0 2
Wimborne to Christchurch	Hyve pass	0 4
from Christchurch to Lemington 9	hampton	2 0
frō Lemington to Bewly 4	ichen pass	0 2
fro Bewly to Hyve 2	. . et Church	0 6

Mr. laden of Wimborne. 16 4

E 2

[MS. f. 13

In preparation for this journey he made copies of Lobel's descriptions of four species of plants which were already recorded as living at Portland. *Anthyllis prior lentifolia peplios effigie maritima*,¹ *Sedum Portlandicum*,² *Hederatium thlaspi*,³ and *Papaver cornutum flore phoeniceo*.⁴ We do not know whether he found them all, but we do know that he noticed as a new species a fourth kind of English Elm, the Narrow-leaved Elm, already described on p. 40.

Eight years later he wrote, 'This kinde I have seene growing but once, and that in the hedges by the high way as I rode betweene Christ Church and Limmington in the New Forest, about the middle of September 1624, from whence I brought some small plants of it, not a foot in length, which now, 1633, are risen up ten or twelve foot high, and grow with me by the first kinde [the common Elme], but are easily to be discerned apart, by any that will looke on both.'

It has been suggested, by Druce, that Goodyer's Narrow-leaved Elm was the Cornish Elm, but Elwes and Henry both refer it to *Ulmus minor* Miller, which they call Goodyer's Elm, and with which they merge Druce's *U. Plotii*. Elwes states that Goodyer's Elm has been lately found near Christchurch by Dr. Moss.

On 10 September he wrote a description of the Shrubby Suaeda (*Suaeda fruticosa* Forsk.) under the name of 'Chamaeptytis vermiculata'.

Johnson quotes him on the subject of the great abundance of the Common Spleenwort in Woolmer Forest. 'Mr. Goodyer saith that in January 1624 he saw enough to lade an horse growing on the bancks in a lane, as he rode betweene Rake and Headly in Hampshire, neere Wollmer Forrest' (*Ger. emac.* 1146).

¹ *Arenaria peploides* L. ² *Euphorbia portlandica* L. ³ *Cochlearia danica* L. ⁴ *Glaucium phoeniceum* Crantz. The 'Sedum' is usually quoted as 'Tithymalus', and the following plant as 'Thlaspi hederaceum'. These records are about a century earlier than those given by Mansell-Pleydell, *Flora of Dorset*.

1625

It was about this time that he learnt the local name (Honewort) and vertues of Corn Parsley, a plant with which he had been long familiar, and to which he had already given the name 'Selinum Siifoliis': but he had no English name for it, until one day he saw Miss Ursula Leigh, servant to Mistress Bilson of Mapledurham, gathering it in the 'wheate ershes' about Mapledurham (where it still grew in 1632, especially in clay grounds). She told him that it was called Honewort, and that her mother 'late of Brading in the Isle of Wight deceased, taught her to use it after the manner heere expressed, for a swelling which she had in her left cheeke, which for many yeeres would once a yere at the least arise there, and swell with great heat, rednesse, and itching, until by the use of this herbe it was perfectly cured, and rose no more nor swelled, being now (5 Martij 1632) about twenty yeeres since, only the scar remaineth to this day. This swelling her mother called by the name of a Hone, but asking whether such tumors were in the said Isle usually called Hones she could not tell, by reason shee was brought from Brading aforesaid young, and not being above twelve yeeres old when shee used this medicine'.

The Vertues.

'Take one handfull of the greene leaves of this Honewort, and stampe them, put to it about halfe a pinte or more of beere, straine it, and drinke it, and so continue to drinke the like quantity every morning fasting till the swelling doth abate, which with or in her was performed in the space of two weekes at the most.'¹

Next we have the discovery of the compact little Knotted Pearlwort (*Sagina nodosa*), described as 'Alsine palustris foliis tenuissimis: sive Saxifraga palustris alsinefolia', on the boggy ground below the Red Well of Wellingborough in Northamptonshire. 'This hath not been described that I finde. I observed it at the place aforesaid 11 August 1625.'

And in the following month he was the first to record the poisonous Cowbane or Water Hemlock (*Cicuta virosa*),

¹ *Ger. emac.* 1017-18. See p. 121.

which he named 'Sium alterum olusatri facie', on 16 September 1625. The locality was 'by Moreparke, and at Denham in Hertfordsheire in standinge water *sine caule*' (f. 58).

It has been suggested that Goodyer, as a staunch Royalist, visited the Red Well because King Charles and Henrietta were residing there. But surely the evidence is of the flimsiest. The legend repeated by Morton, and supported by a misquotation from Laud's Diary,¹ of the King and Queen living for *weeks* in a tent beside the habitat of *Sagina nodosa*, supplies but a sorry explanation for Goodyer's visit to a popular watering-place.

The Northamptonshire flora owes the first notice of the Grass of Parnassus to the same visit.

1627-8

Of his proceedings during the next few years there are but few notes. On 9 February 1627 he 'Rec. of Christopher Potecary of Stockton, 5 myles from Venny Sutton Clother'. It is not clear what he did receive, for after this note follows, though written another way up, a list of fruit trees and plants, days of Assizes, and (upside down) the recipe for an ointment.²

Two entries, dated 23 June 1628, show that he visited the garden of one 'Millaine' in London, and saw there 'Sophia latifolia in horto millaine prope le pest house' and 'Triticum spica multiplici, in horto Millaine'.

The former may have been *Sisymbrium Sophia* (?) and Millen's grass may have been a variety of *Triticum turgidum* L.

The owner of the garden was the 'Master John Millen, dwelling in Old Street, in whose nursery are to be found

¹ The statement in a well-known County History that Laud visited the sovereigns at *Wellingborough*, is based on a misreading of his own entry that the King appointed him Bishop of Bath and Wells, 'Rex Carolus me nominavit in Episcopum Bathon. et *Wellen*.'

² MS. f. 129 v, see p. 384.

the choisest fruits this kingdom yields' ¹ and 'who from John Tradescant and all others that have good fruit hath stored himself with the best only, and he can sufficiently furnish any'.¹ Parkinson mentions the 'Great bearing cherry of Master Millen in 1629'. His name and address also occur on the back of a letter of 1631.²

Ceribum. Jacobus dei
 capita / ^{gra} Jacobus dei gra
 Johannis. Jacobus dei gra
 Gramen ^{gra} Jacobus dei
 littera in ^{gra} ~~gratia~~
 Johannis Goodyer.

GOODYER'S HANDWRITING

1631

In 1631 Goodyer was living at Mapledurham. When he moved, we do not know, but both his letters and notes show that his thoughts were in the planting of a new garden. He journeyed to London and possibly to Oxford, and on 27 May appears to have been at 'Godlemen in Surrey' (Godalming). He would have had a good reason for visiting Oxford, for his favourite nephew Edmund Yalden had gone up to Magdalen College as a Demy in 1630. A note of a botanical station near Oxford was supplied him by Leonard Buckner,³ a London apothecary, and one

¹ Johnson, 1633, and Parkinson, 1629, p. 575.

² MS. f. 133.

³ Leonard Buckner's discoveries are printed by Johnson. 'In a field joyning Witney Parke' in 1632 he found *Stachys germanica* L., and 'in a bog upon a common by the Beacon hill neere Cumner-wood in the end of August 1632', three miles beyond Oxford, a little on this side of Eynsham ferry, he found the Horse-taile Coralline. *Ger. emac.* 1115. This last plant has been variously regarded as *Equisetum sylvaticum* L., Druce, *Flora Oxford*, p. 356, and as *Chara hispida* L., Druce, *Flora Berks.* xcix. In the case of the *Stachys*, Goodyer's note antedates the date printed, and generally quoted from Johnson, by a year and a month.

Leonard's excursions near Oxford suggest kinship with Dr. Thomas Buckner, Fellow of Magdalen, 1618-31.

of the 'loving friends and Fellow travellers' who accompanied Johnson in searching for plants over a great part of Kent.

'Stachys: by windy parke wall on ye west of it, 8 myles from Oxford—July 1631. Leo. Buckner.' [MS. f. 133

Another correspondent, Griffith Hinton, who shows as much familiarity with the movement of Bishops as with the stock-in-trade of nurserymen, addressed two letters to Goodyer in this year.

To his very loving frend

M^r John Goodyear at Maple Derham neare Peeterfyeld in Hampshier
geeve theis.

Mr. Goodyer I rec. the Acquittances, and as sone as I have the Rents together you shall heer from mee. My Lo. Byshop of Wynchester¹ is this day com out of the town for Farnham and how long he wyll stay ther I know not, but as I heard by on of my Lords men hee wyll stay at Farnham 10 dayes. Thus wyth my duety and kynd Remembrance this 13th of July 1631 I rest,

Yors ever loving

GRIFFITH HINTON.

[MS. f. 14

In November 1631 Goodyer was again in London. The notes of this visit are not very easy to read, but there is sufficient to show that he paid 13s. 8d. for 'Diett at Gilford' and that supper at the King's Head cost 10s. The paper is undated, but the year is settled by the days of the month on which he purchased certain books, Dioscorides and Thevett among others, which are still in existence, and are clearly inscribed with a date and price exactly corresponding with the note on the paper and the year 1631. The further entry 'Nov. 8 wyne wth Johnson 6^d' has a special significance, for about this time the two friends would have been discussing their great scheme of producing a second edition of Gerard's *Herbal*. In spite of its great popularity, this *Herbal* had been an unsatisfactory book from the start. Indeed, when we remember its history, it would have been strange had it

¹ In 1631 Walter Curle succeeded Rich^d Neile as Bishop of Winchester.

been otherwise. The printer, John Norton, had commissioned Dr. Priest to translate Dodoens' *Pemptades* (1583) into English. Priest died, and Gerard continued the work. But to mask the fact of his *Herbal* being little else than a mere translation, he altered the arrangement from that of Dodoens to that of Lobel; and flippantly remarking that he had heard of Dr. Priest's labours, but the man being dead his work had perished with him, he had the effrontery to declare that his own researches had produced the work. Wood-blocks used by Tabernaemontanus in his *Eicones* (1590), with some others, were procured from Frankfort by Norton, but Gerard soon showed his slender knowledge, by misapplying many of the figures, and caused so much confusion in the early chapters of the *Herbal*, that the attention of the printer was directed to it by James Garret, the London apothecary. Lobel was therefore invited to correct the work, and by his own account he actually corrected it in a thousand places, but further emendation was stopped by the author, who contended that the *Herbal* was already sufficiently accurate, and that his censor had forgotten the English language.¹

Gerard's book gave a very real impetus to the study of our English flora, but it was avowedly a popular work, 'being principally intended for gentlewomen'²; and in this connexion Mrs. Gerard, who assisted her barber-surgeon husband in his practice, would have been most helpful; for her professional assistance was similar to that which Mrs. Gamp was in the habit of rendering at

¹ An excellent account of the *Herbal* is contained in B. J. Jackson, *Gerard's Catalogue of Plants*, 1876.

² Americans of 'royal descent' may be interested to learn that the copy of Gerard's *Herbal* now in the library of the Botanic Garden at Oxford, originally belonged to a gentlewoman, Dorathie Redmayne (1565-1645) whom I believe to have been identical with the mother of John Rolfe of Heacham, ancestress of all who trace their descent to the Indian Princess Pocahontas. Dorothy Rolfe's second husband was Robert Redmayne, Chancellor of Norwich. She was buried 'at the feet of her two husbands' in Heacham Church. *Rolfe Family Records*, p. 11.

certain periods to her lady clients. The scientific botanist, however, judges a book from another standpoint. Even the genuineness of the Catalogue of plants in Gerard's own garden has been denied by the 'attestor' Lobel himself. In one copy of the work in the British Museum the certificate has been crossed out, and the words, in Lobel's handwriting, 'haec esse falsissima, Matthias de Lobel', are written at the end of it.

Goodyer made the fullest use of the *Herbal*, and in 1620-1 he may have been contemplating a new and improved edition. When or how the same thought came to Dr. Johnson we do not know, but in the next year Goodyer was sending Johnson twenty-seven sheets of manuscript.

The name of the inn, where he put up, is given in a second letter from his loving friend Hinton, addressed :

To his very Loving frend

Mr. John Goodyear at the sygne of the

Angell neere Denma^rk Hous in Strand.

1631

Mr. Goodyer I wrote unto my frend for the trees and this day I spake with him and hee telleth mee that ther is no sure trees growyng about Barn Elmes¹ wher hee dwelleth, but hee hath enquired and found that ther is Malacaton trees at Twycknam in Mydd. and they wyll not be sold under 3^s 6^d a tree, but hee may have an apricok tree for 1^s.² I shall speake with him agayn eyther

¹ The Earl of Essex had a garden at Barne Elmes, *Ger. emac.* 1396, and so had Sir Francis Walsingham (d. 1590), *Ger.* 501.

² The Twickenham fruit garden must have been that of Mr. VINCENT POINTER, who had the greatest variety of plums in England (*Gerard*), and is quoted by Sir Hugh Platt, *Floraes Paradise*, 1608, pp. 117-18, as an authority on grafting. His nurseries are mentioned by Ben Jonson, who states that he was better known 'by Poynter's name than by his owne'. His real name was CORBET, and he was the father of Dr. Richard Corbet (b. 1582), Dean of Christ Church (Aubrey, *Lives*), and was also connected with a garden at Ewell in Surrey.

An Epitaph on Master Vincent Corbet.

Deare Vincent Corbet who so long	His Mind was pure, and neatly kept,
Had wrestled with Diseases strong	As were his Nourceries; and swept
That though they did possesse each limbe,	So of uncleannesse, or offence,
Yet he broke them, e're they could him,	That never came ill odour thence:
With the just Canon of his life,	And adde his Actions unto these
A life that knew nor noise nor strife:	They were as specious as his Trees.

Ben Jonson, *Underwoods*, 1640.

tomorrow or Thursday that I may geve him an answer, therfore I pray you send mee word by this bearer or yf you Gooe to have them about this town bethink yo^r self. I spake with on Crawley who was my Lord Byshopp Androse¹ his gardner after I had written to Barn elmes and Crawley promysed me to help mee to som but at what pryce I know not yf you please to speake with him send mee word and appoynt the tyme and I will speak with him tomorrow or yf you wyll have them out of the contry send me word that they may bee reddy to bee sent down by the carrier tomorrow. serving this with my duety this 15 of november I rest

Yo^{rs} ever

Griffith Hinton.

[MS. f. 133

On the same letter is written in Goodyer's hand :

James	11	10	4	Violett	1	0	Mr. Withowbie	HamsHERE
D.	16	7	4	May cherry	1	8	Audit	
Roberts	-	10	0	Duke cherry	2	0	Draba ger.	
Pashley	-	12	6	Bon crittian	1	8	Gramen alopecuroides spica	
(sic)	28	0	2		6	4	aspera is Gramen cristatum	Baeticum by y ^o adiacent pts of Shepey.
	11	10	4				Stachys. by windy parke wall on ye west of	
	16	7	4				it 8 myles from Oxford in July 1631.	
Buckett	4	10	0				Leo. Buckner.	
	32	7	8				at Mr. John Harrison in pater noster Rowe	

at the Golden Unicorne—a stationer.

[MS. f. 133

Both the letter and Goodyer's notes written upon it show that he was stocking his garden. 'Millain of old Street' is again mentioned, as is 'Hugobert at Ratcliffe'.² The prices paid for some fruit trees are noted on the back of this letter.

In January he was trying to get into touch with foreign herborists with a view to obtaining the seeds of foreign plants in exchange for English seeds. Drafts of two letters are extant.

Sir,

I have made a short Catalogue of some plants which growe for the most p[ar]te wild in Fraunce; you may acquaint anie herborists there that you please yf they will [be pleased] helpe me to seeds of them, or any other, I will by your directions furnish

¹ Lancelot Andrewes, Bishop of Winchester, 1618-26.

² The name Hugobert recalls an apothecary, Abraham Hugobert, who was fined by the Apothecaries Company for not presenting an apprentice in 1650.

them with such as they shall desire which growe wild with us in England or in our gardens; those in my Catalogue are all containd in Pena his *Adversaria*, I have quoted the page that there may be no mistakinge. And I entreate you to desire so much of those that send me any seeds to name the Author and the page of his booke that wrightes of them for every seed—all sort he sends, if not written of, to say so. This way is the triall of an herborist, and will save me a greate deale of paines; by gods helpe I will doe the like to them for those seeds I shall send. Thus making bold to trouble you uppon your kind promise, wishing you a prosperous journey,—I rest

Yours to my power—

[MS. f. 132

[The catalogue contains the names of fifty-one plants with references to Lobel and Pena, *Adversaria*.]

16 January 1631.

Mr. Wray, I have made bold to send this inclosed letter to you, I entreate you to convey him to Mr. Langrish if he be yet in England, if gone over to send him after him if it may be done with convenience, if not pray send him back to me againe. It was his desire when he was with us in the contrey that I should send this letter to you for him. Thus remayning ready to doe as much and more for you as it be in my power, I rest

Your loving friend

[MS. f. 132

Both of these letters are in Goodyer's handwriting. The identity of Mr. Wray is a matter of doubt. In the *D. N. B.* are mentioned two members of a Yorkshire family of the name, of whom Sir J. Wray (1586-1655) is known to have spent 1603-6 in foreign travel, which is rather early for our present purpose, and the Captain Wray, who with John Evelyn in 1646 found rare simples growing on the Euganean Hills, is rather too late.

1632

The doctrine of the fixity of species so tenaciously held in after years by Linnaeus and his school formed no part of the science of Goodyer and his contemporaries, or of the ancients. The artificial production of new varieties was

already known to Shakespeare before 1610, at any rate as far as gillyflowers are concerned. Perdita, having been told the cause of their streakiness, cared not to get slips of them for her rustic garden: they had the reputation of being 'nature's bastards', and she was also aware that it was possible to produce similar piedness by art (inoculation). Polixenes tried to argue with her that as all arts that add to nature are made by nature, the crossing of different races is but natural after all, and that she should therefore make her garden 'rich in gilliflowers'.¹

In Shakespeare's day people were familiar with the idea of two sexes among plants, although the function of pollen was still unrecognized. But it was a remarkable prevision of genius, which we can only describe as Shakespearean, to explain the production of varieties among plants as the result of cross-breeding, at a time when botanists knew nothing about the function of the flower.² The Art to which Perdita refers must be the art of Inoculation, of inserting the buds of one plant into or upon another.³ And Shakespeare's *alter ego*, Bacon, reflects the same thought. 'It is a Curiosity to make Flowers Double. . . . Enquire also, whether *Inoculating* of Flowers (as *Stock*

¹ *Winter's Tale*, iv. 3.

² Dr. Church points out that at the time of which we are speaking, the idea of crossing by grafting was accepted, and the methods of grafting fruit trees were fully set forth in many books. As far as the 'Secretes' of the ordering and care of Gillyflowers are concerned, Th. Hill, *Arte of Gardening*, 1574, divulged the following: 'you may make one stalke to bring forth floures of many colours, if you take the seeds of every colour of the Gilifloure, and put them altogether into a thinne small rede or Terdill of a sheepe or goate, or else tied up in a thinne worne linnen cloth, setting the same in the earth well mixed with dung: which after the watering will cause a plant to come uppe, bearing the like number of colours in one stalke, as there were seedes sowed.

And there be some which write, that if you myxe the Basill seedes with the Gilifloure seedes, and use them (as above sayde) that they will spring together on one stalke'.

It must also be remembered that Hill had many contemporaries, who, had they been acquainted with the method of the production of new varieties by hybridization, would not have imparted their secret to others, so long as they thought that there might be money in it.

³ Fleming's *Virgil*, *Georgics*, ii. 21. 1589.

Gilly-flowers, Roses, Musk-Roses, &c.), doth not make them Double'.¹

The idea of the sudden appearance of a new species as a Mutation would have been quite familiar to Goodyer. We have already quoted a passage² (under 1620) in which he put forward a theory, as worthy of consideration, that the seed of Sweet Marjoram might degenerate and send forth *Acinos odoratissimum*.

Francis Bacon, seven years later, accepted the possibility of such a change, and suggested an experiment for the Transmutation of Flowers. 'The second rule shall be to bury some few seeds of the herb you would change, amongst other Seeds; And then you shall see whether the juyce of those other seeds doe not so qualify the Earth, as it will alter the seed, whereupon you worke. As for Example . . . put *Basill-seed* amongst *Thyme-seed*, and see the change of taste, or otherwise.'³ The effect of a change of environment was illustrated by the classical instance of Lobel,⁴ who sowed 'Papaver nigrum' in Somersetshire and found it to come up changed 'by the sport of Nature and metamorphosis' into 'Papaver album'.⁵ And in 1632 Goodyer found what was believed to be an instance of the partial change of an ear of wheat into oats. Our modern knowledge of the possibilities of plant-breeding will of course not permit us to believe in his explanation of the phenomenon: he was probably misled by some monstrosity in the ear, but the record is of value as showing the frame of mind in which these early botanists tried to describe honestly what they saw. A later generation would have cast the thing aside as being 'against nature' and not worth a serious thought. Johnson records it as 'a rare

¹ Bacon, *Nat. Hist.* § 513. 1627. Here the word 'flower' is used in different senses, first as a bloom, secondly as a plant. Bacon's science was occasionally muddled.

² *Ger. emac.* 65.

³ Bacon, *I. c.* § 527.

⁴ Lobel, *Officina Pharmaceutices Rondellet*, 1605, p. 37.

⁵ A change of colour from blue or yellow to white in the case of Wild Succory and Moth Mullein was attributed by Merrett to a change to a poor soil. *Pinax*, Epistle to reader, 16th page. 1667.

observation, of the transmutation of one species into another, in plants; which though it have been observed of ancient times, as by *Theophrastus, de caus. plant. lib. 3. cap. 6*, whereas amongst others hee mentioneth the change of *Ζεῖα πρὸς τὸν βρόμον*, Spelt into oates: and by *Virgill* in these verses;

*Grandia saepe quibus mandavimus Hordea sulcis,
Infoelix Lolium, & steriles dominantur avenae.*

That is;

In furrowes where great Barley we did sow,
Nothing but Darnel and poore Oats do grow;

yet none that I have read have observed, that two severall graines, perfect in each respect, did grow at any time in one eare: the which I saw this yeare 1632, in an eare of white Wheat, which was found by my very good Friend Master *John Goodyer*, a man second to none in his industrie and searching of plants, nor in his iudgement or knowledge of them. This eare of wheat was as large and faire as most are, and about the middle thereof grew three or foure perfect Oats in all respects: which being hard to be found, I held very worthy of setting downe, for some reasons not to be insisted upon in this place'.¹

The Goodyer crest, a partridge with a *good ear* of wheat in its beak, is said to have been suggested by this exceptional find.

Various lists of exotic plants and of foreign floras which passed between correspondents abroad and their agents in this country, show how eager botanists were at this period to obtain accounts of the floras of foreign countries. Few of these documents are dated, but one, a list of Russian plants, was written in June of this year. At the end of the list the writer adds: 'and many other which I know not and are not to be found in my herball. I will have their effigies drawne, and will hereafter send them you with their leaves and Russe names and vertues.

Mosco, 12 Junij 1632

'Robert Tewe.'

¹ *Ger. emac.* 65.

The note which follows may or may not refer to Tewe's expenses.

'150^l p. ann. 12^{li} 10^s a moneth.'


It is clear that Tewe must have been a member of, or have been helped by, the Muscovia Company, then engaged in active trade with Russia.

The important domestic event of his marriage occurred in this year. The Licence issued by the Faculty Office of the Archbishop of Canterbury bears date 15 November 1632.

'John Goodyer of Beryton co. Southampton, gentleman, bachelor, 40, and Patience Crumpe, spinster, about 30, daughter of Walter Crumpe, late of St. Giles in the Fields, Middlesex, tailor at St. Giles in the Fields or St. Gregory's London.'

There is evidence that John and Miss Crump were friends of long standing. In a letter dated 9 November 1621 Laurence Davis ends with a postscript: 'I praye remind me kindly to yo^r fellows Patience, Mr. Parker, and Henry Henly'. At that time Goodyer and presumably

Patience also, were living at or near Droxford. Soon after the wedding he moved from Mapledurham to a house in the Spain in Petersfield. The Spain is a picturesque open place or square which is said to have received its name from the Spanish merchants who resorted there for



PETERSFIELD.

wool-dealing.¹ The street of approach from the market-place is still named Sheep Street. In the eighteenth century a horse-market was held there.

John Goodyer's house 'is still standing to-day, and is one of the most interesting in the whole of Hampshire. The half of it which I had the privilege of inspecting has been burdened with the ridiculous name of St. Aubyns, but until 1907 it bore legally the title which it bore in

¹ J. Williams, *History of Petersfield*, p. 34.



West side showing the blocked windows



South side

GOODYER'S HOUSE IN THE SPAIN, PETERSFIELD

John Goodyer's time, "The Great House". The word "great" was not synonymous with "big"; it simply meant of chief importance'.¹ In April 1917 a tablet was imbedded in the brickwork of the front of the house :

JOHN GOODYER
Botanist and Royalist
(1592-1664)
lived here.


In August last I made a pilgrimage to Petersfield to see the house. The two-gabelled front facing the Spain was a disappointment: it has been altered out of recognition by the substitution of sash windows for the old casements: Goodyer would not know it. To south and west there are still a few of the original features. On the south an old doorway, with stone jambs and flat-pointed arch-stone dated '1755', leading out into the garden, is almost certainly the original front door to the house. The old mullioned windows, with splayed jambs of red brick, have mostly been covered with wall-tiling, though several, now walled up, are an architectural feature in the small yard at the back of the house. Perhaps the window-tax may have been the cause of the blocking of the windows, and the utility of their embrasures inside, when fitted as cupboards, may have led to some hesitation in reopening them.

Within the house there is much to delight the antiquary. The heavy oak beams crossing the ceilings, the uneven floors of thick and broad oak planks, the broad slanting corbels close under the ceilings, which support the hearth-stones above, and many other details, all help him to reopen in his mind the old windows whose splays and mullions he sees in the walls, and to think away the thin partitions that now subdivide the rooms in which Goodyer had his library, and where he worked and wrote.

Of unusual interest is the old stairway, a square brick and stone built structure at the back of the house, formerly lit on two sides by small windows, now blocked up, placed

¹ M. E. Wotton, *John Goodyer in Hants and Sussex News*, 11 April 1917.

in a spiral to follow the stairs. The window-openings can be seen from the inside, but on the outside they have been concealed by more modern buildings raised against them. The conjectural arrangement of the original windows is shown in the sketch.



GOODYER'S HOUSE IN THE SPAIN.

(The blocked windows of the Stairway have been reconstructed by Mr. A. E. Gunther from photographs taken for the purpose by Mr. Llewellyn Bradley.)

And then there is a long four-light mullioned window up in the wall of the present kitchen, so high above the floor that to one architect it has suggested a chapel, but perhaps the original builder knew the advantage of a top light to his kitchen, as well as of a window through which the neighbours could not peep. Many constructional details,

including an upstairs fire-place in a massive pier, facing a window, and within four feet of it, still need explanation: the well in the cellar and a possible underground passage need exploration. It is greatly to be desired that the house may find an owner who will undertake the intelligent and sympathetic restoration of what is perhaps the most interesting house in Petersfield.

The papers of the next two years bear evidence of his change of abode and change of state. Memoranda and accounts show that he was occasionally commissioned to do shopping for his wife. Notes of one and a half yards of white cotton, of hooks and eyes, of silk and buttons, tell a story as clear as it is brief, while on the other hand we read of the employment of certain labourers on 24 June 1634 for 'digging my grounds'.¹ This evidently refers to a new garden in Petersfield.

On 5 March 1632 he noted the beneficial result of the application of Honewort (*Carum segetum* Benth.) to a swelling or 'Hone' in the cheek of Mrs. Mooring, when a young girl.

1633

The summer after his marriage he found on 2 July the local Starry-headed Small Water Plantain (*Damasonium stellatum* Pers.) between Sandie Chappell and Kingston, having previously found it on Hounslow Heath. Johnson found it a little beyond Ilford in the way to Rumford.² But Goodyer had observed it in 1618.

On 4 July he was busy with the Ferns of the neighbourhood of Petersfield. 'I have observed' he wrote 'fower sorts of Ferne, by most wrighters esteemed to be the male Fern of Dioscorides: by Anguillara, Gesner, Caesalpinus, and Clusius accounted to be the Female, and so indeed doe I thinke them to be, though I call them the Male with

¹ Theophilus Hasted	} 24 Junij	Tho. Bowyer	} digging my grounds.
Tho. Crowcher		Tho. Sylvester	
Andrew Ansell			

[MS. f. 14

² *Ger. emac.* 418.

the multitude. If you looke on these Fernes accordinge to their seuerall growthes and ages, you may make many more sorts of them than I have done; which I am afraid hath beene the occasion of describinge more sorts than indeed there are in nature. These descriptions I made by them when they were in their perfect growthes' (cf. p. 181).

His No. 1 'Filix mas ramosa pinnulis dentatis' is the earliest reference to the Broad Shield-fern (*Aspidium dilatatum*).

'This groweth plentifully in the boggie shadowie moores neare Durford Abbey in Sussex, and also on the moist shadowie rocks by Mapledurham in Hampsheire . . . and I have found it often on the dead putrified bodies and stemms of old rotten okes, in the said moores, neare the old plants I have observed verie manie small yonge plants growinge, which came by the fallinge of the seed from those dusty scales: for I believe all herbes have seeds in themselves to produce their kindes, Gen. i. 11 and 12.'

His second species was the Male Shield-fern (*Aspidium Filix mas*). 'This grows plentifully in most places in shadowie woods and copses.' It had not been previously recorded for Hampshire. The variety *Aspidium Filix mas var. affinis*, was noted 'in many places in the shade'. The Prickly Shield-fern (*Polystichum lobatum*) or the allied Angular Shield-fern (*P. angulare*) was also abundant 'on the shadowie moist rockes by Mapledurham neare Petersfeld in Hampsheire'.

The Marsh Shield-fern (*Aspidium Thelypteris* Sw.) is probably the species described under the name 'Dryopteris Penae et Lobelii' on 6 July; if so, Goodyer's is the earliest British record of it.

'Manie yeres past I found this ferne in a verie wett more or bogge beinge the land of Richard Austen called Whitrowe moore, where Peate is now digged, a mile from Petersfeld in Hampsheire, and this sixt of July 1633 I digged there manie plants, and by them made this description. I never found it growinge in anie other place.'

The descriptions of these Ferns were almost certainly

prepared for his friend Johnson's new edition of Gerard's *Herbal*, which appeared in the winter of 1633, and has hitherto been the principal source of information about Goodyer's plants, many of which are acknowledged in the most exemplary manner. On the other hand many species are just referred to him without further notice of the date or place of discovery, and doubtless other species of Goodyer's, having come to Johnson's knowledge independently, are not acknowledged as his at all. Among the Goodyer MSS. there is a list of descriptions of 113 plants sent by him on 5 March 1632, of five more sent on 12 March, and of two more sent on 19 March, to Johnson for incorporation in the book, and with the exception of seven,¹ all these descriptions are extant.

In the preface to the edition, dated from his house on Snow-hill 22 October 1633, Johnson acquaints the reader with what he has performed, 'either by mending what was amisse or by adding such as formerly were wanting', or by putting out descriptions and words that were not very necessary. Moreover, he conscientiously marked all new figures and text with signs ‡, by which the new work can be readily distinguished from the old. In all this Johnson's work shows much careful thought; and it is here that we would desire to quote from his concluding sentences.

'I must not in silence passe over those from whom I have received any favour or encouragement, whereby I might be the better enabled to performe this Taske. In the first place let me remember the onely Assistant I had in this Worke, which was Mr. *John Goodyer* of Maple-Durham in Hampshire, from whom I received many accurate descriptions, and some other observations concerning plants; the which (desirous to give every man his due) I have caused to be so printed, as they may be distinguished from the rest: and thus you shall know them; in the beginning is the

¹ The seven descriptions which we have not been able to trace are: *Sonchus Africanus Boelii*, *Alsine major repens Clusio*, *A. palustris foliis tenuissimis*, *Ranunculus flammeus aquatilis angustifolius hirsutus flore magno*, *Anagallis erecta floribus albis*, *Radix cava minima viridi flore*, *Draba lutea siliquis longissimis, v. strictissimis C. Bau.* [MS. f. 134.

name of the plant in Latine in a line by it selfe, and at the end his name is inserted; so that the Reader may easily finde those things that I had from him, and I hope together with me will be thankfull to him, that he would so readily impart them for the further increase of this knowledge.'

Dr. Reynolds Green has estimated that the new book contained about 2,850 descriptions of plants, so far the largest number included in any herbal. This made the work the most important and influential of its time, but we cannot agree with Dr. Green in attributing the name by which it is widely quoted, 'Gerard Emaculatus', to Ray, for How in 1650, and John Goodyer still earlier, had already made a practice of referring to it by that name. Nor can we agree that Johnson was the first to depart from the practice of the older botanists in relying on their gardens for the plants they described. Goodyer had long paid special attention to wild plants.

Green remarks, as have some others, on the 'rapidity with which Johnson worked' and that 'he had but little assistance'. But this is a wrong view. He had the assistance of the best English botanist of the day.

The work of the two men was essentially on different lines. Johnson was an M.D., he had translated the surgical works of Ambrose Parey; he discussed the medicinal properties or *vertues* of plants with greater gusto than he displayed for their morphology. Goodyer was a scientific botanist, 'second to none in his industrie and searching of plants, nor in his judgement or knowledge of them'.

Johnson freely availed himself of the archaeological knowledge of Goodyer, as in the case of the figure that was supposed to be the oldest drawing of *Saxifraga*, taken from an illustrated manuscript of Apuleius Platonicus.

In the case of the confusion between *Solidago sarra-cenica* and *Arabis quorundam*, Johnson notes 'My very good friend Mr. Iohn Goodyer was the first, I thinke, that observed this mistake in our Author; for which his observation, together with some others formerly and hereafter to be remembered, I acknowledge myself beholden

to him' (*Ger. emac.* 275). Nor did Goodyer's criticisms cease when the 'emaculated' Gerard appeared. Among his notes are four pages of suggested emendations to the first twenty-two chapters of the book, which are characteristic of the accuracy and carefulness of his work.

Goodyer's Corrections to Johnson's Gerard's Herbal.

The first booke. Ca. 1.

Description of Comon Meadowe Grasse.

p. 1, l. 2 *from bottom.* For *light read smooth.*

p. 2. *Tyme.* I have seene it flowringe in the beginninge of March A^o. 1619.

Names. After ἀγρωϊς *read,* And this particular meadowe grasse is called of Theophrastus πῶα, as Bauhinus hath it in his *Phytopinax*, pag. 4.

Nature. Number that which is spoken for the nature amongst the vertues for indeed Pena affirmeth that the seed of hay beinge beaten forth, many Physicians doe use for the stopping of the inward parts, beinge druncke: and applied to the dissolvinge of hard tumors and wind.

p. 3. *Vertues.* Theis vertues doe properlie and trewlie belonge to *Gramen caninum* pag. 22.

D. It is apparent that Fernelius, li. 4, ca. 4 *de methodo medendi*, meaneth the roots of *Gramen caninum* to have theis vertues.

Ca. 2. *Names.* L'obell calls the first of theis grasses, *Gramen minimum Xerampelinum*, and Xerampelinus color is a color somewhat ruddie, and therefore this name cannott belonge to White dwarfe grasse.

Ca. 3. *Names.* He mistaketh, for that which L'obell calleth *Agrorum venti spica*, and *Gramen agrorum*, is this 2 kind called here called *Gramen harundinaceum*.

He mistaketh the 2 grasse also for it is not L'obell but Tabernaemontanus that calleth it *Gramen harundinaceum*.

The titles over the figures I would amend thus:

1. *Gramen pratense vulgatius.* Common Meddow Grasse.

2. *Gramen minus vulgatius.* Small Common grasse. For the description sheweth it not to grow in Meddowes.

The figures are better in L'obell's *Icones*, p. 1, which are the same with Dodoneus in Latyne of the laste edition, to which you may refer them.

Description. Dodoneus describeth not the particular Meddow grasses but in generall only, therefore you shall doo well to examyn the descripcon both of the first and second, by the *Advs.* & the *Observ.* unlesse you cann add any notorious difference in them from other grasses, out of your own observation.

Ca. 4. Both their descriptions are in Dutch & in no Author that I have, and quere for what grasse the figure under the title of *Gramen maius aquaticum* must serve.

Ca. 5. The description of *Gramen Sorghinum* is also in Duch. Somethinge may be added to the description of the root of *Calamogrostis*, if the worth of the grasse deserve it, & you thinke it pertinent.

Ca. 6. The grasse under the title of *Gramen Panniculatum*, p. 8, seemes to be *Gramen amourettes Clusii*, p. ccxviii. If it be so? howe came it to be *Gramen tomentosum & acerosum* Lobell: in *Icon.* pag. 6. I have two editions of Lobell's *Icons* printed at severall tymes, the first MDLXXXI, printed at Anwerpe, and hath but one table, viz. the lattin, french, duch all together; the other was printed at Anwerpe A^o. MDXCI and hath severall tables, everie language by itself. The first it seems was one Willm. Mounts, a Physician dwellinge at Mallinge (in what Sheere I knowe not) who hath added to some herbes certaine noates, and thus he hath noted of *Gramen paniculosum phalarioides* Lobel, *Icon.* p. 7, 'Grasse called in Surrey branched grasse, in Corne: and in orchards, or shadowie places usuallie mowen. They seeth it in water with Purslaine and small Reysons for wormes, in the somer tyme, and geve it comonly to very younge children'. I must leave this Grasse (which Gerard hath confusedlie written of) to be sett dulie, and in his true place by you.

Gramen sylvaticum, I knowe not where it is written of, in Tabermont I thinke.

Ca. 7. Their descriptions are all in dutch.

Ca. 8. The 1 is described I knowe not where, the 2 you shall find in the *Obs.* p. 10, the eares are not described.

Ca. 9. The 1 is in the *Obs.* p. 10 the 2 I knowe not where, belike in Tabermont.

Ca. 10-16, 18, 19. Their descriptions are in the Teutonick or Duch.

Ca. 16. The 2), the description saith the knobs or buttons growe on both sides of the stalk, the figure hath them but on one side, query whether this be not *Gramen montanum spicatum Clusii*, p. ccxix.

Ca. 17. The 1) I can observe no such cuttinge qualitie in the edges of the leaves of this grasse.—*Vertues.* Many more vertues are spoken of in *Advers.* p. 468, and if you please you may add the best of them.

Ca. 21. I see nothing that I can amend.

Ca. 22. *Cyperus Typhinus* I know not where to find the descriptions.

'Venarum spiracula laxat Cyperus.' Only Turner hath taken notice of these words which I have added. I take it the meaninge of it is that it openeth the small branches of vena porta, called mesentericae venae. If it be so?, it is a speciall vertue not to be omitted. [MS. ff. 149-52]

Goodyer also made several corrections in the text of his copy of the 1633 edition of the *Herbal*, e. g. on p. 567 he notes that the figure given for *Saxifraga anglicana alsinefolia* is really the picture of *Synanchica*, and should be placed on p. 1120. Also that the adjoining figure named *S. palustris* is really an *Arenaria*. This mistake explains Johnson's confused account of *Saxifraga anglicana* on p. 568.

His other additions consist of a few medical notes on the virtues of certain herbs, e. g. *Tithymalus cupressinus*, p. 499, *Elaterium*, p. 913, and *Vicia maior sylvestris*, p. 1229.

1634

After Johnson had sent his *Gerard emaculatus* to the printers, we are left with next to no published news of Goodyer's doings for the next twenty years. Fortunately

many papers among his manuscripts help to bridge the gap. He still does shopping for his wife (p. 381); there is a letter of 24 June showing that he was interested in the date of the Surrey Assizes.

'Mr. Worlidge you were intreated to wright me word, when Surrey Assizes were kept, at what place and what Judge was to sitt on the Nisi prius; which hetherto you have not done. If you can be at Surrey Assizes & our triall goe forward I must goe to the Bp. of Wint¹ & expound his letter to that Judge, which will aske some time to doe, & I must also goe before hand to London to prepare our witnesses. In regard whereof I pray you this weeke wright me word all those things I have spoken of, & if any other things you knowe to be therein necessarie, And so I rest

Your assuredly

24 June 1634.

[MS. f. 14

In August he was away in Sussex on one of the summer excursions, which he generally performed on horseback, having previously, as was his way, made careful notes for the intended journey.

From London to Lewsam	3	fro London to Croydon	7
to Brumley	4	to Godstone	7
to Farnborowe	3	to Lingfield	6 wch is
to Rethered	5	4 miles from East Grinstead.	
to Sen oak [Sevenoaks]	1		
to Tunbridge	8	—at Riverhill	
to Ffant [Frant]	5		
to Mayfield	5		
to Black boyes	6	—34	
to Ringmer	4		
to Lewis	2		

Mayfield Carrier John Manser lyeth at the White Hart in Southwark: comes in Thursday out Friday.

Lewis Abell Tabrett lodgeth at the Tabott, comes in Wed., out Thurs.

W^m Barham of Mallinge halfe a mile from Lewis but goes to Borne 10 myles from Lewis. Mountaine Neppe.

Mr White of West Tarringe.

Mr Ric Relf of Tenterden 4 miles from Rumney marsh.

Putt in the L.T. 8 *li.* 8s. 10*d.* 4 Aug.

[MS. f. 62

Goodyer visited Mayfield at the time of the greatest

¹ Walter Curle.

prosperity of the iron industry in Sussex, and although the connexion between iron smelting or forging and the county flora may not appear very intimate, yet it is far closer than might be supposed. In 1607 there were, or had lately been, nearly 140 hammers and furnaces for iron in Sussex alone, and each of them spent 'in every twenty-four hours two, three, or foure loades of charcoal, which in a yeare amounteth to an infinit quantitie'.¹ About the year 1640 some 1,300 cords of wood were being used at one works alone, and the woodlands were in danger of being lost.

Jove's oak, the warlike ash, vein'd elm, the softer beech,
Short hazel, maple plain, light asp, the bending wych,
Tough holly, and smooth birch, must altogether burn,
What should the builder serve, supplies the forger's turn.
Drayton, *Polyolbion* (1612).

The botanical results of the journey are noted on the same page.

Dentaria baccifera. [Coral-root. *Dentaria bulbifera* L.]

At Mayfield in a wood of Mr. Stephen Penckhurst, called High-wood, and in another wood of his called Foxholes.

Oxyacantha in Rumney Mersh neare the house of Mr. John Snavel the place called Whey street, flours at about Xmas 1605 in November.

[MS. f. 62

Elsewhere he noted '*Dentaria bulbifera* Lo. 687, G. 833, in Foxholes wood in Mayfeilde parish 6 Aug. 1634', which fixes the date of his tour.²

On 9 August at Buttersworth Hill he collected '*Ferrum equinum Germanicum siliquis in summitate*' with ripe seed. (*Hippocrepis comosa* L.) For the year 1636 we have only a List of 'Virginia seeds recd. from Mr. Morrice 18 March' (p. 370), but in the case of this document, which is in the handwriting of John Parkinson, we cannot be sure that it came into Goodyer's possession at so early a date.

¹ Norden, *Surveyor's Dialogue*, *Suss. Arch. Coll.* ii. 192.

² I have suggested elsewhere that the little colony of Coral-root near the Church Meadow at Droxford may have sprung from roots of his planting.

1637-40

On 21 August 1637, when visiting his brother-in-law William Yalden at Sheet, he saw 'Batata Hispanorum, or Common Potatoes'. They were the Sweet Potatoes, *Ipomaea Batatas*, such as could be purchased at the Exchange in London, and were liable to be killed by the first frosts. The tubers 'howsoever they be dressed, they comfort, nourish, and strengthen the body, procuring bodily lust, and that with greedinesse'.

Goodyer was acquainted with 'Potatoes of Virginia', but we do not know that he cultivated them himself.

There is no evidence that he ever accompanied Johnson and his 'socii itinerantes' upon their herborizing excursions, but he certainly took the greatest interest in their discoveries and helped them with their reports. He possessed the accounts of their Kentish tours in 1629 and 1632, and added to the list of plants in the former from his own knowledge. His presentation copy of the *Mercurius botanicus* inscribed '28 Octob. 1634—Ex dono Thomae Johnson', containing the account of the tour in Wales in which Stonehouse of Magdalen College also took part, is similarly annotated, and has many plant names picked out by yellow paint marks—a favourite method of his. The Appendix to this work, a treatise on the Waters of Bath (1634), as well as the lure of new plants, may have determined him to visit Bath in 1638.

Johnson (*Mercurius, pars altera* 1641) is our authority for believing that Goodyer found a new Speedwell on St. Vincent's Rocks near Bristol, and as the statement is left unaltered in Goodyer's own copy of the book we may take it that he did botanize there.

And nothing is more natural, for the grand and precipitous cliffs of St. Vincent's Rocks have always been classic ground to the botanist. Their vegetation is luxuriant: their ledges, crowded with an abundance of good plants in a small area, are comparable only to the similar floral

wealth of Cheddar Gorge. This towering limestone mass and the adjoining Downs, rising to a height of over 300 feet, overlook the tidal Avon in so picturesque a fashion that Clifton's river scenery will ever be famous.¹

And if they are a Nature reserve now, what a paradise these rocks must have been in the days of Lobel and Goodyer when they were clothed with 'millions' of ferns.

The father of English botany, William Turner, discovered *Trinia* or Honewort here. It is one of the choicest indigenous plants in England and of great local interest. In 1562 he recorded '*Peucedanum* . . . I found a root of it at Saynt Vincentis rock a little from Bristow'.

The next visitor, L'Obel, found the 'Mules Fern' about 1569. Gerard 'spent two daies upon the Rocks to seeke for *Meum*', which had been reported to him as growing there, probably in mistake for *Trinia*. In 1634 Johnson and his 'socii itinerantes' guided by John Price, a jovial apothecary of Bristol, directed their steps to the famous Rocks and the precipitous cliffs commanding the banks of the Avon. It is no wonder that Goodyer was drawn thither also.

And he was rewarded by finding a species new to the British flora, the Welsh Spiked Speedwell (*Veronica hybrida* L.), which he called 'Veronica recta mas, Lob. Ger., vulgaris recta Clus.' It is the same as the 'Great Speed-well or Fluellin. Found at Saint Vincents Rocke by Master Goodyer'.² White³ states that it is still abundant on the more inaccessible ledges of the rocks, and occasionally strays on to the riverside masonry below. It is the most beautiful of our native Veronicas, and in the Avon gorge often grows twice as large as on the Great Orme's Head in North Wales.

He verified the occurrence of the Tutsan (*Hypericum Androsaemum* L.), already recorded by Lobel (1570), Lyte (1578), and Johnson (1634), Dropwort, and *Trinia*, and

¹ White, *Flora*, 1912.

² Johnson, *Mercurius, pars altera*, 1641.

³ *Flora*, p. 463 (1912).

added the Horseshoe Vetch to the County flora.¹ His records were :

Androsæmum magnum, the Great Saint John's-wort in a wood over the . . . St. Vincents rocks nigh Bristow,

Filipendula vulgaris, *Oenanthe Fuch. Lob. Dropwort*. 'On St. Vincents rock 30 Aug.' (= *Spiræa filipendula* L., previously noted by Lobel.)

Ferrum equinum siliquis in summitate. 'On St. Vincent's Rock 30 Aug. 1638.' (= Horseshoe Vetch, *Hippocrepis comosa* L.)

Peucedanie facie pusilla planta Lob. = *Selinum montanum pumilum* Clus. = *Peucedanum pumilum*, *Petroselinum* = Dwarf Rock Parsley, 1638. (= *Trinia* or *Apinella glauca* O. Kuntze.)

Rock Stonecrop (*Sedum Rupestre* L.), 'Sedum Divi Vincentii Non Descriptum', is also said to have been first recorded by him.²

The manuscript notes in Goodyer's hand in the Johnson's *Mercurius*, 1634, have unfortunately been cut off by some wretched bookbinder, but one about 'Saxifraga palustris alsinefolia³ *Ger. emac.*' confirms 1638 as the year of his visit to the west of England. He found it 'In the springs about Smocombe wood neare Bath, on the north-east parts of the wood, j. Sept. 1638'. In modern language it is the White Sandwort (*Sagina nodosa*) that he had been the first to add to science in 1626.

Probably at the same time he became acquainted with the Wood Vetch (*Vicia sylvatica* L.), which he called 'Vicia Bathoniensia vel maxima sylvatica' and found 'In Smoak-hall Wood by the Bathe, and at the Devizes in Wiltshire'.⁴

A list of the names of useful Willows bears date 1640.

Salix aquatica lo. o. 137 g. 1203. I have seene 8 kinds of osyars thus called by the baskett makers

- | | |
|-------------------------|--------------------|
| 1. gelderlander | 5. redd willowe |
| 2. goldstone or hornead | 6. privett |
| 3. white withe | 7. black } yelster |
| 4. yealow osier | 8. white } |

¹ Not noted by White, *Flora Bristol*, 1912, p. 256.

² Merrett, p. 111 (1666).

³ *Sagina nodosa* Meyer.

⁴ Merrett, *Pinax*, 1666, p. 125. The same plant is mentioned in How, *Phytologia*, p. 129.

A similar list, on a small scrap of paper, is dated '18 Marcii 1640 of Edw. Greene (? Gawne)'. And two rather longer and later lists enumerate thirteen varieties.

- | | |
|---|-------------------------------------|
| 1. White gorerod | 8. Gilford redd osier |
| 2. Black gore rod | 9. Erlic ta rod [lesser browne rod |
| 3. Gelderlander | 10. Backward ta rod, or Privett, or |
| 4. Yealowest yelster | 11. White osier |
| 5. Hard yelster—the best | 12. Horneyead or golston. |
| 6. Erlic leavinge redd osier | 13. Oxford or Dutch golston |
| 7. Backward Leavinge redd osier-skragged. | [MS. f. 16 v. |

1643-9

The disturbed state of the country during the next few years made peaceful pursuits impossible. Hampshire men were soon in the thick of the Civil War. Lucky were those who were not driven from their homes when the 'wrong' side obtained the ascendancy.

During one of the many phases of the struggle Petersfield was garrisoned for King Charles under the then General in Command, Ralph, Lord Hopton. Goodyer, like all his family, was a strong 'King's man', and his loyalty was acknowledged by a 'Protection Order' which was granted him by Hopton. This document of great interest was accidentally found in 1907, concealed under a floor-board in a dwelling-room of Goodyer's house in the Spain in Petersfield. It runs as follows :

(SEAL) To all Colonells & Lewetennt-Colonells, Serjant-Maijors, Captains & Commanders, Officers & Soldyers of his Ma^{ts} army both of horse & foot And to all other his Ma^{ts} officers and loving Subjects whom these may concerne.

These are in his Ma^{ts} name to will & command you & every and either of you not to fayle upon all occasions to defend and protect John Goodyer of Petersfield in the County of Southton Gent : his house horses servants family goods chattels and estates of all sortes from all damages disturbances & oppressions whatsoevere to the uttermost of yo^r abillitys And that you and every of you forbear to grieve or molest him the sayd John Goodyer or any of his as

aforesayd requiring hereunto yo^r due obedience as you will answere the contrary at yo^r uttermost perill Given under my hand & seale the 9th December, 1643

RALPH HOPTON.¹

Of Goodyer's movements we know nothing for certain, but it is quite likely that a State Paper in the Record Office may refer to him. About 1649 one Daniel Cusick stated that John Goodyer was a malignant and recusant, and was constantly resident in Oxford during the war. Having adventured his own life in the service of the State, and being now in the Lord General's regiment, the informer begged his arrears out of Goodyer's estate.²

If it be true that Goodyer was 'constantly resident' in Oxford during this troubled period, he would have found many botanical friends with tastes congenial to his own.

Walter Stonehouse, now no longer a Fellow of Magdalen, would not have been in residence, but he may have joined his Oxford friends in 1648 when ejected from his Darfield living. And there were other botanists at Magdalen to whom the botanical uncle of Edmund Yalden, Fellow until 1642, would have needed no further introduction.

The senior of them, William Hooper, the arboriculturist, became a Fellow in 1643. He had been 'outed' from his Fellowship, but was allowed a pension of £30 per annum and lived in one of the College houses in the Gravel Walk. 'After he had left the College he went without a gown, and wore constantly a very long coat, like your frocks worn by wagners; and applied himself to gardening with wonderful success, digging himself with a man that he constantly hired. He would carry his spade upon his shoulders, and work hard every working day. He would likewise prune, engraft, and do other things of that kind himself. He raised several nurseries, and planted many orchards; but he did all for nothing, for he would never

¹ Mabel E. Wotton, *Hants and Sussex News*, 11 April 1917.

² Calendar of Commission for Advancement of Money, p. 1178, *State Papers Domestic*, 1649.

take anything of anybody soever. It was his constant practice to give away trees, &c. ; but then he took care it should only be to the poor and such as were in want, not to others. He was near fourscore years of age, a comely, neat, proper, upright man, and beloved and respected by all sorts of people.' He planted (c. 1660) elms in the Gravel Walk by Magdalen College ;¹ and elms on this site are well known all over the world as an essential feature in what was formerly one of the most popular views in Oxford.

In his old age he was one of the characters of Oxford, but when Goodyer might have known him he had just been recommended by Charles I for election to a Fellowship on the ground that he had 'given ample proof of his sober carriage, conformableness, and commendable abilities in the way of his studies'.

Among the younger men then up at Magdalen College were the three contemporaries Browne, Stonehouse, and Drope. William Browne, Demy 1644, was a native of Oxford, who became 'one of the best botanists of his time, and had a chief hand in the composure of a book entitled, *Catalogus Horti Botanici Oxoniensis*', 8vo. Oxon. 1658 ; Walter Stonehouse, Demy 1645, was the son of Goodyer's friend the Rev. Walter Stonehouse of Darfield, whose garden lists are preserved among the Goodyer papers ; and Francis Drope, Demy 1645, was a most enthusiastic lover of trees and author of *A short and sure Guide in the practice of raising and ordering Fruit-trees*. 8vo. Oxford, 1672. And, in any account of the botanists of Magdalen of this early period, should also be mentioned the unknown writer of marginal notes in the Bodleian copy of Lyte's *Herbal*. This book was successively in the possession of

¹ Hearne, *Diary*. According to one account Hooper's Elms were replaced by others in 1680, but, be that as it may, the Gravel Walk elms, after forming for more than two hundred years an incomparably beautiful setting to the grey stone architecture of the Great Tower of Magdalen, were wantonly felled before their time in 1916, when many who would have advocated their retention were away at the War.

a Thomas Gill and of a John Herbert, 1619. The writer of the notes was almost certainly a Magdalen man who had studied at Padua.

At this time Goodyer may have made the acquaintance of Dr. Philip Stephens, Principal of Magdalen Hall and collaborator with Browne, and of William How who came up as a Commoner to St. John's College in 1637, and took his Master's degree in 1645.

Goodyer's other friend Dr. Merrett, the author of the *Pinax rerum Naturalium Britannicum*, had been a student both of Gloucester Hall (1631) and of Oriel College (1633-4). He was created a Doctor of Physic in 1642, and afterwards became one of the original Fellows of the Royal Society. At New College was William Cole, the future author of *Adam in Eden*; and Jacob Bobart was gradually forming the first University collection of plants in the new Physic Garden, the catalogue of which he published in 1648.

And yet Oxford was in the very midst of troubles. At the end of 1642 the Royalists gathered round the King at Oxford, students and citizens alike worked together on the fortifications and barricades: the College plate was being surrendered to the minters: fighting in the neighbourhood was incessant. Goodyer's friend, Thomas Johnson, appears to have been in Oxford on 9 May 1643 to receive the D.Ph. degree, but he must have left soon afterwards, for he is heard of as a Colonel of Horse and one of the defenders of Basing House, which was being strongly fortified by the Marquis of Winchester. And it was there that he met his death in September 1644, fighting for the King. The Royalists were defeated at Alton and Cheriton. Winchester was taken by Cromwell in October 1645, and the King became a fugitive. What followed is matter of history. On Midsummer Day, 24 June 1646, the Royalist garrison of Oxford, 3,000 strong, 'marched out of the town through a guard of the enemy extending from St. Clement's to Shotover Hill'.

In 1648-9 the Parliamentary Visitors, after many delays, expelled all from the University who did not submit to their visitation, including young Stonehouse and Drope. The latter answered 'I cannot submit for fear of perjury'.

1651

After the Civil War Goodyer was unquestionably the best botanist in England. Parkinson had died in 1650; and, until Morison returned in 1660 and Ray had reached his full development, there was no one to approach him in knowledge of our native plants or of the whole range of botanical literature.

The high reputation that he had acquired naturally brought him visitors and correspondence. Elias Ashmole, the antiquary, had not long 'entred upon the Study of Plants'. A note in his diary is to the effect that 6 June 1648 'about three of the clock was the first time I went a Simpling. Dr. Canter of Reading and Mr. Watlington¹ an apothecary there, accompanying me'. And on '19 Oct. 1651 my Father, Backhouse,² and I went to see Mr. Goodier, the great botanist, at Petersfield'. Unfortunately we have no further account of the visit, but possibly Ashmole's recent experiences of the virtues of Bryony may have formed part of the conversation. He had fallen ill of a surfeit occasioned by drinking water after venison at the Astrologer's Feast in London. 'I was', he wrote, 'greatly oppressed in my stomach; and next day Mr. Saunders the astrologian, sent me a piece of bryony root to hold in my hand, and within a quarter of an hour my stomach was freed of that great oppression, which nothing which I took from Dr. Wharton could do before.' It is characteristic of the writings of Goodyer that no hint of this kind of quackery appears. He was evidently able to put Ashmole

¹ John Watlington, buried 2 October 1659.

² Mr. William Backhouse, astrologer of Swallowfield in Berkshire, had communicated so many secrets to Ashmole that he caused his pupil to call him 'Father' (Ashmole, *Diary*, 3 April 1651).

on sounder lines of thought; for after this visit Ashmole 'took a journey into the Peake, in search of plants and other curiosities'; and as there is no entry between the 15th and the 29th of October of this year in the notebook,¹ in which he usually cast his horoscopes, we may assume that he was not encouraged to foretell Goodyer's future by the stars.

1652-6

The books printed in Oxford during the next few years bear witness to the pleasure and profit that many were deriving from their gardens. And in illustration we may cite the works of Ralph Austen on Fruit Trees, various editions of which were published in Oxford in 1653, 1657, 1658, and 1665, of John Beale whose *Treatise on Fruit Trees shewing their manner of Grafting, Pruning, and Ordering, of Cyder and Perry, of Vineyards in England, &c.*, appeared in Oxford in 1653, and of Robert Sharrock, Fellow of New College,² whose *History of the Propagation and Improvement of Vegetables, by the Concurrence of Art and Nature*, 8vo. Oxford, 1660. These works show the natural tendency of the time, a utilitarian tendency that our recent experiences towards the end of the Great War will teach us to connect with the troubles of the forties of the seventeenth century.

During the last ten years of his life Goodyer's occupations appear to become more and more sedentary. When a man is over sixty years of age, he must perforce leave the searching for new plants to the young and active. John Goodyer now devoted himself to his books. The dated entries in the covers of his volumes show that he kept in the closest touch with the London booksellers; indeed, in some cases he appears to have secured a work in advance of the day of publication. Some of his books he pro-

¹ MS. Ashmole, 374, which contains the horoscope of John Tradescant, the younger.

² Sharrock gave several medical books, with his autograph, to New College Library. They are still, we are glad to think, on the shelves: among others a copy of Lower, *de Corde*.

cured through Dr. How, some through Dr. Dale, and they came down to Petersfield by carrier, or in the trunk of Mrs. Heath, presumably the wife of his friend and neighbour the Rev. John Heath.

He was now closely associated with a botanist in the work of preparing a list and a synonymy of all known British plants, incorporating and extending the lists drawn on by How. The greater part of the labour of this work fell on a collaborator who had access to Goodyer's books and made notes in them: his name is not known to us for certain, but we have his manuscript. The evidence all tends to identify him with the Dr. Dale just mentioned, but the matter will be again discussed below, p. 295.

This last period of his life is also marked by a literary labour that remains a record which to this day has never been broken. It was the writing out of the Greek text of the *Materia Medica* of Dioscorides, and the rendering it into English. Goodyer therefore accomplished in the case of Dioscorides a work which not one of the tens of thousands of Greek scholars who have lived before or during the past three centuries have been known to have attempted, a worthy sequel to his translation of Theophrastus.

And neither Theophrastus nor Dioscorides can be considered as of no import. Of Matthioli's *Commentaries on Dioscorides* alone thirty-two thousand copies were sold before 1561, and it passed through seventeen editions. His works have been translated into almost every civilized language, except English, and there is no separate article on Dioscorides in our national Encyclopaedia.

The interlinear translation of Dioscorides fills six quarto volumes of 4,540 pages in all. It is most clearly written, the Greek text being easier to read in Goodyer's manuscript than in the 1499 edition, printed at the Aldine Press at Venice, Goodyer's copy of which has now been restored to its proper place among his books in the Magdalen Library.

3 Martij 1683

2071

Πελαγία Διοσκορίδης Αναξαρχεύς,
of Pedacius Dioscorides y Anazarban,
ὄρει ἰατρικῆς ὕλης τέταρτον
of medicinal matter, 4th

βιβλίον

βιβλίον ὕλης
books 3 23

Ἐν τοῖς τοῦ βιβλίου πρό τούτων, φιλοτάτε ἄρχει

In the three books before this, most lovingly, I have

παροδόντες περὶ ἀρωμάτων, καὶ ἐλαίων,

spoken, of stomatical matters, & oyles,

καὶ μύρων, καὶ δένδρων, καὶ ῥιζῶν, καὶ βιτησῶν

& ointments, & trees, & living creatures, & roots,

καὶ λαβάνων, καὶ εἰλών, καὶ χυλιμάτων, καὶ βοτάνων,

& rockherbs, & roots, & juices, & herbs,

καὶ ὡσεύματων. Ἐν τούτῳ οὐκ ἐλάττω

& of seeds. In this being y^e fourth.

διαλεγόμεθα περὶ τῶν βοτάνων, καὶ εἰλών

wee will discourse of y^e herbs, and roots

λειπομένων

left unsproken off

1420
31
32-5
33
34
35
37
38
39
40
41
42-15
43
44
45-20
46
47
48
49
50
51
52-25
53
54
55
56
57-30
58
59

Gooder's Greek from Latin of Dioscorides in the
Nicolaus manuscript of June 1477
1470
1471
1472
1473
1474
1475
1476
1477-30
1478-40
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600

GOODYER'S INTERLINEAR TRANSLATION OF DIOSCORIDES

Every page bears evidence to his neat and methodical habits. At the beginning and end of every volume he wrote the dates of the beginning and end of his labours, and also of the time that was spent on reading over and revision. The entries are of interest because they show the speed with which the work was accomplished.

	Pages.	English.	Greek.
Vol. I.	1-728	27 Apr. 1652 h. 10 a.	29 Aug. 1655 ho. 2. p.
II.	733-	28 Marcii 1653	13 Nov. 1655
	1470	18 Aug. 1653	18 Jan. 1655
III.	1471-	18 Aug. 1653	21 Jan. 1655 Exam.
	2070	3 Martii 1653	1 Feb. 1655 Exam.
IV.	2071-	3 Marcii 1653	4 Feb. 1655 Exam.
	2658	20 Junii 1654	14 Feb. 1655 Exam.
V.	2659-	20 Junii 1654	15 Feb. 1655 Exam.
	3174	31 Aug. 1654	21 Feb. 1655 Exam.
VI.	3497-	2 Nov. 1654	4 Marcii 1655 Exam.
	4540	29 Aug. 1655	17 Marcii 1655 Exam.

Other entries show that he paid an assistant, probably as a reader :

Vol. II.	‘Received of Mr. John Goodyeere, upon this 28 of March 1653 . . . 20 ^s And 18 of Januarie 1655 . . . 15 ^s ’
IV.	‘3 March 1653 . . . £2 0 0 20 Junij 1654 . . . £0 10 0’
V.	‘20 Junij 1654 . . . £1 0 0 4 Sept. 1654 . . . £0 10 0’
VI.	‘10 Martii 1654 paid . . . 20 ^s 29 Aug. 1655 paid . . . 20 ^s ’

In Vol. V there is an entry for ‘3^s binedinge’ on 14 September 1654, and in the last volume is written :

‘22 Sept. 1655 cariage upp and doone
the binedinge’

A few notes on a duplicate page (p. 2071, MS. f. 17) of the fourth volume, written on 3 March 1653, show that he was acquainted with at least eighteen editions of Dioscorides dated ‘1499 gr., 1500 gr., 1506 gr., 1512 padua-nensis, 1518 gr., 1529 gr., 1529 Herm. Barb., 1549 Gualt. Riffe, 1549 gr.-lat., 1552 Ruellius, 1552 Lacuna, 1554 lat. Mat., 1557 Jan Cornarius, 1558 Amatus Lusitanus, 1566 french, 1581 Alphabeticum Empyricum, 1591 Antonini Pasini, 1598 gr.-lat. Saracen.’ There is also a note that

Theodorus Gaza translated Theophrastus from Greek into Latin and dedicated it to Pope Nicolas: a calculation shows that Nicolas was pope from 1447 to 1455 and that Gaza died in 1478. But of still greater interest are the words:

ιοάννης έαθ Johes έαθ 1 obolon

which evidently refer to Goodyer's coadjutor John Heath, who will be mentioned again presently, and to whom we believe that the payments just recorded were made.

Had Goodyer wished for a Preface to his translation he might well have used the words of his senior, Henry Lyte:

'If perchance any list to picke a quarrell to my translation, as not being either proper or not ful, if I may obtaine of him to beare with me til he himselfe shall have set foorth a better . . . and in the meane while (considering that it is easier to reprehend a mans doing than to amend it) use me as a whetstone to further himselfe withal, I will not much strive: for I seeke not after vayne glorie, but rather how to benefite and profite my countrie.'

In March he continued his studies in the botany of the ancients by translating, doubtless with the help of John Heath, the work of one of the principal commentators on Dioscorides, the *Scholia or Animadversions upon the 5 Bookes of Dioscorides of Physicall matter and upon his two Bookes of Poisons*, of Antonius Saracenus of Lyons, 'In which y^e severall readings of divers Bookes are examined, y^e different opinions of y^e old or new propounded, and some tymes reconciled, and y^e most corrupt, obscure, and difficult places of y^e Author himself, restored, illustrated, and unfolded'.

Goodyer, with his usual punctiliousness, noted that the translation was begun on 20 March 1655 at 11 a.m.

Saracen's dedication is 'To Henry y^e 4th y^e most Christian King of France and Navarre' and is dated 'y^e Calends of March 1598'. This is followed by the Introduction, which is worth quoting as an example of the literary style of the time.

*To the Gentle Reader**Salutation.*

Thou wilt wonder, it may be (Gentle Reader) how it should come into my minde, y^t I should sett about a new interpretation of Dioscorides, especially since y^t soe many famous men, renowned for Eloquence, learning and iudgement, as Hermol. Barbarus, Marc. Virgilius, Janus Cornarius, & Jo. Ruellius, have long since with great contention & aemulation taken abundant if not overabundant paines in translating him, & have donne well herein to their great commendations. But you will then leave off to wonder, when you shall have understood by what reasons I was induced, or drawn heereunto. Dr. Jo. Sambucus of godly memorie the Emperours Physition & Historiographer had formerly often importuned by letters Henry Steevens, Printer, a man most skillful of y^e twoe tongues both Greek and Latin, to sett out in print y^e Greek & Latin text of Dioscorides with y^e most elegant letters of y^e King's stampe, & y^t hee should add to y^e Margent thereof his notes, sent over a little before, or rather y^e divers readings upon y^t Author, gathered by him with great labour, by a faithfull & diligent comparing of divers antient bookes found in y^e severall Libraries of Princes. But while Steevens did prepare himself to goe about this worck, hee was minded to place Ruellius his Translation (which hee did, and not without cause, prefur before the rest) right over against the Greek text, but hee found it to be, as indeed it is, a little too free in many places, that y^e Latin did not sufficiently aunser to the Greeke: therefore for y^e acquaintance sake y^t did passe between him & mee, hee did intreat you & overintreate mee, y^t I would take a diligent review of it, & sett downe in the margent y^e correction of all y^e places in which Ruellius, either following some corrupted copie, or else by his beeing too much addicted to Plinie, as for y^e most part hee is, did not sufficiently expresse y^e sence of Dioscorides, nor y^e force of his wordes. But I had scarce compared a few of the first leaves, when being as it were deterred with y^e difficultie of soe most grievous a burthen, I was compelled to decline from my first resolution, & to change my minde. For I mett further (under favour be it spoken) with soe many places worthy of censure, & animadversion, y^t partly to avoide y^e hatefull labour, & partly to avoide y^e envie, & offence of them, which might have suspected mee to be too injurious against Ruellius his ghost, I thought it better to sett out a new translation, then to correct an others. In which I may truely say, y^t I have performed y^e part,

not of a Paraphrast, but an*Interpreter, as whoe, as much as lay in mee, have not departed a nailes breadth from Dioscorides his meaning, and yet withall have stuck close to y^e puritie of Plinie's style. But I had rather they should judge of this, which shall faithfully compare mine with their translations. Howsoever it be, I had almost made an end of y^e worck, when behold Sambucus an earnest prosecutor heereof was taken away from us by untimely death. Then Steevens began to be, as it were, faint hearted in y^e businesse, yea & although y^t many others were urgent upon y^e worck, to knit many delaies, and to put it off from day to day, either by his due employments, whereby he was some tymes diversly distracted or for y^e travellings, which befell him often in y^e meane space, & were almost continuall soe y^t he could not performe his promise nor be as good as his word. And soe it came to passe that these my Elucubrations upon Dioscorides lay concealed in my desk for many yeares. This then, when y^e heyres of Andrew Wecher did understand, being most studious of helping on learning & did withall desire to satisfie y^e wishes & expectations of some good men, at last they obtained this of mee y^t I should suffer them to be brought out into y^e sight & view of men. These are (y^t I may once say it) y^e causes of this worcke intended, & also of the suppressing of it soe long unto this tyme.

As concerning y^e Greeke text, wee have laboured, y^t as farre as might be, it should be restored to its old splendor. For although wee have relligiously followed y^e Parisian edition of y^e most learned Goupylus, as y^e most sound, & best amended of any that are extant, yet wee have, as much as might be, taken away y^e faults of y^e print, which were yet remaining. But what passages, both y^e matter it self, & y^e certaine faith & authoritie of y^e Copies, whether printed, or manuscript, did persuade to be changed, those wee have boldly changed. Yet wee have noe where yielded soe much to our owne or others conjectures, that wee have dared to doe anything without y^e Creditt of y^e better bookes, being contented only to note them with an interlineary Asterisck, which should direct thee y^e Reader to our Notes. But if, which yet falls out but seldome, wee found some places, soe corrupted & depraved, y^t there could noe sence be made of them, and yet there was no place for any Remedie either out of conjecture, or out of y^e footsteps of antient readings, there wee have marked y^e margent with an Asterisck. But y^e divers readings, both of the most renowned man Dr. Sambucus, as also of others drawne out here & there upon Dioscorides. For both y^e same Dr. Steevens communicated some

unto us, & D. Opsopœus some by comparing of y^e Palatine copies, these, I say, and y^t without choice, I have thought fitt to set upon y^e margent, y^t it might be free for every man to judge of them, and withall some Animadversions & Corrections of Interpretors, adding still y^e names of everie one out of whom wee tooke them, y^t wee might not seem to defraud any one of his desert. But wee have placed our own notes at y^e end of y^e worck, in which for the most part, there is a reason given of our different interpretation from others, as also there are sett downe our divers conjectures upon Dioscorides. And also you may finde many thinges incidently, &, as it were by y^e way, noted upon divers Authors, but especially upon Theophrastus & Plinie, as they came under mine hand.

Finally wee doe propose, velut, ἐξ ἐπιμέτρῳ, i. by way of Additament, the severall, & discrepant opinions, whether of y^e old or new writers concerning Medicinall Matter, & some tymes wee reconcile them, & besides, wee endeavour to resolve & cleare many doubts which wee met with everie where. But yet I would have you to understand, y^t I did chiefly cavell heereat with all my labour, y^t I might restore, illustrate, & explaine y^e most corrupted, obscure, & difficult places of Dioscorides himself. And soe you have been given to understand (Gentle Reader) what thinges have been performed by us in this Edition, out of which if I shall percieve, y^t you have got any fruit, it will encourage mee peradventure to divulge other more great & more profitable matters.

In the meane space, Farewell.

[Goodyer MS. 6*, ff. 5-8

The text of the work is written out on pp. 13-292 of the MS., which ends abruptly with Book 4, chap. 61, the last line being dated '2 Octob. 1656', and then follows an ominous note, which evidently refers to his collaborator.

'Johannes Heath Clericus obiit 25^{to} die Novembris 1656.'

This must have been Goodyer's neighbour, the Rev. John Heath, who was presented by the Earl of Worcester to the rectory of Clanfield,¹ in or soon after 1617. He may have been the John Heath who came up to Christ Church as a Westminster Student in 1607, and who matriculated in 1610 aged 19.²

¹ Clanfield is about three miles south-west of Buriton, and six from Petersfield.

² Foster, *Alumni Oxon.*

In 1654 we have evidence of the return of Goodyer to his old love, field botany. There is the short note

‘Behen album . . . Rotherwort 5 Maij 1654’.

And there is an interesting description of a station where ‘*Dryopteris Tragi*’ used to be found.¹

‘It growes on a bottome called Rogers Deane in y^e parish of Faringdon in Hampshire, about a mile and halfe from y^e church, a furlong from one John Trybes dwelling-house on y^e north-east part of y^e house about 2 miles from Alton above a mile north-east from Dogford Wood. Great antient beeches kept y^e sunne from shining on y^e Plants. Anno 1654 many of those trees were cut downe. The Plants y^e sunne shoane on y^e summer 54 were short y^e leaves growing on short stemms neere y^e earth, as Tabernacmont pictureth it, pag. 501, tom. 2, under y^e title of *Filicula petraca fem. 3*. Those y^t grew under y^e trees were much higher agreeable to *Tragus*’ figure pag. 538.’

In the beginning of August 1654 he recorded a new Crane’s bill (*Geranium columbinum*) in his native county.² The *Rubia sylvestris* described by him on 12 August 1655, if it be the Wild Madder (*R. peregrina* L.), is of historical interest because of its having been one of the first Hampshire plants to be recorded by our first botanist. William Turner, more than a century previously, wrote of it, ‘The most that ever I saw is in the Isle of Wight, but the fairest and greatest that ever I saw groweth in the lane besyde Wynchester, in the way to Southampton’. It is now very rare on the mainland, but that is where Goodyer may have found it.

His botanical labours in the field were almost done. We only note two occasions in the last ten years of his life on which he may possibly have left home and herborized.

Again, the summer of 1656 he found the Marsh Isnardia (*Ludwigia apetala* Walter), ‘*Holosteum perpusillum*’, which he had previously observed near Holburie

¹ In Goodyer MS. 9, under Pin. 358, there is a note ‘*Dryopteris Tragi*, 17 Aug. 1650, J. G. first saw it’.

² Morison, in ignorance of Goodyer’s discovery, attributed this species to Jacob Bobart in 1680. The plant should be called *Goodyer’s long cut Crane’s bill* rather than ‘Bobarts long cut Crane’s bill’.

in the New Forest, growing in a little lake in the east part of a heath near Petersfield, 'The water of this lake this 2 of June 1656 about 4 of y^e clocke in y^e afternoone was well neere as warme as y^e Bath-water at Bath in Summersetshire although y^e day was cloudy'. 'In a hott summer some parts of y^e lake are drie in August, sometimes before, and then the plant, which had been green, all the winter under water, flowered'. Immediately recognizing the novelty of the plant, he described it as '*Anonymos aquatica rubida, foliis Anagallidis flore luteo*'.

1657-9

In June 1657 he described what appears to be our Smooth Tare (*Vicia tetrasperma*), but as no locality is stated, we cannot claim his note as being the first evidence for the occurrence of this plant in Hampshire.

The following information on a scrap of paper was sent him by an acquaintance :

5^o March 1657

At Judge Rumseys 3 miles from Abergevenny croweth the Sweet Willowe, as I remember the plant I saw, was called

Robert Baskett.

To this a note is added in Goodyer's hand :

9^o Apr. 1658—Judge Rumsey lives in Glamorganshire by the relation of Gryffin Morgan of Malmesbury, a glasse carrier. [MS. f. 147

But though able to move about in his own county, where he found the alien *Xanthium Strumarium* in 1659 (his last recorded find), we imagine that he now felt himself too old to herborize in Wales.

But the record would have been valued by him as an addition to the last work on which he is known to have been engaged, the compilation of a new British Flora. How's *Phytologia Britannica*, published in 1650, was very imperfect, as any first attempt at so comprehensive a work is bound to be, and no one would have been in a better position to recognize its many errors and deficiencies than

Goodyer, nor more ready to remedy them. Our information comes through Edward Morgan, the 'very skilful botanist' of Westminster, who was in close touch with what was going on in the botanical world. He told John Ward in 1662 that Dr. Dale, Dr. Merrett, and Mr. Goodyer, 'next Dr. Modesy, the best botanists of their age in London, were about a new *phytologia* 3 or 4 years agoe', but that 'Dr. Modeseye's coming to towne, itt's thought, hindered itt'. Elsewhere Ward¹ also noted that 'Dr. Dale and another had a designe to amend y^e *phytologia Britannica* to adde somewhat and take out somewhat'. This contemporary account is of the greatest interest to us, because it explains the presence of certain excellent catalogues of British plants among the Goodyer manuscripts (Goodyer MS. 8, 9, see p. 296), and it also explains why they were never printed. I have not as yet found any clue to the handwriting, but I strongly suspect it to be that of Doctor John Dale, and the 'another' to have been Goodyer himself. The case will be again considered in our note on Dr. Dale, and in the light of his Will which I have recently discovered at Somerset House.

The 30 April 1659 must have been a red-letter day, for he then received the interleaved and annotated copy of the *Phytologia* to which reference has so often been made, and possibly with it the Lobel manuscripts which are described in a later chapter. Their late owner and part editor, Dr. How, died 30 August 1656.

The greater number of manuscript notes in the *Phytologia* are in How's handwriting: they include information received from William Browne of Magdalen College and from John Goodyer, obviously between 1650 and 1656. Goodyer after acquiring the volume wrote in it the notes on seven plants, printed on p. 194, including his last dated record of a plant, the Common Ragwort (*Senecio Jacobaea* L.) from Ladle Hill (1659).

¹ Ward also noted that 'Mr. Goodyer is good at Insects as well as plants.'

1662-4

The last specimen of his handwriting, which we have, was written when he was seventy years of age. His hand was evidently very shaky. It is a receipt for a Resin ointment similar to that in use until recently in the British Pharmacopoeia.

20 Mar. 1662

Rosen that is blackish

Fresh lard a wallnutt

Crowne sope a wallnutt

Boil till it sets clere (?) H . . and keepe stirringe.

[MS. f. 16

Then follows, in a steadier hand, 'Mar. 22. John Neale in Lippock, the howse is called Gurmes, hadd a third Ague, and hath lost him about a moneth, and now hath a great cough'.

Another note on the same paper is ominous :

‘ The Line of Selborne	{	Goute
		Scurvie
		Dropsie.’

These notes supply the clue to the occupation of his declining years. He was evidently applying his great knowledge of simples to the good of ailing neighbours. The latest medical works were sent him, as soon as they were printed, by his London bookseller, who evidently had a standing order to secure the sheets direct from the press. Thus he acquired Culpeper's *English Physician*, Pemel on *Simples* and on the *Diseases of Children*, Cole's *Art of Simpling*, Cooke's two works on *Chirurgery* (containing the 'Marrow of many good authors on the art of Chyrurgery'), Coghan's *Haven of Health*, and Muffet's *Health's Improvement; or rules comprizing and discovering the nature, method and manner of preparing all sorts of food used in this nation*. But the most striking confirmation of his practising I found in the single word 'phisicke', with a blank space in front of it, which occurs after his name in the opening sentence, written within a few months

of his death, in the Book of Accounts kept by the Trustees of the Weston Charity in Petersfield.

Obviously he was well known as a physician, but the writer left a blank as if in doubt whether to style him 'Doctor of Physic'—probably because he held no such degree, qualification, or licence at all. Moreover, his charitable disposition would have led to doctoring without fees.

His great friend and neighbour, Dr. John Dale, died in May 1662, having appointed Goodyer one of the overseers of his will, but we do not know whether he was able to act.

The exact day of John Goodyer's death is uncertain, but his will is dated 22 April 1664, and was proved 9 May 1664 by the executor, the Rev. Edmond Yalden.¹ He is now described as of Weston, in the parish of Buriton, co. Southampton.

IN THE NAME OF GOD AMEN I John Goodyer of Weston in the Parish of Buriton in the County of Southampton Gentleman being sick and weak in body but of good and perfect mind and memory thanks be given to God therefor revoking all former Wills by me made do this two and twentieth day of April in the sixteenth year² of the reign of our Sovereign Lord Charles the Second by the grace of God of England Scotland France and Ireland King Defender of the Faith &c and in the year of our Lord 1664 make and ordain this my last Will and Testament in manner and form following (that is to say)

First I bequeath my soul into the hands of Almighty God Jesus Christ my only Saviour and Redeemer hoping assuredly by his mercies and merits to receive pardon and remission of all my sins and to enjoy life everlasting And my body to the earth from whence it was taken decently to be buried by my Executor hereafter named in the Church yard of Buriton aforesaid near my late wife and as for such worldly goods and estate as it hath pleased God of his goodness to bestow on me I give and dispose thereof as followeth Item I give to the poor people of the Tything of Weston aforesaid twenty shillings to be distributed with-in a month after my decease at the discretion of my Executor hereafter named

¹ Archdeaconry Court of Winchester Register, 1660-6, fol. 564.

² That the beginning of the reign is reckoned not from the Restoration in 1660, but from the execution of Charles I in 1649, is a further indication that John Goodyer belonged to the Royalist party.

I give and devise unto my honoured friend Leonard Bilson Esq and my Nephew Edmund Yalden in the County of Surrey Clerk and to their heirs and assigns for ever all my messuage dwelling-house together with all the barns stables outhouses and buildings and all the gardens and orchards thereunto belonging situate in Weston aforesaid wherein I now live and in my possession and likewise all those lands in Weston aforesaid called Halfpenny Land now in the possession of Thomas Jacques together with free liberty to water and overflow the said lands as it now and heretofore hath been used for the best improvement thereof to the intent and purpose that they the said Leonard Bilson and Edmund Yalden and the Survivor of them their heirs and assigns shall grant and convey all the said messuage lands and premises with the appurtenances unto six able honest and sufficient persons their heirs and assigns as they or the Survivor of them shall think fit Upon trust and confidence and to the intent and purpose that all the yearly rents issues and profits of the said messuage lands and premises shall be employed and disposed of for ever hereafter for the putting forth and placing abroad of all such poor children of the Tything of Weston aforesaid and the overplus thereof shall be distributed unto the poorest inhabitants of the said Tything of Weston aforesaid as my said Trustees and their assigns shall think fit and if any or either of my said Trustees shall die then the Survivors of them shall convey the premises aforesaid unto the use intents and purposes and upon the trust aforesaid all the rest of my messuages lands tenements meadows and hereditaments whatsoever in Weston aforesaid I give and devise unto my said Nephew Edmund Yalden his heirs and assigns for ever Item I give and bequeath unto the said Leonard Bilson, Osmund Bilson Gentleman, William Bilson Gentleman five pounds a piece to buy each of them a piece of plate Item I give and bequeath unto all the children that shall live unto the age of one and twenty years of Anne Worlidge Widow the sum of Forty pounds of lawful money of England equally to be divided betwixt them at their several and respective ages of one and twenty years and to such of them as shall be of the age of one and twenty years at my decease to be paid unto them within three months after my decease. Item I give unto my servant Mary Blackman the sum of twenty shillings all the rest of my goods chattels household stuff and personal estate whatsoever my debts legacies and funeral expenses being first satisfied paid and discharged except all my books de plantis which I do give and bequeath to Magdalen College in Oxon to be kept

entirely in the library of the said College for the use of the said College unto my said Nephew Edmund Yalden whom I make sole Executor of this my last Will and Testament. IN WITNESS whereof to this my last Will and Testament contained in three sheets of paper together with this being fixed together I have set my seal subscribed my name the day and year above written JOHN GOODYER. Signed sealed and published as the last Will and Testament of the above named John Goodyer in the presence of Arch. Bold, Osmund Bilson,¹ John Westbrook, Richard Goddin, John Winter, William Gammon.

MEMORANDUM that immediately after the signing and sealing and before the publication hereof the above named John Goodyer did declare that his Will was that Susan the daughter of Thomas James should have and enjoy the lease of the houses he holdeth of the Dean and Chapter of Winton during all his term therein and likewise did give unto John Westbrook Gentleman his book of Chirurgery called Ambrose Barry² in the presence of Arch. Bold, Osmund Bilson, William Gammon.

The books came to Magdalen College soon after his death, but some of the manuscripts may have been a year or two later in coming, for Edmund Yalden lent his uncle's descriptions of plants to Dr. Christopher Merret to be used in his 'Pinax' in 1666. The loan is duly acknowledged by Merret in his preface, but by a mistake he refers to Edmund Yalden as Mr. Yalden Goodyer, evidently believing him to have taken the family name on succeeding to his uncle's property. This was certainly not the case.

He was buried, as he directed in his will, in 'the Churchyard of Buriton near his late wife'. No stone marks the spot, and when Canon Vaughan wrote his charming account of him in 1909,³ no memorial commemorated his benefactions to the parish. Through the devotion of Miss Mabel Wotton a sum of money, towards which Magdalen College contributed five pounds, was col-

¹ Sir Thomas Bilson had four sons and two daughters: 1. Thomas Bilson m. Edith da. of Peter Bettesworth of Finning, co. Sussex; 2. Leonard B. m. Eleanor da. of Sir W. Lewis, Kt.; 3. Osmund B.; 4. William B.; 5. Anna; 6. Susanna. (Information from Charles Billson, Esq.)

² Parey.

³ J. Vaughan, *Cornhill*, 1909.

lected and an armorial window to his memory was put up in Buriton Church. Under the Goodyer arms is the inscription 'To the Glory of God and in Memory of John Goodyer of Alton, Mapledurham, Petersfield, 1592-1664, Royalist, Botanist, Founder of Goodyer Charity, Weston'. A water-colour drawing of the window by Mrs. Davis has been hung near his books in the Magdalen Library.

The property which he left for the benefit of the poor of Weston, has already been described in his will. The way in which his Trustees set out to administer the Trust is shown in their first Account Book which we had the pleasure of consulting through the kindness of Mr. Burley, the solicitor to the Trust. The first entries of receipts and payments give some idea of the initial value of the legacy.

A Booke of Receipt and Disbursements of the issues and profitts of the house and lands in Weston late Mr. John Goodyers, [*a blank space*] phisicke, And by his last Will and Testament settled upon two Trustees viz. Leonard Bilson Esq and Edmund Yalden, Clarke, to the end that they should nominate six able persons and Convey the said house and lands to them or ffeoffees in trust for the pro-lating and puttinge forth apprentices of y^e Children of the poorer sort of Inhabitants within the said Tithinge of Weston accordinge to the true intent and purport of the said last Will and Testament. In pursuance whereof they, the said Leonard Bilson and Edmund Yalden have nominated, and by their deed bearinge date the 11th day of June in the sixteenth yeare of the Raigne of King Charles the second, have Conveyed the said house & lands unto these six persons followinge viz. :

William Bilson, gent.	Henry Voake, yeoman.
John James, gent.	Jacob Voake, yeoman.
John Bold, gent.	Robert Cox, yeoman.

The first Receipts.

1664

12 Julii	Rec ^d of John Girdler for a bushell and an halfe of apples	00 02 3
	Rec ^d of Will ^m Budde for a parcell of faggotts	00 01 8
12 Aug.	Rec. of John Girdler for 3 halfe bushell one gallon of apples & one gallon of peares & 3 halfe bushells	00 6 3
23 Sept.	ditto ditto for apples	00 4 0

15 Oct.	Rec. of Mr. Jaques for halfe a yeares rent of halpenney lands ending at Michaelmas last past	7 10 0
1665		
14 Apr.	Rec. of Mr. Bettesworth for his halfe yeares rent ending att Lady day last past (abat- inge nine shillings for Chimney money before hee came to the house	4 11 0

The first Payments.

1664		
24 June	Imprimis to Will. Cox and Robert Tribe for mending the garden hedge	00 00 9
13 Julii	To Will. Cox and Robert Tribe for scour- inge the river	00 02 4
15 Oct.	Paid att the Trustees first meeting to lett the lands the expenses of the house in fire, beare, & tobacco	00 05 0
30 Dec.	Paid John Gamons Bill for repairinge the glasse windowes of the house	00 10 0
1666		
30 March	Paid Doctor Gunter for losse of time in cominge over for a witesse	00 10 0

His 'large house' was afterwards sold for over £1,000, the proceeds of which, invested in Consols, together with the rent of the land, bring in an annual income of some £75, which is a source of considerable benefit to the parish. Part of this sum is yearly expended in gifts of money and clothing, part in making allowances by way of encouragement to servant girls, and part in apprenticing the young lads of the tithing. The people of Buriton have, indeed, much cause for gratitude towards the good botanist of Petersfield, whose very name is now forgotten in the village.

His memorial in modern scientific botany is the customary one of a genus named in his honour. But unfortunately his name has been somewhat inappropriately connected with a rare Orchid, *Goodyera repens*, that is found in certain fir-woods in Cumberland and Scotland, but which could never have lived, where it was thought that he might have found it, in the 'moist meadow named Wood-mead, neere the path leading from Petersfield towards Beryton'. There Goodyer was acquainted with an orchid which he knew by

the name of 'Palma Christi radice repente' or 'Creeping Satyrion', and which we should now call the Marsh Helleborine or *Epipactis palustris*. Robert Brown, who gave the name *Goodyera* to the northern genus in honour of our Hampshire botanist, was misled by Johnson, who in error had attached a figure of *Goodyera repens* to the text relating to the Marsh Helleborine. It is somewhat unfortunate that by a triple error his name should have been attached to a plant which in all probability he could never have seen, but that is only one of the many chance circumstances which have led to the passing from memory of the services of 'an incomparable botanist, of sound judgement, and of immense industry'.

*Ocymoides sempervirens*

A Fern

DRAWINGS BY GOODYER

DESCRIPTIONS OF PLANTS BY JOHN GOODYER

THESE descriptions of plants are for the most part the earliest that are extant in the English language. They are now printed for the first time from Goodyer's original manuscripts, with others reprinted from passages contributed to the second or emaculate edition of Gerard's Herbal, printed in 1633.

In the following Table the extent of Goodyer's botanical labours is indicated by differences in the type.

TABLE OF NAMES WITH MODERN EQUIVALENTS, GROUPED IN NATURAL ORDERS.

Capital letters indicate plants of which descriptions are extant.

Small letters indicate plants of which no descriptions are extant.

Roman type indicates English plants.

Italic type indicates Foreign and Garden plants.

B, H, S, W, &c., denote Record or First Mention for Britain or for the Counties of Hants, Sussex, or Wilts., &c.

Gardens are denoted by the names of their owners.

Goodyer frequently quotes more than one name of a plant: in such cases we have only printed one for reasons of economy of space. Determinations for which I have had the advantage of the experience of Dr. Daydon Jackson, Mr. Britten, and Dr. Stapf are marked with J, B., and S. respectively.

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	PAGE
<i>Ranunculaceae.</i>			
Ranunculus Ficaria L.		Chelidonium minus.	115
„ Flammula L.		Ranunculus flammeus aquatilis angustifolius.	69
Adonis autumnalis L.		Adonis.	136
<i>NIGELLA DAMASCENA</i> L.		Nigella multiplex.	152
„ <i>HISPANICA</i> L. (prob.).	Spain	„ elegans.	153
<i>Papaveraceae.</i>			
<i>Papaver hybridum</i> L. (possibly).		Argemone Pavio.	155
Papaver Argemone, L.	Durford	S Argemone capitulo longiore.	178
„ <i>Rhoeas</i> , β <i>setigerum</i> Boenn. (J.)	Spain	Papaver <i>Rhoeas</i> Baeticum.	155
<i>HYPECOUM PROCUMBENS</i> L.	Coys	Hypecoon Clusii.	129
<i>Fumariaceae.</i>			
Corydalis claviculata DC.	Southsea	H Fumaria clavicularis donata.	47

DESCRIPTIONS OF PLANTS

101

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	<i>PAGE</i>
Cruciferae.			
Sisymbrium Alliaria Scop.	Droxford H	Alliaria recentiorum. 'Herbe John.'	111
„ Irio L.	Whitechapel	Erysimum ii Tab.	191
Nasturtium sylvestre R. Br.		Eruca palustris minor.	192
Cardamine impatiens L.	Bath	Cardamine impatiens.	
Dentaria bulbifera L.	Mayfield B	Dentaria bulbifera.	186
SINAPIS ALBA L. ?	Droxford H	Sinapi sativum alterum Penae.	111
Brassica Sinapistrum Boiss.		Cherlock.	162
Draba sp.		Draba lutea sil. long.	69
IBERIS UMBELLATA L. (J.)	[Crete]	Thlaspi umbellosum marinum fl. alb.	130
Bunias orientale L.	Whitechapel	Rapistrum aliud non bulbosum.	191
Violaceae.			
Viola tricolor L.		Viola tricolor sylv. parva. How.	194
Viola odorata L. fl. pl.	Sheet garden H	Viola martia purpurea multiplex.	109
Frankeniaceae.			
FRANKENIA LAEVIS L.	Haylinge B	An Polygoni marini species.	148
„ „	Bursledon Ferry	Polygonum alterum pusillo.	151
Caryophyllaceae.			
Silene fruticosa L. (J.)		Ocymoides sempervirens.	99
Lychnis dioica L.	Petersfield	'Lychnis sylv. flore carneo odorato' Merrett.	195
„ Githago Scop.	Chichester S	Pseudomelanthium.	112
SAGINA NODOSA Meyer.	Wellingborough B	Alsine palustris foliis tenuissimis. 'Saxifraga pal. alsin.'	179
„ APETALA L.	Bath	[No name.]	186
Stellaria aquatica Scop.		Alsine major repens.	69
Portulacaceae.			
Montia fontana L.	Hants H	'Alsine flosculus conniventibus' Merrett. 'Blinks' Goodyer.	195
Hypericineae.			
Hypericum Androsaemum L.	St. Vincent's Rocks	Androsaemum magnum.	77
Malvaceae.			
Malva moschata L.	Mapledurham H	Alcea vulgaris albo flore.	111
„ stipulacea Cav. ? (J.)	Coys. Spain	Malva flore amplo Baetica aestiva.	134
Linaceae.			
LINUM CATHARTICUM L.	Hants, Purfleet E	Linum silvestre catharticum	
	H	Milmountaine.	109, 112
Geraniaceae.			
Geranium columbinum L.	Hants, Whitechapel B	'Geranium columbinum' How.	191
„ lucidum L.	Guildford B	'Geranium saxatile' Park. 708.	185
ERODIUM GRUINUM Willd. ? (J.)	Coys. Spain	Geranium Baeticum sp. Boelii.	146
Leguminosae.			
ULEX EUROPAEUS L.		Ye great furze.	189
„ „ (whitish fl. var.) (S.)		Genista spinosa flore albo.	189
„ NANUS Forst. (S.)		Ye least furze (not Gen. sp. minor Park. 1003).	189
„ parviflorus. (S.)	Provence	Genista spinosa major brevibus aculeis B. P. 394.	190
?		Genista spinosa minor.	189
MEDICAGO MINIMA L. ? (J.)		Medica anglica minor.	141
MEDICAGO POLYMORPHA L.	Spain	Medica major Baetica sp. 1. spinulis intortis.	142

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	<i>PAGE</i>
<i>MEDICAGO INTERTEXTA</i> L.	Spain	<i>Medica major Baetica altera.</i>	142
„ <i>MARINA</i> L.		„ <i>marina spinosa</i> sp.	142
<i>PISUM SATIVUM</i> L.		<i>Pisum quadratum.</i>	139
<i>MELILOTUS INDICA</i> L.?	Coys	<i>Melilotus Indiae orientalis.</i>	126
<i>ASTRAGALUS HAMOSUS</i> L.	Coys	<i>Securidaca minor.</i>	130
<i>Trifolium ligusticum</i> Balb. ? (J.)	Coys	<i>Lagopus trifolius maior Baeticus.</i>	144
„ <i>Lagopus</i> L. ? (J.)		<i>Lagopus trifolius flore ruberrimo.</i>	144
<i>HEDYSARUM HUMILE</i> L.		<i>Hedysarum clypeatum.</i>	147
<i>Onobrychis sativa</i> Lam.	Langford, Wilts. w	<i>Caput Gallinaceum Belgarum</i>	178
<i>Hippocrepis comosa</i> L.	Buttersworth Hill St. Vincent's Rocks	<i>Ferrum equinum Germanicum.</i>	187
<i>ASTRAGALUS LUSITANICUS</i> Lam. (J.)	Park. Spain	<i>Astragalus marinus Lusitanicus Boelii.</i>	140
<i>VICIA FABA</i> L. var. (J.)	Coys. Spain	<i>Faba veterum serratis foliis Boelii.</i>	140
<i>Vicia tetrasperma</i> Moench.		H <i>Vicia sive Cracca minima.</i>	193
<i>Vicia sylvatica</i> L.	Bath	' <i>Vicia maxima sylvatica spicata Bathoniensis Goodyeri.</i> '	190, 196
<i>VICIA SATIVA</i> L. var.	Coys	<i>Vicia indica fructu albo.</i>	139
<i>LEUCOSPERMA</i> Moench. (J.)			
<i>VICIA SATIVA</i> β <i>LINEARIS</i> Lange? (J.)	Coys. Spain	<i>Aracus maior Baeticus Boelii.</i>	138
<i>Vicia lutea</i> β <i>laevigata</i> Boiss. (J.)	Coys. [Portugal]	<i>Legumen pallidum Vlissiponense Nonii Brandonii.</i>	139
<i>Ervum Lens</i> L.	Droxford	<i>Lens minor.</i>	111
<i>PISUM ARVENSE</i> L.	Park. Spain	<i>Pisum maculatum Boelii.</i>	141
<i>LATHYRUS OCHRUS</i> DC.	Park. Coys	<i>Ervilia silvestris Dodonaei.</i>	141
„ <i>ANNUUS</i> L. or	Spain	<i>Lathyrus aestivus flore luteo.</i>	136
<i>L. OCHRUS</i> DC. ? (J.)			
<i>LATHYRUS CLYMENUM</i> L. ? (J.)		<i>Lathyrus aestivus Baeticus fl. coeruleo Boelii.</i>	136
<i>LATHYRUS SPHAERICUS</i> Retz ? (J.)		<i>Lathyrus aestivus flore miniato.</i>	137
<i>LATHYRUS TUBEROSUS</i> L.	Spain	<i>Lathyrus aestivus dumetorum Baet. Boelii.</i>	138
„ <i>PALUSTRIS</i> L.	Spain	<i>Lathyrus palustris Lusitanicus Boelii.</i>	137
<i>Lathyrus silvestris</i> L.	Hants H	' <i>Lathyrus maior angustifol. fl. pall. rubro</i> ' Merrett.	195
<i>LENS ESCULENTA</i> Moench. ? (J.)		<i>Lathyrus aestivus Baeticus fl. albo Boelii.</i>	137
<i>ORNITHOPUS SCORPIOIDES</i> L.		<i>Scorpioides mathioli.</i>	131
<i>SCORPIURUS SUBVILLOSA</i> L. (J.)		<i>Scorpioides multiflorus Boelii.</i>	151
<i>SCORPIURUS VERMICULATA</i> L.		„ <i>siliqua crassa Boelii.</i>	151
„ <i>laevigata</i> L. (J.)		„ <i>repens Bupleuri folio.</i>	132
Cucurbitaceae.			
<i>CUCURBITA PEPO</i> L. var.		<i>Macocks Virginiani.</i>	165
<i>CITRULLUS VULGARIS</i> Schrad. (J.)		<i>Melones Aquatici.</i>	165
Rosaceae.			
<i>POTENTILLA COMARUM</i> Nutt.		<i>Quinquefolium palustre.</i>	170
<i>Potentilla Anserina</i> L.		<i>Heptaphyllon maius Phyto. 651</i>	155
<i>ALCHEMILLA ALPINA</i> L. ? (J.)		<i>Filipendula vulgaris.</i>	77
<i>Spiraea Filipendula</i> L.	St. Vincent's Rocks	<i>Rubus repens fructu caesio.</i>	114
<i>RUBUS CAESIUS</i> L.		H <i>Cloudberry.</i>	195
„ <i>Chamaemorus</i> L.	? Ingleborough	<i>Rosa cinamomea simpl. fl.</i>	112
<i>Rosa cinnamomea</i> L.	Droxford	' <i>Rosa silvestris odora Eglanteria</i> ' How MS.	
<i>Rosa Eglanteria</i> L.	Bath		

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	<i>PAGE</i>
<i>Rosa gallica</i> L. ¹	.	<i>Rosa holoserica.</i>	110
<i>Pyrus Aria</i> L.	'Sandrish in Kent' K	' <i>Aria Theophrasti</i> ' Merrett.	195
Onagrarieae.			
<i>Epilobium angustifolium</i> L.	Winchester	<i>Lysimachia forte.</i>	111
		<i>Chamaenerion Gesneri.</i>	111
<i>OEŃOTHERA</i> BIENNIS L.		<i>Lysimachia virginiana.</i>	159
LUDWIGIA PALUSTRIS Elliot.	New Forest	<i>Herba aquatica rubescens facie</i>	187
		<i>Anagall. Holosteum.</i>	193, 195
Crassulaceae.			
<i>Sedum rupestre</i> L. var. minus.	St. Vincent's Rocks	' <i>Sedum Divi Vincentii non-descr.</i> ' Merrett.	77, 195
„ sp.		<i>Aizoon.</i>	152
Saxifrageae.			
<i>Chrysosplenium oppositifolium</i> L.	Mapledurham	<i>Saxifraga aurea.</i>	185
<i>Parnassia palustris</i> L.	Wellingborough	<i>Gramen parnassi.</i>	180
Umbelliferae.			
<i>Eryngium maritimum</i> L.	Tichfield Bay	<i>Eryngium marinum.</i>	110
<i>Trinia glauca</i> O. K.	St. Vincent's Rocks	<i>Peucedanum pumilum.</i>	77
<i>Cicuta virosa</i> L.	Denham, Herts.	<i>Sium alterum olusatri facie.</i>	179
APIUM NODIFLORUM Reich.	Droxford Petersfield	{ <i>Sium repens.</i>	114
		{ <i>Sium umbellis ad caulium nodos</i> ' Merrett.	195
APIUM INUNDATUM Reich. b. f. (J.)		<i>Sium pusillum foliis variis.</i>	192
<i>Apium crispum</i> L.	Idsworth	<i>Apium crispum.</i>	172
<i>Aegopodium Podagraria</i> L.		<i>Podagra germanica.</i>	110
<i>Sison Amomum</i> L.		<i>Sium odoratum Tragi.</i>	121
CARUM SEGETUM Benth.		<i>Sium siifoliis. Honewort.</i>	53, 121
SIUM LATIFOLIUM L.	Oxford	<i>Pastinaca aquatica latifolia.</i>	176
SIUM ERECTUM Huds.	Droxford	„ „ minor.	116
<i>Bupleurum rotundifolium</i> L.		<i>Thorow-wax.</i>	
OENANTHE LACHENALII Gmel.	Soberton	<i>Oenanthe angustifolia</i> Lob.	115
„ SILAIFOLIA Bieb. (T.)		„ „	35
„ pimpinelloides L.		„ apii folio.	145
<i>Crithmum maritimum</i> L.	Hurst Castle	<i>Crithmum chrysanthemum</i> G.	193
<i>Peucedanum sativum</i> B. and H.			
<i>Caucalis arvensis</i> Huds.	Petersfield	<i>Caucalis pumila segetum</i> Merrett.	195
„ nodosa Scop.		<i>Caucalis nodosa echinato semi</i> Bauhini.	114
„ LATIFOLIA L. (S.)	Spain	<i>Caucalis major Baetica.</i>	128
<i>Daucus Carota</i> L.	Droxford	<i>Siser erraticum</i> Plinii.	112
<i>Smyrnum Olusatrum</i> L.		<i>Alexanders.</i>	
Caprifoliaceae.			
<i>Adoxa Moschatellina</i> L.	Bunnykens Holworte.	<i>Radix cava minima viridi flore.</i>	69
<i>Sambucus Ebulus</i> L.	Veny Sutton	<i>Ebulus.</i>	111
Rubiaceae.			
RUBIA PEREGRINA L.		<i>Rubia sylvestris.</i>	191
CRUCIANELLA sp.		„ spicata Cretica Clusii.	132
<i>Sherardia arvensis</i> L.		<i>Rubia minor flore rubro.</i>	148
ASPERULA CYNANCHICA L.	Hants	<i>Synanchica.</i>	113
Valerianeae.			
VALERIANA CORNUCOPIAE L.		<i>Valeriana mexicana.</i>	133
„ Locusta L.		<i>Lactuca agnina.</i>	133

¹ Probably. J. notes that the Moss Rose *R. muscosa* Ait. seems not to be catalogued before 1720 by Boerhaave.

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	PAGE
Dipsaceae.			
<i>Jasione montana</i> L.	Sheet H	<i>Scabiosa minima hirsuta</i> G.	
		585. <i>S. media.</i>	110, 164
<i>SCABIOSA ATROPURPUREA</i> L.		<i>Scabiosa flore rubro</i> = <i>S. sexta</i>	
		<i>indica</i> Clusii.	164
„ <i>columbaria</i> L.		<i>Scabiosa minor.</i>	164
Compositae.			
<i>Tussilago Farfara</i> L.			
<i>ERIGERON ACRE</i> L.	Winchester H	<i>Conyza coerulea</i> C. <i>odorata.</i>	162
<i>Jasonia tuberosa</i> DC.		<i>Aster conyzoides</i> Gesneri.	120
<i>ANTHEMIS TINCTORIA</i> L.		<i>Bupththalmum vulgare</i> i Mat-	
		thioli.	154
<i>Anthemis Cotula</i> L.		<i>Cotula foetida.</i>	135
„ <i>nobilis</i> L.		Camomile.	
<i>CHRYSANTHEMUM CORONA-</i>	[Crete]	<i>Chrysanthemum Creticum</i> i	
<i>RIUM</i> L.		Clusii.	134
<i>CHRYSANTHEMUM CORONA-</i>	Spain	<i>Chrysanthemum Baeticum</i>	
<i>RIUM</i> L.		Boellii <i>inscr.</i>	135
<i>CHRYSANTHEMUM CORONA-</i>	Spain	<i>Chrysanthemum tenuifolium</i>	
<i>RIUM</i> L.		Baeticum Boellii.	135
<i>Chrysanthemum segetum</i> L.		<i>Chrys. segetum.</i>	135, 155
<i>HELIANTHUS TUBEROSUS</i> L.		<i>Flos solis pyramidalis.</i>	109, 166
„ <i>annuus</i> L.		„ „ <i>Peruanus.</i>	167
<i>Senecio Doria</i> L.		<i>Herba Doria.</i>	119
<i>SENECIO SARRACENICUS</i> L. (S. J.) Coys.		<i>Herba Doria altera.</i>	119
<i>Senecio paludosus</i> L.	Downham Fen B	<i>Conyza aquatica laciniata.</i>	193
„ <i>Jacobaea</i> L.	Ladle Hill H	<i>Jacobaea angustifolia</i> Pan-	
		nonica 2 Clus.	194
<i>XERANTHEMUM ANNUUM</i> L. (S.)		<i>Parnica imperati.</i>	119
<i>ACHILLEA NOBILIS</i> L.		<i>Achillea sideritis.</i>	155
<i>Achillea Millefolium</i> L.		Common Yarrow.	155
<i>DIOTIS MARITIMA</i> Cass.	Hayling H	<i>Gnaphalium marinum.</i>	148
<i>Gnaphalium margaritaceum</i> L.		<i>Gnaphalium Americanum.</i>	148
[<i>Filago minima</i> Pers. ¹	Petersfield H	' <i>Filago minor</i> Lob.'	
[<i>Xanthium Strumarium</i> L.	Southwick St., Hants ? B	<i>Zanthium.</i>	128
<i>Arctium minus</i> Bernh.?			
„ <i>Lappa</i> L.	Mangerfield, Glos. G	<i>Arctium montanum et Lappa</i>	
		<i>minor</i> Galeni.	194
<i>NOTOBASIS SYRIACA</i> Cass. Coys. Spain		<i>Silibum minus flore nutante</i>	
		Boellii.	145
<i>CNICUS PRATENSIS</i> Willd.		<i>Carduus bulbosus</i> Monspel-	
		liensium.	144, 154
<i>CNICUS ERIOPHORUS</i> Roth.	Holyborne H	<i>Carduus eriocephalus.</i>	109, 146
<i>CARDUUS ACAULIS</i> L.	Hants: Purfleet B	<i>Carduus acaulis septentriona-</i>	
		<i>lium</i> L'Obelii.	118, 145
<i>Carduus crispus</i> L.			
„ <i>nutans</i> L.	[Hants] H	<i>Carduus viarum fl. purp. et alb.</i>	112
<i>Carlina vulgaris</i> L.	Hants: Purfleet HE	<i>Carlina sylvestris.</i>	112
<i>CARLINA LANATA</i> L.	Coys. [Naples]	<i>Acarna flore rubro.</i>	145
<i>CENTAUREA SALMAN-</i>	Montp. Spain	<i>Stoebe Salmantica</i> i Clusii	
<i>TICA</i> L.		<i>foliis Cichorei.</i>	154
<i>Centaurea Scabiosa</i> L.	Chawton H	<i>Jacea albo flore.</i>	110
„ „		„ <i>major.</i>	165
<i>Sonchus tingitanus</i> L.?		<i>Sonchus Africanus</i> Boellii.	69
<i>LACTUCA VIROSA</i> L.	Southampton H	<i>Lactuca sylvestris vera ingrato</i>	
		<i>odore.</i>	111, 158

¹ *Filago minor* is included in How's list, p. 280. He may have obtained the locality, Petersfield, from Goodyer.

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	<i>PAGE</i>
<i>LACTUCA AGRESTIS</i> L.		Lactuca agrestis.	159
<i>RHAGADIOLUS EDULIS</i> Gaertn. (J.)	[Montpelier]	{ Hieratium Narbonense falcata siliqua L'Obelii.	148
<i>TOLPIS BARBATA</i> Gaertn. (J.)	Spain	Hieratium medio nigrum flore maior Boelii.	149
„ <i>UMBELLATA</i> β <i>MINOR</i> Lange (J.)	Spain	Hieratium medio nigrum flore minore Boelii.	149
<i>HIERACIUM INTYBACEUM</i> L.?	[Italy]	Hieratium intybaseum.	149
„ <i>ANDRYA-LOIDES</i> Vill. (J.)	Coys. Spain	„ lanosum.	150
Hieracium murorum L.	Godalming	S Pulmonaria Gallica sive aurea latifolia.	180
„ Pilosella L.		Pilosella repens.	150
Taraxacum officinale Willd.		Dens leonis vulg.	115
Cichorium Intybus L.		'Wild Cicorie.'	150
Lapsana communis L.		Lampsana.	149
<i>LAPSANA ZACINTHA</i> L.?	[Italy: Florence]	Cichorium verrucatum.	151
<i>CENTAUREA NEVADENSIS</i> & Reut. ? (J.)	Bois. Coys	Jacea capitulis hirsutis Boelii.	164
?		„ palustris Baetica Boelii.	164
<i>Primulaceae.</i>			
<i>ANDROSACE MAXIMA</i> L. (J.)		Androsace altera Matthioli Ger. p. 425.	150
<i>Primula veris</i> L.	Sheet	Cowslip 2 in a hose.	109
„ <i>vulgaris</i> Huds.	Sheet	Primrose 2 in a hose.	109
Samolus Valerandi L.	Emsworth	B Anagallis aquatica tertia.	111
Lysimachia nemorum L.		„ flore luteo.	187
<i>Campanulaceae.</i>			
<i>PHYTEUMA ORBICULARE</i> L. (S.)	Maple-durham	B [Rapunculus corniculatus montanus?]	185, 196
„ „	Droxford	B Rapunculum silvestre.	111, 116
<i>Campanula patula</i> L.		'Rapunculus sylvestris fl. rubro albescente' Merrett.	195
<i>CAMPANULA PUMILA</i> L. var.?		Lesser Bell-flower.	132
<i>RESEDA PHYTEUMA</i> L. (J.)		Phyteuma monspeliensium.	134
<i>Reseda Luteola</i> L.		Luteola.	134
<i>Ericaceae.</i>			
<i>MONOTROPA HYPOPITYS</i> L.	Droxford	B Orobanche verbasculi odore.	122
<i>GENTIANA PNEUMONANTHE</i> L.		B Pneumonanthe.	170
<i>Convolvulaceae.</i>			
<i>Convolvulus Soldanella</i> L.	Hants Chichester	H [Soldanella marina].	185
<i>CONVOLVULUS PURPUREUS</i> L. or some var. of it (J.)		S Convolvulus coeruleus Bryoniae nigrae folio flos Noctis non script.	153
„ „	Boel. Coys	Convolvulus coeruleus minor Baeticus.	129
„ <i>ARVENSIS</i> L.		Convolvulus minor.	129
<i>Cuscuta Epithymum</i> L.		H Cuscuta.	112
<i>Asclepiadae.</i>			
<i>ASCLEPIAS PURPURASCENS</i> L. (S.)		Periploca recta virginiana.	131
<i>Boragineae.</i>			
<i>ANCHUSA ANGUSTIFOLIA</i> L.		Buglossum scorpioides an Echii facie Buglossum minimum flore rubente (Lobel).	131
<i>PULMONARIA ANGUSTIFOLIA</i> L.	New Forest	B Pulmonaria foliis Echii.	115, 190

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	PAGE
Solanaceae.			
<i>Hyoscyamus niger</i> L.		<i>Hyoscyamus luteus.</i>	122, 160
<i>NICOTIANA TABACUM</i> L.	Coys	<i>Petum indicum folio pene obtuso.</i>	160
var. <i>BRASILIENSIS</i> Comes. (S.)			
<i>NICOTIANA TABACUM</i> L.	Uvedale	<i>Petum indicum folio Hydro-lapathi acuto.</i>	161
var. <i>FRUTICOSA</i> Hook. f. (S.)			
Orobanchaceae.			
LATHRAEA SQUAMARIA L.		{ <i>Anblatum</i> Dod.	123
		{ <i>Dentaria maior sive αφύλλος</i> Clus.	
Orobanche Purpurea Jacq.?		Orobanche.	122
Scrophulariaceae.			
<i>Verbascum nigrum</i> L.	? Droxford	H <i>Blattaria flo. luteo.</i>	
LINARIA MINOR Desf.		<i>Antirrhinum minus.</i>	115
LINARIA THYMIFOLIA DC. ? (J.)	Coys	<i>Antirrhinum minus flore Linariae luteo inscriptum.</i>	143
LINARIA CYMBALARIA Mill.	Coys.	B <i>Cymbalaria Italica.</i>	17, 163
	Droxford	H	
LINARIA SERPYLLIFOLIA Lange ? (J.)	Coys	<i>Linaria minor aestiva.</i>	143
<i>Linaria Elatine</i> Desf.		Fluellin. <i>Elatine.</i>	163
[<i>Scrophularia vernalis</i> L.	Coys		17
<i>Scrophularia nodosa</i> L.		<i>Common Scrophularia.</i>	156
DIGITALIS FERRUGINEA L.		<i>Digitalis ferruginea.</i>	186
<i>Veronica hybrida</i> L.	St. Vincent's Rocks	B 'Veronica mas recta' Merrett.	76
BARTSIA ODONTITES Huds.		<i>Eufrasia altera Dodo.</i>	117
" " var. alb.	Bellmere	H <i>Euphrasia 2 Dod. flo. albo.</i>	150
MELAMPYRUM SYLVATICUM L.		B <i>Melampyrum luteum latefolium.</i>	118
<i>Pedicularis sylvatica</i> L.	Warwickshire	w 'Pedicularis fl. albo' How MS.	
? CERINTHE MAJOR L.	[Prob. Spain]	<i>Cerinthe flore rubro.</i>	128
CERINTHE MINOR L.	Spain	" minor flore albo veris luteis.	180
" " major L. var. flavo flore.		<i>Yeallow flowered Cerinthe.</i>	128
Labiatae.			
SALVIA VERTICILLATA L.	Coys	<i>Horminum silvestre iii Clusii.</i>	156
<i>Nepeta nuda</i> L.	Droxford	<i>Menthastrum montanum.</i>	111
<i>Origanum Majorana</i> L.	Yalden, Sheet	<i>Sweete Marjoram.</i>	166
OCIMUM BASILICUM L.	" "	<i>Acinos [odoratissimum].</i>	166
<i>Nepeta Cataria</i> L. var. ? (J.)	Parkinson	<i>Nepeta media.</i>	162
NEPETA TUBEROSA L.	Coys	<i>Cattaria tuberosa radice Baetica Boelii non script.</i>	161
<i>Lavandula officinalis</i> L.		<i>Lavender.</i>	157
STACHYS GERMANICA L.	Witney	B <i>Stachys (Buckner).</i>	59
" " "		<i>Stachys Wild Horehound.</i>	177
" " <i>Betonica</i> Benth.		<i>Betony.</i>	120
<i>Lamium Orvala</i> L.	Coys	<i>Lamium Pannonicum 2^{um} exoticum Clusii.</i>	156
<i>Thymus Serpyllum</i> L.	Petersfield	H <i>Serpillum. 'Serpillum foetidum Goodyeri' How MS.</i>	195
<i>Galeopsis Tetrahit</i> L. var. bifida.		H 'Cannabis spuria altera flo. purp.' How.	194
Plantagineae.			
Littorella lacustris L.		B 'Holosteum Junciifolium repens Goodyeri' How MS.	
<i>Plantago Psyllium</i> L.		<i>Psyllium.</i>	157
<i>Plantago Coronopus</i> L.		<i>Cornu cervinum Lobelii.</i>	130, 155

DESCRIPTIONS OF PLANTS

107

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	PAGE
<i>Illecebraceae.</i>			
<i>Scleranthus annuus</i> L. Knawel. (B.)	Tichfield Bay	H Polygonum germanis.	110
<i>Chenopodiaceae.</i>			
SUAEDA FRUTICOSA Forsk.		H Chamaepitys vermiculata.	179
SALICORNIA HERBACEA L.	? Portland	H Kali album, p. 81, Dodo.	152
EMEX SPINOSA Campd. (J.)		Blitum spinosum Creticum.	136
Rumex Acetosa L.		Acetosa maxima.	194
<i>Santalaceae.</i>			
THESIUM HUMIFUSUM DC.	Droxford	B Anthyllis montana, Linaria adulterina.	117
<i>Urticaceae.</i>			
Urtica dioica L.		Nettle.	156
Humulus Lupulus L.		Hop.	114
<i>Ulmaceae.</i>			
ULMUS CAMPESTRIS Sm.		Ulmus vulg. folio lato scabro.	38
„ MONTANA Stokes.		„ folio latissimo scabro.	41
„ GLABRA Miller.	Stubbers	B „ folio glabro.	43
„ MINOR Miller.	New Forest	B „ minor f. angusto scabro.	39
<i>Amentaceae.</i>			
ALNUS GLUTINOSA Gaertn.		Alder.	175
BETULA ALBA L.		Unnamed.	175
Fagus sylvatica L.	Faringdon	H 'Great antient Beeches.	189
„ „ L. var.		Fagus.	188
QUERCUS ROBUR L.		Quercus. Cachryes and Galls.	172
„ „ agg.	Bramshaw, Wilts.	W 'Quercus serotina, procerior. 'Dor-Oak' Merrett.	195
JUGLANS REGIA L.	Longwood	Nux Juglans.	112, 174
CASTANEA SATIVA Mill.		Cachrys Castaneae.	174
Salix viminalis L.		Salix aquatica.	77
<i>Monocotyledons.</i>			
ACORUS CALAMUS L.		? B Acorus legitimum Clus. 231.	177
POTAMOGETON CRISPUS L.	Durford	B Tribulus aquaticus minor floribus uvae.	123
„ DENSUS L.	Droxford	B Tribulus aquaticus minor muscatellae floribus.	126
Butomus umbellatus L.		Butomus.	177
Alisma Plantago L.			
Damasonium stellatum Pers.	Hounslow Heath	B Plantago aquatica stellata.	110
	Between Sandie Chappell and Kingston	Plantago aquatica stellata.	180
Neottia Nidus-avis Rich.		Nidus avis.	127, 195
Epipactis palustris Crantz.	Petersfield	B Palma Christi, radice repente.	184
EPIPACTIS VIOLACEA Bor.	Holiborne	B Nidus avis flore et caule violaceo.	47, 126
TAMUS COMMUNIS L.	Hants	H Bryonia nigra.	127, 153
Allium ursinum L.		Ramsons.	
Colchicum autumnale L.	Warminster	W Colchicum flo. albo e: purpur.	110
Juncus bufonius L.		Gramen holosteum Alp. min.	190
Paris quadrifolia.	Chawton	H Herba Paris.	110
SCIRPUS SYLVATICUS L.	Oxford	B Cyperus gramineus Lobelii.	175
Carex vulpina L. or C. Pseudo Cyperus ?(J.).		'Gramen palustre Cyperoides Lob. Ger. Great Cyperus Grasse' How.	194
Carex pulicaris L.		? O Flea-grass. Ray, <i>Synopsis</i> .	196
Hordeum sylvaticum Huds.	Petersfield	B 'Gramen secalinum maximum' Merrett.	196
Panicum sanguinale L.	Petersfield	'Gramen paniceum procumbens' Merrett.	196

<i>Modern Name.</i>	<i>Locality.</i>	<i>Goodyer's Name.</i>	<i>PAGE</i>
CALAMAGROSTIS EPIGEJOS Roth. ? (<i>not Festuca Myurus</i> L.) (S.)	Winchester	B Calamagrostis. B Gramen inurorum spica longis- sima.	172 171, 190 195
ECHINOCHLOA CRUS-GALLI L. (S.)	Petersfield	'Gramen Paniceum' Merrett.	195
<i>PHALARIS CANARIENSIS</i> L. (S.)	Sheet Spain	Panicum sylvestre. Phalaris minor Baetica Boelii, sem. nigro.	120 133
„ <i>MINOR</i> L. (S.)	Spain	Phalaris minor Baetica Boelii, sem. albo.	133
„ <i>BULBOSA</i> L. (S.)	Spain	Phalaris bulbosa Boelii.	133
CYNOSURUSECHINATUS L. Coys. Sheppey		Gramen cristatum Baet. Boelii.	157
Nardus stricta L.		Spartum or Matweed.	171
BRIZA MAXIMA L.	Coys. Spain	Gramen tremulum maximum.	158
Phragmites communis Trim.		Arundo vallisoria.	176
<i>Coniferae.</i>			
JUNIPERUS COMMUNIS L. ♂	Surrey	S Juniperus sterilis.	123, 195
TAXUS BACCATA L.		Taxus glandifera baccifera.	168
„ „ ♂		„ tantum florens.	169, 196
<i>Filices, etc.</i>			
Ophioglossum vulgatum L.			
Botrychium Lunaria Sw.	Droxford	H Lunaria minor.	110
POLYPODIUM DRYOPTERIS L.	Hants	B Dryopteris Tragi.	189
ASPIDIUM ACULEATUM Sw. var. β LOBATUM.	Mapledurham	H Filix mas non ramosa pinnulis latis auriculis spinosis.	183
DRYOPTERIS THELY- PTERIS Sw.	nr. Petersfield	B Dryopteris Penae et Lobelii.	183
ASPIDIUM FILIX-MAS Sw.	Hants	H Filix mas non ramosa pinnulis latis.	182
„ „ var. AFFINIS.	Hants	H Filix mas non ramosa pinnulis angustis.	183
„ DILATATUM Sm.	Durford ; Mapledurham	B Filix ramosa pinnulis dentatis.	181
Asplenium Trichomanes L.	Wolmer Forest	H	179
„ Ruta-muraria L.		?H Ruta muraria	
Scolopendrium vulgare L. var. multifida	Swaneling	H Phyllitis multifida.	185
Ceterach officinarum Willd.		?H Ceterach	230
Lycopodium clavatum L.	Petersfield	H 'Chamaepeuce foemina seu polyspermos' Merrett.	196
Pilularia globulifera L.	Petersfield	H Gramen piperinum' Merrett.	196
Clathrus cancellatus L. (J.)	Petersfield	H Fungus corallinus' Merrett.	196

The manuscript descriptions are wholly in Goodyer's handwriting: the colour of the ink shows that in several cases the whole of a paragraph was not written at the same time. The concluding sentences, usually mentioning locality and date, were sometimes added later in a browner ink. Such sentences are indicated by the use of the mark ||.

These additions must have been made before 1632, when many of the descriptions were handed to Johnson for his revision of the Herbal. The methodical Goodyer kept a list of descriptions thus lent to Johnson (MS. 11,

f. 134). In all more than two hundred and fifty plants are specially noticed in addition to some hundred which are casually mentioned in his descriptive writings. About a hundred and fifty of Goodyer's descriptions of plants are still extant: sixty were printed by Johnson, ninety are in manuscript. Johnson did not print all the descriptions sent him.

In cases in which both printed and manuscript versions are available, we have followed Goodyer's own manuscript in essential particulars, but have retained the spelling of the printed version, except of the word 'flower', in which case we have adopted Goodyer's spelling.

It may be noted that Goodyer's spelling of English words, though perhaps more uniform than that of many of his contemporaries, was apt to vary. On the whole he favoured such forms as color, devide, flower, yealowe, coople, apece, fower, toppe, ioynt, bignes, ymediatlie, ynch, and a final *e* at the ends of present participles and other words.

The manuscript referred to is Goodyer MS. 11.

Woolly-headed Thistle. *Cnicus eriophorus* L.

Cardus eriocephalus. Corona fratrum quorundam. 1617

[See 29 June 1621 and 13 Aug. 1621.]

Linum silvestre catharticum. Mil-mountaine. Oct. 1617

[See 2 July 1619.]

Jerusalem Artichoke. *Helianthus tuberosus* L.

Heliotropium Indicum vel Virginianum. 25 March 1617

You had lately planted it when I was at your howse. 25 Martii 1617.—*MS.* ff. 48 v., 54.

[Refers to a visit to Coys. See p. 24 and under 17 Oct. 1621.]

Cowslipps 2-in-a-hose. 1617

Double violet. *Viola odorata* L., fl. plen.

Viola martia purpurea multiplex. 9 Apr. 1618

At Sheet.—*MS.* f. 58 v.

Primula veris L. var. and *P. vulgaris* Huds. var.

Primula veris flore gemino. 9 Apr. 1618

Cowslipps 2 in a hose at Sheet.

Primrose 2 in a hose at Sheet.—*MS.* f. 56 v.

- Ash weed. *Ægopodium Podagraria* L.
 Podagra germanica. Lo. 700; Herba Gerardi, 848. 11 Apr. 1618
 Lungwort, good wife hewes.—*MS.* f. 56 v.
- Moonwort. *Botrychium Lunaria* Sw.
 Lunaria minor. 21 Maij & 1 Junii 1618
 I found it in Droxford in a wood by Strugnells in the
 Thetcher.—*MS.* f. 55.
- Velvet Rose. *Rosa gallica* L.
 Rosa holoserica. G. 1085; Lo. o. 207. 3^o Junii 1618
 Ch: Edwards.—*MS.* f. 57.
- Herb Paris. *Paris quadrifolia* L.
 Herba Paris. Lo. 267; G. 328. 10 Junii 1618
 I sawe some with 5 leaves and some with 6 leaves at Chawton.—
MS. f. 54.
 [This is the first notice for Hants, but the discovery of 5- and
 6-leaved forms had been already made by Sir John Salusbury in
 North Wales in 1606.]
- Sheep's bit Scabious. *Fasione montana* L.
 Scabiosa minima hirsuta. G. 585. 7 Julii 1618
 At Sheete.—*MS.* f. 57 v.
- Greater Knapweed. *Centaurea Scabiosa* L.
 Jacea albo flore. G. 589. 18 Julii 1618
 At Chawlton.—*MS.* f. 54 v.
- Sea Holly. *Eryngium maritimum* L.
 Eryngium marinum. G. 999. 20 Julii 1618
 One plant at Tichfield Bay.—*MS.* f. 53 v.
- Knawel. *Scleranthus annuus* L.
 Polygonum germanis. Trag. p. 393. 20 Julii 1618
 In a barren rye feild below Tichfield Bay & is y^e Parsley pert
 of Ger. p. 453.—*MS.* f. 56 v.
- Water Plantain. *Damasonium stellatum* Pers.
 Plantago aquatica stellata. Phitopinax 355; Lo. 301; Deles. 1058.
 30 Julii 1618
 In Hounslowe Heath.—*MS.* f. 56 v.
 [First record for Britain (see 2 July 1633). In 1723 the Duke of
 Argyle took in a large part of the Heath and planted it with a large
 collection of trees and shrubs from the Northern Colonies.]
- Autumn Crocus. *Colchicum autumnale* L.
 Colchicum flo. albo et purpureo. 21 Aug. 1618
 At Warminster in flo: 21 August 1618.—*MS.* f. 59.

? Willow-herb.

Lysimachia forte. 27 Augusti 1618
Wiltshere.—*MS.* f. 55.

Danewort. *Sambucus Ebulus* L.

Ebulus. Lo. o. 164; G. 1238. 27 Augusti 1618

At Venny Sutton in Wiltes: it is called Scotts blood.—*MS.* f. 53.
[‘They call it Danes weede in Suffolk,’ *Bullein* 1562.]

Brookweed. *Samolus Valerandi* L.

Anagallis aquatica terciā. Lo. 467. 1618

By a mill at Emsworth.—*MS.* f. 51.

Willow-herb. *Epilobium angustifolium* L.

Chamaenerion Gesneri. Lo. 343; G. 386. 1618

It is called Willowe at Winchester.—*MS.* f. 52 v.

White Mustard. *Sinapis alba* L.

Sinapi alterum sativum. Lo. ps. 2: pag. 277. 1618

White pepper: Drox.—*MS.* f. 54 v.

Lactuca virosa L.

Lactuca silvestris vera ingrato odore. 1618

[See 13 Sept. 1621.]

Musk Mallow. *Malva moschata* L.

Alcea vulgaris albo flore. [1618]

At Mapledurham.—*MS.* f. 51.

Malva verbenacea. n. d.

‘Mr. Goodyer found the Vervain Mallow with white flowers growing plentifully in a close neere Maple-Durham in Hampshire, called Aldercrofts.’—*Ger. emac.* 931.

[The first record for Hants.]

Jack-by-the-Hedge. *Sisymbrium Alliaria* Scop.

Alliaria recentiorum. Lo. 530; G. 560. [1618]

At Droxford. herbe John.—*MS.* f. 51.

Lentil. *Ervum Lens* L.

Lens minor. Lo. o. 74; G. 1049. [1618]

Droxford.—*MS.* f. 55.

Nepeta nuda L.

Menthastrum montanum. [1618]

Droxford in y^e stone wall.—*MS.* f. 55 v.

Phyteuma orbiculare L.

Rapunculum silvestre. Tragi, p. 726. [1618]

Droxford.—*MS.* f. 57.

Cinnamon Rose. *Rosa cinnamomea* L.
Rosa cinamomea simpliciflora. G. 1086. [1618]
 Droxford.—*MS.* f. 57.

Wild Carrot. *Daucus Carota* L.
Sisyrinchium erraticum Plinii. [1618]
 Droxford.—*MS.* f. 57 v.

Black Mullein. *Verbascum nigrum* L.
Blattaria fl. luteo. G. 633; Lo. 565. [1618]
 I found this wild.—*MS.* f. 47 v.

Musk Thistle. *Carduus nutans* L.
Carduus viarum fl. purpureo. G. 1011. } [1618]
 „ *viarum fl. albo*. G. 1011. }
 We have a kind here that smells like muske.—*MS.* f. 47 v.

Carlina vulgaris L.
Carlina sylvestris. Lo. 14; G. 997; Math. 497. [1618]
 Clusius. p. clvi, hath ye figure of *Carduus vulgariss. viarum*, Lo. o.
 20, for *Carlina sylvestris*. || It growes on our Chalke Downes &
 also at Purflet.—*MS.* ff. 47 v., 52.
 [See 8 July 1620.]

Dodder. *Cuscuta Epithimum* L.
Cuscuta. Lo. 427; G. 462. [1617-18]
 I have seene it on furse, heath, nettles.—*MS.* f. 48.
 „ „ on furse, heath, nettles, fatches.—*MS.* f. 53.

Walnut. *Juglans regia* L.
Nux Juglans. Lo. o. 108; G. 1252. [1618]
 Forked at y^e toppe: at Longwood. Ex relat. Daniel Waite.—
MS. f. 55 v.

[Longwood, in the parish of Owslebury, was in the possession of
 Richard Garth, who died seised of it in 1597.]

Corn Cockle. *Lychnis Githago* Scop.
Pseudomelanthium. G. 926; Lo. 38. [1618]
 Crape. In Sussex about Cheichester.—*MS.* f. 56 v.
 [The name *Crap* was formerly given to various weeds growing
 among corn. Withering applied it to Rye Grass and Buckwheat.]

Purging Flax. *Linum catharticum* L.
Linum silvestre catharticum. Mil-mountaine. 2 July 1619

It riseth up from a small white threddy crooked root, sometime
 with one, but most commonly with five or six or more round stalks,
 about a foot or nine inches high, of a browne or reddish color,

every stalk dividing it selfe neere the top, or from the middle upward into many parts or branches of a greener colour than the lower part of the stalke: the leaves are small, smooth, of colour green, of the bignes of Lentill leaves, and have in the middle one rib or sinew, and no more that may bee perceived, & grow alongst the stalke in very good order by couples, one opposite against the other: at the tops of the small branches grow the flowers, of a white colour, consisting of five small leaves apiece, the nailes whereof are yellow: in the inside are placed small short chives also of a yellow colour, after which come up little knobs or buttons, the top whereof when the seede is ripe divideth it selfe into five parts; wherein is contained small, smooth, flat, slippery, yellow seed: when the seed is ripe the herbe perisheth: the whole herbe is of a bitter taste, and herby smell. It groweth plentifully in the unmanured inclosures of Hampshire, on chalkie downs, & on Purfleet hils in Essex, and in many other places. It riseth forth of the ground at the beginning of the Spring, and flowereth all the sommer.—*Ger. emac.* 559.

[For the rest of Goodyer's description see p. 21.]

Squinancy wort. *Asperula Cynanchica* L.

Synanchica.

3 Aug. 1619

This herbe groweth in y^e inclosures of Hampshire in drie Chalkie grounds. The root is crooked, blackish without, yellow underneath the skinne, white within that and wooddie; about five or six inches long, with many hairy strings: from the root arise many foure-square branches trailing upon the ground, sometimes reddish towards the root: the leaves are small and sharpe pointed, like [those of] *Gallium*, and grow along the stalke, on certaine knees or ioints, foure or 5 together, sometimes fewer: from those knees the stalk divideth it selfe towards the toppe into many parts, whereon grow many flowers, each flower having foure leaves, sometimes white, sometimes of a flesh colour, and every leafe of these flesh coloured leaves is artificially straked in the middle, and neere the sides with three lines of a deeper red, of no pleasant smell: after which commeth the seed something round, growing two together like stones. It flowereth all the sommer.—*MS.* f. 81; *Ger. emac.* 1120.

[In the printed version the date is given as 13 August, and either Goodyer or his editor has added an account of the *Vertue*. 'It dries without biting, and it is excellent against squinancies, either taken inwardly or applied outwardly, for which cause they have called it *Synanchica*, *Hist. Lugd.*']

Water Parsnip. *Apium nodiflorum* Reichb. f.
and *Caucalis nodosa* Scop.

Sium repens.

27 Aug. 1619

Hath longe plaine and smoothe leaves, made and fashioned like the leaves of the Ash havinge comonlie 4 or 5 small leaves growinge on ech side of the midle ribbe directlie one against another, snipt about the edges, amongst which come forth a round chamfered or furrowed hollowe stalk, of the bignes of a thumbe, browne or reddish neare the root, 4 or 5 foot longe, devided at the ioyntes or knees into many p[ar]tes or braunches ymediatlie from the root even to the toppe whereon growe the leaves without order like the former but shorter, nowe and then of a browne colour both above and underneath, of a stronge smell as is the whole plant, the flowers are white and many in number consisting of 5 sharpe pointed leaves apeece, growinge at the ioynts or devidinge of the stalkes, on short stems umbell fashion, after the manner of *Caucalis nodosa echinato semine Bauhini*. The stalks creape or run on the water, river or diches bankes, and hamper or matt them selves fast together, (contrarie to both those before mentioned, which growe upright and beare their flowers and seed at the topps of their stalks and branches). The flowers past, there appeareth the seed, two ioyned together at the first greene when it is ripe of a browne colour like to parsley seed, of a stronge tast but not plesant. The root is compacted of white threddie stringes infinite in number for the most parte as small as haire growinge or creepinge at the bottome of the water within the mire or marish ground, wherby it infinitely increaseth. The leaves of this plant growe greene in or above the water all the yere winter and somer. This groweth plentifullic by the lakes and rivers sides at Droxford in Hampshire.—*MS.* f. 82.

Dewberry. *Rubus caesius* L.

Rubus repens fructu caesio.

6 Sept. 1619

This hath a round stalke set full of small crooked and very sharpe pricking thornes, and creepeth on hedges and low bushes of a great length, on the upper side of a light red colour, and underneath greene, and taketh root with the tops of the trailing branches, whereby it doth mightily encrease: the leaves grow without order, composed of three leaves, and sometimes of five, or else the two lower leaves are divided into two parts, as Hop leaves are now and then, of a light greene colour both above and underneath. The flowers grow on the tops of the branches, *raccematim*, many together, sometimes white, sometimes of a very light purple colour.

every flower containing five leaves, which are crompted or wrinkled, and do not grow plaine: the fruit followes, first green, and afterwards blew, everie berry composed of one or two graines, seldome above foure or five growing together, about the bignesse of corans; wherein is contained a stony hard kernell or seed, and a iuyce of the colour of Claret wine, contrarie to the common *Rubus* or Bramble, whose leaves are white underneath: the berries being ripe are of a shining blacke colour, and every berry contains usually above forty graines closely compacted and thrust together. The root is woody and lasting. This growes common enough in most places, and too common in ploughed fields.—*Ger. emac.* 1271-2.

Lesser Celandine. *Ranunculus Ficaria* L.

Chelidonium minus, Lo. 593, *Ger.* 669. 10 March 1620

Inc. fl. [flowers open] 10 Marcii 1620.—*MS.* f. 52 v.

Dandelion. *Taraxacum officinale* Willd.

Dens leonis vulgi, Lo. 232, *Ger.* 228. 10 March 1620

Inc. fl. [flowers open] 10 Marcii 1620.—*MS.* f. 53.

Marsh Parsley Dropwort. *Oenanthe Lachenalii*
Gmel.

Oenanthe angustifolia Lob. Obs. p. 420. 19 May 1620

[Identified by Druce as the first British Record.]

This 19 of May 1620 I found this wild in East Hoo in ye parish of Subberton about 7 miles from Petersfield in Hampshire in a hedgerowe about a flightshott from ye then dwelling house of Mr. William Browne on ye south part of ye said house and ye 18 of June 1620 I saw it there in flower.—G. quoted by How in *MS.* note to *Phytologia*, p. 81.

[Merrett, *Pinax*, p. 84, gives the locality as East How. Druce gives the date as 28 June, but the figure is more like 18.]

Lungwort. *Pulmonaria angustifolia* L.

Pulmonaria foliis Echii. 25 May 1620

Found, May 25 Anno 1620 flowering in a Wood by Holbury House in the New Forest in Hampshire.—*Ger. emac.* 809.

Linaria minor Desf.

Antirrhinum minus. 20 Junii 1620

The stalks are small, round, hairy & branched, about 4 or 5 ynches high, ye leaves are small, smooth, blunt topped & like to Hissope leaves, y^e flowers are small fashioned like ye greate Antirrhinum, ye upper leaves whereof are of an ill favored purple color, & ye under leaves somewhat whitish, haveinge a very little

taile like to larkes spur, y^e seed followeth containd in a husk which is somethinge longc. The root is small whitish & threddy.—*MS. f. 83.*

Sium erectum Huds.

Pastinaca aquatica minor. *Sium odoratum* Tragi. 2 Julij 1620
figura in hist. lug: p. 701. *Apium palustre* Fuchsii.

The leaves and footstalkes that growe in the water by them selves distant from the stalk are about 2 foot longe, reddish, spongie, smooth, on the upper parte whereof on ech side of the midle ribbe groweth nine or ten broad short smooth sharpe pointed leaves and opposite against another, fast to the midle ribbe without any footstalk notched about the edges, and one alone at the toppe of the footstalk, which leaves are of a browne colour; amongst which cometh uppe a small round hollowe ioynted stalk about 3 foot high, no bigger then a parsley stalk, reddish towards the root, finely straked, not deeply champfered, devided into many partes towards the toppe, on ech ioynt groweth one leafe more notched and devided then the former, and those towards the toppe of the stalk have fewer leaves growinge on the sides of the midle ribbe then the lower, and of a lighter greene. The flowers growe on the topps of the branches in umbells, of colour white, everie flower havinge five very small leaves devided into two pts at the toppe, the flowers past the seed followeth, w^{ch} is small very like to parsley seed. The root is verie full of white hairy threddes, and putteth forth by the sides neue springes or shootes whereby it increaseth. The whole herbe doth yeld a stronge savor, like to Petroleum. This groweth plentifulle in the River by Droxford in Hampshire.—*MS. f. 82 v.*

Phyteuma orbiculare L.

Rapunculum silvestre Trago. p. 726. *Phyto.* 137 (4).

5^{to} Julij & 27 Augusti 1620

This hath 6 or 7 leaves with footstalks square abroad uppon the ground, in forme like those of ye wild March violett, but much smaller, finely indented about ye edges, amongst which riseth uppe a small, round straked stalk, not so bigg as a strawe, sometimes of a browne color, seldome a foot high whereon growe very small, narrowe, sharpe pointed leaves, without footstalks at ye toppe of ye stalk groweth one blewe flower tendinge to purple, almost round and sometimes somewhat longe, like those of ye comon Trefoile, composed of abundance of small crooked flowers, ech crooked flower beinge devided into two parts at ye toppe. After which follow

[] seeds contain'd in small husks. Ye root is white growinge deepe into ye ground, nine or ten ynches longe whereof 2 or 3 ynches of ye upp part is very small, and sometimes divided into more hedds than one. Ye midle part is Rampion-like, as bigge as a goose quill, the whole herbe & root beinge broken or cut doth yeld a white iuyce like milke.—*MS.* f. 84.

[Dr. Stapf points out that *P. spicatum* with which this plant has been identified has white flowers.]

Bartsia odontites Huds.

Eufrasia altera Dodo.

5^{to} Julij & 22 Augusti 1620

This sendeth forth from a small threddie, hard, crooked root, one 4 square, upright hard rough stalk, about 9 ynches or a foot high, divided into many branches, which are sett one opposite against another, ye leaves are small, rough, sharpe pointed, indented about ye edges, comonly hanginge or bowinge downewards, growinge by cooples, also one opposite against another, ye flowers are many hooded and growe but one side of ye branches and stalks, of a reddish color, with yealowish cheives in ye midle, after which cometh a small round seed vessell, neare as bigge as a wheate corne, wherein is contain'd [ends abruptly].—*MS.* f. 84.

Bastard Toadflax. *Thesium humifusum* DC.

Anthyllis montana. hist. lug. p. 1150. Phyto: 403. 15. an Anonymos Clus: p. 324.

5 Julij 1620 et citius et 27 Augusti 1620

[First record for Britain of the only species of the Sandalwood family known in Britain.]

This hath many very small round, cornered branches sometimes 17 or 18 from one root, which are divided into branches, growinge close uppon ye ground, sometimes 7 or 8 ynches longe, whereon growe very small narrowe, thick leaves out of order, one after, not one against another, of a ¹ or yealowish greene color, as are also ye branches, of a salt tast, neare ye topps of ye branches on short foot-stalks growe leaves smaller then ye other, 3 allwaies together whereof one is longer then ye other two, in ye midst of these 3 leaves groweth one small white flower, havinge 5 sharpe pointed leaves, spreadinge wide open starr fashion, in ye midle whereof groweth 7 small short cheives, with pale yealowe topps, after cometh one small long round harde husk, containinge a seed which is white within. The root is small white, crooked, short, divided into branches & threddie and is perennis.—*MS.* f. 84.

¹ The word 'pale' is struck out here, and the word substituted looks like 'nervie'.

Linaria adulterina.

n. d.

Johnson states that 'Mr. Goodyer found it growing wilde on the side of a chalkie hill in an inclosure on the right hand of the way, as you goe from Droxford to Poppie Hill in Hampshire'.—*Ger. emac.* 555.

Dwarf Thistle. *Carduus acaulis* L.

Carduus acaulis septentrionalium L'obelii. 8° Julij & 5 Sept. 1620
Ger. non habet.

It hath many greene short narrowe leaves, somewhat hairie, spread abroad upon the ground, seldome above six ynches longe, parted or gashed even to the midle ribbe, sett with sharpe prickles: amongst which groweth one (sometimes more) scaly head without prickles, with a thrum of purple flowers which are nothinge but small cheives, and after turne into downe, contayninge gray seed within: theis hedds growe close to the ground comonly without any stalk, yet sometimes havinge a small smooth footstalk 3 or 4 ynches longe. The root is small crooked scragged, with many out growinge branches, reddish in the midle, and of longe continewance. || It groweth wild on the Chalkie downes of Hampsheire plentifulle; and also at Purflett in Essex.—*MS.* f. 105.

[See 1618.]

Common Cow wheat. *Melampyrum sylvaticum* L.

Melampyrum luteum latifolium. Phyto. 444 (3). Pin. 234 (a. 4).
Crategeonon, Lob. icon. p. 36. *Parietaria silvestris*, 2 Clus. p. xliiij.
Ger. p. 84 (2). 22 Julij & 22 Augusti 1620

It hath a stalk about a cubit high, round close by ye root 4 square above, spreading it selfe abroad, often of a browne redd color on ye upper side, ioynted, devided ymediatly from ye root into branches, alwaies one branch growinge right against another, under which branches growe ye leaves, also one opposite against another, ye broadest and lowest are about 3 ynches longe, & one yinch broad, smooth nothinge at all notched by the sides, of a darke greene color, of an unpleasent an harsh tast, ye flowers growe neare ye toppes of ye branches, amongst smaller iagged leaves, and opposite against another, yet ye topps hang downwards & ioyne neare together of a yealowe color, which when they begin to wither are whitish, ye mouth not withstanding remayninge yealowe, after which followeth broad, flatt, sharpe pointed seed vessells, wherein is contayned 2 or 3 seeds like wheate cornes.

The root is small, whitish and threedie, and dieth at winter.—*MS.* f. 83 v.

Xeranthemum annuum L.

Ptarmica imperati.

26 Julii 1620

Ptarmica Austriaca species Clusii.

This riseth uppe with a small hard tough cornered whitish woolly stalk, devided into many branches, and those againe devided into other branches like those of *Cyanus* about two foot high, whereon grow long narrow whitish cottonie leaves out of order, of a bitter taste, whiter below than above, of the colour of the leaves of Wormwood, having but one rib or sinew & that in the midle of the leafe, and commonly turne downewards: on the top of each slender branch groweth one small scalie head or knap, like that of *Cyanus*, which bringeth forth a pale purple flower without smell, containing sixe, seven, eight, or more, small hard drie sharp pointed leaves: in the midle whereof groweth many stiffe chives, their tops being of the colour of the flowers; these flowers fall not away till the whole hearbe perisheth, but change into a rustie colour: amongst those chives grow long flat blackish seed, with a little beard at the toppe. The root is small, whitish, hard and threddie, and perisheth when the seed is ripe, and soone springeth up by the fallinge of the seed, and remaineth greene all the Winter, and at the Spring sendeth forth a stalke as aforesaid. The herbe touched or rubbed sendeth forth a pleasant aromaticall smell.—*MS.* f. 92; *Ger. emac.* 606-7.

Senecio sarracenicus L.

Herba Doria altera.

26 July 1620

This herbe groweth uppe with a greene round brittle stalke, very much champhered, sinewed, or furrowed; about 4 or 5 foot high, full of white pith like that of Elder, and sendeth forth small branches: the leaves growe on the stalk out of order, & are smooth, sharpe pointed, in shape resembling those of *Herba Doria*, but much shorter and narrower, the broadest and longest seldome being above 10 or 11 inches long, and scarce 2 inches broad, and are more finely and smally nickt or indented about the edges; their smell being nothing pleasant, but rather when together with the stalke they are broken and rubbed yeeld forth a smell having a small touch of the smell of Hemlocke. Out of the bosomes of these leaves spring other smaller leaves or branches. The flowers are many, and grow on small branches at the tops of the stalkes like those of *Herba Doria*, but more like those of *Iacobaea*, of a yealow colour, as well the middle button, as the small leaves that stand round about, every flower having commonly 8 of those small leaves. Which being past the button turneth into downe and

containeth a very small long seed which flieth away with the winde. The root is nothing else but an infinite of small strings which most hurtfullie spread in the ground, and by their infinite increasinge destroyeth and starveth other herbes that grow neare it. Its naturall place of growing. I know not, || for I had it from M^r. *John Coys*, and yet keep it growing in my garden.—*MS.* ff. 83, 92; *Ger. emac.* 431.

[The sentence after the || is written in a hand that I believe to be that of the editor, Thomas Johnson.]

Fasonia tuberosa DC.

Aster conyzoides Gesneri.

[Goodyer's note in Latin on this plant (*MS.* f. 83 v.) is copied from Lobel, *Observationes*, (1576) p. 189.]

Echinochloa Crus-galli Beauv.

Panicum sylvestre. Sheet.

10 Augusti 1620

Hath very many ioynted stalks, 2 foot high or higher, whereof some growe upright, and some growe sidelonge & leane towards ye earth, the leaves are longe & smooth not hairie. On ye toppe of ye stalks growe spikes or eares, sometimes single but comonly divided into many parts, lesser shorter & thinner than those of Panick, everie one whereof is composed of small short sharpe pointed husks, of a browne redd color sometimes of a greenish color, not [words illegible] ye stem on rib whereon they growe, but growinge on ye outer side thereof; wherein is a whitish seed, somewhat hard, lesse then those of Panick. The root is nothinge els but white strings.—*MS.* f. 83 v.

Basil. *Ocimum Basilicum* L.

Acinos [odoratissimum].

12 Augusti 1620 & longe before.

Phyto: 427 (4). *Ger.* 548. 2. *Math.* 595 (i).

It hath manie fower square hairie stalks proceedinge from the root, sometimes two foot longe or longer, parted into a fewe branches, the leaves growe on the ioynts, in wide distances by cooples one opposite against another, in forme like those of wild Margerom but smaller and are hairie, rough, lightlie snipt or indented about the edges: the flowers are purple & resemble those of Betonie, but of a lighter purple colour and growe forth of rough round whorles or crownetts close above the leaves, and one allwaies at the toppe of the stalk and branches, in forme like those of Horehound. The flowers past the seed followeth inclosed in those whorles. The root is small threedie & lastinge.—*MS.* ff. 83 v., 84.

[This description differs essentially from that printed in Gerard. See 11 Oct. 1621.]

Corn Parsley. *Carum segetum*, Benth.

Sium siifoliis. Hone-wort.

18 August 1620

¶ *The Description.*

This herbe commeth up at the first from seed like Parsley, with two small long narrow leaves, the next that spring are two small round smooth leaves nickt about the edges, and so for two or three couples of leaves of the next growth there are such round leaves growing on a middle rib by couples, and one round one, also at the top; after as more leaves spring up, so the fashion of them also change, that is to say, every leafe hath about eight or nine small smooth greene leaves, growing on each side of a middle rib one opposite against another, and one growing by it selfe at the top, and are finely snipt or indented about the edges, in forme resembling those of *Sium odoratum Tragi*, but not so bigge, long, or at all brownish; amongst which rise up many small round straked stalkes or branches, about two foot long, now and then above twenty from one root, sometimes growing upright, sometimes creeping not farre from the ground, joynted or kneed, and dividing themselves into very many branches, at every joynt groweth one leafe smaller than the former, which together with the lowermost perish, so that there is seldome one greene leafe to be seen on this herbe when the seed is ripe; the flowers are white, and grow most commonly at the tops of the branches, sometimes at most of the joynts even from the earth, in uneven or unorderedly umbells, every flower having five exceeding small leaves, flat, and broad at the toppe, and in the middle very small cheives with purple tops, the whole flower not much exceeding the bignesse of a small pins head, which being past there commeth up in the place of every flower two small gray crooked straked seeds, like Parsley seeds, but bigger, in taste hot and aromaticall. The root is small and whitish, with many threds not so big as Parsley roots. It beginneth to flower about the beginning of Iuly, & so continues flowering a long time; part of the seed is ripe in August, and some scarce in the beginning of October, mean while some falleth wherby it renueth it selfe, and groweth with flourishing greene leaves all the winter.

I tooke the description of this herbe the yeere 1620, but observed it long before, not knowing any name for it: first I refered it to *Sium*, calling it, *Sium terrestre*, and *Sium segetum & agrorum*; afterwards upon sight of *Selinum peregrinum primum Clusii*, which in some respects resembleth this herbe, I named it

Selinum Sij folijs; yet wanting an English name, at length about the yeere 1625 I saw Mistris *Vrsula Leigh* (then servant to Mistris *Bilson* of Mapledurham in Hampshire, and now (5 *Marcij* 1632 wife to Master *William Mooring* Schoolemaster of Petersfield, a Towne neere the said Mapledurham) gather it in the wheate ershes about Mapledurham aforesaid (where in such like grounds it still groweth, especially in clay grounds) who told me it was called Honewort.—*Ger. emac.* 1017–18.

[The remainder of the description has been printed on p. 53.]

Yellow Bird's-nest. *Monotropa Hypopithys* L.

Orobanche verbasculi odore, MS. Good: 22 August 1620

This riseth up with a soft round very brittle stalk, seldome 8 ynches high, sett with thinn small short scaly leaves like skynns growing close to the stalk, at or very near ye top of ye stalk groweth one sometimes 4 or 5 small flowers in fashion like ye flowers of *Hyosciamus luteus* or of y^e Cowslip every flower consisting of 4, but most commonly of 5 leaves, growing all of one height, evenly & of one proportion, & nothing like those of *Orobanche*. In ye middle of every flower groweth a small round umbo, no further out but just even with ye leaves, broad at ye top, with a small hoale in the middle, and ye lower end of which groweth at ye bottome of ye flower round, as bigge as a pease, so that it resembleth y^e suckbottle which children use to suck their drinke out of, having small cheives growing round about it with purplish tops. The root is obtuse, not usually so bigge as ye stalk, with very few threeds growing to it, & groweth at ye very upper face of ye earth. The whole herbe, flowers stalks & leaves are at their first flowring of a whitish yealowe, or strawe colour, and being broken or brused smelleth like to ye roote of a Primrose. This I found in a hedgerowe in a ground belonging to Droxford farme, neare ye foot path that leadeth from Droxford to Waltham, and took this Description ye 22 of August 1620.

[This copy of a lost note of G.'s is written on the back of f. 249 of Bannister's *Herbarium Siccum*, (Herb. Sloane) in the Botanical Department of the British Museum, to which my attention was drawn by Mr. J. Britten.

A similar description evidently derived from this same description and an engraving of the plant are given by Plot, *Nat. Hist. of Oxfordshire*, 1677, p. 146, who was the first to find this plant in Oxfordshire. Plot gives the number of flowers as 'eight or ten': he adds that 'It grows at the bottoms of Trees in the woods near Stoken-Church, and we find it mention'd in some MS. notes of the famous Mr. Goodyer'.]

Toothwort. *Lathraea squamaria* L.

Anblatum Dod. in fol. p. 553; Math. 689 (9). 15 Apr. 1621

Dentaria maior sine ἀφύλλος Clus: p. cxx.

From the root riseth uppe 2 or 3 sometimes more slender brickle stalks, hairy & full of iuyce, like those of Orobanche, 6 or 7 ynches longe, garnished with many flowers thick sett together not much unlike those of Satyrion or Orobanche, which doe all bend or looke that way which the stalk bendeth or leaneth. On ye back side growe 2 rowes of leaves, or rather small whitish skinne-like scales, and also amongst ye flowers there growe the like leaves: the flowers past there come small hedds wherein is containd very small seed the stalk soone perisheth, and leaveth the root in ye ground which is composed of whitish scales like teeth. The stalks and flowers when they growe in darke shadowie woods, are of a purplish color, but when they growe where the sunne cometh on them, they have no purple at all.

The scales on the root are not sharpe pointed as Clusius, Lobel & Dodo: pictureth them, but round topped, as Math: hath best of all by his figure expressed them.—*MS.* f. 87.

[This note is written on the back of an uncompleted order to the Overseers of the Poor, dated Southwick 1621.]

Cachryes.

28 Apr. 1621

[See under 9 May 1622.]

Juniper. *Juniperus communis* L.♂

Juniperus sterilis.

15 Maii 1621


This shrub is in the manner of growing altogether like the Juniper tree that beareth berries, only the upper part of the leaves of the youngest and tenderest bowes and branches are of a more reddish greene colour: the flowers grow forth of the bosoms of the leaves, of a yellowish colour, which never exceed three in one row, the number also of each row of leaves: each flower is like to a small bud, more long than round, never growing to the length of a quarter of an inch, being nothing else but very small short crudely chives, very thicke and close thrust together, fastened to a very small middle stem, in the end turning into small dust, which flieth away with the winde, not much unlike that of *Taxus sterilis*: on this shrub is never found any fruit.—*Ger. emac.* 1629.

Curly Pondweed. *Potamogeton crispus* L.

Tribulus aquaticus minor, quercus floribus [uvæ]. 2 Junii 1621

This water herbe bringeth forth from the root, thin, flat, knottie stalkes, of a reddish colour, two or three cubits long, or longer, according to the depth of the water (which when they are drie, are


pliant and bowing) devided towards the top into many parts or branches, bearing but one leafe at every ioynt, sometimes two ynches long, and halfe an ynch broad, thin, and as it were shining, so wrinckled and crompted by the sides that it seemeth to be



POTAMOGETON CRISPUS.

torne, of a reddish greene colour: the foot-stalkes are something long and thicke, and rise up from amongst those leaves, which alwaies grow two one opposite against another, in a contrarie manner to those that grow below on the stalk: neare the top of which foot-stalke groweth small grape-like huskes, out of which spring

very small reddish flowers, like those of the Oke, every flower having foure very small round topped leaves. After every flower commeth commonly foure sharpe pointed graines growing together, containing within them a little white kernell. The lower part of the stalke hath at every ioint small white threddie roots, somewhat



POTAMOGETON DENSUS.

long, whereby it taketh hold in the mudde, and draweth nourishment unto it. The whole plant is comonlie covered over with water. It flowereth in June and the beginning of July. I found it in the standing pooles or fish-ponds adioyning to a dissolved Abbey called Durford, which ponds divide Hampshire and Sussex, and in other standing waters elsewhere. This description was made]upon

sight of the plant the 2 of Iune, 1622.—*MS.* ff. 120 a, 122; *Ger.* . 823.

[In the fair copy (f. 122) Goodyer altered the name 'floribus uvae' to 'quercus floribus'. The rough copy (*MS.* f. 120 a) has the notes 'about ye beg. of July, or fortnight before Lamas' and 'both begunn to flower 2 Junij 1621' written in the margin. If the year 1622 given in the text be correct, Goodyer would appear to have been at Durford two years running on the 2nd of June.]

Pondweed. *Potamogeton densus* L.

Tribulus aquaticus minor, muscatellae floribus. 2 Junii 1621

This hath not flatt stalkes like the other, but round, kneed, and alwaies bearing two leaves at every ioint, one opposite against another, greener, shorter and lesser then the other, sharpe pointed, not much wrinckled and crumpled by the edges. *Clusius* saith, that they are not at all crompted. I never observed any without crumples and wrinckles. The flowers grow on short small foot-stalkes, of a whitish green colour, like those of *Muscattella Cordi*, called by *Gerard*, *Radix cava minima viridi flore*: viz. two flowers at the top of every foot-stalke, one opposite against another, every flower containing foure small leaves: which two flowers beeing past there come up eight small husks making six several waies a square of flowers. The roots are like the former. This groweth abundantly in the river by Droxford in Hampshire. It flowereth in June and July when the other doth, and continueth covered over with water, greene, both winter and somer.—*MS.* ff. 120 a, 122; *Ger. emac.* 823-4.

[The rough copy (*MS.* f. 120 a) has the note 'in running brooks' and a reference to *Clusius* cclij after the name.]

? *Melilotus indica* L.

Melilotus Indiae orientalis. 11 Junij 1621

This is in stalks, branches, leaves flowers and smell altogether like *Melilotus Italica* Camerarii, but smaller more branched & delicate, not growinge so high, & the stalks are greene and have no redd at all. || The seed is also like, but smaller. The root groweth downe right, and is small white, with a very fewe thredds, and perisheth when the seed is ripe the same yere it is sowen. || This hath not beene written of by any that I find. I received seeds thereof from Mr. William Coys often remembred.—*MS.* f. 97.

Epipactis violacea Dur.

Nidus avis flore et caule violaceo purpureo colore. an *Pseudo-leimodoron* *Clus.* *Hist. Rar. Pl.* p. 270. 29 June 1621

This riseth up with a stalke about nine inches high, with a few

small narrow sharpe pointed short skinny leaves, set without order, very little or nothing at all wrapping or inclosing the stalke; having a spike of flowers like those of *Orobanche*, without tailes or leaves growing amongst them: which fallen, there succeed small seed-vessels. The lower part of the stalke within the ground is not round like *Orobanche*, but slender or long, and of a yellowish white colour, with many small brittle roots growing underneath confusedly, wrapt or folded together like those of the common *Nidus avis*. The whole plant as it appeareth above ground, both stalkes, leaves, and flowers, is of a violet or deepe purple colour. This I found wilde in the border of a field called Marborne, neere Habridge in Haliborne, a mile from a towne called Alton in Hampshire, being the land of one *William Balden*. In this place also groweth wilde the thistle called *Corona fratrum*.—*Ger. emac.* 228; Druce, pp. 6–7.

[In Dillenius' interleaved copy of Ray's *Synopsis*, 1724, 'cum notis MSS. Lightfoot, Yalden, etc.', at the Botanic Garden at Oxford is a note, 'Limodorum Austriacum found in the border of a field called Marborne near Habridge in Haliborne a mile from Alton. Mr. Goodyer. See Camden's Britannia. I have searched for it in vain several years'.

The identity of this plant is very doubtful. See p. 47.]

Woolly Thistle. *Cnicus eriophorus* L.

Corona fratrum herbar. Caput monachorum. 29 June 1621

I found this wild in Hampshire in greate plentie by Haliborne in a feild called Marborne, nere a bridge called Habridge, beinge the land of Wm. Balden, & also in the next feild to it 29 Junii 1621.—*MS.* f. 53.

[See under 1617 and 13 Aug. 1621 for the description.]

Black Bryony. *Tamus communis* L.

Bryonia nigra florens et fructum ferens. Summer 1621

„ florens non fructum ferens.

This is altogether like the first described in roots, branches, and leaves; onely the foot-stalks whereon the flowers grow are about eight or nine inches long: the flowers are something greater, having neither before or after their flowering any berries or shew thereof; but the flowers and footstalks do soone wither and fall away: this I have heretofore, and now this Sommer, 1621, diligently observed, because it hath not bene mentioned or observed by any that I know.—*Ger. emac.* 871.

[The first record for Hants: probably of a male plant.]

Cerintho major L.

Cerintho flore rubro.

9 Julij 1621

The stalks or branches for the most parte growe uppe of 2 or 3 foot high devided into branches even from the root, about which growe leaves out of order, not one against another like the yealowe flowred *Cerintho*, but somethinge lesser, of a greene and blewe color as it were mixed together, also spotted with white spots on the upper side, the topps of the stalks bend downwards, and send forth amonge the leaves, out of cupps whose footstalks (not cupps as Clusius was informed) are of a deepe purple color, longe hollowe flowers like those of the said yealowe flowred *Cerintho* growinge but one in a place, of a redd purple color, which seeme to be sprinckled with a certaine whiteness, which fallen there followeth blackish seed contayned in small seed vessells two usually ioyned together in forme like to the seed of Borage or redd Ciches, but as bigge as a pease. The root is white and short with a few small branches or thredds, and perisheth when the seed is ripe.—*MS.* f. 120.

Caucalis latifolia L.

Caucalis maior Baetica.

9 Julii 1621

An *Caucalis pumila*, Clusio *Cur. post.* p. 37.

The stalks are rough round straked, a cubite high or higher, kneed or ioynted devided into branches, sett with rough iagged leaves, of an herbie smell, on the topps of the stalks and branches growe umbells of flowers, reddish before they open, after of a perfecte white color, the leaves that growe on the out side of ech particular flower are greater and broader then those within; close under ech umbell groweth a rowe of greene leaves cutt and devided into very small sharpe pointed leaves. After the flowers cometh greate prickley burrs half as bigge as those of *Xanthium* sett with rowes of large prickles everie burr when it is ripe partinge into 2 parts contayninge one seed a peece. The root is small white threddie in smell like to a carrott and perisheth when the seed is ripe. The whole plant in stalks, leaves and flowers is verie like to the wild carrott. || The seeds hereof I received from M^r William Coys often remembred, and he from Boelius a Lowe Contry man.—*MS.* f. 95.

Xanthium Strumarium L.

Bardana minor. The lesser Burre Docke.

? 1621

It groweth plentifulle in Southwicke street in Hampshire, as I have been informed by Mr. Goodyer.—*Ger. emac.* 810.

[Not included in Hants *Flora* by Townsend.]

Horned wild Cumīn. *Hypecoon procumbens* L.

Hypecoon Clusii.

11 Julij 1621

The leaves of this plant are devided into many partes, verie like in figure and color to the leaves of Rewe but softer or tenderer, and sometimes longer, spread uppon the ground ; amongst which rise uppe many small stalks, growinge slopewise, about 6 ynches longe, without leaves, but onlie neare the toppe where it devideth itself into 3, 4 or more small branchés and those againe sometimes devided into other little short branches, with little tender footstalks at their topps, ech foot stalk bearinge one small yealowe flower, somewhat sweet, contayninge 6 leaves, but of a strange fashion, and unlike other flowers, for it hath 2 leaves bigger then the rest, with a greene strake or line on the out side, (which is blunt and as it were bowed or folded in) the other fower leaves are very small, round topped and scarce to be seene, unlesse the flower be open, which past there succeed uppon everie of the said footstalks or branches, one longe crooked blunt topped codd, with ioyns knotts or devisions contayning in the space betwene everie bunch or knott, one blackish seed in a manner round, which is hard to be pulled forth. The root is single sometimes parted, somewhat yealowe, añuall, not livinge over yere, the whole plant with us is of little smell or none at all. || It groweth not wild in England. I have scene it growinge plentifullie in Mr. W^m Coyses garden in Essex, who kindlie imparted seeds thereof unto me, Anno 1620.—*MS.* f. 97.

Convolvulus purpureus L.

Convolvulus coeruleus minor Baeticus.

11 Julij 1621

This herbe hath many tender, small round weake hairie branches a foot or a Cubite longe or longer, devidinge it self presently from the root, some leaninge some traylinge on the ground none standinge upright, or windinge them selves about anie thinge, whereon growe rough leaves without footstalks, out of order, like those of *Alsine myosotis*, L'obelij, narrowe neare the stalk, broad towards the toppe, out of the bosomes of those leaves growe the flowers on long footstalks, made of one leafe, like in forme and bignes to those of the common *Convolvulus minor*, folded or plated when it is shut uppe, into five folds of a bright blewe color, with raies of yealowe in the bottom within, which fallen there succeedeth round buttons or knapps, as bigge or little bigger then a pease, wherein in ech button is contayned 2, 3 or 4 three cornered seeds almost as bigge as Radish seed. The root is small white single, and groweth

downeright with a few threddie strings or side branches. Both herbe and root do perish at winter. ||

The seeds hereof were gathered in Spaine by Boelius and communicated by him to my good friend Mr. William Coys who yerlie doth carefullie sowe the same and infinite other seeds of strange herbes, and hath imparted thereof unto me. It hath not bene hetherto written of that I knowe.—*MS.* f. 97.

Astragalus hamosus L.

Securidaca minor. Ob. p. 523. *Adversar.* p. 401. 13 Julij 1621

This plant is like to the greater *Securidaca* in stalks and leaves, but altogether smaller, and of a darker greene color, ech little leafe havinge a small nick at the toppe; the flowers are very small and white, (and not purple with us, as Pena hath it) growinge in small tufts: after which followe round, crooked, sharpe pointed coddcs, so sharpe that they will perce into the flesh, first bowinge downwards to the footstalk, and then turninge uppe againe, like crooked hornes, wherein is contayned two rowes of small seed, in fashion of a kidney almost round, in tast like drie pease or beanes, not bitter; of a darke color drawinge neare an ash color. The root is small white with some threddes, and perisheth when the seed is ripe. || Seeds hereof were sent me from Mr. William Coys a man verie skillfull in the knowledge of symples, my singuler good frend.—*MS.* f. 97.

Iberis umbellata L.

Thlaspi umbellosum marinum flore albo. 14 Julij 1621

An *Thlaspi quartum parvum odorato flore Clusii*, p. cxxxij.

This plant riseth uppe with one small stalk and divideth it self ymediatelie from the root into many branches which are about a foot high, greene, round, rough brittle straked as is the stalk and everie branch devidinge it self into more branches, whereon growe longe narrowe rough leaves, devided neare the toppe like to *Cornu Cervinum L'obelij*, of a bitter tast, which leaves sometimes turne downewards: on the toppe of the stalks & branches growe umbells of pleasant flowers, of a stronge but no pleasant smell, appearinge purple before they be open, (the midle of the umbell alwaies flowringe last) but beinge fullie blowen, they are of a perfect white color, ech flower beinge made of fower blunt topped leaves, sometimes nicked at the toppe, and the two leaves that growe on the out side, are neare fower times as bigge, as those two that growe within, havinge short yealowe cheives in the midle, which fallen there succeedeth || broad flatt powches, seed vessels or husks, forked

or devided into two parts at the toppe, ech huske contayning two flatt yealowish seeds, of a very bitter tast. The root is white and short, with a few side branches. The whole herbe perisheth when the seed is ripe.—*MS.* f. 98.

[A rough copy of the last five lines is written on *MS.* f. 11.]

Asclepias purpurascens L.

Periploca recta Virginiana.

14 Julij 1621

An Apocynum Syriacum Clusii, p. lxxxvij.

The stalk is bluntlie fower squared, upright, strait, not branched fower or 5 foot high, straked, of the bignes of a finger, black spotted, full of spongious white pith within, whereon growe by cooples, one opposite against another, uppon thick short footstalks, large, broad blunt topped leaves, with a whitish softe cotton underneath, of a light greene above, plaine and smooth not notched by the sides, neare the toppe of ech stalk, exactlie against the leaves (not out of the bosomes of the leaves) groweth forth about two short rough footstalks, but one in a place, ech bearinge 4 or more knapps like small buttons, which by nature turne downe, open and appeare like a greenish flower, with five leaves; above which groweth the flower of a purple or reddish color, composed of 5 round small hollowe leaves sharpe pointed on the outside, ech leafe havinge a short sharpe pointed pointell of the color of the flower, turninge into the umbo or midle of the flower.—*MS.* f. 98.

Anchusa angustifolia L.

Buglossum Scorpioides.

14 Julij 1621

An Echij facie Buglossum minimum flore rubente. Ob. p. 310.

This is like the common Buglosse in stalks and leaves, but altogether smaller & lower, the topps of the branches whereon the flowers growe, are more crooked, turninge and bowing inwards, the occasion (as I take it) of the name. The flowers are smaller, the leaves thereof narrower at the toppe, at their first openinge more reddish, and when they are fullie blowne, of a perfecte purple, or black violet color, but in tast, savor, operation and all things else alike.—*MS.* f. 98.

Ornithopus scorpioides L.

Scorpioides Mathioli.

16 Julij 1621

Math. p. 895. Phyto. 568 (5). Ger. hath it not.

This hath many small round firme branches proceedinge from one root, and those againe devided into other branches, whereon growe leaves almost round, comonly 3 sometimes 4 together; whereof the midlemost is 4 times as bigge as anie of the other, both

stalks and leaves, are of a mealie white color, and are some what like the leaves of *Portulaca*: the flowers are small, yealowe, like those of *Scorpioides repens Bupleuri folio L'Obelij*, but smaller, growinge forth of the bosomes of the leaves, on long footstalks, sometimes three, most comonly 4 together. After which cometh on ech footstalk as many longe, small, jointed. slender, sharpe pointed codd, bowinge like the taile of a scorpion or like to birds claws, wherein is containd betweene everie ioynt one longe slender yealowish seed like *Galega*. The root is small, white, with a fewe thredds, and perisheth when the seed is ripe.—*MS.* f. 88.

Crucianella sp.

Rubia spicata Cretica Clusii, p. clxxvii.

19 July 1621

This hath proceeding from the root many knottie foure square rough little stalks, a foot high, devided immediately from the root into many branches, having but one side branch growing forth of one ioint: about which ioints grow spread abroade 4, 5, sometimes 6 narrow, short, sharpe pointed leaves, somewhat rough: the toppes of the stalkes and branches are nothing but long small foure square spikes or eares, made of three leafed greene huskes: out of the top of each huske groweth a very small greenish yellow flower, having foure exceeding smal leaves scarce to be seene: after which followeth in each huske one smal blackish seed, somewhat long, round on the one side, with a dent or hollownesse on the other. The root is small, hard, woodie, crooked or scragged, with many little branches or threds, red without, and white within, and perisheth when the seed is ripe.—*MS.* f. 88; *Ger. emac.* 1119.

Campanula pumila hort. *C. pulla* e.

Lesser Bellflower.

[19 July 1621]?

Ye lesser Bellflower hath small, rough, round, straked stalks, seldom branched, or above a foot or a cubit high. The leaves are rough, of a light greene colour, very lightly indented about y^e edges in form like sloe leaves, or those of ye wild violett, those that grow below have footstalks, & those above are smaller, & grow close to ye stalk without any footstalk at all; the flowers are like small bells, & devided in the toppe into 5 sharpe pointed leaves, which are pointed in y^e middle devided in ye toppe in 3 pts. Some whereof growe alongest ye stalk, and some times 12 together in a bunch or Cluster at very toppe of ye stalk, after which cometh ye seed cod in small rough(?) husks. The root is white, small, strayght, with many threads & perennis.—*MS.* f. 8 v.

[Note on the back of a letter from F. Waller to Sir T. Bilson, bearing date 19 July 1621.]

Phalaris canariensis L.

Phalaris minor Baetica Boëlij, semine nigro. [20 Julij 1621]

Is in stalks, height, leaves and scalie eares, like the greate Phalaris, but twice as small, of a darker greene color, and the seed is much smaller and of a blackish color.—*MS.* f. 88.

Phalaris minor L.

Phalaris minor Baetica semine albo Boëlij. [20 July 1621]

The stalks growe not above a foot high, the leaves are whighter then the other, the eares are also whighter, and scarce growe forth of the greate hose or uppermost leafe, the seed is whitish in all things else like the former.—*MS.* f. 88.

Phalaris bulbosa L.

Phalaris bulbosa Boëlij. [20 July 1621]

Is altogether like the Phalaris with blackish seed in stalks, leaves and spikes, but the roots are bulbus, like those of Catts taile grasse, growinge most comonlie on the upper crust of the earth, with a few small threddie roots hanginge thereat fastened within the earth, the seed is of an Ash color or darke white, the bulbus roots most comonlie live manie yeres.—*MS.* f. 88.

Valeriana Cornucopiae L.

Valeriana mexicana. [20 Julij 1621]
non est Valeriana Indica Clusii.

The stalke is round, tender, brittle, verie much straked, hollowe, greene, yet reddish in some places, & ioynted, devided into branches, about 2 foot high; at everie ioynt groweth two leaves, those on the lower parte of the stalk are crompted, not notched by the sides, round topped, of a light greene color, verie like the leaves of garden lettice (nothings at all like the leaves of *Lactuca agnina*) about 4 ynches longe, and 3 ynches brode, with broad leafed footstalks; those on the upper parte of the stalke, are narrower, shorter, somewhat notched by the sides, without footstalks; on the topps of the branches growe the flowers as it were in umbells, and are longe, of a bright purple color, everie flower contayninge 5 small round topped leaves, whereof 2 are greater then the rest. The seed followeth (sem. 1. Sept.) growinge in chaffie scales or hedds, as bigge or bigger then a wheate corne, blackish, seminge to be nothings but light husks and no seed.

The root is very small for the bignes of the plant, white and threddie, and perisheth when the seed is ripe; the whole plant is without any manifest tast or smell. || Clusius, Gerard & Bauhinus

have written of *Valeriana mexicana* sive *indica*, but this herbe agreeth with neither of their descriptions. I received seeds whereof grewe this described herbe from Mr. William Coys often remembred.—*MS.* f. 89.

Reseda Phyteuma L.

Phyteuma monspeliensium. Ger. p. 918. 21 Julij 1621

The stalks are small, greene, round, straked, rough, creeping or leaninge neare the ground, a foot long, devided into many branches immediatlie from the root; the leaves are small like those of *Lutcola*, round topped, rough on the midle ribbe on the under side, sometimes crompted by the sides; the flowers growe from the midle pte of the stalk upwards spike fashion, not close together, on short rough foot stalks; beinge nothinge but thrumie whitish cheives, like those of *Rheseda*; ech flower havinge growinge close adioyninge under it comonlie 6 small leaves star fashion, the flowers past.—*MS.* f. 89.

? *Malva stipulacea* Cav.

Malva flore amplo Baetica aestiva. 21 Julij 1621

This mallowe hath stalks and leaves altogether like the greate comon high wild mallowe; but the flowers are as bigge againe, composed likewise of five leaves a peece, of a lighter purple color and makinge a braver showe, the seed is blackish like a half moone, not rough [but finely straked on the upper side, twice as bigge as the seed of the comon mallowe, growinge in a round circle closely compact together within or at the bottome of a duple huske, and covered over with a flatt spongiouse cake, blackish on the upper side, so this seed in manner of growinge is contrarie to the comon mallowe, this havinge two husks and a little cake and seed underneath, and that having by one huske a little cake but half so bigge, made only of the seed. The root is very small, white, brittle, with a few small threddes hanginge thereat and perisheth at winter. ||

I cannot find that this mallowe hath been written of heretofore. The seeds were sent to my worthy frend Mr. William Coys by Boelius who gathered them in Spaine, who with many other imparted them unto me Anno 1620.—*MS.* f. 89.

Chrysanthemum coronarium L.

Chrysanthemum Creticum primum Clusii. 28 Julii 1621

The stalkes are round, straked, branched, hard, of a whitish greene, with a very little pith within; neere three foot high: the

leaves grow out of order, devided into many parts, and those again snipt or devided, of the color of the stalkes : at the topps of the stalkes and branches grow great flowers, bigger than any of the rest of the Corne-flowers, forth of scaly heads, consisting of twelve or more broad leaves apeece, notched at the top, of a shining golden colour at the first, which after turne to a pale, whitish, or very light yellow, and grow round about a large yellow ball, of smell somewhat sweet. The flowers past, there commeth abundance of seed closely compact or thrust together, and it is short, blunt at both ends, straked, of a sand color, somewhat flat, & of a reasonable bignes. The root is whitish, neere a fingers bignesse, short, with many threds hanging thereat, and perisheth when the seede is ripe ; and at the Spring groweth uppe againe by the falling of the seed.—*MS.* f. 90 ; *Ger. emac.* 744-5.

Chrysanthemum coronarium L.

Chrysanthemum Baeticum Boelij inscriptum. 28 Julij 1621

The stalks are round, straked, reddish brown, devided into branches, containing a spongius white pith within, a cubite high : the leaves growe out of order, without footstalkes, about 3 inches long, and an inch broad, notched about the edges, not at all devided, of a darke greene colour : the flowers growe at the tops of the stalkes and branches, forth of great scalie heads, containinge twentie leaves a piece or more, notched at the top, of a shining yellow color, growinge about a round yellow ball, of a reasonable good smell, verie like those of the common *Chrysanthemum segetum* : the seede groweth like the other, and is very small, long, round, crooked and whitish : the root is small, whitish, threddie, and perisheth also when the seed is ripe.—*MS.* f. 90 ; *Ger. emac.* 745.

Chrysanthemum coronarium L.

Chrysanthemum tenuifolium Baeticum Boelij. 28 Julij 1621

The stalks are round, small, straked, reddish, somewhat hairie, branched, a cubit high, or higher : the leaves are small, much devided, jagged, and verie like the leaves of *Cotula foetida* : the flowers are yellow, shining like gold, composed of thirteene or fourteene leaves a pece, notched at the top, set about a yealowe ball, also like the common *Chrysanthemum segetum* : the seed groweth amongst white flattish scales, which are closelie compacted in a round head together, and are small, flat, greyish, and broad at the top : the root is small, whitish, with a few threds, and dyeth when the seed is ripe.—*MS.* f. 90 ; *Ger. emac.* 745.

Prickly Beet of Candy. *Emex spinosa* Campd.

Blitum spinosum Creticum.

28 Julij 1621

est Beta Cretica semine aculeata Bauhini. Joynt Flower-topp.

This sendeth forth from one root many round greene straked, trayling, ioynted, small branches, about a foot longe: the leaves are of a light greene colour, and grow at every ioint one, somewhat like the leaves of great Sorrell, but they are round topped without barbes or eares below, or any manifest tast or smell, very like the leaves of Beetes, but much smaller: the flowers grow clustering together about the ioyns, and at the tops of the branches small and greenish, each flower containing 5 or 6 very small blunt topped leaves, and a few dustie cheives in the middle: which past, there cometh greate prickley shrivelled seed, growing even close to the root, and upwards on the ioints, each seed having three sharpe prickes at the top growing sidewaies, which indeed may be more properly called the huske; which huske in the inside is of a darke reddish color, and containeth one seed in forme like the seed of *Adonis*, round at the lower end, 3 cornered towards the toppe, and sharp pointed, covered over with a darke yealowish skyn; which skyn pulled away, the kernell appeareth yealow on the outside, and exceeding white within, and will with a light touch fall into very small powder like meale.—*MS.* f. 90; *Ger. emac.* 1626.

Lathyrus Ochrus DC. or *L. annuus* L.

Lathyrus aestivus flore luteo.

28 Julii 1621

This is like *Lathyrus latiore folio Lobelij*, in stalks, leaves, and branches, but smaller: the stalks are two or three foot long, made flatt with two skyns, with 2 exceeding small leaves growinge on the stalks, one opposite against another: betweene which spring up flat footstalks, an inch long, bearing two exceeding narrow sharpe pointed leaves, three inches long: betweene which grow the tendrels, divided into many parts at the top, and taking hold therwith: the flowers are smal, and grow forth of the bosomes of the leaves, on each footstalk one flower, wholly yealow, with purple strakes. After each flower followeth a smooth cod, almost round, two inches long, wherein is contained seven round Peason, somewhat rough, but after a curious manner, of the bignesse and taste of field Peason, and of a darke sand color.—*MS.* f. 108; *Ger. emac.* 1628.

? *Lathyrus Clymenum* L.

Lathyrus aestivus Baeticus flore coeruleo Boëlii.

[28 Julii 1621]

This is also like *Lathyrus latiore folio Lobelij*, but smaller, yet

greater than that with yellow flowers, having also adjoining to the flat stalkes, two eared sharpe pointed leaves, and also two other slender sharpe pointed leaves, about foure inches long, growing on a flat foot-stalke betweene them, an inch and a halfe long, and one tendrel between them devided into two or three parts: the flowers are large, and grow on long slender foure-square foot-stalkes, from the bosomes of the leaves, on each footstalk one: the upper great covering leafe being of a light blew, & the lower smaller leaves of a deeper blew: which past there come up short flat cods, with two filmes, edges, or skins on the upper side, like those of *Eruilia Lobelij*, containing within foure or five great flat cornered Peason, bigger than field Peason, of a darke sand color.—*MS.* f. 108; *Ger. emac.* 1628.

? *Lens esculenta* Moench.

Lathyrus aestivus edulis Baeticus flore albo Boelii. [28 Julii 1621]

This is in flat skinny stalks, leaves, foot-stalks, and cods, with two skins on the upper side, and in all things else like the said *Lathyrus* with blew flowers; only the flowers of this are milk white: the fruit is also like.—*MS.* f. 108; *Ger. emac.* 1628.

? *Lathyrus sphaericus* Retz.

Lathyrus aestivus flore miniato. [28 Julii 1661]

This is also in skinnie flat stalks and leaves like the said *Lathyrus latiore folio*, but far smaller, not three foot high: it hath also small sharp pointed leaves growing by couples on the stalke, between which grow two leaves, about three inches long, on a flat foot-stalk half an inch long: also between those leaves grow the tendrels: the flowers are of the color of red ledd, but not so bright, growing on smooth short foot-stalks, one on a foot-stalke: after which follow cods very like those of the common field peason, but lesser, an inch and a halfe long, containing foure, five, or sixe cornered Peason, of a sand color, or darke obscure yealowe, as big as common field peason, and of the same taste.—*MS.* f. 108; *Ger. emac.* 1628-9.

Lathyrus palustris L.

Lathyrus palustris Lusitanicus Boelii. [28 Julii 1621]

Hath also flat skinnie stalks like the said *Lathyrus latiore folio*, but the paire of leaves which grow on the stalke are exceeding small as are those of *Lathyrus flore lutco*, and are indeed scarce worthie to be called leaves: the other paire of leaves are about two inches long, above halfe an inch broad, and grow from betweene those small leaves, on flat foot-stalks, an inch long: betweene which

leaves also grow the tendrels: the flowers grow on footstalks which are five inches long, commonly two on a foot-stalke, the great upper covering leaves being of a bright red colour, and the under leaves are somewhat paler: after commeth flat cods, containing seven or eight small round peason, no bigger than a Pepper corne, gray and blacke, spotted before they are ripe, and when they are fully ripe of a blacke colour, in taste like common Peason: the stalks, leaves, foot-stalkes and coddess are somewhat hairy and rough.—*MS.* f. 109; *Ger. emac.* 1629.

Lathyrus tuberosus L.

Lathyrus aestivus dumetorum Baeticus Boelii. [28 Julii 1621]

Hath also flat skinnie stalks like the said *Lathyrus latiore folio*, but smaller, and in the manner of the growing of the leaves altogether contrarie. This hath also two small sharp pointed leaves, adioyning to the stalke: betweene which groweth forth a flat middle rib with tendrels at the top, having on each side (not one against another) commonly three blunt topped leaves, sometimes three on the one side, and two on the other, and sometimes but foure in all, about an inch and a halfe long: the flowers grow on foot-stalks, about two or three inches long, each foot-stalk usually bearing two flowers, the great covering leafe being of a bright red colour; and the two under leaves of a blewish purple colour: after which follow smooth cods, above two inches long, containing, five, sixe, or seven smooth Peason, of a browne Chestnut colour, not round, but somewhat flat, more long than broad, especially those next both the ends of the cod, of the bignesse and taste of common field peason.—*MS.* f. 109; *Ger. emac.* 1629.

cia sativa β *linearis* Lange.

Aracus maior Baeticus Boelii.

30 Julii 1621

It hath small weake foure square straked trayling branches, two foot high, lesser, but like those of Fetches; whereon grow manie leaves without order, and every severall leafe is composed of six, seven, or more small sharpe pointed leaves, like those of Lentils, set on each side of a middle rib, which middle rib endeth with clasping tendrels: the flowers grow forth of the bosomes of the leaves, but one in a place, almost without any foot-stalkes at all, like those of Vetches, but of a whitish colour, with purple strakes, and of a deep colour tendinge to purple towards the nailes of the upper covering leaves: after which follow the cods, which are little above an inch long, not fully so big as those of the wilde beane, almost round, and very hairy: wherein is contained about 4 peason,

seldom round, most commonly somewhat flat, and sometimes cornered, of a blackish colour, neere as big as field peason, and of the taste of Fetches. The whole herbe perisheth when the seed is ripe. || This plant *Boelius* sent to M^r. *William Coys*, who hath care fully preserved the same kind ever since, and friendly imparted seeds thereof to me in *Anno 1620*.—*MS. f. 91*; *Ger. emac. 1627*.

Vicia lutea β *laevigata* Boiss.

Legumen pallidum Vllissiponense Nonii Brandonii. 30 Julii 1621

This plant is very like, both in stalks, leaves, and cods, to *Aracus maior Beticus*, but the flowers of this are of a pale yellow or primrose colour, and the whole herbe smaller, and nothing so hairy. It perisheth also when the seed is ripe. || I received the seeds likewise from M^r. *Coys*.—*MS. f. 91*; *Ger. emac. 1627*.

Vicia sativa var. *leucosperma* Moench.

Vicia indica fructu albo. *Pisum indicum* Gerardo. 30 Julii 1621

This Vetch differeth not in any thinge at all, eyther in stalkes, leaves, coddcs, fashion of the flowers, or colour thereof, from our common manured Vetch, but that it groweth higher, and the fruit is bigger and rounder, and of a very cleare white colour, more like to peason than Vetches. || M^r. *Gerard* was wont to call this Vetch by the name of *Pisum Indicum*, or Indian Pease, gotten by him after the publishing of his Herball, as M^r. *Coys* reported to me. But the said M^r. *Coys* hath (in my iudgment) more properly named it *Vicia fructu albo*: which name I thought most fit to call it by, only addinge *Indica* to it, from whence it is reported to have been gotten.—*MS. f. 91*; *Ger. emac. 1627*.

Pisum sativum L.

Pisum quadratum.

30 Julij 1621

Lotus siliquosus rubello flore. Clus: p. ccxliiij. Phyto. 668 (60).

This hath many round, hairie, branches, proceedinge immediatelie from the root, longe spreadinge for the most pte uppon the ground; yet the topps or ends of the branches liste themselves somewhat upright, whereon growe broad rough hairie blunt-topped leaves at certaine distances 3 allwaies on one foot stalk, out of whose bosomes growe round hairy footstalks, bearinge three other leaves like the former but smaller, and one or two flowers like those of Vetches, of a beautifull color like deepe redd or orange tawnie velvett, after which cometh uppe the coddcs, about 3 ynches longe, havinge 4 wrinkled filmes or skyns growinge alongst them viz. two on the upper side and 2 on the lower side, which make the coddcs (?) appeare fower square when they be drie, wherein in ech

codd is containd about 10 or 11 round seeds, browne when they are ripe, neare of the bignes of feild peason, and of the same tast. The root is white and small, with abundance of thredie small side branches, and perisheth at winter.

It begineth to flower in June, and so contineweth flowringe and bearinge fruite till the extreame froste.—*MS.* f. 91.

[Goodyer sent a description of *Pisum quadratum* to Johnson on 5 March 1632, but the text printed in the *Herbal* differs markedly from this one: cf. *Ger. emac.* 1198.]

Astragalus lusitanicus Lam.

Astragalus marinus lusitanicus Boelii.

31 Julii 1621

This hath five, six, or more round straked reddish hairy stalks or branches, of a reasonable bignesse, proceeding from one root, sometimes creeping or leaning neere the ground, and sometimes standing upright, a cubit high, with many greene leaves, set by certaine distances, out of order like those of *Glaux vulgaris*, but lesser, every leafe being composed of fourteen or more round topped leaves, a little hairy by the edges, set on each side of a long middle rib, which is about nine or ten inches in length, without tendrels: the flowers grow forth of the bosomes of the leaves, neere the tops of the stalkes, on long round streaked hairy foot-stalkes, of a very pale yellow colour, like those of *Securidaca minor*, but bigger, growing close together in short spikes, which turne into spikes of the length of two or three inches, containing many small three cornered cods about an inch long, growing close together like those of *Glaux vulgaris*, each cod containing two rowes of small flat foure cornered seeds, three or foure in each row, of a darke yellowish or leadish colour, like to those of *Securidaca minor*, but three or foure times as big, of little taste: the root is small, slender, white, with a few threds, and groweth downe right, and perisheth when the seed is ripe. || I first gathered seeds of this plant in the garden of my good friend M^r. *John Parkinson* an Apothecary of London, Anno 1616.—*MS.* f. 107; *Ger. emac.* 1627–8.

Vicia Faba L. var.

Faba veterum serratis foliis Boelii.

31 Julii 1621

This is like the other wilde Beane in stalks, flowers, cods, fruit, and clasping tendrels, but it differeth from it in that the leaves hereof (especially those that grow neere the tops of the stalks) are notched or indented about the edges like the teeth of a saw. The root also perisheth when the seed is ripe. || The seeds of this wilde Beane were gathered by *Boelius* a Low-country man. in *Batica* a part of Spaine, and by him sent to M^r. *William Coys*, and by

him carefully preserved, who also imparted seeds thereof to me, in Anno 1620.—*MS.* f. 107; *Ger. emac.* 1628.

Lathyrus Ochrus DC.

Ervilia silvestris Dodonaei, p. 522. 31 Julij 1621

Ervilia sendeth forth 3 or 4 stalks or branches from one root, somewhat like *Lathyrus*, but broader weaker and lyeinge flatt upon the ground: the leaves are about two ynches longe and an ynch broad, with claspinge tendrells at the ends, without footstalks, making the stalks flatt with their two edges, whole bendath on the lower parte of the stalk onlie devided into 2 parts at the ends, but neare the topps of the branches everie leafe is devided at the toppe into 2, 3 or 4 small leaves. The flowers are small of a pale yealowe or primrose color growinge but one in a place on a short footstalk. The coddys are short somewhatt flatt havinge 2 edges or filmes on the upper side wherein is containyd 4 or 5 or 6 round ash colord fruite, verie like feild peason of the same bignes, and verie neare of tast. The root is verie small and threddie and perisheth when the seed is ripe.

|| I first observed this pulse in the garden of Mr. John Parkinson in London Anno 1616, and after 1620 I received seeds hereof from my trewe frend Mr. William Coys, often remembred, with many other.—*MS.* f. 107.

[*Cf. L. Ochrus* DC., 28 July 1621.]

Pisum arvense L.

Pisum maculatum Boelii. 31 Julii 1621

They are like to the small common field Peason in stalkes, leaves, and cods; the difference is, the flowers are commonly smaller, and of a whitish greene colour: the Peason are of a darke gray colour, spotted with blacke spots in shew like to blacke velvet; in taste they are also like, but somewhat harsher. || These peason I gathered in the garden of Mr. *John Parkinson*, a skilfull Apothecarie of London; and they were first brought out of Spaine by *Boelius* a low-countray man.—*MS.* f. 107; *Ger. emac.* 1628.

Medicago minima L.

Medica Anglica minor 2 Augusti 1621

an *Trifolium cochleatum alterum*. Dod. p. 579.

Hath many fower square hairy straked reddish branches, growinge from one root, two or three foot longe, and those also devided againe into other branches, whereon growe smooth leaves three on a footstalk, somewhat indented, very broad at the toppe, and narrowe belowe, of the fashion of a hart, with a crooked black

spott in the middest of ech leafe. The flowers are very small and yealowe, growinge on short footstalks, alongest the branches, forth of the bosomes of the leaves, 3, 4 or 5 on a footstalk ; after cometh small round fruits, writhen or wound together, flatt at the toppe, prickley, no bigger then a small pease, wherein lyeth small yealowe seed, in fashion of a kidney, but no bigger then a flea. The root is small, whitish, groweth downeright with a fewe side branches and small thredds, and perisheth at winter after the seed is ripe.—*MS. f. 104.*

Medicago polymorpha L.

Medicae maioris Baeticae species prima, 2 Aug. 1621
spinulis intortis.

Hath also foure square reddish streaked hairy trailing branches, like the small English *Medica*, but greater and longer, foure or 5 foot long: the leaves are also smooth, growing three together, neither sharpe pointed, nor yet so broad at the top as the said English *Medica*, but blunt topped, with a small blacke spot in the midst, not crooked: the flowers are also yellow, three, foure, or five on a foot-stalke: after commeth a round writhed fruit fully as big as a hasell nut, with small prickles not standing foreright, but lying flat on the fruit, finely wrapped, plaited, folded, or interlaced together, wherein lieth wrapped the seed in fashion of a kidney, very like a kidney beane, but foure times smaller, and flatter, of a shining blacke colour without, like polished Ieat; containing a white kernell within: the root is like the former, and perisheth also at Winter.—*MS. f. 104; Ger. emac. 1200.*

[‘Ieate’ in MS., probably = jet.]

Hedgehog Medick. *Medicago intertexta* L.

Medicae maioris Baeticae spinosae species altera. 2 Aug. 1621

The branches also creepe on the ground, and are straked smooth foure square, reddish here and there, three or foure foot long: the leaves are smooth, finely notched about the edges, sharp pointed, without blacke spots, very like *Medica pericarpio plano*: the flowers are small and yellow like the other: the fruit is round, writhed or turned in, also fully as big as a hasell nut, somewhat cottonic or woolly, with short sharpe prickles: wherein lyeth also wrapped a shining blacke kidney-like seed, so like the last described, that they are not to be discerned apart: the root is also alike, and perisheth at Winter.—*MS. f. 104; Ger. emac. 1200.*

Medicago marina L.

Medicae marinae spinosae species. 2 Aug. 1621

The branches of this are the least and shortest of all the rest,

little exceeding a foot or two in length, and are foure square, greene, somewhat hairie, and trailing on the ground: the leaves are like to those of *Medica pericarpio plano*, not fully so sharpe pointed, without blacke spots, soft, hairy, three on a foot-stalke: the flowers grow alongst the branches, on very small foot-stalkes, forth of the bosomes of the leaves, (not altogether on or neere the tops of the branches) and are very small and yellow, but one on a foot-stalke: after commeth small round writhed fruit, no bigger than a pease, with very short sharpe prickles, wherein is contained yellowish seed of the fashion of a kidney like the former, and is the hardest to be plucked forth of any of the rest: the root is also whitish like the roots of the other, and also perisheth at Winter.—*MS.* f. 104; *Ger. emac.* 1200.

Linaria thymifolia DC.?

Antirrhinum minus flore Linariae luteum inscriptum. 3 Augusti 1621

This hath at the first many very small, round, smooth branches from one root, trayling on the ground, about foure or five inches long, set with many small greene short sharp pointed leaves, like those of *Serpillum*, but that these are longer, smooth, and three or foure growing opposite one against another: amongst which rise up five or six, sometimes ten or twelve upright round smooth little stalks a cubit high, divided into branches bearing small long smooth greene leaves, growing without order, as narrow as the upper leaves of *Oenanthe Angustifolia*: at the toppes of the stalks and branches grow clustering together five six or more small yellow flowers, flowering upwards, leaving a long spike of very small huskes, each huske having a small line or chinke as though two huskes were ioined together, the one side of the huske being a little longer than the other, wherein is contained exceeding small blackish seed. The root is very short, small, and white, with a few threds, and perisheth at winter.

|| This plant is not written of that I can finde. I received seed thereof from M^r. *William Coys* often remembered.—*MS.* f. 103; *Ger. emac.* 1626.

Linaria serpyllifolia Lange?

Linaria minor aetiva.

3 Augusti 1621

The stalkes are round, smooth, of a whitish greene colour, a foot high, weake, not able to stand upright: whereon grow long narrow sharpe pointed leaves, most commonly bending or turning downwards. The flowers grow in spikes at the toppes of the branches, yet not very neere together, and are verie small and yellow, with

a small taylor: the seed of this plant is small, flat, and of a blackish gray colour, inclosed in small round huskes, and you shall commonly have at one time flowers and ripe seed all on a stalk. The whole plant is like to the common *Linaria*, but that it is a great deale lesser, and the flowers are six times as small, and perish at Winter. || I also received seeds thereof from Mr. *William Coys*.—*MS.* f. 103; *Ger. emac.* 1626.

? *Trifolium Lagopus* L.

Lagopus trifolius flore ruberrimo.

4 Augusti 1621

This *Lagopus* sendeth forth many leaves presentlie from the root three growinge together upon a longe footstalk almost round softe hairie of a light greene amongst which groweth uppe a round hairy reddish stalk of a foot or a cubite high divided into certaine branches under which growe the like leaves not so round but a little longer. At the topps of the stalks and branches growe longe heads or spikes about two ynches longe of a finger bignes made of rough husks ech huske divided into five small narrowe parts at the toppe out of which growe small flowers of a delicate bright redd color, which fallen there succedeth in ech of those husks a round thick yealowe seed. || The root is very small, white, short, and perisheth when the seed is ripe.—*MS.* f. 106.

Trifolium ligusticum Balb.

Lagopus trifolius maior Baeticus.

[4 Augusti 1621]

Hath many small round weake hairy branches about nine ynches longe proceedinge from one root traylinge on the ground and those againe divided into other branches whereon growe small greene round topped hairie leaves three together on a round hairy footstalk like those of *Trifolium flore albo* but longer. On the topps of the branches growe round hairie woollie heads amongst which come forth small white flowers like those of *Trifolium flore albo* but smaller, and have allwaies close adioyninge underneath each flower three broad hairy scales and three leaves growinge thereon like the former on a verie short footstalk.

|| The Seeds of this Hares foot were gathered in Baetica by Boëlius, and by him sent to Mr. *William Coys*, who hath ever since carefullie preserved the plant, and imparted seeds to me in Anno 1620.—*MS.* f. 106.

Cnicus pratensis Willd.

Carduus bulbosus Monspelliensium.

n. d.

Hath at the first manie leaves spread abroad upon or neare the ground greene above & somewhat white underneath and cottonie

about two foot longe, sometimes parted or devided into manie parts, sometimes into verie fewe, sett with many small weake prickles in fashion like the leaves of *Carduus viarum*, amongst which riseth uppe one stalke somewhat wollie champhered or straked and devided into branches, whereon growe leaves like the former but much shorter and smaller, on the topps of the branches growe round buttons or hedds, not so bigge as an olive sett full of small scales with verie short innocent pricks, forth of which growe abundance of small purple cheives, and no borderinge flowers like those of *Carduus acaulis Septentrionalium* but lesser, which past the seed followeth inclosed in downe like other thistles. The rootes are bulbus after the manner of *Asphodelus*, the old ones yerelie dieinge, and yonge succeedinge, as doe the roots of *Oenanthe apii folio*.—*MS.* f. 124.

[It is a great pity that no locality is given for this plant. If Goodyer found it growing wild this would be the earliest record for Britain. The plant is now confined to the county of Wiltshire, whence it was recorded for the first time in 1813. Smith.]

Carlina lanata L.

Acarna flore rubro.

11 Augusti 1621

The stalk is round upright, straked full of white within, about 5 foot high devided into many branches espetiallie neare the root; under which growe longe broad leaves deeply notched by the sides, and sett with very sharpe pricks, very full of milk white strakes, smooth above somewhat hairie on the vaines or sinewes underneath: on the topps of the stalks and branches growe 3, 4 or 5 small hedds not farr apart, not fullie an ynch thick full of gentle short crooked pricks by the sides, with purple cheives at the toppe closely compact together, ech havinge growinge close underneath him 3, 4 or more prickly leaves with reddish vaynes, the seed is greate, broad at the toppe, blackish without, with a white kernell within wrapped or inclosed in white downe, of a bitter tast, and is more then twice as bigge, as the seed of *Carduus marie*.

This plant at the first sight is in stalks leaves and purple flowers, verie like to our ladies thistle & is hard to be distinguished from it but by the number of the flowers growinge neare together, and the prickley leaves growinge close underneath them. || Mr. William Coys received the seeds hereof from L'obell by the foresaid name. Mr. Coys imparted of his seeds unto me, Anno 1620.—*MS.* f. 105.

Notobasis syriaca Cass.

Silibum minus flore nutante Boelii.

11 Augusti 1621

This Thistle is in stalkes and leaves much smaller than our Ladies

Thistle, that is to say, the stalkes are round, straked, somewhat woolly, with narrow skinny prickly edges three or foure foot high, divided into many branches, whereon grow long leaves, deeply divided, full of white milke-like strakes and sharpe prickles by the edges: the flowers grow on the tops of the stalks and branches, forth of small heads, commonly turning downwards, of the bignesse of an Olive, set with very small slender sharpe pricks, containing nothing but small purple chives, spreading abroad like those of *Iacca*, with some blewish chives in the middle: the seed followeth, inclosed in downe, and is small and grayish like the seed of other Thistles, but it is as clammy as Bird-lime. || The whole plant perisheth at Winter, and reneweth it selfe by the falling of his seed. I finde not that this is written of. It was first gathered by *Boelius* in Spaine, and imparted unto Mr. *William Coys*, who friendly gave me seeds thereof.—*MS.* f. 105; *Ger. emac.* 1627.

Woolly Thistle. *Cnicus eriophorus* L.

Carduus Eriocephalus, *Corona fratrum quorundam*. 13 Augusti 1621
It is the 6th in *Ger.* p. 900.

This thistle hath many leaves at the beginnunge spread on the ground bigge longe very much devided very pricklie white underneath but greene above and somewhat rough ech leafe havinge as it were fower rowes of small leaves betwene which groweth uppe a stalk 3 or 4 foot high somewhat woollie thick straked devided into many branches sett with leaves like to those which spread uppon the ground but lesser uppon the topps of the branches growe greate hedds with many thornie prickles and so cumpassed about or fraught with woollines like Spiders webbs that the prickles doe only a little appeare, the hedds openinge themselves comes out the flowers consisting of many purple cheives, whereunto succeedeth the seed inclosed with downe, shininge somethinge longe as in many other thistles. The roote is longe with many little thredds, above an ynch thick Russett without. It seedeth not till two or three yeres after the sowinge, and most comonly perisheth after it hath borne seed.

|| I found this wild neare London highwaie on the east parte of Haliborne in Hampsheire, 1617. And also in the highwaie neare Abington leadinge towards Oxford the 2 of July 1622.—*MS.* f. 105.

[See also under 29 June 1621.]

Erodium gruinum Willd.

Geranii Bocticae, species *Boelii*.

14 Aug. 1621

This hath at the beginning many broad leaves, indented about

the edges, somewhat divided, like those of *Geranium Creticum*, but of a lighter greene colour, and smaller: amongst which grow up many round hairy kneed trailing branches, divided into many other branches, bearing leaves like the former, but smaller, and no more divided. The flowers are smal like those of *Geranium Moschatum*, but of a deeper reddish colour, each flower having five small round topped leaves: after followeth small long hairie seed, growing at the lower end of a sharpe pointed beak like that of *Geranium Moschatum*: the whole plant perisheth when the seed is ripe.

|| *Boelius* a Low-countray man gathered the seeds hereof in Bœtica a part of Spaine, and imparted them to M^r. *William Coys*, a man very skilfull in the knowledge of Simples, who hath gotten plants thereof, and of infinite other strange herbes, and friendly gave me seeds hereof, and of many other, Anno 1620.—*MS.* f. 106; *Ger. emac.* 1626.

[This species is probably the *Geranium Alceae vesicariae foliis* or Venice Mallow-leafed Cranesbill concerning which Parkinson wrote, 'This and the third among a number of other seeds were brought me by Guillaume Boel which he gathered in Spaine upon my charge; however Mr. Goodier getting the seeds from Mr. Coys, caused it and divers other things to be published in his name: notwithstanding I told him the charge was mine that procured it and many other'. *Theatrum*, p. 707.]

Hedysarum humile L.

Hedysarum clypeatum.

14 Aug. 1621.

Hath at the first many smooth large leaves, lyeinge on or neare the ground, ech leafe beinge made comonly of 7 or more leaves, usuallie growinge on a cornered straked midle ribbe three one against another, and the biggest at the toppe, sometimes one or 2 growinge by them selves, ech leafe beinge about an ynch broad and 2 ynches longe, and are proportionable round both at the toppe & towards the midle ribbe or footstalk; amonge which from one root growe uppe 4, 5 or more round greene pliant straked stalks, leaninge towards the ground, of a reasonable bignes about three foot high bearinge by longe distances leaves like the former; the flowers growe at the topps of y^e stalks on very longe footstalks in greate spikes, of a delicate redd color, in fashion like those of broome, after followeth rough round flatt burrs, growinge likewise in spikes 4, 5 or more fastened or growinge forthright one uppon another, or one at the toppe of another after a strange fashion, ech burr beinge verie like in fashion and bignes to the burrs of *Cynoglossum* or Hounds tongue, everie burr inclosinge one small seed. The root [MS. ends abruptly].—*MS.* f. 106.

Diotis maritima Cass.

Gnaphalium marinum.

20 Augusti 1621

Gerard pag: 515. The 1 & 2 are both one.

The root is longe slender yealowish woodie and groweth deepe in the sande or gravell, and sendeth forth many small round branches and those againe devided into other branches about half a foot longe, besett with small short blunt topped leaves growinge all alongest the stalk very thick together, without footstalks. The whole herbe both branches & leaves are covered over with a thick white delicate soft cotten like *Gnaphalium Americanum*, called in English 'live ever' much whiter softer and more cottonie then anie of the other Cotton-weeds. Of a bitter tast and pleasant smell, somethinge like to sea wormewood. || The flowers I observed not.

I found one plant hereof growinge on the seashoare on the south parte of the Iland of Haylinge in Hampsheire, the 20th of August 1621. And brought it into my garden where the winter followinge it perished.—*MS.* f. 103.

Sea Heath. *Frankenia laevis* L.

An Polygoni marini species.

20 Augusti 1621

This hath one verie longe whitish root of the bignes of a wheate strawe, sett here and there with small threeds, creapinge farr into the sand or sea baich, which a little within the soile sendeth forth manie weake smooth greene branches sometimes devided into other branches three or fower ynches longe above the grounde, bearinge many smooth grasse greene short small leaves; somewhat round yet sharpe at the toppe, a little resemblinge *Rubia minor flore rubro*. There groweth at the topps of the branches very small whitish flowers¹ contayninge 5 little round topped leaves a peece, no bigger then a pinns head, which perished without bearinge any seed at all.

|| This I found on the sea shoare in the west parte of the Iland of Haylinge in Hampsheire and in other places by sea likewise. I brought it into my garden where it flowred as aforesaid about the midle of May.—*MS.* f. 103.

Rhagadiolus edulis Gaertn.

Hieratium Narbonense falcata siliqua L'obelij.

22 Augusti 1621

Ger. p. 285. (7).

The stalks are round hairy straked a cubite high or higher, devided into many branches: whereon growe broad rough greene leaves, very bluntly indented about the edges; the flowers are small and

¹ Flo: in horto 21 Majj.

yealowe like those of *Lampsana*. The seed is small browne somewhat longe, hairie at the toppe, not inclosed in downe, but within small heads, made of small short crooked hairie sharpe pointed husks, not exceedinge half an ynch in lenght, which open when the seed is ripe and spread abroad like larkes clawes starr fashion, and not before. The root is small hairie, full of milkie iuyce, as is also the whole plant, and perisheth when the seed is ripe.—*MS.* f. 101; *Ger. emac.* 1625.

Rhagadiolus edulis Gaertn.

Hieracium stellatum Boelij.

22 Aug. 1621

This plant is in round, hairy, straked, branched stalks, and longe, rough, blunt indented leaves like to *Hieracium falcatum*, but scarce a foot high: the flowers are also yellow three times smaller: which past, there succeed long crooked slender sharpe pointed cods or huskes, neere an inch long, spreading abroad, star-fashion, wherein a long seed is contained: this hath no heads or woolly down like any of the rest, but onely the said crooked coddess which doe at the first spread abroad. The root is small, threddie, full of milkie iuice, as is also the whole plant and it perisheth when the seed is ripe.—*MS.* f. 101; *Ger. emac.* 1625.

Tolpis barbata Gaertn.

Hieracium medio nigrum flore maiore Boelij.

22 Aug. 1621

This hath at the first spreading upon the ground many long narrow, green, smooth leaves bluntly indented about the edges, like those of *Hieracium falcatum*, but smaller: amongst which rise up three, foure, or more, small, smooth, straked round stalks, divided into other branches, which grow longer than the stalks themselves leaning or trayling neere the ground: the flowers grow on the tops of the stalks, but one together, composed of many pale yellow leaves, the middle of each flower being of a blackish purple colour.—*MS.* f. 101; *Ger. emac.* 1625.

[In last line but one the *MS.* has 'yealowe flowers', doubtless rightly corrected to 'yellow leaves' by the editor.]

Tolpis umbellata β *minor* Lange.

Hieracium medio nigrum flore minore Boelij.

[22 Aug. 1621]

This is altogether like the last before described in stalkes and leaves: the flowers are also of a blackish purple in the middle, but they are three times smaller.—*MS.* f. 101; *Ger. emac.* 1625.

Hieracium intybaceum L.

Hieracium intybaceum.

[22 Aug. 1621]

This sendeth forth from one small root many rough leaves like

those of wild Cicorie very much jagged and devided even to the midle ribbe amongst which rise uppe many short round hairy straked browne redd little stalks a foot high or higher devided into one or two branches bearinge a fewe leaves lesser then the former at the topps of the branches growe greate doble flowers (bearinge many more leaves then Clusius spakes of) sometimes single only havinge the outer border of leaves of a pleasant reddish purple color. The seed is very longe & browne with much white downe at the toppe inclosed in heads made of crooked husks or coddts like those of *Hieratium falcatum*.—*MS.* f. 101.

Hieracium andryaloides Vill.

Hieracium lanosum.

[22 Aug. 1621]

There groweth from one root three, foure or more round upright soft cottonie stalks, of a reasonable bignesse, two foot high, divided into many branches, especially neere the top, whereon groweth at each division one broad sharpe pointed leafe, divided into corners, and very much crumpled, and also very soft cottonie and woolly, as is the whole plant: the flowers are small, double, of a pale yellow colour, very like those of *Pilosella repens*, growing clustering very many together at the tops of the stalkes and branches, forth of small round soft cottonie heads: these foure¹ plants grew from seed which I received from M^r. Coys, 1620, and I made these descriptions by the Plants the 22. of August, 1621.—*MS.* f. 101; *Ger. emac.* 1625-6.

Androsace maxima L.

Androsace altera Mathioli. *Ger.* p. 425.

23 Aug. 1621

Hath many leaves spread abroad uppon the ground like those of plantaine but lesser with three sinewes, of a pale greene color, notched by the sides, full of iuyce, in tast somewhat sharpe: amongst which rise uppe six sometimes more small stalks a span high of an herbie color sometimes purple, naked somewhat hairie, which in the verie toppe hath a little crowne made of 5 smalle leaves hairy and notched like the lower leaves but smaller from where growe forth 5 or more little footstalks bearinge a little herbie hairie huske made also of 5 leaves notched also by the sides, which hath a small white flower in the midle devided into five parts.—*MS.* f. 102.

Bartsia Odontites Huds. var. *alb.*

Euphrasia 2 Dod. flo. albo.

24 Aug. 1621

Apud Bellmere pond.—*MS.* f. 53 v.

¹ The *MS.* has '6', including the two descriptions not printed in *Ger. emac.*

Climbing Fumitory. *Corydalis claviculata* DC.

Fumaria clavicularis donatis. Phyto 245 (4). 30 Aug. 1621

At Southsea Castle in flower 30 Augusti 1621.—*MS.* f. 53 v.

? *Lapsana Zacintha* L.

Cichorium verrucatum. 1 Sept. 1621

Hath at the first many longe jagged greene leaves like those of Cicorie but smaller, a little hairy & very bitter, amongst which springe uppe round straked stalks 2 or 3 foot high, hairy belowe, smooth above, devided into branches, bearinge leaves on the lower parte like the former, but smaller towards the toppe. The flowers are like those of Cicorie but very small, yealowe & growe on very short footstalks, forth of the bosomes, on the sides, and on the topps of the branches, which turne not into downe but into hard hedds, with a little tuft at the toppe, composed commonly of eight knotts, corners or bunches sett orderly round, wherein is the seed.

The root is white and short not much unlike that of *Rapunculus*, full of small branches & little thredds. || The whole herbe perisheth when the seed is ripe.—*MS.* f. 102.

Scorpiurus subvillosa L.

Scorpioides multiflorus Boelii. 1 Sept. 1621

This Plant is in creeping branches and leaves like the common *Scorpioides bupleuri folio*: the flowers are also alike, but a little bigger, and grow foure or five together on one foot-stalke: the cods are rougher, and very much turned round, or folded one within another: in all things else alike.—*MS.* f. 102; *Ger. emac.* 1627.

Scorpiurus vermiculata L.

Scorpioides siliqua crassa Boelii. 1 Sept. 1621

This is also like the other in creeping branches and leaves: the flowers are something bigger than any of the rest, and grow not above one or two together on a foot-stalk: the cods are crooked, without any rough haire, yet finely checkquered, and seven times bigger than any of the rest, fully as big as a great Palmer-worme, wherein is the difference: the seed is almost round, yet extending somewhat in length, almost as big as small field Peason, of a browne or yellowish colour. This also perisheth when the seed is ripe.—*MS.* f. 102; *Ger. emac.* 1627.

Sea Heath. *Frankenia laevis* L.

Polygonum alterum pusillo vermiculato, serpilli folia Penae. *Ger.* 453 (3). 3 Sept. 1621

Hath many small round smooth hard woodie branches, somewhat

reddish, traylinge on the ground 9 ynches or a foot longe : whereon by short distances on small ioynts grow tufts of very small short blunt topped smooth greene leaves, in a manner round, like those of the smallest time, but much smaller and without smell, devidinge it self at the bosomes of those leaves into small branches, on the topps of which branches growe small flowers, one flower on a branch and no more, consistinge of fower small round topped leaves a peece, of a faint or pale purplish color. I observed no seed. The root is woodie blackish without, very bitter with some tast of heate and groweth deepe into the ground. The leaves are nothinge so full of iuyce as Aizoon.

I found it flowringe the 3 of Sept. 1621 on the diches bancks at Burseldon Ferrey by the seaside in Hampsheire.—*MS.* f. 89 and *Ger. emac.* 567.

Glasswort. *Salicornia herbacea* L.

Kali album. Dodo. p. 81. minus, Adversar. p. 170. 3 Sept. 1621
Ger. hath it not.

Hath many small round straked branches sometimes standinge upright, sometimes traylinge on the ground, divided into other smaller branches, sett full of small longe narrowe whitish greene leaves, very full of saltish iuyce. The bosomes of the leaves are thick fraught with very small bunches, husks or little buttons, which opened there appeare very small short pale yealowe cheives, which are the flowers. The seed followeth which is [a blank space]. The root is verie small white threddie and perisheth at winter. This plant doth at the first viewe before you come very neere it appeare like yonge broome.—*MS.* f. 111.

Nigella sp.

Nigella flore albo pleno. [3 Sept. 1621]
is the fourth in Ger. p. 925 accordinge to the description.—*MS.*
f. 111.

Nigella Damascena L.

Nigella multiplex. 5^{to} Sept. 1621
Melanthium Damascenum flore pleno Clus.

The doble damaske *Nigella* hath small round smooth tough stalks, divided into verie many branches, whereon growe the leaves which are exceedinge small like thredds, verie finelie iagged and of a darke greene color : at the toppe of ech stalk and branch groweth one faire doble flower of a pale or whitish blewe color, and close underneath ech flower groweth five iagged leaves altogether like those on the stalks. The seed is inclosed in a head like the

former and is [a blank space]. The root is small and yealowe with some threds. The whole plant (except the doblenes of the flowers) is like the single Damaske Nigella.—*MS.* f. 111.

Nigella hispanica L.

5^{to} Sept. 1621

Nigella elegans ex Hispania.

This beutifull *Nigella* hath small round straked upright stalks, devided into branches, whereon growe iagged leaves of a pale greene color, whose iaggs are not so broad as those of *Nigella flore albo pleno*, nor so narrow or small as those of *Nigella Damascena*. At the toppe of ech stalk and branch groweth one greate pleasant flower, bigger then anie of the other sorts of *Nigella*, made of 5 sharpe pointed wrinckled leaves, everie leafe beinge about an ynch both in lenght and bredth, of a beutifull purple color above, and of a whitish greene with a little shewe of purple underneath. In the midle of the flower groweth the head havinge sometimes but 7 or 8 most comonlie 12 or 13 hornes, at the first small of a deepe murrey or obscure browne redd color, about which groweth manie cheives of the same color, next and close above the leaves of the flower, there groweth spread abroad 8 small forked leaves of a blewe color, with a reddish line crossinge them at the first, and afterwards havinge a white line close adioyninge, and one small short pointell appearinge neare the midle of ech of those leaves. The leaves of the flowers beinge readie to fall away, the heads appeare greater, and are rough sett as it were with fine redd spotts, and the hornes are wound or turned round in the end.—*MS.* f. 111.

[Both of these descriptions of *Nigellas* were sent to Johnson but were not included in the Herbal, cf. p. 1085.]

Convolvulus purpurcus L. var.

Convolvulus coeruleus Bryoniae nigraefolio.

7 Sept. 1621

flos Noctis. non script.

Hath manie small weake round hairie browne redd branches, growinge from one root, windinge wrappinge and turninge them selves against the sunne, round sticks or poles that are sett by them for that purpose: whereon by certaine distances growe greate broad leaves, in a manner round, yet picked at the toppe, without anie corners like Ivie leaves, verie like those of *Bryonia nigra* but rounder, somewhat rough above and smooth underneath. Forth of the bosomes of the leaves growe longe slender hairie footstalks, on the toppe whereof growe 2 or 3 most beutifull flowers, not flowringe all at a time but one after another, those that will open in the morninge make some small shewe overnight, onlie wound together, and not of half their growth, erlie in the morninge they appeare in

their full length, but ioyned close together with 5 corners, which after in a short time open and are round like a little bell, like those of white Bindweed, but of a delicate Azure or as it were a color of blewe and redd mixed together, with five straight strakes or lynes in the inside like redd darke colored crimson velvet. This glorious shewe continueth but awhile, for towards night the same daie that they open, they beginn to vade and fold themselves in together at the toppe, and never open againe, and the next daie fall quite away. Quaere, whether they do always so.—*MS.* f. 110.

Centaurea salmantica L.

Stoebe Salmantica j^{ma} Clusii foliis Cichorei. 8 Sept. 1621

Hath at the first large leaves about a foote and a half longe spread abroad uppon the ground unorderedly iagged even to the midle ribbe and those iaggs are indented about the edges sawe fashion, but not devided into other iaggs, ech small indentinge endinge with a weake sharpe prickley point verie thick sett with fine softe cotton-like hairines somewhat like the devided leaves of Cichorie amongst which riseth uppe a round straked stalk 5 or six foot high, devided into many branches, of a browne reddish color, with a softe hairines like that on the leaves towards the root, whereon growe leaves like the former but lesser and lesser upwards on the stalk, the stalk and branches beinge very little or nothings at all hairy towards their topps, and bearinge very narrowe prickley topped leaves almost without haies, not devided like the former, but onlie deepelie indented, endinge with sharpe but weake pricks. At the toppe of ech stalk and branch groweth one head sett on the outside with smooth scales, ech scale endinge with a very small short harmelisse prickle, out of the toppe whereof groweth abundance of pale purple cheives very like the flowers of *Carduus vulgarissimus*, or *Carduus bulbosus monspel.* and not unlike the flowers of *Jacca*, but havinge no larger spreadinge flowers on the borders like it, or like the flowers of *Cyanus*.—*MS.* f. 92.

Anthemis tinctoria L.

Buphthalmum vulgare. primum Matthioli. 9 Sept. 1621

Hath manie small round straked brittle branches cominge from one root about 2 foot high, and those againe devided into other branches covered with a little thinne white cottonlike woollines whereon growe the leaves of a whitish greene color, spread abroad devided into many parts and those small devisions are finelie iagged or minced like the leaves of Tansie but much smaller. On the topps of the stalks and branches growe the flowers somethinge like those

of *Chrysanthemum segetum* but smaller no bigger then Camomill flowers not onlie havinge a greate yealowe ball or dish in the midle but also those small leaves which compasse the ball and likewise of a bright yealowe color and are finely nickt at the toppe and comonlie two, three, fower or five & twentie in number.—*MS.* f. 110.

? *Papaver hybridum* L.

Argemone Pavio.

9 Sept. 1621

Hath manie round hairie stalks or branches proceeding from one root about 3 foot longe whereon growe longe jagged leaves full of white milk-like iuyce as are also the branches. At the topps of the branches growe greate flowers havinge 4 greate round topped leaves a peece of a light redd or crimson color. The heads or seed vessells are small smooth three quarters of an ynch longe, wherein is contayned plentie of exceedinge small seed of a yealowish redd color. The root is small and perisheth when the seed is ripe. This herbe is like *Papaver Rhoëas* in leaves, stalks, flowers & milkie iuyce, but the stalks are longer, the flower much paler & the seed vessels longer.—*MS.* f. 113.

Papaver Rhoëas β *setigerum* Boenn.

Papaver Rhoëas Baeticum.

[9 Sept. 1621]

I cannot discerne wherein it differeth from our comon Papaver Rhoëas.—*MS.* f. 113.

Achillea nobilis L.

Achillea sideritis. Tanacetum minus Dodo: *Achillea* Math: p. 213.
Achillea, sine millefolium nobile Gerardo, p. 915. 9 Sept. 1621

Hath sometimes 4, 5 or more round hard stiffe straked stalks proceeding from one root, and those sendinge forth even from the root to the toppe many small side branches, whereon growe very manie small jagged leaves, devided into many smaller parts, somethinge like those of *Buphthalmum j^m Matthioli* but not so finelie minced or jagged, the small side branches of the leaves verie well resemblinge *Cornu Cervinum Lobelij*, of a whitish greene color and of a stronge smell but not unpleasent, verie hott and bitter in tast. On the topps of the stalks and branches growe large umbells of small white flowers verie like those of Comon yarrowe in smell like the herbe.—*MS.* f. 113.

? *Alchemilla alpina* L.

Heptaphyllum maius. Phyto. 651. (8).

11 Sept. 1621

The stalks growe upright and are round firme somethinge hairie of a reasonable bignes 2 or 3 foot high devided neare the toppe into many small branches, the leaves growe on longe hairie

footstalks out of order, everie leafe beinge usuallie devided into 7 leaves deeplie notched or indented by the sides, and those that growe next the ground are comonlie 4 or 5 ynches longe and neare 2 ynches broad altogether like in color and fashion to the Comon Cinckfoile. The flowers growe on the branches of a pale yealowe color, made of 5 broad topped leaves, with a blunt nick at the toppe of ech leafe, and are manie in number and flower one after another, whereby it continueth longe in flowringe: the seed is small and browne contayned in leafe husks or hedds. The root is short and small brown without and white within with many strings or small roots growinge from the upper parte thereof, and is perennis.—*MS.* f. 113.

Lamium Orvala L.

Lamium Pannonicum 2^m exoticum Clusij, p. xxxviij. 11 Sept. 1621

This strange *Lamium* the first yere after it is sowed, hath leaves almost round, very like nettle leaves, but for the most parte much bigger, with a fine softe hairines, and whilest they are yonge covered with a fatt clammie matter as it were a dewe, indented by the sides, growinge on longe softe hairie footstalks; amongst which leaves, the springe after the sowinge, there groweth uppe hollowe stalks, 3 or 4 foot high, eyther fower square or with six corners (for there are comonlie of both sorts growinge from one root) also covered with a softe hairines, alongest which by certaine distances, on short footstalks, growe sometimes 2 sometimes 3 leaves, allwaies one right against another lesser and shorter then the other: out of whose bosomes growe sometimes 2, sometimes 3 (accordinge to the number of the leaves) small hairie footstalks, an ynch longe bearinge 3 or 4 flowers or more, of the bignes of a pease, which are of a pale yealowish greene color, hollowe within, with a small hole at the toppe, out of which groweth a fewe small short cheives; after the flowers there succeed small sharpe pointed heads almost like those of flax, but more like those of the Common *Scrophularia*, full of very small black seed. The root is crooked with many small hairie threds like that of the nettle, whose stalks perish in the winter sendinge forth other againe at the springe, and sometimes before winter, and flowringe againe as at the springe.

¶ I received the seeds which produced this plant, with many other from the most worthie English Herborist, my very good frend Mr. William Coys often remembred.—*MS.* f. 112.

Clary. *Salvia Verticillata* L.

Horminum silvestre tercium Clusij, p. xxix. ? [13 Sept. 1621]

Hath at the first hairie leaves spread uppon the ground about

five ynches broad and 5 ynches longe, bluntlie indented by the edges, almost round at the toppe: amongst which rise uppe square hairie kneed stalkes about a foot or cubite high, bearinge leaves by cooples one opposite against another, like the former but lesser and more sharper pointed, sometimes with two little leaves like eares growinge on the footstalks of the leaves, of no ill or stronge smell, the stalks devide them selves into braunches at the bosomes of the leaves. The flowers are of a blewish purple colour (lesser then the flowers of Clarie, and scarce bigger then Lavender flowers) growinge in whorles to the verie toppes of the stalks and braunches makinge longe spikes, which comonlie bowe or turne their toppes downewardes, without anie leaves growinge amongst them. The seed [blank]

The roote is greate of a finger bignes growinge downward into the earth, a foot deepe or more, with a fewe side braunches, black without, hard lastinge, yerelie sendinge forth newe braunches.

It groweth not wild in England, the seed hereof I received from Mr. William Coys with many other, in Anno 1620.—*MS.* ff. 11, 115.

[Rough draft of a description sent to Johnson, March 5, 1633, but not acknowledged by him.]—*MS.* f. 11.

Cynosurus echinatus L.

Gramen cristatum Baeticum Boelij.

13 Sept. 1621

Hath many round ioynted stalks growinge from one root, a cubite or 2 foot high, and greate longe leaves like Barley. At the toppe of ech stalk groweth one eare or rather a one sided bunch, an ynch broad and somethinge above an ynch longe, onlie growinge on one side of the toppe of the stalk, so that it maketh as it were half an eare, with small queveringe cheive-like flowers, like those of other grasse, and, verie thick sett with small whitish haire makinge the whole eare appeare like a cattes beard; the seed is small longe and brownish somewhat like that of *Psyllium*, but nothinge so black, inclosed in very small whitish bearded husks. The root is verie small beinge nothinge but verie small white threddie strings, and perisheth when the seed is ripe. || This grasse is not yet described that I find of. The seeds were gathered by Boelius a lowe Contrey man in Baetica a parte of Spaine and given to that diligent preserver of simples Mr. William Coys often with very good cause remembred, who imparted seeds hereof to me in Anno 1620.—*MS.* f. 99.

[Goodyer himself noted that this species was 'Gramen alopecuroides spica aspera' and that it was found 'by y^e adiacent pts of Shepey'. *MS.* 11, f. 133. See p. 59.]

Briza maxima L.

Gramen tremulum maximum. 13 Sept. 1621
an Phyto. p. 10. No. xxxviii. Gramen lupuli glumis Boelij.

This hath many small round smooth ioyned stalks, 2 or 3 foot high, with longe broade smooth leaves like those of otes or barley. at and neare the toppe of ech stalk groweth scatteringlie or somewhat farr apart, about 7 or 8 flatt eares the topps hanginge downwards, about 3 quarters of an ynch longe, and a quarter of an ynch broad, made of 2 rowes of fine thinn scales, curioslie foldinge one within another, verie like those of the comon *Phalaris pratensis* but 8 times bigger, which eares in a close roome you cannott hold so still, but they will wagge and tremble, their footstalks beinge so longe and small, no bigger then small haire. The seed is verie small, flatt and browne, in a manner round, one seed and no more inclosed at the inner end of ech scalie huske. The root is made of small white thredds, and perisheth when, the seed is ripe. || The seeds also of this grasse were given by Boelius to Mr. William Coys by the name of Gramen lupuli glumis, who afterwards in Anno 1620 sent seeds thereof to me.—*MS.* f. 99.

Lactuca virosa L.

Lactuca silvestris vera ingrato odore. 13 Sept. 1621
Hist. lug. p. 547, optima figura. Phyto. p. 202 N^o xii.

This wild lettise hath at the first many broad leaves spread upon the ground, like to garden lettice leaves, whiter on the underside then above, sometimes a foot longe and 5 ynches broad, broadest neare the toppe, with a greate ribbe or sinewe underneath full of sharpe pricks, and many other small branches sinewes full also of little pricks, very little nicked and crisped by the sides, but nothings at all gashed, yet also full of small pricks, amongst which groweth uppe a greate upright round stiff prickley browne reddish stalk, full of white pith like that of the elder, usuallie 5 or 6 foot high, (sometimes 10 foot high as I observed this yere 1621) devided into many parts or branches, on the lower parte whereof groweth leaves like the former, but upwards on the stalk they are smaller shorter without footstalks, gashed or devided with deepe devisions. The topps of the stalks and branches are garnished with manie small yealowe flowers, like in forme and bignes to those of the common lettice, flowringe one after another, which maketh it longe in flowringe: the flowers past there succeed flatt blackish seed like to lettice seed, with downe at the toppe and is caried away with the wind and reneweth itself by the fallinge thereof,

which quicklie groweth and sendeth forth such broad leaves as aforesaid spread upon the ground and remaine greene all the winter, sendinge then forth such stalks as is aforesaid. The root is hard woodie sometimes devided into branches yealowe without & groweth deepe into the earth. The whole herbe and root is full of clammy white iuyce like milk, and of a very loathsome stinkinge smell, which on the stalks sometimes turneth into a yealowish gummie matter.

|| This wild stinkinge lettuce I found wild on the walls and dry bankes of earth at Southampton. Anno 1618.—*MS.* f. 99.

Lactuca agrestis L.

Lactuca agrestis.

13 Sept. 1621

This hath not leaves spread abroad upon the ground like the former, but riseth uppe presentlie with a small round stiffe stalk prickley only belowe 3, 4 or 5 foot high, devided into very manie branches, whereon growe manie pale greene plaine smooth leaves with many pricks on the midle ribbe on the lower side and also by the edges, the sides or edges are somewhat indented but not at all crisped gashed or devided, sometimes 6 or 7 ynches longe and 3 ynches broad or broader, broadest comonly in the midle and narrowe towards the toppe. The flowers are also yealowe like those of the lettuce, and turne into a small grey seed with downe at the toppe which is caried away by the wind, by the fallinge whereof it increaseth and sendeth forth other yonge plants, with short tender stalks, which remaine greene all the winter, and in the springe growe uppe as aforesaid. The root is hard threddie and yealowe without. The whole herbe is also full of white, milk-like iuyce and of little or no smell at all, and perisheth when the seed is ripe.—*MS.* f. 100.

Oenothera biennis L.

Lysimachia virginiana.

13 Sept. 1621

This riseth uppe with a stalk about 3 foot high, which is round straked firme hard brittle full of pith within, reddish neare the root, devided into manie branches, and spotted with very small purple spots: whereon growe the leaves out of order without footstalks 5 or 6 ynches longe and above an ynch broad, smooth sharpe pointed verie bluntlie indented about the edges, with a whitish midle ribbe, hott in tast and bitinge the tonge. The flowers are yealowe and growe forth of the bosomes of the leaves neare the topps of the stalk flowringe upwards by degrees, whereby it continueth longe in flowringe, and growe upon longe tender stems everie

flower havinge 4 broad topped leaves, and a short yealowe pointell not appearinge above the topps of the leaves of the flower, devided in the toppe into 4 parts, everie part beinge bigger then the lower parte of the pointell, and also yealowe cheives growinge from the nailes of the inner partes of leaves of the lenght of the pointell, which flowers are of a strong fulsome smell, and both they and their tender stems fall away, and there groweth uppe greate longe blunt topped, round straked codd, without anie footstalks, makinge a longe spike of codd, wherein is containd much small round wrinkled seed which when it is ripe the codd openeth into 4 or 5 parts at the toppe and the seed falleth forth and at the next springe groweth uppe whereby it mightelic increaseth.—*MS.* f. 100.

Nicotiana Tabacum var. *brasiliensis* Comes.

Petum indicum folio pene obtuso.

13 Sept. 1621

Ye figure in Hist. Lug. p. 1895 resembleth it well. Peti primum genus Clusii, p. ex: 309.

This groweth uppe with a greate round stalk devided into many branches, five or six foot high, verie hairie fatt and clammie; whereon growe greate broad leaves, somewhat round towards the toppe yet endinge with a sharpe point, narrower & crompted towards the stalkes, without footstalkes, imbracinge or growinge 2 or 3 ynches downe on both sides of the stalk, belowe the fasteninge or growinge of the midle ribbe to the stalke, about 20 ynches longe, and above a foot broad, also verie fatt clammie and rough, of a yealowish greene color, of a good savor and verie sharpe tast. The flowers growe on the toppes of the branches, and are longe, hollowe, in manner of a little pipe or bell, broad at the toppe, endinge with corners most comonlie 5 sometimes 4 somewhat blunt not verie sharpe, of a pale or whitish purple colour; which fallen there cometh small longe round sharpe pointed heddes or seed vessels lesser longer and sharper pointed then those of yealowe Henbane, in which is included abundance of exceedinge small blackish redd seeds, much lesser then poppie seed. The roote is thicke, woodie, branched and yealowish. The whole plant perisheth at the first approach of winter, if it be not planted in an earthen pott or other fitt vessell, and putt into a close place to defend it from the iniurie of the cold.

The scedes of this Tabacco I receaved Anno 1620 from my worthie friend, and most diligent observer and preserver of simples Mr. William Coys of North-okington in Essex.—*MS.* ff. 93, 114.

Nicotiana Tabacum var. *fruticosa* Hook. f.

Petum indicum folio Hydrolapathi acuto. 13 Sept. 1621

The stalkes are also round and branched, somewhat hairie and clammiie, the leaves growe also without footstalkes half compassinge the stalkes, yet sometimes exceedinge narrowe towards the stalkes, generally much narrower and longer then the former, comonly 2 foot, 2 foot and a half and sometimes 3 foot longe, and usuallie 6 or 7 ynches broad or broader, not round towards the toppe like the former, but verie sharpe and slender pointed, smoothe greener and nothings so clammiie as the other, also of a sharpe bitinge tast, and are in forme like the leaves of the greate water docke, the flowers are also of a light purple colour, but smaller endeinge with 5 sharpe pointed corners, much sharper then the former. The seed is like the former inclosed in the like huskes, but somewhat sharper pointed. The roote is also like and must be preserved from the cold as the other.

In Anno 1619 I received the seedes hereof from Mr. Anthony Uvedale who that yere intended to plante greate store hereof, and was hindered of his purpose by a proclamation sett forth by Authoritie.—*MS.* ff. 93, 114.

Nepeta tuberosa L.

Cattaria tuberosa radice Boetica Boelii. non scripta.

14 Sept. 1621

Hath 2, 3 or more square upright stalks, somewhat hairie, two foot high or higher, sometimes leaninge towards or lyinge flatt on the ground, devided into many square branches growing allwaies one right against another, whereon growe leaves by couples one right against another, sometimes with short hairie footstalks, but most comonly espetially on the upper parte of the stalk, with very short or no footstalks at all, full of crumpled vaines or sinewes, and large blunt notches by the sides, like those of the comon neppe, but not so broad, of the same pleasant smell, but not so stronge, and but a little hott and bitinge the tonge, sometimes covered all over with a softe hairie cottonie whitenes, as though a white frost laie thereon. On the upper parte of the stalks and branches growe by certaine distances one from another crownetts or whorles of manie small scalie leaves, somewhat reddish at the topps and by the sides, compassinge the stalks: whereof two that are lowest and biggest, and growe allwaies one right against another, amongst which on the upper parte growe many blewish purple flowers (those small scalie leaves and flowers makinge the entier whorle or crownett) in fashion like those of the comon neppe, but neare as

bigge againe, of verie little smell: the severall whorles makinge a spike sometimes of above a foot longe.

The seeds hereof I received from Mr. Coys in A^o. 1620.—*MS.* f. 96.

Nepeta media.—*MS.* f. 96. (? = *Nepeta Cataria* L. var.)

[No description.]

? *Sinapis alba* L.

Sinapis sativum alterum Penae.

[? 14 Sept. 1621]

Adversar, p. 68. *Lob. icon.* p. 277. pte 2^a.

Hath one stalk growinge from the root, which is round hollowe straked hairie or rough 2 or 3 foot high, without knees devided into verie many branches even from the root. The leaves are greate and growe at the devisions of the stalk, with a verie little roughnes, torne and devided into many partes even to the midle ribbe. The flowers growe on the topps of the branches floweringe upwards and are yealow, very like those of comon musterd but bigger, and so like those of comon cherlock that they are hard to be knowne the one from the other, contayninge 4 broad topped shrivelled leaves apeece: after the flowers cometh rough hairie codd's growinge all alongest the upper parte of the stalk and branches, bigger and more spreading abroad then those of Cherlock, half whereof in lenght towards the toppe is broad and flatt like to the point of a speare yet blunt topped, and in the lower parte thereof towards the stalk is contayned 2 rowes of seedes, havinge 3 or 4 round seeds in ech rowe, either of a white grey or reddish color; three times bigger then comon musterd seed, and sometimes neare as bigge as Radish seed, hott and bitinge the tonge like pepper, bunchinge out the codd where they lie, and when they are ripe doe fall out with a light touch leavinge behind on the stalk the midle (?) devision which is a thinn cleare membrane or skinne and the speare like point of the codd's which continueth on a longe time after. The root is small white threddie and perisheth when the seed is ripe, and reneweth it self yerelie by the fallinge thereof.—*MS.* f. 96.

Blue Fleabane. *Erigeron acre* L.

Conyza coerulea acris, C. Bau. *Conyza odorata*. 20 Sept. 1621

Hath at the first many hairie leaves 2, 3 or 4 ynches longe, spread upon the ground without footstalks, verie slender and narrowe belowe, and half an ynch broad towards the toppe, plaine and not indented by the edges, hott in tast and bitinge the tonge, of a reasonable good smell: amongst which rise uppe 2, 3 or more small round browne redd straked hairie stalks, full of white pith within, a foot or a cubite high seldome devided into branches on

the lower pte, whereon growe hairy leaves out of order without footstalks, like the other but much shorter and smaller, about an ynch and a half longe, and a quarter of an ynch broad comonly turninge downewards, with other very small short leaves growinge forth of their bosomes, or els very short stubbed branches. On the upper pte the stalke is devided into many small short branches, bearinge at the toppe of ech branch one little head, made on the outside of many verie slender narrowe hairie stalks comonly reddish, on the upper parte of the head groweth verie manie small narrowe short leaves no bigger than cheives of a pale or light purple color, seldome spreadinge abroade, but growinge upright, which together with the head are about half an ynch in lenth: next within these leaves and verie neare of their lenth even at their first openinge, appeareth a round circle of pale yealowish downe, which in the midle of the flower is of a dark browne color and in the ende spreadeth abroad and contayneth underneath much small yealowish seed, which together with the downe is caried away with the wind. The whole head rubbed betwene the fingers is of a good smell. The root is nothinge but small white thredds and dyeth not at winter. || I first found this herbe growinge wild on the walls at Winchester in Hampsheire, afterwards in other places on dry bancks, but very seldome.—*MS.* f. 95.

Ivy-leaved Toadflax. *Linaria Cymbalaria* Mill.

Cymbalaria Italica.

20 Sept. 1621

Cymbalaria with us in England, where it is sowen runneth and spreadeth on the ground and clymeth and hangeth on walls even as Ivie or Chickweed doth, the branches are verie small round and smooth, limmer and pliant neare like the hampering threeds of *Cuscuta* devidinge it self plentifullie into other branches and sendinge forth other small threddie roots takinge hold therewith on the earth or walls. The leaves growe on long round footstalks and are like those of Ivie smooth and devided into moe corners, the flowers growe also on longe footstalks forth of the bosomes of the leaves like single violettis but smaller, nearer like the flowers of *Elatine* consistinge of 2 leaves, whereof the uppermost is on the upper parte of a light purple color and devided at the toppe into a parte, the lower leafe is three times bigger, of a pale or whitish color with a verie light dash of purple, and the midle or chappe is yealowe, devided belowe into 3 pts, with a small purple taile behind hanginge downewards, which fallen there succeedeth in the place of the flower some little round knappe or button contayninge small seed.

|| I never saw this growinge but in the garden of my faithfull good frend Mr. William Coys in Northokington in Essex, and in my garden at Droxford of seeds received from him in Anno 1618.—*MS.* f. 95.

[The first mention of English *Cymbalaria* is in Coys' Garden-List of 1617, p. 317. It is a pleasure to note that after the lapse of 245 years the original station for this plant in Great Britain should be noted by Gibson, *Flora of Essex*, 1862, on the authority of Edw. Forster, as 'Old wall at Stubbers', and we ourselves found it there in 1921. Parkinson noted the plant 'about Hatfield' in 1640.]

Scabiosa atropurpurea L.

Scabiosa flore rubro. *Scabiosa sexta Indica Clusii.* 8 Oct. 1621

Hath one round tender stalk proceeding from the root, about 3 or 4 foot high, devided into branches ymediately from the root: whereon growe leaves by cooples much devided as it were leaves sett uppon a midle ribbe and are verie narrowe towards the toppes of the stalk and branches, almost smooth yet havinge a little soft hairines as have also the stalk and branches, in forme like those of *Scabiosa minor* sive *Columbaria lobelii*. The flowers growe on the toppes of the stalk and branches like the said *Scabiosa media*, but of a delicate redd color like to redd velvett, with many small cheives, with quaveringe topps, at the first of a light purple, after of a dustie whitish color, ech head beinge composed of many small flowers closelie thrust together, and ech flower is devided into 5 ptes or leaves those that growe on the outside or border of the head are greater and longer then those in the midle, allwaies having close underneath ech head, a rowe or circle of small short narrowe greene leaves growinge starr fashion. The seed

The root [MS. incomplete].

|| This plant I sawe flowringe and bearinge ripe seed in the garden of my most worthie friend and diligent preserver of plants Mr. William Coys, in Northokington in Essex, the 29 of September 1622. Seeds hercof I also received from him in Anno 1620.—*MS.* f. 94.

— ?

Jacea palustris Baetica Boelii.—*MS.* f. 94. 10 Oct. 1621
[No description.]

Centaurea nevadensis Boiss. & Reut. ?

Jacea capitulis hirsutis Boelii. 10 Oct. 1621

This hath many small cornered straked hairie trayling branches growing from the root, and those again divided into many other branches, trailing or spreading upon the ground three or foure foot

long, imploying or covering a good plot of ground, whereon grow hairy leaves divided or iagged into many parts, like the leaves of *Iacca maior*, or Rocket, of a very bitter taste: at the top of each branch groweth one scaly head, each scale ending with five, six, or seven little weake prickles growing orderly like halfe the rowell of a spurre, but farre lesser: the flowers grow forth of the heads of a light purple colour, consisting of many smal flowers, like those of the common *Iacca*, the bordering flowers, being bigger and larger than those of the middle of the flower, each small flower being divided into five small parts or leaves, not much unlike those of *Cyanus*: the seed is small, and inclosed in downe. The root perisheth when the seed is ripe.

|| This plant hath not been hitherto written of that I can find. Seeds of it I received from M^r. *William Coys*, with whom also I observed the plant, 10 October. 1621. he received it from *Boelius* a Low countrey man.—*MS.* f. 94; *Ger. emac.* 729.

Cucurbita Pepo L. var.

Macoeks Virginiani.

10 Oct. 1621

The Virginian Macocke, or Pompion.

This hath rough cornered straked trailing branches proceeding from the root, eight or nine foot long, or longer, and those againe divided into other branches of a blackish greene colour, trailing, spreading, or running amongst the earth, covering a great deale of ground, sending forth broad cornered rough leaves, on great grosse, long, rough, hairy foot-stalks, like and fully as big as the leaves of the common Pompion, with clasping tendrels and great broad shriveled yellow flowers also like those of the common Pompion: the fruit succeedeth, growing amongst the stalkes, commonly not neere the root, but towards the upper part or toppes of the branches, somewhat round, not extending in length, but flat like a bowle, but not so bigge as an ordinarie bowle, beeing seldome foure inches broad, and three inches long, of a blackish greene colour when it is ripe. The substance or eatable part is of a yellowish white colour, containing in the midst a great deale of pulpe or soft matter, wherein the seed lyeth in certaine rowes also, like the common Pompion, but smaller. The root is made of many whitish branches, creeping far abroad in the earth, and perish at the first approach of Winter.—*Ger. emac.* 919.

Citrullus vulgaris Schrad.

Melones Aquatici. The Virginian Water-Melon.

10 Oct. 1621

This Melon or Pompion is like and fully as bigge as the common

Pompion, in spreading, running, creeping branches, leaves, flowers, and clasping tendrels: the fruit is of a very blackish greene colour and extendeth it selfe in length neere foure inches long, and three inches broad, no bigger nor longer than a great apple, and grow alongst the branches forth of the bosomes of the leaves, not farre from the root even to the toppes of the branches, containing a substance, pulpe, and flat seed, like the ordinary Pompion: the root is whitish, and disperseth it selfe verie farre abroad in the earth, and perisheth about the beginning of Winter.—*Ger. emac.* 921.

Basil. *Ocimum Basilicum* L.

Acinos odoratissimum.

11 Oct. 1621

This herbe hath foure, five, or more, foure square hard woody stalkes growing from one root, divided into many branches, covered with a soft white hairnesse, two or three foot long or longer, not growing upright, but trailing upon the ground; the leaves grow on little-short footstalkes by couples of a light greene colour, somewhat like the leaves of Basill, very like the leaves of *Acinos Lobelij*, but smaller, about three quarters of an inch broad, and not fully an inch long, somewhat sharpe pointed, lightly notched about the edges, also covered with a light soft hoary hairnesse, of a very sweete smell, little inferiour to Garden Marjerome, of a hot biting taste: out of their bosomes grow other smaller leaves, or else branches; the flowers also grow forth of the bosomes of the leaves toward the tops of the stalkes and branches, not in whorles like the said *Acinos*, but having one little short footstalke growing forth of the bosome of each leafe, on which is placed three, foure, or more small flowers, gaping open, and divided into foure unequall parts at the top, like the flowers of Basill, and very neare of the likenesse and bignesse of the flowers of Garden Marjerome, but of a pale blewish colour tending towards a purple. The seed I never observed by reason it flowered late. This plant I first found growing in the garden of M^r William Yalden in Sheete near Petersfield in Hampshire, Anno 1620, amongst sweete Marjerome, and which by chance they bought with the seedes thereof. It is to be considered whether the seedes of sweete Marjerome degenerate and send forth this herbe or not, 11 October 1621.—*Ger. emac.* 677.

[See 1620.]

Jerusalem Artichoke. *Helianthus tuberosus* L.

Heliotropium Indicum.

17 Oct. 1621

Flos solis Pyramidalis, parvo flore, tuberosa radice.

This wonderfull increasing plant hath growing up from one root,

one, sometimes two, three or more round green rough hairy straked stalks, commonly about twelve foot high, sometimes sixteene foot high or higher, as big as a child's arme, full of white spungious pith within. The leaves grow all amongst the stalkes out of order, of a light green color, rough, sharp pointed, about eight inches broad, and ten or eleven inches long, deeply notched or indented about the edges, very like the leaves of the common *flos solis Peruvianus*, but nothing crumpled, and not so broad. The stalkes divide themselves into many long branches even from the roots to their very tops, bearing leaves smaller and smaller toward the tops, making the herbe appeare like a little tree, narrower and slenderer toward the top, in fashion of a steeple or Pyramide. The flowers with us grow onely at the toppes of the stalkes and branches, like those of the said *flos solis*, but no bigger than our common single Marigold, consisting of twelve or thirteene straked sharpe pointed bright yellow bordering leaves, growing forth of a scaly small hairie head, with a small yellow thrummie matter within. These flowers by reason of their late flowering, which is commonly two or three weeks after Michaelmas, never bring their seed to perfection, & it maketh shew of abundance of small heads neere the tops of the stalkes and branches forth of the bosomes of the leaves, which never open and flower with us, by reason they are destroyed with the frosts, which otherwise it seemes would be a goodly spectacle. The stalke sendes forth many small creeping roots, whereby it is fed or nourished, full of hairie threddes even from the upper part of the earth, spreading farre abroad: amongst which from the maine root grow forth many tuberous roots, clustering together, sometimes fastened to the great root it selfe, sometimes growing on long strings a foot or more from the root, raising or heaving up the earth above them, and sometimes appearing above the earth, producing from the increase of one root, thirty, forty, or fifty in number, or more, making in all usually above a pecke, many times neere halfe a bushell, if the soile be good. These tuberous roots are of a reddish colour without, of a soft white substance within, bunched or bumped out many waies, sometimes as big as a mans fist, or not so big, with white noses or peaks where they will sprout or grow the next yeare. The stalkes bowed downe, and some part of them covered over with earth, send forth smal creeping threddie roots, and also tuberous roots like the former, which I have found by experience. These tuberous roots will abide alive in the earth all winter, though the stalkes and rootes by the which they were nourished utterly rot and perish away, and will beginne to spring

up againe at the beginning of May, seldome sooner.—*MS.* f. 116;
Ger. emac. 753-54.

[For Goodyer's descriptions of *The Place* and *The Vertues* which follow see p. 24, also p. 109.]

Yew. *Taxus baccata* L.

Taxus glandifera baccifera.

19 Dec. 1621

The Yew bearing Acornes & berries.

The Yew tree that beareth Acornes and berries is a great high tree remaining alwaies greene, and hath usually an huge trunk or body as big as the Oke, covered over with a scabbed or scaly barke, often pilling or falling off, and a yong smooth barke appearing underneath; the timber hereof is somewhat red, neere as hard as Box, universally covered next the barke with a thicke white sap like that of the Oke, and hath many big limmes divided into many smal spreading branches: the leaves be about an inch long, narrow like the leaves of Rosemary, but smooth, and of a darker greene colour, growing all alongst the little twigs or branches close together, seldome one opposite against another, often having at the ends of the twigs little bunches composed of many leaves like the former, but shorter and broader, closely compact or ioyned together: amongst the leaves are to be seene at all times of the yeare, small slender buds somewhat long, but never any flowers; which at the very beginning of the Spring grow bigger and bigger, till they are of the fashion of little Acornes, with a white kernell within: after they are of this forme, then groweth up from the bottomes of the Acornes a reddish matter, making beautiful reddish berries more long than round, smooth on the out side, very clammy within, and of a sweet taste, covering all the Acorne, onely leaving a little hole at the top, where the top of the Acorne is to be seene: these fallen, or devoured by birds, leave behinde them a little whitish huske made of a few scales, appearing like a little flower, which peradventure may deceive some, taking it to be so indeed: it seemes this tree, if it were not hindred by cold weather, would alwaies have Acornes and berries on him, for he hath alwaies little buds, which so soone as the Spring yeelds but a reasonable heate, they grow into the forme of Acornes: about the beginning of August, seldome before, you shall finde them turned into ripe berries, and from that time till Christmase, or a little after, you may see on him both Acornes and red berries.—*MS.* f. 119; *Ger. emac.* 1370.

Male Yew. *Taxus baccata* L. ♂.

Taxus tantum florens. The Yew wth only flowers.

The Yew which only beareth flowers and no berries, is like the

5

Taxus tantum florens. The Yew wth only flowers. //

The Yew wth only berries, flowered and no berries, is like the Yew wth trunk timber bark and leaves. But at the beginning of November or before, the tree doth beginne to be very thick set or fraught. on the lower side of the twigs or little branches, are small round buds (which were at first) and of the old of Rast, and doth containe all the winter here about the beginning or middle of February upon the open at the toppe (which is forty) one shall find winter, it is the longer the Yew is, and doth into many set, or greenish buds about the toppe, w^{ch} many times doth spring like flowered. If the rods of the Yew be good, they are of the same and the from the flowered an abundance of buds. The buds of the flower containe on the tree till about Easter. The Yew buds are full away, and the buds of the Yew are round buds, as at Easter.

The place.

The Yew is very common in the English, in the Yewes of the Yew of good plants of the Yew. Growing in the English hills, and in the young Yewes of the Yew of the Yew.

The time.

The time is reported in the Description. Dec. 19. 1621. John Goodyer. 120 *

GOODYER'S DESCRIPTION OF THE YEW WITH ADDITIONS IN JOHNSON'S HAND * *

other in trunk, timber, bark, and leaves; but at the beginning of November, or before, this tree doth beginne to be very thick set or fraught on the lower side or part of the twigs or little branches,

with small round buds, verie neere as big, and of the colour of Radish seed, and do so continue all the Winter, till about the beginning or middle of Februarie, when they open at the top, sending forth one small sharpe pointell, little longer than the huske, divided into many parts, or garnished towards the top with many small dusty things like flowers, of the colour of the husks; and if you shall beate or throw stones into this tree about the end of Februarie, or a good space after, there will proceed and fly from these flowers an abundance of dustie smoke. These dusty flowers continue on the trees till about harvest, and then some and some fall away, and shortly after the round buds come up as aforesaid.

The Place.

Theis trees are both very comon in England, in Hampsheire there is good plentie of them growinge wild on the Chalke hills, and in Church yards where they have byn planted.

The Time.

The time is expressed in their descriptions.—*MS.* f. 119; *Gcr. emac.* 1370-71.

[The first record of the Male Yew in Hampshire.]

Calathian Violet. *Gentiana Pneumonanthæ* L.

Pneumonanthæ.

[After 9 Nov. 1621]

hath a small round stalk 3 or 4 ynches high, on wch growe small narrow leaves, half an ynch long, which are curvd in the midle very like y^e leaves of savery, sett thick but orderly by cooples one opposite ye other on y^e toppe of ye stalk groweth one flower an ynch longe of the fashion of a bell, devided at y^e brim into 5 sharpe pointed corners of a perfect blewe color, except 5 plates or strakes wh. are only to be seene before y^e flower openeth, extending from y^e bottome of ye flower to corner, which are not so pleasant a blew color, the seed I observed not. Ye root is small divided at ye upper parte into a fewe small (yet of a sufficient bignes for ye stature of y^e herbe) yealowish lynner branches or fibres. Both leaves and rootes are bitter, and ye root more bitter than the leaves & bitinge ye tonge.—*MS.* f. 9 v.

Marsh Cinquefoil. *Potentilla Comarum* Nutt.

Quinquefolium palustre.

[After 9 Nov. 1621]

The stalks are lynner bendinge pliant round smooth a foot long as bigg as a wheate straw of a reddish color towards ye root, ioynted, at ech ioynt groweth a [leafe] foot-stalk which whooly beclippeth the knott or ioynt, on whose toppe groweth five leaves of

ye fashion of other cinckfoyles nicket about ye edges, of a mealish greene colour above and whitish underneath, ye 5 leaves not growing on the very upper parte of the footstalk like ye other cinckfoyles, but 2 of them growinge lower about a quarter of an ynch from y^e 3 which growe at the extreme part, y^e flowers I observed not. The root creapeth in y^e water & mire, besett with thousand of very small haire, thicker & smaller then the haire of ones head.—*MS.* f. 10 v.

[Roughly drafted descriptions on back of Laurence Davis' letter of 9 Nov. 1621.]

— ?

[c. 1620–1622.]

round buttons or knapps, as bigge or little bigger then the pease wherein in each button is 2, 3 or 4 3-winded seeds almost as bigg as Radish seed. The root is small white, single and groweth downright, with a few threddie shoots with side branches. Both herbe and root doe perish at winter.—*Fragment of a description, MS.* f. 11 v.

'Capon's Tail Grass.' Not *Festuca Myurus* L.¹

Gramen ἀλεκτρούουρος. Alectryonurum. 10 Feb. 1622

[Mentioned with date but without locality.—*MS.* f. 54.]

Gramen murorum spica longissima.

'I cannot omit this elegant Grasse, found by M. *Goodyer* upon the wals of the antient city of Winchester, and not described as yet by any that I know of. It hath a fibrous and stringy root, from which arise leaves long and narrow, which growing old become round as those of *Spartum* or Mat-weed: amongst these grassie leaves there growes up a slender stalke some two foot long, scarce standing upright, but oft times hanging down the head or top of the eare: it hath some two ioints, and at each of these a pretty grassy leafe. The eare is almost a foot in length, composed of many small and slender hairy tufts, which when they come to maturitie looke of a grayish or whitish colour, and do very well resemble a Capons taile; whence my friend, the first observer thereof, gave it the title of *Gramen Αλεκτρούουρος*, or Capons-taile Grasse: by which name I received the seed thereof, which sowed, tooke root, and flourishes". (Johnson)—*Ger. emac.* pp. 30, 29.

¹ The correctness of the determination of this grass as *Festuca Myurus* L. by Druce is doubted by D. Stapf.

Curled Parsley.

Apium crispum.

17 Feb. 1622

At Idsworth, 17 Feb. 1622.—*MS.* f. 51 v.

[Hill in 1574 printed the following prescription for Parsley. 'If you will have the leaves of the Parcely grow crisped, then before the sowing stuffe a tennis ball with the seedes, & beat the same wel against the ground, wherby the seedes may so be a little brused, and then sowe them in the ground, or when the Parcelye is well come up go over the bed with a waighty roller, whereby it may so presse the leaves downe, or els treade the same downe with thy feete.'" Th. Hill, *Arte of Gardening.*]

Calamagrostis Epigejos Roth.

Calamagrostis.

27 Apr. 1622

This sedge sendeth forth many 3 cornered straked stalkes, about 2 foot high, beareinge at ye toppe a spike or eare about 3 ynches longe divided into 12 or 13 lesser scaly eares, and those againe into smaller, the whole spike not spreading abroad, but growinge neare together, so that the thicknes is not above an ynch, of a brownish color before the flowers come forth, which appeare are nothings but an infinite many of small dustie things like cheives, at ye first opening whitish afterwards yealowe like ye flowers of other grasses.

[The seed is small contayned in chaffie scales, of a brownish chestnutt color, neare 3 cornered & broad belowe and sharpe pointed without any manifest tast. 8 Julii 1622.]

The leaves are narrowe about 2 or 3 foot longe growinge ymediatelic from ye root, as it were 3 cornered, and very rough espetially if you slide your fingers from their topps downwards, and so sharpe that they cutt one's fingers even as a knife, as doth also y^e stalk; which usually is without leaves, yet sometimes hath one very small one growinge close belowe ye spike, neare a foot longe. The rootes are infinite of the bignes of ye rootes of *gramen caninum* (couch gras) or bigger of a reddish color without, closely mattinge together, and gathering the mudd or dirt amongst them. Growinge downeright of a greate lenth, and makeinge a greate stronge tuft, no herbe or gras like it, for they are so stiffe & stronge, that growinge in ye midst of water, a man may goe on them & steppe from tuft to tuft, ye water & mud beinge of a good depth betwixt them.—*MS.* f. 121.

Oak. *Quercus robur.*

Cachrys quercus.

28 Apr. 1620 & 9 May 1622

The Cachryes¹ are conceived eyther in a budd with the leaves, or

¹ The term *cachrys* is one that appears to have been used for the young cones

in a budd alone by them selves. Those that are conceived within ye leaves come forth together with them, and growe at the ioynt stoppe or knott betweene the last yeres twigge and the newe shoot, sometimes on the lower part of the newe shoot, those that are conceived by themselves, soe come forth without any leaves or shoot, and these growe by the sides of y^e last yeres shoot. This cachrys is composed of small yealowish cruded bunches or clusters growinge a little asunder, on a footstalk about 2 ynches longe, 3 or 4 forth of one budd, and some wither and fall away, seldome continuinge above a weeke or two. When there are plentie of these cachryes, it is a signe there will followe good store of Acornes.

9 Maii. Forth of the bosomes of the leaves on the newe shoot come forth small foot stalks on y^e toppes whereof growe 2 or 3 or more very small redd flowers: ech flower beinge no bigger then a small pinnes head, and devided into 3 ptes (not worth the name of leues) at the toppe, in the place whereof cometh uppe the Acornes.

4 Maii. In the springe when the leaves first beynn to come forth there often groweth from the topps and sides of the last yeres shootes certaine swellinges as bigge as little aples, not altogether round but bunched out here and there, reddish on that part towards the sunne, contayninge an austere or harsh spongie matter of conifers, for catkins, and for the winter buds of the deciduous trees. Though much used by the early botanists, it appears to have dropped out of use at the end of the seventeenth century, and is not even included in many later lists of technical words. Fuchs in 1542, in his *Explicatio* of difficult words, defined the term as follows:

Cachryes sunt oblonga panicularum modo nucamenta, quae squamatim compacta propendent è ramis. Crescunt hyeme, vere dehiscunt in flavescentes squamulas, et folio prodeunte, decidunt: qualia in abiete, picea et aliis permultis videre licet. Plinius pillulas nominat.—Fuchs, *Hist. Stirp.* f. β 3.

'The Birch, the Nut, the Walnut, and the Plane Tree have on them things in Greek called Cachryes in English Catkines or Catstailes, if I mistake not which are there the most part of the winter. They are of a burning quality in Physick' (Coles, *Art of Simpling*, 1656). And this is the meaning given in the *New English Dictionary*. According to Parkinson (1642) Cachrys is the fifteenth 'Excrecence of the Oke'. It was borrowed from Theophrastus iii. 7 to mean 'a round conception or gathering together of leaves, growing betweene the last yeares shoote, and the young bud for the next to come'. And they are borne on 'the Firre tree, Larch, Pitch, Line, Nut, and Plane trees' as well as on the Oak, all of which 'doe beare a Cachrys after the leaves are falne, abiding on all the winter'. Elsewhere he defines it as 'a scaly tuft of leaves growing in winter, and falling away, say some, in the Spring: but others think that it is but the bud, which spreadeth into branches with leaves after Winter, when the Spring is come on'. (Park. *Theatrum*, 882).

within, and towards Autumne hath eyther maggotts, flies or some livinge creature within it, this hangeth many times two yeres on the twiggs, and then also have had maggotts within them as I have seene.

Gallae foliorum quercus.

There is often to be seene in Autumne on that part of the leaves which is next the ground a round pill or ball without a footstalke, sometimes 8, 10 or more on one leafe, reddish on that parte next the sunne, smooth on the outside, or with little sharpe extuberances, verie like both in forme and bignes to the trewe galls, onlie wantinge hardnes, but contayninge a softe spongious matter within, and often a maggott, and a hole by the side. These beginn to appeare most comonlie about the middest of July, of the bignes of a pease, sometimes by the middest of May of that bignes, as the 15 of May 1622 when I only sawe them.—*MS.* f. 85.

Walnut. *Juglans regia* L.

Cachrys Juglandis.

The Cachryes beginn to appeare in Aprill a little before the leaves are seene, and doe thrust themselves forth of the sides of the last yeres shootes, but one in a place, a little above the place where the footstalks of the leaves were fastened and are at the first closelie compacted and finelie chekered without anie footstalke or leaves cominge about or neare them; afterwards they growe more rare or looser, about two ynches in length neare as bigge as the little finger, not continuinge two weekes before they wither and fall away.—*MS.* f. 85.

Chestnut. *Castanea sativa* Mill.

Cachrys Castancae.

The Cachryes begin to springe about the midle of May together with the newe shoots forth of the bosomes of the yonge and tender leaves, but one out of the bosome of one leafe; composed of a midle ribbe or footstalke about 7, 8 or 9 ynches longe when they are at their full growth about July, which midle ribbe is thick sett by the sides with small scalelie bunches, which about the middest of July open, and there appeareth out of ech bunch many small short cheives of a very pale or light yealowish color as are also their topps, they appeare very slender in respect of their length, and soone after their cheives open or appeare they wither and fall away. These begin to come forth about the beginning of July, like rough hedgehog-like pills or husks, without any flowers, only bearinge at ye toppe a fewe whitish things like cheives, theis

husks growe cluster-wise 7 or more together, on one newe shoot.—
MS. f. 85.

Alder. *Alnus glutinosa* Medic.

Alder.

15 Maii 1622

The strobilus or fruite of the Alder cometh forth at the beginning of the springe when the leaves first appeare, they growe from the toppes of the last yeres twigges, clusterwise though not so neare together as the berries of the vine, about 8 or 9 in a cluster, at the first small, yet of their full length which exceedeth not half an ynch, in shewe like Cachryes, afterwards growinge like in fashion to an olive not fullie so bigge as a sparrowes egge, composed of many brownish scales standinge verie neare one another betwene which the seed lieth. These usuallie hange on a yere or more.—*MS.* f. 86.

Birch.¹ *Betula alba* L.

[Unnamed by Goodyer.]

Undated.

Ye leaves are small, smooth broad below, growing to a large point nicked by the sides, in some like those of the black poplar but smaller.

At y^e very first coming forth of the leaves, there cometh forth with them a round thinge like a Cachryes, but is indeed the fruite, about 3 quarters of an ynch longe, and nere a quarter of an ynch over, composed of flatt scales closely sett together, each scale contayning one seed of a browne color with a white kernell within. The scales with a light touch fall apart when y^e seed is ripe, the seed being neare ripe then cometh forth at y^e toppe and by the sides of y^e same yeres shoot small Cachryes.

The leaves and fruite come forth towards y^e later end of Aprill,
²⁶
the seed is ripe at ye later end of July, the Cachryes begin to appear at ye begining of July.—*MS.* f. 3 v.

Tribulus.

2 June 1622

[The year '1622' may be an error of Johnson. Goodyer's own *MS.* has '2 June 1621', which see.]

Wood Club Rush. *Scirpus sylvaticus* L.

Cyperus gramineus Lobelii.

8 Julij 1622

This hath about 7 or 8 smooth leaves growinge from y^e roote 3 foot longe, half an ynch broad, as it were 3 cornered, sharpe on ye edges especially, if you drawe your fingers on them downewards, amongst which growe uppe one ioynted smooth stalk, belowe neare as bigge as ye little finger, three cornered, but obtuse, blunt or

¹ Identified by A. H. C.

roundish, not sharpe cornered like galangall. Whereon at everie ioynt is placed one leafe cloathing or inclosinge ye stalk upward above ye ioynt an ynch and a half or more, about a foot and a half longe, and somewhat broader then y^e former, at ye toppe of which stalks growe usually 3 like leaves but much shorter, from which spring forth 6 or more little stalks or branches, after ye manner of galangall, 4, 5 or 6 ynches long except ye midle branch which usually growes not an ynch above ye divison, and all these are againe at their toppe diversly devided into many parts, bearinge little knobbs, scales or buttons very like those of Rushes, containyng very small yealowish 3 cornered seeds sharpe pointed at both ends, which together with ye little buttons in ye beg[inninge] of July fall off. The rootes at ye upper parte next ye leaves, are infinite of small white threddy strings, sendinge forth underneath other rootys (?) bigger then wheate strawes like those of *Arundo vallatoria* ioynted, white at ye first after of a brownish yealow, without smell, by which the plant exceedingly increase, growing high in aboundance together.—*MS.* f. 7 v.

On ye west parte of Gloster Hall by Oxford. 5 Julii 1622

The stalk is round rough or hairy, ioynted, neare as bigge on ye lower parte as ye little finger devided towards the toppe into a fewe branches, bearinge at everie ioynt one sharpe-pointed leafe without or with a very little footstalk about 5 ynches long and an ynch and half broad or hairy like ye stalk, not indented by ye sides, but sometimes with small excrescens as may be seen on ye leaves of Plantans standinge farr apart. Ye flowers growe at ye topps of ye stalks & branches on long slender footstalks, of a yealowe color, ech flower beinge composed of 5 greate broad topped leaves, which beinge full blowne is neare 2 ynches broad, containyng within many small yealowe chives. The roots growe forth at certaine ioynts on ye lower parts of ye stalks within ye water and mudd, made of many small hairy strings. This herbe at ye first tastinge seemeth not to be hott, but beinge held in a little space in ye mouth heateth & burneth little inferior to ye rest of his kind.—*MS.* f. 7 v.

Great Water Parsnip. *Sium latifolium* L.

Pastinaca aquatica latifolia at Oxford. 5 Julii 1622

[Rough draft for the next description.]—*MS.* f. 7 v.

Pastinaca aquatica maxima. 5^{to} Julii 1622

Sium maius Gerardo Phyto. p. 270. N^o. 11.

The stalkes are greate upright straight and tall, not inferior to

the garden parsneppe beinge neare as bigge as a mans arme, and as high as a tall man, with greate corners deepe chamfers, and hollowe within, devided into many braunches; whereon growe greate broad leaves like those of the garden parsneppe, composed of 5 or 6 sometimes 7 or 8 indented smooth leaves sett on ech side of a longe midle ribbe, and one at the toppe, of a stronge smell. The flowers were not come forth.

[Two lines left blank, perhaps for a description of the flowers.]

The rootes are infinite growinge at the lower ptes of the stalkes within the water and mudd, white, as bigge as wheatestrawes, and they againe sendinge forth abundance of threddy strings, by which rootes the plant increaseth, usuallie 3 or 4 stalkes together from one tuft. This I found growinge wild plentifully by the Rivers sides and in the water diches about Oxford, 5^{to} Julii 1622.—*MS.* f. 82 v.

Stachys germanica L.

Stachys. Wild Horehound.

[? 8 Julii 1622]

564 Stachys.—Wild Horehound is also like to comon horehound. There rise from the root hereof a greate number of stalks high or ioynted, and out of everie ioynt a couple of leaves, opposite or sett one against another, somewhat hard, a little longer than those of comon horehound and whiter, as also ye stalks are sett with soft haire and of a sweet smell. Ye flowers doe compasse ye stalk about as those doe of comon horehound, but they are yealowe and y^e whorles be narrower. The root is woodie and durable.—*MS.* f. 7.

Calamagrostis.

8 Julii 1622

[See 27 Apr. 1622.]

Sweet Flag. *Acorus Calamus* L.

Acorus legitimum Clus. 231. *Acorus*, Ger. 56. 6 Julij 1623

Acorus hath the leaves almost of the greater narrow leaved *Iris* [*Iris angustifolia*], but much longer, and of a most pleasant greene, the midle nerve or rib somewhat appearinge forth on both sides as in ye leaves of *Xiphium* whose forme they sufficiently resemble; and like them or that moorish plant which brings forth yealowe flowers like ye flowers of *Iris*, which some call *Iris palustris*, the better skilled more trulie *Butomus*. They come one out of another, but the midle leafe is most comonlie longer than the rest; but although they be of a tast somewhat bitter, yet not unpleasant, and beinge crushed they yeld a pleasant and aromaticall smell, which also they retaine

manie yeres after beinge dryed and without iuyce. About the beginning of winter they wither and are dryed, but in the beginninge of springe they shoote anewe as in narrow leaved Iris, moreover it produceth not a stalk betwene the leaves as the *Irides*, but out of the side of y^e root comes a leaf * stalk of the same lenght with y^e other leaves, that is sometimes from the midle unto y^e toppe is plaine and like the rest, but and narrower, and as it were fashioned into a triangular stalk it begins to be extenuated and made plaine it sends forth out (sometimes but very seldome two) not greater in ye beginning appearinge and stanndinge up a pright [little partinge (or cutting) themselves acrossse when it opens itself full of consistinge of fower small leaves, afterwards untill it gett the lenght and thic greene knobbes, in such a comelie order cone of the wild pine. It hath upper parte when it lies hid in ye earth, exceedinge white in the inner parte, distinguished * stronge, of a good smell, of a somewhat bitter and sharper tast, endowed with many and whitish and sweet smellinge threeds growinge to it in ye lower parte, but it creeps and spreads itself on the toppe of the earth, sendinge forth from the sides and almost everie ioynt or knott yonge ones, one after another, obliquely (or crookedlie) so that in a short time it takes uppe a greate space.

21 Julij [1624]

- a. two corners standing so neare together that they make a hollowe like a furrow.
- b. at the upper end of the furrowe.
- c. without any footstalk.—*MS.* f. 125.

* [Page torn.]

[The date shows that this description was made from the *Acorus* 'in flower in Mris. Mervin's garden 6^{to} Julii 1623'.—*MS.* f. 51.]

Papaver Argemone L.

Argemone capitulo longiore Ob. p. 144.
In Durford garden.

24 Julii 1623

Sanfoin. *Onobrychis sativa* Lam.

Caput Gallinaceum Belgarum.

24 Julii 1624

In flower 24 Julii 1624 between Langford & Stapleford in Wiltes by ye way on ye south side of ye river.—*MS.* f. 52.

Shrubby Suaeda. *Suaeda fruticosa* Forsk.

Chamaeypitis vermiculata. 10 Sept. 1624

The stalks are woodie not fuller of a finger bignes, a cubite or 2 foot high of a blackish or dark ash color devided into many branches whereon grow multitudes of small round fatt leaves of y^e fashion of w . . es very like those of comon stone croppe full of iuyce, of a salt tast, of a darke green colour. The rootes are also woodie branched of a blacker color then ye stalks. This plant continueth greene continually as it seemeth, and increaseth by the root growinge in thick tuffets close together. I observed no flowers nor seed.—*MS.* f. 127.

[A roughly written note on the back of the statement of 'Tenthmony' received by Edward Cole in 1608. See p. 373.]

Common Spleenwort. *Asplenium Trichomanes* L.

Trichomanes mas. — Jan. 1624

'Mr. Goodyer saith that in January 1624, he saw enough to lade an horse growing on the bancks in a lane, as he rode between Rake and Headly in Hampshire neere Wollmer Forest.'—*Ger. emac.* 1146.

Cowbane. *Cicuta virosa* L.

Sium alterum olusatri facie. 16 Sept. 1625

Found by Mr. Goodyer in the ponds about Moore Parke (*Ger. emac.* 257) and at Denham in Hertfordshire in standinge motes sine caule.—*MS.* f. 58.

Knotted Pearlwort. *Sagina nodosa* Meyer.

Alsine palustris foliis tenuissimis. 12 Aug. 1626

This hath a great number of very small grasse-like leaves, growing from the root, about an inch long, a great deale smaller and slenderer than small pinnes; amongst which spring up many small slender round smooth firme branches some handfull or handfull and halfe high, from which sometimes grow a few other smaller branches, whereon at certaine ioynts grow leaves like the former, and those set by couples with other shorter comming forth of their bosomes; and so by degrees they become shorter and shorter towards the top, so that toward the top this plant somewhat resembleth *Thymum durius*. The flowers are great for the slenderesse of the plant, growing at the tops of the branches, each flower consisting of five smal blunt roundish topped white flowers, with white chives in the middest. The seed I observed not. The root is small, growing in the myre with a few strings. This groweth plentifully on the boggy ground below the red Well of Welling-

borough in Northampton shire. This hath not beene described that I finde. I observed it at the place aforesaid, August 12 1626.—*Ger. emac.* 568.

Grass of Parnassus. *Parnassia palustris* L.

Gramen parnassi. 12 Aug. 1626

In the boggy ground below the Red Well of Wellingborough in Northamptonshire.—*Ger. emac.* 840.

Cerinthe minor L.

Cerinthe minor flore albo veris luteis. 23 Sept. 1628

This in stalks and leaves differeth verie little from the other, the flowers in shape are like, the color from the midle to the brim is of a whitish or pale yealowe, the brim itself a much deeper yealowe, the midle hath a ringe or circle of a reddish purple from that circle backward that is to the fasteninge of the huske, of a deepe yealowe, the seeds are like the other but as small againe.—*MS. f.* 120.

[The 'other' is *Cerinthe flore rubro*, 9 July 1621.]

Golden Lungwort. *Hieracium murorum* L.

Pulmonaria Gallica sive aurea latifolia. 27 May 1631

'I received some plants of this from Mr. John Goodyer, who first found it May 27, 1631 in flower; and the 3 of the following May not yet flowering in a copse in Godlemen in Surrey, adioyning to the orchard of the Inne whose signe is the Antilope.'—*Ger. emac.* 305.

Triticum vulgare L.

Wheat ear with Oats.—*Ger. emac.* 65. See p. 62. 1632

Water Plantain. *Damasonium stellatum* Thuill.

Plantago aquatica stellata. 2 Julij 1633

The roote is nothinge but a multitude of very small white thredds like hairs growinge in the myre amonge which springe the leaves, their footstalks are about 3 ynches longe, at the toppe of ech groweth igrosse plaine smoth leafe, not indented, an ynch long, a quarter of an ynch broad or somewhat broader, sharpe pointed, with two cares belowe sometimes, but most comonly without.

The stalks growe uppe amongst them, in number 5 or 6 plain smooth round firme not hollowe, small, as bigge as a small straw, on ech stalk groweth an umbell, consisting of 7 footstalks, 6 of them having at the toppe of ech a starr-like fruite, of six sharpe pointed husks like the rowell of a spurr, the seaventh footstalk beareth an other umbell, with 3 or 4, 5 or 6 footstalks & starr-like husks like the former.

Neare London highwaie in the watery plashes at the east end of the greate Comon betweene Sandie Chappell and Kingston, neare the bridge as you ride out of the Comon by a small Cottage there.

I saw no flowers.—*MS.* f. 137.

[A note scribbled on the back of a list of men of the Tithing of . . . [name obliterated], p. 381.]

[Goodyer first found this plant in 1618 growing on Hounslow Heath, the station quoted by Johnson.—*Ger. emac.* p. 418.]

Ferns.

Filix mas varietates & differentiae.

4 Julii 1633

I have observed foure sorts of Ferne, by most writers esteemed to be the male Ferne of *Dioscorides*: by *Anguillaria*, *Gesner*, *Cæsalpinus*, and *Clusius*, accounted to be the female, and so indeed doe I thinke them to be, though I call them the male, with the multitude. If you looke on these Fernes according to their severall growths and ages, you may make many more sorts of them than I have done; which I am afraid hath beene the occasion of describing more sorts than indeed there are in nature. These descriptions I made by them when they were in their perfect growths.—*MS.* f. 138; *Ger. emac.* 1129.

Broad Shield-fern. *Aspidium dilatatum* Sm.

Filix mas ramosa pinnulis dentatis.

4 Julii 1633

The roots are nothing but an abundance of small blacke hairy strings, growing from the lower parts of the maine stalkes (for stalkes I will call them) where those stalkes are ioyned together. At the beginning of the Spring you may perceive the leaves to grow forth of their folding clusters, covered with brownish scales at the superficies of the earth, very closely ioyned together: a young plant hath but a few leaves; an old one, ten, twelve, or more: each stalke at his lower end neere the ioyning to his fellowes, at his first appearing, before he is an inch long having some of those blacke fibrous roots for his sustenance. The leaves being at their full growth hath each of them a three-fold division, as hath that Ferne which is commonly called the female: the maine stalke, the side branches growing from him, and the nerves growing on those side branches bearing the leaves: the maine stalke of that plant I describe was fully foure foot long (but there are usually from one foot to foure in length) full of those brownish scales, especially toward the root, firme, one side flat, the rest round, naked fully one and twenty inches, to the first paire of side branches. The side branches, the longest being the third paire from the root, were nine inches long, and shorter and shorter

towards the top, in number about twenty paire; for the most part towards the root they grow by couples, almost opposite, the neerer the top the further from opposition: the nerves bearing the leaves, the longest were two inches and a quarter long, and so shorter and shorter toward the tops of the side branches; about twentie in number on each side of the longest side branch. The leaves grow for the most part by couples on the nerve, eight or nine paire on a nerve; each leafe being gashed by the sides, the gashes ending with sharpe points, of a deep green on the upper side, on the under side paler, and each leafe having two rowes of dusty red scales, of a browne or blackish colour: toward the top of the maine stalke those side branches change into nerves, bearing only the leaves. When the leaves are at their full growth, you may see in the middest of them at their roots the said scaly folding cluster; and as the old leaves with their blacke threddy roots wholly perish, they spring up; most yeares you may finde many of the old leaves greene all the Winter, especially in warme places. This groweth plentifully in the boggy shadowie moores neere Durford Abbey in Sussex, and also on the moist shadowie rockes by Mapledurham in Hampshire, neere Petersfield; and I have found it often on the dead putrified bodies and stems of old rotten okes, in the said moores; neere the old plants I have observed verie many small yong plants growing, which came by the falling of the seed from those dusty scales: for I beleeve all herbes have seeds in themselves to produce their kindes, *Gen.* i. 11. & 12.—*MS.* f. 138-9; *Ger. emac.* 1129.

[I thought to have called this *Dryopteris*, but that is described by Cordus and Tragus to be a very small tender ferne not above 9 inches high with creeping roots like those of *Polypodium* (*erased*).]—*MS.* f. 138.

The three other have but a twofold division, the many stalks and the nerves bearing the leaves. The roots of them all are blacke fibrous threds like the first, their maine stalks grow many thicke and close together at the root, as the first doth: the difference is in the fashion of their leaves, and manner of growing, and for distinctions sake I have thus called them:

Male Shield-ferne. *Aspidium Filix-mas* Sw.
Filix mas non ramosa pinnulis latis densis minutim dentatis.

4 Julii 1633

The leaves are of a yellowish greene colour on both sides, set

very thicke and close together on the nerve, that you cannot see betweene them, with marvellous small nickes by their sides, and on their round tops: each leafe hath also two rowes of dusty seed scales; the figures set forth by *Lobel*, *Tabern.*, and *Gerard*, under the title of *Felix mas*, do well resemble this Ferne. This growes plentifully in most places in shadowie woods and copses.—*MS.* f. 139; *Ger. emac.* 1129–30.

Nephrodium Filix-mas, Sw., var. *affinis* (Newm.).

Filix mas non ramosa pinnulis angustis, raris, profunde dentatis.

4 Julii 1633

The leaves are of a deepe greene, not closely set together on the nerve, but you may far off see betwixt them, deeply indented by the sides, ending with a point not altogether sharpe: each leafe hath also two rowes of dusty seed scales. I have not seene any figure well resembling this plant. This groweth also in many places in the shade.—*MS.* f. 140; *Ger. emac.* 1130.

Prickly Shield-fern. *Polystichum aculeatum* Sw.
var. *lobatum* Syme; or Angular Shield-fern. *P.*
angulare Presl.

Filix mas non ramosa pinnulis latis auriculatis spinosis.

4 Julii 1633

The leaves are of a deeper greene than either of the two last described, placed on the nerve not very close together, but that you may plainly see between them; each leafe (especially those next the stalke) having on that side farthest off the stalk a large eare or outgrowing ending, with a sharp pricke like a haire, as doth also the top of the leafe: some of the sides of the leaves are also nicked, ending with the like pricke or haire. Each leafe hath two rowes of dusty seed scales. This I take to be *Filix mas aculeata maior Bauhini*. Neither haue I seene any figure resembling this plant. It groweth abundantly on the shadowie moist rockes by Mapledurham neere Petersfield in Hampshire.—*MS.* f. 140; *Ger. emac.* 1130.

Marsh Shield-fern. *Dryopteris Thelypteris* Asa Gray.
Dryopteris Penae & Lobelii.

6 Julii 1633

The roots creepe in the ground or mire, neere the turfe or upper part thereof, and fold amongst themselves, as the roots of *Polypodium* do, almost as big as a wheat straw, and about five, six, or seven inches long, coal blacke without, and white within, of a binding taste inclining to sweetnesse, with an innumerable com-

panie of small blacke fibres like haire growing thereunto. The stalkes spring from the roots in severall places, in number variable, according to the length and encrease of the root; I have scene small plants have but one or two, and some bigger plants have fourteene or fifteene: they have but a two-fold division, the stalke growing from the root, and the nerve bearing the leaves: the stalke is about five, six, or seven inches long, no bigger then a bennet or small grasse stalke, one side flat, as are the male Fernes, the rest round, smooth, and green. The first paire of nerves grow about three inches from the root, and so do all the rest grow by couples, almost exactly one against another, in number about eight, nine, or ten couples, the longest seldome exceeding an inch in length. The leaves grow on those nerves also by couples, eight or nine couples on a nerve, without any nickes or indentures, of a yellowish greene colour. This Ferne may be said to be like *Polypodium* in his creeping root, like the male Ferne in his stalke, and like the female Ferne in his nerves and leaves. I could finde no seed-scales on the backesides of any of the leaves of this Ferne. Many yeares past I found this same in a very wet moore or bog, being the land of *Richard Austen*, called Whitrow Moore, where Peate is now digged, a mile from Petersfield in Hampshire; and this sixth of Iuly, 1633, I digged up there many plants, and by them made this description. I never found it growing in any other place: the leaves perish at Winter, and grow up againe very late in the Spring.—*MS.* f. 140; *Ger. emac.* 1135-6.

[There is a sketch by Goodyer of a fern on the same page as this description].

Ulmus campestris Sm., *U. montana* Stokes, *U. glabra* Miller,
U. minor Miller.

Elms.

1633

[For Goodyer's descriptions of the four species of Elm see p. 38.]

Among the undated notes and species contributed by Goodyer to Johnson's second edition of Gerard were the following. The descriptions must therefore be earlier than 1633 and several may date from *c.* 1621.

Marsh Helleborine. *Epipactis palustris* Crantz.

Palma Christi, radice repente.

[Before 1633]

'It growes also plentifully in Hampshire within a mile of a market Towne called Petersfield, in a moist meadow named Woodmead, neere the path leading from Petersfield, towards Beryton.'

[Johnson (*Ger. emac.* 227) prints this locality without acknowledgement, but it was doubtless communicated to him by Goodyer. The

description of the species is unfortunately illustrated by a wood-block of *Goodyera repens*, an orchid of a genus named by Robert Brown in honour of Goodyer, thus unfortunately connecting his name with a plant which he had probably never seen. Goodyer's 'Palma Christi' or 'Creeping-rooted Satyrion' has been identified by Canon Vaughan as *Epipactis palustris*, a species to be found in boggy situations in several parts of Hampshire. Brown, thinking that it was the same as a rare northern orchid, found in certain fir-woods in Cumberland and Scotland, gave it the name of *Goodyera repens* in honour of our Hampshire botanist, who, it is suggested, might possibly have met with a specimen of this rare northern plant in the low-lying grounds between Petersfield and Maple-Durham.]

Rampions. *Plyteuma orbiculare* L.

Rapunculus corniculatus montanus. [Before 1633]

Johnson 'received seeds and roots hereof from Mr. Goodyer who found it plentifully growing wilde in the inclosed chalkie hilly grounds by Mapledurham'.—*Ger. emac.* 455.

Sea Bindweed. *Convolvulus Soldanella* L.

Soldanella marina. n. d.

[The local Isle of Wight name, 'Scurvy Grass,' given to this plant, supports the truth of the following remark of Johnson:] 'My friend Mr. Goodyer hath told me that in Hampshire, at Chichester, and thereabout, they make use of this for Scurvie-grass, and that not without great errour, as any that know the qualities may easily perceive.—*Ger. emac.* 839.

Golden Saxifrage. *Chrysosplenium oppositifolium* L.

Saxifraga aurea. [Before 1633]

Mr. Goodyer hath also observed it abundantly on the shadowie moist rocks by Maple Durham in Hampshire.—*Ger. emac.* 842.

[The first record for Hants.]

Geranium lucidum L.

Geranium saxatile, Thalii. [Before 1633]

Master Goodyer found it growing plentifully on the banks by the highway leading from Gilford towards London neere unto the Townes end.—*Ger. emac.* 938.

[First record for Britain.]

Phyllitis Scolopendrium, Newm., var. *multifida*.

Phyllitis multifida. Finger Harts-tongue. [Before 1633]

Mr. Goodyer found it wilde in the banks of a lane neere Swaneling, not many miles from Southampton.—*Ger. emac.* 1139.

Digitalis ferruginea L.

Digitalis ferruginea.

21 Sept. 1633

This is a verie comely plant growinge like a pyramide.

The maine stalk is 3 or 4 foot high, greene, smooth, with some edges, not hollowe but filled with a white spungie pith, growinge upright as strait as an arrowe, not farr from the lower part it sendeth forth many branches, not growinge so high as the stalk.

The leaves are many, & spread upon the ground before the stalk groweth uppe, greene & smooth on both sides, without nicks by the sides, usuallie with 5 ribbes, like those of quinquenaria plantayne, about — ynches long & — ynches broad. Also such leaves growe on the stalks and branches, but small and shorter towards the toppe, not by couples but heere and there.

The flowers make a comely spike, & as the stalk and branches growe so they flower still upwards nere their topps hollowe, of the bignes of the toppe of the little finger, the upper side half an ynh long, the lower side hath lipps stickinge forth of a quarter of an ynh, & on ech side of the flower ther is one excrescence or — like the toth of a sawe, in the inside there is usually 4 cheives growinge to the upp part of the flower, the whole is of a rustie iron color, with many purplish strakes, & somewhat hayrie both without & within.

The flowers fallen, the seed vessels playnely appeare makinge a longe spike, besettinge the stalks and branches round about, in forme round and sharpe pointed, but half as bigge as the flower half an ynh large at the lower part of ech seed vessell groweth fower scalie leaves, half as bigge but fully as longe as the nayle of the little finger.—*MS.* f. 137 b.

Tooth cress. *Dentaria bulbifera* L.

Dentaria baccifera v. bulbifera.

6 Aug. 1634

At Mayfield in a wood of Mr. Stephen Penckhurst called Highreed and in another wood of his called ffox-holes.—*MS.* ff. 53, 62.

Procumbent Pearlwort. *Sagina apetala* L.

20 June 1634

Perianthos 4 greene blunt-topped leaves hollowed sherie fashion, spreading open, one opposite against another crossewise.

The flower 4 very small white blunt-topped leaves. not so bigg as a small pinnshead a quarter as bigg as ye perianthos spreading iust between them, making a duble crosse.

- pointell Small white & short open at y^e toppe into 4 pts, one parte lyenge iust over ech flower.
- chives 4 white chives standing just as farr out as the top of ye pointell and ech pointell standinge iust opposite to ech leafe of ye perianthos.
- pericarpium hath 4 greenish leaves in form of those of the periantos, closed fast together till the seed is ripe.
- seed the seed is small, smaller than that of sweet margerom and . . . of colour like it, many lyeng in one pericarpium.—*MS.* f. 14.

[The identification of this unnamed description has been due to my friend Mr. A. H. Church. It is the earliest English account of the plant.]

Horse-shoe Vetch. *Hippocrepis comosa* L.

Ferrum equinum Germanicum siliquis in summitate. 9 Aug. 1634
On Buttersworth Hill, 9 Aug. 1634: ripe seed.—*MS.* f. 53 v.

Marsh Isnardia. *Ludwigia palustris* Ell.

Herba aquatica rubescens, facie Anagallidis flore luteo.

19 Aug. 1645

The stalks are smooth almost round, sometimes some parts of them a little square, reddish, firme within, not hollowe, sometimes allmost a foot longe, of the bignes of those of orgamen.

The leaves growe by coopes, on the stalkes & branches, on short footstalks somethinge like those of *Anagallis flo. luteo*.¹ The biggest are nere 3 quarters of an ynch longe & half an ynch broad, also reddish, smooth, nothinge at all indented by the edges; those on the toppes of the stalkes & branches are shorter and smaller; the ioynts about the middest of the stalkes are allmost an ynch apart but towards the topps of the stalks & branches, they are very neare together. The branches come forth at the bosomes of the leaves, & thereby increase very much. The roots are like small threeds & come forth at the ioynts of the stalks & branches, & take hold in the mudd.

The flowers come forth at the bosomes of the leaves towards the toppes of the stalks & branches, usually forth of ech bosome but one, sometimes two, in one bosome & but one on that opposite to them; they are small, scarce to be called flowers. but are like the huskes of many herbes that containe the flowers before they be opened, this huske or flower is not half a quarter of an ynch longe, and is divided into 4 parts, or leaves at the toppe, ech leafe beinge sharpe pointed & little bigger then a small pinnes head of a greenish

¹ *Lysimachia nemorum* L.

colour, & does not fall offe as the leaves of other flowers doe but continue on till and after the seed is ripe.

The seed is containd in that huske, & is white & as small as dust.

The whole herbe is of a reddish colour, and flotes in or uppon the water and prospers well when all the water is dried from it, & then flowers, seldom before. I could never observe any flowers but on those plants from which the water was dried away, and that in August.

I have long observed this plant, as I found it growinge in the rivulett on the east side of Petersfield, runinge and a heathy comon about the middest thereof. I cannot yet tell what genus it is, nor what name is most proper for it.—*MS.* f. 141.

[The Marsh *Isnardia* is one of the very rarest of Hampshire plants, and Goodyer's description has not been printed before; indeed it was not known that he had noted it about a quarter of a century before the date of the first 'record' given by Townsend. After Merrett had recorded Goodyer's discovery in 1666, the plant does not appear to have been seen again until about 1835? when it was rediscovered by Miss Rickman and J. Barton; 'and in the moist summer of 1848, Dr. Bromfield found it plentifully (he had searched for it unsuccessfully in previous dry summers) in marshy spots, into which expanded at intervals the shallow stream which drains the great pond at Petersfield. I am not aware of the plant having been found at Petersfield since 1848, though it has been repeatedly searched for. The shallow stream above described is now so circumscribed that even during the wet summer of 1879 it expanded into no marshy spots in which *Isnardia* could have a chance of growing. I searched the stream through the Common and along its course downwards for about a mile, but without success. The plant is now extinct in Sussex, the only other county in which it has been found in Great Britain.' 'Mr. Bolton King's patient determination to rediscover the plant [in the Brockenhurst neighbourhood, where it had been found by Borrer in 1843] was rewarded by finding it abundantly in 1878 in another spot in the neighbourhood.']*¹

Beech. *Fagus sylvatica* L.

Fagus.


Before 1650

I found one much varying in his leaves, some were whole as those of the ordinary, others much jagged or divided.—Goodyer quoted by How, *Phytologia*, p. 40.

1650–1656

The notes on the following nine plants occur in HOW'S handwriting in his interleaved copy of the Phytologia Britannica. They must therefore have been written between 1650, when the book was published, and 1656, when he died.

¹ Townsend, *Flora Hampshire*, p. 158.



THE HEATH AT PETERSFIELD

Cardamine impatiens L.

Cardamine flosculus minoribus, sive impatiens.

From rills and ditch sides about Bath.

Dr. Johnson was mistaken in saying yt this was *Sium minimum*, Alp. I have both ye plants. I admonished him of this error but he lived not to amend it. J. Goodyer.—p. 21.

[Johnson died in 1644.]

Polypodium Dryopteris L.

Dryopteris, Trag. Tree-fern.

It growes on a bottome called Rogers Deane in ye parish of Faringdon in Hampshire, about a mile and a half from ye church, a furlong from one John Trybes dwelling house on ye north east part of ye house about 2 miles from Alton about a mile north east from Dogford Wood. Great antient beeches kept ye sunne from shining on ye Plants. Ann. 1654 many of those trees were cut downe. The Plants ye sunne shone on y^t summer '54 were short, ye leaves growing on short stemms neere ye earth as Tabernaemont pictureth it, p. 501 tom 2 under ye title of *Filicula petraea* fem. 3. Those yt grew under ye trees were much higher agreeable to Tragus figure p. 538. John Goodyer.—p. 35.

Ye least Furze. *Ulex nanus* Forst.

Park. des. of *Genista spinosa minor* p. 1003 accords not with ye Least Furze; ours beares no leaves at all. They are but ye first sproutings of ye thornes or prickles, even as of ye great furze (bee hee what hee will y^t willes ye contrary) ye cods have furze, even as ye cods of ye greater furze. I cannot find from whence Park. rec^d his fig. I suppose it was made by imagination. J. Goodyer.—p. 45.

[Dr. Stapf, who has kindly assisted in the determination of Goodyer's furzes, writes that in his opinion and in that of Mr. Sprague *Genista spinosa minor* Park., p. 1003, is entirely dubious. The figure in Parkinson goes back to the 1588 edition of Tabernaemontanus, where it is meant to illustrate *Nepa Theophrasti* of Pena and Lobel, but Tabernaemontanus himself says he is not sure whether it fits that plant. There is no indication where it was drawn from. Lobel's *Nepa Theophrasti* is evidently *Ulex parviflorus*, and it may be that the figure is just a very bad illustration of that species. This *Ulex* would probably not be hardy in England.]

Gorse. *Ulex Europaeus* L.

Genista spinosa flore albo Park. 1003.

[A whitish flowered variety.]

Ulex parviflorus.

Genista spinosa major brevibus aculeis Bauh. Pin. p. 394.

This I suppose groweth not in England. Pena and Lobel in *Adv.* p. 354 had seene it nowhere but in Province wch is a hott country, and Lob. lived time enough in England before ye *Adv.* was written to have observed it if it had growne but half so common as ye lesser Furze.¹ Cam[erarius] in *hort. med.* pag. 106 saith ‘in fichlibus asservanda’ wch argues yt it will not endure abroad in a cold countrie in ye winter. The Icons yt were made for Nepa in *Adv.* p. 354, in Tabern. *Ic.* p. 408, in *Hist. Lugd.* p. 164 agree not with ye lesser Furze. Parkinson sayes yt his *Genista spinosa minor* p. 1003 is ye Nepa of Lob. This duly considered I am confident to affirme yt our lesser Furze is not yet described. John Goodyer.—p. 45.

[Druce (*Goodyer*, p. 26) is all at sea in the transcription of this paragraph. Dr. Stapf writes, ‘The Bauhinian species is evidently a mixture. To judge by the first synonyms (*Anguillaria* and *Pena* and *Lobel*) it was very probably meant for *Ulex parviflorus*, but through the additions (*Tabern. &c.*) it got confused. See my observations sub *U. nanus.*’]

Toad rush. *Fucus bufonius* L.

Gr[amen] holosteum Alpinum minimum. Bauh. Prod.

‘Male a Johnsono *Holosteum pumilum non descriptum*, pervenit in ericetis. Joh. Goodyer.’—p. 53.

—? (*See p 171.*)

Gramen murorum spica longissima. Capons taile-grasse.

‘Mr. Goodyers upon ye walls of Winchester.’—p. 54.

Oenanthe angustifolia.—p. 81. (*See under 18 June 1620.*)

Pulmonaria angustifolia L.

Pulmonaria maculosa.

How substitutes the name of Goodyer for that of Loggins as the authority for the locality. ‘Neer Kings-wood in Hampshire.’—p. 100.

[*See under 25 May 1620.*]

Vicia sylvatica L.

Vicia maxima sylvatica.

Great wood Vetch from a wood nigh Bath.

How has changed name to *V. max. sylvatica* ‘*spicata Bathoniensis Goodyeri*’.—p. 129.

[How’s MS. note in his *Phytologiu.*]

¹ *Ye lesser furze.* Parkinson uses the term ‘lesser furse bush’ for *Genista spinosa minor*. Does not Goodyer use it in the same sense, just as an English translation of that name?—O. S.

Geranium columbinum L.

Geranium columbinum foliis magis dissectis, pediculis longissimis.

Aug. 1654

I found it wild in ye beginning of August 1654. It is not described or pictured yt I find. John Goodyer.

Qu[ery] ye place of growth and des[cription] for this and [the next plant] following.

‘In several places of Hampshire. J. Goodyer.’ Merrett, *Pinax* 45.

[Druce’s reading of the passage giving the locality as White Chapell is not justified.]

Sisymbrium Irio L.

Erysimum ii Tab. Qu[aere] locum. [Before 1656]

Grows in ye streets near White Chappell east from Aldgate, London. J. Goodyer.

[Note in How’s handwriting at end of his *Phytologia*.]

Bunias orientale L.

Rapistrum aliud non bulbosum. P. 862. [Before 1656]

‘In the broad street by White Chappel, Mr. Goodyer.’ Merrett, *Pinax* 103.

[Although this record was not printed before 1666, it is probable that Goodyer found the plant with ‘*Erysimum* ii’ on the same visit to London. If this be so, it must have been before the date of How’s death in 1656.]

Wild Madder. *Rubia peregrina*.

Rubia sylvestris. 12 Aug. 1655

[Recorded by Turner, 1551–68.]

The stalks are fower square, hollowe within smooth, one, two, three foot high, sometimes higher, with ioynts three or fower ynches apart: at the ioynt somethinge about the ground growe forth two side branches & the like side branches at everie ioynt upward on the maine stalk, and those branches at the ioynts send forth other side branches after the same manner. At each ioynt of the greater stalk growe the leaves in a circle which are smooth, 6, 7, 8 sometimes 9 at ech ioynt, the biggest leaves sometimes are about 3 quarters of an ynch longe, and broad. at ech ioynt also of the branches growe such leaves, but are smaller & smaller towards the toppes of the stalkes & branches.

The flowers growe abundantlie neare the toppes of the stalkes and branches, racematim, and are white, everie flower havinge 4 small sharpe pointed leaves. The seeds are small, round, manie times two growinge together. The rootes are small, with some threeds,

creepeth farr in the earth, the barke thereof beinge of a yealowish redd color, & stickie hard in the midle.

The stalks & branches die everie yere, the rootes continue manie yeres.

It usuallie growes in drie chalkie grounds, in barren places the stalks are short, a foot or little longer, & needes no supporter, in richer grownds they are much longer, in hedges & amongst bushes, longest and needes supporters, as doe *Gallium album*.—*MS.* f. 142.

Apium inundatum Reichb. f.

Sium pusillum foliis variis.

2 Junij 1656

The leaves before the plants have stalkes are like those of fennell but much smaller, growinge in abundance in the mudd within the water. The stalkes are hollowe as bigge as a wheate strawe, greene for the most parte sometimes reddish & a foot long or longer & growe uppe amongst the leaves not upright but swimminge sidelonge in the water, the toppes only appearinge above it, wch at the ioynts devides into severall branches. At each ioynt on the stalk within the water growes one leafe, like fennell as the former, but shorter & smaller, towards & on the toppes of the stalkes & branches with eyther and a little above the water, or swimminge on it are leaves much broader then the former, in forme & fashion to those of *Eruca palustris minor* [Water-Rocket¹]. Tab[ernaemontanus] pictured in his Icons p. 447; only these have not above 2 or 3 paire of small leaves on the midle ribbe of ech leafe, & that hath 4, 5 or more.

At the ioynts of the stalks towards their toppes growe the foote-stalks about an ynch of length, ech footstalk for the most parte devided towards the toppe into 2 parts, on ech of which parts comonly groweth 2, 3, 4 or 5 small white flowers, clusterwise together, ech flower havinge 4 small sharpe pointed leaves, ech leafe beinge no bigger then a small pins head.

In their places come 2, 3, 4 or 5 seeds as bigge, and of the forme of Caraway or parsley seeds clusteringe also together.

The rootes are as small as threeds & growe at the ioynts of the stalks & fasten them selves in the mudd whereby it mightelie increaseth. It flowers about the beginninge & midle of May.

This plant growes comonly in small lakes, & water plashes, but not described before that I know of. 2 Junij 1656 I made this description when the plant hadd almost done floweringe, & much of the seed was of its full growth.—*MS.* f. 143.

¹ *Nasturtium sylvestre* R.Br.

Holosteum.

2 Junii 1656

See 'Herba aquatica' 19 Aug. 1645.

Marsh Ragwort. *Senecio paludosus*, L.

Conyza aquatica laciniata.

19 July 1656

Growes in greate plentie in the fenns in Norfolk near Downam marktett neare Linn, by the relation of John Meader of Downam marktett, a grocer.—*MS.* 9, f. 186 a.

Sapphire. *Crithmum maritimum* L.

Crithmum chrysanthemum Ger. em.

[1656]

Grows by Hurst Castle, Hants, by relation of John Meader of Downham.—*MS.* 9, f. 201 a.

? Smooth Tare. *Vicia tetrasperma* Moench.

Viciae sive Craccae minimae species cum siliquis glabris Joh.

Bauhini. Tom 2, p. 315. JG. 626. An *Vicia segetum* singularibus siliquis glabris. Pin. C. Bauh. p. 345 (b. 3). 4 Junii 1657

The stalk, a verie little above the ground & so upward sendeth forth at the bosomes of the leaves severall other stalkes or branches, angular not round, a foot, a foot & half high or longe, little more or lesse.

The Leaves growe on the stalkes or branches, about an ynch apart not by couples, but one in a place, ech leafe beinge composed of a midle ribbe, endinge with a tendrell with which it taketh hold of what groweth neare it and on ech midle riþe 3 or 4 paire of leaves, ech leaf being about half an ynch longe and not fullie one eight part of an ynch broad.

The flowers growe on footstalks forth of the bosomes of the leaves, usuallie one footstalk out of ech bosome & no more, ech footstalk beinge about an ynch longe as small as a small threed. The flowers come forth in May & June & growe on the toppes of the footstalkes, one, two or three at the most, yet seldome above one on ech footstalk, a quarter of an ynche longe, of a blewish or purple violett color, of the fashion of those of *Aracus sive Cracca minima* Lobelij.

The coddcs succeed the flowers, ech codd beinge about half an ynch longe, half a quarter or the eight part of an ynch broad, & are smooth not hairie or woollie.

The seeds are contained in the coddcs, 2, 3 or 4 seeds in ech codd, and are [unfinished].

The roote is small accordinge to the proportion of the plant devided into severall strings or thredds & perisheth when the seed is ripe.—*MS.* f. 145.

[? First evidence for Hants.]

The next seven records are from Goodyer's MS. entries in How's interleaved copy of his *Phytologia* (MS. 18) received by Goodyer on 30 Apr. 1659.

Rumex Acetosa L.

Acetosa maxima After 30 Apr. 1659
Goodyer MS. in How, *Phyt. MS.* 18, p. 2.

Alsine aquatica verna. Springe chickweed.
MS. 18, p. 4.

Arctium Lappa L.

Arctium montanum et Lappa minor Galeni Lob. Button Burre.
Mangerfield in Master Langlie's Yard.¹ MS. 18, p. 10.

Galeopsis Tetrahit L., var. *bifida* Boenn.

Cannabis spuria altera flo. purp. Nettle Hempe.

C. spuria altera sylvestris, Lamium quorundam Lob. Icon. 527.
In agris.—MS. 18, p. 20.

Carex vulpina L.

Gramen palustre Cyperoides Lob. Ger. Great Cyperus Grasse.
MS. 18, p. 54.

Viola tricolor L.

Viola sive Jacea tricolor sylvestris parva. Wild Pansies.
In agris.—MS. 18, p. 130.

Common Ragwort. *Senecio Jacobaea.*

Jacobaea Pannonica 2 Clus. 4 June 1659
Mr. Tho. Bartar² of Petersfeild, schoolemaster gathered this imagined *Pulmonaria Gallica Lobelii*, on Ladle Hill in flower and brought it to J. G. the 4 of June 1659.

¹ Druce, *Goodyer*, p. 25, notes that this record is not included in the *Flora of Hampshire*. Why should it be? The Manor of Mangotsfield was purchased in 1612 by Mr. Philip Langley who lived in Rodway Hill Manor House, still a fine survival of the time of King Henry VIII, three of whose wives may have visited it (Emlyn Jones, *Our Parish: Mangotsfield*). It was here that Johnson and his *socii* were entertained in so grand a manner by 'that truly noble and generous man Philip Langley' when on their herborizing journey in July 1634. Evidently finding his own words inadequate to describe the luxury of the house and the sumptuous meal provided for his company, Johnson fell back on Virgil, and quotes a passage descriptive of some similar occasion when 'laden tables crowned with wines', huge goblets, drinking cups, &c., marked the feast (White, *Flora of Bristol*, p. 54).

² Druce has misread this name as 'Geo. Burton', in the *Rep. Bot. Exch. Club Suppl.* 1916, p. 23.

It is *Jacoboea Pannonica* 2 Clus. C. Bauh. p. 131 (b. 8) & it is *Jacoboea angustifolia* in this booke¹ p. 280.—MS. 18, p. 10.

[Then follows a recipe taken from Parkinson, *Theatrum*, p. 518, for an Alkanet ointment prepared by boiling 20 earthworms in good sallet oil.—MS. in How, *Phytologia*, p. 10.

Ladle Hill, crowned by a circular camp, is near Burghclere in the north of Hampshire.]

Cloudberry. *Rubus Chamaemorus* L.

Cloudberry. 15 Apr. 1663

Mr. Tho. Thornton parson of Sutton aſs Sulton in Sussex in Arundel Rape, borne at Bentham in Yorkshire, 2 miles from Yngleborowe hill, 15 Apr. 1663 promised Cloudberry.—Goodyer's MS. note in his copy of Ray, *Catalogus Plantarum circa Cantabrigiam nasc.*

Species of plants described in Goodyer's MS. and included by MERRETT in his Pinax by permission of Edmund Yalden in 1666.

Alsine flosculis conniventibus. *Montia fontana* L.

Anagallis aquat. flore parvo viridi caule rubro. *Ludwigia palustris* Elliot.

Aria Theophrasti fol. obtusis. *Pyrus Aria* L.

Caucalis pumila segetum. *Caucalis arvensis* Huds.

Geranium columbinum fol. magis dissectis. *Ger. columbinum* L.

Gramen Paniceum. Bearded Panick grass. *Panicum Crus-galli* L.

Juniperus sterilis. *Juniperus communis* L.

Lathyrus maior angustifol. fl. pallidè rubro. *Lathyrus sylvestris* L.

Lychnis sylv. flore carneo odorato. *Lychnis dioica* L.

'Ex Misto Gooderiano.'

Nidus avis. *Neottia Nidus-avis* Rich.

Oenanthe angustifolia, Lob. *Oenanthe Lachenalii* Gmel.

Pulmonaria foliis Echii. *Pulmonaria angustifolia* L.

Quercus serotina, procerior foliis fructuq. minoribus, Dor-Oak.

Linwood Hill, Bramshaw, Wilts. *Quercus robur* L. agg.

Rapistrum aliud non bulbosum. *Bunias orientale* L.

Rapunculus sylvestris flore rubro albescente. *Campanula patula* L.

'In the pastures & hedgesides on the North-west of the Moor not far from the great bog neer *Petersfield*, Mr. Goodyer.'

Sedum Divi Vincentii, ND. *Sedum rupestre* L. var. *minus*.

Serpyllum foetidum. *Thymus Serpyllum* L.

Sium umbellis ad caulium nodos. *Apium nodiflorum* Reichb.

Taxus tantum florens. *Taxus baccata* L.

¹ *Gerard emaculatus*.

- Veronica mas recta. *Veronica officinalis* L.
 Vicia Bathoniensis vel maxima sylvatica. *Vicia sylvatica* L.
- MERRETT also notes the following nine species as occurring in or near Petersfield, a locality possibly supplied by Goodyer.
- Caryophyllus saxatilis Erica fol. umbellatis corymbis, C. B.
 Probably *Arenaria tenuifolia* L.¹
 In the middle way betwixt Lippock and Petersfield.
- Chamaepeuce foemina seu polyspermos. *Lycopodium clavatum* L.
 A mile on this side Lippock in Hampshire.
- Esula minor seu Pithusa G. 502. *Euphorbia* sp.
 In divers corn fields near Petersfield.
- Fungus corallinus ad antiquarum arborum radices.
Clathrus cancellatus L.
 In the Woods near Peters Field.
- Gramen Paniceum procumbens. *Panicum sanguinale* L.
 In a Lane & watery places and Ditches near Petersfield.
- Gramen Piperinum, Pepper-grass. *Pilularia globulifera* L.
 Near Petersfield.
- Gramen Secalinum maximum. The greatest Rye grass. P. 1144.
Hordeum sylvaticum Huds.
 In the Woods a mile west from Petersfield.
- Holosteum repens juncifolium. Probably *Scirpus fluitans* L.¹
 At the bottom of the Moor on the East side of Petersfield.
- Rapunculus corniculatus montanus. Ger. em. 455.
Phyteuma orbiculare L.
 Between Selbury Hill and Beacon Hill in the way to Bath, and
 in the Chalkey hills by Maple-Durham, Hampshire.

Species attributed to Goodyer by Ray, and not included in preceding lists.

- 'Flea-grass.' *Carex pulicaris* L.
 Cyperoides pulicare Merret, *Pinax*, 'observed first by Mr. Goodyer and by him named Flea-grass, from the likeness of its seeds, both for figure and colour'. Ray, *Synopsis*. The locality is given as 'a mile East of Oxford', but whether Goodyer himself found it there is not stated.

¹ Species determined by B. D. J.

GOODYER'S LIBRARY

AT his death in 1664 Goodyer bequeathed his collection of books *de Plantis* to Magdalen College in Oxon, 'to be kept entirely in the library of the said College for the use of the said College'; and one Compton, described in the College Accounts as 'auriga de Petersfield', was paid £2 for bringing the books to Oxford. The librarian,¹ who incorporated the bequest, inscribed the greater number of the volumes with the words 'Ex dono Joh. Goodyer, generosi', and entered a list of them in a Book of Benefactors: a few volumes were, however, left unmarked. At first the books were more or less kept together, but the changing needs and views of successive generations led to their being scattered throughout the library, some being removed to a distant room in the Founder's Tower. In about 1909 Canon Vaughan, the distinguished Hampshire botanist, when preparing an article on John Goodyer, entitled 'A forgotten Botanist of the Seventeenth Century', could get no adequate idea of his predecessor's library. Eventually, however, the original list was discovered and copied. It is headed thus:

'A. D. 1664. Johannes Goodyer generosus idemque Botanicus celeberrimus libros sequentes (qui fere universos de re herbaria tractantes complectuntur) ad valorem plus minus 120^{lib} amoris ergo moriens Collegio Magdalenensi Legavit',

which may be translated as follows: 'John Goodyer, gent., and a most distinguished Botanist, bequeathed at his death to Magdalen College as a token of his affection the following books (which comprise almost all the authorities on botany) to the value of about £120.'

Canon Vaughan at once realized the value of the books. But let him speak for himself. 'The names and descriptions of the volumes, reveal a most splendid legacy of botanical treasures. I had already recognized in Mr. John Goodyer a botanist of high repute, but that he possessed such a library I never for a moment suspected. The discovery came to me as a revelation. With the exception perhaps of the British Museum, the Bodleian Library, the Cambridge University Library, and the Library of the Linnean Society,

¹ In 1664 Trebbecke, a chaplain, was paid £6 10^s 'pro cura Bibliothecae', and John Clitheroe received the customary salary of £1 'pro supervisione Bibliothecae'.

there can be few, if any, such collections of botanical works of the sixteenth and first half of the seventeenth centuries in this country.'

When appointed to the charge of the library in 1919 I was quite unaware of Canon Vaughan's correspondence with my predecessor, and the transcript of the list of the Goodyer books was not in the library. Using the old list in the Book of Benefactors as a guide, the work of bringing the scattered volumes together was begun, and with it the compiling of a new catalogue of those works which are still in the library. It is a matter for congratulation that the collection is far more extensive than a somewhat imperfect list printed in the *Supplement to the Botanical Exchange Club Report for 1916* would lead one to expect, and that with few exceptions most of the volumes mentioned in the original list can be identified. All the books in the collection have now been marked with serial numbers and with the canting crest of the Goodyer family. There are about 239 separate printed treatises bound up into 134 volumes which in size are about equally divided into folio, quartos, and octavos. Of *incunabula* or works printed in the fifteenth century there are a few examples, chiefly from the famous press of Aldus in Venice; about a hundred of the treatises belong to the sixteenth, and the rest to the first half of the seventeenth century.

The wealth and variety of the collection is clearly shown by the catalogue, but the personal associations of particular volumes with contemporary botanists, as well as the marginal annotations, indexes, and notes, which Goodyer so freely added to the works he used, give a unique value to many of the books.

Some sixty-four of his books have the day of their acquisition and the price paid clearly written on the first fly-leaf. It is moreover a matter of interest that of the twenty works published and purchased by him between 1650 and 1660 no less than seven were acquired by Goodyer in the year *before* publication: a clear proof that his enthusiasm for his science led him to keep in the closest touch with the booksellers.

He appears to have started the practice of dating his books on 31 January 1615 when he acquired Bauhin's 1598 folio edition of Matthioli for 20s. The subsequent dated additions to his library were as follows:

<i>Date of acquisition.</i>	<i>Price.</i>	<i>Title and Year of Publication.</i>
13 Novemb. 1616	s d 2 6	Clusius, <i>Curae posteriores</i> , 1611.
12 Decemb. 1616	16 0	{ „ <i>Rariorum plantarum</i> , 1601.
		{ Pona, <i>Monte Baldo</i> , fol. 1601.

<i>Date of acquisition.</i>	<i>Price.</i>	<i>Title and Year of Publication.</i>
	s d	
15 Feb. 1616	15 0	Clusius, Exoticorum, 1605. Paged by G.
28 Feb. 1616	3 6	Bauhin, Phytopinax, 1596. Full of G.'s notes.
10 Marcii 1616	4 0	Lobel, Plantarum, 1605.
12 Mar. 1616	4 0	2 nd part.
17 Mar. 1616	1 6	1 st part
9 April 1620	20 0	the binedinge them together. } 9 ^s 6 ^d
18 May 1623	— —	Dodoens, Stirpium, 1616.
Binding	— —	2 o. Sprecchis, Antabsinthium, 1611.
7 September 1623	15 6	Theophrastus, Opera, 1541.
13 October 1623	16 0	Columna, Minus cogn. stirpium, 1616.
30 Oct. 1623	6 0	Lobel, Kruydtboeck, 1581.
30 Novemb. 1623	received this	booke from bindinge. Bauhin, Pinax, 1623.
26 Junii 1624	3 6	Bauhin, Prodromus, 1620.
17 Novemb. 1627	4 0	Caesalpinus, De Plantis, 1583.
10 Novemb. 1627	10 0	de Passe, Hortus Floridus, 1615.
18 April 1629	9 0	Pona, Monte Baldo, 1617.
10 Novemb. 1631	0 18	Dioscorides, de curationibus, 1565.
14 Novemb. 1631	0 12	Thevet, Amerique, 1558.
13 Novemb. 1632	0 6	Plat, 1608.
2 October 1632	20 0	Matthiolus, Comment. 1583.
10 Octob. 1632	4 0	Matthiolus, Compendium 1571.
18 Maij 1633	4 6	4 ^s 7 ^d . Donati, de simplicibus di Venetia,
bindinge	0 1	1631.
28 Oct. 1634		Johnson, Mercurius, 1634, ex dono Th. Johnson.
30 April 1634	4 0	Dodoens, De Frugum. 1552.
24 Aug. 1640	36 0	39. Parkinson, Theatrum, 1640.
In quires		
The bindinge	3 0	
27 April 1641		Johnson, Mercurius Pars altera.
? 1648		Thurneiser, 1578, bound.
19 Feb. 1651	5 0	Culpeper, English Physician, 1652.
15 March 1651		Bauhin, Historia plantarum.
	li	
22 March 1651	3 2 6	Sent Dr. Dale for Johes Bauhin 3 volumes.
	1 0	for the portage to & from Dr. Dales.
	1 0	to John Symonds to carry up the money.
	1 4	to William Mychell for bringing the bookes down.
	3 5 10	
15 July 1652	3 6	Pemel, Simples, 1652.
1 Decemb. 1652	1 14 0	Hernandez, Planta Mexicanorum, 1651.
		'to a porter for carriage to Dr. Dale.
		Portage down in Mrs Elz. Heathes Trunck.—11 Dec.
11 Aug. 1653	4 6	Renealmus, Specimen, 1611. [Purch. by How.].
10 Feb. 1653	10 0	Bauhin, Theatri botanici, 1658.
7 Septemb. 1654		Duval, Phytologia, 1647.
7 Septemb. 1654	7 6	Laurenberg, Horticultura, 1654.
19 Februarij 1654		Lobel, Stirpium illustrationes, 1655.
15 June 1654	8 6	Dioscorides & Nicander, 1499.
16 Apr. 1654		Lonicer, (imp.).
30 Marcij 1654		Matthiolus, Les Commentaires, 1566.
31 Augusti 1654	2 6	Aristotle & Theophrastus, 1552.
7 Sept. 1654.		Amatus in Diosc. 1558.
14 " "	3 0	Binding Dioscorides MS.
20 Marcij 1655	1 4	Coles, Art of simpling, '56.
4 Aprill 1655	25 0	Matthiolus, Kreutterbuch, 1590.
19 Aprill 1655	3 0	Brunyer, Hort. Blesensis, 1653.

<i>Date of acquisition.</i>	<i>Price.</i>		<i>Title and Year of Publication.</i>
	s	d	
19 April 1655	2	8	Vigna, Animadversiones.
5 April 1655	2	4	Cooke, Chirurgery, 1648.
1 Junij 1655	2	4	" " Supplement, 1655.
10 May 1655	3	6	Muffet, Health, 1655.
25 May 1655	8	0	Troxiten, 1595.
25 May 1655	25	0	Dioscorides, Laguna transl. 1555.
25 May 1655	15	0	Curtius, 1560.
25 May 1655 unbound	4	6	} 5 ^s 8 ^d . Guilandinus, Theon, 1558.
28 Junij " the bindinge	1	2	
25 May 1655 in quires	12	0	} 15 ^s 4 ^d . Lonicer, Kreuterbuch, 1630.
28 Junij 1655 the bindinge	3	4	
6 Sept. 1655	2	14	0 Tabernaemontanus, 1625.
15 Julij 1657	30	0	Theophrastus, ed. Gaza, 1644.
15 Julij 1657	10	0	Moscardo, 1656.
16 Decemb. 1657	4	2	Langham, 1633.
16 Decemb. 1657			Coghan, Health, 1636.
11 Mar. 1658			Neander, Tobacologia, 1626. [Ex dono R. Downes.]
2 Dec. 1658	1	2	Everartus, Panacea, 1659.
30 Apr. 1659	3	0	Dodoens, Herbarius, 1563
30 Apr. 1659	4	0	Dorsten, Botanicon, 1540 } 'Basingstoke.'
1659			[How], Phytologia, 1650.
21 March 1660	6	0	Lovel, Hist. Animals, 1661.
10 May 1660	2	6	Ray, Cat. Pl. Cantabrigiensis, 1657.
26 Sept. 1661	4	0	Binding Theophrastus MS.

SIGNATURES OR PERSONAL MEMORANDA IN GOODYER'S BOOKS.

The numbers are the reference numbers stamped on the backs of the volumes.

<i>Hendrik Alberts.</i>	Dodoens. 11.
<i>Guilielmus Barlous Anglus.</i>	Dioscorides. 113.
<i>JB ex dono CB consang.</i>	Thurneiser. 37.
Evidently a gift from Caspar to Jean Bauhin. ¹	
<i>O. Bilson.</i> ²	Turner. 13.
<i>Henry Blount.</i>	Neander. 67.
<i>Magister Bowden.</i>	Dodoens. 11.
<i>De Brina</i>	Clusius. 106.
<i>Lancelot Browne, M.D., 1598.</i> ³	Caesalpinus. 59.
<i>F. Bust 1577.</i>	Joubert. 128.
<i>Jacob Cole.</i> ⁴	Dodoens. 96.
<i>D. Daile in Long Aker.</i>	Hernandez. 26.
<i>Dr. Dale.</i> ⁵	Hernandez. 26. Bauhin. 43.

¹ JEAN BAUHIN, 1541. CASPAR BAUHIN, 1560-1624. Converted the chaos of plant nomenclature into order in 1623. Goodyer entered biographical notes on Bauhin on a fly-leaf of his *Phytopynax*.

² O. BILSON. See p. 96.

³ LANCELOT BROWNE, M.D. Fellow of Pembroke Hall and of the College of Physicians, 1584; first physician to Queen Elizabeth. He is quoted on the subject of the Balsam tree by Gerard, to whose *Herbal* 1597 he contributed a eulogistic epistle. He died in 1605.

⁴ JAMES COLE. Son-in-law of Lobel, see p. 247.

⁵ DR. DALE. See p. 294.

<i>Ric. Downes</i> . ¹	Neander. 67.
<i>Joannes Freame</i> .	Kyber. 105.
<i>Ricūs Garth</i> . ²	Caesalpinus. 59.
<i>Rob. Garth, Juris Nat. Consult.</i> <i>25 Martij 1598.</i>	Caesalpinus. 59.
<i>Alexis Gaudin</i> .	Dioscorides. 5.
<i>Rich. George, pharm. of Reading.</i>	Dodoens. 11.
<i>Jl. Gilbou[rne]</i> ?.	Porta. 130.
<i>John Gooche</i> .	Lonicer. 14.
<i>Jo: Gæoodier</i> .	Monardus. 126.
<i>Johes Goodyer</i> .	Passe. 81.
<i>E. Gray, 'heboriste'</i> . ³	Turner. 13.
<i>Hen. Harvey et amic.</i>	Stephanus. 95.
<i>Tho. henry</i> .	Cooke. 118.
<i>W. Howe</i> .	{ Matthiolus. 38. Renealmus. 14.
	{ Johnson. 99. Tabernaemontanus. 46. How MS. 18.
<i>R. Huchenson</i> .	Dodoens. 97.
<i>Susan Ironsmith</i> .	Matthiolus. 32.
<i>Thomas Johnson</i> . ⁴	Johnson. 105. Brunfels. 29.
<i>Johan Jul.</i> ?	Fuchs. 35.
<i>Bartho: Kempc</i> .	Dodoens. 12.
<i>G. Le Fevre</i> . ⁵	Hess. 100.
<i>Matth. de Lobel</i> . ⁶	Lobel. 17.
<i>Alexander Massa</i> .	Fuchs. 112.
<i>Geo. Medeleye</i> .	Dorsten. 9.
<i>W. Mount</i> . ⁷	Lobel. 79.
<i>Dr. Martin Ramerius</i> . ⁸	Lobel. 17.
<i>G. Rondelet</i> .	MS. 13.
<i>Anth. Rous</i> .	Soranus. 6.
<i>Antony Swalms</i> .	Fuchs. 8.
<i>H. W.</i> (? Henry Wotton, M.D.) ⁹	Ruellius. 34.
<i>John Yates, barber & chirurgion</i> .	Dodoens. 10.

¹ The SAMUEL DOWNES, M.D., who made the collection of dried plants presented by J. Downes in 1731 to Shrewsbury School, may have been related to Richard Downes.

² RICHARD GARTH. See p. 237.

³ E. GRAY, 'heboriste'. It would be interesting if he should turn out to be Gray the apothecary (fl. 1570), who introduced *Pistacia officinalis* (Lobel, *Adv.* 413) and had trees of *Diospyros Lotus* L. and of *Celtis australis* L. in his garden under London Wall (*Ger.* 1308, 1310).

⁴ THOMAS JOHNSON. See p. 273.

⁵ G. LE FEVRE. At a later date two of this name appear on the Roll of the R. College of Physicians. Sebastian Le Fevre, L.R.C.P., 1684, Joshua Le Feure, F.R.C.P., 1687.

⁶ LOBEL. See p. 246.

⁷ W. MOUNT. See p. 253.

⁸ MARTIN RHAMNEIRUS, M.B., a Spaniard, a native of Cordova and a Bachelor of Medicine, was admitted a Licentiate of the R. College of Physicians, 1584.

⁹ HENRY WOTTON, M.D., Student of Christ Church and Fellow of Corpus Christi College, Oxford, F.R.C.P., 1571-2.

PERSONS FROM WHOM BOOKS WERE ACQUIRED.

<i>Vendor.</i>	<i>Carrier.</i>	<i>Name of work.</i>	
<i>Dr. Dale</i>	<i>Eliz. Heath</i>	Hernandez. 26.	
<i>Dr. Dale</i>	} <i>William Mychell</i>	} Bauhin, Hist. Pl. 43-45.	
<i>Humphrey Robinson</i> ¹			} <i>John Symonds</i>
<i>Bernard Robinson</i>		Bauhin, Theatr. 52.	
<i>John Martin</i>		Tabernaemontanus. 46-47.	
<i>Octavian Pulleyn</i> ²		Matthiolum. 38.	
<i>William Wells</i>		Rencaimus. 68.	
<i>Mr. Allestre</i>			
<i>Donor.</i>			
<i>Richard Downes</i>		Neander. 67.	
		Rencaimus. 68.	
<i>Dr. William How</i>		} Tabernaemontanus. 46-47.	
Perhaps a vendor			Matthiolum. 38.
			How & Lobel. 72.
			How, Phyt. MS. 18.
<i>Thomas Johnson</i>		Johnson. 99.	
		Mercurius. 105.	

BOOKS CONTAINING GOODYER'S NOTES.

Goodyer wrote notes in almost all the books with which he worked. Chief of these were

Tabernaemontanus, *Eicones*, 1590.

[Lobel], *Icones*, 1591.

Bauhin, *Phytopinax*, 1596.

And doubtless his copy of Gerard's *Herbal*, 1597, was similarly annotated and corrected, but it is unfortunately no longer in the Library.³ He numbered the pages or chapters of

Bauhin, *Hist. plantarum*, 1650. G. numbered 6524 columns.

Besler, *Hort. Eystettensis*, 1613. To p. '854'.

,, *Fasc. rariorum*, 1616.

Clusius, *Exotica*, 1605.

Dioscorides, 1499.

Theophrastus, 1497.

He numbered or named plants described or figured in

Bauhin, *Animadversiones*, 1601.

¹ HUMPHREY ROBYNSON of London appears in a book of Richard Napier's astrological practice in 1606. *MS. Ashmole* 181.

² OCTAVIAN PULLEYN described himself as *mercator librorum Italicorum* in an inscription inside the cover of a book (Tower F. 1) given by Sam. Lee to the Wadham library. An Octavius Pullin (? the same man) had been a student at Padua in 1638-9 (Andrich, *De Natione Anglica*, Pataviis 1892). See p. 294.

³ I shall be most grateful for any information which will enable me to trace the whereabouts of the lost volume. It should be readily recognizable by Goodyer's notes.

Johnson, Iter, 1629.	de Passe, Hort. Floridus, 1615.
„ Mercurius, 1634.	Thalius, Sylva Hercynia, 1588.
„ „ 1641.	Tragus, 1552.

Goodyer was an indefatigable indexer. In some cases, in which several works are bound up together in the same cover, he not only corrected the pagination of the separate works, but carried the page-numbers right on. He made indexes to the combined works of Turner (No. 13), to de Bry's *Anthologia*, 1626, and to Clusius, *Rariorum Plantarum* (MS. 8. iii). His index to Gerard's *Herbal* (1597) is in a small 8vo parchment-bound volume (MS. 16). The manuscript indexes to the following works are bound with MS. 11.

Johnson, *Mercurii* 1 and 2, MS. 11, ff. 29-32.
How, *Phytologia*, MS. 11, ff. 33-7.

CATALOGUE OF THE GOODYER LIBRARY.

The notes quoted in small type are mostly in Goodyer's hand.

ALBUCASIS.

- 6 Libri tres Chirurgicorum. fol. Argent. Schott 1532
Bound with HORATIANUS.

ALPINUS, PROSPER.

- 61 { De plantis Aegypti. } 4. Ven. 1592
 { De balsamo dialogus. }
 With notes to plates.
- 62 De plantis exoticis, libri duo; ed. ab Alp. Alpino auctoris
 fil. 4. Ven. 1627
 With G.'s references to Parkinson.
- 92 De rhapontico disputatio. 4. Patav. 1612
 ' Bauhinus transmisit novā Piperā Indicam—Pinax 103 (a 7).'

AMATUS, LUSITANUS.

- 82 In Dioscoridis . . . de Materia Medica. 8. Lugd. 1558
 '7 Septemb. 1654—58.'

APULEIUS, LUCIUS.

- 86 De medicaminibus herbarum, lib. I. Edited with a com-
 mentary by Gabriel Humelberg. 4. Isinae 1537
 De herbarum virtutibus historia; at end of SORANUS, q.v.
 [fol. Basil. 1528]

ARISTOTLE.

- 2 Problemata, Mechanica, Metaphysica. fol. Ald. 1497
 Bound with Theophrastus, De historia plantarum.
- 138 Historia animalium etc. 8vo. Lugd. 1552

ALEXANDER APHRODISIENSIS.

- 2 Problemata. fol. Ald. 1497
 Bound with Theophrastus, De historia plantarum.

BAUHINUS, CASPAR.

- 55 Φυτοπινναξ, seu Enumeratio plantarum ab herbariis nostro saeculo descriptorum cum earum differentiis etc. et cum iconibus; [libri octo]. 4. Basil. 1596
 '28 febr 1616—3^s 6^d.'
 Numerous notes by G., including notes on Bauhin's life.
- 85 { Animadversiones in historiam generalem plantarum Lugduni editam. Catalogus plantarum circiter quadragintarum eo in opere bis terve positarum. 4. Francof. 1601
 { De homine oratio in medicorum Lycaeo 1614. 4. Athen. Rauracis [1614]
 Plants numbered by G.
- 109 De remediorum formulis Graecis, Arabibus et Latinis usitatis. 12. Francof. 1619
 De corporis humani partibus externis. 12. Basil. 1588
- 69 Προδρομος theatri botanici. 4. Francof. 1620
 '26 Junii 1624—3^s 6^d.'
- 56 Πινναξ theatri botanici; sive index in Theophrasti, Dioscoridis, Plinii et aliorum qui de plantis scripserunt; [libri duodecim]. 4. Basil. 1623
 'Received this booke from bindinge 30 Novemb. 1623.'
 Interleaved. A few notes on pp. 2, 52, 96, last leaf.
- 123 Catalogus plantarum circa Basileam sponte nascentium. 8. Basil. 1622
- 52 Theatri Botanici, sive historiae plantarum liber primus. fol. Basil. 1658
 '10 febr. 1653—10^s a Johanne Martin.' Marginal references by G.

BAUHINUS, JOANNES.

- 109 { De plantis a divinis sanctisve nomen habentibus. 8. Basil. 1591
 { De plantis absynthii nomen habentibus. 8. Montis Belg. 1593
- 88 { De aquis medicatis libri quatuor. } 4. Montis Belg. 1612
 { De variis fossilibus, stirpibus et insectis. }
 Reference on p. 194 by G.
- 71 Historiae plantarum generalis prodromus. 4. Ebrod. 1619
 (With CHERLER, J. H.)
 Reference on p. 71 by G.
- 43-5 Historia plantarum universalis, quam recensuit et auxit Dom. Chabraeus, juris publici fecit Fr. Lud. à Graffenried. 3 vols. fol. Ebrod. 1650-1
 'March the 15th 1651
 bought of my Master Humphrey Robinson
 the 3 vollumes of J. Bauhinus which I do warrant
 to be pfect per me Bernard Robinson.'
 Columns numbered 1 to 2064 and 1 to 2172 and 1 to 2288.

Inscription by Goodyer

1651 22 March

sent Dr. Dale for Johes Bauhins 3 volumes	3 ^{li} 2 ^s 6 ^d
for the postage to and from Dr. Dales	o 1 o
to John Symonds to carry up the money	o 1 o
to William Mychell for bringing the bookes down	o 1 4

£3 5 10

BAYFIUS, *see* STEPHANUS, C.

BELLONIUS, P.

41 Observaciones.

Translated & edited by C. Clusius. fol. Plantin, 1605

BELLUS, HONORIUS.

39 Epistolae de rarioribus quibusdam plantis agentes. fol. 1601
(With CLUSIUS, No. 39.)

BESLER, BASIL.

29* Hortus Eystettensis. large fol. s.l. 1613

Paged by G. to 854. Plant names written on some plates, e.g. 540.

80 Fasciculus rariorum et aspectu dignorum, varii generis etc.

fol. Norib. 1616

Pages numbered.

BLESENSIS, *see* BRUNYER.

BOCK, H., *see* TRAGUS.

BRASAVOLUS, ANT. MUSA.

103 Examen omnium simplicium medicamentorum ; with Aristotle's Problemata quae ad stirpium genus et oleracea pertinent. 8. Lugd. 1537

86 De herba Vetonica ; with a commentary by Gabriel Hummelberg ; with L. Apuleii libro De medicaminibus herbarum. 4. Isin. 1537

Two ff. of Latin MS. with marginal notes. No notes by G. A bill of 4 items on p. 1 ' Coyse 3^s 0^d, Fan 3^s 2^d, Apuleius 2^s 0^d, Plato 0^s 6^d '.

BREDA, *see* BROSTERHUSIUS.

BROSTERHUSIUS, JOH.

134 Catalogus plantarum horti medici illustris scholae Auriacae, quae est Bredae. 8. Bredae, 1647

BROSSE, GUY DE LA.

71 Description du jardin royal des plantes médicinales.

4. Par. 1636

108 De la nature, vertu et utilité des plantes, divisé en cinq livres.

8. Par. 1628

On back of title ' 1^{lb} 0 '.

BROWNE, W., *see* STEPHENS, P.

[BROWNE, SIR T.

Hydriotaphia. *Missing*.

London, 1658]

BRUNFELS, O.

- 29 Onomasticon medicinae continens omnia nomina herbarum.
fol. Argent. ap. Jo. Schottum, 1534
'20 Nov. 1634 Tho. Johnson bought him for 3^s.'
Bound with Hildegard, *Physicae*. G. has added names to the list
of writers on fo. k iii.
On the title-page is an inscription in the handwriting that we have
called that of Dr. Dale, who may have owned the book after Johnson's
death in 1644.
- 7 Herbarium. Vol. 1. Herbarium. 2. Novi Herbarii, 1536.
3. Tomus Herbarii III, 1536. Appendix, 1539.
fol. Argent. ap. Jo. Schottum, 1537
'1537 Hen. 8^{vi} 28.'

BRUNYER, ABEL.

- 20 Hortus regius Blesensis. fol. Par. 1653
'19 Aprill 1655—3^s.'

BRUYERINUS, JOANNES.

- 124 De re cibaria, lib. xxii. 8. Francof. 1600

BRY, J. T. DE.

- 25 Anthologia Magna. fol. Francof. 1626
MS. Index by G.

[BUMALDUS, MONTALBANUS.

Bibliotheca Botanica. *Missing*.

12 Bononiae, 1657]

CAESALPINUS, ANDREAS.

- 59 De plantis, lib. 16. 4. Florent. 1583
'17 Nov. 1627—4^s.'

On title:—'Ricū Garth praeciū x^s

Sum Lanceloti Brunii medici Reginei ex dono amicissimi viri
Mri. Rob. Garthii Juris Nationalis Consult 25 Martij 1598.'

2 Titles of books (Durante & Rondelet) noted.

- 65 De metallicis, lib. 3. 4. Romae, 1596
Ref. at end to p. 152 'liber de questionibus Peripat. c. Magnet'.
In binding is a writing relating to — Scott of Sutton, 1608.

CAESAR, *see* CORNARIUS.

CALCEOLARIUS, F.

- 58 Iter Baldi. 8. Francof. 1586
With Matthiolus.

CAMERARIUS, JOACHIMUS.

- 85 Εκλεκτα γεωργικα, sive opuscula de re rustica, partim collecta,
partim composita a J. Camerario. 4. Norib. 1577

- 89 Hortus medicus et philosophicus, in quo plurimarum stirpium breves descriptiones etc. continentur. 4. Francof. 1588
- 89 Icones accurate delineatae praecipuarum stirpium, quarum descriptiones tam in Horto quam in Sylva Hercynia suis locis habentur. 4. Francof. 1588
References to 'Icones'.
Kreutterbuch 1590, see MATTHIOLUS, No. 38.
- [CATO, M. PORCIUS.
De re rustica. *Missing*. 12 Col. 1536]
Probably disposed of as a duplicate of the copy presented by A. Throkorton.
- CIEMNITZ, JOANNES.
92 Index plantarum circa Brunsvigam nascentium: Appendix continens icones plantarum. 4. Brunsvigae, 1652
- CHERLERUS, see BAUHIN, J.
- CLAVENAS, NICOLAUS.
92 { Historia absinthii umbelliferi. }
{ Historia scorzonerae Italici. } 4. Venet. 1610
- CLUSIUS, CAROLUS.
107 Historia rariorum aliquot stirpium per Hispanias observatarum. 8. Ant. 1576
- 106 Historia rariorum aliquot stirpium per Pannoniam, Austriam et vicinas quasdam provincias observatarum. 8. Ant. 1583
Stirpium nomenclator Pannonicus (Beithii). „ 1584
On title, 'huius libri pretium—3^s 4^d. De Brina'. Name of an owner erased. On last p. 'Costa soldi tre, dico 53 moneta sterlinga 1583, 25 Decemb.'
- 41 Exoticorum libri decem, quibus animalium, plantarum, aromatum aliorumque peregrinorum fructuum historiae describuntur; item P. Bellonii observationes C. Clusio interprete. fol. Plantin. 1605
With a copy of the *Curae posteriores*, and *Oratio funebris* at end. '15 feeb 1616—15^s.'
Pages numbered by G., beginning p. '825'. Marginal references.
- 39 { Rariorum plantarum historia. }
{ Fungorum in Pannoniis observatorum historia. } fol. Ant. 1601
This work has been repaged by G., who also compiled an index to it.—Goodyer MS. 8.
- 39 Curae posteriores. fol. Plantin, 1611
With G.'s references to *Phyto*. and *Pinax*.
'13 Novemb. 1616. Cur. poster. 2^s 6^d.
12 Decemb. 1616 16^s.'
- 41 Oratio funebris in obitum C. Clusii (by E. Vorst). fol. Plantin, 1611

COGHAN, THOMAS.

- 132 The Haven of Health, chiefly gathered for the comfort of students; hereunto is added a preservation from the pestilence, with a short censure of a late sickness at Oxford. 4th edit. 8. Lond. 1636
'16 Decemb. 1657.'

COLES, W.

- 121 The art of simpling; an introduction to the knowledge of gathering plants; whereunto is added a discovery of the lesser world. 8. Lond. 1656
'20 Marcij 1655—1^s 4^d.'
In binding a fragment of deed mentioning 'Lady Jane late Queene' and 'Lady Ann' at 'Bower'.

COLLAERT, ADRIAN.

- 80 Florilegium. oblong 4^{to}. n. d.

COLUMELLA, LUCIUS J. M.

- 110 De cultu horticorum. 12. Argent. 1530

COLUMNA, FABIUS.

- 87 *Φυτοβασανος*, sive plantarum aliquot historia; accessit etiam piscium aliquot historia. 8. Neap. 1592
Refs. to *Panax*.

- 70 { Minus cognitarum rariorumque nostro coelo Orientium stirpium *εφραυσις*, item de aquatilibus aliisque animalibus quibusdam paucis libellus. In two parts. 4. Rom. 1616
Purpura; et aliorum aquatiliu[m] observationes. „ „
'13 Octob. 1623—16^s.'

'Fabius Columna was but 25 yere old when he sett forth his *Phutobasanos* p. 210 & he set forth his P. ten yers before he wrote this story, p. 219. Diosc. graecus codex MS. p. 71.

Johannis Carbonarius of Naples hath many old Manuscripts p. 71. Johannis Baptista Raimundi idem p. 72, 142.' G.'s notes.

CONTARINI, *see* PONA.

COOKE, JAMES.

- 118 Melleficium chirurgiae, or the marrow of many good authors on the art of chyrurgery. 8. Lond. 1648
'5 Aprill 1655—2^s 4^d.'
Name of former owner 'Tho' 'Thenry' on front page.
'Frog-spawne, p. 431' } on back cover in G.'s hand.
'Lucatellas Balsam, p. 467,' }
- 119 Supplement to the marrow of chyrurgerie. 8. Lond. 1655
'1 Junij 1655—2^s 4^d.'
'Swellings of the lunges 228.' 'Ricketts—243.'
[Select observations on English bodies. *Missing*. 8. 1657]

- CORDUS, EURICIUS.
 102 Botanologion. 8. Col. ap. J. Gymnicum 1534
- [CORDUS, VALERIUS.
 Annotationes in Pedacii Dioscoridis Anazarbei de materia
 medica libros quinque.
 Historiae stirpium, cum iconibus, libri quatuor.
 Sylva de fossilibus in Germania.
 De artificiosis extractionibus.
 Compositiones aliquot medicinales rariores. fol. Argent. 1561]
 Missing: perhaps duplicated by the copy in the Gibbarde Col-
 lection and sold.
- CORNARIUS, J.
 123 Constantini Caesaris selectarum preceptorum de agricultura
 libri viginti, interprete Jano Cornario. 8. Basil. 1540
 With marginal notes (cut by binder) in early hand.
 'DUUM em C. h. g. o' on last p.
 'the orderig of vynes // liber quint / fo.'
- CORNUTUS, JACOBUS.
 61 Canadensium plantarum aliorumque nondum editarum his-
 toria; Enchiridion botanicum Parisiense. 4 Par. 1635
 Notes to figures: refs. to Parkinson.
- COSTAEUS, JOANNES.
 90 De universali stirpium natura. Libri duo. 8. Aug. Taur. 1578.
- CRESCENTIUS, P.
 1 De plantis, animalibus et agricultura. fol. Lovan. 1473
 In contemporary Oxford stamped leather binding.
 Pagination added by G. MS. list of ten goodwives. See p. 381.
- CUBA, JOHANNES DE.
 9 Hortus sanitatis de animalibus et reptilibus, de avibus et
 volatilibus de piscibus et natatilibus, de gemmis et in
 venis terrae nascentibus.
 fol. Argent. p. Matth. Apiarium 1536
 With Dorsten.
- CULPEPER, NICH.
 23 The English Physician. fol. London 1652
 '19 ffeb. 1651—5^s.'
 'Vale et diu vive.'
- CURTIUS, BENEDICTUS.
 36 Hortorum, libri triginta. fol. Lugd. 1560
 '25 May 1655—15^s.'

DALECHAMPIUS, JACOBUS.

- 135-6 *Historia generalis plantarum.* 2 vols. fol. Lugd. 1586-7
The copy in the Sherardian Collection in the Botanic Garden has the name of Goodyer's great friend *William Coys* written on the title-page.

DIOSCORIDES, PEDACIUS.

- 4 *Opera Gr.* fol. Ven. Ald. 1499 Ed. pr.
'15 Junii 1654—8^s 6^d.'
Paged, and chapters numbered.
- 5 *Opera trad. de lingua Griega en la vulgar Castellana por Andres de Laguna.*¹ fol. Anvers. 1555
'25 May 1655—25^s.'
'Alexis Gaudin' (owner's signature on title). Silver stamp on binding.
- 113 *De materia medica, ab Andr. Matthiolo emendata.* 12. Lugd. 1554
'Guilielmus Barlous Anglus 6½ bats 1567.'
Prescription 'contra pestem D. Cratonis' on last page.
- 83 *De medicinali materia Gr. Lat.* 8 libb. cum castigationibus. 8. Paris 1549
Lat. transl. by J. Ruellius.
With Paragraphs numbered throughout and cross references to Matthiolus by Goodyer.
- 73 { *De simplicibus medicina.* } 4. Lugd. 1512
 { *De naturis et virtutibus aquarum.* }
 With medical prescriptions at end in an old hand.
- 84 *De curationibus morborum per medicamenta paratu facilia* lib. 11. 8. Argent. 1565
 With Latin translation, partly by J. Moibanus, partly by C. Gesner.
 '10 Nov. 1631—18^d.'
 A few marginal notes by G., pp. 258, 593. At end 'Antidotus Saxonica 830. Inflatio stomachi, 533.'
 On title initials of former owner 'RS.' In binding, part of an English theological MS. on parchment.

DODOENS, REMBERT.

- 96 *De Frugum Historia.* 12. Antv. 1552
'30 April 1634—4^s.'
'Jacobi Colei.' His marginal notes on p. 14 v.

¹ In the copy of this work in the Botanical Department of the British Museum I found an inscription, presumably by the translator:—

Doctor Andreas Lagouna hunc librum Johanni
Mutier amico suo, dono dedit anno 1557.

- 10 L'histoire des plantes ; traduite de bas Aleman en François
par Charles de l'Escluse. fol. Anvers. 1557
With G.'s references and many old notes. At end 'John Yates,
barber & chyrrurgion'. 'hope helps hevie harts.'
Of *Lentiscus* he notes:
'I se thys tre at benys marks in London at one Mr. Hennege hys
howse of the previ chamber to quen elizabeth.' p. 547.
'The leves of saven mayd in powder & droken with alle kylleth
worms in the bellye.
'Also y^e sayd powder cast uppon warts in ye yard of man kylleth
ye warts, & if it be swelled it wyll dyssolve it.' p. 538.
- 11 Herbarius oft Cruydt-boeck. fol. Antv. 1563
'30 Apr. 1559—3^s. Basingstoke.'
'Hendrick Alberts—1570' on last page.
'Richardus George pharmacopeus de Reading est verus possessor
huius libri, ex dono Magistri Bowden, Julij 9 die 1619.'
With many MS. translations into English of the Dutch text, which
Mr. George evidently had difficulty in comprehending.
- 12 A neewe herball or history of plantes ; first set forth in the
Doutche or Almaigne tongue, and nowe first translated
out of French into English ; by Henry Lyte.
fol. Lond. Gerard Dewes, 1578
'Bartho: Kempe.'
- 40 Cruydt boeck, met büvoegsels achten elck capittel . . .
Carolus Clusius. fol. Leyden, 1608
- 42 Stirpium historiae pemptades sex, sive libri triginta.
fol. Antv. 1616
'9 April 1620—20^s. the carrier 1^s.
With a few references to Phyto. (p. 125) by Goodyer.
- 97 Purgantium, aliarumque eo facientium, tum et radicum,
convoluulorum. 8. Antv. 1574
'R. Huchenson.'
- DONATI, ANTONIO.
- 92 Trattato de semplici, pietre et pesci marini, che nasceno nel
lito di Venetia. 4. Venetia 1631
'18 Maij 1633—4^s 6^d }
bindinge 1^d } 4^s 7^d.'
- DORSTENIUS, THEODORICUS.
- 9 Botanicon, continens herbarum aliquotque simplicium,
quorum usus in medicinis est, descriptiones et iconas ad
vivum effigiatas. fol. Franc. Christ. Egenolphus 1540
'30 Apr. 1659—4^s.' Basingstoke. Many old notes.
Name of earlier owner 'George Medeleye, vi^s viii^d' on title.
- DURANTE, CASTOR.
- 21 Herbario novo, con figure. fol. Rom. 1585

DUVAL, GUILLELMUS.

- 98 Phytologia ; sive philosophia plantarum. 8. Par. 1647
 '7 Septemb 1654.'

EDER, GULIELMUS.

- 110 Synonyma Plantarum seu Simplicium ut vocant, circa In-
 goldstadium sponte nascentium . . . in usum Scholae
 Medicae Ingolstadiensis collecta. 12. Ingolstadii, 1618

EGENOLPH, C.

- 87 { Plantarum arborum fruticum et herbarum effigies . . . Bäume,
 Stauden, Kreuter, etc., cum Indice sextuplici.
 8. Franc. C. Egen. 1562
 Animantium terrestrium, volatilium et aquatilium effigies.
 19 pp. 8. Franc. C. Egen. 1562
 At end is a Petition of Richard Chambers to Parliament c. 1644.

E[VELYN], J.

- 24 Sylva, or a Discourse of Forest Trees. fol. Lond. 1664

EVERARTUS, AEGIDIUS.

- 120 Panacea, or the universal medicine ; being a discovery of the
 wonderful vertues of tobacco taken in a pipe, with its
 operation and use both in physick and chirurgery ; transl.
 from the Latin with an introductory epistle : by J. R.
 8. Lond. 1659
 '2 Dec. 1658—1^s 2^d.
 Purchased by Goodyer before publication.

FERRARIUS, J. B.

- 74 De florum cultura 4 libb, editio nova accurante Bern.
 Rottendorffio. 4. Amst. 1646

FIERA, J. BAPTISTA.

- 110 Coena de herbarum virtutibus, et ea medicae artis parte,
 quae in victus ratione consistit. 12. Argent. (1530)
 Old notes on p. 22 and at end.

FRAGOSUS, JOH.

- 95 Aromatum, fructuum et simplicium aliquot medicamentorum
 ex India utraque in Europam delatorum, historia brevis,
 Latine reddidit Israel Spachius. 12. Argent. 1601

FUCHS, LEONHART.

- 112 De historia stirpium commentarii, cum imaginibus ad naturae
 imitationem effectis. 12. Lugd. 1555
 'Alexandri Massa' '47^s 6^d.' English names of plants in margins.
 35 Ditto. fol. Basil. Isingrin, 1542
 'Johannis Jul . . .' 'Hen. 8. 34.'

- 8 New Kreüterbüch. fol. Basell. Isingrin, 1542
Coloured copy.
Imperfect, title missing. Inside cover many verses and mottoes
in French, quoted from Rombout Martens and others.
At end 'Sum Antony Swalms et Amicorum'.
- GANSIUS, JOH. LUDOVICUS.
- 102 Corallorum historia. 8. Franc. 1630
'Charle Knolfts' at end.
- GESNER, CONRAD.
- 117 Historia plantarum et vires ex Dioscoride, Paulo Aegineto,
Theophrasto, Plinio etc. 12. Par. Jo. Roigny 1541
- 102 Ditto. 8. Basil. Ro. Wynter 1541
- 101 Apparatus et delectus simplicium medicamentorum.
8. Lugd. 1542
- 93 Catalogus plantarum, Latine Graece Germanice et Gallice.
4. Tiguri 1542
In binding are two copies of an Order to Churchwardens and
Constables of Parishes to assess sum due for relief of maimed
souldiers of City of London according to Act of Parliament 43 Eliz.
and to pay same to W. Antrobus, Treasurer, at his house in Parish
of St. Gregory, S. Pauls Churchyard.
Mizaldus 104 contains a copy of the same document.
- 109 Epistolae aliquot a C. Bauhino editae. 8. Basil. 1591
Praefatio de rei herb. scriptores :
Preface to TRAGUS, *de Stirpium*, q. v.
- GERARD, JOHN.
[Herbal. *Missing*. fol. Lond. 1597]
- 50 The herball, or general historie of plants; very much en-
larged and amended by Thos. Johnson. fol. Lond. 1633
- 110 GUEROALDUS, GUILLERMUS, *see* MACER.
- GUILANDINUS, MELCHIOR
- 91 { Theon, seu apologiae adversus Pet. Andream }
Matthaeolum. } 4. Patav. 1558
{ De Stirpibus aliquot epistolae quinque; item }
{ descriptio manu-co-diatae, seu aviculae Dei. }
- Commentarius in C. Plinii Majoris capita aliquot.
3rd edit. 4. Lausaniae 1576
'25 Majij 1655, vnbound 4^s 6^d }
28 Junij 1655, the bindinge 1^s 2^d } 5^s 8^d.
- 122 Papyrus, hoc est commentarius in tria C. Plinii majoris de
papyrocapita, recensente Henrico Salmuth.
12. Ambergae 1613

- 75 HERBARIUS in Latino cum figuris. 4. s. l. et a.
[? 1485]

In original binding. English plant-names written in.

HERNANDEZ, FRANCISCUS.

- 26 Nova plantarum, animalium et mineralium Mexicanorum historia; in volumen digesta per N. Ant. Recchum, cum iconibus; et cum notis et additt. J. Terentii, J. Fabri, et Fab. Columnae, cumque aliquot tabulis phytosophicis principis Federici Caesii. fol. Romae 1651
 '1 Decemb. 1652 11i 14^a o^d
 to a porter for carriage to Dr. Dale o o o
 11 Decemb. portage downe in Mrs Elz. Heathes Trunck o.
 In another hand:
 'D. Daile in Longe Aker over agst ye freinte ordinary.'

HESS, PAUL.

- 100 Defensio XX Problematum Melchioris Guilandini adversus quae Petr. Andreas Mattheolus ex centum scripsit.
 'G. Le Fevre.' 12. Patav. 1562

HESYCHIUS.

- 3 Dictionarium. fol. Hagenoae 1521

HILDEGARD.

- 29 Physicae. fol. Argent. 1533
 Bound with Brunfels, *Onomasticon*, q.v., and therefore presumably one of T. Johnson's books.

HORSTIUS, JACOBUS.

- 94 { Herbarium Horstianum, seu de selectis
 plantis et radicibus, duobus libris edente
 Greg. Horstio.
 Opusculum de vite vinifera ejusque parti-
 bus.
 Appendix cultori plantarum exoticarum
 necessaria. } 8. Marpurg 1630

HORTO, GARCIA AB. (*G. del Huerto.*)

- 128 Aromatum, et simplicium aliquot medicamentorum apud Indos nascentium historia; ex lingua Lusitanica cum notis per Car. Clusium. 8. Antw. 1574

[How, W.]

- MS. 18 Phytologia Britannica.
 'Rec. 30 Apr. 1659.'
 The Author's interleaved copy with many notes by himself and by G. This work is described in detail on pp. 276-294.

JOHNSON, THOMAS.

- 92 Iter plantarum investigationis ergo susceptum a decem sociis
in agrum Cantianum A.D. 1629. Ericetum Hamstedianum.
4. s. l. (1629)
With list of plants added by Goodyer.
- 99 Descriptio Itineris plantarum investigationis ergo suscepti
a decem sociis in agrum Cantianum A.D. 1632. Ericetum
Hamstedianum, sive plantarum ibi crescentium observatio
habita. 12. s. l. 1632
The author's own copy with his MS. index and additional notes by
How, MS. 19. See pp. 232 and 277.
- 50 The Herball of Gerard 'very much enlarged and amended'.
fol. Lond. 1633
- 105 { Mercurius botanicus.
De thermis Bathonicis tractatus. } 8. Lond. 1634
'28 Octob. 1634.
Ex dono Thomae Johnson.' G.
Plant names picked out with yellow colour.
- 95 Mercurii Botanici, pars altera. 12. Lond. 1641
'27 Aprilis 1641.'
With G.'s notes. Several plant names picked out in colour.

JOUBERT, LAURENT.

- 128 { Medicinae practicae priores libri tres, edit.
tertia. } 8. Lugd. 1577
{ Isagoge therapeutices methodi.
'F Bust 1577° ptium iijs vid.'

KYBERUS, DAVID.

- 105 Lexicon rei herbariae trilingue. 8. Argent. 1553
With C. Gesner's *Tabulae* at end.
'Joannis Freame est verus possessor huius libri.'
Notes in his hand (?) on pp. 302, 303.

LANGHAM, WILLIAM.

- 76 The garden of health ; 2nd edit. 4. Lond. 1633
'16 Decemb. 1657 pretiū 4^s 2^d.'

LAURENBERGIUS, PETRUS.

- 63 Horticultura, duobus libris. 4. Franc. ad Moen. 1654
Apparatus plantarius primus, duobus libris, i. De plantis
bulbosis ; ii. De plantis tuberosis. 4. Franc. ad Moen. 1654
'7 September 1654—7^s 6^d.'
End papers cut from a deed mentioning John Smith, Thomas
Marshall, and Thomas Read of Parish of St. Mary Hill.
Signature of Ja. Ireland.

LEMNIUS, LEVINUS.

- 129 De plantis sacris. See VALLESIIUS. 8 Lugd. 1652

LEONICENUS, NIC.

- 86 De Plinii et aliorum in medicina erroribus liber; accedunt de herbis et fruticibus, animalibus, metallis, serpentibus, tiro seu vipera. 4. Basil. H. Petrus 1529

LERIUS, JOAN.

- 125 Historia navigationis in Brasiliam, quae et America dicitur. 8. Genev. 1586
In original vellum binding, with title on back of binding in Goodyer's hand.

LOBEL, MATTHIAS DE.

- 17 Plantarum seu stirpium historia: cui annexum est adversariorum volumen per M. de L. et P. Penam.

fol. Antv. 1576

'D. Doct. Martino Ramerio veteris amicitiae et perpetuae memoriae ergo d. dedit Matth. de Lobel Insulanus.' On title.

- 18 { Ditto—with Animadversiones in G. Rondeletii methodicam pharmaceuticam officinam. fol. Lond. 1605
Balsami, opobalsami, carpobalsami et xylobalsami cum suo cortice explanatio.

'10 Marcij 1616 Adversa. 2^{da} pars—4^s

12 Mar. 1616 Adversar. 1^a pars—4^s

17 Mar. 1616 the binedinge them together—1^s 6^d.' } 9^s 6^d.'

- 19 Kruydtboeck, oft beschrijuinghe van allerleye ghewassen, kruyderen, hesteren ende gheboomten.

fol. Antv. Plantin. 1581

'30 Oct. 1623—6^s.'

- 79 Plantarum seu Stirpium Icones. 2 vols.

Obl. 4. Antv. Plantin. 1581

Purchased by Gulielmus Mowntius (= William Mount) for 9^s on May 20, 1582, and inscribed with numerous notes in his hand. The binding bears his stamp 'W. M.'. It was acquired by Goodyer before 1633, probably long before.

A few notes in another hand are extracts from W. Bullein's *Bulwarke of Defence against all Sicknesse*, 1562.

- 78 Icones stirpium seu Plantarum tam exoticarum quam indigenarum. In 2 parts, with an index.

Obl. 4. Antv. Plantin. 1591

With full notes in Goodyer's hand, including transcripts of Mount's notes in the 1581 edition (No. 79). On p. 647 Goodyer noted that 'Potatoes' = Batata Hispanorum, Camotes sive Amotes et Iguanes. He acquired this volume before 1633.

- 72 Stirpium illustrationes, plurimas elaborantes inauditas plantas subreptitiis Joh. Parkinsoni rapsodiis sparsim gravatae; ejusdem adjecta sunt ad calcem Theatri botanici ἀμαρτηματα; accurante Guil. How.

4. Lond. 1655

'Rec. 19 february—1654.'

LONITZER, ADAM.

- 14 Naturalis historiae opus novum ; 2 voll. fol. Francof. 1551
'Joh. Gooche.'
- 15 Kreuterbuch. fol. Francof. 1557
'Title missing.
'Rec. 16 Apr. 1654.'
- 16 Kreuterbuch der bäume, stauden, hecken, kräuter, etc. ; item
von den fürnembsten gethieren der erden, vögeln, fischen
und gewürm ; deszgleichen von metallen, ertze, edelge-
steinen ; corrigirt und verbessert durch Pet. Uffenbachium.
fol. Franc. 1630
'25 Majj 1655 in quires 12^s }
28 Junij 1655 the bindinge 3^s 4^d } 15^s 4^d.'

LOVELL, ROBERT.

- 115 A compleat history of animals and minerals. 8. Oxf. 1661
'21 March 1660—6^s.'
[A compleat Herball.] *Missing*.
Probably replaced by the 2nd edit., 1665, pres. by Emes.

LULLY, RAYMOND.

- 101 Secreta secretorum in libros tres divisa. 12^{mo}. Colon. 1592

LYTE, HENRY. *See* DODOENS.

MACER, FLORIDUS.

- 110 De herbarum viribus ; cum commentt. Guillermi Gueroaldi.
12. [Franc. 1540]
- 96 De herbarum viribus ; cum schol. G. Pictorii. 12. Basil. 1559

MANELPHUS, JOANNES.

- 96 De helleboro disceptatio. 8. Rom. 1622

MARANTA, BARTHOLOMAEUS.

- 85 Methodi cognoscendorum simplicium libri tres. 4. Ven. 1559

MARCRAVIUS, GEO. DE LIEBSTADT.

- 53 Tractatus topographicus et meteorologicis Brasiliae cum
eclipsi solari. fol. Lugd. 1648
With W^m Piso, *Hist. Nat. Brasiliae*.

MARONEA, NICOLAUS.

- 65 Commentarius in tractatus Dioscoridis et Plinii, de Amomo.
4. Basil. 1608
- 64 Descrittione dell' Amomo indiano ; trad. dal Latino da
Francesco Pona. 4. Venet. 1617

MATTHIOLUS, PETRUS ANDREAS.

- 32 Commentarii aucti in libros sex Pedacii Dioscorides de materia medica, adjectis quam plurimis plantarum et animalium imaginibus.
De ratione distillandi aquas. fol. Venet. 1583
'2 Octob. 1632—20^s.'
Note on Sig. b. 2. "A harde kernel or impostumi in ye bodye."
At end (erased). . . In yeare . . . lord 1624.
7th of June, be my munday
cost of this booke 4^s . . .
two month after . . .
Susan Ironsmith.
- 31 Les Commentaires de M. Pierre André Matthioli, sur les six livres des Simples de P. Dioscorides; trad. de Latin en Francoys. Edit. 2. fol. Lyon 1566
'Rec. 30 Marcij 1654.'
Figs. crudely coloured.
- 137 Compendium de Plantis omnibus. *With* Calceolarius, F. Iter Baldi montis. 8vo. Venet. 1571
'10 Octob. 1632—4^s.'
This volume was sold out of the Library about 1745 by the Librarian T. W.[est], whose note to that effect is written below the College book-plate on the back of the title. As these pages were in process of being made up, the owner of the volume, Mr. Gilbert R. Redgrave, spontaneously wrote to inform me that it was in his possession. He has since generously restored it to the College to be once again placed among Goodyer's books, after an absence of a century and three-quarters.
- 33 Opera quae extant omnia, cd. Caspar Bauhino. fol. Basil. 1598
'31 Januarij 1615—20^s.'
Goodyer's marginal references throughout, and notes on end papers.
Defensio: *see* HESSUS.
- 38 Kreutterbuch verfertigt durch Joach. Camerarium. fol. Franc. 1590
Figures coloured.
'Aprill ye 4th 1655.'
Sold then to Dr. How this Mathiolus in Dutch and colored for w^{ch} I have twentie five shillings per me William Wells at the Princes Armes in Little Brittain. Bookseller.
William Wells.
Aloyis: Mundillu: 8^o: o : 1 : 6
Fragosa in Span: o : o : 6
Camerarius in marantham o : 1 : o
Ye Camer: wanting the title and Fragosa the last page in ye Index, if they displease, you may returne per next.'
- 58 Epitome utilissima de plantis . . . aucta . . . a Joach. Camerario. 4. Franc. 1586
Perhaps Goodyer's first botany book. The marginal headings and English names may have been added by him as a boy.

MIZALDUS, ANT.

- 104 { Historia hortensium quatuor opusculis contexta ; i. Hortorum
 curam ornatum et secreta ostendit ; ii. insitionum artes
 proponit ; iii. auxiliares et medicas hortensium utilitates
 percurrit ; iv. medicamentorum hortensium olerum, radi-
 cum et artificia explicat.
 Opusculum de sena.
 De hominis symmetria, proportione, et commensuratione.
 An caseus edendo sit salubris. 12. Col. Agr. 1576
 Alexikepus seu auxiliaris et hortus medicus.
 Artificiosa methodus comparandorum hortensium fructuum,
 etc. 12. Col. 1579
 In binding are two copies of the 'Order' described under GESNER
 93, which must have been bound at the same time as this volume,
 which also contains a Churchwarden's Receipt for maimed soldiers.

MONARDUS, NICOLAUS.

- 128 De simplicibus medicamentis ex occidentali India delatis,
 quorum in medicina usus est, interpr. Car. Clusio.
 8. Antv. 1574
 126 Joyfull newes out of the newe founde worlde, wherein is
 declared the rare and singular vertues of diuerse and sundrie
 hearbes, trees, oyles etc. Englished by Iohn Frampton.
 4. Lond. E Alde 1596
 'Jo: Gæoodier' on title.
 With a MS. index of fruits, &c.
 41 { De lapide Bezaar et herba Scorzonera. } at end of CLUSIUS,
 { De ferro dialogus. } *Exotica*, q.v.

MONTANUS, PETRUS.

- 110 De Morborum generibus, ex satyra imprecatoria. At end
 of COLUMELLA de cultu hortorum, q. v. 12^{mo}. n. d.

MOSCARDO, LODOVICO.

- 27 Note overo memorie del museo di Ludovico Moscardo.
 fol. Padoa 1656
 '15 Julij 1657—10^s.'

MUFFET, THOMAS.

- 133 Health's improvement ; or rules comprizing and discovering
 the nature, method and manner of preparing all sorts of
 food used in this nation ; enlarged by Christopher Bennett.
 4. Lond. 1655.
 '10 May 1655—3^s 6^d.'
 'John Mouffetts elder brother of Aldham hall in Essex.'

MULBERRY TREES.

- 92 Instructions for the increasing of Mulberrie-trees, and the breeding of Silke-wormes, for the making of Silke in this kingdome ; Whereunto is annexed his Majesties letters to the Lords Liefetenants of the seuerall Shieres of England tending to that purpose. 4. Lond. 1609

MUSA. *See* BRASAVOLUS.

- MYREPSUS, NICOLAUS, PROPOSITUS. 4. Lugd. 1512
- 73 Dispensatorium ad aromatorios.
 '20 Majj 1639—2^a.
 On title 'Richard Hamon de London ge Will^m Bosone sc psn'
 'maste Hamond prey Hyll mr. chr.'
 Gerrad's Herball. Henry Jones, price iiijs.
 Jos. Quercitanus. [Author of *Diaeteticon*, Paris, 1606.]
 Oratio Salamonis fol xl. W^m Bofn' (? = Bosum = Bossom).

NEANDER, JOHANN.

- 67 Tobacologia. 8. Lugd. 1626
 Richard Downes 1658.
 '11 Mar. 1658 Liber Johis Goodyer ex dono D. Rici Downes.'
 'Si fa zara su'l Dado Henry Blount.'
 'Liber Rici: Downes ; ex dono D. Henrici Blount.'
 A Richard Downes was Vicar of East Meon and Steep in 1681.

NICANDER.

- 4 Theriaca. fol. Ven. 1499
 Bound with DIOSCORIDES, 1499, q.v.

ODONUS, CAESAR.

- 85 Theophrasti sparse de plantis sententiae in continuatam seriem ad propria capita revocatae, nominaque secundum literarum ordinem disposita. 4. Bonon. 1561
 'vide Vanderlinden p. 108.' G.

ÖLHAFIUS, NICOLAUS.

- 92 Elenchus plantarum circa nobile Borussorum Dantiscum sua sponte nascentium. 4. Dantisci 1643
- [PAAW, PETRUS. [8 Lugd. 1601]
 Hortus publicus Academiae Lugduno-Batavae. *Missing*.

[PAREY, AMBROSE. Chirurgery.]

This work was bequeathed by Goodyer to John Westbrook, a witness to his will.

PARKINSON, JOHN.

- 22 Paradisi in sole paradisus terrestris ; or a choice garden of all sorts of pleasant flowers. fol. Lond. 1629
 With notes for the Theatrum, 1640.

- 51 Theatrum botanicum, the theater of plantes; or, an universall and compleate herball. fol. Lond. 1640
 'In quires 36^s }
 The bindinge 3^s } 39^s.'
- In many places, Goodyer has indicated the species described by numbers in the margin, and has marked with a 'P' those which the author claims as 'not having been remembered by any other (author) before'. The chapter on *The Elme* (p. 1403) is marked 'JG', probably to note that it is based on a description by Goodyer, and other passages in which the second edition of Gerard is quoted are indicated by the marginal note 'TJ' for Thomas Johnson, the 'corrigider' as Parkinson styles him of Gerard.
- There are a few notes in the hand which we have referred to Dr. J. Dale.
- At end is part of an account, dated 1572, relating to public preachers attached to the parish of S. Martin in Leicester. Among the names mentioned are Thomas Furner, a benefactor; the mayors Richard Davy, Richard Darker, John Eyrick (a kinsman of Herrick, the poet); the preachers Th. Sparke, Fellow, and L. Humphrey, President of Magdalen, and T. Sampson, Dean of Christ Church.
- PASSE, CRISPIAN DE. [Obl. 4. Utrecht 1615
- 81 Hortus Floridus, or A Garden of Flowers. In two parts.
 '10 Novemb. 1627 10/- Johes. Goodyer.' Figures numbered by G.
- PAYNE, JOHN.
- 80 Flowers, Fruicts, Beastes, Birds and Flies exactly drawne. With their true colours lively described. [28 plates.]
 Sold by Compton Halland over against the Exchange.
 Obl. 4. [?c. 1620]
- PEMELL, ROB.
- 131 A treatise of the nature and qualities of such simples as are most frequently used in medicines. 4. Lond. 1652
 '15 July 1652—3^s 6^d.' 'Cambogia. Kap. 35. Yealowe Jaundice.'
 The second part of the treatise on Simples. 4. Lond. 1653
- 92 A treatise of the diseases of children. 4. Lond. 1653
- PHILOPONUS, JOANNES, GRAMMATICUS.
- 3 Commentaria in Aristotelis libros de anima. fol. Ven. 1535
- PICTORIUS, GEORGIUS.
- 96 Scholia to Æmilius Macer de herbarum viribus.
 Carmen de quadam herba exotica cuius nomen mulier est amara. 12. Basil. 1559
- PILLETERIUS, CASPAR.
- 114 Plantarum tum patriarum tum exoticarum in Walachria, Zeelandiae insula, nascentium synonymia. 12. Mid. 1610
- PISO, GUL.
- 53 De Medicina Brasiliensi. '1^{li} 2^s.' fol. Lugd. 1648

PLAT, HUGH.

- 134 Florae Paradise. 12. Lond. 1608
 '13 Nov. 1632—6^d.'

PLATEARIUS.

- 73 De simplicibus medicina. 4. Lugd. 1512

PONA, JO.

- 39 Plantae seu simplicia, ut vocant, in Baldo monte, et in via
 ab Verona ad Baldum reperiuntur. fol. Ant. 1601
 With Clusius, *Rariorum*, 1601.
- 65 — edit. secunda. 4. Basil. 1608
 With notes to *Pinax*, pp. 87, 89.
 Part of deed dated 1608 bearing name 'Sutton' in binding.
- 64 Monte Baldo descritto da Giov. Pona, trad. dal Latino per
 Fr. Pona. 4. Ven. 1617
 '18 April 1629—9^s.' Notes and references to *Pinax*.

PORTA, JO. BAPTISTA.

- 130 Phytognomonica, octo libris contenta. 8. Franc. 1608
 'J. L. Gilbou[rne].'
- 60 Villae. 4. Franc. 1592
 'Pret vj^s.' Notes, but not by G.

PROPOSITI Dispensarium, *see* MYREPSUS.

RAY, JOHN.

- 98 Catalogus Plantarum circa Cantabrigiam nascentium. 1657
 Index Plantarum agri Cantabrigiensis. 12. Cant. 1660
 Appendix ad Catalogum Plantarum circa Cantabrigiam
 nascentium. 12. Cant. 1662
 '10 May 1660—2^s 6^d.'
 'written { Mr. John Nid, who is dead
 by { Mr. John Wray of Trinitie Colledge in Cambridge.
 'Received this instruction of Mr. John Mapletoft, tutor to the
 Earle of Northumberland's son and of Mr. John Snagge an Apothecarie
 of Petworth 23 July 1659.
 Mr. John Nid
 Mr. John Wray
 Mr. Thomas Pockley
 Francis Willuby Esq.
 Mr. Peter Curthop
 All of Trinitie Colledge in Cambridge.
 Francis Willughby Esq. of Middleton neare Coleshill in Warwick-
 shire, Sr Francis Willughby's son.
 Mr. Peter Courthop of Danny in Sussex on this side Lewes.
 Mr. Tho. Thornton, parson of Sutton als Sulton in Sussex in
 Arundel Rape, borne at Bentham in Yorkshire, 2 miles from
 Yngleborowe hill, 15 Apr. 1663 promised Cloudberry.'

Recd this instruction of
 23 July 1659 Mr John Mapletoft, tutor to the Earle of
 Northumberland's son,
 and of Mr John Snagge an Apothecarie
 of Peterborough.

Mr John Kid

Mr John Wray

Mr Thomas Pockley

Francis Willaby Esq,

Mr Peter Courthop

All of Trinitie Colledge in Cambridge

Francis Willaby^{Esq} of Middleton neare
 Coleshill in warwick shire, Sr Francis
 Willughbys son.

Mr Peter Courthop of Danny in Suffe
 on this side Lewes.

Mr Tho. Thornton parson of Sutton abbs Sutton, in
 Suffe in Arundel Rape, borne at Bentham
 in yockshire, 2 miles from yugleborowe hill
 15 Apr. 1663. promised Cloud berry.

10 May 1660 - 2⁶ 87

written by Mr John Kid, who is dead

by Mr John Wray

of Trinitie Colledge in ~~London~~ Cambridge

RENEALMUS, PAULUS.

- 68 { Specimen historiae plantarum cum iconibus. 4. Par. 1611
 { Crambe, viola, lilium a J. A. Thuano versibus illustrata a
 { P. Renealmo edita. 4. Par. 1611
 '11 Augustj 1653—4^s 6^d.'
 'Dr. William Howe bought him of Mr. Allestre in Pauls
 Churchyard.'

RUAEUS, LAURENTIUS.

- 6 Hippiatria sive Marescalia. fol. Paris 1531
 With MS. notes in an early hand on pp. 136, 139 and at end.
 An early prescription on the first of 12 blank pp. at end.
 'ffor the bargett
 R. ffengreke black ornamet. long pep: annyse seed. rew. ros-
 mary. sage. sethe al y^{se} to gether wth vergys. & iij or iiij heads of
 garlek & put it to y^e herbs & when al the herbes ar sodyn then
 put to y^e juse of ye same for every oon ij eggs shells & all. When
 ye have geven y^e best drynk, dryve hym up & down half a q^{tr} of an
 hour & put to eÿy drÿk of a cove an handful of salt & aft^{er} ye best
 have drÿk rub y^e tÿg wth salt.'
 At end, old prescriptions 'For a hors that ys seke'. 'Pro tussi
 equino'. 'A soveren medicyn for achys, brokynes or swellngs callyd
 y^e . . . oyntment . . . It must be made in Maye or between y^e
 Lady days'.

RUBEUS, HIERONYMUS.

- 89 Disputatio de melonibus; acc. responsum medicinale pro
 asthmate cardinalis Bonifacii, auctore Vincentio Alsario
 a Cruce. 4. Venet. 1607

RUELLIUS, JOANNES.

- 34 De natura stirpium libri tres. fol. Basil. 1537
 'H. Spes nō cōfūdit. W.
 Jy silētio et spe.'
 With old notes in hand of H. W. (?) on pp. 342, 630. A similar
 inscription is written in a copy of Aristotle, *de Hist. Animalium*, and
 Theophrastus, *de Plantis*, fol. Basil 1534, in the library of the Oxford
 Botanic Garden. This copy afterwards passed to 'Robt. Leedes',
 'N. Johnson', and Sherard.

RUEUS, FRANCISCUS.

- 129 De Gemmis, 1652. See VALLESIIUS.

RYFF, GUALTERUS H.

- 102 De memoria artificiali quam memorativam artem vocant;
 item de naturali memoria quomodo medicinae beneficio
 excitanda, augenda, et confirmanda, etc. 8. s. l. 1541

SCALIGER, JULIUS C.

- 100 In libros duos, qui inscribuntur de Plantis, Aristotele
 authore. 12. Marpurg. 1598

SCHWENCKFELT, CASPAR.

- 90 Stirpium et fossilium Silesiae catalogus. 4. Lips. 1600

[SHARROCK, R.

History of the propagation & improvement of vegetables,
by the concurrence of art and nature. *Missing.* 1660]

SORANUS.

- 6 In artem medendi isagoge; Oribasii fragmentum de victus
-
- ratione; C. Plinii Secundi de re medica libri quinque.

fol. Basil. in aed. And. Cratandri 1528

‘R^{ossessor}
ous } 10^s
nthoεισαγωγή
eis in + αγω, duco
αγωγή ductio.’[Anthony Rous, armiger, co. Devon, matric. Broadgates Hall,
18 March 1602.]

[SOWERBY, LEONARD.

The Ladies Dispensatory. *Missing.* 8 Lond. 1652]

SPIGELIUS, ADRIAN.

- 90 Isagoges in rem herbariam libri duo. 4. Patav. 1606

SPRECCHIS, POMPEIUS.

- 92 Antabsinthium Clavenae. 4. Ven. 1611

‘18 May 1623 2 6 } 2 7.
Bindinge 1

STENGEL, C.

- 111 Hortorum, florum et arborum historia in duo tomos dis-
-
- tributa. 8 Aug. Vind. 1650

‘14 Decemb. 1654—3.’

STEPHANUS, CAROLUS.

- 103 { De vasculis libellus, adulescentulorum causa ex Bayfio de-
-
- scriptus; addita vulgari Latinarum vocum interpretatione.
-
- Seminarium sive plantarum earum arborum quae post hortos
-
- conseri solent.
-
- De re vestiaria libellus ex Bayfio excerptus; addita vulgaris
-
- linguae interpretatione; secunda editio.
-
- De re hortensi libellus, vulgaris herbarum, florum ac
-
- fruticum qui in hortis contineri solent nomina Latinis
-
- vocibus efferre docens. 8. Par. Rob. Stephanus. 1536
-
- ‘Curtius’ on last page.

STEPHANUS, ROBERTUS.

[12. Lutet. 1545

- 95 De Latinis et Graecis nominibus arborum, etc., ex Aristotele.

On title: ‘Sum Henrici Harvey et amicorum
Quanto maior iminet nec
cit aut fac. tanto magis viget.’

STEPHENS, P., & BROUNE, GUL.

- 114 Catalogus horti botanici Oxoniensis. 12. Oxon. 1658
 The second part of the Catalogue of the Trees & Plants of
 the Physick Garden. 12. Oxford 1658

SWEERTIUS, EMANUEL.

- 80 Florilegium amplissimum et selectissimum, quo non tantum
 varia diversorum florum genera, sed et raræ quam-
 plurimæ Indicarum plantarum, et radicum formæ, ad
 vivum partibus duabus, quatuor etiam linguis, offeruntur
 et delineantur. fol. Franc. 1612

TABERNAEMONTANUS, JAC. THEODORUS.

- 46-47 Neuw kreuterbuch, mit schönen, künstlichen und leblichen
 figuren und conterfeyten, allerhand gewächs, blumen,
 kräuter, etc. 2 vols. fol. Frankof. 1625
 'Rec^d this 6 of September 1655 of the right worth Dr. How
 the summ of fifty foure shillinge in full for this booke } 2 14 0
 I say Octavian Pulleyn.' With old deeds used as guards.
- 77 Eicones plantarum seu stirpium arborum nempe fructicum,
 herbarum fructuum, lignorum . . . curante Nic. Bassaeo.
 obl. 4. Francof. 1590
 Annotated by Goodyer with modern names throughout, with cross
 references to Lobel.

THALIUS, JOANNES.

- 89 Sylva Hercynia ; sive catalogus plantarum sponte nascentium
 in montibus et locis vicinis Hercyniæ, quæ respicit
 Saxoniam. 8. Francof. 1588
 With refs. to *Pinax*.

THEOPHRASTUS, ERESIUS.

- 2 { De historia plantarum, libri decem.
 De causis plantarum, libri sex
 Metaphysica. } fol. Ven. Ald. 1497
 Contemporary stamped leather Italian binding.
 Chapters numbered by G., who probably used this copy for his
 translation.
- 3 Opera omnia Gr. cum præfatione Joach. Camerarii.
 fol. Basil. Operini 1541
 'Sept^o September 1623 pr. 15^a 6^d.'
 End papers with account for timber and carpentry, signed Thomas
 Herytage.
- 138 De historia plantarum. (With Aristotle.) 8. Lugd. 1552
- 30 De historia plantarum, Gr. Lat. ex interpr. Theod. Gazac ;
 totum opus absolutissimis cum notis, tum commentariis,
 item rariorum plantarum iconibus illustravit Jo. Budaeus

à Stapel ; acc. Julii S. Scaligeri in eosdem libros animad-
versiones, et R. Constantini annotationes. fol. Amst. 1644
' 15^{to} Julij 1657—30^s.'

- 104 De suffruticibus, herbisque ac frugibus libri quatuor, Theod.
Gaza interprete. 8. Argent. Sybold, s. a.
[With *Mizaldus*.]
Pagination added.

THEVET, ANDRÉ.

- 61 Les singularitez de la France antarctique autrement nommée
Amerique ; et de plusieurs terres et isles decouvertes de
nostre temps. 4. Par. 1558
' Novemb. 14 1631—12^d.'
' . . . anguinea vitæ.' On title three names of former owners erased.

THURNEISSER, LEONHARDUS.

- 37 Historia, sive descriptio plantarum omnium, tam domestica-
rum quam exoticarum, earundem virtutes et icones pro-
ponens ; atque una his partium omnium corporis humani,
ut externarum ita internarum, picturas, etc., complectens.
fol. Berlini. 1578

' Vixi satis si Christe sat vixi tibi
JB ex dono CB consang.'

Evidently a gift from Caspar to his brother John Bauhin.

In binding is a folio Proclamation dated 24 May 1648 printed by
Cotes, London. And used as a guard is part of a letter ending
' millitia' and endorsed 'Castle baynard', the name of one of the
Wards of the City of London, to the south of St. Paul's Cathedral.

TRADESCANT, JOHN.

- 134 Plantarum in horto Johannem Tradescanti nascentium cata-
logus. 8. s. l. 1634
A unique copy. See p. 334.

TRAGUS, HIERONYMUS.

- 127 De stirpium, maxime earum, quae in Germania nostra
nascuntur, usitatis nomenclaturis propriisque differentiis,
etc., commentariorum libri tres, ex ling. Germ. in Lat.
conversi D. Kybero interprete. 4. Argent. 1552
With G.'s references to plants, cf. p. 73.

- 48 { Kräutterbuch . . . verbessert durch Melchiorem Sebizium.
{ Teutsche Speisskammer. fol. Strasburg 1630

TREVERIS, PETER.

- 28 { The grete herball.
4. Lond. in Southwarke by Peter Treveris 1526
{ The great herball newly corrected.
4. Lond. in aed. Tho. Gybson 1532

TROXITEN.

- 20 Horn des heyls menschlicher blödigkeit ; oder Kreütterbuch, darinn die Kräuter des Teudschenlands, ausz dem liecht der natur, . . . beshriben durch Philomusum anonymum ; nachmals durch doctorem Troxiten in truck geben.
 '25 May 1655—8^s.' fol. Strasb. 1595

TURNER, WILLIAM.

- 13 The first and second partes of the herbal of William Turner, lately oversene corrected and enlarged with the thirde parte lately gathered and nowe set oute with the names of the herbes, in Greke, Latin, English, Duche, Frenche, and in the apothecaries and herbaries Latin.
 fol. Collen, by Arnold Birckman, 1568
 With a MS. index and notes by Goodyer.
 A booke of the natures and properties, as well of the bathes in England as of other bathes in Germanye, and Italye. fol. Collen, by Arnold Birckman, 1568
 'To Mr. O. Bilson from E. Gray heboriste.'

VALLA, GEORGIUS.

- 134 De simplicium natura liber unus. 8. Argentinae 1528

VALLESIIUS, FRANCISCUS.

- 129 De sacra philosophia, sive de iis, quae in libris sacris physice scripta sunt ; liber singularis. 6th edit. 8. Lugd. 1652
 With Lemnius, de Plantis sacris, and Rueus, de Gemmis.
 End papers are part of a printed Proclamation of the House of Commons concerning spirits who steal children, dated 1661, printed by Rich. Hodgkinson living in Thames Street over against Baynards Castle 1661.

VESLINGIUS, JOANNES.

- 85 Paraeneses ad rem herbariam. p. 85.
 De plantis Aegyptiis observationes et notae ad Prosp. Alpinum ; cum additamento aliarum ejusdem regionis.
 4. Patav. 1638
 Opobalsami veteribus cogniti vindiciae. p. 217.
 With Prosp. Alpinus, de plantis Aegyptis, q. v.

VIGNA, DOMINICUS.

- 66 Animadversiones sive observationes in libros de historia, et de causis plantarum Theophrasti ; addita fuit tabula studio et opera Andreae Checcaccii. 4. Pisis. 1625
 '19 April 1655—18^d.'
 19 Apr. 1655 unbound—1^s 6^d }
 28 Junij 1655 the binding—1^s 2^d } 2^s 8^d.'

WOLPHIUS, CASPAR.

101 De stirpium collectione tabulae. 12. Tiguri 1587

WORMIUS, OLAUS.

54 Museum Wormianum. fol. Lugd. 1655

Two of the titles in the old list have hitherto proved untraceable. They are 'Anatomia Sambuci' and 'Alb. Montani Isagoge Physico-magico-medica'.

THE GOODYER MANUSCRIPTS.

MS. 1-6. GOODYER, J., and HEATH, J. Translation of DIOSCORIDES into English with interlinear Greek text. 4,540 pages 4to, bound in six vols., dated 1653-5.

See p. 85.

MS. 6*. GOODYER, J., and HEATH, J. Translation of SARACEN, Scholia on Dioscorides, into English. 300 pp. folio.

MS. ends abruptly on p. 292 with a note, 'Joh. Heath clericus obiit 25 Nov. 1656'. See p. 89.

MS. 7. GOODYER, J. Translation of THEOPHRASTUS, *περι φυτων*, into English. 494 pages, interleaved, one vol., sm. folio, dated 1623.

De Plantis, pp. 1-238. De Causis Plantarum, pp. 239-494.

In the binding is part of a deed: 'Witherdon de Stone' in Kent and 'Joh. Sharpe' are mentioned. See p. 50.

MS. 8. [? DALE, J.] i. Descriptions of English Grasses. 80 pp. folio.

Not in Goodyer's hand, but enlarged with additions by him.

Reasons for referring these MSS. to John Dale are given on p. 296.

ii. Descriptions of Plants headed 'Ex manuscriptis Turn. Annexis Lobelii Observationibus'. 30 pp. folio, dated Jan. 22, 1651.

In the same handwriting as No. i. The date of Lobel's work was about 1576.

iii. GOODYER, J. Index to Plants described in his copy of C. Clusius, *Rariorum Plantarum Historia* and other works bound up therewith. 20 pp. folio.

MS. 9. [? DALE, J.] i. Index of British Plants, c. 1650-1656.

The work, written on about 250 leaves folio, is based on C. Bauhin's *Pinax Theatri Botanici*, 1623, which Goodyer acquired before 30 Nov. 1623. The handwriting is that of the author of MS. 8. i and ii.

References are given to the following works: Bauhin, *Pinax*, 1623; [Lobel], *Plantarum seu stirpium icones*, 1581; *Icones Stirpium*, 1591; Gerard, *Herball*, 1597; Johnson's *Gerard*, 1633; Parkinson, *Theatrum*, 1640; How, *Phytologia*, 1650; Johnson, *Mercurius Botanicus*, 1634.

The entries from the last book have been marked with yellow paint by Goodyer, who made a few additions to the two last pages and to Pin. 81, 131, 155, 265, 288, the two last being dated 19 July 1656. The following extracts illustrate the author's style and notes:

Rapunculus folio oblongo spica orbiculari. (Pin. 92.)

= *Merc.* 64 *Rapuntium corniculatum montanum*, Col.

'Dns Goodyerius semper suspicabatur haec duo novissime proposita non ipse nisi ejusdem plantae, ratione soli vel Arigosioris vel laeti, variantis diversa nomina.'

Melampyrum latifolium. (Pin. 234.)

'Mali coniunxit emaculator Sideritum pratensem luteum Lugd. et Crataegonon, Lob. quae sunt diversae Plantae. Sideritis pratensis lutea Lugd. non habetur apud Lobelium. Inquirendum an Sideritis lutea Mri. Stonehouse ad hanc Sideritem possit referri.'

Blattaria alba. (Pin. 241.)

'Huc etiam referimus Blattariam fl. viridi et Blattariam fl. albo flavescente, indignus quae novam speciem facerat.'

- ii. List of 153 British Plants. 'Plantae ad Methodum Pinacis reducendae quibus Botanographi nostratis ortum tribuere Britannicum.' Dated the last day of April 1659. By John Dale?

The list is in the same hand as the *Index*, and like it has been annotated by Goodyer, who was evidently in correspondence with the writer, e.g. after No. '8. Gramen Parnassi' Goodyer adds, 'This you have putt into the draught of your Catalogue'.

MS. 10. Hortus siccus.

Among the MSS. received by Magdalen College from Goodyer were two Herbaria, listed as 'Hortus hyemalis fol.' and 'Hortus hyemalis minor fol.'. The former is missing, but the latter may be represented by this small herbarium of 10 leaves folio.

It contained a small collection of Mosses and Ferns, made c. 1620 and mounted on the leaves of a MS. Botanical Glossary, English-Latin, beginning with 'Anise seade Anisum' and ending 'Yerrowe Nose bleede. Stratiotes millefolia. Militaris Millefolium'.

All the dried plants have been removed, but the following names on paper straps are legible:

- f. 3. Lichen arborum. Lichen cinereus. Muscus peltatus.
Muscus pyxidatus. [] sylvestris, 3 Jul. 1620.
4. Phillitis digitata. Ceterach.
5. Filix palustris.
6. Polypodium.
7. Filix spinosa. Chamaefilix marina anglica.

MS. 11. GOODYER'S Miscellaneous Papers.

The more important of these are published or referred to in this volume as MS. 11, f. —, or more briefly as MS. f. —, with the leaf number but without the number of the manuscript.

MS. 12. LOBEL. *Stirpium Illustrationes; plurimas elaborantes inauditas plantas subreptitiis Joh: Parkinsoni rapsodiis (ex codice MS. insalutato) sparsim gravatae*: edited by W^m How. 37 leaves folio.

The original MS. from which the work was printed in 1655. See p. 252.

MS. 13-15. LOBEL. *Stirpium Illustrationes*.

An unpublished work in preparation about 1612. About 835 plants are described. See p. 253.

MS. 16. GOODYER, J. *Index to Gerard's Herbal (1597)*. Sm. 8vo. With localities of British Plants.MS. 17. [STONEHOUSE, W.] *Catalogus Plantarum Horti mei Darfeldiae. Anno 1640*. With plan 12mo.

See p. 348.

MS. 18. HOW, W. *Additional Notes to his Phytologia Britannica 1650*. Written 1650-1656. See p. 276.

An interleaved copy with many corrections, notes, and MS. lists of plants by the author who died 30 Aug. 1656. Goodyer, according to a note inside the cover, received the book on 30 Apr. 1659, and then added marginal references throughout and notes on six plants printed on p. 194.

On the inside of the cover is a receipt dated July 29 and unsigned, probably written after 1659, when the book was in Goodyer's possession. 'Rec. of Mr. Goodier ten pounds for Mr. Bold's use.' The Mr. Bold was probably Arch. Bold, one of the witnesses to Goodyer's will.

The notes in How's handwriting are partly his own, and partly from information received from Goodyer, Hunnibon, and William Browne of Magdalen College. Their source is acknowledged both where they occur and generally on the front page. 'Gaine I was for Goodyers Plants and des. y^e like for Brownes, Lobells [and Pennyes MS. w^{ch} review for names etc.]' (struck out). Inserted are 7 coloured and 3 uncoloured drawings of plants.

Passages from this volume have been frequently quoted by Druce, but we have not found any evidence for his statement that 'on the death of Goodyer the book probably came into the possession of W^m Browne'.¹ It would have passed into the possession of the College direct, with the rest of Goodyer's library in 1664.

¹ Druce, *Flora Berks.*, p. ciii.

MS. 19. JOHNSON, T., and HOW, W. Additions to T. Johnson's
*Descriptio Itineris Plantarum investigationis in agrum
Cantianum* A. D. 1632.

Formerly in the possession of W. How, and previously Johnson's own
copy. See p. 276.

Additions in T. Johnson's handwriting :

On p. 3, l. 10 'Alga membranacea' has been struck out.

On p. 6 to the list of plants obtained in the Isle of Thanet, *Absinthium
vulgare* has been added. On p. 9 'Saxifraga, Dod.' has been
added after 'Serpillum' in l. 3. On p. 37, to the Flora of Hampstead
Heath are added

'Coronopus Ruellii, Cornu Cervi alt. vulg.

'Scabiosa minor sive ovina, Dod. Cam., media, Lob.

'Scabiosa media serrato angustifolio, flore Globularia. Adv., Ra-
puntium alterum leptophyllum capitatum, Col.'

At back of plate is a note of 'Trifolium pumillum . . . White dwarfe
Trefoile'.

All these are printed in the *Phytologia*.

Eighteen leaves are bound in at the end.

ff. 1 and 2. Blank.

ff. 3-6. Index to genera mentioned, including the MS. additions.

In Johnson's hand with additions by How.

f. 6 v. Notes by How.

ff. 7-11. Alphabetical list of about 190 species of English plants
not included in Johnson's lists. In How's hand.

ff. 12-17. Blank.

ff. 17 v., 18. Notes by How.

This little volume has the great sentimental interest of being the germ
from which all British Floras are descended. The first index comprises
all the plants mentioned in Johnson's book, and is written, I believe, by
the author himself. The second list contains names of English plants,
including those quoted by How as derived from 'Dr. Johnsons. *Mf.*', in
How's hand, with additions by him at a later date. Unfortunately these
plants are not localized. How's rough notes on the last two pages are
evidently memoranda used when preparing the *Phytologia*, our first
British Flora.

If my interpretation of the handwritings be correct, this volume would
probably have passed from Johnson (d. 1644) to How, who would have
used it in the compilation of the *Phytologia* (1650), and after How's
death in 1656 it would have passed to Goodyer, perhaps with How's copy
of the *Phytologia*, in 1659.

NOTES ON CONTEMPORARY BOTANISTS

MOSTLY FROM GOODYER'S BOOKS AND PAPERS

IN the England of John Goodyer, the stream of Botanical learning was flowing along a very small channel. So small was it, that viewed from the present time after the lapse of three hundred years, the water appears confined to a few isolated pools and backwaters with no certain channel between them. The number of true men of science, as opposed to herbalists, could be counted on the fingers of one hand, and the untimely death or defection of any one of them might have put back the progress of botany for a century.

The following notes on Goodyer's friends or contemporaries were gleaned from, or suggested by, the study of his own manuscripts. By their publication we may perchance bring to light an occasional fact which reveals the course of the stream of botanical learning.

The names of previous owners of his books have been listed: his manuscripts suggest material for more extended notes on some of the following botanists, the others emerged during our research: they are not to be found in the *Biographical Index of British Botanists*. The identification of the handwritings of some of them was by no means an easy matter: two are still doubtful.

- i. THOMAS PENNY, *c.* 1530-1589.
- ii. THE 1570 BOTANIST OF OXFORD AND WINCHESTER.
- iii. RICHARD GARTH, *d.* 1597.
- iv, v. WILLIAM AND SIR JOHN SALUSBURY, 1567-1613.
- vi. M. LOBEL, 1538-1616 and HOW.
- vii. WM. MOUNT, 1545-1602.
- viii. RICHARD SHANNE, 1561-1627.
- ix. JOHN PARKINSON, 1567-1650.
- x. WALTER STONEHOUSE, 1597-1655.
- xi. THOMAS JOHNSON, *c.* 1600-1644.
- xii. WILLIAM HOW, 1619-1656.
- xiii. JOHN DALE, *d.* 1662.
- xiv. WILLIAM BROWNE, 1629-1678.

i. THOMAS PENNY, c. 1530-89.

Thomas Penny, M.D., of Trinity College, Cambridge, B.A. 1551, F.R.C.P. 1582, contributed to a natural history of Insects, which, begun by Edward Wotton and amplified with extracts from Conrad Gesner, was finally completed by Thomas Muffett of Oxford. Muffett died in 1604, leaving the book in manuscript. It was eventually published in 1634 by Sir Theodore de Mayerne.

Penny's botanical reputation has been rescued 'from an almost total obscurity' by Pulteney, who points him out as 'A second Dioscorides, for his singular knowledge in plants'. He had resided in Switzerland and had visited the island of Majorca. He was personally acquainted with Gesner, Camerarius, and Clusius. From Majorca he brought *Geranium tuberosum*, *Swertia perennis*, and *Hypericum balearicum*, which Clusius named 'Myrtocistus Pennaei' in his honour.¹ Clusius² in 1583 thanked him for a drawing of *Cnicus heterophyllus* Roth., sent in 1581, and noted³ his discovery of *Cornus suecica* L. in the Cheviots. We have Mount's statement that he grew *Acorus Calamus* in his London garden before 1582, Penny communicated the following plants to Camerarius, who describes him as a leading London Physician 'rerum naturalium peritissimus, amicus meus singularis'. *Hort. med.* 1588, p. 36.

Caryophyllata vulgaris or C. altera alpina with white flowers. in France.	Monte Lupo <i>Geum reptans</i> L.
Lactuca sylvestris 'odore prorsus Opii'.	<i>Lactuca virosa</i> L.
Matricaria tertia fl. pl. in Anglia frequens.	<i>Matricaria parthenium</i> L.
Rhodia radix. Ingleborrow.	<i>Sedum Rhodiola</i> DC.

And Lobel, *Adv.* 367, associates 'Myrrhis altera' (*Myrrhis odorata* Scop.) with him.

The mention of the name of Penny by How at a date subsequent to 1650 in an erased passage quoted on p. 280 is of importance, because it may throw new light on his botanical MSS. Pulteney's account of the matter is that 'Dr. Penny died in 1589, and is said by Jungerman to have left his papers to Mouffet and Turner; but, in this account, there is surely a very striking anachronism since Turner himself died in the year 1568'. It is clear, therefore, that Penny's zoological MS. on Insects went to Muffett, and that his botanical MSS. could not have gone to WILLIAM, but to some other Turner. As a possibility, a man who would have valued them would have been the well-known astrological botanist, ROBERT

¹ *Ger. emac.* 434, 946, 1279.

² *Stirp. Pannon. Hist.* 1583.

³ *Rar. Plant. Hist.* i. 59, 1601.

TURNER,¹ whom we have suggested as identical with the 'Turn.' owner of the Lobel MS. mentioned below. Robert lived at Holdshot in the north-east corner of Hampshire: 'Turn.' was in correspondence with Goodyer and Dale (?), either of whom might have shown his MS. to How, without however giving permission for publication by the latter.

ii. THE 1570 BOTANIST OF OXFORD AND WINCHESTER.

[? DR. WALTER BAYLEY, 1529-92.]

On looking through some of the older books in the Botanical Department of the British Museum I was rewarded by finding twenty-nine plant records, some dated 1570-2, in the hand of an unknown botanist, who appears to have lived at Oxford and Winchester. In accordance with a practice very usual in those days, he wrote English names of plants in the margins of his Latin botany book, *Du Pinet*, *Historia Plantarum*, Lugd. 1561, and in a few instances added the names of persons and localities. The names are Watson, Jeames, Barnabye, Norton, Strowde, Heiden, Basket, and Crosse. The localities are mostly the several gardens of these persons; and a few places, all near Winchester, are noted as stations of common Hampshire plants.

When the preceding clause was already in type, I happened to see an autograph inscription in a precious little volume by Dr. Walter Bayley of New College,² printed privately and issued anonymously as a New Year's gift to a friend. The writing at once caught my eye on account of its resemblance to the writing of our unknown botanist. Both writings are in the style of the period; and without further specimens of each, it is impossible to be certain of identity, but Dr. Bayley was certainly the kind of man who might have entered botanical memoranda in a *Du Pinet*.

WALTER BAYLEY was educated at Winchester and New College, becoming a Fellow there in 1550. When Junior Proctor, he demanded the degree of Bachelor of Physic, and supplicated for leave to practise medicine 'per totam Angliam'. He was Queen's Professor of Medicine at Oxford from 1561 to 1582. In 1581 he was appointed Physician in Ordinary to Queen Elizabeth.

The *Du Pinet*, 1561, would therefore have been the newest botany book out at the time of his becoming Professor, and the marginal notes would have been written about the middle of his

¹ A Robert Turner was born at Reading at h. 9.48 a.m. on 30 July 1626 (*MS. Ashm.* 183).

² The property of Sir D'Arcy Power: see his *Dr. Walter Bayley, Med. Chir. Trans.* xc.

tenure of the chair. Besides those printed below, one other entry may be mentioned. Under 'Piper indicum' p. 12 is noted 'a kynd of pepper groweing in India: husked'. It may only have been a coincidence, but Bayley happened to have been greatly interested in Peppers, for some years later he printed, for distribution among his friends, *A short Discourse of the Three Kindes of Peppers in common use and certain Medicines made of the same, tending to the preservation of health*, 1588. The discourse contains one personal note which bears on the author's movements: 'I have often seene at Poole at Dorsetshire and also in London, the whole clusters of pepper preserved in brine and in salt'. It is unnecessary to point out that an old Wykehamist journeying from Oxford to Poole would naturally break the journey at Winchester.

If we accept the dates 1570-2 and the identifications of the species, these brief notes are among the earliest known evidences for the occurrence of eight species of plants in Hampshire, two of which are the first localized notices for Britain.

- Halimus or Atriplex marinus, p. 62. *Atriplex littoralis* L.
 'Upon hāble bankes in great quantitie.'
 [The river Hamble is not far from the locality where Lobel noted the plant.
 His, the first printed record, dates from 1655.]
- Conyza media, *Du Pinet*, p. 390. *Pulicaria dysenterica* Gray.
 'In diches evrywhere about Winchester.'
- Sium, p. 171. *Nasturtium officinale* R. Br.
 'In ye dych towards Nortons.'
- Clematis altera Dioscorides, p. 442. *Clematis vitalba* L.
 'White vyne. In every hedge about Winchester.'
- Thlaspi. *Thlaspi arvense* L.
 'In Mr. Strowdes garden and medoe, 1572.'
- Eupatorium Avicennae, p. 476. *Eupatorium cannabinum* L.
 'In ye dyche toward blak Bridge.'¹
- Tithymalus masculus, p. 605. *Euphorbia amygdaloides* L.
 'In cops by Cathe of ij sorts.'¹
- Personata altera, p. 559. *Petasites vulgaris* Desf.
 'Close by the river sydes.'
- Cynocrambe, p. 635. *Mercurialis perennis* L.
 'Oxon.'
 [An evidence more than 200 years earlier than the first record in the County
Flora.]

For the notices of garden plants see p. 304.

¹ My friend Mr. H. Salter tells me that Black Bridge over the Itchen is near the west end of the Warden's Garden, and suggests that 'Cathe' may be an abbreviation for St. Catherine's Hill.

iii. RICHARD GARTH, *d.* 1597.

Richard Garth was an accomplished botanist whose contributions to science are better remembered in the works of Clusius and other foreign botanists than in his own country. He was the son of Edward Garth, one of the six Clerks in Chancery, and owned a property at Morden in Surrey in 1564. Between 1581 and 1591 his relations with Brazil enabled him to bring several of the plants of that country, including the 'Papyrifera arbor', the 'Juni-pap-peeywa Brasiliorum', the 'Phaseolus Brasiliorum', and some exotic fruits to the notice of Clusius, who described them in his *Libri Exoticorum* in 1605. In return Clusius gave him a Solomon's Seal, a root of which he 'very lovingly imparted' to Gerard, who not unnaturally described him as 'a worshippingfull Gentleman, and one that greatly delighteth in strange plants' (*Ger.* 757). So far as the English flora is concerned his name should be associated with the Great Toothwort (*Lathraea squamaria* L.) which grew on his land at Groutes, not far from Croydon.¹

In 1592 Garth purchased the manor of Drayton from Robert, Earl of Sussex. He died in 1597, having married, firstly, Elizabeth Dixon² and secondly Jane da. of . . . Busher, co. Linc., who survived him, living at Drayton Manor facing Haylinge Island, two miles from Portsmouth. After his death, Lobel appears to have visited her garden there, and to have found '*Alopecurus altera maxima Anglica paludosa sive Gramen Alopecuroides maxima*'³ and a variety of Bindweed, '*Helxine cissampelos alt.*'⁴ growing wild near the house. She seems to have refreshed him with Metheglin of her own brewing, and to have given him her recipe for it, which he printed (*Advers. alt.*, 1605, p. 473).

Lobel speaks of Garth as Senior Clerk in Chancery ('*Diplomatica Curia*'), an office to which he had probably succeeded by inheritance, and as most learned in the natural history of Indian as well as of our native English plants. Hugh Morgan, James Garret, the communicator of the vegetable discoveries of his brother Peter and of those who accompanied Sir 'Walterus Raulaeus' to Guiana,

¹ *Ger. emac.* 1585. In the author's copy of this work there is on p. 762 an old MS. note relating to the Greatest Wolfe-bane, now *Doronicum Pardalianches* L. 'This growes wild in the Orchard of a house called Grouts in the parish of Morden in Surry lately belonging to Mr. Garth, Lord of that Mannour. It floures in Aprill.'

² The *Heralds Visitation* of London mentions a Richard Garth of Moorden, co. Surrey, who m. Dorothy Style.

³ *Polypogon monspeliensis* Desf.

⁴ *Lobel and How*, p. 127; *Park. Theatrum*, p. 173.

and Richard Garth were the principal authorities in this country on tropical plants in the sixteenth century.

His copy of *Caesalpinus* is in the Magdalen Library. The title-page bears his signature, and in the body of the book he underlined passages relating to the properties of Tobacco, p. 344, and of Scorzonera, p. 427, and added the name 'Battato' for 'Castaneae terrestres', on p. 427. The volume passed in 1598 from his son Robert Garth (*d.* 1613) to Dr. Lancelot Browne, the author of a prefatory eulogy in Gerard's *Herbal*, 1597, and then to John Goodyer.

iv. WILLIAM SALUSBURY, 1520?-1600?

v. SIR JOHN SALUSBURY OF LLEWENI, 1567-1612.

Some few years ago I had the satisfaction of finding in the Library of Christ Church a copy of Gerard's *Herbal*, with a few dated marginal notes of plants found in 1606-1608 in North Wales, and with notes on the medicinal properties of others. The *Herbal* is inscribed 'Sir John Salusbury his booke', and the notes are evidently in his own handwriting. They illustrate the manner in which Gerard's work encouraged the practice of recording exact plant-localities at the time when Goodyer was a boy, even in quite remote parts of the country. They are not mentioned by any botanical writer with whom I am acquainted, nor do the recent historical notes on the flora of Denbighshire by Dallman take us farther back than Waring's letter of 1772. Salusbury's date was 1606, and though there is no chance of his ever being forgotten as an historical character, he also deserves to be remembered by compilers of county floras: moreover, unlike his cousin, William, he wrote in English. He received his first education at Oxford at Jesus College.

By all accounts Sir John Salusbury of Lleweni, known as 'the Strong', was no ordinary man. He came of a remarkable family, one member of which, WILLIAM SALUSBURY (1520?-1600?), the first translator of the New Testament into Welsh, is stated to have been the author of a Welsh *Botanologia* which is said to have been an original work showing close observation of plant life in Wales. It is possible, however, that William's great literary reputation has led his biographer to overstate his botanical achievement (*D.N.B.*). I believe the 'Welsh *Botanologia*' of the *D.N.B.* to be the *Llysiuilyfr Meddyginiacthol*, recently (1916) edited by Mr. E. Roberts. It is a *Herbal* in Welsh, a compilation of extracts from Fuchs, Turner, and Dodoens translated into Welsh before 1597 by

William Salusbury, with the Welsh names of the plants, and a few (very few) localities added. Mr. Roberts points out that the more detailed localities are those nearest Llansannan and Llanrwst where William Salusbury is known to have lived. Lleweni is thrice mentioned, and both author and work must have been well known to Sir John Salusbury: the names of plants localized by William are printed below.

John's mother was the celebrated Catherine Tudor of Beraine, popularly known as 'Mam Cymru' or Mother of Wales, celebrated alike for her numerous descendants and her four marriages.¹

Sir John has been stated to have had two thumbs on each hand. His gardening is not remembered as well as is his huge strength. The Denbighshire tradition that he used to 'tear up forest trees by the roots' is reminiscent less of his interest in botany than of a fondness of displaying his physical powers.² He married Ursula, daughter of Henry 14th Earl of Derby, and left three sons and two daughters, the eldest of whom, Sir Henry Salusbury, Bart. (*d.* 1632), also wrote his name in the *Herbal* (in 1627), and entered notes on two plants growing on the Chirk estate of his father-in-law, Sir Thomas Myddleton (1550-1631).³

On a recent pilgrimage to Denbigh to visit the site of Sir John's gardens at Lleweni, chance made me acquainted with Mr. A. Foulkes-Roberts of Denbigh, himself a lineal descendant of Catherine of Berain. He at once took the greatest interest in the quest and drew attention to the fact that Sir John Salusbury

¹ CATHERINE OF BERAINE was a great granddaughter of Henry VII and therefore cousin to Queen Elizabeth. She *m.* 1. Sir John Salusbury. 2. Sir Richard Clough (and from this marriage was descended Mrs. Thrale, the friend of Dr. Johnson). 3. Morris Wyn (as his third wife). 4. Edward Thelwall. The story goes that after the funeral of her first husband she left the church in the company of Mr. Wynn who then and there offered her marriage. She declined on the ground that, on her way to church, she had promised Sir Richard Clough!

² 'A popular tradition credits him with having killed a mythical and much dreaded beast that had its lair in the cliffs below the castle, and having also slain a great white lioness with his naked fist in the Tower of London, thus earning for his estate the name of Lleweni, Llew being the Welsh for the king of beasts. He also overthrew in a wrestling match a famous giant, Edward Shon David, whose walking-stick was the axle-tree of a cart with a crow-bar driven through it. Syr John, too, was accustomed to show off his strength, when he had no worthier object for it, by tearing up forest trees by the roots.'—Bradley, *North Wales*.

³ SIR T. MYDDLETON was elected Lord Mayor of London on the same day that his brother Hugh opened the New River Head. He had purchased the estate of Chirk Castle in 1595.

was already known by his poems, and that the MSS. of some of the poems are in the Library of Christ Church in Oxford.¹

On returning to Oxford, through the kindness of Canon Cooke, I was given the earliest possible opportunity of examining the two volumes known as Christ Church MSS. 183 and 184 and Professor Brown's notes thereon, and then learnt, what has been known to Shakespearean scholars for some years, that Shakespeare, Marston, Chapman, Jonson, and 'Ignoto' contributed 'Diverse Poeticall Essaies' on the subject of the *Phoenix and Turtle*, as a supplement to a poem by Robert Chester entitled *Loves Martyr*, and that the whole collection was dedicated to Sir John Salusbury of Lleweni.

The personal association of Salusbury with the great Elizabethans is also shown by the presence of a poem written in Jonson's own hand among the Salusbury MSS. at Christ Church, and by the lines in which Robert Chester welcomed his patron home from London, where

The swanns that laue their blacke feet in the streames,
Have in their sweetnes sang you golden theames :
Court-bewtefying Poets in their verse,
Homerian like sweete stanzoes did rehearse.

Robert Chester may well have been a member of the Salusbury household, who knew Lleweni and his patron's love of flowers. Indeed, in his *Wynter garland of Sommer fflowes made in manner of A Neweyeaeres gyfte to the Right Worshipfull John Salusbury Esq^r of the body to the Queenes most exclent Maiestye, 1598*,² he may have drawn inspiration from plants actually growing at Lleweni. He mentions twenty-seven in thirteen stanzas of his poem.

Jelliflower.	Venus Looking glasse.
Daphadill.	Hyacinthus.
Organy.	Strawbery.
Orice.	Angellica.
Honysuckle.	Basill.
Rose.	Lauender.
Ladies Nauel.	Virgins bower.
Oleander.	Sweet Marierome.
Stickadoue.	Rosemarye.
Touchmenot.	Arkeangell.
Agnus Castus.	Ladies smock.
Hartsease.	Youthwort.
Ladies Seale.	Lady lacies.
Yooke Elme.	

A selection of Sir John Salusbury's own poems was printed in 1597 in a small volume dedicated to him by Robert Parry, gent.,

¹ Carleton Brown, *Poems by Sir John Salusbury and Robert Chester*. *Bryn Mawr College Monographs* xiv, Pennsylvania, 1913.

² MS. 184, f. 45 a.

of which only one copy is known. His *Certaine Necessary observations For Health*, a poem illustrating his interest in practical hygiene, was printed as a broadside, of which a copy is bound up in his 'booke of notes', MS. 183 (f. 4), and is dated (in MS.) 1596, though a copy of the poem itself, written by a clerkly hand, is dated 1603 (MS. 184, f. 77 b). The volumes of MSS. contain many verses in Welsh, praising various members of the Salusbury family, copies of letters from Sir Henry Sydney, the Earl of Essex, Sir Walter Raleigh, and other notabilities, English poems by Sir John Salusbury and his circle, a few medical recipes, and stray quotations and verses.

Sir John's own muse drew less from a knowledge of plants than did the muse of Robert Chester, though there are, as in all Elizabethan poetry, references to 'choyse and sweetest flowers', sweet Briere and sweet Eglantine. And one of his verses, on Pride, entitled *A Conceite*, ends with the lines

And those that grow of sundry seeds
At last do proue but stinking weeds
And if pure wheat be sowde in tares
The wheat Assuredly it mars.

finis John Salusbury.

but they can hardly be cited as showing exceptional cultural lore.

The names of many herbes are contained in his medical recipes, of which the following is a specimen :

Tacke a certain amand milke mayde Whit these ierbes Tacke plantain, ribbe Whorthe, knott grasse, cheaper purse, confery of evere one a handfull, strabury leaves, sanicula, of evere one halfe a handfull. Let this by boylet in a quantitie of faier Water of this Liquor macke an amand milke.

This is excelent against a consumcion, waste, or runninge of the raynes, or brekinge of a vayne & within the boodie, or anye foule matter wthin manes boodye.

[Christ Church MS. 184, f. 33¹

¹ By another hand in the same volume is *A Dietary for those who have weak backs*, in ten 4-line stanzas :

1. Good sir yf you lack the strengthe in your back
and wolde have a Remediado
Take Eryngo rootes and Marybone tartes
Redde wine and riche Potato.
2. An oyster pie and a Lobsters thighe
hard eggs well drest in Marow
This will ease your backes disease
and make you a good Cocksparrowe.
4. An Apricock or an Artichock
Anchovies oyle and Pepper
These to use doe not refuse
twill make your backe the better.
10. The milke of an Asse will bringe to passe
all thinges in such a matter.
When this is spente you must be contente
with an ounce of Synamon water.

[Christ Church MS. 184, f. 35

Sir John was evidently of a most cheerful disposition, and we wish that a page had not been mutilated on which was written

*A notable sentence wherewith Sir John Salusburie was woonte
to solace him S[elfe].*

Often with a mery thought, do I myselfe well please
it is a thing that coste me nought, yet dooth . . . [paper torn].

[Christ Church MS. 184, f. 1

Three of Sir John's notes referring to plants which he grew in his garden at Lleweni are printed with accounts of other garden plants below, but his other notes all refer to stations of native Welsh plants, and are usually authenticated with the addition of his name in full, 'Sir John Salusbury, Knight'. The records are the earliest known to me for Denbighshire and the adjoining counties.

The plants, though interesting, are not very rare and had mostly been localized by William Turner, or by Gerard, in England. The locality for *Matthiola sinuata* appears, however, to be the earliest given for this plant in Britain.

WILLIAM SALUSBURY'S LOCALITIES, c. 1597.

- Radish (? *Raphanus sativus*). Transplanted by W. Salusbury from meadow adjacent to the Abbey at Maenan, near mouth of the Llugwy, to his garden at Llanrwst.
- Radish sp. Llanefydd; Denbigh; Llantwrog.
- Marsh Mallow (*Althaea officinalis*). Llansannan.
- Rest harrow (*Ononis spinosa*). Plas yn Llewini.
- Gromell (*Lithospermum officinale*). Denbigh; Whitford; near Mostyn.
- Chamomile (*Anthemis sp.*). Llanefydd; Llewini; near Llangollen; Dolgelly.
- Mistletoe (*Viscum album*). 'I saw it with berries in March near the Bontvaen near Chirk, and without berries all the way to Ludlow.'
- Fig. Conway; 'maesglas yn tegeingl, 2 m. from Holywell.'
- Great Cat's tail (*Typha latifolia*). Whittington Castle, 2 m. from Oswestry.
- Great Water Plantain (*Alisma plantago*). In a pool by the great house of Sir John Salusbury [presumably at Lleweni].
- Rye. Plas Llewini 1555.
- Maidenhair Spleenwort (*Asplenium trichomanes*). Bettws y Coed.
- Harts tongue fern. Talacre in Flint; in a wooded glen near Llanefydd.

SIR JOHN SALUSBURY'S LOCALITIES, 1606-8.

Papaver argemone L.

This hearbe is to be found by Llansanan hard by Ryd y Rienn or at Aber in Carnarvonshire.

Crambe maritima L.

English Sea Colewort, 'by my weare at Llanddylas upon the Baich and brine of the Sea, where there is no Earth to be seene but sande & pible stones'. In flower 30 May 1606.

Matthiola sinuata Br.

Purple sea stock Gilloflower, 'by seaside by Sir John Salusbury his weare theare at Llandulas'.

Rubus Idaeus L.

The Ruspie Bush or Hindebery doth growe by Moelvodiari¹ in Rees Tailors tenement & in grove of woodde Kylynllwyn behind the house of Berain, where I Sir John Salusbury found plentie of them growing wild yet naturally there. It is also found in Merionethshire, very comon by hedge ows and in th' topps of old thatched houses, and so likewise in the upp partes of Denbighshire at Spuddy and other places thereabouts.

Rosa spinosissima L.

I Sir John Salusbury found the Rose Pimpernell very comon in Garth snodnay Parke by Denbighe. Also in Merionethshire..

Sedum acre L.

Castle wall of Harden Castle. J. S.

Parnassia palustris L.

Mr. Tho. Willames, Clarke & phision, sent it mee Sir John Salusbury Knight, for another hearb. It groweth in a meade of Sir John Winn Knight.

Silvaus pratensis Bess.

English Saxifrage in the Copic in Llewenny parke next the newe stable over the high stile upon the right hand of the footway that leadeth from Llewenny Hall to Denbighe I found it the 23 Maye 1606.

Daucus Carota L.

Copic of Llewenny 23 Maye 1606.

Scabiosa arvensis L.

Purple flowered Scabious groweth neere St. Michells well or Fynnon Mihangil by Carwys, in the field that is above the well named Blorant,² in many other places neere Skewiog [Ysceifog] Church, in the field adioyning to the highe way upon the left hande as you come from Denbighe to the mountayne as you ride to Chester.

'Conyza maior' ? = *Erigeron acre* L.

Llewenny.

Gnaphalium erectum L.

Coten weede or Comon Cudweed groweth by Llewenny Brewhouse near to the Causaye theare.

Vaccinium myrtillus L.

This groweth in most of the montaynes of Wales.

Lysimachia nummularia L.

Herb twopence. Newburge in Llewenny Park & in wood called Kylynllwyn.

¹ Three miles west of Henllan. I am indebted to Mr. Edwards, the Librarian of Jesus College, for help with the spelling of Welsh place-names and for the loan of the *Herbal* of William Salusbury.

² Blorant in Parish of Aberwheeler near Bodfari.

Erythraea Centaurium Pers.

24 May 1606 Llewenny.

Scrophularia nodosa L.

Groweth in the Orchard at Chirck Castle.

Digitalis purpurea L. var.

Foxglove with white flower growes in a Parke of the Right Worthy Sir Thomas Myddelton called [? Castle] park [of Chirck ?].

Veronica officinalis L.

Fluellen 8 Oct. 1609.

Veronica spicata L.

Veronica serpyllifolia L.

Salvia verbenacea L.

Wild Clarie in great plentie in lital Park by the wall of the Castle of Denbigh.

Prunella vulgaris L.

Selfe heale or ye graite ynnos groweth plentifullie in Llewenny parke & the white flowered Self heale is found in a meadow of John Wyn mathenor of Llewenny green.

Ajuga reptans L.

Bugle or Middle Comfrey also white flowered Bugle I, Sir John Salusbury, Knight, found both in Llewenny park in the Coppice adjoining upon the River Cloyd where the herbes Adders tonge & Twiblade growe I found them the xxth of May 1606 growing there plentifullie.

Plantago coronopus L.

Buckeshorne. Weare at Llandulas on the side of the Bancks neare the weare. 3 May 1606.

Euphorbia paralias L.

Sea Spurge. Llanddylas 'risinge furthe of the Sandes' and Baiche of the Sea in very great plentie. 30 May 1606.

Ophrys ovata L.

Twyblade is likewise found neare Carrwis in a place called Cadnant, where a faire well springeth called St. Michael's well, in Welsh ffynnon Mihangel. [And in the] Castle Park of Chircke in a close next the lower Barnes (belonging to the Rt. Hon. Sir Tho: Myddelton). Twiblade are found by Chambers wood in a field called Ravenscroftes field in Wales neare Denbigh and likewise is found hard by Cloyde in a field of John ap Roberts of Pont Gruffith & upon the banks of the river Wheler, neare the house of the parson of Botuarry. The Herbe Addertonge groweth likewise in the lower end of Ravenscroftes field.

Paris quadrifolia L.

Herbe Paris is found neare Carrewis in a place called Cadnant, where a faire well springeth called St. Michael's well, ffynnon Mihangel within a bolt shot of the well down the spring, one that side of the water as Carewis standeth, where like wise is found the hearb Twyblade and by reason of the ranknes of the place there are found a greate store of herbe paris with five leaves apeece, but the yeare 1606 I found the same with six leaves.

Sir John states that he planted them in his garden in 1608.

Ophioglossum vulgatum L.

The Herbe Addertonge groweth likewise in the lower end of Ravenscroftes field. [See above.]

Botrychium Lunaria Sw.

Lunaria minor is found in Cunnygree of the Right Hon. Sir John Salusburys, Knight, lying betweene Botuarry [Bodfari] & Carewis, and great plenty of them are found in Place y Chambers fielde lying hard by Snodioge parke¹ neare Denbigh being the highest & the next field to the parke on the left hand as you go to Henllan from Place y Chambers, in a place of Llanywith called Ogoyr graig upon the side of the banke theare, and are found in the littel park of Denbigh in the syde of a hill. . . [continuation cut in rebinding].

Poisonous Fungi.

Let my advice perswade thy mynde
not to truste any of that kynde
such as be takenn for the beaste
doe proue as poisnusse as the reste.

J. S.

With these notes are others concerning the medicinal properties of herbs, on which Sir John Salusbury was also an authority. But in none does he show his wisdom as clearly as in two lines in his poem on *Certaine Necessary observations for Heath*, 1603.

Apothecaries shop of drugges let not thy stomack be :
Nor use noe phisick till thou neede, thy frende adviseth thee.

vi. THE MSS. OF MATHIAS DE L'OBEL AND WILLIAM HOW.

Among the papers which came to Magdalen College with the botanical Library bequeathed by John Goodyer in 1664, were some thousands of printed slips cut from Lobel's *Adversaria*, 1576, *Observationes*, 1576, and *Icones Stirpium*, with the author's corrections and MS. additions. These had evidently formed part of a pasted-up copy prepared by himself for a projected work which he did not live to publish, but from which How printed a selection under the title, perhaps the same as the one Lobel himself would have chosen, of *Stirpium Illustrationes*.

The first question that occurs to one is, To what extent do these MSS. throw light on the life and work of Lobel?

Lobel was born in Flanders in 1538. Like d'Aléchamps (1513-1588), Clusius (1526-1609), Pierre Pena and Jean Bauhin (1541) he studied at Montpellier under Guillaume Rondelet, who is said to have taken such a liking to the serious young student, that he

¹ 'Snodiog Park' is marked as a round enclosure between Lleweni and Denbigh in old maps of the county.

bequeathed his botanical manuscripts to him. In the Goodyer collection there is a parchment cover that was used to hold loose papers, which bears evidence to the association, for inside is written *Rondelet de Febribus*. It may have originally served to contain notes taken by Lobel at Rondelet's lectures.



LOBEL.

Some ten years later we find Lobel living in England. Driven from his native land by civil war, he described himself as being unable to make a home on a sea tossed by incessant tempests, or to indulge his peaceful love of gardens and flowers on lands watered with human blood. He came over to England to produce his first great work, the *Adversaria*, and in return for English hospitality he dedicated the work to Queen Elizabeth. The book was printed in London in 1570. It was our first scientific Botany. In Turner's

Herball the plants were arranged in alphabetical order. In Lobel we find the first striving after a natural classification, and for the first time the straight-veined plants, now called Monocotyledons, were partly separated from those with net-veined leaves (Dicotyledons).

Five years later we find him back in his native country and in close alliance with Plantin, printing his *Stirpium* or *Plantarum Historia* illustrated with 1,486 wood blocks, to which the *De Succedaneis* of Rondelet was added. And six years later he moved to Delft to superintend a Dutch translation of his work, the *Kruyd-boeck* printed there in 1581.

At the age of fifty-four he was again in England, superintending a Botanical Garden which had been established by Lord Zouch at Hackney. In this he was probably in friendly rivalry with Gerard, who, seven years his junior, was cultivating eleven hundred kinds of plants in his garden in Holborn. Lobel in fact prefaced the 1596 catalogue of Gerard's garden with a printed letter of eulogy, but a note in his own hand in the copy in the British Museum (N.H.), 'haec esse falsissima M. Lobel,' is distinctly unkind. Dr. Daydon Jackson tells me that this is the only specimen of Lobel's handwriting that was known to him before he saw the Goodyer manuscripts.

About 1606 Lobel was honoured by being appointed King's Botanist to James I, but feeling the weight of his sixty-eight years, retired, it is believed, to live with his son-in-law James Cole at Highgate. The will of JAMES COLE,¹ a document of great interest, presents a graphic picture of the wealth of Lobel's son-in-law, who evidently maintained intimate relations with the Low Countries to his dying day. He was engaged probably in the spinning, and certainly in the importation of silk into this country. He left his house at Highgate to his wife Louisa, then to Abraham Bush, his sister's son. His house in Lyme St., held on lease from the Carpenters' Company, he left to his nephews Henry and Peter Cole, and Henry was also to have a 'gilte cuppard with the arms of Antwerpe graven therein' and his chain of gold. To Abr. Bush

¹ Will (P. C. C. Barrington 42) written on six leaves of paper, dated 31 Dec. 1627, with a codicil witnessed by Eliz. van de Bossche and Louise Cool, proved May 1628. It is to be hoped that one day a more complete account of Lobel's relations will be forthcoming. In addition to James Cole, he mentions another son-in-law Ludovic Myres, an authority on pharmacy, Abraham Hoguebat, pharmacist, son of his second wife, and Michael de Lannoy 'affinis meus'. His second wife may have been related to the Hugobert mentioned by Goodyer, cf. p. 59. Was de Lannoy synonymous with de Laune?

‘all my bookes¹ as well concerning my Latin studies, as picturenge with all my printes, little pictures, shells, marbers, statues, and all my antiquities and old Coines and such like rarities commonly shutt within my Counting House in Lyme St.’ To numerous friends and relations with Dutch names he bequeathed silver cups and pieces of plate.

Under Cole’s roof Lobel probably added the last touches to a compilation, the manuscript of which is now before us, and which we may call the *Stirpium Illustrationes*. The date of the work is settled by the imprimatur which bears the signature of Thomas Moundeford in his official capacity of President of the College of Physicians. He filled this post from 1612 to 1614.

The original testimonials with which Lobel had provided himself for two of his works are still extant. They are both from the College of Physicians, and are signed by the most eminent doctors in London. The names on the earlier document take us back a long way, to 1605, two years before Harvey became a Fellow of the College. The signatures are headed by :

Thomas Langton² as President ; then follow John Craig,³ who is said to have given Napier a hint that led to the invention of logarithms ; Sir William Paddy,⁴ Sir T. Turquet de Mayerne,⁵ Henry Atkins,⁶

¹ Two of James Cole’s books are known to me. One a *Dodoens* in the Goodyer collection ; the other, Clusius, *Per Hispanias*, formerly in the British Museum, but turned out as a duplicate in 1769, and now in the possession of my friend Dr. Daydon Jackson. Both books have his signature *Jacobi Colei* on the title-page.

² THOMAS LANGTON, M.D. Cantab. F.R.C.P. 1581 ; President, 1604, 5, 6. *d.* 1606.

³ JOHN CRAIG, M.D. of Bâle. First Physician to James VI of Scotland. F.R.C.P. 1604. He attended James I in his last illness.

⁴ Sir W. PADDY, M.D., of St. John’s College, Oxford, where a monument records his great benefactions to the College. Physician to James I. President of the College of Physicians, 1609-11, and in 1618.

⁵ Sir THEODORE DE MAYERNE, M.D. *b.* Geneva 1573, *d.* 1655 ; M.D. Montpellier 1597 ; F.R.C.P. 1616. Physician to Anne of Denmark, Charles I, and Charles II.

⁶ HENRY ATKINS. President of the College of Physicians, on seven occasions, 1606-25. He started with the naval expedition of 1597 as physician to the Earl of Essex, but was so bad a sailor that he had to be put ashore at Plymouth, and the College was ordered by the Queen to select another medical man to accompany the expedition. The choice fell on Dr. Moundeford. Dr. Atkins was chosen by James I to fetch his younger son, subsequently Charles I, then an infant from Scotland. *d.* 1635. His will has recently been acquired by the College of Physicians.

Nos infrascripti Medicinae Doctores fidem facimus Lobelij Scripta
& Lucubrations ubique terrarum apud doctissimos Medicos & in celeberrimis
Academijs Valere; quanto magis praevalent, & huius florentissimae
Regij iam decoratae illustriores erunt, in humani generis & Reipublicae
Ierusalensem, si postremam manum in semet ipse sua addiderit, & quae plurimum
Britannicis indolis Plantis, earum virtutibus reclusis, nobilitavit, totius orbis
spectandas dederit, quae hoc expectatissimum opus summo opere videre auct.
Quod hoc nostris chirographis attestamus.

Tho. Langton praesidens Collegij Medicorum London.

Jean. raiques
ordinaire de la Cour

Hs. Aitme Richard Fosb, William Barrow, & Thomas Fran

Willelmus Cassy

Martinus Seanevus

Britanniae Reg. Medicus ordinarius

Commaerme Turquet.

frons Reg. Medicus ordinarius.

gubernator principibus
primarius et ordinarius regis

Will: Dun

Adamus Britanni regis ordinarius
regis regis, et ordinarius

Bellin

Fran. Fleming

Matth. Guime

John. technicus

OLD TESTIMONIAL TO LOBEL

Signed by the President of the College of Physicians,
and by other eminent Doctors

Sonner.

De Dressement, tressage et vertueux f.
Mons^r le Lobel en edec^r de Le^rgall.
De monseigneur de France D'Orange Cos^r.

Le Vray de tes vertus, L'amic mutuelle,
Etouurant tes faiz: quantq^e L'Escluz & tiens
Mier & uels maternel: (L'obel) fait que bilis
S'ens^r vers ton somer ma Raice Chalonnele.
Q'us est vray que d'uroit ta memoire eternelle
Estre, par auech^r d'han^r d'indue, et plus parfaith.
Mais quand venra le Cocu, les actions les faith
De Cil, qui te chanter vray embat se boure.
Te dyz comme haultment Te te p'cut f'edomde
En son Luth mal-poz ta Louange d'indomde:
Diras, d'm, t'air, toy Te cognois ton Couraige
Surpasser ton moy d; alors Te tout k'elz
Le hauct Dieu suppleur Ma sera pas, quantif
Pour au foug d'admitie vous tenir vray songe.

De Vos saiges Docteur
Indigne Ruker
D'ay ma^r

SONNET DEDICATED TO LOBEL BY MAES

Richard Forster,¹ William Baronsdale,² Thomas Frear,³ William Dun,⁴ D. Sellin,⁵ Francis Hering,⁶ Matthew Gwinne,⁷ all in Munk's *Roll* of the College. Gilbert Primrose, though not on the *Roll*, was the father of James Primrose who was; J. Nasmyth,⁸ surgeon to James VI of Scotland, had only recently arrived in London in attendance on his royal patron. Gwinne's *Vertumnus* was acted at Magdalen College in 1607. Nasmyth had already presented Lobel with some plants which the latter had already acknowledged in print. This testimonial was accompanied by a eulogistic Sonnet written by Jan Maes. Neither testimonial nor sonnet appear to have been published.

The later testimonial is signed by Thomas Moundeford,⁹ President, Henry Atkins, Richard Forster, Thomas Friar, Mark Ridley,¹⁰

¹ RICHARD FORSTER, M.D., of All Souls College. F.R.C.P. about 1575; President, 1601-3, and 1615. Author of *Ephemerides Meteorologicae, ad annum 1575, secundum positum Finitoris Londoni*. 8vo Lond. 1575. When out walking with Dr. Herring and Lobel he found a new grass, 'Gramen supinum Monspeliense inter Islington et altam portam, vernaculè Highgate'. (Lobel.)

² WILLIAM BARONSDALE, M.D., of St. John's College, Cambridge. President of the College of Physicians, 1589-1600. He died before 7 June 1608.

³ THOMAS FRYER, M.D. of Padua, of Trinity College, Cambridge. F.R.C.P. 1572. He incorp. Doctor of Physic at Oxford, 28 Feb. 1623.

⁴ WILLIAM DUNNE, of Exeter College. F.R.C.P. 1592. Died before 16 May 1607.

⁵ DANIEL SELIN, M.D., of Magdalene and Christ's Colleges, Cambridge. F.R.C.P. 1599. *d.* 1615.

⁶ FRANCIS HERRING, M.D., of Christ's College, Cambridge. F.R.C.P. 1599. He was present at the finding of 'Gramen supinum Monspeliense' between Islington and Highgate, and wrote a Latin poem *Epigramma in Opera novissima L'Obelii* as a mark of his 'love and friendship', which was printed at the end of Lobel's *Animadversiones in G. Rondelletii Pharmaceuticam Officinam*. Lond. 1605.

⁷ MATTHEW GWINNE, M.D., Fellow of St. John's College, Oxford. F.R.C.P. 1605; First Gresham Professor of Physic, 1598-1607; author of the Latin comedy *Vertumnus*, acted before James I at Magdalen College. *d.* 1627.

⁸ JAMES NASMYTH. Surgeon and Botanist to James I.

Provided Lobel with new drawings of 'Hyacinthus stellatus Peruanus' and 'Narcissus Indicus rubro flore', one of Robin's new plants. Lobel, *Adv. alt.* p. 482. He evidently had a garden in which he grew and flowered

Fritillaria nigra Pyrenaea. 1605. Lobel, *Adv. alt.* p. 496.

Crocus Byzantinus serotinus candidus. p. 498.

Colchicum minimum tenuifolium Gallaecium. p. 500.

Plantago Umbilicata. Staghorn fern. (*Platyserium aethiopicus*). p. 506. 1604-5.

⁹ THOMAS MOUNDEFORD, M.D., of Cambridge. Pres. R. C. P. 1612-14, 1621-3.

¹⁰ MARK RIDLEY, M.A., of Clare Hall. F.R.C.P. 1594. Died before 1624.

Edward Lister,¹ Richard Palmer,² John Argent,³ and Matthew Gwynn. It was printed by How in 1655 together with an original letter from Dr. Argent, who will always be remembered for having obtained from Bermuda the original bunch of Bananas, which was exhibited in Johnson's shop on Snow Hill.

Lobel died in 1616 (four years before the birth of his editor, William How). James Cole inherited his manuscripts, and Parkinson, when collecting all available material for his *Theatrum botani-*

Tho: Alcockford preside:
 H: Atkyn.
 Richardus Forsley Thomas Friar
 Marcus Biddens.

E. In. Lister.
 Ric: Palmer.
 Job. Argent.
 Matth: Gwynn.

SIGNATURES TO LOBEL'S SECOND TESTIMONIAL.

cum, was undoubtedly permitted to see some or all of them. Lobel wrote in Latin: Parkinson would therefore have had to translate, and a strict sense of honesty in the obligation to make acknowledgement may have become blunted in the process: he merely mentioned Lobel's help in general terms, for Lobel was a foreigner, dead, and perhaps no one cared.

¹ EDWARD LISTER, M.D., King's College, Cambridge. F.R.C.P. 1593. Physician in ordinary to Queen Elizabeth and James I. *d.* 1620.

² RICHARD PALMER, M.D., of Christ's College. Took part in the consultation at the death-bed of Henry, Prince of Wales.

³ JOHN ARGENT, M.D., of Peterhouse. F.R.C.P. 1597; President, 1625-7, 29-33. *d.* May 1643. An autograph letter from him to Lobel is among the Goodyer MSS. He is remembered for his enterprise in causing his man to

Ten years later William How, a young doctor of St. John's College, Oxford, was assiduously collecting notes for the first British Flora. Concise in the choice of its information and scrupulous in its duty of quoting authorities, the *Phytologia Britannica* of How was the antithesis to the *Theatrum* of Parkinson. It was printed in 1650, almost certainly before the author had acquired the Lobel MSS. About 22 Jan. 1651 Lobel MSS., or some of them, appear to have been in the hands of a man of learning,¹ who then copied out many Latin descriptions of plants (Goodyer MS. 8, ff. 104-21).

How was a most indefatigable hunter after exact localities of plants. He would have searched the *Theatrum* in vain for information which he afterwards found clearly given in Lobel's MS., and which he would have assuredly included in the *Phytologia*, had only Parkinson quoted authorities: and Parkinson had still further transgressed by taking Lobel's credit of priority to himself.

This appears to be one explanation of the violence of How's criticisms of Parkinson's lapse from the higher standards of literary honesty. My friend Dr. Church has suggested another motive which might well repay a more extended inquiry. Parkinson was a Puritan, whereas How was probably, like Goodyer and his friends, a Royalist.

It is reasonable to suppose that How's first idea after realizing the originality of Lobel's work was to publish it *in toto*. It was, however, in Latin, and the market had already been spoilt by the appearance of two popular works, Johnson's *Gerard emaculatus* in 1633 and Parkinson's *Theatrum* in 1640, and no publisher would undertake a third.

Yet Lobel's 'volumes were compleat, The Title! Epistles! and Diploma affix'd'. How, indignant that Parkinson had, as he put it, 'murdered his (Lobel's) genuine scrutiny in treacherous oblivion,' and perhaps dimly conscious that his time for work through failing

follow the roots of a species of 'Pease' by scrapping away the beach between Orford and Aldborough, 'vntill hee got some equall in length vnto his height, yet could come to no ends of them'. *Ger. emac.* 1251. And during the last year of his Presidency of the College of Physicians, he gave Johnson the first bunch of Bananas that was ever exhibited in a London shop. *Ger. emac.* 1515.

¹ The handwriting of the unknown commentator is characterized by the frequent use of scrolls. We have noted it in Goodyer MSS. 8 and 9 and in Goodyer's copy of Parkinson's *Theatrum*, and have evidence that the writer was closely associated with Goodyer about 1650 to 1659. There is reason to believe that the writer was Goodyer's friend and neighbour, Dr. John Dale of East Meon and of Long Acre in London, who died in 1662.

health was short, made a selection of Lobel's descriptions under the title *Stirpium Illustrationes; plurimas elaborantes inauditas plantas subreptitiis Joh: Parkinsoni rapsodiis (ex codice MS. insalutato) sparsim gravatae*. The work was printed by Thos. Warren for Jos. Kirton of St. Paul's Churchyard in 1655.¹

We have the original manuscript from which the book was printed before us, with the excerpts from Lobel, with How's additions pasted or pinned thereto, and the leaves of the MS. exactly as they were marked for the compositor, and returned by the printers to the editor.

After How's early death on 30 Aug. 1656, his own annotated copy of the *Phytologia* passed to John Goodyer on 30 Apr. 1659, and with it probably the Lobel MSS. as well, but too late for them to be of real use, for Goodyer's working life was drawing to a close. Except for their disarray through having been ungunmed and misplaced, we may assume the papers to be in the state in which How left them.

In this volume there are thirty-seven leaves.

First comes the original imprimatur with the signatures of Tho. Moundeford, the President of the College of Physicians, and of eight other members. Then the Preface, with many lines erased in Lobel's hand, and some eulogistic verses by Alexander Rhedus of which the last eight lines were not printed. Next follow some introductory remarks by How, and his Index and list of erratula in his own hand.

Then Lobel's descriptions of 223 kinds of plants, a large number of which were apparently claimed by Parkinson as his own discoveries. How's notes, which appear in small type near the margin of his printed book, are intercalated in the Lobel MS., and show that the selection and arrangement of the volume was entirely the work of How, and not of Lobel.

How evidently had Lobel's materials for the larger book before him. He cut out the descriptions which appeared to him to be of the greatest importance. There are also included original letters from JOANNES DE MONNEL² and from JOHN ARGENT, dated 'Wood Street 2 June 1608', and also notes on various plants from Montpellier communicated by, and apparently in the handwriting of, Pellisierius. At the end is one leaf of ἀμάρτηματα in How's

¹ John Goodyer received his copy on 19 February 1654.

² JOHN MONNEL of Tournay was a correspondent of Clusius. Parkinson associates him with 'Anagallis tenuifolia fl. coeruleo' which he received from Cadiz and grew in his garden at Tournay. *Theatrum*, p. 559.

autograph, in which more than a score of Parkinson's errors are pointed out.

The remainder of Lobel's materials and the manuscript of his projected *Stirpium Illustrationes* have now been bound in three volumes, for convenience in handling.

Volume 1 contains the classification and description of 223 kinds of grasses. Lobel had evidently become acquainted with a great many more species than the some forty-five he knew when he printed his *Historia plantarum* in 1576. But evidently the very novelty of the newer descriptions has brought about the spoiling of the MS.—from which How cut out accounts of ninety-eight different grasses, for his book printed in 1655.

Lobel had evidently worked hard at his grasses. The different kinds described have been numbered and renumbered in some cases four times over, and many alterations have been made in the MS.

The last fourteen leaves of this volume are from a pasted-up copy of the *Adversaria* which has been much cut about.

Vols. 2 and 3 are built up out of a further portion of the pasted-up copy of the *Adversaria*, the leaves of two copies of the 1576 edition being pasted on leaves of paper. The greater part of the text has been struck out, and marginal references to the pages of the *Observationes*, 1576, have been added. In vol. 2 the leaves have been roughly numbered by How (?) 1 to 134, and in vol. 3 the leaves run from 135 to 251. The plants are numbered up to No. 835.

Between the leaves so prepared he intercalated the leaves of a printed copy of the *Observationes* (edit. 1576), and between them again the illustrations from his 1591 edition of *Icones Stirpium* printed by Plantin. In many cases the page-margins are filled with Lobel's additional MS. notes on the plants.

There are no references to Goodyer, nor traces of his handwriting, on these Lobel MSS.

vii. WILLIAM MOUNT AND HIS RECORDS OF KENTISH
PLANTS IN 1582-4.

Among the books in the Goodyer Library are two editions of the *Icones Stirpium* by Lobel, printed by the Plantin press. The earlier copy of 1581, according to a note on the title-page, was purchased by 'Gulielmus Mowntuus' for 19 shillings on May 20, 1582. This note was written by Mount himself, and his initials, 'W. M.', are stamped upon the leather binding. Several notes,

written in the margins of the illustrations, appear to have been added by him within a few months of his acquiring the volume. In this, the earlier of the two volumes, there are no notes by Goodyer; but in the later edition of 1591, containing no notes in Mount's hand, there are numerous notes by Goodyer, including copies by him of all Mount's notes on plants growing in Kent, taken from the copy of 1581.

It seems likely that Goodyer having first acquired the later edition, copied Mount's notes into it, perhaps borrowing them for the purpose. He subsequently obtained possession of the earlier copy and owned both volumes before 1633.

It has been ascertained¹ that Mount was born at Mortlake in 1545, was educated at Eton and King's College, Cambridge, of which he was admitted scholar in 1563 and fellow in 1566. He resigned his fellowship before Lady Day 1570. It appears that he owed much to the patronage of Secretary Sir Thomas Smith and Lord Burghley.² In a letter addressed to Sir William Cecil on 20 Oct. 1567, he professes his great satisfaction in being placed at the University under the patronage of the former, his 'most honoured Mecaenas'. Medicine was the first object of his studies: later he took orders and was appointed Master of the Savoy in January 1593-4, and died in December 1602.

It was known that Mount had taken considerable interest in the making of distilled waters, an art which he probably learnt as part of his medical studies, and that he had written some Latin verses prefixed to Lobel's *Balsami, Opobalsami, Carpobalsami et Xylobalsami explanatio* in 1598, but he has not as yet received from botanists the credit which he deserves of having been the first to record the provenance of several plants in the county of Kent, nor for his knowledge of the construction of perpetual calendars.

In the collection of medical books which my friend the late Sir William Osler bequeathed to McGill University, there is a manuscript to which Mr. Craster has recently drawn my attention: it throws a new light on the knowledge of the Kentish botanist. It is 'A shorte declaration of the meaning and use of a perpetuall calendare or almanack' by W[illiam] M[ount], in eleven chapters, with dedicatory preface to Sir Thomas Bromley, Lord Chancellor of England, whose chaplain the writer was, and whose arms are blazoned in colour on p. vi. The work, which is the author's holograph, is illustrated by coloured tables and diagrams. Three

¹ Cooper, *Athenae Cantab.* ii, p. 271.

² *Calendar State Papers Domestic*, 1547-80, pp. 294, 301.

of the latter contain the arms respectively of the Universities of Cambridge and Oxford, and of Queen Margaret of Anjou, foundress of Queen's College, Cambridge.

It was written by William Mount in 1583; and the hand is identical with that in which the entries in Goodyer's copy of Lobel's *Icones* at Magdalen College, are written.

Mount's botanical localities were mostly within four miles west and north-west of Maidstone, indicating that he was living in Kent in 1581 at East Malling, where he had an orchard. And even in the case of plants for which Mount gives no localities, it may safely be assumed that he was referring to specimens growing near his Kentish home. His notes appear to have been written between 1582 and 1584. The plants which he may then have observed in Kent, and for which his are the earliest references, are about thirty-three in number. Of these thirty-three plants we find that Gerard, who wrote thirteen years later than Mount, mentions only eight, and to these Johnson adds only eleven more in his various works printed 1629 to 1633, forty-five and more years later. These notes of William Mount, therefore, constitute an important contribution to a first Flora of Kent,¹ with a few notes on the virtues of imported garden and medicinal plants; and it must be remembered that they were compiled in days when the idea of local floras was as yet unknown.

The modern names of some of the plants for which he has recorded dates and localities are included in the following lists.

NATIVE PLANTS.

<i>Modern Names.</i>	<i>Localities recorded by Mount c. 1582.</i>	<i>First printed records.</i>
Poa pratensis, L.		Gerard, 1597.
Poa trivialis, L. (?)		
Eragrostis major, Host.		Ger. 1597.
Phleum pratense, L., var. nodosum.	Mount's 'Alderes'.	Johnson, 1629.
Carex acuta, L.	East Malling.	Curtis, 1670.
Juncus acutiflorus, Ehrh.	Snodland.	Johnson, 1632.

¹ I have found some other references to plants dating from the sixteenth century in a copy of Lyte's *Herbal*, 1578, belonging to the Radcliffe Trustees in Oxford. Two of the notes certainly refer to Kent, possibly the others may too. There was a Smallbrydge in the manor of Horsmonden.

- Ground pyne (*Ajuga Chamaepithys* L.) 'luxuriat in Cantio'.
- Rhus sylvestris Plin. (*Myrica Gale* L.) 'Canterberyie Call and Cole'.
- Rhamnus (*Rhamnus catharticus* L.) 'au pres de la forest de Hatiele' (?).
- Buckthorne (*Hippophae Rhamnoides* L.) 'au pres de small bregge'.
- Plane (*Platanus orientalis* L.). Planted in England 'at Small bregge'.

The signature of the recorder is doubtless on the title-page, but it has been obliterated by over-scribbling.

<i>Modern Names.</i>	<i>Localities recorded by Mount c. 1582.</i>	<i>First printed records.</i>
Hordeum murinum, L.	About London.	Turner, 1548.
Luzula vernalis, DC.	Mount's Orchard.	Ger. 1597.
Eleocharis uniflorus, Reichenb. (?)	Sandwich, Dover.	
Triticum repens.	Addington.	Ger. 1597.
Agropyrum junceum, Beauv. (?)	Addington.	
Cynodon Dactylon, Pers.	} Addington, 1581.	Ray, 1688.
or		
Digitaria glabra.	Leybourne.	Hooker, 1829.
Sparganium ramosum, Huds.	'Very common', prob.	Johnson, 1629.
Narcissus pseudo-narcissus, L.	in Kent.	Ray, 1724.
Scilla autumnalis, L. (?)	Sea-coasts of England.	Park. 1629.
Allium ursinum, L.	Wrotham.	Martyn, 1763.
Bupleurum rotundifolium, L.		Johnson, 1629.
Coronopus Ruellii, All.		Ger. 1597.
Stellaria media, Vill.		Turner, 1735.
Lysimachia Nummularia, L.		Johnson, 1632.
Rhinanthus Crista-galli, L.	Common.	Johnson, 1629.
Digitalis purpurea, L.		Johnson, 1632.
Tussilago Farfara, L.		Johnson, 1629.
Nymphaea lutea, L.		Jacob, 1777.
Bryonia dioica, L.		Johnson, 1629.
Tamus communis, L.		Johnson, 1629.
Potentilla anserina, L.		Johnson, 1629.
Pedicularis palustris, L.	East Malling.	Merrett, 1667.
Fumaria officinalis, L.		Ger. 1597.
Ceterach officinarum, Willd.	Maidstone, 1583.	Park. 1640.
Botrychium Lunaria, Sw.	Blackheath; near Rochester.	Lobel, 1570.
Asplenium Ruta-muraria, L.		Ger. 1597.
Rhamnus catharticus, L.	Dover-Folkestone, 1582.	Ger. 1597.

GARDEN AND EXOTIC PLANTS.

Zea Mays, L.	Morgan's garden.
Acorus Calamus, L.	D. Penny's garden.
Gladiolus communis, L.	{ Morgan's garden, 1578.
	{ Mount's Malling garden, 1583.
Crocus sativus, L.	Saffron Walden.
Colchicum autumnale, L.	Mount's garden, 1583; and at Bath.
Saponaria Vaccaria, L.	Lord Abergavenny's garden, 1584.
Ipomaea Jalapa.	Root used medicinally in England, 1580.
Anthriscus cerefolium.	

THE MANUSCRIPT NOTES OF WILLIAM MOUNT.

On Title-Page.		
	'Gulielmus Mowntuus 19 ^o . Maij 20. 1582.'	
Page.		
1. Gramen pratense.	'Great leavyd Medowe grasse very vulger.'	<i>Poa pratense</i> L. ¹
1. Gramen minus.	'The lesser vulger Grasse.'	<i>Poa trivialis</i> L. (?)

¹ Lobel's illustrations of these grasses have been determined by Dr. Stapf, who notes that they are 'mostly too crude for us to name from them critically, and Mount would not have been in a better position, except perhaps in so far as he may have been supported by some tradition that is lost to us'.

7. Gramen paniculosum Phalarioides. *Eragrostis major* Host.¹
 'Grasse called in Surrey, braunched grasse in Coarne : and in orchardes, & shaddowye places usually mowen. They seathe y^t in water wth purselane and small Peysons for wormes, in the sommer tyme, and gyve y^t commonly to very younge chylderen.'
10. Gramen typhinum. *Phleum pratense* L., var. *nodosum*.
 'An other Sedge Grasse in watery moyste places in my Alderes & (?) muche.'
11. Gramen palustre maius. *Carex acuta* L.
 'The sharpe edge grasse flaggis he in black brookes [in Est mallinge] growinge in Tuffettes, very Common.'
12. Gramen aquaticum alterum. *Juncus acutiflorus* Ehrl.
 'Ponde grasse growinge in a Ponde in Snodelande, w^{ch} ys some tyme allmoste drye.'
13. Panici effigie, Gramen simplici spica. *Hordeum murinum* L.²
 'Barley grasse : because y^t resembleth Barleye in hye wayes, and pathes about London.'
16. Gramen hirsutum nemorosum. *Luzula vernalis* DC.
 'Hearye or hoarye Grasse in my orcharde.'
17. Gramen iunceum marinum dense stipatum. *Eleocharis uniflorus* Reichenb. (?)³
 'Pusshye grasse : in the sandes by the Castles betweene Sandwyche and Douer (in Kent).'
20. Gramen Canarium. *Triticum repens* L.
 'In all places where the wryters use this worde : Gramen : It ys to be understoode y^t they meane Quyceke or Couche grasse, whereof there be dyverse kyndes well knowen. The best in my opynyon hathe longe greate rootes creeping in lengthe ij yeardes greater than wheate strawes full of ioyntes agroinge wth ye shape, fol^o. 23. Yt groweth in Addington in Kent in the sandy drye dustie hye wayes there. The same prevayleth against the straw, and against wormes in my experience.'
22. Gramen caninum longius radicatum marinum alterum. *Agropyrum junceum* Beauv.
 'Of Dogges Grasse, or Couche grasse, thus writeth Poena, and Lobell, fol. 2. . . . I have used the Grasse with ye great roote, growinge in sandye wayes set downe here, fol^o. 23 : growinge in Addington in Kent and doe fynde yt muche better then the usuall Couche grasse. W^m: Mount.'
23. Gramen caninum alterum. *Cynodon Dactylon* Pers., or *Digitaria glabra* Beauv.
 'Quyce grasse growinge in sandye wayes the very trewe Cowche grasse whereof the Phisiciones wryte. It groweth in the sandy hie waye plentifullye betweene Wrothame and Addington
- ¹ Dr. Stapf writes: 'How this came to be noted down for Kent is a puzzle, unless he saw it in a garden. Tabernaemontanus says it was grown in gardens, and as it had an old popular name "amourette", it may not have been unknown in English gardens of the period.'
- ² The figure and name are of *Setaria viridis* Beauv. (O. S., J.).
- ³ The figure is *Scirpus caespitosus* L. (O. S.); the name belongs to *Bromus asper* (J.).

in Addington in Kent. (I there fownde y^t a^o [15]81 and have synce often used it against wormes wth good successe and ye stoane also euen in my selfe for ye stoane.)

33. [Quotation from Pena and Lobel.]
40. Milium indicum rubrum. *Zea Mays* L.
 ‘Redd Indyan Myllitt, which I have seene in M^r Morgan hys house, the Queenes apothecarie in London.’
49. Harundo saccharina Indica. *Saccharum officinarum* L.
 ‘The suger Canes which yielde us our best suger of it selfe suger without compoundinge, or connynges which ys in deede good and comfortable: the other here in Suger houses by pollycie denyed, ys perchance more profitable to the maker or merchaunte, then healthfull to the partie, which ys to use it.’
55. China. *Smilax China* L.
 ‘The diet roote Chinee whereof we have none growinge in Englande and yet muche used: it hathe byn in Englande for the Great pockes, also for all deseases proceedinge from a moiste brayne and the lyuer obstructed. Hereof I referre the reader to Vesalius whoe hath lardge written in his book *de radice Chyna* lardgelye thereof.’
57. Acorus Diosc. & Acorus Theophrasti, Officinis falso Calamus. *Acorus Calamus* L.
 ‘I have hearde M^r D. Pennye often saye that he hath this roote in his garden in London.’
59. Iris nostras vulgaris. *Iris Pseud-acorus* L.
80. Sparganium & Butomus Theophrasti. *Sparganium ramosum* Huds.
 ‘The sharpe edged burre flagge. It groweth in watery dytches about Leybourne in Kent, smale use thereof in Phisick or Chyrurgie.’
98. Gladiolus Narbonensis. *Gladiolus communis* L.
 ‘Coarne flagge or Coarne gladdyn. I have y^t my garden M^r Morgan gave yt to me a^o 1578 at London yt groweth nowe with me a^o 1583, at Mallinge.’
112. Narcissus poeticus. Daffodil. *Narcissus Pseudo-narcissus* L. and others.
 ‘Theis herbes which are set here under the name of Narcissus we calle Daffadowndyllyes: they be very common, and of them (as here they be set) dyuerse sortes of dyuerse colores.’
137. Crocus sine flore. Croci flores. *Crocus sativus* L.
 ‘Safforne withoute flower I have never sene here: safforne with the flower I have sene plentye at Safforne Walden in Cambridge-shyre as I thinck yt ys and in many other places in Englande. Also yt ys concluded amonge all the most approued Authours y^t of Englishe Safforne ys the best and hath the greatest virtues.
 ‘The Qualities of safforne shall followe as the Roöme wyll permytt soe farre as there ys aney mency one by shape or picture made of safforne.’
143. Colchicum sive Strangulatorium Ephemereum Crocifolium. Autumn crocus. *Colchicum autumnale* L.
 ‘Thys we calle Hermodactylus: y^t was gevin me a^o 1578; and groweth now in my garden a^o 1583; very trewly y^t agreeth wyth this shape [cf. figure in Lobel, *Icones*, 1581]. I did neuer see

aney flower but only poddes, as here set downe yet y^t may be that y^t hath the flowers here underset and soe I thinck y^t hath.

D. Symyns¹ tould me, that y^t groweth plentifully about the bathes in Somerset shyre or Wyltshyre when he sawe yt in pastures whereof when casually their catle chaunce to feed, they become daungerus syck untill ye herdsman or keper have well starved them euen allmoste to sweate; and thin y^e swellinge and other accidents doe diminishe, also they kepe y^e cattell from drynck whyle they be yll. Thys ys Mr. D. Symyns observacon of thys herbe in that countrie, as I took the same from hyme by note.²

[Turner (*Herbal*, 1568, p. 156) figures the Widdowe Saffrone from Bath both in flower and seed, but appears not to have known of its dangerous qualities to cattle. He is eloquent about it in relation to man.

'It is good to knowe this herbe that a man maye isschewe it. It will strangell a man and kyll him in the space of one daye, even as some kinde of Tode-stolles do. The roote is swete and provoketh men thereby to eate of it. If anye man by chaunce have eaten anye of thys, the remedye is to drinke a great draught of cowe milke.'

'It stirreth up tossinges, wamlings, windinesse and vomiting' (Lyte, 1578, p. 367).]

150. Cepe.

Onion. *Allium Cepa* L.

'Unyones fynely slysed, and in faier water one night steiped; the water y^s next morninge geven to chylderen which have the wormes wonderfull effectuallye helpeth them. Lonicerus, fol. 193. Tragus also first used yt, fol. 739.'

151. Scillas. Cepamarina. *Scilla verna* Huds. confused with *S. maritima* L. (J.)

'The sea Onyone or purginge onyone plentifull in England uppon the sea coastes.

It ys also knowen, and to be bought at the Apothecaries only, by the names of Scylla, squylla or sea onyone: very muche used in phisicke. The syrupe thereof purgeth very well clammye, rawe, flegme, yet not without suspicione of some daunger and paynes. And therefore yt ys the better used in S^r [? summer] tyme, when all phisick muste be conveyed into our bodyes without offense of taste, and worcke without sense of the least grypinge, or troble that may, and muste be.'

169. [Error for 153.]

Ascalonites antiquorum.

Shallot. *Allium ascalonicum* L.

'Ascalyones muche used by the poore husbandemen and welsh men which love leeks wonderfull well. Theis rude people which be acquaynted with thys harde hotte foode doe fynde noe inconvenience therein, and the opinion of the very learned alloweth the same for them to be right good and holesome that the same also ys to them meate and medicyne, because yt norrisheth and so pryserueth them against all infectiones of the hotte tyme of the somer when they doe moste use yt with cheese. To those which seldome taaste this hotter herbes they are concluded to bee perillouse.'

¹ John Symings, M.D. of Oxford, F.R.C.P. 1555; President of Coll. of Physicians, 1569 and 1572; died at his house in Little St. Bartholomew's Smithfield, 1588. (Munk, *Roll R. C. P.*)

154. Shoeno prasson. Chive. *Allium Schoenoprasum* L.
 'I tak theis to be set for ye Cyues.'
 Porrum vulgare. Leek. *Allium Porrum* L.
 'The vulgar Leëke (Sett).'
155. Porrum tonsile. *Allium Porrum* var.
 'The unset Leeke which they cutt to the potte.'
172. [Error for 156.]
 Allium sylvestre tenuifolium. Crow garlick. *Allium vineale* L.
 'Wylde garlyck.'
158. Allium. *Allium sativum* L.
 'Garlyck.'
159. Allium ursinum latifolium. Ramsons. *Allium ursinum* L.
 'I thinck thys to be set for our Ramsynes, whereof there
 groweth great plentye about fourde in Kent in Wrotham parishe,
 and ys esteameyd very good against the stoane.'
228. Botrys. *Chenopodium Botrys* L.
 'Oake of Jerusalem.
 It heateth attenuateth diuideth or cutteth, openeth and purgeth.
 Matthiolus.
 It pryuayleth against all flegmatycke, mattered or putrifed
 affectiones of the breast, and lunges; also yt healpeth such as
 consume, be trobled with shorte breathinge, and stuffinge or
 makinge noyse in the breathinge pipes; as well the herbe in the
 decoctione of liquirishe druncke as the decoctione of the herbe
 onlye, many dayes taken with violate or Rosate honney.
 Matthiolus lib. 3, cap 119, fol. 852.
 To suche as spyte mattered, humores, putrifed, yt
 marvelously pryuayleth which I my selfe can trewlye testifie.
 The same author in the same place.'
229. Cichorium sativum. coeruleum. *Cichorium Intybus* L.
 'Succorye well knowne.
 'Amonge the residewe of suche as by my very long experience,
 and certen credytt and fidelitie, have been prooued: I commend
 the infusion of Rhewbarbe in Endyue water or Succorye water
 against anye obstructiones or lingeringe agewes. For I neuer
 sawe Agewe (by obstructione) not cured with this Remedie
 if any will continewe the use thereof. Because euen the most
 thicke, clammye, cleaninge and stuffed, choakinge humores,
 obstructiones also, which by weaknes of naturell heats coulde not
 be eased or removed: by the takinge and use of Rhewbarbe
 I have seene cured. I have accustomed therefore to take a pynte
 of Endyue water, wherein I infuse or steepe the weight of fyfteene
 pence of fynelie slyced Rhewbarbe, tyed in a thynne woven linnen
 clothe. After every daye of the same infusione (ye Rhewbarbe
 still beinge pressed or wringe into the same) fower ownces (which
 ys about a dosen sponesfull) I geve in charge to be geven in ye
 mornynge, and this ys the quantitie for childeren. Neither doe
 I discontynewe the same, untill I see the obstructiones and Agewe
 whollye gon and taken awaye. For without all doubt all wil be
 safe, quickly removed, and health regayned yf all thinges be
 herein well don, and convenient to the Cure. Montanus, libro
 de Componendis medicamentis, Consilio pro puero ,
 consilio quinto, fol. 105.'

352. *Isatis sylvestris Vaccaria dicta.* *Saponaria Vaccaria L.*
 'I sawe yt grow in the garden of the very honorable the Lord Abergavenny, his garden in Kent a^o. 1584.'
 [A casual introduction from Continent.]
374. *Aloe. Sempervivum marinum. Aloë succotrina Lam. or A. Perryi Baker.*
 'Aloë or aloë succotrina best knowen by that name. To be bought at th'apothecaries only: a singuler, good and very safe purger. W. & L.
 Amonge all other medicines the use of Aloe marvelous well pleaseth me. Noe man nedeth to feare the heat thereof, euen in the Somer tyme. Although the vulger phisiciones speake their pleasure. Baptista Montanus, Veronensis, *Consultat.* xci. De preservat. a calculo. The weight thereof is 3^{ds} before supper halfe one hower or lesse.'
396. *Perfoliatum vulgatus, flore luteo, fo. umbilicato.*
 Hare's Ear. *Bupleurum rotundifolium L.*
 'Thorowe waxe or Perfoliata. Is unto chylderen broken use to give the seed hereof in mylke meates, also the destilled water thereof with good success. Tragus, fol. 484.'
438. *Coronopus repens Ruellij & Cornu Cervi alterum vulgi.*
 Wart-cress. *Coronopus Ruellii All.*
 'The Englyshe in moste places call this Swynes Cresses and not Harteshornes in aneye place to my knowledge.'
459. *Alsine, sive Hippia major.* *Stellaria aquatica Scop. (J.)*
 'The great Chyckweede.'
 'Of Chyckweede thus writeth Lonicerus, fol^o. 168. *S. media L.*
 The distilled water of Chyckweede in virtew ys equall with Purselane; wyth wine or simple alone wythe good successe yt ys geven to those which pyn, and waste with longe sycknesse. Unto chylderen in immoderate and unnaturall heate yt ys good to give because yt coaleth the inwarde heate and mitigateth or quyte takethe awaye those terrible accidentes whiche chylderen have by suche extreame heates as Crampes, palsies, tortures and schreamynges, schrychynges, crynges, startlinges, bowynges and sudden bendinges sometymes forwardes sometymes backwarde and suche like accidentes which chylderen be subject unto. Lonicerus, fol^o. 168^o.
 The decoction of Chyckweede or y^e Decoctione of the rootes of great grasse or of purslane, or ye rootes of Male fearnie with spotted leaves aney one alone sodden in water with a lytle whyte wyne or male or possit ale ys excellent in my proofs against ye agew and worm in chylderen. Fiorananta lib^o. (Capricei medicinali).'
474. *Nummularia sive Centimorbia.* *Lysimachia Nummularia L.*
 'Herbe twopence (also Woundeworte).
 'In water wth suger, yt ys geuen againste the exulceratione of the Breast and Lunges, yt helpeth the coughe and those which hardely breathe. Chylderen which hardely receaue medicines trobled with a drye coughe are cured herebye. Lonicerus, fol. 208. Tragus, also, fol^o. 808.'
529. *Crista Galli Herbariorum.*
 Yellow Rattle. *Rhinanthus Crista-galli L.*
 'Ratle grasse in meddowes very vulger.'

572. *Digitalis purpurea*. *Digitalis purpurea* L.
 'Foxegloues.'
 'Hereof Lonicerus writeth fol. 74 that yt doth attenuate, clense, purge, loase, cut flegme or grosse humores: and all virtewes and qualities which Gentiane hathe, yt also hathe.'
589. Tussilago, Farfara. Coltsfoot. *Tussilago Farfara* L.
 'Foole foote, horse houe (loote leafe, the father before the sonne, coulte foote).
 It may also be called Coughe worte. It groweth moste in wheate lande, and fallowed feildes. This shape aptlie agreeth with the herbe when yt freshe springeth in March and Aprill.
 Tragus colored hathe the herbe moste trewlie shewed, muche better then this, the one syde hoarlye whyte next the grownde, and the upper syde freshe grene; the leaves then theis more rownde. The roote medicinal against the coughe; and imperfectiones of the lunges.'
594. *Nymphaea lutea*. *Nymphaea lutea* L.
 'The Yellowe Nymphye or water lyllye.
 The roote or seede of ether of theis Nymphies sodden in redde wyne and drunck (noe remedie otherwyse healpinge) stayth ye immoderate courses of women. Lonicerus, fol^o. 177.'
622. *Vitis alba*, Bryonia. White Bryony. *Bryonia dioica* L.
 'Herbe bryane, hedge vyne, agew roote, tetter burye roote, and Bryonie. The great whyte roote.'
625. *Vitis*, vel *Bryonia siluestris*. Black Bryony. *Tamus communis* L.
 'Blacke Bryonie, wylde vyne.'
625. Peruuiana Mechoaca Mechoacae Prouinciaie planta Bryoniae similis. *Ipomaea Jalapa* L.
 'The Mechoane or Mecoacane: from the Indianes muche used in Englande, a^o 1580, untill muche hurte enseeded ye boule indiscrete practize thereof; beinge a simple not without great daunger. Hereof Garzias ab Horto, and Monardus the Spaniarde, also Clusius have wryten. Soe maney other have don, but none doe warraunt yt safe though they commende yt in some cases, beinge prescribed with the advyce of the learner, and those which very well knowe its virtewes and qualities thereof.
 The best ys brought from ye Citie Mexicho: they have 2 sortes, the one lesse daungerous then the other. In my opinione theis 2 doe resemble our 2 Bryonies, and of my mynde, I fynde the moste learned of my tyme in Englande, also diuerse out of Germanye, and Fraunce icompe with us, the difference of Soyle only cause the varietie of effectes in operatione.
 The weight of 7^l or a french crowne in houlder (?), drunck in twelve sponisfull of seck, will aboundantly purge bothe wayes in common but in moste by stoale onlye. The rootes only ys used, halfe soe muche of our Bryonie wyl worck wonders if it be prepared specially, or simple of it selfe.'
693. Argentina, Potentilla. Silver weed. *Potentilla Anserina* L.
 'Wylde Tansie. Anserina Tragi, fol^o. 480.
 The herb sodden in wyne redd or Whyte and drunck healpeth those which have paine in their backes and torments there: also suche women as be troubled with their whytes immoderately this

herbe euen so used cureth : for it byndeth and strengtheneth as Pimpinella Italica. Tragus, fol^o. 481.

The distilled water thereof is very good to cure redd eyes. Idem ibidem.

Against the Dysenterie and lienterie, which is the bloodie flux and passinge of foode by stol undigested and against all fluxes and fluxes this herbe is used in our adge. Idem, fol^o. 480.

Tragus estemeth it drye because it bindeth there.'

748. Pedicularis. L. 431.

Marsh Red-rattle. Lousewort. *Pedicularis palustris* L.

'Lowzye weede because the catle feadynge thereof will become lowzye: thereof very muche groweth in black brookes in Estmallinge in Kent.'

757. Capnos, Fumaria.

Fumitory. *Fumaria officinalis* L.

'Fumytarie; the distilled water thereof wyth Tryacle before the purgatione certaine dayes drunck ys very holesome for suche as labor of the Frenche euyll. Because yt purgeth the infected bloode. Lonicerus, fol^o. 167. The same ys good against the Plague and the desease called the Englishe sweate; also good when aney shall purpose to Bathe and sweate. The same author there.

The Decoctione thereof with Fennyll openeth the obstructions of the Lyuer and forceth oute the Jaundyce by uryne. The Juice thereof and the roote of Esula, eche one drachme which ys in weight 7^d mixed and drunck with hott water dryueth ye Dropsie and . . . (?). In same author, fol^o. 167.'

807. Asplenium. Scolopendria.

Ceterach officinarum Willd.

'Splene worte. Yt groweth uppon the southwest ende of Est Peckham church in Kent: also uppon ye Pallace walles in Maydston, from where I did transerre y^t to my garden walle, where it groweth. a^o. 83.'

807. Lunaria racemosa. L. 470. Moonwort. *Botrychium Lunaria* Sw.

'I haue sene thys lunarie or Moneworte growe in black heathe; also nere Saynt Margaretes, nere Rochester.'

810. Adiantum album et nigri Plinii. A. 361.

Wall Rue. *Asplenium Ruta-muraria* L.

'Thus wryteth Matthiolus of Ruta muraria in the chapter of Paronichia.

Moreover this moueth uryne and grauell.'

Vol. II.

180. Rhamnus primus Diosc. creditus. L. 598.

Evidently intended for Sea-Buckthorn. *Hippophae Rhamnoides* L.

'This Rhamnus I founde betweene Douer and Foulkestone by the sea syde under the Clyffes, a^o. 1582, with reddish berries Orenge colored.'

280. Cerefolium descriptum.

Chervil. *Anthriscus Cerefolium* L.

'Cheruyale. It is used with meate as Perslye of a meane temperature betweene hott and coulde; But principally to force out clotted or brused bloode "as it is called" the same herbe to be of wonderfull efficacie certen sure experience hath approued. Tragus, fol^o. 472. The Juice of the herbe, the distilled water also, dissolueth the congealed blood of contusion, or strype, and is of force against the stoane in the kyndnes. Idem ibidem.'

viii. RICHARD SHANNE, 1561-1627.

In the Goodyer MS. there is a note of the name of Richard Shanne, at the head of a paper upon which are recorded two localities for north country plants, but there is no further indication as to who he might be.

Pyrola groweth in shadowed woods in Craven, in a place called Craggie Close in Lanscale.

Monophyllon groweth in Lancasheir in Dingley wood and in Harwood neare to Blackburne.

Both localities are given in Gerard.

During a visit to the British Museum in June last, a happy chance made me acquainted with 'The Shanne Family Book' (Addit. MS. 38599. 17), from which I discovered that Richard Shanne of Woodrowe was a considerable horticulturist, who was living near Methley¹ in Yorkshire in Goodyer's time, and who certainly deserves to be better remembered. I have no doubt but that he is the authority whose name is quoted by Goodyer. It is not unlikely that Goodyer may have got into correspondence with him through Walter Stonehouse, the friend of John Savile of Methley, whose neighbour Shanne was.

Richard Shanne sonne and heire of William was borne the tenth of Auguste beinge Thursdaie 1561 he married Amy Burton daughter of Richard Burton alias Carver the ixth daie of June 1588 and had issue by hir, Thomas. He married his second wife Marie Chamber . . .

This Richard was of reasonable tallness stright of bodie, he was somewhat pale of complexion, his heire of his head mouse colored, he was verie light and nimble of foote, his chefest delite was in plantinge and grafting all maner of herbes & trees, and had growinge in his gardinge a great number of rare and straunge plants, there was not allmost anie herbe growinge but he did knowe the severall names therof, and the nature and operation of the same, he did practise both in phisicke and specially in Chirurgerie and did cure verie manie daungerous wounds and ulcers. He made three bookes of the Nature and operations of herbes and Trees and drew with his pen the trew picktures of everie plante, set downe in what ground everie herbe and tree was to be found and the tymes of their springinge, florishinge and sedinge. He planted three Orchards of his owne, the first at the Mickle towne which he sett in anno domine 1577. The Springe of Aspe trees he planted 1596. The Orchard at the East more syde he planted in divers yeares, first in 1607 & 1613 (?). The

¹ Methley gardens should be famous in the history of horticulture. For at Methley lived John Savile, the friend of Walter Stonehouse; there also in Mr. Witham's garden, gold-streaked Pansies growing spontaneously 'mightily beautifie the border of an hedge'; in the Wood-close flourished the Blush coloured Bugle (*Ajuga reptans* L.), and at no great distance grew *Lunaria minor* in John Nun's cow-pasture. Witham's son contributed localities of some Oxfordshire plants. Merrett, *P'inax*, pp. 17, 65, 74.

Springe of Elme trees by the Moorsyd was planted in 1613. The litle Orchard in the end of the More house field he compassed about with quicksetts in anno 1616, and the trees was set in anno 1617. He maid two large bookes diologue wyse of Phisicke & Chirurgerie, He delited much in reding Granados meditations, and was verie seldome seene in anie rude companie, but avoyded companie as much as he could and took much pleasure to walke in woods and to be solitarie. He lyved in the daies of Quene Elizabeth, Kynge James & Kinge Charles when he was fiftie & nine yeares oulde he mayd a large booke of prayers & meditations which he did drawe out of sondrie learned authors. The meditations are of all the miseries of man from his verie birth unto the daie of his death. Allso of the torments which the wicked do suffer in hell . . .

He died of consuming consumption at the age of six and three score wanting one moneth. [ff. 83 v, 84.]

[His garden list is printed on p. 310.]

A most interesting extract from Shanne's diary was made by Antony Wood when he consulted it in October 1674, by permission of the then owner. It contains a most lifelike description of the Ruff (*Machetes pugnax* L.), a bird that was not described by Turner in his book on Birds, so that Shanne's is probably the first English description.

Anno 1588, there was taken at Crowley in Lincolnshire in the winter time 5 strange fowles of divers colours, having about their necks as it were great monstrous ruffs, and had underneath those ruffs certaine quills to beare up the same, in such a manner as our gallant dames have now of wier to beare up their ruffs (which they call *supporters*). About their heads they had feathers so curiously set together and frised, altogether like unto our nice gentlewomen who do curle and frisle their haire about their heads. Three of these strang fowles was brought into Sir Henrie Leese, and they would walk up and doune the hall as if they were great states, and sometimes they would stand still and lay their heads together as if they were in a secret counsell. It made the beholders to wonder therat. They cast them come to eat, but they refused to tast of any meat and so at length died. Mr. Richard Shann, of Wodrow in Medley, Yorks., drew a picture of one of them which he placed in his herball. Two men that had set lime twigs to catch birds withall did find them taken therein. The like never seen or heard of before.

ix. JOHN PARKINSON, 1567-1650.

Several pages of notes in the Goodyer collection on American, Bermudan, and Oriental plants are written in a hand that was at first unknown to me, but which I have since been able to identify beyond doubt as that of the well-known herbalist John Parkinson. Certain of these notes are written on the back of a letter, every word of which was so completely scribbled over as to be quite unreadable. By slow and careful erasure with a sharp knife I found it possible to remove sufficient of the ink-scrawling from the

surface to bring into view the following letter, signed by him. It is a unique possession, as being the only signed specimen of his hand-writing known.



PARKINSON.

To the worthy Gentlewoman
Mr^{es} Geeres
geve these.

Seal



GOOD MRS. GEERES, I hav by this messenger sent you the ij trees wherof I tould when we were last together at yor howse. I brought them thither on the Saterdag you were so earnest with me to come to dinner where I thought to have met with you, but missing you I carried them home again & laid them in the ground wherin they have been safe without taking any harme. I do also think that then you would have pleasure, and with that you promised & have long expected it. I pray you doe not weaye me with deseyre which is worse then denyall as you please appoint Mr. Codemer to doe it speedily. So shall ever remaine,

Yors JOHN PARKINSON.

[MS. f. 168 v.]

A comparison of the text of several of the notes with passages printed in the *Theatrum* in 1640 confirmed the view that they are indeed the author's own notes, and were probably the actual notes used by him in the preparation of his book. Some of them are

printed with the lists of exotic plants on p. 363. The following horticultural notes were evidently extracted by Parkinson from Francis Bacon's *Naturall Historie*, Century v, 1627; but another possibility should not be lost sight of, namely that Bacon may have derived part of his horticultural knowledge from the distinguished botanical writer who two years later dedicated his *Paradisus in Sole* to Queen Henrietta Maria.

Notes on Horticulture by Parkinson.

The steepings of Wheate seede in horse dung, cowe dung, pigeon dung, urine, chalke, bay salt, claret wine, soote, ashes, malmesey, & spirite of wine: the urine, soote, ashes, chalk & salt shooting within six dayes, the best & lustiest of them was first from the urine, then dung, next chalke, then soote & then ashes, & last the salt therewith wines not at all except the claret wine. [Bacon, *Sylv. sylv.* § 402.]

The drawing of boughes of a tree or vine to the inside of a roome where fyre is continually kept doth accelerate the fruite a moneth sooner. [§ 405.]

The removing of living plantes into new freshe & loose earth once in a yeare doth accelerate & enlarge them. [§ 406.]

The grafting of Roses in Maye will cause them to beare flowers the same yeare but late. [§ 418.] The binding also of the bark worketh the like effect. [§ 419.] Grafting upon contrary stocks will never thrive long, as Peaches on Cheryes etc., the cause is the cyon overruleth the stock quite, the stock is but passive onely & giveth aliment but no motion to the graft. [§ 421.]

The laying of a heape of flinte or other stones about the roote of a wild tree doth make it prosper twice as much as without them, because the stones retain moisture longer & not to be consumed so soone, it keepeth also the tree from cold blastes & frosts & geveth more warmth at all tymes. [§ 422.]

To boare a hole in a tree that beareth not is usually done to cause it bearing; too much repletion may be the cause opprest with his sap. [§ 428.] As also to cleave 2 or 3 of the cheife rootes & to put a peble into eache to keepe them open for els they will close againe. [§ 429.]

To draw the bough of a tree through a wall to the south sun hath been practised by some to ripen the fruite the better, but it sorted not. [§ 431.]

It were good to trye whether a tree grafted lowe & the lower boughes maintained, the upper ones being continually proined off, would not make larger fruite. [§ 432.]

It is expected that trees will growe greater & beare better fruite if you putt salt, lees of wine or blood to the roote which are more forcible. [§ 457.]

It is also delyvered before that if one take the bough of of a lowe fruit tree newly budded & drawe it gently without hurting it into an earthen pot perforate at the bottome & set in the slant & then cover the pot with earth, it will yield a very large fruite within the ground, the like will be effected by an empty pot without earth in it, put over a fruit propped up with a stake as it hangeth on the tree, the pot being perforate to let in aire. [§ 470.] [MS. 11, f. 168]

On the same paper are several entertaining notes in Parkinson's handwriting relating to Brazilian and Peruvian plants and their 'vertues'.

The Vertues of some Exotic Plants.

Giniber is a tree growing in Brasill about Pernambuco, whose fruite is bigger then those of an Orrenge, but of a darck greene colour from whence with one kind may be crushed out a liquor very like both for colour and consistence unto the whey of milke: this liquor or water hath such a qualitie that what parte of the skin of the bodye be washed therwith it will colour it so black as no black inck or dye can doe more & so likewise the haire in any parte of the bodye, which colour will abide so strongly fixed therin for the space of eight dayes that if all the remedies to whiten the partes againe were applied they would be of none efect; but of itselpe the colour will fade after that tyme & the partes will become as white as they were before. This water is also as it issued is of espetiall propertye to encrease haire. Other things this water doth performe without the least daunger or harne.

The Brasilians have a familiar medicine that is common & well knowen to most of them, beinge a most certaine remedye to staye or stanche the blood that flowes from any part of man or woman.

This is made of a certaine herbe which groweth in the high & rockye mountaines which therupon the Portugalls call Rais de serra that is Radix montium, the Roote of the Mountains, because the roote onely of this herbe hath that effect to staunche blood, the leaves herof are said to be like unto Plantaine leaves & the roote unto a Cicharye roote both for forme and colour which roote being a little burned in a cleane earthen pot or vessell & afterwarde rubbed into partes & a scruple in weight taken fasting with Plantaine or Rosmary doth most certainly staye the fluxe of blood issuing from any parte of the bodye. Petrus de Osma cited by Monardes testifyeth the vertue that many herbes have that hidden propertye to stay bleeding by the example of some Negros that constrayned through hunger to cut of the calves of their legs to eate, & by layinge a leafe of a certaine greene herbe did whollye staye the fluxe that not a droppe of blood was seene to be shed. [Cf. Park. *Theatr.* 1622.]

There is a certaine kinde of Rushe growing on the hilles in Peru, which they there call Jeho or Yoha, very like onto the Spanish rushe called Spartum wherof the fraile are made that they putt Raysins & figs; & wherwith the metalline bodyes are sooner melted by its flame than by much wood, & separated from the Quicksilver that is put into them for that purpose. The fumes also of this Rushe (a wonder to be spoken) taken under a Canopay or close covering, causeth all the Quicksilver, that abideth in the bodies of those that have ben annointed therewith to helpe the Frenche poxe, to sweate out of all partes of the bodyes, without any maner of sense of paine.

Lakeka is a kind of gum gathered in Martaban in the East Indies, whose propertye is to expell drunckeness if pieces therof being put on a thred or stringe & wounde on the bare arme, so that whosoever shall drinck much & strong wine shall not be overtaken therwith.

Lapis Lipis is a kinde of blew minerall stone found on the hills of Potosi called Lipis of the cite neere them. And is of a transparent blew colour, very hard to breake yet brought into small picces, sharpe and bitinge in taste so that being put to the tongue it will with the heate oxalegrate it, it is brought as merchandise being cut into tablets & so sold: it is (saith Zacutus Lusitanus) so like Anil Indica that a form of it might be taken for Indico & being made

into pouter & put . . . water will colour the water blew within an hour. All fowle ulcers eyther of the mouth or privy partes of man or woman beinge washed will presently cleanse them & take away or restrain their virulencye. 4 gr. in p. to ζij aq^a rosa^p hora.

Coque . . . is a certaine tree well knownen to all in Brasill growing plentifully in the woods about Pernambuco very like unto the Lentisk or Mastick tree whose barck being boiled after a pound thereof hath ben cut small in 16 poundes or pintes of water to the halfe, adding therto ζiij of allom in the boiling, we^h when it is strained & settled will become of so pure a purple colour as if it were made of *ostrum* or the purple fishe, or of the colour of the red Feild Poppye. The face washed with this water beside^s that it coloureth the black spotts in the face, it geveth to the face & lips so beautifull a Rose colour that it would be a brave fucus for women: but this colour will not abide above eight dayes although before that tyme it will not be blemished by many washings, but after that tyme the colour fading it will change into a dark Rose colour and after 4 dayes more will be quite consumed; yet those black spots that were in the face before will no more appear, but by using this water againe upon the face etc the fresh colour will be apparant againe: If this barck were brought us it would redound much to the good of women, who to beautifye them selves with Arsenicum sublimatum, Cerussa, Camphire & other such thinges doe exchange their good for wrinkles, fowle ulcers in their gums, blacknes of teeth, rottennes and stincking of the breath.

The Cabbage Tree groweth to an incredible height neere unto 200 foote bare without branches unto the very toppe as most of the kindes of Palme-trees doe, where among the long leaves groweth this round heade or cabbage, to gett which they usually cut downe the tree at the roote. The stemme or bodye therof having not much above 2 inches thickness of wood, the rest being of a pithy substance. From this tree likewise they drawe wine as from the Coco & other Palmito trees by boring a hole & applying a tappe or other convenient thing with a gourd or the like to receive the liquor that droppeth therfrom, & in 12 howres will by dropes fill the vessel & so will it doe from tyme to tyme by boring new holes one after another beginning above & so descending: what fruite this tree doth beare, our men that have had the benefit herof were never so wise or industrious to observe, feeding like swine on the mast but never looking higher.

The water distilled in glasse vessels from Cloves while they are greene, besides the excellent smell they yeld, it hath ben found by good experience not onely to expell windye humours in the bodye, but that disease also called Priapismus and that effectually. The like is thought may worke the distilled water of the fresh flowers taken from the Cinamon trees or from the fresh Cinamon itselfe.

The fruite or nutt called Cola growing in Guiney & like a chesnutt hath ben often used and found very effectual for chapt lipps, and the descomities of the skin, the itching of womans partes, the raggednes of the nailes, the rednes of face, much Dickwek often castings, the falling of the haire, . . . and many other diseases that proceed from the heate and distemper of the liver, as also against fevers & burning agues, very profitable drincking some endive water after it. The tree is like to the Chesnut tree & so is the nut both for forme & greatene^s but of a pale reddish colour on the outside & a little bitter in taste.

Amivil with the Persians is a tree like to the Chesnut tree whose rootes the deaper the better being tied about the neck or worne on the arme so that they touch the flesh doe induce a mightye hatred for wine espetially to those that are geven to love it much & be often drunck therewith. Gesner reciteth 2 espetiall thinges available for that purpose, viz. a greane frog that is found often in the springes of water put alive into wine & there suffocated. And an Eclousus formed also in wines: this hath ben often urged so before, especially if 2 ounces of the blood of Cates (?) be put into 3 measures of wine. Opium also is thought to performe the like cure & so be the more prone & strong to venery.

[Written by John Parkinson on the back of his letter to Mrs. Geeres. MS. f. 168 v.]

Among his other writings are Lists of Foreign plants, see pp. 358-70; a List of 116 plants, including many bulbs (MS. 11, f. 164), grouped under numbers '36 to 69', evidently referring to plates in the *Anthologia magna*, 1626; a list of 48 plants (MS. 11, f. 157 v.) described by Clusius in his *Appendix altera ad Rariorum Plantarum Historiam*, issued with the *Exotica* in 1605; two lists of Evergreens,¹ one headed *Arbores sempervirentes fol.* 44 (MS. 11, f. 156), the other headed *Perpetua coma virentes variis provinciis* (f. 158 v.); and a brief note on plants used for tanning skins by Mediterranean peoples.

Parkinson was appointed Apothecary to King James and also King's Herbarist. He had a garden in Long Acre, where Goodyer no doubt saw the many rare plants of which he has preserved a list. The following account is in a hand that appears to me to be Parkinson's: if so, it may refer to an accident to the wall of his garden in Long Acre in 1636. A search for the workmen's names in the local parish register might settle the question.

The charge of my outer wall blown downe the 4th Novemb^r 1636.

Mr	iiij ^s	
Tho	.	For	$\frac{1}{2}$ daye	$\frac{1}{2}$.	.	.		xij ^d
John	.	.	2 dayes	iiij ^s	
W ^m West	.	.	$2\frac{1}{2}$	v ^s	
Christopher	.	.	3 dayes	vi ^s	
Howse	.	.	$\frac{1}{2}$ daye		xij ^d
The boye	.	.	$3\frac{1}{2}$	v ^s	ij ^d
Labourers :									
Randel	.	.	3 dayes	v ^s	x ^d
Anthonye	.	.	a daye	$\frac{1}{2}$.	.	.		xxij ^d
John	.	.	6 dayes	vij ^s	
2 other labourers		xvij ^d

¹ It may be noted that at a somewhat later period Sir Richard Browne, writing from Paris to Sir Edward Nicholas on 5 July 1658, states that he was then at work on a *Catalogue of evergreenes*, but had lost the help of a Mr. Keipe, who had left for England. He adds 'Alaternes beare a graine like that of privet, which beinge sowed comes upp and prospers without difficulty'.

[*Camden Society*, xxxi, p. 65, 1920.]

eventually formed the basis of that department of the very curious museum formed by Thoresby in his house at Leeds (see Hunter, *South Yorkshire*).

In 1639 Thomas Johnson organized an expedition of the 'Socii Itinerantes' of the Pharmaceutical Society of London to the mountains of North Wales: an account of the expedition is given in his *Mercurii Botanici pars altera* (1641) reprinted in facsimile in *Opuscula omnia botanica Thomae Johnsoni* edited by T. S. Ralph (London, 1647). The constitution of this travelling club is thus stated by Johnson in the preface to his *Iter Plantarum Investigationis* 'susceptum a decem Sociis in Agrum Cantianum: Anno Dom. 1629', and published in the same year: 'Paucis abhinc elapsis annis, consuetudo vero laudibilis inter rei herbariae studiosus crevit, bis aut saepius, quotannis triduum aut quadriduum iter Plantarum investigationis ergo suscipere'. Stonehouse joined the party at Chester, having spent the previous night at Stockport, where he had not been favourably impressed with the inn. Their route took them by Conway, Penmaenmawr, Bangor, and Carnarvon to Glynn-lhivona, where they were the guests of Thomas Glynn, to whom Johnson dedicated his account of the expedition. After discoursing on the perils of climbing Snowdon, Johnson gives a list of the plants found by the party. At Beaumaris they enjoyed the hospitality of Richard Buckley, visited his vivarium, and collected seaweeds. They then recrossed the straits to Lhan-lhechid, climbed Carnedh-lhewellyn in a mist and in fear of nesting eagles, but saw little of botanical interest. After a farewell visit to Glynn-lhivona, the party journeyed to Harlech and Barmouth. Their homeward journey lay through Merionethshire; at Guerndee Stonehouse left them and went home through Shropshire to Darfield. Here he remained for a time in quiet enjoyment of his garden, to the Catalogue of which, drawn up in 1640, reference has already been made; some of the plants in Johnson's list are included in the Catalogue, and were probably obtained on the Welsh expedition.

About 1648 we learn from Walker's *Sufferings of the Clergy* that Stonehouse was forcibly ejected from his living by the Parliamentary Commissioners and imprisoned. On his return, probably in 1652, his spirit as a horticulturist seems to have been broken, for he then wrote in the Catalogue a pathetic note in Latin, to the effect that but a few of his plants had survived—*Novamque despero coloniam*,—'I have no hope of a new colony'. After this he would appear to have lived in London, to have made or renewed acquaintance with the younger Tradescant, and to have written some intro-

ductory verses to the Catalogue of Tradescant's *Museum*, published in 1656 :

To *John Tradescant* the younger, surviving.

Anagr. :

JOHN TRADESCANT.

Cannot hide Arts.

Heire of thy Fathers goods, and his good parts,
Which both preservest, and augment'st his store,
Tracing th' ingenuous steps he trod before :
Proceed as thou begin'st, and win those hearts,
With gentle curt'sie, which admir'd his Arts,
Whilst thou conceal'st thine own, and do'st deplore
Thy want, compar'd with his, thou shew'st them more
Modesty clouds not worth ; but hate diverts,
And shames base envy, ARTS he CANNOT HIDE
That has them. Light through every chink is spy'd.

*Nugas has ego, pessimus Poëta
Plantarum tamen, optimique amici
Nusquam pessimus aestimator, egi.*

GUALTERUS STONEHOUSUS.

Theologus servus natus.

By re-arranging the letters of John Tradescant's name he composed the anagram *Cannot hide Arts*, and by a similar process his own name, Gualterus Stonehousus, became *Theologus servus natus*—words quoted by Macray, who, however, did not grasp their meaning, as occurring on the title-page of a volume of his Sermons in Magdalen College Library.

These verses, his only printed work, were not published until 1656, the year after his death (probably in London) at the age of 58.

xi. THOMAS JOHNSON, c. 1600–1644.

The outlines of the life of this distinguished botanist are well known. Trained as an apothecary, with a business on Snow Hill, he attained to be the best herbalist of his age in England and a trusted physician. In 1643 he had the rank of Lieutenant-Colonel to Sir Marmaduke Rawdon. During the siege of Basing House he led several sorties with success. 'When a dangerous piece of service was to be done, this doctor, who publicly pretended not to valour, undertook and performed it.¹ On one occasion he had a hand to hand struggle with Capt. Clinson, but at last in the affair of 14 Sept., he was shot in the shoulder, whereby contracting a fever he died a fortnight after, his worth challenging funerall tears,

¹ Fuller, *Worthies*, 1662, p. 204.

being no less eminent in the garrison for his valour and conduct as a soldier than famous through the kingdom for his excellency as an herbarist and physician.'

Johnson had a wide circle of friends and helpers in addition to the *socii itinerantes* who took part in his botanical excursions. Among others he mentions GEORGE BOWLES of Chisellhurst in Kent; SIR JOHN TUNSTAL, gentleman usher to the Queen, owner of a garden at Edgcomb, Surrey; HUGH MORGAN, apothecary to the Queen; ROBERT ABBOT of Hatfield, a learned preacher; JOHN REDMAN, 'a skilful herbarist' of the north of England. The manner in which his friends used to make use of his services is shown by a letter that is still extant from the Provost of Eton, SIR HENRY WOTTON.¹

To my very loving and learned friend Mr. Johnson, apothecary, at his house on Snowe Hill, London.

2nd of July 1637.

MY GOOD FRIEND MR. JOHNSON,

I have addressed this my servant unto you at the present with two or three requests. First, that you would direct him where he may buy one of your Gerrards, well and strongly bound; next, where I may have for my money all kinds of coloured pinks to set in a quarter of my garden, or any such flowers as perfume the air. Thirdly, I pray let me consult you whether you know any sick of that fastidious infirmity, which they call *melancholia hypochondriaca*, wherewith I have been troubled of late, but more with a symptom very frequent in that passion (as the great Fernelius describes it) . . .

HENRY WOTTON.

Johnson's early death was a great blow to Goodyer. In his will (P. C. C. 114 Twisse) dated 11 May 1635 he described himself as Citizen and Barber surgeon of London. He left legacies to his aunt, Mary Cave, 5s., to William Parker 40s., to Eliz. Parker 50s., to Richard Parker, son of Wm. P., £5 to set him forth to prentice, and to his wife Alice Johnson, his sole exor., the residue of his estate. The will, witnessed by W. Parker and Mary Vudell, was proved in July 1647.

¹ SIR HENRY WOTTON (1568-1639) of New College, Queen's, and Eton, diplomatist and poet, is best remembered by his exquisite poem *The Character of a Happy Life*. His letters allude to contributions to English horticulture during the second tenure of his office as Ambassador at Venice. They mention:

'Seeds and roots and slips of rare flowers and plants' to Sir R. Winwood. Oct. 1616.

'The most excellent choice of those seeds which his Majesty desireth' to James I. Dec. 1621.

'More melon seeds of all sorts' to James I. Dec. 1622.

'Finocchio' or Italian Fennel (with full directions for cooking and eating) to Tradescant. Probably about the same time, but not recorded until 1656. Parkinson, *Paradisus*.

To James he wrote that he intended to examine some of the best *hortolani* of Chioggia and other places about the manner of cultivation. He was in close correspondence with Lord Zouch, and may have also sent plants to the Hackney garden. Pearsall Smith, *Wotton*.

The intimate relations between Johnson and Goodyer at the time when the former was preparing his second edition of Gerard's *Herbal*¹ have often been alluded to. The Goodyer MSS. throw further light on their intercourse by providing us with a detailed list of the plant-descriptions which Goodyer lent to Johnson and subsequently received back again with the printers' marks upon them.

The copy of Johnson's *Iter Plantarum in agrum Cantianum*, 1629, in the Magdalen Library, appears to have been a presentation copy to Goodyer. It is of historical importance as being the first English local catalogue, and is marked with the following corrections by the author (?).

p. 3, l. 30	for	festinante	read	festinantibus.
32	„	Glycyrhiza	„	Glycyrrhiza.
4, 23	„	audeo	„	ausim.
4, 23	„	possum	„	possim.
6, 5	„	est	„	sit.
6, 11	„	investigemur	„	investigemus
7, 29	„	possumus	„	possimus.
9, 4	„	potuimus	„	potuerimus.
11, 31	After	<i>Androsaemum quorundam</i>	add	<i>Park-leaves.</i>
13, 3	„	sit	„	quod.

At the end Goodyer has added the following list of plants :

Solanum lethale.	Tricomanes seu Capillus Veneris verus.
Helicacabus.	Dryopteris.
Viburnum.	Alleluia.
Antirrhinum minus.	Epicrion.
Mille grana minima.	Alchimilla.
Centaurium fl: albis.	Sanicula.
Lysimachia lutea.	Acer minor.
Alsine serpilli folio.	Betulus.
Osmunda regalis.	Calamintha agre: Belgarum.
Eupatorium Can: mas et femin:	Petasites.
Angelica sylvest:	Gramen Bufonium.
Herba paris.	Cynocrambe.
Eleborine.	Allium ursinum.
Aleca.	Xyris.
Spergula altera flo: purp. minor.	Rhamnus catharticus.
Ascyrum supinum hirsutum.	Rubia flo: carneo angustifolia.
„ palustre rotundifolium.	„ flo: rubello.
Pentaphyllum supinum tormentillae facie.	Chamaepitys.
	Aphaca.

The copy of the 1632 edition of the same work appears to have been the author's own copy with his MS. index, afterwards extended by How and used in the preparation of his *Phytologia* (1650).

¹ A presentation copy of this book to the Apothecaries Company was the first contribution to their Library. With their other books it was probably burnt in the Great Fire of London.

xii. WILLIAM HOW, 1619-1656.

(See p. 251.)

How's botanical reputation rests upon his *Phytologia Britannica natales exhibens indigenarum Stirpium sponte emergentium*, the book in which for the first time all the known plants of Britain were garnered together with their localities. He was a Master of Arts of five years' standing of St. John's College, Oxford, and his list comprising 1,220 plants is a memorable achievement for the time. Apparently owing to his private means being insufficient to permit of much travelling, he had to rely upon information sent by other botanists, with the result, as Ray pointed out, that many exotics and garden-escapes got included in his lists. 'The rare plants were almost wholly communicated by his friends, Mr. Stonehouse, Dr. Bowles, Mr. Heaton, Mr. Loggins, Mr. Goodyer and others, and he drew some from a MS. of Dr. Johnson.'

Goodyer may have become personally acquainted with How at Oxford, and in any case they were in close correspondence both before and after 1650 when How was living in London, either in St. Lawrence Lane or in Milk Street.


The newly discovered writings of How appear to belong to the periods immediately preceding and following the publication of the *Phytologia Britannica*. Though not extensive, they throw valuable light on his relations with contemporary botanists; and if his notes be somewhat slipshod, we must remember that they were not intended for our perusal, and that he had but recently done duty as Captain of a troop of horse in the King's army.

First in time and in importance are his list of plants and notes written at the end of what I believe to have been Johnson's own copy of his *Descriptio Itineris . . . in agrum Cantianum*, A.D. 1632, with MS. index. This may have come into How's possession after the death of Johnson in 1644. He supplemented it with a further list of indigenous plants from the Herbals of Gerard and Parkinson and from other sources; and then no doubt he developed the idea of including all British plants in one comprehensive, alphabetical list. A few rough notes on the last pages almost certainly refer to his programme of work, but as they are mostly crossed out with ink lines, it is impossible to give a complete transcript of them. The meaning of such remarks as 'Review both ye Herballs againe and take all', is clear, and the references to persons may be of historical importance. The names mentioned I believe to be those of T. Johnson, Parkinson, the Rev. Walter Stonehouse then exiled from his Darfield living, Bobart of Oxford, Leonard Buckner and Edward

[Faint, mostly illegible handwritten text at the top of the page, possibly bleed-through from the reverse side.]

[A small handwritten signature or note, possibly "John Parkinson".]

OBLITERATED LETTER BY JOHN PARKINSON



Johnson's Index

How's 'additional plants'

FIRST DRAFT OF A BRITISH FLORA BY HOW

MS. additions to Johnson's Descriptio Itineris in Agrum Cantianum, 1632

Morgan of Westminster. His notes for planning his work, such as 'Remember ye English names to bee last placed', 'Remember to give ye proper name in Latin of ye place where each plant flourisheth', &c., have the appearance of being the advice of some botanical friend of experience.

The following extract gives an idea of his controversial style of writing :

Anchusa Alcibiadion Ἀγχύσα. Fucus herba. Onocleia.

Buglossa Hispanica, Red Alkanet, crescit isola Thanet.

‡ *Ger. em.* : I doubt whether our author found any of these in the place here set downe, for I have sought it but failed of finding; yet if he found any it was only the first described, for I think the other three are strangers.

‡ Our Johnson uses not his usuall charges for heere hee gooes about to confute with a 'for-I-thinke', and seconds it with as good an argument from an Hypothesis (if hee found any, it was onely ye first), and why not as well ye other three, since hee confesses afterwards in both his Catalogues ye Anchusa minor, ye 3rd of these, to bee our countrey plant.

Anchusa lutea. Yellow Alkanet.

[How MS. in Johnson's *Desc. Itin. Cant.* 1632, f. 6 v.]

At the end of the volume are the following memoranda, but so dreadfully scored over as to be almost illegible :

Qu. whether some plants which are given by Johnson in his Itineraries to bee *non descript.*, are not described by Parkinson?

Putt him nondes nere ye additionated plants and ye and see whether they be contained (?) in Mr. Holybyn's Catalogue.

. montanum sylvestre Anglicum ut agagua Cl

Remember ye English names to bee last placed. Consult how many plants wee have added in our treatise by telling ye old, ye remainder, ij ye number.

Mr. Holloway hath Stonhous papers manuscript.

To all Johnsons new plants, put Jonh. MS.

Remember to ye plants to put their titles, as D^r. Bowle, M^r. M, and So every man's name to his plant.

Remember to give ye proper name in Latin of ye place where each plant flourisheth, and to bee carefull to f

Review ye spelling of names of Townes by Cambden's Brittan :

vid. 2 Eriffe.

[1. Wulwich] Kent

[2. Eryth]¹ Oxfords

¹ In Johnson's writing.

Review both ye Herballs againe and take all. Remember to insert againe *Apium sylvestre*. See whether you putt downe ye *Aristolochia*. . . . *Anonis* you must adde to *Anonis* 3 more *Auricula muris repens* . . . *Auricula* see w^{ch} of them growe wild. *Birds' eyes*, see them—none wilde.

Remember to insert Jonhsons (*sic*) Author (?) nere Parkinson's
 Borrowe a Coopers Dictionary. English Catalogue.

Asarina.

Consult ye place amicorum benevolentia.

[4 lines illegible.]

Send to Bobert about ye *Millegrana non descr.* whether to Gr. d. E. . . .

Remember Mr Bucner to send for ye Bupleur and Caucalis. Send to
 Dr. Browne and asist Mr

Abrotanum non antea repertum in descript. Then each plant's place
 of and ye places you have found out, faithfully dealt all ye
 English fruites which are not described, ye pharmacutists about such
 as Bedaguar Fungus sambucus met of ye Parson of St. James.

Ubi sativus inveni vulg. sativus.

Observe ye same letters in our printed peece as there is in ye Welsh Cata-
 logue. Those which have

Send to Stonehouse. Consult Morgan about Orchis. Mr. Crosse.

Reade ye Catalogue Epistle onely . . . all plants yt are to be blotted out
 better be done on ye former Catalogue, because they are worse described
 there. [How MS. at end of Johnson's *Desc. Itin. Cant.* 1632.]

How published his *Phytologia* in 1650: though anonymous it must at once have made his reputation as a botanist, and have brought him into correspondence with other plant-lovers who sent memoranda of plants which he had omitted. The names of these he entered in an interleaved copy of his book which he kept carefully corrected with a view to a revised and enlarged edition. Among those who gave him valuable help were Goodyer, Browne, and Hunnibon,¹ and he for the first time had the advantage of looking through the manuscripts of Lobel and of Dr. Penny. That Goodyer put his knowledge freely at the service of all true workers alike, is proved by the fact that at this time he was also assisting another person, believed to have been Dr. Dale, in the compilation of a very similar Catalogue of British Plants. That the idea of a complete British Flora was very much to the fore in Goodyer's mind is proved by his communications to How (MS. 18), by his abstracts from the *Phytologia* (MS. ff. 33-7), by the Dale (?) - Goodyer *Catalogue of British Plants* (MSS. 8 and 9), and by his Index to the English plants localized in Gerard's *Herbal* (MS. 16).

Further information concerning How's activities in the period immediately following the issue of the *Phytologia* is contained on a small scrap of paper measuring 3 x 4 inches, with his memoranda, characteristically struck out.

¹ 'Dr. How had 2 Apothecaries to help him in composing y^e *Phytologia britannica*.' John Ward's *Diary*. Perhaps Hunnibon was one of them.

My Ld. Hatton's letters with seeds. Morines plants. Dr. Morisons 2 Correspondence. Exper: Phy: Brit: Deliver Anderson's letter. *Qu.* way of Correspondence. Goades buisnesse (?).

Sends Ld. L. at Tibr:

Monday Committee

Kay pro horse, Statim.

Proyne Wild, Anderson Phyt. Str. Ill.

pro Stone,

Ye preparacon of sublimatum Cupum *vide* ye Paper. Seeds sent by Cadell dead since in these partes, which I hope you have. ʒ.

Parthenium

Lilac

All maner of seeds

what plants you think you

Dubble flourd Apple

Black Cherry

Sedum maius arborescens

Jasminum Indicum luteum

Phyllitis multifida foliis crispis

Letter in Milk Street finds mee

Answered both Dr. Morisons letters have

I have all Plant bookes for dressing *Pinax*, what bookes I am about.

[How MS. bound with Goodyer MS. 11, f. 169.]

Dr. MORISON was at this time abroad in charge of the Duke of Orleans' garden at Blois. Full of energy and enthusiasm for his subject, he wrote to How about the *Phytologia*, and sent him a pressing request for a number of plants which we have printed on p. 355. If How ever executed the order, some of the additions then made to the Duke's garden should appear in the Catalogues¹ that were printed in 1653 and 1669.

Lord HATTON transmitted his interest in Botany to his son Charles, who acquired the Boccone MSS. that were edited by Morison in 1674.

MORINE may have been the 'ordinary gardener' of Paris who became 'one of the most skilful and curious persons in France. . . . His garden is of an exact oval figure, planted with cypress cut flat & set as even as a wall: the tulips, anemones, ranunculuses, crocuses, etc., are held to be of the rarest, and draw all the admirers of that kind to his house during the season'.²

*MS. Additions by How to his copy of the 'Phytologia Britannica'.
Between 1650 and 1656.*

A certain number of the plant-names and notes that How wrote in his interleaved *Phytologia* have been quoted by Druce and others. They are of sufficient historical importance to be worth printing in full—or at least as far as they are legible.

¹ Brunyer, *Hortus Blesensis*, 1653; Morison, *Hortus Regius Blesensis*, 1669.

² Evelyn, *Diary*, 2 April 1644. The plants that Tradescant had from him in 1631 are noted on p. 331.

How's MS. notes on f. 1.

Plantae in aspectu in convictu nobis, semperque ad manum, nec fuga, nec ullis proripiuntur latibulis.

Universus fere ager amoenissimum viridarium.

Rariores quaedam et nondum quoad sciam exhibitae, ac a me observatae cum suis descriptionibus plantae commemorantur, ut quae adhuc laterent publico vix donarem, et ut quisque de iis, earumque facultatibus aptius indicaret.

Denique suas cuique plantae vires aut proprietates quam brevi ac dilucide fieri potuit annexi nec quisquam sibi nomina varietati tanquam labyrintho incumberet sed cum iisdem facultates etiam exploratas haberet.

Vid. Phyt. Pion 4^{to}.

Gainé I was for Goodyers Plants and des[criptions]: ye like for Brownes, Lobells [and Pennyes MS. w^{ch} review for names etc. (*erased*)].

Insert fig: iterum in Desc[riptiones].

Herbidos campos perlustratur.

Hanc nostratem Ἀντοκρατ ἀνταρκίαν Botanicae.

Vid. Epis. alt. libr: ad Coll. Med.

Cynoglossum flore albo neere Redding.

Arctium montanum et Lappa minor Gal. Lob. Button burr. Mangersfield in Mr. Langlyes yard, ill left out in yore Catal.

Acinos Anglica flore albo in Salisbury feild neere Basing stoake. D. Dale.

Verbascum syl: sive 4. Math: groweth naturaly about ye seate of Sr Th: Hanson called Toplow neere Maydenhead.

Erica tenuifolia caliculata, Ger: hirsuta Anglica, Bauh: ericeto Hamstediano.

Cardamine maior flore albo.

Bugula flore carneo.

Urtica foliis variegatis.

Speculum veneris maius in a ditch adioyning to St. Georges feildes.

Primula veris flore pleno viridi }? Morg[an], an inser:
 „ „ sive Paralysis fatua }

Arum non maculatum, Park.

Arundo anglica multifida, Park.

Malva syl: flore albo, Morg.

Millefolium pennatum, Bauh. neere Cambridge.

Ballote flore albo neere [Cholsy *erased*] at Tadenham ye south side of Mr. Crofts house Warwickshire.

Filago minor, Lob: neere Petersfield.

[Holosteum non descr. ? ye place Goodyer (*erased*).]

Alsine aquatica Portulacae fac: left out.

- Hieratium montanum at Mangersfield in Mr. Langtons yard.
 Cannabis spuria altera flore purpureo. 1 Cat.
 Saxifraga antiquorum, not inserted.
 Absinthium marinum spica erecta.
 [Bellis flore herbaceo globoso non descr., M. (*erased*).]
 Foeniculum at Rie — Leucii folii. *Qu.* D.
 Primula veris polyanthos, M. at great Wulford Wood.
 Agrifolium folio variegato Hampsted and in ye North.
 Serpillum magnum latifolium neere [Sr. W. Walters Oxfordshire *erased*]
 at Sarsden betweene ye house and ye pond garden of Sr. W. Walter.
 Geranium columbinum flore albo.
 Carduus vulgatissimus fl. albo in little Wulford field, Warwickshire.
 Geranium Rupertianum flore albo in a close betweene little Wulford
 and Bacton, Warwickshire.
 Perfol: purp. vulg. fl. albo by Whichford wood, Warwickshire.
 Consolida maior fl. cinericeo. Redding.
 Chamomelum nudum bullato flore. Tuthill feildes.
 Peryclymenum dissectis foliis. St. John Wood.
 Lamium elegans fl. lut. foliis variegatis neere Wulwich.
 Tubera terrae.
 Polium montanum neere Maidstone.
 Lamium rubrum pumilum.
 Lamium rubrum non serratum. *Qu.* an diff: a. 1
 Lamium rubrum foliis variegatis.
 Beta marina neere Lewis.
 Tapsus barbatus fl. albo at Weston.

Folio 2

- Carduus lanceolatus fl. alb., M. St. James.
 Rosmarinum syl. minus nostras, Park.
 Scabiosa ovilla flore albo., M.
 Sideritis hederulae fol., Park.
 Senecio foliis variegatis.
 Typha.
 Senecio sentilis? } *Qu.* D.
 Vicia bisiliq. *Qu.* G.
 Alsine (*Qu.* Holilas sen) minor. Androsace alterius Mathioli facie, Bauh
 Prod.

[Found by Mr. Halilah, Apothecary, in Lincolnshire. Merret, *Pinax*.]

- G. spartium capillaceo folio minimum, Ger. em. Ericet: Hamp:
 Quercus lapidea, Lithoxylum, Ligna lapidea. Ad. Lob.
 Sium maius angustifolium.
 Serpillum vulgare flore albo.
 Nymphaea lutea minor flore parvo, Catal. 1.
 Pulegium regium vulg. maius, Park.

Lychnis minor Anglica, Park.

Atriplex syl. *Halimi* fol., eiusd.

Pilosella minima, eiusd.

Muscus ex cranio humano.

Caucalis Anglica fl. rubente, Park. 919.¹

Pimpinella saxifraga minor nostras, eiusd. 946.

[*Genista spinosa minor*, Park. (*erased*).]

Rubus saxatilis Alpinus.

Lotus corniculatus frutescens, P. 1102.

Lotus corniculatus minor pilosus, *ibid*.

Hordeum spontaneum elatius, sive *maius*, P. 1147.

G. bromoides maximum hirtum, 1149.

G. avenaceum pratense, *ibid*.

G. bromoides segetum latiore panicula, *ibid*.

G. avenaceum murorum erectum, *ibid*.

G. avenaceum supinum arvense.

G. avenac: sup: *flosculis secalinis*, *ibid*.

G. avenaceum exile mollicellis foliis, *ibid*. All these oaten grasses grow in ye feildes of this land according to Park. ; bold assertions yett hee has no such warrant from Lob. Ill., from whence hee gathered them for rendring this exile mollicellis foliis spontaneall.

G. arundinaceum panicula miliacea, 1153.

G. montanum panicula miliacea sparsa, 50 in *Prod.*, *G. miliaceum alterum*, P. 1153, sine Ic.

G. arund: *Sorghii panicula sparsa*, 52 *Prod.*, *G. Sorghinum alterum*, P. 1153, sine Ic.

G. paniceum syl. *Anglicum*, P. 1155, sine Ic.

G. cristatum *Anglicum*, P. 1156, *cristatum Britannicum*, L. S. Ill.²

G. cristatum spica multiplici, P. 1160.

Phalaris pratensis maior, P. 1164.

G. alopecuroides cuspidatum maximum *Anglicum*, P. 1167, L. S. Ill.

G. alop: *spica aspera brevis*, et *longa*, P. 1169.

G. alopec: *cuspidatum minus et minimum*, *ibid*.

G. cyperoides *Anglicum parum lanosum maius et minus*, P. 1172, neere High gate. ?L. S. Ill. 2 variet.

G. iunceum medium et minus, P. 1190, in ye woode neere highgate, ?L. S. Ill.

Juncus capitulis equiseti alter, P. 1195, L. S. Ill.

Juncellus capitulis equiseti fluitans, P. 1196.

Spartum minimum *Anglicum*, P. 1199.

Equisetum omnium minus tenuifolium, P. 1201.

Anagallis aq; sive *Becabungae maior*, P. 1237. An: aq: ma: fol. subrot. Bauh.

¹ These references are to Parkinson, *Theatrum Botanicum*, 1640.

² Lobel, *Stirpium Illustrationes*.

- Sagittaria minor angustifolia, P. 1246.
- G. Arundin: sericea molliore spica, P. 1273, L. S. Ill. 45.
- G. maritimum vulgato canario simile, P. 1278, L. S. Ill. 24.
- Muscus arboreus nodosus, sive geniculatus, P. 1311.
- Muscus aridus crustatus, P. 1313, vid. loc. Bauh. Pin.
- Fungus Cambro-Brit. Corallii rub. colore multis lineis nigricantibus maculisque luteis orbiculatis insignibus, P. 1221.
- Fungus parvus nigricans crenatus, P. 1322, about Hackney.
- Fungus cinaræ formæ, P. 1224, at Ripton neere Ashford in Kent.
- Acer montanum, P. 1426.
- Primula veris Raii, M. *Qu.* loc.
- Quercus natalitiis Dni virens, ye Christmas greene oake, P. 1646, neere ye Castle of Malwood Hampshire, Kg. J. went to visit and caused it to bee paled about.
- Glycirrhiza syl: alt: fl: puniceis fol. Arachi, Glaux quaedam leguminosa herbariorum Adv: Bauh. Hist. Pl. l. 17 juxta agrorum viarumque margines obviam sese dat in Anglia.
- Hieracium Chondrillæ fol. glabrum, Bauh. Hier. 5, sive Aphacoides, Tab. Ger.
- Ulmus foliis luteis at Fulham.
- Hieracium Alpinum asperum Conyzae facie Bauh. Hierac. Brit., vel 4 Clus. Pan. et Hist. H. latif: mont: Genevense fol. long: maior: Monsp. J. B. H. P. l. 24, p. 1026. Borealibus Britanniae regionibus sponte crescit.
- Lychnis syl: foliis variegatis fl. albo. *Qu.* M.
- Virga aurea subrotundo fol. in Surry.
- Ascyrum, sive Hypericum bifolium glabrum non perforatum, Bauh. Hypericum in dumetis, nascens 1 Trag: Hyper. Ascyron dictum, caule quadrangulo J. B. H. P. l. 29, p. 382, ic. ib.
- Androsemon, Tur. In vinario sionis saepe observavit.
- Gentianella brevi fol., Bauh: non procul Douera.
- Anchusa Echii foliis et floribus, Bauh: arvis argillosis.
- Anthyllis chamaepityides frutescens, Bauh.
- Paralysis inodora calycibus dissectis, Raii: not farre from Kynlett¹ in Worcestershire: P. Par.
- Euphrasia coerulea not far from Canterbury, Mr. Hunnibon.
- Pseudonarc. fl. plen. Ger. Angl. fl. plen. P. P. ser.
- Serpillum foetidum Goodyeri, on ye chalkie downes 2 or 3 miles from Petersfield.
- Papaver cornutum flore phoeniceo, Adv. 241.
- Gentianella brevi fol: Bauh., minima L. ob. copiose provenit Angliae septentrionalis collibus.

¹ Probably Kinlet in Shropshire near Cleobury Mortimer, about a mile from the Worcester border (T.R.G.-P.).

Mollugo mont. angustif: B: vulgatior Herbariorum. L. Ob.
 Corallina fruticosa purpurea, B: rubens Antiphatis faire, L. Ob.
 Imperatoria Ad. Lob: Tab. Ger: about Morpeth in ye north parke,
 Tur. p. 37.
 Ribesium fructu rubro Dod: Ribes Fuch. Turn: by a water's side at
 Clover in Somersetsh:
 Cyclamen in ye west country of Engl: Turn: in his booke intituled Ye
 names of Herbes.
 Jacobaea montana neere Lewis.
 Hieracium lonchites, Bauh. B. Plants.
 Lactuca agnina rosea, neere red Morly.
 Capillus veneris verus, neere Lydberry.
 Cynoglossum fl. carneo, neere Bath.
 Alsine foliis Trissaginis variat fl. albo.
 Chamaedris syl: variat. fl. albo.
 Sedum aestivum minus.
 Plantago latifolia serrata.

Folio 3.

Lagopus maximus fl. rubro, P. nere red Morley.
 Ranunculus fl: lut: galericulato, at Yatton.
 Geranium batrachoides fl. variegato, neere Kidermaster Pitts.
 2. Pin: Gentiana maior ii Clus: fl: coeruleo neere Bath.
 Artemisia foliis variegatis neere Alchurch.
 Anagallis maxima latifolia neere Lidborough.
 Cracca minima variat. flore albo.
 Ascyrum maius neere Worcester.
 Muscus stellatus roseus, Bauh: Park. neere red Morly.
 Daucus proliferus, Qu. lo: Jolyff.
 Verbascum nigr. salvifol: purp. fl. Adv: 241. ii et nigrum latifolium luteum
 eiusd. 242.
 Juncus capitulo lanuginoso, sive Schoenolaguros, in Prod: Juncus Alpinus
 cum cauda Leporina. J. B. H. P. Moss crops. In ericetis nudis,
 Cambro-Britann. et Comit: Salop.

e Lob. MS.

Equisetum minus omnium tenuifolium.
 Synanchia altera Angliae, sive minor.
 Rapistrum aliud syl. non bulbosum, P. 862. Rapistrum fl. lut. alterum
 L. S. III.
 Cardamine alpina sive media, vid. Clus: cxxviii.
 Thlaspi alterum siliquosum.
 Planta juxta Heigate reperta, vid. nom. in Pin. Bauh.
 Sonchus laevis alter Danicus aut Anglicus folio profundis laciniis sinuato
 fl. lut.
 Hier: alt: fol. obtuse laciniatis fibrosa radice, Qu. MS. L. S. III.

- Brassica marina folio suaverubente, L. S. Ill.
 Beta maritima syl. spontanea eiusque varietates.
 Beta maritima syl: minor.
 Atriplex maritima altera Osyridis aut Scopariae fol., sive minima.
 Atripl. maritim. angustifol: 2 et 3.
 Limonium medium Anglicum.
 Hellebor. 2 Clus. sterilis planta.
 Campanula coerulea supina.
 Lychnis arvensis anglica.
 Pulegium vulg: maius minus repens cubitalis altitudinis.
 Scabiosa montana minor capitulo squarroso, Bauh. Pin. 270, in collibus
 Kantii non procul Kingshey.
 G. vulgatiss: pratense elatius tertium.
 G. rudium etiamque arborum, L. S. Ill. 6, multis locis Ag. Lond.
 gaudet.
 G. quoddam modo supinum vulg: latiusculo binum et ternum unciarum
 complicato folio.
 G. maritimum vulgatissimo pratensi gramini congener aut str.
 G. marit. alt. sive 2 elatius.
 G. marit. 3 vulgari str., supinum, exigua avenacea gluma.
 G. dulce udorum, vid. Bauh. pag. 2, Gr. vi.
 Gr. minimum Anglo-Brittan.

Folio 4.

- Gr. exile vicinorum maris aggerum, P. 1278, numerosa gracillimorum latius-
 colorum uncialium foliorum sobole.
 Gr. omnium minimum, Anglo-Britan. alterum.
 G. aquat. longius radicum spicata avenacea gluma.
 G. tenuifol. exile Britan: ex genere Xerampelini graminis
 G. maritimum Vectis Insulae Anglo-Britt.
 G. cuspid: tenui torosa villosa spicata gluma, Panici granulis prodita.
 G. phalaroides tremulum max: comosa elegantius gluma provenit in
 quibusdam collibus Britan.
 G. tremulum minus alterum Aquitanicum et Anglicum.
 G. hirsutum nemorum latioribus maioribusque foliis praecox vernalis.
 G. hirsutum sive exile ferrugineum.
 G. Anglo-Britan: exilis hirsuti graminis differentia Boelii.
 G. cyperoides aculeatum, sive . . (?) natum aquaticum alterum.
 G. cyp: minimum panicula subflava, Lob. Ad: part. 2.
 G. cyp: minimum nigricante panicula, Lob. Ad. p. 2, nullis uliginosis
 pratis Anglo-Br.
 G. cyp: palustre longius spicatum Anglo-Brit: acerosum et echinatum.
 G. cyp: minimum Boel: tenuifol: puis per caulem dissectis torulis. Provenit
 in Angl.-Brittan.
 G. cyp: sparsa panicula altae portae (= Highgate).

- G. cyp: aquat: tenui, triquetro, longoque caule, meduloso, junceo, bicubitali altioreque caule.
- G. cyp. comosa torulis distincta sparsa panicula palustre. Anglo-Britan.
- G. cyp. eleganti multifera congesta spica. Anglo-Britt.
- G. cyp. sylvarum tenuius spicatum.
- G cyp: gracile alterum glomeratis torulis, spatio distantibus.
- G. iuncoides alterum minus granatum, comitat: Kantii.
- Gramines parvus gracilis Juncellus Altae Portae.
- G. iuncoides tenuissimum subfuscis torulis apiculis carentibus proditum.
- G. iunc: minimum An.-Britt. Holosteo Math: congener, aut Bufonis gramine Flandrico et varietas.
- Mentha syl: verticillata tetri odoris.
- G. Xerampelinum exile. Ang. B.
- Angl. B. Panici effigies Gn. syl: L. S. III.
- Verbascum nigrum flore albo betweene Chesilhurst & Greenwich.
- Plantago latifolia incana spicis variis, Bauh. Pin. 789.
- Pl. peregr. latif: Gareti Clus. Hist.
- Plantag. latif. rosea multiformis, P. 494.
- Pl. ros. incana latifol: L. MS. gave ye Engl. name of Park.
- Pl. angustif. rosea, Park. 495.
- Pl. 5-ner. alt: sive angustif: roseo, folioso, sparso radiato fastigio. L. MS.
In borealibus Angliae tractibus utramque latifoliam, incanam roseam et hanc spontaneam esse intellexi.
- Sium foliis dissectis. }
Sium umbellis ad caulem nudis. } Qu. P. ye name place.
- Pimpinella saxifraga maxima. }
Pimpinella saxifraga maior foliis dissectis. } in Hampshire.
- Spina acuta biflora Britannica. Park. 1025, sive Oxyacantha.
- Erynus Math: in Hampshire.
- Juncus triquetrus by ye horsferry.
- Sambucus laciniatis foliis Dr. Jolyff¹ neere Winchester.
- Origanum roseum surculus densis, on Polstead downe in Compton parish nigh Gilford.
- Geranium violaceum at Putinham neere Gilford.
- Orchis foliis sessilibus non maculatis, Bauh.: 5 Clus. rar. Plant. pag. 268.
Similem etiam observabam MDXXCI in pratis urbi Londinensi in Anglia proximis, ubi et aliam eruebam grandiore paulo florum purpuratorum spicâ teterrimi factoris.
- Gentianella angustifolia autumnalis minor floribus ad latera pilosis, Bauh.
Gent. autumnal: fol: Centaur: min: flor: coerul: Eyst:, Gentianella

¹ Probably GEORGE JOYLIFFE, who entered Wadham College in 1637, migrated to Pembroke, and took his M.A. in 1643, being then a lieutenant under Lord Hopton in the Royalist army. He, with Dr. Clayton, discovered the Lymphatics in the liver about 1651. Samuel Pepys was one of his patients. Power, *Trans. Med. Soc.* xl.

autumnalis Centaur. minor fol: Park: pag. 406. Among his names there hee informes you yt it is ye x Gentian of Clus: wch if you compare theire Icons you will find it to bee Gentiana 11 minima Clus: pag. 316, and further affirms it to be Gentiana minima of Lob: wch Plant Bauh: worthily rankes under ye following title, not farre from ye ruines of ye old City Verulam neere St Albans.

Gentianella brevis fol:, Bauh: Gentianella minima, Lob: Gentiana x sive Gent. fugax 4 Clus: rar. Plant. pag. 315. Memini et in Britannia observare, non procul Douera Sept: partim dilutiore colore florentem partim semine pregnantem.

Alsine hirsuta minor, Bauh:, prope Rochester.

Fucus marit. alter tuberculis pauciss:, Bauh. marinus 4 Dod.

Fucus spongiosus ramosus.

Fucus folliculaceus foeniculi folio longiore, Bauh. Ferulaceus, Lob.

Helleborus pratensis latifolius, it growes in meddowes about Lichborough in Northamptonshire. } Br. lett

Serratula flore albo. } non descr.

How's MS. additions on leaves introduced in body of book.

Acorus verus offic. falso Calamus cum Julo, Calamus gromaticus vulgo Ger. em. pag. 63. I received from Mr. Th. Glyn of Glynrhinon in Carnarvonsh: ye pretty Julus or flower of this plant, which I could never see heere about London, though it groweth with us in many gardens and yet in great plenty. I rec^d: divers of them wth elegant flowers in June from Dr. Browne¹ who found them in severall places of Norfolk, these grew in Witton fens.

Alnus nigra baccifera, Tab. Lob. Frangula, Matth.

Est et species Frangulae fructu rubro proveniens in Angliae Comit. Somerset: prope aedes D. Thyn, an Vaccinium Plin: Lugd:

Althaea Ibisus. *Marsh Mallowes.*

Variat maioribus foliis et stipite littoreis circa Gravesend.

Anagallis foemina flore caeruleo. *Female Pimpernell.*

Faemina folia habet paulo quam in mare maiora, ima parte punctis nigricantibus multis insignita. Crescit copiose inter segetes iuxta Stanford.

Anonis foliis maioribus leniter crenatis flore luteo, in viis pratis non procul Bristolia.

Aquilegia flore caeruleo. *Columbine.* In a wood within 10 miles of London Kent.

Arbutus unedo. At Bellamont 3 miles from Dublyn.

Asarina foliis Asari, Bauh: Asarina, Math: Nec dispar foliis et odore Asarina devexa angusta via qua itur Clover versus Mendiep ubi effoditur plumbum. Florem videre non contigit illo tempore quo illac iter faciebam cauliculis serpit.

Ballote, Marrubium nigrum foetidum. *Stinking Horehound.*

Variat flore albo neere Gravesend.

¹ The well-known author of the *Religio Medici*.

Bellis minor vulgaris. *Little Daisies.*

Bellis minor flore viridi globoso, at Mr. Sheldens wood at Weston, Warwickshire.

Betonica vulgaris. *Betony.*

Variat in sylvis angustioribus aut rotundioribus foliis.

Calamogrostis huc referenda pro *Calamog:* Syl. D. J. Bapt: vid. non . . . Pin: Bauh: *Calamus* aromat. vid: *Acorus* verus.

Campanula media. Ista campanulae rotundifoliae similis, sed magna ex parte caulem habet unicum aliquando tamen plures sed hoc rarius, striatum, folia alterna per caulem rapuntii fere, sed latiora nec adeo longa, caules et folia sunt leviter hirsuta. Flores omnino purpurei campanulae praedictae floribus similimi, sed maiores et longiores multo. Radix parva paucis fibris. Provenit copiose inter Herefordiam et Kyneton, ac per totum Wye Flu: tractum.

Cannabis spuria altera flore purpureo. In agris. *Nettle-hemp.*

Cardamine pumila *Bellidis* folio. Alpina, *Ger. emac. Rock-cresses.*

By Shawford neere Gilford abundantly.

Christophoriana. *Herbe Christopher.*

Ad radices montis Ingleborrowe copiose inter saxa.

[A coloured drawing inserted.]

Cirsium Anglicum. *Single-headed Thistle.*

Radix nigricans fibrata quandoque per magnas fibras fere propagans ut in figura.

[A coloured drawing inserted.]

[? The last or] *C. Anglicum* minus, Park.

Crescit in pratis ad radices montis Ingleborrow totius Angliae altissimi in Comit. Eboracensi 12 miliaribus a Lancastria.

Cochlearia rotundifolia sive *Batava*, Lob. *Round-leaved Scurvy-grasse.* It groweth nigh unto a castle in ye Peake of Darbyshire which is 30 miles distant from ye sea.

Cochlearia rotundifolia marina.

About ye walls in Bermonice abundantly.

Echium flore alb. *Vipers Buglosse.*

Et in Cretaceis collibus ad Thamesim prope Greyn-hey tribus miliaribus a Gravesend.

Erica tenuifolia, Ger.

Coris fol. 6 Clus: *rur. Plant:* pag: 43. Nascentem hanc vidi in Anglia supra Windesoram mense Sept. florentem (Angli Boreales *Ling* vocare).¹

Erica coris folio 6 Clus: flore albo.

Erica coris folio 13 Clus: flore albo Saepius reperiuntur in ericetis circa Worplesdowne in Comit. Surriae.

Gentianella alpina.

In ye mountaines betwixt Gort and Galloway abundantly, Mr. Heaton.

¹ In Goodyer's copy of Parkinson's *Theatrum*, in the margin of p. 1480, is a note in the handwriting of the botanical friend of Goodyer (? Dr. J. Dale) who wrote MS. 9. Referring to Fig. 2, *Erica vulgaris* hirsutior, *Common rough Heath*, the writer observes, 'The 2 is the figure of *Erica Coris* folio, Clus. hist. p. 41, and not of the *Ericae myricae folii similis* found by Clusius about Windesor'.

Gramen plumosum elegans. *Fetched grass.*

Vepretis et quibusdam aridis agrorum marginibus Dorcestriensis Anglo-Britan: gaudet priori inutile.

Gramen tremulum. In upland cornefeilds at Hatfeild.

Gramen typhoides maximum spica longissima. *The largest Catstaille-grass.*

With a very large eare at ye entrance into Chelsy feilds.

Hederula aquatica. *Water Ivy.*

In a ditch by Bermondsey house neere London.

In aquis residibus iuxta Petriburgum.

Hieracium alterum foliis obtusa laciniatis fibrosa radice. Londinensis agri pratensibus, via Hackneum, Stepneum, et alibi passim obviam. foliis item humi stantis, hirsutis, duas uncias cum dimidia ternasque longis, obtusis dentatis laciniis: cauliculis dodrantalibus et pedalis floribus luteis papposis Hieracio longius radicato paribus et similibus donatis radice tamen non ita longa, sed fibrata.

Hieracium lactescens frutex est sesquicubitalis quandoque altior, caules multos ab radice fibrata satis, emittens, tenues tristes, intus fungosos et medulla alba plenos. Folia oblonga, serrata leniter, mucronata, parte ima modice pilosa, alterna per caulem, multis vel brevissimis pediculis cauli habentia, obscure virentia. Flores versus summos caules e cavis foliorum exeunt, pediculis longiusculis Tragopog: floribus omnino similes sed minores, lutei cuius foliola in extremis in quatuor partes dissecta. Flores nocte clauduntur mane sese expandunt. Floret Augusto, Septembri vero in pappos abeunt: Semen parvum oblongum sapore est ex dulis subamon, flores foliis sunt amariore paulo. Tota planta lactescit. Crescit in sylvis Comitatus: Hertford.

Horminum sylvestre Lavendulae flore.

Nascitur satis frequens ad Regium oris (?) Grenwicii Hippodromum.

Hyoseris mascula, Ger. *Male Swines Cichory.*

Non diff: ab Hieracio minimo Clus: pag: cxlii. Provenit inter segetes locis parum arenosis, et terra friabili; frequens in Anglia multis in locis.

Jacobaea, sive Senecion minimum. Radice pro plantae proportione maiuscula fibrata; folia Bellidi fere sed breviora parum incana 5, 6 vel 7 supra terram strata, e quorum medio caulis assurgit dodrantalis aut minor singularis in quo una, 4, 5 vel 6 quandoque folia sunt alterna, angusta mucronata. Caulis et folia lanugine candida sunt obducta. In summo caule flores sunt quatuor, 5, 6 rarique 7 breviusculis pediculis Senecii maioris, sive pubae Jacobaeae, pappescentes. Radix fibrata non vaga. Floret Julio et initio Augusti. Crescit in Agro Cantabrigiensi in parvis collibus non procul a Stapleford.

Jacobaea 3 Clus. Pann: latifol: s[essi]llis Planta Flore intense luteo et folio nigriore, vulgari Jacobaeae fere paribus, rivulorum sectatim Anglo-Britaniae.

[With a coloured drawing.]

Juncaria salmaticensis. *Small stone Woodrooffe of Spain.* At ye lower end of Grayes Inne lane by London neere ye water course.

Linariola parva planta dodrantalis aut brevior, caules aliquot a radice teretes et duriusculi, in quibus folia fere lini bina per intervalla. Flosculi parvi lini fere sed albi ex quinque foliolis in summis ramulis. Semina valde minuta lini effigie fusca. Floret Junio, Julio et Augusto in pratis Angliae frequens a nullo descripta.

[With a pen-and-ink drawing.]

Militaris aizoides, Lob. *Water soldier.*

Si non est Stratiotes Dios: certe similima est, sed non sine radice vivit.

Copiose prope pagum Over Jun. Julio et Augusto floret in aquis stagnantibus et fossis limosis Eliensis Insulae.

Muscus coralloides lacustris.

Corallii fere modo crescit, sed plures caules ab una radice, certo tamen spatio a se modice (?) distantes, utpote tres, 4, 5, et quandoque sex caules dodrantaes aut paulo longiores emittit minimi fere digiti crassitudine, teretes, spongiosos, ut et substantia totius plantae spongiosa est et porosa, et spongiae fere consistentiam in aqua habet.

Sicca vero dura et friabilis est, ex caulibus unus vel duo quandoque tres in summo sunt ramosi, in duos vel tres ramos ad cornu cervi similitudinem sunt divisi. Nec folia alia, nec flores nec fructum profert (quantum potui observare). Radix lapidibus in fundo adhaeret, supraque eos sese expandit. Tota planta unius est coloris, in aqua viridis saturate cum quadam nigredine, sicca vero diluta virescit. Invenitur in fluviis ex arborum truncis vel radicibus in aquis exiens ut in fluvio iuxta Petriburgum.

[An early description of the Fresh-water Sponge, *Spongilla fluviatilis*, illustrated with an excellent coloured pen-and-ink drawing.]

Myrrhis. Fuch., Cicutaria, Ges. hort. *Wild Sweete Cherville.*

Napus syl: minimus Montosis agris hoc pusillum Napi syl: genus iuxta pistrinam ventillatam sive ventimolam D. Rich. Garth semiliari ab aedibus antiquis Drayton vocatis e regione Vectis Insulae reperi; tota planta admodum parva est, foliis Napo syl: longe minoribus, angustioribus, parum [4 lines of description follow].

Nidus avis flore et caule violaceo purpureo colore an Pseudo-limodoron, *Clusii, hist. rar. Plant.* pag. 270: A mile from *Alton* in Hampshire. Mr. *Goodyer.*

[To this printed note, How has added in MS.] Limodoron Austriacum *Clus. Hist.* Orchis abortiva violacea, Bauh:

Orchis minor flore carnato Eyst: Pannonia 4 Clus: Cynosorchis minor Pannonica, Ger. em: The lesser *Austins dog-stones.* On Scosby lease. Mr. Stonehouse.

Oxalis major. *Great leaved Sorrel.*

Ad Shackerforth-mill loco uliginoso.

Papaver corniculatum flore rubro. Copiose crescit Vectis Insula.

Paronychia.

Folia radici proxima superne obscure rubent, interne virent. Lobelii mutilam et depravatam iconem Lugd:, Dod:, Ger. em., Park. etc. exhibent Omnes esse sequest (?) maturam e vivis naturae typis exhibeo.

[With a coloured drawing.]

Pedicularis flore albo. *White Rattle.*

Severall places of Warwickshire.

Polygonum minimum.

Plantula vix unciam altitudine excedit, ramosa, ut arbor assurgit, surculosa, erecta magna ea parte, foliola parva binatim coniuncta, e geniculis exeuntia. Flores in summo parvi numerosi, albicantes. Radice nititur minima singulari paucis fibris, crescit locis hyeme udii, aestate veneniatii. An potius ad *Alsines* speciem referenda?

[With a small neat drawing in pen and ink.]

Pulsatilla vulgaris. Purple passe-flower.

About Oxford.

Pirum foliatum.

[Coloured drawing of a pear, with well developed green leaves growing from the apex.]

Pirum supra pirum.

[A fruit with a second pear growing among the leaves at the apex.]

Ranunculus minimus Septentrionalium herbido muscoso flore. Dwarfse Crow-foot, or Small Bunikins Hollwortell.

In Anglia frequens praecipue versus Septentrionem Folia odorem epirant moscho similem, sed valde imbecillum.

Fumaria cubica vel capnos Moschatella Cordi.

[A coloured drawing marked] Haec effigies praeferenda.

Ranunculus montanus, Cam. globosus. Globe Crowfoot.

Floret in fine Jun et Jul: ad radices montis Ingle-borrow. Angli boreales. *Locker-Gowing* vel *locker goling*, a flore clauso non est venenatus.

Ranunculus pumilus floribus deciduis.

The whole plant seldome exceeds 3 inches in compasse and in his full strength and flowring is not above an inch or 2 high, amongst a hundred plants of them yt I found not far from Oxford though it were in ye time theire flowring I could not find one with a whole flower, severall of these had 3, 4 or 5 little yellow leaves of flowers, about a small thrum of yellow pointells, and every plant had fresh yellow pointells, with the leaves as it were new fallen of. [it has not yett flowred in my garden.]¹

Rosa sylvestris odora Eglenteria. Sweet Briar bush.

Oritur in Bathonica Angliae proximis collibus non procul ab aedibus

D: Laur. Hyde. [Quoted from *Lobel, Obs.* 618.]

Rubia sylvestris laevis radice perenni.

Omnino priori similis sed radice vivace ac perenni, humi maxima ex parte procumbit planta et multis surculis brachiata, folia paulo latiore stellatim disposita leniter serrata flores coerulei ex quatuor foliolis mucronatis. Caules quadranguli geniculati, e singulis geniculis folia stellatim decussata quinq, quatuor, sex septem et octo. Radix longa fibrata valde rubescens. Juxta Harwica prope littora maris.

Sagittaria minor. Small Arrow-head.

Radices—multas fibrosas, pallidas vel albicantes, prolixas inter quas aliquas utpote 4, 5 vel 6 maiores radices chordis maioris Lyrae instar reliqui longiores in extremo bulbosas bulbi oblongi mucronati, incurvi, membranulis duabus aut tribus tecti, colore ex albo et coeruleo mixto, bulbi magnitudine inaequales, nucis avellanae magnitudinem aequantes, quorum caro alba, solida.

[A pen-and-ink drawing entitled] *Sagittae minoris radix cum bulbis.*

Sambucus foliis variegatis neere Totnam.

Sedum arborescens Anglicum. Frutex est varius multis lignosis duris, cuius folia sunt magis longa, similiter disposita, non adeo carnosae, nec crassa, *Sedum vulgare* situ et ortu aemulantia, radice satis longa, et crassiuscula non multis fibris. Crescit in Insulis Hooms vocatis sinus Bristoiensis, Oceani Anglici.

¹ Erased.

Sium alterum Olusatri facie. *Long-leaved Water Cresses.*

In fossis prope Petriburgum frequens, nec non procul ab Oxbridge in quibusdam puteis Londinum versus iuxta viam publicam.

Spartum spica secalina. *Great English Mat-weede.*

On ye further side of ye Isle of Tenet.

Trachelium majus Belgarum. *Giant Throat-wort or Bellflower.*

Pratensibus ad radices cretaceosas collium prope Dartfort et Greehyth.

Trifolium flore viridi foliaceo elegans.

In mine owne Orchard at Darfield: it grew with mee one yere plentifully but I have not since observ'd it. I have now sent some of it dried as it was gathered about 9 yeeres since. Mr. Stonehouse.

Turritis major. *Towers Mustard.*

Non procul a meta antiquissima diruta, miliari a Colchestria et in agris prope Deditfort.

Typha minor crescit in fluvio Petriburgum praeterfluens nusquam alias vidi.

Vaccinia nubis. *Cloudberry.*

Chamaemorus Anglica, Park. *Chamaemorus Cambro-Britanniae* sive *Lancastriense vaccinium nubis.* Radix utcunque nodosa, et ex nodis fibras paucas dimittit; radice se propagat (sicut *Cirsium Anglicum* vid. fig.) et quam longissime serpit adeo ut brevi tempore maximum spatium occupet.

Vicia maxima sylvatica nondum descripta, spicata Bathoniensis Goodyeri.

Umbilicus Veneris.

About Bath and North Wales plentifully.

Umbilicus Veneris maximus Anglicus rudentibus foliis maior ac elegantior quam precedens est. *Pilleter. Plant: Synon:* pag. 126.

MS. additions at end between pp. 132 and 133.

Cotula non foetida flore pleno latiore.

Chamaebuxus fl. colutea, Bauh: sive *Rhus* Plin: myrtifol:

Carduus lanceolatus fl. alb et fl. purp: Q. Chyrurg: for ye places of these plants growth from Morgan.

Taxus tantum florens on ye chalky hills in Hampsh. J. Goody:

Persic. minor non urens sine maculis in Tuthill Feilds.

Antirrhinum medium Hispan: flore albo, Hunnibon and what other spontaneally hee hath with all other simples.

[Species of *Cynoglossum*, *Hypericon*, *Orchys*, and *Sium* from Browne See p. 302.]

<i>Oxyacanthus flore rubro</i>	} <i>Qu.</i> Hunnibon.
<i>Alsine foliis variegatis</i>	
<i>Seseli pratense</i>	

Qu. ye *Sium* in ye pott with small leaves per Phy. Br.

{	<i>Gentiana altera dubia Anglica punctato medio flore.</i>
	<i>Helleborines uti superior forte cognata</i> , L. St. III.
	<i>Gentiana dubia Anglica</i> , Park.

Q. Newarkes *Turritis*. *Pentaphyllum*.

Q. ye *Opuntium marinum* on Oyster shells.

Nymphaea foliis hederaceis. *Qu.* Stevens.

[Species of *Carduus*, *Plantago*, and *Echium* from Browne. See p. 302.]

Prunella vulg: fl. al. incarnato et fl. purpuréo.

in Chappell on ye heath. Bobert.

Stevens his Becabungae maior *Plantago Aquat. latif. maior.*

Geranium columbinum foliis magis dissectis, pediculis longissimis flore magno. I found it wild in ye beginning of August 1654, it is not described or pictured yt I find. John Goodyer. *Q.* ye place of growth. vid. *Phyt.* 47.

Erysimum ii Tab. *Q.* locum.

growes in ye streetes neere white chappell east from Algate London. J. Goodyer.

Anonymos aquatica rubida, foliis Anagallidis flore luteo. This growes in a little lake in a heath neere Petersfeild in Hamshire, in a hott summer some parts of ye lake are drie in August, sometimes before, there and then ye flowers are to bee seene.

Holosteum perpusillum growes in ye same lake¹ in ye East part of ye said heath greene all ye winter under water, and flowers when ye water is vanished in August, and sometimes much sooner. I first observed this plant in a pond neare Holburie in ye new Forrest in Hamshire. J. Goodyer.

¹ The waters of this lake this 2 of June 1656 about 4 of ye clocke in ye afternoone was well neere as warme as ye Bathwater at Bath in Summersetshire although ye day was cloudy.

Holosteum juncifolium repens Goodyeri copiose inveni in Comit: Surriae juxta Purbright (an diff: a priore, *Qu.* Goody).

But How's good intentions for a revised edition of his *Flora* were not destined to bear fruit. Chance appears to have cast the Lobel manuscripts in his way: he purchased them with his own money, but his store of vital energy was unequal to the task of producing a second *Phytologia Britannica*.

The circumstances of his dealing with these manuscripts have been often described. A contemporary account of the matter is given by the Rev. J. Ward in his *Commonplace Book*: 'Dr. How hath put out a piece showing what Plants Parkinson stole out of a manuscript of Lobel's wch. never was put out, but came by chance to Dr. Modesy's [= Morison's] hand.' How has been universally reproved for the violence of his language, *re* Parkinson, first by Pulteney and then by later historians, who, however, quote his remarks with some gusto.

If How had found the Lobel MSS. in anything like the disorder in which they were when I first saw them, and if he attributed the mutilations to Parkinson, his expressions might be considered as

well justified. It is more likely that the vigour of his language was partly due to haste consequent on failing health, and perhaps to a presentiment of death and to the knowledge that his time and resources were insufficient to do his author more than scant justice. How died in 1656, a few months after the publication of his book.

We have consulted William Howe's will (P. C. C. Berkeley 315) at Somerset House. He described himself as of Milke St., co. Middlesex, Gent., 'being in perfect memory though much crased in Body'. He willed that his body should be 'very humblie interred in St. Margaret's Church, Westminster, on the left side of my mother. The earth to be taken up att least 6 foote, my funeral I desire may be observed about ten at night. I would not have above 6 of my choyce friends accompaning'. His wife Elizabeth Howe was appointed absolute executor, but as regards his Library he 'would have her advised by some knowing person to putt of, and by no means to part with any particular book from the whole'. Will proved, 22 Sept. 1656.

It is probable that Goodyer then acquired a selection of his books and manuscripts. The most important of these, which are still in the Library at Magdalen College, are the Lobel Manuscripts, How's own annotated copy of his *Phytologia*, Johnson's *Descriptio Itineris* 1632, both with the MS. additions printed below, certain loose papers including Goodyer MS. 11, ff. 169-73, described on p. 355, and a few printed books including editions of *Tabernaemontanus*, *Matthiolus*, and *Renealmus*.

William Cole or Coles of New College gratefully acknowledged the help that How gave him in the preparation of his *Adam in Eden* 1657. 'Furnished' with How's 'best advice and with some of his select and choicest papers for perfecting his design', it is a pity that Coles did not produce a sounder book.

xiii. DR. JOHN DALE, *d.* 1662.

In the Latin introduction to his *Pinax rerum naturalium Britannicarum*, Dr. Christopher Merrett, F.R.S., tells us that the one and only comprehensive Flora of Britain that had ever been printed was almost out of print, and that Pulleyn,¹ a publisher living in St. Paul's Churchyard, had asked him to undertake a new Catalogue

¹ Octavian Pulleyn was the publisher of How's *Phytologia* in 1650; Cave Pulleyn published Merrett's *Pinax* in 1667. Octavian sold to How the copy of *Tabernaemontanus* which was afterwards acquired by Goodyer: his signature is inside the cover. See pp. 202 and 226.

of all the British plants then known, to be followed by a History, and that the work was to be done in association with a DR. DALE, 'Botanologus peritus'.

The only botanist of that name in Merrett's time who is at all well-known, or indeed appears in Messrs. Britten & Boulger's *Index of British Botanists*, is Dr. Samuel Dale of Braintree, who in after years gave such valuable assistance to Ray, and was often mentioned in his botanical works. It is, therefore, but natural for an uncritical reader, on encountering a solitary mention of Dr. Merrett's proposed partner Dr. Dale, to assume unconsciously that the reference was to Dr. Samuel Dale. Merrett was, however, writing at the College of Physicians in August 1666, a few days before the outbreak of the Great Fire, and Samuel, who is believed to have been born in Whitechapel in 1659, could not then have been a distinguished 'Botanologist'. Nor must he be confused with William Dale of Queen's College, the helper of the younger Bobart.¹

John Ward,² writing in 1662, also mentions Dr. Dale, and rather as if he were the originator of the idea of re-editing the *Phytologia Britannica*; and, lastly, John Ward's editor, Sir D'Arcy Power, notes that 'no record of his (Dale's) attainments in botany seems to have survived'. Ward had, however, a high idea of him: 'there are in London but two doctors y^t have any great skill in simpling, y^t is Dr. Moddesey (*i.e.* Morison) and Dr. Dale'; and in 1661 he noted one of his prescriptions.

'White Mullen or Higtaper: wash itt and fume itt and boyl itt with hog's grease, yⁿ add Red Lead and Linseed Oil. It is said to bee excellent good against y^e piles. *Dr. Dale.* I find ye same in a manner in Gerard's *Herbal* with but very little difference.'³

Except in a single instance I have not met with any mention of the name of Dale among the Goodyer papers, but nevertheless there are good grounds for attributing to him the authorship of certain manuscripts, and a few notes in the margins of Goodyer's copies of Parkinson's *Theatrum*; Turner's *Herbal*, p. 10; Thalius

¹ 'Mr. Bobart, the Botanist, was greatly assisted in the II^d vol. of ye Oxford History of Plants, by Mr. Dale of Queen's College, who revised the whole and put it into proper Latin for him. Presently after the Death of the said Mr. Dale, I had a sight of a Folio Book in MS^t drawn up by himself, being Tables & Explications on Aristotle's Rhetorick' (Hearne, *Diary* 1705, Nov. 6).

² The Rev. John Ward of Christ Church had taken his M.A. in 1652. His sixteen commonplace books are now preserved in the library of the Medical Society of London. (D'Arcy Power, *Ann. Med. Hist.* ii, p. 123.)

³ Information from Sir D'Arcy Power, who has been kind enough to transcribe for me the paragraphs relating to Dale in J. Ward's *Diary*.

Sylva Hircynia, p. 51; *Gerard emac.*, p. 800, and other books. At any rate, it appears more likely that Dr. Dale, rather than any one else, should have written them.

Dale's botanical partnership with Goodyer dates at least from 1651, as is shown by the latter's entry in his copy of J. Bauhin's *Historia Plantarum*. This work was purchased of Robinson on 15 March 1651, and sold by Dr. Dale, then living in Long Acre, to Goodyer on the 22nd for £3 2s. 6d.

The Goodyerian Lists and Descriptions of British plants were undoubtedly written during the sixth decade of the seventeenth century. One paper is dated 22 January 1651, another April 1659, and during part of this period the writer was evidently in close touch with Goodyer himself. The four principal manuscripts are:

1. Descriptions of Plants extracted from Lobel's MSS. (in Latin).
2. A Catalogue of Grasses, foreign as well as British, with brief descriptions and synonymy, comprising 203 species, written two on a page, on 101 leaves.
3. A Catalogue of British plants, with synonymy, comprising 922 species (including 70 grasses), written four on a page (with blanks), on 276 leaves.
4. A Supplementary List of British plants comprising 153 names. If these are to be regarded as different from the 922, the British Flora as known to Goodyer and Dale (?) in 1659 would comprise 1,075 species—a figure which comes very near to the 1,050 of Ray in the year 1669. How, by including varieties and exotic plants, had accumulated 1,220 names in the *Phytologia* of 1650.

The work is exactly what would be expected from the author of a new *Phytologia*. None of Dr. Dale's plants are quoted under his own name in the printed *Phytologia* of 1650, but among How's MS. additions (1650-6) we have noted

'Acinos anglica flore albo in Salisbury feild neere Basing stoake, D. Dale.'
 'Foeniculum at Ric. — Leucii folii: Qu. D.'
 'Typha
 'Senetio sentilis } Qu. D.'

He was the first person to point out 'Juncus caule Triangulati' (? the rare Galingale *Cyperus longus*) growing at the Horse ferry at Westminster and to record 'Vicia fol. gramineo siliqua porrectissima' about Tyburn and Maribone Park. Merrett, who printed these records in 1667,¹ designates him as 'insignis Britannicus'.

¹ Merrett, *Pinax*, 67, 125.

There can, therefore, be no doubt that there was such a person: the trouble is to identify him among the numerous Dales who were living at the time. Unfortunately his christian name is nowhere mentioned. I was at first inclined to think that Sir D'Arcy Power, who has the credit of a first attempt, was right in identifying Dr. Dale with the Dr. Robert Dale, said to be of Magdalen College, who was admitted an Extra-Licentiate of the College of Physicians in 1663.¹ But this is rather too late a date for our botanist. Merrett alludes to his death, but not as if it were a recent event. It may have occurred before the Fire and the Plague, probably shortly before Goodyer's own death in 1664, for else it would be hard to explain the presence of Dale MSS. among the Goodyer papers.

While engaged in an unsuccessful hunt for Robert Dale's will at Somerset House, I found the will of Dr. John Dale, which connects many of the clues and is a surer guide to the identity of our botanist.

JOHN DALE, Doctor of Physick, of the parish of St. Martin's in the Fields, bequeaths to poore neighbours of St. Martin's Parish £5; to poore of East Meon £5; to poore of Petersfield £5; to poore of Gosport £5; to Mr. Hunt £5; to Mr. Gray £5; to brother Andrew Vidian and wife £5; to sister Codd 20s., to sister Browne 20s., to Mr. Darlaston and wife 40s., for rings; to wife Blanche Dale, his sole ex^{or}, lease of house in Long Acre and lease of his impropriate parsonage of East Meon. Brother-in-law Andrew Vidian and 'my very good friend John Goodyer of Petersfield' to be overseers and to have a ring of 40s. Will dated 30 Apr. in the presence of W. Darlaston, Mary Robinson, Eliz. Coleman. Proved 27 May 1662, after the death of Blanche his widow.—*Abstract of P. C. C. Laud.* 63.

We hope that some local genealogist will follow the family further. 'Sister Browne' may have been related to the Brownes of East Hoo, already mentioned; and Mr. Gray may have been the herborist friend of O. Bilson (p. 201), but the names are too widespread to be a trustworthy guide.

¹ Munk records that Robert Dale was a Bachelor of Arts of Magdalen College, Oxford, who practised medicine at Stourbridge in Worcestershire and was admitted an Extra-Licentiate of the College of Physicians of London on October 1st 1663. (*Roll of the Royal College of Physicians.*) But I have not as yet been able to find any Robert Dale in the books of Magdalen College. John Ward elsewhere refers to an incident in the practice of a 'Mr' Dale, and, in the next sentence, to 'An ichneumon frequent about Sturbridge'. Unfortunately Ward nowhere mentions his Dales' Christian names. Sir D'Arcy Power, to whom I mentioned the difficulty, has again investigated the matter, and has found among the Prattinton MS., vol. 31, at the Society of Antiquaries, an entry: 'Robert Dale Ludimagister of Free school at Stourbridge, 9 Nov. 1661'.

It seems probable that Goodyer, if not already in possession of Dale's botanical papers, would have annexed them as part of his duty as overseer of the will, and on his death (1664) they may have been among those which Yalden lent to Merrett.

Dr. John Dale, unlike Robert, does not appear to have been licenced by the College of Physicians, of which institution Dr. Merrett was the resident Librarian. The College of Physicians was then situated in Amen Corner—very near the house of Pulleyn, the friendly publisher who persuaded Merrett to co-operate in the proposed work. 'I could not strive against his honourable wishes', as Merrett afterwards put it. How far Dr. Dale did co-operate, we are not told. He died before the day of publication.

Merrett certainly engaged in the work with great energy. He purchased 800 figures of plants, which Johnson had caused to be engraved, with the intention of using them to embellish his book. He engaged Thomas Willisel to search for plants in those distant parts of the kingdom which, owing to his London duties, were beyond his range. His son, Christopher, also made excursions for the same purpose, and he procured the loan of Goodyer's MSS. from Edmund Yalden.

By the inclusion of exotics and many varieties, which he had not the critical acquaintance with the subject to omit, he brought up the list of the British Flora to some 1,400 species, a number which, only three years afterwards, was reduced to 1,050 by the 'accurate Mr. Ray'.

xiv. WILLIAM BROWNE, 1629-1678.

Several of the plant-records of William Browne are quoted in the interleaved copy of How's *Phytologia Britannica* bequeathed to Magdalen College by John Goodyer. They are all in the handwriting of William How.

William Browne was known to Wood as his examiner, and also to Peshall as a native of Oxford, his father being described alternatively as William Brown, a Mercer of Oxford,¹ or as John Browne, a Bailiff of the City.² He had a distinguished career at Magdalen College, becoming B.A. in 1647, M.A. in 1650, and B.D. in 1665. He succeeded to a Fellowship in 1657, was Praelector of Moral Philosophy in 1658, Dean of Divinity in 1659, and Vice-President in 1669-1670. He died suddenly about the age of fifty, and was

¹ Peshall, *City of Oxford*, *add.* 29.

² Wood, *Hist.*, p. 344, ed. Gutch.

buried in the Antechapel of his College. A gravestone of black marble preserves his memory :

H. S. E.
 GULIELMUS BROWNE S. T. B.
 HUIUS COLLEGII SOCIUS
 VIR
 INDUSTRIAE INDEFESSAE,
 ERUDITIONIS PERSPECTAE :
 QUI S. THEOLOGIAE HORAS COMPOSITAS
 REI BOTANICAE SUCCESIVAS
 IMPENDENS,
 IN UTRAQUE EMICUIT
 APOPLEXIA CORREPTUS SUCCUBUIT
 FATO MULTUM LUGENDO
 NISI VIXISSET INDIES MORITURUS.
 OB. MAR. 25, AN. AET. 49°
 MDCLXXVIII.

Canon Vaughan¹ favours the view that the presence of so well-known a botanist at Magdalen College would account for Goodyer's gift of botanical books to the College; and this idea would gain in probability if we could establish a close kinship between William Browne of Magdalen and Goodyer's Hampshire neighbour of the same name. But so far we have not found any evidence on this point.

Mr. Druce (*Flora Berks.*, p. cvi) is not quite right in saying that no certain writing of Browne's has been discovered, for we have specimens of his signature at Magdalen, and for several reasons we believe that Druce is also mistaken in suggesting that the MS. notes in one of the Bodleian copies of Lyte's *Herbal* were made by Browne: they are obviously by an earlier member of Magdalen College who had studied at Padua.

Browne is best known through the *Catalogus Horti Botanici Oxoniensis* prepared by Bobart, Dr. Stephens, the Principal of Magdalen Hall, and himself in collaboration, but both Anthony Wood and Merrett agree 'that he had the chief hand in it', and the MS. copy in the British Museum gives his name as that of the author.

Attention has been drawn to the fact that in this edition the authors have, in every instance where it was possible, not only adopted the scientific appellation given by Gerard and Parkinson

¹ *Cornhill Magazine*, 1909, p. 802.

to each plant, but also quoted the page of their works. Druce¹ claims priority in England for this procedure for the authors jointly. It is quite likely that Browne, rather than Bobart, was responsible for the innovation, but as the quotation of authorities by pages had been the usual everyday practice of Browne's friend Goodyer for the past thirty years, any credit for priority or precept in this matter should be given to the latter.

There are many citations from him printed in Dr. Merrett's *Pinax*, and although he does not appear as one of the contributors to the *Phytologia*, perhaps because How was not acquainted with him before 1650, or because he had not then paid much attention to botany, yet he was able to supplement that work with several valuable additions which How duly entered in his interleaved copy of the *Phytologia* already referred to. Most of these citations are distinguished by a peculiar mark, a Δ in a square.

The Plant Records of William Browne, entered by How in his copy of the 'Phytologia', 1650-1656.

Anagallis aquatica, sive Becabunga flore albo. *Veronica Beccabunga* L.
in fossis aliquibus sub colle Heddington prope Oxonium. Guil.
Browne Oxon:

Atriplex marina latifolia tota rubra. *Atriplex patula* L.

E radice tenui et fibrata caules exit sesquicubitales ramulis donatos frequentissimis folia ferentes in extremitate angusta et acuta in media et una parte lata et acuminata ubique densa et admodum rubra hortis translata ruborem fidelissime servat, flosculi ex viridi rubescunt, semina sunt angularia, et eiusdem ruboris cum Asteris (?) plantae partibus occurrit ad maris littus prope Shoram comitat: Sussexi. Guil. Browne Oxon:

Behen album hispidum. *Silene Cucubalus* Wib.

Planta est ubique obvia adhuc tamen non descripta. Guil.
Browne Oxon.

Colchicum Anglicum purpureum duplici serie foliorum in flore.

Colchicum Anglicum saturationis purpureae.

Colchicum Anglicum florum foliis ex albo et purpureo dimidiatim variegatis. *Colchicum autumnale* L.

Haec tria proveniunt in prato amoenissimo prope pagum vernaculem Combe, Comit: Oxon; ubi flore albo plurima sunt et vulgaris purpureae millia. Guil. Browne Oxon.

¹ Druce, *Flora Oxford*, p. 373.

Colchicum Anglicum, foliis elegantissime striatis.

in prato prope Corneberry, Comitatu Oxon., gramen striatum. pulchritudine multum superat colores horto meo pluribus annis iactavit. Guil. Browne.

Cotula alba. Mayweed.

Anthemis Cotula L.

Et flore pleno ex luteo viridi reperitur [juxta Oxon. *erased*] Dorcestria versus Comit. Oxon. via regia vulgo Honey Fixlong. Guil. Browne Oxon.

Geranium columbinum minus foliis magis dissectis et foliis minus dissectis in agris sterilibus. variat. flore albo. Guil. Br. Oxon.

Geranium molle L. and *G. dissectum* L.

Lysimachia, sive *Gratiola latifolia* flore albo in pagulo vulgo Purbright Comitatu: Surriae. Guil. Browne Oxon. *Scutellaria minor* Huds.

Orchis Antropophora trunco pallido, brachiis et cruribus saturate rubescentibus. The Red Shanke. Roy Satyrion. Hasce Orchides rariores in cretaceis quibusdam collibus observant non procul a via communi qua itur Wallingfordio Reddingam per Comitatum Bercherium. Guil. Browne Oxon. *Orchis simia* Lam.?

Orchis, sive *Cynosorchis militans* holosericea, hanc orchidum merito reginam inveni juxta mediam partem viae communis inter Nettleton et Bathe communicante D.D. Stevens Botanico perito. G. Browne Oxon. *Orchis militaris* L.

Orchis, sive *Cynosorchis Austriaca* flore albo colle Chilswelliensi prope Oxonium. Guil. Browne Oxon: *Orchis ustulata* L.

Periclymenum syl: 3plici serie florum ex luteo virentium, alioqui toto habitu a vulgari non dissimile, colle vulgo Shotover juxta Oxonium.

Lonicera Periclymenum L.

Periclymenum alterum quercinis foliis, perelegans planta; observavi in colle Chilswelliensi prope Oxonium. G. B. Oxon.

Plantago quinquerivaria fimbriis latis ex aureo argenteis, hanc nitidam plantam juxta Corneberry exploravi Comitatu: Oxon. Guil. Browne Oxon. *Plantago major* L. var.

Ranunculus pumilus floribus deciduis. ? *Ranunculus parviflorus*.

The whole plant seldome exceeds 3 inches in compasse and in his full strength and flowring is not above an inch or 2 high, amongst a hundred plants of them y^t I found not far from Oxford though it were in ye time of their flowring I could not find one with a whole flower, severall of these had 3, 4 or 5 little yellow leaves of flowers about a small thrum of yellow pointells, and every plant had fresh yellow pointells, with the leaves as it were new fallen of. [It has not yet flowered in my garden. *Erased.*]

Rosa pimpinella foliis flore suaverubente, in agris sterilibus prope Worcestriam. Guil. Browne Oxon: *Rosa spinosissima* L.

Saxifraga aurea, maior, foliis, pediculis longis insidentibus. Juxta radicem tenuem albam et repentem folia edit rotunda ex luteo virentia innata itaque hirsuta, paediculis paene triuncialibus saniculae guttatae foliis non dissimilia; cauliculos profert plurimos palmares juxta summitatem parum divisos vel plurimum unico tantum folio, aliquando altero ornatos; flosculi aurei foliis plurimis decore lutescentibus longescunt. Hanc saxifragam vere auream copiose inveni in paludoso nemori non procul ab aedibus D. Fanteleroij in pago vulgo Hedley vocato Comitati Hamptonis ubi Dryopteridis Trag. iuxta et Calami Aromatici cum Julio ad humanam altitudinem assurgentis gliscit copia. Guil. Browne Oxon.

Chrysosplenium alternifolium L.

Solanum marinum Dulcamarae congener. *Solanum Dulcamara* L.

E radice longa dura et perenni, caulem profert crassum glabrum atque humi procumbentem, pedalem vel sesquipedalem longitudinem nunquam superantem ubi sponte emergit, folia ferentem Solani lignosi duplo densiora et saturationis multo viriditatis, Augusto floribus ornatur et albis et violaceis, utrius coloris copiose: reperitur ad maris littus iuxta Shoram Sussexio. Guil. Browne Oxon:

Trachelium minus flore albo in pago Worplesdowne vocato, Comitatus

Surriae locus floribus albis admodum insignis. Guil. Browne Oxon:

Campanula glomerata L.

On blank page at end.

Cynoglossum foliis variegatis.

Hypericon flore albicaster.

Orchys Sphegodes flore albo.

Mrs. Yalding about Guilford.

Sium cubitate fol. variis non des[criptum]: a very beautifull plant about Redding plentifully. An diff. a Sio foliis variis Phyt.? *Apium inundatum* Reich. f.

} Qu. Br.

Carduus pratensis caule folioso is a common plant about 3 foote high and has ye same filmy stalk as ye Aster Virg. caule membranaceo Park: wee often find it with white flowers. *C. pratensis* L. var.

Plantago marina foliis tenuissimis, haec singulis partibus alteri non dissimilis est, sed foliolis multo tenuioribus praedita.

Echium scorpioides minus flosculis luteis Bauh. Pin. 254 growes within 3 miles of Redding plentifully. *Myosotis versicolor* Sm.

On blank page, f. 4 v.

Helleborus pratensis latifolius, it growes in meddowes

about Lichborough in Northamptonshire. *H. viridis*.

Serratula flore albo. (*Serratula tinctoria* L.)

} Qu. Br. lett non. descr.

[Goodyer MS. 18.]

EARLY LISTS OF PLANTS GROWN IN ENGLISH GARDENS.

HUGH MORGAN of Coleman St., 1569-87. See under No. xiii

- i. OXFORD AND WINCHESTER GARDENS, 1570-2.
- ii. SIR JOHN SALUSBURY of Lleweni, 1596, 1607-8.
- iii. SIMON FORMAN, 1597-1608.
- iv. LORD SALISBURY, 1611.
- v. RICHARD SHANNE of Woodrowe, Methley, 1615.
- vi. WILLIAM COYS of Stubbers, 1604, 1616, 1621-2.
- vii. FRANQUEVILLES' Garden in Long Acre, 1600-4, 1617.
- viii. JOHN PARKINSON, c. 1618-20.
- ix. JOHN GOODYER, 1622, &c.
- x. JOHN TRADESCANT, the elder. Hatfield, 1611. Lambeth, 1629-34.
- xi. GEORGE GIBBES, ? and 1634.
- xii. WALTER STONEHOUSE of Darfield, 1640-4.
- xiii. EDWARD MORGAN of Westminster, 1662, and HUGH MORGAN, 1569-87.
- xiv. ROBERT MORISON of Blois, c. 1651; of London, 1661-2.

The earliest English garden list that has been published is that of the Holborn garden of John Gerard, which, first printed in 1596 and again in 1599, has been re-edited in a valuable form by my friend Dr. Daydon Jackson in 1876. One of the first results of the re-examination of the Goodyer books and papers was the recognition of one of his manuscripts as a very early list of the garden of the Rev. Walter Stonehouse at Darfield Rectory in Yorkshire, 1640-1644. By the courtesy of the editor of the *Gardeners' Chronicle*, this was printed in full in the numbers of that Journal for May 15, 22, 29 and June 12, 1920. No sooner had this list been identified than in the same collection of papers, several other garden lists were found that successively antedated the Stonehouse list and one another, and finally gave the clue to the finding of the list of a garden that is probably older than that of Gerard.

First in importance is the unique copy of the printed plant list of the Lambeth garden of the elder John Tradescant, in which more than 750 plants are named. It is the only work known to have been printed for the author in his lifetime: it is dated 1634.

Then were found short lists of the gardens of John Franqueville and George Gibbes of Bath, both extending the list of plants

already known to be grown by them, and the highly interesting lists of the gardens of William Coys of Stubbers in Essex and of John Parkinson in Long Acre. Lastly, the mention of the name of Richard Shanne has led to the rediscovery of his, the oldest garden list of all that are still unprinted, in the British Museum.

The Goodyer MSS. have thus provided a richer store of definite horticultural facts relating to special English gardens, and dating from the first half of the seventeenth century, than any that has yet been published. To this material we have added a few other contemporary notes and lists, not previously published, which we came across when searching for matter relating to the early annals of English horticulture.

i. OXFORD AND WINCHESTER GARDENS, 1570-2.

The following notes of plants growing in gardens in Oxford and Winchester are written by a sixteenth-century botanist in his copy of Du Pinet's *Historia Plantarum*, now in the library of the Botanical Department of the British Museum.¹ When no locality is mentioned, we provisionally assume the garden to have been in or near Winchester. In one case only is another county mentioned: the Olive, 'Olea sativa', p. 81, 'at belnys nothers in Suffolk (that is not wild)', showing that in spite of Tacitus's adverse opinion of our climate the olive was being grown in England a full quarter of a century before the date of Gerard's garden list. We have suggested that the writer may have been Dr. Walter Bayley of New College, p. 235.

HENRY CROSSE'S OXFORD GARDEN, 1570.

The friend of the unnamed botanist may be identified with Henry Crosse, Bedell of Theology, who was Registrar of the University from 1566 to 1570. His house and garden are of great historic interest, because there in after years (1654-68) Robert Boyle lodged, had his laboratory, and invented his famous air-pump.² There on the south side of the High Street, and not far from the quarter of the ancient Apothecaries in Oxford, Henry Crosse cultivated simples which may have been of value to his successor (and ? descendant), Crosse the apothecary,³ whose drugs

¹ See p. 235.

² An engraving of the front of Crosse's house is reproduced in Gunther, *Early Science in Oxford*, 1920, p. 11.

³ Perhaps we owe the first evidence of the Lily of the Valley as an Oxfordshire plant to Crosse. 'Lilly of ye Vallies, Crosse, ye Apothecarie, had a basket full of ye flower. They grow about Stokenchurch.' John Ward's *Diary*, 1665. (Information from Sir D'Arcy Power.) The first *printed* 'record' for the Lily of the Valley in Oxfordshire is 'Blackstone, 1746'.

and scientific entourage doubtless proved no small attraction to Robert Boyle.

Our author mentions the garden thrice.

Hedysarum aut Securidaca, *Du Pinet*, p. 402.

Coronilla varia, L.

‘In Crosses Garden at Oxford in 1570.’

‘In Mr. Watson’s booke.’

Aethiopsis, p. 557.

Salvia Aethiopsis, L.

‘In Crosses Garden at Oxford.’

Scorpioides, p. 639.

Ornithopus scorpioides, L.

‘At Oxford in Crosses garden 1570.’

[JOHN] WATSON’S WINCHESTER GARDEN, 1572.

Six of the plant-records are associated with the name of a Watson, who I think must be the John Watson (1520–84) who took an M.D. degree at Oxford, and began life practising as a physician. A native of Evesham, he became a Fellow of All Souls in 1540. ‘At length, about the time Queen Elizabeth came to the throne, if not happily before, he entred into holy orders, was made prebendary of Winchester, archdeacon of Surrey, and Master of the Hospital of St. Cross near Winchester’ (1559). In the fifteenth year of Queen Elizabeth, 1572, the date of our notes, he was made Dean of Winchester. He evidently kept up his early interest in medicine, for in 1575 he was admitted to the degree of Dr. of Physick. He became Bishop of Winchester in 1580.

Some of the notes refer to a ‘booke’ which we take to be a *Hortus siccus*, which may have also included wild plants. His garden is likely to have been at St. Cross.

Myrica s. Tamarix, *Du Pinet*, p. 55.

Tamarix gallica, L.

‘In Mr. Watson’s booke and garden 1572. Of this there is said to be 2 kyndes, the greater bearing fruit like to the lesser oake apples or gâules, the other bearing grayeish leaves without fruite.’

Ornithogallum, p. 207.

Ornithogalum umbellatum, L.

‘In Mr. Watson’s booke.’

Gratiola, p. 290.

Gratiola officinalis, L.

‘A kynd of the lesser Centaurie in Mr. Watson’s and S. Cross, 1570.’

Althaea Ibisc, p. 420.

Althaea officinalis, L.

‘Mr. Watson’s garden.’

Althaea sive Bismalva, p. 421.

Malva Alcea, L.

‘Mr. Watson’s garden.’

Centaurium maius, p. 270.

Centauria Centaureum, L.

‘In Mr. Barnaby’s garden and Mr. Watson’s booke.’

JAMES’ WINCHESTER GARDEN.

Cornus, p. 108.

Cornus mas, L.

‘In Mr. Jeames garden at Winchester.’

BARNABY'S GARDEN.

Cicer, p. 136.	<i>Cicer arietinum</i> , L.
‘Chiche. Mr. Barnbyes garden.’	
Centaureum maius, p. 270.	<i>Centaurea Centaureum</i> , L.
‘In Mr. Barnaby's garden and Mr. Watson's booke.’	
Acanthus, p. 277.	<i>Acanthus mollis</i> , L.
‘Mr. Barnbyes garden.’	
Ammi, p. 329.	<i>Ammi majus</i> , L.
‘Mr. Barabyes garden.’	
Alchimilla, p. 571.	<i>Alchemilla vulgaris</i> , L.
‘Mr. Barabyes garden.’	

HEYDEN'S GARDEN, 1570.¹

Aristolochia longa, p. 268.	<i>Aristolochia longa</i> , L.
‘In Mr. Heden's garden.’	
Stachys, p. 366.	<i>Stachys germanica</i> , L.
‘In Mr. Heiden's garden 1570.’	
Hypecoum, p. 410.	<i>Hibiscus trionum</i> , L.
‘In Mr. Heidens garden.’	
Mandragoras, p. 512.	<i>Mandragora officinalis</i> , L.
‘In Hedans garden.’	
Galega, p. 584.	<i>Scandix Pecten-Veneris</i> , L.
‘Sheperds neydle. In Mr. Heidēs garden.’	
Tithymalus cyparissias, p. 609.	<i>Euphorbia Cyparissias</i> , L.
‘In Mr. Heidēs garden.’	

THE GARDENS OF NORTON, WALLOP AND BASKET, AND THAT OF DUMMER RECTORY.

All names are well known in Hampshire. The Nortons owned land in Nutley, which is near Dummer.

Anemone altera, p. 250.	<i>Anemone sp.</i>
‘In Norton's garden.’	
Trifolium odoratum, p. 372.	<i>Melilotus officinalis</i> , L.
‘In Mr. Basket's garden.’	
Antirrhinum, p. 579.	<i>Antirrhinum majus</i> , L.
‘At Dümer in the p̄stes garden.’	
Cupressus, p. 40.	<i>Cupressus sempervirens</i> , L.
‘Wallops Gardens at hāpton.’ [? Southampton.]	
[Sir Henry Wallop was a very important personage at the time.]	

ii. SIR JOHN SALUSBURY'S GARDEN AT LLEWENI, 1596. 1607-8.

The botanical records entered by Sir John Salusbury in his copy of Gerard's *Herbal* have already been described, p. 243. Two documents in Oxford mention the Lleweni gardens in the seventeenth

¹ A Benjamin Heyden of Hants matric. at New College in 1586, and became a master at Winchester College.

century, and it has been suggested that a passage in a poem by Salusbury's friend and 'Court poet', Robert Chester, described the site.¹ Elsewhere Chester records in verse a failure of fruit due to an unusually cold spring about the end of the sixteenth century.² The lines were written for *A merrimt of christmas at the house of the Right Worshipfull John Salusbury of Leweny, Esq.* After stating that the occasion was one on which 'we of Arcadia sometime frolique swaines'

'should heare present as newe yeares homely gift
peares Apples fild bieres or the hazell nut
or other fruite that this faire clymatt yelds
but nipping winter and a forward spring
blasted our trees and all our summer budds
whose blossomes should have yelded dainty fare'

the poet goes on to propose a hornepipe, songs, and a dance.

Sir John grew three novelties which he considered worthy of mention in his *Herbal*. Two of them he raised from seed, which he may have obtained from Gerard, and to his evident satisfaction they proved true to the pictures in the *Herbal*.

Helianthus annuus, L.

'This galant greate sunflower grewe in Sir John Salusbury's Garden at Llewene & cam to the full perfection of this portraiture the yeare 1607.

Datura Metel, L. The Smooth Thorn Apple introduced by Gerard from Robin.

'This faire herbe grewe to full perfection accordinge to the portraiture in Sir J. S. his Gardeyne at Lleweny in the yeare of our Lord 1607.'

Paris quadrifolia, L., he transplanted to his garden in 1608.

Marks in the margins of the book show that he was well acquainted with a large number of garden plants, but the only one that is

¹ Chester, *Love's Martyr*, p. 11:

Hard by a running streame or crystall fountaine,
Wherein rich *Orient* pearle is often found,
Enuiron'd with a high and steepie mountaine,
A fertill soile and fruitful plot of ground,
There shalt thou find true *Honors* lovely *Squire*,
That for this *Phoenix* keepes *Prometheus* fire.
His bower wherein he lodgeth all the night,
Is fram'd of *Caedars* and high loftie *Pine*.

An ancient well-head is one of the antiquities at Lleweni, and there still remains the stump of a very old cedar that is figured in one of the engravings of the old Hall near the artificial lake.

² It has already been noticed that the weather about this time was most unpropitious. 'For a series of years, wet summers had raised the price of corn, and in 1596 wheat in London reached the famine price of £5 4 0 per quarter; this too when the purchasing power of money was fully six times its present value.' Jackson, *Gerard's Catalogue*, vii.

further described is a variety of Clove Pink: 'At Seaton in Northumberland at the House of the Right Worthy Sir Ralphe Delavale¹ is a Clove Gilloflower almost perfecte blacke.' The production of artificial varieties among Gilliflowers was known to Shakespeare, see p. 61.

Sir John's garden continued to flourish long after his death. I have recently found unexpected evidence as to its state in the year 1680 in a letter contained in the *Hortus siccus* of Edward Morgan in the Bodleian Library (Ashm. MS. 1797). This collection of plants seems to have been begun in 1672, and to have received additions until 1682; unfortunately the plants, which are arranged alphabetically are not localized. The collection includes specimens of *Vicia Bathoniensis Goodyeri* and of the four species of Elms described by him.

The letter is addressed

ffor Mr. Edward Morgan living att

Bodesclen, theis deliver with a Basged with Care.

MR.

MORGAN my humbell servis to you returning you humbell thanks for my plants & seeds I have send you theis seeds folwing which I thought would bee exceptabell to you. And I have send you what plants as I know of what you have send for and if you have a mind of any thing that you know to bee in Leweny gardens I shall bee redy to serve you in other plants or seeds; and so with my humbell serveis to you I rest and remane to serve you whilst I am

THO THORNES.

Lleweny 20th day of Sept. [c. 1680]

Dubell wh popy	Littell snipt Canpion
evear lasting pece	Venus Lookinglas
Dubell throtwort	Pollyanthus out of ye Litell Garden
Sweet William out of ye potts	ffenill flower
Jarman Cachfley	Marsh mallows
Pollyanthus	harey sen Johns wort
Scarlett Popys	hashed fish
Ross Larkheell	Sweet Scabious
Bedford pinke	dubell Collanbine
2 faces under a hood	Vulnearia for yor oneld (oueld?) Cock
blew bodell	

I pray if you have any of theis plants to send mee a few of

Sanicell	Moon wort	Spare wort
Parsly Ston broke	Cotton tree	Laser wort
Barren wort	Balm of Gilead	Iron wort
As bodill	Lavender Cotton	Mony wort
Anons	Claver	Moth wort
Fraxsenella ore Dittany	Corall wort	Pony wort
Dragon wort	Cross wort	
Glass wort	Fig wort	

Mr. Harisson presents his servis to you and Will. Tomas & Coocke.

[At end Morgan's List of Plants in the Appendix to his *Hortus siccus*.]

¹ Craster, *Hist. Northumberland, Tynemouth*, describes his coal-mining.

At the present day, alas, the splendours of Lleweni are no more. The far-famed agricultural wealth of the Vale of Clwyd has not proved sufficient to maintain the larger country houses of the Tudors, let alone their gardens, and the great house of the Salusburys is now only represented by a couple of large farm houses, the one in the stables and the other in the offices of the old mansion. A huge walled orchard still remains, though the trees have run wild and its high walls have been partly removed for building material; and there are scattered fruit trees in the woods.

The original flower gardens seem to have been altered before the demolition of the house, in accordance with the cult of the 'landscape' inculcated by the eighteenth-century school of Kent and Capability Brown, while the architecture of the house was being remodelled in the style of Adam. The present tenant, Mrs. Roberts, informs me that the present garden is largely within the lines of the foundations of the old Hall, but that all tradition of the original gardens has been lost.

iii. DR. SIMON FORMAN'S LAMBETH GARDEN, 1608.

The strictly scientific character of Goodyer's writings, revealing the practical attitude of his mind, is remarkable in an age when so many of his 'scientific' contemporaries were seduced by the specious promises of astrology, alchemy, and divination, legacies from Arabian civilization. The astrological gardening of the time is illustrated by a fragment from the diary of a former member of Magdalen College, which will at any rate be of interest to those who make a detailed study of the topography of London.

Thurneiser, Winckler, and Culpeper have all emphasized the importance of gathering roots and 'enchanted herbs' under particular stars, but Forman evidently believed in the influence of the moon on planting as well.

The first entries evidently refer to the acquisition of the site.

1596. 29 Jan. at 4 I went first to see the garden. On 30 Jan. . . . I went to spek with Mr. Katerins for his garden.

1597. This yere between Michelmas and Christmas I toke Lambeth House and entred yt a moneth after Christmas. Lambeth House at Westminster, *nota*.

1607.

15 July at 7 p.m. I began to cut and bush up the garden of Roses at Lambeth Marsh¹ and I began in the north west part and made som dossen bushes or better.

30 Aug. I put the gilliflowers in the flower (?) garden.

¹ Lambeth Marsh was due east of the end of Westminster Bridge. Seven houses only are shown there in a map of 1560.

Trees planted in 1608.

Anno 1608 the 29 of November post mer. I set the framboy trees under the west pall all along at Lambeth Marshe post mer. at 30 p. 3 in Hauflids garden.

The 7 dai Decemb. ♀ I set the willowes all alonge towarde the Lane, and at afternon at 33 set the first rose tre at the bankes end toward the door by the Lane (in ♂ ○ ω 4

The 8 dai 7 p.m. at 2, I set the rest of the rose tres all along the bank towards the Lane (in ♂ . . pt a ♀ ap Δ ♂ in ✕ & 6 more rose tres on the bank at Davies syd toward the lane.

The 9 dai ♀ I set all the other rosse . . . toward Davies syd from the pryvi Doun toward the Lane but only 6 which wer set the night before those that stand towards Davies syd towards the Lane.

Also the 9 dai December I set the 6 philberts and the 2 pech trees and they stand right against the postes at the ends, vz. toward the lane & towards the hous. The camonill on the banck by the pryvi and the premrose were set the 9 dai at 3 p. merid.

The 16 dai of Decemb. ♀ p.m. I set the roses and 2 apple tres all along by the palle toward Hammons garden & Mr. Walters garden except those west the arb exopt to Mr. Walterers garden (in 17.

The roses by the gose berryes and all the small roses under Waterers pall wer set 1609 the 4 of January a.m. between 9 & 11 of the pmeks (?).

The first beanes I set this year 1609 wer set the 7 dai of January ½ . . . between 8 & 9 & they wer set in Haufilde garden all along from the corner of the pall from the pryvi by the Raspis & so along by John Davies garden to the lane, & all along the banck by the lane in both gardenes, & they wer watered 24 howares befor, 4 daies befor the (was at full.

[MS. Ashm. 2402, f. 37.]

iv. LORD SALISBURY'S GARDEN AT HATFIELD, 1611.

See below under J. TRADESCANT.

v. RICHARD SHANNE'S GARDEN LIST, 1615.

See p. 264.

Richard Shanne of Woodrowe near Methley in Yorkshire was planting woods and orchards as early as 1577. The entries in his diary come down to 1617, so that it is exceedingly probable that many of the plants named in his list were being cultivated by him long before the date of the writing out of the list. His observations on the weather are of great local interest.

THE NAMES OF ALL THE HERBES

*growinge together at this daie in my Garden,
and fyrst of the plants called gramen striatum.*

RICHARD SHANNE. 1615.

Ladies Girdle.
Blewe flower de luce.
Purple " "
Whit " "
Narrow leaved flower de luce.
An other flower de luce.
Velvet " "

Bulbed blewe flower de luce.
Bulbed Chaungable flower de luce.
An other bulbed " "
Stinkinge Gladon. " "
Kings Speare.
Gladeolus cleare red.
" way red.

- Yellowe Lillie with — blades.
 Day " "
 May " "
 Whit " "
 Great red Lillie.
 Small red "
 Pearle red "
 Calcedonian Lillie.
 Mountaine "
 Swete yellowe "
 Imperiall "
 Hyacinthus Whit flo:
 " purple.
 " Ash cullour.
 " Skye collour.
 " Botroides.
 " bush topp.
 Narcissus yellowe Circle.
 " redd Circle.
 " doble flowers.
 " yellowe flowers.
 Tulipa wth yellowe flowers.
 Tulipa precox with yellowe flowers.
 " with flowers redd & yellow.
 " bright red flowers.
 " darke red flowers.
 " whit flowers.
 " apple blome collour
 and thirtie or fortie more
 varieties.
 Fritillaria Chekered Daffodilly.
 Doble Daffodillie.
 Snowe Dropps.
 Saffron three kynds.
 Starr of Jerusalem.
 Starr of Bethlem.
 Scodoraprasum.
 Serpents Molie.
 Raphanus.
 Dittander.
 Tarragon.
 Succorie.
 Endive.
 Orach whit.
 " redd.
 Winter cherries.
 Yellowe henbane.
 Doble poppie divers kynds.
 Anemone scarlet coullour.
 " blewe flowers.
 Munkes Ruberbe.
 Bistorte.
 Scurvie grasse.
 Adders tounge.
 Herbe Trewlove.
 Lunaria minor.
 " maior.
 Rose plantaine.
 Goulden Rodd.
 Sopewort Gentian.
 Bell flowers.
- Dames violets.
 Doble campions.
 Flower Constantinople.
 Lysimachia blewe flower.
 " purple flower.
 " yellow flower.
 Thorowe leafe.
 Tutsan.
 Time of Candie.
 Yellow Time.
 Winter Savourie.
 Sommer "
 Dodder.
 Whit Hyssope.
 Yellow "
 Greene "
 Lavander gentle.
 " spike.
 Maile Lavander.
 Gilliflowers manie kindes.
 Sweet Williams.
 London Tufts.
 Thrift.
 Sneese wort.
 Gromell.
 Everlastinge.
 Cost Marie.
 Maudlaine.
 Doble Tanseie.
 " feverfewe.
 Germander.
 Sweet marjerome.
 Pott "
 Yellow "
 Pennie Riall.
 Mints divers kynds.
 Calamints.
 Balme.
 Motherwort.
 Avicen's Agrimonie.
 Sawwoorte.
 Betanie wth whit flowers.
 Mountaine Scabious.
 Great blewe Bottell.
 Marigoulds all the kynds.
 Sage divers kynds.
 Claray.
 Oculus Christi.
 Doble Cowslipps.
 Double Primrose.
 Bears eare.
 Spotted sanicle.
 Sanicula alpina.
 Ladies mantle.
 Enula campana.
 Borage wth whit flowers.
 " blewe "
 Comfrey wth whit "
 " purple "
 Cowslipps of Jerusalem.
 Dragon.

Assarabacca.	Clounes Allheale.
Whit Double Violets.	Lovage.
Blew " "	Corianders.
Ladies Bower or Virgins Bow.	Smallage.
An other Virgins Bower.	Alexanders.
Periwinkle w ^h whit flowers.	Fennell (giant).
" purple flowers.	Dill.
" blewe flowers.	Chervill.
Asclepias.	Myrhis or Sifilis.
Salamons seale.	Spjgnel.
Double holihoeks sundrie collours.	Filipendula.
Single " divers kynds.	Setwall Valerian.
Mary mallows.	Grekish "
Musled Cranes bill.	Larkes peelee redd. white & blew.
Double Gouldins.	Columbines all collours doble.
Globe flowers.	Romaine wormwood.
Wolves baines two kynds.	Oke of Jerusalem.
Hellibor Blacke.	Lavander Cotton.
Bastard Hellibore.	Asparagus.
Double peonie redd.	Madder.
" " white.	Carduus benedictus.
Female " the lesse.	Ladies thistle.
" " " great.	Artichokes.
Maile "	Cirsium.
Angellica.	Kidnie beanes.
Bastard pellitorie.	Lupines.
Herbe Gerard.	Rewe.

PLANTS OF WODDIE SUBSTANCE

Red Rose.	Yellowe Goosberies.
Whit "	Red Goosberies.
Damask Rose.	Savey Tree.
Provence " "	Tamariske.
Yellowe "	Privitt.
Cynamon "	Chesnut Tre.
Muske "	Sycamor Tre.
Gelderland "	Quince Tre.
Rosmarie.	Medler Tre.
Woodbine thorow leafe.	Mulberie Tre.
Lencholans.	Job's teares.
Pescod tree.	Firr tre.

vi. WILLIAM COYS'¹ GARDEN 1604-1616.

William Coys' garden at Stubbers in Essex has already been described, p. 17. It will always be remembered in horticulture as the classic locality where the *Yucca* first flowered in England,

¹ Mr. C. B. Russell has very kindly supplied extracts from the Parish Register of North Ockendon which show that William Coys was buried on 9 March 1627, having lost his wife Mary ten years previously. His tombstone which is not now visible, is stated in the *Essex Archaeological Society's Journal*, to have borne an inscription, the dates of which do not agree with those given in the Register. It probably ran as follows:

Here under ly buried the Bodies of William Coys of Stubbers in this parish Esquier, who decessed the 6 day of March 1627 and of Marie his wife, the second daughter of Giles Aleyne of Hasleigh Hall in the County

and whence Goodyer obtained so many Spanish plants from the stock recently imported by the traveller William Boel. We now know that it was the station whence the newly imported Ivy-leaved

Nona gloriose clara & opinata YVCA.



Statim Clarissimus & piiffimus V. D. Guil. Coys

COYS' YUCCA, AFTER LOBEL, 1605.

of Essex, Esq who decessed ye 13th Day of March 1617 who had issue 8 sonnes and 6 daughters

And here 's the worst that envious Death could doe
Let loose two soules that long'd to Heaven to goe.

Of his fourteen children, Matthew, bapt. 1594, bur. 1595; Daniel, bur. 1595; Maralah, b. 1596; Matthew, bapt. 1596; and Sylvanus, bur. 1613, are all mentioned in the Register. Giles Coys, whose children were born between 1627 and 1632, was the eldest son. The signatures of both William and Giles Coys are written on the title-page of a copy of Daléchamp's *Historia generalis Plantarum*, 1587, now in the Sherardian Collection in the Botanic Garden at Oxford. Both volumes are inscribed '*Liber Gulielmi Coys, ab M^{ro} Rimee Junii* 18, 1604, *nunc vero Aegidii Coys Sept. 4, 1627*'.

By inquiry at Somerset House I found his will (P.C.C. 37 Skyenner). It was dated 8 Dec. 1618 and proved 2 April 1627, Giles Coys being named as sole executor.

Toadflax spread over England, where Goodyer learnt about a new species of Elm, saw many other plants for the first time, and whence he stocked his Hampshire garden at Droxford.

Yeast and Beer.

If in 1515 so supreme a genius as Leonardo da Vinci did not despise the petty details of the preparation of his national drink, and wrote letters thereon deprecating a bad habit of making wine in uncovered vessels, so that the essence escaped to the air during fermentation, surely a scientific history of cultivators of plants should take cognizance of the first advances in practical brewing. With the name of Coys may be associated some of the earliest recorded experiments of an exact nature in what is the most fundamental operation in the whole of Biochemistry—the production of alcohol from malt. Beer has not always been the national drink of the Englishman. Only a very few years before Coys' childhood, an English author on *Dyetary* thought it necessary to define in print what this new drink, beer, really is. 'Bere is made of malte, of hoppes, and water: it is a naturall drynke for a Dutche man. And nowe of late dayes it is moche used in Englande to the detryment of many Englysshe men.'¹

The older recipes give hardly more detail than this.² Without knowing of its existence as a plant William Coys had made a detailed study of the culture of the Yeast plant, the results of which he communicated to Lobel, who printed them in 1605.³ It is clear that he knew from experience that a small rise or fall of temperature would profoundly alter the working of yeast, and that he introduced improvements in the brewing of beer. Coys may not improbably have received his first lessons in the art of brewing from persons who could recall a time when no hop-brewed beer was made in England, for hops, it is believed, came to England from Flanders on one and the same ship with 'peacocks and heretics', or, more precisely, between 1520 and 1524. An account of brewing as practised in France had recently been published in 1600 in Surflet's translation of the *Maison Rustique* by Charles Stevens (Estienne), but the use of yeast is not as well

¹ Boorde, *Dyetary*, x. 256. 1542.

² Possibly Dr. Walter Bayley's MS. entitled '*Explicatio Galeni de potu . . . et praecipue de nostrae Alae et Biriae paratione*', might be helpful, but though said to have been in the Library of Robert, Earl of Aylesbury, it is not now to be found' (D'A. Power).

³ Lobel, *Adversaria pars altera*, 1605, pp. 471-2.

explained as in Coys' account. Owing to the improved methods of brewing advocated by Coys and others, English Beer made from Barley and Hops, became 'famous in Netherland, for England yields plenty of Hops'.¹

The following is a somewhat free translation of Lobel's Latin version of Coys' recipes.

The fullest and most precise directions in the English language for the brewing of the most agreeable and wholesome of all German & English Cerevisia or Beer, that is at once most suitable for use in warm countries, received from the illustrious Mr. William Coys, the highly skilled Botanist.

Rec. Heat 40 gallons of river water, but do not let it boil. Draw off a third part into a tub and soak in it one measure of the best malt, crushed soft, for a whole hour or longer, but without heating. At the end of this time the liquor, known in England and Belgium as *Wort*, is strained into a suitable vessel by withdrawing a spigot from a hole at the bottom of the tub. While the straining is in progress, heat another third of the water, as described, and pour it on the liquor that has been strained for an hour.

But if you wish to brew a larger quantity of rather stonger beer, pour the second part of the hot water on to the strained malt and let it infuse for an hour or an hour and a half. At the same time heat the strained wort first prepared with 4 ounces of flowers of Hops—and for the same time (1-1½ hours), then strain through a hair sieve and cool by pouring it into a tub. In this tub the wort must not be more than a foot in depth, though in winter it may be a little deeper. When it has got cool in summer (but luke-warm in winter), take up a fair quantity in a convenient vessel. and put to it half a pound or 8 oz. of the best barm of beer, called in English *Yeast*, in Belgian *Gyst*, and in French *Gy*. The yeast should be diluted and thoroughly stirred in another vessel or pitcher with three or four times its quantity of the strained *wort*. The pitcher is stood in the remainder of the wort in the tub, and the scum frothing up in the pitcher is allowed to overflow into the strained wort in which the pitcher is standing, so as to work up the whole of the strained wort into a state of boiling and fermentation. The second infusion of strained wort is then poured in gently, so as not to hinder the boiling. All this is done until the barm has risen sufficiently; it often rises a foot, and without reheating; and the beverage begins to smell like the strained liquor, but not bitter (for Cerevisia rapidly acquires a flavour of vinegar). Finally, the Cerevisia is kept in vessels that are filled and closed after the effervescence is over.

Light Beer. If you want a light beer (such as we usually find agreeable in summer), mix the first and last strained liquors: heat, add a sufficient quantity of hops, and proceed as aforesaid. Pour the third and last part of the hot water on the malt to yield the weakest beer of all: let it soak for two hours, and then proceed as before.

¹ Moryson, *Itinerary*, 1617.

Then follow recipes for making :

CEREVISIA MARTIA (March Beer) *invigorating, most nutritious, very lasting, keeping a year.*

ALE, *an agreeable drink, of the flavour of wine, as used by the first nobility of England.*

Coys' Garden 1604-5.

Several early printed references to plants grown by Mr. Coys show that he was the most enterprising amateur grower of his day of new and rare plants in England. His success with Primulas was praised by Lobel,¹ who noted three new kinds at Stubbers, '*Primula veris gemino flore, altero superiore luteo, alteri inferiore viridi innato*', two varieties of the '*Primula veris ex luteolo subviridi altius laciniata aut fimbriata*', and '*Primula veris flore viridi umbellifera*'. At that time, *c.* 1604-5, Coys had in his garden :

Allium silvestre perpusillum, Iuncifolium moschatum. Rec. from D. Leister² from Montpellier.

Bellis spinosa elatior et fruticosior Herbariorum.

Colchicum minimum tenuifolium Gallaeicum.

Crocus syl. Byzantinus serotinus candidus.

Fritillaria nigra, Pyrenaea. 1605.

Gladiolus minimus. flowered in May.

Narcissus Cyprius luteus polyanthes, flore pleno. Dec. 1604.

Introduced by Lete³ from Cyprus.

Parvulus Hyacinthus stellaris vernus. April 1605.

Received from George le Seigneur.

Yuca gloriosa. July 1604.

The English names in italics, taken from Gerard, are given in the case of those plants which can be identified as having been included in Gerard's garden list of 1599. The plants without English names, were presumably not cultivated by Gerard. The difficult task of the determination of the greater number of the species was undertaken by Dr. Daydon Jackson who in the most friendly way gave up a part of his Christmas holiday to the work.

¹ Lobel, *Stirpium adversaria nova, altera pars.* Lond. 1605.

² Perhaps Dr. EDWARD LISTER, *c.* 1556-1620. Physician to Queen Elizabeth.

³ NICHOLAS LETE a London merchant 'greatly in love with rare and faire flowers' traded in the Levant. In the Bodleian Library there is an *Account* of his, dated 1601-3, to Richard Sandy for £20 worth of drugs, including *Salsaperiglia, Sassafras, Rebarbe, Licoris,* and *Agarick.* There are notes on the ailments of himself and his family in the same book. One of the symptoms of his malady, recorded by his astrologer-physician, was that 'he picketh his nose'. He died 'about a quarter of a year after'. MS. Ashm. 181, f. 66 b.

Mr. Coys his Garden. 24 & 25 of March 1616-1617.

- | | |
|---|---|
| Thymbra Boelii. | <i>The herbe Masticke. Thymus Mastichina L.</i> |
| Cymbalaria italica. | [<i>Linaria Cymbalaria Mill.</i>] |
| Cochlearia mi. | <i>Cochlearia danica L.</i> |
| Chamaelia tricoccos. | <i>Widow waile. Cneorum tricoccos L.</i> |
| Cistus foe. | <i>Female Holly Rose. Cistus salvifolius L.</i> |
| „ ledon. | <i>Cistus ladaniferus L.</i> |
| „ mas. | <i>Male Holly Rose. Cistus parviflorus Lam.</i> |
| Lamium 2 Clus. | <i>Scrophularia vernalis L.</i> |
| Alaternus. | <i>Rhamnus Alaternus L.</i> |
| Cytisus maior semp. virens. | <i>Cytisus canariensis Steud.</i> |
| Arbor Judae cum flo. | <i>Judas tree. Cercis Siliquastrum L.</i> |
| Chamaedris laciniatis fol. | <i>Jagged Germander. Teucrium Botrys L.</i> |
| Eryngium Alpinum. | <i>Eryngium alpinum L.</i> |
| Echium flo. rubro. | <i>Echium violaceum L.</i> |
| „ angustifol. | <i>Echium creticum L.</i> |
| „ omnium maximum. | <i>Echium australe Lam.?</i> |
| Scrophularia lusitanica. | <i>Scrophularia frutescens L.?</i> |
| Heliotropium indicum. | <i>Helianthus tuberosus L.</i> |
| * Jacea capitulis hirsutis. | <i>?Centaurea nevadensis B. & R.</i> |
| „ „ albo flore. | <i>Centaurea scabiosa L.</i> |
| „ repens luteo flo. | <i>Great Knapweed. Centaurea solstitialis L.</i> |
| Caput monacorum. | <i>Cnicus eriophorus Roth.</i> |
| Melilot German alb. flo. | <i>Germain Claver. Melilotus L.</i> |
| „ Italica. | <i>Italian Claver. M. italica Lam.</i> |
| Gratiola. | <i>? Hedge Hyssope. Gratiola officinalis L.</i> |
| 2 Hieratium indicum belgicum an Pylosella Syriaca. | <i>Sweete wilde Horehound. Sideritis syriaca L.?</i> |
| Galeopsis Clus. | <i>Hungarie Dead Nettle. Lamium Orvala L.</i> |
| „ flo. rubeo Clus. | <i>Lamium var.?</i> |
| 3 Cattaria media. | <i>Nepeta Cattaria L.?</i> |
| Cynoglossum boeoticum. | <i>Cynoglossum pictum Ait.?</i> |
| Scorzonera latifolia. E.R. | <i>Vipers grasse. Scorzonera hispanica L.</i> |
| „ angustifolia. | <i>Scorzonera angustifolia L.</i> |
| Valeriana flo. albo. | <i>Centranthus ruber DC. var.</i> |
| Salvia indica. | <i>Indian white Sage. Salvia officinalis L.</i> |
| Tithymalus characias monsp. | <i>Spurge. Euphorbia Characias L.</i> |
| „ serratus. | [<i>Euphorbia serrata L.</i>] |
| Sesamoides argentea. | <i>Astrocarpus Clusii J. Gay?</i> |
| Teucrium arborescens. | <i>Tree Germander. Veronica Teucrium L.</i> |
| Stoebe Sarmantica altera. | <i>Great Silver Knapweede.</i> |
| | <i>Centaurea salmantica L., vel C. splendens L.</i> |
| Cnicus alter Clusii. | <i>Carduncellus caeruleus Less.</i> |
| * Jacea palustris lusitanica. n. d[escriptum]. ¹ | <i>Centaurea Seridis β maritima Lange?</i> |
| Phaleris bulbosa. | <i>Phalaris bulbosa L.</i> |
| Beros ex Hispa. g. Barbarae spec. | <i>Barbarea praecox R. Br.?</i> |
| Cirsium maius. | <i>Cnicus dissectus Willd., vel Cn. heterophyllum Roth.</i> |

¹ Dr. Jackson considers that this must be the 'Jacea marina' of Parkinson.

- Myagrurn thlaspi facie. *Erysimum cheiranthoides* L.
 Oenanthe apii folio. *Oenanthe pimpinelloides* L.
 „ & cicutae folio. *Hemlocke Dropwoort. Oenanthe crocata* L.
 Hysopus flo. albo. *White flowred Hyssope.*
Hyssopus officinalis L. var.
 Millefolium odoratum. *Achillea odorata* L.
 Carduus Aspho. radice monsp. *Cnicus pratensis* Willd.?
 Trachelium gyganteum. *Giants Throatwort. C. latifolia* L.
 Herba Doria. *Captain Doreas Woundwoort. Senecio Doria* L.
 „ „ altera Virg. [aurea?]. *Golden rod. Solidago Virgaurea* L.?
 Absinthie insipida. *Unsavorie Wormwood. Artemisia inodora* Mill.
 Oxalis franca. *Round-leafed Sorrell. Rumex scutatus* L.
 Selicio Italarum. Sea cabbage rom. *Brassica oleracea, var. sabellica* DC.
 Chondrus dens leonis . . . lob. [writing indistinct]. *Leontodon tuberosus* L.?
 Chamaemelum albo duplo. flo. *Double Cammonill.*
Anthemis nobilis L. fl. pleno.
 Betonica maior Danica. *Betonie. Stachys Betonica* Benth.
 „ „ albo flore. *? Stachys Betonica, Benth. var.*
 Colus Iovis. *Jupiter's Distaffe. Salvia glutinosa* L.
 Heptaphillon maius. *Alchemilla alpina* L.?
 Levisticum verum Gerrar. Siler mont. Lob. *True Lovage.*
Laserpitium Siler L.
 Geran. haematodes rubro flore Clus. *Storks bill.*
Geranium sanguineum L.
 Clematis surrecta Pan. Clus. *Ladies Bowre. Clematis Viticella* L.
 Libanotis. *Herbe Franckincense. Seseli Libanotis* Koch.
 Seseli. *Spanish Toothpikes. Anni Visnaga* Lam.
 Mentha Romana. [*Mentha spicata* L.]
 Ranu[nculus] globosus. *Globe Crowfoote or Locker gowlons.*
Trollius europaeus L.
 Anemone nemorum albo pleno flo. *Double Wild White Windflower.*
Anemone nemorosa L. var.
 Crassularia maior Hispanica. *Great Orpin. Sedum Telephium* L.?
 Ptarmica imperati. *Xeranthemum annuum* L.
 Arisarum latifol. *Broad leafed Friers Hood. Arisarum vulgare* Targ.
 Lactuca virgiana. *Lactuca canadensis* L.?
 Hypocoum Clusii luteo flore. *Horned wilde Cumin.*
Hypocoum procumbens L.?
 Rosa sine spinis. *Rosa alpina* L.
 „ semper virens. *Rosa sempervirens* L.
 Hepatica coerulea mult. *Blew Liuerwoorts.*
Hepatica triloba Chaix. var.
 N Asphodelus bulbosus maior. *? Bulbous Asphodill.*
Ornithogalum pyrenaicum L.
 Rhamnus Clusii. *Rhamnus Alaternus* L.
 Cyclamen Rom. } *Cyclamen europaeum* L.
 „ hederac. } *Sowbread with leaves like Iute.*
Cyclamen hederifolium Willd.
 Flos Adonis. *Adonis flower. Adonis autumnalis* L.
 [Melilotus Italica (erased).]

Pulmonaria maculata.	} <i>Cowslips of Jerusalem.</i>	<i>Pulmonaria officinalis L.</i>
„ non maculata.		<i>Scabiosa sylvatica L.</i>
Scabiosa montana.		<i>? Cerinthe major L.</i>
Cerinthe flo. rubro.		<i>Picea excelsa Link.</i>
Abies vel Biota.		<i>Atriplex Halimus L.</i>
Halymus surrect. Clusii.	<i>Bastard Sea Purslane.</i>	<i>Honisuckle. Lonicera alpigena L.</i>
Periclymenum rectum.		<i>Eruca sativa Mill.</i>
Eruca maior flo. albo.		<i>Laburnum anagyroides Medic.</i>
Laburnum.		<i>Argemone mexicana L.</i>
Papaver spinosum.		<i>White Henbane. Hyoscyamus albus L.</i>
Hyosciamus albus.		<i>Blew Willow herbe. Veronica spicata L.</i>
Lysimachia flo. coeruleo.	<i>Bastard Ginnie Pepper.</i>	<i>Solanum pseudo-capsicum L.</i>
Amomum Plinij.		<i>Bladder nut tree. Staphylea pinnata L.</i>
Nux vesicaria.		<i>Carduus lacteus.</i>
Silybum minus flore nutante (hanging downwards).		<i>= Silybum eburneum Coss. & Dur.?</i>

The lists next following are written on the same sheet of paper, though not under the same heading, so we cannot tell for certain whether the plants mentioned were cultivated by Coys or not.

- g.¹ Aster atticus. Some of them are like to *Hieracium*, nomenclature of Milk 392.
Blew Starwoort. Aster Amellus L.
- g. Hieracium atticus,² 1, 2, 3. have milk in y^e roots, whether all have or no.
Inula crithmoides L. ? var.
- g. Rapuntium. *Great Spanish Orpin. Sedum Telephium L. ?*
- g. Ammi vulgare. *Bishops tweede. Ammi majus L.*
- g. Blitum. *Amaranthus Blitum L. vel A. sylvestris Desf.*
- g. Argemone. 300. *Argemone mexicana L. ?*
- g. Pulsatilla. 304. *Purple Passflower. Anemone Pulsatilla L.*
- g. Virga aurea. 348. *Golden rod. Solidago Virgaurea L.*
- g. Perchepier. 453. *Parsley piert. Alchemilla arvensis Scop.*
- g. Colus Jovis. 627. *Jupiter's Distaffe. Salvia glutinosa L.*
- Barbarea. 188. *Winter Cresses. Barbarea vulgaris R. Br.*
- Battata virginiana. *Ipomoea pandurata Mey. ?*
- Pyrethrum. *True Pellitorie of Spaine. Anacyclus Pyrethrum DC.*
- Eriophorus. *Andryale integrifolia L.*
- Stramonium peregrini et Spinosum.

[*S. peregrinum = Smooth Thornapples, Datura Metel L.*]

[*List of Virginian Plants.*]

- Pishimon. *Prunus Virginianus. Diospyros virginiana L.*
- Maracocc. *Passiflora incarnata L.*
- Attamusco bulbus. *Zephyranthus Atamasco Herb.*
- Mosarus (?).
- Aquascomense inter asterem et iaceam. (*Some Composite.*)

¹ g. probably added by Goodyer to indicate plants described in Gerard's *Herbal*.

² Dr. J. notes that the false concord suggests that Goodyer must have written 'Hieracium' in place of 'Aster' from its yellow, not purple, flowers.

anapcor(?). Pepo.		<i>Cucurbita Pepo</i> L.?
Cerastus virginianus. ¹		<i>Prunus virginiana</i> L.?
Vitis virginiana.		<i>Vitis Labrusca</i> L.
Periploca.		<i>Gonolobus suberosus</i> R. Br.?
Sumac Anonymos virginianum.		<i>Rhus typhina</i> L.
Solanum virginianum. _____		<i>Solanum carolinense</i> L.?
Alopecurus altera maxima Anglica paludosa. 469.		<i>Polygomon monspeliensis</i> Desf.
Beane. 471.		<i>Vicia faba</i> L.?
Pabulus Hyacinthus stellatus vernus. 486.		<i>Starrie Iacint.</i>
		<i>Scilla bifolia</i> L.?
Nar. Cyprius. 49.	<i>Italian Daffodil.</i>	<i>Narcissus Tazzetta</i> L.?
Fritillaria nigra Pyrenea. 496.		<i>Fritillaria pyrenaica</i> L.
Bellis spinosa. 508.		<i>Chrysanthemum flosculosum</i> L.
T//acori Clu. 48.		<i>Yucca aloifolia</i> L.
Jucca.		<i>Yucca gloriosa</i> L.
Gladiolus minimus. 511.		<i>Gladiolus imbricatus</i> L.
		[MS. ff. 24 v, 25.

WILLIAM COYS' SECOND AND THIRD GARDEN LISTS.

Later lists in Goodyer's handwriting, dated 1621, 1622, probably include many of Coys' plants described by Goodyer either at Stubbers, or when they flowered in his garden at Droxford.

Yet another list of Coys' plants is contained in a comprehensive list of garden plants known to Goodyer, which we have printed in the form of an index. Coys' plants are marked with the letter c. 324 names are so marked. See p. 387.

In the 1622 list, plants already included in the 1621 list are omitted, but are marked with an asterisk in the 1621 list.

1621.	1622.
Acarna flo: rubro.	
Aethiopsis.	
Alsine maior baccifera.	Antirrhinum.
Arum quorundam.	
*Aracus maior Boet.	
Argemone Pavio.	
*Asperula flo: coeruleo.	
*Astragalus marinus Lusit.	
Astrantia nigra.	
Atractilis flo: luteo.	
Behen rubrum monspel.	
Beros ex Hispania.	
Beta marina.	
„ alba.	
„ rubra.	Beta candida et rubra.
Blattaria flo: albo et luteo.	Borago sempervirens.

¹ Cerastus, probably = *Cerasus*.

1621.

- Blitum spinosum Creticum.
 Brassica monospermos.
 Caltha silvestr. Baet.
 Capons taile gras.
 Caput monachorum.
 Carduus globosus.
 Cerefolium vulgare.
 *Cerinthe flo. rubro.
 *Christophoriana.
 Chrysanthemum inscript. Baeticum.
 " creticum flo: luteo.
 " tenuifolium Baet.
 *Cicer rubrum.
 Cicutaria marina.
 Clymenum Matthioli. Lathyrus.
 *Cochlearia Batavorum.
 Convolvulus coerul. minor Baet.
 Conyza odorata.
 Cucumer agrestis.
 Cyanus varius.
 " flo. albo.
 Delphinium elatius variorum colorum.
 Etrioselinum.
 *Eringium Alpinum flo. albo.
 Ervilia.
 Faba veterum.
 " foliis serratis.
 Flos Adonis.
 Flos Africanus maior multiplex.
 *Foenum graecum.
 Geranii Baet. spēs.
 *Geranium Creticum.
 Gramen cristatum Baet.
 * " lupuli glumis.
 *Heptaphyllon maius.
 Herba Doria Narbonensis.
 Hieratium stellatum.
 * " falcatum.
 " intibaceum.
 Hippolapathum rotundifol.
 Hordeum nudum.
 Hyosciamus luteus. Nicotiana.
 Hypecoum Clusii.
 Lagopus flo. ruberrimo.
 Lamium 2^m Pannonicum exoticum Clus.

1622.

- Botrys.
 Brassica multifida.
 Calamintha mont praestantior.
 Caucalis maior Baet.
 Centaurea ma us.
 Consolida regalis.
 Eringium Alpinum flo. coeruleo.
 Hedysarum clypeatum.
 Holyhock.
 Hormium flo. coeruleo.
 Hyssopus flo. albo.

1621.

1622.

Lapathum sativum Rhabarbarum monachorum.	
Linaria minor aestiva.	
Lathyrus Dumetorum Baet.	
,, flo. miniato.	
,, palustris Lusit.	
,, edulis Baet flo. albo.	
,, Baet. flo. coeruleo.	
,, aestivus flo. luteo.	
Legumen pallidum. vllissiponensis.	
Lens maior.	
* ,, minor.	
Lychnidis Baet. spēs.	
Lychnis Calcedonica flo: carneo.	
,, ,, flo. rubro.	
Lychnis coronaria flo: suaverubente.	
,, ,, flo. rubro.	
*Lysimachia virginiana.	
* ,, flo: coeruleo.	
*Malva flo. amplo Baet. aestiva.	
Medica maxima spinosa spēs.	
*Medica maioris Baet. spēs prima.	
* ,, ,, ,, spēs altera.	
Medica Anglica minor.	
Medica maior pericarpio plano.	
*Melilotus Italica.	
* ,, Indiae orientalis.	
* ,, Germanica.	
* ,, officinarum flo. luteo.	
*Melissa Turcica.	Millium nigrum.
Mill mountaine.	Millefolium odoratum.
Myagrum monospermon.	
* ,, Thlaspi facie.	Myrrhis.
Napellus vulgaris.	
Nasturtium hortorum vulgare.	Good neighbourhood.
,, crispum.	Nigella Damascena.
Oenanthe angustifolia.	
Orobus hebariorum receptus.	
Panax heracleum alterum.	
*Panicum.	
Papaver multiplex variorum colorum.	Papaver cornutum flo: rubro.
,, Rhoeas Baet.	
*Perfoliata.	
Petroselinum virginianum.	
Petum Indicum folio pene obtuso.	
*Phalaris maior.	
,, minor Baet. semine nigro.	Phalaris Baet.
Phu minus.	

1621.	1622.
Pisum maculatum.	Pimpinella Agrimoniae foliis.
„ quadratum.	
*Pomum spinosum flo. albo.	
*Psillium.	
*Ptarmica Imperati.	
Raphanus hortensis radice albo toto.	
„ niger	
*Rubia spicata.	
Scabiosa montana maior flo. luteo.	Sanicula guttata.
-Scorpioides Bupleuri folio.	
„ siliqua crassa.	
„ Matthioli.	
Scorsonera latifolia.	
*Securidica maior.	
„ minor.	
Silibum minus flo. nutante.	Smyrnum Creticum.
Sinapi sativum alterum.	
*Solidago Saracenica.	
Sonchus Africanus Boëlio.	Stachys genuina Gerardo.
Tanacetum inodorum.	Spina solstitialis.
*Thlaspi Drabae folio.	
Trifolium odoratum.	Redd storks bill.
Vaccaria.	Tanacetum incipidum.
Valeriana graeca flo. coeruleo.	Trachelium flo. coeruleo et albo.
* „ mexicana.	Tragopogon foliis laciniatis.
Verbascum 4 ^{tum} Matthioli.	„ flo: purpureo.
*Vicia Indica fructu albo.	Viola Mariana.
Vrtica Romana.	

136.

Lobi a Dnō Zouch.¹

1621 — 136

1622 — 70

—
206

Sent 23 sheets wherein are 103 descriptions.

¹ EDWARD LORD ZOUCH had a fine Physic garden at Hackney, of which Lobel had charge. He had travelled abroad and introduced to English gardens the Thorn Apple from Constantinople, the 'Small Candie Mustard' (*Iberis umbellata* L.) which grew in 'Austria, Candy, Spain, and Italy, in untoiled places and by high waie sides', and the Great Honie woort (*Cerintho major* L.) (Gerard, *Herbal*, pp. 207, 277, 431). Lobel tells us that his plants included 'Sedum Norvegicum minus', 'Linaria lutea minima altera', 'Betonica Danica maxima', 'Phaseolus . . . Indicus', and that in 1605 he had a 'new Physic Garden' in which (presumably) he grew 'Hyacinthus stellaris Bizantinus alter elegantissimus serotinus bullatus' from Constantinople, 'Moly luteum Botanicorum, fl. 1604' and 'Pancratium Indicum alterum vernum, fl. 1605', Lobel, *Adv. altera pars.*, pp. 467, 486, 502.

- Crocus luteus. 3 vars.
 Gladiolus minimus. May.
 „ flore albo. From J. Robin in 1601
 Hyacinthus minimus serotinus elegans.
 „ stellatus Byzantinus v. exoticus Someri.
 Iris bulbosa alba.
 Liliun Corona platycallos, sive laticaulis. 1600.
 Liliun montanum luteum. 3 vars.
 Liliun sylvestre—Martagon imperiale.
 Moly luteum botanicorum. Pyrenees. 1604.
 Narcissi toti lutei praecocis Septentrionalium.
 „ floris lutei multiplicis varietates.
 Narcissus Cyprius luteus polyanthes fl. pleno.
 Ranunculus tripolitanus.

20 Marcii 1617 in Franquevils.

[g. = In Gerard's Garden List.]

- Agnus castus. *Vitex Agnus Castus L*
 hellebo alb. ver. flo. rubente. (See below).
 g. Clem[atis coerulea surrecta] Pan[nonica Clusii]. *Bush Ladies Bowve.*
Clematis integrifolia L.
 Mart[agon] alb. flo.
 Phalan[gium] Allobrogicum. *Anthericum Liliastrum L.*
 Leucoium magnum.
 Hy[acinthus] lilij bulbus.
 Jesi luteum seu Trifolium fruticans. *Yellow Jasmine.*
Jasminum fruticans L.
 g. Anagiris. *Beane Trefoile. Anagiris foetida L.*
 Rhuum (?) plantaginis foliis albo fl.
 g. Tulipa pumila. *Tulipa sp.*
 g. Pseudo-nar[cissus] ampl. calice. *Daffodill. Narcissus.*
 g. Liliun conval[lidium] flore rubente. *May Lillie with red flowers.*
Convallaria majalis L. var.
 g. Lotus arbor. *Nettle Tree. Celtis australis L.*
 [Cf. p. 339.]
 Cerasus pumilus. *Prunus cerasus pumila L.*
 Seseli Peloponense alter.
 g. Condrilla coeruleo flore. *Sowthistle with blew flowers.*
Lactuca perennis L.
 Mentha Danica. *Mint. Mentha sp.*
 g. Ribes . . . ruber. *Red Corraus. Ribes rubrum L.*
 [MS. f. 25 v.]

Of the 8 plant-names marked F in Goodyer's List of Plants (1620?), the following are not included above.

- g. Chamaecerasus. *Dwarfe Cherrie Tree. Lonicera alpigena L.*
 g. Helleborus albus flo. dp. atrorubente. *White Hellebore with flowers of a dark red colour. Veratrum nigrum L.*
 Laurocerasus Clusii.

viii. JOHN PARKINSON'S GARDEN LIST, c. 1620.

There is no separate list of Parkinson's plants that is identifiable, but there are convincing reasons for the belief that Goodyer, in his comprehensive list of Garden Plants, p. 387, marked those which he knew to be growing in Parkinson's garden, or which he had obtained from Parkinson, with a capital P. 250 plants are so marked.

ix. JOHN GOODYER'S GARDEN LISTS.

None of the many separate lists of garden plants preserved among the Goodyer papers can be indubitably produced as that of his own garden at Droxford or at Petersfield. The long list in his handwriting which we print as an index includes names of many plants known to him, and not grown either by Coys or Parkinson, but we cannot be sure that he grew them himself. And the same criticism applies to several short lists of plants in his handwriting to which neither date nor locality is attached. Such are the lists on MS. 11, ff. 23, 26, 27, 28, 45, 83.

The following short list on f. 28 certainly seems to refer to his own plants.

my *Cameline similis* is *Leucoium sylvestre Clusii*.

my *Sium agrorum* is *Petroselinum macedonicum parvo semine*.

Esula rotunda.

Pomum spinosum flo. albo.

Drabis.

Phyteuma mons.

Thysselinum.

Buphthalmum, it came for *Millefolium rubrum*.

Stoebe.

Pseudodictamnus.

Oenanthe angustifolia.

Cochlearia minima.

& *altera*.

Digitalis ferruginea.

Galega.

minor flo. luteo, seed.

Helleborus niger.

Acanthus.

Scorzonera by *Sorbus*.

Mercurialis.

Arbor vitae.

Herniaria.

Geranium Romanum.

Stachys.

Daucus Hispanicus like wild chervile.

Trifolium fruticans.

Tabaco.

Blattaria purp.

Carduus globosus.

Iris Italarum.

Leucoium melancholicum.

[MS. ff. 18, 28.

And the list of 95 names dated 16 Januarii 1620 (MS. 11, f. 83) has every appearance of being a list of the more interesting plants in some private garden, with references to the pages in Gerard where they are described. This list includes 'Trinidado Tabacco'.

More probability attaches to two lists of plants dated 1621 and 1622 and ending with a list of seeds obtained from William Coys in the latter year. We believe that these, or a fair proportion of them, flowered in Goodyer's garden at Droxford, and were described by him then and there.

X. THE GARDEN LISTS OF JOHN TRADESCANT, THE ELDER,
d. 1638.

The Garden of Lord Salisbury at Hatfield, 1611.

In 1611 Tradescant was in the service of Lord Salisbury, who had planted an extensive vineyard at Hatfield on a property which had been given to him in exchange for Theobalds by James I in 1607. The site was on the north side of the River Lea, on a piece of ground sloping to the south, hedged in with privet and sweet briar. The vineyard had been stocked with some 30,000 vines sent by Mme. de la Broderie, wife of the French ambassador, 500 vines from the Queen of France, and the few Muscats mentioned below. In the winter of 1611 Cecil commissioned Tradescant to plant his garden with a selection of good fruit trees and flowers then in cultivation by the best Dutch and French growers. The original bills, still preserved at Hatfield, contain several items of horticultural interest.

John Tradescant his bill for Routes flowers, seedes, trees and plants by him bought for my Lo: [the first Earl of Salisbury].¹ 3 and 5 January, 1611.

Roots of flowers, of Roases and shrubs of strang and rare bought at Leyden in Holland £3 0 0.

	Number.	Price.	Total.
			£ s. d.
Anemones (C. Helin of Harlem)			5 0
Aprycoke, The whit	1	6s.	6 0
„ (the archedukes gardener called Peere Vyens)	10	2s.	20 0
Arbor vita trees (C. Helin)	2	6d.	1 0
[Cherries]			
Rathe ripe cherry trees (C. Helin)	32	4s.	6 8 0
Archedukes cherye (prob. of Peere Vyens)	12	1s.	12 0
Biggandres (Robyns)	24	2s.	2 0 0
Boores cherye, an exceedyng greatt cherye (Harlem)	1		12 0
Currants Great Blacke (Harlem)	12	1d.	1 0
„ Great red (C. Helin)	6	2d.	1 0
[Cypress] Sypris trees (? Robyns) ²	200	1s.	10 0 0
Echatega, double (John Jokkat)			
Frittelarias (C. Helin)	40	3d.	10 0
Fyg trees, Whit (Robyns)	2	2s.	4 0
Genista hispayca (Parrys)			2 0
Gilliflowers, Dubble whit stok } (? Robyns)			3 0
„ Other }			
Irys calsedonye John Jokkat }			
„ susyana „ }			2 0 0
Martygon pompone blanche „ }			
„ pompong orang coller „ }			
Medlar, Great M. of Naples (Brussels)	2	2s. 6d.	5 0

¹ Abstracted from the *History of Gardening*, 1895, by the Hon. Alicia Amherst to whom I am also indebted for the loan of the portrait blocks of Lobel and Parkinson. The items have been arranged in alphabetical order.

² Tradescant's son introduced the American Cypress from Virginia.

	Number.	Price.	Total.
			£ s. d.
Mulbery (C. Helin)	2	3s.	6 0
„ „ Blak (? Robyns)	17	2s.	1 14 0
Myrtill tre (Parrys)	7	2s. 6d.	17 6
Ollyander „	6	2s. 6d.	15 0
Orrang trees of on years growthe grafted „	8	10s.	4 0 0
[Peaches] Alberge (? Robyns)			
„ Male cotton peach „			
„ The Troye ? „	4	2s.	8 0
Pomgranat withe many other small trees at the root (Parrys)			6 0
Quince, Lion's (Brussels)	1	3s.	3 0
„ Rath ripe Portingall „	1	6s.	6 0
Rose, Province (C. Helin)	16	6d.	8 0
Tulips (Harlem)	800		4 0 0
Vines, Muscat (Master Robyns)			4 0

The plants were purchased in France and Holland. Cornellis Helin lived at Haarlem. 'Master Robyns' was Jean Robin (1550-1629), a famous botanist of Paris and first curator of the 'Jardin des Plantes': several of his introductions are mentioned by Gerard to whom he sent *Apocynum rectum* (*Marsdenia erecta*), *A. repens* (*Periploca graeca*), *Christophoriana* (*Actaea spicata*, L.), *Crocus luteus*, *Epimedium alpinum*, *Tropaeolum majus*, *Lepidium sativum*, L. var., *Geranium lucidum*, L., *Datura Metel*, L.; 'peere vyens' we have interpreted as the name of the Archduke's gardener; the name of Vines is still remembered in connexion with the Cambridge School of Botany, but in Tradescant's bill it may denote a particular kind of vine. A re-examination of the MS. might settle the question. Of Mr. John Jokkat we have no further information. A few of the plants grown in a French garden of the period are exquisitely depicted in an album recently exhibited in the S. Kensington Museum for its beautiful late sixteenth-century binding. The volume contains the name *de Morogues*, and may turn out to be a horticultural work of great interest.¹ The watermark of the paper is French, c. 1570.

To this first period may be referred the interesting collection of coloured drawings of fruit-trees, popularly known as 'Tradescant's Orchard', now in the Bodleian Library (MS. Ashmole 1461). It has been suggested that the artist was the Alex. Marshall mentioned in *Musaeum Tradescantianum*, p. 41. Among the names of the fruits, written by some person who was evidently quite as illiterate as the elder Tradescant, is an entry 'The Amber Plum which J. T. as I take it brought out of France and groweth at Hatfeld'. This shows that the writer of the names is not likely to have been Tradescant himself. The collection of trees illustrated, which are arranged in the order of fruiting, may have been similar to the selection chosen by Tradescant for Lord Salisbury: the pictures were certainly painted as a guide-book for the use of visitors to the garden; on the first page is written 'Heare by the figures you may finde each fruite'.

For the sake of facilitating comparison, we have printed the

¹ Since identified by Mr. S. Savage as the work of 'Jaques le Moyne dit de Morgues Paintre', author of *La Clef des Champs*, 1586.

names of the fruits in *italics* in the right-hand column alongside Tradescant's own fruit catalogue of 1634. See p. 343.

In 1618 Tradescant was engaged in 'A Voiaĝ of ambasad' to Russia with Sir Dudlie Diggs,¹ and had, botanically speaking, a most successful campaign. After rounding the North Cape on 6th July, he landed at Archangel on the 16th, and finding a 'bery growing lowe' (the yellow Cranberry), which was eaten by the people 'for a medsin against the skurbi', he proceeded forthwith to dry 'sume of the beryes to get seed whearof' he 'sent par to Robiens of Parris'.

On 20th July he had 'one of the Emporer's boats to cari him from Iland to Iland to see what things growe upon them', and there he found 'pinks growing naturall of the best sort we have heere in England, withe the eges of the leaves deeplie cut or jaged very finely', also the *Rosa Muscovita* that he grew in his Lambeth garden later. None of the other plants observed by him on the Rose Island or elsewhere can be proved to have been introduced to western horticulture by him at this time.

In 1627 Tradescant was botanising in the Island of Rhé, where he went as a member of the Duke of Buckingham's expedition,² and whence he obtained the 'greatest Sea Stocke Gilloflower' (*Matthiola sinuata*).

Lambeth Garden, 1629-33.

Thanks to Goodyer and Ashmole, we have unexpectedly full notes of the plants that the elder Tradescant grew in his garden at Lambeth from 1629 onwards. He appears to have kept notes of additions to the garden on a few blank pages at the end of his copy of Parkinson's *Paradisus*, printed in 1629. This copy was acquired by Ashmole, who also wrote in it, and it has recently been added to the Bodleian collections at a cost of £28.³

¹ MS. Ashmole 824.

² A MS. account of this adventure is in MS. Ashmole 824, ff. 187-192, immediately after Tradescant's autograph account of the Russian Expedition, so that it may also have belonged to him.

³ The volume has been described in the *Bodleian Quarterly Record* and in greater detail by Mr. Boulger in *A seventeenth-century Botanist Friendship*, *J. Bot.* 1918, p. 197. There is stated the evidence for the identification of the MS. additions as the work of Tradescant and Ashmole, but the quotations from them leave much to be desired in point of accuracy. *c*'s are often transcribed as *t*'s, *p* as *q*, 16 as i 6, &c. It should be mentioned that the entry of Elias Ashmole's monogram-signature in the body of the book is dated 1680.

Reseved since the Impression of this Booke.

- In primis.* Sittissos Amarantinum.
Barba Jovis.
Polygolan.
Digitalem lutem maior.
Frittillaria Aquitanica.
Rosam Vittriensem.
Cogciggrum Plinnii.
- from Morine.* The great whyt Renunculus single.
on other sort of whyt Renunculus single.
Renunculus *Drape de Argent.*
Anemone Duple Greene with a littill leafe.
A thrice fayer Duple Anemone whyt Anemone.
on other sort of Dubble whyt Anemone.
- from Mr. Robine.* Tulipa perte maior.
Cardinalis planta 2.
- from Holland.* on Aster.
on Vyola matronallis.
Plattanos arbor
Cogciggra or shumahat } from Mr. Humfry Slaynie.
- from France*
Robyne. Iris Affracanis.
Iris pertyca.
6 Anemones tenuifollio Duple.
4 latifollio Duple Anemones.
- from Mounser*
Robyne. on German Rose of Mr. Parkinson from Mounser Robine,
whiche is called Rosa Austriaca flore phenissio.
4 more Roses whereof Mr. Tuggy hathe two.
on strang vyene.
on Red Honnysoccle.
Two Irisses without name.
Arbutus slipes.
Tragacantha slipe.

Reseved in the yeare 1630 from forrin partes.

- on Narciss.
- from Constanti-*
noble. Sr
Peetter Wyche. on Ciclamen.
4 Renunculuses.
Tullipe Caffu.
Tullipe perte.
4 sortes of Anemones.

Reseved in the yeare 1631.

- W.* on Tulype called the Coronell & on of hir owne.
T. S. on Tulipe Brewer 3 collers sh welcom Hom Best Golyathe.
on Palmer more good Tulipes unknowne.
- Mr. Colfe.* on Tulipe Beau without a Circle.
Blanck swisant.
Unick De Armentier.
Mr. Groves olyas.
Hollias Beu.
- From W. Win.* Blienborgh Admirall of 3 collers olli van Dusport.
flamed Red & Whyt Crowne.
- Mr. Rene.* Two Holliasses.

Reseved in the yeare 1631 from Mr. Rene Morine.¹

- In primis.* Renunculus Asiaticus flore duplicē luteo.
Narcissus Jacobei, Narcissus indecus, Narcissus flore rubro.
Semper eternum flore luteum.
Geranium noctu odorato.

¹ Cf. p. 279.

From Bruxsells. 6 Hiasinthes.
 Narcissus medio luteo,
 „ Narboniensis.
 „ Mussart.¹

In the yeare 1632.

Lotus libica.
 Phillerea.
 Sarsaparilla.
 Smilex aspera.
 Lentiscus.
 Agnus Castus.
 Cittisis maranthe.
 Absinthum arborescente.
 Citisus panonicum Clusii.
 Pseudo Dictamnium.
 Vyburnum.
 Lauristinus folio glabro.
 Cistus folis chrispus.
 „ popelium folyo.
 „ mas.
 Cortusa Americana.
 Thimum verum verum Hispanicum.
 Hisopium tenuifolio.
 Tragacanthum.
 Amanker lobelii.
 Frutex Coronaria flore pleno.
 Mirtis florence.
 Sesely Ethiopicum.
 Caradathe Americana.
 Narcissus Tobago.
 Ornithogalum arabicum.
 Iris percicus.
 Absinthium umbelatum.
 „ folio lavendulum.
 Auriggunum verum Hihipanicum.
 Tumariscus Itallica.
 Lutea Creatica.
 Linaria odorata.
 Feratium Indicum.
 Serpentaria.
 Arum mius.
 Iris gloriossa.
 Coulchicum frittilaria.
 Fraxinello flore rubro.
 Fraxsanela minor.
 Herunde Hispanica.

In the yeare 1632.

Fraxinella flore albo.
 „ minor flore albo.
 Dronicum maior.
 Eupatorem Nove Belgicum.

In the yeare 1633.

Abrotanum unguentaria.
 Androsaca Mathioli.
 Renunculus Lusetanycus odorata luteo.
 Colis Jovis.

Chama Iredis 3 sorts.
 Iris Anglica variagata.
 Dentaria Herundelesie.
 „ trefolia.
 „ setfolia.
 Telethium maius & minus.
 Moluka.
 Sentaurum magnossi.
 Ciclamen flore albo.
 Hulimus.
 Geranium 3 spetius.
 Tithemali caratius.
 Pistolochia.
 Hiasithus flore albo.
 Chamalea tricockos.
 Tulipe Chistmaker.
 Tulipa se bloome.
 Tricolor Nomvull.
 Ratabet.
 The Lyon.
 Brandinburg
 Oudenard.
 De Turbone.
 Tulipe swice.
 Crowne.
 Canadense.
 Otho Demeine.
 Oliva Cappadocia.
From the frencheman.
 The whyt Crown Tulipe.
 Hiasinthus Brumalis.
 Narcissus totus albus.
 „ Constantinopolis.
From Mr. Ploves brother.
 Narcissus virginianus.
 „ totus albis.
 „ De Diverse specie.
 Millefolium flore luteum.
 Virga Aurea virgine.
from Brussells 16 tulipes.

Papaver Reas flore luteo, Radx papoti. (?)
 Pulegium servinom.
 Frutex Canadencis Epimedium folio.
 Aquilegi variagata rubro et albo.

Bellis maior Americanum Arboessente prolifra.
 Cepe Lobellii.
 Aspeck.

¹ Query David Mostart, mentioned by Clusius, *Cur. Post.*, p. 19.

Buglosa minor sempervirente.	Ciclamen Antiochum & withe them 2 others.
Fumaria arboessente flore luteo semper virescente variagata.	Narcissus Virginnianum.
Asarum maius Americana.	" Indicus squamosus.
Absinthium innodorum.	Cicmen flore pleno, albo et rubro.
Polygon Creticum verum luteo.	Denis caninis flore luteo.
Phalangium virginianum flore albo.	Frittilari flore luteo.
Scabiosa Alpina vera.	" Hispanica.
Galliosus panonica Clusii.	Hiasinthus Indicus tuberosa radice.
Teucrium arboessente.	Prumela flore flore purpureo.
Aposinum Americana foliis Ascle- poydes floribus purpureo.	Gladiolus Canadensis.
Viola luteo Americana arboessente.	Alowaye mucronata.
Tordillum maius sive Sesseli Crettica.	Mirtis flore pleno.
Alkamilla pese leonis.	Sedum arboessence.
Stelaria argentina.	Casie Quorundum Clusii.
Fillapedulla altra.	Hipericon arboessence.
Campanella lactencis piramidalis.	Mispilis arona.
Barba hersi Coronopi folio.	Piometum realie.
Cetterache.	Alipum motcsetie.
Millifolium flore luteo.	Annagiris feotida.
Cianus Constantinopilus.	Jugibie Arabum.
Cottila marinum.	Zisipha Capadocia.
Absinthium tridentinum.	Carobie.
Hisopium mirtifolio.	Limonium minus angusti folio.
Mar Rubium.	Mariaranum sempevirence.
Renunculus minor bulbose flore pleno.	Sabina baccisfira.
Sanicula guttata montana.	Phlirea lattifolio.
Aster 4 speties.	Sicorum grumosa radice.
Abrottinum altra.	Capris vera.
Cardus bulbosus monspellesis.	Gnapphalium marinum et Cotonuri vulgi sive Bumbax humlis.
Balsamum sive Osimum oderatum.	Martigon Canadensis.
Ciclamen vernale flore rubro odoratis- simum.	Saldanella Alpina.
	Philex bacciffera.

In 1634, the year of the printing of the Catalogue, Tradescant's Museum and Garden were one of the sights of London, and were described as such by Peter Mundy whose manuscript is preserved in the Bodleian Library.¹

'Having Cleired with the Honourable East India Company whose servant I was, I prepared to goe downe to my freinds in the Country.

In the meane tyme I was invited by Mr. Thomas Barlowe . . . to view some rarieties att John Tredescans, soe went with him and one freind more, where wee spent that whole day in peruseing, and that superficially such as hee had gathered together . . .

Moreover, a little garden with divers outlandish herbes and flowers, whereof some that I had not seene elsewhere but in India, being supplied by Noblemen, Gentlemen, Sea Commaunders etts. with such Toyes as they could bringe or procure from other parts.'²

¹ MS. Rawlinson A. 315.

² Hakluyt Society's reprint of Mundy's Travels.


The existence of a printed list of plants grown by the elder Tradescant in his Lambeth garden appears to have been quite unknown to bibliographers until the recataloguing of the Goodyer Library once more drew attention to this unique possession, which I had the pleasure of describing in the *Journal of Botany* in 1920. It is of course possible that the little book was never published, and that the copy in the Magdalen Library is to be regarded as a proof of a work that was never put into circulation. And this view derives some support from a glaring grammatical mistake on the title-page, which we reproduce. No printer of repute would have permitted such a blunder to issue from his press.



PLANTARVM
IN HORTO
 IOHANNEM TRADE-
 SCANTI *nascentium*
 Catalogus.

NOMINA
SOLVMMODO
Solis vulgata exhibens.

Anno 1634.



Abies.	Acanthus sylvestris.
Abrotanum mas.	" sativus, <i>id est</i> , Branca ursina.
" foemina, <i>id est</i> , Chamae	Acetosa Hispanica major.
cyperissus.	" Franca rotundifolia lobelii.
* Abrotanum montanum.	" Indica.
* Absinthium arborescens.	Aconitum coeruleum, <i>id est</i> , Napellus.
" folio Lavandulae.	* " lycoctonum; Luteum hiemale.
" marinum, <i>id est</i> , Seri-	Aconitum luteum ponticum majus.
" phium. ¹	" luteum ponticum minus.
" umbellatum Clus.	Acorus verus.
" vulgare.	Ageratum.

* *Plant names marked with an * do not appear in the later edition of the catalogue printed in the Museum Tradescantianum in 1656.*

¹ First sent over from L. of Rhé by J. Tradescant. *Ger. emac.* 1c99.

- Agnus castus, *id est*, Vitex.
 Agrimonia.
 Alaternus.
 Alcea Veneta, *id est*, Alcea vesicaria.
 Alkakengi.
 Allium sativum.
 „ maius (*z*) Scorodoprassum.
 * Alopecuros vulgaris.
 „ spica aspera.
 * Alsine repens maxima.
 Althaea arborea flore purpureo.
 „ arborea flo. alb., montis
 Olb[iae].
 Alyssum Clusii, *i.* Alyssum Plinii.
 * Amaranthus spersa Pannicula.
 „ tricolor.
 „ holosericus.
 Ammi vulgatius.
 Amomum Plinii.
 Amygdalus.
 Anagallis tenuifolia flore coeruleo.
 Anemone latifolia Pavot. ma. flo. plen.
 „ latif. Calcedonica flo. plen.
 * „ latif. la Bruyne flo. plen.
 „ latif. rosea flore pleno.
 „ latif. albicans flo. pleno.
 „ latif. coccinea flo. pleno.
 „ latif. potorine flo. pleno.
 „ latif. superisse flo. pleno.
 „ latif. Constantinopol. flo. plen.
 „ latif. aliae diversae species.
 „ tenuifolia rubro flore pleno.
 „ tenuif. albo flo. plen.
 „ tenuif. albo dilut. flore pleno.
 „ tenuif. mutabilis flore pleno.
 „ tenuif. viridis flo. pleno.
 „ tenuif. carnea vivacis. flo. pl.
 „ tenuif. scarlata flo. pleno.
 „ tenuif. *Pink colour*, flo. plen.
 „ tenuif. pl. Coma-amaranthina.
 „ tenuif. flore ple. roseo.
 „ tenuif. flore pleno variegata.
 „ tenuif. flore pleno purpureo.
 „ tenuif. flore pleno purpureo
 dilutor.
 Anemone tenuif. flo. simplici Brancion.
 „ flore simp. diversae species.
 Angelica sativa.
 * Anthora.
 Anthyllis leguminosa erecta flo: rubro.
 Antirrhinum maius flore albo.
 „ minus flore variegato.
 „ minus flore albo.
 * „ minus sylvestre.
 *[Apium, see Petroselinum.]
 Apocynum Americanum.
 „ alterum.
 Aquilegia variegata albo & purpureo.
 „ variegata albo & rubro.
 „ rosea.
 * Aquilegiae magna diversitas.
 Aracus Baeticus.
 „ clematites.
 Arbor vitae vel Thyia.
 „ Judae.
 Arbutus, sive Unedo.¹
 Aristolochia clematidis.²
 „ rotunda radice.
 Armeria holoserica.
 „ flore pleno.
 „ flo. simpl. magna diversitas.
 Arum sylvestre.
 „ maculato folio.
 „ maius.
 Arundo Hispanica, Donax.
 Asarum vulgare.
 Asclepias flore albo.
 „ flore nigro.
 Asparagus sativus.
 Asperula flore albo.
 Asphodelus minor, Clusii.
 „ Lusitanicus.
 Aster caeruleus serotinus fruticans.
 * „ alter minor fruticans & pre-
 cocior.
 Astragalus Baeticus.
 „ Marinus.
 Astrantia nigra.
 Atractylis.
 Atriplex baccifera maior.
 „ minor.
 Atriplicis varietates.
 Avellanae diversitates.
 Avena nuda.
 Auricula ursi flore albo.
 „ „ flore luteo maximo.
 „ „ folio glabro.
 „ „ folio luteo medio.
 „ „ albo & rubro variegata.
 „ „ albo & purpureo varie-
 gata.
 Auricula ursi flore holoserico.
 „ „ flore rubro.
 „ „ flore violaceo.
 „ „ flore fusco.
 „ „ holoserica, Potrine.
 „ „ altera, Potrine.
 „ „ maxima Tradescanti flore
 obsoleto.
 Auricula ursi diversae species.
 Balsamina foemina.
 Barba Jovis frutex.
 Barba Hirci Tragi.³

¹ Two trees, 'the largest I have seen', noted as still living in the garden in 1749 by Sir W. Watson (*Phil. Trans.* xlvi).

² *A. Clematidis* L. reported by Sir W. Watson in 1749, *l.c.*

³ 'Barba capri' was noted by Johnson in 1633.

- Be[he]n album.
 „ rubrum.
 Bellis maior.
 Bellidis minoris magna diversitas.
 Beta sativa.
 „ spinosa Cretica, Bauhinii.
 Betonica maior Danica.
 Bistorta maior.
 Blattaria flore luteo.
 „ flore albo & violaceo.
 „ maxima obsoleta.
 „ maxima odorata flo: luteo.
 Borago flore albo.
 „ semper virens.¹
 „ flore caeruleo minima.
 Botrys.
 Brassica marina latifolia.
 „ perfoliata.
 „ foliis crispis.
 „ striata.
 „ Hispanica.
 „ Sabaudica, varietas.
 Brunella flore albo.
 Bolbocastanum maius.
 Buglossa sativa.
 Buxus maior.
 „ minor.
 „ auratus.

 Calamintha montana praestantior.
 Calceolus Mariae.
 Calendula flore pleno.
 „ prolifera.
 Caltha palustris, flore pleno.
 Canna Indica flore rubro.
 „ Indica flore luteo.
 Cardamine flore pleno.
 „ trifolia.
 „ impatiens.
 Carduorum diversae species.
 Carduus globosus.
 * „ chrysanthemus.
 „ Benedictus.
 *Carlina, i. Chamaelaeon albus.
 Carobe Americana.
 Caryophyllata montana.
 Caryophyllorum elegantium magna
 varietas.
 Caryophyllus globosus latifolius.
 *Cassia quorundam, Clusii.
 Castanea equina.
 „ vulgaris.
 Centaurium maius fol. Helenii.
 „ maius flore luteo.
 „ alterum Clusii.
 Cerasorum diversae species.
 Chamae-cerasus.
 Chamae-irides variae.

 Chamelaea tricoccus.
 Chamaepitys secunda.
 Chelidonium maius.
 „ maius fol. quernis.
 Chondrilla.
 „ purpurea.
 Christophoriana.
 Chrysanthemum Creticum.
 „ segetum bellidis fol.
 Cichorium sativum.
 Cinara sylvestris Boetica.
 Cistus annuus Clusii.
 „ mas.
 „ foemina.
 „ ledum.
 „ flore albo.
 „ flore albo alter.
 „ foliis crispis.²
 „ ledum primum Clusii.
 „ ledum latif: secundum Clusii.
 „ ledum quartum Clusii.
 „ ledum septimum Clusii.
 „ Halimi folio.
 „ quintus Clusii.
 Clematis flore pleno.
 „ flore coeruleo.
 „ flore rubro.
 „ Virginiana.
 Cochlearia Batavorum.
 Coccigria.
 Colchicum atropurpureum.
 „ vernum.
 „ flo: albo.
 „ flore pleno.
 „ fritillariae facie.
 „ Bizantinum.
 „ variegatum.
 „ vulgare.
 „ ex Insula Chios.
 Colutea vesicaria.
 „ scorpoides.
 Coniza major vera.
 Convolvulus minor folio Althaeae.
 Coriandrum.
 Cornus.
 Cornus fructu rubro.
 „ fructu albo.
 „ sylvestris.
 Corona imperialis.
 Coronopus maior.
 Cortusa Matthioli.
 „ Americana.
 Crocus, Neapolitanus.
 „ flore albo.
 „ Maesiacus luteus.
 „ Maesiacus flore albo.
 „ flore luteo.
 „ violaceus maior.

¹ *Anchusa sempervirens* reported by Sir W. Watson in 1749, *l.c.*

² *Cistus laxus*, *crispus*, *populifolius*, and *monspeliensis* were introduced by Tradescant. Loudon, *Arboretum*, p. 50.

Crocus, violaceus minor.	Eryngium marinum vulgare.
„ flore cinereo.	Esula major.
„ Maesiacus argentinus.	„ montana.
„ Maesiacus luteo Duc.	„ minor.
Croci flore variegato diversae species.	Eupatorium Novae Belgiae.
Cruciata Gentiana.	
Crupina.	Faba Americana.
Cucumer asininus.	Fabarum diversae species.
Cyanus maior.	Ferula Galbanifera.
Cyani hortensis varietates.	*Ficus.
Cyclamen flo. albo.	„ Indica.
„ folio Hederæ.	Flamula Jovis.
„ folio Hederæ Italicum.	Flos Africanus.
Cynoglossum minus.	„ „ flo. pleno.
„ non descriptum.	„ Constantinopolitanus fl. Miniato.
*Cyperus longus.	„ Constantinop: flore rubro.
Cypressus. ¹	„ Constantinop: flore pleno.
Cytisus Maranthæ.	„ Passionis.
„ primus Clusii.	„ solis maior.
„ secundus Clusii.	„ solis prolifera.
	Foeniculum.
Delphinium flore pleno.	„ dulce.
Dens caninus flo. albo.	Fraga fructu albo.
„ caninus flo. rubello.	„ fructu maiora.
Dentaria tryphylla.	„ fructu viridi.
„ septifolia.	„ spinosa sive hispida.
Dentillaria Rondeletii.	„ communia.
Digitalis flo. albo.	Fraxinella flo. albo minor.
„ alba maior.	„ purpur. maior.
„ variegata.	„ flore rubro.
„ ferruginea maior & minor. ²	Fritillaria flore rubro.
Doronicum Americanum.	„ flore albo.
Draba flo. albo.	„ vulgaris maior & minor.
Draco herba.	„ Aquitanica.
Dracuntium maius, Serpentaria. ³	Frumenti Turcici variet. triplex.
	Frutex Canadensis Epimediî folio.
Echium.	„ Coronaria flo. pleno, Syringæ
Elleborus albus. ⁵	flore.
„ albus flo. atrorubente.	
„ niger.	Galega flore albo.
Elleboraster.	„ flore carneo.
Endiviae species.	Genista Hispanica.
Epemedium.	„ hortensis.
Equisetum Marinum.	Gentiana maior.
Eruca perigrina Clusii.	„ foliis Asclepiadis.
„ Aragonica.	Gentianella alpina Helvetica.
Eryngium Constantinopolitanum.	Geranium Virginianum.
„ planum.	„ odoratum longius radicatum.
„ flore luteo.	„ muscatum.

¹ Tradescant's Virginian Cypress 'Cupressus Virginiana Tradescanti' in the 1656 catalogue does not appear in 1634. It is *Taxodium distichum* Rich. Parkinson reported English seedlings in 1640.

² Seen in 1632 by Johnson.

³ *Dracunculus vulgaris* L. seen by Sir W. Watson, 1749 (*Phil. Trans.* xlvii).

⁴ 'Gladiolus.' 'Iohn Tradescant assured mee, that hee saw many acres of ground in Barbary spread over with them.' *Parkinson*, 1629, p. 190.

⁵ White Ellebor (*Veratrum album* L.) grew 'in some partes of Russia, in that abundance, by the relation of that worthy, curious, and diligent searcher and preserver of all natures rarities and varieties, my very good friend, Iohn Tradescante, . . . that as hee said, a good ship might be loaden with the rootes hereof, which he saw in an Island there'. *Park.* 1629, p. 346. He accompanied Sir Dudley Digges to Russia in 1618, and was the first man to investigate the flora of that country.

- Geranium Creticum.
 " Indicum nocte odoratum.
 " tuberosa radice.
 " non descriptum Dodonaei.
 Gingidium.
 Gladiolus Byzantinus. [See note ⁴,
 " flo. albo. p. 337.]
 Glaux aestiva supina Lusitanica.
 Gnaphalium flore albo.
 Gramen striatum.
 Graminis diversae species.
 Gratiola.
 Grossularia maxima.
 " maxima longa.
 " coerulea.
 " rubra major rotunda.
 " media species longa.
 " rubra minor.
 " spinosa.
 Guaiacum Patavinum.

 Halimus arborescens.
 Hedysarum clypeatum Lob.
 Helleborus, see Elleborus.
 Hepatica flore albo [twice].
 " flore coeruleo major.
 " flore coeruleo minor.
 " flore albo cum staminibus
 " rubris.
 Hepatica coerulea flo. pleno.
 " coerulea flore pleno altera.
 Herba Doria.
 Hesperus Italica.
 Hieracium medio nigrum.
 " lanuginosum flore luteo.
 " dentis leonis facie.
 Hippoglossum.
 Hippomarathrum Lusitanicum.
 Horminum sylvestre Lusitan: flore albo.
 " sylvestre Lusitan: flore coe-
 " ruleo.
 Hyacinthus botroides flore albo.
 " botroides flo: coeruleo.
 " brumalis.
 " comosus.
 " Orientalis flore albo.
 " Orient: flore coeruleo.
 " Orient: atro-rubiens.
 " Peruvianus flore albo.
 " Peruv: flore coeruleo.
 " paniculatus.
 " Pyrenaeus flore albo.
 " Pyrenaeus flore coeruleo.
 " flore obsoleto, Clusii.
 Hyoscyamus albus.
 " albus medio-purpureus.
 Hypericum latifolium Lusitanicum.
 Hyssopus sativa.
 " prolifera.
 " alba.
 " tenuifolia.
 " comosa.
- Jacea maxima odorata.
 " spinosa.
 " flo. luteo.
 " aestiva elegans.
 Jacobaea.
 " marina.
 " latifolia: Baetica.
 Jasminum Catalonicum flo: albo.
 " flo. albo.
 " flo. luteo.
 " Persicum.
 Hex.
 Impatiens herba Dodonaei. Persicaria
 " siliquosa.
 Irides maiores variae.
 Iris gloriosa.
 " Susiana major.
 " Clusii flore pleno.
 " Clusii flore albo.
 " Clusii flo: coeruleo.
 " bulbosa Anglica maior flore albo.
 " " " minor flore albo.
 " bulbosa Anglica flore coeruleo.
 " Persica.
 " bulbosa Africana.
 " bulbosa Anglica variegata.
 " bulbosa flore luteo.
 Iridis bulbosae aliae diversitates.
 Irides humiles sive Chamaeirides va-
 " riae.
 Juniperus minor.
 Jucca.

 Keyri maius simplex.
 " maius ferrugineo flo. pleno.
 " flore albo simplex.
 " flore pleno pyramidale.
 " flore pleno vulgare.
 " flore pleno auratum.

 Laburnum maius.
 " minus.
 Lagopus flore rubro.
 Lapathum hortense.
 Lathyrorum elegantium variae species.
 Lavendula.
 " multifido folio.
 Laurea cerasus.
 Laurus tinus.
 " tinus folio glabro.
 " Alexandrina.
 " Gallica.
 *Lentiscus.
 Leucojum bulbosum maius.
 " bulbosum minus.
 " arbo. flore pleno rubro.
 " arbo. flo. pleno albo.
 " arbo. flo. pleno purpureo.
 " arbo. flo. pleno rubro varie-
 " gato.
 Leucojum arbo. flo. pleno purpureo
 " variegato.

Leucojum marinum. ¹	Medica doliata echinata.
Libanotis.	" doliata ramosa.
Liliasphodelus flore luteo.	" clypeata.
" flore albo.	Melilotus Italica flore luteo.
Lilium album.	" arborescens flore albo.
" Convallium flore albo.	Melissa.
" Convallium flo: rubro.	Mentha.
" flore luteo.	" crispa.
" Constantinopolitanum.	Meum.
Lilac Matthioli. ²	Mezereon album.
Limonium minus angustifolium.	" rubrum:
Linaria odorata.	Milium nigrum.
Locusta Virginiana arbor. ³	" album.
Lotus arbor. ⁴	Millefolium atro rubente flore.
" Lybica.	Mollugo.
" coronata maxima Hispanica.	Morus.
Lupinus Indicus.	" alba.
" candidus ex Candia.	" Virginiana.
" flore luteo.	Muscari flore luteo.
" flo. coeruleo minor.	" flore albo.
Lutea Cretica.	Myagrum monopermon.
Lychnis sativa rubra flore pleno.	Myrrhys sativa.
" coronaria pleno flo. albo.	Myrtus latifolia.
" coronaria pleno flo. rubro.	" angustifolia.
" sylvestris Pyrenaeus.	" florida.
Maiorana.	Napellus.
Malva arborescens.	Narcissus medio Croceus.
" maxima Hispanica striata.	" Anglic. flo. pleno Wilmot.
" segetum Lusitanica.	" roseus maximus flore pleno
Malum arantium.	Tradescanti. ⁶
" limonium.	Narcissus Africanus odoratus.
" granatum.	" Africanus major praecox.
Mandragoras mas. ⁵	" Indicus Jacobaeus.
Marum.	" Capa bonae spei.
Martagon Panonicum spadaceum.	" tertius Matthioli.
" Pompon.	" Montis Carmeli.
" flore albo punctato.	" Virginianus.
Matricaria.	" medio fimbriatus.
" flore pleno.	" Robinus maior.
Medica spinosa maior.	" juncifolius luteo flore pleno.
" transversis spinis.	" humilis.
" scoparia.	" reflex flo: luteo.
" elegans Catalonica.	" reflex flo: albo.
" minor spinosa.	" oblong: calice flore luteo.

¹ The greatest Sea Stocke Gilloflower (*Matthiola sinuata* L.) was brought out of the Isle of Ree [Rhé] by *Rochel* by Mr. John Tradescant, when the Duke of Buckingham was sent with supplies for Mounseur *Soubise* (1627). Park *Theatrum*, 624.

² The Lilac, *Syringa persica* L., was introduced by Tradescant. Loudon, *Arboretum*, p. 49.

³ *Robinia Pseudacacia* L. 'A very great tree and of exceeding height with Master Tradescant,' in 1640. Parkinson, *Theatrum*, 1550. Mentioned by Ashmole, in 1662 (?), but not by Watson in 1749.

⁴ The Date Plum (*Celtis australis* L.). 'Lotus arbor Virginiana' (*Celtis occidentalis*) was in the garden in 1656.

⁵ The Male Mandrake. Some years previously Parkinson 'saw in my Lord *Woolton* his garden at *Canterbury*, whereof Mr. *J. Tradescant* had then the keeping, an other sort'. *Theatrum*, p. 343.

⁶ 'The greatest double yellow bastard Daffodill, or Iohn Tradescant his great Rose Daffodill. This Prince of Daffodils belongeth primarily to Iohn Tradescant, as the first founder thereof. . . . Whether raised from seed, or gained from beyond Sea, I know not.' Parkinson, 1629.

Narcissus oblongo calice flore albo.	Papas Americana flore albo.
„ calice brevi.	„ Americana flore. purpureo.
„ titesose.	Papaver rhaeas fl. luteo radice perpetua.
„ va Hecueus.	„ rhaeas flore simplici.
„ omnium maximus.	„ rhaeas flore duplici.
„ montanus.	„ nigrum capit. rotundis.
„ Non-parell flore albo.	„ corniculatum flore luteo.
„ Non-parell.	Paralysis flore viridante simplici.
„ Constantinopolitanus.	„ fatua.
„ totus albus.	„ inodora geminata.
„ Matineus.	„ flo. & calice crispo.
Nasturtium Indicum.	Pepo Americanus luteus.
Nigella flo: pleno.	„ Americanus viridis.
„ citrina.	Periclymenum rectum 2. Clusii.
„ flo: simplici.	„ fructu cerasino.
Nux juglans Virginiana	„ hortense.
„ juglans Canadensis.	„ Germanicum flo. rubro.
„ juglans Angliae novae.	Periploca.
„ juglans maior.	Petroselinum crispum.
„ juglans minor.	„ hortense.
„ ¹	„ Virginianum.
Oenanthe bulbosa marina venenosa.	Phalangium Alobrogium Clusii.
Olea sylvestris.	„ Virginianum Tradescanti. ²
Oleander flore albo.	Philyrea angustifolia.
„ flore rubro.	Pimpinella maior.
Ononis non spinosa Pyrenaea.	„ sativa.
„ non spinosa oderata flore luteo.	„ agrimonoides Colum.
„ non spinosa aestiva minor flore luteo.	Pinus.
Origanum verum Hispanicum.	Pinaster.
Ornithogalum Neopolitanum.	Pistachia sativa. ³
„ Arabicum.	Pistolochia smilacis folio.
„ maius flo. albo.	„ Virginiana.
Orobus Venetus.	Pisum indicum.
Paliurus.	„ perennes.
Paeonia mas.	„ bacciferum.
„ foemina flore simplici.	„ maculatum.
„ foemina flore pleno.	„ quadratum.
„ flore pleno incarnato.	Plantago rosea.
„ flore purpureo.	„ serrato folio.
	Platanus. ⁴
	Polium montanum.
	Polygala Valentina 1. Clusii.

¹ 'Nux vesicaria' in the 1656 catalogue may be our common Bladdernut *Staphylea pinnata* L., but it should be remembered that Parkinson, *Theatrum*, p. 1417, says that 'Mr. Tradescant hath brought a sort from Virginia, having divers nuts in the bladder'. Loudon, *Arboretum*, p. 49, attributes the introduction of *Staphylea trifolia* to Tradescant.

² 'Tradescant his Spiderwort. This Spiderwort is of late knowledge, and for it the Christian world is indebted vnto that painfull industrious searcher, and louer of all natures varieties, Iohn Tradescant (sometimes belonging to the right Honourable Lord Robert Earle of Salisbury, . . . and then vnto . . . the Lord Wotton at Canterbury in Kent, and lastly vnto the late Duke of Buckingham [assassinated 1627]), who first receiued it of a friend, that brought it out of Virginia, thinking it to be the Silke Grasse that groweth there, and hath imparted hereof, as of many other things, both to me and others.' *Park.* 1629, p. 152. In 1617, Tradescant (who may have then been in Wotton's service) paid the expense of a passenger to Virginia under 'Captain Argall'. Boulger, *J. Bot.* 1918, p. 200.

³ *Pistacia Terebinthus* L. was introduced by Tradescant acc. to Loudon, *Arboretum*, p. 49.

⁴ *P. orientalis*. 'There are one or two yong ones at this time growing with Mr. Tradescant.' *Ger. emac.* 1633, p. 1489. *P. occidentalis* is believed to have been introduced by Tradescant the younger.

- Polygala Valentina 2. Clusii.
 Polygonatum maius.¹
 " maius angustifolium.
 " minus.
 " racemosum Virginianum.
 " alterum.
 Polygonon marinum Lobelii.
 Pomum amoris medium.
 Primula veris flore albo.
 " veris albo flore pleno.
 " veris flore pleno viride.
 " veris angustifolia flore albo.
 " veris angustifolia flore rubro.
 " veris flo. viride & albo simpl.
 Pseudo dictamnium.
 Ptarmica vulgaris.
 Pulegium regale.
 " cervinum.
 Pyracantha.
 Radix cava maior flore purpureo.
 " cava minor flore albo.
 Ranunculus albus flore pleno.
 " Asiaticus sang. flo. pleno.
 " Asiat. tenuifol. pleno flo.
 luteo.
 Ranunculus Asiat. flore albo.
 " Asiaticus flore rubro.
 " Asiat. flore luteo.
 " Asiat. folio papaveris.
 " Asiat. *Drape de Argentine*.
 tine.
 Ranunculus Illyricus.
 " aliae diversitates.
 Raphanus niger.
 Rapunculus.
 Reseda maior.
 Rhodia radix.²
 Rhus myrtifolia.³
 Ribes fructu rubro.
 " fructu albo.
 " fructu nigro.
 Ribesium dulce.
 Rosa Provincialis.
 " Provincialis flore albo.
 " Provincialis flore rubro.
 " vitriensis flore pleno.
 " incarnata.
 " flore luteo pleno.
 " flore luteo simplici.
 " muscata flore pleno.
 " Italica.
 Rosa cinamomea.
 " cinamomea flore albo.
 " Francofurtiana.
 " Batavica.
 " alba variegata.
 " flo: pleno elegans variegata.
 " flore simplici pomifera.
 " Virginiana.
 " Moscovitica.⁴
 " canina flore pleno.
 " Eglanteria flore pleno.
 " Eglanteria.
 " holoserica.
 " sempervivens.
 " flore rubro.
 " flore albo.
 " Damascena.
 " Austriaca flore Phoeniceo.
 Rosmarinus.
 " auratus.
 " coronarius maximus.
 *Ruta canina.⁵
 Sabina.
 Salvia variegata.
 " hortensis rubra.
 " hortensis viridis.
 " maior foliis crispis.
 " minor odoratissima.
 Sambucus aquatica.
 " foliis laciniatis.
 Sanicula Alpina guttata.
 Saponaria flore pleno.
 Saxifraga aurea.
 Scabiosa Hispanica major.
 " Hispanica Clusii.
 " Indica.
 " Indica Clusii.
 Scorpioides Portulacae folio.
 " bupleurifol. siliq. crassa
 torosa.
 Scorpioides minor.
 " minor elegans.
 Scorzonera.
 Scrophularia Montis-Serrati.
 " Pannonica Clusii.
 Securidaca minor.
 " maior.
 " perigrina Clusii.
 Sedum majus.
 " arborescens.
 " elegans.

¹ Solomon's Seal was found living in the Lambeth garden by Sir W. Watson in 1749 (*Phil. Trans.* xlvj).

² *Rhamnus catharticus* is absent from this list, but is in the 1656 list. In 1749 Sir W. Watson reported a tree 'about 20 feet high and near a foot in diameter' (*l. c.*). Perhaps the younger Tradescant planted it.

³ *Rhus Cotinus* L. was introduced by Tradescant acc. to Loudon, *Arboretum*, p. 49.

⁴ No doubt one of the spoils of Tradescant's Russian expedition of 1618. Cf. Boulger, *First Russian Botanist. J. Bot.* 1895.

⁵ [Probably in error for *Rosa canina*.]

Serpentaria.	Thapsia Neronis Carotif. Lob.
Seseli Aethiopicum frutex.	Thlaspi umbellatum.
Sinapi Castiliae novae.	Thymum verum Hispanicum.
Smilax aspera folio rotundo.	Tithymalus Charachia.
„ aspera folio maculato.	Tithymalorum diversae species.
„ aspera levis.	Trachelium album flo. pleno.
Solanum Lethale.	„ coeruleum flo: pleno.
Spina Solstitialis.	Trifolium barbaricum stellat. Tradesc. ¹
Staechas Arabica.	Triticum spica multiplici.
„ Citrina.	Tuliparum eligant: maxima diversitas.
Stachys Hispanica.	Tulipa Num. 50. diversae species.
„ spuria.	
Stachylodendrum.	Valeriana Graeca flore coeruleo.
Stoebe Salamantica.	„ Graeca Dodon. flore albo.
Stramonia flore albo.	„ Dodonaei.
„ flore purpureo.	Verbascum salvifolium.
Superbae diversae species.	„ blattariae folio.
	Vergae aureae quatuor spec.
Tamariscus Italica.	Veronica mas.
„ minor.	„ foemina.
„ vulgaris.	Vinca pervinca maior.
Tanacetum.	„ minor.
„ crispum.	Viola Matronalis flore pleno.
„ inodorum.	Vite sylvestris.
Telephium maius.	Vlmaria perigrina Clusii.
„ minus.	Vmbilicus Veneris.
Terebinthus vera.	„ „ Hispanicus.
Thalictrum Virginianum.	Vrtica Romana.

A Catalogue of Fruits.²

*Apples.*³

D octor Barchams Apple	Ribon Apple
Pome de Rambure	Pome Mater
Master Williams	Russet Pippin
Yellow Russeting	Puffing Apple
Harry Apple	French Pippen
Dutch Pearmaine	Snouting
Black Apple	Blandrille
Barfolde Queninges	Torne Crab
Smelling Costard	Great Russeting
John apple	Summer Beiliboon
Red master Williams	Quince Crab
Quince apple	Pome de Chastania
Summer Pearmaine	Pome de Renet
Winter Pearemaine	Pome de Carpandū
Gillefloure Apple	Pome de Caluele

¹ Pulteney (*Biogr. Sketches*, i. 176) states that Tradescant brought *Trifolium stellatum* L. from the Island of Formentera. This visit may have been during the Algerian campaign of 1620.

² To the names in the 1634 catalogue, for the most part printed in Roman type, we have added in the right-hand column and in *italics* the names and dates of ripening of the fruits depicted (? by Alex. Marshall, see *Mus. Trad.*, p. 41) in a book known as *Tradescant's Orchard* now in the Bodleian Library (MS. Ashmole 1461).

³ Shakespeare's apples, the John apple, Bitter-sweet or sweeting, Crab, Codling, Leather-coat (= Yellow Russeting), Pippin, and Pomewater are all in this list, but curiously enough, though Tradescant had a Poperin pear, he does not mention the Warden pear by that name.

Violet Apple	Little sweeting
Darling Apple	Yellow Spising
Stoken Apple	Dari Gentles
Sack and Sugar	Livinges
Pidgions bill	Mother Pippin
The Kings apple	Russet Peare apples
M. Molines apple	Keelings
Grey Costard	Ginitings
Winter Belliboorne	Black Pipin

An Early ripe Apple good in taste Aug. 22

Peares.

B Infeild Peare	Poyer Messer Ian
Gergonell	Nutmeg Peare
Sir Nathaniel Bacons great Peare	Bishops Peare
Red Peare	Orenge Burgamott
Rose water Peare	May Peare
Greenefield Peare	Swise Peare
Dego Peare	Summer Burgamot
Scarlet Peare	Hony Peare
French Popering	Mid-summer Peare
Snow Peare	Winter Burgamot
Winter Boon Critian	<i>Poyer de Poydre</i>
Summer Boon Critian	Portingale Peare
Arundell Peare	Sugar Peare
Pallas Peare	Double floure Peare
Prince Peare	Bloud Peare
Greene Peare	<i>Poyer Fran Rial</i>
Hedera De Besa	Winter Winsor
Michaelmas Peare	Summer Winsor
M. Motts Peare	<i>Poyer Irish Madam</i>
Paynted Peare	<i>Poyer Dangobet</i>
Sliper Peare	<i>Poyer de Valet</i>
Greene Rowling	<i>Poyer de Sauoyse</i>
Kings Peare	

[TRADESCANTS ORCHARD.¹]

The greet 16 ounce peere
The grete winter peere
The Jerusalem Pere
A french peare called R. Collanit ripe Aug. 29

Quinces.

The portingegale Quince
The peare Quince Oct. 5

Plums.²

M Oroco Plum	<i>Red plumordin Plume ripe July 12</i>
Spanish Plum	<i>Vilot plum „ July 24</i>
Blew peare plum	<i>Maraco plume „ July 15</i>
Red peascod Plum	<i>Early whight pere plum „ July 30</i>
White Plum	<i>Denny plum „ Aug. 6</i>
Plum Dine	<i>Grene Oysterly plum „ Aug. 9</i>
Rath ripe Damaske violet	<i>Grene mother plum „ Aug. 14</i>

¹ To this is prefixed a pleasing coloured drawing of 'Mertagon vel Leli novae Angliae'.
² Plums. 'The choysest for goodnesse, and rarest for knowledge are to be had of my very good friend Master John Tradescante, who hath wonderfully laboured to obtaine all the rarest fruits he can heare off in any place in Christendome, Turky, yea or the whole world.' Parkinson, *Paradise*, 1629, p. 575.

Damaske Violet	<i>Nutmeg plum</i>	<i>ripe Aug. 18</i>
Verdoch Plum	<i>Frier plum</i>	<i>„ Aug. 16</i>
Friers Plum	<i>Red Mussell plumbe</i>	<i>„ Aug. 20</i>
Bowle Plum	<i>Sheffell Bullis</i>	<i>„ Aug. 22</i>
Nutmeg Plum	<i>Impryall plum</i>	<i>„ Aug. 22</i>
White Rath ripe Plum	<i>Gante plum</i>	<i>„ Aug. 23</i>
Peake Plum	<i>Mussule plum</i>	<i>„ Aug. 24</i>
Apricocke plum	<i>Grene pescod plum</i>	<i>„ Aug. 24</i>
Oreng Plum	<i>Damsonn</i>	<i>„ Aug. 26</i>
Michaelmas damaske Plum	<i>Pruon Damson</i>	<i>„ Aug. 27</i>
Red Mirabolane	<i>Red pescod plum</i>	<i>„ Aug. 28</i>
White mirabolans	<i>Whight Date</i>	<i>„ Aug. 29</i>
The Monsiers Plum	<i>Whight mussell plumbe</i>	<i>„ Sept. 2</i>
The Perdigon Plum	<i>Blacke peare plum</i>	<i>„ Sept. 3</i>
The Kings Plum	<i>Amber plum which J. T.</i>	
The Queenes Plum	<i>as I take it brought</i>	
The white Perdigon	<i>out of France and</i>	
The pruneola Plum	<i>groweth at Hatfeld</i>	<i>„ Sept. 8</i>
The Diapre Plum of Malta	<i>The Turke Plum</i>	<i>„ Sept. 16</i>
The Diapre Plum		
The Imperiall Plum		
The Date Plum		
The Musle Plum		
The Damascene Plum		
The Irish Plum		
White Damaske violet plum		

Cheries.

S wertes Cherie		
Seelinars Cherry		
The great Hart Cherry	<i>Harte Cherry</i>	<i>ripe June 24</i>
The great bearing Cherry	<i>Corone Cherry</i>	<i>„ June 19</i>
The Arch-Dukes Cherry ¹	<i>Naples Cherry</i>	<i>„ July 1</i>
The Spanish cherry		
The Luke Ward Cherry	<i>Luke ward Cherry</i>	<i>„ June 10</i>
The Agriot Cherry		
The Chamelion Chery		
The dwarfe Hungarian chery	<i>Dwarfe Cherry</i>	<i>„ July 1</i>
Tradescants Chery ¹	<i>Tradescant Cherry</i>	<i>„ June 21</i>
The white Chery	<i>Whighte Cherry</i>	<i>„ June 24</i>
The cluster Chery	<i>Cluster Chery</i>	<i>„ June 15</i>
The double floure Chery		
The May Chery	<i>May Cherry</i>	<i>„ [June] 2</i>

Apricocks.

B arbarie Apricocks 2 sorts ²		
Small Holland Apricocke	<i>Round Apricock</i>	<i>ripe Aug. 15</i>
Masculine Apricocke	<i>The Apricocke that is</i>	
Longe muske Apricock	<i>both long and great</i>	<i>„ Aug. 23</i>
The ordinary Apricocke		

¹ 'John Tradescantes Cherie is most usually sold by our Nursery Gardiners, for the Archdukes cherrie, because they haue more plenty thereof, and will better be increased.' Parkinson, *l.c.*, p. 574.

² The Argier Apricocke 'with many other sortes John Tradescante brought with him returning from the Argier voyage, whither hee went voluntary with the Fleete, that went against the Pyrates in the year 1620.' Parkinson, *l.c.*, p. 579.

Nectarins.

The Roman red Nectarine	<i>Roman Reed Nectrion ripe</i>	Sept. 2
Sir Edward Sillards ed Nectarine	<i>Bastard Red Nectrion</i>	„ Sept. 4
The little yellow Nectarine	<i>Cluster Red Nectrion</i>	„ Aug. 22
The white Nectarine	<i>Yellow Nectrion</i>	„ Sept. 13
	<i>Grene N.</i>	„ Sept. 5

Peaches.

Tradescants double floured Peach	<i>Grete early yellowe peeche</i>	ripe Sept. 6
The Queenes Peach	<i>Blake peeche red all within</i>	„ Sept. 20
The White Peach	<i>Whight peeche</i>	„ Sept. 21
The Nutmeg Peach		
Peach de Troas	<i>The peeche Dutroye</i>	
Newington Peach	<i>Nwingetonn Peeche</i>	„ Aug. 4
Carnation Peach	<i>Round Carnation peeche</i>	„ Sept. 24
Spanish Peach	<i>Graunde</i>	„ Sept. 4
Devine Peach	<i>Bell</i>	„ „
Lions Peach	<i>which pealleth like a codling</i>	„ Sept. 3
Roman Peach	<i>The Russet Blud peeche</i>	
Peach Pavi Iaune	<i>or Durosynus</i>	„ Sept. 25
	<i>The Mallycotone peeche</i>	„ Sept. 26
	<i>A late ripe yellow peeche</i>	„ Oct. 10
	<i>but very good firme peeche</i>	

Vines.

The Parsly leaved Vine	<i>The buxtet Grape wich very seildum ripe</i>	
The Fronteneac Vine		
The great blew Grape	<i>The blue grape</i>	ripe Sept. 27
The Potbaker Grape		
The reison Grape	<i>The Grat Reson Grape</i>	„ Oct. 10
The currans Grape	<i>The smalle Reson Grape</i>	„ Sept. 12
with divers other.		

FINIS.

The grete Roman Hasell Nut
The great French Fragara, ripe the 20 of May.

A contemporary notice of this catalogue occurs in the Diary of Georg Christoph Stirn of Nürnberg (MS. Bodl. Add. B 67). Stirn left Dieppe for England on 2 July 1638. His sightseeing in London included the Tower, York House, and the Tradescant Museum, and he wrote in his diary that in the garden were all kinds of foreign plants, the names of which are to be found in a special little book which Mr. Tradescant has had printed about them. This 'special little book' now shares with Gerard's first Catalogue (1596) the honour of being among the scarcest of printed botanical works in the world.

The contents of the Lambeth garden were again listed by the younger Tradescant, who printed the second *Catalogus Plantarum in Horto Johannis Tradescanti nascentium* in 1656. And a few years later Ashmole drew up a list of all the trees still surviving in

the garden, and wrote it out at the end of his copy of Parkinson's *Paradise*, now in the Bodleian Library.

Trees found in Mrs. Tredescants Ground when it came into my possession. [1662.]

Platinus orientalis verus.	<i>Platanus orientalis</i> L.
" occidentalis, aut Virginensis.	" <i>occidentalis</i> L.
Arbor siliquosa Virginensis spinosa, Locus nostratibus dicta.	" <i>Robinia Pseud-acacia</i> L.
Cerasus racemosa putida ¹ Padus Theophrasti dicta.	<i>Prunus Padus</i> L.
Periclymenum rectum ¹ flore rubro.	<i>Lonicera alpigena</i> L.
Nux Vesicaria, altera Virginensis.	<i>Staphylea trifolia</i> L.
Euonymus Theophrasti.	<i>Euonymus europaeus</i> L.
Lotus Arbor.	<i>Celtis australis</i> L.
Sambucus Rosea.	<i>Viburnum Opulus</i> var. <i>sterilis</i> .
Arbor Judae.	<i>Cercis Siliquastrum</i> L.
Cornus Mas.	<i>Cornus Mas</i> L.
" foemina.	<i>Cornus sanguinea</i> L.
Latana, sive Viburnum.	<i>Viburnum Lantana</i> L.
Guaicum Patavinum.	<i>Diospyros Lotus</i> L.
Syringa alba.	<i>Syringa vulgaris</i> L. var. <i>alba</i> .
Pyracantha.	<i>Pyracantha coccinea</i> Roemer.
Alaternus.	<i>Rhamnus Alaternus</i> L.
Arbutus.	<i>Arbutus Unedo</i> L.
Castanea equina.	<i>Æsculus Hippocastaneum</i> L.
Pinaster.	<i>Pinus Pinaster</i> L.
Laurus Tinus.	<i>Viburnum Tinus</i> L.
" " Lusitanicus flore glabro.	<i>Prunus lusitanica</i> L.
Tillia.	<i>Tilia vulgaris</i> Hayne.
Tamariscus.	<i>Tamarix anglica</i> Webb.
Acer majus latifolium.	<i>Acer Pseudo platanus</i> L.
Rhus Virginiana.	<i>Rhus typhina</i> L.
Vitis Virginensis.	<i>Vitis quinquefolia</i> Lam.
Apocynum, sive Periploca repens.	<i>Periploca graeca</i> L.?
Althea arborea flore albo fundo purpureo Montis Olbiae.	<i>Hibiscus syriacus</i> L.
Seseli Æthiopicum frutex.	<i>Bupleurum fruticosum</i> L.

xi. GEORGE GIBBES' GARDEN LISTS, UNDATED AND 1634.

George Gibbes had a garden at Bath which was visited by Thomas Johnson and the *Socii itinerantes* on their tour in the west of England in 1634. There is a short, undated list of Gibbes' plants among Goodyer's papers, though it is not in his handwriting. It is headed 'To have from Mr. Gibbes'. Johnson printed a list of 117 of Gibbes' exotic plants in the *Mercurius Botanicus*, of these only six (marked below with double asterisks) occur in the short list.

It will be noted that none of the plants imported by Boel from Spain occur in either list, whereas they do occur in the Stonehouse

¹ Mr. Boulger, who saw the MS. in 1917 before it had been purchased by the Bodleian, by reading the word 'putida' as 'qubida', and conjecturally amending it as 'quibusdam' (!), has illustrated the danger of not minding one's p's and q's. In the next line he read 'erectum' in error for 'rectum'. I have, however, placed implicit trust in his determination of the modern equivalents of the names.

list of 1640. Parkinson has an interesting note on Gibbes in his note on a Virginian Aster, which was evidently imported after the date of Johnson's list.

Aster Virgineus luteus alter minor.

'We have had scarce time enough to observe it thorowly since we got it from Virginia by the means of Master *George Gibbes* Chirurgion of Bathe, who brought in his returne from thence, a number of seeds & plants he gathered there himselfe, & flowered fully only with Mr. Tradescant.' Parkinson, *Theatrum*; 1640, p. 133.

Gibbes' Garden 1634.

*Plants marked * are in the Goodyer list only.*

*Plants marked ** are in both lists.*

Abrotanum mas.	Cotyledon altera Ger. emac.
Absinthium tenuifolium, quibusdam Romanum, aliis Ponticum.	Crocus vernus.
Ageratum.	**Cucumis asininus, Offic.
Allium max. radice simplici.	Cyanus flo. albo, carneo et.
Allium vulgare.	„ major.
Antirrhinum majus flo. albo.	Doronicum Romanum.
*Aquilegia angustifolia, mult., simpl.	Faba Graecorum.
Arbor vitae.	Fragaria fructu magno, Boëmica, Quorund.
Aristolochia Clematites.	Flos adonis.
*Armerius latifol.	*Flos africanus.
Asclepias flore albo.	Flos solis major.
Aster Italorum.	Gentianella verna flo. amplo.
**Astragalus Lusitanicus.	Geranium longius radiculatum, Lob.
Astrantia nigra.	„ Moschatum.
Auricula ursi.	„ Romanum variegatum.
Blattaria purpurea.	Gladiolus Narbonensis.
* „ alba.	Gnaphalium Americanum.
*Bistorta minor.	Glycyrrhiza vulgaris.
Bolbonach, sive Lunaria Greca.	Gramen Lupuli glumis, sive Tremulum max. Bauh.
Branca Vrsina.	**Helleborus niger verus.
*Buphthalmum.	Hepatica nobilis.
*Buxus versicoloribus foliis.	* „ albida fl. rub.
*Calceolus mariae.	Hipposelinum
Caltha flo. multiplici.	Hyssopus vulgaris.
* „ palustris.	* „ foliis aureis.
Campanula Persicifolio, Lob.	Imperatoria.
Caryophyllorum hortensium variae spec.	Iris bulbosa.
Cerinte.	„ latifolia vulgaris.
Chamaecyparissus.	Iacea tricolor flo. amplo lut.
Chamaedrys.	„ „ vulgaris.
*Chamaeiris latifol.	Iasminum album.
* „ angustifol.	„ luteum, sive Trifolium fruticans, Polemonium Ger.
Chamaexyris, Lob.	Laurus vulgaris.
Chrysanthemum Creticum.	Lavendula vulgaris.
Cochlearia Batava.	„ flo. albo.
Colutea vulgaris.	Leucium bulbosum et alia verna, quae tunc temporis non apparuerunt.
Consolida regalis.	
**Convolvulus coeruleus minor.	
Coriandrum.	
Costus hortorum.	
Cotyledon minus montanum, Palmaria Tab., Umbilicus veneris minor, Gr.	

- Leucoium flo. albis et purp.
 Liliium album.
 " montanum.
 " non bulbosum.
 Lotus hortorum, Lob. sive Trifolium
 odoratum.
 *Lupinus albus, max., coer., aur., et
 luteus.
 Lychnis Chalcedonica.
 " hortensis.
 " syl. flo. multipl.
 Lysimachia coerulea.
 " lutea Virginiana.

 Majorana.
 Marum sive Mastich Gallorum et
 Anglorum.
 Mentha sativa rubra.
 Meum.
 Muscipula.
 Myrrhis sativa.

 Napellus coeruleus.
 Nigella Romana.

 Oxalis rotundifolia.

 Papaver sativum simp. et multipl. flo.
 Parthenium flo. multipl.
 Pentaphyllum surrectum.
 Periclymenum perfoliatum.
 *Peucedanum.
 Phalangium non ramosum.
 " Virginianum.
 Polium luteum.
 Psyllium.

 *Radix cava. flo. carn.
 Raphanus rusticanus, Offic.
 Rha rotundifolium.
 Rhabarbarum Monachorum.
 " verum.
 *Rosmarinum aureum.
 * " latifolium.
 Ruta.

 Sabina.
 Salvia angustifolia.
 " major vulgaris.
 " variegata.
 Saponaria flo. multipl.
 Scabiosa montana max.
 Scordium.
 Scorzonera.
 Serpilli 3. spec.
 Stoechas vulg. Offic.
 Sumach Virginianum.

 Tamariscus Narbonensis.
 Thalictrum majus Hispanicum.
 * " virginianum.
 **Thlaspi Creticum.
 Thymum durius.
 Trachelium majus flo. albo.
 Tragopogon flo. purp.
 Tuliparum varietates plurimae.

 Umbilicus Veneris sive Cotyledon
 altera, Ger. emac. Sedum serratum.

 Valeriana Graeca.
 Vinca pervinca, sive Clematis Daph-
 noides.
 Viola mariana.
 " matronalis flo. pleno.

xii. WALTER STONEHOUSE'S GARDEN LIST, 1640-1644.

The garden of the Rev. Walter Stonehouse at Darfield Rectory is described in a neatly written vellum-bound 12mo volume of 44 leaves, with a leaf with plans of the garden, known as Magdalen College MS. No. 239, or as Goodyer MS. 17. It is entitled *Catalogus Plantarum Horti mei Darfeldiae Quibus is instructus est Anno Domini 1640*, and though anonymous, is convincingly identified as the work of Stonehouse by his anagram 'Theologus servus natus' (= Gualterus Stonchousus) on f. 5 of the MS. The same anagram occurs in a MS. volume of Sermons in the Library of Magdalen College and at the end of his poems in the *Musaeum Tradescantianum* (1656).

Stonehouse's notes and plans show that the rectory grounds comprised a 'best garden', measuring 34 yards by 30 yards, the

Saffron Garth, a long rectangular strip running north and south for 82 yards, and the orchard. The plans are clearly drawn to scale and are marked with numbers which corresponded to his lists of plants. We have even his notes of the exact dimensions of the garden in terms of his own paces, from which we may infer that he was a man of no great stature. '2,624 of my usuall paces made an English mile of 1,056 paces Geometricall: that is 5,280 feete and 1,760 yards.' The garden 15 times half round is just a mile; 16 times is a mile and 57 yards.'

Though we have their orientation, we have no clue as to how the three parts of his garden were disposed, but may conjecture that the Rectory stood in the angle of the best garden. Plums, Peaches, Apricots and a Pomegranate covered the west and north walls; a vine was trained against a wall, possibly of the house. Several rectangles drawn at the ends of paths may represent garden ornaments; an arbour or summer house seems to have stood in the north-east corner.

The beds in the style of the sixteenth century may have been the work of an earlier incumbent. They were laid out in five 'knots', perhaps enclosed with tile, stone or Box edgings, which bordered the 'forthrights', as the broader walks were called. According to the plan the beds in the knots were two and three feet in width, and would, as Parkinson (1629) recommended, have contained the greater part of the herbaceous collection.

'The Safforn-garth 9 times round wants 50 yards of a mile; and 10 times about it is a mile and 140 yards. The long streight walke in the Safforn-garth is 82 yards; so that this gone 22 times single, or 11 times double, is a mile and 44 yards.'

The name recalls the fact that in the sixteenth century 'Crocus', or Saffron, was perhaps the most paying crop that it was possible to raise in a garden. 'Our English Honey and Saffron,' wrote Bullein in 1588, 'is better than any that cometh from any strange or foreign land.'

'A little of ground
brings Saffron a pound,'

and in 1539-40, Doncaster Saffron was sold at more than £1 per pound weight. Saffron Hill and Saffron Walden recall an extinct British horticultural industry, but Darfield Rectory, as the present Rector informs me, is noted for a Crocus border 170 yards long.

Most of Stonehouse's fruit was grown in the Saffron Garth and the Orchard, both of which were walled in. In the former he had 33 wall-trees, planted 6 feet apart, against the walls on the north

and west sides, and a selection of 30 Apples and Pears 'in the open plotte', as well as some more recent acquisitions from the garden of his neighbour, Sir John Reresby.

The best varieties of the original stock were probably obtained from Parkinson's 'very good friend, Master John Tradescante', who had made a speciality of all fruit trees, 'the choysrest for goodness, and rarest for knowledge,' and especially of Plums 'fit for an orchard'.

In the orchard there were fruit walls along the north and east, and a raised walk or 'mount' along the wall on the west side. This 'high walke' probably dated from before Stonehouse's time, for he refers to 'olde plum trees' upon it. The 'open plotte' was stocked with Apples of various kinds, with a row of Pear trees along the north walk.

'The high walke in the orchard, 66 times single gone, or 33 times double (that is backward and forward) is a just mile—5,280 foote.'

'The orchard 9 times round about is a mile and 4 yards.'

In the open the fruit trees were planted about 18 to 20 feet apart. Their positions are clearly indicated upon his numbered plan, from which the illustration in Fig. 2 has been redrawn for purposes of reproduction. (See *Gardeners' Chronicle*, 1920.)

The Sir John Reresby, from whom he obtained several varieties of Apples and Pears, was a neighbour living at Thrybergh, some six miles south of Darfield. His son, of the same name, was the well-known Governor of York, who wrote memoirs containing a secret history of the Courts of Charles II and James II.

The manuscript concludes with an epitaph in Latin verse by Stonehouse, to his favourite cat, Delia, which died when kittening and was buried in the garden.

*IN FELEM DILECTAM, INTER PARTUS
DOLORES EXTINCTAM.*

Delia (sic fertur) coelum invadente Typhoco.

Sub Fele, in terris, condidit ora latens.

*Partubus illa praeest; sed et illa (heu) saepe vocata,
Tardat, parturiens dum mihi Felis obit.*

*Ah, factum, Lucina, male! Ah, ea gratia Divae est,
Praesidii ut possit non memor esse sui.*

The present Rector of Darfield, the Rev. A. E. Sorby, informs me that the house has been greatly enlarged and probably covers the beds at T and the geometrical beds at S. 'The other four beds, V, W, Y, Z, with the star in the centre, are still in existence, though

not in the same geometrical position. The Saffron Garth is, I think, undoubtedly in the walled garden between the Rectory and the School Street. The measurements do not agree with the existing garden, but the shape is the same and the Saffron Crocus borders are still there. To the north of the Rectory and of the walled garden is a large orchard, where the Saffron Crocus up till lately also flourished and bordered the paths. . . . Our soil is very light and most suited to bulbs.'

The list of plants in Stonehouse's own hand was evidently compiled at different times. The first list, in strict alphabetical order, was completed on or shortly before the 27th of August, 1640. The names are very clearly written: ten lines on a page, on alternate lines for additions. The later entries are therefore readily distinguishable. They are in blacker ink, and may with confidence be referred to the next four years. Probably the greater number of new plants was added in 1644. Stonehouse estimates that he had 450 perennials in 1640, or 651 at the later date (probably 1644), as well as 215 annuals and biennials raised from seed, making 866 in all.

Then, no doubt, came the period of his persecution, banishment and imprisonment. A pathetic little note tells us that in 1652 he again visited his garden. It is in Latin and appears to mean that of the 866 plants, 'Alas! but few are there to-day, and I have no hope of founding a new colony'.

The lists enumerate 14 species from Virginia, 5 from Guinea, and 4 from New England, including *Juglans*, *Terrae glandes*, 'Maiz' and *Pepo*. Sir David Prain informed me that he knew of no earlier record of English garden plants from this last locality.

The plant lists have been printed *in extenso* in the *Gardeners' Chronicle* for 15 May 1920 and three following numbers, and reprinted with corrections.¹

xiii. THE WESTMINSTER GARDEN OF EDWARD MORGAN,
c. 1619-c. 1677.

Edward Morgan, 'rei herbariae studiosus,' a Welshman, was one of the *Socii itinerantes* who accompanied Johnson on his famous excursion to North Wales in July 1639, and acted as interpreter.² He then made the acquaintance of Walter Stonehouse to whose garden he contributed a plant of the Leopard's Bane of America '*Doronicum Americanum majus*' (1640). Morgan had a garden at

¹ Gunther, *Garden of the Rev. Walter Stonehouse*, 1920.

² Johnson, *Mercurii Botanici pars altera*, Lond. 1641.

Westminster, where he grew the first Polyanthus, obtained from Great Woolver Wood in Warwickshire, and possibly also *Phlomis purpurea*, which he is stated by Aiton to have been the first to introduce. His name is associated with the following British Plants by How in MS. notes which must have been written between 1650 and 1656. He evidently knew where to find white and purple Spear Thistles in London.

Primula veris fl. pl. viridi.

„ „ sive Paralysis fatua.

Malva syl: flore albo.

[*Bellis* fl. herbaceo globoso non descr. (*erased*).]

Scabiosa ovilla fl. albo.

Carduus lanceolatus fl. alb. St. James.

Lychnis syl: foliis variegatis fl. albo. Qu. M.

Carduus lanceolatus fl. alb. et fl. purp. Q. Chyrurg: for ye places of these plants growth from Morgan.

On 10 June 1658 John Evelyn 'went to see ye Medical Garden at Westminster, well stored with plants under Morgan a very skilful botanist'. In 1662 the garden was visited by the Rev. John Ward, who recorded in his diary¹ that Dr. Morison commended 'Ned Morgan's for ye best collection of plants in England', and the admiration appears to have been mutual, for Morgan affirmed Morison to be the best botanist of his age in London, and next to him are 'Dr. Dale, Dr. Merit, and Mr. Goodyer'. A few of Morgan's plants of special interest are noted in Ward's diary. They included

Leucojum bulbosum. 3 sorts.

Iris persica, a very pretty Iris, 'Ye flowers open with a mouth like Snapdragon'.

Cinara spinosa et aculeata.

Muscus filicinus, 'winged like ferne. Much of it gathered in Hamden woods'.

Gramen innatans and *fluviatile*.

and *Burri* (= *Burre grasse*), 'as it hath a yellow spike on top'.

1. *Ilex coccigera*.

2. „ *glandifera*.

3. „ *aktæ-foliis*.

4. „ ye common.

Medler tree.

Strychnos Nux vomica, ye Physic nut } dead.

Yellow Jessemine

Zizipha, ye Bead tree.

Glaux maritima.

Cruciata marina.

¹ D'Arcy Power, *The Oxford Physic Garden*, *Ann. Med. Hist.* ii, p. 111.

Absynthium arborescens 'July 3'.
 ,, insipidum et inodoratum, 'a very pretty plant, very like common wormwood, July 3'.
 A pretty hedge of Spanish broom.

In 1667 we again get news of the garden through Merrett who added five of Morgan's plant records to the Vegetabilia in his British *Pinax*.

Malva arborea marina nostras. *Lavatera arborea* L.
 'Mr. Morgan received it from the Isle of Wight.'

Plantago aquatica major muricata.
 'In a small pond betwixt Clapham and South Lambeth-Common. Mr. Morgan.'

Primula veris Polyanthos.
 'In great Woolver Wood in Warwickshire, Mr. Morgan, qui transtulit in hortum suum instructissimum Westmonasteriensem.'

Fungus campani formis, niger, parvus multa semina plana in se continens.
 'Mr. Morgan's Garden Westminster, call'd in Wostershire, *Corn bells*, where it grows plentifully.'

Fungus rotundus supernè concavus et translucidus coloris succini.
 'In Mr. Morgans garden, et alter coccinei coloris, in St. James's Park in the Winter time on old decayed Trees.'

Edward Morgan's *Hortus Siccus*,¹ comprising some two thousand dried plants, is contained in three large folio volumes of about 160 leaves each, now preserved in the Bodleian Library. The specimens give us a very complete illustration of the plants, which were probably grown in his Westminster garden in 1672, when the collection is believed to have been begun. At least two of Morgan's garden plants found their way into the Morisonian Herbarium at Oxford. One is the Basil-leaved Red Dead Nettle (*Lamium purpureum* L. var. *ocymifolium* Boulger), and the other is the *Phlomis purpurea* L. mentioned above. His name is also associated with 'Chamaecyse Virginianus E Morgan' (*Euphorbia maculata* L.)²

In 1676 the plants were removed from Westminster to the newly founded garden of the Apothecaries Company at Chelsea; and in 1677, 'Mr. Morgan, the gardener, asked for increased "consideration" for keeping the garden and for his plants'.

The exact site of Morgan's garden is not known. It appears to me that it may have been the same as the one by which RALPH TUGGY, the correspondent of Johnson, had previously made his name as a famous grower of pinks, carnations, and auriculas. Bobart referred to him as 'Tuggie in Westminster, beyond ye

¹ Noticed on p. 308. I am not quite convinced that Edward Morgan of Bodesclen is the same person as Edward Morgan of Westminster.

² Vines & Druce, *Morisonian Herbarium*.

Abbey'.¹ We know that Tuggy died before 1633, when his widow was keeping up the garden, but he is mentioned by Tradescant in 1629-30, see p. 331. A Richard Tuggy and Edward Morgan appear in 1657 as co-signatories to a letter printed by W. Coles at the beginning of his *Adam in Eden*.

On the other hand, Edward Morgan has to some extent been confused² with HUGH MORGAN, 'the Queen's Apothecary' and 'a curious conserver of rare simples', who had a garden near Coleman Street, where he had a tree of *Celtis australis* L. Coleman Street was of course in the City, leading north from Lothbury. Hugh Morgan introduced *Clematis viticella* in 1569, and is quoted by Lobel as having also introduced to English horticulture

Althea arborea Olbia in Galloprovinc. *Adv.* p. 294.

Buphthalmum, *Oculus bovis*, *Millefolium* folio, *Chrysanthemi flore*. *Adv.* p. 343.

Cicercula altera, an *Phaseolus* Diosc? *Adv.* p. 394.

Mount (see p. 256) mentions Maize and the Gladiolus in Morgan's garden in 1578. A panegyric advertisement of Hugh Morgan addressed to Dr. Bayley 'by your assured loving friend B. G.' and dated Alvingham, 14 Aug. 1587, is bound up with the Bodleian copy of Walter Bayley's *Discourse of the three Peppers*. Obviously Hugh belonged to a generation before Edward Morgan.

xiv. LISTS OF ENGLISH GARDEN PLANTS WANTED BY DR. ROBERT MORISON, A CORRESPONDENT OF DR. WILLIAM HOW, FOR THE ROYAL BOTANIC GARDEN OF BLOIS, c. 1651.

These three lists were evidently enclosed in a begging letter from an unnamed correspondent, endorsed 'For his much respected friend Dr. V. How'. They are written on 5 leaves, measuring 11 inches x 4 inches. The writer was evidently exceedingly anxious to obtain the plants. Of the 51, 92, and 70 plants included in three lists, 213 in all, How appears to have been able to supply the 42 indicated by numerals and crosses in his handwriting. It will be noticed that these include a proportion of new plants from Virginia 'not yet described'.

A comparison with the Morison MSS. in the Oxford Botanic Garden has convinced me that these lists are in the hand of Dr. Robert Morison, a pupil of the Frenchman Vespasian Robin

¹ Bobart MS. note in his Cat. of Oxford Garden at Oxford.

² Amherst, *History of Gardening in England*, 1895, p. 223.

who in the year of the publication of the *Phytologia* had been appointed Curator of the Royal Botanic Garden at Blois.¹

A chance presentation to Charles II in February 1660 resulted in his being offered an appointment as Regius Professor of Botany in London, to which he added the Chair of Botany at Oxford in 1669. In a book primarily about John Goodyer and his contemporaries, I have nothing good to say of Morison. He was practically a foreigner, and unlike Ray, he ignored the work of English botanists. In the words of Sir Arthur Shipley 'He seems to have been a somewhat selfish man of science, self-assertive, taking every credit to himself, while allowing little to his predecessors and contemporaries.'²

LIST I.

Semina harum primo quoque tempore ad me mitti desidero plantas vero tempestate idonea.

[Followed by an illegible note in How's hand] :—

Rememb. *Vina max. syl.* by sending in . . .

1. Abrotanum Indorum.
2. „ unguentarium.
3. Absynthium album umbilicatum.
4. Acinos anglica Clus.
5. „ „ fl. albo.
Anonis flore purpureo.
6. Armeria syl. humilis fl. vineo rubello.
7. Beta maxima.
8. Cachrys verior Galeni.
9. Calamintha pulegii odore.
10. Makenbuy valde aneo scire quid sibi vult et planta haec ipsiusque nomen.
11. Oxys lutea virgineana.
12. Chelidonium minus fl. pleno.
13. Juniperus major.
Cochlearia vulgaris Britannica.
„ vera rotundifolia minima.
14. Matricaria fl. Bullato aureo.
15. Mentha crispa.
16. Solanum lignosum virginianum.
17. Lysimachia lutea fl. globoso.

¹ It may be noted that while Morison was cataloguing his garden at Blois, an Englishman, Sir Richard Browne, then resident in Paris, was at work on a *Catalogue of Evergreens* in which he had been helped by a Mr. Keipe. In a letter to Sir E. Nicholas, dated 5 July 1658, he noted that 'Alaternes beare a graine like that of privet, which beinge sowed comes upp and prospers without difficulty'. *Camden Soc.* xxxi, p. 65. Goodyer had noted *Rhannus Alaternus* in Coys' garden in 1616.

² Schuster & Shipley, *Britain's Heritage of Science*, p. 234.

18. *Persicaria virginiana*.
Trifolium pumilum supinum flosculis longis albis non descriptum.
 „ *nodiflorum.*
19. *Salvia germanica non descriptum.*
20. *Saxifraga aurea sane sanicula guttata non descripta.*
Atriplex marina latifolia tota rubra non descripta.
21. *Periclymenum foliis quercinis non desc.*
Plantago quinquenervia fimbriis lat. ex aureo argenti. non descripta.
Acer floescens virginianum non descriptum.
 „ *odoratum virginianum non descriptum.*
 „ *virginiana non desc.*
22. *Apocynum virginianum latifolium majus fl. albicost non desc.*
23. „ *minus virginianum angustifolium fl. albo non desc.*
 „ *virginianum minus fl. elegantissimo miniato non descriptum.*
 „ *virginianum minus roris marini folio non descr.*
24. *Apium virginianum.*
25. *Arbor tinctoria virginiana non desc.*
26. *Aster Novae Angliae.*
27. *Artemisia (?) foliis variegatis.*
28. *Aster foliis (?) et flore minore Novae Angliae.*
29. *Battatas virg.*
Beta foliis variegatis.
30. *Bistorta minor nostrarum.*
Blitum majus repens fl. virid. Novae Angliae.
Calceolus Mariae.
Caucalis apii foliis floribus rubeis.
Digitalis flore variegato.
 „ *palustris hybernica.*
31. „ *virginiana angustifolia.*
32. „ *virginiana fl. luteo.*
33. *Lupinus perenni radice virginianus.*
Mentha variegata.

LIST 2.

Plantae excerpta ex *Phytologia Britannica* quas desidero aut saltem annuaria semina.

[This list contains 92 names, of which How supplied 7 plants only.]

34. *Anonis non spinosa purp. Ger.*
35. *Bardana rosea s. Lappa rosea.*
36. *Bugula s. Consolida media.*
37. *Cardamine pumila Bellidis folio.*
38. *Cerasus syl. fructu minimo cordiformi non descripta.*
39. *Foenum graecum syl.*
40. *Vina maxima syl. nondum descripta.*

LIST 3.

Plantae solis Johannis Park[inson] scriptae quarum communicationem aut seminibus aut radicibus peroptarem.

[A list of 70 plant names, of which How marked two with numbers.]

41. *Lychnis arvensis minor anglica.*
42. *Platanus occidentalis s. virginensis.*

Harum praedictarum tam in primo quam secundo, 3^{io} Catalogo contentarum semina penes te, ad me mitti primo quoque tempore non dubito quia hic apud nos appropinquat serendi tempus.

Quin oratum et exoratum te velim ut ex 400 sponte in Anglia provenientibu adjiciendisque aliquot rariora semina (quorum tibi est copia) ad me mittas.

[MS. f. 169-73.

In 1661 and 1662 Dr. Morison, established in the Keepership of the King's Gardens in London, was visited by John Ward.¹ Under Feb. 1661-2, Ward noted that, unlike Goodyer, Dr. Modesay 'never studied grasses and mosses'. 'Att ye King's Garden Dr. Modesie showed us some of his very rare plants as

Marum Syriacum.	Mastick.
Crithmum spinosum.	Samphire.
Barba Jovis.	Silver bush.

and the conversation turned on *Lysimachia purpurea spicata*, *Chelidonium majus* and minus, the Sycamore, and Rabelais. Ward evidently enjoyed the visit, for he repeated the experience on 1 Sept. 1662, remarking 'itt is a very rare thing to discourse with him'. This time he noted

Jacobaea crithmifoliis
Capparis Fabago
Cappadiis
Capsicum polygala
Valentina Clusii

and many other rare plants.

¹ D'Arcy Power, *Oxford Physic Garden, Ann. Med. Hist.* ii, p. 122.

LISTS OF EXOTIC PLANTS.

THE greater number of Lists of Plants in Goodyer's handwriting are of English indigenous and garden plants, and are written on odd pieces of paper of quarto width. Among them are several lists of foreign plants (now bound with Goodyer MS. 11), very closely written on narrow slips of paper, 3 or 4 inches in width, which have become crumpled, rubbed, and foul with dirt, as if they had been carried about in a pocket. They have all the appearance of hand-lists of desiderata, such as a London merchant might have furnished for the use of travellers visiting foreign regions, who required in compact form, catalogues of local plants with native names. Some of the lists were obviously restricted to tropical produce of economic value: others may be regarded as among the earliest exotic floral lists known in manuscript. The handwriting of certain of the lists appears to be that of John Parkinson, and their date to be about 1636, but several are no doubt earlier.

These early lists with their flavour of foreign travel recall Rauwolf's allusion to 'the great Troubles and Dangers those that have written of Exotick Plants have sustained and incurred in their foreign *Peregrinations*, would not be grievous to me, did I not fear that it would extend this Dedication to too disproportionate and tedious a length'.¹

FRANCE. *List of 51 plants taken from and with references to Lobel's Adversaria, written by Goodyer on the back of the draft of a letter dated 16 Jan. 1631. Not printed. [MS. f. 132 v.*

Glossary of French names of Plants, evidently abstracted by Parkinson from P. Bellonius De neglecta Cultura Stirpium (issued with Clusius Exotica), pp. 224, 226. Not printed. [MS. f. 156 v.

List of Plants, mostly from Italian Localities, abstracted from Camerarius, Hortus medicus et philosophicus 1588, presumably by J. Parkinson.

Abutilon Indicum.

[f. 162

Acacalis. Alisma Matth.

Adonis flore pallido in Thuringis segetibus e Suevia.

Alnus nigra, Frangula. Siler Plinii non multis(?).

Amygdalo persicus pulpa persicum nucleus amygdalum sapit.

¹ Epistle Dedicatory to Rauwolf's Travels in *Ray's Collection of Curious Travels*, 1693.

Antirrhinum pulcherrimum flore luteo, grandi, sponte nascitur inter Savonam et Genuam.

Apocynum verum non repens.

Aristolochia rotunda vera. In montibus Euganeis prope Patavium crescit.

Asphodelus autumnalis sive maior ex Italia missus : hyemem non fert nisi in fictili asservetur.

Camphorata Monsp. ¹[*Carduus bulbosus*.] *Card. pineus* Narbon.

Chamaelaea Germanica flore albo.

Chelidonium minus flore pleno. Francofurt. [ex horto A. Keck].

Cistus flore luteo Ledi [Ladani] folio Italici. Florentia.

Clematis altera flo: albo ; *Flammula repens* Dodon.

Clinopodium Matthioli flore albo.

Colchicum vernum subobscuro flore circa Tigurum Helvet.

Corcorus sive *Melochia*.

Corydalis Fumaria lutea montana. Splith.

Cotinus Coccygria Plinii frequens in Trident. Alpibus.

Cyperus rotundus. *Cyperus dulcis*, Trasi dulce.

Cytisus Columellae, *Glaux* quibusdam. Pisa et Samnitibus oritur et Bononiensi agro.

Cytisus niger sive maior *Pseudocytisus Cordi* elegans planta est, et tota aestate florens topiariis idonea.

Daucus Creticus verus in montibus Vicentinis, Baldo et monte prope Comum.

Elaphoboscus, Florentia.

Ferulago.

Flammula recta.

Galeopsis flore luteo, folio longo, rectis caulibus.

Halicacabum sive *Solanum Indicum*. Florent. [a J. de Casa bona].

Hissopoides ad portam S. Julianae [Justinae] Patavii & intra moenia.

Jacea pumila, seu *Carduus pineus* Narbāe.

Jacea Montana Narbonensis.

Juniperus maior Italica baccis rubris.

Libanotis cachrifera quae rotundis foliis prodit e semine cum pleraeque aliae umbelliferae oblongis oriantur.

Libanotis altera semine lato, nigricante, odorato.

Lilium non bulbosum sive *Arundinaceum* flore subcroceo et alterum luteo odorato admodum. Profert interdum germina in summitate caulis ubi flores sine semine deciderunt, quae terrae comissa statim radices agunt.

Muttelina ex Arba monte Helvetiorum ; inter Carum et Meum ambigua.

Paeonia ochranthemos flore pallescente.

Paeonia hispanica pumila.

Panax Heracleum in Apennino, in Alpibus Scalae dictis. non procul Bononia et in Samnitibus ubi Rampe d'orso veñ.

Panax Asclepias folio *Ferulae* tenuiore et *Foeniculi* crassiore. In Creta *Seseli* vulgo. Sicilia *Peucedanum* vocant. [f. 162 v.

Perfoliata montana *Gesneri* et *Lugdunensis* radice perenni, [foliis quodammodo *Limonii* seu *Pyrolae fruticantis Clusii* parfoliata quasi ca. . lis *Smyrnii*, floribus medio multiplicatis itemque et seminibus dense stipatis.]¹

Perfoliata crispa seu *mucosa* multiplici florum et foliolorum tenuissimorum farctum. Sterilis ut *Cannabis foem.* ex semine vulgari oritur.

¹ Not in Camerarius, *Hort. med.*

Polyacantha, raro in alterum annum durat.

Pulegium Anguillarae, ex Mentastro non multum dissidens foliis rotundis et gratiose odore. Plurimum nascitur Patavii tam extra quam intra urbis moenia.

Quamoclit planta nova ex India allata, et Ducis Florentiae horto Josepho de Casa bona.¹ [= *Ipomoea Quamoclit* L.]

Rosa Hierichuntica pulchrè ex semine nata.

Sagapeni planta vulgo dicta: *Costus spurius* Apuliae Matthioli. *Sagapenum* in Italia gigni. Plinius autor est sed in totum transmarino alienatum.

Saxifraga magna Matthioli; alia ab Hyrcina [sit *Pimpinella saxifraga*].

Saxifraga Bavarica in petris nudis musci more innascens, foliolis thymi sed virentioribus flosculo stellato candido quinque foliolis constante, elegant.

Scammonaea Antiochena aliquot annos apud nos durat asservata in fictilibus per hyemem, et aestatis initio floret, sed semen nunquam absolvit.

Scammonaea Monspeliensis maritima. Crescit etiam copiose prope Tripolim in maritimis ubi incolae vocant Meudheudi.

Scordium secundum Plinii, *Salvia* syl. Tragi. *Scorodonia* Cordi. Odorem habet *Scordii* nonnihil *Teucris* referens.

Scorpioides folio Bupleuri r. *Dodonaei*, *Thelephium Anguillarae*.

Scrophularia peregrina.

Sena Italica apud nos fuit annua planta. Alexandrina ne quidem in Italia ultra annum dura ideoque pro frutice haberi nequit. Contrahit nocte folia sicut multa alia eius generis leguminosa.

Sesamum verum.

Seseli Aethiopicum alterum folia fert *Aquilegiforme* similia sed multo ampliora: semen aliquantum *Cinamomum* representat odore et sapore. ex illustriss. principis Landgr. Wilhelmi Cassellano horto.

Seseli Massiliense. Lucas Chinus² excellens rerum simplicium censor, circa viginti plantas *Seseleos* Massiliensis descriptioni convenientes se novisse affirmavit.

Siler Creticum quo nomine ex Italia missum foliis *Cicutae* sed latioribus et paucioribus, semine incurvo, grandi, hirsuto et fusco.

Sison interdum ex semine provenit, pro *Amomo* habetur in officinis.

Solanum somniferum.

Sphaerocephalus flore albo.

Symphitum petraeum Neapoli missum foliis *Thymi* sed maioribus et longioribus nonnihil hirsutis aliquantum sed minus, odoratis, capitulis *Thymi*, flosculis purpurascens. [f. 163]

Telephium Matth. flore albicante et rubro. Radice instar *aloes* suspensa in aere crescit.

Thalictrum i. *Italicum* *Dodonaeo* tertium.

ii. *Germanicum latifolium*. *Dodon.* primum.

iii. *Angustifolium*.

iv. *Parvum* *Dodonaeo* quintum.

v. *Flore purpureo* *Dodonaeo* quartum.

vi. *Flore albo*.

¹ Joseph de Casa bona, Duke of Tuscany, corresponded with Camerarius. The *Aloc spinosa* sive *Americana* flowered in his garden in 1586, throwing up a spike to a height of over 12 cubits.

² Luca Ghini delivered botanical lectures at Bologna which were attended by William Turner about 1542.

- Thora Italica, Napello congener.
 Thymelaea multis locis Italiae in Monte S. Juliani Hetruriae et per Apuliam.
 Thymus verus capitatus seu Creticus in Italia circa Neapolim et in Syrcilia qui tamen annuus fuit.
 Trifolium Halicacabum sive Vesicarium aliis cicer sylvestre, quibusdam etiam satis inepte Dorychnium, folia plerumque quaterna vel quina gerit ut Loto annumerandum videatur.
 Trifolium hepaticum sive Aureum, Hepatica nobilis, flore rub. circa Corbachium Westphaliae, flore albo Cadelburgi in sylvis.
 Trinciarella ex Italia. Denti leonis planta similis sed tenerior, foliis quoque magis crispis, laciniatis gustu dulcibus, floribus minoribus et elegantioribus, semine oblongo, lato, membranaceo, superiore parte duriusculis aliquot pilis munito, eo ipso anno quo seritur flores et semen producit et si per hyemem in fictili servetur etiam in plures annos perdurat: alioquin radices quae dulces sunt ut universa planta primo frigore leduntur et intereunt.
 Valeriana Alpina multo odoratior, flores enim Citrii pomi odorem referunt [ex Allobrogum montibus ad me missa a C. L. Medico Ioanne Antonio Saraceno].
 Verbascum sylvestre foliis Salviae. Secundum Anguillaram in Italia provenit apud Marsos ad arcem quandam Pisanam dictam Suchamele incolae vocant.
 Viola speciem Bysantii coli Costaeus in Mesuem annotavit caetera Martiae similem, sed flore infinitis foliolis constructo et aequante Rosam Damascenam magnitudine et odore etiam nostratis superante. Huius recentem florem unum atque alterum prima mensa sumptum alvum subducere instar recentis rosae Hyeronimum Capellum Venetum rerum naturalium peritissimum observasse tradit.
 [MS. ff. 162-3.]

Andalusian and other Spanish Plants collected by Boelius and described by Goodyer.

Guillaume Boel in the year 1608 brought to Parkinson two Trefoils which he had gathered in Spain 'with about two hundred other sorts of seeds, besides divers other rare plants, dried and laide betweene papers'. By sowing the seeds Parkinson 'saw the faces of a great many excellent plants, but many came not to maturitie' with him and most 'by unkindly years that fell afterwards perished'.
Theatrum, 1108.

- Aracus maior Baeticus Boelii.
 Astragalus marinus Lusitanicus Boelii.
 Cactaria tuberosa radice Baetica Boelii.
 Caulalis maior Baetica.
 Chrysanthemum Baeticum Boelii inscriptum.
 „ tenuifolium Baeticum Boelii.
 Convolvulus coeruleus minor Baeticus.
 Faba veterum serratis foliis Boelii.
 Geranium Baeticum sp. Boelii.

- Gramen cristatum Boelii.
 „ tremulum maximum.
 Hieratium Narbonense falcata siliqua L'Obelii.
 „ stellatum Boelii.
 „ medio nigrum flore maior Boelii.
 „ „ „ „ minore Boelii.
 „ lanosum (?).
 Jacea capitulis hirsutis Boelii.
 Jacea palustris Baetica Boelii.
 Lagopus trifolius maior Baeticus.
 Lathyrus aestivus Baeticus flore coeruleo Boelii.
 „ „ edulis Baeticus flore albo Boelii.
 „ palustris Lusitanicus Boelii.
 „ aestivus dumetorum Baeticus Boelii.
 Legumen pallidum Vlissiponense Nonii Brandonii.
 Malva flore amplo Baetica aestiva.
 Medica maioris Baeticae species prima, spinulis intortis.
 „ „ „ spinosae, species altera.
 Nigella elegans ex Hispania.
 Papaver rhoeas Baeticum.
 Phalaris minor Baetica Boelii semine nigro.
 „ „ „ „ semine albo.
 Phalaris bulbosa Boelii.
 Pisum maculatum Boelii
 Scorpioides multiflorus Boelii.
 „ siliqua crassa Boelii.
 Silibum minus flore nutante Boelii.
 Sonchus Africanus Boelii.

Robert Tewe's List of 69 Plants from Russia, 1632.

- | | |
|---|------------------------------------|
| Glaux Diosc. Clusii | Coniza media |
| Genuina syl. pasticha | Campanula minor rotundifolia |
| Aster Italarum | Anthillis leguminosa |
| Chamaerododendros mont. | Gnaphalium minus montanum |
| Linum marinum luteum | Muscus clavatus |
| Scabiosa maior satorum | Muscus alter clavatus inscriptus |
| Saginae spergula | Clynopodium |
| Chamaelinum syl. flo. albo | Viorna |
| Eruca nasturtio cognata | Buphthalmion |
| Thlaspi narbonense | Gnaphalium Anglicum et
Belgicum |
| Helxine cissampelos altera attri-
plicis effigie | Cruciata minima |
| Hieratium alterum grandius | Thalictrum |
| Condrilla coerulea | Geranium batrachoides |
| Origanum hieracleoticum | Veronica recta herbariorum |
| Blattaria flo. luteo | Ledon foliis Rorismarini |
| Cannabis syl. spuria | Gramen iunceum marit. |
| Nucula terrestris | Milleum |
| Ranunculus tomentosus | Herba paris |
| Scrophularia maior | Lysimachia lutea |
| Pimpinella maior ffuchsii | Chamaenerion |
| Agriogiganum | Coniza helenitis |

Mellita incana	Anagallis aquatica
Astragaloides altera herbariorum	Ranunculus flo. albo
Hieracium Sabaudicum	Gramen parnassi
Veronica mas Matthioli	Gentiana minima
Helleborine seu Epipactis	Gentianella alpina
Dentaria altera herbar.	Monophyllon
Alabastris altera	Phaseolus ex phaseolis diversus
Rubus Idaeus	Polygonatum
Cicutaria latifolia	Ranunculus min. sept.
Coniza maior vera	Sorbus syl.
Aconitum luteum ponticum	Ribes nigra
Carduus pratensis Tragi	Ribes Arabum
Barbarea	Pyrola

& many other which I know not & are not to be found in my herball, I will have their effigies drawne & will hereafter send them you with their leaves & Russe names & vertues.

Mosco. 12 Junii 1632

Robert Tewe.

150 *li.* p. ann. 12 *li.* 10^s a moneth.

[MS. f. 135.]

Abstract of Adrian Spigelius' Isagoges in rem herbariam, 1606.

In the hand of Parkinson.

Pimpinalla ad radicem quadam in Polonia granula rubra gerit quibus lana purpureo colore inficiuntur. *Isagogis* Spigellii. fol. 15.

Cachrys pilula est quae Latino caret nomina, urendi vim habens in medecina. hanc arbores aliquot praeter proprium fructum pariunt hyeme durantem postquam folia cecidere ut Quercus, Abies, Larix, Pinus, Tilia, Nux Juglans, Platanusque; sed in Pino Juglandaeque saepius quam in aliis arboribus conspicitur. Is crescit hyeme deciditque, vere cum pilula aperitur. fol. 17.

Lysimachia purpurea trifolia Patavina Spigellii. fol. 12.

Sophonia Plinii nonnullis Blitum tricolor. fol. 24.

Ononis non spinosa Senensis capreolos, ut Vicia odore iucundissimo. fol. 39.

Hieratium flore purpureo. [fol. 41.]

Junci odorati caulis rotundus est, foliis alternatim obsitus, in quorum alis floras continentur, folia Gramini similia, cuius potius est species. Et cur Juncum appellatur miratur cum neque caulem, neque folia Junci habeat. [p. 42.]

Linum syl. fl: coeruleo, albido, rubescente et luteo.

Osiris flores edit luteos a fronte latiusculos, clausosque, os ranae imitantes: cauda angusta est et inflexa, ut *Consolidae regalis*; foliis Lini, qualia etiam in *Esula* minori conspiciuntur. [p. 43.]

Linaria odorata et Linaria Valentina Clusii quorum folia Lino non sunt similia sed flores tantum Linariae.

Vrtica foetida seu Galeopsis ideo dicta quia Galei id est mustelae faciem et rictum flores quodammodo referant.

Blattaria flore rubro.

Phlomis a foliorum hirsutiae appellatur.

[MS. f. 159]

[Continuation of MS. (68 lines) is not printed.]

Graccorum Turcarum et Arabum appellationes.

MS. f. 155.

[Abstracted from P. Bellonius, *Observationes*, translated into Latin by Clusius and printed with his *Exotica* in 1605. A Traveller's list of Native names, closely written by Parkinson on a folded paper measuring $7\frac{3}{4}$ in. \times 3 in.]

Plantarum Cretarum indigenarumque rarum nomina.

[This list containing about 125 names of plants was evidently abstracted by Parkinson from Bellonius, *Observationes*, *loc. cit.*, pp. 23-5.]

Acer. Cistus. Coris. Anagyris. Ferula. Ilex. Cupressus. Tithymalus dendroides, 2. hominem alt. Picea. Thapsea. Libanotis. Pinaster. Seseli. Dictamnus. Pseudodictamnus vulgo Cromido filo. Oxycdrus. Ferula, Artica dicitur.

Aristolochiae quartum genus Clematitide similis arbores scandens et premens.

Tria Origani diversa genera. Onitis peculiariter inter scopulos. Heracloticum contra humentibus locis gaudet. Sylvestre neutrius naturae particeps circa sepes crescit.

Tragacantha non gumnifera in Candia.

[MS. f. 157.]

[Fifty lines omitted. This list is followed by lists of about 20 plants growing in LEMNOS and 23 growing on MOUNT ATHOS. Both lists were derived from Belon, pp. 31 and 42-3.]

Plants of Gallipoli, Sinai, and Palestine.

[MS. 11, f. 158 v.]

[Practically a table of contents of the plants described by Belon, very closely and neatly written on the back of the paper with the Cretan List.]

List of Arabian and Indian Plants, Desiderata compiled for the guidance of a Traveller in the East.

The first eight items are not in Parkinson's handwriting.

Per Cayr et Alexandria.

Folium indicum is a leafe of a tree & hath 3 ribbes in every leafe it tasteth somewhat hott like cloves or spice: the Arabians in their language doe call it Gadegi Indi & some doe call them Famalapatra. The freshest in coullour & the quickest in taste is the best but remember the true have ribbes onely on every leafe both on the backside & the lowerside, and none true have none.

[Probably a form of *Cinnamomum Zeylanicum* Breyne.]

Costus is a certaine whitish roote that tasteth very hott & somewhatt bitter withall. The Arabians call it Cost or Cast & some of them Costi. If it be fresh it will smell very sweete. At Guzarate it is called Vylot & in some places Pucho.

[Root of *Aucklandia Costus*.]

Amomum is a certaine fruite that is clustering together having about 7 or 8 or 10 berryes on a stalke, every berry somewhat like a greate grape but whitish or yellowish & is skinnye on the outside & conteyneth divers browne seedes within them that smell very hott & are very sharpe biting the tongue. The Arabians call it Amana.

[One of the many kinds of Cardamons. Clusius received *Amonum Cardamonum* L., in 1605.]

Calamus gromaticus is a certaine drogue not very well knowen to us & therefore can give no true description, yet we understand it is to be had in those East partes: we would faine see the true. The Arabians call it Cassabalderira.

[*Acorus calamus* L., received by Clusius in Western Europe in 1574.]

Squinant is a certaine small rushe that tasteth hott like spice: the whole cuntrye of Arabia aboundeth as plentifully with it as other cuntries with grasse & rushes. The Arabians call it Adhar & Adher. The smaller & finer the rushes are, the hotter in taste & the better: the greater rushes are courser.

[Herba Schoenanthi vel Squinanthi from *Andropogon laniger* Desf.]

Acacia is the iuyce of the fruite of a tree called Akakia growing in Egypt & Arabia, thickened & made into small cakes very astringent & binding in tast. There is also a fine small white gum that is taken from the same tree Akakia wherof we would fayne have some. The fruite of the tree are certaine coddys wherwith they use & dresse lether in Egypt. We wold faine have store of those coddys which are like beane coddys to be sent unto us also.

[Gum of *Acacia arabica* was known in 17th cent. B.C. and is mentioned by Theophrastus, Dioscorides, and Pliny.]

Ameos is a small seed smaller than persly seede & tasteth somewhat hott, it is called Ameos I thinck every where.

[This is doubtless the species described as *Ammi parvum foliis foeniculi* (Park. *Theatrum*, 912) = *Carum ammoides* B. & H.]

Balsamum is a whitish oyle thin & cleere that cometh from Mecha & other places in Arabia gathered distilling from a tree there growing which they call Balessan of a sweete & sharpe sent somewhat like Turpintine but thiner & whiter, & if a droppe be put upon water it spreadeth on the toppe of the water like common oyle, & wil be whitish if it be gathered together from off the water with a little stick or other thinge.

The berryes of the tree likewise we wold have some store they are brownish round berryes with a sharpe pointe or end & smell somewhat stronge like the oyle.

The small sprigges & branches of the same tree Balessan likewise we want with smell of the oyle, if they be fresh are of a darke browne collour on the outside & are a little gumye & will stick to ones fingers espetially if they be fresh and new.

[*Balsamodendron Gileadense*.]

The remainder of the items were written by Parkinson himself.

Cate so called of most both Indians, Arabians and Persians is a thick blackish substance almost like Opium but a little harsh or astringent and bitter withall. The tree is full of thornes and is called by the Indians Hacchhic & we call it Lycium: we would faine have some of the beryes or fruite that growe upon this tree.

[Perhaps the *Lycium* of Dioscorides prepared from Indian species of *Berberis*. Parkinson spells the word Hattych].

Sandal of the Arabians & Chandama of the Indians is that wood called Saunders white and yellow.

[In 1635 the duty levied on the wood of *Santalum album* L., was 1s. per lb. on the white, and 2s. per lb. on the yellow.]

Caxcax is the greate heade of a Poppy that will hold a pinte of liquor.

[*Papaver somniferum* L.]

Imgu & Imgara of the Indians is a kinde of Gum or juice & is called of the Arabians Altiht & Antit: the plante from whence this iuice is taken is called Anguiden & Angeidan & in the Latine tongue Laser. If it were possible to gett some seede of the plante which is broad & flatt almost like unto a short rounde wing of a bird.

[*Ferula asafoetida*. Description taken from Garcias, Park. *Theatrum*, 1569.]

Calambac of the Indians of Malacca is the best Lignum Aloes: and Garro is the worst. The Arabians call this last, which is the worser, Aquilage & Haind.

[The best Aloes wood, for use as incense, comes from *Aquilaria Agallocha*, Roxb. Garo de Malaca is *A. ovata*. 'Selectissimi fragmenta' of this wood were seen by Clusius in England in 1581 in the collections of Hugh Morgan and James Garet.]

[MS. f. 165.]

List of Indian and Arabian, &c., names of Plants, and Vegetable Products of Economic Value.

Aloes, the Arabs & Turks call Cebur. p. 149.¹ Opium. Caphura or Camphora. p. 160.

Lacca called Luc Sumutri of the Indians of Pegu & Martaban that doe not use the Arabian wordes Luc. It is called Trec. p. 158.

Tabaxir is a drogue unknowen to us: it is like a graye gum or blackish sugar. p. 164.

Sambarane is a wood like unto Saunders. p. 174.

Macra is the barke of the rootes of a tree. p. 264.

Coru, Cura, & Curodapala is all but one thing.

Pavate in Malabar, of the Portug. Arbol contra las erisipolas. p. 266.

Cubebe be berries. p. 184. Araca is a fruite. p. 189.

Cocos be greate nuts as big as a childes heade. p. 191.

There be 5 sortes of Myrobalanes called after this manner by the Indians.

Aritiqui be those we call Citrini, some call them Arare. p. 194.

[*Terminalia citrina*.]

Anuale we call Emblici. These 2 sortes are espetially these we gladly would have ayther yong trees if it were possible or els the stoness of the fruite as fresh as maye be brought.

[*Emblica officinalis*.]

Rezanuale are called with us Indi Gotin be Bellerici. Aratca be Chebuli.

[*Terminalia chebulica*.]

Garcias ab Orta doth call them Canarini which inhabit those sea coasts of Decan and Guzarate (aunciently called Gadosia), lib. 1. cap. 2.

Tamarinde is a sowre pulpe of a fruite. p. 196.

Anacardii be certaine big beanes black and fashioned somewhat like an harte. p. 272.

¹ The page numbers refer to Clusius' *Exotica*, 1605.

- Spica Nardii. p. 273. Nardus Indica.
 Costus. Ambargrise. Meske.
 Rhabarbarum, we call it Rubarbe, a purging roote. p. 274.
 Radix China the Chineses call it Lampatan. The Turkes, Persians & Arabians Chophehma.
 [*Smilax China*, L.]
- Curcum of the Arabians & Saroth of the Turkes is that we call heare Turmeric.
 [*Curcuma longa*, L.]
- Galanga is of 2 sortes, one is a small reddish roote hott in tast & is called of the Chineses Lavandou. The other is a farre greater roote blackish on the outside & white within. The Indians call it Calcharu the Arabians call it Calvegian. They of Java Lancuare. *Park.* p. 1585.
- Moringa is a tree & the fruite of the same is so called also, but the Turkes & Arabians call it Morian, they at Guzarate Turiaa : the roote of this tree is a singular remedy against poysons. *Park.* 1650.
- Bezoar is a stone of greate valew if it be true & naturall & not artificiall as a greate many are that are brought frō thence.
- Negundo is a small berrye like pepper : the Turkes, Arabians & Persians call it Bache.
- Jaca or Panax is a fruite as big as a pompion set with thick shorte thornes but not pricking. *P.* 1639.
- Durione or Duriaen the fruite is as big as the Jaca & hath sharpe thornes on the rinde of it very dangerous to touch : it hath divers stones within it like unto Peache stones. *P.* 1640.
- Mangas is called of the Turks & Arabians Amba & is a delicate fruite. Another kinde herof called Mangas bravas is a most strong & violent poyson. *P.* 1631.
- Ananas is a delicate fruite like the heade of an Hartichoke the Portugals & Spaniards call it Pinas.
- Carcapuli is like an Orenge when the red outer side is taken awaye & the meate wthin somewhat like, but will not be separate in y^t manner.
- Carambolas the fruite is of the bignes of an hens egg or bigger, yellow on the outside with fowre ribs on the outside : in the middle it hath seedes that are of a sowre or sharpe tast yet pleasant.
- Jamboli of the Indians & Tupha Indi of the Arabians & Persians, is a most excellent fruite well smelling & of a pleasant taste of the bignes of a greate peare : there are herof 2 sortes one is of a darke red colour allmost black & for the most parte is without kernell, the other is of a pale red colour & containeth a hard white stone of the bignes of a peache stone. It beareth fruite divers tymes in a yeare & is never without flowers & fruite.
- Ambares is a fruite as big as a wallnut and of sowre yet somewhat tarte & pleasant when it is thorough ripe it hath within a hard gristly kernell.
- Bangue is a smaller seede then Hempeseed & of a darker colour.
- Pinones de Maluco so called of the Portugalls.
- Charameis the Arabians, Persians, & Turkes call it Ambela. They are of 2 sortes, one is of the bignes of an haselnutt with ringing corners, of a pleasant sowre tast. The other is bigger.
- Brungara aradua is otherwise called the Herbe of Maluca & is an excellent herbe to make salves to heale any sore. We wold faine have the seedes of it.

Tamalapatra be certaine leaves of greate use in Phisick wch the Arabians call Cadegi Indi & are like unto Pomecitron leaves but narrower towarde the end wth three ribs only running along the leafe & doe somewhat smell like cloves.

[MS. fol. 160.

[MS. continues with several items included in the list on f. 165, printed above. The descriptions of the last ten items agree closely with those in Parkinson's *Theatrum* and with those added by him to the last document].

List of Plants growing in the Bermudas.

In the handwriting of Parkinson.

The poisoned weede is like our English Ivie, being touched causeth rednes itching & lastly blisters wch after a while passe away without further harme. [*Rhus toxicodendron.*]

The red weede a tall plant whose stalke being all over covered with a red rinde therupon termed the red weede, the roote wherof being soked in any licour or a small quantitye of the iuice druncke procureth a very forcible vomitt & is generally used of the people & found very effectuall for the paines & distemper of the stomach.

The purging beane is a kinde of Woodbinde very cōmonly growing by the seaside & rūneth upon trees twining like a Vine: the fruite somewhat resembleth a beane but somewhat flatter, the which anyway eaten worketh excellently in the nature of a purge but very vehemently yet without any perill.

The Costive tree is a small tree.

Another plante like a Bramble bushe which beareth a long yellowe fruite whose shell is very harde & within it a hard berrye that beaten & taken inwardly purgeth gentlye.

Red pepper a fruite like our Barberies whose taste is terrible hot for the tyme if it be chewed betweene the teeth, & therefore it is swallowed downe whole & is founde to be of the same operation with red pepper & thence borrowed the name.

[*Capsicum* sp.]

In the bottome of the Sea upon the Rocks there is found growing a large kinde of plante in the forme of a Vine leafe but farre more spreade with veines of a pale red colour very strangely interlaced one into another which they call the Feather whose vertue is altogether unknowne.

There are since their plantation nourished white red & yellowe Potatos Pineapples, Papaws, Plantains, the America breade, the Cassado roote, the Indian Pumpion, Water Melon, Indicos etc.

Cassado a roote of a wonderfull encrease & maketh very good white breade but the iuice rank poison, yet boiled is better than wine.

[*Manihot utilissima.*]

Pineapples neare as big as Artichokes, the most daintye tast of any fruite. Apples, Prickly Pears & Pease, all differing from ours.

A Pepper groweth in a little red huske as big as a Wallnutt 4 inches long & small cods.

Two sortes of Cotton: the silke Cotton as in the East Indies groweth upon a small stalke as good for beds as downe: the other upon a shrub & beareth a cod bigger then a Wallnutt, full of Cotton wooll.

Locus tree, hard timber, 2 or 3 fadome about, of a greate height beareth a cod full of meale will make breade in necessitye.

[*Hymenaea courbaril.*]

Sope berries like a Musket bullet that washe as white as sope & goode to eate: in the middle of the roote is a thinge like a Sedge, a very good fruite: we call them Pengromes.

[*Sapindus*.]

A Pappaw is as greate as an apple, coloured like an Orrange & good to eate.

[*Carica papaya*.]

A small hard nutt like a hasell nutt groweth close to the grounde & is like that groweth on ye Palmetas: we call them Mucca nuts.

A greate tree whose leaves are used to make a Mustarde.

The Mancinell tree the fruite is poyson, it is an apple of a most pleasant sweate smell of the bignes of a Crab: the Swine & birds eschewe to eate of them.

[*Hippomane mancinella*.]

A tree like a Pine beareth a fruite so greate as a Muske Melon: hath alwayes ripe fruite flowers & greene fruite which will refresh 2 or 3 men very comfortably.

Plum trees many, the fruite greate & yellowe, which but strained into water will in 24 houres make very good drinke.

Wild figge trees the fruite feede the hogs etc.

The Gwane tree beareth a fruite so bigge as a Peare, good & wholesome.

Palmetos of 3 severall sortes.

[*Chamaerops glabra*, *C. excelsa*.]

Cedar trees tall & greate.

[The Pencil Cedar, *Juniperus Bermudiana*.]

Fustick trees the wood yellow, good for dying.

[*Maclura tinctoria*.]

Anotto.

[*Bixa Orellana*.]

[MS. f. 167.]

[This list, both in the order of the plants and in the wording of the descriptions, so closely resembles Parkinson's list of Bermudan plants printed in 1640 at the end of his *Theatrum Botanicum*, p. 1671, that it is probable that the printed version was taken from this copy in an abbreviated form.]

*List of Nuts, Fruits, and Seeds, desiderata 'To have from
Virginia, New England etc.'*

Ciprasse nuttes or young trees.

The ripe Ackornes of their White Oake or red scarlet Oake & Black Oake.

Black & White Walnutt & a greate one.

Cedars the berryes & the leaves, to see of what kinde they be.

Firre & Pine trees young & their apples.

Grapes white & red & a smaller black Grape growing in the Islandes, some ripe & sweeter.

Cherryes in clusters like grapes but not so good.

Black & Yellowe Plumes as big as a Damson.

White Thorne berryes are as big as English Cherryes & esteemed better than Cherryes.

A white Poplar tree with greate leaves broade pointed.
 Sassafras trees young & the berries.
 The Cassado roote or Jocka, Hiucca?. Sarsaparilla, all the sortes.
 The Maracoc Apples.
 The Maye Apples or seedes.
 The Snake-roote freshe plantes & seede.
 Virginia Lentills.
 Divers sortes of Lillias.
 Maidenhaire.
 Penni marva is Silke grasse.
 Musquaspen a small roote of a fingers bignes as red as blood wherewth
 they dye their mattes.
 Poconas is a small roote in the mountaines which they use for swellinges
 and so painte their faces.
 A small lowe tree whose white flower is like a small hedge Honysuckle.
 Puchuming are plumes sweete like dates.
 Red flat cods of a tree whose red flowers they use for salletts.
 Arbor Judae.
 The Locus tree the cods wth seede.
 Checinquamins are small Chesnuttes.
 A plante that beares a Scarlet flower.
 Flower de luces. Lillyes. Colombines.
 Or any other herbe or seede growing there allthough you thinck we have
 the same in England for we finde most thinges to diffarr.
 [Lentills of that countrye.]

[MS. f. 20.]

List of Seeds imported from Virginia, 1636.

Virginia seeds rec^d from Mr. Morrice¹ 18 March 1636.

1. A scarlet flower requiring a moist ground. *Cardinalis forte.*
2. A yellow wood called Locust long flat blackish browne pods, black
 round flatt seede kidney like.
 [*Robinia pseudacacia.*]
3. A poisonous berrye black rugged round berry with blackish seede
 straked like a tick.
4. Arbor Judae forte thin flat browne cods with somewhat flat round
 shining browne seede. Sent for a tree whose gredeline (?) flower
 is an excellent sallat.
 [*Cercis canadensis.*]
5. A black round rugged berry like large peper cornes with 3 square
 black shining seeds sent for a running vine y^t bears no flowers.

¹ Who Mr. Morrice may have been, we do not know. A Richard Morrice was Master of the Barber Surgeons Company in 1634. But there was also a Iohannes Mauritius or John Morris, a friend of Parkinson, who extolled the excellencies of the *Theatrum Botanicum*, and the worth of its author, in three pages of *Éloges*, in two languages. Two of his lines lend colour to our suggestion that he was interested in Virginian plants. Morris addresses Parkinson with the words

. . . neither dost thou spare

T'insert whatsoere the other world doth beare.

A remark which would come naturally from a poetaster who had also given Parkinson seeds of plants from the New World.

6. *Nux vesicaria* forte brownish bladders wth ij round pale coloured shining seeds like ours but farre lesse. Found on small trees in the woods.

[? *Staphylea trifoliata*.]

7. *Euonymus* forte growing on a dogwood tree: a reddish great seed wth huskes like it.
 8. A kind of Hawes much eaten by the Indians blackish flat kernells like the Pishamin: wthin black shining thin huskes with little or no pulpe.
 9. Virginia silke growing in a long white thick cod: flat browne seede with silken dowe *Periploca* forte.
 10. *Palma christi* brought thether & thrives well.

[*Ricinus communis*.]

11. *Maye Apple* seede a blackish browne seede lesse than *Stramonium*, the flower white & the fruit much eaten by them.
 12. The greate Snake roote small long yellowish seed like *Langdebeefe* but scalye & downy among, being greene it hath a milke more bitter than aloes.

[In the Morisonian Herbarium at Oxford is 'The great Rattle Snake root of Virginia, a rare Cordiall, Mr. Willis 1676'. It is *Asarum virginicum* L.]

13. A kinde of *Pescicaria* but higher by much then ours, a brownish black . . . flattish seed in whitish skin, huskes or skins.
 14. A kinde of double *Yarrowe*.
 15. Found in the woods. A pale browne pointed berrye set in a fine long-leaved huske like *Guaiacana* & 3 or 4 halfe round crested or straked kernells or skin.
 16. The berrye of a plante whose juice is good red incke. *Solanum Virginianum* sorte.
 17. Seede of the herbe *Carales*, so called frō a Captain that said it was much eaten in East India & in a tyme of scarsitye there also: a small black shining seade like *Amaranthus purpureus*, not *Milium nigrum*.
 18. Small Snake roote seede. N^o 7.
 19. A duskye roundish forme or solid graine as big as the heade of *Scrophularia*, with a white kernell in it: common in the woods.
 20. A long duskye gray pod small at the lower ende & bigger round & crooked pointed at the other opening in 2 partes with one rugged brownish pease there lying & the shewe of another at the bottome, but emptye. Found on small lowe imbracing vines.
 21. Virginia Lentills small short blackish thin cods somewhat thick on both edges or sides & much smaller black lentill-like seede within.
 22. Water melons flat darck graye seede like *Citrulls*.
 23. Red water melons flat darck graye seade like unto the other but a little longer pointed.
 24. *Stramonium* forte: it came without name.

[MS. f. 21.

APPENDIX

GOODYER'S MISCELLANEOUS PAPERS

GOODYER'S miscellaneous papers and letters have recently been sorted out and bound up in one folio volume (GOODYER MS. II). They include :

1. *Descriptions of new or rare plants*, mostly written on ff. 81-146, including many rough drafts of descriptions scribbled on the backs of other papers. See pages 109 to 193 of the present volume.

2. *Notes of Botanical Excursions*, 1617-1658.

Goodyer rarely allowed a summer to pass without an excursion to some part of south Hampshire, Southampton, the sea-shore at Hayling, or the New Forest. There is evidence of journeys to Essex in 1616, 1620, 1621; London, 1617, 1628, 1654; Wiltshire, 1618, 1624; Oxford, 1622; Northamptonshire, 1625; Surrey, 1634; and Bath, 1638.

3. *Lists of Garden Plants*.

4. *Lists of Exotic Plants*.

5. *Lists of Medicinal Plants and Materia Medica*.

6. *Sundry Medical Prescriptions*.

7. *Lists of Books*.

8. *Letters addressed to GOODYER and drafts of letters written by him*.

1618. Nov. 7. G. at the Red Lion in Fleet Street to some person believed to be Coys, describing the botanical results of his excursion in Wiltshire. p. 29.
On back. Addresses of Nicholas Everenden, Sedlescombe, Samuel Shute, Thomas Coltherste at the albus Lion in Deanside.
Last poem by Sir Walter Raleigh, 'Even such is time . . .' p. 32.
- n. d. GOODIER to person unknown. Concerning payment for Canary Wine on back of a Terrier of Durford Priory. p. 9.
1621. Nov. 9. LAURENCE DAVIS to JOHN GOODIER at Droxford. Concerning gold weights sent by Maye the carrier. p. 48.
1631. July 13. GRIFFITH HINTON to JOHN GOODYEAR at Maple Derham. Concerning the Bishop of Winchester. p. 56.
1631. Nov. 15. GRIFFITH HINTON to JOHN GOODYEAR at the sygne of the Angell near Denmark House in the Strand. Concerning fruit trees. p. 58.
1631. Jan. 16. G. communicating a list of plants indigenous in France to a traveller. p. 59.
- „ G. to MR. WRAY enclosing the above. p. 60.

1632. June 12. ROBERT TEWE in Moscow communicating list of Russian plants. p. 362.

1634. June 24. G. to MR. WORLIDGE about date of Surrey Assizes. p. 73.

9. *Other letters.*

n. d. JOHN PARKINSON to the worthy gentlewoman MRS. GEERES. Scribbled over and almost indecipherable. p. 266.

n. d. [DR. MORISON] to DR. V. HOW, asking for plants, of which he gives a list. p. 355.

n. d. DR. ARGENT to Lobel. p. 252.

n. d. J. DE MONNEL to Lobel. p. 252.

10. *Miscellaneous Papers.*

Of the Tenth mony due at Christmas 1608 received by Edward Cole and left with Roger Cole.

Thomas Hull received to my Lord's use the 22 of January 1608	iiij ^{li}
Mr. Myers received the v th of January 1608 to my Ls use	xxxij ^{li}
Thomas Hull received more upon my Lords noate 15 Feb ^{ry} 1608	iiij ^{li}
Edward Cole hath payd to my Lord in money the 29 of Aprill 1609	lxx ^{li}
Sum cxj ^{li} of Tenth	

Of the sixth payment of the subsedy due the xxvjth day of March 1909 (*sic*) & by him received in Swod.
Edward Cole hath payd to my Lord in mony

lxx^{li}
[MS. f. 127.]

Summons to John Rowland, Tithing-man of Ropley. 1614.

South.

Theis are in his ma^{ty} name to will and require you that ye uppon the receipt hereof you repaire unto me at Bishopps Waltham, or to some other of his maj^{ties} Justices of the peace within this Countie neare adjoyninge to take you Corporall oth for the dewe Execution & performance of the office of Tithingman-shippe for the Tithinge of Roplie, And hereof I require you not to fayle at your perrill dated at Newe Alresford under my hand and seale the xxiii of September 1614.

To John Rowland of Roplie
deliver theis.

[MS. f. 3.]

On back

[Description of a tree bearing Cachryes, p. 175.]

Terrier of Durford Monastery.

. Domum et Scitum omp monasterii de Durford cum ptinent, videlē duo prata, cont. p. estimac. quinq. acr. quorum un̄ vocat. le mill meade & alterum vocat. le kechyn meade; duo alia prata sine clausa vocat Smithes meadowes, cont. p. estimac. octo acras, un̄ aliud pratum sine clausa vocat. le wish meade af le wash meade cont. p. estimac. viginti acr.; un̄ clausū pasture vocat. le Brydgfield cont. p. estimac. duodecim acr.; un̄ aliud clausū pasture vocat. le Smithfield contin. p. estimac. duodecim acr., un̄ aliud clausū pasture vocat. le Rye cont. p. estim. sex acr., un̄ aliud clausū terre pastur vocat. le Conduit field cont. p. estim. viginti quinq. acr., un̄ aliud clausū terr. vocat. le North field cont. p. estim. trigint. quinq. acr., tria crofta sine terre pasture vocat. le Chapell Croftes cont. p. estim. triginta acr., duos campos vocat. le Rydgfields cont. p. estim. trigint. acr., duo alia crofta terre past. vocat. Robyns fields cont. p. estimac. duodecim acr. Ac tota illa terre Vastr. Bruera et rampua in Durford Hethe et Westherting Hethe, cont. p. estim. ducent. acr.; un̄ mesuagium sive Tenementum vocat. Hethe Howse in Petersf. in Com. Sowtht, dcto monasterio spectañ. cont. p. estim. centum acras.

Summary.

	Acres.	
2 prata	5	Myll meade & Kichin meade.
2 „	8	Smith's meadowes.
1 „	20	Wash meade.
1 clausum	12	Brydg field.
1 „	3	Calfes lease.
2 „	20	Dame Annys.
1 „	12	Smythfield.
1 „	6	le Rie.
	25	Conduit field.
<hr/>		
North field	35	
Chapell Croftes	30	
Rydg fields	30	
Robyns fields	12	
<hr/>		
Durford Hethe & West- herthing hethe	200	
Hethehowse	100	
<hr/>		
Total	[518]	

[MS. f. 6.]

[Dureford Abbey was a small house of Premonstratensian canons founded in the reign of Henry II, and suppressed in 1536. Its last days were passed in great poverty, the buildings were in ruins, the canons reduced in number. Layton, in a letter to Cromwell, describes the poverty of Dureford, 'which might better be called Dirteforde—the poorest abbey I have seen, as this bearer, the abbot, can tell you—far in debt and in great decay. This young man, for his time, has done well, and I have licensed him to repair to you for the liberty of himself and his brethren',¹

A seal of the Abbey is in the possession of Magdalen College.]

Draft receipt for £250 received by W. Inkerbie & Richard Bell from Sir T. Bilson. 1620. In Goodyer's handwriting.

Mem. that the xvth day of No: we y^e within mentioned W Inkerbie & Ric Bell have had and received of y^e within-mentioned S^r Tho. Bilson Knight the within-mentioned some of 250^{li} of lawfull Eng. money at his mansion house called Westmapledurham in the Countie of South by virtue of y^e within-mentioned power given us by our mother in law Philipp Bynwyn als Benwyn, and have delivered upp the within-mentioned deed according to y^e said power within given us to the said Sir Th. Bilson Knight: In witness whereof we here unto sett our hands, received the said 250^{li} and d[eliver]ed in ye presence of

Wel I have sent

[11 Januarij 1620.]

[MS. f. 11 v.]

[The INKFORBYE family fairly invaded Magdalen College in the sixteenth century. Of the four children of Andrew Inkerforbye, a Merchant of Ipswich, two gained Fellowships, one married a Fellow, and one married the President of the College, Lawrence Humphrey, by whom she became the mother of three Demies. William Inkerforbie, the brother-in-law of the President, held several College offices, and eventually accepted a Winchester Rectory in 1592, which he held with that of Selborne after 1596. It is difficult not to believe that he or some near kinsman was the William Inkerforbie mentioned above. Inkerforbie's wife is mentioned below, p. 382.]

¹ *Victoria County Hist. Sussex*, ii. 91.

Draft of Commission to Overseers of the Poor. 1621.

Overseers of the poore of the parish of _____ nominated and appointed by us this present yere viz the _____ daie of Aprill Anno dni 1621 shall by vertue hereoff levie all such somes of money as by them shalbe assessed for the releife of the poore within the said parish, And shall distraine and sell the goods of everie person that shall refuse to paie accordinge to their taxation rendringe the overplus to the ptie. Geoven at Southwick under our hands & seales the daie and yere above written.

[MS. f. 87 v.]

John Pitt of Chiddien in Hamelden [= Chidden, a tithing in Hambledon hundred] to have the peace against Elias Pelie of Hameldon [Hambledon was the village where 30 years later Charles II spent the night at the house of his guide, Col. Gunter's brother-in-law].

[MS. f. 87.]

F. Waller's Petition to Sir Thomas Bilson. Before 19 July 1621.

Most humbly Intreating yo^r good worship Si^r Thomas Bilson and to you & yo^r man ma^r Goodyer greeting: the Cause is this. my wryting unto you S^r is for to request & desire yo^r worships ayd and assistance concerning the two Buls and the fower Heffers w^{ch} the sayd Ilesley was committed by yo^r worship for, desiring for two know whether I may not lawfully have them prayed and so for two sell them, and (= if) it may be possibell good yo^r worship aid me in hit, for the Cause is thus I must step asid and away. for beecause freeman of Durley or of Frogmill have forth as I understand an Execution agaynst me for his counter band w^{ch} the sayd freeman hau not bine damnified one penny, but onely in suing for to bring in the bands, & soe forth wth arrested the sayd Ilesley's bease, and I was pledge for two bring him to his answer and could not paye the money being Eightenne pounds full, and tooke up the bands & went to the sayd freeman desiring for to have his Harmeley's band in to, and so to end hit, karying forthwith to hime his sayd Charges before divers witnesses, and hee would not for hee would hand the fantage or forfit of his counter bande being Six Score & Eight pounds and their of S^r I think it mete for two acquaint yo^r worship with hit. Other good men, for I must be droven to put forth my land and so to distribute my goods for a tyme untill I canne take som corse in it, I would yo^r worship should thinke that although I am forst to goe that it is for no misdemannors although I sayet my selfe nor for one penny or pound of debt but onely thus to be the cause meaning the Execution. Further mor sir I would request of yo^r worship for to know whether I may not send to the Sises some lawfull deputi, meaning some honest fend for to frame a bill of inditment against the said Ilesley. Requesting for two send me in two or three words in wryting and I would be much thankfull to yo^r Worship for hit, and so wth wepping teares I wayt these few words wishing yo^r worships health you & yo^{rs}.

Yo^r poore Oreator for to Command
fra Waller.

Deliver this to Sir Thomas Billson Knight.

[MS. f. 8.]

Deposition of Arthur Hyde relating to a distraint levied on Weston Farm in the possession of Sir Thomas Bilson.

Arthur Hyde of Weston in the parish of Buriton in the County of South^t yeoman aged fiftie eight yeares or thereabouts, deposeth that aboute sixteene yeares nowe last past, this deponent being tenant unto the late most gracious Queene Elizabeth of Weston farme, by reason of the recusancie of one Stephen Vachell Esquier deceased, duringe which tyme one of the Servants of the then gen^lall Receivoures of y^e said Queene Eliz. reaveannes, in the said County of Sowtht, came upon a peace of Lande of the said Weston farme, called the

Readons. And have distrained certaine Cattell of his deponente for Arrearage of rent of abowte fyve pownds, supposes to be done unto the said Queen Elizabeth for which this deponent entered into bande in the some of [blank] pounds, upon condicon either to paie the money, the then next Tearme folowinge, or otherwise to discharge yt, whereupon the Corte beinge moved, & fyndinge this deponent to be Tenante unto the said Quene Eliz. of all Weston farme, under a certaine Rent, by reason of the Recuzancie of the said Vachell, and that the said Reasone was but a small parcell of the said Weston farme, And that this deponent, by the covenants of his Lease during the Recuzancie of the said Vachell, was to be discharged of all Rents & charge payinge the Rente rearserved in his Lease, the said Corte therupon detayned the said deponents bande, & toke order for the staye of the leavyinge of the said Arrearage duringe the said Vachell Recuzancie. Sithence w^{ch} tyme the said Vachell beinge deade, And the said Weston farm beinge in the possession of Sir Thomas Bilson Knight, And then the said Weston farm beinge discharged of Recuzancie, there hath been leavied by the Sherif at sundry tymes the some of Nyne pounds & tenne shillings by xx^{ti} shillinge on ye pound only, under coloure of the said deponents bande, which was taken for the said supposed Arrearage. And yet none (notwithstanding so much payed, as aforesaid) theare is a newe seisure, diverted to the said Sherif for the leavyinge of abowte six poundes, owte of the said Lands called Readons, for the same Arrearage.

And this deponent further saieth, that he hath not beane Tenante, unto the said Readons sihence the ende of his said Lease, which ys now about some tenne yeares last past, and yet beinge tenant unto some other part of Weston farme, by reason of the said bande (w^{ch} was only taken for the said Arrearage of Readon) he this deponent with some other of the possessors of Weston farme, hath alreadye payed Nyne poundes & tenne shillings, besides twoe shillinge at any sevall tyme of seisure, to the Sherif his Bailies, w^{ch} cometh unto xviii^s more.

[MS. f. 5.]

[Weston is a tithing in the parish of Buriton which seems to have been, to some extent, co-extensive with the manor of West Mapledurham. According to the authorities quoted in the *Victoria County History of Hampshire*, iii, p. 90, Stephen Vachell was part lord of Weston in 1579. In September 1600 he forfeited two-thirds of his lands and possessions for recusancy, and in December 1600 the queen granted the capital message called Weston farm and lands in the parish of Buriton to Arthur Hide, for a term of 21 years. It is doubtful, however, whether Arthur Hide ever gained possession of the manor, for in 1598 Vachell and others conveyed it to Nicholas Hunt and his wife, from whom in 1607 Thomas Bilson, Bishop of Winchester, purchased it. After the purchase Weston formed part of the manor of West Mapledurham.]

Draft of a Petition from the Tithing of Weston concerning payments towards repair of Redbridge. In handwriting of John Goodyer.

The names of those who were to goe to paie viij^s towards the repairinge of Redbridge beinge required of the Inhabitants of this Tything, after they had paid their proportionable hereinafter said rights as anciently they have don.

S ^r Tho: Bilson	John sen.
Hen: Bl:	Tho: Wyse
Hen: Vok	Tho: Kent
Jo. Tayler	W ^m : Hall [or Hull]

who made with one accord this answer, that the inhabitants of the Upper half hundred of Finchdeane have alwaies hetherto paid two parts and the Lower half hundred but one pte of the said impostes uppon the sd Hundred, towards the repations of the sd Bridge, and that they of the sd Tything of Weston have latelie paid [blank] beinge their pte of the said third parte.

And that nowe they refuse to paie the sd 8^s beinge a some imposed in them, apportionable with the somes imposed on the other parts of the before said lower half hundred, . . . the lower half hundred equall in payment with the *Upp* , which they never did. And that they pray to have their case heard

this next mich^{as} Sessions when they hope to obtain an order that the lower half hundred shall paie but a third part thereof as anciently it did.

And this rate now imposed, being not after the ancient manner, but an increase of charge to the lower half hundred by making them equall in payment to ye upper half hundred, whose lands are almost twice as good, the tything of Weston being a small member of the lower half hundred, humbly pray they may be heard this Sessions what they can say with the rest of the lower half hundred for themselves.

[MS. f. 12.

*Portion of a deed concerning lands at Bramshott and Kingsley
in which John Goodyer was concerned.*

This Indenture made the 15th
God of England, Scotland France and Ireland

[Dorothy his wife of the one part and] ¹ John Goodyer
[pounds] ¹ of good and lawfull money of England unto
the ensembling and delivery of these presents
[Release and Confirmation bearing even date] ¹
[Sambrooke of London knight and Administrator] ¹
[payment receipt and Satisfaction of which said] ¹
[doth acquitt release and discharge the said John Goodyer his heirs] ¹
bargained sold and confirmed and by these presents doeth grant b[argain sell and confirm
and *Parcefeild* with all and singular the rights members and app
gardens Orchards Courtyards and backsides thereunto belong
Bramshott and Kingsley some or one of them in the said County [of Southampton
quantities of acres hereinafter menconed (that is to say) the planted
estimacon Tenn acres three quarters of an acre and eight perches
One quarter of an acre and six perches *Great frith* containing by e[stimacon
acres three quarters of an acre and thirty two perches *affeild*
thirty perches *Garden feild* containing by estimacon eight acres
perches *Great Waterfeild* containing by estimacon Tenn acres &
by estimacon Six acres and halfe an acre and thirty two perches *Gr*
estimacon twelve acres and a quarter of an acre and sixteen perches
two *Heathfeilds* containing by estimacon one and twenty acres and
and Sixteen perches *Upper Wassells* containing by estimacon flour
Lower Mead containing by estimacon eight acres and two perches
three quarters of an acre and twenty perches *The Green* before th
acres *Upper Mead* containing by estimacon three acres *Middle Me* [ad containing by estimacon
acres and an halfe of an acre and eighteen perches and also all
and wood ground thereunto belonging and also all those severall
called *Woolmore* containe together in the whole by estimacon Sixt
Mead and one Cottage with the Land thereunto belonging scituate
Closes and grounds are severally and respectively called or known and
possession of the said Andrew Wall and also all that dwelling house
and all those severall Closes or grounds severally and respectively cal[led or known and
likewise menconed that is to say *The Upper Six acres* containing by e[stimacon
thirty two perches the *four acres* containing by estimacon flour acres
Homefeild containing by estimacon six acres and a quarter of an
containing by estimacon eight acres and an halfe and one and twe
perches *Hillfeild* containing by estimacon flour acres three quar
tenn perches *Outward Heathfeild* containing by estimacon seven
. . . . *Green feild* containing by estimacon seven acres three q

[This portion of a deed, in which Goodyer's name is mentioned, was used as a wrapper of a document belonging to Mrs. Ruck-Keene of Swyncombe, Oxon., who, discovering it at the very moment of the discovery of Goodyer papers at Magdalen, has kindly permitted its publication with them.

In the *Victoria County History for Hants*, ii, p. 493, are notes that Ludshott Manor was acquired by Andrew Wall in 1638 from the Knight family (Feet of F. Hants. Hil. 13 Chas. I), and that in 1704 Samuel Diggle acquired the manor of John Goodyer (Feet of F. Hants., Mich. 2 Anne).

A Sir Jeremiah Sambroke, Knight, Merchant, was on board the Earl of Barkley's ship 31 Jan. 1681 and died 27 Apr. 1705.]

¹ erased.

Goodyer's note relating to Rents in Wiltshire.

Chappel. Crine.

Richard Bedelcombe of Weeke in Dounton parish, aetat. 80 vel ultra, gathered of Henry Davys of Chestgrove in Tisbury parish in Wilts, aetat. 72, who was tenant to Edward Willowby gent, yerely as a Cheif Rent vjs. viij d. He was gatherer as longe as M^r Barrowe was lord. Mr. Willowby never denied the Rent, but denyed y^e service to Mr. Estroud in y^e open Court.

Mr. Willowby sould it to George Stable gent, whether he paid it or not, they know not, he byd M^r Dawley shew wherefore he should paie it.

Mr. Staple sould it M^r Westfeild of Tilsford 7 or 8 mile beyond Warminster. Hester Abell widdowe in the High St. at Salesbury nere to the Close gate.

[MS. f. 5 v.]

Draft of an Order concerning the discharge of the Watch of the Beacons in Portesdown Division. In the hand of J. Goodyer.

Whereas I have received letters from Sir R. T. knight one of ye deputy liutenens of this County of Southh. concerning the watch of the Beacons within this Division of Portesdowne, which Beacons must be well kept & repaired, and the watch of them to be presently discharged. Theis are therefore to require you to cause the watch of ye Beacons within your hundred (if there be any) presently to be discharged, and y^e rest performed according to y^e contents of ye said letters,

And hereof faile not as you will answer the Contrarie (?) at your peril. Dat. 7.

[MS. f. 5 v.]

[The Sir R. T. would have been Sir Robert Tichborne, who was living in 1617.]

Goodyer's rough notes of Accounts 19 July 1621.

James Rabson of Hamelden Carpenter 10^s due onto Katherine his wiffe, for wages long since due.
Stephen Woley.

hog rec ^d y ^e last year	16	} 112	
rec ^d this year	96		
hog spent this year	96		
recd as appeareth	16		Wheate spent 166 ^h
			Otes spent li. 196
has rec ^d ye last yere	137 ^h	} 329 ^h	Joh R
bought this yere	192		
spent this yere	275 ^h		
recd as appeareth	54 ^h		

William Browne de Hoe infra parochia de Subberton ¹ gent, To paie lxxij^h x^s the 11th of November next.

Will^m Hatch de Clanfeild yeoman } bondeth John Parker in 20^h

Simon Hatch Blacksmith

To paie 11^h the 19th of July 1621.

W^m Hatch bondeth (?) S. k: in 40^h to store harweste.

[MS. f. 8 v.]

[Then follows a Description of 'Ye lesser Bellflower'. See p. 132.]

¹ [The Manor of East Hoe, Soberton, passed to W. Browne, senior, of Hoe in 1619, then to W. Browne, junior, his son.]

Note concerning certain inhabitants of Soberton c. July 1622.

Willmus Arnold de Soberton yoman. v^h
 Willmus Perrin de Subberton husb. v^h
 Johes Lower de Soberton husb. v^h
 That Margery Lower to keape y^e peace ags^t Susan Talbott spinster.
 [MS. f. 7.]

Assessment in Goodyer's handwriting. After 9 Nov. 1621.

	[Acres?]	[£	s.]				
S ^r Tho. [Bilson]	433	3	3				
Kent	125	1	0				
Chaw	56		5				
Petter	23		1				
Ra: Tribe	51		4				
T. Deed	20		1				
Wm. Tribe sen.	36	3		winnowed	17	7	3 ... 1 g
„ jun.	15	1		spent	7	1	2
John Tribe	37		3				
Blackman	63		6		10	6	1 ... 1 gall.
Silvester	56		5				
	915	7	3				

Anne the wiffe of John Ilsley of Prelingworth in
 T husb. to have the peace against Tho: Ilsley.
 [On the back of Laurence Davis' letter, dated 9 Nov. 1621.]

[MS. f. 9.]

Note of proceedings at a Court held later than 9 Nov. 1621, affecting persons living at Swanmore and Chcedden.

Edward Collyn de Swanmor husb:
 . . . Tho: Cludlie de eadem Carp. in 8^h
 To paie 4^h 8^s: 20 Sept. 1622 in Tho: Cludlie's house.
 W^m Drewe of Hamelden served Thomas Pina and John Rynd and y^e rest of
 y^e parishioners of Chcedden as their comōn shepherd.
 Nic: Brittaine de Swanmore husb. . . . Jaspē in 20^h.
 To paie 11^h the 20th Sept. 1622 in this C[ourt] house.
 [A Botanical Fragment. Description of *Thlaspi* 14 July 1621.]

[MS. f. 11.]

Descriptions copied from Lobel's Adversaria of four plants. Perhaps in preparation for a botanical excursion to Portland.

Anthyllis
Sedum Portlandicum
Hederacium Thlaspi
Papaver Cornutum.

[MS. f. 13.]

Goodyer's Notes 1625.

The 2 of May to have 100 Be , 8 foot longe & 4 foot 7 ynch lenght.

Robt Westbrooke¹ of Steep husb.

W^m Hinde of Sheet

3 teneri pr. to paie 16^h 13^s 4^d the 22 of July next.
 & 16^h 3^s 4^d the 21 of December 1625.

Ewie sat about 4 a clock to take fagotts.

[MS. f. 13 a.]

[Below: the itinerary of a botanical excursion for Sept. 1624. See p. 51.]

¹ Robert Westbrooke was a tenant of Magdalen College in the Spain Petersfield, about 1580 (Magdalen College, Ledger G. f. 10).

Rough notes made by Goodyer in November 1631.

The names of those w^t were taxed to paie the muster Master's entertaynment.

Bilson 2 corsletts 2 musketts—at 9 ^d 6 ^s	} 12 ^s
Hen. Vok	} for 1 corslett 1 6	
Tho. lee		
Tho. Kent	} for 1 corslett 1 6	
Jo: Trybe		
Jo: Pinke	} for 1 corslett 1 6	
W ^m Hall for 1 corslett 1 6		
Jo: Horwood for 1 muskett 1 6		

Of all those above named I have demanded the somes they were taxed at
And they all do answer me they ought not to paie it.

Dielt at Gilford 13 ^s 8 ^d	} 8 ^s 11 ^d	7 hors at hey at 8 4 8 coch — ib } 4 has — ib } 5 4
Otes—2 ^b 5 4		
hey 2 1		
greas for y ^e Coach 0 6		
gave the hosler 12 ^d		
ferre at lambheath		
Coach 12 ^d	} 3 coch harnes (?) 1 coch bridle	
5 horses 10 ^d		
Supp. at Kings head 10 ^s	} 1 ladie sadle, 3 g, 2 snaffle bridles	
	1 14 5	
	3 8	

rem. 3 3 11

Edyn Haslett? 22^d

2^s
Nutts? 1

[1631]

Nov. 7. 2 bauhins 4^s
8. wyne wth Johnson 6^d
Diosc. 18^d
Bruelc [?] 2^s
Thevett 12^d
14. beane there 2^d

Leo 1597.
Scholzius Aphorismes
Ludovicus Septalius, 1625.
Duncanus Liddelius
Plate[ar]ius
Wekerl Syntaxis 1576
Petri Bayri Enchiridion, quae Veni
mecū vocant. 1578

interest comes in his wife's use.
Mr. Betts—whether Burnetts generall release during his owne life (?), the
[MS. f. 12.]

Roll of Armour-bearers, Pioneers, and Spearmen of Buriton.

(Before 1633. Not in Goodyer's hand.)

Beritun

The earmur bearers.

Robberd Heath with a muscet.
and Jasper Hillis with a muscet both of M^r Watkers.
Randol Jouning for John Benet his muscet.
Richard Searl with a muscet.
Neckolas Greenwood with a muscet.

- ab. Thomas Linter.
- ab. John Pinssers.
Steeven Pit.
- ab. William Barker, the elder.
William Tayler.
- ab. Richard Cheas.

The names of the Peionors.

[Larence Plidger and Richard Paige] (erased).

John Barker, hors post.

John Pitter, foote post.

Vitllers John Luf and Richard Plidger the younger.

Carters Arter Cornwel, Benge Paige and Henry Burch.

Spear Men.

musk. James Poell.	musk. Thomas Paterick.
— Peeter Heath.	Larenc Brasier.
— Umphery Biges.	ab. Larenc Pinssers.
— William Parker.	ab. Robberd Holt.
musk. John Weiet.	ab. John Hide.
ab. William Chrispe. }	— William Barker the younger.
ab. Francis Miles. }	— Neckolas Bokkis.
ab. William Hide. }	— James Bate.
ab. Andrew Hickman.	— Thomas Horn.

[MS. f. 12 v.]

THE NAMES OF THE SPARE MEN WITHIN TYTHINGE OF [SHEETE] (?).

Henry Beale.	Nicholas Polen.
William Whitcher.	Humfrey Deane.
Robert Whicher.	Thomas Jacob.
John Stummer.	Arthure Tanner.
Robert Wells.	Edward Polen.
[William Hide] erased.	John Wiat.
John Inwood.	Thomas Yalden.
Robert Brigger.	Thomas Hunt.
Robert Betsor.	

Rychard Jenman,
Tythingman.

[MS. f. 137 v.]

List of ten Hampshire Women concerned in the payment of certain sums of money (? early 17th cent.).

	ii	
Goodwife Hardinge of Hakle	iiij	of medlee
„ Gaman of Hakle	ij	of medlee
„ Sharpe of Steepe	ij	of medlee
„ Eames of Lesse	iiij	of medlee
„ Tyler of Leese	ij	of medle
„ Chilese	ij	
„ Rose of Leese	ij	
„ Wheatle of Leese	ij	
„ Veryam of Hearting	ij	
„ Bakere of Frokesfeeld	iiij	
Margerie Winnige	ij	

[Written on blank leaf in the binding of Goodyer's copy of *Crescentius*.]

Goodyer's Notes relating to a shopping excursion to Winchester.

Edward Williamson of Lippock Butcher to release to Richard Costen of Shidfield,¹ husb., all actions.

Seaminge lace	ij yards	16 ^d
Edginge lace	1 yard	8 ^d
Edging lace	1 yard $\frac{1}{4}$	10 ^d
Lace for ij Capps	j ell	10 ^d
One ell of Halland for ij Capps		18 ^d
		5 ^s 2 ^d
Caried to Winton		20 ^s
S. money		3 ^s 1 ^d

[Goodyer's rough draft of a description of the Broad-leaved Elm is on the same sheet.]

[MS. f. 4.]

¹ Shedfield, South of Bishops Waltham.

Goodyer's Memoranda for shopping for a lady, probably in London.

c. 1621.

20 yards of galome lace.

4 doz. buttons.

Homs (?) fustian 2 yards 3 qters.

Canvis j ell.

8 skeynes of sticching } half an ounce in all will serve.

8 skeynes of sowinge }

2 oyle skins.

half a yard of Jeynes Fustian.

Cambrick j ell & qter at 3^s 6^d or . . . q^r ell.Mr^{rs} Benwyn at the little Minerier neare Algate. if she wil paie to gev a discharge from Holman, windmil (?) and Inkforbies wife.

The fan of a pinke color. Morgan to doe.

8 yards of 4^d rebon of y^e same pinke color, to desire Mr^{rs} Ison for a patterne of ye bredth & color.6 yards of 2^d rebon of y^e same colour. The fans handell to be doble gilt.Mr. Bull was pd 5^s for 5 q^{ters} ended y^e 30th of Januarie 1621.

Mr. Wilson of Barbican.

50 needles of all sorts, none to bigge.

a girdle of stuff 18^d pise russet color.

a pound of Annis seed comfitts.

half a pound of English liquerish.

[A list of books on same page.]

[MS. f. 46.]

Memorandum relating to Corn. 1631.

1631.			
Wheate		in y ^e lobell	13 ^b
of Hughes	20 ^b	Little lobell	3 ^b 2 ^a
of Voke	14 ^b	(?) Jemox	23 ^b 2 ^a
		Little lobell	3 ^b j ^a
Win	4 ^b 3 ^a	(?) Jmox	2
Oct. 6	4 ^b	Sept. 29 Mill	2
" 17	8 ^b	Oct. 6 Mill	2
50 ^b 3 ^a		Oct. 13 Mill	3
		Oct. 20 Mill	2 — 2 ^b 2 ^a

[MS. f. 132.]

An account of purchases, probably for his wife, in 1634.

1 yard ha. & ha. qter of white Cotton at 12 ^d		1	7 ¹ / ₂
1 skyn		1	1
3 yards of fustian Rebon		0	6
10 paire of hooks & eyes		0	2 ¹ / ₂
Canvis in the wostbud (?) & bustian		0	2
Silk		0	2
6 Buttons		0	1
		<hr/>	
		3	10
Making the hose	1	2	
lengthning the other		8	
		<hr/>	
		1	10
		<hr/>	
		5	8

[MS. f. 14.]

Accounts 1646-50.

Paye 7: y: behind 4ⁿ 13^s 4^d of this
 Rec^d 1646: 6^s 8^d and in May 1647 1ⁿ
 payd all but since Michaelmas 1649.

Stocker oweth 1 18
 besides this half 1 8
 Tomlinson d 13 9
 1ⁿ 18^s
 ded: 1ⁿ 12^s 6

Midsommer 1650.

[4 lines illegible]

Knight 0 4 0
 Wayd 0 1 0
 Bold 0 13 4

A Long (?) ded: 4^s 6^d
 ded: 1
 Received 2 1 7
 owes 4^s 2^d to Keens to
 Paied 1^s 8^d, do a rent 3^s 1 17^s 8^d
 Collins ded. 13 4
 Baker on high 0 14 0
 ded 12 4^d
 Cariage payd 2 2 6
 Rd. Eomes ded: 3^s 2^d &
 4 11 and
 L. Rent 1^s 3^d 5 0 0
 Eomes last ded: 1^s 6^d
 till the R day

This half ded: 10^d 1 0 0
 Hall to ded: 2^l p. annum 0 15 0
 King 0 2 3
 Anthony Corper 0 1 1
 Isaak Knight 0 3 4
 Girdler 0 3 4
 Thomas sen. 0 11 6
 John Pescod 0 6 8
 R. Rogers Life 0 15 0
 J. Roke behind 0 yr
 besides this halfe 0 1 8
 Lady day 1650.

PRESCRIPTIONS AND MEDICINAL NOTES.

Goodyer's views on medicine were evidently those of his senior by nine years, Lord Herbert of Cherbury. Lord Herbert conceived it a 'fine study, and worthy a gentleman to be a good botanic, so that he may know the nature of all herbs and plants'. A gentleman should know how to make medicines himself.

Tonic for a horse: a snuff: purchase of tobacco.

880. e. Anisum
 557 g. calamintha
 702 f. n. violets
 1128 c. Spartium sp: broome:

Richard Skillinge: 1^l 1^o Feb:
 Muscadine 1 pint } pound ye sugar candy
 Sugar candy an ounce } & putt in ye yolks
 Yolks of Eggs iij } with ye musc. & mak it luke warme & geve it your
 horse when you travell.

Euphorbius putt into ye nose to sneese.

Tabaco 4^d
 Pulletts 4^d
 Alresford 4^d

[MS. f. 4.]

¹ The Skillings were a well known Hampshire family. A Richard Skillling sold the Manor of Lainston in 1613 (*Vict. County Hist. Hants*, iii, p. 445).

Ointment for the Spleen, c. 1621.

Unguentum et emplastrum per liene

℥ olei de capparibus }
 de absinthio } anā ʒvj
 amygdal: am: }
 aceti scillini }

Misceantur probe, et inbregatur plen per dies 7 mane

et vespi, dein apponatur emplastrum sequens

℥ gummi amoniaci clari in aceto scillini solut

piperis nigri

olibani puri ana ʒvj

In mortario calido misceantur et applicetur parti calide.

Put a little vinegar in a hot mortar, and you putt the amoniacum and stirr it with a pestell and it will quickly be a plaister.

Mr. Michaell Easton an apothecary on y^e south side of y^e greate conduit neare Cheape Side And pray him to have the trewe Ammi.

[MS. f. 46 v.]

List of plants.

Beta candida	Condrilla rara flore purpurante. ob.116
rough Hyssope. Hyssopus incana	Aster conyzoides Virginiana
Acinos odoratissimum	Helianthemum radice repente Vir-
Angelica	ginianum
Auena nuda	Ephemerum Virginianum
hawes of glassenbury thorne	Sumach vir.
Astrantia nigra	Sium segetum & agrorum, Honewort
Eruca peregrina Clusii	Cuminum Candia
Cochlearia out of y ^e north	Seseli massilense. thapsia.

[MS. f. 46 v.]

Prescription.

℥ conserv. florū fumariae et } ana ʒfs
 Radicum cichorei }
 pulpae cassiae } ʒiij
 confectionis hamech } anā ʒv
 diaprunis solut }
 pulv. subtilissimi foliorum Sennae } ʒiij
 tartari albi } ʒjfs
 pulveris electuarij triasantali } ʒfs

Misc & cū syrupo de citron compos.

cū Rheo fiat electuarium instar opiatae.

[MS. f. 128.]

[Goodyer has copied out this prescription twice over.]

Recipe, c. 9 Feb. 1627.

Hoggs grease } of ech 2ⁿ
 Sallctt oyle }
 Litarge of gold . . . ʒⁿ
 green Amoniac } of ech 1ⁿ
 Opopanax }

Boyled & made uppe by art into a plaister, the gummes must be clarified wth vinegar & made uppe into a true body.

[MS. f. 129.]

*Recipe: fragment of a petition: note on ailments of John Neale:
note on local maladies of Selborne.*

20 Mar. 1662.

Rosen that is blackish
Fresh lard a wallnutt
Crowne sope a wallnutt
Boil till it sets clere. H
keepe stirringe.
Elder
Som

[This recipe is written in a shaky hand.]

Mar. 22. John Neale in Lippock, the howse is called Gurmes, hadd a third
Ague, & hath lost him about a moneth & nowe hath a great cough.

The Line of Selborne Goute
Scurvie
Dropsie.

[These notes are written by Goodyer on a fragment of a Petition of Glasier
and Good of earlier date.]

upon all or any ye said premisses, not
desistinge from youre Crueltyes and insolencyes, for ye entreatyes and
lamentacons as well of yo^r said poore subiects Richard Glasier and ye said
Elizabeth Good, as of strangers and lookers on, intreatinge but forbearance
untill harvest tyme was paste, for ye mowinge of ye grasse,—

[MS. f. 16.

*Note of Localities of northern plants, perhaps supplied by Shanne:
notes of books: method of taking Laudanum.*

Richard Shanne.

Pyrola groweth in shadowed woods in Craven, in a place called Craggie Close
in Lansdale.

Monophyllum groweth in Lancashire in Dingley Wood & in Harwood neare
to Blackburne. [Quotation from *Gerard emac.* p. 409.

Iatronicum medicamentorum Simplicium liber 2 continens eo quos ad probus,
ventrem, interanea que attinent. Otho Brunfelsii authore.

Viaticū novū per Casparū Wolphiū 1565.

Laudanū ꝑ̄s gr. 2 to be given w. a little conserve of Roses or any other
Conserve, and mare ale after it.

[MS. f. 15.

List of Chirurgical Works.

John Vigo wth certayne works published as by Tho: Gale.

Philippe Barone works of Chirurg.

Ambrosius Parrius

Chametans

Tagaultius

Franciscus Arcaeus

Jacobus Ruess de generacione hominis

Variorumque Scotus or Chirurgie magnie

Margarita Chyurgica by S: H: student in Physicke.

Schola Salerni.

Banisters Chirurgerie.

English Phlebotomy by Nicholas Guyer.

Pathway to Health.

Chimicall Phisicke and hermeticall Phisicke written by Josephus Luers-
itanus, translated by Tho: Tyme, minister.

The secrets of Alexis of Piemount.

The secrets of Albertus Magnus.

Bord his brevary of Health.

} Chirur:

I falls boke of Chirurgerie.

Antidotarium speciale a Jo: Jacobo Vueckero.

Bartholomewe Glanfeild translat per Stephen Batman, Chaplayne to the Erle of Oxford.

Cochinelo ground mingled with water gumme water.

Fernelius Judgement of Urines.

[MS. f. 131.

Weather record for the last part of May 1621 (?).

May

- 22 ♂ ♂⁷ ☽ thunder
 23 ♂ ♀ ☽ 9 a. winds & clouds.
 24 ♀ ♀ ☽ temprate
 25 ● ☽ full moon
 26 ☽ var cur.
 27 Rayne
 28 △ ♀ ☽ 10 a.
 29 △ ♀ ☽ 11 a.
 30 △ ☽ ☽ 11 a.
 31 ☽ in apog.

[Written on a paper on which Goodyer afterwards wrote his draft description of the Jerusalem Artichoke on 17 Oct. 1621.]

[MS. f. 116.

Miscellaneous Notes. Lists of Fruit Trees and Plants. Dates of Assizes.

Rec. 9 Feb. 1627 of Christopher Potecary of Stockton 5 myle from Venny Sutton Clothier.

- | | | |
|---------------------------------------|------------------------------|--|
| 1. Burgomaster peare till Easter | Southt Wed. | 5 Mar. Wimbor[ne] |
| 2. Kingekaterine 3 grafts till All S. | Dor. lun. | 10 Mar. Dorchester |
| 3. Painted peares | Somers. wed. | 12 Mar. Chard |
| 4. Norwich peare | Cornub. wed. | 19 Mar. Lanceston |
| | Exon. civit. lune | 24 Mar. Exon Gildhall |
| | Devon. Eadem die | apud Cash . . . |
| 1. Harvey aple | Wiltes martii j ^o | Apr. apud Sarum |
| 2. Quadlinge aple
bunch cherry | 18 Majj rem. | 4 ^h 5 ^s 4 ^d ob. |

[MS. f. 129.

- | | |
|--|---|
| lilli: convall: flo: rubro | Salvia versicolor |
| Aconitum luteum Ponticum | Yuca |
| Lanium Pannonicum | Paliurus? |
| Laburnum leaved trefoile | Pistachio |
| Colutea minor | Periclimum rectum |
| Anisum | Laurus Tinus |
| Digitalis minor lutea | Trachelium |
| Astragaloides | Rosmarinum |
| Balsamita flo. alb. | Nardus montana |
| Aconitum flo. delphinii | Gentianella verna |
| Geranium creticum variegatum | Ranunculus thalictri folio |
| Paralisis flo: viridi simplici | Polygonatum Virginianum |
| Sedum serratum | Ignotum ex Virginia |
| Cotiledon minor | Tanacetum striatum et hissupus |
| Gelsinum luteum | an Meum alterum |
| Fragaria fructu aculeato | Cicer arietinum flo. . . . et fructu albo |
| Geranium gratia dei batrachoides albo,
coerul. et variegat. | Cicutaria max. |
| Thlaspi vaccariae folio | Acanthus syl. |
| Jacea Baetica | Keyri max. |
| Thlaspi Mechlinense | Sorbus legitimus |
| " Creticum | Jacea lutea |
| Hissopus latifolia | Pimpinella fl. albo |
| | Cirsium mont. maius |

[MS. f. 129.

INDEXES

GOODYER'S PLANT LISTS

THE following Alphabetical List (Index I) includes plants with which Goodyer was acquainted, or which happened to interest him. The great majority were in cultivation in English gardens or were those whose native habitat he had discovered. That it was not exclusively a garden-list is indicated by the presence in it of 'Quercus marina' and of other seaweeds, but it may have been a catalogue of one of his Horti sicci. There are two copies of this list in Goodyer's handwriting (MS. 11, ff. 47-59). The fair copy seems to have been written out in the winter of 1617 and it remained in use for many years, for several additions dated 1618, 1625, 1628, 1634, &c., have been made later. A few of these added notes are printed below, in the form of foot-notes, but the greater number, being dated, have already been included among his descriptions of plants on page 109 *et seq.* The names are mostly those used by Gerard and Lobel, with a few taken from Dodoens, Clusius, Deleschamps, and Matthioli, a fact which points to the date of the compilation of the original list being anterior to the appearance of Bauhin's *Pinax*. Goodyer quotes references to pages of Lobel and Gerard (1597) throughout, but these, and his synonyms, have been omitted for the sake of economy of space.

We interpret the letters C., P., or F. in the margin as meaning that the plants so marked were growing in the gardens of Coys, Parkinson, or Franqueville respectively.

These two lists, written out as they were within a few months of one another and including additions up to the year 1634, are therefore of importance for tracing the spread of plants in English gardens. In the earlier list the number of plants marked P (Parkinson) is 239, in the later list 264, and the numbers of plants marked C (Coys) are 147 and 324 in the two lists respectively. Eight are marked as growing in Franqueville's garden.

The indexes do not include the names in Tradescant's lists, printed on pages 331-46, which should also be consulted.

INDEX I

PLANT-NAMES USED BY GOODYER.

Names in ordinary type are printed from Goodyer MS. 11, ff. 47-59. Names in *italics* occur in Goodyer's other MSS., and are probably of later date than the others.

C = grown in the garden of Coys.

P = " " Parkinson.

F = " " Franqueville.

C	Abies.	320	Allium silvestre tenuifolium.
	Abrotanum mas.		260, 316
C	" foemina.	318	Alnus.
	Absinthium marinum album,		" nigra.
	Seriphium vulgare.	281	Aloe.
P	Absinthium folio Lavendulae.		C Alopecurus gramen.
	" vulgare Ponticum.		Alsine corniculata.
CP	" inspidum.	319, 353	CP " repens maior baccifera.
	Acanthium album.		261, 318
C	Acanthus sylvestris aculeatus.		P " maior
		318	" minima
CP	" sativus, Branca ursina.		" hederacea } Morsus galli.
		306, 318	" hederula }
C	Acarna flo. rubro, like Carduus		" foliis trissaginis.
	mariae, but higher & lesser.	145	" myosotis.
	Acer maior. We call this a		" palustris.
	Sycamore tree.		" aquatica.
	<i>Acer minor.</i>	275, 283	" <i>josculis commixtibus.</i>
	<i>Acetosa maxima.</i>	194	" <i>serpillifolio.</i>
	<i>Achillea sideritis.</i>	155	Althea.
	Acinos.	120, 166, 280, 296	C Alyssum Dioscoridis.
P	Aconitum hyemale.		C Amara dulcis.
CP	" luteum Ponticum.	318	CP Amaranthus purpureus.
CP	" lycoctonum flore		CP Ammi vulgatius.
	Delphinii.	318	CP Amomum Plinii.
P	Acorum legittimum.	177	Amygdalus.
	Adiantum album et nigrum.	263	Anagallis.
	<i>Adonis.</i>	136, 319	CP " foemina coerulea.
C	Aegylops.		" lutea.
CP	Aethiopsis.	305	" aquatica.
F	Agnus castus.	326	" aquatica tercia.
	Agrimonia.		" aquatica tercia.
C	Agrioriganum.		" aquatica tercia.
	<i>Aizoon.</i>	152	FP Anagyris.
CP	Alaternus.	270, 317, 346, 355	P Anchusa lutea.
CP	Alcea peregrina.		C " supina lusitanica non
P	" vulgaris.		script.
	" vulgaris albo flore.	111	<i>Androsacc.</i>
	Alchimilla.	275, 306	Androsaeum; Climenon Italo-
	Alleluia offic. Opis alba.	275	rum.
C	" offic. Opis lutea.		P Anemone flo. pallido. I cannot
	Alliaria recentiorum.	111	sett downe the varieties I have
	Allium.		secne.
	" ursinum latifolium.	260, 275	Anemone Cardinals Hatt.

See also Index II and p. 334.

	Anemone pleno rubro.		Armeniaca malus.
	" coeruleo.		Armeria alba et rubra, & of
C	" nemorum albo pleno		severall colours on one stalk.
	flore. 318	C	Armeria varia. 346
	Anemone nemorum albo.		Artemisia. 284
	" nemorum purpurea.		Arum offic. 280
C	Anethum.		Arundo Vallatoria. 176, 280
	Angelica sativa.	P	Asarum.
	" silvestris. 275	CP	Asclepias flore albo.
	Anisum.	CP	" flore nigro.
	Anonis. 277, 287, 355		Ascyron. 275, 283-4
CP	Anthora.		Asparagus sativus.
	Anthyllis leguminosa Belgarum.		Asperula.
	" montana Deles. 1150.	CP	" flore coeruleo.
	Anonymos lini folio Clus. 324;		<i>Asphodelus</i> . 145, 319
	Leguminosa Belgarum Lo. O.		Asplenium. 263
	87. Ger. 1060. Gerard's figure		" silvestris Tragi.
	of Lagopus max. p. 1023 is	CP	Aster Italorum.
	likewise a trewe figure of this.	CP	" conizoides ex Virginia.
	117		120, 347
C	Anthyllis Valentina Lo. 421;	P	Astragalus. 16, 347
	Herniaria Boelii Clus. <i>Cur.</i>	CP	" marinus. 140
	<i>post.</i> p. 37. Ger. p. 452 hath	P	Astrantia nigra. 347
	a false figure. 318	CP	Atractylis flo. luteo.
	Antirrhinum album. 306		Atriplex marina repens. 285, 300
C	" minimum.		" sativa alba.
	" purpureum.		" sativa purpurea.
P	" minus.		" silvestris.
	115, 143, 275, 292	P	" olida.
C	" maius Baeticum		Avena vesca.
	flore rubro.		Avenaria ustilago.
	Aparine.	P	Auricula ursi flore rubro.
	Aphaca. 275		" flore purpureo.
	Apium hortense.		
	" crispum. 172	P	Baccharis Monspeliensium.
C	" Virginianum, Petrose-	C	Balsamina foem.
	linum.		Balsamita mas.
	Aquifolium sive Agrifolium.		" foem.
P	Aquilegia.	C	" foem. flo. albo. Age-
	I have seene of more colors		raturum.
	than I can nowe remember.		Barbarea. 320
	287, 347	C	" species. Beros ex
C	" multiplex.		Hispania. 317
C	Aracus maior Baeticus. 138		<i>Bardana minor</i> . 128
	" <i>minima</i> . 193		Batata Virginiana. 320, 356
	Arantia.		" Hispanorum at Mr. W ^m
P	Arbor vitae. 327		Yaldens 21 Aug. 1637.
C	" Judae. 317, 346		<i>Bellflower, Lesser</i> . 132
	<i>Arctium montanum</i> . 194, 280		Bellis maior.
C	Argemone. 155, 320		" maior silvestris.
	" capitulo longiore. 178		" hortensis multiplex flore
	Aria Theo[phrastii] Lo. o. 167. G.		rubro.
	1146. Why he calleth it a haw-		Bellis hortensis multiplex flore
	thorne I knowe not. Deles.		albo.
	202 hath a very trewe figure of		Bellis hortensis multiplex varie-
	it. We call it quick beame.		gatus.
	195	C	Bellis spinosa. 316, 318, 321
CP	Arisarum latifolium. 319		Berberis. Holythorne.
CP	Aristolochia clematitis. 306		Beta rubra.
P	" rotunda.		" alba.

See also Index II and p. 334.

- Beta marina. 281, 285
 " *Cretica*. 136
 Betonica.
 C " maior Danica. 319, 324
 CP " flo. albo. 319
 " aquatica. 175
 Betula. 275
 Bifolium.
 P Bistorta maior. Reddshancks
 or leggs.
 P Bistorta minor. 347
 CP Blattaria flore luteo. 112
 C " floe albo. 347
 " *purp.* 327
 C " maior lusitanica flo.
 odoratissimo.
 Blitum. 320
 C " seu Amaranthi species
 sem. nigro, like Amaranthus
 tricolor.
 C Blitum spinosum Creticum. 136
 Bonus Henricus.
 C Borago hortensis. 128
 C " hortensis flo. albo.
 " semper virens.
 CP Botrys. 260
 Brassica vulgaris sativa.
 " florida botrytis.
 " marina monospermos.
 " capitata alba. 285
 " capitata rubra.
 " prolifera. Colewort.
 Bromos sterilis.
 Bryonia alba. 262
 " nigra. 127, 262
Buglossa hispanica. 277
 Buglossum angustifolium.
 " echioides luteum.
 CP " scorpioides. 131, 318
 Bugula. 280
 Bulbocastanon.
 P Bupthalmum Dod. 327, 354
 C " primum Mathioli.
 Gerard hath the same fig.
 under the title of Buph-
 thalmum album p. 607.
 154, 318
 Bursa pastoris.
 Butomus. 177, 258
 Buxus. 347
 P Cachris verior. Gerard, p. 858,
 hath the figure of Seseli massi-
 liense Mathi. for it.
Calamagrostis. 172, 288
 CP Calamintha montana praestan-
 tior.
 Calamintha montana vulgaris.
 " aquatica Belgarum. 275
 P Calceolus Mariae. 347
 Calendula simpliciflora.
 " multiflora.
 " prolifera.
 C " silvestris mediofusca
 Poet.
 C Calendula silvestris medio lutea
 Boet.
 Caltha palustris. 347
 P " palustris multiplex.
 CP Campanula persici-folia. flo.
 albo et caeruleo. 347
 Campanula minor rotundifolia.
 Cannabis sylvestris spuria altera.
 " sylvestris tercia.
 " *spuria*. 194
 CP Capnos altera fabacea radice.
 263
 Capsicum.
 Caput gallinaceum Belgarum. 178
 Cardamine altera. Ladysmocks. 280
 " *impatiens*. 189
 P Carliaca.
 Carduus acaulis septentriona-
 lium. 145
 P Carduus Mariae lacteus. 145, 320
 C " " flo. nu-
 tante.
 CP Carduus globosus maior. 327
 CP " globosus minor.
 " stellatus.
 " *eriocephalus*, see Co-
 rona fratrum. 320
 C Carduus stellatus flo. luteo
 monspel.
 Carduus Benedictus.
 " spinosissimus.
 C " bulbosus monspel. 144, 319
 " viarum flo. purpureo }
 " viarum flo. albo }
 112, 145
 We have a kind here that
 smells like Musci.
 Carlina silvestris. 112
 C Caryophyllata maior rotundi-
 folia.
 P Caryophyllata vulgaris.
 Caryophyllus marinus.
 C " proliferus.
 C " repens.
 " multiplex. I have
 seen more varieties of them
 than I can set downe.

See also Index II ant p. 334.

- C Heliotropium Indicum vel Virginianum. Yow had lately planted it when I was at your howse 25 Martii 1617.
109, 166, 317
- CP Helleborastrum.
- P Helleborine. 275, 292
- P Helleborus albus. 337
- FP " albus flo. atrobente. 326
- CP Helleborus niger verus. 327, 347
- Helxine. 237
- C Hepatica trifolia coeruleo pleno flore. 319
- P Hepatica exalbido flore. Heptaphyllon.
- C " maius. 155, 319
Herba aquatica rubescens. 187
- CP Herba Doria. 119, 319
- CP " " altera serratofolio. 119
- P Herba Paris. 110, 275, 307
- P Herniaria. 327
- Hieratium alterum grandius. 284
" longius radicum.
" aphacoides. Succorie hawkweed. I have seen another Hieratium of 4 or 5 foot high having a round smooth stalk. 283
- C Hieracium falcatum maius. 149
- C " falcatum minus.
- C " medio nigrum flore majore. 149
- C Hieracium medio nigrum flore minore. 149
- Hieracium stellatum. 149
" *intybaceum.* 149
" *lanosum.* 150
- CP Hippoglossum bonifacia.
- P Hippolapathum rotundifolium.
" sativum latifolium.
- Hipposelinum. 347
- C " marittimum
Quicki. 318
Holosteum. 193, 280, 293
Honewort. 123
Hop. 114
Hordeum spurium.
" distichon. Lo. 29.
Ger. calls this Polystichum, p. 64, which is false or els his fig. is misplaced.
- C Hordeum nudum.
Hordei distichi ustilago.
Horminum sylvestre sive oculus Christi. 289
- P Horminum sylvestre Pannonicum. 156
- C Horminum flore coeruleo.
Hydrolapathum magnum.
- CP Hyosciamus albus.
" niger.
" luteus.
" albus lutescens medio purpureus.
- C Hyecoum Clusii. 129, 306, 309
- P Hypericum. 302
" Syriacum.
" exiguum Tragi.
" tomentosum.
- Hyssopus offic. flo. coeruleo. 115
- C " flo. albo. 319
- Jacea maior. 165
- C " palustris lusitanica non script. 164, 317
- C Jacea repens luteo flore. 317
- C " albo flore. 110, 317
- C " capitulis hirsutis. 164, 317
- P " marina. 317
- P " marina Baetica.
- C " Baetica aestiva.
Jacobaea aestiva. 284
" *Pannonica.* 194, 289
- Jasminum.
" Americanum.
- C Ignotum ex Virginea flo. Chrysanthemi parvo radice repenti, like to Lysimachia lutea.
- P Ilex glandifera.
Illecebra.
- P Imperatoria. 284
- Iris alba.
" palustris lutea. 177
" vulgaris.
- C Isatis silvestris vaccaria. 261
- Juncus laevis.
" aquaticus maximus.
- Juniperus. 123, 195
- Kali geniculatum. Salicornia.
" minus.
- CP Laburnum. 320
- Lactuca sativa.
P " agnina. 133, 284
P " agrestis. 159
" capitata.
C " Virginiana. 319
C " crispa.
" *sylvestris.* 111, 158
- Lagopus. 144
" flore ruberrimo. 284
- CP Lamium album.
" rubrum. 281
" luteum. 281, 317
- P " Pannonicum. G. 568,
a false figure. 156

See also Index II and p. 334.

C	Lamium Pannonicum. 2 Clusii.	C	Lupinus coeruleo maior.
	Lampsana. Hastie Roger, for cutts. 149	C	" ex coeruleo albo.
	Lapathum sanguineum.	C	" Indicus flo. coeruleo medio purpureo.
	" acutum.		Lupulus salictainus.
	" folio minus acuto.		" sylvestris.
	Lappa maior. Bardana maior.		Luteola. 134
	Lathyrus sylvestris floribus luteis.		Lychnis sylvestris albus multiplex.
	Lathyrus latiore folio 137		Lychnis sylvestris cauliculis striatis albo flore.
C	" palustris Lusit. 137		Lychnis sylvestris rubello flore. 195
C	" Creticus belli.	P	" Chalcedonica.
	" perenni radice.	C	" " fl. carneo.
C	" Orobeus Boeticus.	C	" " flo. albo.
C	" Dumetorum Baetic. 138	C	" " flo. mi-
	" aestivus Baeticus. 137		niato et carneo.
	" major ang. 195		Lychnis coronaria alba.
	Laureola.		" coronaria rubra.
FP	Laurocerasus. 326	C	" coronaria flo. suaveru-
	Laurus mas.		bente.
	" foem.	C	Lychnis marina Anglica. 282, 318
	Legumen pallidum. 139	P	Lysimachia lutea. 275
C	Lens minor.		" campestris.
	" palustris. 111		" galericulata.
	Lens minor. 111	P	" flore Delphinii.
	Lepidium Aeginetae.	C	" coerulea. 320
	Leucojum luteum duplici flore.		" purpurea spicata. 357
C	" luteum simplici flore.		" siliquosa.
	Wall-flower & Hartesease.	CP	" siliquosa ex Boria-
	Ione Grant: Turner calls it so.		libus Virginiae. 159
	Leucojum album.		Lysimachia forte. Wiltshere. 111
	" purpureum.		Majorana nostras. 120
P	" Patavinum marinum.		Malus.
	" melancholicum. 327		" sylvestris. Crabbe tree, wildinge tree.
P	Levisticum. Lo. 703. G. 855		Malva rosea fruticosa.
	hath ye fig. of Siler montanum for it; and hath put the figure of Levisticum mathi. p. 853 for Laserpitium.	C	" rosea multiplex
	Levisticum verum Gerardi. 319		{ albo } flo.
	Lichen. 230		{ rubro } flo.
	" marinus.		{ purpureo }
	Ligustrum.		Malva sylvestris repens pumila.
P	Lilium Convallium.		" sylvestris precocior vulgaris.
	" Convallium rubro flore.		" crispa.
	Limonium. 285		" vesicaria. Vide Alcea peregrina.
	Linaria. 118, 324		Malva aestiva flo. amplo Baet. 134
C	" aestiva minor. 143	CP	Mandragoras mas. 339
	Linum sativum.	P	" foem.
	Linum catharticum. 109, 112	P	Maracoc Virginiana. 165, 320
P	Lithospermum maius.		Marrubium album.
	" minus.		" aquaticum vulgi.
	Lolium album.		" nigrum. 287
F	Lotus arbor. 326, 339, 346		Marum, herbe Mastick.
CP	" siliquosus. Pisum quadratum. 139		Matricaria nostras.
	Lunaria minor. 110, 263, 264		
CP	Lupinus sativus flo. luteo.		
P	" " flo. coeruleo.		

See also Index II and p. 334.

- C *Matricaria grato odore.*
Medica spinosa. 142
 " *anglica.* 141
 " *pericarpio plano.* 143
 C " *minor elegans.*
 C " *marina spinosa.*
 C " *vulgaris.*
Melampyrum. 118
 C *Melilotus Italica.* 126, 317
 CP " *Germanica flo. albo.* 317
 " *coronata.*
 " *Indiæ.* 126
 C *Melissa Turcica.*
 Melo. *Muske Melon.* 165
 P *Mentha Cattaria.*
 C " *Cattaria media.*
 C " *Cattaria minima.*
 C " *Cattaria tuberosa.*
 C " *Romana offic.* 319
 " *sativa rubra.*
 " *Sisymbria aquatica.*
 F " *Danica.* 326
Menthastrum.
 C " *montanum.* 111, 318
 CP *Mercurialis mas.* 327
 CP " *foemina.*
 C " *species forte ex Virginia.*
Mespilus.
 CP *Meum.* 348
 P *Mezereon Germanorum.*
 CP *Millefolium odoratum sive nobile.* 319
 P *Millefolium luteum.*
 " *terrestre purpurcis floribus.*
Millefolium terrestre vulgare.
 " *flore & semine Ranunculi.*
Millegrana minima. 275
Mill mountaine. 109, 112
 CP *Mirabilia Peruviana.*
Moly serpentinum.
 " *Homericum.*
Morsus Diaboli.
 " *Ranae.*
Morus candida.
 " *celsa offic.*
Muscattella Cordi. 126, 291
 CP *Muscipula.* 348
Muscus terrestris vulgaris.
 " *ramosus.*
 " *ramosus floridus.*
 " *pyxidatus.* 230
 " *pellatus.* 230
 " *clavatus.*
 " *scoparius.*
 " *quercus.*
Muscus marinus sive corallina.
 C *Myagrum monospermon.*
 CP " *thlaspi facie.* 319
 CP *Myrrhis.* 290
Myrtus Brabantica.
Napellus verus coeruleus. 348
 C " *minor serotinus.*
Napus sive Bunias.
 " *silvestris.* 290
 P *Nasturtium Indicum.*
 C " *hortorum crispum.*
 " *hortense.*
 " *aquaticum.* The fig. is misplaced in Ger. p. 200. *Sion cratervae Erucae folium* L. o. 209.
Nepeta media. 162
 P *Nerium. Oleander.*
Nidus avis. 126, 195, 290
 C *Nigella Delphinium elatius var. color.*
 C *Nigella trigonum Lusitanicum.*
 CP " *Romana flo. albo.* 152, 348
 " *multiplex.* 152
 " *elegans.* 153
Nummularia. 261
Nux Juglans. 112, 174
Nymphaea alba.
 " *lutea.* 262, 281
Ocymoides semper virens, Clus. 99
Ocymum.
 C *Oenanthe cicute facie.* 319
 C " *filipendula aquatica.*
 " *angustifolia.* 115, 143, 195, 327
 C " *stellata.*
Olea silvestris.
Ophioglossum.
Ornithopodium perpusillum.
Orobanche. 126
 C *Orobis ex Venetiis.*
 P " *recept herbariorum.*
Osmunda. 275
 CP *Oxalis franca.* 319
 " *tenuifolia sinuata ververina.*
Oxis, vide Alleluia.
Oxyacantha. Hawthorne. 286, 292
 CP *Paeonia mas.* I heard of one with double flowers.
 CP *Paeonia foemina.*
 " *foemina multiplex rubro flore.*
Palma. 184

See also Index II and p. 334.

	Paludapium.		C	Phaseolus vulgaris albus.	
C	" selino Italomum.		"	" vulgaris niger.	
	Panax coloni. Lamium ut puto.		"	" vulgaris ruber.	
CP	" heracleum alterum recentiorum.			Phyllitis.	230
C	Panicum vulgare.		P	" multifidofolio.	185, 279
	" <i>sylvestre</i> .	120	P	Phyteuma monspel.	134, 327
P	Papaver spumen.		P	Pilosella maior.	
	" album sativum.		C	" maiore Syriaca.	317
C	" album multiflorum.		"	" repens.	150
	" corniculatum flore luteo.		"	" flo. coeruleo.	
	Papaver cornutum flo. rubro.		"	" flo. carneo.	
		283, 290	P	Pimpinella maior Fuchsii. L. 720.	
	" rhaeas.	155		Ger. fig. 887 agrees not with this herbe.	
CP	" spinosum, Clus.	320	P	Pimpinella saxifraga minor.	
	Parietaria.	118			282, 286
	Paronychia alsine folia.	290	P	" hortensis.	
	" rutaceo folio.		"	" hortensis maior.	
	Pastinaca domestica vulgi.		C	" agrimoniae folio.	318
	" sativa atro rubens.			Pinus.	
	" latifolia sylvestris.		CP	Pishimon prunus Virginiana.	320
	" sativa Diosc.			Pisum.	
	" silvestris tenuifolia.		C	" falcatum Boet. sine membrano.	
	" <i>aquatica minor</i> .	116	C	Pisum Indicum Gerardo. Vicia fructu albo.	139
	" " <i>latifolia</i> .	176		Pisum maculatum Boelii.	16, 141
	" " <i>maxima</i> .	176		" vulgatus maius.	
	Pedicularis.	263, 290		" minus.	
P	Pentaphyllon maius. Quinquifolium.		C	" silvestre Boeticum.	
P	Pentaphyllon minimum alpinum petrosum.			" umbellatum. G. 1045.	
	Pentaphyllon supinum Tormentillae facie.	275		French pease.	
	Pentaphyllon vulgare.			Plantago incana.	
	Pepo oblongus.		CP	" rosea spicata.	
	" rotundus.			" quinque nervia.	186, 301
CP	Perfoliatum vulgatus.	261, 281		" aquatica.	293, 353
CP	Periclymenum rectum.			" aquatica stellata.	110, 180
		281, 301, 320, 346		" marina.	
	" perfoliatum.		P	Plumbago Plinii. Dentaria Rondeletii.	
	" non perfoliatum.			<i>Pneumonanthe</i> .	170
P	Periploca repens folio angustiore.			Podagria germanica.	110
		346		Polemonium monspel.	
	" recta latifolia Virginiana.		P	Polium montanum.	281
	Persica malus.	131, 321		Polygala flore coeruleo.	
	" malacatone.			" flore purpureo.	
	Persicaria Hydropiper.			" flore albo.	
	" mitis maculosa.			" flore rubro.	
	Pes anserinus.			" obsoleto flore.	
	Petasites.	275	C	Polygonatum.	341
	Petroselinum, vide Apium.		C	" angustifolium.	
		318, 327	P	" Virginianum.	
	<i>Petum indicum</i> .	160		Polygonum Germanicum.	110
CP	Peucedanum.	348		" foemina.	
C	Phalaris bulbosa.	133, 317		" mas vulgare.	
	" pratensis.	282		" <i>marinum sp.</i>	148
C	" minor Boet. albo.	133		" <i>alterum</i> .	151
C	" minor nigro.	133		Polypodium.	184, 230
			C	Pomum amoris parvum.	

See also Index II and p. 334.

- C Pomum spinosum. 327
 C " spinosum flo. coeruleo.
 Populus alba.
 " nigra.
 " Lybica.
 Porrum vulgare. 260
 " tonsile. French leeks. 260
 Portulaca domestica.
 Potamogeton.
 Potentilla. Argentina. 262
 Primula hortensis maxima Anglica floribus plenis. 280
 Primula veris, flore gemino. Cowslips 2 in a hose.
 Primrose 2 in a hose. 109, 316
 Primula veris rubro flore. 283
 " pratensis. Cowslips.
 " pratensis inodora. Oxlips.
 Primula silvarum.
 " silvarum obscura virentibus.
 Prunella.
 " flore albo. 293
 Prunus sylvestris.
 " domestica.
 Pseudo-melanthium. 112
 C Psillium. 157, 348
 Ptarmica folio Taraconis.
 CP " Imperati. 119, 319
 Pulegium regium vulgatum. 281
 CP " cervinum.
 C Pulmonaria maculosa. G. 662.
 inc. flo. 10 Marcii. 320
 C Pulmonaria non maculosa. 320
 " officinar.
 " foliis Echii. 115, 195
 " gallica aurea. 180
 Punicum malum.
 Pyra.
 P Pyracantha. 346
 C Pyrethrum offic. 320
 Pyrus silvestris.
 Quadrifolium Phaeum fuscum hortorum.
 Quercus marina. Sea oare.
 " vulgaris. 173, 195, 283
 Quinquefolium. 170
 CP Radix cava flore albo. 348
 Ranunculus albus multiflorus.
 " aquaticus.
 " aquatilis flammeus angustifolius.
 Ranunculus aquatilis angustifolius serratus.
 P Ranunculus arvorum.
 " auricomus.
 Ranunculus bulbosus.
 P " Creticus flo. albo.
 P " gramineus.
 P " Illyricus minor.
 P " Illyricus maior.
 P " omnium maximus.
 " magnus Anglicus polyanthos flo. luteo. 284
 Ranunculus maximus multiplex. palustris rotundiore folio.
 Ranunculus pratensis.
 P " Pyrenaeus folio plantaginis.
 C Ranunculus globosus. 319
 " Apulei quibusdam Clus. p. 233.
 Raphanus silvestris aquaticus.
 P " rusticanus.
 P " niger.
 " sativus.
 Rapistrum arvorum.
 " flo. albo.
 Rapum maius orbiculatum.
 " silvestris non bulbosum. 191
 C Rapunculum silvestre Tragi. 111, 116
 " corniculatus. 185, 230
 P Rapuntium parvum. 320
 CP Rha capitatum folio Enulae. 318
 CP " Ponticum verum. 318
 P Rhamnus primus Clusii. 263, 275
 C " Clusii. 319
 " solutivas.
 P Rheseda. 134
 P Rhodia radix.
 P Rhus Plinii myrtifolia.
 P " coriaria Virginiana. Sumach. 321, 340
 Ribes Arabum.
 " nigra.
 C Ricinus.
 Ros solis.
 Rosa moscata simplici flore.
 " cinamomea flore pleno.
 Rose de Canelle.
 Rosa cinamomea simplici flore. 112
 C " sine spinis. 319
 " alba.
 " silvestris pomifera.
 " canina odorata.
 " canina inodora.
 " Hollandica. Province Rose.
 Rosa holoserica. 110
 " lutea.
 C " semper virens. 319

See also Index II and p. 334.

Rosa rubra.		C	Scorpioides siliqua crassa.	151
„ Damascena.			„ <i>multiflorus</i> .	151
Rosmarinum Coronarium.		CP	Scorzonera latifolia.	317
	168, 281	C	„ angustifolia.	317
Rubia maior tinctorum.	275	C	Scrophularia Lusitanica non	
„ silvestris.	191, 291		script.	317
C „ spicata Clusii.	132	P	Scrophularia lutea Bauhini.	
Rubus.	114		Math. p. 792. ? whether all are	
„ Idaeus.			not Lamium 2 exotic. Clusii.	
Ruscus.			Scrophularia maior.	156
C Ruta hortensis.	129		„ secunda Cretica.	
„ muraria.	263		Secale.	
			Secalina ustilago.	
C Sabina baccifera.			<i>Securidaca minor</i> .	130, 140
„ vulgator, sive sterilis.		C	Sedum maius Lusitanicum.	279
Sagittaria aquatica.	283, 291		„ minus offic. Prick-	
Salix aquatica. I have seene			madam.	284
8 kinds of osyars thus called			<i>Sedum Divi Vincentii</i> .	195
by the baskett-makers.	77		Sedum minus haematodes.	
Salix caprea latifolia.			Stoncroppe.	
„ humilis repens.			Sempervivum maius.	
C Salvia Indica.	317		Serpillum vulgare.	
„ maior.				143, 195, 281, 283
„ maior rubra.			Serratura.	287, 302
„ minor.		C	Sesamoides argentea.	317
„ agrestis.			„ salmanticum.	
Sambucus.		CP	Seseli.	346
„ aquatica		P	„ species a D. Franquville.	
„ rosea.	346			326
C Sana sancta Indorum: a trewe		C	„ species monspel.	319
Tabaco. Petum latifolium.		P	„ Aethiopicum.	346
P Sanicula.	275	CP	„ Aethiopicum Mathioli.	
CP „ guttata.		CP	„ Creticum.	
Saponaria.		C	Sideritis species.	230, 281
Satureia hortensis. Winter-			<i>Silibum</i> .	145
Savorie.			Sinapi sativum.	
Satureia hortensis aestiva.			„ alterum sativum. White	
Saxifraga alba.			pepper.	162
„ aurea lichenis facie.			Sisarum.	
	185	C	Siser erraticum Plinii. Droxford.	
„ Anglica facie Seseli				112
prat. monsp.			Sison Math. 552.	
Scabiosa maior vulgator.			Sium maius = Pastinaca aqua-	
„ maior vulgator flo.			tica.	116, 281
Scabiosa minor.	164		Sium Olusatri folio.	179, 292
„ minima hirsuta.	110		„ odoratum.	121
CP „ montana flo. luteo.			„ repens.	114
	285, 320		„ <i>sifoliis</i> .	121
„ flore rubro.	164		„ <i>pusillum</i> .	192
Scarlea.			„ <i>agrorum</i> .	327
Schoenoprasum. Cives.	260	C	Smilax coeruleus minor Boetic.	
Scolymus Theophrasti.	318		„ laevis.	
„ „ alterum.			„ laevis minor.	
Scordium.		C	Smyrnum Creticum.	318
C Scorpioides repens Bupleuri		P	Solanum heliacabum.	
folio.	151		„ hortense.	
C Scorpioides Mathioli. Telephi-		P	„ somniferum & lethale.	
um Scorpioides.	131, 305		Bella dona.	275
		CP	Solanum Virginianum.	321
		C	„ Boeticum flo. rubro.	

See also Index II and p. 334.

- P Soldanella marina. 185
 CP Solidago Sarracenicagermanica.
 Sonchus aspera vulgi.
 " foliis profundis laciniis
 sinuata hederacea. 284
 C Sonchus Africanus Boelio: like
 a Dandelion.
 Sophia Chirurgorum.
 " latifolia.
 Sorbus tormentalis Plinii.
 " sylvestr. Alpina.
 Sparganium.
 Spartum Anglicanum. 171, 282
 Spergula saginae. 275
 " marina.
 Spica lavendula flo. coeruleo. 157
 " hortensis minima.
 Spinacia.
 Spondilium.
 CP Stachys. 177
 C Staphilodendron Plinii.
 P Stoebe Salmantica. 154
 C " Salmantica altera. 317
 Stoechas.
 Superba maior vulgaris flo. Ar-
 meriae.
 Symphytum alum.
 P " tuberosum.
 Synanchica Deles. 113, 284
 P Syringa Italica alba. 346
 P " coerulea.

Tabaco. 161, 327
 Tamariscus. 305
 Tanacetum millefolii foliis. 154
 CP " non odorum.
 " speciosus cristatum
 Anglicum.
 Tapsus barbatus. 281
 Taxus. 123, 168, 195, 292
 C Telephium legitimum Imperati.
 318
 C Terrae glandes. Chamabalani.
 P Tetrabit. 194
 C Teucrium arborescens. 317
 P Thalictrum pratensis.
 P " tenuifolium.
 CP " Hispanicum aquilinae folio. 348
 P Thapsia.
 CP Thlaspi Diosc.
 P " vulgatissimum: mytri-
 date mustard.
 C Thlaspi candice flo. albo.
 " umbellosum. 130
 C Thymbra Boelii. 317
 Thymum durius. 179
 " latifolium.
 Tilia foemina. 346

 Tithymalus Cupressinus sive
 humipinus. Welcome to our
 howse, or Cypres Spurge.
 C Tithymalus montanus Boet.
 " helioscopius.
 " paralius.
 C " serratus monspel. 317
 Trachelium maius Urticae foliis.
 292
 P " flo. albo.
 CP " gyganteum. 319
 " minus. 302
 P Tragacantha.
 Tragon Mathioli.
 Tragopogon luteum.
 CP " purpureum.
 C " flo. cinereo.
Tribulus. 123-6
 Trichomanes Diosc. 179, 275
 CP Trifolium burgundiarum.
 " pratense flo. albo. 144
 " pratense flo. purpureo.
 C " odoratum. 306
 " corniculatum.
 " cordatum.
 " luteum.
 " vesicaria.
 " paludosum.
 C " falcatum Clusii.
 " *fruticans.* 326-7
 Tripolium vulgare.
 Triticum spica multiplici.
 Tubera terrae. 281
 Turritis. 292
 Tussilago. 262
 Typha. 281, 292, 296

 Ulmaria.
 Ulmus latifolia. Wich hasell. 184
 " Wich elme in Essex. 43
 Umbilicus Veneris. 292
 Urtica minor acrior.
 " Romana.
 " silvestris asperior.
 Uva crispa.
 " " rubro.
 " " coeruleo.

 Vaccinia nigra.
 P " rubra.
 Valeriana Graeca.
 C " Graeca, flore albo. 317
 " maior. Longe sore
 leafe for cuts.
 CP Valeriana rubra.
 " silvestris.
 " silvestris minor.

CP	Valeriana petraea.	318	C	Viola Mariana flo. albo.	
C	„ Mexicana.	133		„ aquatilis.	
	Veneris pecten.			„ lunaris G. 377. Bolbonac.	
C	Verbascum 4 Mathioli.	280		„ canina coerulea inodora.	
C	„ blattariae folio.			Viorna.	
	Verbena communis.		P	Virga aurea serrato folio.	
	Veronica vera et major.				283, 320
	„ minor serpilli folia.			Virga pastoris.	
	„ <i>mas recta</i> .	196		Viscum quercum.	
	Viburnum.	275, 346	C	Visnaga.	
	Vicia silvestris.			Vitis uvifera.	
	„ <i>maxima sylvatica</i> .	190, 292, 308	CP	„ Virginiana.	321, 346
	„ <i>segetum</i> .	193			
P	Viola Matronalis.			<i>Wheat</i> .	180
	„ tricolor.		P	Wosaran Virginianum.	
	„ tricolor silvestris.	194			
	„ Martia precox purpurea et alba.			Xanthium. Bardana minor.	128
	Viola Martia purpurea multiplex.	109		<i>Xiphium</i> .	177
	„ Martia alba multiplex.	318		Xyris.	275
CP	„ Mariana.			[Zea] Erased.	

See also Index II and p. 334.

INDEX II

PLANT-NAMES USED BY OTHER BOTANISTS, SUPPLEMENTARY TO THOSE LISTED IN INDEX I.

- Abrotanum 355.
 Absinthium 353-5.
 Abutilon 358.
 Acacalis 358.
 Acacia 365.
 Acanthus 318.
 Acer 346, 356.
 Achillea 155, 319.
 Acinos 355.
 Aconitum 318.
 Acorus 177, 234, 258, 287, 326, 365.
 Actaea 329.
 Adders tongue 311.
 Adonis 319, 358.
 Adoxa 69, 126.
 Aegopodium 110.
 Aesculus 346.
 Agnus castus 240.
 Agrifolium 281.
 Agrimony 311.
 Agropyrum 257.
 Ajuga 244, 255, 264.
 Alchemilla 155, 319-20.
 Alexanders 312.
 Alisma 242, 358.
 Allium 259.
 Alnus 175, 358.
 Aloe 261, 360.
 Alopecurus 237.
 Alsine 287, 292.
 Althaea 242.
 Amaranthus 320.
 Ameos 365.
 Amivil 270.
 Ammi 319-20, 365.
 Amomum 365.
 Amygdalus 358.
 Anacyclus 320.
 Anagallis 252.
 Anagyris 326.
 Anchusa 131, 283, 336.
 Andropogon 365.
 Androsace 150.
 Andryale 320.
 Anemone 311, 319-20, 328, 331.
 Angelica 240, 312.
 Anonis 355-6.
 Anotto 369.
 Anthemis 135, 154, 242, 301, 318-19.
 Anthericum 326.
 Anthriscus 253.
 Anthyllis 283.
 Antirrhinum 359.
 Apium 114, 172, 192, 195, 302, 356.
 Apocynum 329, 356, 359.
 Apples 342, 386.
 ,, of Love 345.
 Apricots 58, 241, 328, 344.
 Aquilaria 366.
 Arach 31.
 Arbor tinctoria 356.
 ,, vitae 328.
 Arbutus 287, 331, 346.
 Arctium 194.
 Argemone 320.
 Arisarum 319.
 Aristolochia 335, 359.
 Arkangel 240.
 Artemisia 318-19, 356.
 Artichoke 23-5, 241, 312.
 Asarina 287.
 Asclepias 131, 312.
 Ash 5.
 Asparagus 312.
 Aspen 264.
 Asperula 33, 113.
 Asphodelus 359.
 Aspidium 181-3.
 Asplenium 108, 242, 263.
 Assarabacca 312.
 Aster 50, 320, 331, 356.
 Astragalus 130, 140.
 Astrocarpus 317.
 Athamanta 318.
 Atriplex 236, 282, 320, 356.
 Aucklandia 364.
 Ballote 280, 287.
 Balme 311.
 Balsamodendron 365.
 Banana 250.
 Barba Jovis 331, 357.
 Bardana 356.
 Bartsia 117, 150.
 Basil 240.
 Battato 238.
 Bead tree 352.
 Beans, Kidney 312.
 Bears Ears 311.
 Becabunga 293, 300.

See also Index I and p. 334.

- Bellflower 311.
 Bellis 281, 288, 352.
 Berberis 365.
 Beta 355-6.
 Betonica 288.
 Betony 311.
 Betula 175.
 Bindweed 325.
 Bistorta 311, 356.
 Bixa 369.
 Blitum 356.
 Bluebottle 311.
 Borage 311.
 Botrychium 110, 245, 263.
 Brambles 34.
 Brassica 101, 319.
 Briza 158.
 Brookweed 29.
 Bryonia 82, 202.
 Bugle 264.
 Bugula 356.
 Bunias 191.
 Bupleurum 261, 346.
 Burri grass 352.
 Butomus 177.

 Cabbage tree 269.
 Cachrys 355.
 Calamintha 311, 355.
 Calceolus 356.
 Camomile 310.
 Campanula 132, 195, 285, 288, 302, 318.
 Champion 311.
 Camphorata 359.
 Candy Mustard 324.
 Cannabis 281, 288.
 Capillus veneris 284.
 Capparis 357.
 Capsicum 357, 368.
 Cardamine 284, 288, 356.
 Cardamons 365.
 Cardinal 331.
 Carduncellus 317.
 Carduus 21, 112, 118, 145, 154, 281,
 292, 302, 352, 359.
 Carex 194-6, 257.
 Carica 369.
 Carlina 112, 145.
 Carum 53, 121, 365.
 Cassado 368.
 Castanea 174.
 Cattaria 317, 318.
 Caulalis 53, 114, 128, 195, 282, 356.
 Cedar 369.
 Celtis 201, 326, 339, 346, 354.
 Centaurea 110, 154, 164, 317.
 Centranthus 317.
 Cerasus 346, 356.
 Cercis 317, 346, 370.
 Cerinthe 128, 180, 324.

 Ceterach 230, 263.
 Chamaebuxus 292.
 Chamaelaea 359.
 Chamaerops 369.
 Chara 55.
 Chelidonium 355.
 Chenopodium 260.
 Cherry 55, 59, 328, 344.
 Chervil 312.
 Chestnut 34, 312.
 Chrysanthemum 134, 155, 318, 321.
 Chrysosplenium 185, 302.
 Cichorium 150, 260.
 Cicuta 179.
 Cinara 352.
 Cinnamomum 269, 364.
 Cistus 317, 336.
 Citrullus 165.
 Clary 244, 311.
 Clathrus 196.
 Clematis 236, 319, 326, 354, 359.
 Clinopodium 359.
 Cloudberry 222.
 Cloves 269.
 Clown's Allheal 312.
 Cneorum 317.
 Cnicus 21, 49, 109, 144, 146, 154, 234,
 317-19.
 Cochlearia 317, 355.
 Coco 269.
 Cogcigra 331.
 Cola 269.
 Colchicum 110, 249, 258, 300, 316, 326,
 359.
 Columbine 25, 312.
 Comfrey 311.
 Conioselinum 318.
 Convallaria 326.
 Convolvulus 129, 153, 185.
 Coral root 26, 74.
 Corallina 284.
 Corcorus 359.
 Coriander 312.
 Corn parsley 53.
 Cornus 234, 346.
 Corona 326.
 Coronilla 305.
 Corydalis 359.
 Costmary 311.
 Costus 364.
 Cotinus 359.
 Cotton 368.
 Cotula 292.
 Cowslip 27, 311.
 „ of Jerusalem 311.
 Cracca 284.
 Crambe 242.
 Cranberry 330.
 Cranes bill 312.
 Crithmum 193.

See also Index 1 and p. 334.

- Crocus 249, 258, 316, 329, 326.
 Crucianella 132.
 Cucubalus 318.
 Cucurbita 165, 321.
 Currant 328.
 Cuscuta 112.
 Cyclamen 319, 331.
 Cynodon 257.
 Cynoglossum 280, 302, 317.
 Cynosurus 157.
 Cyperus 296, 359.
 Cytisus 317, 331, 359.

 Daffodil 27, 240, 311, 326.
 Dames Violets 311.
 Damasonium 110, 180.
 Datura 307, 320, 329.
 Daucus 112, 243.
 Delphinium 318.
 Dentaria 186.
 Dewberry 33.
 Digitalis 186, 244, 262, 356.
 Digitaria 196, 257.
 Dill 312.
 Diospyros 201, 320, 346.
 Diotis 148.
 Dittander 311.
 Dodder 311.
 Doronicum 237, 351.
 Draba 69.
 Dracunculus 337.
 Dragon 311.
 Dryopteris 183.
 Durien 367.

 Echatega 328.
 Echinochloa 120, 195.
 Echium 302, 317.
 Elaphoboscus 359.
 Eleocharis 257.
 Elm 17, 30-1, 37-43, 52, 265, 308, 314.
 Emblica 366.
 Emex 107.
 Endive 311.
 Enula 311.
 Epilobium 111.
 Epimedium 329.
 Epipactis 47, 107, 126.
 Equisetum 55.
 Eragrostis 257.
 Erigeron 162, 243.
 Erodium 146.
 Eryngium 110, 317.
 Erynus 286.
 Erysimum 319.
 Erythraea 244.
 Eupatorium 236.
 Euphorbia 236, 244, 306, 317, 353.
 Everlasting 311.
 Exotic fruit names 366.

 Fagus 188.
 Feather grass, 325.
 Fennel 274, 312.
 Ferns 67, 90, 181.
 Ferula 366.
 Ferulago 359.
 Feverfew 311.
 Ficus 242, 328.
 Figwort 17.
 Filago 280.
 Filipendula 312.
 Fir 369.
 Flax 156.
 Flower Constantinople 311.
 ,, de luce 310.
 ,, gentle, 325.
 Foeniculum 281, 318.
 Foenum graecum 356.
 Frangula 287.
 Frankenia 148, 151, 318.
 Fritillaria 249, 316, 321, 328.
 Fritillary 311.
 Fruits 267, 312, 329, 342-5.
 ,, exotic 365-71.
 Fucus 287.
 Fumaria 263.
 Fungus 245, 353.
 Fustick 369.

 Galanga 367.
 Galeopsis 194.
 Gallium 113.
 Genista 328.
 Gentiana 20, 170, 292.
 Gentianella 283, 286, 288.
 Geranium 185, 191, 234, 284, 286, 301,
 319, 329, 331.
 Germander 311.
 Geum 234.
 Gilliflower 61, 240, 243, 308-11, 328,
 330.
 Gladiolus 258, 310, 316, 321, 326, 338,
 354.
 Gladon 310.
 Globe flower 312.
 Gnaphalium 148, 243.
 Goldenrod 311.
 Gonolobus 321.
 Goodyera 99, 185.
 Gooseberry 312.
 Gratiola 317.
 Gramen 352.
 Grasses 71.
 Grass of Parnassus 54.
 Gromel 311.
 Guaicum 346, 371.

 Hacchic 365.
 Hazel 345.
 Heartscase 240.

See also Index I and p. 334.

- Hederula 289.
 Hedysarum 147, 318.
 Helianthemum 50.
 Helianthus 23, 109, 166, 307, 317.
 Heliotropium 24.
 Hellebore 312.
 Helleborus 285, 302, 327.
 Henbane 311.
 Hepatica 319.
 Herb Gerard 312.
 Hibiscus 306, 346.
 Hieracium 149, 180, 281, 283, 318.
 Hieracium 289, 317.
 Higtaper 295.
 Hippocrepis 77, 187.
 Hippomane 269.
 Hippophae 255, 263.
 Hissopoides 359.
 Hollias, 331.
 Hollyhock 312.
 Honewort 53, 67, 121.
 Honeysuckle 240.
 Honeywort 180.
 Hops 314.
 Hordeum 257, 282.
 Humulus 107.
 Hyacinths 311.
 Hyacinthus 240, 249, 311, 316, 321,
 325-6.
 Hymenaea 368.
 Hyoscyamus 122, 160, 320.
 Hyoseris 289.
 Hypecoum 129, 319.
 Hypericum 77, 234.
 Hyssopus 311, 319.

 Iberis 130, 324.
 Ilex 352.
 Imgara 366.
 Inula 320.
 Ipomaea 262, 320.
 Iris 177, 258, 326, 328, 331, 352.
 Isnardia, *see* Ludwigia.

 Jaca 367.
 Jacea 359.
 Jacobaea 357.
 Jamboli 367.
 Jasione 110, 164.
 Jasminum 326, 352.
 Jasonia 120.
 Job's tears, 312.
 Joseph and Mary 35.
 Juglans 112, 174.
 Juncaria 289.
 Juncellus 282.
 Juncus 257, 282, 284, 286, 296.
 Juniperus 123, 195, 355, 359.

 Kings spear 310.
 Knot grass 241.

 Lactuca 111, 158, 234, 319, 326.
 Ladies bower 312.
 „ girde 310.
 „ lacies 240.
 „ mantle 311.
 „ navel 240.
 „ seal 240.
 „ smock 240.
 „ thistle 312.
 Lamium 156, 317, 353.
 Lapsana 151.
 Larks peel 308, 312.
 Laserpitium 319.
 Lathraea 123, 237.
 Lathyrus 136, 195.
 Laurus 346.
 Lavandula 106.
 Lavatera 353.
 Lavender 311.
 „ cotton 312.
 Leneholans 312.
 Lens 137.
 Leontodon 319.
 Lepidium 329.
 Leucoium 326, 352.
 Libanotis 359.
 Lignum aloes 366.
 Lilium 326, 359.
 Lily 311.
 Lily of Valley 304.
 Lime 17.
 Linaria 17, 115, 143, 163.
 Linariola 289.
 Linum 109, 112.
 Lithospermum 242.
 Littorella 106.
 Locust tree 346, 368, 370.
 London tuft 311.
 Lonicera 301, 320, 326, 346.
 Lovage 312.
 Ludwigia 90, 187, 193, 195.
 Lupinus 312, 356.
 Lunaria 311.
 Luzula 357.
 Lychnis 283, 285.
 Lycium 365.
 Lycopodium 108.
 Lysimachia 187, 243, 261, 311, 355.

 Madder 312.
 Maize 354.
 Makenbuy 355.
 Malacaton 58, 329.
 Mallow 312.
 Malva 111, 134, 352-3.

See also Index I and p. 334.

- Mancinel 369.
 Mandragora 306.
 Mango 367.
 Manihot 368.
 Maracoc 370.
 Marigold 311.
 Marigold, French 25, 311.
 Marjoram 36, 62, 166, 240, 311.
 Marsdenia 329.
 Martagon 326, 328, 343.
 Marum 356.
 Marvel of Peru 25.
 Matricaria 234, 355.
 Matthiola 242, 330, 339.
 Maudlain 311.
 May apple 371.
 May Lily 326.
 Mechoane 262.
 Medicago 141.
 Medlar 328, 352.
 Melampyrum 118.
 Melilotus 126, 306, 317.
 Melon 274, 369, 371.
 Mentha 286, 326, 355-6.
 Mercurialis 236.
 Militaris 290.
 Mint 311, 326.
 Mollugo 284.
 Moly 325-6.
 Monophyllon 264.
 Monotropa 122.
 Montia 195.
 Moringa 367.
 Motherwort 311.
 Mucca nut 369.
 Mulberry 312, 329.
 Mullen 295.
 Munk's Rhubarb 311.
 Muscus 282-4, 290, 352.
 Muttelina 359.
 Myosotis 302.
 Myrica 255, 305.
 Myrobalan 366.
 Myrrhis 234, 312.
 Myrtil 329.
 Narcissus 258, 311, 316, 321, 326,
 331-3.
 Nardus 171.
 Nasturtium 236.
 Nectarine 345.
 Negundo 367.
 Neottia 107.
 Nepeta 111, 161, 317-18.
 Nettle tree 326.
 Nicotiana 160.
 Nigella 25, 152.
 Notobasis 145.
 Nux vesicaria 371.
 Nymphaea 293.
 Oak 5, 34, 369.
 Ocimum 166.
 Oculus Christi 311.
 Oenanthe 35, 115, 145, 319
 Oenothera 159.
 Olea 304.
 Oleander 240, 329.
 Olive 304.
 Olyas 331.
 Onobrychis 178.
 Ononis 242.
 Ophioglossum 245.
 Ophrys 244.
 Opuntium 292.
 Orach 311.
 Orange 329.
 Orchis 286, 290, 301-2.
 Organy 240.
 Orice 240.
 Origanum 166, 286.
 Ornithogalum 305, 319.
 Ornithopus 131, 305.
 Orobanche 47, 122.
 Osiers 77.
 Oxalis 290.
 Oxys 355.
 Paeonia 359.
 Paeony 312.
 Panax 359.
 Pansy 325.
 Papaver 155, 242, 366.
 Papyrifera 237.
 Paralysis 283.
 Paris 110, 244, 307.
 Parnassia 180, 243.
 Passiflora 320.
 Peach 329, 345.
 Pear 59, 343, 386.
 Pellitory, bastard 312.
 Pedicularis 106, 262.
 Penny royal 311.
 Pepper 236, 368.
 Perfoliata 359.
 Periclymenum 356.
 Periploca 346, 371.
 Periwinkle 312.
 Persicaria 292, 356, 371.
 Pescod tree 312.
 Petasites 234.
 Phalangium 326.
 Phalaris 133, 317.
 Phaseolus 237.
 Phleum 257.
 Phlomis 352-3.
 Phragmites 108.
 Phyteuma 111, 116, 185, 196.
 Picea 320.
 Pilosella 282.
 Pilularia 196.

See also Index I and p. 334.

- Pinus 346.
 Piper 237.
 Pistacia 201, 340.
 Pisum 139, 141.
 Planta 284.
 Plantago 106, 244, 284, 286, 301-2,
 356.
 Plantain, Rose 311.
 Platycerium 249.
 Plum 343.
 Poa 256.
 Poison weed 368.
 Polyacantha 360.
 Polyanthus 281, 352.
 Polygonatum 237.
 Polygonum 290.
 Polypodium 189.
 Polypogon 237, 321.
 Pompion 27.
 Pomegranate 329.
 Poppy 311.
 Potamogeton 123.
 Potato 75, 216.
 „ sweet 28.
 Potentilla 170, 262.
 Poterium 318.
 Primrose 27, 311.
 Primula 109, 281, 316, 318, 352, 353.
 Privet 355.
 Prunella 244.
 Prunus 321, 326, 346.
 Pseudonarcissus 283, 326.
 Pseudodictamnus 327.
 Pulegium 285, 360.
 Pulicaria 236.
 Pulmonaria 115, 190, 320.
 Pulsatilla 290.
 Pyracantha 346.
 Pyrethrum 318.
 Pyrola 264.
 Pyrus 195.

 Quamoclet 360.
 Quercus 172, 195.
 Quince 329, 343.

 Ranunculus 291, 301, 326.
 Raphanus 242, 311.
 Rapistrum 284.
 Raspberry 310.
 Reseda 134.
 Rhagadiolus 148.
 Rhamnus 255, 263, 317, 319, 341, 346.
 Rheum 318.
 Rhinanthus 261.
 Rhubarb 49, 260, 367.
 Rhus 321, 341, 346, 368.
 Ribes 326.
 Ribesium 284.
 Ribwort 241.

 Ricinus 371.
 Robinia 339, 346, 370.
 Rose 110, 112, 243, 291, 301, 312, 319,
 329-31, 341, 360.
 Rosemary 8, 240, 312.
 Rubia 191.
 Rubus 114, 195, 243, 282.
 Rue 312.
 Rumex 194, 319.

 Saccharum 258.
 Saffron 258, 311, 349.
 Sagapeni 360.
 Sage 311.
 Sagina 77, 179, 186.
 Salicornia 152.
 Salix 77.
 Salvia 156, 244, 305, 317, 356.
 Sambucus 87, 111, 286, 291.
 Samolus 111.
 Sanicle 241, 311.
 Santalum 366.
 Saponaria 261.
 Savey tree 312.
 Savory 311.
 Saxifraga 281, 301, 356, 360.
 Sawwort 311.
 Scabiosa 164, 243, 281, 320, 352.
 Scabious 25, 311.
 Scammonia 360.
 Scandix 306.
 Scilla 259, 321.
 Scirpus 175, 257.
 Scleranthus 110.
 Scolopendrium 108.
 Scolymus 318.
 Scordium 360.
 Scorodoprassum 311.
 Scorpioides 324.
 Scorpiurus 132, 151.
 Scurvy grass 311.
 Scutellaria 301.
 Sedum 77, 152, 195, 234, 243, 319-20,
 324.
 Selfheal 244.
 Selicio 319.
 Sena 360.
 Senecio 119, 193-4, 281, 319.
 Serpents Moly 311.
 Scrpillum 281.
 Serratula 302.
 Sesamum 360.
 Seseli 292, 319, 360.
 Sherardia 148.
 Sideritis 317.
 Silaus 243.
 Siler 360.
 Silybum 320.
 Sinapis 111.
 Sison 121, 360.

See also Index I and p. 334.

Sium 116, 176, 286, 302.
 Smallage 312.
 Smilax 258.
 Smyrniium 318.
 Snakeroot 371.
 Sneezewort 311.
 Snowdrop 311.
 Soapwort Gentian 311.
 Solanum 302, 320-1, 355, 360, 371.
 Solidago 319-20.
 Solomon's Seal 312.
 Sonchus 69.
 Sowthistle 326.
 Spartum 292.
 Speculum veneris 280.
 Spignel 312.
 Spiraea 77.
 Spongilla 290.
 Squinant 365.
 Stachys 59, 177, 319.
 Staphylea 320, 340, 346, 371.
 Star of Bethlehem 311.
 „ Jerusalem 311.
 Stellaria 261.
 Stickadove 240.
 Stipa 318.
 Storksbill 319, 324.
 Stramonium 320, 371.
 Strychnos 352.
 Suaeda 179.
 Succory 311.
 Sumach 321.
 Sugar cane 258.
 Sweet William 308, 311.
 Swertia 234.
 Sycamore 312, 357.
 Symphitum 360.
 Syringa 339, 346.
 Tamalapatra 368.
 Tamarix 305, 312, 346.
 Tamus 127, 153, 262.
 Tansy 311.
 Taraxacum 115.
 Tarragon 311.
 Taxodium 337.
 Taxus 168, 196.
 Telephium 318, 360.
 Terminalia 366.
 Thalictrum 360.
 Thesium 117.
 Thlaspi 236, 284.
 Thora 360.
 Thorowleaf 311.
 Thrift 311.
 Thymus 195, 317.
 Tilia 346.
 Tithymalus 306.
 Toadflax 17.
 Tobacco 32, 34, 160, 238, 327, 383.

Tolpis 149.
 Tomatoes 50.
 Touchmenot 240.
 Tragacanth 331.
 Trewlove 311.
 Trefoyle 25, 326.
 Trifolium 292, 342, 356, 360.
 Trinciatella 360.
 Trinia 77.
 Triticum 257.
 Trollius 319.
 Tropaeolum 329.
 Tulips 311, 329, 331.
 Turmerick 367.
 Tussilago 262.
 Tutsan 311.
 Twyblade 244.

Ulex 189.
 Ulmus 38-43, 283.
 Urtica 156, 280.
 Vaccinium 243, 292.
 Valentina 357.
 Valeriana 133, 312, 318, 360.
 Veratrum 326, 337.
 Verbascum 106, 284, 286, 360.
 Venus Looking glass 240.
 Veronica 76, 244, 300, 317, 320.
 Viburnum 346.
 Vicia 77, 139, 190, 193, 196, 281, 321.
 Vicia 296.
 Vina 356.
 Vine 329, 331, 369.
 Viola 109, 194, 360.
 Violet 312.
 Virgin's bower 240.
 Viscum 242.
 Vitex 326.
 Vitis 321, 346.
 Vulnearia 308.

Wheat 63.
 Winter cherry 311.
 Wolfsbane 312.

Xanthium 128.
 Xeranthemum 119, 319.

Yeast 314.
 Yew 5, 48.
 Yooke Elme 240.
 Youthwort 240.
 Yucca 313, 321.

Zea 258.
 Zephyranthus 320.
 Zizipha 352.

INDEX III

PERSONS, PLACES, AND THINGS.

- Abbot 274.
 Abell, H. 378.
 Abergavenny, Lord, garden 261.
 Alberts, H. 200.
 Ale 316.
 Alexandria 364.
 Aleyne 312.
 Allestre 202.
 Aloes 261.
 Alresford 5, 12, 373, 383.
 Alton I, 81.
 America 369.
 Amherst, Hon. A. 328.
 Anagrams 273.
 Anderson 279.
 Andrewes, Bishop 59.
 Annis seed 382.
 Ansell 67.
 Apothecaries 20, 21, 55, 57, 59, 76, 201,
 270, 275, 304, 353, 384.
 Apuleius 70.
 Arabia 364-8.
 Argent 20, 250.
 Armada 12.
 Ashmole 82, 330.
 Assizes 73, 386.
 Astrology 82, 309.
 Athos Mt. 364.
 Atkins, Dr. H. 248.
 Austen 83, 184.
 ,, Jane 4.

 Backhouse 82.
 Bacon, F. 62, 267.
 Baetica 157.
 Balden 47, 127.
 Barber surgeon 274.
 Bargett 224.
 Barlous 200.
 Barlowe, T. 333.
 Baronsdale, Dr. 249.
 Barrowe 378.
 Bartar, T. 194.
 Basing House 81, 273.
 Basingstoke 211.
 Baskett 91.
 Bath 91.
 Bauhin 46, 200.
 Bayley, W. 235, 304, 314.
 Beacon-fire 5, 378.
 Beale, J. 83.
 Bedelcombe, R. 378.

 Bed-fleas 31.
 Beeches Place I, 29.
 Beer 314.
 Bell 374.
 Belon 358.
 Benwyn 374, 382.
 Berain 239.
 Bere Forest 50.
 Bermuda 368.
 Beryton, *see* Buriton.
 Betisworth, Edith 8.
 Bettesworth, P. 96.
 Billson, C. 96.
 Bilson fam. 27, 95.
 ,, Bishop T. 7, 376.
 ,, L. 7, 95.
 ,, Mrs. 9.
 ,, O. 96, 200, 297.
 ,, Sir T. 6, 9, 46, 374-6, 379.
 Binding, cost of 213.
 Bishops of Winchester 7.
 Andrewes 59.
 Bilson q. v.
 Cooper 10.
 Curle 56.
 Neile 56.
 Bishops Waltham 12.
 Blackman 379.
 Blois 279.
 Blount 200.
 Bobart 81, 276, 278, 293, 295.
 Bodleian Library 271.
 Boel 16, 313, 361.
 Bold 231, 383.
 ,, A. 96.
 ,, J. 97.
 Bonham-Carter 6.
 Books 13, 197-232, 380, 385.
 ,, Italian 202.
 Boorde 314.
 Botanical excursions 75.
 Boulger 295, 330, 346.
 Bowden 200.
 Bower 208.
 Bowles 274, 276-7.
 Bowyer 67.
 Boyle 304.
 Bradley 65.
 Bramshott 5, 377.
 Brazil 237, 268.
 Brewing 16, 315.
 Brewning 11.

- British Museum 210.
 Britten, J. 122, 295.
 Broadgates Hall 225.
 Browne, Lancelot 200, 238.
 " Sir R. 270, 355.
 " T. 287.
 " W. 80, 278, 280, 292, 293,
 298-302.
 Browne of Hoe 35, 115, 297, 378.
 Buckett 59.
 Buckley, R. 272.
 Buckner 55, 59, 276-8.
 Bull 382.
 Bullein 216.
 Burghley, Lord 254.
 Buriton 5, 8, 11, 94, 375.
 " armour bearers 380.
 " pioneers 381.
 Burley 97.
 Burton, A. 264.
 " G. 194.
 " R. 264.
 Bush 247.
 Busher 237.
 Bust 200.
 Butser Hill 5.
 Buttersworth Hill 74, 187.

 Cachrys 172, 363.
 Cadel 279.
 Cairo 364.
 Caltherste 31.
 Cambrai 325.
 Camerarius 234, 358.
 Canary Wine 9.
 Canterbury 283.
 Capel 30.
 Carriers 202.
 Casa bona, J. de 360.
 Cassel 360.
 Cave, M. 274.
 Cecil, Sir W. 254.
 Cerevisia 315.
 Chamber, M. 264.
 Charles I 54.
 " II 355.
 Chescombe 5.
 Chester, R. 240, 307.
 Chidden 375, 379.
 Chioggia 274.
 Chocolate 24.
 Christchurch 51.
 Christ's Hospital 12.
 Church, A. H. 187, 251.
 Civil war 78.
 Clanfield 89, 378.
 Clifton 76.
 Clinson, Capt. 273.
 Clough 239.
 Cludlie, T. 379.

 Clusius 14, 234, 237, 245, 364-5.
 Cochineal 386.
 Codemer 266.
 Cole, J. 14, 179, 200, 247.
 " E. 12, 373.
 " W. 81, 294.
 Coleman 297.
 Colfe 331.
 Colson 21.
 Compton 2.
 Constantinople 324.
 Cooke, Canon 240.
 Coppels 9.
 Corbet 58.
 Corn 382.
 Courthope 222.
 Cox 97.
 Coys fam. 313.
 " G. 313.
 " Win. 14, 16, 24, 36, 43, 50, 120,
 128-30, 134, 139-47, 150, 156-8,
 160, 210.
 Craig, Dr. J. 248.
 Craster, H. 254, 308.
 Cratonis 210.
 Crete 364.
 Croft 280.
 Crowcher 67.
 Crumpe 64.
 Cusick 79.

 Dale, Dr. J. 84, 92-4, 199-202, 206, 214,
 229-30, 251, 278, 280, 288, 294-8.
 Dale, R. 297.
 " S. 295.
 " W. 295.
 Dallman 238.
 Darlaston 297.
 Davis 48, 171.
 Davys, H. 378.
 Dawley 378.
 De Brina 200.
 De la Broderie 328.
 De Lannoy 247.
 De Laune 14, 19-20.
 Descriptions of Plants 44, 100-96.
 Digges, Sir D. 330.
 Dioscorides 84-9.
 Dixon 237.
 Dodoens 14.
 Double flowers 62.
 Downes 201, 220.
 Downton 378.
 Drope, F. 80, 82.
 Droxford 3, 25, 28, 29, 33, 35, 74, 114,
 116, 118, 122, 126.
 Druce 47, 52, 231.
 Drugs 19, 383.
 Dunne, W. 249.
 Durford Abbey 5, 373.

- Elizabeth 211, 240, 246.
 Eryngo root 241.
 Essex 219.
 Essex, Earl of 58, 241.
 Eton Coll. 274.
 Evelyn 60, 352.
 Everender 31.
 Evergreens 270, 355.
 Exotic plants 237, 358-70.
- Fauntleroy 302.
 Finchdean 6.
 Florence 359.
 Flower lists 240, 308-12, 317-27, 331, 347.
 Forster, R. 249.
 Foulkes-Roberts 239.
 Frankfort 359.
 Franqueville 22.
 Freame 201.
 Frear, T. 249.
 French plants 59.
 Froxfield 381.
 Fruit lists 328, 342-5, 369, 386.
 Fuchs 45.
 Furner 221.
- Gallipoli 364.
 Galls 35, 49.
 Gambier-Parry 283.
 Gammon, W. 96.
- GARDENS, PERSONS.
 Abergavenny, Lord 261.
 Barnaby 306.
 Basket 306.
 Capel 30.
 Coys 312-25.
 Crosse 304.
 Davies 309.
 Delavale 307.
 Edwards 110.
 Forman 309.
 Franqueville 303, 325-6.
 Gerard 18, 247, 303.
 Gibbes 18, 33, 303, 346-8.
 Goodyer 67, 97, 325-7.
 Helin 328.
 Heneage 211.
 Heyden 306.
 James 305.
 Johnson 273.
 Jokkat 329.
 Lobel 247.
 Millen 54, 59.
 Monnel 252.
 Morgan, E. 351-4.
 " " of Bodesclen 308, 353.
 " H. 354.
 Morine 279, 331.
 Morison 354.
- Mount 256.
 Nasmyth 249.
 Norton 306.
 Orleans, Duke of 279.
 Parkinson 270, 327.
 Pointer 58.
 Reresby, Sir J. 350.
 Robine 326, 328, 331, 354.
 Salisbury, Lord 328.
 Salusbury 306.
 Shanne 310.
 Stonehouse 292, 303, 348-51.
 Thornes 308.
 Tradescant 303, 328-46.
 Tuggy 331, 353.
 Uvedale 34.
 Wallop 306.
 Walters 310.
 Wootton, Lord 339-40.
 Wotton 274.
 Watson 305.
 Yalden 28.
 Zouch 247, 324.
- GARDENS, PLACES.
 Blois 354.
 Brussels 329.
 Chioggia 274.
 Constantinople 6, 324.
 Holland 329.
 Paris 329.
 Tournai 252.
 London.
 Chelsea 353.
 Coleman St. 354.
 Hackney 247, 324.
 Highgate 247.
 Holborn 303.
 Lambeth 303, 309, 330-46.
 Long Acre 270.
 Snow Hill 273.
 Tuthill Fields 292.
 Westminster 352-4.
- Essex.
 Stubbers 312-32.
- Hants.
 Droxford 16, 321, 327.
 Dummer 306.
 Sheet 28.
 Southampton 306.
 Winchester 305.
- Herts.
 Hatfield 328.
- Kent.
 Canterbury 339-40.
 East Malling 256.
- Middlesex.
 Enfield 34.
 Twickenham 58.
- Northumberland.
 Seaton 308.
- Oxford 304.

GARDENS, PLACES (*cont.*).

- Somerset.
 Bath 303, 347.
 Suffolk.
 belnys nothers 304.
 Wales.
 Bodesclen 308.
 Lleweni 306-9.
 Yorks.
 Darfield 272, 292, 303, 348.
 Methley 310.
 Thrybergh 350.
 Garret 237.
 Garth 201, 237, 290.
 Gaudin 201.
 Geeres, Mrs. 266.
 George 201.
 Gerard 16, 69, 139.
 Gerard's Herbal 45, 57, 275, 307.
 Gesner 234.
 Ghini 360.
 Gibbon 4.
 Gilbourne 201.
 Gill 81.
 Girdler 97.
 Glasier 385.
 Glyn, T. 287.
 Glynn-lhivona 272.
 Goade 279.
 Gooch 201.
 Good, E. 385.
 Goodyer fam. I.
 " J. Botanical excursions 372 ;
 crest 63 : descriptions of plants 100-
 96 ; garden 67, 97, 325-7 ; house
 64 ; handwriting 55, 169 ; indexing
 203 ; letters 372 ; manor 377 ; manu-
 scripts 229-31, 275, 296, 373-86 ;
 marriage 64 ; plant records 293 ;
 protection order 78 ; recusancy 79 ;
 shopping 48, 73, 381-2 ; will 94.
 Goodyer Plants, *see* Index I.
 " Reginald I.
 Grafting 267.
 Gray 297.
 Green 46, 70.
 Grey 201.
 Guildford 56, 380.
 Guinea 269.
 Gunter, Dr. 98.
 Gwinne, Dr. M. 249, 250.
 Halilah 281.
 Hamden Woods 352.
 Hamelden 378.
 Hampshire lists of villagers 376, 381.
 Hanbury, Th. 2.
 Hanson, Sir T.
 Harrison 59.
 Harvey 201.
 Hasted 67.
 Hatch 378.
 Hatfield 289.
 Hatton, Ld. 279.
 Heacham 57.
 Heath, Eliz. 199, 202.
 Heath, J. 84, 86, 89.
 Heath House 11, 373.
 Heaton 276, 288.
 Heneage 211.
 Henly 64.
 Henrietta Maria 267.
 Henry 201.
 Herbal 45.
 Herbert, J. 81.
 " of Cherbury 383.
 Herrick 221.
 Herring, Dr. F. 249.
 Herts 179, 289.
 Herytage 226.
 Hill, T. 61.
 Hill 172.
 Hinton 56, 58.
 Hoe 378.
 Holloway 277.
 Holybourne 5.
 Holybyn 277.
 Hooper, W. 79.
 Hopton, Lord 78, 286.
 Horse medicine 224.
 Hort, Sir A. 51.
 Horticulture 267.
 How, W. 81, 188-95, 201, 245-51,
 276-94, 298, 300, 354-6.
 How MS. 231-2, 245.
 Huchenson 201.
 Hugobert 59, 247.
 Hull 373.
 Humphrey, L. 221.
 Hunnibon 278, 283, 292.
 Hyde, A. 9, 375.
 " L. 291.
 Ilsley 379.
 India 366.
 Inkforbie 9, 374, 382.
 Inn-Names :
 albus Leon 31, 372 ; Angell 58, 372 ;
 Antilope 180 ; King's Head, Guild-
 ford 56 ; Red Lion, Fleet St. 30,
 372 ; Tabott 73 ; White Hart, South-
 wark 73.
 Inoculation 61.
 Insects 174.
 Iron works 74.
 Ironsmith 201.
 Jackson, B. D. 100, 303, 316.
 " Sir J. 271.
 James 59.
 James I 32, 247, 274.
 Jane, Lady, late Queen 208.

Johnson, S. 49.
 ,, Alice 274.
 ,, Dr. S. 239.
 ,, T. 8, 10, 30, 49, 56, 58, 69,
 71, 81, 201, 271, 273-7, 351, 380.
 Johnson's handwriting 169.
 ,, MSS. 232.
 Jolyff 284, 286.
 Jonson, Ben 58, 240.

Keipe 355.
 Kempe 201.
 Kingsley 377.

Lagouna 210.
 Lake 21.
 Langrish 60.
 Langton 281.
 ,, T. 248, 271.
 Laud 54.
 Laudanum 385.
 Le Seigneur 316.
 Leese, Sir H. 265.
 Lefevre 201.
 Legge 6.
 Leicester 221.
 Leigh 122.
 Leister 316.
 Le Moyne 329.
 Lete 316.
 Library 13, 93, 197-232.
 Lime 17.
 Lincolnshire 265.
 Liphook 5, 385.
 Liss (Leese) 381.
 Lister, Dr. E. 250, 316.
 ,, M. 318.
 Lleweni 238-45, 307.
 Lobel 14, 33, 52, 58, 201, 234, 237,
 245-53, 280, 282, 379.
 Lobel MSS. 231.

LOCALITIES OF PLANTS.

ENGLAND 91, 232, 298.

Berks.

Abingdon 146.
 Cumnor 55.
 Reading 280-1, 301-2.
 Wallingford 301.
 Windsor 288.

Cambridgeshire.

Cambridge 280.
 Downham 193.
 Ely, Isle of 290.
 Over 290.
 Saffron Walden 258.

Dorset.

Portland 379.

Derbyshire 288.*Essex.*

Colchester 292.

Harwich 291.
 Purfleet 113, 118.
 Rumford 67.

Hants.

Alton 47, 290.
 Basingstoke 280, 296.
 Bishops Waltham 122.
 Buriton 184.
 Bursledon 5, 47, 152.
 Chawton 4, 28.
 Chawton 28.
 Christ Church 52.
 Drayton 33, 237, 290.
 Droxford 30, 122.
 Emsworth 28.
 Habridge 127.
 Haliborne 21, 46, 127, 146.
 Hamble 5, 236.
 Hayling 47, 148.
 Hedley 302.
 Holbury House 35, 115, 293.
 Hurst Castle 193.
 Idsworth 5, 50.
 Isle of Wight 33, 90, 285, 290, 353.
 Ladle Hill 194.
 Liphook 196.
 Lymington 51.
 Marborne 46, 127.
 New Forest 35, 115, 293.
 Petersfield 184, 188, 195-6, 280,
 283, 293.
 Poppie Hill 36.
 Rake 52, 179.
 Southsea 47.
 Southwick 123.
 Tichfield 28.
 Vectis, *see* Isle of Wight.
 Winchester 28.

Hereford 288.

Ledbury 284.

Yatton 284.

Herts.

Denham 179.
 Moore Park 37, 179.
 St. Albans 287.

Ireland.

Dublin 287.
 Galloway 288.

Kent 255-63, 275, 283-7.

Addington 257.
 Ashford 283.
 Black brooks 257.
 Canterbury 283.
 Chiselhurst 286.
 Dartford 292.
 Deptford 292.
 Dover 257, 263, 283.
 East Malling 255, 257, 263.
 East Peckham 263.
 Folkestone 263.
 Gravesend 287.

LOCALITIES OF PLANTS (*cont.*).

- Greenhythe 288, 292.
 Greenwich 286, 289.
 Lewes 284.
 Leybourn 258.
 Maidstone 263.
 Rochester 263, 287.
 Sandwich 257.
 Snodland 257.
 Thanet, I. 277, 292.
 Woolwich 281.
 Wrotham 260.
- Lancashire.*
 Blackburn 264.
 Dingley Wood 264.
- London* 352.
 Aldgate 293.
 Bermondsey 289.
 Bermonice 288.
 Chelsea 289.
 Fulham 283.
 Hackney 247, 283, 289.
 Hamstead Heath 232, 280-1.
 Highgate 284, 286.
 London Fields 285, 289.
 Marylebone 296.
 S. Georges Fields 280.
 S. Johns Wood 288.
 S. James 281, 352.
 Totnam 291.
 Tuthill Fields 281.
 Tyburn 296.
 Westminster 296.
 Whitechapel 191.
- Middlesex.*
 Chapel on the Heath 293.
 Hounslow Heath 28, 29.
 Uxbridge 292.
- Northants* 53, 287.
 Lichborough 287, 302.
 Peterboro 289, 290, 292.
 Wellingborough 53, 179, 180.
- Northumbertand.*
 Morpeth 284.
- Norfolk.*
 Kings Lynn 193.
 Witton Fen 287.
- Oxfordshire.*
 Chilswell 301.
 Combe 300.
 Cornbury 301.
 Dorchester 289, 301.
 Headington 300.
 Honey Fixlong 301.
 Oxford 49, 177, 196, 281, 291.
 Sarsden 281.
 Shotover 301.
 Stokenchurch 122, 304.
 Witney 55.
- Shropshire* 284.
 Kinlet 283.
- Somerset* 284, 287.
 Bath 77, 190, 259, 284, 291-3, 301.
 Bristol 75, 287, 291.
 Clover 284, 287.
 Hooms I. 291.
 Mendip 287.
 Nettleton 301.
 St. Vincents Rocks 76.
 Smoccombe Wood 77.
- Surrey.*
 Kingston 181.
 Godalming 55, 180.
 Guildford 185, 286, 288, 302.
 Purbright 293.
 Worplesdown 302.
- Sussex.*
 Darford 46, 125, 178.
 Mayfield 186.
 Rye 296.
 Shorham 300, 302.
- Warwickshire* 280-1, 288, 290.
 Whichford Wood 281.
 Wolver (Wulford) Wood 281 353.
- Wills.*
 Devizes 77.
 Langford 51, 178.
 Stapleford 51, 178.
 Sutton-Veny 28, 54.
 Warminster 28.
- Worcestershire* 283, 301.
 Alchurch 284.
 Kidderminster 284.
- Yorks.*
 Craven 264.
 Darfield 348.
 Ingleborough 222, 288, 291.
 Methley 310.
- Wales, North* 292.
 Bodfari 243.
 Chirk 239, 244.
 Clwyd 244, 309.
 Denbigh 238-45.
 Glynn-lhivona 272.
 Llandulas 242-4.
 Llanrwst 239, 242.
 Llansannen 239, 242.
 Merioneth 243.
 Snowdon 272.
- Loggins 190, 276.
 London 54, 56.
See under Gardens.
 Addresses 48.
 Baynard's Castle 227.
 Gray's Inn 289.
 Great Fire 295.
 Long Acre 270, 296.
 Militia 227.
 Milk St. 294.
 St. Martin's 297.
 Stepney 289.

- London Highway 146, 181.
 Lymphatics 286.
 Lyte Herbal 80.
 Machetes 265.
 Maes, J. 249.
 Maggotts 174.
 Majorca 234.
 Malacca 367.
 Mallinge 73.
 Mangersfield 281.
 Maple Durham 6, 55, 69, 183, 185.
 Mapletoft 222.
 Marshall 215.
 " A. 329.
 Martin 202.
 Massa 201.
 Materia Medica 19, 93, 241, 257, 316, 368.
 Mayerne, T. de 234, 248.
 Mayfield 34 *n.*, 73.
 Meader 193.
 Medeley 201.
 Medical books 93, 385.
 " recipes 241, 383-5.
 Melancholia 274.
 Meon, East 5, 252 *n.*, 297.
 Merrett 35, 62, 81, 195-6, 294.
 Mervyn 1, 178.
 Metheglin 237.
 Methley 264.
 Monnel 252.
 Mooring 67, 122.
 Morden 237.
 Morgan 48.
 " E. 92, 277-8, 292, 351-4.
 " H. 237, 256, 258, 274, 354, 366.
 Morison 92, 279, 293, 352.
 Morrice 74, 370.
 Mosco 63.
 Mostart, D. 332.
 Mouffet, J. 219, 234.
 Moundeford 248, 250.
 Mount 201, 216, 253-63.
 Mundy, P. 333.
 Mussart 332.
 Myddleton 239, 244.
 Myers 373.
 Myres 247.
 Nasmyth, J. 249.
 Neale, J. 93, 385.
 New Forest 51.
 Newark 202.
 Nicholas, Sir E. 355.
 Nid 222.
 Nortons 236.
 Nun, J. 264.
 Ointment 384.
 Orleans 279.
 Osler, Sir W. 254.
 Oxford 48, 81.
 " Colleges:
 Christ Church 58, 89, 221, 238-45.
 Gloucester Hall 49, 176.
 Jesus 238.
 Magdalen 1, 49, 79, 249, 271, 297, 374.
 New 81-3, 274.
 St. Johns 81, 251, 276.
 Wadham 202, 272, 286.
 Worcester 49.
 Paddy, Sir W. 248.
 Padua 202, 299, 318.
 Palestine 364.
 Palmer, R. 9.
 " Ri. 250.
 Palmer worm 151.
 Papists 10.
 Parey's Surgery 70.
 Parish relief for maimed soldiers 213.
 Parker 64.
 " fam. 274.
 Parkinson 16, 27, 74, 140-1, 164, 173, 195, 250-2, 265-71, 293, 318, 330, 358-70.
 Pashley 59.
 Pawlett, Sir Richard 1.
 Pellisserius 252.
 Penckhurst, S. 74, 186.
 Penny 234, 256, 258.
 Pernambuco 269.
 Persia 367.
 Petersfield 2, 64, 297, 373.
 Pharmaceutical Society 20, 272.
 Physicians 20.
 " College of 298.
 Phytologia 91, 251, 276.
 Pioneers 8.
 Pitt, J. 375.
 Plantin 14.
 Ploves 332.
 Pocahontas 57.
 Portesdowne 378.
 Potecary 54, 186.
 Potosi 268.
 Power, Sir D'A. 235, 295-7, 357.
 Prain, Sir D. 351.
 Pratt 2.
 Prelingworth 379.
 Price 76.
 Prices, Building 270.
 " Corn 307.
 Primrose, Dr. 249.
 Priors Reade 1.
 Proclamation 228.
 Provence 190.
 Pulleyn 202, 294.
 Pulteney 14.
 Quicke, W. 318.

- Rabson, J. 378.
 Raleigh 9, 32, 237, 241.
 " last poem 32.
 Ramerus 15, 201.
 Rauwolff 358.
 Rawdon, Sir M. 273.
 Ray 222, 298.
 Read 215.
 Recusants 11, 79.
 Red bridge 376.
 Red Morley 284.
 Red Well 53.
 Redgrave 218.
 Redmayne 57.
 Relf 73.
 Rhé Island 330, 339.
 Rhedus 252.
 Ridley, M. 249.
 Roberts 59.
 " E. 238.
 " Mrs. 309.
 Robin 249.
 Robinson 202, 204, 296.
 Rolfe 57.
 Romney 73.
 Rondelet 14, 201, 246.
 Ropley 2, 12, 373.
 Rother 3.
 Rous 201.
 Rowland 12, 373.
 Ruff 265.
 Russel, C. B. 16, 312.
 Russia 63, 330, 362.

 St. Mary Hill Parish 215.
 St. Vincents Rocks 75.
 Salisbury 29, 378.
 Salter, H. 236.
 Salusbury, Sir J. 238-45.
 " Wm. 238, 242.
 Sambroke, Sir J. 377.
 Sandy Chapel 181.
 Sandy, R. 316.
 Sanfoin 51.
 Saracen 86, 361.
 Savile of Methley 264.
 Sedlescombe 31.
 Selborne 5, 93.
 " maladies 385.
 Seward, P. 7.
 Shakespeare 17, 61, 240, 342.
 Shanne 264-5, 385.
 Sharrock 83.
 Sheete 2, 27, 28, 120, 379.
 Shelden, 288.
 Shelley 12.
 Shelley, Henry 7.
 Shepey 59.
 Shepherd 379.
 Shipley 355.
 Shrewsbury School 201.


 Shute 31.
 Sinai 364.
 Skilling, R. 383.
 Slaynie, H. 331.
 Smith, J. 215.
 Snave, J. 74.
 Snow Hill 273.
 Snuff 383.
 Soberton 5, 35, 115.
 " inhabitants 379.
 Socii itinerantes 272, 274.
 Southampton 159, 185.
 Southropp 1.
 Southwick 375.
 Spain 64, 361.
 Species 61.
 Spigelius 363.
 Splith 359.
 Sprague 189.
 Stable, G. 378.
 Stanford 287.
 Stapf 47, 171, 189, 256-7.
 Stealing children 228.
 Steep 5, 379.
 Stevens 81, 292, 299, 301.
 Steventon 4.
 Stirn 345.
 Stockport 272.
 Stockton 54.
 Stonehouse 15, 49, 79, 80, 82, 231, 264,
 271-3, 277-8, 290, 348-51.
 Stourbridge 297.
 Stroud Common 5.
 Strowde 236.
 Stubbers 16, 17.
 Surrey Assizes 73.
 Sussex 73.
 Swalms 201.
 Swaneling 185.
 Swanmore 379.
 Sydney, Sir H. 241.
 Sylvester 67, 379.
 Symings, Dr. 259.

 Taplow 280.
 Terrier 373.
 Tewe 63, 362.
 Theophrastus 50.
 Thornton 222.
 Thyn 287.
 Tichborne, Sir R. 378.
 Tithing 373.
 Townsend 34, 35.
 Tradescant 55, 83, 272.
 Tribe 379, 380.
 Trinity College, Cambridge 222.
 Tunstal, Sir J. 274.
 Turkey 366.
 Turn[er] 229, 235.
 Turner, R. 234.
 " Wm. 45, 76, 90, 234.

- Twickenham 58.
 Uvedale 8, 11, 34, 161.
 Vachell 11, 375.
 Vaughan, Canon 26, 35, 96, 197, 299.
 Venice 274.
 Vesalius 258.
 Vidian fam. 297.
 Virgil 63.
 Virginian plants 18, 74, 321, 340, 347,
 351, 369.
 Voake, Vok 97, 380, 382.
 Wales 76, 91, 238-45, 272.
 Wall, A. 377.
 Waller, F. 9, 375.
 Walnut 34.
 Walter, Sir W. 281.
 Walton, I. 3, 26.
 Ward, J. 92, 278, 293-5, 304, 352, 357.
 Warminster 378.
 Warwickshire 280-1, 288.
 Watson, Sir W. 335.
 Weather 307, 386.
 Wells 202.
 West, T. 218.
 Westbrook 96, 379.
 Weston 6, 9, 94, 281, 288, 375-6.
 „ Inhabitants of 376.
 Wharton 82.
 Wheatham Green 5.
 White 73.
 „ Gilbert 3.
 Whitrow Moore 184.
 Wickham 8.
 Willisel 298.
 Willoughby, F. 222.
 Willowby, E. 378.
 Wilmot 339.
 Wilts 28, 29, 178, 378.
 Win, W. 331.
 Winchester 21, 56, 163, 235, 286.
 Winter, J. 96.
 Winwood, Sir R. 274.
 Witham 264.
 Withowbie 59.
 Woodcot 11.
 Woolmer Forest 5, 179, 52.
 Woolwich 281.
 Worlidge, A. 95.
 Wotton, E. 234.
 „ H. 65, 96, 201, 271, 274.
 „ Miss M. 1, 65, 79.
 Wray 60.
 Wyche, Sir P. 331.
 Wye 288.
 Wynn 239.
 Yalden, Crusophilus 2.
 „ Dorothe 2.
 „ Edmund 2, 55, 79, 95-7.
 „ William 2, 27, 49, 166.
 Yates 201.
 Yorkshire 264.
 Zouch, Lord 247, 274.

New York Botanical Garden Library

QK21.G7 G8 gen
Gunther, Robert Wil/Early British botani



3 5185 00096 4591

