







LIBRARY
NEW YORK
BOTANICAL
GARDEN

LIBRARY
NEW YORK
BOTANICAL
GARDEN

THE FLORA OF THE DUTCH WEST INDIAN ISLANDS ST. EUSTATIUS, SABA AND ST. MARTIN.

PROEFSCHRIFT TER VERKRIJGING VAN DEN GRAAD
VAN DOCTOR IN DE PLANT- EN DIERKUNDE AAN DE
RIJKS-UNIVERSITEIT TE UTRECHT, OP GEZAG VAN DEN
RECTOR MAGNIFICUS DR. JAN DE VRIES, HOOGLEERAAR
IN DE FACULTEIT DER WIS- EN NATUURKUNDE. VOL-
GENS BESLUIT VAN DEN SENAAAT DER UNIVERSITEIT
TEGEN DE BEDENKINGEN VAN DE FACULTEIT DER
WIS- EN NATUURKUNDE TE VERDEDIGEN OP DONDER-
DAG 1 JULI 1909, DES NAMIDDAGS TE 4 UREN DOOR
ISAÄC BOLDINGH, GEBOREN TE PURMEREND :-: :-:

BOEKHANDEL EN DRUKKERIJ
voorheen
E. J. BRILL. -- LEIDEN
1909.

THE FLORA OF THE DUTCH
WEST INDIAN ISLANDS ST. EUSTATIUS,
SABA AND ST. MARTIN.

I. BOLDINGH.



The Flora of the
Dutch West Indian Islands St. Eustatius,
Saba and St. Martin.

THE FLORA OF THE DUTCH WEST INDIAN ISLANDS ST. EUSTATIUS, SABA AND ST. MARTIN.

PROEFSCHRIFT TER VERKRIJGING VAN DEN GRAAD
VAN DOCTOR IN DE PLANT- EN DIERKUNDE AAN DE
RIJKS-UNIVERSITEIT TE UTRECHT, OP GEZAG VAN DEN
RECTOR MAGNIFICUS DR. JAN DE VRIES, HOOGLEERAAR
IN DE FACULTEIT DER WIS- EN NATUURKUNDE, VOL-
GENS BESLUIT VAN DEN SENAAAT DER UNIVERSITEIT
TEGEN DE BEDENKINGEN VAN DE FACULTEIT DER
WIS- EN NATUURKUNDE TE VERDEDIGEN OP DONDER-
DAG 1 JULI 1909, DES NAMIDDAGS TE 4 UREN DOOR
ISAËC BOLDINGH, GEBOREN TE PURMEREND : : :—:

LIBRARY
NEW YORK
BOTANICAL
GARDEN.

BOEKHANDEL EN DRUKKERIJ
voorheen
E. J. BRILL. — LEIDEN
1909.

QK 250

.B6

Aan mijne Moeder
en aan de nagedachtenis van mijn Vader.

Gaarne de gewoonte volgend om ook op papier een woord te richten tot hen, die de leiders zijn geweest bij de studie in dat levenstijdperk, dat ligt tusschen de eerste inschrijving als Student aan eene Hoogeschool en de Promotie plechtigheid, is het mij een groot voorrecht hier dank te brengen aan U, Hooggeleerde Heeren Professoren in de Faculteit der Wis- en Natuurkunde aan de Utrechtsche Hoogeschool.

Geldt mijn dank in 't algemeen U allen, ik hecht er veel waarde aan in 't bijzonder U te noemen, Hooggeleerde HUBRECHT en WICHMANN.

Maar bovenal Hooggeachte Promotor, Hooggeleerde WENT, mijn dank aan U, bij wien ik het voorrecht heb gehad de laatste jaren van mijn studietijd assistent te zijn. Ik acht mij gelukkig, dat het einde van mijn student-zijn niet als dadelijk gevolg meesleept het einde van mijn assistentschap.

De herinnering aan den grooten steun, dien ik van U mocht ondervinden zal mij steeds bij blijven, tegelijk met die aan de aangename wijze, waarop Gij mij invoerde in de studie der Botanie.

Met grooten dank voor hunnen steun bij mijne studiereis noem ik hier: de Maatschappij tot bevordering van het Natuurkundig Onderzoek der Nederlandsche Koloniën, de Koninklijke Akademie van Wetenschappen te Amsterdam, de Commissie van het F. W. van Eedenfonds en de Directie van het Provinciaal Utrechtsch Genootschap van Kunsten en Wetenschappen.

CONTENTS.

Preface.	Pages.
Additions and Corrections.	v x

FIRST PART.

SYSTEMATICAL. 1

Pteridophyta 1, Gymnospermae 8, Monocotyledoneae 8, Dicotyledoneae 44.

SECOND PART.

HISTORICAL. 209

CHAPTER I. HISTORICAL SKETCH. 211

St. Eustatius 211, Saba 214, St. Martin 214.

CHAPTER II. NOTES ON THE LARGER COLLECTIONS. 216

Collection W. F. J. Suringar 216, Collections Mrs. J. van Grol—Meyers 217, Collection A. C. W. Lionarons 225, Collection I. Boldingh 229.

THIRD PART.

PHYTOGEOGRAPHICAL. 233

SECTION A. OROLOGICAL, GEOLOGICAL AND METEOROLOGICAL.

CHAPTER I. GEOLOGICAL AND OROLOGICAL NOTES. 235

Introduction 235, St. Eustatius 237, Saba 239, St. Martin 241, General notes 243.

CHAPTER II. METEOROLOGICAL NOTES. 245

SECTION B. DISTRIBUTION OF THE WILD PLANTS ENUMERATED IN THE FIRST PART.

CHAPTER I. DISTRIBUTION OF ALL THE WILD PLANTS ENUMERATED IN THE FIRST PART. 249

	Pages.
CHAPTER II. DISTRIBUTION OF THE PLANTS KNOWN ON THE ANTILLES ONLY, ENU- MERATED IN THE FIRST PART.	255
<i>SECTION C. THE VEGETATION OF ST. EUSTATIUS, SABA AND ST. MARTIN.</i>	
CHAPTER I. NOTES ON THE ASPECT OF THE VEGETATION OF THE THREE DUTCH WEST INDIAN ISLANDS.	265
Introduction 265, St. Eustatius 266, Saba 271, St. Martin 276.	
CHAPTER II. DESCRIPTION OF THE DIFFERENT TYPES OF THE VEGETATION.	282
Introduction 282, Eriodendronvegetation 282, Crotonvegetation 286, Littoralvegetation 288, Vegetation of the Cultivated Region 291.	
Literature.	292
Vernacular Names.	295
Index to the First Part.	304
Maps of the Dutch West Indian Islands.	

STELLINGEN.

I.

Bij de bestudeering van de Flora der West-Indische Eilanden moet meer dan tot nu toe geschied is, gelet worden op de verspreiding der planten op elk eiland in 't bijzonder.

II.

De indeeling der Antillen volgens E. Suess is ook om floristische redenen te verkiezen boven die van P. T. Cleve.

III.

Conocarpus erectus Linn. is geen Mangroveplant.

IV.

Batis maritima Linn. behoort niet tot de Orde der Centrospermae.

V.

De nieuwere onderzoekingen over de embryozak der Phanerogamen verminderen de waarde der phylogenetische beschouwingen, zooals die door O. Porsch zijn gegeven.

VI.

Op phylogenetische gronden is het gebruik van het begrip „Gesammtart” niet te verdedigen; ook heeft het geene andere dan verwarrende uitwerking. (Ascher-son und Gräbner: Synopsis der Mitteleuropäischen Flora).

VII.

Onze kennis van de Bacteriaceae en Cyanophyceae is te onvolledig om daaruit conclusies te trekken omtrent de cytologie van de oercel.

VIII.

De Appendicularia zijn geen neotenische Ascidiënlarven.

IX.

De Cetacea vormen eene monophyletische groep.

X.

Hyolithidae zijn verwant met Cephalopoda.

XI.

De White Wall op St. Eustatius is ontstaan door plaatselijke opheffing.

XII.

In een herbarium is alphabetische rangschikking van de Species eener zelfde Plantenfamilie de beste.

**THE FLORA OF THE DUTCH WEST
INDIAN ISLANDS.**

**I. The Flora of St. Eustatius, Saba and
St. Martin.**

THE FLORA OF THE DUTCH WEST INDIAN ISLANDS.

FIRST VOLUME

St. Eustatius, Saba and St. Martin

by

I BOLDINGH.

BOEKHANDEL EN DRUKKERIJ
voorheen
E. J. BRILL — LEIDEN
1909.

The Flora of St. Eustatius, Saba and St. Martin.

by

I. BOLDING.

LIBRARY
NEW YORK
BOTANICAL
GARDEN.



BOEKHANDEL EN DRUKKERIJ
voorheen
E. J. BRILL — LEIDEN
1909.

PREFACE.

Mrs. J. van Grol-Meyers, wife to Mr. G. J. van Grol, administrator of the Dutch West Indian Island St. Eustatius, sent to the „Koloniaal Museum in Haarlem” a nice collection of dried plants collected in St. Eustatius in the years 1904—1906.

She also gave such a collection to the Herbarium of the Botanical Laboratory of the University of Utrecht.

The latter collection of about 350 numbers gave rise to a renewed investigation to the Flora of the Dutch West Indian Islands. I am glad to say that the collection made by Mrs. J. van Grol-Meyers is a very good one. The name of Mr. van Grol himself is well known to the student of the Melocacti of the Dutch West Indian Islands, and the work Mrs. van Grol did now, proved to be a great help for the study of the Flora of the Dutch West Indies. Prof. Dr. F. A. F. C. Went, Director of the Botanical Laboratory and Garden of the University of Utrecht kindly intrusted to me the determination of the collection of the above mentioned West Indian Plants, and in 1905 he proposed to me to make a voyage to the three Dutch West Indian Islands St. Eustatius, Saba and St. Martin in order to get acquainted with the vegetation there and to get a collection of the plants from the three Islands.

I wish to express my thanks to „De Maatschappij ter Bevordering van het Natuurkundig Onderzoek der Nederlandsche Koloniën” („Society for promoting Scientific Investigation in the Dutch Colonies”), De Koninklijke Akademie van Wetenschappen (Korthalsfonds) te Amsterdam, the „Commissie van het F. W. van Eedenfonds in Haarlem” and the „Directie van het Provinciaal Utrechtsch Genootschap van Kunsten en Wetenschappen” (Provincial Utrecht Society of Art and Science), who subsidised and so made it possible for me to undertake the above mentioned investigations.

Most of all I am indebted to the kindness of Prof. F. A. F. C. Went who, after having taken the initiative for my study of the

Flora of the Dutch West Indies, has helped me so energetically by his personal advice as well as by his influence with the above mentioned Societies.

I am glad to have an opportunity now to express my thanks to all those who have helped me in any way in promoting my studies. As to the Dutch Antilles I thankfully mention the Governor of Curaçao Jhr. J. O. de Jong van Beek en Donk and the administrators of St. Eustatius, G. J. van Grol, and T. G. Groebe, the administrator of Saba H. J. Beaujon, the administrator of St. Martin A. J. C. Brouwer and the maire of the French Part of St. Martin D. Beauperthuy.

Moreover: in St. Eustatius: Mrs. J. van Grol-Meyers, Mr. J. Morgan Griffith, Rev. P. Delgueur, Mr. J. G. C. Every, Messr. Hill and Mr. Hassel; in Saba: Dr. A. C. W. Lionarons, Rev. Mulder and Mr. T. Holms; in St. Martin: Mr. J. A. Gravestein, Mr. L. A. van Romond, Mr. A. A. van Romondt, Mr. D. J. van Romondt, Mr. W. E. van Romondt, Mr. W. Netherwood and Mr. F. Morales.

During my voyage to the West Indies I experienced much friendly help from Sir Daniel Morris, Mr. J. R. Bovell, R. P. Duss, Mr. Stephens and Mr. Shepherd.

I also highly appreciate the assistance I got in Holland where I was informed with the greatest loyalty by Mr. L. A. Bakhuis, Dr. J. Boeke, Dr. M. Greshoff, Dr. G. A. F. Molengraaff, Dr. J. Valckenier Suringar and Dr. C. E. A. Wichmann.

According to the wishes of the „Commissie van het F. W. van Eedenfonds” I paid much attention to the cultivated plants too; in a former publication: *Lijst van Planten die door de bewoners van de drie Nederlandsche Antillen, St. Eustatius, Saba en St. Martin als geneeskrachtig worden beschouwd* (Bull. Kolon. Mus. Haarlem n. 38 (2907) p. 93—112), I wrote on the use made of certain plants and in this Flora now I have enumerated all the cultivated plants I met with and which I was able to determine. I also feel very much indebted to the following Directors of „’s Rijks Herbarium” at Leiden viz: Dr. J. M. Janse, Dr. J. P. Lotsy and Dr. J. W. C. Goethart, who greatly facilitated by their kind assistance my working there on the collection of Dr. F. W. R. Suringar.

But most of all I kindly thank Prof. I Urban for his assistance when I was working in the magnificent Herbarium Krug et Urban in the Königl. Bot. Garten und Museum Steglitz-Dahlem (Berlin.).

Although the time I could be there was only very short, my visit

however has been of great use to me owing to the kind assistance of Prof. Urban in looking after doubtful determinations and undetermined plants and in describing some new species. He has enabled me by his great knowledge to bring out the names of some plants that were collected without flowers or fruits.

I feel inclined to express my thanks to Mr. O. E. Schulz and Dr. F. Niedenzu for the determination of some plants and to Dr. H. Hallier, Dr. A. A. Pulle and Dr. S. H. Koorders for their advices. My best thanks also to Mr. S. Perrin who has helped me in looking after the English text of this Flora.

This Flora is divided in Three Parts; whereas the different Parts, Sections and Chapters need no further explanation a short account of the methods I followed in nomenclature, synonymy and distribution of plants is given here.

The enumeration of Families and Genera is very nearly given after the order of De Dalle Torre et Harms: *Genera siphonogamarum ad systema Englerianum Conscripta* (1908).

In a few cases I thought it better to follow:

Urban: *Symbolae Antillanae* vol. I, II, III, IV (fasc. 1 and 2), V, VI (fasc. 1) and Urban: *Addimenta ad Cognitionem Florae Indiae Occidentalis* 1892-97.

The names of the species used are mostly drawn from the two latter papers, Engler und Prantl.: *Die natürlichen Pflanzenfamilie* up to 1909, Engler: *Regnum Vegetabile* up to 1909, or from other monographs.

Whenever I could not find any name in the above mentioned publications I took the name I thought best after the International rules for Botanical Nomenclature chiefly of Vascular Plants. Wien 1905. (*Verh. Int. Bot. Congr.* 1805 Wien.).

Besides the names thus given I also mention the synonyms that are used in: Grisebach: *Flora of the British West Indian Islands*, Duss: *Flore phanérogamique des Antilles françaises*, Millspaugh: *Flora of the Island of St. Croix*, and Jackson: *Index Kewensis* up to 1909 and in a very few cases still another one.

I tried to get aim at the greatest completeness possible in giving the data about the names thus used, and I also mentioned the pages where the plants are given up in: Grisebach: *Flora of the*

British West Indian Islands, Duss: Flore phanérogamique des Antilles françaises, Millspaugh: Flore of the Island of St. Croix and in a few cases still some other work.

To every plant is added the name of the Dutch West Indian Leeward Island where it is found in this order: St. Eustatius, Saba, St. Martin; whenever I saw or collected it also on Anguilla this Island too is mentioned ¹⁾ and when the plant is also known from St. Croix this Island is named after Anguilla.

I have mentioned as much as I could find about the distribution of the plants in all parts of the world, and these data are taken from the following publications in this order:

Urban: Symbolae Antillanae Vol. I, II, III, IV (fasc. 1 and 2), V, VI (fasc. 1) and Additamenta ad Cognitionem Florae Indiae Occidentalis (1892—97), Engler: Regnum vegetabile up to April 1909, Engler und Prantl: Die Natürlichen Pflanzenfamilien up to 1909, newer Monographs and Jackson: Index Kewensis up to 1909.

For the abbreviated names see under Literature.

Prof. Urban was so kind as to allow me the use of the not yet published data from the Herbar Krug et Urban Berlin (Steglitz-Dahlem). A few data are from „'s Rijksherbarium" at Leiden.

Whenever I could not find that a plant was known in other parts of the world besides the Antilles, then the names of all the Islands where it is found are enumerated.

The Systematic Part consists of the plants belonging to the collections of Suringar, van Grol-Meyers, Lionarons and Boldingh.

The numbers with B. added are from the collection Boldingh, marked G. are from the collection van Grol-Meyers when they belong to the collection in the Herbarium at Utrecht, marked H when they belong to the collection in the Koloniaal Museum at Haarlem; the numbers marked L. are from the collection Lionarons and the numbers where no initials are added are from the collection Suringar.

The numbers of the collections Suringar were not given by Suringar when he collected the plants (as Suringar did not put any number on the labels) but they are only the last ciphers of the long number that is fixed to the paper on which the plants are mounted in „'s Rijks Herbarium" at Leiden.

As I am sure that when giving the name under which the plant

¹⁾ See Literature: Boldingh.

is mounted and these ciphers the plant is sufficiently indicated, I thought it better not to give the long numbers that are used in „'s Rijks Herbarium”.

The plants collected in the different Islands are thus given: collection Boldingh, collection Suringar, both in the order of the day of collecting, collections Van Grol-Meyers and collection Lionarons. Not all the vernacular names people told me are given here, but the names I have given in this Flora are used without any doubt. The heights where the plants were collected are given with the greatest accuracy possible.

Whenever a plant struck my attention although I did not think it necessary to collect it, I made some notes about it; these notes are also to be found in this Flora, printed with small type.

As to the flowering and fruiting time of the plants I did not think it necessary to mention with every plant whether I collected flowers on fruits as the greater part of the plants are collected with flowers, either by Suringar, Van Grol-Meyers or by myself.

I wish to draw the attention to the fact that I have tried to render this Flora of the Dutch West Indian Islands as complete as possible, both by what my own investigations have taught me and what I could find elsewhere.

Only the genus *Melocactus* has not been studied with the same care, as I was sure I could not add anything to the work done by Suringar on this head by the data I could bring together without giving too much time to it. So I refer to the studies of F. W. R. Suringar on *Melocactus* and those of J. Valekenier Suringar.

Botanical Laboratory of the University
in Utrecht.

June 1909.

Additions and Corrections.

Page

- 13 Line 19 from top. *For* 1859 *read* 1889.
- 17 Line 9 from top. *After* Millspauch 437 *insert*, *Panicum lanatum* Rottb. in Act. litt. univ. Hafn. I (1778) 269.
- 47 Line 16 from bottom. *After* 81 *insert* 60a.
- 79 Line 4 from top. *For* **macrantha** *read*, **macracantha**.
- 100 Line 15 a from top. *Insert*, **St. Martin**: Tintamarre: n. 3376 B.
- 106 Line 16 from bottom. *After* (Symb.). *insert*. Baham. (Hitchc.).
- 119 Line 1 from top and line 8 from bottom. *For* 1885 *read* 1883.
- 164 Line 15 from top. *For* Antill., Cuba to French Islands. (Gris. Fl.). *read*: Jamaica, Hispaniola, Portorico, St. Thomas, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Barbados. (Herb. Krug et Urban).
- 178 Line 11 from top. *For* Antill. (Gris. Fl.). *read*: St. Thomas, St. Jan, Guadeloupe, Desirade, Les Saintes, Marie-Galante, Dominica, Martinique, St. Vincent, Barbados. (Herb. Krug et Urban).
- 235 Line 9a from bottom. *Insert*, J. W. W. *Spencer*. Reconstruction of the Antillean Continent. Bull. Geol. Soc. Amer. 6. 1894 p. 103—146.

Map of St. Eustatius: Bengalen is to be put to the N. E. of the Quill.

Map of The Dutch W. I. Islands: *For* Barbada *read* Barbuda.

Map of Saba: *For* Marizot *read* Marigot.

CONTENTS.

Preface.	Pages.
Additions and Corrections.	v x

FIRST PART.

SYSTEMATICAL. 1

Pteridophyta 1, Gymnospermae 8, Monocotyledoneae 8, Dicotyledoneae 44.

SECOND PART.

HISTORICAL. 209

CHAPTER I. HISTORICAL SKETCH. 211

St. Eustatius 211, Saba 214, St. Martin 214.

CHAPTER II. NOTES ON THE LARGER COLLECTIONS. 216

Collection W. F. J. Suringar 216, Collections Mrs. J. van Grol—Meyers 217, Collection A. C. W. Lionarons 225, Collection I. Boldingh 229.

THIRD PART.

PHYTOGEOGRAPHICAL. 233

SECTION A. OROLOGICAL, GEOLOGICAL AND METEOROLOGICAL.

CHAPTER I. GEOLOGICAL AND OROLOGICAL NOTES. 235

Introduction 235, St. Eustatius 237, Saba 239, St. Martin 241, General notes 243.

CHAPTER II. METEOROLOGICAL NOTES. 245

SECTION B. DISTRIBUTION OF THE WILD PLANTS ENUMERATED IN THE FIRST PART.

CHAPTER I. DISTRIBUTION OF ALL THE WILD PLANTS ENUMERATED IN THE FIRST PART. 249

	Pages.
CHAPTER II. DISTRIBUTION OF THE PLANTS KNOWN ON THE ANTILLES ONLY, ENU- MERATED IN THE FIRST PART.	255
<i>SECTION C. THE VEGETATION OF ST. EUSTATIUS, SABA AND ST. MARTIN.</i>	
CHAPTER I. NOTES ON THE ASPECT OF THE VEGETATION OF THE THREE DUTCH WEST INDIAN ISLANDS.	265
Introduction 265, St. Eustatius 266, Saba 271, St. Martin 276.	
CHAPTER II. DESCRIPTION OF THE DIFFERENT TYPES OF THE VEGETATION.	282
Introduction 282, Eriodendronvegetation 282, Crotonvegetation 286, Littoralvegetation 288, Vegetation of the Cultivated Region 291.	
Literature.	292
Vernacular Names.	295
Index to the First Part.	304
Maps of the Dutch West Indian Islands.	

FIRST PART

SYSTEMATICAL

Pteridophyta.

Hymenophyllaceae.

Trichomanes Linn.

Trichomanes alatum Sw. in Schrad. Journ. 1800, II (1801) 77; Gris. Fl. 654.

Saba: Mountain 800 M.: n. 2204 B., 2206 B., 1, 6.
Antill., Amer. austral. (Symb.).

Hymenophyllum Smith.

Hymenophyllum hirtellum Sw. Schrad. Journ. 1800, 2. 100; Gris. Fl. 659.

Saba: Mountain 800 M.: n. 1814 B., 1815 B., 1816 B., 2201 B., 7.
Antill., Amer. centr., Mexico. (Christensen 1906.)

Cyatheaceae.

Cyathea J. E. Sm.

Cyathea arborea J. E. Sm. in Mém. Acad. Scienc. Turin V (1790) 417; Gris. Fl. 704.

Saba: Mountain 600 M.: n. 1755 B.
Antill., Bras. (Symb.).

Cyathea serra Willd. Spec. Plant. V. I (1810) 491; Gris. Fl. 704.

St. Eustatius: Top of the Quill: n. 163, 164, 171, 172, 173, 174, 175.

I only saw two specimens of a Fern-tree in the Quill, in a very deep gult and it was impossible for me to get a part of it: most probably it was *Cyathea serra*.

Saba: Mountain near Windwardside 600 M.: n. 2170 B.; Rendez-vous Mountain: n. 46.

Antill., Bras., Mexico. (Symb.).

Hemitelia R. Br.

Hemitelia horrida R. Br. Prodr. I (1810) 158; Gris. Fl. 706.

Saba: Mountain 800 M.: n. 2221 B., 2225 B., 165, 188 L., 189 L.
Antill., Amer. austr. sept. (Symb.).

Alsophila R. Br.

Alsophila phalerata Mart. Ic. Cr. Bras (1834) 67 t. 30 f. l. t. 42.

Saba: Mountain 800 M.: n. 2220 B., 39, 40, 41, 42, 43, 44, 45, 170.

Amer. trop. (Christensen 1906.).

Polypodiaceae.**Dryopteris** Adans.

Dryopteris opposita Urb. in Symbolae Antillanae IV (1903) 14.

Aspidium conterminum Willd. Spec. Plant. V. I. (1810) 249; Gris. Fl. 691.

Nephrodium oppositum Diels in Engl.-Prantl. Nat. Pflanzenfam. I. 4 (1889) 172 (non Hook). (Symb.).

Saba: Gut near Bottom: n. 1434 B.; Top of the Mountain: n. 1798 B., 1808 B., 2151 B., 2195 B.; Mountain near Hellsgate: n. 2285a B.; Mountain: n. 182, 183, 184, 185; Rendez-vous Mountain: n. 186; Crespeen: n. 253 A; locality unknown: n. 1660a B., 220 L.
Antill., Amer. cont. trop. (Symb.).

Dryopteris parasitica O. Ktze. Rev. Gen. Pl. 2 (1891) 811.

Aspidium molle Sw. Schrad. Journ. 1800². 34 (1801); Gris. Fl. 693; Millspaugh 468.

St. Eustatius: slope of the Quill 350 M.: n. 407 B.; inner slope of the Quill 350 M.: n. 463 B.; Quill: n. 116, 122; locality unknown: n. 258 H., 258a G., 276 G.; Bengalen: n. 16.

Saba: Gut near the Bottom: n. 1423a B., 1431a B.; Booby-hill: n. 1592a B.; Hellsgate to Sulphurmines: n. 1673 B.; Mountain: n. 1750a B., 1761a B.; Mountaintop: n. 1832 B.; Sulphurmines to Marypoint: n. 2063a B.; Mountain near Hellsgate: n. 2282a B.; Mountain: n. 224, 230, 234; Rendez-vous. 4—600 M.: n. 169; Jenkeygut: n. 15.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2644 B.; in an

orchard near Bellevue: n. 2832 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2895 B.; hill on the northern side of Cul de sac: n. 3007a B.; Milldrumhill 300—400 M.: n. 3182 B., 3183 B.; Mount Paradis 300—400 M.: n. 3239 B.; gut from Bethlehem to Cul de sac: n. 3413 B.; Centryhill: n. 76.

St. Croix ex Millspaugh.

Baham., Bermud., Antill., Amer. trop., Africa, Asia austro-orient., Australia, Nova Zeal. (Symb.).

Dryopteris reticulata Urb. Symbokae Antillanae IV (1903) 22.

Meniscium reticulatum Sw. in Schrad. Journ. 1800. II (1801) 10; Gris. Fl. 687.

Saba: Mountain 800 M.: n. 1822 B., 2215 B.

Amer. trop. (Christensen 1906).

Aspidium Sw.

Aspidium Plumierii Presl. Rel. Haenk. I (1830) 29.

Aspidium trifoliatum forma Gris. Flor. W. 1. (1864) 695.

St. Eustatius: Quill: n. 123; Bengalen: n. 82; inner slope of the Quill: n. 437 B.

Saba: Mountain 800 M.: n. 2199 B., 1, 2; locality unknown: n. 212 L.

Portorico, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada, Tobago, Trinidad. (Symb.).

Aspidium trifoliatum Sw. in Schrad. Journ. 1800. II (1801) 303; Gris. Fl. 695 (p.p.). (Symb.).

Saba: Mountain 800 M.: n. 1823 B.; Mountain near Hellsgate 600 M.: n. 2286 B.; Crespeen-Rendez-vous 400 M.: n. 3, 4; Gainbygut: n. 227.

Antill., Amer. centr. and austr. (Symb.).

Polybotrya H. et B.

Polybotrya cervina Klfs. Enum. (1824) 55.

Olfersia cervina Kze. in Regensb. Flora VII. I (1824) 312; Gris. Fl. 678.

Saba: Mountain 800 M.: n. 1819 B., 1824 B., 1826 B., 2118a B., 2202 B., Mountain 6—800 M.: n. 70, 71, 72, 168, 202 L., 204 L.

Antill., Amer. cont. trop. (Symb.).

Nephrolepis Schott.

Nephrolepis exaltata Schott Gen. Fil. (1834) ad t. 3: Millspaugh 468.

Aspidium exallatum Sw. in Schrad. Journ. 1800. II (1801) 32; Gris. Fl. 688.

St. Eustatius: Quill: n. 137; Top of the Quill: n. 138, 141, 142, 143, 144; Gilboohill: n. 1125 B.

Saba: Mountain 800 M.: n. 1802 B., 2194 B., 2197 B.; Mountain near Hellsgate 600 M.: n. 2285 B.; Mountain: n. 62, 63, 64, 65, 66, 67; locality unknown: n. 172 L.

St. Croix ex Millspaugh.

Bermud., Antill., Amer. cont. trop., Afr. trop., Asia austro-orient., Polynes., Australia sept. (Symb.).

Diplazium Sw.

Diplazium semihastatum C. Christensen Index Filicum (1906) 239.

Asplenium bipartitum Spr. Syst. IV (1827) 85.

St. Eustatius: Top of the Quill: n. 100.

Cuba, Jamaica, Portorico. (Christensen 1906).

Diplazium striatum Presl. Tent. pterid. (1836) 114.

Asplenium striatum Linn. Spec. (1753) 1082; Gris. Fl. 686.

Saba: Mountain 800 M.: n. 1833 B., 2203 B., 181, 238, 239.

Antill., Amer. cont. trop. (Symb.).

Asplenium Linn.

Asplenium auritum Sw. in Schrad. Journ. 1800 II (1801) 52; Gris. Fl. 682.

W. I.: Suringar: n. 240.

Antill., Amer. cont. trop., Ins. afrie. orient., India orient. (Symb.).

Asplenium cristatum Lam. Enc. 2. (1786) 310.

Asplenium rhizophorum Linn. γ *cicutarium* Sw.; Gris. Fl. 684.

St. Eustatius: Top of the Quill 400 M.: n. 332 B., 333 B.; inner slope of the Quill 350 M.: n. 419 B.; Bottom of the Quill 300 M.: n. 436 B., 451 B.; slope of the lower part of the Quill near Bengalen: n. 730 B.; Bengalen: n. 96, 97, 98; top of the Quill: n. 94; Quill: n. 106, 111, 112, 113, 114; locality unknown: n. 95.

Saba: Springbaygut: n. 2111 B.; Mount Paradis 300—400 M.: n. 3230 B.; Gainbygut: n. 6; Hellsgategut near Devilshand: n. 7; Mountain: n. 2, 3, 4, 8, 10.

Antill., Amer. cont. trop., Afr. trop. (Symb.).

Asplenium monanthes Linn. Mant. (1767) 130.

Saba: Marypointgut: n. 12.

Amer. trop., Ins. Hawaii, Africa and Atlant. Ins. (Christensen 1906).

Asplenium obtusifolium Linn. Spec. (1753) 1080; Gris. Fl. 681.

St. Eustatius: Bottom of the Quill 300 M.: n. 441 B., 467 B. Antill., Amer. cont. trop. (Symb.)

Asplenium pumilum Sw. Prodr. (1788) 129; Gris. Fl. 683.

Saba: Hellsgate to Marypoint: n. 2041 B.

St. Martin: Milldrumhill 300—400 M.: n. 3178 B., 3181 B.; Mount Paradis 300—400 M.: n. 3213a B.

Antill., from Mexico to Columbia. (Symb.).

Asplenium serratum Linn. Spec. (1753) 1079; Gris. Fl. 680.

St. Eustatius: slope of the Quill 350 M.: n. 387 B.; Bottom of the Quill 300 M.: n. 444 B., 450 B.; Quill: n. 179; Top of the Quill: n. 177, 178; locality unknown: n. 150.

Saba: locality unknown: n. 211 L.

Antill., Amer. centr. and austr., Ins. Societ. (Symb.).

Blechnum Linn.

Blechnum occidentale Linn. Syst. X ed. II (1759) 1322; Gris. Fl. p. 673; Millspaugh 469.

St. Eustatius: slope of the Quill 350 M.: n. 385 B.; inner slope of the Quill 350 M.: n. 1225 B.; Top of the Quill: n. 135; slope of the Quill: n. 78, 79, 136.

Saba: Bottom 200 M.: n. 1449 B.; Hellsgate: n. 1689 B.; Springbaygut: n. 2113 B.; 2114 B.; Mountain near Windward side 600 M.: n. 2168 B., 2172 B.; Mountain: n. 35; Mountain 6'—800 M.: n. 23, 233; Peperpot: n. 25, 26; Crespeen—Rendezvous 400 M.: n. 34; Gainbygut: n. 24.

St. Martin: Marigotroad near Marigot: n. 2803 B.; Mount Paradis 300—400 M.: n. 3342 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Stenochlaena J. Sm.

Stenochlaena sorbifolia J. Sm. in Hook. Journ. of Bot. IV (1842) 149; Gris. Fl. 674.

St. Eustatius: Bottom of the Quill 300 M.: n. 470 B., 453 B.; Top of the Quill: n. 242a B.; Quill: n. 130, 131, 132, 133; Boven: n. 80; Bengalen: n. 81.

Saba: Top of the Mountain: n. 221; locality unknown: n. 181 L.

Antill., Amer. cont. trop., Afr. trop., Asia austro-orient., Ins. Monsun., Polynes. (Symb.).

Hemionitis Linn.

Hemionitis palmata Linn. Spec. (1753) 1077; Gris. Fl. 678; Millspaugh 469.

St. Martin: Nakedboyhill 50—200 M.: n. 2716 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2893 B.; Milldrumhill 200—300 M.: n. 3118 B.

St. Croix ex Millspaugh.

Antill., from Mexico to Peru. (Symb.).

Neurogramme Lk.

Neurogramme calomelanos Diels in Engl.-Prantl Nat. Pflanzenfam. I. 4 (1899) 264.

Gymnogramme calomelanos Kaulf. Enum. Fil. (1824) 76; Gris. Fl. 679; Millspaugh 469.

St. Eustatius: in a garden in Oranjestad: n. 68 B.; top of the Quill 380 M.: n. 270 B., 314 B.; slope of the Quill 350 M.: n. 386 B.; slope of the Quill near Bengalen: n. 744 B.; in a garden near Glassbottle: n. 1208 B.; top of Pangahill: n. 128; seashore: n. 125; Boven: n. 126, 127, 129; Greatgut: n. 124; locality unknown: n. 140 G., 189 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1302 B.; 1367 B.; along the seashore near Forthbaylanding: n. 1649 B.; Saddle: n. 1659 B.; Mountain 800 M.: n. 1795 B.; gut between Hellsgate and Masticgut 300—400 M.: n. 2060 B., 2069 B.; Springbaygut: n. 2089 B.; seashore near Forthbaylanding: n. 2148 B.; Mountain near Windwardside 600 M.: n. 2167 B., 2179 B.; Ladderlanding: n. 232, 240; Greathill 200—400 M.: n. 18; Mountain: n. 167; Peperpot: n. 10; Crespeen-Rendez-vous: n. 17; locality unknown: n. 1473 B., 199 L.

I saw *Neurogramme calomelanos* also: near the Sulphurmines; Crespeenhill.

St. Martin: Milldrumhill 300—400 M.: n. 3177 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Afr. trop., Samoa. (Symb.).

Cheilanthes Sw.

Cheilanthes microphylla Sw. Syn. Fil. (1806) 127; Gris. Fl. 667; Millspaugh 469.

St. Eustatius: along roads in Oranjestad: n. 943a B.; Cultuurvlakte; n. 118; Quill: n. 121; Oranjestad: n. 115.

Saba: between Bottom and Marypoint 100—200 M.: n. 1513 B.; seashore near Forthbaylanding: n. 1651 B.; Laddergut: n. 1996 B.; Bottom: n. 228.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Adiantum Linn.

Adiantum tenerum Sw. Prodr. (1788) 135; Gris. Fl. 666; Millspaugh 469. Maiden hair.

St. Eustatius: Mountain (dedit Mr. Falk): n. 104, 105, 106; Bengalen: n. 110; locality unknown: n. 109.

Saba: Paris 400 M.: n. 1373 B.; lower part of the Bottom Mountain 250 M.: n. 1412 B.; along roads between Bottom and Windwardside 200 M.: n. 1320 B.; Ladder: n. 229; Peperpot: n. 13; Greathill: n. 14, 15, 16; Gainbygut: n. 8.

St. Martin: along roads from Colebay to Simsonbay: n. 3247 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Pteris Linn.

Pteris biaurita Linn. Spec. (1753) 1076; Gris. Fl. 669.

Saba: Gut between Hellsgate and Mastiegut 300—400 M.: n. 2066 B.; Springbaygut: n. 2112 B.; Mountain near Hellsgate 600 M.: n. 2283 B.; Peperpot: n. 30, 36, 37, 38; Gainbygut: n. 27, 28, 29.

Antill., Amer., Afr., Asia trop. (Symb.).

Lonchitis Linn.

Lonchitis hirsuta Linn. Spec. (1753) 1078.

Pteris lucinata Willd. Spec. Plant. V. I (1810) 397; Gris. Fl. 669.

Saba: Top of the Mountain 800 M.: n. 2218 B.

Antill., Americ. cont. trop. (Symb.).

Vittaria Sw.

Vittaria lineata Sw. in Schrad. Journ. 1800. II (1801) 72; Gris. Fl. 671.

St. Eustatius: Top of the Quill 400 M.: n. 335 B.; slope of the Quill 350 M.: n. 405 B.; top of the Quill: n. 157, 158.

Baham., Antill., Amer. cont. trop. untill Florida. Afr. trop. untill Cap, Asia austro orient. untill Japon. (Symb.).

Heteropteris Fee.

cf. *Paltonium* Presl. (Christensen 1906).

Heteropteris lanceolata Fée ex Moore Ind. Fil. (1857) 343; Diels in Engl.-Prantl. Nat. Pflanzenfam. I. 4. (1899) 305 fig. 161 E.

Pallonium lanceolatum Pr. Epim. 156 (1849) 156. (Christensen 1906).

Taenitis lanceolata R. Br. Prodr. (1810) 154; Gris. Fl. 671; Millspaugh 469.

St. Eustatius: Top of the Quill 400 M.: n. 243 B., 316^a B.; slope of the Quill near Bengalen 250 M.: n. 693 B.

St. Martin: Mount Paradis 300—400 M.: n. 3231 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. centr. (Symb.).

Polypodium Linn.

Polypodium aureum Linn. Spec. (1753) 1087; Gris. Fl. 698; Millspaugh 469.

St. Eustatius: along roads in Oranjestad: n. 531 B.; slope of the Quill near Bengalen 250 M.: n. 698 B.

Saba: Windwardside—Hellsgate: n. 1661 B.; lower part of the Bottom Mountain 300 M.: n. 1441 B.; Windwardside: n. 1948 B.; Mountain near Hellsgate 600 M.: n. 2287 B.; Quarter: n. 236.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2903 B.; Milldrumhill 300—400 M.: n. 3134 B.; Mount Paradis 300—400 M.: n. 3223 B., 3233 B., 3235 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop. (Symb.).

Polypodium brasiliense Poir. Enc. 5. (1804) 525.

Polypodium nerüfolium Schkuhr Kr. Gew. 1. 14. t. 15. (1806); Gris. Fl. 698.

St. Eustatius: Top of the Quill 400 M.: n. 291 B., 300 B., 321 B.; Top of the Quill 500 M.: n. 485 B., 83, 140; Quill: n. 134, 139.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1294 B.; Parishill 200—400 M.: n. 225, 226.

St. Martin: Milldrumhill 300—400 M.: n. 3144 B.

Amer. trop. (Christensen 1906.).

Polypodium crassifolium Linn. Spec. (1753) 1083; Gris. Fl. 703.

Saba: Mountain 800 M.: n. 2193 B.; 48, 49, 51; locality unknown: n. 61 L.

Antill., Amer. cont. trop. (Symb.).

Polypodium loriceum Linn. Spec. (1753) 1086; Gris. Fl. 698.

Saba: Mountain 800 M.: n. 2191 B., 232, 237.

Antill., Amer. cont. trop. (Symb.).

Polypodium lycopodioides Linn. Spec. (1753) 1082; Gris. Fl. 702. Bastard catclaw.

St. Eustatius: Top of the Quill 380 M.: n. 259 B., 266 B., 408 B.; slope of the Quill near Bengalen 200 M.: n. 692 B.; Signal-hill 200 M.: n. 1051 B.; Slope of the Quill: n. 151, 153; Quill: n. 145, 152; Bengalen: n. 146, 147; locality unknown: n. 179 G., 261 H.

Saba: Paris 400 M.: n. 1374 B., 54; between Bottom and Marypoint 100—200 M.: n. 1499 B.; Mountain 600 M.: n. 1743 B., 57; Windwardside: n. 1941 B.; Springbaygut: n. 2094 B.; Mountain near Hellsgate: n. 2282 B.; Peperpot: n. 55; Crespeen—Rendez-vous 400 M.: n. 60; locality unknown: n. 56, 49 L.

St. Martin: Nakedboyhill 200—260 M.: n. 2722 B., 2724 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2902 B.; Mount Paradis 300—400 M.: n. 3227 B., 3229 B.; Mount Paradis 200—300 M.: n. 3318 B.; Centryhill: n. 75.

Antill., Amer. cont. trop., Afr., Java, Ins. Sandw. (Symb.).

Polypodium phyllitidis Linn. Spec. (1753) 1083; Gris. Fl. 702; Millspaugh 470.

St. Eustatius: Top of the Quill 400 M.: n. 235 B., 320 B.; inner slope of the Quill 350 M.: n. 462 B.; Top of the Quill 500 M.: n. 484 B.; Slope of the Quill near Bengalen 250 M.: n. 704 B., 705 B.; Slope of the Quill: n. 159; Quill: n. 101, 160, 161; Bengalen: n. 162; locality unknown: n. 183 G., 257 H.

Saba: Windwardside—Hellsgate: n. 1667 B.; Springbaygut: n. 2109 B.; Mountain 400—800 M.: n. 52; Gainbygut: n. 47.

St. Martin: Milldrumhill 300—400 M.: n. 3138 B.; French frontier: n. 74.

St. Croix ex Millspaugh.

Baham., Antill., from Florida until Bras. (Symb.).

Polypodium piloselloides Linn. Spec. (1753) 1080; Gris. Fl. 701; Millspaugh 470.

Lepicystis piloselloides Diels in Engl.-Prantl. Nat. Pflanzenfam. I. 4 (1899) 323.

Saba: Boobyhill 300 M.: n. 1581 B.; Mountain 800 M.: n. 1804 B.; St. Giles: n. 1866 B.; Mountain near Windwardside 600 M.: n. 2171 B.; Crispeen-Rendez-vous 400 M.: n. 61, 235; Hellsgategut near Devilshand: n. 53.

St. Martin: Mount Paradis 300—400 M.: n. 3218 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Polypodium plumula H. B. Willd. Sp. V (1810) 178.

St. Eustatius: Top of the Quill 400 M.: n. 222 B., 237 B., 311 B.; slope of the Quill 350 M.: n. 406 B.; slope of the Quill near Bengalen 250 M.: n. 694 B.; slope of the Quill: n. 89, 90; Quill: n. 85, 86, 88; Top of the Quill: n. 37; Bengalen: n. 84.

Saba: Crispeen: n. 68.

Florida, Amer. trop. (Christensen 1906).

Polypodium serrulatum Mett. Fil. hort. Lips. (1856) 30.*Xiphopteris serrulata* Kaulf. Enum. (1824) 85; Gris. Fl. 703.**Saba:** Mountain 800 M.: n. 1834 B.

Antill., Amer. cont. trop., Afr., Ins. Sandw. (Symb.).

Polypodium squamatum Linn. spec. (1753) 1086.*Lepicystis squamata* J. Sm. Brit. and For. Ferns (1866) 80.**St. Eustatius:** slope of the Quill near Bengalen 50—200 M.: n. 643 B., 663 B.**Saba:** along roads between Bottom and Windwardside 200 M.: n. 1306 B.; Mountain 600 M.: n. 1771 B.; Laddergut 0—200 M.: n. 2021 B.; Paris: n. 32; Peperpot: n. 22; Mountain: n. 9; Gainbygut: n. 21; Laddergut: n. 31; Hellsgategut: n. 33.**St. Martin:** Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2856 B.; Mount Paradis 300—400 M.: n. 3232 B.; French frontier: n. 77.

Antill., Amer. cont. from. Mexico to Peru. (Symb.).

Polypodium trichomanoides Sw. Prodr. (1788) 131; Gris. Fl. 701.**Saba:** Mountain 800 M.: n. 1817 B., 1835 B., 1837 B.

Antill., Amer. austr., Ascencion, Asia austro-orient. (Symb.).

Elaphoglossum Schott.**Elaphoglossum conforme** Schott. Gen. ad. t. 14 (1834).*Acrostichum conforme* Sw. Syn. 10 192 t. 1 f. 1 (1806); Gris. Fl. 677.**St. Eustatius:** Top of the Quill 400 M.: n. 302 B., 327 B., 363 B., 384 B., 180; locality unknown: n. 1147a B.**Saba:** Mountain 800 M.: n. 1820 B., 50.

Amer. trop. (Christensen 1906).

Elaphoglossum latifolium J. Sm. in Hook. Lond Journ. Bot. I (1842) 197.*Acrostichum latifolium* Sw. Prodr. (1788) 128; Gris. Fl. 677.**Saba:** Mountain 800 M.: n. 2208 B.

Antill., Amer. cont. trop. (Symb.).

Elaphoglossum petiolatum Urb. in Symbolae Antillanae V (1903) 61.*Acrostichum viscosum* Vent. ap. Sw. Syn. Fil. (1806) 193; Gris. Fl. 676.**St. Eustatius:** Top of the Quill: n. 192a B., 248a B.

Antill., Amer. cont. trop. until Argentina, Africa trop., Asia austro-orient., Ins. Monsun. (Symb.).

Gleicheniaceae.**Gleichenia** J. E. Smith.**Gleichenia bifida** Spreng. Syst. 1V. 1 (1827) 27.*Mertensia furcata* Gris. Fl. W. I. (1861) 651 (non Spreng.). (Symb.).**Saba:** Mountain 800 M.: n. 2227 B., 69.

Antill., Amer. cont. trop. (Symb.).

Ophioglossaceae.**Ophioglossum** Linn.**Ophioglossum reticulatum** Linn. Spec. (1753) 1063; Gris. Fl. 649.**Saba:** Mountain 6—800 M.: n. 73.

Antill., Amer. cont. trop., Africa. (Symb.).

Lycopodiaceae.**Lycopodium** Linn.**Lycopodium taxifolium** Sw. Prodr. (1788) 138; Gris. Fl. 617.**St. Eustatius:** Top of the Quill: n. 479*a* B.**Saba:** Top of the Mountain: n. 2200*a* B., 27.

Antill., Amer. cont. from Mexico to Brasil. austr. (Symb.).

Lycopodium verticillatum Linn. fil. Suppl. (1781) 448; Gris. Fl. 647.**Saba:** Mountain 650 M.: n. 21, 26.

Antill., Amer. cont. trop., Afr. austr. et orient., Polynesia. (Symb.).

Psilotaceae.**Psilotum** Sw.**Psilotum nudum** Gris. Veg. Kar. (1857) 130.*Psilotum triquetrum* Sw. in Schrad. Journ. 1800 II (1801) 109; Gris. Fl. 648.**Saba:** Crepsen-Rendez-vous: n. 1761*a* B., 30.; Paris: n. 20; Greathill: n. 15; Mountain: n. 25.

Baham., Antill., Trop. countries of both hemisph. (Symb.).

Selaginellaceae.**Selaginella** Spring.**Selaginella flabellata** Spring Mon. Lyc. (1849); Gris. Fl. 646.**Saba:** Mountain: n. 1778*a* B., 22, 24, 29, 214 L.

Antill., Amer. merid. sept. (Pulle.)

Selaginella substipitata Spring. Mon. Lyc. (1849); Gris. Fl. 645.

Saba: Mountain 6–800 M.: n. 1813*a* B., 2206*a* B., 2221*a* B., 2227*a* B., 23, 28.

Antill., Panama to Brasil. and Peru. (Gris. Fl.).

Gymnospermae.

6. Pinaceae.

45. *Juniperus* Linn.

Juniperus bermudiana Linn. Spec. (1753) 1039.

Cultivated [Bermudas, Barbados.] (Ind. Kew.).

Juniperus virginiana Linn. Spec. (1753) 1039.

Cultivated [Amer. septentr.] (Ind. Kew.).

Monocotyledoneae.

11. Potamogetonaceae.

60. *Cymodocea* C. H. König.

Cymodocea manatorum Aschers. in Sitzungsber. Ges. naturforsch. Freunde, Berlin (1868) 19; Duss 473; Millspaugh 470.

St. Martin: near Fort Amsterdam: n. 38; Simsonbay lagoon; n. 3260 B.

Suringar saw the *Cymodocea manatorum* also in Simsonbay lagoon.

St. Croix ex Millspaugh.

Bermud., Key West, Antill. (Symb.).

19. Gramineae.

102. *Zea* Linn.

Zea Mays Linn. Spec. (1753) 971.

Cultivated. [Amer. austr. Peru.?] (Symb.).

111. *Saccharum* Linn.

Saccharum officinarum Linn. Spec. (1753) 54.

Cultivated. [Asia austro-orient.?] (Symb.).

134. **Andropogon** Linn.

Andropogon bicornis Linn. Spec. (1753) 1046; Duss 530.

Anatherum bicornis Beauv. agrost. (1812) 15; Gris. Fl. 559.

Saba: Greathill: n. 214; between Bottom and Marypoint 300 M.: n. 1480a B.

Antill., Amer. cont. trop., Teneriffa. (Symb.).

Andropogon contortus Linn. Sp. (1753) 1045; Duss 528.

Andropogon secundus Willd. ex Nees. Agrost. Bras. (1829) 364, Gris. Fl. 558.

St. Eustatius: Near Boven 200 M.: n. 1238 B., 1239 B.

Domingo, Haiti, Antigna, Guadeloupe. (Herb. Krug et Urban.).

Andropogon glomeratus Britton, Stern. et Poggenb. Prel. Cat. (1888) 67.

Andropogon macrouros Michx. Flor. bor. amer. I (1803) 56.

Anatherum macrourum Gris. Flor. W. I. (1864) 559.

Saba: Mountain near Windwardside 600 M.: n. 2169 B.

Baham., Antill., Amer. centr. and sept. (Symb.).

Andropogon Schoenanthus Linn. Spec. (1753) 1046.

Cultivated. [Ind. orient., Afr. trop., Asia.] (Hackel 1859).

Lemon grass.

Andropogon sorghum Brot. Flor. lusit. I (1804) 88.

Cultivated. Creole corn, Guinea corn.

Andropogon zizanioides Urb. Symb. Ant. IV. (1903) 79.

Andropogon squarrosus Linn. fil. Suppl. (1781) 433; Duss 529.

Vetiveria arundinacea Griseb. Flor. W. I. (1864) 559.

St. Eustatius: locality unknown: n. 287 G.

Antill., Bras., from Ind. orient.? (Symb.).

140. **Antheophora** Schreb.

Antheophora hermaphroditica O. Ktze. Rev. II (1891) 759.

Antheophora elegans Schreb. Besch. Gräs. III (1810) 105 t. 44; Gris. Fl. 556; Duss 527; Millspaugh 471.

St. Eustatius: Along roads in Oranjestad: n. 559 B., 1094 B., 1107 B.

Saba: Along roads between Bottom and Windwardside 200 M.: n. 1342 B.; Ladder: n. 376.

St. Martin: Guanabay: n. 2427 B.

St. Croix ex Millspaugh.

Antill., Curaçao, Aruba, Amer. cont. trop. (Symb.).

143. **Tragus** Hall.

Tragus racemosus Haller. Hist. Stirp. Helv. II (1768) 203.

Lappago aliena Spreng. Neue Entd. III (1822) 15; Gris. Fl. 557.

Nazia aliena Millsp. in Millspaugh 471.

St. Eustatius: Between Oranjestad and Fort de Wind on the top of the Cliff: n. 707 B.; Hill near Concordia: n. 946 B.; near White Wall: n. 1259 B.

Saba: St. Giles: n. 1870 B.

St. Martin: From Philipsburg to Guanabay: n. 2367 B.; Naked-boyhill 50—200 M.: n. 2676 B.; from Philipsburg to Point blanche: n. 3404 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Curaçao, Aruba, trop. and subtrop. countries of both hemisph. (Symb.).

161. **Paspalum** Linn.

Paspalum conjugatum Bergius in Act. helv. VII (1772) 129 t. 8; Gris. Fl. 541; Duss 591; Millspaugh 471. Crabgrass.

Saba: Between Bottom and Marypoint 100—200 M.: n. 1470 B.; Mountain 600 M.: n. 1776 B.; Gainbygut: n. 101.

St. Martin Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2907 B.; Mount Paradis 300—400 M.: n. 3242 B.

St. Croix ex Millspaugh.

Antill., trop. countries of both hemisph. (Symb.).

Paspalum fimbriatum H. B. K. Nov. Gen. I (1815) 93 t. 28; Gris. Fl. 542; Duss 511; Millspaugh 472.

St. Eustatius: Along roads in Oranjestad: n. 1109 B.; along roads near Glassbottle: n. 1213 B.; Slope of the Quill: n. 963.

Saba: Along roads between Bottom and Windwardside 200 M.: n. 1290 B.; between Bottom and Marypoint 100—200 M.: n. 1509 B.; along roads 200—300 M.: n. 1711 B.; Laddergut 0—200 M.: n. 2017 B.; Peperpot: n. 1018; Crespeen-Rendez-vous: n. 988.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Amer. austr. (Symb.).

Paspalum hemisphaericum Poir. in Lam. Enc. V (1804) 31.

Paspalum panniculatum Willd. Spec. plant. I (1798) 331; (non Linn.) Gris. Fl. 543; Duss 512. (Symb.).

St. Eustatius: Along roads in Oranjestad: n. 553 B.; slope of the Quill 200 M.: n. 879 B.; slope of the Quill on the side of White Wall 200 M.: n. 1167 B.

Saba: Between Bottom and Marypoint 100—200 M.: n. 1525 B.; Ladderhut: n. 2009 B.; pastureground on Middle Island: n. 2247.

St. Martin: near Mulletpond: n. 2763 B., 2782 B.; in pasture ground near Bellevue: n. 2827 B.; between Belvedere and Oysterpond: n. 3068 B.; Lowlands: n. 3281 B.

Anguilla Boldingh.

Antill., warmer parts of Amer. (Symb.).

165. **Isachne** Pr. Br.

Isachne arundinaceae Gris. Fl. W. I. (1864) 553; Duss 523.

Saba: Top of the Mountain: n. 2196 B.

Jamaica, Portorico. St. Kitts, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada. (Symb.).

Isachne rigens Trin. Gram. Panic. (1836) 252; Gris. Fl. 553; Duss 524.

Saba: Top of the Mountain 800 M.: n. 1386.

Antill., Venezuela. (Gris. Fl.).

166. **Panicum** Linn.

Panicum brevifolium Linn. Spec. (1753) 59; Gris. Fl. 552; Duss 522.

Saba: Lower part of the Bottom Mountain 300 M.: n. 1433 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2031 B.; Springbaygut: n. 2100 B., 2122 B.

Antill., Cuba and Mexico to Bras. (Gris. Fl.).

Panicum colonum Linn. Syst. X ed. II (1759) 870; Gris. Fl. 545; Duss 515; Millspaugh 472.

Saba: Middle Island: n. 1613 B., 1613a B.

St. Martin: Cul de Sac: n. 2614 B.; in pasture ground near Bellevue: n. 2841 B.; Centry hill: n. 606; Colebayhill: n. 207.

St. Croix ex Millspaugh.

Antill., Curaçao, trop. and warmer countries of both hemisph. (Symb.).

Panicum diffusum Sw. Prodr. (1788) 23; Duss 518.

St. Eustatius: Gut near Fort de Wind 50 M.: n. 839 B.

St. Martin: from Philipsburg to Guanabay: n. 2379B.; Marigotroad: n. 2808 B.; in orchard near Bellevue: n. 2838 B.; Mountain between

Mont des Accords and Centry hill: n. 2864 B.; along roads from Colebayhill to Simsonsbay: n. 3015 B.; between Belvedere and Oysterpond: n. 3077 B.; Tintamarre: n. 3394 B.

Anguilla Boldingh.

St. Croix (Symb.).

Cuba, Jamaica, Hispaniola, St. Thomas, St. Jan. St. Barthélemy, Antigua, Guadeloupe, Martinique. (Symb.).

Panicum divaricatum Linn. Syst. X. ed. II (1799) 871; Gris. Fl. 551; Duss 521; Millspaugh 472. Cane grass, wild cane.

St. Eustatius: Top of the Quill 400 M.: n. 238 B.; inner slope of the Quill 350 M.: n. 417 B.; slope of the Quill near Bengalen 150 M.: n. 676 B.; slope of the Quill near Bengalen 100 M.: n. 750 B.; seashore near White Wall: n. 850 B.; from Little Mountain to Venusbay: n. 990 B.; slope of the Quill: n. 584; Sugarloaf: n. 564 B.; Quill: n. 194 G., 234 H.

Saba: between Bottom and Marypoint 400 M.: n. 1484 B.; gut between Hellsgate and Mastiegut: n. 2070 B.; Peperpot: n. 561, 589; Rendez-vous 400 M.: n. 593; locality unknown: n. 303 L.

St. Martin: Marigothill — Marigot 0—200 M.: n. 2645 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2896 B.; Milldrumhill: n. 3050 B.; near Bellevue: n. 3356a B.; Centryhill: n. 559; near the French frontier: n. 568, 586.

I saw *Panicum divaricatum* in St. Martin also on Mount Paradis 300—400 M., hills on the northern side of Cul de Sac and Nakedboyhill 100—200 M.

St. Croix ex Millspaugh.

Key Ins., Baham., Antill., Amer. cont. trop. (Symb.).

Panicum fuscum Sw. Prod. Veg. Ind. Occ. (1788) 23; Gris. Fl. 546; Duss 516; Millspaugh 472.

St. Eustatius: in deserted plantations at the seashore near Bengalen: n. 598 B.; slope of the lower part of the Quill near Bengalen: n. 724 B.; along roads in Oranjestad: n. 1095 B.; Cultuurvlakte: n. 610.

Saba: between Bottom and Marypoint 100—200 M.: n. 1543 B., 1562 B.; Boobyhill: 1583 B.; locality unknown: n. 147 bis L.

St. Martin: Nakedboyhill: n. 2699 B.; along roads near Bethlehem: n. 2935 B.

St. Croix ex Millspaugh.

Jamaica. (Ind. Kew.)

Panicum geminatum Forsk. Flor. aeg. arab. (1775) 18.

Panicum paspaloides Pers. Syn. I (1805) 81; Gris. Fl. 545; Duss 514; Millspaugh 473.

St. Martin: between Belvedere and Oysterpond: n. 3069 B.

Anguilla Boldingh.

St. Croix ex Millspaugh 473.

Florida, Bermud., Baham., Antill., Aruba, trop countries of both hemisp. (excl. Australia.) (Symb.).

Panicum insulare G. F. W. Mey. Prim. Esseq. (1818) 60.

Tricholaena insularis Gris. Flor. W. I. (1864) 557.

Panicum leucophaeum H. B. K. Nov. Gen. I (1815) 97; Duss 522.

Syntherisma insularis Millsp. in Millspaugh 437.

Long grass.

St. Eustatius: in a garden in Oranjestad: n. 67 B., 1063a B.; along roads in Oranjestad: n. 575 B., 1070 B., 1096 B.; slope of the lower part of the Quill near Bengalen: n. 723 B.; slope of the Quill 300 M.: n. 881 B.; along roads near English Quarter: n. 1000 B., 770; near Boven: n. 1251 B.; Cultuurvlakte: n. 771; Mountain near Bengalen: n. 772.

Saba: Boobyhill 300 M.: n. 1579 B.; Laddergut 200 M.: n. 2003 B.; Ladder: n. 1195; Greathill: n. 1206; locality unknown: n. 294 L.

I saw *Panicum insulare* in great quantities also; Gut between Hells-gate and Mastiegut; Crespeenhill; Flat point.

St. Martin: from Philipsburg to Guanabay: n. 2397 B.; lower part of the Milddrumhill: n. 3175 B.; Mount Paradis 300—400 M.: n. 3331 B.

I also saw *Panicum insulare*: Nakedboyhill; Marigothill—Marigot.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida austr., Baham., Amer. cont. from Texas to Patagonia. (Symb.).

Panicum maximum Jacq. Ic. pl. rar. I (1781) 2, t. 13; Gris. Fl. 549; Duss 519; Millspaugh 472. Guinea grass.

St. Eustatius: in a garden in Oranjestad: n. 1063 B.; slope of the Quill: n. 1404; Cultuurvlakte: n. 1402.

Saba: Boobyhill 300 M.: n. 1580 B.; Bottom: n. 1411; locality unknown: n. 240 L.

St. Martin: Top of the hills on the northern side of Cul de Sac: n. 2988 B.

I also saw *Panicum maximum*, and in great quantities: from Bethlehem to Cul de Sac, Top of Mount Paradis 400 M.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop., Afr. trop. and subtrop. (Symb.).

Panicum molle Sw. Prodr. (1788) 2; Gris. Fl. 547; Duss 518.

Panicum barbinode Trin. in Millspaugh 472. Paragrass,

St. Eustatius: Slope of the Quill near Glassbottle: n. 1201 B.

I did not collect this plant in Saba, but from my notes it appears that I saw the plant cultivated in Windwardside.

I do not remember that I saw this plant in St. Martin, but I believe that, as it is cultivated in St. Eustatius and Saba, I may not hold myself entitled to say that *Panicum molle* might not be found in St. Martin as well.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Curaçao, Amer. cont. trop. (Symb.).

Panicum prostratum Lam. Ill. I (1791) 171; Gris. Fl. 546; Duss 515; Millspaugh 473. Runninggrass.

St. Martin: Nakedboyhill 50—200 M.: n. 2698 B.; between Belvedere and Oysterpond: n. 3071 B.; Mount Paradis 150—400 M.: n. 3190 B., 200—300 M.: n. 3315 B.; lowerpart of Milldrumhill: n. 3165 M.: n. 3165 B.; Tintamarre: n. 3377 B., 3383 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Curaçao, trop. countries of both hemisph. (Symb.).

Panicum sanguinale Linn. Spec. (1753) 57; Duss 522.

Digitaria marginata Lk. Enum. I (1821) 102 et *Digitaria setigera* Roth. ap. R. et Sch. Syst. II (1817) 474?: Gris. Fl. 544.

Syntherisma sanguinale Dulac. Flor. Hautes-Pyrén. (1867) 77; Millspaugh 473. Haygrass.

St. Eustatius: along roads in Oranjestad: n. 530 B., 1097 B.; Bengalen: n. 614 B., 748 B.; slope of the Quill near Bengalen 50 M.: n. 662 B.; slope of the Quill: n. 1860; locality unknown: n. 2600.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1326 B.; lower part of the Bottom Mountain 250 M.: n. 1430 B.; between Bottom and Marypoint 100—200 M.: n. 1560 B.; top of the Cliff near the seaside of Boobyhill: n. 1604 B.; Ladderhut 0—200 M.: n. 2013 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2068 B.; locality unknown: n. 2603, 273 L.

St. Martin: from Philipsburg to Guanabay: n. 2363 B.; along the saltpond near Philipsburg: n. 2450 B.; from Philipsburg to Belvedere: n. 2599 B.; in pastureground near Bellevue: n. 2829 B.; top of the hills on the northern side of Cul de Sac 200—300 M.: n.

2987 B.; pastureground near Belvedere: n. 3051 B.; lower part of Milldrumhill: n. 3114 B.; Lowlands: n. 3289 B.; locality unknown: n. 2601.

I also saw *Panicum sanguinale* in Tintamarre.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., trop. countries of both hemisph. (Symb.).

167. *Ichnanthus* Beauv.

Ichnanthus pallens Munro ex Benth. Fl. Hongk. (1861) 414.

Panicum pallens Sw. Prod. Veg. Ind. Occ. (1788) 23; Gris. Fl. 550; Duss 520.

Saba: Top of the Mountain: n. 1794 B., 2213 B.; locality unknown: n. 206 L.

St. Martin: locality unknown: n. 3430a B.

Antill., Guiana Gallica. (Herb. Krug et Urban.).

169. *Oplismenus* Beauv.

Oplismenus setarius R. et Sch. Syst. II (1817) 481; Duss 514.

Orthopogon setarius Spreng. Syst. I (1825) 306; Gris. Fl. 545. comp. Millspaugh 473. Running mountaingrass.

St. Eustatius: inner slope of the Quill 350 M.: n. 414 B.; Quill: n. 260 H., 274 G.

Saba: Lower part of the Bottom Mountain 250 M.: n. 1408 B.; Bottom 200 M.: n. 1453 B.; Boobyhill 350 M.: n. 1600 B.; Mountain near Windwardside 600 M.: n. 2182 B.; locality unknown: n. 53 L.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2643 B.

I also saw *Oplismenus setarius* near Bellevue under old *Mangifera* trees and on the hill between Mont des Accords and Centryhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Antill., Amer. cont. trop. Asia trop. (Symb.).

171. *Setaria* Beauv.

Setaria glauca Beauv. Agrost. (1812) 51; Gris. Fl. 554; Duss 524.

Characthloa glauca (Linn.) Scribn. in U. St. Dep. of Agr. Div. of Agrost. Bull. n. 4 (1897) 39; Millspaugh 473.

St. Eustatius: in a garden in Oranjestad: n. 523 B.; along roads in Oranjestad: n. 1108 B.; slope of the Quill: n. 435.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1289 B., 1328 B.; Mountain 600 M.: n. 1772 B.; locality unknown: n. 133 L.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., trop. and temp. countries of both hemisph. (Symb.).

Setaria setosa Beauv. Agrost. (1812) 51; Gris Fl. 555; Duss 525.

Chaetochloa setosa (Sw.) Scribn. in U. S. Dep. Agr. Div. Agrost. Bull. 4 (1897) 37; Millspaugh 473.

St. Eustatius: Slope of Signalhill: n. 1054 B.; along roads in Oranjestad: n. 1130 B.; slope of Signalhill: n. 451; seashore: n. 386, 389; along roads in Oranjestad: n. 452; locality unknown: n. 278 G.

Saba: Top of the Cliff near the seaside of Boobyhill: n. 1608 B.; Mountain 800 M.: n. 1812 B.; Bottom: n. 1857 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2064 B.

St. Martin: Nakedboyhill 50—200 M.: n. 2688 B., 2694 B.; between Belvedere and Oysterpond: n. 3070 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Setaria viridis Beauv. Agrost. (1812) 51; Gris. Fl. 554.

St. Martin: from Philipsburg to Guanabay: n. 2404 B.

Jamaica, Europa and Asia sept. (Gris. Fl.).

174. **Cenchrus** Linn.

Cenchrus tribuloides Linn. Spec. (1753) 1050; Gris. Fl. 556; Duss 526. Burrgrass.

St. Eustatius: Slope of the lower part of Signalhill: n. 109 B.; slope of the Quill near Bengalen 100 M.: n. 673 B.; Fort de Wind: n. 802 B.; along roads near Glassbottle: n. 863 B.; lower part of the hills near Concordia: n. 984 B.; along roads near Bengalen: n. 1191 B.; locality unknown: n. 249 G., 265 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1283 B.; between Bottom and Marypoint 100—200 M.: n. 1589 B., 1563 B.; along roads: n. 1710 B.; Ladder: n. 1372.

I also saw *Cenchrus tribuloides* in the gut between Hellsgate and Mastiegut

St. Martin: from Philipsburg to Guanabay: n. 2402 B.; Colebay (coll. George): n. 2518 B.; near Mulletpond: n. 2758 B., 2775 B.; in pasture ground near Bellevue: n. 2830 B.; along roads near Bethlehem: n. 2932 B.; Tintamarre: n. 3385 B.

I also saw *Cenchrus tribuloides*: Top of Mount Paradis 400 M.; Naked-boyhill.

Anguilla Boldingh.

Bermud., Baham., Antill., Aruba, Bonaire, Amer. cont. trop. (Symb.).

175. **Pennisetum** L. Cl. Rich.

Pennisetum setosum Rich. in Pers. Syn. I (1805). 72; Gris. Fl. 555; Duss 525.

St. Eustatius: Slope of the Quill: n. 241.

Trop. countries of both hemisph. (Ind. Kew.).

186. **Pharus** Linn.

Pharus glaber H. B. K. Nov. Gen. I (1815) 196; Gris. Fl. 536; Duss 505; Millspaugh 474.

St. Eustatius: Top of the Quill 400 M.: n. 330 B.; inner slope of the Quill 350 M.: n. 413 B., 415 B., 1221 B.; Bottom of the Quill 300 M.: n. 908 B.; slope of the Quill: n. 1380; Quill: n. 1377, 1385, 1379; Bengalen: n. 1382; locality unknown: n. 1381; Quill: n. 225 G.

Saba: Boobyhill 300 M.: n. 1573 B.; 350 M.: n. 1613 B.; locality unknown: n. 34 L.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2858 B.

St. Croix ex Millspaugh.

Antill., Amer. austr. (Symb.).

208. **Aristida** Linn.

Aristida americana Linn. Syst. X ed. II (1759) 879; Gris. Fl. 534; Duss 503.

Bouteloua juncifolia Lag. Gen. et Spec. nov. (1816). 5.

Bouteloua litigosa Lag. Gen. et Spec. nov. (1816). 5. Millspaugh 475. Mulegrass.

St. Eustatius: between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 784 B.; gut near Fort de Wind: n. 829 B.; slope of the Quill: n. 1043, 1055, 1079; near Sugarloaf: n. 1084; locality unknown: n. 215 H.

Saba: Paris 400 M.: n. 1372 B.; along the Forthgut 150—200 M.: n. 1631 B.; along the seashore near Forthbaylanding: n. 1648 B., 2147 B.

St. Martin: Lowlands: n. 3275 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Bonaire, Curaçao, Aruba, Amer. cont. trop. (Symb.).

230. **Sporobolus** R. Br.

Sporobolus indicus R. Br. Prodr. I (1810) 170; Gris. Fl. 533; Duss 503; Millspaugh 474. Lizard grass.

Saba: Paris 400 M.: n. 1376 B.; between Bottom and Marypoint 100—200 M.: n. 1505 B.; Boobyhill 300 M.: n. 1618 B.; St. Johns 250 M.: n. 1730 B.; Windwardside: n. 1916 B.; Rendezvous 400 M.: n. 725.

St. Martin: Along roads from Marigot to Grande Case: n. 3356 B.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Amer. sept., media, austro-occid., Afr., Australia. (Symb.).

Sporobolus virginicus Kth. Rév. Gram. I (1823) 67; Gris. Fl. 533; Duss 503; Millspaugh 474.

St. Eustatius: Seashore near Oranjestad: n. 12 B., 18 B.; seashore near English Quarter: n. 1008 B.; near Boven 200 M.: n. 1236 B.; near White Wall: n. 1261 B.; Concordiabay: n. 563; Turtlebay: n. 566; Pastureground on the slope of the Quill: n. 305; White Wall: n. 302, 315.

Saba: Ladderlanding: n. 562; Paris: n. 1376 B.; Laddergut 50 M.: n. 1975 B.

St. Martin: Along the Saltpond near Philipsburg: n. 2239 B., 2339a B., 2446 B., 2448 B.; Colebay (Coll. George): n. 2507 B.; Fort Willem: n. 2800 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2905 B.; in pastureground near Colebay: n. 3024 B.; between Belvedere and Oysterpond: n. 3067 B.; Lowlands: n. 3276 B.; Fort Amsterdam: n. 300.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Aruba, Amer. cont., Afr. trop., Australia, Ins. Pacif. (Symb.).

282. **Cynodon** Pers.

Cynodon dactylon Pers. Syn. I (1805) 85; Gris. Fl. 540; Duss 509; Millspaugh 474. Bahama grass.

St. Eustatius: Along roads in Oranjestad: n. 1061 B.; slope of the Quill near Glassbottle: n. 1200 B.

St. Martin: Pastureground near Belvedere: n. 3050 B.; pastureground between Bellevue and Oysterpond: n. 3084 B.; along roads from Grande Case to Marigot: n. 3093 B.

St. Croix ex Millspaugh.

This grass as I was told was introduced by Mr. Every from St. Martin

into *St. Eustatius*; in *St. Eustatius* it is cultivated on a very small scale, but in *St. Martin* I saw many pieces of pastureground that consisted chiefly of *Cynodon dactylon*.

Bermud., Baham., Antill., Curaçao, trop. and warmer countries of both hemisph. (Symb.).

288. *Chloris* Sw.

Chloris ciliata Sw. Prod. Veg. ind. occ. (1788) 25; Gris. Fl. 539; Duss 508; Millspaugh 474.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2877 B.

St. Croix ex Millspaugh.

Antill., Mexico. (Gris Fl.).

Chloris paraguayensis Steud. Gram. (1855) 204.

Chloris barbata Sw. Flor. I (1797) 200; Gris. Fl. 549; Duss 508.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1299 B.; along the seashore near Forthbaylanding: n. 1644 B.; along roads 200—300 M.: n. 1712 B.

St. Martin: Cul de Sac: n. 2617 B.; along roads in Prince-quarter: n. 2913 B.

St. Croix (Symb.).

Baham., Antill., trop. countries of both hemisp. (Symb.).

Chloris radiata Sw. Prodr. (1788) 26; Gris. Fl. 539; Duss 507; Millspaugh 475.

Saba: Mountain 600 M.: n. 1777 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop. (Symb.).

304. *Eleusine* Gärtn.

Eleusine indica Gärtn. Fruct. I (1788) 8; Gris. Fl. 540; Duss 508; Millspaugh 475. Dutch grass.

St. Eustatius: along roads in Oranjestad: n. 554 B.; in a garden near Glassbottle: n. 1211 B.; in a garden in Oranjestad: n. 1278 B.; Cultuurvlakte: n. 786; locality unknown: n. 264 G., 237 H.

Saba: lower part of the Bottom Mountain 250 M.: n. 1390 B.; Windwardside: n. 1890 B., 1917a B.; Laddergut 0—200 M.: n. 2014 B.; Bottom: n. 785; locality unknown: n. 23 L.

St. Martin: Colebay (Coll. George): n. 2521 B.; Marigothill—Marigot 0—200 M.: n. 2655 B.; in pastureground near Bellevue: n. 2823 B.; in pastureground between Belvedere and Oysterpond: n. 3086 B.

I also saw *Eleusine indica* in Tintamarre.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Bonaire, Curaçao, trop. and warmer countries of both hemisph. (Symb.).

305. *Dactyloctenium* Willd.

Dactyloctenium aegyptium Willd. Enum. Hort. Berol. (1809) 1029; Gris. Fl. 540; Duss 508; Millspaugh 475.

Eleusine aegyptiaca Desf. Flor. Atl. I (1798) 85.

St. Eustatius: seashore near Oranjestad: n. 39 B.; White Wall: n. 820 B., 1260 B.

Saba: between Bottom and Marypoint 100—200 M.: n. 1514 B.; along the seashore near Forthbaylanding: n. 1647 B.

St. Martin: Marigot—Marigot 0—200 M.: n. 2656 B.; in pastureground near Colebay: n. 3023 B.; Centryhill: n. 654.

St. Croix ex Millspaugh.

Baham., Antill., Bonaire, Curaçao, Aruba, trop. and warmer countries of both hemisph. (Symb.).

307. *Leptochloa* Beauv.

Leptochloa filiformis Beauv. Agrost. (1812) 71; Duss. 507.

Leptochloa mucronata Kth. Rév. Gram. I (1829) 91; Gris. Fl. 537; Millspaugh 475.

St. Martin: Boobyhill 300 M.: n. 1616 B.

St. Croix ex Millspaugh.

Bermud., Antill., Curaçao, Amer. sept., Nova Granata, Asia and Afr. trop. (Symb.).

Leptochloa virgata Beauv. Agrost. (1812) 71; Gris. Fl. 538; Duss 507; Millspaugh 475.

St. Eustatius: Lower part of the slope of the hills near Concordia: n. 940 B.; near Boven: n. 1250 B.

Saba: between Bottom and Marypoint 100—200 M.: n. 1530 B.; Mountain 1751 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2055 B.; Peperpot: n. 55.

St. Croix ex Millspaugh.

Antill., Amer. trop. cont. (Symb.).

310. *Pappophorum* Schreb.

Pappophorum alopecuroideum Vahl Symb. bot. III (1794) 10 t. 51; Gris. Fl. 537.

Pappophorum laguroideum Schrad. in Schult. Mant. II (1824) 342; Duss 506. Crab grass.

St. Eustatius: Between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 772 B.; seashore near White Wall: n. 856 B.; near Boven 200 M.: n. 1233 B.; near White Wall on the top of the Cliff: n. 1255 B.; Cliff: n. 1728; Concordiabay: n. 2236.

Saba: Laddergut 50 M.: n. 1974 B.; Ladder: n. 2231; at the seashore: n. 2137 B.

St. Martin: Tintanarre: n. 3386 B.

Trinidad, La Tinta, Bras. to Uruguay (Gris. Fl.).

341. *Eragrostis* Host.

Eragrostis ciliaris Lk. Hort-Berol I (1827) 192; Gris. Fl. 532; Duss 502; Millspaugh 475.

St. Eustatius: in a garden in Oranjestad: n. 69 B.; Bengalen: n. 615 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 794 B.; along roads in Oranjestad: n. 1073 B.; Cliff: n. 2144; slope of the Quill: n. 2131; Cultuurvlakte: n. 2145.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1362 B.; between Bottom and Marypoint 100—200 M.: n. 1561 B.; top of the Cliff near the seaside of Boobyhill: n. 1602 B.; Mountain 600 M.: n. 1746 B.; Ladder: n. 2080.

St. Martin: Colebay (Coll. George): n. 2511 B.; Nakedboyhill 50—200 M.: n. 2686 B.; in an orchard near Bellevue: n. 2847 B.; Top of the hills on the northern side of Cul de Sac 200—300 M.: n. 2986 B.; in a plantation of cotton near Belvedere: n. 3409 B.;

St. Croix ex Millspaugh.

Baham., Antill., Amer. trop., Afr. trop., Ind. orient. (Symb.).

Eragrostis pilosa Beauv. Agrost. (1812) 71; Gris. Fl. 532; Duss 502; Millspaugh 475.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1308 B.; Crispeen 400 M.: n. 1767 B.; Windwardside in 1891a B.;

St. Martin: Nakedboyhill 50—200 M.: n. 2678 B.; between Mount Vernon and Orientbay: n. 2942 B.; from Belvedere to Oysterpond: n. 3053 B.; lower part of the Milldrumhill: n. 3157 B.

St. Croix ex Millspaugh.

Antill., warmer countries of both hemisph. (Symb.).

Eragrostis tephrosanthos Schult. Syst. Mant. II (1824) 316.

Eragrostis poaeoides Gris. Fl. W. I. (1864) 532; (p. p.). (Symb.).

St. Martin: Fort Willem 0—50 M.: n. 2550 B.; near Oyster-pond: n. 3053 B.

Antill., Amer. austr. (Symb.).

385. *Festuca* Linn.

Festuca Myurus Linn. Spec. (1753) 74.

St. Eustatius: from Signalhill to Tumbledown-dickbay: n. 129a B.; Signalhill: n. 982a B., 1040 B.; near Boven: n. 1237a B. Europa, Amer. and Asia boreal. (Ind. Kew.).

20. Cyperaceae.

Juncellus C. B. Clarke.

cf. 459 *Cyperus* Linn. sect. 2.

Juncellus laevigatus C. B. Clarke in Hook. f. Fl. Brit. Ind. VI (1893) 596.

Cyperus mucronatus Rottb. Descr. et Ic. (1773) p. 19 t. 8 fig. 4; Gris. Fl. 562.

St. Martin: near the saltponds of Philipsburg: n. 2444 B.

Antill., in almost all warmer countries of both hemisph. (Symb.).

459. *Cyperus* Linn.

Cyperus distans Linn. Fil. Suppl. (1781) 103; Gris. Fl. 565; Duss 540.

St. Eustatius: Slope of the Quill 200 M.: n. 23, 29.

Saba: Top of the Mountain: n. 1745a B.; Rendez-vous Mountain: n. 209.

Antill., warmer countries of both hemisph. (Symb.).

Cyperus elegans Linn. Spec. (1762) 68.

Cyperus viscosus Swartz Prod. (1788) 20; Gris. Fl. 563; Duss 537.

Cyperus confertus Gris. Fl. W. I. (1864) 563 pro magna parte, non Swartz. Millspaugh 476. (Symb.).

St. Eustatius: locality unknown: n. 266 H.

St. Martin: along the saltpond near Philipsburg: n. 2447 B.; Cul de Sac—Marigot-hill 0—200 M.: n. 2625 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2906 B.

St. Croix ex Millspaugh.

Florida, Antill., Bonaire, Curaçao, Amer. centr., Argentina. (Symb.).

Cyperus rotundus Linn. Spec. (1762) 67; Gris. Fl. 564; Millspaugh 477. Nutgrass.

Cyperus purpureo-variegatus Boeck. Cyp. Novae. Heft 2 (1890) 37; Duss 539.

St. Eustatius: along roads in Oranjestad: n. 1074 B.; Cultuurvlakte: n. 35; locality unknown: n. 159 G.

Saba: Top of the Cliff near the seashore of Boobyhill: n. 1603 B.; Mountain 800 M.: n. 2186 B.

St. Martin: in pastureground near Bellevue: n. 2828 B.; along roads near Bethlehem: n. 2936 B.; in pastureground between Belvedere and Oysterpond: n. 3085 B..

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., warmer countries of both hemisph. (Symb.).

Mariscus Gärtn.

cf. 459. *Cyperus* Linn. sect. 6.

Mariscus brizaeus C. B. Clarke in Symbolae Antillanae II (1900) 52.

Cyperus bruneus Swartz Fl. Ind. Occ. I (1797) 116 partim. (Symb.); Duss 539. Bullgrass.

St. Eustatius: slope of the Quill near Bengalen 100 M.: n. 671 B., 699 B.; locality unknown: n. 115 G., 262 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1343 B.; near Flat Point: n. 2129 B.; Greathill: n. 46.

I also saw *Mariscus brizaeus*: Green Island.

St. Martin: near Mulletpond: n. 2786 B.; Tintamarre: n. 3369 B.

St. Croix ex Millspaugh.

Baham., Antill., Florida. (Symb.).

Mariscus capillaris Vahl Enum. II (1806) 372.

Cyperus capillaris (Sw.) Millspaugh 476.

Saba: along the Forthgut 150—200 M.: n. 1632 B.; Mountain 600 M.: n. 1738 B.; Springhaygut: n. 2097 B., 2101 B.

St. Martin: from Philipsburg to Guanabay: n. 2356 B.; along roads from Grande Case to Marigot: n. 3103 B.

Anguilla Boldingh.

St. Croix (Symb.).

Cuba, Jamaica, Hispaniola. (Symb.).

Mariscus flavus Vahl Enum. II (1806) 374.

Cyperus flavus Nees. in Linnaea XIX (1847) 698; Millspaugh 476.

Cyperus flavomaris Gris. Fl. W. I. (1864) 567 partim;
Duss 542. (Symb.).

Saba: Mountain 600 M.: n. 1749 B.

St. Croix ex Millspaugh.

Antill., from Mexico and Texas to Patagonia. (Symb.).

Torulinium Desv.

cf. 459. *Cyperus* sect. 5.

Torulinium confertum Desv. in Ham. Prodr. (1825) 15.

Cyperus ferax L. C. Rich. in Acta Soc. Hist. Nat. Paris I
(1792) 106; Duss 541.

St. Eustatius: slope of the Quill: n. 43.

Saba: Mountain 600 M.: n. 1773 B.

Bermud., Baham., Antill., Aruba, in warmer countries of both
hemisph. (Symb.).

462. **Kyllinga** Rottb.

Kyllinga brevifolia Rottb. Deser. et Ic. (1773) p. 13 t. 4 fig. 3;
Gris. Fl. 568; Duss 543; Millspaugh 475.

Kyllinga monocephala Thunb. Fl. Japon. (1784) 35; Gris.
Fl. 568.

St. Martin: Hill on the northern side of Cul de Sac: n. 2977 B.;
Mount Paradis 200—400 M.: n. 3206 B.; Retreat: n. 42.

St. Croix ex Millspaugh.

Bermud., Antill., warmer countries of both hemisph. (Symb.).

469. **Heleocharis** R. Br.

Heleocharis capitata Boeck. in Vidensk. Meddel. Kjob. (1868—69)
159; Duss 545; Millspaugh 477.

Scirpus capitatus Linn. Spec. (1753) 48; Gris. Fl. 570 (partim.).

St. Martin: along the saltpond near Philipsburg: n. 2451 B.

St. Croix ex Millspaugh.

Bermud., Baham., warmer countries of both hemisph. (Symb.).

471. **Fimbristylis** Vahl.

Fimbristylis spathacea Roth. Nov. Pl. Sp. (1821) 24.

Scirpus obtusifolius Gris. Fl. W. I. (1864) 571.

St. Martin: Lowlands: n. 3298a B.

Antill., in trop. contries of both hemisph. (Symb.).

473. **Dichromena** Vahl.

Dichromena radicans Schlecht et Cham. in Linnaea VI (1831) 38.

Rhynchospora Persooniana Gris. Fl. W. I. (1864) 577.

Rhynchospora Humboldtiana Gris. Fl. W. I. (1864) 577 syn. excl. (Symb.).

Saba: between Bottom and Marypoint 300 M.: n. 1582 B.; Crespeen 400 M.: n. 1765 B.; Mountain 600 M.: n. 1775 B., 37.

Antill., Amer. centr. (Symb.).

489. **Cladium** Crantz.

Cladium restioides Benth. in Benth. et Hook. f. Gen. Pl. III (1883) 1066.

Machaerina restioides Vahl. Enum. II (1806) 238; Gris. Fl. 573.

Saba: Mountain 800 M.: n. 2210 B., 42, 208 L.

Cuba, Hispaniola, St. Kitts, Montserrat, Guadeloupe, Dominica, Martinique, St. Vincent. (Symb.).

492. **Rhynchospora** Vahl.

Rhynchospora polyphylla Vahl. Enum. II (1806) 230; Gris. Fl. 575; Duss 551.

Saba: Mountain 600—800 M.: n. 1790 B., 49.

Antill., Amer. centr., Nova Granata, Venezuela. (Symb.).

515. **Scleria** Berg.

Scleria lithosperma Sw. Prod. (1788) 18; Duss 555; Millspaugh 477. Mountaingrass.

Scleria filiformis Sw. Prod. (1788) 19; Gris. Fl. 579.

St. Eustatius: Top of the Quill 400 M.: n. 277a B., 269 H.

St. Croix ex Millspaugh.

Baham., Antill., warmer countries of both hemisph. (Symb.).

Scleria pterota Presl. in Oken Isis XXI (1828) 268; Millspaugh 477.

Scleria pratensis Nees in Mart. Fl. Bras. II pars I (1843) 179 t. 23; Gris. Fl. 577; Duss 553.

Saba: lower part of the Bottom Mountain 300 M.: n. 1432 B.; Mountain 600 M.: n. 1747 B., 1750 B.; locality unknown: n. 51, 182 L., 184 L.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

21. **Palmae.**

528. **Phoenix** Linn.

Phoenix dactylifera Linn. Spec. (1753) 1188.

Cultivated [Afr. sept.] (Symb.).

536. **Thrinax** Linn. f.**Thrinax** spec.**St. Eustatius:** Gilboohill: n. 1120a B.**Saba:** Windwardside—Bottom: n. 298 L.613. **Oreodoxa** Willd.**Oreodoxa caribaea** Damm. et Urb. in Urban Symbolae Antillanae. IV. (1908) 129.

Cultivated. [Portorico.] (Symb.).

631. **Euterpe** Mart.**Euterpe** spec. (prob. *Oenocarpus* Mart.).**Saba:** Mountain: n. 135, 136, 137.663. **Cocos** Linn.**Cocos nucifera** Linn. Spec. (1753) 1188.

Cultivated. [Amer. austr. sept. occid.?] (Symb.).

23. **Araceae.**689. **Anthurium** Schott.**Anthurium cordatum** G. Don. in Sweet. Hort. brit. ed. III (1839) 633. Monkeytail.**St. Eustatius:** Top of the Quill 400 M.: n. 182 B., 232 B., 234 B.; slope of the Quill 350 M.: n. 409 B.; slope of the Quill near Bengalen 250 M.: n. 697 B., 736 B.; slope of the Quill: n. 31, 33, 34, 35; locality unknown: 195 G.**Saba:** Mountain 600 M.: n. 1741 B.; Paris: n. 32; locality unknown: n. 287 L.**St. Martin:** Nakedboyhill 200—265 M.: n. 2720 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2857 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3415 B.; locality unknown: n. 36.**St. Croix** (Engler 1898).

Jamaica, St. Kitts, Martinique. (Engler 1898).

739. **Philodendron** Schott.**Philodendron dispar** Schott. Syn. Ar. (1856) 79; Gris. Fl. 510; Duss 479.

St. Eustatius: Top of the Quill 400 M.: n. 298 B.; slope of the Quill 350 M.: n. 389 B.; Quill: n. 913 B.; Bottom of the Quill 300 M.: n. 432 B., 468 B., 14, 29, 30; locality unknown: n. 220 G.

Saba: Windwardside—Hellsgate: n. 1674 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2034 B.

Guadeloupe, Martinique, St. Vincent. (Symb.).

Philodendron giganteum Schott Syn. Ar. (1856) 89; Duss 479.

St. Eustatius: I saw *Philodendron giganteum* on the top of the Quill.

Saba: Mountain: n. 2275a B.

Portorico. St. Kitts, Guadeloupe, Martinique, St. Vincent. (Symb.).

Philodendron oxycardium Schott Syn. Ar. (1856) 82 (vel. affin.).

St. Eustatius: Top of the Quill: n. 298 B., 394 B., 389 B.; inner slope of the Quill: n. 457 B.

Saba: Mountain: n. 2076a B.; gut near Devilshand: n. 2086 B. Antill., Guatemala. (Symb.).

755. **Colocasia** Schott.

Colocasia esculenta Schott Melet. I (1832) 18.

Cultivated. [India orient.] (Symb.).

758. **Xanthosoma** Schott.

Xanthosoma spec.

Cultivated.

24. Lemnaceae.

795. **Lemna** Linn.

Lemna paucicostata Hegelm. ex Engelm. in A. Gray Man. Bot. V ed. (Jan. 1868) 681; Duss 484.

Lemna minor Griseb. Fl. W. I. (1864) 512 (p. p.). (Symb.): Millspaugh 478.

St. Martin: Lowlands: n. 3285 B.; near Vineyard: n. 3350a B.

St. Croix ex Millspaugh.

Antill., Amer. cont., Afr., Asia austro-orient., Japon., Australia. (Symb.).

32. Bromeliaceae.

840. **Bromelia** Linn.

Bromelia pinguin Linn. Spec. (1753) 285; Gris. Fl. 591; Millspaugh 478. Pingwin.

St. Eustatius: along roads in Oranjestad: n. 103 G.

I saw *Bromelia pinguin* in gardens in Oranjestad.

Saba: Bottom: n. 2292 B.; Peperpot: n. 79, 80.

I also saw *Bromelia pinguin*: Crespeenhill.

St. Martin: along roads from Colebay to Simonsbay: n. 3250 B.

St. Croix ex Millspaugh.

Antill., Amer. cont., Venezuela. (Symb.): also cultivated.

851. **Ananas** Adans.

Ananas sativus Schult. f. in R. et Sch. Syst. VII (1830) 1283.

Cultivated. [Brasilia.] (Symb.).

860. **Wittmackia** Mez.

Wittmackia lingulata Mez. in Mart. Fl. Bras. III. 3. (1891) 274: Duss 570.

Aechmea lingulata Baker in Journ. Bot. XVII (1879) 164.

Saba: Greathill: n. 1377a B., Paris 300 M.: n. 82; Bottom: n. 85; locality unknown: n. 293 L.

St. Martin: Milldrumhill 300—400 M.: n. 3142 B.; Centryhill: n. 106.

St. Thomas, Virgin. Isl., Viequez, Crabben Isl., St. Lucia, Martinique, Guadeloupe, St. Vincent, Beequia, Antigua, Tobago, Trinidad. (Mez 1896).

878. **Pitcairnia** l'Hérit.

Pitcairnia angustifolia Redouté Liliac. II (1805) tab. 76; Gris. Fl. 594; Millspaugh 478. Klapper.

St. Eustatius: Top of the Quill 400 M.: n. 246 B., 304 B., 366 B., 383 B.; along roads in Oranjestad: n. 529 B.; near Maryglory 150 M.: n. 1134 B.; Quill: n. 89, 97; Bengalen: n. 113; Greatgut: n. 88.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1368 B; top of the Cliff near the seaside of Boobyhill: n. 1609 B.; Laddergut 0—200 M.: n. 1973 B.; Paris: n. 115; Wellgut: n. 101.

I also saw *Pitcairnia angustifolia*: near the Sulphurmines: Hellsgate to Marypoint.

St. Croix ex Millspaugh.

Portorico, Grenada, St. Thomas, Antigua, Dominica, St. Vincent, Martinique, Guadeloupe, St. Croix. (Mez 1896).

890. *Tillandsia* Linn.

Tillandsia excelsa Gris. Flor. W. I. (1864). 597.

Saba: Top of the Mountain: n. 2222 B.

Cuba, Jamaica. (Mez 1896).

Tillandsia fasciculata Sw. Prodr. (1788) 56; Gris. Fl. 595; Duss 572.

Saba: St. John's 250 M.: n. 1728 B.; Paris: n. 116; locality unknown: n. 21 L.

St. Martin: Milldrumhill 300—400 M.: n. 3145 B., Centryhill: n. 105.

Florida austral., Baham., Antill., Mexico, Amer. centr. and austro-sept. (Symb.).

Tillandsia recurvata Linn. Spec. (1762) 410; Gris. Fl. 598; Duss 574; Millspaugh 478. Old man's beard.

St. Eustatius: slope of the lower part of Signalhill: n. 108a B.; along roads in Oranjestad: 581 B.; Signalhill: n. 91. 96; locality unknown: n. 217 H.

Saba: along the Forthgut 150—200 M.: n. 1634 B.; Bottom: n. 93.

St. Martin: Princequarter (Coll. George.): n. 2486 B.; Cul de Sac: n. 2620 B.; Nakedboyhill 280 M.: n. 2736 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2937a B.; hills on the eastern side al Cul de Sac n. 3441 B.

I also saw *Tillandsia recurvata*: gut from Bethlehem to Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., warmer countries of Amer. (Symb.).

Tillandsia usneoides Linn. Spec. (1762) 411; Gris. Fl. 598; Duss 574; Millspaugh 478. Old man's beard.

St. Eustatius: locality unknown: n. 295 G.

I also saw *Tillandsia usneoides* near Boven.

Saba: along the Forthgut 150—200 M.: n. 1633 B.; Peperpot: n. 92, 94; Paris: n. 95.

St. Martin: Gut from Bethlehem to Cul de Sac 100—200 M.: n. 3416 B.; top of Centryhill: n. 107.

St. Croix ex Millspaugh.

Antill., warmer countries of Amer. (Symb.).

Tillandsia utriculata Linn. Spec. (1753) 286; Gris. Fl. 596; Duss 573; Millspaugh 478. Wild pine.

St. Eustatius: Top of Signalhill 225 M.: n. 1031 B.; Top of Pongo: n. 90; Top of the Quill: n. 99.

Saba: Slope of the Mountain near St. John's: n. 1735 B.; Paris 200 M.: n. 83, 84; Jenkygut: n. 114; locality unknown: n. 81.

St. Martin: Nakedboyhill 260 M.: n. 2729 B.

St. Croix ex Millspaugh.

Florida austral., Baham., Antill., Curaçao, Mexico, Venezuela, Guyana. (Symb.).

891. *Vriesea* Lindl.

Vriesea guadalupensis Mez in C. de Candolle Monogr. vol. IX (1896) 596. (vel affin.).

St. Eustatius: Top of the Quill: n. 245 B., 315 B.

Saba: Mountain near Windwardside: n. 2181 B.; St. John's: n. 86.

Guadeloupe. (Mez 1896).

892. *Catopsis* Griseb.

Catopsis nutans Griseb. Fl. W. I. (1864) 599; Duss 577.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 57 B.; rocky slope of the hills near Tumbledown dickbay: n. 152 B.; top of the Quill: n. 303 B., 100, 101.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1358 B.; Boobyhill 300 M.: n. 1622 B.; slope of the Mountain near St. John's.: n. 1734 B.

St. Martin: Nakedboyhill 50—200 M.: n. 2709 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2854 B.; Mount Paradis 300—400 M.: n. 3237 B.

Antill., Amer. centr. (Symb.).

33. Commelinaceae.

896. *Commelina* Linn.

Commelina virginica Linn. Spec. (1762) 62; Millspaugh 478.

Commelina elegans H. B. K. Nov. Gen. I (1815) 259; Gris. Fl. 525; Duss 498. Watergrass.

St. Eustatius: seashore near Tumbledown dickbay: n. 158 B.; along roads in Oranjestad: n. 565 B.; slope of the Quill near Bengalen 50—200 M.: n. 640 B., 653 B., 654 B.; Signalhill 150 M.: n. 1053 B.; Mountain near Bengalen: n. 262; Cultuurvlakte: n. 281; locality unknown: n. 2 H., 2 G.

I also saw *Commelina virginica* at the seashore near Oranjestad.

Saba: between Bottom and Marypoint 100—200 M.: n. 1471 B.; 1531 B.; Mountain 400 M.: n. 1770 B.; Laddergut 0—200 M.: n. 2015 B.; Windwardside: n. 2318 B.; Peperpot: n. 284; Jenkeygut: n. 285; Bottom: n. 280; locality unknown: n. 15 L.

St. Martin: From Filipsburg to Guanabay: n. 2381 B.; Mountain between Mont des Accords and Centryhill 0—200 M.: n. 2853 B.; along roads from Colebay to Simonsbay: n. 3249 B.; Mount Paradis 200—300 M.: n. 3307 B.; roads near Colebayhill: n. 264; John Ednyhill: n. 263.

I also saw *Commelina virginica* near Saline; near Bellevue; Nakedboyhill.

St. Croix ex Millspaugh.

Antill., Curaçao, warmer countries of Amer. cont. (Symb.).

911. *Tradescantia* Linn.

Tradescantia geniculata Jacq. Enum. (1760) 18; Gris. Fl. 523; Duss 495.

St. Martin: Milldrumhill 300—400 M.: n. 3141 B.; Mount Paradis 300—400 M.: n. 3222 B., 3346 B.; locality unknown: n. 3622 B.

Antill., Amer. cont. trop., from Mexico to Paraguay, Ins. Sandw. (Symb.).

912. *Callisia* Linn.

Callisia repens Linn. Spec. (1752) 62; Gris. Fl. 524; Duss 497; Millspaugh 479.

St. Eustatius: cultuurvlakte: n. 261.

Saba: between Bottom and Marypoint 100—200 M.: n. 1459a B.; Springbaygut: n. 2099 B.; Bottom: n. 273; locality unknown: n. 311 L.

St. Martin: from Filipsburg to Guanabay: n. 2380 B.; from Filipsburg to Belvedere: n. 2591 B.; hill on the northern side of Cul de Sac 0—100 M.: n. 3000 B.

I also saw *Callisia repens*: on the hill between Mont des Accords and Centryhill; top of Mount Paradis 400 M.

St. Croix ex Millspaugh.

Antill., Mexico, Amer. centr., Venezuela, Peru, Bras. (Symb.).

916. *Rhoeo* Hance.

Rhoeo discolor Hance Misc. in Walp. Ann. III (1852—'53) 660. Cultivated. [Mexico, Antill.] (Symb.).

918. **Zebrina** Schnizl.

Zebrina pendula Schnizl. in Bot. Zeitg. VII (1849) 870; Millspaugh 479.

Saba: Lower part of the Bottom Mountain 250 M.: n. 1416 B.; along roads between Bottom and Windwardside 200 M.: n. 1335 B.; Windwardside n. 1936 B., 2314 B.: Crespeen—Rendez-vous: n. 282; Gainbygut: n. 283; locality unknown: n. 42 L.

St. Croix ex Millspaugh.

Antill., Mexico, Amer. centr. (Symb.).

38. **Liliaceae.**1026. **Aloe** Linn.

Aloe vera Linn. Spec. (1753) 320.

Aloe vulgaris Lam. Enc. I (1783) 86; Gris. Fl. 582; Duss 566; Millspaugh 479. Aloe.

St. Eustatius: locality unknown: n. 3589 B., 102 G.

I also saw Aloe vera on Signalhill.

St. Martin: I saw Aloe vera: along roads from Philipsburg to Naked-boyhill; along roads near Bethlehem; along roads near Belvedere.

Bermud., Antill., Indig. in mediterr. countries and Ins. Canar. (Symb.).

1049. **Allium** Linn.

Allium ascalonicum Linn. Amoen. Acad. IV (1759) 454.

Cultivated. [Everywhere cultivated.] (Ind. Kew.) Charlotten.

1103. **Yucca** Linn.

Yucca aloifolia Linn. Spec. (1753) 319; Gris. Fl. 582; Duss 557. Adam's needle. Spanish dagger.

St. Martin: Near Orientbay: n. 2922a B., 3358 B.

Cuba, Haïti, Jamaica, Tobago. (Herb. Krug et Urban.).

1110. **Sansevieria** Thunb.

Sansevieria guineensis Willd. Spec. Plant. II (1799) 159 Rhammi.

Cultivated. [Afr. trop.] (Symb.).

1151. **Smilax** Linn.**Smilax coriacea** Spr. Syst. Veg. II (1825) 103.*Smilax havanensis* Gris. Flor. W. I. (1864) 586 (pro parte) — non Jacq. (Symb.). White withe.**St. Eustatius:** inner slope of the Quill 350 M.: n. 412 B. 1227; slope of the Quill near Bengalen 150 M.: n. 678 B., 735 B.; locality unknown: n. 188 G., 263 H.**St. Croix.** (Symb.)

Haiti, Domingo, Portorico. (Symb.).

Smilax guianensis Vitman Summa Plant. V (1791) 422.*Smilax macrophylla* Willd. Spec. Plant. IV 2 (1806) 786; Gris. Fl. 586; Duss 560. Basket hoop.**St. Eustatius:** Top of the Quill: n. 189 B.; Quill: n. 2022; Bengalen 150—200 M.: n. 2019.**Saba:** Lower part of the Bottom Mountain 250 M.: n. 1423 B.; Hellsgate: n. 2023.

Antill., Guiana. (Symb.).

Smilax ilicifolia Kth. Enum. V (1850) 171.*Smilax havanensis* Griseb. Flor. W. I. (1864) 586 (p. p. non L.) (Symb.); Millspaugh 479.**St. Martin:** Top of the hill on the northern side of Cul de Sac 200—300 M.: n. 3013 B.; Lowlands: n. 3262 B., 3277 B.**St. Croix** ex Millspaugh.

Cuba. (Symb.).

40. Amaryllidaceae.1181. **Zephyranthes** Herb.**Zephyranthes carinata** Herb. in Bot. Mag. LII (1825) t. 2594.*Amaryllis carinata* Spreng. Syst. Cur. post (1827) 132; Gris. Fl. 584; Duss 562. White snowdrop.**St. Eustatius:** in a garden in Oranjestad: n. 42 H.I also saw *Zephyranthes carinata*: slope of the Quill near Glassbottle.**Saba:** Bottom: n. 3, 4, 5.**St. Martin:** slope of Centryhill: n. 6.

Antill., Mexico. (Symb.). Also cultivated.

1189. **Crinum** Linn.**Crinum americanum** Linn. Spec. (1753) 292.

Cultivated. [Amer. austr.] (Ind. Kew.).

1194. **Hymenocallis** Salisb.

Hymenocallis caribaea Herb. App. Bot. Reg. (1821) 14; Duss 559; Millspaugh 480.

Pancratium caribaeum Linn. Spec. (1753) 291; Gris. Fl. 583. White lily.

St. Eustatius: Glassgut near Oranjestad: n. 51 B.; slope of the Quill near Bengalen 100 M.: n. 674 B.; near Maryglory 150 M.: n. 1148 B.; Oranjestad: n. 25; locality unknown: n. 298 H., 298a G.

Saba: Boobyhill: n. 1572 B.; along the Forthgut 150—200 M.: n. 1635 B.; Bottom: n. 26.

St. Martin: Guanabay: n. 2429 B.; Mountain between Mont des Accords and Centryhill 0—200 M.: n. 2851 B.

St. Croix ex Millspaugh.

Florida, Baham., Antill. (Symb.). Also cultivated.

1196. **Eucharis** Planch.

Eucharis grandiflora Planch. et Linden in Fl. des Serres. Sér. I (IX) (1853—1854) 255.

Cultivated. [New Granada.] (Ind. Kew.)

Eucharis lily.

1208. **Hippeastrum** Herb.

Hippeastrum puniceum Urb. in Symb. Antill. IV. (1903) 151.

Amaryllis equestris Ait. Hort. Kew. I (1789) 417; Gris. Fl. 584.

Hippeastrum equestre Herb. App. Bot. Reg. (1821) 31; Duss 561; Millspaugh 480.

St. Eustatius: Little Mountain: n. 15; locality unknown: n. 23.

Saba: Peperpot: n. 14; Bottom: n. 24.

St. Croix ex Millspaugh.

Antill., Amer. trop. (Symb.). Also cultivated.

1222. **Fourcroya** Vent.

Fourcroya gigantea Vent. in Bul. Soc. Philom. I (1793) 65; Gris. Fl. 582; Duss 558. Karata, Pita.

St. Eustatius: along roads in Oranjestad: n. 191 G.

I also saw *Fourcroya gigantea* on the top of the Cliff near Gallowsbay; Bergje; slope of Signalhill.

St. Martin: along roads near Belyedere: n. 3359 B.
Antigua, French Islands to Bras. (Gris. Fl.).

1230. **Hypoxis** Linn.

Hypoxis decumbens Linn. Syst. X ed. II (1759) 986; Gris. Fl. 585; Duss 562. (procumbens errore).

Saba: Mountain 600—800 M.: n. 1785 B., 1796 B., 2191 B.;
Mountain near Windwardside 600 M.: n. 2161 B.; Jenckygut: n. 2.
Antill., Amer. cont. trop. (Symb.).

43. **Dioscoreaceae.**

1252. **Dioscorea** Linn.

Dioscorea alata Linn. Spec. (1753) 1033; Gris. Fl. 587. Yams.

St. Eustatius: locality unknown: n. 245 H.

Saba: Mountain near Windwardside: n. 2157 B.

Antill. Patria veris. Asia austro-orient. (Symb.). Also cultivated.

Dioscorea trifida Linn. f. Suppl. (1781) 427; Gris. Fl. 587.

Saba: Top of the Mountain 800 M.: n. 1789 B.; Mountain near Windwardside: n. 2155 B.

Antill., Patria Amer.-austral. (Symb.).

44. **Iridaceae.**

1285. **Belamcanda** Adans.

Belamcanda punctata Moench. Meth. Suppl. (1794) 214.

Cultivated. [China.] (Ind. Kew.).

1292. **Eleutherine** Herb.

Eleutherine pilcata Herb. in Bot. Reg. (1843) t. 57. Red root.

Cultivated. [India occid.] (Ind. Kew.)

45. **Musaceae.**

1318. **Musa** Linn.

Musa paradisiaca Linn. Spec. (1753) 1043.

Musa sapientum Linn. Syst. X ed. II (1759) 1303.

Cultivated. [India orient.] (Symb.).

1321. **Heliconia** Linn.

Heliconia Bihai Linn. Mant. II (1771) 211; *Gris. Fl. 600; Duss 579. Wild banana.

St. Eustatius: Bottom of the Quill 300 M.: n. 449 B.; Quill: n. 1077 B., 295, 223 G.

Saba: Mountain 800 M.: n. 2223 B.

Antill., Amer. cont. trop. (Symb.).

46. **Zingiberaceae.**1357. **Costus** Linn.

Costus speciosus Sm. in Trans. Linn. Soc. I (1791) 249. Shell-plant.

Cultivated. [Ins. Malay.] (Schumann 1904.).

47. **Cannaceae.**1363. **Canna** Linn.

Canna indica Linn. Spec. (1753) 1. Cannon, Indian shot.

Cultivated. [Cosmop. trop.] (Ind. Kew.).

48. **Marantaceae.**1370. **Maranta** Linn.

Maranta arundinacea Linn. Spec. (1753). 2. Arrowroot.

Cultivated. [Baham., Antill., Amer. cont. trop.] (Symb.).

50. **Orchidaceae.**1490. **Spiranthes** L. Cl. Rich.

Spiranthes elata L. Cl. Rich. Orch. Eur. Adn. (1818) 37; Gris. Fl. 641.

St. Martin: Centryhill: n. 72.

Antill., Amer. austr. (Symb.).

1493. **Stenorrhynchus** L. Cl. Rich.

Stenorrhynchus orchioïdes L. Cl. Rich. Orch. Eur. (1817) 37; Gris. Fl. 642.

Stenorrhynchus lanceolatus L. Cl. Rich. l. c.; Gris. Fl. 642.

St. Eustatius: Top of the Quill 400 M.: n. 229a B.

Baham., Antill., Amer. cont. trop. (Symb.).

1498. **Physurus** L. Cl. Rich.

Physurus hirtellus Lindl. Gen. and Spec. Orch. (1840) 501; Gris. Fl. 642; Duss 606.

St. Eustatius: Top of the Quill: n. 65, 66.

Cuba, Jamaica, Portorico, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada, Trinidad. (Symb.).

Physurus plantagineus Lindl. Gen. and Sp. Orch. (1840) 503; Gris. Fl. 643; Duss 605.

St. Eustatius: Top of the Quill 400 M.: n. 229 B.

Saba: Mountain near Hellsgate: n. 228*a* B.; Mountain 600—800 M.: n. 1, 2, 3, 4, 67, 68, 69, 176 L.

Cuba, Jamaica, Portorico, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada, Trinidad. (Symb.).

1527. **Prescottia** Lindl.

Prescottia stachyoides Lindl. in Bot. Reg. (1834) sub t. 1915; Gris. Fl. 639; Duss 603.

St. Eustatius: Top of the Quill: n. 71.

Antill., Amer. centr., Bras. (Symb.).

1553. **Microstylis** Nutt.

Microstylis spicata Lindl. Gen. and Sp. Orch. (1830) 19; Gris. Fl. 612; Duss 590.

Saba: Top of the Mountain 6—800 M.: n. 1828 B., 2224 B., 2228 B.

Cuba, Jamaica, Hispaniola, Martinique. (Symb.).

1565. **Polystachya** Hook.

Polystachya luteola Hook. Exot. Flor. (1825) tab 103; Gris. Fl. 628; Duss 597.

St. Eustatius: Quill: n. 70.

Baham., Antill., Amer. trop. (Symb.).

1614. **Epidendrum** Linn.

Epidendrum anceps Jacq. Sel. (1763) 224 t. 138.

Epidendrum fuscatum Sm. Spic. bot. (1791) 21 t. 23; Gris. Fl. 617; Duss 591.

St. Eustatius: Top of the Quill 400 M.: n. 250 B., 381 B.; top of the Quill: n. 92, 93, 95; Quill: n. 96; locality unknown: n. 94.

Saba: Windwardside—Hellsgate: n. 1666 B.; Crispeen 400 M.:

n. 1763 B.; Mountain 800 M.: n. 1818 B.; Springbaygut: n. 2090 B., 2108 B.; Mountain 600 M.: n. 117; Mountain 800 M.: n. 97; Mastiegut: n. 119; Wellridge: n. 118.

St. Martin: Milldrumhill 300—400 M.: n. 3149 B.

Antill., Amer. cont. trop. (Symb.).

Epidendrum ciliare Linn. Syst. X ed. (1759) 1246; Gris. Fl. 615; Duss 590; Millspaugh 481.

St. Eustatius: Slope of the Quill near Bengalen: n. 715 B.; top, Panga hill: n. 101; top of the Quill: n. 102; Boven: n. 103; locality unknown: n. 281 G., 268 H., 308 G.

Saba: between Bottom and Marypoint 250 M.: n. 1554 B.; Boobyhill 300 M.; n. 1597 B.; Crispeen 400 M.: n. 1727 B.; Paris 2—400 M.: n. 80, 99; Greathill 2—400 M.: n. 81; Rendez-vous 400 M.: n. 100; Hellsgate: n. 98; locality unknown: n. 165 L.

St. Martin: Nakedboyhill 200—260 M.: n. 2727 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2897 B.; Milldrumhill 300—400 M.: n. 3154 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3421 B.; Centryhill: n. 104, 105; French frontier: n. 106.

St. Croix ex Millspaugh.

Antill., Amer. austr-sept. and centr. (Symb.).

Epidendrum difforme Jacq. Enum. (1760) 29; Duss 592.

Epidendrum umbellatum Sw. Prodr. (1788) 121; Gris. Fl. 618.

St. Eustatius: Top of the Quill 400 M.: n. 241 B., 107, 108.

Antill., Amer. trop. (Symb.).

Epidendrum globosum Jacq. Enum. (1760) 29; Gris. Fl. 619; Duss 594.

St. Eustatius: Top of the Quill: n. 83.

Saba: Mountain 800 M.: n. 1827 B.; Mountain near Windwardside 600 M.: n. 2177 B.; Mountain: n. 82; Rendez-vous 400 M.: n. 84.

Antill., Amer. cont. trop. (Symb.).

Epidendrum papilionaceum Vahl in West St. Croix (1793) 303.

St. Eustatius: Signalhill: n. 110 B.; Boven: n. 112; locality unknown: n. 280a G.

Saba: Forthbay 30 M.: n. 5.

St. Martin: near Mulletpond: n. 2784 B.; Lowlands: n. 3254 B., 3278 B.

Baham., Antill. (Symb.).

Epidendrum strobiliferum Reichb. f. in Nederl. Kruidk. Archief. IV. (1859) 333; Gris. Fl. 618; Duss 592.

Saba: Rendez-vous 400 M.: n. 109.

St. Croix ex Millspaugh.

Jamaica, Trinidad, Guiana. (Gris. Fl.).

1619. **Brassavola** R. Br.

Brassavola cucullata R. Br. in Ait. Hort. Kew. ed. II V. (1813) 216; Gris. Fl. 620; Duss 594.

St. Eustatius: Lower part of the slope of the hills near Concordia: n. 941 B.; Boven: n. 90.

Saba: Greathill 2—400 M.: n. 91; Mastiegut: n. 110.

St. Martin: Milldrumhill 300—400 M.: n. 3139 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3419 B.; Centryhill: n. 111.

Antill., Mexico to Venezuela. (Gris. Fl.).

Brassavola rigida (comb. nov.).

Tetramicra rigida Lindl. Gen. et Sp. Orch. (1831) 119; Gris. Fl. 622.

St. Eustatius: Gut near Fort de Wind 50 M.: n. 825 B.; Little Mountain n. 75; Greatgut: n. 73, 74, 78; locality unknown: n. 76, 77, 51 H.

Ind. occid. (Ind. Kew.).

1733. **Ornithidium** Salisb.

Ornithidium coccineum Salisb. in Trans. Hort. Soc. I (1812) 293; Gris. Fl. 626; Duss 596.

Saba: Mountain 800 M.: n. 1801 B.

Antill., Venezuela. (Symb.).

1779. **Oncidium** Sw.

Oncidium Spec. veris: 1)

St. Eustatius: Signalhill 200 M.: n. 134*** B., 1049 B***.; near White Wall: n. 849*** B.; Signalhill—Red Cliff: n. 56*, 58*, 60*; slope of Signalhill: n. 61***; Signalhill: n. 57*; Red Cliff: n. 62*; White Wall: n. 63*; Grandgut: n. 64*; locality unknown: n. 50*** H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1469*** B., 2039*** B.; Luddergut: n. 53***; Peperpot: n. 54**; Paris: n. 55**.

St. Martin: Nakedboyhill 280 M.: n. 2733*** B.; Centryhill: n. 52**; Princequarter: n. 51**, 59; road near Oysterpond: n. 50**.

Antill. (Symb.).

1) *) *Oncidium variegatum* Sw. in Act. Holm. XXI (1800) 240.

**) *Oncidium sylvestre* Lindl. in Ann. and Mag. Nat. Hist. III Ser. I (1858) 332.

***) *Oncidium tetrapetalum* W. (L.) Gris. Fl. 631.

Dicotyledoneae.

51. Casuarinaceae.

1855. **Casuarina** Linn.

Casuarina equisetifolia Linn. Amoen. IV (1759) 143. Casuarine.
Cultivated. [Afr. occid., Australia, Ins. Pacif. and Malay.]. (Symb.).

53. Piperaceae.

1862. **Piper** Linn.

Piper dilatatum L. Cl. Rich. in Act. Soc. Hist. nat. Par. (1792) 105.
Artanthe Bredemeyeri Griseb. Flor. W. Ind. (1859) 170.

Piper Bredemeyeri Jacq. Eclog. (1811—44) 125 f. 84; Millspaugh 481.

St. Eustatius: Inner slope of the Quill 350 M.: n. 416 B., 424 B. top of the Quill 500 M.: n. 491 B., slope of the Quill near Bengalen 200 M.: n. 706; Quill 2566, 2560, 2501; locality unknown: n. 174 G.

Saba: Lower part of the Bottom Mountain 300 M.: n. 1439 B.; Bottom 200 M.: n. 1451 B.; Laddergut 0—200 M.: n. 1964 B.; gut between Masticgut and Hellsgate 300—400 M.: n. 2061; Crespeen—Rendez-vous 400 M.: n. 2511; Hellsgategut: n. 2569.

St. Martin: Hills on the northern side of Cul de Sac 100—200 M.: n. 2981 B.; Mount Paradis 300—400 M.: n. 3219 B.

St. Croix ex Millspaugh.

Antill., Mexico, Costarica, Panama, Columbia, Venezuela, Bras. (Symb.).

Piper marginatum Jacq. Ic. rar. II (1786) 2 t. 215.

Schillera caudata Kunth in Linnaea XIII (1839) 716; Gris. Fl. 170.

Saba: locality unknown: n. 159 L.

Antill., Panama, Surinam., Guiana gallica, Venezuela, Bras., Peru. (Symb.).

Piper medium Jacq. Ic. rar. I (1781) 2 t. 8.

Piper ceanothifolium Kunth. in H. B. K. Nov. Gen. I (1815) 26; Duss 177.

Enckea Sieberi Gris. Flor. W. I. (1859) 177 (p. p.) (Symb.).

St. Eustatius: Quill: n. 459 B.

Antill., Amer. cont. (Symb.).

Piper peltatum Linn. Spec. (1753) 30.*Heckeria peltata* Kunth in Linnaea XIII (1839) 565.*Pothomorphe peltata* Miq. Comm. phyt. (1838) 37; Gris. Fl. 169; Duss 175; Millspaugh 481.**Saba**: Mountain 500 M.: n. 899; Gainbygut: n. 888, 889, 895. Antill., Amer. cont. centr. and austral. (Symb.).**Piper reticulatum** Linn. Spec. (1753) 29.*Enckea smilacifolia* Griseb. Flor. W. Ind. (1859) 169. (excl. syn. Willd.) (Symb.).*Piper smilacifolium* C. DC. Prodr. XVI, 1 (1869) 245; Duss 175: — non H. B. K. (Symb.). Wild cane.**St. Eustatius**: Inner slope of the Quill 350 M.: n. 423 B., 459 B.; Bottom of the Quill 300 M.: n. 911 B.; Quill: n. 78 H.; 2553, 2527, 925, 2529, 922, 926. 939.**Saba**: Boobyhill 300 M.: n. 1617 B.; Springbaygut: n. 2083 B., 2106 B., locality unknown: n. 159 L.

Guadeloupe, Martinique, St. Vincent, Trinidad. (Symb.).

1866. **Peperomia** Ruiz. et Pav.**Peperomia acuminata** C. DC. in Symb. Antill. III (1902) 242.*Peperomia glabella* Griseb. Flor. W. Ind. (1859) 165. (Symb.).*Peperomia glabella* var. *nigropunctata* Duss Fl. phan. (1897) 172. (Symb.).**St. Eustatius**: Top of the Quill 400 M.: n. 178 B., 317 B., near Maryglory 150 M.: n. 1160 B.; Missionhouse road: n. 551.**Saba**: Greathill 400 M.: n. 1382 B.; Paris n. 315, 321; John Ednyhill: n. 302; locality unknown: n. 103 L, 171 L.**St. Martin**: Nakedboyhill: n. 2725 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2898 B.

Antill., Columbia, Guiana. (Symb.).

Peperomia emarginella C. DC. Prodr. XVI, 1 (1869) 437.*Peperomia exilis* Griseb. Flor. W. Ind. (1859) 164. (Symb.); Duss 170.**Saba**: Mountain 800 M.: n. 1813 B; Mountain 6—800 M.: n. 19.

Cuba, Jamaica, Hispaniola, Portorico, St. Kitts, Guadeloupe, Martinique, St. Vincent, Trinidad. (Symb.).

Peperomia glabella A. Dietr. Sp. I (1831) 156; Gris. Fl. 165 p. p. (Symb.). *Orchidspurslane*.

St. Eustatius: Top of the Quill 400 M.: n. 180 B., 334 B.; slope of the Quill near Bengalen 150 M.: n. 679 B.; locality unknown: n. 1023 B., 193 G.; Quill: n. 646, 593, 620, 633, 1054; Boven: n. 1053.

Saba: Along roads between Bottom and Windwardside 200 M.: n. 1352 B.; Gainbygut: n. 1445a B.; Boobyhill 350 M.: n. 1612 B.; along the Forthgut 150—200 M.: n. 1625 B.; Crispeen 400 M.: n. 1723 B.; Springbaygut: n. 2105 B., 2110 B.; Windwardside: n. 2321 B.; Peperpot: n. 1052.

St. Martin: Mountain between Mont des Accords and Centry hill 200—300 M.: n. 2886 B.; top of the hill on the northern side of Cul de Sac 200—300 M.: n. 3008 B., Milldrumhill 300—400 M.: n. 3135 B.; Mount Paradis 150—400 M.: n. 3192 B., 3196 B., 3340 B.

I also saw *Peperomia glabella* on Nakedboyhill 100—200 M.

Antill., Costarica, Amer-austr. (Symb.).

Peperomia hernandifolia A. Dietr. Sp. I (1831) 157; Duss 172.

Saba: Mountain 800 M.: n. 1808 B.; Mountain 6—800 M.: n. 634, 140; locality unknown: n. 174 L.

Antill., Costarica, Venezuela. (Symb.).

Peperomia Langsdorffii Miq. Syst. (1843) 116.

St. Eustatius: Top of the Quill 380 M.: n. 278 B.; lower parts of the hills near Concordia: n. 974 B.

St. Martin: Nakedboyhill 200—265 M.: n. 2717 B., along roads near Colebayhill: n. 3041 B.; gut from Bethlehem to Cul de Sac 0—100 M.: n. 3414 B.; Milldrumhill 100—200 M.: n. 3117 B.; Colebayhill: n. 141.

St. Croix. (Symb.).

Haiti, St. Domingo, Portorico, St. Thomas, St. Jan, Tortola, Antigua. (Symb.).

Peperomia obtusifolia A. Dietr. Sp. I (1831) 154; Duss 173.

Peperomia magnoliaefolia A. Dietr. Sp. I (1831) 153; Duss 173. (Symb.).

St. Eustatius: Top of the Quill 400 M.: n. 242 B., 353 B.; Quill: n. 277, 293, 279, 278, 291, locality unknown: n. 290.

Saba: Mountain 600 M.: n. 1757 B.; Springbaygut: n. 2102 B.; Milldrumhill 300—400 M.: n. 3146 B.

St. Croix. (Symb.).

Antill., Amer. cont. trop., (Florida, Mexico, Venezuela.) (Symb.).

Peperomia petiolaris C. DC. in Seem. Journ. of Bot. IV (1866) 138. Wall purslane.

St. Eustatius: In a garden in Oranjestad: n. 63 B.; at the seashore near Tumbledown Dickbay: n. 155 B.; on walls in Oranjestad: n. 69 G.

Saba: Along roads between Bottom and Windwardside 200 M.: n. 1341 B.; lower part of the Bottom Mountain 250 M.: n. 1410 B.; Windwardside: n. 1893 B.; Bottom: n. 2245 B.; locality unknown: n. 39 L.

Cuba, Paraguay. (Symb.).

Peperomia scandens R. et Pav. Fl. Per I (1798) 32 t. 51 fig b.

Peperomia repens Kth. in H. B. K. Nov. gen. 1 (1815) 65; Gris. Fl. (1859) 165. (Symb.).

St. Eustatius: Bottom of the Quill 300 M.: n. 434 B.

Antill., Amer. cont. austral. (Symb.).

Peperomia urocarpa Fisch. et Mey. Ind. Sem. Hort. Petrop. IV (1837) 42.

Saba: Mountain 800 M.: n. 1825 B., 594, 647.

Antill., Amer. cont. trop. (Symb.).

81. Batidaceae.

2362. **Batis** Linn.

Batis maritima Linn. Syst. X ed. II (1759) 1380; Gris. Fl. 61; Duss 92; Millspaugh 486. Wild banana.

St. Martin: near Philipsburg: n. 2338 B.; road near Fort Amsterdam: n. 47.

I also saw *Batis maritima*: near Simonsbay village; Oysterpond; Green Key.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida, Baham., Antill., Amer. cont. trop. orient., Californ., Ins. Sandw. (Symb.).

63. Ulmaceae.

1898. **Celtis** Linn.

Celtis Iguanacea Sarg. Silva VII (1895) 64.

Celtis aculeata Sw. Prodr. (1788) 53; Gris. Fl. 149; Duss 152; Millspaugh 482. Snake y.

St. Eustatius: Top of the Quill 400 M.: n. 219 B.; Bottom of the Quill 300 M.: n. 895 B.; top of Signalhill 225 M.: n. 1029 B.; Red Cliff—Tumbledowndickbay: n. 229.

Saba: Bottom—Marypoint 300 M.: n. 1476 B.; Windwardside: n. 1950 B.; Springbaygut: n. 2116 B.

St. Martin: from Philipsburg to Guanabay: n. 2389 B., 2399 B.; from Philipsburg to Belvedere: n. 2572 B.; Nakedboyhill 50—200 M.: n. 2680 B.; between Belvedere and Oysterpond: n. 3079 B.; Mount Paradis 100—200 M.: n. 3293 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

1902. **Trema** Lour.

Trema Lamarckianum Bl. in Mus. Bot. Lugd. Bat. II (ca 1853) 58.

Sponia Lamarckiana Dene. in Nouv. Ann. Mus. Paris III (1834) 498; Gris. Fl. 150; Duss 153.

St. Eustatius: Gilboohill 150 M.: n. 1123 B.

Saba: Hellsgate—Marypoint: n. 2040 B.

Bermud., Baham., Antill. (Symb.).

Trema micranthum Bl. in Mus. Bot. Lugd. Bat. II (ca 1853) 58; Millspaugh 482 (?).

Sponia micrantha Dene. in Nouv. Ann. Mus. Paris III (1834) 498; Gris. Fl. 150; Duss 152.

St. Eustatius: Top of the Quill 400 M.: n. 186 B., 308 B.

Saba: Gainbygut: n. 292, 293; Laddergut: n. 130, 294.

St. Croix ex Millspaugh.

Florida austr., Antill., Amer. cont. trop. (Symb.).

64. **Moraceae.**

1931. **Dorstenia** Linn.

Dorstenia contrajerva Linn. Spec. (1753) 121; Gris. Fl. 154; Duss 159.

St. Martin: Mount Paradis 300—400 M.: n. 3344 B.

Antill., from Mexico to Peru. (Symb.).

1946. **Artocarpus** Linn.

Artocarpus incisa Linn. f. Suppl. (1781) 411.

Artocarpus communis Forst. Char. Gen. (1776) 100. Bread fruit tree.

Cultivated. [Ins. Pacif.] (Symb.).

1961. **Ficus** Linn.

Ficus Krugiana Warb. in Urb. Symb. Antill. III (1903) 487.

Ficus laurifolia Duss Fl. Ant. franç. (1897) 153. (haud Lam.). (Symb.).

St. Eustatius: Top of the Quill 400 M.: n. 207 B.; bottom of the Quill 300 M.: n. 907 B.

Saba: Bottom: n. 1875 B.

Guadeloupe, Dominica, Martinique, St. Vincent. (Symb.).

Ficus omphallophora Warb. in Urb. Symb. Antill. III (1903) 466.

Ficus pertusa Gris. (haud L. f.) Flor. W. I. (1859) 152 (p. p.); Duss 155.

Ficus pallida Gris. (haud Vahl) Fl. W. I. (1859) 151 (p. p.); (Symb.).

Saba: Top of the Mountain 800 M.: n. 1827a B.

St. Eustatius, St. Kitts, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada. (Symb.).

Ficus populnea Willd. Sp. Pl. IV (1806) 1141; Gris. Fl. 151; Millspaugh 483.

Ficus laevigata Vahl. Enum. II (1805) 183; Gris. Fl. 151; Millspaugh 483.

St. Eustatius: Slope of the lower part of Signalhill: n. 103 B.; top of the Quill: n. 207 B., 371 B., 378 B.; along roads in Oranjestad: n. 914 B.; along roads near Concordia: n. 1091 B.; near Boven 200 M.: n. 1247 B.; Quill: n. 732, 846, 883; top of the Quill: n. 135, 733; Cliff: n. 738, 788.

Saba: Bottom: n. 2233 B.; from Windwardside to Hellsgate: n. 1681 B.

St. Martin: in a garden in Philipsburg: n. 2552 B.; from Philipsburg to Belvedere: n. 2577 B.; Marigotroad near Marigot: n. 2820 B.; Lowlands: n. 3284 B.; Mount Paradis 100—200 M.: n. 3297 B.; 300—400 M.: n. 3324 B.; Mary's Fancy: n. 705; Centryhill: n. 888; Prince quarter: n. 838.

St. Croix ex Millspaugh.

Baham., Antill. (Symb.).

Ficus Urbaniana Warb. in Urb. Symb. Antill. III (1903) 459.

St. Eustatius: Bottom of the Quill 300 M.: n. 897 B.; Boven: n. 2331, 2371; Quill: n. 2322; locality unknown: n. 2395.

Saba: Gut between Hellsgate and Masticgut 300—400 M.: n. 2026 B.; Springbaygut: n. 2080 B.

St. Martin: Point Blanche (coll. George): n. 2495 B.; Mount Paradis 150—400 M.: n. 3200 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3423 B.; John Ednyhill: n. 2347.

St. Croix. (Symb.).

Guadeloupe, Martinique, St. Vincent. (Symb.).

1971. *Cecropia* Linn.

Cecropia peltata Linn. Syst. X. ed. II (1759) 1286; Gris. Fl. 153; Millspaugh 498. Wild papaw.

St. Eustatius: Quill: n. 254 B.; Bottom of the Quill 300 M.: n. 904 B., 955 B., 105 H.; Quill: n. 243, 219.

Saba: Mountain near Windwardside 600 M.: n. 2158 B. locality unknown: n. 192 L.

St. Croix ex Millspaugh.

Antill., Venezuela, Guiana. (Symb.).

65. Urticaceae.

1978. *Urera* Gaudich.

Urera caracasana Gris. Flor. W. I. (1859) 154; Duss 161.

Urera alceaefolia Gaudich. in Freyc. Voy. (1826) 479.

St. Eustatius: Bottom of the Quill 300 M.: n. 438 B., 903 B., 909 B., 130 H., 222 G.

Saba: Mountain 6—800 M.: n. 1949; Hellsgategut: n. 1950; locality unknown: n. 194 L.

*Antill., Amer. cont. trop. (Symb.).

1982. *Fleurya* Gaudich.

Fleurya aestuans Gaudich. in Freyc. Voy. (1826) 497; Gris. Fl. 154; Duss 160; Millspaugh 483. Stinging nettle.

St. Eustatius: Glassgut near Oranjestad: n. 47 B.; along roads in Oranjestad: n. 1079 B.; Oranjestad: n. 2832; locality unknown n. 135 G.

Saba: near the Sulphurmines: n. 1703 B.; locality unknown: n. 154 L.

I also saw *Fleurya aestuans*: Windwardside.

St. Martin: Milldrumhill in plantations 300 M.: n. 3132 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop., Afr. trop., Ind. Sundan. (Symb.).

1984. *Pilea* Lindl.

Pilea microphylla Liebm. in Vidensk. Selsk. Skr. Kopenh. V. Ser. Vol. II (1851) 286; Gris. Fl. 155; Duss 161.

Adiantum microphyllum O. Ktze. Rev. II (1891) 622; Millspaugh 483.

Pilea muscosa Lindl. Coll. Bot. (1821.) t. 4. Lace plant.

St. Eustatius: Signalhill 200 M.: n. 137 B.; near Maryglory 150 M.: n. 1156 B.; in a garden in Oranjestad: n. 1269 B.; Cultuurvlakte: n. 1059; Signalhill: n. 1037; Boven: n. 1058; locality unknown: n. 150 G., 301 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1330 B.; slope of the Mountain near St. Johns 250 M.: n. 1733 B.; Windwardside: n. 1900 B.; Laddergut 0—200 M.: n. 2020 B.; between Hellsgate and Mastiegut: n. 2072 B.; Mountain near Windwardside 600 M.: n. 2173 B.; Ladder: n. 1038; Greathill: n. 1039; Crespeen-Rendez-vous 400 M.: n. 1026; locality unknown: n. 142 L.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2900 B.; Mount Paradis 300—400 M.: n. 3235a B.

St. Croix ex Millspaugh.

Florida austr., Bermud., Baham., Antill., Amer. cont. trop. (Symb.).

Pilea obtusata Liebm. in Vidensk. Selsk. Skr. Kopenh. V Ser. Vol. II (1851) 300.

Pilea pubescens Gris. Flor. W. I. (1859) 159 (non Liebm.); (Symb.).

Saba: Top of the Mountain 800 M.: n. 1783a B., 1786a B., 2218a B.

Portorico, Montserrat, Martinique. (Symb.).

Pilea parietaria Bl. Mus. Bot. Lugd. Bat. II (1856) 48; Gris. Fl. 156.

Pilea ciliaris Wedd. in Ann. Sc. nat. III Sér. XVIII (1852) 209; Gris. Fl. 156.

Saba: Top of the Mountain 800 M.: n. 1800 B.

Antill., Guatemala, Venezuela. (Symb.).

Pilea semidentata Wedd. in Ann. Sc. nat. III Sér. XVIII (1852) 215; Gris. Fl. 157.

Pilea elegans Wedd. in Ann. Sc. nat. III Sér. XVIII (1852) 215; Gris. Fl. 157.

St. Eustatius: Top of the Quill 400 M.: n. 175 B., 340 B., 351 B., 368 B.; slope of the Quill near Bengalen 150 M.: n. 685 B.; Quill: n. 244, 1102, 192 G., 273 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1338 B.; from Marypoint to Bottom: n. 2063 B.; Gainbygut: n. 1190.

Hispaniola, Portorico, St. Thomas, Guadeloupe, Dominica, Martinique. (Symb.).

1990. **Boehmeria** Jacq.

Boehmeria ramiflora Jacq. Enum. Pl. carib. (1760) 31; Gris. Fl. 160; Duss 163.

St. Eustatius: Bottom of the Quill 300 M.: n. 447 B.; Bengalen: n. 1152, 1172, 1202, 1222, 1242; Quill: n. 1192, 1225, 1233.

Saba: Mountain 800 M.: n. 2219 B.

Guadeloupe, Dominica, Martinique, St. Vincent, Grenada, Barbados. (Herb. Krug et Urban.).

2009. **Rousselia** Gaudich.

Rousselia humilis Urb. in Symb. Antill. V (1905) 205.

Rousselia lupulacea Gaudich. in Bot. Freye. Voy. (1826) 503; Gris. Fl. 160.

Saba: from Hellsgate to the Sulphurmines: n. 1669a B. Baham., Antill. (Symb.).

67. Loranthaceae.

2089. **Phoradendron** Nutt.

Phoradendron trinervium Gris Flor. W. I. (1860) 314; Duss 328.

St. Eustatius: Signalhill 200 M.: n. 135 B., 1048 B.; slope of the Quill near Bengalen 100 M.: n. 672 B.; gut near Fort de Wind 50 M.: n. 843 B.; Cultuurvlakte: n. 26; Red Cliff: n. 14; White Wall: n. 13; locality unknown: n. 209 G.

Saba: Paris 400 M.: n. 1371 B.; Saddle: n. 1660 B.; Paris 2—400 M.: n. 27.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2888 B.; hills on the eastern side of Cul de Sac: n. 3443 B.

Antill., Costarica. (Symb.).

72. Olacaceae.

2129. **Schoepfia** Schreb.

Schoepfia Schreberi J. F. Gmel. Syst. (1791). 376.

St. Eustatius: Top of the Quill 400 M.: n. 201 B., 364 B., 375 B., 377 B.; inner slope of the Quill 350 M.: n. 1226 B.

Saba: Bottom 200 M.: n. 1447 B.; Springbaygut: n. 2085 B., 2125 B.; Wellgut: n. 31.

Antill., Mexico? San Salvador? Venezuela. (Symb.).

74. Aristolochiaceae.

2174. **Aristolochia** Linn.

Aristolochia odoratissima Linn. Spec. (1763) 1362.

Cultivated. [Jamaica.] (Ind. Kew.).

Aristolochia trilobata Linn. Spec. (1753) 960; Gris. Fl. 299; Duss 314; Millspaugh 484.

St. Martin: Mount Paradis 300—400 M.: n. 3337 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. cent. and austr. trop. (Symb.).

77. Polygonaceae.

2209. **Coccoloba** Linn.

Coccoloba diversifolia Jacq. Enum. Syst. (1760) 19; Millspaugh 484.

Coccoloba punctata Gris. (non L.) Flor. W. I. (1859) 163.

Coccoloba barbadiensis Jacq. Enum. Syst. (1760) 37; Duss 166.

Coccoloba diversifolia Gris. (non Jacq.) pr. p. Flor. W. I. (1859) 163 (Symb.). Red wood.

St. Eustatius: Top of the Quill 400 M.: n. 179 B., 210 B., 312 B.; slope of the Quill on the side of White Wall 200 M.: n. 1172 B.; Boven: n. 37; Little Mountain—Red Cliff: n. 36, 38, 156; top of the Quill: n. 28; Gilboohill: n. 25.

Saba: lower part of the Bottom Mountain 250 M.: n. 1418 B.; Windwardside—Hellsgate: n. 1665 B.; St. Giles: n. 1860 B.; Mountain near Windwardside 600 M.: n. 2175 B.; Paris—Greathill: n. 29, 30, 31; Peperpot: n. 24; locality unknown: n. 340 L.

St. Croix ex Millspaugh.

Baham., Antill., Curaçao. (Symb.).

Coccoloba Krugii Lindau Mon. Cocc. (1892) 145.

St. Martin: near Mulletpond: n. 2756 B.

Anguilla Boldingh.

Baham., Portorico, Barbuda. (Symb.).

Coccoloba nivea Jacq. Enum. Syst. (1760) 19; Gris. Fl. 163; Millspaugh 484.

Coccoloba excoriata Linn. Spec. (1762) 524; Duss 168. Sugar-grape.

St. Eustatius: Quill: n. 142; from Signalhill to Venusbay: n. 964 B.

Saba: Boobyhill 300 M.: n. 1582 B.; Windwardside—Hells-gate: n. 1688 B.; Springbaygut: n. 2087 B.; Windwardside: n. 2316 B.; Bottom: n. 187, 188, 189; locality unknown: n. 101 L.

St. Martin: from Philipsburg to Belvedere: n. 2579 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2961a B.; top of the hill at the northern side of Cul de Sac 200—300 M.: n. 2993 B.; Mount Paradis 300—400 M.: n. 3228 B., 3319 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3420 B.; John Ednyhill: n. 3.

I also saw *Coccoloba nivea*: along roads near Grande Case.

St. Croix ex Millspaugh.

Jamaica, Hispaniola, Portorico, St. Thomas, St. Barthélemy, Antigua, Guadeloupe, Martinique, Barbados (cult), Grenada, Tobago, Trinidad. (Symb.).

Coccoloba obtusifolia Jacq. Enum. Syst. 19.

Coccoloba punctata Griseb. (non L.) pr. p. Flor. W. I. (1859) 163.

Coccoloba punctata Griseb. γ *microstachya* in Flor. W. I. (1859) 163; (Symb.).

Saba: Boobyhill: n. 1569 B., 1571 B.

Baham., Antill. (Symb.).

Coccoloba uvifera Jacq. Enum. Syst. 19; Gris. Fl. 161; Duss 165; Millspaugh 484. Sea grape.

St. Eustatius: Seashore near Oranjestad: n. 17 B.; top of the Quill 400 M.: n. 263 B.; slope of the lower part of the Quill near Bengalen: n. 717 B.; along roads in Oranjestad: n. 1057 B.; near White Wall: n. 1257 B.; Concordiabay: n. 183; Venusbay: 185; locality unknown: n. 120 H., 198 G.

I also saw *Coccoloba uvifera*: slope of the Quill on the side of White Wall.

Saba: Paris 400 M.: n. 1370 B.; Bottom: n. 1988 B.; Ladder: n. 186; locality unknown: n. 360 L.

I also saw *Coccoloba uvifera*: Green Island.

St. Martin: Guanabay: n. 2432 B.; at the seashore near Simonsbay village: n. 3032 B.

I also saw *Coccoloba uvifera* in Lowlands; Marigothill to Marigot.

St. Croix ex Millspaugh.

Bermud., Florida, Baham., Antill., Mexico, Amer. centr., Venezuela, Guiana. (Symb.).

2205. **Antigonon** Endl.

Antigonon cinerascens Mart. et Gal. in Bull. Ac. Brux. X. 14. Cultivated. [from Mexico to Guatemala.] (Symb.).

Antigonon leptopus Hook. et Arn. Bot. Beech. (1841) 308 t. 69; Gris. Fl. 164; Duss 612. Coralita.

St. Eustatius: in a garden in Oranjestad: n. 1071 B., 18 H., 21 G.

Saba: Windwardside: n. 1931 B.; Bottom: n. 2242 B.; locality unknown: n. 46 L., 299 L.

St. Martin: near Bethlehem: n. 2670 B.

Jamaica. (Gris. Fl.) Cultivated and escaped.

78. **Chenopodiaceae.**

2223. **Chenopodium** Linn.

Chenopodium ambrosioides Linn. Spec. (1753) 219; Gris. Fl. 60; Duss 51; Millspaugh 485. Worm bush.

St. Eustatius: along roads in Oranjestad: n. 938 B.; in a garden in Oranjestad: n. 1085 B.; locality unknown: n. 5 G., 5 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1921 B., 1301 B.; Bottom: n. 1846 B.; Laddergut 0—200 M.: n. 1955 B.; seashore near Forthbaylanding: n. 2149 B.; Paris: n. 223; Bottom: n. 224; locality unknown: n. 235 L.

St. Martin: along roads from Marigot to Grande Case: n. 3355 B.

St. Croix ex Millspaugh.

Trop. and warmer countries of both hemisph. (Symb.).

Chenopodium murale Linn. Spec. (1753) 219; Gris. Fl. 60; Duss 51; Millspaugh 485. Sprain bush.

St. Eustatius: along roads in Oranjestad: n. 578 B.; locality unknown: n. 9 G., 9 H.; seashore: n. 244; Village: n. 243.

St. Martin: from Philipsburg to Guanabay: n. 2344a B.

St. Croix ex Millspaugh.

Baham., Antill., indig. in the Old world. (Symb.).

2257. **Salicornia** Linn.

Salicornia herbacea Linn. Spec. (1762) 5.

St. Martin: near Mulletpond: n. 2749 B.

Anguilla Boldingh.

Amer. sept., Europa, Afr., Asia. (Symb.).

79. Amarantaceae.2292. **Celosia** Linn.**Celosia argentea** Linn. Spec. (1753) 205.

Cultivated. [Antill., Amer., Afr., Asia trop.] (Symb.).

Celosia nitida Vahl Symb. II (1791) 44; Gris. Fl. 62; Duss 53; Millspaugh 485.**St. Martin:** near Bethlehem: n. 2666 B.; Marigotroad: n. 2803a B.

Florida austral., Antill., Mexico, Calif. peninsula, Amer. centr. (Symb.).

2299. **Amarantus** Linn.**Amarantus gracilis** Desf. Tabl. I ed. (1804) 43.*Euxolus caudatus* Moq. in DC. Prodr. XIII. II (1849) 247; Gris. Fl. 68; Millspaugh 486.*Amarantus caudatus* Linn. Spec. (1753) 990.**St. Eustatius:** in a garden: n. 547.**St. Martin.** (Symb.).**St. Croix** ex Millspaugh.

Baham., Antill., Bonaire, Curaçao, trop. countries of both hemisph. (Symb.).

Amarantus paniculatus Linn. Spec. (1762) 1406. Princeplum. Cultivated. [Baham., Guiana., trop. countries of both hemisph.] (Gris. Fl.).**Amarantus polygonoides** Linn. Pl. Jam. Pugil. II. (1759) 27; Millspaugh 486.*Amblogyne polygonoides* Raf. Fl. Tellur III (1836) 42; Gris. Fl. 68; Duss 58. White polly.**St. Eustatius:** near Maryglory 150 M.: n. 1135 B.; locality unknown: n. 19 G.**Saba:** Bottom: n. 559.**St. Martin:** from Philipsburg to Guanabay: n. 2406 B.; in an orchard near Bellevue: n. 2839 B.; between Mount Vernon and Orientbay: n. 2931 B., 3360 B.; Tintamarre: n. 3384a B., 3393 B.; near Philipsburg: n. 595.**St. Croix** ex Millspaugh.

Florida, Key, Baham., Antill., Aruba, Curaçao. (Symb.).

Amarantus spinosus Linn. Spec. (1753) 991; Gris. Fl. 68; Duss 58; Millspaugh 486.**St. Eustatius:** along roads in Oranjestad: n. 1066 B.; locality unknown: n. 18 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1323 B.; Bottom 200 M.: n. 1454 B.; Windwardside: n. 1918 B., 2327a B.; Bottom: n. 568, 569; locality unknown: n. 252 L.

St. Martin: Mount Paradis 100—200 M.: n. 3295 B.

St. Croix ex Millspaugh.

Bermud., Baham., Amer. cont., Afr. trop., Asia trop. to Japon. and N. Guinea. (Symb.).

Amarantus tristis Linn. Spec. (1753) 989; Gris. Fl. 69; Duss 59; Millspaugh 486. Calaloe.

Amarantus gangeticus Linn. Syst. ed. X. (1759) 1268.

St. Eustatius: in a garden in Oranjestad: n. 84 B.; along roads in Oranjestad: n. 551 B.; Cultuurvlakte: n. 576; in a garden: n. 575; locality unknown: n. 16 H., 17 G.

Saba: between Bottom and Marypoint 100—200 M.: n. 1462 B.; Windwardside: n. 1919 B.; Rendez-vous 400 M.: n. 580; Bottom: n. 577; locality unknown: n. 33 L.

St. Martin: between Mont Chambord and the Saline: n. 2923 B.; lower part of the Milldrumhill: n. 3163 B.; Marigotroad: n. 552.

St. Croix ex Millspaugh.

Antill., Chile. (Symb.).

2328. **Achyranthes** Linn.

Achyranthes obtusifolia Lam. Enc. I (1783) 545.

Achyranthes aspera Linn. var. *obtusifolia* Gris. Flor. W. I. (1860) 220; Millspaugh 485. Man better man.

St. Eustatius: in a garden in Oranjestad: n. 82 B.; along roads in Oranjestad: n. 1067 B.; lower part of the Quill: n. 515; Cliff at the seashore: n. 514; English quarter: n. 513; along roads in Oranjestad: n. 560 B.; locality unknown: n. 40a B., 93 G., 176 H.

I also saw *Achyranthes obtusifolia* on the seashore near Oranjestad; slope of the Cliff.

Saba: Windwardside—Hellsgate: n. 1671 B.; Windwardside: n. 1885 B.; locality unknown: n. 254 L.

I also saw *Achyranthes obtusifolia*: Laddergut.

St. Martin: from Filipsburg to Belvedere: n. 2583 B.; Marigotroad: n. 516.

I also saw *Achyranthes obtusifolia* in Cul de Sac; Marigothill—Marigot: in pastureground near Bellevue.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Afr., Asia-trop., Ins-Paul. (Symb.).

2335. **Alternanthera** Forsk.**Alternanthera crucis** Comb. Nov.

Telanthera crucis Moq. in DC. Prod. XIII. II. (1849.) 362 ;
Gris. Fl. 67.

St. Martin : Pointe Blanche : n. 3407 B.

St. Croix. (Herb. Krug et Urban.).

Alternanthera repens O. Ktze Rev. II (1891) 540.

Alternanthera achyrantha R. Br. Prodr. I (1810) 417 ; Gris.
Fl. 67 ; Duss 57 ; Millspaugh 485.

St. Eustatius : in a garden in Oranjestad : n. 83 B. ; Cultuur-
vlakte : n. 522 ; in a garden : n. 521 ; locality unknown : n. 520.

Saba : between Bottom and Marypoint 100—200 M. : n. 1458 B. ;
Windwardside : n. 1892 B., 1897 B. ; Bottom : n. 524, 525.

St. Martin : along roads from Colebay to Simonsbay : n.
3246 B. ; from Philipsburg to Point Blanche : n. 3407 B. ; gut from
Bethlehem to Cul de Sac 100—200 M. : n. 3421 B. ; Colebayhill :
n. 523.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont., Ins. Canar., Hispania. (Symb.).

2338. **Gomphrena** Linn.**Gomphrena globosa** Linn. Spec. (1753) 224. Bachelor's button.

Cultivated. [Indig. Asia orient. et austr. from Ins. Sundaic. to
Nova Guinea.]. (Symb.).

2339. **Iresine** Linn.

Iresine elatior L. Cl. Rich. ap. Willd. Spec. IV (1806) 766 ; Gris.
Fl. 64 ; Duss 55 ; Millspaugh 485. White snow plant.

St. Eustatius : Cultuurvlakte : n. 591 ; road near Sugarloaf :
n. 594 ; locality unknown : n. 1118 B.

Saba : Peperpot : n. 592.

St. Martin : Colebayhill : n. 593 ; from Philipsburg to Guanabay :
n. 2396 B. ; Nakedboyhill 100 M. : n. 2692 B.

St. Croix ex Millspaugh.

Antill., Venezuela, Nova Granata. (Symb.).

Iresine paniculata O. Ktze. Rev. II (1891) 542.

Iresine celosioides Linn. spec. (1763) 1456 ; Gris. Fl. 64 ;
Duss 55.

St. Eustatius: from Signalhill to Tumbledowndickbay 200 M.: n. 125 B.; deserted plantations near Tumbledowndickbay: n. 151 B.; near Maryglory 150 M.: n. 1164 B.; slope of the Quill: n. 585; Boven: n. 586.

Saba: Bottom: n. 1455 B.; Boobyhill 300 M.: n. 1577 B.; Hellsgate: n. 1693 B.; Ladder: n. 590; Greathill 2—400 M.: n. 589; Rendez-vous: n. 588; locality unknown: n. 146 L.

St. Martin: Marigotroad near Marigot: n. 2817 B.; lower part of the Milldrimhill: n. 3176 B.; Centryhill: n. 587.

Baham., Antill., Amer. cont. from Un. Stat. to Argentina. (Symb.).

Philoxerus R. Br.

cf. 2339. Iresine Linn. sect. I.

Philoxerus vermicularis R. Br. Prodr. I (1819) 410; Gris. Fl. 65; Duss 56.

Lithophila vermiculata Uline in Field Col. Mus. Bot. II (1900) 39; Millspaugh. 485.

St. Martin: Oysterpond: n. 9; locality unknown: n. 2523 B.

St. Croix ex Millspaugh.

Florida austr., Key ins., Baham., Antill., Amer. cont. trop., Afr. occid. (Symb.).

Lithophila Sw.

Lithophila muscoides Sw. Prod. Veg. Ind. Occ. (1788) 14; Gris. Fl. 66; Duss 56; Millspaugh 485.

Alternanthera muscoides Benth et Hook. f. Gen. III (1880) 39.

St. Eustatius: Seashore near Oranjestad: n. 20 B.; deserted plantations near Tumbledowndickbay: n. 167 B.; Concordiabay: n. 534; seashore: n. 533, 586a B.; Cliff near Fort Amsterdam: n. 532; locality unknown: n. 408a B.

Saba: along the seashore near Forthbaylanding: n. 1645 B.; Laddergut: n. 1977 B.; seashore near Flat Point: n. 2136 B.; Ladder: n. 530, 531.

St. Martin: from Philipsburg to Guanabay: n. 2376 B.; along the saltpond near Philipsburg: n. 2449 B.; Green Key: n. 3368 B.; Philipsburg—Fort Amsterdam: n. 536; Guanabay: n. 535.

St. Croix ex Millspaugh.

Baham., Antill., Bonaire, Curaçao, Aruba. (Symb.).

80. Nyctaginaceae.

2347. *Mirabilis* Linn.

Mirabilis jalapa Linn. Spec. (1753) 177; Gris. Fl. 69; Duss 59; Millspaugh 486. Four o'clock.

St. Eustatius: in a garden in Oranjestad: n. 1253 B., 88, 153 H., 154 H., 239 G., 240 G., 247 G.

Saba: Lowerpart of the Bottom Mountain 250 M.: n. 1417 B.; Peperpot: n. 34, 87; Bottom: n. 33; locality unknown: n. 112 L.

St. Martin. (Symb.).

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop. (Symb.). Cultivated and escaped.

2349. *Boerhaavea* Linn.

Boerhaavea erecta Linn. Spec. (1753) 3; Gris. Fl. 69; Duss 60; Millspaugh 486.

St. Eustatius: Oranjestad: n. 25, 26, 27; locality unknown: n. 601 B., 86 G., 223 H., 223 G.

St. Martin: from Philipsburg to Guanabay: n. 2405 B.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Amer. cont. (Symb.).

Boerhaavea hirsuta Willd. Phytogr. (1794) 1; Gris. Fl. 69; Duss 60. Potta potta, Hog meat.

Boerhavia repens Linn. Spec. (1753) 3.

St. Eustatius: Seashore near Oranjestad: n. 40 B.; in a garden in Oranjestad: n. 85 B.; Cultuurvlakte: n. 20, 21; locality unknown: n. 87 G., 175 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1508 B.; Laddergut 200 M.: n. 1995 B., 22.

St. Martin; along roads from Grande Case to Marigot: n. 3098 B.

St. Croix (Symb.).

Antill., Amer. sept. and centr. (Symb.).

Boerhaavea paniculata L. Cl. Rich. in Act. Hist. Nat. Par. 1 (1792) 105; Gris. Fl. 69; Duss 60; Millspaugh 487.

St. Eustatius: in a garden in Oranjestad: n. 524 B.; Bengalen: n. 632 B.; along roads near Glassbottle: n. 872 B.; Fairplay: n. 61; Oranjestad: n. 3; locality unknown: n. 1011 B., 1019 B., 62, 64.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1347 B.; along roads 200 M.: n. 1384 B.; between Bottom and

Marypoint 100—200 M.: n. 1534 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2056 B.; locality unknown: n. 31 L.

St. Martin: Fort Willem 0—50 M.: n. 2537 B.; Fort Amsterdam: n. 39; road to Guanabay: n. 4; locality unknown: n. 2305 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. centr. and Austral. (Symb.).

Boerhaavea scandens Linn. Spec. (1753) 3; Gris. Fl. 69.

St. Martin: Colebay (Coll. George): n. 2516 B.; near Fort Amsterdam: n. 10.

Anguilla Boldingh.

Baham., Antill., Curaçao, Amer. cont. from Arizona to Peru. (Symb.).

2354. **Pisonia** Linn.

Pisonia aculeata Linn. Spec. (1753) 1026; Gris. Fl. 70; Duss 61; Millspaugh 487. Cockspur.

St. Eustatius: from Little Mountains to Vennsbay: n. 957 B.; along roads near Oranjestad: n. 1002 B.; slope of the hills near Concoridiabay: n. 89; Venusbay: n. 35.

Saba: lower part of the Bottom Mountain 300 M.: n. 1425 B.; Boobyhill: n. 1610a B.; Rendez-vous: n. 90; Quarter 300 M.: n. 36; locality unknown: n. 148 L.

I also saw *Pisonia aculeata*: Springbaygut; Flat Point.

St. Martin: Princequarter (Coll. George): n. 2467 B.; from Philipsburg to Belvedere: n. 2556 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2873 B.; Colebayhill: n. 32.

I also saw *Pisonia aculeata* in: Lower part of Milldrumhill; Mount Paradis: along roads near Nakedboyhill.

St. Croix ex Millspaugh.

Florida austral., Baham., Antill., trop. contr. of both hemisph. (excl. Afr. cont.) (Symb.).

Pisonia fragrans Dumont Cours. Bot. Cult. II ed. vol. VII (1814) 114.

Pisonia obtusata Sw. Flor. III (1806) 1960 (non Jacq. 1798); Gris. Fl. 71 (p. p.) (Symb.); Duss 62.

Pisonia inermis Gris. Flor. W. I. (1859) 7 (p. p., non Jacq.) (Symb.); Millspaugh 487. Black Loblobly tree.

St. Eustatius: along the road from Oranjestad to White Wall: n. 4 B.; top of the Quill 400 M.: n. 337 B.; 450 M.: n. 357 B.,

362 B.; inner slope of the Quill 350 M.: 425 B.; top of the Quill 500 M.: n. 478 B., 489 B.; slope of the Quill near Bengalen: n. 707 B.; 200 M.: n. 738 B.; from Signalhill to Venus bay: n. 959a B.; near Boven 200 M.: n. 1236a B.; Cultuurvlakte—Pango: n. 9; Cultuurvlakte: n. 20; locality unknown: n. 180 G., 17.

Saba: Marypoint: n. 37.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2649 B.; Nakedboyhill 50—200 M.: n. 2704 B.; lower part of the hills between Mount Chambord and Mount Vernon: n. 2918 B., 2921 B.; between Mount Vernon and Orientbay: n. 2927 B.; along roads from Grande Case to Marigot: n. 3091 B.; Milldrumhill 300—400 M.: n. 3186 B.; Mount Paradis 200—300 M.: n. 3312 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3418 B.; near Fort Amsterdam: n. 1; John Ednyhill: n. 23.

I also saw *Pisonia fragrans* on the hill between Mont des Accords and Centryhill; near Saline.

St. Croix ex Millspaugh.

Florida austr., Antill., Amer. centr., Columbia, Guiana. (Symb.).

Pisonia subcordata Sw. Prodr. (1788) 60; Gris. Fl. 70; Duss 61; Millspaugh 487. *Ma p p o o*.

St. Eustatius: Seashore near Oranjestad: n. 25 B.; Top of the Quill 400 M.: n. 185 B., 370 B.; in deserted plantations at the seashore near Bengalen: n. 596 B.; slope of the lower part of the Quill near Bengalen: n. 714 B.; locality unknown: n. 181 G., 322 H.

I also saw *Pisonia subcordata*: in Signalhill 200 M.; top of the Cliff between Oranjestad and White Wall; gut near Fort de Wind; Tumbledown-diekbay—Venusbay; Bergje.

Saba: Between Bottom and Marypoint 350 M.: n. 1498 B.

I also saw *Pisonia subcordata*: Forthbaygut; Tentgut; Saddle; Crispeenhill; Bottom—Windwardside; Gilesgut; Springbaygut; Flatpoint.

St. Martin: from Filipsburg to Guanabay: n. 2373 B.; Prince quarter (Coll. George.): n. 2471 B.

I also saw *Pisonia subcordata*: along roads near Bethlehem; along roads from Filipsburg to Nakedboyhill; Marigothill—Marigot; Nakedboyhill; Fort Willem; hills on the northern side of Cul de Sac; along roads from Filipsburg to Simonsbay; along roads near Grande Case; Vallée des Pères; lower part of Milldrumhill; Mount Paradis 300—400 M.; Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Portorico, St. Thomas, St. Barthélemy. St. Kitts, Antigua, Guadeloupe, Désirade, Martinique. (Symb.).

83. Phytolaccaceae.

2370. **Rivina** Linn.

Rivina humilis Linn. Spec. (1753) 121; Millspaugh 486.

Rivina laevis Linn. Mant. I (1767) 41; Gris. Fl. 59; Duss 49. J u m b y p e p p e r.

St. Eustatius: In a deserted plantation near Tumbledown Dickbay: n. 170 B.; Bengalen: n. 611 B., 636 B.; slope of the Quill near Bengalen 50—200 M.: n. 646 B.; along roads in Oranjestad: n. 939 B.; Village: n. 176; locality unknown: n. 161 G., 211 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1339 B.; along roads 200—300 M.: n. 1713 B.; Laddergut 200 M.: n. 1993 B.; Peperpot: n. 173; Bottom: n. 174, 175; locality unknown: n. 32 L.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2904 B.; along roads in Cul de Sac: n. 2997 B.

I also saw *Rivina humilis* in Mount Paradis 100—200 M.; gut from Bethlehem to Cul de Sac.

St. Croix ex Millspaugh.

Florida austr., Baham., Antill., Amer. cont. trop. et subtrop. (Symb.).

Villamilla R. et P.

cf. 2370. *Rivina* Linn. Sect. 2.

Villamilla octandra Hook f. in Benth. et Hook. Gen. Plant III (1880) 81; Millspaugh 486.

Rivina octandra Linn. Cent. plant. II (1756) 9; Gris. Fl. 59; Duss 50. White hoop.

St. Eustatius: deserted plantations near Tumbledown Dickbay: n. 169 B.; slope of the hills near Concordiabay: n. 191, 192, 194; Red Cliff—Tumbledown Dickbay: n. 193; locality unknown: n. 323 H.

Saba: Top of the Cliff near the seaside of Boobyhill: n. 1610 B.; Hellsgate: n. 1698 B.; gut between Hellsgate and Masticgut 300—400 M.: n. 2053 B.; Crispeen Rendez-vous: n. 127; Hellsgate n. 177, 178, 179; locality unknown: n. 236 L.

St. Martin: Cul de Sac along roads: n. 2966 B.; Centryhill: n. 126; French frontier: n. 190.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

2371. **Petiveria** Linn.

Petiveria alliacea Linn. Spec. (1753) 342; Gris. Fl. 59; Duss 50; Millspaugh 486.

St. Eustatius: near Tumbledown Dickbay: n. 139 B.; locality unknown: n. 238 H., 260 G.

Saba: Windwardside: n. 1591 B.; Laddergut: n. 2001 B.; Peperpot: n. 152, 153; Rendez-vous 400 M.: n. 151; Bottom: n. 150; locality unknown: n. 72 L., 328 L.

St. Martin: Marigot Hill: n. 2615 B.; gut from Bethlehem to Cul de Sac: n. 3422 B.

St. Croix ex Millspaugh.

Florida austr., Antill., Amer. cont. trop. (Symb.).

2372. **Microtea** Sw.

Microtea debilis Sw. Prod. Veg. Ind. Occ. (1788). 53; Gris. Fl. 59; Duss 49; Millspaugh 486.

St. Eustatius: in a garden in Oranjestad: n. 89 B.; along roads in Oranjestad: n. 569 B., 1086 B.; Oranjestad: n. 195; locality unknown: n. 43 G., 22.

St. Martin: From Philipsburg to Belvedere: n. 2588 B.

St. Croix ex Millspaugh.

Antill., Panama to Peru, Bras. equat. and Guiana. (Gris. Fl.).

2380. **Phytolacca** Linn.

Phytolacca icosandra Linn. Syst. X. ed. II (1759) 1040; Gris. Fl. 58; Duss 49.

Saba: Mountain 800 M.: n. 2214 B.; locality unknown: n. 169 L., 175 L., 209 L.

St. Martin: Mount Paradis 300—400 M.: n. 3211 B.

Baham., Antill., Amer. cont. trop. (Symb.).

84. **Aizoaceae.**2387. **Mollugo** Linn.

Mollugo verticillata Linn. Spec. (1753) 89; Gris. Fl. p. 56.

St. Eustatius: along roads in Oranjestad: n. 1098 B.; slope of the Quill near Bengalen: n. 196.

Antill., Amer. cont. (Symb.).

2394. **Sesuvium** Linn.

Sesuvium portulacastrum Linn. Syst. X. ed. II (1759) 1058; Gris. Fl. 57; Duss 47; Millspaugh 487.

St. Eustatius: Seashore near Bengalen: n. 585 B.; near White Wall: n. 851 B., 855 B.; near Maryglory 150 M.: n. 1162 B.; seashore near Turtlebay: n. 93, 94.

Saba: near the Sulphurmines: n. 1703a B.; seashore near Laddergut: n. 2132 B.

St. Martin: near Filipsburg: n. 2336a B.; near Mulletpond: n. 2763a B.; Tintamarre: n. 3389 B.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Curaçao, trop. and subtr. countries of both hemisph. (Symb.).

2395. *Trianthema* Linn.

Trianthema portulacastrum Linn. Spec. (1753) 223.

Trianthema monogynum Linn. Mant. I (1767) 69; Gris. Fl. 58; Duss 48; Millspaugh 487.

St. Eustatius: in a garden in Oranjestad: n. 1058a B., 104, 90 G.

St. Martin: Nakedboyhill: n. 2707a B.; road near Simonsbay: n. 3247a B.; Colebayhill: n. 103.

St. Croix ex Millspaugh.

Key, Antill., Curaçao, Amer. cont. trop., Afr., and Asia trop. (Symb.).

85. Portulacaceae.

2406. *Talinum* Adans.

Talinum paniculatum Gärtn. Fruct. II (1791) 219 t. 128.

Talinum patens Willd. Spec. II (1800) 863 (var *z*) (Symb.); Gris. Fl. 57; Duss 46; Millspaugh 487.

Saba: Laddergut 0—200 M.: n. 1952 B.; Peperpot: n. 40; locality unknown: n. 124 L.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.).

Talinum triangulare Willd. Spec. II (1800) 862; Gris. Fl. 56; Duss 46; Millspaugh 487.

St. Eustatius: Bengalen: n. 612 B.; along roads in Oranjestad: n. 1132 B.; near Boven 200 M.: n. 1241 B.; locality unknown: n. 63 G.

Saba: near the Sulphurmines: n. 1701 B.; Mountain 400 M.: n. 1768 B.; Bottom: n. 2232 B.; locality unknown: n. 256 L.

St. Martin: from Filipsburg to Belvedere: n. 2569 B.; from Belvedere to Oysterpond: n. 3044 B.; Prince quarter: n. 42; Cul de Sac: n. 41; locality unknown: n. 3613 B.

St. Croix ex Millspaugh.

Antill., Amer. austr. (Symb.).

2421. **Portulaca** Linn.

Portulaca halimoides Linn. Spec. (1762) 639; Gris. Fl. 57; Duss 47; Millspaugh 488. Silk cotton purslane.

St. Eustatius: in a garden in Oranjestad: n. 86 B.; along roads in Oranjestad: n. 549 B, 1080 B.; seashore near White Wall: n. 853 B.; old Fort: n. 67; White Wall: n. 66; locality unknown: n. 339 H.

St. Martin: Colebay (coll. George): n. 2522 B.; between Mount Vernon and Orientbay: n. 2943 B.; Green Key: n. 3362 B.; Fort Amsterdam: n. 57; Guanabay: n. 58.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Mexico. (Symb.).

Portulaca oleracea Linn. Spec. (1753) 445; Gris. Fl. 57; Duss 46; Millspaugh 487.

St. Eustatius: Seashore near Oranjestad: n. 41 B.; in a garden in Oranjestad: n. 88 B., 521 B.; Bengalen: n. 635 B.; slope of the lower part of the Quill near Bengalen: n. 722 B.; seashore near Venusbay: n. 966 B.; near Boven: n. 200 M.; seashore: n. 72, 73; locality unknown: n. 28 G., 29 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1329 B.; Windwardside—Hellsgate: n. 1664 B.; near Flat Point: n. 2130 B.; Windwardside: n. 2320 B.; locality unknown: n. 80 L.

St. Martin: from Philipsburg to Guanabay: n. 2413 B.; along roads near Prince quarter: n. 2912 B.; between Mount Vernon and Orientbay: n. 2940 B.; from Belvedere to Oysterpond: n. 3052 B.; Tintamarre: n. 3387 B.; locality unknown: n. 68.

I also saw *Portulaca oleracea*: in pastureground near Belvedere; Naked-boyhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., trop. and warmer countries of both hemisph. (Symb.).

Portulaca pilosa Linn. Spec. (1753) 445; Gris. Fl. 57; Duss 47; Millspaugh 487.

Saba: along roads 200 M.: n. 1383 B.; Bottom: n. 2264 B., 2299 B.

St. Martin: I also saw *Portulaca pilosa*: along roads near Bethlehem.

St. Croix ex Millspaugh.

Florida, Antill., Mexico. (Symb.). Also cultivated.

Portulaca poliosperma Urb. in Symbolae Antillanae IV (1905) 232.

St. Martin: Lowlands: n. 3286a B.

Portorico. (Symb.).

Portulaca quadrifida Linn. Mant. I (1767) 73.

St. Eustatius: along roads in Oranjestad: n. 559 B.; locality unknown: n. 1022 B.

St. Martin: along roads from Bethlehem to Mont Chambord: n. 2917 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2917 B.; Marigotroad: n. 79.

Trop. countries of the Old world. (Ind. Kew.).

86. Basellaceae.

2427. **Boussingaultia** H. B. K.

Boussingaultia baselloides H. B. K. Nov. Gen. et Spec. VII (1825) 196 t. 645.

Cultivated. [Amer. trop.] (E. & P.).

87. Caryophyllaceae.

2452. **Drymaria** Willd.

Drymaria cordata Willd. ap. R. et Sch. Syst. V (1819) 406; Gris. Fl. 56; Duss 45; Millspaugh 468. White snow.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1287 B.; Windwardside—Hellsgate: n. 1670 B.; Windwardside: n. 2306 B.; Bottom: n. 200; Peperpot: n. 198; Crispeen—Rendezvous 400 M.: n. 199.

I also saw *Drymaria cordata*: Laddergut.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Afr. and Asia trop. (Symb.).

88. Nymphaeaceae.

2513. **Nymphaea** Linn.

Nymphaea spec.

Cultivated.

94. Menispermaceae.2574. **Cissampelos** Linn.

Cissampelos Pareira Linn. Spec. (1753) 1031; Gris. Fl. 10; Duss 7; Millspaugh 488.

St. Eustatius: along roads in Oranjestad: n. 574 B.; slope of the Quill near Bengalen 100 M.: n. 667 B.; slope of the lower part of the Quill near Bengalen: n. 710 B.; seashore near White Wall: n. 848 B.; slope of the Quill: n. 55, 56; locality unknown: n. 218 H.

Saba: near the Sulphurmines: n. 1702 B.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2885 B.; Mount Paradis 150—400 M.: n. 3191 B.; locality unknown: n. 3620 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Afr., Asia trop. (Symb.).

2622. **Hyperbaena** Miers.

Hyperbaena domingensis Benth. in Journ. Linn. Soc. V. Suppl. II (1861) 50.

Cocculus domingensis DC. Syst. I (1818) 528; Gris. Fl. 10; Duss 7.

Pachygone domingensis Eichl. in Flora XLVII (1864) 394.

St. Eustatius: Bottom of the Quill 300 M.: n. 426 B., 891 B., 896 B.; top of the Quill 500 M.: n. 486 B.; slope of the lower part of the Quill near Bengalen: n. 708 B.; seashore: n. 57; Quill; n. 58, 73.

Saba: Springbaygut: n. 2092 B.

Cuba, Jamaica, Hispaniola, Portorico, St. Thomas, Guadeloupe, Dominica, Martinique. (Symb.).

98. Anonaceae.2729. **Anona** Linn.

Anona montana Macf. Flor. Jam. I (1837) 7; Gris Fl. 4.

St. Eustatius: near Bengalen: n. 1188a B., locality unknown: n. 274 H.; slope of Signalhill: n. 21.

Saba: along the Forthgut n. 1638 B.; Ladder: n. 20.

Antill., Guiana. (Symb.).

Anona muricata Linn. Spec. (1753) 536; Gris. Fl. 4; Duss 2; Millspaugh 488. Sour sap.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 54 B.; along the lower part of Signalhill: n. 121 B.; lower part of the slope of the Quill near Bengalen: n. 745 B.; along roads in Oranjestad: n. 1279 B.; locality unknown: n. 89 H., 173 G.

I also saw *Anona muricata* on the slope of the Quill on the side of White Wall.

Saba: lower part of the Bottom Mountain 300 M.: n. 1438 B.; Bottom: n. 1844 B.; Springbaygut: n. 2078 B.; Marypoint: n. 22; locality unknown: n. 3 L.

I also saw *Anona muricata*: between Bottom and Marypoint; Fortbaygut; Gilesgut.

St. Martin: Prince quarter (coll. George): n. 2463 B.; from Philipsburg to Belvedere: n. 2593 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2880 B.; lower part of Mount Paradis: n. 3354 B.

I also saw *Anona muricata*: Marigothill—Marigot; Nakedboyhill; hill on the northern side of Cul de Sac; from Belvedere to Oysterpond; one of the islands in Simonsbaylagoon; Top of Mount Paradis 400 M.

St. Croix ex Millspaugh.

Antill.; cult. in Amer. trop. Asia trop. and Afr. occ. (Symb.). Also cultivated.

Anona reticulata Linn. Spec. (1757) 537; Gris. Fl. 5; Duss 4; Millspaugh 488. Custard apple.

St. Eustatius: Slope of the Quill 350 M.: n. 1219 B.

Saba: Boobyhill 300 M.: n. 1632a B.; Bottom: n. 1842 B.

I also saw *Anona reticulata*: Laddergut.

St. Croix ex Millspaugh.

Antill.; cult. in Amer., Afr. and Asia trop. (Symb.).

Also cultivated.

Anona squamosa Linn. Spec. (1753) 537; Gris. Fl. 5; Duss 3; Millspaugh 488.

St. Eustatius: in a garden in Oranjestad: n. 525 B.; in deserted plantations at the seashore near Bengalen: n. 595 B.; locality unknown: n. 88 H., 91 G.

I also saw *Anona squamosa* in Signalhill.

Saba: Bottom: n. 1848 B., 2256 B.

I also saw *Anona squamosa*: Laddergut, Forthbaygut.

St. Martin: from Philipsburg to Guanabay: n. 2347 B.; Mountain between Mont des Accords and Centryhill: n. 2869 B.; from Belvedere to Oysterpond: n. 3049 B.; Mount Paradis 200—300 M.: n. 3309 B.

I also saw *Anona squamosa*: along roads from Philipsburg to Nakedboy-hill; Margothill—Marigot: near Bellevue; lower part of Milldrumhill; Vallée des Pères; one of the islands in Simonsbaylagoon.

St. Croix ex Millspaugh.

Antill.; cult. in Amer. cont. trop., Afr., Asia trop. (Symb.).

Also cultivated.

102. Lauraceae.

2783. *Persea* Gärtn.

Persea americana Mill. Gard. Dict. VIII ed. (1768).

Persea gratissima Gärtn. f. Fruct. III (1807) 222 tab. 221; Gris. Fl. 280. Duss 298.

Persea Persea Cock. in Bull. Torr. Bot. Club. XIX (1892) 95; Millspaugh 489. Pear tree, Avocado.

St. Eustatius: in deserted plantations near Tumbledown-dick-bay: n. 141 B., 146 B.; top of the Quill 400 M.: n. 223 B.; slope of the Quill near Bengalen 50—200 M.: n. 652 B.; in a garden near Glassbottle: n. 1210 B.; locality unknown: n. 112 H., 146 G.

Saba: Windwardside: n. 1928 B.; Mountain near Windward-side 600 M.: n. 2153 B.; locality unknown: n. 1565 B., 149 L.

I also saw *Persea americana* in Laddergut.

St. Martin: along roads near Anse des Pères: n. 3107 B.; Mount Paradis 150—400 M.: n. 3193 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham. Antill., trop. countries of both hemisph., Ind. in Mexico. (Symb.). Also cultivated.

2785. *Phoebe* Nees.

Phoebe elongata Nees. Syst. (1836) 116; Duss 298.

St. Eustatius: Quill: n. 256a B., 410 B., 898 B.

Cuba, Hispaniola, Portorico, Guadeloupe, Martinique, St. Vincent, Grenada, Tobago, Trinidad. (Symb.).

2790. *Nectandra* Roland.

Nectandra coriacea Griseb. in Flor. W. I. (1860) 281; Duss 301.

Nectandra Wildenoviana Nees. Syst. (1836) 321. Sweet wood.

St. Eustatius: Top of the Quill 400 M.: n. 221 B., 256 B., 272 B., 288 B., 289 B.; slope of the Quill 350 M.: n. 410 B.; slope of the Quill near Bengalen 50—200 M.: n. 648 B., 680 B., 696 B., 728 B., 751 B.; Bottom of the Quill 300 M.: n. 893 B., 902 B.; lower part of the Hills near Concordia: n. 981 B.; Glassbottle 150 M.: n. 1114 B.; top of the Quill: n. 248; Cliff: n. 243; slope of the Quill: n. 247; Quill: n. 168 G., 116 H.; locality unknown: n. 244, 245.

Saba: Paris 2—400 M.: n. 242; Windwardside: n. 1947 B.

St. Martin: Mountain between Mount des Accords and Centry-hill 200—300 M.: n. 2855 B., 2898a B.; Milddrumhill 300—400 M.: n. 3125 B.; Mount Paradis 200—300 M.: n. 3306 B., 3320 B.; locality unknown: n. 3584 B.

St. Croix. (Symb.).

Florida, Baham., Antill., Yucatan. (Symb.).

Nectandra Krugii Mez in Jahrb. Bot. Gart. Berl. V (1889) 422.

Nectandra discolor Gris. Flor. W. I. (1860) 282 non Nees. (Symb.). Black sweet wood.

St. Eustatius: Top of the Quill 500 M.: n. 477 B., 487 B.; Quill: n. 1, 2, 3, 4, 16, 19, 20, 22; locality unknown: n. 17, 18, 23, 24, 25, 215, 246.

Saba: Boobyhill 300 M.: n. 1575 B.; Windwardside: n. 1946 B.; near Devilshand: n. 2074 B.; near Flat Point: n. 2140 B.

Portorico, Guadeloupe, Dominica. (Symb.).

104. Papaveraceae.

2852. *Argemone* Linn.

Argemone mexicana Linn. Spec. (1753) 508; Gris. Fl. 13; Duss 8; Millspaugh 489. Thistle.

St. Eustatius: Seashore near Oranjestad: n. 43 B.; Cultuur-vlakte: n. 124; locality unknown: n. 20 G., 17 H.

Saba: Bottom: n. 2271 B.; locality unknown: n. 155 L.

St. Martin: I saw this plant: near Philipsburg and in pastureground near Bellevue.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Amer. cont. trop. and from there in trop. countries of the Old world. (Symb.).

105. Cruciferae.

2883. *Lepidium* Linn.

Lepidium virginicum Linn. Spec. (1753) 645; Gris. Fl. 14; Duss 9; Millspaugh 489.

St. Eustatius: along roads near Glassbottle: n. 885 B.; slope of the Quill: n. 340.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1292 B., 1363 B.; Mountain near Windwardside 600 M.: n. 2164 B.; Ladder: n. 342.; Peperpot: n. 341; locality unknown: n. 123 L.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2901 B.; hill on the northern side of Cul de Sac 0—100 M.: n. 3004 B.; Milldrumhill 200—300 M.: n. 3121 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Amer. sept. from Canada to Texas. (Symb.).

2920. *Cakile* Juss.

Cakile lanceolata O. E. Schulz. in Urb. Symb. Antill. III (1903) 504.

Cakile aequalis L'Hérit ap. DC. Syst. II (1821) 430; Gris. Fl. 14; (Symb.); Duss 10; Millspaugh 489.

St. Eustatius: Turtlebay: n. 328.

Saba: between Hellsgate and Marypoint: n. 2044 B.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Columbia. (Symb.).

2949. *Brassica* Linn.

Brassica integrifolia O. E. Schulz in Urb. Symb. Antill. III (1903) 509.

Sinapis brassicata Gris. Flor. W. I. (1859) 14; Millspaugh 489.

Sinapis juncea Eichl. in Mart. Fl. Bras. XIII. 1. (1865) 306. (non L.). (Symb.); Duss 10. Mustard.

St. Eustatius: locality unknown: n. 141 H., 292 G.

Saba: Windwardside: n. 1884 B.; Rendez-vous 400 M.: n. 325, 326; locality unknown: n. 26 L.

St. Croix ex Millspaugh.

Antill.; Indig. in Asia austr. and orient. (Symb.).

107. Capparidaceae.**3087. Gynandropsis DC.****Gynandropsis pentaphylla** DC. Prodr. I (1824) 238.*Cleome pentaphylla* Linn. Spec. (1763) 938; Gris. Fl. 15; Duss 11; Millspaugh 489. Massamby.**St. Eustatius:** along roads in Oranjestad: n. 547 B., 933 B.; Cultuurvlakte: n. 205, 185; locality unknown: n. 29 H., 33 G.**Saba:** Bottom: n. 1841 B.; locality unknown: n. 153 L.; Bottom: n. 186.**St. Martin:** Colebayhill: n. 188; Prince quarter: n. 189.I also saw *Gynandropsis pentaphylla*: Mount Paradis 100–200 M.**Anguilla** Boldingh.**St. Croix** ex Millspaugh.

Baham., Antill., Amer. cont. calid.; Indig. in Afr. and Asia trop. (Symb.).

3090. Polanisia Raf.**Polanisia viscosa** DC. Prod. I. (1824) 242.*Cleome graveolens* Rafin. in Journ. phys. (1819) 98.*Cleome viscosa* Linn. Spec. (1753) 672; Gris. Fl. 16; Duss 12.*Polanisia icosandra* Wight et Arn. Prod. (1834) 22; Millspaugh 490. Wild massamby.**St. Eustatius:** along roads in Oranjestad: n. 535 B.; locality unknown: n. 28 H., 32 G.I also saw *Polanisia viscosa* along the seashore near Oranjestad.**Saba:** lower part of the Bottom Mountain 250 M.: n. 1393 B.**St. Martin:** from Philipsburg to Guanabay: n. 2401 B.; from Philipsburg to Belvedere: n. 2558 B.; Colebayhill: n. 190.**St. Croix** ex Millspaugh.

Antill.; introduced from South. China and Senegamb. (Gris. Fl.).

3101. Capparis Linn.**Capparis Breynia** Jacq. Sel. Amer. (1763) 161 t. 103; Duss 13.*Capparis amygdalina* Lam. Enc. I (1783) 608; Gris. Fl. 17.**St. Eustatius:** Gut near Fort de Wind 50 M.: n. 832 B.; from Little Mountain to Venusbay: n. 995 B.; along roads in Oranjestad: n. 1068 B.; slope of the Quill on the side of White Wall 200 M.: n. 1178 B.; Pangohill: n. 168; Little Mountain—Boven: n. 166, 167.I also saw *Capparis Breynia* on Signalhill,

Saba: between Bottom and Marypoint: n. 1466 B.; Springbay-gut: n. 2120 B.; Windwardside: n. 2319 B.

St. Martin: Prince quarter (coll. George): n. 2460 B., 2481 B.; Nakedboyhill 50—200 M.: n. 2695 B., 200—265 M.: n. 2730 B., 2731 B.; Marigotroad near Marigot: n. 2815 B.; Mountain between Mout des Accords and Centryhill 200—300 M.: n. 2889 B.; gut from Bethlehem to Cul de Sac: n. 3433 B.; near Fort Amstêrdam: n. 143; behind Centryhill: n. 144.

I also saw *Capparis Breynia* on Mount Paradis 100—200 M.

St. Croix. (Symb.).

Antill., Amer. centr., Nova Granata, Venezuela. (Symb.).

Capparis coccolobifolia Mart. ap. Eichl. in Mart. Fl. bras. XIII. I. (1865) 284.

Capparis cynophallophora var. *latifolia* Griseb. Flor. W. I. (1859) 18 (p.p.) (Symb.).

St. Eustatius: in deserted plantations near Tumbledown-dick-bay: n. 145 B.

St. Martin: Prince quarter (coll. George): n. 2478 B.; Mill-drumhill 300—400 M.: n. 3147 B.; near Fort Amstêrdam: n. 109; Centryhill: n. 129; Guanabay: n. 108; locality unknown: n. 3612 B.

St. Croix. (Symb.).

Antill., Curaçao, Venezuela. (Symb.).

Capparis cynophallophora Linn. Spec. (1762) 721; Gris. Fl. 18; Duss 13; Millspaugh 490. Mustard tree, Man of war bush.

St. Eustatius: Slope of the lower part of Signalhill: n. 118 B.; seashore near Oranjestad: n. 27 B.; slope of the Quill near Bengalen 200 M.: n. 691 B., 695 B., 734 B.; along roads in Oranjestad: n. 925 B.; from Little Mountain to Venusbay: n. 988 B.; Bengalen: n. 749 B.; near Maryglory 150 M.: n. 1145 B.; Seinpost: n. 124; Red Cliff: n. 125; Boven: n. 127; locality unknown: n. 46 H., 191 G.

I also saw *Capparis cynophallophora* on Signalhill; in a garden in Oranjestad.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1305 B.; along the Forthgut 150—200 M.: n. 1626 B.; along roads 200—300 M.: n. 1714 B.; slope of the Mountain near St. John's: n. 1736 B.; St. Giles: n. 1859 B.; Bottom: n. 1882 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2059 B.; Windwardside: n. 2310 B.; Great Hill: n. 202; Quarter: n. 128.

I also saw *Capparis cynophallophora* in Springbaygut.

St. Martin: Cul de Sac: n. 2606 B.; near Mulletpond: n. 2772 B.; hills on the eastern side of Cul de Sac: n. 3436 B.

I also saw *Capparis cynophallophora*: Cul de Sac; Marigot—Marigot; Nakedboyhill 100–200 M.; near Saline: hills on the northern side of Cul de Sac; along roads from Philipsburg to Simonsbay; from Belvedere to Oysterpond; lower part of Milddrumhill; Mount Paradis 100–400 M.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida austral., Antill., Panama, Amer. austr. (Symb.).

Capparis frondosa Jacq. Enum. (1760) 24; Gris. Fl. 19; Duss 14; Millspaugh 490. *Church blossom*.

St. Eustatius: in deserted plantations near Tumbledown Dick-bay: n. 140 B.; Bottom of the Quill 300 M.: n. 445 B.; slope of the Quill near Bengalen 50 M.: n. 655 B.; hills near Concordia: n. 967 B.; slope of Signalhill: n. 157; Cultuurvlakte: n. 158; near Seinpost: n. 159; Red Cliff: n. 160; Quill: n. 161, 162, 201; Waterrock: n. 162; locality unknown: n. 189 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1334 B.; lower part of the Bottom Mountain 300 M.: n. 1443 B.; Laddergut 200 M.: n. 2007 B.; Peperpot: n. 148, 163; Cresp.—Rendez-vous: n. 164; Marypointgut: n. 152.

I also saw *Capparis frondosa* in Forthbaygut and Springbaygut.

St. Martin: from Philipsburg to Guanabay: n. 2383 B.; Prince-quarter (coll. George): n. 2461 B., 2482 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2961 B.; from Belvedere to Oysterpond: n. 3058 B.; Centryhill: n. 145; John Ednyhill: n. 146.

I also saw *Capparis frondosa* along roads near Bethlehem; Mount Paradis 100–200 M.; lowerpart of Milddrumhill; Nakedboyhill.

St. Croix ex Millspaugh.

Antill., Amer. centr., Nova Granata, Venezuela. (Symb.).

Capparis jamaicensis Jacq. Enum. (1760) 23; Gris. Fl. 18; Duss 13; Millspaugh 490.

St. Eustatius: Slope of the Quill above White Wall 350 M.: n. 3 B.; slope of the lower part of Signalhill: n. 105 B.; top of the Quill 400 M.: n. 305 B.; top of the Quill 500 M.: n. 488 B., 494 B.; near Maryglory 150 M.: n. 1139 B.; near White Wall: n. 1256a B.; Boven: n. 133, 134, 135; Greatgut: n. 174.

Saba: Boobyhill 300 M.: n. 1621 B.; St. Giles: n. 1862 B.; Windwardside: n. 155; Wellridge: n. 179, 180; Bottom: n. 222; locality unknown: n. 48 L.

St. Martin: from Filipsburg to Guanabay: n. 2368 B., 2382 B., 2412 B., 2417 B.; Prince quarter (coll. George): n. 2466 B., 2476 B., 2480 B.; Fort Willem 0—50 M.: n. 2539 B.; Nakedboyhill 50—200 M.: n. 2712 B.; near Mulletpond: n. 2771 B.; near Fort Willem: n. 2797 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2960 B., 2965 B.; between Belvedere and Oysterpond: n. 3075 B., 3078 B.; Milldrumhill 300—400 M.: n. 3136 B.; Centryhill: n. 178, 170; Oysterpond: n. 177; Prince quarter: n. 171; John Ednyhill: n. 175; road to Guanabay: n. 176.

I also saw *Capparis jamaicensis* along roads from Filipsburg to Simonsbay, in the Lowlands and near Saline.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida austral., Baham., Antill. (Symb.).

3104. *Morisonia* Linn.

Morisonia americana Linn. Spec. (1753) 503. Gris. Fl. 19; Duss 14; Millspaugh 490. Wild misple.

St. Eustatius: Slope of the lower part of Signalhill: n. 114 B.; from Little Mountain to Venusbay: n. 965 B.; Little Mountain: n. 195; Boven: n. 218; locality unknown: n. 217.

Saba: between Bottom and Marypoint 100—200 M.: n. 1556 B.; Cow pasture: n. 196.

St. Martin: Prince quarter (coll. George): n. 2483 B.; from Filipsburg to Belvedere: n. 2568 B.; Colebayhill: n. 219, 197.

St. Croix ex Millspaugh.

Antill., Nova Granata. (Gris. Fl.).

109. *Moringaceae*.

3128. *Moringa* Juss.

Moringa oleifera Lam. Enc. I (1783) 398.

Moringa pterygosperma Gärtner. Fruct. II (1791) 314 t. 147 f. 2; Gris. Fl. 16; Duss 12.

Moringa moringa Millsp. in Field. Col. Mus. Bot. I (1902) 490. Millspaugh 490. Moringo.

St. Eustatius: in a garden in Oranjestad: n. 66 B., 227 G.

Saba: Bottom: n. 1878 B.

St. Martin: from Filipsburg to Belvedere: n. 2576 B.

I also saw *Moringa oleifera* along roads from Filipsburg to Nakedboyhill.

St. Croix ex Millspaugh.

Baham., Antill., trop countries of both hemisph. (Symb.). Indig. in India orient. Also cultivated.

115. Crassulaceae.

3165. *Bryophyllum* Salisb.

Bryophyllum pinnatum S. Kurz. in Journ. As. Soc. Beng. XL. (1871) II 52; Millspaugh 490.

Bryophyllum calycinum Salisb. Parad. Lond. (1805) t. 3; Gris. Fl. 303; Duss 319. Leaf of live.

St. Eustatius: along the lower part of Signalhill: n. 120a B.; Cultuurvlakte: n. 335, 336, 287 H., 297 H.

Saba: Great Hill 400 M.: n. 1379 B.; top of the Cliff near the seaside of Boobyhill: n. 1607 B.; Peperpot: n. 337; locality unknown: n. 288 L.

I also saw *Bryophyllum pinnatum*: gut between Hellsgate and Mastiegut; Crispeñhill; Windwardside; Laddergut; Springbaygut.

St. Martin: I did not collect the plant on this island, but I saw it: along roads from Philipsburg to Nakedboyhill; Marigothill—Marigot; from Belvedere to Oysterpond; Mount Paradis 400 M.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., trop. and subtrop. countries of both hemisph. (Symb.).

126. Rosaceae.

3396. *Prunus* Linn.

Prunus myrtifolia Urb. in Symbolae Antillanae V (1904) 93.

Prunus sphaerocarpa Sw. Prodr. (1788) 80; Gris. Fl. 231.

St. Eustatius: Slope of the Quill 400 M.: n. 272a B.

Antill., Brasilia. (Symb.).

3398. *Chrysobalanus* Linn.

Chrysobalanus icaco Linn. Spec. (1753) 513; Gris. Fl. 229; Duss 257; Millspaugh 490. Cocoa plum, Fat pork.

St. Eustatius: Top of the Quill 400 M.: n. 339 B.; slope of the Quill 300 M.: n. 880 B.; slope of the Quill 150 M.: n. 232, 247; slope of the Quill: n. 255, 256, 257, 258; Grandgut: n. 259; locality unknown: n. 254, 73 H., 155 H., 224 G., 246 G.

Saba: between Bottom and Marypoint 350 M.: n. 1491*a* B.; Mountain 600 M.: n. 1739 B.; Bottom: n. 2290 B.; Long peak: n. 249; locality unknown: n. 18 L., 335 L.

St. Martin: Colebay (coll. George): n. 2514 B.: seashore near Simonsbay: n. 3036 B.; Lowlands: n. 3273 B.

St. Croix ex Millspaugh.

Florida austral., Baham., Antill., Amer. austr., Afr. austr. (Symb.).

3403. *Hirtella* Linn.

Hirtella triandra Sw. Prodr. (1788) 51; Gris. Fl. 230; Duss 258. Hairy plum.

Saba: Mountain near Hellsgate 600 M.: n. 2277 B.

Antill., Amer. austr. (Symb.).

128. Leguminosae.

3436. *Inga* Willd.

Inga laurina Willd. Spec. IV. 2 (1806) 1018; Gris. Fl. 227; Duss 255; Millspaugh 491. Red wood.

St. Eustatius: Top of the Quill 400 M.: n. 183 B., 205 B., 380 B., 481 B., 483 B., 498 B., 863*a* B., 367 B., 395 B.; slope of the Quill near Bengalen: n. 726 B.; Kant: n. 312, 313, 314; locality unknown: n. 51, 100.

Saba: lower part of the Bottom Mountain 250 M.: n. 1421 B.; Boobyhill 300 M.: n. 1578 B.; Windwardside: n. 1611 B., 2142 B.; Peperpot: n. 79, 80.

St. Martin: Mount Paradis: n. 3214 B., 3226 B., 3327*a* B.

St. Croix ex Millspaugh.

Hispaniola, Portorico, St. Thomas, St. Jan, St. Kitts, Antigua, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Barbados, Grenada, Trinidad. (Symb.).

3440. *Enterolobium* Mart.

Enterolobium Timbouwa Mart. Flora XX (1827) 128.

Cultivated. [Antill., Amer. austr.] (E. & P.).

3441. *Pithecolobium* Mart.

Pithecolobium unguis-cati Benth. in Hook. Lond. Journ. Bot. III (1844) 200; Gris. Fl. 226; Duss 254; Millspaugh 491. Crabwood.

St. Eustatius: Slope of the lower part of Signalhill: n. 106 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 768 B.; Signalhill 200 M.: n. 1035 B.; Cultuurvlakte: n. 1250; slope of the Quill: n. 1369; Boven: n. 1363; Gilboohill: n. 1379; road to English Quarter: n. 1372; locality unknown: n. 284 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1541 B.; gut between Hellsgate and Mastigut 300—400 M.: n. 2054 B.; Hellsgategut: n. 1361, 1370; pasture: n. 1368.

St. Martin: From Guanabay to Philipsburg: n. 2394 B.; from Philipsburg to Belvedere: n. 2569a B.; Nakedboyhill: n. 2681 B.; near Mulletpond: n. 2754 B.; hill on the northern side of Cul de Sac 0—100 M.: n. 2971 B.

I also saw *Pithecolobium unguis-cati*: Marigothill—Marigot; from Belvedere to Oysterpond; along roads near Grande Case; Lowlands; Mount Paradis 300—400 M.; Tintamarre; gut from Bethlehem to Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida austr., Key Ins., Antill., Venezuela, Nova Granata. (Symb.).

3443. *Albizzia* Durazz.

Albizzia lebbek Benth. in Hook. Lond. Journ. Bot. III (1844) 87.

Cultivated. [Baham., Antill., Curaçao, Asia austr. and orient., Afric. trop.] (Symb.).

3446. *Acacia* Willd.

Acacia Farnesiana Willd. Spec. IV. 2 (1806) 1083; Gris. Fl. 222; Duss 250; Millspaugh 491. Casha.

St. Eustatius: Red Cliff: n. 390.

Saba: between Bottom and Marypoint 100—200 M.: n. 1460 B.; St. Giles: n. 1864 B.; Paris 2—400 M.: n. 372; Bottom: n. 414.

St. Martin: from Guanabay to Philipsburg: n. 2434 B.; Fort Willem 0—50 M.: n. 2545 B.; near Mulletpond: n. 2778 B.; Green Key: n. 3365 B.; Philipsburg: n. 251; Fort Amsterdam: n. 386.

St. Croix ex Millspaugh.

Baham., Antill., warmer countries of both hemisph. (Symb.).

Acacia macrantha Humb. et Bonpl. ex Willd. Spec. Plant. IV (1805) 1080; Gris. Fl. 221; Duss 249. French casha, Creole casha, Spanish casha.

St. Eustatius: Seashore near Oranjestad: n. 15 B., 38 B.; top

of the Quill 400 M.: n. 216a B., 318 B.; along roads in Oranjestad: n. 543 B., 567 B.; Bengalen: n. 630 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 781 B.; near White Wall: n. 858 B.; near Zeelandia: n. 253; slope of the hills near Concordia: n. 260; top of Pangohill: n. 259; Seafeathersbay: n. 263; locality unknown: n. 36 H., 39 H., 50 G., 304 G.

St. Martin: Prince quarter (coll. George): n. 2454 B.; from Filipsburg to Belvedere: n. 2552a B.; between Belvedere and Oysterpond: n. 3076 B.; lower part of Mount Paradis: n. 3291 B.; Colebayhill: n. 49.

Antill., Mexico, Venezuela, Equador, Peru, Buenos Ayres, Galapagos. (Gris. Fl.).

Acacia riparia H. B. K. Nov. Gen. VI (1823) 276; Duss 248.

Acacia sarmentosa Gris. Kar. (1857) 64; (non Desv.) (Symb.); Gris. Fl. 221.

St. Eustatius: near Bengalen: n. 600a B.

St. Martin: Prince quarter (coll. George): n. 2455 B.; Naked-boyhill 200—265 M.: n. 2732 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2862 B.; Colebey—Centryhill: n. 787.

St. Croix ex Millspaugh.

Antill., Amer. austr. trop. (Symb.).

Acacia tortuosa Willd. Spec. Plant. IV (1805) 1083; Gris. Fl. 222; Millspaugh 491. Dutch casha.

St. Eustatius: in deserted plantations at the seashore near Bengalen: n. 583 B., 590 B., 600 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 780 B.; along roads near English Quarter: n. 1004 B.; Signalhill 150 M.: n. 1024 B., 1026 B.; slope of the Quill on the side of White Wall 200 M.: n. 1175 B.; Cultuurvlakte: n. 383, 395; seashore: n. 384; Red Cliff: n. 378; locality unknown: n. 371.

St. Croix ex Millspaugh.

Antill., Mexico, Venezuela, Equador, Galapagos. (Gris. Fl.).

. 3447. **Leucaena** Benth.

Leucaena glauca Benth. in Hook. Journ. Bot. IV (1842) 416; Gris. Fl. 220; Duss 247; Millspaugh 491. Mimosa, Tantan.

St. Eustatius: Seashore near Oranjestad: n. 23 B.; slope of the lower part of the Quill near Bengalen: n. 719 B.; in a garden in Oranjestad: n. 1274 B.; Cultuurvlakte: n. 317, 318; locality unknown: n. 37 H., 44 G.

I also saw *Leucaena glauca* on the slope of the Cliff; Signallhill; deserted plantations near Bengalen.

Saba: Laddergut: n. 318; locality unknown: n. 129 L., 136 L.

St. Martin: from Philipsburg to Belvedere: n. 2590 B.

I also saw *Leucaena glauca*: hill between Mont des Accords and Centryhill; near Saline; hill on the northern side of Cul de Sac; along roads near Grande Case; lower part of Milldrumhill; Mount Paradis 300—400 M.; hills on the eastern side of Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Curaçao, warmer countries of both hemisph.; in Amer. trop. indig? (Symb.).

3449. *Mimosa* Linn.

Mimosa pudica Linn. Spec. (1753) 518; Gris. Fl. 219; Duss 244; Millspaugh 492. Shame lady shame.

St. Eustatius: Slope of the Quill on the side of White Wall 200 M.: n. 1184 B.; along roads near Bengalen: n. 1192 B.; locality unknown: n. 216 H.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Asia trop. (Symb.).

3450. *Desmanthus* Willd.

Desmanthus depressus Humb. et Bonpl. ex Willd. Spec. Plant. IV (1805) 1046; Gris. Fl. 218; Duss 244.

St. Eustatius: Top of the Quill 450 M.: n. 347 B.; lower part of the slope of the Quill near Bengalen: n. 721 B.; Fort de Wind: n. 797a B.; along roads near English Quarter: n. 999 B.; Cultuurvlakte: n. 282.

St. Martin: Nakedboyhill 50—200 M.: n. 2677 B.; Marigot road near Marigot: n. 2812 B.; lower part of the Milldrumhill: n. 3167 B.; Tintamarre: n. 3375 B., 3402 B.

Florida. Antill., Texas to Peru, Galapagos. (Gris. Fl.).

Desmanthus virgatus Willd. Spec. IV. 2 (1806) 1047; Gris. Fl. 218; Duss 244; Millspaugh 492. Wild tantan.

St. Eustatius: in a garden in Oranjestad: n. 71 B.; seashore near Tumbledowndickbay: n. 161 B.; Bengalen: n. 603 B.; slope of the Quill near Bengalen 200 M.: n. 741 B.; along roads in Oranjestad: n. 1075 B.; near Boven 200 M.: n. 1244 B.; Cultuurvlakte: n. 296; locality unknown: n. 162 G., 302 G.

Saba: Boobyhill 300 M.: n. 1614 B.; Laddergut 0—200 M.: n. 1967 B., 2012 B.; Paris 200—400 M.: n. 99.

St. Martin: from Philipsburg to Guanabay: n. 2371 B.; Cul de Sac—Marigot hill 0—200 M.: n. 2633 B.; between Belvedere and Oysterpond: n. 3080 B.; slope of Centryhill: n. 98; French frontier n. 295.

St. Croix ex Millspaugh.

(Bermud.), Baham., Antill., (Amer. cont. trop. and subtrop. India-orient.). (Symb.).

3451. **Neptunia** Lour.

Neptunia plena Benth. in Hook. Journ. Bot. IV (1842) 355; Gris. Fl. 218; Duss 243. Waterthistle.

St. Martin: near Marigot: n. 2627a B.

Antill., Curaçao, Amer. austr., Asia trop. (introd.) (Symb.).

3459. **Adenanthera** Linn.

Adenanthera pavonina Linn. Spec. (1753) 384.

Cultivated. [Asia trop.] (Symb.).

3499. **Hymenaea** Linn.

Hymenaea courbaril Linn. Spec. (1753) 1192; Gris. Fl. 213; Duss 238; Millspaugh 492. Locust.

St. Eustatius: Slope of the Quill near Bengalen 50—200 M.: n. 649 B.; slope of the Quill 250 M.: n. 889 B.; slope of the Quill: n. 461, 2947; locality unknown: n. 75 H., 175 H., 216 G.

Saba: Windwardside: n. 2332 B.; locality unknown: n. 348 L.

St. Martin: from Philipsburg to Belvedere: 2592 B.; Cul de Sac: n. 2995 B.; Centryhill: n. 472; French frontier: n. 462.

I also saw *Hymenaea courbaril* on the hills on the northern side of Cul de Sac.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop. (Symb.). Also cultivated.

3508. **Tamarindus** Linn.

Tamarindus indica Linn. Spec. (1753) 34; Gris. Fl. 213; Duss 237; Millspaugh 492. Tamarind.

St. Eustatius: Seashore near Oranjestad: n. 29 B.; slope of the lower part of Signalhill: n. 112 B.; in a garden in Oranjestad: n. 1065 B., 40 H., 47 G.

Saba: between Bottom and Marypoint 100—200 M.: n. 1547 B.; Bottom: n. 1873 B., 1874 B.; locality unknown: n. 117 L.

St. Martin: from Philipsburg to Guanabay: n. 2350 B.; Prince-quarter (coll. George): n. 2468 B.

I also saw *Tamarindus indica*: Marigothill—Marigot; Fort Willem; near Saline; along roads from Philipsburg to Simonsbay; Vallée des Peres; one of the islands in Simonsbaylagoon.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Trop. countries of both hemisph.; Ind. veris. in Africa. trop. (Symb.). Also cultivated.

3528. *Bauhinia* Linn.

Bauhinia Kappleri Sagot in Ann. Sc. nat. VI Sér. vol. XIII (1882) 317.

Cultivated. [Asia austro—orient.]. (Symb.).

3536. *Cassia* Linn.

Cassia alata Linn. Spec. (1753) 378.

Cultivated. [Cosmop. trop.]. (Ind. Kew.).

Cassia bicapsularis Linn. Spec. (1753) 376; Gris. Fl. 207; Duss 231; Millspaugh 492. Blydog.

St. Eustatius: Seashore near Oranjestad: n. 26 B., 50 B.; slope of the lower part of Signalhill: n. 113a B.; in deserted plantations at the seashore near Bengalen: n. 591 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 793 B.; gut near Fort de France: n. 846 B.; along roads near Glassbottle: n. 1214 B.; locality unknown: n. 25 G.; Cultuurvlakte: n. 329; near Seinpost: n. 325.

Saba: Windwardside: n. 1891 B.; Peperpot: n. 330; Great-hill: n. 315; locality unknown: n. 28 L.

I also saw *Cassia bicapsularis*: in Laddergut.

St. Martin: from Philipsburg to Guanabay: n. 2370 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2955 B.; Cul de Sac: n. 320.; Guanabay: n. 313.

I also saw *Cassia bicapsularis*: along roads from Philipsburg to Naked-boyhill; Marigothill to Marigot; Nakedboyhill; in pastureground near Bellevue; lower part of Milldrumhill; Mount Paradis 300–400 M.; one of the islands of Simonsbaylagoon; Lowlands.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Curaçao., Amer. cont. trop., Inquil. in India Orient. and Madeira. (Symb.).

Cassia fistula Linn. Spec. (1753) 377. Liquorice plant.

Cultivated. [Asia trop.]. (Symb.).

Cassia glandulosa Linn. Syst. X ed. II (1759) 1017; Gris. Fl. 211; Duss 233.

Chamaecrista glandulosa Greene in Pitton. IV (1899) 28; Millspaugh 493. Wild peas.

St. Eustatius: Top of the Quill 390 M.: n. 499 B.; slope of the Quill near Bengalen 100 M.: n. 666 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 787 B.; near White Wall: n. 1262 B.; Cultuurvlakte: n. 740; Little Mountain: n. 1768; slope of the Quill: n. 992, 1763; locality unknown: n. 81 H., 99 G., 170 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1350 B.; between Windwardside and Hellsgate: n. 1680 B.; Windwardside: n. 2324 B.; Peperpot: n. 1770, 732; Crespeen—Rendez-vous: n. 991.

St. Martin: Marigotbill—Marigot 0—200 M.: n. 2651 B.; Centryhill: n. 260; Marigotroad: n. 993.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Amer. austr. (Symb.).

Cassia obovata Collad.; Gris. Fl. 209.

St. Martin: along the saltpond near Filipsburg: n. 2441 B.; Guanabay: n. 274.

Anguilla Boldingh.

From Africa trop. (Gris. Fl.).

Cassia occidentalis Linn., Spec. (1753) 377; Gris. Fl. 209; Duss 235; Millspaugh 492. Bitterroot.

St. Eustatius: Seashore near White Wall: n. 859 B.; along roads near Glassbottle: n. 864 B.; along roads in Oranjestad: n. 562a B., 1103 B.; locality unknown: n. 4 G., 4 H.

Saba: lower part of the Bottom mountain 250 M.: n. 1394 B.; Bottom: n. 2243 B.; Gainbygut: n. 1035; locality unknown: n. 162 L., 237 L.

St. Martin: along roads near Fort Willem: n. 2525 B.; Mari-gotroad near Marigot: n. 2819 B.; in pastureground near Bellevue: n. 2843 B.; Prince quarter: n. 1034; locality unknown: n. 2524 B.

I also saw *Cassia occidentalis*: Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida austr., Baham., Antill., Amer., Africa, Asia trop. (Symb.).

Cassia polyadena DC. Pl. Rar. Jard. Genève. 2 Rapp. (1825). 12; Gris. Fl. 210.

St. Martin: Lowlands: n. 3272 B., 3274 B., 3288 B.

Baham., Cuba, Domingo, Jamaica, St. Barthélemy, Guadeloupe. (Herb. Krug et Urban).

3543. **Krameria** Loeffl.

Krameria ixina Linn. (1762) 117; Gris. Fl. 31; Wild senna.

St. Eustatius: Signalhill 200 M.: n. 126 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 763 B.; near Maryglory 150 M.: n. 1163 B.; near White Wall on the top of the Cliff: n. 1256 B.; Cultuurvlakte near Golden Rock: n. 181; slope of the hills near Fort Amsterdam: n. 164; Signalhill—Red Cliff: n. 163; locality unknown: n. 200 H.

Hispaniola, St. Thomas, Antigua, Curaçao, Columbia. (Symb.).

3551. **Parkinsonia** Linn.

Parkinsonia aculeata Linn. Spec. (1753) 375; Gris. Fl. 204; Duss 227; Millspaugh 493. Wonder tree, Jeruzalem thorn.

St. Eustatius: along roads in Oranjestad: n. 915 B., 63, 73, 150 H., 73 G.

St. Martin: near Philipsburg: n. 2340 B.

I also saw *Parkinsonia aculeata* on Mount Paradis 100—200 M.

St. Croix ex Millspaugh.

Key ins., Antill., Amer. cont. trop.; escaped in Afr. and Asia trop. (Symb.). Also cultivated.

3552. **Haematoxylon** Linn.

Haematoxylon campechianum Linn. Spec. (1753) 384; Gris. Fl. 204; Duss 226; Millspaugh 493. Log wood.

St. Eustatius: along a road near Oranjestad: n. 1254 B; locality unknown: n. 291 H.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cent. et Austr. sept. (Symb.).

3556. **Poinciana** Linn.

Poinciana regia Boj. ex Hook. Bot. Mag. (1829) tab. 2284. July tree, Flamboyant.

Cultivated. [Madagascar.]. (Symb.).

3559. **Caesalpinia** Linn.

Caesalpinia ciliata Urb. in Symbolae Antillanae VI (1900) 275.

Guilandina Bonduc Gris. Kar. (1857) n. 466 p. 60 et quoad specim. Guad. Flor. p. 240; non Linn. (Symb.).

Guilandina Grisebachiana Kr. et Urb. in Duss Flore. Ant. franc. (1897) 229.

St. Eustatius: Seashore near Oranjestad: n. 21 B.; seashore near Tumbledown Dickbay: n. 156 B.; along roads in Oranjestad: n. 1090 B., 69 H.

Saba: O. Kuntze (Symb.).

St. Martin: Colebay (coll. George): n. 2501 B.; near Mulletpond: n. 2766 B.

Barthélemy, Antigua, Guadeloupe, Desirade, Marie Galante, Martinique. (Symb.).

Caesalpinia coriaria Willd. Spec. Plant. II (1799) 532; Millspaugh 493.

Lebidibia coriaria Schlecht. in Linnaea V (1830) 193; Gris. Fl. 231; Duss 231. Divi divi.

St. Eustatius: in a garden in Oranjestad: n. 64 B., 245 G.

Saba: locality unknown: n. 301 L.; Windwardside: n. 2331 B.

St. Martin: along roads from Colebayhill to Simonsbay: n. 3016 B., 3017 B.

St. Croix ex Millspaugh.

Baham., Antill., Curaçao, Mexico, Venezuela, Nova Granata. (Symb.).

Caesalpinia Crista Linn. Spec. I ed. I (1753) 380; Millspaugh 493.

Guilandina glabra. Griseb. Flor. W. I. (1860) 205 (quoad specim. Martin. a. Cl. Sagot. non Duchassaing missum.); Duss 228; (Symb.).

St. Eustatius: Top of the Quill 400 M.: n. 306 B.; slope of the Quill 50 M.: n. 635; locality unknown: n. 160 G.

Saba: lower part of the Bottom Mountain 250 M.: n. 1409 B.; Bottom: n. 634; locality unknown: n. 354 L.

St. Martin: near Mulletpond: n. 2788 B.

St. Croix ex Millspaugh.

Bermuda., Key ins., Baham., Antill., Amer. cont. trop., Asia austro-orient., Nova Guinea, Australia. (Symb.).

Caesalpinia pulcherrima Sw. Obs. (1791) 166; Gris. Fl. 205; Duss 230; Millspaugh 493. Pride of Barbados.

St. Eustatius: in a garden in Oranjestad: n. 1088 B., 67 H., 68 H., 74 G., 75 G.

Saba: St. Giles: n. 1871 B.; locality unknown: 223 L.

St. Martin: from Philipsburg to Belvedere: n. 2598 B.

I also saw *Caesalpinia pulcherrima* on Marigot hill near Bellevue.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Key ins., Baham., Antill., Amer. cont. trop., Afr. trop., Asia trop. Patria ignot. (Symb.). Also cultivated.

3669. *Crotalaria* Linn.

Crotalaria incana Linn. Spec. (1753) 716; Gris. Fl. 180; Duss 193; Millspaugh 494. Small yellow popbush.

St. Eustatius: along roads: n. 256 H., 596a B.

Saba: Windwardside: n. 2144 B.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2872 B.; roads from Marigot to Philipsburg: n. 3110 B.; French frontier: n. 2294.

St. Croix ex Millspaugh.

Florida austral., Antill., warmer countries of both hemisph. (Symb.).

Crotalaria retusa Linn. Spec. (1753) 715; Gris. Fl. 179; Duss 193; Millspaugh 494. Big yellow popbush.

St. Eustatius: Top of the Quill 400 M.: n. 231 B.; White Wall: n. 810 B.; along roads near Bengalen: n. 1187 B.; Fairplay: n. 625; Cultuurvlakte: n. 635; slope of the Quill: n. 645; along roads: n. 285 G., 286 H.

Saba: lower part of the Bottom Mountain 250 M.: n. 1387 B.; Mountain 600 M.: n. 1753 B.; Laddergut 0—200 M.: n. 1961 B.; locality unknown: n. 66 L.

St. Martin: from Guanabay to Philipsburg: n. 2436 B.; Point Blanche (coll. George): n. 2491 B.; from Philipsburg to Belvedere: n. 2600 B.; in an orchard near Bellevue: n. 2848 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop., Afr. trop., Asia austro-orient., Austral. (Symb.).

Crotalaria verrucosa Linn. Spec. (1753) 715; Gris. Fl. 178; Duss 192; Millspaugh 494. Purple popbush.

St. Eustatius: along roads in Oranjestad: n. 923 B.; Fairplay: n. 1634; slope of the Quill: n. 1614, 1624; locality unknown: n. 285 H.

St. Martin: along roads from Grande Case to Marigot: n. 3108 B. Antill.; from the East Indies. (Gris. Fl.).

3688. **Medicago** Linn.**Medicago sativa** Linn. Spec. (1753) 778.

Cultivated. [Europa]. (Ind. Kew.).

3702. **Indigofera** Linn.**Indigofera suffruticosa** Mill. Gard. Dict. VIII ed. (1768) n. 2.*Indigofera anil* Linn. Mant. II (1771) 272; Gris. Fl. 181; Duss 134; Millspaugh 494. Wild indigo.**St. Eustatius:** in a garden in Oranjestad: n. 73 B.; Bengalen: n. 609 B.; along roads near English Quarter: n. 1001 B.; Fairplay: n. 290; Cultuurvlakte: n. 221; slope of the Quill: n. 231, 241; locality unknown: n. 172 G., 157 H., 219 G.**Saba:** between Bottom and Marypoint 100—200.: n. 1524 B.; along roads 200—300 M.: n. 1709 B.; St. Giles: n. 1867 B.; Hellsgate—Marypoint: n. 2047 B.; gut near Devilshand: n. 2077a B.; Ladder: n. 251; locality unknown: n. 87 L., 251 L.I also saw *Indigofera suffruticosa* on Crespeenhill.**St. Martin:** along roads near Filipsburg: n. 2528 B.I also saw *Indigofera suffruticosa* near Grande Case.**St. Croix** ex Millspaugh.

Baham., Antill., Amer. cont. calidior; inq. in the trop. Old world. (Symb.).

3718. **Tephrosia** Pers.**Tephrosia cathartica** Urb. Symb. Antill. IV (1905) 283. Senna.**St. Eustatius:** Seashore near English Quarter: n. 1010 B.; English Quarter: n. 1497, 1508; road to English Quarter: n. 1493; locality unknown: n. 158 G., 240 H.**Saba:** Bottom: n. 2236 B.; locality unknown: n. 91 L.

Baham., Antill., Bay of Honduras, Cartagena. (Symb.).

Tephrosia cinerea Pers. Syn. II (1807) 328; Gris. Fl. 182; Duss 195. Senna.**St. Eustatius:** Fort de Wind: n. 801 B.; White Wall: n. 812 B., 814 B., 821 B.; near Boven: 200 M.: n. 1243 B.; slope of the Quill: n. 1580.**St. Martin:** Point Blanche (coll. George): n. 2478 B.**St. Croix** (Symb.).

Baham., Antill., Curaçao, Amer. cont. trop. (Symb.).

3745. **Cracca** Benth.

Cracca caribaea Benth. ex Oerst. in Kjob. Vidensk. Medd. (1853) 9; Gris. Fl. 183; Duss 195.

Benthamantha caribaea O. Ktze. Rev. III 2 (1898) 53; Millspaugh 494.

St. Eustatius: along roads in Oranjestad: n. 920 B.; near Maryglory 150 M.: n. 1151 B.

St. Martin: Marigot road near Marigot: n. 2813 B.; lower part of Mount Paradis: n. 3290 B.; hills on the eastern side of Cul de Sac: n. 3444 B.; road near Fort Amsterdam: n. 1133 B.; Colebay-hill: n. 1143; Prince quarter: n. 1098; road to Guanabay: n. 1123.

St. Croix ex Millspaugh.

Antill., Amer. centr., Venezuela, Equador. (Symb.).

3747. **Sesbania** Scop.

Sesbania grandiflora Pers. Syn. II (1807) 316. Tiger tongue. Cultivated. [India orient., Austral. sept.]. (Symb.).

Sesbania sericea P. D.C. Prodr. II (1825) 266; Gris Fl. 184; Duss 196.

St. Martin: Colebay (coll. George): n. 2520 B.; near Simons-baylagoon: n. 3037 B.

I also saw *Sesbania sericea* near Marigot along a Saltpond.

Baham., Antill. (Symb.).

3802. **Stylosanthes** Sw.

Stylosanthes hamata Taubert Mon. Stylos. (Nov. 1889) 22; Millspaugh 495.

Stylosanthes procumbens Sw. Prodr. (1788) 108; Gris. Fl. 188; Duss 202. Wild clover.

St. Eustatius: Signalhill 200 M.: n. 130 B.; in a garden in Oranjestad: n. 755 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 758 B.; hills near Concordia 100 M.: n. 954 B.; lower part of Gilboohill: n. 1306; near Sugarloaf: n. 1310; English Quarter: n. 1386; locality unknown: n. 174 H., 179 H., 252 G., 300 G.

Saba: along Forthgut 150—200 M.: n. 1628 B.; along roads 200—300 M.: n. 1716 B.; seashore near Flatpoint: n. 2138 B.; Marypoint: n. 1328.

St. Martin: from Philipsburg to Guanabay: n. 2358 B.; Point Blanche (coll. George): n. 2490 B.; Cul de Sac—Marigot 0—200 M.:

n. 2623 B.; in an orchard near Bellevue: n. 2449a B.; between Mount Vernon and Orientbay: n. 2939 B.; Colebayhill: n. 1308; road to Guanabay: n. 1316.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Amer. sept., Mexico, Amer. centr., Nova Grenata. (Symb.).

3803. **Arachis** Linn.

Arachis hypogaea Linn. Spec. (1753) 741; Gris. Fl. 189; Duss 203; Millspaugh 495. Pea nut.

Saba: Mountain 600 M.; n. 1778 B.

St. Croix ex Millspaugh.

Cult. in tropical countries. Patria ignota. (Symb.).

3804. **Zornia** Gmel.

Zornia diphylla Pers. Syn. II (1807) 318; Gris. Fl. 185; Duss 198.

St. Eustatius: along roads near Bengalen: n. 1186 B.

St. Croix (Symb.).

Antill., warmer countries of both hemisph. (Symb.).

3807. **Desmodium** Desv.

Desmodium axillare P. DC. Prodr. II (1825) 334; Gris. Fl. 186; Duss 200.

Meibomia axillaris O. Ktze. Rev. I (1891) 195; Millspaugh 495.

Saba: between Bottom and Marypoint 100—200 M.: n. 1501 B.; Mountain 800 M.: n. 1807 B.; Mountain near Windwardside 600 M.: n. 2184 B.; Mountain 600—800 M.: n. 2011; Gainbygut: n. 2018; locality unknown: n. 201 L.

St. Croix ex Millspaugh.

Antill., Costarica, Venezuela, Nova Granata, Chili, Guiana, Brasil. (Symb.).

Desmodium scorpiurus Desv. Journ. Bot. I (1813) 122; Gris. Fl. 187; Duss 201.

Meibomia scorpiurus O. Ktze. Rev. I (1891) 198; Millspaugh 495.

St. Eustatius: Fairplay: n. 1346; seashore: n. 1345; Cultuur-vlakte: n. 1344.

St. Croix ex Millspaugh.

Antill., from Mexico to Peru. (Symb.).

Desmodium spirale P. DC. Prodr. II (1825) 332; Gris. Fl. 188; Duss 201;

Meibomia spiralis O. Ktze. Rev. I (1891) 197; Millspaugh 495.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1327 B.; Laddergut 0—200 M.: n. 1969 B.; locality unknown: n. 262 L.

St. Martin: hills on the eastern side of Cul de Sac: n. 3446 B., 3445a B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Afr. trop., Ind. or. and Ins. pacif. (Symb.).

Desmodium supinum P. DC. Prodr. II (1825) 332.

Desmodium incanum DC. Prodr. II (1825) 332; Gris. Fl. 186; Duss 200. Wild peas.

Meibomia incana Ktze. Rev. I (1891) 197; Millspaugh 495.

St. Eustatius: Top of the Quill 380 M.: n. 273 B., 278 B.; slope of the Quill near Bengalen 50—200 M.: n. 642 B., n. 675 B.; slope of the lower part of the Quill near Bengalen: n. 725 B.; along roads in Oranjestad: n. 1093 B.; near Maryglory 150 M.: n. 1142 B.; slope of the Quill: n. 1786, 1807, 1817; Cultuurvlakte: n. 1778; locality unknown: n. 329 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1281 B.; between Bottom and Marypoint: 100—200 M.: n. 1517 B.; top of the Cliff near the seaside of Boobyhill: n. 1605 B.; Mountain near Windwardside 600 M.: n. 2166 B.; Ladder: n. 1776, 1796; Rendez-vous 400 M.: n. 1806, 1848; Gainbygut: n. 1816; locality unknown: 25 L.

St. Martin: Nakedboyhill 50—200 M.: n. 2702 B.; along roads in Cul de Sac: n. 2998 B.; top of the hill on the northern side of Cul de Sac 200—300 M.: n. 3012 B.; Mount Paradis 150—400 M.: n. 3199 B., 3343 B.; Colebayhill: n. 1777; French frontier: n. 1787; road to Guanabay: n. 1797.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop., Afr. trop. (Symb.).

Desmodium tortuosum P. DC. Prodr. II (1825) 332; Gris. Fl. 188.

Saba: Ladder: n. 171, 181; Gainbygut 191.

Antill., Amer. cent. and austr. sept. (Symb.).

Desmodium triflorum P. DC. Prodr. II (1825) 334; Gris. Fl. 186; Duss 198. Sinternweed.

Meibomia triflora O. Ktze. Rev. I (1891) 197; Millspaugh 495.

St. Eustatius: locality unknown: n. 249 H.

St. Martin: Mountain between Mont des Accords and Centry-hill 200—300 M.: n. 2875 B.; Mount Paradis 150—400 M.: n. 3198 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., Afric. trop., Asia austr.-orient. (Symb.).

Desmodium uncinatum P. DC. Prodr. II (1825) 325. Gris. Fl. 187.

Saba: from Marypoint to Bottom 100—200 M.: n. 1504a B.

Cuba, Jamaica, Hispaniola. (Herb. Krug et Urban). Sonora to Bolivia, Venezuela to Uruguay. (Gris. Fl.).

3810. **Alysicarpus** Neck.

Alysicarpus nummularifolius P. DC. Prodr. II (1825) 353.

Alysicarpus vaginalis P. DC. Prodr. II (1825) 353; Gris. Fl. 185; Duss 199; Millspaugh 494. Sestern peanut.

St. Eustatius: locality unknown: n. 250 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1549 B., 1550 B.; seaside of Tentguthill 100 M.: n. 1653 B.

St. Martin: from Philipsburg to Guanabay: n. 2359 B.; lower part of Milldrumhill: n. 3115.

St. Croix ex Millspaugh.

Antill.; Patria Ind. or. (Symb.).

3834. **Lonchocarpus** H. B. K.

Lonchocarpus violaceus H. B. K. Nov. Gen. et Sp. VI (1823) 383; Gris. Fl. 200; Duss 219.

St. Eustatius: Top of the Quill 400 M.: n. 176 B., 345 B.; Quill: n. 45; slope of the Quill: n. 888 B.

Amer. austr. (Ind. Kew.), Antill. (Herb. Krug et Urban).

3839. **Piscidia** Linn.

Piscidia piscipula Sarg. in Gard. and For. IV. (1891) 436.

Piscidia erythrina Linn. Syst. X ed. II (1759) 1115; Gris. Fl. 200; Duss 220.

Ichthiomethia piscipula Hitchc. in Gard. and For. IV. (1891) 472; Millspaugh 495. Dogwood, Stinkwood.

St. Eustatius: from Tumbledown Dickbay to Signalhill: n. 164 B.; White Wall: n. 823 B.; slope of the hills near Concordiabay: n. 705; Seinpost: n. 715; Boven: n. 706, 725; locality unknown: n. 210a G., 314 H., 724.

St. Martin: Prince quarter (coll. George): n. 2477 B.; Fort Willem: n. 2544 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3424 B.

I also saw *Piscidia piscipula*: Marigot to Marigothill; Nakedboyhill; hill on the northern side of Cul de Sac; near Saline: one of the islands in Simonsbaylagoon.

St. Croix ex Millspaugh.

Florid. austr., Key ins., Baham., Antill., Mexico austr. and Amer. austro-sept. (Symb.).

3841. *Andira* Lam.

Andira jamaicensis Urb. in *Symbolae Antillanae* IV (1905) 298.

Andira inermis H. B. K. Nov. Gen. VI (1823) 385 in obs.; Gris Fl. 202; Duss 224.

St. Eustatius: Quill: n. 442 B., 906 B.; near Sigalhill: n. 1252a B.; locality unknown: n. 42.

Antill., Amer. cont. trop., Afr. occ. (Symb.).

3856. *Abrus* Linn.

Abrus precatorius Linn. Syst. XII ed. II (1767) 472; Gris. Fl. 190; Duss 204; Millspaugh 496. Jumby beans, Liquorice plant.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 55 B.; along roads in Oranjestad: n. 572 B.; slope of the Quill near Bengalen 250 M.; n. 702 B.; along roads: n. 26; slope of the Quill: n. 24; locality unknown: n. 23, 26 H., 235 G.

I also saw *Abrus precatorius*: in deserted plantations near Bengalen; Signalhill 100 M.

Saba: Bottom 200 M.: n. 1450.; Laddergut 200 M.: n. 2006 B.; locality unknown: n. 312 L.

St. Martin: Prince quarter (coll. George): n. 2459 B.; Marigot-hill—Marigot 0—200 M.: n. 2640 B.; along roads from Bethlehem to Mont Chambord: n. 2910 B.; Colebayhill: n. 25; locality unknown: n. 3619 B.

I also saw *Abrus precatorius* on Nakedboyhill; top of Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. calid., Africa, Asia calidior. (Symb.).

3857. *Clitoria* Linn.

Clitoria ternatea Linn. Spec. (1753) 753.

Cultivated. [Afric. orient.], (Symb.).

3858. **Centrosema** Benth.

Centrosema virginianum Benth. in Ann. Wien. Mus. II (1838) 120; Gris. Fl. 193; Duss 209.

Bradburya virginianum O. Ktze. Rev. 1 (1891) 164; Millspaugh 300.

St. Eustatius: Top of the Quill 380 M.: n. 265 B.; slope of the Quill near Bengalen 50--200 M.: n. 639 B., 661 B., 683 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 777 B.; gut near Fort de Wind 50 M.: n. 844 B.; slope of the Quill 200 M.: n. 878 B.; lower part of the hills near Concordia: n. 979 B.; along roads in Oranjestad: n. 1131 B.; along roads near Bengalen: n. 1193 B.; Cultuurvlakte: n. 151, 161; slope of the Quill: n. 211; locality unknown: n. 104 H., 236 G.

Saba: lower part of the Bottom Mountain 250 M.: n. 1414 B.; Bottom 200 M.: n. 1452 B., 152; locality unknown: n. 69 L.

I also saw *Centrosema virginianum* in Ladderhut.

St. Martin: from Philipsburg to Guanabay; n. 2387 B.; Point Blanche (coll. George): n. 2494 B.; from Philipsburg to Belvedere: n. 2574 B.; Nakedboyhill: 2682 B.; lower part of Milldrumhill: n. 3172 B.; gut from Bethlehem to Cul de Sac 0—100 M.: n. 3412 B.; Cul de Sac: n. 213; Centryhill: n. 162.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., Amer. cont. from the South U. S. to Argentina., Afr. trop. occ. (Symb.).

3866. **Teramnus** Sw.

Teramnus labialis Spreng. Syst. III (1826) 235; Gris. Fl. 193; Duss 209; Millspaugh 496.

St. Martin: in pastureground near Bellevue: n. 2824 B.; from Philipsburg to Belvedere: n. 2563 B.; locality unknown: n. 3500a B.

St. Croix ex Millspaugh.

Antill., Africa trop. to the Cape, East Indies. (Gris. Fl.).

3870. **Erythrina** Linn.

Erythrina corallodendron Linn. Spec. (1753) 706; Gris. Fl. 199; Duss 217; Millspaugh 496.

Saba: lower part of the Bottom Mountain 250 M.: n. 1411 B.; St. John's 250 M.: n. 1729 B.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2653 B.

St. Croix ex Millspaugh.

Bermud., Antill., Mexico, Amer. centr. and austro-sept. (Symb.).

3877. **Mucuna** Adans.**Mucuna pruriens** P. DC. Prodr. II (1825) 405.

Cultivated. [trop. countries of both hemisph.] (E. a. P.).

3882. **Galactia** Adans.**Galactia dubia** P. DC. Prodr. II (1825) p. 238.*Galactia filiformis* Griseb. Flor. W. I. (1860) 194 (p. p.). (Symb.); Duss 210 (p. p.).**St. Eustatius**: Top of the hills near Concordia 100 M.: n. 943 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 789 B.; Fort de Wind: n. 795a B.; White Wall: n. 818a B.; gut near Fort de Wind 50 M.: n. 827 B., 850 B.;**St. Martin**: near Mullet pond: n. 2779 B.; along roads in Cul de Sac: n. 2999 B.**Anguilla** Boldingh.

Portorico, St. Thomas, Guadeloupe, Antigua, Desirade, Marie-Galante. (Symb.).

Galactia longifolia Benth. in Ann. Wien. Mus. II (1838) 127.*Galactia angustifolia* Gris. Flor. W. I. (1860) 194; Duss 210; — non Kth. (Symb.).*Galactia tenuiflora* Wight et Arn. var. Benth. in Mart. Flor. Bras. XV. I (1859) 143. (Symb.); Millspaugh 495.**St. Eustatius**: Fairplay: n. 300; slope of the Quill: n. 301, 2755; Cultuurvlakte: n. 302; locality unknown: n. 303.**St. Croix** ex Millspaugh.

Antigua, Guadeloupe, Marie-Galante. (Symb.).

Galactia nummularia Urb. in Symb. Antillanae VI. (1909) 12.**St. Martin** near Guanabay: n. 2428 B.**Galactia rubra** Urb. in Symb. Antillanae, II. (1900) 309.*Galactia longiflora* Gris. Flor. W. I. (1860) p. 194 (p. p.); Duss 210. — non Arn. (Symb.).**St. Eustatius**: Slope of the Quill on the side of White Wall 200 M.: n. 1174 B.; Tree gut: n. 204 H., 282 G.**Saba**: between Bottom and Marypoint 350 M.: n. 1496 B.; slope of the Mountains near St. John's: n. 1737 B.; along roads 200—300 M.: n. 1715 B.; Hellsgate—Marypoint: n. 2042 B.; Ladder-gut: n. 2760.**St. Martin**: Top of the hill on the northern side of Cul de Sac 200—300 M.: n. 2994 B.; French frontier: n. 76, 78.

St. Kitts, Antigua, Guadeloupe, Dominica, Martinique. (Symb.).

3891. **Canavalia** P. DC.

Canavalia obtusifolia P. DC. Prodr. II (1825) 404; Gris. Flor. 197; Duss 215; Millspaugh 496. Sea peas.

St. Eustatius: Seashore near Oranjestad: n. 22 B., 33 B.; near Boven 200 M.: n. 1231 B.; Cultuurvlakte near Golden Rock: n. 381; locality unknown: n. 119 H.

I also saw *Canavalia obtusifolia* on the top of the Cliff near Gallowsbay.

Saba: I saw *Canavalia obtusifolia* near Flat Point.

St. Martin: Point Blanche (coll. George) n. 2489 B.: seashore near Mulletpond: 2767 B.

St. Croix ex Millspaugh.

Florida, Bermud., Baham., Antill., warmer countries of both hemisph. (Symb.).

Canavalia spec.

St. Eustatius: n. 244 G.

3892. **Cajanus** P. DC.

Cajanus indicus Spreng. Syst. III (1826) 248; Gris. Fl. 191; Duss 205.

Cajanus cajan Millsp. in Field. Col. Mus. Bot. II (1900) 53; Millspaugh 497.

St. Eustatius: locality unknown: n. 55 G., 210 H.

Saba: Hellsgate: n. 1696 B.; locality unknown: n. 16 L.

I also saw *Cajanus indicus*: Laddergut.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Warmer countries of both. hemisph. (Symb.): Also cultivated.

3897. **Rhynchosia** Lour.

Rhynchosia minima P. DC. Prodr. II (1825) 385; Gris. Fl. 190; Duss 205.

Dolicholus minimus Medic. Vorles. Churpf. Phys. Ges. II (1787) 354; Millspaugh 496.

St. Eustatius: Top of the Quill 450 M.: n. 354 B.; slope of the Quill: n. 1429, 1449; Cultuurvlakte: n. 1439.

Saba: between Bottom and Marypoint 100—200 M.: n. 1504 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Warmer countries of both hemisph. (Symb.).

Rhynchosia reticulata P. DC. Prodr. II (1825) 385; Gris. Fl. (1859) 190.

Dolicholus reticulatus Millsp. in Field Col. Mus. Bot. II (1900) 53; Millspaugh 496. Pea with e.

St. Eustatius: Top of the Quill 400 M.: n. 215 B., 276 B.; slope of the Quill near Bengalen 50 M.: n. 660 B.; hills near Concordia 100 M.: n. 944 B.; along roads near English Quarter: n. 1005 B.; Red Cliff—Tumbledown dickbay: n. 1549; slope of the Quill: n. 1529, 1552, 1539.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1346 B.; Crispeen 400 M.: n. 1718 B.; Laddergut 200 M.: 1972 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2027 B.; Windwardside: n. 2311 B.; Ladder: n. 1522, 1561; Paris 2—400 M.: n. 1542; Mountain 400—800 M.: n. 1542.

St. Martin: Prince quarter (coll. George): n. 2470 B.; Cul de Sac—Marigothill 0—200 M.: n. 2634 B.; Nakedboyhill 50—200 M.: n. 2693 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2870 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2947 B.; lower part of the Mill-drumhill: n. 3173 B.; hills on the eastern side of Cul de sac: n. 3439 B.; French frontier: n. 1512.

St. Croix ex Millspaugh.

Antill., Amer. austr. (Symb.).

3901. *Phaseolus* Linn.

Phaseolus lathyroides Linn. Spec. (1763) 1018.

Phaseolus semirectus Linn. Mant. I (1767) 100; Gris. Fl. 197; Duss 214; Millspaugh 497.

St. Eustatius: in a garden in Oranjestad: n. 1101 B.; Fairplay: n. 590; Cultuurvlakte: n. 472, 580.

St. Croix ex Millspaugh.

Baham., Antill., Curaçao, Amer. centr. et austr., India orient., Malaya. (Symb.).

Phaseolus lunatus Linn. Spec. (1753) 724. White beans. Cultivated. [All. trop. countries.] (Symb.).

Phaseolus vulgaris Linn. Spec. (1753) 723. Snijboontjes. Cultivated. [Everywhere.] (Ind. Kew.).

3905. *Vigna* Savi.

Vigna repens O. Ktze. Rev. I (1891) 212; Millspaugh 497.

Vigna luteola Benth. in Mart. Flor. Bras XV. I (1859) 194 t. 50 f. II; Gris. Fl. 195; Duss 211.

Vigna glabra Savi. Mem. Phas. III (1882) 8.

St. Eustatius: Slope of the lower part of the Quill near Bengalen: n. 716 B.; locality unknown: n. 1 G., 66 H.; slope of the Quill: n. 1548, 1549, 2005, 2015.

St. Croix ex Millspaugh.

Bermud., Antill., Amer. calid. from U. S. to Argentin., Afr. trop., Asia trop., Australia. (Symb.).

3909. **Lablab** Adans.

Lablab vulgaris Savi Diss. (1821) 19 t. 19 f. 8.

Dolichos lablab Linn. Spec. (1753) 725; Gris. Fl. 196; Duss 212; Millspaugh 497.

St. Eustatius: Cultuurvlakte: n. 1891; slope of the Quill: n. 1937.

Saba: Windwardside: n. 1887 B.; Bottom: n. 2240 B.; Crispeen-Rendez-vous: n. 1927; locality unknown: n. 167 L., 304 L.

St. Croix ex Millspaugh.

Trop. countries of both. hemisph.; Ind. veris. in Afr. trop. (Symb.). Also cultivated.

130. **Oxalidaceae.**

3936. **Oxalis** Linn.

Oxalis corniculata Linn. Spec. (1753) 435; Gris. Fl. 133; Duss 134; Millspaugh 497.

St. Eustatius: Slope of the Quill near Glassbottle: n. 1206 B.; slope of the Quill: n. 131, 132.

Saba: lower part of the Bottom Mountain 250 M.: n. 1400 B.; between Bottom and Marypoint 100—200 M.: n. 1515 B.; Booby-hill 300 M.: n. 1620 B.; locality unknown: n. 50 L.; Crespeen—Rendez-vous: n. 130; St. John's: n. 134; Hellsgate: n. 133.

St. Martin: Mountain between Mont des Accords and Centry-hill 0—200 M.: n. 2868 B.; hill on the northern side of Cul de Sac 0—100 M.: n. 3001 B.

I also saw *Oxalis corniculata* on the top of Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Warmer countries of both hemisph. (Symb.).

Oxalis Martiana Zucc. in Denkschr. Akad. Münch. IX (1823—24) 144.

Cultivated. [Madagascar]. (Ind. Kew.).

134. **Erythroxylaceae.**3956. **Erythroxylum** Linn.**Erythroxylum brevipes** DC. Prodr. I (1824) 573; Millspaugh 497.**St. Martin:** Colebayhill: n. 2014; Princee quarter: n. 2018; John Ednyhill: n. 2015.**St. Croix** (Symb.).

Cuba, Hispaniola, St. Thomas, St. Barthélemy, Guadeloupe. (Symb.).

Erythroxylum havanense Jacq. Enum. Pl. Carib. (1760) 21.*Erythroxylum obtusum* DC. Prodr. I (1824) 574; Gris. Fl. 113. p. p. (Symb.). Bracelet.**St. Eustatius:** Top of the Quill 400 M.: n. 217 B., 329 B.; 500 M.: n. 495 B.; from Little Mountains to Venusbay: n. 958 B.**Saba:** between Bottom and Marypoint 300 M.: n. 1478 B.; Springbaygut: n. 2118 B.; Laddergut: n. 2017.**St. Martin:** Milldrumhill 300—400 M.: n. 3155 B.; Mount Paradis 200—300 M.: n. 3314 B.

Cuba, Isla de Pinos. (Symb.).

135. **Zygophyllaceae.**3968. **Guaiacum** Linn.**Guaiacum officinale** Linn. Spec. (1753) 381; Gris. Fl. 134; Duss 136; Millspaugh 498.**St. Martin:** Tintamarre: n. 3370 B., 3378 B.**St. Croix** ex Millspaugh.

Baham., Antill., Amer. austr. (Symb.).

3978. **Tribulus** Linn.**Tribulus cistoides** Linn. Spec. (1753) 387; Gris Fl. 134; Duss 136; Millspaugh 498.**St. Martin:** near Filipsburg: n. 2388 B.; Fort Willem: n. 2548 B.**St. Croix** ex Millspaugh.

Florida austr., Baham., Antill., trop. countries of both hemisph. (Symb.).

3979. **Kallstroemia** Scop.**Kallstroemia maxima** Wight et Arn. Prodr. I (1834) 145; Millspaugh 498.*Tribulus maximus* Linn. Spec. (1753) 386; Gris. Fl. 134; Duss 136.

St. Eustatius: in a garden in Oranjestad: n. 511 B.; Bengalen: n. 616 B.; Cultuurvlakte: n. 243, 248, 249; locality unknown: n. 30 G.

Saba: Windwardside: n. 1951 B.; Bottom: n. 247.

St. Martin: from Guanabay to Philipsburg: n. 2437 B.; along roads near Philipsburg: n. 2530 B.; Centryhill: n. 246; Marigot-road: n. 244, 245.

St. Croix ex Millspaugh.

Key Ins., Baham., Antill., Amer. from the Southern U. S. to Bolivia. (Symb.).

137. Rutaceae.

3991. **Fagara** Linn.

Fagara flava Kr. et Urb. in Engl. Jahrb. XXI (1896) 571.

Xanthoxylum flavum Vahl. Eclog. III (1807) 48; Duss 140; Millspaugh 499. Yellow saunder.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Key Ins., Baham., Antill. (Symb.).

Fagara martinicensis Lam. Ill. I (1791) 334 n. 1639 t. 811 f. 1.

Xanthoxylum clara Herculis Sw. Obs. (1791) 375 (non Linn.); Gris. Fl. 138; Millspaugh 499.

Xanthoxylum martinicense P. DC. Prodr. I (1824) 726; Duss 141. Yellow prickle.

St. Eustatius: Quill: n. 190 B., 416a B., 1.

Saba: Windwardside: n. 1942 B.; Mountain near Hellsgate: n. 2279 B.

St. Martin: along roads in Cul de Sac: n. 2989a B.

St. Croix ex Millspaugh.

Cuba, Jamaica, Hispaniola, Portorico, St. Thomas, St. Jan, Tortola, Montserrat, Guadeloupe, Desirade, Martinique, St. Lucia, Trinidad. (Symb.).

Fagara spinifex Jacq Fragm. (1809) p. 10 t. 6 f. 2.

Fagara microphylla Desv. Tabl. I ed. (1804) 200; Gris. Fl. 137 (excl. syn. -Br. et L.). (Symb.); Duss 139; Millspaugh 499.

Xanthoxylum spinifex P. DC. Prodr. I (1824) 728.

St. Martin: between Mount Vernon and Orientbay: n. 2928 B.; from Belvedere to Oysterpond: n. 3062 B.; Mount Paradis 100—200 M.: n. 3296 B.; gut from Bethlehem to Cul de Sac 100—200 M.: n. 3425 B.; Prince quarter: n. 2, 3.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Venezuela. (Symb.).

Fagara trifoliata Sw. Prodr. (1788) 33.

Tobinia ternata Desv. in Ham. Prodr. (1825) 57; Gris. Fl. 136.

Tobinia punctata Gris. Flor. W. I. (1859) 137; Duss 138; Millspaugh 499.

Xanthoxylum ternatum Sw. Flor. I (1797) 570.

St. Eustatius: Top of the Quill 400 M.: n. 204 B.

St. Martin: from Philipsburg to Guanabay: n. 2386 B.; Point Blanche (coll. George): n. 2492 B.; Nakedboyhill 50—200 M.: n. 2708 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2953 B.; from Belvedere to Oysterpond: n. 3056 B., 3074 B., 3081 B., 3082 B.; Prince quarter: n. 4, 10, 11, 12, 13; Centryhill: n. 15.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Jamaica, Hispaniola, Portorico, St. Barthélemy, Antigua, Montserrat, Guadeloupe, Marie-Galante, Desirade, Les Saintes, Dominica, Martinique, St. Lucia, St. Vincent, Barbados, Trinidad. (Symb.).

4084. **Amyris** Linn.

Amyris elemifera Linn. Syst. X ed II (1759) 1000; Duss 183.

St. Martin: Tintamarre: n. 3391 B., 3395 B.

St. Croix (Symb.).

Baham., Florida austr., Key West, Antill. (Symb.).

4090. **Murraya** Linn.

Murraya exotica Linn. Mant. II (1771) 563; Duss 132; Millspaugh 499. Boxwood.

St. Eustatius: in a garden in Oranjestad: n. 937 B., 40 G.; Little Mountain—Boven: n. 14.

Saba: Windwardside: n. 1908 B.; locality unknown: n. 281 L.

St. Martin: Hills on the eastern side of Cul de Sac: n. 3446a B.

St. Croix ex Millspaugh.

Antill.; Ind. in Asia austro-orient., Austral. sept., Polynes. (Symb.).

4094. **Triphasia** Lour.

Triphasia trifoliata P. DC. Prodr. I (1824) 536; Duss 132; Millspaugh 499. Myrtle lemon.

Triphasia aurantiola Lour. Fl. Cochinchin. (1790) 153.

St. Eustatius: in a garden in Oranjestad: n. 538 B.; Bengalen: n. 613 B.; seashore: n. 24, 26; locality unknown: n. 226 G.

Saba: Bottom: n. 1856 B.; locality unknown: n. 97 L.

St. Martin: Cul de Sac—Marigot 0—200 M.: n. 2622 B.; hills on the eastern side of Cul de Sac: n. 3447 B.

St. Croix ex Millspaugh.

Antill.; Indig. in India orient. (Symb.). Also cultivated.

4100. **Citrus** Linn.

The author was not yet able to find out with a security so great, to make it possible to mention it in this Flora the names of the different species of Citrus that are cultivated in the gardens.

Only one plantation of Citrus Limonum Risso in Saba be mentioned.

138. **Simarubaceae.**

4106. **Suriana** Linn.

Suriana maritima Linn. Spec. (1753) 284; Gris. Fl. 58; Duss 48; Millspaugh 499.

St. Eustatius: Seashore near Concordiabay: n. 36; near Sea-feathersbay: n. 35;

I also saw *Suriana maritima* at the seashore near English Quarter.

St. Martin: Colebay (coll. George): n. 1643 B; from Belvedere to Oysterpond: n. 3043 B.

I also saw *Suriana maritima* near Mulletpond.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., trop. countries of both hemisph. (Symb.).

4121. **Picrasma** Blume.

Picrasma antillana Urb. in Symbolae Antillanae V (1908) 378.

Picraena excelsa Gris. in Flor. W. I. (1859) 140 (p. p.); Duss 143; (Symb.).

St. Eustatius: Quill: n. 448 B., 461a B.; locality unknown: n. 210a G.

Saba: from Bottom to Marypoint 100—200 M.: n. 1465a B.; near Hellsgate: n. 1697a B.; locality unknown: n. 2073a B., 341 L.; Cresp.—Rendez-vous 400 M.: n. 44.

St. Martin: Mount Paradis: n. 3219a B.

St. Croix (Symb.).

St. Thomas, St. Jan, St. Kitts, Antigua, Guadeloupe, Martinique, St. Vincent, Barbados. (Symb.).

4131. **Picramnia** Sw.

Picramnia pentandra Sw. Flor. 1 (1797) 220; Gris. Fl. 140; Duss 142. Wild coffee, Bastard bough.

St. Martin: Marigot—Marigot 0—200 M.: n. 2639 B.; Naked-boyhill 50—200 M.: n. 2690 B., 2697 B., 2714 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2878 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2956 B., 2962 B.; top of the hill on the northern side of Cul de Sac 200—300 M.: n. 3010 B.; Milldrumhill 300—400 M.: n. 3184 B.; Mount Paradis 150—400 M.: n. 3197 B., 3299 B., 3345 B.; Centryhill: n. 92, 93; Prince quarter: n. 90; French frontier: n. 37, 38, 39; locality unknown: n. 3621 B., 91.

I also saw *Picramnia pentandra*: from Belvedere to Oysterpond.

Florida austr., Key Ins., Baham., Antill. (Symb.).

139. **Burseraceae.**4150. **Bursera** Linn.

Bursera simaruba Sarg. Gard. and For. III (1890) 260.

Bursera gummiifera Jacq. Sel. (1763.) 94 t. 65; Gris. Fl. 173; Duss 181; Millspaugh 500. Gum tree, Balsam tree, Lime tree.

St. Eustatius: near Tumbledowndiekbay: n. 139a B., 146a B.; top of the Quill 400 M.: n. 202 B.; slope of the Quill on the side of White Wall 200 M.: n. 1182 B.; locality unknown: n. 106 H., 200 G.

I also saw *Bursera simaruba* in a gut near Fort de Wind.

Saba: Seashore near Saddle: n. 1652a B.; gut near Devils-hand: n. 2078a B.; locality unknown: n. 290 L.

I also saw *Bursera simaruba*: in Hellsgate, Tentgut, Springbaygut.

St. Martin: Marigotroad: n. 2819a B.; French frontier: n. 71, 72.

I also saw *Bursera simaruba*: Top of Nakedboyhill; near Saline; along roads from Philipsburg to Simonsbay; Vallée des Pères; Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida, Key Ins., Baham., Antill., Curaçao, from Mexico to Nova Granata and Venezuela. (Symb.).

140. **Meliaceae.**4164. **Swietenia** Jacq.

Swietenia mahagoni Jacq. Enum. (1760) 20; Gris. Fl. 131; Duss 130; Millspaugh 500. *M a h a g o n y*.

St. Eustatius: Bottom of the Quill 300 M.: n. 216 B.; in a garden in Oranjestad: n. 1072 B.; locality unknown: n. 82 H.

Saba: Ladder gut 0—200 M.: n. 1978 B.; Greathill: n. 92; locality unknown: n. 12 L.

I also saw *Swietenia mahagoni*: near Marypoint.

St. Croix ex Millspaugh.

Key Ins., Baham., Antill., Mexico, Amer. centr., Peru. (Symb.).

Also cultivated.

4175. **Melia** Linn.

Melia azedarach Linn. Spec. (1753) 384; Millspaugh 500.

Melia sempervirens Sw. Prodr. (1788) 67; Gris. Fl. 128; Duss 126. *L i l a c*.

St. Eustatius: in a garden in Oranjestad: n. 44 H., 139 G.

I also saw *Melia azedarach* in the slope of the Cliff.

Saba: Crispeen 400 M.: n. 1722 B.; Mountain near Windward-side 600 M.: n. 2163 B.; Peperpot: n. 93; Crispeen—Rendez-vous 400 M.: n. 92; locality unknown: n. 125 L.

St. Martin: near Bethlehem: n. 2661 B.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., warmer countries of both hemisph. (Symb.). Also cultivated.

141. **Malpighiaceae.**4226. **Heteropteryx** Juss.

Heteropteryx purpurea H. B. K. Nov. Gen. V (1821) 163; Duss 116; Millspaugh 498.

St. Eustatius: Hills near Concordia 100 M.: n. 950 B.; from Little Mountains to Venusbay: n. 990a B.; slope of the Quill on the side of White Wall 200 M.: n. 1179 B.; Pongo—Signalhill: n. 1965; Red Cliff: n. 1969; Sugarloaf: n. 1966, 1967, 1968.

St. Croix ex Millspaugh.

Antill., Venezuela, Colombia. (Symb.).

4228. **Stigmatophyllon** Juss.

Stigmatophyllon ciliatum A. Juss. in A. St. Hil. Fl. Bras. Mer. III (1827) 49.

Cultivated. [Jamaica, Brasil.], (Gris. Fl.).

Stigmatophyllon periplocifolium A. Juss. Malp. Syn. (1840) 290; Gris. Fl. 119; Duss 116; Millspaugh 498. Goat bush.

St. Eustatius: Seashore near Oranjestad: n. 31 B.; in a garden in Oranjestad: n. 96 B.; deserted plantation near Tumbledown-dick-bay: n. 142 B., 171 B.; top of the Quill 400 M.: n. 191 B., 212 B.; slope of the Quill near Bengalen 150 M.: n. 686 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 791 B.; along roads in Oranjestad: n. 922 B.; lower part of the hills near Concordia: n. 980 B.; Signalhill 150 M.: n. 1042 B., 1052 B.; along roads in Oranjestad: n. 1129 B.; Cultuurvlakte: n. 608, 609, 619, 630; locality unknown: n. 137 G., 144 H.

St. Martin: from Filipsburg to Guanabay: n. 2407 B.; from Filipsburg to Belvedere: n. 2557 B.; Cul de Sac—Marigothill 0—200 M.: n. 2629 B.; lower part of the Milldrumhill: n. 3171 B.; Tintamarre: 3398; Vineyardhill: n. 618; Cul de Sac: n. 639; Colebayhill: n. 94.

I also saw *Stigmatophyllon periplocifolium* on: Marigothill—Marigot; Nakedboyhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Cuba, Jamaica, Hispaniola, Portorico, St. Thomas, St. Jan, Antigua, Martinique, St. Lucia. (Symb.).

Stigmatophyllum sericeum Wright ex. Gris. Cat. Pl. Cub. (1866) 43.

Stigmatophyllum diversifolium A. Juss. in Arch. Mus. Par. III (1843) 381.

St. Martin: Lowlands: n. 3286 B.

Anguilla Boldingh.

Cuba (Herb. Krug et Urban).

4244. **Galphimia** Cav.

Galphimia gracilis Bartl. in Linnaea XIII (1839) 552.

Cultivated. [Mexico.], (Symb.).

4251. **Malpighia** Linn.

Malpighia angustifolia Linn. Spec. (1762) 610; Gris. Fl. 117; Duss 113. Bastard cherry.

St. Martin: lower part of the hills between Mont Chambord and Mount Vernon: n. 2920 B.; along roads from Grande Case to Marigot: n. 3090 B.

I also saw *Malpighia angustifolia* near Saline.

St. Thomas, St. Jan, Antigua, Guadeloupe, St. Lucia. (Niedenau 1899).

Malpighia puniceifolia Linn. Spec. (1762) 609; Gris. Fl. 116; Duss 113. Cherry.

St. Eustatius: Seashore near Oranjestad: n. 28 B.; slope of the lower part of Signalhill: n. 107 B.; Oranjestad: n. 97; Gilboohill: n. 38; along the road to Sugarloaf: n. 87; locality unknown: n. 63 H., 78 G.

I also saw *Malpighia puniceifolia*: top of Signalhill: top of the Cliff between Oranjestad and White Wall.

Saba: between Bottom and Marypoint 100—200 M.: n. 1546 B.; locality unknown: n. 1477 B., 334 L.

St. Martin: from Guanabay to Philipsburg: n. 2435 B.; Marigotroad near Marigot: n. 2809 B., 2810 B.; Prince quarter: n. 47, 57, 58.

Anguilla Boldingh.

St. Croix (Symb.).

St. Thomas, Antigua, St. Barthélemy, Guadeloupe, Martinique, Trinidad, Margarita, Bonaire, Curaçao. (Symb.).

4252. **Bunchosia** L. Cl. Rich.

Bunchosia jamaicensis Urb. et Ndz. Index Lect. in Lyc. Reg. Hos. Brunsb. de Genere *Bunchosia* (1898) 10.

Bunchosia Lindeniana Gris. Flor. W. I. (1860) 115.

Saba: Seaside of Tentguthill 100 M.: n. 1655 B.; Windward-side: n. 1943 B.

Antill., Mexico, Costarica, Guiana. (Gris. Fl.).

Bunchosia nitida L. Cl. Rich. ap. Juss. in Ann. Mus. Par. XVIII (1811) 481.

St. Martin: between Mont Chambord and the Saline: n. 2924 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2963 B.; near Simonsbayvillage: n. 3257 B.; Mount Paradis 300—400 M.: n. 3322 B.

Hispaniola, Montserrat, Guadeloupe, Martinique, St. Vincent, Bequia, Barbados, Grenada. (Symb.). Surinam. (Pulle).

4255. **Byrsonima** L. Cl. Rich.

Byrsonima crassifolia H. B. K. Nov. Gen. V (1821) 149; Gris. Fl. 114. (vel. *B. martinicensis* Kr. et Urb.?)

St. Martin: Lowlands: n. 2381a B.

Antill., Amer. austr. sept. (Symb.).

Byrsonima lucida L. Cl. Rich. ap. Juss. in Ann. Mus. Par. XVIII (1811) 481; Gris. Fl. 115; Duss 111.

St. Martin: Lowlands: n. 3621 B.

Anguilla Boldingh.

Key Ins., Baham., Antill. (Symb.).

Byrsonima spicata L. Cl. Rich. ap. Juss. in Ann. Mus. Par. vol. XVIII (1811) 481; Gris. Fl. 115; Duss 111; Millspaugh 497. Holly hock.

St. Eustatius: Top of the Quill 380 M.: n. 253 B.; inner slope of the Quill 350 M.: n. 1229 B.; Quill: n. 80 H.

Saba: Windwardside—Hellgate: n. 1679 B.; Windwardside: n. 1944 B.

St. Croix ex Millspaugh.

Antill., Panama, Venezuela, Guiana, Brasilia. (Symb.).

147. Euphorbiaceae.4299. **Phyllanthus** Linn.

Phyllanthus distichus Müll. Arg. in DC. Prodr. XV. 2 (1866) 413; Millspaugh 500.

Cicca disticha Linn. Mant. I (1767) 124; Gris. Fl. 32; Duss 20.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 52 B.; in a garden in Oranjestad: n. 90 H., 148 G.

Saba: Bottom: n. 1876 B.; locality unknown: n. 116 L., 238 L.

St. Martin: Top of the hill on the northern side of Cul de Sac 200—300 M.: n. 2992 B.

St. Croix ex Millspaugh.

Antill.; Indig. in India orient? (Symb.). Also cultivated.

Phyllanthus epiphyllanthus Linn. Spec. (1763) 1392; Duss 25.

Phyllanthus falcatus Sw. Flor. II (1800) 1115; Gris. Fl. 35.

St. Martin: Colebay (coll. George): n. 2502 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2744 B.; near Mulletpond: n. 2785 B.

I also saw *Phyllanthus epiphyllanthus*: Lowlands.

Anguilla Boldingh.

Baham., Antill. (Symb.).

Phyllanthus niruri Linn. Spec. (1753) 981; Gris. Fl. 34; Duss 23; Millspaugh 501. Surinaam bitter.

St. Eustatius: in a garden in Oranjestad: n. 76 B.; along roads in Oranjestad: n. 579 B.; along roads near Glassbottle: n. 871 B.; locality unknown: n. 41 G., 34 H.

Saba: lower part of the Bottom Mountain 300 M.: n. 1429 B.;

St. Martin: from Philipsburg to Guanabay: n. 2408 B.; lower part of the Milldrumhill: n. 3166 B.; in a plantation of Cotton near Bethlehem: n. 3410 B.

I also saw *Phyllanthus niruri* in pastureground near Bellevue.

St. Croix ex Millspaugh.

Bermud., Baham., Antill., almost all warmer countries of both hemisph. (Symb.).

Phyllanthus nobilis Müll. Arg. in DC. Prodr. XV 2 (1866) 114; Millspaugh 501

Cicca antillana A. Juss. Tent. Euphorb. (1824) 108 t. 4 f. 13 B.; Gris. Fl. 33; Duss 21. Gooseberry.

Saba: Gainbygut: n. 2074; locality unknown: 43 L.

St. Martin: Nakedboyhill: n. 2735a B.; Mount Paradis: n. 3202a B.

St. Croix ex Millspaugh.

Antill., Guiana. (Symb.).

4348. **Croton** Linn.

Croton astroites Dryand. in Ait. Hort. Kew. 1 ed. vol. III (1789) 375; Gris. Fl. 42; Duss 33; Millspaugh 501. Wild marrow.

St. Eustatius: Slope of the Quill above White Wall: n. 2 B.; slope of the lower part of White Wall: n. 97 B.; Signalhill 200 M.: n. 128 B.; gut near Fort de Wind 50 M.: n. 826 B.; along a road in Oranjestad: n. 930 B.; hill near Concordia 100 M.: n. 949 B.; top of Signalhill 225 M.: n. 1030 B.; Signalhill 200 M.: n. 1047 B.; near Maryglory 150 M.: n. 1147 B.; along roads near Bengalen: n. 1194 B.; near Boven 200 M.: n. 1239a B.; Cultuurvlakte: n. 125; Pangohill: n. 122, 123; Gilboohill: n. 124; Sugarloaf: n. 128; locality unknown: n. 150a B., 269 B., 171 G., 3577 B.

St. Martin: from Philipsburg to Guanabay: n. 2362 B.; Fort Willem 50 M.: n. 2536 B.; locality unknown: n. 3611 B.; near Fort Amsterdam: n. 126, 127.

I also saw *Croton astroites*: Cul de Sac; Nakedboyhill

St. Croix ex Millspaugh.

Portorico, St. Thomas, St. Jan, St. Barthélemy, St. Kitts, Antigua, Guadeloupe, Desirade. (Symb.).

Croton betulinus Vahl Symb. II (1791) 98; Duss 32; Millspaugh 501. Black sage.

St. Martin: Marigotroad near Marigot: n. 2802 B.; in pasture-ground near Bellevue: n. 2826 B.; Mountain between Mont des Accords and Centryhill: n. 2865 B.; along roads near Colebay: n. 3025 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2739 B., 2743 B., 2746 B.; Lowlands: n. 3265 B.; Tintamarre: n. 3384 B.

I also saw *Croton betulinus*: one of the islands in Simonsbaylagoon.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida austr., Antill. (Symb.).

Croton flavens Linn. Syst. X ed. II (1759) 1276; Gris. Fl. 38; (excl. syn. Willd.). (Symb.); Millspaugh 501. Marrow, Yellow balsam.

Croton balsamifer Jacq. Enum. Pl. Carib. (1760); Duss 31.

St. Eustatius: on the slope of the Cliff near Gallowsbay: n. 61 B.; in deserted plantations at the seashore near Bengalen: n. 582 B., 584 B., 594 B., 597 B., 599 B.; slope of the Quill near Bengalen 200 M.: n. 737 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 775 B., 778 B., 785 B., 792 B.; Fort de Wind: n. 795 B.; seashore near Venusbay: n. 968 B.; Gilboohill 150 M.: n. 1124 B.; along roads near Bengalen: n. 1195 B.; slope of the hills near Concordiabay: n. 131; Cultuurvlakte: n. 130, 132; Tumbledown Dickbay: n. 133; Vennsbay: n. 134; Sugarloaf: n. 140; locality unknown: n. 54 H., 120 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1321 B.; along the Forthgut 150—200 M.: n. 1627 B.; along the lower part of the Forthgut: n. 1643a B.; St. Giles: n. 1865 B.; Laddergut 0—200 M.: n. 1971 B.; Ladderlanding: n. 135, 136; Greathill 200 M.: n. 141; Quarter 300 M.: n. 249; Marypoint: n. 250; locality unknown: n. 315 L.

St. Martin: from Philipsburg to Guanabay: n. 2351 B.; Guanabay: n. 2431 B.; Prince quarter (coll. George): n. 2469 B.; Fort Willem 0—50 M.: n. 2531 B., 2543 B.; near Bethlehem: n. 2662 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2741 B.; road near Fort Willem: n. 2796 B.; between Mount Vernon and Orientbay: n. 2937 B.; near Simonsbayvillage: n. 3256 B.; Green Key: n. 3364 B., 3867 B.; Tintamarre: n. 3382 B.; Philipsburg: n. 137; Centryhill: n. 138; Vineyardhill: n. 139; locality unknown: n. 37.

I also saw *Croton flavens*: from Belvedere to Oysterpond; between Mount Vernon and Saline; Nakedboyhill.

Anguilla: Boldingh.

St. Croix ex Millspaugh.

Hispaniola, Portorico, St. Thomas, St. Jan. (Symb.). Baham. (Herb. Leiden).

Croton lobatus Linn. Spec. (1753) 1005; Gris. Fl. 42; Duss 33; Millspaugh 501. Lila c bush.

St. Eustatius: in a garden in Oranjestad: n. 74 B., 1276 B.; along roads in Oranjestad: n. 564 B.; Fairplay: n. 167; slope of the Quill: n. 164, 165, 166; locality unknown: n. 92 G., 177 H.

Saba: Bottom: n. 1839 B., 1843 B.

St. Martin: Colebayhill: n. 161.

I also saw *Croton lobatus* near Belvedere.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Bonaire, Curaçao, Aruba, Amer. cont. trop. (Symb.).

4358. *Ditaxis* Vahl.

Ditaxis fasciculata Vahl ap. A. Juss. Tent. Euph. (1824) p. 110 t. 7 f. 24; Gris. Fl. 44; Millspaugh 501.

St. Martin: Prince quarter: (coll. George) n. 2457 B.; near Nakedboyhill: n. 2566 B., 2733a B.; Point Blanche: n. 3405 B.; Vineyardhill: n. 769; locality unknown: n. 21.

St. Croix ex Millspaugh.

Cuba, Portorico, St. Thomas, St. Jan. (Symb.).

4360. *Argithamnia* Sw.

Argithamnia candicans Sw. Prodr. (1788) 39; Gris. Fl. 44; Millspaugh 502.

St. Martin: Fort Willem 0—50 M.: n. 2540 B.; Nakedboyhill 50—200 M.: n. 2700 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2959 B.; Mount Paradis 200—300 M.: n. 3317 B.; Tintamarre: n. 3401 B.; Colebayhill: n. 2075; Centryhill: n. 767.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., (Hitchc.). Antill.

4374. **Bernardia** Adams.

Bernardia corensis Müll—Arg. in Linnaea XXXIV. (1865, 66) 173; Duss 35.

Polyboea corensis Klotzsch. in End. Gen. Suppl. IV. III (1850) 88. Gris. Fl. 46.

St. Eustatius: Top of the Quill 400 M.: n. 195 B., 226 B., 398 B., 404 B.; slope of the Quill near Bengalen 100 M.: n. 665 B.; Signalhill 200 M.: n. 1050 B.

St. Martin: Mount Paradis 300—400 M.: n. 3224 B.; Mount Paradis 200—300 M.: n. 3303 B.

Haiti, St. Barthélemy, Martinique, Dominica, Guadeloupe, Grenada, St. Vincent, Curaçao. (Herb. Krug et Urban).

4397. **Adelia** Linn.

Adelia ricinella Linn. Syst. X. ed. II (1759) 1298; Gris. Fl. 45; Millspaugh 502.

St. Martin: Philipsburg near the Saltponds: n. 2357 B.; from Philipsburg to Fort Amsterdam: n. 2566 B.; Prince quarter (coll. George): n. 2457 B.; top of Nakedboyhill: n. 2711 B.; Point Blanche: n. 3405 B.; near Fort Amsterdam: n. 750; Colebayhill: n. 751; Prince quarter: n. 752.

St. Croix ex Millspanch.

Cuba, Jamaica, Hispaniola, Portorico, St. Thomas, St. Jan. (Symb.).

4407. **Acalypha** Linn.

Acalypha chamaedrifolia Müll. Arg. in DC. Prodr. XV. 2. (1866) 879; Duss 35.

Acalypha reptans Sw. Prodr. (1788) 99; Gris. Fl. 48; Millspaugh 502. Bastard nettle.

St. Eustatius: along roads in Oranjestad: n. 552 B., 921 B.

Saba: between Bottom and Marypoint 100—200 M.: n. 1519a B., 1520 B., 1540 B.; Hellsgate: n. 1692 B.; Laddergut 0—200 M.: n. 1962 B.; Windwardside: n. 2334 B.

St. Martin: in an orchard near Bellevue: n. 2850 B.; from Grande Case to Marigot: n. 3092 B.; lower part of Milldrums: n. 3159 B.

St. Croix ex Millspaugh.

Florida austral., Antill. (Symb.).

Acalypha Poirerii Spreng Syst. III (1826) 879.

St. Martin: near Grande Case: n. 3092 B.

Anguilla Boldingh.

St. Thomas, Curaçao. (Herb. Krug et Urban).

Acalypha Wilkesiana Müll—Arg. in DC. Prodr. XV. 2. (1866) 817.
Cultivated. [Ins. Pacif.]. (Symb.).

4416. **Tragia** Linn.

Tragia volubilis Linn. Spec. (1753) 980; Gris. Fl. 48; Duss 36; Millspaugh 502.

St. Eustatius: Top of the Quill 400 M.: n. 225 B., 252 B., 282 B.; inner slope of the Quill near Bengalen 350 M.: n. 703 B.

Saba: between Bottom and Marypoint 100—200 M.: n. 1467 B.; Mountain 600—800 M.: n. 1791 B.; Laddergut 0—200 M.: n. 2011 B.

St. Martin: Nakedboyhill 50—200 M.: n. 2710 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2894a B.; Mount Paradis 200—300 M.: n. 3308 B.

St. Croix ex Millspaugh.

Antill., Curaçao, Amer. cont. trop., Afr. oce. (Symb.).

4422. **Dalechampia** Linn.

Dalechampia scandens Linn. Spec. (1753) 1054; Gris. Fl. 51; Duss 40; Millspaugh 502.

St. Eustatius: near Maryglory 150 M.: n. 1143 B.

St. Martin: Nakedboyhill 280 M.: n. 2735 B.

St. Croix ex Millspaugh.

Antill., Amer. cont. trop., (Afr. trop. and Ind. or.) (Symb.).

4424. **Ricinus** Linn.

Ricinus communis Linn. Spec. (1753) 1007; Gris. Fl. 37; Duss 30; Millspaugh 502. Castor oil plant.

St. Eustatius: locality unknown: n. 23 H., 26 G., 27 G., 258 G., 288 H., 289 H.

I also saw *Ricinus communis*: Seashore near Oranjestad.

Saba: Boobyhill 300 M.: n. 1586 B.; locality unknown: n. 64 L.;

I also saw *Ricinus communis*: Laddergut.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Indig. veris. in Africa. (Symb.). Also cultivated.

4428. **Aleurites** Forst.

Aleurites moluccana Willd. Spec. Pl. IV. 1 (1805) 590. Wild date.

Cultivated. [Asia austr. orient. and Ins. pacific.] (Symb.).

4433. **Jatropha** Linn.

Jatropha curcas Linn. Spec. (1753) 1006; Gris. Fl. 36; Duss 27; Millspaugh 502. Grave physic nut.

St. Eustatius: between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 771 B.; along roads near Glassbottle: n. 1209 B.; seashore: n. 242, 2073; Cultuurvlakte: n. 244; locality unknown: n. 52 G., 168 H.

Saba: Bottom: n. 243.

I also saw *Jatropha curcas* in Forthbaygut; Hellsgategut.

St. Martin: from Philipsburg to Belvedere: n. 2595 B., 2601 B.; along roads from Grande Case to Marigot: n. 3106 B.; Cul de Sac: n. 245.

St. Croix ex Millspaugh.

Bermud., Antill., Amer. cont. trop. (Symb.).

Jatropha gossypifolia Linn. Spec. (1753) 1006; Gris. Fl. 36; Duss 25; Millspaugh 502. Physic nut.

St. Eustatius: Seashore near Oranjestad: n. 19 B.; along roads in Oranjestad: n. 570 B., 558 B., 1126 B.; in deserted plantations near Bengalen: n. 589 B.; seashore: n. 247, 249; Cultuurvlakte: n. 246; locality unknown: n. 132 G., 133 G., 196 H., 197 H.

I also saw *Jatropha gossypifolia*: on Signalhill 200 M.; top of the Cliff between Oranjestad and White Wall.

Saba: along roads between Bottom and Windwardside 200 M.; n. 1360 B.; along the Forthgut 0—150 M.: n. 1639 B.; locality unknown: n. 241 L., 313 L.

I also saw *Jatropha gossypifolia* in: Laddergut.

St. Martin: lower part of Milldrumhill: n. 3113 B.

I also saw *Jatropha gossypifolia*: along roads from Philipsburg to Naked-boyhill: in pastureground near Bellevue.

St. Croix ex Millspaugh.

Key Ins., Baham., Antill., Curaçao, Aruba, Amer. cont. trop. Africa occ. (Symb.).

Jatropha multifida Linn. Spec. (1753) 1000. Coral plant.

Cultivated. [Amer. cont. trop.] (Symb.).

4444. **Manihot** Adans.

Manihot utilissima Pohl Plant. Bras. I (1827) p. 32 t. 24. Cassave.

Cultivated. [Amer. cont. trop.]. (Symb.).

4454. **Codiaeum** A. Juss.

Codiaeum variegatum Bl. Bijdr. XII (1825) 606. Croton.

Cultivated. [Ins. Sundaic.]. (Symb.).

4483. **Sapium** Jacq.

Sapium spec. (without flowers or fruits).

Saba: Top of the Mountain 600 M.: n. 2189 B.

4486. **Hippomane** Linn.

Hippomane mancinella Linn. Spec. (1753) 1191; Gris. Fl. 50; Duss 37; Millspaugh 503. Mangeniel tree.

St. Eustatius: Seashore near Oranjestad: n. 16 B.; Concordia-bay: n. 184; seashore: n. 182; locality unknown: n. 1014 B., 324 H.

I also saw *Hippomane mancinella*: seashore near Tumbledown Dickbay.

Saba: Seashore near Flat Point: n. 2133 B.

St. Martin: from Philipsburg to Guanabay: n. 2345 B.; near Fort Willem: n. 2799 B.

I also saw *Hippomane mancinella*: along roads from Philipsburg to Nakedboyhill; Cul de Sac; Fort Willem; near Mont Chambord; near Saline; near Grande Case; Vallée des Pères; one of the islands in the Simsonbay-lagoon; Lowlands; Green Key; Tintamarre.

St. Croix ex Millspaugh.

Florida austr., Baham., Antill., Curaçao, from Mexico to Venezuela. (Symb.).

4488. **Actinostemon** Mart. ex Klotzsch.

Actinostemon caribaeus Gris. in Goett. Abh. VII (1857) 168.

Eccoccaria caribaea Gris. Flor. W. I. (1859) 51; Duss 40.

St. Eustatius: Great gut: n. 748, 749.

St. Martin: Milldrumhill: n. 3143a B., 3148 B.; John Ednyhill: n. 345, 346, 347, 348, 2051.

Antill. (Herb. Krug et Urban); Amer. cont. (Herb. Leiden).

4492. **Gymnanthes** Sw.

Gymnanthes lucida Sw. Prodr. (1788) 96; Millspaugh 503.

Eccoccaria lucida Sw. Flor. II (1800) 122; Gris. Fl. 50; Duss 39.

St. Eustatius: Top of the Quill 450 M.: n. 348 B., 372 B.; Signalhill: n. 1036 B.; near Maryglory 150 M.: n. 1152 B., 1161 B.; locality unknown: n. 333 H.

St. Croix ex Millspaugh.

Florida austr., Baham., Antill. (Symb.).

4493. **Hura** Linn.

Hura crepitans Linn. Spec. (1753) 1008; Gris. Fl. 50; Duss 38; Millspaugh 503. Sandbox tree.

St. Eustatius: along roads in Oranjestad: n. 546 B., 142 H., 147 G.

St. Martin: from Philipsburg to Belvedere: n. 2596 B.

St. Croix ex Millspaugh.

Baham., Antill., Amer. cont. trop. (Symb.).

4098. **Euphorbia** Linn.

Euphorbia articulata Burm. ex Aubl. Pl. Gui. 1 (1775) 480; Gris. Fl. 53; Millspaugh 503.

Euphorbia linearis Retz. Obs. III (1791) 32; Duss 41.

St. Eustatius: between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 767 B.; gut near Fort de Wind 50 M.: n. 847 B.; lower part of the hills near Concordia: n. 976 B., 2071; Tumbledowndiekbay: n. 2070; White Wall: n. 2068, 2069.

Saba: Green Island: n. 2131 B.

St. Croix ex Millspaugh.

Portorico, St. Thomas, Culebra Island, St. Kitts, Guadeloupe, Martinique, St. Vincent. (Herb. Krug. et Urban).

Euphorbia buxifolia Lam. Encyc. II (1786) 421; Gris. Fl. 53; Duss 42; Millspaugh 503.

St. Eustatius: White Wall: n. 811 B.; seashore near White Wall: n. 854 B.; seashore: n. 2065; Seafeatersbay: n. 2076, 2077; White Wall: n. 2064.

St. Martin: Colebay (coll. George): n. 2517 B.; Green Key: n. 3361 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida and Mexico and throughout the West Indies (Hemsley 1884).

Euphorbia heterophylla Linn. Spec. (1753) 453; Gris. Fl. 54; Duss 44.

Euphorbia cyathophora Murr. Comm. Gotting. VII (1786) 81; Millspaugh 503. Joseph's coat.

St. Eustatius: Glassgut at the seashore near Oranjestad: n. 49 B.; in a garden in Oranjestad: n. 576 B., 2096, 32 H., 36 G.; Oranjestad: n. 2098; Cultuurvlakte: n. 2097; locality unknown: n. 2078.

St. Martin: near Bethlehem: n. 2668 B.; along roads from Grande Case to Marigot: n. 3095 B.

St. Croix ex Millspaugh.

Bermudas, Eastern North-America, from Illinois southward and through the West Indies and Mexico to Peru and Brazil. (Hemsley 1884).

Euphorbia hypericifolia Linn. Spec. (1753) 454; Gris. Fl. 54; Duss 43; Millspaugh 503.

St. Eustatius: Bengalen: n. 602 B.; along roads near Glass-bottle: n. 869 B.; slope of the Quill: n. 2079, 2080, 2095.

Saba: lower part of the Bottom Mountain 250 M.: n. 1399 B.; Laddergut 200 M.: n. 2008 B.; Peperpot: n. 2081; Crespeen—Rendez-vous 400 M.: n. 2083 B.; Bottom: n. 2082; locality unknown: n. 128 L., 161 L.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2654 B.

St. Croix ex Millspaugh.

Bermudas, Eastern States of North-America, West Indies, Mexico and South-America. (Hemsley 1884).

Euphorbia petiolaris Curt. Bot. Mag. (1795) 883; Millspaugh 503.

Euphorbia verticillata Poir. Encyc. Supp. II (1811) 611; Duss 45. Black mangeniel.

St. Martin: near Filipsburg: n. 3341 B.; from Filipsburg to Belvedere: n. 2567 B., 2573 B.; Marigothill—Marigot 0—200 M.: n. 2652 B.; near Simonsbaylagoon: n. 3033 B.; along roads from Colebay to Simonsbay: n. 3253 B.; Filipsburg: n. 2031; Vineyard-hill: n. 2032; Cul de Sac: n. 2033, 2034.

I also saw *Euphorbia petiolaris*: along roads near Bethlehem; along roads from Filipsburg to Nakedboyhill; Nakedboyhill; hill between Mont des Accords and Centryhill; near Saline; along roads from Belvedere to Oysterpond; near Grande Case: lower part of Milldrunkhill; Lowlands; lower part of Mount Paradis; top of Mount Paradis.

St. Croix ex Millspaugh.

Portorico, St. Thomas, St. Jan, Guadeloupe. (Herb. Krug. et Urban).

Euphorbia pilulifera Linn. Spec. (1753) 454; Gris. Fl. 54; Duss 42; Millspaugh 503. Chicken weed.

St. Eustatius: in a garden in Oranjestad: n. 75 B.; along roads in Oranjestad: n. 512 B., 577 B.; along roads near Glassbottle: n. 875 B.; locality unknown: n. 8 G., 8 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1459 B.; along the lower part of Forthgut: n. 1640 B.; Laddergut 0—200 M.: n. 1963 B.; Crespeen—Rendez-vous: n. 2037; locality unknown: n. 239 L.

St. Martin: in an orchard near Bellevue: n. 2840 B.; Centry-hill: n. 349.

Anguilla Boldingh.

St. Croix ex Millspaugh.

All tropical countries. (Gris. Fl.).

Euphorbia prostrata Ait. Hort. Kew. ed. I. II. (1789) 139; Gris. Fl. 53; Duss 42; Millspaugh 504.

St. Martin: Guanabay: n. 2433 B.; near Mulletpond: n. 2776 B., 2780 B.

St. Croix ex Millspaugh.

Bermudas, Louisiana and Texas southward to Peru and Brazil; also in tropical Africa and naturalised in Sicily and elsewhere. (Hemsley 1884).

Euphorbia pulcherrima Willd. ex Klotzsch, in Otto et Dietr. Allg. Gartenz. II (1834) 27. Star of Bethlehem.

Cultivated. [Mexico.] (Ind. Kew.).

Euphorbia thymifolia Linn. Spec. (1753) 454; Duss 42; Millspaugh 504; Children chickenweed.

St. Eustatius: locality unknown: n. 229 H., 255 G.

St. Martin: Lowlands near Cherrycap: n. 3259 B.

All tropical countries. (Gris. Fl.).

4501. **Pedilanthus** Neck.

Pedilanthus tithymaloides Poit. in Ann. Mus. Par. XIX (1812) 390. t. 19; Gris. Fl. 52; Duss 41; Millspaugh 504. Parakiet.

St. Eustatius: along the lower part of Signalhill: n. 123 B.; gut near Fort de Wind 50 M.: n. 842 B.; along roads near Signalhill: n. 1128 B.

Saba: Bottom: n. 2250 B.

St. Martin: from Philipsburg to Guanabay: n. 2377 B.; Prince quarter (coll. George): n. 2465 B.; from Belvedere to Oysterpond: n. 3054 B.; near Fort Amsterdam: n. 2072.

I also saw *Pedilanthus tithymaloides*: along roads near Bethlehem; Marigothill to Marigot; Nakedboyhill; near Bellevue; hill between Mont des Accords and Centryhill; near Saline; hill on the northern side of Cul de Sac; Mount Paradis 300—400; Lowlands; lower part of Mount Paradis; Tintamarre; hills on the eastern side of Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Portorico, Guadeloupe, Martinique, St. Vincent, Barbados, Grenada, Tobago. (Herb. Krug et Urban.)

153. *Anacardiaceae*.

4545. *Mangifera* Linn.

Mangifera indica Linn. Spec. (1753) 200; Gris. Fl. 176; Duss 187; Millspaugh 504. *Mango tree*.

St. Eustatius: in a garden in Oranjestad: n. 57 H., 130 G., 3590 B.

Saba: locality unknown: n. 156 L., 196 L., 302 L.

I also saw *Mangifera indica*: Laddergut; Bottom; Marypoint.

St. Martin: I saw *Mangifera indica*: Marigothill—Marigot; from Cul de Sac—Marigot; Vallée des Pères; lower part of Milldrumhill; Mount Paradis 300—400 M.: Lowlands.

St. Croix ex Millspaugh.

India orient, Malaya, Mexico, Amer. centr. et merid—sept. Antill. (Engler 1883). Also cultivated.

4546. *Anacardium* Rotböll.

Anacardium occidentale Linn. Spec. (1753.) 383; Gris. Fl. 176; Duss 189; Millspaugh 505. *Cashew, Cherry*.

St. Eustatius: Slope of the lower part of the Quill near Bengalen: n. 718 B.; locality unknown: n. 45 H., 207 G.

I also saw *Anacardium occidentale*: on the slope of the Quill on the side of White Wall; top of the Quill; top of Signalhill.

Saba: locality unknown: n. 227 H.

I also saw *Anacardium occidentale*: Laddergut; near Marypoint.

St. Martin: from Philipsburg to Belvedere: n. 2597 B.; Lowlands: n. 3267 B.

I also saw *Anacardium occidentale*: along roads from Philipsburg to Nakedboyhill; Nakedboyhill; lower part of Milldrumhill.

St. Croix ex Millspaugh.

In all warmer countries of America. (Engler 1885). Also cultivated.

4552. **Spondias** Linn.

Spondias lutea Linn. Spec. (1753) 613. Yellow plum.

St. Eustatius: Signalhill: n. 125*a* B.; in a garden in Oranjestad: n. 1016 B.

Saba: Bottom: n. 1873*a* B., 1877 B., 2234 B.

Antill., Mexico, Amer. cont. austr., Malay. (Engler 1883). Also cultivated.

Spondias purpurea Linn. Spec. (1753) 613. Red plum, Jamaica plum.

Cultivated. [Antill., Mexico, Peru, Nova Granata.] (Engler 1883).

4590. **Comocladia** Linn.

Comocladia ilicifolia Sw. Prod. Veg. Ind. Occ. (1788) 17; Gris. Fl. 176; Duss 184; Millspaugh 504. Red man, Centepeep plant.

St. Eustatius: White Wall: n. 822 B.; Signalhill 150 M.: n. 1041 B., 99; Gilboohil: n. 100; locality unknown: n. 111 H., 332 H.

Saba: Bottom: n. 1852*a* B.; Springbaygut: n. 2096 B.

St. Martin: from Philipsburg to Guanabay: n. 2369 B.; Centryhill: n. 101, 102; near Point Blanche: n. 104; near Guanabay: n. 103.

I also saw *Comocladia ilicifolia*: along roads near Bethlehem; along roads from Philipsburg to Nakedboyhill; Nakedboyhill 100—200 M.; hill between Mont des Accords and Centryhill; near Saline; from Belvedere to Oysterpond; lower part of Mount Paradis.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Portorico, St. Thomas, St. Domingo, Guadeloupe, Antigua, St. Lucia. (Engler 1885).

158. **Celastraceae.**

4626. **Maytenus** Molina.

Maytenus elliptica Kr. et Urb. in Duss Fl. Ant. franç. (1897) 145 (excl. syn. Griseb.).

St. Eustatius: Top of the Quill 400 M.: n. 233 B., 344*a* B., 369 B.; Quill: n. 412*a* B.; slope of the Quill near Bengalen: n. 744*a* B.; near Signalhill: n. 952*a* B.

St. Martin: Top of Nakedboyhill: n. 2713 B.; Milldrumhill: n. 3152 B., 3182a B.

St. Croix. (Symb.).

St. Domingo, Portorico, St. Thomas, St. Jan, Montserrat, Guadeloupe, Marie Galante, Martinique, St. Vincent, Becquia. (Symb.).

4648. **Gyminda** Sarg.

Gyminda latifolia Urb. in Symbolae Antillanae V (1904) 80.

Myginda latifolia Sw. Prodr. (1788) 39; Gris. Fl. 146; Duss 148.

Myginda pallens Sarg. Forest Trees N. Amer. IX 38; Duss 147 (non Sm.); (Symb.).

Rhacoma latifolia Loesener in Engl.-Prantl. Nat. Pflanzenfam. III. 5 (1892) 217.

St. Eustatius: Top of the Quill 400 M.: n. 375a B., 393 B.

St. Martin: Colebay (coll. George): n. 2512 B.; near Simonsbay: n. 3035a B.

Key ins., Antill., Mexico. (Symb.).

Rhacoma Linn.

cf. 4649. *Myginda* Jacq. syn.

Rhacoma crossopetalum Linn. Syst. X. ed. II (1759) 896.

Myginda rhacoma Sw. Prodr. I (1788) 39; Gris. Fl. 146; Duss 147.

Myginda pallens Smith in Rees Cycl. XXV (1813) n. 4; Gris. Fl. 146; Millspaugh 505.

St. Eustatius: Signalhill 900 M.: n. 129 B.; deserted plantations near Tumbledowndickbay: n. 144 B.; White Wall: n. 816 B.; gut near Fort de Wind 50 M.: n. 841 B.; seashore near White Wall: n. 860 B.; Signalhill 50 M.: n. 1028 B.; Berg Amsterdam: n. 115; top Pongo: n. 113, 114; Pongo—Signalhill: n. 112; White Wall: n. 111; Crookwell: n. 110; Little Mountain—Boven: n. 116; Sugarloaf: n. 117.

Saba: lower part of the Bottom Mountain 250 M.: n. 1391 B.; gut between Masticgut and Hellsgate 300—400 M.: n. 2029 B.; Paris: n. 211.

St. Martin; near Mulletpond: n. 2751 B., 2753 B.; near Bethlehem: n. 2600a B.; Tintamarre: n. 3397 B., 3400 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florid. austral., Key, Baham., Antill. (Symb.).

4653. **Schaefferia** Jacq.

Schaefferia frutescens Jacq. Enum. (1760) 33; Gris. Fl. 146; Duss 148; Millspaugh 505. Boxwood.

St. Eustatius: deserted plantations near Tumbledown Dickbay: n. 147 B.; from Little Mountain to Venusbay: n. 962 B.; Boven: n. 215.

Saba: along the Forthgut 150—200 M.: n. 1636 B.

St. Martin: from Philipsburg to Cul de Sac: n. 2610 B.; Marigothill—Marigot 0—200 M.: n. 2647 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2859 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2964 B.; Colebayhill: n. 210; Centryhill: n. 1, 2, 3; French frontier: n. 5; locality unknown: n. 4, 7.

St. Croix ex Millspaugh.

Florid. austral., Key, Baham., Mexico, Nova Granata, Antill. (Symb.).

165. **Sapindaceae.**4726. **Cardiospermum** Linn.

Cardiospermum halicacabum Linn. Spec. (1753) 366; Gris. Fl. 122; Duss 118; Millspaugh 505. Sprainbushwine.

St. Eustatius: Bengalen: n. 618 B.; along roads in Oranjestad: n. 1106 B.; near the seashore: n. 146; slope of the Quill: n. 148; Cultuurvlakte: n. 147; locality unknown: n. 154 G., 294 H.

Saba: St. Giles: n. 1868 B.; Windwardside: n. 2307 B.

St. Martin: from Philipsburg to Guanabay: n. 2421 B.; Point Blanche (coll. George): n. 2496 B.; Cul de Sac—Marigothill 0—200 M.: n. 2632 B.; along roads from Marigot to Philipsburg: n. 3112 B.; Mount Paradis 200—300 M.: n. 3316 B.; Marigotroad: n. 150, 151, 152.

St. Croix ex Millspaugh.

Trop. and warmer countries of both hemisph. (E. a. P.).

4734. **Allophylus** Linn.

Allophylus occidentalis Radlk. in Sitzungsber. phys. math. Cl. Bayer. Acad. Wiss. XX (1890) 230.

Schmiedelia occidentalis Sw. Fl. Ind. Occ. II (1800) 665; Gris. Fl. 126; Duss 122; Millspaugh 506.

St. Eustatius: Top of the Quill 400 M.: n. 213 B.; inner slope of the Quill 350 M.: n. 455 B., 1215 B., 329 H.; locality unknown: n. 50, 51, 52.

Saba: Boobyhill 300 M.: n. 1591 B.; gut between Hellsgate and Mastiegut 300 M.: n. 2057 B.

St. Croix ex Millspaugh.

Antill., Cuba to Minaes Geraes. (Gris. Fl.).

4760. **Melicocca** Linn.

Melicocca bijuga Linn. Spec. (1762) 495; Gris. Fl. 127; Duss 123; Millspaugh 505. Kinnuptree.

St. Eustatius: Signalhill 200 M.: n. 136 B., 165 B.; Bengalen: n. 633 B.; lower part of the hills near Concordia: n. 978 B.; along the slope of the lower part of Signalhill: n. 143; Oranjestad: n. 144, 86 H., 204 H.

Saba: Bottom: n. 1872 B.; locality unknown: n. 1a L.

I also saw *Melicocca bijuga* in Laddergut.

St. Martin: I saw *Melicocca bijuga*: along roads from Philipsburg to Nakedboyhill; near Bellevue.

St. Croix ex Millspaugh.

Trinidad, Jamaica, Guiana, Nova Granata. (Gris. Fl.).

Also cultivated.

4831. **Dodonaea** Linn.

Dodonaea viscosa Linn. Mant. (1767) 228; Gris. Fl. 127; Duss 123; Millspaugh 506.

St. Eustatius: Slope of the Quill above White Wall 350 M.: n. 1 B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 766 B.; White Wall: n. 804 B.; gut near Fort de Wind: n. 828 B., 835a B.; slope of the Quill on the side of White Wall 200 M.: n. 1199 B.; near Sugarloaf: n. 154, 155, 162; near White Wall: n. 156, 157, 163; locality unknown: n. 3635 B.

St. Martin: at the seashore near Simonsbayvillage: n. 3031 B.; Colebay (coll. George): n. 2504 B.; Simonsbay: n. 159, 161.

St. Croix ex Millspaugh.

In almost all warm countries. (Hemsley 1884).

4833. **Hypelate** Swartz.

Hypelate trifoliata Sw. Prod. Veg. Ind. Occ. (1788) 61; Gris. Fl. 127.

St. Martin: near Simonsbaylagoon: n. 2738a B.

Anguilla Boldingh.

Baham., Antill. (Herb. Krug et Urban).

168. Balsaminaceae.4856. **Impatiens** Linn.**Impatiens Balsamina** Linn. Spec. (1753) 938.

Cultivated. [India orient?]. (Ind. Kew.).

169. Rhamnaceae.4861. **Zizyphus** Adans.**Zizyphus Jujuba** Lam. Encycl. III (1789) 318; Duss 94.**St. Martin:** from Filipsburg to Guanabay: n. 2393a B.; near Filipsburg: n. 2337a B.

India. orient., Malay. (Ind. Kew.).

4875a. **Krugiodendron** Urb.**Krugiodendron ferreum** Urb. Symb. Ant. III (1902) 314; Millspaugh 506.*Condalia ferrea* Gris. Flor. W. I. (1859) 100; Duss 92. Iron-berry.**St. Eustatius:** from Little Mountain to Venusbay: n. 963 B.; Boven: n. 192; locality unknown: n. 337 G.**St. Martin:** near Mulletpond: n. 2773 B.; Mount Paradis: n. 3313 B.; French frontier: n. 191.**St. Croix** ex Millspaugh.

Florida, Key Ins., Baham., Antill. (Symb.).

4882. **Colubrina** L. Cl. Rich.**Colubrina ferruginosa** Brong. in Ann. Sc. Nat. Sér. I X (1827) 386, t. 15 f. 3; Gris. Fl. 100; Millspaugh 506.**Saba:** St. Giles: n. 1866a B.

Florida, Antill. (E. a. P.).

4902. **Gouania** Jacq.**Gouania domingensis** Linn. Spec. (1763) 1663; Gris. Fl. 101; Duss 93; Millspaugh 506. White root.**St. Eustatius:** from Tumbledownickbay to Signalhill; n. 168 B.; near Maryglory 150 M.: n. 1136 B.**Saba:** lower part of the Bottom Mountain 300 M.: n. 1444 B.; Hellsgate: n. 189.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2636 B.; Marigotroad near Marigot: n. 2811 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2954 B.; lower part of Colebayhill: n. 153, 190.

St. Croix ex Millspaugh.

Amer. trop. (E. a. P.).

170. Vitaceae.

4918. **Cissus** Linn.

Cissus obovata Vahl Symb. Bot. III (1764) 19; Gris. Fl. 102.

St. Martin: Top of Mount Paradis: n. 3236 B., 3350 B.

St. Croix. (Planchon 1883).

Portorico. (Planchon 1883).

Cissus sicyoides Linn. Syst. ed. X. (1759). 897; Gris. Fl. 102; Duss 95; Millspaugh 506. Pudding withe.

Vitis sicyoides Miq. in Ann. Mus. Bot. Lugd. Bat. I. (1863) 83.

St. Eustatius: Seashore near Oranjestad: n. 44 B.; slope of the hills near Tumbledowndiekbay: n. 153 B.; top of the Quill 880 M.: n. 284 B.; slope of the Quill near Bengalen 100 M.: n. 668 B., 709 B., 727 B.; Cultuurvlakte: n. 5; seashore: n. 6, 7; locality unknown: n. 129 H., 155 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1344 B.; between Bottom and Marypoint 190—200 M.: n. 1506 B.; gut between Hellsgate and Masticgut 300—400 M.: n. 2051 B., 2076 B.; Springbaygut: n. 2107 B.; Bottom: n. 14, 15; locality unknown: n. 104 L.

I also saw *Cissus sicyoides* in Laddergut.

St. Martin: from Filipsburg to Guanabay: n. 2384 B.; along the saltpond near Filipsburg: n. 2443 B.; from Filipsburg to Belvedere: n. 2589 B.; gut from Bethlehem to Cul de Sac: n. 3434 B.; hills on the eastern side of Cul de Sac: n. 3437 B.; Centryhill: n. 17; Cul de Sac: n. 16.

I also saw *Cissus sicyoides*: in Marigothill—Marigot; Nakedboyhill; in pastureground near Belvedere; hill between Mont des Accords and Centryhill; hill on the northern side of Cul de Sac; from Belvedere to Oysterpond; Mount of Paradis 300—400 M.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Americ. trop. (Planchon 1887).

174. Tiliaceae.

4953. **Corchorus** Linn.**Corchorus acutangulus** Linn. Encyc. (1786) II 104; Gris. Fl. 97; Millspaugh 507.**St. Eustatius**: Bengalen: n. 607 B.; Fort de Wind: n. 803 B.**St. Croix** ex Millspaugh.

In all tropical countries (E. a. P.).

Corchorus hirsutus Linn. Spec. (1753) 530; Gris. Fl. 97; Duss 507.**St. Eustatius**: Slope of the Quill near Bengalen: n. 656a B.; from Little Mountains to Venusbay: n. 994 B.; Signalhill 200 M.: n. 1044 B.; in a garden in Oranjestad: n. 1273 B.**St. Martin**: (coll. George:) n. 2495a B.; near Mulletpond: n. 2762 B.; Mount Paradis 300—400 M.: n. 3245 B.; Fort Amsterdam: n. 252; French frontier: n. 225.**Anguilla** Boldingh.**St. Croix** ex Millspaugh.

Baham., Antill., Aruba, Bonaire. (Herb. Krug et Urban).

Corchorus olitorius Linn. Spec. (1758) 529; Gris. Fl. 97.**St. Eustatius**: locality unknown: n. 269 G.

In all tropical countries. (E. a. P.).

Corchorus siliquosus Linn. Spec. (1753) 529; Gris. Fl. 97; Duss 89; Millspaugh 507.**St. Martin**: Top of the hill on the northern side of Cul de Sac: n. 3005 B.; Marigotroad: n. 248.**Anguilla** Boldingh.**St. Croix** ex Millspaugh.

Antill., Florida to Texas, Panama, Nova Granata. (Gris. Fl.).

4975. **Triumfetta** Linn.**Triumfetta rhomboidea** Jacq. Enum. Pl. Carib. (1760) 22; Gris. Fl. 96; Millspaugh 507. Wild mahoe.**St. Eustatius**: locality unknown: n. 185 H., 210.; slope of the Quill: n. 247.**Saba**: Ladder: n. 246.**St. Croix** ex Millspaugh.

In both hemisph. (E. a. P.).

Triumfetta semitriloba Jacq. Enum. Pl. Carib. (1760) 22; Gris. Fl. 96; Duss 89; Millspaugh 506.**St. Eustatius**: Cliff near Oranjestad: n. 56 B.; top of the Quill 400 M.: n. 198 B.; slope of the Quill near Bengalen 50—200 M.:

n. 645 B.; 711 B.; Cultuurvlakte: n. 155; slope of the Quill: n. 229, 230, 231, 232; Quill: n. 243, 245; locality unknown: n. 244.

Saba: lower part of the Bottom Mountain 250 M.: n. 1398 B., 1415 B.; Windwardside—Hellsgate: n. 1663 B.; Mountain 600—800 M.: n. 1792 B.; gut between Hellsgate and Masticgut 300—400 M.: n. 2062 B.; Mountain near Windwardside 600 M.: n. 2162 B.; Peperpot: n. 236; Gainbygut: 235.

St. Martin: Cul de Sac—Marigothill 0—200 M.: n. 2635 B.; near Bethlehem: n. 2665 B.; Nakedboyhill 50—200 M.: n. 2705 B.; Mount Paradis 300—400 M.: n. 3215 B., 3243 B.; Centryhill: n. 234; Marigotroad: n. 233.

St. Croix ex Millspaugh.

In both hemisph. (E. a. P.).

175. Malvaceae.

4983. **Abutilon** Adans.

Abutilon crispum Sweet. Hort. Brit. ed. I (1827) 53; Gris. Fl. 79; Duss 68.

St. Eustatius: English Quarter: n. 606, 616, 626, 636.

St. Vincent, New Mexico, Texas, Cuba and Mexico to Venezuela, East Indies. (Gris. Fl.).

Abutilon indicum Sweet. Hort. Brit. ed. I. (1827) 54; Gris. Fl. 78; Duss 67; Millspaugh 507.

St. Eustatius: Bengalen: n. 627 B., 1188 B.; along roads in Oranjestad: n. 932 B.; Concordia: n. 255; English Quarter: n. 287; locality unknown: n. 201 H.

Saba: locality unknown: n. 75 L.

St. Martin: along roads from Bethlehem to Mont Chambord: n. 2915 B.; Mount Paradis 200—400 M.: n. 3204 B.

I also saw *Abutilon indicum*: along roads from Filip-burg to Nakedboy-hill; hill between Mont des Accords and Centryhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Panama. Niger, Nubia to Mozambique, East Indies. (Gris. Fl.).

4991. **Althaea** Linn.

Althaea officinalis Linn. Spec. (1753) 686.

Cultivated. [Europa orient.] (Ind. Kew.).

4995. **Malvastrum** A. Gray.

Malvastrum spicatum A. Gray. in Mem. Am. Acad. N. S. IV (1849) 22; Gris. Fl. 72; Duss 62; Millspaugh 508.

St. Eustatius: in a garden in Oranjestad: n. 90 B.; Fort de Wind: n. 799 B.; locality unknown: n. 256 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1304 B.; Hellsgate—Marypoint: n. 2046 B.

St. Martin: from Philipsburg to Guanabay: n. 2346 B.; near Bethlehem: n. 2663 B.

St. Croix ex Millspaugh.

Antill., Texas to Brazil., Cape Verde Islands, East Indies, New Holland. (Gris. Fl.).

Malvastrum tricuspidatum A. Gray Pl. Wright I (1852) 16; Gris. Fl. 72; Duss 63.

Malvastrum coromandelianum Garcke Bonplandia V (1857) 297; Millspaugh 508.

St. Eustatius: Quill: n. 901 B.

St. Croix. (Symb.).

Cosmop. trop. (Ind. Kew.).

4998. **Sida** Linn.

Sida carpinifolia Linn. f. Suppl. (1781) 307; Gris. Fl. 73; Duss 63; Millspaugh 508.

Sida acuta Burm. f. Fl. Ind. (1768) 147. Soap bush.

St. Eustatius: in a garden in Oranjestad: n. 91 B., 568 B.; Bengalen: n. 620 B.; Fairplay: n. 1505; Cultuurvlakte: n. 1502; slope of the Quill: n. 1506; locality unknown: n. 1507, 15 H., 16 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1340 B.; between Bottom and Marypoint 100—200 M.: n. 1503 B.; top of the Cliff near the seaside of Boobyhill: n. 1601 B.; Windwardside: n. 1901 B.; Laddergut 0—200 M.: n. 2018 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2028 B.; locality unknown: 263 L.

St. Martin: from Guanabay to Philipsburg: n. 2439 B.; lower part of Milldrumhill: n. 3170 B.

I also saw *Sida carpinifolia* on the hill between Mont des Accords and Centryhill; between Mount Vernon and Saline; near Grande Case; Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Antill., Florida, Bermud., Cuba and Guatemala to Nova Granata and Brazil., trop. Afr., Canary Islands, East Indies. (Gris. Fl.).

Sida ciliaris Linn. Syst. Ed. X (1759) 1145; Gris. Fl. 73; Duss 64; Millspaugh 509.

St. Eustatius: on the top of the Cliff between Oranjestad and Fort de Wind: n. 756 B.; near Maryglory 150 M.: n. 1153 B.; near Boven 200 M.: n. 1245 B.

St. Martin: from Philipsburg to Belvedere: n. 2562 B.; between Mount Vernon and Orientbay: n. 2938 B.; lower part of Colebay-hill: n. 470 A.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Venezuela. (Gris. Fl.).

Sida cordifolia Linn. Spec. (1753) 684; Gris. Fl. 76; Duss 67; Millspaugh 509. Mash mellow.

St. Eustatius: Bengalen: n. 623 B.; Cultuurvlakte: n. 511, 524; English Quarter: n. 510; locality unknown: n. 3582 B., 182 G.

Saba: Windwardside—Hellsgate: n. 1662 B.; Hellsgate: n. 1695 B.

St. Martin: Prince quarter (coll. George): n. 2462 B.; from Philipsburg to Belvedere: n. 2559 B.; Nakedboyhill 50—200 M.: n. 2679 B.; from Belvedere to Oysterpond: n. 3048 B.; along roads from Grande Case to Marigot: n. 3094 B.; locality unknown: n. 3597 B.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Peru and Brazil., Cape Verde Islands, Niger, Nubia, Cape, East Indies, Canton. (Gris. Fl.).

Sida glomerata Cav. Diss. I (1785) 18 t. 2 f. 6.; Gris. Fl. 73; Duss 64.

St. Eustatius: locality unknown: n. 275 H.

Bermud., Antill., Mexico. (Hemsley 1884).

Sida rhombifolia Linn. Spec. (1753) 684; Gris. Fl. 74; Duss 75; Millspaugh 509.

St. Martin: from Flat Point to Windwardside: n. 2143 B.; Mount Paradis: n. 3203 B.

Trop. countries of both hemisph. (Ind. Kew.).

Sida spinosa Linn. Spec. (1753) 683; Gris. Fl. 74; Duss 64; Millspaugh 509.

St. Eustatius: Bengalen: n. 617 B.; hill near Concordia 100 M.: n. 945 B.; Cultuurvlakte: n. 131, 205.

St. Martin: Cul de Sac: n. 2616 B.; near Bethlehem: n. 2664 B.

St. Croix ex Millspaugh.

Afr., Asia, America. (E. a. P.).

Sida supina L'Hérit. Stirp. Nov. (1784) 109bis, t. 52; Gris. Fl. 75; Millspaugh 509.

St. Eustatius: along roads in Oranjestad: n. 918 B.; near Maryglory 150 M.: n. 1155 B.

Saba: lower part of the Bottom Mountain 250 M.: n. 1396 B.; between Bottom and Marypoint 100—200 M.: n. 1457 B.; Ladder-gut 200 M.: n. 2005 B.; Peperpot: n. 658.

St. Martin: near Fort Willem: n. 2542 B., 2793 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2948 B.; along roads in Cul de Sac: n. 2967 B.; lower part of the Mill-drumhill: n. 3168 B.

St. Croix ex Millspaugh.

Antill., Haiti and Mexico to Venezuela and Lima, Galapagos Islands. (Gris. Fl.).

5006. **Urena** Linn.

Urena lobata Linn. Spec. (1753) 692; Gris. Fl. 81; Duss 70.

Urena americana Linn. Pl. Surin. II (1775); Millspaugh 509.

Saba: between Bottom and Marypoint 300 M.: n. 1559 B., 1472 B.; Mountain 800 M.: n. 1799 B., 1803 B., 19, 31.

St. Croix ex Millspaugh.

In all tropical countries. (E. a. P.).

5007. **Pavonia** Cav.

Pavonia spinifex Cav. Diss. III (1787) 133; Gris. Fl. 82; Duss 72; Millspaugh 508. Ginger bush.

Saba: lower part of the Bottom Mountain 250 M.: n. 1401 B.; Windwardside: n. 1592 B., Crespeen—Rendez-vous 400 M.: n. 29; Middle island: n. 30.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2650 B.; lower part of Mount Paradis: n. 3188 B., 3302 B.

St. Croix ex Millspaugh.

Bermud., Antill., Amer. merid. (Hemsley 1884).

5013. **Hibiscus** Linn.

Hibiscus esculentus Linn. Spec. (1753) 696.

Cultivated. [In all trop. countries]. (Ind. Kew.).

Hibiscus radiatus Cav. Diss. III (1787) 150 t. 54 f. 2. Hawthorn.

Cultivated. [Asia and Austral. trop.]. (Ind. Kew.).

Hibiscus Rosa-sinensis Linn. Spec. (1753) 694. Hibisc.

Cultivated. [Trop. countries of the Old world]. (Ind. Kew.).

Hibiscus Sabdariffa Linn. Spec. (1753) 695.

Cultivated. [Trop. countries of the Old world]. (Ind. Kew.).

Hibiscus schizopetalus Hook. f. Bot. Mag. (1880) t. 6524. Basket hibisc.

Cultivated. [Afr. trop.]. (Ind. Kew.).

Hibiscus tiliaceus Linn. Spec. (1753) 694.

Paritium tiliaceum St. Hill Fl. Bras. Mer. I (1827) 256; Gris. Fl. 86; Duss 79; Millspaugh 508.

St. Eustatius: at the seashore near Concordiabay: n. 1089 B.

St. Martin: Mount Paradis 300—400 M.: n. 3326 B.

St. Croix ex Millspaugh.

All tropical countries. (E. a. P.).

5014. **Abelmoschus** Médik.

Abelmoschus moschatus Médik. Malv. (1787?) 46.

Cultivated. [Trop. countries of the Old world]. (Ind. Kew.).

5018. **Thespesia** Soland.

Thespesia populnea Soland. ex Correa in Ann. Mus. Par. IX (1807) 290; Gris. Fl. 87; Duss 79; Millspaugh 509.

St. Eustatius: Seashore near Oranjestad: n. 30 B.; seashore near Tumbledown Dickbay: n. 154 B.; slope of the Quill 50 M.: n. 35.

Saba: Bottom: n. 2246 B., 36; locality unknown: n. 233 L.

St. Martin: Cul de Sac—Marigothill 0—200 M.: n. 2625a B.

St. Croix ex Millspaugh.

Antill., East Indies, North coast of Austral., New South Wales, Senegambia. (Gris. Fl.).

5020. **Gossypium** Linn.

Gossypium barbadense Linn. Spec. (1753). 693. Creole cotton.

Cultivated. [Amerika]. (E. a. P.).

177. **Bombacaceae.**

5023. **Adansonia** Linn.

Adansonia digitata Linn. Spec. (1753); 1190. Guinea tamarind.

[Africa]. (E. a. P.).

5026. **Ceiba** Médik.

Ceiba pentandra Gärtn. Fruct. II (1791). 244 t. 133.

Eriodendron anfractuosum DC. Prod. I (1824) 479; Gris. Fl. 88; Duss 80; Millspaugh 510. Silk cotton tree.

St. Eustatius: Top of the Quill 400 M.: n. 224 B.; in a garden in Oranjestad: n. 321 H.

I also saw *Ceiba pentandra*: at the seashore near Oranjestad: Bottom of the Quill.

St. Martin: Cul de Sac: n. 2619 B.

I also saw *Ceiba pentandra*: near Grande Case; Vallée des Pères.

St. Croix ex Millspaugh.

Mexico, Antill., Guiana, Afr., Malay. (E. a. P.).

Also cultivated.

5033. **Quararibea** Aubl.

Quararibea turbinata Poir. Encycl. Suppl. IV. 636.

Myrodia turbinata Sw. Prod. Veg. Ind. Occ. (1788) 102; Gris. Fl. 88; Duss 82; Millspaugh 510. Miller wood.

St. Eustatius: Bottom of the Quill: n. 424a B., 896a B., 113 H., 217 G., 118, 119, 120, 121, 122.

St. Croix ex Millspaugh.

Brazil. (Ind. Kew.).

178. **Sterculiaceae.**5057. **Melochia** Linn.

Melochia nodiflora Sw. Prod. Veg. Ind. Occ. (1788) 97; Gris. Fl. 94; Duss 87; Millspaugh 510.

St. Martin: Top of the hill on the northern side of Cul de Sac 200—300 M.: n. 2989 B.; Tintamarre: n. 3392 B.

St. Croix ex Millspaugh.

Antill., Mexico, Panama. (Gris. Fl.).

Melochia pyramidata Linn. Spec. (1753) 674; Gris. Fl. 93; Duss 86; Millspaugh 510.

St. Eustatius: along roads in Oranjestad: n. 1076 B.; slope of the Quill: n. 198, 199; Cultuurvlakte: n. 197, 200; in a garden in Oranjestad: n. 200, 39 G.

St. Croix ex Millspaugh.

Baham., Antill., Mexico, Texas, Venezuela, Guayaquil, Brazil. (Gris. Fl.).

Melochia tomentosa Linn. Syst. ed. X (1759) 1140; Gris. Fl. 93; Duss 86; Millspaugh 510. Black toarch.

St. Eustatius: Slope of the lower part of Signalhill: n. 102 B.; Signalhill 200 M.: n. 127 B.; along roads in Oranjestad: n. 924 B.; along roads near Concordiabay: n. 975 B.; Cultuurvlakte: n. 204; along roads: n. 232 H.

Saba: I saw *Melochia tomentosa* on Crispeenhill.

St. Martin: from Philipsburg to Guanabay: n. 2354 B.; Prince quarter (coll. George): n. 2472a B.; Fort Willem 0—50 M.: 2532 B., 2795 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2945 B.; Colebayhill: n. 205; Prince quarter: n. 116.

I also saw *Melochia tomentosa*: along roads from Nakedboyhill to Philipsburg; Nakedboyhill; from Belvedere to Oysterpond; Tintamarre.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Venezuela and Brazil. (Gris. Fl.).

5059. **Waltheria** Linn.

Waltheria americana Linn. Spec. (1753) 673; Gris. Fl. 95; Duss 87.

Waltheria indica Linn. Spec. (1753) 673; Millspaugh 511. Mash mellow.

St. Eustatius: Seashore near Oranjestad: n. 37 B.; top of the Quill 400 M.: n. 503 B.; in a garden in Oranjestad: n. 509 B.; Bengalen: n. 619 B.; slope of the lower part of the Quill near Bengalen: n. 720 B.; on the Cliff between Oranjestad and Fort de Wind: n. 757 B., 763a B.; Pongohill: n. 43; slope of the Quill: n. 44, 45, 46; Cultuurvlakte: n. 42, 47; along roads in Oranjestad: n. 181 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1528 B.; Windwardside—Hellgate: n. 1685 B.

St. Martin: near Fort Willem: n. 2794 B.; in an orchard near Bellevue: n. 2844 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

In tropical and subtropical countries. (Hemsley 1884).

5063. **Ayenia** Loefl.

Ayenia pusilla Linn. Syst. ed. X (1759) 1247; Gris. Fl. 92; Millspaugh 510.

St. Eustatius: Top of the Cliff between Gallowsbay and White Wall: n. 761 B.

St. Martin: from Filipsburg to Point Blanche: n. 3400a B.

St. Croix ex Millspaugh.

Antill., Peru. (Ind. Kew.).

5068. **Theobroma** Linn.

Theobroma Cacao Linn. Spec. (1753) 782; Gris. Fl. 91; Duss 85; Millspaugh 510. Cocoa.

St. Eustatius: Bottom of the Quill 350 M.: n. 446 B., 1217 B.; Quill: n. 123, 124, 125, 102 H., 214 G., 221 G.

Saba: Windwardside: n. 1937 B.

St. Croix ex Millspaugh.

Amer. cent. and merid., Curaçao(?) (E. a. P.).

5081. **Helicteres** Linn.

Helicteres jamaicensis Lam. Encyc. IV 87; Gris. Fl. 89; Millspaugh 510.

Helicteres Isora Linn. Spec. (1753) 963.

St. Martin: Prince quarter: n. 501.

St. Croix ex Millspaugh.

Antill., Asia et Austral. trop. (Ind. Kew.).

182. Ochnaceae.

5126. **Sauvagesia** Linn.

Sauvagesia erecta Linn. Spec. (1753) 203; Gris. Fl. 112; Duss 108.

Saba: Mountain 800 M.: n. 1821 B., 2209 B., 134; gut between Hellsgate and Mastiegut 300—400 M.: n. 2025 B.

In all tropical countries. (E. a. P.).

184. Marcgraviaceae.

5137. **Marcgravia** Linn.

Marcgravia umbellata Linn. Spec. (1753) 503; Gris. Fl. 110; Duss 106.

Saba: Mountain near Windwardside 600 M.: n. 2159 B., 2160 B.; Mountain 800 M.: n. 2207 B., 58, 59; Mountain near Hellsgate 500 M.: n. 2275 B., 2276 B.

Antill., Guiana and Columbia. (Urban 1896.).

186. Theaceae.

5135. **Ternstroemia** Mutis ex Linn. f.**Ternstroemia peduncularis** DC. in Mem. Soc. Phys. Genève. I (1822) 409.*Ternstroemia elliptica* Sw. Prod. Veg. Ind. Occ. (1788) 81; Gris. Fl. 103; Duss 96; Millspaugh 511.**St. Eustatius**: Top of the Quill 400 M.: n. 5 B., 228 B., 248 B., 264 B., 323 B., 359 B., 401 B., 504 B., 222, 223, 224, 225, 227, 228: inner slope of the Quill 350 M.: n. 1228 B.; locality unknown: n. 130, 226.**St. Martin**: Lowlands: n. 3271 B.**St. Croix** ex Millspaugh.

Antill., Mexico, Panama. (Gris. Fl.).

Freziera Sw.cf. 5157. *Eurya* Thunb. subg. II.**Freziera undulata** Willd. Sp. Plant. II. (1799) 1179; Gris. Fl. 104; Duss 97.**Saba**: Mountain 800 M.: n. 2211 B., 64, 65.

St. Kitts, Montserrat, Guadeloupe, Dominica, Martinique, St. Vincent, St. Lucia, Grenada, Trinidad. (Urban 1896).

187. Guttiferae.

5164. **Marila** Swartz.**Marila racemosa** Sw. Prod. Veg. Ind. Occ. (1788) 84; Gris. Fl. 111; Duss 107.**Saba**: Mountain 800 M.: n. 1829 B.

Guadeloupe, Martinique, Dominica, St. Vincent. (Herb. Krug et Urban).

5176. **Mammea** Linn.**Mammea americana** Linn. Spec. (1753) 512; Gris. Fl. 108; Duss 102; Millspaugh 511. Mammie.**St. Eustatius**: along the lower part of Signalhill: n. 120 B.; inner slope of the Quill 350 M.: n. 461 B.; slope of the Quill near Bengalen 150—200 M.: n. 651 B.; Quill: n. 9; locality unknown: n. 70 H., 184 G.**Saba**: Boobyhill 300 M.: n. 1619 B.; locality unknown: n. 114 L.I also saw *Mammea americana*: Hellsgate.

St. Martin: I saw *Mammea americana*: Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Antill., Guiana, Nova Granata; in warmer parts of Amer. Afr. and Asia cultivated. (Vesque 1893).

Also cultivated.

5181. *Clusia* Linn.

Clusia alba Jacq. Enum. Pl. Carib. (1760) 34; Gris. Fl. 107; Millspaugh 511. Wild mammie.

St. Eustatius: Top of the Quill 400 M.: n. 193 B., 376 B., 8, 10, 100 H., 185 G.

Saba: between Bottom and Marypoint 350 M.: n. 1489 B.; Hellsgate: n. 1705 B.; Mountain near Hellsgate 600 M.: n. 2288 B.; locality unknown: n. 357 L.

I also saw *Clusia alba*: on Gates Hill; in Gut between Hellsgate and Mastiegut.

St. Croix ex Millspaugh.

St. Kitts, Martinique, Dominica, Guadeloupe. (Vesque 1893).

Clusia rosea Jacq. Enum. Pl. Carib. (1760) 34; Gris. Fl. 107; Duss 99; Millspaugh 511.

St. Martin: Nakedboyhill 200—265 M.: n. 2718 B.; Centryhill: n. 12.

St. Croix ex Millspaugh.

Baham., Antill., Venezuela, Panama, Carolina. (Vesque 1893).

Clusia spec. nov. without flowers and fruits.

Saba: Greathill: n. 11.

194. *Bixaceae*.

5249. *Bixa* Linn.

Bixa Orellana Linn. Spec. (1753) 512; Gris. Fl. 20; Duss 14; Millspaugh 511. Roucou.

St. Eustatius: in a garden in Oranjestad: n. 533 B., 140 H., 231 G.

Saba: lower part of the Bottom Mountain 300 M.: n. 1426 B.; Bottom: n. 1850 B., 2241 B.; locality unknown: n. 245 L., 292 L.

St. Croix ex Millspaugh.

Indig. in Amer. trop. (E. a. P.). Also cultivated.

197. *Canellaceae*.

5254. *Canella* Swartz.

Canella alba Murr. Syst. ed. XIV (1784) 443; Gris. Fl. 109; Duss 103; Millspaugh 511. Pepper cinnamom.

St. Martin: Colebay (coll. George): n. 2503 B., 2505 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2740 B.; near Mulletpond: n. 2769 B.; near Vischvijver: n. 2926 B.; near Simonsbaylagoon: n. 3038 B.; Milldrumhill 300—400 M.: n. 3126 B.; Lowlands: n. 3282 B.

St. Croix ex Millspaugh.

Florida, Antill. (E. a. P.).

199. Flacourtiaceae.

5307. **Prockia** Linn.

Prockia Crucis Linn. Syst. ed. X (1759) 1074.

Cultivated. [Amer. trop.l. (E. a. P.).

5320. **Xylosma** Forst.

Xylosma buxifolium A. Gray in Mem. Am. Acad. N. S. VIII. (1860) 155.

Myroxylon buxifolium Krug et Urban in Additamenta ad Cognitionem Florae Indiae Occidentalis I (1893) 300; Duss 15. Cockspur.

St. Eustatius: Top of the Quill 450 M.: n. 373 B., 43.

Baham., Antill. (Urban 1893).

5337. **Samyda** Linn.

Samyda serrulata Linn. Spec. (1762) 558; Gris. Fl. 25; Duss 17; Millspaugh 512. Bansa, Wild guave.

St. Eustatius: Signalhill 150 M.: n. 1025 B.; near Maryglory 150 M.: n. 1146 B.; slope of the Quill on the side of White Wall 200 M.: n. 1169 B.

St. Martin: from Philipsburg to Guanabay: n. 2352 B., 2372 B., 2390 B.; Prince quarter (coll. George): n. 2425a B., 2458 B.; Fort Willem: n. 2532a B.; Cul de Sac: n. 2605 B.; Nakedboyhill 50—200 M.: n. 2687 B.; 200—265 M.: n. 2723 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2951 B.; Mount Paradis 200—400 M.: n. 3207 B.; 100—200 M.: n. 3301 B.; 200—300 M.: n. 3305 B.; locality unknown: n. 3614 B.; Centryhill: n. 30, 88; Colebayhill: n. 32; Vineyardhill: n. 31, 33; French frontier: n. 89.

I also saw *Samyda serrulata*: near Saline; from Belvedere to Oysterpond; along roads near Grande Case; Lowlands.

St. Croix ex Millspaugh.

St. Domingo, Portorico, St. Thomas, Antigua, St. Barthélemy, Trinidad. (Herb. Krug et Urban).

5343. **Casearia** Jacq.

Casearia parvifolia Willd. Sp. Pl. II (1799) 628; Gris. Fl. 23; Duss 16; Crack open.

St. Eustatius: Top of the Quill 400 M.: n. 220 B., 236 B., 299 B.; top of the Quill 500 M.: n. 496 B.; slope of the Quill near Bengalen 150 M.: n. 677 B., 682 B.; slope of the Quill near Glassbottle: n. 867 B.; Bottom of the Quill 300 M.: n. 892 B.; top of the Quill: n. 160, 166; locality unknown: n. 196 G.

I also saw *Casearia parvifolia*: slope of the Quill on the side of White Wall.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1345 B.; Greathill 2—400 M.: n. 163; Paris: n. 161, 162; Gainhygut: n. 164; Cowpasture: n. 165; locality unknown: n. 327 L.

I also saw *Casearia parvifolia*: Forthbaygut: gut between Hellsgate and Mastiegut; Springbaygut.

St. Martin: Marigot hill—Marigot 0—200 M.: n. 2638 B., 2646 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2894 B.; Mount Paradis 200—400 M.: n. 3310 B.; Centryhill: n. 29; John Ednyhill: n. 28; French frontier: n. 27.

I also saw *Casearia parvifolia*: on the hills on the northern side of Cul de Sac; from Belvedere to Oysterpond.

St. Croix ex Millspaugh.

Antill., Guiana, Pernambuco. (Gris. Fl.).

Casearia sylvestris Sw. Fl. Ind. Occ. II. (1800) 752; Gris. Fl. 23; Millspaugh 512.

St. Martin: Hill on the northern side of Cul de Sac 0—100 M.: n. 3005a B.; 200—300 M.: n. 3011 B.; Mount Paradis 300—400 M.: n. 3201 B., 3241 B.

St. Croix ex Millspaugh.

In tropical America. (E. a. P.).

201. **Turneraceae.**5357. **Piriqueta** Aubl.

Piriqueta villosa Aubl. Pl. Gui. I (1775) 298 t. 117.

Piriqueta cistoides G. F. W. Mey ex Steud. Nom. ed. II (1841) 344; Gris. Fl. 298; Duss 314.

St. Eustatius: along roads near English Quarter: n. 997 B., 998a B.; slope of the Quill on the side of White Wall 200 M.: n. 1185 B.; slope of the Quill: n. 237, 238; locality unknown: n. 236.

Antill., Cuba and Panama to Brazil. (Gris. Fl.).

5360. **Turnera** Linn.

Turnera ulmifolia Linn. Spec. (1753) 271; Gris. Fl. 296; Millspaugh 512.

St. Eustatius: White Wall: n. 808 B.; gut near Fort de Wind: n. 835 B.; near Boven 200 M.: n. 1237 B.; White Wall: n. 240; Greatgut: n. 241.

Saba: Flat Point: n. 2134 B.

St. Martin: between Mount Vernon and Orientbay: n. 2925 B.; Tintamarre: n. 3403 B.

St. Croix ex Millspaugh.

From Mexico and the Antill. to Argentina. (E. a. P.).

203. **Passifloraceae**.5372. **Passiflora** Linn.

Passiflora foetida Linn. Spec. (1753) 959; Gris. Fl. 294; Duss 313; Millspaugh 512.

St. Eustatius: Fairplay: n. 2305; seashore: n. 1974, 2331; road near Sugarloaf: n. 2397.

Saba: Gut near Marypoint: n. 2048 B.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

Passiflora laurifolia Linn. Spec. (1753) 956; Gris. Fl. 293; Duss 312; Millspaugh 512. Bell apple.

St. Eustatius: Top of the Quill 400 M.: n. 249 B., 290 B.; in a garden in Oranjestad: n. 526 B.; slope of the Quill near Bengalen 250 M.: n. 701 B.; Cultuurvlakte: n. 1439; slope of the Quill: n. 1426; Boven: n. 85; locality unknown: n. 1429, 1442, 1443, 24 G., 79 H., 144 G.

Saba: Bottom: n. 2272 B.; Jenkygut—Hellsgate: n. 87; locality unknown: n. 2 L., 170 L.

St. Martin: Mount Paradis 300—400 M.: n. 3333 B.

St. Croix ex Millspaugh.

Antill., Nova Granata, Guiana, Brazil. (Gris. Fl.). Also cultivated.

Passiflora maliformis Linn. Spec. (1753) 956; Gris. Fl. 293; Duss 313.

Saba: Mountain 200 M.: n. 1904; locality unknown: n. 305 L. Antill., Haiti to Nova Granata (Gris. Fl.).

Passiflora quadrangularis Linn. Syst. ed. X. (1759) 1248. Granadilla.

Cultivated. [Antill., Panama, Peru, Brasil.]. (Gris. Fl.).

Passiflora rubra Linn. Spec. (1753) 956; Gris. Fl. 292; Duss 312; Millspaugh 512. Snakeberry wine.

Saba: between Bottom and Marypoint 300 M.: n. 1488 B., 1494 B.; Crispeen 400 M.: n. 1764 B.; gut between Hellsgate and Masticnt 300—400 M.: n. 2024 B.; Mountain 800 M.: n. 2217 B.

St. Martin: lower part of Mount Paradis: n. 3292 B.

St. Croix ex Millspaugh.

Antill., Panama, Venezuela, Nova Granata. (Gris. Fl.).

Passiflora suberosa Linn. Spec. (1753) 958; Gris. Fl. 290; Duss 311; Millspaugh 512.

St. Eustatius: Top of the Quill 400 M.: n. 192 B., 313 B., 342 B.; 500 M.: n. 430 B.; slope of the Quill near Bengalen 200 M.: n. 690 B.; slope of the lower part of the Quill near Bengalen: n. 713 B.; gut near Fort de Wind: 50 M.: n. 837 B.; slope of the Quill: n. 2417; locality unknown: n. 188 H., 271 G.

Saba: Windwardside—Hellsgate: n. 1668 B.; Bottom: n. 2546, 2571; locality unknown: n. 314 L.

St. Martin: Nakedboyhill 200—260 M.: n. 2721 B., 2734 B.; near Mulletpond: n. 2757 B., 2764 B.; in an orchard near Bellevue: n. 2834 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2871 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Panama, Venezuela. (Gris. Fl.).

205. Caricaceae.

5377. **Carica** Linn.

Carica Papaya Linn. Spec. (1753) 1026. Papao.

Cultivated. [Amer. trop.]. (E. a. P.).

208. Begoniaceae.

5397. **Begonia** Linn.

Begonia domingensis DC. in Ann. Sc. Nat. Sér IV. XI (1859) 124 (vel. affin.); Gris. Fl. 304.

St. Eustatius: Top of the Quill 450 M.: n. 382 B., 472 B.; Bengalen: n. 193, 194; Quill: n. 192.

Saba: Mountain 600—800 M.: n. 1787 B., 1809 B., 2212 B., 169, 189, 190, 191; locality unknown: n. 173 L.

Dominica, Haiti. (Gris. Fl.)

210. Cactaceae.

5401. **Cereus** Mill.

Cereus grandiflorus Mill. Gard. Dict. ed. VIII (1768) n. II; Gris. Fl. 301; Duss 317; Millspaugh 513.

St. Eustatius: Jenkinsbay; n. 1253a B.

Saba: Bottom: n. 2242a B.

St. Croix ex Millspaugh.

Jamaica, Caraiben, Haiti, Vera Cruz, Mexico, Sonora. (Schumann 1903).

Cereus Peruvianus Mill. Gard. Dict. ed. VIII (1768) n. 4.

St. Eustatius: in a garden in Oranjestad.

St. Martin: near Bethlehem: n. 3438a B.

Antill., Mexico, Bras., Guiana. (Schumann 1903).

Cereus triangularis Haworth. Syn. Pl. Succ. (1812) 180; Gris. Fl. 301; Duss 317; Millspaugh 513.

Cereus compressus Mill. Gard. Dict. ed. VIII (1768) n. 10. Slingereactus, Strawberry.

Saba: St. John's: n. 1871a B.; Windwardside: n. 1917a B.; near Devilshand: n. 2097a B.

St. Martin: near Oysterpond: n. 3089a B.; Mount Paradis: n. 3322a B.

St. Croix ex Millspaugh.

Antill., Florida, Mexico, Rio de Janeiro, Asia. (Schumann 1903).

Cereus spec.

St. Eustatius: n. 141, 153.

Saba: locality unknown: n. 1660a B.

5409. **Melocactus** Link et Otto.

Melocactus communis Link et Otto in Verh. Preuss. Ver. Gartenb. III (1827) 417 t. 11; Gris. Fl. 300; Duss 317; Millspaugh 513. Popehead.

St. Eustatius: I saw *Melocactus communis*: Top of Signahill; near Tumbledownickbay; Gilboohill.

Saba: I saw *Melocactus communis*: near the Sulphurmines.

St. Martin: I saw *Melocactus communis*: near Filipsburg; near Guana-bay; along roads from Filipsburg to Nakedboyhill; top of Nakedboyhill; Lowlands; Green Key.

St. Croix ex Millspaugh.

Antill. (Schumann 1903).

5417. **Opuntia** Mill.

Opuntia Ficus indica Mill. Gard. Dict. ed. VIII (1768) n. 2; Gris. Fl. 302; Sweet prickly.

St. Eustatius: along roads: n. 151 G.

I saw *Opuntia Ficus indica*: in deserted plantations near Bengalen; roads along the lower part of Signallhill; Signallhill.

Saba: I saw *Opuntia Ficus indica*: St. Giles; Laddergut.

St. Martin: I saw *Opuntia Ficus indica*: near Filipsburg; Nakedboy-hill; Fort Willem; Green Key.

Mexico, warmer countries of both hemisph. (Schumann 1903).

Opuntia spinosissima Mill. Gard. Dict. ed. VIII (1768) n. 8; Gris. Fl. 302; Duss 318; Millspaugh 513.

St. Eustatius: I saw *Opuntia spinosissima*: on the top of the Cliff near Signallhill; on the White Wall.

St. Martin: I saw *Opuntia spinosissima*: Top of Nakedboyhill; Tintamarre.

St. Croix ex Millspaugh.

Ind. Occid. (Schumann 1903). Baham. (Hitche.)

Opuntia triacantha Haworth. (1812); Gris. Fl. 302. Spanish lady.

St. Eustatius: English Quarter: n. 1004a B.; Gilboohill: n. 1126a B.; Jenkinsbay: n. 1230a B.

I saw *Opuntia triacantha*: Slope of Signallhill; roads along the lower part of Signallhill; top of the Cliff between Oranjestad and White Wall.

Saba: Paris: n. 1384a B.

I also saw *Opuntia triacantha*: Laddergut.

St. Martin: near Fort Willem: n. 2551a B.; near Nakedboy-hill: n. 2577a B.

I also saw *Opuntia triacantha*: near Filipsburg; along roads near Bethlehem; one of the islands in Simonsbaylagoon.

Ind. Occid. (Schumann 1903).

Opuntia Tuna Mill. Gard. Dict. ed. VIII. (1768) n. 3; Gris. Fl. 302; Duss 318; Millspaugh 513. Sour prickly.

St. Eustatius: along roads: n. 152.

Saba: I saw *Opuntia Tuna*: Giles gut; near the Sulphurmines; Green Island.

St. Martin: I saw *Opuntia Tuna*: Nakedboyhill; Fort Willem; one of the Islands in Simonsbaylagoon.

India Occid., California? (Schumann 1903).

5418. **Nopalea** Salm-Dyck.

Nopalea coccinellifera Salm-Dyck Cath. Hort. Dyck. ed. II (1850) 64. Curaçao prickly.

Cultivated. [Mexico]. (Schumann 1903).

5421. **Peireskia** Mill.

Peireskia aculeata Mill. Gard. Diet. ed. VIII (1768).

Cultivated. [warmer countries of Amerika]. (Schumann 1903).

Peireskia bleo DC. Prod. III (1828) 475.

Cultivated. [Nova Granata, Brasilia]. (Schumann 1903).

214. **Thymelaeaceae.**5449. **Daphnopsis** Mart. et Zucc.

Daphnopsis caribaea Gris. Flor. W. I., (1860) 278; Duss 297. Maho.

St. Eustatius: Top of the Quill 400 M.: n. 197 B., 251 B., 397 B.; slope of the Quill on the side of White Wall 200 M.: n. 1177 B.; Quill: n. 154.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1359 B.; Greathill 400 M.: n. 1386 B.; seaside of Tentguthill 100 M.: n. 1654 B.; Hellsgate: n. 1699 B.; Springbaygut: n. 2077 B.; Paris 2—400 M.: n. 233, 234, 235.

I saw *Daphnopsis caribaea*: Gut between Hellsgate and Mastiegut; Saddle.

St. Martin: Cul de Sac: n. 2604 B.; Milldrumhill 200—300 M.: n. 3119 B., 3124 B.; Mount Paradis 300—400 M.: n. 3240 B.

I also saw *Daphnopsis caribaea*: Top of Nakedboyhill; near Saline.

St. Croix. (Symb.).

Portorico, St. Thomas, St. Kitts, Martinique, Guadeloupe, Margareta, Dominica. (Herb. Krug et Urban).

216. **Lythraceae.**5493. **Lagerstroemia** Linn.

Lagerstroemia indica Linn. Spec. (1762) 734. Queen of Crib-flower.

Cultivated. [Japan, China, Molukk., Australia]. (Koehne 1903).

5494. **Lawsonia** Linn.

Lawsonia inermis Linn. Spec. (1753) 349. Miminet.

Cultivated. [Africa, Asia, Australia]. (Koehne 1903).

218. Punicaceae.5501. **Punica** Linn.

Punica Granatum Linn. (1753) 472. Pomegranate tree.
Cultivated. [Balkan and Himalaya]. (E. a. P.).

220. Rhizophoraceae.5523. **Rhizophora** Linn.

Rhizophora Mangle Linn. Spec. (1753) 443; Gris. Fl. 274; Duss 292; Millspaugh 514.

St. Martin: near Colebay (coll. George): n. 2498a B.; from
Filipsburg to Simonsbay: n. 3027 B.

I saw *Rhizophora Mangle*: at Simonsbaylagoon: Oysterpond.

St. Croix ex Millspaugh.
America. (E. a. P.).

221. Combretaceae.5541. **Quisqualis** Linn.

Quisqualis indica Linn. Spec. (1762) 556. Different coloured jessamine.

Cultivated. [Ind. orient., Ins. Philipp., Afr. trop. orient., Guinea]. (E. a. P.).

5544. **Terminalia** Linn.

Terminalia Catappa Linn. Mant. II (1771) 519. Almond tree.
Cultivated. [Madagascar, Malay arch., Nova Guinea]. (E. a. P.).

5548. **Conocarpus** Gärtn.

Conocarpus erectus Linn. Spec. (1753) 76; Gris. Fl. 277; Duss 295; Millspaugh 516. Button wood.

St. Eustatius: Seashore at Venusbay: n. 970 B.

St. Martin: from Filipsburg to Guanabay: n. 2378 B.; Colebay
(coll. George.): n. 2515 B.; shore of the Simonsbaylagoon near
Cul Piquard: n. 2747 B.; near Simonsbayvillage: n. 3029 B.,
3029 B.; Simonsbay: n. 160.; Lowlands: n. 3266 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Florida to Brazil., Galapagos, Marianne Islands, trop. coast of Africa. (Gris. Fl.).

5551. **Laguncularia** Gärtn.

Laguncularia racemosa Gärtn. f. Fruct. III (1805) 209 t. 217 f. 2; Gris. Fl. 276; Duss 295; Millspaugh 516.

St. Martin: Colebay (coll. George): n. 2497 B.; from Filipsburg to Cul de Sac along the Saltpond: n. 2612 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2748 B.; along the Simonsbaylagoon near the village: n. 3028 B., 3285 B.; Oysterpond n. 376.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Amer. trop. to Florida, trop. Africa. (E. a. P.).

222. Myrtaceae.

5559. **Psidium** Linn.

Psidium Guajava Linn. Spec. (1753) 470; Gris. Fl. 241; Duss 261; Millspaugh 515. Guava tree.

St. Eustatius: in a garden in Oranjestad: n. 512 B., 41 H., 48 G.; slope of the Quill: n. 62, 63, 64; road near English Quarter: n. 41.

Saba: Boobyhill 350 M.: n. 1615 B.; Ladder: n. 40; Longpeak: n. 61; locality unknown: n. 119 L.

I also saw *Psidium Guajava*: in a gut between Hellsgate and Mastiegut; between Bottom and Marypoint; Crespeenhill.

St. Martin: from Filipsburg to Belvedere: n. 2582 B.; Cul de Sac: n. 60.

I also saw *Psidium Guajava*: Marigothill to Maigot; Nakedboyhill; lower part of Milldrumhill; Mount Paradis 3—400 M.; Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill. from Mexico to Brasilia. (Urban 1895).

Amomis Berg.

cf. 5566. *Pimenta* Lindl. sect. I.

Amomis caryophyllata Kr. et Urb. in Urban. Additamenta ad Cognitionem Florae Indiae Occidentalis II (1895) 573; Duss 262.

Pimenta acris Kostel. Allg. med. pharm. Fl. IV (1835) 1526; Gris. Fl. 241; Millspaugh 515. Cinnamom tree, Bayberry.

St. Eustatius: Top of the Quill 500 M.: n. 493 B., 56, 57; Bengalen: n. 53.

Saba: between Bottom and Marypoint 300 M.: n. 1479 B., 1557 B.; Boobyhill 300 M.: n. 1596 B.; Bottom: n. 1987 B.; Great-hill: n. 54; Bottom: n. 51, 52; Mastiegut: n. 55.

St. Martin: Milldrumhill 300—400 M.: n. 3156 B.

St. Croix ex Millspaugh.

Venezuela, Guiana; cult. in Baham. and Ind. Orient. (Urb. 1895).

5573. *Myrcia* DC.

Myrcia paniculata Kr. et Urb. in Urban Additamenta ad Cognitionem Florae Indiae Occidentalis II (1895) 577; Duss 263.

Myrcia coriacea DC. Prodr. III (1828) 243; Gris. Fl. 234; Millspaugh 515. Red wood, Birdsberry.

St. Eustatius: Top of the Quill 400 M.: n. 173 B., 188 B., 260 B., 346 B., 505 B.; locality unknown: n. 114 H.

Saba: Paris 400 M.: n. 1369 B.; Mountain near Windwardside 600 M.: n. 2180 B.; locality unknown: n. 1568 B., 140 L.; Well-gut: n. 43.

St. Martin: Nakedboyhill 50—200 M.: n. 2715 B.; lower part of Mount Paradis: n. 3189 B.; Mount Paradis 300—400; n. 3221 B., 3352 B.; Lowlands: n. 3268 B.

St. Croix ex Millspaugh.

Antill., Brazil. (Urb. 1895).

Myrcia splendens DC. Prodr. III (1828) 244; Gris. Fl. 234; Duss 265. Surinaam cherries.

St. Eustatius: Top of the Quill 400 M.: n. 200 B., 285 B., 358 B.; inner slope of the Quill 350 M.: n. 420 B., 1218 B., 1222 B.; slope of the Quill near Bengalen 50—200 M.: n. 656 B.; slope of the Quill near Bengalen 150 M.: n. 684 B.; slope of the Quill 150 M.: n. 1112 B.; Quill: n. 44, 46, 47; slope of the Quill: n. 45; Little Mountain: n. 44; locality unknown: n. 177 G., 213 G.

Saba: along roads 200—300 M.: n. 1711a B.; Windwardside: n. 1945 B.; Hellsgate—Marypoint: n. 2043 B.; Crespeen—Rendezvous: n. 48.

St. Martin: Mount Paradis 150—400 M.: n. 3195 B.; Tintamarre: n. 3396 B.

Cuba, Jamaica, Haiti, St. Domingo, Portorico, St. Thomas, Guadeloupe, Martinique, St. Lucia, St. Vincent, Tobago, Trinidad. (Urban 1895).

5575. **Calyptranthes** Swartz.**Calyptranthes Boldinghii** Urb. in *Symbolae Antillana* VI (1909) 22.**St. Martin:** Lowlands: n. 3270 B.5578. **Eugenia** Linn.**Eugenia axillaris** Willd. Spec. II (1800) 970; Gris. Fl. 236; Duss 270; Millspaugh 514. Choakberry, Pigeonberry.**St. Eustatius:** Slope of the Quill near Bengalen 50—200 M.: n. 641 B.; Gilboohill 150 M.: n. 1120 B.; Boven: n. 19, 21; top of the Quill: n. 20; locality unknown: n. 55 H.**Saba:** Top of the Cliff near the seaside of Boobyhill: n. 1607a B.; Mountain 400 M.: n. 1740 B.; Windwardside: n. 1883 B.; Laddergut: n. 2000 B.; Springbaygut: n. 2117 B.; Greathill: n. 22.**St. Martin:** from Filipsburg to Guanabay: n. 2392 B.; from Belvedere to Oysterpond: n. 3055 B., 3072 B.; Lowlands: n. 3263 B.; Tintamarre: n. 3388 B.; top of Centryhill: n. 39.**Anguilla** Boldingh.**St. Croix** ex Millspaugh.

Bermud., Florida, Key, Baham., Antill. (Urban 1895).

Eugenia cordata DC. Prodr. III (1828) 272; var. **Sintensii** Kr. et Urb. in *Additamenta ad Cognit. Fl. Ind. Occ.* (1895) 656.*Eugenia sessiliflora* DC. Prodr. III (1828) 273; Gris. Fl. 236.**St. Eustatius:** from Signalhill to Venusbay: n. 960 B.; Gilboohill: n. 1121 B.; near White Wall: n. 1266 B.**St. Croix.** (Symb.).

St. Domingo? Portorico, St. Thomas, St. Jan, Tortola, Antigua, Martinique, St. Lucia, St. Vincent, Mustique Island, Becquia Island. (Urban 1895).

Eugenia floribunda West St. Croix (1793) 290; Gris. Fl. 239; Duss 273; Millspaugh 514. Guaveberry.**St. Eustatius:** locality unknown: n. 200a G.; 40.**St. Martin:** in an orchard near Bellevue: n. 2831 B.; Mount Paradis 300—400 M.: n. 3320a B.**St. Croix.** (Urban 1895).

Antill., Guiana, Brazil. (Urban 1895).

Eugenia fragrans Willd. Spec. (1800) 964.*Ananomis fragrans* Gris. Flor. W. I. (1860) 240.*Ananomis punctata* Gris. Flor. W. I. (1860) 240.**St. Martin:** Top of Nakedboyhill: n. 2715 B.

Florida, Antill. (Urban 1895).

Eugenia Jambos Linn. Spec. (1753) 470.

Jambosa vulgaris DC. Prod. III (1828) 286; Gris. Fl. 235; Duss 274.

Jambosa Jambos Millspaugh 515. Plum rose.

St. Eustatius: Slope of the Quill near Bengalen 50—200 M.: n. 644 B., 26; locality unknown: n. 103 H.

Saba: Crespeen—Rendez-vous 400 M.: n. 27; locality unknown: n. 109 L., 321 L.

I also saw *Eugenia Jambos*: Bottom; Laddergut.

St. Martin: Mount Paradis 300—400 M.: n. 3347 B.

St. Croix ex Millspaugh.

Indig. in Asia austro-orient. and Australia. (Urban 1895).

Also cultivated.

Eugenia ligustrina Willd. Spec. II (1800) 962; Gris. Fl. 239; Duss 268; Millspaugh 514. Blackberry, Blackcherry.

St. Eustatius: deserted plantations near Tumbledown dickbay: n. 148 B.; slope of the Quill 350 M.: n. 392 B.; top of the Quill 500 M.: n. 473 B., 497 B.; hill near Concordia 100 M.: n. 952 B.; from Little Mountain to Venusbay: n. 986 B., 992 B.; slope of Signalhill: n. 28; Quill: n. 29; Venusbay: n. 30; top of the Quill: n. 31, 32; locality unknown: n. 109 H.

St. Martin: from Philipsburg to Guanabay: n. 2411 B.; Prince quarter (coll. George): n. 2484 B.; from Philipsburg to Belvedere: n. 2584 B.; Cul de Sac: n. 2607 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2946 B., 2952 B.; Mill-drumhill 300—400 M.: n. 3137 B.; Lowlands: n. 3280 B.; near Fort Amsterdam: n. 36; Vineyardhill: n. 33, 35; Prince quarter: n. 34; Guanabay: n. 34 b.

St. Croix ex Millspaugh.

Baham., Antill., Brazil. (Urban 1895).

Eugenia monticola DC. Prodr. III (1828) 275.

Eugenia Poirerii Berg. in Linnaea XXVII (1856) 186 (excl. syn. Cand., Poir., Spreng.); Gris. Fl. 236; (ex descr.).

Eugenia obtusata Willd. ex Berg. in Linnaea XXVII (1856) 240; Gris. Fl. 237; (ex descr.).

Eugenia buxifolia Gris. Flor. W. I. (1850) 236; (excl. syn.), — non Willd.

Eugenia pallens Gris. Flor. W. I. (1860) 237; (specim. Guad.), — non DC. (Symb.). Duss 269; Millspaugh 515. Birdsberry.

St. Martin: Philipsburg near the Saltpond: n. 2365 B.; Naked-boyhill 200 M.: n. 2585 B.; Marigothill: n. 2609 B.; near Simsons-baylagoon: n. 2745 B.; near Mulletpond: n. 2768 B.; hill between Mont des Accords and Centryhill: n. 2863 B.; near Oysterpond: n. 3073 B.; Milldrumbill: n. 3116 B.; Mount Paradis: n. 3220 B., 3330 B.; Lowlands: n. 3283 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham. (Hitche.); Antill. (Urban 1899).

Eugenia procera Poir. in Lam. Enc. Suppl. III (1813) 129; Gris. Fl. 238; Millspaugh 515. Bastard black-berry.

St. Eustatius: deserted plantations near Tumbledown-dickbay: n. 149 B.; Signallhill 200 M.: n. 1037 B.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1356 B.; Springbaygut: n. 2098 B.; Peperpot: n. 38.

St. Martin: along roads from Grande Case to Marigot; n. 3100 B.; Mount Paradis 300—400 M.: n. 3321 B.; Colebayhill: n. 37.

St. Croix ex Millspaugh.

Cuba, Jamaica, Haiti, St. Domingo, Portorico, St. Thomas, St. Jan, Antigua, Martinique, Barbados. (Urban 1895).

Eugenia rhombea Kr. et Urb. in Urban. Additamenta ad Cognitionem Florae Ind. Occid. III (1895) 665.

St. Eustatius: Seinpost: n. 39; Wittehoek: n. 40.

Florida, Key West, Baham., Antill. (Urban 1895).

Eugenia uniflora Linn. Spec. (1753) 470; Gris. Fl. 239. Honey-berry, Surinaam cherries.

St. Eustatius: Slope of the lower part of Signallhill: n. 117 B.; deserted plantations near Tumbledown-dickbay: n. 143 B.; top of the Quill 400 M.: n. 172 B., 177 B., 326 B.; in a garden in Oranjestad: n. 861 B., 55a B., 76 H.

Saba: lower part of the Bottom Mountain 250 M.: n. 1424 B.; Crispeen 400 M.: n. 1720 B.; locality unknown: n. 70 L.

St. Croix ex Millspaugh.

Florida, Antill. (Urban 1895).

223. Melastomataceae.

5633. **Tibouchina** Aubl.

Tibouchina strigosa Cogn. in Monogr. Phan. VII. (1891) 266.

Chaetogastra strigosa DC. Prod. III (1818) 134; Gris. Fl. 266.

Chaetogastra chironioides Gris. Flor. W. I. (1860) 267 (p. p.); Duss 288.

Tibouchina ornata Baill. Adansonia XII (1877) 74.

Saba: Top of the Mountain 800 M.: n. 18.

Antill., British Guiana. (Cogniaux 1891).

5749. **Charianthus** D. Don.

Charianthus crinitus Naud. in Ann. Sc. Nat. Sér. 3, XVIII (1852) III; Gris. Fl. 264.

Charianthus nodosus Triana Melastom. in Trans. Linn. Soc. XXVIII (1871) 99, var *crinitus* Naud; Duss 286.

Saba: Mountain 600—800 M.: n. 1784 B.; 187 L., 197 L., 203 L. 1, 2, 3.

Antill. (Cogniaux 1891).

5758. **Tetrazygia** Rich.

Tetrazygia discolor DC. Prod. III (1828) 172; Gris. Fl. 255; Duss 280.

St. Eustatius: Top of the Quill: n. 16, 17.

Saba: lower part of the Bottom Mountain 300 M.: n. 1436 B.; between Bottom and Marypoint 100—200 M.: n. 1551 B.; Gainbygut: n. 14, 15.

Antill., French Guiana (Cogniaux 1891).

5759. **Miconia** Ruiz. et Pav.

Miconia impetiolaris D. Don. in Mem. Wern. Soc. IV (1823) 316; Gris. Fl. 256; Duss 284; Millspaugh 516.

St. Eustatius: Top of the Quill 400 M.: n. 310 B.; Bottom of the Quill 300 M.: n. 469 B.

St. Croix ex Millspaugh.

Antill., Bolivia, Panama, Costarica, Honduras, Guatamala, Mexico. (Cogniaux 1891).

Miconia laevigata DC. Prod. III (1828) 188; Gris. Fl. 257; Duss 281; Millspaugh 516.

St. Eustatius: Top of the Quill 400 M.: n. 208 B., 262 B., 338 B., 344 B., 506 B.; slope of the Quill 350 M.: n. 388 B.; Bottom of the Quill 300 M.: n. 452 B.; slope of the Quill near Bengalen 200 M.: n. 742 B.; slope of the Quill: n. 9, 10, 11.

Saba: lower part of the Bottom Mountain 250 M.: n. 1406 B.; Windwardside—Hellsgate: n. 1682 B., 1683 B.; Mountain 800 M.: n. 1825a B., 2216 B.; Springbaygut: n. 2119 B.; Peperpot: n. 8;

Mountain: n. 5; Crespeen--Rendez-vous 400 M.: n. 6, 7; locality unknown: n. 41 L., 216 L., 306 L.

St. Martin: Top of the hill on the northern side of Cul de Sac 200--300 M.: n. 3006 B.; along roads from Grande Case to Marigot: n. 3100a B.; Milldrumhill: n. 3180 B.; Mount Paradis 300--400 M.: n. 3208 B., 3341 B.

St. Croix ex Millspaugh.

Antill., Mexico, Guatanala, Costarica, Venezuela, Surinam, Guian., Brazil. (Cogniaux 1895).

227. Araliaceae.

5883. **Panax** Linn.

Panax fruticosum Linn. Spec. (1763) 1513.

Cultivated. [Asia trop., Malaya, Polynes.] (Ind. Kew.).

228. Umbelliferae.

5923. **Eryngium** Linn.

Eryngium foetidum Linn. Spec. (1753) 232; Gris. Fl. 308; Duss 324.

Saba: between Bottom and Marypoint 100--200 M.: n. 1537 B.; Mountain near Windwardside 600 M.: n. 2176 B.; locality unknown: n. 37 L.

St. Martin: Top of the hill on the northern side of Cul de Sac 200--300 M.: n. 2985 B.; Mount Paradis 200--400 M.: n. 3205 B.

Antill., Cuba and Panama to Nova Granata and Brazil. (Gris. Fl.).

6062. **Foeniculum** Adans.

Foeniculum vulgare Mill. Gard. Dict. ed. VIII (1768) n. 1.

Cultivated. [Europa]. (Ind. Kew.).

235a. Theophrastaceae.

6282. **Jacquinia** Linn.

Jacquinia armillaris Jacq. Enum. Pl. Carib. (1760) 15; Gris. Fl. 397; Duss 383; Millspaugh 517. Picrous bark.

St. Eustatius: White Wall: n. 807 B.; hill near Concordia: n. 953 B.; seashore near Venusbay: 969 B., 284, 285.

St. Martin: from Filipsburg to Guanabay: 2418 B.; Prince quarter (coll. George): n. 2508 B.; near Mulletpond: n. 2774 B.; Mountain between Mont des Accords and Centryhill: n. 2908 B.; between Mount Vernon and Orientbay: n. 2929 B.; from Belvedere to

Oysterpond: n. 3061 B.; Mount Paradis 200—300 M.: n. 3304 B.; near Colebayhill: n. 65, 137; Prince quarter: n. 606; Guanabay: n. 63, 192.

St. Croix ex Millspaugh.

Cuba, Jamaica, St. Domingo, Portorico, St. Thomas, St. Jan, St. Barthélemy, Montserrat, Guadeloupe, Dominica, St. Vincent, Mustique, Cannouan, Barbados, Grenada, Tobago, Curaçao, Bonaire. (Symb.).

Jacquinia Berteri Spreng. Syst. I (1825) 668.

St. Martin: near Mulletpond: n. 2781 B.; Lowlands: n. 3283a B.

Anguilla Boldingh.

Baham., Antill. (Symb.).

236. Myrsinaceae.

6285. **Ardisia** Swartz.

Ardisia coriacea Sw. Prod. (1788) 48; Millspaugh 517.

Ardisia guadalupensis Duchass. et Walp. apud Gris. Flor. W. I. (1861) 396; Duss 382. Bastard cinnamom.

St. Eustatius: Top of the Quill 400 M.: n. 196 B., 244 B., 399 B., 44; locality unknown: n. 30, 45.

Saba: Hellsgate: n. 1704 B.; gut between Masticgut and Hellsgate 300—400 M.: n. 2032 B.; Jenkygut: n. 46; Hellsgate: n. 43; locality unknown: n. 76 L., 183 L.

St. Croix ex Millspaugh.

Portorico, St. Thomas, St. Jan, Tortola, Guadeloupe, Dominica, Martinique, St. Lucia. (Symb.).

6314. **Rapanea** Aubl.

Rapanea ferruginea Mez in Symbolae Antillanae II (1901) 429.

Myrsine laeta Gris. Flor. W. I. (1861) 392.

Saba: Top of the Mountain 800 M.: n. 1836a B.; Rendez-vous Mountain 600 M.: n. 2150a B.; Mountain near Hellsgate: n. 2278 B. Antill. Amer. trop. from Mexico to Bolivia. (Symb.).

238. Plumbaginaceae.

6342. **Plumbago** Linn.

Plumbago capensis Thunb. Prod. I Pl. Cap. (1794) 33. Quakerblossom.

Cultivated. [Afr. australis]. (Ind. Kew.).

Plumbago scandens Linn. Spec. (1762) 215; Gris. Fl. 390; Duss 380; Millspaugh 517. Blisterbush.

St. Eustatius: Seashore near Oranjestad: n. 13 B., 42 B.; along roads in Oranjestad: n. 561 B.; in deserted plantations at the seashore near Bengalen: n. 592 B.; gut near Fort de Wind: n. 845 B.; Cultuurvlakte: 354; locality unknown: n. 31 H., 35 G.

I also saw *Plumbago scandens*: along the lower part of Signalhill; top of Signalhill.

Saba: Bottom: n. 1852 B.; Laddergut 200 M.: n. 1995a B.; Peperpot: n. 149, 356; Ladder: n. 352; locality unknown: n. 319 L.

I also saw *Plumbago scandens*: Forthbaygut.

St. Martin: along roads in Prince quarter: n. 2933 B.

St. Croix ex Millspaugh.

Antill., Mexico to Brazil and Chili, Galapagos. (Gris. Fl.).

239. Sapotaceae.

6361. *Achras* Linn.

Achras sapota Linn. Syst. X ed. II (1759) 988; Millspaugh 517.

Sapota Achras Mill. Gard. Dict. VIII ed. (1768) n. 1; Gris. Fl. 399; Duss 385. Mispel, Sapodille.

St. Eustatius: lower part of Signalhill: n. 115 B.

St. Croix ex Millspaugh.

Baham., Antill., Curaçao from Mexico to Guiana. (Symb.).

Also cultivated.

6363. *Lucuma* A. DC.

Lucuma multiflora A. DC. Prodr. VIII (1844) 168.

Saba: near Hellsgate: n. 1675 B.; Windwardside: n. 2313 B.

St. Croix. (Symb.).

Portorico, St. Thomas, St. Kitts, Montserrat, Guadeloupe, Martinique, St. Lucia, St. Vincent, Grenada, Trinidad. (Symb.).

6368. *Sideroxylon* Linn.

Sideroxylon foetidissimum Jacq. Enum. (1760) 15.

Sideroxylon mastichodendron Jacq. Coll. II (1788) p. 253 t. 17 f. 5; Gris. Fl. 399; Duss 386; Millspaugh 518. Mastictree.

Saba: Hellsgate: n. 2049 B.; Masticgut: n. 46; Jenkygut: n. 40.

St. Croix ex Millspaugh.

Florida, Key, Baham., Antill. (Symb.).

6373. **Dipholis** A. DC.

Dipholis salicifolia A. DC. Prodr. VIII (1844) 188; Gris. Fl. 401; Duss 388; Millspaugh 518.

St. Eustatius: Slope of the Quill on the side of White Wall 200 M.: n. 1168 B.

St. Croix ex Millspaugh.

Florida, Key, Baham., Antill., Yucatan. (Symb.).

6374. **Bumelia** Sw.

Bumelia obovata A. DC. Prodr. VIII (1844) 191.

Bumelia cuneata Sw. Flor. I (1797) 496; Gris. Fl. 401; Duss 613; Millspaugh 517.

St. Eustatius: from Little Mountain to Venusbay: 985 B.; locality unknown: n. 338 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1535 B.

St. Martin: near Mulletpond: n. 2765 B.; at the seashore of Simonsbay: n. 3035 B.; Colebayhill: n. 49.

St. Croix ex Millspaugh.

Jamaica, St. Domingo, Portorico, St. Thomas, Tortola, St. Barthélemy, Antigua, Montserrat, Guadeloupe, Desirade, Martinique, St. Lucia. (Symb.).

6377. **Chrysophyllum** Linn.

Chrysophyllum argenteum Jacq. Enum. (1760) 15; Duss 385.

Chrysophyllum glabrum Jacq. Enum. (1760) 15; Gris. Fl. 398; Duss 385.

Saba: Gut near Devilshand: n. 2115 B.

Cuba, Haiti, St. Domingo, Portorico, Antigua, Montserrat, Guadeloupe, Marie-Galante, Dominica, Martinique, St. Vincent, Barbados, Grenada, Trinidad. (Symb.).

242. **Symplocaceae,**6418. **Symplocos.** Jacq.

Symplocos martinicensis Jacq. Enum. Pl. Carib. (1760) 24; Gris. Fl. 403; Duss 389. Blue berry.

St. Eustatius: Top of the Quill 400 M.: n. 297 B.; top of the Quill 500 M.: n. 474 B.; locality unknown: n. 126, 127, 128, 129.

Saba: Mountain near Windwardside 600 M.: n. 2152 B., 2185 B.

Portorico, St. Thomas, St. Kitts, Montserrat, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Grenada, Trinidad. (Brand 1901).

243. Oleaceae.

6427. **Forestiera** Poir.

Forestiera Eggersiana Kr. et Urb. in Bot. Jahrb. XV (1893) 339.

St. Martin: Prince quarter: n. 1, 2.

St. Thomas. (Urban 1893).

6428. **Linociera** Swartz.

Linociera compacta R. Br. Prod. (1810) 523; Gris. Fl. 405. Bridgo tree.

Mayepea caribaea O. Ktze Rev. Gen. (1891) 411; Duss 391; Millspaugh 518.

St. Eustatius: Top of the Quill 400 M.: n. 181 B., 261 B., 301 B., 343 B., 352 B.; slope of the Quill near Bengalen 250 M.: n. 700 B.; slope of the Quill near Glassbottle 150 M.: n. 1113 B., 1117 B.; Quill: n. 345; slope of the Quill: n. 343, 344; top of the Quill: n. 336, 337; locality unknown: n. 77 H., 169 G., 40.

I also saw *Linociera compacta*: Gilboohill.

Saba: lower part of the Bottom Mountain 250 M.: n. 1419 B., 1422 B; Jenkeygut: n. 340.

St. Martin: Milldrumhill: n. 3151 B.

St. Croix ex Millspaugh.

Antill., Guiana. (Urban 1893).

6440. **Jasminum** Linn.

Jasminum azoricum Linn. Spec. (1753) 7.

Cultivated. [Ins. Azor. and Madeira]. (E. a. P.).

Jasminum pubescens Willd. Sp. Pl. I (1798) 37.

Cultivated. [Asia trop.] (Ind. Kew.).

Jasminum undulatum Ker. Gawl. in Bot. Reg. t. 436. Orebian jessamine.

Cultivated. [Himalaya, China]. (Ind. Kew.).

245. Loganiaceae.

6453. **Spigelia** Linn.

Spigelia Anthelmia Linn. Spec. (1753) 149; Gris. Fl. 331; Duss 336.

St. Eustatius: I saw *Spigelia Anthelmia* at Bengalen.

Saba: lower part of the Bottom Mountain 300 M.: n. 1427 B.; between Bottom and Marypoint 100—200 M.: n. 1522 B., 1529 B.; gut between Hellsgate and Masticgut: 300—400 M.: n. 2050 B.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2876 B.

Anguilla Boldingh.

Antill., Amer. austral. (E. a. P.).

246. Gentianaceae.

6541. **Leiphaimos** Cham. et Schlecht.

Leiphaimos aphylla Gilg in Engl. u. Prantl. Nat. Pfl. fam. IV. 2. (1895) 104.

Voyria uniflora Pers. Syn. I (1805) 284; Gris. Fl. 425; Duss 402.

Voyria aphylla Pers. Syn. I (1805) 284.

St. Eustatius: Top of the Quill 400 M.: n. 296 B.; 450 M.: n. 365 B., 402 B.; top of the Quill: n. 40.

Saba: Mountain 800 M.: n. 1831 B., 2226 B.

Antill., Guiana, Brazil. (E. a. P.).

247. Apocynaceae.

6560. **Allamanda** Linn.

Allamanda cathartica Linn. Mant. II (1771) 214. Yellow bell. Cultivated. [Antill., Amer. centr. Brazil.] (E. a. P.)

6578. **Plumiera** Linn.

Plumiera alba Linn. Spec. (1753) 210; Gris. Fl. 411; Duss 395; Millspaugh 518. White frangepane, Pigeonwood.

St. Eustatius: in a garden in Oranjestad: n. 49 H., 67 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1354 B.

St. Martin: from Philipsburg to Guanabay: n. 2374 B.; Prince-quarter (coll. George): n. 2464 B.; Fort Willem: n. 2534 B.; near Fort Amsterdam: n. 237, 238.

I also saw *Plumiera alba*: along roads from Philipsburg to Simonsbay; from Belvedere to Oysterpond; lower part of Milldrumhill; Lowlands: Tintamarre; gut from Bethlehem to Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Portorico, St. Thomas, Guadeloupe, Martinique, St. Lucia, Grenada. (Herb. Krug et Urban).

Plumiera rubra Linn. Spec. (1753) 209. Red frangepane.
Cultivated. [Mexico and Venezuela]. (E. a. P.).

6597. **Lochnera** Reichb.

Lochnera rosea Reichb. Comp. (1828) 134.

Vinca rosea Linn. Syst. ed. X (1759) 944; Gris. Fl. 410; Duss 395; Millspaugh 519. Churchyard blossom.

St. Eustatius: Seashore near Oranjestad: n. 36 B.; along roads in Oranjestad: n. 571 B.; slope of the Quill: n. 252; in a garden. n. 38; locality unknown: n. 22 G., 60 H., 330 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1349 B.; Springbaygut: n. 2104 B.; seashore near Forthbaylanding: n. 2146 B.; Ladder: n. 254; locality unknown: n. 74 L., 90 L.

St. Martin: along the saltpond near Filipsburg: n. 2440 B.; Colebay (coll. George): n. 2510 B.

I also saw *Lochnera rosea*: along roads near Bethlehem; along roads near Grande Case.

St. Croix ex Millspaugh.

In all trop. countries. (E. a. P.).

6603. **Tabernaemontana** Linn.

Tabernaemontana citrifolia Jacq. Sel. Am. St. Hist. (1763) t. 175 f. 13. p. 38; Gris. Fl. 409; Duss 395. Milky tree.

St. Eustatius: Top of the Quill 400 M.: n. 234 B.; inner slope of the Quill 350 M.: n. 464 B.; in a garden in Oranjestad: n. 527 B.; slope of the Quill 300 M.: n. 886 B.; slope of the Quill near Glassbottle: n. 1207 B.; top of the Quill: n. 244, 245, 246.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1357 B.; Windwardside—Hellsgate: n. 1676 B.; Springbaygut: n. 2121 B.; Crispeen—Rendez-vous 400 M.: n. 241; Hellsgategut: n. 243; Marypointgut: n. 242.

St. Martin: near Bethlehem: n. 2671 B.; Mountain between Mont des Accords and Centryhill 0—200 M.: n. 2867 B.; Mount Paradis 300—400 M.: n. 3213 B., 3325 B.

Antill., Mexico. (Gris. Fl.).

Tabernaemontana coronaria Willd. Enum. Hort. Berol. (1809) 275.
Cultivated. [India orient.] (Ind. Kew.).

Tabernaemontana Cumingiana A. DC. Prod. VIII (1844) 373.
Cultivated. [Ins. Philipp.] (Ind. Kew.).

6619. **Rauwolfia** Linn.

Rauwolfia Lamarkii DC. Prod. VIII. (1844) 337; Gris. Fl. 408; Duss 393; Millspaugh 519. Snakeberry tree, Billy ache, Antigua balsam.

St. Eustatius: Top of the Quill 500 M.: n. 480 B.; slope of the Quill near Bengalen 50—200 M.: n. 647 B., 739 B.; gut near Fort de Wind 50 M.: n. 840 B.; along roads in Oranjestad: n. 935 B., 1069 B.; near Boven: 200 M.: n. 1235 B., 240; slope of the Quill 200 M.: n. 239; near Roundhill: n. 37; locality unknown: n. 27, 237 G., 203 H.

Saba: between Bottom and Marypoint 300 M.: n. 1468 B.; locality unknown: n. 270 L.

St. Martin: from Philipsburg to Guanabay: n. 2361 B.; Fort Willem 50 M.: n. 2541 B.; Nakedboyhill 50—200 M.: n. 2684 B.; in pastureground near Bellevue: n. 2812a B., 2825 B.; lower part of Milldrumhill: n. 3164 B.

I also saw *Rauwolfia Lamarkii*: along roads from Philipsburg to Nakedboyhill; near Saline; in pasture ground near Belvedere; Lowlands; Mount Paradis 100—200 M.; Tintamarre.

St. Croix ex Millspaugh.
Antill., Equador. (Gris. Fl.).

6632. **Thevetia** Adans.

Thevetia neriifolia Juss. ex Steud. Nom. Ed. II. II (1841) 680.
Cultivated. [Antill. and Mexico to Brazil.] (E. n. P.).

6661. **Urechites** Müll.—Arg.

Urechites suberecta Müll.—Arg. in Linnaea XXX (1859-60) 444.
Echites suberecta Jacq. Enum. Pl. Carib. (1760) 13; Gris. Fl. 415; Millspaugh 518. Bitterpot.

St. Eustatius: on the top of the Cliff between Oranjestad and Fort de Wind: n. 759 B., 776 B.; Signalhill 200 M.: n. 1045 B.; Cultuurylakte: n. 248; near Brookswell: n. 247; near Sugarloaf: n. 249; locality unknown: n. 184 H.

Saba: lower part of the Bottom Mountain 250 M.: n. 1395 B.

St. Martin: Fort Willem 50 M.; n. 2549 B.; near Fort Amsterdam: n. 250.

I also saw *Urechites suberecta*: Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Cuba, Jamaica, Hispaniola, Portorico, St. Thomas, St. Kitts.
(Herb. Krug et Urban).

6687. **Nerium** Linn.

Nerium Oleander Linn. Spec. (1753) 209. Oleander.

Cultivated. [From Portugal to Mesopotamia]. (E. a. P.).

248. Asclepiadaceae.

6736. **Cryptostegia** R. Br.

Cryptostegia grandiflora R. Br. in Bot. Reg. t. 435 (1819).

Cultivated. [Patria ignot.] (Symb.).

6791. **Asclepias** Linn.

Asclepias nivea Linn. Spec. (1753) 215; Gris. Fl. 419.

Asclepias Curassavica Linn. Spec. (1753) 215; Gris. Fl. 419;
Duss 398; Millspaugh 519. Ipecacuanha.

St. Eustatius: Locality unknown: n. 1 H., 134 G.; slope of the
Quill: n. 52, 222, 223.

Saba: between Bottom and Marypoint 100—200 M.: n. 1517a B.;
Paris 200—400 M.: n. 221; locality unknown: n. 160 L.

St. Martin: Milldrumhill 200—300 M.: n. 3120 B.; Mount
Paradis 300—400 M.: n. 3329 B.; Mary's Fancy: n. 220.

St. Croix ex Millspaugh.

Baham., Antill.; widely dispersed in both hemisph. (Symb.).

6792. **Calotropis** R. Br.

Calotropis procera R. Br. in Ait. Hort. Kew. Ed. 2 (1811) II
78; Gris. Fl. 420; Duss 399; Millspaugh 519. Liberty tree,
Sprain leaf.

St. Eustatius: Top of the Quill 500 M.: n. 475 B.; in deserted
plantations near English Quarter: n. 998 B.; locality unknown:
n. 43 H., 54 G.

Saba: along the seashore near Forthbaylanding: n. 1646 B.;
locality unknown: n. 113 L.

I also saw *Calotropis procera*: Flat Point.

St. Martin: near Simonsbayvillage: n. 3255 B.; locality un-
known: n. 3610 B.

I also saw *Calotropis procera*: on Mount Paradis 100—200 M.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Indig. in Africa sept. and Ind. Orient. (Symb.).

6798. **Metastelma** R. Br.

Metastelma parviflorum R. Br. in Wern. Soc. I (1809) 52; Duss 398.

Metastelma Paralias Dene in DC. Prodr. VIII (1844) 514; Gris. Fl. 417.

Metastelma Schlechtendalii Gris. Flor. W. I. (1861) 417; (nec Dene); (Symb.); Millspaugh 519.

St. Martin: juxta salinas prope Philipsbourg, Febr. 1821: Pleë; (Symb.).

Lowlands: n. 3267a B.

Baham. (Hitche.); Antill. (Symb.).

6907. **Hoya** R. Br.

Hoya carnosa R. Br. in Mém. Wern. I (1809) 27.

Cultivated. [Ins. Malay.] (Symb.).

6938. **Ibatia** Dene.

Ibatia maritima Dene. in DC. Prodr. VIII (1844) 599; Millspaugh 519.

Ibatia muricata Gris. Flor. W. I. (1861) 421; Duss 400.

Lachnostoma maritimum Nichols. Dict. Gardn. II 236. Maria-pompoeën.

St. Eustatius: Bengalen: n. 634 B.; near Boven 200 M.: n. 1232 B.; locality unknown: n. 225 H., 277 G.

St. Martin: from Philipsburg to Guanabay: n. 2400 B.; near Fort Willem 50 M.: n. 2789 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2861 B.

I also saw *Ibatia maritima*: Mount Paradis 100—200 M.

St. Croix ex Millspaugh.

Haiti, St. Domingo, St. Thomas, Antigua, Guadeloupe, Dominica, Martinique, St. Vincent. (Symb.).

249. **Convolvulaceae.**

6968. **Cuscuta** Linn.

Cuscuta americana Linn. Spec. (1753) 124; Gris. Fl. 476; Duss 443; Millspaugh 519. Love wine.

St. Eustatius: in deserted plantations at the seashore near Bengalen: 587 B.; Cultuurvlakte: n. 141; seashore: n. 140; locality unknown: n. 99 H., 186 G.

Saba: lower part of the Bottom Mountain 250 M.: n. 1413 B.; locality unknown: n. 106 L.

I also saw *Cuscuta americana* in Laddergut.

St. Martin: from Philipsburg to Guanabay: n. 2349 B.; from Philipsburg to Belvedere: n. 2578 B.; between Mont Chambord and the Saline: n. 2922 B.; Cul de Sac: n. 295.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

Cuscuta umbellata H. B. et K. Nov. Gen. et Spec. III (1818) 121; Gris. Fl. 476.

Saba: near Vischvijver: n. 2922 B.

Jamaica, New Mexico to Brazil (Gris. Fl.).

6973. ***Evolvulus*** Linn.

Evolvulus argyraeus Choisy Conv. Rar. (1838) 153.

St. Martin: Tintamarre: n. 3371 B.

Equador. (Ind. Kew.).

Evolvulus glaber Spreng. Syst. I (1825) 862.

Evolvulus mucronatus Sw. ex Wikst. in Vet. Acad. Handl. Stockholm (1827) 61; Gris. Fl. 475; Millspaugh 520.

St. Martin: along roads from Bethlehem to Mont Chambord: n. 2916 B.; between Belvedere and Oysterpond: n. 3066 B.; near Fort Amsterdam: n. 150; locality unknown: n. 151.

St. Croix ex Millspaugh.

Baham., Antill., Portorico to Peru. (Gris. Fl.).

Evolvulus sericeus Sw. Prod. Veg. Ind. Occ. (1788) 55; Gris. Fl. 475; Duss 442.

Evolvulus incanus Pers. Syn. I (1805) 288.

St. Eustatius: Top of the Quill: n. 380 B.; top of the Cliff between Oranjestad and Fort de Wind: n. 760 B.; White Wall: n. 43.

St. Martin: along roads from Colebayhill to Simonsbay: n. 3014 B.; Fort Willem: n. 2793a B.; Tintamarre: n. 3371 B.

Baham., Antill., Texas to Montevideo. (Gris. Fl.).

6986. ***Porana*** Burm.

Porana paniculata Roxb. Hort. Beng. (1814) 13. White coralita. Cultivated. [India orient. Malay.] (Ind. Kew.).

6991. **Jacquemontia** Choisy.

Jacquemontia pentantha G. Don. Gen. Syst. IV (1838) 283; Millspaugh 521.

Jacquemontia violacea Choisy in Mém. Soc. Phys. Genèv. VIII 1. (1838) 61; Duss 442.

Convolvulus pentanthus Jacq. Coll. IV. 210; Gris. Fl. 474.

St. Eustatius: Seashore near White Wall: n. 852 B.; near Concordia: n. 948 B.; Gilboohill: n. 1119 B., 1128a B.; Little Mountain: n. 1163a B.; Golden Rock: n. 55; locality unknown: n. 222 H.

Saba: Boobyhill: n. 1576 B.

Portorico, Bequia, St. Thomas, Barbuda, Guadeloupe, Martinique, St. Vincent, Grenada, Barbados. (Herb. Krug et Urban).

6999. **Operculina** Silva.

Operculina ventricosa Peter in Engl. u. Prantl. Natürl. Pflanzenfam. IV. 3a. (1891) 32. Hoofdpijnblad.

Cultivated. [Antill.] (Herb. Krug et Urban).

7003. **Ipomoea** Linn.

Ipomoea arenaria Steud. Nom. Ed. II. 1. (1841) 815.

St. Martin: near Mulletpond: n. 2777 B.; Lowlands: n. 3264 B.

St. Croix (Herb. Krug et Urban).

Cuba. Portorico, St. Thomas. (Herb. Krug et Urban).

Ipomoea batatas. Poir. Encycl. VI (1804) 14. Sweet potatoes.

Cultivated. [Amer. centr.] (E. a. P.).

Ipomoea dissecta Pers. in Linn. Syst. ed. XIV (1797) 207 in nota; Fl. 467; Duss 435.

Ipomoea sinuata Orteg. Hort. Matr. Dec. (1800) 84; Millspaugh 520.

St. Eustatius: Slope of the Quill on the side of White Wall 200 M.: n. 1183 B.; Fairplay: n. 1, 3; locality unknown: n. 164 G.

Saba: Bottom: n. 4.

St. Martin: Cul de Sac—Marigothill 0—200 M.: n. 2628 B.; French frontier: n. 2; locality unknown: n. 3598 B.

St. Croix ex Millspaugh.

All trop. countries. (Gris. Fl.).

Ipomoea fastigiata Sweet. Hort. Brit. ed. I. (1823) 288; Gris. Fl. 468; Duss 436; Millspaugh 520. Wild potato wine.

St. Eustatius: Slope of the Quill near Bengalen 200 M.: n. 689 B.; slope of the Quill: n. 177; locality unknown: n. 220 H., 226 G., 270 G.

Saba: lower part of the Bottom Mountain 300 M.: n. 1445 B.; between Bottom and Marypoint 100—200 M.: n. 1464 B.; Mountain 600 M.: n. 1745 B.; Ladder gut 0—200 M.: n. 2010 B.; Windward-side: n. 2281 B.; Crespeen—Rendez-vous: n. 179; Rendez-vous 400 M.: n. 178; locality unknown: n. 68 L., 218 L.

St. Martin: from Philipsburg to Guanabay: n. 2391 B.; along roads in Prince quarter: n. 2913 B.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

Ipomoea Nil Roth. Catalect. Fasc. I (1797) 36; Gris. Fl. 473; Duss 441.

Pharbitis Nil Choisy in Mem. Soc. Phys. Genève. VI (1833) 441; Millspaugh 521.

Ipomoea hederacea Jacq. Coll. I (1786) 124.

St. Eustatius: inner slope of the Quill 350 M.: n. 458 B.

Saba: Slope of the Mountain near St. John's 300 M.: n. 1732 B.

St. Martin: Mount Paradis 100—200 M.: n. 3294 B.

St. Croix ex Millspaugh.

In warmer countries of both hemisph. (Hemsley 1884).

Ipomoea pentaphylla Jacq. Coll. II (1788) 297; Gris. Fl. 467; Duss 436; Millspaugh 520. Noyeaux.

St. Eustatius: Cultuurvlakte: n. 196; locality unknown: n. 19 H., 281 H.

I also saw *Ipomoea pentaphylla*: on the slope of the Cliff.

Saba: Locality unknown: n. 30 L., 363 L.

St. Croix ex Millspaugh.

Antill., Venezuela, Equador, Pacif. Isl., Asia trop. and Africa. (Gris. Fl.).

Ipomoea pes caprae Roth. Nov. Pl. Sp. (1821) 109; Gris. Fl. 470; Duss 438; Millspaugh 520.

Ipomoea biloba Forsk. Fl. Aegypt. Arab. (1775) 44. Sea wine.

St. Eustatius: Seashore near Oranjestad: n. 10 B., 32 B., 152 H.; seashore near White Wall: n. 857 B.; seashore: n. 162, 163.

I also saw *Ipomoea pes caprae*: Seashore near Tumbledown Dickbay.

St. Martin: Guanabay: n. 2425 B.; near Fort Amsterdam: n. 164.

I also saw *Ipomoea pes caprae*: at Simonsbay.

St. Croix ex Millspaugh.

In all trop. and subtrop. countries. (Hemsley 1884).

Ipomoea sphenophylla Urb. in Symbolae Antillanae V (1908) 474.

St. Eustatius: Signalhill: n. 1038 B.

St. Eustatius. (Symb.).

Ipomoea violacea Linn. Spec. (1753) 161.

St. Martin: Tintamarre: n. 3374 B.

Venezuela, Antill. (Herb. Krug et Urban).

7004. **Calonyction** Choisy.

Calonyction speciosum Choisy Conv. Or. (1834) 59.

Ipomoea bona nox Linn. Spec. (1762) 288; Gris. Fl. 466; Duss 434; Millspaugh 520. Lilac bell.

St. Eustatius: Slope of the Quill: n. 165; locality unknown: n. 280 H.

St. Croix ex Millspaugh.

Amer. trop. (Ind. Kew.).

Calonyction tuba Colla. Nov. Spec. Calon. (1840) 15.

Ipomoea tuba G. Don. Gen. Syst. IV. (1838) 271; Gris. Fl. 467; Duss 434; Millspaugh 521.

Ipomoea grandiflora Lam. Tabl. Encyc. I (1791) 467.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 59 B.; seashore near Tumbledown Dickbay: n. 163 B.

St. Croix ex Millspaugh.

Antill., Guiana. (Gris. Fl.).

7005. **Quamoclit** Moench.

Quamoclit vulgaris Choisy Mem. Soc. Phys. Genève. VI (1833) 434.

Cultivated. [India orient.] (E. a. P.).

251. Hydrophyllaceae.

7037. **Hydrolea** Linn.

Hydrolea jamaicensis Raeusch. Nom. Ed. III (1797) 76.

Numa jamaicense Linn. Spec. ed. X (1759) 950; Gris. Fl. 477; Duss 444; Millspaugh 521.

St. Eustatius: near English Quarter: n. 1021 B.

St. Croix ex Millspaugh.

Bermud., Florida, Texas, Mexico, Antill. (Hemsley 1884).

252. Boraginaceae.

7038. **Cordia** Linn.

Cordia salvifolia Juss. in Poir. Encyc. VII (1806) 46; Gris. Fl. 480; Duss 447.

St. Eustatius: Little Mountain: n. 1141 B., 1154 B.; slope of the Quill above White Wall: n. 1173 B., 1176 B.

Saba: Hellsgate: n. 1686 B.; St. Giles quarter: n. 1861 B.; gut near Devilshand: n. 2088 B.

Guadeloupe, Martinique. (Herb. Krug et Urban).

Cordia Sebestena Linn. Spec. (1753) 190. Scarlet accordia. Cultivated. [Baham., Antill., Cuba to Guiana and Nova Grana-
nata]. (Gris. Fl.).

Cordia sulcata DC. Prod. IX. (1845) 488; Gris. Fl. 480; Millspaugh 522. Manjack.

St. Eustatius: Quill: n. 93.

Saba: between Bottom and Marypoint 300 M.: n. 1477 B., 1495 B.; Windwardside: n. 1938 B.; Gainbygut 500 M.: n. 90, 91, 92.

St. Croix ex Millspaugh.

Antill., Cuba to French Islands. (Gris. Fl.).

Cordia tremula Gris. Flor. W. I. (1861) 479.

Saba: from the Sulphurmines to Marypoint: n. 2030a B.

Antigua, Martinique, St. Vincent, Grenada, Barbados, Tobago. (Herb. Krug et Urban).

7042. **Beureria** Jacq.

Beureria succulenta Jacq. Enum. Pl. Carib. (1760) 14; Gris. Fl. 481; Duss 449.

Ehretia Bourreria Linn. Spec. (1762) 275; Millspaugh 522. White chank.

Saba: Bottom: n. 1983 B.; from Sulphurmines to Marypoint: n. 2031a B.; gut near Devilshand: n. 2093a B.; locality unknown: n. 307 L.

St. Martin: Marigotroad near Marigot: n. 2805 B., 2807 B.; lower part of the Milldrumhill: n. 3174 B.; Tintamarre: n. 3379 B., 3380 B.; gut from Bethlehem to Cul de Sac 0—100 M.: n. 3413 B.; Colebayhill: n. 63.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham. (Herb. Leiden).; Antill., Cuba to French Islands, Curaçao. (Gris. Fl.).

7045. **Rochefortia** Swartz.

Rochefortia acanthophora Gris. Flor. W. I. (1861) 482; Millspaugh 523.

St. Eustatius: Little Mountain: n. 1140a B.

St. Martin: Philipsburg near the saltponds: n. 2356*a* B.; Mount Paradis: n. 3244*a* B.

St. Domingo, Portorico, St. Jan. (Herb. Krug et Urban).

7051. **Tournefortia** Linn.

Tournefortia bicolor Sw. Prod. Veg. Ind. Occ. (1788) 40; Gris. Fl. 483; Duss 451.

Saba: Bottom: n. 2237 B.; Quarter: n. 147; Gainby gut: n. 146.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

Tournefortia caribaea Gris. Flor. W. I. (1861) 484.

Saba: Hellsgate: n. 148, 149.

Dominica, Martinique, St. Vincent, Barbados. (Herb. Krug et Urban).

Tournefortia foetidissima Linn. Spec. (1753) 140; Gris. Fl. 483; Duss 450; Millspaugh 523.

St. Eustatius: Bottom of the Quill: 300 M.: n. 443.

Saba: lower part of the Bottom Mountain 300 M.: n. 1431 B.; Mountain 800 M.: n. 1810 B.; gut between Hellsgate and Mastic-gut 300—400 M.: n. 2030 B.; Gainbygut: n. 150, 151; locality unknown: n. 186 L.

St. Martin: Marigothill: n. 2649*a* B.; Marigotroad: n. 2806 B.; hill on the northern side of Cul de Sac: n. 2996 B.; Mount Paradis: n. 3317*a* B.; gut from Bethlehem to Cul de Sac: n. 3411 B.

St. Croix ex Millspaugh.

Hispaniola, Portorico, Dominica, Guadeloupe, Martinique. (Herb. Krug et Urban).

Tournefortia gnaphalodes R. Br. Prod. (1810) 496; Gris. Fl. 483; Duss 450; Millspaugh 523. White lavender, Sea purslane.

St. Eustatius: White Wall: n. 809 B.; Concordiabay: n. 154; seashore: n. 267 H.

Saba: Bay: n. 66.

St. Martin: Guanabay: n. 2426 B.; near Mulletpond: n. 2770 B.; near Fort Amsterdam: n. 152.

I also saw *Tournefortia gnaphalodes*: Green Key.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida and throughout the West Indies. (Hemsley 1884).

Tournefortia hirsutissima Linn. Spec. (1753) 140; Gris. Fl. 483; Duss 450; Millspaugh 523.

St. Martin: Hill on the northern side of Cul de Sac 0—100 M.: n. 2976 B.; top of the hill on the northern side of Cul de Sac 200—300 M.: n. 2990 B.; Mount Paradis 300—400 M.: n. 3330 B.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

Tournefortia volubilis Linn. Spec. (1753) 140; Gris. Fl. 484; Duss 451; Millspaugh 523.

St. Eustatius: between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 762 B., 788 B.; near Maryglory 150 M.: n. 1144 B.

St. Martin: from Philipsburg to Belvedere: n. 2565 B., 2566a B.; near Fort Willem: n. 2791 B.; from Belvedere to Oysterpond: n. 3064 B.; Tintamarre: n. 3381 B.; locality unknown: n. 164, 165.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Venezuela to Brazil. (Gris. Fl.).

7052. **Heliotropium** Linn.

Heliotropium curassavicum Linn. Spec. (1753) 130; Gris. Fl. 486; Duss 453; Millspaugh 523. Sealavendel.

St. Eustatius: Seashore near Oranjestad: n. 9 B.; seashore near Tumbledown dickbay: n. 166 B.; Signalhill: n. 1055 B.; seashore near Fairplay: n. 97.

St. Martin: Guanabay: n. 2422 B.; near Fort Amsterdam: n. 59.

I also saw *Heliotropium curassavicum*: at Simonsbay village; Oysterpond.

St. Croix ex Millspaugh.

Amer. trop. and subtrop., Marokko, the Cape, Australia. (E. a. P.).

Heliotropium indicum Linn. Spec. (1753) 130; Gris. Fl. 485; Duss 452; Millspaugh 523. Eye bright.

St. Eustatius: Bengalen: n. 631 B.; along roads near Glass-bottle: n. 887 B.; slope of the Quill: n. 115; English Quarter: n. 119; locality unknown: n. 14 H., 15 G.

Saba: Crispeen 400 M.: n. 1726 B.

St. Martin: near Bethlehem: n. 2672 B.; between Mount Vernon and Orientbay: n. 2941 B.; Mary's Fancy: n. 74; locality unknown: n. 73.

I also saw *Heliotropium indicum*: Mount Paradis

St. Croix ex Millspaugh.

All trop. countries. (E. a. P.).

Heliotropium microphyllum Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1827) 58; Gris. Fl. 486.

St. Martin: in deserted plantations near Simonsbayshore: n. 3034 B.; Tintamarre: n. 3372 B.

Anguilla Boldingh.

Antigua, Guadeloupe. (Herb. Krug et Urban). Baham. (Hitchc.).

Heliotropium parviflorum Linn. Mant. II (1771) 201; Gris. Fl. 485; Duss 452; Millspaugh 523. Eyebright.

St. Eustatius: in a garden in Oranjestad: n. 80 B.; along roads in Oranjestad: n. 517 B.; Bengalen: n. 622 B.; slope of the lower part of the Quill near Bengalen: n. 732 B.; slope of the Quill: n. 72; in a garden: n. 129; locality unknown: n. 7 G., 7 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1361 B.; Crispeen 400 M.: n. 1721 B.; Laddergut 0--200 M.: n. 1956 B.; Bottom: n. 130.

St. Martin: from Filipsburg to Guanabay: n. 2344 B.; Prièregut: n. 128.

I also saw *Heliotropium parviflorum*: on Nakedboyhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Florida and Texas to Peru and Brazil. (E. a. P.).

Heliotropium peruvianum Linn. Spec. (1762) 187.

Cultivated. [Peru]. (Ind. Kew.).

253. Verbenaceae.

7138. **Verbena** Linn.

Verbena chamaedrifolia Juss. in Ann. Mus. Par. VII (1806) 73. Lilac verbenia.

Cultivated. [Argentin. and South Brazil.] (E. a. P.).

7144. **Lantana** Linn.

Lantana Camara Linn. Spec. (1753) 627; Gris. Fl. 495; Duss 463; Millspaugh 524. Sage.

St. Eustatius: Top of the Quill 400 M.: n. 209 B., 267 B.; near Maryglory 150 M.: n. 1159 B.; slope of the Quill on the side of

White Wall 200 M.: n. 1166 B.; Oranjestad: n. 165, 166; Cultuurvlakte: n. 167, 168, 169, 170; locality unknown: n. 171, 88 G., 110 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1293 B.; Greatbill 400 M.: n. 1380 B.; along the Forthgut 150—200 M.: n. 1624 B.; Laddergut 200 M.: n. 2002 B.; Bottom: n. 2255 B.; Ladder: n. 172; Bottom: n. 173; locality unknown: n. 253 L.

St. Martin: from Philipsburg to Guanabay: n. 2364 B.; Prince-quarter (coll. George): n. 2472 B.; Fort Willem 0—50 M.: n. 2535 B.; from Philipsburg to Belvedere: n. 2564 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2881 B.; from Belvedere to Oysterpond: n. 3046 B., 3083 B.; along roads from Grande Case to Marigot: n. 3105 B.; Vineyardhill: n. 175; Guanabay: n. 174.

I also saw *Lantana Camara* on: Nakedboyhill; lower part of Milldrum-hill; hills on the eastern side of Cul de Sac.

St. Croix ex Millspaugh.

In all warmer countries of America; also in the Old world. (Hemsley 1884).

Lantana involucrata Linn. Cent. Pl. II (1756) 22; Gris. Fl. 496; Duss 464; Millspaugh 524. Sage.

St. Eustatius: Seashore near Oranjestad: n. 14 B.; slope of the lower part of Signalhill: n. 101 B.; seashore near Tumbledown-dickbay: n. 160 B.; top of the Quill 400 M.: n. 218 B., 508 B.; along roads in Oranjestad: n. 580 B., 1100 B.; slope of the Quill near Bengalen 100 M.: n. 664 B., 670 B.; 200 M.: n. 739a B.; between Oranjestad and Fort de Wind on the top of the Cliff 40 M.: n. 773 B., 779 B., 786 B.; slope of the Quill: n. 194, 195, 196; Cultuurvlakte: n. 197, 198; English Quarter: n. 199, 200; locality unknown: n. 56 H., 89 G.

Saba: along the Forthgut 150—200 M.: n. 1637 B.; Hellsgate—Marypoint: n. 2036 B.; Marypoint: n. 201.

St. Martin: Prince quarter (coll. George.): n. 2556 B., 2485 B.; Fort Willem 0—50 M.: n. 2538 B.; from Philipsburg to Belvedere: n. 2560 B.; shore of the Simonsbaylagoon near Cul Piquard: n. 2742 B.; near Mulletpond: n. 2761 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2882 B.; between Mount Vernon and Orientbay: n. 2930 B.; Mount Paradis 100—200 M.: n. 3353 B.; Green Key: n. 3363 B.; Tintamarre: n. 3399 B.; hills on the eastern side of Cul de Sac: n. 3440 B.; near Fort Amsterdam: n. 202; Colebayhill: n. 203; French frontier: n. 204, 212.

I also saw *Lantana involucrata*: on one of the Islands in Simonsbaylagoon.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida and Antill., Mexico and northern part of Amer. austr. (Hemsley 1884).

7145. **Lippia** Linn.

Lippia reptans H. B. et K. Nov. Gen. et Sp. II (1817) 263; Gris. Fl. 495; Duss 462.

St. Martin: near Philipsburg: n. 2336 B.; from Philipsburg to Guanabay: n. 2419 B.; Guanabay: n. 186.

Anguilla Boldingh.

Antill., Mexico to Brazil. (Gris. Fl.).

7148. **Bouchea** Cham.

Bouchea Ehrenbergii Cham. in Linnaea VII (1832) 253; Gris. Fl. 493; Millspaugh 523. Vervine.

St. Eustatius: Locality unknown: n. 299 G., 228 H.

Saba: Windwardside: n. 1922 B.

St. Croix ex Millspaugh.

Amer. and Afric. (E. a. P.).

7151. **Stachytarpheta** Vahl.

Stachytarpheta jamaicensis Vahl. Enum. I (1804) 206; Gris. Fl. 494; Duss 461.

Stachytarpheta indica Vahl. Enum. I (1804) 206.

Valerianoides jamairensis Medic. Phil. Bot. I (1789) 177; Millspaugh 525. Purple wormbush.

St. Eustatius: in a garden in Oranjestad: n. 92 B.; Bengalen: n. 605 B.; slope of the Quill near Bengalen 150 M.; n. 685a B.; White Wall: n. 817 B.; Cultuurvlakte: n. 227, 229; slope of the Quill: n. 228; locality unknown: n. 293 H., 305 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1331 B.; top of the Cliff near the seaside of Boobyhill: n. 1606 B.; Windwardside: n. 2323 B.; Ladder: n. 230; locality unknown: n. 346 L.

I also saw *Stachytarpheta jamaicensis*: Crispeen.

St. Martin: from Philipsburg to Belvedere: n. 2581 B.

I also saw *Stachytarpheta jamaicensis*: Margothill—Marigot; in pasture-ground near Bellevue; hill between Mont des Accords and Centryhill; lower part of the Milldrumbill; Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida and Mexico to Brazil.; also in Afr. trop. and Asia. (Hemsley 1884).

7153. **Priva** Adans.

Priva echinata Juss. in Ann. Mus. Par. VII (1806) 69; Gris. Fl. 493; Duss 461.

Priva lappulacea Pers. Syn. II (1807) 139; Millspaugh 524.

St. Eustatius: along roads in Oranjestad: n. 557 B.; along roads near Glassbottle: n. 873 B.; in a garden in Oranjestad: n. 1275 B.; Quill: n. 219; locality unknown: n. 220, 221, 222, 38 G.

Saba: Boobyhill: n. 1594 B.; Laddergut 0—200 M.: n. 2019 B.; 223; Peperpot: n. 224; Bottom: n. 225, 226.

St. Martin: Marigotroad near Marigot: n. 2814 B.

St. Croix ex Millspaugh.

Bermud., Florida and Mexico to Brazil. (Hemsley 1884).

7161. **Citharexylum** Mill.

Citharexylum quadrangulare Jacq. Enum. Pl. Carib. (1760) 26; Gris. Fl. 497; Duss 465; Millspaugh 524. Susannaberry.

St. Eustatius: on the slope of the Cliff near Oranjestad: n. 53 B.; seashore: n. 152; Quill: n. 153; top of the Quill 400 M.: n. 271 B., 641a B.; slope of the Quill near Bengalen: n. 746 B.; hills near Concordia: n. 984a B.; locality unknown: n. 107 H., 121 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1314 B.; between Bottom and Marypoint 100—200 M.: n. 1527 B.; Paris: n. 154; locality unknown: n. 182 L., 190 L., 336 L.

St. Martin: Prince quarter (coll. George): n. 2473a B.; from Filipsburg to Guanabay: n. 2385 B., 2409 B.; Marigotroad near Marigot: n. 2818 B.; Centryhill: n. 155.

St. Croix ex Millspaugh.

Antill., Guiana. (Gris. Fl.).

7162. **Duranta** Linn.

Duranta repens Linn. Spec. (1753) 637; Millspaugh 524.

Duranta Plumieri Jacq. Select. Am. (1763) 186 t. 176 f. 76; Gris. Fl. 498; Duss 465. Adonis.

St. Eustatius: in a garden in Oranjestad: n. 1058 B.; near Maryglory 150 M.: n. 1137 B., 1157 B.; locality unknown: n. 139 H., 298 G., 1013 B.

Saba: Springbaygut: n. 2095 B.; Windwardside: n. 2308 B.; locality unknown: n. 364 L.

St. Martin: Point Blanche (coll. George): n. 2488 B.; along roads from Grande Case to Marigot: n. 3097 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

From Bolivia and Brazil. to Mexico and the Antill. (E. a. P.).
Also cultivated.

7178: *Aegiphila* Jacq.

Aegiphila martinicensis Jacq. Obs. Bot. II (1767) 3; Gris. Fl. 500; Duss 466; Millspaugh 523. Bastard white root.

St. Eustatius: Slope of the Quill: n. 246; top of the Quill: n. 386a B., Quill: n. 894 B.; locality unknown: n. 137 H., 212.

St. Croix ex Millspaugh.

Antill., Cuba and Panama to Guiana. (Gris. Fl.).

7191. *Clerodendron* Linn.

Clerodendron aculeatum Gris. Flor. W. I. (1861) 500; Duss 467; Millspaugh 524. Haguebush.

St. Eustatius: Seashore near Oranjestad: n. 24 B., 60 B.; Bengalen: n. 637 B.; Cultuurvlakte: n. 158, 159, 161; Cliff near Oranjestad: n. 162; locality unknown: n. 95 G., 147 H., 226 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1351 B.

I also saw *Clerodendron aculeatum*: along Forthbaygut

St. Martin: Fort Willem 0—500 M.: n. 2547 B., 2792 B.; Fort Amsterdam: n. 160.

I also saw *Clerodendron aculeatum* near Mont Chambord.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Amer. trop. (E. a. P.).

Clerodendron fallax Lindl. Bot. Reg. (1844) t. 19.

Cultivated. [Java]. (Ind. Kew.).

Clerodendron fragrans Willd. Enum. Hort. Berol. (1809) 659.

Cultivated. [China]. (E. a. P.).

Clerodendron Thomsonae Balf. Edinb. New. Phil. Journ. N. S. XV (1862) 233.

Cultivated. [Afr. trop.] (Ind. Kew.).

7205. **Avicennia** Linn.

Avicennia nitida Jacq. Enum. Pl. Carib. (1760) 25; Gris. Fl. 502; Duss 470; Millspaugh 523. Saltpond tree.

St. Martin: Colebay (coll. George): n. 2498 B.; along roads from Bethlehem to Mont Chambord: n. 2911 B.; along Simonsbaylagoon near the village: n. 3026 B.; Oysterpond: n. 135; Prince quarter: n. 136.

I also saw *Avicennia nitida*: near Fort Amsterdam; one of the Islands in Simonsbaylagoon.

St. Croix ex Millspaugh.

Antill., Cuba to Brazil., Afr. trop. (Gris. Fl.).

254. **Labiatae.**7264. **Leonotis** R.Br.

Leonotis nepetaefolia R. Br. in Ait. Hort. Kew, ed. II. III. (1811) 409; Gris. Fl. 492; Duss 460; Millspaugh 525. Adonis abbot.

St. Eustatius: in a garden in Oranjestad: n. 93 B., 1105 B.; Bengalen: n. 621 B.; Cultuurvlakte: n. 56; locality unknown: n. 138 G., 172 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1300 B.; Ladder: n. 216; Bottom: n. 57; locality unknown: n. 247 L.

St. Martin: along roads near Philipsburg: n. 2529 B.

St. Croix ex Millspaugh.

Antill. to Brazil. Afr. and Asia trop. (Gris. Fl.)

7268. **Leucas** R. Br.

Leucas martinicensis R. Br. Prodr. (1810) 504; Gris. Fl. 491; Duss 459; Millspaugh 525.

Saba: Bottom: n. 61.

St. Martin: from Philipsburg to Guanabay: n. 2342 B.

St. Croix ex Millspaugh.

Antill., Haiti to Brazil., Afr. trop. to the Cape and East Indies. (Gris. Fl.).

7373. **Leonurus** Linn.

Leonurus sibiricus Linn. Spec. (1753) 584; Gris. Fl. 525; Duss 459; Millspaugh 525.

St. Eustatius: in a garden in Oranjestad: n. 94 B.; along roads near Glassbottle: n. 865 B.; n. 865 B.; seashore: n. 58; locality unknown: n. 136 G., 194 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1312 B.; Ladder: n. 60.

St. Martin: from Philipsburg to Belvedere: n. 2587 B.; Mount Paradis 300—400 M.: n. 3238 B.; Prince quarter: n. 214.

St. Croix ex Millspaugh.

Europa and Siberia and naturalised in many other countries. (Hemsley 1884).

7290. *Salvia* Linn.

Salvia occidentalis Sw. Prod. Veg. Ind. Occ. (1788) 14; Gris. Fl. 490; Duss 458; Millspaugh 526. Rabbit meat.

St. Eustatius: along roads in Oranjestad: n. 536 B.; Cultuurvlakte: n. 74, 75, 78; Quill: n. 76, 77; Oranjestad: n. 72, 73; locality unknown: n. 292 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1313 B.; between Bottom and Marypoint 300 M.: n. 1480 B.; gut between Hellsgate and Mastiegut 300—400 M.: n. 2065 B.; Peperpot: n. 80; Crespeen—Rendez-vous 400 M.: n. 79; locality unknown: n. 35 L.

St. Martin: I saw *Salvia occidentalis*: hill between Mont des Accords and Centryhill; Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Bermud., Florida, Antill., and Mexico to Peru. (Hemsley 1884).

Salvia serotina Linn. Mant. 1 (1767) 25; Gris. Fl. 490; Millspaugh 526. Cat nip.

St. Eustatius: in a garden in Oranjestad: n. 95 B.; Cultuurvlakte n. 81; locality unknown: n. 33 H., 37 G.; along roads near Glassbottle: n. 868 B.

Saba: Seaside of Tentguthill 100 M.: n. 1652 B.; Springbaygut: n. 2103 B.; Bottom: n. 2257 B.; Greathill: n. 83; locality unknown: n. 35 L., 320 L.

St. Martin: along roads to Fort Willem: n. 2526 B.; Simonsbayvillage: n. 2838 B.; near Fort Amsterdam: n. 84; Prince quarter: n. 82;

I also saw *Salvia serotina*: on a hill between Mont des Accords and Centryhill.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida austr. and throughout the West Indies.
(Hemsley 1884).

7342. **Hyptis** Jacq.

Hyptis pectinata Poit. in Ann. Mus. Par. VII (1806) 474 t. 30;
Gris. Fl. 489; Duss 457.

Mesosphaerum pectinatum O. Ktze. Rév. Gen. (1891); Millspaugh 525. Holly stalk.

St. Eustatius: Bengalen: n. 628 B.; Cultuurvlakte: n. 42;
slope of the Quill: n. 43, 2010; locality unknown: n. 241 H.

Saba: between Bottom and Marypoint 300 M.: n. 1481 B.,
100—200 M.: n. 1536 B.; St. Giles: n. 1858 B.; gut between Hells-
gate and Masticgut 300—400 M.: n. 2052 B.; Ladder n. 44; Gainby-
gut: n. 45.

I also saw *Hyptis pectinata* at Flat Point.

St. Martin: from Philipsburg to Belvedere: n. 2603 B.; hills on
the northern side of Cul de Sac 0—100 M.: n. 2980 B.; between
Belvedere and Oysterpond: n. 3088 B.; locality unknown: n.
3599 B.

St. Croix ex Millspaugh.

From Mexico to Brazil, and in the Old world. (E. a. P.).

7355. **Coleus** Lour.

Coleus amboinicus Lour. Fl. Cochinch. II (1790) 372; Gris. Fl.
487; Duss 455; Millspaugh 525.

Coleus aromaticus Benth. in Wall. Pl. As. Rar. II (1832) 16.
Stinging thyme.

St. Eustatius: on the road along the lower part of Signalhill:
n. 119 B., 1133 B., 37; Cultuurvlakte: n. 38.

Saba: lower part of the Bottom Mountain 250 M.: n. 1392 B.;
Laddergut 200 M.: n. 1994 B.; Peperpot: n. 39; Bottom: n. 2009;
locality unknown: n. 286 L.

St. Martin: along roads near Colebayhill: n. 3042 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Indig. in the East Indies. (Gris. Fl.).

Coleus scutellarioides Benth. in Wall. Pl. As. Rar. II (1832) 16.
Cultivated. [Malaya, Australia]. (Ind. Kew.).

7366. **Ocimum** Linn.**Ocimum basilicum** Linn. Spec. (1753) 597. Basil.

Cultivated. [Asia occid. and trop.] (Ind. Kew.).

Ocimum gratissimum Linn. Spec. (1753) 597.

Cultivated [India orient.] (Ind. Kew.).

Ocimum micranthum Willd. Enum. Hort. Berol. (1813) 630; Gris. Fl. 487; Duss 454; Millspaugh 525. Mosquito basil.**St. Eustatius**: White Wall: n. 805 B.; slope of the Quill 200 M.: n. 877 B.; in a garden in Oranjestad: n. 1271 B.; slope of the Quill: n. 62; locality unknown: n. 63, 64, 131 G., 233 G., 233*a* G., 296 H.**Saba**: Laddergut 0—200 M.: n. 2016 B.; Bottom: n. 2254 B.; locality unknown: n. 327 L.**St. Martin**: Cul de Sac: 2613 B.; Nakedboyhill 50—200 M., n. 2701 B.; Marigotroad near Marigot: n. 2816 B.; near Colebayhill: n. 66; Marigotroad: n. 65.**St. Croix** ex Millspaugh.

Antill., Cuba and Mexico to Brazil. (Gris. Fl.).

256. **Solanaceae.**7379. **Lycium** Linn.**Lycium americanum** Jacq. Stirp. Am. (1763) 50.**St. Martin**: Guanabay: n. 2430 B., 2438 B.; near Fort Willem: n. 2790 B.; near Fort Amsterdam: n. 105, 106.**Anguilla** Boldingh.

Hispaniola, Cuba, St. Domingo. (Herb. Krug et Urban).

7381. **Acnistus** Schott.**Acnistus arborescens** Schlecht. in Linnaea VII (1832) 67; Gris. Fl. 435; Duss 410.**Saba**: along roads between Bottom and Windwardside 200 M.: n. 1297 B.; Rendez-vous 400 M.: n. 107, 108.

Antill., Mexico, Venezuela, Nova Granata, Brazil., Peru. (Gris. Fl.).

7401. **Physalis** Linn.**Physalis angulata** Linn. Spec. (1753) 183; Gris. Fl. 436; Duss 411; Millspaugh 527.**St. Eustatius**: Slope of the Quill: n. 206, 207.**St. Martin**: near Bethlehem: n. 2659 B.; between Belvedere and Oysterpond: n. 3065 B.

St. Croix ex Millspaugh.

Antill., from U. S. A. to Brazil., East Indies. (Gris. Fl.).

Physalis pubescens Linn. Spec. (1753) 183; Gris. Fl. 435; Millspaugh 527.

Saba: Bottom: n. 205; from Windwardside to Bottom: n. 2058a B.; gut near Bottom: n. 1430a B.

St. Martin: Marigotroad: n. 208.

St. Croix ex Millspaugh.

Amer. (natural. in other trop. countries). (Gris. Fl.).

7404. **Capsicum** Linn.

Capsicum ceratocarpum Fing. Monogr. (1832) 22.

Cultivated. (Patria ignot.).

Capsicum dulce Hort. ex Dun. in DC. Prod. XIII. 1 (1852) 428. Birdpepper.

Cultivated. [Trop countries]. (Ind. Kew.).

Capsicum frutescens Linn. Spec. (1753) 189; Gris. Fl. 436; Duss 411; Millspaugh 526.

St. Eustatius: Locality unknown: n. 51 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1336 B.; locality unknown: n. 7 L., 8 L.

St. Martin: lower part of the Milldrumhill: n. 3158 B.; Mount Paradis 300—400 M.: n. 3217 B.

St. Croix ex Millspaugh.

Trop. countries. (Gris. Fl.). Also cultivated.

7407. **Solanum** Linn.

Solanum aculeatissimum Jacq. Coll. I (1786) 100; Gris. Fl. 442; Millspaugh 527.

St. Eustatius: Locality unknown: n. 190 H., 263 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1309 B.; Bottom: n. 1855 B., 209; Windwardside: n. 1923 B.; locality unknown: n. 244 L.

St. Croix ex Millspaugh.

Widely spread in Amer. trop. and subtrop. (Hemsley 1884).

Solanum lanceaefolium Jacq. Coll. II (1788) 286; Gris. Fl. 440; Duss 414.

St. Eustatius: Top of the Quill 400 M.: n. 227 B., 268 B., 295 B.; slope of the Quill: n. 470a B.; top of the Quill 300 M.: n. 882 B.; Quill: n. 216.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1316 B., 1296 B.; Springbaygut: n. 2093 B.; Hellsgategut: n. 215; Crespeen—Rendez-vous: n. 129; Hellsgate: n. 218.

St. Martin: Mount Paradis 300—400: n. 3216 B.
Antill., Venezuela. (Gris. Fl.).

Solanum lycopersicum Linn. Spec. (1753) 185.

Lycopersicum esculentum Mill. Gard. Dict. ed. VIII (1768) n. 2; Gris. Fl. 436; Duss 412; Millspaugh 527. Tomato.

St. Eustatius: Seashore: n. 196; in a garden in Oranjestad: n. 96 H., 143 G., 227 H., 230 G.

Saba: Locality unknown: n. 145 L.

St. Croix ex Millspaugh.

Indig. in Amer. austral. (E. a. P.). Also cultivated.

Solanum Melongena Linn. Spec. (1753) 186. Boulanger.

Cultivated. (Patria ignota).

Solanum nigrum var. **nodiflorum**. Linn. Spec. (1753) 186.

Solanum nodiflorum Dunal Hist. Sol. (1813) 151; non Jacq.; Gris. Fl. 437; Duss 412.

Solanum nigrum nodiflorum A. Gray; Millspaugh 527. Wild pepper.

St. Eustatius: in a garden in Oranjestad: n. 81 B.; along roads in Oranjestad: n. 1078 B.; slope of the Quill near Glassbottle: n. 1202 B.; Cultuurvlakte: n. 75; slope of the Quill: n. 219; near Roundhill: n. 220; locality unknown: n. 282 G.

Saba: lower part of the Bottom Mountain 250 M.: n. 1400a B.; Mountain 600 M.: n. 1779 B.; Windwardside: n. 1904 B.; Ladder-gut 0—200 M.: n. 1957 B.; Bottom: n. 2247 B.; Crespeen—Rendez-vous 400 M.: n. 222; locality unknown: n. 251 L.

St. Martin: Marigotroad near Marigot: n. 2804 B.; hill on the northern side of Cul de Sac 0—100 M.: n. 2979 B., 2983 B., 2983 B.; Marigotroad: n. 76.

St. Croix ex Millspaugh.

All trop. countries. (Gris. Fl.).

Solanum racemosum Jacq. Enum. Pl. Carib. (1760) 15; Gris. Fl. 439; Duss 414; Millspaugh 527. Cankerberry.

St. Eustatius: Seashore near Oranjestad: n. 11 B.; slope of the lower part of Signalhill: n. 98 B., 113 B.; along roads in Oranjestad: n. 556 B.; in deserted plantations at the seashore near Bengalen: n. 586 B., 593 B.; Bengalen: n. 608 B.; Fort de Wind: n. 797 B.; hill near Concordia 100 M.: n. 955a B.; from Little Moun-

tain to Venusbay: n. 987 B.; Signalhill 150 M.: n. 1043 B.; slope of the Quill on the side of White Wall 200 M.: n. 1180 B.; Venusbay: n. 223.

Saba: Seashore near Forthbaylanding: n. 1650 B.; Springbay-gut: n. 2084 B.

St. Martin: Guanabay: n. 2424 B.; along the saltpond near Philipsburg: n. 2452 B.; lowerpart of Milldrumhill: n. 3160 B.; Mount Paradis 300—400 M.: n. 3210 B.

St. Croix ex Millspaugh.

Antill. (Gris. Fl.).

Solanum racemosum Jacq. var **igneum** (Linn.) O. E. Schulz.¹⁾

Solanum igneum Linn. Spec. (1762) 270; Gris. Fl. 440; Millspaugh 527.

St. Eustatius: Seashore near Oranjestad: n. 35 B., 46 B.; along roads in Oranjestad: n. 566 B., 573 B., 926 B.; in deserted plantations at the seashore near Bengalen: n. 588 B.; top of the Cliff between Fort de Wind and Oranjestad: n. 774 B.; gut near Fort de Wind: n. 833 B.; Cultuurvlakte: n. 213; slope of Signalhill: n. 114; seashore: n. 211, 212; locality unknown: n. 46 G., 47 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1468 B.; Bottom: n. 1838 B.; Ladder: n. 214; locality unknown: n. 141 L., 362 L.

St. Martin: near Philipsburg: n. 2337 B.; along the saltpond near Philipsburg: n. 2442 B.; near Bethlehem: n. 2667 B.; near Mulletpond: n. 2787 B.; in pastureground near Bellevue: n. 2821 B.; lower part of the hills between Mont Chambord and Mount Vernon: n. 2958 B.; along roads from Colebay to Simonsbay: n. 3251 B.; gut from Bethlehem to Cul de Sac 200 M.: n. 3428 B., 3430 B.; Cul de Sac: n. 210.

St. Croix ex Millspaugh.

Antill., Cuba to Guiana. (Gris. Fl.).

Solanum torvum Sw. Prod. Veg. Ind. Occ. (1788) 47; Gris. Fl. 441; Duss 415; Millspaugh 527.

St. Eustatius: Slope of the Quill: n. 226.

Saba: lower part of the Bottom Mountain: n. 1402 B.; between Bottom and Marypoint 100—200 M.: n. 1533 B.; Windwardside—Hellsgate: n. 1678 B.; Laddergut 0—200 M.: n. 1959 B., 224.

St. Martin: Cul de Sac: n. 2618 B.; along roads from Bethle-

¹⁾ Det. O. E. Schulz.

hem to Mont Chambord: n. 2909 B.; hills on the eastern side of Cul de Sac: n. 3448 B.; Prince quarter: n. 225.

St. Croix ex Millspaugh.

Bermud., Antill., Cuba and Mexico to Brazil and Equador. East Indies. (Gris. Fl.).

Solanum tuberosum Linn. Spec. (1753) 185. Potatoes.

Cultivated. [Amer. austr.] (E. a. P.).

7415. **Datura** Linn.

Datura fastuosa Linn. Syst. ed. X (1799) 932; Gris. Fl. 434; Duss 409; Millspaugh 526. Fire weed.

Saba: Crispeen 400 M.: n. 1766 B.; locality unknown: n. 260 L.

St. Martin: near Bethlehem: n. 2660 B.

St. Croix ex Millspaugh.

Antill., Afr. trop., East Indies. (Gris. Fl.).

Datura Metel Linn. Spec. (1753) 179; Gris. Fl. 434; Duss 408; Millspaugh 526. Fire weed.

St. Eustatius: along roads near Oranjestad: n. 563 B.; in a garden in Oranjestad: n. 1015 B., 1081 B., 190, 12 H., 13 G.

St. Martin: along roads near Philipsburg: n. 2527 B.; Guana-bay: n. 189.

St. Croix ex Millspaugh.

Antill., Haiti and Mexico to Brazil and in the Old world. (Gris. Fl.).

Datura Stramonium Linn. Spec. (1753) 179; Gris. Fl. 434; Millspaugh 526. Fire weed.

St. Eustatius: in a garden in Oranjestad: n. 542 B.; Cultuur-vlakte: n. 194; locality unknown: n. 12 G., 14 G., 13 H., 3592 B.

Saba: Bottom: n. 1991 B., 195; locality unknown: n. 83 L., 260 L.

St. Martin: Mary's Fancy: n. 193.

St. Croix ex Millspaugh.

Europa. Asia, Africa and Amer. sept. (E. a. P.).

7420. **Cestrum** Linn.

Cestrum laurifolium l'Hér. Stirp. Nov. (1784) 69. t. 34; Gris. Fl. 444; Duss 416; Millspaugh 526.

St. Eustatius: Top of the Quill 400 M.: n. 211 B., 239 B., 240 B., 374 B.; Quill: n. 102, 103, 113; Boven: n. 104; locality unknown: n. 3638 B.

Saba: lower part of the Bottom Mountain 300 M.: n. 1442 B.; along roads between Bottom and Windwardside 200 M.: n. 1288 B.; Crispeen 400 M.: n. 1723a B.; Springbaygut: n. 2124 B.; Bottom: n. 2252 B.; Paris 300 M.: n. 109; Greathill 2—400 M.: n. 112; Rendez-vous: n. 110; Crispeen—Rendez-vous: n. 111; Rendez-vous Mountain: n. 67; locality unknown: n. 68.

St. Croix ex Millspaugh.

Cuba, Hispaniola, Portorico, St. Thomas, St. Jan, Antigua, Montserrat, Guadeloupe, Dominica, Martinique, Barbados, Trinidad. (Herb. Krug et Urban).

7434. **Nicotiana** Linn.

Nicotiana Tabacum Linn. Spec. (1753) 180.

Cultivated. [Amer. austr.] (E. a. P.).

7436. **Petunia** Juss.

Petunia violacea Lindl. Bot. Reg. t. 1626.

Cultivated. [Argentina]. (Ind. Kew.).

7450. **Brunfelsia** Linn.

Brunfelsia americana Linn. Spec. (1753) 191. Lady of the night.

Cultivated. [Antill.] (Gris. Fl.).

257. Scrophulariaceae.

7470. **Angelonia** Humb. et Bonpl.

Angelonia angustifolia Benth. in DC. Prodr. X (1846) 254. Snapdragon.

Cultivated. [Mexico]. (Gris. Fl.).

7487. **Maurandia** Ortega.

Maurandia semperflorens Jacq. Hort. Schoenb. III (1800) 20.

Saba: Gut near Bottom: n. 1388 B.

Antill. [Mexico]. (Ind. Kew.).

7496. **Russelia** Jacq.

Russelia juncea Zuccar in Flora XV (1832) II Beibl. 99. Bugal plant.

Cultivated. [Mexico]. (Ind. Kew.).

7574. **Capraria** Linn.

Capraria biflora Linn. Spec. (1753) 628; Gris. Fl. 427; Duss 403; Millspaugh 527. Wild tea.

St. Eustatius: Seashore near Oranjestad: n. 34 B.; in a garden in Oranjestad: n. 70 B.; seashore near Tumbledown Dickbay: n. 162 B.; Bengalen: n. 624 B.; along roads in Oranjestad: n. 929 B.; Cultuurvlakte: n. 39; locality unknown: n. 11 G., 11 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1319 B.; between Bottom and Marypoint 100—200 M.: n. 1510 B.; Laddergut 0—200 M.: n. 1953 B., 54; Bottom: n. 55.

St. Martin: from Philipsburg to Guanabay: n. 2348 B.; Philipsburg—Cul de Sac: n. 2611 B.; Mount Paradis 200—400 M.: n. 3202 B.

St. Croix ex Millspaugh.

Bermud., Florida and Mexico to Brazil and Peru. (Hemsley 1884).

7576. *Scoparia* Linn.

Scoparia dulcis Linn. Spec. (1753) 116; Gris. Fl. 427; Duss 403; Millspaugh 528. Sweet broom.

St. Eustatius: along roads near Glassbottle: n. 883 B., 1205 B.; Cultuurvlakte: n. 85; slope of the Quill: n. 86, 87; locality unknown: n. 10 G., 10 H.

Saba: between Bottom and Marypoint 100—200 M.: n. 1523 B.; Ladder: n. 78; Peperpot: n. 79; Mountain: n. 80; Gainbygut: n. 81.

St. Croix ex Millspaugh.

All tropical countries. (Gris. Fl.).

258. Bignoniaceae.

7705. *Bignonia* Linn.

Bignonia Unguis-cati Linn. Spec. (1753) 623; Gris. Fl. 448; Duss 421; Millspaugh 528. Cat claw.

St. Eustatius: Top of the Quill 400 M.: n. 194 B., 281 B., 283 B., 350 B.; Bottom of the Quill 350 M.: n. 454 B.; Red Cliff—Tumbledown Dickbay: n. 202; Quill: n. 41, 42, 102; slope of the Quill: n. 196, 197; locality unknown: n. 178 G., 259 H.

I also saw *Bignonia Unguis-cati* in deserted plantations near Bengalen.

Saba: Boobyhill 300 M.: 1599 B.; Laddergut 0—200 M.: n. 1966 B.; between Hellsgate and Marypoint: n. 2071 B.; pasture-ground: n. 40; Middle Island: n. 149.

St. Martin: Hill on the northern side of Cul de Sac 0—150 M.: n. 2970 B.; gut from Betlehem to Cul de Sac 200 M.: n. 3435 B.; John Ednyhill: n. 198, 199, 200.

I also saw *Bignonia Unguis-cati* on Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Antill. (Gris. Fl.); Amer. cont. (Herb. Leiden.).

Bignonia spec. without flowers and fruits.

St. Eustatius: Quill: n. 438a B.

7733. **Tecoma** Juss.

Tecoma capensis Lindl. Bot. Reg. t. 1117.

Cultivated. [Afr. austr.] (Ind. Kew.).

Tecoma leucoxydon Mart. ex DC. Prod. IX (1845) 219; Gris. Fl. 447; Duss 420; Millspaugh 528. White cedar.

St. Eustatius: in a garden in Oranjestad: n. 72 B.; slope of the lower part of Signalhill: n. 104 B.; top of the Quill 400 M.: n. 214 B., 331 B., 355 B.; between Oranjestad and Fort de Wind: n. 782 B.; along roads in Oranjestad: n. 919 B.; along roads near English Quarter: n. 1003 B.; slope of the Quill near Glassbottle: n. 1116 B.; slope of the hills near Concordiabay: n. 206; Tumble-down-dickbay: n. 216; near Sugarloaf: n. 217; locality unknown: n. 52 H., 109 G.

I also saw *Tecoma leucoxydon*: on Gilboohill; near Jenkinsbay.

Saba: between Bottom and Marypoint 100—200 M.: n. 1500 B.; Boobyhill: n. 1570 B.; Green Island: n. 2145 B.; Windwardside: n. 2333 B.; locality unknown: n. 284 L.

I also saw *Tecoma leucoxydon*: Sulphurmines; Bottom—Windwardside; Hellsgate.

St. Martin: Colebay (coll. George): n. 2513 B.; from Philipsburg to Belvedere: n. 2555 B.; Nakedboyhill 50—200 M.: n. 2675 B.

I also saw *Tecoma leucoxydon*: Marigothill to Marigot; hill between Mont des Accords and Centryhill: hill on the northern side of Cul de Sac; Mount Paradis 300—400 M.; Lowlands.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Cuba to Guiana. (Gris. Fl.).

7740. **Stenolobium** D. Don.

Stenolobium stans Seem. Journ. Bot. I (1863) 88.

Tecoma stans Juss. Gen. (1789) 139; Gris. Fl. 477; Duss 420; Millspaugh 529. Yellow blossom.

St. Eustatius: Gut near Fort de Wind 50 M.: n. 834 B.; hill near Concordia 100 M.: n. 956 B.; Cultuurvlakte: n. 209; locality unknown: n. 100 G., 178 H.

I also saw *Stenolobium stans*: slope of the Cliff; slope of the Quill on the side of White Wall.

Saba: lower part of the Bottom Mountain 800 M.: n. 1440 B.; Laddergut 200 M.: n. 1998 B., 211; locality unknown: n. 134 L.

I also saw *Stenolobium stans*; between Bottom and Marypoint; Forthbaygut; Bottom—Windwardside.

St. Martin: Mount Paradis 300—400 M.: n. 3323 B.; Cul de Sac: n. 210.

I also saw *Stenolobium stans*: on Marigothill—Marigot; Nakedboyhill; hill between Mont des Accords and Centryhill; hill on the northern side of Cul de Sac; along roads near Grande Case; lower part of Milkdrumhill.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Tucuman. (Gris. Fl.).

7759. *Crescentia* Linn.

Crescentia Cujete Linn. Spec. (1753) 626; Gris. Fl. 445; Duss 418; Millspaugh 528. Calabash.

St. Eustatius: in a garden in Oranjestad: n. 149 G., 255 H.

I saw *Crescentia Cujete* in deserted plantations near English Quarter.

Saba: Windwardside: n. 1920 B.; locality unknown: n. 65 L.

St. Martin: from Philipsburg to Belvedere: n. 2554 B.; Mount Paradis 300—400 M.: n. 3338 B.

I also saw *Crescentia Cujete*: near Saline.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill. and Amer. austr. (E. a. P.).

262. Gesneriaceae.

7839. *Besleria* Linn.

Besleria lutea Linn. Spec. (1753) 619; Gris. Fl. 463; Duss 431.

St. Eustatius: Top of the Quill 500 M.: n. 471a B., 479a B.

Saba: Gut near Bottom: n. 1424a B.; Mountain 600 M.: n. 1782 B., 41, 42, 43, 208.

Jamaica, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Grenada. (Symb.).

7880. *Kohleria* Regel.

Kohleria hirsuta Reg. in Regensb. Flor. XXXI (1848) 250.

Isoloma hirsuta Reg. in Bot. Zeit. IX (1851) 583; Gris. Fl. 459.

Saba: Locality unknown: n. 60 L., 331 L.
Cultivated?
Trinidad, Venezuela. (Symb.).

7891. **Gesneria** Linn.

Gesneria ventricosa Sw. Prod. Veg. Ind. Occ. (1788) 89.

Pentarhaphia longiflora Lindl. Bot. Reg. sub. t. 1110; Gris. Fl. 460; Duss 430.

St. Eustatius: Top of the Quill 400 M.: n. 471 B.; top of the Quill 500 M.: n. 479 B.; top of the Quill: n. 325, 326, 327, 328.

Saba: Mountain 600 M.: n. 1758 B., 1783 B., 332, 333.

Jamaica, St. Kitts, Montserrat, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent. (Symb.).

264. **Lentibulariaceae.**

7901. **Utricularia** Linn.

Utricularia alpina Jacq. Enum. Pl. Carib. (1760) 11.

Utricularia montana Jacq. Select. Am. (1763) 7 t. 6; Gris. Fl. 390.

Saba: Mountain 800 M.: n. 2229 B.

Antill. and Amer. austral. (E. a. P.).

266. **Acanthaceae.**

7914. **Thunbergia** Retz.

Thunbergia affinis Moore var. **pulvinatus** Moore in Journ. Bot. XVIII (1880) 431.

Cultivated. [Africa]. (Herb. Berlin).

Thunbergia alata Boj. in Hook. Exot. Fl. 1823 t. 17; Duss 428.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1311 B.; between Bottom and Marypoint 100--200 M.: n. 1548 B.; locality unknown: n. 1593 B., 36 L., 10; Village: n. 283.

St. Croix. (Symb.).

Indig. in Africa orient. (Symb.).

Thunbergia fragans Roxb. Pl. Corom. I (1795) 47 t. 67; Gris. Fl. 458; Duss 428; Millspaugh 530.

St. Eustatius: along roads in Oranjestad: n. 1092 B., 183 H.

Saba: Bottom: n. 284; locality unknown: n. 17 L.

St. Martin: along roads from Grande Case to Marigot: n. 3102 B.

St. Croix ex Millspaugh.

Indig. in Asia trop. (Symb.).

7919. **Sanchezia** R. et Pav.

Sanchezia nobilis Hook. Bot. Mag. t. 5594.

Cultivated. [Equador]. (Ind. Kew.).

7930. **Blechnum** Juss.

Blechnum Brownei Juss. in Ann. Mus. Par. IX (1807) 270; Gris. Fl. 453; Duss 424.

Blechnum Blechnum Millspaugh 529.

St. Eustatius: Locality unknown: n. 304 H.; slope of the Quill: n. 258.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1295 B.; between Bottom and Marypoint 100—200 M.: n. 1507 B.; Mountain 600 M.: n. 1744 B.; Bottom: n. 1847 B., 1984 B.; Ladder: n. 260; Laddergut: n. 260 bis; locality unknown: n. 316 L.

St. Martin: Cul de Sac—Marigothill 0—200 M.: n. 2626 B.; in an orchard near Bellevue: n. 2846 B.

St. Croix ex Millspaugh.

Amer. trop. (Symb.).

7965. **Ruellia** Linn.

Ruellia tuberosa Linn. Spec. (1753) 635; Gris. Fl. 452; Duss 423; Millspaugh 529. Devil's bit, Fever root.

St. Eustatius: along roads near Glassbottle: n. 862 B.; Cultuurvlakte: n. 272; Concordia: n. 274; locality unknown: n. 83 H., 129 G.

I also saw *Ruellia tuberosa* on Signalhill.

Saba: between Bottom and Marypoint 100—200 M.: n. 1512 B.; Hellsgate: n. 1697 B.; locality unknown: n. 297 L.; Bottom: n. 273.

I also saw *Ruellia tuberosa* in the Laddergut.

St. Martin: from Philipsburg to Guanabay: n. 2375 B.; from Philipsburg to Belvedere: n. 2553 B.; along roads near Bethlehem: n. 2934 B.; Centryhill: n. 12.

I also saw *Ruellia tuberosa*: on Nakedboyhill: hill between Mont des Accords and Centryhill; in pastureground near Belvedere; in pastureground near Bellevue; Lowlands.

St. Croix ex Millspaugh.

Antill., from Guiana, Venezuela, Peru and Columbia to Mexico. (Symb.).

7990. **Stenandrium** Nees.

Stenandrium rupestre Nees. in DC. Prodr. XI. (1847) 283.

St. Martin: in an orchard near Bellevue: n. 2835 B.; lower part of the hills on the eastern side of Cul de Sac: n. 3442 B.

I also saw *Stenandrium rupestre* on the hill between Mont des Accords and Centryhill.

Baham., Antill. (Symb.).

8012. **Graptophyllum** Nees.

Graptophyllum pictum Griff. Notul. IV. (1851) 139.

Cultivated. [Nova Guinea]. (Symb.).

8094. **Justicia** Linn.

Justicia sessilis Jacq. Enum. (1760) 11.

Dianthera sessilis Gris. Flor. W. I. (1861) 455; Duss 426. Prickly balsam.

St. Eustatius: Slope of the lower part of Signalhill: n. 99 B.; lower part of the hills near Concordia: n. 977 B.; near Maryglory 150 M.: n. 1150 B.; near Boven 200 M.; n. 1246 B.; Cultuurvlakte: n. 13, 262, 263, 264; slope of the Quill: n. 265, 266; locality unknown: n. 74 H., 111 G., 141 G.

Saba: Windwardside—Hellsgate: n. 1667 B.; Laddergut 200 M.: n. 2004 B.; Paris: n. 267.

St. Martin: along roads near Bellevue: n. 2842 B.; Mountain between Mont des Accords and Centryhill 0—200 M.: n. 2866 B.; from Belvedere to Oysterpond: n. 3057 B.

I also saw *Justicia sessilis*: on the lower part of the Milldrums Hill; Mount Paradis.

St. Croix. (Symb.).

Hispaniola, Portorico, St. Thomas, Antigua, Guadeloupe. (Symb.).

8098. **Beloperone** Nees.

Beloperone eustachiana Benth. in Benth. et Hook. Gen. Pl. II (1876) 1111.

Justicia eustachiana Jacq. Enum (1760) 11; Gris. Fl. 456; Duss 426.

St. Eustatius: Slope of the Quill near Bengalen 200 M.: n. 752 B.; top of Signalhill 225 M.: n. 1033 B.; near Maryglory 150 M.: n. 1138 B., 1149 B.; locality unknown: n. 166 G., 248 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1286 B.; Windwardside—Hellsgate: n. 1684 B.; Paris 2—400 M.: n. 256; Greathill: n. 255; Rendez-vous 400 M.: n. 257; Bottom: n. 1; locality unknown: n. 24 L.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2860 B.; along roads from Colebay to Simonsbay: n. 3248 B.; Green Key: n. 3366 B.

St. Barthélemy, Antigua, Montserrat, Guadeloupe. (Symb.).

267. Myoporaceae.

8111. **Bontia** Linn.

Bontia daphnoides Linn. Spec. (1753) 638.

Cultivated. [Antill.] (E. a. P.).

269. Plantaginaceae.

8116. **Plantago** Linn.

Plantago major Linn. Spec. (1753) 112; Gris. Fl. 389; Duss 379; Millspaugh 530.

Saba: Boobyhill: n. 1566 B.; Windwardside: n. 2327 B.; locality unknown: n. 178 L.

St. Croix ex Millspaugh.

In almost all parts of both hemisph. (E. a. P.).

270. Rubiaceae.

8136. **Oldenlandia** Linn.

Oldenlandia herbacea DC. Prod. IV. (1830) 425; Gris. Fl. 330; Duss 335; Millspaugh 531.

St. Martin: from Philipsburg to Belvedere: n. 2580 B.; along roads from Grande Case to Marigot: n. 3104 B.

St. Croix ex Millspaugh.

Amer. trop., Afric., Asia, Malay, Kaiser Wilhelmsland. (E. a. P.).

8208. **Hillia** Jacq.

Hillia longiflora Sw. Prod. Veg. Ind. Occ. (1788) 58; Gris. Fl. 325; Duss 334.

Hillia parasitica Jacq. Enum. Pl. Carib. (1760) 18.

St. Eustatius: Top of the Quill: n. 247 B., 68, 69, 70.

Saba: Mountain 800 M.: n. 2192 B., 2198 B.; locality unknown: n. 127 L.

Antill., Amer. austral. (E. a. P.).

8219. **Exostemma** L. C. Rich.

Exostemma caribaeum Roem. et Schult. Syst. V (1819) 18; Gris. Fl. 324; Duss 333; Millspaugh 531.

St. Martin: Colebay (coll. George): n. 2500 B.; Nakedboyhill 200—260 M.: n. 2726 B.; near Mulletpond: n. 2783 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2891 B.; from Belvedere to Oysterpond: n. 3045 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill. (Herb. Krug er Urban); Key West (Melvill.); Baham. (Hitche.).

Exostemma spec.

Saba: locality unknown: n. 107 L.

8238. **Mussaenda** Linn.

Mussaenda frondosa Linn. Spec. (1753) 177.

Cultivated. [India orient.] (Ind. Kew.).

8239. **Gonzalagunia** Ruiz. et Pav.

Gonzalagunia hirsuta Schumann. in Mart. Fl. Bras. VI 6 (1889) 291.

Gonzalea spicata DC. Prodr. IV (1830) 437; Gris. Fl. 321; Duss 332.

Saba: between Bottom and Marypoint 350 M.: n. 1493 B., 1552 B.; Mountain near Hellsgate: n. 2284 B.; Mountain: n. 171, 172; Devilshand: n. 173.

Antill., Guiana, Bahia. (Gris. Fl.).

8283. **Randia** Linn.

Randia aculeata Linn. Spec. (1753) 1192; Gris. Fl. 318; Duss 330; Millspaugh 531. Fishing rod, Black cherry.

St. Eustatius: Slope of the lower part of Signalhill: n. 100 B.; top of the Quill 400 M.: n. 280 B.; slope of the lower part of the Quill near Bengalen: n. 712 B.; between Oranjestad and Fort de Wind on the top of the Cliff: n. 790 B.; slope of Signalhill: n. 125, 182; Quill: n. 181; White Wall: n. 127; Little Mountain: n. 126; Gilboohill: n. 124; locality unknown: n. 176 G., 195 H.

I also saw *Randia aculeata*: on the slope of the Quill on the side of White Wall; Tumbledowndiekbay—Venusbay.

Saba: between Bottom and Marypoint 100—200 M.: n. 1519 B.; Saddle: n. 1658 B.; Hellsgate: n. 120; locality unknown: n. 123.

St. Martin: from Philipsburg to Guanabay: n. 2416 B.; Prince quarter (coll. George): n. 2474 B.; near Fort Willem: n. 2798 B.; top of the hill on the northern side of Cul de Sac 200—300 M.: n. 2991 B.; Milldrumhill 300—400 M.: n. 3140 B.; Vineyardhill: n. 121, 122.

I also saw *Randia aculeata*: Marigotthill—Marigot; top of Nakedboyhill; near Bellevue; hill between Mont des Accords and Centryhill; along roads near Grande Case; Mount Paradis 300—400 M.; Lowlands.

St. Croix ex Millspaugh.

Bermud., Florida, Antill. (Hemsley 1884).

8361. *Guettarda* Linn.

Guettarda parvifolia Sw. Fl. Ind. Occ. III (1806) 1958; Gris. Fl. 333; Duss 337; Millspaugh 531. Wild cherry.

St. Eustatius: from Little Mountain to Venusbay: n. 961 B.; slope of the Quill on the side of White Wall 200 M.: n. 1181 B.

Saba: between Bottom and Marypoint 100—200 M.: n. 1464 B.; Springbaygut: n. 2079 B.

St. Martin: from Philipsburg to Belvedere: n. 2571 B.; Naked-boyhill 280 M.: n. 2737 B.; mountain between Mont des Accords and Centryhill 200—300 M.: n. 2887 B.; lower part of the mountains between Mount Vernon and Mont Chambord: n. 2950 B.; Milldrumhill 200—300 M.: n. 3122 B.; hills on the eastern side of Cul de Sac: n. 3438 B.

St. Croix ex Millspaugh.

Antill., Venezuela. (Gris. Fl.).

Guettarda scabra Lam. Tabl. Encycl. II (1793) 218. t. 154 f. 3; Gris. Fl. 332; Duss 337; Millspaugh 531. Candle wood, Wild guave.

St. Eustatius: Top of the Quill 380 M.: n. 255 B., 319 B., 502 B.; from Little Mountain to Venusbay: n. 993 B.; Signalhill: n. 76; Quill: n. 78.

I also saw *Guettarda scabra* on the slope of the Quill on the side of White Wall.

Saba: between Bottom and Marypoint 100—200 M.: n. 1463 B.; Greathill: n. 75 A.; Hellsgate: n. 75; locality unknown: n. 77.

I also saw *Guettarda scabra*: in the gut between Hellsgate and Mastic-gut: Crespeenhill.

St. Martin: Mount Paradis 300—400 M.: n. 3224a B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill., Yucatan, Brazil. (Gris. Fl.).

8371. *Erithalis* Linn.

Erithalis fruticosa Linn. Syst. ed. X (1759) 930; Gris. Fl. 336; Duss 338; Millspaugh 530. Flambeau.

St. Eustatius: from Little Mountain to Venusbay: n. 973 B.; top of Signalhill 225 M.: n. 1032 B.; White Rock: n. 87; slope of the Quill near White Wall: n. 86; White Wall: n. 83, 151; Sugarloaf: n. 152; Boven: n. 84; Oranjestad n. 85.

St. Martin: Colebay (coll. George): n. 2499 B.; near Mullet-pond: n. 2752 B.; from Belvedere to Oysterpond: n. 3047 B.; Tintamarre: n. 3390 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill. (Herb. Krug et Urban); Key West (Melvill.); Baham. (Hitche.)

8376. *Chiococca* Linn.

Chiococca racemosa Jacq. Select. Am. (1763) 68; Gris. Fl. 336; Duss 339; Millspaugh 530. Buckroot.

St. Eustatius: Top of the Quill 400 M.: n. 199 B., 336 B., 400 B.; slope of the Quill near Bengalen 200 M.: n. 688 B.; from Little Mountain to Venusbay: n. 972 B.; slope of the Quill on the side of White Wall 200 M.: n. 1170 B.; top of the Quill: n. 141, 142, 143, 152; Quill: n. 156; locality unknown: n. 157, 187 G., 271 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1355 B.; Greathill 400 M.: n. 1385 B.; between Bottom and Marypoint 350 M.: n. 1497 B.; Boobyhill 300 M.: n. 1574 B.; Hellsgate—Marypoint: n. 2038 B.; Springbaygut: n. 2081 B., 2082 B.; Paris: n. 154, 155, 159; Rendez-vous: n. 158; Middle Island: n. 153; Wellgut: n. 151; locality unknown: n. 137 L., 309 L.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2637 B.; mountain between Mont des Accords and Centryhill 200—300 M.: n. 2879 B.; top of the hills on the northern side of Cul de Sac 200—

300 M.: n. 3009 B.; from Belvedere to Oysterpond: n. 3060 B.; Milldrumhill 200—300 M.: n. 3123 B.; 300—400 M.: n. 3153 B.; Mount Paradis 300—400 M.: n. 3225 B.; 100—200 M.: n. 3298 B.; gut from Bethlehem to Cul de Sac 200 M.: n. 3429 B.

St. Croix ex Millspaugh.

Antill., Mexico, Amer. austr. (E. a. P.).

8381. *Coffea* Linn.

Coffea arabica Linn. Spec. (1753) 172; Gris. Fl. 338; Duss 341. Coffee.

St. Eustatius: Bottom of the Quill 300 M.: n. 900 B.; inner slope of the Quill 350 M.: n. 1220 B., 101 H., 215 G.

Saba: Windwardside: n. 1935 B.; Gainbygut: n. 98; Hells-gategut: n. 99.

Indig. in Abyssinia, Angola and Mosambique. (E. a. P.).

8384. *Ixora* Linn.

Ixora coccinea Linn. Spec. (1753) 110.

Cultivated. [Asia trop.] (Ind. Kew.).

Ixora jucunda Thwaites Enum. Pl. Zeyl. (1864) 155. Riceplant.

Cultivated. [Zeylan.] (Ind. Kew.).

8391. *Strumpfia* Jacq.

Strumpfia maritima Jacq. Enum. Pl. Carib. (1760) 28; Gris. Fl. 336; Duss 338.

St. Eustatius: White Wall: n. 813 B., 1264 B., 138.

St. Martin: Colebay (coll. George): n. 2519 B.

Anguilla Boldingh.

Baham., Antill., Aruba, Bonaire, Curaçao. (Herb. Krug et Urban).

8399. *Psychotria* Linn.

Psychotria horizontalis Sw. Prod. Veg. Ind. Occ. (1788) 144; Gris. Fl. 344; Duss 344; Millspaugh 531.

St. Eustatius: Slope of the Quill near Bengalen 50—200 M.: n. 609a B., 657 B.; slope of the Quill above White Wall: n. 1191a B.; locality unknown: n. 218 G.

Saba: near the Sulphurmines: n. 1701a B.; near Devilshand: n. 2075 B., 2076 B.; Springbaygut: n. 2123 B.

St. Martin: from Philipsburg to Belvedere: n. 2575 B., 2586 B.; Marigothill—Marigot 0—200 M.: n. 2642 B.; Nakedboyhill 50—200 M.: n. 2689 B.; mountain between Mont des Accords and Centryhill

200—300 M.: n. 2883 B.; along roads in Cul de Sac: n. 2968 B.; from Belvedere to Oysterpond: n. 3063 B.; lower part of Mill-drumhill: 3161 B.; along roads from Colebay to Simonsbay: n. 3252 B.; Mount Paradis 100—200 M.: n. 3300 B.; French frontier: n. 149.

St. Croix ex Millspaugh.

Antill., Guiana. (Gris. Fl.).

Psychotria pendula Urb. Symb. Ant. I (1900) 445.

Psychotria parasitica Sw. Prodr. (1788) 44; Gris. Fl. 344 (p. p.): Duss 344.

Mapouria parasitica K. Sch. in Engl. u. Prantl. Nat. Pfl. Fam. IV. 4. (1891) 112.

Psychotria crassa Gris. Flor. W. I. (1861) 344 (p. p.); Duss 344. (Symb.).

Saba: Mountain 800 M.: n. 148, 153.

Cuba, St. Kitts, Montserrat, Guadeloupe, Dominica, Martinique, St. Vincent, Grenada. (Symb.).

Psychotria rufescens H. B. et K. Nov. Gen. et Sp. III (1818) 364; Gris. Fl. 342. Bastard cankerberry.

St. Eustatius: Top of the Quill 400 M.: n. 174 B., 349 B., 403 B., 500 B., 117; locality unknown: n. 116, 175 G., 213 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1353 B.; Bottom: n. 1456 B.; Crispeen 400 M.: n. 1762 B.; Mary-point: n. 118; Jenkinsbay: n. 114.

St. Martin: Centryhill: n. 115.

Antill., Cuba and Mexico to Nova Granata. (Gris. Fl.).

Psychotria uliginosa Sw. Prodr. Veg. Ind. Occ. (1788) 43; Gris. Fl. 340; Duss 342.

St. Eustatius: Bottom of the Quill 300 M.: n. 899 B.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1306a B., 1365 B.; Bottom: n. 1448 B.; Mountain: n. 119; Gainbygut: n. 150.

Antill., Amer. centr., Guiana. (Gris. Fl.).

8401. **Palicourea** Aubl.

Palicourea Pavetta DC. Prodr. IV (1830) 525; Gris. Fl. 346; Duss 345; Millspaugh 531.

St. Eustatius: inner slope of the Quill 350 M.: n. 1216 B., 1224 B.; Quill: n. 146, 147.

Saba: along roads from Bottom to Windwardside: n. 1313a B., 1364 B.; Mountain: n. 1755a B.

St. Croix ex Millspaugh.

Cuba, Jamaica, St. Thomas, Guadeloupe. (Herb. Krug et Urban).

8456. **Faramea** Aubl.

Faramea odoratissima DC, Prod. IV (1830) 496; Gris. Fl. 338; Duss 341; Millspaugh 531.

Saba: from Marypoint to Bottom 300 M.: n. 2071a B.

Antill., Mexico. (E. a. P.).

8468. **Ernodea** Swartz.

Ernodea litoralis Sw. Prod. Veg. Ind. Occ. (1788) 29; Gris. Fl. 347; Duss 347; Millspaugh 530.

St. Eustatius: White Wall: n. 806 B., 1263 B.; White rock: n. 96; locality unknown: n. 97.

Saba: Hellsgate—Marypoint: n. 2037 B.

St. Martin: near Mulletpond: n. 2759 B.; Lowlands: n. 3279 B.; near Orientbay: n. 3357 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham., Antill. (Gris. Fl.).

8471. **Diodia** Linn.

Diodia rigida Cham. Schlecht. in Linnaea III (1828) 341; Gris. Fl. 348.

St. Eustatius: Top of the Cliff between Gallowsbay and White Wall: n. 765 B.

Amer. austr. (Ind. Kew.).

8472. **Hemidiodia** K. Schum.

Hemidiodia ocimifolia K. Schum. in Mart. Fl. Bras. IV. 4 (1888) 29.

St. Eustatius: Slope of the Quill near Bengalen 50—200 M.: n. 650 B.; slope of the Quill 200 M.: n. 876 B., 72, 73.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1337 B.; Hellsgate: n. 1691 B.; Mountain 800 M.: n. 1806 B.; Mountain near Windwardside 600 M.: n. 2174 B., 2183 B.; Ladder: n. 74a; Gainbygut: n. 74; locality unknown: n. 138 L., 205 L.

St. Martin: Marigothill—Marigot 0—200 M.: n. 2648 B.; hill on the northern side of Cul de Sac: n. 2982 B.

Antill., Mexico. (E. a. P.).

8473. **Borreria** G. F. Mey.**Borreria ocymoides** DC. Prodr. IV (1830) 544.**Saba:** Hellsgategut: n. 196.

Amer., Africa, Asia. (E. a. P.).

Borreria podocephala DC. Prodr. IV (1830) 542; Gris. Fl. 350; Duss 349.**St. Eustatius:** White Wall: n. 815 B., 818b B.; near Signalhill: n. 943a B., 1039 B.; seashore near English Quarter: n. 1009 B.

Cuba, Pinos, Desirade, Antigua, Guadeloupe. (Herb. Krug et Urban).

8475. **Spermacoce** Gärtn.**Spermacoce tenuior** Linn. Spec. (1753) 102; Gris. Fl. 349; Duss 347; Millspaugh 532.**St. Eustatius:** Top of the Quill 400 M.: n. 230 B.; Bengalen: n. 625 B.; in a garden in Oranjestad: n. 1084 B.; along roads near Glassbottle: n. 1115 B.; slope of the Quill: n. 133 A.; locality unknown: n. 301 G.**Saba:** lower part of the Bottom Mountain 300 M.: n. 1428 B.; between Bottom and Marypoint 100—200 M.: n. 1521 B.; along the Forthgut 150—200 M.: n. 1630 B.**St. Martin:** Nakedboyhill 50—200 M.: n. 2683 B.; in an orchard near Bellevue: n. 2849 B.; Mount Paradis 300—400 M.: n. 3332 B.; Colebayhill: n. 133.**Anguilla** Boldingh.**St. Croix** ex Millspaugh.

Antill., Amer. centr. (E. a. P.).

8479. **Mitracarpus** Zucc.**Mitracarpus polycladus** Urb. in Symbolae Antillanae III (1903) 389.**Saba:** along roads from Bottom to Windwardside n. 1322 B.; Paris: n. 1377 B.; Laddergut: n. 1966a B., 175; locality unknown: n. 86 L.

Portorico. (Symb.).

Mitracarpus villosus Cham. et Schlecht. in Linnaea III (1828) 363; Gris. Fl. 350.*Mitracarpus hirtus* DC. Prodr. IV (1830) 572; Duss 349.**St. Eustatius:** Slope of the Quill: n. 164; road in Oranjestad: n. 173.

Antill., Cuba and Mexico to Ecuador and Venezuela. (Gris. Fl.).

271. Caprifoliaceae.8515. **Sambucus** Linn.**Sambucus canadensis** Linn. Spec. (1753) 269. Elder.

Cultivated. [Amer. sept.] (E. a. P.).

8523. **Lonicera** Linn.**Lonicera japonica** Thunb. Fl. Jap. (1784) 89. White honey-suckle.

Cultivated. [Himalaya]. (Ind. Kew.).

275. Cucurbitaceae.8562. **Melothria** Linn.**Melothria fluminensis** Gard. in Hook. Lond. Journ. Bot. I (1824) 173.**St. Eustatius:** Quill: n. 433 B.; inner slope of the Quill 350 M.: n. 418 B.**Saba:** Windwardside: n. 1933 B.; Gainbygut: n. 35.

Antill., Amer. cent. and austral. (Cogniaux 1881).

8591. **Momordica** Linn.**Momordica Charantia** Linn. Spec. (1753) 1009; Gris. Fl. 287; Duss 307; Millspaugh 533. Maidenapple.**St. Eustatius:** on the slope of the Cliff near Oranjestad: n. 58 B.; along roads in Oranjestad: n. 562 B., 35 H., 42 G.**Saba:** along roads between Bottom and Windwardside 200 M.: n. 1325 B., 1706; locality unknown: n. 163 L.**St. Croix** ex Millspaugh.

In all trop. and subtrop. countries. (Cogniaux 1881).

8599. **Cucumis** Linn.**Cucumis Anguria** Linn. Spec. (1753) 1011; Gris. Fl. 288; Duss 309; Millspaugh 532. Pumpkin.**St. Eustatius:** Locality unknown: n. 156 G.**Saba:** along roads: n. 1707 B.; locality unknown: n. 276 L.**St. Croix** ex Millspaugh.

Florida austr., Antill., Texas, Amer. centr. and austr. sept. (Cogniaux 1881).

8610. **Lagenaria** Ser.

Lagenaria vulgaris Ser. in Mem. Soc. Phys. Genève. III. 1 (1825) 25; Gris. Fl. 288; Duss 308; Sweet gourd.

Saba: Mountain 800 M.: n. 2230 B.; locality unknown: n. 275 L., 308 L.

St. Croix ex Millspaugh.

Indig. in Afr. trop. and India orient. (Cogniaux 1881). Also cultivated.

8622. **Cucurbita** Linn.

Cucurbita maxima Duchesne in Lam. Encyc. II (1786) 151.

Cultivated. [Asia merid.] (Cogniaux 1881).

8636. **Sechium** Juss.

Sechium edule Sw. Fl. Ind. Occ. II (1800) 1150; Gris. Fl. 286; Duss 306.

Saba: Boobyhill: n. 1567 B.; Windwardside: n. 1595 B.; Mountain 800 M.: n. 2198a B.

Antill., Mexico, Amer. centr., Panama., Nova Granata, Venezuela, Brasilia. (Cogniaux 1881). Also cultivated.

277. **Goodeniaceae**8716. **Scaevola** Linn.

Scaevola Plumieri Vahl Symb. II (1791) 36; Gris. Fl. 388; Duss 378; Millspaugh 533.

Scaevola Lobelia Murr. Syst. ed. XIII (1774) 178.

St. Martin: Colebay (coll. George): n. 2503a B.; near Simons-baylagoon: n. 3035b B.

Bermud., Baham., Antill. (Herb. Krug et Urban).

280. **Compositae.**8751. **Vernonia** Schreb.

Vernonia punctata Sw. ex Wikstr. in Vet. Acad. Handl. Stockh. (1827) 72; Gris. Fl. 353; Duss 351.

Vernonia longifolia Pers. Syn. II (1807) 404. Wild tobacco.

St. Eustatius: Signalhill 200 M.: n. 138 B.; top of the Quill 400 M.: n. 206 B., 309 B., 328 B.; slope of the Quill near Bengalen 50—200 M.: n. 658 B., 669 B.; along roads near English Quarter: n. 1006 B.; along roads near Bengalen: n. 1197 B.; slope of the Quill: n. 55, 57; slope of Signalhill: n. 56; locality unknown: n. 198 H., 272 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1280 B.; Crespeen 400 M.: n. 1719 B.; Laddergut 0—200 M.: n. 1968 B.; Ladder: n. 54.

St. Martin: from Filipsburg to Guanabay: n. 2420 B.; Marigot-hill—Marigot 0—200 M.: n. 2641 B.; near Mulletpond: n. 2755 B.; hill on the northern side of Cul de Sac: n. 2975 B.; from Belvedere to Oysterpond: n. 3059 B.; Mount Paradis 150—400 M.: n. 3194 B.; 3348 B.; Centryhill: n. 319.

Antill., Mexico, Honduras. (Gris. Fl.).

8775. *Elephantopus* Linn.

Elephantopus spicatus Aubl. Pl. Guian. II (1775) 808; Duss 352.

Distreptus spicatus Cass. in Bull. Soc. Philom. (1817) 66; Gris. Fl. 355; Millspaugh 534.

St. Eustatius: Slope of the Quill near Bengalen: n. 711a B.

Saba: Boobyhill: n. 1590a B.; Crespeen—Rendez-vous: n. 199; Rendez-vous—Mountain 4—600 M.: n. 198.

St. Croix ex Millspaugh.

Antill., Cuba and Mexico to Venezuela and Peru. (Gris. Fl.).

8795. *Ageratum* Linn.

Ageratum conyzoides Linn. Spec. (1753) 839; Gris. Fl. 356; Duss 353; Millspaugh 533. White cap.

St. Eustatius: in a garden near Oranjestad: n. 77 B.; along roads in Oranjestad: n. 545 B.; slope of the Quill: n. 290, 291, 292; locality unknown: n. 6 G., 6 H., 230 H.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1318 B.; Greathill 400 M.: n. 1378 B.; Mountain 600 M.: n. 1781 B.; Laddergut 0—200 M.: n. 2022 B.; Crespeen—Rendez-vous: n. 289; Gainbygut: n. 288; locality unknown: n. 20 L., 135 L.

St. Martin: Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2899 B.; Milldrumhill in plantations 300 M.: n. 3130 B.

St. Croix ex Millspaugh.

In all warmer countries of both hemisph. (E. a. P.).

8816. *Eupatorium* Linn.

Eupatorium canescens Vahl. in West Bidr. St. Croix (1793) 302; Gris. Fl. 360; Duss 355.

Eupatorium sinuatum Lam. Encycl. II (1786) 407; Millspaugh 534.

St. Eustatius: Gut near Fort de Wind: n. 841a B.; White Wall: n. 814a B.

St. Croix ex Millspaugh.

Cuba, Domingo, Portorico, Antigua, Guadeloupe. (Herb. Krug et Urban).

Eupatorium conyzoides Vahl. Symb. Bot. III (1807) 96; Gris. Fl. 358.

St. Eustatius: Top of the Quill 400 M.: n. 184 B., 258 B., 501 B.; along roads near Bengalen: n. 1196 B.; slope of the Quill: n. 25, 26, 288; road to Roundhill: n. 24.

Saba: Crispeen 400 M.: n. 1717 B.; Laddergut: n. 1999 B.; Flatpoint: n. 2126 B.; Peperpot: n. 23; Mountain: n. 290; Gainbygut: n. 289.

St. Martin: lower part of the Milldrumhill: n. 3162 B.

Bermud., Florida austr., Antill., Mexico to Brazil. (Hemsley 1884).

Eupatorium iresinoides H. B. et K. Nov. Gen. et Spec. IV (1820) 105. t. 340; Gris. Fl. 360; Duss 355.

Saba: Bottom: 200 M.: n. 1446 B.; from Bottom to Marypoint 100—200 M.: n. 1515a B.; Hellsgate: n. 1690 B.; Crispeen 400 M.: n. 1725 B.; Peperpot: n. 29; Crispeen Rendez-vous: 400 M.: n. 28.

Antill., Panama to Venezuela. (Gris. Fl.).

Eupatorium macranthum Sw. Fl. Ind. Occ. III (1806) 1315.

St. Eustatius: Top of the Quill 400 M.: n. 307 B., 32, 30, 293.

Saba: Mountain near Windwardside 600 M.: n. 2154 B., 2178 B.; Crespeen—Rendez-vous 400 M.: n. 31, 293.

St. Kitts. (Herb. Krug et Urban).

Eupatorium macrophyllum Linn. Spec. (1763) 1175; Millspaugh 534.

Hebeclinium macrophyllum DC. Prodr. V (1836) 136; Gris. Fl. 356; Duss 353.

Saba: between Bottom and Marypoint 300 M.: n. 1487 B.; 100—200 M.: n. 1511 B.; Laddergut 0—200 M.: n. 1965 B.; Gainbygut: n. 33.

St. Martin: Mount Paradis 300—400 M.: n. 3349 B.

St. Croix ex Millspaugh.

Bermud., throughout the Antill. and southward to Brazil. (Hemsley 1884).

Eupatorium paniculatum Schrad. Ind. Sem. Hort. Gotting. (1832) 2; Gris. Fl. 362; Duss 356.

Eupatorium guadalupense Spreng. Syst. III (1826) 414.

Saba: Top of the Mountain 800 M.: n. 1806 B.; Hellsgate: n. 294; Marypointgut: n. 27.

Antill., Honduras, Peru, Venezuela, Brazil. (Gris. Fl.).

8901. **Erigeron** Linn.

Erigeron bonariensis Linn. Spec. (1753) 863; Gris. Fl. 365; Duss 360.

Saba: Mountain 6—800 M.: n. 11.

Antill., Amer. centr. to Buenos Ayres. (Gris. Fl.).

Erigeron canadensis Linn. Spec. (1753) 863; Gris. Fl. 365; Duss 360. Foxtail.

St. Eustatius: Slope of the Quill: n. 13; Cultuurvlakte: n. 12; locality unknown: n. 145 G., 173 H.

Saba: Hellsgate—Marypoint: n. 2033 B.; Rendez-vous 400 M.: n. 296; locality unknown: n. 248 L., 259 L.

St. Martin: near Mulletpond: n. 2760 B.; lower part of Mill-drumhill: n. 3169 B.

Bermud., Canada to Brazil, and naturalised in many other countries of both hemisph. (Hemsley 1884).

Erigeron spathulatus Vahl, in West. Bidr. St. Croix (1793) 303; Gris. Fl. 365; Duss 359; Millspaugh 534.

Saba: between Bottom and Marypoint 100—200 M.: n. 1502 B., 1558 B.; Laddergut 0—200 M.: n. 1958 B., 286; Peperpol: n. 20; Crespeen—Rendez-vous: n. 19; Gainbygut: n. 287.

St. Martin: Milldrumhill in plantations 300 M.: n. 3128 B.

St. Croix ex Millspaugh.

Hispaniola, Portorico, Martinique, Guadeloupe, Trinidad. (Herb. Krug et Urban).

8915. **Gundlachia** A. Gray.

Gundlachia domingensis A. Gray Bot. Contr. in Proc. Amer. Acad. XVI (1880) 100.

Baccharis dioica Gris. Kar. (1857) 85; Gris. Fl. 366; Duss 361.

Saba: near the Sulphurmines: n. 1700 B.

Baham., Antill. (Symb.).

8933. **Baccharis** Linn.

Baccharis cotinifolia Urb. in Urb. Symbol. Antill. III (1903) 406; Gris. Fl. 366; Duss 360.

Saba: Top of the Mountain: n. 274.

Antill., Costarica. (Symb.).

8941. **Pluchea** Cass.

Pluchea odorata Cass. in Dict. Sc. Nat. XLII (1826) 3; Gris. Fl. 366; Duss 361; Millspaugh 535. Wild tobacco.

St. Eustatius: Slope of the Cliff near Gallowsbay: n. 62 B.; seashore: n. 30; locality unknown: n. 61 H., 116 G.

Saba: between Bottom and Marypoint 100—200 M.: n. 1526 B.; Crispeen—Rendez-vous 400 M.: n. 303; locality unknown: n. 228 L., 234 L.

St. Martin: Milldrumhill in plantations 300 M.: n. 3127 B.

St. Croix ex Millspaugh.

Antill., Guiana., Amer. centr. (Hemsley 1884).

Pluchea purpurascens DC. Prod. V (1836) 452; Gris. Fl. 367; Duss 362; Millspaugh 535.

St. Martin: Cul de Sac—Marigothill 0—200 M.: n. 2621 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Bermud., Florida and Mexico, Antill., to Nova Granata. (Hemsley 1884).

8992. **Gnaphalium** Linn.

Gnaphalium purpureum Linn. Spec. (1753) 854. vel. affin.

Gnaphalium americanum Mill. Gard. Dict. ed. VIII (1768) n. 17; Duss 373.

Saba: Paris 400 M.: n. 1375 B.; between Bottom and Marypoint 100—200 M.: n. 1502a B.; Greathill: n. 37; Jenkeygut: n. 36.

Amer. and warmer parts of Africa, Asia and Australia. (E. a. P.).

9101. **Lagascea** Cav.

Lagascea mollis Cav. in Anal. Cienc. Nat. VI (1803) 333 t. 44.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1310 B.; along roads 200—300 M.: n. 1708 B.; Windwardside: n. 2091 B., 67.

St. Martin: from Filipsburg to Guanabay: n. 2403 B., 2415 B.; Cul de Sac—Marigothill 0—200 M.: n. 2631 B.; hill on the northern side of Cul de Sac: n. 2972 B.; French frontier: n. 68; Marigotroad: n. 69, 299.

Anguilla Boldingh.

Indig. in Amer. centr. (E. a. P.).

9117. **Clibadium** Linn.

Clibadium erosum DC. Prod. V (1836) 506; Gris. Fl. 368; Duss 362.

Saba: Mountain 6—800 M.: n. 179.

Portorico, Dominica, Martinique, St. Vincent. (Herb. Krug et Urban).

9130. **Acanthospermum** Schrank.

Acanthospermum humile DC. Prod. V (1836) 522; Gris. Fl. 369; Millspaugh 533.

St. Martin: Cul de Sac—Marigot 0—200 M.: n. 2657 B.; along roads from Philipsburg to Marigot: n. 3109 B.

St. Croix ex Millspaugh.

Jamaica, Cuba, Haiti, Panama. (Gris. Fl.).

9138. **Parthenium** Linn.

Parthenium Hysterophorus Linn. Spec. (1753) 988; Gris. Fl. 369; Duss 365; Millspaugh 535.

St. Eustatius: in a garden in Oranjestad: n. 79 B., 10; along roads in Oranjestad: n. 520 B., 3 G., 3 H.

Saba: Laddergut: n. 1954 B., 11; Mountain near Windward-side 600 M.: n. 2165 B.; locality unknown: n. 56 L., 95 L.

St. Martin: from Philipsburg to Guanabay: n. 2398 B.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Antill., Amer. sept. and centr., and in other parts of both hemisph. (E. a. P.).

9146. **Ambrosia** Linn.

Ambrosia hispida Purch. Fl. Am. Sept. Suppl. (1814) 743.

Ambrosia crithmifolia DC. Prod. V (1836) 525.

St. Eustatius: in a garden in Oranjestad: n. 157 G., 1102 B.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1303 B.; Windwardside: n. 1949 B.

St. Martin: from Philipsburg to Belvedere: n. 2561 B.

Florida. (Ind. Kew.). Also cultivated.

9148. **Xanthium** Linn.

Xanthium orientale Linn. Spec. (1763) 1400; Duss 364.

Saba: Bottom: n. 62.

In almost all warmer countries of both hemisph. (E. a. P.).

9155. **Zinnia** Linn.**Zinnia elegans** Jacq. Coll. III 152.

Cultivated. [Patria ignota.].

9166. **Eclipta** Linn.**Eclipta alba** Hassk. Pl. Jav. Rar. (1848) 528; Gris. Fl. 370; Duss 366; Millspaugh 534.*Eclipta erecta* Linn. Mant. II (1771) 286.**St. Martin**: lower part of Mount Paradis: n. 3187 B.**St. Croix** ex Millspaugh.

In the warmer countries of both hemisph. (E. a. P.).

9191. **Borrichia** Adans.**Borrichia arborescens** DC. Prod. V (1836) 489; Gris. Fl. 371; Duss 366; Millspaugh 533.**Saba**: Peperpot: n. 278.**St. Martin**: Point Blanche (coll. George): n. 2493 B.; Colebay (coll. George): n. 2506 B.; near Mulletpond: n. 2750 B.; near Simonsbayvillage: n. 3030 B.; Tintamarre: n. 3373 B.; Oysterpond: n. 306; Guanabay: n. 304, 305.**St. Croix** ex Millspaugh.

Antill., Bermud., Key of Florida, seashore of the Carib. Sea to Peru. (Hemsley 1884; E. a. P.).

9192. **Wedelia** Jacq.**Wedelia bupthalmoides** Gris. in Goett. Abh. VII (1857) 235; Gris. Fl. 372; Duss 367; Millspaugh 536. Yellow sage.**St. Eustatius**: Top of the Quill 400 M.: n. 274 B., 286 B.; slope of the Quill near Bengalen 150 M.: n. 681 B., 687 B.; gut near Fort de Wind 50 M.: n. 836 B.; along roads near Glassbottle: n. 866 B.; from Little Mountain to Venusbay: n. 996 B.; slope of the Quill on the side of White Wall 200 M.: n. 1171 B.; along roads near Bengalen: n. 1198 B.; Gilboohill: n. 58; Sugarloaf: n. 317; locality unknown: n. 165 G., 233 H.**Saba**: along roads between Bottom and Windwardside 200 M.: n. 1284 B., 1298 B.; between Bottom and Marypoint 200 M.: n. 1553 B.; Helsgate: n. 1694 B.; Laddergut 0—200 M.: n. 1960 B.; gut between Hellsgate and Masticgut 300—400 M.: n. 2058 B.; Windwardside: n. 2315 B.; Peperpot: n. 318; Paris 2—400 M.: n. 61; Crespeen—Rendez-vous: n. 60; locality unknown: n. 316, 29 L.**St. Martin**: from Filipsburg to Guanabay: n. 2410 B.; Naked-

boyhill 50—200 M.: n. 2703 B., 200—260 M.: n. 2719 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2884 B.; hills on the northern side of Cul de Sac: n. 2984 B.; lower part of Mount Paradis: n. 3291a B.; Colebayhill: 59; John Ednyhill: n. 315.

I also saw *Wedelia bupthalamoides*: on the top of Mount Paradis 400 M.; hills on the eastern side of Cul de Sac.

Anguilla Boldingh.

St. Croix ex Millspaugh.

Baham. (Hitche.); Antill. (Herb. Krug et Urban).

9196. **Tithonia** Desf.

Tithonia tagetiflora Desf. in Ann. Mus. Par. I (1802) 46. t. 4. Cultivated. [Mexico]. (E. a. P.).

9207. **Spilanthes** Jacq.

Spilanthes uliginosa Sw. Prod. Veg. Ind. Occ. (1788) 110; Gris. Fl. 376; Duss 370.

St. Eustatius: near English Quarter: n. 1018 B.

Warmer countries of both hemisph. (E. a. P.).

9224. **Synedrella** Gärtn.

Synedrella nodiflora Gärtn. Fruct. II (1791) 456. t. 171; Gris. Fl. 377; Duss 371.

Ucacoua nodiflora Hitch. in Missouri. Bot. Gard. Rep. IV (1893) 100; Millspaugh 535.

St. Eustatius: Slope of the Quill: n. 47; locality unknown: n. 242 H.

Saba: along roads between Bottom and Windwardside 200 M.: 1332 B.; lower part of the Bottom Mountain 250 M.: n. 1397 B.; along the Forthgut 150—200 M.: n. 1629 B.; Mountain 600 M.: n. 1754 B.

St. Martin: Nakedboyhill 50—200 M.: n. 2706 B.; mountain between Mont des Accords and Centryhill 0—200 M.: n. 2852 B., 200—300 M.: n. 2892 B.

St. Croix ex Millspaugh.

Indig. in Amer. trop.; dispersed in the Old world. (E. a. P.).

9237. **Bidens** Linn.

Bidens bipinnatus Linn. Spec. (1753) 832; Gris. Fl. 373; Duss 369; Millspaugh 533.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1333 B., 1366 B.; Rendez-vous: n. 275; Rendez-vous—Crespeen: n. 276; locality unknown: n. 255 L.

St. Martin: Hill on the northern side of Cul de Sac 0—100 M.: n. 3003 B.; along roads from Grande Case to Marigot: n. 3099 B.; French frontier: n. 295.

St. Croix ex Millspaugh.

Indig. in Amer., dispersed in the warmer countries of the Old world. (E. a. P.).

Bidens leucantha Willd. Sp. Pl. III (1800) 1719; Gris. Fl. 373; Duss 368; Millspaugh 533.

Bidens pilosa Linn. Spec. (1753) 832.

St. Eustatius: Slope of the Quill: n. 302.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1282 B.; Windwardside—Hellsgate: n. 1672 B.; seashore near Flat Point: n. 2139 B.; Crespeen—Rendez-vous n. 301; Gainbygut: n. 303.

St. Martin: Hill on the northern side of Cul de Sac 0—100 M.: n. 2978 B.; Marigotroad: n. 279; locality unknown: n. 3602 B.

St. Croix ex Millspaugh.

Indig. in Amer., dispersed in the warmer countries of the Old world. (E. a. P.).

9238. **Cosmos** Cav.

Cosmos sulfureus Cav. Ic. I (1791) 56 t. 79. Poppy.

Cultivated. from Cuba and Mexico to Brazil.] (Gris. Fl.).

9306. **Gaillardia** Fouger.

Gaillardia aristata Purch. Fl. Am. Sept. II (1814) 573.

Cultivated. [Amer. bor.] (Ind. Kew.).

9316. **Porophyllum** Adans.

Porophyllum ellipticum Cass. Dict. XLIII (1826) 56.

Porophyllum ruderales Gris. Flor. W. I. (1861) 379; Duss 372. (p. p.). (Symb.).

St. Eustatius: along roads in Oranjestad: n. 1111 B.; slope of the Quill: n. 36, 37, 38; locality unknown: n. 277 H., 284 G.

Saba: lower part of the Bottom Mountain 300 M.: n. 1435 B.; between Bottom and Marypoint 100—200 M.: n. 1555 B.; locality unknown: n. 269 L.

St. Martin: Locality unknown: n. 3603 B.
Baham., Antill., Amer. cont. trop. (Symb.).

9319. **Pectis** Linn.

Pectis febrifuga van Hall. in Ann. Hort. et Bot. ou Flore Jard. des Pays-Bas vol. IV (1861) 33 cum tab.

Pectis linifolia Less. in Linnaea VI (1831) 709. (excl. syn.); (Symb.).

Pectis Swartziana Borg et Pauls. Veget. Dansk-vestind. Oer (1898) 110 — non Less; (Symb.).

Saba: Seashore near Forthbay: n. 1641 B.

Antill., Bonaire, Curaçao, Aruba, Columbia. (Symb.).

Pectis humifusa Sw. Prod. Veg. Ind. Occ. (1788) 114; Gris. Fl. 378; Duss 372; Millspaugh 535.

St. Eustatius: White Wall: n. 824 B.; Gilboohill 150 M.: n. 1122 B.; near Boven 200 M.: n. 1249 B.; near White Wall: n. 1258 B.; Venusbay: n. 20; locality unknown: n. 3586 B.

Saba: along the lower part of Forthgut: n. 1642 B.; St. Giles: n. 1869 B.

St. Martin: from Philipsburg to Guanabay: n. 2366 B., 2414 B., 19; near Fort Willem: n. 2801 B.; near Fort Amsterdam: n. 18.

Anguilla Boldingh.

St. Croix ex Millspaugh.

St. Domingo, Portorico, St. Thomas, St. Jan, St. Barthélemy, St. Kitts, Antigua, Guadeloupe, Desirade, Marie—Galante, Dominica, Martinique, St. Lucia, St. Vincent, Mustique, Barbados. (Symb.).

9376. **Neurolaena** R. Br.

Neurolaena lobata R. Br. in Trans. Linn. Soc. XV (1817) 120; Gris. Fl. 381; Duss 373.

St. Eustatius: Slope of the Quill above White Wall: n. 1188 B.

Saba: Top of the Mountain 800 M.: n. 1836 B.; locality unknown: n. 210 L.

Antill., Mexico, Columbia. (E. a. P.).

9389. **Erechthites** Raf.

Erechthites hieracifolia Raf. ex DC. Prodr. VI (1837) 294; Gris. Fl. 381; Duss 374; Millspaugh 534.

Erechthites praeulta Raf. Fl. Ludov. (1817) 65.

St. Eustatius: Slope of the Quill near Bengalen: n. 9.

Saba: between Bottom and Marypoint 300 M.: n. 1485 B.; Gainbygut: n. 10.

St. Croix ex Millspaugh.

In all warmer parts of Amer., also in the Old world. (E. a. P.).

Erechthites valerianaefolia DC. Prod. VI (1837) 295.

Saba: between Bottom and Marypoint 350 M.: n. 1491 B.; Mountain 600—800 M.: n. 1786 B.; locality unknown: 166 L., 193 L. Amer. centr. and sept., Java. (Herb. Berlin).

9411. **Senecio** Linn.

Senecio sonchifolius Moench. Meth. Suppl. (1802) 231.

Emilia sonchifolia DC. Prod. VI (1837) 302; Gris. Fl. 381; Duss 374; Millspaugh 534.

St. Eustatius: near Boven 200 M.: n. 1234 B.; locality unknown n. 126 G.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1285 B., 1348 B.; Windwardside: n. 2309 B.

I also saw *Senecio sonchifolius* in the Laddergut.

St. Martin: Nakedboyhill 50—200 M.: n. 2707 B.; Mountain between Mont des Accords and Centryhill 200—300 M.: n. 2890 B.; Milldrumhill in plantations 300 M.: n. 3129 B.

I also saw *Senecio sonchifolius*: Mount Paradis 300—400 M.

St. Croix ex Millspaugh.

Indig. in the trop. of the Old world. (E. a. P.).

9529. **Chaptalia** Vent.

Chaptalia nutans Hemsl. Biol. Centr. Am. Bot. II 255; Millspaugh 533.

Leria nutans DC. Ann. Mus. Par. XIX (1812) 68; Gris. Fl. 383; Duss 375.

St. Eustatius: Slope of the Quill: n. 310.

Saba: along roads between Bottom and Windwardside 200 M.: n. 1315 B.; Mountain 400 M.: n. 1769 B., 1774 B., 308; Crespeen—Rendez-vous: n. 307, 309.

St. Martin: Milldrumhill 300 M.: n. 3179 B.; Mount Paradis 300—400 M.: n. 3339 B.

St. Croix ex Millspaugh.

In all warmer countries of Amer. (E. a. P.).

9595. **Sonchus** Linn.

Sonchus oleraceus Linn. Spec. (1753) 794; Gris. Fl. 384; Duss 375; Millspaugh 535. Wild salad.

St. Eustatius: Glassgut at the seashore near Oranjestad: n. 48 B.; Bengalen: n. 606 B.; along roads near Glassbottle: n. 884 B.; slope of the Quill: n. 41; Roundhill: n. 302.

Saba: lower part of the Bottom Mountain 300 M.: n. 1430 B.; Mountain 600 M.: n. 1780 B.; Bottom: n. 40; locality unknown: n. 40 L.

St. Martin: Milldrumhill in plantations 300 M.: n. 3133 B.

St. Croix ex Millspaugh.

Widely dispersed in both hemisph. (E. a. P.).

9596. **Lactuca** Linn.

Lactuca intybacea Jacq. Ic. Pl. Rar. I (1781) t. 162; Millspaugh 534.

Brachyrhamphus intybaceus DC. Prod. VII (1838) 177; Gris. Fl. 384; Duss 376.

St. Eustatius: Cultuurvlakte: n. 57; seashore: n. 307.

St. Croix ex Millspaugh.

Antill., Nicaragua, East Indies, Africa trop. (Gris. Fl.).

SECOND PART

HISTORICAL

HISTORICAL

CHAPTER I

HISTORICAL SKETCH.

The following persons have given informations about the Flora of the Dutch West Indian Islands. *Liter. Urban Symbolae Antillanae* Vol. I and Vol. III.

St. Eustatius.

J. B. Labat, between 1694 and 1705.

J. B. Labat in his: *Nouveau voyage aux isles de l'Amérique...* Editio 1724. La Haye, does not give any compendium of the vegetation of the Dutch Antilles.

N. J. Jacquin between 1755 and 1757.

N. J. Jacquin in his: *Selectarum stirpium americanum historia*, in qua ad Linnaeanum systema determinatae descriptaeque sistuntur plantae illae, quas in insulis Martinica, Jamaica, Domingo aliisque et in vicinae Continente parte observavit rariores; adjectis iconibus in solo natali delineastis. Vindobonae 1763, mentions only 6 plants found in St. Eustatius, of which I point out only those quoted by *Grisebach* in his: *Systematische Untersuchungen über die Vegetation der Karaihen*, insbesondere der Insel Guadeloupe: *Justicia sessilis*.

Justitia eustachyana. (*Beloperone eustachiana* Benth.).

Heliotropium gnaphaloides. (*Tournefortia gnaphalodes* R. Br.).

Coccoloba nivea.

Cactus triangularis. (*Cereus triangularis* Haw.).

(Herb. Banks. Brit. Mus. London; Herb. Willd. Berlin).

F. Masson 1780.

I only saw *Masson* mentioned by *O. Swartz* in his: *Flora Indiae occidentalis*, concerning *Microtea debilis*.

(Brit. Mus. London).

S. Fahlberg between 1785 and 1834.

I only saw *Fahlberg* mentioned by *O. Swartz* in his *Flora Indiae occidentalis* with reference to a few plants from *St. Barthé-lémy*.

(Herb. Swartz. Stockholm; herb. Thunberg Upsala).

Placide Duchassaing de Fontbressis about 1848.

P. D. de Fontbressis collected only a few plants in *St. Eustatius*; I saw the following mentioned in *Grisbach*: *Systematische Untersuchungen über die Vegetation der Karaiben, insbesondere der Insel Guadeloupe*:

no. 16. *Lepidium virginicum* Linn.

no. 22. *Polanisia viscosa* DC.

no. 24. *Capparis cynophallophora* Linn. ? *attenuata*.

no. 162. *Sida spinosa* Linn.

no. 286. *Tribulus maximus* Linn. (*Kallstroemia maxima* Wight et Arn.).

no. 386. *Stylosanthes procumbens* Sw. (*Stylosanthes hamata* Taubert).

no. 942. *Cyphiacanthus barbadensis* Ns. (*Ruellia tuberosa* Linn.).

no. 1047. *Heliophytum indicum* DC. (*Heliotropium indicum* Linn.).

no. 1074. *Lantana camara* Linn.

no. 1076. *Lantana involucrata* Linn.

(Herb. Göttingen).

B. A. Euphrasen 1788.

B. A. Euphrasen in his: *Beskrifning öfver Svenska vestindiska ön St. Barthélémi samt öarne St. Eustache och St. Christopher. Stockholm 1798*, makes also mention of a few plants from *St. Eustatius*. I did not think it necessary to quote them here as, according to the statement of *Urban* (*Symbolae Antillanae* I p. 49) it is impossible to control the determination of *Euphrasen*.

F. L. l'Herminier 1815.

M. D. Feenstra about 1836.

The few botanical data I could take from his: *De Nederlandsche West-indische eilanden in derzelver tegenwoordigen toestand, Amsterdam. 1836—1837* are of no use to the *Systematic Part*. In describing the vegetation in the *Third Part* I shall say something about this paper of *Feenstra*.

W. F. R. Suringar 1885.

W. F. R. Suringar made a botanical investigation on the Dutch West Indian Islands in 1885.

Further details about this expedition will be found in Chapter II of the Historical Part of this Flora.

(Herb. Leiden, Herb. Krug et Urban Berlin).

J. J. Walch about 1889.

In the Report of the 60 meeting of the Br. Ass. f. Adv. Sc. Leeds 1890 p. 448 the following is said about the collection *Walch*: A small collection of plants numbering 143 specimens, was received from *Mr. J. J. Walch, R. N.* This collections includes plants from Dominica, St. Martin, St. Eustatius, St. Kitts, St. Lucia and Grenada. Most of the plants consisted of common West Indian species, presumably such as would be met with in the more accessible spots in the various places visited.

(Herbar. Kew.).

F. A. F. C. Went 1902

F. A. F. C. Went is his: Rapport omtrent den toestand van land- en tuinbouw op de Nederlandsche Antillen, bijlage V van het Koloniaal verslag van Curaçao van 1902, gives some observations on the vegetation of St. Eustatius; in so far as these observations will be interesting to us I shall deal with them in the Third Part of this Flora.

J. van Grol-Meyers 1904—1906.

Two collections of plants gathered by *Mrs. van Grol-Meyers* will be treated more fully in Chapter II of the Historical Part of this Flora.

(Herb. Utrecht; Herb. Koloniaal Museum Haarlem).

I. Boldingh 1906.

The botanical investigations I made in 1906 will more largely be dwelt on in Chapter II of the Historical Part of this Flora.

Besides the above mentioned data I found in *Grisebach*: Flora of the British W. I. Islands that the following plants were also collected in St. Eustatius:

p. 66. *Lithophila muscoides* Sw.

p. 75. *Sida ulmifolia* Cav.

And in *Grisebach*: Systematische Untersuchungen über die Vegetation der Karaien, insbesondere der Insel Guadeloupe I found: 1472. *Aspidium macrophyllum* Sw. (*Aspidium martinicense* Spr.).

Saba.

J. H. Labat between 1694 and 1705.

Conformable to what I mentioned under St. Eustatius.

J. E. Forsström 1805—1812.

I saw *Forsström* mentioned only in reference to a few plants in *Urban*, *Symbolae Antillanae*.
(Herb. Swartz and Herb. Stockholm).

M. D. Feenstra about 1836.

Conformable to what I mentioned under St. Eustatius.

Placide Duchassaing de Fontbressis about 1848.

According to *Grisebach*: Systematische Untersuchungen über die Vegetation der Karaiben, insbesondere der Insel Guadeloupe, *P. D. de Fontbressis* collected in Saba: n. 187. *Guazuma tomentosa* Kth.

O. Kuntze 1874.

Besides at *Caesalpinia ciliata* Urb. I did not see *O. Kuntze* mentioned at the label of any plant in the Herbarium of Krug and Urban in Berlin. In his: Um die Erde. Reisebericht eines Naturforschers, Leipzig 1881, *Kuntze* mentions only *Melocactus communis* Link et Otto.

(Herb. Krug et Urban Berlin).

W. F. R. Suringar 1885.

F. A. F. C. Went 1902.

J. Boldingh 1906.

For these three persons I refer to what was said under St. Eustatius.

A. C. W. Lionarons 1907.

The plants gathered by *Lionarons* will fully be treated in Chapter II of the Historical Part of this Flora.

St. Martin.

J. B. du Tertre about 1650—1656.

J. B. Labat between 1694 and 1705.

Conformable to what I mentioned under St. Eustatius.

N. J. Jacquin between 1755 and 1757.

N. J. Jacquin in his: Select. Stirp. Amer., does not mention but 8 plants found in St. Martin. I will quote only the plants enu-

merated in *Grisebach* Systematische Untersuchungen über die Vegetation der Karaiben, insbesondere der Insel Guadeloupe:

Myginda uragoga cf. *Symb. Antill.* V. p. 80. (*Rhacoma crosso-petalum* Linn.).

Rhamnus colubrinus (*Colubrina ferruginosa* Brong.).

Malpighia linearis (*Malpighia angustifolia* Linn.).

Corchorus hirsutus Linn.

Lobelia Plumieri (*Scaevola Plumieri* Vahl).

A. Plé 1821.

After a discussion I had with *Prof. I. Urban*, I think it superfluous to quote the few plants that are collected by *Mr. Plé*. (*Herb. Mus. Paris*).

M. D. Feenstra about 1836.

W. F. R. Suringar 1885.

J. J. Walch about 1889.

F. A. F. C. Went 1902.

I. Boldingh 1906.

For these 5 persons I refer to what was said under *St. Eustatius*.

According to *Grisebach*: Systematische Untersuchungen über die Vegetation der Karaiben, insbesondere der Insel Guadeloupe, also the following plants were collected in *St. Martin*:

n. 868. *Metastelma paralias* Decs. (*Metastelma parviflorum* R. Br.).

n. 1009. *Cuscuta americana* Linn.

According to the notes I made about the persons and collections afore said I did not think it desirable to treat any of the smaller collections. The difficulties in getting all these plants would not be in proportion to the small amount of greater completeness of the plants then mentioned in this *Flora*.

So I have treated in the Systematic Part only the collections made by *Suringar*, *van Grol-Meyers*, *Boldingh* and *Lionarons*.

It will appear from these collections that I have about 5000 numbers of plants at my disposal, a big number when we take into consideration that the surface of the 3 Islands does not surpass 140 □ km.

CHAPTER II

NOTES ON THE LARGER COLLECTIONS.

Collection W. F. R. Suringar.

Dr. J. Valckenier Suringar was so kind as to allow me to make use of the diary written by his father *Prof. W. F. R. Suringar* during his botanical investigations of the Dutch West Indian Islands in 1885.

From these notes together with the data drawn from the labels attached to the collected plants I was enabled to make the following list of the localities visited by *Suringar* and the numbers of plants collected there.

St. Eustatius: April 8th—18th and May 7th—18th.

April 8th.	Concordiabay, Turtlebay.	}	coll. 70 nos.
" 15th.	Fairplay, English Quarter.		
May 7th.	Bengalen.		
April 9th.	Signalhill and environment.	}	coll. 50 nos.
" 11th.	Little Mountain, Tumbledowndiekbay.		
" 17th.	Boven, Little Mountain.		
May 8th.	Fort Amsterdam.		
" 9th.	Gilboohill.	}	coll. 60 nos.
April 8th.	Seashore.		
" 10th.	Seashore.		
" 11th.	Sugarloaf and White Wall.	}	coll. 35 nos.
May 11th.	Sugarloaf.		
April 13th.	}		Quill. coll. 170 nos.
" 14th.			
" 16th.			
May 18th.			
" 13th.			

On his 16 excursions through St. Eustatius *Suringar* has evidently collected 385 nos.

All the localities visited by *Suringar* were also investigated by me.

In proportion to the other two islands St. Eustatius was investigated by *Suringar* best of all, thanks to a longer stay there.

Saba: April 21—28th.

April 22th. Paris and Greathill and surroundings of the Bottom.
coll. 50 nos.

" 22th. Peperpot near Springbaygut. coll. 35 nos.

" 23th. Top of the Mountain. coll. 100 nos.

" 24th. |

" 26th. | Surroundings of the Bottom. coll. 50 nos.

" 28th. |

" 25th. Gainbygut. coll. 30 nos.

" 27th. Round the Island. coll. 90 nos.

" 27th. Laddergut. coll. 40 nos.

Suringar has evidently collected during these 9 excursions
395 nos.

I myself saw all the localities visited by *Suringar*.

St. Martin: April 30th—May 5th.

April 30th. Saltpond and Fort Amsterdam. coll. 40 nos.

May 1st. Centryhill and Surroundings. coll. 45 nos.

" 2nd. from Philipsburg to Simonsbay. coll. 50 nos.

" 3rd. Nakedboyhill, Princequarter. coll. 25 nos.

" 4th. from Philipsburg to Cul de Sac and French frontier.
coll. 60 nos.

" 5th. from Philipsburg to Guanabay. coll. 20 nos.

Suringar has collected during these 6 excursions 240 nos.

The short time *Suringar* could spend on St. Martin is the reason why this island was explored but on a very small scale; the French part of St. Martin has not been studied; all the localities named by *Suringar* were also visited by me.

The collection *Suringar* is preserved is „'s Rijks-herbarium" at Leiden; duplicates are to be found in the collection Herb. Krug et Urban in Berlin (K. Bot. Gart. und Museum Steglitz-Dahlem). The greater part of the *Suringar* plants were determined by *I. Urban* before I took up the study of the West Indian Flora and another part was determined by him afterwards.

Collections Mrs. J. van Grol-Meyers.

The two collections of dried plants given by *Mrs. J. van Grol-Meyers* to the Koloniaal Museum at Haarlem and the Her-

barium of the Botanical Laboratory of the University of Utrecht consist both of about 350 numbers of well dried plants with ample notes on the colour of flowers and on the way the inhabitants of St. Eustatius make use of these plants.

In my paper: *Lijst van Planten, die door de Bewoners van de drie Nederlandsche Antillen St. Eustatius, Saba en St. Martin als geneeskrachtig worden beschouwd*. Bull. Kolon. Mus. Haarlem n. 38 (1907) p. 93—112, these notes have been of great use to me.

As might be expected the greater part of the plants in these collections are common tropical plants growing in the more accessible parts of the Island; there is moreover a fine collection of plants from the Quill. The examination these plants before I entered upon my voyage, has rendered it easy for me to recognise them in their natural state.

The cultivated plants sent in by *Mrs. van Grol* facilitated my paying attention also to these plants, and this was the more agreeable for me as I could thus the better comply with the wishes of the „Commissie van het F. W. van Eeden-fonds.”

Thanks to the great number of flowers and fruits occurring in the collection van Grol-Meyers it was not too difficult for me, though beginning with the study of the West Indian Flora to find out the names.

The following list contains the names of the plants collected by *Mrs. J. van Grol-Meyers*.

1. Collection J. van Grol-Meyers of the Koloniaal Museum at Haarlem.

- | | |
|-------------------------------------|---------------------------------------|
| 1 <i>Asclepias nivea</i> . | 15 <i>Sida carpinifolia</i> . |
| 2 <i>Commelina virginica</i> . | 16 <i>Amarantus tristis</i> . |
| 3 <i>Parthenium Hysterophorus</i> . | 17 <i>Argemone mexicana</i> . |
| 4 <i>Cassia occidentalis</i> . | 18 <i>Antigonon leptopus</i> . |
| 5 <i>Chenopodium ambrosioides</i> . | 19 <i>Ipomoea pentaphylla</i> . |
| 6 <i>Ageratum conyzoides</i> . | 20 <i>Triphasia trifoliata</i> . |
| 7 <i>Heliotropium parviflorum</i> . | 21 <i>Passiflora quadrangularis</i> . |
| 8 <i>Euphorbia pilulifera</i> . | 22 <i>Cassia bicapsularis</i> . |
| 9 <i>Chenopodium murale</i> . | 23 <i>Ricinus communis</i> . |
| 10 <i>Scoparia dulcis</i> . | 24 <i>Capsicum dulce</i> . |
| 11 <i>Capraria biflora</i> . | 26 <i>Abrus precatorius</i> . |
| 12 <i>Datura Metel</i> . | 28 <i>Polanisia viscosa</i> . |
| 13 <i>Datura Stramonium</i> . | 29 <i>Gynandropsis pentaphylla</i> . |
| 14 <i>Heliotropium indicum</i> . | 30 <i>Poinciana regia</i> . |

- | | |
|--------------------------------------|---|
| 31 <i>Plumbago scandens</i> . | 76 <i>Eugenia uniflora</i> . |
| 32 <i>Euphorbia heterophylla</i> . | 77 <i>Linociera compacta</i> . |
| 33 <i>Salvia serotina</i> . | 78 <i>Piper reticulatum</i> . |
| 34 <i>Phyllanthus niruri</i> . | 79 <i>Passiflora laurifolia</i> . |
| 35 <i>Momordica Charantia</i> . | 80 <i>Byrsonima spicata</i> . |
| 36 <i>Acacia macracantha</i> . | 81 <i>Cassia glandulosa</i> . |
| 37 <i>Leucaena glauca</i> . | 82 <i>Byrsonima spicata</i> . |
| 38 <i>Terminalia Catappa</i> . | 83 <i>Ruellia tuberosa</i> . |
| 39 <i>Acacia macracantha</i> . | 85 <i>Opuntia Ficus indica</i> . |
| 40 <i>Tamarindus indica</i> . | 86 <i>Melicocca bijuga</i> . |
| 41 <i>Psidium guajava</i> . | 87 <i>Citrus spec.</i> |
| 42 <i>Zephyranthes carinata</i> . | 88 <i>Anona squamosa</i> . |
| 43 <i>Calotropis procera</i> . | 89 <i>Anona muricata</i> . |
| 44 <i>Melia azedarach</i> . | 90 <i>Phyllantus distichus</i> . |
| 45 <i>Anacardium occidentale</i> . | 91 <i>Moringa oleifera</i> . |
| 46 <i>Capparis cynophallophora</i> . | 93 <i>Foeniculum vulgare</i> . |
| 47 <i>Solanum racemosum igneum</i> . | 95 <i>Carica Papaya</i> . |
| 48 <i>Plumiera rubra</i> . | 96 <i>Solanum lycopersicum</i> . |
| 49 <i>Plumiera alba</i> . | 97 <i>Solanum lycopersicum</i> . |
| 50 <i>Oncidium spec.</i> | 98 <i>Solanum esculentum</i> . |
| 51 <i>Brassavola rigida</i> . | 99 <i>Cuscuta americana</i> . |
| 52 <i>Tecoma leucoxydon</i> . | 100 <i>Clusia alba</i> . |
| 53 <i>Picrasma antillana</i> . | 101 <i>Coffea arabica</i> . |
| 54 <i>Croton flavens</i> . | 102 <i>Theobroma Cacao</i> . |
| 55 <i>Eugenia axillaris</i> . | 103 <i>Eugenia Jambos</i> . |
| 56 <i>Lantana involucrata</i> . | 104 <i>Centrosema virginianum</i> . |
| 57 <i>Mangifera indica</i> . | 105 <i>Cecropia peltata</i> . |
| 58 <i>Punica Granatum</i> . | 106 <i>Bursera simaruba</i> . |
| 59 <i>Achras sapota</i> . | 107 <i>Citharexylum quadrangu-</i>
<i>lare</i> . |
| 60 <i>Lochnera rosea</i> . | 108 <i>Artocarpus incisa</i> . |
| 61 <i>Pluchea odorata</i> . | 109 <i>Eugenia ligustrina</i> . |
| 63 <i>Malpighia puniceifolia</i> . | 110 <i>Lantana Camara</i> . |
| 66 <i>Vigna repens</i> . | 111 <i>Comocladia ilicifolia</i> . |
| 67 <i>Caesalpinia pulcherrima</i> . | 112 <i>Persea americana</i> . |
| 68 <i>Caesalpinia pulcherrima</i> . | 114 <i>Myrcia paniculata</i> . |
| 69 <i>Caesalpinia ciliata</i> . | 116 <i>Nectandra coriacea</i> . |
| 70 <i>Mammea americana</i> . | 117 <i>Heliconia Bihai</i> . |
| 71 <i>Spondias lutea</i> . | 118 <i>Tillandsia utriculata</i> . |
| 72 <i>Spondias purpurea</i> . | 119 <i>Canavalia obtusifolia</i> . |
| 73 <i>Chrysobalanus icaco</i> . | 120 <i>Coccoloba uvifera</i> . |
| 74 <i>Justicia sessilis</i> . | 121 <i>Costus speciosus</i> . |
| 75 <i>Hymenaea Courbaril</i> . | |

- | | |
|--|--------------------------------------|
| 123 <i>Aristolochia odoratissima</i> . | 168 <i>Jatropha curcas</i> . |
| 124 <i>Verbena chamaedrifolia</i> . | 169 <i>Hibiscus radiatus</i> . |
| 125 <i>Lawsonia inermis</i> . | 170 <i>Nymphaea spec.</i> |
| 126 <i>Lawsonia inermis</i> . | 171 <i>Jatropha multifida</i> . |
| 127 <i>Brunfelsia americana</i> . | 172 <i>Leonotis nepetaefolia</i> . |
| 128 <i>Amarantus paniculatus</i> . | 173 <i>Erigeron canadensis</i> . |
| 129 <i>Cissus sicyoides</i> . | 174 <i>Stylosanthes hamata</i> . |
| 130 <i>Urera caracasana</i> . | 175 <i>Hymenaea Courbaril</i> . |
| 131 <i>Codiaeum variegatum</i> . | 175a <i>Boerhaavea hirsuta</i> . |
| 132 <i>Phaseolus lunatus</i> . | 176 <i>Achyranthes obtusifolia</i> . |
| 133 <i>Phaseolus vulgaris</i> . | 177 <i>Croton lobatus</i> . |
| 135 <i>Antigonon cinerascens</i> . | 178 <i>Stenolobium stans</i> . |
| 137 <i>Aegiphila martinicensis</i> . | 179 <i>Stylosanthes hamata</i> . |
| 138 <i>Althaea officinalis</i> . | 181 <i>Waltheria americana</i> . |
| 139 <i>Duranta repens</i> . | 182 <i>Sida cordifolia</i> . |
| 140 <i>Bixa Orellana</i> . | 183 <i>Thunbergia fragans</i> . |
| 141 <i>Brassica integrifolia</i> . | 184 <i>Urechites suberecta</i> . |
| 142 <i>Hura crepitans</i> . | 185 <i>Triumfetta rhomboidea</i> . |
| 143 <i>Jasminum spec.</i> | 188 <i>Passiflora suberosa</i> . |
| 144 <i>Stigmatophyllum periploci-</i>
folium. | 189 <i>Neurogramme calomelanos</i> . |
| 145 <i>Cordia Sebestena</i> . | 190 <i>Solanum aculeatissimum</i> . |
| 146 <i>Fourcroya gigantea</i> . | 191 <i>Lonicera japonica</i> . |
| 147 <i>Clerodendron aculeatum</i> . | 192 <i>Clerodendron fallax</i> . |
| 149 <i>Plumbago capensis</i> . | 193 <i>Sesbania grandiflora</i> . |
| 150 <i>Parkinsonia aculeata</i> . | 194 <i>Leonurus sibiricus</i> . |
| 151 <i>Quisqualis indica</i> . | 195 <i>Randia aculeata</i> . |
| 152 <i>Ipomoea pes caprae</i> . | 196 <i>Jatropha gossypifolia</i> . |
| 153 <i>Mirabilis Jalapa</i> . | 197 <i>Jatropha gossypifolia</i> . |
| 154 <i>Mirabilis Jalapa</i> . | 198 <i>Vernonia punctata</i> . |
| 155 <i>Chrysobalanus icaco</i> . | 200 <i>Krameria ixina</i> . |
| 156 <i>Picrasma antillana</i> . | 201 <i>Abutilon indicum</i> . |
| 157 <i>Indigofera suffruticosa</i> . | 203 <i>Rauwolfia Lamareckiana</i> . |
| 158 <i>Coleus scutellarioides</i> . | 204 <i>Galactia rubra</i> . |
| 159 <i>Cosmos sulphureus</i> . | 210 <i>Cajanus indicus</i> . |
| 160 <i>Sambucus canadensis</i> . | 211 <i>Rivina humilis</i> . |
| 161 <i>Hibiscus Rosa-sinensis</i> . | 212 <i>Aegiphila martinicensis</i> . |
| 162 <i>Nopalea coccinellifera</i> . | 213 <i>Psychotria rufescens</i> . |
| 163 <i>Opuntia Tuna</i> . | 214 <i>Andropogon Schoenanthus</i> . |
| 164 <i>Gomphrena globosa</i> . | 215 <i>Aristida americana</i> . |
| 165 <i>Codiaeum variegatum</i> . | 216 <i>Mimosa pudica</i> . |
| 167 <i>Codiaeum variegatum</i> . | 217 <i>Tillandsia recurvata</i> . |
| | 218 <i>Cissampelos Pareira</i> . |

- | | |
|--|--|
| 220 <i>Ipomoea fastigiata</i> . | 265 <i>Cenchrus tribuloides</i> . |
| 221 <i>Porana paniculata</i> . | 266 <i>Cyperus elegans</i> . |
| 222 <i>Jacquemontia pentantha</i> . | 267 <i>Tournefortia gnaphalodes</i> . |
| 223 <i>Boerhaavea erecta</i> . | 268 <i>Epidendrum ciliare</i> . |
| 224 <i>Operculina ventricosa</i> . | 269 <i>Scleria lithosperma</i> . |
| 225 <i>Ibatia maritima</i> . | 271 <i>Chiococca racemosa</i> . |
| 227 <i>Solanum lycopersicum</i> . | 272 <i>Allium ascalonicum</i> . |
| 228 <i>Bouchea Ehrenbergii</i> . | 273 <i>Pilea semidentata</i> . |
| 229 <i>Euphorbia thymifolia</i> . | 274 <i>Anona montana</i> . |
| 230 <i>Ambrosia hispida</i> . | 275 <i>Sida glomerata</i> . |
| 231 <i>Russelia juncea</i> . | 277 <i>Porophyllum ellipticum</i> . |
| 232 <i>Melochia tomentosa</i> . | 278 <i>Allamanda cathartica</i> . |
| 233 <i>Wedelia bupththalmoides</i> . | 280 <i>Calonyction speciosum</i> . |
| 234 <i>Panicum divaricatum</i> . | 281 <i>Ipomoea pentaphylla</i> . |
| 235 <i>Clitoria ternatea</i> . | 282 <i>Maranta arundinacea</i> . |
| 237 <i>Eleusine indica</i> . | 283 <i>Cucurbita maxima</i> . |
| 238 <i>Petiveria alliacea</i> . | 284 <i>Pithecolobium unguis-cati</i> . |
| 239 <i>Allophylus occidentalis</i> . | 285 <i>Crotalaria verrucosa</i> . |
| 240 <i>Tephrosia cathartica</i> . | 286 <i>Crotalaria retusa</i> . |
| 241 <i>Hyptis pectinata</i> . | 287 <i>Bryophyllum pinnatum</i> . |
| 242 <i>Synedrella nodiflora</i> . | 288 <i>Ricinus communis</i> . |
| 243 <i>Medicago sativa</i> . | 289 <i>Ricinus communis</i> . |
| 244 <i>Gossypium barbadense</i> . | 290 <i>Manihot utilisima</i> . |
| 245 <i>Dioscorea alata</i> . | 291 <i>Haematoxylon campechi-</i>
<i>anum</i> . |
| 248 <i>Beloperone eustachyana</i> . | 292 <i>Salvia occidentalis</i> . |
| 249 <i>Desmodium triflorum</i> . | 293 <i>Stachytarpheta jamaicensis</i> . |
| 250 <i>Alysicarpus nummularifo-</i>
<i>lius</i> . | 294 <i>Cardiospermum halicacabum</i> . |
| 251 <i>Hibiscus esculentus</i> . | 295 <i>Aloe vera</i> . |
| 252 <i>Hibiscus esculentus</i> . | 296 <i>Ocimum micranthum</i> . |
| 253 <i>Canna indica</i> . | 297 <i>Bryophyllum pinnatum</i> . |
| 254 <i>Canna indica</i> . | 298 <i>Hymenocallis caribaea</i> . |
| 255 <i>Crescentia Cujete</i> . | 299 <i>Eucharis grandiflora</i> . |
| 256 <i>Crotalaria incana</i> . | 300 <i>Sansevieria guineensis</i> . |
| 257 <i>Polypodium phyllitidis</i> . | 301 <i>Pilea microphylla</i> . |
| 258 <i>Dryopteris parasitica</i> . | 302 <i>Pilea microphylla</i> . |
| 259 <i>Bignonia unguis-cati</i> . | 304 <i>Blechum Brownei</i> . |
| 260 <i>Oplismenus setarius</i> . | 305 <i>Jasminum pubescens</i> . |
| 261 <i>Polypodium lycopodioides</i> . | 306 <i>Panax fruticosum</i> . |
| 262 <i>Mariscus brizaeus</i> . | 307 <i>Panax fruticosum</i> . |
| 263 <i>Smilax coriacea</i> . | 308 <i>Codiaeum variegatum</i> . |
| 264 <i>Xanthosoma spec.</i> | 309 <i>Codiaeum variegatum</i> . |

310 <i>Codiaeum variegatum</i> .	324 <i>Hippomane mancinella</i> .
311 <i>Codiaeum variegatum</i> .	325 <i>Musa paradisiaca</i> .
312 <i>Gaillardia aristata</i> .	326 <i>Verbena chamaedrifolia</i> .
314 <i>Piscidia piscipula</i> Sarg.	329 <i>Desmodium supinum</i> .
316 <i>Angelonia angustifolia</i> .	330 <i>Lochnera rosea</i> .
317 <i>Hibiscus schizopetalus</i> .	331 <i>Panax fruticosum</i> .
318 <i>Bromelia pinguin</i> .	332 <i>Comocladia ilicifolia</i> .
321 <i>Ceiba pentandra</i> .	334 <i>Musa paradisiaca</i> .
322 <i>Pisonia subcordata</i> .	336 <i>Nerium Oleander</i> .
323 <i>Villamilla octandra</i> .	339 <i>Portulaca halimoides</i> .

**Collection J. van Grol-Meyers of the Herbarium of the
Botanical Institute of the University in Utrecht.**

1 <i>Vigna repens</i> .	29 <i>Portulaca oleracea</i> .
2 <i>Commelina virginica</i> .	30 <i>Kallstroemia maxima</i> .
3 <i>Parthenium Hysterophorus</i> .	31 <i>Jasminum azoricum</i> .
4 <i>Cassia occidentalis</i> .	32 <i>Polanisia viscosa</i> .
5 <i>Chenopodium ambrosioides</i> .	33 <i>Gynandropsis pentaphylla</i> .
6 <i>Ageratum conyzoides</i> .	34 <i>Poinciana regia</i> .
7 <i>Heliotropium parviflorum</i> .	35 <i>Plumbago scandens</i> .
8 <i>Euphorbia pilulifera</i> .	36 <i>Euphorbia heterophylla</i> .
9 <i>Chenopodium murale</i> .	37 <i>Salvia serotina</i> .
10 <i>Scoparia dulcis</i> .	38 <i>Priva echinata</i> .
11 <i>Capraria biflora</i> .	39 <i>Melochia tomentosa</i> .
12 <i>Datura Stramonium</i> .	40 <i>Murraya exotica</i> .
13 <i>Datura Metel</i> .	41 <i>Phyllanthus niruri</i> .
14 <i>Datura Stramonium</i> .	42 <i>Momordica Charantia</i> .
15 <i>Heliotropium indicum</i> .	43 <i>Microtea debilis</i> .
16 <i>Sida carpinifolia</i> .	44 <i>Leucaena glauca</i> .
17 <i>Amarantus tristis</i> .	45 <i>Terminalia Catappa</i> .
18 <i>Amarantus spinosus</i> .	46 <i>Solanum racemosum igneum</i> .
19 <i>Amarantus polygonoides</i> .	47 <i>Tamarindus indica</i> .
20 <i>Argemone mexicana</i> .	48 <i>Psidium Guajava</i> .
21 <i>Antigonon leptopus</i> .	49 <i>Allamanda cathartica</i> .
22 <i>Lochnera rosea</i> .	50 <i>Acacia macracantha</i> .
23 <i>Capsicum dulce</i> .	51 <i>Capsicum frutescens</i> .
24 <i>Passiflora quadrangularis</i> .	52 <i>Jatropha curcas</i> .
25 <i>Cassia bicapsularis</i> .	53 <i>Cordia speciosa</i> .
26 <i>Ricinus communis</i> .	54 <i>Calotropis procera</i> .
27 <i>Ricinus communis</i> .	55 <i>Cajanus indicus</i> .
28 <i>Portulaca oleracea</i> .	56 <i>Clerodendron fallax</i> .

- | | |
|---------------------------------------|--|
| 57 <i>Brunfelsia americana</i> . | 101 <i>Fourcroya gigantea</i> . |
| 58 <i>Ixora Jucunda</i> . | 102 <i>Aloe vera</i> . |
| 59 <i>Solanum torvum</i> . | 103 <i>Bromelia pinguin</i> . |
| 61 <i>Lagerstroemia indica</i> . | 104 <i>Phoenix dactylifera</i> . |
| 62 <i>Canna indica</i> . | 105 <i>Amarantus paniculatus</i> . |
| 63 <i>Talinum triangulare</i> . | 106 <i>Achras sapota</i> . |
| 64 <i>Plumbago capensis</i> . | 107 <i>Codiaeum variegatum</i> . |
| 65 <i>Quisqualis indica</i> . | 108 <i>Nerium Oleander</i> . |
| 66 <i>Jatropha multifida</i> . | 109 <i>Tecoma leucoxydon</i> . |
| 67 <i>Clusia alba</i> . | 110 <i>Artocarpus incisa</i> . |
| 68 <i>Plumiera rubra</i> . | 112 <i>Russelia juncea</i> . |
| 69 <i>Peperomia petiolaris</i> . | 113 <i>Hibiscus Rosa-sinensis</i> . |
| 70 <i>Peireskia aculeata</i> . | 114 <i>Hippeastrum puniceum</i> . |
| 71 <i>Eleutherine plicata</i> . | 115 <i>Mariscus brizaeus</i> . |
| 72 <i>Verbena chamaedrifolia</i> . | 116 <i>Pluchea odorata</i> . |
| 73 <i>Parkinsonia aculeata</i> . | 117 <i>Coleus scutellarioides</i> . |
| 74 <i>Caesalpinia pulcherrima</i> . | 119 <i>Angelonia angustifolia</i> . |
| 75 <i>Caesalpinia pulcherrima</i> . | 120 <i>Croton flavens</i> . |
| 76 <i>Ixora coccinea</i> . | 121 <i>Althaea officinalis</i> . |
| 77 <i>Punica Granatum</i> . | 122 <i>Foeniculum vulgare</i> . |
| 78 <i>Malpighia puniceifolia</i> . | 123 <i>Begonia spec.</i> |
| 79 <i>Sambucus canadensis</i> . | 124 <i>Codiaeum variegatum</i> . |
| 81 <i>Panax fruticosum</i> . | 125 <i>Codiaeum variegatum</i> . |
| 82 <i>Panax fruticosum</i> . | 126 <i>Senecio sonchifolius</i> . |
| 83 <i>Panax fruticosum</i> . | 127 <i>Cassia fistula</i> . |
| 84 <i>Citrus spec.</i> | 128 <i>Citharexylum quadrangulare</i> . |
| 85 <i>Tabernaemontana coronaria</i> . | 129 <i>Ruellia tuberosa</i> . |
| 86 <i>Boerhaavia erecta</i> . | 130 <i>Mangifera indica</i> . |
| 87 <i>Boerhaavia hirsuta</i> . | 131 <i>Ocimum micranthum</i> . |
| 88 <i>Lantana Camara</i> . | 132 <i>Jatropha gossypifolia</i> . |
| 89 <i>Lantana involucrata</i> . | 133 <i>Jatropha gossypifolia</i> . |
| 90 <i>Trianthema portulacastrum</i> . | 134 <i>Asclepias nivea</i> . |
| 91 <i>Anona squamosa</i> . | 135 <i>Fleurya aestuans</i> . |
| 92 <i>Croton lobatus</i> . | 136 <i>Leonurus sibiricus</i> . |
| 93 <i>Achyranthes obtusifolia</i> . | 137 <i>Stigmatophyllum periplocifolium</i> . |
| 94 <i>Lonicera japonica</i> . | 138 <i>Leonotis nepetaefolia</i> . |
| 95 <i>Clerodendron aculeatum</i> . | 139 <i>Melia azedarach</i> . |
| 96 <i>Bauhinia Kappleri</i> . | 140 <i>Neurogramme calomelanos</i> . |
| 97 <i>Jasminum undulatum</i> . | 141 <i>Justicia sessilis</i> . |
| 98 <i>Gaillardia aristata</i> . | 142 <i>Nicotiana Tabacum</i> . |
| 99 <i>Cassia glandulosa</i> . | 143 <i>Solanum lycopersicum</i> . |
| 100 <i>Stenolobium stans</i> . | |

- | | |
|---------------------------------------|---|
| 144 <i>Passiflora laurifolia</i> . | 186 <i>Cuscuta americana</i> . |
| 145 <i>Erigeron canadensis</i> . | 187 <i>Chiococca racemosa</i> . |
| 146 <i>Persea americana</i> . | 188 <i>Smilax coriacea</i> . |
| 147 <i>Hura crepitans</i> . | 189 <i>Capparis frondosa</i> . |
| 148 <i>Phyllanthus distichus</i> . | 190 <i>Capparis cynophallophora</i> . |
| 149 <i>Crescentia Cujete</i> . | 191 <i>Capparis cynophallophora</i> . |
| 150 <i>Pilea microphylla</i> . | 192 <i>Pilea semidentata</i> . |
| 151 <i>Opuntia Ficus indica</i> . | 193 <i>Peperomia glabella</i> . |
| 152 <i>Opuntia Tuna</i> . | 194 <i>Panicum divaricatum</i> . |
| 154 <i>Cardiospermum halicabum</i> . | 195 <i>Anthurium cordatum</i> . |
| 155 <i>Cissus sicyoides</i> . | 196 <i>Casearia parvifolia</i> . |
| 156 <i>Cucumis Anguria</i> . | 198 <i>Coccoloba uvifera</i> . |
| 157 <i>Ambrosia hispida</i> . | 199 <i>Carica Papaya</i> . |
| 158 <i>Tephrosia cathartica</i> . | 200 <i>Bursera simaruba</i> . |
| 159 <i>Cyperus rotundus</i> . | 201 <i>Mussaenda frondosa</i> . |
| 160 <i>Caesalpinia crista</i> . | 202 <i>Eucharis grandiflora</i> . |
| 161 <i>Rivina humilis</i> . | 203 <i>Codiaeum variegatum</i> . |
| 162 <i>Desmanthus virgatus</i> . | 204 <i>Codiaeum variegatum</i> . |
| 163 <i>Medicago sativa</i> . | 205 <i>Carica Papaya</i> . |
| 164 <i>Ipomoea dissecta</i> . | 206 <i>Allium ascalonicum</i> . |
| 165 <i>Wedelia buphthalmoides</i> . | 207 <i>Anacardium occidentale</i> . |
| 166 <i>Beloperone eustachyana</i> . | 209 <i>Phoradendron trinervium</i> . |
| 167 <i>Maranta arundinacea</i> . | 210a <i>Piscidia piscipula</i> . |
| 168 <i>Nectandra coriacea</i> . | 213 <i>Myrcia splendens</i> . |
| 169 <i>Linociera compacta</i> . | 214 <i>Theobroma Cacao</i> . |
| 170 <i>Cassia glandulosa</i> . | 215 <i>Caesalpinia coriaria</i> . |
| 171 <i>Croton astroites</i> . | 216 <i>Hymenaea Courbaril</i> . |
| 172 <i>Indigofera suffruticosa</i> . | 217 <i>Quararibea turbinata</i> . |
| 173 <i>Anona muricata</i> . | 218 <i>Tabernaemontana citrifolia</i> . |
| 174 <i>Piper dilatatum</i> . | 218a <i>Psychotria horizontalis</i> . |
| 175 <i>Psychotria rufescens</i> . | 219 <i>Indigofera suffruticosa</i> . |
| 176 <i>Randia aculeata</i> . | 220 <i>Philodendron dispar</i> . |
| 177 <i>Myrcia splendens</i> . | 221 <i>Theobroma Cacao</i> . |
| 178 <i>Bignonia ungius-cati</i> . | 222 <i>Urera caracasana</i> . |
| 179 <i>Polypodium lycopodioides</i> . | 223 <i>Boerhaavea erecta</i> . |
| 180 <i>Pisonia fragans</i> . | 223a <i>Heliconia Bihai</i> . |
| 181 <i>Pisonia subcordata</i> . | 224 <i>Chrysobalanus icaco</i> . |
| 182 <i>Sida cordifolia</i> . | 225 <i>Pharus glaber</i> . |
| 183 <i>Polypodium phyllitidis</i> . | 226 <i>Triphasia trifoliata</i> . |
| 184 <i>Mammea americana</i> . | 227 <i>Moringa oleifera</i> . |
| 185 <i>Clusia alba</i> . | 228 <i>Gomphrena globosa</i> . |
| | 229 <i>Hibiscus radiatus</i> . |

- | | |
|---|---|
| 230 <i>Solanum lycopersicum</i> . | 270 <i>Ipomoea fastigiata</i> . |
| 231 <i>Bixa Orellana</i> . | 271 <i>Passiflora suberosa</i> . |
| 232 <i>Boussingaultia baselloides</i> . | 272 <i>Vernonia punctata</i> . |
| 233 <i>Ocimum micranthum</i> . | 274 <i>Oplismenus setarius</i> . |
| 234 <i>Lawsonia inermis</i> . | 276 <i>Dryopteris parasitica</i> . |
| 235 <i>Abrus precatorius</i> . | 277 <i>Ibatia maritima</i> . |
| 236 <i>Centrosema virginianum</i> . | 278 <i>Setaria setosa</i> . |
| 237 <i>Rauwolfia Lamarkiana</i> . | 279 <i>Oplismenus setarius</i> . |
| 239 <i>Mirabilis Jalapa</i> . | 280 <i>Epidendrum papilionaceum</i> . |
| 240 <i>Mirabilis Jalapa</i> . | 281 <i>Epidendrum ciliare</i> . |
| 241 <i>Codiaeum variegatum</i> . | 282 <i>Galactia rubra</i> . |
| 243 <i>Cosmos sulphureus</i> . | 283 <i>Capsicum dulce</i> . |
| 244 <i>Canavalia spec.</i> | 284 <i>Porophyllum ellipticum</i> . |
| 245 <i>Caesalpinia coriacea</i> . | 285 <i>Crotalaria retusa</i> . |
| 246 <i>Chrysobalanus icaco</i> . | 286 <i>Citrus spec.</i> |
| 247 <i>Mirabilis Jalapa</i> . | 287 <i>Andropogon zizanioides</i> . |
| 248 <i>Antigonon cinerascens</i> . | 289 <i>Andropogon Schoenanthus</i> . |
| 249 <i>Cenchrus tribuloides</i> . | 290 <i>Sesbania grandiflora</i> . |
| 251 <i>Gossypium barbadense</i> . | 291 <i>Sesbania grandiflora</i> . |
| 252 <i>Stylosanthes hamata</i> . | 292 <i>Brassica integrifolia</i> . |
| 253 <i>Operculina ventricosa</i> . | 294 <i>Lawsonia inermis</i> . |
| 254 <i>Melicocca bijuga</i> . | 295 <i>Tillandsia usneoides</i> . |
| 255 <i>Euphorbia thymifolia</i> . | 296 <i>Celosia argentea</i> . |
| 256 <i>Malvastrum spicatum</i> . | 297 <i>Porana paniculata</i> . |
| 258 <i>Ricinus communis</i> . | 298 <i>Hymenocallis caribaea</i> . |
| 259 <i>Cassia alata</i> . | 299 <i>Bouchea Ehrenbergii</i> . |
| 260 <i>Petiveria alliacea</i> . | 300 <i>Stylosanthes hamata</i> . |
| 261 <i>Ocimum Basilicum</i> . | 301 <i>Spermacoce tenuior</i> . |
| 263 <i>Solanum aculeatissimum</i> . | 302 <i>Desmanthus virgatus</i> . |
| 264 <i>Eleusine indica</i> . | 303 <i>Crotalaria verrucosa</i> . |
| 265 <i>Ipomoea fastigiata</i> . | 304 <i>Acacia macracantha</i> . |
| 266 <i>Ipomoea fastigiata</i> . | 305 <i>Stachytarpheta jamaicensis</i> . |
| 267 <i>Cucurbita maxima</i> . | 306 <i>Musa paradisiaca</i> . |
| 268 <i>Achyranthes obtusifolia</i> . | 307 <i>Phaseolus lunatus</i> . |
| 269 <i>Corchorus olitorius</i> . | 308 <i>Epidendrum ciliare</i> . |

Collection Lionarons.

A. C. W. Lionarons, gov. physician of Saba, made in 1907 a collection of about 350 plants and gave it to the Koloniaal Museum at Haarlem. *Dr. M. Greshoff*, Director of this Museum was so kind as to give me an opportunity to study these plants.

The greater part of the plants are cultivated and another part has evidently been collected on the top of the Mountain.

There are some interesting plants, which I myself saw but once during my rambles through Saba.

The following list gives the names of the plants collected by *Lionarons*.

- | | |
|------------------------------------|--------------------------------------|
| 1 <i>Anona muricata</i> . | 49 <i>Polypodium lycopodioides</i> . |
| 1a <i>Melicocca bijuga</i> . | 50 <i>Oxalis corniculata</i> . |
| 2 <i>Passiflora laurifolia</i> . | 53 <i>Oplismenus setarius</i> . |
| 4 <i>Jasminum azoricum</i> . | 55 <i>Canna indica</i> . |
| 7 <i>Capsicum frutescens</i> . | 56 <i>Parthenium Hysterophorus</i> . |
| 8 <i>Capsicum frutescens</i> . | 57 <i>Sambucus canadensis</i> . |
| 9 <i>Capsicum ceratocarpum</i> . | 61 <i>Polypodium crassifolium</i> . |
| 12 <i>Swietenia Mahagoni</i> . | 63 <i>Carica Papaya</i> . |
| 15 <i>Commelina virginica</i> . | 64 <i>Ricinus communis</i> . |
| 16 <i>Cajanus indicus</i> . | 65 <i>Crescentia Cujete</i> . |
| 17 <i>Thunbergia fragans</i> . | 66 <i>Crotalaria retusa</i> . |
| 18 <i>Chrysobalanus icaco</i> . | 67 <i>Manihot utilissima</i> . |
| 21 <i>Tillandsia fasciculata</i> . | 68 <i>Ipomoea fastigiata</i> . |
| 22 <i>Carica Papaya</i> . | 69 <i>Centrosema virginianum</i> . |
| 23 <i>Eleusine indica</i> . | 72 <i>Petiveria alliacea</i> . |
| 24 <i>Beloperone eustachyana</i> . | 74 <i>Lochnera rosea</i> . |
| 25 <i>Desmodium supinum</i> . | 75 <i>Abutilon indicum</i> . |
| 26 <i>Brassica integrifolia</i> . | 76 <i>Ardisia coriacea</i> . |
| 28 <i>Cassia bicapsularis</i> . | 78 <i>Allium ascalonicum</i> . |
| 29 <i>Wedelia buphthalmoides</i> . | 79 <i>Allium ascalonicum</i> . |
| 30 <i>Ipomoea pentaphylla</i> . | 80 <i>Portulaca oleracea</i> . |
| 31 <i>Boerhaavea paniculata</i> . | 82 <i>Operculina ventricosa</i> . |
| 32 <i>Rivina humilis</i> . | 85 <i>Plumbago capensis</i> . |
| 33 <i>Amarantus tristis</i> . | 87 <i>Indigofera suffruticosa</i> . |
| 34 <i>Pharus glaber</i> . | 88 <i>Aleurites moluccana</i> . |
| 35 <i>Salvia occidentals</i> . | 90 <i>Lochnera rosea</i> . |
| 36 <i>Thunbergia alata</i> . | 91 <i>Tephrosia cathartica</i> . |
| 37 <i>Eryngium foetidum</i> . | 92 <i>Zinnia elegans</i> . |
| 39 <i>Peperomia petiolaris</i> . | 93 <i>Jasminum azoricum</i> . |
| 40 <i>Sonchus oleraceus</i> . | 95 <i>Parthenium Hysterophorus</i> . |
| 42 <i>Zebrina pendula</i> . | 96 <i>Nicotiana Tabacum</i> . |
| 44 <i>Foeniculum vulgare</i> . | 97 <i>Triphasia trifoliata</i> . |
| 46 <i>Antigonon leptopus</i> . | 99 <i>Punica Granatum</i> . |
| 47 <i>Gossypium barbadense</i> . | 101 <i>Coccoloba nivea</i> . |
| 48 <i>Capparis jamaicensis</i> . | 103 <i>Peperomia acuminata</i> . |

- | | |
|---------------------------------------|---|
| 104 <i>Cissus sicyoides</i> . | 162 <i>Cassia occidentalis</i> . |
| 106 <i>Cuscuta americana</i> . | 163 <i>Momordica Charantia</i> . |
| 109 <i>Eugenia Jambos</i> . | 164 <i>Foeniculum vulgare</i> . |
| 110 <i>Quisqualis indica</i> . | 165 <i>Epidendrum ciliare</i> . |
| 111 <i>Hibiscus Rosa-sinensis</i> . | 168 <i>Belamcanda punctata</i> . |
| 112 <i>Mirabilis Jalapa</i> . | 169 <i>Phytolacca icosandra</i> . |
| 113 <i>Calotropis procera</i> . | 170 <i>Passiflora laurifolia</i> . |
| 114 <i>Mammea americana</i> . | 171 <i>Peperomia acuminata</i> . |
| 117 <i>Tamarindus indica</i> . | 172 <i>Nephrolepis exaltata</i> . |
| 118 <i>Manihot utilisima</i> . | 174 <i>Peperomia hernandifolia</i> . |
| 119 <i>Psidium guajava</i> . | 175 <i>Phytolacca icosandra</i> . |
| 123 <i>Lepidium virginicum</i> . | 176 <i>Physurus plantagineus</i> . |
| 124 <i>Talinum paniculatum</i> . | 177 <i>Casuarina equisetifolia</i> . |
| 125 <i>Melia azedarach</i> . | 178 <i>Plantago major</i> . |
| 127 <i>Hillia longiflora</i> . | 181 <i>Stenochlaena sorbifolia</i> . |
| 128 <i>Euphorbia hypericifolia</i> . | 182 <i>Citharexylum quadrangulare</i> . |
| 129 <i>Leucaena glauca</i> . | 183 <i>Ardisia coriacea</i> . |
| 132 <i>Scleria pterota</i> . | 184 <i>Scleria pterota</i> . |
| 133 <i>Setaria setosa</i> . | 186 <i>Leucaena glauca</i> . |
| 134 <i>Stenolobium stans</i> . | 187 <i>Chorizanthe crinitus</i> . |
| 135 <i>Ageratum conyzoides</i> . | 188 <i>Hemitelia horrida</i> . |
| 136 <i>Leucaena glauca</i> . | 189 <i>Hemitelia horrida</i> . |
| 137 <i>Chiococca racemosa</i> . | 192 <i>Cecropia peltata</i> . |
| 140 <i>Myrcia paniculata</i> . | 194 <i>Urera caracasana</i> . |
| 142 <i>Pilea microphylla</i> . | 196 <i>Neurogramme calomelanos</i> . |
| 143 <i>Hibiscus schizopetalus</i> . | 196a <i>Mangifera indica</i> . |
| 145 <i>Solanum lycopersicum</i> . | 197 <i>Chorizanthe crinitus</i> . |
| 146 <i>Iresine paniculata</i> . | 201 <i>Desmodium axillare</i> . |
| 147 <i>Panicum brevifolium</i> . | 202 <i>Polybotrya cervina</i> . |
| 147b <i>Panicum fuscum</i> . | 203 <i>Chorizanthe crinitus</i> . |
| 148 <i>Pisonia aculeata</i> . | 204 <i>Polybotrya cervina</i> . |
| 149 <i>Persea americana</i> . | 206 <i>Ichnanthus pallens</i> . |
| 151 <i>Tecoma capensis</i> . | 208 <i>Cladium restioides</i> . |
| 153 <i>Gynandropsis pentaphylla</i> . | 209 <i>Phytolacca icosandra</i> . |
| 154 <i>Fleurya aestuans</i> . | 210 <i>Neurolaena lobata</i> . |
| 155 <i>Argemone mexicana</i> . | 211 <i>Asplenium serratum</i> . |
| 156 <i>Mangifera indica</i> . | 212 <i>Aspidium Plumierii</i> . |
| 158 <i>Maranta arundinacea</i> . | 214 <i>Selaginella flabellata</i> . |
| 159 <i>Piper reticulatum</i> . | 216 <i>Miconia laevigata</i> . |
| 159a <i>Piper marginatum</i> . | 218 <i>Ipomoea fastigiata</i> . |
| 160 <i>Asclepias nivea</i> . | 220 <i>Dryopteris opposita</i> . |
| 161 <i>Euphorbia hypericifolia</i> . | 221 <i>Colocasia esculenta</i> . |

- | | |
|--------------------------------|---------------------------------|
| 222 Nerium Oleander. | 287 Anthurium cordatum. |
| 223 Caesalpinia pulcherrima. | 288 Bryophyllum pinnatum. |
| 224 Codiaeum variegatum. | 290 Bursera simaruba. |
| 225 Codiaeum variegatum. | 291 Solanum tuberosum. |
| 227 Anacardium occidentale. | 292 Bixa Orellana. |
| 229 Quisqualis indica. | 293 Wittmackia lingulata. |
| 230 Russelia juncea. | 294 Panicum insulare. |
| 231 Nicotiana Tabacum. | 297 Ruellia tuberosa. |
| 232 Gossypium barbadense. | 298 Thrinax spec. |
| 233 Thespesia populnea. | 299 Antigonon leptopus. |
| 234 Pluchea odorata. | 301 Caesalpinia coriaria. |
| 236 Villamilla octandra. | 302 Mangifera indica. |
| 237 Cassia occidentalis. | 303 Panicum divaricatum. |
| 238 Phyllanthus distichus. | 303a Colocasia esculenta. |
| 239 Euphorbia pilulifera. | 306 Miconia laevigata. |
| 240 Panicum maximum. | 309 Chiococca racemosa. |
| 241 Jatropha gossypifolia. | 311 Callisia repens. |
| 245 Bixa Orellana. | 312 Abrus precatorius. |
| 247 Leonotis nepetaefolia. | 313 Jatropha gossypifolia. |
| 251 Solanum nigrum nodiflorum. | 315 Croton flavens. |
| 251a Indigofera suffruticosa. | 316 Blechnum Brownei. |
| 252 Amarantus spinosus. | 320 Salvia serotina. |
| 253 Lantana camara. | 321 Eugenia Jambos. |
| 254 Achyranthes obtusifolia. | 322 Colocasia esculenta. |
| 256 Talinum triangulare. | 324 Colocasia esculenta. |
| 257 Jatropha multifida. | 325 Colocasia esculenta. |
| 260 Datura fastuosa. | 328 Petiveria alliacea. |
| 262 Desmodium spirale. | 330 Codiaeum variegatum. |
| 265 Lawsonia inermis. | 330a Lochnera rosea. |
| 269 Mussaenda frondosa. | 335 Chrysobalanus icaco. |
| 269a Porophyllum ellipticum. | 336 Citharexylum quadrangulare. |
| 270 Rauwolfia Lamarekii. | 337 Casearia parvifolia. |
| 273 Panicum sanguinale. | 338 Aleurites moluccana. |
| 276 Cucumis Anguria. | 339 Codiaeum variegatum. |
| 278 Quisqualis indica. | 340 Coccoloba diversifolia. |
| 279 Galphimia gracilis. | 343 Panicum divaricatum. |
| 280 Brunfelsia americana. | 345 Codiaeum variegatum. |
| 281 Murraya exotica. | 346 Stachytarpheta jamaicensis. |
| 282 Sanchezia nobilis. | 348 Hymenaea Courbaril. |
| 284 Tecoma leucoxydon. | 354 Caesalpinia crista. |
| 285 Rhoeo discolor. | 356 Poinciana regia. |
| 286 Coleus amboinicus. | |

357 *Clusia alba*.361 *Canna indica*.359 *Albizzia lebbek*.363 *Ipomoea pentaphylla*.360 *Coccoloba uvifera*.**Collection I. Boldingh.**

I do not think it necessary to give more details about my own excursions; the different data, which are found in the Systematic Part are all taken from my notes and collections made during my rambles.

The following list may give an idea of the localities visited by me. In the Third Part Section C. I give a detailed description of the vegetation also taken from my personal observations.

St. Eustatius: May 25th—July 5th.

May 25th. Signalhill, Tumbledownickbay.

" 26th. round the Quill.

" 27th. Quill.

" 28th. Seashore near Oranjestad.

" 29th. Seashore near Oranjestad, Gallowsbay.

" 30th. Tumbledownickbay.

" 31st—June 5th. Quill.

June 6th—8th. Quill.

" 9th—10th. Cultuurvlakte.

" 12th. Bengalen—seashore.

" 13th—14th. Quill near Bengalen.

" 15th—16th. White Wall and seashore.

" 17th. Quill.

" 19th. Cultuurvlakte.

" 21st. Schotsenhoek, Concordia, Venusbay.

" 22nd. English Quarter and seashore.

" 23th. Signalhill.

" 25th. Cultuurvlakte.

" 26th. Gilboohill, Venusbay.

" 27th. Schotsenhoek, Maryglory, Bergje, Jenkinsbay.

" 28th. Quill.

" 30th. Jenkinsbay, Bergje.

July 2nd. White Wall.

" 3rdth. Zeelandia, Concordiabay.

" 5th. Bengalen.

Saba : July 6th—29th.

- July 6th. St. John's.
 „ 7th. Windwardside, Paris, Great hill.
 „ 8th. Bottom.
 „ 9th. Bottom to Marypoint.
 „ 10th. Boobyhill.
 „ 12th. Bottom, Forthbaygut, Saddle.
 „ 13th. Hellsgate, Sulphurmines.
 „ 14th. Mountain, Crespeenhill.
 „ 16th. Mountain.
 „ 17th. St. John's, Giles gut.
 „ 18th. Windwardside.
 „ 19th. Laddergut.
 „ 20th. Bottom, Windwardside, Hellsgate, Marypoint.
 „ 21st. Springbaygut.
 „ 23th. Sulphurmines, Green Island, Flat Point.
 „ 24th. Mountain, Windwardside.
 „ 25th. Mountain.
 „ 26th. Bottom.
 „ 27th. Mountain, Hellsgate.
 „ 29th. Windwardside, Bottom.

St. Martin : August 14th—September 14th.

- August 4th. Philipsburg, Marigot.
 „ 6th. Philipsburg, Guanabay.
 „ 8th. Philipsburg, Belvedere, Mont Chambord, Grande Case,
 Marigot, Philipsburg.
 „ 17th. Fort Willem.
 „ 18th. Prince quarter.
 „ 19th. Philipsburg, Cul de Sac, Marigothill, Marigot, Colebay.
 „ 20th. Bethlehem.
 „ 21st. Nakedboyhill, Fort Willem.
 „ 22nd. Simonsbay, Cherry cap, Anse des Bourgaux.
 „ 23th. Fort Willem.
 „ 24th. Bellevue.
 „ 25th. Hill between Centryhill and Marigothill.
 „ 27th. Mont Chambord, Orientbay.
 „ 28th. Hills on the north s.de of Cul de Sac.
 „ 29th. Philipsburg, Simonsbay.
 „ 30th. Oysterpond.

September 4th. Philipsburg, Grande Case, Milldrumhill.

” 5th—7th. Anguilla.

” 8th. Grande Case, Anse des Pères, Marigot, Philipsburg.

” 10th. District Colombier, Mount Paradis.

” 11th. Lowlands, Simonsbaylagoon.

” 12th. Bethlehem, Mount Paradis.

” 13th. Green Key, Tintamarre.

” 14th. Point Blanche, Bethlehem, Cul de Sac.

THIRD PART

PHYTOGEOGRAPHICAL

**Section A. Orological, Geological
and Meteorological**

PHYTOGEOGRAPHICAL

A. Orological, Geological and Meteorological

CHAPTER I

GEOLOGICAL AND OROLOGICAL NOTES.

Introduction.

In making notes on the Geology and Orology of the above mentioned Islands I have made use of the following treatises:

P. T. Cleve. On the Geology of the Northeastern West Indian Islands, Kongl. Vetensk. Akad. Handl. 1870. Bd. 9. no. 12. p. 19 a. f.

A. G. Högbom. Zur Petrographie der Kleinen Antillen. Bull. Geol. Institution of Upsala. VI. 1905. p. 228—229.

E. O. Hovey. Volcanoes of Martinique, Guadeloupe, and Saba. Report 8th Internat. Geogr. Congress. 1904. London, p. 447—451.

E. O. Hovey. Volcanoes of St. Vincent, St. Kitts, and Statia. *ibid.*, p. 452—454.

G. A. F. Molengraaff. De Geologie van St. Eustatius, 1886, Leiden [Dissertat. Utrecht].

G. A. F. Molengraaff. Het Geologisch verband tusschen de West-Indische eilanden, Handelingen van het eerste Nederlandsche Natuur- en Geneeskundig Congres 1888, p. 288.

G. A. F. Molengraaff. Über vulkanischen Schwefel aus West-Indien. Zeitschr. f. Krystallogr. XIV. 1888, p. 43—48.

J. W. W. Spencer. On the Geological and Physical Development of Anguilla, St. Martin, St. Bartholomew, and Sombbrero. Quarterly Journal of the Geological Society of London. 1901. vol. 57. p. 520 a. f.

J. W. W. Spencer. On the Geological and Physical Development of the St. Christopher Chain and Saba Banks. Quarterly Journal of the Geological Society of London, 1901. vol. 57. p. 534. a. f.

K. Sapper. Ein Besuch von St. Eustatius und Saba. Centralblatt für Mineralogie etc. 1903, p. 314—318.

K. Sapper. In den Vulkangebieten Mittel Amerika's und West-Indien. Stuttgart 1905.

It will appear that our geological and orological information about St. Eustatius is sufficient, whereas Saba is all but absolutely unknown in literature, so that almost all that I can communicate rests on my personal observation with the addition of some information borrowed from maps.

St. Martin is somewhat better known than Saba but a good description has not yet appeared. The maps from which I borrowed different data are the following:

St. Eustatius: *Molengraaff* 1886 l. c. pl. I.

Saba: 1. Admiralty Chart surveyed by Lieut. *G. B. Lawrance* 1850. London 1864.

2. West Indies. den Haag. ed. Nov. 1903 by the Navy-Department.

3. A map of Saba borrowed from one of the inhabitants, evidently made by *G. B. Lawrance* afore said.

St. Martin: 1. West Indies. den Haag. ed. Nov. 1903 by the Navy Department.

2. A map of St. Martin borrowed from Mr. *A. J. C. Brouwer*, Administrator of St. Martin.

3. *Molengraaff* 1888 l. c. p. 288.

4. *C. J. Hudig*, the Saltworks in the Isle of St. Martin. de Ingenieur May 24. 1905.

From these maps it appeared to me that there is reason for calling to mind the words spoken by *Kan*¹⁾ in 1888: It can „hardly be called superfluous to refer to the value of a good „map.... both for science and practice....

„In time of peace such a map might of course render invaluable „services in agricultural and industrial enterprises, in making

1) *C. M. Kan*. Onze aardrijkskundige kennis van de Molukken. Handelingen van het Eerste Ned. Nat. en Geneesk. Congres. Haarlem 1888. p. 285:

„Het kan schier overbodig geacht worden... op de waarde eener goede kaart... te wijzen. In vreedstijd kan zulk een kaart natuurlijk onschatbare diensten bewijzen: bij cultuurs en industriele ondernemingen, bij den aanleg van wegen, mijnontginningen, regeling van grondbezit en belastingen, aanleg van openbare en waterstaatswerken, enz. enz.

Doch evengroot is het nut van zulk een kaart voor verschillende beoefenaars der wetenschap, dus niet enkel voor den geograaf. Hebben niet de *H. H. Martin* en *Suringar* op meer dan een der W. I. eilanden het gemis van goede kaarten gevoeld?"

„roads in working mines, in settling landed property and taxes,
„and in public works.

„Still not only for the geographer but also for those who apply
„themselves to any science, such maps are of great use.

„Have not Messrs *Martin* and *Suringar* in more than one of the
„West Indian Islands felt the want of good maps”?

In my rambles I also greatly felt the want of really good maps. By means of panorama photos I tried to see my way in different points of the islands. Probably these photos may not be of any use in making a topographic map, still I dare say they have had for me such results as to hold myself entitled to deviate sometimes from the maps consulted by me. In this way I perhaps contributed a little to obtain a somewhat clearer description of Saba and St. Martin.

St. Eustatius.

The Isle of St. Eustatius extends from $17^{\circ} 28' 12''$ to $17^{\circ} 31' 35''$ N. and from $62^{\circ} 56' 25''$ to $62^{\circ} 59' 48''$ W.

The greatest length runs from N.W. to S.E. whereas the breadth from S. to W. except a small narrowing in the middle gradually decreases: the length of St. Eustatius amounts to 7.8 km. whereas the breadth in the southern part amounts to 3.7 km. in the middle decreases to 2.6 km. and in the northern part it does not exceed 2.7 km. The Surface is 28 □ km.

The structure of St. Eustatius is occasioned by rocks belonging to one and the same geological period.

Still from their relative age we can sharply trace out two parts viz. the northern hills on one side and the principal volcano with its eruptions on the other side. These two groups of mountains are clearly separated from each other by a wide plain. The hilly soil taking up the whole northern part of the islands consists of remnants of old volcanoes.

The southern part chiefly consists of a single very regularly built mountain which has the form of a truncated cone. The highest point of this mountain, named Quill is 581 m. above the level of the sea. On the top of it is found a circular valley shut in on all sides and of which the bottom is 297 m. above the surface of the sea.

The inner slopes of the crater are very steep and in most parts inaccessible on account of numerous perpendicular cliffs; the

slope has here a declivity of 34° to 36° the average slope in other places being 67° to 78° .

The bottom of the crater is in general tolerably flat but uneven on account of the numerous pieces of rock that have fallen down from the sides. The whole surface is covered with a bed of humus.

The fragments of rock that have served in the structure of the crater are for the greater part Augite—Audesite only here and there occurs Hornblende—Augite—Andesite.

The Quill is bordered on all sides by very steep, not seldom perpendicular rocks. The browns of these rocks form a ridge which has a width of only a few m. and the height of which varies between 391 and 581. On the outside the slope of the hill is at first, in general, very steep, but it is gradually declining to shelve away at last in a plain in a northwestern direction or to be suddenly cut off by the steep sea coast. On the southern side only inaccessible rocks called the White Wall, which with a declivity of 45° rising from the sea disturb the regularity of the profile of the mountains.

The second group of mountains taking up the N. W. part of the island consists of a great number of hills of which the highest called „Boven” has a height of 295 m.

The very steep coast that begins near Heiligenbaai bordering the further northern side of the island, is formed by a gigantic, perpendicular wall of lava (Augite—Andesite.)

By a deep ravine Boven is separated from Gilboohill. This hill 174 m high, is formed partly from ashes and partly from solid volcanic matter. The other hilly ground consists out of a number of more or less sharp tops which are connected by a ridge. This ridge runs close to the seashore from Boven to Tumbledown-dickbay and extends from there in eastern direction as far as Mary Glory, the highest top of the range. Going eastward from there we perceive that the ground is considerably sloping.

To the northern hills of St. Eustatius belongs also a crater, the exact form of a horse shoe lying south of the Tumbledown-dickbay and facing the town. The highest eastern top of this crater is Signalhill (226 m.) the northwestern one is called Pilothill.

As to petrography the northern part of St. Eustatius offers but little variety; the mineral most occurring is Augite—Andesite. Almost the whole northern part of the island is covered with a kind of good humus, produced from volcanic ashes mostly mixed

with lapilli and pumice. The slopes of these hills especially on the seaside are generally very steep, as a rule even inaccessible. Of the plains that are to be found in St. Eustatius only the large one that connects the two mountainous parts is of importance.

If we include in the plain that part only of which the general declivity does not surpass 4 degrees, the extent of it may be estimated at 5 □ km. A small plain at 0.5 □ km. may moreover be found in the Quill whereas there are some level parts in the Concordiavalley and the Venusvalley.

The formation of dells takes place on a small scale. In the northern part of the island are of importance the two large valleys that open upon the Vennsbay, the Concordiabay and Tumbledown-dickbay.

These valleys are bordered by rocks suddenly rising from the plain. The average declivity of these rocks is seldom less than 30° and often exceeds 50°.

In the southern part of the island few dells are to be found. The outside of the principal volcano is furrowed by numerous, narrow and deep ravines, the so called „guts” which show a beginning of dell formation by erosion.

Running water is found nowhere in St. Eustatius but during the heavy showers that will sometimes fall in the rainy season all aforesaid valleys and ravines are turned into beds of strong currents which by their havy fall and temporary considerable masses of water possess a great force in carrying off.

The coast of St. Eustatius is partly a steep coast partly a flat one bordering the precipices.

A steep coast occurs in the whole northern part except in places where valleys face the sea. In the southern part a steep coast is found on the eastern border of the White Wall as far as Gallowsbay and moreover in some places east of the White Wall. A flat coast bordering the precipices occurs near Oranje Bay and on the other side of the island near Concordiabay, Turtlebay and further in most places on the south-eastern and southern side. An easily practicable sandy beach, nowhere broader than 50 m. is to be found only near Oranjebay the Concordiabay and Turtlebay.

The beach on the south-eastern side is no more than 10 to 20 m. wide and is covered by pieces of rock worn by the seawater.

Saba.

The isle of Saba extends from 17° 37' 10" to 17° 39' 18" N. and from 63° 12' 25" to 63° 15' 5" N.

Round the island may be drawn a parallelogram of which the longer diagonal going from Ladderpoint to Flat Point has a length of 5.25 km.; the other from Torrents Bay to South-East Point 4.5 km. The surface is 16 □ km.

In order to get a better insight in the geological condition I collected a number of minerals during my rambles on this island. I am greatly indebted to Prof. Dr. C. E. A. Wichmann at Utrecht for his determination of these minerals. It proved to me that I brought only Andesites except some stones containing sulphur, which, especially with a view to the geological condition of St. Eustatius, was however to be expected.

As far as may result from my observations the rocks rising steeply from the sea consist of Andesites.

The conical mountain (of which Mountain Peak is 880 m. above the sea-level), rising steeply from the sea, perfectly commands the form of the island. Its top lies in the intersection of the mentioned diagonals and ends in an almost sharp summit that, viewed from St. Barthelemy seems somewhat level. To the North this mountain is sloping almost down to the seashore, higher up at an angle of 30° lower it is less steep; only near Flat point there is a tolerably large plain where the sea may be reached.

Also on the other sides of the island the rocks rise from the sea at a steep angle but do not go up to the Mountain top. They rise as far as 300, 400 or 500 m. and end in different cones which seem to rest on the central part of the island at 200—300 m. above the level of the sea.

Southwestward of Mountain Peak there is the small plain where the chief village. The Bottom, is to be found. I estimate its surface at 0.5 □ km.

South East of Mountain Peak lies another plain of still smaller surface where the little village Windwardside is situated about 300 m. above the level of the sea.

On one of the least steep parts of the conical mountain Mary-point, a knot of houses is built N.W. of the Peak at about 400 m. above the sea level whereas a group of houses, Hellsgate is built on a gentle slope N. E. of the Mountain Peak.

So the Bottom is surrounded on all sides by mountain tops and the situation is such as might easily be taken for the bottom of the crater.

In the sulphurmines that extend in the vicinity of Hellsgate near Flat Point, occurs Andesite, changed into Opal.

Here as well as in St. Eustatius there is but little dell formation; very narrow dells, so called „guts” extend in all directions from the mountain top towards the sea. So there are two issuing from the Bottom viz. Laddergut and Forthbaygut. Hellsgategut is one of the most characteristic and especially at the seashore wider than the others; all ravines abound in big stones.

Except near Flat Point and the immediate vicinity there is everywhere a steep coast in front of which are found boulder stones to a height of 2 m.

The Mountain Peak and neighbouring tops are covered with a thick bed of humus; going downward it is getting gradually thinner and on the proper arable land on the steep slopes it must be heaped up in many places, whereas in more favorable places the immense quantity of stones must be removed. In the neighbourhood of the sulphurmines the soil consists of very fine tuff.

St. Martin.

The isle of St. Martin extends from 18° to $18^{\circ} 7' 20''$ N. and from $63^{\circ} 1' 20''$ to 63° W. The greatest distance in the direction N. to S. is 13.5 km. and from W. to E. 15 km. The surface (the large lagoons not included) is about 90 □ km.

The form of St. Martin is chiefly commanded by a number of ridges that extend in the direction from N. to S.; the eastern ridge is running pretty near the shore from Oysterpond to Point Blanche (Nackedboyhill 280 m.).

A valley 2.5 km. wide separates this ridge from the other mountainous land of the Island. This mass of hills reaches in Mount Paradis (412 m.) its highest point and in all directions extend ridges of which Mont Chambord, Mont Vernon and Mill-drumhill (380 m.) end very near the seashore in the northwestern part of the Island; separated from it by a cross valley there is somewhat lower hilly country in the northern part of St. Martin. Two ridges extend from Mount Paradis to the District Colombier, whereas the two largest ranges separate Cul de Sac from the large plain and Colebay District. The ridge of hills between Cul de Sac and the plain ends in Fort Willem and Fort Amsterdam: the highest tops are 250 m. above the sea level. The largest ridge of hills east of Cul de Sac rises in Centryhill to 408 m.

Of great importance in the structure of St. Martin is a sedimentary formation, visible to the naked eye (Pointe Blanche for-

mation) consisting alternately out of Breccia, Sandstone and Limestone with a large proportion of Flint.

In general the Pointe Blanche formation covers the northern slope of the large range of hills and the south eastern slope of the small range. These older sedimentary layers are probably of Cretaceous origin. Later sediments have but a short extension on St. Martin. On the promontary near Simonsbay a hard limestone is found which perhaps is Miocene.

The Lowlands separated by Simonsbaylagoon from the other part of the Island is almost flat with some hills of which the highest rises to about 90 m. and consist for the greater part of a horizontal layer of soft marly limestone: it is of Miocene age and quite resembles the Miocene formation of Anguilla.

The minerals I brought from this island were also determined by Prof. Wichmann and proved to contain nothing particular for this island; I only mention gypsum, which I found on Green Key. Plains of some importance are found between the eastern and western ridges of hills, in the southern part (Prince quarter) however bordered by a few isolated conical hills. By a gentle slope the plain gradually passes into the western ridge of hills. Cul de Sac as an important plain is bounded on the east-northern- and western side by rather steep hills; especially Centryhill and surrounding is much more steep than the other part of the hilly country of St. Martin.

The principal rocks of the Island are composed of Quartz-diorite, Mica-diorite or Phonolite; the part that is exposed to view is generally found unchanged only on the seacoasts. In the large plain it is broken up 1 m. deep into a coarse-grained sand, containing a large proportion of mica. It was not however regularly crumbled down for there are a great many conical parts that resisted the breaking up. These continued to stand out when the wasted part was carried down by the water. So the plain contains for the greater part an abundance of fragments of rocks of considerable dimension. They form a real „Felsenmeer“ as is frequently met with on granitic soil.

Further there are some pieces of flat land between the ends of the different ridges of the hilly country between Grande Case and Marigot.

In some places from Bethlehem to Mount Paradis deep valleys are found whereas the valley in the District Colombier is one of the few spots in St. Martin where a typical tropical vegetation is met with. Especially here in the District Columbiere and also near

Mont Chambord are some small currents when the weather is not too dry.

The larger valleys open on a bay and there the coast is flat whereas almost everywhere else a rather steep coast is to be found.

The size of these bays is in direct proportion to the plains that end in them. The larger bays are separated from the sea by shoals and the inner seas thus formed are made use of as Saltponds.

The largest bays are found near the transversal valley Grande Case, near Orientbay and at Filipsburg, which is built on a sandbank.

General notes.

Whereas the map added indicates the distances between the islands, the following may serve to inform.

Cleve ¹⁾ says: „The islands north of Guadeloupe form two parallel chains from North-west to South-east. The western chain commences with Saba and consists of St. Eustatius, St. Kitts, Nevis, Redonda and Montserrat. All of those islands are volcanoes and if the line were extended farther to the north it would reach the island of Amegada, of postpliocene date and all the volcanoes seem to be of the same or nearly the same geological time. East of the volcanic range is another completely different range of islands. They are not volcanic and commence with Sombrero comprising Anguilla, St. Martin, St. Bartholomew, Barbuda and Antigua. All of these islands are of the tertiary age, eocene, miocene and pliocene.”

Although I do not agree with *Cleve* here, who wants to separate the group of St. Martin from the Virgin Islands and denies as such, that a connection might have existed between the great Antilles and the group St. Martin I want to state that we see clearly mentioned here how wide a difference there exists between the two groups the St. Martin plateau and the Christoffer chain.

About the connection between Anguilla and St. Martin *Spencer* ²⁾ says:

„The same limestone (we find in St. Martin) also occurs a short

¹⁾ *Cleve*: On the Geology of the Northeastern West Indian Islands p. 19.

²⁾ *Spencer*: On the Geological and Physical Development of Anguilla, St. Martin, St. Bartholomew and Sombrero p. 526.

„distance away in the island of Anguilla, which was once evidently a part of the coastal plain extending from the mountains of St. Martin. Tintamarre, an adjacent outlier of St. Martin, represents a fragment of the dissected coastal plain. It is underlain by the same white limestones.”

According to Spencer the various heights and depths of the different groups of St. Martin plateau and St. Christoffer chain seem to agree. As to the connection between the Caribbean isles he says ¹⁾: „... if this inference be true, then the modern islands of the West Indies formed an elevated plateau bridge between the two Americas during the two epochs of elevation, namely in the Pliocäne and Pleistocäne period.”

¹⁾ Spencer: Reconstruction of the Antillean Continent p. 132.

CHAPTER II

METEOROLOGICAL NOTES.

There is not much known about the climate of the Dutch West Indian Islands.

As they form part of the Caribbean Islands it will be sufficient for me to give all I found about the climate of these Islands in Hann, Handbuch der Meteorologie, 2e Aufl. Stuttgart 1897. The direction of the winds does not change very much and throughout the year there is generally the N. E. tradewind.

The amount of rain is mostly about 1200—2000 mm. a year; the maxima of rainfall are in May and October; the dry months are February and March. The mean temperature varies from 24—26° in January and 26—28° in July.

The following list shows the quantity of rain that will fall on the Dutch West Indian Islands and the neighbouring Islands St. Croix and St. Kitts.

**Rainfall in mm. of the West Indian Islands St. Croix,
St. Kitts, St. Eustatius, Saba and St. Martin.**

	Average quantity.					1906. 4)		
	St. Croix 1)	St. Kitts. 2)	St. Eustatius. 3)	Saba. 3)	St. Martin. 3)	St. Martin.	Saba.	St. Eustatius.
January	72	103	63	61	60	56	100	31
February	61	48	43	28	41	16	71	46
March	24	55	40	21	33	55	67	31
April	77	90	60	32	70	17	38	36
May	79	98	106	89	118	23	18	27
June	126	92	96	78	80	123	53	124
July.	191	105	118	121	85	70	65	66
August	112	144	126	143	116	83	30	61
September	123	154	139	189	145	368	465	419
October	156	173	121	123	128	48	66	66
November	181	134	131	162	139	137	88	98
December	82	96	82	72	69	214	223	148
Total	1284	1294	1125	1119	1084	1213	1282	1159

1) and 2) from Hann l. c.

3) from Went Rapport l. c.

4) from Inspectie van den Landbouw in West-Indië, Meteorologische Waarnemingen gedaan op de Meteorologische stations in de kolonien Suriname en Curaçao in het jaar 1906.

As appears from the preceding list there is a great harmony about the data concerning the rainfall on the different Islands and especially when we take in consideration that the first data about the rainfall in the Dutch West Indian Islands are most probably not so accurate as the last, there is almost the same amount of rain on each of them.

Moreover I am sure (and here I quite agree with what Went says in his Report) that the quantity of rain that falls on the higher parts of the Mountain in St. Eustatius and Saba is much more than elsewhere on these Islands.

THIRD PART

Section B. Distribution of the Wild Plants enumerated in the First Part

B. Distribution of the Wild Plants enumerated in the First Part

CHAPTER I

DISTRUBUTION OF ALL THE WILD PLANTS ENUMERATED IN THE FIRST PART.

	Both hemisph.	Continent of America.	Southern Florida.	Atlantic Islands.	Antilles.	Total.
Hymenophyllaceae		2				2
Cyatheaceae		4				4
Polypodiaceae	13	22			2	37
Gleicheniaceae		1				1
Ophioglossaceae	1					1
Lycopodiaceae	1	1				2
Psilotaceae	1					1
Selaginellaceae		2				2
Potamogetonaceae				1		1
Gramineae	23	19			4	46
Cyperaceae	8	5	1		2	16
Araceae					4	4
Lemnaceae	1					1
Bromeliaceae		6			4	10
Commelinaceae	1	3				4
Liliaceae	1	1			3	5
Amaryllidaceae		4	1			5
Dioscoreaceae	1	1				2
Musaceae		1				1
Orchidaceae		11		1	4	16
Piperaceae		11			3	14
Batidaceae	1					1
Ulmaceae		2		1		3
Moraceae		2		1	3	6
Urticaceae	1	3		1	3	8
Loranthaceae		1				1

	Both hemisph.	Continent of America.	Southern Florida.	Atlantic Islands.	Antilles.	Total.
Olacaceae	1	1				1
Aristolochiaceae	1	1				1
Polygonaceae	1	1		3	2	6
Chenopodiaceae	3					3
Amarantaceae	5	4	1	1	1	12
Nyctaginaceae	1	6			1	8
Phytolaccaceae		5				5
Aizoaceae	2	1				3
Portulacaceae	2	4			1	7
Caryophyllaceae	1					1
Menispermaceae	1				1	2
Anonaceae	1	3				4
Lauraceae	1	1			2	4
Papaveraceae	1					1
Cruciferae	1	2				3
Capparidaceae	2	6				8
Moringaceae	1					1
Crassulaceae	1					1
Rosaceae	1	2				3
Leguminosae	32	22		1	5	60
Oxalidaceae	1					1
Erythroxylaceae					2	2
Zygophyllaceae	1	2				3
Rutaceae	3	1		2	1	7
Simarubaceae	1			1	1	3
Burseraceae		1				1
Meliaceae	1	1				2
Malpighiaceae		5		1	4	10
Euphorbiaceae	9	9	3	3	8	32
Anacardiaceae	3				1	4
Celastraceae		2		1	1	4
Sapindaceae	2	2		1		5
Rhamnaceae	1	1	2			4
Vitaceae		1			1	2
Tiliaceae	4	1		1		6
Malvaceae	11	5				16

	Both hemisph.	Continent of America.	Southern Florida.	Atlantic Islands.	Antilles.	Total.
Bombacaceae	1	1				2
Sterculiaceae	2	5				7
Ochnaceae	1					1
Marcgraviaceae		1				1
Theaceae		1			1	2
Guttiferae		2			2	4
Bixaceae		1				1
Canellaceae			1			1
Flacourtiaceae		2		1	1	4
Turneraceae		2				2
Passifloraceae		5				5
Begoniaceae					1	1
Cactaceae	1	3		1	2	7
Thymelaeaceae					1	1
Rhizophoraceae		1				1
Combretaceae	2					2
Myrtaceae	1	5	4	1	4	15
Melastomataceae		4			1	5
Umbelliferae		1				1
Theophrastaceae				1	1	2
Myrsinaceae		1			1	2
Plumbaginaceae		1				1
Sapotaceae		2	1		3	6
Symplocaceae					1	1
Oleaceae		1			1	2
Loganiaceae		1				1
Gentianaceae		1				1
Apocynaceae	1	2			2	5
Asclepiadaceae	2				2	4
Convolvulaceae	4	9			3	16
Hydrophyllaceae		1				1
Borraginaceae	2	4	1	2	6	15
Verbenaceae	4	7				11
Labiatae	5	2	1			8
Solanaceae	9	4			3	16
Scrophulariaceae	1	2				3

	Both hemisph.	Continent of America.	Southern Florida.	Atlantic Islands.	Antilles.	Total.
Bignoniaceae	. . .	4	. . .	. . .	. . .	4
Gesneriaceae	. . .	. . .	. . .	. . .	2	2
Lentibulariaceae	. . .	1	. . .	. . .	. . .	1
Acanthaceae	2	2	. . .	1	2	7
Plantaginaceae	1	. . .	. . .	. . .	. . .	1
Rubiaceae	3	13	1	4	4	25
Cucurbitaceae	2	3	. . .	. . .	. . .	5
Goodeniaceae	. . .	. . .	. . .	1	. . .	1
Compositae	17	14	1	2	6	40
Total	207	301	18	34	114	674
	508			148		
			166			

I have enumerated in the Systematic Part 806 species; 132 of these are cultivated, and as will be seen from the list before given there are 674 wild plants, 166 of these are West Indian plants, of which 114 only are known on the Antilles. Another 34 are distributed in the Antilles and the other Atlantic Islands, whereas 18 are to be found both in the Antilles, the other Atlantic Islands and Southern Florida. From the 508 plants that are not restricted to the West Indies, there are 301 that occur on the Continent of America, whereas another 207 are to be found as well in the Continent of America as in the Old world.

From the plants enumerated in the Systematic Part there are 550 named in *Grisebach*: Flora of British West Indian Islands. I could make out by what I found in *Grisebach*: Die geographische Verbreitung der Pflanzen West Indiens, that from these 550 plants there are about 180 indigenous in the Antilles and the Bahamas.

When the results given by *Grisebach* in 1865 had not been changed by more recent investigations on the Flora of the West Indian Islands than the number of indigenous plants I

enumerate is this Flora ought to be between 180 and $304 = (180 + 124)$. As it appears that my number is but 148, it is evident that the data we had in 1865, about the West Indian plants have changed by the newer collections we have got from the West Indies and the neighbouring countries.

Whereas I draw this conclusion from our knowledge about the Dutch West Indian Islands so I believe that a similar study of the other Antilles would show us that there would be a still greater difference between what we know now and what *Grisebach* stated as the result of his studies on the West Indies in 1865.

In order to show that there is probably not a very large number of plants of the Dutch Antilles St. Eustatius, Saba and St. Martin that might have been overlooked by *Suringar* and myself I give the next list where I compare the numbers of the species from the larger families of the Island of St. Croix (*Eggers*: Flora of St. Croix and the Virgin Islands, 1879) and the Dutch West Indian Islands above named.

	Dutch West Indian Islands.	St. Croix.
Leguminosae	60	57
Gramineae	46	39
Compositae	40	36
Polypodiaceae	37	34
Euphorbiaceae	32	33
Rubiaceae	25	26
Moraceae	17	14
Ulmaceae		
Urticaceae		
Convolvulaceae	16	27
Cyperaceae	16	18
Malvaceae	16	25
Solanaceae	16	13
Borraginaceae	15	20
Myrtaceae	15	22
Caryophyllaceae	12	10
Portulacaceae		
Amarantaceae		

The following list may give any idea about the percentage of the species of the larger families in the Flora of the Islands St. Eustatius, Saba and St. Martin.

Leguminosae	9 %
Gramineae	6.5 %
Compositae	6 %
Polypodiaceae	5.5 %
Euphorbiaceae	4.5 %
Rubiaceae	3.5 %
Borraginaceae	2.5 %
Convolvulaceae	2.5 %
Cyperaceae	2.5 %
Malvaceae	2.5 %
Myrtaceae	2.5 %
Orchidaceae	2.5 %
Piperaceae	2.5 %
Solanaceae	2.5 %

As may be seen from the following list there is no great difference between the numbers of plants known from each of the Islands St. Eustatius, Saba and St. Martin.

St. Eustatius	Saba	St. Martin	249
St. Eustatius	Saba		91
St. Eustatius		St. Martin	65
	Saba	St. Martin	26
St. Eustatius			59
	Saba		100
		St. Martin	84
464	456	424	674

CHAPTER II

DISTRIBUTION OF THE PLANTS KNOWN ON THE ANTILLES ONLY ENUMERATED IN THE FIRST PART.

	Cuba.	Jamaica.	Haiti.	Portorico.	St. Thomas.	St. Jan.	St. Croix.	Anguilla.	St. Martin.	Saba.	St. Barthelémy.	St. Eustatius.	St. Kitts.	Barbuda.	Montserrat.	Antigua.	Guadeloupe.	Dominica.	Martinique.	St. Lucia.	St. Vincent.	Barbados.	Grenada.	Tobago.	Trinidad.
<i>Aspidium Plumieri</i>	.	.	.	.	.	.	.	.	.	X	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X
<i>Diplazium semihastatum</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
<i>Andropogon contortus</i>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X
<i>Isachne arundinacea</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Panicum diffusum</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Marsicus capillaris</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Cladium restioides</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Anthurium cordatum</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Philodendron dispar</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Philodendron giganteum</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Wittmackia lingulata</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Pitcairnia angustifolia</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Tillandsia excelsa</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Vriesea guadalupensis</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Yucca aloifolia</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
<i>Smilax coriacea</i>	.	.	.	.	.	.	.	.	.	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X

1. 2. 3. 4.

	Cuba.	Jamaica.	Haiti.	Portorico.	St. Thomas.	St. Jan.	St. Croix.	Anguilla.	St. Martin	Saba.	St. Barthélemy.	St. Eustatius,	St. Kitts.	Barbuda.	Montserrat.	Antigua.	Guadeloupe.	Dominica.	Martinique.	St. Lucia.	St. Vincent.	Barbados.	Grenada.	Tobago.	Trinidad.
<i>Smilax ilicifolia</i>	X	—	—	—	—	—	—	—	X	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Physurus hirtellus</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Physurus plantagineus</i>	—	X	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Microstylis spicata</i>	—	—	X	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Piper reticulatum</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Peperomia emarginella</i>	X	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Peperomia Langsdorffii</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Ficus Krugiana</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Ficus omphalophora</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Ficus Urbaniana</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pilea obtusata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pilea semidentata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Boehmeria ramiflora</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Coccoloba nivea</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Alternanthera crucis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pisonia subcordata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Portulaca poliosperma</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Hyperbaena dominicensis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Phoebe elongata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Nectandra Krugii</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Inga laurina</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Caesalpinia ciliata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

b, f.

a.

	Cuba.	Jamaica.	Haiti.	Porto Rico.	St. Thomas.	St. Jan.	St. Croix.	Anguilla.	St. Martin.	Saba.	St. Barthelémy.	St. Eustatius.	St. Kitts.	Barbuda.	Montserrat.	Antigua.	Guadeloupe.	Dominica.	Martinique.	St. Lucia.	St. Vincent.	Barbados.	Grenada.	Tobago.	Trinidad.
<i>Galactia dubia</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Galactia longifolia</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Galactia nummularia</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Galactia rubra</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Erythroxylum brevipes</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Erythroxylum havanense</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Fagara martinicensis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Fagara trifoliata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Picrasma antillana</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Stigmatophyllon periplocifolium</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Stigmatophyllon sericeum</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Malpighia angustifolia</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Croton astroites</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Ditaxis fasciculata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bernardia corensis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Adella ricinella</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Acalypha Poirerii</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Euphorbia articulata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Euphorbia petiolaris</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Pedilanthus tithymaloides</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Comocladia ilicifolia</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Maytenus elliptica</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

	Cuba.	Jamaica.	Haiti.	Portorico.	St. Thomas.	St. Jan.	St. Croix.	Anguilla.	St. Martin.	Saba.	St. Barthélemy.	St. Enstatius.	St. Kitts.	Barbuda.	Montserrat.	Antigua.	Gadeloupe.	Dominica.	Martinique.	St. Lucia.	St. Vincent.	Barbados.	Grenada.	Tobago.	Trinidad.
<i>Cissus obovata</i>	—	—	—	×	—	—	×	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Freziera undulata</i>	—	—	—	—	—	—	—	—	—	×	—	—	×	—	—	—	—	—	×	—	—	—	—	—	—
<i>Marila racemosa</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Clusia alba</i>	—	—	—	—	—	—	×	—	—	×	—	—	×	—	—	—	—	—	×	—	—	—	—	—	—
<i>Samyda serrulata</i>	—	—	—	×	—	—	—	—	×	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	×
<i>Begonia domingensis</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Daphnopsis caribaea</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Myrcia splendens</i>	×	×	×	×	—	—	—	—	×	—	—	—	—	—	—	—	—	—	×	—	—	—	—	—	×
<i>Calyphanthes Boldinghii</i>	—	—	—	—	—	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Eugenia cordata</i>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Eugenia procera</i>	×	×	×	×	×	×	×	—	×	—	—	×	—	—	—	×	—	—	×	—	—	×	—	—	—
<i>Jacquinia armillaris</i>	×	×	×	×	×	×	×	—	×	—	×	—	×	—	—	×	—	—	×	—	—	×	—	—	—
<i>Ardisia coriacea</i>	×	×	×	×	×	×	×	—	×	—	×	—	×	—	—	×	—	—	×	—	—	×	—	—	—
<i>Lucuma multiflora</i>	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Bumelia obovata</i>	—	—	—	×	—	—	—	—	×	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Chrysophyllum argenteum</i>	×	×	×	×	—	—	—	—	×	—	×	—	—	—	—	—	—	—	×	—	—	×	—	—	—
<i>Symplocos martinicensis</i>	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Forestiera Eggersiana</i>	—	—	—	×	—	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Plumiera alba</i>	—	—	—	×	—	—	×	—	×	—	×	—	×	—	—	—	—	—	—	—	—	—	—	—	—
<i>Urechites suberecta</i>	—	—	—	×	—	—	×	—	×	—	×	—	×	—	—	—	—	—	—	—	—	—	—	—	—
<i>Ibattia maritima</i>	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Jacquemontia pentantha</i>	—	—	—	×	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

4.

5, 6.

5.
10, 13, 14, 15.

4, 5, 13.

12.

	Cuba	Jamaica	Haiti	Portorico	St. Thomas	St. Croix	Anguilla	St. Martin	Saba	St. Barthélemy	St. Eustatius	St. Kitts	Barbuda	Montserrat	Antigua	Guadeloupe	Dominica	Martinique	St. Lucia	St. Vincent	Barbados	Grenada	Tobago	Trinidad
<i>Ipomoea arenaria</i>	X					X		X																
<i>Ipomoea sphenophylla</i>											X													
<i>Cordia salvifolia</i>									X			X				X		X						
<i>Cordia sulcata</i>		X		X	X	X					X						X	X	X	X	X			
<i>Cordia tremula</i>		X		X											X		X	X				X		
<i>Rochefortia acanthophora</i>				X				X									X	X						
<i>Tournefortia caribaea</i>																								
<i>Tournefortia foetidissima</i>		X	X	X		X		X		X						X	X	X						
<i>Lycium americanum</i>	X	X	X	X		X	X	X		X														
<i>Solanum racemotum</i>				X		X		X		X				X	X	X	X	X		X	X			
<i>Cestrum laurifolium</i>	X	X	X	X		X			X	X					X	X	X	X			X	X		6, 7, 9.
<i>Besleria lutea</i>		X		X					X	X					X	X	X	X	X	X	X	X	X	
<i>Gesneria ventricosa</i>		X								X		X		X										
<i>Justicia sessilis</i>				X		X		X	X	X					X									
<i>Beloperone eustachiana</i>								X	X	X					X									
<i>Psychotria pendula</i>								X	X	X		X		X	X	X	X	X		X	X	X		
<i>Palicourea Pavetta</i>	X	X		X		X			X	X														
<i>Borreria podoccephala</i>	X	X		X		X			X	X														
<i>Mitracarpus polycladus</i>	X			X					X	X					X									6, 8.
<i>Eupatorium canescens</i>	X			X		X				X					X									
<i>Eupatorium macranthum</i>				X		X			X	X					X									
<i>Erigeron spathulatus</i>						X		X							X			X			X		X	
<i>Clibadium erosum</i>				X					X						X			X			X			
<i>Pectis humifusa</i>						X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	X	6, 7, 13.

1 = Virgin Islands.
2 = Vieques.
3 = Crabben Island.
4 = Bequia.
5 = Tortola.
6 = Desirade.

7 = Marie Galante.
8 = Isla de Pinos.
9 = Les Saintes.
10 = Curacao.
11 = Culebra Island.
12 = Margareta.

13 = Mustique Island.
14 = Canouan.
15 = Bonaire.

The list given in this chapter needs no further explanation.

As may be seen from the great amount of plants occurring in such a small number of different Islands it will be clear that our knowledge of the Flora of the West Indian Islands is far from being complete.

That is the reason why I will not try to draw any conclusion about the distribution of the different species except a single one and this with all kinds of reserve.

I draw the attention to the data that may be gathered from the following lists.

On the Islands only to the **north** of the Dutch West Indian Islands together with these Islands are collected:

	St. Martin.	Saba.	St. Eustatius.
Diplazium semihastatum	—	—	×
Mariscus capillaris	×	×	—
Tillandsia excelsa	—	×	—
Yucca aloifolia	×	—	—
Smilax coriacea	—	—	×
Smilax ilicifolia	×	—	—
Alternanthera crucis	×	—	—
Portulaca poliosperma	×	—	—
Erythroxylum havanense	×	×	×
Stigmatophyllum sericeum	×	—	—
Ditaxis faxiculata	×	—	—
Adelia ricinella	×	—	—
Acalypha Poiretii	×	—	—
Cissus obovata	×	—	—
Forestiera Eggersiana	×	—	—
Ipomoea arenaria	×	—	—
Rochefortia acanthophora	×	—	×
Lycium americanum	×	—	—
Mitracarpus polycladus.	—	×	—

On the Islands only to the **south** of the Dutch West Indian Islands together with these Islands are collected:

	St. Martin.	Saba.	St. Eustatius.
<i>Philodendron dispar</i>	—	×	×
<i>Vriesea guadalupensis</i>	—	×	×
<i>Piper reticulatum</i>	—	×	×
<i>Ficus Krugiana</i>	—	×	×
<i>Ficus omphallophora</i>	—	×	×
<i>Boehmeria ramiflora</i>	—	×	×
<i>Caesalpinia ciliata</i>	×	×	×
<i>Galactia rubra</i>	×	×	×
<i>Freziera undulata</i>	—	×	—
<i>Marila racemosa</i>	—	×	—
<i>Cordia salvifolia</i>	—	×	×
<i>Cordia tremula</i>	—	×	—
<i>Tournefortia caribaea</i>	—	×	—
<i>Beloperone eustachiana</i>	×	×	×
<i>Eupatorium macranthum</i>	—	×	×

As may be seen from these two lists the greater number of the plants found on the northern Islands only are but to be met with on St. Martin, the greater number of the plants found on the southern Islands only are but to be met with on Saba and St. Eustatius.

In conclusion, I believe the different indigenous plants of the Antilles make it sufficiently clear that the vegetation of St. Martin has been influenced by some other circumstances than the vegetation of Saba and St. Martin.

Species only known from the Dutch West Indians Islands are:

<i>Galactia nummularia</i>	St. Martin.
<i>Calypttranthes Boldinghii</i>	St. Martin.
<i>Ipomoea sphenophylla</i>	St. Eustatius.

THIRD PART

.

Section C. The vegetation of St. Eustatius, Saba and St. Martin

.

C. The vegetation of St. Eustatius, Saba and St. Martin

CHAPTER I

NOTES ON THE ASPECT OF THE VEGETATION OF THE THREE DUTCH WEST INDIAN ISLANDS.

Introduction.

The vegetation of St. Eustatius, Saba and St. Martin is that of a tropical zone, without any pronounced dry season. As may be seen from the meteorological data the quantity of rain is almost sufficient to make possible the existence of a tropical rainforest vegetation.

The N. E. tradewinds however are of great influence upon the vegetation; where these winds are not broken by mountains we may meet with the typical vegetation of trees blown in a certain direction as well as with the Cactaceae and thornplants, characteristics of a dry climate.

From the edaphic factors that exert influence upon the character of the different Plantformations I will in this chapter only draw the attention to the amount of humus that covers the rocks. It is one of the principal factors, as there are a great number of localities where but a small quantity of humus is to be found. Only there were no strong wind is blowing, where the atmosphere is humid and where the quantity of humus has surpassed a minimum the tropical vegetation shows itself in its wellknown typical features.

But the above named factors have not exclusively determined the plantcovering of the Islands; one of the greatest factors has been the influence of man, who has been destroying the woods and introducing many plants in cultivating the soil. So there are a great number of plants introduced that have superseded the primitive vegetation.

When taking the vegetation in general and without entering into details we may consider it to be identical on the three Islands. From the short notes however I give in this chapter on the aspect of the different Islands it will be obvious that on each of the three Islands a particular Formation gives a predominant character to each of them.

St. Eustatius.

The monotony of the level part of St. Eustatius is relieved by some plantations surrounded by trees and by roads also bordered by trees and shrubs.

The formation that predominates in the „Cultuurvlakte” is a dry shrubby vegetation of a greyish aspect that I will call the Croton vegetation, following the nomenclature of *Eggers* in his Flora of St. Croix and the Virgin Islands.

The greater part of the hills near Signalhill and Concordia are also covered with the same vegetation.

Only the higher parts of these hills as well as the Quill are covered with a greenish wood that I will call the Eriodendron vegetation also in analogy with *Eggers* l. c.

These two parts of the vegetation are not wholly separated and both of them are intermixed with those plants which form the vegetation of the cultivated region.

The impracticable parts of the rocky cliffs covered only with few plants form a typical vegetation with thicker or more coriaceous leaves in close relation with the vegetation of the seashore; the latter is not very well developed in St. Eustatius as there is but a small beach and the Cultuurvlakte is about 40 m. above the sealevel.

Without entering into details we can practically divide the vegetation of St. Eustatius in:

1. Vegetation of the higher parts of the mountains.
2. Vegetation of the Cultuurvlakte and the lower parts of the mountains.
3. Vegetation of the seashore and the rocky localities of the Island.
4. Vegetation of the cultivated region.

Whereas a survey of the vegetation of St. Eustatius based on these principles is given in the following lists it may be mentioned here that an almost synomical list could be given, viz:

1. Eriodendron vegetation.

2. Croton vegetation.

3. Littoral vegetation.

4. Eriodendron- and Croton vegetation intermixed with and partly replaced by plants of the cultivated region.

We will see in Chapter II of this Section that these two series of names do not quite agree and this is the reason why I use the first mentioned names when in the following lists I give a survey of the distribution of the plants on the different parts of St. Eustatius. In these lists as well as in the other ones I give in this Chapter the West Indian plants are printed in italics whereas the plants that are cultivated on the Dutch West Indian Islands are printed in small type.

1. Vegetation of the Higher Parts of the Mountains of St. Eustatius.

Acacia macracantha, *Acacia tortuosa*, *Aegiphila martinicensis*, *Allophylus occidentalis*, *Anomis caryophyllata*, *Anacardium occidentale*, *Andira jamaicensis*, *Anona muricata*, *Anona reticulata*, *Anthurium cordatum*, *Aspidium Plumieri*, *Asplenium cristatum*, *Asplenium obtusifolium*, *Asplenium serratum*, *Bernardia corensis*, *Besleria lutea*, *Bignonia unguis cati*, *Blechnum occidentale*, *Bumelia obovata*, *Bursera simaruba*, *Byrsonima spicata*, *Calotropis procera*, *Capparis Breynia*, *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Casearia parvifolia*, *Cassia glandulosa*, *Catopsis nutans*, *Cecropia peltata*, *Ceiba pentandra*, *Celtis Iguanacea*, *Centrosema virginianum*, *Chiococca racemosa*, *Chrysobalanus icaco*, *Cissus sicyoides*, *Citharelyxum quadrangulare*, *Clusia alba*, *Coccoloba diversifolia*, *Coccoloba nireia*, *Coccoloba uvifera*, *Coffea arabica*, *Corchorus hirsutus*, *Cordia salvifolia*, *Crotalaria retusa*, *Croton astroites*, *Cyathea serra*, *Daphnopsis caribaea*, *Desmanthus depressus*, *Desmodium supinum*, *Dipholis salicifolia*, *Dodonaea viscosa*, *Elaphoglossum conforme*, *Elaphoglossum petiolatum*, *Epidendrum anceps*, *Epidendrum difforme*, *Erithalis fruticosa*, *Erythroxylum havanense*, *Eugenia axillaris*, *Eugenia cordata*, *Eugenia ligustrina*, *Eupatorium conyzoides*, *Eupatorium macranthum*, *Eugara martinicensis*, *Eugara trifoliata*, *Ficus Krugiana*, *Ficus populnea*, *Ficus Urbaniana*, *Galactia dubia*, *Gesneria ventricosa*, *Guettarda parvifolia*, *Guettarda scraba*, *Gyminda latifolia*, *Gymnanthes lucida*, *Heliconia Bihai*, *Heteropteris lanceolata*, *Heteropteryx purpurea*, *Hillia longifolia*, *Hyperbaena domingensis*, *Inga laurina*, *Impomoea dissecta*, *Jacquemontia pentantha*, *Krugiodendron ferreum*, *Lantana camara*, *Lantana involu-*

crata, *Leiphaimos aphylla*, *Linociera compacta*, *Lonchocarpus violaceus*, *Lycopodium taxifolium*, *Mammea americana*, *Maytenus ellipticus*, *Melothria fluminensis*, *Miconia impetiolaris*, *Miconia laevigata*, *Mimosa pudica*, *Morisonia americana*, *Myrcia paniculata*, *Myrcia splendens*, *Nectandra coriacea*, *Nectandra Krugii*, *Neurogramme calomelanos*, *Oplismenus setarius*, *Palicourea Pavetta*, *Panicum divaricatum*, *Panicum insulare*, *Paspalum hemisphaericum*, *Passiflora laurifolia*, *Passiflora suberosa*, *Pectis humifusa*, *Peperomia acuminata*, *Peperomia glabella*, *Peperomia Langsdorffii*, *Peperomia obtusifolia*, *Peperomia scandens*, *Persea americana*, *Pharus glaber*, *Philodendron dispar*, *Philodendron oxycardium*, *Phoebe elongata*, *Physurus plantagineus*, *Picrasma Antillana*, *Pilea semidentata*, *Piper dilatatum*, *Piper medium*, *Piper reticulatum*, *Piriqueta villosa*, *Pisonia aculeata*, *Pisonia fragrans*, *Pisonia subcordata*, *Pitcairnia angustifolia*, *Polypodium brasiliense*, *Polypodium lycopodioides*, *Polypodium phyllitides*, *Polypodium plumula*, *Polystachya luteola*, *Prescotria stachyoides*, *Prunus myrtifolius*, *Psychotria horizontalis*, *Psychotria rufescens*, *Psychotria uliginosa*, *Randia aculcata*, *Rauwolfia Lamarekii*, *Rhynchosia reticulata*, *Samyda serrulata*, *Schaefferia frutescens*, *Schoepfia Schreberi*, *Scleria lithosperma*, *Smilax coriacea*, *Smilax guianensis*, *Solanum lanceaefolium*, *Solanum racemosum*, *Solanum racemosum igneum*, *Spermacece tenuior*, *Stenochlaena sorbifolia*, *Stenolobium stans*, *Stenorrhynchus orchioides*, *Stigmatophyllon periplocifolium*, *Swietenia mahagoni*, *Symplocos martinicensis*, *Tabernaemontana citrifolia*, *Tecoma leucoxydon*, *Ternstroemia peduncularis*, *Theobroma Cacao*, *Tournefortia foetidissima*, *Tragia volubilis*, *Trema Lamarckianum*, *Trema micranthum*, *Triumfetta rhomboidea*, *Urera Caracasana*, *Vernonia punctata*, *Vittaria lineata*, *Vriesea guadalupensis*, *Waltheria americana*, *Wedelia buphthalmoides*, *Xylosma buxifolium*.

2. Vegetation of the Cultuurvlakte and the Lower Parts of the Mountains of St. Eustatius.

Acacia macracantha, *Acacia tortuosa*, *Amarantus polygonoides*, *Andira jamaicensis*, *Andropogon contortus*, *Aristida americana*, *Ayena pusilla*, *Beloperone eustachyana*, *Bernardia corensis*, *Borreria podoccephala*, *Brassavola cucullata*, *Canavalia obtusifolia*, *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Cassia bicapsularis*, *Cassia glandulosa*, *Catopsis nutans*, *Celtis Iguanacea*, *Cenchrus tribuloides*, *Centrosema virginianum*, *Citharexylum quadrangulare*, *Coccoloba uvifera*, *Commelina virginica*, *Comocladia ilicifolia*, *Corchorus hirsutus*, *Cordia salvifolia*, *Cracca caribaea*,

Croton astroites, *Croton flavens*, *Cuscuta americana*, *Dactyloctenium aegyptium*, *Dalechampia scandens*, *Desmanthus depressus*, *Desmanthus virgatus*, *Desmodium supinum*, *Diodia rigida*, *Dodonaea viscosa*, *Duranta repens*, *Epidendrum papilionaceum*, *Eragrostis ciliaris*, *Erithalis fruticosa*, *Ernodea litoralis*, *Eugenia cordata Sintensis*, *Eugenia ligustrina*, *Eugenia procera*, *Eugenia uniflora*, *Euphorbia articulata*, *Evolvulus sericeus*, *Fourcroya gigantea*, *Galactia dubia*, *Gouania domingensis*, *Gymnanthes lurida*, *Heteropteryx purpurea*, *Hymenocallis caribaea*, *Ipomoea sphenophylla*, *Jacquemontia pentantha*, *Jacquinia armillaris*, *Jatropha curcas*, *Jatropha gossypifolia*, *Justicia sessilis*, *Krameria Ixina*, *Lantana camara*, *Lantana involucrata*, *Malpighia puniceifolia*, *Malvastrum spicatum*, *Maytenus elliptica*, *Melicocca bijuga*, *Melochia tomentosa*, *Melocactus communis*, *Morisonia americana*, *Opuntia Ficus indica*, *Opuntia triacantha*, *Opuntia Tuna*, *Panicum insulare*, *Pappophorum alopecuroideum*, *Pectis humifusa*, *Phoradendron trinervium*, *Phyllanthus distichus*, *Pilea microphylla*, *Pisonia fragrans*, *Pisonia subcordata*, *Pitcairnia angustifolia*, *Pithecolobium unguis cati*, *Polypodium lycopodioides*, *Portulaca oleracea*, *Randia aculeata*, *Rauwolfia Lamareckii*, *Rhacoma crossopetalum*, *Rhynchosia reticulata*, *Rochefortia acanthophora*, *Samyda serrulata*, *Schaefferia frutescens*, *Setaria setosa*, *Sida ciliaris*, *Sida supina*, *Solanum racemosum*, *Solanum racemosum igneum*, *Sporobolus virginicus*, *Stenolobium stans*, *Stigmatophyllon periplocifolium*, *Strumpfia maritima*, *Stylosanthes hamata*, *Talinum triangulare*, *Tamarindus indica*, *Tecoma leucoxyton*, *Tephrosia cinerea*, *Tournefortia volubilis*, *Tragus racemosus*, *Turnera ulmifolia*, *Urechites suberecta*, *Vernonia punctata*, *Waltheria americana*.

3. Vegetation of the Seashore and Rocky Parts of St. Eustatius.

Amarantus polygonooides, *Andropogon contortus*, *Argemone mexicana*, *Aristida americana*, *Borreria podocephala*, *Caesalpinia crista*, *Calonyction tuba*, *Canavalia obtusifolia*, *Cassia bicapsularis*, *Comocladia ilicifolia*, *Conocarpus erectus*, *Ernodea litoralis*, *Eupatorium canescens*, *Euphorbia articulata*, *Euphorbia buxifolia*, *Evolvulus sericeus*, *Heliotropium curassavicum*, *Hibiscus tiliaceus*, *Hippomane mancinella*, *Ipomoea pescaprae*, *Jacquinia armillaris*, *Lithophila muscodes*, *Malpighia puniceifolia*, *Melocactus communis*, *Opuntia Ficus indica*, *Opuntia spinosissima*, *Opuntia triacantha*, *Opuntia Tuna*, *Panicum diffusum*, *Pappophorum alopecuroideum*, *Rhacoma crossopetalum*, *Sesuvium portulacastrum*, *Setaria setosa*, *Strumpfia*

maritima, *Tephrosia cinerea*, *Thespesia populnea*, *Tournefortia gnaphalodes*.

4. Vegetation of the Cultivated Region of St. Eustatius.

Abrus precatorius, *Abutilon indicum*, *Acacia macracantha*, *Acacia riparia*, *Acacia tortuosa*, *Acalypha chamaedrifolia*, *Achras sapota*, *Achyranthes obtusifolia*, *Ageratum conyzoides*, *Alternanthera repens*, *Amarantus spinosus*, *Amarantus tristis*, *Ambrosia hispida*, *Anona montana*, *Anona muricata*, *Antheophora hermaphroditica*, *Antigonon leptopus*, *Boerhaavea hirsuta*, *Boerhaavea paniculata*, *Bryophyllum pinnatum*, *Calotropis procera*, *Capparis Breynia*, *Capparis cynophallophora*, *Capraria biflora*, *Cardiospermum halicacabum*, *Cassia occidentalis*, *Cenchrus trihuloides*, *Centrosema virginiana*, *Cheilanthes microphylla*, *Chenopodium ambrosioides*, *Chenopodium murale*, *Cissampelos Pareira*, *Clerodendron aculeatum*, *Coccoloba uvifera*, *Coleus amboinicus*, *Commelina virginica*, *Corchorus acutangulus*, *Corchorus hirsutus*, *Cracca caribaea*, *Crotalaria verrucosa*, *Croton astroites*, *Croton flavens*, *Croton lobatus*, *Cynodon dactylon*, *Cyperus rotundus*, *Datura Metel*, *Datura Stramonium*, *Desmanthus depressus*, *Desmanthus virgatus*, *Desmodium supinum*, *Eleusine indica*, *Eragrostis ciliaris*, *Eupatorium conyzoides*, *Euphorbia heterophylla*, *Euphorbia hypericifolia*, *Euphorbia pilulifera*, *Festuca Myurus*, *Fleurya aestuans*, *Gynandropsis pentaphylla*, *Haematoxylon campechianum*, *Heliotropium indicum*, *Heliotropium parviflorum*, *Hura crepitans*, *Hyptis pectinata*, *Ibatia maritima*, *Indigofera suffruticosa*, *Jacquemontia pentantha*, *Jatropha gossypifolia*, *Kallstroemia maxima*, *Lantana involucrata*, *Leonotis nepetaefolia*, *Leonurus sibiricus*, *Leucaena glauca*, *Lochnera rosea*, *Malvastrum spicatum*, *Melicocca bijuga*, *Melocactus communis*, *Melochia pyramidata*, *Melochia tomentosa*, *Microtea debilis*, *Mimosa pudica*, *Mirabilis Jalapa*, *Mollugo verticillata*, *Momordica Charantia*, *Hydrolea jamaicensis*, *Neurogramme calomelanos*, *Ocimum micranthum*, *Opuntia Ficus indica*, *Opuntia triacantha*, *Opuntia Tuna*, *Panicum insulare*, *Panicum maximum*, *Panicum sanguinale*, *Parthenium Hysterophorus*, *Paspalum fimbriatum*, *Paspalum hemisphaericum*, *Passiflora laurifolia*, *Pedilanthus tithymaloides*, *Peperomia petiolaris*, *Phaseolus lathyroides*, *Phyllanthus niruri*, *Pilea microphylla*, *Piriqueta villosa*, *Plumbago scandens*, *Polanisia viscosa*, *Polypodium aureum*, *Porphyllum ellipticum*, *Portulaca halimoides*, *Portulaca oleracea*, *Portulaca quadrifida*, *Priva echinata*, *Rauwolfia Lamareckii*, *Rhynchosia reticulata*, *Rivina humilis*, *Salvia occidentalis*, *Salvia serotina*, *Setaria glauca*, *Setaria setosa*, *Sida carpinifolia*, *Sida cordifolia*,

Sida supina, *Solanum nigrum nodiflorum*, *Solanum racemosum*, *Solanum racemosum igneum*, *Sonchus oleraceus*, *Spermacoce tenuior*, *Stachytarpheta jamaicensis*, *Stigmatophyllon periplocifolium*, *Stylosanthes hamata*, *Talinum triangulare*, *Tamarindus indica*, *Tecoma leucoxydon*, *Tephrosia cathartica*, *Thunbergia fragans*, *Tillandsia recurvata*, *Vernonia punctata*, *Waltheria americana*, *Wedelia buphthalmoides*, *Zornia diphylla*.

Saba.

The Island of Saba makes a very different impression as St. Eustatius does, especially by the total absence of a level country like the Cultuurvlakte on the latter Island.

Owing to the conical form of Saba there are only steep slopes of hills and these hills are covered by the same vegetation as those in St. Eustatius, viz.: the *Eriodendron* vegetation. Moreover there is a good quantity of almost naked rock where a very poor flora is to be found.

According to the fact that almost in all the localities that are apt for cultivation the soil is occupied by the inhabitants, there is a well developed vegetation of the cultivated region intermixed with the *Eriodendron* vegetation.

The *Croton* vegetation is only to be seen in a very few localities and nowhere on Saba it is so predominant as in St. Eustatius.

That part of the Mountain higher than 500 m. above the sea-level, though also greatly influenced by ruderal plants has a very well developed tropical rainforest which, without losing the typical character of the St. Eustatius forests is characterised by its great number of Fern trees.

The vegetation of the seashore is very poor and has no influence on the general aspect of the Saba vegetation.

Whereas it is not so easy to make a sharp difference between the four Plant formations still I think it better also to trace out here the four following typical groups:

1. Vegetation of the Mountain top and the greater part of all the slopes of the Island.
2. Vegetation of some arid hills.
3. Vegetation of the seashore and rocky localities.
4. Vegetation of the cultivated region.

Referring to page 267 we might take the same series of names I gave there for the first three divisions whereas the

fourth division could be replaced by: *Eriodendron* vegetation intermixed with and partly replaced by ruderal plants.

1. Vegetation of the Mountaintop and the Greater Part of the Slopes of Saba.

Ageratum conyzoides, *Alsophila phalerata*, *Amomis caryophyllata*, *Andropogon glomeratus*, *Anona muricata*, *Anthurium cordatum*, *Aspidium Plumierii*, *Aspidium trifoliatum*, *Asplenium cristatum*, *Baccharis cotinifolia*, *Begonia domingensis*, *Besleria lutea*, *Beureria succuleata*, *Blechnum occidentale*, *Blechnum Brownei*, *Boehmeria ramiflora*, *Bromelia pinguin*, *Bryophyllum pinnatum*, *Bursera simaruba*, *Callisia repens*, *Capparis Breynia*, *Capparis cynophallophora*, *Casearia parvifolia*, *Cecropia peltata*, *Celtis Ignanacea*, *Cestrum laurifolium*, *Chaptalia nutans*, *Chiococca racemosa*, *Chloris radiata*, *Chrysobalanus icaco*, *Chrysophyllum argenteum*, *Cissus sicyoides*, *Cladium restioides*, *Clusia alba*, *Coccoloba diversifolia*, *Coccoloba nivea*, *Commelina virginica*, *Comocladia ilicifolia*, *Cordia salrifolia*, *Crotalaria retusa*, *Cyathea arborea*, *Cyathea serra*, *Cyperus distans*, *Cyperus rotundus*, *Daphnopsis caribaea*, *Desmodium axillare*, *Desmodium supinum*, *Dichromena radicans*, *Dioscorea alata*, *Dioscorea trifida*, *Diplazium striatum*, *Dryopteris opposita*, *Dryopteris reticulata*, *Duranta repens*, *Elaphoglossum conforme*, *Elaphoglossum latifolium*, *Epidendrum anceps*, *Epidendrum ciliare*, *Epidendrum globosum*, *Eragrostis ciliaris*, *Eragrostis pilosa*, *Erechthites valerianaeifolia*, *Eryngium foetidum*, *Erythroxylon havanense*, *Eugenia axillaris*, *Eugenia procera*, *Eupatorium iresinoides*, *Eupatorium macrantum*, *Eupatorium paniculatum*, *Fagara martinicensis*, *Ficus omphalophora*, *Ficus Urbaniana*, *Freziera undulata*, *Gesneria ventricosa*, *Gleichenia bifida*, *Gonzalagunia hirsuta*, *Guettarda parvifolia*, *Heliconia Bihai*, *Heliotropium indicum*, *Heliotropium parviflorum*, *Hemidiodia ocimifolia*, *Hemitelia horrida*, *Hillia longifolia*, *Hirtella triandra*, *Hyperbaena domingensis*, *Hypoxis decumbens*, *Ichnanthes pallens*, *Indigofera suffruticosa*, *Ipomoea fastigiata*, *Isachne arundinacea*, *Isachne rigens*, *Lagascea mollis*, *Lagenaria vulgaris*, *Leiphaimos aphylla*, *Lepidium virginicum*, *Leptochloa virgata*, *Lochnera rosea*, *Lycopodium taxifolium*, *Marcgravia umbellata*, *Mariscus capillaris*, *Mariscus flavus*, *Melia Azedarach*, *Melochia tomentosa*, *Miconia laevigata*, *Microstylis spicata*, *Myrcia coriacea*, *Nectandra Krugii*, *Nephrolepis exaltata*, *Neurogramme calomelanos*, *Neurolaena lobata*, *Oplismenus setarius*, *Opuntia triacantha*, *Ornithidium coccineum*, *Panicum brevifolium*, *Palicourea Pavetta*, *Paspalum conjugata*

tum, *Passiflora foetida*, *Passiflora rubra*, *Peperomia acuminata*, *Peperomia emarginella*, *Peperomia glabella*, *Peperomia hernandifolia*, *Peperomia obtusifolia*, *Peperomia urocarpa*, *Philodendron giganteum*, *Philodendron oxycardium*, *Physurus plantagineus*, *Phytolacca icosandra*, *Picrasma antillana*, *Pilea microphylla*, *Pilea obtusata*, *Pilea parietaria*, *Piper reticulatum*, *Pisonia aculeata*, *Pisonia subcordata*, *Polypodium cervina*, *Polypodium aureum*, *Polypodium crassifolium*, *Polypodium lycopodioides*, *Polypodium phyllitidis*, *Polypodium piloselloides*, *Polypodium squamatum*, *Polypodium serrulatum*, *Polypodium trichomanoides*, *Psilotum nudum*, *Psychotria horizontalis*, *Psychotria rufescens*, *Pteris biaurita*, *Rapanea ferruginea*, *Rhynchosia reticulata*, *Rhynchospora polyphylla*, *Sauvagesia erecta*, *Salvia serotina*, *Schoepfia Schreberi*, *Scleria pterota*, *Secchium edule*, *Selaginella flabellata*, *Selaginella substipitata*, *Setaria glauca*, *Setaria setosa*, *Solanum lanceaefolium*, *Solanum nigrum nodiflorum*, *Solanum racemosum*, *Sonchus oleraceus*, *Stachytarpheta jamaicensis*, *Synedrella nodiflora*, *Symplocos martinicensis*, *Tabernaemontana citrifolia*, *Talinum triangulare*, *Tecoma leucoxydon*, *Tibouchina strigosa*, *Tillandsia excelsa*, *Torulinum confertum*, *Tournefortia foetidissima*, *Tragia volubilis*, *Trichomanes alatum*, *Triumfetta semitriloba*, *Urena lobata*, *Utricularia alpina*, *Vernonia punctata*, *Vriesea guadalupensis*, *Wedelia bupththalmoides*.

2. Vegetation of Dry Hills of Saba.

Acacia Farnesiana, *Anona muricata*, *Capparis cynophallophora*, *Capparis jamaicensis*, *Cardiospermum halicacabum*, *Cissampelos Pareira*, *Coccoloba diversifolia*, *Colubrina ferruginosa*, *Cordia salvifolia*, *Croton flavens*, *Ernodea litoralis*, *Gundlachia domingensis*, *Indigofera suffruticosa*, *Melocactus communis*, *Opuntia Ficus indica*, *Opuntia triacantha*, *Opuntia Tuna*, *Pectis humifusa*, *Pisonia subcordata*, *Polypodium piloselloides*, *Psychotria horizontalis*, *Sesuvium portulacastrum*, *Talinum triangulare*, *Tecoma leucoxydon*, *Tragus racemosus*.

3. Vegetation of the Seashore and Rocky Parts of Saba.

Aristida americana, *Bursera simaruba*, *Calotropis procera*, *Canavalia obtusifolia*, *Capraria biflora*, *Cheilanthes microphylla*, *Chenopodium ambrosioides*, *Chloris paraguayensis*, *Coccoloba uvifera*, *Dactyloctenium aegyptium*, *Eupatorium conyzoides*, *Euphorbia articulata*, *Hippomane mancinella*, *Lithophila muscoides*, *Lochnera rosea*, *Mariscus brizaeus*, *Melocactus communis*, *Neurogramme calomelanos*, *Opuntia Ficus indica*, *Opuntia triacantha*, *Opuntia Tuna*.

Panicum insulare, *Pappophorum alopecuroideum*, *Pectis febrifuga*, *Pisonia aculeata*, *Pisonia subcordata*, *Portulaca oleracea*, *Sesuvium portulacastrum*, *Sida rhombifolia*, *Solanum racemosum*, *Stylosanthes hamata*, *Tecoma leucoxylon*, *Turnera ulmifolia*.

4. Vegetation of the Cultivated Region of Saba.

Abrus precatorius, *Acacia Farnesiana*, *Acalypha chamaedrifolia*, *Acnistus arborescens*, *Achyranthes obtusifolia*, *Adiantum tenerum*, *Ageratum conyzoides*, *Alternanthera repens*, *Alysicarpus nummularius*, *Amarantus paniculatus*, *Amarantus spinosus*, *Amarantus tristis*, *Ambrosia hispida*, *Anacardium occidentale*, *Anona montana*, *Anona muricata*, *Anona reticulata*, *Anona squamosa*, *Antheophora hermaphrodita*, *Antigonon leptopus*, *Argemone mexicana*, *Aristida americana*, *Asclepias nivea*, *Asplenium pumilum*, *Beloperone eustachiana*, *Besleria lutea*, *Beureria succulenta*, *Bidens bippinatus*, *Bidens leucantha*, *Bignonia unguis-cati*, *Bixa Orellana*, *Blechnum occidentale*, *Blechnum Brownei*, *Boerhaavea hirsuta*, *Boerhaavea paniculata*, *Borreria podocephala*, *Bouchea Ehrenbergii*, *Boussingaultia baselloides*, *Brassica integrifolia*, *Bromelia pinguin*, *Bryophyllum pinnatum*, *Bumelia obovata*, *Bursera simaruba*, *Byrsonima spicata*, *Caesalpinia crista*, *Cajanus indicus*, *Callisia repens*, *Capparis Breynia*, *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Capraria biflora*, *Cardiospermum halicacabum*, *Casearia parvifolia*, *Cassia bicapsularis*, *Cassia glandulosa*, *Cassia occidentalis*, *Catopsis nutans*, *Celtis Iguanacea*, *Cenchrus tribuloides*, *Centrosema virginianum*, *Cestrum laurifolium*, *Chaptalia nutans*, *Cheilanthes microphylla*, *Chenopodium ambrosioides*, *Chiococca racemosa*, *Chloris paraguayensis*, *Chrysobalanus icaco*, *Cissus sicyoides*, *Citharexylum quadrangulare*, *Clerodendron aculeatum*, *Clusia alba*, *Coccoloba diversifolia*, *Coccoloba nivea*, *Coccoloba obtusifolia*, *Coccoloba uvifera*, *Coleus amboinicus*, *Commelina virginica*, *Comocladia ilicifolia*, *Cordia salvifolia*, *Cordia sulcata*, *Cordia tremula*, *Crescentia Cujete*, *Crinum americanum*, *Crotalaria incana*, *Crotalaria retusa*, *Croton flavens*, *Croton lobatus*, *Cucumis Anguria*, *Cuscuta americana*, *Cyperus rotundus*, *Dactyloctenium aegyptium*, *Daphnopsis caribaea*, *Desmanthus virgatus*, *Desmodium axillare*, *Desmodium spirale*, *Desmodium supinum*, *Desmodium uncinatum*, *Dichromena radicans*, *Drymaria cordata*, *Dryopteris opposita*, *Duranta repens*, *Eleusine indica*, *Epidendrum anceps*, *Epidendrum ciliare*, *Epidendrum papilionaceum*, *Eragrostis ciliaris*, *Eragrostis pilosa*, *Erechthites hieracifolia*, *Erechthites valerianaeifolia*, *Erigeron canadensis*, *Erigeron spathulatus*, *Eryngium*

foetidum, *Erythroxylum havanense*, *Eugenia axillaris*, *Eugenia Jambos*, *Eugenia procera*, *Eugenia uniflora*, *Eupatorium conyzoides*, *Eupatorium iresinoides*, *Eupatorium macrophyllum*, *Euphorbia hypericifolia*, *Euphorbia pilulifera*, *Euphorbia pulcherrima*, *Fagara martinicensis*, *Faramea odoratissima*, *Ficus Krugiana*, *Ficus omphalophora*, *Ficus populnea*, *Ficus Urbaniana*, *Fleurya aestuans*, *Galactia rubra*, *Gonzalagunia hirsuta*, *Gouania domingensis*, *Guettarda parvifolia*, *Guettarda scabra*, *Gynandropsis pentaphylla*, *Heliotropium parviflorum*, *Hemidiodia ocimifolia*, *Hymenaea Courbaril*, *Hymenocallis caribaea*, *Hyptis pectinata*, *Inga laurina*, *Ipomoea fastigiata*, *Ipomoea Nil*, *Indigofera suffruticosa*, *Iresine paniculata*, *Jacquemontia pentantha*, *Jatropha curcas*, *Jatropha gossypifolia*, *Justicia sessilis*, *Kallstroemia maxima*, *Lagascea mollis*, *Lantana Camara*, *Lantana involucrata*, *Leonotis nepetaefolia*, *Leonurus sibiricus*, *Lepidium virginicum*, *Leptochloa virgata*, *Linociera compacta*, *Lochnera rosea*, *Malpighia puniceifolia*, *Malvastrum spicatum*, *Mammea americana*, *Mariscus brizaeus*, *Mariscus capillaris*, *Melothria fluminensis*, *Miconia laevigata*, *Mitracarpus polycladus*, *Momordica charantia*, *Morisonia americana*, *Myrcia coriacea*, *Myrcia splendens*, *Nectandra coriacea*, *Nectandra Krugii*, *Neurogramme calomelanos*, *Ocimum micranthum*, *Oncidium spec.*, *Oplismenus setarius*, *Oxalis corniculata*, *Palicourea Pavetta*, *Panicum brevifolium*, *Panicum divaricatum*, *Panicum insulare*, *Panicum maximum*, *Panicum molle*, *Panicum sanguinale*, *Pappophorum alopecuroideum*, *Parthenium Hysterophorus*, *Paspalum conjugatum*, *Paspalum fimbriatum*, *Paspalum hemisphaericum*, *Passiflora laurifolia*, *Passiflora rubra*, *Passiflora suberosa*, *Pavonia spinifex*, *Pectis humifusa*, *Pedilanthus tithymaloides*, *Peperomia glabella*, *Peperomia petiolaris*, *Petiveria alliacea*, *Pharus glaber*, *Philodendron dispar*, *Philodendron oxycardium*, *Phoradendron trinervium*, *Phyllanthus niruri*, *Physalis pubescens*, *Picrasma antillana*, *Pilea microphylla*, *Pilea semidentata*, *Piper dilatatum*, *Piper reticulatum*, *Pisonia aculeata*, *Pisonia subcordata*, *Pitcairnia angustifolia*, *Pithecolobium unguis-cati*, *Plantago major*, *Pluchea purpurascens*, *Plumbago scandens*, *Plumiera alba*, *Polanisia viscosa*, *Polypodium aureum*, *Polypodium brasiliense*, *Polypodium lycopodioides*, *Polypodium phyllitidis*, *Polypodium piloselloides*, *Polypodium squamatum*, *Porophyllum ellipticum*, *Portulaca oleracea*, *Portulaca pilosa*, *Priva echinata*, *Psidium Guajava*, *Psychotria rufescens*, *Psychotria uliginosa*, *Pteris biaurita*, *Randia aculeata*, *Rauwolfia Lamarekii*, *Rhacoma crossopetalum*, *Rhynchosia reticulata*, *Rivina humilis*, *Ruellia tuberosa*, *Salvia occidentalis*, *Salvia serotina*, *Sauvagesia erecta*, *Schaefferia frutescens*, *Schoepfia*

Schreberi, *Scleria pterota*, *Scoparia dulcis*, *Sechium edule*, *Senecio sonchifolius*, *Setaria glauca*, *Setaria setosa*, *Sida carpinifolia*, *Sida cordifolia*, *Sida supina*, *Sideroxylon foetidissimum*, *Smilax guianensis*, *Solanum aculeatissimum*, *Solanum lanceaefolium*, *Solanum nigrum nodiflorum*, *Solanum racemosum igneum*, *Solanum torvum*, *Sonchus oleraceus*, *Spermacece tenuior*, *Spigelia Anthelmia*, *Sporobolus indicus*, *Stachytarpheta jamaicensis*, *Stenolobium stans*, *Stylosanthes hamata*, *Synedrella nodiflora*, *Tabernaemontana citrifolia*, *Talinum paniculatum*, *Talinum triangulare*, *Tamarindus indica*, *Tecoma leucoxylon*, *Tephrosia cathartica*, *Tetrazygia discolor*, *Thespesia populnea*, *Thunbergia alata*, *Tillandsia fasciculata*, *Tillandsia recurvata*, *Tillandsia usneoides*, *Tillandsia utriculata*, *Tournefortia foetidissima*, *Tragia volubilis*, *Trema Lamarckianum*, *Triumfetta semitriloba*, *Urechites suberecta*, *Urena lobata*, *Vernonia punctata*, *Villamilla octandra*, *Waltheria americana*, *Wedelia buphthalmoides*, *Wittmackia lingulata*, *Zebrina pendula*.

St. Martin.

Widely different from the former two Islands the vegetation of St. Martin is chiefly composed of a *Croton* vegetation much more developed than on St. Eustatius, whereas the tropical woods that are to be found on St. Eustatius and Saba are not to be met with on St. Martin.

The greater part of the Island being cultivated the ruderal plants are dispersed in almost the whole Island, along the roads and on the plantations and meadows, except in the typical *Croton* vegetation which on the outer parts of St. Martin is gradually changing into a well developed Littoral flora.

This Littoral flora together with the *Croton* vegetation commands the whole aspect of St. Martin.

Only where the hills are a little higher the *Eriodendron* vegetation like that one of the northern part of St. Eustatius is to be found.

The vegetation in the neighbourhood of the Saltponds with species that are not to be seen on the other Islands has also a typical aspect.

According to the results I got in studying the vegetation of St. Martin more minutely I dare say that we can also trace out here the following formations.

1. Vegetation of the tops of the hills.
2. Vegetation of the level parts of the Island and of the lower slopes of the hills.
3. Vegetation of the seashore and the rocky parts.
4. Vegetation of the cultivated region.

Referring to page 267 we could substitute for the first three divisions the same names I gave there, whereas the fourth division is almost synonym with: Crotonvegetation and in some cases also Eriodendronvegetation intermixed with, and partly replaced by ruderal plants.

1. Vegetation of the Tops of the Hills of St. Martin.

Abrus precatorius, *Acacia macracantha*, *Acacia riparia*, *Actinostemon caribaeum*, *Adelia ricinella*, *Ageratum conyzoides*, *Amarantus spinosus*, *Amomis Caryophyllata*, *Anona muricata*, *Anona squamosa*, *Anthurium cordatum*, *Argythamnia candicans*, *Aristolochia trilobata*, *Asclepias nivea*, *Asplenium cristatum*, *Asplenium pumilum*, *Beloperone eustachiana*, *Bernardia corensis*, *Bignonia unguis-cati*, *Blechnum occidentale*, *Brassavola cucullata*, *Bryophyllum pinnatum*, *Bunchosia nitida*, *Bursera simaruba*, *Callisia repens*, *Canella alba*, *Capparis Breynia*, *Capparis coccolobifolia*, *Capparis cynophallophora*, *Capparis jamaicensis*, *Capraria biflora*, *Capsium frutescens*, *Cardiospermum halicacabum*, *Casearia parvifolia*, *Casearia sylvestris*, *Cassia bicapsularis*, *Catopsis nutans*, *Cenchrus tribuloides*, *Chaptalia nutans*, *Chiococca racemosa*, *Chloris ciliata*, *Cissampelos Pareira*, *Cissus obovata*, *Cissus sicyoides*, *Clusia rosea*, *Coccoloba nivea*, *Commelina virginica*, *Comocladia ilicifolia*, *Corchorus hirsutus*, *Corchorus siliquosus*, *Crotalaria incana*, *Croton betulinus*, *Cyperus elegans*, *Dalechampia scandens*, *Daphnopsis caribaea*, *Desmodium supinum*, *Desmodium triflorum*, *Dorstenia contrajerva*, *Epidendrum anceps*, *Epidendrum ciliare*, *Erigeron spathulatus*, *Eryngium foetidum*, *Erythroxyllum havanense*, *Eugenia fragrans*, *Eugenia Jambos*, *Eugenia monticola*, *Eugenia procera*, *Eupatorium macrophyllum*, *Euphorbia petiolaris*, *Exostemma caribaeum*, *Ficus populnea*, *Ficus Urbaniana*, *Fleurya aestuans*, *Guettarda parvifolia*, *Guettarda scabra*, *Hemionitis palmata*, *Heteropteris lanceolata*, *Hymenocallis caribaea*, *Inga laurina*, *Krugiodendron ferreum*, *Kyllinga brevifolia*, *Lantana Camara*, *Lantana involucrata*, *Leonurus sibirica*, *Lepidium virginicum*, *Leucaena glauca*, *Loniciera compacta*, *Mammea americana*, *Mangifera indica*, *Maytenus ellipticus*, *Miconia laevigata*, *Myrcia paniculata*, *Myrcia splendens*, *Neurogramme calomelamos*, *Oncidium spec.*, *Oplis-*

menus setarius, *Oxalis corniculatus*, *Panicum diffusum*, *Panicum divaricatum*, *Panicum insulare*, *Panicum maximum*, *Panicum prostratum*, *Panicum sanguinale*, *Paspalum conjugatum*, *Passiflora laurifolia*, *Passiflora suberosa*, *Pedilanthus tithymaloides*, *Peperomia acuminata*, *Peperomia glabella*, *Peperomia obtusifolia*, *Pharus glaber*, *Phoradendron trinervium*, *Phyllanthus nobilis*, *Phytolacca icosandra*, *Pierammia pentandra*, *Picrasma antillana*, *Pilea microphylla*, *Piper dilatatum*, *Pisonia aculeata*, *Pisonia fragrans*, *Pisonia subcordata*, *Pithecolobium ungius-cati*, *Pluchea purpurascens*, *Polypodium aureum*, *Polypodium brasiliense*, *Polypodium lycopodioides*, *Polypodium phyllitidis*, *Polypodium piloselloides*, *Polypodium squamatum*, *Psidium Guajava*, *Psychotria horizontalis*, *Randia aculeata*, *Rhynchosia reticulata*, *Rivina humilis*, *Rochefortia acanthophora*, *Ruellia tuberosa*, *Samyda serrulata*, *Schaefferia frutescens*, *Senecio sonchifolius*, *Sida carpinifolia*, *Sida rhombifolia*, *Smilax ilicifolia*, *Solanum lanceaefolium*, *Solanum racemosum*, *Sonchus oleraceus*, *Spermacoce tenuior*, *Spigelia Anthelmia*, *Spiranthes elata*, *Sporobolus virginicus*, *Stenolobium staus*, *Synedrella nodiflora*, *Tabernaemontana citrifolia*, *Tecoma leucoxydon*, *Tillandsia fasciculata*, *Tillandsia recurvata*, *Tillandsia utriculata*, *Tournefortia fortidissima*, *Tournefortia hirsutissima*, *Tradescantia geniculata*, *Tragia volubilis*, *Triumfetta rhomboidea*, *Vernonia punctata*, *Wedelia buphthalmoides*, *Wittmackia lingulata*.

2. Vegetation of the Level Parts and of the Lower Parts of the Hills of St. Martin.

Abrus precatorius, *Abutilon indicum*, *Amarantus polygonoides*, *Amarantus tristis*, *Amyris elemifera*, *Anacardium occidentale*, *Anona muricata*, *Anona squamosa*, *Argythamnia candidans*, *Beloperone eustachiana*, *Beureria succulenta*, *Boerhaavea paniculata*, *Bunchosia nitida*, *Bursera simaruba*, *Calotropis procera*, *Calyptroanthus Boldinghii*, *Canella alba*, *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Cardiospermum halicacabum*, *Cassia bicapsularis*, *Cassia occidentalis*, *Cenchrus tribuloides*, *Centrosema virginianum*, *Chloris paraguayensis*, *Chrysobalanus icaco*, *Cissus sicyoides*, *Clusia rosea*, *Coccoloba uvifera*, *Commelina virginica*, *Comocladia ilicifolia*, *Conocarpus erectus*, *Crotalaria retusa*, *Croton astroites*, *Croton belulinus*, *Croton flavens*, *Cuscuta americana*, *Cyperus elegans*, *Cyperus rotundus*, *Desmanthus depressus*, *Ditaxis fasciculata*, *Dodonaea viscosa*, *Eleusine indica*, *Epidendrum papilionaceum*, *Eragrostis ciliaris*, *Eragrostis pilosa*, *Eragrostis tephrosanthos*, *Erigeron canadensis*,

Erithalis fruticosa, *Eugenia axillaris*, *Eugenia monticola*, *Euphorbia buxifolia*, *Euphorbia petiolaris*, *Euphorbia thymifolia*, *Evolvulus glaber*, *Evolvulus sericeus*, *Exostemma caribaeum*, *Fagara spinifex*, *Fagara trifoliata*, *Ficus populnea*, *Ficus Urbaniana*, *Gouania dominicensis*, *Guajacum officinale*, *Guettarda parvitolia*, *Gyminda latifolia*, *Heliotropium indicum*, *Heliotropium microphyllum*, *Hymenocallis caribaea*, *Ipomoea fastigiata*, *Jacquinia armillaris*, *Krugiodendron ferreum*, *Lantana Camara*, *Lantana involucrata*, *Lemna paucicostata*, *Leucaena glauca*, *Lochnera rosea*, *Malpighia angustifolia*, *Malpighia puniceifolia*, *Mangifera indica*, *Melochia nodiflora*, *Melochia tomentosa*, *Myrcia paniculata*, *Panicum diffusum*, *Panicum prostratum*, *Panicum sanguinale*, *Paspalum hemisphaericum*, *Passiflora suberosa*, *Pectis humifusa*, *Pedilanthus tithymaloides*, *Picramnia pentandra*, *Piscidia piscipula*, *Pisonia aculeata*, *Pisonia fragrans*, *Pisonia subcordata*, *Pithecolobium unguis-cati*, *Plumiera alba*, *Portulaca halimoides*, *Portulaca oleracea*, *Portulaca poliosperma*, *Portulaca quadrifida*, *Psidium Guayava*, *Randia aculeata*, *Rauwolfia Lamarkii*, *Rhacoma crossopetalum*, *Rhynchosia reticulata*, *Rochefortia acanthophora*, *Ruellia tuberosa*, *Salvia serotina*, *Samyda serrulata*, *Schaefferia frutescens*, *Sida carpinifolia*, *Sida ciliaris*, *Sida cordifolia*, *Sida supina*, *Smilax ilicifolia*, *Solanum racemosum*, *Solanum racemosum igneum*, *Solanum torvum*, *Sporobolus virginicus*, *Stachytarpheta jamaicensis*, *Stigmalophyllum periplocifolium*, *Stigmalophyllum sericeum*, *Stylosanthes hamata*, *Tamarindus indica*, *Tecoma leucoxydon*, *Ternstroemia peduncularis*, *Tillandsia recurvata*, *Tournefortia volubilis*, *Tragus racemosus*, *Tribulus cistoides*, *Vernonia punctata*, *Waltheria americana*.

3. Vegetation of the Seashore and Rocky Parts of St. Martin.

Acacia Farnesiana, *Alternanthera crucis*, *Antheophora hermaphrodita*, *Aristida americana*, *Avicennia nitida*, *Batis maritima*, *Borrichia arborescens*, *Bumelia obovata*, *Byrsonima crassifolia*, *Caesalpinia ciliata*, *Caesalpinia crista*, *Canavalia obtusifolia*, *Cassia obovata*, *Cassia polyadena*, *Clerodendron aculeatum*, *Coccoloba Krugii*, *Coccoloba uvifera*, *Conocarpus erectus*, *Ernodea littoralis*, *Euphorbia prostrata*, *Fimbristylis spathacea*, *Galactia dubia*, *Heleocharis capitata*, *Heliotropium curassavicum*, *Hippomane mancinella*, *Hypelate trifoliata*, *Ipomoea arenaria*, *Ipomoea pes caprae*, *Jacquinia Berteri*, *Juncellus laevigatus*, *Laguncularia racemosa*, *Lithophila muscoides*, *Lycium Americanum*, *Mariscus brizaeus*, *Melocactus communis*, *Metastelma parviflorum*, *Opuntia Ficus indica*, *Opuntia triacantha*, *Opuntia Tuna*, *Pappophorum alopecuroideum*, *Phyllanthus epiphyll-*

lanthus, *Plumbago scandens*, *Rhizophora Mangle*, *Salicornia herba-cea*, *Scaevola Plumieri*, *Sesbania sericea*, *Sesuvium portulacastrum*, *Suriana maritima*, *Tephrosia cinerea*, *Tournefortia gnaphalodes*, *Turnera ulmifolia*, *Urechites suberecta*, *Yucca aloifolia*.

4. Vegetation of the Cultivated Region of St. Martin.

Abrus precatorius, *Abutilon indicum*, *Acacia macracantha*, *Acacia riparia*, *Acalypha chamaedrifolia*, *Acalypha Poirerii*, *Acanthospermum humile*, *Achyranthes aspera*, *Adelia ricinella*, *Adiantum tenerum*, *Alternanthera repens*, *Alysicarpus nummularifolius*, *Amarantus polygonoides*, *Amarantus tristis*, *Ambrosia hispida*, *Anacardium occidentale*, *Anona muricata*, *Anona squamosa*, *Antigonon leptopus*, *Ayenia pusilla*, *Beloperone eustachiana*, *Beureria succulenta*, *Bidens bipinnatus*, *Blechnum occidentale*, *Blechnum Brownei*, *Boerhaavea erecta*, *Boerhaavea hirsuta*, *Boerhaavea paniculata*, *Boerhaavea scandens*, *Brassavola cucullata*, *Bromelia pinguin*, *Bryophyllum pin-natum*, *Bursera simaruba*, *Caesalpinia coriaria*, *Caesalpinia pulcherrima*, *Callisia repens*, *Calotropis procera*, *Capparis Breynia*, *Capparis coccolobifolia*, *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Capraria biflora*, *Capsicum frutescens*, *Cardiospermum halicacabum*, *Casearia parvifolia*, *Cassia bicapsularis*, *Cassia glandulosa*, *Cassia occidentalis*, *Catopsis nutans*, *Ceiba pentandra*, *Celosia nitida*, *Celtis Iguanacea*, *Cenchrus tribuloides*, *Centrosema virginianum*, *Chenopodium ambrosioides*, *Chenopodium murale*, *Chiococca racemosa*, *Chloris paraguayensis*, *Chrysobalanus icaco*, *Cissus sicyoides*, *Citharexylum quadrangulare*, *Coccoloba nivea*, *Coccoloba uvifera*, *Coleus amboinicus*, *Commelina virginica*, *Comocladia ilici-folia*, *Corchorus hirsutus*, *Cracca caribaea*, *Crotalaria incana*, *Crotalaria retusa*, *Crotalaria verrucosa*, *Croton astroites*, *Croton betulinus*, *Croton flavens*, *Croton lobatus*, *Cuscuta americana*, *Cyperus elegans*, *Cyperus rotundus*, *Dactyloctenium aegyptium*, *Datura fastuosa*, *Datura Metel*, *Desmanthus depressus*, *Desmanthus virgatus*, *Desmodium spirale*, *Desmodium supinum*, *Ditaxis fasciculata*, *Duranta repens*, *Eleusine indica*, *Eragrostis ciliaris*, *Eragrostis tephrosanthos*, *Erigeron canadensis*, *Erithalis fruticosa*, *Erythroxylon brevipes*, *Erythrina corallodendron*, *Eugenia axillaris*, *Eugenia monticola*, *Eugenia procera*, *Eupatorium conyzoides*, *Euphorbia buxifolia*, *Euphorbia heterophylla*, *Euphorbia hypericifolia*, *Euphorbia petiolaris*, *Euphorbia pilulifera*, *Evolvulus glaber*, *Evolvulus sericeus*, *Exostemma caribaeum*, *Fagara martinicensis*, *Fagara spinifex*, *Fagara trifoliata*, *Ficus populnea*, *Galactia rubra*, *Gouania domingensis*, *Guettarda parvi-*

folia, *Gyminda latifolia*, *Gynandropsis pentaphylla*, *Heliotropium indicum*, *Heliotropium parviflorum*, *Hemidioda ocymifolia*, *Hippomane mancinella*, *Hura crepitans*, *Hymenaea Courbaril*, *Hyptis pectinata*, *Ipomoea dissecta*, *Ipomoea fastigiata*, *Indigofera suffruticosa*, *Iresine elatior*, *Jacquinia armillaris*, *Jatropha curcas*, *Jatropha gossypifolia*, *Justicia sessilis*, *Kallstroemia maxima*, *Krugiodendron ferreum*, *Kyllingia brevifolia*, *Lagascea mollis*, *Lantana Camara*, *Lantana involucrata*, *Leonotis nepetaefolia*, *Leonurus sibiricus*, *Lepidium virginicum*, *Leucaena glauca*, *Leucas martinicensis*, *Lippia reptans*, *Lithophila muscoides*, *Lochnera rosea*, *Malpighia augustifolia*, *Malpighia puniceifolia*, *Malvastrum spicatum*, *Mangifera indica*, *Mariscus capillaris*, *Melia azedarach*, *Melicocca bijuga*, *Melochia tomentosa*, *Microtea debilis*, *Moringa oleifera*, *Morisonia americana*, *Murraya exotica*, *Neptunia plena*, *Ocimum micranthum*, *Oldenlandia herbacea*, *Oplismenus setarius*, *Panicum colonum*, *Panicum diffusum*, *Panicum geminatum*, *Panicum insulare*, *Panicum prostratum*, *Panicum sanguinale*, *Parkinsonia aculeata*, *Parthenium Hysterophorus*, *Paspalum hemisphaericum*, *Pavonia spinifex*, *Pectis humifusa*, *Pedilanthus tithymaloides*, *Peperomia Langsdorffii*, *Persea americana*, *Petiveria alliacea*, *Phloxeris vermicularis*, *Phyllanthus niruri*, *Physalis angulata*, *Picramnia pentandra*, *Piscidia piscipula*, *Pisonia aculeata*, *Pisonia fragrans*, *Pisonia subcordata*, *Pithecolobium unguis cati*, *Plumiera alba*, *Plumiera rubra*, *Polanisia viscosa*, *Portulaca halimoides*, *Portulaca oleracea*, *Priva echinata*, *Psidium Guajava*, *Psychotria horizontalis*, *Randia aculeata*, *Rauwolfia Lamarekii*, *Rhacoma crossopetalum*, *Rhynchosia reticulata*, *Rivina humilis*, *Ruellia tuberosa*, *Salvia serotina*, *Samyda serrulata*, *Schaefferia frutescens*, *Setaria setosa*, *Setaria viridis*, *Sida carpinifolia*, *Sida ciliaris*, *Sida cordifolia*, *Sida supina*, *Solanum nigrum nodiflorum*, *Solanum racemosum igneum*, *Solanum torvum*, *Spermacoce tenuior*, *Sporobolus indicus*, *Sporobolus virginicus*, *Stachytarpheta jamaicensis*, *Stenandrum rupestre*, *Stenolobium stans*, *Stigmatophyllum periplocifolium*, *Stylosanthes hamata*, *Talinum triangulare*, *Tamarindus indica*, *Tecoma leucoxydon*, *Teramnus labialis*, *Thespesia populnea*, *Thunbergia fragrans*, *Tillandsia recurvata*, *Tournefortia foetidissima*, *Tournefortia volubilis*, *Tragus racemosus*, *Tribulus cistoides*, *Triumfetta rhomboidea*, *Vernonia punctata*, *Villanilla octandra*, *Waltheria americana*, *Wedelia buphthalmoides*, *Zizyphus Jujuba*.

CHAPTER II

DESCRIPTION OF THE DIFFERENT TYPES OF THE VEGETATION.

Introduction.

In chapter I of the Third Part I divided the vegetation of the Dutch West Indian Islands into three regions: 1. the higher parts, 2. the level plains and the lower part of the slope of the hills, 3. the slope of the hills joining the seashore together with the seashore. Whereas I refer for further details to the Geographical Part I draw the attention to the fact that on St. Eustatius the three regions are well developed. In Saba, however there is almost no level plain to be seen, whereas in St. Martin there is but a very small part of the Island that belongs to the first region.

There is a close connection between the above given data and the development of the three typical forms of the vegetation.

1. ERIODENDRONVEGETATION.

2. CROTONVEGETATION.

3. LITTORALVEGETATION.

The first two names are taken from *Eggers* Flora of St. Croix and the Virgin Islands.

ERIODENDRONVEGETATION.

This type of vegetation is best characterised by the following trees: *Capparis cynophallophora*, *Capparis frondosa*, *Capparis jamaicensis*, *Ardisia coriacea*, *Rauwolfia Lamarkiana*, *Linociera compacta*, *Pisonia fragrans*, *Daphnopsis caribaea*, *Coccoloba diversifolia*, *Clusia alba*, *Ternstroemia peduncularis*, *Nectandra coriacea*, *Inga laurina*, *Picrasma antillana*, *Ficus populnea*, *Ficus Urbaniana*, *Ficus Krugiana*, *Allophylus occidentalis*, whereas *Eugenia axillaris*, *Eugenia cordata*, *Eugenia ligustrina*, *Myrcia splendens*, *Palicourea Pavetta*, *Psychotria horizontalis*, *Psychotria rufescens*, *Chiococca*

racemosa, *Samyda serrulata*, *Casearia parvifolia*, *Byrsonima spicata*, *Erythroxylon havanense*, *Vernona punctata*, are the most predominant shrubs. In very humid localities the soil is covered by *Peperomia glabella*, *Peperomia obtusifolia* and *Pilea semidentata*.

St. Eustatius.

The *Eriodendron* vegetation is well developed on the higher parts of the Quill and on those parts of the northern hilly country that are shut in between the higher tops, viz the ravines.

Quill. The higher parts of the Quill are covered by a forest of evergreen plants and interspersed between these evergreens there are some species with deciduous leaves; all the plants named on pag. 282 and 283 as typical for the *Eriodendron* vegetation are to be seen here.

The lower parts of this region show some plants that occur also in the *Croton* vegetation viz. *Acacia*, *Croton*, *Lantana*, *Solanum*. Shrubs and smaller trees: *Chrysobalanus icaco*, *Symplocos martinicensis*, *Rauwolfia Lamarckiana*, *Tabernaemontana citrifolia*, *Lonchocarpus violacea*, *Casearia parvifolia*, are forming a dense mixture and gradually pass into the forest of the higher tops; there we see a real tropical rainforest with its typical features. The trees that dominate on this top are: *Linociera compacta*, *Neetandra coriacea*, *Inga laurina*, *Myrcia paniculata*, *Myrcia splendens*, *Psychotria rufescens*, *Eugenia ligustrina*, *Micoma laevigata*, *Byrsonima spicata*, *Clusia alba*, *Ceiba pentandra*, *Pisonia fragans*, *Coccoloba diversifolia*, *Ardisia guadalupensis*, *Daphnopsis carabaea*, *Bursera simaruba*, *Ficus Krugiana*, *Cestrum laurifolium*, *Allophylus occidentalis*, *Tecoma leucoxyton*, *Celtis Iguanacea*, *Casearia parvifolia*, *Persea gratissima*, *Ternstroemia peduncularis*, *Tabernaemontana citrifolia*, *Guettarda scabra*, *Ficus laevigata*. On humid localities we see the following plants: *Peperomia glabella*, *Pilea semidentata*, *Anthurium oxycardium*. Especially the eastern part of the Top of the Quill is very humid and there *Clusia alba*, *Linociera compacta* and *Myrtaceae* are covered with *Tillandsia recurvata*. The inner slope of the Quill has the same plants, which occur on the outer slope, but moreover the following plants are to be seen here: *Ceiba pentandra*, *Piper reticulatum* and *Piper dilatatum*, *Urera Caracasana*, *Pharus glaber*, *Panicum divaricatum*, *Oplismenus setarius*, *Smilax coriacea*, *Stenochlaena sorbifolia*, *Cyathea serra*, *Begonia domingensis*, *Phoebe elongata*, *Hyperbaena dominicensis*, *Pisonia aculeata*, *Psychotria uliginosa*, *Palicourea Pavetta*,

Heliconia Bihai, *Asplenium obtusifolium*, *Asplenium cristatum*, *Asplenium serratum*, *Stenochlaena sorbifolia*, *Polypodium Phyllitidis*, *Polypodium lycopodioides*, *Dryopteris parasitica*, whereas *Boehmeria ramiflora*, *Peperomia scandens* together with *Araceae* cover humid rocks. There occur in the Quill also *Coffea arabica*, *Theobroma Cacao*, *Spondias lutea* and *Mammea americana*.

Northern hilly country. In the northern hilly country the *Eriodendron* vegetation is to be found in the ravines and other parts which are not exposed to the heavy winds and where a certain amount of humus is covering the rocks.

It may easily be recognised here by the great amount of smaller trees and shrubs mentioned on page 282 and 283, whereas the higher trees that are to be seen in the Quill are almost absent. Moreover there is a great quantity of trees, which certainly are not wild here, intermixed with the lower shrubs viz: *Tamarindus indica*, *Melicocca bijuga*, *Moringa oleifera*, *Spondias purpurea*, *Persea americana*.

The species that predominate here in the northern *Eriodendron* vegetation are: *Rhacoma crossopetalum*, *Eugenia ligustrina*, *Eugenia procera*, *Samyda serrulata*, *Eugenia axillaris*, *Trema Lamarckiana*, *Pithecolobium unguis-catis*, *Celtis Iguanacea*, *Guettarda parvifolia*, *Schaefferia frutescens*, *Krugiodendron ferreum*, *Chiococca racemosa*, *Randia aculeata*, *Erythroxylon brevipes*, *Erythroxylon havanense*, *Pisonia aculeata*, *Guettarda scabra*, *Bumelia obovata*, *Anona muricata*, *Anona squamosa*, *Rauwolfia Lamarckiana*.

Here in the northern hilly country no sharp difference can be traced between the *Eriodendron* vegetation and the *Croton* vegetation, nor between the latter and the Vegetation of the rocky slopes of the hills.

Where this *Croton* vegetation begins to change into the *Eriodendron* vegetation we see the following plants: *Lantana Camara*, *Lantana involucrata*, *Croton flavens*, *Croton astroites*, *Bernardia corensis*, *Wedelia buphthalmoides*, *Melochia tomentosa*, *Fourcroya gigantea*, *Vernonia punctata*, *Stylosanthes hamata*, *Portulaca oleracea*, *Panicum divaricatum*, *Melocactus communis*, *Tragus racemosus*, *Ruellia tuberosa*, *Pectis humifusa*, *Desmanthus virgatus*, *Sida ciliaris*, *Justicia sessilis*, *Commelina virginica*, *Setaria setosa*, *Jatropha gossypifolia*, *Opuntia triacantha*, *Cordia sulcata*, *Gouania domingensis*, *Cissus sicyoides*, *Rhynchosia reticulata*, *Stigmatophyllon periplo-cifolium*, *Tillandsia utriculata*, *Phoradendron trinervium*.

There are very few herbaceous plants; only in humid localities some species of *Peperomia* are to be seen.

Saba.

On Saba the Eriodendronvegetation is well developed in the guts and those parts of the lower slopes of the hills which are not used for cultivation. The plants typical for this vegetation are *Eugenia axillaris*, *Pisonia obtusata*, *Anona muricata*, *Stenolobium Stans*, *Lantana Camara*, *Tournefortia foetidissima*, *Chiococca racemosa*, *Trema Lamarckiana*, *Pithecolobium unguis-cati*, *Allophylus occidentalis*, *Nectandra Krugii*, *Schoepfia Schreberi*, *Capparis cynophallophora*, *Casearia parvifolia*, *Hyperbaena domingensis*, *Pisonia aculeata*, *Celtis Iguanacea*, *Symplocos martinicensis*, *Coccoloba diversifolia*, *Myrcia coriacea*, *Tabernaemontana citrifolia*, *Coccoloba nivea*, *Capparis frondosa*, *Chrysobalanus icaco*, *Inga laurina*, *Erythroxylon havanense*, *Picrasma antillana*, *Daphnopsis caribaea*, *Amomis caryophyllata*, *Clusia alba*, *Miconia laevigata*, *Ardisia coriacea*, *Citharexylum quadrangulare*, *Cestrum laurifolium*, *Tecoma leucoxylon*, *Psychotria rufescens*.

It must however be understood that there are but few parts of Saba where the Eriodendronvegetation has the same feature as that of the northern hilly country of St. Eustatius. Whereas the shrubs of the latter contain some Myrtaceae, these plants do not occur in Saba in a great quantity; the trees that give a typical aspect to the Quill vegetation viz. *Linociera compacta*, *Nectandra coriacea*, do not occur on Saba except sporadically.

There is however another part of the Eriodendronvegetation in Saba covering the higher tops of the Mountain, where the typical plants of a real tropical rainforest occur.

The greater part of the plants we meet in this part of Saba are not known from St. Eustatius and St. Martin and chiefly occur in Guadeloupe and other Antilles with high mountains. The feature of this rainforest is given to it by the Treeferns: *Hemitelia horrida*, *Alsophila phalerata*, *Cyathea serra*, *Cyathea arborea* together with *Euterpe*.

On the most humid localities occur: *Psychotria pendula*, *Hillia parasitica*, *Marcgravia umbellata*, *Heliconia Bihai*, *Fagara martinicensis*, *Nectandra Krugii*, *Psychotria rufescens*, *Besleria lutea*, *Gesneria ventricosa*, *Charianthus crinitus*, *Peperomia emarginella*, *Peperomia urocarpa*, *Peperomia hernandifolia*, *Hymenophyllum hirtellum*, *Trichomanes alatum*, *Polypodium trichomanoides*, *Utricularia alpina*, *Ornithidium coccineum*. The soil is densely covered by: *Dryopteris parasitica*, *Dryopteris reticulata*, *Aspidium Plumierii*, *Aspidium trifoliatum*, *Polybotrya cervina*, *Neptrolepis exal-*

tata, *Diplazium striatum*, *Blechum occidentale*, *Stenochlaena sorbifolia*, *Neurogramme calomelanos*, *Lonchitis hirsuta*, *Polypodium brasiliense*, *Polypodium crassifolium*, *Polypodium loricum*, *Polypodium piloselloides*, *Elaphoglossum conforme*, *Elaphoglossum latifolium*, *Gleichenia bifida*.

St. Martin.

The well developed *Eriodendron* vegetation of Saba and St. Eustatius with its typical plants is not so pronounced in St. Martin. Small remnants of an in former times well developed *Eriodendron* vegetation are to be found on the tops of the Milldrumhill, Mount Paradis, Nakedboyhill, Centryhill and on the higher parts of the road from Filipsburg to Marigot. The following plants make it sure that these parts are covered by the *Eriodendron* vegetation: *Panicum divaricatum*, *Peperomia obtusifolia*, *Peperomia glabella*, *Polypodium aureum*, *Daphnopsis caribaea*, *Miconia laevigata*, *Tabernaemontana citrifolia*, *Samyda serrulata*, *Inga laurina*.

Moreover there are to be seen the following plants: *Guettarda scabra*, *Myrcia coriacea*, *Casearia parvifolia*, *Pisonia fragrans*, *Peperomia Langsdorffii*. Especially typical for St. Martin are the many individuals that may be found of *Samyda serrulata*, *Coccoloba nivea*, *Pisonia fragrans*, *Comocladia ilicifolia*, *Schaefferia frutescens* and of the following ones unknown from St. Eustatius and Saba: *Picramnia pentandra*, *Bunchosia nitida*, *Casearia sylvestris*, *Eugenia monticola*, *Tournefortia hirsutissima*.

Coccoloba diversifolia, *Allophylus occidentalis*, *Ardisia coriacea*, *Linociera compacta*, *Cestrum laurifolium* do not occur on St. Martin. Where the *Croton* vegetation comes in connection with the *Eriodendron* vegetation, especially in the northern parts of St. Martin we find another number of plants not known on the other Dutch West Indian Islands viz. *Malpighia augustifolia*, *Bunchosia nitida*, whereas there occur also in great quantity, *Samyda serrulata*, *Pisonia subcordata*, *Capparis frondosa* and *Capparis jamaicensis*.

CROTONVEGETATION.

The *Croton* vegetation is characterised by *Croton*, *Acacia*, *Lantana*, *Solanum racemosum*, *Melochia tomentosa*, *Opuntia triacantha*, *Opuntia Tuna*, *Opuntia Ficus indica* and *Melocactus*.

St. Eustatius.

The level country and the lower parts of the slopes of the hills

which are not used by man for agricultural purposes, are covered with the Crotonvegetation. Moreover this vegetation is to be seen also in the northern hilly country as well on the slopes of the hills as on the higher localities viz. Boven.

Especially in the hilly country there is not a great difference between a poorly developed Crotonvegetation and that of the rocky slopes. The Crotonvegetation on St. Eustatius has not the same aspect over the whole Island.

Where Croton flavens is predominant and but a few trees like Pisonia subcordata are growing among the lower shrubs, the Crotonvegetation has a very monotonous aspect and it is not very difficult to walk through it; we find this form of the Crotonvegetation near Boven.

In the lower parts of St. Eustatius the Croton is much more intermixed with other plants such as *Acacia tortuosa*, *Solanum racemosum* and *igneum*, *Pisonia subcordata*, *Tecoma leucoxydon*, *Crescentia Cujete*, *Ceiba pentandra*, *Leucaena glauca*, *Plumbago scandens*, *Randia aculeata*, *Jatropha gossypifolia*, *Lochnera rosea*, *Cassia bicapsularis* and other ruderalplants. From the Cactaceae there are to be seen *Opuntia Tuna*, *Opuntia Ficus Indica*, *Opuntia triacantha* and *Melocactus communis*.

Moreover there is a great quantity of *Lantana Camara*, *Lantana involucrata*, *Waltheria americana*, *Melochia tomentosa*, *Stigmatophylon periplocifolium* and *Tournefortia volubilis*. This type of the Crotonvegetation, is to be seen especially near Bengalen and between Oranjestad and Fort de Wind.

Still another feature of the Crotonvegetation is to be mentioned where *Krameria ixina* is predominant; this typical Crotonvegetation is only to be seen near Fort de Wind; here the plants are much smaller and the soil is covered by ruderalplants.

Saba.

The Crotonvegetation of Saba is not well developed; from the plants typical for this vegetation on St. Eustatius *Acacia Farnesiana* is not to be seen in Saba, whereas the other plants although they occur in Saba are nowhere in such quantities as to predominate and to form a real Crotonvegetation.

St. Martin.

The level part of St. Martin as far as it is not used for agricultural purposes and the slopes of the surrounding hills are covered with the Crotonvegetation.

This Crotonvegetation has another aspect as the one of St. Eustatius, owing to different other plants that are intermixed with the St. Eustatiusplants viz. *Croton betulinus*, *Euphorbia petiolaris*, *Plumiera alba*, *Fagara spinifex*, *Fagara trifoliata*, *Ditaxis fasciculata*, *Argithamnia candicans*, *Adelia ricinella*, *Comocladia ilicifolia*, *Melochia tomentosa*, *Phyllanthus epyphyllanthus*, *Coccoloba Krugii*, *Jacquinia Berteri*.

Here also the Crotonvegetation shows some likeness with the Eriodendronvegetation and especially in the Lowlands and near Orientbay we see the Crotonvegetation gradually intermixed with or replaced by the Eriodendronvegetation; there we find the following plants: *Eugenia axillaris*, *Capparis frondosa*, *Chiococca racemosa*, *Picramnia pentandra*, *Capparis cynophallophora*, *Pithecolobium ungius-cati*, *Comocladia ilicifolia*, *Casearia parvifolia*, *Samyda serrulata*, *Croton flavens*, *Melochia tomentosa*, *Plumiera alba*, *Acacia Farnesia*, *Lantana Camara*, *Lantana involucrata*, *Argithania candicans*, *Rauwolfia Lamarkiana*, *Clerodendron aculeatum*, *Comocladia ilicifolia*, *Capparis jamaicensis*, *Byrsonima lucida*, *Eugenia axillaris*, *Myrcia coriacea*, *Calypttranthes Boldinghii*, *Terntroemia peduncularis*, *Chrysobalanus Icaco*, *Eugenia ligustrina*, *Ficus laevigata*, *Clusia alba*, *Psidium Guajava*, *Randia aculeata*, *Pisonia subcordata*, *Bursera simaruba*, *Euphorbia petiolaris*, *Cassia polyadena*, *Paspalum paniculatum*, *Stigmatophyllon sericeum*, *Croton betulinus*, *Ruellia tuberosa*, *Cassia occidentalis*, *Stachytarpheta jamaicensis*, *Melocactus communis*.

LITTORAL VEGETATION.

The Littoralvegetation may easely be divided into:

1. **Vegetation of the beach.**
2. **Vegetation of the rocky slopes near the seashore.**
3. **Vegetation of the lagoons and the saltponds.**

I. Vegetation of the Beach.

This vegetation is characterised by:

Hippomane mancinella, *Erithalis fruticosa*, *Exostemma caribaeum*, *Suriana maritima*, *Canella alba*, *Tournefortia gnaphalodes*, *Scaevola Plumieri*, *Coccoloba uvifera*, *Ipomoea pescaprae*, *Heliotropium curasavicum*, *Sporobolus virginicus*.

St. Eustatius.

This vegetation is only to be seen at the different Bays.

The trees which mostly occur are *Hippomane mancinella* and *Coccoloba uvifera*. The smaller plants are *Tournefortia gnaphalodes*, *Suriana maritima* and *Clerodendron aculeatum*, *Ipomoea pes-caprae*, *Canavalia obtusifolia*, *Sporobolus virginicus*, *Heliotropium curassavicum* and *Lithophila muscoides*.

On localities where the ruderal plants have also invaded into the seashore we see the following plants: *Caesalpinia ciliata*, *Jatropha gossypifolia*, *Lantana Camara*, *Acacia macracantha*, *Plumbago scandens* and *Solanum racemosum igneum*. Where the seashore does not consist of sand but only of stones the vegetation is still less developed and we only find *Portulaca halimoides*, *Euphorbia buxifolia*, *Tephrosia cinerea* and *Borreria podocephala*.

Saba.

The seashore and accordingly also the Seashorevegetation is almost not developed on Saba: only at Flatpoint there is a small beach and the plants to be found there are: *Lithophila muscoides*, *Sesuvium portulacastrum*, *Stylosanthes hamata* and *Turnera ulmifolia*.

St. Martin.

Here the beach is very well developed along the southern part of the Island near Grande Case and near Orientbay; the typical plants to be met with are: *Erithalis fruticosa*, *Exostemma caribaeum*, *Canella alba*, *Borrchia argentea*, *Jacquinia armillaris*, *Suriana maritima*, *Chrysobalanus icaco*, *Conocarpus erectus*, *Strumpfia maritima*, *Jacquinia Berteri*, *Phyllanthus epiphyllanthus*, *Caesalpinia ciliata*, *Caesalpinia crista*, *Tournefortia gnaphalodes*, *Scaevola Plumierii*, *Bumelia obovata*, *Coccoloba uvifera*, *Ernodea litoralis*, *Ipomoea pes-caprae*, *Heliotropium curassavicum*, *Sporobolus virginicus*.

II. Vegetation of the Rocky Slopes near the Seashore.

I refer to the Geographical and Orological Section where may be seen where these rocks are to be found.

The plants typical for this vegetation are: *Coccoloba uvifera*, *Jacquinia armillaris*, *Tecoma leucoxydon*, *Rhacoma crossopetalum*, *Conocarpus erectus*, *Erythalis fruticosa*, *Melocactus communis*, *Pe-*

dilanthus tithymaloides, *Dactyloctenium aegyptium*, *Aristida americana*.

St. Eustatius.

This vegetation is formed here by the following plants: *Pedilanthus tithymaloides*, *Fourcroya gigantea*, *Opuntia spinosissima*, *Lochnera rosea*, *Croton flavens*, *Jacquinia armillaris*, *Erythalis fruticosa*, *Melocactus communis*, *Opuntia triacantha*, *Opuntia Tuna*, *Opuntia Ficus indica*, *Jacquinia armillaris*, *Conocarpus erecta*, *Acacia macracantha*, *Tecoma leucoxylon*, *Dactyloctenium aegyptium*, *Pappophorum alopecuroides*, *Euphorbia articulata*, *Eragrostis ciliaris*, *Aristida americana*, *Pectis humusa*. *Ernodea littoralis* and *Strumpfia maritima* are to be seen on the White Wall only.

Saba and St. Martin.

Whereas the plants occurring on the rocky parts of these Islands do not differ from those on St. Eustatius this vegetation needs no further description.

I will only mention that *Jacquinia armillaris* has not been seen on Saba. The following summary of the plants found on the small Islands Green Island, Green Key and Tintamarre may give an idea of the vegetation there.

On Green Island I collected the following plants:

Euphorbia articulata, *Coccoloba uvifera*, *Tecoma leucoxylon*, *Mariscus brizaeus*, *Opuntia Funa*.

On Green Key I collected the following plants: *Euphorbia buxifolia*, *Portulaca halimoides*, *Lantana involucrata*, *Croton flavens*, *Acacia Farnesiana*, *Beloperone eustachiana*, *Lithophila muscoides*, *Tournefortia gnaphalodes*, *Batis maritima*, *Hippomane mancinella*, *Opuntia Ficus indica*, *Melocactus communis*, *Coccoloba uvifera*. The vegetation is a mixture of Croton- and Littoralvegetation.

The Island of Tintamarre that partly is used for cultivation of Cotton and consists also out of limestone I saw the following plants: *Mariscus brizaeus*, *Guajacum officinale*, *Evolvulus argyraeus*, *Heliotropium microphyllum*, *Borrchia arborescens*, *Jacquemontia pentantha*, *Desmanthus depressus*, *Fagara flava*, *Panicum prostratum*, *Beureria succulenta*, *Tournefortia volubilis*, *Plumiera alba*, *Pedilanthus tithymaloides*, *Bignonia unguis-cati*, *Hippomane mancinella*, *Rauwolfia Lamarekii*, *Melochia tomentosa*, *Croton flavens*, *Croton betulinus*, *Amarantus polygonoides*, *Cenchrus tribuloides*, *Pappophorum alopecuroides*, *Sesuvium portulacastrum*, *Erithalis fruti-*

cosa, *Myrcia splendens*, *Rhacoma crossopetalum*, *Stigmatophyllon periplocifolium*, *Argithamnia caudicans*, *Turnera ulmifolia*.

On this Island as may be seen from the above mentioned species there is a typical Littoral- and Crotonvegetation which wholly resembles the one on St. Martin.

III. Vegetation of the Lagoons and the Saltponds.

This Vegetation is to be seen on St. Martin only; the typical plants are *Rhizophora* Mangle, *Laguncularia racemosa*, *Avicennia nitida*. Accordingly we find here a Mangrovevegetation whereas in the neighbourhood of the Saltponds are to be seen *Batis maritima*, *Pycereus flavescens*, *Eleocharis capitata* together with other small plants of the beach.

VEGETATION OF THE CULTIVATED REGION.

This vegetation needs no further explanation as it is of no phytogeographical interest; for further details I refer to page 270, 274, 275, 280 and 281.

From the data given by *Feenstra* in his: *De Nederlandsche West-Indische eilanden* etc. as well as from the data gathered by myself when collecting the plants on the different parts of the Islands it is evident that in former time a much greater part of St. Eustatius was in cultivation; I refer also to the Rapport of *Went*; not only the northern hilly country but also the inner part of the Quill shows some plants that certainly were introduced there by man.

The greater part of the localities of Saba that are apt for cultivation is used for that purpose whereas on St. Martin the greater part of the level plains and the lower slopes of the hills are in cultivation; the top of Mount Paradis certainly was used in former times for the same purposes.

The well developed meadows in St. Martin take up a great part of this Island.

LITERATURE.

- Boldingh = *I Boldingh*: A Contribution to the Knowledge of the Flora of Anguilla (B. W. I) Rec. d. Trav. Botan. Néerlandais, vol. VI (1909) p. 1—36.
- I. Boldingh*: Lijst van Planten, die door de Bewoners van de drie Nederlandsche Antillen St. Eustatius, Saba en St. Martin als geneeskrachtig worden beschouwd. Bull. Kolon. Mus. Haarlem n. 38 (1907) p. 93—112.
- Brand 1901 = *A. Brand*: Symplocaceae in *Engler*: Regnum vegetabile IV, 242, 1901.
- Christensen 1906 = *C. Christensen*: Index Filicum 1906.
- Cogniaux 1881 = *A. Cogniaux*: Cucurbitaceae in *De Candolle Monographiae Phanerogamarum*, vol. III 1881 p. 325—954.
- Cogniaux 1891 = *A. Cogniaux*: Melastomaceae in *De Candolle Monographiae Phanerogamarum*, vol. VII, 1891.
- Duss* = *R. P. Duss*: Flore phanérogamique des Antilles françaises (Guadeloupe et Martinique). Avec annotations du professeur *Dr. Edouard Heckel* sur l'emploi de ces plantes. [*Heck. Ann. de l'inst. colon. de Marseille*, 4-ème année 1896, vol. III]. Macon 1897.
- H. F. A. Eggers*: Flora of St. Croix and the Virgin Islands, West Indies. (Bull. of the United States National Museum no. 13. Washington 1870, 133 p. 8^o).
- E. a. P. = *A. Engler und K. Prantl*: Die natürlichen Pflanzenfamilien, up to April 1909.
- Engler 1883 = *A. Engler*: Burseraceae et Anacardiaceae in *De Candolle Monographiae Phanerogamarum*, vol. IV, 1883.
- Engler 1898 = *A. Engler*: Araceae in *De Candolle Monographiae Phanerogamarum*, vol. II, 1889.
- B. A. Euphrasen*: Beskrifning öfver Svenska vestindiska ör St. Barthélémi samt öarne St. Eustache och St. Christopher. Stockholm 1759.
- M. D. Feenstra*: De Nederlandsche West-indische eilanden in derzelver tegenwoordigen toestand, Amsterdam 1836—1837.
- A. H. R. Grisebach*: Systematische Untersuchungen über die Vegetation der Karaiben, insbesondere der Insel Guadeloupe. [Abh. Kgl. Gesellsch. Wiss. Göttingen. VII (1857) p. 151—286].
- Gris. Fl. = *A. H. R. Grisebach*: Flora of the British West Indian

- Islands. London. p. 1—192: 1859; p. 193—315: 1860; p. 315—506: 1861; p. 507—789, Tit. et Index: 1864.
- A. H. R. Grisebach: Die geographische Verbreitung der Pflanzen Westindiens. [Abh. Kgl. Gesellsch. Wiss. Göttingen, vol. XII (1865) p. 3—80] — *Gesamm. Abhandl.* Leipzig 1880. 8°, p. 222—285.
- Hackel 1889 = *E. Hackel*: *Andropogoneae* in *De Candolle Monographiae Phanerogamarum*, vol. VI, 1889.
- Hemsley 1884 = *W. Botting Hemsley*: Report on the Botany of the Bermudas and various other Islands of the Atlantic and Southern Oceans. The Bermudas. [Rep. on the scient. results of the Voy of H. M. S. Challenger. London, 4°. Botany vol. I part I (1884) p. 1—135, tab. I—XIII et introduction (1885) p. 48—49.
- Hitchc. = *Albert S. Hitchcock*: List of plants collected in the Bahamas, Jamaica and Grand Cayman. [Fourth annual rep. of the Missouri Bot. Gard. vol. IV (Mart. 1893) p. 47—179, tab. XI—XIV. 8°].
- Ind. Kew. = *Index Kewensis Plantarum Phanerogamarum* by *B. D. Jackson* 1893—1905.
- Nicolaus Joseph Jacquin*: *Selectarum stirpium americanarum historia*, in qua ad Linnaeanum systema determinatae descriptaeque sistuntur plantae illae, quas in insulis Martinica, Jamaica, Domingo aliisque et in vicinae Continentis parte observavit rariores, adjectis iconibus in solo natali delineatis. Vindobonae 1763.
- Koehne 1903 = *E. Koehne*: *Lythraceae* in *Engler*: *Regnum vegetabile* IV, 216, 1903.
- Otto Kuntze*: *Um die Erde. Reiseberichte eines Naturforschers.* Leipzig 1881.
- Melvill = *J. Cosmo Melvill*: List of the phanerogams of Key West, South Florida, mostly observed there in March 1872. [Memoirs of the Manchester Liter. and Philos. Society III ser. vol. VIII p. 138—154, London 1884.
- Mez 1896 = *C. Mez*: *Bromeliaceae* in *De Candolle Monographiae Phanerogamarum*, vol. IX, 1896.
- Millspaugh = *Charles Frederick Millspaugh*: *Flora of the Island of St. Croix.* [Field Columbian Museum. Chicago. 8°. Publication 68. Botanical Series. Vol. I. Nr. 7 (Nov. 1902) p. 441—546 with map of St. Croix.].
- Nieden zu 1899 = *F. Niedenzu*: De genere *Malpighia* in *Ind. lect. in lyceo Hosiano aest. instit. Brunsbergae* 1899.
- Planchon 1883 = *J. E. Planchon*: *Ampelidae* in *De Candolle Monographiae Phanerogamarum*, vol. V. 1883—1887.
- Pulle = *A. Pulle*: *An Enumeration of the Vascular Plants known*

- from Surinam, together with their Distribution and Synonymy. 1906.
- Report of the 60 meeting of the Br. Ass. f. Adv. Sc. Leeds 1890 p. 448.
- Schumann 1903 = *K. Schumann*: Gesamtbeschreibung der Kakteen (Monographia Cactacearum) Neudamm 1903.
- Olof Swartz*: Flora Indiae occidentalis aucta atque illustrata sive descriptiones plantarum in Prodomo recensitarum. Erlangae. 1797—1806. 3 vol. 8°: vol. I. 1797. VIII et 640 p., tab. 1—15; vol. II. 1800 p. 641—1230, tab. 16—24; vol. III. 1806. p. 1231—2018, ind. X p., tab. 25—29.
- W. F. R. Suringar*: Melocacti novi ex insulis archipelagi indici-occidentalis neerlandicis Curaçao, Aruba et Bonaire (Bijdrage tot de kennis der Melocacti. V in Versl. d. Natuurk. Afd. Kon. Akad. Wetensch. Amsterdam. Deel VI (1897) p. 178.
- Symb. = *I. Urban*: Symbolae Antillanae seu Fundamenta Florae Indiae Occidentalis. Berolini, Parisiis, Londini 8°.
- Vol. I. fasc. I 1898. p. 1—192, fasc. I 1899. p. 384—471.
- Vol. II. fasc. I 1900. p. 1—160, fasc. II 1900. p. 161—336, 1901, p. 344—388, fasc. III 1901. p. 337—508.
- Vol. III. fasc. I. 1902. p. 1—160, fasc. II. 1902. p. 161—352, fasc. III. 1903. p. 352—546.
- Vol. IV. fasc. I. 1903. p. 1—192, fasc. II. 1905. p. 193—352.
- Vol. V. fasc. I. 1904. p. 1—176, fasc. II. 1907. p. 177—352, fasc. III 1908. p. 353—552.
- Vol. VI. fasc. I. 1909.
- Urban 1893 = *I. Urban*: Additamenta ad cognitionem florae Indiae occidentalis in Engler's Botan. Jahrb. I. in vol. XV (1892) p. 286—361.
- Urban 1895 = *I. Urban*: Additamenta ad cognitionem florae Indiae occidentalis in Engler's Botan. Jahrb. II. in vol. XIX (1894—95) p. 562—681.
- Urban 1896 = *I. Urban*: Additamenta ad cognitionem florae Indiae occidentalis in Engler's Botan. Jahrb. III. in vol. XXI (1896) p. 514—638.
- Vesque 1893 = *J. Vesque*. Guttiferae in *De Candolle* Monographiae Phanerogamarum, vol. VIII. 1893.
- F. A. F. C. Went*: Rapport omtrent den toestand van land- en tuinbouw op de Nederlandsche Antillen, bijlage V van het Koloniaal verslag van Curaçao van 1902.
-

VERNACULAR NAMES.

- Adam's needle = *Yucca aloifolia* Linn. 36.
Adonis = *Duranta repens* Linn. 170.
Adonisabbot = *Leonotis nepetaefolia* R. Br. 172.
Almond tree = *Terminalia Catappa* Linn. 143.
Aloe = *Aloe vera* Linn. 36.
Antigua balsam = *Rauwolfia Lamarkii* DC. 157.
Arrowroot = *Maranta arundinacea* Linn. 40.
Avocado = *Persea americana* Mill. 70.
Bachelor's button = *Gomphrena globosa* Linn. 58.
Bahama grass = *Cynodon dactylon* Pers. 22.
Balsam tree = *Bursera simaruba* Sarg. 103.
Banso = *Samyda serrulata* Linn. 136.
Basil = *Ocimum basilicum* Linn. 175.
Basket hibisc = *Hibiscus schizopetalus* Hook. f. 130.
Basket hoop = *Smilax guianensis* Vitman. 37.
Bastard black berry = *Eugenia procera* Poir. 148.
Bastard bough = *Picramnia pentandra* Sw. 103.
Bastard cancker berry = *Psychotria rufescens* H. B. et K. 192.
Bastard catclauw = *Polypodium lycopodioides* Linn. 8.
Bastard cherry = *Malpighia augustifolia* Linn. 105.
Bastard cinnamom = *Ardisia coriacea* Sw. 151.
Bastard nettle = *Acalypha chamaedrifolia* Müll-Arg. 110.
Bastard white root = *Aegiphyla martinicensis* Jacq. 171.
Bayberry = *Amomis caryophyllata* Kr. et Urban. 144.
Bellapple = *Passiflora laurifolia* Linn. 138.
Big yellow popbush = *Crotalaria retusa* Linn. 87.
Billyache = *Rauwolfia Lamarkii* DC. 157.
Bitterroot = *Cassia occidentalis* Linn. 84.
Birdsberry = *Eugenia monticola* DC. 147.
= *Myrcia paniculata* Kr. et Urban. 145.
Bitterpot = *Urechites suberecta* Müll-Arg. 157.
Blackberry = *Eugenia ligustrina* Willd. 147.

- Blackcherry = *Eugenia ligustrina* Willd. 147.
 = *Randia aculeata* Linn. 188.
- Black lobloly tree = *Pisonia fragrans* Dumont, 61.
- Black mangeniel = *Euphorbia petiolaris* Curt. 116.
- Black sage = *Croton betulinus* Vahl. 109.
- Black sweetwood = *Nectandra Krugii* Mez. 71.
- Black toarch = *Melochia tomentosa* Linn. 132.
- Blisterbush = *Plumbago scandens* Linn. 152.
- Birdpepper = *Capsicum dulce* Hort. 176.
- Blueberry = *Symplocos martinicensis* Jacq. 153.
- Blydog = *Cassia bicapsularis* Linn. 83.
- Boulanger = *Solanum Melongena* Linn. 177.
- Boxwood = *Murraña exotica* Linn. 101.
 = *Schaefferia frutescens* Jacq. 121.
- Bracelet = *Erythroxylum havanense* Jacq. 99.
- Breadfruit tree = *Artocarpus incisa* Linn. f. 48.
- Bridgo tree = *Linociera compacta* R. Br. 154.
- Bruckroot = *Chiococca racemosa* Jacq. 190.
- Bugal plant = *Russelia juncea* Zuccar. 180.
- Bullgrass = *Mariscus brizaeus* C. B. Clarke. 27.
- Burrgrass = *Cenchrus tribuloides* Linn. 20.
- Button wood = *Conocarpus erectus* Linn. 143.
- Calabash = *Crescentia Cujete* Linn. 183.
- Calaloe = *Amarantus tristis* Linn. 57.
- Canckerberry = *Solanum racemosum* Jacq. 177.
- Candle wood = *Guettarda scabra* Lam. 189.
- Cane grass = *Panicum divaricatum* Linn. 16.
- Cannon = *Canna indica* Linn. 40.
- Casha = *Acacia Farnesiana* Willd. 79.
- Cashew = *Anacardium occidentale* Linn. 118.
- Cassave = *Manihot utilissima* Pohl. 114.
- Castoroil plant = *Ricinus communis* Linn. 112.
- Casuarine = *Casuarina equisetifolia* Linn. 44.
- Cat claw = *Bignonia unguis-cati* Linn. 181.
- Cat nip = *Salvia serotina* Linn. 173.
- Centepleeplant = *Comocladia ilicifolia* Sw. 119.
- Charlotten = *Allium ascalonicum* Linn. 36.
- Cherry = *Anacardium occidentale* Linn. 118.
 = *Malpighia puniceifolia* Linn. 106.
- Choaky berry = *Eugenia axillaris* Willd. 146.
- Chickenweed = *Euphorbia pilulifera* Linn. 116.
- Children chickenweed = *Euphorbia thymifolia* Linn. 117.

- Churchblossom = *Capparis frondosa* Jacq. 75.
 Churchyardblossom = *Lochnera rosea* Reichb. 156.
 Cinnamon tree = *Amomis caryophyllata* Kr. et Urban. 144.
 Cockspur = *Pisonia aculeata* Linn. 61.
 = *Xylosma buxifolium* A. Gray 136.
 Cocoa = *Theobroma Cacao* Linn. 133.
 Cocoaplum = *Chrysobalanus icaco* Linn. 77.
 Coffee = *Coffea arabica* Linn. 191.
 Coralita = *Antigonon leptopus* Hook. et Arn. 55.
 Coralplant = *Jatropha multifida* Linn. 113.
 Crabgrass = *Pappophorum alopecuroideum* Vahl. 24.
 = *Paspalum conjugatum* Bergius. 14.
 Crabwood = *Pithecolobium unguis-cati* Benth. 78.
 Crack open = *Casearia parvifolia* Willd. 137.
 Creole casha = *Acacia macracantha* Humb. et Bonpl. 79.
 Creole corn = *Andropogon sorghum* Brot. 13.
 Creole cotton = *Gossypium barbadense* Linn. 130.
 Croton = *Codiaeum variegatum* Bl. 114.
 Curaçao prickly = *Nopalea coccinellifera* Salm-Dyck. 142.
 Custard apple = *Anona reticulata* Linn. 69.
 Devil's bit = *Ruellia tuberosa* Linn. 185.
 Different coloured jessamine = *Quisqualis indica* Linn. 143.
 Divi divi = *Caesalpinia coriaria* Willd. 86.
 Dogwood = *Piscidia piscipula* Sarg. 92.
 Dutch casha = *Acacia tortuosa* Willd. 80.
 Dutch grass = *Eleusine indica* Gärt. 23.
 Elder = *Sambucus canadensis* Linn. 195.
 Eucharis lily = *Eucharis grandiflora* Planch. et Linden. 38.
 Eye bright = *Heliotropium indicum* Linn. 166.
 = *Heliotropium parviflorum* Linn. 167.
 Fat pork = *Chrysobalanus icaco* Linn. 77.
 Fire weed = *Datura fastuosa* Linn. 179.
 = *Datura Metel* Linn. 179.
 = *Stramonium* Linn. 179.
 Fishing rod = *Randia aculeata* Linn. 183.
 Flambeau = *Erithalis fruticosa* Linn. 190.
 Flamboyant = *Poinciana regia* Boy. 85.
 Four o' clock = *Mirabilis Jalapa* Linn. 60.
 Fox tail = *Erigeron canadensis* Linn. 199.
 French casha = *Acacia macracantha* Humb. et Bonpl. 79.
 Gingerbush = *Pavonia spinifex* Cav. 129.
 Goatbush = *Stigmatophyllum periplocifolium* A. Juss. 105.

- Lizardsgrass = *Sporobolus indicus* R. Br. 22.
 Locust = *Hymenaea Courbaril* Linn. 82.
 Logwood = *Haematoxylon campechianum* Linn. 85.
 Longgras = *Panicum insulare* G. F. W. Mey 17.
 Love wine = *Cuscuta americana* Linn. 159.
 Mahagony = *Swietenia mahagoni* Jacq. 104.
 Maho = *Daphnopsis caribaea* Gris. 142.
 Maidenapple = *Momordica Charantia* Linn. 195.
 Maidenhair = *Adiantum tenerum* Sw. 7.
 Mammie = *Mammea americana* Linn. 134.
 Man better man = *Achyranthes obtusifolia* Lam. 57.
 Man of war bush = *Capparis cynophallophora* Linn. 74.
 Mangueniel tree = *Hippomane mancinella* Linn. 114.
 Mango tree = *Mangifera indica* Linn. 118.
 Man jack = *Cordia sulcata* DC. 164.
 Mappoo = *Pisonia subcordata* Sw. 62.
 Maria pompoe = *Ibatia maritima* Dcne. 159.
 Marrown = *Croton flavens* Linn. 109.
 Mash mellow = *Sida cordifolia* Linn. 128.
 = *Waltheria americana* Linn. 132.
 Massamby = *Gynandropsis pentaphylla* DC. 73.
 Milky tree = *Tabernaemontana citrifolia* Jacq. 156.
 Mastic tree = *Sideroxylon foetidissimum* Jacq. 152.
 Millerwood = *Quararibea turbinata* Poir. 131.
 Miminet = *Lawsonia inermis* Linn. 142.
 Mimosa = *Leucana glauca* Benth. 80.
 Mispel = *Achras sapota* Linn. 152.
 Monkeytail = *Anthurium cordatum* G. Don. 30.
 Moringo = *Moringa oleifera* Lam. 76.
 Mosquito basil = *Ocimum micranthum* Willd. 175.
 Mountaingrass = *Scleria lithosperma* Sw. 29.
 Mulegrass = *Aristida americana* Linn. 21.
 Mustard = *Brassica integrifolia* O. E. Schulz 72.
 Mustard tree = *Capparis cynophallophora* Linn. 74.
 Myrtle lemon = *Triphasia trifoliata* P. DC. 101.
 Noyeaux = *Ipomoea pentaphylla* Jacq. 162.
 Nutgrass = *Cyperus rotundus* Linn. 27.
 Old man's beard = *Tillandsia recurvata* Linn. 33.
 = *Tillandsia usneoides*. Linn. 33.
 Oleander = *Nerium Oleander* Linn. 158.
 Orchids purslaue = *Peperomia glabella* A. Dietz. 45.
 Orebian jessamine = *Jasminum undulatum* Ker. Gawl. 154.

- Papao = *Carica Papaya* Linn. 139.
 Paragrass = *Panicum molle* Sw. 18.
 Parakiet = *Pedilanthus tithymaloides* Port. 117.
 Pea nut = *Arachis hypogaea* Linn. 90.
 Pea withe = *Rhynchosia reticulata* P. DC. 97.
 Pear tree = *Persea americana* Mill. 70.
 Pepper cinnamon = *Canella alba* Murr. 135.
 Physicnut = *Jatropha gossypifolia* Linn. 113.
 Picrous bark = *Jacquinia armillaris* Jacq. 150.
 Pigeonberry = *Eugenia axillaris* Willd. 146.
 Pigeonwood = *Plumiera alba* Linn. 155.
 Pingwing = *Bromelia pinguin* Linn. 31.
 Pita = *Fourcroya gigantea* Vent. 38.
 Plum rose = *Eugenia Jambos* Linn. 147.
 Pomegranate tree = *Punica Granatum* Linn. 143.
 Popehead = *Melocactus communis* Link et Otto 140.
 Poppy = *Cosmos sulfureus* Cav. 204.
 Potatoes = *Solanum tuberosum* Linn. 179.
 Potta potta = *Boerhaavea hirsuta* Willd 60.
 Prickly balsam = *Justicia sessilis* Jacq. 186.
 Pride of Barbados = *Caesalpinia pulcherrima* Sw. 86.
 Prince plum = *Amarantus paniculatus* Linn. 56.
 Pudding withe = *Cissus sicyoides* Linn. 124.
 Pumpkin = *Cucumis Anguria* Linn. 195.
 Purple popbush = *Crotalaria verrucosa* Linn. 87.
 Purple wormbush = *Stachytarpheta jamaicensis* Vahl. 169.
 Quaker blossom = *Plumbago capensis* Thunb. 151.
 Queen of Cribflower = *Lagerstroemia indica* Linn. 142.
 Rabbitmeat = *Salvia occidentalis* Sw. 173.
 Red frangepane = *Plumiera rubra* Linn. 156.
 Red man = *Comocladia ilicifolia* Sw. 119.
 Red plum = *Spondias purpurea* Linn. 119.
 Red wood = *Coccoloba diversifolia* Jacq. 53.
 = *Inga laurina* Willd. 78.
 = *Myrcia paniculata* Kr. et Urban 145.
 Rhamni = *Sansevieria guineensis* Willd. 36.
 Rice plant = *Ixora Jucunda* Twaithes 191.
 Roucou = *Bixa Orellana* Linn. 135.
 Runninggrass = *Panicum prostratum* Lam. 18.
 Running mountaingrass = *Oplismenus setarius* R. et Sch. 19.
 Sage = *Lantana Camara* Linn. 167.
 = *Lantana involucrata* Linn. 168.

- Saltpond tree = *Avicennia nitida* Jacq. 172.
 Sandbox tree = *Hura crepitans* Linn. 115.
 Sapodille = *Achras sapota* Linn. 152.
 Scarlet accordia = *Cordia Sebestena* Linn. 164.
 Sea grape = *Coccoloba uvifera* Jacq. 54.
 Sea lavendel = *Heliotropium curassavicum* Linn. 166.
 Sea peas = *Canavalia obtusifolia* P. DC. 96.
 Sea purslane = *Tournefortia gnaphalodes* R. Br. 165.
 Sea wine = *Ipomoea pescaprae* Roth. 162.
 Senna = *Tephrosia cathartica* Urb. 88.
 = *Tephrosia cinerea* Pers. 88.
 Shame lady shame = *Mimosa pudica* Linn. 81.
 Shellplant = *Costus speciosus* Sw. 40.
 Silk cotton purslane = *Portulaca halimoides* Linn. 66.
 Silk cotton tree = *Ceiba pentandra* Gärt. 131.
 Sintern pea nut = *Alysicarpus nummularifolius* P. DC. 92.
 Sintern weed = *Desmodium triflorum* P. DC. 91.
 Slingeractus = *Cereus triangularis* Haworth 140.
 Small yellow popbush = *Crotalaria incana* Linn. 87.
 Snakeberry tree = *Rauwolfia Lamareckii* DC. 157.
 Snaky = *Celtis Iguanacea* Sarg. 47.
 Snap dragon = *Angelonia angustifolia* Benth 180.
 Snijboontjes = *Phaseolus vulgaris* Linn. 97.
 Soapbush = *Sida carpinifolia* Linn. f. 127.
 Sour prickle = *Opuntia Tuna* Mill. 141.
 Sour sap = *Anona muricata* Linn. 68.
 Spanish casha = *Acacia macracantha* Humb. et Bonpl. 79.
 Spanish dagger = *Yucca aloifolia* Linn. 36.
 Spanish lady = *Opuntia triacantha* Haworth. 141.
 Sprainbush = *Chenopodium murale* Linn. 55.
 Sprainbush wine = *Cardiospermum halicacabum*. Linn 121.
 Sprain leaf = *Calotropis procera* R. Br. 158.
 Star of Bethlehem = *Euphorbia pulcherrima* Willd. 117.
 Stinging nettle = *Fleurya aestuans* Gaudich. 50.
 Stinging thyme = *Coleus amboinicus* Lour. 174.
 Stinkwood = *Piscidia piscipula* Sarg. 92.
 Strawberry = *Cereus triangularis* Haworth. 140.
 Sugarygrape = *Coccoloba nivea* Jacq. 53.
 Surinaam bitter = *Phyllanthus niruri* Linn. 108.
 Surinaam cherries = *Eugenia uniflora* Linn. 148.
 = *Myrcia splendens* DC. 145.
 Susanna berry = *Citharexylum quadrangulare* Jacq. 170.

- Sweet broom = *Scoparia dulcis* Linn. 181.
 Sweet gourd = *Lagenaria vulgaris* Ser. 196.
 Sweet potatoes = *Ipomoea batatas* Poir. 161.
 Sweet prickle = *Opuntia Ficus indica* Mill. 141.
 Sweetwood = *Nectandra coriacea* Griseb. 70.
 Tamarind = *Tamarindus indica* Linn. 82.
 Tantan = *Leucaena glauca* Benth. 80.
 Thistle = *Argemone mexicana* Linn. 71.
 Tomato = *Solanum lycopersicum* Linn. 177.
 Tiger tongue = *Sesbania grandiflora* Pers. 89.
 Vervine = *Bouchea Ehrenbergii* Cham. 169.
 Wall purslane = *Peperomia petiolaris* C. DC. 47.
 Waterthistle = *Neptunia plena* Benth. 82.
 Watergrass = *Commelina virginica* Linn. 34.
 White beans = *Phaseolus lunatus* Linn. 97.
 White cap = *Ageratum conyzoides* Linn. 197.
 White cedar = *Tecoma leucoxydon* Mart. 182.
 White coralita = *Porana paniculata* Roxb. 160.
 White frangepane = *Plumiera alba* Linn. 155.
 White honeysuckle = *Lonicera japonica* Thunb. 195.
 White hoop = *Villamilla octandra* Hook. f. 63.
 White lavender = *Tournefortia gnaphalodes* R. Br. 165.
 White polly = *Amarantus polygonoides* Linn. f. 56.
 White root = *Gouania domingensis* Linn. 123.
 White chunk = *Beureria succulenta* Jacq. 164.
 White snow = *Drymaria cordata* Willd. 67.
 White snowdrop = *Zephyranthes carinata* Herb. 37.
 White snowplant = *Iresine elatior* L. Cl. Rich. 58.
 White withe = *Smilax coriacea* Spr. 37.
 Wild banana = *Batis maritima* Linn. 40.
 = *Heliconia Bihai* Linn. 40.
 Wild cane = *Panicum divaricatum* Linn. 16.
 = *Piper reticulatum* Linn. 45.
 Wild cherry = *Guettarda parvifolia* Sw. 189.
 Wild clover = *Stylosanthes hamata* Taubert. 89.
 Wild coffee = *Picramnia pentandra* Sw. 103.
 Wild date = *Aleurites moluccana* Willd. 113.
 Wild guave = *Guettarda scabra* Lam. 189.
 = *Samyda serrulata* Linn. 136.
 Wild indigo = *Indigofera suffruticosa* Mill. 88.
 Wild maho = *Triumfetta rhomboidea* Jacq. 125.
 Wild mammie = *Clusia alba* Jacq. 135.

- Wild marrow = *Croton astroites* Dryand. 108.
Wild massamby = *Polanisia viscosa* DC. 73.
Wild misple = *Morisonia americana* Linn. 76.
Wild papaw = *Cecropia peltata* Linn. 50.
Wild peas = *Cassia glandulosa* Linn. 84.
 = *Desmodium supinum* P. DC. 91.
Wild pepper = *Solanum nigrum nodiflorum* Linn. 177.
Wild pine = *Tillandsia utriculata* Linn. 33.
Wild potatowine = *Ipomoea fastigiata* Sweet 161.
Wild salad = *Sonchus oleraceus* Linn. 207.
Wild senna = *Krameria ixina* Linn. 85.
Wild tantan = *Desmanthus virgatus* Willd. 81.
Wild tea = *Capraria biflora* Linn. 180.
Wild tobacco = *Vernonia punctata* Sw. 196.
Wonder tree = *Parkinsonia aculeata* Linn. 85.
Wormbush = *Chenopodium ambrosioides* Linn. 55.
Yellow balsem = *Croton flavens* Linn. 109.
Yellow bell = *Allamanda cathartica* Linn. 155.
Yellow blossom = *Stenolobium stans* Seem. 182.
Yellow plum = *Spondias lutea* Linn. 119.
Yellow prickle = *Fagara martinicensis* P. DC. 100.
Yellow sage = *Wedelia buphthalmoides* Gris. 202.
Yellow saunder = *Fagara flava* Kr. et Urb. 100.
-

INDEX TO THE FIRST PART.

- **Abelmoschus moschatus* Médik. 130.
Abrus precatorius Linn. 93.
Abutilon crispum Sweet. 126.
Abutilon indicum Sweet. 126.
Acacia Farnesiana Willd. 79.
Acacia macracantha Humb. et Bonpl. 79.
Acacia riparia H. B. K. 80.
Acacia sarmentosa Gris. 80.
Acacia tortuosa Willd. 80.
Acalypha chamaedrifolia Müll-Arg. 111.
Acalypha Poirerii Spreng. 111.
Acalypha reptans Sw. 111.
**Acalypha Wilkesiana* Müll-Arg. 112.
Acanthaceae 184.
Acanthospermum humile DC. 201.
Acnistus arborescens Schlecht. 175.
**Achras sapota* Linn. 152.
Achyranthes aspera var. *obtusifolia* Gris. 57.
Achyranthes obtusifolia Lam. 57.
Acrostichum conforme Sw. 10.
Acrostichum latifolium Sw. 10.
Acrostichum viscosum Vent. 10.
Actinostemon caribaenm Gris. 114.
**Adansonia digitata* Linn. 130.
Adelia pedunculosa A. Rich. 111.
Adelia ricinella Linn. 111.
**Adenanthera pavonina* Linn. 82.
Adiantum tenerum Sw. 7.
Adicea microphylla O. Ktze. 51.
Aechmea lingulata Baker. 32.
Aegiphila martinicensis Jacq. 171.
Ageratum conyzoides Linn. 197.
Aizoaceae 64.
**Albizia lebbek* Benth. 79.
**Aleurites moluccana* Benth. 113.
**Allamanda cathartica* Linn. 155.
**Allium ascalonicum* Linn. 36.
Allophylus occidentalis Radek. 121.
Aloe vera Linn. 36.
Aloe vulgaris Lam. 36.
Alternanthera achyrantha R. Br. 58.
Alternanthera crucis. Comb. nov. 58.
Alternanthera muscoides Bent. et Hook. 59.
Alternanthera repens O. Ktze. 58.
Alsophila phalerata Mart. 2.
**Althaea officinalis* Linn. 126.
Alysicarpus nummularifolius P. DC. 92.
Alysicarpus vaginalis P. DC. 92.
Amarantaceae 56.

- Amarantus caudatus* Linn. 56.
Amarantus gangeticus Linn. 57.
Amarantus gracilis Desf. 56.
 **Amarantus paniculatus* Linn. 56.
Amarantus polygonoides Linn. 56.
Amarantus spinosus Linn. 56.
Amarantus tristis Linn. 57.
Amaryllidaceae 37.
Amaryllis carinata Spreng. 37.
Amaryllis equestris Ait. 38.
Amblogyne polygonoides Raf. 56.
Ambrosia crithmifolia DC. 201.
 **Ambrosia hispida* Purch 201.
Amomis caryophyllata Kr. et Urb. 144.
Amymis elemifera Linn. 101.
Anacardiaceae 118.
 **Anacardium occidentale* Linn. 118.
Anamomis fragrans Gris. 146.
Anamomis punctata Gris. 146.
 **Ananas sativus* Schult. 32.
Anatherum bicornne Beauv. 13.
Anatherum macrourum Gris. 13.
Andira inermis H. B. K. 93.
Andira jamaicensis Urb. 93.
Andropogon bicornis Linn. 13.
Andropogon contortus Linn. 13.
Andropogon glomeratus Britton. 13.
Andropogon macrouros Michx. 13.
 **Andropogon Schoenanthus* Linn. 13.
Andropogon secundus Willd. 13.
 **Andropogon sorghum* Brot. 13.
Andropogon squarrosus Linn. f. 13.
Andropogon zizanoides Urb. 13.
 **Angelonia angustifolia* Benth. 180.
Anonaceae 68.
Anona montana Macf. 68.
 **Anona muricata* Linn. 68.
 **Anona reticulata* Linn. 69.
 **Anona squamosa* Linn. 69.
Antheophora elegans Schreb. 13.
Antheophora hermaphroditica O. Ktze. 13.
Anthurium cordatum G. Don. 30.
 **Antigonon cinerascens* Mart. et Gal. 55.
 **Antigonon leptopus* Hook. et Arn. 55.
Apocynaceae 155.
Araceae 30.
 **Arachis hypogaea* Linn. 90.
Ardisia coriacea Sw. 151.
Ardisia guadalupensis Duchuss. 151.
Argemone mexicana Linn. 71.
Argithamnia candicans Sw. 110.
Aristida americana Linn. 21.
Aristolochiaceae 53.
 **Aristolochia odoratissima* Linn. 53.
Aristolochia trilobata Linn. 53.
Artanthe Bredemeyeri Gris. 44.
Artocarpus communis Forst. 48.
Artocarpus incisa Linn. fil. 48.
Asclepiadaceae 158.
Asclepias Curassavica Linn.
Asclepias nivea Linn. 158.
Aspidium conterminum Willd. 2.
Aspidium exaltatum Sw. 3.
Aspidium molle Sw. 2.
Aspidium Plumierii Presl. 3.
Aspidium trifoliatum Sw. 3.
Asplenium auritum Sw. 4.
Asplenium bipartitum Spr. 4.
Asplenium cristatum Lam. 4.
Asplenium monanthes Linn. 4.
Asplenium obtusifolium Linn. 4.
Asplenium pumilum Sw. 5.

- Asplenium rhizophorum* Linn. 7.
cicutarium Sw. 4.
Asplenium serratum Linn. 5.
Asplenium striatum Presl. 4.
Avicennia nitida Jacq. 172.
Ayenia pusilla Linn. 132.
Baccharis cotinifolia Urb. 199.
Baccharis dioica Gris. 199.
Balsaminaceae 123.
Basellaceae 67.
Batidaceae 47.
Batis maritima Linn. 47.
**Bauhinia Kappleri* Sagot. 83.
Begoniaceae 139.
Begonia domingensis DC. 139.
**Belamcanda punctata* Moench. 39.
Beloperone eustachiana Benth. 186.
Benthamantha caribaea Benth. 89.
Bernardia corensis Müll.-Arg. 111.
Besleria lutea Linn. 183.
Beureria succulenta Jacq. 164.
Bidens bipinnatus Linn. 203.
Bidens leucantha Willd. 204.
Bidens pilosa Linn. 204.
Bignoniaceae 181.
**Bignonia Unguis-cati* Linn. 181.
Bignonia spec. 182.
Bixaceae 135.
**Bixa Orellana* Linn. 135.
Blechnum occidentale Linn. 5.
Blechnum Blechnum Millsp. 185.
Blechnum Brownei Juss. 185.
Boehmeria ramiflora Jacq. 52.
Boerhaavea erecta Linn. 60.
Boerhaavea hirsuta Willd. 60.
Boerhaavea paniculata L. Cl. Rich. 60.
Boerhaavea repens Linn. 60.
Boerhaavea scandens Linn. 61.
Bombacaceae 130.
**Bontia daphnoides* Linn. 187.
Borraginaceae 164.
Borreria ocymoides DC. 194.
Borreria podoccephala DC. 194.
Borrichia arborescens DC. 202.
Bouchea Ehrenbergii Cham. 169.
**Boussingaultia baselloides* H. B. K. 67.
Bouteloua juncifolia Lag. 21.
Bouteloua litigosa Lag. 21.
Brachyrrhamphus intybaceus DC. 207.
Bradburya virginianum O. Ktze. 94.
Brassavola cucullata R. Br. 43.
Brassavola rigida comb. nov. 43.
Brassica integrifolia O. E. Schulz. 72.
Bromeliaceae 31.
**Bromelia pinguin* Linn. 31.
**Brunfelsia americana* Linn. 180.
Bryophyllum calycinum Salisb. 77.
Bryophyllum pinnatum S. Kurz. 77.
Bumelia cuneata Sw. 153.
Bumelia obovata A. DC. 153.
Bunchosia jamaicensis Urb. et Niedenz. 106.
Bunchosia nitida L. Cl. Rich. 106.
Burseraceae 103.
Bursera gummifera Jacq. 103.
Bursera simaruba Sarg. 103.
Byrsonima crassifolia H. B. K. 107.
Byrsonima lucida L. Cl. Rich. 107.
Byrsonima martinicensis Kr. et Urb. 107.
Byrsonima spicata L. Cl. Rich. 107.
Cactaceae 140.
Caesalpinia ciliata Urb. 86.

- Caesalpinia coriaria Willd. 86.
 Caesalpinia crista Linn. 86.
 *Caesalpinia pulcherrima Sw. 86.
Cajanus cajan Millsp. 96.
 *Cajanus indicus Spreng. 96.
Cakile aequalis L'Herit. 72.
 Cakile lanceolata O. E. Schulz. 72.
 Callisia repens Linn. 35.
 Calotropis procera R. Br. 158.
 Calonyction speciosum Choisy. 163.
 Calonyction tuba Colla. 163.
 Calyptranthes Boldinghii Urb. 146.
 Canavalia obtusifolia P. DC. 96.
 Canavalia spec. 96.
Canellaceae 135.
 Canella alba Murr. 135.
Cannaceae 40.
 Canna indica Linn. 40.
Capparidaceae 73.
Capparis amygdalina Lam. 73.
 Capparis Breynia Jacq. 73.
 Capparis coccolobifolia Mart. 74.
 Capparis cynophallophora Linn. 74.
Capparis cynophallophora var. *latifolia* Gris. 74.
 Capparis frondosa Jacq. 75.
 Capparis jamaicensis Jacq. 75.
 Capraria biflora Linn. 180.
Caprifoliaceae 195.
 *Capsicum ceratocarpum Fing. 176.
 *Capsicum dulce Hort. 176.
 *Capsicum frutescens Linn. 176.
 Cardiospermum halicacabum Linn. 121.
Caricaceae 139.
 *Carica Papaya Linn. 139.
Caryophyllaceae 67.
 Casearia parvifolia Willd. 137.
 Casearia sylvestris Sw. 137.
 *Cassia alata Linn. 83.
 Cassia bicapsularis Linn. 83.
 *Cassia fistula Linn. 84.
 Cassia glandulosa Linn. 84.
 Cassia obovata Collad. 84.
 Cassia occidentalis Linn. 84.
 Cassia polyadena DC. 85.
Casuarinaceae 44.
 *Casuarina equisetifolia Linn. 44.
 Catopsis nutans Griseb. 34.
 Cecropia peltata Linn. 50.
 *Ceiba pentandra Gärtn. 131.
Celastraceae 119.
 *Celosia argentea Linn. 56.
 Celosia nitida Vahl. 56.
Celtis aculeata Sw. 47.
 Celtis Iguanacea Sarg. 47.
 Cenchrus tribuloides Linn. 20.
 Centrosema virginianum Benth. 94.
Cereus compressus Mill. 140.
 Cereus grandiflorus Mill. 140.
 Cereus Peruvianus Mill. 140.
 Cereus triangularis Haw. 140.
 Cereus spec. 140.
 Cestrum laurifolium L'Hér. 179.
Chaetochloa glauca Linn. 19.
Chaetochloa setosa Sw. 20.
Chaetogastra chironioides Gris. 148.
Chaetogastra strigosa Cogn. 148.
Chamaecrista glandulosa Linn. 84.
 Chaptalia nutans Hemsl. 206.
 Charianthus crinitus Naud. 148.
Charianthus nodosus crinitus Naud. 149.
 Cheilanthes microphylla Sw. 6.
Chenopodiaceae 55.
 Chenopodium ambrosioides Linn. 55.

- Chenopodium murale* Linn. 55.
Chiococca racemosa Jacq. 190.
Chloris ciliata Sw. 23.
Chloris paraguayensis Steud. 23.
Chloris radiata Sw. 23.
Chrysobalanus icaco Linn. 77.
Chrysophyllum argenteum Jacq. 153.
Chrysophyllum glabrum Jacq. 153.
Cicca antillana A. Juss. 108.
Cicca disticha Linn. 107.
Cissampelos Pareira Linn. 68.
Cissus obovata Vahl. 124.
Cissus sicyoides Linn. 124.
Citharexylum quadrangulare Jacq. 170.
**Citrus limonum* Risso. 102.
Cladium restioides Benth. 29.
Cleome graveolens Rafn. 73.
Cleome pentaphylla Linn. 73.
Cleome viscosa Linn. 73.
Clerodendron aculeatum Gris. 171.
Clerodendron fallax Lindl. 171.
Clerodendron fragans Willd. 171.
Clerodendron Thomsonae Balf. 171.
Clibadium erosum DC. 201.
**Clitoria ternatea* Linn. 93.
Clusia alba Jacq. 135.
Clusia rosea Jacq. 135.
Clusia spec. 135.
Coccoloba barbadensis Jacq. 53.
Coccoloba diversifolia Gris. 53.
Coccoloba diversifolia Jacq. 53.
Coccoloba excoriata Linn. 53.
Coccoloba Krugii Lindau. 53.
Coccoloba nivea Jacq. 53.
Coccoloba obtusifolia Jacq. 54.
Coccoloba punctata Gris. 53, 54.
Coccoloba punctata var. microstachya Gris. 54.
Coccoloba uvifera Jacq. 54.
Cocculus domingensis DC. 68.
**Cocos nucifera* Linn. 39.
**Codaeum variegatum* Bl. 114.
Coffea arabica Linn. 191.
Coleus amboinicus Lour. 174.
Coleus aromaticus Benth. 174.
**Coleus scutellarioides* Benth. 174.
**Colocasia esculenta* Schott. 31.
Colubrina ferruginosa Brong. 123.
Combretaceae 143.
Commelinaceae 34.
Commelina elegans H. B. K. 34.
Commelina virginica Linn. 34.
Comocladia ilicifolia Sw. 119.
Compositae 196.
Condalia ferrea Gris. 123.
Conocarpus erectus Linn. 143.
Convolvulaceae 159.
Convolvulus pentanthus Jacq. 161.
Corchorus acutangulus Lam. 125.
Corchorus hirsutus Linn. 125.
Corchorus olitorius Linn. 125.
Corchorus siliquosus Linn. 125.
Cordia salvifolia Juss. 164.
**Cordia Sebestena* Linn. 164.
Cordia sulcata DC. 164.
Cordia tremula Gris. 164.
**Cosmos sulphureus* Cav. 204.
**Costus speciosus* Sm. 40.
Cracca caribaea Benth. 89.
Crassulaceae 77.
Crescentia Cujete Linn. 183.
**Crinum americanum* Linn. 37.
Crotalaria incana Linn. 87.
Crotalaria retusa Linn. 87.
Crotalaria verrucosa Linn. 87.
Croton astroites Dryand. 108.
Croton balsamifer Jacq. 109.
Croton betulinus Vahl. 109.
Croton flavens Linn. 109.

- Croton lobatus Linn. 110.
Cruciferae 72.
 *Cryptostegia grandiflora R. Br. 158.
 Cucumis Anguria Linn. 195.
Cucurbitaceae 195.
 *Cucurbita maxima Duchesne. 196.
 Cuscuta americana Linn. 159.
 Cuscuta umbellata H. B. et K. 160.
Cyatheaceae 1.
 Cyathea arborea J. E. Sm. 1.
 Cyathea serra Willd. 1.
 Cymodocea manatorum Aschers. 12.
 Cynodon dactylon Pers. 22.
Cyperaceae 26.
 Cyperus bruncus Sw. 27.
 Cyperus capillaris Sw. 27.
 Cyperus confertus Gris. 26.
 Cyperus distans Linn. fil. 26.
 Cyperus elegans Linn. 26.
 Cyperus ferox L. C. Rich. 28.
 Cyperus flavus Nees. 27.
 Cyperus flavomariscus Gris. 28.
 Cyperus mucronatus Rottb. 26.
 Cyperus purpureo-variegatus Boeck. 27.
 Cyperus rotundus Linn. 27.
 Cyperus viscosus Sw. 26.
 Dactyloctenium aegyptium Willd. 24.
 Dalechampia scandens Linn. 112.
 Daphnopsis caribaea Gris. 142.
 Datura fastuosa Linn. 179.
 Datura Metel Linn. 179.
 Datura Stramonium Linn. 179.
 Desmanthus depressus Humb. et Bonpl. 81.
 Desmanthus virgatus Willd. 81.
 Desmodium axillare P. DC. 90.
 Desmodium incanum P. DC. 91.
 Desmodium scorpiurus Desv. 90.
 Desmodium spirale P. DC. 91.
 Desmodium supinum P. DC. 91.
 Desmodium tortuosum P. DC. 91.
 Desmodium triflorum P. DC. 91.
 Desmodium uncinatum P. DC. 92.
 Dianthera sessilis Gris. 186.
 Dichromena radicans Schlecht. 28.
Dicotyledoneae 44.
 Digitalia marginata Lk. 18.
 Digitalia setigera Roth. 18.
 Diodia rigida Cham. 193.
Dioscoreaceae 39.
 *Dioscorea alata Linn. 39.
 Dioscorea trifida Linn. 39.
 Dipholis salicifolia A. DC. 153.
 Diplazium semihastatum C. Christensen. 4.
 Diplazium striatum Presl.
 Distreptus spicatus Cass. 197.
 Ditaxis fasciculata Vahl. 110.
 Dodonaea viscosa Linn. 122.
 Dolicholus minimus Medic. 96.
 Dolicholus reticulatus Millsp. 97.
 Dolicholus lablab Linn. 98.
 Dorsteria contrajerva Linn. 48.
 Drymaria cordata Willd. 67.
 Dryopteris opposita Urb. 2.
 Dryopteris parasitica O. Ktze. 2.
 Dryopteris reticulata Urb. 3.
 Duranta Plumieri Jacq. 170.
 Duranta repens Linn. 170.
 Echites suberecta Jacq. 157.
 Eclipta alba Hassk. 202.
 Eclipta erecta Linn. 202.
 Ehretia Bourreria Linn. 164.
 Elaphoglossum conforme Schott. 10.
 Elaphoglossum latifolium J. Sm. 10.

- Elaphoglossum petiolatum* Urb. 10.
Elephantopus spicatus Aubl. 197.
Eleusine aegyptiaca Desf. 24.
Eleusine indica Gärtn. 23.
**Eleutherine plicata* Herb. 39.
Emilia sonchifolia DC. 206.
Enckea Sieberi Gris. 44.
Enckea smilacifolium C. DC. 45.
**Enterolobium Timbouwa* Mart. 78.
Epidendrum anceps Jacq. 41.
Epidendrum ciliare Linn. 42.
Epidendrum difforme Jacq. 42.
Epidendrum fuscatum Sm. 41.
Epidendrum globosum Jacq. 42.
Epidendrum papilionaceum Vahl 42.
Epidendrum strobiliferum Reichb. f. 43.
Epidendrum umbellatum Sw. 42.
Eragrostis ciliaris Lk. 25.
Eragrostis pilosa Beauv. 25.
Eragrostis poaeoides Gris. 25.
Eragrostis tephrosanthos Schult. 25.
Erechthites hieracifolia Raf. 205.
Erechthites praealta Raf. 205.
Erechthites valerianaefolia DC. 206.
Erigeron bonariensis Linn. 199.
Erigeron canadensis Linn. 199.
Erigeron spathulatus Vahl 199.
Eriodendron anfractuosum DC. 131.
Erithalis fruticosa Linn. 190.
Ernodea litoralis Sw. 193.
Eryngium foetidum Linn. 150.
Erythrina corallodendron Linn. 94.
Erythroxylaceae 99.
Erythroxylum brevipes P. DC. 99.
Erythroxylum havanense Jacq. 99.
Erythroxylum obtusum DC. 99.
**Eucharis grandiflora* Planch. 38.
Eugenia axillaris Willd. 146.
Eugenia buxifolia Gris. 147.
Eugenia cordata var *Sintensi* Kr. et Urb. 146.
Eugenia floribunda West. 146.
Eugenia fragrans Willd. 146.
Eugenia Jambos Linn. 147.
Eugenia ligustrina Willd. 147.
Eugenia monticola DC. 147.
Eugenia obtusata Willd. 147.
Eugenia pallens Gris. 147.
Eugenia Poirerii Berg. 147.
Eugenia procera Poir. 148.
Eugenia rhombea Kr. et Urb. 148.
Eugenia sessiliflora DC. 146.
Eugenia uniflora Linn. 148.
Eupatorium canescens Vahl 197.
Eupatorium conyzoides Vahl 198.
Eupatorium guadalupense Spreng. 198.
Eupatorium iresinoides H. B. et K. 198.
Eupatorium macranthum Sw. 198.
Eupatorium macrophyllum Linn. 198.
Eupatorium paniculatum Schrad. 198.
Eupatorium sinuatum Lam. 197.
Euphorbiaceae 107.
Euphorbia articulata Burm. 115.
Euphorbia buxifolia Lam. 115.
Euphorbia cyathophora Murr. 115.
Euphorbia heterophylla Linn. 116.

- Euphorbia hypericifolia* Linn. 116.
Euphorbia linearis Retz. 115.
Euphorbia petiolaris Curt. 116.
Euphorbia pilulifera Linn. 116.
Euphorbia prostrata Ait. 117.
Euphorbia pulcherrima Willd. 117.
Euphorbia thymifolia Linn. 117.
Euphorbia verticillata Poir. 116.
Euterpe spec. 30.
Euxolus caudatus Moq. 56.
Evolvulus argyraeus Choisy 160.
Evolvulus glaber Spreng. 160.
Evolvulus incanus Pers. 160.
Evolvulus mucronatus Sw. 160.
Evolvulus sericeus Sw. 160.
Excoecaria caribaea Gris. 114.
Excoecaria lucida Sw. 114.
Exostemma caribaeum Roem. et Schult. 188.
Exostemma sp. 188.
Fagara flava Kr. et Urb. 100.
Fagara martinicensis Lam. 100.
Fagara microphylla Dess. 100.
Fagara spinifex Jacq. 100.
Fagara trifoliata Sw. 101.
Faramea odoratissima DC. 193.
Festuca Myurus Linn. 26.
Ficus Krugiana Warb. 49.
Ficus laurifolia Duss. 49.
Ficus laevigata Vahl. 49.
Ficus omphalophora Warb. 49.
Ficus pallida Gris. 49.
Ficus pertusa Gris. 49.
Ficus populnea Willd. 49.
Ficus Urbaniana Warb. 49.
Fimbristylis spathacea Roth. 28.
Flacourtiaceae 136.
Fleurya aestuans Gaudich. 50.
**Foeniculum vulgare* Mill. 150.
Forestiera Eggersiana Kr. et Urb. 154.
Fourcroya gigantea Vent. 38.
Freziera undulata Willd. 134.
**Gaillardia aristata* Purch. 204.
Galactia angustifolia Griseb. 95.
Galactia dubia P. DC. 95.
Galactia filiformis Griseb. 95.
Galactia longiflora Griseb. 95.
Galactia longifolia Benth. 95.
Galactia nummularia Urb. 95.
Galactia rubra Urb. 95.
Galactia tenuiflora Wight. et Arn. 95.
**Galphimia gracilis* Bartl. 105.
Gentianaceae 155.
Gesneriaceae 183.
Gesneria ventricosa Sw. 184.
Gleicheniaceae 11.
Gleichenia bifida Spreng. 11.
Gnaphalium americanum Mill. 200.
Gnaphalium purpureum Linn. 200.
**Gomphrena globosa* Linn. 58.
Gonzalagunia hirsuta Schumann 188.
Gonzalea spicata DC. 188.
Goodeniaceae 196.
**Gossypium barbadense* Linn. 130.
Gouania domingensis Linn. 123.
Gramineae 12.
**Graptophyllum pictum* Griff. 186.
Guaiacum officinale Linn. 99.
Gundlachia domingensis A. Gray 199.
Guetarda parvifolia Sw. 189.
Guetarda scabra Lam. 189.
Guilandina Bonduc Gris. 86.
Guilandina glabra Gris. 86.
Guilandina Grisebachiana Kr. et Urb. 86.

Guttiferae 134.

Gyminda latifolia Urb. 120.

Gymnanthes lucida Sw. 114.

*Gymnogramme calomelanos**Kaulf.* 6.**Gymnospermae** 12.

Gynandropsis pentaphylla DC. 75.

Haematoxylon campechianum

Linn. 85.

Hebeclinum macrophyllum DC.

198.

Heckeria peltata Kunth. 45.

Helecharis capitata Boeck. 28.

Heliconia Bihai Linn. 40.

Helicteres Isora Linn. 133.

Helicteres jamaicensis 133.

Heliotropium curassavicum

Linn. 166.

Heliotropium indicum Linn. 166.

Heliotropium microphyllum Sw.

167.

Heliotropium parviflorum Linn.

167.

*Heliotropium peruvianum Linn.

167.

Hemidiodia ocimifolia K. Schum.

193.

Hemionitis palmata Linn. 5.

Hemitelia horrida R. Br. 2.

Heteropteris lanceolata Fee. 7.

Heteropteryx purpurea H. B. K.

104.

*Hibiscus esculentus Linn. 129.

*Hibiscus radiatus Cav. 129.

*Hibiscus Rosa-sinensis Linn.

129.

*Hibiscus Sabdariffa Linn. 129.

*Hibiscus schizopetalus Hook.

130.

Hibiscus tiliaceus Linn. 130.

Hillia longiflora Sw. 187.

Hillia parasitica Jacq. 187.*Hippeastrum equestre* Herb. 38.

*Hippeastrum puniceum Urb. 38.

Hippomane mancinella Linn. 114.

Hirtella triandra Sw. 78.

*Hoya carnosa R. Br. 159.

Hura crepitans Linn. 115.

Hydrolea jamaicensis Raeusch.

163.

Hydrophyllaceae 163.

Hymenaea Courbaril Linn. 82.

Hymenocallis caribaea Herb. 38.

Hymenophyllaceae 1.

Hymenophyllum hirtellum

Sw. 1.

Hypelate trifoliata Sw. 122.

Hyperbaena domingensis

Benth. 68.

Hypoxis decumbens Linn. 39.

Hyptis pectinata Poit. 174.

Ibatia maritima DCne 159.

Ibatia muricata Gris. 159.

Ichnanthus pallens Munro. 19.

Ichthiomethia piscipula Hitchc. 92.

*Impatiens Balsamina Linn. 123.

Indigofera anil Linn. 88.

Indigofera suffruticosa Mill. 88.

Inga laurina Willd. 78.

Ipomoea arenaria Steud. 161.

*Ipomoea batatas Poir. 161.

Ipomoea biloba Forsk. 162.*Ipomoea bona nox* Linn. 163.

Ipomoea dissecta Pers. 161.

Ipomoea fastigiata Sweet. 161.

Ipomoea grandiflora Lam. 163.*Ipomoea hederacea* Jacq. 162.

Ipomoea Nil Roth. 162.

Ipomoea pentaphylla Jacq. 162.

Ipomoea pes caprae Roth. 162.

Ipomoea sinuata Ortega. 161.

Ipomoea sphenophylla Urb. 163.

Ipomoea tuba G. Don. 163.

Ipomoea violacea Linn. 163.

Iresine celosioides Linn. 58.
Iresine elatior L. Cl. Rich. 58.
Iresine paniculata O. Ktze 58.
Iridaceae 39.
Isachne arundinacea Gris. 15.
Isachne rigens Trin. 15.
Isoloma hirsuta Reg. 184.
**Ixora coccinea* Linn. 191.
**Ixora Jucunda* Thwaites 191.
Jacquemontia pentantha G. Don.
 161.
Jacquemontia violacea Choisy.
 161.
Jacquinia armillaris Jacq. 150.
Jacquinia Berteri Spreng. 151.
Jambosa Jambos Millsp. 147.
Jambosa vulgaris DC. 147.
**Jasminum azoricum* Linn. 154.
**Jasminum pubescens* Willd. 154.
**Jasminum undulatum* Ker.
 Gawl. 154.
Jatropha curcas Linn. 113.
Jatropha gossypifolia Linn. 113.
**Jatropha multifida* Linn. 113.
Juncellus laevigatus C. B. Clarke
 26.
**Juniperus bermudiana* Linn. 12.
**Juniperus virginica* Linn. 12.
Justicia eustachiana Jacq. 186.
Justicia sessilis Jacq. 186.
Kallstroemia maxima Wight et
 Arn. 99.
**Kohleria hirsuta* Reg. 183.
Krameria ixinia Linn. 85.
Krugiodendron ferreum Urb.
 123.
Kyllinga brevifolia Rottb. 28.
Kyllinga monocephala Thunb. 28.
Labiatae 172.
Lablab vulgaris Savi 98.
Lachnostoma maritimum Nichols.
 159.

Lactuca intybacea Jacq. 207.
Lagascea mollis Cav. 200.
Lagenaria vulgaris Ser. 196.
**Lagerstroemia indica* Linn. 142.
Laguncularia racemosa Gärtn.
 144.
Lantana Camara Linn. 167.
Lantana involucrata Linn. 168.
Lappago aliena Spreng. 14.
Lauraceae 70.
Lawsonia inermis Linn. 142.
Lebidibia coriaria Schlecht. 86.
Leguminosae 78.
Leiphaimos aphylla Gilg. 155.
Lemnaceae 31.
Lemna minor Griseb. 31.
Lemna paucicostata Hegelm. 31.
Lentibulariaceae 184.
Leonotis nepetaefolia R. Br. 172.
Leonurus sibiricus Linn. 172.
Lepicystis piloselloides Diels 9.
Lepicystis squamata J. Sm. 10.
Lepidium virginicum Linn. 72.
Leptochloa filiformis Beauv. 24.
Leptochloa mucronata Kth. 24.
Leptochloa virgata Beauv. 24.
Leria nutans DC. 206.
Leucaena glauca Benth. 80.
Leucas martinicensis R. Br. 172.
Liliaceae 36.
Linociera compacta R. Br. 154.
Lippia reptans H. B. et K. 169.
Lithophila muscoides Sw. 59.
Lithophila vermiculata Uline. 59.
**Lochnera rosea* Reichb. 156.
Loganiaceae 154.
Lonchitis hirsuta Linn. 7.
Lonchocarpus violaceus H. B.
 et K. 92.
**Loniceria japonica* Thunb. 195.
Loranthaceae 52.
Lucuma multiflora A. DC. 152.

- Lycium americanum Jacq. 175.
Lycopodiaceae 11.
 Lycopodium taxifolium Sw. 11.
 Lycopodium verticillatum Linn.
 fil. 11.
Lythraceae 142.
Machaerina restioides Vahl. 29.
Malpighiaceae 104.
 Malpighia angustifolia Linn. 105.
 Malpighia puniceifolia Linn. 106.
Malvaceae 126.
Malvastrum coromandelianum
 Garcke 127.
 Malvastrum spicatum A. Gray
 126.
 Malvastrum tricuspidatum A.
 Gray 127.
 *Mammea americana Linn. 134.
 *Mangifera indica Linn. 118.
 *Manihot utilisima Pohl. 114.
Mapouria parasitica K. Sch. 192.
Marantaceae 40.
 *Maranta arundinacea Linn. 40.
Marcgraviaceae 133.
 Marcgravia umbellata Linn. 133.
 Marila racemosa Sw. 134.
 Mariscus brizaens C. B. Clarke
 27.
 Mariscus capillaris Vahl. 27.
 Mariscus flavus Vahl. 27.
 Maurandia semperflorens Jacq.
 180.
Mayepea caribaea O. Ktze. 154.
 Maytenus elliptica Kr. et Urb.
 119.
 *Medicago sativa Linn. 88.
Meibomia axillaris O. Ktze 90.
Meibomia incana O. Ktze 91.
Meibomia scorpiurus O. Ktze 90.
Meibomia spiralis O. Ktze 91.
Meibomia triflora O. Ktze 91.
Melastomataceae 148.
Meliaceae 104.
 *Melia areolaris Linn. 104.
Melia sempervirens Sw. 104.
 *Melicocca bijuga Linn. 122.
 Melocactus communis Link. et
 Otto 140.
 Melochia nodiflora Sw. 131.
 Melochia pyramidata Linn. 131.
 Melochia tomentosa Linn. 132.
 Melothria fluminensis Gard. 195.
Meniscium reticulatum Sw. 3.
Menispermaceae 68.
Mertensia furcata Gris. 11.
Mesosphaerum pectinatum O.
 Ktze 174.
Metastelma Paralias Dcne 159.
 Metastelma parviflora R. Br. 159.
Metastelma Schlechtendalii Gris.
 159.
 Miconia impetiolaris D. Don. 149.
 Miconia laevigata DC. 149.
 Microstylis spicata Lindl. 41.
 Microtea debilis Sw. 64.
 Mimosa pudica Linn. 81.
 *Mirabilis Jalapa Linn. 60.
Mitracarpus hirtus DC. 194.
 Mitracarpus polycladus Urb. 194.
 Mitracarpus villosus Cham. et
 Schlecht. 194.
 Mollugo verticillata Linn. 64.
 Momordica Charantia Linn. 195.
Monocotyledoneae 12.
Moraceae 48.
Moringaceae 76.
Morinda moringa Millsp. 76.
 *Moringa oleifera Lam. 76.
Moringa pterygosperma Gärtn.
 76.
 Morisonia americana Linn. 76.
 *Mucuna pruriens P. DC. 95.
 *Murraya exotica Linn. 101.
Musaceae 39.

Musa sapientium Linn. 39.
 **Musa paradisiaca* Linn. 39.
 **Mussaenda frondosa* Linn. 188.
Myginda latifolia Sw. 120.
Myginda pallens Sarg. 120.
Myginda pallens Smith 120.
Myginda rhacoma Sw. 120.
Myoporaceae 187.
Myrcia coriacea DC. 145.
Myrcia paniculata Kr. et Urb. 145.
Myrcia splendens DC. 145.
Myrodia turbinata Sw. 130.
Myroxylon buxifolium Kr. et Urb. 136.
Myrsinaceae 151.
Myrsine laeta Gris. 151.
Myrtaceae 144.
Nama jamaicensis Linn. 163.
Nazia aliena Millsp. 14.
Nectandra coriacea Gris. 70.
Nectandra discolor Gris. 71.
Nectandra Krugii Mez. 71.
Nectandra Wildenoviana Nees. 70.
Nephrodium oppositum Diels. 2.
Nephrolepis exaltata Schott. 3.
Neptunia plena Benth. 82.
 **Nerium Oleander* Linn. 158.
Neurogramme calomelanos Diels. 6.
Neurolaena lobata R. Br. 205.
 **Nicotiana Tabacum* Linn. 180.
 **Nopalea coccinellifera* Salm-Dyck 142.
Nyctaginaceae 60.
Nymphaeaceae 67.
Nymphaea spec. 67.
Ochnaceae 133.
 **Ocimum basilicum* Linn. 175.
 **Ocimum gratissimum* Linn. 175.
Ocimum micranthum Willd. 175.
Olacaceae 52.
Oldenlandia herbacea DC. 187.

Oleaceae 154.
Olfersia cervina O. Ktze. 3.
Oncidium silvestre Lindl. 43.
Oncidium tetrapetalum W. 43.
Oncidium variegatum Sw. 43.
 **Operculina ventricosa* Peter. 161.
Ophioglossaceae 11.
Ophioglossum reticulatum Linn. 11.
Oplismenus setarius R. et Sch. 19.
Opuntia Ficus indica Mill. 141.
Opuntia spinosissima Mill. 141.
Opuntia triacantha Haworth. 141.
Opuntia Tuna Mill. 141.
Orchidaceae 40.
 **Oreodoxa caribaea* Damm. et Urb. 30.
Ornithidium coccineum Salisb. 43.
Orthopogon setarius Spreng. 19.
Oxalidaceae 98.
Oxalis corniculata Linn. 98.
 **Oxalis Martiana* Zucc. 98.
Pachygone domingensis Eichl. 68.
Palicourea Pavetta DC. 192.
Palmae 29.
Paltonium lanceolatum Pr. 7.
 **Panax fruticosum* Linn. 150.
Pancreatium caribaeum Linn. 38.
Panicum barbinode Trin. 18.
Panicum brevifolium Linn. 15.
Panicum colonum Linn. 15.
Panicum diffusum Sw. 15.
Panicum divaricatum Linn. 16.
Panicum fuscum Sw. 16.
Panicum geminatum Forsk. 16.
Panicum insulare G. F. W. Mey. 17.
Panicum lanatum Rottb. 17.
Panicum leucophaeum H. B. K. 17.
Panicum maximum Jacq. 17.

- Panicum molle* Sw. 18.
Panicum pallens Sw. 19.
Panicum paspaloides Pers. 16.
Panicum prostratum Lam. 18.
Panicum sanguinale Linn. 18.
Papaveraceae 71.
Pappophorum alopecuroideum
Vahl. 24.
Pappophorum laguroideum
Schrud. 25.
Paritium tiliaceum St. Hill. 130.
**Parkinsonia aculeata* Linn. 85.
Parthenium Hysterophorus
Linn. 201.
Paspalum conjugatum Bergius.
14. —
Paspalum fimbriatum H. B. K.
14.
Paspalum hemisphaericum Poir.
14.
Paspalum paniculatum Willd. 14.
Passifloraceae 138.
Passiflora foetida Linn. 138.
**Passiflora laurifolia* Linn. 138.
Passiflora maliformis Linn. 138.
**Passiflora quadrangularis* Linn.
139.
Passiflora rubra Linn. 139.
Passiflora suberosa Linn. 139.
Pavonia spinifex Cav. 129.
Pectis febrifuga van Hall. 205.
Pectis humifusa Sw. 205.
Pectis linifolia Less. 205.
Pectis Swartziana Borg et Pauls.
205.
Pedilanthus tithymaloides Poit.
117.
**Peireskia aculeata* Mill. 142.
**Peireskia bleo* DC. 142.
Pennisetum setosum Rich. 21.
Pentarrhaphia longiflora Lindl.
184.
Peperomia acuminata C. DC. 45.
Peperomia emarginella C. DC. 45.
Peperomia exilis Griseb. 45.
Peperomia glabella A. Dietr. 45.
Peperomia glabella Griseb. 45.
Peperomia glabella var. *nigropunctata* Duss. 45.
Peperomia hernandifolia A. Dietr.
46.
Peperomia Langsdorffii Miq. 46.
Peperomia magnoliaefolia A.
Dietr. 46.
Peperomia obtusifolia A. Dietr.
46.
Peperomia petiolaris C. DC. 47.
Peperomia repens. Kth. 47.
Peperomia scandens R. et Pav.
47.
Peperomia urocarpa Fisch. et
Mey. 47.
**Persea americana* Mill. 70.
Persea gratissima Gärtn. 70.
Persea Persea Cock. 70.
Petiveria alliacea Linn. 64.
**Petunia violacea* Lindl. 180.
Pharbitis Nil Choisy. 162.
Pharus glaber H. B. K. 21.
Phaseolus lathyroides Linn. 97.
**Phaseolus lunatus* Linn. 97.
Phaseolus semierectus Linn. 97.
**Phaseolus vulgaris* Linn. 97.
Philodendron dispar Schott. 30.
Philodendron giganteum Schott.
31.
Philodendron oxycardium
Schott. 31.
Philoxerus vermicularis R. Br.
59.
Phoebe elongata Nees. 70.
**Phoenix dactylifera* Linn. 29.
Phoradendron trinervium Gris.
52.

Phyllanthus distichus Müll.-Arg.
107.

Phyllanthus epiphyllanthus
Linn. 107.

Phyllanthus falcatus Sw. 107.

Phyllanthus niruri Linn. 108.

Phyllanthus nobilis Mull.-Arg.
108.

Physalis angulata Linn. 175.

Physalis pubescens Linn. 176.

Physurus hirtellus Lindl. 41.

Physurus plantagineus Lindl. 41.

Phytolaccaceae 63.

Phytolacca icosandra Linn. 64.

Picraena excelsa Gris. 102.

Picramnia pentandra Sw. 103.

Picrasma antillana Urb. 102.

Pilea ciliaris Wedd. 51.

Pilea elegans Wedd. 51.

Pilea microphylla Liebm. 50.

Pilea muscosa Lindl. 51.

Pilea obtusata Liebm. 51.

Pilea parietaria Bl. 51.

Pilea pubescens Gris. 51.

Pilea semidentata Wedd. 51.

Pimenta acris Kostel. 144.

Pinaceae 12.

Piperaceae 45.

Piper Bredemeyeri Jacq. 44.

Piper ceanothifolium Kunth. 44.

Piper dilatatum L. Cl. Rich. 44.

Piper marginatum Jacq. 44.

Piper medium Jacq. 44.

Piper peltatum Linn. 45.

Piper reticulatum Linn. 45.

Piper smilacifolium P. DC. 45.

Piriqueta cistoides G. F. W.

Mey 137.

Piriqueta villosa Aubl. 137.

Piscidia erythrina Linn. 92.

Piscidia piscipula Sarg. 92.

Pisonia aculeata Linn. 61.

Pisonia fragans Dumont 61.

Pisonia inermis Gris. 61.

Pisonia obtusata Sw. 61.

Pisonia subcordata Sw. 62.

Pitcairnia angustifolia Redouté
32.

Pithecolobium ungius-cati
Benth. 78.

Plantaginaceae 187.

Plantago major Linn. 187.

Pluchea odorata Cass. 200.

Pluchea purpurascens DC. 200.

Plumbaginaceae 151.

**Plumbago capensis* Thunb. 151.

Plumbago scandens Linn. 152.

Plumiera alba Linn. 155.

**Plumiera rubra* Linn. 156.

**Poinciana regia* Boj. 85.

Polanisia icosandra Wight et
Arn. 73.

Polanisia viscosa DC. 73.

Polyboea corensis Klotsch. 111.

Polybotrya cervina Klfs. 3.

Polygonaceae 53.

Polypodiaceae 2.

Polypodium aureum Linn. 8.

Polypodium brasiliense Poir. 8.

Polypodium crassifolium Linn. 8.

Polypodium loriceum Linn. 8.

Polypodium lycopodioides

Linn. 8.

Polypodium neriifolium
Schkuhr. 8.

Polypodium phyllitidis Linn. 9.

Polypodium piloselloides Linn. 9.

Polypodium plumula H. B.
Wild. 9.

Polypodium serrulatum Mett. 10.

Polypodium squamatum
Linn. 10.

Polypodium trichomanoides
Sw. 10.

- Polystachya luteola* Hook. 41.
 **Porana paniculata* Roxb. 160.
Porophyllum ellipticum Cass. 204.
Porophyllum ruderale Gris. 204.
Portulacaceae 65.
Portulaca halimoides Linn. 66.
 **Portulaca oleracea* Linn. 66.
Portulaca pilosa Linn. 66.
Portulaca poliosperma Urb. 67.
Portulaca quadrifida Linn. 67.
Potamogetonaceae 12.
Pothomorphe peltata Miq. 45.
Prescottia stachyoides Lindl. 41.
Priva echinata Juss. 170.
Priva lappulacea Pers. 170.
 **Prockia Crucis* Linn. 136.
Prunus myrtifolia Urb. 77.
Prunus sphaerocarpa Sw. 77.
Psidium Guajava Linn. 144.
Psilotaceae 11.
Psilotum nudum Gris. 11.
Psilotum triquetrum Sw. 11.
Psychotria crassa Gris. 192.
Psychotria horizontalis Sw. 191.
Psychotria parasitica Sw. 192.
Psychotria pendula Urb. 192.
Psychotria rufescens H. B. et K. 192.
Psychotria uliginosa Sw. 192.
Pteridophyta 1.
Pteris biauaria Linn. 7.
Pteris laciniata Willd. 7.
Punicaceae 143.
 **Punica Granatum* Linn. 143.
 **Quamoclit vulgaris* Choisy 163.
Quararibea turbinata Poir. 131.
 **Quisqualis indica* Linn. 143.
Randia aculeata Linn. 188.
Rapanea ferruginea Mez. 151.
Rauwolfia Lamarekii DC. 157.
Rhacoma crossopetalum Linn. 120.
Rhacoma latifolia Loesener 120.
Ramnaceae 123.
Rhizophoraceae 143.
Rhizophora Mangle Linn. 143.
 **Rhoeo discolor* Hance 35.
Rhynchosia minima P. DC. 96.
Rhynchosia reticulata P. DC. 97.
Rhynchospora Humboldtiana Gris. 29.
Rhynchospora Persooniana Gris. 29.
Rhynchospora polyphylla Vahl. 29.
 **Ricinus communis* Linn. 112.
Rivina humilis Linn. 63.
Rivina laevis Linn. 63.
Rivina octandra Linn. 63.
Rochefortia acanthophora Gris. 165.
Rosaceae 77.
Rousselia humilis Urb. 52.
Rousselia lappulacea Gaudich. 52.
Rubiaceae 187.
Ruellia tuberosa Linn. 185.
 **Russelia juncea* Zuccar. 180.
Rutaceae 100.
 **Saccharum officinarum* Linn. 12.
Salicornia herbacea Linn. 55.
Salvia occidentalis Sw. 173.
Salvia serotina Linn. 173.
 **Sambucus canadensis* Linn. 195.
Samyda serrulata Linn. 136.
 **Sanchezia nobilis* Hook. 185.
 **Sansevieria guineensis* Willd. 36.
Sapindaceae 121.
Sapotaceae 152.
Sapota Achras Mill. 152.
Sauvagesia erecta Linn. 133.
Scaevola Lobelia Murr. 196.
Scaevola Plumieri Vahl. 196.
Schaefferia frutescens Jacq. 121.

Schmidelia occidentalis Sw. 121.
Schoepfia Schreberi J. F. Gmel.
 52.

Scirpus capitatus Linn. 28.
Scirpus obtusifolius Gris. 28.
Scleria filiformis Sw. 29.
Scleria lithosperma Sw. 29.
Scleria pratensis Nees. 29.
Scleria pterota Presl. 29.
Scoparia dulcis Linn. 181.

Scrophulariaceae 180.

Sechium edule Sw. 196.

Selaginellaceae 11.

Selaginella flabellata Spring. 11.
Selaginella substipitata Spring.
 12.

Senecio sonchifolius Moench.
 206.

**Sesbania grandiflora* Pers. 89.
Sesbania sericea P. DC. 89.
Sesuvium portulacastrum Linn.
 64.

Setaria glauca Beauv. 19.
Setaria setosa Beauv. 20.
Setaria viridis Beauv. 20.
Sida acuta Burm. 127.
Sida carpinifolia Linn. 127.
Sida ciliaris Linn. 127.
Sida cordifolia Linn. 128.
Sida glomerata Cav. 128.
Sida rhombifolia Linn. 128.
Sida spinosa Linn. 128.
Sida supina l'Herit. 128.
Sideroxylon foetidissimum

Jacq. 152.

Sideroxylon mastichodendron
 Jacq. 152.

Simarubaceae 102.

Sinapis brassicata Gris. 72.
Sinapis juncea Eichl. 72.
Smilax coriacea Spr. 37.
Smilax guianensis Vitmann 37.

Smilac havanensis Gris. 37.

Smilax ilicifolia Kth. 37.

Smilac macrophylla Willd. 37.

Solanaceae 175.

Solanum aculeatissimum Jacq.
 176.

Solanum igneum Linn. 178.

Solanum lanceaefolium Jacq. 176.

**Solanum lycopersicum* Linn. 177.

**Solanum Melongena* Linn. 177.

Solanum nigrum nodiflorum A.
 Gray 177.

Solanum nigrum nodiflorum
 Linn. 177.

Solanum nodiflorum Dunal 177.

Solanum racemosum Jacq. 177.

Solanum racemosum Jacq. igneum (Linn.) O. E. Schulz. 178.

Solanum torvum Sw. 178.

**Solanum tuberosum* Linn. 179.

Sonchus oleraceus Linn. 207.

Spermacoce tenuior Linn. 194.

Spigelia Anthelmia Linn. 154.

Spilanthus uliginosa Sw. 203.

Spiranthes elata L. Cl. Rich. 40.

**Spondias lutea* Linn. 119.

**Spondias purpurea* Linn. 119.

Sponia Lamarckiana Dcne 48.

Sponia micrantha Dcne 48.

Sporobolus indicus R. Br. 22.

Sporobolus virginicus Kth. 22.

Stachytarpheta indica Vahl 169.

Stachytarpheta jamaicensis

Vahl 169.

Stenandrium rupestre Nees 186.

Stenochlaena sorbifolia J. Sm. 5.

**Stenolobium stans* Seem. 182.

Stenorrhynchus lanceolatus L. Cl.
 Rich. 40.


Stenorrhynchus orchioides L. Cl.
 Rich. 40.

Sterculiaceae 131.

- Stigmatophyllon ciliatum* A. Juss. 105.
Stigmatophyllon periplocifolium A. Juss. 105.
Stigmatophyllon sericeum Wright 105.
Stigmatophyllum diversifolium A. Juss. 105.
Strumpfia maritima Jacq. 191.
Stylosanthes hamata Taubert 89.
Stylosanthes procumbens Sw. 89.
Suriana maritima Linn. 102.
 **Swietenia mahagoni* Jacq. 104.
Symplocaceae 153.
Symplocos martinicensis Jacq. 153.
Synedrella nodiflora Gärtn. 203.
Syntherisma insularis Millsp. 17.
Syntherisma sanguinale Dulac. 18.
Tabernaemontana citrifolia Jacq. 156.
 **Tabernaemontana coronaria* Willd. 156.
 **Tabernaemontana Cumingiana* A. DC. 156.
Taenitis lanceolata R. Br. 7.
Talinum paniculatum Gärtn. 65.
Talinum patens Willd. 65.
Talinum triangulare Willd. 65.
 **Tamarindus indica* Linn. 82.
 **Tecoma capensis* Lindl. 182.
Tecoma stans Juss. 182.
 **Tecoma leucoxydon* Mart. 182.
Telanthra crucis Moq. 58.
Tephrosia cathartica Urb. 88.
Tephrosia cinerea Pers. 88.
Teramnus labialis Spreng. 94.
 **Terminalia Catappa* Linn. 143.
Ternstroemia elliptica Sw. 134.
Ternstroemia peduncularis DC. 134.
Tetramicra rigida Lindl. 43.
Tetrazygia discolor DC. 149.
Theaceae 134.
Theobroma Cacao Linn. 133.
Theophrastaceae 150.
Thespesia populnea Soland. 130.
 **Thevetia neriifolia* Juss. 157.
Thrinax sp. 30.
 **Thunbergia affinis pulvinatus* Moore 184.
 **Thunbergia alata* Boj. 184.
 **Thunbergia fragans* Roxb. 184.
Thymelaeaceae 142.
Tibouchina ornata Baill. 149.
Tibouchina strigosa Cogn. 148.
Tiliaceae 125.
Tillandsia excelsa Gris. 33.
Tillandsia fasciculata Sw. 33.
Tillandsia recurvata Linn. 33.
Tillandsia usneoides Linn. 33.
Tillandsia utriculata Linn. 33.
 **Tithonia tagetiflora* Desf. 203.
Tobinia punctata Gris. 101.
Tobinia ternata Desv. 101.
Torulinium confertum Desv. 28.
Tournefortia bicolor Sw. 165.
Tournefortia caribaea Gris. 165.
Tournefortia foetidissima Linn. 165.
Tournefortia gnaphalodes R. Br. 165.
Tournefortia hirsutissima Linn. 166.
Tournefortia volubilis Linn. 166.
Tradescantia geniculata Jacq. 35.
Tragia volubilis Linn. 112.
Tragus racemosus Haller 14.
Trema Lamarckianum Bl. 48.
Trema micranthum Bl. 48.
Trianthema monogynum Linn. 65.
Trianthema portulacastrum Linn. 65.


Tribulus cistoides Linn. 99.
Tribulus maximus Linn. 99.
Tricholacna insularis Gris. 17.
Trichomanes alatum Sw. 1.
Triphasia aurantiola Lour. 102.
 **Triphasia trifoliata* P. DC. 101.
Triumfetta rhomboidea Jacq. 125.
Triumfetta semitriloba Jacq. 125.
Turneraceae 137.
Turnera ulmifolia Linn. 138.
Ucacoua nodiflora Hitchc. 203.
Ulmaceae 47.
Umbelliferae 150.
Urechites suberecta Müll.-Arg. 157.
Urena americana Linn. 129.
Urena lobata Linn. 129.
Urera alceaefolia Gaudich. 50.
Urera caracasana Gris. 50.
Urticaceae 50.
Utricularia alpina Jacq. 184.
Utricularia montana Jacq. 184.
Valerianoides jamaicensis Medic. 169.
Verbenaceae 167.
 **Verbena chamaedrifolia* Juss. 167.
Vernonia longifolia Pers. 196.
Vernonia punctata Sw. 196.
Vetiveria arundinacea Gris. 13.
Vigna glabra Sav. 89.
Vigna luteola Benth. 97.
Vigna repens O. Ktze 97.

Villamilla octandra Hook. 63.
Vinca rosea Linn. 156.
Vitaceae 124.
Vitis sicyoides Miq. 124.
Vittaria lineata Sw. 7.
Voyria aphylla Pers. 155.
Voyria uniflora Pers. 155.
Vriesea guadalupensis Mez. 34.
Waltheria americana Linn. 132.
Waltheria indica Linn. 132.
Wedelia buphthalmoides Gris. 202.
Wittmackia lingulata Mez. 32.
Xanthium orientale Linn. 201.
 **Xanthosoma* spec. 31.
Xanthoxylum clava Herculis Sic. 100.
Xanthoxylum flavum Vahl. 100.
Xanthoxylum martinicense P. DC. 100.
Xanthoxylum spinifex P. DC. 100.
Xanthoxylum ternatum Sic. 101.
Xiphopteris serrulata Kaulf. 10.
Xylosma buxifolium A. Gray 136.
Yucca aloifolia Linn. 36.
 **Zea Mays* Linn. 12.
Zebrina pendula Schnizl. 36.
 **Zephyranthes carinata* Herb. 37.
Zingiberaceae 40.
 **Zinnia elegans* Jacq. 202.
Zizyphus Jujuba Lam. 123.
Zornia diphylla Pers. 90.
Zygophyllaceae 99.





*The Dutch W.I Islands
and the Surrounding Leeward Islands.*







Boldingh. — Flora Dutch, West Ind., Islands.





Boldingh. — Flora Dutch, West Ind., Islands.





New York Botanical Garden Library
OK 238 .B6
Boldingh, Isaac/The flora of the Dutch W
ge
3 5185 00101 4065

