

## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



1  
F31F  
Ent. cop.

# FOREIGN PLANT DISEASES

A HANDBOOK OF ECONOMIC PLANT DISEASES WHICH  
ARE NEW TO OR NOT WIDELY DISTRIBUTED  
IN THE UNITED STATES

U. S. DEPARTMENT OF AGRICULTURE  
PLANT INDUSTRY BOARD

~~Bureau of Entomology and Plant Quarantine~~

Bu

UNITED STATES  
DEPARTMENT OF AGRICULTURE  
LIBRARY



BOOK NUMBER 1

F31F

~~Ent. & P.Q. Library copy.~~

1  
F31 F

Property of the  
Bureau of Entomology and Plant Quarantine  
U. S. Department of Agriculture

Library



*Handwritten signature*

UNITED STATES DEPARTMENT OF AGRICULTURE

OFFICE OF THE SECRETARY

CONTRIBUTION FROM THE FEDERAL HORTICULTURAL BOARD  
C. L. MARLATT, CHAIRMAN, IN COLLABORATION WITH  
THE BUREAU OF PLANT INDUSTRY  
W. A. TAYLOR, CHIEF

FOREIGN PLANT DISEASES

A MANUAL OF ECONOMIC PLANT DISEASES WHICH  
ARE NEW TO OR NOT WIDELY DISTRIBUTED IN  
THE UNITED STATES

PREPARED BY

JOHN A. STEVENSON

Chief Pathological Inspector, Federal Horticultural Board

*Library*  
Property of the  
Bureau of Entomology and Plant Quarantine  
U. S. Department of Agriculture



WASHINGTON  
GOVERNMENT PRINTING OFFICE

1926

ADDITIONAL COPIES

OF THIS PUBLICATION MAY BE PROCURED FROM  
THE SUPERINTENDENT OF DOCUMENTS  
GOVERNMENT PRINTING OFFICE  
WASHINGTON, D. C.

AT

25 CENTS PER COPY



## INDEX OF COMMON NAMES

Abaca.....	Musa.	Betal palm.....	Areca.
Aconite.....	Aconitum.	Betony.....	Stachys.
Aconite, winter.....	Eranthis.	Bindweed.....	Convolvulus.
Adder's-tongue.....	Erythronium.	Birch.....	Betula.
Adder's-tongue.....	Ophioglossum.	Bird-of-paradise flower.....	Strelitzia.
African cherry orange.....	Citropsis.	Bishop's cap.....	Mitella.
African lily.....	Agapanthus.	Bitter cress.....	Cardamine.
African millet.....	Eleusine.	Bittersweet.....	Celastrus.
African oil palm.....	Elaeis.	Blackberry.....	Rubus.
Agrimony.....	Agrimonia.	Blackberry lily.....	Belamacanda.
Aguacate.....	Persea.	Black bryony.....	Tamus.
Air potato.....	Dioscorea.	Black locust.....	Robinia.
Alder.....	Alnus.	Black salsify.....	Scorzonera.
Alder, white.....	Clethra.	Bladder fern.....	Cystopteris.
Alfalfa.....	Medicago.	Bladder nut.....	Staphylea.
Alfilaria.....	Erodium.	Bladder senna.....	Colutea.
Alkanet.....	Anchusa.	Bladderwort.....	Utricularia.
Alligator pear.....	Persea.	Blazing star.....	Mentzelia.
Almendro.....	Terminalia.	Bleeding heart.....	Dicentra.
Almond.....	Prunus.	Blister cress.....	Erysimum.
Almond, Spanish.....	Terminalia.	Blood leaf.....	Iresine.
Alumroot.....	Heuchera.	Blood lily.....	Haemanthus.
Alumroot, false.....	Tellima.	Blueberry.....	Vaccinium.
Amaranth.....	Amaranthus.	Blue-curls.....	Trichostema.
Amazon lily.....	Eucharis.	Blue-eyed grass.....	Sisyrinchium.
Amazon vine.....	Stigmaphyllon.	Blue grass.....	Poa.
Anatto.....	Bixa.	Bog asphodel.....	Narthecium.
Angel's-trumpet.....	Datura.	Boneset.....	Eupatorium.
Anise tree.....	Illicium.	Borage.....	Borago.
Apple.....	Malus.	Bottle brush.....	Callistemon.
Apricot.....	Prunus.	Bottle tree.....	Brachychiton.
Arborvitae.....	Thuja.	Bottle tree.....	Sterculia.
Arborvitae, false.....	Thujiopsis.	Box.....	Buxus.
Arrowhead.....	Sagittaria.	Bowstring hemp.....	Sansevieria.
Arrowroot.....	Maranta.	Boxwood.....	Maclura.
Artichoke, globe.....	Cynara.	Brake.....	Pteris.
Artichoke, Jerusalem.....	Helianthus.	Breadfruit.....	Artocarpus.
Ash.....	Fraxinus.	Bridal wreath.....	Spiraea.
Aspen.....	Populus.	Brome grass.....	Bromus.
Asphodel.....	Asphodelus.	Broom.....	Cytisus.
Aster, China.....	Callistephus.	Broom.....	Genista.
Autumn crocus.....	Colchicum.	Broomcorn millet.....	Panicum.
Avens.....	Geum.	Brussels sprouts.....	Brassica.
Avocado.....	Persea.	Bryony.....	Bryonia.
Azalea.....	Rhododendron.	Bryony, black.....	Tamus.
		Bugloss.....	Anchusa.
Baby's breath.....	Gypsophila.	Buckwheat.....	Fagopyrum.
Bachelor's-button.....	Centaurea.	Buckeye.....	Aesculus.
Bael fruit.....	Aegle.	Buckthorn.....	Rhamnus.
Balloon-flower.....	Platycodon.	Bugbane.....	Cimicifuga.
Balloon vine.....	Cardiospermum.	Bugle.....	Ajuga.
Balm.....	Melissa.	Bugle lily.....	Watsonia.
Balsam.....	Impatiens.	Burnet.....	Poterium.
Balsam apple.....	Momordica.	Burnet.....	Sanguisorba.
Bamboo.....	Bambuseae.	Bur, New Zealand.....	Acaena.
Bamboo.....	Phyllostachys.	Burning bush.....	Euonymus.
Banana.....	Musa.	Bur reed.....	Sparganium.
Baneberry.....	Actaea.	Burstwort.....	Herniaria.
Banyan.....	Ficus.	Bush clover.....	Lespedeza.
Barberry.....	Berberis.	Bush honeysuckle.....	Diervilla.
Barberton daisy.....	Gerbera.	Butcher's-broom.....	Ruscus.
Barley.....	Hordeum.	Butcher's-broom, climbing.....	Semele.
Barwood.....	Baphia.	Butterbur.....	Petasites.
Barnyard grass.....	Echinochloa.	Buttercup.....	Ranunculus.
Basil.....	Ocimum.	Butterfly bush.....	Buddleia.
Basswood.....	Tilia.	Butterfly flower.....	Schizanthus.
Bayberry.....	Myrica.	Butterfly pea.....	Clitoria.
Bay, sweet.....	Laurus.	Butterfly weed.....	Asclepias.
Beach grass.....	Ammophila.	Butternut.....	Juglans.
Bean.....	Phaseolus.	Butterwort.....	Pinguicula.
Beard grass.....	Andropogon.	Buttonbush.....	Cephalanthus.
Beardtongue.....	Pentstemon.		
Bear's-breech.....	Acanthus.	Cabbage.....	Brassica.
Beauty berry.....	Callicarpa.	Cacao.....	Theobroma.
Bedstraw.....	Galium.	Cainito.....	Chrysophyllum
Beech.....	Fagus.	Calla.....	Zantedeschia.
Beefwood.....	Casuarina.	Camomile.....	Anthemis.
Beet.....	Beta.	Camphor tree.....	Cinnamomum.
Belladonna.....	Atropa.	Campion.....	Lychnis.
Bellflower.....	Campanula.	Campion.....	Silene.
Bent grass.....	Agrostis.	Canary grass, reed.....	Phalaris.
Bermuda grass.....	Cynodon.	Candlenut.....	Aleurites.

Candytuft.....	Iberis.	Cotton.....	Gossypium.
Canistel.....	Lucuma.	Cotton thistle.....	Onopordon.
Canterbury bells.....	Campanula.	Cottonweed.....	Diotis.
Cape jasmine.....	Gardenia.	Cottonwood.....	Populus.
Carambola.....	Averrhoa.	Cow parsnip.....	Heracleum.
Caraway.....	Carum.	Cowpea.....	Vigna.
Cardamom.....	Elettaria.	Crab.....	Malus.
Cardoon.....	Cynara.	Cranberry.....	Oxycoccus.
Carnation.....	Dianthus.	Crane's bill.....	Geranium.
Carob.....	Ceratonia.	Crape jasmine.....	Tabernaemontana.
Carrot.....	Daucus.	Crape myrtle.....	Lagerstroemia.
Cashew.....	Anacardium.	Cress, water.....	Nasturtium.
Cassava.....	Manihot.	Crested dogtail.....	Cynosurus.
Cassia bark tree.....	Cinnamomum.	Crocus, autumn.....	Colchicum.
Castor bean.....	Ricinus.	Crosswort.....	Crucianella.
Catbrier.....	Smilax.	Croton.....	Codiaeum.
Catchfly.....	Silene.	Crowberry.....	Empetrum.
Catnip.....	Nepeta.	Crowfoot.....	Ranunculus.
Cat-tail.....	Typha.	Crownbeard.....	Verbesina.
Cauliflower.....	Brassica.	Crown imperial.....	Fritillaria.
Cayenne pepper.....	Capsicum.	Crown vetch.....	Coronilla.
Ceara rubber.....	Manihot.	Cubeb.....	Piper.
Cedar.....	Cedrus.	Cuckoo-flower.....	Cardamine.
Cedar, white.....	Chamaecyparis.	Cucumber.....	Cucumis.
Cedar of Lebanon.....	Cedrus.	Cupang.....	Parkia.
Celandine poppy.....	Chelidonium.	Cupflower.....	Nierembergia.
Celery.....	Apium.	Currant.....	Ribes.
Century plant.....	Agave.	Custard apple.....	Annona.
Ceriman.....	Monstera.	Cypress.....	Cupressus.
Chamomile.....	Anthemis.	Daffodil.....	Narcissus.
Chaste tree.....	Vitex.	Daisy, Barberton.....	Gerbera.
Chayote.....	Sechium.	Daisy, English.....	Bellis.
Cherimoya.....	Annona.	Daisy, Michaelmas.....	Aster.
Cherry.....	Prunus.	Daisy, Swan River.....	Brachycome.
Cherry, ground.....	Physalis.	Daisy tree.....	Olearia.
Cherry orange.....	Citropsis.	Dammar pine.....	Agathis.
Chestnut.....	Castanea.	Dame's rocket.....	Hesperia.
Chestnut, water.....	Trapa.	Dandelion.....	Taraxacum.
Chick-pea.....	Cicer.	Darnel.....	Lolium.
Chickweed, mouse-ear.....	Cerastium.	Dasheen.....	Colocasia.
Chicory.....	Chicorium.	Date palm.....	Phoenix.
Chilean crocus.....	Tecophilaea.	Dayflower.....	Commelina.
Chilean lily.....	Alstroemeria.	Day lily.....	Hemeroallis.
Chile bells.....	Lapageria.	Dead nettle.....	Lamium.
China aster.....	Callistephus.	Desert candle.....	Eremurus.
Chinaberry.....	Melia.	Devil's-tongue.....	Amorphophallus.
China fir.....	Cunninghamia.	Dewberry.....	Rubus.
Chinese laurel.....	Antidesma.	Dishcloth gourd.....	Luffa.
Chinese tallow tree.....	Sapium.	Dock.....	Rumex.
Chinese tallow tree.....	Stillingia.	Dogwood.....	Cornus.
Chinquapin.....	Castanopsis.	Douglas fir.....	Pseudotsuga.
Chive.....	Allium.	Dovewood.....	Alchornea.
Chocolate tree.....	Theobroma.	Dracena.....	Dracaena.
Christmas fern.....	Polystichum.	Dragonhead.....	Dracocephalum.
Christmas rose.....	Helleborus.	Dragon-root.....	Arisaema.
Cigar-box cedar.....	Cedrela.	Dragon tree.....	Dracaena.
Cinnamon.....	Cinnamomum.	Dropseed.....	Sporobolus.
Cinquefoil.....	Potentilla.	Dune grass.....	Elymus.
Citron.....	Citrus.	Durian.....	Durio.
Citronella grass.....	Cymbopogon.	Dutchman's-pipe.....	Aristolochia.
Cliff brake.....	Pellaea.	East Indian trumpet flower.....	Oroxylum.
Climbing hempweed.....	Mikania.	Ebony.....	Diospyros.
Clock vine.....	Thunbergia.	Eddoe.....	Colocasia.
Clove tree.....	Eugenia.	Edelweiss.....	Leontopodium.
Clover.....	Trifolium.	Eggplant.....	Solanum.
Clover, bush.....	Lespedeza.	Elder.....	Sambucus.
Clover, holy.....	Onobrychis.	Elephant's-ear.....	Colocasia.
Clover, sweet.....	Melilotus.	Elm.....	Ulmus.
Club moss.....	Selaginella.	Endive.....	Cichorium.
Cocaine tree.....	Erythroxyton.	English daisy.....	Bellis.
Cockseggs.....	Salpichroa.	Eryngio.....	Eryngium.
Cocoa plum.....	Chrysobalanus.	Eugenia.....	Caryophyllus.
Cockscomb.....	Celosia.	Evening primrose.....	Oenothera.
Coconut.....	Cocos.	Everlasting.....	Helichrysum.
Coffee.....	Coffea.	Everlasting, pearl.....	Naphalis.
Colocynth.....	Citrullus.	Fairy bells.....	Disporum.
Coltsfoot.....	Tussilago.	False alumroot.....	Tellinia.
Columbine.....	Aquilegia.	False arborvitae.....	Thujaopsis.
Comfrey.....	Symphytum.	False camomile.....	Matricaria.
Conehead.....	Strobilanthes.	False hellebore.....	Veratrum.
Congo pea.....	Cajanus.	False indigo.....	Amorpha.
Copaiba.....	Copaifera.	False mallow.....	Malvastrum.
Copper leaf.....	Acalypha.	False olive.....	Elaeodendron.
Coral blow.....	Russelia.	False Solomon's-seal.....	Smilacina.
Coral tree.....	Erythrina.	False spirea.....	Sorbaria.
Coriander.....	Coriandrum.	False tamarix.....	Myricaria.
Cork tree.....	Phellodendron.	Fan palm.....	Livistona.
Corn, Indian.....	Zea.	Featherfoil.....	Hottonia.
Corn cockle.....	Lychnis.		
Corn flower.....	Centaurea.		

Feather grass.....	Stipa.	Harebell.....	Campanula.
Fennel.....	Foeniculum.	Hat palm.....	Carludovica.
Fennel, giant.....	Ferula.	Hawaiian algarroba.....	Prosopis.
Fenugreek.....	Trigonella.	Hawkweed.....	Hieracium.
Fescue.....	Festuca.	Hawthorn.....	Crataegus.
Fig.....	Ficus.	Hazelnut.....	Corylus.
Fig marigold.....	Mesembryanthemum.	Heal-all.....	Brunella.
Figwort.....	Scrophularia.	Heartseed.....	Cardiospermum.
Filaree.....	Erodium.	Heath.....	Erica.
Filbert.....	Corylus.	Heather.....	Calluna.
Filmy fern.....	Hymenophyllum.	Hellebore, false.....	Veratrum.
Finger grass.....	Chloris.	Hellebore.....	Helieborus.
Fir.....	Abies.	Heliotrope.....	Heliotropium.
Fir, Douglas.....	Pseudotsuga.	Hemlock.....	Tsuga.
Fish poison tree.....	Piscidia.	Hempweed, climbing.....	Mikania.
Flamboyant.....	Poinciana.	Hemp.....	Cannabis.
Flat sedge.....	Cyperus.	Henbane.....	Hyoscyamus.
Flax.....	Linum.	Henequen.....	Agave.
Flax lily.....	Phormium.	Heronbill.....	Erodium.
Flax, New Zealand.....	Phormium.	Himalaya honeysuckle.....	Leycesteria.
Fleabane.....	Erigeron.	Hoarhound.....	Marrubium.
Flowering rush.....	Butomus.	Hoary pea.....	Tephrosia.
Foamflower.....	Tiarella.	Holly.....	Ilex.
Forget-me-not.....	Myosotis.	Hollyfern.....	Polystichum.
Four-o'clock.....	Mirabilis.	Hollygrape.....	Berberis.
Foxglove.....	Digitalis.	Hollyhock.....	Althaea.
Frangipani.....	Plumeria.	Holy clover.....	Onobrychis.
Fringe bell.....	Schizocodon.	Honesty.....	Lunaria.
Fringe tree.....	Chionanthus.	Honey locust.....	Gleditsia.
Fritillary.....	Fritillaria.	Honeysuckle.....	Lonicera.
Fumitory.....	Fumaria.	Honeysuckle, bush.....	Diervilla.
Gama grass.....	Tripsacum.	Honeywort.....	Cerithe.
Gamboge.....	Garcinia.	Hop bush.....	Dodonaea.
Gandul.....	Cajanus.	Hop.....	Humulus.
Gambuzo.....	Cicer.	Hop tree.....	Ptelea.
Garlic.....	Allium.	Hop hornbeam.....	Ostrya.
Gas plant.....	Dictamnus.	Horn poppy.....	Glaucium.
Gentian.....	Gentiana.	Horned rampion.....	Phyteuma.
Germander.....	Teucrium.	Hornbeam.....	Carpinus.
Geranium.....	Pelargonium.	Horse-radish.....	Radicula.
Giant fennel.....	Ferula.	Horse-radish tree.....	Moringa.
Giant reed.....	Arundo.	Horse-chestnut.....	Aesculus.
Gilliflower.....	Matthiola.	Houseleek.....	Sempervivum.
Ginger.....	Zingiber.	Huckleberry.....	Gaylussacia.
Ginger lily.....	Hedychium.	Husk tomato.....	Physalis.
Ginger, wild.....	Asarum.	Hyacinth.....	Hyacinthus.
Ginseng.....	Panax.	Hyacinth bean.....	Dolichos.
Globe amaranth.....	Gomphrena.	Icaco.....	Chrysobalanus.
Globe daisy.....	Globularia.	India almond.....	Terminalia.
Globeflower.....	Trollius.	India mulberry.....	Morinda.
Globe mallow.....	Sphaeralcea.	Indigo.....	Indigofera.
Globe thistle.....	Echinops.	Indigo, false.....	Amorpha.
Glory bower.....	Clerodendrum.	Iron tree.....	Metrosideros.
Glory bush.....	Tibouchina.	Ironweed.....	Vernonia.
Glory-of-the-snow.....	Chionodoxa.	Ivy.....	Hedera.
Glory pea.....	Clianthus.	Jaboticaba.....	Myrciaria.
Goatsbeard.....	Aruncus.	Jack bean.....	Canavalia.
Goat's-rue.....	Galega.	Jack fruit.....	Artocarpus.
Golden bell.....	Forsythia.	Jack-in-the-pulpit.....	Arisaema.
Golden chain.....	Laburnum.	Jacob's ladder.....	Polemonium.
Golden-rain tree.....	Koelreuteria.	Jacob's rod.....	Asphodeline.
Golden saxifrage.....	Chrysosplenium.	Jasmine.....	Jasminum.
Goldenrod.....	Solidago.	Jasmine, cape.....	Gardenia.
Gold-eye grass.....	Hypoxis.	Jasmine, crape.....	Tabernaemontana.
Gooseberry.....	Ribes.	Jasmine, orange.....	Chalcas.
Goosefoot.....	Chenopodium.	Jerusalem artichoke.....	Helianthus.
Gourd.....	Cucurbita.	Jerusalem sage.....	Phlomis.
Gourd.....	Lagenaria.	Jessamine.....	Jasminum.
Goutweed.....	Aegopodium.	Jet bead.....	Rhodotypos.
Gram.....	Cicer.	Jicama.....	Pachyrhizus.
Grape.....	Vitis.	Jimson weed.....	Datura.
Grapefruit.....	Citrus.	Jobo.....	Spondias.
Grape hyacinth.....	Muscari.	Job's-tears.....	Coix.
Grass of Parnassus.....	Parnassia.	Johnson grass.....	Holcus.
Grass tree.....	Xanthorrhoea.	Judas tree.....	Cercis.
Greek valerian.....	Polemonium.	Joint fir.....	Ephedra.
Greenbrier.....	Smilax.	Jujube.....	Zizyphus.
Gromwell.....	Lithospermum.	June berry.....	Amelanchier.
Ground cherry.....	Physalis.	Juniper.....	Juniperus.
Ground ivy.....	Nepeta.	Jupiter's-beard.....	Centranthus.
Groundsel.....	Senecio.	Jute.....	Corchorus.
Guava.....	Psidium.	Kafir lily.....	Clivia.
Guinea grass.....	Panicum.	Kaki.....	Diospyros.
Guinea-hen flower.....	Fritillaria.	Kale.....	Brassica.
Gum.....	Eucalyptus.	Kapok.....	Ceiba.
Gum myrtle.....	Angophora.	Kauri pine.....	Agathis.
Hackberry.....	Celtis.	Kidney vetch.....	Anthyllis.
Hair grass.....	Aira.	Knotweed.....	Polygonum.
		Kohlrabi.....	Brassica.

Kudzu bean.....	Pueraria.	Meadow rue.....	Thalictrum.
Kumquat.....	Fortunella.	Meadowsweet.....	Filipendula.
Labrador tea.....	Ledum.	Medick.....	Medicago.
Lady bell.....	Adenophora.	Medlar.....	Mespilus.
Lady's-slipper.....	Cypripedium.	Melic grass.....	Melica.
Ladies-tresses.....	Spiranthes.	Melon.....	Cucumis.
Lancepod.....	Lonchocarpus.	Mesquite.....	Prosopis.
Lansa.....	Lansium.	Mexican poppy.....	Argemone.
Larch.....	Larix.	Mexican rubber tree.....	Castilla.
Larkspur.....	Delphinium.	Michaelmas daisy.....	Aster.
Laurel.....	Laurus.	Mignonette.....	Reseda.
Lavender.....	Lavandula.	Milk vetch.....	Astragalus.
Lavender cotton.....	Santolina.	Milkweed.....	Asclepias.
Lead plant.....	Amorpha.	Millet, broomcorn.....	Panicum.
Leadwort.....	Plumbago.	Millet, pearl.....	Pennisetum.
Leek.....	Allium.	Millet.....	Setaria.
Lemon.....	Citrus.	Millet grass.....	Milium.
Lemon grass.....	Cymbopogon.	Mint.....	Mentha.
Lemon verbena.....	Lippia.	Mist flower.....	Eupatorium.
Lentil.....	Lens.	Moccasin flower.....	Cypripedium.
Leopard's-bane.....	Doronicum.	Mock cypress.....	Kochia.
Leopard flower.....	Belamacanda.	Mock orange.....	Philadelphus.
Lettuce.....	Lactuca.	Monkey comb.....	Pithecoctenium.
Lettuce, water.....	Pistia.	Monkey-face tree.....	Mallotus.
Licorice.....	Glycyrrhiza.	Monkey puzzle.....	Araucaria.
Lilac.....	Syringa.	Monkshood.....	Aconitum.
Lily.....	Lilium.	Moonflower.....	Calonyction.
Lily, African.....	Agapanthus.	Moonseed.....	Menispermum.
Lily, Amazon.....	Eucharis.	Morea.....	Moraea.
Lily, blackberry.....	Belamacanda.	Morning-glory.....	Ipomoea.
Lily, blood.....	Haemanthus.	Mosquito trap.....	Cynanchum.
Lily, Chilean.....	Alstroemeria.	Mountain ash.....	Sorbus.
Lily, day.....	Hemerocallis.	Mountain mahogany.....	Cercocarpus.
Lily, ginger.....	Hedychium.	Mourning bride.....	Scabiosa.
Lily of the valley.....	Convallaria.	Mouse-ear chickweed.....	Cerastium.
Lily, Peruvian.....	Alstroemeria.	Mulberry.....	Morus.
Lily, plantain.....	Hosta.	Mullein.....	Verbascum.
Lily, spider.....	Hymenocallis.	Muskmelon.....	Cucumis.
Lily, St. Bernard.....	Anthericum.	Mustard.....	Brassica.
Lily, water.....	Nymphaea.	Myrrh.....	Myrrhis.
Lima bean.....	Phaseolus.	Myrtle.....	Myrtus.
Lime.....	Citrus.	Naibel.....	Hesperethusa.
Lime.....	Tilia.	Napier grass.....	Pennisetum.
Linden.....	Tilia.	Nasturtium.....	Tropaeolum.
Lion's-ear.....	Leonotis.	Necklace tree.....	Ormosia.
Litchi.....	Nephelium.	Nectarine.....	Prunus.
Lizard's-tail.....	Saururus.	New Zealand bur.....	Acaena.
Locust.....	Robinia.	New Zealand flax.....	Phormium.
Locust, honey.....	Gleditsia.	New Zealand spinach.....	Tetragonia.
Loosestrife.....	Lysimachia.	Night jasmine.....	Nyctanthes.
Loosestrife, purple.....	Lythrum.	Nightshade.....	Solanum.
Loquat.....	Eriobotrya.	Nightshade, Malabar.....	Basella.
Lotus.....	Nelumbo.	Ninebark.....	Physocarpus.
Love grass.....	Eragrostis.	Niter bush.....	Nitraria.
Love-in-a-mist.....	Nigella.	Norfolk Island pine.....	Araucaria.
Lovage.....	Levisticum.	Nutmeg.....	Myristica.
Lousewort.....	Pedicularis.	Oak.....	Quercus.
Lucerne.....	Medicago.	Oat grass.....	Arrhenatherum.
Lungwort.....	Pulmonaria.	Oats.....	Avena.
Lupine.....	Lupinus.	Oil palm, African.....	Elaeis.
Lychee.....	Nephelium.	Oleander.....	Nerium.
Madar bush.....	Calotropis.	Olive.....	Olea.
Madeira vine.....	Boussingaultia.	Onion.....	Allium.
Mahogany.....	Swietenia.	Orach.....	Atriplex.
Mahogany, mountain.....	Cercocarpus.	Orange.....	Citrus.
Maidenhair fern.....	Adiantum.	Orange jasmine.....	Chalcas.
Maidenhair tree.....	Ginkgo.	Orchard grass.....	Dactylis.
Maize.....	Zea.	Oregon grape.....	Berberis.
Malabar plum.....	Caryophyllus.	Osage orange.....	Maclura.
Malabar nightshade.....	Basella.	Otaheite gooseberry.....	Phyllanthus.
Malanga.....	Xanthosoma.	Oxeye.....	Buphthalmum.
Mallow.....	Malva.	Painted cup.....	Castilleja.
Mallow, false.....	Malvastrum.	Palm.....	Palmae.
Mamey colorado.....	Lucuma.	Palmyra palm.....	Borassus.
Mandarin orange.....	Citrus.	Pansy.....	Viola.
Mandioca.....	Manihot.	Papaya.....	Carica.
Mango.....	Mangifera.	Paper mulberry.....	Broussonetia.
Mangosteen.....	Garcinia.	Paper tree.....	Edgeworthia.
Maple.....	Acer.	Papyrus.....	Cyperus.
Marigold.....	Tagetes.	Pampas grass.....	Gynerium.
Marjoram.....	Origanum.	Para grass.....	Panicum.
Marram grass.....	Ammophila.	Para rubber.....	Hevea.
Marshmallow.....	Althaea.	Paraguay tea.....	Ilex.
Marsh marigold.....	Caltha.	Parrotbeak.....	Clanthus.
Masterwort.....	Astrantia.	Parsley.....	Petroselinum.
Maté.....	Ilex.	Parsnip.....	Pastinaca.
Matrimony vine.....	Lycium.	Passion flower.....	Passiflora.
May apple.....	Podophyllum.		

Patchouli.....	Pogostemum.	Rhubarb.....	Rheum.
Pea.....	Lathyrus.	Rice.....	Oryza.
Pea.....	Pisum.	Rice, wild.....	Zizania.
Pea, butterfly.....	Clitoria.	Rock brake.....	Cryptogramma.
Pea, chick.....	Cicer.	Rock cress.....	Arabis.
Pea, scurf.....	Psoralea.	Rock jasmine.....	Androsace.
Pea shrub.....	Caragana.	Rock purslane.....	Calandrinia.
Pea, sweet.....	Lathyrus.	Rockrose.....	Cistus.
Peach.....	Prunus.	Rosary bean.....	Rhynchosia.
Peanut.....	Arachis.	Rosary pea.....	Abrus.
Pear.....	Pyrus.	Rose.....	Rosa.
Pear, alligator.....	Persea.	Rose apple.....	Jambosa.
Pearl everlasting.....	Anaphalis.	Rose mallow.....	Hibiscus.
Pearl millet.....	Pennisetum.	Rosemary.....	Rosmarinus.
Pearlwort.....	Sagina.	Rose myrtle.....	Rhodomyrtus.
Pelican flower.....	Aristolochia.	Rosewood.....	Dalbergia.
Pennycress.....	Thlaspi.	Rouge plant.....	Rivina.
Pentstemon.....	Pentstemon.	Royal fern.....	Osmunda.
Peony.....	Paeonia.	Royal poinciana.....	Poinciana.
Pepper.....	Piper.	Rubber, ceara.....	Manihot.
Pepper, cayenne.....	Capsicum.	Rubber, Mexican.....	Castilla.
Pepper, red.....	Capsicum.	Rubber, Para.....	Hevea.
Pepper tree.....	Schinus.	Rubber vine.....	Cryptostegia.
Periwinkle.....	Vinca.	Rue.....	Ruta.
Persimmon.....	Diospyros.	Russian olive.....	Elaeagnus.
Peruvian bark.....	Cinchona.	Rutabaga.....	Brassica.
Peruvian lily.....	Alstroemeria.	Rye.....	Secale.
Physic nut.....	Jatropha.	Rye grass.....	Lolium.
Pickernel weed.....	Pontederia.		
Pigeon pea.....	Cajanus.	St. Bernard's lily.....	Anthericum.
Pimpernel.....	Anagallis.	St. John's-wort.....	Hypericum.
Pine.....	Pinus.	Safflower.....	Carthamus.
Pineapple.....	Ananas.	Sage.....	Salvia.
Pink.....	Dianthus.	Sago palm.....	Cycas.
Pinkroot.....	Spigelia.	Salal.....	Gaultheria.
Pistache.....	Pistacia.	Salsify, black.....	Scorzonera.
Pitcher plant.....	Nepenthes.	Saltbush.....	Atriplex.
Planetree.....	Platanus.	Sandalwood.....	Santalum.
Plantain.....	Musa.	Sand-box tree.....	Hura.
Plaintain lily.....	Hosta.	Sandwort.....	Arenaria.
Plum.....	Prunus.	Sapodilla.....	Achras.
Plume poppy.....	Bocconia.	Sapote.....	Lucuma.
Plumeria.....	Plumeria.	Sarsaparilla.....	Aralia.
Poinsettia.....	Euphorbia.	Saxifrage.....	Saxifraga.
Pokeberry.....	Phytolacca.	Saxifrage, golden.....	Chryso-splenium.
Pokeweed.....	Phytolacca.	Screw pine.....	Pandanus.
Polypody.....	Polypodium.	Scurf pea.....	Psoralea.
Pomegranate.....	Punica.	Sea buckthorn.....	Hippophae.
Poplar.....	Populus.	Sea grape.....	Coccoloba.
Poplar, yellow.....	Liriodendron.	Sea holly.....	Eryngium.
Poppy.....	Papaver.	Sea onion.....	Urginea.
Poppy, celandine.....	Chelidonium.	Sea pink.....	Armeria.
Poppy, horn.....	Glaucium.	Sea pink.....	Statice.
Poppy, Mexican.....	Argemone.	Self-heal.....	Brunella.
Potato.....	Solanum.	Sensitive fern.....	Onoclea.
Pot marigold.....	Calendula.	Sensitive plant.....	Mimosa.
Prayer bead.....	Abrus.	Serpent root.....	Scorzonera.
Prickle poppy.....	Argemone.	Service berry.....	Amelanchier.
Prickly ash.....	Zanthoxylum.	Sesame.....	Sesamum.
Prickly pear.....	Opuntia.	Shad blow.....	Amelanchier.
Prickly thrift.....	Acantholimon.	Shadbush.....	Amelanchier.
Primrose.....	Primula.	Shallot.....	Allium.
Primrose willow.....	Jussiaea.	Shinleaf.....	Pyrola.
Privet.....	Ligustrum.	Siberian pea tree.....	Caragana.
Pumpkin.....	Cucurbita.	Silk cotton.....	Bombax.
Purple loosestrife.....	Lythrum.	Silk cotton.....	Ceiba.
Purslane.....	Portulaca.	Silk oak.....	Grevillea.
		Silk-tassel bush.....	Garrya.
Quaking grass.....	Briza.	Silk vine.....	Periploca.
Quince.....	Cydonia.	Silver climber.....	Argyrea.
Quinine.....	Cinchona.	Silver tree.....	Leucadendron.
Quinoa.....	Chenopodium.	Sisal.....	Agave.
		Skullcap.....	Scutellaria.
Rabbit-tail grass.....	Lagurus.	Sky flower.....	Duranta.
Radish.....	Raphanus.	Smartweed.....	Polygonum.
Raisin tree.....	Hovenia.	Snail seed.....	Cocculus.
Ramie.....	Boehmeria.	Snakebeard.....	Ophiopogon.
Rape.....	Brassica.	Snake gourd.....	Trichosanthes.
Raspberry.....	Rubus.	Snake palm.....	Amorphophallus.
Rattlebox.....	Crotalaria.	Snapdragon.....	Antirrhinum.
Ravenna grass.....	Erianthus.	Snapweed.....	Impatiens.
Redbud.....	Cercis.	Sneezewort.....	Achillea.
Red cedar.....	Juniperus.	Snowbell.....	Styrax.
Red pepper.....	Capsicum.	Snowberry.....	Symphoricarpus.
Redtop.....	Agrostis.	Snowdrop.....	Galanthus.
Red valerian.....	Centranthus.	Snowflake.....	Leucojum.
Reed.....	Phragmites.	Snow-in-summer.....	Cerastium.
Reed canary grass.....	Phalaris.	Soapberry.....	Sapindus.
Reed, giant.....	Arundo.	Soapwort.....	Saponaria.
Rest-harrow.....	Ononis.	Solomonseal.....	Polygonatum.
Rhodes grass.....	Chloris.		

Solomonseal, false	Smilacina.	Toothwort	Dentaria.
Sorghum	Holcus.	Touch-me-not	Impatiens.
Sorrel	Rumex.	Towel gourd	Luffa.
Soursop	Annona.	Traveler's-tree	Ravenala.
Sow thistle	Sonchus.	Tree fern	Alsophila.
Soy bean	Soja.	Tree fern	Cyathea.
Spanish almond	Terminalia.	Treemallow	Lavatera.
Spanish moss	Tillandsia.	Tree-of-heaven	Ailanthus.
Spanish oyster plant	Scolymus.	Tree tomato	Cyphomandra.
Spanish plum	Spondias.	Trout lily	Erythronium.
Spatter-dock	Nymphaea.	Trumpet bush	Tecoma.
Speedwell	Veronica.	Trumpet creeper	Bignonia.
Spiderflower	Cleome.	Tuft root	Dieffenbachia.
Spider lily	Hymenocallis.	Tulip	Tulipa.
Spiderwort	Tradescantia.	Tulip tree	Liriodendron.
Spiked loosestrife	Lythrum.	Tunic flower	Tunica.
Spikenard	Aralia.	Turmeric	Curcuma.
Spinach	Spinacia.	Turnip	Brassica.
Spinach, New Zealand	Tetragonia.	Twayblade	Liparis.
Spiraea	Spiraea.	Twinleaf	Jeffersonia.
Spleenwort	Asplenium.	Twin spur	Melasmaerula.
Spring beauty	Claytonia.	Udo	Aralia.
Spruce	Picea.	Unicorn plant	Martynia.
Spurge	Euphorbia.	Valerian	Valeriana.
Spurry	Spergula.	Vegetable oyster	Tragopogon.
Squash	Cucurbita.	Venus-looking-glass	Specularia.
Squill	Scilla.	Vetch	Vicia.
Squirreltail grass	Hordeum.	Vine spinach	Basella.
Star apple	Chrysophyllum.	Violet	Viola.
Star flower	Trientalis.	Virginia creeper	Ampelopsis.
Star glory	Quamoclit.	Wahoo	Euonymus.
Star gooseberry	Phyllanthus.	Wallflower	Cheiranthus.
Star grass	Aletris.	Walnut	Juglans.
Star grass	Hypoxis.	Wand flower	Sparaxis.
Star-of-Bethlehem	Ornithogalum.	Water chestnut	Trapa.
Sting lily	Blumenbachia.	Water cress	Nasturtium.
Stock	Matthiola.	Water cress	Radicula.
Stone cress	Aethionema.	Water lettuce	Pistia.
Stonecrop	Sedum.	Water lily	Nymphaea.
Stone mint	Cunila.	Watermelon	Citrullus.
Strawberry	Fragaria.	Water soldier	Stratiotes.
Strawberry tree	Arbutus.	Wattle	Acacia.
Sudan grass	Holcus.	Wax plant	Hoya.
Sugar apple	Annona.	Weaversbroom	Spartium.
Sugar beet	Beta.	Wheat	Triticum.
Sugar cane	Saccharum.	Wheat grass	Agropyron.
Sugar palm	Arenga.	Weigelia	Diervilla.
Sunflower	Helianthus.	White alder	Clethra.
Sumac	Rhus.	White cedar	Chamaecyparis.
Summer cypress	Kochia.	White sapote	Casimiroa.
Sunn hemp	Crotolaria.	Whitlow grass	Draba.
Sun rose	Helianthemum.	Wild calla	Calla.
Surinam cherry	Eugenia.	Wild ginger	Asarum.
Swan-River daisy	Brachycome.	Wild rice	Zizania.
Sweet alyssum	Alyssum.	Wild rye	Elymus.
Sweet bay	Laurus.	Willow	Salix.
Sweet clover	Melilotus.	Willow weed	Epilobium.
Sweet flag	Acorus.	Windflower	Anemone.
Sweetleaf	Symplocos.	Wing nut	Pterocarya.
Sweet pea	Lathyrus.	Winter creeper	Euonymus.
Sweet potato	Ipomoea.	Wisteria	Wisteria.
Sweet shrub	Calycanthus.	Winter aconite	Eranthis.
Sword bean	Canavalia.	Winter cress	Barbarea.
Sword fern	Nephrolepis.	Witch-hazel	Hamamelis.
Sycamore	Platanus.	Wintergr n	Gaultheria.
Syringa	Philadelphus.	Woodbetony	Pedicularis.
Tail grape	Artabotrys.	Wood fern	Aspidium.
Tallow tree, Chinese	Sapium.	Wood fern	Dryopteris.
Tamarack	Larix.	Wood fern	Phegopteris.
Tamarind	Tamarindus.	Wood-oil tree	Aleurites.
Tangerine	Citrus.	Woodruff	Asperula.
Tansy	Tanacetum.	Wood rush	Luzula.
Tea	Thea.	Wormwood	Artemisia.
Teak	Tectona.	Yam	Dioscorea.
Teasel	Dipsacus.	Yam bean	Pachyrhizus.
Teff	Eragrostis.	Yautia	Xanthosoma.
Teosinte	Euchlaena.	Yautia	Colocasia.
Thatch palm	Thrinax.	Yarrow	Achillea.
Thistle, sow	Sonchus.	Yellow mombin	Spondias.
Thistle, globe	Echinops.	Yew	Taxus.
Thoroughwort	Eupatorium.	Yellow poplar	Liriodendron.
Thrift	Statice.	Yellow rocket	Barbarea.
Thyme	Thymus.	Yellowwood	Cladrastis.
Tick clover	Meibomia.	Zephyr lily	Zephyranthes.
Timothy	Phleum.		
Toadflax	Linaria.		
Tobacco	Nicotiana.		
Tomato	Lycopersicum		
Tomato, husk	Physalis.		

## FOREIGN PLANT DISEASES

### A MANUAL OF ECONOMIC PLANT DISEASES WHICH ARE NEW TO OR NOT WIDELY DISTRIBUTED IN THE UNITED STATES

---

This publication has been prepared for the purpose of supplying such available information as is required by the officers of the Federal Horticultural Board and others in the enforcement of plant quarantines and in the safeguarding of this country against foreign plant diseases. It is hoped that it will prove of value not only to inspectors in actually preventing entry of specific diseases, but that it will aid in a general way in pointing out the magnitude of the quarantine problem by presenting the very great number of plant diseases reported from foreign countries which are not known in the United States.

The manual is based on the data contained in the host index of foreign plant diseases maintained in the Office of Foreign Plant Quarantines of the Federal Horticultural Board. This file, which has now reached a total of several hundred thousand entries, is planned to cover the plant pathological literature of the world from the host standpoint, and, while by no means complete as yet, is of such extent as to give an excellent basis for the work in hand. This host index was started in 1914 by R. Kent Beattie, pathologist in charge, Office of Foreign Plant Quarantines, and has been continued by him and his assistants since that time.

As a matter of convenience, an alphabetical arrangement according to their scientific names has been followed both for the hosts and the causal agents of the diseases thereunder.

The selection of host genera has in large part been based on the Standard Cyclopedia of Horticulture, those genera being included which contain one or more species of economic importance. A number of genera are included, for instance *Chenopodium* and *Amaranthus*, which consist mostly of weeds, but since some members of such genera are of value, all the foreign diseases of the entire genus are given in each case, since these diseases are in large part capable of attacking both the economic and noneconomic species.

Common names have been taken in large part from Standardized Plant Names and include only those in common and general use or those most likely to be met with by inspectors. Common names of generic significance are given in caps and small caps immediately following the generic name. Other common names of specific value only then follow in ordinary type. Common names of only local significance are omitted. The characterization of each host genus is drawn in large part from the cyclopedia already referred to.

Throughout the course of the work an attempt has been made to select the included material on a uniform basis. Species of parasitic fungi or bacteria known to occur more or less widely in the United

States have been excluded. Species not so excluded and belonging to genera the members of which are parasitic for the most part, form the basis of the present work, together with species of other more or less saprophytic genera where specific mention of parasitism has been suggested. Some parasites are included which are reported from limited localities in the United States, it being advisable to guard against their further importation. A few fungi causing diseases of major importance and of comparatively recent introduction, such as *Endothia parasitica* which causes the chestnut blight, and *Cronartium ribicola*, which causes the white pine blister rust, are also listed. These still have interest or are of importance to inspectors. The records of the plant disease survey of the Bureau of Plant Industry have been helpful in determining established American plant diseases.

Under each fungus or bacterium listed there is given a brief summary of the symptoms of the disease produced by it, together with its specific hosts and the countries from which it has been reported as occurring. Diseases due to nematodes and to unknown causes supposed to be of an infectious nature are included. Limitations of space have made it impossible to cite the literature from which the information has been drawn. The citations are in the above-mentioned host index and are available for reference to anyone interested. For like reasons it has not been feasible to include illustrations.

It has not been possible to attempt a critical examination of the fungus species listed, although a few suggestions as to synonymy are made. It is realized that probably many fungi have been included which would prove harmless if introduced into America and that similarly many have been omitted which should properly find a place. The literature is so fragmentary as to make uniform treatment exceedingly difficult, if not impossible. The nomenclature used has been for the most part that of current usage. Certain monographic works, including Salmon on Erysiphaceae, Theissen and Sydow on Dothidiaceae, Sydow on Uredineae, Gauman on Peronospora, and others have been followed.

Because of the alphabetical arrangement of the material, indices other than the common-name index have been considered unnecessary.

The manuscript has been reviewed by the offices of Plant Disease Survey, Laboratory of Plant Pathology, Pathological Collections, Fruit Disease Investigations, Forest Pathology, Sugar Plant Investigations, Cotton, Truck, and Forage Crop Disease Investigations, and Cereal Investigations of the Bureau of Plant Industry. The nomenclature of the host genera has been checked by George B. Sudworth, dendrologist of the Forest Service and by F. V. Colville and S. F. Blake, of the Office of Economic and Systematic Botany. It has been necessary for the most part to use the host names given in the literature. Obviously it was not possible to handle the subject critically. Where the generic plant name thus used differs from that now recognized by the Department of Agriculture, the department name, together with the abbreviation Ag., follows it; for example, ABERIA. DOVYALIS. Ag.

- ABERIA. DOVYALIS** Ag. Small ornamental trees or shrubs of Africa and Ceylon.  
**Asterostromella aberiae** Petch. On leaves of *A. gardneri* in Ceylon.  
**Phyllachora aberiae** P. Henn. Tar spot on leaves of *A. caffra* in the Union of South Africa.
- ABIES.** Fir. Ornamental coniferous evergreens and timber trees.  
**Barclayella deformans** (Barel.) Diet. See Picea.  
**Camarosporium abietis** Wilson. Attacks twigs and small branches of *A. lowiana* in Scotland.  
**Cucurbitaria pithyophila** Fr. See Pinus.  
**Dasycephala abieticola** Henn. and Shir. Produces a canker on trunks and limbs of *A. homolepis* (*A. brachyphylla*) and *A. firma* in Japan.  
**Dasycephala willkommii** Hart. See Larix.  
**Diplodia abiegna** Maubl. On living needles of *A. concolor* in France.  
**Fusicoccum abietinum** (Hart.) Prill. and Delacr. (*Dothiorella pithya* Sacc.) This fungus produces cankers on the branches, which die as a result of girdling. Black stromata are produced on the cankered areas. On *A. alba* (*A. pectinata*) and *Abies* sp. in France and Germany. Reported from Ohio on *A. veitchii*, an introduced ornamental type.  
**Lophodermium abietis** Rostr. See Picea.  
**Melampsora abieti-caprearum** Tub. See Salix.  
**Melampsorella symphyti** Bub. See Symphytum.  
**Mycosphaerella abietis** (Rostr.) Lind. On leaves and shoots of *A. alba*, *A. arizonica*, *A. cephalonica*, *A. nordmanniana*, and *A. pinsapo* in Denmark.  
**Oospora abietum** Oudem. A single row of minute greenish-gray tufts on each side of the median line and on both surfaces of needles of *A. nordmanniana*, *A. pinsapo*, *Picea abies*, and *Pseudotsuga taxifolia* in Holland. Defoliation may result.  
**Phomopsis pseudotsugae** Wilson. See Pseudotsuga.  
**Physalospora abietina** Prill. and Delacr. On *A. alba* (*A. pectinata*) and *Picea abies* in France.  
**Phytophthora fagi** R. Htg. See Fagus.  
**Rehmielopsis bohemica** Bub. and Kab. (*Phoma bohemica* Bub. and Kab.) The leaves of the current year are attacked soon after expanding and rapidly shriveled. They turn red at first, then become darker, and remain hanging about a year. The fungus spreads into the shoots, which die back and with some host species twist and curve. The disease is so severe that planting of silver fir (*A. pectinata*) has been discontinued to a large extent in Scotland. Other hosts are *A. alba*, *A. cephalonica*, *A. nobilis*, *A. pindrow*, and *A. pinsapo* in Scotland, Bohemia, and probably other European countries.  
**Rhizothyrium abietis** Naoumoff. On needles of *A. sibirica* in Russia.  
**Sclerotopsis piceana** (Karst.) Diet. See Picea.
- ABRUS.** ROSARY PEA. Leguminous vine common throughout the Tropics, sometimes called "prayer bead."  
**Phyllachora perforans** (Rehm.) Sacc. and Syd. Tar spot on leaves of *A. precatorius* in Porto Rico.  
**Ravenelia ornata** Syd. A rust producing yellow-brown to black pustules on lower surface of leaves of *A. precatorius* and *A. pulchellus* in India, Ceylon, and the Philippines.  
**Uredo karnbachii** P. Henn. This rust produces yellow-brown pustules on the lower (rarely on the upper) leaf surfaces of *A. precatorius* in New Guinea.
- ABUMON.** See Agapanthus.
- ABUTILON.** Malvaceous shrubs or herbs, some species cultivated.  
**Ascochyta abutilonis** Höll. Causing a leaf spot of *A. theophrasti* (*A. avicennae*) in Hungary.  
**Gloeosporium leptostromoides** Bub. Small dark-brown pustules on stems of *Abutilon* sp. in Austria.  
**Meliola molleriana** Wint. Black sooty mold on leaves of *A. striatum* and *Abutilon* sp. in Brazil and Argentina.  
**Phyllosticta abutilonis** P. Henn. Circular, gray leaf spots with brown margins on *Abutilon* sp. in Brazil.  
**Puccinia abutili** B. and Br. This rust appears as powdery brown pustules on both leaf surfaces of *A. graveolens*, *A. indicum*, *A. muticum*, and *A. sonneratianum* in Ceylon, Abyssinia, Somali coast, central and south Africa.
- ACACIA.** Tropical and subtropical ornamental leguminous trees and shrubs. Several species of economic importance for gums, resins, tanning extracts, and gum arabic. A number of species are known as "wattle."  
**Aecidium esculente** Barel. This rust distorts and renders succulent the young shoots of *A. eburnea* in India.  
**Aecidium immersum** P. Henn. The aecia of this rust occur on flesh-colored galls on young twigs of *A. abyssinica* in Eritrea.  
**Aecidium schweinfurthii** P. Henn. A rust causing galls on twigs of *A. seyal* in Somali.  
**Aecidium torquens** McAlp. Leaf rust on *A. abyssinica* in Abyssinia.  
**Ascochyta borjomi** Bond. Circular, white leaf spots with dark brown margins on *Acacia* sp. in Russia.  
**Catacauma acaciae** Theiss. and Syd. Forms numerous black, circular, shiny spots on upper surface of leaves of *A. leucophloea* in India.  
**Cercospora alemquerensis** Speg. Leaf spot on *A. alemquerensis* in Brazil.  
**Cercosporella theae** Petch. See Thea.  
**Corticium salmonicolor** B. and Br. See Citrus.  
**Hapalophragmium acaciae** Bacc. Leaf rust on *A. nubica* in Abyssinia.  
**Hapalophragmium ponderosum** Syd. and Butl. The brown telial sori of this rust occur on woody galls, 1 to 5 centimeters in diameter on twigs of *A. leucophloea* in India.  
**Marsonia acaciae** Cke. and Mass. Produces irregular spots on the phyllodes, which are pale-brown to whitish, brown-margined, 0.5 to 1 centimeter in diameter and often confluent. On *Acacia* sp. in Australia.  
**Phyllachora acaciae** P. Henn. Small, circular, shiny black spots on both sides of leaves of *A. ammentacea*, *A. farnesiana*, and *A. tortuosa* in Ecuador, Mexico, and St. Thomas.  
**Phyllachora indica** Theiss. and Syd. A similar tar-spot disease on leaves of *A. penninervis* in India, Ceylon, and adjoining islands.  
**Phyllachora parvicapsa** (Cke.) Theiss. and Syd. This fungus produces black convex stromata on leaves of *Acacia* sp. in Australia.  
**Phyllosticta acaciicola** P. Henn. Amphigenous, dark brown spots, becoming lighter and drying out on leaves of *A. ramosissima* in Germany.  
**Phyllosticta pedicellata** Speg. Leaf spot on *A. pedicellata* in Brazil.  
**Phyllosticta phyllodiorum** Sacc. On phyllodes of *Acacia* sp. in Australia.  
**Physalospora phyllodii** Cke. and Mass. On phyllodes of *A. longifolia* and *A. suaveolens* in Australia.  
**Ravenelia acaciae-melliferae** Bacc. Rust on *A. millifera* in Abyssinia.  
**Ravenelia acaciae-micranthae** Diet. A rust producing dark-brown pustules on both surfaces of leaves of *A. micrantha* in Mexico.

## ACACIA—Continued.

- Ravenelia acaciae-pennatulae** Diet. Powdery, cinnamon-brown to blackish-brown pustules on upper surfaces of leaves of *A. cochliacantha* and *A. pennatula* in Mexico.
- Ravenelia australis** Diet. and Neg. Dark-brown pustules on leaves of *A. cavenia* in Chile. What is probably the same species is reported from Texas on *A. farnesiana*.
- Ravenelia deformans** (Maubl.) Diet. This rust produces yellow to brown pustules on branchlets of *Acacia* sp. in Portuguese East Africa.
- Ravenelia escharoides** Syd. Yellow-brown to black rust pustules on leaves of *A. burkei* in the Union of South Africa.
- Ravenelia evansii** Syd. Rust similar to above on leaves of *A. robusta* in the Union of South Africa.
- Ravenelia expansa** Diet. and Holw. Cinnamon-brown to chestnut-brown rust pustules on both leaf surfaces of *A. tequilana* in Mexico.
- Ravenelia farlowiana** Diet. Chestnut-brown to dark-brown pustules on leaves of *A. anisophylla*, *A. crassifolia*, and *A. micrantha* in Mexico.
- Ravenelia formosana** Syd. Rust on *A. farnesiana* in Formosa.
- Ravenelia inornata** Diet. Yellow to chestnut-brown pustules on lower leaf surfaces of *A. horrida* in the Union of South Africa.
- Ravenelia inquirenda** Arth. and Holw. Rust on leaves of *A. bursaria* in Guatemala.
- Ravenelia macowaniana** Pazschke. A rust, the cluster-cup stage of which deforms the pods and young branches, and the other stages of which form brown pustules on the leaves of *A. horrida* and *A. seyal* in Central and South Africa.
- Ravenelia natalensis** Syd. and Evans. A rust on the branches of *A. hirtella* in the Union of South Africa.
- Ravenelia papillosa** Speg. See Albizzia.
- Ravenelia peglerae** Pole-Evans. Rust on *A. eriadenia* in the Union of South Africa.
- Ravenelia pretoriensis** Syd. Yellow-brown to black rust pustules on leaves of *Acacia* sp. in the Union of South Africa.
- Ravenelia stevensii** Arth. Rust on the leaves of *A. riparia* in Porto Rico.
- Ravenelia volkensis** P. Henn. Chestnut-brown rust pustules in diseased areas on branches of *Acacia* sp. and *A. seyal* in Central Africa.
- Septobasidium acaciae** Saw. Brown patches of feltlike fungus growth on trunks and twigs of *A. richi*, *Citrus* sp., *Glochidion obovatum*, *Melia azedarach*, *Prunus persica*, *P. salicina*, *Salix glandulosa*, and *Thea sinensis* in Formosa. Reported to have killed entire trees at times.
- Septoria martiniana** Sacc. On phyllodes of *A. longifolia* in Australia.
- Septoria melanoxyli** Wint. A leaf spot of *A. melanoxylin* in Portugal.
- Septoria mortolensis** Penz. and Sacc. On leaves of *Acacia* sp., *Eucalyptus amygdaloides*, and *Hedera algeriensis* in Italy. Reported from California.
- Sphaerophragmium silveirae** Speg. Leaf rust on *A. pedicellata* in Brazil.
- Uredo alemquercensis** Speg. Leaf rust on *A. alemquercensis* in Brazil.
- Uredo puttermansii** P. Henn. A rust appearing as brown, powdery pustules on dark-brown leaf spots of *Acacia* sp. in Brazil.
- Uromyces bicinetus** McAlp. This rust produces ruddy-brown, raised pustules on phyllodes and pods of *A. fasciculifera* in Queensland.
- Uromyces discoidens** Racib. On phyllodes of *Acacia* sp. in Java.
- Uromyces fusisporus** Cke. and Mass. Produces dark-brown to black erumpent rust pustules, solitary or in groups, on phyllodes of *A. neriiifolia*, *A. salicina*, and *Acacia* sp. in Australia.
- Uromyces hyalosporus** Saw. A rust reported from Japan and Formosa as causing a serious disease of *A. confusa* by deforming phyllodes and twigs.
- Uromyces phyllodiorum** (B. and Br.) McAlp. Rust-brown pustules on phyllodes of *A. aulacocarpa*, *A. binervata*, *A. crossicarpa*, *A. cunninghami*, *A. dallachiana*, *A. dealbata*, *A. excelsa*, *A. gonoclada*, *A. holosericea*, *A. leptocarpa*, *A. linifolia*, *A. microbotrya*, *A. neriiifolia*, *A. notabilis*, *A. penninervis*, and *A. pruinosa* in Australia.
- Uromyces schweinfurthii** P. Henn. Ferruginous rust pustules on slightly deformed areas on branches of *A. ehrenbergiana* and *A. flava* in Arabia and Abyssinia.
- Uromycladium acaciae** (Cke.) Syd. Brown rust pustules on branches, leaves and pods of *A. dealbata* in Australia.
- Uromycladium alpinum** McAlp. Rusty-brown rust pustules on both leaf surfaces, phyllodes and pods of *A. buxifolia*, *A. dallachiana*, *A. dealbata*, *A. implexa*, and *A. linifolia* in Australia.
- Uromycladium maritimum** McAlp. Dark-brown rust pustules on phyllodes and stems of *A. longifolia* in Australia and Tasmania.
- Uromycladium notabile** (Ludw.) McAlp. This rust produces large, swollen, distorted galls, yellow-brown to chocolate-brown in color, on branches, phyllodes, and pods of *A. binervata*, *A. dealbata*, *A. decurrens*, *A. elata*, *A. notabilis*, and *A. pruinosa* in Australia and Tasmania.
- Uromycladium robinsoni** McAlp. Rust pustules brown, on both surfaces of phyllodes of *A. melanoxylin* in Australia and Tasmania.
- Uromycladium simplex** McAlp. A rust which produces red-brown to dark-brown pustules on both surfaces of phyllodes and on branches of *A. pycnantha* in Australia.
- Uromycladium tepperianum** (Sacc.) McAlp. A rust-producing gall-like masses along entire length of leaves or phyllodes, and large, nearly spherical galls or broadly effused areas on branches coated with brown, powdery spores. On *A. acuminata*, *A. armata*, *A. calamifolia*, *A. diffusa*, *A. erioclada*, *A. exiophylla*, *A. flavescens*, *A. glaucoptera*, *A. hakeoides*, *A. inflexa*, *A. juniperina*, *A. longifolia*, *A. melanoxylin*, *A. myrtifolia*, *A. pendula*, *A. pycnantha*, *A. rigens*, *A. salicino*, *A. sicutiformis*, *A. spinescens*, *A. stricta*, *A. torulosa*, *A. verniciflua*, *A. verticillata*, and *A. vomeriformis*, in Australia and Tasmania, and on *Albizzia montana* in Java.
- ACAENA.** New Zealand bur. Herbaceous or partly woody trailing ornamentals.
- Catacauma acaenae** (P. Henn.) Theiss. and Syd. Black convex areas on the upper surfaces of leaves of *A. ovalifolia* in Chile.
- Ustilago acaenae** Diet. and Neg. A smut reported on the leaves of *Acaena* sp. in Chile.
- ACALYPHA.** COPPER LEAF. Tropical or greenhouse ornamental shrubs or herbs, many varieties with highly colored foliage.
- Accidium acalyphae** P. Henn. Rust on leaves of *Acalypha* sp. in the Congo.
- Cercospora profusa** Syd. Causes spots on lower surfaces of leaves of *A. australis* in Japan.
- Melampsora acalyphae** Petch. Brown rust on leaves of *A. fruticosa* in Ceylon.
- Phyllosticta briosiana** Trav. More or less circular gray spots, 2 to 8 millimeters in diameter, on upper leaf surfaces of *A. virginica* in Italy.
- Puccinia evansii** P. Henn. A rust on leaves and stems, often deforming the parts attacked. Occurs on *Acalypha* sp. in the Union of South Africa.
- Rosellinia bunodes** B. and Br. See Citrus.
- Septoria cavarae** Scalia. Olive-colored subcircular spots on leaves of *Acalypha* sp. in Sicily.
- Uredo paulistana** Speg. Rust on leaves of *Acalypha* sp. in Brazil.

- ACANTHOLIMON.** PRICKLY THRIFT. Hardy evergreen perennials.  
**Uromyces acantholimonis** Syd. Brown to black rust pustules on leaves of *A. schirasianum* in Persia and Turkestan.
- ACANTHOPANAX.** Hardy shrubs or trees with ornamental foliage.  
**Aecidium acanthopanacis** Diet. Rust on leaves of *A. spinosum* in Japan.  
**Mycosphaerella acanthopanacis** Syd. Leaf spot of *A. ricinifolium* in Japan.  
**Phyllosticta acanthopanacis** Syd. Leaf spot of *A. ricinifolium* in Japan.  
**Triphragmium thwaitesii** B. and Br. Black, powdery rust pustules occurring on yellow to brown spots, 1 to 2 centimeters in diameter on both leaf surfaces of *A. innovans*, *A. sciadophylloides*, *Akebia* sp., *Hedera stellata*, and *Heptapleurum ellipticum* in Ceylon, Java and Japan.
- ACANTHOPHOENIX.** See Palmae.
- ACANTHUS.** BEARS-BREECH. Hardy herbaceous perennials.  
**Aecidium acanthi** Lott.—Sm. Rust on leaves of *Acanthus* sp. in east Africa.  
**Cercospora acanthi** Pass. Brown spots on both leaf surfaces of *A. mollis*, *A. niger*, *A. spinosus*, and *A. spinulosus* in France, Italy, and Algeria.  
**Cercosporella compacta** Trav. Brown leaf spots on *A. spinosus* in Italy.  
**Septoria acanthi** Thuem. Circular, white spots, concentrically zoned on the upper sides of leaves of *A. mollis* in Algeria and Italy.  
**Septoria acanthina** Sacc. and Magn. Subcircular, brown leaf spots on *A. mollis* in Italy.
- ACER.** MAPLE. Shade and timber trees.  
**Ascochyta acericola** Massal. On *A. campestre* in Italy.  
**Ascochyta pallida** Kab. and Bub. Circular to irregular brown spots on both sides of leaves of *A. platanoides* in Bohemia.  
**Ascochyta velata** Kab. and Bub. On leaves of *A. platanoides* in Bohemia.  
**Cercospora acerina** Hart. Attacks the cotyledons and stems of seedlings, often killing the plants. On *A. pseudoplatanus* in France and Germany.  
**Cercosporella acerina** (Hart.) Arn. Brown spots on cotyledons of *Acer* sp. in Europe, causing death of seedlings.  
**Cylindrosporium acerellum** (Sacc.) Died. (*Septoria acerella* Sacc.) Minute, angular, white spots on upper surfaces of leaves of *A. campestre* in France and Denmark.  
**Cylindrosporium platanoidis** (Allesch.) Died. (*Septoria apatela* Allesch.) (*Septoria samarigens* Bub. and Krieg.) (*Septoria seminalis* Sacc.) Small brown leaf spots on *A. campestre* and *A. platanoides* in Europe.  
**Cylindrosporium pseudoplatani** (R. and D.) Died. (*Septoria incondita* Desm.) (*S. pseudoplatani* R. and D.) Subcircular to irregular reddish or brown spots without definite margins on leaves and cotyledons of *A. campestre*, *A. platanoides*, and *A. pseudoplatanus* in Italy, France, Denmark, and Germany.  
**Dermatea acerina** Karst. On *Acer* sp. in Europe.  
**Diaporthe longirostis** Sacc. (*Gloeosporium acerinum* West.) 'On leaves of *A. platanoides*, *A. pseudoplatanus*, and *Acer* sp. in Great Britain, Belgium, Denmark, and Austria.  
**Discomycopsis rhytismoides** J. Muell. Said to produce black spots on the leaves of *A. pseudoplatanus* in Germany. Probably *Rhytisma*.  
**Discosia artocreas** (Tode) Fr. On leaves of *A. campestre* in Italy.  
**Exoascus acericola** Mass. and f. **pseudoplatani** Mass. Occurs on leaves of *A. campestre* and *A. pseudoplatanus* in Italy.  
**Exoascus acerinus** Eliasson. Forms roughened discolored areas on leaves and branches of *A. platanoides* in Sweden.  
**Exoascus confusus** Jacz. On leaves and branches of *A. campestre* in Russia.  
**Exoascus nikkoensis** (Kus.) Sacc. and Trott. Forms grayish scurfs on pale-red or violet spots on under leaf surfaces of *A. purpurascens* in Japan.  
**Gnomonia inclinata** (Desm.) Auersw. On *A. campestre* in Italy.  
**Guignardia acerifera** (Cke.) Lind. (*Laestadia acerifera* [Cke.] Sacc.) On *A. campestre* in Great Britain.  
**Guignardia rhytismophila** Rehm. On upper surface of leaves of *A. pseudoplatanus* in Germany.  
**Laestadia pseudoplatani** Pass. On leaves of *A. pseudoplatanus* in France.  
**Leptothyrium acerinum** (Kze.) Cda. Small, circular, black fruiting bodies on leaves of *A. californicum*, *A. campestre*, *A. opulifolium*, *A. palmatum*, *A. pictum*, and *A. platanoides* in middle and western Europe, the Balkan Peninsula, Great Britain, and Japan.  
**Marsonia acerina** (West.) P. Magn. 'On *A. pseudoplatanus* in Europe.  
**Marsonia decolorans** Kab. and Bub. On leaves, causing them to die back from the tips. On *A. negundo* in Bohemia and Denmark.  
**Marsonia truncatula** Sacc. (*Marssonina truncatula* [Sacc.] Magn.) Ochraceous leaf spots on *A. campestre*, *A. monspessulanum*, *A. negundo*, and *A. tataricum* in Europe.  
**Mycosphaerella latebrosa** (Cke.) Schroet. (*Phyllosticta platanoidis* Sacc.) Leaf spot on *A. campestre*, *A. negundo*, *A. opalus*, and *A. platanoides* in Europe. Possibly occurs to a limited extent in the United States.  
**Mycosphaerella maculiformis** (Pers.) Schroet. On leaves of *A. campestre*, *A. opulifolium*, *A. platanoides*, *A. pseudoplatanus*, and *Morus alba* in France, Switzerland, Denmark, Esthonia, and Germany.  
**Myxosporium devastans** Rostr. See *Betula*.  
**Parodiella aceris** Rac. On leaves of *A. laurinum* in Java. The systematic position of this fungus is uncertain.  
**Phyllosticta aceris** Sacc. Subcircular, pale-yellow spots on leaves of *A. campestre*, *A. monspessulanum*, *A. negundo*, and *A. platanoides* in France, Denmark, Italy, and Russia. Reported from Virginia and Minnesota.  
**Phyllosticta apatela** All. and var. **perniciosa** Kab. and Bub. Large indefinite, pale-brown to gray spots on leaves of *A. platanoides* and *A. pseudoplatanus* in Bohemia and Germany.  
**Phyllosticta campestris** Pass. Small, red to red-brown leaf spots on *A. campestre* and *A. pseudoplatanus* in France and Austria.  
**Phyllosticta negundicola** Sacc. Small pale-yellow to white spots on leaves of *A. negundo* in France.  
**Phyllosticta obtusula** Sacc. and Br. On leaves of *A. campestre* in Germany.  
**Phyllosticta platanoidis** Sacc. On leaves of *A. campestre* and *A. platanoides* in Europe.  
**Phyllosticta pseudoplatani** Sacc. and var. **fallax** Sacc. and Roum. On leaves of *A. campestre*, *A. pseudoplatanus* in Italy, France, Portugal, and Germany.  
**Phyllosticta tambowiensis** Bub. and Sere. On leaves of *A. platanoides* in Russia.  
**Phytophthora fagi** Hartig. See *Fagus*.  
**Phytophthora syringae** Kleb. See *Syringa*.  
**Schizothyrium annuliforme** Syd. Large, irregular, discolored areas on leaves of *A. oblongum* in India.

**ACER**—Continued.

**Sclerotium scutellatum** A. S. Orbicular, dark-brown sclerotia, on leaves of *Acer*, *Fraxinus*, and *Populus* in northern Europe.

**Septogloeum hartigianum** Sacc. The fungus causes the young twigs to dry up, preventing the buds from swelling. The mycelium spreads in the twigs during the summer, the effects not appearing until the following spring. This maple blight attacks *A. campestre* and *A. platanoides* in Italy, Russia, and Germany.

**Septoria schirajewskii** Bub. and Sere. On leaves of *A. platanoides* in Russia.

**Syngonium insigne** Penz. and Sacc. Forms black, cushion-shaped, carbonaceous fruiting bodies on leaves of *A. laurinum* in Java.

**Taphrina polyspora** (Sorok.) Johans. This fungus produces irregular, shiny, reddish-brown to black patches on leaves of *A. pseudoplatanus* and *A. tataricum* in Bulgaria, Serbia, Hungary, Scandinavia, Russia, and Germany.

**Uncinula aceris** (DC.) Sacc. and var. **tulasnei** Fckl. Powdery mildew on leaves of *A. campestre*, *A. monspessulanum*, *A. pictum*, *A. platanoides*, *A. pseudoplatanus*, *A. rubrum*, *A. spicatum*, and *A. tataricum* in Europe, Transcaucasia, and Japan.

**ACHILLEA**. YARROW. SNEEZEWORT. Herbaceous perennials.

**Ascochyta millefolii** Oud. On stems of *A. millefolium* in Holland.

**Cercospora achilleae** Jasp. Circular to oblong brown spots on leaves of *A. macrophylla* and *A. microphylla* in Switzerland.

**Entyloma achilleae** P. Magn. A smut producing spots on leaves of *A. millefolium* in Switzerland, Sweden, Denmark, and Germany.

**Peronospora radii** De By. See Chrysanthemum.

**Laestadia ptarmicae** Karst. and Starb. On *A. ptarmica* in Finland.

**Puccinia achilleae** Cke. Rust on stems and leaves of *A. albicaulis* in Kurdistan.

**Puccinia ptarmicae** Karst. Yellow to yellow-brown rust pustules on leaves of *A. cartilaginea* and *A. ptarmica* in Belgium, Denmark, Sweden, Finland, Russia, and Germany.

**Puccinia santolinae** P. Magn. Rust on stems and leaves of *A. santolina* in Egypt.

**Ramularia ptarmicae** Lind. Small, subcircular, dark-brown leaf spots on *A. ptarmica* in Germany.

**Schizothyrium ptarmicae** Desm. This fungus appears as small, flat, black fruiting bodies on the leaves of *A. ptarmica*, bringing about a stunting of infected plants in Great Britain, France, Belgium, Denmark, Finland, Siberia, and Germany.

**Septoria ptarmicae** Passer. Causes a leaf spot of *A. ptarmica* in Italy.

**Synchytrium globosum** Schroet. See Potentilla.

**Tuberculina microstigma** Sacc. On leaves of *A. millefolium* in Germany.

**Tylenchus millefolii** Loew. This nematode produces numerous small galls and hypertrophied areas on stems of *A. millefolium* and *A. tanacetifolium* in Germany.

**ACHRAS**. Sapodilla. Fruit and ornamental tree. Furnishes the chicle of commerce.

**Phyllosticta sapotae** Sacc. Deep-brown, circular to irregular spots appearing on both leaf surfaces with ashen-brown margins. On *A. zapota* in Ceylon and the Bahamas.

**Uredo sapotae** Arth. and Johnston. Rust on leaves of *A. zapota* in Cuba and the Bahamas.

**ACIDANTHERA**. See Tritonia.**ACNISTUS**. Ornamental tropical shrubs.

**Puccinia acnisti** Arth. Rust on leaves of *A. aggregatus* in Trinidad and Costa Rica.

**ACONITUM**. MONKSHOOD. Aconite. Hardy perennial herbs.

**Colosporium aconiti** Thuem. Rust on leaves of *A. barbatum* in Siberia.

**Fusicladium aconiti** Bres. On leaves of *A. clusium* in Hungary.

**Phyllosticta aconitina** Petr. Large, usually irregular, dark olive-green or brownish-green spots on leaves and sometimes on stems of *A. moldavicum* in Bohemia.

**Puccinia lycoctoni** Fckl. Yellow to yellow-brown rust pustules on leaves of *A. lycoctonum* in France, Italy, Switzerland, Norway, Austria, Hungary, India, and Germany.

**Ramularia monticola** Speg. On leaves of *A. lycoctonum*, *A. moldavicum*, and *A. napellus* in Italy and Austria.

**Septoria aconiti** Bacc. Circular, dark-brown leaf spots with definite margins on *Aconitum* sp. in China.

**Septoria fuscomaculans** Kab. and Bub. Oblong to irregular, often confluent, dark-brown spots on lower leaf surfaces of *A. paniculatum* in Tyrol.

**Septoria lycoctoni** Speg. including var. **sibirica** Sacc. and var. **macrospora** C. Mass. Ashen, irregular spots, with dark-brown margins, on leaves of *A. lycoctonum* and *Aconitum* sp. in Italy, Russia, and Siberia.

**Septoria napelli** Speg. On leaves of *A. napellus* in Italy and Yugoslavia.

**ACOEUS**. SWEET FLAG. Aquatic plants.

**Ascochyta acori** Oud. On leaves of *A. calamus* in Holland.

**Eriospora leucostoma** B. and Br. On leaves of *A. gramineus* in Europe.

**Leptosphaeria microscopica-calami** Karst. On leaves of *A. calamus* in Italy.

**Phyllosticta acorella** Sacc. and Penz. Linear spots on both surfaces of leaves of *A. calamus* in France.

**Phyllosticta acori** Oud. Elongate black spots on leaves of *A. calamus* in Holland.

**Physoderma calami** Krieg. Subcircular to oblong brown spots on leaves of *A. calamus* in Germany.

**Physoderma tenue** Nowak. See Nymphaea.

**Uredo acori** Racib. Brown-rust pustules on leaves of *A. calamus* and *A. terrestris* in Java, India, and the Philippines.

**ACRISTA**. See Palmae.**ACROCOMIA**. See Palmae.**ACTAEA**. BANE BERRY. Herbaceous perennials.

**Actinonema actaeae** Allesch. Irregular, brown to black spots on leaves of *A. spicata* in Germany.

**Colosporium actaeae** Karst. Yellow rust pustules on leaves of *A. erythrocarpa* and *A. rubra* in Siberia.

**Didymascus kitmanoffi** Sacc. On leaves of *A. spicata* in Siberia.

**Marsonia actaeae** Bres. Leaf spot of *A. spicata* in Denmark and Germany.

**ACTINIDIA**. Ornamental woody climbers.

**Aecidium actinidiae** Syd. Rust on leaves of *A. arguta* in Japan.

**Gloeosporium** sp. Anthracnose on leaves of *A. arguta* in Japan.

**ACTINOSTEMMA**. Ornamental vines.

**Marsonia actinostemmae** F. Tassi. Circular, dirty-brown, zonate spots on both surfaces of leaves of *A. paniculatum* in Italy.

**ADENOCARPUS**. Ornamental woody shrubs.

**Aecidium bubakii** Gz. Frag. On leaves of *A. intermedius* in Spain.

- ADENOPHORA.** LADYBELL. Hardy herbaceous perennials.  
**Accidium adenophorae** Jacz. Yellow rust pustules on yellow or brown leaf spots on *A. latifolia* and *A. verticillata* in Japan and Russia.  
**Accidium adenophorae-verticillatae** Syd. Leaf rust on *A. verticillata* in Japan.  
**Septoria adenophorae** Thuem. Gray spots on leaves of *A. tricuspidata* in Siberia.
- ADEOSTYLES.** Herbaceous perennials.  
**Ascochyta adenostylis** Kab. and Bub. Large, irregular, dirty-gray to dark-brown spots on leaves of *A. albifrons* in Tyrol.  
**Cercospora septorioides** Sacc. Angular gray spots on leaves of *A. albifrons* and *A. alliariae* in Switzerland and Austria.  
**Coleosporium cacaliae** Otth. See *Cacalia*.  
**Phyllosticta adenostylis** Allesch. Subcircular, often confluent, dark-brown spots on leaves of *A. alpina* in Germany.  
**Ramularia filaris** Fres. Dull yellow, irregular leaf spots on *A. albifrons*, *A. alliariae*, *A. alpina*, and *Doronicum austriacum* in Italy, Bohemia, and Austria.  
**Uromyces cacaliae** (DC.) Ung. Yellow-brown rust pustules on *A. albifrons*, *A. alliariae*, *A. alpini*, *A. auriculata*, *A. glabra*, *A. kernerii*, and *Cacalia* spp. in Japan, Rumania, Italy, Switzerland, Austria, Hungary, and Germany.  
**Uromyces veratri** (DC.) Schroet. (*Accidium adenostylis* Syd.) See *Veratrum*.
- ADESMIA.** Herbs or shrubs.  
**Protomyces vagabundus** Speg. On stems of *A. punctata* and *Medicago denticulata* in Argentina.  
**Puccinia adesmiae** P. Henn. Rust-producing globular galls on branches of *A. trijuga* in Argentina.  
**Puccinia bergii** Speg. Cinnamon-colored rust pustules on leaves of *A. punctata* in Uruguay.  
**Uredo solitaria** Diet. and Neg. Dark-brown rust pustules on leaves of *A. radicefolia* in Chile.
- ADIANTUM.** MAIDENHAIR FERN.  
**Aphelenchus olesistus** Ritz. Bos. See *Begonia*.  
**Catharinia pazschkeana** Rehm. Greenish-gray to whitish irregular spots on leaves of *A. curvatum* in Brazil.  
**Hyalopora adianti-capilli-veneris** Syd. Yellow rust pustules on leaves of *A. capillus-veneris* in Spain, France, Italy, Austria, and Dalmatia.  
**Mycosphaerella rehmanniana** Jaap. On leaves of *A. capillus-veneris* in Dalmatia.  
**Phyllosticta adianticola** Young. Brown leaf spots on *A. tenerum* in Porto Rico.  
**Septoria filicum** (Desm.) Auersw. On leaves of *A. capillus-veneris* in Italy.  
**Uredinopsis adianti** Kom. Rust pustules on leaves of *Adiantum* sp. in Manchuria.  
**Uredo gymnogrammes** P. Henn. See *Dryopteris*.
- ADONIS.** Hardy annual and perennial herbs.  
**Accidium solms-laubachii** Boy. and Jacz. Rust on leaves of *A. aestivalis* in France.  
**Urocystis leimbachii** Oertel. A smut forming galls at the base of the stems of *A. aestivalis* in Germany
- AEGLE.** Bael fruit. Tropical fruit tree.  
**Bacterium citri** Hasse. See *Citrus*.  
**Cercospora undulata** (Ch. Bern.) Sacc. Leaf spots, blackish above, yellow beneath, on *A. odorata* and *Citrus* sp. in Java.
- AEGOPODIUM.** GOUTWEED. Herbaceous perennials.  
**Ascochyta podagrariae** Bres. On leaves of *A. podagraria* in Germany.  
**Mycosphaerella aegopodii** A. Pot. Whitish or pale spots on leaves of *A. podagraria*, widespread in Europe.  
**Peronospora podagrariae** Otth. On leaves of *A. podagraria* in Switzerland.  
**Phyllosticta podagrariae** Oud. Irregular, indefinite dark-brown spots on leaves of *A. podagraria* in Holland.  
**Protomyces macrosporus** Unger. See *Coriandrum*.  
**Puccinia aegopodii** (Schum.) Mart. Dark-brown rust pustules on leaf blades and petioles of *A. podagraria* in Europe.  
**Puccinia leioderma** Lindr. Brown powdery rust pustules on under side of leaves of *A. alpestre* and *A. tenera* in Turkestan, Siberia, and Japan.  
**Septoria aegopodii** (Pr.) Sacc. Causes spotting of leaves of *A. podagraria* in Russia.  
**Septoria aegopodina** Sacc. Angular, brown, finally white, leaf spots on *A. podagraria* in Italy and Serbia.
- AERIDES.** See *Orchidaceae*.
- AESCULUS.** HORSECHESTNUT. Buckeye. Shade trees.  
**Ascochyta aesculi** Kab. and Bub. Large, irregular, yellow to brown spots, which coalesce, destroying the leaves of *A. hippocastanum* in Bohemia.  
**Ascochyta grandimaculans** Kab. and Bub. Large, dark-yellow, irregular spots with reddish-purple margins on leaves of *A. hippocastanum* in Bohemia.  
**Phleospora parcissima** v. Hoehn. Numerous subcircular brown spots on leaves of *A. hippocastanum* in Austria and Germany.  
**Phyllosticta aesculana** Oud. Leaf spots on *A. hippocastanum* in Holland.  
**Phyllosticta aesculina** Sacc. Ochraceous leaf spots on *A. hippocastanum* in France.  
**Phyllosticta hippocastani** Oud. Indefinite brown areas on *A. hippocastanum* in Holland.  
**Phyllosticta socialis** Bub. and Kab. Small, angular, purple-brown spots on leaves of *A. hippocastanum* in Bohemia.  
**Phytophthora syringae** Kleb. See *Syringa*.  
**Septoria aesculi** (Lib.) West. Leaf spots on *A. hippocastanum* in Belgium, France, Italy, Russia, and Austria. Of limited distribution in the United States.  
**Septoria aesculina** Thuem. Leaf spots on *A. hippocastanum* in France and Austria. Said to be serious in California on *A. californica*.  
**Stereum purpureum** Pers. See *Prunus*.
- AETHIONEMA.** STONE CRESS. Herbaceous perennials.  
**Accidium aethionematis** P. Magn. Rust pustules on leaves of *A. buzbaumi* in Asia Minor.
- AEXTOXICON.** South American trees.  
**Hendersonia aetoxici** Speg. Circular, grayish spots with purple areas surrounding, on the upper surface of leaves of *A. punctatum* in Chile.  
**Phyllosticta aetoxici** Speg. Whitish spots, with narrow dark-purple borders, on leaves of *A. punctatum* in Chile.
- AFZELIA.** INTSIA. Ag. Tropical timber trees.  
**Fomes lignosus** Klotzsch. See *Hevea*.  
**Hyaloderma afzeliae** V. Keissl. Small black globular fruiting bodies in groups on leaves of *Afzelia* sp. in Samoa and the Solomon Islands.  
**Phyllachora afzeliae** Syd. Shining black convex areas on leaves of *A. bijuga* in the Philippines.  
**Trabutia stephaniae** Banc. On leaves of *A. retusa* in the Straits Settlements.

- AGAPANTHUS. ABUMON** Ag. Herbaceous perennials with tuberous root stalks. The name African lily is sometimes applied to *A. umbellatus*.  
**Ascochyta hyacinthi** Tassi. See Hyacinthus.  
**Mycosphaerella agapanthi** (K. and Cke.) Lind. Large, irregular, dark leaf spots on *A. umbellatus* in south Africa.
- AGATHIS. DAMMARA** Ag. DAMMAR PINE. Karui pine. Coniferous trees.  
**Aecidium balansae** Cor. On *A. alba* in Straits Settlements and New Caledonia.  
**Dielsiella pritzelii** P. Henn. Circular, black tar spots on leaves of *A. palmerstoni* in Queensland.
- AGAVE.** Century plant. Sisal. Henequen. Ornamental and fiber plants.  
**Bótryosphaeria agaves** (P. Henn.) Butl. Dark-brown spots, becoming pale at the center, on leaves of *A. sisalana*, in India and Tanganyika.  
**Gloeosporium agaves** Syd. Irregular, black, diseased areas which finally coalesce on leaves of *Agave* sp. in Brazil.  
**Hendersonia agaves** Maubl. Dirty-white spots with brown margins on upper surface of leaves of *Agave* sp. in Algeria.  
**Leptosphaeria agaves** Syd. and Butl. On leaves of *A. sisalana* in India.  
**Marsonia agaves** Earle. Sunken yellow spots on leaves of *Agave* sp. in Colombia.  
**Phyllosticta agaves** Maubl. Yellowish spots, with brown margins, on upper surfaces of leaves of *Agave* sp. in Algeria.  
**Septoria megaspora** Speg. On leaves of *A. americana* in Argentina.  
**Tubercularia agaves** Pat. On leaves of *Agave* sp. in Costa Rica.  
**Tuberculina concentrica** Mont. and Fr. Large brown spots on leaves of *Agave* sp. in France.
- AGERATUM.** Garden annuals.  
**Aecidium agerati** P. Henn. Rust pustules on brown leaf spots on *A. conyzoides* in central Africa.  
**Albugo brasiliensis** Speg. On leaves of *A. conyzoides* in Colombia and Brazil.  
**Puccinia rosea** (Diet. and Holw.) Arth. Brown rust pustules on leaves of *A. conyzoides*, *A. corymbosum*, *A. strictum*, *Eupatorium deltoideum*, *E. glabratum*, *E. gonzalesii*, *E. polyodon*, *E. trinervium*, *E. tubiflorum*, and *Stevia rhombifolia* in Mexico, Porto Rico, and Guatemala.  
**Uredo agerati** Mayor. Rust on leaves of *A. conyzoides* in Colombia.
- AGLAONEMA.** Variegated-leafed aroid.  
**Gloeosporium graffii** Syd. Anthracnose on leaves of *A. dorsinervium* in the Philippines.
- AGONIS.** See Leptospermum.
- AGRIMONIA. AGRIMONY.** Perennial herbs.  
**Discosia artocreas** (Tode) Fr. On leaves of *A. eupatoria* in Austria.  
**Phyllosticta agrimoniae** Lasch. On leaves of *A. eupatoria* in Denmark.  
**Ramularia agrimoniae** Sacc. White, moldlike growth on under sides of leaves of *A. eupatoria* in Siberia.
- AGROPYRON. WHEAT GRASS.**  
**Ascochyta graminum** Lasch. Pale-brown spots on leaves of *Agropyron* sp. in Germany.  
**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.  
**Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.  
**Uromyces agropyri** Barcl. Yellow-brown to brown rust pustules on leaves of *Agropyron* sp. in India.  
**Uromyces fragilipes** Tranzsch. Chestnut-brown rust pustules on leaves of *A. squarrosus* in Transcaspi (Russia).
- AGROSTIS. BENT GRASS. REDTOP.** Forage grasses.  
**Ascochyta graminicola** Sacc. See Bromus.  
**Dilophia graminis** (Fekl.) Sacc. See Triticum.  
**Endodothella helvetica** (Fekl.) Theiss. and Syd. Black, slightly raised areas on leaves of *Agrostis* sp. in Switzerland and Germany.  
**Endodothella tosenensis** (P. Henn.) Theiss. and Syd. Black fruiting bodies on leaves of *A. perennans* and *A. tenuiflora* in Japan.  
**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.  
**Leptosphaeria sparsa** (Fekl.) Sacc. See Poa.  
**Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.  
**Phyllachora fuscescens** Speg. Black, linear fruiting bodies on leaves of *Agrostis* sp. in Argentina.  
**Physoderma agrostidis** Lagerh. Oblong spots on leaves of *A. alba* in Sweden.  
**Puccinia agrostidis** Plowr. (*Aecidium aquilegiae* Pers.) Yellow to brown rust pustules on leaves of *Aquilegia alpina*, *A. atrata*, *A. atroviolacea*, *A. castellana*, *A. glandulosa*, *A. haenkeana*, *A. nigricans*, *A. pyrenaica*, *A. sternbergi*, *Agrostis alba*, *A. perennans*, *A. tenuiflora*, and *A. vulgaris* in Europe, India, and Madeira. Reported from New York.  
**Puccinia borealis** Juel. Rust with its aecial stage on *Thalictrum alpinum*, and the uredo and telia stages on *A. borealis*, *Anthoxanthum odoratum*, *Calamagrostis stricta*, and *Hierochloë odorata* in Sweden, Norway, and Iceland.  
**Puccinia moyanoi** Speg. Brown rust pustules on leaves of *A. moyanoi* in Patagonia.  
**Septoria triseti** Speg. Small, linear ashen-colored spots on leaves of *A. magellanica* and *Trisetum phleoides* in Chile.  
**Tilletia decipiens** (Pers.) Koern. A smut forming black spore masses in the ovaries and stunting the plants of *A. alba*, *A. stolonifera*, and *A. vulgaris* (*A. pumila*) in Denmark, Italy, Sweden, and Germany.  
**Tilletia separata** Kunze. Forms black spore masses in ovaries of *A. spica-venti* and *A. vulgaris* in France, Switzerland, Great Britain, Russia, and Germany.  
**Uromyces poae** Rabh. See Poa.
- AILANTHUS.** Ornamental trees, *A. altissima*, sometimes called "tree-of-heaven."  
**Ascochyta ailanthi** Boud. and Fautr. Ashen-colored spots on leaves of *A. altissima* (*A. glandulosa*) in France.  
**Cercospora ailanthi** Syd. Circular black spots, changing to brown, on both leaf surfaces of *A. altissima* in Germany.  
**Phyllosticta ailanthi** Sacc. Irregular, pale-yellow spots with reddish margins, on leaves of *A. altissima* in Italy, Japan, and China.  
**Uncinula delavayi** Pat. Powdery mildew on leaves of *Ailanthus* sp. in China.
- AIRA. HAIRGRASS.** Ornamental annual grasses.  
**Entyloma catenulatum** Rost. Oblong gray spots on leaves of *A. caespitosa* in Denmark.  
**Leptosphaeria insignis** Karst. On leaves of *A. alpina* and *Alopecurus ovatus* in Spitzbergen.  
**Leptosphaeria nigrans** (Desm.) Ces. and De N. On leaf sheaths of *A. caespitosa*, *Alopecurus pratensis*, *Avena pratensis*, *Dactylis glomerata*, and *Milium effusum* in Finland. France, Italy, and Great Britain.  
**Mycosphaerella recutita** (Fr.) Johans. Black fruiting bodies on leaves of *A. caespitosa*, *Bromus benekeni*, *B. ramosus*, *Dactylis glomerata*, and *Glyceria fluitans* in Sweden, Denmark, Great Britain, and Germany.  
**Puccinia borealis** Juel. See Agrostis.

**AIRA**—Continued.

**Tilletia airae** Blytt. Black spore masses in ovaries of *A. caespitosa* in Sweden.

**Tilletia airae-caespitosae** Lindr. Black, powdery spore masses produced in discolored areas in leaves of *A. caespitosa* in Finland.

**Tilletia flectens** Lagh. Dark-brown fruiting areas in leaves of *A. flexuosa* in Sweden.

**Uredo airae** Lagh. Rust pustules on leaves of *A. caespitosa* and *A. flexuosa* in Denmark.

**Uredo airae-flexuosae** Liro. Brown rust pustules on leaves of *A. flexuosa* in Finland.

**Ustilago grammica** B. and Br. Black spore masses occurring in short parallel transverse areas on leaves of *A. caespitosa* and *Glyceria aquatica* in Great Britain.

**AJUGA**. BUGLE. Herbaceous perennials.

**Aecidium ajugae** Syd. Yellow rust pustules in circular brown spots on leaves of *Ajuga* sp. in India.

**Phyllosticta ajugae** Sacc. and Speg. Dull-yellow spots, with brown margins, on leaves of *Ajuga* sp. and *A. reptans* in France and Italy.

**Ramularia ajugae** (Niess.) Sacc. and var. **ajugae-pyramidalis** Sacc. Subcircular brown spots on leaves of *A. genevensis*, *A. lazmanni*, *A. multiflora*, *A. pyramidalis*, and *A. reptans* in France, Switzerland, Russia, Denmark, Italy, Yugoslavia, Bulgaria, Bohemia, Austria, and Germany.

**AKEBIA**. Twining woody plants.

**Aecidium akebiae** P. Henn. Yellow rust pustules in yellow or brown leaf spots on *A. lobata* and *A. quinata* in Japan.

**Ascochyta akebiae** Bres. Circular to angular brown spots, fading out to white with black borders, on leaves of *A. quinata* in Germany.

**Triphragmium thwaitesii** B. and Br. See *Acanthopanax*.

**ALANGIUM**. Tropical trees or shrubs.

**Chaetosphaeria indica** Niessl. On leaves of *A. decapetalum* in India.

**ALBIZZIA**. Leguminous ornamental woody plants.

**Ascochyta julibrissin** F. Tassi. On pods of *A. julibrissin* in Italy.

**Diplodia cacaicola** P. Henn. See *Theobroma*.

**Endodothella albizziae** Syd. Black stromata on leaves of *A. marginata* in the Philippines

**Fomes lamaoensis** Murr. See *Hevea*.

**Fomes pseudoferreus** Wakef. See *Hevea*.

**Helminthosporium albizziae** Petch. Small spots on leaves of *A. lebbek* in Ceylon.

**Microstroma albizziae** Syd. On living leaves of *A. fastigiata* in the Union of South Africa.

**Phragmocaula viventis** (Cke.) Theiss. and Syd. Black concave areas on upper surfaces of leaflets of *Albizzia* sp. and *A. fastigiata* in the Union of South Africa.

**Phyllosticta divergens** Sacc. Subcircular pale-yellow spots, with darker margins, on leaves of *A. anthelminthica* and *A. lebbek* in Abyssinia and Porto Rico.

**Ravenelia albizziae** Diet. Brown rust pustules on leaves of *A. anthelminthica* in Abyssinia.

**Ravenelia albizziae-amarae** Bacc. Brown leaf rust on *A. amara* in Abyssinia.

**Ravenelia japonica** Diet. and Syd. Cinnamon-brown to dark-brown rust pustules on leaves and branches of *A. julibrissin* in Japan.

**Ravenelia minima** Cke. Brown rust pustules on leaves of *A. fastigiata* in Uganda and the Union of South Africa.

**Ravenelia papillosa** Speg. Rust on leaves of *A. julibrissin* and *Acacia platensis* in Argentina.

**Ravenelia sessilis** Berk. Yellow-brown to black rust pustules on leaves and pods of *A. lebbek* in Ceylon and India.

**Sphaerophragmium acaciae** (Cke.) P. Magn. Brown rust pustules on leaves of *A. lebbek* in China and India.

**Sphaerophragmium luzonicum** Yates. Rust on *A. procera* in the Philippines.

**Uredo ngamboensis** P. Henn. Rust on leaves of *A. lebbek* in east Africa and the Philippines.

**Uromycladium tepperianum** (Sacc.) McAlp. See *Acacia*.

**ALBUCA**. Bulbous plants.

**Uromyces albucacae** Kalchbr. and Cke. Cinnamon-brown rust pustules on leaves of *A. altissima*, *A. aurea*, *A. juncifolia*, and *A. minor* in Angola and the Union of South Africa.

**Ustilago albucacae** Syd. Dark-brown, powdery masses of smut spores replacing the ovaries and anthers of *Albuca* sp. in Central Africa.

**ALCHEMILLA**. Hardy herbaceous perennials.

**Gymnoconia alchemillae** Bacc. Rust on leaves of *A. pedata* in Abyssinia.

**Laestadia alchemillae** Rostr. On leaves of *A. vulgaris* in Greenland.

**Ovularia alpina** Mass. On leaves of *A. hoppeana* in Switzerland.

**Ovularia haplospora** (Speg.) Magn. Subcircular, often confluent, dull-brown spots, with brownish-purple margins, on leaves of *A. alpestris*, *A. fissa*, *A. montana*, *A. pratensis*, *A. pubescens*, *A. subcrenata*, and *A. vulgaris* in Russia, Finland, Sweden, Serbia, Bulgaria, Switzerland, Italy, Austria, Denmark, and Germany.

**Ovularia schroeteri** (Kuehn.) Sacc. Ashen leaf spots on *A. vulgaris* in France, Switzerland, Austria, and Germany.

**Peronospora alchemillae** Oth. Downy mildew on leaves of *A. vulgaris* in Europe and Siberia.

**Phyllosticta alchemillae** (Vgr.) Allesch. On leaves of *A. vulgaris* in Denmark.

**Puccinia aliena** Syd. Dark-brown powdery rust pustules on leaves of *A. pedata* in tropical Africa.

**Uromyces alchemillae** (Pers.) Lev. Yellow to brown powdery rust pustules on under sides of leaves of *A. acutangula*, *A. acutiloba*, *A. alpestris*, *A. fissa*, *A. montana*, *A. pastoralis*, *A. pratensis*, *A. pubescens*, *A. splendens*, and *A. vulgaris* in all of Europe, Greenland, and Asia Minor.

**Uromyces melosporus** (Therry.) Syd. Dark-brown rust pustules on leaves of *A. alpina*, *A. hoppeana*, *A. pedata*, *A. pentaphylla*, and *A. saratatis* in Uganda, Switzerland, Austria, and Germany.

**Venturia alchemillae** (Grev.) B. and Br. Pale spots on leaves of *A. vulgaris* in Italy, Belgium, France, Scotland, Finland, and Germany.

**ALCHORNEA**. DOVEWOOD. Tropical shrubs or trees.

**Aecidium alchorneae** Sacc. Leaf rust on *Alchornea* sp. and *A. rugosa* in the Philippines.

**Gloeosporium alchorneae** Syd. Anthracnose on leaves of *A. javanica* and *A. rugosa* in the Philippines and Amboina.

**Oliva capituliformis** (P. Henn.) Arth. Rust on leaves of *A. iricurana* and *A. latifolia* in Porto Rico, Tortola, and Brazil.

**ALETRIS**. STAR GRASS. Small hardy herbs.

**Colletotrichum aletridis** P. Henn. Anthracnose on leaves of *A. japonica* in Japan.

**Gloeosporium thuenenii** Sacc. See *Anthurium*.

**ALEURITES**. CANDLENUT. WOOD-OIL TREE. Oil-producing trees.

**Cercospora aleuritidis** Miy. Leaf spot of *A. cordata* in China.

**Gloeosporium aleuriticum** Sacc. Anthracnose of *A. moluccana* in the Philippines.

- ALLAMANDA.** Tropical shrubs and climbers.  
**Septoria allamanda** Wint. On leaves of *A. hendersoni* in Portugal.
- ALLIUM.** ONION. GARLIC. LEEK. CHIVE. SHALLOT. Bulbous herbaceous perennials, cultivated for the most part as food plants, but with some ornamental species.  
**Bacillus ceptivorus** Delacr. Said to cause a soft rot of bulbs of *A. cepa* in France.  
**Bacillus croci** Mizu. See Crocus.  
**Cercospora duddiae** Welles. Yellow, circular to irregular spots, on leaves of *A. cepa* and *A. sativum* in the Philippines. The lesions are numerous and rapidly coalesce, killing leaves, with resulting losses to the crop often as high as 50 per cent.  
**Cercospora victorialis** Thuem. Large, subcircular, greenish-black leaf spots on *A. victorialis* in Siberia.  
**Chlorospora vastatrix** Speg. Produces a "very serious" decay of bulbs of *A. cepa* and *A. sativum* in Argentina.  
**Fusariella atro-virens** (Berk.) On bulbs of *A. sativum* in Great Britain.  
**Fusarium cepae** Hanz. Found in connection with root rot and bulb decay of *Allium* in Japan.  
**Heterosporium allii-cepae** Ranoj. Produces gray, elongated spots, with greenish margins over the surface of scapes of *A. cepa* in Yugoslavia.  
**Hypochnus cucumeris** Frank. See Cucumis.  
**Melampsora allii-fragilis** Kleb. See Salix.  
**Melampsora allii-populina** Kleb. See Populus.  
**Melampsora salicis-albae** Kleb. See Salix.  
**Mycosphaerella allicina** (Auersw.) Lind. On leaves of *A. asperum* in Yugoslavia.  
**Mycosphaerella schoenoprasii** (Rabenh.) Schroet. On leaves of *A. fistulosum*, *A. moschatum*, *A. porrum*, and *A. schoenoprasum* in China, Japan, and throughout Europe.  
**Mystrosporium alliorum** Berk. Forms dark patches on bulbs of *A. sativum* in Great Britain.  
**Physoderma allii** Krieger. On leaves of *A. schoenoprasum* and *Allium* sp. in Germany.  
**Phytophthora allii** K. Saw. This fungus forms cottony, white, mycelial masses on leaves and flower stalks of *A. fistulosum* in Japan. A soft rot ensues.  
**Puccinia allii-japonici** Diet. Dark-brown, powdery, rust pustules on leaves of *A. japonicum* in Japan.  
**Puccinia permixta** Syd. Rust on leaves of *A. decipiens*, *A. moschatum*, *A. rotundum*, and *A. sphaerocephalum* in Russia.  
**Puccinia porri** (Sow.) Wint. Yellow to dark-brown rust pustules on leaves of *Allium* spp. (over 40 species are reported as hosts) in New Zealand, Formosa, Japan, Syria, and Europe. Sparingly introduced into the United States.  
**Puccinia winteriana** P. Magn. See Phalaris.  
**Sclerotinia bulborum** (Wak.) Rehm. On *A. sativum* and other species in Great Britain. Probably not distinct from the following species.  
**Sclerotium cepivorum** Berk. This fungus causes a serious rot of onion bulbs. Small black fruiting bodies (sclerotia) appear on the surface of diseased bulbs. The disease attacks *A. ascalonicum*, *A. cepa*, *A. porrum*, and *A. sativum* in New Zealand, Argentina, Great Britain, Spain, France, Italy, Holland, and Germany. Confused in American literature with *Botrytis allii* Munn. Definitely reported from Oregon and Virginia.  
**Septoria alliacea** Cke. On leaves of *Allium* sp. in India.  
**Septoria alliiicola** Baum. On leaves of *A. flavum* and *A. sphaerocephalum* in Spain.  
**Septoria allii-striatelli** Speg. Pale, indefinite spots on leaves of *A. striatellum* in Argentina.  
**Septoria ranojevicii** Bub. On leaves of *A. moschatum* in Yugoslavia.  
**Stemphylium allii** Oud. On leaves of *Allium* sp. in Holland.  
**Tylenchus dipsaci** Kuehn. See Narcissus.  
**Urocystis colchici** (Schlecht) Rab. See Colchicum.  
**Urocystis magica** Pass. Black masses of smut spores formed in leaves of *A. magicum* in Italy.  
**Uromyces ambiguus** (DC.) Lev. Yellow to black powdery rust pustules on leaves and stems of *A. rotundum*, *A. schoenoprasum*, *A. scorodoprasum*, and *A. sphaerocephalum* in Turkey, Yugoslavia, Russia, Finland, Sweden, Denmark, Holland, France, Austria, and Germany.  
**Uromyces durus** Diet. Rust-forming hard, black, crustlike masses of telial spores on leaves of *A. nipponicum* in Japan.  
**Uromyces primaverilis** Speg. Linear brown rust pustules on leaves of *A. striatellum* in Argentina.  
**Uromyces reticulatus** (Thuem.) Bub. Yellow to dark-brown rust pustules on *A. victorialis* in Japan, Siberia, Spain, Portugal, France, Switzerland, Austria, and Hungary.  
**Ustilago allii** McAlp. Black spore masses in pustules on bulbs of *A. cepa* in Australia.  
**Ustilago ceparum** Glow. Black spore masses in bulbs of *A. cepa* in Germany.
- ALNUS.** ALDER. Ornamental woody plants and timber trees.  
**Exoascusalni-glutinosa** Tub. On *A. glutinosa* in Italy, Denmark, and Sweden.  
**Exoascus epiphyllus** Sad. Causes a witches'-broom and forms a gray fruiting layer on leaves, which are distorted, of *A. glutinosa* and *A. incana* in Russia, Switzerland, Portugal, Sweden, Bulgaria, Denmark, Italy, and Austria.  
**Gloeosporium alneum** West. Anthracnose on leaves of *A. glutinosa* and *A. incana* in Albania, Yugoslavia, and Austria.  
**Gnomonia setacea** (Pers.) Ces. and De N. On leaves of *A. glutinosa* in Italy.  
**Gnomoniella albomaculans** Neger. Chalky-white areas on leaves of *A. incana* in Norway.  
**Gnomoniella tubiformis** (Tode.) Sacc. On leaves of *A. glutinosa* in Italy, Russia, and Germany.  
**Hyphoderma roseum** (Pers.) Fr. See Populus.  
**Leptothyrium alneum** (Lév.) Sacc. On leaves of *A. glutinosa*, *A. incana*, and *A. viridis* in Europe.  
**Melampsoridiumalni** (Thuem.) Kleb. Yellow to brown rust pustules on lower leaf surfaces of *A. acuminata*, *A. firma*, *A. glauca*, *A. incana*, *A. jorullensis*, and *A. viridis* in Ecuador, Guatemala, Great Britain, Japan, and Siberia. Also reported from California.  
**Mycosphaerella alnobetulae** Jaap. On leaves of *A. viridis* (*A. alnobetula*) in Switzerland.  
**Passalora alnobetulae** Jaap. On leaves of *A. viridis* in Austria.  
**Passalora bacilligera** (Mont.) Fr. Subcircular spots, yellow-brown above, gray-green below, on leaves of *A. viridis* and *A. glutinosa* in Austria.  
**Phyllosticta alnea** Oud. Pale rust-colored spots on leaves of *A. glutinosa* in Holland.  
**Phyllosticta alnicola** C. Mass. Small, crowded, angular, often confluent, rust-brown spots on leaves of *A. glutinosa* in Russia and Italy.  
**Phyllosticta alnigena** Thuem. Large, yellow-brown, circular, concentrically zoned spots on leaves of *A. cordifolia* and *A. glutinosa* in Italy and Portugal.  
**Phyllostictaalni-glutinosa** Syd. Irregular, yellow-brown spots, 0.5 to 1 centimeter in diameter, on leaves of *A. glutinosa* in Italy, Russia, and Germany.  
**Phyllosticta alniperda** Oud. Circular to angular brown spots, with black margins, on leaves of *A. glutinosa* in Holland.  
**Phytophthora syringae** Kleb. See Syringa.

## ALNUS—Continued.

- Ramularia alnicola** Cke. Circular gray leaf spots on *A. glutinosa* in Russia and Great Britain.
- Rhytisma nervale** (Alb. and Schw.) Rehm. Linear rough, black fruiting areas on leaves of *Alnus* sp., *Betula* sp., and *B. alba* in Portugal.
- Sclerotinia alni** Maul. Forms sclerotia in place of seed in catkins of *A. glutinosa* and *A. incana* in Denmark, Germany, and Russia.
- Septoria alni** Sacc. Subcircular ochraceous spots on upper leaf surfaces of *A. glutinosa* in France and Italy.
- Septoria alnicola** Cke. Brown circular leaf spots on *A. glutinosa* in Great Britain and Italy.
- Septoria alnigena** Sacc. On leaves of *A. cordifolia* and *A. glutinosa* in Italy.
- Septoria cariscolensis** Kab. and Bub. Circular to angular, often confluent, purplish-brown spots with red-brown margins on leaves of *A. viridis* in Italy.
- Taphrina japonica** Kus. Produces a witches'-broom effect on *A. japonica* in Japan. The diseased branches are triangular in cross section. Leaves on the broomed branches appear early but are soon checked in their growth and remain small and pale in color until they wither and fall. The mycelium of the fungus overwinters in the buds.
- Taphrina sadebeckii** Johans. Circular yellowish or white leaf spots on *A. glutinosa*, *A. incana*, and *A. tinctoria* in Europe.
- Taphrina viridis** Sadeb. Forms leaf spots on *A. viridis* in Switzerland.
- Uncinula miyabei** (Salm.) Sacc. and Syd. White powdery mildew on leaves of *A. incana*, *A. japonica*, *A. maritima*, and *A. pendula* in Japan.
- ALOCASIA.** Tropical aroids used as decorative foliage plants.
- Caecoma anthurii** Har. var. **alocasiae** Racib. Rust on leaves of *A. metallica* in Java.
- Gloeosporium thumenii** (Von Th.) Sacc. See Anthurium.
- Mycosphaerella alocasiae** Syd. Leaf spot on *A. indica* and *A. macrorhiza* in Java and the Philippines.
- Phyllosticta portiana** Sacc. On leaves of *A. portei* in the Philippines.
- Uredo alocasiae** P. Henn. Yellow rust pustules on leaves of *Alocasia* sp. in New Guinea.
- ALOE.** Acaulescent perennial succulents.
- Dothidea alocicola** P. Henn. On leaves of *Aloe* sp. in Abyssinia.
- Gloeosporium affine** Sacc. See Hoya.
- Macrophoma aloës** Scalia. On leaves of *A. latifolia* in Italy.
- Montagnella maxima** Mass. Raised (6 to 8 mm.) circular black areas, 4 to 5 centimeters broad, on leaves of *A. ferox* and *A. platylepis* in the Union of South Africa.
- Montagnella uberata** Sacc. Gall-like fruiting areas on leaves of *A. abyssinica* and *A. flavovirens* in Abyssinia and Eritrea.
- Phyllosticta aloës** Kalchbr. On leaves of *A. latifolia*, *A. purpurascens*, and *A. volkensi* in Spain and the Union of South Africa.
- Septoria bellynckii** West. Leaf spot of *A. variegata* in Belgium.
- Uromyces aloës** (Cke.) P. Magn. Brown rust pustules on large sunken spots on leaves of *A. abyssinica*, *A. arborescens*, *A. baumi*, *A. eru.*, *A. maculata*, *A. saponaria*, *A. schimperii*, *A. spicata*, *A. transvaalensis*, and *A. vera* in India and Africa.
- ALPINIA.** Tropical decorative plants of the ginger family.
- Catacauma alpiniae** (Sacc. and Berl.) Theiss. and Syd. Black, slightly raised stromata on long elliptical whitish spots on leaves of *A. coerula* in Australia.
- Catacauma renalmiae** (Rehm.) Theiss. and Syd. Tar spot on leaves of *A. antillarum* and *Renealmia* sp. in Porto Rico and Brazil.
- Cercospora alpiniae** Syd. Leaf spot on *Alpinia* sp. in the Philippines.
- Leptosphaeria alpiniae** Maubl. Leaf spot on *A. nutans* in Brazil.
- ALSINE.** Herbaceous annuals, mostly weeds, but a few species cultivated in rock gardens. See, also, *Stellaria* and *Arenaria*.
- Septoria alsines** Rostr. On leaves and stems of *A. verna* in Denmark.
- Septoria vandassii** Bub. Blackened areas on branches, leaves, bracts, and calyces of *A. glomerata* in Bulgaria.
- Uromyces alsines** Tranzsch. Rust on leaves of *A. setacea* in Russia.
- ALSOPHILA.** TREE FERN.
- Monorhizina filicina** (B. and Br.) Theiss. and Syd. Large black circular to irregular crustlike fruiting areas on leaves of *Alsophila* sp. in Ceylon and Queensland.
- Pachypatella alsophilae** (Rac.) Theiss. and Syd. See *Cyathea*.
- Phyllosticta alsophilae** Syd. Subcircular brown spots, becoming pale at the center, on leaves of *Alsophila* sp. in Brazil.
- Polystomella pulchella** (Speg.) Theiss. On leaves of *Alsophila* and *Blechnum* in Brazil and Australia.
- ALSTONIA.** PALA Ag. Ornamental trees.
- Ascochyta alstoniae** P. Henn. Circular to oblong pale spots, with brown margins, on leaves of *A. scholaris* in Brazil.
- ALSTROEMERIA.** Tuberos-rooted herbs, sometimes called "Peruvian lily" and "Chilean lily."
- Puccinia alstroemeriae** Syd. Produces light-brown to dark-brown rust pustules on dead irregular areas on both surfaces of leaves of *A. revoluta* in Chile.
- Scolecotrichum alstroemeriae** Allesch. On the lower surfaces of leaves of *Alstroemeria* sp. in Brazil.
- Uromyces alstroemeriae** (Diet.) P. Henn. Yellow sunken, circular to elliptic spots on both leaf surfaces, in which appear yellow-brown to deep-brown rust pustules. The hosts are *A. aurantiaca*, *A. isabellana*, *A. revoluta*, and *A. subrosulacea* in Brazil and Chile.
- ALTERNANTHERA.** Herbaceous bedding plants.
- Laestadia cephalariae** (Awd.) Sacc. and var. **alternantherae** Sacc. Small, pale, sunken spots on leaves of *A. sessilis* and *Cephalaria alpina* in Germany.
- ALTHAEA.** MARSHMALLOW HOLLYHOCK. Annual, biennial, or perennial herbs.
- Cercospora nebulosa** Sacc. On leaves and stems of *A. rosea* in Italy.
- Erysiphe taurica** Lévl. Powdery mildew on leaves of *Acanthophyllum glandulosum*, *Alhagi camelorum*, *A. maurorum*, *Althaea ficifolia*, *A. kurdica*, *Arctium*, *Artemisia dracuncululus*, *Astragalus* sp., *Capparis spinosa*, *Carduus crispus*, *Carlina corymbosa*, *C. lanata*, *Carthamus lanatus*, *Carinthe major*, *Chondrilla juncea*, *Cicer*, *Clematis songorica*, *Cnicus arvensis*, *C. cardunculus*, *C. lanceolatus*, *Coccinea dubia*, *Cynara cardunculus*, *Daucus maximus*, *Eryngium campestre*, *E. noëanum*, *Euphorbia lanata*, *Ezochorda alberti*, *Foeniculum vulgare*, *Hedysarum falconeri*, *Helianthemum oelandicum*, *Inula nervosa*, *Nepeta podostachys*, *Odontospermum aquaticum*, *Phlomis herba-venti*, *P. tuberosa*, *Picris hieracioides*, *Psoralea drupacea*, *Saussurea*, *Scutellaria multicaulis*, *Taraxacum montanum*, *Teucrium chamaedrys*, *Verbascum blattaria*, *V. phlomooides*, *V. speciosum*, *Vicia tenuifolia*, and *Zygophyllum fabago* in Europe, Algeria, Turkey, Syria, Persia, Turkestan, and India.
- Phyllosticta althaeicola** Passer. Definite subcircular to irregular, often confluent, whitish spots on leaves of *A. officinalis* in France.

**ALTHAEA**—Continued.

**Phyllosticta althaeina** Sacc. Irregular brown spots on leaves of *A. rosea* in France and Italy. Reported from Ohio.

**Phyllosticta destructiva** Desm. Subcircular pale-yellow leaf spots, with darker margins, on *A. rosea*, *Euonymus europaeus*, *Euonymus* sp., *Lycium barbarum*, *Malva neglecta*, *M. nigrescens*, *M. sylvestris*, *Malva* sp., and *Menyanthes* sp. in Great Britain, France, Denmark, Belgium, Italy, Russia, Austria, and Germany. The diseased areas fall out, causing a "shot-hole" effect.

**Phyllosticta puccinospila** Mass. Small whitish spots on leaves of *A. rosea* in Italy.

**Puccinia heterogena** Lagh. See *Malva*.

**Septoria althaeae** Thuem. Pale brown spots on leaves of *A. rosea* in Bohemia.

**Septoria heterochroa** Desm. See *Malva*.

**Septoria parasitica**. Fautr. White spots on upper surfaces of leaves of *A. rosea* and *Althaea* sp. in France.

**ALYSSUM**. SWEET ALYSSUM. Lowherbs, cultivated in gardens.

**Peronospora alyssi calycini** Gäum. Downy mildew on leaves of *A. calycinum* in Switzerland and Germany.

**Peronospora alyssi incani** Gäum. As above in Bohemia.

**Puccinia alyssi** Lindr. Dark-brown rust pustules on leaves and stems of *A. halimifolium* and *A. spinosum* in Italy and Spain.

**Septoria alyssi** Brcs. Small, subcircular to oblong pale spots on leaves of *A. saxatilis* in Germany.

**AMARANTHUS**. AMARANTH. Coarse annuals; some species grown for colored foliage and showy flower clusters.

**Ascochyta amaranthi** Allesch. Subcircular to irregular pale or white spots with brown margins on leaves of *Amaranthus* sp. in Germany.

**Coniothyrium amaranti** (Fl. Tassi.) Sacc. and D. Sacc. Circular spots on leaves of *A. caudatus* in Italy.

**Peronospora amaranthi** Gäum. Downy mildew on leaves of *A. bliti* in Switzerland and Holland.

**Phyllosticta moderiana** Thuem. Large dull-yellow leaf spots, with dark-red margins, on *A. melancholicus* in Portugal.

**Phyllostictiella amaranti** Fl. Tassi. On leaves of *A. caudatus* in Italy.

**Rhizoctonia napi** West. See *Brassica*.

**AMARYLLIS**. Bulbous plants.

**Accidium amaryllidis** Syd. and Butl. Yellow rust pustules appear on long sunken spots on the lower surfaces of leaves, more rarely on the upper, of *Amaryllis* sp. (cultivated forms) in India.

**Phyllosticta amaryllidis** Brcs. Small black spots which become elongated or confluent are produced on the leaves of *Amaryllis* sp. in Brazil.

**AMBROSINIA**. Dwarf perennial bulbous herbs.

**Entyloma dietelianum** Bub. Black spore masses in red-brown areas on leaves of *A. bassii* in Sardinia.

**AMELANCHIER**. SHAD BLOW. JUNE BERRY. SHADBUSH. SERVICE BERRY. Trees and shrubs grown for ornament and fruit.

**Coleopuccinia sinensis** Pat. See *Cotoneaster*.

**Gymnosporangium amelanchieris** (DC.) Ed. Fisch. See *Juniperus*.

**Ochropsora ariae** (Fekl.) Syd. See *Sorbus*.

**Phyllosticta mespili** Sacc. var. **amelanchieri** Jacz. On leaves of *A. rotundifolia* (*A. vulgaris*) in Russia.

**AMMOPHILA**. BEACH GRASS. MARRAM GRASS. Sand binding grasses.

**Ascochyta perforans** (Rob.) Sacc. On leaves of *A. arenaria* in Belgium.

**Heterosporium graminis** McAlp. On leaves and culms of *A. arenaria* in Australia.

**Leptosphaeria littoralis** Sacc. On leaves and stems of *A. arenaria* and *Calamagrostis* sp. in Belgium and Germany.

**Mycosphaerella lineolata** (Desm.) Schroet. See *Calamagrostis*.

**Septoria ammophila** Syd. Long, narrow spots, limited by the veins, on leaves of *A. arenaria* in Germany.

**Thecaphora ammophilae** Oud. Black powdery spore masses on leaves of *A. arenaria* in Holland.

**Uredo ammophilae** Syd. Golden-yellow rust pustules over entire leaf surfaces of *A. arenaria* in Germany.

**Uredo ammophilina** Kleb. Brown rust pustules on leaves of *A. arenaria* in Europe.

**AMORPHA**. FALSE INDIGO. LEAD PLANT. Ornamental shrubby plants, grown for flowers and foliage.

**Systemma amorphae** (Rabh.) Theiss. and Syd. On *A. fruticosa* in Germany.

**AMORPHOPHALLUS**. DEVIL'S-TONGUE. Sometimes known as "snake palm." Giant aroids.

**Cercospora amorphophalli** P. Henn. Dull-brown circular spots on leaves of *A. campanulatus* and *A. morphophallus* sp. in Java and the Philippines.

**Cercospora chevalieri** Sacc. Circular to angular, scattered or confluent, white spots on leaves of *A. morphophallus* sp. in Uganda.

**Hypochnus cucumeris** Frank. See *Cucumis*.

**Puccinia pauluia** Syd. Rust pustules on leaves of *A. campanulatus* in the Philippines.

**Rhizoctonia destruens** Tass. See *Brassica*.

**AMPELOPSIS**. VIRGINIA CREEPER. Ornamental woody vines.

**Cercospora vitis-heterophyllae** P. Henn. On leaves of *A. heterophylla* in Japan.

**Phakopsora ampelopsidis** Diet. and Syd. Golden-yellow to brown rust pustules on under surfaces of leaves of *A. heterophylla*, *A. japonica*, *A. lecoides*, *Vitis flexuosa*, and *V. inconstans* in Japan and China.

**Phyllosticta allescheri** Syd. Circular, fuscous-brown spots on leaves of *A. quinquefolia* and *A. tricuspidata* (*A. veitchii*) in Germany.

**Phyllosticta quinquefoliae** Allesch. Irregular gray spots, becoming confluent on both surfaces of leaves of *A. hederacea* in Germany.

**Phyllosticta speschnewiana** Sacc. and Syd. Small circular to irregular spots on both surfaces of leaves of *A. hederacea* in Transcaucasia.

**AMSONIA**. Perennial herbs.

**Phyllosticta amsoniae** Fl. Tass. Small subcircular to irregular spots on leaves of *A. salicifolia* in Italy.

**AMYGDALUS**. See *Prunus*.**ANACARDIUM**. CASHEW. Tropical fruit and shade trees.

**Atichia millardeti** Racib. See *Cinnamomum*.

**Dendrodochium paraense** Vinc. On leaves of *A. occidentale* in Brazil.

**Fusarium udum** Butl. See *Dianthus*.

**Parodiella melioides** (B. & C.) Wint. On leaves of *Anacardium* sp. in Brazil.

- ANAGALLIS.** PIMPERNEL. Annual or perennial herbs.  
**Peronospora anagallidis** Schroet. Downy mildew on lower sides of leaves of *A. arvensis* and *A. coerulea* in Germany.  
**Peronospora candida** Fckl. White effused fungus patches (downy mildew) on lower surfaces of leaves of *A. arvensis*, *A. coerulea*, and *A. phoenicea* in Tunis, Great Britain, France, Switzerland, Denmark, Belgium, Italy, and Germany.
- ANANAS.** PINEAPPLE.  
**Asterinella stuhlmanni** (P. Henn.) Theiss. Brown spots on leaves extending rapidly to cover entire area with black superficial fruiting bodies. On *A. sativus* in Tanganyika, Philippines and India.  
**Bacterial rot.** A rot of pineapple (*A. sativus*) slips and suckers, thought to be due to bacteria, has caused considerable loss in plantings in Costa Rica.  
**Diplodia cacaoicola** P. Henn. See Theobroma.  
 "New disease." A disease of *A. sativus* in the Philippines and Queensland, due to an unknown cause, is characterized by an hypertrophy of the tissues, giving a rough corrugated appearance. Many downward suckers develop a heart rot. A similar disease is known in Hawaii.  
**Thielaviopsis paradoxa** (De Seyn.) v. Hoeh. See Saccharum.
- ANAPHALIS.** PEARL EVERLASTING. Hardy, herbaceous border plants.  
**Miyagia anaphalidis** Miy. Leaf rust on *A. margaritacea* and *A. yedoensis* in Japan.
- ANARRHINUM.** Biennial and perennial herbs.  
**Septoria anarrhini** Syd. Small circular or elongated spots on leaves of *A. bellidifolium* in Portugal.
- ANCHUSA.** BUGLOSS. Sometimes, but erroneously, called "alkanet." Hardy biennial or perennial herbs.  
**Ramularia anchusae** Massal. Circular fuscous spots on leaves of *A. barrelieri*, *A. italica*, *A. officinalis*, and *Echium vulgare* in Serbia, Bulgaria, Italy, Denmark, Russia, and Germany.  
**Ramularia anchusae-officinalis** Elias. Irregular dark-brown spots, with indefinite sunken margins, on leaves of *A. officinalis* in Sweden.
- ANDIEA.** VOUCAPOUA Ag. ANGELIN. Tropical shade trees.  
**Cercospora stevensii** Young. Small leaf spots, reddish-brown above, black beneath, on leaves of *Andira* sp. in Porto Rico.  
**Physalospora andirae** Stevens. Conspicuous, irregular, tan-colored spots on both surfaces of leaves of *A. inermis*, (*A. jamaicensis*) in Porto Rico.  
**Polystigma pusillum** Syd. Small circular to irregular yellowish spots on leaves of *A. excelsa* in Guatemala.  
**Ravenelia goyazensis** P. Henn. Dark-brown rust pustules on leaves of *A. pisonis* in Brazil.
- ANDROMEDA.** See Pieris and Zenobia.
- ANDROPOGON.** BEARD GRASS. Ornamental grasses. See also Holcus and Cymbopogon.  
**Balansia andropogonis** Syd. Abortive inflorescences of *A. aciculatus* in India.  
**Balansia sessilis** P. Henn. Forming black sessile subglobose or pulvinate stromata on culms of *Andropogon* sp. in Africa.  
**Cerebella andropogonis** Ces. Black spore masses in spikelets of *A. pertusus* in Ceylon.  
**Cintraetia columellifera** (Tul.) McAlp. Dark-brown spore masses in ovaries, enclosed at first by glumes and ovary walls, finally exposed. On *A. australis*, *A. hirtus*, and *Pennisetum cenchroides* in Australia, Mauritius, and Madeira.  
**Claviceps pusilla** Ces. Ergot on *Andropogon* sp. in Italy.  
**Endodothella andropogonis** (P. Henn.) Theiss. and Syd. Black elliptical stromata on both leaf surfaces of *Andropogon* sp. in the Congo.  
**Endodothella dispar** Syd. Black elliptical stromata on both surfaces of leaves of *A. contortus* in India.  
**Entyloma obesum** Syd. Elongate smut pustules on lower leaf surfaces of *A. annulatum* in India.  
**Ephelis pallida** Pat. Gray or black stromata, deforming spikelets of *A. aciculatus* in Tonkin and the Philippines.  
**Leptosphaeria herpotrichoides** De Not. See Triticum.  
**Phaeodothis fallax** (Sacc.) Theiss. and Syd. Long black stromata on both leaf surfaces of *A. gryllus* and *A. ischaemum* in Italy, Yugoslavia and Austria.,  
**Phyllachora andropogonicola** Speg. Small black erumpent fruiting bodies on leaves of *A. sec-caratus* in Argentina.  
**Phyllachora andropogonis** Karst. and Har. Tar spot on leaves of *A. candolleana*, *Andropogon* sp., and *Cymbopogon confertiflorus* in Timor, Ceylon, and the Philippines.  
**Phyllachora assimilis** Theiss. and Syd. Linear black stromata on both leaf surfaces of *A. assimilis* in India.  
**Physalospora festucae** (Lib.) Sacc. On leaves of *Andropogon* sp., *Arundo* sp., *Festuca* sp., *Holochaenus* sp., and *Phragmites communis* in France, Italy, and Germany.  
**Puccinia andropogonicola** Har. and Pat. Linear brown to black rust pustules on both leaf surfaces of *Andropogon* sp. in the Congo.  
**Puccinia andropogonis-macranthi** Diet. Rust on leaves of *A. macranthus* in Japan.  
**Puccinia andropogonis-micranthi** Diet. Brown rust pustules on leaves of *A. micranthus* in Japan.  
**Puccinia cesatii** Schroet. Cinnamon-brown to dark-brown rust pustules on leaves of *A. angustifolius*, *A. annulatus*, *A. gryllus*, *A. hirtus*, *A. ischaemum* and *A. pubescens* in Albania, Yugoslavia, Dalmatia, Rumania, Bulgaria, Russia, Portugal, France, Switzerland, Italy, Austria, Hungary, and Germany.  
**Puccinia chrysopogi** Barcl. Yellow rust pustules in sunken yellow spots on leaves of *Jasminum humilis* and brown pustules on leaves of *A.* (*Chrysopogon*) *gryllus* in India.  
**Puccinia citrata** Syd. Brown leaf rust on *A. citratus* in the Philippines.  
**Puccinia duthiae** Ell. and Tr. Rust pustules on leaves of *A. intermedius* and *A. pertusus* in India.  
**Puccinia erythraeensis** Pазschke. Rust on leaves of *A. commutatum* and *Andropogon* sp. in Abyssinia.  
**Puccinia hookeri** Syd. Brown rust pustules on both sides of leaves of *A. echinulatus* in north India.  
**Puccinia kojukensis** Diet. Brown to black rust pustules on leaves of *Andropogon* sp. in Japan and India.  
**Puccinia nakanishikii** Diet. Brown, linear, powdery rust pustules on leaves of *A. intermedius*, *Cymbopogon confertiflorus*, *C. schoenanthus*, and *C. nardus* in Ceylon, India, Japan, and the Philippines.  
**Puccinia posadensis** Sacc. and Trott. Linear brown rust pustules on leaves, culms and sheaths of *A. condensatus* in Argentina.  
**Puccinia propinqua** Syd. and Butl. Rust on leaves of *Andropogon* sp. in India.  
**Puccinia prunicolor** Syd. and Butl. Rust pustules on purple sunken spots on leaves and sheaths of *A. seriatus* in India.  
**Puccinia pusilla** Syd. Black rust pustules on yellow or brown indefinite spots on leaves of *A. assimilis* in India.

**ANDROPOGON**—Continued.

- Puccinia venestula** Arth. Brown leaf rust on *A. brevifolius* in Porto Rico and Costa Rica.
- Puccinia versicolor** Diet. and Holw. Brown to black rust pustules on leaves of *A. contortus* and *A. melanocarpus* in Argentina, Mexico, and India.
- Septoria grylli** Sacc. On leaves of *A. gryllus* in Italy.
- Sorosporium andropogonis-aciculati** Petch. Smut forming black spore masses in inflorescence of *A. aciculatus* in Ceylon and the Philippines.
- Sorosporium dembianense** Bacc. Smut on *A. rufus*, *A. papillipes*, and *A. arrhenobasis* in Abyssinia.
- Sorosporium geminellum** Syd. and Butl. Elongate dark-brown smut sori replacing ovaries of *Andropogon* sp. in India.
- Sorosporium icosiense** Maire. Smut on *A. hirtus* in French north Africa.
- Sorosporium maranguensis** P. Henn. Black powdery spore masses in inflorescences of *A. lepidus* in tropical Africa.
- Sorosporium tembuti** P. Henn. and Evans. Black spore masses in ovaries of *A. cymosum* in the Union of South Africa.
- Sorosporium wildemanianum** P. Henn. Smut sori, with ashen coverings, formed in ovaries and stamens of *Andropogon* sp. in the Congo.
- Thecaphora berkeleyana** Fisch. Spherical black smut sori in spikelets of *A. perforatus* in Ceylon.
- Tilletia calospora** Pass. Smut on *A. agrestis* in Italy.
- Tolyposporium philippinense** Syd. Black powdery smut sori replacing ovaries of *A. contortus* in the Philippines.
- Uredo andropogonis-hirti** R. Maire. Brown rust pustules on small sunken brown spots on leaves of *A. hirtus* and *A. pubescens* in France, Spain, and Greece.
- Uredo andropogonis-lepidi** P. Henn. Elongate yellow rust pustules on leaves of *A. lepidus* in tropical Africa.
- Uromyces andropogonis-annulati** Syd. and Butl. Brown rust pustules on leaves of *A. annulatus* in India.
- Uromyces elignyi** Pat. and Har. Brown rust pustules on leaves of *A. hirtiflorus*, *A. proximus*, and *Andropogon* sp. in Niger Territory, Abyssinia, and Guatemala.
- Ustilago amadelpa** Syd. and Butl. Black, powdery, smut sori in panicles of *Andropogon* sp. in India.
- Ustilago andropogonis-aciculati** Petch. Smut on *A. aciculatus* in the Philippines and Ceylon.
- Ustilago andropogonis-annulati** Bref. Powdery, dark brown, spore masses in ovaries of *A. annulatus* in India.
- Ustilago andropogonis-finitimi** Maubl. Black spore masses in inflorescences of *A. finitimus* in east Africa.
- Ustilago andropogonis-hirtifolii** P. Henn. Smut sori in the spikelets, protected at first by long cylindrical, cinnamon-colored membranes which rupture, exposing the black spore masses. On *A. hirtifolius* in Mexico.
- Ustilago andropogonis-tuberculati** Bref. Compact, hard, black sori in ovaries of *A. tuberculatus* in India.
- Ustilago bicornis** P. Henn. Sori black, in panicles of *A. bicornis* in Brazil.
- Ustilago congensis** Syd. Powdery, black smut sori, 1 to 2 centimeters long, in ovaries of *Andropogon* sp. in the Congo.
- Ustilago culmiperda** Schroet. Smut sori in culms of *A. bicornis* in Brazil.
- Ustilago dinteri** Syd. Sori developing in inflorescences, 1 to 5 centimeters long, inclosed by gray or brown membranes, which rupture, exposing dark fuscous or black spore masses, on *A. papillosus* in southwest Africa.
- Ustilago duthiei** Ricker. Smut sori, olive-brown, deforming the spikelets of *A. bladhii* in India.
- Ustilago effusa** Syd. Black, smut sori in sheaths, culms, and unexpanded leaves of *A. muricatus* in India.
- Ustilago filiformis** P. Henn. Long smut sori (up to 4 centimeters) destroying ovaries of *A. contortus* in east Africa.
- Ustilago furcata** Pat. and Har. Powdery, dark chestnut-colored sori, protected by white membranes, in ovaries of *Andropogon* sp. in west Africa.
- Ustilago globulifera** Sacc. and Trott. Hard, black, smut sori in inflorescences of *Andropogon* sp. in the Congo.
- Ustilago guaranitica** Speg. Smut on *A. condensatus* in Brazil and Argentina.
- Ustilago ischaemonoides** P. Henn. Black smut sori, covered at first with ashen membranes, destroying the inflorescences of *Andropogon* sp. in the Congo.
- Ustilago lanigeri** P. Magn. Black smut sori destroying inflorescences of *A. lanigerus* in Persia.
- Ustilago leucostachys** P. Henn. Sori in panicles of *A. leucostachys* in Brazil, yellowish at first, then black, due to the rupturing of the protecting membranes.
- Ustilago nyassal** Syd. Dark-brown smut sori in inflorescences of *Andropogon* sp. in Nyassa.
- Ustilago occulta** P. Henn. Black smut masses in culms of *Andropogon* sp. in Brazil.
- Ustilago stuhlmanni** P. Henn. Powdery, black smut masses on *Andropogon* sp. in central Africa.
- Ustilago superflua** Syd. Smut sori destroying ovaries of *A. foveolatus* in India.
- Ustilago tenuis** Syd. Long, dark brown sori in inflorescences of *A. pertusus* in India.
- Ustilago tonkinensis** P. Henn. Sori in red-brown spots on both surfaces of leaves of *Andropogon* sp. in Tonkin.
- Ustilago tumefaciens** P. Henn. Smut sori gall-like, causing abortion of inflorescences of *A. rufus* in tropical Africa.
- Ustilago vanderysti** P. Henn. Black, powdery sori destroying spikelets of *Andropogon* sp. in the Congo.
- Ustilago warneckeana** P. Henn. Dark-brown smut sori, deforming and destroying inflorescences of *A. contortus* in Camerun and India.

**ANDROSACE**. ROCK JASMINE. Rock plants.

- Peronospora agrorum** Gäum. Downy mildew on leaves of *A. septentrionalis* in Sweden.
- Puccinia volkartiana** Ed. Fisch. Dark-brown rust pustules on leaves and stems of *A. chamaejasma* in Switzerland.
- Puccinia dubyi** Muell. Rust on leaves of *A. alpina*, *A. glacialis*, and *A. laggeri* in Switzerland.
- Septoria androsaces** Pat. Sunken black spots on leaves of *A. rotundifolia* in China.
- Ustilago androsaces** Karst. Powdery, dark-brown smut sori in ovaries of *Androsace* sp. in Siberia.

**ANEMONE** (including *Pulsatilla*). Hardy perennials, sometimes called "windflower."

- Aphelenchus olesistus** Ritz. Bos. See Begonia.
- Ascochyta anemones** Kab. and Bub. Circular to irregular, yellow to dark-brown leaf spots on *A. nemorosa* in Bohemia.
- Coleosporium pulsatillae** Lev. Golden rust pustules in brown sunken spots on leaves of *A. cernua*, *A. halleri*, *A. montana*, *A. patens*, *A. pratensis*, *A. vulgaris*, and on the needles of *Pinus silvestris* in Japan, Siberia, Russia, Switzerland, Denmark, France, Italy, Austria, Hungary, and Germany.

**ANEMONE**—Continued.

**Micropeltis wettsteinii** v. Hoehn. Black fruiting bodies on upper surfaces of leaves of *A. wettsteinii* in Brazil.

**Ochropsora ariae** (Fekl.) Syd. See Sorbus.

**Phyllosticta anemones** P. Brun. Circular to irregular, gray or gray-brown, spots on leaves of *A. japonica* in France.

**Phyllosticta nigro-maculans** Sacc. Black spots, often involving entire leaf area, on *A. nemorosa* in France, Spain, and Italy.

**Physoderma deformans** Rostr. On *A. nemorosa* and *A. ranunculoides* in Denmark.

**Puccinia japonica** Diet. Chestnut-colored rust pustules on leaves and petioles of *A. flaccida* in Japan.

**Puccinia schelliana** Thuem. Powdery, dark-brown rust pustules on yellow leaf spots of *A. narcissiflora* in Russia.

**Puccinia singularis** P. Magn. Powdery, cinnamon-brown rust pustules, on pale circular sunken spots of *A. altaica* and *A. ranunculoides* in Japan, Yugoslavia, Rumania, Spain, France, Denmark, Austria, and Hungary.

**Puccinia vesiculosa** Schlecht. Powdery, brown rust pustules, in circular spots, on leaves of *A. narcissiflora* in Alaska.

**Ramularia pulsatillae** Höll. Dull yellow, indefinite spots, generally near the tips of leaves of *A. (Pulsatilla) nigricans* in Hungary.

**Sclerotinia tuberosa** (Hedw.) Fekl. Attacks the roots of *A. nemorosa* in Great Britain, Bohemia, Russia, and Austria, transforming them into hard, black sclerotia resembling the roots.

**Septocylindrium anemones** Delacr. Large, dirty, white spots on leaves of *A. sylvestris* and *A. sylvatica* in France.

**Septoria altaica** Thuem. On leaves of *A. altaica* in Siberia.

**Septoria brunaudii** Sacc. and Syd. Large, subcircular to irregular, gray or ashen spots, with narrow brown margins, on leaves of *A. japonica* in France.

**Septoria ficariae** Desm. Ashen spots with brown margins on leaves of *A. nemorosa* and *A. ranunculoides* in Denmark, France, Belgium, and Great Britain.

**Septoria silvicola** Desm. On leaves of *A. nemorosa* in Russia, France, and Belgium.

**Synchytrium anemones** Wor. Small galls on under sides of leaves and stems of *A. nemorosa*, *A. ranunculoides*, *A. silvestris*, and *A. virginiana* in Europe and North Africa. Also collected once in Iowa on *A. cylindrica*.

**ANGELICA**. Perennial herbs.

**Mycosphaerella angelicae** (Fr.) A. Pot. On *A. montana* in Switzerland and Italy.

**Phyllachora angelicae** (Fr.) Fekl. On leaves of *A. decursiva* and *A. sylvestris* in Japan, Siberia, Switzerland, France, and Finland. A doubtful species reported by Theissen as probably only an immature *Mycosphaerella*.

**Puccinia angelicae-edulis** Miy. Powdery brown to black rust pustules, on blades and petioles of leaves of *A. matsumurae*, *A. miqueliana*, *A. polyclada*, *A. polymorpha*, *A. shikokiana*, *A. ursina*, and *A. vulgata* in Japan.

**Puccinia angelicae-mamillata** Kleb. Brown rust pustules on leaves and petioles of *A. sylvestris*, *Polygonum bistorta* and *P. viviparum* in Helvetia and Italy.

**Puccinia angelicicola** P. Henn. Black rust pustules on yellow spots on leaves of *A. miqueliana* in Japan.

**Puccinia karstenii** Lindr. Brown rust pustules on leaves and petioles of *A. sylvestris* in Sweden, Denmark, and Finland.

**Puccinia miyabeana** Miy. Dark-brown rust pustules on leaves of *A. hakonensis*, *A. kiusiana*, and *A. utilis* in Japan.

**Puccinia psoroderma** Lindr. Brown rust pustules on *A. purpurascens* in the Caucasus.

**Ramularia angelicae** v. Hoeh. On leaves of *A. sylvestris* in Spain, Austria, and Denmark.

**Ramularia archangelica** Lindr. White spots with brown borders on leaves of *A. archangelica* in Lapland.

**Septoria bondarzewii** P. Henn. Yellow leaf spots on *A. sylvestris* in Russia.

**ANGOPHORA**. GUM MYRTLE. Trees or shrubs.

**Gloeosporium angophorae** F. Tassi. Anthracnose on branches and petioles of *A. costata* (*A. lanceolata*) in Australia.

**ANGRAECUM**. See Orchidaceae.**ANISACANTHUS**. Ornamental shrubs.

**Puccinia anisacanthi** Diet. and Holw. Brown rust pustules on leaves of *Anisacanthus* sp. in Mexico.

**ANNONA**. SOURSOP. CUSTARD APPLE. CHERIMOYA. SUGAR APPLE. Tropical fruit trees.

**Aecidium anonae** P. Henn. Yellow rust pustules on leaves of *Annona* sp. in Brazil.

**Ascochyta cherimolae** Thuem. Irregular white or yellow leaf spots, on *A. cherimola* in Ecuador and Portugal.

**Colletotrichum anonicola** Speg. Subcircular, gray spots, with fuscous margins, on leaves of *A. cherimola* in Argentina.

**Corticium salmonicolor** B. and Br. See Citrus.

**Fomes lamaoensis** Murr. See Hevea.

**Phyllachora atro-maculans** Syd. Dark-brown, circular to irregular, sunken-leaf spots on *Annona* sp. in Costa Rica. Black opaque stromata are produced on the spots.

**Phyllosticta anonae** P. Henn. Circular to oblong, pale spots, with dark-brown margins, on leaves of *Annona* sp. in Brazil.

**Phyllosticta cherimoliae** Alm. and Cam. Elongate gray spots, with brown margins, on leaves of *A. cherimola* in Portugal.

**Phyllosticta insularum** Sacc. Causes leaves to die, beginning at margins and extending irregularly inward. On *A. muricata* in the Philippines.

**Pocosphaeria anonae** Rangel. On leaves of *A. reticulata* in Brazil.

**Uredo cherimoliae** Lagh. Light-brown, effused spore masses on lower surfaces of leaves of *A. cherimola*, *A. reticulata*, *A. squamosa*, and *Rollinia multiflora* in Ecuador, Cuba, Trinidad, and sparingly in Florida.

**ANOMATHECA**. See Lapeyrouisia.**ANTHEMIS**. CAMOMILE, sometimes spelled "chamomile." Annual, biennial and perennial herbs.

**Peronospora anthemidis** Gäum. Downy mildew on leaves of *A. austriaca* and *A. cotula* in Norway, Holland, Denmark, Bohemia, Austria, and Germany.

**Peronospora radii** De By. See Chrysanthemum.

**Puccinia anthemidis** Syd. Dark-brown, rust pustules on stems and leaves of *A. altissima* in France.

**Puccinia baumleriana** Bub. Dark-brown, rust pustules on leaves of *A. tinctoria* in Hungary.

**Ramularia anthemidis** Höll. Indefinite spots on leaves of *A. ruthenica* in Hungary.

**ANTHERICUM.** ST. BERNARD'S LILY.

**Accidium antherici** P. Henn. and Evans. Yellow pustules on effuse oblong yellow spots on leaves of *Anthericum* sp. in the Union of South Africa.

**Puccinia asphodeli** Duby. See *Asphodelus*.

**ANTHOLYZA.** Cormous plants related to *Gladiolus*.

**Accidium antholyzae** Bub. Yellow rust pustules on leaves of *A. aethiopica* in the Union of South Africa.

**Uromyces antholyzae** Syd. This rust produces dark-brown pustules on both leaf surfaces of *A. abyssinica* in Abyssinia.

**ANTHOXANTHUM.** Aromatic grasses.

**Dilophia graminis** (Fekl.) Sacc. See *Triticum*.

**Helminthosporium dematioideum** Bub. and Wrobl. On leaves of *A. odoratum* in Bohemia. Reported from New York.

**Puccinia anthoxanthi** Fekl. Brown to dark-brown rust pustules on leaves, leaf-sheaths and stems of *A. odoratum* in Great Britain, Belgium and Germany.

**Puccinia borealis** Juel. See *Agrostis*.

**Puccinia fujiensis** S. Ito. Brown rust pustules on leaves of *A. japonicum* in Japan.

**Septoria oxyspora** Penz. and Sacc. On leaves of *A. odoratum*, *Arundo donax*, and *Hierochloa borealis* in Italy and Denmark.

**Uredo anthoxanthina** Bub. Powdery, yellow rust pustules on leaves of *A. odoratum* in Great Britain, Switzerland, and Germany.

**ANTHURIUM.** Ornamental tropical aroids, much cultivated in greenhouses.

**Colletotrichum anthurii** Delacr. Dark-brown spots on leaves of *Anthurium* sp. in France.

**Gloeosporium anthurii** Allesch. On leaves and petioles of *A. martianum* in Germany. Probably the same as the following species.

**Gloeosporium thumenii** (von Th.) Sacc. Black, oblong acervuli on leaves of *Aletris fragrans*, *Anthurium leuconeurum*, *Alocasia cucullata*, *A. violacea*, *Carludovica atrovirens*, *Dracaena draco*, *Dieffenbachia contorta* and *Smilax atrovirens* in France, Denmark, and Austria.

**Mycosphaerella anthurii** Miles. Large, irregular, white areas on leaves of *A. acaule* in Porto Rico. The diseased areas become dry and papery, often dropping out.

**Phyllachora engleri** Speg. Black, circular, scattered, slightly raised, shining stromata on leaves of *A. dominicense*, *A. scandens*, *Philodendron* sp. and *Spathycarpa* sp. in the West Indies, Central and South America.

**Phyllosticta cavarae** Trinch. Irregular, yellow to brown spots, with ashen brown or dirty white distinct margins, on leaves of *A. crassinervium* in Italy.

**Physalospora inanis** (Schw.) Sacc. On leaves of *Anthurium* sp. and other aroids in Ecuador and Surinam.

**Uredo anthurii** Har. Rust on leaves of *A. scandens* in Porto Rico and France.

**ANTHYLLIS.** KIDNEY VETCH. Perennial herbs, including ornamental and forage plants.

**Ascochyta vulnerariae** Fekl. Purple spots on leaves of *A. vulneraria* in Portugal and Germany.

**Cercospora radiata** Fekl. Dull-brown spots on leaves of *A. dilleni* var. *dalmatica* and *A. vulneraria* in Spain, Italy, Switzerland, Denmark, Russia, and Germany.

**Ramularia schulzeri** Baeuml. Dull-yellow to reddish spots, on leaves of *A. vulneraria*, *Lotus corniculatus*, and *L. uliginosus* in France, Austria, Hungary, and Germany.

**Septoria anthyllidis** Sacc. Brown spots, becoming whitish, on leaves of *A. dilleni* and *A. vulneraria* in Dalmatia, Sardinia, Russia, Austria, and Hungary.

**Septoria henryana** Trav. Circular to irregular, grayish-white spots with yellow-gray margins, on leaves of *A. vulneraria* in Italy.

**Typhula trifolii** Rostr. See *Trifolium*.

**Uromyces anthyllidis** Grev. Cinnamon-brown to dark-brown rust pustules on leaves of *A. dilleni*, *A. maritima*, *A. polyphylla*, *A. tetraphylla*, *A. vulneraria*, *Hippocrepis bicontorta*, *H. ciliata*, *H. multisiliquosa*, *H. unisiliquosa*, *Lotus aegyptiacus*, *L. cytisoides*, *L. ornithopodioides*, *Lupinus albus*, *L. luteus*, *Medicago arabica*, *M. hispida*, *M. orbicularis*, *Trigonella foenum-graecum*, *T. monspeliaca*, and *T. occulta* in Madeira, Tripoli, Tunis, Morocco, Egypt, India, and Europe.

**ANTIDESMA.** CHINESE LAUREL. Tropical trees.

**Cronartium antidesmae-dioicae** Syd. Leaf rust on *A. ghaesembilla* in the Philippines.

**Guignardia fusco-coriacea** Rehm. On leaves of *A. bunius* and *Antidesma* sp. in the Philippines.

**Phyllosticta ghaesembillae** Koord. Subcircular, solitary or confluent, brown spots, becoming whitish, on leaves of *A. ghaesembilla* in Java.

**Puccinia aequalis** P. Henn. Yellow-brown to dark-brown rust pustules on leaves of *A. venosum* and *Antidesma* sp. in tropical Africa.

**Uredo antidesmae** Rac. Golden-yellow rust pustules on leaves of *A. bunius* in Java.

**Uredo antidesmae-dioicae** Rac. Brown rust pustules on circular sunken spots on leaves of *A. dioica* and *A. ghaesembilla* in Java and the Philippines.

**ANTIRRHINUM.** SNAPDRAGON. Flower-garden and greenhouse herbs.

**Ascochyta decipiens** Pass. Dark-brown spots on stems and branches of *A. majus* in Italy.

**Cercospora** sp. Shot-hole disease. Minute, buff-colored spots appear on the lower leaves, increasing in size rapidly and spreading to the upper leaves, calyces, seed pods and stems of younger shoots, the latter being girdled causing them to die back rapidly. The spots are more or less circular in shape, with a well defined margin, and are surrounded by a ring of discolored tissue. The central areas, which are slightly sunken, dry out and fall, giving a shot hole effect. On cultivated snapdragon (*A. majus*) in Great Britain.

**Cercospora antirrhini** Wakef. Small, sunken patches on leaves and stems of *A. majus* in Great Britain. These patches have a dark-green, water-soaked appearance at first, later becoming dull yellow.

**Fusarium udum** Butl. See *Dianthus*.

**Peronospora antirrhini** Schroet. Downy mildew on leaves of *A. majus* and *A. orontium* in Denmark, Switzerland, and Germany.

**Phyllosticta linariae** Sacc. Subcircular brown spots, becoming whitish, with narrow reddish margins on leaves of *A. hispanicum* and *Linaria clatine* in France.

**Pseudomonas antirrhini** Takimoto. Bacterial leaf spot of *A. majus* in Japan.

**Septoria antirrhini** Desm. Yellow spots on stems and leaves of *A. majus* in Great Britain, France, Italy, Portugal, Dalmatia and Australia.

**AOTUS.** Leguminous shrubs.

**Cronartium jacksoniae** P. Henn. See *Jacksonia*.

**APHELANDRA.** Tropical ornamental shrubs.

**Accidium aphilandrae** P. Henn. Circular dull-brown spots, with dark margins, on leaves of *Aphelandra* sp. in Brazil.

**APIUM. CELEBI** Ag. Celery.**Bacillus apii** Mig. Bacterial rot of *A. graveolens* in Italy.**Bacillus apiovorus** Worm. Reported as the cause of a heart rot of celery (*A. graveolens*) in Great Britain. Probably not distinct from *B. caratovorvus* Jones.**Cercospora pastinacae** Karst. On leaves of *A. graveolens* in Finland.**Phlyctaena magnusiana** (Allesch.) Bres. On leaves of *A. graveolens* in Russia, Italy and Germany. Reported from New York.**Puccinia apii** Desm. Powdery, cinnamon-brown to dark-brown rust pustules on leaves and petioles of *A. graveolens*, *A. prostratum*, and *Selinum japonicum* in Tasmania, India, Japan, Spain, Italy, Finland, Sweden, Denmark, Switzerland, France, Holland, Belgium, Great Britain, Austria, Hungary, and Germany.**Puccinia thumeni** (Thuem.) McAlp. Cinnamon-brown to dark-brown rust pustules on both surfaces of leaves of *A. graveolens* and *A. prostratum* in Victoria and Tasmania.**Septoria apicola** Speg. Leaf spot on *A. australe* in Chile and Argentina.**Uredo cundinamarcensis** Mayor. Brown rust pustules on leaves of *A. ternatum ranunculifolium* in Colombia.**APOSERIS.** See Hyoseris.**AQUILEGIA.** COLUMBINE. Hardy herbaceous perennials.**Actinonema pallens** Sacc. and Cav. On leaves of *A. vulgaris* in Italy.**Haplobasidium pavoninum** v. Hoeh. Large, solitary or marginal yellow spots, with purple-brown margins on leaves of *A. vulgaris* in Spain and Austria.**Marssonina aquilegiae** (Rbh.) Rostr. Subcircular, ashen colored spots, with dark margins, on leaves of *A. vulgaris* in Great Britain, France, and Denmark.**Phyllosticta aquilegicola** Brun. Dull-brown spots on leaves of *A. vulgaris* in Italy.**Puccinia agrostidis** Plowr. See Agrostis.**Puccinia melasmoides** Tranzsch. Brown rust pustules on indefinite spots on leaves and petioles of *A. akimensis* and *A. vulgaris* in Japan and Turkestan.**Septoria penzigii** Cav. and Mor. Whitish leaf spots, with broad brown margins, on *A. vulgaris* in Italy and Russia.**ARABIS.** ROCK CRESS. Perennial or annual herbs, mostly rock plants.**Peronospora arabis alpinae** Gäum. Downy mildew on leaves of *A. alba* and *A. alpina* in Switzerland.**Peronospora arabis hirsutae** Gäum. As above on *A. arenosa* and *A. hirsuta* in France, Switzerland, Norway, Denmark, Austria, and Germany.**Peronospora arabis turritae** Gäum. As above on *A. turrita* in Switzerland.**Phyllosticta alpina** Allesch. and var. *helvetica* Jaap. Large, circular or elongate definite spots on leaves of *A. alpina* in Bavaria and Switzerland.**Puccinia thlaspeos** Schubert. Brown rust pustules on leaves of *A. halleri*, *A. hirsuta*, *A. thaliana*, *Thlaspi alpestre*, *T. arvense*, *T. calaminaris*, *T. montanum*, *T. perfoliatum*, *T. praecox*, and *T. rotundifolium*, in Alaska, Russia, Sweden, Switzerland, Holland, Belgium, France, Austria, Hungary, and Germany.**Septoria arabis Sacc.** On leaves of *A. ciliata* in Italy.**Ustilago thlaspeos** (Beck.) Lagh. A smut forming black spore masses in the ovules in pods of *A. hirsuta*, *A. petraea* and *Thlaspi alpestre* in Switzerland and Austria.**ARACHIS.** PEANUT.**Cercospora** sp. A leaf spot, due to a *Cercospora* sp., said to be distinct from *C. personata*, has caused considerable losses to the peanut (*A. hypogaea*) in India.**Hypochnus cucumeris** Frank. See Cucumis.**Mosaic** or **Curl**. An obscure disease of *A. hypogaea*, probably of the mosaic group of diseases, occurs in the Union of South Africa, Java, and other parts of the Orient. It is characterized by a dwarfing of infected plants and the yellowing and crinkling of the foliage. The disease is called "crinkle leaf" or rosette in south Africa.**Rhizoctonia destruens** Tassi. See Solanum.**Sclerotinia arachidis** Hanzawa. Stem rot of *A. hypogaea* in Japan.**Sclerotinia miyabeana** Hanzawa. Stem rot of *A. hypogaea* in Japan.**Uredo arachidis** Lagh. Numerous dull-brown rust pustules on the under sides of leaves of *A. hypogaea* throughout the West Indies, Central America, and South America. This disease has proven itself the limiting factor of production in some of the West Indies. It has been found in one locality in Florida, but is thought to have been eradicated.**ARALIA.** Spikenard. Sarsaparilla. Udo. Herbs, shrubs, and trees, often spiny.**Cercospora araliae** P. Henn. Effuse, dull-brown spots on leaves of *A. sinensis* and *A. spinosa* in Japan.**Phyllosticta araliae** Sacc. and Berl. Large yellow to ashen spots, with dull yellow margins, on leaves of *Aralia* sp. in Algeria.**Physalospora araliae** Pat. On leaves of *Aralia* sp. in Ecuador.**Puccinia araliae-cordatae** P. Henn. Brown to black rust pustules on brown spots on leaves of *A. cordata* in Japan.**ARAUCARIA.** MONKEY PUZZLE. NORFOLK ISLAND PINE. Ornamental coniferous trees.**Coniothyrium pallido-fuscum** Sacc. See Chamaecyparis.**Gloeosporium auracariae** Karst. and Har. Anthracnose on leaves and cones of *A. rulei* in New Caledonia.**Haplodothis auracariae** (Rehm.) v. Hoeh. Wart-like stromata on needles of *A. brasiliana* in Brazil.**Phyllosticta auracariae** Wor. On needles of *A. imbricata* in Russia.**ARBUTUS.** STRAWBERRY TREE. Broad-leaved evergreen trees and shrubs.**Ascochyta unedonis** Sacc. (*Ascochyta unedonis* [Sacc.] Fl. Tass.) Ashen colored spots with redish margins on leaves of *A. unedo* in France and Spain.**Cercospora molleriana** Wint. Circular to irregular, often confluent, gray leaf spots on *A. andrachne*, *A. longifolia*, and *A. unedo* in Russia, Portugal, and Italy.**Coryneum microstictum** B. and Br. On *A. unedo* in Italy.**Exobasidium unedonis** Maire. The fungus deforms the branches and destroys the leaves of *A. unedo* in Algeria. It fruits on the under surface of the leaves and branches.**Mycosphaerella unedonis** Jaap. On leaves of *A. unedo* in Dalmatia.**Phyllosticta arbuti** (Desm.) Sacc. Small fuliginous-colored spots on leaves of *A. unedo* in France**Phyllosticta arbuti-unedonis** Pass. On leaves of *Arbutus* sp. in Portugal.**Septoria arbuti** Pass. On leaves of *A. unedo* in Italy.**Septoria hiascens** Sacc. Small dirty-white to dark-purple spots, with slightly raised margins, on leaves of *Arbutus* sp. in Mexico.**Septoria unedonis** Rob. and Desm. Small, irregular, numerous whitish spots with purple margins on leaves of *A. canariensis* and *A. unedo* in Algeria, Albania, Yugoslavia, Portugal, France, Italy, Great Britain, and Austria.

- ARDISIA. ICACOREA** Ag. Ornamental shrubs.  
**Helminthosporium flagellatum** Yates. On *A. disticha* in the Philippines.  
**Phyllachora ardisiae** P. Henn. Black tar spot on leaves of *A. candolleana* in the Philippines.  
**Phyllosticta ardisiae** Trinch. On leaves of *A. humilis* in Italy.  
**Physalospora circinana** Pat. Small fuscous circular spots on leaves of *Ardisia* sp. in Brazil.  
**Uromyces myrsines** Diet. Brown rust pustules on leaves of *A. compressa* and *Myrsine* sp. in Brazil and Costa Rica.
- ABECA.** Betal palm. See *Palmae*.
- AREGELIA.** Epiphytic hothouse plants of the *Bromeliaceae*.  
**Lembosia bromeliacearum** Rehm. Black fruiting bodies on dark-brown circular spots on leaves of *A. compacta* and *A. cruenta* in Brazil.
- ARENARIA.** SANDWORT. See, also, *Alsine*. Low herbs, suitable for borders and rockwork.  
**Helminthosporium nubigenum** Speg. On leaves of *A. tetraquetra* in Italy.  
**Laestadia arctica** Rostr. On leaves of *A. (Honckenia) peploides* in Greenland.  
**Peronospora campestris** Gaum. Downy mildew on lower surfaces of leaves of *A. ciliata*, *A. serylifolia*, and *A. trinervia* in France, Switzerland, Belgium, Denmark, Russia, Austria, and Germany.  
**Peronospora honckenya** Syd. Downy mildew on leaves of *A. (Honckenia) peploides* in Scotland.  
**Puccinia modica** Holw. Brown to black powdery rust pustules on leaves of *A. alsinoides*, *A. lanuginosa*, *A. peyritschii*, and *A. reptans* in Mexico and Guatemala.  
**Uredo arenariicola** P. Henn. Yellow-brown to brown rust pustules on leaves of *A. diffusa* in Argentina.  
**Uromyces arenariae** Tranzsch. Linear dark-brown rust pustules on leaves of *A. capillaris* and *A. longifolia* in Russia.  
**Uromyces arenariae-grandiflora** Mayor. Leaf rust on *A. grandiflora* in Switzerland.
- ARENGA.** Sugar palm. See *Palmae*.
- ARGEMONE.** PRICKLE POPPY. Mexican poppy. Prickly herbs.  
**Peronospora indica** Gäum. Downy mildew on leaves of *A. mexicana* in India. The name is occupied by *P. indica* Syd. on *Calceolaria*.  
**Rhizoctonia napi** West. See *Brassica*.  
**Septoria chelidonii** Desm. See *Chelidonium*.
- ARGYREIA.** SILVER CLIMBER. Ornamental climbers.  
**Aecidium argyreae** B. and Br. Yellow rust pustules on leaves of *A. argentea*, *A. elliptica*, *A. pomacea*, *A. populifolia*, and *A. speciosa* in India and Ceylon.  
**Cercospora cordobensis** Speg. Yellow circular spots on leaves of *A. megapotamica* in Ceylon and Argentina.  
**Uredo argyreae** Petch. Dark-brown rust pustules on leaves of *A. tiliaefolia* in Ceylon.
- ARISAEMA.** DRAGON-ROOT. JACK-IN-THE-PULPIT. Aroids with tuberous roots.  
**Septoria arisaemae** Petch. Leaf spots on *A. leschenaultii* in Ceylon.
- ARISARUM.** Ornamental aroids.  
**Phyllosticta arisari** Bres. Subcircular dull-brown spots on leaves of *A. vulgare* in Portugal.  
**Phyllosticta arisaricola** Sacc. and Syd. Circular brown spots, with light-brown margins, on leaves of *A. vulgare* in Tunis.  
**Septoria arisari** (Dur. and Mont.) Sacc. Dark irregular spots on leaves of *A. simorhinum* and *A. vulgare* in France and Algeria.  
**Septoria arisaricola** Pat. Circular scattered or confluent whitish spots on leaves of *A. vulgare* in Tunis.
- ARISTEA.** Greenhouse herbs.  
**Puccinia dehiscens** Syd. Brown rust pustules on leaves of *Aristea* sp. in the Union of South Africa.
- ARISTOLOCHIA.** DUTCHMAN'S-PIPE. PELICAN FLOWER. Ornamental perennial vines.  
**Ascochyta aristolochiae** Sacc. Subcircular dark-brown to black leaf spots on *A. clematitis* in Yugoslavia and Italy.  
**Ascochyta aristolochicola** Hóll. On capsules of *A. clematitis* in Hungary.  
**Ascochyta siphonis** Allesch. Subcircular to irregular black spots, becoming ashen, with black margins, on both sides of leaves of *A. siphon* in Germany.  
**Ascochyta versicolor** Bub. Circular to angular dark-brown spots, with raised black margins, on leaves of *A. clematitis* in Tyrol.  
**Cercospora olivascens** Sacc. Olive-gray spots on leaves of *A. clematitis* and *A. longa* in France and Italy.  
**Meliola atricapilla** Starb. Black mold on leaves of *Aristolochia* sp. in Brazil.  
**Mycosphaerella aristolochiae** Syd. On leaves of *A. tagala* in the Philippines.  
**Phyllosticta aristolochiae** F. Tassi. Round or angular to irregular dirty-white spots, with brown margins, on leaves of *A. clematitis*, *A. sempervirens*, and *A. siphon* in Italy and Germany. Reported once from the United States.  
**Phyllosticta siphonis** Kab. and Bub. Small angular, finally confluent, spots on upper surfaces of leaves of *A. siphon* in Bohemia.  
**Puccinia aristolochiae** ((D. C.) Wint. Yellow to dark-brown rust pustules on leaves of *A. botta*, *A. clematitis*, *A. longa*, *A. maurorum*, *A. pallida*, *A. poicilantha*, and *A. rotunda* in Russia, Kurdistan, Anatolia, Syria, Spain, Italy, France, Austria, and Germany.  
**Sclerotium zeylanicum** (B. and Br.) Petch. Root and collar rot of *A. leuconeura* in Ceylon. See, also, *Thea*.  
**Septoria aristolochiae** Sacc. On leaves of *A. clematitis* in Italy.
- ARISTOTELIA.** Evergreen trees or shrubs.  
**Cercospora aristoteliae** Cke. Subcircular brown spots, with red margins, on leaves of *A. racemosa* in New Zealand.  
**Phyllosticta aristoteliae** Speg. Indefinite angular dull-brown spots on leaves of *A. macqui* in Chile.  
**Septoria maqui** P. Henn. Dark-brown spots, with red margins, on leaves of *A. macqui* in Chile and Germany.
- ARMERIA.** STATICE Ag. THRIFT. Sometimes called "sea pink." Perennial herbs, used as border and rock garden plants.  
**Gloeosporium armeriae** Allesch. Anthracnose on scapes of *A. vulgaris* (*Statice armeria*) in Greenland.  
**Phyllosticta armeriae** Allesch. On leaves of *A. vulgaris* in Greenland.  
**Septoria armeriae** Allesch. and P. Henn. On leaves of *A. (Statice) plantaginea* in Greenland, Italy, and France.  
**Uromyces armeriae** (Schlecht.) Lev. Yellow to dark-brown rust pustules on leaves of *A. alpina*, *A. elongata*, *A. longibracteata*, *A. martinia*, *A. plantaginea*, *A. splendens*, *A. stenophylla*, and *A. vulgaris* in Europe. Reported from California.
- ARMORACIA.** See *Radicula*.

**ARNICA.** Perennial herbs.**Entyloma calendulae** (Oud.) De By. See *Calendula*.**Phyllosticta arnicae** Fckl. Indefinite yellow spots on upper surfaces of leaves of *A. montana* in Switzerland. Reported from Colorado.**Septoria arnicae** Fckl. Dark-brown spots on leaves of *A. montana* in Switzerland.**ARACACIA.** Umbelliferous herbs. Roots of one species eaten.**Puccinia arracacharum** Lindr. Yellow to brown powdery pustules on leaves of *A. bracteata* in Ecuador and Guatemala.**Puccinia imperspicua** Syd. Black rust pustules on leaves of *A. multifida* in Mexico.**ARRHENATHERUM.** OATGRASS. Cultivated as a meadow grass.**Dilophia graminis** (Fckl.) Sacc. See *Triticum*.**Exobasidium graminicolum** Bres. A doubtful species reported as causing indefinite yellow spots on leaves of *A. elatius* and *Bromus* sp. in Germany.**Phyllosticta stomaticola** Baeuml. Circular to irregular brown spots, becoming whitish, with red margins, on leaves of *A. elatius* in Hungary.**Puccinia arrhenatheri** (Kleb.) Erikss. Yellow aecial stage producing witches'-broom of *Berberis heteropoda*, *B. hispanica*, and *B. vulgaris*; brown uredo and telial stages on *Arrhenatherum elatius* in Turkestan, Algeria, Spain, France, Sweden, Austria, Hungary, and Germany.**Septoria culmifida** Lind. See *Phleum*.**Uromyces seseli-graminis** Ed. Fisch. Brown rust pustules on leaves of *Arrhenatherum elatius*, *Melica ciliata*, *Poa* sp., and *Seselis glaucus* in Europe.**Ustilago dura** App. and Gassn. Smut in inflorescences of *A. elatius* in Europe.**ARTABOTRYS.** TAIL GRAPE. Woody tropical climbers.**Uredo artabotrydis** Syd. Brown rust pustules on leaves of *A. odoratissima* in Formosa.**ARTEMISIA.** WORMWOOD. Aromatic and bitter herbs and shrubs.**Cercospora ferruginea** Fckl. On leaves of *A. vulgaris* in Japan, Poland, Bohemia, Italy, Denmark, Austria, and Germany.**Cercospora fuscescens** Niessl. On stems of *A. vulgaris* in Europe.**Cercospora olivacea** Oth. On leaves of *A. absinthium* in Switzerland.**Erysiphe taurica** Lev. See *Althaea*.**Nematostoma artemisiae** Syd. On living leaves of *A. vulgaris* in Japan.**Puccinia artemisiae-keiskeana** Miura. Rust on leaves of *A. keiskeana* in Japan.**Puccinia artemisiella** Syd. Black rust pustules on leaves of *A. vulgaris* in Japan, India, Russia, and Germany.**Puccinia artemisiicola** Syd. Brown rust pustules on leaves and stems of *A. austriaca* and *A. campestris* in Denmark, Russia, Austria, Hungary, and Germany.**Puccinia ferruginosa** Syd. Powdery, dull-brown rust pustules on small, sunken, brown leaf spots on *A. vulgaris* var. *japonica* and *Artemisia* sp. in India and Japan.**Septoria artemisiae** Pass. Dull-brown spots on leaves of *A. vulgaris* in Italy.**Septoria kriegieriana** Bres. On leaves of *A. vulgaris* in Saxony.**Uredo artemisiae-japonica** Diet. Brown rust pustules on leaves of *A. japonica* in Japan.**ARTOCARPUS.** BREADFRUIT. JACKFRUIT. Tropical fruit trees.**Catacauma microcentum** Theiss. and Syd. See *Ficus*.**Cercospora artocarpi** Syd. On leaves of *A. communis* (*A. incisa*) in the Philippines.**Corticium salmonicolor** B. and Br. See *Citrus*.**Diplodia artocarpi** Sacc. Causes twig die-back and fruit rot of *Artocarpus* sp. in Java and the Philippines. Probably not distinct from *D. cacaoicola*.**Fomes lamaoensis** Murr. See *Hevea*.**Fomes lignosus** Klotzsch. See *Hevea*.**Marchalia constellata** (B. and Br.) Sacc. Circular, flat, black stromata on leaves of *Artocarpus* sp. in the Philippines and *A. integra* (*A. integrifolia*) in Ceylon.**Phyllosticta** sp. On leaves of *A. communis* in India.**Phyllosticta** (?) **artocarpi** Speg. Circular, determinate white spots, with broad dark margins, on leaves of *A. integra* (*A. integrifolia*) in Argentina.**Phytophthora** sp. Causes fruit rot of *A. incisa* in Ceylon.**Phytophthora faberi** Maubl. See *Theobroma*.**Rhizopus artocarpi** Rac. Causes a soft rot of fruit and male inflorescences of *A. integra* in the Philippines.**Rosellinia pepo.** Pat. See *Citrus*.**Septoria artocarpi** Cke. Circular, brown sunken spots on leaves of *A. integrifolia* in India.**Septoria eburnea** v. Hoeh. Circular to angular or irregular leaf spots on *A. communis* in Samoa.**Sphaerostilbe repens** B. and Br. See *Hevea*.**Uredo artocarpi** B. and Br. (*Physopella artocarpi* [B. and Br.] Arth.) Numerous dull-brown rust pustules covering lower surfaces of leaves of *A. camansi*, *A. communis*, *A. lacoocha*, and *Castilla elastica* in Ceylon, Cuba, and Porto Rico.**ARUM.** Variegated leafed, herbaceous aroids.**Ascochyta arophila** Bub. Circular to elliptical brown spots, becoming white at the centers, on leaves of *A. italicum* in Montenegro.**Colletotrichum montemartini** Togn. Anthracnose on *A. italicum* in Italy.**Phyllosticta aricola** Bub. Brown, circular, or elongate leaf spots, becoming white at the centers, on *A. italicum* in central Europe.**Phyllosticta scrophulariae-bosniacae** Bub. On leaves of *A. italicum* and *A. bosnia* in Central Europe.**Phyllosticta tuzsonii** Bub. Elongate to irregular yellow spots with white centers, on leaves of *A. italicum* in Hungary.**Ramularia ari** Fautr. Gray leaf spots on *A. italicum* and *A. maculatum* in Spain, France, and Dalmatia.**Septoria ari** Desm. Subcircular brown spots on leaves of *A. italicum* and *A. maculatum* in Algeria, Yugoslavia, Bulgaria, France, and Italy.**Ustilago plumbea** Rostr. A smut producing black spore masses on leaves and petioles of *A. maculatum* in Denmark.**ARUNCUS.** GOATSBEARD. Ornamental, tall perennial herbs.**Ascochyta arunci** Sacc. On stems of *A. sylvester* in Italy.**Colletotrichum exiguum** Penz. and Sacc. Numerous, small, brown leaf spots on *A. sylvester* in Italy and Austria.**Ochropsora ariae** (Fckl.) Syd. See *Sorbus*.**Phyllosticta aruncina** Sacc. Small, grayish-white leaf spots on *A. sylvester* in Italy.**Phyllosticta griseo-fusca** Bub. Small, angular to confluent leaf spots dull green, then brown above, gray beneath on *A. sylvester* in Bohemia.**Ramularia spiraeae-arunci** (Sacc.) Allesch. On leaves of *A. sylvester* in Switzerland and Austria.**Septoria arunci** Pass. Gray-white leaf spots with dark-red margins on *A. sylvester* in Italy.

**ARUNDINA.** See Orchidaceae.

**ARUNDINARIA.** See Bambuseae.

**ARUNDO.** GIANT REED. Tall perennial grasses.

**Apiospora montagni** Sacc. Linear black stromata on leaves of *A. donax* and *A. mauritanica* in Algiers, Portugal, and France.

**Asochyta donacina** Sacc. On culms of *A. donax* in Italy and France.

**Phyllachora atrofigurans** Rehm. Black stromata on leaves of *A. cannaeformis* in the Philippines.

**Phyllachora cyperi** Rehm. var. **donacis** Berl. and F. Sacc. "Tar spot" disease on culms of *A. donax* in India and Portugal.

**Phyllachora donacina** Rehm. Black, convex, orbicular or ellipsoid stromata in parallel series on leaves of *A. cannaeformis* in the Philippines.

**Physalospora festucae** (Lib.) Sacc. See Andropogon.

**Puccinia isiacae** (Thuem.) Wint. Rust-brown to dark-brown rust pustules on leaves of *A. isiacae* and *Phragmites communis* in Egypt, Asia Minor, and Russia. Aecia may possibly occur on *Raphanus sativus*.

**Puccinia torosa** Thuem. Brown rust pustules on leaves of *A. donax* in the Union of South Africa.

**Septoria donacina** Sacc. Indefinite spots on culms of *A. donax* in Italy.

**Septoria donacis** Pass. Small dry spots on leaves of *A. donax* in Dalmatia, Italy, France, and Portugal.

**Septoria oxyspora** Penz. and Sacc. See Anthoxanthum.

**ASARUM.** WILDGINGER. Low, nearly stemless herbs.

**Septoria asari** Sacc. Circular to angular whitish spots, with black borders, on leaves of *A. europaeum* in Italy.

**Septoria asaricola** Allesch. Subcircular to angular, brown then gray spots, often confluent, on leaves of *A. europaeum* in Russia and Bavaria.

**ASCLEPIAS.** MILKWEEED. BUTTERFLY WEED. Perennial, milky-juiced herbs.

**Cercospora asclepiadis** P. Henn. Numerous circular leaf spots on *Asclepias* sp. in Brazil.

**Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.

**Mycosphaerella asclepiadis** Siem. On leaves of *A. incarnata* in Caucasia.

**Puccinia conrescens** E. and E. Leaf rust on *A. curassavica* and *A. nivea* in Porto Rico, Cuba, Bahamas, and South America.

**Septoria schlechteriana** P. Henn. Pale leaf spots, with dark brown margins, on *A. fruticosa* in the Union of South Africa.

**Septoria syriacae** Kab. and Bub. Numerous circular to angular often confluent, spots on leaves of *A. syriaca* in Bohemia.

**ASPARAGUS.** Herbaceous perennials and tender woody shrubs and vines. One species used for food.

**Aecidium schlechterianum** P. Henn. Round yellow spots in which yellow rust pustules appear, are formed on the leaves of *A. medeoloides* in south Africa.

**Botryosphaeria dispersa** De Not. On *A. officinalis* in Europe.

**Cercospora asparagi** Sacc. Dirty-white to gray spots on false leaves and branches of *A. officinalis* in Brazil, Cuba, Argentina, and Italy.

**Cercosporina asparagicola** Speg. On *A. officinalis* in Argentina.

**Colletotrichum** sp. Anthracnose on stems of *Asparagus* sp. in Brazil.

**Coniothyrium** sp. Produces stem cankers on stems of *Asparagus* sp. in Brazil.

**Hysterostomella tenella** Syd. On leaves and stems of *A. striatus* and *Asparagus* sp. in the Union of South Africa.

**Leptosphaeria circinans** (Fekl.) Sacc. On roots and stems of *A. officinalis*, *Beta vulgaris*, *Medicago sativa*, *Phaseolus* sp., *Solanum tuberosum*, *Trifolium pratensis*, and *T. repens* in Italy, Russia, and Germany.

**Leptosphaeria indica** Syd. and Butl. On *Asparagus* sp. in India.

**Melanospora asparagi** Arnaud. Attacks the base of the stalks of *A. officinalis* in France, preventing further development.

**Ophiobolus compar** Karst. Purple sunken spots, in which small, black, fruiting bodies appear, are formed on stems of *A. officinalis* in Finland.

**Puccinia asparagi-lucidi** Diet. Chestnut-colored rust pustules, becoming dark brown to black on both leaf surfaces and on stems of *A. lucidus* and *A. schobertoides* in Japan.

**Puccinia phyllocladiae** Cke. This rust produces powdery brown pustules on the lower leaf surfaces and on stems of *A. falcatus* in Natal and Ceylon.

**Zopfia variospora** (Boud.) Arnaud (*Z. rhizophila* Rabenh.) On *A. acutifolius*, *A. horridus*, and *A. officinalis* in France, Italy, and Algeria.

**ASPERULA.** WOODRUFF.

**Mycosphaerella asperula** Roum. and Fautr. On leaves of *A. tinctoria* in Esthonia.

**Peronospora calotheca** De By. Downy mildew on leaves of *A. odorata* and *A. tinctoria* in central and north Europe.

**Phyllosticta asperulae** Sacc. Indefinite, yellow spots on leaves of *A. odorata* in France and Hungary.

**Pseudopeziza autumnalis** (Fr.) Karst. See Galium.

**Puccinia asperulae-cynanchicae** Wurth. Rust on leaves of *A. cynanchica* and *A. odorata* in Spain, Yugoslavia, Switzerland, and Germany.

**Puccinia asperulae-odoratae** Wurth. On leaves of *A. odorata* in Denmark, Sweden, Switzerland, Yugoslavia, and Japan.

**Puccinia asperulina** (Juel) Lagh. Golden-yellow rust pustules, becoming brown, causing distortion of leaves of *A. ciliata* and *A. tinctoria* in Sweden, Russia, and Hungary.

**Puccinia helvetica** Schroet. Brown rust pustules on leaves of *A. taurina* in Switzerland, Italy, and Hungary.

**Puccinia oligantha** McAlp. Brown to dark-brown rust pustules on leaves and stems of *A. oligantha* in Victoria (Australia).

**Puccinia spillogena** Lindr. Yellow-brown to dark-brown rust pustules on leaves of *A. mollugo* in Persia.

**Septoria asperulae** Baeuml. and var. **italica** Mass. Irregular gray spots with fuscous margins, soon involving the entire leaf, on *A. odorata* and *A. taurina* in Denmark, Italy, and Hungary.

**Septoria asperulae-arvensis** Maire. On leaves of *A. arvensis* in French north Africa.

**Septoria asperulae-taurinae** Bub. Dirty-white spots on upper surfaces of leaves of *A. taurina* in Hungary.

**Thekopsora guttata** (Schroet.) Syd. Yellow to dark brown rust pustules on leaves of *A. galioides*, *A. odorata*, *Galium aristatum*, *G. hercynicum*, *G. mollugo*, *G. rotundifolium*, *G. saxatile*, *G. schultesium*, *G. silvaticum*, *G. sylvestris*, *G. spurium*, *G. uliginosum*, *G. verum*, and *Sherardia arvensis* in Europe. Reported from Oregon.

**Uromyces asperulae** McAlp. Cinnamon-brown to dark-brown rust pustules on leaves and stems of *A. oligantha* in Australia.

**ASPHODELINE.** JACOB'S ROD. Hardy, herbaceous plants.**Septoria asphodelines** Syd. On leaves of *Asphodeline taurica* in Bulgaria.**ASPHODELUS.** ASPHODEL. Hardy, herbaceous, stemless plants.**Cladochytrium asphodeli** De By. (*Physoderma asphodeli* [De By.] Vesterg.) On leaves of *Asphodelus* sp. causing dark brown or black irregular spots, in Tunis and Algeria.**Diplodia uncinjensis** Bub. On stems of *A. microcarpus* in Montenegro.**Phoma asphodeli** Sacc. (*P. asphodelina* Thuem.) On stems and leaves of *Asphodelus* sp. in Spain.**Phomopsis asphodelina** (Thuem.) Trav. and Sp. On scapes and capsules of *A. microcarpus* in Spain.**Phyllosticta caulicola** Pat. Broad, brown spots with black margins are produced on the stems of *A. microcarpus* in Algeria.**Puccinia asphodeli** Moug. Yellow to yellow-brown rust pustules, becoming darker on both surfaces of the leaves of *Anthericum ramosum*, *Asphodelus albus*, *A. cerasiferus*, *A. microcarpus*, *A. occidentalis*, *A. ramosus*, *A. subalpinus*, and *A. tenuifolius* in France, Italy, Portugal, Spain, Greece, Yugoslavia, Syria, Algeria, Canary Islands, Corsica, and Balearic Islands.**Puccinia barbeyi** (Roum.) P. Magn. Sunken elongate spots occur on both leaf surfaces, and on the stems, in which yellow and brown rust pustules appear. On leaves of *A. fistulosus*, *A. microcarpus*, and *A. tenuifolius* in France, Greece, Syria, Algeria, Spain, and India.**Puccinia unamunoi** Gz. Frag. Yellow to brown rust sori on leaves and scapes of *A. albus* in Spain.**Septoria asphodelina** Sacc. On leaves of *A. albus*, *A. fistulosus* and *A. microcarpus* in Spain and Belgium.**Septoria asphodeli-ramosi** Pat. Ashen elongate spots on stems of *A. ramosus* in Tunis.**ASPIDISTRA.** Florist's plant grown for its stiff, shining foliage which is often variegated.**Ascochyta aspidistrae** Mass. Large, irregular, bleached areas on leaves of *A. lurida* in Great Britain. Reported from Minnesota.**Phyllosticta aspidistrae** Oud. Large, rust-colored spots on leaves of *A. elatior* in Holland.**Septoria transversalis** Sacc. On leaves of *A. elatior* in Italy.**ASPIDIUM.** DRYOPTERIS Ag. WOOD FERN.**Exoascus cornu-cervi** Sadeb. On leaves of *A. aristatum* in New Caledonia.**Exoascus fuscus** (Giesen.) Sacc. and Syd. Produces fleshy, dull-brown galls on leaves of *A. pallidum* in Sicily and Albania.**Exoascus verstergrenii** (Giesen.) Sacc. and Syd. Brown galls on both surfaces of leaves of *A. filix-mas* in the Swedish Baltic Islands, and Switzerland.**Exobasidium brevieri** Boud. On leaves of *A. filix-mas* in Esthonia.**Glœosporium filicinum** Rostr. Anthracnose of leaves of *A. filix-mas* in Sweden.**Hyalopora filicum** Diet. See *Asplenium*.**Milesina carpatica** A. Wrób. Yellow rust pustules on leaves of *A. filix-mas* in Austria.**Milesina kriegeriana** P. Magn. Yellow rust pustules on leaves of *A. aculeatum*, *A. dilatatum*, *A. filix-mas* and *A. spinulosum* in Dalmatia, Finland, Denmark, Austria, Hungary and, Germany.**Milesina vogesiaca** Syd. Yellow rust pustules on leaves of *A. lobatum* in Austria and Germany.**Mycosphaerella aquilina** (Fr.) Schroet. and f. **aspidiorum** Sacc. On leaves of *A. filix-mas* and *Asplenium adiantum* in Switzerland and Esthonia.**Mycosphaerella thelypteridis** Syd. On *A. thelypteris* in Germany.**Trabutiella filicina** (Sacc. and Sc.) Theiss. and Syd. Black stromata on leaves of *A. lonchitis* in Alaska.**ASPLENIUM.** SPLEENWORT. Ferns.**Aphelenchus olesistus** Ritz. Bos. See *Begonia*.**Cercospora aspleni** Jaap. On leaves of *A. trichomanes* in Dalmatia.**Exobasidium brevieri** Boud. Small, irregular white spots, often confluent, on leaves of *A. filix-femina* in France.**Hyalopora asplenii-wichuriae** Diet. Brown rust pustules on indefinite dirty-brown spots on leaves of *A. wichuria* in Japan.**Hyalopora filicum** Diet. Yellow rust pustules on leaves of *A. decursivae-pinnatum*, *A. japonicum*, *A. patens*, *Athyrium nipponicum*, *Dryopteris laza*, and *Pteris semipinnata* in Japan.**Milesina feurichii** P. Magn. Yellow rust pustules on leaf blades and petioles of *A. septentrionalis* in Finland, Switzerland, and Germany.**Milesina magnusiana** Jaap. Rust pustules on yellow-brown spots on leaves of *A. nigrum* in Dalmatia, Italy, and Corsica.**Milesina murariae** (P. Magn.) Grove. Yellow rust pustules on leaves of *A. ruta-muraria* in Scotland, Switzerland, Austria, and Germany.**Mycosphaerella asplenii** (Auersw.) Lind. On leaves of *A. septentrionale* in Europe.**Mycosphaerella filicum** (Desm.) Starb. See *Aspidium*.**Phyllosticta asplenii** Jaap. On leaves of *A. ruta* in Switzerland.**Ramularia aspleni** Jaap. On leaves of *A. ruta* in Switzerland and Germany.**Scirrhodopsis confluens** (Starb.) Theiss. and Syd. Black stromata on gray leaf spots on *Asplenium* sp. in Sweden.**ASSONIA.** See *Dombeya*.**ASTER.** MICHAELMAS DAISY. Perennial herbs.**Aecidium microsporum** Diet. Rust on leaves of *A. divaricatus* in Brazil.**Bacillus asteracearum** Pava. Small, scattered, ocher-colored spots, first appear on the lower leaves, increasing in size and spreading up the stalk to the flowers which, together with the leaves, turn brown and wither on *Aster* spp. in Italy.**Cephalosporium asteris** Dow. A serious wilt disease on cultivated *Aster* in Great Britain is attributed to this fungus. Infected plants show bright-lemon or orange-colored mottling of the lower leaves which finally become brown and shrivelled. An excessive suckering is also in evidence.**Cercospora asterina** Speg. Ashen-brown spots, with darker margins, on leaves of *A. linifolius* in Argentina.**Coleosporium asterum** (Diet.) Syd. Yellow rust pustules on leaves of *A. glenki*, *A. hispidus*, *A. pinnatifidus*, *A. scaber*, *A. tataricus*, *A. trinervius*, *Boltonia cantoniensis*, *B. incisa*, *B. indica*, *Callistephus chinensis*, *A. sinensis*, and *Pinus densiflora* in China, Formosa, and Japan. Reported from California.**Didymaria asteris** Oud. Broad, irregular white spots on leaves of *Aster* sp. in Holland.**Entyloma calendulae** (Oud.) De By. See *Calendula*.**Puccinia asteris-alpini** Syd. Black rust pustules on small brown sunken spots on leaves of *A. alpinus* in Switzerland.**Ramularia asteris-tripolii** Jaap. Subcircular to oblong, often indefinite, yellow, finally gray or brown spots on leaves of *A. tripolium* in Denmark.**Septoria tatarica** Syd. Greenish-gray, circular then confluent, leaf spots on *A. tataricum* in Japan.

**ASTILBE.** Ornamental perennial herbs.

**Pucciniostele clarkiana** (Barcl.) Diet. Yellow rust pustules on circular or elongate spots, on leaves and petioles of *A. congesta*, *A. philippinensis*, *A. rivularis*, *A. sinensis*, and *A. thunbergii* in India, Japan, and the Philippines.

**Pucciniostele mandshurica** Diet. Powdery, yellow rust pustules on lower surfaces of leaves and on petioles of *A. chinensis* in Japan, Korea, and Manchuria.

**Septoria hotejiae** F. Tassi. On leaves of *A. japonica* in Italy.

**Triphragmium nishidanum** Diet. Chestnut-colored rust pustules on leaves of *A. chinensis* var. *albiflora* and *A. sinensis* in Japan.

**ASTRAGALUS.** MILKVEGET. Hardy herbs or subshrubs.

**Cylindrosporium astragali** E. Rostr. On leaves of *A. oroboides* in Norway.

**Erysiphe taurica** Lev. See *Althaea*.

**Euryachora tragacanthae** (Lev.) Theiss. and Syd. Black, irregular stromata on thorax and leaves of *A. tragacantha* in Persia.

**Microsphaera astragali** (DC.) Trev. Powdery mildew on leaves of *A. cicer*, *A. glycyphyllos*, and *A. onobrychis* in Europe.

**Mycosphaerella magnusiana** Jaap. On leaves of *A. alpinus* in Austria.

**Ovularia sphaeroidea** Sacc. On leaves of *A. sinicus* in Japan.

**Ovularia tuberculiniformis** v. Hoeh. Small, often confluent, dirty-yellow leaf spots on *A. cicer* in Yugoslavia and Austria.

**Peronospora astragalina** Syd. Downy mildew on leaves of *A. alpinus* and *A. oroboides* in Norway, Lapland and Austria.

**Phyllachora melaena** (Rabh.) Sacc. Black stromata on brown definite leaf spots on *A. glumaceus* in Kurdistan.

**Phyllosticta astragalicola** Mass. Olivaceous spots, with brown margins, on leaves of *A. glycyphylloides* in Italy.

**Phyllosticta desertorum** Sacc. Subcircular small brown spots on both surfaces of leaves of *A. alopecurus* in Siberia.

**Phyllosticta exscapi** Höll. On leaves of *A. exscapus* in Hungary.

**Physalosporina astragalina** (Rehm. Wor. On leaves of *A. cicer* in Russia and Germany.

**Physalosporina obscura** (Juel.) Wor. On leaves of *A. alpinus* and *A. arenarius* in Denmark and Russia.

**Placosphaerella tragacanthae** Pat. Black, rough, fruiting areas on leaves of *A. fontanesius* and *A. tragacantha* in Persia and Arabia.

**Polystigma obscurum** Juel. Stromata occupying most of leaf surface, white beneath, yellowish-red, then gray-brown above. On *A. alpinus* and *A. oroboides* in Sweden and Norway.

**Septoria astragali** Desm. On leaves of *A. alpinus* and *A. glycyphyllos* in Spain, Denmark, Russia, Bohemia, and Austria.

**Septoria henningsiana** Wint. On leaves of *A. onobrychis* in Asia Minor and Yugoslavia.

**Septoria serebriankowii** Sacc. Causes yellowing of leaves of *A. onobrychis* in Russia.

**Uromyces carneus** (Nees.) Har. Brown, powdery, rust pustules covering entire leaf surfaces of *A. alpinus* and *A. australis* in Norway, Sweden, Switzerland, and Austria.

**Uromyces gordianus** Bub. Brown rust pustules on leaves of *A. exscapus* in Switzerland, Austria, Hungary and Germany.

**Uromyces patagonicus** Speg. Cinnamon-brown rust pustules on leaves of *Astragalus* sp. in Patagonia.

**Uromyces persicus** Syd. Dark-brown rust pustules on leaves of *A. remotijugus* in Persia.

**Uromyces splendens** Blytt. Yellow to brown rust pustules on leaves of *A. oroboides* in Norway.

**ASTRANTIA.** MASTER WORT. Perennial garden plants.

**Ascochyta astrantiae** Roum. On leaves of *A. major* in France.

**Fabraea astrantiae** (Ces.) Rehm. On *Astrantia* sp. in Switzerland.

**Phyllosticta astrantaecola** Gz. Frag. On leaves of *A. major* in Spain.

**Pseudopeziza saniculae-astrantiae** Niessl. On leaves of *A. major* in Germany.

**Puccinia astrantiae** Kalchbr. Dark-brown rust pustules on leaves and petioles of *A. major* and *A. minor* in Portugal, France, Belgium, Austria, Hungary and Germany.

**Puccinia astrantiae-vivipari** Semad. Rust pustules on leaves of *A. minor* and *Polygonum viviparum* in Switzerland.

**Ramularia oreophila** Sacc. Angular, dull yellow-brown spots on leaves of *A. major* and *A. minor* in Italy, Spain, Switzerland, Russia and Austria.

**ASTROCARYUM.** See *Palmae*.**ASYSTASIA.** Hothouse evergreen herbs and shrubs.

**Puccinia phaeosticta** Pat. and Har. Brown rust pustules on circular brown spots on leaves of *Asystasia* sp. in Indo-China.

**Uredo asystasiae** P. Henn. Brown rust pustules on yellow spots on leaves of *A. gangetica* in east Africa.

**ATALANTIA.** Small, usually spiny trees and shrubs related to *Citrus*.

**Bacterium citri** Hasse. See *Citrus*.

**Phyllosticta disciformis** Penz. See *Citrus*.

**ATAMOSCO.** See *Zephyranthes*.**ATRIPLEX.** SALT BUSH. Orach. Ornamental shrubs, garden vegetables and many weedy species.

**Ascochyta atriplicis** Diet. Subcircular, often confluent, dull-yellow spots, with pale centers and dark margins, on leaves of *A. hastata* in Germany.

**Ascochyta chenopodii** Rostr. Circular, pale-brown spots, with yellow margins, on leaves of *A. bonushenricus*, *A. littoralis*, *Chenopodium album*, *C. glaucum*, and *Spinacia oleracea* in Denmark, China, Russia, and Germany.

**Cercospora dubia** (Riess.) Wt. On leaves of *A. hastata*, *A. hortensis*, *A. laciniata*, *A. littoralis*, *A. nitens*, and *A. patula* in France, Bohemia, Russia, Denmark, and Germany. Reported from Montana.

**Peronospora minor** (Casp.) Gäum. Downy mildew on leaves of *A. nitens*, *A. patula* and *A. roseum* in England and north Europe.

**Phyllosticta atriplicicola** Speg. Circular, ashen-colored spots on leaves of *A. hastata* in Argentina.

**Phyllosticta halophila** Speg. Circular, often confluent, white spots, with purplish margins, on leaves of *A. hastata* in Argentina.

**Ramularia dubia** Wint. Spots on leaves of *A. hastata*, *A. nitens* and *A. patula* in Russia and Germany.

**Septoria atriplicis** (West.) Fekl. Greenish or dirty yellow to whitish leaf spots on *A. patula*, *Chenopodium murale* and *C. vulvaria* in Siberia, Tunis, Bulgaria, France, Spain, Italy, Belgium, Germany, and Central America. Reported from a few localities in the United States.

**Uromyces atriplicis** McAlp. Pale-brown to dark-brown rust pustules on leaves of *A. semibaccata* in Victoria.

**ATRIPLEX**—Continued.

**Urophycitis pulposa** (Wallr.) Schroet. Swellings on stems and leaves of *A. hastata*, *A. patula*, *Beta vulgaris*, *Chenopodium album*, *C. cubicum*, *C. glaucum*, *C. rubrum* and *C. urticum* in Argentina, Denmark, Siberia, and Germany.

**ATROPA. BELLADONNA.** Medicinal herbs.

**Ascochyta atropae** Bres. Pale-brown spots with darker margins on leaves of *A. belladonna* in Germany.

**Phyllosticta atropa** e F. Tassi. Circular, white spots, with yellow borders, on leaves of *A. belladonna* in Italy.

**Phytophthora erythroseptica** Pethyb. See Solanum.

**Puccinia atropae** Mont. Yellow to black rust pustules on leaves of *A. aristata* in the Canary Islands.

**Ramularia atropae** Allesch. Irregular, dull-brown spots on both leaf surfaces of *A. belladonna* in Switzerland and Germany.

**AUCUBA.** Ornamental evergreen shrubs, often with variegated leaves, and with bright scarlet fruit.

**Ascochyta aucubae** Sacc. and Speg. On leaves of *A. japonica* in Italy and France. Var. *brunaudiana* Sacc. differs in the more definite spots with black margins.

**Ascochyta aucubicola** Wint. Irregular, whitish spots with fuscous margins on leaves of *A. japonica* in Portugal.

**Colletotrichum pollaccii** Magn. Anthracnose on *A. japonica* in Japan.

**Phyllosticta ampla** P. Brun. Large, irregular brownish-green or gray-green spots on leaves of *A. japonica* in France.

**Phyllosticta aucubae** Sacc. and Speg. On leaves of *A. japonica* in Italy.

**Phyllosticta aucubicola** Sacc. Pale, indefinite spots on leaves of *A. japonica* in Denmark and Italy.

**Phyllosticta indica** Roum. and Karst. Large, whitish spots, with dull-brown margins, on leaves of *A. japonica* and *A. lanceolata* in Indo-China.

**Pleospora infectoria** Fekl. var. **aucubicola** Mont. Black patches appear on tips and along edges of the leaves. These areas increase in size, soon involving the entire blade which withers and dies. New leaves are attacked as they unfold. On *A. japonica* in Italy.

**Septoria aucubae** West. Large, irregular, pale brown spots with darker margins on leaves of *A. japonica* in Italy and Belgium.

**Sphaerulina aucubae** Shir. and Har. On leaves of *A. japonica* in Japan.

**AURICULA.** See Primula.**AVENA.** OATS. Annual grasses cultivated for grain and forage.

**Acremoniella verrucosa** Togn. Producing black spots on culms and sheaths of *A. sativa* and *Triticum* sp. in Italy.

**Ascochyta graminicola** Sacc. var. **ciliolata** Sacc. On *A. filifolia* and *A. pratensis* in Spain.

**Cladochytrium graminis** Buesgen. See Festuca.

**Heterosporium cerealinum** Oud. On leaves of *Avena* sp. in Holland and Belgium.

**Leptosphaeria nigrans** (Desm.) Ces. and De N. See Aira.

**Metasphaeria avenae** (Auersw.) Sacc. On leaves of *Avena* sp. and *Calamagrostis* sp. in France and Germany.

**Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.

**Puccinia pratensis** Blytt. Powdery, brown pustules on leaves of *A. pratensis* in Sweden and Norway.

**Rhizoctonia napi** West. See Brassica.

**Sclerospora macrospora** Sacc. See Triticum.

**Septoria planiuscula** Diet. Leaf spots on *A. planiuscula* in Germany.

**Tylenchus hordei** Schoeyen. This nematode is said to cause small galls on the roots of *A. sativa*, *Elymus arenarius*, *Hordeum vulgare* and *Poa pratensis* in northern Europe. Possible should be referred to *Heterodera radicola*.

**Tylenchus tritici** Bast. See Triticum.

**AVERRHOA.** Carambola. Tropical fruit trees.

**Cercospora averrhiae** Petch. Circular, white leaf spots with purple-red margins, on *A. carambola* in Ceylon.

**Cercospora averrhoi** Welles. Irregular yellow lesions on leaves 3 to 5 millimeters in diameter, which may coalesce to form large irregular spots. Center of diseased areas die, becoming grayish-brown, surrounded by band of yellow tissue. Black lesions form on the fruit. On *A. carambola* in the Philippines. Probably not distinct from the preceding species.

**Phyllosticta** sp. On leaves of *A. bilimbi* in Cuba.

**AZALEA.** See Rhododendron. The diseases of *Azalea* and *Rhododendron* are interchangeable to such an extent that it does not appear feasible to separate them, just as the host genera themselves are botanically inseparable.**AZARA.** Small, ornamental evergreen trees or shrubs.

**Uredo azarae** Neg. Golden-yellow rust pustules on leaves of *A. integrifolia* in Chile.

**BACKHOUSIA.** Evergreen shrubs or trees.

**Ascochyta apiospora** Cke. and Mass. Brown spots with purple borders on leaves of *Backhousia* sp. and *Myrtus* sp. in Australia.

**BACTRIS.** See Palmae.**BAHIA.** Herbaceous perennials.

**Phragmodothis minutissima** (Starb.) Theiss. and Syd. On stems of *Bahia* sp. in Brazil.

**BAMBUSA.** See Bambuseae.**BAMBUSEAE.** BAMBOO. Large, ornamental, and economic perennial grasses. It has seemed desirable to consider under one heading the diseases of the various genera comprising the tribe *Bambuseae* of the *Gramineae*, not only since the diseases are, to a considerable extent, interchangeable, but because the host plants themselves are so difficult of determination that in many cases disease reports give only the common name bamboo, making exact host determination impossible, and a separation of the diseases, therefore, out of the question.

**Aciculosporium take** Miy. Produces witches'-brooms of *Phyllostachys* sp. in China and Japan.

**Ascochyta arundinariae** Gz. Frag. On leaves of *Arundinaria nitida* in Spain.

**Ascopolyporus polychrous** A. Moell. On living twigs of *Bambusa* sp. in Brazil.

**Balansia chusqueicola** P. Henn. Black, carbonaceous crusts, about one centimeter long, surrounding stems of *Chusquea* sp. in Costa Rica.

**Balansia claviceps** Speg. On *Bambusa* sp. in Costa Rica and Brazil.

**Balansia regularis** Moell. Producing witches'-broom effects on *Bambusa* sp. in Brazil.

**Cenangella bambusicola** Rick. White, then greenish-yellow, sessile fruiting bodies on branches of *Bambusa* sp. in Brazil.

**Chaetophoma pellicula** Sacc. and Syd. Black effused areas on leaves of *Chusquea* sp. in Chile.

**Coccodiella arundinariae** Hara. Fleshy, disk-shaped, scattered fruiting bodies on leaves of *Arundinaria simoni* and *Sasa borealis* in Japan.

## BAMBUSEAE—Continued.

- Colletotrichum septorhoides** Sacc. On culms of *Bambusa* sp. in China.
- Cylindrosporium bambusae** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys bambusoides* in Japan.
- Dothidella gigantochloae** (Rehm.) Theiss. and Syd. On stems and leaves of *Gigantochloa* sp., *G. levis*, and *G. scribneriana* in the Philippines. The exact position of this fungus is uncertain.
- Endodothella bambusae** (Rabh.) Theiss. and Syd. Linear black stromata on rust-brown leaf spots on *Bambusa spinosa* and *Bambusa* sp. in India.
- Epichloe bambusae** Pat. Fleshy, black, effused (8 to 10 centimeters) stromata on leaves of *Bambusa* sp. and *Sasa spiculosa* in Japan, Java, and India.
- Epichloe sasae** Hara. Causes witches' brooms on *Sasa spiculosa* in Japan.
- Fomes lignosus** Klotzsch. See Hevea.
- Gilletiella chusqueae** (Pat.) Sacc. and Syd. Small, black, circular or elliptical stromata on brown leaf spots on *Chusquea* sp. in Ecuador.
- Gloeosporium sphaeosporum** Hara. Anthracnose on *Phyllostachys reticulata* in Japan.
- Gloniella chusqueae** P. Henn. Leaf spot on *Chusquea* sp. in Chile.
- Gloniella chusqueicola** P. Henn. Black, erumpent fungus pustules on young twigs of *Chusquea* sp. in Chile.
- Guignardia bambusae** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys* sp. in Japan.
- Guignardia blumeanae** (Sacc.) Yates. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Helminthosporium bdellomorphum** Speg. On culms of *Chusquea valdiviensis* in Chile.
- Helminthosporium blumeanum** Sacc. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Homostegia fusispora** Syd. Small, yellow, indefinite spots on leaves of *Bambusa* sp. in the Philippines.
- Hypocreopsis phyllostachydis** (Syd.) M. and H. In the ovaries of *Phyllostachys puberula* in Japan.
- Kusanobotrys bambusae** P. Henn. Round, black, often confluent, spots on upper surfaces of leaves of *Bambusa veitchii* in Japan.
- Leptosphaeria schnelderiana** Rick. Small black spots on leaves of *Bambusa* sp. in Brazil.
- Loculistroma bambusae** Pat. and Charles. Sessile, dark-green or black fruiting bodies, resembling sclerotia of *Claviceps purpurea*, produced at the nodes. Diseased branches show a witches'-broom effect due to shortening of internodes. The fungus attacks *Phyllostachys* sp. in China.
- Malmeomyces pulchellus** Starb. On young branches of *Bambusa blumeana* and *B. spinosa* in Brazil and the Philippines.
- Melasmia phyllostachydis** Hara. On leaves of *Phyllostachys edulis* and *Phyllostachys* sp. in Japan.
- Miyoshiia fusispora** Kawamura. Elliptical brown spots on stems of *Arundinaria narihira* in Japan.
- Mosaic.** A typical Mosaic disease has been found on *Phyllostachys* plants imported from China. The disease is related to, and may be identical with, the sugar-cane mosaic.
- Mycosphaerella bambusae** Pat. On culms of *Phyllostachys reticulata* and *Bambusa* sp. in Japan and Venezuela.
- Mycosphaerella bambusicola** Miy. and Hara. Circular, elliptical, or irregular dull-brown spots, with black borders, often coalescing and causing death of surrounding areas, on leaves of *Phyllostachys bambusoides* and *P. puberula* in Japan and India.
- Mycosphaerella bambusifolia** Miy. and Har. Circular to irregular, fuscous, often coalescent, leaf spots on *Phyllostachys bambusoides*, *P. puberula*, and *Phyllostachys* sp. in Japan.
- Ophlobothis linearis** Rehm. Black stromata formed on lower leaf surfaces which die and split open. On *Chusquea* sp. in Brazil.
- Phalosphaeria bambusae** Miy. and Hara. Brown, later grayish, spots along veins of leaves, which are ultimately killed. On *Arundinaria simonii*, *Bambusa* sp. and *Sasa paniculata* in Japan.
- Phillipsiella graminicola** v. Hoehn. Black stromata on leaves of *Chusquea wetsteinii* in Brazil.
- Phyllachora bambusae** Syd. and Butl. Circular, black, slightly raised stromata on linear brown leaf spots on *Bambusa spinosa* and *B. blumeana* in the Philippines and Malabar.
- Phyllachora bambusina** Speg. Black, linear, stromata on leaves of *Bambusa* sp. in Brazil.
- Phyllachora caespitiola** Theiss. and Syd. Black stromata on upper surfaces of leaves of *Bambusa* sp. in Indo-China.
- Phyllachora chusqueae** P. Henn. and Lind. Black, linear, cushion-shaped stromata on leaves of *Chusquea quila* in Chile.
- Phyllachora eximia** Syd. Black convex stromata on sunken yellow or strawberry colored spots on leaves of *Arundinaria alpina* in Tanganyika.
- Phyllachora gracilis** Speg. Black stromata on lower surfaces of leaves of *Bambusa* sp. in Brazil.
- Phyllachora maculans** (Karst.) Theiss. and Syd. On leaves of *Bambusa* sp. in Indo-China.
- Phyllachora malabarensis** Syd. and Butl. Black stromata on yellowish-white spots on upper surfaces of leaves of *Bambusa* sp. in India.
- Phyllachora megastroma** Pat. On leaves of *Bambusa* sp. in Congo.
- Phyllachora orbicula** Rehm. Black stromata on small, round, yellowish spots on leaves of *Bambusa blumeana* and *B. spinosa* in China and the Philippines.
- Phyllachora phyllostachydis** Hara. Black stromata on leaves of *Phyllostachys reticulata* and *P. shiraiana* in Japan.
- Phyllachora shiraiana** Syd. Oblong to irregular black stromata on upper surfaces of leaves of *Arundinaria japonica*, *A. simonii*, *Bambusa* sp., *Phyllostachys bambusoides*, *Sasa paniculata*, *S. spiculosa*, *Schizostachyum acutiflorum*, and *S. diffusum* in India, the Philippines, and Japan.
- Phyllachora sinensis** Sacc. On *Bambusa* sp. in China.
- Phyllachora tjankorreh** Rac. Black circular, slightly convex, stromata on leaves of *Bambusa* sp., *Dinochloa scandens*, *D. tjankorreh*, and *Schizostachyum rotundifolium* in Java and the Philippines.
- Phyllosticta phyllostachydis** Siem. Leaf spot on *Phyllostachys kumazaca* in southern Russia.
- Phyllosticta take** Miy. and Hara. On leaves of *Bambusa* sp. and *Phyllostachys bambusoides* in Japan.
- Puccinia bambusoides** S. and Z. Brown rust pustules on leaves of *Phyllostachys bambusoides* in Japan.
- Puccinia corticioides** B. and Br. A rust producing yellowish-brown leathery crusts, up to 10 centimeters long, on stems of *Arundinaria hindsii*, *A. japonica*, *A. narihira*, *A. simoni*, and *Phyllostachys bambusoides* in Japan.
- Puccinia gracilentia** Syd. and Butl. Brown rust pustules on large, dull-brown sunken leaf spots on *Bambusa* sp. in India.
- Puccinia ignava** Arth. Brown powdery rust pustules on leaves of *Bambusa vulgaris*, *Bambusa* sp., and *Dendrocalamus giganteus* in Trinidad, Porto Rico, and Cuba.
- Puccinia kusanoi** Diet. Brown to black rust pustules on *Arundinaria fortunei*, *A. japonica*, *A. narihira*, *A. simoni*, *A. variabilis*, *Bambusa veitchii*, *Phyllostachys bambusoides*, *P. nigra*, *P. nigro-punctata*, *Sasa nipponica*, and *S. ramosa* in Japan.

**BAMBUSEAE**—Continued.

- Puccinia longicornis** Pat. and Har. Yellow-brown to dark-brown rust pustules on leaves of *Arundinaria japonica*, *A. simoni*, *Bambusa spinosa*, *B. tessellata*, *B. vulgaris*, *Sasa albo-marginata*, and *S. paniculata* in Japan, Ceylon, China, Indo-China, and the Philippines.
- Puccinia melanocephala** Syd. Brown to dark-brown rust pustules, covering under surfaces of leaves of *Arundinaria* sp., *Bambusa* sp., and *Phyllostachys* spp. in Japan and India. This rust has been introduced into Florida and Georgia.
- Puccinia mitrifomis** S. Ito. Dark-brown to black rust pustules on leaves of *Sasa paniculata* in Japan.
- Puccinia phyllostachydis** Kusano. Powdery rust-brown to dark-brown or black rust pustules on leaves of *Phyllostachys aurea*, *P. bambusoides*, *P. henonis*, and *P. marliacea* in Japan.
- Puccinia sasa** Kusano. Powdery rust-brown pustules on leaves of *Sasa borealis* in Japan.
- Puccinia xanthosperma** Syd. Dark-brown rust pustules on yellow, then gray, sunken leaf spots on *Bambusa* sp. in India and Japan.
- Puccinospora chusqueae** Speg. Elliptical dark brown spots, with yellow margins, on leaves of *Chusquea tenuiglumis* in Brazil.
- Pycnoderma bambusinum** Syd. More or less thickened circular spots on upper surfaces of leaves of *Bambusa vulgaris* and *Schizostachyum* sp. in the Philippines.
- Rosellinia rachidis** Rehm. On rachides of *Gigantochloa scribneriana* in the Philippines.
- Scirrhia bambusae** Malusio. (*Melanconium bambusae* Mal.) Brown spots and streaks appear on the branches and twigs and extend rapidly, soon involving the entire area. These spots turn whitish and dry out, finally being covered by the characteristic small black fruiting bodies. On *Bambusa gracilis* and *B. mitis* in Italy.
- Septoria bambusae** Brun. Pale-brown irregular spots, with broad black margins, on leaves of *Bambusa arundinacea* and *Phyllostachys puberula* in Japan.
- Septoria bambusella** Speg. Elliptical, indefinite, dark-brown spots on leaves of *Chusquea tenuiglumis* in Brazil.
- Shiraia bambusicola** P. Henn. Fleshy stromata on culms of *Bambusa* sp., *Phyllostachys bambusoides*, *P. puberula*, and *Sasa spiculosa* in Japan.
- Trichonectria bambusicola** Rehm. On leaves of *Bambusa blumeana* and *B. spinosa* in the Philippines.
- Uredo arundinariae** Syd. Brown rust pustules on leaves of *Arundinaria fortunei* and *A. japonica* in Japan and Ceylon.
- Uredo bambusarum** P. Henn. Powdery brown rust pustules, on round to linear spots, covering both leaf surfaces, of *Bambusa* sp. in Brazil.
- Uredo dendrocalami** Petch. Leaf rust on *Dendrocalamus strictus* in Ceylon.
- Uredo inflexa** Ito. Small powdery yellow-brown rust pustules on leaves of *Bambusa* sp. and *Sasa* sp. in Japan.
- Uredo ochlandrae** Petch. Brown rust pustules on leaves of *Ochlandra stridula* in Ceylon.
- Ustilago phyllostachydis** Syd. Black sclerotia formed in place of ovaries of *Phyllostachys bambusoides* in Japan.
- Ustilago shiraiana** P. Henn. This smut attacks the growing points of young branches and causes a witches'-broom effect. Infected branches are somewhat swollen and their growth stopped, the sori developing beneath the leaf sheaths and bracts, the powdery deep-brown spore masses finally breaking through. The hosts known for this very serious bamboo disease are *Arundinaria simonii*, *Arundinaria* sp., *Bambusa paniculata*, *B. veitchii*, *Phyllostachys bambusoides*, *P. henonis*, *P. puberula*, *P. quilloi*, *Sasa paniculata*, and *S. ramosa* in Japan and China. The disease has appeared at several points in California and Florida on imported plants but is now thought to have been eradicated.
- BANKSIA. SIEMUELLERA** Ag. Ornamental Australian evergreen shrubs or trees.
- Parodiella banksiae** Sacc. and Bizz. Numerous globose black aggregated perithecia on leaves of *B. marginata* in New South Wales.
- Phyllosticta banksiae** P. Henn. Brown spots, becoming pale, with yellow-brown margins, on leaves of *B. verticillata* in Germany.
- BAPHIA.** Barwood. Shrubs or small trees.
- Uredo baphiae** P. Henn. Black powdery rust pustules on leaves of *B. nitida* in central Africa.
- Uredo kampuluvensis** P. Henn. Powdery brown rust pustules on leaves of *B. cornifolia* and *Combretum baumium* in the Union of South Africa.
- BARBAREA. BARBAREA** Ag. WINTER CRESS. Yellow-flowered hardy biennials, sometimes called yellow rocket.
- Cercospora nasturtii** Pass. subsp. *barbareae* Sacc. On leaves of (*Campe barbarea*) *B. vulgaris* in France, Italy, and Russia.
- Peronospora barbareae** Gäum. Downy mildew on leaves of *Campe stricta* (*B. stricta*) and *C. barbarea* (*B. vulgaris*) in central and north Europe.
- Ramularia cochleariae** Cke. Circular white spots on leaves of *C. stricta* (*B. stricta*) and *Armoracea rusticana* (*Radicula officinalis*) in Great Britain and Germany.
- BARRINGTONIA.** Broad-leaved evergreen trees.
- Cercospora barringtoniae** Syd. Circular yellow to yellow-brown spots on leaves of *B. luzonensis* in the Philippines.
- Phyllachora barringtoniae** (B. and Br.) Sacc. On *B. speciosa* in Ceylon. A doubtful species.
- Phyllosticta microstegia** Syd. On leaves of *B. racemosa* in the Philippines.
- Phybaspora barringtoniae** Syd. On *Barringtonia* sp. in the Philippines.
- BASELLA.** VINE-SPINACH. Annual or biennial tropical pot herbs, sometimes called Malabar nightshade.
- Ascochyta basellae** P. Henn. Circular dull-brown spots, drying out lighter in color, on leaves of *B. rubra* in Brazil.
- Phyllosticta basellae** Rangel. Associated with *Stagonospora basellae* Rangel.
- Stagonospora basellae** Rangel. Circular to elliptical white leaf spots, with dark-purple margins, on *B. rubra* in Brazil.
- Uromyces basellae** Syd. Yellow to brown rust pustules on leaves of *B. rubra* in the Philippines.
- BAUHINIA.** Ornamental tropical trees and shrubs.
- Cercospora bauhiniae** Syd. On leaves of *B. malabarica* in the Philippines.
- Cercospora latimaculans** Wakef. Circular to irregular, often confluent, pale-brown leaf spots on *B. reticulata* and *Bauhinia* sp. in Gold Coast.
- Corticium salmonicolor** B. and Br. See Citrus.
- Phyllachora bauhiniae** (Wint.) Theiss. and Syd. "Tar-spot" disease on leaves of *B. vahlii* in India.
- Phyllachora phanerae** Racib. Circular black stromata on small yellow-green spots on leaves of *B. glauca* in Java.
- Phyllachora tenuis** (Berk.) Sacc. Black stromata on upper surfaces of leaves of *B. vahlii* and *Bauhinia* sp. in Mexico, Nicaragua, and Java.

**BAUHINIA**—Continued.

- Phyllosticta bakeri** Syd. On leaves of *B. malabarica* in the Philippines.
- Phyllosticta bauhiniae** Cke. Irregular brown spots on leaves of *Bauhinia* sp. in Vevezuela. Var. *europaea* Allesch. occurs on leaves of *Bauhinia* sp. in Germany.
- Phyllosticta bauhiniae-reticulatae** P. Henn. Circular to angular light-brown leaf spots on *B. reticulata* in the Union of South Africa.
- Phyllosticta bauhinicola** P. Henn. Circular dark fuscous spots on leaves of *Bauhinia* sp. in Brazil. Rangel has also described what is apparently the same species as new under the same name.
- Phyllosticta candicans** Pass. Irregular brown to whitish spots, with red-brown margins, on leaves of *B. aculeata* in Italy.
- Phyllosticta juruana** P. Henn. Circular red-brown spots, with surrounding yellow zones, on leaves of *Bauhinia* sp. in Brazil.
- Phyllosticta mayilae** Petch. Irregular red-brown leaf spots on *Bauhinia* sp. in Ceylon.
- Phyllosticta missionum** Speg. Large definite ashen-colored spots, with narrow purplish borders, on leaves of *Bauhinia* sp. in Argentina.
- Phyllostictiella globulifera** (Rabh.) F. Tassi. On leaves of *B. vahlii* in India.
- Ramularia bauhiniae** E. and E. Fruiting as a white powdery mildew on lower surfaces of leaves of *B. divaricata* in Jamaica.
- Rhizisma bauhiniae** Nees. Black fruiting bodies on leaves of *Bauhinia* sp. in tropical America.
- Uredo amazonensis** P. Henn. Rust on leaves of *Bauhinia* sp. in Brazil.
- Uredo bauhiniicola** P. Henn. Brown rust pustules on young twigs and on leaves of *B. heterophylla* and *B. rubiginosa* in Cuba and Brazil.
- Uromyces anthemophilus** Vestergr. Powdery brown pustules on calyces of *B. longifolia* in Brazil.
- Uromyces bauhiniicola** Arth. Powdery dark-brown rust pustules on both leaf surfaces of *B. pringlei* and *Bauhinia* sp. in Mexico.
- Uromyces congoensis** Syd. Powdery dark-brown rust pustules on leaves of *Bauhinia* sp. in the Congo.
- Uromyces dietelianus** Pazschke. Powdery brown rust pustules on leaves of *Bauhinia* sp. in Brazil.
- Uromyces fiebrigii** P. Henn. and Vestergr. Dark-brown rust pustules on leaves of *Bauhinia* sp. in Paraguay.
- Uromyces floralis** Vestergr. Powdery chestnut-brown rust pustules on leaves, flowers, calyces, and pedicels of *B. cuyabensis*, *B. hiemalis*, and *B. holophylla* in Brazil.
- Uromyces foveolatus** Juel. Chestnut-brown to black rust pustules on leaves of *Bauhinia* sp. in Brazil.
- Uromyces goyazensis** P. Henn. Powdery dull-brown rust pustules on branches and flowers of *Bauhinia* sp. in Brazil.
- Uromyces guatemalensis** Vestergr. Powdery dark-brown rust pustules on leaves of *B. inermis* and *Bauhinia* sp. in Guatemala and Costa Rica.
- Uromyces hemmendorffii** Vestergr. Powdery yellow-brown to chestnut-brown rust pustules on leaves of *B. forficata* in Brazil.
- Uromyces imperfectus** Arth. (*U. bauhiniae* [B. and C.] Vestergr. Brown rust pustules on leaves of *Bauhinia* sp. in Nicaragua and Jamaica.
- Uromyces jamaicensis** Vestergr. Powdery chestnut-brown rust pustules on leaves of *B. divaricata*, *B. paulletia*, and *B. porrecta* in Porto Rico, Cuba, Jamaica, Trinidad, and Mexico.
- Uromyces pannosus** Vestergr. Yellow-brown powdery rust pustules on leaves of *Bauhinia* sp. in Brazil.
- Uromyces peraffinis** Diet. Powdery chestnut-brown rust pustules on leaves of *Bauhinia* sp. in Brazil.
- Uromyces perlebiae** Vestergr. Powdery brown to dark brown rust pustules on leaves of *B. pentandra* in Brazil.
- Uromyces praetextus** Vestergr. Rust on leaves of *B. bongardi*, *B. cuyabensis*, and *B. hiemalis* in Brazil.
- Uromyces postulatus** Wakef. Leaf rust on *B. fassoglensis* in tropical Africa.
- Uromyces regius** Vestergr. Brown powdery rust pustules on leaves of *B. candicans* in Brazil.
- Uromyces superfixus** Vestergr. Rust on leaves of *B. mollis* in Argentina and Bolivia.
- Uromyces vestergreni** Syd. Ferruginous to dark-brown rust pustules on leaves of *B. tomentosa* in Ceylon and India.

**BEGONIA**. Succulent herbs grown mostly as potted plants for both flowers and variegated foliage.

- Aphelenchus olesistus** Ritz. Bos. (*A. ormerodis* Ritz. Bos.) This nematode or eel-worm forms brown streaks or patches on the leaves of *Acrostichum flagelliferum*, *Adiantum capillus-veneris*, *Ane-mone japonica*, *A. sylvestris*, *Asplenium* spp., *Aragena alpina*, *Begonia* spp., *Blechnum brasiliense*, *Calceolaria* sp., *Chrysanthemum indicum*, *Coleus* sp., *Crassula* sp., *Cypripedium* sp., *Cystopteris fragilis*, *Epipactis palustris*, *Erygium alpinum*, *Ficus* spp., *Fragaria* sp., *Gloxinia* sp., *Gymnogramme calo-melanos*, *Hepatica triloba*, *Heuchera sanguinea*, *Lomaria* sp., *Pteris* spp., *Ranunculus* spp., *Saintpaulia ionantha*, *Salvia* sp., *Scabiosa* sp., *Spiraea astilboides*, and other ornamentals, and on the fronds of various species of cultivated ferns. The disease is reported from England and Holland. It occurs in greenhouses in the United States and out of doors in the South to some extent. Further importations would be most undesirable.
- Coleosporium begoniae** Arth. Yellow powdery rust pustules on the lower surfaces of leaves of *Begonia* spp. in Mexico and Italy.
- Gloeosporium begoniae** Magnaghi. Produces reddish-brown spots on leaves of *Begonia* spp. in Italy.
- Mycosphaerella begoniae** Pat. Causes spots on the lower surfaces of leaves of *Begonia* spp. in Ecuador.
- Myxosporium sordidum** F. Tassi. On *B. argyrostigma* in South America and Europe.
- Phyllachora begoniae** Pat. Produces black, shiny, more or less irregular spots on the upper surfaces of the leaves of *Begonia* spp. in Ecuador.
- Phyllosticta begoniae** Brun. Circular light olive-colored leaf spots on *B. credneri*, *B. metallica*, and *B. tuberosa* in Italy, France, and Portugal.
- Phyllosticta begoniae** Rangel. Reported from Brazil as the cause of a leaf spot of cultivated *Begonias*. May be the same species as the preceding but, if not, should be renamed.
- Puccinia granularis** Kalchbr. & Cke. See Pelargonium.
- BELAMACANDA**. BLACKBERRY LILY. Hardy herbaceous perennials, sometimes called leopard flower.
- Puccinia belamacandae** (P. Henn.) Diet. Powdery brown to black rust pustules on brown spots with red-brown margins, on leaves of *B. chinensis* and *B. punctata* in Japan.
- Uredo belamacandae** P. Henn. Brown rust pustules on leaves of *B. chinensis* in India.
- BELIS**. See Cunninghamia.
- BELLEVALIA**. Bulbous plants. Now referred to *Hyacinthus* (q. v.).
- Septoria bellevaliae** Pat. Produces elongate (2 to 4 cm.) white spots with red margins on leaves of *B. dubia* in Algeria.

**BELLIDIASTRUM.** Herbaceous perennials.

**Entyloma bellidiastrum** Maire. Circular, whitish spots, turning brown, on leaves of *B. michelii* in Tyrol.

**Entyloma calendulae** (Oud.) De By. See *Calendula*.

**Puccinia firma** Diet. Brown rust pustules on leaves of *B. michelii* and *Carex firma* in Tyrol and Germany.

**Septoria bellidiastrum** Allesch. On leaves of *B. michelii* in Bavaria.

**BELLIS.** ENGLISH DAISY. Low perennial herbs.

**Entyloma bellidis** Krieger. Circular white or yellowish spots, which are often confluent so as to occupy the entire area of leaf blades of *B. perennis* in Dalmatia, Denmark, and Germany.

**Puccinia distincta** McAlp. Orange-yellow and brown rust pustules on leaves of *B. perennis* in Australia and Tasmania.

**Ramularia bellidis** Sacc. Subcircular dull yellow spots on leaves of *B. perennis* in Italy.

**Septoria bellidicola** Rob. & Desm. White spots with red margins on leaves of *B. perennis* in Dalmatia and Italy.

**BERBERIS.** (Including *Mahonia*, Oregon grape, holly grape.) BARBERRY. Deciduous or evergreen shrubs cultivated for their ornamental leaves, flowers, and fruits.

**Accidium aridum** Diet. and Neg. Rust on leaves of *B. buxifolia* and *B. heterophylla* in Chile.

**Accidium berberis-ruscifolia** P. Henn. Yellow rust pustules covering the under sides of leaves of *B. ruscifolia* in Argentina.

**Accidium berberidis-thunbergii** P. Henn. Yellow rust pustules on round red spots on under surfaces of leaves of *B. thunbergii* in Japan.

**Accidium haussknechtianum** P. Henn. Golden-yellow rust pustules on leaves, petioles, and fruits of *B. crataegina* in Persia.

**Accidium levileanum** P. Magn. Rust on leaves of *B. buxifolia* in Chile.

**Accidium montanum** Butl. Rust pustules on red leaf spots on *B. aristata*, *B. coriaria*, and *B. lycium* in India.

**Accidium tubiforme** Diet. and Neg. Rust on leaves of *B. buxifolia* in Chile.

**Ascochyta australis** Speg. Circular to irregular gray-brown spots, with yellow margins on leaves of *B. glauca* in Argentina.

**Ascochyta berberidina** Sacc. On branches of *B. vulgaris* in France and Russia.

**Dictyodthis berberidis** (Rehm.) Theiss. and Syd. On branches of *B. buxifolia* in Chile.

**Didymosphaeria epidermidis** (Fr.) Fekl. var. **macrospora** Elias. On branches of *B. vulgaris* in Sweden.

**Dothichiza carneofusca** v. Hoeh. On branches of *B. vulgaris* in Austria.

**Gambleola cornuta** Mass. On leaves of *B. nepalensis* in India.

**Gloeosporium japonicum** Hemmi. Anthracnose on *Mahonia japonica* in Japan.

**Heterosporium berberidis** Ranoj. On branches of *B. vulgaris* in Serbia.

**Laestadia berberidis** Delacr. On branches of *B. vulgaris* in France.

**Melasmia berberidis** Thuem and Wint. Small brown spots on leaves of *B. vulgaris* in Austria and Germany.

**Microspheera berberidis** (DC.) Lév. Powdery mildew on leaves of *B. aquifolium* and *B. vulgaris* in Europe, Cyprus, Transcaucasia, Turkestan, and Japan.

**Mycosphaerella australis** (Speg.) Lind. On leaves of *B. ilicifolia* in Chile.

**Mycosphaerella berberidis** (Auersw.) Lind. On leaves of *B. vulgaris* in Belgium and Switzerland.

**Phoma berberidicola** Vesterg. On branches of *B. vulgaris* in Sweden and Switzerland.

**Phragmedothis berberidis** (Alm. and Cam.) Theiss. and Syd. On branches of *B. vulgaris* in Portugal.

**Phyllosticta aquifolii** Allesch. On leaves of *Mahonia japonica* in Denmark.

**Phyllosticta asiatica** Cke. Circular to irregular dull-brown spots, with dark, purplish-brown margins, on leaves of *B. asiatica* in Great Britain.

**Phyllosticta berberidicola** Speg. Spots on leaves of *B. laurinea* in Uruguay.

**Phyllosticta berberidis** Rbh. Irregular gray spots, becoming whitish, on leaves of *B. neubertii* and *B. vulgaris* in France, Italy, Denmark, Russia, Austria, and Germany.

**Phyllosticta mahoniae** (Thuem.) Keissl. On leaves of *Mahonia aquifolium* in Italy, France, Russia, and Denmark.

**Phyllosticta mahoniaecola** Pass. Large brown indefinite spots on leaves of *Mahonia aquifolium*, *M. beali*, and *M. japonica* in Italy, France, and Great Britain. Var. *aquifolii* P. Brun. is the same as the species, except that the leaf spots have definite brown margins.

**Phyllosticta spegazziniana** Keissl. On leaves of *Mahonia aquifolium* in Argentina.

**Phyllosticta westendorpii** Thuem. Angular white spots, with red margins, on leaves of *B. altaica* and *B. vulgaris* in Yugoslavia, Siberia, Belgium, and Germany.

**Puccinia antarctica** Speg. Dark-brown rust pustules on leaves of *B. buxifolia* (*B. dulcis*) in Patagonia.

**Puccinia arrhenatheri** (Kleb.) Eriks. See *Arrhenatherum*.

**Puccinia barri-aranae** Diet. and Neg. Chestnut-colored rust pustules on leaves of *B. buxifolia* in Chile.

**Puccinia berberidis** Mont. Yellow to yellow-brown rust pustules on circular or irregular leaf spots on *B. crassifolia*, *B. darwinii*, *B. empetrifolia*, *B. glauca*, *B. heteropoda*, and *B. spinulosa* in Chile.

**Puccinia berberidis-trifoliae** Diet. and Holw. Black, powdery, rust pustules on small brown sunken spots on leaves and petioles of *B. trifolia* in Mexico.

**Puccinia droogensis** Butl. Rust on leaves of *B. aristata* in Ceylon and India.

**Puccinia meyeri-alberti** P. Magn. Brown rust pustules on circular leaf spots on *B. buxifolia*, *B. congestifolia*, *B. darwini*, *B. linearifolia*, and *B. pearcei* in Chile.

**Puccinia naumanniana** (P. Magn.) Diet. Black powdery rust pustules on leaves of *B. buxifolia* in Chile.

**Puccinia stolpiana** (P. Magn.) Diet. and Neg. Yellow to black powdery pustules on leaves of *B. buxifolia* and *B. ilicifolia* in Chile.

**Septoria berberidis** Niessl. Subcircular dull-brown spots, with dark purple margins on leaves of *B. vulgaris* in Yugoslavia, Russia, Italy, Switzerland, and Austria.

**Septoria mahoniae** Passer. Small brown spots on leaves of *Mahonia aquifolium* in Russia, Italy, and Denmark.

**Uropyxis quitensis** Lagh. Rust on leaves of *Berberis* sp. in Ecuador.

**BETA.** BEET. Sugar beet. Cultivated for thick edible roots, edible leaves, and for sugar production.

**Actinomyces** spp. Several species of *Actinomyces*, including *A. albus* (R. D.) Gasp., *A. intermedius* (Krug.) Wr., and *A. xanthostroma* Wr., said to cause scab of beet roots (*B. vulgaris*) have been reported from Europe. Their relation to *A. scabies* is uncertain.

**Aphanomyces laevis** de By. Causes a soft rot of sugar beet (*B. vulgaris*) roots in Europe.

**Ascochyta betae** Prill. and Delacr. On petioles of *B. vulgaris* in France.

**BETA**—Continued.

- Ascochyta beticola** Prill. and Delacr. On petioles of *B. vulgaris* in France. Probably not distinct from above species.
- Bacillus** spp. Bacteriosis has been reported from a number of countries, particularly in Europe, as a serious disease of the beet (*B. vulgaris*) under various names, including *B. betae* Mig., *B. bussei* Mig., *B. lacerans* Mig., *B. mycoides* Mig., and *B. tabificans* Delacr. The status of these species and their relation to *B. carotovorus* is most uncertain.
- Bacillus** sp. Bacterial scab of *B. vulgaris* in Korea.
- Entyloma betiphilum** Bub. Leaf smut on *B. vulgaris* in Austria. A doubtful species.
- Helicobasidium mompa** Tan. See Morus.
- Leptosphaeria circinans** (Fekl.) Sacc. See Asparagus.
- Moniliopsis aderboldi** Ruhl. See Solanum.
- Peronospora schachtii** Fekl. The downy mildew of beet (*B. maritima* and *B. vulgaris*) is serious in Europe, Egypt, Japan, and Argentina. The disease occurs to a limited extent in New York, Minnesota, and California. Delicate gray patches of mildew appear on lower leaf surfaces. Young central leaves are most frequently attacked and the growth of the plants checked. Seedlings may be killed outright.
- Pionnotes betae** Sacc. Yellow spots coalescing to form crustlike areas and cankers of roots of *B. vulgaris* in Great Britain.
- Pleospora putrefaciens** (Fekl.) Frank. Causes a serious disease of beets (*B. vulgaris*) in Germany.
- Ramularia betae** Rostr. Numerous subcircular grayish spots with reddish margins on both surfaces of leaves of *B. trigyna* and *B. vulgaris* in Bulgaria, Belgium, Denmark, Holland, and Russia.
- Rhizoctonia napi** West. See Brassica.
- Septeria betae** Westd. Pale-brown spots with white centers and dark-brown margins, on leaves of *B. vulgaris* in Australia and Belgium. Reported from Indiana.
- Scorpidium betae** Nemeec. On roots of *B. vulgaris* in Europe.
- Typhula betae** Rostr. Attacks seedlings of *B. vulgaris* in Sweden, Denmark, Holland, Great Britain, and the Azores and is often destructive.
- Uredo marmoxiae** Speg. Rust on leaves and stems of *B. procumbens* in the Canary Islands.
- Uromyces betae** (Pers.) Lév. Yellow to brown rust pustules on leaves of *B. cicla*, *B. maritima*, and *B. vulgaris* in Argentina, Uruguay, Union of South Africa, Australia, New Zealand, and Europe. Found sparingly in California and Colorado.
- Urophlyctis leproides** (Trab.) P. Magn. Forms nodular outgrowths from the upper portion of the roots, which often become as large as walnuts. These outgrowths are modified roots and remain attached by narrow bases. On *B. vulgaris* in New Zealand, Algeria, Argentina, Great Britain, France, Belgium, and Sweden. Occurs scatteringly in California and has been found in Iowa.
- Urophlyctis pulposa** (Walbr.) Schroet. See Atriplex.
- A disease characterized by orange-colored spots, irregular swellings, and distortion of the foliage of beet has been reported from northern France, the cause being unknown.
- BETULA. BIRCH.** Forest and shade trees with handsome green foliage and ornamental bark.
- Didymosphaeria latebrosa** Mont. On leaves of *B. alba* in Belgium.
- Exoascus alpinus** (Johans.) Sacc. On leaves and branches of *B. nana* in Scandinavia.
- Exoascus betulae** Fekl. Fruiting areas on lower leaf surfaces of *B. alba*, *B. odorata*, *B. pubescens* and *B. verrucosa* (*B. pendula*) in middle Europe.
- Exoascus betulinus** (Rostr.) Sadeb. Produces witches'-brooms on *B. alba*, *B. odorata* and *B. pubescens* in north and middle Europe. Reported from Wisconsin.
- Exoascus janus** Thomas. Large, circular, pale rose-colored spots on both leaf surfaces of *B. verrucosa* in Switzerland and Germany.
- Exoascus nanus** (Johan.) Sacc. Produces witches'-brooms and yellow-green leaf spots on *B. nana* and *B. odorata* in Norway and Sweden.
- Exoascus turgidus** Sadeb. On leaves of *B. alba*, *B. odorata*, *B. pubescens*, and *B. verrucosa* in Finland, Russia, Sweden, Denmark, Switzerland, Great Britain, Austria, and Germany.
- Guignardia betulae** (Auersw.) Sacc. and Trott. On leaves of *B. verrucosa* in Denmark and Germany.
- Macrophoma sydowiana** (Bres.) Sacc. and Syd. On leaves of *B. alba* in Germany.
- Marsonia betulae** (Lib.) Sacc. Irregular, radiate, dull-brown spots on leaves of *B. alba* and *B. verrucosa* in Europe.
- Myxosporium devastans** Rostr. Causes withering and die-black of young twigs of *B. alba*, *B. verrucosa*, and *Acer pseudoplatanus* in Denmark.
- Phyllosticta betulae** Oud. On leaves of *B. alba* in Holland.
- Phyllosticta betulina** Sacc. On leaves of *B. alba* in Italy and France. Reported from New York.
- Plowrightia virgultorum** Sacc. Black knot of birch. Shoots and branches from 1 to 4 years old are attacked, narrow elongated stromata forming, which are olive to black in color and carbonaceous in texture. Leaves on infected branches wilt and die and the twigs die back. A witches'-broom effect is produced finally. On *B. alba*, *B. nana*, *B. pubescens*, and *B. verrucosa* in Great Britain, Sweden, Finland, Switzerland, Russia, Denmark, and Germany. Theissen and Sydow delete this species.
- Rhytisma nervale** (Alb. and Schw.) Rehm. See Alnus.
- Sclerotinia betulae** Wor. Attacks the catkins of *B. alba*, *B. papyrifera*, and *B. verrucosa* in Japan, Russia, and Denmark.
- Septoria betulina** Pass. Irregular, often confluent, gray spots on leaves of *B. alba* and *B. pubescens* in Russia, Denmark, Tyrol, and Italy. Reported from Wisconsin.
- Taphrina aurea** (Pers.) Fr. Attacks seedlings of *Betula* sp. in Great Britain.
- BIGNONIA. TRUMPET CREEPER.** Woody climbers.
- Ascochyta tweediana** Penz. and Sacc. On capsules of *B. tweediana* in Italy.
- Cercospora bignoniaceola** Speg. Irregular, often confluent, gray spots on leaves of *Bignonia* sp. in Brazil.
- Cercospora unguis-cati** Speg. Subcircular dark-purple spots on both leaf surfaces of *B. unguis-cati* (*Doxantha unguis-cati*) in Argentina.
- Colletotrichum bignoniae-igneae** Rangel. Anthracnose on leaves of *Bignonia* sp. in Brazil.
- Phyllachora amphigena** Speg. (*P. nitidula* Pat. and Gaill.) Black stromata on pale brown spots on leaves of *Bignonia* sp. and *Clytostoma callistegioides* in Venezuela, Paraguay and Brazil.
- Phyllachora nitidula** Pat. Black stromata on leaves of *Bignonia* sp. in Venezuela. A doubtful species.
- Phyllachora tenuis** Speg. Tar-spot on leaves of *Bignonia* sp. in Argentina. This species should be renamed, since *P. tenuis* (B. and C.) Sacc. has priority.
- Phyllosticta tweediana** Penz. and Sacc. On capsules of *B. tweediana* in Italy.
- Pleospora briosiana** Luigi. Large hazel-colored blotches, with black borders, on leaves of *B. buccinatoria* (*Phaedranthus buccinatorius*) in Italy.
- Puccinia appendiculatoides** P. Henn. Yellow rust pustules on round or effuse circular leaf spots on leaves of *Bignonia* sp. in Brazil.

**BIGNONIA**—Continued.

- Puccinia bignoniacearum** Speg. Brown rust pustules on leaves of *Bignonia* sp. in Brazil.  
**Uredo cuticulosa** E. and E. Brown rust pustules on leaves of *Bignonia* sp. in Nicaragua.  
**Uropyxis rickiana** P. Magn. Irregular, woody galls, on which brown rust pustules occur on trunks of *Bignonia* sp. in Brazil.
- BIHAI.** See Heliconia.  
**BIKUKULLA.** See Dicentra.  
**BISCHOFIA.** Ornamental tropical trees.  
**Phyllachora bischofiae** Syd. Irregular, black, shining stromata on leaves of *B. javanica* (*B. trifoliata*) in India.  
**BIXA.** ANATTO. Ornamental tropical trees. Red coloring matter produced from the pulp surrounding the seeds.  
**Cercospora bixae** Allesch. Large, irregular, brown spots, with surrounding yellow zones, on leaves of *B. orellana* in Brazil.  
**Ovularia bixae** Rac. On leaves of *B. orellana* in Java and Ceylon.  
**Phyllosticta bixina** Young. Subcircular to irregular pale yellow spots, surrounded by purplish-black borders, on leaves of *B. orellana* in Porto Rico.  
**Rhizoctonia lamellifera** Small. See Grevillea.  
**Uredo bixae** Arth. Brown powdery rust pustules on leaves of *B. orellana* in Porto Rico.
- BLECHNUM.** Ferns.  
**Aphelenchus olesistus** Ritz. Bos. See Begonia.  
**Milesina blechni** Syd. Brown rust pustules on leaves of *B. occidentale*, *B. spicant*, and *Lygodium polymorphum* in Colombia, Trinidad, Great Britain, Denmark, France, Spain, Switzerland, Austria, and Germany.  
**Polystomella paichella** (Speg.) Theiss. See Alsophila.  
**Uredinopsis mayoriana** Diet. Rust on leaves of *B. blechnoides* in Colombia.  
**Uredo blechnicola** P. Henn. Powdery yellow rust pustules on leaves of *B. volubile* in Brazil.  
**Uredo orientalis** Racib. Brown, erumpent, rust pustules deforming the leaves of *B. orientale* in Java.
- BLETIA.** Terrestrial orchids. See Orchidaceae.  
**BLUMENBACHIA.** SALOA Ag. STING LILY. Ornamental annuals.  
**Septoria blumenbachiae** Speg. Indefinite light-brown spots on leaves of *B. urens* in Argentina.  
**BOCCONIA.** PLUME POPPY. Tall garden herbs.  
**Puccinia bocconiae** Mayor. Rust on leaf blades and petioles of *B. frutescens* in Colombia.  
**BOEHMERIA.** Ramie. Herbs, shrubs, or trees. Tropical fiber and ornamental plants.  
**Cercospora boehmeriana** Wor. On leaves of *B. nivea* in Russia.  
**Colletotrichum boehmeriae** K. Saw. Scattered ashen-brown circular spots on leaves and fusiform cankers on stems of *B. nivea* in Japan and Formosa.  
**Puccinia boehmeriae** P. Henn. Yellow-brown rust pustules on leaves of *Boehmeria* sp. in Brazil.  
**Pucciniastrum boehmeriae** Syd. Yellow-brown rust pustules on leaves of *B. biloba*, *B. grandiflora*, *B. japonica*, *B. longifolia*, *B. sieboldiana*, *B. spicata*, and *B. tricuspis* in Japan and the Philippines.  
**Ramularia boehmeriae** Fuji. Causes a root and rhizome rot of *B. nivea* in Japan.  
**Septoria boerhaviae** Pat. On leaves of *B. verticillata* in Abyssinia.  
**Uredo boehmeriae** Diet. Rust on leaves of *B. platyphylla* in Ceylon.
- BOLTONIA.** Asterlike herbs.  
**Coleosporium asterum** (Diet.) Syd. See Aster.
- BOMAREA.** Twining, herbaceous plants of the *Amaryllidaceae*.  
**Aecidium bomareae** Mayor. Yellow rust pustules on leaves and stems of *B. caldasiana* and *B. potacocensis* in Colombia, South America.  
**Heterosporium allii-bomareae** Pat. On leaves of *Bomarea* sp. in Ecuador.  
**Puccinia bomareae** P. Henn. (*Uredo bomareae* Lagerh.) This rust produces elongated lemon-yellow pustules, which later become dark-brown, on lower surfaces of leaves of *B. acutifolia*, *B. edulis*, *B. ovata*, and other species in Ecuador, Mexico, and Brazil.  
**Puccinia pallor** Arth. and Holw. Yellow to brown powdery rust pustules on lower surfaces of leaves of *B. acutifolia* in Costa Rica and Guatemala.  
**Urocystis bomareae** Diet. and Neg. Black powdery spore masses form in swellings on the stems of *B. salsilla* in Chile.  
**Uromyces bomareae** P. Henn. Bright yellow, then brown, rust pustules on the lower surfaces of the leaves of *Bomarea* sp. in Brazil.
- BOMBAX.** Tropical shrubs and trees, supplying a useful floss from the cottony contents of the pods, sometimes known as silk cotton tree.  
**Chrysonyxa bombacis** Petch. Rust on leaves of *B. malabaricum* in Ceylon.  
**Cladotrichum folicola** (Niessl.) Ferro. Effuse black areas on leaves of *B. malabaricum* in India.  
**Fomes lignosus** Klotzsch. See Hevea.  
**Gloeosporium bombacis** Putte. Irregular, then confluent, dull-brown areas on leaves of *B. insignis* in Brazil.  
**Puccinia bombacis** Diet. Brown rust pustules on leaves of *Bombax* sp. in Colombia.  
**Uredo bombacis** Petch. Rust on leaves of *B. malabaricum* in Ceylon.
- BORAGO.** BORAGE. Coarse, annual herbs, sometimes cultivated for culinary purposes.  
**Cercospora** sp. Leaf spot on *B. officinalis* in Cuba.  
**Cylindrosporium myosotidis** Sacc. See Myosotis.  
**Entylessa serotinum** Schroet. Smut sori in numerous white, then brown, circular leaf spots on *B. officinalis*, *Pulmonaria officinalis*, *Symphytum officinalis*, and *S. tuberosum* in Denmark, France, Austria, and Germany. Ciferri has described the form on *Borago* as *E. borraginis* Cif.
- BORASSUS.** Palmyra palm. See Palmae.  
**BORONIA.** Ornamental shrubs.  
**Puccinia boroniae** P. Henn. Dark cinnamon-colored rust pustules on branches of *B. spinescens* in Australia.
- BOTOR.** See Psophocarpus.  
**BOUGAINVILLE.** BUGINVILLE Ag. Tropical woody climbers with brilliant colored bracts.  
**Puccinia bougainvilleae** (Speg.) Schroet. Yellow to dark-brown rust pustules, on circular yellow spots, on leaves and branches of *B. frondosa* and *B. stipitata* in Argentina.
- BOUSSINGAULTIA.** Madeira vine. Herbaceous perennial vines.  
**Cercospora boussingaultiae** Roum. On leaves of *B. baselloides* in France.  
**Phyllosticta boussingaultiae** Speg. Small whitish spots, with purple borders, on leaves of *B. baselloides* in Argentina.
- BOUVARDIA.** Small flowering shrubs.  
**Uromyces bouvardiae** Syd. Yellow and brown to black rust pustules on leaves of *B. hirtella*, *B. leiantha*, *B. versicolor* and *Bouvardia* sp. in Mexico and Guatemala.

**BRACHYCHITON.** BOTTLE TREE. Australian trees.

**Phyllosticta sterculiæ brachychiti** Wint. On leaves of *B. diversifolium* (*B. populneum*) in Portugal.  
**Septoria ochracea-maculans** Theum. Irregular, dull yellow spots, with narrow dark purple borders, on leaves of *B. diversifolium* (*B. populneum*) in Portugal.

**BRACHYCOME.** SWAN RIVER DAISY. Australian herbs.

**Puccinia brachycomes** McAlp. Yellow and black rust pustules on swollen and distorted portions of leaves, and on stems and branches of *B. ciliaris*, *B. diversifolia*, *B. pachyptera* and *B. scapiformis* in Australia.

**BRASSAVOLA.** Epiphytic orchids. See Orchidaceae.**BRASSICA.** Cabbage. Mustard. Turnip. Rutabaga. Cauliflower. Kale. Kohlrabi. Brussels sprouts. Annual and perennial herbs.

**Ascochyta brassicæ** Thuem. Large, irregular, indefinite, yellow-gray spots on leaves of *B. oleracea capitata* and other varieties in Europe.

**Bacillus brassicaevorus** Del. Bacteriosis of *B. oleracea capitata* and *botrytis* in France. Probably not distinct from *B. carotovorus*.

**Bacillus** sp. (?) Bacterial leaf spot of *B. oleracea capitata* and *B. rapa* in Guam.

**Cercospora brassicæ-campestris** Rangel. On leaves of *B. campestris* in Brazil.

**Cercospora brassicicola** P. Henn. Circular, numerous pale spots, with dull-brown surrounding zones, on leaves of *B. chinensis* and *B. pekinensis* in Japan and the Philippines.

**Colletotrichum brassicæ** Schulz. and Sacc. Anthracnose on stems of *B. oleracea* in Yugoslavia.

**Cylindrosporium brassicæ** Fautr. and Roum. Numerous, large, irregular, dirty white spots on leaves of *B. campestris* and *B. napus* in Yugoslavia and France. Reported from Missouri and Georgia.

**Gloeosporium concentricum** (Grev.) B. and Br. Circular bleached spots on leaves of *B. napus*, *B. oleracea* and *B. rapa* in Japan, Great Britain, Denmark, and Germany.

**Monilopsis aderholdi** Ruhl. See Solanum.

**Mycosphaerella brassicicola** (Duby.) Lind. (*Phyllosticta brassicæ* [Curr.] West.) The "ring-spot" disease of *B. oleracea* var. *botrytis*, *gemmifera*, and *capitata* occurs in Europe, Australia, New Zealand, Argentina, Uruguay, and California. Numerous definite, circular, light brown to gray spots occur on the leaves, surrounded by olive-green or blue-green borders. Affected leaves turn yellow. The disease develops rapidly after harvesting and is often serious enough to cause heavy losses in marketing.

**Olpidiaster radicis** (De Willd.) Pascher. See Linum.

**Olpidium brassicæ** Wor. Attacks the roots of seedlings of *B. oleracea* var. *capitata* and *Brassica* spp., causing a "damping off," in Russia, Finland, Sweden, and Belgium. Reported from Wisconsin.

**Olpidium radiceolum** De Willd. On roots of *B. oleracea* and *Bursa bursapastoris* in Russia and Belgium.

**Ovularia brassicæ** Bres. and Allesch. Subcircular or irregular, often confluent, white spots on leaves of *B. napus* and *B. oleracea* in Germany.

**Phoma napa-brassicæ** Rostr. Causes a dry rot of roots of *B. napa* and *B. oleracea* in New Zealand, Great Britain, Denmark, and Holland. Reported from Canada.

**Phyllosticta brassicina** Sacc. Circular to angular, dull, greenish-gray spots on leaves of *B. oleracea* in Italy and Malta.

**Phyllosticta napi** Sacc. Indefinite spots on leaves of *B. napa* and *B. oleracea* in France and Italy.

**Pseudomonas (Bacterium) destructans** Potter. Causes a white rot of *B. napus* and *B. oleracea* in Great Britain and *Raphanus sativus* in Japan. Probably not distinct from *B. carotovorus* Jones.

**Ramularia rapæ** Pim. Irregular spots on leaves of *B. napa* in Great Britain.

**Rhizoctonia napi** West. Covers stems, pods, leaves, and roots with white mycelium in which sclerotia develop, white at first, turning black on the outside. Causes a dry rot of infected areas, seriously diseased plants breaking over at the ground level. The recognized hosts are *Amaranthus tristis*, *Argemone mexicana*, *Avena sativa*, *Beta bengalensis*, *Brassica campestris*, *Calamintha*, *Cannabis sativa*, *Chenopodium album*, *Cicer arietinum*, *Cnicus arvensis*, *Fumaria parviflora*, *Hordeum vulgare*, *Lathyrus sativus*, *Lens esculenta*, *Leucas* sp., *Linum usitatissimum*, *Medicago lupulina*, *Pisum sativum*, *Scoparia dulcis*, *Triticum vulgare*, and *Vicia hirsuta*. Shaw states that this species should be transferred to *Botrytis*. Other workers consider it synonymous with *Sclerotinia sclerotiorum* Mass.

**Sporidesmium brassicæ** Mass. Indefinite olive-gray or green spots, on leaves and pods of *B. campestris* in India and China.

**Sporidesmium exitiosum** Kuehn. On stems and pods of *B. napa*, *B. oleracea*, *B. rapa*, and *B. arvensis* in Japan, Sweden, Denmark, and Germany. Also reported from Canada. *Leptosphaeria napi* (Fckl.) Sacc. is said to be the perfect stage of this species.

**Urocystis coralloides** Rostr. Affected plants are stunted and galls up to one and one-half inches in diameter form on the roots. On *Brassica* sp. (mustard) in India.

**Uromyces brassicæ** Niessl. On *B. oleracea* in France. A very doubtful species.

**BRICKELLIA.** COLEOSANTHUS Ag. Herbs or small shrubs.

**Puccinia decora** Diet. Dull-brown rust pustules on leaves of *Brickellia* sp. in Mexico.

**Puccinia praemorsa** Diet. and Holw. Rust on leaves of *B. veronicaefolia* in Mexico.

**Puccinosira brickelliae** Diet. and Holw. Brown rust pustules on leaf blades and petioles and stems of *B. adenocarpa*, *B. cavanillesii*, *B. secundiflora*, and *Coleosanthus canavillesius* in Mexico and Guatemala.

**BRIZA.** QUAKING GRASS. Annual or perennial grasses.

**Dilophia graminis** (Fckl.) Sacc. See Triticum.

**Septoria crispiniana** Gz. Frag. On leaves of *B. maxima* in Spain.

**Tilletia brizæ** Ule. Brown striations on leaves which finally rupture exposing powdery black masses of smut spores. On *B. media* in Germany.

**BROMELIA.** Fleshy leafed Bromeliads.

**Perisporium bromeliæ** Stevens. Forms smoky patches on leaves of *B. pinguin* in Porto Rico.

**Phyllosticta bromeliæ** Alm. and Cam. Whitish spots on leaves of *B. acanga* in Portugal.

**BROMUS.** BROMEGRASS.

**Exobasidium graminicolum** Bres. See Arrhenatherum.

**Helminthosporium fragosi** Bub. On leaves of *B. sterilis* in Spain.

**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.

**Leptosphaeria culmicola** (Fr.) Karst. On culms of *B. erectus*, *Festuca arundinacea*, *F. sylvatica*, *Phalaris arundinacea*, *Secale cereale*, and *Triticum sativum* in Denmark, Italy, Sweden, France, and Germany.

**Mycosphaerella longissima** Fckl. On leaves of *B. asper* and *B. maximus* in Italy, Switzerland, Denmark, and Germany.

**Mycosphaerella recutita** (Fr.) Johans. See Aira.

**Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.

**Phyllachora bromi** Fckl. Tar spot on leaves and leaf sheaths of *B. japonicus*, *B. unioides*, *Brachypodium* sp., and *Dactylis* sp. in Argentina, Japan, Algeria, Italy, France, and Germany.

**BROMUS**—Continued.

- Phyllosticta bromi** Poteb. On leaves of *B. patulus* in Russia.
- Puccinia brachypus** Speg. Linear, brown, rust pustules on leaves of *B. auleticum*, *Lolium perenne*, and *Triticum sativum* in Argentina.
- Puccinia bromina** Eriks. Yellow and brown to black rust pustules on leaf blades and petioles of *Pulmonaria montana*, *Symphytum officinalis*, and nearly 50 species of *Bromus* in Argentina, Australia, Asia Minor, Japan, and all of Europe.
- Puccinia bromi-japonica** S. Ito. Yellow-brown to black linear rust pustules on leaves of *B. japonicus* in Japan.
- Septoria affinis** Sacc. Linear whitish spots with rufous margins, on leaves of *Brachypodium pinnatum*, *Bromus inermis*, and *B. mollis* in Italy and Russia.
- Septoria bromicola** Speg. On leaves of *B. unioloides* in Chile.
- Septoria bromivora** Speg. On leaves of *Bromus* sp. in Argentina.
- Tilletia belgradensis** P. Magn. Smut in inflorescences of *B. secalinus* in Yugoslavia.
- Tilletia holci** (West.) Rostr. See *Holcus*.
- Tilletia velenovskyi** Bub. Smut sori in ovaries of *B. arvensis* and *B. secalinus* in Bulgaria. Said to be the same as *T. guyotiana* Har. which is of limited occurrence in the western United States.
- Uredo bromi-pauciflorae** S. Ito. Yellow-brown rust pustules on leaves of *B. pauciflorus* in Japan.
- BROUSSONETIA PAPHYRIUS** Ag. PAPER MULBERRY. Ornamental trees or shrubs, bark used for paper making.
- Gibberella moricola** (Ces. and De N.) Sacc. See *Morus*.
- Phyllosticta broussonetiae** Trav. and Migli. Irregular whitish areas, often involving entire leaves of *B. papyrifera* in Italy.
- BROWNEA**. Small, tropical evergreen trees.
- Fomes lamaoensis** Murr. See *Hevea*.
- Phyllachora juruensis** P. Henn. Black stromata on small brown spots scattered irregularly over leaves of *Brownea* sp. in Brazil.
- BRUNELLA**. **PRUNELLA** Ag. Selfheal. Hardy herbaceous perennials. Also called heal-all.
- Ophiobolus rostrupii** Ferd. and Winge. On leaves of *B. vulgaris* in the Aleutian Islands.
- Puccinia brunellarum-moliniae** Cruchet. Rust with yellow aecial stage on leaves of *B. grandiflora* and *B. vulgaris*, and brown uredinal and telial stages on *Molinia caerulea* in Europe.
- Ramularia brunellae** E. and E. f. *pyrenaica* Gz. Frag. On leaves of *B. hybrida* in Spain.
- Ramularia harioti** Sacc. More or less circular, grayish white spots on both leaf surfaces of *B. vulgaris* in France.
- Septoria greschikii** Bres. On *B. grandiflora* in Europe.
- Septoria tralliana** Sacc. and var. *italica* Ferr. On leaves of *B. vulgaris* in Yugoslavia, Italy, Russia, and Great Britain.
- BRUNFELSIA**. Ornamental tropical shrubs.
- Fomes lamaoensis** Murr. See *Hevea*.
- BRUNSVIGIA**. Summer or autumn flowering bulbs.
- Aecidium brunswigiae** P. Henn. Leaf rust on *Brunswigia* sp. in the Union of South Africa.
- Tilletia sydowii** Sacc. and Trott. Smut sori 0.5 inch long in place of ovaries, protected at first by ashen-colored membranes which rupture to expose powdery black spore masses. On *B.* (*Ammochloa subacaulis* in Algeria.
- BRYONIA**. **BRYONY**. Herbaceous perennial climbers.
- Ascochyta bryoniae** Bub. and Kab. Irregular, grayish-yellow or brown spots, often confluent, on leaves of *B. alba* in Bohemia.
- Ascochyta tirolensis** Bub. Small circular to oblong whitish spots with irregular surrounding zones, on leaves of *B. dioica* in Tyrol.
- Ramularia bryoniae** Fautr. and Roum. Large, irregular, brown spots on leaves of *B. dioica* in France.
- BRYOPHYLLUM**. Succulent foliage plants.
- Stilbella flavida** (Cke.) Lind. See *Coffea*.
- BUDDLEIA**. **BUTTERFLY BUSH**. Ornamental shrubs or trees cultivated for their showy flowers.
- Diplodia buddleiae** Pat. On leaves of *Buddleia* sp. in Ecuador.
- Phyllosticta auriculata** Kalch. and Cke. Circular pale-brown spots, with purple margins, on leaves of *B. auriculata* and *B. globosa* in the Union of South Africa and Denmark.
- Phyllosticta montemartini** Ciferri. Irregular, yellow-brown spots on leaves of *B. variabilis* (*B. davidi*) in Italy.
- Septoria buddleiae** Kalch. and Cke. Irregular, dark brown spots on leaves of *B. salvifolia* in the Union of South Africa.
- Septoria merrillii** Syd. On leaves of *B. asiatica* in the Philippines.
- Septoria thümenii** Sacc. Red-brown spots on leaves of *B. auriculata* in the Union of South Africa.
- BUGINVILLEA**. See *Bougainvillea*.
- BULBINE**. Bulbous and nonbulbous African and Australian plants.
- Aecidium bulbines** Henn. and Evans. Brown spots on both leaf surfaces in which yellow aecia appear. On *Bulbine* sp. in the Union of South Africa.
- Uromyces bulbinis** Thuem. Brown rust pustules are arranged concentrically in large circles on sunken spots on both leaf surfaces of *B. aloidis*, *B. bulbosa*, and *B. latifolia* in New South Wales and Victoria.
- Uromyces semibarbatæ** Osborn. Leaf rust on *B. semibarbatæ* in Australia.
- BULBOCODIUM**. Crocuslike bulbous plants.
- Aecidium bulbocodii** Koern. Yellow rust pustules on leaves of *B. vernum* in Asia.
- Urocystis colchici** (Schlecht.) Rab. See *Colchicum*.
- BULBOPHYLLUM**. See *Orchidaceae*.
- BUPHTHALMUM**. **ОХЕYE**. Perennial Asian herbs.
- Phyllosticta diphthalmi** Allesch. Irregular spots, often occupying entire leaf area, on *B. salicifolium* in Germany.
- Ramularia buphtalmi** Allesch. Subcircular or irregular, often confluent, dark-violet spots, drying out to brown, on leaves of *B. salicifolium* in Germany.
- Septoria buphtalmi** Allesch. Small, irregular, purple or olive-colored spots on leaves of *B. salicifolium* in Germany.
- BUPLEURUM**. Shrubs or herbs.
- Ascochyta bupleuri** Thuem. (Including var. *bupleurum-fruticosum* Gz. Frag. Gray spots on leaves of *B. falcatum*, *B. fruticosum*, and *B. obovatum* in Spain, Portugal, and Germany.
- Cercospora bupleuri** Pass. Reddish spots on stems and branches of *B. tenuissimum* in Italy.
- Didymella fruticosa** v. Hoeh. Circular or oblong purple-brown spots with black margins on leaves of *B. fruticosum* in Corsica.
- Entyloma bupleuri** Lindr. Smut sori in brown spots on leaves and stems of *B. glaucum* in France.
- Laestadia bupleuri** (Dur. and Mont.) Sacc. On leaves of *B. spinosum* in Algeria.

**BUPLEURUM**—Continued.

- Phyllosticta asteromoides** Bub. Irregular, confluent, purple-brown or black leaf spots on leaves of *B. falcatum* in Bohemia.
- Phyllosticta bupleuri** (Fckl.) Sacc. Gray spots on leaves of *B. falcatum*, *B. fruticosum*, and *B. gibraltarium* in Algeria and Germany.
- Puccinia bupleuri-falcata** (DC.) Wint. This rust in all its stages occurs on a wide range of species of *Bupleurum* in Japan, Indo-China, India, China, Siberia, Asia Minor, north Africa, and all of Europe.
- Septoria amphigena** Miy. On leaves of *B. falcatum* in China.
- Septoria bupleuri** Desm. Small subcircular whitish spots with black borders on leaves and buds of *B. frutescens*, *B. fruticosum*, and *B. semicompositum* in Spain, Portugal, France, Italy, and Austria.
- Septoria bupleuricola** Sacc. Circular, dark brown spots, becoming gray with dark-brown margins on leaves of *B. aureum* and *B. longifolium* in Siberia.
- Septoria bupleuri-falcata** Died. Subcircular or angular spots, white above gray below, with black margins, on leaves of *B. falcatum* in Germany.
- Septoria bupleurina** de Lamarl. Black, then brown, circular or angular spots with narrow greenish borders on leaves of *B. longifolium* in France.
- Septoria diffusa** F. Tassi. Diffuse dark-brown spots on leaves of *B. fruticosum* in Italy.
- Uromyces bupleuri** P. Magn. Chestnut-brown rust pustules on leaves and stems of *B. frutescens* and *B. graminifolium* in Persia and Spain.
- BURCHARDIA.** Herbaceous perennials, allied to *Veratrum*.
- Puccinia burchardiae** Sacc. Yellow-brown to dark-brown rust pustules on both leaf surfaces of *B. umbellata* in Australia.
- BUTOMUS.** FLOWERING RUSH. Hardy perennial aquatic plants.
- Doassansia niessli** De Toni. Smut sori in gray-brown spots on leaves of *B. umbellatus* in Poland, Austria, and Germany.
- Physoderma butomi** Schroet. On leaves of *B. umbellatus* in Finland, Denmark, Norway, and Germany.
- Ramularia butomi** Lind. Small, oblong or angular, scattered black spots on leaves of *B. umbellatus* in Sweden.
- BUXUS.** Box. Ornamental evergreen, small trees or shrubs.
- Ascochyta buxina** Sacc. Indefinite light-brown spots on leaves of *B. sempervirens* in France and Italy. Possibly *Phyllosticta buxina* Sacc., which occurs in the United States, is the same species.
- Dothidella depazeoides** (Desm.) Theiss. and Syd. Black stromata on white spots on leaves of *B. sempervirens* in France, Belgium, Switzerland, and Austria.
- Gloeosporium louisiae** Baeuml. Anthracnose on leaves of *B. sempervirens* in Belgium and Hungary.
- Guignardia buxi** Desm. Yellow spots on lower surfaces of leaves of *B. sempervirens* in Europe.
- Helminthosporium martinicense** Thuem. Black effuse spots on leaves of *B. citrifolia* in Martinique.
- Phyllosticta auerswaldii** Allesch. On leaves of *B. sempervirens* in Germany. Reported from New Jersey.
- Phyllosticta buxina** Sacc. Variable, pale-brown leaf spots with distinct dark purple borders on *B. sempervirens* in Great Britain and Italy.
- Phyllosticta phacidioides** (Sacc.) Allesch. On leaves of *B. sempervirens* in Yugoslavia.
- Puccinia buxi** DC. Brown rust pustules on leaves of *B. sempervirens* in Madeira, Persia, Dalmatia, Spain, Portugal, Greece, Russia, Italy, Switzerland, France, Great Britain, Austria, and Germany.
- Septoria phacidioides** Desm. On leaves of *B. sempervirens* in France and Belgium.
- Uromyces ambiens** Cke. Yellow-brown rust pustules on leaves of *B. sempervirens* in India.
- BYRSONIMA.** Tropical American trees and shrubs.
- Accidium byrsonimae** Kern. & Kellerm. Leaf and stem rust on *B. crassifolia* in Brazil and Guatemala.
- Accidium byrsonimaticola** P. Henn. Leaf rust on *Byrsonima* sp. in Brazil.
- Accidium byrsonimatis** P. Henn. Leaf and stem rust on *B. verbascifolia* and *Byrsonima* sp. in Brazil and Trinidad.
- Accidium singulare** (Diet and Holw.) Arth. Leaf rust on *B. crassifolia* in Guatemala.
- Cronartium byrsonimatis** P. Henn. Leaf rust on *B. coccolobifolia* in Brazil.
- Cronartium notatum** Arth. Brown leaf rust on *B. crassifolia* in Cuba and Porto Rico.
- Uredo byrsonimatis** P. Henn. Leaf rust on *Byrsonima* sp. in Brazil.
- Uredo uberabensis** P. Henn. Leaf rust on *Byrsonima* sp. in Peru and Brazil.
- CACALIA.** Perennial herbs.
- Coleosporium cacaliae** Otth. Golden-yellow rust pustules on needles of *Pinus montana*, *P. silvestris*, and on leaves of *Adenostyles albifrons*, *A. alliariae*, *A. alpina*, *Cacalia atriplicifolia*, *C. farfaraefolia*, *C. hastata*, *C. pubescens*, and *C. suaveolens* in Japan, Siberia, Russia, Spain, Bohemia, Montenegro, Denmark, Great Britain, France, Italy, Switzerland, Austria, Hungary, and Germany.
- Puccinia senecionicola** Arth. See Senecio.
- Puccinia tranzschelii** Diet. Dark-brown rust pustules on leaves of *C. hastata* in Russia.
- Rhytisma hysteroioides** Fr. On stems of *C. hastata* in Siberia.
- Uromyces cacaliae** (DC.) Wint. See *Adenostyles*.
- Uromyces veratri** (DC.) Schroet. See *Veratrum*.
- CACARA.** See *Pachyrhizus*.
- CAESALPINIA.** Ornamental tropical shrubs or trees grown for showy flowers and attractive leaves. Some species yield tanning material and dye stuffs.
- Helminthosporium bonducellae** P. Henn. Leaf spot of *C. bonducella* in Brazil.
- Laestadia caesalpiniae** Pat. Pale-brown spots on leaves of *Caesalpinia* sp. in Java.
- Ravenelia cohniiana** P. Henn. Cinnamon-brown rust pustules on leaves and stems of *Caesalpinia* sp. in Brazil.
- Ravenelia humphreyana** P. Henn. Cinnamon-brown to dark-brown rust pustules on leaves of *C. bahamensis* and *C. pulcherrima* in Porto Rico, Jamaica, Mexico, and Guatemala.
- Ravenelia inconspicua** Arth. Rust on leaves of *Caesalpinia* sp. in Mexico and *C. exostemma* in Guatemala.
- CAJANUS.** PIGEON PEA. Sometimes called Congo pea or gandul. Cultivated for the edible peas.
- Accidium cajani** Petch. Rust producing small galls on under surfaces of leaves and on petioles and stems of *C. indicus* in Ceylon. Probably should be referred to *Woroninella umbilicata*.
- Cercospora cajani** P. Henn. (*Velloziella cajani* Rangel.) Small circular to irregular dark-brown spots on leaves of *C. indicus* in Porto Rico, Cuba, and Brazil.
- Cercospora instabilis** Rangel. Small angular, then confluent, dark-brown spots on leaves, branches and pods of *C. indicus* in Brazil.
- Colletotrichum cajani** Rangel. Anthracnose on leaves of *C. indicus* in Brazil.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Diplodia cacaoicola** P. Henn. See *Theobroma*.

**CAJANUS**—Continued.

**Fusarium udum** Butl. This fungus produces a gradual, or at times, sudden wilting and dying of infected plants. Tissues at the base of the stems and along the main roots of diseased plants are blackened either entirely or in streaks. A serious disease in India on the pigeon pea (*C. indicus*). See also *Dianthus*.

**Heterosporium lagunense** Syd. On *C. indicus* in the Philippines.

**Phoma cajani** Rangel. On stems of *C. indicus* in Brazil.

**Phyllosticta cajani** Rangel. On leaves of *C. indicus* in Brazil.

**Rosellinia bunodes** B. and Br. See *Citrus*.

**Uredo cajani** Syd. Rust on leaves of *C. indicus* in India.

**Woroninella umbilicata** (B. and Br.) v. Hoeh. Minute, red swellings on the stems and leaves which burst, exposing orange-red powdery spore masses. On *C. indicus* in Ceylon.

**CALACINUM**. See *Muehlenbeckia*.**CALADIUM**. Large-leaved aroids grown for their variegated foliage. Herbaceous perennials.

**Cercospora caladii** Cke. Circular dull-brown leaf spots on *Caladium* sp. in India.

**Gloeosporium arecarum** P. Henn. Circular to irregular brown spots, with darker brown surrounding zones, on leaves of *Caladium* sp. and *Philodendron bipinnatifidum* in Japan and Brazil. Hemmi in Japan has described what is probably the same species under the same name.

**Helminthosporium caladii** Stevens. Large (up to 1.5 to 2 centimeters), circular or oval ashen white spots, with tan-colored borders, on leaves of *C. bicolor* in Porto Rico.

**Macrophoma surinamensis** (B. and C.) Berl. and Vogl. On leaf blades and petioles of *Caladium* spp. in Surinam.

**CALAMAGROSTIS**. Hay and forage grasses.

**Dilophia graminis** (Fekl.) Sacc. See *Triticum*.

**Leptosphaeria sparsa** (Fekl.) Sacc. See *Poa*.

**Metasphaeria avenae** (Auersw.) Sacc. See *Avena*.

**Mycosphaerella calamagrostidis** Volk. On leaves of *C. varia* in Switzerland.

**Mycosphaerella lineolata** (Desm.) Schroet. On leaves of *Ammophila* sp., *Calamagrostis epigejos*, and *Phragmites communis* in Great Britain, Denmark, France, Italy, and Germany.

**Mycosphaerella perforans** Desm. On leaves of *C. arenaria* in Denmark, France, and Holland.

**Mycosphaerella pusilla** (Auers.) Johans. On leaves of *C. arenaria* in Finland, Denmark, and Germany.

**Puccinia borealis** Juel. See *Agrostis*.

**Puccinia brevicornis** S. Ito. Powdery golden-yellow and black rust pustules on leaves of *C. canadensis* and *C. villosa* in Japan.

**Puccinia epigejos** Ito. As in the preceding species on *C. arundinacea* and *C. epigejos* in Japan.

**Puccinia erythropus** Diet. Brown to black rust pustules on purple leaf spots on *C. sciuroides* and *Miscanthus sinensis* in Japan.

**Puccinia ishikawai** S. Ito. Golden-yellow and black rust pustules, on brown sunken spots, on leaves of *C. epigejos* var. *densiflora* in Japan.

**Puccinia pertenuis** Ito. Rust on leaves of *C. arundinacea* in Japan.

**Puccinia pygmaea** Erikss. Golden-yellow and dark-brown rust pustules on leaves of *C. arundinacea*, *C. epigejos*, and *C. halleriana* in Sweden, Finland, and Germany.

**Puccinia rangiferina** Ito. Powdery black rust pustules on leaves and sheaths of *C. arundinacea*, *C. littorea*, and *Melica* sp. in Japan.

**Puccinia stichosora** Diet. Rust on leaves of *C. sciurois* in Japan.

**Septoria arenariae** Rostr. On leaves of *C. arenaria* in Denmark.

**Septoria calamagrostidis** (Lib.) Sacc. White linear spots on leaves of *C. arundinacea*, *C. epigeios* and *C. sylvatica* in Russia, Denmark, and France.

**Septoria epigeios** Theum. On leaves of *C. epigeios* in Russia and Denmark.

**Septoria nebulosa** Rostr. Indefinite areas on leaves and culms of *Calamagrostis* sp., *Poa* sp., and *Trisetum* sp. in Greenland.

**Septoria rhizodes** Bres. and Krieg. On leaves of *C. halleriana* in Germany.

**Tilletia calamagrostidis** Fekl. Long linear smut pustules on the leaves which break open exposing black spore masses. On *C. arundinacea*, *C. epigeios*, *C. halleriana*, and *C. lanceolata* in Germany.

**Uredo paulensis** P. Henn. Rust on leaves of *Calamagrostis* sp. in Brazil.

**CALAMUS**. See *Palmae*.**CALANDRINIA**. ROCK PURSLANE. Fleishy, trailing plants used in borders or rockeries.

**Heterosporium calandriniae** Mass. Irregular pale olive-colored spots on leaves and inflorescences of *C. potentilloides* in Argentina.

**Peronospora calandriniae** Speg. Downy mildew on leaves of *C. portulacoides* in Argentina.

**Ustilago calandrinicola** Speg. Forms dark violet powdery spore masses in inflorescences of *C. affinis* in Argentina.

**CALATHEA**. Tropical perennial foliage plants.

**Niptera calathea** P. Henn. On leaves of *Calathea* sp. in Peru.

**CALCEOLARIA**. **FAGELIA**. Ag. Showy-flowered herbs and shrubs.

**Aphelenchus olesistis** Ritz. Bos. See *Begonia*.

**Entyloma calceolariae** Lagerh. Smut producing sori in circular whitish spots on leaves of *Calceolaria* sp. in Ecuador.

**Peronospora indica** Syd. Downy mildew on leaves of *C. scabiosaefolia* in India.

**Phyllosticta calceolariae** Speg. Elliptical, definite whitish spots on leaves of *C. foliosa* in Argentina.

**CALENDULA**. POT MARGOLD. Flower garden annuals or perennials.

**Cercospora calendulae** Sacc. Subcircular, dull-brown leaf spots on *C. officinalis* in Italy and Portugal.

**Entyloma calendulae** (Oud.) De By. Smut sori in circular, greenish or white, finally brown, spots on leaves of *Arnica montana*, *Arnoseris minima*, *Aster alpinus*, *Bellidiastrum michelii*, *Calendula arvensis*, *C. officinalis*, *C. pluvialis*, *Cineraria papposa*, *Crepis biennis*, *Erigeron acer*, *Hieracium murorum*, *H. pilosella*, *H. schmidtii*, *H. silvaticum*, *H. stygium*, *H. sudeticum*, *H. tridentatum*, *H. villosum*, *H. vulgatum*, *Leontodon hispidus*, and *Parthenium hysterophorus* in Argentina, Algeria, and throughout Europe. American references to this species apparently are referable to *E. arnicale* E. and E. and other species.

**Puccinia calendulae** McAlp. Rust on leaves, stems, and fruits of *C. officinalis* in Australia.

**CALIMERIS**. Daisylike border plants.

**Puccinia calimeris** Syd. Dark-brown rust pustules on leaves of *C. altaica* in Siberia.

**CALLA**. Wild calla. Bog plants.

**Ascochyta pellucida** Bub. Large irregular spots on leaves of *C. palustris* in Bohemia.

**CALLIANDRA**. Evergreen shrubs and trees.

**Ravenelia affinis** Syd. Yellow-brown and deep-brown rust pustules on leaves of *C. turbinata* in Brazil.

**Ravenelia armata** Syd. Brown leaf rust on *Calliandra* sp. in Brazil.

**CALLIANDRA**—Continued.

**Ravenelia bizonata** Arth. and Holw. Powdery brown rust pustules on leaves of *C. houstoni* and *Calliandra* sp. in Guatemala.

**Ravenelia dieteliana** P. Henn. Brown rust pustules on leaves of *C. macrocephala* in Formosa and Brazil.

**Ravenelia echinata** Lagerh. and Diet. Cinnamon-brown to black rust pustules on leaves of *Calliandra* sp. in Ecuador.

**Ravenelia ectypa** Arth. and Holw. Brown leaf rust on *C. gracilis* and *Calliandra* sp. in Costa Rica and Guatemala.

**Ravenelia lagerheimiana** Diet. Brown rust pustules on leaves of *Calliandra* sp. in Ecuador.

**Ravenelia mexicana** Tranzsch. Brown rust pustules on leaves of *C. grandiflora* in Mexico.

**Ravenelia pазschkeana** Diet. Brown to black rust pustules on leaves of *Calliandra* sp. in Brazil.

**Uredo longipedis** P. Henn. Dull yellow-brown rust pustules on brown subcircular leaf spots on *C. longipes* in Brazil.

**CALLICARPA**. BEAUTY BERRY. Ornamental woody plants cultivated chiefly for their bright-colored fruit.

**Uromyces callicarpae** (Petch.) Fuji. Brown rust pustules on leaves of *C. formosana*, *C. lanata*, and *C. tomentosa* in Ceylon and Japan.

**CALLISTEMMA**. See Callistephus.**CALLISTEMON**. BOTTLE BRUSH. Hardy ornamental shrubs.

**Phyllosticta flavidula** Sacc. On leaves of *Callistemon* sp. in France.

**Septoria callistemonis** F. Tassi. On leaves of *Callistemon* sp. in Italy.

**CALLISTEPHUS**. **CALLISTEMMA** Ag. CHINA ASTER. Garden annuals.

**Coleosporium asterum** (Diet.) Syd. See Aster.

**Phyllosticta asteris** Bres. Subcircular to angular brown spots on leaves of *C. chinensis* in Europe.

**Phytophthora** sp. The "black neck" or wilt disease of China asters in Great Britain. Plants of all ages wilt and die, the lower parts of the stems becoming brown or black. The causal organism is said to be close to *P. omnivora*, but is considered distinct.

**CALLUNA**. HEATHER. Low evergreen shrubs.

**Metasphaeria callunae** Fautr. On branches of *C. vulgaris* in France.

**Plowrightia polyspora** (Bref.) Sacc. Small subcircular stromata on branches of *C. vulgaris* in Germany.

**CALONYCTION**. MOONFLOWER. Twining perennial herbs.

**Septocylindrium platense** Speg. Indefinite brown leaf spots on *C. aculeatum* (*C. speciosum*) in Argentina.

**CALOTROPIS**. MODAR-BUSH. Milkweedlike shrubs or small trees.

**Ascochyta tripolitana** Sacc. and Trott. Circular, definite, whitish spots with reddish margins, on leaves of *C. procera* in north Africa.

**Cercospora calotropidis** Speg. Indefinite yellow leaf spots on *C. procera* in Senegal. This species should be renamed, the name being occupied by the following species.

**Cercospora calotropidis** E. and E. Large dirty-brown spots on leaves of *C. procera* in the Bahamas and Central America.

**Cercospora inconspicua** Pat. and Har. Circular spots on leaves of *Calotropis* sp. in French North Africa.

**Cercospora patouillardii** Sacc. and D. Sacc. On leaves of *C. procera* in Guadalupe.

**Napcladium calotropidis** Mors. Circular, often confluent, yellow-green to olive-green spots on leaves and stems of *C. procera* in Tanganyika.

**CALPURNIA**. Tropical shrubs and trees.

**Ascochyta calpurniae** Wint. Angular to irregular, white or dirty-gray spots, with dull purple margins, on leaves of *C. sylvatica* in the Union of South Africa.

**Cercospora calpurniae** Petch. Diffuse yellow areas on leaves of *C. aurea* in Ceylon.

**Ravenelia glabra** Kalchbr. and Cke. Brown rust pustules on leaves of *C. sylvatica* in the Union of South Africa.

**CALTHA**. MARSH MARIGOLD. Hardy, blooming marsh plants.

**Cercospora calthae** Cke. Circular dull-brown leaf spots on *Caltha* sp. in Great Britain and Esthonia.

**Physoderma bohemicum** Sacc. Small brown galls on leaf blades and petioles of *C. palustris* in Bohemia.

**Puccinia zopfii** Wint. Chestnut-brown to dark-brown rust pustules on leaves of *C. leptosepala* and *C. palustris* in Argentina, Japan, Great Britain, France, Denmark, Belgium, Holland, Russia, Switzerland, Italy, Austria, and Germany.

**Ramularia calthae** Lindr. (*R. calthae* Gz. Frag.) Irregular whitish spots, with brown margins on leaves of *C. palustris* in Spain, Bohemia, Denmark, Finland, Austria, and Germany.

**CALYCANTHUS**. SWEET SHRUB. Ornamental shrubs with fragrant flowers.

**Ascochyta calycanthi** Sacc. and Speg. and var. **occidentalis** Fl. Tass. Leaf spots on *C. floridus* and *C. occidentalis* in Italy.

**Phyllosticta calycanthi** Sacc. and Speg. Brown spots, becoming white, on leaves of *C. orientalis* and *Meratia praecox* (*C. praecox*) in Denmark and Italy.

**Septoria calycanthi** Sacc. and Speg. Dull-yellow to brown spots on leaves of *C. floridus* and *C. laevigatus* in Italy and Portugal.

**CAMELLIA**. Woody plants grown for their showy white or red flowers and handsome evergreen leaves.

**Ascochyta minutissima** Pass. On young branches of *C. japonica* in Italy.

**Ascochyella camelliae** (Pass.) Fl. Tass. Oblong or irregular, grayish-white leaf spots on *C. japonica* in France.

**Exobasidium assamense** Syd. and Butl. Circular, then confluent and irregular, white areas on leaves of *C. drupifera* in India.

**Exobasidium camelliae** Shirai. See Thea.

**Helminthosporium polyphragmium** Syd. On stems of *C. japonica* in India.

**Leptosphaeria camelliae** Cke. and Mass. Brown indefinite spots on leaves of *Camellia* sp. in Australia.

**Phyllosticta amphigena** D'Alm. Brown leaf spots on *C. japonica* in Portugal.

**Phyllosticta camelliae** West. and var. **meranensis** Bub. On leaves of *C. japonica* in Denmark, France, and Austria, and on *Thea* (tea) in Japan.

**CAMPANULA**. BELLFLOWER. HAREBELL. CANTERBURY BELLS. Hardy herbaceous perennials.

**Ascochyta bohémica** Kab. and Bub. Circular to angular, often confluent, small yellowish or brown spots with purple margins on leaves of *C. trachelium* in Bohemia.

**Ascochyta carpathica** (Allesch.) Keissl. Small oblong or irregular whitish spots with brown margins on leaves of *C. carpathica* in Germany.

**Cliathrosorus campanulae** Ferd. and Winge. Small swellings on roots of *C. rapunculoides* in Denmark.

**CAMPANULA**—Continued.

**Marsonia campanulae** Bres. and Allesch. Small subcircular dull-yellow leaf spots on *C. latifolia* and *C. morettiana* throughout Europe.

**Peronospora corallae** Tranzsch. Downy mildew attacking the flowers of *C. persicifolia* in Russia and Sweden.

**Phyllosticta alliariaefolia** Allesch. Large subcircular to angular ashen spots, with indefinite margins, on leaves of *C. alliariaefolia* in Germany.

**Phyllosticta campanulae** Sacc. and Speg. Yellowish-brown spots with darker margins on leaves of *C. glomeratum* and *C. trachelium* in Italy and Siberia.

**Phyllosticta campanulina** Moesz. Whitish spots with red-brown margins on leaves of *C. persicifolia* in Hungary.

**Phyllosticta michauxioides** P. Magn. Large leaf spots, yellow above, black beneath, on *C. michauxioides* in Asia Minor.

**Phyllosticta rehmi** Bub. On leaves of *C. trachelium* in Hungary.

**Placosphaeria campanulae** (DC.) Bäuml. Black stromata on leaves of *C. bononiensis*, *C. rapunculoides*, and *C. trachelium* in France, Yugoslavia, Switzerland, Austria, and Germany.

**Pseudopeziza radians** Sacc. On leaves of *C. patula*, *C. rapunculoides*, and *C. trachelium* in Great Britain, Finland, France, Belgium, and Denmark.

**Puccinia heideri** Wettst. Black rust pustules on leaves of *C. barbata* in Austria.

**Ramularia campanulae-barbatae** Jaap. and Lind. Subcircular brown spots on leaves of *C. barbata* in Switzerland.

**Ramularia campanulae-rotundifoliae** Lindr. Small gray or reddish spots on leaves of *C. rotundifolia* in Finland.

**Ramularia coleosporii** Sacc. and var. **campanulae** Car. On leaves of *C. rapunculoides*, *C. trachelium*, *Campanula* sp., *Hypericum androsaenum*, *Melampyrum* sp., *Senecio* sp., and *Telekia speciosa* in Europe.

**Ramularia macrospora** Fres. and var. **major** Lindr. Subcircular or irregular greenish-gray spots with ochraceous margins on leaves of *C. glomerata*, *C. latifolia*, *C. media*, *C. persicifolia*, *C. pyramidalis*, *C. rapunculoides*, and *C. trachelium* in Alaska and Europe.

**Ramularia michauxioides** P. Magn. On leaves of *C. michauxioides* in Asia Minor.

**Septoria campanulae** (Lév.) Sacc. On leaves of *Campanula* sp. in Persia.

**Septoria obscura** Trail. Dull brown circular or irregular leaf spots on *C. rotundifolia* in Scotland.

**Septoria trachelii** Allesch. Irregular spots on leaves of *C. trachelium* in Switzerland and Russia.

**CAMPANUMAEAE**. Twining perennial herbs.

**Coleosporium campanumaeae** Diet. Yellow rust pustules on leaves of *C. javanica* in Japan.

**Puccinia campanumaeae** Pat. Yellow-brown rust pustules on lower surfaces of leaves of *C. javanica* and *Campanumaeae* sp. in China and Java.

**CAMPE**. See Barbarea.**CANAVALIA**. JACK BEAN. SWORD BEAN. Prostrate trailing herbs used for green manure crops.

**Cercospora canavaliae** Syd. Dull-red circular to irregular areas on leaves of *C. ensiformis* and *C. gladiata* in the Philippines.

**Crotelium canavaliae** Arth. Brown rust pustules on red-brown leaf spots on *C. ensiformis*, *C. gladiata*, and *Canavalia* sp. in Porto Rico and French Congo.

**Elsinoe canavaliae** Racib. This fungus causes distortion of the leaves and stems of *C. ensiformis* and *C. gladiata* in Ceylon and the Philippines.

**Gloeosporium canavaliae** Syd. Anthracnose on stems of *C. ensiformis*, *C. gladiata* and *C. turgida* in Hawaii and the Philippines.

**Mycosphaerella canavaliae** Syd. Indefinite brown leaf spots on *C. gladiata* in the Philippines.

**Physalospora guignardioides** Sacc. On leaves of *C. ensiformis* and *C. gladiata* in the Philippines.

**Septoria canavaliae** Lyon. Small circular to irregular spots on leaves causing them to drop prematurely. The spots are dark brown to black, becoming confluent, with ashy-gray centers on *C. ensiformis* in Hawaii.

**Septoria molleriana** Bres. Circular pale-brown spots with reddish margins on leaves of *C. lineata* and *C. obtusifolia* in the Philippines and island of St. Thomas (Africa).

**CANDOLLEA**. See Stylidium.**CANNA**. Tall ornamental plants cultivated as bedding plants for the foliage and showy flowers. One species with edible root stocks.

**Ascochyta cannae** Rangel. On leaves of *Canna* sp. (cult.) in Brazil.

**Bunchy-top**. See *Musa*.

**Puccinia cannae** (Wint.) P. Henn. Yellow to black rust pustules covering the under sides of leaves of *Maranta arundinacea*, *Thalia geniculata*, *Canna coccinea*, *C. glauca*, *C. indica*, and other cultivated species and varieties throughout Central America, the West Indies and South America. Occurs also to a limited extent in south Florida. This disease destroys the ornamental value of the canna to such an extent that its culture has been abandoned in some countries where the rust is prevalent. A bacterial stem disease of *Canna* has been reported from Ceylon.

**CANNABIS**. HEMP. Fiber plants and occasionally used as ornamentals.

**Cercospora cannabina** Wakef. Circular to elongate, often confluent, pale-brown leaf spots on *C. sativa* in Uganda.

**Dendrophoma marconii** Cav. On *C. sativa* in Italy.

**Leptosphaeria cannabina** Ferr. and Mass. On leaves of *C. sativa* in Italy.

**Peronospora cannabina** Oth. Downy mildew causing yellowing of leaves of *C. indica* and *C. sativa* in Japan, Switzerland, Russia, and Italy.

**Phyllachora cannabis** P. Henn. On *C. sativa* in Brazil. Referred to the Clypeosphaeriaceae by Theissen and Sydow.

**Phyllosticta cannabis** (Kirch.) Speg. Subcircular dark-brown spots on leaves of *C. indica* and *C. sativa* in Yugoslavia, Bohemia, Italy, and Russia.

**Rhizoctonia napi** West. See Brassica.

**Septoria cannabis** Sacc. Indefinite dirty-yellow spots on leaves of *C. sativa* in Russia, Italy, Belgium, and Germany. Probably the imperfect stage of *Leptosphaeria cannabina* Ferr. and Mass.

**Uredo kriegeriana** Syd. Brown rust pustules on leaves of *C. sativa* in Germany.

**CAPNOIDES**. See Corydalis.**CAPRIOLA**. See Cynodon.**CAPSICUM**. RED PEPPER. Sometimes called Cayenne pepper. Herbs or shrubs cultivated for their fruit.

**Acrothecium capsici** Turc. Hazel-colored circular to oval depressed areas with brown margins on fruit of *C. annuum* in Italy.

**Bacillus capsici** Pavar. and Tarc. Bacterial fruit rot and wilt of plants of *C. annuum* in Italy.

**Phomopsis capsici** (Magn.) Sacc. On harvested fruit of *C. annuum* in Italy and the Philippines.

**Phyllosticta capsici** Speg. Circular grayish leaf spots with definite deep-brown margins on *C. annuum* and *Capsicum* sp. in Argentina.

**CAPSICUM**—Continued.

**Puccinia capsici** Mayor. Rust on leaves of *C. baccatum* in Colombia. This species has priority over the following, which is probably not distinct.

**Puccinia capsici** Averna-Sacca. Rust pustules on leaves of *C. frutescens*, *C. microcarpum*, *C. odoriferum*, and *C. pendulum* in Brazil.

**Puccinia gonzalezi** Mayor. Rust on leaves of *Capsicum* sp. in Colombia.

**Puccinia paulensis** Rangel. (*P. capsici* Averna?). Rust on leaves and twigs of *C. annuum* in Brazil.

**Rhizoctonia palida** Matz. See Saccharum.

**Rosellinia arcuata** Petch. See Thea.

**Vermicularia capsici** Syd. (*Steirochaete capsici* [Syd.] Sacc.). This fungus causes the most serious disease known of pepper in India, Ceylon, China, and the Philippines. Butler describes the disease as follows: "At first the flowers droop and dry up, the disease then spreading, if weather conditions are favorable, through the flower stalks to the stem. The bark turns brown and finally white. The upper portion of the plant may be killed. Round, sunken, anthracnose spots occur on the fruit, dark at first then pale at the center and coalescing. Diseased fruits shrivel and dry up." Known hosts are *Capsicum annuum*, *C. frutescens*, *Dolichos biflorus*, *D. lablab*, *Lycopersicon esculentum*, *Solanum melongena*, and *Vigna catjang*.

**CARAGANA**. PEA SHRUB. Siberian pea tree. Ornamental shrubs grown chiefly for their bright-yellow flowers, also as hedge plants.

**Ascochyta bondarzewii** P. Henn. Dull-brown spots on leaves of *C. arborescens* in Russia.

**Hendersonia septem-septata** Vesterg. On leaves of *C. arborescens* in Russia.

**Melasmia caraganae** Thuem. Black areas on leaves and branches of *C. arborescens* in Siberia.

**Mycosphaerella jaczewskii** A. Pot. (*Phyllosticta borszczowii* Thuem.). On leaves of *C. arborescens* in Siberia, Russia, and Germany.

**Peronospora lagerheimii** Gäum. Downy mildew on leaves of *C. arborescens* in Norway.

**Phleospora caraganae** Jacz. Indefinite yellow spots on leaves of *C. arborescens* in Russia.

**Phyllachora crebia** Syd. Black stromata on rust-colored spots on leaves of *Caragana* sp. in India.

**Phyllosticta caraganae** Syd. Whitish spots with brown margins, often at the top or along the margins of leaves of *C. chamlaga* in Germany.

**Phyllosticta caraganae** Fl. Tass. On leaves of *C. arborescens* in France. This species should be renamed, the name being occupied as above.

**Phyllosticta gallarum** Thuem. On leaves of *C. arborescens* in Siberia.

**Phyllosticta spaethiana** All. and Syd. Small irregular white spots with dull-brown margins on leaves of *C. arborescens* in Russia, Austria, and Italy.

**Septoria caraganae** P. Henn. Yellow then brown leaf spots on *C. arborescens* and *Caragana* sp. in Russia and Germany.

**Uromyces genistae-tinctoriae** (Pers.) Wint. Powdery cinnamon-brown rust pustules on under leaf surfaces of *C. arborescens*, *C. chamlaga*, *C. frutescens*, *C. fruticosa*, *C. pygmaea*, *Colutea arborescens*, *C. orientalis*, *Cytisus alpinus*, *C. austriacus*, *C. biflorus*, *C. capitatus*, *C. decumbens*, *C. falcatus*, *C. heuffeli*, *C. hirsutus*, *C. laburnius*, *C. liberus*, *C. linifolius*, *C. nigricans*, *C. proliferus*, *C. ramentaceus*, *C. ratisbonensis*, *C. sagittalis*, *C. sessilifolius*, *C. supinus*, *Galega officinalis*, *G. orientalis*, *Genista angelica*, *G. germanica*, *G. pilosa*, *G. sagittalis*, *G. sericea*, *G. tinctoria*, *Laburnum anagyroides*, *Sarothamnus scoparius*, and *Tetragolobus siliquosus* in Argentina, Japan, Siberia, Canary Islands, and Europe.

**CARDAMINE**. BITTER CRESS. CUCKOO-FLOWER. Small leafy-stemmed perennials.

**Phyllosticta cardamines** Allesch. On leaf blades and petioles of *C. bellidifolia* in Greenland.

**Puccinia cardamines-bellidifoliae** Diet. Chestnut-brown rust pustules on leaf blades and petioles of *C. bellidifolia* in Norway and Greenland.

**Puccinia cardamines-cordatae** Diet. and Neg. Yellow-brown rust pustules on leaf blades and petioles of *C. cordata* and *C. reniformis* in Chile.

**Puccinia cruciferarum** Rud. Red-brown rust pustules on leaves of *C. alpina*, *C. asarifolia*, *C. bellidifolia*, *C. gelida*, *C. pratensis*, and *C. resedifolia* in Iceland, Spain, Switzerland, France, Belgium, Austria, and Hungary.

**Ramularia cardamines** Syd. Circular to irregular spots on leaves of *C. amara* and *C. pratensis* in Switzerland, Denmark, Bohemia, and Germany.

**Septoria cardamines** Fckl. Dark-brown effuse spots on leaves and stems of *C. amara* and *C. pratensis* in Germany.

**Septoria cardamines-resedifoliae** Heim. Small brown leaf spots, becoming whitish, on *C. resedifolia* in Tyrol.

**Septoria cardamines-trifoliae** v. Hoehn. Subcircular whitish spots with dull brown margins on both leaf surfaces of *C. trifolia* in Austria.

**CARDIOSPERMUM**. HEARTSEED. BALLOON VINE. Tendril climbing tropical herbs.

**Aecidium cardiospermi** Cke. Yellow rust pustules on yellow-brown leaf spots on *C. halicacabum* (*C. microcarpum*) in Zanzibar.

**Cercospora cardiospermi** Petch. On leaves of *C. halicacabum* in Ceylon.

**Phyllosticta cardiospermi** Speg. Circular spots on leaves of *C. velutinum* in Argentina.

**CARICA**. PAPAYA. Small rapid-growing unbranched trees of the Tropics with milky juice. One species with edible fruit.

**Ascochyta caricae** Pat. Dark linear or ovate spots on petioles becoming ashen on *C. papaya* in Ecuador.

**Asterina caricarum** Rehm. On leaves of *C. papaya* in Brazil.

**Cercospora caricae** Speg. Small, pale, indefinite spots on leaves of *C. papaya* in Brazil.

**Diplodia cacaoicola** P. Henn. See Theobroma.

**Helminthosporium papayae** Syd. Definite circular ochraceous concentrically zoned leaf spots on *C. papaya* in the Philippines.

**Mycosphaerella caricae** Syd. On leaves of *C. papaya* in the Philippines and Ceylon.

**Ovulariopsis papayae** Van der Bijl. Powdery mildew on lower leaf surfaces of *C. papaya* in the Union of South Africa. A powdery mildew, which may be the same, occurs in Florida and other countries where the papaya is grown.

**Phyllosticta caricae-papayae** Allesch. On leaves of *C. papaya* in Brazil and the Union of South Africa.

**Phyllosticta papayae** Sacc. Small subcircular whitish spots on fruit of *C. papaya* in Abyssinia.

**Phytophthora faberi** Maubl. See Theobroma.

**Sphaerostilbe repens** B. and Br. See Hevea.

**CARISSA**. Tropical spinose shrubs cultivated for ornament and hedges as well as for the edible berrylike fruit.

**Leveillina arduinae** (K. and C.) Theiss. and Syd. Black irregular stromata on upper leaf surfaces of *C. arduina* in the Union of South Africa.

**Phyllosticta carissae** Kalchb. and Cke. Subcircular pale-brown leaf spots with brown margins on *C. arduina* in south and central Africa.

**Puccinia carissae** Cke. and Mass. Dark-brown rust pustules on leaves of *C. ovata* in Australia.

- CARLUDOVICA.** HAT PALM. Palmlike plants of tropical America, used in making Panama hats.  
*Microthyrium carludovicæ* P. Henn. Dark-brown effuse spots on leaves of *Carludovica* sp. in Brazil.
- CARPINUS.** HORNBEAN. Shade and timber trees.  
*Asechyta carpinea* Sacc. Subcircular or sinuous dull-yellow leaf spots on *C. betulus* in Italy.  
*Exoascus carpini* Rostr. Produces yellowing and distortion of leaves and witches'-brooms of *C. caroliniana* and *C. betulus* in Europe.  
*Gloeosporium carpini* (Lib.) Desm. Irregular indefinite olive-brown anthracnose spots on leaves of *C. betulus* in Italy, France, Austria, and Germany.  
*Melampsorium carpini* (Fckl.) Diet. Yellow to yellow-brown rust pustules on lower leaf surfaces of *C. betulus*, *C. cordifolius*, and *C. yedoensis* in Japan, Russia, Italy, Switzerland, Austria, Hungary, and Germany.  
*Phyllosticta carpinea* Sacc. Subcircular or sinuous dull-yellow leaf spots with brown margins on *C. betulus* and *C. orientalis* (*C. duinensis*) in Italy.  
*Phyllosticta carpini* Schulz. and Sacc. On leaves of *C. betulus* in Russia.  
*Sclerotium carpini* West. Round black sclerotia on living leaves of *Carpinus* sp. in Belgium.  
*Uncinula geniculata* Ger. var. *carpinicola* K. Hara. Powdery mildew on leaves of *Carpinus* sp. in Japan.
- CARTHAMUS.** SAFFLOWER. Hardy annuals.  
*Cercospora carthami* Syd. On leaves of *C. tinctorius* in Russia.  
*Cercosporina carthami* Syd. Foliage frequently entirely destroyed and plants killed by this leaf disease of *C. tinctorius* in the Philippines. The spots are circular to irregular and gray-brown in color.  
*Erysiphe taurica* Lév. See *Althaea*.  
*Gloeosporium carthami* Hori and Hemmi. Brown elliptic to irregular sunken areas on leaves and stems, in which pink spore masses appear. The tips of diseased plants wilt and fall over. On *C. tinctorius* in Japan.  
*Puccinia carthami* (Hutzelm.) Cda. Brown to dark-brown rust pustules on leaves of *C. oxyacanthus* and *C. tinctorius* in Japan, India, Egypt, Austria, and Germany.
- CARUM.** CARAWAY. Annual or perennial herbs, some yielding aromatic and edible garden products.  
*Cercospora cari* Westerd. and Luyk. Sunken brown spots with dark-brown margins occur on petioles and flower stalks of *C. carvi* in Holland.  
*Protomyces macrosporum* Ung. See *Coriandrum*.  
*Puccinia dictyospora* Tranzsch. The yellow aecia of this rust form elongate swellings along the veins of the leaves with brown telial pustules following on *Carum* sp. and *Scaligeria hirtula* in Turkestan. Possibly the same as the following species.  
*Puccinia microsphineta* Lindr. Black rust pustules on leaves and stems of *C. atosanguineum* in Turkestan.  
*Urophlyctis kriegeriana* P. Magn. Small pearl-like galls formed on the surface of leaves, stems, and floral parts. These galls are depressed at the apex and hollow. On *C. carvi* and *C. persicum* in Switzerland and Germany.
- CARYOPHYLLUS.** EUGENIA. Malabar plum. Rose apple. Tropical ornamental and fruit trees.  
*Puccinia jambosae* P. Henn. Brown rust pustules on leaves of *C. jambos* in Brazil.  
*Puccinia jambulana* Rangel. Brown rust pustules on leaves and fruit of *C. jambos* in Brazil.  
*Puccinia psidii* Wint. See *Psidium*.  
*Rosellinia bunodes* B. and Br. See *Citrus*.  
*Uredo flavidula* Wint. Rust on leaves of *C. jambos* in Brazil.
- CARYOTA.** See *Palmae*.
- CASIMIROA.** WHITE SAPOTE. Tropical evergreen trees. One species cultivated for its fruit.  
*Bacterium citri* Hasse. See *Citrus*.  
*Cercospora coleroides* Sacc. Subcircular then confluent gray leaf spots with dark-brown margins on *C. edulis* in Mexico.
- CASTANEA.** CHESTNUT. Nut and timber trees.  
*Actinopelte japonica* Sacc. On leaves of *C. vesca* in Japan. Reported from New Jersey.  
*Bacterium castanicolum* Carvara. Said to cause galls at the base of stems of seedlings of *Castanea* sp. in Italy and to be distinct from *B. tumefaciens*.  
*Blepharospora cambivora* Petri. See *Ink disease*.  
*Cylindrosporium castanicolum* (Desm.) Berl. Dark-brown indefinite spots on both leaf surfaces of *C. vesca* (*C. sativa*) in Europe.  
*Diplodina castaneae* Prill. and Delacr. Forms cankerous swellings on branches of *C. vesca* in France and Italy.  
*Endothia parasitica* (Murr.) Ander. and Ander. This serious disease, native of Japan and China, has been introduced into the United States and Belgium. The chestnut forests of the eastern United States have been nearly exterminated, entailing a loss of many millions of dollars by this fungus, which in its native home is a comparatively harmless wound parasite. *C. mollissima*, the Chinese chestnut, is very resistant. *C. dentata*, the American chestnut, and other species of chestnut and chinquapin, are extremely susceptible. The disease appears as sunken cankers on the bark of trunks and limbs, which increase rapidly in size until girdling is effected and the limb, or the entire tree, dies. The wood is not attacked, the fungus working in the inner bark, where its presence can be noted by the characteristic fan-shaped mycelium. Diseased areas soon become covered with yellow, orange, or reddish-brown fruiting pustules. In damp weather spores are extruded in long irregular yellow or greenish horns.  
*Fomes korthalsii* (Lév.) Cke. Causes heart and butt rot of *C. argentea* and *C. javanica* in Java.  
*Guignardia diffusa* (Crié.) Sacc. and Trott. On leaves of *Quercus* sp. and *Castanea* sp. in France.  
*Lanomyces tjibodensis* Güm. On leaves of *C. argentea* in Java.  
*Melanconis modonia* Tul. (*Coryneum perniciosum* Br. and Farn.) On dead branches of *C. vesca* in France, Great Britain, Italy, and Germany, but reported as "a very destructive" parasite in the Caucasus. Associated with the so-called ink disease, (q. v.).  
*Monochaetia pachyspora* Bub. Large circular brown leaf spots on *Castanea* sp. in Bohemia.  
*Mycosphaerella punctiformis* (Pers.) Starb. (*Phyllosticta punctiformis* Sacc.) Irregular, dull, yellowish-red leaf spots on *C. vesca* in Italy and Dalmatia.  
*Phoma endogena* Sacc. Causes a hard rot of the kernels of the nuts of *C. vesca* in Italy.  
*Phyllosticta nubecula* Pass. On leaves of *C. vesca* in Italy.  
*Plowrightia noxia* (Ruhl.) Sacc. On *Castanea* sp., *Fagus* sp., and *Quercus* sp. in Germany. Said to be not a *Plowrightia*, and its systematic position uncertain.  
*Pucciniastrum castaneae* Diet. Yellow rust pustules on leaves of *Castanopsis javanica* and *Castanea vesca* in Japan, India, and the Philippines.  
*Sclerotinia candolleana* (Lév.) Fckl. See *Quercus*.  
*Scolecodothis castaneae* Reeb. Brown-black stromata on the upper sides of leaves, with corresponding yellowish-red spots beneath, on *C. argentea* in Java.  
*Septoria castaneae* Lév. On leaves of *C. vesca* in France and Austria.  
*Tubercularia confluens* Pers. On leaves of *C. vesca* (*C. vulgaris*) in Spain.

**CASTANEA**—Continued.

**Uredo castaneae** P. Henn. Brown rust pustules on leaves of *C. vesca* in the Philippines.

**Black spot.** A disease of unknown origin attacking the fruit of *Castanea vesca* in Italy. The nuts are normal in appearance until cut open, when the kernel is found dark brown or black in color.

**Ink disease.** The so-called ink disease of European chestnut (*C. vesca*) has caused heavy losses in France, Italy, Portugal, and Spain. A number of causes, including several fungi and various environmental influences, have been assigned by various workers. Diseased trees wilt and die, and upon close examination the roots are found dead and rotted, the bark at the base of the trunk being similarly affected. There is a black liquid exudate, rich in tannin, from the roots and trunk which gives the characteristic name to the disease. The disease spreads in irregular circles from the original infection points, trees dying in from 2 to 3 years after being attacked. Some workers have assigned *Melanconis modonia* Tul. (or its imperfect stage *Coryneum perniciosum* Br. and Farn.) as the cause through destruction of the bark. More recently a phycomycete, *Blepharospora cambivora* Petri, has been considered as the causative agent. This fungus is described as bringing about the formation of dark cankerous bands in the bark in the region of the root collar and out along the larger roots. This type of disease would be similar to the collar rot of *Citrus* (*Phytophthora terrestris* Sherb.).

**A root rot,** "gangrene humide," said to be due to bacteria, is reported from Portugal.

**CASTANOPSIS.** Chinquapin. Ornamental trees or shrubs.

**Pucciniastrum castaneae** Diet. See Castanea.

**CASTILLA (CASTILLOA).** MEXICAN RUBBER TREE.

**Corticium salmonicolor** B. and Br. See Citrus.

**Diplodia cacaoicola** P. Henn. See Theobroma.

**Fomes lamaoensis** Murr. See Hevea.

**Rosellinia bunodes** B. and Br. See Citrus.

**Uredo artocarpi** B. and Br. See Artocarpus.

**CASTILLEJA.** PAINTED CUP. Herbs with showy bracts in terminal heads or spikes.

**Puccinia nesodes** Arth. and Holw. Brown rust pustules on leaves of *C. communis*, *C. tenuiflora*, and *Lamourouxia* sp. in Guatemala and Costa Rica.

**CASUARINA.** BEEFWOOD. Ornamental trees and shrubs with equisetum like leaves.

**Botryosporium diffusum** Cda. On branches of *C. leptoclada* in Russia.

**Corticium salmonicolor** B. & Br. See Citrus.

**Rhizoctonia lamellifera** Small. See Grevillea.

**Sphaerostilbe repens** B. & Br. See Hevea.

**Trichosporium vesiculosum** Butl. Causes a root disease of *C. equisetifolia* in India and possibly in Mauritius.

**CATALPA.** Ornamental trees cultivated for their flowers and foliage.

**Ascochyta catalpae** F. Tassi. Circular ashen-colored leaf spots with dull-brown margins on *C. speciosa* in Italy.

**Phyllosticta bacillispora** Kab. and Bub. Circular to angular brown spots becoming gray and often increasing so as to occupy entire leaf area on *C. syringifolia* in Bohemia.

**Septoria catalpae** Sacc. On pods of *C. bignonioides* and *C. syringifolia* in Italy.

**CATASETUM.** Epiphytic terrestrial orchids. See Orchidaceae.**CATTLEYA.** Epiphytic orchids. See Orchidaceae.**CEDRELA.** CIGAR-BOX CEDAR. Ornamental trees grown for their handsome foliage and as timber trees.

**Cercospora** sp. On leaves of *C. odorata* in Cuba.

**Fomes lamaoensis** Murr. See Hevea.

**Phyllachora balansae** Speg. Small black shiny stromata on leaf spots 0.5 to 2 centimeters in diameter on *C. brasiliensis*, *C. fissilis* and other species in Brazil, Argentina, and Paraguay.

**Phyllosticta agnostoica** Speg. Leaf spot of *Cedrela* sp. in Brazil.

**Triphragmium cedrelae** Hori. Yellow to black powdery pustules on both leaf surfaces of *C. chinensis* and *C. sinensis* in Japan.

**Uncinula delavayi** Pat. Powdery mildew on under leaf surfaces of *Ailanthus* sp. and *Cedrela sinensis* in China and Japan.

**Uromyces cedrelae** P. Henn. Dull-brown rust pustules on lower leaf surfaces of *C. serrata* in Java.

**CEDRUS.** CEDAR. CEDAR OF LEBANON. Valuable timber trees. Also grown for their handsome foliage and striking habit.

**Peridermium cedri** Barcl. A rust producing witches'-brooms with golden-yellow pustules on the needles of *C. deodara* in Northern India.

**CEIBA.** A fiber of great economic importance (silk cotton or kapok), is obtained from the pods. Also grown as shade trees.

**Cercospora** sp. A leaf spot on *C. pentandra* in Cuba.

**CELASTRUS.** BITTERSWEET. Woody climbers grown for their brightly colored fruit and handsome foliage.

**Pucciniastrum celastri** Syd. Brown rust pustules on leaves of *C. paniculatus* in India.

**Uncinula sengokui** Salmon. Powdery mildew on leaves of *C. articulatus* in Japan.

**CELERI.** See Apium.**CELOSIA.** COCKSCOMB. Garden annuals grown for their showy agglomerated flower heads, and sometimes colored foliage.

**Cercospora gilbertii** Speg. Circular gray leaf spots on *C. trigyna*, *Iresine celosioides*, and *I. paniculata* in Ecuador, Uruguay, and Argentina.

**Phyllosticta celosiae** Thuem. On leaves of *C. cristata* in Italy and Portugal.

**Uromyces celosiae** Diet and Holw. Brown rust pustules on leaves of *C. latifolia*, *Iresine calea*, and *I. canescens* in Mexico, Cuba, Guatemala, and Costa Rica.

**CELTIS.** HACKBERRY. Shade trees.

**Exoascus aemiliae** Pass. Grayish-white spots on leaves of *C. australis* in Italy. Probably the same as the following species.

**Exoascus celtidis** (Sad.) Sacc. Brown spots or small excrescences on leaves of *C. australis* in middle and south Europe.

**Mycosphaerella castagnei** (Har. and Br.) Jaap. On leaves of *C. australis* in Dalmatia.

**Napcladium celtidis** Cav. Subcircular to angular, olivaceous spots on leaves of *C. latifolia* in Italy.

**Phyllosticta celtidicola** Bub. and Kab. Angular to irregular, dull-yellow or brown leaf spots on *C. australis* in Austria.

**Phyllosticta destruens** Desm. Whitish spots on leaves of *C. australis*, *Prunus laurocerasus* and *P. lusitanica* in France, Denmark, Russia, Italy, Portugal, and Austria. Reported from Wisconsin and Kansas.

**Phyllosticta gei** Bres. On leaves of *C. australis* in Europe.

**Phyllosticta immersa** Bub. Small circular sinuate greenish or yellow leaf spots on *C. australis* in Hungary.

**CELTIS**—Continued.

- Phyllosticta talae** Speg. Circular ashen-colored leaf spots on *C. tala* in Argentina.
- Septoria celtidis** Sacc. On leaves of *C. australis* in France and Italy.
- Sporodesmium celtidis** Syd. Small, definite circular gray leaf spots on *C. australis* in Yugoslavia.
- Uncinula kusanoi** Syd. Powdery mildew on under sides of leaves of *C. sinensis* in Japan.
- Uncinula shiraiana** P. Henn. Powdery mildew on lower leaf surfaces of *C. sinensis* in Japan.
- Uredo celtidis** Pазsch. Brown powdery rust pustules on lower leaf surfaces of *Celtis* sp. in Peru and Brazil.
- Uromyces celtidis** Diet. Rust on leaves of *Celtis* sp. in Brazil.
- CENTAUREA**. CORN FLOWER. BACHELOR'S-BUTTON. Annuals or half-hardy perennials, used for bedding, borders, etc.
- Ascochyta cyani** Cruch. On leaves of *C. cyanus* in Switzerland.
- Ascochyta schelliana** Thuem. On leaves and stems of *C. glastifolia* in Siberia.
- Bremia centaureae** Syd. Downy mildew on leaves of *C. jacea* and *C. montana* in Europe.
- Cercospora centaureae** Diet. Irregular ashen-colored spots on both leaf surfaces of *C. phrygia* in Germany.
- Cercosporella centaureae** Syd. Small circular, or somewhat irregular, blackish leaf spots on *C. scabiosa* in Denmark and Germany.
- Cercosporella triboutiana** Sacc. and Let. Small angular leaf spots on *C. amara*, *C. nigrescens*, *C. phrygia*, and *C. pratensis* in Spain and France.
- Heterosporium centaureae** Ranoj. On stems of *C. spinulosa* in Yugoslavia.
- Phyllosticta tagana** (Thuem.) Fl. Tass. On leaves of *C. tagana* in Portugal.
- Puccinia arenariicolor** Plowr. Yellow to brown rust pustules on leaves of *C. nigra* and *Carex arenaria* in Great Britain.
- Puccinia beltrani** Gz. Frag. Rust on leaves of *C. lingulata* in Spain.
- Puccinia caricis-montanae** Ed. Fisch. Yellow rust pustules of *C. montana*, *C. nemoralis*, *C. nigra*, *C. nigrescens*, *C. plumosa*, and *C. scabiosa* in Russia, France, Switzerland, Austria, Hungary, and Germany.
- Puccinia centaureae** Mart. Brown rust pustules on many species of *Centaurea* spp. in India, Siberia, Asia Minor, Tripoli, and throughout Europe. Reported from several localities in the United States
- Puccinia leuzeae** Syd. Cinnamon-brown powdery rust pustules on leaves of *Leuzea* (*Centaurea*) *carthamoides* in Siberia.
- Puccinia montana** Fekl. Orange-yellow, then chestnut-brown or dark-brown, rust pustules on *C. cana*, *C. montana*, and *C. phrygia* in Anatolia, Belgium, Italy, Switzerland, Yugoslavia, Austria, Hungary, and Germany.
- Puccinia obducens** Syd. Dark-brown powdery rust pustules on leaves of *C. ruthenica* in Turkestan.
- Puccinia pencana** Syd. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *C. bulbosa* in Chile.
- Puccinia persica** Wettst. Yellow-brown to black powdery rust pustules on leaves of *C. belangerana*, *C. carduiformis*, and *C. kermanensis* in Persia.
- Puccinia tenuistipes** Rostr. Rust on leaves of *C. jacea* and *Carex muricata* in Yugoslavia, Sweden, Russia, Denmark, Italy, Austria, and Germany.
- Puccinia verruca** Thuem. Rust on leaves of *C. jacea*, *C. maculosa*, *C. montana*, *C. napifolia*, *C. nicaeensis*, *C. orientalis*, *C. pullata*, *C. romana*, *C. scabiosa*, *C. sonchifolia* and *C. spinulifolia* in Egypt, Algeria, Corsica, Yugoslavia, Spain, Italy, Russia, Denmark, Austria, Hungary, and Germany.
- Puccinia vestita** Syd. Yellow-brown rust pustules on leaves of *C. myriocephala* in Kurdistan and Asia Minor.
- Pyrenochaete centaureae** Vogl. Indefinite gray or ashen-brown leaf spots on *C. candidissima* in Italy.
- Ramularia centaureae** Lindr. Small whitish spots, with dull-brown margins on leaves of *C. dealbata*, *C. nigra*, *C. phrygia*, *C. salicifolia*, and *C. scabiosa* in Denmark, Finland, Russia, France, Switzerland, and Germany.
- Ramularia centaureae atro-purpureae** Bub. Subcircular, dull-brown leaf spots with yellow centers on *C. atro-purpurea* in Hungary.
- Septoria aderholdi** Vogl. Indefinite, ashen-brown leaf spots on *C. candidissima* and *C. pseudo-phrygia* in Italy and Austria.
- Septoria centaureae** (Roum.) Sacc. Small olivaceous leaf spots on *C. jacea* and *C. nigra* in France.
- Septoria centaureicola** Brun. Circular, purple-brown leaf spots with white centers on *C. phrygia*, *C. scabiosa*, and *C. spinulifolia* in Russia, Yugoslavia, and France.
- Septoria collinae** Gz. Frag. On leaves of *C. collina* in Spain.
- Septoria cyani** Höll. Indefinite brown leaf spots on *C. cyanus* in Hungary.
- CENTAURIUM**. See Erythraea.
- CENTRANTHUS**. JUPITER'S BEARD. Red valerian. Annual or biennial herbs.
- Aecidium centranthi** Thuem. Rust on leaves of *C. calcitrapa* in Spain, France, Switzerland, and Belgium.
- Puccinia valerianae** Carest. See Valeriana.
- Ramularia centranthi** Brun. Subcircular to irregular brown leaf spots on *C. ruber* in France.
- Septoria centranthicola** Brun. Angular to irregular, dirty-white leaf spots with narrow dull-green margins on *C. ruber* in France.
- CEPHALANTHUS**. BUTTONBUSH. Shrub grown for its attractive white flower heads.
- Aecidium cephalanthi-peruviani** P. Henn. Chestnut-brown rust pustules on red-brown leaf spots on *C. peruvianus* in Peru.
- Fusicladium cephalanthi** Speg. Superficial dense olivaceous fungal layers on branches, leaves, and young inflorescences, all of which as a result are more or less distorted. On *C. sarandus* in Uruguay and Argentina.
- CEPHALARIA**. Coarse annual or perennial herbs.
- Laetadia cephalariae** (Awd.) Sacc. See Alternanthera.
- Peronospora cephalariae** Vincens. Downy mildew on leaves of *C. alpina*, *C. leucantha*, and *C. transylvanica* in France.
- Phyllosticta cephalariae** Wint. Circular to irregular and finally confluent white leaf spots with dark purple margins on *C. attenuata* in the Union of South Africa.
- Septoria amicabile** Boy. and Jacz. Brown leaf spots on *C. leucantha* in France.
- Septoria cephalariae-alpinae** Roum. Small, numerous, often confluent, whitish leaf spots with purple margins on *C. alpina* in France.
- Septoria scabiosicola** Desm. See Scabiosa.
- Venturia cephalariae** Kalchbr. and Cke. Dull-brown, circular leaf spots on *C. attenuata* in the Union of South Africa.
- CERASTIUM**. SNOW-IN-SUMMER. MOUSE-EAR CHICKWEED. Decumbent annuals or perennials, used in rockeries or for borders.
- Aecidium balearicum** Gz. Frag. Rust on leaves of *Cerastium* sp. in the Balearic Islands.
- Ascochyta alpina** Rostr. On leaves of *C. alpinum* in Norway.

**CERASTIUM**—Continued.

- Ascochyta silenes** E. and E. f. *cerastii* Sacc. On leaves of *C. arvense* in Italy.
- Fabraea cerastiorum** (Fr.) Rehm. Yellow spots on leaves and stems of *Cerastium* spp. and *Stellaria* sp. in Argentina, Australia, and throughout Europe.
- Isariopsis alborosella** (Desm.) Sacc. See *Stellaria*.
- Peronospora helvetica** Gäum. Downy mildew on leaves of *C. latifolium* in Switzerland.
- Peronospora tornensis** Gäum. Downy mildew on leaves of *C. alpinum* in Norway, Sweden, and Iceland.
- Peronospora trivialis** Gäum. As above on *C. arvense*, *C. semidecandrum*, and *C. triviale* in Europe and Argentina.
- Septoria celanensis** D. Sacc. Discoloring leaves of *C. tomentosum* in Italy.
- Septoria cerasticola** Rostr. On leaves of *C. alpinum* in Denmark.
- Septoria cerastii** Desm. and Rob. Pale-yellow leaf spots, turning ashen-gray, on *C. pumilum*, *C. triviale* and *C. vulgatum* in Italy, Spain, France, Belgium, Russia, and Austria.
- Septoria commersoniana** Speg. On leaves of *C. humifusum* in Argentina.
- Ustilago duriaeana** Tul. A smut producing brown powdery spore masses in ovaries of *C. arvense*, *C. brachypetalum*, *C. glomeratum*, and *Stellaria media* (*Alsine media*) in French and Italian north Africa, Great Britain, and Germany.
- CERASUS.** See Prunus.
- CERATONIA.** CAROB. A leguminous evergreen tree, the pods of which are used for forage and to some extent for human food.
- Cercospora ceratonia** Pat. and Trab. Small circular or angular brown spots on leaves of *Ceratonia* sp. in Algeria and Malta.
- Phyllosticta calaritana** Briosi. and Cav. Irregular marginal brown blotches on leaves of *C. siliqua* in Italy.
- Phyllosticta ceratoniae** Berk. Irregular black marginal spots on leaves of *C. siliqua* in Tunis, Russia, Italy, Malta, and Portugal.
- Ramularia australis** Sacc. Leaf spot on *C. siliqua* in Italy.
- Septoria carruhi** Pass. Dark-brown, irregular spots on leaves and pods of *C. siliqua* in France and Malta. Said to "menace the existence of the trees in this part of the country."
- Septoria ceratoniae** Pass. On leaves of *C. siliqua* in Italy and France.
- CERCIDIPHYLLUM.** Tree grown for its handsome foliage and habit.
- Polythrincium shiraianum** P. Henn. Circular to angular dull-brown leaf spots on *C. japonicum* in Japan.
- CERCIS.** REDBUD. Judas tree. Trees or shrubs grown for their pink flowers.
- Aecidium lipskianum** Jacz. A rust on leaves and pods of *C. siliquastrum* in Turkestan and Russia.
- Ascochyta siliquastrum** Pass. On pods of *C. siliquastrum* in Italy.
- Ovularia cercidis** S. Cam. Irregular brown spots with deeper colored borders on leaves of *C. siliquastrum* in Portugal.
- Phaeosphaerella japonica** Shir. and Har. On leaves of *C. chinensis* in Japan.
- Phyllosticta siliquastrum** Sacc. and Speg. Reddish leaf spots on *C. siliquastrum* in Italy and Portugal.
- Septoria cercidis** Fr. On leaves of *C. canadensis* and *C. siliquastrum* in Italy, France, Albania, Yugoslavia, and Russia.
- CERCOCARPUS.** Mountain mahogany. Small trees or shrubs cultivated for their attractive foliage and peculiar feathery-tailed achenes.
- Ascochyta zonata** Syd. Circular brown leaf spots (0.5 to 1 centimeter in diameter) on *C. parvifolius* in Germany.
- Phyllosticta cercocarpi** Syd. Small circular pale leaf spots with red margins on *C. parvifolius* in Germany.
- CEREUS.** Arborescent, columnar forms of cactus.
- Aecidium cerei** P. Henn. Rust on *Cereus* sp. in Argentina.
- Cytospora cerei** Póll. On *Cereus* sp. in Italy.
- Gloeosporium josephinae** D. Sacc. Anthracnose on branches of *C. nycicalis* in Italy.
- Leptothyrium parasiticum** Pollacci. Large gray or ochraceous spots on stems of *C. stellatus* and *C. triangularis* in Italy.
- Montagnella opuntiarum** Speg. On *Cereus* sp. and *Cactus peruvianus* in Brazil and Paraguay.
- Mycosphaerella cerei** P. Henn. On phyllodes of *Cereus macrogenus* in Brazil.
- Septoria cacticola** P. Henn. Pale-brown, indefinite spots, often confluent, on trunks of *C. pentagonus* in Germany.
- Stagonospora assans** Pass. On *Cereus* sp. and *Echinocactus* sp. in Italy.
- CERINTHE.** HONEYWORT. Annual or perennial herbs, with showy purple bracts.
- Coleosporium cerinthes** Schroet. Golden-yellow rust pustules on leaves of *C. minor* in Russia and Germany.
- Ramularia cerinthes** Hóll. Circular, black leaf spots, with dull-yellow margins on *C. minor* in Hungary.
- CEROPEGIA.** Tropical vines cultivated in greenhouses.
- Aecidium nummulare** Berk. Rust on leaves of *C. biflora*, *C. candelabrum*, and *C. cumingiana* in Ceylon and the Philippines.
- Phyllachora ajrekari** Syd. Irregular black stromata on both leaf surfaces of *C. hirsuta* in India and Japan.
- CESTRUM.** Ornamental tropical shrubs.
- Didymosporium stromaticum** Pat. On leaves of *Cestrum* sp. in Ecuador.
- Phyllachora cestri** Pat. Black stromata on leaves of *Cestrum* sp. in Ecuador.
- Phyllosticta cestri** Speg. Ashen leaf spots with reddish borders on *C. pubescens* in Argentina.
- Phyllosticta humerispora** Speg. On leaves of *Cestrum* sp. in Brazil.
- Puccinia cestri** Diet. and P. Henn. Brown rust pustules on lower leaf surfaces of *Cestrum* sp. in Brazil and Ecuador.
- Septoria cestri** (Mont.) Sacc. Circular white leaf spots with dull-brown margins on *C. parqui* in Chile.
- Stigmatea cestri** Pat. Black, circular spots on leaves of *Cestrum* sp. in Costa Rica.
- Tuberculina jaffueli** Speg. On leaves of *C. parqui* in Chile.
- Uromyces cestri** Mont. Yellow and powdery black rust pustules on leaves of *C. athroxanthum*, *C. aurantiacum*, *C. auriculatum*, *C. lanatum*, *C. laurifolium*, *C. macrophyllum*, *C. pallidum*, *C. parqui*, and *C. pseudoquina* in Chile, Peru, Bolivia, Brazil, Argentina, Uruguay, Porto Rico, Virgin Islands, Jamaica, Mexico, Costa Rica, and Guatemala.
- Uromyces maculans** Arth. Brown leaf rust on *C. nocturnum* in Costa Rica.
- Uromyces venustus** Diet. and Holw. Rust on leaves of *C. nitidum* in Mexico.

- CHAEROPHYLLUM.** Scented annual, biennial, or perennial herbs, often with tuberous roots.
- Protomyces macrosporus** Ung. See Coriandrum.
- Puccinia aromatica** Bub. and Lindr. Golden-yellow and deep-brown rust pustules on leaves of *C. aromaticum* in Russia, Austria, Hungary, and Germany.
- Puccinia chaerophylli** Purt. Cinnamon-brown to dark-brown rust pustules on leaves, petioles, and stems of *C. aureum*, *C. coloratum*, *C. hirsutum*, *C. temulum*, *Anthriscus sylvestris*, *A. nemerosa*, *A. tenerrima*, and *Myrrhis odorata* in Europe and Siberia.
- Puccinia enormis** Fekl. Cinnamon-brown, powdery rust pustules on leaf blades, and petioles and on stems of *C. villarsium* in Switzerland and Austria.
- Puccinia pozzii** Sem. Leaf rust on *C. hirsutum* in Europe.
- Puccinia prescottii** Lindr. Dark-brown or black rust pustules on leaf blades and petioles of *C. prescottii* in Russia.
- Puccinia retifera** Lindr. Yellow and cinnamon-brown to dark-brown rust pustules on leaves of *C. bulbosum* in Russia, Hungary, and Germany.
- Septoria chaerophylli** Bres. Leaf spot on *C. aromaticum* in Germany.
- CHAETOCHELOA.** Annual grasses, mostly weeds. See also *Setaria*.
- Puccinia cameliae** (Mayor) Arth. Rust on leaves of *C. setosa* in Porto Rico and Colombia.
- Sphaeclotheca pamparum** (Speg.) Clint. Sori infecting the ovaries of the spikes, extending between the glumes as oblong bodies, 3 to 7 millimeters long, inclosed by a false membrane which ruptures irregularly at the apex, disclosing the powdery, olive-black spore masses. On *C. geniculata*, *C. imberbis*, and *Chaetochloa* sp. in Argentina, Cuba, and the Bahamas.
- CHALCAS.** ORANGE JASMINE. Small evergreen-leaved shrubs.
- Bacterium citri** Hasse. See Citrus.
- Corticium salmonicolor** B. and Br. See Citrus.
- CHAMAECYPARIS.** WHITE CEDAR. Trees or shrubs cultivated for their handsome evergreen foliage. Timber trees.
- Asterula chamaecyparisii** Shir. and Hara. On needles of *C. obtusa* in Japan.
- Coniothyrium pallido-fuscum** Sacc. On needles of *C. pisifera squarrosa* (*Retinospora*) and *Araucaria brasiliensis* in France.
- Gymnosporangium solenoides** (Diet.) Kern. (*G. miyabei* Yam. and Miy.). A rust with the aecial (cluster-cup) stage on leaves of *Sorbus alnifolia* and *S. aria*, and the telial stage on branches of *Chamaecyparis pisifera* (*C. plumosa*) in Japan. The telial stage is characterized by elongated reddish-brown swellings with rough surfaces, on which red-brown spore masses appear.
- Helicobasidium mompa** Tanaka. See Morus.
- Lophodermium chamaecyparisii** Shir. and Hara. On needles of *C. obtusa* in Japan.
- CHAMAEDOREA.** See Palmae.
- CHAMAEROPS.** See Palmae.
- CHEIRANTHUS.** WALLFLOWER. Flower-garden perennials.
- Ascochyta cheiranthi** Bres. Circular to oblong yellow or brown spots on leaves of *C. cheiri* in Alaska and Germany.
- Bacterium** sp. A bacterial disease of *C. annuus* is reported from Holland, characterized by a constriction of the upper part of the tap root of diseased plants, followed by yellowing and dropping of the leaves. The vascular bundles in the roots are stained brown or black.
- Cercospora cheiranthi** Sacc. White leaf spots on *C. cheiri* in Italy.
- Peronospora cheiranthi** Gäum. Downy mildew on leaves of *C. cheiri* in Europe.
- Phyllosticta cheiranthicola** Bub. and Zimm. Circular, greenish then yellowish spots on leaves of *C. cheiri* in Bohemia.
- Phyllosticta cheiranthorum** Desm. Scattered irregular white or gray-green leaf spots on *C. cheiri* in France.
- Septoria cheiranthi** Rob. and Desm. Circular or irregular leaf spots on *C. cheiri* in France.
- CHELIDONIUM.** CELANDINE POPPY. Garden herbs.
- Ascochyta chelidonii** Kab. and Bub. Irregular brown to black leaf spots on *C. major* in Bohemia.
- Didymaria chelidonii** Jacz. Indefinite brown leaf spots on *C. uniflorum* in Siberia.
- Melampsora magnusiana** G. Wagn. See Populus.
- Peronospora chelidonii** Miy. Downy mildew on leaves of *C. major* in Japan.
- Phyllosticta chelidonii** Bres. Irregular yellow to dull-brown leaf spots on *C. major* in Germany.
- Septoria chelidonii** Desm. Circular to angular greenish-olive or brown leaf spots with white centers on *C. major* and *Argemone mexicana* in Porto Rico, China, Japan, Siberia, and throughout Europe.
- CHENOPODIUM.** GOOSEFOOT. Quinoa. Weedy herbs.
- Ascochyta boni-henrici** Ranof. On leaves of *C. bonus-henricus* in Yugoslavia.
- Ascochyta chenopodii** Rostr. See Atriplex.
- Ascochyta nebulosa** Sacc. and Berl. and f. **foliicola** Gz. Frag. On stems and leaves of *C. glaucum* in Siberia, Italy, and Spain.
- Cercospora chenopodii** Fres. On leaves of *Chenopodium* sp. in Denmark, Russia, and Japan.
- Cercospora chenopodiicola** Bres. Small subcircular pale-brown leaf spots on *C. polyspermum* in Russia and Germany.
- Cercospora macularis** (Schroet.) Magn. On leaves of *C. bonus-henricus* in Austria.
- Hypochnus peronosporoides** Speg. Thin white fungus layers on lower leaf surfaces of *C. ambrosioides* in Argentina.
- Ovulariopsis erysiphoides** Pat. and Har. On leaves of *C. ambrosioides* in Spain.
- Peronospora bohemica** Gäum. Downy mildew on leaves of *C. ficifolium* in Bohemia.
- Peronospora bonihenrici** Gäum. Downy mildew on leaves of *C. bonus-henricus* in Europe.
- Peronospora chenopodii-ficifolia** Saw. Circular to irregular pale-yellow spots, often covering entire leaf area of *C. ficifolium* in Japan.
- Peronospora chenopodii polyspermi** Gäum. Downy mildew on leaves of *C. polyspermum* in central and north Europe.
- Peronospora chenopodii rubri** Gäum. As above on *C. rubrum* in Holland, Yugoslavia, and Germany.
- Peronospora muralis** Gäum. Downy mildew on leaves of *C. muralis* in Europe and Argentina.
- Phragmidium ambrosioides** Thuem. Leaf rust on *C. album* in Yugoslavia.
- Phyllosticta dimorphospora** Speg. Circular pale-brown indefinite leaf spots on *C. hircinum* in Argentina.
- Ramularia chenopodii** Speg. On leaves of *C. album* in Argentina.
- Ramularia dubia** Reiss. On leaves of *C. album* and *C. patula* in Russia and Germany.
- Ramularia macularis** Schroet. Angular pale-yellow then white leaf spots on *C. bonus-henricus* in Switzerland, Poland, Austria, and Germany.
- Rhizoctonia napi** West. See Brassica.
- Septoria ambrosioides** Speg. On leaves of *C. ambrosioides* in Argentina.
- Septoria atriplicis** (West.) Fekl. See Atriplex.

**CHENOPODIUM**—Continued.

**Septoria chenopodii** West. Greenish-white than yellow leaf spots on *C. album*, *C. bonus-henricus*, *C. murale*, *C. rubrum*, and *C. viride* in Belgium, Portugal, Italy, and Germany.  
**Septoria undullospora** Bub. On leaves of *C. murale* and *C. vulvaria* in Spain and Hungary.  
**Tolyposporium leptideum** Syd. Cinnamon-brown smut sori in ovaries of *Chenopodium* sp. in Germany.

**Urophlyctis pulposa** (Wallr.) Schroet. See *Atriplex*.

**CHILOGLOTTIS**. See Orchidaceae.**CHIONODOXA**. GLORY-OF-THE-SNOW. Spring bulbs.

**Ustilago vaillantii** Tul. See *Hyacinthus*.

**CHIONANTHUS**. FRINGE TREE. Woody plants grown for their profuse white flowers.

**Aecidium chionanthi** B. and Br. Rust on leaves of *Chionanthus* sp. in Ceylon.

**Phyllosticta chionanthi** Thuem. Irregular yellowish-white leaf spots with narrow red-brown margins on *C. virginica* in Portugal.

**Septoria velespora** Sacc. On leaves of *C. virginica* in France.

**CHLOBIS**. FINGER GRASS. Rhodes grass. Annual or perennial forage grasses. Some species grown as ornamentals.

**Balansia discoidea** P. Henn. On culms of *C. distichophylla* and *Chloris* sp. in Peru and Brazil.

**Heterosporium chloridis** Speg. On leaves of *Chloris* sp. in Brazil.

**Phyllachora chloridicola** Speg. Black stromata on leaves of *C. distichophylla* and *C. radiata* in Argentina.

**Puccinia chloridicola** P. Henn. Dull yellow-brown to black elongate rust pustules on linear dull-brown leaf spots on *Chloris* sp. in the Congo.

**Puccinia chloridina** Bacc. Black linear rust pustules on leaves of *Chloris* sp. in Abyssinia.

**Sorosporium chloridicola** Beeli. Smut on *C. polydactyla* in the Congo Free State.

**Tolyposporium chloridis** P. Henn. Spikelets of *C. abyssinica* and *C. gayana* become black smutty masses in central and south Africa.

**Ustilago deserticola** Speg. Brown-black powdery smut sori in culms and rhizomes of *Chloris* sp. in Argentina.

**Ustilago ulei** P. Henn. Dark-brown or dark olivaceous spore masses covering and often distorting the leaves of *Chloris* sp. in Brazil.

**CHLOROPHYTUM**. Rhizomatous herbaceous plants.

**Aecidium chlorophyti** Har. and Pat. Brown rust pustules on large leaf spots on *Chlorophytum* sp. in African Congo.

**CHORIZEMA**. Small shrubs grown for their showy pea-like yellow, orange, and red flowers.

**Phyllosticta chorizemae** P. Henn. Brown confluent leaf spots on *Chorizema* sp. in Germany.

**Septoria straussiana** P. Henn. Dull-brown then paler leaf spots with red-brown margins on *Chorizema* sp. in Germany.

**CHRYSANTHEMUM**. Herbaceous and subshrubby plants cultivated in greenhouses and out of doors for their flowers.

**Aphelenchus olesistus** Ritz. Bos. See *Begonia*.

**Aphelenchus ritzei-bosi** Schwartz. This nematode is parasitic on the leaves of *Chrysanthemum* (cult.), causing them to turn brown and fall prematurely. Blossom buds also shrivel up and the entire plant ultimately dies. The disease occurs in England and Germany.

**Cercospora chrysanthemi** Puttem. On leaves of *C. indicum* in Brazil. Heald and Wolf have used the same name for a species found in Texas.

**Corticium chrysanthemi** Plow. Said to cause a basal rot of stems of *Chrysanthemum* (cult.) in Great Britain.

**Entyloma leucanthemi** Syd. Small circular yellow-green leaf spots on *C. leucanthemum* in Switzerland and Austria.

**Peronospora danica** Gäum. Downy mildew on leaves of *C. segetum* in Denmark.

**Peronospora leptosperma** (De By.) Gäum. As above on *C. inodorum*, *Matricaria chamomilla*, and *M. discoidea* in Europe.

**Peronospora radii** De By. Downy mildew attacking leaves and deforming flower heads of *Achillea ptarmica*, *Anthemis arvensis*, *A. austriaca*, *Chrysanthemum leucanthemum*, *Matricaria chamomilla*, and *M. inodora* in Switzerland, France, Sweden, Denmark, Belgium, Italy, Finland, Bohemia, Austria, and Germany.

**Phoma chrysanthemi** Vogl. On stems and leaves of *Chrysanthemum* sp. in Italy, causing a wilt.

**Phyllosticta leucanthemi** Speg. Grayish-white circular leaf spots on *C. leucanthemum* in Italy.

**Protomyopsis chrysanthemi** Büren. Flat yellowish-white callosities on lower leaf surfaces of *C. alpinum* in Switzerland.

**Protomyopsis leucanthemi** Büren. On leaves of *C. leucanthemum* in Switzerland.

**Puccinia aecidii-leucanthemi** Ed. Fisch. Brown to black rust pustules on leaves of *C. leucanthemum* in Europe. The aecial stage occurs on species of *Carex*.

**Puccinia horiana** P. Henn. Dark-brown to black rust pustules on lower leaf surfaces of *C. chinense*, *C. decaisneanum*, and *C. morifolium* (*C. sinense*) in Japan.

**Puccinia leucanthemi** Pass. Rust pustules on leaf blades and petioles of *C. leucanthemum* in Italy.

**Puccinia proximella** Syd. Brown rust pustules on leaves of *C. millefoliatum* in Russia.

**Puccinia pyrethri** Rabh. Pale brown to black rust pustules on leaves of *C. chinense*, *C. corymbosum*, *C. parthenifolium*, *C. parthenium*, and *C. morifolium* in Japan, Russia, Switzerland, Italy, Spain, Portugal, Rumania, Austria, Hungary, and Germany.

**Septoria cercosporoides** Trall. Irregular indefinite spots on leaves of *C. leucanthemum* in Scotland.

**Septoria chrysanthemella** Cav. Circular dull red-brown leaf spots of varying size and with yellow centers on *C. morifolium* and *Chrysanthemum* (cult.) in Denmark, Italy, and Japan. Also said to cause a damping-off of seedlings.

**Septoria chrysanthemi-indici** Bub. and Kab. Circular, rarely irregular, dull-brown to dark-brown leaf spots on *C. indicum* in Bohemia.

**Septoria chrysanthemi-rotundifolii** Namys. Brown circular or irregular, often confluent, leaf spots on *C. rotundifolium* in Poland.

**Septoria leucanthemi** Sacc. and Speg. Yellow-brown circular or irregular leaf spots with white centers on *C. (Leucanthemum) maximum* in Italy and Portugal.

**Septoria obesa** Syd. Circular to irregular leaf spots on *C. arcticum* in Japan.

**Septoria rostrupii** Sacc. and Syd. Circular dark-brown leaf spots on *C. indicum* in Russia and Denmark.

**Septoria socia** Pass. and f. *catalaunica* Gz. Frag. Subcircular to irregular dull-brown leaf spots on *C. leucanthemum* in Italy, Spain, and Portugal.

**Uredo autumnalis** Diet. Yellow-brown rust pustules on leaves of *C. boreale*, *C. indicum*, and *C. sinense* in Japan.

- CHRYSOBALANUS.** COCOA PLUM. Icaco. Shrubs or trees with edible fruit. Sometimes planted for ornament.
- Helminthosporium chrysobalani** P. Henn. Large indefinite leaf spots on *Chrysobalanus* sp. in the Congo.
- Mycosphaerella chrysobalani** Miles. Irregular spots, generally confluent and occupying most of the leaf surface on *C. icaco* in Porto Rico.
- CHRYSOPHYLLUM.** STAR APPLE. Cainito. Handsome broad-leaved evergreen trees grown for fruit and ornamental effect.
- Helminthosporium chrysophylli** P. Henn. Dark-brown circular leaf spots on *Chrysophyllum* sp. in East Africa.
- Phyllosticta chrysophylli** Syd. Indefinite gray leaf spots on *C. imperialis* in Germany.
- Septoria evansii** Syd. Small circular white leaf spots with dull-purple margins on *C. magalis-montana* in the Union of South Africa.
- Uredo chrysophyllicola** P. Henn. Brown rust pustules on dark-brown leaf spots on *Chrysophyllum* sp. in Brazil.
- CHRYSOSPLENIUM.** GOLDEN SAXIFRAGE. Semiaquatic plants.
- Entyloma chrysosplenii** (B. and Br.) Schroet. Smut sori in small circular white or yellow leaf spots on *C. alternifolium* in Great Britain, Denmark, and Germany.
- Peronospora chrysosplenii** Fckl. Downy mildew on leaves of *C. alternifolium* in central and northern Europe.
- Puccinia chrysosplenii** Grev. Brown rust pustules on leaves of *C. alternifolium*, *C. oppositifolium*, *C. trachyspermum*, and *Mitella longiscapa* in Japan, eastern Asia, Russia, Finland, Belgium, Great Britain, Austria, Hungary, and Germany.
- Septoria posoniensis** Bauml. Circular or irregular greenish-gray or ashen leaf spots with narrow brown margins on *C. alternifolium* and *C. oppositifolium* in Great Britain and Hungary.
- CHUSQUEA.** See Bambuseae.
- CICER.** CHICK-PEA. Garbanzo. Gram. Herbaceous legumes.
- Erysiphe taurica** Lévl. See Althaea.
- Phyllosticta rabiei** (Pass.) Trott. (*Phyllosticta cicerina* Prill. and Delacr.) Dull-yellow to gray leaf spots on *C. arietinum* in France, Italy, and Spain. The stems, petioles, and rachides are also attacked, diseased plants often breaking over at the lesions and dying.
- Rhizoctonia napi** West. See Brassica.
- Uromyces ciceris-arietini** (Grogn.) Jacz. and Boy. Small round or oval cinnamon-brown powdery rust pustules, which tend to coalesce, on both leaf surfaces of *C. arietinum* in India, East Africa, Yugoslavia, Spain, Portugal, France, and Italy.
- CICORIUM.** CHICORY. Endive. Herbs cultivated for their fleshy roots and as salad plants.
- Puccinia cichorii** (DC.) Bell. Cinnamon-brown to dark-brown rust pustules on leaves and stems of *C. endivia* and *C. intybus* in India, Argentina, and Europe. Reported from Oregon.
- Puccinia endiviae** Pass. Cinnamon-brown to dark-brown powdery rust pustules on leaves and stems of *C. endivia* in Argentina, Italy, Spain, France, Holland, and Denmark. Reported from Connecticut.
- Ramularia lampsanae** (Desm.) Sacc. On leaves of *C. intybus*, *Lampsana communis*, and *Taraxacum officinale* in Europe.
- Septoria endiviae** Thuem. On leaves of *C. endivia* in Austria and Holland.
- Septoria intybi** Pass. On branches of *C. intybus* in Italy.
- CIMICIFUGA.** BUGBANE. Tall ornamental herbaceous perennials.
- Aecidium shiraianum** Syd. Rust on leaves of *C. biternata*, *C. foetida*, *C. intermedia*, and *C. japonica* in Japan and Siberia.
- Coleosporium cimicifugatum** Thuem. Yellow or orange rust pustules on leaves of *C. foetida*, *C. intermedia*, and *C. simplex* in Siberia and Japan.
- CINCHONA.** Tropical trees cultivated for their medicinal bark, which is sometimes called Peruvian bark and from which quinine is made.
- Colletotrichum cinchonae** Koord. Anthracnose on leaves of *Cinchona* sp. in Java.
- Corticium salmonicolor** B. and Br. See Citrus.
- Dasyscypha warburgiana** P. Henn. Causes a serious canker disease of limbs of *C. ledgeriana* in Java.
- Guignardia yersini** Vinc. On *C. ledgeriana* and *C. succirubra* in Indo-China.
- Moniliopsis aderholdi** Ruhl. See Solanum.
- Phlyctaena cinchonae** Vinc. Causes roughened, cankered areas on bark of branches of *Cinchona* spp. in Indo-China.
- Phyllosticta cinchonae** Koord. Irregular yellowish spots on leaves of *Cinchona* sp. in Java. The name of this species is untenable, the following having priority.
- Phyllosticta cinchonae** Pat. Angular dull-yellow leaf spots on *Cinchona* sp. in Ecuador.
- Phyllosticta cinchonae** Vinc. Circular deep-brown patches with violet borders on leaves of *C. ledgeriana* and *C. succirubra* in Indo-China. On old leaves a shot hole effect is produced; on young leaves the numerous spots cause deforming. Developing shoots are also involved, stunting the plants.
- Phyllosticta honbaensis** Vinc. On leaves of *C. ledgeriana* and *C. succirubra* in Indo-China.
- Phyllosticta yersini** Vinc. Attacks young leaves and shoots of *Cinchona* spp. in Indo-China.
- Physalospora cinchonae** Vinc. On *C. ledgeriana* and *C. succirubra* in Indo-China.
- Tulasnella cinchonae** Racib. Thin pale-rose fungus layers on lower leaf surfaces of *C. ledgeriana*, *C. officinalis*, and *C. succirubra* in Java.
- Uredo cinchonae** P. Henn. Brown rust pustules on leaves of *Cinchona* sp. in Java.
- CINERARIA.** Herbs cultivated for their flowers. See also Senecio.
- Aecidium cinerariae** Rostr. Rust on leaves and stems of *C. cruenta* (*Senecio cruentus*) and *C. palustris* in Siberia, Denmark, Austria, and Germany. Said to be a stage of *Puccinia eriophori* Thuem. which occurs on *Eriophorum* spp.
- Ascochyta cinerariae** Fl. Tass. On leaves of *C. maritima* in Italy.
- Ascochyta fibricola** Sacc. On stems of *C. cruenta* and *C. maritima* in France.
- Coleosporium senecionis** Fr. See Senecio.
- Entyloma calendulae** (Oud.) De By. See Calendula.
- Phytophthora cryptogea** Pethybr. and Laff. See Lycopersicum.
- Puccinia cinerariae** McAlp. Orange-yellow and chestnut-brown to black rust pustules on leaves of *Cineraria* (cult.) in Australia.
- Ramularia senecionis** (B. and Br.) Sacc. See Senecio.
- CINNAMOMUM.** CINNAMON. CAMPHOR TREE. CASSIA BARK TREE. Evergreen trees and shrubs with aromatic leaves and wood, yielding aromatic and medicinal products.
- Aecidium cinnamomi** Rac. Rust on leaves, petioles, and stems of *C. iners* and *C. zeylanicum* in Java.
- Ascochyta camphorae** Turc. Whitish spots with purple borders on leaves of *C. glandulifera* in Italy.
- Atichia millardeti** Racib. Black, superficial, radiate fruiting bodies on leaves of *Anacardium occidentale*, *Cinnamomum zeylanicum*, *Coffea arabica*, *Styrax benzoin*, and *Syzygium jambolanum* in Java and the Malay Peninsula.

**CINNAMOMUM**—Continued

- Colletotrichum briosii** Turc. Anthracnose on leaves *C. burmanni* in Italy.
- Corticium salmonicolor** B. and Br. See Citrus.
- Cytospora cinnamomi** Turc. Circular to irregular white leaf spots with dull-brown margins on *C. burmanni* in Italy.
- Desmotascus cinnamomi** Petch. Attacks the trunks and branches of *C. zeylanicum* in Ceylon.
- Diplodia cacaicola** P. Henn. See Theobroma.
- Exobasidium cinnamomi** Petch. Rough excrescences up to 8 centimeters long and of pale-brown color on branches of *C. cassia* and *C. zeylanicum* in Ceylon.
- Fomes carophylli** (Rac.) Bres. Causes a root rot and wood rot of *C. zeylanicum*, *Citrus* spp., *Dalbergia latifolia*, *Mesua ferrea*, and other economic plants in Java, Ceylon, Mauritius, Japan, and Central Africa.
- Fomes lamaoensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Gloeosporium camphorae** Sacc. Anthracnose on leaves of *C. camphora* (*C. officinarum*) in Italy. Reported from Texas and Alabama.
- Glomerella cinnamomi** Yosh. (*Guignardia cinnamomi* [Yosh.] Hara.) Anthracnose on leaf blades and petioles, buds, and young shoots of nursery stock of *C. camphora* in Japan and Formosa. The diseased spots are circular, elliptical or fusiform, reddish-brown at first and finally fading out to a light brown. The stems are girdled by a brown rot, the plants dying.
- Hypochnus cucumeris** Frank. See Cucumis.
- Hypochnus sasakii** Shirai. Leaf blight of *C. camphora* in Japan.
- Leptosphaeria cinnamomi** Shir. and Hara. Branch canker and leaf spot of *C. camphora* in Japan.
- Macrophoma cinnamomi-glanduliferi** Luigi. Large irregular blotches on leaves of *C. glanduliferum* in Italy.
- Phyllosticta aromatophila** Sacc. Small gray leaf spots on *C. camphora* (*C. officinarum*) in Italy.
- Phyllosticta cinnamomi** Delacr. Irregular dull-yellow leaf spots on *C. zeylanicum* in India.
- Phyllosticta cinnamomi-glanduliferi** P. Henn. Circular, finally confluent, dull-brown leaf spots on *C. camphora* and *C. glanduliferum* in Russia and Germany.
- Phytophthora cinnamomi** Rands. Irregular vertical stripes of dead bark (1 to 5 centimeters wide) appear on the trees, extending upward from the ground for a distance of 1 to 10 meters. At the upper end of each canker drops of wine or amber colored exudate appear. The host is *C. burmanni* in Sumatra.
- Rosellinia arcuata** Petch. See Thea.
- Rosellinia bunodes**. B. and Br. See Citrus.
- Scolecotrichum cinnamomi** Rac. Circular gray leaf spots from 1 to 4 centimeters in diameter on *C. zeylanicum* in Java.
- CISSAMPELOS**. Twining herbs or shrubs.
- Cercospora pareirae** Speg. Circular dull-brown leaf spots on *C. pareira* in Argentina.
- Phyllosticta cissampeli** Speg. Small brown leaf spots with white centers on *C. pareira* in Argentina.
- Uromyces cissampelidis** Diet. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *Cissampelos* sp. in Colombia.
- CISSUS**. Tropical woody climbers.
- Aecidium cissigenum** Welw. Leaf rust on *C. adenanchus* in Abyssinia.
- Cercospora riachuclii** Speg. On leaves of *C. palmata* in Argentina.
- Cronartium wilsonianum** Arthur. Straw-colored to dark-brown powdery rust pustules on leaves of *C. rhombifolia* in Cuba, Costa Rica, and Ecuador.
- Endophyllum guttatum** (Kunz.) Syd. (*Endophyllum circumscriptum* [Schw.] W. and O.) Yellow rust pustules on large swollen areas on leaves of *C. quadrangularis*, *C. rhombifolia*, and *C. sicyoides* in Italian north Africa, Central and South America, Trinidad, and the West Indies.
- Entyloma cissigenum** P. Henn. Smut sori in small, circular, black, often confluent, leaf spots on *Cissus* sp. in tropical Africa.
- Phyllosticta cissicola** Speg. Small, circular, whitish leaf spots with brown borders on *C. sicyoides* in Argentina and Porto Rico.
- Schizonella colemani** Iyen and Nar. This smut causes abnormal growths resembling witches' brooms at the nodes of *C. quadrangularis* in India. The diseased branches are pale-green in color and contain dark-green elongate swellings within which the spores form as black powdery masses.
- Schroeteria arabica** P. Henn. Smut on *C. quadrangularis* in Arabia and India. Probably the same as the following species.
- Schroeteria cissi** (DC.) De T. (*Mykosyrinx cissi* [DC.] G. Beck.) Smut sori in the peduncles and pedicels of inflorescences distorting the infected parts somewhat and filling them with a dusty-purple spore mass. On *C. acida*, *C. erosa*, *C. quadrangularis*, and *C. sicyoides* in Uganda, Cameroon, Mexico, South America, and the West Indies. Also reported from Florida.
- Uromyces cladomanes** Traverso. Rust causing coralloidal formations on *Cissus* sp. in Somalia.
- CISTUS**. ROCKROSE. Low shrubs grown for their red or white flowers.
- Ovulariopsis cisti** Jaap. On leaves of *C. albidus*, *C. monspeliensis*, and *C. salvifolius* in Dalmatia.
- Phyllosticta cistina** Thuem. Small circular purplish black, finally ashen-colored leaf spots with purple borders on *C. crispus*, *C. laurifolius*, and *C. salvifolius* in Greece, France, and Portugal.
- Uredo helianthemii** Rbh. See Helianthemum.
- CITHAREXYLUM**. Shrubs or trees cultivated for their ornamental effect.
- Phyllachora citharexylifera** (Rehm.) v. Hoeh. Black stromata on leaves of *C. suberosum* and *Citharexylum* sp. in Brazil.
- CITROPSIS**. AFRICAN CHERRY ORANGE. Trees allied to *Citrus*.
- Bacterium citri** Hasse. See Citrus.
- CITRULLUS**. WATERMELON. Colocynth. Annual or perennial vines.
- Hypochnus cucumeris** Frank. See Cucumis.
- Puccinia citrulli** Syd. and Butl. Powdery red-brown rust pustules on leaves of *C. colocynthus* in India.
- Scolecotrichum melophthorum** Prill. and Delacar. See Cucumis.
- Septoria citrullicola** Poteb. Small circular white leaf spots on *C. vulgaris* in Russia.
- CITRUS**. ORANGE. LEMON. GRAPEFRUIT. CITRON. LIME. MANDARIN ORANGE. TANGERINE. Tropical and subtropical fruit trees.
- Ascochyta citri** Penz. Nearly circular dark-brown, finally grayish, leaf spots with definite margins on *C. limonia* and *C. sinensis* in Italy and Brazil.
- Ascochyta corticola** McAlp. This "bark-blotch" disease attacks the lemon (*C. limonia*) and orange (*C. sinensis*) in Australia and New Zealand. The disease begins as dark-brown spots on the lower portion of the trunk and spreads rapidly until the trunk is girdled, death of the tree resulting.
- Ascochyta destruens** (McAlp.) F. Tass. Minute black pustules on yellowish-brown scab-like patches with red-brown margins on leaves of *Citrus sinensis* in Australia. Diseased leaves wither and the scabby areas fall out.

## CITRUS—Continued.

- Bacillus citrimaculans** Doidge. Bacterial spot. This bacterial disease produces circular to irregular slightly sunken spots on the fruit which are yellowish at first but soon become brown, varying in shade, and often with red margins. The skin only may be affected, or the disease may penetrate into the pulp with a peculiar accompanying odor. Under moist conditions yellow viscid drops appear. Branches are attacked, infection taking place at points of attachment of the leaves, water-soaked areas of 6 to 10 millimeters in diameter appearing which increase in size. Leaves and twigs within these areas wither and die. On the larger branches irregular brown areas occur. Practically all *Citrus* spp. are attacked, including orange, lemon, citron, mandarin, tangerine, lime, and grapefruit in the Union of South Africa.
- Bacterium citri** Hasse. Citrus canker attacks all parts of the tree above ground. On the leaves the disease first appears as minute yellow-brown translucent spots, which soon become visible on both sides. These spots increase rapidly in size, and the tissues at the center soon rupture, giving the characteristic corky appearance, white or pinkish at first, later brown. The corky raised areas are surrounded by a yellowish-brown, oily, translucent area, and are inclosed by yellowish-green zones which shade into the normal green. The spots vary in size from a quarter to half an inch at maturity and are circular in shape, occurring singly or in groups. Similar corky areas occur on the twigs, thorns, branches, and fruit, tending to be more irregular and more prominent, and with a tendency to develop cracks or fissures. The fruit is attacked only while green, and may ripen prematurely and fall. While the canker spots do not penetrate deeply, the cracking of the spots opens a way for decay organisms.
- This disease occurs in China, Japan, the Philippines, Hawaii, Mauritius, Ceylon, Siam, Dutch East Indies, Borneo, and Formosa. It has been introduced into the Union of South Africa, northern territory of Australia, and the Gulf States of the United States, where eradication campaigns are in progress. Practically all species and varieties are susceptible, although to a greatly varying degree, the grapefruit (*C. grandis*), *Citrus (Poncirus) trifoliata*, lime (*C. aurantifolia*) and round orange (*C. sinensis*), being especially so, and the Satsuma, King orange and kumquat rather resistant. In addition to the large number of natural hosts of the genus *Citrus* and its segregates, many *Citrus* relatives have been found susceptible to a greater or less extent through artificial inoculations. Among these are *Aegle marmelos*, *Atalantia citrioides*, *A. ceylonica*, *A. disticha*, *Casimiroa edulis*, *Chaetosperrum glutinosum*, *Chalcas exotica*, *Citropsis schweinfurthii*, *Claucaena lansium*, *Eremocitrus glauca*, *Evodia ridleyi*, *E. latifolia*, *Feronia limonia*, *F. lucida*, *Hesperethusa crenulata*, *Melicope triphylla*, *Microcitrus australasica*, *M. garrowayi*, *M. australis*, *Paramignya longipedunculata*, *Poncirus trifoliata*, and *Toddalia asiatica*.
- Bacterium gummi** Comes. This bacterium is reported from Italy, Paraguay, and Brazil as the cause of a gumming disease of twigs, branches, and fruit of orange (*C. sinensis*) and lemon (*C. limonia*). In Brazil the type of disease known as mal-di-gomma or foot rot is also attributed to this organism. The disease is apparently the same as the foot rot and gummosis of Florida, now attributed to other causes.
- Bark rot.** This disease attacks the calomondin (*Citrus mitis*), grapefruit (*C. grandis*), mandarin (*C. nobilis* var.), and orange (*C. sinensis*) in the Philippine Islands and China, its cause being as yet unknown. It is marked by the oozing out of sap from the bark on the trunk and branches, a putrid sore from 0.5 to 3.5 centimeters in diameter forming. These sores run together and finally girdle the infected tree. The attempts of the tree to heal the spots causes a rough, scaly appearance. Gum pockets form between the wood and cambium. Any fruit present matures prematurely and falls.
- Botrytis citricola** Briozzi. Oranges (*C. sinensis*) and lemons (*C. limonia*) are rotted and mummified by this fungus in Italy and Brazil.
- Cercospora** sp. Circular dark-brown leaf spots on *C. sinensis* in Japan.
- Cercospora fumosa** Penz. Gray irregular areas on the lower leaf surfaces of *Citrus* spp. in Algeria, Argentina, and Italy.
- Cercospora penzigii** Sacc. On leaves of *C. limonia* in Italy.
- Cercospora undulata** (Bern.) Sacc. See *Aegle*.
- Cladosporium brunneo-atrum** McAlp. Minute brown, generally circular, raised spots on fruit, leaves, and young shoots of *C. sinensis* characterize the false melanose of Australia. On the fruit these spots usually run together into long, irregular curves and become black.
- Cladosporium elegans** Penz. Dry, bleached, subcircular to oval spots on orange (*C. sinensis*) leaves in Italy and Australia. This species has been confused in the literature with *C. citri* Mass.
- Cladosporium furfuraceum** McAlp. The lemon-scurf disease, forming dingy-white patches on the surface of lemon (*C. limonia*) fruits in Australia. The discolored areas peel off in small flakes.
- Cladosporium subfusoidium** McAlp. Forms gray-brown scabs on lemons (*C. limonia*), which crack and become covered with darker fruiting patches. Occurs in Australia and the Philippines.
- Coniothecium citri** McAlp. Forms numerous small brown patches with raised brown or black margins on the upper sides of leaves of the orange (*C. sinensis*) in Australia.
- Coniothecium scabrum** McAlp. Causes "black scurf" of citrus (*Citrus* spp.) fruits in Australia and the Union of South Africa. Small black depressions, surrounded by yellowish-green areas appear, which enlarge to form large, sooty-black patches. The black masses become detached in flakes, leaving dirty-gray scurfy depressed areas.
- Coniothyrium cervinum** McAlp. Irregular gray to fawn-colored leaf spots on *C. limonia* in New South Wales.
- Coniothyrium citricolum** McAlp. Grayish-white leaf spots with red-brown margins on *C. sinensis* in New South Wales.
- Corticium koleroga** (Cke.) v. Hoeh. See *Coffea*.
- Corticium salmonicolor** B. and Br. This fungus produces the well-known and widespread "pink disease" of the Tropics, which attacks practically all woody economic plants. It occurs on the trunk and branches, first being noted as dirty-white to pinkish pustules, which push through the dead bark. Following this stage, a layer of mycelium spreads out over the bark, turning to a characteristic pink color, constituting the fruiting areas of the fungus. This layer which may extend for several feet, and even girdle the trunk and one or more limbs, is at first smooth and velvety, but with age becomes roughened and breaks into patches, which fade out to a dirty-white. The bark beneath is at first watery and somewhat darker than normal, soon, however, becoming dried out, and is finally reduced to a dry mass of fiber. The outermost wood layers are also penetrated and killed. The disease has been reported from the following countries: West Indies, including Porto Rico, Trinidad, British Guiana, Surinam, Brazil, Caucasus, Ceylon, India, Java, Guam, Philippines, Borneo, Malaya, Sumatra, Burma, Mauritius, Fiji, Formosa, west Africa, Portuguese India, and the Cameroons. Among the known hosts are *Amherstia nobilis*, *Annona muricata*, *A. squamosa*, *Artocarpus integrifolia*, *Averrhoa* sp., *Bauhinia* sp., *Cajanus indicus*, *Castilla elastica*, *Casuarina equisetifolia*, *Chalcas exotica*, *Cinchona ledgeriana*, *C. succirubra*, *Cinnamomum camphora*, *C. zeylanicum*, *Citrus aurantifolia*, *C. grandis*, *C. hystrix*, *C. limonia*, *C. medica*, *C. micrantha*, *C. nobilis*, *C. sinensis*, *C. webberi*, *Clitoria cajanifolia*, *Coffea arabica*, *C. liberica*, *C. robusta*, *Corchorus* sp., *Duranta*

## CITRUS—Continued.

**Corticium salmonicolor**—Continued.

*plumieri*, *Durio zibethinus*, *Eriobotrya japonica*, *Erythroxyton coca*, *Eucalyptus* sp., *Feronia elephantum*, *Feroniella lucida*, *Ficus* spp., *Gardenia* sp., *Gliricidia maculata*, *Hevea brasiliensis*, *Hibiscus* sp., *Indigofera arrecta*, *Lansium domesticum*, *Mangifera indica*, *Myristica fragrans*, *Photinia japonica*, *Piper* sp., *Pithecolobium dulce*, *Prunus persica*, *Randia macrantha*, *Shorea* sp., *Strobilanthes* sp., *Thea* sp., and *Theobroma cacao*.

**Cytospora citri** May. On *Citrus* sp. in Italy.

**Cytosporina citriperda** Camp. Circular depressed brownish-red, then black, patches on fruit of *C. sinensis* in Italy.

**Dendrophoma valispora** Penz. On lemon (*C. limonia*) leaves in Italy.

**Diaporthe citricola** Rehm. Found in connection with *Citrus* die-back in the Philippines.

**Didymella citri** Noack. Produces a canker on branches and twigs of *C. sinensis* (orange) in Brazil.

**Diplodia cacaicola** P. Henn. See *Theobroma*.

**Diplodia citricola** McAlp. On fruit, leaves, twigs, and branches of *Citrus* spp. in Australia.

**Diplodia destruens** McAlp. Said to cause scabby, dirty-gray patches with red-brown margins on leaves of *C. sinensis* in Australia.

**Dry rot.** See *Nematospora* sp.

**Eutypella citricola** Speg. On branches of *C. sinensis* in Brazil and Argentina.

**Exobasidium citri** Jaec. On young fruit of *C. nobilis* in the Caucasus.

**Fusarium** sp. A blossom end-rot of orange (*C. sinensis*) is produced by this fungus in Porto Rico, marked by a brown staining of the blossom ends and an internal pink rot. Affected fruits turn a deep abnormal yellow and fall prematurely.

**Gloeosporium aurantiorum** West. Anthracnose on *Citrus* spp. in Belgium and Argentina.

**Gloeosporium citri** Mass. This species, reported on leaves of *C. sinensis* from Trinidad and Australia, does not seem to be other than *Colletotrichum gloeosporioides* Penz.

**Gloeosporium citricolum** Hori. Irregular indefinite brown patches on leaves, causing defoliation. Young twigs are also attacked and caused to die-back. Brown anthracnose spots appear on the fruit. Attacks *C. nobilis* var. and *C. sinensis* in Japan. *G. citricolum* Masee is referred to *Colletotrichum gloeosporioides* Penz.

**Gloeosporium follicolum** Nish. Rapidly spreading brown areas around the blossom end of fruit of Satsuma and Mandarin (*C. nobilis* var.) in Japan. These spots become grayish and sunken at the center with dark-brown surrounding areas.

**Gloeosporium hesperidearum** Catt. Anthracnose on leaves of *Citrus* spp. in Italy and Argentina. Referred by Grove to *G. aurantiorum* West.

**Gloeosporium intermedium** Sacc. and var. **subramulosum** Sacc. Anthracnose on *Citrus* spp., *Coryncarpus laevigata*, and *Hoya carnosa* in Italy, France, and Australia. Also referred to *G. aurantiorum* West.

**Gloeosporium spegazzini** Sacc. Anthracnose on *C. limonia* and *C. sinensis* in Brazil and Argentina. Like most of the other *Gloeosporiums* of *Citrus* not definitely distinct from *Colletotrichum gloeosporioides*.

**Hainesia aurantii** P. Henn. On leaves of *C. sinensis* in Costa Rica.

**Hendersonia citri** McAlp. On leaves of *C. aurantium* in Australia.

**Hendersonia socia** McAlp. On leaves and bark of *C. limonia* in Australia.

**Laestadia socia** Penz. On leaves of *C. limonia* in Italy.

**Lasiodiplodia citri** Avena. Produces a fruit rot, die-back of twigs, and yellow to brown leaf spots of *C. sinensis* and *C. limonia* in Brazil. Said to be one of the worst of Brazilian *Citrus* diseases. The fungus is apparently not distinct from *D. cacaicola* P. Henn.

**Leaf drop.** This disease, due to an unknown cause, defoliates Satsumas (*C. nobilis* var. *unshiu*) in Japan, and is considered serious.

**Leptosphaeria citricola** Penz. On *Citrus* sp. in Italy.

**Leptosphaeria coniothyrium** Sacc. On twigs of *C. limonia* and *C. sinensis* in Brazil.

**Leptosphaeria diana** Sacc. and Speg. On leaves of *C. aurantium* in Italy.

**Macrophoma aurantii** Scalia. On leaves of *C. sinensis* in Italy.

**Macrophoma mantegazziana** (Penz.) Berl. and Vogl. On leaves of *Citrus* spp. in Italy and Ceylon.

**Meliola butleri** Syd. Circular black superficial fungus patches on leaves of *C. medica* and *C. grandis* in India.

**Meliola citricola** Syd. Superficial black fungus patches on leaves of *C. grandis*, *C. nobilis*, and *C. sinensis* in China.

**Microthyrium citri** Penz. On leaves of *C. limonia* in Italy.

**Mycosphaerella citricola** McAlp. On leaves of *C. limonia* in Ceylon and Australia.

**Mycosphaerella gibelliana** Pass. Subcircular, brown leaf spots on *Citrus* spp. in Italy, Portugal, Spain, and Austria.

**Mycosphaerella horii** K. Hara. Circular to irregular definite reddish-brown to gray leaf spots with raised blackish-brown margins on *C. sinensis* and *C. nobilis* in Japan. The "small round brown spot disease."

**Mycosphaerella inflata** Penz. On living branches of *C. aurantium* in Italy.

**Mycosphaerella loefgreni** Noack. Small, circular to irregular, brown to dirty-white spots on leaves, fruit, and twigs of *C. sinensis* in Brazil. The spots on the twigs are raised, but sunken on leaves and fruit. The disease may cause premature fall of leaves and fruit.

**Nematospora** sp. This fungus is reported as the cause of the "dry-rot" disease of fruit of *C. sinensis*, *C. aurantium*, and *C. nobilis* in the Philippine Islands, China, Japan, Formosa, and possibly Barbados. There are no external symptoms. Internally infected fruits are more or less dry and free of juice, the sections becoming atrophied and falling apart. The pulp at first has an unpleasant sour taste, but soon becomes dry and tasteless.

**Nothopatella lecanidium** (Speg.) Sacc. On trunks and branches of *C. sinensis* in Brazil.

**Oidium** sp. Forms a white powdery layer on leaves and twigs, causing yellowing and premature fall of leaves and die-back of twigs of *Citrus* spp. in India and Ceylon. In the latter country said to be so serious as to prevent the growing of *Citrus* on any scale. *O. tingitaninum* Carter, described from California, is probably the same species.

**Oospora aurantii** Petch. Causes a rotting of fruit of *C. sinensis* in Ceylon.

**Ovularia aurantii** McAlp. Erumpent, effused, grayish-green patches on fruit of *C. sinensis* in Australia and Ceylon.

**Ovularia citri** Br. and Farn. Attacks orange fruits (*C. sinensis*) in Italy causing a disease known as "white rust."

**Phoma aurantiorum** (Rabenh.) Sacc. On branches of *Citrus* sp. in middle Europe.

**Phoma citricarpa** McAlp. The black-spot disease causes a spotting of fruit of *C. limonia*, *C. maxima*, *C. mitis*, *C. nobilis*, and *C. sinensis* in China, Australia, and New Zealand. The spots are small, irregular, from 1 to 9 millimeters in diameter, reddish-brown at first, later much darker and sunken.

## CITRUS—Continued.

- Phoma flaccida** McAlp. Brown, apical leaf spots on *C. sinensis* in Australia.
- Phoma omnivora** McAlp. This fungus forms irregular gray to black scabby patches on the fruit and leaves. Leaves and twigs die-back, gray blotches appearing on the latter. Dark-brown cankers occur on the roots. The hosts are *C. limonia*, *C. medica*, and *C. sinensis* in Australia and Ceylon.
- Phoma stigmaea** Dur. and Mont. On leaves of *C. aurantium* in Algeria.
- Phomopsis caribaea** Horne. Causes a stem-end rot of *C. grandis* in the Isle of Pines. Probably not distinct from *P. citri* Fawc.
- Phyllosticta arethusa** Bub. On leaves of *C. aurantium* in Austria.
- Phyllosticta beltranii** Penz. On leaves of *C. limonia* in Italy.
- Phyllosticta circumsepta** Sacc. On fruit of *C. nobilis* in the Philippines.
- Phyllosticta citricola** Hori. Circular to irregular brown leaf spots on *C. nobilis* and *C. sinensis* in Japan.
- Phyllosticta deliciosa** Pass. Brown leaf spots with raised margins on *Citrus* sp. in Italy.
- Phyllosticta disciformis** Penz. and var. **brasilienis** Speg. Circular to irregular ashen-white leaf spots with dull-brown margins on *Citrus* spp. and *Atalantia buxifolia* in Russia, Ceylon, Italy, and Brazil.
- Phyllosticta fuliginosa** Massal. Irregular olivaceous to dull-brown leaf spots on *Citrus* sp. in Italy.
- Phyllosticta hesperidearum** Penz. Small, irregular, brown, then yellow, and finally ashen leaf spots, and dull-brown corky areas on twigs of *C. limonia* and *C. sinensis* in Brazil, Algeria, and Australia.
- Phyllosticta lenticularis** Pass. Leaf spot on *Citrus* sp. in Portugal.
- Phyllosticta longispora** McAlp. Brown to gray apical and marginal leaf spots on *C. sinensis* in Australia.
- Phyllosticta marginalis** Penz. Irregular white spots with brown margins on leaves of *C. medica* in Italy.
- Phyllosticta micrococcoides** Penz. Dark-brown, irregular spots on young leaves of *Citrus* sp. in Italy.
- Phyllosticta scabiosa** McAlp. On leaves of *C. limonia* in Australia.
- Phytophthora** sp. Causes a rot of *Citrus* fruits in Porto Rico.
- Phytophthora citri** Ven. Causes a soft rot of fruit of *C. medica* in India.
- Pleospora batumensis** Naou. On petioles of *C. sinensis* in south Russia.
- Pleospora citrorum** Sacc. On *Citrus* sp. in Italy and Argentina.
- Pleospora disrupta** McAlp. Disorganizes the tissues of orange (*C. sinensis*) leaves in Australia.
- Pleospora hesperidearum** Catt. Causing a scabbing of fruit of *Citrus* sp. in Italy and Uruguay.
- Polyporus caryophylli** Racib. See *Cinnamomum*.
- Pyrenochaeta destructiva** McAlp. Circular to oval gray leaf spots with brown margins on *C. aurantium* and *C. grandis* in Australia.
- Ramularia citri** Penz. On leaves of *C. sinensis* in Italy.
- Ramularia scabiosa** McAlp. and Tryon. Circular definite light-brown, then nearly black, raised spots on leaf blades and petioles of *C. limonia* and *C. sinensis* in Australia.
- Ramularia undulata** Bern. On leaves of *Citrus* sp. and *Aglaia odorata* in Java.
- Rhabdospora flexuosa** (Penz.) Sacc. On bark of *C. sinensis* in Italy and Brazil.
- Rhynchodiplodia citri** Br. and Farn. This fungus, reported as the cause of a serious lemon (*C. limonia*) disease in Italy and Brazil, is probably the same again as *D. cacaoicola* P. Henn.
- Rosellinia** sp. An undetermined species of *Rosellinia* causes a root rot of *Citrus* spp., *Erythrina* sp., *Theobroma cacao*, and possibly other economic plants in the West Indies. Diseased trees die-back slowly, suckering freely, and finally dying. The rot in the roots and crowns resembles that produced by *R. pepo*. A *Rosellinia* sp. is reported as the cause of a root disease of *C. sinensis* in Spain and the Balearic Islands.
- Rosellinia bunodes** B. and Br. The "black root disease" attacks and destroys a wide range of tropical plants, including *Acalypha* sp., *Artocarpus incisa*, *Cajanus indicus*, *Castilla elastica*, *Cinnamomum camphora*, *Citrus* spp., *Coffea arabica*, *Dryobalanops aromatica*, *Eugenia jambos*, *Ficus dubia*, *Hevea brasiliensis*, *Hibiscus rosa-sinensis*, *Mangifera indica*, *Maranta arundinacea*, *Miconia* sp., *Panax plumatum*, *Persea gratissima*, *Petiveria alliacea*, *Piper nigrum*, *Sterculia caribaea*, and *Theobroma cacao* in the West Indies, Trinidad, India, Malaya, Java, and Ceylon. The disease is apparently most prevalent and most serious in the West Indies, the *Rosellinia* root disease of tea and other woody plants in the oriental tropics being attributed for the most part to other species.
- The fungus attacks the roots and crowns of the hosts, working through the bark to the wood. Black fungus strands or rhizomorphs appear on the surface and may grow together to form a dense layer. From the lower surface black strands, with a white core, grow out, penetrating the bark and finally the wood. The fungus fruits on the surface layer which extends up the trunk of the host a short distance, a conidial stage of black bristle-like stalks (*Graphium*) appears first, followed by the black globose carbonaceous perithecia. Infected plants die back gradually as the disease progresses, until girdled, when wilting and death occurs.
- What is thought to be the same species has been found in the British West Indies, causing a root disease and rot of tubers or rhizomes of arrowroot (*Maranta*), *Colocasia*, *Dioscorea*, and other herbaceous plants. The external rhizomorphs are lacking, but the internal conditions are similar to those of woody plants attacked by *R. bunodes*.
- Rosellinia pepo** Pat. The root disease attributed to this species is called white root rot and attacks *Cajanus indicus*, *Citrus* spp., *Coffea arabica*, *Hymenaea courbaril*, *Manihot utilisima*, *Theobroma cacao*, and other woody plants in Porto Rico, Martinique, Guadeloupe, Trinidad, and the British Antilles. Herbaceous plants, such as dasheens (*Colocasia*), *Musa*, and *Canavalia*, are also attacked at times.
- The fungus forms an irregular gray, then black, coating over infected roots. White strands grow out from this layer through the bark and form another much-branched white fungus layer or "fan" between the wood and the bark. The white strands penetrate the wood along the medullary rays. The fungus is black only when exposed. The effect on diseased plants is similar to that produced by *R. bunodes*.
- Septobasidium acaciae** Saw. See *Acacia*.
- Septoria arethusa** Penz. Ochraceous leaf and fruit spots on *C. limonia* and *Citrus* sp. in Brazil, Argentina, Italy, France, and Algeria. *Podocarpus* sp. has also been given as a host, but this reference seems doubtful.
- Septoria cattanei** Thuem. Small ochraceous leaf spots on *C. medica* in India and the Canary Islands.
- Septoria cinerescens** (Dur. and Mont.) Sacc. On leaves of *C. aurantium* in Algeria.
- Septoria citri** Pass. On leaves of *Citrus* sp. in Italy, Spain, Portugal, and France.
- Septoria depressa** McAlp. Brown to black circular to irregular sunken spots on fruit of *C. limonia*, *C. medica*, and *C. sinensis* in Australia.
- Septoria flaccescens** McAlp. Apical brown areas on leaves of *C. sinensis* in Australia.
- Septoria glaucescens** Trab. "Spot disease" of mandarin (*C. nobilis* var.) in French North Africa.

**CITRUS**—Continued.

- Septoria limonum** Pass. On leaves and fruit of *Citrus* spp. in Italy.  
**Septoria sicula** Penz. On leaves of *C. limonia* in Italy.  
**Septoria tibia** Penz. Brown leaf spots on *C. limonia* in Italy.  
**Septoria westraliensis** McAlp. Circular to irregular ashen-gray leaf spots with brown margins on *C. sinensis* in Australia.  
**Septoriopsis citri** Paul. and Gz. Frag. On fruit of *C. sinensis* in Spain.  
**Sphaeropsis pseudo-diplodia** (Fckl.) Delacr. On twigs and branches of *Citrus* sp. in Brazil. Apparently not distinct from *Diplodia cacaicola*.  
**Sphaeropsis tumefaciens** Hedges. Knots or galls are produced on the twigs and branches, varying from 0.37 to 0.66 inch in diameter. A witches'-broom effect may appear, but ultimately the portion above the gall dies. The disease occurs on *C. aurantifolia* and *C. sinensis* in Cuba, British Guiana, and Jamaica.  
**Sphaerostilbe repens** B. and Br. See Hevea.  
**Sporidesmium griseum** McAlp. Dirty-gray scabby patches, cracking and breaking into irregular areas on fruit and leaves of *C. limonia* and *C. sinensis* in Australia and New Zealand.  
**Trichoseptoria alpei** Cav. Causes a spotting of fruit of *C. limonia* in Italy.
- CLADRASTIS.** YELLOWWOOD. Trees grown for their white flowers and handsome foliage.  
**Ascochyta cladrastidis** Kab. and Bub. On *C. lutea* in Bohemia.  
**Cercospora cladrastidis** Jacz. Subcircular brown leaf spots with red margins on *C. (Maackia) amurensis* in Japan and Siberia.  
**Uromyces amurensis** Kom. Yellow and cinnamon-brown to black rust pustules on leaves of *C. (Maackia) amurensis* in Siberia.  
**Uromyces cladrastidis** Kusano. Black rust sori on leaves of *C. shikokiana* in Japan.  
**Uromyces shikokianus** Kusano. Yellow-brown to black rust sori on leaves of *C. shikokiana* in Japan.
- CLARKIA.** Flower-garden annuals.  
**Phytophthora parasitica** Dastur. See Ricinus.
- CLAYTONIA.** SPRING BEAUTY. Succulent herbs cultivated for their flowers.  
**Puccinia claytoniae** Thuem. Rust on leaves of *C. arctica* in Siberia.
- LEMATIS.** Climbing or erect perennial garden plants with handsome, showy flowers.  
**Acidium englerianum** P. Henn. and Lind. Leaf rust on *C. wightiana* in Abyssinia and Tanganyika.  
**Acidium orbiculare** Barcl. Rust causing hypertrophy of stems of *C. grata*, *C. orientalis*, and *C. puberula* in India.  
**Acidium otagense** Limds. A rust deforming the leaves, peduncles, and flowers of *C. colensoi*, *C. hexapetala*, *C. indivisa*, and *C. marata* in New Zealand. Possibly a stage of *Puccinia clavata*.  
**Ascochyta indusiata** Bres. Subcircular to ovoid dull-brown leaf spots on *C. recta* in France and Germany.  
**Ascochyta vitalba** Br. and Har. On leaves and branches of *C. vitalba* in France.  
**Coleosporium clematidis** Barcl. Yellow to orange rust pustules on leaves of *Clematis* spp. in China, Japan, Siberia, India, Australia, Russia, Abyssinia, central and south Africa, and Australia.  
**Coleosporium clematidis-apiifoliae** Diet. Yellow rust pustules on leaves of *C. apiifolia* and *C. parviloba* in Japan and China.  
**Coleosporium elongatum** Syd. Leaf rust on *C. hedysarifolia* in Japan.  
**Erysiphe taurica** Lév. See Althaea.  
**Marsonia clematidis** All. Large irregular indefinite dull-brown zoned leaf spots on *C. vitalba* and *Thalictrum minus* in Germany.  
**Phyllosticta bacteriosperma** Pass. Irregular angular dull-brown leaf spots on *C. vitalba* in Italy.  
**Phyllosticta beguinotiana** Sacc. Small subcircular light-brown, then white, leaf spots on *C. flammula* in Italy.  
**Phyllosticta corrodens** Pass. Indefinite gray leaf spots with dull-brown margins on *C. vitalba* in Italy and Denmark.  
**Phyllosticta intermedia** Allesch. Small subcircular to angular whitish leaf spots with brown margins on *C. recta* in France and Germany.  
**Phyllosticta vitalba** Cke. Leaf spots on *Clematis* sp. in Great Britain.  
**Physalospora disseminata** Sacc. On leaves of *C. glauca* in Siberia.  
**Puccinia clavata** Syd. Brown rust pustules on small sunken brown leaf spots on *C. hexapetala* in New Zealand.  
**Puccinia exhausta** Diet. Brown powdery rust pustules on leaves of *C. heracleaefolia*, *C. javanica*, *C. taiwaniana*, and *C. tuberosa* in Java, Japan, and the Philippines.  
**Puccinia insidiosa** Berk. Leaf rust on *C. nutans* in India.  
**Puccinia wattiana** Barcl. Brown rust pustules on sunken black leaf spots on *C. buchianiana*, *C. gouriana*, and *C. puberula* in India.  
**Septoria aciduicola** Pat. On leaves of *C. cirrhosa* in Tunis.  
**Septoria cirrhosae** Sacc. Circular dull-yellow leaf spots with dark-purple margins on *C. cirrhosa* in Sicily.  
**Septoria clematidis** Rob. and Desm. Circular to angular grayish leaf spots with dull-brown margins on *C. flammula*, *C. glauca*, and *C. vitalba* in Ecuador, Siberia, and Europe.  
**Septoria clematis-flammulae** Roum. Small subcircular gray, then white, leaf spots with brown margins on *C. flammula* in Dalmatia, France, and Italy.  
**Septoria clematis-rectae** Sacc. Grayish-white leaf spots on *C. recta* in Italy.  
**Septoria viticellae** Pass. Leaf spots on *C. viticella* in Italy.
- CLEOME.** Spiderflower. Mostly weedy herbs.  
**Cercospora conspicua** Earle. Leaf spot on *C. pentaphylla* in Porto Rico.
- CLERODENDRUM.** GLORY BOWER. Shrubs, trees, or climbers grown for their showy flowers.  
**Acidium clerodendri** P. Henn. Leaf rust on *C. calamatosum*, *C. cyrtophyllum*, *C. fragrans*, *C. intermedium*, and *C. paniculatum* in Java, Japan, and the Philippines.  
**Acidium clerodendricola** P. Henn. Leaf rust on *C. buchnerum* in Tanganyika.  
**Acidium clerodendronis** P. Henn. Leaf rust on *Clerodendrum* sp. in Indo-China.  
**Cercospora bakeri** Syd. On leaves of *C. intermedium* in the Philippines.  
**Cercospora clerodendri** Miy. Leaf spot on *Clerodendrum* sp. in China.  
**Chrysomyxa peregrina** Syd. and Butl. Yellow-brown rust pustules on leaves of *Clerodendrum* sp. in India.  
**Coleosporium clerodendri** Diet. Yellow to orange rust pustules on lower leaf surfaces of *C. cyrtophyllum*, *C. fragrans*, and *C. trichotomum* in Java, Japan, and Formosa.  
**Hemileia scholzii** Syd. Yellow powdery rust pustules on leaves of *Clerodendrum* sp. in central and south Africa and Portuguese East Africa.  
**Physalospora clerodendri** Syd. On leaves of *Clerodendrum* sp. in the Congo.  
**Puccinia erebia** Syd. Leaf rust on *C. commersonii* and *C. minahassae* in the Philippines.  
**Synchytrium collapsus** Syd. and Butl. Small swellings on both leaf surfaces of *Clerodendrum* sp. in India.  
**Uredo clerodendricola** P. Henn. Leaf rust on *C. inerme* in Java, Ceylon, and China.

- CLETHRA.** WHITE ALDER. Shrubs or small trees grown for their spikes of white fragrant flowers.  
**Phyllachora rubefaciens** Rehm. Shiny black stromata on red-brown leaf spots on *C. laevigata* in Brazil.  
**Phyllosticta clethrae** Syd. On leaves of *C. scabra* in Austria.  
**Pucciniastrum kusanoi** Diet. Leaf rust on *C. barbinervis* in Japan.
- CLIANTHUS.** PARROTBEAK. Sometimes called Glory pea. Tender half trailing shrubs with large showy flowers.  
**Ascochyta clianthi** Tassi. Dull-brown to whitish irregular leaf spots on *C. arboreus* in Italy.
- CLINTONIA.** Low-growing liliaceous herbs.  
**Puccinia clintoniae-udensis** Bub. Cinnamon-colored rust pustules on the upper surfaces of leaves of *C. udensis* in Amur Province, Siberia.
- CLITORIA.** BUTTERFLY PEA. Hardy perennial vines with pealike flowers.  
**Cercospora pantoleuca** Syd. Dark-brown blotches on leaves of *C. ternatea* in India and the Philippines.  
**Cercospora ternatea** Petch. Circular black spots with gray centers, which become gray with yellow or pale-greenish margins, on leaves and pods of *C. ternatea* in Ceylon.  
**Corticium salmonicolor** B. and Br. See *Citrus*.  
**Melasmia juruana** P. Henn. Black shiny stromata on leaves of *Clitoria* sp. in Brazil.  
**Uromyces clitoriae** Arth. Powdery brown rust pustules on lower leaf surfaces of *C. mexicana* in Mexico.  
**Uromyces neurocarpi** Diet. Powdery brown rust pustules on leaves of *C. cajanifolia* and *C. rubiginosa* in Porto Rico, Cuba, Jamaica, Trinidad, and Brazil.  
**Uromyces yurimaguasensis** P. Henn. Brown rust pustules in circular dull-brown leaf spots on *C. arboreas* and *Clitoria* sp. in Peru and Panama.
- CLIVIA (Himantophyllum).** KAFIR LILY. Tender bulbous plants with evergreen foliage and showy red or red and yellow flowers.  
**Ascochyta cliviae** Magnaghi. On leaves of *C. nobilis* in Italy.  
**Colletotrichum cliviae** Oud. Small irregular red to brown leaf spots on *C. miniata* and *Clivia* sp. in Bohemia, Holland, and France.  
**Physalospora himantophylli** Gz. Frag. On leaves of *Himantophyllum (Clivia) miniata* in Spain.
- CLUSIA.** Tropical trees used to some extent for shade.  
**Coccomyces clusiae** (Lév.) Sacc. Black fruiting bodies on leaves of *Clusia* sp. in Costa Rica and northern South America.  
**Phyllosticta clusiae** Allesch. On leaves of *Clusia* sp. in Brazil.  
**Septogloeum clusiae** Karst. and Har. On leaves of *Clusia* sp. in Colombia.  
**Uredo clusiae** Arth. Dark-brown rust pustules on leaves of *C. rosea* in Porto Rico.
- CLYTOSTOMA.** Ornamental vines, grown for their beautiful flowers.  
**Phyllachora amphigena** Speg. See *Bignonia*.
- COBAEA.** Climbers grown for their large bell-shaped flowers.  
**Phyllosticta cobaeae** Tassi. Leaf spot on *C. scandens* in Italy.  
**Septoria oligocarpa** Nannizzi. Small, subcircular leaf spots with narrow purple margins on *C. scandens* in Italy.
- COCCOLOBA.** SEA GRAPE. Tropical shrubs and trees, some species grown for their fruit or foliage.  
**Cercospora bicolor** Wint. Leaf spots on *C. sagittaeifolia* in Portugal.  
**Fabraea coccolobae** P. Henn. Effuse dull-brown areas on leaves of *Coccoloba* sp. in Brazil.  
**Phyllachora coccolobae** Speg. Black stromata on leaves of *Coccoloba* sp. in Argentina.  
**Phyllachora simplex** Starb. Small scattered black stromata on leaves of *C. laurifolia* and *Coccoloba* sp. in Paraguay and Porto Rico.  
**Phyllosticta coccolobae** E. and E. Circular, purplish leaf spots, 3 to 5 millimeters in diameter on *C. wifera* in Porto Rico and the Bahamas.  
**Physalospora coccolobae** Rehm. Circular yellowish leaf spots on *Coccoloba* sp. in Brazil.  
**Uredo coccolobae** P. Henn. Powdery light-brown rust pustules on leaves of *C. populifolia* and *C. wifera* in Brazil, Porto Rico, and Cuba. Also reported from Florida.
- COCCOTHEINAX.** See *Palmae*.
- COCCULUS.** SNAIL SEED. Shrubs grown for their handsome foliage and ornamental fruit.  
**Phomopsis cocculi** Luigi. Light-ashen spots with irregular black borders on leaves of *C. laurifolius* in Italy.  
**Phyllosticta cocculi** Speg. Leaf spot on *C. laurifolius* in Argentina.  
**Phyllosticta thunbergii** Wint. Large, indefinite, irregular, pale-yellow leaf spots on *C. thunbergii* in Japan.
- COCHLEARIA.** Small fleshy seaside herbs.  
**Puccinia cochleariae** Lindr. Powdery black rust pustules on leaf blades and petioles of *C. fenestrata*, *C. groenlandica*, and *C. pyrenaica* in France and Greenland.
- COCOS.** Coconut. See *Palmae*.
- CODIAEUM (PHYLLAUREA Ag.).** Tropical shrubs grown for their variegated foliage.  
**Ascochyta banosensis** Syd. On leaves of *C. variegatum* in the Philippines.  
**Corticium koleroga** (Cke.) v. Hoeh. See *Coffea*.  
**Corticium stevensii** Burt. Leaf blight of *C. variegatum* in Trinidad.  
**Fomes lamaoensis** Murr. See *Hevea*.  
**Gloeosporium sorauerianum** Allesch. Large irregular indefinite yellow to ashen leaf spots on *C. variegatum* in France, Holland, and Germany.  
**Laestadia minuscula** (Lév.) Sacc. On leaves of *C. variegatum* in Java.  
**Phyllosticta codiaei** Diet. On leaves of *Codiaeum* sp. in the Philippines.  
**Phyllosticta reyesii** (Sacc.) Yates. On leaves of *C. variegatum* in the Philippines.
- CODONOPSIS.** Twining or decumbent perennials with showy flowers.  
**Coleosporium horianum** P. Henn. Golden-yellow to brown rust pustules on leaves of *C. lanceolata* in Japan.
- CODONORCHIS.** See *Orchidaceae*.
- COELOGLOSSUM.** See *Orchidaceae*.
- COELOGYNE.** See *Orchidaceae*.
- COFFEA.** COFFEE. Tropical shrubs or small trees.  
**Anthostomella coffeae** Delacr. Found in connection with die-back of twigs of *C. arabica* in Mexico.  
**Ascochyta coffeae** P. Henn. Subcircular brown leaf spots with darker margins on *C. arabica* in Brazil.  
**Ascospora coffeae** Faber. A wound parasite destroying large areas of bark of *C. robusta* trees in Java. Leaves of infected plants turn yellow and fall.  
**Atichia millardeti** Racib. See *Cinnamomum*.  
**Cephalosporium** sp. Concentrically ringed brown leaf spots on *C. arabica* in Porto Rico.

**COFFEA**—Continued.

- Cercospora coffeae** Zimm. Subcircular dark-brown leaf spots on *C. arabica*, *C. laurina*, *C. robusta*, and *C. stenophylla* in Malaya and Tanganyika.
- Cercospora coffeicola** B. and C. Circular brown leaf spots, 6 to 10 millimeters in diameter, sometimes causing defoliation. Young twigs are also attacked to some extent. The disease is serious on the fruit, appearing first as small brown spots which enlarge to cover about half the fruit, generally the upper half. The diseased pulp turns black and dries down, adhering to the parchment, making the process of preparation more difficult, and lowering the grade of the final product. The disease occurs in practically all coffee-growing countries, including Porto Rico, Cuba, Jamaica, Trinidad, Surinam, Guatemala, Costa Rica, Brazil, Hawaii, Dutch East Indies, India, Java, New Caledonia, Queensland, the Philippines, and central Africa. The hosts are *C. arabica* and *C. buxobensis*.
- Cercospora herreraana** Farn. Circular chestnut-brown leaf spots with dark-purple margins on leaves and fruit of *C. arabica* in Mexico. Probably the same as the preceding species.
- Cobweb disease.** A Javan disease similar to the "koleroga" disease of coffee (*C. arabica*), but due to a different fungus. White, thick, branching fungus threads grow along the surface of twigs and leaves. Infected leaves turn yellow, wilt, and die, accompanied by die-back of the twigs.
- Colletotrichum coffeanum** Noack. Circular to irregular, large, brown, and finally gray spots on leaves of *C. arabica* in Porto Rico, Costa Rica, Brazil, central Africa, Madagascar, Reunion, Java, and India. A die-back of twigs is often associated with this or closely related species. The berries are also attacked, the spotting reducing the grade of infected material. *Colletotrichum incarnatum* Zimm. from Ceylon is probably not distinct.
- Corticium koleroga** (Cke.) v. Hoeh. (*Pellicularia koleroga* Cke.) The "koleroga" disease of coffee (*C. arabica*) occurs in Porto Rico, Cuba, Jamaica, Surinam, Venezuela, Guatemala, India, Java, Malaya, Queensland, Congo, and possibly in Brazil. The fungus threads grow up along the undersides of the twigs and out onto the lower leaf surfaces. The leaves turn black, wither up or drop, or remain suspended hanging by the fungus filaments. These strands are at first white and finally deep brown. Diseased twigs may die back for considerable distances. The berries are attacked to some extent, infected berries showing blackened grains. Additional hosts are *Citrus* spp., *Codiaeum variegatum*, *Cucumis anguria*, *Garcinia mangostana*, *Hevea brasiliensis*, and *Luffa aegyptiaca*.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Dietyothyriella mucosa** Syd. Small, black, superficial fruiting bodies on *C. liberica* in Indo-China and the Philippines.
- Diplodia cacaofcola** P. Henn. See *Theobroma*.
- Euryachora coffeicola** Averna. Black stromata on leaves of *C. arabica* in Brazil.
- Euryachora liberica** Oud. A doubtful species on branches of *C. arabica* and *C. liberica* in Java.
- Fomes lamaoensis** Murr. See *Hevea*.
- Fomes lignosus** Klotzsch. See *Hevea*.
- Fusarium coffeicola** P. Henn. Found in connection with die back of *C. arabica* and *C. liberica* in central Africa.
- Gloeosporium coffeanum** Delacr. Anthracnose on leaves of *C. arabica* in Reunion and Ceylon. Probably the same as *Colletotrichum coffeanum* Noack.
- Gloeosporium coffeicolum** Tass. On *C. arabica* in Italy. Not distinct from *G. cingulata*.
- Helminthosporium coffeae** Mass. On leaves of *C. liberica* in Gold Coast Colony.
- Helminthosporium ubangiense** P. Henn. On leaves of *Coffea* sp. in tropical Africa.
- Hemileia vastatrix** B. and Br. This very destructive rust has destroyed the coffee industry of some countries and made it unprofitable in others. The characteristic yellow rust pustules develop during wet weather on the lower leaf surfaces, and more rarely on fruits and tips of young branches. Small yellow spots first appear, which increase in size, merge, and finally involve the entire leaf blade, which becomes brown and falls. Plants of all ages are attacked, defoliated, and killed within a season or two. *C. arabica* is very susceptible. Other species (*C. bengalensis*, *C. laurina*, *C. liberica*, *C. robusta*, *C. trawan-korensis*) vary in their resistance, some varieties being practically immune. The rust occurs in Burma, Ceylon, India, Indo-China, China, Malaya, Philippines, Union of South Africa, east Africa, Malacca, Sumatra, Java, Fiji, Samoa, Mauritius, Reunion, New Hebrides, New Caledonia, Gold Coast, and Madagascar. The disease does not occur in the western Tropics.
- Laestadia coffeicola** Speg. Circular whitish leaf spots on *C. arabica* in Costa Rica.
- Leptosphaeria coffeigena** (B. and C.) Sacc. On leaves of *C. arabica* in Brazil.
- Leptosphaeria tonduzi** Speg. On leaves of *Coffea* sp. in Costa Rica.
- Metasphaeria bifoveolata** Speg. On *C. arabica* in Costa Rica.
- Mosaic** (*Phloem necrosis*). This disease, apparently of the group of infectious mosaics, attacks *C. liberica*, and occasionally *C. arabica*, in Surinam. In the acute form the leaves wilt rapidly and hang down. The roots become black and the entire plant is dead within a few weeks. In other cases the disease progresses more slowly. The leaves turn pale-green, then yellow, and finally fall, being replaced by small abnormal ones.
- Muratella coffeae** Bunting. Attacks *C. liberica* on Gold Coast Colony.
- Mycosphaerella coffeae** Noack. On leaves of *C. arabica* in Brazil.
- Mycosphaerella coffeicola** Cke. Subcircular, brown leaf spots with narrow gray margins on *C. arabica* in Jamaica, Ecuador, Venezuela, Brazil, Uganda, and India.
- Ophionectria foliicola** A. Zimm. On leaves of *C. liberica* in Java.
- Phoma coffeicola** Tassi. On branches and twigs of *C. arabica* in Italy.
- Phyllosticta camoensis** Delacr. White thickened leaf spots on *C. camoensis* in tropical Africa.
- Phyllosticta coffeicola** Delacr. Circular whitish leaf spots on *C. arabica* in Malaya, Java, Mexico, Costa Rica, and Brazil. Probably synonymous with the Spegazzini species, which has priority.
- Phyllosticta coffeicola** Speg. On leaves of *Coffea* sp. in Costa Rica and Brazil.
- Phyllosticta usteri** Speg. Circular, then confluent, leaf spots on *C. arabica* and *C. robusta* in Ceylon and Brazil.
- Placosphaeria citricola** Berl. and Roum. On leaves of *C. arabica* in Indo-China.
- Pleonectria coffeicola** A. Zimm. On leaves of *C. liberica* in Java.
- Ramularia goldiana** Sacc. Subcircular zonate spots on leaves and stems of *C. arabica* in Brazil.
- Rhizoctonia lamellifera** Small. See *Grevillea*.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Rosellinia pepo** Pat. See *Citrus*.
- Rostrella coffeae** Zimm. On *C. arabica* in Guatemala.
- Saccardinula tahitensis** Pat. On leaves of *Coffea* sp. in Tahiti.
- Sclerotium** sp. Causes a damping-off of coffee (*C. arabica*) seedlings in the Philippines.
- Sclerotium coffeicolum** Stahel. The "coremium" or sclerotial disease of *C. abeocuta*, *C. arabica*, *C. excelsa*, and *C. liberica* in Surinam is characterized by dark-brown concentric spots on leaves and fruits. Small, greenish-brown or orange-brown sclerotia develop on the diseased areas.
- Septoria berkeleyi** Sacc. and Trav. Subcircular pale-brown leaf spots on *C. arabica* in South America.

**COFFEA**—Continued.

**Septoria coffeae** Wakef. On leaves of *C. arabica* in Uganda.

**Septoria coffeicola** P. Henn. Circular white leaf spots with dark-brown margins on *C. liberica* in Kamerun.

**Stilbella flavida** (Cke.) Lind. Circular dark-brown and finally light-brown leaf spots on *C. arabica*, *C. excelsa*, *C. perrieri*, and *C. robusta* in Porto Rico, Jamaica, Cuba, British West Indies, Central America, Mexico, Trinidad, Ecuador, Venezuela, Colombia, British Guiana, Dutch Guiana, and Java. Other plants, including *Commelina* sp., *Bryophyllum pinnatum*, *Inga vera*, and *Mangifera indica*, growing under or adjacent to coffee are attacked. The spots are from 6 to 12 millimeters in diameter, and are often so numerous as to destroy the leaves. On the upper surfaces of the spots and, to some extent, on the lower surfaces yellow, hairlike projections, 1 to 4 millimeters long, with a small red terminal spore mass occur. Young twigs and the fruit are also attacked to some extent. The disease may be sufficiently serious to reduce yields as much as 75 per cent.

**Trachysphaera fructigena** Tabor. and Bunting. The "mealy pod" disease attacks *C. liberica* and *Theobroma cacao* in Gold Coast Colony. Coffee fruits of all ages are attacked, but the young berries are most subject to the disease. Their development is stopped and they shrivel up and become hard, a dark purplish-brown discoloration appearing, which is later covered by a whitish or pinkish-brown mealy incrustation. On cacao pods small brown areas appear which increase rapidly until the entire fruit is involved. The same mealy layer ultimately appears.

**Venturia coffeicola** Averna-Sacca. On leaves of *C. arabica* in Brazil.

**COIX.** JOB'S-TEARS. Tall, broad-leaved grasses with beadlike inflorescences.

**Phyllachora coicis** P. Henn. Shiny black elongate stromata on lower leaf surfaces of *C. agrestis* and *C. lachryma-jobi* in India, Indo-China, and the Philippines.

**Sclerospora maydis** (Rac.) Butl. See Zea.

**Uredo operta** Syd. and Butl. Straw-colored rust sori on leaves of *C. lachryma-jobi* in India and Ceylon.

**Ustilago coicis** Bref. Grain transformed into a black mass of smut spores, surrounded by a membrane and confined within the glumes. On *C. agrestis* and *C. lachryma-jobi* in India, Java, and Japan.

**COLCHICUM.** AUTUMN CROCUS. Autumn-flowering bulbous plants.

**Ascochyta juellii** Bub. Round or elliptical leaf spots on *C. autumnale* in Austria.

**Cladochytrium mauryi** Hariot. Small brown spots on leaves of *C. autumnale* in France.

**Cylindrosporium colchici** Pass. On leaves of *C. alpinum* and *C. officinale* in France and Holland.

**Perisporium colchici** Rbh. Black superficial fungus on leaves of *C. autumnale* in Germany.

**Phyllosticta latemarensis** Kab. and Bub. Small irregular black, often confluent, leaf spots on *C. autumnale* in Italy.

**Septoria colchici** Pass. Long light-brown spots on leaves of *C. alpinum* and *C. officinale* in France, Italy, and Germany.

**Septoria gallica** Sacc. and Syd. Oval to elongate, often confluent, spots on leaves of *C. autumnale* and *C. officinale* in France and Austria.

**Urocystis colchici** (Schlecht) Rab. Long rows of black, powdery streaks occur on the leaves of *C. autumnale*, *Allium cepa*, *A. magicum*, *A. rotundum*, *Bulbocodium vernum*, *Muscari comosum*, *M. racemosum*, *Narcissus* spp., *Paris quadrifolia*, *Scilla bifolia*, and *Tulipa* spp., in England, Yugoslavia, Spain, and Dalmatia. A smut referred to this species occurs to a limited extent in the United States on *Polygonatum* spp. and *Vagnera* spp., but if the European and American forms are conspecific different strains are most certainly involved.

**Uromyces colchici** Mass. Brown powdery rust pustules on both leaf surfaces of *C. autumnale*, *C. bavaricum*, *C. speciosum*, and *C. spectabilis* in Great Britain.

**COLEOSANTHUS.** See Brickellia.**COLEUS.** Showy-leaved herbs used as bedding plants.

**Aphelenchus olesistus** Ritz. Bos. See Begonia.

**Meliola cavitensis** Yates. Black superficial fungus patches on leaves of *Coleus* sp. in the Philippines.

**COLLETTIA.** Spiny shrubs grown in warm regions or greenhouses.

**Dothidella colletiae** (P. Henn. and Lind.) Theiss. and Syd. On branches and cladophylls of *C. spinosa* in Chile.

**Montagnella curumamuel** Speg. Black circular stromata on leaves of *C. cruciata* and *C. ferox* in Argentina.

**Placostroma lorentzianum** (Speg.) Theiss. and Syd. On branches of *C. ferox* in Argentina.

**COLLOMIA.** See Gillia.**COLOCASIA.** DASHEEN. ELEPHANT'S-EAR. Yautia. Eddoe. Taro. Tropical aroids grown for their ornamental leaves (bedding plants) and for edible roots.

**Cercospora caladii** Cke. var. **colocasiae** v. Hoeh. Subcircular brown zoned leaf spots on *Colocasia* sp. in Samoa.

**Heterosporium colocasiae** Mass. On leaves of *Colocasia* sp. in Jamaica.

**Mycosphaerella colocasiae** K. Hara. Circular, then confluent, brown, zoned, more or less sunken leaf spots with dark-brown margins on *C. antiquorum* in Japan. The disease greatly reduces the yields of tubers.

**Peronospora trichotoma** Mass. This supposed species of downy mildew, said to cause a tuber rot of *C. esculenta* in the West Indies, is now considered a hyphomycete overgrowing *Phytophthora colocasiae* Rac.

**Phyllosticta colocasiae** v. Hoeh. Circular brown zoned leaf spots on *Colocasia* sp. and *Dieffenbachia* sp. in Samoa and Porto Rico.

**Phyllosticta colocasicola** v. Hoeh. Probably not distinct from the preceding species.

**Phytophthora colocasiae** Rac. This downy mildew causes a serious blight of *C. antiquorum* in India, Java, Ceylon, Formosa, Philippines, and Hawaii. Circular to oval dark-brown spots appear on the leaves and, becoming irregular, may occupy most of the leaf surface. They become yellow-brown in color, and are often marginally zoned in shades of brown, green, and yellow. The petioles may be soft rotted, so that the leaves fall over. Inflorescences and the corms are attacked in turn, although if the plants are attacked while young no corms are formed. Where corms form they are destroyed by a soft rot.

**COLQUHOUNIA.** Tender plants grown for their whorls of scarlet and yellow flowers.

**Puccinia leucophaea** Syd. and Butl. Leaf rust on *C. coccinea* in India.

**COLUTEA.** BLADDER SENNA. Shrubs grown for their attractive flowers and ornamental bladder-like pods.

**Ascochyta coluteae** Lambr. and Fautr. On leaves of *C. orientalis* in Yugoslavia.

**Ovularia vogeliapa** Sacc. and Syd. White fungus layers over leaf surfaces of *C. arborescens* in Germany.

**Uromyces genistae-tinctoriae** (Pers.) Wint. See Caragana.

- COMARUM.** Decumbent herbs. The genus is usually placed in *Potentilla*.  
*Mycosphaerella innumerella* (Karst.) Starb. On leaves of *C. palustre* in Europe.  
*Phyllosticta argentinae* Desm. See *Potentilla*.  
*Physothermum comari* (Berk. and White.) Lagh. Black smut sori in leaves of *C. palustre* in Great Britain and Scandinavia.  
*Septogloeum comari* Allesch. and Bres. Dull yellow leaf spots on *C. palustre* in Sweden, Denmark, Esthonia, and Germany.  
*Septoria comari* Lasch. On leaves of *C. palustre* in Germany.  
*Venturia palustris* Sacc. On leaves of *C. palustre* in Belgium and Germany.
- COMBRETUM.** Tropical shrubs and trees.  
*Hyphaster kutuensis* P. Henn. Superficial black stromata on leaves of *C. baumii* in the Union of South Africa.  
*Phyllosticta combreticola* P. Henn. Irregular ashen leaf spots on *C. argenteum* in Germany.  
*Uredo kampuluvensis* P. Henn. See *Baphia*.
- COMMELINA.** DAYFLOWER. Perennial or annual herbs.  
*Cercospora commelinae* Kalchbr. and Cke. Circular, dull-brown leaf spots on *C. benghalensis* in the Union of South Africa.  
*Cylindrosporium kilimandscharicum* Allesch. Small, definite, brown, often confluent leaf spots on *Commelina* sp. in Tropical Africa.  
*Rhizosporium commelinae* Stevens. Circular dull-yellow leaf spots on *C. longicaulis* and *C. virginica* in Porto Rico.  
*Kordyana celebensis* Racib. Causes leaf spots on *C. bengalensis* in Java.  
*Phyllosticta commelinicola* Young. Indefinite leaf spots on *C. nudiflora* in Porto Rico.  
*Puccinia commelinae* Holw. Brown rust pustules on brown irregular leaf spots on *Commelina* sp. in Mexico.  
*Stilbella flavida* (Cke.) Lind. See *Coffea*.  
*Tilletia* (?) *commelinae* Kom. Doubtful species of smut in ovaries of *C. communis* in Manchuria and Japan.  
*Uredo commelyneae* Kalchbr. Leaf rust on *C. elegans* and *C. virginica* in Porto Rico, St. Croix, and Trinidad.  
*Uredo ochracea* Diet. Brown rust pustules on leaves of *C. bengalensis*, *C. communis*, *C. longicaulis*, and *C. obliqua* in Brazil, India, Japan, Ceylon, and the Philippines.  
*Uredo pedicellata* Rac. Rust on leaves and bracts of *Commelina* sp. in Java.  
*Uromyces commelinae* Cke. Leaf rust on *C. bengalensis*, *C. communis*, *C. forskalii*, *C. longicaulis*, *C. nudiflora*, *C. subulata*, and *Tradescantia cumanensis* in the Union of South Africa, India, Japan, Portuguese East Africa, Uganda, Porto Rico, Cuba, Jamaica, Guatemala, and South America.  
*Uromyces tosenis* P. Henn. Brown rust pustules on leaves of *C. communis* in Japan.
- CONVALLARIA.** LILY OF THE VALLEY. Herbaceous perennials cultivated for their racemes of fragrant flowers.  
*Ascochyta majalis* Massal. Causes red-brown oblong to irregular spots on leaves of *C. majalis* in Italy.  
*Dendrophoma convallariae* Cav. Dark-red elongated spots are produced on the leaves of *C. majalis* in Great Britain, Denmark, Italy, and Germany.  
*Gloeosporium convallariae* Allesch. On leaves of *C. majalis* in Germany.  
*Heterosporium ornithogali* Klotzsch. Oval to irregular brown spots on the leaves of *Convallaria* sp. and other species of the family *Liliaceae* in New Zealand.  
*Lophodermium herbarum* (Fr.) Fckl. On leaves of *C. majalis* in France, Belgium, Switzerland, Sweden, Finland, and Germany.  
*Mycosphaerella asteroma* (Fr.) Lind. On leaves of *Convallaria* sp., *Maianthemum* sp. and *Polygonatum* sp. in Europe.  
*Mycosphaerella brunneola* (Fr.) All. and Schnab. On leaves of *C. majalis* in Denmark, Belgium, and Austria.  
*Phyllachora melanoplaca* Sacc. Forms black, shiny, slightly concave spots on the leaves of *Convallaria majalis*, *Veratrum album*, *V. nigrum*, and *V. viride* in Switzerland and Austria.  
*Sclerotium convallariae* West. On leaves of *Polygonatum verticillatum* (*C. verticillata*) in Belgium.  
*Septoria brunneola* (Fr.) Niessl. (*S. convallariae* West.) Causes brown leaf spots on *C. majalis* and *Polygonatum* sp. in Siberia, Great Britain, Belgium, Sweden, Italy, and Moravia. Probably the imperfect stage of *Mycosphaerella brunneola*.  
*Septoria majalis* Aderh. Large brown spots without definite margins, often involving entire leaves of *C. majalis* in Germany.  
*Septoria subradians* (Fr.) Karst. A leaf-spot disease of *C. majalis* in Denmark.
- CONVOLVULUS.** BINDWEED. Annual and perennial climbing herbs, mostly weeds.  
*Aecidium congoam* P. Henn. Leaf rust on *Convolvulus* sp. in the Kongo.  
*Aecidium convolvulinum* Speg. Leaf rust on *Convolvulus* sp. in Argentina.  
*Cercospora sorokinii* Sacc. On leaves of *Convolvulus* sp. in central Asia.  
*Entyloma convolvuli* Bres. Smut sori in circular to oblong brown leaf spots, on *C. soldanella* in Portugal.  
*Puccinia macrocephala* Speg. See *Ipomoea*.  
*Septoria calystegiae* West. Small subcircular brown leaf spots on *C. arvensis* and *C. soldanella* in Europe.  
*Septoria convolvulina* Speg. Circular white leaf spots with reddish margins on *C. arvensis* in Argentina.  
*Septoria longispora* A. Bond. Small circular brown to gray leaf spots on *C. arvensis* and *C. lineatus* in Russia.  
*Septoria obesipora* Oud. Leaf spot on *C. sepium* in Holland.  
*Thecaphora hyalina* Fingerh. A smut reducing ovaries of *C. arvensis*, *C. sepium*, and *Soldanella* sp. to brown, powdery, spore masses in Europe. What is apparently this species has been collected in California.  
*Uromyces gemmatus* B. and C. Leaf rust on *Convolvulus* sp. in Cuba.
- COPAIFERA.** Brazilian trees, producing a balsam called *copaiba*.  
*Phyllachora copaiferae* Speg. Black stromata on irregular brown leaf spots on *C. langsdorffii* in Paraguay and Brazil.  
*Uredo copaiferae* P. Henn. Leaf rust on *Copaifera* sp. in Brazil.
- COPEBNICIA.** See *Palmae*.
- COPROSMA.** Shrubs or small trees grown for their ornamental fruit and variegated leaves.  
*Phyllosticta coprosmae* McAlp. Ashen leaf spots, with narrow, raised, dark-red margins on *Coprosma* sp. in Australia.  
*Puccinia coprosmae* Cke. Rusty-brown rust pustules on irregular leaf spots on *C. billardieri*, *C. hirtella*, and *C. lucida* in New Zealand.  
*Septoria coprosmae* Cke. On leaves of *Coprosma* sp. in New Zealand.

**CORCHORUS.** Jute. Shrubs or herbs, some species yielding fiber.

**Corticium salmonicolor** B. and Br. See Citrus.

**Diplodia corchori** Syd. The "black band" disease of jute (*C. capsularis* and *C. olitorius*) is serious in India. Dense black bands form on the stems of infected plants up to the height of 2 or 3 feet, the bark splits, and finally only the brown, dry fibers are left, the tissue between having decayed away. The leaves of infected plants turn yellow and drop.

**Hymenula nigra** Saw. On *Corchorus* sp. in Formosa.

**Hypochnus centrifugus** (Lév.) Ful. On *Corchorus* sp. in Formosa.

**Macrophoma corchori** Saw. Indefinite spots and cankers on stems of *Corchorus* sp. (jute) in Formosa, causing die-back. Considered serious.

**Phyllosticta corchori** Saw. On leaves of *Corchorus* sp. in Formosa.

**Rhizoctonia** sp. See Vigna.

**CORDIA.** Tropical shrubs and trees, some species cultivated for fruit and ornament.

**Aecidium brasiliensis** Diet. Leaf rust on large round yellow spots on *C. cylindrostachya*, *C. macrophylla*, *C. nodosa*, and *C. rothii* in Brazil, Trinidad, and India.

**Aecidium cordiae** P. Henn. Leaf rust on *C. bullata* and *C. urticifolia* in the Dominican Republic.

**Alveolaria andina** Lagh. Chestnut-brown rust pustules on yellow-brown circular leaf spots of *Cordia* sp. in Ecuador.

**Alveolaria cordiae** Lagh. Rust on sunken spots on leaf blades and petioles of *C. cylindrostachya*, *C. ferruginea*, *C. laxiflora*, and *C. riparia* in Costa Rica, Guatemala, Jamaica, Ecuador, and Colombia.

**Dimargium stevensii** Garm. Indefinite yellow spots on leaves of *C. corymbosa* in Porto Rico.

**Guignardia javanica** Koord. Indefinite dark-brown leaf spots on *C. suareolens* in Java.

**Helminthosporium cordiae** Welw. and Curr. On leaves of *Cordia* sp. in west Africa.

**Phyllachora caffra** Syd. Black stromata on rust-brown indefinite leaf spots on *C. caffra* in the Union of South Africa.

**Placosphaeria cordiae** Allesch. Subcircular black stromata on leaves of *Cordia* sp. in Brazil.

**Puccinia cordiae** Arth. Brown rust pustules on leaves of *C. alliodora* and *C. gerascanthus* in Porto Rico and Guatemala.

**Puccinia corticola** Arth. and Rorer. Rust attacking leaves, twigs, branches, and trunks of *C. gerascanthus* in Trinidad. Large lesions are formed on trunks and branches.

**Uredo cordiae** P. Henn. A rust causing gall-like swellings on leaves and shoots and witches'-broom effect on *Cordia* sp. in Peru.

**Uromyces cordiae** P. Henn. Brown rust pustules on leaves of *Cordia* sp. in Brazil.

**CORDYLINE.** See Dracaena.

**COREOPSIS.** Annual or perennial herbs or shrubs, some species cultivated in flower gardens.

**Puccinia coreopsidis** Jacks. and Holw. Brown rust pustules on leaves of *C. mutica* (*C. mexicana*) in Guatemala.

**Puccinia coreopsidis** Wakef. Leaf rust on *Coreopsis* sp. in Uganda. The preceding species has priority.

**CORIANDRUM.** CORIANDER. Strong-smelling umbelliferous herbs the seeds of which are used in seasoning.

**Protomyces macrosporus** Ung. Galls up to half an inch long by an eighth of an inch wide are produced on flower stalks, petioles, and young stems of *Aegopodium podagraria*, *Carum carvi*, *Chaerophyllum hirsutum*, *Coriandrum sativum*, *Daucus carota*, *Heracleum sphondylium*, *Meum athamanticum*, *M. mutellina*, and *Pastinaca sativa* in Europe, north Africa, India, Japan, and Australia.

**CORIARIA.** Shrubs or herbs grown for their ornamental fruit.

**Phyllosticta coriariicola** Speg. Angular, then confluent, dull-brown leaf spots with purple margins on *C. ruscifolia* in Chile.

**Pucciniastrum coriariae** Diet. Yellow to brown rust pustules on leaves of *C. japonica* and *C. nepalensis* in India and Japan.

**Septoria coriariae** Pass. Irregular brown leaf spots on *C. myrtifolia* in Spain and Italy.

**CORNUS.** DOGWOOD. Woody plants grown for their attractive flowers and fruit.

**Coryneum corni-albi** (Roum.) Sacc. On leaves of *C. alba* and *C. mas* in France, Bohemia, Austria, and Germany.

**Erysiphe tortilis** (Wallr.) Fr. Powdery mildew on leaves of *C. alba* and *C. sanguinea* in Europe.

**Helminthosporium phylophilum** Karst. On leaves of *C. alba* in Finland.

**Phoma thallina** Sacc. On stems of *C. sanguinea* in Europe.

**Pucciniastrum corni** Diet. Yellow to brown rust pustules on leaves of *C. kousa* and *C. officinalis* in Japan.

**Septoria corni-maritima** Sacc. On leaves of *C. sanguinea* in northern Europe.

**Sphaeropsis suspecta** Vestreg. Stem canker of *C. sanguinea* in Russia.

**COROKIA.** Evergreen shrub.

**Septoria corockeae** P. Henn. Dull-brown spots, occupying large portions of leaf surfaces of *C. budleyoides* in Germany.

**CORONILLA.** Crown vetch. Shrubs and herbs grown in gardens for their flowers.

**Ascochyta emeris** Sacc. Dirty-white, often marginal, leaf spots with red margins on *C. emeris* in Italy.

**Cercospora ratuensis** C. Mass. Small subcircular pale-gray leaf spots with rufous margins on *C. varia* in Italy.

**Helminthosporium bornmuelleri** P. Magn. On leaves of *C. coronata*, *C. montana*, *C. scorpioides*, and *C. vaginalis* in Spain, Dalmatia, and Austria.

**Myosphaerella ariadna** (Sacc.) Lind. On leaves of *C. emeris* in Italy.

**Peronospora coronillae** Gäum. Downy mildew on leaves of *C. scorpioides* and *C. varia* in Dalmatia and Silesia.

**Puccinia coronillae** Wor. Leaf rust on *C. cappadocica* in Russia.

**Ramularia coronillae** Bres. Small pale-brown leaf spots on *C. varia* in Yugoslavia and Austria.

**Septoria emeris** Sacc. White leaf spots with dull-brown margins on *C. emeris* in Italy.

**Uromyces anthyllidis** (Grev.) Schroet. See Anthyllis.

**CORREA.** Australian shrubs.

**Phyllosticta correae** McAlp. Elongate brown leaf spots on *C. speciosa* in Australia.

**Puccinia correae** McAlp. Brown rust pustules on leaves of *C. lawrenciana* in Tasmania.

**CORYDALIS.** CAPNOIDES Ag. Hardy erect or prostrate herbaceous perennials.

**Entyloma corydalis** De By. Smut sori in white, then brown, leaf spots on *C. cava* and *C. solida* in Russia, France, Denmark, and Germany.

**Entyloma corydalis-luteae** Vogl. Smut sori in circular yellow-brown spots on leaves and stems of *C. lutea* in Switzerland and Italy.

**Melampsora magnusiana** G. Wagn. See Populus.

**Peronospora bulbocapni** Beck. Downy mildew on leaves of *C. ambigua*, *C. cava*, *C. fabacea*, and *C. laxa* in Japan and Europe.

**Urocystis corydalis** Niessl. Smut sori in brown circular leaf spots on *C. cava* and *C. solida* in Bohemia.

- CORYLUS.** HAZELNUT. FILBERT. Shrubs grown for nuts and foliage.  
**Ascochyta coryli** Sacc. and Speg. Subcircular to irregular white leaf spots on *C. avellana* in Italy.  
**Bacterium** sp. A bacterial disease of *Corylus* is reported as serious in Oregon. It causes a blighting of buds and new shoots, a brown speckling of the leaves, a girdling and dying-back of smaller branches, and formation of large cankers on larger branches and trunks.  
**Bacterium coryli** Brzez. This bacterium is reported as the cause of a canker disease of *C. avellana* and *C. colurna* in central Europe.  
**Cryptosporella anomala** (Pk.) Sacc. The disease due to this fungus, known as "filbert blight," is serious in the eastern part of the United States on *C. avellana*, and an effort is being made to prevent its spread to the Pacific coast. The fungus is native on *C. americana*. Sunken cankers of varying size and shape occur on branches and trunks, girdling the stems and soon killing outright many plants, particularly those of the European varieties.  
**Labrella coryli** (Desm. and Rob.) Sacc. Irregular rufous leaf spots on *C. avellana* in Great Britain and France.  
**Mamiania coryli** (Batsch.) Ces. and De N. On leaves of *C. avellana*, *C. heterophylla*, and *C. rostrata* in Japan, Denmark, and Austria.  
**Nematospora coryli** Pegl. Causes malformation of nuts of *Corylus* sp. in Italy.  
**Phyllohendersonia corylaria** (Sacc.) Tass. On leaves of *C. avellana* in Italy.  
**Phytophthora syringae** Kleb. See Syringa.  
**Pucciniastrum coryli** Kom. Ochraceous rust pustules on leaves of *C. heterophylla* and *C. rostrata* in Russia, Manchuria, and Japan.  
**Septoria avellanae** B. and Br. On leaves of *C. avellana* in Great Britain and Italy.
- CORYNOCARPUS.** Ornamental evergreen trees.  
**Gloeosporium intermedium** Sacc. See Citrus.  
**Phyllosticta corynocarpi** Alm. and Cam. On leaves of *C. laevigata* in Portugal.  
**Septoria corynocarpi** Thuem. Circular to elliptical white leaf spots on *C. laevigata* in Portugal.
- CORYPHA.** See Palmae.
- COSMOS.** Annual or perennial herbs grown in flower gardens.  
**Entyloma holwayi** Syd. Smut sori in circular to irregular yellow-brown, then deep-brown, leaf spots on *C. sulphureus* in Mexico.  
**Uromyces bidentis** Lagerh. Brown rust pustules on leaves of *C. caudatus* and *Bidens* spp. in Porto Rico, Jamaica, and northern South America.
- COSTUS.** Perennial thick-rooted tropical herbs.  
**Cercospora costina** Syd. Leaf spot on *C. speciosus* in the Philippines.  
**Dactylaria costi** Saw. Leaf spot on *C. speciosus* in Japan.  
**Puccinia costi** (P. Henn.) Syd. Powdery brown rust pustules on indefinite sunken yellow leaf spots on *C. pumilus* in Brazil.  
**Uredo costina** Syd. Leaf rust on *C. speciosus* in the Philippines.
- COTONEASTER.** Shrubs grown for their ornamental red or black fruits and for the foliage.  
**Aecidium cunninghamianum** Bardl. Leaf rust on *C. bacilleris* in India.  
**Coleopuccinia sinensis** Pat. Brown rust pustules on leaves of *Cotoneaster* sp. and *Amelanchier* sp. in China.  
**Entomospodium mespili** (DC.) Sacc. Brown leaf spots on *C. nigra*, *Eriobotrya japonica*, *Mespilus germanica*, and *Pyrus silvestris* in Europe and Argentina. Probably not distinct from *E. maculatum* Lév.  
**Fusicladium pyracanthae** (Othth.) Rostr. On fruit and peduncles of *C. pyracantha* in Denmark and Russia. Probably the same as *F. pirinum*.  
**Gymnosporangium mespili** (DC.) Kern. See Juniperus.  
**Phyllosticta cotoneastri** Allesch. Small subcircular, often confluent, brown, then gray, leaf spots with red-brown margins on *Cotoneaster* sp. in Germany.
- COTYLEDON.** Succulent herbs or shrubs grown mostly for their oddity.  
**Aecidium umbilici** Trotter. Leaf rust on *C. umbilicus* in Portugal.  
**Ascochyta cotyledonis** H. Zimm. Large gray leaf spots with purple-black margins on *C. gibbiflora* in Austria.  
**Microsphaera umbilici** Kom. Powdery mildew on leaves of *C. semenowii* in Turkestan.  
**Septoria dalmatica** Jaap. Circular gray leaf spots on *C. chlorantha* in Dalmatia.  
**Septoria zimmermanni-hugonis** Bub. Circular to elliptical brown leaf spots with gray margins on *C. desmetiana*, *C. gibbiflora*, and *C. pachyphytum* in Bohemia.
- CRACCA.** See Tephrosia.
- CRASSULA.** Fleshy-leaved greenhouse shrubs and herbs grown for the grotesque appearance of some species.  
**Aphelenchus olesistus** Ritz. Bos. See Begonia.  
**Puccinia exanthematica** McO. Leaf rust on *C. cordata* and *C. spathulata* in the Union of South Africa.
- CRATAEGUS.** HAWTHORN. Trees grown for their flowers and decorative fruit.  
**Aecidium patulum** Syd. Rust on leaf blades, petioles, and fruit of *C. coccinea* in India.  
**Ascochyta crataegi** Fekl. On leaves of *C. oxyacantha* in Russia, Italy, and Germany.  
**Ascochyta crataegicola** Allesch. On leaves of *C. oxyacantha* in Russia and Germany.  
**Ascochyta crus-galli** P. Brun. Circular to subcircular white leaf spots with brown margins on *C. crus-galli* in France.  
**Ascochyta misera** Oud. On leaves of *C. monogyna* in Holland.  
**Cercospora crataegi** Sacc. and C. Mass. Small angular yellow leaf spots on *C. oxyacantha* in Italy.  
**Coryneum foliocolum** Fekl. Ochraceous leaf spots on *C. oxyacantha*, *Quercus* sp., *Rubus discolor*, and *R. fruticosus* in Argentina, Russia, Italy, and Germany. Reported from Indiana.  
**Exoascus crataegi** (Sad.) Sacc. Causes a crumpling and red spotting of the leaves and sometimes a deformation of twigs of *C. monogyna*, *C. oxyacantha*, and *C. sanguinea* in Europe.  
**Fusicladium crataegi** Aderh. (*Venturia crataegi* Aderh.) Black spots on fruit of *C. monogyna* and *C. oxyacantha* in Denmark and Germany.  
**Gymnosporangium mespili** (D. C.) Kern. See Juniperus.  
**Gymnosporangium orientale** Syd. Aecia on fruit and branches of *C. ararella* and *C. insigna* in Russia, Greece, Thessaly, Asia Minor, and Persia.  
**Monilia crataegi** Died. Effuse brownish leaf spots on *C. oxyacantha* in Germany.  
**Mycosphaerella crataegi** (Fekl.) Oud. On leaves of *C. monogyna* in Great Britain, Denmark, and Germany.  
**Mycosphaerella crataegicola** (Fekl.) Bond. and Tranz. On leaves of *C. monogyna* and *C. sanguinea* in Russia.  
**Mycosphaerella oxyacanthae** Jaap. On leaves of *C. oxyacantha* in Switzerland, Austria, and Germany.  
**Phleospora oxyacanthae** (Kunze. and Schm.) Walb. Crowded yellowish leaf spots on *C. oxyacantha* in Russia, Great Britain, Italy, and Germany. "Hawthorn leaf scorch."  
**Phyllosticta casinalbensis** D. Sacc. White leaf spots on *C. azarolus* in Italy.

**CRATAEGUS**—Continued.

- Phyllosticta michailorskensis** Elenk. and Ohl. Leaf spots on *C. sanguinea* in Russia.  
**Phyllosticta monogyna** Allesch. On leaves of *C. monogyna* and *C. oxyacantha* in Russia.  
**Phyllosticta phaea** Sacc. On leaves of *C. oxyacantha* in Moravia.  
**Phytophthora syringae** Kleb. See *Syringa*.  
**Septoria crataegicola** Bond. and Tranz. On leaves of *Crataegus* sp. in Russia.  
**Septoria crataegophia** Ranoj. On *C. monogyna* in Yugoslavia.  
**Stereum purpureum** Pers. See *Prunus*.  
**Trichoseptoria fructigena** Maubl. See *Cydonia*.  
**Uncinula prunastri** (DC.) Sacc. See *Prunus*.

**CRATAEVA**. Tropical trees and shrubs.

**Acidium crataevae** Syd. Rust pustules on large, circular to subcircular, sunken, yellow-brown leaf spots on *C. religiosa* in India.

**Fusidium crataevae** B. and Br. On leaves of *C. roxburghii* in Ceylon.

**Napidadium crataevae** Syd. On leaves of *C. religiosa* in India.

**CREPIS**. Annual, biennial, or perennial herbs, some species grown in flower gardens.

**Bremia ovata** Saw. Downy mildew on leaves of *C. japonica* in Japan and Formosa.

**Entyloma calendulae** (Oud.) De By. See *Calendula*.

**Entyloma crepidicola** Trot. and var. **crepidis-rubrae** Jaap. A smut forming galls on the fibrous roots and rhizomes of *C. bulbosa* and *C. rubra* in Italy and Dalmatia. The variety has been described as a new species under the name *Tolyposporium crepidis-rubri* (Jaap) Cif.

**Phragmidium eximia** Bub. Leaf rust on *C. grandiflora* in Yugoslavia.

**Phyllosticta crepidis-paludosae** Petr. Subcircular to irregular, ocher-brown leaf spots on *C. paludosa* in Austria.

**Phyllosticta eximia** Bub. Angular dark-brown to black leaf spots on *C. viscidula* in Yugoslavia.

**Protomyces crepidicola** Buren. On leaves of *C. biennis* in central Europe.

**Protomyces crepidis** (Jaap.) Sacc. and Trott. Small circular dirty yellow-gray then yellow-brown pustules on *C. aurea*, *C. biennis*, *C. incarnata*, and *C. montana* in Switzerland and Austria.

**Protomyces crepidis-paludosae** Buren. On leaves of *C. paludosa* in Switzerland.

**Protomyces inouei** P. Henn. Effuse, elongate sori in peduncles of *Crepis* sp. in Japan.

**Puccinia alpestris** Syd. Yellow and brown rust pustules on leaves of *C. alpestris* in Switzerland and Austria.

**Puccinia aschersoniana** P. Henn. Brown rust pustules on leaves of *C. rueppellii* in Arabia.

**Puccinia barkhousiae-rhoeadifolia** Bub. Yellow and dark brown to black rust pustules on leaves of *C. foetida* and *C. rhoeadifolia* in Bohemia and Spain.

**Puccinia crepidicola** Syd. Brown powdery rust pustules on leaves of *Crepis* spp. in Europe and Asia Minor.

**Puccinia crepidis** Schroet. Leaf rust on *Crepis* spp. in Europe.

**Puccinia crepidis-aureae** Syd. Leaf rust on *C. aurea* in Switzerland and Austria.

**Puccinia crepidis-blattarioides** Kasl. Leaf rust on *C. albida*, *C. alpestris*, *C. blattarioides*, *C. tectus*, and *C. virens* in Switzerland and Spain.

**Puccinia crepidis-grandiflorae** Hasl. Rust on leaves of *C. bellidifolia*, *C. grandiflora*, *C. nicaeensis*, and *C. tectus* in Switzerland.

**Puccinia crepidis-japonicae** Diet. Powdery, chestnut-brown rust pustules on leaves of *C. japonica* in Japan.

**Puccinia crepidis-leontodontoidis** R. Maire. Powdery, cinnamon-brown to dark-brown rust pustules on leaves of *C. leontodontoides* in Corsica.

**Puccinia crepidis-pygmaeae** Gaill. Dark-brown rust pustules on leaf blades and petioles of *C. pygmaea* in France and Austria.

**Puccinia crepidis-sibiricae** Lindr. Brown powdery rust pustules on leaves of *C. sibirica* in Siberia, Russia, and Finland.

**Puccinia crucheti** Hasl. Rust on leaves of *C. succisaefolia* in Switzerland.

**Puccinia intybi** (Juel.) Syd. Leaf rust on *C. praemorsa* in Great Britain, Sweden, Esthonia, Denmark, and Germany.

**Puccinia major** Diet. Cinnamon to dark-brown rust pustules on leaves of *C. conyzifolia*, *C. grandiflora*, and *C. paludosa* in Europe.

**Puccinia praecox** Bub. Powdery yellow and brown rust pustules on leaves of *C. biennis*, *C. foetida*, and *C. rubra* in Russia, Yugoslavia, Switzerland, Bohemia, Denmark, and Austria.

**Puccinia scaliana** Syd. Leaf rust on *C. biennis* and *C. bursifolia* in Spain and Sicily.

**Puccinia silvatica** Schroet. See *Taraxacum*.

**Ramularia eximia** Bub. Angular to irregular yellow leaf spots on *C. grandiflora* and *C. viscidula* in Yugoslavia.

**Septoria crepidis** Vestergr. Irregular, often confluent, leaf spots on *C. biennis* and *C. tectorum* in Russia, Esthonia, and Sweden.

**Stagonospora crepidis** Hóll. Circular, dull-brown leaf spots with black margins on *C. biennis* in Hungary.

**Taphrina rhaetica** Volk. On leaves of *C. blattarioides* in Switzerland.

**Uredo crepidis-integrae** Lindr. Leaf rust on *C. integra*, *C. japonica*, *C. keiskeana*, and *C. platyphylla* in Japan and Australia.

**Uredo crepidis-japonicae** Lindr. Leaf rust on *C. japonica* in Ceylon and Australia.

**CRINUM**. Large, showy, flowering bulbs.

**Acidium crini** Kalchbr. Leaf rust on *C. longifolium* (*C. capense*) and *C. cooperi* in the Union of South Africa.

**Acidium mangaranga** P. Henn. Rust on large, effuse, yellow leaf spots on *Crinum* sp. in the Union of South Africa.

**Gloeosporium crini** Sacc. Leaf anthracnose of *Crinum* sp. in Italy.

**Mycosphaerella crini** Siem. On leaves of *Crinum* sp. in the Caucasus.

**Phyllosticta crini** Sacc. Subcircular yellow leaf spots on *Crinum* sp. in Italy.

**Phyllosticta crinicola** Siem. On leaves of *Crinum* sp. in southern Russia.

**Septoria crini** Tassi. On leaves of *C. asiaticum* in Italy.

**CROCUS**. Low, spring and autumn flowering bulbs.

**Bacillus croci** Miz. This bacterium causes a serious rot of *Crocus* spp. in Japan, attacking bulbs, roots, sheaths, leaves, and flower stalks. *Narcissus* sp., *Hyacinthus* sp., and *Allium* spp. have been infected by inoculation. The lower leaves become rotten, turn yellow above, and are readily detached from the corm, which is in turn soft-rotted.

**Perisporium crocophila** (Mont.) Sacc. Brown spots appear on corms of *C. sativus* in France, increasing in size, and finally bring about rot of the corm. A black fungus layer covers the rotted areas.

**Sclerotinia tuliparum** (Wakk.) Rehm. See *Tulipa*.

**Uromyces croci** Pass. Elongate, powdery brown rust pustules on both leaf surfaces of *C. biflorus* and *C. susianus* in Russia, Italy, France, and Austria.

- CROTALARIA.** Rattlebox. Sunn-hemp. Annual leguminous herbs and shrubs, some species grown as green-manure crops.
- Aecidium crotalariae** P. Henn. Leaf rust on *C. engleri*, *C. grandibracteata*, and *Crotalaria* sp. in Uganda and west Africa.
- Aecidium crotalariae** P. Henn. Rust on dull-brown leaf spots on *Crotalaria* sp. in Brazil.
- Aecidium dielsii** P. Henn. Ochraceous rust pustules on leaves of *Crotalaria* sp. in Tanganyika.
- Cercospora crotalariae** Sacc. Leaf spots on *C. striata* in Ceylon.
- Colletotrichum crotalariae** Petch. Anthracnose on *C. striata* in Ceylon.
- Mycosphaerella crotalariae** Petch. Pale-brown zoned leaf spots on *C. striata* in Ceylon.
- Phyllosticta crotalariae** Sacc. On leaves of *C. striata* and *Crotalaria* sp. in Uganda and Ceylon.
- Poria hypobrunnea** Petch. See Hevea.
- Uredo crotalariae** Diet. Ochraceous rust pustules on leaves of *C. incana* and *C. vitellina* in Brazil.
- Uredo crotalariae** P. Henn. Leaf rust on *C. lachnoclada* in central Africa.
- Uredo harmslana** P. Henn. Leaf rust on *C. lanceolata* in the Union of South Africa.
- Uredo theresiae** Neger. Brown rust pustules on leaves of *C. anagyroides* in Colombia.
- Uromyces decoratus** Syd. Powdery brown to black rust sori on yellow leaf spots on *C. albida*, *C. ferruginea*, *C. juncea*, and *C. vitellina* in India, Ceylon, and Costa Rica.
- Uropyxis crotalariae** Arth. Brown rust pustules on leaves of *C. maypurensis* and *Crotalaria* sp. in Guatemala.
- CROTON.** Shrubs and trees, a few species of economic importance.
- Cercospora crotonophila** Speg. On leaves of *Croton* sp. in Paraguay.
- Cercospora manaensis** P. Henn. On leaves of *Croton* sp. in Brazil.
- Cercospora tigilii** P. Henn. Circular, black leaf spots on *C. tigilium* in the Philippines.
- Cercospora crotalariae** P. Henn. Leaf spots, often marginal, causing curling on *Croton* sp. in Brazil.
- Gloeosporium crotonis** Delacr. Anthracnose of *Croton* sp. in France.
- Phakopsora crotonis** (Cke.) Arth. Leaf rust on *C. gossypifolius* and *C. hirtus* in Trinidad.
- Phyllachora crotonis** (Cke.) Sacc. "Tar spot" on leaves of *C. arborescens*, *C. chamaedryfolius*, *C. floribundus*, and *C. silvaticus* in South America and the Union of South Africa.
- Phyllachora globispora** Speg. Black stromata on leaves of *Croton* sp. in Argentina.
- Phyllachora tragiae** (B. and C.) Sacc. Circular black stromata on leaves of *C. arborescens*, *C. chamaedryfolium*, *C. flavens*, and *C. lucidus* in Brazil, Porto Rico, French Guiana, Argentina, and Brazil.
- Phyllosticta portoricensis** Young. Circular yellowish leaf spots on *C. lucidus* in Porto Rico.
- Schroeteriaster argentinensis** (Speg.) Syd. Rust on leaf blades, petioles, and stems of *C. hirtus* in Argentina.
- Schroeteriaster mexicanus** (Arth.) Syd. Cinnamon-brown rust pustules on leaves of *C. calvescens* in Mexico.
- Schroeteriaster stratosus** (Cke.) Syd. Powdery brown rust pustules on leaves of *C. oxypetalus* and *C. silvaticus* in the Union of South Africa and the Congo.
- Septoria crotonis** Bres. On leaves of *C. macrostachya* and *Croton* sp. in Denmark and Abyssinia.
- Uredo crotonicola** P. Henn. Leaf rust on *C. glandulosus* and *Croton* sp. in Brazil and Argentina.
- Uredo crotonis** P. Henn. Leaf rust on *Croton* sp. in Brazil.
- CRUCIANELLA.** CROSSWORT. Hardy rock plants.
- Puccinia crucianellae** Desm. Brown rust pustules on leaves and stems of *C. angustifolia*, *C. herbacea*, and *C. maritima* in Tripoli, Palestine, Portugal, Spain, Italy, and France.
- Puccinia monopera** Lindr. Rust on leaves and stems of *C. (Asperula) glomerata* in Greece.
- Puccinia syriaca** Syd. Powdery dark-brown rust pustules on leaves and stems of *C. macrostachya* and *C. syriaca* in Syria.
- Uredo mediterraneae** Lindr. Rust on stems of *C. maritima* in France and Portugal.
- CRYPTOGRAMMA.** ROCK BRAKE. Hardy subalpine ferns.
- Hyalospora cryptogrammes** Diet. Brown rust pustules on leaves of *C. japonica* in Japan.
- CRYPTOMERIA.** Ornamental coniferous trees.
- Fusarium blasticola** Rostr. See Pinus.
- Helicobasidium mompa** Tanak. See Morus.
- Valsa cryptomeriae** Kitajima. The "blister" disease of seedlings of *C. japonica* is characterized by blackish-brown or reddish-brown areas on the stems which become sunken and cracked. These cankers cause hypertrophy and bending of diseased seedlings. The disease occurs in Japan.
- CRYPTOSTEGIA.** RUBBER VINE. Tropical climbers.
- Ramularia cryptostegiae** Pim. On leaves and fallen seeds of *Cryptostegia* sp. in Ireland.
- CRYPTOSTEMMA.** Composite herbs.
- Ascochyta cryptostemmatis** McAlp. Irregular, deep brown leaf spots on *C. calendulaceum* in Australia.
- Septoria perforans** McAlp. Circular, then confluent, pale gray-brown leaf spots on *C. calendulaceum* in Australia.
- CUCUMIS.** MELON. CUCUMBER. Herbaceous vines cultivated for their fruit.
- Ascochyta melonis** Poteb. On fruit, stems, and leaf blades and petioles of *C. melo* and *C. sativus* in Russia.
- Corticium koleroga** (Cke.) v. Hoeh. See Coffea.
- Corynespora melonis** (Cke.) Sacc. (*Cercospora melonis* Cke.). Small pale-green translucent spots on leaves and young fruits, becoming gray, on *C. melo* and *C. sativus* in Great Britain, Sweden, Russia, Denmark, and Germany.
- Helminthosporium cucumerinum** Jarb. Pale-yellow leaf spots up to 1.5 cm. in diameter, with definite margins, on *C. sativus* in south Russia. Infected plants are so weakened that death finally ensues.
- Hypochnus cucumeris** Frank. This fungus attacks its hosts at the ground level, rotting the stems and forming a thin, grayish film of hyphae. Diseased plants turn yellow and collapse. The hosts are *Amorphophallus konja*, *Arachis hypogaea*, *Cinnamomum camphora*, *Citrullus edulis*, *C. vulgaris*, *Cucumis melo*, *C. sativus*, *Daucus carota*, *Morus alba*, *Phaseolus mungo*, *P. vulgaris*, *Sesamum indicum*, *Soja max*, *Trifolium pratense*, *Vicia faba*, and *Vigna sinensis* in Japan, Great Britain, Denmark, and Germany.
- Puccinia cucumeris** P. Henn. Yellow and black rust pustules on leaves of *C. ficifolius* in Abyssinia.
- Rhizoctonia** sp. See Vigna.
- Scolecotrichum melophthorum** Prill. and Delacr. Sunken brown spots on leaves and young stems, causing a rot of *C. melo*, *C. sativus*, and *Cucurbita vulgaris* in Great Britain, Italy, France, Belgium, Holland, and Russia.
- Ustilago cucumeris** Graff. A doubtful species of root smut on *C. sativus* in Great Britain.
- CUCURBITA.** PUMPKIN. SQUASH. GOURD. Cucurbitaceous vines.
- Cercospora cucurbiticola** P. Henn. Circular brown leaf spots on *Cucurbita* sp. in Brazil.
- Scolecotrichum melophthorum** Prill. and Delacr. See Cucumis.

- CUDRANIA.** Woody foliage plants used for hedges.  
**Hymenopsis cudraniae** Mass. On leaves of *C. javanensis* in Queensland, India, and the Philippines.  
**Melasma cudraniae** (Mass.) v. Hoeh. On *C. javanensis* in the Philippines.  
**Physopella sinense** Syd. Leaf rust on *Cudrania* sp. in China.  
**Rhytisma hypoxanthum** B. and Br. Black, irregular stromata on leaves of *C. javanensis* in Queensland.  
**Uredo cudraniae** Petch. Leaf rust on *C. javanensis* in Ceylon.
- CUMMINGIA.** Tender summer-blooming bulbs.  
**Phyllosticta santiaguina** Speg. Linear, red leaf spots on *C. campanulata* in Chile.
- CUNILA.** STONE MINT. Low-tufted hardy perennials.  
**Puccinia cunilae** Diet. Brown rust pustules on leaves of *C. angustifolia* in Brazil.  
**Puccinia fuscata** Arth. and Holw. Leaf rust on *C. leucantha* in Guatemala.
- CUNNINGHAMIA.** BELIS Ag. CHINA FIR. Coniferous trees cultivated for their handsome foliage.  
**Mycosphaerella cunninghamiae** Wor. On leaves of *C. sinensis* in Russia.
- CUNONIA.** Trees and shrubs.  
**Cercospora capensis** (Thuem.) Sacc. Subcircular, depressed, black leaf spots on *C. capensis* in the Union of South Africa.
- UPHEA.** PARSONSIA Ag. Tropical and subtropical herbs and shrubs.  
**Didymariopsis cuphaeicola** Speg. Circular, then irregular, leaf spots on *C. mesostemum* in Argentina.  
**Puccinia cuphaeae** Holw. Dark-brown rust pustules on leaves of *C. aequipetala*, *C. cyanea*, *C. hookeriana*, *C. nitidula*, *C. procumbens*, and *C. squamuligera* in Mexico and Guatemala.  
**Puccinia jaliscensis** Holw. Brown leaf rust on *C. cyanea*, *C. hookeriana*, *C. nitidula*, and *C. squamulifera* in Mexico.  
**Septoria cuphaeae** Tassi. Brown leaf spots with wide rosy margins on *C. ilavea* in Italy.  
**Uredo cuphaeae** P. Henn. Brown rust pustules on confluent, violet-brown leaf spots on *C. micrantha*, *C. parsonsia*, and *C. serphyllifolia* in Porto Rico, Colombia, and Brazil.
- CUPRESSUS.** CYPRESS. Ornamental evergreens and timber trees.  
**Gymnosporangium cunninghamianum** Barcl. Brown, slightly raised areas on smaller branches of *C. torulosa* in India. The aecial stage of this rust occurs on leaves of *Pyrus variolosa*.
- CURATELLA.** Small trees or scandent herbs.  
**Phyllosticta curatellae** P. Henn. Circular red-brown to dull-brown leaf spots on *C. americana* in Peru.
- CURCULIGO.** Stemless herbs of palmlike habit.  
**Puccinia curculigonis** Racib. Circular to irregular dark-brown spots, in which appear brown powdery rust pustules, on both leaf surfaces of *C. latifolia*, *C. orchoides* and *C. recurvata* in Java, India, and the Philippines.
- CURCUMA.** Turmeric. Tropical gingerlike plants.  
**Taphrina maculans** Butl. Numerous dirty-yellow to bay, often confluent, spots on both leaf surfaces of *C. amada*, *C. angustifolia*, *C. longa*, *Hedychium* sp., *Zingiber casumunar*, *Z. mioga*, and *Z. zerumbet* in Japan and India.
- CYANOTIS.** Creeping or ascending woody herbs much like *Tradescantia*.  
**Uredo davaoensis** Syd. Powdery brown rust pustules on leaves of *C. tuberosa* and *C. zeylanica* in India, Ceylon, and the Philippines.  
**Uromyces commelinae** Cke. See Commelina.
- CYATHEA.** TREE FERN.  
**Griggsia cyathea** Stev. Numerous irregular slightly elevated black spots on leaves of *C. arborea* in Porto Rico.  
**Pachypatella alsophilae** (Rac.) Theiss. and Syd. On leaves of *C. caudata* and *Alsophila contaminans* in Java and the Philippines.  
**Phyllachora hieronymi** P. Henn. Shiny black stromata on leaves of *C. dregei* in Nyassa and the Union of South Africa.
- CYBELE.** See Stenocarpus.
- CYCAS.** FERN PALM. SAGO PALM. Cycads.  
**Cladosporium cycadis** Marc. Small circular yellow leaf spots with dark margins and with a greenish mold forming beneath, on *C. revoluta* in Italy.  
**Dendrophoma clypeata** D. Sacc. Irregular, brown leaf spots on *C. revoluta* in Italy.  
**Hendersonia togniniana** Pöhl. On leaves of *C. revoluta* in Italy.  
**Pestalozzia cycadis** Allesch. On leaves of *C. revoluta* in Monaco.  
**Phyllosticta cycadina** Pass. Brown leaf spots on *C. revoluta* in Italy.  
**Septoria montemartinii** Pöhl. On leaves and petioles of *C. revoluta* in Italy.
- CYCLAMEN.** Herbaceous plants with flat tubers or corms, grown under glass for their flowers.  
**Phyllosticta cyclaminella** Bub. On leaves of *C. neapolitanum* in Yugoslavia.  
**Phyllosticta cyclaminis** Brun. Circular brown spots finally destroying plants of *C. europaeum* and *C. persicum* in France and Italy.  
**Septoria cyclaminis** Dur. and Mont. Large irregular smoky leaf spots with rufous margins on *C. europaeum*, *C. hederifolium*, and *C. repandum* in Algeria, Dalmatia, and Italy.
- CYCLANTHERA.** Annual and perennial herbs.  
**Cercospora cucubitina** Speg. Small, circular to irregular, then confluent, white leaf spots on *Cyclanthera* in Brazil.
- CYDISTA.** Ornamental vines cultivated for their flowers. (See also Bignonia).  
**Puccinia adenocalymnatis** (P. Henn.) Arth. Leaf rust on *C. aequinoctialis* in Cuba and Trinidad.  
**Puccinia aequinoctialis** Holw. Cinnamon-brown rust pustules on circular, yellow leaf spots on *C. aequinoctialis* in Cuba.  
**Puccinia cuticulosa** (E. and E.) Arth. Brown leaf rust on *C. aequinoctialis* in Porto Rico, Cuba, Brazil, and Nicaragua.
- CYDONIA.** QUINCE. Fruit trees.  
**Cercospora cydoniae** Rangel. On leaves of *C. oblonga* (*C. vulgaris*) in Brazil.  
**Cercospora tomenticola** (Thuem.) Sacc. On leaves of *C. vulgaris* in Brazil and Italy.  
**Exoascus bullatus** (B. and Br.) Fekl. See *Pyrus*.  
**Gymnosporangium koreaense** (P. Henn.) Jacks. See *Juniperus*.  
**Gymnosporangium mespili** (DC.) Kern. See *Juniperus*.  
**Gymnosporangium photiniae** (P. Henn.) Syd. See *Juniperus*.  
**Gymnosporangium spiniferum** Syd. Rust on leaves of *C. vulgaris* in Japan.  
**Helicobasidium mompa** Tan. See *Morus*.  
**Mycosphaerella pomacearum** Sacc. See *Malus*.  
**Ovularia necans** (Pass.) Sacc. Large spots causing premature leaf fall of *C. vulgaris* and *Mespilus germanica* in Italy and France.

**CYDONIA**—Continued.

- Phyllosticta cydoniae** (Desm.) Sacc. Circular to irregular brown leaf spots on *Chaenomelea japonica* (*Cydonia japonica*) and *C. vulgaris* in Italy, France, and Denmark.
- Phyllosticta cydoniaecola** Allesch. Large irregular leaf spots on *C. (Chaenomelea) japonica* in Italy and Germany.
- Phyllosticta cydonicola** P. Henn. Circular, then confluent, dull-brown leaf spots on *C. vulgaris* in Brazil.
- Phyllosticta velata** Bub. Circular to ellipsoid, often confluent, yellow-brown to red-brown leaf spots on *C. vulgaris* in Hungary.
- Ramularia tenuior** Fautr. and Brun. On leaves of *C. vulgaris* in France.
- Sclerotinia linhartiana** Prill. and Delacr. Brown leaf spots, twig die-back, and brown rot of fruit of *C. vulgaris* in Europe.
- Sclerotinia mespili** Woron. See *Mespilus*.
- Septoria cydoniae** Fekl. On leaves of *Chaenomelea japonica* (*Cydonia japonica*) and *C. vulgaris* in Italy, Austria, and Germany.
- Septoria cydonicola** Thuem. Pale-gray, irregular leaf spots on *C. vulgaris* in Italy and Malta.
- Trichoseptoria fructigena** Maubl. Causes a brown rot of fruit of *C. vulgaris*, *Crataegus* sp., and *Malus* (apple) in France and Germany.

**CYMBIDIUM**. See *Orchidaceae*.**CYMBOPOGON**. CITRONELLA GRASS. LEMONGRASS. Oil-producing grasses.

- Cerebella nardi** Butl. Black fungus areas in ovaries of *C. nardus* in India.
- Ophiodothis sclerotica** (Pat.) P. Henn. Black sclerotia in inflorescences of *C. nardus* and *C. schoenanthus* in India and Indo-China.
- Phyllachora andropogonis** Karst. and Har. See *Andropogon*.
- Puccinia cesatii** Schroet. See *Andropogon*.
- Puccinia cymbopogonis** Mass. Leaf rust on *C. citratus* in Uganda.
- Puccinia nakanishikii** Diet. See *Andropogon*.
- Uredo cymbopogonis-polyneuri** Petch. Leaf rust on *C. polyneuros* in Ceylon.
- Uromyces schoenanthi** Syd. Yellow to brown rust pustules on *C. schoenanthus* in India.
- Ustilago bengalensis** Syd. and Butl. Linear dark-brown smut sori in ovaries of *C. pendulus* in India.
- Ustilago mildbraedii** Syd. Powdery black smut masses replacing the ovaries of *C. schoenanthus* in tropical Africa.
- Ustilago nardi** Syd. Dark-brown elongate smut sori in ovaries of *C. nardus* in India.
- Ustilago schoenanthi** Syd. and Butl. Smut sori in inflorescences of *C. schoenanthus* in India.
- Ustilago spermoidea** B. and Br. Elongate smut sori in inflorescences of *C. martii*, *C. nardus*, and *C. venustum* in Ceylon.

**CYNANCHUM**. MOSQUITO TRAP. Herbaceous or semiwoody twiners.

- Cercospora bellyneckii** (West.) Sacc. On leaves of *C. acutum* and *C. vincetoxicum* in Italy, Belgium, China, and Spain.
- Cercospora miurae** Syd. On leaves of *C. caudatum* in Japan.
- Cercospora punctiformis** Sacc. and Roum. Yellow leaf spots on *C. acutum* in Russia and Algeria.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See *Paeonia*.
- Melampsora cynanchi** Thuem. Leaf rust on *C. sibiricum* in Siberia.
- Mycosphaerella albescens** Rabh. On leaves of *C. vincetoxicum* in Europe.
- Phyllosticta asclepiadeorum** West. and var. **minor** R. Rosal. Gray-white leaf spots on *C. vincetoxicum* and *Hoya carnosa* in Italy and Victoria.
- Phyllosticta atro-maculans** Speg. Large pale-yellow areas on leaves of *C. vincetoxicum* in Italy.
- Phyllosticta cynanchi** Brun. Circular to angular white leaf spots with brown margins on *C. acutum* in France.
- Phyllosticta viridi-tingens** Ferr. Small subcircular leaf spots on *C. vincetoxicum* in Italy.
- Phyllostictiella vincetoxici** (Sacc.) Tass. Sinuous white leaf spots with red margins on *C. vincetoxicum* in Denmark and Italy.
- Puccinia cynanchi** Lagh. Leaf rust on *C. parviflorum* and *Cynanchum* sp. in Martinique and Venezuela.
- Septoria asclepiadea** Sacc. On leaves of *C. vincetoxicum* and *Marsdenia erecta* in Italy, France, and Denmark.
- Septoria cynanchica** Sacc. On stems of *C. vincetoxicum* in Belgium.
- Septoria maculosa** Lévy. Black marginal leaf spots on *C. ercctum* in France.
- Septoria rajkoffi** Bub. On leaves of *C. acutum* in Bulgaria.
- Septoria vincetoxici** (Schub.) Awd. On leaves of *C. nigrum* and *C. vincetoxicum* in Russia and Denmark.

**CYNARA**. Artichoke. Cardoon. Perennial herbs.

- Ascochyta cynarae** Maffei. Circular ochraceous to gray-white leaf spots on *C. scolymus* in Italy.
- Phyllosticta cynarae** West. Subcircular zoned ashen-brown leaf spots on *C. scolymus* in Argentina, Italy, and Belgium.
- Ramularia cynarae** Sacc. Circular to irregular, gray blotches on leaves of *C. scolymus* and *Silybum marianum* in Argentina, French North Africa, Denmark, Russia, Italy, France, and Portugal. The leaves are browned and destroyed by the fungus, and heads fail to form.
- Septoria cardunculi** Pass. On leaves of *C. cardunculus* in Italy.

**CYNODON**. CAPRIOLA Ag. BERMUDA GRASS. Lawn and pasture grasses.

- Balansia** sp. On *C. dactylon* in the Union of South Africa.
- Cerebella cynodontis** Syd. Black, corrugated fungus masses in ovaries of *C. dactylon*, *Panicum distachyum*, and *P. prostratum* in Ceylon, India, the Philippines, and the Union of South Africa.
- Phyllachora cynodontis** (Sacc.) Niessl. (*Physalospora cynodontis* Del.) Black stromata on leaves of *C. dactylon* and *C. incompletus* in Japan, China, the Philippines, India, French North Africa, Egypt, Union of South Africa, Madeira, Yugoslavia, Bulgaria, Italy, Spain, France, Austria, and Germany.
- Puccinia cynodontis** Desm. Rust on leaves, sheaths, and culms of *C. dactylon* in Porto Rico, Guatemala, Ceylon, China, Japan, India, Turkestan, Persia, Australia, Algeria, Asia Minor, Russia, Spain, Portugal, France, Yugoslavia, Bulgaria, Italy, Austria, and Germany. Also reported from California and Florida.
- Puccinia varians** Diet. Powdery black rust pustules on leaves of *C. dactylon* in Japan.
- Septoria cynodontis** Fekl. On leaves of *C. dactylon* in Italy and Germany.
- Ustilago cynodontis** P. Henn. Black powdery smut masses destroying the inflorescences of *C. dactylon* and *C. glabratum* in Japan, China, Australia, Uganda, French North Africa, Abyssinia, Bulgaria, Portugal, Spain, Italy, and India.
- Ustilago dregeana** Tul. Black smut sori deforming peduncles and inflorescences of *C. dactylon* in the Union of South Africa.
- Ustilago paraguariensis** Speg. Smut in culms of *C. dactylon* in France, Brazil, and Paraguay.

- CYNOSURUS.** CRESTED DOG-TAIL. Ornamental lawn and pasture grasses.  
**Phyllosticta cynosuri** Gz. Frag. Leaf spots on *C. echinatus* in Spain.  
**Uromyces phyllachoroides** P. Henn. Brown to black rust sori on leaves of *C. cristatus*, *C. echinatus*, and *C. elegans* in French North Africa.
- CYPHERUS.** FLAT SEDGE. Papyrus. Aquatic or moist-land plants.  
**Ascochyta papyricola** Tass. On leaves of *C. papyrus* in Italy.  
**Cintractia cyperi-polystachyi** P. Henn. Smut in peduncles of *C. polystachyus* in the Philippines.  
**Cintractia javanica** Racib. Black powdery smut sori in inflorescences of *Cyperus* sp. in Java.  
**Cintractia peribebuyensis** Speg. Smut in peduncles of *C. distans* in India and Ceylon.  
**Cintractia tangensis** P. Henn. Black powdery smut masses in leaf axils of *Cyperus* sp. in Tanganyika.  
**Cintractia togoensis** P. Henn. Smut on *Cyperus* sp. in central Africa.  
**Helminthosporium cyperi** Bacc. On leaves of *C. dichostachys* in Abyssinia.  
**Kawakamia cyperi** (Miy. and Ideta.) Miy. Downy mildew causing browning of leaves and stems of *C. tegetiformis*, one of the matting sedges in Japan. The disease has also been found in Texas on imported material.  
**Phyllachora cyperi** Rehm. Black stromata on leaves and culms of *C. giganteus* and *C. longus* in Porto Rico, Spain, and Portugal. Also reported from Wisconsin.  
**Puccinia abrepta** Kern. Leaf rust on *C. ferax* in Costa Rica.  
**Puccinia antioquiensis** Mayor. Leaf rust on *C. diffusus* in Panama, Trinidad, and Colombia.  
**Puccinia concluda** Thuem. Black rust pustules on leaves of *C. longus* in Portugal.  
**Puccinia juncelli** Diet. Leaf rust on *C. serotinus* in Japan.  
**Puccinia philippinensis** Syd. Brown leaf rust on *C. compressus*, *C. rotundus*, and *C. polystachyus* in Japan and the Philippines.  
**Puccinia subcoronata** P. Henn. Ochraceous to dark-brown rust pustules on leaves of *C. malaccensis* in Brazil and Japan.  
**Schinzia aschersoniana** Magn. (*S. cypericola* [Magn.] De T.) Attacking roots of *C. flavescens* in Algeria.  
**Uredo cyperi** P. Henn. Rust on leaves and stems of *Cyperus* sp. in Abyssinia.  
**Uredo cypericola** P. Henn. Leaf rust on *C. capitatus*, *C. distans*, *C. rotundus*, and *C. tuberosus* in India and tropical Africa.  
**Uredo philippinensis** Syd. Cinnamon-brown rust pustules on leaves of *C. polystachyus* and *Cyperus* sp. in China and the Philippines.
- CYPHOMANDRA.** TREE TOMATO. Trees or shrubs, one species cultivated for its fruit.  
**Ascochyta cyphomandrae** Petch. Subcircular black, then gray, leaf spots on *C. betacea* in Ceylon.  
**Cronartium uleanum** Syd. Leaf rust on *Cyphomandra* sp. in Brazil.
- CYPRIPEDIUM.** LADY'S-SLIPPER. MOCCASIN FLOWER. Hardy terrestrial orchids. See Orchidaceae.
- CYRTOPODIUM.** See Orchidaceae.
- CYSTOPTERIS.** (*Filix*.) BLADDER FERN.  
**Aphelenchus olesistis** Ritz. Bos. See Begonia.
- CYTISUS.** BROOM. Shrubs cultivated for their profuse yellow or purple flowers.  
**Ascochyta cytisi** Lib. On *C. laburnum* (*Laburnum vulgare*) in Switzerland.  
**Ascochyta laburni** Sacc. On branches of *C. laburnum* (*Laburnum vulgare*) in France and Italy. *A. laburni* Kab. and Bub. is described on *C. laburnum* from Bohemia.  
**Ceratophorum setosum** Kirchner. Small brown, then dark-brown to black, spots enlarging rapidly and often confluent to form large, irregular, often concentric, ringed areas on leaves of *Cytisus* sp., *Lupinus cruihshanksii*, *L. mutabilis*, *L. polyphyllus*, and other species in Japan and Europe.  
**Cucurbitaria laburni** Pers. On twigs of *Cytisus* sp. in Europe.  
**Heterosporium laburni** Oud. Brown leaf spots on *C. laburnum* in Denmark and Holland.  
**Leptosphaeria corrugans** Rehm. Subcircular, oblong, or irregular yellow leaf spots with red-brown margins on *C. alpinus* (*Laburnum alpinum*) in Austria.  
**Marsonia carnea** Vestegr. Large circular leaf spots on *C. laburnum* (*Laburnum vulgare*) in Europe.  
**Mycosphaerella laburni** (Pass.) Lind. On leaves of *C. laburnum* and *C. ramentaceus* in Italy and Dalmatia.  
**Peronospora cytisi** Rostr. Downy mildew, causing brown leaf spots and destroying young plants of *C. alpinus* (*Laburnum alpinum*) and *C. laburnum* in Europe.  
**Phyllosticta coniothyrioides** Sacc. Subcircular ashen leaf spots on *C. laburnum* and *C. nigricans* in France and Denmark.  
**Phyllosticta cytisella** Sacc. Angular white leaf spots on *C. nigricans* in Italy.  
**Phyllosticta cytisi** Desm. Subcircular dull-brown leaf spots on *C. laburnum* in France, Great Britain, Italy, Belgium, and Austria.  
**Phyllosticta cytisorum** Pass. On leaves of *C. laburnum* in France.  
**Phyllosticta laburni** Oud. Irregular gray-white leaf spots on *C. laburnum* in Holland.  
**Phyllosticta laburnicola** Sacc. On leaves of *C. laburnum* in Italy.  
**Septoria cytisi** Desm. Numerous small subcircular white leaf spots with brown margins on *C. alpinus*, *C. capitatus*, *C. hirsutus*, *C. laburnum*, *C. nigricans*, and *Genista tinctoria* in Italy, France, Switzerland, Denmark, and Austria.  
**Septoria laburni** Pass. On leaves of *C. sessilifolius* in Italy.  
**Uromyces fulgens** (Hazsl.) Bub. Brown rust pustules on leaves and branches of *C. austriacus* and *C. hirsutus* in Yugoslavia, Bulgaria, Italy, Austria, and Germany.  
**Uromyces genistae-tinctoriae** (Pers.) Wint. See Caragana.  
**Uromyces pallidus** Nicssl. Yellow-brown rust pustules on leaves of *C. capitatus*, *C. hirsutus*, and *C. prostratus* in Europe.
- DACTYLIS.** ORCHARD GRASS. Forage grasses.  
**Apiospora lloydii** (Crouan) Sacc. See Holcus.  
**Aplanobacter rathayi** Sm. A viscid lemon-yellow slime envelops the upper leaves, stems, and inflorescences of *D. glomerata* in Denmark and Austria, causing a dwarfing and premature drying out of infected plants.  
**Cladochytrium graminis** Büsg. See Festuca.  
**Dilophia graminis** (Fckl.) Sacc. See Triticum.  
**Leptosphaeria nigrans** (Desm.) Ces. and De N. See Aira.  
**Mycosphaerella recutita** (Fr.) Johans. See Aira.  
**Ovularia pulchella** (Ces.) Sacc. and var. **lolii-italici** Sacc. Red leaf spots on *D. glomerata* and *Lolium italicum* in France and Italy.  
**Phyllachora bromi** Fckl. See Bromus.  
**Phyllosticta dactyloides** Gz. Frag. On leaves of *D. glomerata* in Spain.  
**Puccinia dactylidina** Bub. Powdery brown to black rust pustules on leaves of *D. glomerata* in India, Hungary, and Bohemia.  
**Septoria culmifida** Lind. See Phleum.  
**Sphaerodopsis dactylidis** (Delacr.) Theiss. and Syd. Black stromata on leaves of *D. glomerata* in France and Spain.

**DACTYLIS**—Continued.

- Uromyces dactylidis** Oth. Yellow and brown to black rust pustules on leaves of *D. glomerata* and *Ranunculus* spp. in Europe and New Zealand. Reported from Virginia.
- Ustilago dactylidis** Maire. Smut on *D. glomerata* in French North Africa.
- DAEDALACANTHUS.** Tropical shrubs with blue or rose-colored flowers.
- Isariopsis acanthacearum** Cke. Leaf spot of *D. nervosus* and *Eranthemum* sp. in Great Britain.
- Puccinia polystegia** Syd. Brown rust pustules on brown sunken leaf spots on *D. nervosus* in China.
- DAEMONOROPS.** See *Palmae*.
- DAHLIA.** Stout perennial herbs with fleshy roots and profuse autumn bloom.
- Accidium dahliae** Syd. Leaf rust on *D. variabilis* in Mexico.
- Bacillus dahliae** Hori. and Bakis. Bacterial rot of *Dahlia* sp. in Japan.
- Cercospora** sp. A leaf spot of *D. variabilis* in Argentina.
- Cercospora grandissima** Rangel. Leaf spot on *D. variabilis* in Brazil.
- Coleosporium dahliae** Arth. A rust producing yellow powdery pustules on the undersides of leaves of *D. variabilis* in Mexico.
- Cytospora dahliae** Peyl. On stems of *Dahlia* sp. in Bohemia.
- Entyloma calendulae dahliae** Sternon. Minute spots appear on the leaves at flowering time. These spots are at first few in number, 5 to 7 millimeters in diameter, limited by the principal veins and visible only on the upper leaf surfaces. The spots soon enlarge, becoming confluent, brown in color with dark-brown margins, and finally involve the entire leaf area as well as the petioles. The original dead areas fall out, giving a shot-hole appearance to the leaves. The loss of all or most of the leaf area reduces or even entirely prevents the development of flowers. On *Dahlia* (cult.) in Belgium. See, also, *Calendula*.
- Entyloma dahliae** Syd. Smut sori in pale circular to elliptical leaf spots on *D. variabilis* in the Union of South Africa.
- Ophiobolus georginae** (Fekl.) Sacc. On stems of *Dahlia* sp. in Germany.
- Phoma fuscata** Sacc. On stems of *Dahlia* sp. in Germany.
- Phyllosticta dahliaeicola** Brun. Produces large brown spots on leaves of cultivated varieties of *D. variabilis* in France, Switzerland, Italy, Austria, and Germany.
- Sporotrichum dahliae** Thuem. On stems of *D. variabilis* in Siberia.
- Urophlyctis** sp. Causes a leaf spot of *Dahlia* [cult.] in Kenya Colony.
- Verticillium dahliae** Kleb. Causes a wilt disease of the *Dahlia* in Germany. The fungus is said to differ from *V. alboatrum*, the cause of a serious wilt disease of the potato and other economic plants. The symptoms of the *Dahlia* disease are typical of wilt diseases, namely, a preliminary drooping of the leaves which finally involves the entire plant. The disease can be carried in the roots.
- DALBERGIA.** ROSEWOOD. Tropical trees and shrubs.
- Catacauma dalbergiae** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *D. acanthophylla*, *D. armata*, *D. ferruginea*, and *D. variabilis* in Brazil, the Union of South Africa, and the Philippines.
- Endodothella natalensis** Doidge. On leaves of *D. armata* in the Union of South Africa.
- Fomes caryophylli** (Rac.) Bres. See *Cinnamomum*.
- Fomes lamaoensis** Murr. See *Hevea*.
- Phyllachora dalbergiae** Niessl. Black stromata on leaves of *D. ferruginea*, *D. lactuca*, *D. sissoo*, and *D. variabilis* in India, the Philippines, and Brazil.
- Phyllachora lactea** Theiss. and Syd. Black stromata on red-brown leaf spots on *D. lactea* in Tanganyika.
- Phyllachora perforans** (Rehm.) Sacc. and Syd. Tar spot on leaves of *D. acanthophylla* in Brazil.
- Phyllachora spissa** Syd. Circular shiny black stromata on brown leaf spots on *D. sissoo* in India.
- Phyllosticta dalbergiae** Syd. Circular to irregular, often confluent, leaf spots with pale purple margins on *Dalbergia* sp. in Brazil.
- Phyllosticta dalbergicola** Syd. Circular pale-brown leaf spots on *D. variabilis* in Brazil.
- Phyllosticta nivea** Syd. Subcircular brown leaf spots on *Dalbergia* sp. in Brazil.
- Puccinia periodica** Racib. Leaf rust on *Dalbergia* sp. and *Derris* sp. in Java.
- Sphaerophragmium dalbergiae** Diet. Yellow to brown rust pustules on leaves of *D. amerimum* and *D. armata* in Cuba and Natal.
- Uredo dalbergiae** P. Henn. Ochraceous rust sori on *D. variabilis* in Brazil.
- Uredo mararyensis** P. Henn. Leaf rust on *Dalbergia* sp. in Brazil.
- Uredo nidulans** Syd. Powdery ochraceous rust pustules on *D. foliolosa* in Bolivia.
- Uredo sissoo** Syd. and Butl. Powdery, yellowish rust pustules on leaves of *D. sissoo* in India and Ceylon.
- Uromyces achrous** Syd. Leaf rust on *D. sissoo* in India.
- A serious canker disease due to an as yet undetermined fungus, attacks *D. latifolia* in Java.
- DALEA. PARSLEY.** Ag. Leguminous herbs and small shrubs, grown for their various-colored flowers.
- Puccinia daleae** Diet. and Holw. Cinnamon-brown to black rust pustules on leaves of *D. citriodora* in Mexico.
- DALECHAMPIA.** Climbing tropical shrubs, cultivated for their ornamental bracts.
- Accidium cornu-cervi** P. Henn. A rust, forming woody galls up to 10 centimeters in diameter on branches of *Dalechampia* sp. in Brazil.
- Accidium dalechampiicola** P. Henn. Leaf rust on *Dalechampia* sp. in Brazil.
- Caecoma theissenii** Syd. Leaf rust on small subcircular spots on *Dalechampia* sp. in Brazil.
- Didymella sphaerelloides** Sacc. and Syd. Indefinite pale brown sunken leaf spots on *D. ficifolia* in Brazil.
- DAMMARA.** See *Agathis*.
- DAPHNE.** Ornamental woody plants with handsome foliage and sweet-scented flowers.
- Colletotrichum daphnes-ponticae** Siem. Anthracnose on *D. pontica* in Caucasia.
- Dothidella mezerei** (Fr.) Theiss. and Syd. On stems of *D. alpina* in Europe.
- Gloeosporium mezerei** Cke. Anthracnose on leaves on *D. mezereum* in Great Britain and Australia.
- Marssonina daphne** (Rob. and Desm.) Magn. Irregular greenish, then dull-brown, leaf spots on *D. mezereum* in France, Italy, Switzerland, Denmark, Austria, and Holland.
- Marssonina obtusata** Maubl. Small raised ochraceous, then dull dark-brown leaf spots on *D. laureola* in France.
- Mycosphaerella gnidii** Maire. On leaves of *D. gnidius* in Algeria.
- Mycosphaerella laureolae** Desm. Subcircular, yellowish leaf spots on *D. laureola* in France, Switzerland and Germany.
- Phyllohendersonia daphnes** (Pass.) Tass. On leaves of *D. indica* in France.
- Phyllosticta carniolica** Voss. On leaves of *D. blagayana* in Austria.
- Phyllosticta daphnes ponticae** Siem. Leaf spot on *D. pontica* in Caucasia.

**DAPHNE**—Continued.

**Phyllosticta laureolae** Desm. Subcircular to irregular leaf spots on *D. laureola* in Denmark, France, and Portugal.

**Septoria daphnes** Desm. Irregular indefinite greenish leaf spots on *D. mezereum* in France and Germany.

**Uredo daphnicola** Diet. Brown leaf rust on *Daphne* sp. in Hongkong.

**DALENYPHYLLUM**. Broadleaved evergreen, hardy or semitropical shrubs, and small trees.

**Leptothyrium yoshinagai** P. Henn. On leaves of *D. glaucescens* in Japan.

**Mycosphaerella daphniphylli** Syd. and Hara. Circular to irregular spots on leaves of *D. glaucescens* in Japan.

**DATISCA**. Tall perennial herbs.

**Coleosporium datiscæ** Tranzsch. Golden rust pustules on leaves of *D. cannabina* in Kashmir and Caucasia.

**Phyllosticta datiscæ** Syd. Irregular, then confluent, often marginal, dull brown leaf spots on *D. cannabina* in Russia.

**DATURA**. ANGEL'S-TRUMPET. JIMSON WEED. Rank-growing annual or perennial herbs or shrubs mostly weeds.

**Ascochyta daturæ** Sacc. Subcircular gray-white leaf spots on *D. arborea* and *D. stramonium* in Russia, Denmark, Italy, and Portugal.

**Septoria daturæ** Speg. Definite circular gray-white leaf spots on *D. inermis* and *D. stramonium* in Argentina and Italy.

**DAUCUS**. CARROT. Annual and biennial herbs.

**Aecidium carotinum** Bub. Leaf rust on *D. carota* in Bohemia.

**Aphelenchus modestus** J. M. Attacks the roots of *D. carota* in Europe.

**Cercospora pastinacæ** Karst. See Pastinaca.

**Erysiphe taurica** Lév. See Althaea.

**Helicobasidium mompa** Tan. See Morus.

**Hypochnus cucumeris** Frank. See Cucumis.

**Leptosphaeria rostrupii** Lind. Gray-brown cankers on the roots, which work up into the stems of *D. carota* in Sweden and Denmark.

**Mycosphaerella sagedioides** (Wint.) Lind. On stems of *Daucus* sp. and *Dipsacus* sp. in Switzerland.

**Phyllachora pastinacæ** Rostr. (*Septoria pastinacæ* West.) Leaf spot and stem canker of *D. carota* in Great Britain and Denmark.

**Protomyces macrosporus** Ung. See Coriandrum.

**Ramularia pastinacæ** Bub. See Pastinaca.

**Septoria carotæ** P. Nag. On leaves of *D. carota* in Russia.

**Septoria daucina** Brun. Subcircular to irregular, then confluent, brown leaf spots on *D. carota* in France.

**DEERINGIA**. Climbing herbs or subshrubs.

**Aecidium deeringiae** Cke. and Mass. Leaf rust on *D. celosioides* in Australia.

**Puccinia calosperma** Syd. and Butl. Rust on leaf blades, petioles, stems, and flowers of *D. celosioides* in India.

**Uromyces deeringiae** Syd. Brown leaf rust on *D. indica* in Java, Japan, and the Philippines.

**DELPHINIUM**. LARKSPUR. Hardy herbs cultivated for their flowers.

**Cercospora delphinii** Thuem. Dull-yellow subcircular leaf spots on *D. elatum* in Siberia. Also reported from Colorado.

**Coleosporium martianoffianum** Syd. Golden-yellow rust pustules on leaves of *D. intermedium* in Siberia.

**Phyllosticta ajacis** Thuem. Irregular, often marginal, gray-white leaf spots on *D. ajacis* in Austria.

**Rhizoctonia destruens** Tassi. See Solanum.

**Ramularia delphinii** Jaap. Leaf spot on *D. elatum* in Switzerland.

**Septoria staphysagriae** Wint. Circular to irregular definite brown leaf spots on *D. staphysagria* in Portugal and Italy.

**DENDROBIUM**. See Orchidaceae.**DENDROCALAMUS**. See Bambuseae.**DENDROPANAX**. Tropical trees and shrubs.

**Mycosphaerella didymopanicis** Miles. Light-brown to gray, circular, sometimes confluent, leaf spots with narrow orange-brown margins on *D. arboreum* in Porto Rico.

**Phyllosticta araliana** Young. Leaf spot on *D. arboreum* in Porto Rico and Cuba.

**DENTARIA**. TOOTHWORT. Small, early-flowering herbs.

**Phyllosticta dentariae** Kab. and Bub. Greenish, then pale-brown to white, irregular leaf spots on *D. enneaphylla* in Bohemia.

**Peronospora dentariae macrophyllae** Gäum. Downy mildew on leaves of *D. macrophylla* in Japan.

**Puccinia dentariae** (Alb. and Schw.) Fckl. Leaf and stem rust on *Dentaria* spp. in Europe. Also reported from Oregon.

**DERRIS**. Tall, tropical trees or climbers.

**Cercospora pumila** Syd. On leaves of *Derris* sp. in the Philippines.

**Diorcidium koordersi** Wurth. Leaf rust on *D. elliptica* in Java.

**Dothidella derridis** (P. Henn.) Theiss. On leaves of *Derris* sp. and *D. elliptica* in central Africa and the Philippines.

**Fomes lignosus** Klotzsch. See Hevea.

**Hapalophragmium derridis** Syd. Brown rust pustules on leaves of *D. uliginosa* in Ceylon and central Africa.

**Phyllachora affinis** Theiss. and Syd. Black stromata on leaves of *D. philippinensis* in the Philippines.

**Phyllachora derridis** Syd. Black stromata on indefinite brown leaf spots on *D. glabrata* in Angola.

**Phyllachora luzonensis** P. Henn. See Milletia.

**Phyllachora ramosii** (P. Henn.) Theiss. and Syd. Numerous, black stromata on small, leaf spots on *Derris* sp. in the Philippines.

**Phyllachora yapensis** (P. Henn.) Syd. Black circular to irregular stromata on leaves of *D. diadelpa*, *D. elliptica*, and *D. philippinensis* in the Philippines, Java, and the island of Yap.

**Phyllosticta derridis** P. Henn. Circular or effuse, pale-brown, marginal leaf spots on *Derris* sp. in the Congo.

**Placosphaeria derridis** P. Henn. Circular leaf spots on *Derris* sp. in the Congo.

**Placosphaeria merrillii** P. Henn. Circular effused yellow or dull brown leaf spots on *Derris* sp. in the Philippines.

**Puccinia periodica** Racib. See Dalbergia.

**Triphragmium pulchrum** Racib. Brown rust pustules on leaves of *D. elliptica* in Java.

**Uredo derridis** P. Henn. Leaf rust on *D. bantamensis* in Java.

**DESCHAMPSIA.** See also Aira.

**Tilletia schenckiana** P. Henn. Smut sori deforming and destroying the ovaries of *D. antarctica* in Kerguelen Island.

**DEUTZIA.** Ornamental shrubs, grown for their showy, white or bluish flowers.

**Aecidium deutziae** Diet. Leaf rust on *D. gracilis*, *D. scabra*, *D. sieboldiana*, and *D. staminea* in Japan and India.

**Ascochyta deutziae** Bres. Irregular, dull-yellow leaf spots on *D. scabra* in Germany.

**Cylindrosporium deutziae** Syd. Indefinite confluent yellow-brown leaf spots on *D. scabra* in Japan.

**Leptosphaeria dichroa** Pass. On branches of *D. scabra* in Italy.

**Mycosphaerella deutziae** Syd. On leaves of *D. lemoinei* in Germany.

**Phyllosticta deutziicola** Petr. Large irregular brown leaf spots on *Deutzia* sp. in Bohemia and Austria.

**Septoria phyllostictoides** Sacc. Indefinite gray-white leaf spots on *D. scabra* in France.

**DIANELLA.** Tender perennial rhizomatous plants.

**Uredo dianella** Diet. Leaf rust on *D. ensifolia* and *D. nemorosa* in China, Hongkong, Ceylon, Java, and Japan.

**DIANTHUS.** PINK. CARNATION. Herbs cultivated for their showy flowers.

**Ascochyta dianthi** (A. and S.) Berk. Large, subcircular, brown patches on leaves of *D. caryophyllus*, *Lychnis* sp. and *Saponaria* sp. in Great Britain, Italy, Holland, and Germany.

**Fusarium udum** Butl. This fungus causes a wilt of seedlings of *Dianthus* sp., *Anacardium occidentale*, *Antirrhinum* sp., *Eriobotrya japonica*, *Eugenia jambos*, and *Grevillea robusta* in Uganda. See also Cajanus.

**Gloeosporium dianthi** Cke. Small circular purple confluent leaf spots with brown centers on *D. caryophyllus* in Great Britain.

**Phomopsis caryophylli** Grove. On calyces, peduncles, and stems of *D. caryophyllus* in Japan and Great Britain.

**Phyllosticta dianthi** West. Irregular gray-white blotches on leaves of *D. barbatus* in Belgium and France.

**Phyllosticta dubia** Sacc. On leaves of *D. chinensis* (*D. sinensis*) in Malaya.

**Placosphaeria stellariae** (Lib.) Sacc. See *Stellaria*.

**Pseudodiscosia dianthi** Host. and Laub. Large brown, then light gray spots on leaves, flower stalks and stems of *D. caryophyllus* in Germany.

**Puccinia dianthi-japonici** P. Henn. Brown rust pustules on circular yellow leaf spots on *D. japonicus* and *D. nipponicus* in Japan.

**Puccinia fastidiosa** Sacc. and De T. Powdery, ochraceous rust pustules on leaves and stems of *D. chinensis* in Siberia.

**Ramularia dianthi** Lind. Large leaf spots with brown margins on *D. carthusianorum* in Germany.

**Septoria carthusianorum** West. Irregular, dull-yellow spots on calyces of *D. carthusianorum* in Belgium.

**Septoria caryophylli** Scalia. Irregular elongate ochraceous spots with dull brown margins on leaves and stems of *D. caryophyllus* in Italy.

**Septoria dianthicola** Sacc. On leaves of *D. barbatus*, *D. caryophyllus*, and *Tunica prolifera* in Italy, Switzerland, and Portugal.

**Septoria dianthophila** Speg. and f. **hispanica** Gz. Frag. On leaves of *D. caryophyllus* and *D. (Tunica) prolifera* in Brazil and Spain.

**Septoria sinarum** Speg. Large subcircular white leaf blotches on *D. chinensis* in Italy and Great Britain.

**Uredo dianthicola** Har. Leaf rust on *D. caryophyllus* in France. (Doubtful species.)

**Urocystis purpurea** Hazsl. Purple masses of smut spores replacing the ovaries of *D. deltooides* and *D. (Tunica) prolifera* in Hungary.

**Uromyces formosus** Syd. Brown rust pustules on stems and leaves of *D. crinitus*, *D. libanotis*, and *D. macranthoides* in Persia.

**DIAPENSIA.** Small, compact, tufted, woody perennials.

**Phyllosticta diapensiae** Pat. Leaf spots on *D. lapponica* in China.

**Physalospora diapensiae** Rehm. On leaves of *D. lapponica* in Lapland.

**Septoria diapensiae** Karst. On leaves of *D. lapponica* in Lapland.

**DIASIA.** See *Melasphaerula*.

**DICENTRA. BIKUKULLA.** Ag. BLEEDINGHEART. Herbaceous perennials grown for their flowers and cut foliage. Sometimes called *Dielytra* or *Dicylra*.

**Ascochyta dicentrae** Oud. On branches of *D. spectabilis* in Holland.

**DICHAEA.** See *Orchidaceae*.

**DICHORISANDRA.** Tropical perennial herbs, grown for their handsome flowers and foliage.

**Colletotrichum dichorisandrae** Rangel. Anthracnose on *D. thyrsiflora* in Brazil.

**Uromyces dichorisandra** P. Henn. Leaf rust on *Dichorisandra* sp. in Brazil.

**DICHROA.** Hydrangealike shrubs.

**Endophyllum dichroae** Rac. Rust causing hypertrophy of leaves of *D. cyanitis* in Java.

**DICLIPTERA.** Annual or perennial herbs or subshrubs with red or blue bracted flowers.

**Puccinia diclipterae** Syd. Brown rust on leaves of *D. longiflora* in Formosa.

**Uromyces tweediana** (Speg.) Arth. Yellow-brown to chestnut-brown rust pustules on leaves and stems of *D. maculata*, *D. squarrosa*, *D. tweediana*, and *Dicliptera* sp. in Mexico, Brazil, Argentina, Abyssinia, and India.

**DICTAMNUS.** GAS PLANT. Semiwoody plants.

**Ascochyta nobilis** Kab. and Bub. Circular to angular, often confluent, yellow-white or gray leaf spots with red margins on *D. albus* (*D. frazinella*) in Russia and Bohemia.

**Colletotrichum dictamni** Höll. Circular to irregular dull-brown leaf spots on *D. albus* in Hungary.

**Septoria dictamni** Fekl. Discolored leaf patches on *D. albus* (*D. frazinella*) in Russia and Italy.

**DICTYOSPERMA.** See *Palmae*.

**DIEFFENBACHIA.** TUFT ROOT. Large-leaved tropical aroids.

**Gloeosporium thuenenii** Sacc. See *Anthurium*.

**Phyllosticta colocasiae** v. Hoeh. See *Colocasia*.

**DIERAMA.** South African cormous plants with large spikes of flowers.

**Puccinia dieramae** Syd. Leaf rust on *D. ensifolia* in the Union of South Africa.

**DIERVILLA.** BUSH HONEYSUCKLE. Ornamental deciduous shrubs with showy flowers. Certain Asiatic species sometimes placed under *Weigela*.

**Ascochyta diervillae** Kab. and Bub. Circular to irregular, then confluent, brown leaf spots on *D. loniceria* (*D. canadensis*) in Bohemia.

**Ascochyta weigeliae** Sacc. and Speg. Angular gray-white leaf spots on *D. florida* (*D. rosea*) in Italy.

**Phyllosticta weigeliae** Sacc. and Speg. Gray-white leaf spots on *D. florida* (*D. rosea*) in Denmark, Italy and France.

**D. ERVILLA**—Continued.

- Phyllosticta weigeliina** Bub. and Kab. Leaf spots on *D. florida* in Bohemia.
- Ramularia weigeliae** Speg. On leaves of *D. florida* in Italy.
- Septoria weigeliae** Kab. and Bub. Leaf spots on *D. florida* in Bohemia.
- DIGITALIS**. FOXGLOVE. Hardy herbaceous perennials with long racemes of flowers. The leaves of some species used medicinally.
- Ascochyta digitalis** Fekl. Purplish leaf spots on *D. purpurea* in Portugal and Germany.
- Gloeosporium digitalidis** Rostr. Large dull-brown leaf spots on *D. purpurea* in Denmark.
- Mycosphaerella mariae** Sacc. and Bomm. On leaves and stems of *D. ferruginea* and *D. lutea* in Italy.
- Peronospora digitalidis** Gäum. Downy mildew on leaves of *D. ambigua*, *D. lutea*, and *D. purpurea* in central Europe.
- Phyllosticta digitalis** Bell. On leaves of *D. lutea* and *D. purpurea* in Italy and Alaska.
- Ramularia variabilis** Fekl. Irregular brown leaf spots starting on the basal leaves and progressing upward on *D. purpurea* in Italy, Denmark, Austria, and Germany. The medicinal value of the leaves is reduced or destroyed.
- Septoria digitalis** Pass. Irregular, dull-brown leaf spots on *D. ferruginea*, *D. grandiflora*, *D. lanata* and *D. lutea* in Russia, the Balkans, Italy, Great Britain, and Austria.
- DILLENIA**. Tall tropical trees.
- Cercospora dilleniae** Petch. Gray-brown leaf spots on *D. retusa* in Ceylon and India.
- DINOCHLOA**. See Bambuseae.
- DIOCLEA**. Tender woody twiners.
- Phyllachora diocleae** P. Henn. Black stromata on circular to irregular brown leaf spots on *Dioclea* sp. in Brazil.
- DIOSCOREA**. YAM. AIR POTATO. Tropical climbers with edible roots.
- Bagniopsis dioscoreae** Wakef. Produces a reduction in length of the internodes and in size of the leaves, giving a witches'-broom effect on *D. prehensilis* in Nigeria. The stromata are produced on stems and petioles, but not on the leaf blades which are, however, chlorotic.
- Catacauma glaziovii** (P. Henn.) Theiss. and Syd. Circular, shiny black stromata on leaves of *D. heptaneura*, *D. pachycarpa* and *D. polygonoides* in Brazil.
- Cercospora brasiliensis** Averna. Leaf spot on *Dioscorea* sp. in Brazil.
- Cercospora carbonacea** Miles. Angular, black leaf spots 1 to 1.5 centimeters in diameter on *D. alata* in Porto Rico and Cuba.
- Cercospora pachyderma** Syd. On leaves of *D. alata* and *D. esculenta* in the Philippines.
- Cercospora ubi** Rac. Circular leaf spots, yellow-brown above, gray-brown below, on *D. alata*, *D. daemona*, *D. esculenta*, and *D. glabra* in Java, China, Malaya, and the Philippines.
- Cercospora contraria** Syd. Gray-brown leaf spots on *Dioscorea* sp. in the Congo.
- Colletotrichum dioscoreae** Averna. Anthracnose on *Dioscorea* sp. in Brazil.
- Cylindrosporium dioscoreae** Miy. and I. Ito. Small, yellowish to brown spots on leaf blades, petioles and stems of *D. batatas* and *D. japonica* in Japan, Defoliation results.
- Ellisiodothis rehmaniae** Theiss. and Syd. Black stromata on stems of *D. esculenta* in the Philippines.
- Gloeosporium pestis** Mass. Anthracnose spots on leaves of *Dioscorea* sp. in Fiji and Russia.
- Hemileia dioscoreae-aculeatae** Rac. Yellow leaf rust on *D. aculeata* and other species in Uganda and Java.
- Laestadia perusta** (B. and Br.) Sacc. Circular dull-brown leaf spots on *D. tomentosa* and *Dioscorea* sp. in Ceylon and India.
- Mycosphaerella dioscoricola** Syd. Leaf spot on *D. esculenta* in the Philippines.
- Phyllachora ulei** Wint. Shiny black circular stromata on leaves of *Dioscorea* sp. in Brazil.
- Phyllosticta dioscoreacearum** Bacc. Subcircular gray leaf spots on *Dioscorea* sp. in India.
- Phyllosticta dioscoreacola** P. Brun. Irregular red-brown leaf spots on *D. batatas* in France.
- Phyllosticta dioscoreae-daemona** P. Henn. Circular pale-brown leaf spots with dark-brown margins on *D. daemona* in Brazil.
- Phyllosticta graffiana** Sacc. On leaves of *D. esculenta* and *D. pentaphylla* in the Philippines.
- Puccinia valida** Arth. Cinnamon-brown rust pustules on lower leaf surfaces of *D. convolvulacea* in Mexico.
- Rostrupia dioscoreae** (Kom.) Syd. Cinnamon-brown to black rust pustules on leaves of *D. quinqueloba* in Manchuria and Japan.
- Septoria versicolor** Pat. Ashen-white leaf spots with brown margins on *Dioscorea* sp. in Ecuador.
- Uredo dioscoreae** (B. and Br.) Petch. (*Uredo dioscoreae* P. Henn.) Leaf rust on *D. alata*, *D. bulbifera*, *D. esculenta*, *D. grandiflora*, *D. piperifolia* and *D. polygonoides* in Ceylon, the Philippines, Porto Rico, Cuba, Brazil, India, and Uganda.
- Uredo dioscoreae-alatae** Rac. Brown rust sori on leaf blades, petioles, and stems of *D. alata* and other species in Java and the Philippines.
- Uredo dioscoreae-filiformis** Rac. Rust pustules on dull-brown leaf spots on *D. filiformis* in Java.
- Uredo dioscoreae-pentaphyllae** Petch. Leaf rust on *D. pentaphylla* in Ceylon.
- Uredo dioscoreae-quinquelobae** P. Henn. Leaf rust on *D. quinqueloba* in Japan.
- Uredo dioscoreae-sativae** Syd. Brown rust pustules on leaves of *D. sativa* in India.
- Urocystis dioscoreae** Syd. Black, elongate, smut pustules on leaf blades, petioles and stems of *D. quinqueloba* and *D. tokoro* in Japan and Russia.
- DIOSPYROS**. PERSIMMON. Kaki. Ebony. Fruit and timber trees.
- Aecidium atro-album** P. Henn. Leaf rust on *Diospyros* sp. in the Union of South Africa and the Congo.
- Aecidium atrocrustaceum** Syd. Leaf rust on *D. discolor* in the Philippines.
- Aecidium calosporum** Juel. Rust on circular leaf spots on *Diospyros* sp. in Brazil.
- Aecidium diospyri** A. L. Sm. Leaf rust on *D. mespiliformis* in Angola.
- Aecidium melaenum** Syd. Leaf rust on *Diospyros* sp. in the Philippines.
- Aecidium ramosii** Syd. Leaf rust on *Diospyros* sp. in the Philippines.
- Aecidium reyesii** Syd. Leaf rust on *D. discolor* in the Philippines.
- Aecidium rhytismoideum** B. and Br. and var. **mabae** P. Henn. Rust on black leaf spots on *D. discolor*, *D. embryopteris*, *D. mespiliformis*, *D. ovalifolia*, *D. tomentosa*, and *Maba abyssinica* in Java, India, Ceylon, Abyssinia, and the Philippines.
- Aecidium ulei** P. Henn. Leaf rust on *Diospyros* sp. in Brazil.
- Botrytis diospyri** Brizi. A fruit rot of *D. kaki* in Italy, Russia, and Japan. The fungus also causes a leaf spot and die-back of twigs.
- Colletotrichum kaki** Maffei. Anthracnose on leaves of *D. kaki* in Italy.
- Cylindrosporium kaki** Syd. Circular dull-brown or gray leaf spots with black limiting lines on *D. kaki* in Japan.
- Fusicladium diospyrae** Hori. and Yosh. Circular, black spots with definite margins on leaves, fruit and buds of *D. kaki* in Japan, producing a curl of young leaves. On young shoots the spots are fusiform to oblong, dark-brown to black with sunken centers. Diseased twigs die-back and spotted fruit falls.

**DIOSPYROS**—Continued.

- Fusicladium kaki** Hori. and Yosh. On leaves of *D. kaki* in Japan.
- Fusicladium levieri** P. Magn. Circular brown leaf spots with black margins on *D. lotus* in Caucasia.
- Gloeosporium kaki** Ito. Small black spots, increasing in size and becoming sunken, on fruit of *D. kaki* in Japan. The spots may fuse and crack irregularly. Complete rot of the fruit ensues. Infections also occur on young shoots, showing typical blackish spots of elliptical shape, usually depressed. On stems of young plants all parts above the infected area die. Infected leaves fall, complete defoliation often resulting.
- Helicobasidium mompa** Tan. See Morus.
- Lophodermium diospyri** Pat. Long, black, superficial fruiting bodies on circular, often confluent, leaf spots on *D. ebenum* in Indo-China.
- Melasmia falcata** Syd. Black stromata on leaves of *Diospyros* sp. in Brazil.
- Myxosporium kaki** Hara. Dark-brown, more or less sunken areas, on stems, branches and rarely on fruit of *D. kaki* in Japan.
- Pestalozzia diospyri** Syd. Circular to irregular reddish-brown spots on leaves of *D. kaki* in China and Japan. Smaller brown spots, generally regular, also occur on the fruit.
- Plaeosaccardinula diospyricola** P. Henn. Black flat fruiting bodies on leaves of *Diospyros* sp. in Brazil.
- Phyllosticta erythraea** Bacc. Brown leaf spots with black margins on *D. mespilifolius* in Abyssinia.
- Septoria diospyri** McAlp. On leaves of *D. carginia* in New South Wales.
- Uredo peteloti** Pat. Brown rust pustules on circular leaf spots on *D. siamensis* in Indo-China.
- DIOTIS.** COTTONWEED. Perennial seaside plants.
- Puccinia diotidis** Pat. and Roum. Leaf rust on *D. candidissima* in France.
- DIPCADI.** Tender, bulbous, scapose plants.
- Aecidium dipcadi** Har. and Pat. Leaf rust on *D. udellensis* in the Congo.
- DIPHYSA.** Shrubs or trees.
- Calliospora diphysae** Arth. Powdery, black rust pustules on leaves and stems of *D. robinoides* and *D. suberosa* in Guatemala, Costa Rica, and Mexico.
- DIPLOGLOTTIS.** Australian trees.
- Uromyces diploglottidis** Cke. and Mass. Leaf rust on *D. cunninghamii* in Queensland.
- DIPLOSTEPHIUM.** Composites resembling *Aster*.
- Rhynchostoma biolleyana** Bomm. and Rouss. On leaves and stems of *D. costaricense* (*D. rupestre*) in Costa Rica.
- DIPLOTHEMIUM.** See Palmae.
- DIPSACUS.** TEASEL. Biennial or perennial herbs, used as fuller's teasels.
- Ascochyta dipsaci** Bub. Subcircular small ochraceous leaf spots with dull-brown margins on *D. pilosus* in Asia Minor.
- Mycosphaerella sagedioides** (Wint.) Lind. See *Daucus*.
- Peronospora violacea** Berk. Downy mildew on leaves of *Dipsacus pilosus*, *Knautia silvatica*, *Scabiosa columbaria*, and *Succisa pratensis* in Europe.
- Phyllosticta dipsaci** Br. and Fautr. Gray leaf spots on *D. pilosus* in France.
- Phyllosticta vandae** Namys. On leaves of *D. sylvestris* in Poland and Galicia.
- Ramularia silvestris** Sacc. On leaves of *D. fullonum* and *D. sylvestris* in Spain, Yugo-Slavia, Denmark, France, Bohemia, Belgium, and Germany.
- Septoria dipsaci** West. Small circular brown, then white leaf spots on *D. azureus* and *D. fullonum* in Belgium and Italy.
- Septoria fullonum** Sacc. Gray-white leaf spots on *D. fullonum* in Austria.
- Uredo involucrorum** Rabh. Leaf rust on *D. pilosus* in France, Belgium, and Switzerland.
- DISCHIDIA.** Evergreen trailing plants cultivated for their white flowers.
- Phyllachora dischidiae** Syd. Circular black shiny stromata on leaves of *D. rosca* in Chile and the Philippines.
- Uredo dischidiae** P. Henn. Yellow rust pustules on leaves of *Dischidia* sp. in New Guinea.
- DISPORUM.** FAIRY BELLS. Small perennial rhizomatous herbs.
- Aecidium dispori** Diet. Leaf rust on *D. sessile* in Japan.
- DISSOTIS.** Bristly-hairy herbs or shrubs.
- Puccinia dissotidis** P. Henn. Yellow-brown rust sori on circular dull-brown leaf spots on *Dissotis* sp. in the Congo.
- Puccinosira dissotidis** Wakef. (*Aecidium dissotidis* Cke.) Brown leaf rust on *D. incana* and *D. princeps* in Uganda, the Congo, and the Union of South Africa.
- Uredo dissotidis-longicaudae** P. Henn. Brown leaf rust on *D. longicaudata* in the Union of South Africa.
- DIURIS.** See Orchidaceae.
- DODONAEA.** HOP BUSH. Ornamental trees and shrubs.
- Uredo dodonaeae** Koord. Leaf rust on *D. viscosa* in Java.
- DOLICHOLUS.** See Rhynchosia.
- DOLICHOS.** HYACINTH BEAN. Tropical twiners.
- Cercospora wildemanii** Syd. Circular brown leaf spots with purple margins on *Dolichos* sp. in the Congo.
- Helminthosporium accedens** Syd. On leaves of *D. baumii* in the Union of South Africa.
- Helminthosporium dolichi** Syd. On leaves of *D. euryphyllus* in the Union of South Africa.
- Phyllachora dolichogena** (B. and Br.) Sacc. and var. **samoensis** Theiss. and Syd. Black stromata on leaves of *D. lablab* and *Dolichos* sp. in Ceylon and Samoa.
- Phyllosticta dolichi** Brun. Brown leaf spots with dark margins on *D. lablab* and *D. myoides* in Russia and France.
- Physopella concors** Arth. See Phaseolus.
- Puccinia dolichi** Arth. Powdery brown rust pustules on leaves of *D. reticulatus* in Cuba.
- Rhizoetonia** sp. See Vigna.
- Septoria lablabina** Sacc. Circular dull-brown leaf spots on *D. lablab* in tropical Africa and the Philippines.
- Septoria lablabis** P. Henn. Circular, then confluent, dull-brown spots on leaves of *D. lablab* in Tropical Africa and the Philippines.
- Uromyces dolichi** Cke. Brown rust pustules on leaves of *D. axillaris* and *D. gibbosus* in the Union of South Africa.
- Uromyces kisantuensis** P. Henn. Powdery dark-brown rust pustules on leaves of *Dolichos* sp. in the Congo.
- Vermicularia capsici** Syd. See Capsicum.
- Woroninella dolichi** (Cke.) Syd. Orange galls on leaves of *D. lablab* and *Rhynchosia volubilis* in China and the Philippines.

- DOMBEYA. ASSONIA** Ag. Shrubs and small trees with large loose umbels of showy flowers.  
**Phyllachora dombeyae** Syd. Black stromata on leaves of *D. rotundifolia* and *D. schimperiana* in the Union of South Africa.
- DONAX.** See Arundo.
- DOREMA.** Large perennial herbs yielding gum and resins.  
**Puccinia doremae** Spechn. Leaf rust on *Dorema* sp. in Turkestan and Transcaspia.
- DORONICUM. LEOPARD'S-BANE.** Hardy herbs with yellow many-flowered heads.  
**Ascochyta daronici** Allesch. Large ovoid subochraceous to ashen-brown leaf spots on *D. caucasicum* in Germany.  
**Coleosporium daronici** Namysl. Yellow leaf rust on *D. austriacum* in Austria.  
**Mycosphaerella aronici** (Fekl.) Volk. On leaves of *D. austriacum*, *D. caucasicum*, *Aronicum scorpioides*, and *Carduus defloratus* in north Africa, Switzerland, Italy, and Austria.  
**Phyllosticta austriaca** Sacc. Brown leaf spots on *D. austriacum* in Italy.  
**Phyllosticta doromicella** Maire. Black leaf spots on *D. thirkeum* in Asia Minor.  
**Phyllosticta doromicigena** Bub. On leaves of *D. cordatum* in Hungary.  
**Puccinia daronicella** Syd. Brown rust pustules on leaves of *D. austriacum* and *D. cordatum* in Austria and Hungary.  
**Puccinia daronici** Niessl. Black rust sori on leaves of *D. austriacum* and *D. macrophyllum* in Bulgaria and Austria.
- Ramularia daronici** (Sacc.) Grove. On leaves of *D. pardalianches* in Great Britain, Denmark, and France.  
**Ramularia daronicum** Vogl. Subcircular to oblong, often marginal, yellow, then white, leaf spots on *D. clusium*, *D. cordatum*, and *D. scorpioides* in Italy and Spain.  
**Ramularia filaris** Fres. See Adenostyles.  
**Septoria czarnohorica** Nam. Circular to irregular and confluent brown leaf spots on *D. cordifolium* in France and Poland.
- DORSTENIA.** Tropical herbs and small shrubs.  
**Accidium bertonii** Speg. Leaf rust on *D. brasiliensis* in Paraguay.  
**Accidium buchwaldii** P. Henn. Rust on brown leaf spots on *Dorstenia* sp. in Tanganyika.  
**Accidium dorsteniae-holstii** P. Henn. Rust on circular dull-brown leaf spots on *D. holsti* in Tanganyika.  
**Puccinia dorsteniae** Lagh. Leaf rust on *D. psilurus* in Angola.  
**Uredo consanguinea** Syd. Brown leaf rust on *D. multiformis* in Brazil.  
**Uredo rubescens** Arth. Powdery brown rust pustules on leaves of *D. contrajerva* and *D. houstoni* in Porto Rico, Guatemala, and Trinidad.
- DOYALIS.** See Aberia.
- DRABA. WHITLOW GRASS.** Spring-flowering annuals.  
**Ascochyta drabae** Oud. On leaves of *D. alpina* in Russia.  
**Peronospora norwegica** Gäum. Downy mildew on leaves of *D. hirta* in Norway.  
**Puccinia drabae** Rud. Brown rust pustules on leaves, stems, and flowers of *Draba* spp. in Europe and Asia Minor. Reported from Utah and New York.
- DRACAENA. (including Cordyline.) DRACENA. DRAGON TREE.** Variegated-leaved shrubs.  
**Cercospora cordylines** P. Henn. On leaves of *C. dracaenoides* in Argentina and Brazil.  
**Colletotrichum cordylines** Pöhl. Anthracnose on leaves of *C. indivisa* in Italy.  
**Colletotrichum dracaena** Allesch. On leaves of *D. latifolia* in Germany.  
**Colletotrichum dracaenicola** Sacc. and Trott. Anthracnose on leaves of *D. fragrans* in Italy.  
**Gloeosporium polymorphum** Trinch. Large dull-brown to ashen leaf spots on *D. fragrans* in Italy.  
**Gloeosporium thuenenii** Sacc. See Anthurium.  
**Graphiola disticha** (Ehrbg.) Lévl. Raised brown hard fruiting bodies on leaves of *D. draca* in India.  
**Lembosia orbicularis** Pat. Elongate black fruiting bodies disfiguring the leaves of *Dracaena* sp. in Zanzibar.  
**Lembosia patouillardii** Sacc. and Syd. On leaves of *Dracaena* sp. in east Africa.  
**Macrophoma bakeri** Syd. Discolored areas on leaves of *Dracaena* sp. in Brazil.  
**Melanconium stictoides** Sacc. and Paol. On leaves of *Dracaena* sp. in Malacca.  
**Meliola dracaenicola** Pat. and Har. Superficial black fungus patches on leaves of *Dracaena* sp. in the Congo.  
**Meliola subdentata** Pat. Circular black superficial fungus patches on leaves of *Dracaena* sp. in Indo-China.  
**Mycosphaerella dracaenae** Tassi. On leaves of *D. repera* in Italy.  
**Myxosporium dracaenicolum** B. and Br. Destroys leaves of *Dracaena* sp. in Great Britain.  
**Phyllachora nervisequia** Wint. Black stromata on leaves of *D. cannifolia* in Queensland.  
**Phyllosticta cordylines** Sacc. and Berl. On leaves of *D. australis* and *D. terminalis* in Queensland.  
**Phyllosticta dracaenae** Griff. and Maubl. Large irregular leaf spots on *Dracaena* sp. in France, working injury to greenhouse plants.  
**Phyllosticta draconis** Berk. Irregular pale-brown leaf spots with purple margins on *D. cooperi*, *D. draca*, and *D. terminalis* in Portugal, France, Great Britain, and Finland.  
**Physalospora amphididyma** Syd. Subcircular white leaf spots with red-purple raised margins on *D. papahii* in Tanganyika.  
**Placoasterella schweinfurtheii** (P. Henn.) Theiss. and Syd. On leaves of *D. ombetis* in Abyssinia.  
**Rosellinia echinata** Mass. See Ficus.
- Ustilago dracaenae** S. da Cam. Leaf smut on *D. draco* in Portugal.  
**Ustilago trabutiana** Sacc. Black powdery masses of smut spores in ovaries of *D. draco* in Algeria.
- DRACOCEPHALUM. MOLDAVICA** Ag. DRAGONHEAD. Hardy herbaceous annuals or perennials.  
**Peronospora rossica** Gäum. Downy mildew on leaves of *D. thymiflora* in Russia.  
**Septoria dracocephali** Thuem. On leaves of *D. peregrinum* in Siberia.
- DRIMYS.** Broad-leaved evergreen trees and shrubs.  
**Actinothyrium drimydis** Speg. Circular pale-brown leaf spots on *D. winteri* in Chile.  
**Gloeosporium walteri** McAlp. Anthracnose on leaves of *D. aromatica* in Australia.  
**Helminthosporium orbiculare** Lévl. On leaves of *D. chilensis* and *D. winteri* in Chile.  
**Munkia drimydis** (Lévl.) Speg. On leaves of *D. winteri* in Chile.  
**Mycosphaerella drimydis** (Berk.) Sacc. Circular brown leaf spots on *D. winteri* in Chile and Brazil.  
**Phyllosticta drimydis** Speg. Circular gray-white leaf spots with purple margins on *D. winteri* in Chile.  
**Phyllosticta winterii** Speg. Circular brown leaf spots on *D. winteri* in Chile.  
**Septoria drimydicola** Speg. Circular to irregular leaf spots on *Drimys* sp. in Brazil.  
**Septoria drymios** Mont. On leaves of *D. chilensis* in Chile.  
**Septoria winterii** Speg. On leaves of *D. winteri* in Chile.

**DRYAS.** Dwarf, hardy, tufted evergreen shrubs.

**Didymella dryadis** Speg. Indefinite leaf spots on *D. octopetala* in Italy.

**Guignardia rhytismoides** (Berk.) Trav. On *D. octopetala* in Spain.

**Mycosphaerella ootheca** (Sacc.) On leaves of *D. octopetala* in Alaska.

**Septoria semilunaris** Johans. On peduncles of *D. octopetala* in Scandinavia.

**Venturia tirolensis** v. Hoeh. On leaves of *D. octopetala* in Austria.

**DRYOPTERIS.** WOOD FERNS. Wood ferns with much dissected leaves.

**Hyalospora filicum** Diet. See Asplenium.

**Milesia columbiana** Diet. Rust on fronds of *D. patens* and *Nephrolepis pendula* in Porto Rico and Colombia.

**Uredo gymnogrammes** P. Henn. Rust on fronds of *Adiantum latifolium*, *D. mollis*, *D. poiteana*, *Pityrogramma calomelanos*, and *Tectaria martenicensis* in Porto Rico, Cuba, Jamaica, Trinidad, and Brazil.

**DURANTA.** SKY FLOWER. Tropical shrubs cultivated for their flowers.

**Corticium salmonicolor** B. and Br. See Citrus.

**Phyllachora durantae** Rehm. Black stromata on small yellowish to reddish leaf spots on *Duranta* sp. in Ecuador.

**Phyllachora fusicarpa** Seaver. Black, irregular, slightly raised stromata on leaves of *D. repens* in Porto Rico and the Bahamas. This fungus causes a serious leaf fall.

**DURIO.** Durian. Tropical fruit trees.

**Colletotrichum durionis** Koord. Anthracnose on leaves of *D. zibethinus* in Java.

**Corticium salmonicolor** B. and Br. See Citrus.

**Gloeosporium zibethinum** Sacc. Anthracnose on leaves of *D. zibethinus* in Malaya.

**Homostedia durionis** Racib. Circular black stromata on leaves of *D. zibethinus* in Java.

**Phoma durionis** Petch. Attacks fruit of *D. zibethinus* in Ceylon.

**Phyllachora macrospora** A. Zimm. Circular black stromata on leaves of *D. zibethinus* in Java.

**Phyllosticta durionis** A. Zimm. Pale-brown leaf spots on *D. zibethinus* in Java.

**Phyllosticta nephelii** Delacr. Dull-brown indefinite leaf spots on *D. zibethinus* and *Nephelium lappaceum* in the Congo and France.

**Placosphaeria durionis** Syd. On leaves of *D. zibethinus* in the Philippines.

**ECHEVERIA.** Fleshy-leaved plants.

**Endophyllum sempervivi** (A. and S.) De B. See Sempervivum.

**ECHINOCACTUS.** A genus of the cactus family.

**Stagonospora assans** Pass. See Cereus.

**ECHINOCHLOA.** BARNYARD GRASS. A few species sometimes cultivated for grain or forage.

**Claviceps balansoides** A. Moell. Irregular black-brown sclerotia in spikes of *Echinochloa* sp. in Brazil.

**ECHINOPS.** GLOBE THISTLE. Coarse thistle-like plants with flowers in globose masses.

**Puccinia echinopsis** DC. Powdery brown to black rust pustules on leaves of *E. banaticus*, *E. cyaneus*, *E. ritro*, and *E. sphacrocephalus* in Europe.

**Puccinia hellenica** Trotter. Leaf rust on *E. microcephalus* in Greece.

**Puccinia pulvinata** Rabh. Brown powdery rust pustules on leaves of *E. amplexicaulis*, *E. chamaecephalus*, *E. cornigerum*, *E. echinatus*, *E. heldreichii*, *E. macrochaetus*, *E. spinosus*, and *E. viscosus* in India, Abyssinia, Uganda, Asia Minor, and French North Africa.

**ECHITES.** Tropical American twining shrubs.

**Puccinia balansae** Speg. Leaf rust on *E. fusiformis* in Argentina.

**Septoria echitis** Syd. Small, gray-white leaf spots with purple margins on *E. tweediana* in Argentina.

**ECHIUM.** Coarse, rough herbs and shrubs with flowers in spikes.

**Accidium asperifolii** Pers. Leaf rust on *E. italicum*, *E. lusitanicum*, and *E. rubrum* in Dalmatia, Russia, and Portugal.

**Accidium echii** Thuem. Leaf rust on *E. lusitanicum* in Portugal.

**Cercospora echii** Wint. Angular to irregular brown leaf spots on *E. tuberculatum* in Portugal.

**Cercospora euchlora** Maire. On leaves of *E. australe* in North Africa.

**Ramularia anchusae** Mass. See Anchusa.

**Synchytrium echii** Speg. Small galls on pale-brown leaf spots on *E. plantagineum* and *E. violaceum* in Argentina.

**EDGEWORTHIA.** PAPER TREE. Ornamental shrubs grown for their yellow flowers.

**Helicobasidium mompa** Tan. See Morus.

**EHRETIA.** Tender trees and shrubs.

**Schroeteriaster ehretiae** (Hir.) Syd. and Butl. Leaf rust on *E. acuminata* in Japan and Formosa.

**Uredo ehretia** Barel. Leaf rust on *E. acuminata*, *E. macrophylla*, and *E. serrata* in India and Japan.

**ELAEAGNUS.** RUSSIAN OLIVE. Shrubs and small trees grown chiefly for their handsome foliage and ornamental fruits.

**Accidium elaeagni** Diet. Leaf rust on *E. glabra* and *E. pungens* in Japan.

**Accidium elaeagni-latifolia** Petch. Ochraceous rust pustules on small, definite leaf spots on *E. latifolia* in Ceylon.

**Accidium elaeagni-umbellatae** Diet. Rust on circular to irregular leaf spots on *E. macrophylla* and *E. umbellata* in Japan.

**Accidium minoense** Syd. Leaf rust on *E. japonicus* in Japan.

**Accidium quintum** Syd. Leaf rust on *E. umbellata* in Japan.

**Ascochyta elaeagni** Sacc. Ochraceous leaf spots on *E. argentea*, *E. multiflora* (*E. edulis*), and *E. gussonii* in Russia, Italy, and France.

**Ochropsora nambuana** (P. Henn.) Diet. Leaf rust on *E. multiflora* (*E. longipes*), *E. macrophylla*, and *E. umbellata* in Japan.

**Puccinia achora** Syd. Black rust sori on leaves of *E. macrophylla* in Japan.

**Puccinia elaeagni** Yosh. Leaf rust on *E. pungens* in Japan.

**Septocylindrium olivascens** Thuem. Subcircular, olivaceous leaf spots on *E. angustifolia* and *E. rhamnoides* in France and Denmark.

**Septoria argyrea** Sacc. Ochraceous leaf spots on *E. angustifolia* and *E. argentea* in Italy. Reported from North Dakota.

**ELAEIS.** African oil palm. See Palmae.

**ELAEODENDRON.** FALSE OLIVE. Tropical shrubs and small trees.

**Septoria elaeodendri** P. Henn. Circular dull-brown leaf spots on *E. xylocarpum* in Germany.

**ELETTARIA.** Cardamom. Perennial hothouse herbs.

**Placostroma elettariae** (B. and Br.) Theiss. and Syd. Shiny black stromata on leaves of *E. floribunda* in Ceylon.

**Schroeteriaster elettariae** Rac. Rust sori on narrow, long, gray or brown leaf spots on *Elettaria* sp. in Java.

- ELEUSINE.** AFRICAN MILLET. Annual grasses, some grown as ornamentals.
- Acrothecium lunatum** Wakk. See Panicum.
- Phyllachora eleusines** Speg. Black, smooth, subcircular stromata on leaves of *E. coracana*, *E. indica*, and *E. tristachya* in Argentina, Brazil, Uganda, and the Congo.
- Ustilago eleusines** Kulkarni. Circular to elongate green, then chocolate-brown to black, smut sori in single grains or groups of grains of *E. coracana* in India, the sori rupture exposing the deep brown to black spore masses.
- ELSHOLTZIA.** Herbs or shrubs cultivated for their spikes of blue or lilac flowers.
- Coleosporium perillae** Syd. See Perilla.
- ELYMUS.** WILD RYE. DUNE GRASS. Erect perennial grasses, sometimes cultivated as ornamentals or for forage.
- Cercospora elymi** Rostr. Oblong dull-brown leaf spots on *E. arenarius* in Denmark.
- Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.
- Puccinia elymi-sibiricae** S. Ito. Powdery, brown rust pustules on leaves of *E. sibiricus* in Japan.
- Rostrupia elymi** (West.) Lagh. Brown leaf rust on *E. arenarius*, *E. mollis*, and *E. sibiricus* in Japan and northern Europe.
- Septoria elymi** E. Rostr. On leaves of *E. arenarius* in Denmark.
- Septoria elymicola** Died. On leaves of *E. arenarius* in Europe.
- Septoria elmyi-europaei** Jaap. Yellow, then brown, sunken leaf spots on *E. europaeus* in Switzerland.
- Tilletia bornmülleri** Magn. Brown smut sori in ovaries of *E. crinitus* in Asia Minor.
- Tilletia controversa** Kuehn. See Triticum.
- Tilletia serbica** Ranoj. Smut sori in spikelets of *E. crinitus* in Yugoslavia.
- Tylenchus hordei** Schoeyen. See Avena.
- Uredo elmyi-capitis-medusae** Gz. Frag. Leaf rust on *E. caput-medusae* in Spain.
- Ustilago phrygia** Magn. Brown spore masses in spikes of *E. crinitus* in Asia Minor.
- EMBOTHRIUM.** South American trees and shrubs.
- Phyllosticta embotryi** Speg. Indefinite dull-brown leaf spots on *E. coccineum* in Chile.
- EMILIA.** Annual or perennial herbs with orange or scarlet flower heads.
- Accidium emiliae** Petch. Leaf rust on *E. sonchifolia* in Ceylon.
- Accidium formosanum** Syd. Leaf rust on *E. sonchifolia* in Formosa.
- Puccinia synedrellae** P. Henn. Brown rust sori on lower-leaf surfaces of *E. sagittata*, *E. sonchifolia*, and *Synedrella nodiflora* in the West Indies and Trinidad.
- Uredo emiliae-zeylanicae** Petch. Leaf rust on *E. zeylanica* in Ceylon.
- EMPETRUM.** CROWBERRY. Low shrubs, sometimes grown for their evergreen foliage and attractive fruit.
- Melasmia empetri** Magn. Destroys young shoots of *E. nigrum* in Russia.
- Metasphaeria empetri** (Fr.) Sacc. On leaves of *E. nigrum* in Europe and Alaska.
- Phyalospora empetri** Rostr. On leaves of *E. nigrum* in Denmark.
- Rhynisma empetri** White. Black smooth stromata on stems of *E. nigrum* in Great Britain, Italy, Denmark, and Sweden.
- ENCELIA.** Herbs or subshrubs.
- Accidium enceliae** Diet. and Holw. Leaf rust on *E. canescens* in Peru.
- Puccinia enceliae** Diet. and Holw. Powdery dark-brown rust sori on leaves of *Simsia amplexicaulis* (*E. mexicana*) in Mexico.
- Uredo enceliae-tomentosae** Maire. Leaf rust on *E. tomentosa* in Chile.
- ENDYMION.** See Scilla.
- ENKIANTHUS.** Shrubs grown for their handsome flowers.
- Accidium enkianthi** Diet. Leaf rust on *E. japonicus* in Japan.
- ENTADA.** Tropical shrubs.
- Ravenelia entadae** Lagh. and Diet. Brown rust pustules on leaves of *E. polystachya* in Panama and Guatemala.
- Ravenelia schweinfurthii** Syd. Brown leaf rust on *E. sudanica* in central Africa.
- ENTEROLOBIUM.** Tropical trees.
- Phyllachora enterolobii** Speg. Circular black stromata on leaves of *E. timbouva* in Brazil.
- Ravenelia hassleri** Speg. Leaf rust on *E. timbouva* in Paraguay.
- Ravenelia oligotheles** Speg. Dull-brown rust pustules on leaves of *E. timbouva* in Argentina.
- EPHEDRA.** JOINT FIR. Shrubs.
- Accidium ephedrae** Speg. Rust on leaves and branches of *E. tweediana* in Argentina.
- Epiclinium negerianum** Sacc and Syd. Black, crust-like areas on twigs of *E. andina* in Chile.
- EPIDENDRUM.** See Orchidaceae.
- EPILOBIUM.** WILLOW WEED. Herbs with willow-like foliage and large showy flower spikes.
- Ascochyta epilobii** Oud. Elongate dull-brown leaf spots on *E. angustifolium* in Holland.
- Cercospora epilobii** Schn. On leaves of *E. hirsutum* and *E. montanum* in Europe.
- Cylindrosporium epilobianum** Sacc. and Faut. Confluent leaf spots on *E. hirsutum* in France.
- Fusicladium heterosporum** V. Hoehn. Brown, often confluent, leaf spots on *E. parviflorum* in Austria.
- Gloeosporium epilobii** Pass. Brown leaf spots on *E. angustifolium* in France.
- Marsonia chamaenerii** (Rostr.) P. Magn. and var. **germanica** Syd. Irregular ochraceous brown leaf spots on *E. angustifolium* and *E. hirsutum* in Greenland and Germany.
- Mycosphaerella adusta** Fckl. On *E. boreale*, *E. bongardi*, and *Epilobium* sp. in Alaska.
- Mycosphaerella microspila** B. and Br. On leaves of *E. dodonaei*, *E. montanum*, and *E. palustre* in Denmark, Great Britain, Belgium, and Italy.
- Ovularia epilobiana** Sacc. and Fautr. Pale-brown leaf spots on *E. hirsutum* in France.
- Ovularia epilobii** Lindr. Large, irregular, confluent gray or gray-brown leaf spots on *E. palustre* in Finland.
- Phyllosticta chamaenerii** Allesch. Small angular to irregular, ochraceous then white, leaf spots with red margins on *E. angustifolium* in Germany.
- Phyllosticta epilobii** Brun. Small circular gray-white leaf spots with brown margins on *E. hirsutum* in France.
- Phyllosticta epilobii-rosei** Krieg. Circular to irregular, often confluent, brown leaf spots on *E. roseum* in Germany.
- Plasmopara epilobii** (Rhb.) Schroet. Downy mildew on leaves of *E. hirsutum*, *E. latifolium*, and *E. parviflorum* in Alaska, Denmark, and Germany.
- Puccinia epilobii** DC. Brown rust sori on lower leaf surfaces of *Epilobium* spp. in Australia and Europe.
- Puccinia epilobii-fleischeri** Ed. Fisch. Yellow to brown rust pustules on leaves of *E. fleischeri* in Switzerland.
- Puccinia gigantea** Karst. Rust on brown sunken spots on leaves and stems of *E. angustifolium* in Switzerland and Scandinavia.

**EPILOBIUM**—Continued.

- Puccinia krookii** P. Henn. Powdery, dark-brown rust sori on leaves of *E. flavescens* in the Union of South Africa.
- Ramularia enecans** Magn. Gray-white to yellow leaf spots on *E. angustifolium* in Germany.
- Ramularia epilobii** All. On leaves of *E. palustre* in Germany.
- Ramularia epilobii-parviflori** Lindr. Subcircular to oblong gray-brown leaf spots with purple margins on *E. parviflorum* in Denmark and Finland.
- Ramularia epilobii-rosei** Lind. Subcircular or oblong grayish-green to pale-brown leaf spots on *E. roseum* in Denmark and Austria.
- Ramularia hornemanni** Lindr. Subcircular brown, yellow, or red leaf spots on *E. hornemanni* in Lapland.
- Ramularia punctiformis** (Schlecht.) V. Hoeh. Leaf spot on *Epilobium* spp. in Europe. Reported from Wisconsin.
- Septoria alpicola** Sacc. On leaves of *E. alpinum* in Italy.
- Venturia maculaeformis** (Desm.) Sacc. On leaves of *E. hirsutum* in Austria.
- EPIMEDIUM.** Herbs used in rock gardens.
- Cercospora epimedii** Jach. On leaves of *E. pinnatum* in Russia.
- Mycosphaerella epimedii** (Sacc.) Jaap. On leaves of *E. alpinum* in Dalmatia, Italy, and Switzerland.
- Phyllosticta epimedii** Sacc. Irregular ochraceous leaf spots with red margins on *E. alpinum* in Italy.
- Puccinia epimedii** (Henn. and Shir.) Miy. and Ito. Rust sori on circular yellow-brown leaf spots on *E. macranthum* in Japan.
- EPIPACTIS.** Hardy terrestrial orchids. See Orchidaceae.
- ERAGROSTIS.** LOVE GRASS. Teff. Annual or perennial grasses.
- Epichloe eragrostis** Pole-Evans. Attacks *E. plana* in the Union of South Africa.
- Helminthosporium eragrostidis** P. Henn. Black, velvety fungus patches in the glumes of *Eragrostis* sp. in the Congo.
- Sorosporium turneri** McAlp. Dense black masses of smut spores in the ovaries of *E. nigra* in Australia.
- Sphaeciotheca kusanoana** P. Henn. (*Ustilago kusanoana*, P. Henn.) Smut in inflorescences of *E. ferruginea* in Japan.
- Uromyces pedicellata** Pole-Evans. Yellow-brown to black rust sori on leaves and culms of *E. abyssinica* and *E. curvula* in the Union of South Africa.
- Ustilago egenula** Syd. and Butl. Olivaceous smut sori replacing the ovaries of *E. nutans* in India.
- ERANTHEMUM.** Tropical shrubs, some species cultivated for the foliage or flowers.
- Isariopsis acanthacearum** Cke. See Daedalacanthus.
- ERANTHIS.** WINTER ACONITE. Low perennial herbs.
- Peronospora eranthidis** (Pass.) Fekl. Downy mildew on leaves of *E. hyemalis* in Switzerland.
- Septoria cajadensis** Speg. Ashen white or ochraceous circular to angular, often confluent, leaf spots on *E. hyemalis* in Italy.
- Tubercinia eranthidis** (Pass.) Liro. Smut on *E. hyemalis* in southern Europe.
- EREMOCITRUS.** Small trees allied to *Citrus*.
- Bacterium citri** Hasse. See *Citrus*.
- EREMURUS.** DESERT CANDLE. Hardy liliaceous desert plants with tall flowering stalks.
- Puccinia eremuri** Kom. Dark-brown rust pustules on both leaf surfaces of *E. inderiensis*, *E. kaufmanni*, *E. robustus*, and *E. spectabilis* in Asia Minor and Persia.
- Rhabdospora eremuri** Ohl. Dark-brown spots on stems of *Eremurus* sp. in Russia.
- ERIA.** See Orchidaceae.
- ERIANTHUS.** RAVENNA GRASS. Tall, reedlike, perennial grasses with large woolly plumelike inflorescences.
- Cintractia pulverulenta** Cke. and Mass. Black smut sori in ovaries of *Erianthus* sp. in India.
- Puccinia damloii** Bub. Brown to black powdery rust pustules on leaves of *E. hostium* in Yugoslavia.
- Uredo ravennae** Maire. Leaf rust on *E. ravenna* in French north Africa and Spain.
- Ustilago erianthi** Syd. Smut sori destroying the ovaries and forming dark-olive spore masses on *E. ravenna* in Russia.
- Ustilago sacchari** Rabh. See *Saccharum*.
- ERICA.** HEATH. Ornamental shrubs.
- Gibbera salisburgensis** Niessl. On leaves of *E. carnea* in Germany.
- Phyllosticta ericae** Allesch. Red-brown patches on leaves of *E. carnea* in Germany.
- Stemphylium ericoctonum** B. and De B. On *Erica* spp. in Europe.
- Venturia straussii** Sacc. On branches and leaves of *E. scoparia* in Italy and France.
- ERIGERON.** FLEABANE. Mostly weeds, a few species cultivated in border plantings like *Aster*.
- Aecidium spegazzinii** Pat. Leaf rust on *E. albus*, *E. bonariensis* and *E. maximus* in South America.
- Coleosporium erigerontis** Syd. Powdery golden rust pustules on leaves of *E. linifolius* in Formosa.
- Entyloma calendulae** (Oud.) De B. See *Calendula*.
- Puccinia doloris** Speg. Brown rust pustules on small brown sunken leaf spots of *E. bonariensis*, *E. deamii*, and *Erigeron* sp. in Argentina, Chile, Colombia, Costa, Rica and Guatemala.
- Puccinia dovrensis** Blytt. Dark-brown rust pustules on leaves of *E. alpinus* and *E. uniflorus* in Norway, Switzerland, and India.
- Septoria chanousii** Ferr. Irregular brown leaf spots on *E. uniflorus* in Italy.
- Septoria erigeronata** Thuem. On leaves of *E. elongatus* in Siberia.
- Venturia bonariensis** Speg. Brown indefinite leaf spots on *E. bonariensis* in Argentina.
- Venturia tucumanensis** Speg. On leaves of *E. canadense* in Argentina.
- ERIOBOTRYA.** LOQUAT. Small trees grown for ornament and edible fruit.
- Ascochyta eriobotryae** Vogl. Circular to oblong dull-brown to white leaf spots on *E. japonica* in Italy.
- Coleopuccinia simplex** Diet. Leaf rust on *E. japonica* in Japan.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Coryneum eriobotryae** Scalia. Circular, gray-white leaf spots with dark-purple margins on *E. japonica* in Sicily.
- Entomosporium** sp. Raised shiny spots on fruit and leaves surrounded by yellowish rings and never confluent on *E. japonica* in the Union of South Africa.
- Entomosporium mespili** (DC.) Sacc. See *Cotoneaster*.
- Fomes lamaoensis** Murr. See *Hevea*.
- Fusicladium** sp. Circular brown, then velvety dark olive-green, often confluent, spots on fruit and leaves of *E. japonica* in the Union of South Africa. Probably the same as the following species. The leaves are often distorted.
- Fusicladium eriobotryae** Cav. Scabby areas on leaves, stems and fruit of *E. japonica* in Australia, Russia, and Italy. Reported from California.

**ERIOBOTRYA**—Continued.**Fusarium udum** Butl. See *Dianthus*.**Gloeosporium eriobotryae** Speg. Circular, then confluent, ashen leaf spots on *E. japonica* in Argentina.**Leptosphaeria eriobotryae** Syd. and Butl. On leaves of *E. japonica* in India.**Phyllosticta eriobotryae** Thuem. Subcircular to irregular and confluent gray-brown leaf spots on *E. japonica* in Argentina, Brazil, Japan, Formosa, Australia, Russia, and Italy.**Phyllosticta uleana** Syd. Large circular, rarely confluent, dull-yellow to brown leaf spots with purple margins on *E. japonica* in Brazil.**Septoria eriobotryae** Maffei. Irregular, black leaf spots on *E. japonica* in Italy.**ERIOSTEMON**. Australian evergreen shrubs.**Puccinia eriostemonis** McAlp. Dark-orange rust pustules on leaves of *E. myoporoides* in Australia.**ERLANGEA**. Vernonia-like shrubs.**Mycosphaerella erlangeae** Bacc. On leaves of *E. abyssinica* in Abyssinia.**Puccinia erlangeae** Grove. Leaf rust on *E. tomentosa* in British east Africa.**ERODIUM**. HERON'S-BILL. Alfilaria. Filaree. Annual or perennial herbs grown in flower gardens.**Gloeosporium schweinfurthianum** Thuem. On leaves of *E. glaucophyllum* in Egypt.**Peronospora erodii** Fekl. Downy mildew on leaves of *E. ciconium*, *E. cicutarium*, and *E. malachoides* in Sardinia, Denmark, Russia, Austria, and Germany.**Phyllosticta erodii** Speg. Subcircular brown leaf spots with broad purple margins on *E. malachoides* in Argentina.**Ramularia erodii** Bres. Small irregular leaf spots on *E. cicutarium* in Spain, Denmark, and Germany.**Ramularia geranii** (West.) Fekl. See Geranium.**Synchytrium papillatum** Farl. Small galls on dull-purple leaf spots on *E. cicutarium* in Guadeloupe, Canary Islands, St. Croix, and the Union of South Africa. Also in California on imported plants.**ERYNGIUM**. ERYNGIO. Sea holly. Annual and perennial herbs valued for the steel-blue cast of their stems.**Aphelenchus olesistus** Ritz. Bos. See Begonia.**Ascochyta phomoides** Sacc. On stems of *Eryngium* sp. in France.**Entyloma eryngii** (Cda.) De By. Smut pustules on circular to irregular, often confluent, brown leaf spots on *E. campestre* and *E. planum* in France, Italy, Austria, and Germany. Reported from Iowa.**Erysiphe taurica** Lévl. See Althaea.**Leptosphaeria woodrowi wilsoni** Garb. On leaves of *E. campestre* in Russia.**Mycosphaerella eryngii** (Fr.) Lind. and var. **libanotis** Fekl. On leaves of *E. campestre*, *E. maritimum*, and *Libanotis montana* in Europe.**Phyllosticta eryngiana** Sacc. and Fautr. Dull yellow-brown leaf spots on *E. campestre* in France.**Phyllosticta eryngiella** Bub. Gray-white angular leaf spots on *E. campestre* in Hungary.**Phyllosticta eryngii** Syd. Large indefinite black, often confluent, leaf spots on *E. maritimum* and *E. pandanifolium* in Denmark and Germany.**Phyllosticta eryngicola** Bub. On leaves of *E. campestre* in Hungary.**Puccinia eryngii** DC. Leaf rust on *E. bovei*, *E. campestre*, *E. congestum*, *E. creticum*, *E. glomeratum*, and *E. virens* in Europe, Egypt, French North Africa, and Asia Minor.**Ramularia eryngii** Jachw. On leaves of *E. planum* in Russia.**Septoria eryngicola** Oud. and Sacc. On leaves of *E. campestre* in Italy and Spain.**Septoria eryngii** West. Angular brown to gray-white leaf spots on *E. maritimum* in Denmark and Belgium.**ERYSIMUM**. BLISTER CRESS. Early-blooming hardy annuals with yellow and orange flowers.**Septoria erysimi** Niessl. Pale-brown leaf spots on *E. cheiranthoides* in Bohemia.**Septoria repanda** Bub. Circular to irregular, yellow or white leaf spots on *E. banaticum* and *E. repandum* in Yugoslavia and Bohemia.**ERYTHRAEA**. CENTAURIUM. Ag. Low, annual, biennial or perennial herbs.**Cercospora erythraeae** Höll. Dull-yellow areas on leaves and stems of *E. linariaefolia* in Hungary.**Peronospora erythraeae** (Kuehn) Gäum. Downy mildew on leaves, peduncles and stems of *E. centaurium*, *E. linariaefolia*, *E. litoralis*, *E. pulchella*, and *E. ramosissima* in central and north Europe.**Phyllosticta erythraeae** Sacc. and Speg. On leaves of *E. centaurium* in Italy.**Pyrenopeziza plantaginis** Fekl. var. **erythraeae** Pat. Brown spots on lower leaf surfaces of *E. centaurium* in Tunis and Algeria.**Septoria gentianae** Thuem. var. **erythuseae** Gz. Frag. On leaves of *E. chloodis* in Spain.**ERYTHRINA**. CORAL TREE. Herbs, shrubs, and trees cultivated for ornament and as coffee shade.**Coleosporium erythrinae** Petch. Leaf rust on *E. lithosperma* in Ceylon.**Colletotrichum erythrinae** Koord. Anthracnose on leaves of *E. lithosperma* in Ceylon and Java.**Diclierinia binata** (Berk.) Arth. (*Uredo cabreriana* Kern. and Kellerm.) Powdery deep-brown rust pustules on lower leaf surfaces of *E. glauca* and *E. umbrosa* in Porto Rico, Cuba, Guatemala, and Trinidad.**Diplodia cacaoicola** P. Henn. See Theobroma.**Fomes lamaoensis** Murr. See Hevea.**Fomes lignosus** Klotzsch. See Hevea.**Helminthosporium extensum** Petch. On leaves of *E. lithosperma* in Ceylon.**Helminthosporium inversum** Sacc. On leaves of *E. indica* in the Philippines.**Mycosphaerella erythrinae** Koord. Circular gray-white leaf spots on *E. lithosperma* and *E. ovalifolia* in Java and Ceylon.**Phyllosticta australis** Speg. Indefinite grayish-white leaf spots on *E. crista-galli* in Argentina.**Phyllosticta erythrinae** Petch. Spots on branches and leaves of *E. lithosperma* in Ceylon.**Phyllosticta erythrinicola** Young. Numerous small, circular, sordid-white leaf spots on *E. micropteryx* in Porto Rico.**Ravenelia erythrinae** Gäum. Leaf rust on *E. microcarpa* and *E. velutina* in Java.**Ravenelia platensis** Speg. Powdery brown rust sori on pyriform galls on leaf blades and petioles and on branches of *E. crista-galli* in Argentina and Uruguay.**Rosellinia arcuata** Petch. See Thea.**Rosellinia pepo** Pat. See Citrus.**Septoria bonanseaana** Sacc. Numerous subcircular gray-white leaf spots with ochraceous margins on *E. breviflora* in Mexico.**Sphaerostilbe repens** B. and Br. See Hevea.**Telimena erythrinae** Racib. Subcircular yellow-green, then yellow, finally brown, leaf spots on *E. lithosperma* in Java.**Trabutia erythrina** Rick. Circular black, often confluent, stromata on *E. crista-galli* in Brazil.

**ERYTHRINA**—Continued.

**Uredo erythrinae** P. Henn. Rust pustules on circular, then confluent, yellow-brown leaf spots on *E. indica*, *E. tomentosa* and *Erythrina* sp. in the Philippines, the Congo, and Uganda.

**Uredo erythrinae-ovalifoliae** Petch. Leaf rust on *E. ovalifolia* and *E. velutina* in Ceylon.

**Uromyces erythrinae** Lagh. Brown rust pustules on leaves of *Erythrina* sp. in Ecuador.

**ERYTHRONIUM**. TROUT LILY ADDER'S-TONGUE. Small, spring-flowering, hardy bulbous plants.

**Ascochyta erythronii** Sacc. and Speg. Indeterminate white leaf spots on *E. dens-canis* in Italy.

**Septoria erythronii** Sacc. and Speg. Subcircular to irregular white spots on leaves of *E. dens-canis* in Italy.

**Septoria falsipora** Bub. Circular to elongate dirty-white spots without definite margins on leaves of *E. dens-canis* in Montenegro.

**Uromyces erythronii** (DC.) Pass. Yellow rust pustules, which later become brown, on both sides of the leaves and on the petioles of *E. dens-canis* in France, Spain, Italy, Switzerland, Bulgaria, Yugoslavia, Dalmatia, Austria, Hungary, Japan, Siberia, and Germany.

**ERYTHROXYLON**. COCAINE TREE. Woody medicinal plants.

**Corticium** sp. Black rot of *E. coca* in Ceylon. Similar to *Corticium theae* on *Thea* (q. v.).

**Corticium salmonicolor** B. and Br. See Citrus.

**Cronartium gilgianum** P. Henn. Red-brown rust pustules on leaves of *Erythroxyton* sp. and *Euclea* sp. in central and east Africa.

**Fomes lamaoensis** Murr. See Hevea.

**Phyllachora asteriana** Speg. Shiny black stromata on leaves of *E. suberosum* in Brazil.

**Phyllosticta erythroxyli** Graz. Small circular to irregular gray-white leaf spots with yellow to dull-brown margins on *E. coca* in Bolivia and Peru.

**Uredo erythroxytonis** Graz. Leaf rust on *E. areolatum* and *E. coca* in Porto Rico, Cuba, Peru, Bolivia, and Brazil.

**ESCALLONIA**. South American evergreen shrubs or small trees.

**Phyllachora escalloniae** Pat. Subcircular black stromata on leaves of *Escallonia* sp. in Ecuador.

**Physalospora escalloniae** P. Henn. Black spots on leaves of *E. chlorophylla* in Brazil.

**Trabutia escalloniae** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *E. rubra* in Argentina.

**EUCALYPTUS**. Valuable ornamental and timber trees, natives of Australia, many species of which are called gum trees, e. g., blue gum, spotted gum.

**Aulographium eucalypti** Cke. and Mass. On leaves of *E. obliqua* in Australia and Tasmania.

**Camarosporium eucalypti** Wint. Angular to irregular pale-brown to ashen leaf spots on *Eucalyptus* sp. in Australia.

**Cercospora epicoccoides** Cke. and Mass. Small purple, then confluent, leaf spots on *Eucalyptus* sp. in Australia.

**Cercospora eucalypti** Cke. and Mass. Subcircular, then confluent, pale-brown leaf spots with rosy margins on *Eucalyptus* sp. in Australia.

**Cercospora theae** Petch. See Thea.

**Coniothecium eucalypti** Thuem. On branches of *E. globulus* and *Eucalyptus* sp. in Spain, Portugal, and Ecuador.

**Corticium salmonicolor** B. and Br. See Citrus.

**Coryneum eucalypti** d'Alm. and da Cam. Small irregular white spots with brown margins on *E. globulus* in Portugal.

**Cylindrosporium eucalypti** McAlp. Circular definite gray leaf spots with red-brown margins on *E. melliodora* in Victoria.

**Diaporthe cubensis** Bruner. Produces a canker of trunks and limbs of *Eucalyptus* spp. in Cuba. The bark is destroyed, often with a gummy exudation, and lesions are formed which permit the entry of other fungi.

**Endothia havanensis** Bruner. On bark of twigs and branches of *E. botryoides*, *E. microphylla*, *E. occidentalis*, *E. robusta*, *E. rostrata*, *Mangifera indica*, *Persea gratissima*, and *Spondias mombin* in Cuba.

**Gloeosporium eucalypti** McAlp. Leaf anthracnose on *E. corynocalyx* in Australia.

**Gloeosporium nigricans** Cke. and Mass. Anthracnose on leaves of *E. pauciflora* in Australia.

**Gloeosporium rhipidium** Speg. Anthracnose on leaves of *E. globulus* in Uruguay and Argentina.

**Harknessia uromycoides** Speg. On leaves of *Eucalyptus* sp. and *E. globulus* in west Australia, Portugal, and California.

**Hendersonia grandispora** McAlp. Attacks leaves and twigs of *Eucalyptus* sp. in Victoria.

**Hypospila eucalypti** Wakef. Red-brown spots on leaves of *Eucalyptus* sp. in Queensland.

**Lembosia orbicularis** Wint. Black elongate superficial perithecia on subcircular leaf spots on *E. pilularis* in Australia.

**Leptostromella eucalypti** Cke. and Mass. Subcircular red-brown leaf spots on *Eucalyptus* sp. in Australia.

**Melasmia eucalypti** Cke. and Mass. On *Eucalyptus* leaves in Queensland.

**Mycosphaerella cryptica** Cke. On leaves of *Eucalyptus* sp. in Australia.

**Mycosphaerella molleriana** Thuem. and var. **megalospora** da Cam. Irregular dull-brown leaf spots on *E. ficifolia*, *E. globulus* and *Eucalyptus* sp. in Brazil, Algeria, and Portugal.

**Mycosphaerella nubilosa** Cke. On leaves of *Eucalyptus* sp. in Australia.

**Oidium** sp. Powdery mildew causing crinkling and distortion of leaves on young shoots of *Eucalyptus* spp. in Brazil.

**Oidium eucalypti** Rostr. Powdery mildew on leaves of *Eucalyptus* seedlings in Denmark.

**Phoma australis** Cke. Elliptical, then confluent, dull-brown leaf spots on *Eucalyptus* sp. in Australia.

**Phoma eucalyptidea** Thuem. On leaves of *E. globulus* in Victoria and Portugal.

**Phyllachora eucalypti** (Cke. and Mass.) Theiss. and Syd. Black stromata on leaves of *E. viminalis* in Tasmania.

**Phyllachora maculata** Cke. Shiny black stromata on upper leaf surfaces of *Eucalyptus* sp. in Australia.

**Phyllosticta eucalypti** Thuem. Large irregular dull-brown, then gray-white, leaf spots on *E. giganteus* and *E. globulus* in Russia, Denmark, Spain, Portugal, Algeria, and Australia.

**Phyllosticta eucalyptina** Pat. On leaves of *E. globulus* in Tunis.

**Phyllosticta globuli** Pass. On leaves of *E. globulus* in Italy.

**Phyllostictiella phomatoides** (Cke. and Mass.) Tass. On leaves of *Eucalyptus* sp. in Victoria.

**Physalospora latitans** Sacc. Attacks leaves and twigs, causing a die-back of *E. colosseae* and *E. rostrata* in Portugal and Brazil. Reported from California.

**Placostroma inaequale** (Cke.) Theiss. and Syd. Black shiny stromata on brown leaf spots on *Eucalyptus* sp. in Victoria.

**Raderiella mirabilis** Syd. Stromata on circular, sunken, dull-brown leaf spots on *E. capitellata* in Victoria.

**EUCALYPTUS**—Continued.

**Rhytisma eucalypti** P. Henn. Black opaque stromata on dull-brown, circular leaf spots on *E. santalifolia* in Australia.

**Schizothyrium eucalyptorum** Cke. and Mass. Circular reddish leaf spots on *E. obliqua* in Australia.

**Septoria eucalypti** Wint. and Roum. On leaves of *E. amygdalina* in Algeria.

**Septoria mortoleusis** Penz. and Sacc. See Acacia.

**Sphaerostilbe repens** B. and Br. See Hevea.

**Stictus emarginata** Cke. and Mass. On leaves of *Eucalyptus* spp. in Ceylon.

**Ustilago vriesiana** Vuill. A smut said to cause galls on stems and roots of seedlings from which witches'-broomlike growths develop. On *E. amygdalina* in France and Holland.

**Canker.** An undetermined fungus attacks *Eucalyptus* spp., *Mangifera indica*, *Persea gratissima*, and other trees in Hawaii and possibly in Brazil. It works inward from the leaves and young shoots, as well as through the bark, forming cankers and ultimately killing infected trees.

**EUCHARIS.** AMAZON LILY. Bulbous plants.

**Accidium delicatulum** Arth. Leaf rust on *Eucharis* sp. in Trinidad.

**Uredo eucharidis** P. Henn. Brown leaf rust on *E. candia* in Peru and Ecuador.

**EUCHLAENA.** TEOSINTE. Stout annual or perennial grasses grown for forage and ornament.

**Helminthosporium euchlaenae** Zimm. Elongate brown leaf spots on *E. mexicana* in Tanganyika.

**Phyllosticta euchlaena** Sacc. Turns leaves of *E. luxurians* white, destroying their forage value. In the Philippines.

**Sclerospora maydis** (Rac.) Butl. See Zea.

**Sclerospora philippinensis** Weston. See Zea.

**Sclerospora sacchari** Miy. See Saccharum.

**Sclerospora spontanea** Weston. See Zea.

**Ustilago kellermanii** Clint. Smut sori involving the terminal and nodal growths of *E. luxurians* in Guatemala. The brown-black dusty spore masses are inclosed by the leaf sheaths.

**EUGENIA.** SURINAM CHERRY. CLOVE TREE. A large group of tropical trees and shrubs, many species cultivated for ornament or fruit.

**Actinothecium callicola** Speg. Small circular pale-brown leaf spots on *Eugenia* sp. in Brazil.

**Campotrimum eugeniae** Pat. Circular dull-brown leaf spots on *Eugenia* sp. in Indo-China.

**Catacauma peglerae** Doidge. Black stromata on leaves of *E. capensis* in the Union of South Africa.

**Coniothyrium trigonicolum** Rangel. On leaves of *E. uniflora* in Brazil.

**Entyloma eugeniaram** Cke. and Mass. Dark-brown smut sori on circular to angular leaf spots on *Eugenia* sp. in Queensland.

**Entyloma paradoxum** Syd. Leaf smut on *E. aquea* in Java.

**Fusarium udum** Butl. See Dianthus.

**Helminthosporium asterinoides** Sacc. and Syd. Black superficial fungus layers on lower leaf surfaces of *Eugenia* sp. in Brazil.

**Lasmenia balansae** Speg. Small black irregular stromata on leaves of *Eugenia* sp. in Paraguay and Brazil.

**Melasmia pulchella** Speg. On leaves of *Eugenia* spp. in Paraguay and Brazil.

**Metasphaeria incompleta** Rehm. On *Eugenia* sp. in the Philippines.

**Microcyclus labens** Sacc. and Syd. On leaves of *Eugenia* sp. in Chile.

**Mycosphaerella eugeniae** Rehm. On leaves of *E. uniflora* (*E. micheli*) in Brazil.

**Myiocopron valdivianum** Speg. On leaves of *Eugenia* sp. in Chile.

**Napidadium fumago** Speg. Thin black superficial fungus layers on leaves of *Eugenia* sp. and *Myrtus* sp. in Chile.

**Napidadium myrtacearum** Speg. As above on *Eugenia* sp. in Brazil.

**Phaeophleospora eugeniae** Rangel. On leaves of *E. uniflora* in Brazil.

**Phyllachora ambigua** Syd. Black irregular stromata on leaves of *E. jambolana* in India.

**Phyllachora angustispora** Speg. Black stromata on leaves of *Eugenia* sp. in Argentina.

**Phyllachora biareolata** Speg. Black stromata on circular brown leaf spots on *E. guabiju* in Argentina.

**Phyllachora phylloplaca** (Kze.) Theiss. and Syd. Shiny black stromata on leaves of *Eugenia* sp. in Brazil.

**Phyllachora vimulosa** Speg. Black stromata on leaves of *Eugenia* sp. in Costa Rica.

**Phyllachora wetzeli** Chardon. Small black circular stromata on brown leaf spots on *Eugenia* sp. in Porto Rico.

**Phyllosticta eugeniae** Young. Circular dark-brown leaf spots with raised light-brown margins on *E. buxifolia* in Porto Rico.

**Phyllosticta icarahunensis** Rangel. Leaf spots on *E. uniflora* in Brazil.

**Phyllosticta myrticola** Speg. Circular white leaf spots with definite narrow dull-red margins on *E. smithii* and *Eugenia* sp. in Brazil and Victoria.

**Polystomella granulosa** (Kl.) Theiss. and Syd. Dull-black stromata on leaves of *Eugenia* sp. in Chile.

**Puccinia barbaccensis** Rangel. Leaf rust on *Eugenia* sp. in Brazil.

**Puccinia eugeniae** Rangel. Leaf rust on *E. grandis* in Brazil.

**Puccinia grumixamal** Rangel. Leaf rust on *E. brasiliensis* in Brazil.

**Septoria eugeniaram** P. Henn. On leaves of *Eugenia* sp. in Argentina.

**Septoria eugenicola** Speg. Sanguineous leaf spots with white centers on *Eugenia* sp. in Argentina.

**Sphaerodothis balansae** (Tass.) V. Hoeh. Black stromata on lower leaf surfaces of *Eugenia* sp. in Paraguay.

**Uncinula australis** Speg. Powdery mildew on leaves of *Eugenia* sp. in Paraguay.

**Uredo eugeniaram** P. Henn. Leaf and stem rust on *E. uvalha* and *Eugenia* sp. in Brazil.

**Uredo goeldiana** P. Henn. Yellow to golden rust pustules on fruit of *Eugenia* sp. in Brazil.

**Uredo myrtacearum** Pazsch. Yellow-brown rust pustules on leaves of *E. grandis* in Brazil.

**EULOPHIA.** Terrestrial orchids. See Orchidaceae.**EUONYMUS.** BURNING BUSH. WINTER CREEPER. WAHOO. Deciduous or evergreen shrubs and small trees.

**Ascochyta evonymi** Oud. Circular gray-white leaf spots with brown margins on *E. europaea*, *E. japonica*, and *E. vulgaris* in Bohemia and Holland.

**Ascochyta evonymicola** Allesch. Irregular red-brown, then gray-white, spots on leaves of *E. europaea* in Italy and Germany.

**Cercospora evonymi** Erikss. Subcircular to angular brown leaf spots with purple margins on *E. europaea* in Scandinavia.

**Gloeosporium euonymicolum** Hemmi. Anthracnose causing leaf fall of *E. japonica* and *E. radicans* in Japan.

**Gloeosporium evonymi** Br. and Cav. Leaf anthracnose on *E. japonica* in Italy.

**EUONYMUS**—Continued.

- Gloeosporium frigidum** Sae. Anthraenose on leaves of *E. japonica* in Italy.
- Macrophoma cylindrospora** (Desm.) Sacc. See *Hedera*.
- Marssonia thomasiana** Sacc. Subcircular dull-red leaf spots on *E. europaea* and *E. latifolia* in Europe. Reported from Wisconsin.
- Melampsora evonymi-capraearum** Kleb. See *Salix*.
- Microsphaera euonymi** (DC.) Saec. Powdery mildew on leaves of *E. europaea* and *E. verrucosus* in Europe.
- Mycosphaerella evonymi** (Kunze.) Sehroet. On leaves of *E. europaea* in Europe.
- Oidium euonymi-japonici** (Areang.) Salm. Dense white patches of mildew on leaves of *E. japonica* and *E. radicans* in Japan, Argentina, and Europe. Also known from Louisiana.
- Phyllosticta aliena** (Fr.) Sae. Elongate spots on branches of *E. europaea* in France.
- Phyllosticta bolleana** Sae. On leaves of *E. japonica* in Italy and Austria.
- Phyllosticta destructiva** Desm. See *Althaea*.
- Phyllosticta evonymella** Sae. Angular olive leaf spots on *E. europaea*, *E. japonica*, and *E. latifolia* in Spain, Italy, France, and Germany.
- Phyllosticta evonymicola** Togn. Gray-white leaf spots with ochraceous margins on *E. europaea* in Italy.
- Phyllosticta nemoralis** Saec. Indefinite pale-brown leaf spots on *E. europaea* in Spain and France.
- Phyllosticta pustulosa** S. and R. Rufous gray-white leaf spots on *E. japonica* in Algeria.
- Phyllosticta sardoa** Pass. Large gray-white leaf spots on *E. japonica* in Italy.
- Septoria evonymella** Pass. On leaves of *E. japonica* in Italy and Austria. Reported from South Carolina.
- Septoria evonymi** Rbh. Large pale-brown leaf spots on *E. europaea* in Italy, Russia, and Germany. Reported from Virginia.
- Septoria evonymi-japonicae** Pass. On leaves of *E. japonica* in Italy.
- Septoria evonymina** Sever. On *E. japonica* in Italy.
- Septoria japonica** Oud. Large pale-brown leaf spots on *E. japonica* in Holland.
- Septoria semicircularis** Saec. and Scalia. On leaves of *E. fimbriatus* in Portugal.
- EUPATORIUM**. THOROUGHWORT. BONESET. MIST FLOWER. Perennial herbs, the tropical species shrubby.
- Accidium ampliatrium** Jaeks. and Holw. Leaf rust on *Eupatorium* sp. in Costa Rica.
- Accidium eupatorii** Diet. Rust pustules on large yellow circular leaf spots on *Eupatorium* sp. in Brazil.
- Accidium heteromorphum** Speg. Leaf rust on *Eupatorium* sp. in Argentina.
- Accidium paramense** Mayor. Rust on leaf blades and petioles and stems of *E. obscurifolium* in Colombia.
- Accidium roseum** Diet. and Holw. Golden rust pustules on red or purple leaf spots on *Eupatorium* sp. in Mexico.
- Baeodromus eupatorii** Arth. Brown rust pustules on leaves and stems of *E. aschenbornianum* and *E. pazcuarensis* in Mexico and Guatemala.
- Cionothrix praelonga** (Wint.) Arth. Yellow rust pustules on leaves of *Eupatorium* spp. in Mexico, Guatemala, Costa Rica, Trinidad, Colombia, Ecuador, and Brazil.
- Coleosporium eupatorii** Arth. Golden rust pustules on leaves of *E. chinense*, *E. collinum*, *E. macrophyllum*, and *E. oerstedianum* in Porto Rieo, Cuba, Guatemala, Costa Rica, Nicaragua, Colombia, and Formosa.
- Cronartium andinum** Lagh. Brown rust pustules on leaves of *E. glutinosum* in Ecuador.
- Cronartium eupatorinum** Speg. Leaf rust on *Eupatorium* sp. in Argentina.
- Mycosphaerella eupatoriicola** Petr. On leaves of *E. cannabinum* in Bohemia and Silesia.
- Phyllosticta eupatoriicola** Kab. and Bub. Irregular, often confluent, dull-brown leaf spots on *E. cannabinum* and *E. odoratum* in Porto Rieo and Bohemia.
- Phyllosticta eupatorii** Allesch. Small subcircular dull-brown to ashen, often confluent, leaf spots with dull-purple margins on *E. cannabinum* in Germany.
- Phyllosticta eupatorina** Thuem. Subcircular ashen leaf spots with purple borders on *E. cannabinum* in Italy and Portugal. Reported from Illinois and New Jersey.
- Puccinia aegopogonis** Arth. and Holw. Brown leaf rust on *E. mairtianum*, *Aegopodium cenchroides*, and *A. tenellus* in Guatemala.
- Puccinia basisporula** Jaeks. and Holw. Leaf rust on *E. mairtianum* and *E. rafaense* in Guatemala.
- Puccinia cacheutensis** Speg. Brown rust pustules on leaves and branches of *E. patens* in Argentina.
- Puccinia espinosarum** Diet. and Holw. Brown to black rust pustules on leaves of *E. espinosarum*, *E. smithii*, *E. sordidum*, and *E. spiraeae-folium* in Mexico.
- Puccinia eupatorii** Diet. Leaf rust on *E. ballotae-folium*, *E. iresinoides*, *E. macrocephalum*, and *E. viridiflorum* in Trinidad.
- Puccinia eupatoriicola** Mayor. Leaf rust on *E. pycnocephalum* and *E. schiedeanum* in Colombia.
- Puccinia eupatorii-columbiani** Mayor. Leaf rust on *E. columbianum* and *E. involucratum* in Colombia and Trinidad.
- Puccinia hodgsoniana** Kern. Leaf rust on *E. phoenicolepis* and *E. schultzei* in Guatemala.
- Puccinia horrida** Lagh. Cinnamon-brown rust pustules on leaves of *Eupatorium* sp. in Ecuador.
- Puccinia inanipes** Diet. and Holw. Brown to black rust pustules on leaves of *E. brevipedis*, *E. hirsutum*, and *E. tubiflorum* in Mexico.
- Puccinia inermis** Jacks. and Holw. Powdery brown rust pustules on leaves of *Eupatorium* sp. in Costa Rica.
- Puccinia noackii** Syd. Dark-brown rust pustules on leaves of *Eupatorium* sp. in Brazil.
- Puccinia pachyspora** Diet. Black rust pustules on leaves and stems of *E. oblongifolium* in Brazil.
- Puccinia rosea** Roth. See *Ageratum*.
- Puccinia solidipes** Jacks. and Holw. Brown leaf rust on *E. tubiflorum* in Guatemala and Mexico.
- Puccinia tinctoria** Speg. Brown rust pustules on leaves of *E. prasifolium* and *E. tinctorium* in Argentina and Paraguay.
- Puccinosira eupatorii** Lagerh. Leaf rust on *E. aschenbornianum* and *Eupatorium* sp. in Guatemala and Ecuador.
- Septoria albo-maculans** Syd. Circular to irregular leaf spots with raised margins on *E. nubigenum* and *E. pomaderrifolium* in Colombia and Guatemala.
- Septoria eupatorii** Rob. and Desm. Numerous small subcircular gray-white leaf spots on *E. cannabinum* in Italy, France, and Germany. Reported from the lower Mississippi Valley.
- Syncarpella castagnei** (Speg.) Theiss. and Syd. Black stromata on brown, indistinct leaf spots on *E. tinctorium* in Brazil.
- Uredo eupatoriicola** P. Henn. Brown leaf rust on *Eupatorium* sp. in Brazil.
- Uredo scopigena** P. Henn. Rust on leaves and stems of *Eupatorium* sp. in Brazil.
- Uredo suspecta** Jaeks. and Holw. Brown rust pustules on leaves of *E. daleoides* in Costa Rica.

- EUPHORBIA.** SPURGE. Poinsettia. Plants with milky juice and very diverse habit, from succulent cactus-like plants to low or prostrate herbaceous weeds.
- Cercospora euphorbiae** Pat. Circular gray leaf spots on *Euphorbia* sp. in Ecuador.
- Colletotrichum euchroum** Syd. Anthracnose, destroying the leaves of *E. antiquorum* and *E. neriiifolia* in China and the Philippines.
- Endophyllum euphorbiae-silvaticae** (DC.) Wint. Yellow rust pustules on leaves of *E. amygdaloides* and *E. silvatica* in Europe.
- Erysiphe taurica** Lév. See Althaca.
- Melampsora euphorbiae** (Schub.) Cast. Yellow and brown to black rust pustules on leaves and stems of *Euphorbia* spp. in Europe, Egypt, and Tripoli. Sparingly introduced into the United States on *E. cyparissias*.
- Melampsora euphorbiae-amygdaloides** Muell. Leaf rust on *E. amygdaloides* in Europe.
- Melampsora euphorbiae-dulcis** Otth. Golden and brown to dark-brown rust pustules on leaves and stems of *Euphorbia* spp. in Japan and Europe.
- Melampsora euphorbiae-engleri** P. Henn. Leaf rust on *E. engleri* in Tanganyika.
- Melampsora euphorbiae-gerardiana** W. Muell. Rust on stems and leaves of *Euphorbia* spp. in Asia Minor, Tripoli, Persia, India, and Europe. Also known from Indiana.
- Melampsora gelmii** Bres. Yellow and dark-brown rust pustules on leaves of *Euphorbia* spp. in Persia, north Africa, and Europe.
- Melampsora helioscopiae** Wint. Yellow and dark-brown to black rust pustules on leaves and stems of *Euphorbia* spp. (over 30 reported species) in Europe, Egypt, Abyssinia, Italian North Africa, Tunis, Asia Minor, India, Japan, and Ceylon.
- Mycosphaerella canariensis** Tass. On stems of *E. canariensis* in Italy.
- Ovulariopsis erysiphoides** Pat. and Har. White erysiphoid spots on leaves of *E. balsamifera* in central Africa.
- Peronospora andina** Speg. Downy mildew deforming the leaves of *E. rhytisperma* in Argentina.
- Peronospora cyparissiae** De By. Downy mildew on *E. cyparissias* in central Europe.
- Peronospora esulae** Gäum. Downy mildew on leaves of *E. esula* in central Europe.
- Peronospora valesiaca** Gäum. Downy mildew on leaves of *E. gerardiana* in Switzerland.
- Phoma baldratii** Bacc. On stems of *E. tirucalli* in Eritrea.
- Phragmodothis asperata** Syd. On *Euphorbia* sp. in the Union of South Africa.
- Phyllosticta euphorbiaecola** P. Brun. Subcircular brown leaf spots on *E. amygdaloides* in France.
- Phyllostictella euphorbiae** (Roum.) Tass. On leaves of *E. silvatica* in France.
- Puccinia euphorbiae** P. Henn. and var. **longipes** Syd. Leaf rust on *E. agowensis* and *E. eritrea* in Abyssinia and *E. calyculata*, *E. cotinifolia*, and *Euphorbia* sp. in Mexico. Reported from Minnesota.
- Puccinia intumescens** (Syd.) Holw. Yellow aecial and black powdery telial sori on leaves of *E. calyculata* in Mexico.
- Ramularia euphorbiae** (Cast.) Sacc. Whitish areas on leaves of *E. peplus* in France.
- Septoria bractearum** Mont. Rufous spots on bracts on *E. serrata* in France and Spain.
- Septoria euphorbiae** Guep. Small subcircular olivaceous leaf spots on *E. amygdaloides*, *E. angulata*, *E. esula*, and *E. peplus* in Russia and France.
- Septoria euphorbicola** Höll. Circular olive-brown leaf spots on *E. procera* in Hungary.
- Septoria hariotiana** Sacc. White leaf spots with narrow dark-purple margins on *E. palustris* and *E. pilosa* in France and Spain.
- Septoria kalchbrenneri** Sacc. On leaves of *E. aspera*, *E. palustris*, and *E. silvatica* in Austria and Germany.
- Septoria media** Sacc. and Brun. Subcircular, then confluent, brown leaf spots with gray-white centers on *E. amygdaloides* and *E. palustris* in France and Yugoslavia.
- Septoria thuemeniana** Pass. Attacks leaves, stems, and twigs of *E. exigua* in Denmark and Italy.
- Uredo cornui** Har. Brown leaf rust on *Euphorbia* sp. in Ecuador.
- Uredo euphorbiae-nudiflorae** P. Henn. Dull-brown rust pustules on yellow leaf spots on *E. nudiflora* in Jamaica.
- Uredo velata** E. and E. Brown leaf rust on *E. cordata* in Hawaii.
- Uromyces alpestris** Tranzsch. Brown to black rust pustules on leaves and stems of *E. cyparissias* in Europe.
- Uromyces andinus** P. Magn. Powdery chestnut-brown rust pustules on leaves of *E. chilensis*, *E. collina*, and *E. portulacoides* in Chile and Argentina. Reported from Utah.
- Uromyces astragali** (Opiz.) Sacc. See Astragalus.
- Uromyces bresadolae** Tranzsch. Brown rust pustules on leaves of *E. angulata* in Austria.
- Uromyces eristulatus** Tranzsch. Dark-brown to black rust pustules on leaves of *E. girardiana* and *E. petrophila* in Bohemia.
- Uromyces excavatus** (DC.) Lév. Leaf rust on *E. angulata*, *E. dulcis*, *E. gloriosa*, *E. pubescens*, and *E. verrucosa* in Europe.
- Uromyces fischeri-eduardi** P. Magn. See Vicia.
- Uromyces haussknechtii** Tranzsch. Brown rust pustules on leaves of *E. spinosa* and *E. thamnoides* in Syria, Italy, Dalmatia, and France.
- Uromyces hermonis** P. Magn. Leaf rust on *E. caudiculosa* in Asia Minor.
- Uromyces kalmusii** Sacc. Chestnut-brown rust pustules on leaves of *E. cyparissias* and *E. esula* in Manchuria and central Europe.
- Uromyces kawakamii** Syd. Leaf rust on *E. serrulata* in Formosa.
- Uromyces laevis** Koernicke. Dark-brown rust sori on leaves of *E. gerardiana* in Russia, France, Switzerland, Holland, Austria, and Germany.
- Uromyces loti** Blytt. See Lotus.
- Uromyces mayerii** Tranzsch. Leaf rust on *E. orbiculata* in Colombia.
- Uromyces monspessulanus** Tranzsch. Brown rust pustules on leaves of *E. serrata* in France and Spain.
- Uromyces natalensis** P. Magn. Powdery brown rust pustules on leaves of *E. gueinzii* in the Union of South Africa.
- Uromyces pisi** (Pers.) Wint. See Pisum.
- Uromyces scutellatus** (Schrank.) Lév. Leaf rust on *Euphorbia* spp. in Europe. Reported from Colorado.
- Uromyces striatellus** Tranzsch. Brown leaf rust on *E. hebecarpa* in Persia.
- Uromyces striolatus** Tranzsch. Leaf rust on *E. cyparissias* and *Euphorbia* sp. in Turkestan, Italy, France, and Switzerland.
- Uromyces sublevis** Tranzsch. Dark-brown to black rust pustules on leaves of *Euphorbia* spp. in Caucasasia, Syria, and Europe.
- Uromyces tinctoriicola** P. Magn. Leaf rust on *Euphorbia* spp. in Kurdistan, Persia, Syria, Anatolia, Spain, France, Russia, Macedonia, Austria, and Germany.

**EUPHORBIA**—Continued.

**Uromyces tordillensis** Speg. Powdery brown rust pustules on leaves of *E. ovalifolia* and *E. serpens* in Argentina and Uruguay.

**Uromyces tuberculatus** Fckl. Cinnamon-brown to black rust pustules on leaves and stems of *E. exigua*, *E. hissarica*, and *E. platyphyllos* in Turkestan, Spain, France, Switzerland, and Germany.

**Uromyces uleanus** Diet. Brown, powdery, rust pustules on leaves of *E. stenophylla* in Brazil.

**Uromyces undulatus** Tranzsch. Leaf rust on *Euphorbia* sp. in Turkestan.

**Uromyces winteri** Wettst. Black rust pustules on leaves of *E. falcata* and *Euphorbia* sp. in Tripoli, Asia Minor, and Turkestan.

**EURYA**. Shrubs.

**Aulographum euryae** Syd. Large irregular leaf spots on *E. sinensis* in Japan.

**Catacauma euryae** (Racib.) Theiss. and Syd. Black stromata on circular, yellow-green leaf spots on *E. acuminata* and *E. japonica* in the Philippines and Java.

**Exobasidium euryae** Syd. and Butl. Causes hypertrophy of flowers and fruit of *E. acuminata* in India.

**Phyllachora transiens** Syd. and Butl. Black stromata on minute indefinite yellow leaf spots on *E. acuminata* in India.

**Septoria euryae** P. Henn. Dull-brown, then white leaf spots on *E. canescens* in Ecuador.

**EVELYNA**. See Orchidaceae.**EVODIA**. Ornamental shrubs grown for their handsome foliage.

**Bacterium citri** Hasse. See Citrus.

**Cercospora evodiae** Syd. Small yellow leaf spots on *E. meliaefolia* in Formosa.

**Coleosporium evodiae** Diet. Powdery golden rust pustules on leaves of *E. meliaefolia* in Japan and Formosa.

**Phyllosticta evodiae** Cke. Leaf spots on *E. accedens* in Australia.

**EVOLVULUS**. Annual or perennial prostrate or erect herbs.

**Cercospora balansae** Speg. On leaves of *Evolvulus* sp. in Brazil.

**Helminthosporium balansae** Speg. Indefinite, olivaceous spots, often involving entire leaf blades of *Evolvulus* sp. in Brazil.

**Puccinia desertorum** Syd. Yellow-brown rust pustules on leaves and stems of *E. alsinoides* in tropical Africa.

**Puccinia enecta** Speg. Dark-brown rust pustules on leaves and stems of *Evolvulus* sp. in Argentina.

**Puccinia tuyutensis** Speg. Brown rust pustules on leaves and stems of *E. falcatus* in South America.

**Uredo evolvuli** Speg. Powdery pale-brown rust pustules on leaves and stems of *E. nummularius* and *E. sericeus* in Argentina and Venezuela.

**EXACUM**. Herbs, cultivated for their blue, white, or lilac flowers.

**Coleosporium exaci** Syd. Golden rust pustules on leaves of *E. tetragonum* in the Philippines.

**FABA**. See Vicia.**FAGELIA**. See Calceolaria.**FAGOPYRUM**. BUCKWHEAT.

**Ascochyta fagopyri** Theum. and var. *italica* Trav. On leaves of *F. vulgare* (*F. esculentum*) in Russia, Italy, and Germany. Reported from Vermont.

**Fusicladium fagopyri** Oud. On leaves of *F. vulgare* (*F. esculentum*) in Holland.

**Phyllosticta polygonorum** Sacc. See Polygonum.

**Phytophthora parasitica** Dast. See Ricinus.

**Puccinia fagopyri** Barel. Brown to black rust pustules on leaves of *F. vulgare* (*F. esculentum*) in India.

**Ramularia curvula** Fautr. On leaves of *F. vulgare* in Europe.

**Sclerotinia fagopyri** Hori. Sclerotia form in diseased seed of *F. vulgare* in Japan.

**Sphaelotheca fagopyri** Syd. and Butl. Smut forming dark-violet sori in ovaries of *F. vulgare* in India.

**FAGUS**. BEECH. Timber and ornamental trees.

**Actinonema fagicola** Allesch. Subcircular spots causing premature leaf-fall and swelling of buds of *F. sylvatica* in Germany.

**Ascochyta fagi** Wor. On *Fagus* sp. in Russia.

**Clasterosporium epiphyllum** (Lév.) Sacc. On leaves and fruit of *F. sylvatica* in France.

**Cryptostictus niesslii** Oud. On leaves of *F. sylvatica* in Holland.

**Cyttaria berterii** Berk. Yellow to golden obovate or turbinate fruiting bodies on branches of *F. obliqua* in Patagonia.

**Cyttaria darwinii** Berk. Causes a wood rot of branches of *F. antarctica* and *F. betuloides* in Chile.

**Cyttaria gunnii** Berk. Rotting branches of *F. cunninghamii* in Tasmania and Victoria and fruiting in dense clusters.

**Cyttaria harioti** Fisch. Subglobose brown fruiting bodies on rotted branches of *F. antarctica* and *F. betuloides* in Chile.

**Cyttaria hookeri** Berk. Yellowish to cinnamon-colored fruiting bodies on rotted branches of *F. antarctica* and *F. obliqua* in Patagonia.

**Discosia maculiformis** Syd. On *F. sylvatica* in Japan.

**Laestadia faginea** (Cke. and Plowr.) Sacc. On leaves of *F. sylvatica* in Great Britain.

**Leptosphaeria valdobbiae** Ferr. On leaves of *F. sylvatica* in Italy.

**Linospora faginea** Sacc. Discolored areas on leaves of *F. sylvatica* in Italy and France.

**Melasmia antarctica** Speg. On branches of *F. antarctica* in Patagonia.

**Mikronegeria fagi** Diet. and Neg. Yellow to rufous rust pustules on leaves of *F. obliqua* and *F. procera* in Chile.

**Mycosphaerella fagi** (Awd.) Lind. On leaves of *F. sylvatica* in Europe.

**Phyllosticta fagi** Oud. Pale-brown, often confluent, leaf spots on *F. sylvatica* in Holland.

**Phyllosticta faginea** Bres. Small subcircular dull-brown leaf spots on *F. sylvatica* in Germany.

**Phytophthora fagi** R. Hartig. Downy mildew attacking the cotyledons of seedlings causing a serious "damping-off" disease in Europe. Among the hosts are *F. grandifolia*, *F. sylvatica*, *Abies* spp., *Acer palmatum*, *Acer* sp., *Fraxinus* spp., *Larix* sp., *Picea* sp., *Pinus montana*, *P. sylvestris*, and *Robinia pseudoacacia*. The cotyledons and basal portions of diseased stems turn brown.

**Plowrightia noxia** (Ruhl.) Sacc. See Castanea.

**Septoria fagi** Awd. and f. *amphigena* D. Sacc. On leaves of *F. sylvatica* in Italy, Germany, and Austria.

**Septoria fagicola** Speg. On leaves of *F. betuloides* in Patagonia.

**Sphaeronaema fagi** Oud. On leaves of *F. sylvatica* in Holland.

**Sphaeronaema fagineum** Oud. On leaves of *F. sylvatica* in Holland.

**Uncinula curvispora** K. Hara. Powdery mildew on leaves of *F. sylvatica* in Japan.

**FARADAYA**. Climbing shrubs allied to *Clerodendron*.

**Phyllosticta faradayae** Sacc. On leaves of *F. papuana* in Malaya.

- FATSIA.** Shrubs or small trees. Some species are put under *Tetrapanax*.  
**Aecidium fatsiae** Syd. Leaf rust on *Tetrapanax* (*F. papyrifera*) in Formosa.
- FEDIA.** See Patrinia.
- FEIJOA.** Tropical fruit trees.  
**Catacauma feijoa** (Rehm.) Theiss. and Syd. Shiny black stromata on leaves of *Feijoa* sp. in Brazil  
**Schizothyrium hypodermoides** Rehm. On leaves of *Feijoa* sp. in Brazil.
- FERONIA.** Spiny deciduous trees with hard-shelled fruits, related to *Citrus*.  
**Bacterium citri** Hasse. See Citrus.  
**Corticium salmonicolor** B. & Br. See Citrus.
- FERONIELLA.** Small spiny trees related to *Feronia*.  
**Bacterium citri** Hasse. See Citrus.  
**Corticium salmonicolor** B. & Br. See Citrus.
- FERULA.** GIANT FENNEL. Hardy herbs.  
**Aecidium ferulae** Rouss. and Dur. Rust pustules on leaves and petioles of *F. communis* and *F. tingitana* in French North Africa, Sardinia, France, Switzerland and Belgium.  
**Puccinia elliptica** Lindr. Leaf rust on *F. longifolia* in Russia.  
**Puccinia ferulae** Rud. Powdery yellow and dark-brown rust pustules on leaves and stems of *F. communis* and *Ferulaga galbanifera* in Italy, France, and Austria.  
**Puccinia libani** P. Magn. Rust on leaves and stems of *F. nodiflora*, *Cachrys gonicarpa*, *C. odonto calyx*, and *Ferulago* spp. in Asia Minor, Persia, Turkestan, Russia, Finland, Italy, and Greece.  
**Puccinia sogdiana** Kom. Brown rust pustules on leaves of *F. jaeschkeana* and *F. rigidula* in Turkestan.  
**Uromyces ferulae** Juel. Yellow and powdery brown rust pustules on leaf blades, petioles, and stems of *F. assoi* and *F. communis* in Algeria, Sicily, Corsica, and Spain.
- FESTUCA.** FESCUE. Annual or perennial grasses, some species cultivated for ornament or as pasture grass.  
**Cladochytrium graminis** Busgen. This fungus attacks and destroys the roots of *F. ovina*, *Dactylis glomerata*, *Panicum* sp., *Poa annua*, and other grasses in Great Britain and Germany. The disease kills out the grass in patches, which are small and yellow at first, finally becoming of considerable size, irregular in shape, and brown in color.  
**Dilophia graminis** (Fckl.) Sacc. See Triticum.  
**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.  
**Leptosphaeria culmicola** (Fr.) Karst. See Bromus.  
**Leptosphaeria kerguelensis** P. Henn. On leaves of *F. kerguelensis* in Kerguelen Island.  
**Ophiobolus cariceti** (B. & Br.) Sacc. See Triticum.  
**Phyllachora sylvatica** Sacc. Ovate to oblong dark-brown stromata on leaves of *F. duriuscula* in Spain and Italy.  
**Phyllosticta paleicola** P. Henn. On leaves and palea of *F. erecta* in Kerguelen Island.  
**Physalospora festucae** (Lib.) Sacc. See Andropogon.  
**Puccinia gibberosa** Lagerh. Dark-brown to black rust pustules on leaves of *F. sylvatica* in Switzerland and Germany.  
**Puccinia mellea** Diet. and Neg. Brown leaf rust on *F. muralis* in Chile.  
**Sclerospora macrospora** Sacc. See Triticum.  
**Septoria festucae** Died. On leaves of *F. gigantea* in Central Europe.  
**Septoria festucae-sylvaticae** Died. On leaves of *F. sylvatica* in Central Europe.  
**Tilletia sterilis** Ule. Leaf smut on *F. ovina* and *Koeleria cristata* in Germany.  
**Uredo abscondita** Fautr. Brown linear rust pustules on leaves of *F. heterophylla* in France.  
**Uredo kerguelensis** P. Henn. Brown rust pustules on yellow leaf spots on *F. kerguelensis* in Kerguelen Island.  
**Urocystis festucae** Ule. Smut on culms and leaves of *F. ovina* in Germany.  
**Uromyces cuspidatus** Wint. Elongate dark-brown rust sori on leaves of *F. commersonii* in Chile.  
**Uromyces festucae** Syd. Black rust pustules on leaves of *F. arundinacea*, *F. ovina*, and *F. rubra* in Spain, Bohemia, Russia, Sweden, and Germany. The aecial stage occurs on *Ranunculus bulbosus* and *R. illyricus*.  
**Uromyces fuegianus** Speg. Leaf rust on *F. purpurascens* in Chile.  
**Uromyces ranunculi-festucae** Jaap. Powdery golden to brown rust pustules on leaves of *F. ovina* in Germany. The aecial stage occurs on *Ranunculus bulbosus*.  
**Ustilago festucae-tenellae** P. Henn. Rachides and flowers deformed and converted into olivaceous-black powdery masses of smut spores. On *F. tenella* in Mexico.  
**Ustilago sphaerocarpa** Syd. Black powdery smut sori replacing ovaries of *F. amplissima* in Mexico.
- FICARIA.** See Ranunculus.
- FICUS.** FIG. Banyan. A large genus of tropical trees, shrubs, and climbers. Many species of economic importance for fruit, ornament, rubber, and other products.  
**Acrotheciella javanica** Koord. On leaves of *F. elastica* in Java.  
**Alteritaria fici** Farneti. Causes atrophy of fruit of *F. carica* in Italy.  
**Anisochora topographica** (Speg.) Theiss. and Syd. Black irregular stromata on leaves of *Ficus* sp. in Brazil.  
**Aphelenchus olesistus** Ritz. Bos. See Begonia.  
**Ascochyta caricae** Rabh. Red-brown leaf spots on *F. carica* in France and Austria.  
**Atichia treubii** V. Hoeh. Superficial fruiting bodies on leaves of *F. elastica* in Java.  
**Bacterium fici** Cav. This bacterium is said to cause a browning of the tissues of twigs of *Ficus* sp in Italy.  
**Catacauma apoense** Syd. Scattered irregular dull-black stromata on leaves of *F. apoensis*, *F. nervosa*, and *F. sibulanensis* in the Philippines.  
**Catacauma aspideum** (Br.) Theiss. and Syd. and ff. **fici-albae** and **spinifera** Theiss. and Syd. Black stromata on leaves of *F. alba*, *F. banahaensis*, *F. fastigiata*, *F. fulva*, *F. heterophylla*, *F. minahassae*, *F. odorata*, *F. repens*, *F. riedeli*, *F. scandens*, *F. spinifera*, *F. tomentosa*, *F. ulmifolia*, and *F. validicaudata* in Java, India, Ceylon, Malaya, the Philippines, and the Union of South Africa.  
**Catacauma circinatum** Syd. Circular black stromata on leaves of *F. chrysolepis* and *F. odorata* in the Philippines.  
**Catacauma decaisneanum** (Lév.) Theiss. and Syd. Black stromata on leaves of *F. laeta* in Timor.  
**Catacauma elmeri** Syd. Small scattered black stromata on leaves of *F. blepharostoma*, *F. manilensis*, *F. minahassae*, and *F. ulmifolia* in the Philippines.  
**Catacauma fici-fulvae** Koord. Black stromata on leaves of *F. sinuosa* and *F. validicaudata* in the Philippines.  
**Catacauma fici-obscurae** (Koord.) Theiss. and Syd. Shiny-black stromata on brown leaf spots on *F. obscura* in Java.

## FICUS—Continued.

- Catacauma garciae** Theiss. and Syd. Black stromata on small greenish leaf spots on *F. garciae* in the Philippines.
- Catacauma gramicum** (P. Henn.) Theiss. and Syd. Black linear stromata on yellow-brown leaf spots on *F. capensis* in the Congo and the Union of South Africa.
- Catacauma infectorum** (Cke.) Theiss. and Syd. Shiny-black irregular stromata on leaves of *F. infectoria*, *F. religiosa*, and *F. ulmifolia* in India and the Philippines.
- Catacauma irregulare** (W. and C.) Theiss. and Syd. Black stromata on leaves of *Ficus* sp. in Angola.
- Catacauma karnbachii** (P. Henn.) Theiss. and Syd. Dull-black stromata on circular brown leaf spots on *F. flavocortica*, *F. heterophylla*, and *F. ulmifolia* in New Guinea and the Philippines.
- Catacauma lagunense** Syd. Shiny-black stromata on leaves of *F. hauili* in the Philippines.
- Catacauma microcentrum** (B. and Br.) Theiss. and Syd. and var. **graphica** Theiss. and Syd. Circular black stromata on leaves of *F. mysorensis*, *Ficus* sp., and *Artocarpus* sp. in Ceylon, India, and the Philippines.
- Catacauma merrillii** Syd. Black stromata on leaves of *F. lucbanensis* in the Philippines.
- Catacauma microplicatum** Syd. Tar-spot disease of leaves of *Ficus* sp. in Amboina.
- Catacauma nipponicum** Syd. Black circular shiny stromata on leaves of *F. nipponica* in Japan.
- Catacauma repens** (Cd.) Theiss. and Syd. Shiny-black stromata on leaves of *F. gossypina* and *F. religiosa* in Uganda, the Congo, and India.
- Catacauma robinsonii** Syd. Tar-spot on leaves of *Ficus* sp. in Amboina.
- Catacauma sanguineum** Theiss. and Syd. Small black stromata on reddish-yellow leaf spots on *F. heterophylla* and *F. odorata* in the Philippines.
- Catacauma schweinfurthii** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *F. palmata* in Abyssinia.
- Catacauma ulcerata** (Mass.) Theiss. and Syd. Dull-black stromata on leaves of *F. ovata* in Uganda.
- Catacauma urophyllum** (V. Hoeh.) Theiss. and Syd. Linear black stromata on leaves of *F. urophylla* in Java.
- Catacauma valsiforme** (Rehm.) Theiss. and Syd. Black stromata on brown leaf spots on *F. crassiflora* in the Philippines.
- Cercospora annulata** Cke. Circular, concentrically zoned, pale-brown leaf spots on *F. hispida* in India.
- Cercospora elastica** A. Zimm. Gray-brown leaf spots on *F. elastica* in Java.
- Cercospora rubro-cincta** Pat. On leaves of *Ficus* sp. in Indo-China.
- Cercospora rufula** Syd. Large circular rufous leaf spots on *Ficus* sp. in British North Borneo.
- Cercospora urostigmatis** P. Henn. Circular to angular brown leaf spots on *Ficus* (*Urostigma*) sp. in Brazil.
- Cladosporium sicophilum** Farneti. On fruit of *F. carica* in Italy.
- Clasterosporium elasticae** Koord. On bark of *F. elastica* in Java.
- Clasterosporium javanicum** Koord. On leaves of *F. elastica* in Java.
- Coccodothella placida** Syd. Black stromata on upper leaf surfaces on *F. oreodryadum* in Tanganyika.
- Coleroa elasticae** Koord. On leaves and branches of *F. elastica* in Java.
- Corticium salmonicolor** B. and Br. See Citrus.
- Diplodia cacaoicola** P. Henn. See Theobroma.
- Eutypa erumpens** Mass. See Theobroma.
- Fomes lamaoensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Fusicladium elasticae** Koord. On leaves of *F. elastica* in Java.
- Fusicladium fici** Bacc. Circular brown leaf spots with red-brown borders on *Ficus* sp. in Abyssinia.
- Gloeosporium piuggarii** Speg. On leaves of *Ficus* (*Urostigma*) sp. in Brazil.
- Gloeosporium sycophilum** Trinch. Anthracnose on leaves of *F. elastica* in Italy.
- Helminthosporium elasticae** Koord. On leaves of *F. elastica* in Java.
- Helminthosporium ficinum** Sacc. On *F. wassa* and *Ficus* sp. in the Philippines and Amboina.
- Helminthosporium ficuum** Rostr. Ashen leaf spots with dull-brown margins on *F. retusa* and *Ficus* sp. in Siam.
- Helminthosporium ficuum** Yates. On leaves of *F. caudatifolia* and *F. ulmifolia* in the Philippines. This species, if distinct, should be renamed, the name being occupied.
- Mycosphaerella elasticae** Koord. On leaves of *F. elastica* in Java.
- Napicladium elasticae** Koord. On leaves of *F. elastica* in Java.
- Phaeosaccardinula ficicola** P. Henn. Effuse dull-brown leaf spots on *Ficus* sp. in Tanganyika.
- Phyctaena ficuum** P. Henn. Circular yellow leaf spots on *Ficus* sp. in Brazil.
- Phomopsis cinerescens** (Sacc.) Trav. Large cankers occur on the branches, rotting away the bark and sapwood and girdling and destroying involved limbs of *F. carica* in Great Britain, France, and Italy.
- Phyllachora amamensis** P. Henn. Circular to irregular groups of dull-black stromata on leaves of *F. capensis* and *Ficus* sp. in Tanganyika and the Union of South Africa.
- Phyllachora aspideoides** Sacc. and Berl. Dull-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora catervaria** (Br.) Sacc. Round black stromata on leaves of *F. hispida*, *F. merrillius*, and *F. oppositifolia* in India, Indo-China, Java, the Philippines, and Amboina.
- Phyllachora cayennensis** (DC.) Theiss. and Syd. A doubtful species on leaves of *F. roxburghi* and *Psidium guajava* in Brazil, Surinam, and India.
- Phyllachora dawci** Mass. Tar-spot on leaves of *Ficus* sp. in Uganda.
- Phyllachora devriesei** Koord. Black stromata on red-brown leaf spots on *F. leucantatoma* in Java.
- Phyllachora effigurata** Syd. Dull-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora ficicola** Allesch. and P. Henn. Round to elliptical shiny-black stromata on leaves of *Ficus* sp. in Brazil.
- Phyllachora pseudos** Rehm. Black stromata on leaves of *F. minahassae* and *F. nota* in the Philippines.
- Phyllachora vinosa** Speg. Smooth black stromata on violet leaf spots on *Ficus* sp. in Brazil.
- Phyllosticta ambiguella** Sacc. Large irregular gray-white leaf spots with narrow rufous margins on *F. rubiginosa* in Italy.
- Phyllosticta caricae** C. Mass. On leaves of *F. carica* in Italy.
- Phyllosticta elasticae** Koord. On leaves of *F. elastica* in Java.
- Phyllosticta fici** Bres. Circular gray-white leaf spots with dull-brown margins on *F. macrophylla* in Portuguese St. Thomas.
- Phyllosticta ficicola** Pat. Circular gray-white leaf spots on *F. carica* in Tunis and Italy.

**FICUS**—Continued.

- Phyllosticta roberti** Boy. and Jacz. Pale-brown leaf spots on *F. aurea* and *F. elastica* in France and the Bahamas. Reported from Florida.
- Phyllosticta sycina** Trav. On leaves of *F. heterophylla* in Italy.
- Phyllosticta sycophila** Thuem. Large brown leaf spots on *F. carica* in Victoria and Italy.
- Phyalospora atractina** Syd. Small gray-brown leaf spots on *Ficus* sp. in Brazil.
- Phyalospora clasticæ** Koord. Attacks branches of *F. elastica* in Java as a wound parasite.
- Phytophthora carica** (Hara) Hori. A downy mildew on fruit of *F. carica* in Japan. Small, sunken areas appear which soon involve the entire fruit in a soft brown rot. A thick cottony mycelium covers the surface of rotted fruits, accompanied by a disagreeable odor. Fruits remaining on the trees mummify.
- Phytophthora fici** Ven. Causes a soft rot of fruit of *F. carica* in India.
- Puccinia sepulta** B. and C. Black leaf rust on *Ficus* sp. in Nicaragua.
- Ramularia sycina** Sacc. and D. Sacc. On leaves of *Ficus* sp. in Italy.
- Rosellinia bunodes** Sacc. See Citrus.
- Rosellinia echinata** Mass. Causes a serious decay of roots, bringing about death of infected trees of *F. dubia* in Malacca and Malaya.
- Schizochora elmeri** Syd. Circular to elliptical dull-black stromata on lower leaf surfaces of *F. guyeri* in the Philippines.
- Septogloeum elasticæ** Koord. Subcircular light-brown leaf spots on *F. elastica* in Java.
- Septoria arcuata** Cke. White circular leaf spots with purple margins on *F. benghalensis* and *Ficus* sp. in India and Abyssinia.
- Septoria brachyspora** Sacc. Pale-brown leaf spots with ochraceous margins on *F. elastica* in Denmark and Italy.
- Septoria elastica** Koord. On leaves of *F. elastica* in Java.
- Septoria pipulæ** Cke. Irregular purple-brown leaf spots on *F. religiosa* in India.
- Septoria pirottae** Tassi. Brown leaf spots with rufous margins on *F. pumila* (*F. repens*) in Italy.
- Septoriella conformis** Sacc. On leaves of *F. alba* in Malaya.
- Sporodesmium sicynum** Thuem. Spots on leaves and stems of *F. carica* in Italy and Germany.
- Trabutia abyssinica** (P. Henn.) Theiss. and Syd. Irregular shiny-black stromata on leaves of *F. sycamorus* and *Ficus* sp. in Abyssinia and India.
- Trabutia amboiensis** Syd. On leaves of *F. hasskarlii* in Amboina.
- Trabutia bengetensis** Yates. On leaves of *F. benquetensis* in the Philippines.
- Trabutia butleri** Theiss. and Syd. Black stromata on leaves of *Ficus* sp. in India.
- Trabutia chinense** Yates. Black stromata on leaves of *Ficus* sp. in the Philippines.
- Trabutia elmeri** Theiss. and Syd. Scattered irregular black stromata on leaves of *F. banahaensis* in the Philippines.
- Trabutia evansii** Theiss. and Syd. Small circular black stromata on yellow or brown leaf spots on *Ficus* sp. in Portuguese East Africa.
- Trabutia fici-dekdekenæ** Bacc. Black stromata on leaves of *F. dekdekena* in Abyssinia.
- Trabutia fici-hochstetteri** Bacc. Tar spot on leaves of *F. hochstetteri* in Abyssinia.
- Trabutia ficuum** (Niessl.) Theiss. and Syd. Shiny-black irregular stromata on leaves of *F. foveolata*, *F. garciae*, *F. infectoria*, *F. mysorensis*, *F. nipponica*, and *F. riedelii* in Japan, India, Portuguese East Africa, and the Union of South Africa.
- Trabutia incrustans** Racib. Black irregular stromata on leaves of *Ficus* sp. in Java.
- Trabutia nervisequens** (Lingl.) Theiss. and Syd. Black stromata on yellow leaf spots on *F. hochstetteri* and *Ficus* sp. in Abyssinia and the Union of South Africa.
- Trabutia novoguineensis** Theiss. and Syd. Black stromata on rust-brown leaf spots on *Ficus* sp. in New Guinea.
- Trabutia vernicosa** Theiss. and Syd. Irregular black stromata on leaves of *F. heterophylla* and *F. ulmifolia* in the Philippines.
- Trabutiella microthryoides** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *F. erecta* in Japan.
- Uncinula pirottiana** Bacc. Powdery mildew on leaves of *Ficus* sp. in Abyssinia.
- Volutella** sp. Causes a trunk and branch canker of *F. elastica* in Italy.
- FILIPENDULA.** MEADOWSWERT. Hardy herbs grown for their showy panicles of varicolored flowers.
- Ascochyta obducens** Fckl. Large brown irregular spots destroying the leaves of *F. ulmaria* in Siberia and Germany.
- Colletotrichum volutella** Sacc. and Malbr. On stems of *F. ulmaria* in France.
- Cylindrosporium filipendulæ** Thuem. Pale-yellow areas on leaves of *F. ulmaria* and *Spiraea salicifolia* in Denmark and Austria. Reported from Washington.
- Mycosphaerella maculans** Sacc. and Roum. On leaves of *F. ulmaria* in Russia, France, Belgium, and Germany.
- Phyllosticta filipendulæ** Sacc. Subcircular gray-white leaf spots on *F. hexapetala* in Italy.
- Phyllosticta filipendulina** Sacc. and Syd. and var. **ulmariae** Sacc. Subcircular dull-brown leaf spots on *F. ulmaria* in Italy and France.
- Phyllosticta ulmariae** Thuem. Small irregular gray-white leaf spots on *F. ulmaria* in Scotland and Siberia.
- Ramularia ulmariae** Cke. Subcircular white spots on leaves of *F. hexapetala* and *F. palmata* in France, Denmark, and Germany.
- Septoria quevillensis** Sacc. Small dark-red leaf spots with pale centers on *S. ulmaria* in France.
- Triphragmium anomalum** Tranz. Leaf rust on *F. palmata* in Russia.
- Triphragmium filipendulæ** (Lasch.) Pass. Orange-yellow rust pustules on leaf blades and petioles of *F. hexapetala* and *Ulmaria pentaphylla* in central and northern Europe.
- Urocystis filipendulæ** Fckl. Black powdery smut sori on leaf petioles and nerves of *F. hexapetala* in Estonia, Sweden, Denmark, and Germany.
- FLA COURTIA.** Tropical shrubs, one species (Governor plum) cultivated for its fruit.
- Uredo uguressæ** Petch. Rust on fruit of *F. ramontchi* in Ceylon.
- FLEMINGIA.** Shrubs used as ornamentals.
- Uromyces flemmingiae** P. Henn. Brown powdery rust pustules on leaves of *Flemingia* sp. in the Congo.
- FLUGGÆA.** Tropical shrubs, sometimes cultivated.
- Nothoravenia japonica** Diet. See Securinega.
- FOENICULUM.** FENNEL. Annual, biennial, or perennial herbs, sometimes cultivated.
- Ascochyta foeniculina** McAlp. On fruit of *F. vulgare* in Australia.
- Cercospora foeniculi** P. Magn. On leaves of *F. officinale* in Austria.
- Erysiphe taurica** Lév. See Althæa.
- Mycosphaerella himantia** (Pass.) Died. Leaf spot of *Foeniculum* sp. in Dalmatia.

- FORSYTHIA.** GOLDEN BELL. Shrubs grown for their abundance of yellow spring flowers.
- Marsonia forsythiae** Lind. Large ashen leaf spots with dull brown margins on *F. suspensa* (*F. fortunei*) in Denmark.
- Phyllosticta forsythiae** Sacc. Subcircular ochraceous leaf spots on *F. suspensa* in Denmark, Spain, France, and Italy.
- Phytophthora syringae** Kleb. See Syringa.
- FORTUNELLA.** KUMQUAT. Small trees, related to *Citrus*, grown for their yellow ornamental and edible fruit.
- Bacterium citri** Hasse. See Citrus.
- FRAGARIA.** STRAWBERRY.
- Aphelenchus fragariae** Ritz. Bos. This nematode causes a fasciation of *Fragaria* sp. in Europe known as the "brush" or "cauliflower" disease. Fleshy malformations take the place of leaves and flowers, or the flowers may appear free of the deformed leaves and be in turn caused to assume odd shapes. The diseased areas assume a deep blood-red color. Infested plants are stunted and fail to form fruit.
- Aphelenchus olesistus** Ritz. Bos. See Begonia.
- Peronospora fragariae** Roze. and Cornu. Downy mildew on leaves of *F. vesca* in France, Switzerland, Italy, and Russia. Also reported from Iowa.
- Pheospora fragariae** (Br. and Har.) Petr. On leaves of *F. vesca* and *Potentilla caulescens* in Europe.
- Phragmidium fragariastris** (DC.) Schroet. See Potentilla.
- Phyllosticta grandimaculans** Bub. and Krieg. Brown leaf spots on *F. vesca* in Denmark and Germany.
- Puccinia fragariae** Barcl. Yellow to brown rust pustules on leaves of *F. vesca* in India.
- Septoria fragariae** Desm. Large circular brown leaf spots with red margins on *Fragaria* spp. in Brazil, Chile, Siberia, and Europe. Possibly only another name for the imperfect stage of *Mycosphaerella fragariae*.
- Thecaphora pallescens** Fingerh. Doubtful species of leaf smut on *F. collina* in Germany.
- Tylenchus dipsaci** Kuehn. See Narcissus.
- FRANKENIA.** Low perennial evergreen shrubs.
- Puccinia frankeniae** Link. Leaf rust on *F. hirsuta*, *F. laevis*, and *F. pulverulenta* in Egypt, Tunis, Portugal, France, Belgium, Switzerland, and Russia.
- FRAXINUS.** ASH. Ornamental, shade, and timber trees.
- Ascochyta fraxini** Kab. and Bub. (*A. fraxini* Oud.) On branches of *F. excelsior* and *F. ornus* in Europe.
- Ascochyta fraxinicola** Brun. Circular to oblong brown spots on twigs of *F. excelsior* in France.
- Ascochyta metulispora** B. and Br. Circular brown leaf spots on *Fraxinus* sp. in Scotland.
- Ascochyta orni** Sacc. and Speg. On leaves of *F. ornus* in Italy.
- Cercospora fraxini** (DC.) Sacc. Large brown areas on leaves of *F. excelsior* and *F. ornus* in Russia and Italy.
- Cercospora lumbricoides** Turc. and Maffei. Subcircular to oblong brown leaf spots with red-brown margins on *Fraxinus* sp. in Italy.
- Laestadia fraxini** Fautr. On leaves of *Fraxinus* sp. in France.
- Phyllosticta ambigua** Scalia. On leaves of *F. ornus* in Italy.
- Phyllosticta diedickei** Bub. and Syd. On leaves of *F. excelsior* in Germany.
- Phyllosticta fraxinicola** Carr. Subcircular to irregular dull-brown leaf spots on *F. excelsior* in France, Russia, and Germany. Reported from several localities in the United States.
- Phyllosticta orni** Bub. Circular dull-yellow leaf spots with narrow brown margins on *F. ornus* in Hungary.
- Phyllosticta osteospora** Sacc. See Morus.
- Phyllosticta trappenii** Oud. Large ochraceous leaf spots on *F. juglandifolia* in Holland.
- Phytophthora fagi** R. Htg. See Fagus.
- Puccinia fraxini** Kom. Powdery dark-brown rust pustules on leaves of *F. longicuspis* and *F. rhynchophylla* in Manchuria and Japan.
- Sclerotium scutellatum** A. S. See Acer.
- Scolecotrichum fraxini** Pass. Attacks the leaves of *F. ornus* in Italy.
- Septoria elaeospora** Sacc. Indefinite ochraceous leaf spots with brown margins on *F. excelsior*, *F. ornus*, and *F. pubescens* in Italy and Portugal.
- Septoria orni** Pass. Dull-brown effuse leaf spots on *F. excelsior* and *F. ornus* in Denmark, France, and Italy.
- Uncinula fraxini** Miy. Powdery mildew on leaves of *F. longicuspis* in Japan.
- Uncinula salmoni** Syd. Powdery mildew on leaves of *F. bungeana* in Japan.
- Uncinula sengokui** Salm. Powdery mildew on leaves of *F. bungeana* in Japan.
- Venturia fraxini** (Fr.) Aderh. On leaves of *F. excelsior* in northern Europe.
- FREESIA.** Cormous plants, cultivated for their flowers.
- Phoma bulbicola** Tassi. On bulbs of *F. odorata* in Italy.
- Uromyces ecklonii** Bub. Yellow-brown to black rust pustules on leaves of *F. refracta* in the Union of South Africa.
- Uromyces freesiae** Bub. Brown leaf rust on *F. odorata* in the Union of South Africa.
- FREYCINETIA.** Climbing shrubs.
- Sphaerodothis merrillii** (P. Henn.) Theiss. and Syd. Circular black stromata, shiny above, dull below, on leaves of *F. ensifolia* and *F. williamsii* in the Philippines.
- Uredo freycinetiae** Rac. Rust on yellow, sunken leaf spots on *F. imbricata* in Java.
- FRITILLARIA.** FRITILLARY. Crown-imperial. Guinea-hen flower. Hardy, low-growing, spring-blooming bulbous plants.
- Uromyces aecidiiformis** (Schlecht.) Thuem. See Lilium.
- Uromyces fritillariae** (Schlecht.) Thuem. Yellow-brown to chestnut-brown rust pustules on leaf blades and petioles and stems of *F. meleagris* in Europe.
- Uromyces lilii** (Link.) Fckl. See Lilium.
- Uromyces miurae** Syd. Leaf rust on *F. kamtschatensis* in Japan.
- Uromyces mogianensis** Bub. Brown powdery rust pustules in sunken yellow leaf spots on *F. bucharica* and *F. guicciardia* in Greece, Morocco, and Turkestan.
- FUCHSIA.** Shrubs or small trees, cultivated for their flowers.
- Coleosporium fuchsiae** Cke. Yellow rust pustules on leaves of *F. excorticata* in New Zealand.
- Endomyces meliolicola** Rehm. On leaves of *Fuchsia* sp. in Brazil.
- Phomopsis fuchsiae** (Brun.) Sacc. On stems of *Fuchsia* sp. in Bohemia.
- Phyllosticta fuchsicola** Speg. Gray-white circular leaf spots with broad purple margins on *F. coccinea* in Chile.
- Puccinia fuchsiae** Syd. and Holw. Leaf rust on *F. microphylla*, *F. thymifolia*, and *Lopezia hirsuta* in Guatemala, Mexico, Costa Rica, and Panama.
- Septoria fuchsiae** Roum. Irregular pale-brown leaf spots on *F. coccinea* in France.

**FUCHSIA**—Continued.

- Septoria fuchsicola** Syd. Subcircular to irregular brown leaf spots with reddish-purple raised margins on *F. coccinea* in Germany.
- Uredo fuchsiae** Arth. and Holw. Leaf rust on *F. splendens* in Guatemala.
- FUMARIA**. FUMITORY. Annuals cultivated for their flowers.
- Ascochyta fumariae** Höll. Indefinite brown areas on leaves of *F. schleicheri* in Hungary.
- Peronospora affinis** Rossm. Downy mildew on leaves of *F. capreolata*, *F. macrocarpa*, *F. officinalis*, *F. parviflora*, and *F. vaillanti* in Tunis and Europe.
- Ramularia fumariae** Speg. Small circular ashen leaf spots with violet borders on *F. capreolata* in Argentina.
- Rhizoctonia napi** West. See Brassica.
- FUNCKIA**. See Hosta.
- FUNTUMIA**. Tropical trees, one species cultivated for rubber.
- Colletotrichum funtumiae** Petch. Anthracnose on leaves of *F. elastica* in Ceylon.
- Fomes lamaoensis** Murr. See Hevea.
- Nectria funtumiae** Mass. Said to cause cankers on trunks of *F. elastica* in Uganda.
- FURCRAEA**. Succulent desert plants resembling *Agave*.
- Physalospora fourcroyae** P. Henn. On leaves of *F. gigantea* in Tanganyika.
- GAGEA**. Liliaceous herbs with white, yellow, or rose flowers.
- Heterosporium ornithogali** Klotzsch. See Ornithogalum.
- Puccinia megatherium** Syd. Cinnamon-colored rust pustules on both leaf surfaces of *G. reticulata* in the Caucasus.
- Puccinia pachyderma** Wettst. Powdery cinnamon-colored rust pustules on both leaf surfaces of *G. caucasica*, *G. damascena*, *G. graeca*, *G. lutea*, *G. persica*, and *G. pusilla* in Assyria, Russia, and Karpatos Island (Asiatic Turkey).
- Septoria commutata** Bub. Indefinite yellow, then grayish, leaf spots on *G. lutea* and *G. pratensis* in Bohemia and Galicia.
- Synchytrium laetum** Schroet. (*Pycnochytrium laetum* Schroet.) Forms small sulphur to golden-yellow punctiform galls on leaves, sepals and peduncles of *G. arvensis*, *G. fascicularis*, *G. lutea*, *G. silvatica*, *G. minima*, *G. pratensis*, and *Tulipa sylvestris* in Japan, Norway, Denmark, Russia, Austria, and Germany.
- Synchytrium punctatum** Schroeter. Forms small dirty-white to yellowish swellings or gall-like structures on leaves of *G. pratensis* in Austria and Germany.
- Uromyces gageae** G. Beck. Brown powdery rust pustules on both surfaces of leaves of *G. lutea*, *G. pratensis*, and *G. silvatica* in Europe.
- Uromyces ornithogali** Lévy. Linear brown to dark-brown rust pustules on leaves of *G. arvensis*, *G. bohémica*, *G. lutea*, *G. minima*, *G. pratensis*, *G. pusilla*, *G. saxatilis*, *G. stenopetala*, *Muscari racemosum*, *Ornithogalum nanum*, *O. nutans*, and *O. umbellatum* in France, Great Britain, Spain, Portugal, Hungary, Persia, Austria, and Germany.
- Ustilago ornithogali** (Schm. and Kze.) Kuehn. Olive-brown, powdery smut pustules on leaves and scapes of *G. arvensis*, *G. bohémica*, *G. fibrosa*, *G. liotardi*, *G. pratensis*, *G. lutea*, *G. minima*, *G. pusilla*, *G. saxatilis*, *G. spathacea*, and *G. stenopetala* in Denmark and Italy. **U. ornithogali-erythronii** Clint. has been reported on *Erythronium* sp. in the United States.
- GAILLARDIA**. Showy annual and perennial herbs grown in flower gardens and borders.
- Septoria gaillardiae** Speg. Ashen leaf spots with dull-brown margins on *G. doniana* in Argentina.
- GALACTIA**. Prostrate or twining leguminous, perennial herbs or shrubs, mostly tropical.
- Phyllachora galacteae** Earle. Irregular black stromata on leaves of *G. rudolphoides*, *G. striata*, and *G. tenuiflora* in the Bahamas and Porto Rico.
- GALANTHUS**. SNOWDROP. Spring-blooming bulbs with solitary white flowers.
- Botrytis galanthina** (B. and Br.) Sudw. Attacks the host plants as they appear above ground, distorting and finally rotting the leaves and flowers. A gray mold followed by black sclerotia forms over the diseased areas. The host is *G. nivalis* in England, Denmark, Holland, and Germany.
- The species of *Botrytis* and *Sclerotium* attacking the various ornamental bulbous plants have never been carefully studied as to their interrelationships and it is not clear whether few or many species are involved, or whether one or several occur in the United States. With this fact in mind, importations of bulbs should be carefully watched.
- Melampsora galanthi-fragilis** Kleb. A rust characterized by sunken yellow spots up to 1.5 cm. in diameter on both surfaces of the leaves of *G. nivalis*, *Salix fragilis* and other *Salix* spp. in Russia, Bulgaria, Spain, Switzerland, Austria, Hungary, and Germany.
- Puccinia galanthi** Unger. Rust sori, powdery, round to elliptical, up to 6 mm. in diameter, brown to black in color. On *G. nivalis* in Austria, Hungary, and Germany.
- Urocystis galanthi** Pape. Smut sori in blue-gray linear spots on leaf blades, sheaths, and flower stalks of *G. nivalis* in Germany.
- GALEGA**. GOAT'S-RUE. Bushy perennial leguminous plants.
- Ascochyta galegae** Höll. Irregular dull-brown leaf spots on *G. officinalis* in Hungary.
- Cercospora galegae** Sacc. Oblong white leaf spots on *G. officinalis* in Italy.
- Ramularia galegae** Sacc. Subcircular sordid-white leaf spots on *G. officinalis* in Italy and France.
- Uromyces galegae** (Opiz.) Sacc. Leaf rust on *G. officinalis* in Italy and the Balkans.
- Uromyces genistae-tinctoriae** (Pers.) Wint. See Caragana.
- GALEOPSIS**. Annual weeds of the mint family.
- Phyllosticta galeopsidis** Sacc. Ochraceous leaf spots on *G. versicolor* in Italy.
- Septoria cotylea** Pat. and Har. Indefinite brown leaf spots on *G. ceratonia* and *G. tetrahit* in Spain, France, and Denmark.
- GALIUM**. BEDSTRAW. Weak, slender herbs with square stems and whorled leaves.
- Aecidium molluginis** Wurth. Leaf rust on *G. mollugo* in Switzerland.
- Calloria quitensis** Pat. On *Galium* sp. in Ecuador.
- Fusicladium ruthenicum** Petr. On leaves of *G. schultesii* in Europe.
- Peronospora galii** Fekl. Downy mildew on leaves of *G. mollugo*, *G. palustre*, and *G. uliginosum* in central and north Europe.
- Peronospora galii veri** Gäum. As above on *G. verum*.
- Peronospora silvatica** Gäum. Downy mildew on leaves of *G. schultesii* and *G. silvaticum* in Switzerland, Hungary, Austria, and Germany.
- Phakopsora punctiformis** (Barel. and Diet). Diet. Yellow-brown rust pustules on leaves of *G. aparine* in India.
- Phyllosticta decipiens** C. Mass. On leaves of *G. mollugo* in Italy.
- Physalospora molluginis** (Oth.) Sacc. On leaves and stems of *G. mollugo* in Switzerland.
- Placosphaeria galii** Sacc. On stems of *Galium* spp. in Europe.
- Pseudopeziza autumnalis** (Fr.) Karst. Black spots on leaves and stems of *G. boreale*, *G. mollugo*, *G. uliginosum*, *Asperula odorata*, and *Rubia tinctorum* in Siberia, Great Britain, France, Belgium, and Germany.

**GALIAM**—Continued

- Puccinia celakovskyana** Bub. Powdery brown rust pustules on leaves and stems of *G. cruciata* in Europe.
- Puccinia eximia** Arth. and Holw. Brown leaf rust on *G. mexicanum* in Guatemala.
- Puccinia galii-elliptici** Maire. Powdery brown rust pustules on leaves of *G. ellipticum* in Corsica.
- Puccinia galii-silvatici** Wurtl. Leaf rust on *G. sylvaticum* in Switzerland.
- Puccinia lagerheimii** Lindr. Dark-brown rust pustules on leaves and stems of *G. sylvestre* in Switzerland.
- Puccinia pallidifaciens** Lindr. Dark-brown rust pustules on leaves and stems of *G. boreale* in Russia.
- Septoria cruciatae** Rob. and Desm. Dull-brown leaf spots with brown borders on *G. boreale*, *G. chersonense*, *G. cruciata*, *G. pedemontanum*, and *G. rotundifolium* in Europe.
- Septoria galii-borealis** P. Henn. (*S. galii-borealis* Bub. and Kab.) On leaves of *G. boreale* in Japan and Bohemia.
- Septoria relicta** Bub. On leaves of *G. relicta* in Bohemia.
- Septoria urens** Pass. On leaves of *G. tricornis* in Italy.
- Synchytrium globosum** Schroet. See *Potentilla*.
- Thekopsora guttata** (Schroet.) Syd. See *Asperula*.
- Uromyces galii** Diet. Rust on leaves and stems of *G. aparine* in Japan.
- GALPHIMIA.** Shrubs.
- Uromyces galphimiae** Diet. and Holw. Powdery cinnamon-brown to black rust pustules on *G. humboldtiana* in Mexico.
- GARCINIA.** MANGOSTEEN. GAMBOGE. Tropical fruit trees.
- Ceuthospora garciniae** Syd. On leaves of *G. mangostana* in the Philippines.
- Corticium koleroga** (Cke.) V. Hoeh. See *Coffea*.
- Gamboge disease.** A fruit disease due to an unknown cause is reported on *G. mangostana* in Ceylon.
- Gloeosporium garciniae** Koord. Irregular dull-brown leaf spots on *G. dulcis* in Java.
- Helminthosporium garciniae** Petch. On leaves of *G. mangostana* in Ceylon.
- Micropeltis garciniae** P. Henn. Black radiate perithecia on leaves of *Garcinia* sp. in Tanganyika.
- Niptera garciniae** P. Henn. On leaves of *Garcinia* sp. in Tanganyika.
- Pirostoma garciniae** P. Henn. On leaves of *Garcinia* sp. in Tanganyika.
- Zignoella garciniae** P. Henn. Forms cankers on woody portions in which appear superficial, black, carbonous perithecia. Attacks *G. mangostana* in Malaya and Tanganyika, death of infected plants often resulting.
- GARDENIA. WARNERIA.** Ag. CAPE JASMINE. Tropical shrubs or small trees cultivated for foliage and flowers.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Hemileia woodii** Kalchbr. and Cke. Yellow rust sori as broad patches on leaves of *G. edulis*, *G. jasminoides*, and *G. radicans* in Java, China, and Queensland.
- Phyllosticta gardeniae** Tassi. Subcircular to irregular pale-yellow leaf spots on *G. jasminoides* (*G. florida*) in Italy.
- Ramularia gardeniae** C. Mass. On leaves of *G. grandiflora* in Italy.
- Septoria gardeniae** Sacc. On leaves of *Gardenia* sp. in Italy.
- Uredo gardeniae-thunbergiae** P. Henn. Rust pustules on yellow effuse leaf spots on *G. thunbergia* in the Union of South Africa.
- GARRYA.** SILK-TASSEL BUSH. Ornamental shrubs.
- Ascochyta garryae** Sacc. Brown leaf spots with dark borders on *G. elliptica* in France.
- Phyllosticta garryae** Pass. Subcircular brown leaf spots on *G. elliptica* in France.
- Septoria garryae** Roum. Subcircular to irregular, often marginal, brown, then ashen-gray, leaf spots on *G. elliptica* in France.
- Septoria thuretii** Brun. Pale-brown leaf spots with dull-brown margins on *G. thuretii* in France.
- GARUGA.** Tropical fruit trees.
- Fomes lamaoensis** Murr. See *Hevea*.
- Kuehneola garugae** Syd. Powdery yellow rust pustules on leaves of *G. abilo* in the Philippines.
- GASTROLOBIUM.** Australian shrubs.
- Achorella gastrolobii** (P. Henn.) Theiss. and Syd. On *G. spinosum* in Australia.
- Aecidium stowardii** Har. Leaf rust on *G. calycinum* in Australia.
- Puccinia gastrolobii** Diet. Cinnamon-brown rust pustules on leaves of *G. calycinum* in Australia.
- Puccinia stowardii** Har. Rust on leaves and branches of *G. calycinum* in Australia.
- Uredo stowardii** Har. Leaf rust on *G. calycinum* in Australia.
- GAULTHERIA.** WINTERGREEN. SALAL. Evergreen, erect or procumbent plants cultivated for their leaves, flowers, and fruit.
- Pucciniastrum gaultheriae** Syd. Rust on leaves of *G. nummularioides* in India.
- GAYLUSSACIA.** HUCKLEBERRY. Shrubs grown for their fruit. See also *Vaccinium*.
- Cercospora gaylussaci** Speg. Angular spots on leaves of *Gaylussacia* sp. in Brazil.
- Exobasidium gaylussaciae** P. Henn. Irregular fleshy swellings occur on leaf blades and petioles and stems of *G. brasiliensis* in Brazil.
- Phaeochora densa** (B. and Br.) Theiss. and Syd. Irregular black stromata on leaves of *Gaylussacia* sp. in Brazil and Costa Rica.
- GEISSORRHIZA.** Half-hardy Ixialike bulbous plants.
- Uromyces geissorrhizae** P. Henn. Yellow rust pustules, becoming dark-brown, on both leaf surfaces of *G. rupestris*, *G. secunda*, and *Moraea ramosa* in the Union of South Africa.
- GEITONOPLESIIUM.** Woody twiners in Australia.
- Uredo geitonoplesii** McAlp. Brown rust sori on leaves of *G. cymosum* in Australia.
- GENIPA.** West Indian shrubs and trees, one species grown for its fruit.
- Phyllachora genipae** Arth. and Dal. Black stromata on leaves of *G. americana* in Porto Rico.
- GENISTA.** BROOM. Ornamental shrubs grown for their handsome flowers.
- Diaporthe inaequalis** (Curr.) Nke. On *Genista* sp. in Europe.
- Gloeosporium garganicum** Sacc. and D. Sacc. On branches of *G. michelii* in Italy.
- Septoria cytisi** Desm. See *Laburnum*.
- Septoria nitida** Ferr. On leaves of *G. germanica* in Italy.
- Uromyces genistae-tinctoriae** (Pass.) Fckl. See *Caragana*.
- Ustilago albida** Bub. Smut in anthers of *G. spathulata* in Yugoslavia.
- Venturia genista** (Fckl.) Wint. On branches of *G. germanica* in Austria and Hungary.
- GENTIANA.** GENTIAN. Herbs, cultivated for their blue flowers.
- Cronartium delawayi** Pat. Yellow to brown rust pustules on leaves of *G. picta* and *G. yunnanensis* in China.
- Cronartium gentianeum** Thuem. Brown rust pustules on leaves and stems of *G. asclepiadea* in Russia, Bulgaria, and Austria.
- Mycosphaerella gentianae** Niessl. On stems of *G. asclepiadea* in China and Austria. Reported from Colorado.

**GENTIANA**—Continued.

- Peronospora carniolica** Gäum. Downy mildew on leaves of *G. amarellum* in Austria.
- Phyllosticta gentianellae** Mass. Brown leaf spots with yellow margins on *G. asclepiadea* in Italy.
- Ramularia evanida** (Kuehn.) Sacc. Yellow-brown leaf spots on *G. asclepiadea* in Switzerland, Bulgaria, Yugoslavia, and Germany.
- Septoria gentianae** Thuem. Dull-brown, irregular leaf spots on *G. adscendens*, *G. macrantha*, and *G. scabra* in Japan, Siberia, and Russia.
- Septoria microspora** Speg. Large brown leaf spots on *G. asclepiadea* and *G. nivalis* in Italy and Japan.
- Septoria montana** Trav. Subcircular ochraceous leaf spots on *G. acaulis* in Italy.
- Septoria tosevi** Bub. Circular to oblong, often confluent, dull-yellow leaf spots on *G. cruciata* in Bulgaria and Austria. Reported from Colorado.
- GERANIUM.** CRANE'S BILL. Annual, biennial, or perennial herbs, some species cultivated.
- Aecidium infrequens** Barcl. Leaf rust on *G. nepalense* in Japan.
- Cercospora geranii-sanguinei** P. Henn. Dull-brown spots at tips of leaves of *G. sanguineum* in Norway.
- Cercospora magnusiana** Allesch. Ochraceous-olivaceous leaf spots on *G. sylvaticum* in Switzerland, Austria, and Germany.
- Coleosporium geranii** Pat. Brown rust pustules on leaves of *Geranium* sp. in China.
- Euryachora geranii** (Fr.) Schroet. Fungus of uncertain position on leaves of *G. sylvaticum* in Germany.
- Graphium geranii** Vogl. Circular to elliptical, often confluent, dull-brown leaf spots on *G. molle* in Italy.
- Peronospora conglomerata** Fckl. Downy mildew on leaves of *G. columbinum*, *G. dissectum*, *G. molle*, *G. phaeum*, *G. pusillum*, and *G. robertianum* in French North Africa and Europe.
- Phyllosticta trailii** Sacc. Brown leaf spots on *G. sylvaticum* in Norway.
- Plasmopara pusilla** (De By.) Schroet. Downy mildew on leaves of *G. palustre*, *G. phaeum*, *G. pratense*, *G. pusillum*, and *G. sylvaticum* in Europe.
- Puccinia callaquiensis** Neger. Rust pustules on leaves and stems of *G. berterianum* in Chile.
- Puccinia cuneata** Diet. Leaf rust on *Geranium* sp. in Japan.
- Puccinia escharoides** Syd. Powdery dark-brown rust pustules on leaf blades and petioles of *G. sinense* in central Africa.
- Puccinia geranii-pilosi** McAlp. Brown to black rust pustules on leaf blades and petioles of *G. pilosum* in Australia.
- Puccinia morthieri** Koern. Brown-black rust pustules on circular yellow leaf spots on *G. macrorrhizum*, *G. pyrenaicum*, *G. robertianum*, and *G. sylvaticum* in Europe and Siberia.
- Puccinia saniniensis** P. Magn. Powdery dark-brown rust pustules on leaves of *G. crenophilum* in Syria.
- Ramularia dolomitica** Kab. and Bub. On leaves of *G. phaeum* in Finland.
- Ramularia geranii** (West.) Fckl. Subcircular dull-brown leaf spots on *Erodium cicutarium* and *Geranium* spp. in Europe and Siberia. Reported from Louisiana.
- Ramularia geranii-phaei** (Mass.) Magn. On leaves of *G. phaeum* in Austria and Bohemia.
- Ramularia geranii-sanguinei** Massal. Rufous leaf spots on *G. sanguineum* in Italy.
- Ramularia geranii-silvatici** Vesterg. Angular-brown leaf spots on *G. sanguineum* and *G. sylvaticum* in Sweden and Esthonia.
- Septoria daniloi** Bub. On leaves of *G. columbinum*, *G. lucidum*, and *G. pallens* in Spain, Yugoslavia, and Russia.
- Septoria expansa** Niessl. Indefinite areas on leaves of *G. dissectum*, *G. palustre*, and *G. sanguineum* in France, Russia, and Bohemia.
- Septoria geranii** Rob. and Desm. Irregular olivaceous-brown leaf spots with purple margins on *G. molle*, *G. pusillum*, *G. robertianum*, *G. sanguineum*, *G. sylvaticum*, and *Pelargonium* sp. in Spain, France, Italy, Tunis, the Union of South Africa, and Germany.
- Septoria geranii-nodosi** Massal. Brown leaf spots on *G. nodosum* in Italy.
- Uromyces geranii** (DC.) Otth. and Wartm. Brown rust pustules on leaf blades and petioles of *Geranium* spp. (about 30 species) in Colombia, Europe, Japan, and the Union of South Africa. Several doubtful American references.
- Uromyces kabatianus** Bub. Yellow to brown rust pustules on circular yellow leaf spots on *G. molle* and *G. pyrenaicum* in Denmark, Sweden, Esthonia, Austria, and Germany.
- Uromyces scariosus** Berk. Brown leaf rust on *G. dissectum* and *G. potentilloides* in New Zealand.
- Uromyces trunculatus** Trott. Yellow-brown rust pustules on leaves of *G. striatum* in Italy.
- Venturia circinans** (Fr.) Sacc. On leaves of *G. molle*, *G. rotundifolium*, and *Geranium* sp. in Australia, Italy, France, and Germany.
- GERARDIA.** See *Stenandrium*.
- GERBERA. PERDICUM** Ag. BARBERTON DAISY. Perennial herbs grown for their yellow or pink flower heads.
- Aecidium crypticum** Kalch. and Cke. Leaf rust on *G. lanuginosa* in India and the Union of South Africa.
- Ascochyta gerberae** Maffei. Circular spots, enlarging to occupy entire area of leaves of *G. jamesoni* in Italy.
- Septoria gerberae** Syd. Large purplish-black blotches on leaves of *G. jamesonii* and other species in the Union of South Africa. The leaves are often destroyed.
- GESNERIA.** Shrubs grown for their showy tubular flowers.
- Cocconia gesneraceae** P. Henn. Circular black leaf spots on *Gesneria* sp. in Brazil.
- Puccinia gesneriacearum** Diet. Leaf rust on *Gesneria* sp. in Brazil.
- Rostronitschkia nervincola** (Rehm.) Fitzp. Black superficial rough fruiting bodies on yellow leaf spots on *G. albiflora* in Porto Rico.
- GEUM.** AVENS. Herbs, sometimes cultivated for their yellow flowers or long plummy fruit.
- Cercospora gei** Fckl. On leaves of *G. rivale* in Austria.
- Depazea geicola** Fr. On leaves of *G. strictum* in Siberia.
- Gloeosporium gei** Trail. On stems of *G. urbanum* in Scotland.
- Phragmidium circumvallatum** P. Magn. Yellow rust sori on leaves of *G. heterocarpum* and *G. kokanicum* in Asia Minor, Turkestan, Algeria, Spain, and Russia.
- Phyllosticta gei** Thuem. On leaves of *G. urbanum* and *Geum* sp. in Siberia, Bohemia, Austria, and Germany.
- Puccinia gei** McAlp. Brown rust pustules on yellow-green sunken leaf spots on *G. renifolium* in Tasmania.
- Ramularia gei** (Elias.) V. Hoeh. Subangular ochraceous to gray-brown spots on leaves of *G. rivale* and *G. urbanum* in Europe. Reported from Missouri.
- Ramularia trotteriana** Sacc. Small subcircular gray leaf spots with dark purple margins on *G. montanum* and *G. urbanum* in Italy.

- GIGANTOCHLOA.** See Bambuseae.
- GILIA.** Annual, biennial, and perennial herbs.  
**Phytophthora parasitica** Dastur. See Ricinus.  
**Puccinia patagonica** Speg. Rust pustules on leaves, stems, and flowers of *G. gracilis* in Argentina and Chile.  
**Urocystis giliae** Speg. Smut on roots of *G. foetida* in Argentina.
- GINKGO.** MAIDENHAIR TREE. Ornamental trees.  
**Phyllosticta ginkgo** Brun. On leaves of *G. biloba* in France and Russia. Reported from the District of Columbia.  
**Phyllosticta salisburiae** Tassi. Large irregular brown leaf spots on *G. biloba* in Italy.
- GLADIOLUS.** Summer and autumn flowering cormous plants.  
**Ascochyta gladioli** Trav. and Spess. Dead areas at base of stems of *Gladiolus* sp. in Portugal.  
**Bacterium gladioli** Sever. (*Pseudomonas gladioli* Sever.) This bacterium is reported from Italy and Holland as the cause of a soft rot of *Gladiolus* corms. The disease is first noted as a dying of the tops, beginning with the leaf tips, but soon involving the entire leaf which droops and finally falls over. *G. colvillei* is given as the host.  
**Mycosphaerella fusca** Pass. On *Gladiolus* sp. in Italy.  
**Puccinia gladioli** Cast. Produces brown, linear spots, limited by the veins, on both surfaces of leaves in which brown or black rust pustules appear. The host species are *G. aleppicus*, *G. byzantinus*, *G. communis*, *G. eckloni*, *G. illyricus*, *G. reuteri*, and *G. segetum*. The disease occurs in Italy, France, Spain, Portugal, Dalmatia, Algeria, and Asia Minor.  
**Uredo gladioli-buettneri** Bub. Brown rust pustules on leaves of *G. buettneri* in Togo.  
**Uredo pulchra** Syd. Rust pustules on leaves of *Gladiolus* sp. in the Himalaya Mountains of India.  
**Urocystis gladioli** (Req.) Sm. This smut fungus attacks and destroys the corms of *Gladiolus* (*G. communis*, *G. imbricatus*, and *G. segetum*), a black spore mass forming within. It occurs in England, France, Holland, and Germany.  
**Uromyces gladioli** P. Henn. Yellow, finally brown, rust pustules on both surfaces of leaves of *G. angustus*, *G. blandus*, *G. maculatus*, *G. orchidiflorus*, and *G. quartianus* in the Union of South Africa.  
**Uromyces nyikensis** Syd. Dark-brown to black rust pustules are produced on both surfaces of leaves of *G. nyikensis* on the Nyika Plateau, Africa. The rust greatly disfigures the leaves.  
**Uromyces transversalis** Thuem. Round to elongate dead spots on both surfaces of the leaves in which appear brown to deep-brown or black rust pustules. The hosts are *G. psittacinus*, *G. saundersii*, *Tritonia* (*Montbretia*) *lineata*, *T. securigera*, and *T. squalida* in the Union of South Africa.
- GLAUCIUM.** HORN POPPY. Perennial herbs with large poppylike flowers and glaucous blue foliage.  
**Entyloma fuscum** Schroet. Smut sori in gray-white, then brown, leaf spots on *G. flavum*, *G. luteum*, *Papaver somniferum*, and *Rhoeas* sp. in France, Hungary, and Germany. Also known from Maine and Iowa.  
**Entyloma glaucii** Dang. Smut sori in subcircular leaf spots on *G. flavum* and *G. luteum* in France, Denmark, and Germany.
- GLEDITSIA.** HONEY LOCUST. Trees grown for their finely divided foliage and fragrant flowers.  
**Phyllosticta triacanthi** Sacc. Irregular buff leaf spots with narrow brown margins on *G. triacanthos* in Italy.
- GLIRICIDIA.** Small, ornamental, leguminous trees.  
**Cercospora gliricidiae** Syd. On leaves of *G. sepium* in the Philippines.  
**Colletotrichum gliricidiae** Syd. Anthracnose on *G. sepium* in the Philippines.  
**Corticium salmonicolor** B. and Br. See Citrus.
- GLOBULARIA.** GLOBE DAISY. Herbs and shrubs with small blue flowers in heads.  
**Phyllosticta globulariae** West. Subcircular dull-brown leaf spots on *G. vulgaris* in Belgium.  
**Puccinia grisea** (Strauss.) Wint. Rust sori in sunken brown leaf spots on *G. cordifolia*, *G. nudicaulis*, *G. vulgaris*, and *G. willkommii* in Europe.  
**Septoria globulariae** Sacc. Subcircular gray-white leaf spots on *G. nudicaulis* and *G. vulgaris* in Italy.
- GLOXINIA.** Stemless cormous plants with showy bell-like flowers.  
**Aphelenchus olesistus** Ritz. Bos. See Begonia.
- GLYCERIA.** PANICULARIA Ag. Perennial grasses, sometimes grown for ornament.  
**Entyloma spectabile** Karst. Smut sori in elongate yellow to brown leaf spots on *G. spectabilis* in Finland.  
**Fusicladium hariotianum** Sacc. On leaves of *G. borrieri* in France.  
**Mycosphaerella recutita** (Fr.) Johans. See Aira.  
**Physoderma gerhardti** Schroet. Brown areas on leaves and sheaths of *G. aquatica*, *G. fluitans*, *Alopecurus pratensis*, and *Phalaris arundinacea* in Germany.  
**Puccinia glyceriae** Ito. Powdery golden to brown rust pustules on leaves of *G. aquatica* and *G. tonglensis* in Japan.  
**Ustilago grammica** B. and Br. See Aira.
- GLYCINE.** See Soja.
- GLYCYRRHIZA.** LICORICE. Perennial herbs with fleshy roots.  
**Cercospora cavarae** Sacc. and D. Sacc. Brown indefinite leaf spots on *G. glabra* in Italy.  
**Phyllosticta glycyrrhizae** Brun. Brown leaf spots on *G. glabra* in France.  
**Uromyces ellipticus** Diet. and Neg. Brown rust pustules on leaves and stems of *G. astragalina* in Argentina and Chile.
- GOETHEA.** Evergreen shrubs.  
**Phyllosticta goetheae** Magn. Irregular gray-white leaf spots with dull-brown borders on *G. cauliflora* (*G. strichiflora*) in Italy.
- GOMESA.** See Orchidaceae.
- GOMPHOCARPUS.** Perennial herbs or subshrubs.  
**Puccinia schlechteri** P. Henn. Leaf rust on *G. schinzianus* in the Union of South Africa.  
**Septoria gomphocarpi** P. Henn. Circular brown leaf spots on *G. fruticosus* in Italy and Australia.
- GOMPHOLOBIUM.** Australian shrubs.  
**Cronartium jacksoniae** P. Henn. See Jacksonia.
- GOMPHRENA.** GLOBE AMARANTH. Erect or prostrate herbs.  
**Cercospora gomphrenicola** Speg. Indefinite pale-yellow leaf spots on *G. glauca* in Argentina.  
**Phyllosticta gomphrenae** Sacc. and Speg. Large indefinite yellow leaf spots with dull-brown margins on *G. globosa* in Italy.  
**Puccinia stuckerti** Speg. Black rust pustules on leaves of *Gomphrena* sp. in Argentina.  
**Septoria gomphrenae** Sacc. and D. Sacc. Subcircular to angular gray-white leaf spots with reddish margins on *G. globosa* in Italy.  
**Uromyces bonariensis** Speg. Leaf rust on *G. elegans* and *G. gracilis* in Argentina.
- GONGORA.** See Orchidaceae.

- GONOLOBUS. VINCETOXICUM** Ag. Trailing or climbing herbs or shrubs.  
**Septoria gonolobicola** P. Henn. Circular white leaf spots on *G. stephanotrichus* in Germany.
- GOODENIA.** Australian herbs and shrubs.  
**Puccinia saccardoi** Ludw. Golden to black rust pustules on leaves of *G. albiflora*, *G. amplexans*, *G. geniculata*, *G. glauca*, *G. hederacea*, *G. ovata*, and *G. pinnatifida* in Australia.  
**Synchytrium succisae** De B. and Wor. On leaf blades and petioles of *Goodenia* sp. and *Succisa pratensis* in Australia, Russia, Scandinavia, Bohemia, and Germany.
- GOODIA.** Australian shrubs with pea-like flowers.  
**Aecidium soleniiforme** Berk. Leaf rust on *G. latifolia* in Australia.
- GOODYERA.** See Orchidaceae.
- GOSSYPIUM.** COTTON. Herbs and shrubs.  
**Alternaria macrospora** Zimm. Leaf spot on *Gossypium* sp. in India.  
**Ascochyta gossypii** Syd. On *Gossypium* sp. in Russia. Also reported from Arkansas.  
**Bacterial.** A bacterial disease of bolls of *G. hirsutum*, said to be distinct from that caused by *B. malvacearum*, is reported from India. The disease is insect carried, black shiny spots appearing on the surface of the bolls where the punctures are made. Immature bolls when infected turn black and are shed. In all cases infected bolls, seed, and lint are discolored, and often a slimy soft rot is present.  
**Cercospora gossypii** Syd. On *Gossypium* sp. in Uganda and Brazil.  
**Club leaf (Cyrtois).** This disease, attacking cultivated *Gossypium* in China, is apparently one of the group of virus diseases. It produces a witches'-broom effect due to a shortening of the nodes and development of additional shoots. The leaves are small, crinkled, and often colored different shades of green, yellow, and red.  
**Doassansia gossypii** Lagerh. Leaf smut of *Gossypium* sp. in Ecuador.  
**Fomes lamaoensis** Murr. See Hevea.  
**Giberella gossypina** Averna. Causes a boll rot of *Gossypium* sp. in Brazil.  
**Internal boll disease.** The disease is characterized by a brown staining of the lint in green, unopened bolls and is due to the growth of a number of species of fungi which are transmitted by sucking insects. Young bolls when infected drop prematurely, and in older bolls the lint is more or less completely rotted. The disease is serious in the British West Indies and Trinidad. Similar diseases occur in Egypt and other cotton-growing countries, including probably the United States.  
**Kuehneola gossypii** (Lagerh.) Arth. (*K. desmium* B. and Br.) Powdery cinnamon-brown rust pustules on purplish or brown leaf spots on *G. acuminatum*, *G. arboreum*, *G. brasiliense*, *G. herbaceum*, *G. hirsutum*, *G. mexicanum*, *G. microcarpum*, and *G. peruvianum* in Cuba, Porto Rico, South America, the Philippines, India, Java, Ceylon, India, Indo-China, New Guinea, west Africa, and Trinidad. Also known from Florida.  
**Oulariopsis gossypii** Wakef. White powdery mildew on yellow or red irregular patches on leaves of *G. barbadense* in Barbados.  
**Peronospora gossypina** Averna. Downy mildew causing rot of bolls of *Gossypium* sp. in Brazil.  
**Pestalozzia gossypii** Hori. Ochraceous to brown blotches on leaves of *G. herbaceum* in Japan.  
**Phoma roumii** Fron. On stems and lateral branches of *Gossypium* sp. in Dahomey and west Africa, causing the leaves to wither.  
**Phyllosticta malkoffii** Bub. Large brown to gray-white leaf spots with brown margins on *G. herbaceum* in Bulgaria.  
**Phytophthora** sp. Causes a soft rot of bolls of *Gossypium* (cult.) in the British West Indies. Several species are involved, including *P. faberi*.  
**Rhizoctonia** sp. See Vigna.  
**Sporidesmium longipedicellatum** Reich. Causes a leaf spot of *Gossypium* (cult.) in Egypt.  
**Stilbum nanum** Mass. f. **gossypina** Averna. Causes a boll rot of *Gossypium* sp. in Brazil.
- GOUANIA.** Tropical shrubs.  
**Puccinia gouaniae** Holw. Brown leaf rust on *G. lupuloides* and *G. polygama* in Porto Rico, Cuba, Costa Rica, Trinidad, Panama, Guatemala, Santo Domingo, and Uganda.  
**Puccinia invaginata** Arth. and Johnst. Leaf rust on *G. lupuloides* and *G. polygama* in Porto Rico, Cuba, Guatemala, St. Croix, Trinidad, and South America.  
**Puccinia paraënsis** Diet. Cinnamon-brown rust pustules on leaves of *G. pyrifolia* in Brazil.  
**Uredo gouaniae** Ell. and Kels. Leaf rust on *G. lupuloides* and *G. polygama* in St. Croix and Porto Rico.  
**Uromyces gouaniae** Kern. Brown leaf rust on *G. lupuloides* in Guatemala.
- GRABOWSKIA.** Spiny tropical shrubs.  
**Puccinia paradoxopoda** Speg. Dark-brown rust pustules on leaves of *G. obtusa* in Argentina.  
**Puccinia penningtonii** Syd. Black leaf rust on *G. duplicata* in Argentina.
- GRAPTOPHYLLUM.** Tropical shrubs with variegated foliage.  
**Rosellinia bunodes** B. and Br. See Citrus.
- GRATIOLA.** Low herbs.  
**Phyllosticta gratiolae** Höll. Ochraceous leaf spots on *G. officinalis* in Hungary and Russia.  
**Septoria gratiolae** Sacc. and Speg. Dull-brown leaf spots on *G. officinalis* in Italy.
- GREVILLEA.** SILK OAK. Timber and ornamental trees.  
**Dimerium orbiculatum** McAlp. Numerous circular, then confluent, brown to black leaf spots on *G. victori* in Australia.  
**Diplodia cacaoicola** P. Henn. See Theobroma.  
**Fomes lamaoensis** Murr. See Hevea.  
**Fusarium udum** Butl. See Dianthus.  
**Phyllachora grevilleae** (Lév.) Sacc. Circular black stromata on leaves of *G. burifolia* and *G. occidentalis* in Australia.  
**Rhizoctonia lamellifera** Small. Infected trees show a browning and wilting of the foliage followed by a stag-headed condition and ultimate death. Blackened areas occur on the lower portions of the trunks from which there are resinous or gummy exudates. Diseased tissues are brown and filled with gum. The roots rot and may be partly covered with thin superficial fungus layers. Numerous small black sclerotia occur in the rotted wood, cortex, and bark, together with sheets of black, irregular fungus tissue. The disease attacks *G. robusta* in Uganda as well as *Casuarina equisetifolia*, *Bixa orellana*, *Coffea robusta*, and *Thea* to a lesser extent.  
**Rosellinia arcuata** Petch. See Thea.
- GREWIA.** Shrubs.  
**Aecidium warneckeanum** P. Henn. Galls up to 20 centimeters long deforming branches of *G. carpinifolia* in Tanganyika.  
**Helminthosporium grewiae** P. Henn. On leaves of *Grewia* sp. in the Congo.  
**Ravenelia atrides** Syd. Yellow to black rust pustules on leaves of *G. caffra* and *G. occidentalis* in the Union of South Africa.  
**Rhytisma grewiae** Kalchbr. Black appanate stromata on leaves of *G. occidentalis* in the Union of South Africa.

**GREWIA**—Continued.

**Stigmatea grewiae** P. Henn. Circular brown leaf spots on *G. caffra* in the Union of South Africa.

**Uredo grewiae** Pat. and Har. Leaf rust on *G. ferruginea* in Senegal.

**Ustilago grewiae** (Pass.) P. Henn. Smut sori in branches of *G. columnaris*, *G. microcarpa*, and *G. mollis* in Abyssinia, tropical Africa, and Ceylon.

**GRIFFITHIA**. Rubiaceous shrubs.

**Endophyllum griffithiae** (P. Henn.) Racib. Leaf rust on *G. fragrans*, *G. latifolia*, and *Randia scandens* in Java.

**GRISELINIA**. Shrubs with large, glossy laurel-like foliage.

**Puccinia griseliniae** Pазschke. Dark-brown rust pustules on small circular sunken leaf spots on *G. ruscifolia* in Brazil.

**GROSSULARIA**. GOOSEBERRY. See *Ribes*.**GUAZUMA**. Tropical trees.

**Trabutia guazumae** Chardon. (*Phyllachora guazumae* P. Henn.) Black circular stromata on leaves of *G. ulmifolia* in Brazil and Porto Rico.

**GUETTARDA**. Tropical and subtropical shrubs.

**Septoria guettardae** Garman. Large irregular dark-red leaf spots with white centers on *G. ovalifolia* in Porto Rico.

**GUNNERA**. Perennial herbs with large leaves used for ornamental effects.

**Ovularia gunnerae** Speg. Circular to angular purplish leaf spots on *G. chilensis* in Chile.

**Phyllosticta gunnerae** Speg. On leaves of *G. magellanica* in Patagonia.

**GUZMANIA**. Tropical American bromeliads.

**Lembosia bromellacearum** Rehm. On leaves of *G. roezlii* in Brazil.

**GYMNADENIA**. See *Orchidaceae*.**GYMNOLOMIA**. Yellow-flowered herbs or shrubs.

**Accidium gymnomiae** Mayor. Leaf rust on *G. quitensis* in Colombia.

**Puccinia gymnomiae** Arth. Powdery brown leaf rust on *G. brachypoda*, *G. patens*, *G. subflexuosa*, and *Hymenostephium microcephalum* in Mexico, Guatemala, and Costa Rica.

**Puccinia semota** Jack. and Holw. Dark-brown leaf rust on *G. subflexuosa* in Guatemala.

**GYMNOSPORA**. Shrubs and small trees, often spiny.

**Englerulaster gymnosporiae** (P. Henn.) Th. On leaves of *Gymnosporia* sp. in the Union of South Africa.

**Septoria gymnosporiae** Syd. Small dark-brown leaf spots on *G. deflexa* in the Union of South Africa.

**GYMNOSTACHYUM**. White-flowered herbs from Ceylon.

**Puccinia leochroma** Syd. Leaf rust on *G. subcordatum* in the Philippines.

**GYNERIUM**. PAMPAS GRASS. Perennial grass.

**Ustilago quitensis** Lagh. Powdery dark olivaceous smut sori destroying panicles of *Cortaderia argentea* (*G. argenteum*) in Ecuador.

**GYNURA**. Tropical herbs with showy foliage.

**Accidium gynurae** Petch. Leaf rust on *G. lycopersicifolia* in Ceylon.

**Uredo gynurae** Petch. Leaf rust on *G. lycopersicifolia* in Ceylon.

**GYP SOPHILA**. BABY'S BREATH. Herbs with a profusion of small flowers used in borders.

**Mycosphaerella gypsophilae** Fckl. On leaves and stems of *G. gmelini*, *G. muralis*, and *G. scleranthus* in Siberia and Germany.

**Septoria lychnidis** Lasch. See *Lychnis*.

**Uromyces gypsophilae** Cke. Black powdery rust pustules on leaves and stems of *G. anatolica*, *G. haussknechtii*, *G. paniculata*, *G. polyclada*, *G. pulchra*, and *Saponaria vaccaria* in Persia, Kurdistan, and Russia.

**HABENARIA**. Terrestrial leafy orchids. See *Orchidaceae*.**HABRANTHUS**. Showy bulbous plants. Generally included with *Hippeastrum*.

**Puccinia habranthi** Diet. and Neg. Rust sori, round to irregular, yellow, becoming dark-brown or black in circular spots with purple margins. On both surfaces of leaves of *H. andicola* in Chile.

**Puccinia reichei** Diet. and Neg. Yellow to brown rust pustules on both surfaces of leaves of *H. chilensis* in Chile.

**HAEMODORUM**. Perennial Australian herbs.

**Puccinia haemodori** P. Henn. Black rust pustules on leaves of *Haemodorum* sp. in Australia.

**HAEMANTHUS**. BLOOD LILY. African bulbous plants.

**Cercospora haemanthi** Kalch. Large, elliptical brown spots on leaves of *H. puniceus* in the Union of South Africa.

**HAKEA**. Australian ornamental evergreen shrubs.

**Phyllachora hakeae** P. Henn. Black stromata on both leaf surfaces of *H. myrtoides* in Australia.

**Uredo angiosperma** Thuem. Powdery brown rust pustules on leaves of *Hakea* sp. in Australia.

**HALLERIA**. Shrubs or small trees.

**Septoria halleriae** P. Henn. Pale, effuse, zoned leaf spots on *H. lucida* in Italy and Germany.

**Septoria hallericola** Sacc. and D. Sacc. On leaves of *H. lucida* in Portugal.

**HAMMAMELIS**. WITCH-HAZEL. Ornamental woody plants with yellow flowers late in fall or in the winter.

**Accidium hammamelidis** P. Magn. Rust pustules on purplish leaf spots on *H. japonica* in Japan.

**HAMELIA**. Tropical ornamental woody plants grown for their flowers and berries.

**Uredo hameliae** Arth. Numerous light-brown powdery rust pustules on lower leaf surfaces of *H. erecta* in Porto Rico and Costa Rica.

**HARDENBERGIA**. Vines grown for their handsome flowers. See also *Kennedia*.

**Phyllosticta hardenbergiae** Cke. and Mass. Irregular dull-brown leaf spots on *Hardenbergia* sp. in Australia.

**Septoria hardenbergiae** Sacc. Pale leaf spots with dull-brown margins on *Kennedia monophylla* (*H. monophylla*) in Australia.

**Uromyces hardenbergiae** McAlp. Brown rust pustules on lower leaf surfaces of *H. monophylla* in Australia.

**HEDERA**. IVY. Ornamental woody vines, grown for their handsome persisting foliage.

**Ascochyta diploidina** Berl. and Bres. Pale, irregular, confluent leaf spots on *H. colchica* and *H. helix* in Russia.

**Bacillus** sp. A bacterial disease on *H. helix* in France, producing brown patches on the leaves and killing-back the young growth.

**Bacterium hederiae** Arn. Transparent leaf spots on *H. helix* in France.

**Colletotrichum hedericola** Laubert. Anthracnose on leaves of *H. helix* in Austria.

**Leptosphaeria hedericola** (Desm.) Sacc. On leaves of *H. helix* in France.

**Leptosphaeria hellicola** (Desm.) Niessl. On leaves of *Hedera* sp. in France.

**Macrophoma cylindrospora** (Desm.) Berl. and Vogl. On leaf blades and petioles of *Euonymus japonicus*, *Hedera helix*, *Quercus ilcx*, and *Vinca* sp. in France, Belgium, Italy, Austria, and Germany.

**HEDERA**—Continued.

- Macrophoma ulcinjensis** Bub. Large yellow to dull-gray leaf spots on *H. helix* in Yugoslavia.
- Meliola dichotoma** B. and C. Black superficial fungus patches on leaves of *Hedera* sp. in Japan.
- Meliola kusanoi** P. Henn. Circular black fungus patches on leaves of *Hedera* sp. in Japan.
- Mycosphaerella hederæ-helicis** Siem. On leaves of *H. helix* in Russia.
- Mycosphaerella hedericola** (Desm.) Lind. Subcircular to irregular brown leaf spots on *H. colchica* and *H. helix* in Italy, Yugoslavia, Russia, Great Britain, Denmark, and France.
- Phyllosticta belgradensis** Bub. and Ranaj. Irregular, often marginal, gray to ashen leaf spots with brown margins on *H. helix* in Yugoslavia and Spain.
- Phyllosticta destructiva-hederæ** Desm. On leaves of *Hedera* sp. in France.
- Phyllosticta hederæ** (Arc.) Allesch. On leaves of *H. helix* in Germany.
- Phyllosticta hederæ** Sacc. and Roum. On leaves of *H. helix* in Europe.
- Phyllosticta hedericola** Dur. and Mont. Subcircular whitish leaf spots with brown margins on leaves of *H. colchica* and *H. helix* in Algeria and Europe. Reported from New Jersey.
- Phyllosticta milenæ** Bub. Large irregular gray to dull-yellow leaf spots on *H. helix* in Bulgaria and Yugoslavia.
- Phyllosticta ulcinjensis** Bub. Large yellowish-white to greenish leaf spots with narrow brown margins on *H. helix* in Yugoslavia.
- Septoria desmazierii** Sacc. On leaves of *H. helix* in France.
- Septoria hederæ** Desm. On leaves of *H. colchica* and *H. helix* in Europe.
- Septoria insularis** B. A. Br. Large brown leaf spots on *H. helix* in Germany.
- Septoria mortolensis** Penz. and Sacc. See *Acacia*.
- Triphragmium thwaitesii** Berk. and Br. See *Acanthopanax*.
- HEDYCHIUM**. GINGER LILY. Rhizomatous herbs allied to *Zingiber*.
- Taphrina maculans** Butl. See *Curcuma*.
- HEDYSARUM**. Herbaceous perennials.
- Anthostomella sullæ** Montem. On leaves of *H. coronarium* in Italy.
- Cercospora ariminensis** Cav. Circular to ovate or elliptical dull chestnut-brown leaf spots on *H. coronarium* in Italy and Brazil.
- Erysiphe taurica** Lév. See *Althaea*.
- Placosphaeria onobrychidis** Sacc. See *Lupinus*.
- Ustilago kusanoi** Syd. Smut on *H. esculentum* in Japan.
- HELIANTHEMUM**. SUN ROSE. Ornamental woody or herbaceous plants grown for their showy flowers.
- Cercospora cistinearum** Sacc. Subcircular gray leaf spots with red-purple margins on *H. vulgare* in Italy.
- Cercospora helianthemi** Cav. Small circular gray-brown leaf spots on *H. polifolium* in Italy.
- Erysiphe taurica** Lév. See *Althaea*.
- Peronospora alpestris** Gäum. Downy mildew on leaves of *H. alpestre* and *H. niloticum* in Algeria and Switzerland.
- Peronospora leptoclada** Sacc. Dirty-white areas on lower leaf surfaces of *H. guttatum* and *H. vulgare* in France, Italy, and Switzerland.
- Phyllosticta helianthemi** Roum. and var. **major** P. Brun. Subcircular white leaf spots on *H. vulgare* in France.
- Phyllosticta helianthemicola** Allesch. and var. **marginatum** Syd. Irregular dull-yellow to gray indefinite leaf spots on *H. vulgare* in Great Britain and Germany.
- Septoria chamaecisti** Vestergr. Brown, then dirty-white, leaf spots with dull-brown borders on *H. chamacustum* and *H. vulgare* in Sweden.
- Septoria helianthemi** Gz. Frag. On leaves of *H. montanum* in Spain.
- Uredo helianthemi** Rbh. Yellow to brown rust pustules on leaves of *Cistus* sp. and *Helianthemum guttatum* in Italy.
- HELIANTHUS**. SUNFLOWER. JERUSALEM ARTICHOKE. Hardy herbaceous annuals and perennials.
- Mycosphaerella compositarum** (Auersw.) Schroet. On leaves of *H. annuus* in Yugoslavia.
- Puccinia minuscula** Arth. Brown leaf rust on *H. hypargyreus* in Ecuador.
- HELICHRYSUM**. EVERLASTING. Shrubs or herbs, some species grown for their flowers.
- Entyoloma aschersonii** (Ule.) Wor. A smut producing swellings on the stems and roots and causing young shoots to die-back, of *Gnaphalium* sp. and *Helichrysum arenarium* in Germany.
- Puccinia kalchbrenneri** Det. Yellow to brown rust pustules on leaves of *H. globosum*, *H. nudatum*, *H. nudifolium*, and *H. undulatum* in Australia and south and central Africa.
- Puccinia mac-owani** Wint. Yellow to fuscous rust pustules on leaves of *H. petiolatum* and *Helichrysum* sp. in the Union of South Africa.
- Septoria helichrysi** Syd. Irregular, then confluent, brown leaf spots on *Helichrysum* sp. in the Union of South Africa.
- Uromyces helichrysi** Lagh. Dark-brown rust pustules in sunken purple spots on leaves and stems of *H. stoechas* in France.
- HELICONIA**. BIHAI Ag. A foliage plant allied to *Musa*.
- Pyrenobotrys heliconiae** (P. Henn.) Theiss. and Syd. Black stromata on brown to black irregular leaf spots on *Heliconia* sp. in Brazil.
- Puccinia heliconiae** (Diet.) Arth. Brown powdery rust pustules on leaves of *H. borinquena*, *H. latispatha*, *H. psittacorum*, and *Heliconia* sp. in Porto Rico, Martinique, Brazil, Panama, and Trinidad.
- HELIOPSIS**. Hardy herbaceous plants cultivated for their profuse yellow flowers.
- Aecidium heliopsidis** Mayor. Rust on leaves of *H. buphthalmoides* in Colombia.
- Puccinia bimbergi** Mayor. Rust on leaves of *H. buphthalmoides* in Colombia.
- HELIOTROPIMUM**. HELIOTROPE. Herbaceous perennials cultivated for their fragrant flowers.
- Aecidium caspicum** Jacz. Rust pustules on large circular yellow to purplish leaf spots on *H. europaeum* in Russia.
- Aecidium guatemalense** Kern. and Kellerm. Rust pustules on discolored areas on leaves of *H. indicum* in Guatemala.
- Cercospora heliotropii-bocconii** Scalia. Subcircular, then confluent and irregular, dirty-yellow to gray leaf spots on *H. bocconi* in Sicily.
- Cercospora taurica** Tranzsch. Large gray leaf spots on *H. europaeum* in Russia and Yugoslavia.
- Puccinia gilva** Arth. and Holw. Leaf rust on *H. physocalycium* in Guatemala.
- Puccinia heliotropii** Kern. and Kellerm. Powdery chestnut-brown rust pustules on leaves of *H. indicum* in Guatemala.
- Puccinia heliotropicola** Speg. Powdery dull-brown rust pustules on leaves of *H. campestris* in Argentina.
- Uromyces heliotropii** Sred. Cinnamon to chestnut-brown rust pustules on leaves of *H. europaeum* in Russia, France, and Bulgaria.

- HELLEBORUS.** HELLEBORE. Christmas rose. Hardy herbaceous perennials cultivated for their flowers and attractive leaves.
- Accidium hellebori** Ed. Fisch. Rust on leaves of *H. viridis* in Spain and Switzerland.
- Ascochyta hellebori** Sacc. Angular whitish leaf spots with dark borders on *H. viridis* in Italy.
- Coniothyrium hellebori** Cke. and Mass. Large irregular brown leaf spots without definite margins on *H. corsicus*, *H. foetidus*, *H. niger*, *H. olympicus*, and *H. viridis* in Great Britain, France, Corsica, Italy, Switzerland, Denmark, Albania, Yugoslavia, Austria, and Germany. Reported from Oregon.
- Peronospora pulveracea** Fekl. Gray mold on leaves of *H. foetidus*, *H. niger*, *H. odoratus*, and *H. viridis* in Yugoslavia, France, Switzerland, Denmark, Austria, and Germany.
- Phyllosticta atro-zonata** Voss. Circular ashen-colored leaf spots with black centers on *H. altifolius* and *H. viridis* in Austria.
- Phyllosticta helleboriana** Brun. Small circular gray leaf spots with brown margins on *H. foetidus* in France.
- Phyllosticta helleborella** Sacc. (*Mycosphaerella hermione* Sacc.) Large, black, then lighter, angular to irregular blotches without distinct margins on upper surfaces of leaves of *H. foetidus*, *H. niger*, and *H. viridis* in Great Britain, Italy, and Austria.
- Phyllosticta helleboricola** C. Mass. Pale angular leaf spots on *H. odoratus* and *H. viridis* in Spain, Italy, and Bulgaria.
- Phyllostictiella olympica** (Allesch.) Tass. On leaves of *H. olympicus* in Germany.
- Ramularia hellebori** Fekl. Small circular whitish spots with black or purple margins on bracts and floral organs of *H. foetidus* and *H. viridis* throughout Europe.
- Ramularia recognita** Massal. On leaves of *H. multifidus* and *H. viridis* in Spain and Austria.
- Ramularia nigricans** (Mass.) Ferr. On leaves of *H. foetidus* in Spain.
- HELYGIA.** See *Parsonsia*.
- HEMEROCALLIS.** DAY LILY. Yellow and orange flowered, stout-rooted perennials with basal leaves.
- Colletotrichum liliacearum** Fen. Anthracnose on leaves of *H. fulva* in Italy.
- Mycosphaerella hemerocallidis** (Pass.) Lind. On leaves of *H. fulva* in Yugoslavia.
- Puccinia hemerocallidis** Thuem. Pale-yellow rust pustules, becoming darker, on sunken yellow leaf spots on *H. aurantiaca*, *H. dumortieri*, *H. lilio-asphodelus*, (*H. flava*), *H. fulva*, and *H. middendorfi* in Siberia and Japan.
- Sporotrichum ochraceum** (Cda.) Sacc. On stems of *Hemerocallis* sp. and *Allium* sp. in Bohemia.
- HEMIGRAPHIS.** Diffuse or prostrate herbs grown for their foliage and flowers.
- Uromyces macintirianus** Barel. Black rust pustules on leaves of *H. latebrosa* in India.
- HEPATICA.** Stemless low perennials.
- Aphelenchus olesistus** Ritz. See *Begonia*.
- Ascochyta molleriana** Wint. On leaves of *H. nobilis* (*H. triloba*) in Hungary.
- Ascochyta vodakii** Bub. On leaves of *H. nobilis* in Bohemia.
- Phyllosticta hepaticae** Brun. Leaf spots on *H. triloba* in France.
- Septoria hepaticae** Desm. Brown to black circular to irregular, then confluent, leaf spots on *H. triloba* in Belgium, France, Denmark, Russia, Switzerland, Italy, Spain, Austria, and Germany.
- HERACLEUM.** COW PARSNIP. Perennial or biennial herbs.
- Ascochyta heraclei** Bres. Deep-brown, circular, often confluent, leaf spots on *H. sibiricum* and *H. spondylium* in Bulgaria and Germany.
- Mycosphaerella caulicola** (Karst.) Lind. On stems of *Chamaenerium angustifolium*, *Heracleum spondylium*, and *Telekia speciosa* in Yugoslavia, Denmark, and Finland.
- Protomyces macrosporus** Unger. See *Coriandrum*.
- Puccinia cortezi** Ranaj. Leaf rust on *H. minimum* in Europe.
- Puccinia heraclei** Grev. Chestnut-brown to deep-brown rust pustules on leaf blades and petioles of *H. minimum*, *H. sibiricum*, and *H. spondylium* in Europe.
- Septoria heracleicola** Kab. and Bub. Small circular brown leaf spots, becoming whitish, on *H. sibiricum* in Italy and Austria.
- Septoria heraclei-palmati** R. Maire. Yellow, then brown, confluent leaf spots on *H. palmatum* in France.
- Taphridium umbelliferarum** (Rostr.) Lagerh. and Juel. Irregular gray leaf spots on leaves of *H. montanum*, *H. sibiricum*, *H. spondylium*, *Oreoselinum* sp., and *Peucedanum palustris* in Europe.
- HERMANNIA.** Shrubs.
- Septoria hermanniae** F. Tassi. Brown marginal leaf spots on *H. micans* in Italy.
- HERNIARIA.** BURSTWORT. Trailing small-leaved herbs.
- Peronospora herniariae** De By. Downy mildew on leaves and stems of *H. glabra* and *H. hirsuta* in Sweden, France, Russia, and Germany.
- Puccinia herniariae** Unger. Powdery red-brown to dark-brown rust pustules on leaves and stems of *H. glabra*, *H. hirsuta*, *H. lenticulata*, and *H. odorata* in Europe.
- Puccinia montagnei** Diet. Dark-brown rust pustules on leaves of *H. glabra*, *H. latifolia*, and *H. maritima* in France and Portugal.
- HERPESTIS.** Herbs, sometimes cultivated.
- Albugo quadrata** Kalch. and Cke. Small white sori (spore masses) on leaves of *H. verticillaris* in the Union of South Africa.
- HESPERETHUSA.** Naibel. Spiny trees or shrubs related to *Citrus*.
- Bacterium citri** Hasse. See *Citrus*.
- HESPERIS.** Dames rocket. Biennial herbs.
- Phyllosticta hesperidiarum** (Catt.) Pez. Circular to irregular brown, then dull-brown or ashen, leaf spots on *H. matronalis* in Italy.
- Ramularia matronalis** Sacc. On leaves of *H. matronalis* in France.
- HETEROPAPPUS.** Hardy herbaceous perennials with blue aster-like flowers.
- Uredo heteropappi** P. Henn. Brown rust pustules on circular dull-brown leaf spots on *H. hispidus* in Japan.
- HEUCHERA.** ALUMROOT. Herbaceous perennials.
- Aphelenchus olesistus** Ritz. See *Begonia*.
- Phyllosticta heucherae** P. Brun. Circular to irregular gray leaf spots on *H. glabra* and *H. sanguinea* in France and Great Britain.
- HEVEA.** PARA RUBBER TREE. Tropical trees furnishing rubber. This account of the diseases of *Hevea brasiliensis* is drawn for the most part from "The Diseases and Pests of the Rubber Tree," by T. Petch, of Ceylon.
- Aposphaeria ulei** P. Henn. Yellow to dark-brown, circular or confluent, leaf spots on *Hevea* sp. in Peru.
- Ascochyta heveae** Petch. Marginal white or brownish-white leaf spots with red-brown borders on *H. brasiliensis* in Ceylon.
- Catacauma huberi** (P. Henn.) Theiss. and Syd. Tar-spot disease, characterized by shining black crusts up to 1 centimeter in diameter on the lower leaf surfaces, with corresponding pale-green patches above. On *H. brasiliensis* in Brazil.

**HEVEA**—Continued.

**Cercospora heveae** Vinc. Leaf spot on *H. brasiliensis* in Brazil.

**Colletotrichum heveae** Petch. Irregular spots to large patches, dry, gray brown, with purple-brown margins on old leaves of *H. brasiliensis* in Ceylon and Brazil. Diseased areas often fall out, producing a shot-hole effect.

**Coniothyrium** sp. Produces a canker on *H. brasiliensis* in Uganda.

**Corticium koleroga** (Cke.) V. Hoehn. See Coffea.

**Corticium salmonicolor** B. and Br. See Citrus.

**Diplodia cacaoicola** P. Henn. See Theobroma.

**Dothidella ulei** P. Henn. (*Melanopsammopsis heveae* Stahel.) (*Fusicladium macrosporium* Kuyper.). This disease, the so-called South American leaf disease, is the most serious rubber disease in the Western Hemisphere. It has made rubber culture unprofitable in Surinam and has caused heavy losses in British Guiana, Trinidad, Brazil, and Peru. Other species of *Hevea* (*H. confusa* and *H. guyanensis*) are attacked, in addition to *H. brasiliensis*.

The disease attacks the young leaves within a few days after their unfolding, causing translucent olive-green to blackish-green spots, which may be so numerous that the leaves blacken and shrivel. Young trees are particularly susceptible, the disease destroying the tops. In some cases the diseased areas dry up and fall out, producing a shot-hole effect. This type of injury is most prevalent in old trees. Petioles and young green twigs are also attacked, swollen areas resulting, which later on crack and form cankers. The fungus produces a velvety mycelial layer on the diseased areas. Other stages develop on old leaves. No method of controlling this disease has yet been found.

**Fomes lamaoensis** Murr. (*Hymenochaete noxia* Berk.). The brown root disease is the most widely spread root disease of economic plants in the eastern tropics. The principal hosts are *Albizia stipulata*, *Annona muricata*, *Artocarpus communis*, *A. integrifolia*, *Brownea grandiceps*, *Brunfelsia Americana*, *Castilla elastica*, *Cedrela odorata*, *Cinnamomum camphora*, *C. cassia*, *C. iners*, *C. zeylanicum*, *Codiaeum variegatum*, *Coffea arabica*, *C. robusta*, *C. stenophylla*, *Cocos nucifera*, *Dalbergia latifolia*, *Eriobotrya japonica*, *Erythrina* sp., *Erythrozyllon coca*, *Ficus glomerata*, *F. infectoria*, *Funtumia elastica*, *Garuga pinnata*, *Gossypium* spp., *Grevillea robusta*, *Hevea brasiliensis*, *Hura crepitans*, *Jacaranda mimosae-folia*, *Manihot* spp., *Myristica magnifica*, *Persea gratissima*, *Poinciana regia*, *Soymida febrifuga*, *Spondias wightii*, *Terminalia bellerica*, *Thea* (cult.), *Theobroma cacao*, and *Thespesia populnea* in India, Ceylon, Java, Sumatra, Malaya, Samoa, New Guinea, east and west Africa, British Guiana, and possibly Brazil.

The disease makes slow progress, one tree dying at a time. The roots of diseased trees are incrustated with a mass of sand, earth, and stones, intermingled with and held together by brown mycelium. The mycelium later becomes black. The bark beneath is brown, or brown mottled with white patches. The wood may be soft and friable, with a network of fine brown lines composed of hyphae, or the rot may be comparatively hard and transversed by rather broad brown bands. In advanced stages the wood is honeycombed. The hard, purple-brown sporophore rarely develops. The minute velvety, sterile brown patches sometimes found ringing the stems of diseased plants at the crown have caused the fungus to be determined as *Hymenochaete noxia*, a conclusion shown by Petch to be erroneous.

**Fomes lignosus** Klotzsch. This fungus causes the most serious losses and is the most widely distributed of all the root diseases of rubber. It occurs in India, Ceylon, Java, Sumatra, Borneo, west Africa, the Congo, and Brazil, attacking in addition to *H. brasiliensis*, *Azelia palembanica*, *Artocarpus integrifolia*, *Berrya ammonilla*, *Bombax malabaricum*, *Cinnamomum camphora*, *Cocos nucifera*, *Coffea liberica*, *C. robusta*, *Dendrocalamus giganteus*, *Derris dalbergioides*, *Erythrina umbrosa*, *Ficus* spp., *Livistona cochinchinensis*, *Koompassia malaccensis*, *Manihot glaziovii*, *M. utilisissima*, *Oncosperma filamentosa*, *Shorea* sp., *Thea* (cult.), *Theobroma cacao*, and many other woody plants.

The fungus spreads from dead roots and old stumps to the living roots of the various hosts, either where they come in contact or by growing out through the soil. As a rule, smooth white, or yellowish-white to reddish rhizomorphs are produced on the exterior of diseased roots. These run longitudinally along the roots, uniting here and there to form a network. The younger growing parts of the fungus consist of a fan-shaped mycelium. The fungus penetrates both bark and wood, causing a soft friable rot, without distinguishing characteristics. Occasionally the fungus develops internally only, producing a soft watery decay without external rhizomorphs. The fruiting body is red brown above with a bright-yellow margin and bright-orange lower surface. The colors become duller with age.

**Fomes pseudo-ferreus** Wakef. (*Poria hypolateritia* Berk.). This fungus produces a wet rot of roots of *H. brasiliensis*, *Albizia* sp., *Symplocos spicata*, *Tephrosia* sp., *Thea* (cult.), and *Theobroma cacao* in Malaya, Ceylon, and Java. A dark-red superficial mycelium is sometimes formed over diseased roots, or occasionally snow-white patches of mycelium on small roots, but no rhizomorphs are produced. Workers in Java refer this species to *Fomes* (*Ganoderma*) *ferreus* Berk.

**Fusicladium** sp. A die-back disease, starting on pruned branches, has been reported from Java on *H. brasiliensis*, but is considered by Petch identical with the disease caused by *Diplodia*.

**Gloeosporium alborubrum** Petch. This fungus is one of the causes of abnormal leaf fall of *H. brasiliensis* in Ceylon, Sumatra, Malaya, and Uganda. On young leaves large, irregular, blackish-green, watery-looking patches occur. Older leaves turn yellow, then brown, black spots with gray centers appearing on veins and midribs. Petioles turn dark brown or black and the leaves fall. Young twigs may also be affected, dying back. Diseased fruit rot and finally become greenish gray and wrinkled.

**Gloeosporium heveae** Petch. This anthracnose fungus attacks young plants of *H. brasiliensis* in the nurseries of Ceylon and Brazil. The leaves turn yellow green, then yellow, and fall. The fruiting pustules appear on both leaf surfaces.

**Guignardia heveae** Syd. Tips and margins of leaves become pale brown with purple-brown margins. On *H. brasiliensis* in Ceylon, Malaya, and the Philippines.

**Helicobasidium mompa** Tan. See Morus.

**Helminthosporium heveae** Petch. The "bird's-eye spot" disease of *H. brasiliensis* attacks for the most part small plants in the nurseries, but is occasionally found on older trees. The spots are at first purple and very small, then increasing in size and becoming white and semitransparent, with narrow purple-brown borders. The disease occurs in Ceylon, India, Malaya, Java, Sumatra, the Philippines, and Gold Coast.

**Hymenochaete noxia** Berk. See *Fomes lamaoensis* Murr.

**Marasmius** spp. See Theobroma.

**Meliola heveae** Vinc. Small black superficial fungus patches on leaves of *H. brasiliensis* in Brazil.

**Mycosphaerella heveae** Petch. Small yellow leaf spots on *H. brasiliensis* in Ceylon, which become purple or purplish brown in color and confluent.

**Ophiobolus heveae** P. Henn. Circular to effuse and confluent gray leaf spots on *Hevea* sp. in Brazil.

**Phacophomopsis heveae** Grove. Causes a bark disease of *H. brasiliensis* in Uganda.

**HEVEA**—Continued.

**Phyllosticta heveae** Zimm. Marginal or apical brown patches on leaves of *H. brasiliensis* in Java, Trinidad, Brazil, and Malaya. Found on young trees in shaded nurseries.

**Phyllosticta ramicola** Petch. This fungus attacks young twigs from 6 to 18 inches back from the tip. Diseased areas are dark brown, progressing up and down from the point of infection. On *H. brasiliensis* in Ceylon, Java, and Malaya.

**Phytophthora faberi** Maubl. This fungus attacks practically all parts of the rubber tree, causing the diseases known as "abnormal leaf fall," pod or fruit disease, die-back of shoots, and "claret-colored canker" or "patch canker." These diseases are prevalent in India, Ceylon, Java, Malaya, Fiji, and other rubber-growing countries.

Diseased fruits are first watery green, then black and soft. They remain hanging on the tree. The fungus passes from the fruit to the leaves and twigs, causing an abnormal leaf fall and dying back of young twigs. The leaves may first turn yellow or mottled, or they may fall while still green. The leaflets often separate from the petioles. In the case of the trunk canker no external symptoms are in evidence other than a slight exudation of a rusty-colored liquid. The layer beneath the outer brown bark is black, and the cortex is first yellow gray, then claret colored. The cankered areas are surrounded by black lines. The disease may spread around the trunk and so destroy the tree. See also *Theobroma* for additional hosts and geographical distribution.

**Phytophthora meadii** McRae. This species causes fruit rot, leaf fall, and black-thread or black-stripe disease. The fruit rot and leaf fall caused by this fungus bring about heavy reductions in latex yield, which is not the case with the preceding species. The symptoms are similar, however. The black-thread phase of this *Phytophthora* disease occurs on the tapped surface, beginning as a series of narrow, vertical, parallel black lines, which may broaden out and coalesce so as to form a continuous wound parallel to the tapping cut. These black lines may extend through to the wood. In addition to *H. brasiliensis*, *Theobroma cacao* is attacked by this species. It is reported from Ceylon, India, Java, and Malaya.

**Poria hypobrunnea** Petch. A root disease of *H. brasiliensis*, *Crotalaria fulva*, *Hibiscus* sp., *Panax* sp., *Tephrosia candida*, and *Thea* (cult.) in Ceylon and Java. The mycelium forms stout red strands on the exterior of the root, which may unite into a continuous red sheet. These rhizomorphs are smooth and tough, bright red to brownish red, white within, turning black with age. Diseased wood is soft, friable, and permeated with red mycelial sheets, which often follow the lines of the annual rings. In advanced cases a soft wet rot is present, with a network of white threads between the bark and wood. The fruiting body is first yellowish white or ochraceous, then red brown, and finally dark-slate color.

**Rosellinia bunodes** B. and Br. See Citrus.

**Sclerotium** sp. Destroys bark on tapped surfaces of *H. brasiliensis* in Sumatra.

**Scolecotrichum heveae** Vinc. Small brown spots, the centers of which drop out on *H. brasiliensis* in Brazil.

**Sphaeronema** sp. Attacks the tapped surface of trunks of *H. brasiliensis* in Malaya, rotting and destroying the exposed cortex tissues and making further tapping for some years impossible.

**Sphaerostilbe repens** B. and Br. Black or red flattened fungous strands or rhizomorphs occur under the cortex of diseased roots and the wood is stained deep blue or purple. There is often a foul smell connected with the disease. Infected plants gradually die. Fruiting bodies of the fungus are of two types, the conidial occurring as short, erect, red, hairy stalks 2 to 8 millimeters high, surmounted by a pink head of conidia. The second consists of small, dark-red perithecia at the base of the conidial stalks. In addition to *H. brasiliensis* the disease attacks *Artocarpus integrifolia*, *Carica papaya*, *Casuarina equisetifolia*, *Citrus* spp., *Erythrina* spp., *Hibiscus* sp., *Mangifera indica*, *Maranta arundinacea*, and *Thea sinensis* in Ceylon, India, Malaya, Uganda, Fiji, Gold Coast, and the British West Indies.

**Thread blight.** A number of species of "thread blights" have been reported from India, Ceylon, Malaya, Java, and other tropical countries on *H. brasiliensis* and other hosts, some at least of which are parasitic. They are all characterized by white strands of mycelium, which run over stems and leaves of the hosts. This type of disease is described in detail for *Coffea*. (See *Corticium koleroga*.)

**Top canker.** This disease of *H. brasiliensis* in Ceylon is thought to be due to *Phytophthora faberi* and attacks the upper branches and stems. The bark splits longitudinally in lines a foot or more long, forming long, narrow scales which in time split transversely. The disease may penetrate to the wood, causing an open wound. The stems ultimately become thickened in the region of the cankers.

**Venturia emergens** Petch. On branches of *H. brasiliensis* in Ceylon.

**White stem blight.** This disease occurs in Ceylon on *H. brasiliensis* and *Thea* (cult.) and is caused by an undetermined fungus. The larger branches are attacked, white patches often 6 feet in length appearing on the bark. A fine white thread runs along the whitened patch, and the upper margin of the patch may show a fringe of hyphae.

**Witches'-broom.** A disease characterized by globose woody galls on the branches of *H. brasiliensis* is reported from Java. The cause has not yet been determined.

**Xylaria thwaitesii** Cke. This fungus appears as black, flat, irregular bands or patches on the roots of *H. brasiliensis* in Ceylon. A rot of the bark and wood results and diseased trees ultimately die. The rotten wood is dark brownish gray and moist but hard at the center of a root, light brown and decayed outside of this region. Black lines may be present. The inner cortex becomes brown or yellow brown and friable.

**HIBBERTIA.** Shrubs or subshrubs with yellow or white showy flowers.

**Puccinia hibbertiae** McAlp. Reddish-brown to black rust pustules on lower leaf surfaces of *H. sericea* and *H. stricta* in Australia.

**HIBISCUS.** ROSE MALLOW. Shrubs or small trees cultivated for their showy flowers and ornamental foliage.

**Accidium erythrobasis** B. and Br. Rust pustules on red leaf spots on *H. collinus* and *H. eriocarpus* in Ceylon and Uganda.

**Accidium garckeanum** P. Henn. Rust on leaves of *H. crassinervius* and *H. micranthus* in Abyssinia and the Congo.

**Bacterium hibisci** Nak. and Tak. Small circular black spots appear on the cotyledons and young leaves, enlarging gradually and becoming irregular to angular, with dull white to yellow outer zones. On *Hibiscus* sp. in Korea.

**Cercospora brachypoda** Speg. On leaves of *Hibiscus* sp. in Argentina. Reported from Louisiana.

**Cercospora hibisci-manihotis** P. Henn. Circular to effuse dull-brown spots on leaves of *H. manihot* in Japan.

**Colletotrichum hibisci** Póll. Circular to irregular brown leaf spots on *H. palustris* in Italy.

**Colletotrichum hibiscicolum** Rangel. On leaves of *H. tiliaceus* in Brazil.

**Corticium salmonicolor** B. and Br. See Citrus.

**Fomes lamaoensis** Murr. See Hevea.

**Phyllachora hibisci** Rehm. Shiny-black stromata on indefinite brown leaf spots on *H. furcatus* and *H. tiliaceus* in Ceylon, Samoa, and Brazil.

**HIBISCUS**—Continued.

- Phyllachora minuta** P. Henn. Black stromata on leaves of *H. tiliaceus* and *Thespesia populnea* in Porto Rico, Java, Australia, and the Philippines.
- Phyalospora hibisci** Racib. On leaves of *H. tiliaceus* in Java.
- Phytophthora melongenae** K. Saw. See Solanum.
- Poria hypobrunnea** Petch. See Hevea.
- Puccinia exilis** Syd. var. **hibisci** Grove. Brown rust pustules on leaves of *Hibiscus* sp. in Uganda.
- Puccinia hibisci** P. Henn. Dull-brown rust pustules on leaves of *H. rostellatus* in the Congo.
- Rhizoctonia destruens** F. Tassi. See Solanum.
- Rosellinia bunodes** B. and Br. See Citrus.
- Septoria hibisci** Sacc. Subcircular to irregular whitish leaf spots with dull-brown margins on *H. syriacus* in Italy.
- Septoria rosae-sinensis** Brun. Olive-brown leaf spots on *H. rosa-sinensis* in France.
- Sphaerostilbe repens** B. and Br. See Hevea.
- Tubercularia hibisci** Petch. On leaves of *H. sabdariffa* in Ceylon.
- Uromyces heterogenens** Cke. Brown rust pustules on leaves of *H. syriacus* in India.
- Xylaria** sp. This fungus causes a root disease of the ornamental *Hibiscus* in Hawaii.
- Wilt.** A wilt disease of *Hibiscus* spp., cause undetermined, is reported from Hawaii. Possibly the same as the preceding disease.
- HIERACIUM.** HAWKWEED. Hardy herbaceous perennials.
- Ascochyta hieracii** Lasch. On leaves of *H. pilosella* in Germany.
- Cercospora hieracii** Jaap. Large subcircular to angular brown leaf spots on *H. prenanthoides* in Switzerland.
- Entyloa calendulae** (Oud.) De By. See Calendula.
- Entyloa hieracii** Syd. Leaf smut on *Hieracium* spp. in Europe. A form of *E. calendulae*.
- Phyllosticta corcontica** Kab. and Bub. Subcircular to angular zoned yellow-brown to brown leaf spots becoming gray on *H. alpinum* in Austria.
- Phyllosticta hieracicola** E. Rostr. Dark-brown leaf spots on *H. umbellatum* in Norway.
- Phyllosticta hieracii** Lasch. Subcircular to irregular, often confluent, zoned dull yellow-brown leaf spots on *H. pilosella* and *H. pseudocerinthe* in Spain and Germany.
- Puccinia argentina** Speg. Cinnamon-brown rust pustules on leaves and stems of *Hieracium* sp. and *Picrosia longifolia* in Argentina.
- Ramularia conspicua** Syd. Circular olive-brown leaf spots on *H. murorum* in Bohemia and Germany.
- Ramularia corcontica** Bub. and Kab. Subcircular to angular dull-yellow to brown leaf spots becoming gray on *H. alpinum* and varieties in Bohemia.
- Ramularia helvetica** Jaap. and Lind. Irregular greenish-gray or brown leaf spots on *H. albidum* in Switzerland.
- Ramularia hieracii** (Baum.) Jaap. On leaves of *H. boreale*, *H. schmidtii*, *H. sylvaticum*, *H. umbellatum*, and *H. vulgatum* in France, Switzerland, Austria, and Germany.
- Ramularia subalpina** Bub. Angular dirty-yellow leaf spots with narrow, dark-brown margins on *H. lanatum* in Yugoslavia.
- Septoria guardarronica** Gz. Frag. On leaves of *H. pilosella* in Spain.
- Septoria mougeoti** Sacc. and Roum. Large yellow leaf spots on *H. praealtum* and *Hieracium* sp. in France and Russia.
- Septoria palustris** (Ces.) Sacc. On leaves of *Hieracium* sp. in Italy.
- HIEROCHLOA. TORRESIA** Ag. Fragrant perennial grasses.
- Puccinia borealis** Juel See Agrostis.
- Puccinia hierochloae** Ito. Yellow to chestnut-brown rust pustules on leaves and sheaths of *H. borealis* and *Stipa siberica* in Japan.
- Septoria oxyspora** Penz. and Sacc. See Anthoxanthum.
- HIPPEASTRUM.** Showy bulbous plants.
- Mosaic.** An infectious mosaic, characterized by a mottling of the leaves of *Hippeastrum* spp., has been reported from Hawaii.
- HIPPOCREPIS.** Herbs or subshrubs.
- Cercospora hippocrepidis** Jaap. Subcircular gray to whitish leaf spots with brown margins on *H. comosa* in Switzerland.
- Uromyces anthyllidis** (Grev.) Schroet. See Anthyllis.
- Uromyces hippocrepidis** E. Mayor. Brown rust pustules on leaves of *H. ciliata* in Europe.
- HIPPOPHAE.** SEA BUCKTHORN. Ornamental woody plants grown for their silvery-gray foliage and bright-colored berries.
- Dothidella hippophaeos** (Pers.) Theiss. and Syd. Circular black stromata on leaves of *H. rhamnoides* in Italy and Switzerland.
- Septocylindrium olivascens** Thuem. See Elaeagnus.
- Septoria hippophaes** Desm. and Rob. On leaves of *H. rhamnoides* in France.
- HIPPURIS.** Perennial aquatic herbs.
- Physoderma hippuridis** Rostr. On leaves of *H. vulgaris* in northern Europe, Greenland, and Iceland.
- HOHERIA.** Small shrubs or trees, native of New Zealand.
- Puccinia hoheriae** Wakef. Brown rust on leaves and stems of *H. populnea* in New Zealand.
- HOLCUS.** SORGHUM. Johnson grass. Sudan grass. Annual or perennial grasses, cultivated for production of grain, forage, sirup, and broomcorn.
- Acrothecium lunatum** Wakker. On leaves and heads of *H. sorghum* in Ceylon and Java.
- Aecidium evansii** P. Henn. Rust on leaves of *H. sorghum* in the Union of South Africa.
- Apiospora lloydii** (Crouan.) Sacc. On *Holcus* sp. and *Dactylis* sp. in France.
- Ascochyta sorghina** Sacc. Long dull-brown leaf spots with blood-red margins on *H. sorghum* in Italy.
- Bacillus omelianskii** Serbinov. Said to cause a soft black rot of tops of plants of *H. sorghum* in Russia.
- Beniowskia penniseti** Wakef. See Pennisetum.
- Botryodiplodia sorghi** P. Henn. On culms of *H. sorghum* in Tanganyika.
- Cercospora acerosa** Dickh. and Arendt. H. Black spots on leaves of *H. sorghum* in Java.
- Cerebella sorghi-vulgaris** Subra. In ovaries of *H. sorghum* in India.
- Colletotrichum andropogonis** A. Zimm. Dark-red leaf spots on leaves of *H. sorghum* in Tanganyika.
- Dilophia graminis** (Fckl.) Sacc. See Triticum.
- Helminthosporium caryopsidium** Sacc. Attacks the grain of various varieties of *H. sorghum*, causing "wholesale destruction of seed" in China and the Philippines.
- Leptosphaeria eustonii** (Fr.) Sacc. See Secale.

**HOLCUS**—Continued.

- Microbasidium sorghi** (Pass.) Bub. and Ran. Elliptical red to dark-brown leaf spots on *H. sorghum* var. *saccharatum* in Yugoslavia.
- Mycosphaerella ceres** Sacc. Pale leaf spots with reddish margins on *H. sorghum* in Russia and Italy.
- Mycosphaerella tassiana** De N. On leaves of *H. lanatus* in Austria.
- Ovularia holci-lanati** Car. Small dull dark-brown leaf spots on *H. (Notholcus) lanatus* in Italy.
- Phyllachora sorghi** V. Hoeh. Black stromata on yellow to white leaf spots with red margins on *H. sorghum* in var. in Java and the Philippines.
- Phyllosticta sorghina** Sacc. Irregular leaf spots with red borders on *H. effusum*, *H. halepensis*, and *H. sorghum* in Russia, India, Italy, Portugal, and Argentina. Reported from Texas.
- Puccinia holcina** Erikss. Yellow-brown to brown rust pustules on leaves of *H. lanatus*, *H. mollis*, and *H. setiglumis* in Europe.
- Ramulispora andropogonis** Miuri. Leaf spot on *H. sorghum* in Manchuria.
- Sclerospora graminicola** (Sacc.) Schroet. var. **andropogonis sorghi** Kulk. This downy mildew attacks plants of *H. sorghum* in all stages of growth in India. The leaves of diseased seedlings are narrow, pale-yellow, and covered with a fine white down. Later on white streaks appear on the upper leaves and the tissues tear along these streaks. Diseased plants remain stunted and sterile. Older plants may also be attacked, singly or in groups, long, narrow streaks appearing on the leaves at first pale-yellow, then orange, and finally dark-brown in color.
- Sclerospora philippinensis** Weston. See Zea.
- Septoria holci** Pass. Small gray subcircular leaf spots on *H. (Notholcus) lanatus* in Italy.
- Sorosporium ehrenbergii** Kuhn. Smut sori in the ovaries, covered at first by a false membrane. On *H. cernuus* in Egypt and Tunis.
- Sorosporium simii** Pole-Evans. Smut attacking and destroying inflorescences of *H. halepense* in the Union of South Africa.
- Tilletia holci** (West.) Rostr. (*T. rauwenhoffii* Fisch de W.) Irregular, black smut sori in the ovaries of *Bromus arvensis*, *B. mollis*, *Holcus (Notholcus)*, *lanatus* and *H. mollis* in Denmark, Belgium, and Great Britain. Reported from Washington and Oregon.
- Tolyposporium filiferum** Busse. Individual grains of the heads of *H. sorghum* in India, Egypt, and east Africa are transformed into smut sori. The sori are longer than the normal grain and light brown in color, soon rupturing, exposing the powdery spore masses.
- Tolyposporium volkensis** P. Henn. Black smut sori in the ovaries of *H. sorghum* in tropical Africa.
- Ustilago bulgarica** Bub. Greenish-brown smut sori in the ovaries of *H. sorghum* in Bulgaria.
- Ustilago sorghicola** Speg. Subcylindrical smut sori in the ovaries of *H. sorghum* in Argentina and Italy. The sori project beyond the glumes, reaching a length often of 8 millimeters.

**HOMERIA.** Bulbous plants from south Africa.

**Uredo homeriae** Bub. Rust on leaves of *Homeria* sp. in the Union of South Africa.

**HOMOGYNE.** Hardy perennials grown for their attractive white or purple heads of flowers.

- Ascochyta homogyne** Ranoj. Brown leaf spots on *H. alpina* in Yugoslavia.
- Puccinia conglomerata** (Strauss.) Kze. and Schm. Dark-brown rust pustules on leaves of *H. alpina* in France, Switzerland, Yugoslavia, Austria, and Germany.
- Ramularia cervina** Speg. On leaves of *H. alpina* in Yugoslavia, Italy, and Austria.
- Septoria prostrata** Kab. and Bub. Subcircular brown, then gray, leaf spots with dark-brown margins on *H. alpina* in Yugoslavia and Austria.
- Uromyces veratri** (DC.) Schroet. See Veratrum.

**HONCKENYA.** See Arenaria.**HORDEUM.** BARLEY. SQUIRRELTAIL GRASS. Annual grasses.

- Bacillus cerealinum** Gentner. Brown spots or blotches appear on the nodes, upper internodes, and leaves of *H. vulgare* (*H. sativum*), *Secale cereale*, and *Triticum* sp. in Europe. The upper leaves become prematurely yellow and dry, the heads fill out poorly, and the glumes may split.
- Cercospora elymi** Rostr. Oblong dull-brown leaf spots on *H. (Elymus) arenarium* in Denmark.
- Fusarium hordearium** Duc. Attacks the leaves and stems of *H. distichon* in France.
- Fusarium nivale** (Fr.) Sor. See Secale.
- Heterosporium hordei** Bub. Rhombic or elliptic-oblong, often confluent, dull-yellow leaf spots with dark-purple margins on *H. distichum* and *H. sativum* in Denmark and Yugoslavia.
- Micrococcus tritici** Prill. See Triticum.
- Mycosphaerella basicola** Frank. On leaves of *H. sativum* and *Secale cereale* in Sweden, Denmark, and Germany.
- Mycosphaerella exitialis** Morini. On culms of *H. sativum* and *Triticum* sp. in Japan, France, Denmark, Sweden, and Italy.
- Mycosphaerella hordicola** Hara. On leaf blades and culms of *Hordeum* sp. and *Triticum* sp. in Japan.
- Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.
- Ophiobolus herpotrichus** (Fr.) Sacc. See Triticum.
- Ramularia hordei** McAlp. Subcircular, then confluent, brown to dark-brown leaf spots on *Hordeum* sp. in Australia.
- Rhizoctonia napi** West. See Brassica.
- Septoria elymi** Rostr. On leaves of *H. arenarium* in Denmark.
- Septoria halophila** Speg. Large, ashen-colored blotches on leaves of *H. halophilum* in Argentina.
- Tilletia hordei** Koern. Dark-green smut sori in the ovaries, concealed at first by the glumes, then black, with a foetid odor. On *H. fragile* and *H. murinum* in Turkey, Algeria, Persia, and Australia.
- Tilletia hordeina** Ranoj. Yellow-brown spore masses in ovaries of *H. maritimum* in Yugoslavia.
- Tilletia panicii** Bub. and Ranoj. Dark violet-brown smut sori in the ovaries of *H. sativum* in Yugoslavia.
- Tilletia trauti** Jacz. A smut forming brown spore masses in the spikelets of *H. murinum* in Algeria.
- Tylenchus hordei** Schöyen. See Avena.
- Typhula graminum** Karst. See Lolium.

**HOSTA.** NIOBE Ag. PLAINTAIN LILY. Hardy perennials.

- Aecidium hostae** Diet. Rust pustules on yellow, circular leaf spots on *H. sieboldiana* in Japan.
- Ascochyta herreana** P. Henn. On leaves of *H. ovata* in Germany.
- Ascochyta hortensis** Kab. and Bub. Circular to angular brown leaf spots with purple-brown margins on *H. albomarginata* in Bohemia.
- Cercospora hostae** Hori. On leaves of *H. sieboldiana* in China.
- Fusarium versiforme** Kab. and Bub. Irregular brown spots on both leaf surfaces of *H. albomarginata* in Bohemia.
- Mycosphaerella hostae** Syd. On leaves and petioles of *H. japonica* in Japan.
- Phyllosticta funkiae** Fen. Pale irregular leaf spots with brown margins on *H. ovata* in Italy.
- Puccinia funkiae** Diet. Black rust pustules on the lower leaf surfaces of *H. longipes*, *H. ovata*, *H. plantaginea*, and *H. sieboldiana* in Japan and other Asiatic countries.
- Uredo hostae** P. Henn. Brown rust pustules on yellow leaf spots on *Hosta* sp. in Japan.

**HOTTONIA.** FEATHERFOIL. Perennial aquatic herbs.

**Doassansia hottoniae** Rostr. Smut pustules in rufous leaf spots on *H. palustris* in Denmark.

**HOVENIA.** RAISIN TREE. Ornamental shrubs, grown for their handsome foliage.

**Microsphaera yamadai** Syd. Powdery mildew on leaves of *H. dulcis* in Japan.

**HOYA.** WAX PLANT. Tropical climbers or trailing evergreen shrubs.

**Gloeosporium affine** Sacc. Anthracnose on leaves of *Aloe hanburyana*, *Citrus aurantium*, and *Hoya carnosa* in Great Britain, Italy, Gibraltar, and the Philippines.

**Gloeosporium hoyae** Syd. Anthracnose on leaves of *H. luzonensis* in the Philippines.

**Gloeosporium intermedium** Sacc. See Citrus.

**Guignardia albicans** Rehm. On leaves of *H. luzonensis* in the Philippines.

**Phyllosticta asclepiadearum** West. See Cynanchum.

**Phyllosticta thuemenii** Tass. On leaves of *H. carnosa* in Austria and Gibraltar.

**Physalospora hoyae** v. Hoehn. On leaves of *H. luzonensis* in the Philippines.

**Puccinia variiformis** Pat. Dark-brown rust pustules on leaves of *Hoya* sp. in Indo-China.

**Septoria hoyae** Sacc. Ivory-white irregular leaf spots with dull-brown margins on *H. carnosa* in Italy.

**HUMULUS.** HOP. Twining vines.

**Ascochyta humuli** Lasch. Pale-brown leaf spots on *H. lupulus* in Germany.

**Ascochyta humuli** Bub. and Kab. Circular to irregular greenish, then dirty-ochraceous to ashen, leaf spots on *H. lupulus* in Bohemia. This species if not synonymous with the preceding must be renamed.

**Cercospora cantariensis** Salm. and Worm. Circular gray leaf spots with narrow purple-brown margins and surrounding yellow zones on *H. lupulus* in Great Britain.

**Cercospora humuli** Hori. Angular dark-brown spots, often coalescing, on *H. japonicus* and *H. lupulus* in Japan.

**Fusoma parasiticum** Salm. This fungus causes a serious wilt disease of hops in Great Britain. Above ground its presence is shown by a wilting of the stems, which break away easily or fall over, having been rotted just below the ground line. The bark and wood at this point are brown and dead. Pink pustules of the causative fungus appear on the basal parts of diseased plants.

**Mosaic.** A mosaic disease of *H. lupulus* is reported from Great Britain. The stems of diseased plants grow only to a height of 4 to 6 feet, the leaves are curled, more or less mottled, green and yellow in color, and very brittle. Diseased plants are sterile.

**Peroplasmodium humuli** Miy. and Tak. This serious downy mildew of the hop (*H. lupulus*) occurs in Japan and Great Britain. It has also been reported from Wisconsin. Small angular dark-green spots, with a water-soaked aspect, first appear on the leaves, becoming finally dark-brown. Diseased leaves dry up and die.

**Phyllosticta bractearum** Oud. On bracts of *H. lupulus* in Holland.

**Phyllosticta japonica** Fautr. Brown irregular leaf spots on *H. japonicus* in Japan and France.

**Phyllosticta lupulina** Kab. and Bub. Circular to angular, confluent, ashen-colored leaf spots with purple-brown borders on *H. lupulus* in Austria and Bohemia.

**Plasmodiophora humuli** Kirk. Causes a disease of *H. lupulus* in New Zealand and Tasmania which resembles clubroot of the crucifers.

**Septoria divergens** Bub. and Kab. Small subcircular to angular ochraceous-brown leaf spots with narrow brown margins on *H. lupulus* in Bohemia.

**Septoria humuli** West. On leaves of *H. lupulus* in Europe.

**Septoria humulina** Bond. Pale to dull-brown leaf spots on *H. lupulus* in Russia.

**HURIA.** SAND-BOX TREE. Tropical trees.

**Cercospora hurae** Stevens. Small brown leaf spots on *H. crepitans* in Porto Rico.

**Fomes lamaoensis** Murr. See Hevea.

**Helminthosporium hurae** P Henn. Circular, then confluent, dull-brown leaf spots on *H. crepitans* in Brazil.

**HUTCHINSIA.** Low, annual or perennial herbs.

**Puccinia hutchinsiae** Rud. Dark-brown rust pustules on leaves of *H. alba* in Turkestan.

**HYACINTHUS.** HYACINTH. Hardy bulbous plants cultivated for their showy flowers.

**Ascochyta hyacinthi** Tassi. On leaves of *Agapanthus umbellatus* and *H. orientalis* in Italy and Australia.

**Bacillus croci** Mizus. See Crocus.

**Bacillus hyacinthus septicus** Heinz. Bulbs, leaves, and flower stalks are attacked by this disease, which starts as a soft rot of the bulb, rapidly extending into the above-ground parts of the plant. Leaves and flower buds wither and drop off. Diseased tissues soon break up, forming a foul-smelling slime. The disease has been reported on cultivated *Hyacinthus* in Great Britain and Germany. By some workers this species is considered synonymous with *B. caratovorvus*.

**Bacterium hyacinthi** Wakke. The disease known as Wakker's hyacinth disease, or the yellow-slime disease, is characterized by yellow stripes beginning near the apex of the leaf and extending downward rapidly, but only slowly sidewise, so that finally the leaf will have a central dead stripe along its entire length with the margins still green. In the bulb the vascular bundles become yellow and full of bacterial slime. The yellow slime extends upward into the bundles of other scales and sidewise slowly into the parenchyma until finally the entire bulb is destroyed. In advanced stages other bacteria enter and aid in destroying infected bulbs. All varieties of *Hyacinthus* are attacked. The disease has been reported from France, Great Britain, Holland, Italy, and Germany. There is one report of its occurrence in the United States.

**Peziza vesiculosa** Bull. This fungus is reported as attacking entire plants, with a growth of mold externally; in Germany on *Hyacinthus* in variety.

**Puccinia liliacearum** Duby. See Ornithogalum.

**Septoria podolica** Bub. and Wrobl. On leaves of *H. leucophaeus* in Galicia.

**Sclerotinia tuliparum** (Wakk.) Rehm. See Tulipa.

**Tylenchus dipsaci** Kuehn. See Narcissus.

**Uromyces scillarum** (Grev.) Wint. A common and widespread rust which produces brown pustules on both leaf surfaces, disfiguring the affected plants. The hosts are *Bellevalia comosa*, *B. fuliginosa*, *B. mauritanica*, *Endymium nutans*, *Hyacinthus ciliatus*, *H. fastigiatus*, *H. orientalis*, *H. pratensis*, *Leopoldia bouvieriana*, *Muscari botryoides*, *M. comosum*, *M. monstrosum*, *M. neglectum*, *M. polyanthum*, *M. racemosum*, *M. tenuiflorum*, *Ornithogalum comosum*, *Scilla autumnalis*, *S. bifolia*, *S. bertolonii*, *S. campanulata*, *S. italica*, *S. obtusifolia*, *S. peruviana*, *S. pratensis*, *S. pumila*, *S. ramburii*, *S. verna*, *Simethis bicolor*, *Urginea maritima*, and *U. scilla*. It occurs in Great Britain, France, Spain, Portugal, Italy, Switzerland, Holland, Russia, Austria, Hungary, Corsica, Montenegro, Serbia, Rumania, Greece, Morocco, Algeria, Tunis, Egypt, Dalmatia, the Union of South Africa, and Germany.

**HYACINTHUS**—Continued.

**Ustilago vaillantii** Tul. One of the smuts producing black powdery masses of spores in place of the anthers and ovaries in the flowers. It is perennial in the bulbs and hence easily carried from place to place. The flowers are disfigured and their commercial value lowered. The hosts are *Bellevalia romana*, *B. trifoliata*, *Chionodoxa luciliae*, *Chionodoxa* sp., *Gagea* spp., *Hyacinthus* spp., *Muscari comosum*, *M. racemosum*, *M. schliemanni*, *M. tenuiflorum*, *Scilla anthericoides*, *S. bifolia*, *S. cernua*, *S. lilio-hyacinthus*, *S. maritima*, *S. trifolia*, *Urginea anthericoides*, and *U. maritima*, reported from Great Britain, France, Corsica, Italy, Belgium, Serbia, Egypt, Algeria, Austria, Germany, and from a few localities in the United States.

**HYALIS**. See *Ixia*.**HYDRANGEA**. Ornamental woody plants grown for their showy flowers.

**Aecidium hydrangeae** Pat. Rust on leaves of *H. davidi* in Thibet.

**Aecidium hydrangeae-paniculatae** Diet. Rust pustules on large, irregular, yellow, or reddish leaf spots on *H. paniculata* in Japan.

**Aecidium hydrangicolum** P. Henn. Rust on leaves of *H. hirta* and *H. thunbergi* in Japan.

**Ascochyta hydrangeae** Arn. Circular to irregular greenish-gray spots on leaves and stems of *H. hortorum* in France. The spots may spread to include the entire area of leaves.

**Cercospora hydrangeicola** Speg. Circular, then confluent, purplish leaf spots on *H. hortensis* in Argentina.

**Cercospora obtegans** Syd. On leaves of *H. hortensis* in Japan.

**Clasterosporium hydrangeae** Sacc. On *H. hortensis* in Italy and Madeira.

**Rhytisma** sp. On leaves of *H. paniculata* in Japan.

**Septoria anthophila** Fl. Tass. On flowers of *H. hortensis* in Italy.

**Septoria hydrangeae** Bizz. Irregular, then confluent, dull-brown leaf spots with blood-red margins on *H. hortensis* in Germany. Said to cause serious damage. Reported from Ohio.

**HYMENAEA**. Tropical trees.

**Rosellinia pepo** Pat. See Citrus.

**Uredo hymenaeae** Mayor. Powdery light-brown rust pustules covering the under leaf surfaces of *H. courbaril* in Porto Rico, Cuba, and northern South America.

**HYMENOCALLIS**. SPIDER LILY. Bulbous plants.

**Tubercinia javanica** Koord. Large dark-brown areas on leaves of *Hymenocallis* sp. in Java.

**HYMENOPHYLLUM**. Filmy fern.

**Micropeltis hymenophylli** Pat. Small black fruiting bodies on leaves of *Hymenophyllum* sp. in Tahiti.

**Sporotrichum niveum** Allesch. and Henn. On leaves of *H. bridgesii* in Chile.

**HYOSCYAMUS**. HENBANE. Herbs cultivated for medicinal purposes.

**Ascochyta hyoscyami** Pat. Circular dull-brown leaf spots on *H. niger* in Tunis, Russia, Hungary, and Germany.

**Ascochyta pinzolensis** Kab. and Bub. Irregular and confluent brown or dull-yellow leaf spots, becoming white with brown margins on *H. niger* in Italy.

**Peronospora hyoscyami** De By. Downy mildew on leaves of *H. niger* in Bohemia, Silesia, and and Russia. See also *Nicotiana*.

**Puccinia hyoscyami** P. Magn. Dark-brown rust pustules on leaf blades and petioles of *H. senecionis* in Persia.

**Septoria hyoscyami** Hóll. Dull-yellow leaf spots with brown margins on *H. niger* in Hungary.

**Septoria pinzolensis** Kab. and Bub. Circular to angular brown or dull-yellow leaf spots, becoming white with brown margins on *H. niger* in Italy.

**HYOSERIS**. Stemless herbs.

**Entyloma aposeridis** Jaap. Smut sori in small dull-brown leaf spots on *H. foetida* in Austria.

**Phyllosticta aposeridis** Allesch. Irregular dull-yellow to brown leaf spots on *H. foetida* in Germany.

**Protomyces hyoseridis** Syd. Circular yellow leaf spots on *H. baetica* in Italy.

**Puccinia hyoseridis-radiatae** R. Maire. Cinnamon-brown to dark-brown rust pustules on leaves of *H. radiata* and *H. scabra* in Tripoli, Spain, and Dalmatia.

**Septoria hyoseridis** Maire. Brown leaf spots with purple margins on *H. radiata* in tropical Africa.

**HYPERANTHERA**. See *Moringa*.**HYPERICUM**. ST.-JOHN'S-WORT. Annual herbs.

**Ascochyta hyperici** Lasch. Dull-brown leaf spots on *H. perforatum* in Italy and Germany.

**Melampora hypericorum** Wint. Powdery orange-yellow then reddish-brown rust pustules on leaves of *Androsaemum* spp. and *Hypericum* spp. in Europe, Asia, and Africa.

**Melampora kusanoi** Diet. Chestnut-brown, then black, rust pustules on yellow sunken leaf spots on *H. ascyron* in Japan.

**Melampora sancti-johannis** Barel. Rust on leaves of *H. cernuum* and *H. patulum* in India.

**Metasphaeria cryptospila** (Berk.) Sacc. On leaves of *Hypericum* sp. in New Zealand.

**Mycosphaerella hyperici** (Auersw.) Starb. On stems of *Hypericum* spp. in Europe.

**Ovularia minutissima** Syd. Circular to elliptical brown leaf spots with purple margins on *H. quadrangulum* in Germany.

**Ramularia coleosporii** Sacc. See *Campanula*.

**Septoria carestiana** Ferr. Pale-yellow or reddish leaf spots on *H. montanum* in Italy.

**Septoria hyperici** Desm. and var. **hyperici-quadranguli** C. Mass. Circular oblong, or indefinite, red-brown leaf spots with dull-yellow margins on *H. burserum*, *H. hirsutum*, *H. montanum*, *H. perforatum*, *H. pulchrum*, and *H. quadrangulum* in Europe.

**Septoria hyperici-burseri** R. Maire. On leaves of *H. burserum* in France.

**Uredo hyperici-japonici** Petch. Rust on leaves of *H. japonicum* in Ceylon.

**Uredo hyperici-mysorensis** Petch. Rust on leaves of *H. mysorensis* in Ceylon.

**HYPOXIS**. GOLD-EYE GRASS. STAR GRASS. Herbaceous plants with hard root stocks or corms.

**Cylindrosporium guttatum** Wint. Round to irregular dull-brown leaf spots on *Hypoxis* sp. in Brazil.

**Puccinia expallens** Syd. Gray-brown rust pustules on leaves of *H. aurea* in India.

**Puccinia hypoxidis** McAlp. Dark-brown to black rust pustules on both leaf surfaces of *H. hygrometrica* in Victoria, Australia.

**Uromyces hypoxidis** Cke. Brown rust pustules on both leaf surfaces of *H. oligotricha* in Natal and Uganda.

**IBERIS**. CANDYTUFT. Small flower-garden and border plants.

**Helminthosporium iberidis** Póll. Circular dull-brown leaf spots on *Iberis* sp. in Italy.

**Puccinia iberidis** Duby. Rust on leaves of *I. sempervirens* in France, Switzerland, and Belgium.

**ICACOREA**. See *Ardisia*.**ICHTHYOMETHIA**. See *Piscidia*.**IDESIA**. Ornamental trees, grown for their handsome foliage and attractive berries.

**Melampora idesiae** Miy. Yellow to brown rust pustules on leaves of *I. polycarpa* in Japan.

- ILEX. HOLLY.** Maté, sometimes called Paraguay tea. Evergreen or deciduous woody plants, grown for their attractive leaves and berries. The leaves of several South American species are used for beverage purposes.
- Boydia inculpta** (Oud.) Grove. Destroys twigs of *I. aquifolium* in Great Britain and probably continental Europe.
- Cercospora ilicicola** Maubl. On leaves of *I. paraguariensis* in Brazil. A doubtful report from Texas.
- Cercospora yerbac** Speg. Dull-brown circular leaf spots on *I. amara* in Argentina.
- Cercosporina mate** Speg. Circular ashen-colored leaf spots with purple margins on *I. paraguariensis* in Argentina.
- Colletotrichum yerbac** Speg. Subcircular whitish or ashen leaf spots on *I. paraguariensis* in Argentina and Brazil.
- Cylindrosporium pollaccii** Ture. Circular whitish leaf spots with broad, dark-purple margins on *I. furcata* in Italy.
- Depazea ilicicola** Fr. On leaves of *I. aquifolium* in France.
- Exobasidium giseckiae** Allesch. White to reddish swellings on leaves and flowers of *Giseckia pharnacioides* and *I. aquifolium* in Italy and tropical Africa.
- Gloeosporium alutaceum** Sacc. Anthracnose on leaves of *I. aquifolium* in Italy.
- Gloeosporium aquifolii** Penz. and Sacc. Anthracnose on leaves of *I. aquifolium* in Italy.
- Gloeosporium orthosporum** Sacc. White indefinite leaf spots on *I. aquifolium* in Italy.
- Laestadia ilicis** Jacz. On leaves of *I. aquifolium* in Switzerland.
- Leptosphaeria paraguariensis** Maubl. On leaves of *I. paraguariensis* in Brazil.
- Peckia mate** Speg. Fruiting bodies occur on lower leaf surfaces of *I. paraguariensis* in Argentina and cause serious damage.
- Phyllosticta azevinhi** Torr. Brown leaf spots on *I. azevinhi* in Madeira.
- Phyllosticta haynaldi** Roum. Indefinite dull-red leaf spots on *I. aquifolium* in Denmark and France.
- Phyllosticta iliciperda** Oud. Irregular gray leaf spots on *I. aquifolium* in Holland.
- Phyllosticta ilicis** Oud. Pale irregular leaf spots on *I. aquifolium* in Holland.
- Phyllosticta mate** Speg. Circular, then confluent, silvery leaf spots on *I. paraguariensis* in Argentina and Brazil.
- Phyllosticta yerbac** Speg. Definite circular dull-brown leaf spots on *I. paraguariensis* in Argentina.
- Rhynisma himalense** Syd. and Butl. Black, slightly raised areas on leaves of *Ilex* sp. in India.
- Rhynisma ilicis-latifoliae** P. Henn. Black patches on subcircular leaf spots on *I. integra*, *I. latifolia*, *I. othera*, and *I. peduncularis* in Japan.
- Septoria orthospora** Lévl. On leaves of *I. aquifolium* in France.
- Trabuttia lagerheimiana** (Rehm.) Theiss. and Syd. Elliptical shiny black stromata on leaves of *I. scopulorum* in Ecuador.
- Venturia ilicifolia** Cke. On leaves of *I. aquifolium* and *I. balearica* in Europe.
- ILLICIUM. ANISE TREE.** Small trees or shrubs.
- Gloeosporium ilicium** Hemmi. Anthracnose on *I. anisatum* in Japan.
- Phyllosticta aromatica** F. Tassi. Circular to elliptical gray leaf spots on *I. floridanum* in Italy.
- IMPATIENS. SNAPWEED. TOUCH-ME-NOT. BALSAM.** Flower-garden annuals.
- Ascochyta impatientis** Bres. Subcircular to angular dull-yellow to brown leaf spots on *I. parviflora* in Germany.
- Cercospora campi-silii** Speg. Circular to angular brown leaf spots on *I. nolitantere* in Denmark, Italy, and Austria.
- Cercospora impatientis** Baeuml. Circular, then irregular, gray or whitish leaf spots with dull-brown margins on *I. noli-tangere* and *I. parviflora* in Hungary.
- Crönartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- Gloeosporium impatientis** Petch. Soft rot of stems of *I. balsamina* in Ceylon.
- Puccinia komarovi** Tranzsch. Powdery cinnamon-brown to chestnut-brown rust pustules on leaves of *I. amphorata* and *I. parviflora* in Turkestan and India.
- Uredo balsaminiae** Cke. Rust on leaves of *I. oppositifolia* in Ceylon.
- IMPERATA. Large perennial tropical grasses.**
- Cercospora imperata** Syd. Leaf rust on *Imperata* sp. in the Philippines.
- Phyllachora cyperi** Rehm. var. **donacis** Berl. and Sacc. Tar spot on leaves of *I. arundinacea* in India.
- Phyllachora imperatae** Syd. Black stromata on leaves of *I. cylindrica* in the Philippines. Probably identical with the preceding species.
- Puccinia fragosoana** Beltram. Leaf rust on *I. cylindrica* in Spain.
- Puccinia infuscans** Arth. and Holw. Brown powdery leaf rust on *I. braziliensis* in Guatemala.
- Puccinia rufipes** Diet. Dull-yellow to black rust pustules on leaves of *I. arundinacea* and *I. cylindrica* in the Philippines, Japan, Ceylon, China, India, and Egypt.
- Sphacelotheca schweinfurthiana** (Thuem.) Sacc. Smut sori replacing ovaries of *I. arundinacea* and *I. cylindrica* in Uganda, Egypt, Tunis, Tripoli, and Spain.
- INDIGOFERA. INDIGO.** Shrubs or perennial herbs sometimes grown for ornament. Several species furnish dyestuff.
- Corticium salmonicolor** B. and Br. See Citrus.
- Diplodia cacaocicola** P. Henn. See Theobroma.
- Gloeosporium incarpi** Sacc. Anthracnose on leaves of *I. arrecta* in Malaya.
- Pythium indigoferae** Butl. On leaves of *I. arrecta* in India.
- Ravenelia indigoferae** Tranzsch. Powdery brown rust pustules on leaves and stems of *I. anil*, *I. conzattii*, *I. cuernavacana*, *I. mucronata*, *I. palmeri*, and *I. suffruticosa* in Mexico, Central America, northern South America, West Indies, Trinidad, and Bermuda.
- Ravenelia laevis** Diet. and Holw. Powdery brown rust pustules on leaves of *I. densiflora* and *I. jalisensis* in Mexico.
- Ravenelia schroeteriana** P. Henn. Powdery cinnamon-brown rust pustules on leaves of *Indigofera* sp. in Argentina.
- Uredo maranguensis** P. Henn. Ashen-brown rust pustules on leaves of *I. arrecta* in tropical Africa.
- Uromyces mbelensis** P. Henn. Brown rust pustules on *Indigofera* sp. in the Congo.
- Uromyces orientalis** Syd. Cinnamon-brown rust pustules on leaves of *I. cordifolia*, *I. glandulosa*, and *I. linifolia* in India.
- Uromyces sphacrocarpus** Syd. Brown rust pustules on leaves of *I. pseudotinctoria* and *I. tinctoria* in Japan.
- INGA. Tropical trees and shrubs, commonly used as coffee shade.**
- Linospora guaranitica** Speg. Indefinite spots on leaves of *Inga* sp. in Brazil.
- Ophidothella ingae** (P. Henn.) Theiss. and Syd. Dull-black irregular stromata on large yellow or yellow-brown leaf spots on *I. strigillosa* and *Inga* sp. in Brazil.
- Ravenelia ingae** (P. Henn.) Arth. Cinnamon-brown rust pustules on leaves and stems of *I. edulis*, *I. inicuil*, *I. laurina*, and *I. vera* in Mexico, South and Central America, and the West Indies.

**INGA**—Continued.

- Stilbella flavida** (Cke.) Lind. See Coffea.  
**Uredo curvata** Arth. Brown leaf rust on *I. vera* in Cuba.  
**Uromyces ingicola** P. Henn. Brown rust pustules on leaves of *Inga* sp. in Brazil.  
**Uromyces porcensis** Mayor. Rust on *I. ingoides* in Colombia.

**INTSIA**. See Afzelia.**INULA**. Herbs with yellow or orange flower heads.

- Accidium inulae-crithmoidis** Pat. Rust on leaves of *I. crithmoides* in Dalmatia  
**Accidium inulae-helenii** Const. Rust on yellow irregular leaf spots on *I. helenium* in Italy.  
**Ascochyta inulicola** Petr. On *I. conyza* in Bohemia.  
**Coleosporium inulae** Rabh. Yellow rust pustules on *Pinus sylvestris*, *Inula aschersoniana*, *I. candida*, *I. coppa*, *I. ensifolia*, *I. graveolens*, *I. helenium*, *I. heterolepis*, *I. hirta*, *I. media*, *I. salicina*, *I. vaillantium*, and *I. viscosa* in Europe, Asia Minor, northern Africa, the Congo, and the Canary Islands.  
**Erysiphe taurica** Lévl. See Althaea.  
**Ovularia inulae** Sacc. and **forma major** P. Brun. Angular to irregular leaf spots with purple margins on *I. dysenterica* in France.  
**Phyllosticta inulae** Allesch. Circular whitish to dull brown leaf spots on leaves of *I. britannica* in Germany.  
**Ramularia cupulariae** Pass. On leaves of *I. conyza* and *I. viscosa* in Denmark, Spain, and Germany.  
**Ramularia inula-britannicae** Allesch. Subcircular yellow, then brown, leaf spots on *I. britannica* in Germany.  
**Septoria dysentericae** P. Brun. Irregular to angular small white leaf spots with narrow brown margins on *I. dysenterica* in France.  
**Septoria inulae** Sacc. and Speg. Rufous-brown leaf spots on *I. britannica* and *I. salicina* in China and Italy.  
**Uredo inulae-candidae** Trott. Yellow rust pustules on leaves of *I. candida* in Greece and Dalmatia.
- IPOMOEA**. MORNING-GLORY. SWEET POTATO. Annual or perennial twining herbs.  
**Accidium kaernbachii** P. Henn. Rust on leaves of *Ipomoea* spp. in Japan, Indo-China, Ceylon, New Guinea, Amboina, and the Philippines.  
**Cercospora batatae** Zimm. Irregular dark-brown leaf spots with paler centers on *I. batatas* in Tanganyika, Indo-China, Japan, Formosa, and the Philippines.  
**Cercospora stuckertiana** Syd. Subcircular gray leaf spots on *Ipomoea* sp. in Argentina.  
**Cercospora timorensis** Cke. Circular brown leaf spots on *I. cymosa* in Molucca.  
**Cylindrosporium bakeri** Syd. On *Ipomoea* sp. in the Philippines.  
**Helicobasidium mompa** Tan. See Morus.  
**Marsonia ipomoea** Cke. and Mass. Brown areas on stems and leaves of *Ipomoea* sp. in England.  
**Phyllosticta stuckertii** Speg. Small leaf spots with purple borders on *Ipomoea* sp. in Argentina.  
**Puccinia batatae** Syd. Brown rust pustules on leaves of *I. paniculata* in Natal and the Congo.  
**Puccinia holosericea** Cke. Brown rust pustules on leaves of *I. holosericea* in Tanganyika and Natal.  
**Puccinia insignis** Holw. Rust on leaves of *Ipomoea* sp. in Mexico.  
**Puccinia macrocephala** Speg. Black rust pustules on leaves of *Ipomoea* sp. and *Convolvulus* sp. in South America.  
**Puccinia nocticolor** Holw. Yellow, then black, powdery rust pustules on leaves of *I. fistulosa*, *I. intrapilosa*, and *I. murucoides* in Mexico and Guatemala.  
**Puccinia rubicunda** Holw. Rust on leaves of *Ipomoea* sp. in Mexico.  
**Puccinia superflua** Holw. Black powdery rust pustules on yellow leaf spots on *I. murucoides* in Mexico.  
**Ramularia batatae** Rac. Angular brown to dark-brown leaf spots on *I. batatas* in Java.  
**Sorosporium ipomoeae** Spesch. Smut on *Ipomoea* sp. in Russia.  
**Uredo ipomoeae-pentaphyllae** P. Henn. Rust on leaves of *I. pentaphylla* in Brazil.  
**Uredo speschnewii** S. and S. Brown powdery rust pustules on leaves of *Ipomoea* sp. in Transcaucasia.  
**Uromyces comptus** Syd. Dark-brown to black rust pustules on leaves of *I. bipinnatipartita* in Tanganyika and the Union of South Africa.  
**Uromyces ipomoeae** (Thuem.) Berk. Yellow, then black, rust pustules on leaves of *I. argyreoides* in the Union of South Africa.  
**Uromyces pachycephs** Lagh. Rust on leaves of *Ipomoea* sp. in Ecuador.
- IRESINE**. BLOOD LEAF. Ornamental-leafed bedding plants.  
**Cercospora gilbertii** Speg. See Celosia.  
**Puccinia macropoda** Speg. Cinnamon-brown to black rust pustules on leaves of *I. angustifolia*, *I. celosia*, *I. celosioides*, and *I. elatior* in South America and the West Indies.  
**Puccinia striolata** (Speg.) Arth. Powdery brown rust pustules on leaves of *I. angustifolia* and *I. celosia* in Trinidad, Porto Rico, Cuba, St. Thomas, and Argentina.  
**Septoria iresines** Speg. Circular or elliptical definite whitish leaf spots with dull-brown margins on *I. celosioides* in Argentina.  
**Uromyces celosiae** Diet. and Holw. See Celosia.  
**Uromyces iresines** Lagh. Dull-brown rust pustules on leaves of *I. celosia*, *I. elatior*, and *I. paniculata* in Guatemala, St. Thomas, Ecuador, and Colombia.
- IRIS**. Perennial rhizomatous or bulbous plants.  
**Ascochyta iridis** Oud. Leaf spot of *I. pseudacorus* in Holland.  
**Ascochyta pseudacori** All. On leaves of *Iris* (cultivated var.) in France.  
**Bacterium iridis** v. Hall. Infected leaves turn yellow and dry up. The underground portions are soft rotted. The disease is generally at its worst during the flowering season. It is readily carried on rhizomes or bulbs, either as small lesions or by means of the bacteria themselves in soil or debris or on the bulb scales. Soft rot of *Iris* bulbs and rhizomes is common in the United States, but the causal agents mentioned have been other than the above. *Bacillus omnivorus* is also given as the cause of an *Iris* rot in Europe. The disease occurs on many varieties and species of cultivated *Iris* in Holland, Great Britain, and France.  
**Cladochytrium (Physoderma) iridis** De By. Causes a leaf spot and blighting of *Iris* sp. in Germany.  
**Heterosporium montenegrinum** Bub. Forms pale spots with purple-brown margins on leaves of *I. graminea* in Montenegro. Said to be distinct from *H. gracile*.  
**Leptosphaeria heterospora** (De Not.) Niessl. Leaf spot and lesions on *Iris* rhizomes in Europe.  
**Leptosphaeria iridicola** Lamb. and Fautr. (*L. vectis* [B. and Br.] Ces. and De Not.) On *I. foetidissima* in France and Italy.  
**Macrophoma rhabdosporioides** Lamb. and Fautr. Reported as the cause of a leaf spot of *I. foetidissima* in France.  
**Metasphaeria iridicola** (Desm.) Sacc. On leaves and stems of *I. foetidissima* in Algeria.  
**Microdiplodia iridicola** Gz. Frag. On *I. filifolia*, *I. longepetala*, *I. pumila*, and *I. spuria* in Spain.  
**Mycosphaerella iridis** (Auersw.) Schroet. Causes a leaf spot on *I. filifolia*, *I. pseudacorus*, *I. pumila*, and *I. sysirinchium* in Great Britain, Denmark, France, Italy, Spain, and Germany.

**IRIS**—Continued.

- Mystrosporium adustum** Mass. This fungus forms large black crusty patches on the outer bulb scales, the mycelium reaching to the heart of the bulb. The disease is readily carried on the bulbs of the host (*I. reticulata*). It is said "to carry off bulbs by the hundreds, making it necessary to lift the bulbs every year and sort out the diseased ones for destruction and dry and store the remainder." The disease occurs in England and Denmark.
- Nectria paludosa** (Fekl.) Sacc. On leaves of *Iris* sp. and *Typha* sp. in Europe.
- Ophiobolus peduncularis** Feltg. At the base of stems of *I. pseudacorus* in Luxemburg. *O. bactrosporus* Feltg. and *O. pseudacori* Feltg. are reported on the same host and from the same region.
- Phoma agapanthi** Thuem. Attacks the base of *Iris* plants in Europe.
- Phoma iridis** Cke. On leaves of *I. foetidissima* in Europe.
- Phyllosticta iridicola** Gz. Frag. Reported from Spain as the cause of leaf spot of *I. pseudacorus*.
- Physoderma tenue** (Nowk.) Karst. See Nymphaea.
- Puccinia capensis** Diet. This rust fungus produces brown pustules on the leaves of *Iris* sp. in the Union of South Africa.
- Puccinia caucasica** Martino. Large stroma-like rust patches on leaves of *I. flavescens* in the Caucasus.
- Puccinia melanopsis** Syd. Black rust pustules are produced on both leaf surfaces of *I. sisyrinchium* in Asia Minor.
- Ramularia rolliandi** Faut. On leaves of *I. pseudacorus* in France.
- Schinzia (Naegelia) cellulicola**. Causes swellings on the roots of *Iris* sp. in Switzerland.
- Sclerotium crustuliforme** Rob. At the base of plants of *I. germanica* in Belgium.
- Sclerotium durum** Pers. At the base of plants and on rhizomes and bulbs of *Iris* spp. in Europe.
- Sclerotium iridis** Thuem. Attacks the rhizomes of cultivated *Iris* varieties, causing a soft rot distinct from bacterial rot. Several shipments of *Iris* rhizomes from Germany have been completely destroyed by what is probably this species. Doubtless also occurs in Holland and France. At least two other species of *Sclerotium* have been reported on *Iris* spp. in Europe, and it is not clear how many species really are involved. What has been referred to as *S. semen* has been intercepted on *I. tingitana*.
- Scolecotrichum cladosporioideum** R. Maire. On leaves of *I. foetidissima* in Algeria.
- Scolecotrichum iridis** Fautr. and Roum. On leaves of *Iris* sp. in Europe.
- Septoria iridina** Sacc. On leaves of *I. foetidissima* in Portugal.
- Septoria iridis** Massal. Circular, brown spots with light-brown margins on leaves and flower stalks of *I. florentina* and *I. germanica* (in var.) in Italy and Dalmatia.
- Stagonospora iridis** C. Mass. Causes a leaf spot disease on cultivated *Iris* in Great Britain and *I. florentina* in Tunis.
- Trabutia atroinquinans** (Wint.) Theiss. and Syd. On *Iris* sp. in Portugal.
- Trabutia molleriana** Cavara. On leaves of *Iris* sp. in Portugal.
- Uredo iridis** (Thuem.) Plowr. Forms narrow elliptical chestnut-colored rust pustules on both surfaces of leaves of *I. flavissima*. Said to be distinct from *Puccinia iridis*, which is present in the United States.
- ISATIS.** Herbs cultivated for ornament and for dyestuff.
- Aecidium isatidis** P. Har. Rust on leaves of *I. tinctoria* in France.
- Peronospora isatidis** Gäum. Downy mildew on leaves of *I. tinctoria* in Switzerland.
- ISCHNOSIPHON.** Calathea-like perennial herbs.
- Uredo ischnosiphonis** P. Henn. Powdery brown rust pustules on leaves of *I. leucophaeus* in Brazil.
- ISOPYRUM.** Annual or perennial dwarf stemless plants.
- Aecidium isopyri** Schroet. Rust on leaves of *I. thalictroides* in France, Switzerland, Belgium, and Germany.
- Cercospora isopyri** v. Hoeh. Irregular dark-brown areas on leaf blades and petioles and on stems of *I. thalictroides* in Austria.
- Peronospora parvula** Schneid. Downy mildew on leaves of *I. fumarioides* in Siberia.
- Triphragmium isopyri** Moug. and Nestl. Black rust pustules on leaf blades and petioles and on stems of *I. thalictroides* in Europe.
- IXIA. HYALIS.** Ag. Spring flowering cormous plants from south Africa with grasslike foliage.
- Bacillus ixiae** Sever. Appearing first as a browning and dying back of the tops of the plants, followed by a soft rot of the bulbs of *I. maculata* in Italy and Holland.
- Uromyces ixiae** (Lév.) Wint. Small yellow rust pustules, becoming brown on leaves of *Acidanthera exscapa*, *Ixia axillaris*, *I. coerulea*, *I. erecta*, *I. paniculata*, *I. patens*, *I. polystachya*, *Sparaxis grandiflora*, and *Watsonia densiflora* in south Africa and Australia.
- Uromyces zeyheri** Bub. Yellow to brown rust pustules on both leaf surfaces of *I. scillaris* in south Africa.
- IXIOLIRIUM.** Hardy spring-flowering bulbs.
- Aecidium ixiolirii** Komarov. Rust on leaves of *I. tataricum* in central Asia.
- Aecidium tataricum** Rostr. Rust on leaves of *I. tataricum* in Transcaasia.
- IXORA.** Woody plants with evergreen leaves and showy flower clusters.
- Aecidium ixorae** Arth. Rust on leaves of *I. ferrea* in Cuba.
- Colletotrichum ixorae** Griff. and Maubl. Large, pale anthracnose spots on leaves of *I. alba* in France.
- Endophyllum ixorae** Gäum. Leaf rust on *I. javanica* in Guam.
- Phyllachora ixorae** Theiss. and Syd. Shiny black stromata on leaves of *I. parvifolia* in India.
- Phyllosticta ixorae** Rangel. On leaves of *Ixora* sp. in Brazil.
- Stagonospora ixorae** Rangel. On leaves of *I. coccinea* and *Ixora* sp. (cult.) in Brazil.
- JACARANDA.** Trees and shrubs with handsome tubular flowers.
- Aecidium circinatum** Wint. Rust on leaves of *Jacaranda* sp. in Brazil.
- Aecidium jacarandae** P. Henn. Rust on leaves of *Jacaranda* sp. in Brazil.
- Aecidium puttiansianum** P. Henn. Rust on leaves of *Jacaranda* sp. in Brazil.
- Fomes lamaoensis** Murr. See Hevea.
- JACKSONIA.** Stiff, leafless shrubs and subshrubs.
- Cronartium jacksoniae** P. Henn. Brown rust pustules on stems of *Aotus villosum*, *Bossiaea cinerea*, *Gompholobius latifolius*, *Jacksonia scoparia*, and *Platylobium formosum* in Australia.
- Uromyces politus** (Berk.) McAlp. Bright-orange rust pustules, becoming brown to black, on dark-purple, elongated patches on stems of *Muehlenbeckia (Jacksonia) cunninghami* in Australia.
- JACQUEMONTIA.** Tropical climbing herbs.
- Aecidium jacquemontiae** E. and E. Rust on leaves of *J. pentantha* in Mexico.
- Uredo jacquemontiae** P. Henn. Powdery dull-yellow rust pustules on leaves of *Jacquemontia* sp. in New Guinea.
- Uromyces gemmatus** B. and C. Rust on leaves of *J. nodiflora* in Porto Rico, Cuba, Jamaica, and St. Croix.

**JACQUINIA.** Tropical American trees and shrubs.

**Phyllachora conspicua** Ferd. and Winge. Dull-black stromata on leaves of *J. armillaris* in the American Virgin Islands.

**Phyllachora inclusa** (B. and C.) Sacc. Small black stromata on leaves of *J. aurantiaca*, *J. berterii*, and *Jacquinia* sp. in Porto Rico, Guatemala, Nicaragua, and Paraguay.

**JAMBOSA.** See Caryophyllus.**JASMINUM.** JASMINE. JESSAMINE. Climbing or erect shrubs, with attractive, often fragrant, flowers.

**Accidium jasminicola** P. Henn. Rust distorting leaves and branches of *J. floribundum* in Abyssinia.

**Accidium longaense** P. Henn. Rust on yellow-brown leaf spots on *J. microphyllum* in the Union of South Africa.

**Blastospora butleri** Syd. Brown rust pustules on leaves of *J. malabaricum* in India.

**Laestadia jasminicola** (Desm.) Sacc. Black sunken leaf spots on *J. officinale* in France.

**Meliola jasminicola** P. Henn. Black superficial fungus patches on upper leaf surfaces of *J. auriculatum* and *J. sambac* in India and Indo-China.

**Mycosphaerella jasmini-officinalis** Siem. On leaves of *J. officinale* in Southern Russia.

**Phyllosticta jasminorum** Togn. On leaves of *J. officinale* in Italy.

**Phytophthora syringae** Kleb. See Syringa.

**Puccinia abyssinica** (P. Henn.) Syd. Brown rust pustules on circular to irregular sunken leaf spots on *J. abyssinicum* in Abyssinia.

**Puccinia chrysopogi** Barcl. See Andropogon.

**Puccinia exhauriens** Thuem. Cinnamon-brown rust pustules on leaves and branches of *J. flexile* and *J. tortuosum* in Ceylon and the Union of South Africa.

**Puccinia jasmini** DC. Dark-brown rust pustules on leaves and stems of *J. fruticans* in Europe and northern Africa.

**Puccinia zimmermanniana** P. Henn. Brown rust pustules on leaves of *J. mauritianum* in east Africa.

**Septoria aitchisoni** Syd. Circular dull-brown leaf spots with raised purple margins on *J. humile* in Afghanistan.

**Septoria jasmini** Roum. Circular, then confluent, brown leaf spots, becoming white with brown margins, on *Jasminum* sp. in France.

**Septoria sambac** Pass. Subcircular whitish leaf spots on *J. sambac* in Italy.

**Uromyces comedens** Syd. Yellow to dark-brown rust pustules on leaves of *J. pubescens* in India.

**Uromyces hobsoni** Vize. Brown rust pustules on leaves, stems, and flower stalks of *J. floribundum*, *J. grandiflorum*, and *Jasminum* sp. in India, Somali, and Abyssinia.

**JATROPHA.** PHYSIC NUT. Tropical trees and shrubs with milky juice.

**Cercospora ajrekari** Syd. Circular yellow-brown leaf spots on *J. nana* in India.

**Cercospora jatropharum** Speg. Definite circular leaf spots, concentrically zoned, on *J. macrocarpa* in Argentina.

**Cercospora peronosporioides** Speg. Dull-yellow indefinite leaf spots on *J. anisophylla* and *J. macrocarpa* in Argentina.

**Cercosporina jatrophicola** Speg. Subcircular to angular ashen-colored leaf spots with dull-brown margins on *J. macrocarpa* in Argentina.

**Uredo jatrophicola** Arth. Brown rust pustules on leaves of *J. curcas* and *J. gossypifolia* in Porto Rico, Cuba, and Dominican Republic.

**Uromyces jatrophae** Diet. and Holw. Powdery cinnamon to chestnut-brown rust pustules on leaves of *J. multifida* in Mexico.

**Uromyces jatrophicola** P. Henn. Powdery yellow to black rust pustules on leaves of *Jatropha* sp. in Brazil.

**JEFFERSONIA.** TWINLEAF. Attractive, hardy perennial herbs.

**Triphragmiopsis jeffersoniae** Naoumoff. Rust on leaf blades and petioles of *J. dubia* in Russia.

**JOANNESIA.** Tropical trees with milky juice, useful for medicinal, ornamental, and timber purposes.

**Puccinia joannesiae** P. Henn. Pale-brown to chestnut-brown rust pustules on leaves of *J. brasiliensis* in Brazil.

**Uredo maceiensis** P. Henn. Brown rust pustules on irregular dull-brown leaf spots and on stems of *J. princeps* in Central America.

**JUBAEA.** PALM. See Palmae.**JUGLANS.** WALNUT. Butternut. Nut and timber trees.

**Ascochyta juglandis** Bolts. Subcircular gray-brown leaf spots, the centers of which fall out. On *J. regia* in Switzerland, Yugoslavia, and Austria.

**Gummosis.** A gumming disease of walnut (*Juglans* spp.) trunks and branches has been reported from France and other parts of Europe. Several distinct diseases are probably included, some due to nonparasitic causes, others possibly to bacteria or fungi not present in the United States.

**Helicobasidium tanakae** Miy. See Morus.

**Hypoderma roseum** (Pers.) Fr. See Populus.

**Marsonia manshurica** Naoum. Numerous sub-circular ashen-white leaf spots on *J. manshurica* in Russia.

**Mycosphaerella saccardoana** Jaap. On old leaves of *J. regia* in Dalmatia.

**Phyllosticta juglandina** Sacc. Indefinite whitish leaf spots with dull-brown borders on *J. regia* in Italy and Portugal.

**Phyllosticta juglandis** (DC.) Sacc. Indefinite whitish leaf spots on *J. regia* in Russia, Italy, France, and Austria.

**Septoria epicarpis** Thuem. On epicarp of nuts of *J. regia* in Europe.

**Septoria letendreana** Sacc. On leaves of *J. regia* in France.

**Septoria nigromaculans** Thuem. On epicarp of nuts of *J. regia* in Austria.

**JUNCOIDES.** See Luzula.**JUNIPERUS.** JUNIPER. Redcedar. Coniferous ornamentals and timber trees.

**Asteridium juniperinum** Cke. On needles of *Juniperus* sp. in Great Britain.

**Clasterosporium glomerulosum** Sacc. On leaves of *J. communis* in Germany.

**Gymnosporangium amelanchieris** (DC.) Ed. Fisch. The aecial stage of this rust forms small galls on the lower leaf surfaces of *Amelanchier ovalis*. The telia stage occurs in indefinite swollen areas along the branches of *J. communis* and *J. nana* in Europe.

**Gymnosporangium koreaense** (P. Henn.) Jacks. The aecial stage occurs on leaves of *Cydonia japonica*, *C. oblonga*, and *Pyrus sinensis*, the telia on *J. chinensis* in Japan and Chosen. This rust has been introduced with nursery stock into Oregon and Connecticut.

**Gymnosporangium mespili** (DC.) Kern. Aecia on leaves and stems of *Cotoneaster tomentosa*, *C. vulgaris*, *Crataegus grandiflora*, *C. laciniata*, *C. monogyna*, *C. orientalis*, *C. oxyacantha*, *C. pinnatifida*, *C. tanacetifolia*, *Cydonia oblonga*, *Mespilus germanica*, and *Pyrus communis*. Telia dark chestnut-brown on fusiform enlargements on branches of *J. oxycedrus*, *J. phoenicea*, *J. sabina*, and *J. virginiana* in Europe and Asia.

**JUNIPERUS**—Continued.

- Gymnosporangium photiniae** (P. Henn.) Kern. The aecia occur on thickened discolored spots on the lower leaf surfaces of *Pourthiaea villosa*. The telia occur on fusiform enlargements on the branches of *J. chinensis* in Japan and Chosen. Reported from Connecticut on imported stock.
- Gymnosporangium sabiniae** (Dicks.) Wint. This rust produces swellings on the twigs, deformed fruit, and leaf spots on which the aecia occur. The aecial hosts are *Pyrus betulæfolia*, *P. communis*, *P. eleagnifolia*, *P. michauxii*, *P. nivalis*, *P. salicifolia*, and *P. tomentosa*. The dark-brown telia occur on swellings on twigs and branches of *Juniperus chinensis*, *J. japonica*, *J. oxycedrus*, *J. phoenicea*, *J. sabina*, *J. sphaerica*, *J. tripartita*, and *J. virginiana* in Europe.
- Gymnosporangium terminali-juniperinum** Ed. Fisch. Aecia in small groups on brown leaf spots on *Sorbus latifolia* and telia on *J. communis* in Europe.
- Gymnosporangium yamadæ** Miy. Aecia on leaves of *Malus* sp. (apple), *M. spectabilis* and *M. toringa* and the telia on *J. chinensis* in China and Japan.
- Herpotrichia mucilaginosa** Start. and Grev. On needles of *J. communis* in Sweden.
- Sarcosoma thwaitesii** (B. and Br.) Petch. Produces witches'-brooms on the stems and main branches of *J. bermudiana* in Ceylon.
- Stemphylium juniperinum** Karst. Kills the needles of *Juniperus* sp. in Finland.
- Venturia furcata** Fautr. On leaves of *J. sabina* in France.
- JUSSIAEA**. PRIMROSE WILLOW. Herbs.
- Puccinia jussiaeae** Speg. Cinnamon-brown rust pustules on leaves of *J. lanceolata* and *J. longifolia* in Argentina and Uruguay.
- Puccinia sphaeroidea** P. Henn. Brown rust pustules on stems and leaves of *Jussiaea* sp. in Japan.
- Septoria obsidionis** Speg. Small circular, often confluent, white leaf spots with broad purple borders on *J. longifolia* in Argentina.
- Septoria platensis** Speg. Small definite white leaf spots with purple margins on *J. longifolia* in Argentina.
- Septoria thomasiana** Sacc. Subcircular light-buff leaf spots with brown-purple margins on *J. acuminata* in the Portuguese Island of St. Thomas.
- JUSTICIA**. Herbs cultivated for their showy flowers.
- Aecidium acanthacearum** Cke. Rust on leaves of *J. procumbens* and *J. uncinulata* in Ceylon and Uganda.
- Peronospora wildemaniana** P. Henn. and var. **macrospora** K. Saw. Downy mildew attacking the leaves of *J. procumbens* in Japan and India, causing irregular light yellowish-green to white spots, which finally occupy the entire leaf area.
- Puccinia fuhrmanni** Mayor. Rust on leaves of *J. secunda* in Colombia.
- Puccinia peraffinis** Syd. Brown rust pustules on yellow-brown circular leaf spots on *J. diffusa* in India.
- Puccinia shiraiana** Syd. Dark-brown rust pustules on leaves and stems of *J. procumbens* in Japan and Ceylon.
- Puccinia thwaitesii** Berk. Dark-brown rust pustules on large sunken brown leaf spots on *J. gendarussa* in Ceylon, Indo-China, Java, Sumatra, Celebes, Malaya, New Guinea, and the Philippines.
- Uromyces kwangensis** P. Henn. Brown rust pustules on leaves of *Justicia* sp. in the Congo.
- Ustilago thwaitesii** Berk. A doubtful smut species reported on *J. gendarussa* in Malaya.
- KADSURA**. Tropical Asian woody climbers.
- Septoria kadsurae** Tassi. On leaves of *K. japonica* in Italy.
- KAGENECKIA**. Small evergreen trees from South America.
- Phyllachora negeriana** P. Henn. and Lind. Shiny black stromata on leaves of *K. oblonga* in Chile.
- KENNEDIA**. Woody trailers or twiners. See also Hardenbergia.
- Cercospora kennedyae** Cke. and Mass. Irregular and confluent cinnamon-brown leaf spots on *K. prostrata* in Australia.
- Phyllosticta kennedyae** Wint. Irregular, often confluent, dull-brown, then whitish, leaf spots on *K. ovata* in Portugal.
- Septoria kennedyae** Trav. and Spessa. On leaves of *Kennedia* sp. in Portugal.
- KENTIA**. See Palmae.
- KERRIA**. Shrubs.
- Helicobasidium tanakae** Miy. See Morus.
- Phomopsis striaeformis** Grove. On stems of *K. japonica* in Algeria.
- Septoria kerriae** Syd. Circular, pale-brown leaf spots on *K. japonica* in Japan.
- KIGELIA**. **KIGELKEIA** Ag. Ornamental African trees.
- Leptosphaeria dryadea** Sacc. On *K. pinnata* in the Philippines.
- Phyllosticta kigelia** Died. On leaves of *K. pinnata* in the Philippines.
- KIGELKEIA**. See Kigelia.
- KOCHIA**. SUMMER CYPRESS. MOCK CYPRESS. Herbs.
- Puccinia kockiae** Mass. Golden-brown to dark-brown rust pustules on leaves of *K. sedifolia*, *K. villosa*, and *Enchylaena tomentosa* in Australia.
- Uromyces kochiae** Syd. Rust on leaves of *K. prostrata* in Russia.
- KOELREUTERIA**. GOLDEN-RAIN TREE. Ornamental trees grown for their panicles of yellow flowers and handsome foliage.
- Phyllosticta koelreuteriae** Hóll. Oblong ashen leaf spots with brown borders on *K. paniculata* in Hungary.
- Triphragmium koelreuteriae** Syd. Leaf rust on *K. paniculata* in Japan.
- Uncinula koelreuteriae** Miy. Powdery mildew on leaves of *K. bipinnata* in China.
- KRAUNHIA**. See Wistaria.
- LABURNUM**. GOLDEN CHAIN. Ornamental trees or shrubs, grown for their showy racemes of yellow flowers. See also Cytisus.
- Microsphaera guarinonii** Br. and Cav. Powdery mildew on leaves of *L. anagyroides* (*L. vulgare*) in Italy.
- Peronospora cytisi** Magn. See Cytisus.
- Septoria cytisi** Desm. Numerous subcircular whitish to brown leaf spots on *L. anagyroides* in Italy, France, and Hungary.
- Stereum purpureum** Pers. See Prunus.
- Uromyces genistae-tinctoriae** (Pers.) Fckl. See Caragana.
- LACTUCA**. LETTUCE. Hardy annual or perennial herbs, several species grown as vegetables; others weedy.
- Ascochyta lactucae** Rostr. On leaves and stems of *L. sativa* in Denmark.
- Cercospora lactucae** P. Henn. Dark areas on leaves of *L. raddeana* in Japan.
- Cercospora longissima** Trav. (*C. lactucae* Stevenson) (*C. lactucae* Welles). Circular to angular, often confluent, whitish leaf spots with rufous or brown margins on *L. sativa* in Italy, Brazil, Porto Rico, Cuba, China, and the Philippines.

**LACTUCA**—Continued.

- Puccinia lactucae** Diet. Brown rust pustules on leaves of *L. albiflora*, *L. brevirostris*, *L. denticulata*, *L. laciniata*, *L. raddeana*, *L. squarrosa*, and *L. thunbergiana* in Japan.
- Puccinia lactucae-debilis** Diet. Dull-brown powdery rust pustule on leaves of *L. debilis* and *L. stolonifera* in Japan.
- Puccinia lactucae-denticulatae** Diet. Powdery cinnamon-brown to chestnut-brown rust pustules on leaves of *L. denticulata* and *L. stolonifera* in Japan.
- Puccinia lactucae-repentis** Miy. Rust on leaves of *L. repens* in Japan.
- Puccinia lactucina** Syd. Yellow-brown to brown rust pustules on leaves of *L. scarioloidea* in Persia.
- Puccinia lactucarum** Syd. Brown rust pustules on leaves and stems of *L. altissima*, *L. perennis*, *L. quercina*, *L. sagittata*, and *L. scariola* in Europe.
- Puccinia prenanthis** (Pers.) Lindr. Orange aecia and brown uredinia and telia on leaf blades and petioles of *L. muralis*, *L. sativa*, *L. stricta*, *L. virosa*, and *Prenanthes* spp. in Alaska, Europe, and Australia.
- Ramularia lactucae** Jaap. On leaves of *L. muralis* in Yugoslavia.
- Ramularia lactucosa** Lamb. and Fautr. Numerous small circular leaf spots on *L. sativa* in France.
- Ramularia lumpsanae** (Desm.) Sacc. var. **lactucae-muralis** Jaap. Leaf spots on *L. muralis* in Germany.
- Septoria fernandezii** Unam. On leaves of *L. virosa* in Spain.
- Uredo proximella** Arth. Brown rust pustules on leaves of *L. intybacea* in Porto Rico, Cuba, and Dominican Republic.
- LAELIA**. Epiphytic orchids. See Orchidaceae.
- LAGENARIA**. GOURD. Cucurbitaceous vines.
- Phyllosticta lagenariae** Pass. On leaves of *L. leucantha* (*L. vulgaris*) in Italy.
- Phyllostictiella cucurbitacearum** Tass. On leaves of *L. leucantha* (*L. vulgaris*) in Italy.
- LAGENOPHORA**. Small perennial scapose herbs.
- Puccinia lagenophorae** Cke. Brown rust pustules on leaves of *L. billardieri* in Australia.
- Septoria lagenophorae** McAlp. Circular to irregular ashen to dull-brown leaf spots on *L. billardieri* in Australia.
- Thecaphora lagenophorae** McAlp. Smut sori developing in the ovaries, converting entire flower heads into powdery, brown spore masses. On *L. emphyosopus* in Australia.
- LAGERSTROEMIA**. CRAPE MYRTLE. Showy-flowered shrubs and trees.
- Cercospora lagerstroemiae** Syd. Circular to irregular gray-brown to white leaf spots on *L. speciosa* in the Philippines.
- Mollisia ravida** Syd. Effuse, pale, indefinite leaf spots on *L. speciosa* in the Philippines.
- Rhizisma lagerstroemiae** Rabh. Irregular black stromata on leaves of *L. indica*, *L. lanceolata*, and *L. speciosa* in India and the Philippines.
- Septoria lagerstroemia** Sacc. and Scalia. Brown leaf spots on *L. indica* in Portugal.
- Uncinula australiana** McAlp. Powdery mildew on leaves and inflorescences of *L. indica* and *L. ovalifolia* in Japan and Australia.
- LAGURUS**. RABBIT TAIL GRASS. Hardy annual grasses.
- Puccinia laguri** Jaap. Rust on leaves of *L. ovatus* in Dalmatia.
- Septoria caruaniana** Sacc. Oblong pale ochraceous spots on leaves of *L. ovatus* in Malta.
- LAMIUM**. DEAD NETTLE. Annual and perennial herbs, some species cultivated as hardy border plants.
- Ascochyta lamiorum** Sacc. Ochraceous leaf spots on *L. album* in Italy.
- Ovularia lamii** (Fekl.) Sacc. On leaves of *L. album*, *L. amplexicaule*, and *L. purpureum* in Russia and Denmark.
- Peronospora lamii** A. Braun. Broadly effused dingy lilac, downy mildew patches on under leaf surfaces of *L. album*, *L. amplexicaule*, *L. maculatum*, *L. purpureum*, and *L. rubrum* in Europe and Tunis. Reported from New York.
- Phyllosticta albanica** Bub. Circular to irregular gray leaf spots with brown margins on *L. galeobdolon* in Yugoslavia.
- Phyllosticta lamii** Sacc. Whitish leaf spots with broad ochraceous surrounding zones on *L. album* and *L. ovata* in Siberia and Italy.
- Phyllosticta venziana** Mart. Small circular gray leaf spots on *Lamium* sp. in Italy.
- Ramularia exilis** Syd. Indefinite yellowish leaf spots on *L. galeobdolon* in Denmark and Germany.
- Ramularia lamiiicola** C. Mass. Angular, then confluent, leaf spots on *L. album* in Europe.
- Septoria diedickei** Sacc. Circular to irregular, then confluent, white leaf spots with brown or dark-purple margins on *L. galeobdolon* in Denmark and Germany.
- Septoria lamiiicola** Sacc. and var. **intermedia** C. Mass. Whitish leaf spots with red margins on *L. album*, *L. amplexicaule*, *L. bithynicum*, *L. maculatum*, *L. ovata*, and *L. purpureum* in Europe.
- LANDOLPHIA**. PACURIA. Ag. Tropical shrubs mostly climbers.
- Colletotrichum paucipilum** Delacr. Brown anthracnose spots, often marginal, on leaves of *L. kleinii* in France and the Congo.
- Phyllosticta landolphiae** P. Henn. Marginal brown leaf spots on *L. kirkii* in Germany.
- Phyllosticta owariensis** Maubl. Small irregular whitish leaf spots with brown borders on *L. owariensis* in central Africa.
- LANSIUM**. LANSIA. Oriental trees, one species cultivated for its edible fruit.
- Bacterium citri** Hasse. See Citrus.
- Corticium salmonicolor** B. and Br. See Citrus.
- LANTANA**. Shrubs, some species cultivated for the verbena-like flowers.
- Acanthostigma lantanae** Theiss. On leaves of *Lantana* sp. in Brazil.
- Aecidium lantanae** Mayor. Rust on leaves of *L. hispida* in Colombia.
- Phyllachora sororcula** Speg. Shiny black stromata on leaves of *Lantana* sp. in Paraguay and Brazil.
- Phyllosticta lantanae** Pass. Small irregular whitish leaf spots on *L. camara* and *Viburnum lantana* in Porto Rico and Italy.
- Prospodium tuberculatum** (Speg.) Arth. Powdery cinnamon-brown to black rust pustules on leaves of *L. camara*, *L. horrida*, *L. involucrata*, *L. lilacina*, *L. robusta*, *L. salvifolia*, and *L. tiliaefolia* in Mexico, South and Central America, and Cuba.
- Puccinia natalensis** Diet. and Syd. Dark-brown rust pustules on leaves of *L. salvifolia* and *L. trifolia* in Uganda and the Union of South Africa.
- Puccinia schimperiana** Syd. Rust on leaves of *L. citrifolia* in Abyssinia.
- Septoria lantanae** Gar. Small, irregular leaf spots on *L. camara* in Porto Rico.
- Uromyces dubiosus** P. Henn. Dark-brown powdery rust pustules on leaves of *Lantana* sp. in Brazil.
- LAPAGERIA**. CHILE BELLS. Half-hardy evergreen twiner.
- Phyllosticta jaffueli** Speg. On leaves of *L. rosea* in Chile.

**LAPEYROUSIA.** African summer-flowering bulbs.

**Uromyces anomathecae** Cke. Brown elliptical rust pustules on both leaf surfaces of *L. cruenta* in Natal.

**Uromyces delagoënsis** Bub. Round to oblong golden-yellow to black rust pustules on both surfaces of leaves of *L. delagoënsis* in Mozambique and Portuguese East Africa.

**LAPORTEA.** Perennial herbs, shrubs or trees.

**Aecidium laporteeae** P. Henn. Rust pustules on circular yellow-brown leaf spots on *L. bulbifera* in Japan.

**LARDIZABALA.** Evergreen climbers of Chile.

**Physalospora lardizabala** Speg. On leaves of *L. biternata* in Chile.

**Septoria lardizabala** P. Henn. Effuse brown leaf spots on *L. biternata* in Germany.

**LARIX.** LARCH. TAMARACK. Timber and ornamental trees.

**Cercospora exosporioides** Bub. On needles of *L. decidua* (*L. europaea*) in Bohemia.

**Dasyscypha calycina** (Schum.) Fekl. (*D. willkommii* Hartig.) This fungus causes the most serious disease of the larch known and is widespread throughout Europe. It has also been reported from Newfoundland, and there is grave danger of the introduction of the parasitic strain of the fungus into the United States. The fungus occurs commonly on dead branches, the fruiting bodies appearing as small orange or yellow cups. It attacks living parts of the tree through wounds, and develops large, perennial cankers. The bark becomes black, resin oozes out and a flattened cavity forms, which increases year by year. Young trees may be girdled and killed, older trees persist often for years, but their merchantable value is lowered or destroyed, particularly when several or more cankers occur along the trunk. The hosts are *L. decidua*, *L. occidentalis*, *Abies pectinata*, *Pinus austriaca*, *P. laricio*, *P. pumilo*, and *P. sylvestris*.

**Exosporina laricis** Oud. Very injurious to leaves of *Larix* sp. in Holland.

**Hartigella laricis** (Hart.) Syd. (*Meria laricis* Vuill.) Brown spots on needles, particularly those on young shoots of *Larix* sp. in Russia, France, and Germany. Leaf fall results, and often the death of young trees.

**Heterosporium laricis** Cke. and Mass. On leaves of *L. decidua* in Great Britain.

**Hypodermella laricis** Tub. On needles of *L. decidua* in Switzerland and Germany, causing a dwarfing of shoots and dying of foliage.

**Melampsora larici-capraearum** Kleb. See Salix.

**Melampsora larici-epitea** Kleb. See Salix.

**Melampsora larici-pentandrae** Kleb. See Salix.

**Melampsora larici-populina** Kleb. See Populus.

**Melampsora laricis** R. Hart. See Populus.

**Melampsora larici-urbaniana** Mat. See Salix.

**Melampsorium betulae** (Schum.) Arth. This rust attacks the needles of *L. europaea* in Europe, Siberia, and Japan. The alternate stages occur on *Betula* spp. and have been reported from the United States.

**Mycosphaerella laricina** R. Hartig. Brown patches on needles of *L. decidua* in Germany, causing premature leaf fall.

**Mycosporium abietinum** Rostr. On bark of *L. decidua*, *Picea sitchensis*, *Pinus strobus*, and *Pseudotsuga taxifolia* in Denmark.

**Phoma laricis** Oud. On leaves of *L. decidua* in Denmark.

**Phoma lineolata** Desm. On branches and cone scales of *L. decidua* in Europe.

**Phomopsis pseudotsugae** Wils. See Pseudotsuga.

**Phyllosticta lineolata** Desm. On *L. decidua* in Germany.

**Phytophthora fagi** R. Hart. See Fagus.

**LASIA.** Perennial herbs, woody at the base.

**Mycosphaerella lasiana** Sacc. On leaves of *L. heterophylla* in Malaya.

**LATANIA.** Palm. See Palmae.

**LATHYRUS.** (Including *Orobus*). PEA. Sweet pea. Annual and perennial climbing or upright herbs and shrubs.

**Ascochyta lathyri** Trail. and var. **lathyri-odorati** Bub. and Kab. Large brown areas on leaves of *L. annuus*, *L. nissolia*, *L. odoratus*, and *L. sylvestris* in Scotland, Denmark, Bohemia, Italy, and Spain.

**Ascochyta orobi** Sacc. Irregular whitish leaf spots with dull-brown margins on *L. niger*, *L. (Orobus) vernus*, *Orobus lathyroides*, and *O. vernus* in Siberia, Russia, Italy, and Austria.

**Cladosporium album** Dowson. Irregular sunken faint-buff spots on both leaf surfaces and wings of stems of *L. odoratus* in Great Britain.

**Cylindrosporium lathyri** Bub. and Kab. Irregularly circular to angular ochraceous leaf spots with brown-purple margins on *L. vernus* in Italy and Austria.

**Cylindrosporium orbicolum** (Sacc.) Bub. Large indefinite brown-margined leaf spots on *L. variegatus*, *Orobus venetus*, and *O. vernus* in Russia, Bohemia, and Italy.

**Entyloma feurichii** Krieg. Smut sori in confluent brown leaf spots on *L. sylvestris* in Germany.

**Gloeosporium orobi** Karst. Irregular dull-brown leaf spots on *Orobus vernus* in Finland.

**Isariopsis carnea** Oud. Leaf spot on *L. pratensis* in Italy and Holland.

**Mycosphaerella lathyri** A. Pot. Indefinite irregular ochraceous leaf spots on *L. pisiformis* in Siberia and Russia.

**Ovularia deusta** (Fekl.) Sacc. Large dark-brown areas on leaves of *L. pratensis*, *L. tuberosus*, and *Tetragonolobus siliquosus* in Siberia, Sweden, Italy, France, and Germany.

**Peronospora fulva** Syd. Downy mildew on leaves of *L. pratensis* and *L. vernus* in central and northern Europe.

**Peronospora lathyri palustris** Gäum. Downy mildew on leaves of *L. heterophyllus*, *L. palustris*, and *L. sativus* in Sweden, Switzerland, and India.

**Peronospora orobi** Gäum. Downy mildew on leaves of *L. montanus* and *L. tuberosus* in central and northern Europe.

**Peronospora senneniana** Sacc. Downy mildew on leaves of *L. nigrus* in Spain, Switzerland, Bohemia, Denmark, Norway, Austria, Poland, and Germany.

**Phyllachora lathyri** (Lév.) Theiss. and Syd. Black, crust-like stromata on both leaf surfaces of *L. grandiflorus*, *L. latifolius*, *L. pratensis*, *L. tuberosus*, *Onobrychis sativa*, *O. viciaefolia*, and *Vicia tenuifolia* in Siberia, Russia, Asia Minor, and Austria.

**Phyllosticta lathyricola** Bub. and Krieg. Small circular to irregular dull purple-brown leaf spots on *L. sylvestris* in Germany.

**Phyllosticta lathyrina** Sacc. and Wint. Pale ochraceous leaf spots with brown margins on leaves of *L. sylvestris* in Russia, Italy, and Switzerland.

**Phyllosticta orobella** Sacc. On leaves of *L. vernus* in Russia and Italy.

**Physalospora lathyri** (D. and M.) Sacc. On leaves of *L. amphicarpus* in Algeria.

**Placosphacteria onobrychidis** Sacc. See Lupinus.

**Ramularia lathyri** Höll. Oblong yellowish leaf spots on *L. hirsutus* in Hungary.

**LATHYRUS**—Continued.

- Rhizoctonia napi** West. See Brassica.  
**Rhytisma onobrychis** Sacc. See Lupinus.  
**Septogloeum lathyri** Lind. Large confluent pale-reddish leaf spots on leaves of *L. sylvestris* in Denmark.  
**Septoria fatreyana** Sacc. On leaves of *L. sylvestris* in Italy.  
**Septoria fulvescens** Sacc. Brown confluent leaf spots on *L. maritimus*, *L. pisiformis* and *L. sylvestris* in Russia, Bohemia, Italy, and Germany.  
**Septoria podgoricensis** Bub. Circular to elliptical yellowish spots on leaves and stems of *L. angulatus* and *L. annuus* in Spain and Yugoslavia.  
**Septoria silvestris** Pass. Leaf spots on *L. pratensis*, *L. sylvestris*, *L. tuberosus* and *Vicia sepium* in Sweden, Russia, Denmark, and Italy.  
**Septoria stipularis** Pass. See Lotus.  
**Uromyces clavatus** Diet. Yellow and cinnamon-brown rust pustules on leaves of *L. magellanicus*, *L. multiceps*, and *Vicia tenuifolia* in Chile, Argentina, and Brazil.  
**Uromyces lathyrinus** Speg. Yellow and dark-brown rust pustules on leaves and stems of *L. clymenum* in Argentina.  
**Uromyces orobi** (Pers.) Lévy. Brown rust pustules on *L. cicera*, *L. latifolius*, *L. montanus*, *Orobis tuberosus*, and *Vicia unijuga* in Siberia and Europe.  
**Uromyces pisi** (Pers.) De By. See Pisum.
- LAUREL. LAUREL. SWEET BAY. Trees.**  
**Ascochyta laurina** Tass. Marginal white leaf spots with rufous margins on *L. mollis* in Italy.  
**Cercospora unicolor** Sacc. and Penz. On leaves of *L. nobilis* in France.  
**Exobasidium lauri** Geyl. Long sulcate excrescences on trunks of *L. canariensis* and *L. nobilis* in Portugal, Italy, Brazil, and the Canary Islands.  
**Gleosporium nobile** Sacc. Circular dull-yellowish anthracnose spots on leaves of *L. nobilis* in Italy and Austria.  
**Metasphaeria nobilis** Sacc. Whitish leaf spots with dull-brown margins on *L. nobilis* in Italy and Portugal.  
**Phragmonaevia lauri** Pat. Circular, then confluent, rufous to whitish leaf spots with brown margins on *L. nobilis* in Tunis.  
**Phyllosticta allantella** Sacc. On leaves of *L. nobilis* in Spain and France.  
**Phyllosticta laurina** d'Alm. Irregular dull-brown leaf spots with darker borders on *L. nobilis* in Portugal.
- LAVANDULA. LAVENDER. Perennial herbs, subshrubs, and shrubs.**  
**Phoma lavandulae** Brierley. Diseased buds and shoots turn brown, the leaves wither and fall and the epidermis of the twigs splits away in shreds. On *L. spica* (*L. officinalis*) in Great Britain.  
**Septoria lavandulae** Desm. Circular to irregular white leaf spots with raised purple margins on *L. spica*, and *L. stoechas* in Great Britain, France, Italy, Denmark, and Madeira.
- LAVATERA. TREEMALLOW. Annual and perennial flower-garden herbs and some ornamental shrubs.**  
**Ramularia daniloi** Bule. Angular yellowish leaf spots becoming white on *L. rotundata* and *L. thuringiaca* in Spain and Montenegro.
- LEDUM. LABRADOR TEA. Shrubs cultivated for their white flowers and evergreen foliage.**  
**Ascochyta ledicola** Oud. On leaves of *L. palustre* in Holland.  
**Coccomyces ledi** Rehm. On living branches of *L. palustre* in Sweden.  
**Phyllosticta ledi** Rostr. White to gray leaf spots with purple borders on *L. groenlandicum* in Greenland.  
**Sclerotinia heteroica** Wor. A heteroecious species, one stage attacking the flowers and ovaries of *L. palustre*, the other shoots of *Vaccinium uliginosum* in Finland, Sweden, Russia, and Germany.
- LENS. LENTIL. Herbs.**  
**Peronospora lentis** Gäum. Downy mildew on leaves of *L. esculenta* in Germany.  
**Rhizoctonia napi** West. See Brassica.  
**Uredo lentis** Lagerh. Dark brown, elongate rust sori on leaves and stems of *L. esculenta* in Ecuador.  
**Uromyces viciae-craccae** Const. See Vicia.
- LEONOTIS. LION'S-EAR. Shrubs cultivated for their scarlet or orange flowers.**  
**Accidium leonotidis** P. Henn. Rust on leaves of *L. velutina* in central Africa.  
**Puccinia leonotidicola** P. Henn. Brown rust pustules on leaves of *L. nepetifolia* in Porto Rico, Cuba, Bahamas, Jamaica, central and southern Africa.  
**Uredo cancerina** P. Henn. A rust producing large cancerous swellings on the stems, which break irregularly and finally become woody. On *L. velutina* in Abyssinia.  
**Uredo leonoticola** P. Henn. Rust on leaves of *Leonotis* sp. in Brazil.
- LEONTODON. See Taraxacum.**  
**LEONTOPIDIUM. EDELWEISS. Perennial herbs.**  
**Puccinia leontopodii** Vogl. Rust on leaves of *L. alpinum* in Switzerland.
- LEPIDAGATHIS. Tropical herbs and shrubs.**  
**Accidium lepidagathis** Syd. Rust on brown leaf spots on *L. hyalina* in India.
- LEPTODERMIS. Low shrubs with small white or purple flowers.**  
**Coleosporium leptodermidis** (Barel.) Syd. Golden rust pustules on leaves of *L. lanceolata* in India.
- LEPTOSPERMUM. AGONIS Ag. Ornamental woody plants grown for their flowers and myrtlelike foliage.**  
**Phylachora callistemoni** Rodw. Tar spot on leaves of *P. scoparium* in Tasmania.
- LESCHENAULTIA. Herbs and heathlike Australian plants.**  
**Puccinia gilgiana** P. Henn. Yellow and brown to black rust pustules on petioles, stems, and calyces of *L. linarioides* in Australia.
- LESPEDEZA. BUSH CLOVER. Forage plants.**  
**Uromyces rugulosus** Pat. Black rust pustules on leaves of *L. yunnanensis* in China.  
**Woroninella vulcanica** Wor. On *L. cytoidis* in Java.
- LEUCADENDRON. SILVER TREE. Shrubs or trees, the leaves densely covered with white silky hairs.**  
**Cercospora protearum** Cke. Circular dark-brown leaf spots on *L. argenteum* in the Union of South Africa.
- LEUCAENA. Shrubs with Acacialike foliage.**  
**Exosporium leucaenae** Stev. and Dal. Circular dull-yellow leaf spots on *L. glauca* in Porto Rico.  
**Ravenelia leucaenae** Long. Brown powdery rust sori on leaves of *L. diversifolia* and *L. esculenta* in Mexico.  
**Ravenelia leucaenae-microphyllae** Diet. Brown rust pustules on leaves of *L. angustissima* and *L. microphylla* in Mexico and Guatemala.  
**Ravenelia verrucosa** Cke. and Ell. Yellow-brown to brown rust pustules on leaves of *L. lanceolata* in Mexico.
- LEUCANTHEMUM. See Chrysanthemum.**

**LEUCOJUM.** SNOWFLAKE. Hardy bulbous plants.

**Puccinia schmidiana** Diet. See Phalaris.

**Ramularia ucrainica** Petr. Brown to gray-brown leaf spots on *L. vernum* in southeastern Europe.

**Septoria malisorica** Bub. Elongate (up to 3 centimeters) gray-purple leaf spots with brown margins on *L. aestivum* in Yugoslavia.

**LEUCOTHOE.** Ornamental shrubs grown for their handsome evergreen leaves and white flowers.

**Coniothyrium leucothoës** P. Henn. Circular, then confluent, whitish leaf spots with brown margins on *Leucothoë* sp. in Brazil.

**Exobasidium leucothoës** P. Henn. Deforms leaves and branches of *Leucothoë* sp. in Brazil.

**Rhytisma leucothoës** P. Henn. Black stromata on leaves of *Leucothoë* sp. in Brazil.

**LEUZE.** See Centaurea.

**LEVISTICUM.** LOVAGE. Perennials grown for their aromatic seed.

**Bacterium levistici** Osterw. Small, somewhat sunken, brown spots which are confluent, forming large irregular areas with yellow margins. On leaf blades and petioles and stems of *L. officinale* in Germany.

**Puccinia bornmülleri** P. Magn. Powdery dark-brown rust pustules on leaf blades and petioles and stems of *L. persicum* in Persia.

**Ramularia levistici** Oud. Leaf spots on *L. officinale* in Holland.

**Ramularia schroeteri** P. Syd. Subcircular ochraceous leaf spots with whitish margins on *L. officinale* in Yugoslavia and Austria.

**Ramularia vestergreniana** Allesch. Large brown marginal leaf spots on *L. officinale* in Russia and Finland.

**Septoria levistici** West. Circular to irregular broad brown, then whitish, leaf spots on *L. officinale* in Belgium and Spain.

**LEYCESTERIA.** HIMALAYA HONEYSUCKLE. Small Chinese shrubs with small flowers in whorls.

**Septoria leycesteriae** Sacc. and Speg. Subcircular to irregular ochraceous leaf spots with brown borders on *L. formosa* in Italy and Germany.

**LICUALA.** Dwarf fan palms. See Palmae.

**LIGHTFOOTIA.** African shrubs or herbs with small white, blue, or pink flowers.

**Uredo campanularum** Cke. and Mass. Rust on leaves of *Lightfootia* sp. in Madagascar.

**LIGULARIA.** Perennial herbs.

**Aecidium nikkense** P. Henn. and Shir. Yellow rust pustules on leaves of *L. stenocephala* in Japan.

**Coleosporium ligulariae** Thuem. Golden rust pustules on leaves of *L. sibirica* in Russia, Finland, Siberia, and Denmark.

**Puccinia paulsenii** Syd. Brown leaf rust on *L. altaica* in central Asia.

**LIGUSTICUM.** Hardy herbs used for naturalizing.

**Puccinia aphanicondra** Lindr. Powdery dark-brown rust pustules on *L. alatum* in Armenia.

**Puccinia ligusticicola** S. Miy. Brown to black rust pustules on leaf blades and petioles of *L. acutilobum*, *L. ibukiensis*, and *L. scoticum* in Japan.

**LIGUSTRUM.** PRIVET. Ornamental woody plants cultivated for their handsome foliage and as hedge plants.

**Aecidium klugkistianum** Diet. Rust on leaves of *L. japonicum* in Japan.

**Ascochyta ligustri** Sacc. and Speg. (*Ascochyta ligustrina* [Pass.] Tass.) Dull-yellow leaf spots on *L. vulgare* in Italy and France.

**Cercospora ligustri** Roum. Numerous circular to irregular yellow leaf spots on *L. vulgare* in France.

**Phyllosticta ligustri** Sacc. Leaf spots on *L. vulgare* in Denmark, Italy, Portugal, and Austria.

**Phyllosticta ligustrina** Sacc. and Speg. Leaf spot on *L. vulgare* in France and Italy.

**Phytophthora syringae** Kleb. See Syringa.

**Puccinia obtusata** (Oth.) Ed. Fisch. See Phragmites.

**Ramularia ligustrina** Maubl. Circular to oval dull-yellow leaf spots with dark-brown margins on *Ligustrum* sp. in France.

**Septoria japonica** Thuem. On leaves of *L. japonicum* and *L. vulgare* in Italy and Germany.

**Septoria ligustri** (Desm.) Kicks. Pale ochraceous irregular leaf spots with reddish margins on *L. vulgare* in Belgium and France.

**Septoria ligustrina** Sacc. On leaves of *L. vulgare* in Bohemia.

**LILIUM.** Lily. Scaly bulbous plants with showy flowers.

**Aecidium lillii-cordifolii** Diet. Circular to irregular spots are produced on both leaf surfaces up to 1 centimeter in diameter, in which yellow rust pustules appear. On *L. cordifolium* in Japan.

**Aecidium saffianoffianum** Thuem. Yellow rust pustules appear on leaves of *L. martagon* in Siberia.

**Bacillus lillii** Uyda. This bacterial disease attacks the leaves and bulb scales. The spots are deep brown in color, oval to elongate or more or less irregular on the bulb scales. The entire leaf blade soon becomes involved, turns brown, and droops. When young bulbs are attacked flowering shoots fail to develop. On *L. auratum*, *L. cordifolium*, *L. elegans*, *L. hansonii*, *L. longiflorum*, *L. meaeoloides*, *L. rownii*, *L. speciosum*, *L. tigrinum*, and *L. venustum* in Japan.

**Botrytis lillorum** Fujikuro. This fungus attacks mostly the leaves, first appearing as small spots about 1 millimeter in diameter, rapidly spreading over the entire surface, causing total decay of the host plant. "As bad as any three other lily diseases in Japan." On *L. longiflorum* in Japan.

**Cercospora lilliiicola** (Rich.) Sacc. On stems of *L. candidum* in France.

**Cylindrosporium inconspicuum** Wint. (*Cercospora lillii inconspicuum* [Wint.] v. Hoeh.) (*Cercospora hungarica* Baeume.) Causes irregular, often confluent, spots on the leaves which are dark-brown, becoming paler at the center. On *L. martagon* and other species in France, Switzerland, and Austria.

**Mycosphaerella cixia** Sacc. (*Sphaeria matorina* Sacc.) Causes a leaf blight of *L. candidum* in Italy.

**Phyllosticta lilliiicola** Sacc. A leaf-spot disease of *L. candidum* in Italy.

**Puccinia** sp. Intercepted on *Lilium* sp. from Ecuador.

**Rhizopus necans** Mass. A wound parasite, causing a soft rot of the bulbs, particularly during shipment. On *L. auratum*, *L. lancifolium*, *L. speciosum*, and other species in Japan, Great Britain, and Denmark.

**Uromyces aecidiiformis** (Strauss.) Rees. (*Uromyces lillii* Kunze.) This rust produces yellow-brown to chestnut-brown rust pustules on dead spots on leaves, petioles, and stems. On *L. bulbiferum*, *L. candidum*, *L. carniolicum*, and *L. croceum* throughout central Europe and on *Fritillaria involucreta* in France.

**Uromyces holwayi** Lagerh. Leaf rust on *L. dauricum* and *L. maximowiczii* in Japan.

**Uromyces japonicus** Syd. Cinnamon-colored powdery rust pustules, later becoming darker on the upper surfaces of leaves of *L. maximowiczii* and *L. tigrinum* in Japan.

**Uromyces lillii** (Lk.) Fekl. Brown rust pustules are produced in sunken spots on both surfaces of the leaves of *Fritillaria delphinensis*, *F. graeca*, *F. meleagris*, *F. messanensis*, *F. minor*, *F. ruthenica*, *Lilium auratum*, *L. bulbiferum*, *L. canadense*, *L. candidum*, *L. carniolicum*, and *L. croceum* in France, Switzerland, Italy, Yugoslavia, Russia, Montenegro, Spain, Japan, Austria, Hungary, Dalmatia, and Germany. Reported from Washington.

**LINARIA.** TOADFLAX. Low herbs.

**Contractia cingens** (Berk.) De T. Smut sori on stems and leaves of *L. genistifolia* and *L. vulgaris* in Austria.

**Peronospora fiava** Gäum. Downy mildew on leaves of *L. vulgaris* in central and north Europe.

**Peronospora linariae** Fckl. Downy mildew on leaves of *L. arvensis* and *L. minor* in central and north Europe.

**Phyllosticta linariae** Sacc. See Antirrhinum.

**Puccinia trabutii** Roum. and Sacc. See Phragmites.

**Septoria cymbalariae** Sacc. and Speg. Circular dull-brown leaf spots on *L. cymbalaria* and *L. nivea* in Italy, France, and Spain.

**Septoria gandulphi** Sacc. and D. Sacc. Small orbicular to angular leaf spots on *L. commutata* in Italy.

**Septoria lapadensis** Jaap. Yellow-brown leaf spots on *L. commutata* in Dalmatia.

**Ustilago hypogaea** Tul. Doubtful smut species on *L. spuria* in the Isle of Wight.

**LINUM.** FLAX. Fiber plant. Several species cultivated as ornamentals.

**Ascochyta lini** Rostr. On stems of *L. catharticum* in Denmark.

**Cercospora lini** Wor. On leaves of *L. nervosum* in Russia.

**Colletotrichum lini** (Westerd.) Tochnal. (*C. linicolum* Pethybr. and Laff.) Irregular, shrunken, discolored cankers occur on seedlings of *L. usitatissimum*, involving the cotyledons and causing damping off. Similar dark rusty-brown areas occur on stems, leaves, calyces, buds, and capsules of older plants. The disease is carried in the seed. It has been reported from Ireland, Great Britain, Japan, and Formosa. It is not clear whether the flax anthracnose reported from the United States is identical.

**Cylindrosporium melitense** Sacc. On leaves and flowers of *L. strictum* in Italy.

**Entyloma lini** Oud. Smut sori on yellowish circular leaf spots on *L. usitatissimum* in Holland.

**Fusicladium lini** Sor. On leaves of *L. usitatissimum* in Belgium and Germany.

**Olpidiaster radices** (Willd.) Pascher. (*Asterocystis radices* Willd.) This fungus attacks the roots of *L. usitatissimum*, *Brassica* spp., *Plantago* spp., and *Veronica* spp. in Europe, Argentina, and Morocco, causing a yellowing and stunting of the plants.

**Peronospora lini** Schroet. Downy mildew on leaves of *L. catharticum* in Scandinavia, Finland, Russia, Austria, Switzerland, and Germany.

**Phlyctaena linicola** Speg. Yellowish areas on leaves and stems of *L. usitatissimum* in Argentina. Recently reported from several localities in the United States as a serious disease.

**Phoma** sp. On stems of *L. usitatissimum* in Ireland.

**Pleosphaerulina lini** Lebed. On leaves and stems of *L. perenne* in Russia.

**Polyspora lini** Pethybr. The browning or stem-break disease of flax (*L. usitatissimum*) is serious in Ireland and is a seed-borne disease. The cotyledons are first attacked, the disease spreading successively to leaves, stems, branches, and seed bolls. Stem lesions may cause half-grown plants to break over and die. On older plants numerous brown spots cover the various organs, diseased areas in a field taking on a brown appearance rather than the golden-yellow color produced by healthy plants. Stems develop a brittleness, indicating injury to the fiber. The disease also occurs in Great Britain.

**Ramularia lini** Lebed. On leaves and stems of *L. perenne* in Russia.

**Rhizoctonia napi** West. See Brassica.

**Septogloeum linicola** Speg. Pale effuse areas on leaves and stems of *L. usitatissimum* in Argentina.

**Tylenchus dipsaci** Kuehn. See Narcissus.

**La brûlure** or vlasbrand disease of flax is a widely distributed disease of flax (*L. usitatissimum*) in Belgium, France, and Ireland, due to an unknown cause.

**LIPARIS.** TWAYBLADE. Terrestrial and epiphytic orchids. See Orchidaceae.**LIPPIA.** LEMON VERBENA. Herbs and shrubs grown for their ornamental flowers. One species used for ground cover or lawns.

**Accidium evansi** P. Henn. Rust on leaves of *L. asperifolia* in the Union of South Africa.

**Phyllosticta aloysiae** Speg. Subcircular dull-brown leaf spots with white centers on *L. citriodora* in Argentina.

**Prospodium lippiae** (Speg.) Arth. Powdery golden-yellow rust pustules on *L. asperifolia*, *L. calli-carpaefolia*, *L. dulcis*, *L. lycioides*, *L. myriocephala*, *L. strigosa*, and *L. umbellata* in Argentina, Mexico, Guatemala, Cuba, and Costa Rica.

**Puccinia accedens** Syd. Powdery brown rust pustules on leaves of *L. aristata* and *L. arvensis* in Brazil and Uganda.

**Puccinia conjuncta** Diet. and Holw. Brown rust pustules on leaf blades and petioles and on stems of *L. pringlei* in Mexico.

**Puccinia elatipes** Arth. and Holw. Rust on leaves of *L. myriocephala* in Costa Rica and Guatemala.

**Puccinia lippicola** Pat. and Har. Brown rust pustules on circular brown spots on *L. adoensis* in central Africa.

**Puccinia lippivora** Syd. Rust on leaves of *Lippia* sp. in the Congo.

**Puccinia paraguayensis** Speg. Powdery cinnamon-brown rust pustules on *L. lycioides* in Brazil.

**Puccinia permagma** Arth. and Holw. Brown leaf rust on *L. myriocephala* in Costa Rica.

**Puccinia peruviana** Syd. Powdery dark-brown rust pustules on leaves of *L. urticoides* in Peru.

**Puccinia senilis** Arth. Yellow to brown powdery rust pustules on leaves of *L. myriocephala* in Guatemala.

**Puccinia vungunteni** Mayor. Rust on leaves of *L. americana* in Colombia.

**Uredo lippiae** Diet. and Holw. Brown, powdery rust pustules on leaves of *L. pringlei* in Mexico.

**LIRODENDRON.** TULIP TREE. YELLOW-POPLAR. Timber and ornamental trees.

**Diaporthe delitescens** Bomm., Rouss., and Sacc. On *L. tulipifera* in Belgium.

**Mycosphaerella elatior** Sacc. and Speg. See Magnolia.

**Myxosporium tulipiferae** Died. On twigs of *L. tulipifera* in Germany.

**LISIANTHUS.** Herbs or shrubs cultivated for their flowers.

**Phyllosticta lisianthi** Syd. Circular to subcircular, brown leaf spots with raised margins on *L. exsertus* in Jamaica.

**Uredo lisianthi** Pat. Rust on leaves of *L. elegans* in Brazil.

**LISTERA.** Terrestrial orchids. See Orchidaceae.**LITHOCARPUS.** See Pasania.**LITHOSPERMUM.** GROMWELL. Herbaceous, hardy perennials, used in rock gardens.

**Accidium lithospermi** Thuem. Rust on leaves of *L. arvense* and *L. purpureocaeruleum* in Austria and Macedonia.

**Peronospora lithospermi** Gäum. Downy mildew on leaves of *L. arvense* in central and north Europe.

**Puccinia arnaudi** Har. and Diet. Rust on leaves, stems, and sepals of *L. fruticosum* in France.

**Ramularia lithospermi** Lebed. On leaves of *L. officinale* in Russia.

**LITSEA.** Trees or shrubs.

**Asteroconium saccardoi** Syd. Forms yellow galls on leaves of *L. glaucesens* in Mexico.

**Cercospora litseae** P. Henn. Circular dull-brown leaf spots on *L. glauca* in Japan.

**Cercospora litseae-glutinosa** Syd. On leaves of *L. glutinosa* in the Philippines.

**Endodothella litseae** Racib. Black shiny irregular to circular stromata on leaves of *L. javanica* in Java.

**Gloeosporium litseae** Petch. Anthracnose on *Litsea* sp. in Ceylon.

**Helminthosporium maculosum** Sacc. On *L. perrottetii* in the Philippines.

**Laestadia litseae** (B. and Br.) Cke. Small pale-brown leaf spots on *Litsea* sp. in Australia.

**Oligostroma apiculatum** (Sacc. and Berl.) Theiss. and Syd. Shiny black somewhat irregular stromata on leaves of *L. dealbata* in Australia.

**Phyllachora cantonensis** Syd. Black stromata on leaves of *L. glutinosa* in China.

**Phyllachora laurinearum** Rac. Circular to elliptical black stromata on yellow-green subcircular leaf spots on *L. chrysocoma* and *Tetranthera* sp. in Java.

**Phyllachora lepida** Syd. Black stromata on leaves of *L. tayabensis* in the Philippines.

**Phyllachora litseae** Koord. Black stromata on indefinite yellowish spots on *L. polyantha* in Java.

**Puccinia litsea** (Pat.) Diet. and P. Henn. Chestnut-brown rust pustules on leaves of *L. glauca* in Japan.

**Teichosporella negeriana** Sacc. and Syd. Black fungus layers on leaves of *L. caustica* in Chile.

**Venturia litseae** Syd. On leaves of *L. glutinosa* in the Philippines.

**LIVISTONA.** Fan palms. See *Palmae*.**LLOYDIA.** Small bulbous plants.

**Asteroma lloydiae** Cruchet. Pale-brown, then darker, areas on stems and leaves of *L. serotina* in Switzerland.

**Puccinia bessei** Cruchet. Dull-brown rust pustules on leaves and stems of *L. serotina* in Switzerland.

**LOASA.** Flower-garden annuals.

**Aecidium rehderianum** Magn. Rust on leaves of *L. aurantiaca* and *L. papaverifolia* in Germany.

**Septoria loasae** F. Tassi. On leaves of *L. vulcanica* in Italy. Spegazzini has given the same name to a species on *Loasa* sp. in Argentina.

**LOBELIA.** Annual and perennial flower-garden and border plants.

**Ascochyta lobeliae** Petch. On *L. nicotianaefolia* in Ceylon.

**Cercospora ochracea** Sacc. and Malbr. On leaves of *L. urens* in France.

**Cercospora** (?) **tupae** Speg. Pale, indefinite spots on leaves of *L. tupa* in Chile.

**Gloeosporium hawaiiense** Thuem. Anthracnose on leaves of *L. macrostachys* in Hawaii.

**Heterosporium lobeliae** Speg. Indefinite leaf spots on *L. salicifolia* in Chile.

**Heterosporium tupae** Speg. Indefinite leaf spots on *L. bridgesii* in Chile.

**Mycosphaerella lobeliae** Petch. On leaves of *L. nicotianaefolia* in Ceylon.

**Phoma devastatrix** Berk. On *Lobelia* (cult.) in Great Britain.

**Phyllosticta bridgesii** Speg. Irregular whitish leaf spots on *L. bridgesii* in Chile.

**Puccinia aucta** Berk. and Muell. Yellow to brown rust pustules on leaves of *L. anceps*, *L. pedunculata*, and *L. platycalyx* in Australia.

**Puccinia vacua** Diet. and Holw. Brown rust pustules on leaves of *Lobelia* sp. in Mexico.

**Rhizoctonia destruens** Tass. See *Solanum*.

**LOESELIA.** Greenhouse plants related to *Gilia*.

**Puccinia fumosa** Holw. Brown to black rust pustules on leaves of *L. ciliata* and *L. coccinea* in Costa Rica and Guatemala.

**LOLIUM.** DARNEL. RYE GRASS. Pasture grasses.

**Cladochytrium caespitis** Griff. and Maubl. Attacks young plants of *Lolium* sp. in France, causing stems and leaf sheaths to turn brown, accompanied by a rotting of the roots.

**Fusarium loliaceum** Duc. Attacks the leaves of *L. italicum* in France.

**Fusarium nivale** (Fr.) Sor. See *Secale*.

**Hypochnus fuciformis** (Berk.) McAlp. Produces bright-pink glutinous fungus layers on spikelets of *Agropyron scabrum*, *Agrostis alba*, *Bromus mollis*, *B. sterilis*, *Danthonia pilosa*, *Festuca bromoides*, *Lolium perenne*, *Medicago denticulata*, and *Silybum marianum* in Great Britain and Australia.

**Ovularia lolii** Volk. Subcircular red-brown leaf spots on *L. italicum* and *L. perenne* in France and Switzerland.

**Ovularia pulchella** (Ces.) Sacc. and var. **lolii-italici** Ferr. See *Dactylis*.

**Puccinia brachypus** Speg. See *Bromus*.

**Sclerospora macrospora** Sacc. See *Triticum*.

**Septoria lolii** (Cast.) Sacc. Small subcircular blackish spots on glumes of *L. perenne* in France.

**Septoria passerinii** Sacc. See *Hordeum*.

**Sorosporium lolii** Thuem. A smut forming black spore masses in ovaries of *L. perenne* in Austria.

**Thecaphora westendorpii** Fisch. Dark-brown, irregular smut sori in spikelets of *L. perenne* in Belgium and Germany.

**Tilletia lolii** Awd. Stinking smut in ovaries of *L. arvense*, *F. linicolum*, *L. multiflorum*, *L. perenne*, *L. remotum*, and *L. temulentum* in Europe.

**Typhula graminum** Karst. Forms small black sclerotia in yellowed leaves, resulting at times in destruction of seedlings of *Hordeum* spp., *L. multiflorum*, *L. perenne*, *L. temulentum*, and *Triticum* sp. in Great Britain, Sweden, and Germany.

**Ustilago lolii** Magn. A smut deforming the inflorescences of *L. temulentum* in Egypt.

**Urocystis bolivari** Bub. and Frag. A smut of *L. perenne* in Spain, producing systemic infections and preventing the development of normal heads.

**LONCHOCARPUS.** LANCEPOD. Tropical trees or climbing shrubs.

**Aecidium menyharthi** P. Henn. Rust on leaves of *Lonchocarpus* sp. in South Africa.

**Diorchidium manaosense** P. Henn. Dull-brown rust pustules on leaves of *L. rariflorus* in Brazil.

**Endodothella lonchocarpicola** (P. Henn.) Theiss. and Syd. Dull-black stromata on lower leaf surfaces of *Lonchocarpus* sp. in Brazil.

**Ophiodothella atromaculans** (P. Henn.) v. Hoehn. Black stromata on both leaf surfaces of *Lonchocarpus* sp. in Brazil.

**Phyllachora lonchocarpi** Pat. and Har. Tar spot on leaves of *Lonchocarpus* sp. in the Congo.

**Phyllachora lungusaensis** P. Henn. Dull-black stromata on leaves of *Lonchocarpus* sp. in Tanganyika.

**Ravenelia bakeriana** Syd. Brown rust pustules on leaves of *Lonchocarpus* sp. in Brazil.

**Ravenelia lonchocarpi** Lagh. and Diet. Cinnamon-brown rust pustules on leaves of *L. campestris* and *L. latifolius* in Brazil, Cuba, and Guatemala.

**LONGICERA.** HONEYSUCKLE. Ornamental shrubs and climbers.

**Actinonema loniceriae-alpigenae** Allesch. Small circular to irregular dark-brown leaf spots on *L. alpigena* in Germany.

**LONICERA**—Continued.

- Ascochyta periclymeni** Thuem. Circular dull-yellow to gray leaf spots on *L. periclymenum* in Portugal.
- Ascochyta sarmenticia** Sacc. Subcircular to angular whitish leaf spots on *L. caprifolium* and *L. xylosteum* in France.
- Ascochyta tatarica** Allesch. On *L. tatarica* in Germany.
- Ascochyta vulgaris** Kab. and Bub. Circular to angular yellow-brown leaf spots with purple-brown margins on *L. xylosteum* in Austria.
- Cercospora periclymeni** Wint. On leaves of *L. caprifolium* and *L. periclymenum* in Russia, Denmark, and Germany.
- Criella lonicerae** P. Henn. and E. Nym. Brown concave spots on leaves of *Lonicera* sp. in Java.
- Dothidella periclymeni** (Fckl.) Theiss. and Syd. On *L. periclymenum* in Germany.
- Kabatia latemarensis** Bub. Dull-yellow to white leaf spots with broad purple margins on *L. caerulea* and *L. xylosteum* in Italy and Austria.
- Kabatia mirabilis** Bub. Angular dull-yellow to whitish leaf spots with black margins on *L. alpigena* and *L. nigra* in Austria.
- Lasiobotrys lonicerae** Kunz. and Schm. Circular black spots on leaves and branches of *Lonicera* spp. in India, Siberia, and Europe. What has been referred to this species in California is distinct.
- Leptosphaeria periclymeni** Oud. On branches of *L. periclymenum* in Russia.
- Marsonia staritzii** Bres. Small, then confluent, black leaf spots on *L. tatarica* in Germany.
- Melasmia lonicerae** Jacz. Black stromata on leaves of *L. maackii* and *L. maximowiczii* in Austria.
- Mycosphaerella clymenia** Sacc. Subcircular ochraceous to gray leaf spots on *L. caprifolium* and *L. periclymenum* in Denmark, Italy, and France.
- Mycosphaerella implexicola** (R. Maire.) Jaap. Gray leaf spots with brown margins on *L. implexa* in Dalmatia and Spain.
- Ophiobolus minor** Bub. On branches of *L. xylosteum* in Bohemia.
- Phyllosticta alpigena** Sacc. Black leaf spots on *L. nigra* in Italy.
- Phyllosticta caprifolii** (Op.) Sacc. Subcircular gray leaf spots on *L. caprifolium*, *L. implexa*, and *L. pallasii* in Siberia, Russia, Dalmatia, France, and Italy.
- Phyllosticta nitidula** Dur. and Mont. Large brownish leaf spots on *L. implexa* in Algeria.
- Phyllosticta sydowii** Bres. On leaves of *L. latifolia* in Germany.
- Puccinia longirostris** Kom. Brown rust pustules on small sunken yellow spots on *L. altmanni* and *L. turkestanica* in Turkestan.
- Ramularia lonicerae** Vogl. and f. **hispanica** Gz. Frag. Circular to irregular chestnut-brown leaf spots on *L. alpigena* and *L. hispanica* in Italy and Spain.
- Rhabdospora xylostei** Lamb. and Fautr. On branches of *L. xylosteum* in France.
- Rhytisma lonicericola** P. Henn. Carbonous black stromata on leaves of *L. japonica*, *L. maackii*, and *L. tschonoskii* in Japan and Russia.
- Rhytisma xylostei** Naoumoff. On leaves of *L. xylosteum* in Russia.
- Septoria obscurata** Thuem. Circular ashen-colored leaf spots with dark purple margins on *L. periclymenum* and *L. tatarica* in Russia and Portugal.
- Septoria xylostei** Sacc. and Wint. Small whitish leaf spots with black margins on *L. nigra*, *L. tatarica*, and *L. xylosteum* in Russia and Switzerland.
- Trabutia lonicerae** Rac. Circular black stromata on leaves of *L. javanica* in Java.
- LOPEZIA.** Greenhouse plants, cultivated for their attractive flowers. Grown outdoors in the South.
- Puccinia fuchsiae** Syd. and Holw. See Fuchsia.
- LOPHANTHUS.** Erect plants with clusters of blue flowers.
- Aecidium lophanthi** P. Henn. Rust on leaves of *L. rugosus* in Japan.
- LOTUS.** Herbs or shrubs, grown for their various-colored flowers.
- Cercospora loti** Hóll. Circular red-brown leaf spots on *L. siliquosus* in Hungary.
- Mitrla sclerotiorum** Rostr. See Medicago.
- Ovularia sphaeroidea** Sacc. On leaves of *L. corniculatus*, *L. uliginosus*, *Vicia sativa*, and *V. villosa* in France, Denmark, Sweden, Italy, and Austria.
- Peronospora lotorum** Syd. Downy mildew on leaves of *L. corniculatus* and *L. uliginosus* in central and northern Europe.
- Ramularia loticola** Massal. On leaves of *L. corniculatus* in Italy.
- Ramularia schulzeri** Bäuml. See Anthyllis.
- Septoria stipularis** Pass. On leaves, stems, and stipules of *Lathyrus aphaca* and *Lotus corniculatus* in Denmark and Italy.
- Uromyces anthyllidis** (Grev.) Schroet. See Anthyllis.
- Uromyces libycus** Trott. Rust on leaves of *L. pusillus* in Tripoli.
- Uromyces loti** Blytt. Brown rust pustules on leaves of *Euphcrbia cyparissia*, *Lotus angustissimus*, *L. corniculatus*, *L. palustris*, and *L. uliginosus* in Japan and Europe.
- LUCUMA.** SAPOTE. CANISTEL. MAMEY COLORADO. Tropical trees grown for fruit.
- Phyllosticta lucumae** Syd. Circular to subcircular whitish leaf spots with raised purple-brown margins on *L. neriiifolia* in Uruguay.
- Uredo lucumae** Arth. and Johnst. Rust on leaves of *L. nervosa* in Cuba.
- Uromyces lucumae** Diet. Powdery brown leaf spots on *Lucuma* sp. in Brazil.
- LUFFA.** TOWEL GOURD. DISCLOTH GOURD.
- Corticium koleroga** (Cke.) v. Hoehn. See Coffea.
- LUNARIA.** HONESTY. Herbaceous annuals and perennials.
- Helminthosporium lunariae** Póll. Circular leaf spots on *L. annua* (*L. biennis*) in Italy.
- Peronospora lunariae** Gäum. Downy mildew on leaves of *L. annua* and *L. rediviva* in central Europe.
- LUPINUS.** LUPINE. Herbaceous legumes.
- Ascochyta lupinicola** Petr. On leaves of *Lupinus* sp. in Bohemia and Austria.
- Bacillus elegans** Hegyi. Yellow, then brown, spots on leaves of *L. albus*, *L. angustifolius*, and *L. luteus* in Hungary.
- Ceratophorum setosum** Kirch. See Cytisus.
- Chalaropsis thielavioides** Peyr. Said to cause a root rot of *L. albus* in Italy.
- Chrysocelis lupini** Lagerh. and Diet. Rust on leaves of *L. clarkii*, *L. paniculatus*, and *Lupinus* sp. in Colombia, Costa Rica, and Peru.
- Gloeosporium lupinus** Bondar. Anthracnose on *Lupinus* sp. in Brazil.
- Rhytisma onobrychidis** Sacc. Black stromata on leaves and stems of *Lathyrus latifolius*, *Lupinus* sp., and *Hedysarum coronarium* in Italy, France, and Germany.
- Uromyces anthyllidis** Grev. See Anthyllis.
- Uromyces elatus** Syd. Yellow and brown rust pustules on leaves of *L. ramosissimus*, *L. saxatilis*, and *L. tomentosus* in Bolivia and Peru.
- Uromyces lupinicolus** Bub. Powdery brown rust pustules on leaves of *L. albus*, *L. angustifolius*, and *Lupinus* sp. in Spain and Bohemia.

**LUPINUS**—Continued.

**Uromyces montanus** Arth. Cinnamon-brown rust pustules on leaves of *L. mexicanus* and *L. montanus* in Mexico and Guatemala.

**Uromyces renovatus** Syd. Powdery brown rust pustules on leaves of *L. albus*, *L. angustifolius*, *L. hispanicus*, *L. luteus*, and *L. termis* in Europe, northern Africa, and Madeira.

**Uromyces rugosus** Arth. Powdery cinnamon-brown to dark-brown rust pustules on leaves of *Lupinus* sp. in Mexico.

**LUZULA. JUNCOIDES** Ag. WOOD-RUSH. Grass or rushlike perennial herbs.

**Ascochyta teretiuscula** Sacc. and Roum. On leaves of *L. campestris* and *L. vernalis* in France and Yugoslavia.

**Phyllachora luzulae** (Rabh.) Cke. Dull-black stromata on both leaf surfaces of *Luzula* sp. in Germany.

**Puccinia luzulae-maximae** Diet. Leaf rust on *L. maxima* in Europe.

**Puccinia luzulina** Syd. Cinnamon-brown rust pustules on leaves of *L. alopecurus* in South America.

**Puccinia oblongata** (Lk.) Wint. Dull-brown to black rust pustules on red-brown irregular leaf spots on *L. campestris*, *L. forsteri*, *L. maxima*, *L. multiflora*, *L. nivea*, *L. pilosa*, and *L. sylvatica* in Europe.

**Puccinia tenuispora** McAlp. Yellow-brown to dark-brown rust pustules on leaves and stems of *L. campestris* and *L. oldfieldii* in Australia and Tasmania.

**Septoria chanousiana** Ferr. Purple-brown leaf spots on *L. lutea* in Italy.

**Septoria luzulae** Schrcet. On leaves of *L. forsteri* in Yugoslavia.

**Uredo antarctica** Berk. Rust on leaves of *L. crinita* in the Campbell Islands.

**Urocystis luzulae** (Schroet.) Wt. Black smut pustules on leaves of *L. multiflora* and *L. pilosa* in Denmark and Germany.

**Ustilago hyperborea** Blytt. Smut in fruit of *L. hyperborea* in Norway.

**Ustilago liebmanni** P. Henn. A smut transforming the ovaries of *Luzula* sp. in Mexico into brown spore masses.

**Ustilago vujiickii** Oud. and Beyer. A smut forming black spore masses in ovaries of *L. campestris* in Holland.

**LYCASTE.** Terrestrial and epiphytic orchids. See Orchidaceae.**LYCHNIS. CAMPION. CORN COCKLE. MALTESE CROSS.** Flower-garden herbs.

**Ascochyta cookei** Mass. Blotches on leaves of *L. alba* (*L. vespertina*) in Great Britain.

**Ascochyta dianthi** (A. and S.) Berk. See Dianthus.

**Fabraea inoplexa** Bres. and Cap. On leaves of *L. flos-jovis* in Italy.

**Gloeosporium lychnidis** Oud. Anthracnose on leaves of *L. dioica* (*L. diurna*) in Holland.

**Marsonia delastrei** (Delacr.) Sacc. Dull-yellow indefinite areas on leaves and stems of *L. chalconica*, *L. dioica*, *Silene latifolia*, *S. noctiflora*, and *Agrostemma githago* in Europe and Siberia. Reported from Wisconsin.

**Peronospora lychnitis** Gäum. Downy mildew on leaves of *L. coronaria* in Germany.

**Phyllosticta punctiformis** (Desm.) Allesch. On leaves of *L. dioica* in Scotland.

**Puccinia lychnidis-miqueliana** Diet. Rust on leaves of *L. miqueliana* in Japan.

**Ramularia chalconica** Allesch. Ashen leaf spots on *L. chalconica* in Germany.

**Ramularia lychnicola** Cke. On leaves of *L. chalconica*, *L. dioica*, and *Silene latifolia* in Great Britain, Austria, and Germany.

**Septoria lychnidicola** P. Brun. Circular to oblong ochraceous leaf spots with brown margins on *L. flos-cuculi* in France.

**Septoria lychnidis** Desm. Irregular red-brown leaf spots on *L. chalconica*, *L. dioica*, *L. flos-jovis* and *Gypsophila allissima* in Siberia, Italy, France, and Great Britain.

**Septoria melandrii** Pass. Subcircular reddish leaf spots with dull-brown margins on *L. diurna* and *L. alba* (*L. vespertina*) in Italy. Reported from Wisconsin.

**Uromyces behenis** (DC.) Ung. Rust on leaves of *L. indica* in India.

**Uromyces crassivertex** Diet. Cinnamon-brown to black rust pustules on leaves of *L. miqueliana* in Japan.

**Uromyces lychnidicola** Speg. Rust on leaves of *L. magellanica* and *L. patagonica* in Argentina.

**LYCIUM. MATRIMONY VINE.** Ornamental shrubs and vines.

**Ascochyta destructiva** Kab. and Bub. Subcircular, often confluent, ochraceous leaf spots on *L. barbarum* in Bohemia.

**Ascochyta lycii** Rostr. Ashen leaf spots with dull-brown margins on *L. barbarum* and *L. halimifolium* in Denmark.

**Microsphaera mougeotii** Lévl. Powdery mildew on leaves of *L. barbarum*, *L. europaeum*, *L. ovatum*, and *L. ruthenicum* in Europe.

**Phyllosticta destructiva** Desm. See Althaea.

**Puccinia afra** Wint. Powdery brown rust pustules on leaves, peduncles, and sepals of *L. afrum* in Spain and the Union of South Africa.

**Puccinia lycii** Kalchbr. Brown rust pustules on leaves of *L. tubulosum* in the Union of South Africa. Reported from Ohio.

**Puccinia lyciicola** Speg. Brown rust pustules on leaves of *L. argenteum* and *L. patagonicum* in Argentina.

**Puccinia turgida** Syd. Powdery brown to dark-brown rust pustules on leaves of *L. europaeum* in Palestine.

**Septoria lycii** Paul. and Gz. Frag. On leaves of *L. helimifolium* (*L. vulgare*) in Spain.

**Septoria lyciicola** Speg. Circular definite leaf spots on *L. cestroides* in Argentina.

**Uredo megalospora** Speg. Rust on leaves of *Lycium* sp. in Argentina.

**LYCOPERSICUM. (LYCOPERSICON.)** TOMATO. Herbaceous plants, cultivated for their fruit.

**Bacillus caulivorus** Prill. and Delacr. See Solanum.

**Bacterium briosii** Pavar. Bacteriosis of leaves, stalks, and fruits of *L. esculentum* in Italy and France. The leaves wither up, new buds are deformed, and long brown spots appear on the stems.

**Bacterium lycopersici** Burgwitz. This bacterium is reported as the cause of "blossom-end rot" of fruit of *L. esculentum* in Russia.

**Chrysothlyctis endobiotica** Schilb. See Solanum.

**Diplodina lycopersicola** (v. Bond.) Mont. Large black spots on tops or sides of both green and ripe fruit of *L. esculentum* in Russia. The spots are somewhat depressed, warty, black at the centers, and brown at the margins, with surrounding dirty-green areas. Probably not distinct from *Phoma destructiva* Plowr.

**Fusarium erubescens** A. and v. Ov. Small black sunken spots on fruit of *L. esculentum* in Germany.

**Hainesia lycopersici** Speg. On leaves of *L. esculentum* in Argentina.

**Phoma ferrarisii** Ciferri. Causes a fruit rot of *L. esculentum* in Italy. *Ramularia ferrarisii* Cif. is described as a stage of the same fungus. Probably not distinct from *P. destructiva* Plowr.

**LYCOPERSICUM**—Continued.

- Phytophthora cryptogea** Pethyb. and Laffer. This fungus has been found as one of the causes of damping off of seedlings of *L. esculentum* in Great Britain. It also attacks *Cineraria* seedlings. Diseased plants show brown areas at the soil level and collapse, breaking over at the point of infection.
- Phytophthora melongenae** K. Saw. See Solanum.
- Phytophthora mexicana** Hots. and Hartge. This downy mildew is said to cause a rot of fruit of *L. esculentum* in Mexico and to be capable of attacking other plant parts as well. Infected fruit have been imported into the United States.
- Phytophthora parasitica** Dastur. See Ricinus.
- Puccinia pitteriana** Arth. See Solanum.
- Rhizoctonia** sp. See Vigna.
- Sclerotium setosum** Bewley and Shearn. Affected plants stop growing; the lower leaves turn yellow and die prematurely, and the terminal growths are pale and weak. The stems become hollow and yellow, and a dark-brown discoloration spreads up the stems from the ground level. Finally all the leaves wither and the plants die. The roots decay and shrivel. Numerous minute black globular sclerotia occur in the rotted tissues of the roots and the bases of the stems. On *L. esculentum* in England. This Sclerotium has been connected with *Colletotrichum tabificum* (Hallier) Pethybr. *Vermicularia varians* Duc. is also considered synonymous.
- Spongospora subterranea** (Walls.) Lang. See Solanum.
- Vermicularia capsici** Syd. See Capsicum.
- Vermicularia varians** Duc. See Solanum.
- Verticillium pulverulentum** Gouwen. Circular, sharply delimited, dark spots on fruit of *L. esculentum*, causing premature fall in Holland.
- LYGODIUM.** CLIMBING FERN.
- Aphelenchus olesistus** Ritz. Boz. See Begonia.
- Helminthosporium filicicola** P. Henn. Effuse leaf spots on *Lygodium* sp. and *Selaginella* sp. in Peru and Brazil.
- Milesia blechni** Syd. See Blechnum.
- Mycosphaerella ditissima** Syd. On fronds of *L. flexuosum* in the Philippines.
- Puccinia lygodii** (Har.) Arth. (*Uredo lygodii* Har.) Yellow-brown rust pustules on fronds of *Lygodium* sp. in Brazil.
- LYONIA.** Ornamental shrubs grown for their white flowers and dense foliage.
- Exobasidium fawcettii** Mass. On leaves of *L. jamaicensis* in Jamaica. Probably not distinct from *E. vaccinii* Fckl.
- LYSILOMA.** Trees and shrubs of the American Tropics.
- Ravenelia lysilomae** Arth. Yellow-brown to chestnut-brown rust pustules on leaves of *L. tergemina* in Mexico.
- Ravenelia sololensis** Arth. and Holw. Rust on leaves of *L. acapulcensis* and *L. bahamensis* in Mexico, Cuba, and Guatemala.
- LYSIMACHIA.** LOOSESTRIFE. Erect or creeping leafy herbs.
- Aecidium lysimachiae-japonicae** Diet. Leaves of *L. japonica* covered with rust pustules in Japan.
- Phyllosticta lysimachiae** Allesch. Irregular, then confluent, greenish-ashen, then pale ochraceous, leaf spots on *L. vulgaris* in Russia and Germany.
- Puccinia dieteliana** Syd. Yellow and dark-brown rust pustules on leaves of *L. clethroides* and *L. fortunei* in Japan.
- Ramularia lysimachiae** Thuem. On leaves of *L. nummularia*, *L. thyrsoiflora*, and *L. vulgaris* in Siberia, Yugoslavia, Denmark, Austria, Bohemia, and Germany. Reported from Wisconsin.
- Ramularia lysimachiarum** Lindr. Brown to dark-green, then yellow, leaf spots on *L. nummularia* in Sweden, Finland, Denmark, and Great Britain.
- Septoria bresadoleana** Krieger. Reddish leaf spots, becoming grayish white at the centers, on *L. vulgaris* in Germany.
- Septoria lysimachiae** West. Brown leaf spots on *L. nummularia*, *L. thyrsoiflora*, and *L. vulgaris* in Siberia and Europe.
- Septoria nambuana** P. Henn. Red-brown leaf spots on *L. brachystachys* in Japan.
- LYTHRUM.** Spiked or purple loosestrife. Herbs or subshrubs.
- Aecidium lythri** Diet. and Neg. Rust on leaves of *L. hyssopifolium* in Chile.
- Cercospora lythri** (West.) Niessl. On leaves of *L. salicaria* in Spain, Belgium, Italy, and Germany.
- Doassansia winteriana** (Wint.) Magn. A smut producing small globose punctiform galls on both leaf surfaces of *L. hyssopifolium* in Australia.
- Septoria brissaceana** Sacc. and Let. Small dull-yellow leaf spots with dark margins on *L. salicaria* in Denmark, France, Russia, and Germany.
- MAACKIA.** Trees grown for their handsome foliage and panicles of white flowers.
- Uromyces amurensis** Kom. Leaf rust on *M. amurensis* in Japan.
- MABA.** Tropical trees and shrubs with hard ebonylike wood.
- Aecidium bicolor** Sacc. Rust on leaves of *M. buxifolia* and *M. natalensis* in India and the Union of South Africa.
- Aecidium melaleucum** Syd. Leaf rust on *M. buxifolia* in India.
- Aecidium rhytismoideum** B. and Br. var. **mabae** P. Henn. See Diospyros.
- Ascochyta mabiana** Sacc. Dull-brown leaf spots with darker margins on *M. abyssinica* in Abyssinia.
- Exobasidium symploci-japonicae** Kusano. See Symplocos.
- Phyllachora mabae** (P. Henn.) Theiss. and Syd. Circular shiny-black stromata on leaves of *M. inconstans* in Brazil.
- MACARANGA.** Trees or shrubs.
- Cercospora macarangae** Syd. Circular ochraceous leaf spots on *M. tanaria* in the Philippines.
- Marsonia pavonina** Syd. Leaf spots on *M. bicolor* in the Philippines.
- Phyllachora macarangae** P. Henn. Black stromata on circular brown leaf spots on *Macaranga* sp. in the Philippines.
- MACFADYENA.** Tall climbing tropical shrubs.
- Leptothyrium heterospermum** Speg. Shield-shaped black fruiting bodies on yellow leaf spots on *M. cynanchoides* in Argentina.
- MACLURA.** TOXYLON Ag. OSAGE ORANGE. BOW WOOD. Cultivated as a hedge plant and for its evergreen foliage and orange-like fruit.
- Ectostroma maclurae** Thuem. On leaves of *M. aurantiaca* (*Toxylon pomiferum*) in Italy and Portugal.
- Septoria maclurae** P. Brun. Subcircular to irregular olive or dull-yellow leaf spots with brown margins on *M. aurantiaca* (*T. pomiferum*) in France.
- Uredo maclurae** Speg. Rust on leaves of *M. aurantiaca* (*T. pomiferum*) and *M. ucora* in Argentina.
- MACROZAMIA.** Cycaslike plants.
- Cryptosporella macrozamiaie** P. Henn. On leaf blades and petioles of *M. fraseri* in Australia.

**MAESA.** Tropical and subtropical shrubs.

**Amazonia peregrina** Syd. On *M. indica* and *M. laxa* in Ceylon and the Philippines.

**Fusidium maesae** P. Henn. Pale or ashen leaf spots on *M. lanceolata* in Tanganyika.

**Physalospora chaenostoma** Sacc. On leaves of *Maesa* sp. in Natal.

**Septoria maesae** F. Tassi. Circular yellowish, then dull-brown, leaf spots on *M. indica* in Italy.

**MAGNOLIA.** Woody plants cultivated for their handsome foliage and showy flowers.

**Ascochyta magnoliae** Thuem. On leaves of *M. grandiflora* in Italy.

**Gloeosporium haynaldianum** Sacc. and Roum. Anthracnose on leaves of *M. grandiflora* and *M. soulangeana* in Denmark, Italy, and France.

**Hendersonia magnoliae** Sacc. Irregular whitish leaf spots with brown margins on *M. grandiflora* in France and Italy.

**Mycosphaerella elatior** Sacc. and Speg. and var. **apula** Sacc. On leaves of *M. grandiflora* and *Liriodendron tulipifera* in Spain and Italy.

**Phyllosticta kobus** P. Henn. Irregular dull-brown leaf spots on *M. kobus* in Japan.

**Phyllosticta yulan** Tassi. Large pale-brown leaf spots on *M. denudata* (*M. yulan*) in Italy.

**Septoria caerulea** Tassi. Fuscous leaf spots on *M. denudata* in Italy.

**Sphaeropsis magnoliae** Magnag. On *Magnolia* sp. in Italy.

**MAGYDARIS.** Perennial herbs.

**Puccinia magydaridis** Pat. and Trab. Cinnamon-brown to black rust pustules on leaves of *M. panacina* and *M. tomentosa* in Algeria, Tunis, and Spain.

**MAHONIA.** See Berberis.

**MAIANthemum. UNIFOLIUM** Ag. Small spring-blooming woody perennials.

**Cercospora majanthemi** Fckl. On leaves of *M. bifolium* in Denmark, Belgium, Sweden, Russia, Siberia, and Austria. Reported from Ohio.

**Mycosphaerella asteroma** (Fr.) Lind. See Convallaria.

**Puccinia majanthemi** Diet. Brown rust pustules on both leaf surfaces and on petioles of *M. bifolium* in Japan.

**Ramularia rubicunda** Bres. On leaves of *M. bifolium* in Russia, Germany, and Austria.

**Septoria majanthemi** West. Dull-gray indefinite leaf spots on *M. bifolium* in Belgium and Germany.

**MALLOTUS.** MONKEY-FACE TREE. Tropical trees or shrubs.

**Cercospora laguensis** (Sacc.) Yates. On leaves of *M. moluccanus* in the Philippines.

**Cronartium malloti** Racib. Golden-yellow to red-brown rust pustules on leaves of *M. moluccanus* in Java.

**Helminthosporium insigne** Sacc. On *M. philippinensis* in the Philippines.

**Phyllosticta marmorta** Cke. Confluent leaf spots on *M. philippinensis* in India and the Philippines.

**Uromyces malloti** P. Henn. Brown rust sori on leaves of *M. moluccanus* in New Guinea and the Philippines.

**MALUS.** APPLE. CRAB.

**Asteroma mali** Desm. Circular brown leaf spots on *Malus* sp. in Germany.

**Bacterium mali** Brzez. Reported as the cause of a canker disease of trunks and limbs of *Malus* (apple) in central Europe.

**Bacterium** sp. Blossom blight. See *Pyrus*.

**Botryosphaeria mali** Putt. Causes a cankering of branches and trunks of *Malus* (apple) in the Union of South Africa. Death of infected branches, and even of entire trees, may be brought about through girdling.

**Cercospora porrigo** Speg. Causes a drying of young fruits of *Malus* sp. in Argentina.

**Colletotrichum piri** Noack. See *Pyrus*.

**Coniothecium chromatoporum** Cda. This fungus is reported as the cause of the blister and fruit-cracking disease of *Pyrus* and *Malus* in the Union of South Africa, Australia, Tasmania, New Zealand, and India. Dark specks appear in groups on the twigs and branches, finally becoming reddish-brown blisters. Diseased twigs die back. On the fruit the fungus produces a scurfy condition or russetting. Young fruit when attacked crack on expanding.

**Dendrodochium pulchrum** Marchal. On *Malus* (apple) in Belgium.

**Diaporthe mali** Miura. Pale, discolored areas, from 1 to 2 centimeters in diameter, on the leaves of *Malus* (apple) in Japan, producing curling and defoliation. Young shoots and twigs also show irregular brown areas which cause die-back. Spots on the fruit are circular to irregular, coalescing and involving the entire fruit in a brown rot. A serious disease throughout Japan. *Diaporthe mali* Bres., reported from Australia, is apparently distinct.

**Diaporthe perniciosa** Marchal. See *Prunus*.

**Diplodia griffoni** Sacc. and Trav. This fungus has been confused with *Physalospora cydoniae* but is considered distinct. It causes a serious canker disease of twigs of *Malus* (apple) and *Pyrus* (pear) in Europe and New Zealand. The cankers are slightly depressed, reddish brown, zonate, and with the surfaces studded with the pycnidia. Old cankers are dull brown.

**Dothiorella vinosa** Marchal. See *Prunus*.

**Fusarium** sp. A blossom blight of *Malus* sp. has been attributed to this fungus in Great Britain.

**Gymnosporangium sabiniae** (DC.) Wint. See *Juniperus*.

**Gymnosporangium yamadai** Miy. See *Juniperus*.

**Helicobasidium tanakae** Miy. See *Morus*.

**Hydnum schiedermayeri** Heufler. A wound parasite of *Malus* sp. in Europe, causing a rot of bark and wood of trunk and limbs. Reported from Pennsylvania.

**Labrella piricola** Bres. and Sacc. See *Pyrus*.

**Laestadia mali** (Fckl.) Sacc. On leaves of *Malus* sp. in Belgium and Germany.

**Leptosphaeria vagabunda** Sacc. Causes a fruit spot and rot of *Malus* sp. in Great Britain.

**Marsonia mali** P. Henn. Subcircular, then confluent, dull-brown leaf spots with purple margins on *Malus* in Japan.

**Melanobasidium mali** Maubl. Oval or elongate whitish leaf spots with brown margins on *Malus* in Spain.

**Melasmia mali** Oud. On leaves of *Malus* in Holland.

**Mycosphaerella pomacearum** Sacc. On leaves of *Malus* sp. and *Cydonia* sp. in China and France.

**Mycosphaerella pomei** (Pass.) Lind. On leaves of *Malus* in Italy.

**Myxosporium mali** Bres. Brown spotting and rot of fruit of *Malus* in Great Britain and on dead twigs in Germany.

**Ochropsora pallida** Rostr. See *Sorbus*.

**Phacidiella discolor** (Mont. and Sacc.) A. Poteb. Causes a fruit rot and trunk canker of *Malus* (apple) and *Pyrus* (pear) in Europe.

**Phyllosticta mali** Prill. and Delacr. Small elongate or circular ochraceous, then dirty-gray, leaf spots with fuscous margins on *Malus* sp. in Denmark, Russia, Bulgaria, and France. Reported from West Virginia.

**Phyllosticta briardi** Sacc. Brown or ochraceous leaf spots with darker borders on *Malus* and *Pyrus* sp. in Russia, Great Britain, France, Malta, and Hungary, often causing defoliation.

**MALUS**—Continued.

- Phytophthora syringae** Kleb. Produces a dark-brown rather firm and elastic rot of fruit of *Malus* in Europe. See, also, *Syringa*.
- Plectodiscella piri** Woro. See *Pyrus*.
- Pleospora pomorum** Horne. Causes a brown spotting and rot of fruit of *Malus* in Great Britain.
- Polypeus purpureus** Horne. Associated with fruit spotting and rotting of *Malus* in Great Britain. *P. aureus* Horne, *P. pomi* Horne, and *P. recurvatus* Horne have been found in similar situations.
- Polystigma ochraceum** (Wahlb.) DC. See *Prunus*.
- Sclerotinia mali** Tak. Produces brown areas on leaf blades and petioles, flower stalks, and branches and a brown rot of young fruit of *Malus* in Japan.
- Sclerotium stellatum** Horne. Fruit spot and rot of *Malus* in Great Britain.
- Spilocaea pomi** Fr. Round sunken brown spots on ripe fruit of *Malus*, especially during storage, in Russia, Great Britain, and Germany. Called the "apple-speck" disease.
- Sporodesmium cerebriforme** McAlp. "Eye-scale" disease of *Malus* in Australia.
- Stagonospora mali** Delacr. See *Pyrus*.
- Stereum purpureum** Pers. See *Prunus*.
- Trichoseptoria fructigena** Maubl. See *Cydonia*.
- Trichothecium candidum** Walbr. "White-apple blotch" is marked by small round spots with brown-black borders, causing a brownish rot and mummification of fruit of *Malus* in Europe. Probably not distinct from *T. roseum*.
- Valsa mali** Miy. and Yam. Oblong to irregular slightly elevated brown spots on branches of *Malus* in Japan. These areas dry out, become sunken, more or less darkened, and the surface cracks.
- A trunk canker of apple (*Malus*), with which bacteria and a *Cytospora*-like fungus are associated, has been reported from Great Britain. Large, rapidly extending, sunken cankers form on the lower part of the trunk, soon girdling and killing infected trees.
- MALVA**. MALLOW. Annual or perennial herbs.
- Ascochyta malvae** H. Zimm. Subcircular ochraceous, then white, leaf spots on *M. neglecta* and *M. vulgaris* in Austria and Germany.
- Ascochyta malvicola** Sacc. Subcircular or sinuous whitish leaf spots on *M. rotundifolia* and *M. sylvestris* in Italy and Austria.
- Ascochyta montenegrina** Bub. Subcircular, often confluent, yellow leaf spots on *M. sylvestris* in Yugoslavia.
- Cercospora malvarum** Sacc. Large olive-colored leaf spots on *M. moschata* in Spain and Denmark. Reported from Kansas.
- Cercospora polymorpha** Bub. Subcircular, oblong, or angular yellowish leaf spots with narrow dark-brown borders on *M. sylvestris* and *M. verticillata* in China and Yugoslavia.
- Phyllosticta destructiva** Desm. See *Althaea*.
- Puccinia heterogenea** Lagerh. Chestnut-colored rust pustules on leaves and stems of *Althaea rosea*, *M. crispa*, *M. nicaeensis*, and *M. peruviana* in Ecuador.
- Ramularia malvae** Fekl. Circular to elongate white leaf spots on *M. moschata*, *M. rotundifolia*, and *M. sylvestris* in Denmark and Germany.
- Septoria heterochroa** Desm. Small subcircular gray-brown, then whitish leaf spots on *M. neglecta*, *Althaea officinalis*, and *Sida* sp. in Costa Rica, Russia, Denmark, Italy, Belgium, and Great Britain.
- MALVASTRUM**. FALSE MALLOW. Mallowlike herbs.
- Aecidium malvastricola** P. Henn. Rust on leaves of *M. spicatum* in Argentina.
- MANDEVILLA**. Tall tropical climbers.
- Phyllosticta gelsemii** E. and E. var. **mandevilleae** Sacc. and Scal. Subcircular gray leaf spots on *M. suaveolens* in Portugal.
- Uredo mandevillae** Mayor. Rust on leaves of *M. (Echites) mollissima* and *M. tomentosa* in Colombia and Trinidad.
- MANETTIA**. Twinning plants grown for their flowers.
- Puccinia niederleinii** P. Henn. Cinnamon-brown rust pustules on leaves of *M. bicolor*, *M. ignita*, and *M. leianthiflora* in Argentina and Paraguay.
- Uromyces tener** Schroet. Powdery brown rust pustules on leaves of *M. gracilis* in Brazil.
- MANGIFERA**. MANGO. Tropical fruit and shade trees.
- Bacillus mangiferae** Dodge. This bacterial blight first appears on mango (*M. indica*) fruits as small water-soaked areas which rapidly enlarge, turn brown, and crack, corky tissue developing. Gumming generally occurs. Similar irregular brown spots occur on leaf blades and petioles, young twigs, peduncles, and pedicels. Diseased fruits fall. The disease is serious in Portuguese east Africa and the Union of South Africa.
- Cercospora mangiferae** Koord. Small circular numerous brown leaf spots on *M. indica* in Java, Philippines, and Haiti.
- Coccomyces vilis** Syd. and Butl. Irregular spots with purple margins on leaves of *M. indica* in India.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Cronartium kemangae** Racib. Brown rust pustules on irregular leaf spots on *M. kemanga* in Java.
- Diplodia cacaoicola** P. Henn. See *Theobroma*.
- Endothia havenensis** Brun. See *Eucalyptus*.
- Eutype erumpens** Mass. See *Theobroma*.
- Gloeosporium mangiferae** P. Henn. (*Gloeosporium mangae* Noack.). These anthracnose fungi, reported as causing bloom blight, fruit rot, and leaf spotting of the mango (*M. indica*) in Trinidad, Brazil, and elsewhere, are certainly synonymous with *Colletotrichum gloeosporioides*, widespread in the United States.
- Leptothyrium circumscissum** Syd. Large irregular or angular leaf spots on *M. indica* in the Philippines. Especially destructive to seedlings.
- Macrophoma** sp. Attacks the fruit of *M. indica* in Brazil, causing them to blacken and fall to the ground, rotting or mummifying.
- Meliola mangiferae** Earle. Black circular to irregular superficial fungus patches on leaves of *M. indica* in the West Indies, the Philippines, and Malaya.
- Oidium** sp. A powdery mildew attacking the leaves of *M. indica* has been reported from India and Cuba.
- Phyllachora** sp. Tar spot on leaves of *M. indica* in the Philippines.
- Phyllosticta mortoni** Fairm. Numerous small deep-brown angular leaf spots on *M. indica* in the Bahamas, Mexico, and Florida.
- Rosellinia bunodes** B. and Br. See *Citrus*.
- Sphaerostilbe repens** B. and Br. See *Hevea*.
- Stilbella flavida** (Cke.) Lind. See *Coffea*.
- Zimmermaniella trispora** P. Henn. Black stromata on leaves of *M. indica* in Java.
- Eucalyptus disease**. See *Eucalyptus*.

- MANIHOT.** CASSAVA. Sometimes called mandioca. Ceara rubber tree. Tropical trees and shrubs with many economic uses.
- Ascochyta manihotis** P. Henn. Circular pale-yellow-brown leaf spots on *M. utilissima* in Tanganyika.
- Bacillus manihotis** Arthaud. and Ber. Young shoots rot internally, the vascular bundles decompose, wither, and die. On older shoots elongate gum pockets form and a yellow gum exudes. On *M. palmata* and *M. utilissima* in Brazil.
- Cercospora cearae** Petch. Irregular, then confluent, red-brown, then gray, spots on *M. glaziovii* and *M. piuhyensis* in India, Ceylon, Malaya, and Uganda, causing the leaves to crumple up and fall.
- Cercospora henningsii** Allesch. Subcircular to irregular pale-ochraceous leaf spots on *M. palmata* and *M. utilissima* in Porto Rico, Dominican Republic, Cuba, Ceylon, Siam, Indo-China, the Philippines, and tropical Africa.
- Cercospora manihotis** P. Henn. Circular or effuse dark-fuscous leaf spots on *M. utilissima* in Brazil, China, the Philippines, and central Africa.
- Cercosporella pseudoidium** Speg. Subcircular pale-brown leaf spots on *M. utilissima* in Paraguay and Brazil.
- Colletotrichum lussoniense** Sacc. Anthracnose on *M. utilissima* in the Philippines.
- Colletotrichum manihotis** P. Henn. Circular whitish leaf spots with reddish-brown surrounding zones on *M. utilissima* in Brazil.
- Diplodia cacaoicola** P. Henn. See Theobroma.
- Fomes lamaeensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Haplographium manihoticola** Vinc. Circular brown, zoned leaf spots on *M. glaziovii* in Brazil.
- Macrophoma manihotis** P. Henn. On leaves of *M. utilissima* in Tanganyika.
- Mycosphaerella manihotis** Syd. Circular brown leaf spots on *M. utilissima* in Brazil.
- Phyllosticta manihoticola** Syd. On leaves of *M. dichotoma*, *M. glaziovii*, and *M. heterophylla* in the Philippines.
- Rosellinia arcuata** Petch. See Thea.
- Septogloeum manihotis** Zimm. Brown leaf spots on *M. utilissima* in Java, India, Ceylon, and east Africa. Probably the same as *Cercospora henningsii* Allesch.
- Stagonospora cassavae** Wolk. A wound parasite attacking rooted cuttings of *M. utilissima* in Java.
- Uromyces carthagenensis** Speg. Yellow and dark-brown rust pustules on leaves and stems of *M. carthagenensis* and *M. tweediana* in Argentina.
- Uromyces janiphae** (Wint.) Arth. Rust on leaves of *M. utilissima* in Porto Rico, Cuba, Dominican Republic, Trinidad, and South America.
- Uromyces manihoticola** P. Henn. Powdery dark-brown rust pustules on leaves of *Manihot* sp. in Brazil.
- Uromyces manihotis** P. Henn. Brown to black rust pustules on leaf blades and petioles, stems, and flowers of *M. glaziovii* and *M. utilissima* in Brazil. Produces a witches'-broom effect, as well as galls, on twigs and branches.
- Uromyces manihotis-catingae** P. Henn. Powdery brown rust pustules on leaves of *M. catingae* and *M. glaziovii* in Brazil.
- MAPANIA.** Tropical plants of the sedge family.
- Puccinia mapaniae** Racib. Powdery dark-brown rust pustules on *Mapania* sp. in Java.
- MAPROUNEA.** Tropical trees and shrubs.
- Acidium maprouneae** P. Henn. Rust on irregular, fuscous leaf spots on *Maprounea* sp. in Peru and Brazil.
- Phyllachora maprouneae** P. Henn. Black stromata on yellowish to violet indefinite leaf spots on *M. surinamensis* in Peru.
- Uredo maprouneae** P. Henn. Rust on leaves of *M. guianensis* in Peru.
- MARANTA.** ARROWROOT. Foliage plants.
- Puccinia cannae** (Wint.) P. Henn. See Canna.
- Rosellinia bunodes** B. and Br. See Citrus.
- Sclerotium** sp. See Saccharum.
- Sphaerostilbe repens** B. and Br. Causes a soft rot of the underground stems of *M. arundinacea*. See also Hevea.
- MARATTIA.** Tropical ferns.
- Leptosphaeria caffra** Thuem. Large, more or less circular, leaf spots on *M. salicifolia* in the Union of South Africa.
- MARIANA.** See Silybum.
- MARKHAMIA.** Tropical evergreen trees and shrubs.
- Phragmidiella markhamiae** P. Henn. Brown rust pustules on leaves of *M. zansibarensis* in Tanganyika.
- MARRUBIUM.** HOARHOUND. Aromatic herbs.
- Phyllosticta marrubii** McAlp. Pale-brown irregular leaf spots on *M. vulgare* in Australia.
- Ramularia marrubii** Mass. Small rufous angular leaf spots on *M. album*, *M. creticum*, and *M. vulgare* in Italy, Esthonia, and Yugoslavia.
- Uredo marrubii** Rabh. Rust on leaves of *Marrubium* sp. in Italy.
- MARSDENIA.** Twining shrubs.
- Puccinia aequatoriensis** Syd. Gray-brown rust pustules on leaves of *Marsdenia* sp. in Ecuador.
- Puccinia marsdeniae** Diet. and Holw. Yellow-brown to black rust pustules on leaves of *M. mexicana* in Mexico and Guatemala.
- Septoria asclepiadea** Sacc. See Cynanchum.
- MARTINEZIA.** See Palmae.
- MARTYNIA.** UNICORN PLANT. Annual and perennial plants sometimes cultivated for the peculiar seed pods.
- Cercospora decolor** Pass. Circular gray leaf spots on *M. lutea* in Argentina and Italy.
- Phyllosticta martyniae** Thuem. More or less circular leaf spots on *M. lutea* in Portugal.
- MASDEVALLIA.** See Orchidaceae.
- MATRICARIA.** FALSE CAMOMILE. Annual or perennial weedy herbs.
- Cylindrosporium matricariae** Diet. On leaves of *M. chamomilla* in Germany.
- Entyloma matricariae** Rostr. Smut sori on leaves of *M. discoidea*, *M. inodora*, and *Tripleurospermum disciformis* in Denmark and Sweden.
- Entyloma trailii** Mass. Smut sori in small subcircular to ovate spots on stems and leaves of *M. inodora* in Great Britain.
- Peronospora leptosperma** (De By.) Gäum. See Chrysanthemum.
- Septoria matricariae** Höll. On leaves of *M. chamomilla* and *M. discoidea* in Hungary and Germany.

**MATTHIOLA.** STOCK. Sometimes called gilliflower. Flower-garden plants.

**Ascochyta matthiolae** Oud. On pods of *M. incana* in Holland.

**Bacterium matthiolae** Br. and Pav. Pale-green irregular spots first appear, becoming brown and causing deforming and curling of the leaves. Inflorescences are checked in their development. On *M. annua* in Italy.

**Helminthosporium roridum** Tode. On leaves of *M. incana* in Europe.

**Peronospora matthiolae** Gäum. Downy mildew on leaves of *M. incana* in central and north Europe.

**Puccinia oudemansii** Tranzsch. Powdery brown rust pustules on leaf blades and petioles of *M. nudicaulis* and *Parrya pinnatifida* in Turkestan.

**Sclerotinia matthiolae** Lendner. Causes a stem rot and blight of *M. vallesiaca* in Switzerland. Long black sclerotia form in the rotted stems.

**Septoria henriquesii** Thuem. and f. **santonensis** P. Brun. On leaves of *M. incana* in France and Portugal.

**Urocystis coralloides** Rostr. Irregular gray coral-shaped smut masses on roots of *M. sinuata* and *Turritis glabra* (*Arabis perfoliata*) in France and Denmark.

**MAURITIA.** Fan palms. See Palmae.

**MAXILLARIA.** Epiphytic orchids. See Orchidaceae.

**MAYTENUS.** Trees and shrubs.

**Accidium mayteni** Pazsch. Rust on leaves of *M. brasiliensis* and *Maytenus* sp. in Brazil.

**Heterobotrys paradoxa** Sacc. and subsp. **chilensis** Sacc. and Syd. On leaves of *M. magellanica* in Chile.

**Phyllosticta mayteni** Speg. Leaf spots on *M. magellanica* in Chile.

**Septoria mayteni** Wint. On leaves of *M. magellanica* in Tierra del Fuego.

**Syncarpella missionum** (Speg.) Theiss. and Syd. Black carbonous stromata on living twigs of *M. ligustrina* in Argentina.

**MAZUS.** Low annual or perennial rock-garden plants.

**Accidium foetidum** Diet. Rust on leaves of *M. rugosus* in Japan.

**MEDICAGO.** MEDICK. Alfalfa. Herbaceous perennials cultivated as forage plants.

**Ascochyta affinis** Jaap. Circular to elliptical white leaf spots with brown borders on *M. arabica* in Dalmatia.

**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.

**Laestadia insidiosa** Mass. On leaves of *M. sativa* in Tasmania.

**Leptosphaeria circinans** (Fekl.) Sacc. See Asparagus.

**Marsonia medicaginis** Voss. Circular to elliptical dull-yellow leaf spots on *M. lupulina* in Europe.

**Mitrella sclerotiorum** Rostr. Forms black sclerotia in the roots, infected plants dying. On *M. sativa*, *Lotus corniculatus*, *L. uliginosus*, and *Lotus* sp. in Europe.

**Mycosphaerella destructiva** B. and Br. On leaves of *M. sativa* in Australia.

**Ovularia medicaginis** Br. and Cav. On leaves of *M. sativa* in Italy.

**Peronospora aestivalis** Syd. Downy mildew on leaves of *M. denticulata*, *M. falcata*, *M. lupulina*, *M. media*, *M. minima*, and *M. sativa* in Europe and Australia. Occurs sparingly in the United States.

**Phyllosticta bonanseaana** Sacc. Indefinite yellow areas on leaves of *M. arborea* in Mexico.

**Rhizoctonia destruens** Tassi. See Solanum.

**Rhizoctonia napi** West. See Brassica.

**Septoria medicaginis** Rob. and Desm. Subcircular to irregular whitish leaf spots on *M. lupulina* and *M. sativa* in Argentina, Russia, Sweden, Denmark, Italy, and France.

**Tylenchus dipsaci** Kuehn. This nematode is a serious pest of alfalfa (*M. sativa*), greatly reducing yields by stunting the plants. It has been reported from several localities in the western United States. See Narcissus.

**Typhula trifolii** Rostr. See Trifolium.

**Uredo medicaginicola** Speg. Rust on stems and petioles of *M. sativa* in Argentina.

**Uredo medicaginis** Speg. Rust on leaves of *M. denticulata* in Argentina.

**Uromyces anthyllidis** (Grev.) Schroet. See Anthyllis.

**Urophlyctis alfalfae** (Lagerh.) Magn. The crown-wart of alfalfa (*M. sativa*) is a serious disease in Europe, India, and South America. The disease has been found in limited areas in the United States. It is characterized by warts or galls at the base of the stems. These bodies are irregular in shape, varying in size from a fraction of an inch to 3 or 4 inches, and of a corallike appearance. Affected plants show dead and dying stems with yellow leaves.

**MEDINILLA.** Evergreen shrubs with showy flowers.

**Colletotrichum medinillae** Rangel. Anthracnose on leaves of *M. magnifica* in Brazil.

**Laestadia medinillae** Rangel. Large concentrically zoned leaf spots on *M. magnifica* in Brazil.

**Phyllosticta lageniformis** Rangel. On leaves of *M. magnifica* in Brazil.

**Phyllosticta medinillae** Rangel. On leaves of *M. magnifica* in Brazil.

**MEIBOMIA.** TICK CLOVER. Mostly weedy annual and perennial herbs, partly woody. Some species cultivated for forage.

**Accidium amagense** Mayor. Rust on leaves of *M. tortuosa* in Colombia.

**Cercospora atropunctata** Rac. Circular to irregular, often confluent, brown spots on leaves of *M. umbellata* in Java.

**Phyllachora desmodii** P. Henn. Black stromata on leaves of *M. scalpe* and *Meibomia* sp. in India and central Africa.

**Physopella meibomiae** Arth. Rust on leaves of *M. supina* in Trinidad.

**Uredo amagensis** Mayor. Rust on *M. tortuosa* in Colombia.

**Uredo desmodii-heterocarpi** Petch. Rust on leaves of *M. heterocarpa* in Ceylon.

**Uredo desmodii-parvifolii** Petch. Rust on leaves of *M. parvifolia* in Ceylon.

**Uredo desmodii-pulchelli** Syd. Powdery yellow-brown rust pustules on leaves of *M. pulchella* in Hongkong.

**Uredo desmodii-triquetri** Petch. Rust on leaves of *M. triquetra* in Ceylon.

**Uromyces capitatus** Syd. Rust on leaves of *M. yunnanensis* in China.

**Uromyces castaneus** Syd. Brown rust pustules on leaves of *M. incana* in Brazil.

**Uromyces huallagensis** P. Henn. Brown rust pustules on leaves of *Meibomia* sp. in Peru.

**Uromyces mexicanus** Diet. and Holw. Yellow-brown to black rust pustules on leaf blades and petioles and stems of *M. uncinata* and *Meibomia* sp. in Mexico and Costa Rica.

**Uromyces orbicularis** Diet. Chestnut-brown rust pustules on leaves of *Meibomia* sp. in Brazil.

**Uromyces tenuistipes** Diet. and Holw. Brown rust pustules on leaves of *M. strobilacea* and *Meibomia* sp. in Mexico.

**MELALEUCA.** Ornamental shrubs and trees.

**Laestadia melaleucæ** (Berk.) Sacc. Circular dull-brown leaf spots on *Melaleuca* sp. in Australia.

**Phyllachora melaleucæ** Syd. Shiny black circular to irregular stromata on leaves of *M. spinosa* in Australia.

**Physalospora melaleucæ** (Lév.) Sacc. On leaves of *M. leucadendron* in France.

**Phyllosticta leucadendri** P. Henn. Circular brown leaf spots on *M. leucadendron* in Australia.

**MELASPHAERULA. DIASIA** Ag. TWIN SPUR. Small bulbous plants.

**Uromyces melasphaerulæ** Syd. Brown to black rust pustules on orbicular to irregular spots on both surfaces of leaves of *M. graminea* in the Union of South Africa.

**MELIA. CHINABERRY.** Ornamental trees, also used for windbreaks.

**Cercospora congoensis** Syd. Small olivaceous leaf spots on *Melia* sp. in the Congo.

**Cercospora meliicola** Speg. Small subcircular whitish leaf spots on *M. azedarach* in Argentina.

**Cercospora subsessilis** Syd. Leaf spot on *M. azedarach* in Ceylon, India, China, and the Philippines.

**Gloeosporium meliicola** Speg. On leaves of *M. azedarach* in Argentina.

**Peronoplasmodium portoricensis** Lamk. Irregular, diffuse yellow areas on leaves of *M. azedarach* in Porto Rico, causing premature leaf fall.

**Phyllachora meliæ** Pat. Small circular shiny black stromata on leaves of *M. azedarach* in Indo-China.

**Septobasidium acaciæ** Saw. See Acacia.

**Septoria meliæ** Syd. Small circular white leaf spots on *M. azedarach* in the Union of South Africa.

**MELICA. MELIC GRASS.** Perennial grasses.

**Ascochyta phyllachoroides** Sacc. and Malbr. On leaves of *M. altissima* in France.

**Cintractia melicæ** (Sorok.) Det. Smut on ovaries of *M. ciliata* in Central Asia.

**Phyllachora meliicola** Speg. Tar spot on *M. violacea* in Argentina.

**Physalospora montana** Sacc. On leaves and stems of *Melica* sp. and *Sesleria caerulea* in Italy, Spain, and Switzerland.

**Puccinia heimerliana** Bub. Yellow-brown to black rust pustules on leaves of *M. ciliata* and *M. cupani* in Russia, Italy, and Switzerland.

**Puccinia rangiferina** S. Ito. Black rust on leaves and sheaths of *Melica* sp. and *Calamagrostis arundinacea* in China and Japan.

**Puccinia trebouxii** Syd. Rust on leaves of *M. ciliata* in Russia.

**Septoria melicæ** Pass. Rufous leaf spots on *M. picta* and *M. uniflora* in Europe.

**Septoria simplex** (Schroet.) Sacc. On *M. nutans* in Lapland and Sweden.

**Uredo jozankensis** Ito. Golden rust sori on dark-brown leaf spots on *M. nutans* in Japan.

**Urocystis bornmulleri** Magn. Smut on leaves and inflorescences of *M. cupani* in Syria.

**Uromyces graminis** (Niessl.) Diet. See Seseli.

**Uromyces seseli-graminis** Ed. Fisch. See Arrhenatherum.

**Ustilago trebouxii** Syd. Dark-brown linear smut sori in leaves of *M. ciliata* and *Triticum (Agropyron) cristatum* in Russia.

**MELILOTUS. SWEET CLOVER.** Forage plants.

**Cercospora meliloti** Oud. Circular to oblong white leaf spots on *M. officinalis* in Russia, Italy, and Holland.

**Entyloma meliloti** McAlp. Small circular slightly raised pustules producing large, often confluent, patches on leaves of *M. indica* in Australia.

**Peronospora meliloti** Syd. Downy mildew on leaves of *M. alba*, *M. altissima*, *M. indica*, *M. officinalis*, and *M. parviflora* in India and Europe.

**Septoria meliloti** (Lasch.) Sacc. Small subcircular pale leaf spots on *M. alba*, *M. officinalis*, and *M. vulgaris* in Italy, Russia, and Germany.

**Stagonospora carpathica** Baeuml. See Vicia.

**Uromyces baeumlerianus** Bub. Brown rust pustules on leaves and stems of *M. neapolitana* and *M. officinalis* in Spain and Russia.

**MELIOSMA.** Trees and shrubs grown for their handsome foliage and large panicles of white flowers.

**Aecidium meliosmae-myrianthæ** P. Henn. and Shir. Rust on leaves of *M. myriantha* and *M. simplicifolia* in Japan and India.

**Aecidium meliosma-pungentis** P. Henn. and Shir. Rust deforming leaf blades and petioles of *M. pungens* in Japan.

**Gophana mirabilis** Rac. Rust on leaves of *Meliosma* sp. in Java.

**Phakopsora meliosmae** Kusano. Yellow to brown rust pustules on leaves of *M. myriantha* in Japan and the Philippines.

**Phyllachora meliosmae** Racib. Circular black stromata on leaves of *Meliosma* sp. in Java.

**Uredo meliosmae** Petch. Leaf rust on *M. wightii* in Ceylon.

**MELISSA. BALM.** Hardy perennial herbs.

**Cercospora sennensis** Gz. Frag. Leaf spot on *M. officinalis* in Spain.

**Cylindrosporium melissæ** C. Mass. Irregular leaf spots on *M. officinalis* in Spain and Italy.

**Phleospora melissæ** (Desm.) Parisi. Numerous small angular to irregular brown spots causing leaves to curl up and fall. On *M. officinalis* in Italy, France, and Spain.

**Phyllosticta latemarensis** Kab. and Bub. On leaves of *M. officinalis* in Hungary.

**Phyllosticta melissæ** Bub. Small circular brown leaf spots on *M. officinalis* in Hungary.

**MELITTIS.** Perennial herbs, used for border plantings.

**Phyllosticta montellica** Sacc. Circular or angular dark-red leaf spots, becoming whitish, on *M. melissophyllum* in Italy.

**Septoria melittidis** Sacc. and Speg. Whitish indefinite leaf spots on *M. grandiflora* and *M. melissophyllum* in France and Italy.

**MELOTHRIA.** Slender herbaceous annual or perennial vines.

**Puccinia melothriicola** Syd. Leaf rust on *M. mucronata* in the Philippines.

**Uredo cantonensis** Yates. Rust on leaves of *M. indica* in China.

**Uredo melothriæ** P. Henn. Rust on leaves of *M. tomentosa* in Abyssinia.

**Uromyces hellerianus** Arth. Brown rust sori on *M. guadalupensis* and *M. scabra* in Porto Rico, Cuba, and Guatemala.

**MENISPERMUM. MOONSEED.** Hardy, attractive semiwoody climbers.

**Ascochyta jenissensis** Sacc. Subcircular pale-brown leaf spots on *M. dahuricum* in Siberia.

**Gloeosporium menispermum** Allesch. Anthracnose on leaves of *M. canadense* in Germany and Monaco.

**Guignardia menispermum** (Speg.) Theiss. and Syd. On leaves of *Menispermum* sp. in Brazil and Paraguay.

**Phyllosticta maculiformis** var. **menispermum** Sacc. On leaves of *M. canadense* in France.

**Septoria menispermum** Thuem. Ochraceous leaf spots with gray-brown margins on *M. dahuricum* in Siberia and Russia.

- MENTHA.** MINT. Strong-scented perennial herbs.  
**Peronospora stigmaticola** Raunk. Downy mildew on flowers of *M. aquatica* and *M. arvensis* in Denmark, Sweden, and Russia.  
**Phyllosticta menthae** Bres. On leaves of *M. arvensis* in Germany.  
**Physoderma menthae** Schroet. On stems and occasionally leaves of *M. aquatica* in Germany.
- MENTZELIA.** BLAZING STAR. Showy flowered plants, sometimes cultivated.  
**Coleosporium mentzeliae** (Diet. and Holw.) Arth. Powdery golden rust sori on leaves of *M. hispida* in Mexico.
- MENZIESIA.** Low deciduous shrubs allied to Rhododendron.  
**Chrysomyxa menziesiae** Diet. Numerous brown rust pustules on leaves of *M. pentandra* in Japan.
- MERIANA.** See *Watsonia*.
- MESEMBRYANTHEMUM.** FIG MARIGOLD. Low-growing succulents.  
**Peronospora mesembryanthemi** Verwoerd. Downy mildew on leaves, stems, flower stalks, and calyces of *Mesembryanthemum* spp. in the Union of South Africa.  
**Puccinia mesembryanthemi** McOwan. Golden and black rust pustules on leaves of *M. micranthum* in the Union of South Africa.  
**Septoria confluens** McAlp. Gray-white leaf spots on *M. aequilaterale* in Australia.
- MESPILUS.** MEDLAR. Woody plants cultivated for fruit.  
**Ascochyta mespili** Pass. Irregular brown, then gray or gray-brown, leaf spots on *M. germanica* in France.  
**Entomosporium mespili** (DC.) Sacc. See *Cotoneaster*.  
**Gymnosporangium mespili** (DC.) Kern. See *Juniperus*.  
**Laestadia mespili** Fautr. On leaves of *M. germanica* in France.  
**Ovularia necans** (Pass.) Sacc. See *Cydonia*.  
**Phyllosticta mespilicola** Rota-Rossi. Circular to irregular large dull-brown leaf spots on *M. germanica* in Italy.  
**Phyllostictiella mespili** (Sacc.) Tass. Subcircular, ochraceous leaf spots with rufous margins on *M. germanica* in Denmark, Malta, and Italy.  
**Sclerotinia mespili** Schell. This fungus causes dark-brown leaf blotches from 1 centimeter in diameter up, as well as brown rot and mummification of the fruit of *M. germanica* and *Cydonia* sp. in Switzerland and Great Britain.  
**Septoria mespili** Sacc. Ochraceous leaf spots with rufous margins on *M. germanica* in France, Malta, and Italy.
- METROSIDEROS.** IRON TREE. Trees and shrubs planted for their showy flowers.  
**Phyllosticta metrosideri** P. Brun. Circular pale gray-brown leaf spots with red-brown margins on *M. villosa* in Tahiti.  
**Phyllosticta salicifolia** P. Brun. Round to irregular gray leaf spots with brown borders on *M. salicifolia* in France.  
 A serious disease, due to an unknown cause, destroyed large areas of *M. polymorpha* in Hawaii.
- MEUM.** Perennial herbs with delicate ornamental foliage.  
**Accidium mei** Schroet. Rust on leaf blades and petioles of *M. athamanticum* and *M. mutellinum* in central Europe.  
**Protomyces macrosporus** Ung. See *Coriandrum*.
- MEZONEURUM.** Large, usually armed, climbing shrubs.  
**Ravenelia aculeifera** Berk. Rust on leaves of *M. enneaphyllum* in Ceylon.
- MICHELIA.** Trees and shrubs with magnoliolike foliage and fragrant flowers.  
**Phyllachorella micheliae** Syd. Black circular stromata on leaves of *M. nilagirica* in India.  
**Phyllostictiella micheliae** (P. Henn.) Tassi. On leaves of *M. fuscata* in Germany.
- MICONIA.** Tropical trees and shrubs with large, showy, strongly-veined leaves.  
**Catacauma pululahuensis** (Pat.) Theiss. and Syd. Tar spot on leaves of *Miconia* sp. in Brazil.  
**Catacaumella miconiae** (P. Henn.) Theiss. and Syd. Shiny black stromata on leaves of *Miconia* sp. in Brazil.  
**Cronartium egenulum** Syd. Yellowish-red rust sori on leaves of *M. theezans* in Japan and Brazil.  
**Cyclothea miconiae** (Syd.) Theiss. Fruiting bodies form on red-brown, then black, spots on lower leaf surfaces of *M. rigidiuscula* in Brazil.  
**Dothidina fiebrigii** (P. Henn.) Theiss. and Syd. Black fruiting bodies on small brown spots on *Miconia* sp. in Paraguay.  
**Dothidina miconiae** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *M. laevigata*, *M. prasina*, and *M. sintenisii* in Porto Rico and Brazil.  
**Dothidina peribebuyensis** (Speg.) Chardon. Small hemispherical dark-brown fruiting bodies on brown spots on leaves of *M. calvescens*, *M. laevigata*, *M. lepidota*, *M. prasina*, *M. sintenisii*, *Lasiandra obscura*, and *Tibouchina multiceps* in Porto Rico, Colombia, and Brazil.  
**Hysterostromina miconiae** (P. Henn.) Theiss. and Syd. Thick dull-black stromata on leaves of *Miconia* sp. in Brazil.  
**Phaeofabraea miconiae** Rehm. On leaves of *Miconia* sp. in Brazil.  
**Phylachora miconiicola** Speg. On leaves of *Miconia* sp. in Costa Rica.  
**Physalospora miconiae** (Duby.) Sacc. On leaves of *M. calvescens* in Brazil.  
**Rosellinia bunodes** B. and Br. See *Citrus*.  
**Septoria miconiae** Garm. Brown leaf spots on *M. impetiolearis* and *M. laevigata* in Porto Rico.
- MICROCITRUS.** Small spiny trees allied to *Citrus*.  
**Bacterium citri** Hasse. See *Citrus*.
- MICROGLOSSA.** Shrubs and climbers.  
**Uredo microglossae** Petch. Leaf rust on *M. Zeylanica* in Ceylon.
- MICROSTYLIS.** See *Orchidaceae*.
- MIKANIA.** Climbing hempweed. Shrubs or twining herbs.  
**Accidium mikaniae** P. Henn. Round dull-brown leaf spots on *M. confertissima* in Brazil.  
**Apiospora pachyspora** Rehm. On leaves of *Mikania* sp. in Brazil.  
**Ascomycetella punctoidea** Rehm. Sessile black stromata on leaves of *M. vismiaefolia* in Brazil.  
**Cercospora** sp. Leaf spot on *M. scandens* in Fiji.  
**Cercospora mikaniaeicola** Stevens. Small circular white leaf spots with dark-brown margins on *Mikania* sp. in Porto Rico.  
**Chrysopsora mikaniae** Arth. Yellow rust pustules on leaves of *M. buddleiaefolia* and *Mikania* sp. in Brazil and Bolivia.  
**Endophylloides portoricensis** Whetz. and Olive. Yellow rust sori on leaves of *M. cordifolia*, *M. odoratissima*, and *M. scandens* in Porto Rico, Guatemala, Trinidad, and Brazil.  
**Gibbera mikaniae** (P. H.) Rick. and Theiss. Black stromata on leaves of *Mikania* sp. in Brazil.  
**Mollisia mikaniae** Rehm. Indefinite yellow leaf spots on *M. discolor* in Brazil.  
**Septoria mikaniae** Wint. Circular to irregular leaf spots on *Mikania* sp. in Porto Rico and Brazil.  
 Reported from Texas.

**MILIUM.** MILLET GRASS. Perennial grass.

**Laestadia microspora** (Awd.) Sacc. On leaves of *M. effusum* in Germany.

**Leptosphaeria nigrans** (Desm.) Ces. and De N. See Aira.

**Puccinia milii** Erikss. Rust on leaves of *M. effusum* and *M. paradoxum* in Sweden, Norway, Denmark and Spain.

**Puccinia piptatheri** Laegerh. Yellow-brown to dark-brown rust pustules on leaves of *M. multiflorum* in Portugal.

**MILLETTIA.** Tropical trees producing hard, durable timber.

**Bacillus milletiae** Kawakami. Irregular, hard, woody knots, 5 to 10 millimeters in diameter and up to 10 centimeters long, for the most part on old stems of *M. floribunda* var. *brachybotrys* in Japan.

**Diorchidium woodii** (Kalchbr. and Cke.) Syd. Black powdery rust sori on leaves of *M. caffra* in Natal.

**Microcyclus angolensis** Sacc. and Syd. Irregular black stromata on leaves of *M. thonningii* in Angola.

**Phyllachora luzonensis** P. Henn. Tar spot on leaves of *M. cavitiesis*, *M. merrillii*, and *Derris philippinensis* in the Philippines.

**Phyllachora milletiae** P. Henn. Black stromata on red-brown leaf spots on *Millettia* spp. in the Congo and Tanganyika.

**Phyllosticta milletiae** P. Henn. Circular to angular and confluent pale-brown leaf spots on *Millettia* sp. in the Congo.

**Physalospora congensis** P. Henn. Circular effuse pale-brown leaf spots on *Millettia* sp. in the Congo.

**MILTONIA.** Orchid. See Orchidaceae.

**MIMOSA.** SENSITIVE PLANT. Woody or herbaceous tropical plants grown for their showy flowers and feathery foliage.

**Cercospora mimosae** Pat. Circular dull-brown leaf spots on *M. floribunda* in Ecuador.

**Cercospora sensitivae** Speg. Rufous spots on leaves of *M. sensitiva* in Argentina.

**Mycosphaerella mimosicola** P. Henn. On leaves of *M. asperata* in Brazil.

**Phyllachora michelii** Speg. Black stromata on indefinite brown leaf spots on *M. procurrens* in Brazil.

**Phyllachora rhytmoides** (Cda.) Sacc. Tar spot on leaves of *Mimosa* sp. in India.

**Puccinia mimosae** (Speg.) Syd. Dark cinnamon rust sori on young twigs and inflorescences of *Mimosa* sp. in Argentina.

**Ramularia mimosae** Stev. and Dal. Indefinite spots, often involving entire area of leaflets of *M. pudica* in Porto Rico.

**Ravenelia bahiensis** P. Henn. A rust deforming the branches of *M. remansoana* in Brazil.

**Ravenelia mainsiana** Arth. and Holw. Brown to black rust pustules on leaves of *M. albida* in Guatemala.

**Ravenelia mimosae-albidae** Diet. Rust on leaves of *M. albida* in Mexico, Costa Rica, and Guatemala.

**Ravenelia mimosae-caeruleae** Diet. Cinnamon to dark-brown rust pustules on leaves of *M. caerulea* in Mexico.

**Ravenelia mimosae-sensitivae** P. Henn. Brown rust pustules on leaves of *M. sensitiva* in Colombia and Argentina.

**Ravenelia mimosicola** Arth. Brown rust pustules on leaves of *M. polyanthoides* and *M. stipitata* in Mexico.

**Uromycladium cubense** Arth. and Johnst. Leaf rust on *M. asperata* in Cuba.

**MIMUSOPS.** Tropical trees with milky juice, some producing edible fruit.

**Phyllosticta mimusopsidis** P. Henn. Red-brown leaf spots on *M. schimperii* in Abyssinia.

**Phyllosticta mimusopsis** Cufino. Circular to irregular white leaf spots on *M. crassifolia* in Italy.

**Physalospora placida** Syd. Circular gray-brown leaf spots with raised purple-brown margins on *Mimusops* in Portuguese East Africa.

**Uromyces mimusops** Cke. Brown rust pustules on leaves of *M. caffra* and *M. elengi* in India, Portuguese East Africa, and the Union of South Africa.

**MIRABILIS.** FOUR-O'CLOCK. Erect herbs.

**Aecidium mirabilis** Diet. and Holw. Rust sori on irregular yellow leaf spots on *Mirabilis* sp. in Mexico.

**Gloeosporium mirabilis** Bres. Anthracnose on stems of *M. jalapa* in Portugal.

**MISCANTHUS.** Tall perennial grasses with large terminal, feathery, fan-shaped panicles.

**Didymosphaeria striatospora** Syd. On stems and leaf sheaths of *M. sinensis* in Japan.

**Ephelis japonica** P. Henn. Black stromata forming between the compressed spikelets, deforming them. On *M. tinctorius* and *Paspalum thunbergii* in Japan.

**Phyllachora miscanthi** Syd. Black stromata on leaves of *M. sinensis* and *Miscanthus* sp. in China and the Philippines.

**Puccinia erythropus** Diet. See Calamagrostis.

**Puccinia eulaliae** Barcl. Purple-brown rust pustules on leaves of *M. condensatus*, *M. sacchariflorus*, *M. sinensis*, and *Pollinia japonica* in Japan and India.

**Puccinia miyoshiana** Diet. Brown to black rust pustules on leaves of *M. cotulifer* and *M. sibericus* in Japan and Siberia.

**Sclerospora** sp. See Saccharum.

**Sclerospora philippinensis** Weston. 'See Zea.

**Sphacelia miscanthi** P. Henn. Black elongate sclerotia formed in ovaries of *M. sinensis* in Japan.

**Ustilago kusanoi** Syd. Dark-olive smut sori destroying the panicles and rachides of *M. sacchariflora* and *M. sinensis* in Japan and the Philippines.

**MITELLA.** BISHOP'S CAP. Low perennial herbs with creeping rootstocks.

**Puccinia asiatica** (Kom.) Syd. Rust on leaves of *M. nuda* and *Tiarella polyphylla* in Asia.

**MOLDAVICA.** See Dracoecephalum.

**MOMORDICA.** BALSAM APPLE. Annual or perennial tropical climbing herbs, cultivated for ornament and fruit.

**Aecidium momordicae** Juel. Rust on leaves of *Momordica* sp. in Brazil.

**Ascochyta elaterii** Sacc. Indefinite yellow spots on leaves of *Ecballium elaterium* (*M. elaterium*) in Italy.

**Cercospora elaterii** Sacc. On leaves of *M. (Ecballium) elaterium* in Italy.

**Leandria momordicae** Rangel. Circular, then confluent, gray leaf spots with narrow pale-brown margins on *M. charantia* in Brazil.

**Phyllosticta momordicae** F. Tassi. Effuse grayish-white leaf spots on *M. involucrata* in Italy.

**Puccinia momordicae** Kalchbr. and Cke. Powdery brown rust pustules on leaves of *M. cordifolia* in the Union of South Africa.

**Uredo momordicae** Petch. Rust on leaves of *M. charantia* in Ceylon.

**MONADENIA.** Orchid. See Orchidaceae.

- MONTANOA.** Shrubs, used for subtropical bedding.  
**Accidium montanoae** Diet. and Holw. Rust on petioles and branches of *Montanoa* sp. in Mexico.  
**Uromyces montanoae** Arth. and Holw. Cinnamon-brown rust pustules on leave of *M. dumicola*. and *Montanoa* sp. in Costa Rica.
- MONSTERA.** Ceriman. Root climbing evergreen tropical aroids with edible fruit.  
**Neohenningsia brasiliensis** P. Henn. On leaves of *Monstera* sp. in Brazil.
- MONTIA.** Small annual and perennial herbs, one species used as a salad plant.  
**Cercospora montiae** Rostr. Blackened areas, often over entire surface of leaves of *M. minor* in Denmark.  
**Sorosporium montiae** Rostr. Black smut sori in leaves and stems of *M. minor* in Denmark.  
**Tolyposporium junci** (Schroet.) Wor. Smut in ovaries, peduncles, and culms of *M. minor* and *Juncus* spp. in Denmark and Germany.
- MORAEA.** MOREA. Bulbous plants resembling *Iris*.  
**Phyllosticta macrothecia** (Thuem.) Gz. Frag. A leaf spot on *M. grandiflora* in Spain.  
**Puccinia moraeae** P. Henn. Small brown rust pustules on both surfaces of leaves of *M. edulis* in the Union of South Africa.  
**Uromyces geissorhizae** P. Henn. See Geissorhiza.  
**Uromyces moraeae** Syd. Rust on leaves of *M. spathacea* in the Union of South Africa.
- MORINDA.** INDIA MULBERRY. Woody plants.  
**Cercospora morindae** Syd. Circular dull-yellow leaf spots on *M. tinctoria* in India.  
**Physalospora morindae** Koord. Circular or oblong yellow leaf spots on *M. citrifolia* in Java.
- MORINGA.** HYPERANTHERA Ag. HORSE-RADISH TREE. Tropical ornamental trees.  
**Puccinia moringae** Koord. Rust on pale-yellow leaf spots on *M. oleifera* (*M. pterygosperma*) in Java.
- MORUS.** MULBERRY. Ornamental trees cultivated for their fruit and the leaves used for silkworm rearing.  
**Accidium mori** Syd. and Butl. Yellow rust sori on leaves of *M. alba* in the Philippines, India, Java, Japan, Formosa, and China.  
**Accidium moricola** P. Henn. Rust on leaves of *M. indica* in Java.  
**Ascochyta miyabei** Tan. On branches of *M. alba* in Japan.  
**Ascochyta mori** Maire. On leaves of *M. alba* in France.  
**Cercospora snelliana** Reich. Attacks leaves of *M. alba* in Egypt.  
**Clasterosporium mori** Syd. Brown leaf spots on *M. alba* in China and Japan.  
**Clasterosporium putrefaciens** Frank. On leaves of *M. alba* and *Vitis vinifera* in the Caucasus Mountains.  
**Coryneum mori** Mon. Cankers are produced on trunks and branches, causing a die-back and stunting of trees of *Morus* spp. in Japan and India. It is particularly serious on small trees in nurseries.  
**Cryptosporium mori** Fekl. On leaves of *M. alba* in Germany.  
**Cylindrosporium maculans** Bereng. On leaves of *M. alba* and *M. nigra* in Russia, Poland, Yugoslavia, Albania, and Denmark.  
**Cytodiplospora mori** Miy. Parasitic on twigs of *M. alba* in Japan.  
**Didymella mori** K. Hara. On twigs of *M. alba* in Japan.  
**Gibberella moricola** (Ces. and De N.) Sacc. On branches of *M. alba*, *M. nigra*, and *Broussonetia* sp. in China, Italy, Austria, Hungary, and Germany.  
**Helicobasidium monpa** Tan. and **forma macrosporum** Hara. The fungus attacks the roots which become covered with a network of purple-brown mycelial strands. Growth of infected plants is stopped, twigs and branches die back, and any new leaves formed are small. Death of the infected plants finally results. The hosts are *Beta vulgaris*, *Chamaecyparis obtusa*, *C. pisifera*, *Cryptomeria japonica*, *Cydonia japonica*, *Daucus carota*, *Diospyros kaki*, *Edgeworthia chrysantha*, *Hevea brasiliensis*, *Ipomoea batatas*, *Morus* spp., *Photinia glabra*, *Prunus domestica*, *P. pseudocerasus*, *Pyrus japonica*, *Quercus acuta*, *Q. glandulifera*, *Q. serrata*, *Rhus vernicifera*, *Salix* spp., *Sambucus racemosa*, and *Thuya* sp. in Japan, Uganda, and Straits Settlements.  
**Helicobasidium tanakae** Miy. Circular, then large and irregular, velvety, thick brown fungus layers on trunks and twigs of *Firmiana platanifolia*, *Grossularia*, *Juglans*, *Kerria japonica*, *Malus*, *Morus*, *Paulownia tomentosa*, *Pittosporum undulatum*, *Prunus armeniaca*, *P. mume*, *P. salicina*, *Pyrus sinensis*, *Salix*, *Thea sinensis*, *Vitis*, and *Zanthoxylum* in Japan. Said to be parasitic.  
**Hyopchnus cucumeris** Frank. See Cucumis.  
**Massaria phorcioides** I. Miy. On twigs of *M. alba* in Japan.  
**Mycosphaerella maculiformis** (Pers.) Schroet. See Acer.  
**Mycosphaerella mori** (Pass.) Fekl. Irregular pale-brown leaf blotches, with red-brown margins and fruit spots on *M. alba*, *M. nigra*, and *M. rubra* in Europe, South America, Japan, India, Uganda, and Canada. Infected leaves turn yellow and fall.  
**Mycosphaerella morifolia** Pass. On leaves of *M. alba* in China and Italy. Reported from Indiana.  
**Ovulariopsis moricola** Delacr. Effuse whitish leaf spots on *M. alba* in Madagascar and China.  
**Phoma mororum** Sacc. Attacks twigs of *M. alba* in Italy, "often causing great injury."  
**Phyllosticta berlesiana** Allesch. and var. **socialis** Ferr. Brown leaf spots on *M. alba* in Italy.  
**Phyllosticta consors** Sacc. Brown leaf spots on *M. alba* in Mexico.  
**Phyllosticta kuwacola** K. Hara. The "circle blotch of mulberry" (*M. alba*) in Japan is characterized by minute brown leaf spots, circular to irregular, increasing in size and becoming rufous, then ashen, and concentrically zoned. The leaves are finally destroyed. Diseased twigs become brown and die. On older twigs reddish spots occur, which become black and sunken.  
**Phyllosticta langarum** Ferr. Round to irregular, whitish leaf spots with brown margins on *M. alba* in Italy.  
**Phyllosticta morifolia** Pass. Dirty-gray leaf spots with dull chestnut-brown margins on *M. alba* and *M. nigra* in Russia and Italy.  
**Phyllosticta osteospora** Sacc. Rufous leaf spots on *M. alba*, *M. nigra*, *Frazinus* sp., *Populus nigra*, and *Rhamnus* sp. in Italy and France.  
**Physalospora minuta** Miy. Slightly raised spots on living twigs of *M. alba* in Japan.  
**Sclerotinia shiraiana** P. Henn. Forms sclerotia in fruit of *M. alba* and *M. rubra* in Japan and Formosa.  
**Sphaeropsis mori** Berl. Attacks *Morus* in Italy.  
**Stagonospora mori** I. Miy. This fungus attacks twigs of *M. alba* in Japan, causing them to disintegrate.  
**Thyrococcum sirakoffi** Bub. Causes a die-back of twigs and cankers on trunk and limbs of *Morus* spp. in central Europe.  
**Uncinula mori** Miy. Powdery mildew on lower leaf surfaces of *M. alba* in Japan and China.  
**Ustilago haesendockii** West. A very doubtful species of smut reported on roots of *M. alba*, *M. nigra*, and *M. rubra* in Belgium, Italy, and Argentina.  
**A disease** due to an unknown cause appears yearly on *Morus* in Indo-China, causing the curling, drying up, and premature fall of the leaves, and forming swellings on the trunk.

**MOSCHOSMA.** Showy-flowered small shrubs or tall herbs.

**Aecidium moschosmatis** P. Henn. Rust on petioles and stems of *M. multiflora* in east Africa.

**MUCUNA.** Annual beanlike vines, grown as forage and green manure crops.

**Cercospora mucunae** Syd. Brown circular to irregular leaf spots on *M. urens* and *Mucuna* sp. in Brazil and the Philippines. Reported from Alabama.

**Sphaerophragmium mucunae** Rac. Brown rust pustules on leaves of *Mucuna* sp. in Java.

**Uromyces illotus** Arth. and Holw. Dark-brown powdery rust pustules on leaves of *M. andreana* in Guatemala.

**Uromyces mucunae** Rabh. (*Uredo mucunicola* P. Henn.). Brown to black powdery rust pustules on lower leaf surfaces of *M. capitata*, *M. pruriens*, *Stizolobium deeringianum*, *S. lyoni*, and *S. niveum* in the Congo, India, Ceylon, Japan, Formosa, and the Philippines.

**MUEHLENBECKIA. CALACINUM** Ag. Climbing or erect plants.

**Cercospora platensis** Speg. On leaves of *M. sagittata* in Argentina.

**Gloeosporium muehlenbeckiae** Briosi and Cav. Anthracnose on pseudo-leaves of *M. platyclada* in Italy.

**Puccinia muehlenbeckiae** (Cke.) Syd. Brown rust pustules on leaves of *M. adpressa* and *M. gracillima* in Australia.

**Septoria muehlenbeckiae** F. Tassi. Leaf spot on *M. complexa* in Italy.

**Uromyces politus** (B. and Br.) McAlp. See Jacksonia.

**MULGEDIUM.** Annual, biennial, or perennial herbs, allied to *Lactuca*.

**Aecidium minussense** Thuem. Rust on leaves of *Mulgedium* sp. in Siberia.

**Ectostroma mulgedii** Thuem. Black, often confluent, spots on stems of *M. sibiricum* in Siberia.

**Ovularia mulgedii** Bub. Indistinct, irregular yellow leaf spots with brown centers on *M. alpinum* in Yugoslavia.

**Puccinia minussensis** Thuem. Brown rust pustules on leaves of *M. sibiricum* and *M. tataricum* in Siberia, Russia, Finland, and Sweden.

**Puccinia mulgedii** Syd. Yellow and brown rust pustules on leaves of *Mulgedium* spp. in Europe and Siberia.

**Puccinia tatarica** Tranzsch. Rust on stems and leaves of *M. tataricum* in Turkestan and Tibet.

**Septoria mulgedii** Thuem. Large irregular gray leaf spots with gray-brown margins on *M. sibiricum* in Siberia.

**MUSA. BANANA. ABACA. PLANTAIN.** Tropical fruit and fiber plants.

**Bacillus** sp. (?). A bacterial stem and bud-rot of banana and abacá has been reported from the Philippines. It is characterized by a soft black rot of the terminal bud, which extends down the central cylinder and has a disagreeable odor. Entire fields are reported to have been destroyed by this disease.

**Bacillus musae** Rorer. Causes a serious bacterial disease of the banana and plantain in Trinidad and probably in British Guiana. Lower leaves are first attacked, the petioles giving way, causing the blades to droop and turn yellow. The disease progresses upward, finally involving the terminal bud, after which the plants rot down to the ground. Infected plants show discolored vascular bundles at all stages. Even plants not severely diseased fail to produce fruit.

**Bacillus musarum** Zeman. Another bacterial disease reported as serious on varieties of bananas in Argentina, Paraguay, and probably Brazil. Leaves of infected plants turn yellow and droop, the petioles lose their turgidity and turn black. In severe cases further terminal growth ceases, and the stem dries out and dies from the apex downward. Suckers may start, but soon succumb. Vascular bundles are discolored and exude a yellow liquid when cut, which turns to red and finally black upon exposure. This disease may be identical with the preceding.

**Bacterium celebensis** Gäum. This bacterial disease, called the blood disease, is reported as a very serious enemy of the banana (*Musa*) in the Celebes of the Dutch East Indies. Its progress is at first slow, showing as a discoloration of one of the younger leaves, with yellow-brown stripes spreading out from the midrib. The remainder of the plant is attacked suddenly at fruiting time, and within a few days the leaves turn yellow, droop, and the bunch sags down and decays. A reddish slime exudes from the bundles when they are cut across. The fruits turn yellow and become filled with a slimy brownish-red fluid.

**Bacterium musae** Gäum. A second bacterial disease of the banana (*Musa*) reported from the Dutch East Indies, in which the heart leaf stops growing and the outer leaf sheaths split longitudinally, followed by a premature breaking down and wilting of the leaves. The vascular bundles are discolored.

**Bunchy-top.** A disease of the banana, plantain, and abacá due to an unknown cause, possibly of the group of infectious mosaics, reported as serious in Australia, Fiji, Bonin Islands, Ceylon, and Egypt. Other hosts are *Canna* and *Saccharum officinarum*. The disease checks the growth of infected plants, the leaves bunching together at the top, due to the stunting of the leaf stalks. Diseased plants never produce fruit, remaining stunted. The leaves are smaller and lighter in color than normal and very brittle, becoming ridged or corrugated with age. The bulbs show small brown or yellow flecks internally. The roots are for the most part dead, except immediately adjoining the base of the plant.

**Cercospora musae** A. Zimm. Indefinite brown leaf spots on *Musa* in Fiji, Java, and Ceylon. Reported as a "dangerous disease."

**Cercospora musarum** Ashby. Found on leaves of *Musa* in Jamaica in connection with a black spotting disease.

**Coccomyces musae** (Lév.) Sacc. On leaves of *Musa* in Central America.

**Fusarium cubense** E. F. Sm. The "Panama" or wilt disease of *Musa* spp. is the most serious disease known of this important tropical fruit. While only a few varieties, including the Gros Michel, are subject to attack, they are unfortunately the most important ones economically. The disease occurs in Central and South America, Mexico, the West Indies, Hawaii, the Philippines, and probably in India, Java, and Australia. Thousands of acres of banana land have been abandoned, due to the ravages of this disease in Central America. The entire export banana industry of Surinam was destroyed.

A diseased plant first shows a yellowing of the lower or outer leaves and petioles. The transition from normal green to yellow is sudden and proceeds from the margin inward. Diseased leaves wilt rapidly and break over a few inches from the stem and hang downward. Finally, all the leaves are involved, the innermost folded leaf resisting longest. The plant remains with dry brown, hanging leaves for a time until blown over. In severe cases young plants are attacked and remain stunted, finally dying, as do older plants. A longitudinal splitting of the outer leaf sheaths accompanies this stage. Fruit is rarely produced and is always abnormal, small, yellow, and bitter tasting. The vascular bundles are discolored, yellow at first, gradually darkening, and in advanced stages black, the blackening involving all tissues. The disease is commonly carried in offshoots.

**Guignardia musae** Racib. Irregular reddish-brown leaf spots on *M. paradisiaca* in Java.

**Hainesia tallingii** Koord. Purple, irregular, sunken leaf spots on *Musa* sp. in Java.

**MUSA**—Continued.

- Heart rot.** The abaca plant (*M. textilis*) is seriously attacked by a heart-rot disease which threatens the Manila hemp industry in parts of the Philippines. This disease has doubtfully been attributed to *Fusarium cubense* Sm. The terminal bud of infected plants develops a black rot with a disagreeable saline odor. Various varieties of edible bananas are also subject to the disease.
- Laestadia musae** Syd. On leaves of *Musa* sp. in the Congo.
- Leptosphaeria musarum** Sacc. and Berl. On leaves of *M. paradisiaca* in Ceylon and Portuguese St. Thomas.
- Marasmius semiustus** B. and C. This gill fungus has been reported from the West Indies, Trinidad, Hawaii, and Mauritius as the cause of a root disease of the banana, and an extensive literature has developed. Its status is much the same as *M. sacchari* on *Saccharum*. (q. v.).
- Mycosphaerella musae** (Speg.) Syd. Leaf spot on *M. cavendishii*, *M. sapientum* and *M. textilis* in Guam and the Philippines.
- Phoma musae** Carpenter. The "freckle" disease is serious on *M. cavendishii* in Hawaii and has been reported on *Musa* from the Philippines. The fruit spots are at first minute, circular, gray, with indefinite borders, becoming larger and slightly erumpent. Similar spots occur on the leaves, at times causing defoliation. Diseased fruit are undersized, discolored, and ripen unevenly, and even when only slightly attacked are rendered unfit for export.
- Phoma musarum** Cke. On *Musa* in Bermuda and Spain.
- Phyllachora musae** (Kl.) Sacc. On leaves of *Musa* sp. in India and the Congo. Referred by Theissen to *Macrophoma musae* (Sacc.) Berl. and Vogl.
- Phyllostictiella gastoni** (Roum.) Tassi. On leaves of *M. sapientum* in Haiti.
- Sphaerostilbe musarum** Ashby. The so-called "bonnygate" disease of Jamaican bananas is attributed to this fungus. Leaves of diseased plants show a narrow zone of brown along the margins, bordered by a yellow band. Inner leaves usually bear broad yellow bands. Suckers grow slowly and seldom form fruit. Old stems may break over, a black-rotted area having formed at the base upon which the yellow fruiting bodies of the fungus appear.
- Thielaviopsis paradoxa** (De Sey.) v. Hoeh. See *Saccharum*.
- Tylenchus musicola** Cobb. Plants of *Musa* of all ages are attacked in the British West Indies, the outer leaves dying and the fruit, if developed, drying up. The roots die, becoming black in color. The outer part of the bulb from which the roots develop likewise becomes black and rotted.
- Tylenchus similis** Cobb. This nematode attacks the roots of *Musa* spp. and *Saccharum officinarum* in Jamaica, Hawaii, and Fiji. The roots die back, diseased tissue becoming cinnabar-red and finally dark-purple.
- Uromyces musae** P. Henn. Powdery brown to black elongate rust sori on under leaf surfaces of *Musa* sp. in the Congo.
- MUSCARI.** GRAPE HYACINTH. Hardy spring flowering bulbs.
- Accidium muscari** Link. Forms yellow rust pustules on leaves of *M. comosum* and *M. racemosum* in England, Italy, France, Switzerland, Dalmatia, and Hungary. Said to be distinct from *Uromyces scillarum*.
- Heterosporium ornithogali** Klotzsch. See *Ornithogalum*.
- Physoderma muscari** Poir. On leaves of *M. comosum* in France.
- Puccinia liliacearum** Duby. See *Ornithogalum*.
- Puccinia lojkaiana** Thuem. See *Ornithogalum*.
- Sclerotium** sp. A storage rot due to a *Sclerotium* (*Botrytis*) commonly infects imported Holland bulbs of *Muscari* spp. Infected bulbs are matted together by a web of white mycelium in which small brown to black sclerotia appear. Other related bulbous plants may also be attacked.
- Septoria muscari** P. Brun. Large irregular spots with brown margins on leaves of *M. comosum* in France. This species is probably identical with *S. scillae* West.
- Septoria muscari-neglecti** Bub. On leaves of *M. neglectum* in Yugoslavia.
- Septoria scillae** West. Causes a leaf spot of *Muscari* sp., *Scilla bifolia*, and *S. nutans* in Belgium, Italy, Portugal, Spain, and Germany.
- Urocystis colchici** (Schlecht.) Rab. See *Colchicum*.
- Uromyces ornithogali** Lévl. See *Gagea*.
- Uromyces scillarum** (Grev.) Wint. See *Hyacinthus*.
- Ustilago vaillantii** Tul. See *Hyacinthus*.
- MUTISIA.** Showy flowered shrubs.
- Phyllachora mutisiae** Speg. A doubtful species on *Mutisia* sp. in Argentina.
- Puccinia crassicutis** Syd. Yellow and black rust pustules on *M. clematis* in Bolivia.
- Puccinia mutisiae** Lagerh. Brown rust pustules on leaves and stems of *Mutisia* sp. in Ecuador
- Puccinia mutisiicola** Speg. Brown rust pustules on leaves of *M. subspinosa* in Argentina.
- Septoria jaffueli** Speg. On leaves of *M. latifolia* in Chile.
- Septoria mutisiae** Speg. Ashen-colored leaf spots on *M. subspinosa* in Argentina.
- MYOPORUM.** Heathlike shrubs, grown for their small white or purplish flowers.
- Bagnisiella endopyria** Sacc. On leaves of *M. platycarpum* in Australia.
- Septoria myopori** Cke. and Mass. Circular whitish leaf spots with dull-brown margins on *M. insulare* in Australia.
- MYOSOTIS.** FORGET-ME-NOT. Low perennial or annual herbs.
- Accidium kabatianum** Bub. Rust on leaves of *M. sparsiflora* and *M. stricta* in central Europe.
- Cylindrosporium myosotidis** Sacc. and var. **boraginis** Sacc. Leaf spot on *M. palustris*, *M. scorpioides*, *Borago officinalis* and *Symphytum* sp. in Poland, France, and Spain.
- Entyloma canescens** Schroet. Smut sori in gray circular leaf spots on *M. caespitosa*, *M. hispida*, *M. intermedia*, and *M. sylvatica* in France, Denmark, Italy, and Germany.
- Peronospora myosotidis** De By. Downy mildew on leaves of *M. arenaria*, *M. arvensis*, *M. hispida*, *M. intermedia*, *M. palustris*, *M. sylvatica*, *M. sparsiflora*, *M. stricta*, and *M. versicolor* in Europe.
- Puccinia myosotidis** Tranzsch. Black rust sori on leaves and stems of *M. sylvatica* in Turkestan.
- Puccinia novo-zelandica** Bub. Dark-brown rust pustules on leaves of *M. capitata* in New Zealand.
- Synchytrium globosum** Schroet. See *Potentilla*.
- MYRCIA.** Tropical shrubs.
- Catacauma myrciae** (Lévl.) Theiss. and Syd. Dull-black stromata on leaves of *Myrcia* sp. in Brazil.
- Puccinia sanguinolenta** P. Henn. Dark-brown rust pustules on leaves of *Myrcia* sp. in Brazil.
- Uredo myrciae** Mayor. Rust on leaves of *M. acuminata* in Colombia.
- MYRCIARIA.** Jaboticaba. Brazilian fruit trees and shrubs.
- Laestadia cambucae** Rangel. On leaves of *M. plicati-costata* in Brazil.
- Uredo rochiai** Putt. Rust on fruit and leaves of *M. jaboticaba* and *M. plicati-costata* in Brazil.
- MYRICA.** BAYBERRY. Ornamental woody plants with attractive fruits, some edible.
- Dothidella koordersii** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *M. germanica* in Java.
- Dothidina disciformis** (Wint.) Theiss. and Syd. Black stromata on leaves of *Myrica* sp. in the Union of South Africa.

**MYRICA**—Continued.

- Microcyclus koordersii** P. Henn. Black discoid stromata on lower leaf surfaces of *M. javanica* in Java.
- Peronospora rufibasis** B. and Br. Downy mildew on brown leaf spots on *M. gale* in Great Britain.
- Rhytisma nitidum** Lév. Black tar spots on *M. aethiopica* in Abyssinia.
- MYRICARIA.** FALSE TAMARIX. Shrubs allied to Tamarix.
- Marsonia myricariae** Rostr. On leaves of *M. germanica* in Norway.
- Phyllosticta germanicae** Speg. On leaves of *M. germanica* in Italy.
- Puccinia involvens** (Voss.) Syd. Yellow-brown to black elongate rust pustules on *M. germanica* in France, Belgium, Switzerland, and Austria.
- Pyrenopeziza tamaricis** (Roum.) Sacc. On branches of *M. germanica*, *Tamarix africana*, and *T. narbonensis* in Argentina, Great Britain, France, and Germany.
- MYRISTICA.** NUTMEG. Tropical trees with aromatic fruit.
- Corticium salmonicolor** B. and Br. See Citrus.
- Eutype erumpens** Mass. See Theobroma.
- Fomes lamaoensis** Murr. See Hevea.
- Marasmius** spp. See Theobroma.
- Perisporium myristicae** P. Henn. Black superficial globular perithecia on leaves of *Myristica* sp. in Java.
- Phomopsis** sp. Causes a light-brown rot of fruit of *M. fragrans*, preventing seed germination, in Grenada.
- Phytophthora** sp. An undetermined species of this genus has been reported from Java, attacking the leaves and twigs of *Myristica* (nutmeg).
- MYRRHIS.** MYRRH. Perennial herbs.
- Puccinia chaerophylli** Purt. See Chaerophyllum.
- MYRSINE.** Tropical shrubs and trees with coriaceous leaves and white or yellow flowers.
- Chaetothrium punctiforme** Rick. On leaves of *Myrsine* sp. in Brazil.
- Corynelia fruticola** (Pat.) v. Hoehn. Black carbonous superficial fruiting bodies, more or less completely enveloping fruit of *M. africana* and *Rapanea melanophleas* in China, India, and the Union of South Africa.
- Napicladium piriforme** Speg. Circular brown leaf spots on *M. floribunda* in Argentina.
- Phyllachora myrsinicola** Doidge. Black stromata on leaves of *M. (Rapanea) melanophleas* in the Union of South Africa.
- Phyllachora pittieri** Theiss. and Syd. Tar spot on leaves of *Myrsine* sp. and *Xylosma salzmanni* in Costa Rica.
- Phyllachora sinik-lagaraik** Speg. Circular shiny black stromata on leaves of *M. floribunda* in Brazil.
- Phyllohendersonia nitida** (E. and E.) Tass. On leaves of *Myrsine* sp. in Hawaii.
- Scolecodothis circularis** (Bres.) Theiss. and Syd. Shiny black stromata on brown leaf spots on *Myrsine* sp. in Brazil.
- Uromyces myrsines** Diet. See Ardisia.
- MYRTUS.** MYRTLE. Shrubs grown for their aromatic qualities and attractive foliage, flowers and fruit.
- Ascochyta apiospora** Cke. and Mass. See Backhousia.
- Ascochyta myrticola** Maire. and Sacc. Subcircular reddish leaf spots with dark-purple margins on *M. communis* in Italy and Corsica.
- Cercospora myrti** Erikss. Subcircular purplish leaf spots on *M. communis* and *M. italica* in Scotland, Sweden, Denmark, Spain, and Austria.
- Cercospora saccardiana** Scalia. Indefinite dark-purple leaf spots on *M. communis* in Sicily.
- Depazea myrticola** Rabenh. On leaves of *M. communis* in Italy.
- Macrophoma nuptialis** Bub. Large yellow-brown leaf spots on *M. communis* in Yugoslavia.
- Mycosphaerella myrticola** Speg. On leaves of *M. chequen* in Chile.
- Napicladium fumago** Speg. See Eugenia.
- Phyllosticta costesi** Speg. On leaves of *M. chequen* in Chile.
- Phyllosticta nuptialis** Thuem. Subcircular dull-yellow to white leaf spots with violet margins on *M. communis* and *M. italica* in Albania, Dalmatia, Yugoslavia, Madeira, and Portugal.
- Rehmiodothis myrticola** (Rehm.) Theiss. and Syd. Shiny black stromata on leaves of *Myrtus* sp. in Ecuador.
- NAGEIA.** See Podocarpus.
- NANDINA.** Small shrubs with red or white berries.
- Phyllosticta nandinae** Tassi. Irregular dull-yellow leaf spots on *N. domestica* in Italy.
- NAPOLEONA.** Shrubs of tropical Africa.
- Phyllosticta napoleonae** Thuem. Irregular whitish leaf spots on *N. imperialis* in Portugal.
- NARCISSUS.** Daffodil. Hardy spring flowering bulbous plants.
- Bacillus croci** Miuza. See Crocus.
- Botrytis parasitica** Cav. The black sclerotia of this fungus are common on imported *Narcissus* bulbs, and it has doubtless been introduced into all parts of the United States. See under Tulipa.
- Cercospora narcissi** Boud. Causes a spotting on both surfaces of the leaves of *N. poeticus* in France.
- Coleosporium narcissi** Grove. Brown rust pustules on both surfaces of leaves of *N. poeticus* in England.
- Fusarium bulbigenus** Cke. and Mass. Reported by Masee as a very serious disease of *Narcissus* in Great Britain. The disease has also been reported from Holland. It is characterized by small yellowish to brown leaf spots which finally become covered with pale salmon-colored spore masses. The bulbs also ultimately become infected. Work by recent investigators, however, has quite clearly shown that *Fusarium bulbigenum* is a secondary organism following nematode attack and not the cause of a separate disease.
- Phyllosticta narcissi** Aderh. Large brown spots without definite margins on leaves of *N. poeticus* in Germany.
- Phyllosticta oudemansii** Sacc. and Syd. On leaves of *Narcissus* (cultivated varieties) in Holland.
- Puccinia schroeteri** Pass. Large oblong spots are produced on both surfaces of the leaves, with dull-violet margins. Brown to black rust pustules appear on these spots. The disease occurs in Great Britain, Italy, Austria, Belgium, and Germany on *N. pseudonarcissus* and *N. radiiflorus*.
- Ramularia narcissi** Chittenden. This disease is characterized by yellow stripes down the foliage and a dwarfing of the entire plant, involving also the flowerstalks. Yellow-brown spots up to 1 centimeter in length also occur on leaves, which later become covered with a fine white powder. The host is *N. poeticus* in variety in England. The fungus may be synonymous with the following species as well as with *Cercospora narcissi* Boud.
- Ramularia vallambrosae** Br. and Cav. A brown spotting of leaves and flower stalks in Italy of *N. biflorus*, *N. odoratus*, *N. poeticus*, and *N. pseudonarcissus*.
- Septoria narcissi** Pass. Attacks tips of living leaves of *Narcissus* (cultivated varieties) in Italy, Belgium, and Holland.

**NARCISSUS**—Continued.

- Stagonospora narcissi** Höll. Produces large irregular brown spots on living leaves of *N. poeticus* in Hungary.
- Tylenchus dipsaci** Kuehn. (*T. devastatrix* Kuehn.) The eelworm or nematode disease of bulbs (*Narcissus* and *Hyacinthus*) is beyond any doubt the most serious bulb disease known at present. On growing plants the disease is characterized by twisted leaves, with swollen yellow spots and stripes. Affected leaves as a rule remain stunted and may even die. The minute worms causing the disease can be found in these diseased areas. In the bulbs the presence of the nematode is noted by dark rings. The disease occurs in Holland, Great Britain, and France. It has been introduced into the United States with Dutch bulbs and it is known from a few localities. It has proved a serious pest of *Allium* spp., *Avena sativa*, *Fragaria* sp., *Linum usitatissimum*, *Medicago sativa*, *Pisum sativum*, *Secale cereale*, *Solanum tuberosum*, *Trifolium pratense*, and other economic plants in Europe and Japan.
- Urocystis colchici** (Schlecht.) Rab. See Colchicum.
- NARTHECIUM.** BOG ASPHODEL. Perennial rhizomatous herbs.
- Entyoma ossifragi** Rostr. Smut sori on gray, angular leaf spots on *N. ossifragum* in Denmark.
- Heterosporium ossifragi** Rostr. Produces spots on leaves of *N. ossifragum* in Denmark.
- NASTURTIUM.** WATER-CRESS. Semiaquatic herbs. Now referred to *Radicula*. (q. v.)
- Aecidium nasturtii** Hazsl. Rust on leaves of *Radicula (Nasturtium)* sp. in Hungary.
- Peronospora nasturtii-montani** Gäum. Downy mildew on leaves of *N. montanum* in Japan.
- Phyllosticta anceps** Sacc. Small subcircular ochraceous-yellow leaf spots on *N. amphibium* and *N. anceps* in England and France.
- Physoderma magnusiana** Krieg. Small oblong spots on leaf blades and petioles of *N. palustre* in Germany.
- Septoria uliginosa** Speg. Definite subcircular whitish leaf spots on *N. bonaërense* in Argentina.
- NELUMBO** (*Nelumbium*). LOTUS. Water lilies.
- Cercosporina nelumbi** Hori. On leaves of *N. nucifera (N. speciosum)* in Japan and China.
- NEMESIA.** Flower-garden annuals.
- Albugo evansii** Syd. White rust on leaves of *Nemesia* sp. in the Union of South Africa.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- NEPENTHES.** PITCHER PLANT. Aquatic pitcher plants.
- Phyllachora nepenthidis** Rac. Circular to elliptical shining black stromata on leaves of *N. melamphora* in Java.
- Phyllosticta nepenthacearum** F. Tassi. Subcircular dull-brown leaf spots on *N. sanguinea* in Italy.
- NEPETA.** CATNIP. GROUND IVY. Hardy perennial herbs.
- Erysiphe taurica** Lév. See Althaea.
- Septoria bornmuelleri** Syd. Irregular dirty yellow-brown leaf spots, often destroying entire leaf, on *N. teucrifolia* in Persia.
- Septoria catariae** Bub. Yellow leaf spots, often with greenish-gray surrounding zones, on *N. beltrani* and *N. cataria* in Spain and Hungary.
- NEPHELIUM.** LYCHEE. LITCHI. Tropical fruit trees.
- Exoascus** sp. Leaf curl of *N. litchi (Litchi chinensis)* in India.
- Phyllosticta nephelii** Delaer. See Durio.
- NEPHROLEPIS.** SWORD FERN.
- Entyoma nephrolepidis** Rac. Smut sori in fronds of *N. biserrata (N. acuta)* in Java.
- Milesina columbensis** (Diet.) Arth. Rust on fronds of *N. pendula* and *N. rivularis* in Colombia and Porto Rico.
- NERIUM.** OLEANDER. Glauous, evergreen-leaved shrubs.
- Ascochyta oleandri** Sacc. and Speg. White leaf spots with brown margins on *N. oleander* in Denmark, France, Italy, and Portugal.
- Cercospora neriella** Sacc. On leaves of *N. oleander* in Italy. Reported from Louisiana.
- Phyllosticta glaucispora** Delacr. White leaf spots with dull-brown margins on *N. oleander* in Spain.
- Phyllosticta neriicola** Brun. Pale-ochraceous leaf spots on *N. oleander* in France.
- Phyllosticta neri-oleandri** Stem. Leaf spot on *N. oleander* in the Caucasus.
- Septoria neriicola** Pass. On leaves of *N. oleander* in Italy.
- Septoria oleandrina** Sacc. On *N. oleander* in Bermuda, France, and Italy.
- NICOTIANA.** TOBACCO. Erect, rank-growing herbs with narcotic or poisonous properties.
- Ascochyta nicotianae** Pass. Irregular dull-brown leaf spots on *N. affinis* and *N. tabacum* in Italy, Sweden, Russia, and possibly some of the West Indies.
- Bacillus caulivorus** Prill. and Delacr. See Solanum.
- Bacillus maculicola** Delacr. Bacterial leaf spot of *N. tabacum* in France.
- Bacterium pseudozoogloeae** Honing. Said to cause black rust disease of leaves of *N. tabacum* in the Dutch East Indies.
- Cercospora raciborskii** Sacc. and Syd. Circular brown to whitish zoned leaf spots on *N. tabacum* in Java, India, Ceylon, Fiji, Australia, and Uganda.
- Colletotrichum nicotianae** Averna. Irregular confluent dark-brown anthracnose spots on stems and branches of *N. tabacum* in Brazil.
- “**Corcova.**” This disease of tobacco (*N. tabacum*), due to an unknown cause, is reported from Argentina under the native name. Numerous short (1 to 3 millimeters) dark lines, resembling insect burrows, appear on the leaves, uniting to form much longer ones. The leaf tissue between dies and the leaves wilt. Similar lesions from 2 to 6 centimeters long occur on the stems. Unaffected portions of leaves continue to expand, causing swelling and distortion of the leaf blades. The pith of the stem darkens and dries up.
- Cytospora nicotianae** Averna. Irregular yellow areas on stems of *N. tabacum* in Brazil.
- Diplodia cacaicola** P. Henn. See Theobroma.
- Macrophoma tabaci** Averna. Irregular gray spots on the stems of *N. tabacum* in Brazil.
- Oidium tabaci** Thuem. A powdery mildew forming circular white powdery spots on leaves of *N. tabacum* in the Union of South Africa, Italy, and Portugal.
- Peronospora hyoseyami** De By. (*Peronospora nicotianae* Speg.) A downy mildew attacking cotyledons, leaves, and roots of *N. glauca*, *N. tabacum*, and *Hyoseyamus niger* in Europe, Australia, Mexico, and in limited areas in the United States. Causes the serious blue mold disease in Australia.
- Phoma solanicola** Prill. and Delacr. Irregular yellow, then brown, areas on stems and branches of *N. tabacum* and *Solanum tuberosum* in Brazil, Spain, Italy, and France. The leaves wither when the areas are extensive. Averna-Sacca reports *Plowrightia solanicola* Averna as a perfect stage.
- Phyllosticta capsulicola** Sacc. and Speg. On capsules of *N. dilatata* in Italy.
- Phyllosticta nicotianicola** Speg. Ashen leaf spots on *N. acutiflora* in Argentina.
- Phyllosticta tabaci** Pass. Irregular, often confluent, ashen-colored leaf spots on *N. tabacum* in Italy and Yugoslavia.
- Phytophthora melongenae** K. Saw. See Solanum.

**NICOTIANA**—Continued.

- Phytophthora nicotianae** v. B. de Haan. (*Peronospora nicotianae* Speg.) This downy mildew causes a damping-off of tobacco seedlings. In addition it attacks leaves, roots, and stems of older plants and is destructive in Ceylon, Sumatra, Borneo, Java, Brazil, and Argentina.
- Placosphaeria nicotianae** Averna. Irregular dark-brown or black patches at bases of stems of *N. tabacum* in Brazil.
- Rhizoctonia** sp. See Vigna.
- Sclerotinia nicotianae** Oud. and Konig. Rotten areas, covered with gray mold on leaves and stems, in which black sclerotia form on *N. tabacum* in Holland.
- Septoria diversa** Sacc. and Syd. Rufous leaf spots with ashen-colored centers on *N. longiflora* in Argentina.
- Septoria nicotianae** Pat. Brown to white zoned leaf spots on *N. affinis*, *N. rustica*, and *N. tabacum* in Russia, Ecuador, and Jamaica. The frog-eye disease.
- Septoria tabacina** McAlp. Leaf spot on *N. tabacum* in Australia.
- Uredo nicotianae** Anast. and Splend. Powdery brown rust sori on leaves of *N. quadrivalvis*, *N. sylvestris*, and *N. tabacum* in Brazil and Italy.
- NIDULARIUM.** Epiphytic bromeliads related to the pineapple.
- Uredo nidularii** P. Henn. Linear brown rust pustules on leaves of *N. longiflorum* in Brazil.
- NIEREMBERGIA.** CUPFLOWER. Tender perennial herbs.
- Entyoma nierembergiae** Lagh. Smut sori on pale circular leaf spots on *N. spathulata* in Ecuador.
- Puccinia nierembergiae** Lév. Powdery black rust pustules on leaves of *Nierembergia* sp. in Peru.
- NIGELLA.** LOVE-IN-A-MIST. Hardy annuals grown for their interesting flowers and seed pods.
- Cercospora nigellae** Höll. On leaves of *N. arvensis* in Hungary.
- NIOBE.** See Hosta.
- NIPA.** Oriental thatch palm with edible fruit. See Palmae.
- NITRARIA.** NITER BUSH. Low, rigid shrubs.
- Accidium nitrariae** Pat. Rust on leaves of *N. tridentata* in Algeria.
- NOTHOFAGUS.** Ornamental South American trees.
- Cytaria darwini** Berk. Causes a rot of the wood of branches of *Nothofagus* sp. in Chile.
- Exoascus entomosporus** (Thaxt.) Sacc. and Trot. Fruiting layer on lower leaf surfaces of *N. antarctica* var. *bicrenata* and var. *uliginosa* in Patagonia. The leaves are deformed.
- Uncinula magellanica** Thaxt. Powdery mildew on leaves of *N. antarctica bicrenata* in Patagonia.
- Uncinula nothofagi** Thaxt. Similar to above species.
- NYCTANTHES.** NIGHT JASMINE. Trees or shrubs with numerous white fragrant flowers.
- Cercospora puttemansii** P. Henn. Dull-brown leaf spots on *N. arbor-tristis* in Brazil.
- Ramularia** sp. Areolate mildew on leaves of *N. arbor-tristis* in India.
- NYMPHAEA.** WATER LILY (including *Nuphar*). SPATTER-DOCK. Herbaceous aquatics.
- Accidium nymphoides** DC. Rust on leaves of *N. alba*, *N. luteum*, and *Limnanthemum nymphoides* in France, Italy, Switzerland, Belgium, and Germany.
- Ovulariella nymphaearum** (Allesch.) Kab. and Bub. (*Gloeosporium nymphaearum* Allesch.) Sub-circular to irregular, often confluent, reddish to gray-brown leaf spots, pale at the centers, on *N. alba*, *N. bouchiana*, *N. capensis*, *N. flava*, *N. luteum*, and *N. ortgiensiana* in Europe.
- Phyllosticta hydrophila** Speg. Subcircular sunken dull-brown leaf spots on *N. alba* in Italy.
- Phyllosticta nymphaeae** Pass. On leaves of *N. alba* in Italy.
- Physoderma tenue** (Nowak.) Karst. On petioles of *Nymphaea* sp., *Acorus calamus*, and *Iris pseudacorus* in Denmark, France, Silesia, Finland, and Germany.
- Pythium undulatum** H. E. Peters. Downy mildew on leaf blades and petioles of *N. luteum* and *N. alba* in Denmark.
- Ramularia nymphaeae** Bres. Black circular leaf spots on *N. alba* in England and Germany.
- Rhamphospora nymphaeae** Cunn. Smut on leaves of *N. lotus*, *N. rubra*, and *N. stellata* in India.
- Septoria nupharis** Ranoj. On leaves of *N. luteum* in Yugoslavia.
- OCHLANDRA.** See Bambuseae.
- OCHNA.** Woody plants.
- Cocconia capensis** Doidge. On *O. arborea* in the Union of South Africa.
- Phyllachora ochnae** Pat. and Har. Tar spot on leaves of *Ochna* sp. in Madagascar and the Philippines.
- OCIMUM.** BASIL. Annual and perennial herbs.
- Accidium leiocarpum** Syd. Leaf rust on *O. canum* in India.
- Accidium ocimi** P. Henn. Golden-yellow rust sori on circular brown leaf spots on *O. canum*, *O. graveolens*, and *O. suave* in India and Abyssinia.
- Coleosporium plectranthi** Barcl. See Plectranthus.
- Phyllosticta basilici** Brun. Gray-brown leaf spots on *O. basilicum* in France.
- OCOTEA.** Tropical trees.
- Cephalosporium tumefaciens** Wint. On *O. tristis* in Brazil.
- Oligostroma archavaletae** (Speg.) Theiss. and Syd. Black tar spot on leaves of *O. acutifolia* in Uruguay.
- Phyllachora ocoteae** P. Henn. Irregular black stromata on leaves of *Ocotea* sp. in Brazil.
- Phyllachora ocoteicola** Stev. and Dalby. Angular dull-black stromata on leaves of *O. leucoxydon* in Porto Rico.
- ODONTOGLOSSUM.** Epiphytic orchids. See Orchidaceae.
- OENANTHE.** Aquatic perennial herbs.
- Entyoma oenanthum** Sacc. Smut sori on stems of *O. globulosa* in Malta.
- Entyoma oenanthes** R. Maire. White smut pustules on leaves of *O. apiifolia* in Corsica.
- Physoderma vagans** Schroet. See Ranunculus.
- Puccinia oenanthes** (Diet.) T. Miy. Dark-brown rust pustules on leaves of *O. stolonifera* in Japan.
- OENOTHERA.** EVENING PRIMROSE. Flower-garden plants with bright-colored flowers.
- Cercospora oenotherae** Speg. Circular dull-brown leaf spots on *O. mollissima* in Argentina.
- Puccinia luxurians** Diet. and Neg. Brown rust pustules on leaves of *O. mutica* in Chile.
- Uredo oenothericola** Speg. Powdery, dark ferruginous rust pustules on leaves of *O. mollissima* in Argentina.
- OLEA.** OLIVE. Fruit and ornamental trees.
- Ascochyta oleae** Scal. On leaves of *O. europaea* in Sicily.
- Bacterium olivae** Cif. Found in connection with dieback of twigs of *O. europaea* in Italy.
- Cercospora cladosporioides** Sacc. On leaves of *O. europaea* in Italy and Algeria.
- Ceuthospora oleae** Kalchbr. and Cke. Circular dull-brown leaf spots on *O. capensis* in the Union of South Africa.
- Ceuthospora phacioides** Grev. and var. *oleae* Scalia. On leaves of *O. europaea* in Sicily and Austria.
- Coniothyrium oleae** Pol. On leaves of *O. europaea* in Italy.

**OLEA**—Continued.

- Cylindrosporium olivae** Petri. Large sunken yellowish-purple spots with dark purple margins on fruit of *O. europaea* in Italy.  
**Cystospora oleae** Butl. Rust on pale-brown leaf spots on *O. dioica* in India.  
**Gloeosporium olivarum** d'Alm. Anthracnose on fruit of *O. europaea* in France, Italy, Spain, Portugal, and Algeria.  
**Macrophoma dalmatica** (Thuem) Berl. and Vogl. Subcircular dull-brown spots on fruit of *O. europaea* in Spain, Italy, and Austria.  
**Perischizon oleifolium** (Kalch. and Cke.) Syd. On leaves of *O. capensis* in the Union of South Africa.  
**Phyllosticta insulana** Mont. Small pale leaf spots with narrow brown margins on *O. europaea* in Spain, Italy, and France.  
**Phyllosticta oleae** Petri. Leaf spots on *O. europaea* in Italy.  
**Septoria oleae-chrysophyllae** Pass. On leaves of *O. chrysophylla* in Abyssinia.  
**Septoria oleaginea** Thuem. Irregular pale-ochraceous leaf spots on *O. europaea* in Austria.  
**Septoria olivae** Pass. and Thuem. On leaves of *O. europaea* in Portugal.  
**Stictis panizzae** De N. Causes the leaf disease of *O. europaea* known as Brusca in Spain and Italy.

**OLEARIA. SHAWIA** Ag. DAISY TREE. Australasian trees and shrubs.

- Accidium oleariae** McAlp. Rust on leaves and stems of *O. axillaris* in Australia.  
**Puccinia oleariae** McAlp. Bright orange then brown rust pustules on leaves of *O. argophylla* in Tasmania.  
**Uredo brownii** Syd. Rust on leaves of *O. angustifolia* in New Zealand.

**ONCIDIUM.** Epiphytic orchids. See Orchidaceae.**ONCOSPHERMA.** See Palmae.**ONOBRYCHIS.** HOLY CLOVER. Perennial forage plants.

- Anthostomella** sp. Black leaf spots, often confluent, with yellow surrounding zones, on *O. sativa* in Italy.  
**Ascochyta orobi** Sacc. var. **onobrychidis** Prill. and Délacr. On leaves of *O. sativa* in France and Yugoslavia.  
**Cylindrosporium onobrychidis** (Syd.) Died. On *O. sativa* in Europe.  
**Ovularia bornmülleriana** P. Magn. Yellow leaf spots on *O. tournefortii* in Asia Minor.  
**Peronospora ruegeriae** Gäum. Downy mildew on leaves of *O. sativa* in Switzerland.  
**Phyllachora lathyri** (Lév.) Theiss. and Syd. See Lathyrus.  
**Physalosporina ducellieri** Maire. On leaves of *O. caputgallis* in French North Africa.  
**Ramularia onobrychidis** Allesch. Clear light-brown leaf spots with darker margins on *O. sativa* and *O. viciaefolia* in Russia, Germany, Sweden, Yugoslavia, France, and Denmark.  
**Rhytisma onobrychidis** DC. Irregular warty dull-black patches, on leaves of *O. sativa* and *Vicia cracca* in Siberia and France.  
**Uromyces onobrychidis** (Desm.) Lév. Powdery brown rust pustules on *O. arenaria*, *O. gracilis*, *O. montana*, *O. sativa*, and *O. viciaefolia* in Europe.

**ONOCLEA.** Sensitive fern.

- Taphrina hiratsuka** Nish. Fruiting areas on leaves of *O. sensibilis* in Japan.

**ONONIS.** REST-HARROW. Annual, biennial, and perennial border and rock-garden plants.

- Napicladium ononidis** (Auersw.) Sacc. On leaves of *O. repens* in France.  
**Peronospora ononidis** Wils. Downy mildew on leaves of *O. hircina*, *O. repens*, and *O. spinosa* in Russia, Denmark, Switzerland, Austria, and Germany.  
**Ramularia winteri** Thuem. On leaves of *O. arvensis*, *O. hircina*, *O. repens*, and *O. spinosa* in Switzerland, Denmark, and Germany.  
**Uromyces ononidis** Pass. Cinnamon to dark brown rust pustules on leaves of *O. alopecuroides*, *O. antiquorum*, *O. campestris*, *O. columna*, *O. hircina*, *O. procurrens*, *O. repens*, *O. rotundifolia*, and *O. spinosa* in Europe and Asia Minor.  
**Uromyces sphaeropleus** Cke. Dark-brown rust pustules on *Ononis* sp. in India.

**ONOPORDON.** COTTON THISTLE. Coarse, woolly, old-world herbs.

- Puccinia acanthii** Syd. Powdery brown rust pustules on leaves of *O. acanthium* in Spain and Germany.  
**Puccinia onopordi** Syd. Brown rust pustules on leaves of *O. cardunculum* in Syria.  
**Ramularia onopordi** Massal. Subcircular yellow leaf spots on *O. acanthium* and *O. ceratonicum* in Spain, Italy, Yugoslavia, and Germany.  
**Septoria onopordonis** P. Nag. Leaf spot on *O. acanthium* in Russia.

**ONOSMA.** Herbs grown for their golden-colored flowers.

- Accidium onosmatis** Thuem. Rust on leaves of *O. gmelini* and *O. simplicissima* in Siberia.

**OPHIOGLOSSUM.** ADDER'S-TONGUE.

- Helminthosporium crepini** West. On leaves of *O. vulgatum* in France, Belgium, and Denmark.  
**Helminthosporium diedickei** P. Magn. On *O. vulgatum* in Germany.

**OPHIOPOGON.** SNAKE'S-BEARD. Herbs grown for their small bluish or white flowers.

- Eusciadium transversum** Sacc. Greenish to black areas on leaves of *O. japonicus* in Italy.  
**Septoria aemula** Tassi. Brown leaf spots on *O. spicatus* in Italy.  
**Septoria ophiopogonis** Pass. Brown areas at the tips of leaves of *O. japonicus* in Italy.  
**Uredo ophiopogonis** Syd. Cinnamon-brown rust sori on leaves of *O. japonicus* in Italy.

**OPHRYS.** Terrestrial orchids. See Orchidaceae.**OPULASTER.** See Physocarpus.**OPUNTIA.** PRICKLY PEAR. Sometimes cultivated for ornament or forage.

- Accidium opuntiae** P. Magn. On *Opuntia* sp. in Bolivia.  
**Ascochyta opuntiae** Scalia. On *O. ficus-indica* in Italy.  
**Didymosphaeria opulenta** Sacc. Causes yellow-brown blotches on joints of *O. ficus-indica* in Italy and Malta.  
**Phyllosticta opuntiae** Sacc. and Speg. and var. **microscopicae** Cifer. Whitish spots on *O. ficus-indica* and other species in Italy and Bermuda.  
**Sclerotium opuntiarum** Speg. On *Opuntia* spp. in Argentina.  
**Septoria ficus-indicae** Vogl. On *O. ficus-indica* in Italy.

**ORCHIDACEAE.**

The diseases attacking the various genera of the *Orchidaceae* are considered under the latter head for reasons similar to those given under *Bambuseae*.

- Accidium alaskanum** Trel. This rust forms small, pale to light brown spots on both surfaces of the leaves of *Habenaria* sp. and other orchids in Alaska.  
**Amerosporium rhodospermum** McAlp. On leaves of *Diuris pedunculata* in Australia.  
**Amerosporium vanillae** P. Henn. On leaves of *Vanilla aromatica* in Java.  
**Ascochyta orchidis** Rabenh. On leaves of *Orchis* sp. in Germany.  
**Ascospora** sp. On living leaves of *Evelyna discolor* in Peru.  
**Asterina liparidis** Rac. On leaves of *Liparis* sp. in Java.

## ORCHIDACEAE—Continued.

- Asterinella epidendri** (Rehm.) Theiss. The fungus produces small, black, shield-shaped fruiting bodies on the upper surfaces of leaves of *Epidendrum* sp. in Brazil.
- Atichia vanillae** (Pat.) V. Hoeh. On leaves of *Vanilla planifolia* in Tahiti.
- Bacillus cyripedii** S. Hori. This bacterium causes the brown spot disease of orchids which is described as follows: Dirty-cinnamon or light-umber colored spots appear on the leaf-blades. The spots enlarge with great rapidity, so that within a few days entire green leaves can become discolored. After some days the diseased parts become brownish and finally deep chestnut-brown; their surfaces become more or less wrinkled, with loss of luster; their margins diffused, faint in color, and depressed. The lower surfaces of the leaves beneath the spots assume more or less rapidly a pale color and only gradually develop the same color as the upper surfaces. When the spots develop on the lower portions of a leaf, the upper part soon becomes yellowish and dies off without being directly attacked by the bacteria. The rotting also spreads downward into the stem, and if the diseased leaf is not cut off in time, the entire stock will soon be destroyed.
- A similar orchid disease called the "brown spot" produces circular brown sunken spots on the upper leaf surfaces. As the spots enlarge the shape becomes somewhat irregular and the periphery more or less undulated, but it is well defined by the deeply sunken area from the surrounding healthy part. The color of the spots differs somewhat according to the different hosts; some spots assume a yellowish-brown, others a deep chestnut-brown color. Inoculation experiments demonstrated the two diseases to be identical, differences being due to varying texture of the leaves of the hosts. Brown rot occurred on succulent leaved species and brown spot on firm, tough-leaved species.
- Hori states that the disease is greatly feared by gardeners, since it occurs on the most valuable orchids and spreads very rapidly during the summer season. The host species are: *Cypripedium godeffroyae*, *C. raynaldianum*, *C. philippinense*, *Phalaenopsis amabilis*, *P. aphrodite*, and *P. schilleriana*. The disease has been found in Japan and Formosa on plants in hothouses as well as on plants collected in the wild.
- Bacillus farnetianus** Pavar. Pavarino has reported this disease as attacking the leaves and pseudobulbs of *Oncidium ornithorhynchum* and *Cattleya crispa* in the botanical garden, Pavia, Italy. The disease is characterized by small, indefinite areas showing as transparent spots which increase in size, becoming visible on both sides of the leaves as brown to black dry, depressed areas.
- Bacillus krameriani** Pavar. This disease attacks *Oncidium kramerianum* in Italy. At first the diseased areas are transparent, later becoming irregular and finally brown and dry. Leaves and pseudobulbs are attacked.
- Bacillus poliacii** Pavar. This disease causes irregular, black, depressed spots on leaves of *Odontoglossum citrosimum* in Italy.
- Bacterium briosianum** Pavar. The leaves of *Vanilla planifolia* are said to be seriously attacked by this bacterium in the botanical garden at Pavia, Italy. Small, irregular, pitch-black spots are first seen which soon become lighter in color. The disease progresses rapidly, drying out and killing the leaves and in time involving the entire plant. This disease and the three preceding ones have all been reproduced by artificial inoculations, according to Pavarino.
- Bacterium cattleyae** Pavar. This disease is said to produce brown spots on the leaves and pseudobulbs of *Cattleya warneri* and *C. loddigesii* in the botanical garden of Pavia, Italy.
- Bacterium dendrobii** Pavar. Said to cause a withering of leaves and shoots of *Dendrobium nobile* in Italy.
- Bacterium oncidii** Peglion. This disease produces violet-brown spots on the leaves of *Oncidium* spp. in Italy. The entire leaf area is soon involved, the leaf losing its turgor and falling. Hori states that the causative organism should be known as *Bacillus oncidii* (Peg.) Hori. and that it is closely related to his *B. cyripedii*.
- Black rot.** A black rot of *Cattleya* leaves and bulbs has been reported from England. It is stated to be "A dreadful disease which carries off plants in a short time." Cause unknown.
- Botryodiplodia digitata** Maubl. On pseudobulbs of *Cattleya mossiae* in hothouses in France.
- Botryosphaeria egenula** Syd. and Butl. On leaves of *Cymbidium* sp. in Assam.
- Calonectria copelandii** P. Henn. On leaves of *Orchidaceae* in the Philippines.
- Calospora vanillae** Mass. Anthracnose on leaves of *Dendrobium* spp., *Oncidium* spp., and *Vanilla planifolia* in Madagascar, Tahiti, Colombia, West Indies, Seychelles, Reunion, and Mauritius. Said to be the same as *Gnomoniopsis vanillae* Stonem. on *Vanilla* in greenhouses in the United States.
- Cercospora angreci** Gaill. and Feuill. On leaves of *Angraecum fragrans*, *Cattleya* sp., and *Odontoglossum crispum* in Mauritius.
- Cercospora epipactidis** C. Mass. On leaves of *Epipactis palustris* in Italy.
- Cercospora odontoglossi** Prill. and Delacr. On the under sides of leaves of *Odontoglossum* spp., including *Odontoglossum crispum*, in greenhouses in France. This species is said to cause much damage.
- Ceuthospora cattleyae** Sacc. and Syd. (*C. minima* Delacr.) This fungus produces dark-brown spots on the leaves of *Cattleya intermedia* in Holland. The leaves ultimately die as a result of the fungus attack.
- Chaetodiplodia vanillae** A. Zimm. On rotting leaves of *Vanilla* sp. in Java. Probably not distinct from *Diplodia cacaoicola* P. Henn. See Theobroma.
- Chytridiaceae** (?). A leaf-spot disease of *Cattleya*, *Dendrobium*, *Odontoglossum*, and other cultivated orchids has been briefly described by Brierley. It is characterized by a series of concentric zones of green to pale-straw color and brown through purple to black. These vary in diameter from  $\frac{1}{8}$  to  $1\frac{1}{2}$  inches, occurring either as complete circles or segments of circles. The fungus has not yet been identified, but inoculations with diseased material readily reproduce the disease. It is thought that the fungus concerned belongs in the *Chytridiaceae*.
- Ciliella epidendri** (Rehm) Sacc. and Syd. On leaves of *Epidendrum* sp. in Brazil.
- Cladochytrium olivieri** Har. Dark leaf spots on *Orchis incarnata* and *O. laxiflora* in France.
- Cladosporium** sp. (*C. orchidis* Cke. and Mass. [?]). Forms olive-green blotches, which turn to a brownish-purple on living leaves of *Dendrobium* spp., *Coelogyne* spp., *Oncidium* spp., *Phalaenopsis* sp., and other cultivated orchids in England.
- Coleosporium arundinae** Syd. This rust produces orange-yellow pustules on the under sides of leaves of *Arundina chinensis* and *A. speciosa* in Java and Formosa.
- Coleosporium bletiae** Diet. Characterized by brown to golden-yellow rust pustules which fade out to white on the under sides of leaves of *Bletia hyacinthina* in Japan. The fungus has also been reported from one locality in central California on *Phajus wallichii* (*P. grandifolius*), it doubtless having been introduced from Japan.
- Coleosporium merrillii** P. Henn. Bright-yellow rust pustules on the under sides of leaves of *Spathoglottis chrysantha* and other undetermined orchids in the Philippines.
- Colletotrichum dichaeae** P. Henn. On leaves of cultivated *Dichaea vaginata* in the botanical Garden at Berlin.

## ORCHIDACEAE—Continued.

- Colletotrichum effiguratum** Syd. Irregular, yellow-brown leaf spots with distinct purple-brown margins on *Paphiopedilum roezlii* in Germany.
- Colletotrichum macrosporum** Sacc. On *Orchidaceae* in Brazil.
- Colletotrichum orchidearum** Allesch. On leaves of *Bulbophyllum lobbii*, *B. longiflorum*, *Cattleya eldorado*, *Cymbidium* sp., *Coelogyne mayeriana*, *Epidendrum macrostachyum*, *Eria ornata*, *E. stellata*, *Eulophia saundersiana*, *Laelia crispa*, *Oncidium pulvinatum*, *Physosiphon loddigesii*, *Pholidota imbricata*, *Pleurothallis tribulooides*, *Spathoglottis plicata*, and *Sarcanthus pugioniformis* in the botanical garden at Munich, as well as in Colombia, Brazil, and the Philippines on native plants.
- Colletotrichum orthianum** A. Kostlan. On leaves and leaf sheaths of *Cattleya* sp., *Coelogyne cristata*, *Cypripedium insigne*, *Dendrobium* sp., *Dendrobium nobile*, and *Vanda coerulea* in the botanical garden at Berlin.
- Colletotrichum roseolum** P. Henn. On pseudobulbs of *Stanhopea oculata* in the botanical garden at Berlin.
- Colletotrichum vanillae** Scalia. Forms brown spots on leaves of *Vanilla odorata* and *V. planifolia* in Sicily and Ceylon.
- Colletotrichum vinosum** P. Henn. On pseudobulbs of *Stanhopea oculata* in the botanical garden at Berlin.
- Cylindrina delavarji** Pat. On leaves of *Liparis liliifolia* in China.
- Diplodina dendrobii** Cooke and Mass. On leaves of *Dendrobium speciosum* in Australia.
- Diplodia henriquesiana** Trav. and Spessa. On pseudobulbs of *Cattleya labiata* in Italy.
- Diplodia sobraliae** (P. Henn.) Taub. On leaves of *Sobralia sessilis* in the botanical garden at Berlin.
- Fusarium** sp. Attacks the bases of leaves of *Vanilla planifolia* in Brazil, extending into the stems and forming small, clear, irregular brown spots, which finally become confluent.
- Fusicladium vanillae** A. Zimm. Appears as a brown fungus layer growing over living leaves of *Vanilla* sp. in Java.
- Gloeosporium affine** Sacc. This fungus produces white spots on both surfaces of leaves and pseudobulbs of *Bulbophyllum lobbii*, *Cattleya mendelii*, *Coelogyne cristata*, *Masdevallia psittacina*, *Odontoglossum crispum*, *Pleurothallis lansbergii*, *Vanilla* sp., and other cultivated orchids in Italy and Germany.
- Gloeosporium beyrodtii** Klitz. Produces dark-brown blotches which enlarge rapidly, soon causing the death of leaves of *Vanda coerulea* in Germany.
- Gloeosporium bidgoodi** Cke. On *Odontoglossum* spp. and *Oncidium* sp. in hothouses in England.
- Gloeosporium bussei** P. Henn. On leaves of *Vanilla* sp. in the botanical garden at Berlin.
- Gloeosporium cattleyae** Sacc. On leaves of *Cattleya mossiae* in France.
- Gloeosporium coelogyne** Syd. Forms pale spots on leaves of *Coelogyne viscosa* in the botanical garden at Berlin.
- Gloeosporium dendrobii** Maubl. Characterized by large, pale, ocher-yellow spots on leaves of *Dendrobium farmerii* in France.
- Gloeosporium epidendri** P. Henn. Large black spots on flower stems of *Epidendrum* sp. in the botanical garden at Berlin and on *Epidendrum ciliaris* in Spain.
- Gloeosporium intermedium** Sacc. var *epidendri* Sacc. On leaves of *Orchidaceae* including *Epidendrum* spp., in France.
- Gloeosporium laeliae** P. Henn. Yellow-brown spots covering entire leaf surfaces of leaves of *Laelia* sp. in the botanical garden at Berlin.
- Gloeosporium maxillariae** Allesch. On leaves of *Maxillaria rufescens* in the botanical garden at Berlin.
- Gloeosporium noackianum** Allesch. On leaves of *Orchidaceae* in Brazil.
- Gloeosporium oncidii** Oud. Produces brown spots on the leaves and leaf-sheaths of *Maxillaria infestans*, *M. rufescens*, and *Oncidium ansiferum* in Holland and Germany, and on *Oncidium sphacelatum* in Mexico.
- Gloeosporium ornithidii** Allesch. Produces large, irregular spots which are ocher-colored and often confluent, on leaves of *Ornithidium densum* in greenhouses in Germany.
- Gloeosporium pallidum** (Karst.) Hariot. f. *physosiphonis-loddigesii* Allesch. On leaves of *Physosiphon loddigesii* and *Liparis longipes* in greenhouses in Germany.
- Gloeosporium phaji** Maubl. Produces round spots, which are white at first, becoming black and finally whitish at the centers, on living leaves of *Phajus* spp. and *Phajus wallichii* in greenhouses in France.
- Gloeosporium sobraliae** Maubl. Forms spots near the tips of the leaves, with a brown-black line of demarcation between healthy and diseased tissue, on *Sobralia* sp. in greenhouses in France.
- Gloeosporium stanhopeae** Allesch. On leaves of *Stanhopea* sp. in greenhouses in Germany.
- Gloeosporium stanhopeicolum** P. Henn. On leaves of *Stanhopea* sp. in the botanical garden at Berlin.
- Gloeosporium vandopsisidis** Keissler. On leaves of *Vandopsis* sp. in the Solomon Islands.
- Gloeosporium vanillae** Cooke and Mass. On leaves of *Vanilla planifolia* in Colombia, Mauritius, Ceylon, and other parts of the world and on *Vanda sanderiana* in the Philippines.
- Guignardia microsticta** Sacc. On *Cattleya* sp. in France.
- Guignardia traversii** (Car.) Lind. On leaves of *Vanilla planifolia* in Italy.
- Hemileia americana** Mass. This rust has been reported on the under sides of leaves of *Cattleya dowiana* and *Oncidium cavendishianum* in greenhouses in England, the plants having come originally from Central America. Powdery golden-yellow pustules are produced. The form on *O. cavendishianum* should be referred to *Hemileia oncidii* Griff. and Maubl.
- Hemileia oncidii** Griff. and Maubl. (*Uredo behnickiana* P. Henn.) This species produces bright-yellow pustules on the lower leaf surfaces of *Epidendrum vitellinum*, *Lycaste skinneri*, *Oncidium crispum*, *O. dasystalix*, *O. forbesii*, *O. marshallianum*, and *O. varicosum* in greenhouses in Brazil, France, Ireland, and Germany.
- Hemileia phaji** Syd. Characterized by yellow rust pustules on the lower surfaces of leaves of *Phajus* spp. including *P. blumei* and *P. wallichii* in Java. Grove states that these fungi (*Hemileia* spp.) might easily become dangerous parasites if allowed to spread.
- Hendersonia epidendri** Keissler. On leaves of *Epidendrum bifidum* in the island of St. Croix.
- Hypodermium orchidearum** Cke. and Mass. This fungus produces spots in groups which are often 1 to 2 inches long, causing the leaves of *Cymbidium eburneum* in Great Britain to turn yellow and die.
- Leptostroma orchidearum** Mntg. On living leaves of *Cypripedium macranthum* in Siberia.
- Leucothyridium (?) vanillae** Avena. Irregular black fungus crusts on leaves and stems of *Vanilla planifolia* in Brazil.
- Macrophoma cattleyicola** P. Henn. Characterized by numerous large, brown-black spots on pseudobulbs of *Cattleya labiata* and *C. percivaliana* in the botanical garden at Berlin.
- Macrophoma cattleyicola** P. Henn. var. *brassavolae* Keissler. Forms spots on living leaves of *Brassavola* sp. at the Kew gardens, England.

## ORCHIDACEAE—Continued.

- Macropoma miltoniae** DaCam. On leaves of *Miltonia candida* in the botanical garden, Coimbra, Portugal.
- Macropoma oncidii** P. Henn. Said to cause the death of leaves of *Oncidium pulvinatum* in the botanical garden at Berlin. Also reported on leaves of *O. sphacelatum* and *Cattleya* sp. from Mexico.
- Macrophoma vanillae** Averna. Large irregular, then confluent, ashen leaf spots with dark margins on *Vanilla planifolia* in Brazil.
- Melampsora repentis** Plowr. (*Caecoma orchidis* Wint.) This rust produces pale yellow spots 1 to 2 centimeters long and generally somewhat elongated, on the under sides of the leaves of *Gymnadenia conopsea*, *G. albida*, *Habenaria bifolia*, *H. chlorantha*, *H. conopsea*, *H. solstitialis*, *Listera ovata*, *Ophrys aranifera*, *O. muscifera*, *Orchis angustifolia*, *O. bifolia*, *O. incarnata*, *O. latifolia*, *O. maculata*, *O. mascula*, *O. militaris*, *O. palustris*, *O. sambucina*, *O. undulata*, and *Phalaenopsis esmeralda* in British Guiana, Spain, Austria, Denmark, Great Britain, Hungary, Russia, Sweden, Switzerland, and Germany. The ur- and telial stages occur on *Salix repens*, *S. aurita* and *S. triandra*.
- Me anospora coemansii** West. On leaves of *Oncidium* sp. in Belgium.
- Metola** sp. On living leaves of *Pleurothallis ruscifolia* in Porto Rico and *Laelia albida* in Mexico.
- Myiocopron (?) vanillae** Averna. Black fruiting bodies on extensive confluent ashen leaf spots on *Vanilla planifolia* in Brazil.
- Nectria bulbicola** P. Henn. On bulbs of *Cyrtopodium* sp., *Gomezia planifolia*, *Gongora galeata*, *Maxillaria rufescens*, *M. variabilis*, *Oncidium pulvinatum*, *Polystachya* sp., and *Sophonitis cernua* in the botanical garden at Berlin. The species has been referred to *N. ochroleuca* (Schw.) Bark., a common tropical species.
- Nectria bulbophylli** P. Henn. On *Bulbophyllum lobbii* in the botanical garden at Berlin.
- Nectria goroshankiniana** Wahrlich. On roots of *Vanda tricolor* in orchid houses in Russia.
- Nectria vandae** Wahrlich. On roots of *Vanda suavis* in orchid houses in Russia.
- Nectria vanillae** A. Zimm. Brown spots on stems and leaves of cultivated *Vanilla* in Java.
- Nectria vanillicola** P. Henn. On leaves of *Vanilla aromatica* in Java.
- Pestalozzia vanillae** Averna. Irregular, often marginal, brown leaf spots on *Vanilla planifolia* in Brazil.
- Phoma obtrudens** F. Tassi. On leaves of *Phalaenopsis longifolia* in Italy.
- Phoma oncidii** Speg. (*Phoma corrientina* Speg.) Forms indefinite white spots on leaves of *Oncidium* sp. in Argentina and *O. papilio* in hothouses in Denmark.
- Phoma oncidii-sphacelati** F. Tassi. On living leaves of *Oncidium sphacelatum* in Mexico.
- Phyllosticta bletiae** A. Zimm. On living leaves of cultivated *Bletia hyacinthina* in Austria.
- Phyllosticta decidua** Ferraris. Characterized by pale ochre-colored spots, circular in shape, with brown margins, on leaves of *Goodyera repens* in Italy.
- Phyllosticta donckelaarii** Westendorp. Forms large grayish spots with red-brown margins on the under sides of leaves of *Oncidium* sp. in hothouses in Belgium.
- Phyllosticta epipactidis** Died. Produces dark-brown spots, becoming lighter at the centers, 1 to 1½ centimeters in diameter, on leaves of *Epipactis violacea* in Germany.
- Phyllosticta laeliae** Keissler. The fungus produces light-colored spots on the leaves and leaf sheaths of *Laelia albida* and *L. furracea* in Mexico.
- Phyllosticta masdevalliae** P. Henn. On leaves of *Masdevallia chimaera* in the botanical garden at Berlin.
- Phyllosticta nigramaculans** Sacc. On leaves of *Orchidaceae* in Brazil.
- Phyllosticta pleurothallidis** Keissler. On living leaves of *Pleurothallis longissima* in Costa Rica.
- Phyllosticta renantherae** Keissler. Forms indistinct, light-colored, subcircular spots on living leaves of *Renanthera storiei* in the Philippine Islands.
- Phyllosticta stanhopeae** Allesch. The fungus does not form definite spots but attacks the whole leaf, which soon dies. The hosts are *Stanhopea* spp. in Germany.
- Phyllosticta valparadisica** Speg. Forms circular brown spots 1 to 3 millimeters in diameter, on both sides of leaves of undetermined orchids from Chile.
- Phyllosticta vanillae** P. Henn. Produces large pale-colored spots on leaves of *Vanilla aromatica* in Java.
- Physalospora camptospora** Sacc. On leaves of *Orchidaceae* in Brazil.
- Physalospora cattleyae** Maubl. and Lasn. On *Cattleya mossiae* in greenhouses in France.
- Physalospora orchidearum** P. Henn. On stems of *Laelia schilleriana* and *Tainia stellata* in the botanical gardens at Berlin.
- Physalospora vanillae** A. Zimm. On leaves of *Vanilla planifolia* in Java.
- Physalospora wildemaniana** Sacc. On leaves of *Orchidaceae* in Brazil.
- Phytophthora** sp. A very serious disease of the cultivated vanilla (*Vanilla planifolia*) due to a *Phytophthora* species or related fungus is known to occur in Seychelles and various South Sea Islands. There are no published records of the disease and no studies have been made of it.
- Phytophthora faberi** Maubl. This fungus, which causes one of the most serious diseases known, the black rot, and pod canker of rubber (*Hevea brasiliensis*), cacao (*Theobroma cacao*) and other economic plants, has been reported as attacking *Dendrobium macarthiae* in Ceylon.
- Puccinia aurea** Wint. Produces yellow to brown rust pustules on both leaf surfaces of *Monadenia rufescens* in south Africa.
- Puccinia cinnamomea** Diet. and Holw. Produces scattered cinnamon-colored rust pustules on the lower leaf surfaces of unknown species of *Orchidaceae* in Mexico.
- Puccinia habenariae** P. Henn. Produces brown rust pustules on both sides of the leaves and on stems of *Habenaria* spp. in east Africa.
- Puccinia orchidearum-phalaridis** Kleb. Characterized by yellow, then brown, rust pustules on the lower surfaces of leaves of *Gymnadenia conopsea*, *Habenaria bifolia*, *H. chlorantha*, *H. montana*, *Listera* sp., *L. ovata*, *Orchis incarnata*, *O. latifolia*, *O. maculata*, *O. majalis*, *O. mascula*, *O. militaris*, *O. morio*, *O. purpurea*, and *Phalaris arundinacea* in Denmark, France, Great Britain, Holland, Switzerland, Austria, and Germany.
- Puccinia satyrii** Syd. This rust forms yellow-brown rust pustules on the under sides of the leaves of *Satyrium carneum* in south Africa.
- Sclerotium orchidearum** P. Henn. Forms globose sclerotia on the stems 0.8 to 1.2 millimeter in diameter, white at first, then yellow, and finally brown-black, smooth, white within. Found on *Dichaea vaginata* and *Vanda tricolor* in the botanical gardens at Berlin.
- Septoria codonorchidis** P. Henn. On leaves of *Codonorchis poeppigii* in Chile.
- Septoria epipactidis** Sacc. Characterized by long light-colored spots on the leaves of *Epipactis violacea* and *Epipactis* sp. in Italy and Germany.
- Septoria gymnadeniae** Thuem. On leaves of *Gymnadenia conopsea* in Siberia.
- Septoria listerae** Allesch. On leaves of *Listera ovata* in Germany.
- Septoria orchidearum** West. Produces brown spots on leaves and leaf sheaths of *Coeloglossum viride*, *Habenaria bifolia*, *Listera ovata*, *Orchis latifolia*, *O. maculata*, *O. mascula*, and *O. morio* in Belgium, France, and Italy.

**ORCHIDACEAE**—Continued.

- Septoria posekensis** Sacc. On leaves of Orchidaceae in Siberia.
- Septoria thelymitrae** McAlp. On leaves of *Thelymitra aristata* in Australia.
- Stigmatodochis palawanensis** Syd. On living leaves of *Dendrobium* sp. in the Philippines.
- Stilbella bulbicola** P. Henn. On leaf bases of *Epidendrum* sp., *Gomezia planifolia*, *Oncidium pulvinatum*, *Sarcanthus pugioniformis*, and *Stanhopea* sp. in the botanical gardens at Berlin.
- Tubercularia** sp. Chlorotic or livid irregular areas on stems of *Vanilla planifolia* in Brazil.
- Tylenchus olesistus** Ritz-Bos. See Begonia.
- Uredo aurantiaca** Mortem. Yellow rust pustules on the leaves of *Oncidium cavendishianum* in greenhouses in France and Italy.
- Uredo carnosa** Speg. Rust on *Catasetum fimbriatum* in Brazil.
- Uredo cyrtopodii** Syd. Brown rust pustules on the under surfaces of the leaves of *Cyrtopodium* sp. and *Bletia* sp. in Brazil, Cuba, and Germany; in the latter case on imported plants.
- Uredo epidendri** P. Henn. Brown rust pustules on leaves of *Epidendrum* sp. in Brazil.
- Uredo guacae** Mayor. On *Epidendrum rigidum* and *E. umbellatum* in Ecuador.
- Uredo gynandrearum** Cda. Rust on leaves of *Habenaria maculosa* and other Orchidaceae in Cuba, Porto Rico, Trinidad, and Central and South America.
- Uredo lynckii** Berk. Rust on tropical species of *Spiranthes* growing in greenhouses in England.
- Uredo nigropunctata** P. Henn. Forms yellow rust pustules on the under sides of leaves of *Bletia patula* and *B. purpurea* in Cuba, Porto Rico, Haiti, Bahamas, South America, and Florida and on the leaves of *Stanhopea* spp. in Brazil.
- Uredo oncidii** P. Henn. Rust on leaves of *Oncidium lanceanum* in Brazil.
- Uredo phaji** Racib. Forms golden-yellow rust pustules on the lower sides of leaves of *Phaius blumei* in Java. This species is probably identical with *Hemileia phaji* Rac.
- Uredo pleurothallidis** Keissl. Rust found on plants of *Pleurothallis dinotherii* in greenhouses in Germany.
- Uredo pustulata** P. Henn. Characterized by brown rust pustules on both surfaces of the leaves of *Stenorrhynchus* spp., including *S. lanceolatus* in Brazil.
- Uredo satyrii** Mass. Rust on *Satyrium coriifolium* in the Union of South Africa.
- Uredo scabies** Cooke. Rust on small, irregular spots on both sides of the leaves of *Vanilla planifolia* in Colombia.
- Uredo wittmackiana** P. Henn. and Klitz. Rust on round yellow or brownish spots on the upper surfaces of leaves of *Epidendrum* sp. in Mexico.
- Uromyces citrifloris** Bab. (*U. thelymitrae* McAlp. [?]) Rust on leaves of *Microstylis* sp. or *Thelymitra* sp. in New Zealand.
- Uromyces joffrini** Delacr. Reported on the fruits of *Vanilla planifolia* from Tahiti. A doubtful species.
- Uromyces microtidis** Cooke. Produces irregular groups of yellowish-brown rust pustules on living leaves of *Microstylis perrifolia* and *M. pulchella* in New South Wales, Tasmania, and Chatham Island.
- Uromyces orchidearum** Cke. and Mass. This rust is characterized by golden-brown rust pustules, mostly on the upper surfaces of the leaves (aecia hypophyllous), of *Chiloglottis diphylla* and *C. gunnii* in New South Wales, Victoria, and Tasmania.
- Uromyces stenorrhynchi** P. Henn. Brown rust pustules on the lower surfaces of the leaves of *Stenorrhynchus* sp. in Peru.
- Uromyces thelymitrae** McAlp. The golden-yellow to chestnut-brown pustules are produced on the leaves and leaf sheaths, either in scattered groups or gregarious, on *Thelymitra aristata*, *T. antennifera*, *T. crinita*, and *T. flexuosa* in Victoria, Australia, and *T. javanica* in Java.
- ORCHIS.** Terrestrial orchids. See Orchidaceae.
- ORIGANUM.** MARJORAM. Sweet herbs cultivated for their ornamental flowers and foliage.
- Phyllosticta origani** F. Tassi. Small circular to irregular leaf spots on *O. heracleoticum* in Italy.
- Puccinia caulicola** Schneid. See Thymus.
- Puccinia rubsaameni** P. Magn. Rust on leaves of *O. vulgare* in Denmark, Switzerland, Germany, and Austria.
- Septoria origanicola** Allesch. and var *majoranae* Bres. Large, irregular, often confluent, brown leaf spots on *O. majorana* and *O. vulgare* in Germany.
- ORMOSIA.** NECKLACE TREE. Tropical trees.
- Puccinia ormosiae** Arth. Rust on leaves of *O. krugii* in Porto Rico.
- ORNITHIDIUM.** Epiphytic orchids. See Orchidaceae.
- ORNITHOGALUM.** STAR-OF-BETHLEHEM. Dwarf, hardy bulbous plants.
- Aecidium ornithogalum** Bub. Yellow rust pustules on the leaves of *O. tenuifolium* in Moravia. Possibly the same as *Puccinia liliacearum*.
- Heterosporium ornithogali** Klotzsch. On leaves of *Muscari comosum*, *M. neglectum*, *Ornithogalum* sp., and *Gagea* sp. in Great Britain, Yugoslavia, and Germany. Reported from Washington.
- Puccinia kalchbrenneriana** De Toni. Yellow rust pustules turning to brown on both leaf surfaces of *O. altissimum* in South Africa.
- Puccinia liliacearum** Duby. Rufous-brown rust pustules on the leaves of *Bellerophila romana*, *Endymion nutans*, *Hyacinthus ramosus*, *Muscari comosum*, *M. racemosum*, *Ornithogalum narbonneense*, *O. nutans*, *O. pyrenaicum*, *O. tenuifolium*, *O. umbellatum*, and *Scilla bifolia* in Great Britain, France, Russia, Italy, Belgium, Holland, Hungary, Austria, and Germany.
- Puccinia lojkaiana** Thuem. Black linear rust pustules on both leaf surfaces of *O. boucheanum*, *O. chloranthum*, *O. nutans*, *O. pyrenaicum*, *O. refractum*, *O. prasadrum*, *O. umbellatum*, *Muscari botryoides*, and *M. racemosum* in France, Italy, Hungary, Austria, India, and Germany.
- Puccinia ornithogali-thyrsoideis** Diet. Brown rust pustules on stems and leaves of *O. thyrsoideis* in the Union of South Africa.
- Septoria ornithogalea** Oud. Pale linear spots on stems and capsules of *O. pyrenaicum* and *O. umbellatum* in Holland.
- Septoria ornithogali** Pass. On leaves of *O. umbellatum* in Italy.
- Septoria ornithogalicola** Hóll. Large yellowish spots with indefinite margins on both surfaces of the leaves of *O. boucheanum* in Hungary.
- Synchytrium niesslii** Bub. Small galls, which are dirty-white in color with a brown band, are produced on the leaves of *O. umbellatum* in Austria.
- Urocystis ornithogali** Koern. Black powdery smut sori on the leaves of *O. umbellatum* in Germany. This species is said not to be distinct from *U. colchici*, which has been reported sparingly on other hosts in the United States.
- Uromyces maireanus** Syd. This rust forms yellow-brown pustules, which later become dark brown, on irregular pale spots on both leaf surfaces of *O. sessiliflorum* in Algeria.
- Uromyces ornithogali** Lév. See Gagea.
- Ustilago peglerae** Bub. A smut destroying the anthers of *O. lacteum* in the Union of South Africa.
- Uromyces scillarum** (Grev.) Wint. See Hyacinthus.

- ORNITHOPUS.** Leguminous plants, one species cultivated for forage.  
**Peronospora ornithopi** Gäum. Downy mildew on leaves of *O. perpusillum* in Denmark and Germany.  
**Uromyces ornithopodioides** Gz. Frag. Rust on leaves of *O. ichomocarpus* in north Africa.
- OROBUS.** See Lathyrus.
- OROXYLUM.** EAST INDIAN TRUMPET FLOWER. Ornamental Asiatic trees.  
**Phyllosticta oroxylonis** P. Henn. On leaves of *O. indicum* in Brazil.
- ORTHOSIPHON.** Annual or perennial herbs.  
**Puccinia pallida** Syd. Rust on leaves of *Orthosiphon* sp. in the Union of South Africa.
- ORYZA.** RICE. Annual grasses cultivated for grain.  
**Achlya proliferata** (Nees) De By. Seeds of *O. sativa* in nursery beds are covered by a cottony mycelium and rotted. The disease may also attack young shoots and roots of seedlings; in Formosa.  
**Ascochyta oryzae** Catt. On leaves of *O. sativa* in Italy, Japan, and Brazil.  
**Cercospora oryzae** Miy. Small brown linear spots on leaves and glumes of *O. sativa* in Japan, China, Brazil, Porto Rico, and Cuba. It is not certain that the western form is the same as the oriental. *Cercospora* sp. common throughout Louisiana may be the same.  
**Chaetophoma glumarum** Miy. Whitish areas on glumes of *O. sativa*, hindering formation of grain in Japan.  
**Chaetophoma oryzae** Cav. On sheaths, peduncles, and glumes of *O. sativa* in Italy.  
**Coniothyrium anomale** Miy. White spots with dark-brown borders, mostly at tips and along margins of leaves of *O. sativa* in Japan.  
**Coniothyrium brevisporum** Miy. Irregular white spots finally involving entire leaf area of *O. sativa* in Japan and China.  
**Coniothyrium japonicum** Miy. Brown spots, becoming whitish, commonly at tips and along margins of leaves of *O. sativa* in Japan.  
**Diplodia oryzae** Miy. On leaves and glumes of *O. sativa* in Japan.  
**Diplodiella oryzae** Miy. On leaves and glumes of *O. sativa* in Japan.  
**Entyloma oryzae** Syd. Leaf smut on *O. sativa* in the Philippines.  
**Gnomonia oryzae** Miy. On glumes of *O. sativa* in Japan.  
**Helminthosporium oryzae** v. B. de Haan. The "sesame" spot disease or seedling blight is one of the most serious diseases reported on rice, 10 to 60 per cent of the seedlings often being destroyed. It is known to occur in Japan, Formosa, Java, India, and the Philippines. A similar, if not identical, disease is known from Portuguese East India, Federated Malay States, Italy, and restricted areas in Louisiana. The disease is most destructive to seedlings, but also attacks leaves, culms, and panicles of older plants. In the case of seedlings, the tips of the cotyledons become dark brown and the fungus spreads to new leaves. Numerous small brown spots edged with yellow occur on the leaves. These spots coalesce, becoming irregular, and as a result the leaves die. Diseased culms are yellow or pale brown and the heads either can not push out at all or are bent and distorted. Brown or grayish spots may also occur on the various parts of the head. The disease has been transferred by inoculation to a number of other economic grasses. It has been reported from a restricted area in Louisiana, but it will be desirable to prevent further introductions, especially since other and more virulent strains may gain entrance.  
**Helminthosporium sigmoideum** Cav. Effuse black areas on sheaths, leaves, and stems of *O. sativa* in Italy.  
**Hendersonia oryzae** Miy. Brown spots on glumes and leaves of *O. sativa* in Japan. This disease is said to prevent the development of the grain.  
**Leptosphaeria iwamotoi** Miy. On leaves of *O. sativa* in Japan, Chosen, and Uganda.  
**Leptosphaeria salvinii** Catt. On leaf sheaths of *O. sativa* in Italy.  
**Melanomma glumarum** Miy. Brown, then paler, spots on glumes and culms of *O. sativa* in Japan, China, India, and the Philippines. Causes abortion of the grain.  
**Metasphaeria albescens** Thuem. Small brown-black spots on the leaves of *O. sativa*, increasing rapidly in size and becoming yellowish-brown, in Japan.  
**Metasphaeria oryzae** (Catt.) Sacc. On leaves of *O. sativa* in Italy.  
**Mosaic.** The mosaic disease of sugar-cane has been transmitted to rice experimentally, and might become of importance under favorable conditions.  
**Mycosphaerella hondai** Miy. White spots on leaves of *O. sativa* in Japan and Chosen.  
**Mycosphaerella malinverniana** Catt. On leaves of *O. sativa* in Italy and India.  
**Mycosphaerella oryzae** (Catt.) Sacc. This fungus attacks the leaves of *O. sativa*, checking the development of the grain. It occurs in Japan, Italy, Portugal, and Austria.  
**Mycosphaerella shiraiana** Miy. Gray to black spots on both leaf surfaces and occasionally on glumes of *O. sativa* in Japan and China.  
**Napicladium jenseanum** Rac. Linear red-brown spots on leaves of *O. sativa* in Java.  
**Nigrospora panici** Zimm. See Triticum.  
**Oospora oryzae** Sacc. This disease attacks the heads when they are near maturity, a whitish bloom spreading over the panicles. Diseased heads are largely blasted or destroyed. On *O. sativa* in the Philippines.  
**Ophiobolus cariceti** (B. and Br.) Sacc. (*Ophiochaeta graminis* Hara). The take-all disease which attacks wheat and other cereals so seriously in many parts of the world has been reported on rice (*O. sativa*) in Japan and Italy. See Triticum.  
**Ophiobolus oryzae** Miy. On leaves and glumes of *O. sativa* in Japan.  
**Phaeosphaeria cattanii** (Thuem.) Miy. On leaves and stems of *O. sativa* in Italy.  
**Phaeosphaeria oryzae** Miy. Brown areas beginning at the margins and tips of leaves and finally involving entire leaf area. Also on glumes of *O. sativa* in Japan.  
**Phyllosticta japonica** Miy. Dark-brown, then whitish, spots on both sides of leaves and on glumes of *O. sativa* in Japan. This disease finally destroys infected leaves.  
**Phyllosticta miurai** Miy. See Triticum.  
**Phyllosticta necatrix** Thuem. On culms, leaves, and leaf sheaths of *O. sativa* in Italy.  
**Phyllosticta oryzae** (Cke. and Mass.) Miy. Dark-brown irregular spots on leaf sheaths in Formosa, Japan, and India.  
**Pleosphaerulina oryzae** Miy. Large areas on leaves of *O. sativa* whitened, in Japan.  
**Pseudomonas oryzae** Ishiyama. This bacterium attacks the leaves, causing the "white speck" disease of *O. sativa* in Japan. The disease also attacks the stems and causes losses up to 20 per cent.  
**Puccinia graminis** Pers. forma *oryzae* Risso. A rust on rice (*O. sativa*) has been reported from Spain and other parts of Europe and named as above. A rust referred to as *Puccinia oryzae* has also been reported, but is doubtless the same. The rust involved here is a strain of the common and widespread black stem rust of cereals. This strain is unknown in the United States.  
**Pyrenochaeta oryzae** Shir. Diseased areas occur on the leaves and glumes of *O. sativa* in Japan. The leaves are greatly weakened and the plants stunted, since root development is poor.  
**Rhizoctonia destruens** Tass. See Solanum.

**ORYZA**—Continued.

- Sclerospora** sp. An undetermined species of *Sclerospora* has been reported on *O. sativa* in Japan. The species involved may possibly be *S. macrospora* which occurs to a limited extent in the United States, but this is probably not the case and this downy mildew should be as carefully guarded against as the similar oriental diseases on corn and sugar cane.
- Sclerospora macrospora** Sacc. See *Triticum*.
- Sclerotium** sp. A serious stem-rot disease due to an undetermined species of *Sclerotium* is reported from Java, Japan, China, and the Philippines. From India a similar disease is reported, attacking the leaf-sheaths and causing light-brown areas which ultimately spread over and kill infected plants. These diseases may be referable to one of the following species.
- Sclerotium glumale** Ces. Gray to black spots on glumes of *O. sativa* in Borneo.
- Sclerotium irregulare** Miy. Irregular to globular or oval rust-brown sclerotia produced on greenish-gray spots with dark brown margins on leaf sheaths of *O. sativa* in Japan.
- Sclerotium oryzae** Catt. The bases of stems of diseased plants are discolored at the lowest internodes, with dark-gray mycelium present in the hollow stems, in which appear numerous small, round, black, shining sclerotia. The disease may begin at a node some distance up the stem, involving the leaf sheaths at the same time. Diseased plants are stunted, the culms lodge and little or no grain is formed. This is said to be the most serious of known rice diseases. It occurs in Japan, Italy, India, Ceylon, Portugal, British Guiana, the Philippines, and to a very limited extent in Louisiana and Arkansas. The disease is carried for the most part on seed grain and gained entrance to the United States in this manner. Prohibition of further importations should be effective in preventing continued spread of the disease, provided proper steps are taken to eliminate present centers of infection.
- Septoria curvula** Miy. On leaves of *O. sativa* in Japan.
- Septoria longispora** Miy. Gray areas on leaves of *O. sativa* in Japan. Often sufficiently serious to check development of grain.
- Septoria miyake** Sacc. and Trav. On glumes of *O. sativa* in Japan and the Philippines.
- Septoria oryzae** Catt. On leaves, leaf sheaths, and glumes of *O. sativa* in Japan, China, Brazil, and Italy.
- Septoria poae** Catt. On leaves of *O. sativa* in Brazil.
- Sphaeronema oryzae** Miy. Produces white spots on glumes of *O. sativa* in Japan.
- Sphaeropsis japonicum** Miy. On glumes of *O. sativa* in Japan.
- Sphaeropsis oryzae** (Catt.) Sacc. On leaves and leaf sheaths of *O. sativa* in Italy and Brazil.
- Tilletia horrida** Tak. This smut forms black spore masses in the ovaries, generally concealed by the glumes, but at times forcing them apart. The smut is usually not visible until the grain is threshed. The disease occurs in Burma, India, British Guiana, Indo-China, Siam, Japan, China, and the Dutch Indies. It has been introduced into the United States on seed rice and was at one time serious in the Carolinas, but is said to have been eradicated. It also occurs sparingly in other Southern States, but not in California.
- Tylenchus angustus** Butl. The "ufra" disease of rice, known only from India, is caused by a nematode and is considered one of the most serious diseases of this crop. The worm attacks the aboveground parts of the plant. The ends of the leaves turn brown and this browning continues until the entire plant is involved. Plants are often stunted, with either withered or normal appearing leaves. Brown stains or lesions appear at one or more of the upper nodes just below the base of the peduncles, the stalks of which become deep brown or black, flaccid, and shrunken. The upper part of the stalk may be swollen into a spindle-shaped swelling composed of the ear and inclosing sheaths or the ear may push its way out. Normal grains as a rule are not produced. The nematodes may be carried on seed grain, or on empty glumes and other diseased plant parts.
- Ustilagoidea vires** (Cke.) Tak. The so-called green smut or false smut disease of rice attacks the individual grains which are transformed into hard green masses, more or less twice the size of normal grains. These sclerotia are orange-yellow near the surface and white at the center. The disease occurs in India, Ceylon, Japan, Formosa, China, Sumatra, Malacca, Indo-China, Portuguese East India, Philippines, and Straits Settlements. It has been sparingly introduced into the southern United States, but is not known in California.
- ORYZOPSIS.** Tufted perennial grasses.
- Puccinia oryzopsidis** Syd. and Butl. Yellow and dark-brown rust pustules on leaves of *O. molinoides* in India.
- Septoria trapezuntica** Bub. On leaves of *O. miliacea* in Asia Minor.
- OSBECKIA.** Tropical herbs and shrubs.
- Rehmiodothis osbeckiae** (B. and Br.) Theiss. and Syd. Shining black convex stromata on leaves of *O. virgata* in Ceylon.
- OSMANTHUS.** Ornamental woody plants grown for their handsome foliage and white, fragrant flowers.
- Accidium osmanthi** Syd. and Butl. Rust on leaves of *Osmanthus* sp. in India.
- Phyllosticta osmanthi** Tass. White leaf spots on *O. aquifolius* in Italy.
- Phyllosticta osmanthicola** Trinch. On leaves of *O. fragrans* in Italy.
- Septoria osmanthi** Tass. Irregular white spots on leaves of *O. aquifolius* in Italy.
- OSMORHIZA.** Aromatic herbs with edible roots.
- Puccinia philippii** Diet. and Neg. Yellow and brown rust sori on leaf blades and petioles of *O. berterii* and *O. glabrata* in Chile and Argentina.
- OSMUNDA.** Royal fern.
- Mycosphaerella collista** Syd. On fronds of *O. regalis* in Germany.
- OSTEOSPERMUM.** Yellow-flowered shrubs.
- Ascochyia atro-punctata** Wint. Circular, often confluent, yellow leaf spots on *O. moniliferum* in the Union of South Africa.
- OSTRYA.** HOP HORNBEEAM. Ornamental and timber trees.
- Exoascus ostryae** Mass. Brown spots on leaves of *O. carpinifolia* in Italy.
- Gloeosporium ostryae** Thuem. Irregular dark-brown anthracnose spots on leaves of *O. virginiana* in Portugal.
- Gnomonia ostryae** DeN. On leaves of *O. carpinifolia* in Italy.
- Mamiania ostryae** (P. Henn.) Theiss. and Syd. On *O. japonica* in Japan.
- OURISIA.** Small herbs used in border plantings.
- Puccinia apus** Diet. and Neg. Brown rust sori on leaves of *O. racemosa* in Chile.
- Puccinia ourisiae** Diet. and Neg. Rust on leaves of *O. alpina*, *O. coccinea*, and *O. pygmaea* in Chile.
- Puccinia quillensis** Diet. and Neg. Powdery rust pustules on leaves of *Ourisia* sp. in Argentina.
- OXALIS.** Bulbous or tuberous herbs with cloverlike leaves; a few species cultivated for their edible roots or tubers.
- Cercospora oxalidis** Grove. Small circular leaf spots on *O. acetosella* in Great Britain.
- Cylindrosporium oxalidis** Traill. Circular brown leaf spots on *O. acetosella* in Scotland.

**OXALIS**—Continued.

- Mycosphaerella depazeaeformis** (Ces. and DeN.) Wakef. Circular, rufous leaf spots on *O. acetosella*, *O. corniculata*, and *O. stricta* in Uganda and Germany.
- Ovularia oxalidis** Oud. On leaves of *O. tropaeoloides* in Holland.
- Phyllosticta oxalidicola** Speg. On leaves of *Oxalis* sp. in Argentina. A doubtful species.
- Phyllosticta oxalidicola** P. Henn. Yellow-brown leaf spots on *Oxalis* sp. in Jamaica.
- Puccinia oxalidis** Diet. and E. Brown rust pustules on leaves of *O. newwiedii* and *Oxalis* sp. in Mexico and Brazil.
- Septoria oxalidis** Lind. Circular ashen leaf spots with dull-brown margins on *O. acetosella* in Denmark.
- Septoria oxalidis-japonicae** Pat. Sunken rufous to dark-brown leaf spots on *O. japonica* in Tibet.
- Stagonospora hygrophila** Sacc. On leaves of *O. acetosella* in Great Britain.
- Stigmatea depazeaeformis** (Auersw.) Schroet. On leaves of *O. acetosella* in Austria and Germany. Possibly a form of *Mycosphaerella depazeaeformis*.
- Uredo oxalidis** Lévl. Leaf rust on *O. corniculata*, *O. corymbosa*, *O. martiana* and *O. stricta* in Ecuador, Argentina, France, Switzerland, and Belgium.
- Uromyces oxalidis** Pазschke. Rust on petioles and scapes of *Oxalis* sp. in Brazil.
- OXYCOCCUS.** Cranberry. See also *Vaccinium*.
- Ascochyta oxycocci** P. Henn. On leaves of *O. macrocarpus* in Germany.
- Sclerotinia oxycocci** Woron. Gray mold on branches and leaves and soft rot of fruit of *O. palustris* and *O. quadrifidus* in Denmark and Finland.
- OXYPETALUM.** Herbs and subshrubs.
- Puccinia lagoensis** P. Henn. Black rust pustules on leaves of *Oxypetalum* sp. in Brazil.
- Puccinia oxypetali** P. Henn. Stems shortened, swollen, and completely deformed, dark-brown rust pustules developing on the swollen areas and on leaves and flower buds on *O. banksii* in Brazil.
- Septoria oxypetali** F. Tassi. Small white leaf spots with black margins on *O. caeruleum* in Italy.
- Uromyces cabo-friensis** P. Henn. Black rust pustules on leaves of *Oxypetalum* sp. in Brazil.
- OXYTROPIS.** Perennial herbs and shrubs.
- Ovularia oxytropidis** Jacz. On leaves of *O. pilosa* in Russia.
- Peronospora oxytropidis** Gaum. Downy mildew on leaves of *O. campestris* in Switzerland.
- Phyllosticta oxytropidis** Höll. Circular ochraceous leaf spots on *O. pilosa* in Hungary.
- Septoria oxytropidis** Died. On leaves of *O. pilosa* in Germany.
- Septoria oxytropidis** E. Rostr. Large ashen leaf spots on *O. lapponica* in Norway.
- Uromyces astragali** (Opiz.) Sacc. See *Astragalus*.
- PACHYRHIZUS.** CACARA. Ag. Yam bean. Jicama. Climbing herbs with large tuberous roots used for food and starch production.
- Phakopsora pachyrhizi** Syd. Dull-brown rust sori on leaves of *P. angulatus* and *P. erosus* in Japan, Formosa, and the Philippines.
- PACOURIA.** See *Landolphia*.
- PAEONIA.** PEONY. Showy spring and autumn blooming herbaceous perennials.
- Aecidium paeoniae** Kom. Leaf rust on *P. albiflora* in Russia.
- Botrytis paeoniae** Oud. This fungus causes a blighting of leaves and buds and a rotting of stems. Black sclerotia are formed on old stems which enable the fungus to live over from season to season. The disease is reported as serious in Great Britain, Holland, Germany, and other parts of Europe. All varieties of cultivated paeonies are subject to attack as well as *Convallaria*. The connection of this disease with the American *Botrytis* blight is in doubt.
- Cronartium flaccidum** (Alb. and Schw.) Wint. (*C. paeoniae* Cast.) A rust producing yellow-brown to red-brown pustules on the leaves. Infected leaves present the appearance of being studded with long hairs on the under surfaces. One stage of the rust occurs on *Pinus silvestris*. It also occurs on *Asclepias pneumonanthis*, *A. speciosa*, *Cynanchum fuscatum*, *C. nigrum*, *C. purpurascens*, *Crammatocarpus volubilis*, *Impatiens balsamina*, *Nemesia versicolor*, *Paeonia albiflora*, *P. angustifolia*, *P. anomala*, *P. arborea*, *P. arietina*, *P. broterii*, *P. corallina*, *P. daurica*, *P. decora*, *P. edulis*, *P. fimbriata*, *P. formosa*, *P. grandiflora*, *P. herbacea*, *P. hybrida*, *P. moutan*, *P. officinalis*, *P. perigrina*, *P. rosea*, *P. sessiliflora*, *P. sinensis*, *P. splendens*, *P. tenuifolia*, *Pedicularis palustris*, *Schizanthus grahami*, *Tropaeolum canariensis*, *T. lobbianum*, *T. major*, *T. minor*, *Verbena erinoides*, *V. teucrioides*, *V. officinalis*, *Vincitoxicum laxum*, and *V. officinalis* in Europe, Siberia, and Japan.
- Leptothyrella paeoniae** P. Henn. On leaves of *P. obovata* in Japan.
- Leptothyrium paeoniae** Vogl. Circular to oblong pale-yellow spots on leaves of *P. arborea* in Italy.
- Mycosphaerella moutan** Siem. Leaf spot on *P. moutan* in Russia.
- Phyllosticta baldensis** Massal. A leaf spot producing fungus on *P. decora*, *P. officinalis*, and *P. peregrina* in Denmark, Switzerland, Bulgaria, and Italy.
- Phyllosticta moutan** Pass. Dark-brown spots on leaves of *P. moutan* in Italy.
- Phyllosticta paeoniae** S. and P. Produces indefinite spots on leaves of cultivated *Paeonia* varieties, including *P. corallina* in Great Britain.
- Puccinia martianoffiana** Thuem. Small sunken spots in which appear brown-black rust pustules on both surfaces of the leaves of *P. anomala* in Siberia.
- Ramularia paeoniae** Vogl. On leaves of *P. peregrina* in Italy.
- Ramularia sardoa** Sacc. and Trav. On leaves of *P. corallina triternata* in Italy.
- Septoria macrospora** Sacc. On leaves of *P. officinalis* in Italy and Russia.
- Septoria martianoffiana** Thuem. Produces a leaf spot on *P. anomala* in Siberia.
- Septoria servica** Syd. Suborbicular to irregular brown spots with black margins on both surfaces of leaves of *P. decora* in Yugoslavia.
- PALA.** See *Alstonia*.
- PALIURUS.** Woody plants, sometimes grown for their attractive foliage and curiously-shaped fruits.
- Ascochyta paliuri** Sacc. Yellow-brown leaf spots on *P. aculeatus* in Italy.
- Phyllosticta camusiana** Sacc. Leaf spots on *P. aculeatus* and *P. australis* in France and Italy.
- Phyllosticta paliuri** (Lévl.) Cke. Black leaf spots on *P. australis* in Russia.
- Septoria ascochyella** Sacc. Olivaceous or dull-yellow leaf spots on *P. aculeatus* in Italy.
- PALMAE.** PALMS. The diseases of palms are considered in one group for reasons similar to those explained under *Bambuseae*.
- Aphelenchus cocophila** Cobb. The red-ring disease of the coconut (*Cocos nucifera*) is serious in British Guiana, Trinidad, Tobago, Grenada, and Panama. It doubtless occurs in other parts of the West Indies and Central America. Externally it is marked by a yellowing and browning of leaves which works from below upwards. This phase takes place fairly rapidly, involving even the unfolded leaves. Green nuts of all sizes and the flowers are shed, the branches of the inflorescences withering. The bud rots. Internally a red zone appears in the stem, extending upward several feet, then breaking into longitudinal streaks, and finally into scattered dots. Similar red spots or streaks occur in the leaf petioles. The roots are discolored and finally die. The reddened tissues are filled with the nematodes which cause the disease.

## PALMAE—Continued.

- Ascochyta depazeoides** (Dur. and Mont.) Tass. On leaves of *Phoenix dactylifera* and *Chamaerops humilis* in Portugal and Algeria.
- Ascochyta kentiae** Maubl. Elongate pale-brown leaf spots with dark-brown borders on *Kentia* sp. in Algeria.
- Ascochyella passeriniana** (Thuem.) Tass. On leaves of *Chamaerops* sp. and *Phoenix* sp. in Italy.
- Ascochyella pinnarum** (Pass.) Tass. On leaves of *Phoenix dactylifera* in Italy.
- Auerswaldia decipiens** Rehm. A doubtful species of tar-spot fungus on *Arenga mindorensis* and *A. tremula* in the Philippines.
- Bacillus coli** Esch. See Bud-rot.
- Bagnisiopsis bactridis** (Rehm.) Theiss. and Syd. Black stromata on leaves of *Bactris* sp. in Brazil.
- Bagnisiopsis diplothemii** (Rehm.) Theiss. and Syd. Dull-black stromata on leaves of *Diplothemium littorale* in Paraguay.
- Botryosphaeria palmigena** (B. and C.) Bomm. and Rouss. On leaves of *Chamaedorea* sp. in Costa Rica.
- Brachysporium arecae** (B. and Br.) Sacc. Leaf spot on *Areca catechu* in India and Ceylon.
- Bud rot.** A disease of coconuts and other palms, characterized by a soft rot of the bud which destroys the tree, has been reported from nearly all tropical palm-growing regions and constitutes the most serious menace to these valuable plants. Workers on the problem do not agree as to the causative organism, and in fact several are undoubtedly concerned. The form of the disease on *Cocos nucifera* and *Elaeis guineensis* in Cuba, Jamaica, Santo Domingo, Cayman Islands, Trinidad, British Guiana, Surinam, British Honduras, and possibly also in the eastern tropics has been attributed to a strain of *Bacillus coli* Esch.
- The disease is first noted through the falling of immature nuts, staining of opening flower spikes to a rich chocolate brown and the dying and bending over of center leaves. A dark wet rot develops at the base of affected spikes and spreads around the leaf sheaths. Finally the entire bud is destroyed by a foul-smelling soft rot, the leaves are brown and dead, hanging downward and all nuts have fallen. The entire top of a diseased tree ultimately falls away, leaving the bare dead trunk. Losses have been very large. For further discussion of bud-rot diseases see *Phytophthora*.
- Camarotella astrocaryae** (Rehm.) Theiss. and Syd. Black stromata on both leaf surfaces of *Astrocaryum* sp. in Brazil.
- Catacauma mucosum** (Speg.) Theiss. and Syd. (*Phyllachora cocoicola* P. Henn.) Tar spot on leaves of *Cocos australis* in Brazil.
- Cercospora acrocomiae** Stevenson. Oval red-brown, then gray, leaf spots with dark-brown margins on *Acrocomia media* in Porto Rico.
- Cercospora calamicola** P. Henn. Subcircular, black leaf spots on *Calamus caryotoides* in Queensland.
- Cercospora licualae** Syd. On leaves of *Licuala spinosa* in the Philippine Islands.
- Cercospora palmicola** Speg. Elliptical, often confluent, leaf spots on *Cocos australis* in Brazil.
- Cercospora preisii** Bub. On leaves of *Phoenix* sp. in Russia.
- Cercospora virens** Sacc. On leaves of *Licuala* sp. in Malaya.
- Colletotrichum arecae** Syd. Anthracnose on leaves of *Areca catechu* in the Philippines.
- Colletotrichum paucisetum** Petch. Anthracnose on leaves of *Cocos nucifera* in Ceylon.
- Coniothyrium cocois** P. Henn. Dull-brown leaf spots on *Cocos nucifera* in east Africa.
- Coryneum cocoës** P. Henn. Dull-brown leaf spots with darker brown margins on *Cocos nucifera* in east Africa.
- Didymosphaeria astrocaryi** v. Hoeh. On leaves of *Astrocaryum vulgare* in Guiana.
- Diplodia cacaicola** P. Henn. See Theobroma.
- Dothidina palmicola** (Speg.) Theiss. and Syd. (*Auerswaldia palmicola* Speg.) Irregular to linear brown-black stromata on leaves of *Acrista monticola*, *Cocos yatay*, *Diplothemium littorale*, *Phoenix dactylifera*, and *Trithrinax brasiliensis* in Porto Rico, Brazil, Argentina, Paraguay, and Uruguay.
- Exosporium durum** Sacc. The leaves of *Cocos nucifera* in the Philippines turn yellow and become covered with small spots on which occur numerous dark-colored raised fruiting bodies.
- Exosporium pulchellum** Sacc. Numerous small, dark-colored, raised fruiting bodies on leaves of *Areca catechu* in the Philippines.
- Fomes lamaoensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Fusicladium livistonae** Karst. Black fungus layers on petioles of *Livistona sinensis* in Denmark.
- Gloeosporium catechu** Syd. Anthracnose on leaves of *Areca catechu* in the Philippines.
- Gloeosporium coccophilum** Wakef. Anthracnose on leaves of *Cocos nucifera* in Ceylon.
- Gloeosporium ptychospermatidis** P. Henn. Large white spots on leaves of *Ptychosperma* sp. in Java.
- Graphiola applanata** Syd. and Butl. Small black, hard, raised fruiting bodies on leaves of *Phoenix sylvestris* in India.
- Graphiola arengae** Rac. Hard raised fruiting bodies on circular yellow leaf spots on *Arenga ambong* and *A. saccharifera* in Java and the Philippines.
- Graphiola borassi** Syd. and Butl. On leaves of *Borassus flabelliferus* in India.
- Graphiola cocoina** Pat. Small black, raised, hard fruiting bodies on small brown-leaf spots on *Cocos nucifera* in Fiji and Gambier Islands.
- Graphiola cylindrospora** Syd. On leaves of *Livistona chinensis* in the Philippines.
- Graphiola macrospora** Penz. and Sacc. Globular black, hard fruiting bodies on leaves of *Calamus* sp. and *Plectocomia* sp. in Java.
- Haplosporella asterocaryi** P. Henn. On leaves of *Astrocaryum* sp. in Brazil.
- Helminthosporium coryphae** Syd. On leaves of *Corypha elata* in the Philippines.
- Helminthosporium incurvatum** Ch. Bern. On leaves of *Cocos nucifera* in Ceylon and Java.
- Helminthosporium philippinum** Sacc. On *Arenga tremula* in the Philippines.
- Heterosporium chamaeropsis** Oud. On leaves of *Chamaerops excelsa* in Holland.
- Heterosporium minutulum** C. and M. On leaves of *Chamaerops humilis* in Great Britain.
- Hysterostomella elaeicola** Maubl. Irregular black carbonous fruiting bodies in pale sunken leaf spots on *Elaeis guineensis* in Dahomey.
- Laestadia cocophila** (Cke.) Sacc. On leaves of *Cocos nucifera* in British Guiana.
- Lembosia diplothemii** P. Henn. On leaves of *Diplothemium maritimum* in Brazil.
- Leptosphaeria briosiana** Pöhl. On leaves of *Jubaea spectabilis* in Italy.
- Leptosphaeria molleriana** d'Alm. and da Cam. On leaves of *Cocos romanzoffiana* in Portugal.
- Leptostroma pheenicis** Oud. and Fautr. On leaves of *Phoenix dactylifera* in France.
- Meliola furcata** Lévl. Black superficial fungus patches on leaves of *Acrista monticola* and *Coccothrinax alta* in Porto Rico.
- Melittosporiopsis gigantospora** Rehm. On leaves of *Rhaphidophyllum* sp. in the Philippines.
- Melittosporiopsis pachycarpa** Rehm. On leaves of *Rhaphidophyllum* sp. in Chile.
- Merrilllopettis hoehnelii** Rehm. On petioles of *Arenga mindorensis* and *A. saccharifera* in the Philippines.

## PALMAE—Continued.

- Metasphaeria cocogena** (Cke.) Sacc. On leaves of *Cocos nucifera* in Demerara.
- Mollisia copelandi** Rehm. On leaves of *Caryota* sp. in the Philippines.
- Mycosphaerella** sp. Leaf spot on *Roystonea regia* in Cuba.
- Mycosphaerella chamaeropsis** Trav. Irregular ochraceous-gray leaf spots with broad purple margins on *Chamaerops humilis* in Italy.
- Mycosphaerella elaeidis** Beeli. Large irregular gray leaf spots on *Elaeis* sp. in Belgian Congo.
- Mycosphaerella gastonis** (Sacc.) Lind. On leaves of *Cocos nucifera* in Samoa, Tahiti, Ceylon, and the Philippines.
- Mycosphaerella ptychospermatis** Rehm. On leaves of *Ptychosperma macarthurii* in the Philippines.
- Ovularia palmicola** Pantanelli. On *Chamaerops* sp. and *Pritchardia* sp. in Italy and Tripoli.
- Palawania grandis** (Niessl.) Syd. Black stromata on leaves of *Calamus* sp. and *Oncosperma horrida* in India, Java, and the Philippines.
- Parmulineae rehmii** (Maubl.) Theiss. and Syd. Dull-black stromata on yellow indefinite leaf spots on *Bactris* sp. in Brazil.
- Pestalozzia brevipes** Prill. and Delacr. Irregular yellow leaf spots on *Corypha australis* and *Kentia chamaerops* in France.
- Phaeochora acrocomiae** (Mont.) Theiss. and Syd. Black stromata on leaves of *Acrocomia sclerocarpa* and *A. yatay* in French Guiana, Brazil, and Paraguay.
- Phaeochora chamaeropsis** (Cke.) v. Hoeh. Shining black stromata on brown leaf spots on *Chamaerops humilis* in Morocco and Spain.
- Phaeotrabiella perisporioides** (Sacc.) Theiss. and Syd. Black stromata on circular to irregular leaf spots on *Phoenix dactylifera* in Oceania.
- Phomatospora migrans** Rehm. Large irregular, concentrically zoned areas on leaves of *Arenga saccharifera* in the Philippines.
- Phomopsis palmicola** (Wint.) Sacc. and var. **arecae** Sacc. Extensive irregular dead areas on leaves of *Areca catechu* in the Philippines.
- Phyllachora calamigena** (B. and Br.) Sacc. Brown stromata on brown leaf spots on *Calamus rudentum* in Ceylon.
- Phyllachora palmicola** Speg. Black stromata on long yellowish leaf spots on *Copernicia cerifera* and *Trithrinax brasiliensis* in Brazil and Argentina.
- Phyllachora roystoneae** Johnst. and Brun. Conspicuous black carbonaceous stromata on midribs of leaves of *Roystonea regia* in Cuba.
- Phyllosticta** sp. Leaf spot on *Caryota urens* in India.
- Phyllosticta chamaeropsis** Pöhl. Oblong brown leaf spots on *Chamaerops* sp. in Italy.
- Phyllosticta cocoës** Allesch. Small oblong whitish leaf spots on *Cocos* sp. in Brazil.
- Phyllosticta cocoina** Sacc. Subcircular leaf spots on *Cocos nucifera* in Tahiti.
- Phyllosticta cocophila** Pass. Oblong to irregular whitish leaf spots with dull-brown margins on *Cocos flexuosa* and *C. nucifera* in Italy and the Philippines.
- Phyllosticta cocos** Cke. Irregular pale-brown leaf spots on *Cocos nucifera* in India and Italy.
- Phyllosticta daemonoropsis** Sacc. On leaves of *Daemonorops* sp. in Malaya.
- Phyllosticta palmigena** Sacc. On leaves of *Dictyosperma alba* in Malaya.
- Phyllosticta sabalioia** Szabo. On petioles of *Sabal blackburniana* in Hungary.
- Physalospora calami** Syd. Oblong, white leaf spots with brown margins on *Calamus tenuis* in India.
- Phytophthora arecae** (Colem.) Pethyb. The "koleroga" disease of the betel palm (*Areca catechu*) is serious in India. The fungus has also by inoculation been found capable of causing a rot of potato (*Solanum tuberosum*) tubers. The leaves of diseased palms turn yellow and droop, the flowers and immature fruit fall, and the flower stalks blacken and rot. This rot often spreads to the leaf sheaths and apical bud. The disease is one of the bud-rot type.
- Phytophthora faberi** Maubl. This downy mildew fungus has been found in Porto Rico, Jamaica, Ceylon, and the Philippines, causing typical bud rot of *Cocos nucifera* and other palms. The symptoms given for bud rot under that heading above apply here. The fungus has also been recently reported from limited areas in Florida in connection with the death of coconut palms. See also *Theobroma* for a further account of the fungus.
- Phytophthora melongenae** K. Saw. See *Solanum*.
- Phytophthora palmivora** Butler. The bud-rot disease of coconut (*Cocos nucifera*) and Palmyra palm (*Borassus flabellifer*) in India, Ceylon, Fiji, and Jamaica is attributed to this species. With this disease the inner leaves turn brown and break over and the heart leaf can usually be drawn out, having been rotted off at the base. All the young leaves turn yellow and the center column falls out, leaving a ring of the older leaves which may persist for some time. Young nuts cease growing. During this early stage brown sunken spots occur at the bases of young leaves and extend through to the bud, which soon becomes a pasty mass with an offensive odor. The inner and outer faces of the spots on leaf bases may show a white mildew. Development of the disease continues, the leaves all falling off and the tree dying.
- Phytophthora parasitica** Dastur. See *Ricinus*.
- Piacostroma diplothemii** Syd. Black shining stromata on leaves of *Diplothemium maritimum* in Brazil.
- Pleospora kentiae** Maubl. On leaves of *Kentia* sp. in Algeria.
- Pucciniopsis guaranitica** Speg. Small black fruiting pustules on brown leaf spots on *Cocos yatay* in Brazil.
- Ramularia thrinciae** Rth. On leaves of *Thrinax nudicaulis* in Madeira.
- Rhipidocarpon javanicum** (Pat.) Theiss. and Syd. On leaves of *Nipa fruticans* in Java and the Philippines.
- Septoria calami** P. Henn. Pale indefinite leaf spots on *Calamus caryotoides* in Queensland.
- Septoria cocoës** Petch. On leaves of *Cocos nucifera* in Ceylon.
- Septoria martiniezeae** Thum. Irregular dull-brown leaf spots on *Martinezia caryotaefolia* in Portugal.
- Septoria palmarum** Sacc. On leaves of *Corypha elata*, *C. umbraculifera*, and *Latania borbonica* in Italy and the Philippines.
- Sphaerodothis arengae** (Racib.) Shear. Black, shiny, irregular stromata on leaves of *Arenga obtusifolia*, *A. saccharifera*, *Caryota propinqua*, and *Caryota* sp. in Java and the Philippines.
- Stagonospora kentiae** Maubl. Whitish leaf spots with brown margins on *Kentia* sp. in Algeria.
- Thelaviopsis paradoxa** (DeSeyn.) v. Hoeh. Causes the "stem-bleeding" disease of *Cocos nucifera* in Ceylon and other parts of the oriental Tropics. See *Saccharum* for further discussion.
- Trabuttia atroquinans** (Wint.) Theiss. and Syd. Black circular to elliptical stromata on leaves of *Mauritia flexuosa* in Brazil.

## PALMAE—Continued.

**Ustilago dactylicola** Speg. In fruit of *Phoenix dactylifera* in central Africa. Beyond doubt only *Aspergillus niger* again. *Ustilago phoenicis* Cda. reported from Egypt on the same substratum is doubtless also the same.

**Zukalia stuhlmanniana** P. Henn. Black superficial fungus layers (sooty mold) on leaves of *Cocos nucifera*, *Phoenix dactylifera*, and *Ravenala madagascariensis* in Zanzibar and Tanganyika.

The "baïoud" or "white" disease attacks *Phoenix dactylifera* in Algeria, often destroying the trees. The cause is unknown. Other diseases of the trunks, leaves, and fruit of this host are also reported from Algeria, the causes likewise being unknown.

## PANAX. GINSENG. Hardy perennials with aromatic roots.

**Bacterium araliavorus** Uyeda. Causes a red rot of stems and roots of *P. quinquefolium* in Korea.

**Bacterium panaxi** Nak. and Tak. Produces an amber-colored rot in roots and a brown rot of stems of *P. quinquefolium* in Japan and Korea.

**Colletotrichum panacicola** Nak. and Tak. Leaf blight of *P. quinquefolium* in Korea.

**Phoma panacicola** Nak. and Tak. Black dry rot of roots of *P. quinquefolium* in Korea.

**Phoma panacis** Nak. and Tak. Produces brown to silvery-gray spots and cankers on stems of *P. quinquefolium* in Korea.

**Phyllosticta panax** Nak. and Tak. Brown leaf spots on *P. quinquefolium* in Korea; called the "snake-eye" disease.

**Poria hypobrunnea** Petch. See Hevea.

**Rosellinia arcuata** Petch. See Thea.

**Rosellinia bunodes** B. and Br. See Citrus.

**Uredo panacis** Syd. Dull-yellow rust pustules on leaves of *P. pseudoginseng* in India.

## PANCRATIUM. Summer and winter flowering bulbs.

**Aecidium amaryllidis** Syd. and Butl. Leaf rust on *Pancratium* sp. in India.

## PANDANUS. SCREW PINE. Tropical woody plants and small trees remarkable for their aerial roots and the spiral arrangement of the long leaves.

**Anthostomella pandani** (Rabh.) Sacc. On leaves of *P. furcatus* in India.

**Ellisiodothis pandani** Syd. Black opaque stromata on leaves of *P. luzonensis* in the Philippines.

**Leptosphaeria pandani** Tass. On leaves and sheaths of *P. utilis* in Italy.

**Phyllosticta pandanicola** Young. Grayish-white to light-brown spots on leaves of *Pandanus* sp. in Porto Rico.

**Pirostoma farnetianum** Póll. On leaves of *P. utilis* in Italy, causing much damage.

**Plenodomus inaequalis** Sacc. and Trott. On leaves of *P. utilis* in Italy.

**Stigmatea pandani** Pat. Circular dull-brown leaf spots on *P. odoratissimus* in Gambier Islands.

## PANICULARIA. See Glyceria.

## PANICUM. GUINEA GRASS. PARA GRASS. BROOMCORN MILLET. A very large genus of annual and perennial grasses. Many are weeds, but others of importance as grain and forage plants.

**Acrothecium lunatum** Wak. Small brown leaf spots on *P. frumentaceum*, *Eleusine coracana*, and *Setaria italica* in India.

**Balansia claviceps** Speg. Forms sclerotia in spikelets of *P. carinatum* and *P. nodosum* in the Philippines and Argentina.

**Balansia paspali** P. Henn. Forms sclerotia in spikelets of *P. amplexicaule* in the Philippines.

**Balansia trinitensis** Cke. and Mass. Hard black globose stromata in place of ovaries on *P. palmifolium* in Trinidad.

**Balansiella pulvinula** (B. and Br.) Petch. Forms white to honey-colored stromata in young shoots of *Panicum* sp. in Ceylon, arresting their development.

**Beniowskia graminis** Rac. On leaves of *P. nepalense* in Java.

**Chaetodiplodina graminicola** Speg. On leaves of *P. laxum* in Argentina.

**Chlorochytrium graminis** Buesgen. See Festuca.

**Cerebella antidotale** Subra. Black corrugated fungus masses replacing ovaries of *P. antidotale* in India.

**Cerebella burmanensis** Subra. Black corrugated fungus masses in ovaries of *P. setigerum* in India.

**Cerebella cynodontis** Syd. See Cynodon.

**Cerebella inodantis** (B. and Br.) Petch. On *P. javanicum* and *Paspalum longiflorum* in India, *P. royleanum*, *P. (Syntherisma) sanguinale*, and *P. scrobiculatum* in India and Ceylon.

**Mycosphaerella panicicola** P. Henn. On leaves of *Panicum* sp. in Brazil.

**Phaeodothis stenostoma** (E. and E.) Theiss. and Syd. Black stromata on leaves of *P. brizanthum* in central Africa.

**Phyllachora acutispora** Speg. Dull-black stromata on upper leaf surfaces of *Panicum* sp. in Brazil.

**Phyllachora apiculata** Speg. Small linear black stromata on leaves of *P. grumosum* in Argentina.

**Phyllachora bokensis** P. Henn. Dull-black stromata on oblong dull-brown leaf spots on *Panicum* sp. in the Congo.

**Phyllachora bonariensis** Speg. Small, subcircular, smooth black stromata on leaves of *P. bambusoides* in Venezuela and Argentina.

**Phyllachora congruens** Rehm. Black stromata on leaves of *P. carinatum* in the Philippines.

**Phyllachora heterospora** P. Henn. Shiny black stromata on large yellow leaf areas on *P. maximum* and *Panicum* sp. in Ecuador, the Congo, and the Union of South Africa.

**Phyllachora panici** (Rehm.) Theiss. and Syd. Dull-black stromata on brown leaf spots on *Panicum* sp. in Brazil.

**Phyllachora panici-sulcati** (P. Henn.) Theiss. and Syd. Black stromata on leaves of *P. sulcatum* in Costa Rica.

**Phyllachora pazschkeana** Syd. Circular or elliptical black stromata on leaves of *P. pilosum* and *P. sciurotes* in Brazil and Colombia.

**Phyllachora raciborskii** Theiss. and Syd. Black stromata on leaves of *P. nepalense* in Java.

**Phyllachora sanguinolenta** Theiss. and Syd. Shiny black circular stromata on leaves of *P. sanguinolentum* in the Congo.

**Phyllachora seriata** Theiss. and Syd. Black shiny oblong stromata on yellow leaf spots on *P. palmaefolium* and *Panicum* sp. in the Philippines.

**Phyllachora stenospora** (B. and Br.) Sacc. Tar spot on leaves of *P. patens* and *P. trigonum* in Ceylon and the Philippines.

**Phyllachora vanderystii** Theiss. and Syd. Black elliptical stromata on leaves of *P. plicatum* in Ceylon and the Congo.

**Phyllachora urvilleana** Speg. Tar spot on leaves of *P. urvilleanum* in Argentina.

**Phyllosticta panici** Young. Indefinite brown leaf spots on *P. maximum* in Porto Rico.

**Phylospora panici** Rehm. On *Panicum* sp. in Brazil.

**Puccinia goyazensis** (P. Henn.) Syd. Rust on leaves of *Panicum* sp. in Brazil.

**Puccinia huberi** P. Henn. Brown to black rust pustules on leaves of *P. fasciculatum*, *P. ovalifolium*, *P. trichoides*, *P. paniculatum*, *P. tenellum*, *P. ulowanaceum*, and *Paspalum virgatum* in Porto Rico, Cuba, Mexico, and Brazil.

**PANICUM**—Continued.

- Puccinia insolita** Syd. Powdery brown elongate rust sori on leaves of *P. maximum* in the Congo.
- Puccinia negrensis** P. Henn. Brown rust pustules on dull-brown leaf spots on *P. negrensis* in Brazil.
- Puccinia oahuensis** E. and E. Powdery yellow-brown rust pustules on leaves of *Panicum* sp. and *Syntherisma sanguinale* in Hawaii and Japan.
- Puccinia orientale** Syd. and Butl. Yellow-brown rust sori on leaves of *P. prostratum*, *P. trigonum*, and *Panicum* sp. in India and the Philippines.
- Puccinia panicophila** Speg. Brown rust sori on leaves of *P. insulare* in Argentina.
- Puccinia puttemansii** P. Henn. Leaf rust on *Panicum* sp. in Brazil.
- Septoria tandilensis** Speg. On leaves and sheaths of *P. nitidum* in Argentina.
- Sorosporium africanum** Syd. Black powdery smut masses destroying the ovaries of *P. trichops* in Portuguese East Africa.
- Sorosporium cryptum** McAlp. Smut sori concealed by the glumes and converting the ovaries into black spore masses. On *P. bicolor* and *P. effusum* in Australia.
- Sorosporium ovarium** Griff. Gray-brown smut sori in ovaries of *P. caespitosum* in Mexico, rupturing to expose black powdery spore masses.
- Sorosporium panici** MacKin. and var. **kinshasaensis** Beeli. Smut in ovaries of *P. flavidum*, *P. kinshasaensis*, and *Panicum* sp. in New South Wales and the Congo.
- Sphacelotheca chudaei** Har. and Pat. Long ovoid smut sori in inflorescences of *P. turgidum* in Morocco.
- Telimeia graminis** (v. Hoeh.) Theiss. and Syd. Circular to oblong black stromata on leaves of *Panicum* sp. in Samoa.
- Telimeia panici** Theiss. and Syd. Dull-black stromata on leaves of *P. patens* in the Philippines.
- Tilletia ayresii** Berk. Buff spore masses produced in the ovaries of *P. maximum* in Mauritius.
- Tilletia courtetiana** Har. and Pat. Smut sori consisting of olive-brown spore masses, deforming ovaries of *P. proliferum* in the Congo.
- Tilletia magnusiana** F. de W. Black smut masses in ovaries of *P. geniculatum* in the West Indies.
- Tilletia tumefaciens** Syd. Powdery smut sori deforming leaves and culms of *P. antidotale* in India.
- Tilletia verrucosa** Cke. and Mass. Smut in ovaries of *P. coloratum* and *P. miliare* in South Africa.
- Tolyposporium bogoriense** Racib. Smut sori in flowers of *Panicum* sp. in Java.
- Tolyposporium minus** Schroet. Ashen smut sori in ovaries of *Panicum* sp. in Brazil, rupturing to expose black spore masses.
- Uredo panici-montani** Petch. Leaf rust on *P. montanum* in Ceylon.
- Uredo panici-prostrati** Syd. Powdery brown rust pustules on leaves of *P. prostratum* in India.
- Uredo panici-villosi** Petch. Leaf rust on *P. villosum* in Ceylon.
- Uromyces leptodermus** Syd. Powdery brown rust pustules on leaves of *P. barbinode*, *P. isachne*, *P. javanicum*, and *P. prostratum* in Porto Rico, Cuba, Peru, Guatemala, and India. Also known from southern Florida.
- Uromyces linearis** B. and Br. Brown, then black, linear rust sori on leaves of *P. flavidum*, *P. miliare*, *P. proliferum*, and *P. repens* in India, Ceylon, China, Japan, and the Philippines.
- Uromyces puttemansii** Rangel. Leaf rust on *P. mellinum* in Brazil.
- Uromyces superfluus** Syd. Leaf rust on *P. antidotale* in India.
- Ustilaginoidea congensis** P. Henn. Globose to subtrubinate olivaceous sclerotia in ovaries of *P. brizanthum* and *P. maximum* in the Congo.
- Ustilaginoidea ochraceae** P. Henn. Sclerotia in ovaries of *P. auritum* in the Philippines.
- Ustilaginoidea usambarensis** P. Henn. Ashen-olivaceous sclerotia deforming the ovaries of *P. laxum* in Porto Rico and Tanganyika.
- Ustilago balansae** Speg. Powdery dark-brown smut masses in panicles of *Panicum* sp. in Argentina.
- Ustilago bonariensis** Speg. Powdery dark-olive smut sori covered by a whitish pseudomembrane in spikelets of *P. elephantipedis*, *P. spectabile*, and *Pennisetum tristachyon* in Argentina.
- Ustilago cacheutensis** Speg. Black powdery smut masses destroying the panicles of *P. leucophaeum* in Argentina.
- Ustilago confusa** Mass. Smut in ovaries of *P. paradoxum* in Australia.
- Ustilago digitariae** (Kze.) Rbh. Flowers, rachides, and culms destroyed and replaced by black smut sori on *P. glabrum*, *P. repens*, *P. teneriffae*, and *Syntherisma sanguinale* in Ceylon, India, Formosa, Portuguese East Africa, Sudan, Italy, and Germany.
- Ustilago formosana** K. Saw. Linear smut sori, up to 14 centimeters long, inclosed at first by a grayish-white membrane, which ruptures, exposing the black spore masses; in inflorescences and culms of *P. proliferum* in Formosa.
- Ustilago globigena** Speg. Smut sori in spikelets of *P. spectabile* in Argentina.
- Ustilago heterospora** P. Henn. Powdery olivaceous spore masses deforming the ovaries of *P. maximum*, *Setaria aurea*, and *S. glauca* in tropical Africa.
- Ustilago manilensis** Syd. Smut sori in the spikes of *P. indicum* in the Philippines.
- Ustilago negeriana** Diet. Black spore masses in ovaries of *P. urvilleanum* in Chile.
- Ustilago operta** Syd. and Butl. Smut in ovaries of *P. prostratum*, *P. ramosum*, and *P. villosum* in India.
- Ustilago panici-frumentacei** Bref. Black smut sori covered with a persistent gray membrane involving inflorescences, stems, nodes, and young shoots and causing abortion of fruiting heads. On *P. (Echinochloa) frumentaceum* in India.
- Ustilago panici-gracilis** MacKim. Smut in ovaries of *P. gracile* in New South Wales.
- Ustilago panici-latifolii** P. Henn. Black powdery smut sori in ovaries of *P. latifolium* in Brazil.
- Ustilago panici-leucophaei** Bref. Spikelets of *P. leucophaeum* in Brazil and Guatemala transformed into black powdery smut masses.
- Ustilago panici-petrosi** Syd. Dark-brown powdery smut sori destroying the inflorescences of *P. petrosum* in Brazil.
- Ustilago panici-proliferi** P. Henn. Smut sori in inner folded leaves, aborting the inflorescences and covered by thin transparent membranes which soon rupture, exposing dusty olive-black spore masses, on *P. paspaloides* and *P. proliferum* in Mexico.
- Ustilago paradoxa** Syd. and Butl. Smut sori attacking the individual ovaries of *P. frumentaceum* in India. The sori are round, gray, and covered by persistent membranes, which are broken only by handling, exposing the black spore masses.
- Ustilago pretoriensis** Pole-Evans. Long powdery dark olivaceous smut sori destroying the ovaries of *P. helopodis* in the Union of South Africa.
- Ustilago rickerii** Clint. Olive-black dusty smut masses in ovaries of *P. paspaloides* in Cuba.
- Ustilago trichophora** (Link.) Kze. Smut sori in ovaries of *P. colonum* and *Tricholaena (Panicum) teneriffae* in India and north Africa.
- Ustilago ugandensis** P. Henn. and var. **macrospora** Beeli. Smut sori consisting of olivaceous powdery spore masses at apex of culms, in rachides and flowers of *Panicum* sp. in tropical Africa.
- Ustilago vesiculosa** P. Henn. Smut sori in panicles of *Panicum* sp. in Brazil.

- PAPAVER.** POPPY. Biennial or perennial herbs with milky juice. Economic and flower-garden plants. Oil produced from the seed, opium and its derivatives from the juice.
- Ascochyta papaveris** Oud. On leaves of *P. nudicaule* in Nowaja Semla.
- Entyloma fuscum** Schroet. See Glaucium.
- Gloeosporium** sp. Anthracnose on leaves of *P. somniferum* in Japan.
- Helminthosporium papaveri** K. Saw. Large irregular brown leaf spots on *P. somniferum* in Formosa. Lesions also occur on the stems and circular yellow to brown spots on the capsules. Infected plants are often destroyed. "The most dangerous disease of the cultivated poppy in Formosa."
- Helminthosporium papaveris** P. Henn. On *Papaver* sp. in Tanganyika.
- Macrosporium papaveris** Parisi. On green capsules and leaves of *P. somniferum* in Italy.
- Peronospora arborescens** (Berk.) De By. A downy mildew forming pale-brown leaf spots, more especially near the tips and margins. These spots dry up and fall out and entire leaves may wither. On the lower surfaces a gray-violet fungus layer forms. On *P. dubium*, *P. rhoeas*, *P. somniferum* and *Meconopsis cambrica* in Japan, India, Siberia, Algeria, Egypt, and Europe.
- Peronospora argemones** Gäum. Downy mildew on leaves of *P. argemone* in Scotland, Scandinavia, Austria, and Germany.
- Peronospora cristata** Tranz. As above on *P. hybridum* in Asia Minor.
- Septoria rhoeadis** F. Tassi. Violet-brown circular and confluent leaf spots on *P. rhoeas* in Italy.
- PAPHIOPEDILUM.** Terrestrial or epiphytic orchids. See Orchidaceae.
- PAPYRIUS.** See Broussonetia.
- PARAMIGNYA.** Evergreen climbing shrubs.
- Accidium paramignyae** Racib. Rust on circular leaf spots on *P. blumei* and *P. monophylla* in Java, Ceylon, and India.
- Bacterium citri** Hasse. See Citrus.
- Helminthosporium makilingense** Syd. On branches of *P. monophylla* in the Philippines.
- PARIS.** Small, hardy, rhizomatous plants.
- Cercospora paridis** Erikss. Dark-brown leaf spots on *P. quadrifolia* in Russia, Sweden, Switzerland, Bohemia, Austria, and Denmark.
- Septoria paridis** Passer. Small irregular whitish leaf spots with dull-brown margins on *P. quadrifolia* in Russia and Italy.
- Tubercinia paridis** (Ung.) Vestergr. Smut on leaves and stems of *P. quadrifolia* in Russia, Sweden, Germany, and Austria.
- Urocystis colchici** (Schlecht.) Rab. See Colchicum.
- PARKIA.** CUPANG. Tall, unarmed leguminous trees.
- Endodothella parkiae** (P. Henn.) Theiss. and Syd. Black stromata on brown leaf spots on *Parkia* sp. in Brazil.
- Leptothyrella manaosensis** P. Henn. Black fruiting bodies on leaves of *P. auriculata* in Brazil.
- Phyllachora parkiae** P. Henn. Circular shiny black stromata on indefinite yellowish leaf spots on *P. roxburghi* and *P. timoriana* in the Philippines.
- PARNASSIA.** Sometimes called grass of Parnassus. Low growing, hardy perennial herbs.
- Puccinia uliginosa** Juel. Leaf rust on *P. palustris* and *Carex vulgaris* in Europe.
- PAROSELA.** See Dalea.
- PARRYA.** Low perennial herbs with thick caudices and scapelike peduncles.
- Puccinia oudemansii** Tranzsch. See Matthiola.
- PARSONSIA.** HELYGIA Ag. Twining apocynaceous shrubs.
- Accidium parsonsi** Petch. Leaf rust on *P. spiralis* in Ceylon.
- PARSONSIA.** See Cuphea.
- PARTHENIUM.** Annual or perennial herbs or shrubs, mostly heavily pubescent.
- Entyloma calendulae** (Oud.) De By. See Calendula.
- Uredo parthenii** Speg. Leaf rust on *P. hysterophorus* in Argentina.
- PASANIA.** LITHOCAEPUS Ag. Ornamental trees grown for their handsome foliage.
- Taphrina kusanoi** Ikeno. Fruiting areas on lower leaf surfaces of *P. cuspidata* in Japan.
- PASCALIA.** See Wedelia.
- PASPALUM.** Perennial grasses, including valuable forage grasses as well as weeds.
- Balanisia gigas** Racib. Globose yellow-brown stromata, white within, on *Paspalum* sp. in Java.
- Balanisia paspali** P. Henn. Black rugulose stromata in spikes of *Paspalum* sp. in Oceania and the Philippines.
- Cerebella inquinans** (B. and Br.) Petch. See Panicum.
- Claviceps deliquescens** (Speg.) Haum. Sclerotia in ovaries of *P. dilatatum*, *P. distichum*, *P. hieronymi*, *P. larranagai*, *P. plicatulum*, and *P. quadrifarium* in Argentina. Probably not distinct from *C. paspali*, which is widespread in the United States.
- Claviceps lutea** A. Möll. Yellow oblong curved sclerotia in spikelets of *Paspalum* sp. in Brazil.
- Endodothella platensis** (Speg.) Theiss. and Syd. Elliptic to linear black stromata on leaves of *Paspalum* sp. in Argentina.
- Ephelis japonica** P. Henn. Inflorescences deformed by blackish stromata which form between the compressed spikelets of *P. scrobiculatum*, *P. thunbergium*, and *Miscanthus* sp. in Japan.
- Helminthosporium mayaguezense** Miles. Oval yellow spots with narrow dark-brown margins on leaves and culms of *P. conjugatum* in Porto Rico.
- Phyllachora acuminata** Starb. Tar-spot on leaves of *Paspalum* sp. in Argentina.
- Phyllachora infuscans** Wint. Irregular linear black stromata on *Paspalum* sp. in Brazil.
- Phyllachora paspalicola** P. Henn. Dull-black stromata on leaves of *P. compressum* and *P. conjugatum* in Brazil and Colombia.
- Phyllachora winkleri** Syd. Tar spot on leaves of *P. scrobiculatum* in Ceylon and Tanganyika.
- Puccinia huberi** P. Henn. See Panicum.
- Puccinia macra** Arth. and Holw. Leaf rust on *P. candidum* in Guatemala.
- Puccinia maublancii** Rangel. Leaf rust on *P. densum* in Brazil.
- Puccinia pilgeriana** P. Henn. Dark-brown rust sori on leaves of *Paspalum* sp. in Brazil.
- Puccinia tubulosum** (Pat. and Gaill.) Arth. Aecia in circular yellow areas 1 to 2 centimeters in diameter on leaves of *Solanum tequilense* and *S. torvum* in Mexico, Porto Rico, Cuba, Jamaica, and Central America. Brown powdery uredo and telial sori on leaves of *P. compressum*, *P. conjugatum*, *P. humboldtianum*, *P. paniculatum*, *P. plicatulum*, *Syntherisma digitata*, *S. sanguinale*, and *Valota insularis* in Mexico, Central America, the West Indies, and Trinidad. Also known from Texas.
- Sorosporium paspali** McAlp. Smut sori in inflorescences of *P. scrobiculatum*, destroying them and leaving the elongated remains surrounded by a black spore mass. Occurs in Ceylon, India, Australia, and the Philippines.
- Telimeia graminella** Syd. Opaque black stromata on leaves of *Paspalum* sp. in the Philippines.
- Thecaphora inquinans** B. and Br. Black smut sori in seed of *P. scrobiculatum* in Ceylon.
- Tilletia ulei** Schroet. Black smut sori destroying the ovaries of *P. scrobiculatum* in Brazil.
- Uredo paspali-longiflorae** Petch. Leaf rust on *P. longiflorum* in Ceylon.
- Uredo paspalina** Syd. Brown leaf rust on *P. scrobiculatum* in the Philippines.

**PASPALUM**—Continued.

- Uredo paspali-perrottetii** Petch. Leaf rust on *P. perrottetii* in Ceylon.  
**Uredo paspali-scribiculari** Syd. Brown rust sori on upper leaf surfaces of *P. scribicularum* in India and Ceylon.  
**Ustilago inoidea ochracea** P. Henn. On *Paspalum* sp. in Polynesia.  
**Ustilago microspora** Schroet. and P. Henn. Smut sori reducing panicles of *P. dilatatum* and *Paspalum* sp. in Brazil and Uruguay to olivaceous powdery spore masses.  
**Ustilago paspali** Speg. Black powdery spore masses destroying ovaries and glumes of *Paspalum* sp. in Uruguay and Argentina.  
**Ustilago paspali-dilatati** P. Henn. Black smut masses replacing inflorescences of *P. dilatatum* in Brazil.  
**Ustilago paspali-thunbergii** P. Henn. Black powdery smut masses in the inflorescences of *P. thunbergium* in Japan.  
**Ustilago rabenhorstiana** J. Kuehn. Smut sori in inflorescences of *P. (Syntherisma) sanguinale* in India.  
**Ustilago royleani** Syd. and Butl. Black smut sori in spikes of *P. royleanum* in India.  
**Ustilago schroeteriana** P. Henn. Black powdery smut sori in spikes of *Paspalum* sp. in Brazil.  
**Ustilago subnitens** Schroet. and P. Henn. Smut on *Paspalum* in Brazil.  
**Ustilago venezuelana** Syd. Smut on *Paspalum* sp. in Brazil.  
**Ustilago verrucosa** Schroet. Black smut in ovaries of *P. distichum* in Brazil.

**PASSERINA**. Heathlike shrubs.

**Marsonia andurnensis** (Ces.) P. Magn. On stems of *P. annua* in Europe.

**PASSIFLORA**. PASSION FLOWER. Herbs, shrubs, and trees, many species climbers cultivated for fruit and as ornamentals.

- Aecidium passiflorae** P. Henn. Leaf rust on *Passiflora* sp. in Tanganyika.  
**Ascochyta passiflorae** Penz. and Sacc. On peduncles of *P. hybrida* and *P. incarnata* in Italy.  
**Asterina megalospora** B. and C. Black superficial fungus patches on *Passiflora* spp. in Porto Rico, Cuba, Ecuador, Peru and, Brazil.  
**Colletotrichum passiflorae** Siem. Anthracnose on *P. edulis* in Russia.  
**Didymella passiflorae** v. Hoeh. On leaves of *Passiflora* sp. in Samoa.  
**Didymosphaeria innumerabilis** Wint. Circular to irregular whitish leaf spots on *Passiflora* sp. in Brazil.

**Helminthosporium stahlii** Stevens. Small irregular to large and diffuse brown leaf spots on *P. foetida* in Porto Rico.

**Melampsora passiflorae** Har. Rust on leaves of *P. lutea* in France.

**Mycosphaerella passiflorae** Rehm. Leaf spot on *Passiflora* sp. in South America.

**Ovulariopsis** sp. On leaves of *Passiflora* sp. in India.

**Phyllosticta passiflorae** McAlp. Large irregular leaf spots on *P. edulis* in Australia.

**Phyllosticta superficiale** Stevens. Irregular, brown leaf spots on *P. sexflora* in Porto Rico.

**Puccinia scleriae** (Paz.) Arth. The aecial stage of this rust occurs on circular, yellow-brown leaf spots on *P. rubra* and *P. tuberosa* in Porto Rico, Cuba, Trinidad, Jamaica, and Peru.

**Uromyces appelianus** Gassn. Leaf rust on *P. foetida* in Brazil and Uruguay.

**PASTINACA**. PARSNIP. Fleshy edible roots.

**Cercospora pastinacae** Karst. Small circular brown spots with white centers on leaf blades and petioles of *P. sativa* and *Daucus carota* in Great Britain, Finland, Russia, and Germany. Reported from Indiana.

**Cylindrosporium pimpinellae** C. Mass. var. **pastinacae** Sacc. Small dry spots on leaves of *P. sativa* and *P. urens* in Bulgaria, Italy, and Germany.

**Entyloma pastinacae** Jaap. Smut sori in grayish-white leaf spots on *P. sativa* in Dalmatia.

**Gloeosporium achaenicola** Rostr. See Petroselinum.

**Phyllachora pastinacae** Rostr. Indefinite, finally confluent, pale-green, then brown, leaf spots on which stromata occur on *P. sativa* in France, Belgium, Sweden, Denmark, Russia, and Italy.

**Protomyces macrosporus** Ung. See Coriandrum.

**Ramularia pastinacae** Bub. Circular to irregular dark grayish-green, then brown, leaf spots with white centers on *P. sativa* and *Daucus carota* in Yugo-slavia, Dalmatia, Great Britain, Finland, and Austria. Reported from North Dakota.

**PATRINIA**. FEDIA. Ag. Hardy herbaceous perennials with yellow or white Valerian-like flowers.

**Aecidium patriniae** P. Henn. Leaf rust on *P. scabiosaefolia* and *P. villosa* in Japan.

**Puccinia melanoplaca** Syd. Black rust sori on leaves of *P. palmata* in Japan.

**Puccinia patriniae** P. Henn. Chestnut-brown rust sori on leaves of *P. villosa* in Japan.

**Puccinia patriniae-gibbosae** Miura. Leaf rust on *P. gibbosa* in Japan.

**PAULLINIA**. Twining tropical shrubs.

**Phaeoapiospora nectrioides** (Rehm.) Sacc. and Syd. Rust-brown galls on leaves of *Paullinia* sp. in Brazil.

**PAULOWNIA**. Ornamental trees grown for their showy flower panicles and large leaves.

**Gloeosporium kawakamii** Miy. Causes witches'-brooms on *P. tomentosa* (*P. imperialis*) in Japan. The leaves on these brooms remain small, pale green in color, and curled, with red-brown spots on blades and petioles. Seedlings often die.

**Helicobasidium tanakae** Miy. See Morus.

**Mycosphaerella paulowniae** Shir. and Hara. Leaf spots on *P. tomentosa* in Japan.

**Phyllosticta paulowniae** Sacc. Sinuous and confluent ochraceous leaf spots on *P. tomentosa* in France, Italy, and Russia.

**Septoria paulowniae** Thuem. More or less circular gray leaf spots on *P. tomentosa* in Italy and France.

**Valsa paulowniae** Miy. and Hemmi. Attacks the twigs of *Paulownia* sp. and *P. tomentosa* in Japan, spreading into the branches and finally destroying the entire tree. Considered serious, being compared with chestnut blight in virulence.

**PAVETTA**. Tropical shrubs and small trees allied to Ixora.

**Aecidium pavettae** Berk. Leaf rust on *P. hispidula* and *P. indica* in Ceylon, India, and Uganda.

**Aecidium transvaaliae** P. Henn. and Evans. Leaf rust on *Pavetta* sp. in the Union of South Africa.

**Cocconia placenta** (B. and Br.) Sacc. See Symplacos.

**Endophyllum griffithsiae** Rac. See Randia.

**Hemileia mildbraedii** Syd. Powdery yellow rust pustules on leaves of *P. oliveriana* in central Africa.

**Phoma pavettae** F. Tassi. On branches of *P. indica* in Italy.

**Woodiella natalensis** Sacc. and Syd. Small leaf spots on *P. obovata* in Natal.

**PAVONIA**. Malvaceous herbs or shrubs.

**Caecoma pavoniae** Diet. Golden rust sori on leaves of *Pavonia* sp. in Brazil.

**Phyllachora pusilla** Syd. Tar spot on leaves of *P. schrankii* in Brazil.

**Puccinia exilis** Syd. Yellow-brown rust pustules on leaves of *P. leucantha* and *P. rosea* in Guatemala and Brazil.

**PAVONIA**—Continued.

- Puccinia pavoniae** P. Henn. Ochraceous-brown rust pustules on leaves of *Pavonia* sp. in Tanganyika.
- Puccinosira pallidula** (Speg.) Lagh. Yellow rust sori on leaves of *P. paniculata* and *Triumfetta* spp. in Central and South America and the West Indies.
- Ramularia aguierei** Speg. Indefinite areas on leaves of *P. sepium* in Argentina.
- Kostrupia praelonga** Speg. Linear yellow-brown rust pustules on leaves and branches of *P. polymorpha* in Argentina.
- Uromyces pavoniae** Arth. Leaf rust on *P. racemosa* in Porto Rico.
- PEDICULARIS.** WOOD BETONY. Lousewort. Mostly perennial herbs with finely cut foliage.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- Melampsora pedicularis** Vogl. Leaf rust on *P. verticillata* in Switzerland.
- Peronospora pedicularis** Palm. Downy mildew on leaves of *P. lapponica* in Sweden and Norway.
- Plasmopara densa** (Rabh.) Schroet. Downy mildew on leaves of *P. palustris*, *P. silvatica*, *Alecatorolophus alpinus*, *Bar. sia alpina*, *Euphrasia officinalis*, *E. pratensis*, and *Odontites rubra* in Europe.
- Puccinia oederi** Blytt. Brown rust sori on leaves of *P. oederi* in Norway.
- Puccinia paludosa** Plowr. The yellow aecia of this rust occur on *P. mexicana* and *P. palustris*, the other stages on *Carex* spp. in Europe.
- Puccinia pedicularis** Thuem. Leaf rust on *Pedicularis* sp. in Siberia.
- Ramularia filiformis** Lindr. Leaf spots on *P. silvatica* in Sweden.
- Ramularia obducens** Thuem. On leaves of *P. foliosa* and *P. palustris* in Austria and Switzerland. Reported from California.
- Rhytisma pedicularis** (DC.) Rehm. Black confluent stromata on leaves of *P. incarnata* and *P. alpina* in Italy.
- Septoria pedicularium** Fr. On stems of *Pedicularis* sp. in Europe.
- PELARGONIUM.** GERANIUM. Herbaceous plants used as potted plants for indoor planting and for bedding.
- Cercospora** sp. On leaves of *Pelargonium* sp. in China.
- Coniothyrium trabuti** Riza. Indefinite yellow patches on leaves of *P. pellatum* in Morocco.
- Fusarium pelargonii** Crouan. Causes a black rot of stems, the leaves turning yellow, on *Pelargonium* sp. in Great Britain, France, and Denmark.
- Gloeosporium pelargonii** Cke. and Mass. Anthracnose, causing leaves of *Pelargonium* sp. in Great Britain to wither and droop.
- Leptosphaeria pelargonii** Rehm. Irregular whitish spots on leaves of cultivated *Pelargonium* in Brazil.
- Macrosporium macalpineanum** Sacc. and Syd. Large irregular and confluent yellow-brown leaf spots with purple margins on *P. zonale* in Australia and Ceylon.
- Puccinia granularis** Kalch. and Cke. Brown rust sori in circles on lower leaf surfaces of *P. aconitophyllum*, *P. alchemilloides*, and other species, as well as *Begonia* sp., in the Union of South Africa. Serious in greenhouses.
- Puccinia morrisoni** McAlp. Orange and brown to black rust pustules on leaves of *P. australe* in Australia.
- Puccinia pelargonii** (Thuem.) Syd. Yellow and brown rust sori on leaves of *P. alchemilloides* in the Union of South Africa.
- Septoria geranii** Rob. and Desm. See Geranium.
- Septoria pelargonii** Syd. Circular to irregular yellow-brown to red-brown leaf spots on *P. cucullatum* in the Union of South Africa.
- PELLAEA.** CLIFF BRAKE. Small rock-loving ferns.
- Uredo pellaeae** Diet. and Neg. Rust on leaves of *P. ternifolia* in Chile.
- PELTOPHORUM.** Tropical leguminous trees.
- Phyllachora peltophori** Syd. Shiny black stromata on leaves of *P. africanum* in the Union of South Africa.
- PENNISETUM.** PEARL MILLET. NAPIER GRASS. Stout grasses with bristly spike-like inflorescences.
- Acrothecium penniseti** Mitra. The fungus attacks the leaves, leafsheaths, and ears of *P. typhoideum* in India. The leaf spots are small, yellow brown, and oval to oblong. The lower leaves are destroyed and infected heads fail to set grain.
- Balansia claviceps** Speg. Black stromata in spikes of *Pennisetum* sp. and *Setaria* sp. in Argentina.
- Beniowskia penniseti** Wakef. Small pale-red leaf spots on *P. purpureum* and *Holcus sorghum* in Uganda.
- Cerebella cenchroidis** Subra. Black corrugated fungus masses in ovaries of *P. cenchroides* in India.
- Contractia columellifera** (Tul.) McAlp. See Andropogon.
- Didymella penniseti** Syd. On leaves of *P. tristachyon* in Columbia.
- Leptostroma penniseti** P. Henn. Effuse dull-brown leaf spots on *P. japonicum* in Japan.
- Neovossia barclayana** Bref. Smut sori in ovaries of *P. triflorum* in India.
- Phyllachora minutissima** (Welw. and Curr.) Sm. Small elliptical stromata on leaves of *Pennisetum* sp. in Angola.
- Phyllachora penniseti** Syd. Black stromata on leaves of *P. benthami* and *P. sphacelatum* in Congo and the Union of South Africa.
- Phyllachora pennisetina** Syd. Tar spot on leaves of *P. alopecuroides* in China.
- Puccinia apoda** Har. and Pat. Brown to black rust sori on rufous leaf spots on *P. setosum* in the Congo.
- Puccinia arthuri** Syd. Leaf rust on *P. crinitum* and *P. mexicanum* in Mexico.
- Puccinia burmeisteri** Speg. Leaf rust on *P. tristachyon* in Argentina.
- Puccinia penniseti** Zimm. Brownish-yellow to black rust sori on brown leaf spots on *P. spicatum* and *P. typhoideum* in India and east Africa.
- Tolyposporium pencillariae** Bref. Smut sori bright green to black, projecting beyond the glumes and finally breaking to expose deep-brown spore masses. Attacking individual grains or small groups of grain of *P. spicatum* and *P. typhoideum* in India, Egypt, and other parts of Africa.
- Ustilago inoidea penniseti** Miy. Sclerotia in ovaries of *P. compressum* in China.
- Ustilago bonariensis** Speg. See Panicum.
- Ustilago kamerunensis** Syd. Smut sori destroying inflorescences of *Pennisetum* sp. in Congo.
- Ustilago pappiana** Bacc. Black smut sori in ovaries of *P. orientalis* and *P. ruppeli* in Abyssinia.
- Ustilago penniseti** Rabenh. Black smut sori in ovaries of *P. asperifolium*, *P. cenchroides*, *P. dichotomum*, *P. fasciculatum*, *P. ruppeli*, *P. vulpinum*, and *P. villosum* in Madeira, China, Arabia, Abyssinia, Egypt, Algeria, and Tripoli.
- Ustilago penniseti-japonici** P. Henn. Smut sori destroying flowers of *P. compressum*, *P. japonicum*, and *P. purpurascens* in China and Japan.
- Ustilago scheffleri** Syd. Smut sori up to 3 centimeters long, covered by ashen membranes which rupture, exposing black powdery spore masses in inflorescences of *P. inclusum* in tropical Africa.

**PENTAS.** Tender herbs and subshrubs.

*Puccinia pentadicola* Grove. Leaf rust on *P. verticillata* in Uganda.

*Puccinia pentadis* P. Henn. Brown rust sori on leaves of *P. mombassana* in tropical Africa.

*Puccinia pentadis-carneae* Wakef. Leaf rust on *P. carnea* in Uganda.

**PENTSTEMON.** Sometimes called beardtongue. Tubular-flowered bedding and border plants.

*Phyllosticta pentastemonis* Cke. and f. *pentastemonis-azurei* Allesch. Irregular brown leaf spots on *P. azureus*, *P. grandiflorus*, and *P. hybridus* in Great Britain and Germany.

*Puccinia mexicana* Diet. and Holw. Brown rust pustules on sunken, circular leaf spots on *P. campanulatus* in Mexico.

*Puccinia pentastemonum* Lév. Brown rust sori on leaves of *Pentstemon* sp. in France.

**PEPEROMIA.** Tropical and subtropical herbs, a few species cultivated as foliage plants.

*Albugo tropica* Lagh. White rustlike pustules on leaves of *P. pellucida* in the West Indies and Ecuador.

*Uredo peperomiae* P. Henn. Irregular, dull yellow rust sori on leaves of *Peperomia* sp. in Brazil.

*Uredo piperis* P. Henn. See Piper.

**PERDICIUM.** See Gerbera.**PERESKIA.** Shrubby or clambering vines with edible fruits.

*Uromyces peireskiae* Diet. Light-brown rust pustules on *Pereskia* sp. in Brazil.

**PERILLA.** Herbs, some species with colored foliage.

*Coleosporium perillae* Syd. Golden-yellow rust pustules on leaves of *P. arguta*, *P. crispa*, *P. nankinensis*, *P. ocimoides*, *Elscholtzia cristata*, *Keiskea japonica*, *Mosla formosana*, and *M. punctata* in Japan, Formosa, Manchuria, and India.

**PERIPLOCA.** SILK VINE. Ornamental vines grown for their handsome glossy foliage.

*Ascochyta periplocae* Kab. and Bub. Large round to irregular dark-brown, then ashen, leaf spots with brown-purple margins on *P. graeca* in Bohemia and Hungary.

*Melampsora periplocae* Miy. Leaf rust on *Periploca* sp. in China.

*Phyllosticta periplocae* F. Tassi. Circular, ashen leaf spots on *P. graeca* in Italy.

*Septoria periplocae* Höll. Subcircular, brown leaf spots on *P. graeca* in Hungary.

*Uredo periplocae* P. Henn. Ochraceous rust sori on irregular yellow leaf spots on *P. linearis* in tropical Africa.

**PERISTROPHE.** Greenhouse plants grown for their flowers.

*Accidium peristrophes* Syd. Leaf rust on *Peristrophe* sp. in India.

*Synchytrium rytzii* Syd. Small leaf galls on *Peristrophe* sp. in India.

**PERNETTYA.** Plants grown for their attractive berries, evergreen foliage, and white or pinkish flowers.

*Hypocrella pernettyae* Pat. On leaves of *P. repens* in Java.

*Stereum purpureum* Pers. See Prunus.

**PERSEA.** Avocado. Sometimes called alligator pear. Tropical and subtropical fruit trees.

*Cercospora (?) lingue* Speg. On leaves of *P. lingue* in Chile.

*Endothia havanensis* Bruner. See Eucalyptus.

*Fomes lamaecensis* Murr. See Hevea.

*Mycosphaerella perseae* Miles. Large irregular leaf spots, ashen above, brown beneath on *P. americana* in Porto Rico.

*Phyllachora gratissima* Rehm. Black stromata on irregular red-brown leaf spots on *P. americana* in Porto Rico, Costa Rica, and Ecuador.

*Physalospora perseae* Doidge. Produces a canker disease of *P. americana* in the Union of South Africa. Sunken discolored areas appear, gradually girdling twigs and branches. The dead bark dries out, cracks, and finally falls away. Brown, deeply cracking corky growths on the lower ends of the fruit are attributed to the same fungus.

*Pionnotes capillacea* Sacc. Causes a die back of *P. americana* in the Philippines.

*Rhytisma perseae* Gandara. Black stromata on large, irregular brown leaf spots on *P. americana* in Mexico. Probably not distinct from *Phyllachora gratissima* Rehm.

*Rosellinia* spp. (*R. pepo* Pat.). See Citrus.

*Eucalyptus* disease. See Eucalyptus.

**PERTYA.** Oriental shrubs.

*Accidium pertyae* P. Henn. Rust pustules on circular dull-brown leaf spots on *P. ovata* and *P. scandens* in Japan.

**PETASITES.** BUTTERBUR. Hardy perennial herbs.

*Accidium petasitidis* Syd. Leaf rust on *P. albus*, *P. japonicus*, *P. nivus*, *P. officinalis*, and *P. tomentosus* in Japan and central Europe.

*Coleosporium petasites* Lév. See Pinus.

*Ramularia cervina* Speg. and var. *petasitis* Bauml. On leaves of *P. albus* in Hungary.

*Venturia petasitidis* (Fckl.) Sacc. On leaves of *P. officinalis* in Europe.

**PETIVERIA.** Shrubby herbs.

*Rosellinia bunodes* B. and Br. See Citrus.

**PETROSELINUM.** PARSLEY. Biennial herbs cultivated for ornament and edible foliage.

*Bacillus nelliae* Welles. Bacterial wilt of *P. sativum* (*P. hortense*) in the Philippines.

*Bacillus petroselini* Pot. Small yellow leaf spots with brown margins on *P. sativum* in Russia.

*Fusicladium depressum* (B. and Br.) Sacc. and var. *petroselini* Sacc. On leaves of *P. sativum* in France and Spain.

*Gloeosporium achaenicola* Rostr. Ashen-colored areas on fruit and peduncles of *P. sativum* and *Pastinaca sativa* in Denmark.

*Macrosporium cheiranthi* (Lib.) Fr. Circular, gray leaf spots on *P. sativum* in France.

*Puccinia petroselini* (DC.) Lindr. Cinnamon to dark-brown rust pustules on leaf blades and petioles and on stems of *P. sativum*, *Aethusa cynapioides*, and *Anethum graveolens* in Argentina and Europe.

*Septoria petroselini* Desm. and var. *segetum* P. Brun. Leaf spots on *P. hortense*, *P. sativum*, and *P. segetum* in Australia, Barbados, Argentina, Union of South Africa, and Europe.

**PETUNIA.** Small herbs cultivated for their showy flowers.

*Entyloma petuniae* Speg. Smut sori in pale indefinite spots, often occupying entire area of leaf blades of *P. nyctaginiflora* in Argentina.

*Phyllosticta petuniae* Speg. Circular, then angular, dull-brown leaf spots on *Petunia* sp. in Great Britain.

*Ramularia petuniae* Cke. Large, somewhat circular, dull-yellow leaf spots on *Petunia* sp. in Great Britain.

**PHAJUS.** Epiphytic and terrestrial orchids. See Orchidaceae.**PHALAENOPSIS.** See Orchidaceae.

- PHALARIS.** REED CANARY GRASS. Stout, hardy, ornamental grasses.  
**Entyloma brefeldi** Krieger. Smut sori in leaf spots on *P. arundinacea* in Germany.  
**Heterosporium phragmites** (Opiz.) Sacc. and var. **inflorescentiae** Bub. On leaves of *P. arundinacea* and *Phragmites communis* in France and Bohemia.  
**Leptosphaeria culmicola** (Fr.) Karst. See Bromus.  
**Mycosphaerella chlorina** (Cke.) Lind. On leaves of *P. arundinacea* in Great Britain.  
**Ophiobolus cariceti** (B. and Br.) Sacc. See Triticum.  
**Phyllosticta crastophila** Sacc. Dull-yellow areas on leaves of *P. arundinacea* and *Setaria verticillata* in Italy.  
**Physalospora phalaridis** Zem. On *P. bulbosa* in Argentina.  
**Physoderma gerhardtii** Schroet. See Glyceria.  
**Puccinia brevicornis** Ito. Leaf rust on *P. arundinacea* in Japan.  
**Puccinia orchidearum-phalaridis** Kleb. See Orchidaceae.  
**Puccinia schmidiana** Diet. Yellow and powdery brown to black rust pustules on leaves of *P. arundinacea*, *Leucojum aestivum*, and *L. vernum* in Italy, Austria, and Germany.  
**Puccinia winteriana** P. Magn. Yellow and brown to black rust sori on leaves of *P. arundinacea* and *Allium ursinum* in Spain, France, Switzerland, Great Britain, Denmark, Holland, and Germany.  
**Sclerospora kriegeriana** Magn. Downy mildew on leaves of *P. arundinacea* in Germany.  
**Sclerospora macrospora** Sacc. See Triticum.  
**Septoria phalaridis** Cocc. and Mor. Pale-yellow spots on leaves and sheaths of *P. brachystachys* in Italy.  
**Tilletia menieri** Har. and Pat. Smut sori turning the ovaries of *P. arundinacea* into brown spore masses. Reported from France.
- PHASEOLUS.** BEAN. Lima-bean. Herbs cultivated as vegetables and forage plants.  
**Acanthostigma heterochaeta** Syd. and Butl. Brown or yellow-brown leaf spots on *Dumassia villosa*, *Dumassia* sp., *P. mungo*, and *P. radiatus* in India.  
**Ascochyta boltshauseri** Sacc. Brown spots concentrically zoned and causing premature death of the leaves of *P. lunatus*, *P. vulgaris*, and *Vicia faba* in New Zealand, Russia, Bohemia, Switzerland, Holland, Norway, and Denmark.  
**Ascochyta hormullerii** Syd. On leaves of *P. acutifolius* in Mexico.  
**Ascochyta phaseolorum** Sacc. Indefinite ochraceous leaf spots on *P. mungo*, *P. multiflorus*, *P. nanus*, *P. radiatus*, *P. vulgaris*, *Vigna catjang*, and *Vicia faba* in Japan, Uganda, and Europe.  
**Asteroma phaseoli** Brun. Felty-black spots on pods of *P. vulgaris* in the Philippines.  
**Cercospora lussoniense** Sacc. Leaf spot on *P. lunatus* in the Philippines.  
**Cercospora stublmanni** P. Henn. Irregular dull-brown leaf spots on *P. vulgaris* in Tanganyika.  
**Cercospora zonata** Wint. See Vicia.  
**Cercosporina caracallae** Speg. Angular ashen leaf spots on *P. caracalla* in Argentina.  
**Cylindrosporium phaseoli** Rabh. On leaves of *Phaseolus* sp. in Germany.  
**Diplodia phaseolina** Sacc. Small black spots on ripening pods of *P. lunatus* and *P. vulgaris* in the Philippines.  
**Hypochnus cucumeris** Frank. See Cucumis.  
**Leptosphaeria circinans** (Fckl.) Sacc. See Asparagus.  
**Macrophoma phaseolina** F. Tassi. Irregular pale-brown leaf spots on *P. ornithopus* in Italy.  
**Mosaic.** A mosaic disease, said to resemble a similar disease on soy (*Soja*), is reported from Japan and may be the same as the mosaic of legumes occurring in the United States.  
**Mycosphaerella phaseolicola** Desm. Pale-reddish leaf spots on *P. vulgaris* in Japan and France.  
**Mycosphaerella phaseolorum** Siem. On leaves of *P. mungo* and *Soja max* in Russia.  
**Phyllachora phaseoli** (P. Henn.) Theiss. and Syd. Shiny black stromata on leaves of *P. adenanthus* and *Phaseolus* sp. in Porto Rico and Brazil.  
**Phyllachora phaseolina** Syd. Circular black stromata on rust-brown leaf spots on *P. calcaratus* and *Vigna* sp. in the Philippines and Amboina.  
**Phyllosticta noackiana** Allesch. Subcircular ochraceous leaf spots on *Phaseolus* sp. in Brazil.  
**Phyllosticta phaseolorum** Sacc. and Speg. See Vigna.  
**Phytopella concors** Arth. Brown powdery rust pustules on leaves of *P. lunatus*, *Dolichos lablab*, and *Teramnus uncinatus* in Porto Rico.  
**Pleosphaerulina phaseoli** Syd. On leaves of *P. semirectus* in the Philippines.  
**Rhizoctonia** sp. See Vigna.  
**Rhizoctonia dimorpha** Matz. Attacks leaves and stems of *Phaseolus* sp. and *Vigna catjang* in Porto Rico.  
**Rhizoctonia macrosclerotia** Matz. Reported on petioles and stems of *Phaseolus* sp. in Porto Rico.  
**Sclerophoma phaseoli** Karak. On pods of *P. multiflorus* in Russia.  
**Septoria phaseoli** Maubl. Irregular greenish leaf spots on *P. mungo* and *Phaseolus* sp. in Russia and Brazil.  
**Uredo vignae** Bres. See Vigna.  
**Uromyces phaseolicola** Speg. Leaf rust on *P. prostratus* in Argentina.
- PHEGOPTERIS.** WOOD FERN.  
**Exoascus filicinus** (Rostr.) Sacc. See Polystichum.  
**Uredinopsis filicina** P. Magn. Yellow-brown rust pustules on leaves of *P. polypodioides* (*Dryopteris polypodioides*) in Europe.
- PHELLODENDRON.** CORK TREE. Ornamental trees.  
**Ascochyta phellodendri** Kab. and Bub. Round to irregular brown leaf spots with white centers on *P. amurense* in Bohemia.  
**Coleosporium phellodendri** Kom. Fg. Golden rust sori on leaves of *P. amurense* in Chosen, Manchuria, and Japan.
- PHILADELPHUS.** MOCK ORANGE. Syringa. Ornamental shrubs.  
**Aecidium philadelphia** Diet. Leaf rust on *P. coronarius* in Japan.  
**Ascochyta fuscescens** Kab. and Bub. Indefinite dull-brown to ashen areas on leaves of *P. coronarius* in Bohemia.  
**Ascochyta philadelphia** Sacc. and Speg. Subcircular ochraceous leaf spots on *P. coronarius* in Europe.  
**Phragmidium philadelphia** Pass. Leaf rust on *P. coronarius* in Yugoslavia.  
**Phyllosticta coronaria** Pass. Large brown leaf spots on *P. coronarius* in Russia, Italy, Belgium, and Denmark.  
**Stereum purpureum** Pers. See Prunus.
- PHILIBERTIA.** Climbing shrubs.  
**Aecidium philibertiae** P. Henn. Yellow rust sori on leaf blades and petioles of *P. flava* in Argentina.  
**Cercospora gillesii** Speg. On leaves of *P. gillesii* in Argentina.  
**Napicladium asclepiadinum** Speg. Pale indefinite leaf spots on *P. rotata* in Argentina.  
**Phyllosticta gillesii** Speg. Circular definite whitish leaf spots on *P. gillesii* in Argentina.

- PHILLYREA.** Ornamental woody plants grown for their handsome evergreen foliage.
- Aecidium phillyreae** DC. Leaf rust on *P. latifolia* and *P. media* in Dalmatia and Portugal.
- Ascochyta bacilligera** Wint. Irregular dull-brown to gray leaf spots on *P. angustifolia* in Portugal.
- Fusicladium radiosum** (Lib.) Lind. var. **microsporum** Sacc. On leaves of *P. media* in Spain.
- Phyllosticta goritiensis** Sacc. Circular dull-brown leaf spots on *P. angustifolia* and *P. latifolia* in Italy.
- Phyllosticta phillyreae** Sacc. Gray-ochraceous leaf spots with rufous margins on *P. latifolia* and *P. media* in Italy and France.
- Phyllosticta phillyricola** Rabh. On leaves of *Phillyrea* sp. in Germany.
- Phyllosticta romana** D. Sacc. Subcircular to oblong ashen leaf spots on *P. media* in Italy.
- Septoria crateriformis** (Dur. and Mont.) Sacc. On leaves of *P. latifolia* and *P. media* in France and Algeria.
- Septoria erythrostroma** (Dur. and Mont.) Sacc. On leaves of *Phillyrea* sp. in Algeria.
- Septoria phillyreae** Thuem. Small gray leaf spots on *P. latifolia* in Portugal.
- Zaghouania phillyreae** Pat. Golden-yellow rust sori on leaf blades and petioles and on stems of *P. angustifolia*, *P. intermedia*, *P. latifolia*, *P. media*, and *P. vilmoriniana* in Great Britain, France, Corsica, Italy, Sicily, Dalmatia, Spain, Portugal, Austria, Tunis, and Algiers.
- PHILODENDRON.** Shrubby or treelike ornamental aroids, usually climbing.
- Colletotrichum philodendri** P. Henn. Circular whitish leaf spots with dark brown margins on *P. krebsii* and *Philodendron* sp. in Brazil and Porto Rico.
- Gloeosporium aracearum** P. Henn. See Caladium.
- Mycosphaerella philodendri** (Pat.) Lind. Sunken white leaf spots with brown margins on *P. pertusum* in Venezuela.
- Phyllachora engleri** Speg. See Anthurium.
- Phyllosticta philodendri** Allesch. On leaves of *P. pertusum* in Brazil.
- Septoria aracearum** Sacc. Pale-brown leaf spots on *P. pertusum* in Italy.
- PHLEUM.** TIMOTHY. Hay and pasture grass.
- Dilophia graminis** (Fekl.) Sacc. See Triticum.
- Entyloma camusianum** P. Har. Smut sori in numerous circular to irregular dull-brown leaf spots on *P. arenarium* in France.
- Septoria culmifida** Lind. Small gray-white leaf spots with broad purple margins on *P. pratense*, *Arrhenatherum avenaceum*, *Dactylis glomerata*, and *Poa trivialis* in Denmark and Great Britain.
- PHLOMIS.** JERUSALEM SAGE. Stout herbs grown for their dense axillary whorls of large flowers.
- Ascochyta phlomidis** Jaap. On stems of *P. fruticosa* and *P. tuberosa* in Dalmatia and Russia.
- Erysiphe taurica** Lév. See Althaea.
- Ovularia phlomidis** Bub. and Wrobl. On leaves of *P. tuberosa* in Galicia and Russia.
- Phyllosticta cuestae** Gz. Frag. On leaves of *P. herba-venti* in Spain.
- Puccinia excelsa** Barcl. Dark-brown rust pustules on leaves of *P. lamiifolia* in India.
- Puccinia phlomidis** Thuem. Powdery dull-brown rust sori on leaves of *P. armeniaca*, *P. brevilabris*, *P. canescens*, *P. herba-venti*, *P. nissoli*, and *P. tuberosa* in Siberia, Russia, Turkestan, Asia Minor, Italy, Spain, and Yugoslavia.
- Septoria barrasii** Gz. Frag. On leaves of *P. purpurea* in Spain.
- PHLOX.** Showy-flowered annual and perennial garden herbs.
- Phyllosticta decussata** Syd. Circular to irregular pale-yellow leaf spots with purple margins on *P. decussata* in Germany.
- Phyllosticta phlogis** Vestergr. Subcircular dirty-white leaf spots with brown margins on *P. drummondii* in Sweden.
- Pyrenochaeta phloxidis** Mass. The stems of cultivated *Phlox* are attacked just above the ground line, cankers resulting, with consequent death of the stems. Occurs in Great Britain.
- Septoria vogliniana** Sacc. and Trott. Irregular, often confluent, chestnut-brown rust sori on leaves of *P. drummondii* in Italy.
- PHOENIX.** DATE PALM. See Palmae.
- PHORMIUM.** FLAX LILY. NEW ZEALAND FLAX. Fiber plant.
- Heterosporium** sp. Large irregular sooty blotches on lower leaf surfaces of *P. tenax* in New Zealand rotting the fiber.
- Melampsora** sp. Leaf rust on *Phormium* sp. in New Zealand.
- Phaeoseptoria** sp. On leaves of *P. tenax* in New Zealand, destroying the fiber.
- Phyllosticta haematocycla** Berk. Broad pale-brown leaf spots with red borders on *P. tenax* in Germany and Portugal.
- Physalospora phormii** Schroet. On leaves of *P. tenax* in Germany.
- Ramularia phormii** Waters. The most serious disease of *P. tenax* in New Zealand is the so-called yellow-leaf disease attributed in part to the above fungus, although other causes have been assigned in times past. Diseased plants show a yellowing and dying of the leaves which soon involves and destroys the entire plant. The seat of the trouble is in the roots, which rot off.
- Septoria** sp. Longitudinal purple stripes on leaves of *P. tenax* in New Zealand, ruining the fiber.
- Septoria melanogramma** Tass. On *P. tenax* in Italy.
- PHOTINIA.** Ornamental woody plants grown for their attractive flowers and fruits.
- Corticium salmonicolor** B. and Br. See Citrus.
- Gloeosporium tremelinum** Gäum. Anthracnose on leaves of *P. notoniana* in Java.
- Guignardia photiniae** Alm. and Da Cam. On leaves of *Photinia* sp. in Portugal.
- Gymnosporangium photiniae** (P. Henn.) Syd. See Juniperus.
- Phyllosticta photiniae** Thuem. Grayish-white leaf spots with red margins on *P. glabra* in Italy.
- Helicobasidium mompa** Tan. See Morus.
- PHRAGMITES.** Reed. Large grasses.
- Ascochyta arundinis** Fautr. and Lam. On leaves of *P. communis* in France.
- Ascochyta donacina** Sacc. On culms of *P. communis* and *Donax* sp. in Italy and France.
- Brachysporium phragmitis** Miy. On leaves of *P. communis* in China.
- Heterosporium phragmites** (Opiz.) Sacc. See Phalaris.
- Leptosphaeria eustoma** (Fr.) Sacc. See Secale.
- Leptostromella phragmitis** Brun. On leaves of *P. communis* in Japan and France.
- Mycosphaerella lineolata** (Desm.) De N. See Calamagrostis.
- Napicladium arundinaceum** (Cda.) Sacc. On *P. communis* in Japan, Dalmatia, Switzerland, and Austria. Reported from Nebraska.
- Napicladium laxum** Bub. Long dark leaf spots with yellow surrounding areas on *P. communis* in Austria.
- Phyllosticta phragmitis** Nagorni. On leaves of *P. communis* in Russia.
- Physalospora festucae** (Lib.) Sacc. See Andropogon.
- Pirostoma circinans** Fr. On leaves, leaf sheaths, and stems of *P. communis* in Mongolia, France, Italy, Belgium, Sweden, and Germany.
- Puccinia invenusta** Syd. Yellow-brown to black rust sori on leaves of *P. karka* in India.
- Puccinia isicae** (Thuem.) Wint. See Arundo.

## PHRAGMITES—Continued.

- Puccinia moriokaensis** S. Ito. Black rust pustules on leaves of *P. communis* and *P. longivalvis* in Japan.
- Puccinia obtusata** (Oth.) Ed Fisch. Yellow and black rust pustules on leaves of *P. communis*, *Ligustrum vulgare*, and *L. ihota* in Japan and Germany.
- Puccinia okatamaensis** S. Ito. Black linear rust sori on leaves of *P. communis* and *P. longivalvis* in Japan.
- Puccinia tepperi** Ludw. Leaf rust on *P. communis* in Australia.
- Puccinia trabutii** Roum. and Sacc. Leaf rust on *P. gigantea* and *Linaria reflexa* in Persia and French North Africa.
- Puccinia trailii** Plowr. Red-brown to black rust sori on leaves of *P. communis* and *Rumex acetosa* in Australia and Europe.
- Sclerospora macrospora** Sacc. See Triticum.
- Septoria arundinacea** Sacc. Oblong ochraceous leaf spots on *P. communis* in Italy and France.
- Septoria curva** Karst. On culms of *P. communis* in Finland.
- Septoria littoralis** Speg. and var. **culmicola** Gz. Frag. On leaves of *P. communis* in Spain and Italy.
- Septoria paludosa** Kab. and Bub. On leaves of *P. communis* in Bohemia.
- Septoria phragmites** Sacc. and var. **minor** Sacc. Subovate whitish leaf spots with dull-brown margins on *P. communis* in France, Spain, and Italy.
- Ustilago grandis** Fr. Smut sori at the nodes as black powdery spore masses, causing constrictions of the nodes and shortening of the internodes, on *P. communis* and *P. longivalvis* in Europe.
- Ustilago mirabilis** Sorok. Black powdery smut sori on *P. communis* in Russia.
- PHYLLANTHUS.** STAR GOOSEBERRY. OTAHEITE GOOSEBERRY. Tropical trees and shrubs, some species cultivated for ornament and fruit.
- Accidium albicans** Arth. and Holw. Leaf rust on *P. acuminatus* in Guatemala and Costa Rica.
- Accidium luzoniense** P. Henn. Rust on circular yellow leaf spots on *Phyllanthus* sp. in the Philippines.
- Accidium phyllanthi** P. Henn. Leaf rust on *P. flexuosus* and *Phyllanthus* sp. in Indo-China, New Guinea, Japan, and Uganda.
- Accidium phyllanthinum** Syd. Leaf rust on *P. reticulatus* in India and the Philippines.
- Catacauma egregium** (Rehm.) Theiss. and Syd. Black stromata on leaves of *Phyllanthus* sp. in Brazil.
- Catacauma phyllanthophilum** P. Henn. Black stromata on leaves of *P. brasiliensis* and *Phyllanthus* sp. in Peru and the Philippines.
- Phakopsora phyllanthi** Diet. Brown leaf rust on *P. acidus* (*P. distichus*) and *Phyllanthus* sp. in India and the Philippines.
- Ravenelia appendiculata** Lagh. and Diet. Rust on leaves of *P. acuminatus*, *P. galeottianus*, and *Phyllanthus* sp. in Ecuador, Mexico, and Guatemala.
- Ravenelia emblicae** Syd. Brown rust pustules on leaves of *P. emblica* and *P. polyphyllus* in India, Burma, and Ceylon.
- Ravenelia pygmaea** Lagh. and Diet. Dark-brown rust pustules on deformed petioles and stems of *Phyllanthus* sp. in Ecuador.
- Schroeteriaster fenestrala** Arth. Brown rust pustules on leaves of *P. acidus* (*P. distichus*) in Porto Rico.
- Schroeteriaster ulei** Syd. Leaf rust on *Phyllanthus* sp. in Brazil.
- Uredo phyllanthi** P. Henn. Leaf rust on *P. conami* and *Phyllanthus* sp. in Trinidad and Brazil.
- Uredo phyllanthi-longifolii** Petch. Leaf rust on *P. longifolius* in Ceylon.
- Uredo phyllanthi-reticulati** Petch. Leaf rust on *P. reticulatus* in Ceylon.
- PHYLLAUREA.** See Codiaeum.
- PHYLLOSTACHYS.** Bamboo. See Bambuseae.
- PHYMOSIA.** See Sphaeralcea.
- PHYSALIS.** GROUND CHERRY. HUSK TOMATO.
- Ascochyta alkekengi** Massal. Subcircular to angular dull-brown concentrically zoned leaf spots on *P. alkekengi* in Italy.
- Ascochyta physalicola** Oud. On stems of *P. alkekengi* in Holland.
- Ascochyta physalina** Sacc. Dull-yellow leaf spots on *P. alkekengi* in Italy.
- Cercosporina physalidicola** Speg. Indefinite, pale-brown leaf spots on *P. viscosa* in Argentina.
- Phyllosticta physaleos** Sacc. and var. **calycicola** Speg. Subcircular, ochraceous leaf spots with brown margins on *P. alkekengi*, *P. peruviana*, and *P. viscosa* in Ceylon, Argentina, Russia, and Italy.
- Vermicularia varians** Duc. See Solanum.
- PHYSOCARPUS. OPULASTER.** Ag. NINEBARK. Ornamental shrubs with white flowers, attractive inflated pods, and bright green foliage.
- Phyllachora physocarpi** Jacz. Black stromata on small red leaf spots on *P. amurensis* in Siberia.
- PHYSOSIPHON.** Epiphytic orchids. See Orchidaceae.
- PHYTEUMA.** HORNED RAMPION. Hardy herbaceous perennials used in borders and alpine gardens.
- Cercospora phyteumatis** (Frank.) Sacc. Black leaf spots with white centers on *P. spicatum* in Denmark and Germany.
- Montagnellina stellaris** (Pers.) Theiss. and Syd. Black stromata on leaves of *P. scheuchzeri* and *P. spicatum* in Europe.
- Peronospora phyteumatis** Fekl. Downy mildew on leaves of *P. betonicaefolium*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Switzerland, Norway, Denmark, Austria, Russia, and Germany.
- Phyllosticta phyteumatis** Gz. Frag. Leaf spot on *P. spicatum* in Spain.
- Pyrenopeziza phyteumatis** Fekl. On leaves of *Phyteuma* sp. in Switzerland.
- Ramularia phyteumatis** Sacc. and Wint. Ochraceous leaf spots with brown margins on *P. betonicaefolium*, *P. halleri*, *P. hedraianthifolium*, *P. michelii*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Switzerland, Italy, Spain, Bohemia, Austria, and Germany.
- Septoria phyteumatis** Siegm. Pale leaf spots with dull-brown margins on *Phyteuma* sp. in Italy.
- Septoria phyteumatis** Sacc. On leaves of *Phyteuma* sp. in Austria.
- Uromyces caricis-empervirentis** Ed. Fisch. Aecial stage on *P. betonicaefolium* and *P. orbiculare* in Switzerland, Austria, and Germany. Other stages on *Carex* spp.
- Uromyces phyteumatis** (DC.) Ung. Powdery brown rust pustules on leaves of *P. betonicaefolium*, *P. halleri*, *P. hemisphaericum*, *P. michelii*, *P. nigrum*, *P. orbiculare*, and *P. spicatum* in Europe.
- PHYTOLACCA.** POKEBERRY. POKEWEED. Herbs and woody plants, some species ornamental.
- Ascochyta phytolacca** Sacc. and Scalia. Marginal dull-brown leaf spots on *P. americana* (*P. decandra*) in Portugal.
- Heterosporium dalmaticum** Jaap. On stems of *P. americana* in Dalmatia.
- Phyllosticta phytolaccicola** Brun. Small circular brown leaf spots on *P. acinosa* in France.
- Ramularia harai** P. Henn. Effuse dull-brown leaf spots on *P. acinosa* in Japan.
- Septoria patouillardii** Sacc. and Syd. Circular dull-brown leaf spots on *Phytolacca* sp. in Ecuador.
- Septoria phytolaccae** Cav. Yellow leaf spots on *P. americana* (*P. decandra*) in Russia, Yugoslavia, and Italy.

**PICEA.** SPRUCE. Ornamental and timber trees.

- Ascochyta piniperda** Lindau. Attacks leaves and twigs of *P. abies*, *P. sitchensis*, and *Tsuga* sp. in Belgium, Russia, Germany, and Austria. The fungus also causes a damping-off of seedlings. Reported from North Carolina on *Picea* spp.
- Barclayella deformans** (Barcl.) Diet. Elongate yellow rust pustules on needles and young twigs of *P. smithiana* (*P. morinda*) in India.
- Chrysomyxa abietis** (Walbr.) Ung. Small elongate yellow rust pustules on yellow or reddish spots on needles of *P. engelmannii*, *P. abies*, *P. hondoensis*, and *P. pungens* in Japan and Europe.
- Chrysomyxa rhododendri** DeBy. See *Rhododendron*.
- Coccomyces piceae** (Fckl.) Rehm. On needles of *P. abies* in Germany.
- Cucurbitaria piceae** Borthw. Thin black crustlike stromata enveloping buds of *P. pungens* in Scotland.
- Dasyscypha abietis** Sacc. On branches of *P. abies* in Finland.
- Dasyscypha calycina** (Schum.) Fckl. See *Larix*.
- Fabraea abietina** Sacc. On the upper surfaces of needles of *P. abies* in Italy.
- Gloeosporium conigenum** Sacc. and Roum. Anthracnose on cone scales of *P. abies* in France.
- Hypodermella macrosporum** (Hart.) Lagerh. Black linear fruiting bodies on needles of *P. abies* in Switzerland, Bohemia, and Germany.
- Lophodermium abietis** Rostr. Black areas on needles of *P. glauca* (*P. canadensis*), *P. abies*, *P. smithiana* (*P. morinda*), *P. sitchensis*, *Pseudotsuga taxifolia*, and *Taxus baccata* in Sweden and Denmark.
- Macrophoma excelsa** (Karst.) Berl. and Vogl. var. **infestans** Ohl. On needles of *P. abies* in Russia.
- Myxosporium abietinum** Rostr. See *Larix*.
- Oospora abietum** Oudem. See *Abies*.
- Peridermium piceae** Barcl. Rust on needles of *P. smithiana* in India.
- Peridermium piceae-hondoensis** Diet. Rust on needles of *P. hondoensis* in Japan.
- Peridermium thomsoni** Berk. Rust on needles of *P. smithiana* in India.
- Phlyctaena strobilina** Karst. and Har. On cone scales of *P. abies* in France.
- Phomopsis occulta** Trav. On twigs of *P. abies* in Denmark, France, and Great Britain.
- Phyalospora abietina** Prill. and Delacr. See *Abies*.
- Phytophthora fagi** R. Hart. See *Fagus*.
- Sclerotopsis piceana** (Karst.) Died. On twigs and needles of *Picea* sp., *Abies alba* and *A. nordmanniana* in Europe.
- Thekopsora areolata** (Walbr.) P. Magn. See *Prunus*.
- PIERIS.** Ornamental shrubs grown for their handsome white flowers.
- Exobasidium pieridis** P. Henn. Forms concave galls on leaves of *P. ovalifolia* in Japan and India.
- Phyllosticta andromedae** West. On leaves of *P. floribunda* and *Zenobia cassinefolia* in Belgium.
- Rhytisma piceum** Berk. Circular black stromata on leaves of *P. ovalifolia* in Indo-China and India.
- Rhytisma pierdis** Pat. Black irregular fruiting areas on leaves of *P. ovalifolia* in Japan.
- PILEA.** Mostly weedy herbs, one or two species cultivated as curiosities.
- Uredo pileae** Barcl. Leaf rust on *P. trinervia* in India.
- PILOCARPUS.** Shrubs or small trees.
- Phyllosticta pilocarpi** Pat. Circular white leaf spots with red margins on *P. pinnatifolius* in Brazil.
- Puccinia pilocarpi** Cke. Black rust pustules on leaves of *P. pinnatus*, *P. pinnatifolius*, and *P. seloanus* in Brazil, Argentina, and Paraguay.
- PINGUICULA.** BUTTERWORT. Small acaulescent herbs.
- Ustilago pinguiculae** Rostr. Smut sori in anthers of *P. alpina* and *P. vulgaris* in Denmark, Estonia, and Switzerland.
- PINUS.** PINE. Ornamental and timber trees.
- Bacterium** sp. Bacteria have been found associated with a witches' broom disease of *P. pinaster* (*P. maritima*) in France.
- Bacillus pini** Vuill. Causes galls on branches of *P. halepensis* in France.
- Brachysporium pini-insularis** P. Henn. On needles of *P. insularis* in the Philippines.
- Cercospora pini-densiflorae** Hori. and Nambu. Yellow-brown spots, usually on the upper half of needles of *P. densiflora* in Japan. The disease starts in the upper part of a plant and works down, killing it in a short time. Serious in nursery beds.
- Coleosporium asterum** (Diet.) Syd. See *Aster*.
- Coleosporium cacaliae** Oth. See *Cacalia*.
- Coleosporium euphrasiae** (Schum.) Wint. Rust on needles of *P. montana* and *P. sylvestris* in Tibet, Siberia, and Europe. The alternate stage occurs on *Alectrolophus* spp. and *Euphrasia* spp.
- Coleosporium inulae** Rabh. See *Inula*.
- Coleosporium melampyri** Tul. Rust on needles of *P. montana* and *P. sylvestris* in Japan and Europe. Alternate stage on *Melampyrum* spp.
- Coleosporium petasites** Lév. Yellow rust pustules on needles of *P. sylvestris* in Japan, Siberia, and Europe. Alternate stage on leaves of *Petasites* spp.
- Coleosporium pulsatillae** Lév. See *Anemone*.
- Coleosporium senecionis** Fr. See *Senecio*.
- Coleosporium sonchi** Lév. See *Sonchus*.
- Coleosporium tussilaginis** (Pers.) Lév. Yellow rust pustules on needles of *P. sylvestris* in Europe. Alternate stage on *Tussilago farfara*.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See *Paeonia*.
- Cronartium ribicola** Fisch. The blister rust of white pine (*P. strobus*) was introduced with nursery stock into the United States from 1900 to 1910 and now threatens the existence of this important timber tree in all parts of the country. All five-needled pines, including *P. albicaulis*, *P. armandi*, *P. aristata*, *P. balfouriana*, *P. cembra*, *P. griffithii* (*P. excelsa*), *P. flexilis*, *P. koraiensis*, *P. lambertiana*, *P. monticola*, and *P. peuce*, are susceptible. The uredo and telial stages occur on species of *Ribes* and *Grossularia*, practically all species being subject to attack. The disease occurs throughout Europe, in Japan, Siberia, and western Canada.
- On the pine the fungus causes swellings on the bark of trunk, branches, and twigs, and after several years breaks through to the surface as blisterlike orange spore masses. The areas of infection may vary from one to many hundred, often so numerous as to form long continuous cankers. Trees of all ages are susceptible and are destroyed by girdling when the infections become numerous. Small trees are killed in from 4 to 6 years. The long incubation period of the mycelium in the tissues without external symptoms, other than the swelling of the bark, makes difficult or impossible the elimination of diseased plants by inspection. The uredo and telial stages occur on the lower leaf surfaces of the *Ribes*, the former appearing as powdery yellow pustules, the latter consisting of numerous small brown, hairlike projections.

## PINUS—Continued.

- Cucurbitaria pithyophila** (Schm. et Kunze) De N. Parasitic on branches of *P. sylvestris* and *Abies alba* (*A. pectinata*) in Great Britain and continental Europe. In Japan the fungus causes a serious disease of *A. firma* called millet-canker. Cankers form on trunks and larger branches and finally take on a gnarled appearance. The bark over these areas cracks and numerous small granules form beneath.
- Cylindrosporium acicolum** Bres. On needles of *P. sylvestris* in Germany.
- Cytosporina septospora** Dor. Yellow blotches on needles of *P. montana* in Russia.
- Dasyscypha calycina** (Schuem.) Fekl. See Larix.
- Diplodia pinea** Kickx. A wound parasite attacking terminal shoots of *Pinus* spp. in Europe and the Union of South Africa.
- Dothichiza pini** Sacc. On bark of *P. sylvestris* in Germany.
- Fusarium blasticola** Rostr. Causes a damping-off of coniferous seedlings, including *P. montana* and *Cryptomeria japonica* in Formosa, Denmark, and Germany. Dark patches with a water-soaked appearance occur on the stems, followed by collapse of the infected plants.
- Hypoderma pinicola** Brunch. Brownish or yellow-gray areas on needles of *P. sylvestris* in Europe, often causing defoliation.
- Hypoderma brachysporium** (Rostr.) Tub. (*Hypoderma strobilicola* Brunch.) Black linear fruiting bodies causing yellowing and premature fall of needles of *P. strobus* in Europe. Reported from Pennsylvania.
- Hypodermella sulcigena** (Duby.) Tub. Causes browning and premature shedding of needles of *P. pinaster*, *P. montana* (*P. pumila*), and *P. sylvestris* in northern Europe.
- Lachnella pini** Brun. Injures twigs of *Pinus* sp. in Scandinavia.
- Leptostroma pinastri** Desm. Needles turn red and fall. The disease is reported capable of destroying whole plantations of *P. pinaster*, *P. montana*, and *P. sylvestris* in Italy, France, Belgium, Austria, and Germany.
- Leptothyrella mougeotiana** Sacc. and Roum. On needles of *Pinus* sp. in France.
- Lophodermium gilvum** Rostr. On needles of *P. nigra* (*P. austriaca*) in Denmark.
- Melampsora pinitorqua** Rostr. See Populus.
- Mycosphaerella pinifolia** Duc. On needles of *P. pinaster* in France.
- Myxosporium abietinum** Rostr. See Larix.
- Peridermium brevius** Barcl. Needle rust on *P. griffithii* (*P. excelsa*) in India.
- Peridermium complanatum** Barcl. Two forms of this rust attacking *P. excelsa* and *P. longifolia* in India are recognized; f. *acicola* is a leaf form, while f. *corticola* occurs on the trunk and limbs causing cankers similar to those produced by *Peridermium strobii* on *P. strobus*. It is thought that the alternate host may be *Crataegus* or *Rosa*.
- Peridermium corticola** Link. Rust on branches of *P. halepensis* in Dalmatia.
- Peridermium guatemalense** Arth. and Kern. Rust on needles of *P. filifolia* in Guatemala.
- Peridermium japonicum** Syd. Needle rust on *P. thunbergii* in Japan.
- Peridermium kurilense** Diet. A rust producing fusoid swellings on branches of *P. pumila* in Japan.
- Peridermium orientale** Cke. Needle rust on *P. griffithii* and *P. longifolia* in India.
- Peridermium pini-thunbergii** Diet. Yellow masses of rust spores on needles of *P. massoniana* and *P. thunbergii* in Japan.
- Peridermium praelongum** Syd. Needle rust on *P. thunbergii* in Japan.
- Phacidium infestans** Karst. Black stromata on needles of *Pinus* spp. in northern Europe. Premature needle fall results.
- Phomopsis pitya** (Sacc.) Grove. See Pseudotsuga.
- Phyllosticta strobiligena** Desm. On cone scales of *P. strobus* in Germany.
- Phytophthora fagi** R. Htg. See Fagus.
- Phytophthora omnivora** DeBy. See Fagus.
- Septoria acuum** Oud. On needles of *P. sylvestris* in Russia and Holland.
- Sphaeropsis necatrix** Petri. and Adan. Destroys seeds in the cones of *P. pinea* in Italy.
- PIPER. PEPPER. CUBEB.** Trees, shrubs, and a few herbs, erect or climbing, with aromatic or pungent properties.
- Actinodothis piperis** Syd. Black carbonaceous stromata on leaves of *P. retrofractum*, *P. sarmentosum*, and *Piper* sp. in the Philippines and Amboina.
- Cercospora pipericola** Sacc. and Syd. Subcircular spots on leaves of *P. hispidum* in Jamaica.
- Cercospora piperis** Pat. Circular brown leaf spots on *Piper* sp. in Ecuador.
- Cercospora portoricensis** Earle. Numerous dull-brown angular leaf spots on *P. aduncum*, *P. hispidum*, *P. peltatum*, and *P. umbellatum* in Salvador, Porto Rico, Cuba, and Haiti.
- Colletotrichum necator** Mass. Anthracnose on *Piper* sp. in Malaya.
- Colletotrichum piperis** Petch. Anthracnose on *P. betle* and *P. nigrum* in Ceylon.
- Corticium salmonicolor** B. and Br. See Citrus.
- Cyclodothis pulchella** Syd. Stromata in circles on brown leaf spots on *P. carylistachyum* and *P. medium* in Porto Rico and the Philippines.
- Oidium** sp. Powdery mildew on leaves of *P. betle* in Ceylon.
- Phyllosticta pipericola** Syd. Angular pale-brown leaf spots on *P. nigrum* in Java.
- Phyllosticta piperis** Tassi. Dull-brown leaf spots on *P. betle* and *P. longum* in Italy and Ceylon. The same name has been given to a species on *P. nigrum* in Java by Hennings.
- Physalospora piperina** Syd. On stems of *P. nigrum* in India.
- Ramularia piperis** P. Henn. Circular to angular brown to black leaf spots on *Piper* sp. in Congo.
- Rhizoctonia destruens** Tassi. See Solanum.
- Rosellinia bunodes** B. and Br. See Citrus.
- Septoriopsis piperis** Stev. and Dal. Irregular dirty-white leaf spots with purple-brown borders on *P. medium* in Porto Rico.
- Stigmatea piperis** Rehm. On leaves of *Piper* sp. in Brazil.
- Uredo piperis** P. Henn. Uredo rust pustules on circular dark-brown leaf spots on *Piper* sp. and *Peperomia hernandifolia* in Porto Rico, Peru, and Brazil.
- PIPTADENIA.** Leguminous trees similar to Acacia.
- Endodothella piptadeniae** (P. Henn.) Theiss. and Syd. Black shiny stromata on leaves of *Piptadenia* sp. in Peru and Brazil.
- Phyllachora piptadeniicola** Speg. Tar spot on leaves of *P. rigida* in Argentina.
- Puccinia papillifera** Syd. Leaf rust on *P. latifolia* in Brazil.
- Puccinia piptadeniae** P. Henn. Brown rust pustules on leaves of *Piptadenia* sp. in Brazil.
- Ravenelia cebil** Speg. Leaf rust on *P. macrocarpa* and *P. peregrina* in Porto Rico and Argentina.
- Ravenelia henningsiana** Diet. Yellow-brown to dark-brown rust pustules on leaves of *Piptadenia* sp. in Brazil.
- Ravenelia simplex** Diet. Leaf rust on *P. communis* in Brazil.
- Ravenelia vilis** Syd. Leaf rust on *Piptadenia* sp. in Brazil.

- FISCIDIA. ICHTHYOMETHIA** Ag. Fish poison tree.  
**Cercospora piscidiae** P. Henn. Dull yellowish-brown spots on leaves of *P. erythrina* in Jamaica.
- ISONIA.** Erect, armed trees or shrubs.  
**Aecidium pisoniae** Arth. and Johnst. Leaf rust on *P. aculeata* in Cuba.
- ISTACIA. PISTACHE.** Trees or shrubs.  
**Coryneum pistaciae** Pat. Dull-brown leaf spots on *P. lentiscus* in Tunis.  
**Mycosphaerella pistaciae** Cke. On leaves of *P. lentiscus* in Italy.  
**Phyllosticta lentisci** (Pass.) Allesch. and var. **maculicola** Bub. Small, ashen leaf spots on *P. lentiscus* in Italy and Montenegro.  
**Phyllosticta terebinthi** Pass. Large irregular dull-brown to gray leaf spots on *P. terebinthus* in Italy.  
**Pileolaria terebinthi** (DC.) Cast. Leaf rust on *P. atlantica* and *P. terebinthus* in Dalmatia and French North Africa.  
**Septoria pistaciae** Desm. (*Phleospora pistaciae* Petr.). Black irregular leaf spots on *P. lentiscus* and *P. vera* in Asia Minor, Russia, Albania, Yugoslavia, and Italy.  
**Uromyces terebinthi** (DC.) Wint. Leaf rust on *P. falcata*, *P. lentiscus*, *P. mutica*, *P. terebinthus*, and *P. vera* in Europe, Asia Minor, Persia, and Kurdistan.
- PISTIA. WATER LETTUCE.** Tropical duckweed.  
**Phyllosticta stratiotis** Tassi. Ochraceous indefinite leaf spots on *P. stratiotes* in Italy.
- PISUM. PEA.** Annual or perennial herbs cultivated as food and forage plants.  
**Aecidium pisi-formosi** Syd. Leaves and stems of *P. formosum* in Persia covered with yellow rust sori.  
**Bacillus leguminiperdus** V. Oven. Produces brown, sunken, irregular areas on pods of *P. sativum* in Germany. Has been found capable by inoculation of attacking pods of *Lupinus* spp. and other legumes.  
**Bacterium seminum** Cayley. This bacterial disease of the pea (*P. sativum*) reported from Great Britain, is carried in the seed and is present in most parts of the plant without visible signs until flowering time. The stems then become brown with a water-soaked appearance, brown linear streaks appear at the bases of petioles, the leaves are spotted yellow, and the pods have an abnormal roughened appearance.  
**Brachysporium pisi** Oudem. Black moldy patches occur on the leaves of *P. sativum*, which turn yellow and die, causing a seedling blight, in Holland.  
**Cercospora pisa-sativae** Stevenson. Small circular yellow to brown leaf spots with darker margins on *P. sativum* in Porto Rico and Cuba.  
**Marsonia deformans** Cke. and Mass. Discolored spots on leaf blades, stipules and petioles of *P. sativum* in Australia.  
**Peronospora pisi** Syd. Downy mildew on leaves of *P. arvense* and *P. sativum* in India, France, Holland, Switzerland, Russia, Scandinavia, Bohemia, and Germany.  
**Phoma** sp. Causes a leaf spot on *P. sativum* in China.  
**Phyllosticta pisi** West. Subcircular to oblong dull-brown or yellow leaf spots with darker margins on *P. sativum* in Belgium.  
**Rhizoctonia napi** West. See Brassica.  
**Tylenchus dipsaci** Kuehn. See Narcissus.  
**Uromyces pisi** (Pers.) DeBy. Cinnamon-brown to dark-brown rust pustules on leaves of *P. arvense*, *P. sativum*, *Lathyrus angustifolius*, *L. heterophyllus*, *L. latifolius*, *L. nissolia*, *L. pisiformis*, *L. platyphyllus*, *L. pratensis*, *L. sativus*, *L. setifolius*, and *L. sylvestris* in Europe, Japan, Siberia, India, and New Zealand. The aecial stage occurs on species of *Euphorbia*. Occurs sparingly in the Middle West.  
**Ustilago entorrhiza** Schroet. A doubtful species of root smut reported on *P. sativum* in Germany.
- PITCAIRNIA.** Bromeliads.  
**Puccinia pitcairniae** Lagh. Rust on leaves of *P. palmerum* and *Pitcairnia* sp. in Ecuador and Brazil. Reported from New Mexico.  
**Septoria pitcairniae** Syd. Yellow-brown leaf spots on *Pitcairnia* sp. in Brazil.
- PITHECOCTENIUM. MONKEY COMB.** Ornamental vines cultivated for their showy flowers.  
**Prosopodium amphiphilii** (Diet. and Holw.) Arth. Leaf rust on *P. echinatum*, *P. hexagonum*, *P. muricatum*, and *Amphilophium molle* in Mexico, Isle of Pines, and Costa Rica.  
**Puccinia cordobensis** P. Henn. Rust on branches of *P. cynanchoides* in Argentina.  
**Puccinia depallens** Arth. and Holw. Leaf rust on *P. muricatum* in Costa Rica and Guatemala.  
**Puccinia medusa** Speg. Powdery rust pustules on leaves of *P. clematideum* in Argentina.  
**Puccinia phlyctopus** Syd. Brown to black powdery rust pustules on leaves of *P. hexagonum* in Mexico.  
**Puccinia pithecoctenii** Pazschke. Dull-brown to dark-brown rust pustules on leaves of *Pithecoctenium* sp. in Brazil.
- PITHECOLOBIUM.** Tropical trees and shrubs planted for ornament and shade.  
**Corticium** sp. See Thea.  
**Corticium salmonicolor** B. and Br. See Citrus.  
**Cylindrodadium pithecolobii** Petch. On leaves of *P. saman* in Ceylon.  
**Diorchidium acanthostephum** Syd. Leaf rust on *Pithecolobium* sp. in Brazil.  
**Gloeosporium pithecolobii** Koord. Large irregular gray-brown spots on leaves of *P. lobatum* in Java.  
**Maravalia pallida** Arth. and Thaxt. Leaf rust on *P. latifolium* in Trinidad.  
**Microstoma pithecolobii** Lamk. White irregular spots on lower leaf surfaces of *P. saman* in Porto Rico.  
**Montagnellina pithecolobii** (Racib.) v. Hoeh. Black stromata on yellowish leaf spots on *P. lobatum* in Java.  
**Phyllosticta pithecolobii** Young. (*P. pithecolobii-monensis* Young.). More or less circular pale-yellow leaf spots with dark-brown slightly raised margins on *P. unguis-cati* in Porto Rico.  
**Ravenelia amazonica** Syd. Leaf rust on *P. corymbosum* in Brazil.  
**Ravenelia minuta** Syd. Leaf rust on *Pithecolobium* sp. in Brazil.  
**Ravenelia pileolarioides** Syd. Leaf rust on *Pithecolobium* sp. in Brazil.  
**Septoria pithecolobii** Roum. Leaf spot on *Pithecolobium* sp. in Paraguay.  
**Stigmachora controversa** (Starb.) Theiss. and Syd. Shiny, black stromata on leaves of *P. scalare* and *Pithecolobium* sp. in Bolivia and Argentina.  
**Uredo bomfimensis** P. Henn. Brown rust sori on yellow circular leaf spots on *Pithecolobium* sp. in Brazil.  
**Uredo pithecolobii** P. Henn. Rust deforming twigs of *Pithecolobium* sp. in Brazil.  
**Uredo raciborskii** Sacc. and Syd. Brown leaf rust on *P. lobatum* in Java.  
**Uromyces albescens** Syd. Brown rust pustules on light-brown leaf spots on *P. glomeratum* in Brazil.

**PITTOSPORUM.** Hardy or half-hardy evergreen woody plants.**Helicobasidium tanakae** Miy. See *Morus*.**Mycosphaerella rubiginosa** (Cke.) Lind. On leaves of *P. rubiginosum* in Australia.**Phyllosticta pittospori** P. Brun. Subcircular to angular light-brown leaf spots on *P. tobira* in France and Italy.**Physalospora pittospori** d'Alm. and Cam. Brown apical leaf spots on *Pittosporum* sp. in Portugal.**Septoria pittospori** P. Brun. Circular deep-brown leaf spots on *P. tobira* in France and Italy.**Trabutia pittospori** (P. Henn.) Theiss. and Syd. Irregular black opaque stromata on leaves of *P. abyssinicum* in Abyssinia.**Uredo pittospori** P. Henn. Rust on leaves of *P. abyssinicum* in Abyssinia.**PLACEA.** Amaryllislike bulbous plants from Chile.**Puccinia placeae** Diet. and Neg. Yellow to dark-brown rust pustules on leaves of *Placea* sp. in Chile.**PLAGIANTHUS.** Trees or shrubs.**Puccinia plagianthis** McAlp. Brown rust pustules on leaves and flower stalks of *P. sidoides* in Tasmania.**PLATANUS.** PLANE TREE. Erroneously called sycamore. Shade trees.**Guignardia cylindrica** (Sacc. and Speg.) Lind. On leaves of *P. orientalis* in Italy and Denmark.**Phyllosticta platani** Sacc. and Speg. On leaves of *P. orientalis* in Russia and Italy.**Microstroma platani** Eddel. and Eng. Small circular grayish-white leaf spots on *P. occidentalis* in Sweden and Germany.**PLATYCODON.** BALLOON FLOWER. Perennial, glaucous, erect herbs used in border plantings.**Septoria platycodonis** Syd. Circular to irregular, white leaf spots on *P. grandiflorum* in Japan.**PLECTOCOMIA.** East Indian climbing palms. See *Palmae*.**PLECTRANTHUS.** Herbs and subshrubs with blue to lilac flowers.**Aecidium plectranthi** Barcl. Rust on leaves of *P. coetsa*, *P. glaucocalyx*, and *P. scrophularioides* in Japan and India.**Coleosporium plectranthi** Barcl. Yellow to golden rust pustules on leaves of *P. excisus*, *P. gerardianus*, *P. glaucocalyx*, *P. inflexus*, *P. longitubus*, *P. trichocarpus*, and *Ocimum* sp. in Japan and India.**Puccinia luandensis** Syd. Brown powdery rust pustules on leaves of *P. laxiflorus* and *P. nudiflorus* in the Union of South Africa.**Puccinia plectranthi** Thuem. Brown leaf rust on *P. laxiflorus* in the Union of South Africa.**PLECTRONIA.** Woody plants cultivated for their flowers.**Aecidium baumianum** P. Henn. Leaf rust on *P. abbreviata* in the Union of South Africa.**Aecidium busseanum** P. Henn. Rust on red-brown leaf spots on *P. hispida* in Tanganyika.**Aecidium incomparabile** Syd. Rust forming galls on leaves of *P. arnoldiana* in the Congo.**Aecidium plectroniae** Cke. Leaf rust on *P. barbata* and *P. gueinzii* in Natal and Queensland.**Hemileia canthii** B. and Br. Golden rust pustules on leaves of *P. campanulata*, *P. ciliata*, *P. huillensis*, *P. parviflora*, and *P. peduncularis* in Ceylon, India, Philippines, and the Union of South Africa.**Uredo detergibilis** (Thuem.) P. Henn. Rust on leaves of *P. huillensis* in the Union of South Africa.**PLEUROSPIERMUM.** Biennial or perennial hardy herbs.**Phyllosticta pleurospermi** Died. Angular gray leaf spots on *P. austriacum* in Germany.**PLEUROTHALLIS.** See *Orchidaceae*.**PLUMBAGO.** LEADWORT. Subshrubs or herbs cultivated for their phloxlike flowers.**Ascochyta plumbaginicola** P. Henn. On stems of *P. europaea* in Italy and Germany.**Ascochyta plumbaginis** Sacc. Ochraceous-brown leaf spots on *P. europaea* in Italy and Germany.**Cercospora plumbaginea** Sacc. and D. Sacc. Subcircular olivaceous, then light-buff, leaf spots on *P. europaea* in Italy.**Phyllosticta larpentae** Tassi. Subcircular ashen-white leaf spots on *P. larpentae* (*Ceratostigma plumbaginoides*) in Italy.**PLUMERIA.** FRANGIPANI. Tropical trees with milky juice grown for their showy flowers.**Ascochyta plumeria** P. Henn. On *Plumeria* sp. in Brazil.**Coleosporium domingensis** (Berk.) Arth. Golden rust pustules on leaves of *P. lutea*, *P. rubra*, and *Plumeria* sp. in Guatemala, Peru, and the West Indies.**Coleosporium plumeriae** Pat. Golden rust pustules on leaves of *P. alba*, *P. emarginata*, *P. krugii*, *P. obtusa*, and *P. rubra* in Cuba, Porto Rico, Mexico, Guatemala, and Guadeloupe.**Phyllosticta plumeriae** Tassi. Small gray to white leaf spots on *P. alba* in Italy.**Septoria plumeriae** Sacc. and Syd. Circular light-brown zoned leaf spots on *P. obtusa* in Central America and the Bahamas.**POA.** BLUE GRASS. Perennial grasses, some species cultivated as lawn, forage, and ornamental grasses.**Ascochyta kerguelensis** P. Henn. On leaves of *P. kerguelensis* (*Festuca scoparia*) in the Kerguelen Islands.**Cladochytrium graminis** Buesgen. See *Festuca*.**Claviceps setulosa** Sacc. Sclerotia in ovaries of *Poa* sp. in Switzerland and France.**Dilophia graminis** (Fckl.) Sacc. See *Triticum*.**Entyloma crepinianum** Sacc. and Roum. Doubtful smut species on *Poa* sp. in France.**Hadrotrichum virescens** Sacc. and Roum. var. *poae* Sacc. Oblong brown leaf spots on *Poa* sp. in France, Germany, and Austria.**Leptosphaeria sparsa** (Fckl.) Sacc. On leaves and culms of *Poa* sp., *Agrostis* sp., and *Calamagrostis* sp. in Switzerland.**Phyllachora poae** (Fckl.) Sacc. Black stromata on leaves of *P. alpina*, *P. nemoralis*, *P. pratensis*, and *P. sudetica* in central Europe.**Phyllosticta kerguelensis** P. Henn. On leaves of *P. cookii* in the Kerguelen Islands.**Protomyces rhizobius** Trail. Small root nodules on *P. annua* in Scotland.**Puccinia cognatella** Bub. Yellow to dark-brown rust pustules on elongate yellow spots on *P. nemoralis* in Bohemia.**Puccinia exigua** Diet. Leaf rust on *P. annua* in Brazil.**Puccinia poae-trivialis** Bub. Yellow to black rust pustules on leaves of *P. trivialis* in Bohemia.**Puccinia subandina** Speg. Leaf rust on *P. chorizantheae* in Argentina.**Roumegueria gangraena** (Fr.) Sacc. On leaves of *P. bulbosa*, *P. nemoralis*, and *P. pratensis* in Spain, Italy, and Tripoli.**Scolecotrichum compressum** Allesch. On leaves of *P. compressa* in Germany.**Septoria culmifida** Lind. See *Phleum*.**Septoria macrosperma** Speg. On leaves of *P. forsteri* in Chile.**Septoria nebulosa** Rostr. See *Calamagrostis*.**Septoria oudemansii** Sacc. On culms of *P. flexuosa* and *P. nemoralis* in Holland.**Septoria poae-annuae** Bres. and var. *sepulta* Gz. Frag. On leaves and glumes of *P. annua*, *P. bulbosa*, and *P. pratensis* in Tripoli, Spain, and Germany.

**POA**—Continued.

- Septoria poae-trivialis** Cocc. Small subcircular yellow sunken spots on leaves and culms of *P. nemoralis* and *P. trivialis* in Russia, Italy, and Germany.
- Tilletia hordei** Schoeyen. See Avena.
- Uredo porophila** Speg. Rust on leaves of *P. lanigera* in Argentina.
- Uromyces chubutensis** Speg. Leaf rust on *P. chubutensis* in Argentina.
- Uromyces poae** Rabh. Leaf rust on *P. annua*, *P. bulbosa*, *P. compressa*, *P. fertilis*, *P. nemoralis*, *P. palustris*, *P. pratensis*, *P. trivialis*, *P. violacea*, and *Agrostis alba* in Europe, Asia Minor, Nova Scotia, and north Africa. The aecial stage occurs on species of *Ranunculus*.
- Uromyces poae-alpinae** W. Rytz. Brown to black rust pustules on leaves of *P. alpina* in Switzerland.
- Uromyces seseli-graminis** Ed. Fisch. See Arrhenatherum.
- PODOCARPUS. NAGEIA** Ag. Ornamental trees grown for their evergreen foliage.
- Corynelia bispora** Fitzp. Small black stromata on lower leaf surfaces of *P. milaniana* in central Africa.
- Corynelia brasiliensis** Fitzp. Parasitic on leaves of *Podocarpus* sp. in Brazil.
- Corynelia jamaicensis** Fitzp. Circular to elliptical black stromata on leaves of *P. purdieana* in Jamaica.
- Corynelia nipponensis** Fitzp. Black stromata on leaves of *P. macrophylla* in Japan.
- Corynelia oreophila** (Speg.) Starb. Black stromata on leaves of *P. augustifolia*, *P. sellowii*, *P. chilina*, and *P. macrostachya* in Brazil, Chile, Colombia, Bolivia, and Costa Rica.
- Corynelia portoricensis** Fitzp. Raised black stromata on leaves of *P. coriacea* in Porto Rico.
- Corynelia tropica** (Auers. and Rabh.) Starb. Irregularly elongated black stromata on leaves and stems of *P. andina*, *P. chilina*, *P. costalis*, and *P. saligna* in Chile and the Philippines.
- Corynelia uberata** Fr. Circular to irregular black stromata on leaves, stems, and fruit of *P. costalis*, *P. elongata*, *P. falcata*, *P. gracilior*, *P. latifolia*, *P. nagi*, *P. nulanganianus*, and *P. thunbergii* in Japan, Philippines, New Zealand, and central Africa.
- Septoria arethusa** Penz. See Citrus.
- Septoria podocarpi** Niessl. Irregular ashen leaf spots with narrow dull-brown margins on *P. thunbergii* in Italy and the Union of South Africa.
- Thielaviopsis podocarpi** Petri. A doubtful species on roots of *Podocarpus* sp. in Italy.
- Tripospora tripos** (Cke.) Lind. Parasitic on leaves and green parts of stems of *P. elongata* and *P. thunbergii* in the Union of South Africa and on *P. lamberti* in Brazil.
- PODOLPIS.** Australian herbs cultivated for their yellow, pink, or purple flower heads.
- Puccinia podolepidis** McAlp. Bright-orange and black rust sori on leaves of *P. longipedata* in Australia.
- PODOPHYLLUM. MAY APPLE.** Herbs.
- Phomopsis podophylli** Grove. On leaves of *P. peltatum* in Great Britain.
- POGOSTEMUM. PATCHOULI.** Herbs or shrubs.
- Puccinia princeps** Syd. Rust causing large tumors (1 to 3 centimeters in diameter) on branches of *Pogostemum* sp. in India.
- POINCIANA. Royal poinciana. Flamboyant.** Ornamental-leguminous trees and shrubs.
- Cercospora bakeriana** Sacc. On twigs and leaves of *P. (Caesalpinia) pulcherrima* in the Philippines.
- Fomes lamaoensis** Murr. See Hevea.
- Ravenelia humphreyana** P. Henn. Powdery brown rust pustules causing defoliation of *P. pulcherrima* in Mexico, Costa Rica, Guatemala, Cuba, Jamaica, and Porto Rico.
- Septogloeum poincianae** Syd. Indefinite yellow areas on leaves of *P. alata* in India.
- POLEMIUM. GREEK VALERIAN. JACOB'S LADDER.** Flower-garden herbs with blue or purple flowers.
- Ascochyta polemonii** Rostr. On stems and leaves of *P. coeruleum* in Italy and Denmark.
- POLLIA.** Perennial herbs.
- Ustilago nawaschini** Racib. Smut destroying flowers of *Pollia* sp. in Java.
- POLYGALA. Annual or perennial herbs or subshrubs.**
- Aecidium palustre** Mont. Rust on leaves and stems of *Polygala* sp. in Brazil.
- Cercospora polygalae** P. Henn. On leaves of *P. paniculata* in Brazil.
- Phyllosticta chamaebuxi** A. Lesch. Ochraceous leaf spots with red margins on leaves of *P. chamaebuxus* in Germany.
- Ramularia polygala** (Schroet.) Sacc. and Syd. Light-brown spots often occupying most of leaf surface on *P. vulgaris* in Silesia.
- Septoria polygalicola** Höll. Large ochraceous leaf spots on *P. comosa* in Hungary.
- Uredo peribuyensis** Speg. Brown leaf rust on *P. americana*, *P. acicularis*, and *Polygala* sp. in Guatemala, Argentina, and Mexico.
- Uredo polygalae** Diet. Dull-yellow rust sori on yellow leaf spots on *P. japonica* and *P. ohlendoriana* in Japan.
- Uromyces polygalae** Grove. Leaf rust on *P. persicariaefolia* and *Polygala* sp. in Uganda.
- POLYGONATUM. SOLOMON'S-SEAL.** Perennial herbs with simple stems and creeping rootstocks.
- Cercospora polygonati** Rostr. Subcircular pale-brown leaf spots with dark-purple margins on *P. multiflorum* in Denmark.
- Cylindrosporium komarowi** Jacz. On *P. humile* in Russia.
- Heterosporium allii** Ell. and Mart. var. **polygonati** Oud. On leaves of *P. multiflorum* in Holland.
- Mycosphaerella asteroma** (Fr.) Lind. See Convallaria.
- Phyllosticta polygonati** Bauml. Brown leaf spots on *P. multiflorum* and *P. officinale* in Hungary and Yugoslavia.
- Phyllosticta woronowii** Woron. On leaves of *P. officinale* in Russia.
- Rhytisma punctum** Chev. Black stromata on leaves of *P. officinale* in France.
- Sclerotinia richteriana** P. Henn. Causes a blighting of *P. multiflorum* in Germany.
- Septoria brunneola** Fr. See Convallaria.
- Septoria polygonati** Kab. and Bub. Subcircular to elliptical gray leaf spots with purple-brown margins on *P. multiflorum* in Bohemia.
- POLYGONUM. SMARTWEED. Knotweed.** Erect or twining herbs, a few species grown for ornament.
- Ascochyta polygalicola** Kab. and Bub. Circular to irregular, often confluent, brown zoned leaf spots with yellow margins on *P. lapathifolium* in Bohemia.
- Cercospora paludicola** Speg. Small angular ochraceous-brown leaf spots on *Polygonum* sp. in Argentina.
- Cercospora polygoni** P. Henn. and Shir. Circular brown leaf spots with red-brown margins on leaves of *P. multiflorum* in Japan.
- Entyloma schweinfurthii** P. Henn. Smut sori on black leaf spots on *P. monspeliense* in Egypt.
- Ovularia bistortae** (Fekl.) Sacc. Brown-gray leaf spots with purple margins on *P. bistorta* and *P. viviparum* in Japan and Europe.
- Phyllosticta melanogena** Sacc. Circular to angular dark-brown leaf spots on *Polygonum* sp. in Siberia.

**POLYGONUM**—Continued.

- Phyllosticta polygoni-aviculare** Petr. Small yellowish-brown leaf spots on *P. aviculare* in Austria.
- Phyllosticta polygonorum** Sacc. Subcircular whitish leaf spots with reddish margins on *P. multiflorum*, *P. persicaria*, and *Fagopyrum esculentum* in Japan, Formosa, Russia, and Italy.
- Phyllosticta tokutaroi** Speg. Small irregular or circular brown leaf spots on *P. multiflorum* in Japan.
- Pseudopeziza bistortae** (Lib.) Fekl. On leaves of *P. bistorta* and *P. viviparum* in Greenland and Europe.
- Puccinia angelicae-mamillata** Kleb. See Angelica.
- Puccinia astantiae-vivipari** Semad. See Astrantia.
- Puccinia calumnata** Syd. Leaf rust on *P. weyrichi* in Japan.
- Puccinia congesta** B. and Br. Leaf rust on *P. chinense* and *P. tomentosum* in China and the Philippines.
- Puccinia mammillata** Schroet. Leaf rust on *P. bistortum* and *P. viviparum* in Russia, Sweden, Hungary, and Germany. Reported from New Jersey.
- Puccinia monticola** Kom. Brown rust pustules on leaves of *P. alpinum* and *P. polymorphum* in Turkestan.
- Puccinia nitida** Barcl. Brown to black rust pustules on leaves of *P. amplexicaulis* and *P. chinense* in India.
- Puccinia polygoni-sachalinensis** Pat. and Har. Powdery brown rust pustules on leaves of *P. sachalinense* in Japan.
- Puccinia septentrionalis** Juel. Leaf rust on *P. viviparum* and *Thalictrum alpinum* in Greenland, Iceland, and northern Europe. Reported from California.
- Puccinia solmsii** P. Henn. Leaf rust on *P. acuminatum*, *P. chinense*, *P. nipponense*, and *P. posumbu* in India, Indo-China, Japan, and Brazil.
- Puccinia sommerfeltii** Johans. See Thalictrum.
- Ramularia bistortae** Fekl. On leaves of *P. bistorta* in Siberia.
- Septoria polygonicola** (Lasch.) Sacc. Circular pale-ochraceous leaf spots on *P. convolvulum*, *P. orientale*, and *P. persicaria* in China, Russia, Yugoslavia, Austria, and Germany.
- Sphaelotheca alpina** Schellenb. Brown smut sori in sheaths and peduncles of *P. alpinum* in Switzerland.
- Ustilago bosniaca** Beck. A smut producing powdery, dark-violet spore masses in peduncles and inflorescences of *P. alpinum* in Yugoslavia.
- Ustilago emodensis** Berk. A smut forming clustered outgrowths up to 1 inch long on stems of *P. chinense* in Ceylon, Java, India, Australia, and the Philippines.
- Ustilago koordersiana** Bref. Smut in ovaries of *P. barbatum* and *Polygonum* sp. in China, Java, and the Philippines.
- Ustilago marginalis** (Link.) Lévl. Smut sori as swellings on leaves of *P. bistorta* in central Europe.
- Ustilago moelleri** Bref. Smut forming swollen masses of dark-brown spores in fruit of *P. hispidum* in Brazil.
- Ustilago rosulata** Syd. Powdery violet smut sori in ovaries of *P. chinensis* in the Philippines.
- Ustilago tuberculiformis** Syd. Smut sori with powdery yellow-brown spore masses in purple-brown leaf spots on *P. chinense* and *P. runcinatum* in China and India.
- POLYMNIA**. Coarse viscid composites.
- Phyllachora perlata** Syd. Irregular black stromata on leaves of *P. glabrata* in Colombia.
- Uredo banistericola** P. Henn. Cinnamon-brown rust sori on leaves of *P. sylphoides* in Brazil.
- Uromyces polyminiae** (P. Henn.) Diet. and Holw. Brown rust pustules on leaves of *P. glabrata*, *P. maculata*, and *P. sylphoides* in Mexico, Colombia, and Brazil.
- POLYPODIUM**. POLYPODY. Ferns.
- Hyalopora japonica** Diet. Rust on leaves of *P. senanense* in Japan.
- Laestadia polypodii** Sacc. and Magn. On *P. vulgare* in Italy.
- Milesina dietel'ana** (Syd.) P. Magn. Yellow-brown rust sori on small sunken spots on leaves of *P. vulgare* in Europe.
- Mycosphaerella polypodii** (Rabh.) Magn. On *P. vulgare* in Denmark, Switzerland, and Germany.
- Phyllosticta polypodii-australis** P. Henn. Irregular pale-gray leaf spots on *P. australum* in Kerguelen Islands.
- Septoria polypodii** Grove. On leaves of *P. phegopteris* in Great Britain.
- Sorica maxima** (B. and C.) Giesenh. Small black stromata on leaves of *P. crassifolium*, *P. phyllitidis*, *P. punctatum*, and *P. schomburgkianum* in Porto Rico, Cuba, St. Domingo, Ecuador, Venezuela, and Brazil.
- POLYSTACHYA**. See Orchidaceae.
- POLYSTICHUM**. HOLLYFERN. Christmas fern.
- Exoascus filicina** (Rostr.) Sacc. Small definite leaf spots on *P. spinulosum* and *Phegopteris vulgaris* in Sweden and Russia.
- Magnusiella lutescens** (Rostr.) Sadeb. Yellow leaf spots on *P. thelypteris* in Denmark.
- Metasphaeria polystichi** Feltg. On petioles of *P. filix-mas* in Luxemburg.
- PONGAMIA**. Shrubs native of the Orient.
- Cryptomyces pongamiae** (B. and Br.) Sacc. Black irregular pitchlike leaf spots on *P. glabra* in India.
- Fusicladium pongamiae** Syd. On leaves of *P. glabra*, *P. mitis*, and *P. pinnata* in India and the Philippines.
- Phyllachora pongamiae** (B. and Br.) Petch. Shiny black stromata on leaves of *P. glabra* and *P. pinnata* in India, Ceylon, Java, and the Philippines.
- Ravenelia hobsoni** Cke. Brown powdery rust pustules on leaves of *P. glabra* in India and Ceylon.
- PONTERERIA**. PICKEREL WEED. Perennial aquatic herbs.
- Phyllosticta pontederiae** Syd. Irregular dull-brown leaf spots on *Pontederia* sp. in Brazil.
- POPULUS**. POPLAR. ASPEN. Cottonwood. Timber, pulp, and shade trees.
- Ascochyta populi** Delacr. On leaves of *P. canadensis* in Italy.
- Ascochyta populicola** Kab. and Bub. On leaves of *P. alba* in Bohemia.
- Ascochyta populina** Sacc. Angular white leaf spots with dark borders on *P. nigra* in Italy.
- Ascochyta tremulae** Thuem. Subcircular small grayish leaf spots with dull-brown margins on *P. tremula* in Austria.
- Bacillus populi** Brizi. This bacterium is reported from Italy and France as the cause of a canker disease of trunks and branches of *P. alba*, *P. nigra*, and *P. tremula*.
- Didymosphaeria populina** Vuill. (*Fusicladium tremulae* Fr.). Tips of young shoots are killed and blackened, the seasonal recurrence of which destroys the crowns of infected trees. Blackened patches occur on the leaves, particularly at the tips and along margins. On *P. alba pyramidalis*, *P. canescens*, *P. nigra*, *P. tremula*, and *P. tremuloides* in Russia, Sweden, France, Denmark, Austria, and Germany. Common in Canada.

**POPULUS**—Continued.

- Diplodia gongrogena** Tem. Causes hypertrophies of bark and wood of *P. tremula* in Germany.
- Hyalopus populi** Nypels. Said to cause a canker of *P. canadensis* in Belgium.
- Hypoderma roseum** (Pers.) Fr. On bark of *Populus* sp., *Alnus* sp., and *Juglans* sp. in Great Britain, Sweden, Italy, and Germany.
- Laestadia niessli** Kze. On leaves of *P. alba pyramidalis* in Germany.
- Leptosphaeria livida** Vogl. Irregular grayish-white leaf spots on *P. canadensis* in Italy.
- Leptothyrium populi** Fekl. On leaves of *P. nigra*, *P. alba pyramidalis*, and *P. tremula* in Italy, Austria, and Germany.
- Macrophoma populi-nigrae** (Allesch.) Tass. On leaves of *P. nigra* in Germany.
- Marsonia curvata** Bub. and Kab. Irregular and confluent brown leaf spots on *P. nigra* in Bohemia. Reported from Utah.
- Marsonia piriformis** (Riess.) Sacc. Dark-brown leaf spots on *P. alba* in Europe. Reported from Utah.
- Marsonia populina** Schnabl. Leaf spots on *P. nigra* in Europe.
- Melampsora allii-populina** Kleb. Yellow or golden to brown rust pustules in yellow leaf spots on *P. alba*, *P. angulata*, *P. balsamifera*, *P. canadensis*, *P. deltoides*, and *P. nigra*, in Argentina and Europe. Alternate stage on *Allium asconicum*, *A. cepa*, *A. oleraceum*, *A. sativum*, *A. schoenoprasum*, *A. ursinum*, and *A. vinealis*.
- Melampsora ciliata** Barcl. Leaf rust on *P. ciliata* in northern India.
- Melampsora laricis** R. Hartig. Yellow, then brown, rust pustules on leaves of *P. alba*, *P. balsamifera*, *P. canescens*, *P. tremula*, *Larix decidua*, *L. griffithii*, and *L. sibirica* in Europe and Japan.
- Melampsora larici-populina** Kleb. Leaf rust on *Larix decidua* and practically all species of *Populus* in Japan, Argentina, and Europe.
- Melampsora magnusiana** G. Wagn. Golden and brown rust pustules on leaves of *Chelidonium major*, *Corydalis cava*, *C. digitata*, *C. fabacea*, *C. incisa*, *C. intermedia*, *C. laxa*, *C. pallida*, *C. solida*, *Populus alba*, *P. canescens*, and *P. tremula* in Japan, China, Siberia, and Europe. Apparently not distinct from *M. tremulae* Tul., which occurs in the United States.
- Melampsora pinitorqua** Rostr. Yellow powdery aecia on *Pinus montana* and *P. silvestris*. Telia on *Populus tremuloides*, *P. tremula*, and *P. alba pyramidalis* in Russia, Denmark, Italy, and Spain.
- Melampsora pruinosa** Tranzsch. Pale-yellow to red-brown rust sori on leaves of *P. pruinosa* in Siberia.
- Melampsora pulcherrima** (Bub.) R. Maire. Leaf rust on *P. alba* in Spain.
- Melampsora rostrupii** G. Wagn. Yellow and dark-brown rust sori on leaves of *P. alba*, *P. alba pyramidalis*, *P. balsamifera*, *P. canadensis*, *P. canescens*, *P. tremuloides*, *P. nigra*, *P. tremula*, and *Mercurialis perennis* in Europe.
- Micrococcus dendroporthos** Ludw. Small gum-producing cankers on bark of young trees of *Populus* sp. in Argentina.
- Micrococcus populi** Delacr. Bacterial cankers on stems and branches of seedlings and on trunks of older trees of *P. deltoides* and *P. nigra* in France and Italy.
- Mycosphaerella macularis** (Fr.) Schroet. On leaves of *P. tremula* in Europe. Reported from Kansas.
- Mycosphaerella populi** (Auersw.) Schroet. On leaves of *P. deltoides*, *P. nigra*, and *P. alba pyramidalis* in Europe.
- Napicladium asteroma** (Fekl.) Allesch. On leaves of *P. alba* in Germany.
- Phoma populi-nigrae** Allesch. On leaves of *P. nigra* in Germany.
- Phyllohendersonia alcidis** (Sacc.) Tass. On leaves of *P. nigra* in Italy.
- Phyllohendersonia foliorum** (Fekl.) Tass. On leaves of *P. nigra* in France, Italy, and Germany.
- Phyllosticta adjuncta** Bub. and Sere. On leaves of *P. euphratica* in Russia.
- Phyllosticta bacteriiformis** (Pass.) Sacc. On leaves of *P. nigra* in Italy.
- Phyllosticta cinerea** Pass. Ashen-white leaf spots on *P. alba* in Italy.
- Phyllosticta ostrospora** Sacc. See *Morus*.
- Phyllosticta populea** Sacc. Irregular white leaf spots with dull-brown margins on *P. alba*, *P. tremula*, and *Populus* sp. in Italy and China.
- Phyllosticta populina** Sacc. Angular white leaf spots with black margins on *P. nigra* and *P. alba pyramidalis* in Yugoslavia, Albania, Italy, and Denmark. Reported from Wisconsin.
- Phyllosticta populorum** Sacc. and Roum. On leaves of *P. balsamifera* and *P. canadensis* in Italy, France, Portugal, and Germany.
- Phyllosticta prominens** Oud. Brown leaf spots on *P. balsamifera* in Holland.
- Sclerotium scutellatum** A. S. See *Acer*.
- Septoria aegerina** Pass. On leaves of *P. nigra* in Italy.
- Septoria atosanguinea** Bub. and Sere. On leaves of *P. tremula* in Russia and Poland.
- Septoria butuliformis** Bub. and Sere. On leaves of *P. euphratica* in Russia.
- Septoria candida** (Fekl.) Sacc. White leaf spots on *P. alba* in Italy, France, and Germany.
- Septoria marmorata** Kab. and Bub. Subcircular to irregular brown, then pale-yellow, leaf spots on *P. tremula* in Denmark and Italy.
- Septoria osteospora** Bri. On leaves of *P. nigra* in France.
- Septoria tremulae** Pass. Indefinite dull-brown confluent leaf spots on *P. tremula* in Italy.
- Tifaeosporina tremulae** (Lib.) Luyk. Anthracnose on leaves of *P. alba* and *P. tremula* in Europe. Reported in a few cases from the United States under the name *Gloeosporium tremulae* (Lib.) Pass.
- PORTULACA**. Purslane. Low, fleshy, annual or perennial herbs.
- Phyllosticta portulacae** Sacc. and Speg. Subcircular white leaf spots with raised dull-brown margins on *P. oleracea* in Italy.
- Tuberculina portulacarum** Speg. Dull-brown leaf spots on *P. plan-operculata* in Argentina.
- POSOQUERIA**. Glabrous shrubs and small trees.
- Aecidium posoqueriae** Diet. Leaf rust on *P. latifolia* in Brazil.
- POTENTILLA**. CINQUEFOIL. Perennial herbs and shrubs.
- Entyoma pustulosum** Sacc. and Fautr. Leaf smut on *P. anserina* in France.
- Laestadia potentillae** Rostr. On leaves of *P. maculata* and *P. nivea* in Iceland.
- Peronospora potentillae anserinae** Gäum. Downy mildew on leaves of *P. anserina* in France and Germany.
- Peronospora potentillae reptantis** Gäum. Downy mildew on leaves of *P. recta*, *P. reptans*, and *P. supina* in northern Europe.
- Peronospora potentillae sterilis** Gaum. As above on *P. sterilis* in Switzerland and Germany.
- Phleospora fragariae** (Br. and Har.) Petr. See *Fragaria*.
- Phragmidium fragariastrum** (DC.) Schroet. Yellow to black rust pustules on leaves of *P. alba*, *P. alchemilloides*, *P. carniolica*, *P. fragariastrum*, *P. hybrida*, *P. micrantha*, *P. splendens*, *P. sterilis*, *Fragaria collina*, and *F. indica* in Europe.
- Phragmidium laceianum** Barcl. Yellow rust pustules on leaves of *P. argyrophylla* in India.
- Phragmidium nepalense** Barcl. Yellow to black rust sori on leaves of *P. nepalensis* in India.

**POTENTILLA**—Continued.

- Phragmidium papillatum** Diet. Yellow to brown rust pustules on lower leaf surfaces of *P. strijosa* in Siberia.
- Phyllosticta argentiniae** Desm. On leaves of *P. anserina*, *P. argentea*, and *Comarum palustre* in Siberia and Austria.
- Phyllosticta potentillae** Desm. On leaves of *P. fragarioides* in Siberia.
- Phyllosticta tormentillae** Sacc. Irregular pale ochraceous leaf spots on *P. tormentilla* in Italy.
- Physalospora potentillae** Rostr. On stems of *P. maculata* in Greenland.
- Physoderma vagans** Schroet. See *Ranunculus*.
- Ramularia anserina** Allesch. Subcircular, then confluent, ochraceous leaf spots on *P. anserina* in Germany.
- Ramularia martianofflana** Thuem. Large irregular dull-violet leaf spots on *P. strigosa* in Siberia.
- Septoria corcontica** Kab. and Bub. Circular to angular dull-green, then yellowish, leaf spots with brown-purple margins on *P. procumbens* and *P. tormentilla* in Bohemia.
- Septoria fragariae** Desm. See *Fragaria*.
- Septoria tormentillae** Rob. and Desm. Irregular, often oblong, brown leaf spots with rose-colored margins on *P. erecta*, *P. reptans*, *P. sylvestris*, and *P. tormentilla* in Russia, France, Denmark, Italy, Belgium, and Germany.
- Synchytrium globosum** Schroet. Small rough galls on leaves and stems of *P. reptans*, *Achillea millefolium*, *Cirsium oleraceum*, *Galium mollugo*, *Myosotis palustris*, *Sonchus asper*, *Veronica anagallis*, *V. beccabunga*, *V. chaenadrys*, *V. scutellata*, *Viola canina*, *V. odorata*, *V. persicifolia*, *V. silvatica*, and *V. stagnina*, in northern Europe.
- Synchytrium pilificum** Thomas. Small galls on leaves and stems of *P. tormentilla* in Germany.
- Uredo nervicola** Tranz. Leaf rust on *P. fragarioides* in Manchuria.
- Venturia potentillae** (Fr.) Cke. On leaves of *P. anserina*, *P. bifurca*, *P. cinerea*, *P. reptans*, and *P. tormentilla* in Siberia and Europe.
- POTERIUM**. BURNET. Rosaceous shrubs. (See also *Sanguisorba*.)
- Ovularia bulbifera** (Fekl.) Sacc. On leaves of *Sanguisorba minor* (*P. sanguisorba*) in France.
- Phragmidium sanguisorbae** (DC.) Schroet. Yellow to golden rust sori on leaves of *P. eriocarpum*, *P. magnolium*, *P. mauritanicum*, *P. muricatum*, *P. myriophyllum*, *P. polygamum*, *P. verrucosum*, *P. villosum*, *Sanauisorba dictyocarpa*, *S. duriae*, *S. media*, and *S. minor* in Algeria, Tunis, Asia Minor, Turkestan, and Europe.
- POURTHIAEA**. See *Phytinia*.
- PRATIA**. Slender prostrate or creeping herbs, sometimes grown as ornamentals.
- Uromyces pratiae** Speg. Powdery dark-brown rust pustules on leaves and stems of *P. longiflora* and *P. repens* in South America.
- PREMNA**. Shrubs.
- Cronartium premnae** Petch. Golden-yellow to brown rust pustules on leaves of *P. cordifolia* and *P. corymbosa* in Ceylon.
- Mycosphaerella oculata** Syd. Small generally circular, leaf spots on *P. odorata* in the Philippines.
- Phyllachora premnae** Syd. Black stromata on leaves of *P. cumingiana* in the Philippines.
- Puccinia premnae** P. Henn. Powdery black rust pustules on leaves of *P. macrophylla* and *P. microphylla* in Japan.
- Uredo philippinensis** Syd. Leaf rust on *P. tomentosa* in the Philippines.
- Uredo premnae** Koord. Leaf rust on *P. cumingiana*, *P. odorata*, and *P. tomentosa* in Java and the Philippines.
- PRENANTHES**. Tall perennial herbs.
- Puccinia altaica** Syd. Black rust pustules on leaves and stems of *P. diversifolia* in central Asia.
- Puccinia prenanthis** (Pers.) Lindr. See *Lactuca*.
- Puccinia prenanthis-purpureae** (DC.) Lindr. and var. **himalensis** Barcl. Powdery yellow and brown rust pustules on leaves of *P. alba*, *P. brunoniana*, *P. purpurea*, and *P. tenuifolia* in Europe and India.
- Ramularia prenanthis** Jaap. Subcircular ochraceous or brown leaf spots, which become confluent, occupying entire leaf blades of *P. purpurea* in Austria and Switzerland.
- PRESTONIA**. Tall, climbing, pubescent or glabrous shrubs.
- Helminthosporium prestoniae** P. Henn. Circular to confluent dark-brown leaf spots on *Prestonia* sp. in Peru.
- PRIMULA**. PRIMROSE. Low herbaceous plants with attractive many-colored flowers. (Including *Auricula*.)
- Cercospora primulae** Fautr. Grayish-white leaf spots with yellow margins on *P. elatior* in France.
- Cercosporella primulae** Allesch. Subcircular ochraceous leaf spots on *P. acaulis* and *P. veris* (*P. officinalis*) in Russia and Germany.
- Heterosporium auriculae** Cke. Leaves of *P. auricula* in Great Britain are disfigured by smoky patches due to this fungus.
- Mycosphaerella primulae** (Auersw. and Heufl.) Schroet. On leaves of *P. minima* in Yugoslavia.
- Ovularia primulana** Karst. Yellowish patches on lower leaf surfaces of *P. acaulis*, *P. elatior*, *P. officinalis*, *P. veris*, and *Primula* sp. in Europe.
- Peronospora oerteliana** Kuehn. Downy mildew on brown spots on lower leaf surfaces of *P. acaulis*, *P. elatior*, and *P. officinalis* in Switzerland, Bohemia, Sweden, Denmark, and Germany.
- Phyllosticta primulicola** Desm. Circular white leaf spots with tawny margins on *P. acaulis*, *P. elatior*, *P. officinalis*, *P. vulgaris*, and *P. veris* in Europe.
- Puccinia arctica** Lagh. Golden and yellow-brown rust sori on leaves of *P. sibirica* in Norway, Lapland, and northern Asia.
- Puccinia primulae** (DC.) Duby. Brown rust sori on leaf spots on *P. acaulis*, *P. cartusoides*, *P. elatior*, *P. grandiflora*, *P. integrifolia*, *P. officinalis*, and *P. vulgaris* in Japan and Europe.
- Ramularia primulae** Thuem. Circular to angular pale-ochraceous leaf spots on *P. acaulis*, *P. auricula*, *P. elatior*, *P. japonica*, *P. macrocalyx*, *P. officinalis*, *P. sinensis*, and *P. suaveolens* in Australia, Siberia, and Europe.
- Ramularia tirolensis** Maire. Subcircular, then confluent, brown leaf spots on *P. imbricata* in Tyrol.
- Septoria primulae** Buch. Circular brown leaf spots on *P. veris* and *Primula* sp. in Great Britain and Italy.
- Septoria primulicola** Rostr. Numerous ashen circular leaf spots with dull-brown margins on *P. acaulis* and *P. grandiflora* in Denmark.
- Urocystis primulicola** Magn. Smut sori changing seed capsules into masses of black spores. On *P. farinosa* and *P. veris* in Great Britain, Italy, Silesia, and Germany.
- Uromyces apiosporus** Hazsl. Dark-brown rust sori on leaves of *P. minima* in Italy, Austria, Hungary, and Germany.
- Uromyces ovirensis** Jaap. Leaf rust on *P. wulfeniana* in Austria.

**PRIMULA**—Continued.

- Uromyces primulae** Fekl. Yellow and brown rust pustules on leaves of *P. auricula*, *P. balbisii*, *P. hirsuta*, *P. latifolia*, *P. minima*, *P. pedemontana*, *P. siberica*, and *P. villosa* in Italy, Switzerland, Austria, and Germany.
- Uromyces primulae-integrifoliae** (DC.) Niessl. Leaf rust on *P. deorum*, *P. integrifolia*, *P. minima*, *P. muretiana*, and *P. viscosa* in Bulgaria, Switzerland, Denmark, and Austria.
- PROSOPIS**. MESQUITE. Hawaiian algarroba. Leguminous trees and shrubs used for shade and forage.
- Ascochyta prosopidicola** Speg. Large leaf spots on *Prosopis* sp. in Argentina.
- Phyllosticta prosopidicola** Speg. Definite white leaf spots with purple borders on *P. alba* in Argentina.
- Phyllosticta prosopidis** P. Henn. Oblong white leaf spots with brownish margins on *P. ruscifolia* in Argentina.
- Uncinula prosopidis** Speg. Powdery mildew on leaves of *P. campestris* in Argentina.
- Uredo prosopidis** Jacz. Powdery brown rust pustules on leaves of *P. stephaniana* in Transcaucasia.
- PROSTANTHERA**. Shrubs, commonly strongly scented.
- Phyllosticta prostantherae** Cke. Brown leaf spots on *P. lasianthos* in Australia.
- PROTEA**. Tender shrubs, small trees or acaulescent perennial herbs.
- Leptosphaeria protearum** Syd. Subcircular, then irregular, sordid-white leaf spots on *P. madiensis* and *P. melaleuca* in Uganda and the Union of South Africa.
- Oligostroma maculiformis** (Wint.) Doidge. Black stromata on leaves of *P. abyssinica*, *P. flanaganii*, *P. grandiflora*, and *P. neriifolia* in the Union of South Africa.
- Oligostroma proteae** Syd. On *P. flanaganii* in the Union of South Africa.
- Phaeosphaerella senniana** Sacc. On *P. abyssinica*, *P. acaulis*, and *P. melaleuca* in the Union of South Africa and Abyssinia.
- Phyllachora proteae** Wakef. Tar spot on leaves of *P. mellifera* in the Union of South Africa.
- Pseudomonas proteamaculans** Paine and Stans. This bacterium produces numerous dome-shaped blisters, 1 to 3 mm. in diameter, on upper leaf surfaces of *P. cynaroides* in Great Britain. On young leaves the areas are larger, red to reddish brown with vermilion borders and depressed.
- PRUNELLA**. See Brunella.
- PRUNUS**. PLUM. CHERRY. APRICOT. (Including *Amygdalus*, PEACH, ALMOND, and NECTARINE). Fruit and nut trees.
- Ascochyta chlorospora** Speg. A shot-hole disease of leaves and fruit spot of *P. armeniaca*, (*Amygdalus communis*) *P. cerasus*, *P. divaricata*, *P. domestica*, and *P. (Amygdalus) persica* in Russia and Australia.
- Ascochyta crystallina** McAlp. Leaf spot on *P. amygdalus* in Australia.
- Ascochyta ovalispora** McAlp. On leaves and twigs of *P. cerasus* and *P. domestica* in Australia.
- Ascochyta pruni** Kab. and Bub. Circular, often indefinite, brown, then ashen-gray, leaf spots on *P. padus* in Bohemia.
- Ascospora padi** Grev. Defoliates *P. cerasus* in Europe.
- Auerswaldiella puccinioides** (Speg.) Theiss. and Syd. Dull-black stromata on leaves of *P. argentinensis* and *P. sphaerocarpa* in Central and South America.
- Bacillus spongiosus** Aderh. and Ruhl. Large cankers with heavy gum exudate on trunks of *P. cerasus* in Germany. This is apparently one of the gumming diseases of *Prunus* spp., the exact causes of which have for the most part not been worked out.
- Bacterium** sp. (Blossom-blight.) See *Pyrus*.
- Caecoma makinoi** Kus. A rust attacking the buds of *P. domestica*, *P. grayana*, *P. mume*, and *P. pseudocerasus* in Japan, causing hypertrophy and distortion of the young shoots as they develop.
- Camarsporium prunifolium** McAlp. Elongate narrow gray leaf spots with raised margins on *P. armeniaca* in Australia.
- Cercospora amygdali** Ali Riza. Subcircular, often confluent, gray spots on leaves of *P. amygdalus* in Turkey.
- Cercospora guliana** Sacc. Subcircular yellow leaf spots on *P. amygdalus* in Malta.
- Clasterosporium persicum** (Sacc.) Tsuji (*Cercosporella persicum* Sacc.) On leaves of *P. persica* in Japan and Italy.
- Coniothecium albo-cinctum** Preuss. Light gray patches on shoots of *P. domestica*, *P. persica* and *Malus* in Australia and Germany.
- Coniothyrium pruni** McAlp. Circular to irregular dirty-gray spots with raised dark-red margins on leaves and fruit of *P. armeniaca* and *P. domestica* in Australia.
- Corticium salmonicolor** B. and Br. See *Citrus*.
- Cucurbitaria pruni-mahaleb.** Allesch. Attacks and kills buds and small twigs, a witches'-broom effect resulting. On *P. mahaleb* in Europe. Reported from New York.
- Cylindrosporium tubeufianum** Allesch. Attacks green fruit of *P. avium* and *P. padus* in Germany.
- Dermatella prunastri** Pers. Attacks and destroys terminal twigs of *P. domestica* in Great Britain.
- Diaporthe perniciosus** Marchal. This fungus attacks and destroys trees of all ages, but more particularly the young ones. Slightly sunken, reddish, elongate cankers appear, increasing rapidly in size and bringing about wilting and browning of the leaves which fall prematurely. Under some circumstances there is also a rot of the fruit. The hosts are *P. armeniaca*, *P. persica*, *P. cerasus*, *P. domestica*, *Malus* (apple), and *Pyrus* (pear) in Belgium, France, and Great Britain.
- Didymaria prunicola** Car. Brown leaf spots on *P. domestica* in Italy, causing premature leaf fall.
- Dothiorella vinosa** Marchal. Forms cankers on trunks and branches, spots on leaves, and rot of fruit of *Malus*, *Prunus domestica*, *Pyrus*, and *Ribes* in Belgium.
- Exoascus andinus** (Palm.) Sacc. and Trott. Indefinite reddish, often swollen, areas on leaves of *P. salicifolia* in Ecuador.
- Exoascus rostrupianus** Sadeb. Forms "fruit pockets" on *P. spinosa* in Europe.
- Fusarium gemmiperda** Aderh. Attacks and destroys buds of *P. cerasus* in Germany.
- Fusicladium amygdali** Duc. Small circular to angular brown leaf spots which enlarge to include the entire leaf blade, defoliation resulting. Young fruit and blossoms drop prematurely due to the weakening of the twigs. On the twigs dark-green, then brown, rough spots occur on shaded surfaces. On *P. amygdalus* in Italy and France.
- Fusicladium cerasi** Sacc. (*Venturia cerasi* Aderh.) Olive-black minute velvet blotches on fruit of *P. cerasus* in Europe. Diseased fruits mummify. Leaves and twigs are rarely attacked. Probably not distinct from *Cladosporium carpophilum*. Reported from Ohio and New Mexico.
- Fusicladium pruni** Duc. Gray or brown leaf spots on *P. domestica* in France.
- Fusicoccum amygdali** Delacr. Causes die-back of *P. amygdalus* in France.
- Gloeosporium armeniaca** Speg. Anthracnose on *P. armeniaca* in Argentina.
- Gloeosporium cerasi** Lind. Anthracnose on *P. avium* in Japan.
- Glomerella mume** (Hori.) Hemmi. Anthracnose on *P. mume* in Japan.
- Gnomonia circumscissa** McAlp. Circular ochraceous to gray leaf spots on *P. amygdalus*, *P. armeniaca*, *P. cerasus*, *P. domestica*, and *P. persica* in Australia: Possibly not distinct from the following species.

## PRUNUS—Continued.

- Gnomonia erythrostoma** (Pers.) Auersw. Small yellow leaf spots which become brown and much enlarged, causing defoliation in part. The fruit withers or ripens unevenly, with poor color and flavor. On *P. cerasus* in Europe and New Zealand.
- Graphium ulmi** Schwartz. See Ulmus.
- Haplosporella pruni** McAlp. Causes large warted swellings on branches of *P. armeniaca* and *P. domestica* in Australia.
- Helicobasidium mompa** Tan. See Morus.
- Helicobasidium tanakae** Miy. See Morus.
- Kellermannia pruni** McAlp. On leaves of *P. armeniaca* and *P. persica* in Australia.
- Laestadia circumscissa** Sacc. Subcircular to oblong red to brown leaf spots on *P. spinosa* in Italy. One of the shot-hole diseases.
- Microstoma tonellianum** Ferr. Irregular white leaf spots on *P. domestica* in Italy.
- Mycosphaerella cerasella** Aderh. (*Cercospora cerasella* Sacc.) Brown shot-hole disease on *P. avium*, *P. cerasus*, *P. itosakura*, and *P. yamasakura* in Japan, Russia, Bulgaria, Yugoslavia, and Italy.
- Myxosporium pruni-mahaleb** Moreillon. Attacks and destroys buds and small twigs of *P. mahaleb* in Europe, witches'-brooms developing ultimately.
- Nacmospora crocea** (Bon.) Sacc. Two-year-old shoots are attacked, the young leaves wilting, turning brown and dying about blossom time. The flowers also turn brown and the shoots themselves shrivel and die. On *P. persica* in Great Britain.
- Ovularia cerasi** McAlp. Scabby patches on fruit of *P. cerasus* in Australia.
- Ovularia circumscissa** Sorok. Subcircular reddish leaf spots on *Prunus* sp. in the Caucasus.
- Phoma persicae** Sacc. Grayish areas on twigs of *P. persica* in Australia, apparently causing die-back.
- Phyllohendersonia foliorum** (Fckl.) Tass. On leaves of *P. domestica* in Italy, France, and Germany.
- Phyllosticta albomaculans** Kab. and Bub. Irregular leaf spots, white above, yellow-brown beneath, on *P. padus* in Bohemia.
- Phyllosticta armenicula** Farneti. Irregular spots on fruit of *P. armeniaca* in Italy.
- Phyllosticta chlorospora** McAlp. Elongated irregular leaf spots dirty-white above, pale red-brown beneath, on *P. domestica* in Australia.
- Phyllosticta destruens** Desm. See Celtis.
- Phyllosticta macrospora** McAlp. Irregular gray leaf spots with reddish margins, the centers falling out, on *P. armeniaca*, *P. divaricata*, and *P. domestica* in Australia.
- Phyllosticta mahaleb** Thuem. Circular ashen leaf spots on *P. mahaleb* in Italy and Portugal.
- Phyllosticta matthiolana** (Sacc. and Matt.) McAlp. Circular to elliptical or irregular brown leaf spots on *P. amygdalus*, *P. armeniaca*, *P. cerasus*, *P. communis*, *P. domestica*, *P. lauro cerasus*, and *P. persica* in Italy and Australia.
- Phyllosticta minutissima** Kab. and Bub. Circular to irregular, then confluent, pale ashen-brown leaf spots on *P. spinosa* in Bohemia.
- Phyllosticta padi** P. Brun. Brown or ochraceous subcircular to irregular spots on leaves of *P. padus* in Italy and France.
- Phyllosticta persicophila** Trav. and Migliardi. Marginal irregular ochraceous leaf spots with white centers on *P. persica* in Italy.
- Phyllosticta pruni-avium** Allesch. Yellow-brown leaf spots on *P. avium* in Germany.
- Phyllosticta pruni-domesticae** Vogl. Gray leaf spots, becoming white and somewhat swollen, on *P. domestica* in Italy.
- Phyllosticta serebriankowii** Bub. Circular to elliptical gray, more or less concentrically zoned, leaf spots on *P. padus* in Russia.
- Phyllostictiella pruni-spinosae** (Allesch.) Tass. Ashen leaf spots on *P. spinosa* in Italy and Germany.
- Phytophthora syringae** Kleb. See Syringa.
- Polystigma ochraceum** (Wahlenb.) DC. Stromata on subangular leaf spots on *P. cerasus*, *P. amygdalus*, *P. padus*, and *Malus* sp. in Japan, India, and Europe.
- Puccinia cerasi** (Bereng.) Cast. Golden-yellow to brown rust pustules on leaves of *P. avium*, *P. cerasus*, *P. domestica*, *P. persica*, and *P. spinosa* in Europe.
- Puccinia radiata** Shirai. Rust pustules on leaves of *P. pseudo-cerasus* in Japan.
- Ramularia cerasorum** Marchal. On leaves of *P. cerasus* in Belgium.
- Ramularia lata** Sacc. White leaf spots on *P. laurocerasus* in France. A doubtful species.
- Rhodoseptoria ussuriensis** Naoum. Destroys leaves and fruits of *Prunus* sp. in Russia.
- Sclerotinia kusanoi** P. Henn. Ashen-white areas on leaves of *P. pseudo-cerasus* in Japan.
- Sclerotinia laxa** Aderh. and Ruhl. Rot and mummification of fruit of *P. armeniaca* and *P. communis* in Australia, Russia, France, Hungary, and Germany.
- Septobasidium acaciae** Saw. See Acacia.
- Septoria amygdali** McAlp. On leaves of *P. amygdalus* and *Prunus* sp. in Australia and Russia.
- Septoria anomala** Sacc. Marginal leaf spots on *P. spinosa* in France.
- Septoria cerasi** Pass. Dark sanguineous leaf spots on *P. avium* and *P. cerasus* in France.
- Septoria disseminata** Desm. Leaf spot on *P. laurocerasus* and *P. lusitanica* in France.
- Septoria effusa** (Lib.) Desm. On leaves of *P. amygdalus* and *P. cerasus* in France, Italy, Austria, and Australia.
- Septoria lauro-cerasi** Desm. and Grog. Small brown leaf spots on *P. laurocerasus* in France.
- Septoria lauro-cerasina** Pass. On leaves of *P. laurocerasus* in Italy.
- Septoria myrobalanae** Brun. Circular to oblong brown leaf spots on *P. cerasifera* (*P. myrobalana*), in France.
- Septoria pruni-mahaleb** Therry. Small circular, then confluent, brownish-red leaf spots on *P. mahaleb* in France.
- Septoria staganosporioides** Mass. On leaves of *P. laurocerasus* in Italy.
- Sporotrichum persicae** Póil. White incrustations or patches on fruit of *P. persica* in Italy. The fungus in itself does not cause a rot but lowers the market value of the fruit.
- Stereum purpureum** Pers. The silver-leaf disease is considered a serious disease of *P. armeniaca*, *P. cerasus*, *P. communis*, *P. domestica*, *P. japonica*, *P. lusitanicus*, *P. persica*, *P. spinosa*, *P. serotina*, *Malus*, and *Pyrus* in Europe, New Zealand, Canada, and the Union of South Africa. It has also been reported on *Aesculus* sp., *Pernettya mucronata*, *Crataegus* spp., *Philadelphus* sp., *Ribes* sp., and *Rosa sericca*. The fungus considered as the cause of the disease is widespread throughout the United States, but apparently never occurs, or certainly only rarely, in connection with the silver-leaf type of disease, except in Oregon and Washington.
- Silver-leaf is characterized by a silvering of the foliage, which spreads gradually until the entire tree is affected. The limbs bearing these silvered leaves die back and ultimately the tree itself dies. The fungus does not attack the leaves directly, but causes a brown heart rot of roots and trunk and is supposed to secrete some deleterious substance which causes the external signs of the disease.
- Stigmia briesiana** Far. On *P. armeniaca* in Europe.

**BRUNUS**—Continued.

- Taphrina mexicana** Syd. Small witches'-brooms are formed on *P. microphylla* in Mexico by this fungus.
- Taphrina minor** Sadeb. Reddish spots on leaves and twigs of *P. avium*, *P. cerasus*, and *P. chamaecerasus* in New Zealand, Denmark, Holland, Russia, Great Britain, and Germany. Diseased twigs become malformed.
- Taphrina mume** Nish. On *P. armentaca*, *P. domestica*, and *P. mume* in Japan.
- Taphrina pseudo-cerasi** Shir. Causes a witches'-broom effect on *P. subhirtella* (*P. miqueliana*) and *P. pseudo-cerasus* in Japan.
- Taphrina trunicola** S. Kus. This fungus attacks the buds of *P. incisa* and *P. pseudocerasus* in Japan. It causes hypertrophy and distortion of twigs and petioles. Leaf blades are commonly not changed, but the flowers become succulent pink masses of abnormal tissue.
- Thekopsora areolata** (Fr.) P. Magn. Dark-brown crustlike rust sori on leaves of *P. padus*, *P. serotina*, and *P. virginiana* in Japan and Europe. The aecial stage forms red-brown to dark-brown pustules on cone scales of *Picea ajanensis*, *P. excelsa*, *P. glehnii*, *P. hondoensis*, and *P. obovata*.
- Thekospora pseudocerasi** Hirotsuka. Leaf rust on *P. pseudocerasus* in Japan.
- Uncinula prunastri** (DC.) Sacc. Powdery mildew on leaves of *P. dasyphylla*, *P. domestica*, *P. insititia*, *P. pumila*, *P. spinosa*, and *Crataegus* sp. in Europe and Transcaucasia.
- Uredo pruni-maximowiczii** P. Henn. Rust sori in young branches and petioles of *P. maximowiczii* in Japan.
- Uredo persicae** Speg. Leaf rust on *P. persica* in Brazil.
- Valsa japonica** Miy. and Henn. A wound parasite of trunks and branches of *P. cerasus*, *P. koidzumii*, *P. kurilensis*, *P. mume*, *P. persica*, *P. sachalinensis*, *P. serrulata*, and *P. yedoensis* in Japan.
- Verticillium** sp. A wilt disease of *P. cerasus* is reported as serious in Holland, but details are lacking.
- PSEUDOTSUGA**. DOUGLAS FIR. Timber and ornamental coniferous trees.
- Lophodermium abietis** Rostr. See *Picea*.
- Myxosporium abietinum** Rostr. See *Larix*.
- Oospora abietum** Oudem. See *Abies*.
- Phomopsis pseudotsugae** Wils. This a new and apparently threatening disease of *P. taxifolia* in Great Britain and Holland. It also attacks *Larix kaempferi* (*L. leptolepsis*), *L. decidua*, *Abies grandis*, *A. pectinata*, and *Tsuga heterophylla*.
- Cankers appear on the branches or stems, causing constrictions and the death of the portion of branch or stem beyond the canker. The cankers may occur along one side only, in which case death of the branch is delayed. The minute black fruiting bodies form on the cankers and enlarged gum pockets appear in the swollen portions of the limbs or trunk just above the diseased areas. Diseased trees have an unthrifty appearance, due to the yellowing of the foliage of diseased branches, and younger trees, particularly nursery stock, are killed.
- Phomopsis pithya** (Sacc.) Grove. This fungus, related to the preceding one, occurs on *Pinus*, *Pseudotsuga*, and other conifers in Europe but is generally considered as saprophytic only.
- PSIDIUM**. GUAVA. Tropical and subtropical fruit trees and shrubs.
- Aithaloderma clavatisporum** Syd. On leaves of *P. guajava* in Siam.
- Catacauma goyazense** (P. Henn.) Theiss. and Syd. Tar spot on leaves of *P. guajava* in Paraguay and Brazil.
- Catacauma subcircinans** (Speg.) Theiss. and Syd. Circular black shiny stromata on leaves of *P. cattleianum* in Brazil and Argentina.
- Cercospora psidii** Rangel. On leaves of *P. araca* in Brazil.
- Linhartia höhnelii** Rehm. Small circular leaf spots on *Psidium* sp. in Brazil.
- Meliola psidii** Wint. Superficial black circular to irregular fungus patches on leaves of *P. guajava* and *P. pomiferum* in Central and South America and the West Indies.
- Phyllachora cayennensis** (DC.) Theiss. and Syd. See *Ficus*.
- Phyllachora tropicalis** Speg. Subcircular black stromata on leaves of *P. thea* in Brazil and Argentina.
- Phyllosticta psidiella** Tassi. Small irregular dirty-white leaf spots on *P. montanum* in Italy.
- Phyllosticta psidii** Tassi. Dull-brown leaf spots on *P. araca* in Italy.
- Puccinia psidii** Wint. Yellow to brown rust pustules on dull-brown or brown-purple leaf spots on *P. araca*, *P. guajava*, *P. pomiferum*, and *Eugenia* (*Caryophyllus*) *jambos* in Porto Rico, Cuba, Ecuador, and Brazil.
- PSOPHOCARPUS**. BOTOR. Ag. Tall, twining herbs with large tuberous roots.
- Woroninella psophocarpi** Racib. Forms numerous orange galls on leaves, stems, and fruit of *P. longipedunculatus*, *P. palustris*, and *P. tetragonolobus* in Java, India, the Philippines, and west tropical Africa.
- PSORALEA**. SCURF-PEA. Herbs and shrubs used somewhat in border plantings.
- Erysiphe taurica** Lévl. See *Althaea*.
- Phlyctaena psoraleae** (Cast.) Karst. and Har. On stems of *Psoralea* sp. in France.
- Woroninella aequatoriensis** Syd. On leaves of *P. mutisi* in Ecuador.
- PSYCHOTRIA**. Tropical shrubs and small trees.
- Aecidium iquitosense** P. Henn. Yellow rust sori deforming leaves and stems of *Psychotria* sp. in Peru and Brazil.
- Aecidium micranthum** Syd. Leaf rust on *P. elongata* in Ceylon.
- Aecidium psychotriae** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- Inocyclus psychotriae** Syd. On leaves of *P. luzonensis* in the Philippines.
- Phyllachora psychotriae** Rehm. Scattered black stromata on circular yellowish leaf spots on *Psychotria* sp. in Brazil.
- Phyllosticta psychotriae** P. Henn. On leaves of *Psychotria* sp. in Brazil.
- Puccinia fallaciosa** Arth. Brown leaf rust on leaves of *Palicourea crocea*, *P. riparia*, and *Psychotria patens* in Porto Rico.
- Puccinia psychotriae** P. Henn. Brown rust pustules on leaves of *Psychotria* sp. in Brazil.
- Uredo holstii** P. Henn. Leaf rust on *Psychotria* sp. in tropical Africa.
- Uredo mkusiensis** P. Henn. Leaf rust on *Psychotria* sp. in Uganda and Tanganyika.
- Uredo psychotriae-volkensii** P. Henn. Yellow rust pustules on circular, then confluent, dull-brown leaf spots on *P. volkensis* in tropical Africa.
- Uredo psychotriicola** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- Uromyces psychotriae** P. Henn. Leaf rust on *Psychotria* sp. in Brazil.
- PTELEA**. HOP TREE. Deciduous shrubs and small trees with handsome foliage and attractive green fruit.
- Ascochyta pteleae** Kab. and Bub. Circular leather-brown, often zoned, leaf spots with narrow dark-brown margins on *P. trifoliata* in Bohemia.
- Phyllosticta pteleae** Höll. Large irregular, then confluent, brown leaf spots on *P. trifoliata* in Hungary.

- PTERIS.** BRAKE. A large genus of ferns, many forms cultivated.
- Aphelenchus olesistus** Ritz. Bos. See Begonia.
- Ascochyta pteridis** Bres. On leaves of *Pteridium aquilinum* (*P. aquilinum*) in Germany.
- Dothidella pterodophila** Speg. Black stromata covering leaves of *Pteris* sp. in Brazil.
- Hyalospora filicum** Diet. See Asplenium.
- Mycosphaerella prominula** (Speg.) Lind. On leaves of *P. aquilinum* in Europe.
- Phyllachora rhopographoides** Wint. Doubtful species on *P. aquilinum* in Brazil.
- Rhopographus malmei** Starb. Linear black stromata on leaves of *Pteris* sp. in Brazil.
- Rhytisma piceum** Berk. Black opaque stromata on leaves of *Pteris* sp. in India.
- Septoria aquilina** Pass. Brown spots on leaves of *P. aquilina* in Italy. Reported from Wisconsin.
- Septoria pteridicola** Kab. and Bub. Pale-brown or dull-yellow irregular, then confluent, areas on leaves of *P. aquilina* in Italy.
- Taphrina laurencia** Giesenh. Causes much-branched leafy outgrowths on *P. quadriaurita* in Ceylon and India.
- Taphrina rhomboidalis** Syd. and Butl. Yellow to brown areas on leaves of *P. quadriaurita* in India.
- Taphrina tonduziana** P. Henn. Circular to angular spots on leaves of *P. aculeata* in Costa Rica.
- PTEROCARPUS.** Leguminous trees or woody climbers.
- Aldona stela-nigra** Rac. On leaves of *P. indicus* in Java and the Philippines.
- Catacauma pterocarpi** Syd. Small irregular black stromata on white leaf spots on *P. angolensis*, *P. indicus*, and *P. saxatilis* in Java, the Philippines, and the Union of South Africa.
- Helminthosporium obovatum** Mass. On leaves of *P. indicus* in Malaya.
- Placostroma pterocarpi** (Mass.) Theiss. and Syd. Black stromata on yellow-green leaf spots on *P. indicus* in Malaya and the Philippines.
- Pseudothrips pterocarpi** Syd. On leaves of *P. indicus* in the Philippines.
- Uncinula incrassata** Salm. Powdery mildew on leaves of *P. melliferus* in central Africa.
- PTEROCARYA.** WING NUT. Ornamental trees grown for their handsome pinnate foliage.
- Phyllosticta pterocaryae** Thuem. Irregular, often marginal dull-yellow to gray leaf spots on *P. caucasica* in Portugal.
- PTYCHOSPERMA.** See Palmae.
- PUERARIA.** KUDZU BEAN. Twining herbs or shrubs.
- Cercospora puerariae** Syd. Olive-brown leaf spots on *P. phaseoloides* in the Philippines.
- Synchytrium puerariae** Miy. (*Woroninella puerariae* [Henn.] Syd.). Small galls on leaves and stems of *P. hirsuta*, *P. phaseoloides*, and *P. sericea* in Japan, China, Java, India, New Guinea, and the Philippines.
- PULMONARIA.** LUNGWORT. Perennial herbs with creeping root-stocks, grown in gardens.
- Entyloma serotinum** Schroet. See Borago.
- Phyllosticta pulmonariae** (Fckl.) Sacc. Small, dull-brown, leaf spots on *Pulmonaria* sp. in Switzerland.
- Puccinia bromina** Erikss. See Bromus.
- Ramularia cylindroides** Sacc. Dull-yellow leaf spots on *P. mollissima*, *P. officinalis*, *P. styriaca*, *P. tuberosa* and *Verbascum* sp. in Europe.
- Septoria pulmonariae** Sacc. Subcircular dark-brown leaf spots on *P. officinalis* and *Pulmonaria* sp. in Russia and Italy.
- PUNICA.** POMEGRANATE. Small trees with edible fruit.
- Aposphaeria punicina** Sacc. Found in connection with die-back of twigs of *P. granatum* in southern Europe.
- Cercospora punicae** P. Henn. Circular dark-fuscous leaf spots on *P. granatum* in China and Japan.
- Dothiorella sanninii** Ciferri. Surface of fruit of *P. granatum* in Italy covered with small black patches which disfigure and lower the commercial value.
- Megalonectria caespitosa** Speg. Forms cankers on trunk and branches of *P. granatum* in Brazil and Argentina.
- Phyllosticta granati** Rangel. Circular to angular brown leaf spots on *P. granatum* in Brazil.
- Phyllosticta punica** Sacc. and Speg. Small subcircular white leaf spots with red margins on *P. granatum* in Italy.
- PYCNOSTACHYS.** Erect perennial herbs.
- Uredo pycnostachydis** Kalchbr. Leaf rust on *P. dawei* and *P. reticulata* in Uganda.
- PYRETREUM.** See also Chrysanthemum.
- Puccinia balsamitae** (Strauss.) Rabh. Leaf rust on *P. (Chrysanthemum) balsamita*, *P. tanacetifolium*, and *Tanacetum* sp. in Europe.
- Puccinia proximella** Syd. Leaf rust on *P. millefoliatum* in Russia.
- Septoria pyrethri** Bres. and Krieg. Irregular buff leaf spots on *P. macrophyllum* and *P. (Chrysanthemum) parthenium* in Russia and Germany.
- PYROLA.** Shinleaf. Low, smooth, perennial herbs with subterranean shoots.
- Actinonema pirolae** Allesch. Large irregular black leaf spots on *P. secunda* in Germany.
- Phyllosticta pyrolae** (Ehrbg.) Allesch. Circular, often confluent, purple-brown leaf spots on *P. rotundifolia* in Germany.
- Sclerotinia pyrolae** Grosse. Forms black sclerotia in pods of *P. chlorantha*, *P. media*, *P. minor*, *P. rotundifolia* and *P. uniflora* in Russia.
- PYRUS.** PEAR. Trees and shrubs cultivated for fruit and ornament.
- Ascochyta pirina** Pegl. Irregular brown spots on fruit and leaves of *P. communis* in Russia, Malta, and Italy.
- Bacillus** sp. (Blossom Blight.) This bacterium causes a blighting of flowers and sometimes a die-back of twigs of *Pyrus*, *Malus*, and *Prunus* spp. in England. The sepals turn gray and blacken, the discoloration involves the calyx and flower stalk, and the entire blossom turns black, sometimes falling.
- Bacterium nectarophilum** Doidge. Causes a blackening of flowers, peduncles, and young fruit. of *Pyrus* in the Union of South Africa. Distinct from fire blight.
- Bacterium pyri** Brzez. This bacterium is reported as the cause of a trunk and branch canker of *Pyrus* (pear) in central Europe.
- Cercospora picicola** K. Saw. Angular, then coalescent, brown and finally ashen leaf spots on *P. communis* and *P. sinensis* in Japan.
- Cercospora porrigo** Speg. Leaf spot on *P. communis* in Argentina.
- Colletotrichum piri** Noack and f. *tirolense* Bub. Anthracnose on *Pyrus* and *Malus* in Brazil and Austria.
- Coniothecium chromatosporum** Cda. See Malus.
- Coryneum foliicolum** Fckl. See Crataegus.
- Diaporthe pernicioso** Marchal. See Prunus.
- Dothiorella vinosa** Marchal. See Prunus.
- Entomosporium mespili** (DC.) Sacc. See Cotoneaster.

**PYRUS**—Continued.

- Exoascus bullatus** (B. and Br.) Fckl. Small, then confluent, reddish blisterlike swellings on leaves of *P. amygdaliformis*, *P. communis*, *P. japonica*, *P. sinensis*, *P. ussuriensis*, and *Cydonia japonica* in Australia, Chile, and Europe.
- Gymnosporangium cunninghamianum** Barcl. See Cupressus.
- Gymnosporangium koreaense** (P. Henn.) Jacks. See Juniperus.
- Gymnosporangium mespili** (DC.) Kern. See Juniperus.
- Gymnosporangium sabinæ** (Dicks.) Wint. See Juniperus.
- Gymnosporangium yamadæ** Miy. See Juniperus.
- Hadrotrichum piri** Montem. Small gray circular, sometimes confluent, leaf spots on *Pyrus* sp. in Italy.
- Helicobasidium mompa**. See Morus.
- Helicobasidium tanakæ** Miy. See Morus.
- Labrella piricola** Bres. and Sacc. On leaves of *P. communis* and *Malus* in Germany.
- Laestadia parmensis** Pass. On leaves of *P. communis* in Italy.
- Ochropsora sorbi** Diet. See Sorbus.
- Oospora piricola** Nob. Gray to white patches on leaves and branches of *Pyrus* in France.
- Phacidia discolor** (Mont. and Sacc.) A. Poteb. See Malus.
- Phyllosticta briardi** Sacc. See Malus.
- Phyllosticta mali** Prill. and Delacr. See Malus.
- Phyllosticta piricola** Sacc. and Speg. On leaves of *Pyrus* in Europe.
- Phyllosticta tirolensis** Bub. Small subcircular to irregular gray leaf spots on *P. communis* in Austria.
- Phytophthora syringæ** Kleb. See Syringa.
- Plectodiscella piri** Wor. Circular to oval grayish-white leaf spots on *P. communis* and *Malus* sp. in southern Russia.
- Ramularia magnusiana** Sacc. On leaves of *Pyrus* in Belgium.
- Septoria nigerrima** Fckl. On leaves of *P. communis* in Yugoslavia, France, and Italy. Reported from New York.
- Septoria ralszii** B. and Br. Black irregular spots on fruit of *Pyrus* in Great Britain, with subsequent rot.
- Stagonospora mali** Delacr. On leaves of *Pyrus* and *Malus* in France.
- Stereum purpureum** Pers. See Prunus.
- Taphrina piri** S. Kus. Circular to irregular yellow-green leaf spots on which yellowish or white scurfs appear. These areas may spread over most of the leaf surface and cause distortion. On *P. miyabei* in Japan.
- QUAMOCLIT**. STAR GLORY. Annual or perennial twining vines.
- Phyosticta quamoclit** Thuem. Dull-yellow to brown irregular leaf spots with gray margins on *Q. coccinea* in Portugal.
- QUERCUS**. OAK. Ornamental and timber trees.
- Coccochorella quercicola** (P. Henn.) v. Hoeh. Shiny black circular stromata on leaves of *Q. thalassica* in Japan.
- Coccosdiscus quercicola** P. Henn. Black stromata on leaves of *Q. thalassica* in Japan.
- Cocchohora kusanoi** (P. Henn.) v. Hoehn. Irregular black shiny stromata on leaves of *Q. glauca* in Japan.
- Cocchoidea quercicola** P. Henn. and Shir. Small black stromata on leaves of *Q. glabra* and *Q. glauca* in Japan.
- Coryneum foliicolum** Fckl. See Crataegus.
- Coryneum umbonatum** Nees. Kills twigs and branches of *Q. castaneaefolia* in France.
- Cystotheca wrightii** B. and C. Superficial brown perithecia on leaves of *Q. acuta*, *Q. glauca*, *Q. myrsinaefolia*, and *Q. vibrayeana* in Japan.
- Dermatea cinnamomea** (Pers.) Rehm. Wound parasite on *Quercus* spp. in Europe.
- Diaportha taleola** (Fr.) Sacc. On branches of *Quercus* spp. in Europe.
- Exoascus kruckii** Vuill. Causes a witches'-broom effect on *Q. ilex* in France and Italy.
- Gloeosporium shiraianum** Syd. Circular ochraceous to brown leaf spots with pale centers on *Q. glandulifera* in Japan.
- Gloeosporium umbrinellum** B. and Br. Small irregular brown leaf spots on *Q. robur*, *Q. sessiliflora*, and *Quercus* sp. in Russia, Switzerland, Great Britain, and Germany.
- Gnomonia quercus-ilicis** Berl. On leaves of *Quercus* spp. in Italy.
- Guignardia contecta** (Desm.) Lind. Subcircular to irregular pale-rufous leaf spots with brown margins on *Q. coccifera* in France.
- Guignardia diffusa** (Crié) Sacc. and Trott. See Castanea.
- Guignardia punctoidea** (Cke.) Schroet. On branches of *Quercus* sp. in northern Europe.
- Hadronema orbiculare** Syd. Small circular, velvety fungus patches on leaves of *Q. glauca* and *Q. luzoniensis* in Japan and the Philippines.
- Helicobasidium mompa** Tan. See Morus.
- Heterosporium proteus** Starb. Brown sunken leaf spots on *Quercus* sp. in Sweden.
- Hydnum obrutans** Burt. Causes heart rot of *Quercus* sp. in Java.
- Japonia quercus** v. Hoehn. On leaves of *Q. glauca* in Japan.
- Laestadia cerris** Pass. Pale-brown leaf spots on *Q. cerris* in Italy.
- Laestadia cookeana** (Auersw.) Sacc. On leaves of *Q. ilex* and *Q. robur* in Europe.
- Lasioderma flavovirens** Dur. and Mont. On leaves of *Quercus* spp. in Spain and Algeria.
- Macrophoma cylindrospora** (Desm.) Berl. and Vogl. See Hedera.
- Macrophoma fusispora** Bub. Subcircular to elongate whitish leaf spots on *Q. conferta* in Hungary.
- Macrophoma mexicana** Sacc. On leaves of *Q. glauca* in Mexico.
- Melogramma henriquetii** Br. and Cav. On *Q. suber* in southern Europe.
- Microporella quercus** v. Hoeh. On leaves of *Q. glauca* in Japan.
- Microstoma album** (Desm.) Sacc. and var. **japonicum** P. Henn. On leaves of *Q. cerris*, *Q. glauca*, *Q. grosseserrata*, and *Q. sessiliflora* in Europe, Japan, Argentina, and the Union of South Africa.
- Monochaetia pachyspora** Bub. Subcircular to irregular pale grayish-yellow leaf spots on *Q. ilex* in Austria.
- Mycoderma cuticularia** Duc. Brown leaf spots on *Q. suber* in France.
- Phoma cupulicola** Gz. Frag. On *Quercus* sp. in Spain.
- Phyllohendersonia foliorum** (Fckl.) Tass. On leaves of *Q. ilex* and *Quercus* sp. in Italy, France, and Germany.
- Phyllosticta associata** Bub. On leaves of *Q. robur* (*Q. pedunculata*) in Austria.
- Phyllosticta bresadoleana** Bub. and Kab. Brown or yellow-gray leaf spots with narrow brown-purple margins on *Q. pubescens* in Austria.
- Phyllosticta globulosa** Thuem. Irregular gray-white leaf spots on *Q. robur* (*Q. pedunculata*) in Austria.

**QUERCUS**—Continued.

- Phyllosticta hranicensis** Petr. On leaves of *Q. robur* in Bohemia.  
**Phyllosticta ilicicola** Pass. Circular brown, then ashen-gray, leaf spots on *Q. ilex* in France and Italy.  
**Phyllosticta ilicina** Sacc. Pale-brown leaf spots on *Q. ilex* in Italy, Portugal, and Austria.  
**Phyllosticta iliciseda** Sacc. Broad, irregular leaf spots with black borders on *Q. ilex* in Italy.  
**Phyllosticta macrocarpae** Monte. Brown irregular leaf spots on *Q. macrocarpa* in Italy.  
**Phyllosticta phaeospora** Scalia. On leaves of *Q. ilex* in Italy.  
**Phyllosticta quercicola** Oud. On leaves of *Quercus* sp. in northern Europe.  
**Phyllosticta quercus-cocciferae** Bub. Circular to irregular white leaf spots on *Q. coccifera* in Yugoslavia.  
**Phyllosticta roboris** Oud. Small angular, often confluent, pale-brown leaf spots on *Q. robur* in Bohemia and Holland.  
**Phyllosticta shiraiana** Syd. Circular ochraceous to white leaf spots on *Q. acuta* and *Q. glauca* in Japan.  
**Flowrightia noxia** (Ruhl.) Sacc. See *Castanea*.  
**Rosellinia quercina** Hartig. Attacks roots of young *Quercus* seedlings in nurseries in Germany. Infected plants show an unhealthy browning of the foliage, followed by withering of young shoots and ultimate death.  
**Sclerotinia candolleana** (Lév.) Fekl. Leaves of *Quercus* and *Castanea* in Great Britain, France and Germany turn brown and fall prematurely due to this fungus.  
**Septoria dryophila** Sacc. White leaf spots with narrow dark margins on *Q. ilex* in Italy.  
**Septoria dubia** Sacc. and Syd. Small yellowish leaf spots with red margins on *Q. pedunculata* and *Q. robur* (*Q. pubescens*) in France, Russia, Yugoslavia, and Germany.  
**Septoria quercicola** Sacc. White leaf spots on *Q. humilis*, *Q. pedunculata*, and *Q. sessiliflora* in France, Italy, and Portugal. Reported from Minnesota.  
**Septoria quercina** Desm. Small circular white leaf spots on *Q. robur*, *Q. sessiliflora*, and *Quercus* sp. in Japan, Russia, France, Italy, Denmark, Austria, and Germany.  
**Septoria quercus** Thuem. Subcircular ochraceous leaf spots on *Q. pedunculata* and *Q. sessiliflora* in Portugal.  
**Sphaerotheca kusanoi** P. Henn. and Shir. Powdery mildew on leaves of *Q. glandulifera* in Japan.  
**Trabutia konzattiana** Sacc. Reddish-brown circular, sometimes confluent, leaf spots on *Quercus* sp. in Mexico.  
**Typhulochaeta japonica** Ito and Hara. On leaves of *Q. glandulifera* in Japan.  
**Uncinula septata** Salm. Powdery mildew on leaves of *Q. glandulifera* in Japan.  
**Uredo quercus** Brond. Small circular rust pustules on lower leaf surfaces of *Quercus* spp. in Europe.  
**Yoshinagaia quercus** P. Henn. Black subcarbonaceous stromata on upper leaf surfaces of *Q. glauca* in Japan.  
**Yoshinagella japonica** v. Hoeh. Irregular black stromata on upper leaf surfaces of *Q. glauca* in Japan.

**RADICULA**. HORSE-RADISH. WATER CRESS. (Including *Roripa* and *Armoracia*.)

- Ascochyta armoraciae** Fekl. Scattered pale-brown leaf spots on *Armoracia rusticana* (*R. armoracia*) in Europe.  
**Ascochyta rusticana** Kab. and Bub. Ochraceous or brown leaf spots varying in size from 0.5 to 2 centimeters, on *R. armoracia* in Bohemia.  
**Peronospora nasturtii aquatici** Gäum. Downy mildew on leaves of *Sisymbrium nasturtium-aquaticum* (*R. nasturtium-aquaticum*) in Germany.  
**Phyllosticta armoraciae** Cke. Circular to angular dirty-white leaf spots with narrow brown margins on *R. armoracia* in Great Britain.  
**Ramularia cochleariae** Cke. See *Barbarea*.  
**Septoria armoraciae** Sacc. Irregular ochraceous leaf spots on *R. armoracia* in Italy.

**RANDIA**. Erect or climbing tropical trees or shrubs.

- Aecidium abscondens** Arth. Yellow rust pustules on leaves of *R. aculeata* in Porto Rico and Costa Rica.  
**Aecidium pulverulentum** Arth. Leaf rust on *Randia* sp. in Mexico.  
**Aecidium randiae** P. Henn. Yellow rust pustules on dull-brown, swollen leaf spots on *Randia* sp. in Brazil.  
**Corticium salmonicolor** B. and Br. See *Citrus*.  
**Endophyllum griffithsiae** Rac. Yellow rust sori on circular leaf spots on *R. scandens*, *R. sinensis*, and *Pavetta indica* in Java and China.  
**Lembosia philippinensis** Syd. On leaves of *Randia* sp. in the Philippines.  
**Pestalozzia pauciseta** Sacc. Gray leaf spots on *R. reticulata* in the Philippines.  
**Taphrina randiae** Rehm. Elliptical black, swollen leaf spots on *Randia* sp. in Brazil.  
**Trabutia randiae** (Rehm.) Theiss. and Syd. and var. *aculeatae* Ferd. and Winge. Black stromata on leaves of *R. aculeata* and *R. pubescens* in Bolivia and the American Virgin Islands.

**RANUNCULUS**. (Including *Ficaria*.) BUTTERCUP. Crowfoot. Annual and perennial herbs.

- Aphelenchus olesistus** Ritz. Bos. See *Begonia*.  
**Ascochyta carinthiaca** Jaap. On leaves of *R. thora* in Austria.  
**Cercospora ranunculi** Jaap. Circular to elongate brown leaf spots on *R. muricatus* in Dalmatia.  
**Cylindrosporium ficariae** Berk. On leaves of *R. ficaria* in France and Great Britain.  
**Cylindrosporium ranunculi** Sacc. and f. *sclerati* P. Brun. On leaves of *R. acris*, *R. bulbosus*, and *R. scleratus* in France and Italy.  
**Entyloma ameghinii** Speg. Leaf smut on *R. cymbalaria* in Patagonia.  
**Fabraea litigiosa** (Rob. and Desm.) Sacc. On leaves of *Ranunculus* spp. in Alaska, Europe, and north Africa.  
**Fusidium eburneum** Schroet. Subcircular, often confluent, white leaf spots with yellow centers on *R. acris* and *R. repens* in Siberia and Austria.  
**Heterosporium ferox** Bub. On leaves of *R. arvensis* in Bohemia.  
**Leptosphaeria thorae** Jaap. On leaves of *R. thora* in Austria.  
**Ovularia decipiens** Sacc. On leaves of *R. acris* in Russia, Dalmatia, and Italy. Reported from New York.  
**Peronospora alpicola** Gäum. Downy mildew on leaves of *R. aconitifolius*, *R. pyrenaicus*, and *R. sequieri* in France, Switzerland, Austria, and Germany.  
**Peronospora ficariae** (Nees.) Tul. As above on *R. ficaria* in Europe.  
**Peronospora gigantea** Gäum. Downy mildew on leaves of *R. lingua* in Denmark and Russia.  
**Peronospora glacialis** (Blytt.) Gäum. Downy mildew on leaves of *R. glacialis* in Switzerland and Norway.  
**Peronospora illyrica** Gäum. As above, on *R. illyricus* in Russia.  
**Phyllosticta ficariae** Maire. On leaves of *R. calthaeifolius* in French north Africa.  
**Phyllosticta pygmaea** Allesch. On leaves of *R. pygmaeus* in Greenland.

**RANUNCULUS**—Continued.

- Phyllosticta ranunculi** (Fckl.) Sacc. On leaves of *R. acris* in Germany.
- Phyllosticta ranuncolorum** Sacc. and Speg. Dull-brown leaf spots on *R. repens* in Italy.
- Physoderma vagans** Schroet. Small tubercles on leaves and stems of *R. acris*, *R. flammula*, *R. repens*, *Oenanthe phellandrium*, *Potentilla anserina*, *Selinum lineare*, *Silaus* sp., and *Sium latifolium* in France and Germany.
- Pseudopeziza ranunculi** (Wallbr.) Fckl. Black circular fruiting bodies in dark-brown sunken leaf spots on *R. cassubicus*, *R. chaerophyllus*, *R. nemorosus*, and *R. polyanthemos* in Belgium, France, and Germany.
- Puccinia andina** Diet. and Neg. Leaf rust on *R. peduncularis* in South America.
- Puccinia blyttiana** Lagh. Brown rust pustules on leaf blades and petioles of *R. alpestris* and *R. auricomus* in Norway and Switzerland.
- Puccinia gibberulosa** Schroet. Powdery brown rust pustules on leaf blades and petioles of *R. lancipetalus* and *Ranunculus* sp. in France and Argentina.
- Puccinia magnusiana** Koern. See Phragmites.
- Puccinia nubigena** Speg. Rust on leaf blades and petioles of *Ranunculus* sp. in Argentina.
- Puccinia ustalis** Berk. Dark-brown rust pustules on *R. hirtellus*, *R. pulchellus*, and *R. songaricus* in Mongolia, Turkestan, and India.
- Ramularia acris** Lindr. Large irregular yellow-brown to gray-brown leaf spots on *R. acris* in Finland and Sweden.
- Ramularia aequivoca** (Ces.) Sacc. On leaves of *R. auricomus*, *R. cassubicus*, *R. lanuginosus*, *R. polyanthemos*, *R. repens*, and *R. trachycarpus* in Europe. Reported from Wisconsin.
- Ramularia lapponica** Lindr. Irregular gray or dark-brown leaf spots on *R. lapponicus* in Finland.
- Ramularia repentis** Oud. Elliptical to irregular leaf spots on *R. repens* in Holland.
- Ramularia scelerata** Cke. Elongate brown spots on lower leaves of *R. sceleratus* in Great Britain.
- Ramularia scerbica** Ranoj. Irregular brown leaf spots on *R. montanus* in Yugoslavia.
- Sclerotinia ficariae** Rehm. On leaves of *R. ficaria* in Italy.
- Septoria cymbalarina** Thuem. On leaves of *R. cymbalaria* in Siberia.
- Septoria ficariae** Desm. Ashen leaf spots on *R. ficaria* and *R. vernus* in Europe.
- Septoria oreophila** Sacc. Small white leaf spots on *R. aconitifolius* in Italy.
- Septoria polaris** Karst. White leaf spots on *R. flammula* and *R. parviflorus* in Great Britain and Scandinavia.
- Septoria ranunculacearum** Lévy. Brown leaf spots on *R. acris*, *R. cymbalaria*, and *R. pusillus* in France, Italy, and Siberia.
- Septoria ranunculi** West. Circular greenish-yellow leaf spots on *R. sceleratus* in Belgium.
- Synchytrium andinum** Lagerh. Red-brown warty areas on leaves of *Ranunculus* sp. in Ecuador.
- Uromyces dactylidis** Otth. See Dactylis.
- Uromyces festucae** Syd. See Festuca.
- Uromyces ficariae** (Schum.) Lévy. Leaf rust on *R. ficaria*, *R. grandiflora*, and *R. vernus* in Europe.
- Uromyces fischerianus** E. Mayor. Yellow-brown rust pustules on leaf blades and petioles of *R. glacialis* in Switzerland.
- Uromyces poae** Rabh. See Poa.
- Uromyces ranunculi-distichophylli** Semad. Leaf rust on *R. parnassifolius* in Switzerland.
- Uromyces ranunculi-festucae** Jaap. See Festuca.
- RAPHANUS**. RADISH. Vegetable and forage plants.
- Moniliopsis aderholdi** Ruhl. See Solanum.
- Pseudomonas destructans** Potter. See Brassica.
- Puccinia isiacae** (Thuem.) Wint. See Arundo.
- RAUWOLFIA**. Ornamental trees or shrubs.
- Aecidium rauwolfiae** P. Henn. Leaf rust on *Rauwolfia* sp. in the Congo.
- RAVENALA**. TRAVELER'S-TREE. Large *Musa*-like plants.
- Catacauma ravenalae** (Pat. and Har.) Theiss. and Syd. Black stromata on leaves of *R. madagascariensis* in Madagascar.
- Zukalia stuhlmanniana** P. Henn. See Palmae.
- RENANTHERA**. Tall climbing epiphytic orchids. See Orchidaceae.
- RENEALMIA**. Perennial aromatic herbs.
- Catacauma renealmiae** (Rehm.) Theiss. and Syd. Circular black stromata on brown leaf spots on *Renealmia* sp. in Brazil.
- Endodothella renealmiae** (Rehm.) Theiss. and Syd. Circular to irregular black stromata on leaves of *Renealmia* sp. in Brazil.
- Septoria renealmiae** Tassi. Large brown leaf spots on *R. cinnamomea* in Italy.
- RESEDA**. MIGNONETTE. Garden annuals.
- Peronospora crispula** Fckl. Downy mildew on leaves of *R. lutea* and *R. luteola* in France, Denmark, Switzerland, and Germany.
- Phyllosticta resedae** Petch. On leaves of *R. odorata* in Ceylon.
- RETINOSPORA**. See Chamaecyparis.
- RHAMNUS**. BUCKTHORN. Woody plants grown for their foliage and some species for their attractive fruit and medicinal bark.
- Ascochyta frangulina** Kab. and Bub. On leaves of *R. cathartica* and *R. frangula* in Russia and Bohemia.
- Endophyllum elegans** (Diet.) Pole-Evans. Leaf rust on *R. prinoides* in the Union of South Africa.
- Mycosphaerella vogelii** Syd. On leaves of *R. cathartica* in Germany.
- Phyllosticta advena** Sacc. Oval to indefinite ashen-white leaf spots with dull-brown margins on *R. (Sageretia) corymbosa* in Italy.
- Phyllosticta alaterni** Pass. Circular grayish-white leaf spots with dark-brown margins on *R. alaternus* and *R. frangula* in France and Italy.
- Phyllosticta cathartici** Sacc. Subcircular ochraceous leaf spots with dull-red margins on *R. cathartica* in Russia, Italy, Denmark, and Austria.
- Phyllosticta desmazieri** Tass. On leaves of *R. cathartica* in France.
- Phyllosticta jahniana** Petr. and Sacc. Subelliptical to irregular black leaf spots on *R. cathartica* in Bohemia.
- Phyllosticta osteospora** Sacc. See Morus.
- Phyllosticta rhamnicola** Desm. Circular ashen leaf spots on *R. alpina* in France and Belgium.
- Phyllostictiella rhamnigna** (Sacc.) Tass. Subcircular dull-white leaf spots with dull-brown margins on *R. alaternus* and *R. cathartica* in Portugal, Italy, and France.
- Puccinia himalensis** (Barcl.) Diet. Brown to black rust pustules on leaves and twigs of *R. dahurica* in India. Aecial stage on leaf blades and sheaths of *Brachypodium silvaticum*.

**RHAMNUS**—Continued.

**Puccinia schweinfurthii** (P. Henn.) P. Magn. Brown crustlike rust pustules deforming the leaves of *R. prinoides* and *R. stado* in Abyssinia and east Africa.

**Ramularia alaterni** Thuem. Large gray spots on leaves of *R. alaternus* in Spain and France.

**Rostrupia schweinfurthii** P. Henn. On leaves of *Rhamnus* sp. in Abyssinia.

**Septoria alaterni** Pass. and var. *hispanica* Gz. Frag. Gray leaf spots on *R. alaternus* in Spain and Italy.

**Septoria cathartica** Pass. Irregular gray leaf spots on *R. cathartica* and *R. saxatilis* in Denmark and Italy.

**Septoria frangulae** Guep. Small dark-rufous leaf spots on *R. frangula* in France, Bohemia, Silesia, Poland, and Germany.

**RHAPHIDOPHYLLUM**. See *Palmae*.**RHAPONTICUM**. See, also, *Centaurea*.

**Puccinia rhapontici** Syd. Powdery dark-brown rust pustules on leaves of *R. acaule* and *R. pusillum* in north Africa.

**RHEUM**. RHUBARB. Herbs, grown for their foliage and edible leaf stocks.

**Bacterium rhaponticum** Millard. This bacterium is reported as the cause of crown rot of *R. rhaponticum* in Great Britain. A soft brown rot attacks the crowns and sheathing leaves are discolored and swollen at the base where joined to the crowns. The leaves turn dark-brown and in the absence of terminal buds only spindling shoots are produced from lateral buds.

**Cercospora rhei** Grog. On leaves of *R. officinale* in France.

**Peronospora jaapiana** Magn. Downy mildew forming large irregular gray patches on leaves of *R. officinale*, *R. palmatum*, and *R. undulatum* in Switzerland and Germany.

**Ramularia russi** Allesch. Circular, then confluent, red-brown leaf spots on *R. undulatum* and *Rheum* (cult.) in Russia, Denmark, and Germany.

**Uredo rhei-undulati** Diet. Brown leaf rust on *R. undulatum* in Japan.

**RHIPSALIS**. Epiphytic succulent plants.

**Phaeospora caticola** Stevens. On *R. cassytha* in Porto Rico.

**RHODODENDRON**. (Including *Azalea*.) Woody plants cultivated for their flowers and foliage.

**Ascochyta rhododendri** Lind. Angular brown leaf spots on *Rhododendron* (cult.) in Denmark.

**Cercospora handelii** Bub. Irregular, then confluent, ochraceous spots destroying leaves of *R. ponticum* in Asia Minor.

**Chrysomyxa dietelii** Syd. Yellow rust pustules on leaves of *R. arboreum* in India.

**Chrysomyxa expansa** Diet. Yellow rust pustules on red-brown leaf spots on *R. metternichii* in Japan.

**Chrysomyxa himalensis** Barcl. Rust on leaf blades and petioles of *R. arboreum*, *R. campanulatum*, and *R. hodgsonum* in India.

**Chrysomyxa rhododendri** (DC.) De By. Yellow to red-brown rust pustules on *R. brachycarpum*, *R. dauricum*, *R. ferrugineum*, *R. hirsutum*, *R. intermedium*, *R. kotschyum*, *R. myrtifolium*, and *R. suave* in Japan, Siberia, and Europe. The acial stage occurs on *Picea abies*, *P. obovata*, and *P. pungens*.

**Exobasidium butleri** Syd. Fleshy growths on lower leaf surfaces of *R. arboreum* in India.

**Exobasidium hemisphaericum** Shir. Fleshy pyriform to globose galls on lower leaf surfaces of *R. metternichii* and *R. pentamerum* in Japan.

**Exobasidium japonicum** Shir. Fleshy, smooth, globose, greenish, then reddish-white, galls on leaves of *R. indicum* in Japan.

**Exobasidium vexans** Mass. See *Thea*.

**Exobasidium yoshinagai** P. Henn. Circular to effuse dull-brown leaf spots on *R. tosaense* and *Azalea* sp. in Japan.

**Exobasidium zeylanicum** Petch. Large spherical white or greenish-white, then powdery white galls on lower leaf surfaces of *R. arboreum* in Ceylon.

**Gloeosporium rhododendri** Br. and Cav. Large irregular blotches on leaves of *R. ponticum* in Italy and Australia.

**Gloeosporium succineum** Sacc. Anthracnose on leaves of *R. chrysanthemum* in Siberia.

**Leptosphaeria rhododendri** P. Henn. Large brown, then gray, leaf spots on *R. falconeri* in Germany.

**Melasmia rhododendri** P. Henn. and Shir. On leaves of *R. indicum*, *R. kaempferi*, and *R. tschonoskii* in Japan.

**Mycosphaerella occulta** Bub. Circular gray leaf spots on *R. ponticum* in Bohemia.

**Naemosphaera japonica** Sacc. and Syd. On branches of *R. indicum* in Japan.

**Phacidium falconeri** P. Henn. Gray leaf spots with broad red-brown margins on *R. falconeri* in Germany.

**Phyllohendersonia bicolor** (Pat.) Tass. On *Rhododendron* sp. in China.

**Phyllohendersonia rhododendri** (Thuem.) Tass. On leaves of *R. hirsutum* in Italy.

**Phyllosticta berolinensis** P. Henn. Dull-brown to gray leaf spots on *R. falconeri* in Germany.

**Phyllosticta cunninghami** Allesch. On leaves of *R. cunninghami* in Germany.

**Phyllosticta falconeri** P. Henn. Dull-brown to gray circular leaf spots with dark-red margins on *R. falconeri* in Germany.

**Phyllosticta rhododendricola** Brun. Large oblong to irregular ashen-gray leaf spots with narrow brown margins on *Rhododendron* sp. in France.

**Phyllosticta rhododendri-flavi** Bub. and Kab. Large circular to irregular red-brown, then dark-brown, leaf spots on *R. flavum* in Bohemia.

**Phyllosticta rhodorae** (Cke.) Tass. On leaves of *Rhododendron* sp. in England.

**Phyllosticta saccardoi** Thuem. Brown leaf spots on *R. ponticum* in Russia, France, and Portugal. Reported from New York on *Rhododendron* sp.

**Physalospora alpina** Speg. On leaves of *R. ferrugineum* in Italy.

**Physalospora rhododendri** Naum. Die-back of twigs and browning of leaves of *Rhododendron* sp. in Germany.

**Placosphaeria rhododendri** P. Henn. Circular black stromata on leaves of *Rhododendron* sp. in Japan.

**Puccinia rhododendri** Fekl. Leaf rust on *R. ferrugineum* in Austria.

**Rhytisma rhododendri** Fr. Black irregular areas on leaves of *Rhododendron* sp. in Siberia.

**Sclerotinia rhododendri** Fisch. Forming sclerotia in fruit of *R. ferrugineum* and *R. hirsutum* in Europe.

**Septoria azaleae** Vogl. Die-back of twigs and browning of leaves of *Azalea indica* in Italy and Germany.

**Septoria azaleae-indicae** Maubl. Circular white leaf spots with black margins on *A. indica* and *Azalea* sp. in Brazil and Germany.

**RHODOMYRTUS**. ROSE MYRTLE. Trees or shrubs native to southern Asia.

**Helminthosporium rhodomirti** Syd. On leaves of *R. tomentosa* in China.

**RHODOTYPOS**. JET BEAD. Shrubs grown chiefly for their large white flowers and bright-green foliage

**Septoria rhodotypi** Höll. Irregular rufous leaf spots on *R. kerrioides* in Hungary.

- RHUS.** SUMAC. Ornamental woody plants, some species of economic importance.
- Aecidium foetidum** Diet. Leaf rust on *Rhus* sp. in Japan.
- Aecidium inercassatum** Syd. Leaf rust on *R. sylvestris* in Japan.
- Bagnisiopsis rhoïna** Syd. and Hara. Dull-black circular to elliptical stromata on leaves of *R. sylvestris* in Japan.
- Colletotrichum rhoïnum** F. Tassi. Anthracnose attacking young shoots and leaf blades and petioles of *R. javanica* (*R. semi-alata*) and *R. verniciflua* in Japan. The lesions are long, elliptical, black, and sunken.
- Exobasidium hesperidium** Maire. Leaf spots, brown-orange on the lower surfaces, yellow-green, then dark-brown, on the upper, depressed and causing rolling of the leaves of *R. oxyacantha* in Algeria.
- Helicobasidium mompa** Tan. See *Morus*.
- Phyllachora marginalis** Pat. Irregular black stromata on lower leaf surfaces of *Rhus* sp. in Ecuador.
- Phyllosticta glabra** Brun. Brown angular spots on leaves of *R. glabra* in France.
- Phyllosticta rhoïis** West. Angular to irregular pale brown leaf spots on *R. coriaria* and *R. cotinus* in Denmark, France, and Belgium.
- Pileolaria dicteliana** Syd. Leaf rust on *R. hypoleuca* in China.
- Septorhema henningsii** Bresad. On leaves of *R. abyssinica* in Abyssinia.
- Septoria acuriana** P. Henn. Yellow leaf spots on *R. retinorrhæa* in Abyssinia.
- Septoria cotini** C. Mass. Irregular, then confluent, reddish leaf spots on *R. cotinus* in Italy.
- Septoria rhoïis** Lév. Dull-brown leaf spots on *R. hirta* (*R. typhina*) in Russia.
- Uncinula vernicifera** P. Henn. Powdery mildew on leaves of *R. javanica* (*R. semi-alata*) and *R. verniciflua* in Japan.
- Uredo rhoïna** Syd. Yellow-brown leaf rust on *Rhus* sp. in central Africa.
- Uromyces barbeyanus** P. Henn. Dark-brown rust pustules on leaves of *R. falcata* in Abyssinia.
- Uromyces klugkistianus** Diet. Leaf rust on *R. javanica* and *R. succedanea* in Japan.
- Uromyces propinquus** Syd. Powdery brown leaf rust on *R. mollis* in Mexico.
- Uromyces shiraianus** Diet. and Syd. Powdery brown rust pustules on leaves of *R. silvatica*, *R. succedanea*, *R. sylvestris*, and *R. trichocarpa* in Japan.
- RHYNCHOSIA.** **DOLICHOLUS** Ag. ROSARYBEAN. Twining, prostrate, rarely erect leguminous herbs and shrubs.
- Aecidium erioseomatis** P. Henn. Leaf rust on *Rhynchosia* and *Eriosema* sp. in Brazil.
- Uredo pamparum** Speg. Powdery dark-ochraceous rust pustules on leaves of *Rhynchosia* sp. in Argentina.
- Woroninella dolichi** (Cke.) Syd. See *Dolichos*.
- RIBES.** CURRANT. (Including *Grossularia*, GOOSEBERRY.) Shrubs cultivated for fruit and ornament.
- Alternaria ribis** Bub. and Ranoj. Effuse gray leaf spots with brown margins on *R. rubrum* in Yugoslavia.
- Aphelenchus ribes** (Taylor.) Goodey. This nematode attacks the buds of *R. nigrum* in Great Britain. The buds are destroyed and adjacent unfolding leaves may also be attacked.
- Ascochyta monachorum** Bub. On *Grossularia* in Bulgaria.
- Ascochyta ribesia** Sacc. and Fautr. Numerous circular to angular white leaf spots with dull-purple margins on *R. nigrum* and *Grossularia* sp. in Great Britain, France, and Austria.
- Ascochyta ribis** Massal. On *R. aureum* and *R. rubrum* in Italy and Russia.
- Bacterium** sp. A blossom-blight of *Grossularia* sp. due to a bacterium is reported from Great Britain.
- Cercospora magellanica** Speg. Small reddish leaf spots on *R. magellanicum* in Chile.
- Colletotrichum grossulariae** Jacz. Brown spots on fruit of *Grossularia* sp. in Russia and Italy.
- Coniothyrium vagabundum** Sacc. Said to cause premature fall of leaves of *Grossularia* sp. in Europe.
- Cronartium ribicola** Fisch. See *Pinus*.
- Cytosporina ribis** Magn. Causes a dying back and stem cankering of *R. grossularia*, *R. nigrum*, and *R. rubrum* in England and Holland.
- Dothiorella vinosa** Marchal. See *Prunus*.
- Dothidella ribesia** (Pers.) Theiss. and Syd. On twigs of *R. nigrum*, *R. rubrum*, *R. vulgare*, and *Ribes* sp. in Great Britain, middle Europe, and Alaska.
- Helicobasidium tanakae** Miy. See *Morus*.
- Marsonia grossulariae** Oud. Black spots on branches of *Grossularia* sp. in Europe.
- Melampsora ribesii-purpureae** Kleb. See *Salix*.
- Melampsora ribesii-viminalis** Kleb. See *Salix*.
- Phyllosticta pallidocarpa** Bub. and Sereb. On leaves of *R. aureum* in Russia.
- Phyllosticta ribesicida** Speg. Definite circular white leaf spots on *R. rubrum* in Argentina.
- Phyllosticta ribicola** Sacc. Large yellowish leaf spots on *R. alpinum*, *R. aureum*, *R. nigrum*, *R. rubrum* and *Grossularia* sp. in Europe.
- Phyllosticta ribiseda** Bub. and Kab. Circular, zoned, red-brown, to gray leaf spots on *R. rubrum* in Bohemia.
- Phyllosticta ribis-rubri** Vogl. Circular, then confluent, chestnut-brown leaf spots on *R. rubrum* in Italy.
- Puccinia depressa** Diet. and Neg. Brown rust pustules on leaf blades and petioles of *R. glandulosum* in Chile.
- Puccinia jaffueliana** Speg. Leaf rust on *R. punctatum* in Chile.
- Puccinia ribesii-caricis** Kleb. Brown leaf rust on *R. aureum* in Europe.
- Puccinia ribis-japonici** P. Henn. Brown rust pustules on leaves of *R. japonicum* in Japan.
- Septoria grossulariicola** C. Mass. Irregular brown leaf spots on *Grossularia* sp. in Italy.
- Stereum purpureum** Pers. See *Prunus*.
- Venturia grossulariae** (Awd. and Fleisch.) Sacc. On leaves of *Grossularia* sp. in Germany.
- RICINUS.** CASTOR BEAN. Tall annuals, or small trees in the Tropics.
- Colletotrichum ricini** Bub. and Frag. Anthracnose on leaves of *R. communis* in Spain.
- Macrosporium cavaræ** Parisi. Circular dark-yellow to brown spots on leaves and cotyledons preventing development of plants of *R. communis*, *R. gibsoni*, and *R. viridis* in Italy.
- Melampsorella ricini** (Biv.-Bern.) Det. Powdery orange-yellow rust pustules on small circular yellow leaf spots on *R. communis* and *R. spectabilis* in India, Ceylon, Madeira, north, south, and east Africa, and southern Europe.
- Phyllosticta ricini** Rost. Subcircular brown leaf spots with purple borders on *R. communis* in Denmark.
- Physalospora propinqua** Sacc. On *R. communis* in the Philippines.
- Phytophthora melongenæ** K. Saw. See *Solanum*.

**RICINUS**—Continued.

**Phytophthora parasitica** Dastur. Circular dull-green infection areas over both surfaces of cotyledons, destroying them and spreading down the stems causing damping off. Irregular spots occur on older leaves. On *R. communis*, *Clarkia* sp., *Fagopyrum* sp., *Gilia* sp., *Lycopersicum esculentum*, *Salpiglossis* sp., *Sesamum indicum*, *Solanum melongena*, *S. tuberosum*, and *Vinca rosea* in India.

A leaf-stalk rot of *Cocos nucifera* is attributed to this fungus in Jamaica. Dark-brown, somewhat sunken spots, one to several inches in diameter, occur on the leaf stalks causing a yellowing and wilting of the leaves, progressing successively until all are attacked and drooping. An internal brown rot of the leaves follows the spotting. The bud and central tissues as a rule are not affected. The larger nuts drop as a result of support from the leaves.

This species is now considered identical with *P. terrestria* Sherb. which occurs in the southern United States and the West Indies as the cause of a foot rot or crown rot of *Citrus*, *Lycopersicum*, and other economic plants.

**Sclerotinia ricini** Godf. This serious blight of *R. communis*, characterized by a rot of the various parts of the plant, was introduced into the United States on seeds during the late war. It also occurs in Ceylon and India.

**RIVINA**. ROUGE PLANT. Erect tropical herbs.

**Endophyllum rivinae** (B. and C.) Arth. Yellow to brown rust pustules on branches, inflorescences and leaf blades and petioles of *R. humilis* and *Trichostigma octandrum* in Cuba and Argentina.

**Phyllosticta rivinae** Speg. Circular grayish-white leaf spots on *R. humilis* in Argentina.

**Puccinia rivinae** (B. and C.) Speg. Brown rust pustules on stems and leaf blades and petioles of *R. humilis* and *Trichostigma octandrum* in Porto Rico, Cuba, St. Thomas, and Argentina.

**Septoria rivinae** Pat. Brown, then white, irregular leaf spots on *R. humilis* and *Trichostigma octandrum* in Guadeloupe.

**ROBINIA**. LOCUST. Timber and ornamental trees cultivated for flowers and graceful foliage.

**Ascochyta robiniae** Sacc. and Speg. Grayish-white leaf spots on *R. pseudoacacia* in Italy.

**Mycosphaerella robiniae** Siem. On leaves of *R. pseudoacacia* in Russia.

**Oidium orbiculare** Nannizzi. Powdery mildew on leaves of *R. hispida* in Italy.

**Phleospora robiniae** (Lib.) v. Hoeh. (*Septoria robiniae* Desm.) Leaf spot on *R. pseudoacacia* in Europe.

**Phyllosticta advena** Pass. Oval to vague ashen-white leaf spots on *R. corymbosa* and *R. pseudoacacia* in France, Italy, and Bohemia.

**Phyllosticta robiniae** Sacc. Irregular rufous leaf spots on *R. pseudoacacia* in Italy.

**Phytophthora fagi** R. Hart. See *Fagus*.

**Septoria curvata** (Rbh. and Braun.) Sacc. Circular to oblong brown leaf spots on *R. pseudoacacia* in Russia, Italy, and Germany. Reported from Ohio.

**ROLLINIA**. Tropical American fruit trees and shrubs.

**Ovulariopsis monospora** (Pass.) Sacc. and D. Sacc. Ashen leaf spots on *R. hispida* in Italy.

**Uredo cherimoliae** Lagerh. See *Annona*.

**ROMULEA**. Crocuslike bulbs.

**Phyllosticta romuleae** Gz. Frag. Leaf spots on *R. bifrons* in Spain.

**RONDELETIA**. Evergreen tropical trees and shrubs.

**Phyllosticta rondeletiae** Fl. Red, then ashen-white, leaf spots on *R. speciosa* in Italy.

**Uredo rondeletiae** Arth. and Holw. Leaf rust on *R. cordata* in Guatemala.

**RORIPA**. See *Radicula*.**ROSA**. ROSE. Ornamental shrubs.

**Aecidium rosae-abbyssinicae** P. Henn. Yellow rust pustules on leaves of *R. abyssinica* in Abyssinia.

**Apiospora rhodophila** Sacc. On branches and spines of *R. alpina* in Italy.

**Apiospora rosae** Oud. On branches of *R. canina* in Holland.

**Ascochyta rosicola** Sacc. This fungus produces more or less circular brown spots with distinct red margins on the upper surfaces of leaves of *R. mucosa* in Italy.

**Botryosphaeria dothidea** (Fr.) Ces. and de Not. (*B. diplobia* Moug.) This disease, sometimes known as brier scab, has been epidemic at times in England on certain varieties of roses. Large, slightly raised black scabs are formed on the stems, which crack more or less concentrically. The hosts are *R. arvensis*, *R. canina*, and *R. centifolia* in Great Britain, continental Europe, and South America.

**Catacauma dothidea** (Moug.) v. Hoeh. Black stromata on twigs of *Rosa* sp. in Albania and Yugoslavia.

**Cercospora hypophylla** Cav. Causes a leaf spot of *R. gallica* in Italy.

**Cercospora rosae** (Fckl.) v. Hoehn. Reported as causing a leaf spot of *R. agrestis*, *R. alpina*, *R. arvensis*, *R. indica*, and *R. pendulina* in Madeira, Italy, Switzerland, and Austria.

**Cercospora rosae-alpinae** C. Mass. Causes a spotting of the under sides of leaves of *R. alpina* in Italy.

**Coniothyrium wernsdorffiae** G. Koch. This species, which is said by several writers to differ from *C. fuckelii*, causes a serious canker disease of canes of *Rosa* spp. in Denmark, Austria, and Germany. Diseased areas are more or less oval to circular, frequently girdling the stem, and brown in color with golden-brown margins.

**Cryptosporium minimum** Lan. Forms yellow-gray to brown spots, 1 to 2 centimeters in diameter on stems of *R. multiflora* in Germany.

**Cryptostictis caudata** (Preuss.) Sacc. Said to form cankers on branches of *Rosa* spp. in Europe.

**Cryptostictis cynosbati** (Fckl.) Sacc. Found on fruit of *R. pimpinellifolia* in Germany.

**Diaporthe incarcerationata** (B. and Br.) Nke. (*Phoma incarcerationata* [Nke.] Sacc.) Forms dead areas on twigs of *R. banksiae*, *R. canina* and *R. centifolia* in Europe.

**Dicoccum rosae** Bon. Leaf spot of *Rosa* spp. in Europe.

**Didymella sepincolaeformis** (DeNot.) Sacc. Minute fruiting bodies on dead areas on branches of *R. alpina* in Italy.

**Diplodia rosarum** Fr. (*D. centrophila* Pass.) Attacks the branches and spines of *R. banksiae*, *R. canina*, and *R. centifolia* in France and Austria.

**Exosporium rosae** Fckl. Forms spots on the under sides of living leaves of *Rosa* sp. in Switzerland and Germany.

**Guignardia rosae** (Auersw.) Petr. (*Laestadia rosae* Auersw.) Brown spots on the upper surfaces of leaves of *R. canina*, *R. centifolia*, and *R. rubiginosa* in Germany.

**Gymnoconia rosae** (Barcl.) Liro. (*Puccinia rosae* Barcl.) Powdery red-brown to dark-brown rust pustules on both sides of the leaves and on young twigs. Since the mycelium is perennial in the twigs, inspection of dormant plants would fail to reveal the presence of the disease. The hosts are *R. acicularis*, *R. cinnamomea*, *R. macrophylla*, *R. pimpinellifolia*, *R. platyacantha*, and *R. rugosa* in northern Europe, Turkestan, Kamchatka, Caucasus, India, and Japan.

**Hendersonia canina** Brun. (*H. lichenicola* [Cda.] Fr.) On branches of *R. canina* in France.

**Hendersonia rosicola** Petch. On twigs of *Rosa* sp. in Ceylon.

**ROSA**—Continued.

- Hendersonia vulgaris** Desm. var. *rosae* Vesteg. Produces irregular spots on leaves of *R. alpina* and *R. pimpinellifolia* in Sweden.
- Kuehneola japonica** Diet. (*Phragmidium japonicum* Diet.) Brown rust pustules on both leaf surfaces of *R. lucia*, *R. multiflora*, and *R. wichuriana* in Japan.
- Kuehneola rosae** Sawada. A disease characterized by reddish-orange rust pustules on both leaf surfaces and on stipules, young twigs, and fruit of *R. indica* var. *formosana* in Formosa.
- Leptosphaeria coniothyrium** (Fckl.) Sacc. On *Rosa* sp. in Tasmania.
- Metasphaeria canina** Pass. (*M. constricta* Bres.) Attacks branches and spines of *R. canina* in Italy.
- Metasphaeria persistens** (B. and Br.) Sacc. On stems of *Rosa* sp. in England.
- Myosporium rosae** Fckl. Produces blackened areas on branches of *R. canina* in Austria and Germany.
- Phoma aculeorum** Sacc. Reported on twigs of *R. canina* in Denmark, France, Italy, Switzerland, Austria, and Germany.
- Phoma canina** P. Brun. (*P. sepincola* [Kichx.] Sacc.) On *R. canina* in France, Switzerland, Austria, and Germany.
- Phoma rubiginosa** Brun. On fruit of *R. canina* and *R. rubiginosa* in Denmark and France.
- Phomopsis incarcerata** v. Hoeh. (*Phoma incarcerata* Sacc.) On branches of *R. canina* and *R. banksiae* in Great Britain. See also *Diaporthe incarcerata*.
- Phomopsis rosae** Grove. (*Phoma rosae* Schulz. and Sacc.) On *R. canina* in Great Britain, Albania and Yugoslavia.
- Phragmidium butleri** Syd. Yellow to deep-brown or black rust pustules on the lower leaf surfaces of *R. macrophylla* in the Himalaya Mountains in India.
- Phragmidium devastatrix** Sor. Brown to black powdery rust pustules on both leaf surfaces and on petioles and young twigs of *R. lutea* in central Asia.
- Phragmidium egenulum** Syd. and Butl. Yellow or dark brown rust pustules on the leaves of *R. webbiana* in Kashmir (India).
- Phragmidium fusiforme** Schroet. (*P. rosae alpinae* Wint.) Powdery rust pustules on the under sides of leaves and on petioles and fruits. *P. rosae-acicularis* on *R. acicularis* in Japan is said to be identical. The hosts are *R. acicularis*, *R. alpina*, *R. canina*, *R. nipponensis*, *R. pseudulina*, and *R. silvatica* in Europe.
- Phragmidium rosae-lacerantis** Diet. Yellow rust pustules changing to brown on the lower sides of leaves of *R. lacerans* in Persia.
- Phragmidium rosae-moschatae** Diet. Yellow to brown rust pustules on the under surfaces of leaves of *R. moschata* and *R. webbiana* in India.
- Phragmidium rosae-multiflorae** Diet. Small yellow swellings occur on the petioles and larger leaf veins with yellow rust pustules on the lower leaf surfaces of *R. laevigata* and *R. multiflora* in Japan.
- Phragmidium rosae-rugosae** Kasai. This rust forms gall-like pustules on petioles, fruit and twigs and often deforms the leaf blades of *R. rugosa* in Japan. The pustules form on the lower leaf surfaces and are yellow at first, becoming brown to black.
- Phragmidium rosae-sempervirentis** Maire. Orange-yellow rust pustules on the leaves and black powdery spore masses in cankered areas on the stems of *R. sempervirens* in Algeria.
- Phragmidium rranjense** Bub. and Ranoj. Bright-yellow to deep brown rust pustules on the lower leaf surfaces of *R. spreta* in Albania and Yugoslavia.
- Phragmidium tuberculatum** J. Mueller. Brown to black rust pustules on *R. arvensis*, *R. canina*, *R. cinnamomea*, *R. inodora*, *R. lucida*, *R. micrantha*, *R. mollissima*, *R. rubiginosa*, *R. rubrifolia*, *R. scandens*, *R. sempervirens*, *R. sepium*, *R. spinosissima*, *R. tomentosa*, and *R. trachyphylla* in Europe, Siberia, and north Africa.
- Phragmidium yezoense** Kasai. Brown to black rust pustules on the lower surfaces of leaves and on petioles of *R. rugosa* in Japan.
- Phyllosticta argillaceae** Bres. Causes a leaf spot of cultivated roses in Europe.
- Phyllosticta rosarum** Pass. A dark-red to black leaf spot of *R. canina* and other cultivated roses in Russia, Italy, Switzerland, Austria, New South Wales, and Germany.
- Phyalospora rosicola** (Fckl.) Sacc. On branches of *R. canina* in Germany.
- Protoventuria rosae** (DeNot.) Berl. On branches and twigs of *R. alpina* in Italy.
- Ramularia banksiana** (Pass.) Sacc. On leaves of *R. banksiae* in Europe.
- Septoria rosae** Desm. This widespread European disease, known as leaf scorch, is characterized by pale-brown or dirty-white spots, irregular in shape, bounded by a broad purple margin. The centers of the diseased areas tend to fall out. The disease is especially serious on nursery stock, causing practical defoliation, and when this occurs over several seasons, the cumulative effect is serious. The hosts are *R. canina*, *R. pomifera*, *R. pumila*, *R. sancta*, *R. scandens*, and other species of cultivated roses in Europe, New Zealand, Abyssinia, and Porto Rico.
- Septoria rosae-arvensis** Sacc. Causes a leaf spot of *R. arvensis* and *R. sempervirens* in Spain and other European countries. Probably synonymous with the preceding.
- Septoria rosarum** West. Produces numerous white rounded spots with broad, purple borders on the upper surfaces of leaves of *R. cinnamomea*, *R. collina*, and *R. coriifolia*. Reported from Russia, Belgium, Italy, Great Britain, India, and Brazil. Probably not specifically distinct from *S. rosae* Desm.
- Stereum purpureum** Pers. See Prunus.
- Stilbum** sp. This fungus produces "small purplish spots on the rose stems which gradually increase in size, sometimes even girdling the branch. The bark becomes shrivelled and sunken, and in the later stages is covered with the bright red stalks of the fungus surmounted by red globules." Cultivated roses in Trinidad are subject to this disease.
- Uncinula simulans** Salm. A powdery mildew on the leaves and young twigs of *R. multiflora* in Japan.

**ROSMARINUS**. ROSEMARY. Hardy evergreen shrubs with aromatic leaves.

**Ascochyta rosmarini** Tassi. Circular to angular white leaf spots with red-brown borders on *R. officinalis* in Italy.

**ROTTBOELLIA**. Annual or perennial robust tropical grasses.

**Cintractia densa** McAlp. Smut sori in compact, dark-brown masses along rachides, destroying the florets of *R. compressa* in Australia.

**Meliola parenchymatica** Gaill. Black, superficial fungus patches on leaves of *R. exaltata* in the Philippines.

**Phyllachora rottboelliae** Syd. and Butl. Dull-black stromata on leaves of *R. exaltata* in India and the Philippines.

**Puccinia cacao** McAlp. Chocolate-brown to dark-brown rust pustules on leaves of *R. compressa* and *R. japonica* in India, Japan, and Australia.

**ROTTBOELLIA**—Continued.

- Puccinia microspora** Diet. Brown rust pustules on purple to brown leaf spots on *R. compressa* in Japan.
- Puccinia rottboelliae** Syd. Linear powdery black rust sori on leaves of *R. arundinacea* in Abyssinia and the Congo.
- Schizothyrium congœnsis** Beeli. On *Rottboellia* sp. in the Congo.
- Uromyces rottboelliae** Arth. Leaf rust on *R. compressa* and *R. speciosa* in India and Japan.
- Ustilago flagellata** Syd. and Butl. Elongate powdery dark-brown smut sori in spikes and rachides of *R. exaltata* and *Rottboellia* sp. in the Philippines and the Congo.
- Ustilago rottboelliae** Syd. and Butl. Black powdery smut sori, deforming and destroying rachides and flowers of *R. compressa* in China, Japan, and India.
- ROUPALA**. Tropical American trees.
- Catacauma rhopalinum** (Mont.) Theiss. and Syd. Black stromata on leaves of *Roupala* sp. in Cayenne, Brazil and Peru.
- Melasmia roupalae** Allesch. On leaves of *Roupala* sp. in Brazil.
- ROYENA**. Evergreen trees and shrubs.
- Accidium royenae** C. and M. Leaf rust on *R. pallens* in the Union of South Africa.
- ROYSTONEA**. See *Palmae*.
- RUBIA**. Hardy herbs.
- Accidium archavaletae** Speg. Yellow stem rust on *Rubia* sp. in Uruguay.
- Accidium rubiae** Diet. Leaf rust on *R. cordifolia* in Japan.
- Ascochyta rubiae** Rub. Gray circular leaf spots with purple-brown areoles on *R. peregrina* in Yugoslavia.
- Mycosphaerella peregrina** (Cke.) Lind. On stems and leaves of *R. peregrina* in England.
- Peronospora rubiae** Gäum. Downy mildew on leaves of *R. tinctorum* in Holland, France and Dalmatia.
- Pseudopeziza autumnalis** (Fr.) Karst. See *Galium*.
- Pseudopeziza komarovii** Jacz. Brown fruiting disks on yellow-brown sunken leaf spots on *R. cordifolia* in Manchuria.
- Puccinia algerica** Pat. Brown to black rust pustules on leaves of *R. laevis* in Algeria.
- Puccinia collettiana** Barcel. Leaf rust on *R. cordifolia* in India.
- Puccinia dimorpha** Syd. Chestnut-brown rust pustules on leaves and stems of *R. petiolaris* in the Union of South Africa.
- Puccinia rubicola** Syd. Brown rust pustules on leaves of *R. discolor* in Abyssinia.
- Puccinia rubivora** P. Magn. Leaf rust on *R. fruticosa* in the Canary Islands.
- Thecopsora rubiae** Kom. Yellow and dark-brown rust pustules on leaves of *R. cordifolia* in Japan, China, Manchuria, Ceylon, Siberia, and Russia.
- RUBUS**. BLACKBERRY. RASPBERRY. DEWBERRY. Herbs and shrubs cultivated for their fruit.
- Ascochyta fenilleauboisiana** Sacc. and Roum. White leaf spots with narrow dark borders on *Rubus* sp. in France.
- Ascochyta rubi** Sacc. Small subcircular deep red-brown leaf spots on *R. caesius*, *R. fruticosus* and *R. saxatilis* in Siberia, Russia, Italy, and Germany.
- Cercospora garbiniana** C. Mass. Grayish-white leaf spots on *R. fruticosus* in Italy.
- Cercospora rubicola** Thuem. Subcircular dull yellow-brown leaf spots with purplish margins on *R. fruticosus* in Russia and Portugal.
- Cladosporium** sp. Said to cause a stunting of plants of *Rubus* sp. (raspberry) in Switzerland, due to cankers formed on the stems. Fruit fails to form and the leaves are discolored.
- Coccomyces rubi** (Fr.) Karst. On leaves of *R. caesius*, *R. fruticosus*, and *R. idaeus* in Europe.
- Coleroa chaetomium** (Kze.) Rabh. On leaves of *Rubus* spp. in Europe.
- Coniothyrium tumaefaciens** Guess. Causes galls as large as walnuts on canes of *R. fruticosus* in Great Britain.
- Coryneum follicolum** Fckl. See *Crataegus*.
- Didymella applanata** (Niessl.) Sacc. Irregular gray cankered areas on canes of *Rubus* spp. (raspberry and loganberry), causing wilting as well as witches'-brooms in Great Britain and continental Europe.
- Gerwasia rubi** Rac. Golden yellow rust pustules on leaves of *Rubus* sp. in Java.
- Gnomoniella cercosporae** Pass. On leaves of *R. glandulosus* in Italy.
- Hamaspora acutissima** Syd. Leaf rust on *R. moluccanus*, *R. nantoensis*, *R. rolfeii*, and *Rubus* sp. in Java, Formosa, Australia, and the Philippines.
- Hamaspora engleriana** (Diet.) Syd. Yellow rust pustules on leaves of *R. volkensianus* in East Africa.
- Hamaspora gedeana** Rac. Yellow leaf rust on *R. alpestris* in Java.
- Hamaspora longissima** (Theum.) Koern. Yellow leaf rust on *R. moluccanus*, *R. pinnatus* and *R. rigidus* in India, Australia, Uganda, and the Union of South Africa.
- Hapalosphaeria deformans** Syd. Attacks flowers of *R. dumetorum* in Germany.
- Helminthosporium cymbispermum** Pat. On leaves of *Rubus* sp. in Ecuador.
- Hendersonia rubi** West. White patches on canes, causing sterility and finally wilting of *R. fruticosus*, *R. idaeus*, and *R. vitifolius* in Great Britain, France, Italy, and Belgium.
- Hyalothetes dimerosperma** Speg. On leaves of *R. urticaefolius* in Brazil.
- Hypomyces rubi** (Osterw.) Wr. Attacks roots of *R. idaeus* in Switzerland.
- Kuehneola andicola** Diet. Leaf rust on *R. geoides* in Chile.
- Kuehneola uleana** Syd. Rust on leaves of *Rubus* sp. in Brazil.
- Mycosphaerella chamaemori** (Karst.) Lind. On leaves of *R. chamaemorus* in Finland.
- Mycosphaerella minoensis** Syd. On leaves of *Rubus* sp. in Japan.
- Mycosphaerella winteri** (Pass.) Sacc. On leaves of *R. corylifolius* in Italy.
- Ovularia rubi** Bub. Circular to irregular, often confluent, brown leaf spots on *Rubus* sp. in Hungary.
- Phragmidium arcticum** Lagh. Yellow and black rust pustules on leaves of *R. arcticus* in Sweden and Finland.
- Phragmidium assamense** Syd. Leaf rust on *R. lasiocarpus* in India.
- Phragmidium barclayi** Diet. Yellow and brown rust pustules on red-brown leaf spots on *R. koehleri* and *R. lasiocarpus* in India.
- Phragmidium barnardi** Plowr. and Wint. Rust on leaf blades and petioles and on peduncles of *R. parvifolius* in Australia.
- Phragmidium burmanicum** Syd. Leaf rust on *R. lasiocarpus* in Burma.
- Phragmidium griseum** Diet. Yellow and black rust sori on irregular leaf spots on *R. incisus* in Japan.
- Phragmidium heterosporium** Diet. Leaf rust on *R. trifidus* in Japan.
- Phragmidium incompletum** Barcel. Yellow leaf rust on *R. paniculatus* in India.
- Phragmidium nambrianum** Diet. Powdery black rust pustules on sunken violet-brown leaf spots on *R. kinashii* and *R. occidentalis* in Japan.

**RUBUS**—Continued.

- Phragmidium octoloculare** Barcl. Black leaf rust on *R. rosaefolius* in Japan.
- Phragmidium orientale** Syd. Golden rust pustules on leaves of *R. ellipticus* and *R. moluccanus* in India and Ceylon.
- Phragmidium pauciloculare** Syd. Powdery rust pustules on leaf blades and petioles and on peduncles of *R. parvifolius*, *R. phoenicolasius*, and *R. triphyllus* in Japan.
- Phragmidium quinqueloculare** Barcl. Rust on leaves and stems of *R. biflorus* in India.
- Phragmidium rubi** (Pers.) Wint. Yellow and powdery black rust pustules on lower leaf surfaces of *Rubus* spp. (nearly 40 species are recorded as hosts) in Japan and Europe.
- Phragmidium rubi-fraxinifolii** Syd. Leaf rust on *R. fraxinifolius* and *R. taiwanianus* in Formosa.
- Phragmidium rubi-idaei** (DC.) Karst. Yellow to black rust pustules on leaves of *R. idaeus* in Japan, Alaska, Siberia, and Europe. Probably occurs in the United States.
- Phragmidium rubi-japonici** Kasai. Black rust pustules on leaves of *R. japonicus* in Japan.
- Phragmidium rubi-saxatilis** Liro. Yellow and powdery black rust pustules on leaves of *R. saxatilis* in northern Europe.
- Phragmidium rubi-sieboldii** Kaw. (*Hamaspora rubi-sieboldii* [Kaw.] Diet.) Leaf rust on *R. sieboldii* in Japan.
- Phragmidium rubi-thunbergii** Kus. Leaf rust on *R. idaeus* and *R. thunbergii* in Japan.
- Phragmidium violaceum** (Schultz.) Wint. Golden-yellow and black rust pustules on leaves of *Rubus* spp. (60 species are listed as hosts) in China, Indo-china, Egypt, and Europe.
- Phragmidium yoshinagai** Diet. Leaf rust on *R. crataegifolius*, *R. morifolius*, and *R. sorbifolius* in Japan.
- Phragmidium zeylanicum** Petch. Leaf rust on *R. lasiocarpus* in Ceylon.
- Phyllosticta argillacea** Bres. On leaves of *R. idaeus* in Bohemia and Silesia.
- Phyllosticta cryptocarpa** Kab. and Bub. Brown, often confluent, leaf spots with purple margins on *R. glandulosus* in Bohemia.
- Phyllosticta fusco-zonata** Thuem. Subcircular dull-brown leaf spots on *R. idaeus* in Algeria, Russia, and Italy.
- Phyllosticta pallor** (Berk.) Oud. On *Rubus* (raspberry) in Great Britain and Holland.
- Phyllosticta rubi** P. Henn. Circular leaf spots with red-brown margins on *Rubus* sp. in Brazil.
- Phyllosticta rubicola** Rabenh. Subcircular small white leaf spots with blood-red margins on *R. caesius* and *R. idaeus* in Russia, Dalmatia, Italy, and Germany.
- Phyllosticta rubi-odorati** Bub. and Kab. On leaves of *R. odoratus* in Bohemia.
- Rhabdospora ramealis** (Desm. and Rob.) Sacc. and var. **macrospora** App. and Laub. Causes small purple-red areas on stems of *Rubus* spp. in Europe.
- Sclerotinia rubi** Carm. White, then brown, sclerotia on leaves of *R. fruticosus* in Great Britain.
- Septoria campoi** Speg. On leaves of *R. sanctus* in Chile.
- Uredo chinensis** Diet. Leaf rust on *R. buergeri* and *R. reflexus* in Japan.
- Uredo imperialis** Speg. Golden-yellow leaf rust on *R. imperialis* in Argentina.
- Uredo sinensis** Diet. Brown rust pustules on leaves of *R. reflectus* in China.
- Uromyces arthuri** Syd. Leaf rust on *R. guyanensis* and *R. schiedeana* in Guatemala.
- Uromyces cundimarcensis** Mayor. Leaf rust on *R. peruvianus* in Colombia.
- Uromyces lagerheimii** P. Magn. Yellow leaf rust on *R. glaucus* and *Rubus* sp. in Ecuador and Colombia.
- Uromyces loesenerianus** (P. Henn.) Syd. Yellow leaf and stem rust on *R. bogotensis*, *R. braziliensis*, *R. trichomallus*, and *R. urticaefolius* in Central and South America.
- Uromyces pittierianus** P. Henn. Yellow leaf rust on brown sunken spots on *R. adenotrichos* and *Rubus* sp. in Costa Rica.
- Uromyces quitensis** Lagh. Leaf rust on *Rubus* sp. in Ecuador.
- Uromyces rubi** Diet. and Holw. Rust on sunken leaf spots on *R. adenotrichos*, *R. glaucus*, *R. laxus*, *R. poliophyllus*, *R. pringlei*, and *R. trichomallus* in Mexico, Guatemala, and Costa Rica.
- Uromyces rubi-urticifolii** Mayor. Rust on leaves of *R. urticaefolius* in Colombia.
- Uromyces variabilis** Mayor. Leaf rust on *Rubus* sp. in Colombia.
- Venturia kunzei** Sacc. On leaves of *R. caesius*, *R. idaeus*, and *R. pedatus* in Europe and Alaska.
- RUDGEA**. Shrubs or small trees.
- Phyllachora rudgeae** Syd. Black stromata on yellow-brown leaf spots on *R. coriacea* in Brazil.
- RUELLIA**. Acanthaceous herbs or shrubs.
- Aecidium tuberosa** P. Henn. Leaf rust on *R. tuberosa* in Peru.
- Puccinia mussoni** McAlp. Yellow and dark-brown rust sori on leaves of *R. australe* in Australia.
- Puccinia paranahybae** P. Henn. Dark-brown leaf rust on *R. longifolia* in Brazil and Argentina.
- Puccinia ruelliae-bourgaei** Diet. and Holw. Powdery yellow and brown rust pustules on leaves of *R. bourgaei* in Mexico.
- Uromyces ruelliae** Holw. Brown to black rust pustules on leaves of *Ruellia* sp. in Mexico.
- RUMEX**. DOCK. SORREL. Perennial herbs, mostly weeds.
- Ascochyta rumicis** Bub. and Malk. On *Rumex* sp. in Bulgaria.
- Ascochyta vicina** Sacc. On stems of *R. acetosella* and *Salvia officinalis* in Spain, France, and Italy.
- Cylindrosporium pulchrum** Speg. On leaves of *R. obtusifolius* and *R. pulcher* in Argentina.
- Depazea acetosa** Op. On leaves of *R. acetosa* in Siberia, Austria, and Germany.
- Mycosphaerella insulana** Bub. and Syd. On leaves of *Rumex* sp. in Germany.
- Mycosphaerella rumicis** Desm. Circular brown leaf spots on *R. conglomeratus*, *R. crispus*, *R. hydro-lapathum*, and *R. nemolapathum* in China and Europe.
- Ovularia canaegricola** P. Henn. Pale-yellow or brown leaf spots with white centers on *R. hymenosepalus* in Germany.
- Ovularia rubella** (Bon.) Sacc. On leaves of *R. aquaticus* in Germany.
- Ovularia rumicis** Eliasson. Circular to elliptical ochraceous leaf spots on *R. crispus* in Sweden.
- Pellicularia chilensis** Speg. Leaf blight of *R. crispus* in Chile.
- Peronospora rumicis** Cda. Downy mildew on leaves, stems, and panicles of *R. acetosa*, *R. acetosella*, *R. arifolius*, *R. auriculatus*, *R. crispus*, *R. lunaria*, *R. scutatus*, *R. thyrsiflorus*, and *R. vesicarius* in India, Tunis, and Europe.
- Phyllosticta acetosae** Sacc. Subcircular brown leaf spots on *R. acetosa* in Italy and Russia.
- Phyllosticta straminella** Bres. On leaves of *R. acetosa* in Germany.
- Physoderma acetosellae** Rostr. Spore masses in deformed ovaries of *R. acetosella* in Denmark.
- Puccinia biformis** Lagh. Dark-brown rust pustules on leaves and stems of *R. bucephalophorus* in Portugal.
- Puccinia dissiliens** Cke. Brown leaf rust on *Rumex* sp. in northern India.
- Puccinia ludwigii** Tepper. Rust on leaves of *R. brownii* in Australia.
- Puccinia nepalensis** Barcl. and Diet. Brown rust pustules on leaves of *R. nepalensis* in India.
- Puccinia pachyphloea** Syd. Black rust pustules on leaf blades and petioles and stems of *R. tuberosus* in Kurdistan.
- Puccinia rumescicola** Gz. Frag. Leaf rust on *R. papillaris* in Spain.

**RUMEX**—Continued.

- Puccinia rumicis-scutati** (DC.) Wint. Brown leaf and stem rust on *R. scutatus* in Europe and possibly Australia.
- Puccinia trailii** Plowr. See Phragmites.
- Ramularia rumicis** K. and Cke. Large dull-brown leaf spots on *R. obtusifolius* in the Union of South Africa.
- Ramularia rumicis-scutati** Allesch. Circular leaf spots with white centers and broad ochraceous margins on *R. scutatus* in Spain, Switzerland, and Germany.
- Schroeteriaster alpinus** (Schroet.) P. Magn. Brown rust pustules on leaves of *R. alpinus* in central Europe.
- Septoria acetosae** Oud. Brown leaf spots with purple borders on *R. acetosa* and *Rumex* sp. in Denmark and Holland.
- Septoria glacialis** Ferr. Ochraceous circular zoned leaf spots on *R. scutatus* in Italy.
- Stigmatea rumicis** (Desm.) Schroet. On leaves of *R. alpinus* in Switzerland.
- Uredo purpurascens** Diet. Leaf rust on *R. romassa* in Chile.
- Uromyces acetosae** Schroet. Yellow and brown rust pustules on leaf blades and petioles of *R. acetosa*, *R. acetosella*, and *R. arifolius* in Europe.
- Uromyces argaeus** Maire. Powdery brown rust pustules on leaves of *R. tuberosus* in Asia Minor.
- Uromyces borealis** Liro. Leaf rust on *R. arifolius* in Finland.
- Uromyces crassipes** Diet. and Neg. Brown leaf rust on *R. cuneifolius* in South America.
- Uromyces tingitanus** P. Henn. Leaf rust on *R. tingitanus* in Algeria and Tunis.
- Urophlyctis rübsaameni** Magn. Tuberosus excrescences on roots of *R. scutatus* in Switzerland and Germany.
- Ustilago domestica** Bref. Dull-violet masses of smut spores in stems, leaves, and inflorescences of *R. domesticus* in Norway.
- Ustilago kühneana** (Wolf) F. v. W. Smut sori in leaves and panicles of *R. acetosa*, *R. acetosella*, and *R. tuberosus* in north Africa and Europe.
- Ustilago lagerheimii** Bref. Powdery dark-violet masses of smut spores in stems and petioles of *Rumex* sp. in Ecuador.
- Ustilago warmingii** Rostr. Leaf smut on *R. crispus* and *R. domesticus* in Finland.
- Venturia rumicis** (Desm.) Wint. On leaves of *R. alpinus*, *R. crispa*, *R. nemorosus*, and *R. obtusifolius* in French North Africa, Denmark, and Bulgaria.
- RUSCUS**. BUTCHER'S-BROOM. Erect shrubs.
- Colletotrichum erumpens** Sacc. On *R. aculeatus* in France.
- Leptosphaeria rusci** (Walbr.) Sacc. On stems of *R. aculeatus* and *R. hypoglossum* in Switzerland, Spain and Italy.
- Phyllosticta dancöes** Pass. Large irregular gray-white spots with brown margins on pseudo leaves of *R. racemosus* in France.
- Phyllosticta rusciicola** Dur. and Mont. Subcircular white leaf spots with brown margins on *R. aculeatus*, *R. androgynus*, *R. hypophyllum*, and *R. hypoglossum* in Spain, Portugal, Italy, and France.
- Septoria hypoglossi** Massal. Leaf spots on *R. hypoglossum* in Italy.
- Urocystis jaapiana** Sacc. Smut on *R. aculeatus* in France, Spain, and Italy.
- RUSSELLIA**. CORAL BLOW. Shrubs.
- Phyllosticta armitageana** Sacc. On leaves of *R. juncea* in Malta.
- RUTA**. RUE. Perennial glandular herbs.
- Accidium rutae** Har. Yellow rust on leaf blades and petioles of *R. chalepensis* in France and Sardinia.
- Ovulariopsis haplophylla** (P. Magn.) Trav. On leaves of *R. graveolens* in Italy.
- SABAL**. See Palmae.
- SACCHARUM**. Sugar cane. Tall-growing, perennial, sugar-yielding grasses.
- Allantospora radiceicola** Wakk. On roots of *S. officinarum* in Java and Hawaii.
- Bacillus flavidus** Fawc. Reported as one of the causes of top rot or "polvillo" of sugar cane (*S. officinarum*) in Argentina.
- Bacillus sacchari** Speg. This bacterium is considered by Spegazzini as the cause of the "polvillo" disease of sugar cane (*S. officinarum*) in Argentina. Fawcett, however, in the most recent work on the disease, assigns *B. flavidus* and two other unnamed species as the causes. The disease is typical of diseases discussed under the head of top rot.
- Bacterium** sp. Causes a leaf-scald disease on *S. officinarum* in Australia and probably New Guinea.
- Bacterium vascularum** Cobb. The gumming disease due to this bacterium is a serious sugar-cane (*S. officinarum*) trouble in New South Wales, Queensland, Mauritius, New Guinea, Borneo, Fiji, Reunion, Java and possibly in Brazil. What is apparently the same disease has appeared in Porto Rico within the last few years. Externally diseased canes are stunted, the leaves show red longitudinal stripes, and there may be top rotting. On cutting across infected canes the bundles show discolored and a yellow gumlike substance oozes out. Many of the bundles are stained red, the discoloration extending out into the leaves. The gumming disease of Java is probably not due to *B. vascularum*, since the typical gummy exudate from the cut end of infected canes is entirely absent. Etiological studies have not yet been carried out.
- Bakerophoma sacchari** Died. Small brown leaf spots with red margins on leaves and leaf sheaths of *S. officinarum* in the Philippines.
- Bunchy top**. See Musa.
- Cephalosporium sacchari** Butl. This fungus causes a wilt of sugar cane in India, the Philippines, British West Indies, and the Union of South Africa. Affected stools become stunted, the leaves drying up and stems becoming light and hollow. Diseased canes internally are diffuse purple or dirty-red in color. These areas are distinct from the red-bordered white patches characteristic of red rot (*Colletotrichum falcatum*).
- Cercospora acerosum** Dick. and Hein. Black spots on leaves of *S. officinarum* in Java and the Philippines.
- Cercospora kopkei** Krueg. Dull-yellow, often confluent, leaf spots becoming red at the centers on *S. edule*, *S. glongong*, *S. officinarum*, and *S. spontaneum* in Java, Japan, Reunion, Formosa, Queensland, the Philippines, Argentina, and Cuba.
- Cercospora longipes** Butl. Small oval red leaf spots on *S. officinarum* in Porto Rico and India. The spots increase in size, becoming brown with yellow areolae and finally straw-colored, with oval, deep brown rings surrounding.
- Contractia pulverulenta** Cke. and Mass. Smut in inflorescences of *S. arundinaceum* in India.
- Cytospora sacchari** Butl. The leaf sheaths take on a dark dull-red color and are bound firmly together by the mycelium of the fungus. The leaf blades turn brown and hang parallel to the stalk. Young stalks are killed outright and on older ones which have formed cane, brown lesions appear on the internodes, triangular at first when originating at the nodes, irregular if starting from wounds or cracks in the internodes. The fungus fruits on the outer leaf sheaths, the black, elongated necks of the pycnidia being rough to the touch. The disease occurs on *S. officinarum* in Porto Rico, Barbados, Formosa, India, and Brazil.

## SACCHARUM—Continued.

- Dinemasporium sacchari** P. Henn. Circular to oblong brown leaf spots with red-brown margins on *S. officinarum* in Peru.
- Diplodia cacaoicola** P. Henn. Common as a wound parasite of cuttings and on dead canes and occasionally causing damage to susceptible varieties of *S. officinarum*. Affected canes wither and die, and are internally reddened, turning brown on exposure to air. See also Theobroma.
- Eriosphaeria sacchari** (v. Breda) Went. Dark-red circular leaf spots up to 1 centimeter in diameter on *S. officinarum* in Java and Trinidad.
- Euryachora sacchari** Avena. Stromata on leaves of *S. officinarum* in Brazil.
- Fiji disease.** Linear galls are produced on the leaves and leaf sheaths from 2 millimeters to 6 centimeters long, light-green in color at first and finally brown, bursting open, exposing a brownish mass. Diseased leaves are darker green than normal, as well as smaller and more slender. Infected plants, due to the shortening of the internodes are small, stunted, and have a bunched growth of leaves at the top. The roots are small, bunched, and slightly rotted. New shoots arising from the bases of heavily infected plants have a yellow streak running down the unfolded leaves which become distorted and wrinkled. A brown rot sometimes occurs in advanced cases. The cause of the disease is uncertain, but has recently been attributed to a protozoan (*Phytamoeba sacchari* McWhorter). The disease occurs in Australia, New Guinea, Fiji, and the Philippines on *S. officinarum*.
- Gnomonia iliau** Lyon. Iliau disease of sugar cane (*S. officinarum*) is characterized by a binding of the leaf sheaths firmly together by the mycelium of the causative fungus. The leaf blades die, turn brown, and hang down. The leaf sheaths are pinkish-brown in color and the stalks, if any cane is produced at all, are deep bluish-gray. The disease commonly attacks young shoots and prevents their further development. The imperfect or Melanconium stage of the fungus is produced internally in leaf sheaths and stalks, but the perithecia occur on the outer leaf sheaths, the beaks projecting. The disease is prevalent in Hawaii and is known also from limited areas in Louisiana.
- Helminthosporium sacchari** Butler. The eye-spot disease occurs as linear brown leaf spots with definite margins on *S. officinarum* in Porto Rico, Jamaica, Cuba, Barbados, St. Domingo, Hawaii, the Philippines, Java, Reunion, India, and the Union of South Africa. On some varieties the spots are oval, small, zoned, and yellow-brown in color. Susceptible varieties are checked in their growth and often stunted. The disease is known from Florida and Georgia.
- Hendersonia sacchari** Butl. This fungus causes withering of *S. officinarum* in India and Ceylon. The leaves die from the center outward and downward. Individual leaves die back from the tip and along the margins. Internally the diseased canes are dry and pithy, with a cavity in each internode. The roots are rotted.
- Leptosphaeria sacchari** v. B. H. Numerous oval dull-gray spots with definite red or red-brown margins occur on the leaves of *S. officinarum*. There may be an indistinct yellow halo surrounding each spot. In many varieties of cane the fungus causes premature withering and death of the leaf blades. The fungus is known from practically all the sugar-cane growing countries of the world. It is reported from Alabama and Florida.
- Leptosphaeria saccharicola** P. Henn. Pale-brown leaf spots with darker margins on *S. officinarum* in Brazil. Probably not distinct from the preceding species.
- Leptosphaeria spagazzinii** Sacc. and Syd. and var. **minor** Speg. On leaf blades and sheaths of *S. officinarum* in Argentina.
- Leptosphaeria tucumanensis** Speg. On leaves of *S. officinarum* in Argentina.
- Linospora sacchari** Avena. On leaves of *S. officinarum* in Brazil.
- Lophodermium sacchari** Lyon. Black linear fruiting bodies on midribs and sheaths of "sereh" cane in Hawaii, and "may be mildly parasitic."
- Marasmius sacchari** Wakker. Root disease is one of the most serious, if not the most serious, disease of sugar cane (*S. officinarum*) and occurs in every sugar-cane growing country. There is a very extensive literature on the subject, the disease being attributed in large part to the above fungus. Most recent studies indicate, however, that other fungi, such as *Rhizotonia* spp., *Pythium*, etc., are the initial agents and *Marasmius*, *Odontia*, and other hymenomycetes secondary forms, only mildly parasitic at best. *M. sacchari* is commonly found in connection with root disease, the mycelium binding the lower leaf sheaths together by typical mycelial fans. The small white sporophores are produced in abundance in and around diseased stools during wet weather. Other species occur less commonly in similar manner.
- Melanconium sacchari** Mass. The rind disease of sugar cane (*S. officinarum*) occurs in practically every cane-growing country. The fungus causes a souring of the juice and a soft white rot of the tissues, affected stalks withering and drying up. Fruiting pustules form in great numbers on rotted stalks and at the base of leaf sheaths as small black areas from which the spore mass exudes, often as long, threadlike processes. The fungus is for the most part a wound parasite and seldom serious, except on overmature cane or cane first weakened by other agencies. Occurs as a saprophyte in Louisiana and Florida.
- Meliola arundinis** Pat. Superficial black fungus patches on leaves of *S. officinarum* in China and the Philippines.
- Meliola sacchari** Syd. Similar to the preceding species, in the Philippines.
- Mosaic (Mottling).** The mosaic or yellow-stripe disease of sugar cane (*S. officinarum*) occurs in Porto Rico, Cuba, Jamaica, Guadeloupe, Barbados, Trinidad, Santo Domingo, Argentina, Egypt, Hawaii, the Philippines, Java, China, and Formosa. The disease has been found also on other grasses, including *Chaetochloa magna*, *Brachiaria platyphylla*, *Holcus sorghum*, *Paspalum boschianum*, *Pennisetum glaucum*, *Syntherisma sanguinale*, and *Zea mays*. It has been especially serious on maize (*Zea*). By inoculation other grasses including *Miscanthus sinensis*, *Saccharum narenga*, *Chaetochloa lutescens*, *C. magna*, and *Echinochloa crusgalli* have been proved susceptible. This disease was introduced into Louisiana some years ago and has been disseminated from there into Florida, Georgia, Alabama, and Mississippi, and has already caused considerable losses.
- The disease is characterized by a mottling of the leaves, varying in shading and patterns, but always very characteristic. In early infections there are no other outstanding symptoms. As the disease progresses in susceptible varieties, and generally after one or two seasons, cankers appear on the stalks, at first as discolored or water-soaked patches or streaks on the internodes, later sunken and brown to grayish white in color. On very susceptible varieties the cankers are very numerous and cause the canes to dry out. In such cane there is also a marked shortening of the joints and premature root development beneath the leaf sheaths. Resistant varieties may show no more than the leaf symptoms over long periods, with relatively slight reductions in yield, while on the other hand fields of susceptible types may be rendered worthless within two seasons.
- Mycosphaerella sacchari** Speg. On leaves of *S. officinarum* in Cuba, Porto Rico, and Argentina.
- Mycosphaerella striatiformans** Cobb. This fungus is said to attack the leaf tissue between the veins, causing a characteristic yellow striping. In the final stage the leaves break up into as many strings as there are stripes, the resulting shreds hanging down, while the mid-rib remains erect. The disease occurs in Hawaii on *S. officinarum*.

**SACCHARUM**—Continued.

- Odontia saccharicola** Burt. The granular leaf-sheath fungus binds together the lower leaf sheaths and develops fruiting areas as uniform thin white patches with a granular, somewhat powdery surface encircling the stalk from the ground level to a height of about a foot. The stellate-crystal fungus (*Himantia stellifera* Johnston), which often occurs in connection with root disease, is undoubtedly the vegetative stage of this fungus. The exact relation of this fungus to root disease is entirely problematical and its status is approximately the same as *Marasmius*. *O. saccharicola* is reported from Porto Rico and Santo Domingo, although the stellate crystal form is known from Cuba, Barbados, St. Croix, British Guiana, Trinidad, Hawaii, Zululand, and the Union of South Africa on *S. officinarum*.
- Pestalozzia fuscescens** Sor. var. **sacchari** Wakk. Irregular gray areas with brown margins on leaves of *S. officinarum* in Java and the Philippines.
- Phoma heterospora** Speg. Pale-brown indefinite spots on leaves of *S. officinarum* in Argentina.
- Phyllachora sacchari** P. Henn. Linear dull-black stromata on leaves of *S. officinarum* and *S. spontaneum* in Java, India, and the Philippines.
- Phyllachora sacchari-aegyptiaci** Br. and Cav. Oblong to irregular, black stromata on leaves of *S. aegyptiacum* in Italy.
- Phyllachora sacchari-spontanei** Syd. Black stromata on yellowish to red-violet leaf spots on *S. spontaneum* in India and the Philippines.
- Phyllosticta hawaiiensis** Caum. On leaves of *S. officinarum* in Hawaii.
- Phyllosticta sacchari** Speg. Linear gray-white leaf spots with purple margins on *S. officinarum* in Porto Rico and Argentina.
- Phyllosticta saccharicola** P. Henn. Oblong gray-white leaf spots with dull-brown margins on *S. officinarum* in the Congo.
- Plasmodiophora vascularum** Matz. This organism is said to inhabit the vascular areas of cane stalks (*S. officinarum*) causing a stunting and top rot in Porto Rico and Barbados.
- Polvillo** (*Gangrena humida*). See *Bacillus sacchari*.
- Rhizoctonia ferrugena** Matz. Isolated from diseased roots of *S. officinarum* in Porto Rico.
- Rhizoctonia palida** Matz. Found on roots of *S. officinarum* in Porto Rico, as well as on *Capsicum* sp. and *Zea mays*.
- Sclerospora** sp. Oogonial stages of *Sclerospora* occur in the Philippines on *S. officinarum*, *S. spontaneum*, and *Miscanthus japonicus*, the relationships of which to the conidial forms on *Zea* (q. v.) and other hosts are uncertain.
- Sclerospora philippinensis** Weston. See *Zea*.
- Sclerospora sacchari** Miy. This downy mildew of sugar cane (*S. officinarum*) is very serious in Formosa, Australia, and Fiji and has recently been found in the Philippines. Yellow stripes parallel to the veins appear on the leaves, increasing in number until the greater part of the leaf becomes pale and is marked by long, reddish-brown spots, on the under surfaces of which fluffy fungus outgrowths occur. Diseased leaves die and dry up, often becoming much torn or shredded before dying. Leaf sheaths are also affected. With some varieties the joints are abnormally long and greater in number than in healthy canes. If the cane is attacked while young, shoots arise from buds along the stalk, giving a broomlike appearance. It has been possible to infect *Euchlaena* and *Zea mays*.
- Sclerospora spontanea** Weston. See *Zea*.
- Sclerotium** sp. Yellowish to white irregular patches, broader than long, and with narrow purple or red-brown surrounding zones on leaves of *S. officinarum*. These patches are separated by narrow green or brown gaps, the result being a variegated appearance which is very striking. Irregular white to brown sclerotia form on the diseased areas. The disease is prevalent and serious in Java, India, and the Philippines.
- Sclerotium griseum** Stevenson. Rounded gray sclerotia on leaf sheaths of *S. officinarum* in Porto Rico. This form has recently been connected with root disease in common with one or more species of *Rhizoctonia*.
- Sereh**. Sereh has been a serious disease of sugar cane (*S. officinarum*) in Java, although it is now fairly well controlled by the use of resistant varieties and by growing the seed cane in high altitudes. The disease or one very similar is known from Malacca, Borneo, Siam, Hawaii, Malaya, and the Philippines. The cause remains unknown, although the disease is probably to be classed with the infectious mosaics.
- Diseased stools fail to produce upright stems and the shortening of the internodes close to the ground gives the stunted plants a fan-shaped appearance. The leaves are smaller than normal and yellow-striped, dying in severe cases. The roots are weakly developed and soon die. The vascular bundles are reddened and show some gumming. The disease is readily carried in seed pieces without its presence being in evidence.
- Sphaeronema adiposum** Butl. A weak parasite of cane (*S. officinarum*) cuttings in India and Java. Diseased cuttings are dull-red within, finally drying up, with a black fungus growth on the exterior.
- Sphaerulina sacchari** P. Henn. Circular to oblong, then confluent, pale brown leaf spots with red-brown margins on *S. officinarum* in Peru.
- Stigmia sacchari** Speg. On leaves of *S. officinarum* in Argentina.
- Thielaviopsis paradoxa** (DeSeyn.) v. Hoeh. The pineapple or black rot disease of sugar cane is a disease of cuttings only, acting as a wound parasite when the plantings are made in wet soils or under other unfavorable conditions. Diseased cuttings fail to germinate or young shoots die back, the interior of the cutting turning black and giving off an odor of rotting pineapple. The fungus also causes the common soft rot or shipping rot of pineapples (*Ananas*) which has caused heavy losses to pineapple growers in the West Indies. Circular to irregular brown sunken leaf spots also occur on the pineapple plant due to this fungus. The stem-bleeding disease of coconut (*Cocos nucifera*) and betel palm (*Areca catechu*) is likewise attributed to it. In this instance it acts as a wound parasite, discoloring the tissues as it advances and causing a gummy exudate. The fungus is known to occur in the West Indies from Cuba to Trinidad, British Guiana, Hawaii, the Philippines, Mauritius, Ceylon, Java, Malaya, India, and Gold Coast. It has been isolated from coconut and sugar cane (*S. officinarum*) from Florida and probably occurs on other hosts in the southern United States.
- Top rot**. Top rot, characterized by the death of the terminal bud and inclosing leaves, is a common disease of sugar-cane (*S. officinarum*) in all cane-growing countries. A number of fungi and probably bacteria are undoubtedly involved, some at least of which do not occur in the United States. The "polvilla" of Argentina and dry top rot (*Plasmodiophora*) of Porto Rico and Cuba are two diseases of this class.
- Tylenchus similis** Cobb. See *Musa*.
- Uromyces kühni** Krueg. Elongated brown rust pustules on long orange spots on both leaf surfaces of *S. arundinaceum*, *S. fuscum*, *S. officinarum*, and *S. spontaneum* in Java, Japan, India, Burma, Ceylon, the Philippines, and Australia.

**SACCHARUM**—Continued.

- Ustilaginoidea sacchari-narengae** K. Saw. Top-shaped dark-olive fruiting bodies in ovaries of *S. narenga* in Japan, the protecting membranes of which rupture at maturity, exposing the dark-olive spore masses.
- Ustilago sacchari** Rabh. From the growing point there is produced a long, whiplike, black shoot without leaves, often several feet in length and much curved or twisted. In the earlier stages these smutted shoots are protected by silvery-white membranes, which rupture, exposing dense black spore masses. Secondary lateral shoots may also be infected. On *S. cylindricum*, *S. officinarum*, *S. spontaneum*, and *Erianthus asper* in British Guiana, Trinidad, Mauritius, Reunion, Union of South Africa, the Philippines, Japan, Formosa, China, India, Java, Queensland, and Italy.
- Ustilago sacchari-ciliaris** Bref. Powdery black masses of smut spores in ovaries of *S. ciliare* in India.
- SAGERETIA**. Armed or unarmed shrubs.
- Aecidium sageretiae** P. Henn. Leaf rust on *Sageretia* sp. in China.
- SAGINA**. PEARLWORT. Annual or perennial herbs.
- Septoria nivalis** Rostr. On peduncles of *S. nivalis* in Greenland.
- Septoria velenovskyi** Bub. Small circular yellow leaf spots on *S. tenuifolia* in Yugoslavia.
- SAGITTARIA**. ARROWHEAD. Hardy perennial aquatic herbs.
- Cercospora macrospora** Bres. Subcircular dull-brown leaf spots on *S. sagittifolia* in Germany.
- Doassansia horiana** P. Henn. Black smut sori deforming leaf blades and petioles of *S. sagittifolia* in Japan.
- Doassansia limosellae** (Kze.) Schroet. Smut sori in circular brown leaf spots on *S. sagittifolia* and *Limosella aquatica* in Europe.
- Mycosphaerella sagittariae** Tass. On leaf blades, petioles and sheaths and on stems of *S. montevidensis* in Italy.
- Phyllosticta sagittifoliae** Brun. Subcircular to oblong ochraceous leaf spots with dark brown margins on *S. sagittifolia* in Siberia and France.
- Ramularia sagittariae** Bres. Small dull-brown leaf spots on *S. sagittifolia* in Germany.
- SALIX**. WILLOW. Dioecious trees and shrubs.
- Ascochyta martianoffiana** Thuem. Subcircular ochraceous to gray leaf spots on *S. pyrolaefolia* in Siberia.
- Ascochyta salicicola** Pass. Angular reddish to dull brown leaf spots on *S. alba* in France.
- Ascochyta translucens** Kab. and Bub. Circular to angular, then confluent, broad gray leaf spots on *S. caprea* in Bohemia.
- Ascochyta vitellinae** Pass. On leaves of *A. vitellina* in Europe.
- Bacillus hari** Hori. and Miy. Bacteriosis of *Salix* sp. in Japan.
- Bacterium salicis** Day. This bacterium is reported as the cause of the "watermark" disease of *S. alba* and *S. caerulea* in Great Britain. Leaves of infected trees wither and die, turning brown and remaining attached. New shoots and finally entire branches are infected, the disease ultimately reaching the trunk and killing the tree after one or more years. Infected trees develop adventitious shoots and a gummy exudate from dying branches. There is an internal gray staining of the wood.
- Botryosphaeria gregaria** Sacc. Causes the bark to turn brown, crack and die on twigs and branches of *Salix* sp. (basket willow) in Ireland.
- Dinemasporium purpurascens** Rich. On branches of *Salix* sp. in France.
- Dipodina salicina** Cke. and Mass. Causes a die-back of *Salix* spp. in Great Britain.
- Discella carbonacea** (Fr.) B. and Br. Small bark cankers causing die-back of *Salix* Sp. in Holland and Sweden.
- Entyloma salicis** Karst. Leaf smut on *S. caprea* in Finland.
- Fusicladium saliciperduum** (All. and Tub.) Tub. This disease attacks *S. alba*, *S. aurita*, *S. caprea*, *S. cinerea*, *S. cuspidata*, *S. fragilis*, *S. mollissima*, *S. nigricans*, and *S. pentandra* in Scotland, Russia, Denmark, Holland, and Germany. *Venturia chlorospora* (Ces.) Karst. is considered the perfect stage. Dark olive, velvety patches occur on the leaves and spread down along the petioles to the twigs which as a result die back. Long black patches, sharply demarcated from the healthy yellow bark, form on the branches.
- Gloeosporium deformans** (Schroet.) J. Lind. On catkins of *S. caprea* and *S. caprea-cinerea* in Sweden, Finland, Denmark, Silesia, and France.
- Graphium ulmi** Schwarz. See Ulmus.
- Haplothecium amenti** (Rostr.) Theiss. and Syd. On catkins of *S. reticulata* in Norway.
- Helicobasidium mompa** Tan. See Morus.
- Helicobasidium tanakae** Miy. See Morus.
- Hendersonia foliorum** Fekl. Parasitic on leaves of *S. caprea* in Europe.
- Marsonia didyma** (Fekl.) Sacc. Small dull-brown leaf spots on *S. daphnoides*, *S. fragilis*, and *S. amygdalina* (*S. triandra*) in Russia, Poland, France, and Germany.
- Marsonia kriegeriana** (Bres.) P. Magn. Circular black leaf spots on *S. amygdalina* and *S. triandra* in Germany.
- Marsonia obscura** Rom. Dull-red leaf spots on *Salix* sp. in Scandinavia.
- Marsonia pyrenaica** Gz. Frag. On leaves of *S. caprea* in Spain.
- Marsonia salicicola** (Bres.) Magn. Indefinite rufous leaf spots on *S. alba* and *S. caprea* in Bohemia, Denmark, and Germany.
- Marsonia salicis** Trail. On leaves of *Salix* sp. in Norway.
- Marsonia santonensis** (Pass.) Bub. On leaves of *S. pentandra* in Austria.
- Melampsora abietis-capraearum** Tub. Yellow and brown rust pustules on needles of *Abies alba* and *A. nordmanniana* and on the leaves of *S. caprea* in Denmark, France, Spain, and Germany.
- Melampsora allii-fragilis** Kleb. Yellow leaf rust on *Allium ascalonicum*, *A. cepa*, *A. oleraceum*, *A. porrum*, *A. sativum*, *A. schoenoprasum*, *A. ursinum*, *A. vineale*, *Salix fragilis*, *S. pentandra*, *S. triandra*, and *S. viminalis* in Europe.
- Melampsora amygdalinae** Kleb. Leaf rust on *S. amygdalina*, *S. hippophaefolia*, and *S. pentandra* in Europe.
- Melampsora coleosporioides** Diet. Yellow to brown rust pustules on leaves of *S. babylonica* and *S. glandulosa* in Japan.
- Melampsora epiphylla** Diet. Yellow to dark-brown rust pustules on leaves of *S. shikokiana* in Japan.
- Melampsora euonymi-capraearum** Kleb. Yellow to brown rust pustules on sunken, yellow circular to irregular leaf spots on *S. aurita*, *S. caprea*, *S. cinerea*, *S. incana*, *S. viminalis*, *Euonymus europaeus*, *E. latifolia*, and *E. verrucosa* in Europe and Japan.
- Melampsora galanthi-fragilis** Kleb. See Galanthus.
- Melampsora humboldtiana** Speg. Leaf rust on *S. humboldtiana* in Uruguay and Argentina.
- Melampsora humilis** Diet. Yellow to brown rust pustules on leaves of *S. multinervis* in Japan.
- Melampsora lapponum** Lindf. Leaf rust on *S. lapponum* and *Viola epipsila* in Sweden.

## SALIX—Continued.

- Melampsora larici-caprearum** Kleb. Yellow leaf rust on *S. aurita*, *S. caprea*, *S. lapponum*, *S. pentandra*, *S. phyllifolia*, *S. pyrolaeifolia*, *S. repens*, *S. sieboldiana*, *S. smithiana*, *Larix decidua*, *L. occidentalis*, and *L. sibirica* in Europe and Japan.
- Melampsora larici-epitea** Kleb. Leaf rust on *Salix* spp. (33 species listed) and *Larix decidua* in Siberia, India, Ceylon, and Europe.
- Melampsora larici-pentandrae** Kleb. Yellow to brown leaf rust on *S. amygdalina*, *S. frogilis*, *S. humboldtiana*, *S. jaspidea*, *S. pentandra*, *S. purpurea*, *Larix decidua*, and *L. sibirica* in Uruguay and Europe.
- Melampsora larici-urbaniana** Mat. Leaf rust on *S. urbaniana* and *Larix* sp. in Japan.
- Melampsora microsora** Diet. Yellow to brown rust pustules on leaves of *S. nipponica* in Japan.
- Melampsora repentis** Plowr. See Orchidaceae.
- Melampsora reticulatae** Blytt. Leaf rust on *S. reticulata* and *Saxifraga aizoides* in Sweden, Norway, Switzerland, Austria, and Hungary.
- Melampsora ribesii-purpureae** Kleb. Rust on leaves of *Salix* spp. and *Ribes* spp. in Europe, Japan, and Utah and Colorado on *Ribes*.
- Melampsora ribesii-viminalis** Kleb. Golden-yellow to brown rust pustules on leaves of *S. viminalis*, *purpurea*, *Ribes alpinum*, *R. aureum*, *R. nigrum*, *R. rubrum*, *R. sanguineum*, and *Grossularia* in Europe.
- Melampsora salicis-albae** Kleb. Yellow and brown rust pustules on leaves of *Allium cepa*, *A. porrum*, *A. schoenoprasum*, *A. ursinum*, *A. vinealis*, and on leaves and twigs of *Salix alba* in India and Europe.
- Melampsora salicina** Lév. Black stromata on leaves of *S. purpurea* and *S. viminalis* in Italy.
- Mycosphaerella genuflexa** Auersw. On leaves of *S. alba* in Germany.
- Mycosphaerella salicicola** (Fr.) Fckl. On leaves of *S. caprea*, *S. nigricans*, and *S. triandra* in France, Italy, and Germany.
- Phoma intricans** Schwarz. Causes a die-back of twigs, with resulting death of leaves of *S. alba* var. *vitellina-pendula* in Holland.
- Phyllohendersonia foliorum** (Fckl.) Tass. On *S. caprea* in France, Italy, and Germany.
- Phyllosticta iserana** Kab. and Bub. Circular to angular brown-gray leaf spots on *S. fragilis* in Bohemia.
- Phyllosticta salicicola** Thuem. Subcircular gray-white leaf spots on *S. alba* and *S. triandra* in Denmark, Russia, Poland, and France.
- Phyllosticta translucens** Bub. and Kab. Angular gray-white leaf spots with yellow-brown margins on *S. nigricans* in Austria.
- Physalospora miyabeana** Fuku. Elliptical whitish-gray stem lesions, 5 to 30 millimeters long surrounded by black borders, on *S. purpurea* in Japan. These lesions may unite and girdle infected branches. Circular to irregular, dark-brown, often concentrically zoned areas occur on the leaves.
- Podosphaera schlechtendalii** Lév. Powdery mildew on leaves of *S. alba* and *S. viminalis* in France and Russia.
- Ramulaspera salicina** (Vestergr.) Lind. Violet leaf spots on *S. cinerea* and *S. hastata* in Sweden, Finland, and Austria.
- Septobasidium acaciae** Saw. See Acacia.
- Septoria blennorioides** (Karst.) Berl. and Vogl. On leaves of *S. pendula* in Finland.
- Septoria capreae** West. Angular gray-white leaf spots with rufous borders on *S. caprea*, *S. cinerea*, and *S. atrocinerea* in Belgium and Portugal.
- Septoria jennisseica** Thuem. Circular, then confluent, dull-brown to gray leaf spots on *Salix* sp. in Siberia.
- Septoria salicicola** (Fr.) Sacc. Circular brown leaf spots on *S. caprea*, *S. cinerea*, *S. grandifolia*, *S. pyrolaeifolia*, and *S. viminalis* in Russia, Siberia, Denmark, Sweden, France, Italy, and Austria.
- SALOA.** See Blumenbachia.
- SALPICHROA.** Cockseeds. Herbs and shrubs, sometimes under cultivation.
- Puccinia pampeana** Speg. Rust on leaves and stems of *S. rhomboidea* in Argentina, Brazil, and Uruguay.
- Septoria bonariensis** Speg. Circular to irregular gray-white leaf spots on *S. rhomboidea* in Argentina.
- SALPIGLOSSIS.** Flower-garden annuals.
- Phytophthora parasitica** Dast. See Ricinus.
- SALVIA.** SAGE. Herbs and subshrubs, with some economic and ornamental species.
- Aphelenchus oleisistus** Ritz. Bos. See Begonia.
- Ascochyta vicina** Sacc. See Rumex.
- Cercospora salviae** Pat. Brown leaf spots on *Salvia* sp. in Ecuador.
- Coleosporium salviae** Diet. Orange rust pustules on leaves of *S. bipinnata* and *S. japonica* in Japan.
- Puccinia badia** Holw. Leaf rust on *S. albicans* and *S. chrysantha* in Mexico.
- Puccinia bithynica** P. Magn. Leaf rust on *S. bithynica*, *S. grandiflora*, and *S. pomifera* in Russia and Crete.
- Puccinia conspersa** Diet. Brown rust pustules on small yellow sunken leaf spots on *Salvia* sp. in Brazil.
- Puccinia delavayana** Pat. and Har. Black rust sori on leaves of *Salvia* sp. in China.
- Puccinia delicatula** (Arth.) Sacc. and Trott. Leaf rust on *S. cinnabarina*, *S. elegans*, *S. holwayi*, and *S. pulchella* in Mexico and Guatemala.
- Puccinia diutinum** Mains. and Holw. Brown rust pustules on leaves of *S. chrysantha*, *S. pittieri*, and *S. scordoniaefolia* in Mexico and Costa Rica.
- Puccinia filiola** Mains. and Holw. Leaf rust on *S. involucrata* and *S. pulchella* in Guatemala.
- Puccinia gentilis** Arth. Cinnamon-brown to dark-brown rust pustules on leaves of *S. alamosana* in Mexico.
- Puccinia gilliesii** Speg. Brown rust pustules on leaf blades and petioles of *S. gilliesii* in Argentina.
- Puccinia griseola** Lagh. Leaf rust on *Salvia* sp. in Ecuador.
- Puccinia impedita** Mains. and Holw. Leaf rust on *S. hyptioides*, *S. occidentalis*, and *S. tiliaefolia* in Guatemala, Costa Rica, and Trinidad.
- Puccinia infrequens** Holw. Brown rust pustules on leaves of *S. cinnabarina* in Mexico and Guatemala.
- Puccinia mitrata** Syd. Dark-brown powdery rust pustules on leaves of *S. fluvialis*, *S. mexicana*, *S. polystachya*, *S. purpurea*, *S. sessiliflora*, and *S. tiliaefolia* in Mexico and Guatemala.
- Puccinia nigrescens** Kirchn. Brown to black rust pustules on elongated sunken spots on leaves and petioles of *S. judaica*, *S. verticillata*, and *S. virgata* in Europe and Palestine.
- Puccinia nipponica** Diet. Leaf rust on *S. nipponica* in Japan.
- Puccinia nevadensis** Syd. Leaf rust on *S. lavandulaefolia* in Spain.
- Puccinia nivea** Holw. Brown rust pustules on leaves of *S. purpurea* in Mexico.
- Puccinia paramensis** Mayor. Leaf rust on *S. cernua* in Colombia.

**SALVIA**—Continued.

- Puccinia prospera** Arth. Brown rust pustules on leaves of *S. microphylla* in Mexico.  
**Puccinia roesteliiformis** Lagh. Yellow and brown rust pustules on leaves of *Salvia* sp. in Ecuador.  
**Puccinia salviae** Ung. Brown leaf rust on *S. glutinosa* and *S. lavandulaefolia* in China and Europe.  
**Puccinia soledadensis** Mayor. Leaf rust on *S. pauciserrata* in Colombia.  
**Septoria salviae** Pass. Dull-brown leafspots on *S. horminum*, *S. pratensis*, *S. sylvestris*, and *S. sclarea* in Russia, Italy, Dalmatia, and France.  
**Ustilago betonica** Beck. Anther smut of *Betonica* spp. and *Salvia* sp. in Europe.
- SAMBUCUS**. ELDER. Shrubs grown for their foliage, flowers, and black and red berries.  
**Ascochyta fernandi** Bub and Malk. Circular to irregular gray-brown leaf spots on *S. ebulus* in Bulgaria.  
**Ascochyta sambucella** Pass. On branches of *S. ebulus*, *S. nigra*, and *S. racemosa* in Germany and Italy.  
**Ascochyta sambuci** (Sacc.) Tass. On leaves of *S. nigra* in Italy and France.  
**Cercospora ticinensis** Cav. Gray-brown leaf spots on *S. nigra* in Italy.  
**Exosporium depazeoides** Desm. On leaves of *S. nigra* in France.  
**Gloeosporium ebuli** Allesch. Subcircular to irregular, then confluent, brown leaf spots on *S. ebulus* in Germany.  
**Helicobasidium mompa** Tan. See *Morus*.  
**Marsonia sambuci** E. Rostr. Large brown leaf spots on *S. nigra* in Europe.  
**Mycosphaerella ebulina** Petr. On leaves of *S. ebulus* in Bohemia and Austria.  
**Phyllosticta ebuli** Fekl. On leaves of *S. ebulus* and *S. racemosa* in France, Italy, Hungary, and Germany.  
**Phyllosticta sambucina** Allesch. On leaves of *S. nigra* in Switzerland.  
**Septoria ebuli** Desm. and Rob. Small subcircular pale-rufous to purple, often confluent, leaf spots on *S. ebulus* and *S. nigra* in Europe.
- SANCHEZIA**. Strong, erect herbs or half shrubby plants grown for their flowers and foliage.  
**Pyrenochaete vexans** Syd. Large confluent brown leaf spots on *S. nobilis* in Brazil.
- SANGUISORBA**. See also *Poterium*. BURNET. Perennial herbs grown in hardy borders and as salad plants.  
**Marsonia sennensis** Gz. Frag. On leaves of *S. officinalis* in Spain.  
**Ovularia bulbiger** (Fekl.) Sacc. Ochraceous leaf spots with dark-purple margins on *S. minor* in Alaska, Switzerland, and Germany.  
**Peronospora sanguisorbae** Gäum. Downy mildew on leaves of *S. officinalis* in central Europe.  
**Phragmidium carbonarium** (Schlecht.) Wint. Golden and black rust pustules on leaves of *S. canadensis*, *S. carnea*, *S. officinalis*, and *S. tenuifolia* in Japan, Siberia, and Europe.  
**Phragmidium minor** (Arth.) Syd. Dark-brown rust pustules on leaf blades and petioles of *S. latifolia* in Alaska.  
**Phragmidium sanguisorbae** (DC.) Schroet. See *Poterium*.
- SANSEVIERIA**. BOWSTRING HEMP. Herbaceous perennials cultivated as ornamentals and sometimes as fiber plants.  
**Chaetophoma sansevieriae** Tass. On leaves of *S. zeylanica* in Italy.  
**Leptosphaeria baldratiana** Bacc. Small brown spots on leaves of *S. ehrenbergia* in Abyssinia.  
**Phyllachora pappiana** Bacc. Small subcircular black stromata on leaves of *S. ehrenbergia* in Abyssinia.  
**Septogloeum concentricum** Syd. On leaves of *S. guineensis* in Abyssinia and central Africa.
- SANTALUM**. SANDALWOOD. Evergreen trees with aromatic properties.  
**Spik disease**. This disease, characterized by die-back, unthriftness, and final death of infected trees, is a serious disease of sandalwood (*S. album*) in India. The cause is unknown, but the disease is apparently one of the infectious mosaics.
- SANTOLINA**. LAVENDER COTTON. Shrubs and herbs, sometimes cultivated.  
**Puccinia chamaecyparissi** Trott. Rust on leaves and stems of *S. chamaecyparissus* in Italy.
- SAPINDUS**. SOAPBERRY. Trees and shrubs cultivated for ornament and economic value.  
**Mycosphaerella reyesii** Syd. On leaves of *S. saponaria* in the Philippines.  
**Phyllosticta patouillardii** Sacc. and D. Sacc. Circular to angular, bright-brown leaf spots with dull-brown margins on *Sapindus* sp. in Algeria.  
**Phyllosticta raimundi** Sacc. Leaf spot on *Sapindus* sp. in the Philippines.  
**Phyllosticta sapindi** P. Henn. Circular pale-brown leaf spots on *S. saponaria* in Brazil.  
**Uredo cristata** Speg. Leaf rust on *S. bifoliatus* and *Sapindus* sp. in Ceylon and Argentina.
- SAPIUM**. CHINESE TALLOW TREE. Tropical trees and shrubs cultivated for their economic products.  
**Cercospora micromera** Syd. On leaves of *S. sebiferum* in China.  
**Cercospora sapiicola** Speg. Angular brown leaf spots on *S. aucuparium* and *S. mannianum* in Argentina and Uganda.  
**Exosporium lateritium** Syd. On leaves of *S. abyssinicum* in Tanganyika.  
**Helminthosporium sapii** Miy. On leaves of *S. sebiferum* in China.  
**Uromyces cisnerosanus** Speg. Dark-brown to black rust pustules on leaves of *S. aucuparium*, *S. biglandulosum*, and *S. salicifolium* in Argentina and Paraguay.  
**Uromyces globosus** Diet. and Holw. Powdery black rust pustules on leaves of *S. biglandulosum* in Mexico.  
**Uromyces vestitus** Diet. Brown rust pustules on leaves and stems of *Sapium* sp. in Brazil.
- SAPONARIA**. SOAPWORT. Hardy, often coarse, annual or perennial, erect or decumbent herbs, sometimes cultivated.  
**Ascochyta dianthi** Berk. See *Dianthus*.  
**Ascochyta saponariae** Fekl. Small dull-brown leaf spots on *S. officinalis* in Germany.  
**Cylindrosporium saponariae** Roum. Large circular, often confluent, gray leaf spots on *S. officinalis* in France.  
**Phyllosticta saponariae** (Fekl.) Sacc. On leaves of *S. officinalis* in Belgium, Italy, and Germany.  
**Septoria saponariae** (DC.) Sav. and Becc. Pale-brown circular to irregular leaf spots on *S. officinalis* and *Silene latifolia* in Bulgaria, Yugoslavia, Great Britain, Denmark, and Belgium.  
**Uromyces gypsophilae** Cke. See *Gypsophila*.
- SARCANTHUS**. See *Orchidaceae*.
- SASA**. See *Bambuseae*.
- SATYRIUM**. See *Orchidaceae*.
- SAURURUS**. LIZARD'S-TAIL. Perennial marsh herb.  
**Uromyces saururi** P. Henn. Leaf rust of *S. loureiri* in Japan.
- SAUSSUREA**. Annual, biennial, or perennial herbs, sometimes planted in gardens.  
**Aecidium saussureae-affinis** Diet. Leaf rust on *S. affinis* in Japan.  
**Bremia saussureae** Saw. Downy mildew on pale-yellow polygonal leaf spots on *S. carthamoides* in Formosa.

## SAUSSUREA—Continued.

- Coleosporium saussureae** Thuem. Golden-yellow rust pustules on leaves of *S. grandifolia*, *S. japonica*, *S. latifolia*, *S. maximowiczii*, *S. triangulata* and *S. ussuriensis* in Siberia, Japan, and China.
- Erysiphe taurica** Lév. See *Althaea*.
- Phyllosticta saussureae** Thuem. On leaves of *S. salicifolia* in Siberia.
- Puccinia rupestri** Juel. Rust on leaves of *S. alpina* and *Carex rupestris* in Sweden and Norway.
- Puccinia saussurea** Thuem. Powdery dark-brown rust pustules on leaves of *S. glomerata* in Siberia and India.
- Puccinia saussureae-alpinae** Lindr. Brown rust sori on leaves of *S. alpina* in Finland.
- Puccinia vaginatae** Juel. Brown to black rust pustules on leaves of *S. alpina*, *Carex panicea*, and *C. vaginata* in Sweden.
- Puccinia vomica** Thuem. Leaf rust on *S. tanakae* and *Saussurea* sp. in Siberia.
- Ramularia saussureae** Thuem. Subcircular gray leaf spots on *S. glomerata* in Siberia.
- Septoria saussureae** Thuem. Circular gray leaf spots with dull-brown margins on *S. glomerata* in Siberia and China.
- Thecaphora traili** Cke. Powdery purple-brown smut sori in flowers of *S. alpina* and *Carduus heterophyllus* in Scotland.
- Uromyces saussureae** Karst. Leaf rust on *S. circa*, *S. loureiri*, *S. nipponica*, and *Saussurea* sp. in Japan and Siberia.
- SAXIFRAGA. SAXIFRAGE.** Perennial herbs cultivated in gardens and rockeries.
- Caecoma cernuae** Th. Lind. Yellow leaf rust on *S. cernua* in Lapland.
- Cylindrosporium microspermum** (Speg.) Sacc. Indefinite circular leaf spots on *S. rotundifolia* in Italy.
- Diplochorella stromatica** (Rehm.) Theiss. and Syd. On leaves of *Saxifraga* sp. in Austria.
- Entyloma schinzianum** (P. Magn.) Bub. Leaf smut on *S. heuffeliana* and *S. rotundifolia* in Asia Minor, Switzerland, Yugoslavia, Austria, and Hungary.
- Leptosphaeria monotis** Rehm. On stems of *Saxifraga* sp. in Austria.
- Melampsora hirculi** Lindr. Rust on stems and leaves of *S. hirculus* in Russia, Finland, and Switzerland.
- Melampsora reticulata** Blytt. See *Salix*.
- Melampsora vernalis** Niessl. Golden, then dark-brown, rust pustules on leaves of *S. granulata* in Europe.
- Peronospora minima** G. W. Wils. Downy mildew on leaves of *S. cernua* in Norway.
- Peronospora saxifragae** Bub. Downy mildew on leaves of *S. granulata* in Bohemia and Germany.
- Phyllosticta groenlandica** Allesch. On leaves of *S. aizoides*, *S. oppositifolia*, and *S. stellaris* in Greenland.
- Phyllosticta saxifragae** P. Brun. Circular gray-white leaf spots with brown margins on *S. geum* in France.
- Phyllosticta saxifragicola** P. Brun. Circular leaf spots with brown margins on *S. cordifolia* in France.
- Puccinia fischeri** Crut. and Mayor. Leaf rust on *S. biflora* in Switzerland.
- Puccinia laurentiana** Trel. Leaf rust on *S. neglecta stolonifera* in Alaska.
- Puccinia huteri** Syd. Powdery brown leaf rust on *S. mutata* in Austria.
- Puccinia jueliana** Diet. Dark-brown rust pustules on leaves of *S. aizoides* and *S. oppositifolia* in Norway and Austria.
- Puccinia pazschkei** Diet. Powdery dull-brown rust sori on leaves of *S. aizoon*, *S. elatior*, *S. longifolia*, and *S. mutata* in Switzerland and Austria.
- Puccinia saxifragae** Schlecht. Powdery brown leaf rust on *Saxifraga* spp. in Japan and Europe. Also reported from Oregon.
- Puccinia saxifragae-ciliatae** Barel. Brown rust pustules on circular brown sunken leaf spots on *S. ciliata* and *S. lingulata* in India.
- Puccinia saxifragae-micranthae** Barel. Brown rust sori on leaves of *S. micrantha* in India.
- Puccinia saxifragae-tricuspidatae** P. Henn. Yellow-brown leaf rust on *S. tricuspidata* in Greenland.
- Ramularia saxifragae** Syd. Large brown leaf spots on *S. granulata* in Europe.
- Ramularia sennensis** Gz. Frag. On leaves of *S. rotundifolia* in Spain.
- Septoria saxifragae** Pass. and forma *arctica* Allesch. On leaves of *S. aizoides*, *S. hirculus*, and *S. rotundifolia* in Greenland and Italy.
- Synchytrium rubrocinctum** P. Magn. Small galls on leaves of *S. granulata* in Germany.
- Synchytrium groenlandicum** Allesch. On leaves of *S. cernua* in Greenland.
- SCABIOSA.** Annual or perennial herbs, cultivated in gardens for their showy flowers. Sometimes called mourning bride.
- Aphelenchus olesistus** Ritz. Bos. See *Begonia*.
- Cercospora scabiosicola** Rangel. On leaves of *S. atropurpurea* in Brazil.
- Peronospora knautiae** Fekl. Downy mildew on leaves of *S. columbaria*, *Knautia arvensis*, and *K. silvatica* in Switzerland, France, Scandinavia, Austria, and Germany.
- Peronospora violacea** Berk. See *Dipsacus*.
- Ramularia bosniaca** Bub. Small circular to oblong, often confluent, dark-violet, then gray-white, leaf spots on *S. columbaria* in Yugoslavia.
- Ramularia knautiae** (Mas.) Bub. On leaves of *S. columbaria* and *S. lucida* in Dalmatia and Austria.
- Ramularia scabiosa** Lind. (*R. scabiosae* Jaap.) Subcircular leaf spots on *S. columbaria* and *S. lucida* in Denmark and Switzerland. Jaap has also described what is apparently the same species under the same name.
- Septoria scabiosicola** Desm. Circular purple-brown leaf spots on *S. arvensis*, *S. atropurpurea*, *S. balansae*, *S. columbaria*, *S. kladnikiana*, *S. ochroleuca*, and *S. succisa* in Europe.
- Stigmatea scabiosae** Lebed. On leaves of *S. ucranica* in Russia.
- Ustilago intermedia** Schroet. A smut destroying flowers of *S. arvensis* and *S. columbaria* in Denmark, Belgium, Great Britain, and Germany.
- Ustilago scabiosa** (Sow.) Wint. A smut forming powdery spore masses in anthers of *S. columbaria* and *Knautia arvensis* in Europe.
- SCAEVOLA.** Herbs or shrubs, sometimes cultivated.
- Phyllosticta scaevolae** E. and E. Circular dull-brown leaf spots on *S. chamissoniana* in Hawaii.
- Ramularia scaevolae** Rac. Circular pale-brown leaf spots on *S. koenigii* in Java.
- Uromyces puccinioides** Berk. Rust on leaves and flower stalks of *Scaevola* sp. and *Selliera radicans* in Queensland.
- SCANDIX.** Umbelliferous annual herbs cultivated for their flowers and foliage.
- Uredo petroselini** DC. Leaf rust on *S. pecten-veneris* in France and Great Britain.
- SCHINUS. PEPPER TREE.** Dioecious, resinous, ornamental trees.
- Cercospora schini** Syd. On leaves of *S. dependens* in Argentina.
- Meliola ludibunda** Speg. Black fungus patches on leaves of *S. dependens* in Brazil.
- Phyllosticta schini** Thuem. Large irregular ochraceous to dull-brown leaf spots on *S. latifolius* and *S. molle* in Italy and Portugal.
- Septoria schini** Tassi. Ashen leaf spots with black borders on *S. latifolius* in Italy.

- SCHIZANTHUS.** BUTTERFLY FLOWER. Erect, half hardy, annual herbs grown for their showy flowers.  
**Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- SCHIZOCODON.** Fringe bell. Glabrous perennial herbs.  
**Puccinia schizocodonis** Pat. Brown to black rust pustules on leaves of *S. soldanelloides* in Japan.
- SCHIZOLOBIUM.** Tall leguminous trees.  
**Phyllachora schizolobii** P. Henn. Black stromata on leaves of *S. parahybum* (*S. excelsum*) in Brazil.
- SCHIZONOTUS.** See Sorbaria.
- SCHIZOPHRAGMA.** Ornamental vines grown for their handsome green foliage and showy clusters of white flowers.  
**Mycosphaerella exigua** Syd. On leaves of *S. hydrangeoides* in Japan.
- SCHIZOSTACHYUM.** See Bambuseae.
- SCILLA.** SQUILL. Perennial, bulbous plants with blue, rose, or white flowers. (Including *Endymion*)  
**Accidium doidgei** Syd. Leaf rust on *S. saturata* in the Union of South Africa.  
**Accidium scillae** Fckl. Yellow rust pustules on leaves of *S. bifolia* in England, France, and Switzerland. Said to be distinct from *Uromyces scillarum*, a common rust on Scillas.  
**Accidium scillinum** Dur. and Mont. This rust, reported on leaves of *S. autumnalis*, is probably only one stage of one of the *Scilla* rusts recorded elsewhere.  
**Physoderma debeauxii** (Bub.) Syd. Ellipsoidal spots on the leaves of *S. maritima* (*Urginea scilla*) in Italy. The leaves are injured and bulb development is prevented.  
**Puccinia ficilhoana** Lagh. Elongated black rust pustules on both leaf surfaces of *S. companulata* (*Endymion companulatum*) in Portugal.  
**Puccinia liliacearum** Duby. See Ornithogalum.  
**Puccinia rossiana** (Sacc.) Lagh. Rufous-brown rust pustules on both leaf surfaces of *S. bifolia* and *S. cernua* in Switzerland, Italy, Russia, and Hungary.  
**Sclerotinia tuliparum** (Wakk.) Rehm. See Tulipa.  
**Septoria scillae** West. See Muscari.  
**Urocystis colchici** (Schlecht.) Rab. See Colchicum.  
**Uromyces algeriensis** Syd. Circular to irregular cinnamon-colored rust pustules on both leaf surfaces of *S. obtusifolia* and other species of *Scilla* in Tunis and Algeria.  
**Uromyces circinalis** Kalchbr. and Cke. This rust produces broad spots from 2 to 8 centimeters in diameter in which powdery, cinnamon-colored pustules form. On both leaf surfaces of *S. prasina* in the Union of South Africa.  
**Uromyces scillarum** (Grev.) Wint. See Hyacinthus.  
**Ustilago vailantii** Tul. See Hyacinthus.
- SCOLYMUS.** Spanish oyster plant. Fleshy, edible roots.  
**Phyllosticta scolymi** Pat. Angular ashen leaf spots on *S. hispanicus* in Tunis.  
**Puccinia scolymi** Syd. Powdery brown rust sori on leaves of *S. grandiflorus* and *S. hispanicus* in north Africa, Syria, Spain, and Sicily.  
**Ramularia cynaræ** Sacc. Gray spots with brown margins on leaves of *S. hispanicus* in Chile, France, and French North Africa.  
**Ustilago scolymi** Roum. Powdery black masses of smut spores in flowers and receptacles of *S. grandiflorus* and *S. hispanicus* in Algeria and Tunis.
- SCOPOLIA. SCOPOLINA.** Ag. Erect, hardy, perennial herbs cultivated for their lurid-purple flowers.  
**Accidium pascheri** Bub. Rust on large round gray leaf spots on *S. japonica* in Japan.  
**Ramularia scopaliae** Vos. Irregular brown spots on leaves of *S. atropoides* in Austria.
- SCOPOLINA.** See Scopolia.
- SCORPIURUS.** Hardy stemless or decumbent herbs.  
**Cercospora scorpiuri** Thuem. Subcircular reddish-gray leaf spots with dull-brown margins on *S. muricata* in Portugal.
- SCORZONERA.** SERPENT ROOT. BLACK SALSIFY. Herbs cultivated for the fleshy, edible roots.  
**Ascochyta scorzonerae** Rostr. Irregular brown-leaf spots on *S. hispanica* in Denmark.  
**Cercospora scorzonerae** v. Hoeh. Oblong pale-brown leaf spots with broad purple margins on *S. humilis* in Austria.  
**Phyllosticta scorzonerae** Pass. Small brown leaf spots on *S. humilis* in France and Bohemia.  
**Puccinia angustifolia** McAlp. Yellow and dark-brown rust pustules on leaves of *S. angustifolia* in Australia.  
**Puccinia scorzonerae** (Schum.) Jacky. Golden to brown rust pustules on leaves of *S. austriaca*, *S. hispanica*, *S. humilis*, *S. purpurea*, and *S. rosea* in China and Europe.  
**Ramularia scorzonerae** Jaap. Circular to elongate dark-brown leaf spots on *S. aristata* and *S. humilis* in Austria and Spain.  
**Sporidesmium scorzonerae** Aderh. Attacks stems and leaves of *Scorzonera* sp. in Germany.  
**Ustilago receptaculorum** Tul. A smut destroying the flowers of *Scorzonera* sp. in France.  
**Ustilago scorzonerae** (Alb. and Schw.) Schroet. Dark-violet smut spores in flowers of *S. aristata*, *S. hispanica*, *S. humilis*, and *S. purpurea* in Europe.
- SCROPHULARIA.** FIGWORT. Herbs, occasionally cultivated.  
**Ascochyta scrophulariae** Kab. and Bub. Round to irregular zoned brown leaf spots on *S. nodosa* in Bulgaria and Bohemia.  
**Phyllosticta ehrhartii** Sacc. Gray-white sinuous leaf spots on *S. aquatica* and *S. ehrhartii* in Portugal and Italy.  
**Phyllosticta scrophulariae** Sacc. Ochraceous leaf spots on *S. nodosa* in Denmark, Russia, and Italy.  
**Phyllosticta scrophulariae-bosniacae** Bub. Subcircular to ellipsoid yellow-brown leaf spots on *S. bosniaca* in Yugoslavia.  
**Phyllosticta scrophularinea** Sacc. Small angular white leaf spots on *S. kakudensis* and *S. nodosa* in Japan and Italy.  
**Ramularia nicolai** Bub. (*Ovularia carneola* Sacc.). Angular to irregular leaf spots on *S. bosniaca*, *S. ehrhartii*, *S. nodosa*, and *S. vernalis* in Yugoslavia, Switzerland, Spain, Denmark, and Austria.  
**Ramularia scrophulariae** Fautr. and Roum. Purple leaf spots on *S. aquatica* and *S. nodosa* in France and Bulgaria.  
**Septoria rohlenae** Bub. Angular, often confluent, gray leaf spots with brown margins on *S. scopoli* in Yugoslavia and Albania.  
**Uromyces scrophulariae** (DC.) Fckl. Yellow and brown rust pustules on leaf blades, petioles, and stems of *S. aquatica*, *S. bosniaca*, *S. ebullifolia*, *S. ehrhartii*, *S. nodosa*, and *S. scorodonia* in Europe.
- SCUTELLARIA.** SKULLCAP. Annual and perennial herbs.  
**Accidium scutellariae** Syd. Leaf rust on *S. angulosa* and *S. repens* in India.  
**Accidium scutellariae-indicae** Diet. Leaf rust on *S. indica* in Japan.  
**Erysiphe taurica** Lév. See Althaea.

**SEBASTIANA.** Tropical shrubs.

**Exoascus sebastianae** Sadeb. Fruiting on lower surfaces and often deforming leaves of *S. brasiliensis* in Brazil.

**Phyllachora subtropica** Speg. Black stromata on leaves of *Sebastiana* sp. in Brazil and Paraguay.

**Puccinia sebastianae** Syd. Brown rust pustules on leaves of *S. klotzschiana* in Brazil.

**Uredo sebastianae** Wint. Leaf rust on *S. microstachys* in Brazil.

**SECALE.** RYE.

**Acremoniella occulta** Cav. Brownish-yellow flakes on culms of *Secale* and *Triticum* in Great Britain and Italy.

**Bacillus cerealinum** Gentner. See *Hordeum*.

**Dilophia graminis** (Fekl.) Sacc. See *Triticum*.

**Fusarium nivale** (Fr.) Sor. (*Calonectria graminicola* B. and B.) Attacks the culms of *S. cereale*, *Hordeum*, *Lolium*, and *Triticum* in Europe, causing the so-called "snow disease."

**Gloeosporium secalis** Rostr. Gray-white leaf spots on *S. cereale* in Denmark.

**Leptosphaeria eustoma** (Fr.) Sacc. On culms and peduncles of *S. cereale*, *Holcus sorghum*, and *Phragmites* sp. in Europe.

**Leptosphaeria herpotrichoides** DeN. See *Triticum*.

**Mycosphaerella basicola** Frank. See *Hordeum*.

**Ophiobolus cariceti** (B. and Br.) Sacc. See *Triticum*.

**Ophiobolus herpotrichoides** (Fr.) Sacc. See *Triticum*.

**Sclerotinia temulenta** Prill. and Delacr. (*Hymenoscypha temulenta* P. and D.) Forms sclerotia in place of grain in *S. cereale* in France. Flour made from grain contaminated with these sclerotia is poisonous.

**Septocylindrium secales** Oud. Pale-brown leaf spots on *S. cereale* in Holland.

**Sphaeroderma damnosum** Sacc. See *Triticum*.

**Tilletia secalis** (Cda.) Kueh. Stinking smut of rye (*S. cereale*) in Europe. The smut sori take the place of the grain and are composed of a dense black mass of spores. Reported from Washington and New York.

**Tylenchus dipsaci** Kuehn. See *Narcissus*.

**Tylenchus tritici** Bast. See *Triticum*.

**SECHIUM.** CHAYOTA Ag. CHAYOTE. Cucurbitaceous tropical vines grown for their edible single-seeded fruit.

**Cercospora sechiae** Stevenson. Yellow subcircular, then irregular and indefinite, areas on leaves of *S. edule* (*C. edulis*) in Porto Rico and Cuba.

**Helminthosporium sechiicolum** Stevenson. Small angular brown spots on leaves of *S. edule* (*C. edulis*) in Porto Rico.

**Phyllosticta sechii** Young. Irregular, then confluent, yellowish-white to gray-white leaf spots on *S. edule* (*C. edulis*) in Porto Rico.

**SECURINEGA.** Shrubs with bright-green foliage.

**Nothoravenelia japonica** Diet. Brown rust pustules on leaves of *S. flueggeoides* and *Flueggea japonica* in Japan and China.

**SEDUM.** STONECROP. Hardy, perennial, succulent herbs grown in rockeries.

**Aecidium koreaense** P. Henn. Leaf rust on *S. maximowiczii* in Chosen.

**Aecidium sedi-aizoonis** Tranzsch. Leaf rust on *S. aizoon* in Russia.

**Ascochyta sedi** Allesch. On stems of *S. affine* and *S. maximum* in Austria and Germany.

**Ascochyta telephii** Vestergr. Circular to irregular brown leaf spots with dark-purple margins on *S. telephium* in Sweden.

**Endophyllum sedi** (DC.) Lév. Brown leaf rust on *S. acre*, *S. nicaeense* (*S. altissimum*), *S. boloniense*, *S. maximum*, *S. pruinatum*, *S. purpureum*, *S. reflexum*, and *S. sexangulare* in Europe.

**Euryachora thoracella** (C. Rust.) Schroet. Shiny black stromata, circular on the leaves and elongate on the stems of *S. fabaria*, *S. maximum*, *S. purpureum*, and *S. telephium* in Italy, Switzerland, Austria, Hungary, and Germany.

**Helminthosporium rhopaloides** Fres. and f. *sedi* P. Brun. On stems of *S. fabaria* in France.

**Mycosphaerella sedi** Jacz. On leaves of *S. telephium* in Russia.

**Phyllosticta aizoi** Cke. Small subcircular brown leaf spots on *S. aizoon* in Great Britain.

**Puccinia australis** Koern. Leaf rust on *S. boloniense*, *S. cepaca*, *S. mite*, *S. reflexum*, and *Molinia serotina* in Dalmatia, Italy, and Austria.

**Puccinia benkei** Kusano. Leaf rust on *S. alboroseum* and *S. telephium* in Japan.

**Puccinia longissima** Schroet. Leaf rust on *S. acre*, *S. boloniense*, *S. maximum*, *S. purpurascens*, *S. reflexum*, and *Koeleria* spp. in Sweden, Austria, Hungary, and Germany.

**Puccinia sedi** Koern. Dark-brown rust pustules on leaves and stems of *S. elegans* and *S. glanduliferum* in Italy, France, Switzerland, Belgium, and Germany.

**SELAGINELLA.** CLUB MOSS.

**Helminthosporium filicicola** P. Henn. See *Lygodium*

**Phyllosticta helvetica** Sacc. On leaves of *S. helvetica* in Dalmatia.

**Phyllosticta selaginellae** Sacc. Indefinite pale-brown leaf spots on *S. helvetica* in Italy.

**Sclerotium selaginellae** Peyl. On leaves and stems of *Selaginella* sp. in Bohemia.

**Synchytrium selaginellae** Sorok. Doubtful species on *Selaginella* sp. in Russia.

**SELINUM.** Umbelliferous branched perennial herbs.

**Aecidium selini** Lindr. Yellow leaf rust on *S. lineare* in Finland.

**Physoderma vagans** Schroet. See *Ranunculus*.

**Puccinia apii** Desm. See *Apium*.

**SEMELE.** CLIMBING BUTCHER'S-BROOM. Ornamental shrubs.

**Ascochyta semeles** Sacc. Gray-white oblong leaf spots with rufous margins on *S. androgyna* in Italy.

**Phyllosticta semeles** Ohl. Numerous white leaf spots with red borders on *S. androgyna* in Russia. Possibly only a stage of the preceding species.

**SEMPERVIVUM.** HOUSELEEK. Perennial stemless herbs or subshrubs with thick fleshy leaves.

**Ascochyta sempervivi** Fautr. On leaves and stems of *S. tectorum* in Denmark and France.

**Endophyllum sempervivi** (A. and S.) De By. Rust on leaves of *Sempervivum* spp. (about 20 species so far reported) in Europe. Introduced into New York City, where it has done considerable damage in public plantings.

**SENECIO.** GROUNDSEL. Composites with yellow flower heads.

**Aecidium dubiosum** Syd. Leaf rust on *S. mikanioides* in the Union of South Africa.

**Aecidium halophilum** Speg. Leaf rust on *S. alicornis* in Chile.

**Aecidium hualtatium** Speg. Yellow leaf rust on *S. gilliesii*, *S. glabrum*, *S. halorrhagis*, *S. hieracioides*, *S. hualtata*, and *S. thermarum* in Chile and Argentina.

**Aecidium kilimandscharicum** P. Henn. Yellow leaf rust on *S. multicorymbosus* in central Africa.

**Aecidium margueryanum** Maire. Rust on leaves of *S. aquaticus* in France.

**Aecidium permultum** Syd. Leaf rust on *S. inornatus* in the Union of South Africa.

**Aecidium senecionis-stenocephali** Diet. Leaf rust on *S. stenocephalus* in Japan.

**SENECIO**—Continued.

- Aecidium serrae** Syd. Leaf rust on *S. serra* in the Union of South Africa.
- Ascochyta senecionis** Fekl. On leaves of *S. sarracenicus* in Germany.
- Baeodromus holwayi** Arth. Leaf rust on *S. cinerarioides* in Mexico.
- Baeodromus senecionis** Syd. Leaf rust on *Senecio* sp. in Ecuador.
- Bremia tulasnei** (Hoffm.) Syd. Downy mildew on leaves of *S. vulgaris* in Germany.
- Cercospora jacquiniana** Thuem. Irregular ochraceous leaf spots on *S. jacquinianus* in Germany.
- Coleosporium senecionis** Fr. Yellow rust pustules on leaves of *Senecio* spp. and *Cineraria* spp. in Europe, Brazil, Uruguay, Chile, Siberia, Manchuria, Japan, and the Canary Islands. Collected once in Rhode Island. The aecial state occurs on needles of *Pinus austriaca*, *P. insignis*, *P. echinata*, *P. halepensis*, *P. mughus*, *P. montana*, *P. pinaster*, and *P. sylvestris*.
- Cylindrosporium cleonorae-reginae** Bub. and Malk. On leaves of *S. transsilvanicus* in Bulgaria.
- Peronospora senecionis** Fekl. Downy mildew on leaves of *S. cordatus* in Germany.
- Phyllachora nidulans** Pat. Dull-black stromata on leaves of *Senecio* sp. in Ecuador.
- Phyllosticta albo-brunnea** Bub. and Wrobl. On leaves of *S. umbrosus* in Russia and Poland.
- Phyllosticta huallatae** Speg. Subcircular gray-white leaf spots on *S. huallata* in Argentina.
- Phyllosticta jacobaeae** Sacc. Indefinite ochraceous leaf spots on *S. jacobaea* in Italy.
- Phyllosticta senecionis-cordati** Allesch. Dull-yellow leaf spots on *S. cordatus* and *S. fuchsii* in Bohemia, Silesia, and Germany.
- Puccinia glomerata** Grev. Brown rust pustules on brown leaf spots on *S. jacobaea* in Great Britain and Belgium.
- Puccinia ligericae** Syd. Leaf rust on *S. viscosus* and *Carex ligerica* in Germany.
- Puccinia oedipus** Cke. Yellow-brown to black rust pustules on leaves of *S. pandurifolius* in the Union of South Africa.
- Puccinia schoeleriana** Plowr. and Magn. Leaf rust on *S. jacobaea* and *Carex* spp. in Europe.
- Puccinia senecionicola** Arth. Brown rust sori on leaves of *Cacalia amplifolia*, *C. ampullacea*, *C. calotricha*, *C. pringlei*, *C. obtusiloba*, *C. sinuata*, *Senecio angulifolius*, *S. petasitoides*, *S. sinuatus*, and *S. warszewiczii* in Mexico and Guatemala.
- Puccinia senecionis** Lib. Brown to black rust pustules on leaves of *Senecio* spp. in central Europe and Australia.
- Puccinia senecionis-acutiformis** Hasl. and Mayor. Leaf rust on *Senecio* spp. and *Carex* spp. in Switzerland.
- Puccinia senecionis-ochrocarpi** Bacc. Leaf rust on *S. ochrocarpus* in Abyssinia.
- Puccinia silvatica** Schroet. See Taraxacum.
- Puccinia tasmanica** Diet. Black rust pustules on leaves and stems of *S. vulgaris* in Tasmania.
- Puccinia uralensis** Tranzs. Leaf rust on *S. fuchsii*, *S. nemorensis* and *S. scandens* in Ceylon, Yugoslavia, Russia, and Hungary.
- Ramularia chlorina** Bres. On leaves of *S. fuchsii* in Germany.
- Ramularia coeosporii** Sacc. See Campanula.
- Ramularia pruinosa** Speg. Small circular ochraceous leaf spots which finally involve the entire leaf area of *S. jacobaea* in Great Britain and Italy.
- Ramularia senecionis** (B. and Br.) Sacc. On leaves of *S. carniolicus*, *S. jacobaea*, *S. nebrodensis*, *S. (Cineraria) palustris*, and *S. subalpinus* in Great Britain, the Balkans, and Austria. Reported from Colorado.
- Septoria anaxaea** Sacc. Ochraceous leaf spots on *S. praealtus* in Italy.
- Septoria martiniae** Cke. Gray confluent leaf spots on *S. bedfordii* in Victoria.
- Septoria selloi** Speg. Small circular gray-white leaf spots on *S. selloi* in Argentina.
- Septoria senecionis** West. Circular to irregular gray-black leaf spots with white centers on *Senecio* spp. in Siberia and Europe. Reported from California.
- Uromyces ameghinoi** Speg. Leaf rust on *S. laevicaulis* in Patagonia.
- Uromyces araucanus** Diet. and Neg. Brown leaf rust on circular spots on *S. otites* in Chile.
- Uromyces kurtzii** P. Henn. Rust deforming leaves and stems of *S. salsus* in Argentina.
- Uromyces psamathonophilus** Speg. A rust deforming leaves, branches, and inflorescences of *Senecio* sp. in Argentina.
- Uromyces senecionicola** Arth. Brown leaf rust on *S. roldana* in Mexico and Guatemala.
- SEBJANIA**. Climbing or twining tropical shrubs.
- Aecidium serjaniae** P. Henn. Ochraceous rust pustules on yellow leaf spots on *S. fulva* and *Serjania* sp. in Brazil and Argentina.
- Dothidella serjaniae** P. Henn. Brown stromata on stems of *Serjania* sp. in Brazil.
- Phyllachora duplex** Rehm. Shiny black stromata on brown spots on leaf blades and petioles of *S. caracasana* in Brazil and Argentina.
- Phyllachora intermedia** Speg. Small brown galls on leaves and stems of *S. glabrata* in Brazil. The fungus is said to be more properly a *Valsaria*.
- Phyllachora subrepens** Speg. Black stromata on leaves of *Serjania* sp. in Paraguay and Brazil.
- Physalospora serjaniae** Rehm. Circular red to dull-brown leaf spots on *Serjania* sp. in Brazil.
- SERBATULA**. Thistlelike perennial herbs with purple or violet flower heads.
- Ovularia serratulae** Sacc. Leaf spot on *S. tinctoria* in Spain.
- Puccinia heterophyllae** Cke. Powdery brown rust pustules on leaves and stem of *S. cerinthefolia* and *S. heterophylla* in Kurdistan and Asia Minor.
- Puccinia miurae** Syd. Leaf rust on *S. atriplicifolia* in Japan.
- Puccinia schirajewskii** Tranzsch. Leaf rust on *S. pallida* in India.
- Puccinia schroeteriana** Kleb. Leaf rust on *S. radiata*, *S. tinctoria*, and *Carex* spp. in Russia and central Europe.
- Puccinia serratulae** Thuem. Brown leaf rust on *S. tinctoria* and *Serratula* sp. in Siberia and Russia.
- Puccinia serratulae-oligocephalae** Syd. Powdery brown to black rust pustules on leaves of *S. oligocephala* in Syria.
- Puccinia serratulae-pinnatifidae** Gz. Frag. Leaf rust on *S. pinnatifida* in Spain.
- Puccinia tinctoriicola** P. Magn. Leaf rust on *S. centauroides*, *S. coronata*, *S. heterophylla*, *S. sloanei*, and *S. tinctoria* in Siberia and Europe.
- Septoria cirsii** Niessl. Subcircular brown leaf spots on *S. glauca* and *Cirsium arvense* in Italy.
- Septoria tinctoriae** Brun. On leaves of *S. tinctoria* in Germany.
- SESAMUM**. SESAME. Tropical herbs producing oil-bearing seeds.
- Ascochyta** sp. On *S. orientale* in south Russia.
- Bacillus sesami** Malkoff. Attacks *S. orientale* (*S. indicum*) in connection with *Pseudomonas sesami* in Bulgaria.
- Cercospora sesami** A. Zimm. Numerous small light-brown spots with dark-brown margins on leaf blades, petioles, stems, and capsules of *S. orientale* (*S. indicum*) in Porto Rico, Cuba, Brazil, Tanganyika, the Philippines, Ceylon, China, and Japan. Reported from Florida.
- Fusarium** sp. Causes a serious wilt of *S. orientale* in India. The disease is seed-borne.

**SESAMUM**—Continued.

**Helminthosporium sesami** Miy. On leaves of *S. orientale* in China.

**Hypochnus cucumeris** Frank. See Cucumis.

**Phytophthora parasitica** Dast. See Ricinus.

**Pseudomonas sesami** Malkoff. Produces brown to black spots on leaves and stems of *S. orientale* in Bulgaria. Diseased plants turn black and rot.

**Rhizoctonia** sp. See Vigna.

**SESBANIA**. Leguminous herbs or shrubs.

**Cercospora agatidis** Foex. On leaves of *S. grandiflora* in Indo-China.

**Cercospora sesbaniae** P. Henn. Effuse yellowish leaf spots on *S. grandiflora* in India and the Congo

**Uredo fimbriata** Speg. Brown leaf rust on *Sesbania* sp. in Brazil.

**Uredo sesbaniae** P. Henn. Rust pustules on yellow leaf spots on *S. aegyptiaca* and *Sesbania* sp. in India and the Congo.

**SESELI**. Perennial and biennial herbs.

**Puccinia phymatospora** Lindr. Brown to black rust pustules on leaves of *S. coloratum* in Italy, France, Belgium, and Switzerland.

**Septoria desciscens** Sacc. On leaves of *S. strictum* in Siberia.

**Septoria seseli** Höll. On leaves of *S. glaucum* in Hungary.

**Uromyces graminis** (Niessl.) Diet. Powdery brown to black rust pustules on leaves of *S. glaucum*, *Laserpitium siler*, and *Melica ciliata* in Europe.

**Uromyces seseli-graminis** Ed. Fisch. See Arrhenatherum.

**SETARIA**. **CHAETOCHELOA** Ag. **MILLET**. Annual and perennial grasses, mostly weeds. See, also Chaetochloa.

**Acrothecium lunatum** Wak. See Panicum.

**Balansia claviceps** Speg. See Pennisetum.

**Phyllachora evansii** Syd. Dull-black stromata on yellowish leaf spots on *S. aurea*, *S. nigrirostris*, and *S. sulcata* in the Union of South Africa.

**Phyllachora setariaecola** Speg. Black stromata on leaves of *Setaria* sp. in Brazil and Ecuador.

**Phyllosticta crastophila** Sacc. See Phalaris.

**Phyllosticta glumarum-setariae** P. Henn. Pale-brown spots on glumes of *S. aurea* in the Congo.

**Phyllosticta setariae** Ferr. Irregular gray-white leaf spots with purple margins on *S. glauca* in Italy.

**Puccinia atra** Diet. and Holw. Black rust sori on leaves of *S. grisebachii* in Mexico.

**Puccinia camelia** (Mayor) Arth. Leaf rust on *S. scandens* and *Chaetochloa setosa* in Porto Rico and Colombia.

**Puccinia setariae** Diet. and Holw. Brown rust pustules on leaves of *S. caudata* and *S. imberbis* in Chile and Mexico.

**Scirrhia setariae** (Sacc.) Theiss. and Syd. On *S. glauca* in Italy.

**Sorosporium setariae** McAlp. Ovaries converted into black powdery masses of smut spores. On *S. glauca* in Queensland.

**Sphacelia grisea** Speg. On culms of *Setaria* sp. in Argentina.

**Sphaelotheca pamparum** (Speg.) Clint. Smut sori in the ovaries as oblong bodies protected by false membranes which break irregularly, disclosing dusty olive-black spore masses. On *S. geniculata* and *S. imberbis* in Mexico, Cuba, Bahamas, and Chile.

**Sporotrichum peribueyense** Speg. On leaves of *Setaria* sp. in Brazil.

**Tolyposporium pampeanum** Speg. Dark powdery smut sori in spikelets of *Setaria* sp. in Argentina.

**Tolyposporium setariacolum** Syd. Smut on *S. aurea* in Central Africa.

**Uromyces setariae-italicae** (Diet.) Yosh. Cinnamon-brown to black rust pustules on leaves of *S. glauca*, *S. intermedia*, *S. italica*, *S. verticillata*, and *S. viridis* in India, Japan, Ceylon, China, Manchuria, the Philippines, and north Africa.

**Ustilago oidea setariae** Bref. Powdery smut sori in ovaries of *S. crus-ardea* in Brazil.

**Ustilago evansii** P. Henn. Smut on *S. aurea* in Africa.

**Ustilago heterospora** P. Henn. See Panicum.

**Ustilago setaria** Rbh. A smut destroying the panicles of *S. glauca*, *S. viridis*, and *Syntherisma sanguinale* in Argentina and Germany.

**Ustilago setariae-aureae** P. Henn. Black powdery smut sori destroying the ovaries of *S. aurea* in the Congo and the Union of South Africa.

**Ustilago verruculosa** Wakef. Smut on *S. aurea* in tropical Africa.

**SHAWIA**. See Olearia.**SHOREA**. Tropical trees.

**Corticium salmonicolor** B. and Br. See Citrus.

**Fomes lignosus** Klotzsch. See Hevea.

**Polyperus shoreae** Wakef. Rot of trunk and branches of *S. robusta* in India.

**SIDA**. Malvaceous herbs or shrubs, mostly weeds, one species cultivated for fiber.

**Cercospora densissima** Speg. Large indefinite pale-brown leaf spots on *S. rhombifolia* in Porto Rico and Argentina.

**Cercospora sidae** P. Henn. Pale-brown leaf spots on *S. cordifolia* in the Congo.

**Dietelia verruciformis** P. Henn. Yellow-brown rust pustules on leaves of *S. macrodon* in Argentina.

**Phyllosticta sidaecola** Cke. Irregular, then confluent, brown leaf spots on *S. napaea* in Great Britain.

**Septoria heterochroa** Desm. See Malva.

**SIDEROXYLON**. Tropical trees and shrubs, some of value for timber.

**Phyllachora baumii** P. Henn. Black stromata on gray to brown leaf spots on *Sideroxylon* sp. in the Union of South Africa.

**Puccinia johnstonii** Arth. Brown leaf rust on *Dipholis salicifolia* and *S. foetidissimum* in Cuba.

**SILENE**. **CAMPION**. **CATCHFLY**. Erect annual, biennial, or perennial herbs.

**Marsonia delastrei** (Delacr.) Sacc. See Lychnis.

**Mycosphaerella silenis** v. Hoeh. Numerous small ochraceous spots on leaves and calyces of *S. latifolia* in Austria.

**Peronospora vexans** Gäum. Downy mildew on leaves of *Silene* spp. in Europe.

**Phacidium pumilum** Desm. On stems and leaves of *S. conica* and *S. rubella* in France and Tunis.

**Phyllosticta nebulosa** Sacc. Pale-brown leaf spots on *S. armeria*, *S. pendula*, and *S. splendens* in north Africa, Denmark, and Italy.

**Phyllosticta otites** Brun. Olivaceous leaf spots on *S. otites* in France.

**Phyllosticta zahlbrukneri** Bauml. Subcircular gray-white leaf spots on *S. latifolia* and *S. nutans* in Russia, Hungary, and Austria.

**Puccinia silenens** Schroet. Brown rust pustules on leaves of *Silene* spp. and *Melandryum* sp. in Siberia and Europe.

**Ramularia didymarioides** Br. and Sacc. Indefinite ochraceous leaf spots on *S. latifolia* in Spain and France.

**SILENE**—Continued.

**Ramularia lychnicola** Cke. See Lychnis.

**Ramularia silenes** Karst. Ochraceous leaf spots on *Silene* sp. in Finland.

**Septoria apetalae** P. Magn. On leaves of *S. apetalae* in Palestine.

**Septoria dimera** Sacc. Indefinite ochraceous leaf spots on *S. dichotoma*, *S. latifolia*, *S. livida*, and *S. nutans* in Europe.

**Septoria doehlii** Syd. Circular gray-white leaf spots on *S. nutans* in Germany.

**Septoria dominii** Bub. and f. **calycicola** Gz. Frag. Small circular dull-yellow leaf spots on *S. latifolia* in Yugoslavia and Spain.

**Septoria silenes-nutantis** C. Mass. Subcircular yellow-white leaf spots on *S. nutans* in Italy.

**Septoria saponariae** (DC.) Sav. and Becc. See Saponaria.

**Sorosporium saponariae** Rud. See Cerastium.

**Uromyces behensis** (DC.) Ung. Dark-brown to black rust pustules on leaves of *Silene* spp. in Europe.

**Uromyces inaequialtus** Lasch. Yellow to dark-brown rust pustules on leaves of *Silene* spp. in Europe, Siberia, Asia Minor, and Japan.

**Ustilago major** Schroet. Dark-violet smut sori in anthers of *S. otites* in Europe.

**SILYBUM. MARIANA** Ag. Erect thistlelike plants.

**Aecidiolum marianum** Gz. Frag. Leaf rust on *S. marianum* in Spain.

**Hypochnus fuciformis** (Berk.) McAlp. See Lolium.

**Ramularia cynarae** Sacc. See Cynara.

**Septoria silybi** Pass. Subcircular dull-brown to gray leaf spots on *S. marianum* in Italy.

**Ustilago cardui** F. v. Wald. Powdery brown-violet masses of smut spores replacing ovaries of *S. marianum* and *Carduus acanthoides* in Germany.

**SIPHOCAMPYLUS.** Herbs or shrubs, sometimes climbing.

**Uredo siphocampyli** P. Henn. Powdery brown rust pustules on leaves of *Siphocampylus* sp. in Argentina.

**Uromyces cystopiformis** Lagerh. Leaf rust on *Siphocampylus* sp. in Ecuador.

**SIRMUELLERA.** See Banksia.**SISYRINCHIUM.** BLUE-EYED GRASS. Iridaceous fibrous-rooted perennials.

**Aecidium bunsteri** Neg. Rust on leaves of *S. andinum* in Chile.

**Phyllosticta chilensis** Allesch. Leaf spots on *S. pedunculatum* in Chile.

**Puccinia angulata** Diet. & Neg. Yellow rust pustules, becoming dark-brown to black, on both leaf surfaces and on scapes of *S. pedunculatum* in Chile.

**Puccinia sisyrinchii** Mont. (*Uromyces sisyrinchii* Mont.) Produces brown to black rust pustules on both leaf surfaces and on scapes of *S. graminifolium* and *S. palmifolium* in Chile and Argentina. Reported from Washington.

**Puccinia straminea** Diet. Brown to dark-brown powdery rust pustules on both surfaces of leaves of *Sisyrinchium* sp. in Brazil.

**Septoria sisyrinchii** Speg. Leaf spot on *S. bonariense* in Argentina.

**Uredo nominata** Arth. Leaf rust on *S. bermudianum* in Bermuda.

**SMILACINA. VAGNERA** Ag. FALSE SOLOMON'S SEAL. Perennial herbs.

**Protomyces purpureo-tingens** Mass. Elongated or broadly effused red or purple patches on cotyledons and young leaves of *Smilacina* sp. in Great Britain.

**SMILAX. GREENBRIER. CATBRIER.** Herbaceous or shrubby climbers.

**Blastospora smilacis** Diet. Yellow rust pustules on small sunken leaf spots on *S. herbacea* and *S. sieboldii* in Japan.

**Cercospora miyakei** P. Henn. Circular to angular, then confluent, leaf spots on *S. herbacea* in Japan.

**Cercospora smilacina** Speg. Small brown leaf spots on *Smilax* sp. in Brazil. Saccardo has given the same name to a species on leaves of *S. aspera* and *S. mauritanica* in Italy, Spain, Portugal, and the United States.

**Dothidella smilacicola** Rehm. Black stromata on leaves of *Smilax* sp. in Brazil.

**Gloeosporium thuemeri** Sacc. See Anthurium.

**Marsonia smilacina** Thuem. Irregular ochraceous leaf spots on *S. mauritanica* in Spain and Portugal.

**Phaeochorella clypeata** (Wint.) Theiss. and Syd. Shiny black stromata on leaves of *Smilax* sp. in Brazil.

**Phyllosticta smilacina** Speg. Rufous leaf spots on *S. campestris* in Argentina.

**Puccinia citrina** Syd. Leaf rust on *S. gaudichaudiana*, *S. stenopetala* and *Smilax* sp. in China, Japan, and the Philippines.

**Puccinia ferruginea** Lév. Dull brown rust pustules on leaves of *S. aspera* and *S. zeylanica* in Macao and Ceylon.

**Puccinia henryana** Syd. Leaf rust on *S. china*, *S. herbacea*, and *S. menispermoides* in Japan and China.

**Puccinia kraussiana** Cke. Brown rust pustules on leaves of *S. kraussiana* in central and south Africa.

**Puccinia merrillii** P. Henn. Leaf rust on *S. bracteata*, *S. latifolia*, and *S. vicaria* in the Philippines.

**Puccinia prainiana** Barcl. Brown rust pustules on leaves of *S. aspera*, *S. elegans*, and *S. maculata* in India.

**Puccinia smilacis-chinae** P. Henn. Leaf rust on *S. chinense* and *S. herbacea* in Japan and the Philippines.

**Sphenospora pallida** (Wint.) Diet. Waxy rust pustules on leaves of *Smilax* sp. in Guatemala, Costa Rica, and South America.

**Uredo yurimaguasensis** P. Henn. Leaf rust on *Smilax* sp. in Peru.

**SOBRALIA.** See Orchidaceae.**SOJA. SOYBEAN.** Herbaceous leguminous plants, with a wide range of economic uses. (Including *Glycine*.)

**Aecidium glycines** P. Henn. Leaf rust on *S. max* in Tanganyika and Uganda.

**Asochyta** sp. On *S. max* in Japan.

**Bacterium** sp. A leaf-spotting disease of *S. max*, said to be due to a bacterium distinct from other species on this host, is reported from Japan.

**Cercospora daizii** Miura. Leaf spot on *S. max* in Manchuria.

**Cercospora glycines** Cke. Definite dark-brown leaf spots on *Glycine clandestina* in Australia.

**Colletotrichum glycines** Hori. Anthracnose on stems and pods of *S. max* in Japan and Chosen.

**Fusarium** sp. Causes ring spot disease of *S. max* in Manchuria.

**Hypochnus centrifugus** Tul. Causes cankers on stems, blighting infected plants of *S. max* in Manchuria.

**Hypochnus cucumeris** Frank. See Cucumis.

**Mosaic.** Mosaic disease of *S. max* reported from Japan is probably the same as the mosaic and crinkling reported from the United States.

## SOJA—Continued.

**Mycosphaerella phaseolarum** Siem. See Phaseolus.

**Peronospora manshurica** (Naoum.) Syd. (*P. trifoliorum* De B. var. *manshurica* Naoum.) Downy mildew on leaves of *S. max* in Siberia, India, Formosa, and Manchuria, causing premature leaf fall.

**Phyllosticta sojaecola** Massal. (*Phacosphaerulina sojaecola* Miura.) Subcircular, dull-brown leaf spots on *S. max* in Japan, Russia, Manchuria, and Italy.

**Pseudomonas glycines** Nak. Circular yellow leaf spots, becoming brown to dark-brown with yellow margins, on *S. max* in Japan. The spots may be as numerous as 70 to 80 per square centimeter, causing death of infected leaves.

**Septoria glycines** T. Hem. Brown leaf spots on *S. max* in Japan and Manchuria.

**Septoria sojae** v. Thuem. Irregular yellowish leaf spots on *S. max* in Italy, Japan, and Austria.

**Trotteria venturioides** Sacc. Black mildew on leaves of *S. max* in the Philippines.

**Uredo vignae** Bres. See Vigna.

**Uromyces sojae** (P. Henn.) Syd. Brown rust pustules on leaves of *S. max* in Japan, Java, China, India, Formosa, Manchuria, and the Philippines.

**SOLANUM. NIGHTSHADE. POTATO. EGGPLANT.** Temperate and tropical herbs and shrubs, many species of economic, ornamental, or medicinal value.

**Actinomyces** spp. A number of species of *Actinomyces* have been described from Germany, causing various types of scab on potato (*S. tuberosum*) tubers. Whether these forms differ from *A. scabies* the common American species, is problematical as yet. These forms include *A. albus* (R. D.) Gasp. (girdle scab); *A. incanescens* Wr. (deep scab); *A. intermedius* (Krug.) Wr. (shallow scab); *A. nigricans* (Krug.) Wr. (girdle scab); *A. aeruginosus* Wr. (knobby scab), and *A. tricolor* Wr. (shallow scab).

**Aecidium habunguense** P. Henn. Leaf rust on *S. melongena* in India.

**Aecidium solani** Beeli. Leaf rust on *Solanum* sp. in Belgian Congo.

**Aecidium solani-unguiculati** P. Henn. Leaf rust on *S. unguiculatum* in Abyssinia.

**Ascochyta dulcamara** Bub. On leaves of *S. dulcamara* in Bohemia and Germany.

**Ascochyta solani-nigri** Died. Circular to ovate white leaf spots with dark margins on *S. nigrum* in Germany.

**Bacillus** spp. Bacteria, the occurrence of which is unknown in the United States, cause the rotting of potato (*S. tuberosum*) tubers in India and Malaya.

**Bacillus caulivorus** Prill. and Delacr. Said to cause cankers on the stems of *S. tuberosum*, *Lycopersicon esculentum*, and *Nicotiana tabacum* in Italy and France.

**Bacillus tubifex** Dale. Irregular brown spots, generally along veins or margins of leaves of *S. tuberosum* in Great Britain.

**Bacterium** sp. An undescribed species of *Bacterium* is reported from Morocco as causing a serious potato (*S. tuberosum*) disease. The disease begins at the apex of the plant as brown patches on the leaflets, which rapidly turn black and wilt. The disease works down the petioles into the stems and finally to the tubers. The tops of the plants are covered with elongate black patches and finally die. Small violet-black spots appear on the tubers, enlarging until the entire tuber surface is involved, a soft rot ensuing.

**Cercospora aratai** Speg. Circular ashen-brown leaf spots on *S. glaucum* in Argentina.

**Cercospora dulcamaricola** Hóll. Circular brown leaf spots on *S. dulcamara* in Hungary.

**Cercospora heterosperma** Bres. On leaves of *S. tuberosum* in Poland.

**Cercospora incarnata** P. Henn. On leaves of *Solanum* sp. in Brazil.

**Cercospora melongena** Welles. Irregular, then confluent, zoned, chlorotic areas on leaves of *S. melongena* in the Philippines.

**Cercospora solanacea** Sacc. and Berl. Subcircular brown leaf spots on *S. verbascifolium* in Australia.

**Cercospora tosenensis** P. Henn. Circular, then confluent, dull-brown spots on leaves of *S. biflorum*, *S. nigrum*, and *Solanum* sp. in the Philippines and Japan.

**Chrysophlyctis endobiotica** Schilb. (*Synchytrium endobioticum* [Schilb.] Perc.) The potato (*S. tuberosum*) wart disease or potato canker is one of the most serious diseases of this important crop. It occurs in Great Britain, Holland, Norway, Poland, Luxemburg, the Union of South Africa, and Germany. In the United States the disease also occurs in limited areas in Maryland, West Virginia, and Pennsylvania. The disease is characterized by wartlike outgrowths on the tubers, which vary in size and distribution. They commonly develop from the eyes of tubers, more rarely on stolons or base of stems. The warts are at first whitish, turning black with age. Exposed warts take on a green color. In size they vary from small nodules to masses larger than the tubers on which they are borne. The organism causing the disease can persist in the soil for many years and is readily carried from place to place in this medium. The tomato (*Lycopersicon esculentum*) has been found capable of harboring the disease. By inoculation *S. dulcamara* and *S. nigrum* have been found susceptible.

**Didymopora solani** Diet. Leaf rust on *Solanum* sp. in Brazil.

**Didymopora solani-argentei** (P. Henn.) Diet. Rust pustules on circular dull-brown leaf spots on *S. argenteum* in Brazil.

**Guignardiella nervisequia** (Rehm.) Sacc. and Syd. On leaves of *S. auriculatum* in Brazil.

**Helminthosporium solani** McAlp. On leaf blades and petioles and stems of *S. viride* in New South Wales.

**Helminthosporium solanium** Sacc. and Syd. Leaf spots on *S. argenteum* in Brazil.

**Leptosphaeria circinans** (Fekl.) Sacc. See Asparagus.

**Moniliopsis aderholdi** Ruhl. The "vermerhrungs Pilze" attacks the tubers of *S. tuberosum* and also causes a damping-off of seedlings of many species of cultivated plants, including *Beta vulgaris*, *Brassica* spp., *Cinchona* sp., and *Raphanus* in Europe and Java. The fungus forms black sclerotia similar to those of *Rhizoctonia solani*, with which it is sometimes considered identical.

**Oospora pustulans** Owen and Wakef. The skin spot disease of potatoes (*S. tuberosum*) occurs in Great Britain and in limited areas in Washington and British Columbia. Numerous small circular, slightly raised areas occur on the tubers. The possible connection of this disease with immature stages of powdery scab (*Spongospora*) has not yet been definitely settled.

**Phellomyces sclerotiphorus** Frank. Reported as the cause of a scab and dry rot of potato (*S. tuberosum*) tubers in Russia and Great Britain.

**Phoma solanicola** Prill. and Delacr. See *Nicotiana*.

**Phyllachora fluminensis** Theiss. Shiny black stromata on brown leaf spots on *Solanum* sp. in Brazil.

**Phyllohendersonia dulcamarae** (Sacc.) Tass. On leaves of *S. dulcamara* in Italy.

**Phyllosticta concentrica** Th. On leaves of *Solanum* sp. in Brazil.

**Phyllosticta dulcamarae** Sacc. Brown, often marginal, leaf spots on *S. dulcamara* in France, Italy, and Denmark. Reported from Kansas.

**Phytophthora arecae** (Colem.) Pethybr. See *Palmae*.

**SOLANUM**—Continued.

**Phytophthora erythroseptica** Pethybr. This downy mildew fungus causes a wilting of potato (*Solanum tuberosum*) plants and a moist rot of the tubers in Ireland, Switzerland, Holland, and Dutch East Indies. Cut surfaces of infected tubers turn pink rapidly upon exposure to air, becoming purplish-brown to black after several hours. Diseased tubers remain firm, but of rubbery consistency, and if pressed exude a quantity of juice, becoming finally completely rotted. *Atropa belladonna* is reported as an additional host in Poland.

**Phytophthora melongenae** K. Saw. Downy mildew attacking primarily fruit of *S. melongena* in Japan but also capable of infecting *S. tuberosum*, *Areca catechu*, *Hibiscus esculentus*, *Lycopersicum esculentum*, *Nicotiana* sp., and *Ricinus* sp.

**Phytophthora parasitica** Dastur. See *Ricinus*.

**Pseudomonas solaniolens** Paine. This bacterium has been described as the cause of internal brown spot of potato (*S. tuberosum*) tubers in Europe.

**Puccinia araucana** Diet. and Neg. Yellow and brown rust pustules on deformed leaves and branches of *S. crytopodium* and *S. valdivianum* in Chile.

**Puccinia hieronymi** P. Henn. Leaf rust on *S. incisum* and *S. nodiflorum* in Argentina.

**Puccinia huallagensis** P. Henn. Cinnamon-brown rust sori on dark-brown leaf spots on *S. torvum* in Peru and Colombia.

**Puccinia imitans** Syd. Leaf rust on *Solanum* sp. in Ecuador.

**Puccinia negeriana** Diet. Dark-brown rust pustules on leaf blades and petioles of *S. furcatum* in Chile.

**Puccinia pitteriana** P. Henn. Numerous brown rust pustules on lower leaf surfaces of *S. tuberosum* and *Lycopersicum esculentum* in Costa Rica and Colombia. This rust appears to be capable of causing heavy damage to the host plants through drying and premature fall of leaves.

**Puccinia solanacearum** Sacc. and Syd. Black rust pustules on leaves of *Solanum* sp. in India.

**Puccinia solanita** (Schw.) Arth. Brown leaf rust on *Solanum* sp. and *S. donnellsmithii* in Trinidad, Panama, and Surinam.

**Puccinia solani-tristis** P. Henn. Powdery black rust pustules on leaves of *S. triste* in Brazil.

**Puccinia tijbodensis** Gäum. Brown leaf rust on *S. biflorum* in Java.

**Puccinia tubulosum** (Pat. and Gaill.) Arth. See *Paspalum*.

**Puccinia weberbaueri** P. Henn. Dark-brown rust pustules on yellow-brown leaf spots on *Solanum* sp. in Peru.

**Puccinosira solani** Lagerh. Yellow leaf rust on *Solanum* sp. in Ecuador.

**Rhizoctonia destruens** Tassi. This fungus attacks its hosts at the ground level, setting up a rot which destroys the plants, the numerous brown sclerotia appearing on the rotted areas. Potato tubers are soft rotted, thick white strands of mycelium growing over the surface. The fungus is reported from Italy, India, and Java, on *S. melongena*, *S. tuberosum*, *Amorphophallus campanulatus*, *Arachis hypogaea*, *Delphinium decorum*, *D. grandiflorum*, *D. magnificum*, *D. machayanum*, *D. montanum*, *Dianthus* sp., *Hibiscus rosa-sinensis*, *Lobelia laxiflora*, *Medicago sativa*, *Oryza sativa*, *Piper betel*, and other hosts. The fungus is not distinct from *Sclerotium rolfsii*, but may represent a strain not present in the United States.

**Sarcinella raimundoi** Sacc. On leaves of *S. melongena* in the Philippines.

**Septoria dulcamarae** Desm. Small subcircular brown, then gray, leaf spots on *S. dulcamara* in Europe.

**Septoria pseudo-quinae** Pat. Dull-brown to gray leaf spots with black borders on *S. pseudo-quina* in Ecuador.

**Septoria solanina** Speg. Small circular dull-brown leaf spots on *S. gracile* in Argentina.

**Septoria solani-nigri** Scalia. Circular, then confluent, dull-brown leaf spots on *S. nigrum* in Italy.

**Septoria solanophila** Speg. Circular gray-white leaf spots with purple margins on *S. verbascifolium* in Argentina.

**Spongospora subterranea** (Walls.) Lang. Powdery scab is a disease introduced within recent years into the United States from Europe, and now established in the northern potato (*S. tuberosum*) growing regions. It does not seem to be able to persist south of the New England and Lake States. It occurs in northern Europe, Algeria, northern South America, and New Zealand. Other hosts are *S. ciliatum*, *S. commersoni*, *S. haematocladum*, *S. mammosum*, *S. marginatum*, *S. warscewiczii*, and *Lycopersicum esculentum*.

The disease closely resembles common scab, but is distinguished by the smaller, more definite lesions, which on maturity break open, revealing brown, powdery masses, surrounded by the torn skin of the tubers.

**Sporidesmium exitiosum** Kuehn. var. *solani* Fckl. On stems of *S. tuberosum* in Russia. See also *Brassica*.

**Sporidesmium melongenae** Thuem. On leaves of *S. melongena* in Portugal and Brazil.

**Tylenchus dipsaci** Kuehn. A biological strain of this nematode attacks potato (*S. tuberosum*) tubers in Great Britain. Brown patches appear on the skin followed by internal rot. See also *Narcissus*.

**Uredo minitans** Speg. Brown rust pustules on circular leaf spots on *S. commersoni* in Argentina.

**Urocystis hieronymi** Schroet. Ochraceous smut sori on branches and peduncles of *Solanum* sp. in Bolivia.

**Uromyces solani** Diet. and Holw. Yellow-brown to dark-brown rust pustules on leaves of *S. appendiculatum* and *S. nudum* in Mexico.

**Vermicularia capsici** Syd. See *Capsicum*.

**Vermicularia varians** Duc. The black dot disease or "dartrose" is considered a serious disease of *S. tuberosum* in France and some other European countries. It is also reported from the Union of South Africa, Australia, Brazil, and Canada. The fungus undoubtedly occurs in the United States, but has not been associated with a definite disease of the potato. In Europe the disease attacks the stems, roots, and tubers. The fungus also occurs on *Lycopersicum esculentum* and *Physalis peruviana*. It is more properly referable to *Colletotrichum*.

**SOLARIA**. Bulbous plants.

**Uromyces solariae** Diet. Chestnut colored powdery rust pustules on both leaf surfaces of *S. miersioides* in Chile.

**SOLDANELLA**. Small, glabrous, perennial herbs with short rhizomes.

**Puccinia soldanellae** (DC.) Fckl. Yellow and dark-brown rust pustules on leaves of *S. alpina*, *S. minima*, *S. montana*, and *S. pusilla* in Europe.

**Septoria soldanellae** Speg. and var. *pirolaeifoliae* Vogl. Brown spots, often occupying the entire leaf area, on *S. alpina*, *S. montana*, and *S. pyrolaeifolia* in Italy.

**Septoria versicolor** Bub. Circular to irregular yellow-white leaf spots on *S. montana* in Bohemia.

**Thecaphora hyalina** Fingerh. See *Convolvulus*.

**SOLIDAGO.** GOLDENROD. Perennial herbs.

**Cercospora fulvescens** Sacc. Small yellow-brown leaf spots on *S. virgaurea* in Denmark, Italy, and Japan. Reported from Colorado and Kansas.

**Mycosphaerella virgaureae** Krieg. On leaves of *S. virgaurea* in Germany.

**Ovularia virgaureae** (Thuem.) Sacc. On leaves of *S. virgaurea* in Denmark, Switzerland, and Germany.

**Phyllosticta solidaginis** Bres. On leaves of *S. virgaurea* in Germany.

**Puccinia solidaginicola** Diet. Leaf rust on *Solidago* sp. in Chile.

**Puccinia solidaginis-microglossae** Diet. Chestnut-brown rust pustules on yellow or brown leaf spots on *S. microglossa* in Argentina.

**Puccinia virgaureae** (DC.) Lib. Black rust pustules on leaves of *S. microglossa* and *S. virgaurea* in Japan and Europe.

**Septoria virgaureae** Desm. On leaves of *S. virgaurea* in Yugoslavia, Italy, and Switzerland.

**Uromyces komarovii** Bub. Brown rust pustules on leaves of *S. virgaurea* in Japan and Manchuria.

**SONCHUS.** SOW THISTLE. Composites, mostly weeds, some species used for foliage effects.

**Bremia sonchi** K. Saw. Downy mildew causing small polygonal spots on stems and leaves of *S. oleraceus* in Formosa.

**Coleosporium sonchi** Lév. Leaf rust on *S. asper*, *S. oleraceus*, *S. palustris*, and *S. uliginosus* in Europe. The aecial stage occurs on *Pinus sylvestris*. Introduced into Wisconsin and found locally on *S. asper* and *Pinus sylvestris*.

**Phyllosticta sonchi** Sacc. Gray-white leaf spots on *S. oleraceus* in Russia and Italy.

**Puccinia sonchi** Rob. Yellow to black rust pustules on leaves of *Sonchus* spp. in Ceylon, Japan, Algeria, Canary Islands, and Europe.

**Puccinia sonchina** Syd. Brown rust pustules on leaves of *S. oleraceus* in Japan and Portugal.

**Ramularia sonchi-arvensis** Fautr. On leaves of *S. oleraceus* in Spain and France.

**Septoria modonia** Sacc. Small subcircular spots on leaves and stems of *S. arvensis* in France.

**Septoria sonchi** Sacc. Oblong greenish-gray leaf spots on *S. oleraceus* in Italy.

**Septoria sonchicola** Höll. Circular ochraceous leaf spots with black borders on *S. uliginosus* in Hungary.

**Septoria sonchina** Thuem. Leaf spots on *S. oleraceus* in Siberia.

**Synchytrium globosum** Schroet. See *Potentilla*.

**SOPHRONITIS.** See *Orchidaceae*.**SOPHORA.** Shrubs or herbs grown for their attractive flowers and handsome foliage.

**Aecidium sophorae** Kus. Leaf rust on *S. platycarpa* in Japan.

**Ascochyta sophorae** Allesch. On branches of *S. japonica* in Germany.

**Macrosporium sophorae** Turconi. Circular brown leaf spots on *S. japonica* in Italy.

**Phyllosticta sophoricola** Höll. Irregular ashen leaf spots with dull-brown borders on *S. japonica* in Hungary.

**Uromyces shikokianus** Kus. Leaf rust on *S. shikokiana* in Japan.

**Uromyces sophorae-flavescentis** Kus. Brown leaf rust on *S. flavescens* in Japan.

**Uromyces sophorae-japonicae** Diet. Brown rust pustules on leaf blades and petioles of *S. japonica* in Japan.

**Uromyces trunicola** P. Henn. and Shir. Dark-brown rust pustules on branches and trunks of *S. japonica* in Japan.

**SORBARIA.** SCHIZONOTUS. Ag. FALSE SPIREA. Ornamental woody plants related to *Spiraea*.

**Leptosphaeria spiraeae** Karst. On branches of *S. sorbifolia* in France.

**Phyllosticta spiraeina** Brun. See *Spiraea*.

**SORBUS.** MOUNTAIN ASH. Trees grown for their handsome foliage, white flowers and ornamental red fruit.

**Cercospora ariae** Fckl. On leaves of *S. aria* in Germany.

**Cercospora kriegeriana** Bres. Rufous-brown leaf spots on *S. aucuparia* in Germany.

**Fusicladium orbiculatum** (Desm.) Thuem. Brown scabby areas on leaves of *S. aria*, *S. aucuparia*, *S. domestica*, and *S. torminalis* in Europe.

**Gloeosporium aucupariae** P. Henn. Dark-brown anthracnose spots on fruit of *S. aucuparia* in Russia.

**Gloeosporium sorbi** Rostr. Numerous subcircular ashen leaf spots on *S. fennica* in Finland.

**Gymnosporangium miyabei** Yam. and Miy. See *Chamaecyparis*.

**Gymnosporangium solenoides** (Diet.) Kern. See *Chamaecyparis*.

**Gymnosporangium torminali-juniperinum** Ed. Fisch. See *Juniperus*.

**Laestadia radiata** (Walbr.) Sacc. On leaves of *S. torminalis* in Germany.

**Leptosphaeria sorbi** Jacz. Subcircular gray leaf spots on *S. aucuparia* in Siberia.

**Marsonia sorbi** P. Magn. Indefinite, yellow-brown leaf spots on *S. aria* in Europe.

**Mycosphaerella cinerascens** (Fckl.) Vgr. On leaves of *S. aria* and *S. scandica* in Denmark and Germany.

**Mycosphaerella topographica** (Sacc.) Vgr. On leaves of *S. aucuparia* and *S. torminalis* in Denmark and Italy.

**Ochropsora ariae** (Fckl.) Syd. Yellow leaf rust on *S. americana*, *S. amicana*, *S. aria*, *S. aucuparia*, *S. fennica*, *S. intermedia*, *S. latifolia*, *S. scandica*, *S. torminalis*, *Amelanchier canadensis*, *Aruncus sylvestris*, *Pyrus communis*, and *Malus* in Europe and Formosa. The aecial stage occurs on *Anemone nemorosa*.

**Phyllohendersonia torminalis-ariae** (Br. and Har.) Tass. On *S. aria* and *S. torminalis* in France and Italy.

**Phyllosticta aucupariae** Thuem. Subcircular ashen leaf spots with dull-brown margins on *S. aucuparia* in Siberia, Russia, Italy, and Hungary.

**Phyllosticta sorbi** West. Subcircular, then confluent, ashen leaf spots with dark-purple margins on *S. aucuparia*, *S. domestica*, *S. lanuginosa*, and *S. scandica* in Belgium and Portugal. Reported from Missouri.

**Sclerotinia aucupariae** Woron. Brown rot of fruit, which finally are mummified. On *S. aucuparia* in Finland.

**Septoria aucupariae** Bres. Dull-yellow leaf spots on *S. aucuparia* in Denmark and Germany.

**Septoria aucuparicola** Oud. Circular to irregular, then confluent, dull-brown to ashen leaf spots on *S. aucuparia* in Holland.

**Septoria hyalospora** (Mont. and Ces.) Sacc. On leaves of *S. aria*, *S. domestica*, and *S. torminalis* in Italy and Germany.

**Septoria inaequalis** Sacc. and Roum. On *S. aucuparia* in France.

**Septoria sorbi** Lasch. Small circular brown spots on leaves of *S. aucuparia* in Austria.

**Septoria sorbi-hybridae** Pass. On leaves of *S. hybrida* in Italy.

**SORGHUM.** See *Holcus*.

- SPARAXIS.** WAND FLOWER. Iridaceous, bulbous or cormous plants.  
*Uromyces sparaxidis* Syd. Yellow to dark-brown rust pustules on both leaf surfaces of *S. lineata* in Natal.
- SPARGANIUM.** BUR REED. Marsh or aquatic plants.  
*Ascochyta quadriguttulata* Kab. and Bub. Irregular linear, then confluent, yellow-brown leaf spots on *S. ramosum* in Bohemia.  
*Cylindrosporium aquaticum* (Fautr. and Roum.) Sacc. On leaves of *Sparganium* sp. in France.  
*Physoderma sparganii-ramosi* (Buesg.) Schroet. On *S. ramosum* in Germany.  
*Ramularia frutescens* Kab. and Bub. On leaves of *S. ramosum* in Bohemia.  
*Ramularia sparganii* Lindr. Circular brown leaf spots on *S. glomeratum* and *S. simplex* in Sweden, Finland, Silesia, and Austria.  
*Septoria flexuosa* Oud. On leaves of *S. ramosum* in Holland.  
*Septoria sparganii* Pass. On leaves of *S. erectum* and *S. ramosum* in Italy.
- SPARTINA.** Perennial marsh grasses.  
*Phyllosticta spartinae* Brun. Subcircular small white spots with brown margins on leaf sheaths of *S. stricta* in France.  
*Uredo spartinae-strictae* Pat. and Har. Linear brown rust pustules on leaves of *S. stricta* in France.
- SPARTIUM.** WEAVERS'-BROOM. Ornamental hardy shrubs grown for their bright-yellow flowers.  
*Physalospora euganea* Sacc. (*P. albanica* Petr.) On stems of *S. junceum* in Albania and Yugoslavia.  
*Septoria spartii* Rob. and Desm. Circular olivaceous leaf spots on *S. junceum* in France.  
*Uromyces spartii-juncei* Syd. Powdery brown rust sori on leaves of *S. junceum* in Dalmatia France, and Portugal.
- SPATHOGLOTTIS.** See Orchidaceae.
- SPECULARIA.** VENUS'S LOOKING-GLASS. Erect or decumbent annual herbs.  
*Ramularia prismatocarpi* Oud. Forms grayish areas on lower leaf surfaces of *S. speculum* in Dalmatia and Holland.  
*Septoria prismatocarpi* Desm. On leaves of *S. hybrida* and *S. speculum* in Italy and France.
- SPERGULA.** SPURRY. Annual herbs.  
*Mycosphaerella alsines* Pass. On stems and leaves of *S. campestris* in Denmark.  
*Peronospora lepigoni* Fekl. Downy mildew on leaves of *S. campestris* and *S. marina* in north Europe.  
*Peronospora vernalis* Gäum. Downy mildew on leaves of *S. pentandra* and *S. vernalis* in Germany.  
*Puccinia spergulae* DC. Brown to dark-brown rust pustules on leaf blades and petioles and stems of *S. arvensis*, *S. pentandra*, *S. vernalis*, and *Spergularia rubra* in Europe. Reported from New York.  
*Septoria spergulae* West. On leaves and stems of *S. arvensis* in Sweden and Belgium.  
*Uromyces sparsus* (Kze. and Schm.) Lév. Leaf and stem rust on *S. marginata*, *S. media*, *S. (Spergularia) rubra*, and *S. salina* in French North Africa and Europe.
- SPHAERALCEA.** PHYMOSIA Ag. GLOBE MALLOW. Shrubs and herbs.  
*Aecidium sphaeralcearum* (Speg.) Sacc. and Trott. Leaf rust on *S. bonariensis* and *S. cisplatina* in Argentina.  
*Uromyces costesianus* Speg. Leaf rust on *S. velutina* in Chile.  
*Uromyces platysporus* Speg. Dark-brown leaf rust on *S. miniata* and *S. rhombifolia* in Argentina.
- SPIGELIA.** PINKROOT Annual and perennial herbs.  
*Coleosporium spigeliae* Arth. Yellow powdery rust pustules on leaves of *S. humboldtiana* in Salvador and Guatemala.  
*Puccinia spigeliae* Syd. Leaf rust on *Spigelia* sp. in Brazil.  
*Septoria spigeliae* P. Henn. Subcircular dull-brown leaf spots with dark-brown borders on *S. anthelmintica* in Jamaica and Brazil.
- SPILANTHES.** Herbs, mostly weeds, a few species cultivated for their yellow flower heads.  
*Entyloma spilanthis* Speg. Smut sori on leaves of *S. leptophylla* in Argentina.  
*Puccinia africana* Cke. Dark-brown rust sori on brown sunken spots on *S. acmella*, *S. africana*, and *S. oleracea* in Abyssinia, Madagascar, and tropical and south Africa.  
*Puccinia barranquillae* Mayor. Leaf rust on *S. urens* in Colombia.  
*Puccinia melampodii* Diet. and Holw. See Zinnia.  
*Puccinia spilanthis* Mayor. Leaf rust on *S. americana* and *S. ciliata* in Colombia.  
*Puccinia spilanthis* P. Henn. Dark-brown rust sori on leaves of *S. ciliata*, *S. oleracea*, *S. salzmanni*, and *S. uliginosa* in Martinique, Brazil, Argentina, and Peru.
- SPINACIA.** SPINACH. Annual herbs, cultivated for "greens."  
*Aecidium spinaciae* Rostr. Leaf rust on *S. tetrandra* in Russia.  
*Alternaria spinaciae* Allesch. and Noack. On living leaves of *S. oleracea* in Brazil.  
*Ascochyta chenopodii* Rostr. See Atriplex.  
*Cercospora spinaciae* Oud. Pale greenish or yellowish, often confluent, leaf spots on *S. oleracea* in Holland.  
*Gloeosporium spinaciae* Ell. and Fautr. Subcircular, then confluent, brown leaf spots on *S. oleracea* in France.  
*Phyllosticta spinaciae* A. Zimm. Large irregular leaf spots on *S. oleracea* in Belgium and Austria.  
*Ramularia spinaciae* Nypels. Brown leaf spots on *S. oleracea* in Belgium.  
*Septoria spinaciae* West. Circular leaf spots on *S. oleracea* in France, Russia, and Belgium.
- SPIRAEA.** SPIREA. Bridal wreath. Shrubs grown for their profuse, handsome white or pink flowers.  
*Aphelenchus olesistus* Ritz. Bos. See Begonia.  
*Cercospora spiraeae* Thuem. Irregular, often confluent, dull-brown leaf spots on *S. chamaedryfolia* and *Opulaster opulifolius* (*S. opulifolia*) in Siberia.  
*Cercospora callosa* Allesch. Subcircular brown, then ashen-white, leaf spots with brown margins on *S. japonica* (*S. callosa*) in Germany.  
*Cylindrosporium filipendulae* Thuem. See Filipendula.  
*Excipula spiraeae* Thuem. On leaves of *S. thalictroides* in Siberia.  
*Gloeosporium spiraeae* Bres. Indefinite dull-brown leaf spots on *Opulaster opulifolius* (*S. opulifolia*) in Germany.  
*Heterosporium spiraeae* Syd. Dull-brown irregular, then confluent, spots on leaves of *Spiraea* sp. in Austria.  
*Ochropsora ariae* (Fekl.) Syd. See Sorbus.  
*Phyllosticta ariaefolia* Allesch. Subcircular to irregular dull-brown leaf spots on *S. (Holodiscus) ariaefolia* in Germany.  
*Phyllosticta crenatae* Brun. Oblong to angular leaf spots, black above, grayish-brown beneath, on *S. crenata* in France.  
*Phyllosticta spiraeae-salicifoliae* Kab. and Bub. Circular to oblong dull-yellow to dark-brown leaf spots on *S. salicifolia* in Bohemia.

**SPIRAEA**—Continued.

- Phyllosticta spiraeina** Brun. Subcircular brown leaf spots with red-brown margins on *S. bumalda* and *Sorbaria sorbifolia* in France and Germany.
- Septoriaascochytoidea** Sacc. Subcircular to oblong ochraceous to dull-brown leaf spots on *S. chamaedryfolia* and *S. decumbens* in Siberia.
- Triphragmium filipendulae** (Lasch.) Pass. See *Filipendula*.
- SPIRANTHES**. LADY'S-TRESSES. See *Orchidaceae*.
- SPONDIAS**. YELLOW MOMBIN. JOBO. SPANISH PLUM. Tropical fruit trees.
- Ascochyta spondiacearum** A. L. Sm. White leaf spots with dull-brown margins on *S. mombin* in Angola.
- Endothia havanensis** Bruner. See *Eucalyptus*.
- Fomes lamaensis** Murr. See *Hevea*.
- Kuchneola aliena** Syd. and Butl. Leaf rust on *S. mangifera* in India and Ceylon.
- Uredo spondiadiis** Petch. Leaf rust on *S. mangifera* in Ceylon.
- SPOBOBOLUS**. DROPEED. Grasses, a few species cultivated for forage.
- Entyloma majus** Har. and Pat. Smut sori on small gray leaf spots on *S. spicatus* in the Congo.
- Phyllachora sporoboli** Pat. Elongate black stromata on *S. pungens* in Algeria and the Philippines.
- Puccinia spegazziniella** Sacc. and Trav. Dark-brown rust pustules on leaves of *S. asperifolius* in Argentina.
- Tilletia zonata** Bref. Black powdery smut masses in flowers of *S. ligularis* in Ecuador.
- Uredo egenula** Arth. Brown powdery leaf rust on *S. argutus* in Jamaica.
- Uromyces major** Arth. Brown leaf rust on *S. diander* and *S. indicus*, in India, Ceylon, and Trinidad.
- Uromyces tenuicutis** McAlp. Yellow to brown rust pustules on leaves and stems of *S. indicus* in Australia and Japan.
- Ustilaginoidea ochracea** P. Henn. In spikes of *S. elongatum* in North Borneo.
- Ustilago schlechteri** P. Henn. Black smut sori distorting the panicles and splitting the stems of *Sporobolus* sp. in the Union of South Africa.
- STACHYS**. BETONY. Tall perennial herbs or diffuse winter annuals.
- Ovularia betonicae** Mass. On leaves of *S. alopecuros* in Austria.
- Ovularia stachydis** Bres. On leaves of *S. palustris* in Germany.
- Peronospora stachydis** Syd. Downy mildew on leaves of *S. arvensis* in Europe.
- Phyllosticta stachydis** Brun. and var. *arvensis* Allesch. Angular brown leaf spots on *S. arvensis* and *S. sylvatica* in France and Germany.
- Phyllosticta variicolor** Bub. Irregular brown, then yellow-white, leaf spots on *S. germanica* in Hungary.
- Puccinia aethiopica** Kalchbr. and Cke. Brown leaf rust on *S. aethiopica* and *S. grandifolia* in the Union of South Africa.
- Puccinia albida** Diet. and Neg. Leaf rust on *S. grandidentata* and *S. mayorii* in Chile.
- Puccinia harioti** Lagh. Yellow and dark-brown rust pustules on leaves of *S. setifera* and *S. spectabilis* in Persia and Anatolia.
- Puccinia pallidissima** Speg. Brown leaf rust on *S. arvensis* and *S. lindeni* in Argentina, Brazil, Ecuador, and Guatemala.
- Puccinia stachydis** DC. Brown rust pustules on leaves, stems, and calyces of *S. recta* in Russia, France, Italy, Austria, and Germany.
- Puccinia vossii** Koern. Powdery brown rust pustules on leaves of *S. recta* and *S. setifera* in Europe.
- Ramularia cardiaca** Sacc. and Penz. Oblong to irregular spots on leaves and bracts of *S. annua* and *S. recta* in Italy and Switzerland.
- Ramularia stachydis** (Pass.) Mass. On leaves of *S. annua* and *S. recta* in Yugoslavia.
- Ramularia stachydis-alpinae** Allesch. Irregular ochraceous-brown leaf spots on *S. alpina* in Germany.
- Septoria stachydis** Rob. and Desm. Irregular olivaceous, then pale-brown, leaf spots on *S. annua*, *S. palustris*, and *S. sylvatica* in Europe.
- Ustilago betonicae** Berk. Smut in anthers of *A. alopecuros* in Austria.
- STACHYTARPHETA**. VALERIANOIDES Ag. Tropical and subtropical herbs and shrubs.
- Cercospora stachytarphetae** E. and E. Small circular white leaf spots on *S. jamaicensis* in the Bahamas and Porto Rico.
- Endophyllum stachytarphetae** (P. Henn.) W. and O. Small yellow rust pustules on leaves of *S. cayennensis* and *S. dichotoma* in Porto Rico, Brazil, and Colombia.
- Puccinia urbaniana** P. Henn. Dark-brown rust pustules on small circular brown spots on *S. cayennensis* and *S. jamaicensis* in Porto Rico, Jamaica, and Trinidad.
- STACHYURUS**. Ornamental shrubs grown for their early flowers and handsome foliage.
- Uredo stachyuri** Diet. Brown leaf rust on *S. praecox* in Japan.
- STANGERIA**. A subtropical cycad.
- Pacosphaeria stangeriae** (Zimm.) Bub. Large irregular brown leaf spots on *S. paradoxa* in Bohemia.
- STANHOPEA**. See *Orchidaceae*.
- STAPHYLEA**. BLADDER NUT. Shrubs cultivated for their white flowers, handsome foliage, and inflated podlike fruit.
- Ascochyta staphyleae** Syd. Circular brown leaf spots with purple margins on *S. trifolia* in Germany.
- Laestadia staphyleae** Hazsl. Attacks young branches of *S. pinnata* in Hungary.
- Phyllosticta osteospora** Sacc. var. *staphyleae* C. Mass. Irregular brown leaf spots on *S. pinnata* in Italy and Austria.
- Phyllosticta staphyleicola** Oud. On leaves of *S. pinnata* in Holland.
- Septoria staphyleae** Pass. Rufous leaf spots on *S. pinnata* in Italy.
- STATICE**. THRIFT. Sometimes called "sea pink." Annuals, biennials, and perennials cultivated for their flowers and foliage. See also *Armeria*.
- Ascochyta staticis** P. Nag. On leaves of *Statice* sp. in Russia.
- Phyllosticta staticis** Petr. Large brown leaf spots, lighter at the center, and with red-brown margins on *Statice* sp. in Bohemia.
- Ramularia statices** E. Rostr. Large circular to oblong ochraceous leaf spots with dull-brown margins on *S. (Limonium) behusiensis* and *S. gmelini (Limonium gmelini)* in Russia and Norway.
- Uromyces guayacurii** Speg. Leaf rust on *S. brasiliensis (Limonium brasiliense)* in Argentina.
- STELLARIA**. ALSINE Ag. Annual or perennial herbs, mostly weeds.
- Accidium stellariae** Kirch. Leaf rust on *S. graminea* in Bohemia.
- Ascochyta stellariae** Fautr. On leaves of *S. graminea* in France.
- Fabraea cerastiorum** (Fr.) Rehm. See *Cerastium*.
- Isariopsis alborosella** (Desm.) Sacc. On leaves of *S. aquatica*, *S. graminea*, *S. media*, *S. nemorum*, *Cerastium pumilum*, and *C. vulgatum* in Japan and Europe.

**STELLARIA**—Continued.

- Isariopsis stellariae** Trail. Indefinite brown spots on *S. graminea* in Scotland.
- Mycosphaerella isariphora** (Desm.) Johan. On leaves of *Stellaria* spp. in Siberia and Europe.
- Omphalospora stellariae** (Lib.) Theiss. and Syd. Shiny black stromata on leaves and stems of *S. holostea*, *S. moehringia*, and *S. nemorum* in Siberia and Europe.
- Ovularia stellariae** (Rabh.) Sacc. On leaves of *S. nemorum* in Russia, France, Switzerland, Yugoslavia, and Germany.
- Peronospora media** Gäum. Downy mildew on leaves of *S. media* and *S. nemorum* in Europe and Japan.
- Peronospora parva** Gäum. Downy mildew on leaves of *S. graminea*, *S. holostea*, and *S. uliginosa* in Europe.
- Phyllosticta holosteeae** Allesch. On leaves of *S. holostea* and *S. uliginosa* in Great Britain and Germany.
- Phyllosticta holosteiicola** Oud. Gray leaf spots on *S. holostea* in Holland.
- Placosphaeria stellariae** (Lib.) Sacc. On leaves and stems of *S. holostea* and *Dianthus monspessulanus* in Denmark, France, Italy, and Germany.
- Puccinia detonsa** Arth. and Holw. Leaf rust on *S. ovata* in Costa Rica and Guatemala.
- Ramularia stellariae** Rab. On leaves of *S. nemorum* in France.
- Septoria henslowiana** Sacc. Indefinite pale-ochraceous leaf spots on *S. media* in Malta.
- Septoria paraphysoides** Speg. On leaves and stems of *S. debilis* in Chile.
- Synchytrium stellariae** Fckl. Galls on leaves and stems of *S. media* and *S. nemorum* in Brazil, Argentina, Dalmatia, Denmark, and Germany. Reported from Mississippi.
- Uromyces stellariae** Syd. Dark-brown leaf rust on *S. kotschyana* in Persia.
- Ustilago duriaeana** Tul. See Cerastium.
- STENANDRIUM. GERARDIA** Ag. Stemless or short-stemmed acanthaceous herbs.
- Aecidium cystopoides** Speg. Leaf rust on *S. dulce* in Argentina.
- Puccinia stenandrii** Diet. and Neg. Golden and brown rust pustules on leaves of *S. dulce* in Chile.
- Uredo stenandrii** Speg. Leaf rust on *S. trinerve* in Argentina.
- STENOCAEPUS. CYBELE** Ag. Tropical trees.
- Phyllosticta stenocarpi** Tassi. Irregular brown leaf spots on *S. sinuatus* in Italy.
- STENORRHYNCHUS.** See Orchidaceae.
- STEPHANIA.** Climbing shrubs.
- Trabutia stephaniae** Rac. Circular shiny black stromata on leaves of *S. capitata* in Java.
- STEPHANOTIS.** Twining, glabrous tropical shrubs.
- Phyllosticta stephanotidis** Grove. Circular to irregular gray-white leaf spots with dark-brown margins on *S. floribunda* in Great Britain.
- STERCULIA. BOTTLE TREE.** Ornamental and economic trees.
- Ascochyta sterculiae** Tassi. Angular to irregular gray-white to dull-brown leaf spots on *S. diversifolia* (*Brachychiton diversifolium*) in Italy.
- Dothidasteroma maculosum** (B. and Br.) v. Hoeh. Circular brown stromata on leaves of *S. alata* (*Pterygata alata*) in Ceylon.
- Gloeosporium rhodospermum** Delac. Brown leaf spots on *S. acuminata* in central Africa.
- Guignardia sterculiae** Rehm. On *S. foetida* in the Philippines.
- Phyllosticta sterculiae** Wint. Angular to irregular ochraceous leaf spots on *S. acerifolia* (*Brachychiton acerifolium*) and *S. heterophylla* in Queensland and Portugal.
- Phyllosticta sterculicola** Trav. Large gray-white leaf spots on *S. frondosa* in Italy.
- Rosellinia bunodes** B. and Br. See Citrus.
- STEREOSPERMUM.** Tropical evergreen trees with large begonialike flowers.
- Phakopsora erythraea** Sacc. Black leaf rust on *S. dentatum* in Abyssinia.
- Uredo stereospermi** Syd. Leaf rust on *S. chelonioides* in Ceylon.
- STEVIA.** Tropical shrubs.
- Aecidium steviae** P. Henn. Leaf rust on *S. urticifolia* in Brazil.
- Coleosporium reichel** Diet. Leaf rust on *Stevia* sp. in Mexico.
- Coleosporium steviae** Arth. Golden rust pustules on leaves of *S. lucida*, *S. monardaefolia*, *S. reglensis*, *S. rhombifolia*, *S. salicifolia*, *S. subpubescens*, *S. trachelioides*, and *S. viscida* in Mexico and Guatemala.
- Puccinia rosea** Roth. See Ageratum.
- STIGMAPHYLLON. AMAZON VINE.** Tropical woody vines.
- Mycosphaerella stigmaphylli** Rangel. On leaves of *S. ciliatum* in Brazil.
- Phyllosticta stigmaphylli** Speg. Circular brown leaf spots on *S. littorale* in Argentina.
- Puccinia inflata** Arth. Brown leaf rust on *S. lingulatum*, *S. periplocifolium*, and *S. sagraeanum* in Porto Rico and Cuba.
- Puccinia insueta** Wint. Brown to black rust pustules on leaves of *S. jatrophaeifolium* and *S. littorale* in Brazil, Uruguay, and Ecuador.
- STILLINGIA. CHINESE TALLOW TREE.** Tropical American shrubs and herbs.
- Cercospora sebiferae** Pat. On leaves of *S. sebifera* (*Sapium sebiferum*) in Indo-China.
- STIPA. FEATHER GRASS.** Perennial grasses.
- Puccinia flavescens** McAlp. Brown to black rust pustules on leaves of *S. flavescens*, *S. pubescens*, and *S. scabra* in Australia.
- Puccinia hierochloae** S. Ito. See Hierochloa.
- Puccinia oligocarpa** Syd. and Butl. Leaf rust on *Stipa* sp. in India.
- Puccinia stipae-sibiricae** S. Ito. Leaf rust on *S. sibirica* in Japan.
- Puccinia volgensis** Nawasch. Powdery brown rust pustules on leaves of *S. barbata* and *S. pennata* in Russia.
- Septoria stipae** Died. On leaves of *S. capillata* in Germany.
- Sorosporium tumefaciens** McAlp. Olivaceous to dark-brown smut sori involving entire panicles of *S. pubescens* and *Stipa* sp. in Australia.
- Sphaelotheca macrochloae** (Pat.) Maire. Smut on *S. tenacissima* in Tripoli.
- Tilletia hypsophila** Speg. Powdery brown masses of smut spores in ovaries of *S. caespitosa* and *S. tenuissima* in Argentina.
- Uredo pencana** Diet. and Neg. Brown leaf rust on *S. manicata* in Chile.
- Urocystis stipae** McAlp. Smut sori as elongated dark-colored streaks on leaves, sheaths, and stems of *S. luehmanni* in Australia.
- Uromyces argentinus** Speg. Yellow to cinnamon-brown rust pustules on leaves of *S. neesiana* in Argentina.
- Uromyces mussooriensis** Syd. Brown leaf rust on *S. sibirica* in India.
- Ustilago comburens** Ludw. Powdery black smut sori in culms and inflorescences of *Stipa* sp. in Australia.
- Ustilago macrochloae** Pat. Inflorescences of *S. tenacissima* in Algeria converted into masses of smut spores.
- Ustilago nummularia** Speg. Black smut sori in inflorescences of *Stipa* sp. in Argentina.

**STIPA**—Continued.

**Ustilago stipae-barbatae** Maire. Smut on *S. barbata* and *S. gigantea* in French North Africa.

**Ustilago stiparum** Speg. Olivaceous masses of smut spores in panicles of *Stipa* sp. in Chile.

**Ustilago stipicola** Speg. Powdery black smut sori in panicles of *S. filiculmis* and *S. setigera* in Argentina and Uruguay.

**STIZOLOBIUM**. See *Mucuna*.**STOBÆA**. Shrubs or herbs.

**Puccinia stobæae** McO. and var. **woodii** Syd. Yellow and dark-brown rust pustules on leaves of *S. membranifolia* and *S. speciosa* in the Union of South Africa.

**STRANVAESIA**. Shrubs grown for their handsome foliage and attractive fruit and flowers.

**Aecidium stranvaesiae** Syd. Leaf rust on *S. glaucescens* in India.

**STRATIOTES**. WATER SOLDIER. Perennial aquatic herbs.

**Leptosphaeria stratiotis** Oud. On leaves of *S. aloides* in Holland.

**Phyllosticta aloidis** Oud. On leaves of *S. aloides* in Holland.

**Phyllosticta stratiotis** Oud. Dull-brown leaf spots on *S. aloides* in Holland.

**Septoria stratiotis** Oud. On leaves of *S. aloides* in Holland.

**STRELITZIA**. BIRD-OF-PARADISE FLOWER. Perennial herbs.

**Endothella strelitziae** (Cke.) Theiss. and Syd. Circular dull-black stromata on leaf spots which are wine-red beneath, on *S. regina* in the Union of South Africa.

**STROBILANTHES**. CONEHEAD. Acanthaceous ornamental herbs and shrubs.

**Aecidium strobilanthes** Barcl. Leaf rust on *Strobilanthes* sp. in the Philippines and India.

**Corticium salmonicolor** B. and Br. See *Citrus*.

**Puccinia aggregata** Syd. Brown rust pustules on sunken brown leaf spots on *S. barbatus* in India.

**Puccinia polliniae** Barcl. Yellow and brown to black rust pustules on leaf blades and petioles and stems of *S. cuspidatus*, *S. dalhousianus*, and *Pollinia nuda* in India and Indo-China.

**STROPHANTHUS**. Tropical shrubs.

**Hemileia strophanthi** Rac. Yellow rust pustules on *S. dichotomus* and *S. hispidus* in Gold Coast Colony.

**STRYCHNOS**. Shrubs and trees of economic importance, cultivated for their fruit and medicinal properties.

**Meliola steniospora** Hint. Subcircular black superficial fungus patches on leaves of *S. nux-vomica* in India.

**Phyllosticta strychni** Allesch. Subcircular ochraceous leaf spots with dark-brown margins on *S. stuhlmanni* in tropical Africa.

**Puccinia lindaviana** P. Henn. Dark-brown rust pustules on leaves of *S. henningsii* in tropical Africa and the Union of South Africa.

**Uredo strychni** P. Henn. Yellow leaf rust on *Strychnos* sp. in the Congo.

**STYLIDIUM**. CANDOLLEA Ag. Herbs or subshrubs.

**Puccinia stylidii** McAlp. Orange-yellow and dark-brown to black rust pustules on leaves of *S. graminifolium* in Tasmania.

**STYRAX**. SNOWBELL. Ornamental woody plants grown for their flowers.

**Atichia millardeti** Racib. See *Cinnamomum*.

**Monopus pulverulentus** (B. and C.) Theiss. and Syd. Gray-black stromata on leaves of *Styrax* sp. in Cuba.

**Pucciniastrum stryacinum** Hirats. Leaf rust on *S. japonica* and *S. obassia* in Japan.

**Schneepia guaranica** Speg. On *S. leprosum* and *S. parvifolia* in Brazil.

**SUTHERLANDIA**. Tropical shrubs grown for their flowers.

**Stigmatula sutherlandiae** (K. and Cke.) Syd. On leaves of *Sutherlandia* sp. in the Union of South Africa.

**SWERTIA**. Erect perennial and annual herbs grown in borders and rock gardens.

**Albugo swertiae** Berl. and Kom. Small white powdery pustules on leaves of *S. connata* in Manchuria and Russia.

**Septoria swertiae** Pat. Circular rufous leaf spots on *Swertia* sp. in China.

**Uredo opheliae** Syd. Brown rust on leaves and stems of *S. angustifolia* in India.

**SWIETENIA**. MAHOGANY. Tropical timber trees.

**Tylenchus mahogani** Cobb. This nematode is found in the bark tissues of *S. mahogani* in Barbados, probably parasitic and capable of causing injury.

**SYMPHORICARPOS**. SNOWBERRY. Ornamental shrubs.

**Ascochyta grandispora** Kab. and Bub. Subcircular ochraceous to brown leaf spots on *S. orbiculatus* in Bohemia.

**Ascochyta symphoricae** Kab. and Bub. Circular to irregular spots on leaves of *Symphoricarpos* sp. in Bohemia and France. *A. symphoricae* Br. and Har. has priority.

**Ascochyta symphoricarpi** Pass. On branches of *S. albus* (*S. racemosus*) in Italy and France.

**Heterosporium symphoricarpi** Ranoj. On branches of *S. albus* (*S. racemosus*) in Yugoslavia.

**Phyllosticta symphoriella** Sacc. and March. Leaf spots on *S. cinerascens* in Belgium.

**SYMPHYSTEMUM**. Bulbous plants.

**Uromyces symphyostemi** Speg. Brown linear rust pustules on leaves of *S. narcissoides* in Patagonia.

**SYMPHYTUM**. COMFREY. Erect, often hispid, herbs.

**Aecidium asperifolii** Pers. Leaf rust on *S. tuberosum* in Austria.

**Aecidium symphyti** Thuem. Leaf rust on *S. tuberosum* in Spain and Dalmatia.

**Cercospora agnostoica** Speg. Subcircular pale-brown leaf spots on *S. asperifolium* in Brazil.

**Cylin drosporium myosotidis** Sacc. See *Myosotis*.

**Entyloma serotinum** Schroet. See *Borago*.

**Melampsorella symphyti** Bub. Powdery brown rust pustules on leaves of *S. bulbosum*, *S. cordatum*, *S. ibericum*, *S. officinale*, *S. tauricum*, and *S. tuberosum* in Europe. The acial stage occurs on the needles of *Abies pectinata*.

**Ovularia asperifolii** Sacc. and var. **symphyti-tuberosi** Allesch. On leaves of *S. officinale* and *S. tuberosum* in Germany.

**Ovularia farinosa** (Bon.) Sacc. On leaves of *S. officinale* in Germany and Russia.

**Peronospora symphyti** Gäum. Downy mildew on leaves of *S. bulbosum*, *S. cordatum*, and *S. tuberosum* in southern Europe.

**Physoderma speciosum** Schroet. Brown areas on leaf blades and petioles and stems of *S. officinale* in Silesia and Germany.

**Puccinia bromina** Eriks. See *Bromus*.

**Ramularia symphyti-tuberosi** (Allesch.) Jaap. On leaves of *S. tuberosum* in Dalmatia. Possibly the same as *Ovularia farinosa*.

- SYMPLOCOS.** SWEETLEAF. Shrubs grown chiefly for their attractive flowers and fruits.
- Cocconia placentia** (B. and Br.) Sacc. Dull-black stromata on brown leaf spots on *S. spicata* and *Pavetta indica* in Ceylon.
- Exobasidium indicum** Syd. and Butl. Sordid white galls on leaves of *S. theaeifolia* and *Symplocos* sp. in India and Ceylon.
- Exobasidium symploci-fasciculatae** Rac. Circular to ovate spots on *S. fasciculata* in Java.
- Exobasidium symploci-japonicae** Kus. Distorts young buds of *S. japonica*, *S. lucida*, and *Maba buxifolia* in Japan.
- Fomes pseudoferreus** Wakef. See Hevea.
- Mycosphaerella bhauria** (Cke.) Lind. On leaves of *S. spicata* in India and Malaya.
- Phacidium symplocinum** Syd. On leaves of *Symplocos* sp. in India.
- Phyllachora symploci** Pat. Black stromata on leaves of *S. japonica*, *S. lucida*, *S. neriifolia*, and *S. thwaitesii* in Indo-China, Japan, and Australia.
- Phyalospora symploci** Rac. Circular to irregular yellow leaf spots on *S. fasciculata* in Java.
- Rhytisma austro-caledonicum** Crié. Black irregular stromata on leaves of *S. arborea*, *S. caerulea*, *S. gracilis*, *S. lenormandiana*, *S. mandiana*, and *S. vielillardii* in New Caledonia.
- Rosellinia arcuata** Petch. See Thea.
- Septoria sydowii** Henn. and Sacc. Small subcircular to irregular gray-white leaf spots on *S. crataegoides*, *S. myrtacea*, and *S. prunifolia* in Japan and Brazil.
- Septoria symploci** Allesch. and P. Henn. On *Symplocos* sp. in Brazil.
- Trochila symploci** P. Henn. On leaves of *S. japonica* in Japan.
- Uredo inouyei** P. Henn. and Shir. A rust deforming leaves and branches of *Symplocos* sp. in Japan.
- SYRINGA.** LILAC. Shrubs cultivated for their flowers.
- Ascochyta orientalis** Bond. Confluent gray leaf spots with dark-brown margins on *S. vulgaris* in Russia.
- Ascochyta syringae** Bres. On leaves of *S. chinensis* and *S. vulgaris* in Russia, Denmark, Italy, and Germany.
- Ascochyta syringicola** Bub. and Kab. Circular to angular dull-yellow leaf spots on *S. vulgaris* in Bohemia.
- Ascochyta syringae** Jaap. On fruit capsules of *S. vulgaris* in Germany.
- Gloeosporium syringae** Allesch. Irregular ochraceous to brown leaf spots, often marginal, or involving entire leaf blades on *S. alba* and *S. vulgaris* in Russia and Germany.
- Helminthosporium syringae** Kleb. Irregular brown areas on leaves, gradually involving entire area of the blades on *S. vulgaris* in Great Britain and Germany.
- Heterosporium syringae** Oud. Large irregular gray-brown blotches on leaves of *S. vulgaris* in Spain, Italy, France, Holland, and Germany.
- Phyllosticta superflua** Oud. Circular, then confluent, pale ashen-brown leaf spots on *S. vulgaris* in Holland.
- Phyllosticta syringophila** Oud. On leaves of *S. vulgaris* in Holland.
- Phytophthora syringae** Kleb. This downy mildew causes a leaf-blight and die-back of *S. persica*, *S. vulgaris*, and other woody plants in Europe, including *Acer* sp., *Aesculus* sp., *Alnus* sp., *Corylus* sp., *Crataegus oxyacantha*, *Forsythia viridissima*, *Jasminum nudiflorum*, *Ligustrum vulgare*, *Malus*, *Prunus cerasus*, *P. domestica*, *Pyrus communis*, and *Tilia* sp.
- Pseudomonas syringae** v. Hall. Large black blotches on young leaves and twigs of *S. vulgaris* in Holland, Great Britain, and Germany. The organism is said to be able to attack also *Malus*, *Populus*, *Prunus*, and *Pyrus*.
- Septoria syringae** Sacc. and Speg. Dull-yellow leaf spots on *S. chinensis* and *S. vulgaris* in Italy and France.
- SYZGIUM.** Tropical trees and shrubs.
- Atichia millardeti** Racib. See Cinnamomum.
- Lizoniella syzgiai** (Rac.) Sacc. and Trott. Subcircular black leaf spots on *Syzgium* sp. in Java.
- TABEBUIA.** Ornamental evergreen trees grown for their showy flowers.
- Mycosphaerella tabebuiae** Miles. Amphigenous small white transparent leaf spots with brown-purple margins on *T. haemantha* in Porto Rico.
- Phyllachora sordida** Speg. Black subcircular stromata on pale, indefinite leaf spots on *Tabebuia* sp. in Brazil.
- Phyllachora tabebuiae** (Rehm.) Theiss and Syd. Shiny black stromata on small dark-brown leaf spots on *T. leucoxydon* in Brazil.
- Physalospora tabebuiae** Rehm. On leaves of *T. leucoxydon* in Brazil.
- Prospodium bahamense** Arth. Leaf rust on *T. bahamensis* and *T. leucoxydon* in the Bahamas.
- Prospodium suppressum** Arth. Leaf rust on *Tabebuia* sp. in Trinidad.
- TABERNAEMONTANA.** CRAPE JASMINE. Evergreen trees and shrubs.
- Aecidium ceraceum** B. and Br. Leaf rust on *T. dichotoma* in Ceylon, Brazil, and Argentina.
- Aecidium ochraceum** Speg. Indefinite pale-ochraceous leaf spots on *T. australis* in Paraguay.
- Caeoma apocyni** McAlp. Brownish-yellow rust pustules on leaves of *T. orientalis* in Australia.
- Cercospora tabernaemontanae** Syd. On leaves of *T. pandocahui* and *Tabernaemontana* sp. in Malaya and the Philippines.
- Fomes pectinatus** (Kl.) Cke. Attacks *T. sphaerocarpa* in Java.
- Gloeosporium holstii** P. Henn. Anthracnose on leaves of *T. dichotoma* in Ceylon.
- Hemileia jurensis** Syd. Pale-yellow leaf rust on *Tabernaemontana* sp. in Brazil.
- Puccinia cookei** DeT. Yellow to brown rust pustules on leaves of *T. ventricosa* in India and the Union of South Africa.
- Puccinia engleriana** P. Henn. Leaf rust on *T. heyneana* and *T. pandocahui* in India and the Philippines.
- Puccinia tabernaemontanae** B. and Br. Leaf rust on *T. dichotoma* in Ceylon.
- Seynesia fusco-paraphysata** P. Henn. On leaves of *Tabernaemontana* sp. in Tanganyika.
- Uredo manilensis** Syd. Brown leaf rust on *T. coronaria* and *T. polygama* in the Philippines.
- TACCA.** Perennial herbs with tuberous or creeping rhizomes.
- Cercospora** sp. On leaves of *T. pinnatifida* in Fiji.
- Cercosporina taccae** Syd. Circular brown leaf spots on *T. palmata* in the Philippines.
- TACSONIA.** Tendril-climbing vines resembling *Passiflora*.
- Seynesia lagerheimii** Rehm. On leaves of *Tacsonia* sp. in Ecuador.
- TAGETES.** MARIGOLD. Garden annuals.
- Puccinia tagetica** Diet. and Holw. Powdery brown to black rust pustules on leaves of *T. erecta*, *T. filifolia*, *T. lucida*, *T. micrantha*, *T. patula*, *T. microglossa*, and *T. tenuifolia* in Porto Rico, Mexico, Costa Rica, and northern South America.
- TAINIA.** See *Orchidaceae*.

- TAMARINDUS.** TAMARIND. Tropical leguminous tree. Cultivated for ornament and fruit.  
**Exosporium tamarindi** Syd. On leaves of *T. indica* in India.  
**Gloeosporium tamarindi** P. Henn. Anthracnose on leaves of *T. indica* in Tanganyika.  
**Mycosphaerella tamarindi** P. Henn. Dull-brown leaf spots on *T. indica* in Tanganyika.
- TAMARIX.** Ornamental shrubs or trees, cultivated for their graceful foliage and showy clusters of white or pink flowers.  
**Pyrenopeziza tamaricis** (Roum.) Sacc. See Myricaria.
- TAMONEA.** Tropical shrubs.  
**Gloeosporium cyanophylli** P. Henn. Dull-brown leaf spots on *T. (Cyanophyllum) magnifica* in Germany.
- TAMUS.** Black bryony. Herbaceous perennials with tuberous roots related to Dioscorea.  
**Cercospora scandens** Sacc. and Wint. and var. **longissima** Gz. Frag. Small subcircular dull-brown leaf spots with yellow margins on *T. communis* in Italy, Spain, and Russia.  
**Cercospora tamicola** Lamb. and Fautr. Irregular gray leaf spots on *T. communis* in Italy and France.  
**Phyllosticta tami** Sacc. Subcircular to oblong brown leaf spots with red margins on *T. communis* in Italy.  
**Septoria sarmenticia** Sacc. On stems of *T. communis* in France.  
**Septoria tami** West. Subcircular to angular brown leaf spots with purple-brown margins on *T. communis* in Belgium.
- TANACETUM.** TANSY. Annual or perennial herbs with small yellow flower heads.  
**Peronospora tanacetii** Gäum. Downy mildew on leaves of *T. vulgare* in Europe.  
**Puccinia balsamitae** (Strauss.) Rabh. See Pyrethrum.  
**Puccinia seriata** Syd. Black rust sori on leaves and stems of *Tanacetum* sp. in Kurdistan.  
**Puccinia tanacetii** DC. Dark-brown to black rust pustules on leaves of *T. corymbosum* and *T. vulgare* in Europe.  
**Ramularia tanacetii** Lind. Brown spots, occupying most of leaf area of *T. vulgare* in Denmark and Spain.  
**Septoria tanacetii** Niessl. Indefinite irregular often confluent, dull-brown leaf spots on *T. vulgare* in Bulgaria, Bohemia, and France.  
**Septoria tanacetii-macrophyli** Bub. Elongate or irregular, often confluent, dull-yellow to brown leaf spots on *T. macrophyllum* in Central Europe.
- TARAXACUM.** LEONTODON. Ag. DANDELION. Weedy perennials, sometimes cultivated as a vegetable crop.  
**Cercospora augustana** Ferr. Irregular ochraceous leaf spots on *T. officinale* in Italy.  
**Erysiphe taurica** Lévl. See Althaea.  
**Phyllosticta taraxaci** Höll. Irregular ashen black-bordered leaf spots on *T. officinale* in Hungary.  
**Protomyces pachydermus** Thuem. On peduncles and leaves of *T. officinale* in Japan, Scandinavia, Italy, Switzerland, Austria, and Germany.  
**Puccinia decipiens** Mass. Powdery black rust pustules on leaves of *T. canescens* and *T. montanum* in Persia and Kurdistan.  
**Puccinia kurdistani** Cke. Brown leaf rust on *T. glaucum* in Kurdistan.  
**Puccinia silvatica** Schroet. Leaf rust on *T. dens-leonis*, *T. officinale*, *Crepis biennis*, *Lappa officinalis*, *Senecio fuchsii*, *S. nemorensis*, *S. sarracenicum*, and *Carex* spp. in Siberia and Russia.  
**Puccinia taraxaci-bithynici** R. Maire. Brown leaf rust on *T. bithynicum* in Asia Minor.  
**Puccinia variabilis** Grev. Yellow to dark-brown rust pustules on leaves of *T. collinum*, *T. officinale* and *T. palustre* in Siberia and Europe.  
**Ramularia lampsana** (Desm.) Sacc. See Cichorium.  
**Septoria taraxaci** Höll. Circular ashen, black-bordered leaf spots on *T. officinale* in Hungary. Sydow has given the same name to a species on *T. officinale* in Japan.
- TAXUS.** YEW. Ornamental coniferous trees.  
**Gloeosporium cavarae** (Mont.) Sacc. and Syd. On needles of *T. baccata* in Italy.  
**Gloeosporium taxicolum** Allesch. On needles of *T. baccata* in Denmark, France, and Germany.  
**Lophodermium abietis** Rostr. See Picea.  
**Macrophoma taxi** Berl. and Vogl. Causes leaves of *T. baccata* in Europe to turn brown.  
**Phyllosticta taxi** Höll. Ochraceous spots on needles of *T. baccata* in Hungary.
- TECOMA.** TRUMPET BUSH. Ornamental tropical shrubs and small trees grown for their showy flowers. See also Tabebuia.  
**Accidium simplicius** Arth. and Johnst. Leaf rust on *T. pentaphylla* in Cuba.  
**Cercospora leprosa** Speg. On leaves of *T. arborea* in Brazil.  
**Cercosporina stenolobii** Speg. Subcircular definite dull-rufous leaf spots on *T. (Stenolobium) stans* in Argentina.  
**Cylindrosporium aureum** Speg. Indefinite dull-brown leaf spots on *Tecoma* sp. in Brazil.  
**Munkiella guaranitica** Speg. On leaves of *Tecoma* sp. in Brazil and Paraguay.  
**Ovulariopsis obclavata** Wakef. On leaves of *T. leucoxydon* in Barbados.  
**Phyllosticta crysiphoides** Sacc. On leaves of *T. (Bignonia) radicans* in France.  
**Phyllosticta henriquesii** Thuem. On leaves of *T. (Bignonia) radicans* in Italy.  
**Phyllosticta tecomae** Sacc. Dull-brown to ochraceous leaf spots on *T. (Bignonia) radicans* in Italy and Portugal.  
**Prospodium appendiculatum** (Wint.) Arth. Leaf rust on *T. mollis* and *T. (Stenolobium) stans* in Porto Rico, Cuba, Costa Rica, Mexico, and Trinidad.  
**Prospodium bahamensis** Arth. See Tabebuia.  
**Prospodium plagiopus** (Mont.) Arth. Brown rust pustules on leaves of *T. lepidota* in Cuba.  
**Puccinia exitiosa** Syd. and Holw. Brown leaf rust on *T. mollis* in Mexico.  
**Puccinia maligna** Diet. Leaf rust on *T. stans* in Mexico.  
**Uredo longiaculeata** P. Henn. Powdery dull-brown to ochraceous rust pustules on leaves of *Tecoma* sp. in Brazil.
- TECOPHILAEA.** CHILIAN CROCUS. Bulbous plants.  
**Puccinia roseanae** Arth. Leaf rust on *T. roseana* in Peru.
- TECTARIA.** Ferns.  
**Uredo gymnogrammes** P. Henn. See Dryopteris.
- TECTONA.** TEAK. Valuable timber trees.  
**Accidium effusum** Niessl. Leaf rust on *T. grandis* in India.  
**Heart-rot.** A heart-rot of *T. grandis* due to an undetermined fungus is reported from Java. The heartwood is stained dark brown and finally rots. The tops of diseased trees die.  
**Uncinula tectonae** Salm. Powdery mildew on upper leaf surfaces of *T. grandis* in India.  
**Uredo tectonae** Racib. Brown leaf rust on *T. grandis* in Java, Indo-China, Ceylon, and India.
- TELLIMA.** FALSE ALUMROOT. Erect, hardy, herbaceous perennials.  
**Phyllosticta tellimae** Tassi. Subcircular, brown leaf spots on *T. grandiflora* in Italy.

- TEPHROSIA. CRACCA** Ag. HOARY PEA. Perennial herbs or shrubs, some species cultivated for forage.
- Corticium salmonicolor** B. and Br. See Citrus.
- Fomes pseudoferreus** Wakef. See Hevea.
- Ophiodothella edax** (Br. and B.) v. Hoeh. Shiny black stromata on leaves of *T. suberosa* in Ceylon and the Union of South Africa.
- Poria hypobrunnea** Petch. See Hevea.
- Ravenelia caulicola** Arth. Brown to black powdery rust pustules on stems of *T. cinerea* in the Bahamas.
- Ravenelia irregularis** Arth. Brown rust pustules on leaves of *T. macrantha* in Mexico.
- Ravenelia stictica** B. and Br. Yellow to brown rust pustules on leaves of *T. suberosa* in India, Ceylon, and the Union of South Africa.
- Ravenelia talpa** (Long) Arth. Brown leaf rust on *T. talpa* in Mexico.
- Ravenelia tephrosiae** Kalchbr. Yellow-brown to dark-brown rust pustules on leaves of *T. macropoda* in the Union of South Africa.
- Rosellinia arcuata** Petch. See Thea.
- Uredo tephrosiae** Rabh. Leaf rust on *T. purpurea* in Ceylon.
- Uredo tephrosiicola** P. Henn. Brown leaf rust on *T. toxicaria* and *Tephrosia* sp. in Peru and Brazil.
- TERAMNUS.** Tropical trees.
- Cercospora maricaoensis** Young. Diffuse, brown leaf spots on *T. uncinatus* in Porto Rico.
- Physopella concors** Arth. See Phaseolus.
- Uredo teramni** Mayor. Leaf rust on *T. uncinatus* in Colombia.
- Uromyces cologaniae** Arth. Brown leaf rust on *T. uncinatus* in Porto Rico.
- Uromyces vanderystii** P. Henn. Yellow-brown leaf rust on *T. labialis* in Congo.
- TERMINALIA.** INDIA ALMOND. ALMENDRO. SPANISH ALMOND. Tropical trees with many economic uses.
- Cercospora catappae** P. Henn. Circular dark-brown leaf spots on *T. catappa* in Zanzibar and the Philippines.
- Fomes lamaoensis** Murr. See Hevea.
- Gnomoniella catappae** Koord. Large indefinite angular dull-brown leaf spots on *T. catappa* in Java.
- Phyllosticta terminaliae** P. Henn. Circular brown to olivaceous leaf spots with red margins on *T. baumii* and *T. catappa* in Brazil and the Union of South Africa.
- Ramularia catappae** Rac. Circular ashen leaf spots with dull-red margins on *T. catappa* in the Philippines.
- Uredo terminaliae** P. Henn. Ochraceous rust pustules on leaves of *T. argentea* in Brazil.
- TETRAGONIA.** NEW ZEALAND SPINACH. Herbs cultivated as vegetables.
- Cercosporina tetragoniae** Speg. Subcircular leaf spots on *T. expansa* in Argentina and Russia.
- Puccinia tetragoniae** McAlp. Yellow to brown rust pustules on leaves of *T. implexicoma* in Australia.
- Tubercularia cacao** McAlp. On leaf blades and petioles of *T. implexicoma* in Australia.
- TETRAGONOLOBUS.** Leguminous shrubs related to Lotus.
- Ovularia deusta** (Fekl.) Sacc. See Lathyrus.
- Peronospora tetragonolobi** Gäum. Downy mildew on leaves of *T. siliquosus* in central Europe.
- Uromyces genistae-tinctoriae** Pers. See Caragana.
- TETRAPANA.** See Fatsia.
- TEUCRIUM.** GERMANDER. Herbs and shrubs.
- Ascochyta teucrii** Lasch. On leaves of *T. scordium* in Germany.
- Erysiphe taurica** Lévy. See Althaea.
- Ovulariopsis teucrii** Jaap. Diffuse yellow areas on leaves of *T. chamaedrys* in Dalmatia.
- Peronospora teucrii** Gäum. Downy mildew on leaves of *T. botrys* in Bohemia.
- Phyllosticta infuscata** Wint. Subcircular to irregular dull-brown leaf spots on *T. scordium* in Portugal.
- Phyllosticta obliqua** Tassi. Dull-brown zoned leaf spots on *T. flavum* and *T. pulverulentum* in Italy.
- Phyllosticta teucrii** Sacc. and Speg. Grayish-white leaf spots on *T. chamaedrys* in Italy.
- Puccinia annularis** (Strauss.) Schlecht. Brown rust pustules on leaves of *Teucrium* spp. in Europe.
- Puccinia constricta** (Lagh.) Bub. Brown leaf rust on *T. montanum* in Russia and Germany.
- Puccinia istriaca** Syd. Powdery dark-brown rust pustules on leaves and twigs of *T. aureum* and *T. polium* in Spain and the Balkans.
- Puccinia majoricensis** R. Maire. Leaf rust on *T. capitatum* in the Balearic Islands.
- Puccinia teucrii** Biv. Bernh. Dark-brown rust pustules on leaves of *T. fruticans* and *T. polium* in Morocco, Algeria, Syria, Spain, and Italy.
- Ramularia microspora** Thuem. Brown spots on leaves of *T. chamaedrys* in Austria and Italy.
- Septoria scorodoniae** Pass. Small subcircular reddish to ochraceous leaf spots on *T. scorodonia* in Belgium, France, and Italy.
- Septoria teucrii** Sacc. Circular brown to gray leaf spots with dark-purple margins on *T. chamaedrys* in France.
- Taphrina candicans** Sacc. On leaves of *T. chamaedrys* in Europe.
- THALIA.** Perennial marsh or aquatic herbs.
- Puccinia cannae** (Wint.) P. Henn. See Canna.
- Puccinia thaliae** Diet. Powdery brown to black rust pustules on leaves of *T. dealbata* in Brazil.
- THALICTRUM.** MEADOW RUE. Herbs, sometimes cultivated.
- Cercospora thalictri** Thuem. Irregular, often confluent, violet-purple leaf spots on *T. flavum*, *T. glaucum*, *T. jacquinianum*, and *T. speciosum* in Portugal.
- Haplobasidium thalictri** Eriks. Dark-brown leaf spots on *T. flavum* in Sweden.
- Marsonia clematidis** All. See Clematis.
- Oedemium thalictri** Jaap. Gray spots on leaves of *T. minus* in Germany.
- Phyllosticta leucosticta** C. Mass. Small angular spots on leaves of *T. aquilegifolium* in Italy.
- Phyllosticta thalictri** West. Angular to irregular leaf spots on *T. flavum* and *T. minus* in Belgium and Germany.
- Piasmopara alpina** (Johans.) Blytt. Downy mildew on leaves of *T. alpinum* in Scandinavia.
- Pseudopeziza repanda** Sow. On *T. minus* in Japan.
- Puccinia borealis** Juel. See Agrostis.
- Puccinia castagnei** Schroet. Leaf rust on *T. angustifolium* in France, Belgium and Switzerland.
- Puccinia rhytismoides** Johans. Black rust pustules on leaves and stems of *T. alpinum* in Sweden and Norway.
- Puccinia septentrionalis** Juel. See Polygonum.
- Puccinia sommerfeltii** Johans. Dark-brown, powdery rust pustules on leaves of *T. alpinum* and *Polygonum viviparum* in Alaska, Newfoundland, and Europe.

- THEA.** TEA. Broad-leaved evergreen shrubs cultivated for their leaves and flowers.
- Aglaospora aculeata** Petch. Causes blight of stems of *Thea* (cult.) in Ceylon.
- Ascochyta theae** Hara. On stems of *Thea* (cult.) in Japan and Ceylon.
- Bacillus theae** Hori and Bokura. The "red scaled" disease of *Thea* (cult.) is known only from Japan. On fully developed leaves small pale-brown circular spots occur, which rapidly increase in size to form large confluent, irregular, reddish-brown to black patches. Infected leaves wither and fall. The disease spreads from the leaves to the buds and twigs. The buds turn red-brown and then black. Green twigs turn black, wither, and die back for a length of 4 to 5 inches.
- Cercospora theae** v. B. de H. Circular purplish-red leaf spots with indefinite yellow-green borders on *Thea* (cult.) in Java, India, and Ceylon. The spots finally become white with narrow purple-red borders.
- Cercospora theae** Petch. Numerous, small, dark-brown to black, subcircular then irregular spots on young tea leaves. The spots coalesce, destroying the leaves. On older leaves large diffuse chocolate-brown, finally gray, patches occur with narrow purplish-black margins. Sunken purple patches appear on green twigs. Additional hosts are *Acacia decurrens*, *Eucalyptus dealbata*, *E. diversicolor*, *E. melanoxylon*, *E. paniculata*, and *E. robusta* in Ceylon.
- Chaetophoma penzigi** Sacc. On leaves of *Thea* (cult.) in Italy and the Caucasus.
- Colletotrichum camelliae** Mass. Brown blight or anthracnose on leaves and twigs of *Thea* (cult.) in Japan, India, Ceylon, Malaya, Java, Formosa, Caucasia, and Uganda. The leaf spots are at first yellowish-green, becoming red-brown or dark brown with first yellow-green margins, and finally gray. Young leaves are blackened and sometimes soft rotted. Twigs of unthrifty plants are at times attacked, becoming hard, gray, and stunted.
- Corticium** sp. Black rot of tea (*Thea*) in Ceylon is caused by a *Corticium* distinct from *C. theae* of Java. The Ceylon fungus or similar fungi also attack *Calophyllum burmanii*, *Hemidesmus indicus*, *Oxyanthus tubiflorus*, *Pithecolobium dulce*, and *Zinnia* sp.
- Corticium salmonicolor** B. and Br. See Citrus.
- Corticium theae** Bern. Black rot of tea (*Thea*) in Java, India, and Sumatra is characterized by the drying out and blackening of the leaves and die-back of the twigs due to the fungus, which forms brown fungus strands over infected organs. The disease is one of the thread-blight diseases.
- Didymostilbe coffeae** Zimm. Causes a die-back of plucked *Thea* branches in Ceylon.
- Diplodia cacaoicola** P. Henn. See Theobroma.
- Discosia theae** Cav. Circular definite, often confluent, black leaf spots on *Thea* (cult.) in Italy and the Caucasus.
- Exobasidium camelliae** Shir. and var. *gracilis* Shir. Leaves and leaf-shoots are attacked, becoming thick and fleshy or gall-like. Inflorescences are also deformed and reduced to irregular spherical masses. On *T. (Camellia) sasanqua* and *T. japonica* in Japan.
- Exobasidium reticulatum** Ito and Saw. Attacks *Thea* (cult.) in Japan and Formosa in a similar manner to the following species, except that the leaves are not blistered.
- Exobasidium vexans** Mass. The blister blight is a serious disease of cultivated *Thea* in India and probably in Formosa. It has also been found on *Rhododendron* spp. Numerous translucent spots appear on the leaves from one-fifth to 1 inch in diameter, red below, pale-green, yellowish, or pink above. The upper surfaces gradually become depressed into shallow cavities. The upper concave surfaces are smooth, shining, and pale-green, the lower surfaces dull-gray to white and powdery. Petioles and stems may also be attacked, the latter dying back. As many as 20 blisters may occur on a single leaf, remaining separate or at times coalescing.
- Fomes lamaoensis** Murr. See Hevea.
- Fomes lignosus** Klotzsch. See Hevea.
- Fomes pseudoferreus** Wakef. See Hevea.
- Fusicladium theae** K. Hara. Velvety-black fungus patches on leaves of *T. sinensis* in Japan.
- Gloeosporium theae** Zimm. Large red-brown, then ashen, leaf spots on *Thea* (cult.) in Tanganyika.
- Gloeosporium theae-sinensis** Miy. Large reddish-brown, then gray, leaf spots on *Thea* (cult.) and *T. sasanqua* in Japan. Probably identical with *Colletotrichum camelliae*.
- Guignardia camelliae** (Cke.) Butler. (*Laestadia theae* Rac.) This fungus causes the so-called "copper blight" of tea (*Thea*) in India, Ceylon, Java, and Malaya. In the early stages of the disease a peculiar coppery-colored sheen forms on the lower surfaces of the leaves, leaves so affected bending over, with the undersides outermost. Yellow-brown patches then appear, becoming finally gray and very brittle. Possibly the perfect stage of *Colletotrichum camelliae* Mass.
- Helicobasidium tanakae** Miy. See Morus.
- Helminthosporium gigasporum** B. and Br. On *Thea* (cult.) in Ceylon.
- Hendersonia theae** K. Hara. Dark-brown, then gray, leaf spots on *Thea* (cult.) in Japan.
- Hendersonia theicola** Cke. On leaves of *Thea* in Ceylon and the Caucasus.
- Hypodermopsis theae** K. Hara. Light reddish-brown circular to irregular, often confluent, patches on trunk and branches of *Thea* (cult.) in Japan.
- Leptosphaeria hottai** K. Hari. Circular to irregular large brown spots on trunks of *T. sinensis* in Japan.
- Macrophoma theae** Speschn. Subcircular gray-brown leaf spots on *Thea* (cult.) in the Caucasus.
- Macrophoma theicola** Petch. This fungus is reported as the cause of a canker of *Thea* (cult.) in Ceylon. Small, slightly sunken, dark patches first appear on the branches and develop into irregular sunken cankers.
- Marasmius** spp. Horsehair blight. See Theobroma.
- Massaria theicola** Petch. This fungus attacks the branches of the tea (*Thea*) plant in Ceylon, killing them gradually one by one. Individual branches may die slowly, the leaves withering and falling, or because of girdling a branch may be killed suddenly.
- Mycosphaerella theae** K. Hara. Large irregular dark-brown, then ashen, leaf spots on *Thea* (cult.) in Japan.
- Nectria cancri** Rutg. A wound parasite causing callused cankers on twigs, branches, and trunks of *Thea* sp. in India and Java.
- Neottispora theae** Saw. Irregular brown to ashen leaf spots with raised purplish-black margins on *Thea* (cult.) in Formosa.
- Pestalozzia theae** Saw. The gray blight of tea (*Thea sinensis*) occurs in India, Ceylon, Java, Formosa, and the Caucasus. The fungus produces subcircular to oval leaf spots which vary in size from one centimeter to almost the entire area of the leaf. The spots are dark-brown and concentrically zoned above, brighter beneath, with narrow greenish-yellow surrounding zones. Old spots become gray. Very young leaves when attacked are blackened and distorted. Twigs are also attacked at times, a die-back resulting.
- Phaeosphaerella theae** Petch. Small yellow leaf spots, circular, then angular, on *Thea* (cult.) in Ceylon. The centers of the spots turn gray or gray-brown and fall out, leaving irregular holes bordered by narrow brown zones.
- Phoma theicola** Petch. Subcircular to oval or angular bright red-brown leaf spots on *Thea* (cult.) in Ceylon.

## THEA—Continued.

**Phyllohendersonia theicola** (Cke.) Tass. On leaves of *Thea* (cult.) in Caucasia.

**Phyllosticta camelliae** West. See *Camellia*.

**Phyllosticta theae** Speschn. Subcircular gray-white leaf spots on *T. assamica* and *Thea* (cult.) in Caucasia.

**Physalospora neglecta** Petch. Causes cankers on branches of *Thea* (cult.) in Ceylon.

**Poria hypobrunnea** Petch. See *Hevea*.

**Protomyces theae** Zimm. On roots of *Thea* (cult.) in Java.

**Pseudocommis theae** Speschn. Indefinite, often confluent, gray-brown leaf spots on *T. viridis* (*Thea sinensis*?) in Russia.

**Rhizoctonia lamelifera** Small. See *Grevillea*.

**Rosellinia arcuata** Petch. This fungus causes a rotting of the bark around the collar and on the larger roots, forming a white mycelial layer between the bark and wood. Above ground the fungus forms a layer of purple-gray, then black, mycelium around the stems up to a height of about 6 inches. Some plants, although girdled, continue growth, resulting in peculiar enlargements of stems down to the point of girdling. Diseased plants often wilt and die. The hosts in addition to tea (*Thea*) are *Capsicum annuum*, *Cinnamomum camphora*, *Erythrina* sp., *Grevillea robusta*, *Manihot glaziovii*, *Panax fruticosum*, *Strobilanthes* sp., *Symplocos obtusa*, and *Tephrosia candida* in Ceylon and India.

**Rosellinia bunodes** B. and Br. See *Citrus*.

**Sclerotium zeylanicum** (B. and Br.) Petch. Attacks seedlings of *Thea* (cult.), *Caladium* spp., and other economic plants in Ceylon.

**Septobasidium acaciae** Saw. See *Acacia*.

**Septoria theae** Cav. On leaves of *T. viridis* and *Thea* (cult.) in Italy and Russia.

**Sillia theae** K. Hara. Small dark-pink or gray spots appear on trunks and branches of *T. sinensis* in Japan. These spots increase until the infected trunk or branch is girdled. The disease is known as the "shark-skin" disease because of the shape of the stromata scattered over the cankered area.

**Sphaerostilbe repens** B. and Br. See *Hevea*.

**Stilbella theae** Bern. (*Stilbum nanum* Mass.) Causes a die-back of branches of *Thea* (cult.) in Java.

**Valsa theae** K. Hara. Acts as a wound parasite of *Thea* (cult.) in Japan.

**Venturia speschnewii** Sacc. and D. Sacc. Indistinct brown leaf spots on *Thea* (cult.) in Caucasia.

**White-stem blight.** See *Hevea*.

**Zukalia nantoensis** K. Saw. This species of sooty mold and others, including *Z. theae* Saw., *Scorias capitata* Saw., and *Capnodium footii* Berk. and Desm., occur on tea (*Thea*) leaves, injuring the commercial value of the leaves.

THELYMITRA. See *Orchidaceae*.

## THEOBROMA. CACAO OR CHOCOLATE TREE. Tropical trees.

**Aspergillus delacroixii** Sacc. and Syd. Hastens the hardening and rot of pods of *T. cacao* in Colombia and the Philippines.

**Ceratocarpia theobromae** von Faber. Thick black fungus crust on leaves of *T. cacao* in Cameroon.

**Colletotrichum brachytrichum** Delacr. Broad gray-white leaf spots with brown margins on *T. cacao* in Trinidad and Ceylon.

**Colletotrichum cradwickii** Bancr. Anthracnose of pods of *T. cacao* in Jamaica and Porto Rico. Probably not distinct from *C. gloeosporioides*.

**Colletotrichum incarnatum** Zimm. Anthracnose on pods of *T. cacao* in Ceylon, Java, Belgian Congo, and Uganda, following *Phytophthora*.

**Colletotrichum luxificum** v. Hall. and Drost. This fungus was at one time assigned as the cause of the Surinam witches'-broom disease. See *Marasmius perniciosus*.

**Colletotrichum theobromae** Appel. and Str. Small black sunken areas on pods of *T. cacao* in central Africa.

**Colletotrichum theobromicolum** Delacr. Anthracnose on pods of *T. cacao* in the West Indies and Uganda. Most, if not all, of the various species of *Colletotrichum* reported as attacking leaves and pods of *T. cacao* can be referred to *C. gloeosporioides*. They are all more or less secondary in their parasitism, attacking trees weakened by adverse cultural conditions or following primary parasites such as *Phytophthora*.

**Corticium salmonicolor** B. and Br. See *Citrus*.

**Diplodia cacaoicola** P. Henn. This fungus is found on a wide range of tropical and subtropical plants and has been described under many names, among which are *D. rapax* Mass., *Botryodiplodia theobromae* Pat., *B. elastica* Petch., *Lasiodiplodia theobromae* (Pat.) Griff. and Maubl., *L. nigra* App. and Laub., and *Chaetodiplodia vanillae* Maubl. *Thyridaria tarda* Bancr. has been reported as the perfect stage. Among the many hosts are *T. cacao*, *Hevea brasiliensis*, *Vanilla planifolia*, *Ananas sativa*, *Albizia* sp., *Carica papaya*, *Cajanus indicus*, *Citrus* spp., *Castilloa elastica*, *Cocos nucifera*, *Erythrina* sp., *Cinnamomum camphora*, *Coffea arabica*, *Ficus elastica*, *Grevillea robusta*, *Indigofera* spp., *Mangifera indica*, *Manihot utilissima*, *Nicotiana tabacum*, *Saccharum officinarum*, and *Thea sinensis*. Countries reporting the fungus are Porto Rico, Cuba, French and British West Indies, Trinidad, Brazil, northern South America, Java, India, Ceylon, Malaya, the Philippines, Fiji, and central Africa.

Circular brown spots appear on cacao pods, most frequently at the basal end, and, spreading rapidly, soon bring about complete rot of the pods. The fungus acts as a wound parasite for the most part. Twigs and branches are also attacked, causing a typical die-back and finally stag-headed condition of infected trees. In this phase of the disease, also, the fungus is not virulently parasitic, attacking through wounds or infecting trees weakened by overshadowing or other unfavorable cultural conditions. Where the progress of the fungus is slow, cankers of various sizes form. The action of the fungus on other hosts is the same as on *Theobroma*, a brown rot or soft rot (stem-end rot of *Citrus* fruits) of the fruit and a die-back or cankering of twigs and branches. *D. natalensis* Evans, which causes a typical stem-end rot and die-back of *Citrus* spp. in the Union of South Africa, Porto Rico, Cuba, and southern Florida, is probably not distinct.

**Epicoccum theobromae** Petch. On leaves of *T. cacao* in Ceylon.

**Eutype erumpens** Mass. Causes a rot of bark and wood of branches and trunks of *T. cacao*, *Ficus* spp., *Mangifera indica*, and *Myristica* sp. in Trinidad and the British West Indies. The fungus forms black shining carbonaceous, raised stromatic patches on the rotted areas. Probably a wound parasite only.

**Exoascus bussei** v. Fab. This fungus causes a witches'-broom disease of *T. cacao* in Cameroon (Africa.) The brooms arise from an infected bud, which forms an abnormal fleshy branch with shortened internodes. From other buds at the base of this branch other abnormal branches arise, forming the broom. Such leaves as form are small and soon wither and die, remaining attached to the twigs. The brooms may persist for years, never flowering.

**Exoascus theobromae** Ritz. Bos. Assigned at one time as the cause of the Surinam witches'-broom disease. See *Marasmius perniciosus*.

**Fomes lamaoensis** Murr. See *Hevea*.

**Fomes lignosus** Klotzsch. See *Hevea*.

**Fomes pseudo-ferreus** Wakef. See *Hevea*.

**Gloeosporium theobromicolum** Vinc. Anthracnose on leaves and pods of *T. cacao* in Brazil.

**THEOBROMA**—Continued.

- Helminthosporium theobromae** Turc. On leaves of *T. cacao* in Italy.
- Macrophoma vestita** Prill. and Delacr. Said to attack the roots of *T. cacao* in tropical America.
- Marasmius** spp. The so-called "horse-hair" blights and thread blights occur on *T. cacao*, *Thea* (cult), *Hevea brasiliensis*, *Myristica fragrans*, and many other tropical shrubs and trees. This type of disease has been reported from Ceylon, west Africa, the West Indies, Trinidad, British Guiana, and Brazil, and doubtless occurs in all tropical regions. Among the species of fungi involved are *M. equicrinis* Mull., *M. pulcher* (B. and Br.) Petch, and *M. sarmentosus* Berk. Most of these fungi form thin (hairlike) polished black mycelial cords which grow over branches and leaves, adhering at certain points. They are only weakly parasitic, if at all, in contrast to fungi such as *Corticium koleroga* (see *Coffea*), with which class of parasites they are often confused in the literature.
- Marasmius perniciosus** Stahel. The "krulloten" or Surinam witches'-broom disease which has made the growing of *T. cacao* commercially almost impossible in Surinam is now attributed by Stahel to the above fungus, *Colletotrichum* and *Exoascus* having been previously assigned as causes by other workers. The disease has also been found in British Guiana. The disease is characterized by somewhat curved broomlike growths brought about by an excessive development of lateral shoots, together with a shortening of the internodes of infected twigs. Diseased shoots are thicker than normal and the surfaces ridged. The leaves are always small and soft and pliant, while the petioles are always much enlarged. The witches'-brooms die in from two to three weeks and fall from the trees. Very few pods are formed and these remain small and misshapen or become very hard and black. Continued attacks of the disease not only prevent production of healthy pods but eventually destroy the trees.
- Melanomma henriquesianum** Bres. and Roum. On bark of *T. cacao* in San Thomé and Kamerun.
- Meliola guianensis** Stev. and Dowell. Black fungus patches on brown leaf spots, 3 to 10 millimeters in diameter on *T. cacao* in British Guiana.
- Meliola theobromae** von Faber. Superficial black fruiting patches on leaves of *T. cacao* in Cameroon.
- Monilia** sp. The watery or *Monilia* pod disease of *T. cacao* occurs in Ecuador, having been found there also on the wild species, *T. bicolor* and *T. balaensis*. The disease is first noted on young pods as small protuberances or discolored areas varying from yellow to brown or even bluish. A white powder consisting of conidia of the fungus, often appears on the diseased areas. The interior of diseased pods is dark-brown, dark-brown or black strands and spots also occurring throughout the husk itself. A watery fluid forms in the interior in considerable abundance. Seeds in diseased pods are rendered worthless.
- Nectria** spp. A number of species of *Nectria*, including *N. bainii* Mass., *N. camerunensis* App. and Str., *N. jungeri* P. Henn., *N. theobromae* Mass., *N. ditissima* Tul., and *N. striatospora* Zimm. have been reported from the various cacao-growing countries as attacking branches, trunks and fruit of *T. cacao* as well as other tropical economic woody plants. There is a voluminous literature describing these fungi and the symptoms of the diseases resulting from their activities, but recent studies indicate that the Nectrias are for the most part saprophytic or at best weak parasites following other fungi, notably *Phytophthora faberi* Maubl. Some of the species may act as wound parasites.
- Phyllosticta theobromae** d'Alm. and S. da Cam. Large irregular leaf spots on *T. cacao* in Portuguese St. Thomas, Java, and Ceylon.
- Phyllosticta theobromicola** Vinc. Irregular brown areas on leaves of *T. cacao* in Brazil.
- Physalospora affinis** Sacc. On branches of *T. cacao* in the Philippines.
- Physalospora theobromae** Turc. On leaves of *T. cacao* in Italy.
- Phytophthora faberi** Maubl. This downy mildew fungus causes the most serious known disease of *T. cacao*. Other tropical economic plants, including *Artocarpus integrifolia*, *Carica papaya*, *Cocos nucifera*, and *Hevea brasiliensis* are also attacked. The fungus is reported from Cuba, Jamaica, Dominican Republic, Guadeloupe, Lesser Antilles, Trinidad, Surinam, Ceylon, Java, India, the Philippines, Malaya, Fiji, Siam, Samoa, New Guinea, Gold Coast, Uganda, Cameroon, and Belgian Congo. The pod-rot phase of the disease begins as brown spots at the end of the pods, which rapidly spread, soon involving all of the rind and seeds in a brown rot. Diseased pods may be covered with a white mycelium and may either fall to the ground or harden and hang on the tree for a long time. The disease is more serious in moist, shady situations. Seedlings are blighted and destroyed by the fungus. Cankers very similar to those described for *Hevea* occur on the trunks and branches and are often extensive enough to kill the trees. See also *Hevea*.
- Phytophthora meadii** McRae. See *Hevea*.
- Rosellinia bunodes** B. and Br. See Citrus.
- Rosellinia paraguayensis** Starb. Found in connection with root disease of *T. cacao* in the British West Indies.
- Rosellinia pepo** Pat. See Citrus.
- Ramularia necator** Mass. White mold appears on the cotyledons of seedlings of *T. cacao* in the British West Indies and west Africa. A damping-off results.
- Sphaeronema** sp. This fungus is reported as the cause of a black spot and bark rot disease of *T. cacao* in Ecuador. In the first phase of the disease small dark spots appear on nearly mature pods and increase in size up to 4 centimeters. Diseased pods are invaded by a dry rot and are lighter in weight than normal ones. The fungus gains entrance through wounds and causes large cankers, from which a dark wine-colored liquid exudes in damp weather. The diseased bark tissues become yellow or reddish, a dark line separating healthy tissue from diseased.
- Stachyldium theobromae** Turc. On leaves of *T. cacao* in Italy.
- Trachysphaera fructigena** Tabor and Bunting. See *Coffea*.
- THERMOPSIS.** Hardy perennial herbs.
- Phyllosticta thermopsidis** Thuem. Large irregular dull-brown leaf spots on *T. lanceolata* in Siberia.
- Uredo thermopsidis** Thuem. Leaf rust on *T. lanceolata* in Siberia.
- THESPESIA.** See Hibiscus.
- THEVETIA.** Small, glabrous trees and shrubs.
- Accidium thevetiae** Sacc. Leaf rust on *T. cuneifolia* in Mexico.
- THLASPI.** PENNYCRESS. FRENCH WEED. Annual or perennial herbs.
- Accidium thlaspianum** Syd. Leaf rust on *T. ceratocarpon* in Austria.
- Peronospora thlaspeos alpestris** Gäum. Downy mildew on leaves of *T. alpestre* in central Europe.
- Peronospora thlaspeos arvensis** Gäum. As above on *T. arvensis* in Europe.
- Peronospora thlaspeos perfoliati** Gäum. As above on *T. perfoliatum* in Yugoslavia, Austria, and Germany.
- Puccinia thlaspeos-glaucophylli** P. Henn. Brown rust pustules on leaves of *T. glaucophyllum* in Chile.
- Ustilago thlaspeos** (Beck.) Lagh. See *Arabis*.
- THRINAX.** THATCH PALM. See *Palmae*.
- THUJA.** ARBORVITAE. Timber trees, also grown as ornamentals.
- Caoma deformans** (B. and Br.) Tub. A rust causing witches'-brooms, 4 to 6 inches in diameter, on *T. japonica* and *Thujaopsis dolabrata* in Japan.
- Helicobasidium mōmpa** Tan. See *Morus*.

- THUJOPSIS.** FALSE ARBORVITAE. Ornamental coniferous trees and shrubs grown for their handsome foliage and formal habit.
- Caeoma deformans** (B. and Br.) Tub. See Thuja.
- THUNBERGIA.** CLOCK VINE. Acanthaceous, tall, perennial climbers, with a profusion of variously colored flowers.
- Puccinia thunbergiae** Cke. Brown leaf rust on *T. natalensis* in Japan.
- Puccinia thunbergiae-alatae** P. Henn. Powdery brown rust pustules on leaves of *T. alata* in tropical Africa.
- THYMUS.** THYME. Small shrubs and subshrubs, grown in borders and rockeries.
- Puccinia caulicola** Schneid. Dark-brown rust pustules on leaf blades, petioles, and stems of *T. angustifolium*, *T. chamaedrys*, *T. pannonicum*, *T. serpyllum*, and *Origanum vulgare* in Europe.
- THYRSACANTHUS.** Acanthaceous erect herbs.
- Uromyces hariotanus** Lagerh. Leaf rust on *T. strictus* in Costa Rica and Ecuador.
- TIARELLA.** FOAMFLOWER. Slender, hardy, perennial herbs.
- Puccinia asiatica** (Kom.) Syd. See Mitella.
- TIBOUCHINA.** GLORY BUSH. Melastomaceous shrubs, some species climbers.
- Dothidina peribebuyensis** (Speg.) Chardon. See Miconia.
- Phyalospora tibouchinae** P. Henn. On *Tibouchina* sp. in Brazil.
- TILIA.** LINDEN. Sometimes called basswood and lime. Timber and ornamental trees.
- Actinonema tiliae** Allesch. Causes premature leaf fall and swelling of buds in autumn of *Tilia* sp. in Belgium, Austria, and Germany.
- Ascochyta tiliae** Kab. and Bub. Subcircular to irregular, often confluent, yellow-gray, then ashen, leaf spots with purple-brown margins on *T. dasystyla* in Bohemia.
- Ectostroma tiliae** Fr. On leaves of *Tilia* sp. in Sweden.
- Gloeosporium tiliae** Oud. and var. **maculicolum** Allesch. On leaves of *T. cordata*, *T. platyphyllos*, and *T. cordata* (*T. ulmifolia*) in Europe.
- Gloeosporium tilaeolum** Allesch. Irregular yellow to yellow-brown patches on leaf blades, petioles, and young twigs of *T. platyphyllos*, *T. parvifolia*, and *T. cordata* (*T. ulmifolia*) in Great Britain and Germany.
- Gloeosporium vogelii** Syd. On leaves of *T. cordata* (*T. ulmifolia*) in Germany.
- Helminthosporium tiliae** Fr. On leaves of *T. platyphyllos* and *T. cordata* in France.
- Leptosphaeria vagabunda** Sacc. On branches of *Tilia* sp. in Europe.
- Phyllosticta bacterioides** Vuill. Dark-brown leaf spots on *T. cordata* in France.
- Phyllosticta bracteophila** Ferr. Small ashen-white leaf spots on *T. platyphyllos* in Italy.
- Phyllosticta praetervisa** Bub. Numerous small angular leaf spots on *T. cordata* (*T. parvifolia*) in Bohemia.
- Phyllosticta tillicola** Oud. Small angular brown leaf spots on *T. cordata* (*T. ulmifolia*) in Holland.
- Phyllosticta vogelii** (Syd.) Died. On leaves of *T. platyphyllos* in Poland.
- Phytophthora syringae** Kleb. See Syringa.
- Placosphaeria tiliae** Bub. Circular to elliptical brown, then yellowish, leaf spots on *T. cordata* in Hungary.
- Pucciniastrum tiliae** Hirats. Yellow to brown rust pustules on leaves of *T. cordata*, *T. manshurica*, and *T. miqueliana* in Japan, China, Manchuria, and possibly Russia.
- Pyrenochaeta filarszkyi** Bub. Circular to irregular yellow leaf spots on *T. cordata* in Hungary.
- Pyrenochaete pubescens** E. Rostr. Circular to oblong purplish, then ashen, areas on branches of *Tilia* sp. in Denmark.
- Septoria tiliae** West. On leaves of *T. europaea* and *T. cordata* (*T. ulmifolia*) in Europe.
- Uncinula miyabei** Salm. Powdery mildew on leaves of *T. miqueliana* in Japan.
- TILLANDSIA.** SPANISH MOSS. Epiphytic bromeliads.
- Phyllosticta tillandsiae** Speg. On leaves of *Tillandsia* sp. in Argentina.
- Ustilago tillandsiae** Patters. Powdery black smut sori destroying the inflorescences of *T. leiboldiana* and *Tillandsia* sp. in Mexico and Costa Rica.
- TINNEA.** Tall perennial herbs and subshrubs.
- Aecidium tinneae** P. Henn. Yellow leaf rust on *T. ericalyx* in the Union of South Africa.
- TIPUANA.** TIPU TREE. Ornamental leguminous trees.
- Phyllosticta tipuanae** Tassi. On leaves of *T. tipu* (*T. speciosa*) in Argentina.
- TITEONIA.** Tropical composite herbs.
- Puccinia tithoniae** Diet. and Holw. Brown to black rust pustules on leaves of *T. rotundifolia*, *T. scaberrima*, and *T. tubaeformis* in Mexico, Costa Rica, and Guatemala.
- TODDALIA.** Ornamental subtropical shrubs.
- Aecidium toddaliae** Petch. Yellow leaf rust on *T. aculeata* in Ceylon.
- Bacterium citri** Hasse. See Citrus.
- Puccinia kentaniensis** Pole-Evans. Rust on *T. lanceolata* in the Union of South Africa.
- Puccinia teclae** Pass. Black leaf rust on *T. nobilis* in Abyssinia.
- Puccinia toddaleae** Racib. Leaf rust on *T. aculeata* in Java.
- TOFIELDIA.** Perennial rhizomatous herbs.
- Phyllosticta tofieldiae** Rostr. Leaf spot on *T. borealis* in Europe.
- TOLPIS.** Composite herbs with showy yellow flower heads.
- Puccinia heribaudiana** Har. Leaf rust on *T. barbata* in France.
- TORENIA.** Annual or perennial herbs, sometimes grown in gardens.
- Cercospora toreniae** P. Henn. On leaves of *Torenia* sp. in Brazil.
- Puccinia toreniae** Rac. Brown leaf rust on *T. asiatica* in Java.
- TORRESIA.** See Hierochloa.
- TOXYLON.** See Maclura.
- TRADESCANTIA.** SPIDERWORT. Perennial hardy herbs.
- Kordyana tradescantiae** (Pat.) Rac. On leaves of *T. virginiana* and *Tradescantia* sp. in Java and Ecuador.
- Uromyces commelinae** (Speg.) Cke. See Commelina.
- TRAGOPOGON.** VEGETABLE OYSTER. Erect biennial or perennial herbs with narrow grasslike leaves and heads of yellow or purple flowers.
- Ascochyta tragopogonis** Boud. On leaves and stems of *T. major* in Russia.
- Entyloma tragopogi** Lagerh. Smut sori in elongate yellow, then brown, leaf spots on *Tragopogon* sp. in France.
- Puccinia tragopogi** (Pers.) Cda. Yellow and powdery dark-brown rust pustules on leaves and stems of *T. coloratus*, *T. dubius*, *T. floccosus*, *T. major*, *T. orientalis*, *T. palastinus*, *T. porrifolius*, and *T. pratisensis* in Asia Minor, Algeria, and Europe. Reported from Idaho.
- Ustilago tragopogonis-pratisensis** (Pers.) Wint. Smut sori distorting flower heads of *T. major*, *T. orientalis*, and *T. parvifolius* in Argentina and Europe.

- TRAPA.** WATER CHESTNUT. Floating aquarium plants.  
*Septoria trapae-natantis* Wisn. Subcircular leaf spots with dark-purple margins on *Trapa* sp. in Galicia.
- TREMANDRA.** Australian shrubs.  
*Puccinia tremandrae* B. and Br. Brown leaf rust on *T. oppositifolia* and *T. stelligera* in Australia.
- TRICALYSIA.** Erect or climbing shrubs.  
*Hemileia evansii* Syd. Leaf rust on *Tricalysia* sp. in the Union of South Africa.  
*Phyllosticta tricalysiae* A. L. Sm. Large gray leaf spots with red-brown margins on *T. griseiflora* in Angola.
- TRICHILIA.** Ornamental trees and shrubs.  
*Campotrichum cladosporioides* Sacc. Large subcircular leaf spots with dark-olive margins on *T. emetica* in tropical Africa.  
*Coccinia concentrica* Syd. Circular crustaceous black stromata on leaves of *T. emetica* in Portuguese East Africa.  
*Mycosphaerella asunciensis* Starb. On leaves of *Trichilia* sp. in Paraguay.  
*Phyllachora explanata* (Lév.) Sacc. Circular black stromata on leaves of *T. havanensis* in Cuba.  
*Phyllosticta pertundens* Sacc. Subcircular dull-yellow leaf spots on *T. emetica* in Africa.  
*Uredo trichiliae* Arth. Leaf rust on *T. pallida* and *T. trinitensis* in Porto Rico and Trinidad.
- TRICHLORIS.** Perennial grasses.  
*Puccinia trichloridis* Speg. Linear black rust sori on leaves of *T. mendocina* in Argentina.
- TRICHOSANTHES.** SNAKE GOURD. Annual or perennial climbing herbs.  
*Irene confragosa* Syd. Superficial black patches on leaves of *T. quincangularis* in the Philippines.  
*Macrophoma trichosanthis* Syd. Causes a fruit rot of *T. anguina* in the Philippines.  
*Uredo trichosanthis* Petch. Leaf rust on *T. palmata* in Ceylon.
- TRICHOSTEMA.** BLUECURLS. Herbs, sometimes cultivated in gardens or rockeries.  
*Cercospora trichostemmatis* P. Henn. Pale-brown irregular leaf spots on *T. volkensis* in Tanganyika.
- TRICHOSTIGMA.** Ornamental climbing shrubs.  
*Cercospora trichostigmae* Stevens. Leaf spot on *T. octandra* in Porto Rico.  
*Endophyllum rivinae* (B. and C.) Arth. See Rivina.  
*Puccinia rivinae* (B. and C.) Speg. See Rivina.  
*Septoria rivinae* Pat. See Rivina.
- TRIDAX.** Hardy perennial composite herbs.  
*Puccinia melampodii* Diet. and Holw. See Zinnia.
- TRIENTALIS.** STAR FLOWER. Small, glabrous perennials.  
*Accidium trientalis* Tranzsch. Leaf rust on *T. europaea* in Russia.  
*Puccinia karelica* Tranzsch. Leaf rust on *T. europaea* and *Carex limosa* in Russia. Reported from New York.  
*Ramularia magnusiana* (Sacc.) Lind. On leaves of *T. europaea* in Denmark, Austria, and Germany.  
*Septoria trientalis* (Lasch.) Sacc. Small circular gray-white leaf spots on *T. europaea* and *Trientalis* sp. in Russia, Sweden, Denmark, and Germany.  
*Sorosporium trientalis* Woron. Smut on *T. europaea* in Finland.  
*Tubercinia trientalis* (B. and Br.) Wor. Thickened leaf spots and crustlike growths on stems of *T. europaea* in Japan and northern Europe.
- TRIFOLIUM.** CLOVER. Herbaceous perennials grown for lawns and as forage crops.  
*Ascochyta trifolii* A. Bond and Trus. On *T. pratense* in Russia.  
*Ascochyta volkartii* Bub. On *T. repens* in Bulgaria.  
*Bacillus trifolii* Vogl. Attacks stems, leaves, and peduncles of *T. repens* in Italy.  
*Bacterium* sp. A leaf curl and wilt of stems of *T. pratense* in Italy is thought to be due to a *Bacterium*.  
*Botrytis anthophila* A. Bond. Gray mold destroying flowers and preventing seed production of *T. pratense* in Russia.  
*Cercospora stolziana* Magn. Yellow swollen leaf spots on *T. repens* in Bohemia and Austria.  
*Hypochnus cucumeris* Fr. See Cucumis.  
*Leptosphaeria circinans* (Fckl.) Sacc. See Asparagus.  
*Mycosphaerella carinthiaca* Jaap. (*Ramularia trifolii* Jaap.). Leaf spots, brown above, gray-green below, on *T. medium* in Great Britain, Switzerland, Austria, and Germany.  
*Ovularia exigua* (W. Sm.) Sacc. On leaves of *Trifolium* sp. in Great Britain.  
*Peronospora pratensis* Syd. Downy mildew on leaves of *T. incarnatum*, *T. medium*, and *T. pratense* in Europe.  
*Peronospora trifolii alpestris* Gäum. Downy mildew on leaves of *T. alpestre* in Switzerland, Denmark, and Germany.  
*Peronospora trifolii arvensis* Syd. As above, on *T. arvensis* in Switzerland, Russia, Denmark, and Germany.  
*Peronospora trifolii hybridi* Gäum. Downy mildew on leaves of *T. hybridum*, *T. spadiceum*, and *T. striatum* in Europe.  
*Peronospora trifolii minoris* Gäum. As above on *T. agrarium*, *T. badium*, *T. minus*, *T. patens* and *T. procumbens* in Europe.  
*Peronospora trifolii repentis* Syd. Downy mildew on leaves of *T. monatum*, *T. repens*, and *T. rubens* in central and northern Europe.  
*Phyllosticta trifolii* Rich. On leaves of *T. pratense* and *T. repens* in Sweden and Russia.  
*Pseudovularia trifolii* Speg. Indefinite subcircular brown spots with subshen centers on *T. platensis* in Argentina.  
*Rhabdospora alexandrina* Chrest and Maire. Irregular black spots on stems and circular brown leaf spots on *T. alexandrinum* in Egypt.  
*Septoria trifolii* Cav. On leaves of *T. repens* in Italy.  
*Sphaerulina trifolii* E. Rostr. Circular brown leaf spots with pale-purple surrounding zones on *T. pratense* and *T. repens* in Denmark and Germany.  
*Tylenchus dipsaci* Kuehn. See Narcissus.  
*Typhula trifolii* Rostr. Forms small black sclerotia in rotted areas in stems and leaves of *T. pratense*, *T. repens*, *Anthyllis vulneraria*, and *Medicago lupulina* in northern Europe. The parasitism of this fungus is uncertain.  
*Uromyces flectens* Lagh. Brown powdery rust pustules on leaf blades and petioles of *T. repens* in Japan and Europe. Also reported from Wisconsin.  
*Urophlyctis trifolii* (Pass.) Magn. Pustules on leaf blades and petioles of *T. montanum*, *T. pratense*, and *T. repens* in Great Britain, Denmark, Italy, and Germany.

**TRIGONELLA.** FENUGREEK. Annual or perennial herbs.

*Cercospora traversiana* Sacc. Subcircular brown leaf spots on *T. foenum-graecum* in Italy.

*Cercospora trigonellae* A. Maubl. On leaves of *T. foenum-graecum* in Brazil.

*Peronospora trigonellae* Gäum. Downy mildew on leaves of *T. foenum-graecum* and *T. polycerata* in Algeria, Russia, and India.

*Uromyces anthyllidis* (Grev.) Schroet. See Anthyllis.

*Uromyces trigonellae* Pass. Leaf rust on *T. foenum-graecum* and *T. occulta* in Egypt, India, Italy and France. Said to be synonymous with the above species.

**TRIPETALEIA.** Shrubs cultivated for their flowers.

*Uredo tripetaleiae* Diet. Leaf rust on *T. bracteata* in Japan.

**TRIPSACUM.** GAMA GRASS. Perennial grasses with stout culms, used as fodder plants.

*Puccinia pallescens* Arth. Brown powdery rust pustules on leaves of *T. dactyloides*, *T. latifolium*, *T. lanceolatum*, and *Zea mays* in Mexico, Porto Rico, Guatemala, Nicaragua, Salvador, and Trinidad.

*Ustilago dieteliana* P. Henn. Dark, olivaceous smut sori in panicles of *T. dactyloides* in Mexico.

**TRISETUM.** Hardy perennial grasses.

*Puccinia triseti* Erikss. Yellow-brown to black rust pustules on leaves of *T. barcinonensis*, *T. deyeuxioides*, *T. flavescens*, *T. neglectum*, and *T. ovatum* in Guatemala and Europe. Also reported from Utah.

*Septoria caballeroi* Gz. Frag. On *T. ovatum* and *T. paniceum* in Spain.

*Septoria nebulosum* Rostr. See Calamagrostis.

*Septoria triseti* Speg. See Agrostis.

**TRISTANIA.** Evergreen trees or shrubs.

*Ascochyta brunnea* Cke. and Mass. Ochraceous to dull-brown leaf spots on *T. conferta* in Australia.

*Helminthosporium puccinioides* Sacc. and Berl. On leaves of *T. laurina* in Australia.

*Mycosphaerella tristaniae* Wakef. On leaves of *T. griffithii* in the Federated Malay States.

*Phyllosticta tristaniae* D. Sacc. Subcircular gray-white leaf spots with red margins on *T. conferta* in Italy.

*Septoria hanburyana* Tassi. On leaves of *T. nereifolia* in Italy.

*Septoria tristaniae* P. Henn. Brown leaf spots on *T. laurina* in Germany.

**TRITELEIA.** Bulbous plants.

*Uromyces reichei* Diet. Brown rust pustules on leaves of *T. gaudichaudiana* in Chile.

*Uromyces triteleiae* Diet. and Neg. Rust on leaves and scapes of *T. porrifolia* in Chile.

**TRITHINAX.** South America fan palms. See Palmae.

**TRITICUM.** WHEAT.

*Acremoniella occulta* Cav. See Secale.

*Acremoniella verrucosa* Togn. See Avena.

*Bacillus cerealinum* Gentner. See Hordeum.

*Dilophospora graminis* Desm. (*Mastigosporium album* Reiss.) (*Dilophia graminis* Fekl.) Small light-colored circular to oblong spots appear on both leaf surfaces, becoming larger and dark in color to form blotches and stripes. Infected leaves soon assume a light-yellow color and die. Leaf sheaths are also attacked, particularly the upper ones, and generally before heading out, so that the growing point is firmly inclosed by the young leaves bound together by a mass of mycelium. The continued growth of the tip causes characteristic distortion or buckling of the stems. When infection occurs after heading out deformation of the spikes may be partial or complete, affected areas covered at first with a dirty-white fungus growth which later becomes black and gives a charred appearance. This disease occurs in connection with the nematode *Tylenchus tritici*. Many grasses are attacked, including *Triticum* spp., *Agrostis alba*, *A. stolonifera*, *A. vulgaris*, *Alopecurus agrestis*, *A. pratensis*, *Arrhenatherum elatius*, *Anthoxanthum odoratum*, *Calamagrostis arundinacea*, *C. canadensis*, *C. lanceolata*, *C. epigeios*, *Dactylis glomerata*, *Festuca ovina*, *F. pratensis*, *F. rubra*, *Hordeum lanatum*, *H. mollis*, *Phleum pratense*, *Poa trivialis*, and *Secale cereale*. in England, Holland, France, Belgium, Switzerland, Norway, Denmark, and Germany. Reported on *Calamagrostis canadensis* from Wisconsin.

*Fusarium nivale* (Fr.) Sor. See Secale.

*Gibellina cerealis* Pass. Round or elongate black patches on leaf sheaths of *Triticum* in Italy and Hungary. Infected plants turn yellow, wilt, and finally die.

*Helminthosporium tritici* P. Henn. Brown areas on leaves, culms, and spikes of *Triticum* in Tanganyika.

*Leptosphaeria culmicola* (Fr.) Karst. See Bromus.

*Leptosphaeria herpotrichoides* De N. Brown spots appear on the leaf sheaths at the base of infected plants, enlarging so as to encircle the culms and penetrating to the stem. Mycelial mats form between and bind the leaf sheaths together. Diseased plants fail to produce grain and infected stems lodge. The hosts are *Triticum*, *Andropogon* sp., and *Koeleria cristata* in Argentina, France, Italy, Finland, Sweden, Denmark, and Germany. One of the foot-rot or take-all group of diseases. See Ophiobolus.

*Macrophoma hennebergii* (Kuehn) Berl. and Vogl. Brown spots on glumes and leaves of *Triticum* in Italy, Russia, Sweden, and Germany. The disease causes the grain to shrivel up.

*Micrococcus tritici* Prill. Grains develop poorly, shrivel, and turn red. On *Triticum* (cult.) and *Hordeum vulgare* in Brazil, Argentina, Italy, France, and Denmark.

*Mycosphaerella exitialis* Mor. See Hordeum.

*Mycosphaerella hordicola* Hara. See Hordeum.

*Mystrosporium abrodens* Neum. Dark patches on the lower nodes and leaves of *Triticum* in France. Diseased plants fail to form grain.

*Nigrospora panici* Zimm. Black fungus patches on stems and leaf sheaths of *Triticum*, *Oryza sativa*, and *Zea mays* in Java.

*Ophiobolus cariceti* (B. and Br.) Sacc. (*O. graminis* Sacc.) The take-all, pietin, or foot-rot disease of cereals has been the cause of heavy losses for many years. A number of fungi have been found associated with the disease, but *O. cariceti* is apparently the most common and most virulent species present. The hosts are *Triticum*, *Secale cereale*, *Agropyron* sp., *Avena sativa*, *Agrostis palustris*, *Hordeum sativum*, *H. murinum*, *Oryza sativa*, *Bromus mollis*, *B. sterilis*, *Elymus*, *Festuca*, *Lolium*, *Hystrix*, and *Phalaris* in Australia, Japan, Argentina(?), Great Britain, Sweden, Holland, France, Belgium, Italy, and Germany. The disease has also been found recently in a number of the States.

The symptoms of the disease are very characteristic. There is a browning or blackening of the base of the stems with the development of an enveloping brown mycelial mat in which the perithecia develop. The roots turn black and rot off. Diseased plants turn yellow and finally die, always failing to set grain. The disease kills out irregular patches in the field which may become very extensive.

*Ophiobolus herpotrichus* (Fr.) Sacc. This species attacks the stems and leaves at or near the base of the plant, causing a disease similar to the take-all described above. It attacks *Triticum*, *Hordeum* sp., *Cynodon* sp., and *Agrostis* sp. in Italy, France, Denmark, Belgium, Holland, Hungary, and Germany.

**TRITICUM**—Continued.

- Phoma hennebergii** Kuehn. Large irregular gray-brown spots on glumes of *Triticum* in Sweden.
- Phyllachora tritici-gracilis** (Cast.) Sacc. Black stromata on leaves of *T. gracile* in France.
- Pseudomonas tritici** Hutch. The inflorescences and stems are covered with a bright primrose-yellow slime which forms sticky layers between the glumes and between the stems and sheaths. Such heads as form are distorted. The disease is readily carried on seed grain. On *Triticum* in India.
- Puccinia brachypus** Speg. See Bromus.
- Puccinia megalopotamica** Speg. Powdery brown rust pustules on leaves and sheaths of *Triticum* in Argentina.
- Pyroctonum sphaericum** Prunet. Attacks all parts of *Triticum* plants, causing a yellowing and shriveling of the leaves, abortion of the grain, and dwarfing of the plants in France.
- Rhizoctonia napi** West. See Brassica.
- Sclerospora macrospora** Sacc. This downy mildew gives infected plants a scalded appearance, the leaves dry up, and the heads abort or are deformed. A white mold growth appears on lower leaf surfaces. In addition to *Triticum*, *Alopecurus* sp., *Avena sativa*, *Festuca elatior*, *Lolium temulentum*, *Oryza sativa*, *Phragmites communis*, *Phalaris canariensis*, and *Zea mays* are attacked in Australia, Japan, Abyssinia, Russia, France, Spain, and Italy. The disease is present in the United States in a few localities.
- Septoria agrestis** Sacc. On leaves of *Triticum* in Russia and France.
- Septoria cristati** Hóll. Leaf spots on *T. (Agropyron) cristatum* in Hungary.
- Septoria neglecta** Sacc. On leaves of *Triticum* in France.
- Septoria nymanniana** Sacc. Elongate ochraceous areas on *Triticum* in Malta.
- Sphaeroderma damnosum** Sacc. The fungus develops a white mycelium beneath the sheaths and stems at the base of infected plants, causing a yellowing and dwarfing of the plants and abortion of the spikes. On *Triticum* and *Secale* in Italy. Considered by some workers as identical with *Fusarium culmorum*.
- Tilletia controversa** Kuehn. Black powdery, fetid masses of smut spores in ovaries of *T. vulgare*, *T. (Agropyron) repens*, *T. glaucum*, and *Elymus aralensis* in Turkestan, Russia, Denmark, Bohemia, Italy, Albania, Austria, and Germany.
- Tylenchus tritici** Bast. This nematode attacks young wheat (*Triticum*) plants, causing a wrinkling and distortion of the upper leaves. Such plants may die or remain alive and produce dwarfed, diseased heads. In place of normal grain dark hard galls form, which are shorter and thicker than wheat grains and filled with the parasites. The nematode affects *Triticum* primarily, but by inoculation has been found capable of attacking spelt, emmer, oats, and rye. The disease is known from Brazil, India, China, Turkestan, Australia, and Europe. In the United States it is reported from Virginia, Maryland, West Virginia, Georgia, and California.
- Typhula graminum** Karst. See Lolium.
- Urocystis tritici** Koern. The flag smut of wheat (*Triticum*) is characterized by long gray sori on the leaves and sometimes on the sheaths and stems, which finally rupture, exposing black powdery spore masses. Diseased leaves and stems become twisted, assume a drooping habit, and finally wither. If grain is formed at all, it shrivels up and is valueless. Diseased plants are much stunted. The disease occurs in India, Japan, China, Australia, Spain, Italy, and the Union of South Africa. Within recent years it has been found in Illinois, Missouri, and Kansas in limited areas.
- Ustilago trebouxii** Syd. See Melica.
- TRITONIA.** Cormous or bulbous plants.
- Septoria tritoniae** Pat. Elongate ashen-colored spots on stems of *Tritonia* sp. in Ecuador.
- Uromyces bonae-spei** Bub. Yellow to chestnut-brown rust pustules on both leaf surfaces of *T. scillaris* and *Acidanthera pallida* in the Union of South Africa.
- Uromyces transversalis** Thuem. See Gladiolus.
- TROCHODENDRON.** Evergreen trees with aromatic bark and foliage.
- Phyllosticta trochodendri** Alm. and Cam. Leaf spot of *T. aralioides* in Portugal.
- TROLLIUS.** GLOBEFLOWER. Hardy herbaceous perennials.
- Accidium trollii** Blytt. Leaf rust on *T. europaeus* in Russia.
- Ascochyta trollii** Thuem. Irregular gray-white leaf spots with dark-brown margins on *T. asiaticus* in Siberia.
- Didymaria trollii** Jacz. On leaves of *T. europaeus* in Russia.
- Metasphaeria trollii** Karst. On stems of *T. europaeus* in Finland.
- Phyllosticta trollii** Trail. Irregular brown leaf spots on *T. europaeus* in Great Britain, Italy, and Spain.
- Puccinia dietrichiana** Tranz. Leaf rust on *T. europaeus* in Russia.
- Puccinia trollii** Karst. Brown rust pustules in irregular sunken spots on leaf blades and petioles of *T. europaeus* in Italy, Switzerland, France, Belgium, Norway, Lapland, and Germany.
- Ramularia trollii** (Jacz.) Lind. On leaves of *T. europaeus* in Spain, Switzerland, Russia, and Finland.
- Septoria trollii** Sacc. and Wint. Ochraceous leaf spots on *T. europaeus* in Yugoslavia, Switzerland, Spain, and the Union of South Africa.
- TROPAEOLUM.** NASTURTIUM. Climbing or erect herbs, cultivated in gardens.
- Coleosporium tropaeoli** Palm. Rust on leaves of *T. peregrinum* in Germany.
- Cronartium flaccidum** (Alb. and Schw.) Wint. See Paeonia.
- Phyllosticta tropaeoli** Sacc. and Speg. On leaves of *T. major* in Italy, France, Portugal, and Austria.
- Uredo tropaeolum** Desm. Bright-orange rust pustules on lower leaf surfaces of *Tropaeolum* sp. in Europe.
- Uromyces tropaeoli** Ranaj. Leaf rust on *T. major* in Yugoslavia.
- TSUGA.** HEMLOCK. Coniferous timber and ornamental trees.
- Ascochyta piniperda** Lindau. See Picea.
- Phomopsis pseudotsugae** Wils. See Pseudotsuga.
- TULIPA.** TULIP. Many-colored spring-flowering hardy bulbs.
- Accidium tulipae** Komarov. Yellow rust pustules on the leaves of *T. turkestanica* in central Asia. Possibly one stage of *Uromyces tulipae* Diet.
- Helminthosporium tenuissimum** Nees. Dead spots on scapes of *T. heteropetala* in Siberia.
- Puccinia prostii** Moug. This rust produces brown oblong convex rust pustules on both surfaces of the leaves of *T. celsiana* and *T. silvestris*, including many cultivated varieties in Italy, Great Britain, and France. The disease checks flowering or may even prevent it entirely.
- Puccinia tulipae** Schroet. Brown leaf rust on *T. gesneriana* and *T. suaveolens* in Austria and Germany.
- Rhizoctonia tuliparum** (Kleb.) Whet. and Arth. (*Sclerotium tuliparum* Kleb.) This disease has been very destructive at times to tulip plantings in Holland and has also been reported from England, Switzerland, and Germany. There are several reports of occurrence in America. Infected plants rot at the ground level, wither, and die. Diseased bulbs show a dry grayish or reddish-gray rot internally, with abundant grayish-white mycelium externally and between the diseased scales. Numerous more or less globose, dark-brown to black sclerotia occur on the rotted scales and neck of the bulb and occasionally between the scales.

**TULIPA**—Continued.

**Sclerotinia tuliparum** (Wakk.) Rehm. This fungus causes the so-called black rot of hyacinth bulbs in Italy, Great Britain, Holland, France, and Germany. Bulbs of *Crocus*, *Scilla* spp., and *Tulipa* spp. are also attacked.

**Sclerotium** spp. (*Botrytis* spp.) Several *Sclerotium* and *Botrytis* diseases on tulips and other bulbs have been listed and studied by various workers. It is far from clear how many distinct diseases are involved or which ones have reached the United States. *Botrytis tulipae* (Lib.) Hopk., the so-called *Botrytis* blight of tulips, is widespread in this country, having been introduced times without number on Dutch bulbs. See *Rhizoctonia*, *Sclerotinia*.

**Synchytrium laetum** Schroet. See Gagea.

**Urocystis colchici** (Schlecht.) Rab. See Colchicum.

**Uromyces tulipae** Diet. A rust known only from Japan causing yellow to brown rust pustules on both surfaces of leaves of *T. edulis*.

**Ustilago tulipae** Wint. Elliptical convex pustules appear on the leaves and split longitudinally exposing the black dusty spore masses. *Tulipa* spp., including *T. sylvestris* in France, Austria, and Germany, are the hosts. *Ustilago heufleri* Fekl., which is considered synonymous by some workers with the above, occurs in America on *Erythronium americanum*.

**TUMBOA**. See Welwitschia.**TUNICA**. TUNIC FLOWER. Annual or perennial hardy herbs.

**Septoria dianthicola** Sacc. See Dianthus.

**Urocystis purpurea** Hazl. See Dianthus.

**TURBAEA**. Ornamental trees and shrubs.

**Acidium ugandense** Syd. Rust pustules in circular to irregular yellow to yellow-brown leaf spots on *Turraea* sp. in British tropical Africa.

**Pseudographis volkensii** P. Henn. On branches of *T. volkensii* in tropical Africa.

**TUSSILAGO**. COLTSFOOT. Acaulescent perennial herbs with solitary yellow flower heads.

**Ascochyta tussilaginis** Oud. On leaves of *T. farfara* in Holland.

**Coleosporium tussilaginis** (Pers.) Lév. See Pinus.

**Phyllosticta caballeri** Gz. Frag. On leaves of *T. farfara* in Spain.

**Phyllosticta farfarae** Sacc. and var. **major** P. Brun. Irregular gray-white leaf spots on *T. farfara* in Italy, France, and Denmark.

**Ramularia purpurascens** Wint. Grayish-white leaf spots with indefinite purplish surrounding zones on *T. fragrans* and *Nardosmia fragrans* in Portugal.

**Septoria farfarae** Pass. Gray-brown leaf spots on *T. farfara* in Italy.

**Septoria tussilaginis** West. Irregular brown to olivaceous, finally gray-white, leaf spots on *T. farfara* and *T. fragrans* in Italy, Belgium, and Germany.

**Stagonospora tussilaginis** (Fekl.) Died. On leaves of *T. farfara* in Italy, Denmark, Austria, and Germany.

**TYPHA**. CAT-TAIL. Hardy perennial marsh herbs.

**Hainesia corallina** Sacc. and Fautr. On leaves of *T. latifolia* in France.

**Heterosporium typharum** Cke. and Mass. On leaves of *T. angustifolia* in Great Britain.

**Ophiobolus typhae** Feltg. On leaves of *T. latifolia* in Luxemburg.

**Phyllosticta coralliobola** Bub. and Kab. Irregular elongate ochraceous to brown leaf spots on *T. angustifolia* in Bohemia.

**Phyllosticta typha** (Pass.) Oud. On leaves of *T. angustifolia* in Holland.

**Phyllosticta typhina** Sacc. and Malbr. Oblong reddish leaf spots on *T. japonica* and *T. latifolia* in Japan and Europe.

**Septoria filispora** (Cke.) Sacc. On leaves of *T. latifolia* in France.

**Septoria menispora** B. and Br. On leaves of *T. latifolia* in Belgium and Great Britain.

**Ustilago grandis** Fr. See Phragmites.

**ULMARIA**. See Spiraea.**ULMUS**. ELM. Timber and ornamental trees.

**Acremoniella pallida** Cke. and Mass. On *Ulmus* sp. in Europe.

**Actinonema ulmi** Allesch. Irregular yellow, then brown, leaf spots on *U. campestris* in Austria.

**Ascochyta ulmella** Sacc. Pale-brown leaf spots on *U. campestris* in Italy.

**Asteroma fuckelii** Sacc. On leaves of *Ulmus* sp. in Europe.

**Asteroma ulmi** Klotzsch. On leaves of *U. campestris*, *U. parvifolia*, and *U. glabra*, (*U. scabra*) in Japan, Great Britain, France and, Portugal.

**Cercospora ulmicola** v. Hoeh. Subcircular ashen-brown leaf spots with dark margins on *Ulmus* sp. in Austria.

**Cucurbitaria naucosa** Fr. Causes swellings on the branches of *U. glabra* in Switzerland, followed by withering of the leaves and death of infected branches.

**Exoascus campester** Sacc. Ochraceous leaf spots on *U. campestris* and *U. glabra* (*U. montana*) in France.

**Exosporium ulmi** Eriks. Found in connection with die-back of twigs of *U. campestris*, *U. pedunculata* (*U. effusa*), and *U. glabra* in Sweden.

**Laestadia comedens** (Pass.) Sacc. Subcircular to irregular leaf spots on *U. campestris* in Italy.

**Leptosphaeria ulmicola** Mass. Irregular brown leaf spots on *U. campestris* in Italy.

**Mycosphaerella insularis** Walbr. On leaves of *Ulmus* sp. in Germany.

**Mycosphaerella oedema** (Fr.) Fekl. On leaves of *U. campestris* and *U. effusa* in central Europe.

**Phyllosticta bellunensis** Mart. Large irregular brown leaf spots on *U. campestris* (*U. elliptica*) and *U. glabra* in Russia, Poland, and Italy.

**Phyllosticta lacerans** Pass. Gray confluent leaf spots, the centers of which tear and drop out, on *U. campestris* in Russia and Italy.

**Phyllosticta ulmi** West. Subcircular, then confluent, ashen leaf spots on *U. americana*, *U. montana* and *U. fulva* in Italy, Bulgaria, and Belgium.

**Placosphaeria ulmi** P. Henn. Circular to angular yellow leaf spots on *U. parvifolia* in Japan.

**Septoria ulmicolum** (Biv. Bern.) Elenk. and Ohl. On *Ulmus* sp. in Russia.

**Sphaeropsis nervisequa** Lang. Causes a die-back of twigs of *U. glabra* in Germany. The leaves are first attacked, the fungus working down through the petioles.

**Sporidesmium ulmi** Fekl. On leaves of *U. pedunculata* in Germany.

**Stagonospora ulmifolia** (Pass.) Sacc. Small leaf spots on *U. campestris* in Italy.

**Systemma ulmi** (Schleich.) Theiss. and Syd. (*Phyllachora ulmi* [Duv.] Fekl.) Subcircular scattered, raised black stromata on leaves of *U. campestris*, *U. glabra*, *U. pedunculata*, and *U. suberosa* in Algeria and Europe. Often confused with a similar-appearing American elm-leaf disease caused by *Gnomonia ulmea* (Schw.) Thuem.

**Taphrina ulmi** (Fekl.) Johans. Pellucid, blisterlike spots on leaves of *U. alata*, *U. campestris*, and *U. montana* in Europe.

**Teratosperma singulare** Syd. Black fruiting pustules on leaves of *U. parvifolia* in Japan.

**ULMUS**—Continued.

**Uncinula clandestina** (Biv. Bern.) Schroet. Powdery mildew on leaves of *U. campestris*, and *U. montana* in Japan, Algeria and, Europe.

A serious disease of elms (*U. americana*, *U. campestris*, *U. pedunculata*, *U. monumentalis* and *U. rupellii*), characterized by a dying back of the twigs and branches, occurs in Holland and adjacent countries. Young trees frequently die while older trees are permanently affected, the leaves remaining small and falling prematurely. A number of fungi, including *Graphium ulmi* Schwartz have been found in connection with the disease, but their parasitic relationships have not yet been worked out. Brussoff attributes the disease to bacteria and considers the fungi secondary.

**UMBILICUS**. Herbs related to *Cotyledon*.

**Accidium umbilici** Trott. Leaf rust on *U. erectus* and *U. pendulinus* in Portugal and Spain.

**Phyllosticta umbilici** Brun. Indefinite brown leaf spots on *U. erectus* and *U. pendulinus* in Spain, Portugal, and France.

**Puccinia umbilici** Guép. Powdery red-brown rust pustules on leaf blades and petioles of *U. erectus* and *U. pendulinus* in Great Britain, Belgium, France, Spain, and Portugal.

**Septoria umbilici** Maire. On leaves of *U. horizontalis* in French North Africa.

**UNIFOLIUM**. See *Maianthemum*.**URECHITES**. Tropical and subtropical twining vines.

**Phyllosticta glaucispora** Delacr. On leaves of *U. lutea* in Porto Rico.

**URGINEA**. SEA ONION. Bulbous herbs.

**Accidium urginæ** P. Henn. and Evans. Yellow rust pustules on leaves of *Urginea* sp. in the Union of South Africa.

**Cladochytrium urginæ** Pat. and Trab. Oval or subcircular spots, 1 to 2 centimeters in diameter on leaves of *U. maritima* in Algeria.

**Septoria urginæ** Pass. and Beltr. A leaf spot of *U. scilla* in Sicily.

**Uromyces scillarum** (Grev.) Wint. See Hyacinthus.

**Ustilago vaillantii** Tul. See Hyacinthus.

**UROSTIGMA**. See *Ficus*.**UTRICULARIA**. BLADDERWORT. Aquatic or terrestrial plants.

**Doassansia utriculariæ** P. Henn. Smut sori in subcircular, then confluent, black leaf spots on *U. reniformis* in Brazil.

**UVARIA**. Tropical climbers.

**Actinonema uvariae** (Lév.) Sacc. On leaves of *U. parviflora* in central Africa.

**Accidium uvariae-rufae** P. Henn. Circular red-brown leaf spots on *U. rufa* in the Philippines.

**Phyllachora incareerata** (Berk.) Sacc. Shiny black stromata on leaves of *Uvaria* sp. in Ceylon.

**Phyllosticta uvariae** Berk. Subcircular to irregular brown leaf spots on *U. triloba* in Great Britain.

**VACCINIUM**. BLUEBERRY. CRANBERRY. Shrubs cultivated for their fruit.

**Gloeosporium myrtilli** Allesch. Anthracnose on *V. myrtillus* in Europe.

**Mycosphaerella stemmatea** (Fr.) Rom. On leaves of *V. vitis-idaea* in Sweden, Denmark, Esthonia, and Austria.

**Phacidium vaccinii** Fr. On leaves of *V. vitis-idaea* in Europe.

**Phyllachora japonica** Cke. and Mass. Doubtful species on *V. japonicum* in China.

**Phyllosticta lepidea** (Fr.) Allesch. On leaves of *V. vitis-idaea* in France.

**Phyllosticta vaccinii-hirti** P. Henn. Effuse red-brown leaf spots, ashen-brown beneath, on *V. hirtum* in Japan.

**Physalospora leptidea** (Hazsl.) Sacc. and Trott. On leaves of *V. vitis-idaea* in Hungary.

**Physalospora vitis-idaeae** Rehm. On leaves of *V. vitis-idaea* in Germany.

**Ramularia tumescens** (Fckl.) Sacc. On branches of *V. uliginosum* in Belgium and Germany.

**Sclerotinia baccarum** (Schroet.) Rehm. Fruit rot of *V. myrtillus* in Russia, Switzerland, Austria, and Germany.

**Sclerotinia heteroica** Wor. See *Ledum*.

**Sclerotinia urnula** Rehm. White mildew on stems, leaves, and fruit of *V. vitis-idaea* in Finland, Switzerland, and Germany. The fruits become hard and dry.

**Septoria lagerheimi** Pat. On leaves of *Vaccinium* sp. in Ecuador.

**Septoria oleae-vaccinii** P. Henn. Circular gray-white leaf spots with dark-brown margins on *V. bracteatum* in Japan.

**Septoria stemmatea** (Fr.) Berk. Subcircular dull-brown leaf spots on *V. uliginosum* in Siberia, Sweden, Great Britain, Russia, and Italy.

**Septoria vaccinii** P. Henn. On leaves of *V. bracteatum* and *V. hirtum* in Japan.

**Sphaerodothis circumscripta** (Berk.) Theiss. and Syd. Irregular black shiny stromata on leaves of *Vaccinium* spp. in Colombia and Peru.

**VAGNERA**. See *Smilacina*.**VALERIANA**. VALERIAN. Hardy herbs or shrubs, grown in gardens.

**Accidium valdivianum** Sacc. and Syd. Rust on leaf blades and petioles of *V. valdiviana* in Chile.

**Cylindrosporium valerianæ** Speg. On leaves of *V. heterophylla* in Siberia.

**Endophyllum valerianæ-tuberosae** R. Maire. Yellow rust sori on leaves of *V. tuberosa* in France and Macedonia.

**Peronospora valerianæ** Trail. Indefinite violet-gray downy-mildew patches on leaves of young plants of *V. morisonii*, *V. officinalis*, and *V. olitoria* in Scotland, France, Esthonia, and Switzerland.

**Puccinia commutata** Syd. Golden and brown rust pustules on leaf blades, petioles, and stems of *V. officinalis*, *V. sambucifolia*, and *V. tripteris* in Europe. Reported from New York and Oregon.

**Puccinia valerianæ** Carest. Dark-brown rust pustules on leaves of *V. capitata*, *V. celtica*, *V. officinalis*, *V. sambucifolia*, *V. saxatilis*, *V. tripteris*, and *Centranthus calcitrapa* in Alaska, Italy, Switzerland, and Austria.

**Septeria valerianæ** Sacc. and Fautr. Ovate to angular dull-rufous leaf spots on *V. dioica*, *V. major*, *V. montana*, *V. officinalis*, and *V. tripteris* in France, Switzerland, Denmark, and Austria.

**Uredo valerianæ-wallichii** Diet. Brown leaf rust on *V. leschenaultii* and *V. wallichii* in India.

**Uromyces valerianæ** (Schum.) Fckl. Brown leaf rust on *Valeriana* spp. in Europe, Siberia, and the Union of South Africa.

**VALERIANELLA**. Herbs grown in rock gardens, one species a salad plant.

**Accidium velenovskyi** Bub. Leaf rust on *V. membranacea* in Bulgaria.

**Accidium valerianellæ** Biv. Leaf rust on *V. campanulata*, *V. discoidea*, *V. eriocarpa*, *V. microcarpa*, *V. olitoria*, *V. pumila*, and *V. truncata* in Europe and north Africa.

**Peronospora valerianellæ** Fckl. Downy mildew on leaves of *V. carinata*, *V. dentata*, *V. morisonii*, *V. olitoria*, and *V. ramosa* in Europe.

**VALERIANOIDES**. See *Stachytarpheta*.**VANDA**. See *Orchidaceae*.**VANDOPSIS**. See *Orchidaceae*.

- VANGUERIA.** Tropical shrubs or trees.  
*Accidium vangueriae* Cke. and var. *abyssinica* P. Henn. Leaf rust on *V. edulis*, *V. infausta*, and *V. latifolia* in Abyssinia, Uganda, and the Union of South Africa.  
*Hemileia woodii* Kalchbr. and Cke. Orange-yellow rust pustules on leaves of *V. edulis*, *V. euonymoides*, *V. infausta*, *V. latifolia*, *V. madagascarensis*, and *V. pubescens* in Abyssinia, central, and south Africa.
- VANILLA.** Climbing orchids. Cultivated for their aromatic pods. See Orchidaceae.
- VERATRUM.** FALSE HELLEBORE. Hardy perennial herbs grown in gardens.  
*Cylindrosporium veratrinum* Sacc. and Wint. On leaves of *V. album* in Switzerland. Reported from Utah and New York.  
*Gloeosporium veratrinum* Allesch. Anthracnose on leaves of *V. album* (*V. lobelianum*) in Germany.  
*Mycosphaerella veratri* v. Hoeh. On *V. nigrum* in Europe.  
*Phyllachora melanocephala* Sacc. See Convallaria.  
*Phyllosticta albina* Bub. and Kab. Oblong dark-brown leaf spots on *V. album* and *V. lobelianum* in Hungary and Austria.  
*Phyllosticta macrothecia* (Thuem.) Gz. Frag. On leaves of *V. album* in Spain.  
*Phyllosticta melanocephala* Thuem. Linear black leaf spots on *V. album* and *V. nigrum* in Siberia, Russia, Switzerland, Italy, and Bulgaria. Reported from Utah.  
*Phyllosticta veratrina* Sacc. and Paol. On leaves and stems of *V. album* in Siberia.  
*Puccinia pachycephala* Diet. Brown to black rust pustules on brown or yellow leaf spots on *V. maximowiczii* and *V. nigrum* in Japan and Russia.  
*Septoria sublineolata* Thuem. Linear gray-white leaf spots with dull-brown margins on *V. album* and *V. nigrum* in Siberia.  
*Uromyces veratri* (DC.) Schroet. Yellow and powdery chestnut-brown rust pustules on leaves of *V. album*, *V. nigrum*, *V. stamineum*, *Adenostyles albifrons*, *A. alpina*, *Cacalia delphinifolia*, *C. krameri* and *Homogyne alpina* in Japan, Siberia, and Europe.
- VERBASCUM.** MULLEIN. Hardy biennial herbs, mostly weeds.  
*Ascochyta verbasci* Sacc. and Speg. Ochraceous leaf spots on *V. phlomoides* and *V. sinuatum* in Italy.  
*Erysiphe taurica* Lév. See Althaea.  
*Peronospora verbasci* Güm. Downy mildew on leaves of *V. lychnites*, *V. montanum*, *V. nigrum*, *V. phlomoides*, *V. thapsiforme*, and *V. thapsus* in central and northern Europe.  
*Phyllosticta banatica* Bub. Large irregular leaf spots on *V. banaticum* in Hungary.  
*Phyllosticta verbasci* Sacc. Small, pale-brown leaf spots on *Verbascum* sp. in France. Reported from New York and New Jersey.  
*Ramularia cylindroides* Sacc. See Pulmonaria.  
*Septocylindrium bellocense* C. Mass. and Sacc. On leaves of *V. nigrum* in Italy.  
*Uromyces thapsi* (Opiz.) Bub. Yellow and black rust pustules on leaves of *V. lychnites*, *V. neilreichii*, *V. nigrum*, *V. orientale*, *V. phlomoides*, *V. schraderei*, *V. thapsus*, and *V. thapsiformis* in Europe.
- VERBENA.** Annual or perennial herbs or subshrubs, cultivated in gardens.  
*Accidium spegazzinianum* Sacc. and Trott. Leaf rust on *V. tenera* in Argentina.  
*Cronartium flaccidum* (Alb. and Schw.) Wint. See Paeonia.  
*Microsphaera ferruginea* Erikss. Powdery mildew on leaves of *V. hybrida* in Sweden.  
*Phyllosticta verbenae* Sacc. Ashen leaf spots on *V. hastata* and *V. officinalis* in France and Denmark.  
*Puccinia elongata* Speg. Dark-brown rust pustules on stems of *V. litoralis* in Argentina.
- VERBESINA.** CROWNBEARD. Annual or perennial herbs.  
*Cercospora tubercularioides* Speg. On leaves of *Verbesina* sp. in Argentina.  
*Coleosporium anceps* Diet. and Holw. Golden rust pustules on leaves of *V. sphaerocephala* in Mexico.  
*Coleosporium verbesinae* Diet. and Holw. Golden rust pustules on leaves of *Verbesina* spp. in Mexico, Costa Rica, Guatemala, and Jamaica.  
*Puccinia affinis* Syd. Powdery black rust pustules on leaves of *V. trilobata* and *V. perymenioides* in Mexico and Guatemala.  
*Puccinia eundimarcensis* Mayor. Leaf rust on *V. verbascifolia* in Colombia.  
*Puccinia diaziana* Arth. Chestnut-brown rust pustules on leaves of *V. encelioides* in Mexico.  
*Puccinia ferox* Diet. and Holw. Powdery brown rust pustules on *V. diversifolia* and *V. myriocephala* in Mexico and Costa Rica.  
*Puccinia invelata* H. S. Jack. Leaf rust on *V. montanoifolia* in Mexico.  
*Puccinia irregularis* Diet. Brown leaf rust on *V. pallens* and *V. subcordata* in Nicaragua and Brazil.  
*Puccinia spegazziniana* Det. Leaf rust on *V. montevidensis* in Argentina.  
*Puccinia verbesinicola* Diet. Leaf rust on *Verbesina* sp. in Ecuador.  
*Septoria balansae* Speg. Indefinite leaf spots on *Verbesina* sp. and *Bidens helianthoides* in Brazil.  
*Sphaerodothis pirifera* (Speg.) Theiss. and Syd. Small, black stromata on leaves of *Verbesina* sp. in Brazil and Venezuela.  
*Uredo verbesinae-dentatae* Syd. Brown rust sori on circular leaf spots on *V. dentata* in Ecuador.
- VERNONIA.** IRONWEED. Perennial herbs, shrubs, and trees; some ornamentals.  
*Accidium banosense* Syd. Leaf rust on *Vernonia* sp. in the Philippines.  
*Accidium tarapotense* P. Henn. Dull-brown rust sori on circular leaf spots on *Vernonia* sp. in Peru.  
*Accidium vanderystianum* P. Henn. Leaf rust on *Vernonia* sp. in Congo.  
*Accidium vernoniae* P. Henn. Yellow leaf rust on *Vernonia* sp. in Brazil.  
*Accidium vernoniae-cinereae* Petch. Rust on leaves of *V. cinerea* in Ceylon.  
*Accidium vernoniae-hookerianae* Petch. Yellow leaf rust on *V. hookeriana* in Ceylon.  
*Cercospora sublateralis* P. Henn. Leaf spot on *Vernonia* sp. in Congo.  
*Endophyllum vernoniae* Arth. Powdery ochraceous rust pustules on leaves of *Vernonia* sp. in Mexico.  
*Melampsora ruspoliana* P. Henn. Yellow-brown rust sori on leaves of *Vernonia* sp. in central Africa.  
*Phyllachora vernoniicola* P. Henn. and var. *microspora* Theiss. and Syd. Small, scattered shiny black stromata on brown leaf spots on *V. tweediana* and *Vernonia* sp. in Brazil.  
*Puccinia arthuriana* Jacks. Leaf rust on *V. albicaulis*, *V. arbuscula*, *V. bahamensis*, *V. borinquensis*, *V. canescens*, *V. phyllostachys*, and *V. sericea* in Porto Rico, the Bahamas, and Costa Rica.  
*Puccinia becki* Mayor. Leaf rust on *V. cotoneaster* and *V. divaricata* in Colombia and Jamaica.  
*Puccinia discreta* Jacks. and Holw. Brown leaf rust on *V. deppeana* and *V. stellaris* in Guatemala and Costa Rica.  
*Puccinia egregia* Arth. Leaf rust on *V. uniflora* in Mexico.  
*Puccinia erratica* Jacks. and Holw. Leaf rust on *V. schiedeana* in Guatemala.

## VERNONIA—Continued.

- Puccinia fraterna** Jacks. Rust on leaves of *V. pluvialis* in Jamaica.
- Puccinia fuscella** Arth. and Johnst. Brown leaf rust on *V. menthaefolia* in Cuba.
- Puccinia hyalina** Diet. Leaf rust on *V. scariosa* in Ceylon.
- Puccinia idonea** Jacks. and Holw. Brown leaf rust on *V. triflosculosa* in Guatemala and Costa Rica.
- Puccinia inaequata** Jacks. and Holw. Rust on leaves of *V. patens* in Guatemala.
- Puccinia insulana** (Arth.) Jacks. Brown rust pustules on leaves of *V. albicaulis*, *V. arborescens*, *V. divaricata*, and *V. longifolia* in Porto Rico. St. Croix, Antigua, Guatemala, and Jamaica.
- Puccinia kuntzii** Jacks. Leaf rust on *V. kuntzei* in Bolivia.
- Puccinia le-testii** Maubl. Leaf rust on *Vernonia* sp. in central Africa.
- Puccinia lorentzii** P. Henn. Yellow-brown to dark-brown rust pustules on leaves of *V. cinerea*, *V. lorentzii*, *V. mollissima*, and *V. scorpioides* in Brazil, Argentina, and Ceylon.
- Puccinia membranacea** Diet. Yellow and cinnamon-brown rust pustules on leaves of *V. cauloni* in Brazil.
- Puccinia notha** Jacks. and Holw. Leaf rust on *V. leiocarpa* and *V. shannoni* in Guatemala.
- Puccinia pinguis** Diet. Powdery chestnut-brown rust pustules on leaves of *V. platensis* in Brazil.
- Puccinia rota** Jacks. and Holw. Brown leaf rust on *V. leiocarpa* in Guatemala.
- Puccinia rotundata** Diet. Cinnamon-colored rust pustules deforming leaf blades, petioles, and stems of *V. patens*, *V. scabra*, and *V. tweediana* in Costa Rica, Panama, Colombia, and Brazil.
- Puccinia semiinsculpta** Arth. Powdery cinnamon brown to dark-brown rust sori on leaves of *V. alamani*, *V. dictyophlebia*, *V. karvinskiana*, *V. serratuloides*, and *V. umbellifera* in Mexico.
- Puccinia tonduziana** Speg. Leaf rust on *V. triflosculosa* in Costa Rica and Guatemala.
- Puccinia vernoniae-mollis** Mayor. Leaf rust on *V. mollis* in Colombia.
- Puccinia vernoniae-scariosae** Petch. Brown leaf rust on *V. scariosa* in Ceylon.
- Puccinia vernoniicola** P. Henn. Ochraceous rust pustules on leaves of *Vernonia* sp. in tropical Africa.
- Puccinia vernoniphila** Speg. Leaf rust on *V. flexuosa* in Argentina.
- Uredo pachystegia** Diet. Brown leaf rust on *Vernonia* sp. in Brazil.
- Uredo vernoniae** P. Henn. Dull-brown rust pustules on circular leaf spots on *Vernonia* sp. in Congo.
- Uredo vernoniae-hookeriana** Petch. Leaf rust on *V. hookeriana* in Ceylon.
- Uredo vernoniicola** Petch. Brown leaf rust on *V. cinerea*, *V. setigera*, and *V. wightiana* in Ceylon.
- Uromyces pressus** Arth. and Holw. Leaf rust on *V. deppeana* in Guatemala and Costa Rica.
- VERONICA.** SPEEDWELL. Annual or biennial herbs or shrubs.
- Aecidium disciforme** McAlp. Rust distorting and thickening leaves of *V. calycina* and *V. gracilis* in Australia and Tasmania.
- Aecidium veronicae** Berk. Yellow rust pustules on lower leaf surfaces of *V. sibirica* and *Veronica* sp. in Japan and Australia.
- Ascochyta veronica** Rostr. On leaves of *V. saxatilis* in Denmark.
- Clypeostroma hemisphaericum** (Berk.) Theiss. and Syd. Small black stromata on leaves of *V. elliptica* and *V. odora* in New Zealand.
- Gloeosporium pruinatum** Baeml. and var. **tirolense** Kab. and Bub. Small subcircular, then confluent, brown spots on leaves, stems and capsules of *Veronica* spp. in Spain, Bulgaria, Denmark, and Austria.
- Laestadia veronicae** Rostr. On leaves of *V. alpina* in Iceland.
- Olpidiaster radialis** (Willd) Pascher. See Linum.
- Ovularia chamaedrys** Lindr. Subcircular gray-brown, then black, leaf spots on *V. chamaedrys* in Finland.
- Ovularia veronicae** (Fckl.) Sacc. and var. **microsticta** Sacc. On leaves of *Veronica* spp. in Ceylon, Sweden, Denmark, Italy, Esthonia, and Germany.
- Peronospora agrestis** Gäum. Downy mildew on leaves of *V. agrestis* and *V. polita* in Switzerland, Galicia, and Russia.
- Peronospora arvensis** Gäum. Downy mildew on leaves of *V. hederifolia* and *V. triphyllos* in Europe.
- Peronospora grisea** Unger. Downy mildew on leaves of *V. beccabunga* in Europe.
- Peronospora palustris** Gäum. Downy mildew on leaves of *V. scutellata* in Europe and Switzerland.
- Peronospora saxatilis** Gäum. Downy mildew on leaves of *V. fruticans* in Europe.
- Peronospora silvestris** Gäum. Downy mildew on leaves of *V. officinalis* and *V. urticifolia* in Europe.
- Phyllosticta prostrata** Brun. Gray-white leaf spots with dull-brown margins on *V. prostrata* in France.
- Phyllosticta veronicae** Thuem. On leaves of *V. longifolia* in Siberia.
- Puccinia rhaetica** Ed. Fisch. Powdery brown rust pustules on leaves of *V. bellidioides* in Switzerland. Reported from Washington.
- Puccinia veronicae** Schroet. Yellow-brown to brown rust pustules on leaves of *V. lutea* and *V. montana* in Europe.
- Ramularia anagallidis** Lindr. Circular to oblong pale-brown leaf spots on *V. anagallis* and *V. moretti* in Italy, Spain, Denmark, and Finland.
- Ramularia beccabungae** Fautr. Circular leaf spots on *V. beccabunga* in Denmark, France, and Austria.
- Ramularia caruaniana** Sacc. Subcircular, often concentrically zoned, leaf spots on *V. anagallis* in Malta.
- Ramularia coccinea** (Fckl.) Vester. (*R. pygmaea* Lindr.) Rufous-brown leaf spots on *V. chamaedrys*, *V. pygmaea*, *V. officinalis*, and *V. serpyllifolia* in Finland, Denmark, Sweden, Esthonia, Austria, and Germany.
- Ramularia nivea** Kab. and Bub. Subcircular, often confluent, dull yellow to brown leaf spots on *V. anagallis* in Bohemia.
- Ramularia pseudococcinea** Liro. Rufous to brown leaf spots with dark margins on *V. chamaedrys* in northern Europe and Spain. Said to be the same as *R. coccinea*.
- Septoria exotica** Speg. and var. **andersoni** Tassi. Confluent brown leaf spots on *V. andersoni*, *V. elliptica*, *V. salicifolia*, and *V. speciosa* in New Zealand, Argentina, Ecuador, Italy, and France.
- Septoria triphylli** Höll. On leaves of *V. triphyllos* in Hungary.
- Septoria veroneicola** Karst. On leaves of *V. for-mosa* in Italy.
- Sorosphaera veronicae** Schroet. Small irregular or cylindrical smut sori as outgrowths from leaves of *V. chamaedrys*, *V. hederifolia*, and *V. triphyllos* in Sweden, Denmark, and France.
- Synchytrium globosum** Schroet. See Potentilla.
- VESICARIA.** Annual or perennial herbs cultivated in gardens.
- Septoria cercosperma** Rostr. On leaves of *V. arctica* in Greenland.
- VIBURNUM.** Ornamental shrubs grown for their attractive flowers, fruit, and foliage.
- Aecidium viburni** Henn. and Shir. Leaf rust on *V. burejeticum*, *V. dilatatum*, *V. formosanum* and *V. sargentii* in Russia and Japan.
- Aecidium viburnophilum** Syd. Leaf rust on *V. opulus* in Japan.
- Ascochyta tini** Sacc. Ochraceous leaf spots on *V. tinus* in Italy.

**VIBURNUM**—Continued.

- Ascochyta viburni** (Roum.) Sacc. Brown spots with brown-purple margins on branches of *V. lantana* and *V. opulus* in France, Denmark, Austria, and Germany.
- Cercospora penicillata** Fekl. On leaves of *V. opulus* in Siberia, Italy, Portugal, and Germany.
- Cercospora tinea** Sacc. On leaves of *V. tinus* in Portugal.
- Laestadia tuscula** Pass. Subcircular to irregular gray-white leaf spots on *V. tinus* in Italy.
- Meliola viburni** Syd. Superficial black fungus patches on leaves of *V. odoratissimum* in China and the Philippines.
- Mycosphaerella crepidophora** (Mont.) Rehm. Circular dull-brown leaf spots on *V. tinus* in Corsica, Dalmatia, Portugal, and Tunis.
- Mycosphaerella viburni** (Nitsch) Fekl. On leaves of *V. lantana* and *V. opulus* in Italy and Germany.
- Phyllosticta lantanae** Pass. See Lantana.
- Phyllosticta lantanicola** Sacc. Irregular ochraceous, then gray-white, leaf spots on *V. lantana* in Bohemia.
- Phyllosticta opuli** Sacc. On leaves of *V. opulus* in Europe.
- Phyllosticta roumeguerii** Sacc. Indefinite gray leaf spots on *V. opulus* and *V. tinus* in Europe.
- Phyllosticta tinea** Sacc. Subcircular to irregular gray-white leaf spots on *V. tinus* in Italy.
- Phyllosticta viburni** Pass. Large irregular gray or rose-gray leaf spots on *V. tinus* in France and Portugal.
- Placosphaeria viburni** P. Henn. Black subcarbonaceous stromata on leaves of *V. dilatatum* in Japan.
- Pucciniastrum miyabeanaum** Hirats. Yellow rust pustules on leaves of *V. furcatum* in Japan.
- Rhynisma viburni** P. Henn. Black stromata on leaves of *V. dilatatum* and *V. luzonicum* in the Philippines.
- Septoria tini** (Arc.) Sacc. Irregular ashen-brown leaf spots with purple margins on *V. tinus* in Italy.
- Septoria viburni** West. On leaves of *V. lantana* and *V. opulus* in Great Britain, Denmark, Belgium, and Italy.
- VICIA**. VETCH. Annuals cultivated for forage and flowers.
- Ascochyta bolithauseri** Sacc. See Phaseolus.
- Ascochyta fabae** Speg. Circular to irregular, then confluent, dull-tan leaf spots on *V. faba* in Argentina.
- Ascochyta phaseolorum** Sacc. See Phaseolus.
- Ascochyta viciae-lathyroides** Syd. Large gray-white areas on leaves of *V. lathyroides* in Germany.
- Ascochyta viciae-pisiformis** Bub. Subcircular to irregular, then confluent, dull-yellow leaf spots with rufous margins on *V. pisiformis* in Bohemia.
- Cephalotheca francisci** Sacc. On stems of *V. faba* in Italy.
- Cercospora fabae** Fautr. Dark-purple, zoned, then confluent, leaf spots with gray centers on *V. faba*, *V. gigantea*, and *V. narbonensis* in China, Russia, Italy, Bohemia, and France.
- Cercospora zonata** Wint. On leaves of *V. faba* and *Phaseolus vulgaris* in Brazil, Portugal, Russia, Italy, and Germany.
- Cercosporella viciae** Siem. Leaf spot on *V. aurantia* in Poland.
- Didymaria lindaviana** Jaap. Large pale-brown leaf spots on *V. cracca* in Germany.
- Erysiphe taurica** Lév. See Althaea.
- Gloeosporium tricolor** Lind. Circular dark-brown leaf spots with paler margins on *V. cracca* in Sweden and Denmark.
- Hypochnus cucumeris** Frank. See Cucumis.
- Microsphaera bäumleri** P. Magn. Powdery mildew on leaves of *V. cassubica* and *V. sylvatica* in Great Britain, Russia, Italy, Austria, Hungary, and Germany.
- Ovularia fallax** (Bon.) Sacc. On leaves of *V. cracca*, *V. cassubica*, *V. pyrenaica*, and *V. villosa* in Spain and Germany.
- Ovularia schwarziiana** Magn. Brown leaf spots on *V. villosa* in Denmark and Germany.
- Ovularia sphaeroidea** Sacc. See Lotus.
- Ovularia viciae** (Frank.) Sacc. Brown leaf spots on *V. cassubica* and *V. tenuifolia* in Denmark and Germany.
- Ovularia villiana** P. Magn. On leaves of *V. cassubica* in Germany.
- Peronospora sepium** Gäum. Downy mildew on leaves of *V. sepium* in Scotland, central and northern Europe.
- Peronospora viciae** (Berk.) Gäum. Downy mildew on leaves of *V. grandiflora*, *V. hirsuta*, *V. lathyroides*, *V. pisiformis*, *V. tenuifolia*, and *V. tetrasperma* in Europe.
- Phyllachora lathyri** (Lév.) Theiss. and Syd. See Lathyrus.
- Phyllosticta fabae** West. Dull-brown leaf spots on *V. faba*, *V. sativa*, and *V. villosa* in Russia, Sweden and Belgium.
- Phyllosticta viciae** (Lib.) Cke. Subcircular to oblong ochraceous to brown leaf spots with dark margins on *V. faba* and *V. sepium* in Russia, France, and Belgium.
- Rhizoctonia napi** West. See Brassica.
- Rhynisma onobrychidis** DC. See Onobrychis.
- Septoria silvestris** Pass. See Lathyrus.
- Septoria viciae** West. Dull-brown leaf spots on *V. angustifolia*, *V. pyrenaica*, and *V. sativa* in Denmark, Russia, Spain, and Italy.
- Stagonospora carpathica** Baeuml. Small circular to subcircular red-brown spots appear on the leaves, petioles, stems, and pods, changing to dark-brown with red-brown margins, and finally to gray-brown. Infected plants often die. The hosts are *Vicia faba* in Japan and *Melilotus alba* and *M. officinalis* in central Europe.
- Uromyces briardi** Har. Brown rust pustules on leaves of *V. sativa* in France.
- Uromyces clavatus** Diet. See Lathyrus.
- Uromyces corrugatus** Speg. Powdery brown rust pustules on leaves of *V. patagonica* in Patagonia.
- Uromyces fischeri-eduardi** P. Magn. Powdery brown rust pustules on leaves of *V. cracca* and *Euphorbia cyparissia* in Europe.
- Uromyces heimerlianus** P. Magn. Brown rust pustules on leaves of *V. hirsuta* and *V. pannonica* in Yugoslavia, Hungary, and Austria.
- Uromyces johowii** Diet. and Neg. Yellow to brown rust pustules on leaves of *V. nigricans* in Chile.
- Uromyces nordenskjöldii** Diet. Yellow-brown rust pustules on leaves of *Vicia* sp. in South America.
- Uromyces orobi** (Pers.) Lév. See Lathyrus.
- Uromyces unijugae** Ito. Leaf rust on *V. unijuga* in Japan.
- Uromyces valesiacus** Ed. Fisch. Yellow and brown leaf rust on *V. onobrychioides* in Macedonia and Switzerland.
- Uromyces viciae-cracciae** Const. Powdery brown rust pustules on leaves of *V. cracca*, *V. tenuifolia*, and *Lens esculenta* in Europe.
- Uromyces viciae-unijugae** S. Ito. Leaf rust on *V. unijuga* in Japan.

- VIGNA. COWPEA.** Herbs grown as vegetables, forage, and green-manure crops.
- Aecidium caulicola** P. Henn. Rust on stems of *Vigna* sp. in Central Africa.
- Aecidium nigro-cinctum** Pat. and Har. Leaf rust on *Vigna* sp. in Indo-China.
- Aecidium vignae** Cke. Leaf rust on *V. catjang*, *V. marginata*, and *V. sinensis* in Ceylon, Uganda, and Union of South Africa.
- Ascochyta phaseolorum** Sacc. See Phaseolus.
- Ascochytopsis vignae** P. Henn. Dull-brown leaf spots on *V. vexillata* in east Africa.
- Auerswaldia vignae** P. Henn. Clear brownish spots on leaves of *V. vexillata* in Tanganyika.
- Cercospora vanderysti** P. Henn. Effuse brown leaf spots on *V. venulosa* in the Congo.
- Hypochnus cucumeris** Frank. See Cucumis.
- Mycosphaerella phaseolorum** Jachw. On leaves of *V. rubra* in Russia.
- Phyllachora phaseolina** Syd. See Phaseolus.
- Phyllosticta phaseolorum** Sacc. and Speg. Ochraceous leaf spots on *V. sinensis* and *Phaseolus vulgaris* in Japan, Formosa, and Italy.
- Phyllosticta vignae** Speg. Circular pale dull-brown leaf spots on *V. luteola* in Argentina.
- Puccinia** sp. Leaf rust on *V. angustifolia* in the Union of South Africa.
- Rhizoctonia** sp. Diseased plants wilt and die, the stems at the ground level being blackened and covered with numerous tiny black sclerotia. Among the hosts are *V. catjang*, *V. unguiculata*, *Corchorus capsularis*, *C. olitorius*, *Dolichos lablab*, *D. biflorus*, *Cucurbita* sp., *Citrullus* sp., *Gossypium* sp., *Lycopersicum esculentum*, *Medicago sativa*, *Nicotiana tabacum*, *Phaseolus lunatus*, *P. radiatus*, *Sesamum indicum*, and *Solanum tuberosum* in India.
- Rhizoctonia dimorpha** Matz. See Phaseolus.
- Septoria vignae** P. Henn. Circular dull-brown leaf spots on *Vigna* sp. in the Congo.
- Uredo purpurascens** P. Henn. Rust on circular to effuse purple spots on leaves and stems of *Vigna* sp. in the Congo.
- Uredo vignae** Bres. Brown leaf rust on *V. lutea*, *Glycine hispida*, *Phaseolus mungo*, and *Phaseolus* sp. in the Philippines and St. Thomas.
- Uromyces pазschkeanus** P. Henn. Black rust sori on leaves of *V. membranacea* and *V. triloba* in Abyssinia and Arabia.
- Uromyces punctiformis** Syd. Powdery black rust pustules on leaves of *V. strobiliphora* in Mexico.
- Uromyces vignae** Barel. Brown to black rust pustules on leaves of *V. vexillata* in India.
- Vermicularia capsici** Syd. See Capsicum.
- VIGUIERA.** Shrubby or herbaceous composites.
- Coleosporium viguieriae** Diet. and Holw. Golden rust pustules on leaves of *V. helianthoides* in Mexico.
- Puccinia ensenadensis** Speg. Powdery black rust pustules on leaves of *Viguiera* sp. in Argentina.
- Puccinia inopinata** Syd. Dark-brown rust pustules on leaves of *V. stenophylla* in Argentina.
- Puccinia nanomitra** Syd. Powdery black rust pustules on leaves of *V. dentata*, *V. eriophora*, and *V. helianthoides* in Mexico.
- Puccinia punctoidea** Syd. Black rust pustules on leaves of *V. pringlei* in Mexico.
- Puccinia subglobosa** Diet. and Holw. Brown leaf rust on *V. palmeri* in Mexico.
- VINCA. PERIWINKLE.** Erect and procumbent herbs or subshrubs cultivated for their flowers.
- Colletotrichum vincae** Speg. Circular ashen leaf spots on *V. major* in Argentina.
- Macrophoma cylindrospora** Desm. See Hedera.
- Metasphaeria vincae** (Fr.) Sacc. On leaves of *V. minor* in Italy, France, and Germany.
- Peronospora vincae** Schroet. Downy mildew on leaves of *V. major* and *V. minor* in France and Russia.
- Phytophthora parasitica** Dast. See Ricinus.
- Phyllosticta vincae** Thuem. and var. **pedrosensis** Gz. Frag. Large irregular dull-brown to gray leaf spots on *V. media* and *V. minor* in Spain, Portugal, and France.
- Phyllosticta vincae-majoris** Allesch. Subcircular ochraceous leaf spots on *V. major* in Germany.
- Phyllosticta vincae-minoris** Bres. and Krieg. Dull-brown leaf spots on *V. minor* in Albania, Yugoslavia, and Germany.
- Phyllosticta vincicola** Oud. Irregular black leaf spots on *V. major* in Holland.
- Puccinia vincae** (DC.) Berk. Brown to black rust pustules on leaves of *V. acutiflora*, *V. herbacea*, *V. major*, *V. media*, and *V. minor* in Europe and Algeria. Reported from Massachusetts and Michigan.
- Ramularia vincae** Sacc. and var. **vincae-mediae** Maire. Yellow indefinite leaf spots on *V. major* and *V. media* in Spain and the Balearic Islands.
- VIOLA. VIOLET.** Pansy. Perennial herbs, cultivated for their flowers.
- Aphelenchus olesistus** Ritz. Bos. var. **longicollis** Schwartz. This nematode attacks the bases of the leaves and flower stalks of cultivated *Viola*, causing hypertrophy, with consequent stunting and fasciation and finally production of galls. The disease occurs in Europe.
- Ascochyta violae-hirtae** Bub. White subcircular leaf spots with brown margins on *V. hirta* in Yugoslavia.
- Ascochyta violicola** McAlp. Yellow to gray-white circular to ovate leaf spots on *V. odorata* in Australia.
- Cercospora** ff Trail. Indefinite dull-brown leaf spots on *V. palustris* in Scotland.
- Cercospora lilacina** Bres. On leaves of *V. palustris* in Germany.
- Cercospora violae-sylvatica** Oud. Subcircular pale-brown leaf spots on *V. sylvatica* in Holland.
- Cercospora violae-tricoloris** Br. and Cav. Large subcircular ashen leaf spots on *V. tricolor* in Italy and Spain.
- Cladochytrium violae** Berl. On leaves of *V. odorata* in Europe.
- Entyloma anzianum** Pass. Smut sori on subcircular yellow leaf spots on *V. biflora* in Italy.
- Fusisporium lacteum** Desm. On leaves of *V. odorata* in France.
- Hendersonia triseptata** Da Cav. Subcircular leaf spots on *V. alba* in Portugal.
- Laestadia violae** (Lib.) Sacc. On leaves of *V. odorata* in Belgium.
- Melampsora lapponum** Lindf. See Salix.
- Mycosphaerella violae** A. Pot. On leaves of *V. hirta* in Russia. Possibly the perfect stage of *Ramularia lactea* Sacc.
- Ovularia acutata** (Bon.) Sacc. Small ashen leaf spots with dull-brown margins on *V. odorata* and *V. sylvatica* in Sweden and Germany.
- Peronospora violae** De By. A downy mildew destroying the leaves of *V. biflora*, *V. tricolor*, and other species in Europe. American collections referred to this species should be referred to *Bremiella megasperma* (A. Berl.) G. W. Wils.
- Phyllosticta libertianae** Sacc. and March. Circular black leaf spots on *V. biflora* and *V. odorata* in Great Britain, France, Belgium, and Italy.
- Phyllosticta tricoloris** Sacc. On leaves of *V. odorata* and *V. tricolor* in Russia.
- Puccinia cingens** Bomm. and Rouss. Yellow and black rust pustules on leaves of *Viola* in Patagonia.

## VIOLA—Continued.

- Puccinia depauperans** (Vize.) Syd. Yellow and brown to black rust pustules on leaves and stems of *V. cornuta*, *V. lutea*, *V. tricolor*, and *V. willkommii* in Europe.
- Puccinia hederaceae** McAlp. Powdery black rust pustules on leaf blades, petioles, and stems of *V. betonicifolia* and *V. hederacea* in Australia and Tasmania.
- Puccinia violae-glabbellae** Miura. Powdery brown rust pustules on leaves of *V. glabella* in Japan.
- Ramularia acutata** Bon. Subcircular, then confluent, spots occupying entire surface of leaf blades of *V. canina*, *V. riviniana*, *V. silvatica*, and *V. sylvestris* in Denmark, Russia, Scotland, and Austria.
- Ramularia agrestis** Sacc. Small subcircular olivaceous leaf spots on *V. altaica* and *V. tricolor* in Denmark, Esthonia, and Italy.
- Ramularia biflorae** P. Magn. Subcircular greenish brown, then gray-white, irregularly zoned leaf spots on *V. biflora* in Austria.
- Ramularia deflectens** Bres. On leaves of *V. tricolor* in Russia, Denmark, and Germany.
- Ramularia lactea** (Desm.) Sacc. Circular gray-white leaf spots with brown margins on *V. canina*, *V. hirta*, *V. odorata*, *V. sylvestris*, and *V. tricolor* in Europe and Argentina. Reported from Montana and Colorado.
- Septoria australiae** McAlp. Circular to irregular red-brown leaf spots with dark-brown margins on *V. betonicifolia* in Australia.
- Septoria violae-palustris** Died. On leaves of *V. palustris* in Austria.
- Septoria violicola** Sacc. On leaves of *V. biflora* in Great Britain, Switzerland, and Germany.
- Synchytrium alpinum** Thomas. Small galls on leaves and stems of *V. biflora* in Italy and Switzerland.
- Synchytrium globosum** Schroet. See *Potentilla*.
- Uredo alpestris** Schroet. Leaf rust on *V. biflora* in Japan, Switzerland, and Austria.
- Urocystis violae** Fisch. Smut sori causing blistering and distortion of leaf blades and petioles of *V. hirta*, *V. odorata*, and *V. tricolor* in Great Britain, France, Italy, Denmark, and Germany. The pustules finally split irregularly, exposing black sooty spore masses. Also reported from Utah and Minnesota on wild species.
- VITEX.** CHASTE TREE. Ornamental shrubs and trees with white or blue flowers.
- Aecidium viticis** Jul. Leaf rust on *Vitex* sp. in Trinidad.
- Helminthosporium viticis** Syd. Gray-brown leaf spots on *V. flavens* in Brazil.
- Phyllachora taruma** Speg. Shiny black circular stromata on leaves of *V. montevidensis* and *Vitex* sp. in Paraguay and Brazil.
- Phyllachora viticicola** P. Hent. Shiny black stromata on clear brown leaf spots on *Vitex* sp. in Brazil.
- Ramularia viticis** Syd. Circular to irregular brown leaf spots on *V. negundo* in India.
- Uredo raciborskiana** Sacc. and Trott. Yellow leaf rust on *V. leucoxydon* in Java.
- Uredo vitexi** Rac. Leaf rust on *V. leucoxydon* in Java.
- Uredo viticis** Juel. Leaf rust on *Vitex* sp. in Paraguay. Hennings gives the same name to a species on *V. polygama* in Brazil.
- Uredo viticis-polygamae** P. Henn. Dull-brown rust pustules on circular leaf spots on *V. polygama* in Brazil.
- VITIS.** GRAPE. Woody vines.
- Aecidium vitis** A. L. Sm. Rust pustules on dull-brown leaf spots on *V. vulpina* in Tanganyika.
- Alternaria viticola** Brun. Ashen leaf spots on *V. vinifera* and *V. vulpina* in Italy.
- Ascochyta baecicola** P. Brun. Spots on mature fruit of *V. vinifera* in France.
- Ascochyta chlorospora** Speg. On leaves of *V. vinifera* in Australia and Argentina.
- Aureobasidium vitis** Viala and Boyer. Causes a dying back and blackening of twig tips, irregular red patches on leaves which fall prematurely and a spotting and shrivelling of the fruit of *V. vinifera* in Australia, Japan, Russia, France, Denmark, and Italy.
- Bacillus ampelopsorae** Trev. Said to cause small spongy galls on the roots and at the base of stems of *Vitis* in Argentina, France, Italy, and Germany. Probably crown-gall (*Bacterium tumefaciens*).
- Bacillus uvae** Cug. and Macc. Bacteriosis of *V. vinifera* in Italy.
- Bacillus vitivorus** Bacc. Said to cause malformation of the leaves, stripes on young branches, and dark lines in the wood of *V. vinifera* in Argentina and Italy.
- Cercospora roesleri** (Catt.) Sacc. Indefinite irregular olivaceous leaf spots on *V. vinifera* in Europe and Argentina.
- Cercospora vitis-heterophyllae** P. Henn. On leaves of *V. heterophylla* in Japan.
- Chrysomyxa burnati** Butl. Numerous yellow to brown powdery rust pustules on leaves of *V. latifolia* in India, causing defoliation.
- Clasterosporium putrefaciens erucipes** N. Spes. See *Morus*.
- Coryneum vitiphyllum** N. Spes. Numerous subcircular dull-brown leaf spots with purple-brown surrounding zones on *V. vinifera* in Russia.
- Exobasidium clandestinum** N. Spes. On leaves of *V. vinifera* in Russia.
- Fusidium peronosporae** Fautr. and Lamb. On leaves of *V. vinifera* in France.
- Fusicoccum bulgaricum** Bub. On *V. vinifera* in Austria.
- Helicobasidium tanakae** Miy. See *Morus*.
- Hendersonia rubi** West. See *Rubus*.
- Hypochnus burnati** Lendn. On trunks of *V. vinifera* in Switzerland.
- Kuehneola vitis** (Butl.) Syd. Powdery yellow and brown rust pustules on leaves of *V. latifolia* in India.
- Leptosphaeria vinealis** Pass. (*L. appendiculata* Pir.) On leaves and stems of *V. vinifera* in Italy.
- Marsonia viticola** Miy. On leaves of *V. vinifera* in China.
- Mycosphaerella rathayi** Nyp. On *Vitis* sp. in Belgium.
- Napicladium pusillum** Cav. On fruit of *V. vinifera* in Italy.
- Ovularia vitis** Rich. On leaves of *V. vinifera* in France, Austria, and Germany.
- Phakopsora ampelopsidis** Diet. and Syd. See *Ampelopsis*.
- Phakopsora cronartiiformis** Diet. Yellow and brown rust pustules on leaves of *V. himalayana* in India. Causes leaf-fall.
- Phyllochersonia vitiphylla** (N. Spes.) Tassi. Brown leaf spots on *V. vinifera* in Transcaucasia.
- Phyllosticta badbami** Cke. Indefinite elongate brown leaf spots on *V. vinifera* in Great Britain.
- Phyllosticta bizzzeriana** C. Mass. Small irregular brown leaf spots on *V. vinifera* in Europe.
- Phyllosticta dzumajensis** Bub. On leaves of *V. vinifera* in Bulgaria.
- Phyllosticta frankiana** Sacc. and Syd. Brown leaf spots with red margins on *V. vinifera* in Transcaucasia.
- Phyllosticta microspila** Pass. Small irregular dark-brown leaf spots on *V. vinifera* in Italy.
- Phyllosticta negeriana** (Thuem.) Allesch. Gray leaf spots, dark-brown beneath, on *V. vinifera* in Italy and Central Europe.
- Phyllosticta neurospileae** Sacc. and Berl. Red to ochraceous leaf spots on *V. antarctica* in Australia.
- Phyllosticta pilispora** N. Spes. Irregular dull-brown leaf spots on *V. vinifera* in Transcaucasia.

**VITIS**—Continued.

- Phyllosticta succedanea** (Pass.) Allesch. Dark-brown leaf spots on *V. vinifera* in Italy.
- Phyllosticta viticola** Sacc. and Speg. Gray-white leaf spots on *Vitis* sp. in Italy.
- Phyllosticta vitis** Sacc. Irregular gray leaf spots with brown borders on *V. vinifera* in Italy and central Europe.
- Pseudoopeziza tracheiphila** Miller-Thurg. Said to attack the leaves of *V. vinifera* in Europe, causing the disease known as "rotbrenner."
- Pseudocercospora vitis** (Lév.) Sacc. On leaves of *V. rupestris* in Argentina and Brazil.
- Septocylindrium dissiliens** (Duby.) Sacc. Causes a drying of the leaves of *Vitis* sp. in Italy.
- Septoria melanopsis** Pat. Small numerous angular blackish leaf spots on *V. vinifera* in Tunis, Brazil, England, and Italy.
- Septoria tassiana** Syd. (*S. antarctica* Tass.) Red-ochraceous leaf spots on *V. antarctica* in Italy.
- Valsaria ampleina** Avena. A wound parasite causing cankers of stems and twig die-back of *V. rupestris* and *V. vinifera* in Brazil.
- VITTADINIA.** Composite herbs with white, yellow, or blue flower heads.
- Puccinia vittadeniae** McAlp. Yellow and brown rust pustules on leaves of *V. australis* in Victoria.
- VOCHYSIA.** Resinous tropical trees.
- Acidium vochysiae** P. Henn. Leaf rust on *Vochysia* sp. in Brazil.
- Phyllachora congregata** Theiss. and Syd. Dull-black stromata on leaves of *V. elliptica* in Brazil.
- Phyllachora granulosa** Lév. Black stromata on large leaf spots, yellow beneath, gray-brown above, on *Vochysia* sp. in Brazil.
- Phyllachora lehmanniana** P. Henn. Black stromata on leaves of *V. lehmanni* in Colombia.
- Phyllachora vochysiae** P. Henn. Black stromata on leaves of *V. haenkeana* in Brazil.
- VOUACAPOUA.** See Andira.
- WARNERIA.** See Gardenia.
- WATSONIA. MERIANA** Ag. BUGLE LILY. Summer-flowering bulbous herbs resembling *Gladiolus*.
- Uromyces watsoniae** Syd. Brown to dark-brown or black rust pustules on both leaf surfaces of *W. densiflora* in the Union of South Africa.
- WEDELIA.** Composite annual or perennial herbs or subshrubs.
- Acidium wedeliae-hispidae** Diet. Leaf rust on *W. hispida* in Mexico.
- Endophyllum wedeliae** (Earle) W. and O. Yellow rust pustules on leaves of *W. trilobata* in Porto Rico, Haiti, and Trinidad.
- Laestadia lorentzii** Speg. Circular to angular pale-yellow leaf spots with ashen centers on *W. glauca* in Argentina.
- Puccinia ecuadorensis** Arth. Brown leaf rust on *Wedelia* sp. in Ecuador.
- Puccinia wedeliae** Mayor. Leaf rust on *W. trichostephia* in Colombia.
- Septoria lorentzii** Speg. Circular to angular leaf spots on *W. glauca* in Argentina.
- Uredo nerviseda** Syd. Leaf rust on *W. biflora* in the Philippines.
- Uredo vicina** Arth. Leaf rust on *W. jacquini* and *W. lanceolata* in Porto Rico and Trinidad.
- Uredo wedeliae-biflorae** Syd. Cinnamon-brown rust pustules on leaves of *W. biflora* in the Philippines.
- Uromyces baccarinii** Syd. Powdery dark-brown rust pustules on leaves of *Wedelia* sp. in Abyssinia.
- Uromyces piahyensis** P. Henn. Brown rust pustules on leaves of *W. reticulata*, *W. rugosa*, and *Wedelia* sp. in Brazil, Porto Rico, and Cuba.
- Uromyces wedeliae** P. Henn. Brown rust pustules on leaves of *W. prostrata* and *Wedelia* sp. in Japan, the Philippines, and Abyssinia.
- WELWITSCHIA. TUMBOA** Ag. Peculiar low, woody coniferous desert plants.
- Ustilago welwitschiae** Bres. Powdery dull-brown smut sori in the cone scales of *W. mirabilis* in south Africa.
- WIGANDIA.** Tall perennial herbs or subshrubs, used as bedding plants or grown in greenhouses.
- Phyllosticta wigandiae** Thuem. On leaves of *W. imperialis* in south Europe.
- WIKSTROEMIA.** Ornamental shrubs and trees.
- Melampsora yoshinagai** P. Henn. Powdery yellow, then dark-brown, rust pustules on leaves of *W. canescens*, *W. indica*, *W. japonica*, and *W. sikokiana* in Japan, Formosa, China, and India.
- WISTERIA. KRAUNHIA** Ag. Attractive woody climbers with pea-shaped flowers.
- Ascochyta wisteriae** Tassi. Leaf spots on *W. sinensis* in Italy.
- Bacterium montemartinii** Pav. Causes large irregular hollow spots on branches of *Wisteria* sp. in Italy.
- Mycosphaerella wistaricola** Turc. Causes a leaf spot of *W. chinensis* in Italy. Probably the perfect stage of *Ascochyta wisteriae* Fl. Tass.
- Ochropora kraunhiae** Diet. Yellow and brown rust pustules on leaves of *W. sinensis* and *W. floribunda* in Japan.
- Phyllosticta affinis** Tass. On leaves of *W. sinensis* in Italy.
- Phyllosticta wistariae** Sacc. Ochraceous leaf spots on *W. floribunda* (*W. brachybotrys*) and *W. sinensis* in Italy and France. Reported from Texas.
- Septoria wistariae** Brun. Irregular brown leaf spots on *W. sinensis* in France. Tharp has given the same name to a form on *W. sinensis* in Texas.
- XANTHORRHOEA.** GRASS TREE. Perennials with a thick woody caudex.
- Coniothyrium xanthoraceae** P. Henn. Pale-brown leaf spots on *X. gracilis* in Australia.
- XANTHOSOMA. MALANGA.** Yautia. Ornamental-leaved aroids, corms of some species edible.
- Phyllosticta xanthosomatis** Sacc. Subcircular dull-yellow leaf spots on *Xanthosoma* sp. in Mexico.
- Phytophthora** sp. A downy mildew is considered the cause of a wilt and soft white rot of the corms of *Xanthosoma* sp. in Hawaii and the Philippines. A similar parasite may be concerned in the wilt disease of *Xanthosoma* in Porto Rico. These diseases have not been studied in detail and the fungus present may be *P. colocasiae* Rac. See Colocasia.
- Vasculomyces xanthosomae** Ashby. This fungus was found in Jamaica in connection with a wilting and tuber rot of *X. sagittifolium*. The fungus discolors the vascular bundles and sets up a brown rot, cavities forming in the tissues. The leaves turn yellow.
- Vermicularia xanthosomatis** Sacc. Leaf spot on *X. sagittifolium* in China and the Philippines.
- XANTHOXYLUM.** See Zanthoxylum.
- XERANTHEMUM.** Annual erect tomentose herbs.
- Puccinia xeranthemi** Mori. Brown rust pustules on leaves and stems of *X. annuum* in Italy.
- XYLOPIA.** Shrubs and trees, grown for their fruit and other products.
- Acidium xylopiæ** P. Henn. Leaf rust on *Zylopiæ* sp. in Brazil.
- Diplochorella fertilissima** Syd. Black stromata on leaves of *X. aethiopica* in Tanganyika.
- Phyllachora zylopiæ** P. Henn. Irregular black stromata on large brown leaf spots on *X. grandiflora* in Peru.
- Phyllosticta xylopiæ** Sacc. Subcircular white leaf spots with black raised margins on *Zylopiæ* sp. in Mexico.
- Puccinia foveolata** (B. and C.) P. Henn. (*Puccinia gregaria* Kze.) Dark-brown rust pustules on leaves of *X. grandiflora* and *Xylopiæ* sp. in Guatemala, Peru, Surinam, and Brazil.

**XYLOSMA.** Tropical trees and shrubs.

*Phyllachora pittieri* Speg. See Myrsine.

*Phyllachora xylosmatis* Speg. Smooth black stromata on leaves of *X. pubescens* in Argentina.

*Phyllosticta tonduzi* Speg. On leaves of *X. salzmanni* in Costa Rica.

*Uredo recondita* Speg. Leaf rust on *X. salzmanni* in Costa Rica.

**YUCCA.** Ornamental stiff-leaved plants.

*Colletotrichum yuccae* Pöhl. Anthracnose on leaves of *Y. filamentosa* in Italy.

*Gloeosporium victoriosum* Sacc. Leaf anthracnose of *Y. gloriosa* in Spain and Italy.

*Sclerotium yuccae* Roum. On *Yucca* sp. in France.

*Septoria macrospora* Alm. and Cam. On leaves of *Y. aloifolia* in Portugal.

**ZANTEDESCHIA.** CALLA. Perennial herbs grown for their ornamental corolla-like spathes.

*Bacillus* sp. A bacillus said to be distinct from *B. aroidae* Town. attacks *Zantedeschia* in Great Britain. Diseased plants stop developing, the tops turning yellow and wither, while roots and corms are attacked by a soft watery brown rot.

*Gloeosporium callae* Oud. Large yellow, then brown, leaf spots on *Z. aethiopica* in Holland.

*Ramularia richardiae* Kalchbr. On leaves of *Z. albomaculata* in the Union of South Africa.

*Septoria callae* (Lasch.) Sacc. Small, olivaceous leaf spots on *Z. palustris* in Germany.

**ZANTHOXYLUM.** PRICKLY ASH. Trees and shrubs grown for their handsome foliage and ornamental fruit.

*Acanthohecium mirabile* Speg. On leaves of *Zanthoxylum* sp. in Brazil.

*Accidium spissum* Syd. Leaf rust on *Zanthoxylum* sp. in India.

*Accidium zanthoxyli-schinifolii* Diet. Rust on yellow or brown spots on leaf blades and petioles of *Z. schinifolium* in Japan.

*Bacterium citri* Hasse. See Citrus.

*Coleosporium xanthoxyli* Diet. and Syd. Yellow rust pustules on leaves of *Z. ailanthoides*, *Z. bungei*, *Z. piperitum*, and *Z. schinifolium* in Japan, Formosa, and China.

*Helicobasidium tanakae* Miy. See Morus.

*Phyllachora applanata* Wint. Subcircular to elliptic, black stromata on leaves of *Zanthoxylum* sp. in Brazil.

*Phyllachora tijucensis* (Rehm.) Theiss. and Syd. Circular black stromata on dark-brown leaf spots on *Zanthoxylum* sp. in Brazil.

*Phyllachora winteri* Sacc. and Syd. Black stromata on leaves and leaf-sheaths of *Z. hiemale*, *Z. rhoifolium*, and *Zanthoxylum* sp. in Brazil.

*Phyllachora zanthoxyli* (Lév.) Cke. Black stromata on leaves of *Zanthoxylum* sp. in Java and Brazil.

*Sorokina uleana* Rehm. On leaves of *Zanthoxylum* sp. in Brazil.

*Uredo asperata* B. and C. Brown rust pustules on leaves of *Z. piperitum* in Japan.

*Uredo fagarae* Syd. Leaf rust on *Z. (Fagara) nitidum* in Japan and Formosa.

**ZEAE.** MAIZE. INDIAN CORN.

*Dothiorella zeae* Foex and Berth. Grain, cobs, and glumes overgrown and rotted by mycelium of the fungus. On *Z. mays* in Indo-China.

*Fusarium maydiperdum* Bub. Causes a rot of cobs of *Z. mays* in Bulgaria.

*Nigrospora panici* Zimm. See Triticum.

*Phyllachora maydis* Maubl. Circular to elongate slightly raised black stromata on leaves of *Z. mays* in Porto Rico, Cuba, and Mexico.

*Physoderma maydis* Miy. Numerous circular to elliptical or linear, then confluent, brown spots on culms, mid ribs, and base of husks of *Z. mays* in Japan. Possibly not distinct from *P. zeae-maydis* Shaw, which occurs in India and the southern United States.

*Puccinia pallescens* Arth. See Tripsacum.

*Rhizoctonia palida* Matz. See Saccharum.

*Sclerospora* sp. A downy mildew of *Z. mays*, characterized by broad, pale-green stripes on the leaves, is reported from the Union of South Africa.

*Sclerospora javanica* Palm. Young plants remain stunted, with narrow, yellow or greenish-yellow leaves, with poorly developed root systems. Older plants show the yellow-striped leaves and in more advanced cases narrow, sharply defined yellow or brown stripes. The host is *Z. mays* in Java.

*Sclerospora macrospora* Sacc. See Triticum.

*Sclerospora maydis* (Rac.) Butl. The upper parts of infected plants turn white due to the disappearance of the chlorophyll in long streaks along the leaves. Growth of the plants is checked and the internodes are frequently shortened so as to give a bunched appearance. No grain is produced, the plants withering and dying before harvest time. In addition to *Zea*, *Coix lachrymi-jobi*, and *Euchlaena mexicana* are attacked. This species is known only from India, although it has been confused with *S. javanica*, the Javan species, and reported from that country.

*Sclerospora philippinensis* Weston. This downy mildew is a very serious enemy of *Zea mays* in the Philippines, and together with *S. spontanea* would appear to be the limiting factor in corn production. Additional hosts are *Euchlaena luxurians* and *Holcus sorghum* and by inoculation *Saccharum spontaneum* and *Miscanthus japonicus*.

Plants of all ages are attacked. Young plants, when attacked, show at the base of the upper leaves two or three narrow linear stripes pale-yellow to white in color. Leaves developing above these are almost completely yellow or whitish, remain narrow, and become rigid instead of bending in normal manner. The leaf sheaths continue to develop and may inclose or grow beyond the stunted tassel. The plants remain dwarfed. The roots also become stunted. Such plants are either destroyed by secondary organisms (*Pythium*) or turn brown, wither, and die. Older plants when attacked show the same stripes, generally on the fourth or fifth leaves, but more extensive and broader. Subsequent leaves show the striping to an even greater degree, the last leaves being striped throughout their length. The leaves themselves are not altered in size but at times the midribs become brittle, break at the union with the sheaths, and hang straight down along the stems. The root systems are not affected. The tassels may appear normal, but there is a decreased pollen production. Ears are only partly produced, and then always abnormal. Malformations of many kinds occur, including fasciations and reduplications, torsions of the stem, and other abnormalities.

*Sclerospora sacchari* Miy. See Saccharum.

*Sclerospora spontanea* Weston. This species of downy mildew causes a disease not distinguishable macroscopically from that due to *S. philippinensis*. It appears to be limited to the southern part of the Philippine Islands, where it occurs naturally on *Z. mays*, *Saccharum officinarum*, and *S. spontaneum*. By artificial inoculations the fungus has been found capable of attacking in addition *Euchlaena luxurians* and *Miscanthus japonicus*.

*Septoria maydis* Schulz. and Sacc. On leaf sheaths of *Z. mays* in Spain and Bohemia.

*Sphaerulina maydis* P. Henn. Circular to angular brown leaf spots on *Z. mays* in Brazil.

*Ustilago abortifera* Speg. A smut involving ovaries and often rachides in dark-brown powdery spore masses. On *Z. mays* in Argentina. Probably not distinct from *U. maydis*.

[Undet.] An undetermined fungus is reported from South Africa as destroying grain of *Z. mays* during germination.

**ZENOBLIA.** Ornamental evergreen shrubs.

*Phyllosticta andromedae* West. See *Pieris*.

**ZEPHYRANTHES. ATAMOSCO** Ag. ZEPHYR LILY. Tunicate bulbous herbs.

*Accidium zephyranthis* Shear. Yellow leaf rust on *Zephyranthes* sp. in Mexico.

*Septoria psittacina* F. Tassi. Circular to irregular yellow leaf spots, which often become confluent, on both leaf surfaces of *Z. lindleyana* in Italy.

**ZINGIBER.** GINGER. Perennial rhizomatous herbs.

*Bacillus zingiberi* Uyeda. Reported as the cause of a soft rot of the rhizomes of *Z. officinale* in Japan. Infected plants turn yellow and wither.

*Mycosphaerella zingiberi* Shir. and Hari. On leaves of *Z. mioga* in Japan.

*Phyllosticta zingiberi* Shir. and Hari. On leaves of *Z. officinale* in Japan.

*Piricularia zingiberi* Nish. Leaf spot on *Z. mioga* and *Z. officinale* in Japan.

*Pythium butleri* Subra. The leaves of infected plants turn yellow and die, hanging down parallel to the stem, due to a soft rot which works into the bases of the shoots and down into the rhizomes. The latter are completely destroyed. On *Z. officinale* in India.

*Taphrina maculans* Butl. See *Curcuma*.

*Vermicularia zingiberæ*. Small circular to oval yellow spots on both leaf surfaces of *Z. officinale* in India. The spots increase in size and become coalescent, forming large discolored patches. The centers often dry up and drop out. The leaf sheaths and scaly portions of the stems are similarly attacked. The disease causes reductions in yield.

**ZINNIA.** Annuals grown in gardens for their flowers.

*Ascochyta zinniae* Allesch. Irregular, often confluent, dark-brown leaf spots on *Z. elegans* in Germany.

*Corticium* sp. See *Thea*.

*Phyllosticta zinniae* P. Brun. Angular to irregular brown, then olivaceous, leaf spots on *Z. elegans* in France.

*Puccinia melampodii* Diet. and Holw. Powdery gray-black rust sori on irregular brown sunken leaf spots on *Z. tenuiflora*, *Eleutheranthera ruderalis*, *Synedrella nodiflora*, *Tridax procumbens*, and *Spilanthes oleracea* in Mexico, West Indies, and Central America.

**ZIZANIA.** WILD RICE. Aquatic grasses.

*Uromyces coronatus* Yosp. Brown rust pustules on leaves of *Z. aquatica* in Japan.

*Ustilago esculenta* P. Henn. Smut sori in the heads of *Z. aquatica* and *Z. latifolia* in China, Japan, Formosa, and Russia. These smut masses in the young condition are sold for food in China.

**ZIZYPHUS.** JUJUBE. Shrubs and small trees grown for their handsome foliage and some species for edible fruit.

*Ascochyta zizyphi* Putt. Circular, then irregular dark-brown leaf spots on *Z. spina-christi* in Brazil.

*Cercospora zizyphi* Petch. Yellow, then brown, leaf spots on *Z. oenoptia* in Ceylon.

*Cladosporium zizyphi* Karst. and Roun. Leaf spot on *Z. jujuba* in India and Amboina.

*Cronartium zizyphi* Syd. and Butl. Dull-brown rust pustules on leaves of *Z. oenoptia* and *Z. rugosa* in Ceylon, India, and Indo-China.

*Hyalodema evansi* P. Magn. Galls on stems and petioles of *Z. baclei* and *Z. mucronata* in the Union of South Africa. The fruits are also deformed.

*Phakospora zizyphi-vulgaris* (P. Henn.) Diet. Black rust pustules on leaves of *Z. jujuba*, *Z. rotundifolia*, and *Z. sativa* (*Z. vulgaris*) in India, Formosa, and Japan.

*Phyllosticta zizyphi* Thuem. Irregular gray-ochraceous leaf spots on *Z. vulgaris* in Portugal.

*Septoria capensis* Wint. Gray to grayish-white leaf spots on *Z. mucronata* in the Union of South Africa.

*Septoria jujubæ* Trav. and Spessa. Small irregular ochraceous to dull-brown leaf spots on *Z. jujuba* in Portugal.

*Septoria zizyphi* Sacc. Pale ochraceous to gray-white leaf spots on *Z. vulgaris* in Spain and Italy.

*Uredo belameandæ* P. Henn. Brown leaf rust on *Z. jujuba* in India.

*Uredo zizyphi* Pat. Brown leaf rust on *Z. spina-christi* and *Zizyphus* sp. in Indo-China, Abyssinia, and Brazil.

**ZOYSIA.** Low perennial grasses used for lawns and putting greens.

*Puccinia zoysiae* Diet. Powdery black rust pustules on leaves of *Z. pungens* in Japan.

**ZYGOPHYLLUM.** Shrubs, sometimes cultivated.

*Accidium kuntzii* Kaernb. and Wint. Leaf rust on *Z. atriplicoides* in Turkestan.

*Accidium lagena* Sorok. A rust deforming leaves of *Zygophyllum* sp. in Central Asia.

*Cercospora zygophylli* S. Szembel. On leaves of *Z. fabago* in Russia.

*Erysiphe taurica* Lév. See *Althaea*.

*Septoria zygophylli* Syd. Subcircular dull-brown, then gray-white, leaf spots with red-brown margins on *Z. fabago* in Germany.

*Uredo zygophylli* P. Henn. Brown rust pustules on leaves and stems of *Z. cornutum* and *Z. decumbens* in Algeria, Tunis, and Egypt.

*Uromyces trollipi* Kalchbr. and McO. Cinnamon-brown to black rust pustules on leaves of *Z. foetidum* in the Union of South Africa.

*Uromyces vesiculosus* Wint. Cinnamon-brown to black rust pustules on leaves, stems, and fruit of *Z. billardieri*, *Z. ammophilum*, and *Z. glaucescens* in Australia.













NATIONAL AGRICULTURAL LIBRARY



1022450896